

Computer Algebra Independent Integration Tests

Summer 2024

5-Inverse-trig-functions/5.3-Inverse-tangent/280-5.3.5

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 4:39am

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	47
3	Listing of integrals	50
3.1	$\int x^3 \arctan(a + bx) dx$	53
3.2	$\int x^2 \arctan(a + bx) dx$	60
3.3	$\int x \arctan(a + bx) dx$	67
3.4	$\int \arctan(a + bx) dx$	74
3.5	$\int \frac{\arctan(a+bx)}{x} dx$	79
3.6	$\int \frac{\arctan(a+bx)}{x^2} dx$	86
3.7	$\int \frac{\arctan(a+bx)}{x^3} dx$	93

3.8	$\int \frac{\arctan(ax+bx)}{x^4} dx$	100
3.9	$\int (ce+dex)^3(a+b\arctan(c+dx)) dx$	108
3.10	$\int (ce+dex)^2(a+b\arctan(c+dx)) dx$	117
3.11	$\int (ce+dex)(a+b\arctan(c+dx)) dx$	124
3.12	$\int \frac{a+b\arctan(c+dx)}{ce+dex} dx$	131
3.13	$\int \frac{a+b\arctan(c+dx)}{(ce+dex)^2} dx$	136
3.14	$\int \frac{a+b\arctan(c+dx)}{(ce+dex)^3} dx$	143
3.15	$\int (ce+dex)^3(a+b\arctan(c+dx))^2 dx$	150
3.16	$\int (ce+dex)^2(a+b\arctan(c+dx))^2 dx$	162
3.17	$\int (ce+dex)(a+b\arctan(c+dx))^2 dx$	171
3.18	$\int \frac{(a+b\arctan(c+dx))^2}{ce+dex} dx$	179
3.19	$\int \frac{(a+b\arctan(c+dx))^2}{(ce+dex)^2} dx$	186
3.20	$\int \frac{(a+b\arctan(c+dx))^2}{(ce+dex)^3} dx$	194
3.21	$\int \frac{(a+b\arctan(c+dx))^2}{(ce+dex)^4} dx$	203
3.22	$\int \frac{(a+b\arctan(c+dx))^2}{(ce+dex)^5} dx$	212
3.23	$\int (ce+dex)^2(a+b\arctan(c+dx))^3 dx$	224
3.24	$\int (ce+dex)(a+b\arctan(c+dx))^3 dx$	234
3.25	$\int \frac{(a+b\arctan(c+dx))^3}{ce+dex} dx$	243
3.26	$\int \frac{(a+b\arctan(c+dx))^3}{(ce+dex)^2} dx$	252
3.27	$\int \frac{(a+b\arctan(c+dx))^3}{(ce+dex)^3} dx$	260
3.28	$\int \frac{(a+b\arctan(c+dx))^3}{(ce+dex)^4} dx$	269
3.29	$\int \frac{\arctan(1+x)}{2+2x} dx$	280
3.30	$\int \frac{\arctan(ax+bx)}{\frac{ad}{b}+dx} dx$	285
3.31	$\int (a+bx)^2 \sqrt{\arctan(a+bx)} dx$	291
3.32	$\int (e+fx)^3(a+b\arctan(c+dx)) dx$	296
3.33	$\int (e+fx)^2(a+b\arctan(c+dx)) dx$	307
3.34	$\int (e+fx)(a+b\arctan(c+dx)) dx$	316
3.35	$\int (a+b\arctan(c+dx)) dx$	323
3.36	$\int \frac{a+b\arctan(c+dx)}{e+fx} dx$	328
3.37	$\int \frac{a+b\arctan(c+dx)}{(e+fx)^2} dx$	335
3.38	$\int \frac{a+b\arctan(c+dx)}{(e+fx)^3} dx$	344
3.39	$\int (e+fx)^2(a+b\arctan(c+dx))^2 dx$	354
3.40	$\int (e+fx)(a+b\arctan(c+dx))^2 dx$	363
3.41	$\int (a+b\arctan(c+dx))^2 dx$	370
3.42	$\int \frac{(a+b\arctan(c+dx))^2}{e+fx} dx$	377
3.43	$\int \frac{(a+b\arctan(c+dx))^2}{(e+fx)^2} dx$	383

3.44	$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx$	393
3.45	$\int (e + fx) (a + b \arctan(c + dx))^3 dx$	403
3.46	$\int (a + b \arctan(c + dx))^3 dx$	411
3.47	$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx$	419
3.48	$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx$	426
3.49	$\int (e + fx)^m (a + b \arctan(c + dx)) dx$	436
3.50	$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx$	443
3.51	$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx$	448
3.52	$\int \frac{\arctan(a + bx)}{c + dx^3} dx$	454
3.53	$\int \frac{\arctan(a + bx)}{c + dx^2} dx$	464
3.54	$\int \frac{\arctan(a + bx)}{c + dx} dx$	471
3.55	$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx$	478
3.56	$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx$	485
3.57	$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx$	494
3.58	$\int \frac{\arctan(a + bx)}{1 + x^2} dx$	503
3.59	$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx$	510
3.60	$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx$	517
3.61	$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx$	525
3.62	$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx$	535
3.63	$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx$	543
3.64	$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx$	548
3.65	$\int \frac{\arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx$	554
3.66	$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx$	559
3.67	$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx$	564
3.68	$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx$	573
3.69	$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx$	579
3.70	$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx$	586
3.71	$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx$	591

4	Appendix	597
4.1	Listing of Grading functions	597
4.2	Links to plain text integration problems used in this report for each CAS615	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [71]. This is test number [280].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	98.59 (70)	1.41 (1)
Maple	98.59 (70)	1.41 (1)
Mathematica	95.77 (68)	4.23 (3)
Maxima	50.70 (36)	49.30 (35)
Mupad	42.25 (30)	57.75 (41)
Reduce	40.85 (29)	59.15 (42)
Fricas	39.44 (28)	60.56 (43)
Giac	39.44 (28)	60.56 (43)
Sympy	33.80 (24)	66.20 (47)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	88.732	0.000	0.000	11.268
Mathematica	61.972	12.676	16.901	8.451
Maple	53.521	12.676	22.535	11.268
Fricas	23.944	7.042	0.000	69.014
Maxima	22.535	19.718	0.000	57.746
Giac	11.268	15.493	2.817	70.423
Sympy	7.042	4.225	15.493	73.239
Mupad	0.000	32.394	0.000	67.606
Reduce	0.000	30.986	0.000	69.014

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	3	100.00	0.00	0.00
Rubi	1	100.00	0.00	0.00
Maple	1	100.00	0.00	0.00
Maxima	35	94.29	0.00	5.71
Mupad	41	0.00	100.00	0.00
Fricas	43	97.67	0.00	2.33
Reduce	42	100.00	0.00	0.00
Giac	43	95.35	0.00	4.65
Sympy	47	44.68	55.32	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.16
Reduce	0.21
Rubi	0.71
Maxima	0.92
Maple	1.15
Mathematica	1.35
Mupad	1.76
Sympy	5.95
Giac	12.35

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Fricas	140.75	1.45	81.00	1.25
Mupad	178.37	1.75	102.50	1.37
Rubi	253.66	1.02	153.00	1.00
Sympy	253.71	2.66	172.50	2.20
Mathematica	288.56	1.45	164.00	1.11
Giac	378.21	2.92	118.00	1.29
Maxima	667.69	4.28	149.50	1.45
Maple	672.39	2.43	192.00	1.06
Reduce	2253.83	104.94	128.00	1.74

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{31, 50, 51, 65, 66, 70, 71}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {65, 66, 70, 71}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {10, 15, 67}

Mathematica {39, 43, 44, 53, 55, 62, 65, 66, 67, 70, 71}

Maple {18, 23, 25, 26, 28, 42, 44, 45, 47, 48, 52, 57, 61, 67}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	47

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 60, 61, 62, 63, 64, 67, 68, 69 }

B grade { }

C grade { }

F normal fail { 59 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 4, 5, 9, 10, 11, 12, 13, 15, 16, 17, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 30, 35, 36, 40, 41, 43, 45, 46, 49, 52, 53, 54, 56, 57, 58, 59, 60, 61, 62, 63, 64, 67, 68, 69 }

B grade { 18, 25, 39, 44, 55, 65, 66, 70, 71 }

C grade { 1, 2, 3, 6, 7, 8, 14, 32, 33, 34, 37, 38 }

F normal fail { 42, 47, 48 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 20, 22, 29, 30, 33, 34, 35, 36, 37, 38, 40, 41, 43, 54, 55, 56, 58, 63, 64, 68, 69 }

B grade { 19, 21, 24, 27, 32, 39, 46, 53, 62 }

C grade { 18, 23, 25, 26, 28, 42, 44, 45, 47, 48, 52, 57, 59, 60, 61, 67 }

F normal fail { 49 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 6, 7, 8, 11, 13, 14, 17, 20, 32, 33, 34, 35, 37 }

B grade { 9, 10, 15, 22, 38 }

C grade { }

F normal fail { 5, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 36, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { 31 }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 13, 32, 33, 34, 35, 37, 38, 55 }

B grade { 9, 10, 11, 14, 15, 17, 20, 22, 29, 30, 53, 54, 56, 58 }

C grade { }

F normal fail { 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 36, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 57, 59, 60, 61, 63, 64, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { 31, 62 }

Giac

A grade { 1, 2, 3, 4, 6, 7, 8, 35 }

B grade { 9, 10, 11, 13, 15, 17, 22, 32, 33, 34, 37 }

C grade { 14, 38 }

F normal fail { 5, 12, 16, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 36, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 62, 63, 64, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { 60, 61 }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 13, 14, 15, 17, 20, 22, 29, 32, 33, 34, 35, 37, 38 }

C grade { }

F normal fail { }

F(-1) timedout fail { 5, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 30, 36, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 67, 68, 69 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 35 }

B grade { 9, 11, 14 }

C grade { 6, 7, 8, 10, 13, 15, 17, 20, 32, 33, 34 }

F normal fail { 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 39, 40, 41, 44, 45, 46, 58, 63 }

F(-1) timedout fail { 5, 22, 36, 37, 38, 42, 43, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 62, 64, 67, 68, 69 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 13, 14, 15, 17, 20, 22, 32, 33, 34, 35, 37, 38 }

C grade { }

F normal fail { 5, 12, 16, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 36, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 67, 68, 69 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	99	95	131	104	87	155	103	122	133
N.S.	1	0.93	0.90	1.24	0.98	0.82	1.46	0.97	1.15	1.25
time (sec)	N/A	0.307	0.051	0.387	0.114	0.109	0.798	0.117	0.209	0.979

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	82	114	102	85	66	117	82	94	102
N.S.	1	1.04	1.44	1.29	1.08	0.84	1.48	1.04	1.19	1.29
time (sec)	N/A	0.297	0.036	0.089	0.117	0.102	0.605	0.117	0.202	1.868

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	59	90	63	68	52	78	60	60	61
N.S.	1	0.98	1.50	1.05	1.13	0.87	1.30	1.00	1.00	1.02
time (sec)	N/A	0.274	0.024	0.075	0.112	0.098	0.450	0.119	0.193	1.748

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	31	39	30	31	39	46	31	45	42
N.S.	1	0.94	1.18	0.91	0.94	1.18	1.39	0.94	1.36	1.27
time (sec)	N/A	0.204	0.011	0.055	0.029	0.105	0.216	0.121	0.188	0.099

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	171	94	134	0	0	0	12	0
N.S.	1	1.00	1.42	0.78	1.12	0.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.478	0.008	0.168	0.163	0.000	0.000	0.000	0.223	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	60	67	61	77	57	168	70	71	63
N.S.	1	0.97	1.08	0.98	1.24	0.92	2.71	1.13	1.15	1.02
time (sec)	N/A	0.241	0.044	0.096	0.115	0.120	0.785	0.123	0.193	1.433

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	102	92	84	112	95	382	115	126	232
N.S.	1	1.06	0.96	0.88	1.17	0.99	3.98	1.20	1.31	2.42
time (sec)	N/A	0.303	0.071	0.129	0.113	0.140	1.114	0.116	0.183	1.629

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	136	128	115	165	135	760	177	211	288
N.S.	1	1.05	0.99	0.89	1.28	1.05	5.89	1.37	1.64	2.23
time (sec)	N/A	0.336	0.100	0.144	0.106	0.128	1.470	0.126	0.197	1.455

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	59	56	64	370	151	231	253	164	371
N.S.	1	0.82	0.78	0.89	5.14	2.10	3.21	3.51	2.28	5.15
time (sec)	N/A	0.282	0.015	0.293	0.126	0.120	1.126	0.235	0.208	1.063

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	C	B	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	51	54	61	238	129	182	198	128	144
N.S.	1	0.76	0.81	0.91	3.55	1.93	2.72	2.96	1.91	2.15
time (sec)	N/A	0.275	0.013	0.181	0.119	0.115	1.174	0.190	0.192	1.662

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	45	40	51	120	60	95	121	71	73
N.S.	1	0.94	0.83	1.06	2.50	1.25	1.98	2.52	1.48	1.52
time (sec)	N/A	0.233	0.011	0.124	0.127	0.103	0.609	0.152	0.187	1.946

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	52	52	65	0	0	0	0	35	0
N.S.	1	0.83	0.83	1.03	0.00	0.00	0.00	0.00	0.56	0.00
time (sec)	N/A	0.286	0.014	0.251	0.000	0.000	0.000	0.000	0.192	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	53	52	58	92	75	223	175	106	88
N.S.	1	0.87	0.85	0.95	1.51	1.23	3.66	2.87	1.74	1.44
time (sec)	N/A	0.259	0.017	0.229	0.042	0.112	2.222	0.172	0.195	1.096

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	A	B	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	52	51	57	120	70	314	156	100	103
N.S.	1	0.83	0.81	0.90	1.90	1.11	4.98	2.48	1.59	1.63
time (sec)	N/A	0.254	0.016	0.404	0.112	0.103	3.750	0.150	0.178	1.164

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	C	B	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	136	216	152	597	337	583	610	421	633
N.S.	1	0.87	1.38	0.97	3.80	2.15	3.71	3.89	2.68	4.03
time (sec)	N/A	0.750	0.130	0.384	1.073	0.131	3.220	0.443	0.177	3.999

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	183	150	163	276	0	0	0	0	323	0
N.S.	1	0.82	0.89	1.51	0.00	0.00	0.00	0.00	1.77	0.00
time (sec)	N/A	0.789	0.335	0.829	0.000	0.000	0.000	0.000	0.193	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	87	107	113	218	150	240	240	191	216
N.S.	1	0.92	1.13	1.19	2.29	1.58	2.53	2.53	2.01	2.27
time (sec)	N/A	0.478	0.088	0.223	0.911	0.107	1.338	0.207	0.202	2.127

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	183	172	381	1154	0	0	0	0	62	0
N.S.	1	0.94	2.08	6.31	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.778	0.280	2.392	0.000	0.000	0.000	0.000	0.185	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	119	115	135	325	0	0	0	0	425	0
N.S.	1	0.97	1.13	2.73	0.00	0.00	0.00	0.00	3.57	0.00
time (sec)	N/A	0.555	0.252	0.918	0.000	0.000	0.000	0.000	0.188	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	98	194	129	268	209	1107	0	336	232
N.S.	1	0.84	1.66	1.10	2.29	1.79	9.46	0.00	2.87	1.98
time (sec)	N/A	0.568	0.191	0.450	0.209	0.124	9.218	0.000	0.195	3.492

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	163	163	368	0	0	0	0	1382	0
N.S.	1	0.84	0.84	1.90	0.00	0.00	0.00	0.00	7.12	0.00
time (sec)	N/A	0.807	0.616	2.451	0.000	0.000	0.000	0.000	0.210	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	156	245	164	534	448	0	836	679	438
N.S.	1	0.92	1.44	0.96	3.14	2.64	0.00	4.92	3.99	2.58
time (sec)	N/A	0.865	0.315	0.984	0.199	0.152	0.000	0.339	0.191	4.373

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	271	229	349	1277	0	0	0	0	592	0
N.S.	1	0.85	1.29	4.71	0.00	0.00	0.00	0.00	2.18	0.00
time (sec)	N/A	1.389	0.521	4.103	0.000	0.000	0.000	0.000	0.185	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	151	196	321	0	0	0	0	325	0
N.S.	1	0.92	1.20	1.96	0.00	0.00	0.00	0.00	1.98	0.00
time (sec)	N/A	0.876	0.164	0.648	0.000	0.000	0.000	0.000	0.194	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	279	258	562	2313	0	0	0	0	89	0
N.S.	1	0.92	2.01	8.29	0.00	0.00	0.00	0.00	0.32	0.00
time (sec)	N/A	1.077	0.498	1.713	0.000	0.000	0.000	0.000	0.206	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	163	158	263	2104	0	0	0	0	949	0
N.S.	1	0.97	1.61	12.91	0.00	0.00	0.00	0.00	5.82	0.00
time (sec)	N/A	0.816	0.498	1.648	0.000	0.000	0.000	0.000	0.195	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	180	163	225	423	0	0	0	0	975	0
N.S.	1	0.91	1.25	2.35	0.00	0.00	0.00	0.00	5.42	0.00
time (sec)	N/A	0.944	0.308	1.515	0.000	0.000	0.000	0.000	0.210	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	287	249	360	2446	0	0	0	0	0	0
N.S.	1	0.87	1.25	8.52	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.698	0.958	5.858	0.000	0.000	0.000	0.000	0.270	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	35	31	26	44	0	0	0	14	25
N.S.	1	1.13	1.00	0.84	1.42	0.00	0.00	0.00	0.45	0.81
time (sec)	N/A	0.265	0.005	0.114	0.168	0.000	0.000	0.000	0.185	0.081

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	39	34	38	123	0	0	0	21	0
N.S.	1	0.95	0.83	0.93	3.00	0.00	0.00	0.00	0.51	0.00
time (sec)	N/A	0.279	0.007	0.232	0.196	0.000	0.000	0.000	0.196	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	F(-2)	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	16	0	0	17	3	224	18
N.S.	1	1.00	1.11	0.89	0.00	0.00	0.94	0.17	12.44	1.00
time (sec)	N/A	0.189	3.037	0.136	0.000	0.000	10.913	121.043	0.196	1.334

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	A	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	233	230	157	464	346	317	654	885	494	787
N.S.	1	0.99	0.67	1.99	1.48	1.36	2.81	3.80	2.12	3.38
time (sec)	N/A	0.524	0.185	0.354	0.139	0.128	12.080	1.225	0.215	0.731

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	163	118	265	220	199	376	512	293	411
N.S.	1	1.05	0.76	1.71	1.42	1.28	2.43	3.30	1.89	2.65
time (sec)	N/A	0.427	0.104	0.198	0.132	0.116	4.716	0.451	0.200	1.177

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	103	163	113	116	103	177	250	137	136
N.S.	1	1.10	1.73	1.20	1.23	1.10	1.88	2.66	1.46	1.45
time (sec)	N/A	0.335	0.045	0.131	0.131	0.109	1.945	0.200	0.181	1.609

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	49	35	36	48	51	36	53	49
N.S.	1	1.00	1.29	0.92	0.95	1.26	1.34	0.95	1.39	1.29
time (sec)	N/A	0.177	0.010	0.059	0.033	0.097	0.178	0.126	0.183	1.543

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	162	180	160	198	0	0	0	0	32	0
N.S.	1	1.11	0.99	1.22	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.545	0.079	0.277	0.000	0.000	0.000	0.000	0.197	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	148	121	160	177	190	0	3974	248	127
N.S.	1	0.98	0.80	1.06	1.17	1.26	0.00	26.32	1.64	0.84
time (sec)	N/A	0.432	0.153	0.270	0.127	0.254	0.000	0.352	0.184	2.385

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-1)	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	227	245	175	245	409	682	0	1561	1110	399
N.S.	1	1.08	0.77	1.08	1.80	3.00	0.00	6.88	4.89	1.76
time (sec)	N/A	0.594	0.565	0.595	0.127	1.218	0.000	3.587	0.186	8.601

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	382	383	801	1018	0	0	0	0	771	0
N.S.	1	1.00	2.10	2.66	0.00	0.00	0.00	0.00	2.02	0.00
time (sec)	N/A	0.749	7.632	0.670	0.000	0.000	0.000	0.000	0.198	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	222	228	264	413	0	0	0	0	345	0
N.S.	1	1.03	1.19	1.86	0.00	0.00	0.00	0.00	1.55	0.00
time (sec)	N/A	0.608	0.857	0.397	0.000	0.000	0.000	0.000	0.216	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	109	139	0	0	0	0	107	0
N.S.	1	1.00	1.07	1.36	0.00	0.00	0.00	0.00	1.05	0.00
time (sec)	N/A	0.486	0.075	0.303	0.000	0.000	0.000	0.000	0.189	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	261	288	0	1877	0	0	0	0	59	0
N.S.	1	1.10	0.00	7.19	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.427	0.000	4.290	0.000	0.000	0.000	0.000	0.200	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	480	586	419	776	0	0	0	0	0	0
N.S.	1	1.22	0.87	1.62	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.568	4.545	1.441	0.000	0.000	0.000	0.000	0.335	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	564	554	1844	5843	0	0	0	0	1362	0
N.S.	1	0.98	3.27	10.36	0.00	0.00	0.00	0.00	2.41	0.00
time (sec)	N/A	1.132	13.515	12.274	0.000	0.000	0.000	0.000	0.394	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	337	334	592	8296	0	0	0	0	575	0
N.S.	1	0.99	1.76	24.62	0.00	0.00	0.00	0.00	1.71	0.00
time (sec)	N/A	0.860	4.564	3.860	0.000	0.000	0.000	0.000	0.195	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	141	212	279	0	0	0	0	172	0
N.S.	1	0.99	1.48	1.95	0.00	0.00	0.00	0.00	1.20	0.00
time (sec)	N/A	0.680	0.106	0.822	0.000	0.000	0.000	0.000	0.206	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	372	408	0	3817	0	0	0	0	86	0
N.S.	1	1.10	0.00	10.26	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.471	0.000	4.720	0.000	0.000	0.000	0.000	0.221	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	693	1265	0	3708	0	0	0	0	0	0
N.S.	1	1.83	0.00	5.35	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.353	0.000	5.973	0.000	0.000	0.000	0.000	0.747	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	177	237	162	0	0	0	0	0	1309	0
N.S.	1	1.34	0.92	0.00	0.00	0.00	0.00	0.00	7.40	0.00
time (sec)	N/A	0.476	0.274	0.000	0.000	0.000	0.000	0.000	0.202	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	504	36	0	3	17260	22
N.S.	1	1.00	1.10	1.00	25.20	1.80	0.00	0.15	863.00	1.10
time (sec)	N/A	0.293	3.867	0.197	9.070	0.096	0.000	108.316	0.354	0.846

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	659	52	0	3	42351	22
N.S.	1	1.00	1.10	1.00	32.95	2.60	0.00	0.15	2117.55	1.10
time (sec)	N/A	0.303	0.454	0.210	12.192	0.111	0.000	107.155	0.527	0.853

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	725	869	701	380	0	0	0	0	18	0
N.S.	1	1.20	0.97	0.52	0.00	0.00	0.00	0.00	0.02	0.00
time (sec)	N/A	1.768	0.593	0.899	0.000	0.000	0.000	0.000	44.469	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	319	541	409	542	8520	0	0	0	18	0
N.S.	1	1.70	1.28	1.70	26.71	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.974	0.234	1.138	4.489	0.000	0.000	0.000	0.192	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	170	231	186	284	0	0	0	16	0
N.S.	1	1.12	1.52	1.22	1.87	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.529	0.016	0.339	0.274	0.000	0.000	0.000	0.192	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	244	254	771	295	284	0	0	0	17	0
N.S.	1	1.04	3.16	1.21	1.16	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.537	7.959	0.348	0.230	0.000	0.000	0.000	0.190	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	668	678	660	647	8518	0	0	0	21	0
N.S.	1	1.01	0.99	0.97	12.75	0.00	0.00	0.00	0.03	0.00
time (sec)	N/A	1.211	0.420	2.136	0.805	0.000	0.000	0.000	0.211	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	933	951	896	511	0	0	0	0	21	0
N.S.	1	1.02	0.96	0.55	0.00	0.00	0.00	0.00	0.02	0.00
time (sec)	N/A	1.791	0.869	1.093	0.000	0.000	0.000	0.000	0.192	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	179	304	283	226	328	0	0	0	16	0
N.S.	1	1.70	1.58	1.26	1.83	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	0.568	0.039	0.854	0.181	0.000	0.000	0.000	0.187	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	706	0	631	372	0	0	0	0	78	0
N.S.	1	0.00	0.89	0.53	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.000	22.834	0.272	0.000	0.000	0.000	0.000	0.317	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	673	665	604	364	0	0	0	0	50	0
N.S.	1	0.99	0.90	0.54	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	1.219	0.359	0.165	0.000	0.000	0.000	0.000	0.267	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	770	792	770	388	0	0	0	0	53	0
N.S.	1	1.03	1.00	0.50	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	1.383	0.331	0.170	0.000	0.000	0.000	0.000	0.266	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	398	427	494	967	0	0	0	0	1038	0
N.S.	1	1.07	1.24	2.43	0.00	0.00	0.00	0.00	2.61	0.00
time (sec)	N/A	1.230	0.542	2.076	0.000	0.000	0.000	0.000	0.245	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	127	67	135	0	0	0	0	29	0
N.S.	1	0.96	0.51	1.02	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.289	0.068	0.375	0.000	0.000	0.000	0.000	0.202	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	216	155	95	176	0	0	0	0	34	0
N.S.	1	0.72	0.44	0.81	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.374	0.078	0.494	0.000	0.000	0.000	0.000	0.181	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	B	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	163	26	28	28	29	28	28	28
N.S.	1	1.00	5.82	0.93	1.00	1.00	1.04	1.00	1.00	1.00
time (sec)	N/A	0.252	0.292	0.194	0.138	0.104	0.924	0.135	0.180	0.860

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	B	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	165	31	33	33	31	33	32	33
N.S.	1	1.00	5.00	0.94	1.00	1.00	0.94	1.00	0.97	1.00
time (sec)	N/A	0.273	0.146	0.154	0.141	0.100	7.692	0.144	0.219	0.884

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	1199	2244	1807	981	0	0	0	0	419	0
N.S.	1	1.87	1.51	0.82	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	4.778	5.249	3.033	0.000	0.000	0.000	0.000	0.365	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	187	177	145	180	0	0	0	0	104	0
N.S.	1	0.95	0.78	0.96	0.00	0.00	0.00	0.00	0.56	0.00
time (sec)	N/A	0.440	0.521	1.060	0.000	0.000	0.000	0.000	0.190	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	281	215	189	222	0	0	0	0	109	0
N.S.	1	0.77	0.67	0.79	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.603	0.213	0.865	0.000	0.000	0.000	0.000	0.216	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	B	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	181	33	35	44	36	35	101	35
N.S.	1	1.00	5.17	0.94	1.00	1.26	1.03	1.00	2.89	1.00
time (sec)	N/A	0.341	4.384	0.313	0.335	0.122	3.869	0.154	0.229	0.932

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	B	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	225	38	40	49	37	40	105	40
N.S.	1	1.00	5.62	0.95	1.00	1.22	0.92	1.00	2.62	1.00
time (sec)	N/A	0.402	0.693	0.295	0.352	0.130	72.286	0.170	0.226	0.940

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [3] had the largest ratio of [.750000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	6	0.93	10	0.600
2	A	6	5	1.04	10	0.500
3	A	7	6	0.98	8	0.750
4	A	4	3	0.94	6	0.500
5	A	8	7	1.00	10	0.700
6	A	8	7	0.97	10	0.700
7	A	6	5	1.06	10	0.500
8	A	7	6	1.05	10	0.600
9	A	6	5	0.82	21	0.238
10	A	7	6	0.76	21	0.286
11	A	6	5	0.94	19	0.263
12	A	5	4	0.83	21	0.190
13	A	8	7	0.87	21	0.333
14	A	6	5	0.83	21	0.238
15	A	12	11	0.87	23	0.478
16	A	12	11	0.82	23	0.478
17	A	7	6	0.92	21	0.286
18	A	7	6	0.94	23	0.261
19	A	7	6	0.97	23	0.261
20	A	11	10	0.84	23	0.435
21	A	11	10	0.84	23	0.435

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	16	15	0.92	23	0.652
23	A	13	12	0.85	23	0.522
24	A	11	10	0.92	21	0.476
25	A	8	7	0.92	23	0.304
26	A	8	7	0.97	23	0.304
27	A	10	9	0.91	23	0.391
28	A	17	16	0.87	23	0.696
29	A	5	4	1.13	12	0.333
30	A	5	4	0.95	19	0.211
31	N/A	1	0	1.00	18	0.000
32	A	6	5	0.99	18	0.278
33	A	6	5	1.05	18	0.278
34	A	6	5	1.10	16	0.312
35	A	1	1	1.00	10	0.100
36	A	7	6	1.11	18	0.333
37	A	10	9	0.98	18	0.500
38	A	6	6	1.08	18	0.333
39	A	5	4	1.00	20	0.200
40	A	5	4	1.03	18	0.222
41	A	7	6	1.00	12	0.500
42	A	4	3	1.10	20	0.150
43	A	7	6	1.22	20	0.300
44	A	5	4	0.98	20	0.200
45	A	5	4	0.99	18	0.222
46	A	7	6	0.99	12	0.500
47	A	4	3	1.10	20	0.150
48	A	7	6	1.83	20	0.300
49	A	5	4	1.34	18	0.222
50	N/A	3	0	1.00	20	0.000
51	N/A	3	0	1.00	20	0.000
52	A	3	3	1.20	16	0.188
53	A	3	3	1.70	16	0.188

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	7	6	1.12	14	0.429
55	A	3	3	1.04	16	0.188
56	A	3	3	1.01	16	0.188
57	A	3	3	1.02	16	0.188
58	A	3	3	1.70	14	0.214
59	F	0	0	N/A	0.000	N/A
60	A	5	4	0.99	18	0.222
61	A	6	5	1.03	18	0.278
62	A	2	2	1.07	23	0.087
63	A	3	2	0.96	28	0.071
64	A	4	3	0.72	33	0.091
65	N/A	3	0	1.00	28	0.000
66	N/A	3	0	1.00	33	0.000
67	A	2	2	1.87	25	0.080
68	A	5	4	0.95	35	0.114
69	A	6	5	0.77	40	0.125
70	N/A	3	0	1.00	35	0.000
71	N/A	3	0	1.00	40	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \arctan(a + bx) dx$	53
3.2	$\int x^2 \arctan(a + bx) dx$	60
3.3	$\int x \arctan(a + bx) dx$	67
3.4	$\int \arctan(a + bx) dx$	74
3.5	$\int \frac{\arctan(a+bx)}{x} dx$	79
3.6	$\int \frac{\arctan(a+bx)}{x^2} dx$	86
3.7	$\int \frac{\arctan(a+bx)}{x^3} dx$	93
3.8	$\int \frac{\arctan(a+bx)}{x^4} dx$	100
3.9	$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$	108
3.10	$\int (ce + dex)^2 (a + b \arctan(c + dx)) dx$	117
3.11	$\int (ce + dex) (a + b \arctan(c + dx)) dx$	124
3.12	$\int \frac{a+b \arctan(c+dx)}{ce+dex} dx$	131
3.13	$\int \frac{a+b \arctan(c+dx)}{(ce+dex)^2} dx$	136
3.14	$\int \frac{a+b \arctan(c+dx)}{(ce+dex)^3} dx$	143
3.15	$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$	150
3.16	$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx$	162
3.17	$\int (ce + dex) (a + b \arctan(c + dx))^2 dx$	171
3.18	$\int \frac{(a+b \arctan(c+dx))^2}{ce+dex} dx$	179
3.19	$\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^2} dx$	186
3.20	$\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^3} dx$	194
3.21	$\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^4} dx$	203
3.22	$\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^5} dx$	212
3.23	$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx$	224
3.24	$\int (ce + dex) (a + b \arctan(c + dx))^3 dx$	234
3.25	$\int \frac{(a+b \arctan(c+dx))^3}{ce+dex} dx$	243

3.26	$\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^2} dx$	252
3.27	$\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^3} dx$	260
3.28	$\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^4} dx$	269
3.29	$\int \frac{\arctan(1+x)}{2+2x} dx$	280
3.30	$\int \frac{\arctan(a+bx)}{\frac{ad}{b}+dx} dx$	285
3.31	$\int (a+bx)^2 \sqrt{\arctan(a+bx)} dx$	291
3.32	$\int (e+fx)^3 (a+b \arctan(c+dx)) dx$	296
3.33	$\int (e+fx)^2 (a+b \arctan(c+dx)) dx$	307
3.34	$\int (e+fx) (a+b \arctan(c+dx)) dx$	316
3.35	$\int (a+b \arctan(c+dx)) dx$	323
3.36	$\int \frac{a+b \arctan(c+dx)}{e+fx} dx$	328
3.37	$\int \frac{a+b \arctan(c+dx)}{(e+fx)^2} dx$	335
3.38	$\int \frac{a+b \arctan(c+dx)}{(e+fx)^3} dx$	344
3.39	$\int (e+fx)^2 (a+b \arctan(c+dx))^2 dx$	354
3.40	$\int (e+fx) (a+b \arctan(c+dx))^2 dx$	363
3.41	$\int (a+b \arctan(c+dx))^2 dx$	370
3.42	$\int \frac{(a+b \arctan(c+dx))^2}{e+fx} dx$	377
3.43	$\int \frac{(a+b \arctan(c+dx))^2}{(e+fx)^2} dx$	383
3.44	$\int (e+fx)^2 (a+b \arctan(c+dx))^3 dx$	393
3.45	$\int (e+fx) (a+b \arctan(c+dx))^3 dx$	403
3.46	$\int (a+b \arctan(c+dx))^3 dx$	411
3.47	$\int \frac{(a+b \arctan(c+dx))^3}{e+fx} dx$	419
3.48	$\int \frac{(a+b \arctan(c+dx))^3}{(e+fx)^2} dx$	426
3.49	$\int (e+fx)^m (a+b \arctan(c+dx)) dx$	436
3.50	$\int (e+fx)^m (a+b \arctan(c+dx))^2 dx$	443
3.51	$\int (e+fx)^m (a+b \arctan(c+dx))^3 dx$	448
3.52	$\int \frac{\arctan(a+bx)}{c+dx^3} dx$	454
3.53	$\int \frac{\arctan(a+bx)}{c+dx^2} dx$	464
3.54	$\int \frac{\arctan(a+bx)}{c+dx} dx$	471
3.55	$\int \frac{\arctan(a+bx)}{c+\frac{d}{x}} dx$	478
3.56	$\int \frac{\arctan(a+bx)}{c+\frac{d}{x^2}} dx$	485
3.57	$\int \frac{\arctan(a+bx)}{c+\frac{d}{x^3}} dx$	494
3.58	$\int \frac{\arctan(a+bx)}{1+x^2} dx$	503
3.59	$\int \frac{a+b \arctan(c+dx)}{e+f\sqrt{x}} dx$	510
3.60	$\int \frac{\arctan(a+bx)}{c+d\sqrt{x}} dx$	517

3.61	$\int \frac{\arctan(a+bx)}{c+\frac{d}{\sqrt{x}}} dx$	525
3.62	$\int \frac{a+b \arctan(c+dx)}{e+fx+gx^2} dx$	535
3.63	$\int \frac{\arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx$	543
3.64	$\int \frac{\arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx$	548
3.65	$\int \frac{\arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx$	554
3.66	$\int \frac{\arctan(a+bx)}{\sqrt[3]{(1+a^2)c+2abcx+b^2cx^2}} dx$	559
3.67	$\int \frac{a+b \arctan(c+dx)}{e+fx^2+gx^4} dx$	564
3.68	$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx$	573
3.69	$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx$	579
3.70	$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx$	586
3.71	$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{(1+a^2)c+2abcx+b^2cx^2}} dx$	591

3.1 $\int x^3 \arctan(a + bx) dx$

Optimal result	53
Mathematica [C] (verified)	53
Rubi [A] (verified)	54
Maple [A] (verified)	56
Fricas [A] (verification not implemented)	57
Sympy [A] (verification not implemented)	57
Maxima [A] (verification not implemented)	58
Giac [A] (verification not implemented)	58
Mupad [B] (verification not implemented)	59
Reduce [B] (verification not implemented)	59

Optimal result

Integrand size = 10, antiderivative size = 106

$$\int x^3 \arctan(a + bx) dx = \frac{(1 - 6a^2)x}{4b^3} + \frac{a(a + bx)^2}{2b^4} - \frac{(a + bx)^3}{12b^4} - \frac{(1 - 6a^2 + a^4) \arctan(a + bx)}{4b^4} + \frac{1}{4}x^4 \arctan(a + bx) - \frac{a(1 - a^2) \log(1 + (a + bx)^2)}{2b^4}$$

output

```
1/4*(-6*a^2+1)*x/b^3+1/2*a*(b*x+a)^2/b^4-1/12*(b*x+a)^3/b^4-1/4*(a^4-6*a^2+1)*arctan(b*x+a)/b^4+1/4*x^4*arctan(b*x+a)-1/2*a*(-a^2+1)*ln(1+(b*x+a)^2)/b^4
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.90

$$\int x^3 \arctan(a + bx) dx = \frac{6(1 - 6a^2)bx + 12a(a + bx)^2 - 2(a + bx)^3 + 6b^4x^4 \arctan(a + bx) + 3i(-i + a)^4 \log(i - a - bx) - 3i(i - a - bx)^4}{24b^4}$$

input `Integrate[x^3*ArcTan[a + b*x],x]`

output $(6*(1 - 6*a^2)*b*x + 12*a*(a + b*x)^2 - 2*(a + b*x)^3 + 6*b^4*x^4*ArcTan[a + b*x] + (3*I)*(-I + a)^4*Log[I - a - b*x] - (3*I)*(I + a)^4*Log[I + a + b*x])/(24*b^4)$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5570, 25, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \arctan(a + bx) dx \\
 & \quad \downarrow \text{5570} \\
 & \frac{\int x^3 \arctan(a + bx) d(a + bx)}{b} \\
 & \quad \downarrow \text{25} \\
 & - \frac{\int -x^3 \arctan(a + bx) d(a + bx)}{b} \\
 & \quad \downarrow \text{27} \\
 & - \frac{\int -b^3 x^3 \arctan(a + bx) d(a + bx)}{b^4} \\
 & \quad \downarrow \text{5387} \\
 & - \frac{\frac{1}{4} \int \frac{b^4 x^4}{(a+bx)^2+1} d(a + bx) - \frac{1}{4} b^4 x^4 \arctan(a + bx)}{b^4} \\
 & \quad \downarrow \text{478} \\
 & - \frac{\frac{1}{4} \int \left(6a^2 - 4(a + bx)a + (a + bx)^2 + \frac{a^4 - 6a^2 + 4(1-a^2)(a+bx)a+1}{(a+bx)^2+1} - 1 \right) d(a + bx) - \frac{1}{4} b^4 x^4 \arctan(a + bx)}{b^4}
 \end{aligned}$$

↓ 2009

$$-\frac{\frac{1}{4}(-1-6a^2)(a+bx) + 2a(1-a^2)\log((a+bx)^2+1) + (a^4-6a^2+1)\arctan(a+bx) + \frac{1}{3}(a+bx)^3 - 2a(a+bx)}{b^4}$$

input `Int[x^3*ArcTan[a + b*x],x]`

output `-((-1/4*(b^4*x^4*ArcTan[a + b*x]) + (-((1 - 6*a^2)*(a + b*x)) - 2*a*(a + b*x)^2 + (a + b*x)^3/3 + (1 - 6*a^2 + a^4)*ArcTan[a + b*x] + 2*a*(1 - a^2)*Log[1 + (a + b*x)^2])/4)/b^4)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 478 `Int[((c_) + (d_.)*(x_))^(n_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5387 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.)*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 5570

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^(m*(a + b*ArcTan[x]))^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.24

method	result
parallelrisch	$\frac{3 \arctan(bx+a)x^4b^4 - b^3x^3 + 3ab^2x^2 - 3 \arctan(bx+a)a^4 + 6a^3 \ln(b^2x^2 + 2abx + a^2 + 1) - 9a^2bx + 18 \arctan(bx+a)a^2 + 15a}{12b^4}$
parts	$\frac{x^4 \arctan(bx+a)}{4} - \frac{b \left(\frac{\frac{1}{3}b^2x^3 - abx^2 + 3a^2x - x}{b^4} + \frac{(-4a^3b + 4ab) \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} + \frac{(-3a^4 - 2a^2 + 1 - \frac{(-4a^3b + 4ab)a}{b})}{b^4} \right)}{4}$
derivativedivides	$\frac{\arctan(bx+a)a^4}{4} - \arctan(bx+a)a^3(bx+a) + \frac{3 \arctan(bx+a)a^2(bx+a)^2}{2} - \arctan(bx+a)a(bx+a)^3 + \frac{\arctan(bx+a)(bx+a)^4}{4} - 3a}{b^4}$
default	$\frac{\arctan(bx+a)a^4}{4} - \arctan(bx+a)a^3(bx+a) + \frac{3 \arctan(bx+a)a^2(bx+a)^2}{2} - \arctan(bx+a)a(bx+a)^3 + \frac{\arctan(bx+a)(bx+a)^4}{4} - 3a}{b^4}$
risch	$-\frac{ix^4 \ln(1+i(bx+a))}{8} + \frac{ix^4 \ln(1-i(bx+a))}{8} - \frac{x^3}{12b} - \frac{a^4 \arctan(bx+a)}{4b^4} + \frac{ax^2}{4b^2} + \frac{a^3 \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^4} -$

input

```
int(x^3*arctan(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
1/12*(3*arctan(b*x+a)*x^4*b^4 - b^3*x^3 + 3*a*b^2*x^2 - 3*arctan(b*x+a)*a^4 + 6*a^3*ln(b^2*x^2 + 2*a*b*x + a^2 + 1) - 9*a^2*b*x + 18*arctan(b*x+a)*a^2 + 15*a^3 - 6*a*ln(b^2*x^2 + 2*a*b*x + a^2 + 1) + 3*b*x - 3*arctan(b*x+a) - 9*a)/b^4
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.82

$$\int x^3 \arctan(a + bx) dx = \frac{b^3 x^3 - 3ab^2 x^2 + 3(3a^2 - 1)bx - 3(b^4 x^4 - a^4 + 6a^2 - 1) \arctan(bx + a) - 6(a^3 - a) \log(b^2 x^2 + 2abx + a^2 + 1)}{12b^4}$$

input `integrate(x^3*arctan(b*x+a),x, algorithm="fricas")`output `-1/12*(b^3*x^3 - 3*a*b^2*x^2 + 3*(3*a^2 - 1)*b*x - 3*(b^4*x^4 - a^4 + 6*a^2 - 1)*arctan(b*x + a) - 6*(a^3 - a)*log(b^2*x^2 + 2*a*b*x + a^2 + 1))/b^4`**Sympy [A] (verification not implemented)**

Time = 0.80 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.46

$$\int x^3 \arctan(a + bx) dx = \begin{cases} -\frac{a^4 \operatorname{atan}(a+bx)}{4b^4} + \frac{a^3 \log(a^2+2abx+b^2x^2+1)}{2b^4} - \frac{3a^2x}{4b^3} + \frac{3a^2 \operatorname{atan}(a+bx)}{2b^4} + \frac{ax^2}{4b^2} - \frac{a \log(a^2+2abx+b^2x^2+1)}{2b^4} + \frac{x^4 \operatorname{atan}(a+bx)}{4} \\ \frac{x^4 \operatorname{atan}(a)}{4} \end{cases}$$

input `integrate(x**3*atan(b*x+a),x)`output `Piecewise((-a**4*atan(a + b*x)/(4*b**4) + a**3*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*b**4) - 3*a**2*x/(4*b**3) + 3*a**2*atan(a + b*x)/(2*b**4) + a*x**2/(4*b**2) - a*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*b**4) + x**4*atan(a + b*x)/4 - x**3/(12*b) + x/(4*b**3) - atan(a + b*x)/(4*b**4), Ne(b, 0)), (x**4*atan(a)/4, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.98

$$\int x^3 \arctan(a + bx) dx = \frac{1}{4} x^4 \arctan(bx + a) - \frac{1}{12} b \left(\frac{b^2 x^3 - 3 abx^2 + 3(3a^2 - 1)x}{b^4} + \frac{3(a^4 - 6a^2 + 1) \arctan\left(\frac{b^2 x + ab}{b}\right)}{b^5} - \frac{6(a^3 - a) \log(b^2 x^2 + 2 abx + a^2 + 1)}{b^5} \right)$$

input `integrate(x^3*arctan(b*x+a),x, algorithm="maxima")`output `1/4*x^4*arctan(b*x + a) - 1/12*b*((b^2*x^3 - 3*a*b*x^2 + 3*(3*a^2 - 1)*x)/b^4 + 3*(a^4 - 6*a^2 + 1)*arctan((b^2*x + a*b)/b)/b^5 - 6*(a^3 - a)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^5)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.97

$$\int x^3 \arctan(a + bx) dx = \frac{1}{4} x^4 \arctan(bx + a) - \frac{1}{12} b \left(\frac{3(a^4 - 6a^2 + 1) \arctan(bx + a)}{b^5} - \frac{6(a^3 - a) \log(b^2 x^2 + 2 abx + a^2 + 1)}{b^5} + \frac{b^4 x^3 - 3 ab^3 x^2 + 9 a^2 b^2 x - 3 b^2 x}{b^6} \right)$$

input `integrate(x^3*arctan(b*x+a),x, algorithm="giac")`output `1/4*x^4*arctan(b*x + a) - 1/12*b*(3*(a^4 - 6*a^2 + 1)*arctan(b*x + a)/b^5 - 6*(a^3 - a)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^5 + (b^4*x^3 - 3*a*b^3*x^2 + 9*a^2*b^2*x - 3*b^2*x)/b^6)`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.25

$$\int x^3 \arctan(a + bx) dx = \frac{x^4 \operatorname{atan}(a + bx)}{4} - \frac{\operatorname{atan}(a + bx)}{4b^4} + \frac{x}{4b^3} - \frac{x^3}{12b} + \frac{a^3 \ln(a^2 + 2abx + b^2x^2 + 1)}{2b^4} + \frac{3a^2 \operatorname{atan}(a + bx)}{2b^4} - \frac{a^4 \operatorname{atan}(a + bx)}{4b^4} + \frac{ax^2}{4b^2} - \frac{3a^2x}{4b^3} - \frac{a \ln(a^2 + 2abx + b^2x^2 + 1)}{2b^4}$$

input `int(x^3*atan(a + b*x),x)`output $(x^4 \operatorname{atan}(a + bx))/4 - \operatorname{atan}(a + bx)/(4b^4) + x/(4b^3) - x^3/(12b) + (a^3 \log(a^2 + b^2x^2 + 2abx + 1))/(2b^4) + (3a^2 \operatorname{atan}(a + bx))/(2b^4) - (a^4 \operatorname{atan}(a + bx))/(4b^4) + (ax^2)/(4b^2) - (3a^2x)/(4b^3) - (a \log(a^2 + b^2x^2 + 2abx + 1))/(2b^4)$ **Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.15

$$\int x^3 \arctan(a + bx) dx = \frac{-3 \operatorname{atan}(bx + a) a^4 + 18 \operatorname{atan}(bx + a) a^2 + 3 \operatorname{atan}(bx + a) b^4 x^4 - 3 \operatorname{atan}(bx + a) + 6 \log(b^2 x^2 + 2abx + a^2)}{12b^4}$$

input `int(x^3*atan(b*x+a),x)`output $(-3 \operatorname{atan}(a + bx) a^4 + 18 \operatorname{atan}(a + bx) a^2 + 3 \operatorname{atan}(a + bx) b^4 x^4 - 3 \operatorname{atan}(a + bx) + 6 \log(a^2 + 2abx + b^2x^2 + 1) a^3 - 6 \log(a^2 + 2abx + b^2x^2 + 1) a - 9 a^2 b x + 3 a b^2 x^2 - b^3 x^3 + 3 b x)/(12 b^4)$

3.2 $\int x^2 \arctan(a + bx) dx$

Optimal result	60
Mathematica [C] (verified)	60
Rubi [A] (verified)	61
Maple [A] (verified)	63
Fricas [A] (verification not implemented)	63
Sympy [A] (verification not implemented)	64
Maxima [A] (verification not implemented)	64
Giac [A] (verification not implemented)	65
Mupad [B] (verification not implemented)	65
Reduce [B] (verification not implemented)	66

Optimal result

Integrand size = 10, antiderivative size = 79

$$\int x^2 \arctan(a + bx) dx = \frac{ax}{b^2} - \frac{(a + bx)^2}{6b^3} - \frac{a(3 - a^2) \arctan(a + bx)}{3b^3} + \frac{1}{3}x^3 \arctan(a + bx) + \frac{(1 - 3a^2) \log(1 + (a + bx)^2)}{6b^3}$$

output

```
a*x/b^2-1/6*(b*x+a)^2/b^3-1/3*a*(-a^2+3)*arctan(b*x+a)/b^3+1/3*x^3*arctan(b*x+a)+1/6*(-3*a^2+1)*ln(1+(b*x+a)^2)/b^3
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.44

$$\int x^2 \arctan(a + bx) dx = \frac{\frac{1}{3}b\left(-\frac{a}{b} + \frac{a+bx}{b}\right)^3 \arctan(a + bx) - \frac{1}{3}b\left(-\frac{3ax}{b^2} + \frac{(a+bx)^2}{2b^3} - \frac{(1+ia)^3 \log(i-a-bx)}{2b^3} - \frac{(1-ia)^3 \log(i+a+bx)}{2b^3}\right)}{b}$$

input

```
Integrate[x^2*ArcTan[a + b*x],x]
```

output

$$\frac{((b*(-a/b) + (a + b*x)/b)^3 \text{ArcTan}[a + b*x])/3 - (b*((-3*a*x)/b^2 + (a + b*x)^2/(2*b^3) - ((1 + I*a)^3 \text{Log}[I - a - b*x])/(2*b^3) - ((1 - I*a)^3 \text{Log}[I + a + b*x])/(2*b^3)))/3)/b}$$
Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5570, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \arctan(a + bx) dx \\ & \quad \downarrow \text{5570} \\ & \frac{\int x^2 \arctan(a + bx) d(a + bx)}{b} \\ & \quad \downarrow \text{27} \\ & \frac{\int b^2 x^2 \arctan(a + bx) d(a + bx)}{b^3} \\ & \quad \downarrow \text{5387} \\ & \frac{\frac{1}{3} \int -\frac{b^3 x^3}{(a+bx)^2+1} d(a + bx) + \frac{1}{3} b^3 x^3 \arctan(a + bx)}{b^3} \\ & \quad \downarrow \text{478} \\ & \frac{\frac{1}{3} \int \left(2a - bx - \frac{a(3-a^2) - (1-3a^2)(a+bx)}{(a+bx)^2+1} \right) d(a + bx) + \frac{1}{3} b^3 x^3 \arctan(a + bx)}{b^3} \\ & \quad \downarrow \text{2009} \\ & \frac{\frac{1}{3}(-a(3-a^2) \arctan(a + bx) + \frac{1}{2}(1-3a^2) \log((a + bx)^2 + 1) - \frac{1}{2}(a + bx)^2 + 3a(a + bx)) + \frac{1}{3} b^3 x^3 \arctan(a + bx)}{b^3} \end{aligned}$$

input

$$\text{Int}[x^2 \text{ArcTan}[a + b*x], x]$$

output
$$\frac{((b^3 x^3 \operatorname{ArcTan}[a + b x])/3 + (3 a (a + b x) - (a + b x)^2/2 - a(3 - a^2)) \operatorname{ArcTan}[a + b x] + ((1 - 3 a^2) \operatorname{Log}[1 + (a + b x)^2])/2)/3}{b^3}$$

Defintions of rubi rules used

rule 27
$$\operatorname{Int}[(a_*)(F x_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[F x, (b_*)(G x_)] /; \operatorname{FreeQ}[b, x]$$

rule 478
$$\operatorname{Int}[((c_*) + (d_*)(x_))^{(n_*)}/((a_*) + (b_*)(x_)^2), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(c + d x)^n/(a + b x^2), x], x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \operatorname{IGtQ}[n, 1]$$

rule 2009
$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 5387
$$\operatorname{Int}[((a_*) + \operatorname{ArcTan}[(c_*)(x_)]*(b_*)) * ((d_*) + (e_*)(x_))^{(q_*)}, x_Symbol] \rightarrow \operatorname{Simp}[(d + e x)^{(q + 1)} * ((a + b \operatorname{ArcTan}[c x]) / (e (q + 1))), x] - \operatorname{Simp}[b * (c / (e (q + 1))) \operatorname{Int}[(d + e x)^{(q + 1)} / (1 + c^2 x^2), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, q\}, x] \ \&\& \ \operatorname{NeQ}[q, -1]$$

rule 5570
$$\operatorname{Int}[((a_*) + \operatorname{ArcTan}[(c_*) + (d_*)(x_)]*(b_*))^{(p_*)} * ((e_*) + (f_*)(x_))^{(m_*)}, x_Symbol] \rightarrow \operatorname{Simp}[1/d \operatorname{Subst}[\operatorname{Int}[(d * e - c * f) / d + f * (x/d)]^m * (a + b \operatorname{ArcTan}[x])^p, x], x, c + d x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ \operatorname{IGtQ}[p, 0]$$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.29

method	result
parallelrisc	$-\frac{-2x^3 \arctan(bx+a)b^3+b^2x^2-2 \arctan(bx+a)a^3+3a^2 \ln(b^2x^2+2abx+a^2+1)-4abx+6a \arctan(bx+a)+7a^2-1-\ln(b^2x^2+2abx+a^2+1)}{6b^3}$
derivativedivides	$\frac{-\frac{\arctan(bx+a)a^3}{3}+\arctan(bx+a)a^2(bx+a)-\arctan(bx+a)a(bx+a)^2+\frac{\arctan(bx+a)(bx+a)^3}{3}+(bx+a)a-\frac{(bx+a)^2}{6}+\frac{(-3a^2-1)\ln(b^2x^2+2abx+a^2+1)}{6}}{b^3}$
default	$\frac{-\frac{\arctan(bx+a)a^3}{3}+\arctan(bx+a)a^2(bx+a)-\arctan(bx+a)a(bx+a)^2+\frac{\arctan(bx+a)(bx+a)^3}{3}+(bx+a)a-\frac{(bx+a)^2}{6}+\frac{(-3a^2-1)\ln(b^2x^2+2abx+a^2+1)}{6}}{b^3}$
parts	$\frac{x^3 \arctan(bx+a)}{3} - \frac{b \left(-\frac{1}{2} \frac{x^2 b + 2ax}{b^3} + \frac{(3a^2 b - b) \ln(b^2 x^2 + 2abx + a^2 + 1)}{2b^2} + \frac{\left(2a^3 + 2a - \frac{(3a^2 b - b)a}{b} \right) \arctan\left(\frac{2b^2 x + 2ab}{2b} \right)}{b^3} \right)}{3}$
risc	$-\frac{ix^3 \ln(1+i(bx+a))}{6} + \frac{ix^3 \ln(1-i(bx+a))}{6} + \frac{a^3 \arctan(bx+a)}{3b^3} - \frac{x^2}{6b} - \frac{a^2 \ln(b^2x^2+2abx+a^2+1)}{2b^3} + \frac{2ax}{3b^2} - \frac{1}{6}$

input `int(x^2*arctan(b*x+a),x,method=_RETURNVERBOSE)`output
$$-1/6*(-2*x^3*\arctan(b*x+a)*b^3+b^2*x^2-2*\arctan(b*x+a)*a^3+3*a^2*\ln(b^2*x^2+2*a*b*x+a^2+1)-4*a*b*x+6*a*\arctan(b*x+a)+7*a^2-1-\ln(b^2*x^2+2*a*b*x+a^2+1))/b^3$$
Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int x^2 \arctan(a + bx) dx = \frac{b^2 x^2 - 4 abx - 2(b^3 x^3 + a^3 - 3a) \arctan(bx + a) + (3a^2 - 1) \log(b^2 x^2 + 2abx + a^2 + 1)}{6b^3}$$

input `integrate(x^2*arctan(b*x+a),x, algorithm="fricas")`output
$$-1/6*(b^2*x^2 - 4*a*b*x - 2*(b^3*x^3 + a^3 - 3*a)*\arctan(b*x + a) + (3*a^2 - 1)*\log(b^2*x^2 + 2*a*b*x + a^2 + 1))/b^3$$

Sympy [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.48

$$\int x^2 \arctan(a + bx) dx = \begin{cases} \frac{a^3 \operatorname{atan}(a+bx)}{3b^3} - \frac{a^2 \log(a^2+2abx+b^2x^2+1)}{2b^3} + \frac{2ax}{3b^2} - \frac{a \operatorname{atan}(a+bx)}{b^3} + \frac{x^3 \operatorname{atan}(a+bx)}{3} - \frac{x^2}{6b} + \frac{\log(a^2+2abx+b^2x^2+1)}{6b^3} & \text{for } b \\ \frac{x^3 \operatorname{atan}(a)}{3} & \text{other} \end{cases}$$

input `integrate(x**2*atan(b*x+a),x)`output `Piecewise((a**3*atan(a + b*x)/(3*b**3) - a**2*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*b**3) + 2*a*x/(3*b**2) - a*atan(a + b*x)/b**3 + x**3*atan(a + b*x)/3 - x**2/(6*b) + log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(6*b**3), Ne(b, 0)), (x**3*atan(a)/3, True))`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.08

$$\int x^2 \arctan(a + bx) dx = \frac{1}{3} x^3 \arctan(bx + a) - \frac{1}{6} b \left(\frac{bx^2 - 4ax}{b^3} - \frac{2(a^3 - 3a) \arctan\left(\frac{b^2x+ab}{b}\right)}{b^4} + \frac{(3a^2 - 1) \log(b^2x^2 + 2abx + a^2 + 1)}{b^4} \right)$$

input `integrate(x^2*arctan(b*x+a),x, algorithm="maxima")`output `1/3*x^3*arctan(b*x + a) - 1/6*b*((b*x^2 - 4*a*x)/b^3 - 2*(a^3 - 3*a)*arctan((b^2*x + a*b)/b)/b^4 + (3*a^2 - 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^4)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04

$$\int x^2 \arctan(a + bx) dx = \frac{1}{3} x^3 \arctan(bx + a) + \frac{1}{6} b \left(\frac{2(a^3 - 3a) \arctan(bx + a)}{b^4} - \frac{(3a^2 - 1) \log(b^2 x^2 + 2abx + a^2 + 1)}{b^4} - \frac{b^2 x^2 - 4abx}{b^4} \right)$$

input `integrate(x^2*arctan(b*x+a),x, algorithm="giac")`output `1/3*x^3*arctan(b*x + a) + 1/6*b*(2*(a^3 - 3*a)*arctan(b*x + a)/b^4 - (3*a^2 - 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^4 - (b^2*x^2 - 4*a*b*x)/b^4)`**Mupad [B] (verification not implemented)**

Time = 1.87 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.29

$$\int x^2 \arctan(a + bx) dx = \frac{\ln(a^2 + 2abx + b^2 x^2 + 1)}{6b^3} + \frac{x^3 \operatorname{atan}(a + bx)}{3} - \frac{x^2}{6b} - \frac{a^2 \ln(a^2 + 2abx + b^2 x^2 + 1)}{2b^3} + \frac{a^3 \operatorname{atan}(a + bx)}{3b^3} - \frac{a \operatorname{atan}(a + bx)}{b^3} + \frac{2ax}{3b^2}$$

input `int(x^2*atan(a + b*x),x)`output `log(a^2 + b^2*x^2 + 2*a*b*x + 1)/(6*b^3) + (x^3*atan(a + b*x))/3 - x^2/(6*b) - (a^2*log(a^2 + b^2*x^2 + 2*a*b*x + 1))/(2*b^3) + (a^3*atan(a + b*x))/(3*b^3) - (a*atan(a + b*x))/b^3 + (2*a*x)/(3*b^2)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.19

$$\int x^2 \arctan(a + bx) dx$$

$$= \frac{2 \operatorname{atan}(bx + a) a^3 - 6 \operatorname{atan}(bx + a) a + 2 \operatorname{atan}(bx + a) b^3 x^3 - 3 \log(b^2 x^2 + 2abx + a^2 + 1) a^2 + \log(b^2 x^2 + 1) + 4abx - b^2 x^2}{6b^3}$$

input `int(x^2*atan(b*x+a),x)`output `(2*atan(a + b*x)*a**3 - 6*atan(a + b*x)*a + 2*atan(a + b*x)*b**3*x**3 - 3*log(a**2 + 2*a*b*x + b**2*x**2 + 1)*a**2 + log(a**2 + 2*a*b*x + b**2*x**2 + 1) + 4*a*b*x - b**2*x**2)/(6*b**3)`

3.3 $\int x \arctan(a + bx) dx$

Optimal result	67
Mathematica [C] (verified)	67
Rubi [A] (verified)	68
Maple [A] (verified)	70
Fricas [A] (verification not implemented)	70
Sympy [A] (verification not implemented)	71
Maxima [A] (verification not implemented)	71
Giac [A] (verification not implemented)	72
Mupad [B] (verification not implemented)	72
Reduce [B] (verification not implemented)	73

Optimal result

Integrand size = 8, antiderivative size = 60

$$\int x \arctan(a + bx) dx = -\frac{x}{2b} + \frac{(1 - a^2) \arctan(a + bx)}{2b^2} + \frac{1}{2}x^2 \arctan(a + bx) + \frac{a \log(1 + (a + bx)^2)}{2b^2}$$

output

```
-1/2*x/b+1/2*(-a^2+1)*arctan(b*x+a)/b^2+1/2*x^2*arctan(b*x+a)+1/2*a*ln(1+(b*x+a)^2)/b^2
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.02 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.50

$$\int x \arctan(a + bx) dx = \frac{-2bx + 2b^2x^2 \arctan(a + bx) + i(-i + a)^2 \log(i - a - bx) + i \log(i + a + bx) + 2a \log(i + a + bx) - ia^2}{4b^2}$$

input

```
Integrate[x*ArcTan[a + b*x],x]
```

output

$$\frac{(-2bx + 2b^2x^2 \operatorname{ArcTan}[a + bx] + I(-I + a)^2 \operatorname{Log}[I - a - bx] + I \operatorname{Log}[I + a + bx] + 2a \operatorname{Log}[I + a + bx] - I a^2 \operatorname{Log}[I + a + bx])}{(4b^2)}$$
Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5570, 25, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan(a + bx) dx \\ & \quad \downarrow \text{5570} \\ & \frac{\int x \arctan(a + bx) d(a + bx)}{b} \\ & \quad \downarrow \text{25} \\ & -\frac{\int -x \arctan(a + bx) d(a + bx)}{b} \\ & \quad \downarrow \text{27} \\ & -\frac{\int -bx \arctan(a + bx) d(a + bx)}{b^2} \\ & \quad \downarrow \text{5387} \\ & -\frac{\frac{1}{2} \int \frac{b^2 x^2}{(a+bx)^2+1} d(a + bx) - \frac{1}{2} b^2 x^2 \arctan(a + bx)}{b^2} \\ & \quad \downarrow \text{478} \\ & -\frac{\frac{1}{2} \int \left(1 - \frac{-a^2+2(a+bx)a+1}{(a+bx)^2+1}\right) d(a + bx) - \frac{1}{2} b^2 x^2 \arctan(a + bx)}{b^2} \\ & \quad \downarrow \text{2009} \\ & -\frac{\frac{1}{2}(-(1 - a^2) \arctan(a + bx) + a(-\log((a + bx)^2 + 1)) + a + bx) - \frac{1}{2} b^2 x^2 \arctan(a + bx)}{b^2} \end{aligned}$$

input `Int[x*ArcTan[a + b*x],x]`

output `-((-1/2*(b^2*x^2*ArcTan[a + b*x]) + (a + b*x - (1 - a^2)*ArcTan[a + b*x] - a*Log[1 + (a + b*x)^2])/2)/b^2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 478 `Int[((c_) + (d_)*(x_)^(n_))/((a_) + (b_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5387 `Int[((a_) + ArcTan[(c_)*(x_)*(b_)])*((d_) + (e_)*(x_)^(q_)), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 5570 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)*(b_)])^(p_)*((e_) + (f_)*(x_)^(m_)), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.05

method	result
derivativedivides	$\frac{\frac{\arctan\left(\frac{bx+a}{2}\right)(bx+a)^2}{2} - \arctan(bx+a)(bx+a)a - \frac{bx}{2} - \frac{a}{2} + \frac{a \ln\left(1+(bx+a)^2\right)}{2} + \frac{\arctan(bx+a)}{2}}{b^2}$
default	$\frac{\frac{\arctan\left(\frac{bx+a}{2}\right)(bx+a)^2}{2} - \arctan(bx+a)(bx+a)a - \frac{bx}{2} - \frac{a}{2} + \frac{a \ln\left(1+(bx+a)^2\right)}{2} + \frac{\arctan(bx+a)}{2}}{b^2}$
parallelrisch	$\frac{\arctan(bx+a)b^2x^2 - \arctan(bx+a)a^2 + a \ln(b^2x^2 + 2abx + a^2 + 1) - bx + \arctan(bx+a) + 2a}{2b^2}$
parts	$\frac{x^2 \arctan(bx+a)}{2} - \frac{b \left(\frac{x}{b^2} + \frac{-\frac{a \ln(b^2x^2 + 2abx + a^2 + 1)}{b} + \frac{(a^2 - 1) \arctan\left(\frac{2b^2x + 2ab}{b}\right)}{b^2} \right)}{2}$
risch	$-\frac{ix^2 \ln(1+i(bx+a))}{4} + \frac{ix^2 \ln(1-i(bx+a))}{4} - \frac{a^2 \arctan(bx+a)}{2b^2} + \frac{a \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} - \frac{x}{2b} + \frac{\arctan(bx+a)}{2b^2}$

input `int(x*arctan(b*x+a),x,method=_RETURNVERBOSE)`output `1/b^2*(1/2*arctan(b*x+a)*(b*x+a)^2-arctan(b*x+a)*(b*x+a)*a-1/2*b*x-1/2*a+1/2*a*ln(1+(b*x+a)^2)+1/2*arctan(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.87

$$\int x \arctan(a + bx) dx$$

$$= -\frac{bx - (b^2x^2 - a^2 + 1) \arctan(bx + a) - a \log(b^2x^2 + 2abx + a^2 + 1)}{2b^2}$$

input `integrate(x*arctan(b*x+a),x, algorithm="fricas")`output `-1/2*(b*x - (b^2*x^2 - a^2 + 1)*arctan(b*x + a) - a*log(b^2*x^2 + 2*a*b*x + a^2 + 1))/b^2`

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.30

$$\int x \arctan(a + bx) dx$$

$$= \begin{cases} -\frac{a^2 \operatorname{atan}(a+bx)}{2b^2} + \frac{a \log(a^2 + 2abx + b^2x^2 + 1)}{2b^2} + \frac{x^2 \operatorname{atan}(a+bx)}{2} - \frac{x}{2b} + \frac{\operatorname{atan}(a+bx)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atan}(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atan(b*x+a),x)`output `Piecewise((-a**2*atan(a + b*x)/(2*b**2) + a*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*b**2) + x**2*atan(a + b*x)/2 - x/(2*b) + atan(a + b*x)/(2*b**2), Ne(b, 0)), (x**2*atan(a)/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.13

$$\int x \arctan(a + bx) dx$$

$$= \frac{1}{2} x^2 \arctan(bx + a)$$

$$- \frac{1}{2} b \left(\frac{x}{b^2} + \frac{(a^2 - 1) \arctan\left(\frac{b^2x + ab}{b}\right)}{b^3} - \frac{a \log(b^2x^2 + 2abx + a^2 + 1)}{b^3} \right)$$

input `integrate(x*arctan(b*x+a),x, algorithm="maxima")`output `1/2*x^2*arctan(b*x + a) - 1/2*b*(x/b^2 + (a^2 - 1)*arctan((b^2*x + a*b)/b)/b^3 - a*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\begin{aligned} \int x \arctan(a + bx) dx \\ &= \frac{1}{2} x^2 \arctan(bx + a) \\ &\quad - \frac{1}{2} b \left(\frac{x}{b^2} + \frac{(a^2 - 1) \arctan(bx + a)}{b^3} - \frac{a \log(b^2 x^2 + 2 abx + a^2 + 1)}{b^3} \right) \end{aligned}$$

input `integrate(x*arctan(b*x+a),x, algorithm="giac")`

output `1/2*x^2*arctan(b*x + a) - 1/2*b*(x/b^2 + (a^2 - 1)*arctan(b*x + a)/b^3 - a*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^3)`

Mupad [B] (verification not implemented)

Time = 1.75 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02

$$\begin{aligned} \int x \arctan(a + bx) dx &= \frac{x^2 \operatorname{atan}(a + bx)}{2} \\ &\quad + \frac{\frac{\operatorname{atan}(a+bx)}{2} - \frac{bx}{2} - \frac{a^2 \operatorname{atan}(a+bx)}{2} + \frac{a \ln(a^2 + 2 abx + b^2 x^2 + 1)}}{b^2} \end{aligned}$$

input `int(x*atan(a + b*x),x)`

output `(x^2*atan(a + b*x))/2 + (atan(a + b*x)/2 - (b*x)/2 - (a^2*atan(a + b*x))/2 + (a*log(a^2 + b^2*x^2 + 2*a*b*x + 1))/2)/b^2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\int x \arctan(a + bx) dx$$

$$= \frac{-\operatorname{atan}(bx + a) a^2 + \operatorname{atan}(bx + a) b^2 x^2 + \operatorname{atan}(bx + a) + \log(b^2 x^2 + 2abx + a^2 + 1) a - bx}{2b^2}$$

input `int(x*atan(b*x+a),x)`output `(- atan(a + b*x)*a**2 + atan(a + b*x)*b**2*x**2 + atan(a + b*x) + log(a**2 + 2*a*b*x + b**2*x**2 + 1)*a - b*x)/(2*b**2)`

3.4 $\int \arctan(a + bx) dx$

Optimal result	74
Mathematica [A] (verified)	74
Rubi [A] (verified)	75
Maple [A] (verified)	76
Fricas [A] (verification not implemented)	76
Sympy [A] (verification not implemented)	77
Maxima [A] (verification not implemented)	77
Giac [A] (verification not implemented)	77
Mupad [B] (verification not implemented)	78
Reduce [B] (verification not implemented)	78

Optimal result

Integrand size = 6, antiderivative size = 33

$$\int \arctan(a + bx) dx = \frac{(a + bx) \arctan(a + bx)}{b} - \frac{\log(1 + (a + bx)^2)}{2b}$$

output

```
(b*x+a)*arctan(b*x+a)/b-1/2*ln(1+(b*x+a)^2)/b
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.18

$$\int \arctan(a + bx) dx = -\frac{-2(a + bx) \arctan(a + bx) + \log(1 + a^2 + 2abx + b^2x^2)}{2b}$$

input

```
Integrate[ArcTan[a + b*x],x]
```

output

```
-1/2*(-2*(a + b*x)*ArcTan[a + b*x] + Log[1 + a^2 + 2*a*b*x + b^2*x^2])/b
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5562, 5345, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(a + bx) dx$$

$$\downarrow 5562$$

$$\frac{\int \arctan(a + bx) d(a + bx)}{b}$$

$$\downarrow 5345$$

$$\frac{(a + bx) \arctan(a + bx) - \int \frac{a+bx}{(a+bx)^2+1} d(a + bx)}{b}$$

$$\downarrow 240$$

$$\frac{(a + bx) \arctan(a + bx) - \frac{1}{2} \log((a + bx)^2 + 1)}{b}$$

input `Int[ArcTan[a + b*x], x]`

output `((a + b*x)*ArcTan[a + b*x] - Log[1 + (a + b*x)^2]/2)/b`

Defintions of rubi rules used

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 5345 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 5562

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d
  Subst[Int[(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d},
  x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{(bx+a) \arctan(bx+a) - \frac{\ln(1+(bx+a)^2)}{2}}{b}$	30
default	$\frac{(bx+a) \arctan(bx+a) - \frac{\ln(1+(bx+a)^2)}{2}}{b}$	30
parallelrisc	$-\frac{-2x \arctan(bx+a)b^2 - 2 \arctan(bx+a)ba + \ln(b^2x^2 + 2abx + a^2 + 1)b}{2b^2}$	49
parts	$x \arctan(bx+a) - b \left(\frac{\ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} - \frac{a \arctan\left(\frac{2b^2x + 2ab}{b^2}\right)}{b^2} \right)$	60
risc	$-\frac{ix \ln(1+i(bx+a))}{2} + \frac{ix \ln(1-i(bx+a))}{2} + \frac{a \arctan(bx+a)}{b} - \frac{\ln(b^2x^2 + 2abx + a^2 + 1)}{2b}$	66

input

```
int(arctan(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
1/b*((b*x+a)*arctan(b*x+a)-1/2*ln(1+(b*x+a)^2))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.18

$$\int \arctan(a + bx) dx = \frac{2(bx + a) \arctan(bx + a) - \log(b^2x^2 + 2abx + a^2 + 1)}{2b}$$

input

```
integrate(arctan(b*x+a),x, algorithm="fricas")
```

output

```
1/2*(2*(b*x + a)*arctan(b*x + a) - log(b^2*x^2 + 2*a*b*x + a^2 + 1))/b
```

Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \arctan(a + bx) dx = \begin{cases} \frac{a \operatorname{atan}(a+bx)}{b} + x \operatorname{atan}(a + bx) - \frac{\log(a^2 + 2abx + b^2x^2 + 1)}{2b} & \text{for } b \neq 0 \\ x \operatorname{atan}(a) & \text{otherwise} \end{cases}$$

input `integrate(atan(b*x+a), x)`

output `Piecewise((a*atan(a + b*x)/b + x*atan(a + b*x) - log(a**2 + 2*a*b*x + b**2 *x**2 + 1)/(2*b), Ne(b, 0)), (x*atan(a), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \arctan(a + bx) dx = \frac{2(bx + a) \arctan(bx + a) - \log((bx + a)^2 + 1)}{2b}$$

input `integrate(arctan(b*x+a), x, algorithm="maxima")`

output `1/2*(2*(b*x + a)*arctan(b*x + a) - log((b*x + a)^2 + 1))/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \arctan(a + bx) dx = \frac{2(bx + a) \arctan(bx + a) - \log((bx + a)^2 + 1)}{2b}$$

input `integrate(arctan(b*x+a), x, algorithm="giac")`

output `1/2*(2*(b*x + a)*arctan(b*x + a) - log((b*x + a)^2 + 1))/b`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.27

$$\int \arctan(a + bx) dx = x \operatorname{atan}(a + bx) - \frac{\ln(a^2 + 2abx + b^2x^2 + 1) - 2a \operatorname{atan}(a + bx)}{2b}$$

input `int(atan(a + b*x),x)`

output `x*atan(a + b*x) - (log(a^2 + b^2*x^2 + 2*a*b*x + 1) - 2*a*atan(a + b*x))/(2*b)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.36

$$\int \arctan(a + bx) dx = \frac{2 \operatorname{atan}(bx + a) a + 2 \operatorname{atan}(bx + a) bx - \log(b^2x^2 + 2abx + a^2 + 1)}{2b}$$

input `int(atan(b*x+a),x)`

output `(2*atan(a + b*x)*a + 2*atan(a + b*x)*b*x - log(a**2 + 2*a*b*x + b**2*x**2 + 1))/(2*b)`

3.5 $\int \frac{\arctan(a+bx)}{x} dx$

Optimal result	79
Mathematica [A] (verified)	80
Rubi [A] (verified)	80
Maple [A] (verified)	83
Fricas [F]	83
Sympy [F(-1)]	84
Maxima [A] (verification not implemented)	84
Giac [F]	85
Mupad [F(-1)]	85
Reduce [F]	85

Optimal result

Integrand size = 10, antiderivative size = 120

$$\int \frac{\arctan(a + bx)}{x} dx = -\arctan(a + bx) \log\left(\frac{2}{1 - i(a + bx)}\right) + \arctan(a + bx) \log\left(\frac{2bx}{(i - a)(1 - i(a + bx))}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i(a + bx)}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(i - a)(1 - i(a + bx))}\right)$$

output

```
-arctan(b*x+a)*ln(2/(1-I*(b*x+a)))+arctan(b*x+a)*ln(2*b*x/(I-a)/(1-I*(b*x+a)))+1/2*I*polylog(2,1-2/(1-I*(b*x+a)))-1/2*I*polylog(2,1-2*b*x/(I-a)/(1-I*(b*x+a)))
```


Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.42

$$\int \frac{\arctan(a + bx)}{x} dx = -\frac{1}{2}i \log(1 + i(a + bx)) \log\left(\frac{i(-\frac{a}{b} + \frac{a+bx}{b})}{-\frac{1}{b} - \frac{ia}{b}}\right) + \frac{1}{2}i \log(1 - i(a + bx)) \log\left(-\frac{i(-\frac{a}{b} + \frac{a+bx}{b})}{-\frac{1}{b} + \frac{ia}{b}}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, \frac{i(1 - i(a + bx))}{i + a}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, -\frac{i(1 + i(a + bx))}{-i + a}\right)$$

input

```
Integrate[ArcTan[a + b*x]/x,x]
```

output

```
(-1/2*I)*Log[1 + I*(a + b*x)]*Log[(I*(-(a/b) + (a + b*x)/b))/(-b^(-1) - (I*a)/b)] + (I/2)*Log[1 - I*(a + b*x)]*Log[((-I)*(-(a/b) + (a + b*x)/b))/(-b^(-1) + (I*a)/b)] + (I/2)*PolyLog[2, (I*(1 - I*(a + b*x)))/(I + a)] - (I/2)*PolyLog[2, ((-I)*(1 + I*(a + b*x)))/(-I + a)]
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5570, 25, 27, 5381, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{x} dx$$

↓ 5570

$$\int \frac{\arctan(a+bx)}{x} d(a + bx)$$

↓ 25

$$\begin{aligned}
& - \int \frac{-\frac{\arctan(a+bx)}{x} d(a+bx)}{b} \\
& \quad \downarrow 27 \\
& - \int -\frac{\arctan(a+bx)}{bx} d(a+bx) \\
& \quad \downarrow 5381 \\
& \int \frac{\log\left(\frac{2}{1-i(a+bx)}\right)}{(a+bx)^2+1} d(a+bx) - \int \frac{\log\left(\frac{2bx}{(i-a)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx) - \arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right) + \arctan(a+bx) \log\left(\frac{2bx}{(-a+i)(1-i(a+bx))}\right) \\
& \quad \downarrow 2849 \\
& i \int \frac{\log\left(\frac{2}{1-i(a+bx)}\right)}{1-\frac{2}{1-i(a+bx)}} d\frac{1}{1-i(a+bx)} - \int \frac{\log\left(\frac{2bx}{(i-a)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx) - \arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right) + \arctan(a+bx) \log\left(\frac{2bx}{(-a+i)(1-i(a+bx))}\right) \\
& \quad \downarrow 2752 \\
& - \int \frac{\log\left(\frac{2bx}{(i-a)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx) - \arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right) + \arctan(a+bx) \log\left(\frac{2bx}{(-a+i)(1-i(a+bx))}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right) \\
& \quad \downarrow 2897 \\
& - \arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right) + \arctan(a+bx) \log\left(\frac{2bx}{(-a+i)(1-i(a+bx))}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2bx}{(i-a)(1-i(a+bx))}\right)
\end{aligned}$$

input `Int[ArcTan[a + b*x]/x,x]`

output `-(ArcTan[a + b*x]*Log[2/(1 - I*(a + b*x))]) + ArcTan[a + b*x]*Log[(2*b*x)/((I - a)*(1 - I*(a + b*x)))] + (I/2)*PolyLog[2, 1 - 2/(1 - I*(a + b*x))] - (I/2)*PolyLog[2, 1 - (2*b*x)/((I - a)*(1 - I*(a + b*x)))]`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 2752 $\text{Int}[\text{Log}[(\text{c}_)*(\text{x}_)]/((\text{d}_) + (\text{e}_)*(\text{x}_)), \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{e}^{(-1)})*\text{PolyLog}[2, 1 - \text{c}*x], \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}, \text{e}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{e} + \text{c}*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(\text{c}_)/((\text{d}_) + (\text{e}_)*(\text{x}_))]/((\text{f}_) + (\text{g}_)*(\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[-\text{e}/\text{g} \quad \text{Subst}[\text{Int}[\text{Log}[2*\text{d}*x]/(1 - 2*\text{d}*x), \text{x}], \text{x}, 1/(\text{d} + \text{e}*x)], \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}, \text{e}, \text{f}, \text{g}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{c}, 2*\text{d}] \ \&\& \ \text{EqQ}[\text{e}^2*\text{f} + \text{d}^2*\text{g}, 0]$
- rule 2897 $\text{Int}[\text{Log}[\text{u}_]*(\text{Pq}_)^{(\text{m}_.)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{C} = \text{FullSimplify}[\text{Pq}^{\text{m}}*((1 - \text{u})/\text{D}[\text{u}, \text{x}]]]\}, \text{Simp}[\text{C}*\text{PolyLog}[2, 1 - \text{u}], \text{x}] \text{ ; FreeQ}[\text{C}, \text{x}] \text{ ; IntegerQ}[\text{m}] \ \&\& \ \text{PolyQ}[\text{Pq}, \text{x}] \ \&\& \ \text{RationalFunctionQ}[\text{u}, \text{x}] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[\text{u}, \text{x}][[2]], \text{Expon}[\text{Pq}, \text{x}]]$
- rule 5381 $\text{Int}[(\text{a}_.) + \text{ArcTan}[(\text{c}_)*(\text{x}_)]*(\text{b}_.)]/((\text{d}_) + (\text{e}_)*(\text{x}_)), \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{a} + \text{b}*\text{ArcTan}[\text{c}*x])*(\text{Log}[2/(1 - \text{I}*c*x)]/\text{e}), \text{x}] + (\text{Simp}[(\text{a} + \text{b}*\text{ArcTan}[\text{c}*x])*(\text{Log}[2*c*((\text{d} + \text{e}*x)/((\text{c}*d + \text{I}*e)*(1 - \text{I}*c*x)))]/\text{e}), \text{x}] + \text{Simp}[\text{b}*(\text{c}/\text{e}) \quad \text{Int}[\text{Log}[2/(1 - \text{I}*c*x)]/(1 + \text{c}^2*x^2), \text{x}], \text{x}] - \text{Simp}[\text{b}*(\text{c}/\text{e}) \quad \text{Int}[\text{Log}[2*c*((\text{d} + \text{e}*x)/((\text{c}*d + \text{I}*e)*(1 - \text{I}*c*x)))]/(1 + \text{c}^2*x^2), \text{x}], \text{x}]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{c}^2*\text{d}^2 + \text{e}^2, 0]$
- rule 5570 $\text{Int}[(\text{a}_.) + \text{ArcTan}[(\text{c}_) + (\text{d}_)*(\text{x}_)]*(\text{b}_.)]^{(\text{p}_.)}*((\text{e}_.) + (\text{f}_)*(\text{x}_))^{(\text{m}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[1/\text{d} \quad \text{Subst}[\text{Int}[(\text{d}*e - \text{c}*f)/\text{d} + \text{f}*(\text{x}/\text{d})]^{(\text{m}*(\text{a} + \text{b}*\text{ArcTan}[\text{x}])^{\text{p}}), \text{x}], \text{x}, \text{c} + \text{d}*x], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}, \text{p}\}, \text{x}] \ \&\& \ \text{IntegerQ}[\text{p}, 0]$

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.78

method	result
risch	$\frac{i \operatorname{dilog}\left(-\frac{ixb}{ia-1}\right)}{2} + \frac{i \ln(-ibx-ia+1) \ln\left(-\frac{ixb}{ia-1}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{ixb}{-ia-1}\right)}{2} - \frac{i \ln(ibx+ia+1) \ln\left(\frac{ixb}{-ia-1}\right)}{2}$
parts	$\ln(x) \arctan(bx+a) - b \left(-\frac{i \ln(x) \left(\ln\left(\frac{-bx-a+i}{i-a}\right) - \ln\left(\frac{bx+a+i}{i+a}\right) \right)}{2b} - \frac{i \left(\operatorname{dilog}\left(\frac{-bx-a+i}{i-a}\right) - \operatorname{dilog}\left(\frac{bx+a+i}{i+a}\right) \right)}{2b} \right)$
derivativedivides	$\ln(-bx) \arctan(bx+a) - \frac{i \ln(-bx) \ln\left(\frac{bx+a+i}{i+a}\right)}{2} + \frac{i \ln(-bx) \ln\left(\frac{-bx-a+i}{i-a}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{bx+a+i}{i+a}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{-bx-a+i}{i-a}\right)}{2}$
default	$\ln(-bx) \arctan(bx+a) - \frac{i \ln(-bx) \ln\left(\frac{bx+a+i}{i+a}\right)}{2} + \frac{i \ln(-bx) \ln\left(\frac{-bx-a+i}{i-a}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{bx+a+i}{i+a}\right)}{2} + \frac{i \operatorname{dilog}\left(\frac{-bx-a+i}{i-a}\right)}{2}$

input `int(arctan(b*x+a)/x,x,method=_RETURNVERBOSE)`

output `1/2*I*dilog(-I*x*b/(I*a-1))+1/2*I*ln(1-I*a-I*b*x)*ln(-I*x*b/(I*a-1))-1/2*I*dilog(I*x*b/(-I*a-1))-1/2*I*ln(1+I*a+I*b*x)*ln(I*x*b/(-I*a-1))`

Fricas [F]

$$\int \frac{\arctan(a + bx)}{x} dx = \int \frac{\arctan(bx + a)}{x} dx$$

input `integrate(arctan(b*x+a)/x,x, algorithm="fricas")`

output `integral(arctan(b*x + a)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{x} dx = \text{Timed out}$$

input `integrate(atan(b*x+a)/x,x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.12

$$\begin{aligned} \int \frac{\arctan(a + bx)}{x} dx = & -\frac{1}{2} \arctan\left(\frac{bx}{a^2 + 1}, -\frac{abx}{a^2 + 1}\right) \log(b^2x^2 + 2abx + a^2 + 1) \\ & + \frac{1}{2} \arctan(bx + a) \log\left(\frac{b^2x^2}{a^2 + 1}\right) \\ & + \arctan(bx + a) \log(x) - \arctan\left(\frac{b^2x + ab}{b}\right) \log(x) \\ & - \frac{1}{2}i \operatorname{Li}_2\left(\frac{ibx + ia + 1}{ia + 1}\right) + \frac{1}{2}i \operatorname{Li}_2\left(\frac{ibx + ia - 1}{ia - 1}\right) \end{aligned}$$

input `integrate(arctan(b*x+a)/x,x, algorithm="maxima")`output `-1/2*arctan2(b*x/(a^2 + 1), -a*b*x/(a^2 + 1))*log(b^2*x^2 + 2*a*b*x + a^2 + 1) + 1/2*arctan(b*x + a)*log(b^2*x^2/(a^2 + 1)) + arctan(b*x + a)*log(x) - arctan((b^2*x + a*b)/b)*log(x) - 1/2*I*dilog((I*b*x + I*a + 1)/(I*a + 1)) + 1/2*I*dilog((I*b*x + I*a - 1)/(I*a - 1))`

Giac [F]

$$\int \frac{\arctan(a + bx)}{x} dx = \int \frac{\arctan(bx + a)}{x} dx$$

input `integrate(arctan(b*x+a)/x,x, algorithm="giac")`

output `integrate(arctan(b*x + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{x} dx = \int \frac{\operatorname{atan}(a + bx)}{x} dx$$

input `int(atan(a + b*x)/x,x)`

output `int(atan(a + b*x)/x, x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{x} dx = \int \frac{\operatorname{atan}(bx + a)}{x} dx$$

input `int(atan(b*x+a)/x,x)`

output `int(atan(a + b*x)/x,x)`

3.6 $\int \frac{\arctan(a+bx)}{x^2} dx$

Optimal result	86
Mathematica [C] (verified)	86
Rubi [A] (verified)	87
Maple [A] (verified)	89
Fricas [A] (verification not implemented)	90
Sympy [C] (verification not implemented)	90
Maxima [A] (verification not implemented)	91
Giac [A] (verification not implemented)	91
Mupad [B] (verification not implemented)	92
Reduce [B] (verification not implemented)	92

Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \frac{\arctan(a + bx)}{x^2} dx = -\frac{ab \arctan(a + bx)}{1 + a^2} - \frac{\arctan(a + bx)}{x} + \frac{b \log(x)}{1 + a^2} - \frac{b \log(1 + (a + bx)^2)}{2(1 + a^2)}$$

output `-a*b*arctan(b*x+a)/(a^2+1)-arctan(b*x+a)/x+b*ln(x)/(a^2+1)-b*ln(1+(b*x+a)^2)/(2*a^2+2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.08

$$\int \frac{\arctan(a + bx)}{x^2} dx = -\frac{\arctan(a + bx)}{x} + \frac{b(2 \log(x) + i(i + a) \log(i - a - bx) + (-1 - ia) \log(i + a + bx))}{2(1 + a^2)}$$

input `Integrate[ArcTan[a + b*x]/x^2,x]`

output `-(ArcTan[a + b*x]/x) + (b*(2*Log[x] + I*(I + a)*Log[I - a - b*x] + (-1 - I*a)*Log[I + a + b*x]))/(2*(1 + a^2))`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5568, 896, 25, 479, 452, 216, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(a + bx)}{x^2} dx \\
 & \quad \downarrow \text{5568} \\
 & b \int \frac{1}{x((a + bx)^2 + 1)} dx - \frac{\arctan(a + bx)}{x} \\
 & \quad \downarrow \text{896} \\
 & b \int \frac{1}{bx((a + bx)^2 + 1)} d(a + bx) - \frac{\arctan(a + bx)}{x} \\
 & \quad \downarrow \text{25} \\
 & -b \int -\frac{1}{bx((a + bx)^2 + 1)} d(a + bx) - \frac{\arctan(a + bx)}{x} \\
 & \quad \downarrow \text{479} \\
 & b \left(\frac{\log(-bx)}{a^2 + 1} - \frac{\int \frac{2a+bx}{(a+bx)^2+1} d(a + bx)}{a^2 + 1} \right) - \frac{\arctan(a + bx)}{x} \\
 & \quad \downarrow \text{452} \\
 & b \left(\frac{\log(-bx)}{a^2 + 1} - \frac{a \int \frac{1}{(a+bx)^2+1} d(a + bx) + \int \frac{a+bx}{(a+bx)^2+1} d(a + bx)}{a^2 + 1} \right) - \frac{\arctan(a + bx)}{x} \\
 & \quad \downarrow \text{216}
 \end{aligned}$$

$$b \left(\frac{\log(-bx)}{a^2 + 1} - \frac{\int \frac{a+bx}{(a+bx)^2+1} d(a+bx) + a \arctan(a+bx)}{a^2 + 1} \right) - \frac{\arctan(a+bx)}{x}$$

↓ 240

$$b \left(\frac{\log(-bx)}{a^2 + 1} - \frac{a \arctan(a+bx) + \frac{1}{2} \log((a+bx)^2 + 1)}{a^2 + 1} \right) - \frac{\arctan(a+bx)}{x}$$

input `Int[ArcTan[a + b*x]/x^2,x]`

output `-(ArcTan[a + b*x]/x) + b*(Log[-(b*x)]/(1 + a^2) - (a*ArcTan[a + b*x] + Log[1 + (a + b*x)^2])/2)/(1 + a^2))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 452 `Int[((c_) + (d_.)*(x_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[c Int[1/(a + b*x^2), x], x] + Simp[d Int[x/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c^2 + a*d^2, 0]`

rule 479 `Int[1/(((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^2)), x_Symbol] := Simp[d*(Log[RemoveContent[c + d*x, x]]/(b*c^2 + a*d^2)), x] + Simp[b/(b*c^2 + a*d^2) Int[(c - d*x)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 896

```
Int[((a_) + (b_)*(v_)^(n_))^(p_)*(x_)^(m_), x_Symbol] := With[{c = Coefficient[v, x, 0], d = Coefficient[v, x, 1]}, Simp[1/d^(m + 1) Subst[Int[SimplifyIntegrand[(x - c)^m*(a + b*x^n)^p, x], x], x, v], x] /; NeQ[c, 0]] /; FreeQ[{a, b, n, p}, x] && LinearQ[v, x] && IntegerQ[m]
```

rule 5568

```
Int[((a_) + ArcTan[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.98

method	result
derivativedivides	$b \left(-\frac{\arctan(bx+a)}{bx} + \frac{\ln(-bx)}{a^2+1} - \frac{\frac{\ln(1+(bx+a)^2)}{2} + a \arctan(bx+a)}{a^2+1} \right)$
default	$b \left(-\frac{\arctan(bx+a)}{bx} + \frac{\ln(-bx)}{a^2+1} - \frac{\frac{\ln(1+(bx+a)^2)}{2} + a \arctan(bx+a)}{a^2+1} \right)$
parts	$-\frac{\arctan(bx+a)}{x} + b \left(-\frac{b \left(\frac{\ln(b^2x^2+2abx+a^2+1)}{2b} + \frac{a \arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b} \right)}{a^2+1} + \frac{\ln(x)}{a^2+1} \right)$
parallelrisc	$\frac{-2x \arctan(bx+a)a^2b^2+2b^2 \ln(x)ax-b^2 \ln(b^2x^2+2abx+a^2+1)ax-2 \arctan(bx+a)a^3b-2 \arctan(bx+a)ba}{2xab(a^2+1)}$
risc	$\frac{i \ln(1+i(bx+a))}{2x} - \frac{i(a^2 \ln(1-i(bx+a))+\ln(1-i(bx+a))-i \ln((-a^2b+3iab)x-3a+2ia^2-a^3)bx-i \ln((-a^2b-3iab)x-3a+2ia^2-a^3))}{2x}$

input

```
int(arctan(b*x+a)/x^2,x,method=_RETURNVERBOSE)
```

output

```
b*(-arctan(b*x+a)/b/x+1/(a^2+1)*ln(-b*x)-1/(a^2+1)*(1/2*ln(1+(b*x+a)^2)+a*arctan(b*x+a)))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

$$\int \frac{\arctan(a + bx)}{x^2} dx$$

$$= -\frac{bx \log(b^2x^2 + 2abx + a^2 + 1) - 2bx \log(x) + 2(abx + a^2 + 1) \arctan(bx + a)}{2(a^2 + 1)x}$$

input `integrate(arctan(b*x+a)/x^2,x, algorithm="fricas")`

output `-1/2*(b*x*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*b*x*log(x) + 2*(a*b*x + a^2 + 1)*arctan(b*x + a))/((a^2 + 1)*x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.79 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.71

$$\int \frac{\arctan(a + bx)}{x^2} dx$$

$$= \begin{cases} -\frac{ib \operatorname{atan}(bx-i)}{2} - \frac{\operatorname{atan}(bx-i)}{x} - \frac{i}{2x} & \text{for } a = -i \\ \frac{ib \operatorname{atan}(bx+i)}{2} - \frac{\operatorname{atan}(bx+i)}{x} + \frac{i}{2x} & \text{for } a = i \\ -\frac{2a^2 \operatorname{atan}(a+bx)}{2a^2x+2x} - \frac{2abx \operatorname{atan}(a+bx)}{2a^2x+2x} + \frac{2bx \log(x)}{2a^2x+2x} - \frac{bx \log(a^2+2abx+b^2x^2+1)}{2a^2x+2x} - \frac{2 \operatorname{atan}(a+bx)}{2a^2x+2x} & \text{otherwise} \end{cases}$$

input `integrate(atan(b*x+a)/x**2,x)`

output `Piecewise((-I*b*atan(b*x - I)/2 - atan(b*x - I)/x - I/(2*x), Eq(a, -I)), (I*b*atan(b*x + I)/2 - atan(b*x + I)/x + I/(2*x), Eq(a, I)), (-2*a**2*atan(a + b*x)/(2*a**2*x + 2*x) - 2*a*b*x*atan(a + b*x)/(2*a**2*x + 2*x) + 2*b*x*log(x)/(2*a**2*x + 2*x) - b*x*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*a**2*x + 2*x) - 2*atan(a + b*x)/(2*a**2*x + 2*x), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.24

$$\int \frac{\arctan(a + bx)}{x^2} dx$$

$$= -\frac{1}{2} b \left(\frac{2 a \arctan\left(\frac{b^2 x + ab}{b}\right)}{a^2 + 1} + \frac{\log(b^2 x^2 + 2 abx + a^2 + 1)}{a^2 + 1} - \frac{2 \log(x)}{a^2 + 1} \right)$$

$$- \frac{\arctan(bx + a)}{x}$$

input `integrate(arctan(b*x+a)/x^2,x, algorithm="maxima")`output `-1/2*b*(2*a*arctan((b^2*x + a*b)/b)/(a^2 + 1) + log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^2 + 1) - 2*log(x)/(a^2 + 1)) - arctan(b*x + a)/x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(a + bx)}{x^2} dx$$

$$= -\frac{1}{2} b \left(\frac{2 a \arctan(bx + a)}{a^2 + 1} + \frac{\log(b^2 x^2 + 2 abx + a^2 + 1)}{a^2 + 1} - \frac{2 \log(|x|)}{a^2 + 1} \right)$$

$$- \frac{\arctan(bx + a)}{x}$$

input `integrate(arctan(b*x+a)/x^2,x, algorithm="giac")`output `-1/2*b*(2*a*arctan(b*x + a)/(a^2 + 1) + log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^2 + 1) - 2*log(abs(x))/(a^2 + 1)) - arctan(b*x + a)/x`

Mupad [B] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02

$$\int \frac{\arctan(a + bx)}{x^2} dx = -\frac{\arctan(a + bx)}{x} - \frac{\frac{bx \ln(a^2 + 2abx + b^2x^2 + 1)}{2} - bx \ln(x) + abx \arctan(a + bx)}{x(a^2 + 1)}$$

input `int(atan(a + b*x)/x^2,x)`output `- atan(a + b*x)/x - ((b*x*log(a^2 + b^2*x^2 + 2*a*b*x + 1))/2 - b*x*log(x) + a*b*x*atan(a + b*x))/(x*(a^2 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.15

$$\int \frac{\arctan(a + bx)}{x^2} dx = \frac{-2\operatorname{atan}(bx + a)a^2 - 2\operatorname{atan}(bx + a)abx - 2\operatorname{atan}(bx + a) - \log(b^2x^2 + 2abx + a^2 + 1)bx + 2\log(x)bx}{2x(a^2 + 1)}$$

input `int(atan(b*x+a)/x^2,x)`output `(- 2*atan(a + b*x)*a**2 - 2*atan(a + b*x)*a*b*x - 2*atan(a + b*x) - log(a**2 + 2*a*b*x + b**2*x**2 + 1)*b*x + 2*log(x)*b*x)/(2*x*(a**2 + 1))`

3.7 $\int \frac{\arctan(a+bx)}{x^3} dx$

Optimal result	93
Mathematica [C] (verified)	93
Rubi [A] (verified)	94
Maple [A] (verified)	96
Fricas [A] (verification not implemented)	96
Sympy [C] (verification not implemented)	97
Maxima [A] (verification not implemented)	98
Giac [A] (verification not implemented)	98
Mupad [B] (verification not implemented)	99
Reduce [B] (verification not implemented)	99

Optimal result

Integrand size = 10, antiderivative size = 96

$$\int \frac{\arctan(a + bx)}{x^3} dx = -\frac{b}{2(1 + a^2)x} - \frac{(1 - a^2)b^2 \arctan(a + bx)}{2(1 + a^2)^2} - \frac{\arctan(a + bx)}{2x^2} - \frac{ab^2 \log(x)}{(1 + a^2)^2} + \frac{ab^2 \log(1 + (a + bx)^2)}{2(1 + a^2)^2}$$

output

```
-1/2*b/(a^2+1)/x-1/2*(-a^2+1)*b^2*arctan(b*x+a)/(a^2+1)^2-1/2*arctan(b*x+a)/x^2-a*b^2*ln(x)/(a^2+1)^2+1/2*a*b^2*ln(1+(b*x+a)^2)/(a^2+1)^2
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.96

$$\int \frac{\arctan(a + bx)}{x^3} dx = \frac{-2 \arctan(a + bx) + \frac{bx(-4abx \log(x) - i(i+a)^2 bx \log(i-a-bx) + (-i+a)(-2(i+a) + (1+ia)bx \log(i+a+bx)))}{(1+a^2)^2}}{4x^2}$$

input

```
Integrate[ArcTan[a + b*x]/x^3,x]
```

output

$$\frac{(-2*\text{ArcTan}[a + b*x] + (b*x*(-4*a*b*x*\text{Log}[x] - I*(I + a)^2*b*x*\text{Log}[I - a - b*x] + (-I + a)*(-2*(I + a) + (1 + I*a)*b*x*\text{Log}[I + a + b*x])))}{(1 + a^2)^2}/(4*x^2)$$
Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5568, 896, 480, 657, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{x^3} dx$$

$$\downarrow 5568$$

$$\frac{1}{2}b \int \frac{1}{x^2((a + bx)^2 + 1)} dx - \frac{\arctan(a + bx)}{2x^2}$$

$$\downarrow 896$$

$$\frac{1}{2}b^2 \int \frac{1}{b^2x^2((a + bx)^2 + 1)} d(a + bx) - \frac{\arctan(a + bx)}{2x^2}$$

$$\downarrow 480$$

$$\frac{1}{2}b^2 \left(\frac{\int -\frac{2a+bx}{bx((a+bx)^2+1)} d(a+bx)}{a^2+1} - \frac{1}{(a^2+1)bx} \right) - \frac{\arctan(a+bx)}{2x^2}$$

$$\downarrow 657$$

$$\frac{1}{2}b^2 \left(\frac{\int \left(\frac{a^2+2(a+bx)a-1}{(a^2+1)((a+bx)^2+1)} - \frac{2a}{(a^2+1)bx} \right) d(a+bx)}{a^2+1} - \frac{1}{(a^2+1)bx} \right) - \frac{\arctan(a+bx)}{2x^2}$$

$$\downarrow 2009$$

$$\frac{1}{2}b^2 \left(\frac{-\frac{(1-a^2)\arctan(a+bx)}{a^2+1} - \frac{2a \log(-bx)}{a^2+1} + \frac{a \log((a+bx)^2+1)}{a^2+1}}{a^2+1} - \frac{1}{(a^2+1)bx} \right) - \frac{\arctan(a+bx)}{2x^2}$$

input `Int[ArcTan[a + b*x]/x^3,x]`

output `-1/2*ArcTan[a + b*x]/x^2 + (b^2*(-1/((1 + a^2)*b*x)) + (-(((1 - a^2)*ArcTan[a + b*x])/(1 + a^2)) - (2*a*Log[-(b*x)])/(1 + a^2) + (a*Log[1 + (a + b*x)^2])/(1 + a^2))/(1 + a^2))/2`

Defintions of rubi rules used

rule 480 `Int[((c_) + (d_)*(x_)^(n_))/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[d*((c + d*x)^(n + 1)/((n + 1)*(b*c^2 + a*d^2))), x] + Simp[b/(b*c^2 + a*d^2) Int[(c + d*x)^(n + 1)*((c - d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[n, -1]`

rule 657 `Int[(((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_)))/((a_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^(m+1)*((f + g*x)^n/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g, m}, x] && IntegersQ[n]`

rule 896 `Int[((a_) + (b_)*(v_)^(n_))^(p_)*(x_)^(m_), x_Symbol] := With[{c = Coefficient[v, x, 0], d = Coefficient[v, x, 1]}, Simp[1/d^(m + 1) Subst[Int[SimplifyIntegrand[(x - c)^m*(a + b*x^n)^p, x], x], x, v], x] /; NeQ[c, 0] /; FreeQ[{a, b, n, p}, x] && LinearQ[v, x] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5568 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_)^(m_)), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.88

method	result
derivativedivides	$b^2 \left(-\frac{\arctan(bx+a)}{2b^2x^2} - \frac{1}{2(a^2+1)bx} - \frac{a \ln(-bx)}{(a^2+1)^2} + \frac{a \ln(1+(bx+a)^2) + (a^2-1) \arctan(bx+a)}{2(a^2+1)^2} \right)$
default	$b^2 \left(-\frac{\arctan(bx+a)}{2b^2x^2} - \frac{1}{2(a^2+1)bx} - \frac{a \ln(-bx)}{(a^2+1)^2} + \frac{a \ln(1+(bx+a)^2) + (a^2-1) \arctan(bx+a)}{2(a^2+1)^2} \right)$
parts	$\frac{\arctan(bx+a)}{2x^2} + \frac{b^2 \left(\frac{a \ln(b^2x^2+2abx+a^2+1)}{b} + \frac{(a^2-1) \arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b} \right)}{(a^2+1)^2} - \frac{1}{(a^2+1)x} - \frac{2ab \ln(x)}{(a^2+1)^2}$
parallelrisch	$-\frac{x^2 \arctan(bx+a) a^2 b^2 + 2a b^2 \ln(x) x^2 - a b^2 \ln(b^2 x^2 + 2abx + a^2 + 1) x^2 + \arctan(bx+a) b^2 x^2 - 2a b^2 x^2 + \arctan(bx+a)}{2x^2(a^4 + 2a^2 + 1)}$
risch	$\frac{i \ln(1+i(bx+a))}{4x^2} - \frac{i(a^4 \ln(1-i(bx+a)) + 2a^2 \ln(1-i(bx+a)) + \ln(1-i(bx+a)) - \ln((a^6 b - 4ia^5 b + 9a^4 b + 8ia^3 b - 9a^2 b - 4a))}{4x^2}$

input `int(arctan(b*x+a)/x^3,x,method=_RETURNVERBOSE)`output `b^2*(-1/2*arctan(b*x+a)/b^2/x^2-1/2/(a^2+1)/b/x-1/(a^2+1)^2*a*ln(-b*x)+1/2/(a^2+1)^2*(a*ln(1+(b*x+a)^2)+(a^2-1)*arctan(b*x+a)))`**Fricas [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.99

$$\int \frac{\arctan(a + bx)}{x^3} dx$$

$$= \frac{ab^2x^2 \log(b^2x^2 + 2abx + a^2 + 1) - 2ab^2x^2 \log(x) - (a^2 + 1)bx + ((a^2 - 1)b^2x^2 - a^4 - 2a^2 - 1) \arctan(bx+a)}{2(a^4 + 2a^2 + 1)x^2}$$

input `integrate(arctan(b*x+a)/x^3,x, algorithm="fricas")`

output

```
1/2*(a*b^2*x^2*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*a*b^2*x^2*log(x) - (a^2 + 1)*b*x + ((a^2 - 1)*b^2*x^2 - a^4 - 2*a^2 - 1)*arctan(b*x + a))/((a^4 + 2*a^2 + 1)*x^2)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.11 (sec) , antiderivative size = 382, normalized size of antiderivative = 3.98

$$\int \frac{\arctan(a + bx)}{x^3} dx$$

$$= \begin{cases} -\frac{b^2 \operatorname{atan}(bx-i)}{8} - \frac{b}{8x} - \frac{\operatorname{atan}(bx-i)}{2x^2} - \frac{i}{8x^2} \\ -\frac{b^2 \operatorname{atan}(bx+i)}{8} - \frac{b}{8x} - \frac{\operatorname{atan}(bx+i)}{2x^2} + \frac{i}{8x^2} \\ -\frac{a^4 \operatorname{atan}(a+bx)}{2a^4x^2+4a^2x^2+2x^2} + \frac{a^2b^2x^2 \operatorname{atan}(a+bx)}{2a^4x^2+4a^2x^2+2x^2} - \frac{a^2bx}{2a^4x^2+4a^2x^2+2x^2} - \frac{2a^2 \operatorname{atan}(a+bx)}{2a^4x^2+4a^2x^2+2x^2} - \frac{2ab^2x^2 \log(x)}{2a^4x^2+4a^2x^2+2x^2} + \frac{ab^2x^2 \log(a^2+2a^2x^2+1)}{2a^4x^2+4a^2x^2+2x^2} \end{cases}$$

input

```
integrate(atan(b*x+a)/x**3,x)
```

output

```
Piecewise((-b**2*atan(b*x - I)/8 - b/(8*x) - atan(b*x - I)/(2*x**2) - I/(8*x**2), Eq(a, -I)), (-b**2*atan(b*x + I)/8 - b/(8*x) - atan(b*x + I)/(2*x**2) + I/(8*x**2), Eq(a, I)), (-a**4*atan(a + b*x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) + a**2*b**2*x**2*atan(a + b*x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - a**2*b*x/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - 2*a**2*atan(a + b*x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - 2*a*b**2*x**2*log(x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) + a*b**2*x**2*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - b**2*x**2*atan(a + b*x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - b*x/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2) - atan(a + b*x)/(2*a**4*x**2 + 4*a**2*x**2 + 2*x**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.17

$$\int \frac{\arctan(a + bx)}{x^3} dx$$

$$= \frac{1}{2} \left(\frac{(a^2 - 1)b \arctan\left(\frac{b^2x+ab}{b}\right)}{a^4 + 2a^2 + 1} + \frac{ab \log(b^2x^2 + 2abx + a^2 + 1)}{a^4 + 2a^2 + 1} - \frac{2ab \log(x)}{a^4 + 2a^2 + 1} - \frac{1}{(a^2 + 1)x} \right) b$$

$$- \frac{\arctan(bx + a)}{2x^2}$$

input `integrate(arctan(b*x+a)/x^3,x, algorithm="maxima")`output `1/2*((a^2 - 1)*b*arctan((b^2*x + a*b)/b)/(a^4 + 2*a^2 + 1) + a*b*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^4 + 2*a^2 + 1) - 2*a*b*log(x)/(a^4 + 2*a^2 + 1) - 1/((a^2 + 1)*x))*b - 1/2*arctan(b*x + a)/x^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.20

$$\int \frac{\arctan(a + bx)}{x^3} dx$$

$$= \frac{1}{2} \left(\frac{ab \log(b^2x^2 + 2abx + a^2 + 1)}{a^4 + 2a^2 + 1} - \frac{2ab \log(|x|)}{a^4 + 2a^2 + 1} + \frac{(a^2b^2 - b^2) \arctan(bx + a)}{(a^4 + 2a^2 + 1)b} - \frac{1}{(a^2 + 1)x} \right) b$$

$$- \frac{\arctan(bx + a)}{2x^2}$$

input `integrate(arctan(b*x+a)/x^3,x, algorithm="giac")`output `1/2*(a*b*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^4 + 2*a^2 + 1) - 2*a*b*log(abs(x))/(a^4 + 2*a^2 + 1) + (a^2*b^2 - b^2)*arctan(b*x + a)/((a^4 + 2*a^2 + 1)*b) - 1/((a^2 + 1)*x))*b - 1/2*arctan(b*x + a)/x^2`

Mupad [B] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.42

$$\int \frac{\arctan(a + bx)}{x^3} dx = \frac{a b^2 \ln(a^2 + 2 a b x + b^2 x^2 + 1)}{2(a^2 + 1)^2} - \frac{\frac{bx}{2} + \operatorname{atan}(a + bx) \left(\frac{a^2}{2} + \frac{1}{2}\right) + \frac{b^2 x^2 \operatorname{atan}(a + bx)}{2} + \frac{x^3 (b^3 - 3 a^2 b^3)}{2(a^4 + 2 a^2 + 1)} - \frac{a b^4 x^4}{(a^2 + 1)^2} + a b x \operatorname{atan}(a + bx)}{a^2 x^2 + 2 a b x^3 + b^2 x^4 + x^2} - \frac{\operatorname{atan}\left(\frac{2 x b^2 + 2 a b}{2 \sqrt{b^2 (a^2 + 1) - a^2 b^2}}\right) (b^3 - a^2 b^3)}{\sqrt{b^2} (2 a^4 + 4 a^2 + 2)} - \frac{a b^2 \ln(x)}{(a^2 + 1)^2}$$

input `int(atan(a + b*x)/x^3,x)`output
$$\left(\frac{a^2 b^2 \log(a^2 + b^2 x^2 + 2 a b x + 1)}{2(a^2 + 1)^2} - \left(\frac{b x}{2} + \operatorname{atan}(a + b x) \left(\frac{a^2}{2} + \frac{1}{2} \right) + \frac{b^2 x^2 \operatorname{atan}(a + b x)}{2} + \frac{x^3 (b^3 - 3 a^2 b^3)}{2(a^4 + 2 a^2 + 1)} - \frac{a b^4 x^4}{(a^2 + 1)^2} + a b x \operatorname{atan}(a + b x) \right) \right) / (x^2 + a^2 x^2 + b^2 x^4 + 2 a b x^3) - \frac{\operatorname{atan}\left(\frac{2 a b + 2 b^2 x}{2 \sqrt{b^2 (a^2 + 1) - a^2 b^2}}\right) (b^3 - a^2 b^3)}{\sqrt{b^2} (4 a^2 + 2 a^4 + 2)} - \frac{a b^2 \log(x)}{(a^2 + 1)^2}$$
Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.31

$$\int \frac{\arctan(a + bx)}{x^3} dx = \frac{-\operatorname{atan}(bx + a) a^4 + \operatorname{atan}(bx + a) a^2 b^2 x^2 - 2 \operatorname{atan}(bx + a) a^2 - \operatorname{atan}(bx + a) b^2 x^2 - \operatorname{atan}(bx + a) + \log(a^2 + 2 a b x + b^2 x^2 + 1) a b^2 x^2 - 2 \log(x) a b^2 x^2 - a^2 b x - b x}{2 x^2 (a^4 + 2 a^2 + 1)}$$

input `int(atan(b*x+a)/x^3,x)`output
$$\left(-\operatorname{atan}(a + b x) a^4 + \operatorname{atan}(a + b x) a^2 b^2 x^2 - 2 \operatorname{atan}(a + b x) a^2 - \operatorname{atan}(a + b x) b^2 x^2 - \operatorname{atan}(a + b x) + \log(a^2 + 2 a b x + b^2 x^2 + 1) a b^2 x^2 - 2 \log(x) a b^2 x^2 - a^2 b x - b x \right) / (2 x^2 (a^4 + 2 a^2 + 1))$$

3.8 $\int \frac{\arctan(a+bx)}{x^4} dx$

Optimal result	100
Mathematica [C] (verified)	100
Rubi [A] (verified)	101
Maple [A] (verified)	103
Fricas [A] (verification not implemented)	104
Sympy [C] (verification not implemented)	104
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	106
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	107

Optimal result

Integrand size = 10, antiderivative size = 129

$$\int \frac{\arctan(a+bx)}{x^4} dx = -\frac{b}{6(1+a^2)x^2} + \frac{2ab^2}{3(1+a^2)^2x} + \frac{a(3-a^2)b^3 \arctan(a+bx)}{3(1+a^2)^3} - \frac{\arctan(a+bx)}{3x^3} - \frac{(1-3a^2)b^3 \log(x)}{3(1+a^2)^3} + \frac{(1-3a^2)b^3 \log(1+(a+bx)^2)}{6(1+a^2)^3}$$

output

```
-1/6*b/(a^2+1)/x^2+2/3*a*b^2/(a^2+1)^2/x+1/3*a*(-a^2+3)*b^3*arctan(b*x+a)/(a^2+1)^3-1/3*arctan(b*x+a)/x^3-1/3*(-3*a^2+1)*b^3*ln(x)/(a^2+1)^3+1/6*(-3*a^2+1)*b^3*ln(1+(b*x+a)^2)/(a^2+1)^3
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.99

$$\int \frac{\arctan(a+bx)}{x^4} dx = \frac{-2(1+a^2)^3 \arctan(a+bx) + 2(-1+3a^2)b^3x^3 \log(x) + i(i+a)^3b^3x^3 \log(i-a-bx) - (-i+a)bx((i+a)^3 - (i-a)^3)}{6(1+a^2)^3x^3}$$

input `Integrate[ArcTan[a + b*x]/x^4,x]`

output $(-2*(1 + a^2)^3*ArcTan[a + b*x] + 2*(-1 + 3*a^2)*b^3*x^3*Log[x] + I*(I + a)^3*b^3*x^3*Log[I - a - b*x] - (-I + a)*b*x*((I + a)*(1 + a^2 - 4*a*b*x) + I*(-I + a)^2*b^2*x^2*Log[I + a + b*x]))/(6*(1 + a^2)^3*x^3)$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5568, 896, 25, 480, 657, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{x^4} dx$$

$$\downarrow 5568$$

$$\frac{1}{3}b \int \frac{1}{x^3((a + bx)^2 + 1)} dx - \frac{\arctan(a + bx)}{3x^3}$$

$$\downarrow 896$$

$$\frac{1}{3}b^3 \int \frac{1}{b^3x^3((a + bx)^2 + 1)} d(a + bx) - \frac{\arctan(a + bx)}{3x^3}$$

$$\downarrow 25$$

$$-\frac{1}{3}b^3 \int -\frac{1}{b^3x^3((a + bx)^2 + 1)} d(a + bx) - \frac{\arctan(a + bx)}{3x^3}$$

$$\downarrow 480$$

$$\frac{1}{3}b^3 \left(-\frac{\int \frac{2a+bx}{b^2x^2((a+bx)^2+1)} d(a+bx)}{a^2+1} - \frac{1}{2(a^2+1)b^2x^2} \right) - \frac{\arctan(a+bx)}{3x^3}$$

$$\downarrow 657$$

$$\frac{1}{3}b^3 \left(-\frac{\int \left(\frac{2a}{(a^2+1)b^2x^2} - \frac{3a^2-1}{(a^2+1)^2bx} + \frac{-a(3-a^2)-(1-3a^2)(a+bx)}{(a^2+1)^2((a+bx)^2+1)} \right) d(a+bx)}{a^2+1} - \frac{1}{2(a^2+1)b^2x^2} \right) - \frac{\arctan(a+bx)}{3x^3}$$

↓ 2009

$$\frac{1}{3}b^3 \left(-\frac{-\frac{(3-a^2)a \arctan(a+bx)}{(a^2+1)^2} - \frac{2a}{(a^2+1)bx} + \frac{(1-3a^2) \log(-bx)}{(a^2+1)^2} - \frac{(1-3a^2) \log((a+bx)^2+1)}{2(a^2+1)^2}}{a^2+1} - \frac{1}{2(a^2+1)b^2x^2} \right) - \frac{\arctan(a+bx)}{3x^3}$$

input `Int[ArcTan[a + b*x]/x^4,x]`

output `-1/3*ArcTan[a + b*x]/x^3 + (b^3*(-1/2*1/((1 + a^2)*b^2*x^2) - ((-2*a)/((1 + a^2)*b*x) - (a*(3 - a^2)*ArcTan[a + b*x])/(1 + a^2)^2 + ((1 - 3*a^2)*Log[-(b*x)])/(1 + a^2)^2 - ((1 - 3*a^2)*Log[1 + (a + b*x)^2])/(2*(1 + a^2)^2))/(1 + a^2))/3`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 480 `Int[((c_) + (d_)*(x_))^(n_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[d*((c + d*x)^(n + 1)/((n + 1)*(b*c^2 + a*d^2))), x] + Simp[b/(b*c^2 + a*d^2) Int[(c + d*x)^(n + 1)*((c - d*x)/(a + b*x^2)), x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[n, -1]`

rule 657 `Int[(((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_))/((a_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g, m}, x] && IntegersQ[n]`

```
rule 896 Int[((a_) + (b_.)*(v_)^(n_))^(p_.)*(x_)^(m_.), x_Symbol] := With[{c = Coefficient[v, x, 0], d = Coefficient[v, x, 1]}, Simp[1/d^(m + 1) Subst[Int[SimplifyIntegrand[(x - c)^m*(a + b*x^n)^p, x], x], x, v], x] /; NeQ[c, 0]] /; FreeQ[{a, b, n, p}, x] && LinearQ[v, x] && IntegerQ[m]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5568 Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.89

method	result
derivativedivides	$b^3 \left(-\frac{\arctan(bx+a)}{3b^3x^3} - \frac{(-3a^2+1)\ln(-bx)}{3(a^2+1)^3} - \frac{1}{6(a^2+1)b^2x^2} + \frac{2a}{3(a^2+1)^2bx} - \frac{(3a^2-1)\ln(1+(bx+a)^2)}{2 \cdot 3(a^2+1)^3} + \frac{a^3-3a}{3(a^2+1)^3} \right)$
default	$b^3 \left(-\frac{\arctan(bx+a)}{3b^3x^3} - \frac{(-3a^2+1)\ln(-bx)}{3(a^2+1)^3} - \frac{1}{6(a^2+1)b^2x^2} + \frac{2a}{3(a^2+1)^2bx} - \frac{(3a^2-1)\ln(1+(bx+a)^2)}{2 \cdot 3(a^2+1)^3} + \frac{a^3-3a}{3(a^2+1)^3} \right)$
parts	$-\frac{\arctan(bx+a)}{3x^3} + \frac{b^3 \left(\frac{(3a^2b-b)\ln(b^2x^2+2abx+a^2+1)}{2b^2} + \frac{\left(4a^3-4a-\frac{(3a^2b-b)a}{b}\right)\arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b} \right)}{(a^2+1)^3} - \frac{1}{2(a^2+1)a}$
parallelrisc	$\frac{-2x^3 \arctan(bx+a)a^3b^3+6\ln(x)x^3a^2b^3-3\ln(b^2x^2+2abx+a^2+1)x^3a^2b^3+6x^3 \arctan(bx+a)ab^3-7x^3a^2b^3-2b^3\ln(x)}{3}$
risc	Expression too large to display

```
input int(arctan(b*x+a)/x^4,x,method=_RETURNVERBOSE)
```


output

```
b^3*(-1/3*arctan(b*x+a)/b^3/x^3-1/3*(-3*a^2+1)/(a^2+1)^3*ln(-b*x)-1/6/(a^2+1)/b^2/x^2+2/3/(a^2+1)^2*a/b/x-1/3/(a^2+1)^3*(1/2*(3*a^2-1)*ln(1+(b*x+a)^2)+(a^3-3*a)*arctan(b*x+a)))
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.05

$$\int \frac{\arctan(a + bx)}{x^4} dx = \frac{(3a^2 - 1)b^3x^3 \log(b^2x^2 + 2abx + a^2 + 1) - 2(3a^2 - 1)b^3x^3 \log(x) - 4(a^3 + a)b^2x^2 + (a^4 + 2a^2 + 1)b^2x - 2(a^3 - 3a)b^2x + a^6 + 3a^4 + 3a^2 + 1}{6(a^6 + 3a^4 + 3a^2 + 1)x^3}$$

input

```
integrate(arctan(b*x+a)/x^4,x, algorithm="fricas")
```

output

```
-1/6*((3*a^2 - 1)*b^3*x^3*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*(3*a^2 - 1)*b^3*x^3*log(x) - 4*(a^3 + a)*b^2*x^2 + (a^4 + 2*a^2 + 1)*b*x + 2*((a^3 - 3*a)*b^3*x^3 + a^6 + 3*a^4 + 3*a^2 + 1)*arctan(b*x + a))/((a^6 + 3*a^4 + 3*a^2 + 1)*x^3)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.47 (sec) , antiderivative size = 760, normalized size of antiderivative = 5.89

$$\int \frac{\arctan(a + bx)}{x^4} dx = \text{Too large to display}$$

input

```
integrate(atan(b*x+a)/x**4,x)
```

output

```
Piecewise((I*b**3*atan(b*x - I)/24 + I*b**2/(24*x) - b/(24*x**2) - atan(b*x - I)/(3*x**3) - I/(18*x**3), Eq(a, -I)), (-I*b**3*atan(b*x + I)/24 - I*b**2/(24*x) - b/(24*x**2) - atan(b*x + I)/(3*x**3) + I/(18*x**3), Eq(a, I)), (-2*a**6*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - a**4*b*x/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 6*a**4*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 2*a**3*b**3*x**3*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) + 4*a**3*b**2*x**2/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) + 6*a**2*b**3*x**3*log(x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 3*a**2*b**3*x**3*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 2*a**2*b*x/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 6*a**2*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) + 6*a*b**3*x**3*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) + 4*a*b**2*x**2/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 2*b**3*x**3*log(x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) + b**3*x**3*log(a**2 + 2*a*b*x + b**2*x**2 + 1)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - b*x/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3) - 2*atan(a + b*x)/(6*a**6*x**3 + 18*a**4*x**3 + 18*a**2*x**3 + 6*x**3), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.28

$$\int \frac{\arctan(a + bx)}{x^4} dx =$$

$$-\frac{1}{6} \left(\frac{2(a^3 - 3a)b^2 \arctan\left(\frac{b^2x + ab}{b}\right)}{a^6 + 3a^4 + 3a^2 + 1} + \frac{(3a^2 - 1)b^2 \log(b^2x^2 + 2abx + a^2 + 1)}{a^6 + 3a^4 + 3a^2 + 1} - \frac{2(3a^2 - 1)b^2 \log(x)}{a^6 + 3a^4 + 3a^2 + 1} \right) - \frac{\arctan(bx + a)}{3x^3}$$

input

```
integrate(arctan(b*x+a)/x^4,x, algorithm="maxima")
```

output

```
-1/6*(2*(a^3 - 3*a)*b^2*arctan((b^2*x + a*b)/b)/(a^6 + 3*a^4 + 3*a^2 + 1) + (3*a^2 - 1)*b^2*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^6 + 3*a^4 + 3*a^2 + 1) - 2*(3*a^2 - 1)*b^2*log(x)/(a^6 + 3*a^4 + 3*a^2 + 1) - (4*a*b*x - a^2 - 1)/((a^4 + 2*a^2 + 1)*x^2))*b - 1/3*arctan(b*x + a)/x^3
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.37

$$\int \frac{\arctan(a + bx)}{x^4} dx =$$

$$-\frac{1}{6} b \left(\frac{(3a^2b^2 - b^2) \log(b^2x^2 + 2abx + a^2 + 1)}{a^6 + 3a^4 + 3a^2 + 1} - \frac{2(3a^2b^2 - b^2) \log(|x|)}{a^6 + 3a^4 + 3a^2 + 1} + \frac{2(a^3b^3 - 3ab^3) \arctan(bx + a)}{(a^6 + 3a^4 + 3a^2 + 1)b} \right) - \frac{\arctan(bx + a)}{3x^3}$$

input `integrate(arctan(b*x+a)/x^4,x, algorithm="giac")`output `-1/6*b*((3*a^2*b^2 - b^2)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^6 + 3*a^4 + 3*a^2 + 1) - 2*(3*a^2*b^2 - b^2)*log(abs(x))/(a^6 + 3*a^4 + 3*a^2 + 1) + 2*(a^3*b^3 - 3*a*b^3)*arctan(b*x + a)/((a^6 + 3*a^4 + 3*a^2 + 1)*b) + (a^4 + 2*a^2 - 4*(a^3*b + a*b)*x + 1)/((a^2 + 1)^3*x^2)) - 1/3*arctan(b*x + a)/x^3`**Mupad [B] (verification not implemented)**

Time = 1.46 (sec) , antiderivative size = 288, normalized size of antiderivative = 2.23

$$\int \frac{\arctan(a + bx)}{x^4} dx =$$

$$\frac{\frac{bx}{6} + \operatorname{atan}(a + bx) \left(\frac{a^2}{3} + \frac{1}{3} \right) + \frac{b^2 x^2 \operatorname{atan}(a + bx)}{3} + \frac{x^3 (b^3 - 7a^2 b^3)}{6(a^4 + 2a^2 + 1)} - \frac{ab^2 x^2}{3(a^2 + 1)} - \frac{2ab^4 x^4}{3(a^2 + 1)^2} + \frac{2abx \operatorname{atan}(a + bx)}{3}}{a^2 x^3 + 2abx^4 + b^2 x^5 + x^3}$$

$$-\frac{\ln(x) \left(\frac{b^3}{3} - a^2 b^3 \right)}{a^6 + 3a^4 + 3a^2 + 1} - \frac{b^3 \ln(a^2 + 2abx + b^2 x^2 + 1) (3a^2 - 1)}{6(a^6 + 3a^4 + 3a^2 + 1)}$$

$$-\frac{a \operatorname{atan}\left(\frac{2xb^2 + 2ab}{2\sqrt{b^2(a^2 + 1) - a^2 b^2}}\right) (a^2 - 3) (b^2)^{3/2}}{3(a^6 + 3a^4 + 3a^2 + 1)}$$

input `int(atan(a + b*x)/x^4,x)`

output

```
- ((b*x)/6 + atan(a + b*x)*(a^2/3 + 1/3) + (b^2*x^2*atan(a + b*x))/3 + (x^
3*(b^3 - 7*a^2*b^3))/(6*(2*a^2 + a^4 + 1)) - (a*b^2*x^2)/(3*(a^2 + 1)) - (
2*a*b^4*x^4)/(3*(a^2 + 1)^2) + (2*a*b*x*atan(a + b*x))/3)/(x^3 + a^2*x^3 +
b^2*x^5 + 2*a*b*x^4) - (log(x)*(b^3/3 - a^2*b^3))/(3*a^2 + 3*a^4 + a^6 +
1) - (b^3*log(a^2 + b^2*x^2 + 2*a*b*x + 1)*(3*a^2 - 1))/(6*(3*a^2 + 3*a^4
+ a^6 + 1)) - (a*atan((2*a*b + 2*b^2*x)/(2*(b^2*(a^2 + 1) - a^2*b^2)^(1/2)
)))*(a^2 - 3)*(b^2)^(3/2))/(3*(3*a^2 + 3*a^4 + a^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.64

$$\int \frac{\arctan(a + bx)}{x^4} dx$$

$$= \frac{-2\operatorname{atan}(bx + a)a^6 - 6\operatorname{atan}(bx + a)a^4 - 2\operatorname{atan}(bx + a)a^3b^3x^3 - 6\operatorname{atan}(bx + a)a^2 + 6\operatorname{atan}(bx + a)ab^3x^3}{x^3(a^6 + 3a^4 + 3a^2 + 1)}$$

input

```
int(atan(b*x+a)/x^4,x)
```

output

```
( - 2*atan(a + b*x)*a**6 - 6*atan(a + b*x)*a**4 - 2*atan(a + b*x)*a**3*b**
3*x**3 - 6*atan(a + b*x)*a**2 + 6*atan(a + b*x)*a*b**3*x**3 - 2*atan(a + b
*x) - 3*log(a**2 + 2*a*b*x + b**2*x**2 + 1)*a**2*b**3*x**3 + log(a**2 + 2*
a*b*x + b**2*x**2 + 1)*b**3*x**3 + 6*log(x)*a**2*b**3*x**3 - 2*log(x)*b**3
*x**3 - a**4*b*x + 4*a**3*b**2*x**2 - 2*a**2*b*x + 4*a*b**2*x**2 - b*x)/(6
*x**3*(a**6 + 3*a**4 + 3*a**2 + 1))
```

3.9 $\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$

Optimal result	108
Mathematica [A] (verified)	108
Rubi [A] (verified)	109
Maple [A] (verified)	111
Fricas [B] (verification not implemented)	111
Sympy [B] (verification not implemented)	112
Maxima [B] (verification not implemented)	113
Giac [B] (verification not implemented)	114
Mupad [B] (verification not implemented)	115
Reduce [B] (verification not implemented)	116

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx = \frac{1}{4}be^3x - \frac{be^3(c + dx)^3}{12d} - \frac{be^3 \arctan(c + dx)}{4d} + \frac{e^3(c + dx)^4(a + b \arctan(c + dx))}{4d}$$

output

```
1/4*b*e^3*x-1/12*b*e^3*(d*x+c)^3/d-1/4*b*e^3*arctan(d*x+c)/d+1/4*e^3*(d*x+c)^4*(a+b*arctan(d*x+c))/d
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.78

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx = \frac{e^3 \left(-\frac{1}{4}b(-dx + \frac{1}{3}(c + dx)^3 + \arctan(c + dx)) + \frac{1}{4}(c + dx)^4(a + b \arctan(c + dx)) \right)}{d}$$

input

```
Integrate[(c*e + d*e*x)^3*(a + b*ArcTan[c + d*x]),x]
```

output

$$\frac{(e^{3*(-1/4*(b*(-d*x) + (c + d*x)^{3/3} + \text{ArcTan}[c + d*x]))} + ((c + d*x)^{4*(a + b*\text{ArcTan}[c + d*x]))/4))/d$$
Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {5566, 27, 5361, 254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (ce + dex)^3 (a + b \arctan(c + dx)) dx \\ & \quad \downarrow \text{5566} \\ & \frac{\int e^3 (c + dx)^3 (a + b \arctan(c + dx)) d(c + dx)}{d} \\ & \quad \downarrow \text{27} \\ & \frac{e^3 \int (c + dx)^3 (a + b \arctan(c + dx)) d(c + dx)}{d} \\ & \quad \downarrow \text{5361} \\ & \frac{e^3 \left(\frac{1}{4} (c + dx)^4 (a + b \arctan(c + dx)) - \frac{1}{4} b \int \frac{(c+dx)^4}{(c+dx)^2+1} d(c + dx) \right)}{d} \\ & \quad \downarrow \text{254} \\ & \frac{e^3 \left(\frac{1}{4} (c + dx)^4 (a + b \arctan(c + dx)) - \frac{1}{4} b \int \left((c + dx)^2 + \frac{1}{(c+dx)^2+1} - 1 \right) d(c + dx) \right)}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{e^3 \left(\frac{1}{4} (c + dx)^4 (a + b \arctan(c + dx)) - \frac{1}{4} b (\arctan(c + dx) + \frac{1}{3} (c + dx)^3 - c - dx) \right)}{d} \end{aligned}$$

input

$$\text{Int}[(c*e + d*e*x)^3*(a + b*\text{ArcTan}[c + d*x]), x]$$

output $(e^{3*(-1/4*(b*(-c - d*x + (c + d*x)^{3/3} + \text{ArcTan}[c + d*x])) + ((c + d*x)^4 * (a + b*\text{ArcTan}[c + d*x]))/4))/d$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 254 $\text{Int}[(x_)^(m_)/((a_) + (b_)*(x_)^2), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^2, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[m, 3]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 5361 $\text{Int}[(a_ + \text{ArcTan}[c_)*(x_)^(n_)]*(b_)^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*(a + b*\text{ArcTan}[c*x^n])^{p/(m+1)}, x] - \text{Simp}[b*c*n*(p/(m+1)) \text{Int}[x^{(m+n)}*(a + b*\text{ArcTan}[c*x^n])^{(p-1)/(1+c^2*x^{2*n})}], x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \& \ \& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 5566 $\text{Int}[(a_ + \text{ArcTan}[c_ + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{e^3 a (dx+c)^4 + e^3 b \left(\frac{(dx+c)^4 \arctan(dx+c)}{4} - \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} - \frac{\arctan(dx+c)}{4} \right)}{d}$
default	$\frac{e^3 a (dx+c)^4 + e^3 b \left(\frac{(dx+c)^4 \arctan(dx+c)}{4} - \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} - \frac{\arctan(dx+c)}{4} \right)}{d}$
parts	$\frac{e^3 a (dx+c)^4}{4d} + \frac{e^3 b \left(\frac{(dx+c)^4 \arctan(dx+c)}{4} - \frac{(dx+c)^3}{12} + \frac{dx}{4} + \frac{c}{4} - \frac{\arctan(dx+c)}{4} \right)}{d}$
orering	$\frac{(2d^5 x^5 + 10c d^4 x^4 + 20c^2 d^3 x^3 + 19c^3 d^2 x^2 + 8c^4 dx - 2d^3 x^3 + c^5 - 3c d^2 x^2 - 4dx - c)(dex+ce)^3 (a+b \arctan(dx+c))}{4d(dx+c)^4} - \frac{x(a}{d}$
parallelrisc	$3d^5 e^3 b \arctan(dx+c)x^4 + 3x^4 a d^5 e^3 + 12bc d^4 e^3 \arctan(dx+c)x^3 + 12x^3 ac d^4 e^3 + 18x^2 \arctan(dx+c) b c^2 d^3 e^3 - x^3 b d^4 e^3$
risch	$\frac{ie^3 d^3 b x^4 \ln(1-i(dx+c))}{8} + \frac{ie^3 d^2 bc x^3 \ln(1-i(dx+c))}{2} + \frac{ie^3 b c^3 x \ln(1-i(dx+c))}{2} + \frac{3ie^3 db c^2 x^2 \ln(1-i(dx+c))}{4} -$

input `int((d*e*x+c*e)^3*(a+b*arctan(d*x+c)),x,method=_RETURNVERBOSE)`output `1/d*(1/4*e^3*a*(d*x+c)^4+e^3*b*(1/4*(d*x+c)^4*arctan(d*x+c)-1/12*(d*x+c)^3+1/4*d*x+1/4*c-1/4*arctan(d*x+c)))`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. 2(64) = 128.

Time = 0.12 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.10

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{3ad^4 e^3 x^4 + (12ac - b)d^3 e^3 x^3 + 3(6ac^2 - bc)d^2 e^3 x^2 + 3(4ac^3 - bc^2 + b)de^3 x + 3(bd^4 e^3 x^4 + 4bcd^3 e^3 x^3}{12d}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c)),x, algorithm="fricas")`

output

```
1/12*(3*a*d^4*e^3*x^4 + (12*a*c - b)*d^3*e^3*x^3 + 3*(6*a*c^2 - b*c)*d^2*e^3*x^2 + 3*(4*a*c^3 - b*c^2 + b)*d*e^3*x + 3*(b*d^4*e^3*x^4 + 4*b*c*d^3*e^3*x^3 + 6*b*c^2*d^2*e^3*x^2 + 4*b*c^3*d*e^3*x + (b*c^4 - b)*e^3)*arctan(d*x + c))/d
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 231 vs. $2(61) = 122$.

Time = 1.13 (sec) , antiderivative size = 231, normalized size of antiderivative = 3.21

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$$

$$= \begin{cases} ac^3 e^3 x + \frac{3ac^2 de^3 x^2}{2} + acd^2 e^3 x^3 + \frac{ad^3 e^3 x^4}{4} + \frac{bc^4 e^3 \operatorname{atan}(c+dx)}{4d} + bc^3 e^3 x \operatorname{atan}(c + dx) + \frac{3bc^2 de^3 x^2 \operatorname{atan}(c+dx)}{2} - \dots \\ c^3 e^3 x (a + b \operatorname{atan}(c)) \end{cases}$$

input

```
integrate((d*e*x+c*e)**3*(a+b*atan(d*x+c)),x)
```

output

```
Piecewise((a*c**3*e**3*x + 3*a*c**2*d*e**3*x**2/2 + a*c*d**2*e**3*x**3 + a*d**3*e**3*x**4/4 + b*c**4*e**3*atan(c + d*x)/(4*d) + b*c**3*e**3*x*atan(c + d*x) + 3*b*c**2*d*e**3*x**2*atan(c + d*x)/2 - b*c**2*e**3*x/4 + b*c*d**2*e**3*x**3*atan(c + d*x) - b*c*d*e**3*x**2/4 + b*d**3*e**3*x**4*atan(c + d*x)/4 - b*d**2*e**3*x**3/12 + b*e**3*x/4 - b*e**3*atan(c + d*x)/(4*d), Ne(d, 0)), (c**3*e**3*x*(a + b*atan(c)), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 370 vs. $2(64) = 128$.

Time = 0.13 (sec) , antiderivative size = 370, normalized size of antiderivative = 5.14

$$\int (ce + dex)^3(a + b \arctan(c + dx)) dx = \frac{1}{4} ad^3 e^3 x^4 + acd^2 e^3 x^3 + \frac{3}{2} ac^2 de^3 x^2 + \frac{3}{2} \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^3} - \frac{c \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) bc^2 de^3 + \frac{1}{2} \left(2x^3 \arctan(dx + c) - d \left(\frac{dx^2 - 4cx}{d^3} - \frac{2(c^3 - 3c) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^4} + \frac{(3c^2 - 1) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^4} \right) \right) bc^2 de^3 + \frac{1}{12} \left(3x^4 \arctan(dx + c) - d \left(\frac{d^2 x^3 - 3cdx^2 + 3(3c^2 - 1)x}{d^4} + \frac{3(c^4 - 6c^2 + 1) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^5} - \frac{6(c^3 - c) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^5} \right) \right) bc^2 de^3 + ac^3 e^3 x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) bc^3 e^3}{2d}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output `1/4*a*d^3*e^3*x^4 + a*c*d^2*e^3*x^3 + 3/2*a*c^2*d*e^3*x^2 + 3/2*(x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3))*b*c^2*d*e^3 + 1/2*(2*x^3*arctan(d*x + c) - d*((d*x^2 - 4*c*x)/d^3 - 2*(c^3 - 3*c)*arctan((d^2*x + c*d)/d)/d^4 + (3*c^2 - 1)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^4))*b*c*d^2*e^3 + 1/12*(3*x^4*arctan(d*x + c) - d*((d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 - 1)*x)/d^4 + 3*(c^4 - 6*c^2 + 1)*arctan((d^2*x + c*d)/d)/d^5 - 6*(c^3 - c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^5))*b*d^3*e^3 + a*c^3*e^3*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b*c^3*e^3/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. $2(64) = 128$.

Time = 0.24 (sec) , antiderivative size = 253, normalized size of antiderivative = 3.51

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{6bd^4e^3x^4 \arctan(dx + c) + 6ad^4e^3x^4 + 24bcd^3e^3x^3 \arctan(dx + c) + 24acd^3e^3x^3 + 36bc^2d^2e^3x^2 \arctan(dx + c) + 36ac^2d^2e^3x^2 - 2bd^3e^3x^3 + 24b^2c^3d^2e^3x^2 \arctan(dx + c) + 3\pi b^2c^4e^3 \operatorname{sgn}(dx + c) - 3\pi b^2c^4e^3 + 24a^2c^3d^2e^3x - 6b^2c^4d^2e^3x^2 - 6b^2c^4e^3 \arctan(1/(dx + c)) - 6b^2c^2d^2e^3x + 6b^2d^2e^3x - 3\pi b^2e^3 \operatorname{sgn}(dx + c) + 3\pi b^2e^3 + 6b^2e^3 \arctan(1/(dx + c))}{d}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output `1/24*(6*b*d^4*e^3*x^4*arctan(d*x + c) + 6*a*d^4*e^3*x^4 + 24*b*c*d^3*e^3*x^3*arctan(d*x + c) + 24*a*c*d^3*e^3*x^3 + 36*b*c^2*d^2*e^3*x^2*arctan(d*x + c) + 36*a*c^2*d^2*e^3*x^2 - 2*b*d^3*e^3*x^3 + 24*b*c^3*d*e^3*x*arctan(d*x + c) + 3*pi*b*c^4*e^3*sgn(d*x + c) - 3*pi*b*c^4*e^3 + 24*a*c^3*d*e^3*x - 6*b*c*d^2*e^3*x^2 - 6*b*c^4*e^3*arctan(1/(d*x + c)) - 6*b*c^2*d*e^3*x + 6*b*d*e^3*x - 3*pi*b*e^3*sgn(d*x + c) + 3*pi*b*e^3 + 6*b*e^3*arctan(1/(d*x + c)))/d`

Mupad [B] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 371, normalized size of antiderivative = 5.15

$$\begin{aligned}
& \int (ce + dex)^3 (a + b \arctan(c + dx)) dx \\
&= \operatorname{atan}(c + dx) \left(bc^3 e^3 x + \frac{3bc^2 d e^3 x^2}{2} + bcd^2 e^3 x^3 + \frac{bd^3 e^3 x^4}{4} \right) \\
&\quad - x^3 \left(\frac{d^2 e^3 (b - 20ac)}{12} + \frac{2acd^2 e^3}{3} \right) \\
&\quad + x^2 \left(\frac{c \left(\frac{d^2 e^3 (b - 20ac)}{4} + 2acd^2 e^3 \right)}{d} + \frac{de^3 (10ac^2 - bc + a)}{2} - \frac{ade^3 (4c^2 + 4)}{8} \right) \\
&\quad + x \left(\frac{ce^3 (20ac^2 - 3bc + 6a)}{2} + \frac{(4c^2 + 4) \left(\frac{d^2 e^3 (b - 20ac)}{4} + 2acd^2 e^3 \right)}{4d^2} \right. \\
&\quad \left. - \frac{2c \left(\frac{2c \left(\frac{d^2 e^3 (b - 20ac)}{4} + 2acd^2 e^3 \right)}{d} + de^3 (10ac^2 - bc + a) - \frac{ade^3 (4c^2 + 4)}{4} \right)}{d} \right) \\
&\quad + \frac{ad^3 e^3 x^4}{4} \\
&\quad - \frac{be^3 \operatorname{atan} \left(\frac{\frac{bc e^3 (c^2 + 1)(c - 1)(c + 1)}{4} + \frac{bd e^3 x (c^2 + 1)(c - 1)(c + 1)}{4}}{\frac{be^3}{4} - \frac{bc^4 e^3}{4}} \right) (c^2 + 1)(c - 1)(c + 1)}{4d}
\end{aligned}$$

input

```
int((c*e + d*e*x)^3*(a + b*atan(c + d*x)),x)
```

output

```
atan(c + d*x)*((b*d^3*e^3*x^4)/4 + b*c^3*e^3*x + (3*b*c^2*d*e^3*x^2)/2 + b
*c*d^2*e^3*x^3) - x^3*((d^2*e^3*(b - 20*a*c))/12 + (2*a*c*d^2*e^3)/3) + x^
2*((c*((d^2*e^3*(b - 20*a*c))/4 + 2*a*c*d^2*e^3))/d + (d*e^3*(a - b*c + 10
*a*c^2))/2 - (a*d*e^3*(4*c^2 + 4))/8) + x*((c*e^3*(6*a - 3*b*c + 20*a*c^2)
)/2 + ((4*c^2 + 4)*((d^2*e^3*(b - 20*a*c))/4 + 2*a*c*d^2*e^3))/(4*d^2) - (
2*c*((2*c*((d^2*e^3*(b - 20*a*c))/4 + 2*a*c*d^2*e^3))/d + d*e^3*(a - b*c +
10*a*c^2) - (a*d*e^3*(4*c^2 + 4))/4))/d) + (a*d^3*e^3*x^4)/4 - (b*e^3*ata
n(((b*c*e^3*(c^2 + 1)*(c - 1)*(c + 1))/4 + (b*d*e^3*x*(c^2 + 1)*(c - 1)*(c
+ 1))/4)/((b*e^3)/4 - (b*c^4*e^3)/4))*(c^2 + 1)*(c - 1)*(c + 1))/(4*d)
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 164, normalized size of antiderivative = 2.28

$$\int (ce + dex)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{e^3 (3 \operatorname{atan}(dx + c) b c^4 + 12 \operatorname{atan}(dx + c) b c^3 dx + 18 \operatorname{atan}(dx + c) b c^2 d^2 x^2 + 12 \operatorname{atan}(dx + c) b c d^3 x^3 + 3 a d^3 x^4 - 3 a d^2 x^3 + 3 a d x^2 - 3 a x + 3 a)}{(12 d)}$$

input

```
int((d*e*x+c*e)^3*(a+b*atan(d*x+c)),x)
```

output

```
(e**3*(3*atan(c + d*x)*b*c**4 + 12*atan(c + d*x)*b*c**3*d*x + 18*atan(c +
d*x)*b*c**2*d**2*x**2 + 12*atan(c + d*x)*b*c*d**3*x**3 + 3*atan(c + d*x)*b
*d**4*x**4 - 3*atan(c + d*x)*b + 12*a*c**3*d*x + 18*a*c**2*d**2*x**2 + 12*
a*c*d**3*x**3 + 3*a*d**4*x**4 - 3*b*c**2*d*x - 3*b*c*d**2*x**2 - b*d**3*x*
*3 + 3*b*d*x))/(12*d)
```

3.10 $\int (ce + dex)^2(a + b \arctan(c + dx)) dx$

Optimal result	117
Mathematica [A] (verified)	117
Rubi [A] (warning: unable to verify)	118
Maple [A] (verified)	120
Fricas [B] (verification not implemented)	120
Sympy [C] (verification not implemented)	121
Maxima [B] (verification not implemented)	121
Giac [B] (verification not implemented)	122
Mupad [B] (verification not implemented)	123
Reduce [B] (verification not implemented)	123

Optimal result

Integrand size = 21, antiderivative size = 67

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx = -\frac{be^2(c + dx)^2}{6d} + \frac{e^2(c + dx)^3(a + b \arctan(c + dx))}{3d} + \frac{be^2 \log(1 + (c + dx)^2)}{6d}$$

output `-1/6*b*e^2*(d*x+c)^2/d+1/3*e^2*(d*x+c)^3*(a+b*arctan(d*x+c))/d+1/6*b*e^2*ln(1+(d*x+c)^2)/d`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.81

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx = \frac{e^2(\frac{1}{3}(c + dx)^3(a + b \arctan(c + dx)) - \frac{1}{6}b((c + dx)^2 - \log(1 + (c + dx)^2)))}{d}$$

input `Integrate[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x]),x]`

output

$$\frac{(e^{2*((c+dx)^3*(a+b*\text{ArcTan}[c+dx]))/3} - (b*((c+dx)^2 - \text{Log}[1+(c+dx)^2]))/6))/d}$$
Rubi [A] (warning: unable to verify)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5566, 27, 5361, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (ce + dex)^2 (a + b \arctan(c + dx)) dx \\ & \quad \downarrow \text{5566} \\ & \frac{\int e^2 (c + dx)^2 (a + b \arctan(c + dx)) d(c + dx)}{d} \\ & \quad \downarrow \text{27} \\ & \frac{e^2 \int (c + dx)^2 (a + b \arctan(c + dx)) d(c + dx)}{d} \\ & \quad \downarrow \text{5361} \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx)) - \frac{1}{3} b \int \frac{(c+dx)^3}{(c+dx)^2+1} d(c + dx) \right)}{d} \\ & \quad \downarrow \text{243} \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx)) - \frac{1}{6} b \int \frac{(c+dx)^2}{(c+dx)^2+1} d(c + dx)^2 \right)}{d} \\ & \quad \downarrow \text{49} \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx)) - \frac{1}{6} b \int \left(1 + \frac{1}{-c-dx-1} \right) d(c + dx)^2 \right)}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx)) - \frac{1}{6} b ((c + dx)^2 - \log(c + dx + 1)) \right)}{d} \end{aligned}$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x]),x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTan[c + d*x]))/3 - (b*((c + d*x)^2 - Log[1 + c + d*x]))/6))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 5566 `Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_.)), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{\frac{e^2 a (dx+c)^3}{3} + e^2 b \left(\frac{(dx+c)^3 \arctan(dx+c)}{3} - \frac{(dx+c)^2}{6} + \frac{\ln(1+(dx+c)^2)}{6} \right)}{d}$
default	$\frac{\frac{e^2 a (dx+c)^3}{3} + e^2 b \left(\frac{(dx+c)^3 \arctan(dx+c)}{3} - \frac{(dx+c)^2}{6} + \frac{\ln(1+(dx+c)^2)}{6} \right)}{d}$
parts	$\frac{e^2 a (dx+c)^3}{3d} + \frac{e^2 b \left(\frac{(dx+c)^3 \arctan(dx+c)}{3} - \frac{(dx+c)^2}{6} + \frac{\ln(1+(dx+c)^2)}{6} \right)}{d}$
parallelrisch	$\frac{2d^4 e^2 b x^3 \arctan(dx+c) + 2x^3 a d^4 e^2 + 6d^3 e^2 c b x^2 \arctan(dx+c) + 6x^2 a c d^3 e^2 + 6b c^2 e^2 x \arctan(dx+c) d^2 - x^2 b d^3 e^2 + 6x}{6d^2}$
risch	$-\frac{ie^2(dx+c)^3 b \ln(1+i(dx+c))}{6d} + \frac{ie^2 d^2 b x^3 \ln(1-i(dx+c))}{6} + \frac{ie^2 d b c x^2 \ln(1-i(dx+c))}{2} + \frac{ie^2 b c^2 x \ln(1-i(dx+c))}{2}$

input `int((d*e*x+c*e)^2*(a+b*arctan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(1/3*e^2*a*(d*x+c)^3+e^2*b*(1/3*(d*x+c)^3*arctan(d*x+c)-1/6*(d*x+c)^2+1/6*ln(1+(d*x+c)^2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(61) = 122.

Time = 0.12 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.93

$$\int (ce + dex)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{2ad^3e^2x^3 + (6ac - b)d^2e^2x^2 + 2(3ac^2 - bc)de^2x + be^2 \log(d^2x^2 + 2cdx + c^2 + 1) + 2(bd^3e^2x^3 + 3bcd^2e^2x + bcd^2e^2)}{6d}$$

input `integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c)),x, algorithm="fricas")`

output `1/6*(2*a*d^3*e^2*x^3 + (6*a*c - b)*d^2*e^2*x^2 + 2*(3*a*c^2 - b*c)*d*e^2*x + b*e^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 2*(b*d^3*e^2*x^3 + 3*b*c*d^2*e^2*x^2 + 3*b*c^2*d*e^2*x + b*c^3*e^2)*arctan(d*x + c))/d`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.17 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.72

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx$$

$$= \begin{cases} ac^2e^2x + acde^2x^2 + \frac{ad^2e^2x^3}{3} + \frac{bc^3e^2 \operatorname{atan}(c+dx)}{3d} + bc^2e^2x \operatorname{atan}(c + dx) + bcde^2x^2 \operatorname{atan}(c + dx) - \frac{bce^2x}{3} + \frac{b^2c^2e^2x^2}{6} \\ c^2e^2x(a + b \operatorname{atan}(c)) \end{cases}$$

input `integrate((d*e*x+c*e)**2*(a+b*atan(d*x+c)),x)`

output `Piecewise((a*c**2*e**2*x + a*c*d*e**2*x**2 + a*d**2*e**2*x**3/3 + b*c**3*e**2*atan(c + d*x)/(3*d) + b*c**2*e**2*x*atan(c + d*x) + b*c*d*e**2*x**2*atan(c + d*x) - b*c*e**2*x/3 + b*d**2*e**2*x**3*atan(c + d*x)/3 - b*d*e**2*x**2/6 + b*e**2*log(c/d + x - I/d)/(3*d) - I*b*e**2*atan(c + d*x)/(3*d), Ne(d, 0)), (c**2*e**2*x*(a + b*atan(c)), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. 2(61) = 122.

Time = 0.12 (sec) , antiderivative size = 238, normalized size of antiderivative = 3.55

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx = \frac{1}{3} ad^2e^2x^3 + acde^2x^2$$

$$+ \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2x+cd}{d}\right)}{d^3} - \frac{c \log(d^2x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) bcde^2$$

$$+ \frac{1}{6} \left(2x^3 \arctan(dx + c) - d \left(\frac{dx^2 - 4cx}{d^3} - \frac{2(c^3 - 3c) \arctan\left(\frac{d^2x+cd}{d}\right)}{d^4} + \frac{(3c^2 - 1) \log(d^2x^2 + 2cdx + c^2 + 1)}{d^4} \right) \right) bcde^2$$

$$+ ac^2e^2x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1))bc^2e^2}{2d}$$

input `integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output

```
1/3*a*d^2*e^2*x^3 + a*c*d*e^2*x^2 + (x^2*arctan(d*x + c) - d*(x/d^2 + (c^2
- 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3
))*b*c*d*e^2 + 1/6*(2*x^3*arctan(d*x + c) - d*((d*x^2 - 4*c*x)/d^3 - 2*(c^
3 - 3*c)*arctan((d^2*x + c*d)/d)/d^4 + (3*c^2 - 1)*log(d^2*x^2 + 2*c*d*x +
c^2 + 1)/d^4))*b*d^2*e^2 + a*c^2*e^2*x + 1/2*(2*(d*x + c)*arctan(d*x + c)
- log((d*x + c)^2 + 1))*b*c^2*e^2/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. $2(61) = 122$.

Time = 0.19 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.96

$$\int (ce + dex)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{2bd^3e^2x^3 \arctan(dx + c) + 2ad^3e^2x^3 + 6bcd^2e^2x^2 \arctan(dx + c) + 6acd^2e^2x^2 + 6bc^2de^2x \arctan(dx + c) + \dots}{d}$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c)),x, algorithm="giac")
```

output

```
1/6*(2*b*d^3*e^2*x^3*arctan(d*x + c) + 2*a*d^3*e^2*x^3 + 6*b*c*d^2*e^2*x^2
*arctan(d*x + c) + 6*a*c*d^2*e^2*x^2 + 6*b*c^2*d*e^2*x*arctan(d*x + c) - p
i*b*c^3*e^2*sgn(d*x + c) + pi*b*c^3*e^2*sgn(-d*x - c) + 6*a*c^2*d*e^2*x -
b*d^2*e^2*x^2 + b*c^3*e^2*arctan(d*x + c) - b*c^3*e^2*arctan(-d*x - c) - 2
*b*c*d*e^2*x + b*e^2*log((d*x + c)^2 + 1))/d
```

Mupad [B] (verification not implemented)

Time = 1.66 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.15

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx = \frac{ad^2e^2x^3}{3} - \frac{bce^2x}{3} + \frac{be^2 \ln(c^2 + 2cdx + d^2x^2 + 1)}{6d} + ac^2e^2x - \frac{bde^2x^2}{6} + bc^2e^2x \operatorname{atan}(c + dx) + acde^2x^2 + \frac{bc^3e^2 \operatorname{atan}(c + dx)}{3d} + \frac{bd^2e^2x^3 \operatorname{atan}(c + dx)}{3} + bcde^2x^2 \operatorname{atan}(c + dx)$$

input `int((c*e + d*e*x)^2*(a + b*atan(c + d*x)),x)`output `(a*d^2*e^2*x^3)/3 - (b*c*e^2*x)/3 + (b*e^2*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(6*d) + a*c^2*e^2*x - (b*d*e^2*x^2)/6 + b*c^2*e^2*x*atan(c + d*x) + a*c*d*e^2*x^2 + (b*c^3*e^2*atan(c + d*x))/(3*d) + (b*d^2*e^2*x^3*atan(c + d*x))/3 + b*c*d*e^2*x^2*atan(c + d*x)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.91

$$\int (ce + dex)^2(a + b \arctan(c + dx)) dx = \frac{e^2(2 \operatorname{atan}(dx + c)bc^3 + 6 \operatorname{atan}(dx + c)bc^2dx + 6 \operatorname{atan}(dx + c)bc d^2x^2 + 2 \operatorname{atan}(dx + c)bd^3x^3 + \log(d^2x^2 + 2cdx + c^2))}{6d}$$

input `int((d*e*x+c*e)^2*(a+b*atan(d*x+c)),x)`output `(e**2*(2*atan(c + d*x)*b*c**3 + 6*atan(c + d*x)*b*c**2*d*x + 6*atan(c + d*x)*b*c*d**2*x**2 + 2*atan(c + d*x)*b*d**3*x**3 + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b + 6*a*c**2*d*x + 6*a*c*d**2*x**2 + 2*a*d**3*x**3 - 2*b*c*d*x - b*d**2*x**2))/(6*d)`

3.11 $\int (ce + dex)(a + b \arctan(c + dx)) dx$

Optimal result	124
Mathematica [A] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	127
Fricas [A] (verification not implemented)	127
Sympy [B] (verification not implemented)	128
Maxima [B] (verification not implemented)	128
Giac [B] (verification not implemented)	129
Mupad [B] (verification not implemented)	129
Reduce [B] (verification not implemented)	130

Optimal result

Integrand size = 19, antiderivative size = 48

$$\int (ce + dex)(a + b \arctan(c + dx)) dx = -\frac{1}{2}bex + \frac{be \arctan(c + dx)}{2d} + \frac{e(c + dx)^2(a + b \arctan(c + dx))}{2d}$$

output `-1/2*b*e*x+1/2*b*e*arctan(d*x+c)/d+1/2*e*(d*x+c)^2*(a+b*arctan(d*x+c))/d`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int (ce + dex)(a + b \arctan(c + dx)) dx = \frac{e(b(-dx + \arctan(c + dx)) + (c + dx)^2(a + b \arctan(c + dx)))}{2d}$$

input `Integrate[(c*e + d*e*x)*(a + b*ArcTan[c + d*x]),x]`

output `(e*(b*(-(d*x) + ArcTan[c + d*x]) + (c + d*x)^2*(a + b*ArcTan[c + d*x]))) / (2*d)`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {5566, 27, 5361, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ce + dex)(a + b \arctan(c + dx)) dx \\
 & \quad \downarrow \text{5566} \\
 & \frac{\int e(c + dx)(a + b \arctan(c + dx))d(c + dx)}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{e \int (c + dx)(a + b \arctan(c + dx))d(c + dx)}{d} \\
 & \quad \downarrow \text{5361} \\
 & \frac{e \left(\frac{1}{2}(c + dx)^2(a + b \arctan(c + dx)) - \frac{1}{2}b \int \frac{(c+dx)^2}{(c+dx)^2+1} d(c + dx) \right)}{d} \\
 & \quad \downarrow \text{262} \\
 & \frac{e \left(\frac{1}{2}(c + dx)^2(a + b \arctan(c + dx)) - \frac{1}{2}b \left(- \int \frac{1}{(c+dx)^2+1} d(c + dx) + c + dx \right) \right)}{d} \\
 & \quad \downarrow \text{216} \\
 & \frac{e \left(\frac{1}{2}(c + dx)^2(a + b \arctan(c + dx)) - \frac{1}{2}b(- \arctan(c + dx) + c + dx) \right)}{d}
 \end{aligned}$$

input `Int[(c*e + d*e*x)*(a + b*ArcTan[c + d*x]),x]`

output `(e*(-1/2*(b*(c + d*x - ArcTan[c + d*x])) + ((c + d*x)^2*(a + b*ArcTan[c + d*x]))/2))/d`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 216 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$
- rule 262 $\text{Int}[((c_*)(x_))^{(m_)*((a_) + (b_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)*((a + b*x^2)^{(p+1})/(b*(m+2*p+1))}, x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \text{ Int}[(c*x)^{(m-2)*(a + b*x^2)^p}, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 5361 $\text{Int}[((a_.) + \text{ArcTan}[(c_*)(x_)^{(n_.)}]*(b_.))^{(p_.)*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)*((a + b*\text{ArcTan}[c*x^n])^p/(m+1))}, x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)*((a + b*\text{ArcTan}[c*x^n])^{(p-1)/(1+c^2*x^{2*n}))}, x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 5566 $\text{Int}[((a_.) + \text{ArcTan}[(c_) + (d_*)(x_)]*(b_.))^{(p_.)*((e_.) + (f_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06

method	result
derivativedivides	$\frac{ea(dx+c)^2 + eb\left(\frac{(dx+c)^2 \arctan(dx+c)}{2} - \frac{dx}{2} - \frac{c}{2} + \frac{\arctan(dx+c)}{2}\right)}{d}$
default	$\frac{ea(dx+c)^2 + eb\left(\frac{(dx+c)^2 \arctan(dx+c)}{2} - \frac{dx}{2} - \frac{c}{2} + \frac{\arctan(dx+c)}{2}\right)}{d}$
parts	$ea\left(\frac{1}{2}dx^2 + cx\right) + \frac{eb\left(\frac{(dx+c)^2 \arctan(dx+c)}{2} - \frac{dx}{2} - \frac{c}{2} + \frac{\arctan(dx+c)}{2}\right)}{d}$
parallelrisch	$\frac{d^3 eb x^2 \arctan(dx+c) + x^2 a d^3 e + 2ceb \arctan(dx+c)x d^2 + 2xac d^2 e + \arctan(dx+c)b c^2 de - xb d^2 e - 5a c^2 de + eb \arctan(dx+c)}{2d^2}$
risch	$-\frac{ieb(dx^2+2cx)\ln(1+i(dx+c))}{4} + \frac{iedbx^2\ln(1-i(dx+c))}{4} + \frac{iebcx\ln(1-i(dx+c))}{2} + \frac{ade x^2}{2} + \frac{e \arctan(dx+c)b}{2d}$
orering	$\frac{(2d^3 x^3 + 5c d^2 x^2 + 4c^2 dx + c^3 + 2dx + c)(dex + ce)(a + b \arctan(dx+c))}{2(dx+c)^2 d} - \frac{x(d^2 x^2 + 2cdx + c^2 + 1)(de(a + b \arctan(dx+c)))}{2d(dx+c)}$

input `int((d*e*x+c*e)*(a+b*arctan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(1/2*e*a*(d*x+c)^2+e*b*(1/2*(d*x+c)^2*arctan(d*x+c)-1/2*d*x-1/2*c+1/2*arctan(d*x+c)))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.25

$$\int (ce + dex)(a + b \arctan(c + dx)) dx$$

$$= \frac{ad^2 ex^2 + (2ac - b)dex + (bd^2 ex^2 + 2bcdex + (bc^2 + b)e) \arctan(dx + c)}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c)),x, algorithm="fricas")`

output `1/2*(a*d^2*e*x^2 + (2*a*c - b)*d*e*x + (b*d^2*e*x^2 + 2*b*c*d*e*x + (b*c^2 + b)*e)*arctan(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(41) = 82$.

Time = 0.61 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.98

$$\int (ce + dex)(a + b \arctan(c + dx)) dx$$

$$= \begin{cases} acex + \frac{adex^2}{2} + \frac{bc^2e \arctan(c+dx)}{2d} + bcex \operatorname{atan}(c + dx) + \frac{bdex^2 \operatorname{atan}(c+dx)}{2} - \frac{bex}{2} + \frac{be \operatorname{atan}(c+dx)}{2d} & \text{for } d \neq 0 \\ cex(a + b \operatorname{atan}(c)) & \text{otherwise} \end{cases}$$

input `integrate((d*e*x+c*e)*(a+b*atan(d*x+c)),x)`

output `Piecewise((a*c*e*x + a*d*e*x**2/2 + b*c**2*e*atan(c + d*x)/(2*d) + b*c*e*x*atan(c + d*x) + b*d*e*x**2*atan(c + d*x)/2 - b*e*x/2 + b*e*atan(c + d*x)/(2*d), Ne(d, 0)), (c*e*x*(a + b*atan(c)), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. $2(42) = 84$.

Time = 0.13 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.50

$$\int (ce + dex)(a + b \arctan(c + dx)) dx = \frac{1}{2} adex^2$$

$$+ \frac{1}{2} \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2x+cd}{d}\right)}{d^3} - \frac{c \log(d^2x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) bde$$

$$+ acex + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) bce}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output `1/2*a*d*e*x^2 + 1/2*(x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3)*b*d*e + a*c*e*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b*c*e/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(42) = 84$.

Time = 0.15 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.52

$$\int (ce + dex)(a + b \arctan(c + dx)) dx$$

$$= \frac{2bd^2ex^2 \arctan(dx + c) + 2ad^2ex^2 + 4bcdex \arctan(dx + c) + \pi bc^2 \operatorname{esgn}(dx + c) - \pi bc^2e + 4acdex - 2bd^2ex^2}{4d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output `1/4*(2*b*d^2*e*x^2*arctan(d*x + c) + 2*a*d^2*e*x^2 + 4*b*c*d*e*x*arctan(d*x + c) + pi*b*c^2*e*sgn(d*x + c) - pi*b*c^2*e + 4*a*c*d*e*x - 2*b*c^2*e*arctan(1/(d*x + c)) - 2*b*d*e*x + pi*b*e*sgn(d*x + c) - pi*b*e - 2*b*e*arctan(1/(d*x + c)))/d`

Mupad [B] (verification not implemented)

Time = 1.95 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.52

$$\int (ce + dex)(a + b \arctan(c + dx)) dx = acex - \frac{bex}{2} + \frac{be \operatorname{atan}(c + dx)}{2d}$$

$$+ \frac{adex^2}{2} + \frac{bc^2e \operatorname{atan}(c + dx)}{2d}$$

$$+ bce \operatorname{atan}(c + dx) + \frac{bdex^2 \operatorname{atan}(c + dx)}{2}$$

input `int((c*e + d*e*x)*(a + b*atan(c + d*x)),x)`

output `a*c*e*x - (b*e*x)/2 + (b*e*atan(c + d*x))/(2*d) + (a*d*e*x^2)/2 + (b*c^2*e*atan(c + d*x))/(2*d) + b*c*e*x*atan(c + d*x) + (b*d*e*x^2*atan(c + d*x))/2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.48

$$\int (ce + dex)(a + b \arctan(c + dx)) dx$$

$$= \frac{e(\operatorname{atan}(dx + c)bc^2 + 2\operatorname{atan}(dx + c)bcdx + \operatorname{atan}(dx + c)bd^2x^2 + \operatorname{atan}(dx + c)b + 2acdx + ad^2x^2 - bd)}{2d}$$

input `int((d*e*x+c*e)*(a+b*atan(d*x+c)),x)`

output `(e*(atan(c + d*x)*b*c**2 + 2*atan(c + d*x)*b*c*d*x + atan(c + d*x)*b*d**2*x**2 + atan(c + d*x)*b + 2*a*c*d*x + a*d**2*x**2 - b*d*x))/(2*d)`

3.12 $\int \frac{a+b \arctan(c+dx)}{ce+dex} dx$

Optimal result	131
Mathematica [A] (verified)	131
Rubi [A] (verified)	132
Maple [A] (verified)	133
Fricas [F]	134
Sympy [F]	134
Maxima [F]	134
Giac [F]	135
Mupad [F(-1)]	135
Reduce [F]	135

Optimal result

Integrand size = 21, antiderivative size = 63

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \frac{a \log(c + dx)}{de} + \frac{ib \operatorname{PolyLog}(2, -i(c + dx))}{2de} - \frac{ib \operatorname{PolyLog}(2, i(c + dx))}{2de}$$

output

```
a*ln(d*x+c)/d/e+1/2*I*b*polylog(2,-I*(d*x+c))/d/e-1/2*I*b*polylog(2,I*(d*x+c))/d/e
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \frac{a \log(c + dx) + \frac{1}{2}ib \operatorname{PolyLog}(2, -i(c + dx)) - \frac{1}{2}ib \operatorname{PolyLog}(2, i(c + dx))}{de}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])/(c*e + d*e*x),x]
```

output

$$\frac{(a \cdot \text{Log}[c + d \cdot x] + (I/2) \cdot b \cdot \text{PolyLog}[2, (-I) \cdot (c + d \cdot x)] - (I/2) \cdot b \cdot \text{PolyLog}[2, I \cdot (c + d \cdot x)])}{(d \cdot e)}$$
Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {5566, 27, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \arctan(c + dx)}{ce + dex} dx \\ & \quad \downarrow \text{5566} \\ & \frac{\int \frac{a + b \arctan(c + dx)}{e(c + dx)} d(c + dx)}{d} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{a + b \arctan(c + dx)}{c + dx} d(c + dx)}{de} \\ & \quad \downarrow \text{5355} \\ & \frac{\frac{1}{2} ib \int \frac{\log(1 - i(c + dx))}{c + dx} d(c + dx) - \frac{1}{2} ib \int \frac{\log(i(c + dx) + 1)}{c + dx} d(c + dx) + a \log(c + dx)}{de} \\ & \quad \downarrow \text{2838} \\ & \frac{a \log(c + dx) + \frac{1}{2} ib \text{PolyLog}(2, -i(c + dx)) - \frac{1}{2} ib \text{PolyLog}(2, i(c + dx))}{de} \end{aligned}$$

input

$$\text{Int}[(a + b \cdot \text{ArcTan}[c + d \cdot x]) / (c \cdot e + d \cdot e \cdot x), x]$$

output

$$\frac{(a \cdot \text{Log}[c + d \cdot x] + (I/2) \cdot b \cdot \text{PolyLog}[2, (-I) \cdot (c + d \cdot x)] - (I/2) \cdot b \cdot \text{PolyLog}[2, I \cdot (c + d \cdot x)])}{(d \cdot e)}$$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 2838 $\text{Int}[\text{Log}[(c_*)((d_) + (e_)*(x_)^{(n_)})]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c*d, 1]$

rule 5355 $\text{Int}[(a_.) + \text{ArcTan}[(c_)*(x_)]*(b_.)]/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] + (\text{Simp}[I*(b/2) \text{ Int}[\text{Log}[1 - I*c*x]/x, x] - \text{Simp}[I*(b/2) \text{ Int}[\text{Log}[1 + I*c*x]/x, x], x]) /; \text{FreeQ}[\{a, b, c\}, x]$

rule 5566 $\text{Int}[(a_.) + \text{ArcTan}[(c_) + (d_)*(x_)]*(b_.)^{(p_)}*((e_.) + (f_)*(x_))^{(m_)}], x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^{m*(a + b*\text{ArcTan}[x])^p}, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

method	result
risch	$-\frac{ib \operatorname{dilog}(-idx-ic+1)}{2ed} + \frac{a \ln(-idx-ic)}{ed} + \frac{ib \operatorname{dilog}(idx+ic+1)}{2ed}$
derivativedivides	$\frac{\frac{a \ln(dx+c)}{e} + \frac{b \left(\ln(dx+c) \arctan(dx+c) + \frac{i \ln(dx+c) \ln(1+i(dx+c))}{2} - \frac{i \ln(dx+c) \ln(1-i(dx+c))}{2} + \frac{i \operatorname{dilog}(1+i(dx+c))}{2} - \frac{i \operatorname{dilog}(1-i(dx+c))}{2} \right)}{d}}{e}$
default	$\frac{\frac{a \ln(dx+c)}{e} + \frac{b \left(\ln(dx+c) \arctan(dx+c) + \frac{i \ln(dx+c) \ln(1+i(dx+c))}{2} - \frac{i \ln(dx+c) \ln(1-i(dx+c))}{2} + \frac{i \operatorname{dilog}(1+i(dx+c))}{2} - \frac{i \operatorname{dilog}(1-i(dx+c))}{2} \right)}{d}}{e}$
parts	$\frac{a \ln(dx+c)}{de} + \frac{b \left(\ln(dx+c) \arctan(dx+c) + \frac{i \ln(dx+c) \ln(1+i(dx+c))}{2} - \frac{i \ln(dx+c) \ln(1-i(dx+c))}{2} + \frac{i \operatorname{dilog}(1+i(dx+c))}{2} - \frac{i \operatorname{dilog}(1-i(dx+c))}{2} \right)}{ed}$

input $\text{int}((a+b*\arctan(d*x+c))/(d*e*x+c*e), x, \text{method}=_RETURNVERBOSE)$

output $-1/2*I/e/d*b*\operatorname{dilog}(1-I*c-I*d*x)+1/e/d*a*\ln(-I*d*x-I*c)+1/2*I*b/e/d*\operatorname{dilog}(1+I*c+I*d*x)$

Fricas [F]

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \int \frac{b \arctan(dx + c) + a}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e),x, algorithm="fricas")`

output `integral((b*arctan(d*x + c) + a)/(d*e*x + c*e), x)`

Sympy [F]

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \frac{\int \frac{a}{c+dx} dx + \int \frac{b \operatorname{atan}(c+dx)}{c+dx} dx}{e}$$

input `integrate((a+b*atan(d*x+c))/(d*e*x+c*e),x)`

output `(Integral(a/(c + d*x), x) + Integral(b*atan(c + d*x)/(c + d*x), x))/e`

Maxima [F]

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \int \frac{b \arctan(dx + c) + a}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e),x, algorithm="maxima")`

output `2*b*integrate(1/2*arctan(d*x + c)/(d*e*x + c*e), x) + a*log(d*e*x + c*e)/(d*e)`

Giac [F]

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \int \frac{b \arctan(dx + c) + a}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)/(d*e*x + c*e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \int \frac{a + b \operatorname{atan}(c + dx)}{ce + dex} dx$$

input `int((a + b*atan(c + d*x))/(c*e + d*e*x),x)`

output `int((a + b*atan(c + d*x))/(c*e + d*e*x), x)`

Reduce [F]

$$\int \frac{a + b \arctan(c + dx)}{ce + dex} dx = \frac{\left(\int \frac{\operatorname{atan}(dx+c)}{dx+c} dx \right) bd + \log(dx + c) a}{de}$$

input `int((a+b*atan(d*x+c))/(d*e*x+c*e),x)`

output `(int(atan(c + d*x)/(c + d*x),x)*b*d + log(c + d*x)*a)/(d*e)`

3.13 $\int \frac{a+b \arctan(c+dx)}{(ce+dex)^2} dx$

Optimal result	136
Mathematica [A] (verified)	136
Rubi [A] (verified)	137
Maple [A] (verified)	139
Fricas [A] (verification not implemented)	139
Sympy [C] (verification not implemented)	140
Maxima [A] (verification not implemented)	140
Giac [B] (verification not implemented)	141
Mupad [B] (verification not implemented)	141
Reduce [B] (verification not implemented)	142

Optimal result

Integrand size = 21, antiderivative size = 61

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx = -\frac{a + b \arctan(c + dx)}{de^2(c + dx)} + \frac{b \log(c + dx)}{de^2} - \frac{b \log(1 + (c + dx)^2)}{2de^2}$$

output

$$-(a+b*\arctan(d*x+c))/d/e^2/(d*x+c)+b*\ln(d*x+c)/d/e^2-1/2*b*\ln(1+(d*x+c)^2)/d/e^2$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.85

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx = \frac{\frac{-a-b \arctan(c+dx)}{c+dx} + b(\log(c + dx) - \frac{1}{2} \log(1 + (c + dx)^2))}{de^2}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])/(c*e + d*e*x)^2,x]
```

output

$$((-a - b*ArcTan[c + d*x])/(c + d*x) + b*(Log[c + d*x] - Log[1 + (c + d*x)^2])/2)/(d*e^2)$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5566, 27, 5361, 243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx \\
 & \quad \downarrow \text{5566} \\
 & \frac{\int \frac{a+b \arctan(c+dx)}{e^2(c+dx)^2} d(c + dx)}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a+b \arctan(c+dx)}{(c+dx)^2} d(c + dx)}{de^2} \\
 & \quad \downarrow \text{5361} \\
 & \frac{b \int \frac{1}{(c+dx)((c+dx)^2+1)} d(c + dx) - \frac{a+b \arctan(c+dx)}{c+dx}}{de^2} \\
 & \quad \downarrow \text{243} \\
 & \frac{\frac{1}{2} b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c + dx)^2 - \frac{a+b \arctan(c+dx)}{c+dx}}{de^2} \\
 & \quad \downarrow \text{47} \\
 & \frac{\frac{1}{2} b \left(\int \frac{1}{(c+dx)^2} d(c + dx)^2 - \int \frac{1}{(c+dx)^2+1} d(c + dx)^2 \right) - \frac{a+b \arctan(c+dx)}{c+dx}}{de^2} \\
 & \quad \downarrow \text{14} \\
 & \frac{\frac{1}{2} b \left(\log((c + dx)^2) - \int \frac{1}{(c+dx)^2+1} d(c + dx)^2 \right) - \frac{a+b \arctan(c+dx)}{c+dx}}{de^2} \\
 & \quad \downarrow \text{16} \\
 & \frac{\frac{1}{2} b (\log((c + dx)^2) - \log((c + dx)^2 + 1)) - \frac{a+b \arctan(c+dx)}{c+dx}}{de^2}
 \end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])/(c*e + d*e*x)^2,x]`

output `((-(a + b*ArcTan[c + d*x])/(c + d*x)) + (b*(Log[(c + d*x)^2] - Log[1 + (c + d*x)^2]))/2)/(d*e^2)`

Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 47 `Int[1/(((a_) + (b_)*(x_))*((c_) + (d_)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 5361 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & IntegerQ[m])) && NeQ[m, -1]`

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.95

method	result
derivativdivides	$-\frac{a}{e^2(dx+c)} + \frac{b \left(-\frac{\arctan(dx+c)}{dx+c} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{d e^2}$
default	$-\frac{a}{e^2(dx+c)} + \frac{b \left(-\frac{\arctan(dx+c)}{dx+c} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{d e^2}$
parts	$-\frac{a}{d e^2(dx+c)} + \frac{b \left(-\frac{\arctan(dx+c)}{dx+c} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{e^2 d}$
parallelrisc	$\frac{6 \ln(dx+c) x b c d^2 - 3 \ln(d^2 x^2 + 2 c d x + c^2 + 1) x b c d^2 + 6 \ln(dx+c) b c^2 d - 3 \ln(d^2 x^2 + 2 c d x + c^2 + 1) b c^2 d + 2 x a d^2 - 6 b \arctan(dx+c)}{6(dx+c)c d^2 e^2}$
risc	$\frac{i b \ln(1+i(dx+c))}{2 d e^2(dx+c)} - \frac{-2 \ln(-dx-c) b d x + \ln(-d^2 x^2 - 2 c d x - c^2 - 1) b d x + i b \ln(1-i(dx+c)) - 2 \ln(-dx-c) b c + \ln(-d^2 x^2 - 2 c d x - c^2 - 1) b c}{2 e^2(dx+c) d}$

input

```
int((a+b*arctan(d*x+c))/(d*e*x+c*e)^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-a/e^2/(d*x+c)+b/e^2*(-1/(d*x+c)*arctan(d*x+c)+ln(d*x+c)-1/2*ln(1+(d*x+c)^2)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx = \frac{2 b \arctan(dx + c) + (bdx + bc) \log(d^2 x^2 + 2 c d x + c^2 + 1) - 2 (bdx + bc) \log(dx + c) + 2 a}{2 (d^2 e^2 x + c d e^2)}$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^2,x, algorithm="fricas")`

output
$$-1/2*(2*b*arctan(d*x + c) + (b*d*x + b*c)*\log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 2*(b*d*x + b*c)*\log(d*x + c) + 2*a)/(d^2*e^2*x + c*d*e^2)$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.22 (sec) , antiderivative size = 223, normalized size of antiderivative = 3.66

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx$$

$$= \begin{cases} -\frac{a}{cde^2+d^2e^2x} + \frac{bc \log\left(\frac{c}{d}+x\right)}{cde^2+d^2e^2x} - \frac{bc \log\left(\frac{c}{d}+x-\frac{i}{d}\right)}{cde^2+d^2e^2x} + \frac{ibc \operatorname{atan}(c+dx)}{cde^2+d^2e^2x} + \frac{bdx \log\left(\frac{c}{d}+x\right)}{cde^2+d^2e^2x} - \frac{bdx \log\left(\frac{c}{d}+x-\frac{i}{d}\right)}{cde^2+d^2e^2x} + \frac{ibdx \operatorname{atan}(c+dx)}{cde^2+d^2e^2x} \\ \frac{x(a+b \operatorname{atan}(c))}{c^2e^2} \end{cases}$$

input `integrate((a+b*atan(d*x+c))/(d*e*x+c*e)**2,x)`

output `Piecewise((-a/(c*d*e**2 + d**2*e**2*x) + b*c*log(c/d + x)/(c*d*e**2 + d**2*e**2*x) - b*c*log(c/d + x - I/d)/(c*d*e**2 + d**2*e**2*x) + I*b*c*atan(c + d*x)/(c*d*e**2 + d**2*e**2*x) + b*d*x*log(c/d + x)/(c*d*e**2 + d**2*e**2*x) - b*d*x*log(c/d + x - I/d)/(c*d*e**2 + d**2*e**2*x) + I*b*d*x*atan(c + d*x)/(c*d*e**2 + d**2*e**2*x) - b*atan(c + d*x)/(c*d*e**2 + d**2*e**2*x), Ne(d, 0)), (x*(a + b*atan(c))/(c**2*e**2), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.51

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx$$

$$= -\frac{1}{2} \left(d \left(\frac{\log(d^2x^2 + 2cdx + c^2 + 1)}{d^2e^2} - \frac{2 \log(dx + c)}{d^2e^2} \right) + \frac{2 \arctan(dx + c)}{d^2e^2x + cde^2} \right) b$$

$$- \frac{a}{d^2e^2x + cde^2}$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^2,x, algorithm="maxima")`

output
$$-1/2*(d*(\log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2) - 2*\log(d*x + c)/(d^2*e^2)) + 2*arctan(d*x + c)/(d^2*e^2*x + c*d*e^2))*b - a/(d^2*e^2*x + c*d*e^2)$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 175 vs. $2(59) = 118$.

Time = 0.17 (sec) , antiderivative size = 175, normalized size of antiderivative = 2.87

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx = -\frac{a}{(dex + ce)de} + \frac{\left(\arctan\left(\frac{dex+ce}{e}\right) \tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)^2 + \log\left(\frac{16 \tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)^2}{\tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)^4 + 2 \tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)^2 + 1}\right) \tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)}{2 de^2 \tan\left(\frac{1}{2} \arctan\left(\frac{dex+ce}{e}\right)\right)}$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^2,x, algorithm="giac")`

output
$$-a/((d*e*x + c*e)*d*e) + 1/2*(\arctan((d*e*x + c*e)/e)*\tan(1/2*\arctan((d*e*x + c*e)/e))^2 + \log(16*\tan(1/2*\arctan((d*e*x + c*e)/e))^2/(\tan(1/2*\arctan((d*e*x + c*e)/e))^4 + 2*\tan(1/2*\arctan((d*e*x + c*e)/e))^2 + 1))*\tan(1/2*\arctan((d*e*x + c*e)/e)) - \arctan((d*e*x + c*e)/e)*b/(d*e^2*\tan(1/2*\arctan((d*e*x + c*e)/e)))$$

Mupad [B] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.44

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx = \frac{b \ln(c + dx)}{de^2} - \frac{b \operatorname{atan}(c + dx)}{x d^2 e^2 + c d e^2} - \frac{b \ln(c^2 + 2 c d x + d^2 x^2 + 1)}{2 d e^2} - \frac{a}{x d^2 e^2 + c d e^2}$$

input `int((a + b*atan(c + d*x))/(c*e + d*e*x)^2,x)`

output `(b*log(c + d*x))/(d*e^2) - (b*atan(c + d*x))/(d^2*e^2*x + c*d*e^2) - (b*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(2*d*e^2) - a/(d^2*e^2*x + c*d*e^2)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.74

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^2} dx$$

$$= \frac{-2 \operatorname{atan}(dx + c) bc - \log(d^2 x^2 + 2cdx + c^2 + 1) b c^2 - \log(d^2 x^2 + 2cdx + c^2 + 1) bcdx + 2 \log(dx + c) b c}{2cd e^2 (dx + c)}$$

input `int((a+b*atan(d*x+c))/(d*e*x+c*e)^2,x)`

output `(- 2*atan(c + d*x)*b*c - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c**2 - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d*x + 2*log(c + d*x)*b*c**2 + 2*log(c + d*x)*b*c*d*x + 2*a*d*x)/(2*c*d*e**2*(c + d*x))`

3.14 $\int \frac{a+b \arctan(c+dx)}{(ce+dex)^3} dx$

Optimal result	143
Mathematica [C] (verified)	143
Rubi [A] (verified)	144
Maple [A] (verified)	146
Fricas [A] (verification not implemented)	146
Sympy [B] (verification not implemented)	147
Maxima [B] (verification not implemented)	147
Giac [C] (verification not implemented)	148
Mupad [B] (verification not implemented)	148
Reduce [B] (verification not implemented)	149

Optimal result

Integrand size = 21, antiderivative size = 63

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx = -\frac{b}{2de^3(c + dx)} - \frac{b \arctan(c + dx)}{2de^3} - \frac{a + b \arctan(c + dx)}{2de^3(c + dx)^2}$$

output `-1/2*b/d/e^3/(d*x+c)-1/2*b*arctan(d*x+c)/d/e^3-1/2*(a+b*arctan(d*x+c))/d/e^3/(d*x+c)^2`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx = -\frac{a + b \arctan(c + dx) + b(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -(c + dx)^2\right)}{2de^3(c + dx)^2}$$

input `Integrate[(a + b*ArcTan[c + d*x])/(c*e + d*e*x)^3,x]`

output

$$-1/2*(a + b*\text{ArcTan}[c + d*x] + b*(c + d*x)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, -(c + d*x)^2])/(d*e^3*(c + d*x)^2)$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {5566, 27, 5361, 264, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx \\ & \quad \downarrow \text{5566} \\ & \frac{\int \frac{a+b \arctan(c+dx)}{e^3(c+dx)^3} d(c + dx)}{d} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{a+b \arctan(c+dx)}{(c+dx)^3} d(c + dx)}{de^3} \\ & \quad \downarrow \text{5361} \\ & \frac{\frac{1}{2}b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c + dx) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2}}{de^3} \\ & \quad \downarrow \text{264} \\ & \frac{\frac{1}{2}b \left(- \int \frac{1}{(c+dx)^2+1} d(c + dx) - \frac{1}{c+dx} \right) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2}}{de^3} \\ & \quad \downarrow \text{216} \\ & \frac{\frac{1}{2}b \left(- \arctan(c + dx) - \frac{1}{c+dx} \right) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2}}{de^3} \end{aligned}$$

input

$$\text{Int}[(a + b*\text{ArcTan}[c + d*x])/(c*e + d*e*x)^3, x]$$

output $((b*(-(c + d*x)^{-1}) - \text{ArcTan}[c + d*x])/2 - (a + b*\text{ArcTan}[c + d*x])/(2*(c + d*x)^2))/(d*e^3)$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 216 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 264 $\text{Int}[(c_)*(x_)^m*((a_) + (b_)*(x_)^2)^p], x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}*((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^2*(m+1))) \text{Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 5361 $\text{Int}[(a_) + \text{ArcTan}[c_)*(x_)^{n_}]*b_)^p*(x_)^m], x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTan}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{Int}[x^{(m+n)}*((a + b*\text{ArcTan}[c*x^n])^{(p-1)})/(1 + c^2*x^{(2*n)}), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 5566 $\text{Int}[(a_) + \text{ArcTan}[c_] + (d_)*(x_)]*b_)^p*(e_) + (f_)*(x_)^m], x_Symbol] \rightarrow \text{Simp}[1/d \ \text{Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90

method	result
derivativdivides	$-\frac{a}{2e^3(dx+c)^2} + \frac{b\left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{1}{2(dx+c)} - \frac{\arctan(dx+c)}{2}\right)}{e^3}$
default	$-\frac{a}{2e^3(dx+c)^2} + \frac{b\left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{1}{2(dx+c)} - \frac{\arctan(dx+c)}{2}\right)}{e^3}$
parts	$-\frac{a}{2e^3(dx+c)^2 d} + \frac{b\left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{1}{2(dx+c)} - \frac{\arctan(dx+c)}{2}\right)}{e^3 d}$
parallelrisch	$\frac{-4b d^4 \arctan(dx+c)x^2 c - 8b c^2 \arctan(dx+c)x d^3 + b d^4 x^2 - 4 \arctan(dx+c)b c^3 d^2 - 2x b c d^3 - 4b \arctan(dx+c)c d^2 - 3b c^2 d^3}{8(dx+c)^2 e^3 c d^3}$
oring	$-\frac{2(d^3 x^3 + 3c d^2 x^2 + 3c^2 dx + c^3 + dx+c)(a+b \arctan(dx+c))}{d(dx+ce)^3} - \frac{(d^2 x^2 + 2cdx + c^2 + 1)(dx+c)^2 \left(\frac{bd}{(1+(dx+c)^2)(dex+ce)^3}\right)}{2d^2}$
risch	$\frac{ib \ln(1+i(dx+c))}{4d e^3(dx+c)^2} - \frac{i \ln(-dx-c-i)b d^2 x^2 - i \ln(-dx-c+i)b d^2 x^2 + 2i \ln(-dx-c-i)b c d x - 2i \ln(-dx-c+i)b c d x + i \ln(-dx-c-i)b c d x + i \ln(-dx-c+i)b c d x}{4e^3(dx+c)^2 d}$

input

```
int((a+b*arctan(d*x+c))/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-1/2*a/e^3/(d*x+c)^2+b/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)-1/2/(d*x+c)-1/2*arctan(d*x+c)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx = -\frac{bdx + bc + (bd^2x^2 + 2bcdx + bc^2 + b) \arctan(dx + c) + a}{2(d^3e^3x^2 + 2cd^2e^3x + c^2de^3)}$$

input

```
integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^3,x, algorithm="fricas")
```

output

```
-1/2*(b*d*x + b*c + (b*d^2*x^2 + 2*b*c*d*x + b*c^2 + b)*arctan(d*x + c) + a)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 314 vs. $2(54) = 108$.

Time = 3.75 (sec) , antiderivative size = 314, normalized size of antiderivative = 4.98

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx$$

$$= \left\{ \begin{array}{l} -\frac{a}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} - \frac{bc^2 \operatorname{atan}(c+dx)}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} - \frac{2bcdx \operatorname{atan}(c+dx)}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} - \frac{bc}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} - \frac{bd^2x^2}{2c^2de^3 + 4cd^2e^3x + 2d^3e^3x^2} \\ \frac{x(a+b \operatorname{atan}(c))}{c^3e^3} \end{array} \right.$$

input `integrate((a+b*atan(d*x+c))/(d*e*x+c*e)**3,x)`

output

```
Piecewise((-a/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*c**2*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - 2*b*c*d*x*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*c/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*d**2*x**2*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*d*x/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2), Ne(d, 0)), (x*(a + b*atan(c))/(c**3*e**3), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. $2(57) = 114$.

Time = 0.11 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.90

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx$$

$$= -\frac{1}{2} \left(d \left(\frac{1}{d^3e^3x + cd^2e^3} + \frac{\arctan\left(\frac{d^2x+cd}{d}\right)}{d^2e^3} \right) + \frac{\arctan(dx + c)}{d^3e^3x^2 + 2cd^2e^3x + c^2de^3} \right) b$$

$$- \frac{a}{2(d^3e^3x^2 + 2cd^2e^3x + c^2de^3)}$$

input `integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^3,x, algorithm="maxima")`

output

$$-1/2*(d*(1/(d^3*e^3*x + c*d^2*e^3) + \arctan((d^2*x + c*d)/d)/(d^2*e^3)) + \arctan(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3))*b - 1/2*a/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)$$
Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.48

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx = \frac{bd^2x^2 \log(ix + ic + 1) - bd^2x^2 \log(-ix - ic + 1) + 2bcdx \log(ix + ic + 1) - 2bcdx \log(-ix - ic + 1)}{-4id^3e^3x^2 - 8icd^2e^3x - 4c^2de^3}$$

input

`integrate((a+b*arctan(d*x+c))/(d*e*x+c*e)^3,x, algorithm="giac")`

output

$$(b*d^2*x^2*\log(I*d*x + I*c + 1) - b*d^2*x^2*\log(-I*d*x - I*c + 1) + 2*b*c*d*x*\log(I*d*x + I*c + 1) - 2*b*c*d*x*\log(-I*d*x - I*c + 1) + b*c^2*\log(I*d*x + I*c + 1) - b*c^2*\log(-I*d*x - I*c + 1) + 2*I*b*d*x + 2*I*b*c + 2*I*b*\arctan(d*x + c) + 2*I*a)/(-4*I*d^3*e^3*x^2 - 8*I*c*d^2*e^3*x - 4*I*c^2*d*e^3)$$
Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.63

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx = -\frac{\frac{a+bc}{d} + bx}{2c^2e^3 + 4cde^3x + 2d^2e^3x^2} - \frac{b \operatorname{atan}\left(\frac{bc+bdx}{b}\right)}{2de^3} - \frac{b \operatorname{atan}(c + dx)}{2d^3e^3\left(x^2 + \frac{c^2}{d^2} + \frac{2cx}{d}\right)}$$

input

`int((a + b*atan(c + d*x))/(c*e + d*e*x)^3,x)`

output

```
- ((a + b*c)/d + b*x)/(2*c^2*e^3 + 2*d^2*e^3*x^2 + 4*c*d*e^3*x) - (b*atan(
(b*c + b*d*x)/b))/(2*d*e^3) - (b*atan(c + d*x))/(2*d^3*e^3*(x^2 + c^2/d^2
+ (2*c*x)/d))
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.59

$$\int \frac{a + b \arctan(c + dx)}{(ce + dex)^3} dx$$

$$= \frac{-2 \operatorname{atan}(dx + c) b c^3 - 4 \operatorname{atan}(dx + c) b c^2 dx - 2 \operatorname{atan}(dx + c) b c d^2 x^2 - 2 \operatorname{atan}(dx + c) b c - 2 a c - b c^2 + b^2}{4 c d e^3 (d^2 x^2 + 2 c d x + c^2)}$$

input

```
int((a+b*atan(d*x+c))/(d*e*x+c*e)^3,x)
```

output

```
( - 2*atan(c + d*x)*b*c**3 - 4*atan(c + d*x)*b*c**2*d*x - 2*atan(c + d*x)*
b*c*d**2*x**2 - 2*atan(c + d*x)*b*c - 2*a*c - b*c**2 + b*d**2*x**2)/(4*c*d
*e**3*(c**2 + 2*c*d*x + d**2*x**2))
```

3.15 $\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$

Optimal result	150
Mathematica [A] (verified)	151
Rubi [A] (warning: unable to verify)	151
Maple [A] (verified)	154
Fricas [B] (verification not implemented)	155
Sympy [C] (verification not implemented)	155
Maxima [B] (verification not implemented)	156
Giac [B] (verification not implemented)	157
Mupad [B] (verification not implemented)	159
Reduce [B] (verification not implemented)	160

Optimal result

Integrand size = 23, antiderivative size = 157

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx = \frac{1}{2}abe^3x + \frac{b^2e^3(c + dx)^2}{12d} + \frac{b^2e^3(c + dx) \arctan(c + dx)}{2d} - \frac{be^3(c + dx)^3(a + b \arctan(c + dx))}{6d} - \frac{e^3(a + b \arctan(c + dx))^2}{4d} + \frac{e^3(c + dx)^4(a + b \arctan(c + dx))^2}{4d} - \frac{b^2e^3 \log(1 + (c + dx)^2)}{3d}$$

output

```
1/2*a*b*e^3*x+1/12*b^2*e^3*(d*x+c)^2/d+1/2*b^2*e^3*(d*x+c)*arctan(d*x+c)/d
-1/6*b*e^3*(d*x+c)^3*(a+b*arctan(d*x+c))/d-1/4*e^3*(a+b*arctan(d*x+c))^2/d
+1/4*e^3*(d*x+c)^4*(a+b*arctan(d*x+c))^2/d-1/3*b^2*e^3*ln(1+(d*x+c)^2)/d
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.38

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{e^3((c + dx)(b^2(c + dx) + 3a^2(c + dx)^3 - 2ab(-3 + c^2 + 2cdx + d^2x^2)) + 2b(-b(-3c + c^3 - 3dx + 3c^2d$$

input `Integrate[(c*e + d*e*x)^3*(a + b*ArcTan[c + d*x])^2,x]`output
$$\frac{(e^3((c + d*x)*(b^2*(c + d*x) + 3*a^2*(c + d*x)^3 - 2*a*b*(-3 + c^2 + 2*c*d*x + d^2*x^2)) + 2*b*(-(b*(-3*c + c^3 - 3*d*x + 3*c^2*d*x + 3*c*d^2*x^2 + d^3*x^3)) + 3*a*(-1 + c^4 + 4*c^3*d*x + 6*c^2*d^2*x^2 + 4*c*d^3*x^3 + d^4*x^4))*ArcTan[c + d*x] + 3*b^2*(-1 + c^4 + 4*c^3*d*x + 6*c^2*d^2*x^2 + 4*c*d^3*x^3 + d^4*x^4)*ArcTan[c + d*x]^2 - 4*b^2*Log[1 + (c + d*x)^2]))/(12*d)$$
Rubi [A] (warning: unable to verify)Time = 0.75 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.87, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$, Rules used = {5566, 27, 5361, 5451, 5361, 243, 49, 2009, 5451, 2009, 5419}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$\downarrow \text{5566}$$

$$\frac{\int e^3 (c + dx)^3 (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow \text{27}$$

$$\frac{e^3 \int (c + dx)^3 (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow \text{5361}$$

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \int \frac{(c+dx)^4(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) \right)}{d}$$

↓ 5451

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(\int (c+dx)^2(a+b \arctan(c+dx)) d(c+dx) - \int \frac{(c+dx)^2(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) \right) \right)}{d}$$

↓ 5361

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(- \int \frac{(c+dx)^2(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{1}{3}b \int \frac{(c+dx)^3}{(c+dx)^2+1} d(c+dx) + \frac{1}{3} \right) \right)}{d}$$

↓ 243

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(- \int \frac{(c+dx)^2(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{1}{6}b \int \frac{(c+dx)^2}{(c+dx)^2+1} d(c+dx)^2 + \frac{1}{3} \right) \right)}{d}$$

↓ 49

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(- \int \frac{(c+dx)^2(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{1}{6}b \int \left(1 + \frac{1}{-c-dx-1} \right) d(c+dx) \right) \right)}{d}$$

↓ 2009

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(- \int \frac{(c+dx)^2(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) + \frac{1}{3}(c+dx)^3(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 5451

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(- \int (a+b \arctan(c+dx)) d(c+dx) + \int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) \right) \right)}{d}$$

↓ 2009

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b \arctan(c+dx))^2 - \frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{3}(c+dx)^3(a+b \arctan(c+dx)) - a \right) \right)}{d}$$

↓ 5419

$$\frac{e^3 \left(\frac{1}{4}(c+dx)^4(a+b\arctan(c+dx))^2 - \frac{1}{2}b \left(\frac{1}{3}(c+dx)^3(a+b\arctan(c+dx)) + \frac{(a+b\arctan(c+dx))^2}{2b} - a(c+dx) - \log[1+(c+dx)^2] \right) \right)}{d}$$

input `Int[(c*e + d*e*x)^3*(a + b*ArcTan[c + d*x])^2,x]`

output `(e^3*(((c + d*x)^4*(a + b*ArcTan[c + d*x])^2)/4 - (b*(-(a*(c + d*x)) - b*(c + d*x)*ArcTan[c + d*x] + ((c + d*x)^3*(a + b*ArcTan[c + d*x]))/3 + (a + b*ArcTan[c + d*x])^2/(2*b) - (b*((c + d*x)^2 - Log[1 + c + d*x]))/6 + (b*Log[1 + (c + d*x)^2])/2))/2))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 49 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & IntegerQ[m])) && NeQ[m, -1]`

- rule 5419 $\text{Int}[\left((a_{.}) + \text{ArcTan}[(c_{.})*(x_{.})]*(b_{.})\right)^{(p_{.})}/\left((d_{.}) + (e_{.})*(x_{.})^2\right), x_Symbol]$ $\rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^{(p + 1)}/(b*c*d*(p + 1)), x]$ /; $\text{FreeQ}\{a, b, c, d, e, p\}, x$ && $\text{EqQ}[e, c^2*d]$ && $\text{NeQ}[p, -1]$
- rule 5451 $\text{Int}[\left(\left((a_{.}) + \text{ArcTan}[(c_{.})*(x_{.})]*(b_{.})\right)^{(p_{.})}*((f_{.})*(x_{.}))^{(m_{.})}\right)/\left((d_{.}) + (e_{.})*(x_{.})^2\right), x_Symbol]$ $\rightarrow \text{Simp}[f^2/e \text{ Int}[(f*x)^{(m - 2)}*(a + b*\text{ArcTan}[c*x])^p, x], x] - \text{Simp}[d*(f^2/e) \text{ Int}[(f*x)^{(m - 2)}*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x]$ /; $\text{FreeQ}\{a, b, c, d, e, f\}, x$ && $\text{GtQ}[p, 0]$ && $\text{GtQ}[m, 1]$
- rule 5566 $\text{Int}[\left((a_{.}) + \text{ArcTan}[(c_{.}) + (d_{.})*(x_{.})]*(b_{.})\right)^{(p_{.})}*((e_{.}) + (f_{.})*(x_{.}))^{(m_{.})}, x_Symbol]$ $\rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^{(m)}*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x]$ /; $\text{FreeQ}\{a, b, c, d, e, f, m\}, x$ && $\text{EqQ}[d*e - c*f, 0]$ && $\text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.97

method	result
derivativedivides	$\frac{e^3 a^2 (dx+c)^4 + e^3 b^2 \left(\frac{(dx+c)^4 \arctan(dx+c)^2}{4} - \frac{(dx+c)^3 \arctan(dx+c)}{6} + \frac{(dx+c) \arctan(dx+c)}{2} - \frac{\arctan(dx+c)^2}{4} + \frac{(dx+c)^2}{12} - \frac{\ln(1-i(dx+c))}{4} \right)}{d}$
default	$\frac{e^3 a^2 (dx+c)^4 + e^3 b^2 \left(\frac{(dx+c)^4 \arctan(dx+c)^2}{4} - \frac{(dx+c)^3 \arctan(dx+c)}{6} + \frac{(dx+c) \arctan(dx+c)}{2} - \frac{\arctan(dx+c)^2}{4} + \frac{(dx+c)^2}{12} - \frac{\ln(1-i(dx+c))}{4} \right)}{d}$
parts	$\frac{e^3 a^2 (dx+c)^4}{4d} + \frac{e^3 b^2 \left(\frac{(dx+c)^4 \arctan(dx+c)^2}{4} - \frac{(dx+c)^3 \arctan(dx+c)}{6} + \frac{(dx+c) \arctan(dx+c)}{2} - \frac{\arctan(dx+c)^2}{4} + \frac{(dx+c)^2}{12} - \frac{\ln(1-i(dx+c))}{4} \right)}{d}$
parallelrisch	$-\frac{18e^3 d c^2 a^2 + 5e^3 d c^2 b^2 - 24x^3 \arctan(dx+c) abc d^4 e^3 - 24x \arctan(dx+c) ab c^3 d^2 e^3 - 36x^2 \arctan(dx+c) ab c^2 d^3 e^3 + e^3}{3d}$
risch	$-\frac{e^3 b^2 \ln(d^2 x^2 + 2cdx + c^2 + 1)}{3d} + \frac{ab e^3 x}{2} + i e^3 ab c^3 x \ln(1 - i(dx + c)) + \frac{i e^3 d^3 ab x^4 \ln(1 - i(dx + c))}{4} - \dots$

input $\text{int}((d*e*x+c*e)^3*(a+b*\arctan(d*x+c))^2,x,\text{method}=_RETURNVERBOSE)$

output

```
1/d*(1/4*e^3*a^2*(d*x+c)^4+e^3*b^2*(1/4*(d*x+c)^4*arctan(d*x+c)^2-1/6*(d*x+c)^3*arctan(d*x+c)+1/2*(d*x+c)*arctan(d*x+c)-1/4*arctan(d*x+c)^2+1/12*(d*x+c)^2-1/3*ln(1+(d*x+c)^2))+2*e^3*a*b*(1/4*(d*x+c)^4*arctan(d*x+c)-1/12*(d*x+c)^3+1/4*d*x+1/4*c-1/4*arctan(d*x+c))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 337 vs. $2(143) = 286$.

Time = 0.13 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.15

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{3a^2d^4e^3x^4 + 2(6a^2c - ab)d^3e^3x^3 + (18a^2c^2 - 6abc + b^2)d^2e^3x^2 + 2(6a^2c^3 - 3abc^2 + b^2c + 3ab)de^3x - \dots}{d^5}$$

input

```
integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")
```

output

```
1/12*(3*a^2*d^4*e^3*x^4 + 2*(6*a^2*c - a*b)*d^3*e^3*x^3 + (18*a^2*c^2 - 6*a*b*c + b^2)*d^2*e^3*x^2 + 2*(6*a^2*c^3 - 3*a*b*c^2 + b^2*c + 3*a*b)*d*e^3*x - 4*b^2*e^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 3*(b^2*d^4*e^3*x^4 + 4*b^2*c*d^3*e^3*x^3 + 6*b^2*c^2*d^2*e^3*x^2 + 4*b^2*c^3*d*e^3*x + (b^2*c^4 - b^2)*e^3)*arctan(d*x + c)^2 + 2*(3*a*b*d^4*e^3*x^4 + (12*a*b*c - b^2)*d^3*e^3*x^3 + 3*(6*a*b*c^2 - b^2*c)*d^2*e^3*x^2 + 3*(4*a*b*c^3 - b^2*c^2 + b^2)*d*e^3*x + (3*a*b*c^4 - b^2*c^3 + 3*b^2*c - 3*a*b)*e^3)*arctan(d*x + c))/d
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.22 (sec) , antiderivative size = 583, normalized size of antiderivative = 3.71

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$= \begin{cases} a^2c^3e^3x + \frac{3a^2c^2de^3x^2}{2} + a^2cd^2e^3x^3 + \frac{a^2d^3e^3x^4}{4} + \frac{abc^4e^3 \operatorname{atan}(c+dx)}{2d} + 2abc^3e^3x \operatorname{atan}(c+dx) + 3abc^2de^3x^2 \operatorname{atan}(c+dx) \\ c^3e^3x(a + b \operatorname{atan}(c))^2 \end{cases}$$

input `integrate((d*e*x+c*e)**3*(a+b*atan(d*x+c))**2,x)`

output `Piecewise((a**2*c**3*e**3*x + 3*a**2*c**2*d*e**3*x**2/2 + a**2*c*d**2*e**3*x**3 + a**2*d**3*e**3*x**4/4 + a*b*c**4*e**3*atan(c + d*x)/(2*d) + 2*a*b*c**3*e**3*x*atan(c + d*x) + 3*a*b*c**2*d*e**3*x**2*atan(c + d*x) - a*b*c**2*e**3*x/2 + 2*a*b*c*d**2*e**3*x**3*atan(c + d*x) - a*b*c*d*e**3*x**2/2 + a*b*d**3*e**3*x**4*atan(c + d*x)/2 - a*b*d**2*e**3*x**3/6 + a*b*e**3*x/2 - a*b*e**3*atan(c + d*x)/(2*d) + b**2*c**4*e**3*atan(c + d*x)**2/(4*d) + b**2*c**3*e**3*x*atan(c + d*x)**2 - b**2*c**3*e**3*atan(c + d*x)/(6*d) + 3*b**2*c**2*d*e**3*x**2*atan(c + d*x)**2/2 - b**2*c**2*e**3*x*atan(c + d*x)/2 + b**2*c*d**2*e**3*x**3*atan(c + d*x)**2 - b**2*c*d*e**3*x**2*atan(c + d*x)/2 + b**2*c*e**3*x/6 + b**2*c*e**3*atan(c + d*x)/(2*d) + b**2*d**3*e**3*x**4*atan(c + d*x)**2/4 - b**2*d**2*e**3*x**3*atan(c + d*x)/6 + b**2*d*e**3*x**2/12 + b**2*e**3*x*atan(c + d*x)/2 - 2*b**2*e**3*log(c/d + x - I/d)/(3*d) - b**2*e**3*atan(c + d*x)**2/(4*d) + 2*I*b**2*e**3*atan(c + d*x)/(3*d), Ne(d, 0)), (c**3*e**3*x*(a + b*atan(c))**2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 597 vs. $2(143) = 286$.

Time = 1.07 (sec) , antiderivative size = 597, normalized size of antiderivative = 3.80

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx = \frac{1}{4} a^2 d^3 e^3 x^4 + a^2 c d^2 e^3 x^3 + \frac{3}{2} a^2 c^2 d e^3 x^2 + 3 \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2 x + cd}{d}\right) - c \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) abc^2 de^3 + \left(2x^3 \arctan(dx + c) - d \left(\frac{dx^2 - 4cx}{d^3} - \frac{2(c^3 - 3c) \arctan\left(\frac{d^2 x + cd}{d}\right) + (3c^2 - 1) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^4} \right) \right) abc^2 de^3 + \frac{1}{6} \left(3x^4 \arctan(dx + c) - d \left(\frac{d^2 x^3 - 3cdx^2 + 3(3c^2 - 1)x}{d^4} + \frac{3(c^4 - 6c^2 + 1) \arctan\left(\frac{d^2 x + cd}{d}\right) - 6(c^3 - 3c) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^5} \right) \right) abc^2 de^3 + a^2 c^3 e^3 x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) abc^3 e^3}{d} + \frac{b^2 d^2 e^3 x^2 + 2b^2 c d e^3 x - 4b^2 e^3 \log(d^2 x^2 + 2cdx + c^2 + 1) + 3(b^2 d^4 e^3 x^4 + 4b^2 c d^3 e^3 x^3 + 6b^2 c^2 d^2 e^3 x^2 + 6b^2 c d e^3 x - 4b^2 e^3 \log(d^2 x^2 + 2cdx + c^2 + 1))}{d^5}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/4*a^2*d^3*e^3*x^4 + a^2*c*d^2*e^3*x^3 + 3/2*a^2*c^2*d*e^3*x^2 + 3*(x^2*a \\ & \text{rctan}(d*x + c) - d*(x/d^2 + (c^2 - 1)*\text{arctan}((d^2*x + c*d)/d)/d^3 - c*\log(\\ & d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3))*a*b*c^2*d*e^3 + (2*x^3*\text{arctan}(d*x + c) \\ & - d*((d*x^2 - 4*c*x)/d^3 - 2*(c^3 - 3*c)*\text{arctan}((d^2*x + c*d)/d)/d^4 + (3* \\ & c^2 - 1)*\log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^4))*a*b*c*d^2*e^3 + 1/6*(3*x^4 \\ & *\text{arctan}(d*x + c) - d*((d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 - 1)*x)/d^4 + 3*(c^4 \\ & - 6*c^2 + 1)*\text{arctan}((d^2*x + c*d)/d)/d^5 - 6*(c^3 - c)*\log(d^2*x^2 + 2*c* \\ & d*x + c^2 + 1)/d^5))*a*b*d^3*e^3 + a^2*c^3*e^3*x + (2*(d*x + c)*\text{arctan}(d*x \\ & + c) - \log((d*x + c)^2 + 1))*a*b*c^3*e^3/d + 1/12*(b^2*d^2*e^3*x^2 + 2*b^ \\ & 2*c*d*e^3*x - 4*b^2*e^3*\log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 3*(b^2*d^4*e^3* \\ & x^4 + 4*b^2*c*d^3*e^3*x^3 + 6*b^2*c^2*d^2*e^3*x^2 + 4*b^2*c^3*d*e^3*x + (b \\ & ^2*c^4 - b^2)*e^3)*\text{arctan}(d*x + c)^2 - 2*(b^2*d^3*e^3*x^3 + 3*b^2*c*d^2*e^ \\ & 3*x^2 + 3*(b^2*c^2 - b^2)*d*e^3*x + (b^2*c^3 - 3*b^2*c)*e^3)*\text{arctan}(d*x + \\ & c))/d \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 610 vs. $2(143) = 286$.

Time = 0.44 (sec) , antiderivative size = 610, normalized size of antiderivative = 3.89

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{3b^2d^4e^3x^4 \arctan(dx + c)^2 + 6abd^4e^3x^4 \arctan(dx + c) + 12b^2cd^3e^3x^3 \arctan(dx + c)^2 + 3a^2d^4e^3x^4 + \dots}{\dots}$$

input `integrate((d*e*x+c*e)^3*(a+b*arctan(d*x+c))^2,x, algorithm="giac")`

output

```

1/12*(3*b^2*d^4*e^3*x^4*arctan(d*x + c)^2 + 6*a*b*d^4*e^3*x^4*arctan(d*x +
c) + 12*b^2*c*d^3*e^3*x^3*arctan(d*x + c)^2 + 3*a^2*d^4*e^3*x^4 + 24*a*b*
c*d^3*e^3*x^3*arctan(d*x + c) + 18*b^2*c^2*d^2*e^3*x^2*arctan(d*x + c)^2 +
12*a^2*c*d^3*e^3*x^3 + 36*a*b*c^2*d^2*e^3*x^2*arctan(d*x + c) - 2*b^2*d^3
*e^3*x^3*arctan(d*x + c) + 12*b^2*c^3*d*e^3*x*arctan(d*x + c)^2 + 18*a^2*c
^2*d^2*e^3*x^2 - 2*a*b*d^3*e^3*x^3 + 24*a*b*c^3*d*e^3*x*arctan(d*x + c) -
6*b^2*c*d^2*e^3*x^2*arctan(d*x + c) + 3*b^2*c^4*e^3*arctan(d*x + c)^2 + 3*
pi*a*b*c^4*e^3*sgn(d*x + c) - 3*pi*a*b*c^4*e^3 + 12*a^2*c^3*d*e^3*x - 6*a*
b*c*d^2*e^3*x^2 - 6*b^2*c^2*d*e^3*x*arctan(d*x + c) - 6*a*b*c^4*e^3*arctan
(1/(d*x + c)) - pi*b^2*c^3*e^3*sgn(d*x + c) + pi*b^2*c^3*e^3 - 6*a*b*c^2*d
*e^3*x + b^2*d^2*e^3*x^2 + 2*b^2*c^3*e^3*arctan(1/(d*x + c)) + 2*b^2*c*d*e
^3*x + 6*b^2*d*e^3*x*arctan(d*x + c) + 3*pi*b^2*c*e^3*sgn(d*x + c) - 3*pi*
b^2*c*e^3 + 6*a*b*d*e^3*x - 3*b^2*e^3*arctan(d*x + c)^2 - 6*b^2*c*e^3*arct
an(1/(d*x + c)) - 3*pi*a*b*e^3*sgn(d*x + c) + 3*pi*a*b*e^3 + 6*a*b*e^3*arc
tan(1/(d*x + c)) - 4*b^2*e^3*log((d*x + c)^2 + 1))/d

```

Mupad [B] (verification not implemented)

Time = 4.00 (sec) , antiderivative size = 633, normalized size of antiderivative = 4.03

$$\begin{aligned}
& \int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx \\
&= x \left(\frac{ce^3 (20a^2c^2 + 6a^2 - 6abc + b^2)}{2} + \frac{(6c^2 + 6) \left(2a^2cd^2e^3 + \frac{ad^2e^3(b-10ac)}{2} \right)}{6d^2} \right. \\
&\quad \left. - \frac{2c \left(\frac{2c \left(2a^2cd^2e^3 + \frac{ad^2e^3(b-10ac)}{2} \right)}{d} + \frac{de^3(60a^2c^2 + 6a^2 - 12abc + b^2)}{6} - \frac{a^2de^3(6c^2 + 6)}{6} \right)}{d} \right) \\
&\quad + x^2 \left(\frac{c \left(2a^2cd^2e^3 + \frac{ad^2e^3(b-10ac)}{2} \right)}{d} + \frac{de^3(60a^2c^2 + 6a^2 - 12abc + b^2)}{12} \right. \\
&\quad \quad \left. - \frac{a^2de^3(6c^2 + 6)}{12} \right) - x^3 \left(\frac{2a^2cd^2e^3}{3} + \frac{ad^2e^3(b-10ac)}{6} \right) \\
&\quad + \operatorname{atan}(c + dx)^2 \left(b^2c^3e^3x - \frac{b^2e^3 - b^2c^4e^3}{4d} + \frac{b^2d^3e^3x^4}{4} + \frac{3b^2c^2de^3x^2}{2} \right. \\
&\quad \quad \left. + b^2cd^2e^3x^3 \right) - d^2 \operatorname{atan}(c + dx) \left(x^3 \left(\frac{b^2e^3}{6} - 2abce^3 \right) \right. \\
&\quad \quad \left. - \frac{x(-b^2c^2e^3 + b^2e^3 + 4abc^3e^3)}{2d^2} + \frac{x^2(b^2ce^3 - 6abc^2e^3)}{2d} - \frac{abd^3e^3x^4}{2} \right) \\
&\quad + \frac{a^2d^3e^3x^4}{4} - \frac{b^2e^3 \ln(c^2 + 2cdx + d^2x^2 + 1)}{3d} \\
&\quad + \frac{be^3 \operatorname{atan} \left(\frac{\frac{bce^3(-3ac^4 + bc^3 - 3bc + 3a)}{6} + \frac{bde^3x(-3ac^4 + bc^3 - 3bc + 3a)}{6}}{-\frac{b^2c^3e^3}{6} + \frac{b^2ce^3}{2} + \frac{abc^4e^3}{2} - \frac{ab^2e^3}{2}} \right)}{6d} (-3ac^4 + bc^3 - 3bc + 3a)
\end{aligned}$$

input `int((c*e + d*e*x)^3*(a + b*atan(c + d*x))^2,x)`

output

```
x*((c*e^3*(6*a^2 + b^2 + 20*a^2*c^2 - 6*a*b*c))/2 + ((6*c^2 + 6)*(2*a^2*c*d^2*e^3 + (a*d^2*e^3*(b - 10*a*c))/2))/(6*d^2) - (2*c*((2*c*(2*a^2*c*d^2*e^3 + (a*d^2*e^3*(b - 10*a*c))/2))/d + (d*e^3*(6*a^2 + b^2 + 60*a^2*c^2 - 12*a*b*c))/6 - (a^2*d*e^3*(6*c^2 + 6))/6))/d + x^2*((c*(2*a^2*c*d^2*e^3 + (a*d^2*e^3*(b - 10*a*c))/2))/d + (d*e^3*(6*a^2 + b^2 + 60*a^2*c^2 - 12*a*b*c))/12 - (a^2*d*e^3*(6*c^2 + 6))/12) - x^3*((2*a^2*c*d^2*e^3)/3 + (a*d^2*e^3*(b - 10*a*c))/6) + atan(c + d*x)^2*(b^2*c^3*e^3*x - (b^2*e^3 - b^2*c^4*e^3)/(4*d) + (b^2*d^3*e^3*x^4)/4 + (3*b^2*c^2*d*e^3*x^2)/2 + b^2*c*d^2*e^3*x^3) - d^2*atan(c + d*x)*(x^3*((b^2*e^3)/6 - 2*a*b*c*e^3) - (x*(b^2*e^3 - b^2*c^2*e^3 + 4*a*b*c^3*e^3))/(2*d^2) + (x^2*(b^2*c*e^3 - 6*a*b*c^2*e^3))/(2*d) - (a*b*d*e^3*x^4)/2) + (a^2*d^3*e^3*x^4)/4 - (b^2*e^3*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(3*d) + (b*e^3*atan(((b*c*e^3*(3*a - 3*b*c - 3*a*c^4 + b*c^3))/6 + (b*d*e^3*x*(3*a - 3*b*c - 3*a*c^4 + b*c^3))/6))/((b^2*c*e^3)/2 - (b^2*c^3*e^3)/6 - (a*b*e^3)/2 + (a*b*c^4*e^3)/2))*(3*a - 3*b*c - 3*a*c^4 + b*c^3))/(6*d)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 421, normalized size of antiderivative = 2.68

$$\int (ce + dex)^3 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{e^3 (24 \operatorname{atan}(dx + c) ab c^3 dx + 36 \operatorname{atan}(dx + c) ab c^2 d^2 x^2 + 24 \operatorname{atan}(dx + c) abc d^3 x^3 + b^2 d^2 x^2 + 3 \operatorname{atan}(dx + c) ab c^3 dx + 36 \operatorname{atan}(dx + c) ab c^2 d^2 x^2 + 24 \operatorname{atan}(dx + c) abc d^3 x^3 + b^2 d^2 x^2 + 3 \operatorname{atan}(dx + c) ab c^3 dx)}{e^3 (24 \operatorname{atan}(dx + c) ab c^3 dx + 36 \operatorname{atan}(dx + c) ab c^2 d^2 x^2 + 24 \operatorname{atan}(dx + c) abc d^3 x^3 + b^2 d^2 x^2 + 3 \operatorname{atan}(dx + c) ab c^3 dx)}$$

input

```
int((d*e*x+c*e)^3*(a+b*atan(d*x+c))^2,x)
```

output

```
(e**3*(3*atan(c + d*x)**2*b**2*c**4 + 12*atan(c + d*x)**2*b**2*c**3*d*x +
18*atan(c + d*x)**2*b**2*c**2*d**2*x**2 + 12*atan(c + d*x)**2*b**2*c*d**3*
x**3 + 3*atan(c + d*x)**2*b**2*d**4*x**4 - 3*atan(c + d*x)**2*b**2 + 6*ata
n(c + d*x)*a*b*c**4 + 24*atan(c + d*x)*a*b*c**3*d*x + 36*atan(c + d*x)*a*b
*c**2*d**2*x**2 + 24*atan(c + d*x)*a*b*c*d**3*x**3 + 6*atan(c + d*x)*a*b*d
**4*x**4 - 6*atan(c + d*x)*a*b - 2*atan(c + d*x)*b**2*c**3 - 6*atan(c + d*
x)*b**2*c**2*d*x - 6*atan(c + d*x)*b**2*c*d**2*x**2 + 6*atan(c + d*x)*b**2
*c - 2*atan(c + d*x)*b**2*d**3*x**3 + 6*atan(c + d*x)*b**2*d*x - 4*log(c**
2 + 2*c*d*x + d**2*x**2 + 1)*b**2 + 12*a**2*c**3*d*x + 18*a**2*c**2*d**2*x
**2 + 12*a**2*c*d**3*x**3 + 3*a**2*d**4*x**4 - 6*a*b*c**2*d*x - 6*a*b*c*d*
**2*x**2 - 2*a*b*d**3*x**3 + 6*a*b*d*x + 2*b**2*c*d*x + b**2*d**2*x**2))/(1
2*d)
```

3.16 $\int (ce + dex)^2(a + b \arctan(c + dx))^2 dx$

Optimal result	162
Mathematica [A] (verified)	163
Rubi [A] (verified)	163
Maple [A] (verified)	167
Fricas [F]	167
Sympy [F]	168
Maxima [F]	168
Giac [F]	169
Mupad [F(-1)]	170
Reduce [F]	170

Optimal result

Integrand size = 23, antiderivative size = 183

$$\int (ce + dex)^2(a + b \arctan(c + dx))^2 dx = \frac{1}{3}b^2e^2x - \frac{b^2e^2 \arctan(c + dx)}{3d} - \frac{be^2(c + dx)^2(a + b \arctan(c + dx))}{3d} - \frac{ie^2(a + b \arctan(c + dx))^2}{3d} + \frac{e^2(c + dx)^3(a + b \arctan(c + dx))^2}{3d} - \frac{2be^2(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{3d} - \frac{ib^2e^2 \text{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{3d}$$

output

```
1/3*b^2*e^2*x-1/3*b^2*e^2*arctan(d*x+c)/d-1/3*b*e^2*(d*x+c)^2*(a+b*arctan(
d*x+c))/d-1/3*I*e^2*(a+b*arctan(d*x+c))^2/d+1/3*e^2*(d*x+c)^3*(a+b*arctan(
d*x+c))^2/d-2/3*b*e^2*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/d-1/3*I*b^2*
e^2*polylog(2,1-2/(1+I*(d*x+c)))/d
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.89

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{e^2 (a^2 (c + dx)^3 + ab(-(c + dx)^2 + 2(c + dx)^3 \arctan(c + dx) + \log(1 + (c + dx)^2)) + b^2 (c + dx - \arctan(c + dx)))}{3d}$$

input

```
Integrate[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x])^2,x]
```

output

```
(e^2*(a^2*(c + d*x)^3 + a*b*(-(c + d*x)^2 + 2*(c + d*x)^3*ArcTan[c + d*x]
+ Log[1 + (c + d*x)^2]) + b^2*(c + d*x - ArcTan[c + d*x] - (c + d*x)^2*Arc
Tan[c + d*x] + I*ArcTan[c + d*x]^2 + (c + d*x)^3*ArcTan[c + d*x]^2 - 2*Arc
Tan[c + d*x]*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + I*PolyLog[2, -E^((2*I)*A
rcTan[c + d*x])])))/(3*d)
```

Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.82, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$, Rules used = {5566, 27, 5361, 5451, 5361, 262, 216, 5455, 5379, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx$$

$$\downarrow \text{5566}$$

$$\frac{\int e^2 (c + dx)^2 (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow \text{27}$$

$$\frac{e^2 \int (c + dx)^2 (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow \text{5361}$$

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \int \frac{(c+dx)^3(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) \right)}{d}$$

↓ 5451

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(\int (c+dx)(a+b \arctan(c+dx)) d(c+dx) - \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) \right) \right)}{d}$$

↓ 5361

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(- \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}b \int \frac{(c+dx)^2}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 262

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(- \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}b \left(- \int \frac{1}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) \right) \right) \right)}{d}$$

↓ 216

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(- \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 5455

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(\int \frac{a+b \arctan(c+dx)}{-c-dx+i} d(c+dx) + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) + \frac{i}{2}(c+dx)(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 5379

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(-b \int \frac{\log\left(\frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 2849

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b \arctan(c+dx))^2 - \frac{2}{3}b \left(ib \int \frac{\log\left(\frac{2}{i(c+dx)+1}\right)}{1-\frac{2}{i(c+dx)+1}} d_{i(c+dx)+1} \frac{1}{i(c+dx)+1} + \frac{1}{2}(c+dx)^2(a+b \arctan(c+dx)) \right) \right)}{d}$$

↓ 2752

$$\frac{e^2 \left(\frac{1}{3}(c+dx)^3(a+b\arctan(c+dx))^2 - \frac{2}{3}b \left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx)) + \frac{i(a+b\arctan(c+dx))^2}{2b} + \log \left(\frac{2}{1+i(c+dx)} \right) \right) \right)}{d}$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x])^2,x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTan[c + d*x])^2)/3 - (2*b*(-1/2*(b*(c + d*x - ArcTan[c + d*x])) + ((c + d*x)^2*(a + b*ArcTan[c + d*x]))/2 + ((I/2)*(a + b*ArcTan[c + d*x])^2)/b + (a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] + (I/2)*b*PolyLog[2, 1 - 2/(1 + I*(c + d*x))]))/3))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5379

```
Int[((a_.) + ArcTan[(c_.)*(x_)*(b_.)]^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
:= Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(
p/e Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2))
, x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0
]
```

rule 5451

```
Int[(((a_.) + ArcTan[(c_.)*(x_)*(b_.)]^(p_.)*((f_.)*(x_)^(m_)))/((d_) + (e
_.)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTan[c*x]
)^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/(d
+ e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

rule 5455

```
Int[(((a_.) + ArcTan[(c_.)*(x_)*(b_.)]^(p_.)*(x_))/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Si
mp[1/(c*d Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)*(b_.)]^(p_.)*((e_.) + (f_.)*(x_)^(m
_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.51

method	result
derivativedivides	$\frac{e^2 a^2 (dx+c)^3}{3} + b^2 e^2 \left(\frac{(dx+c)^3 \arctan(dx+c)^2}{3} - \frac{(dx+c)^2 \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{dx}{3} + \frac{c}{3} - \frac{\arctan(dx+c)}{3} \right)$
default	$\frac{e^2 a^2 (dx+c)^3}{3} + b^2 e^2 \left(\frac{(dx+c)^3 \arctan(dx+c)^2}{3} - \frac{(dx+c)^2 \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{dx}{3} + \frac{c}{3} - \frac{\arctan(dx+c)}{3} \right)$
parts	$\frac{e^2 a^2 (dx+c)^3}{3d} + \frac{b^2 e^2 \left(\frac{(dx+c)^3 \arctan(dx+c)^2}{3} - \frac{(dx+c)^2 \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{dx}{3} + \frac{c}{3} - \frac{\arctan(dx+c)}{3} \right)}{d}$
risch	Expression too large to display

input `int((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/3*e^2*a^2*(d*x+c)^3+b^2*e^2*(1/3*(d*x+c)^3*arctan(d*x+c)^2-1/3*(d*x+c)^2*arctan(d*x+c)+1/3*arctan(d*x+c)*ln(1+(d*x+c)^2)+1/3*d*x+1/3*c-1/3*arctan(d*x+c)+1/6*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))-1/6*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I))))+2*e^2*a*b*(1/3*(d*x+c)^3*arctan(d*x+c)-1/6*(d*x+c)^2+1/6*ln(1+(d*x+c)^2))`

Fricas [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^2 dx$$

input `integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")`

output

```
integral(a^2*d^2*e^2*x^2 + 2*a^2*c*d*e^2*x + a^2*c^2*e^2 + (b^2*d^2*e^2*x^2 + 2*b^2*c*d*e^2*x + b^2*c^2*e^2)*arctan(d*x + c)^2 + 2*(a*b*d^2*e^2*x^2 + 2*a*b*c*d*e^2*x + a*b*c^2*e^2)*arctan(d*x + c), x)
```

Sympy [F]

$$\begin{aligned} & \int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx \\ &= e^2 \left(\int a^2 c^2 dx + \int a^2 d^2 x^2 dx + \int b^2 c^2 \operatorname{atan}^2(c + dx) dx + \int 2abc^2 \operatorname{atan}(c + dx) dx \right. \\ & \quad + \int 2a^2 c dx dx + \int b^2 d^2 x^2 \operatorname{atan}^2(c + dx) dx + \int 2abd^2 x^2 \operatorname{atan}(c + dx) dx \\ & \quad \left. + \int 2b^2 c dx \operatorname{atan}^2(c + dx) dx + \int 4abcdx \operatorname{atan}(c + dx) dx \right) \end{aligned}$$

input

```
integrate((d*e*x+c*e)**2*(a+b*atan(d*x+c))**2,x)
```

output

```
e**2*(Integral(a**2*c**2, x) + Integral(a**2*d**2*x**2, x) + Integral(b**2*c**2*atan(c + d*x)**2, x) + Integral(2*a*b*c**2*atan(c + d*x), x) + Integral(2*a**2*c*d*x, x) + Integral(b**2*d**2*x**2*atan(c + d*x)**2, x) + Integral(2*a*b*d**2*x**2*atan(c + d*x), x) + Integral(2*b**2*c*d*x*atan(c + d*x)**2, x) + Integral(4*a*b*c*d*x*atan(c + d*x), x))
```

Maxima [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^2 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")
```

output

```

3/4*b^2*c^4*e^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 1/4*(3*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*b^2*c^4*e^2 + 1/3*a^2*d^2*e^2*x^3 + 36*b^2*d^4*e^2*integrate(1/48*x^4*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^2*d^4*e^2*integrate(1/48*x^4*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 144*b^2*c*d^3*e^2*integrate(1/48*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 4*b^2*d^4*e^2*integrate(1/48*x^4*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*c*d^3*e^2*integrate(1/48*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 216*b^2*c^2*d^2*e^2*integrate(1/48*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 16*b^2*c*d^3*e^2*integrate(1/48*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 18*b^2*c^2*d^2*e^2*integrate(1/48*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 144*b^2*c^3*d*e^2*integrate(1/48*x*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 24*b^2*c^2*d^2*e^2*integrate(1/48*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*c^3*d*e^2*integrate(1/48*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*c^3*d*e^2*integrate(1/48*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^2*c^4*e^2*integrate(1/48*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1),...

```

Giac [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^2 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="giac")
```

output

```
sage0*x
```

Mupad [F(-1)]

Timed out.

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx = \int (ce + dex)^2 (a + b \operatorname{atan}(c + dx))^2 dx$$

input `int((c*e + d*e*x)^2*(a + b*atan(c + d*x))^2,x)`

output `int((c*e + d*e*x)^2*(a + b*atan(c + d*x))^2, x)`

Reduce [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^2 dx$$

$$= \frac{e^2 \left(\operatorname{atan}(dx + c)^2 b^2 c^3 + 3 \operatorname{atan}(dx + c)^2 b^2 c^2 dx + 3 \operatorname{atan}(dx + c)^2 b^2 c d^2 x^2 + \operatorname{atan}(dx + c)^2 b^2 c + \operatorname{atan}(dx + c) \right)}{3d}$$

input `int((d*e*x+c*e)^2*(a+b*atan(d*x+c))^2,x)`

output `(e**2*(atan(c + d*x)**2*b**2*c**3 + 3*atan(c + d*x)**2*b**2*c**2*d*x + 3*atan(c + d*x)**2*b**2*c*d**2*x**2 + atan(c + d*x)**2*b**2*c + atan(c + d*x)**2*b**2*d**3*x**3 + 2*atan(c + d*x)*a*b*c**3 + 6*atan(c + d*x)*a*b*c**2*d*x + 6*atan(c + d*x)*a*b*c*d**2*x**2 + 2*atan(c + d*x)*a*b*d**3*x**3 - atan(c + d*x)*b**2*c**2 - 2*atan(c + d*x)*b**2*c*d*x - atan(c + d*x)*b**2*d**2*x**2 - atan(c + d*x)*b**2 + 2*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*d**2 + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b + 3*a**2*c**2*d*x + 3*a**2*c*d**2*x**2 + a**2*d**3*x**3 - 2*a*b*c*d*x - a*b*d**2*x**2 + b**2*d*x))/(3*d)`

3.17 $\int (ce + dex)(a + b \arctan(c + dx))^2 dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [A] (verified)	174
Fricas [A] (verification not implemented)	174
Sympy [C] (verification not implemented)	175
Maxima [B] (verification not implemented)	176
Giac [B] (verification not implemented)	176
Mupad [B] (verification not implemented)	177
Reduce [B] (verification not implemented)	178

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx = -abex - \frac{b^2e(c + dx) \arctan(c + dx)}{d} + \frac{e(a + b \arctan(c + dx))^2}{2d} + \frac{e(c + dx)^2(a + b \arctan(c + dx))^2}{2d} + \frac{b^2e \log(1 + (c + dx)^2)}{2d}$$

output

```
-a*b*e*x-b^2*e*(d*x+c)*arctan(d*x+c)/d+1/2*e*(a+b*arctan(d*x+c))^2/d+1/2*e*(d*x+c)^2*(a+b*arctan(d*x+c))^2/d+1/2*b^2*e*ln(1+(d*x+c)^2)/d
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.13

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx = \frac{e(a(c + dx)(-2b + ac + adx) + 2b(-b(c + dx) + a(1 + c^2 + 2cdx + d^2x^2)) \arctan(c + dx) + b^2(1 + c^2 + d^2x^2))}{2d}$$

input `Integrate[(c*e + d*e*x)*(a + b*ArcTan[c + d*x])^2,x]`

output `(e*(a*(c + d*x)*(-2*b + a*c + a*d*x) + 2*b*(-(b*(c + d*x)) + a*(1 + c^2 + 2*c*d*x + d^2*x^2))*ArcTan[c + d*x] + b^2*(1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTan[c + d*x]^2 + b^2*Log[1 + (c + d*x)^2]))/(2*d)`

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.92, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5566, 27, 5361, 5451, 2009, 5419}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ce + dex)(a + b \arctan(c + dx))^2 dx \\
 & \quad \downarrow \text{5566} \\
 & \frac{\int e(c + dx)(a + b \arctan(c + dx))^2 d(c + dx)}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{e \int (c + dx)(a + b \arctan(c + dx))^2 d(c + dx)}{d} \\
 & \quad \downarrow \text{5361} \\
 & \frac{e \left(\frac{1}{2} (c + dx)^2 (a + b \arctan(c + dx))^2 - b \int \frac{(c + dx)^2 (a + b \arctan(c + dx))}{(c + dx)^2 + 1} d(c + dx) \right)}{d} \\
 & \quad \downarrow \text{5451} \\
 & \frac{e \left(\frac{1}{2} (c + dx)^2 (a + b \arctan(c + dx))^2 - b \left(\int (a + b \arctan(c + dx)) d(c + dx) - \int \frac{a + b \arctan(c + dx)}{(c + dx)^2 + 1} d(c + dx) \right) \right)}{d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{e \left(\frac{1}{2} (c + dx)^2 (a + b \arctan(c + dx))^2 - b \left(- \int \frac{a + b \arctan(c + dx)}{(c + dx)^2 + 1} d(c + dx) + a(c + dx) + b(c + dx) \arctan(c + dx) \right) \right)}{d}
 \end{aligned}$$

↓ 5419

$$\frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^2 - b\left(-\frac{(a+b\arctan(c+dx))^2}{2b} + a(c+dx) + b(c+dx)\arctan(c+dx) - \frac{1}{2}b\log\left(\frac{1+(c+dx)^2}{2}\right)\right)\right)}{d}$$

input `Int[(c*e + d*e*x)*(a + b*ArcTan[c + d*x])^2,x]`

output `(e*(((c + d*x)^2*(a + b*ArcTan[c + d*x])^2)/2 - b*(a*(c + d*x) + b*(c + d*x)*ArcTan[c + d*x] - (a + b*ArcTan[c + d*x])^2/(2*b) - (b*Log[1 + (c + d*x)^2])/2)))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m+1)*((a + b*ArcTan[c*x^n])^p/(m+1)), x] - Simp[b*c*n*(p/(m+1)) Int[x^(m+n)*((a + b*ArcTan[c*x^n])^(p-1)/(1+c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & IntegerQ[m])) && NeQ[m, -1]`

rule 5419 `Int[((a_) + ArcTan[(c_)*(x_)])*(b_)^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTan[c*x])^(p+1)/(b*c*d*(p+1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]`

rule 5451 `Int[((a_) + ArcTan[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m-2)*(a + b*ArcTan[c*x])^p, x], x] - Simp[d*(f^2/e) Int[(f*x)^(m-2)*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 5566

```
Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{e a^2 \frac{(dx+c)^2}{2} + e b^2 \left(\frac{(dx+c)^2 \arctan(dx+c)^2}{2} + \frac{\arctan(dx+c)^2}{2} - (dx+c) \arctan(dx+c) + \frac{\ln(1+(dx+c)^2)}{2} \right) + 2eab \left(\frac{(dx+c)^2 \arctan(dx+c)}{2} \right)}{d}$
default	$\frac{e a^2 \frac{(dx+c)^2}{2} + e b^2 \left(\frac{(dx+c)^2 \arctan(dx+c)^2}{2} + \frac{\arctan(dx+c)^2}{2} - (dx+c) \arctan(dx+c) + \frac{\ln(1+(dx+c)^2)}{2} \right) + 2eab \left(\frac{(dx+c)^2 \arctan(dx+c)}{2} \right)}{d}$
parts	$e a^2 \left(\frac{1}{2} d x^2 + c x \right) + \frac{e b^2 \left(\frac{(dx+c)^2 \arctan(dx+c)^2}{2} + \frac{\arctan(dx+c)^2}{2} - (dx+c) \arctan(dx+c) + \frac{\ln(1+(dx+c)^2)}{2} \right)}{d} + \dots$
parallelrisch	$\frac{d^3 e b^2 \arctan(dx+c)^2 x^2 + 2x^2 \arctan(dx+c) a b d^3 e + 2c e b^2 \arctan(dx+c)^2 x d^2 + x^2 a^2 d^3 e + 4x \arctan(dx+c) a b c d^2 e + \dots}{4d}$
risch	$-\frac{e b^2 (d^2 x^2 + 2cdx + c^2 + 1) \ln(1+i(dx+c))^2}{8d} + \frac{e b (-2ia d^2 x^2 + b d^2 x^2 \ln(1-i(dx+c)) - 4iacxd + 2bcdx \ln(1-i(dx+c)))}{4d} + \dots$

input

```
int((d*e*x+c*e)*(a+b*arctan(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(1/2*e*a^2*(d*x+c)^2+e*b^2*(1/2*(d*x+c)^2*arctan(d*x+c)^2+1/2*arctan(d*x+c)^2-(d*x+c)*arctan(d*x+c)+1/2*ln(1+(d*x+c)^2))+2*e*a*b*(1/2*(d*x+c)^2*arctan(d*x+c)-1/2*d*x-1/2*c+1/2*arctan(d*x+c))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.58

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx$$

$$= \frac{a^2 d^2 e x^2 + 2(a^2 c - ab) dex + b^2 e \log(d^2 x^2 + 2cdx + c^2 + 1) + (b^2 d^2 e x^2 + 2b^2 c dex + (b^2 c^2 + b^2) e) \arctan(dx+c)}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")`

output
$$\frac{1}{2}(a^2d^2e^x + 2(a^2c - ab)dex + b^2e \log(d^2x^2 + 2cdx + c^2 + 1) + (b^2d^2e^x + 2b^2cdex + (b^2c^2 + b^2)e) \arctan(dx + c)^2 + 2(abd^2e^x + (2abc - b^2)dex + (abc^2 - b^2c + ab)e) \arctan(dx + c))/d$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.34 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.53

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx$$

$$= \begin{cases} a^2cex + \frac{a^2dex^2}{2} + \frac{abc^2e \operatorname{atan}(c+dx)}{d} + 2abce x \operatorname{atan}(c + dx) + abdex^2 \operatorname{atan}(c + dx) - abex + \frac{abe \operatorname{atan}(c+dx)}{d} \\ cex(a + b \operatorname{atan}(c))^2 \end{cases}$$

input `integrate((d*e*x+c*e)*(a+b*atan(d*x+c))**2,x)`

output `Piecewise((a**2*c*e*x + a**2*d*e*x**2/2 + a*b*c**2*e*atan(c + d*x)/d + 2*a*b*c*e*x*atan(c + d*x) + a*b*d*e*x**2*atan(c + d*x) - a*b*e*x + a*b*e*atan(c + d*x)/d + b**2*c**2*e*atan(c + d*x)**2/(2*d) + b**2*c*e*x*atan(c + d*x)**2 - b**2*c*e*atan(c + d*x)/d + b**2*d*e*x**2*atan(c + d*x)**2/2 - b**2*e*x*atan(c + d*x) + b**2*e*log(c/d + x - I/d)/d + b**2*e*atan(c + d*x)**2/(2*d) - I*b**2*e*atan(c + d*x)/d, Ne(d, 0)), (c*e*x*(a + b*atan(c))**2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(89) = 178$.

Time = 0.91 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.29

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx = \frac{1}{2} a^2 dex^2 + \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2x + cd}{d}\right)}{d^3} - \frac{c \log(d^2x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) abde + a^2 cex + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) abce}{d} + \frac{b^2 e \log(d^2x^2 + 2cdx + c^2 + 1) + (b^2 d^2 ex^2 + 2b^2 c dex + (b^2 c^2 + b^2) e) \arctan(dx + c)^2 - 2(b^2 dex + b^2 c e) \arctan(dx + c)}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*a^2*d*e*x^2 + (x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3)*a*b*d*e + a^2*c*e*x + (2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*a*b*c*e/d + 1/2*(b^2*e*log(d^2*x^2 + 2*c*d*x + c^2 + 1) + (b^2*d^2*e*x^2 + 2*b^2*c*d*e*x + (b^2*c^2 + b^2)*e)*arctan(d*x + c)^2 - 2*(b^2*d*e*x + b^2*c*e)*arctan(d*x + c))/d`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(89) = 178$.

Time = 0.21 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.53

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx = \frac{b^2 d^2 ex^2 \arctan(dx + c)^2 + 2abd^2 ex^2 \arctan(dx + c) + 2b^2 c dex \arctan(dx + c)^2 + a^2 d^2 ex^2 + 4abcdex a + \dots}{2d}$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^2,x, algorithm="giac")`

output

```
1/2*(b^2*d^2*e*x^2*arctan(d*x + c)^2 + 2*a*b*d^2*e*x^2*arctan(d*x + c) + 2
*b^2*c*d*e*x*arctan(d*x + c)^2 + a^2*d^2*e*x^2 + 4*a*b*c*d*e*x*arctan(d*x
+ c) + b^2*c^2*e*arctan(d*x + c)^2 + 2*a^2*c*d*e*x + a*b*c^2*e*arctan(d*x
+ c) - 2*b^2*d*e*x*arctan(d*x + c) - a*b*c^2*e*arctan(-d*x - c) - 2*a*b*d*
e*x - b^2*c*e*arctan(d*x + c) + b^2*e*arctan(d*x + c)^2 + b^2*c*e*arctan(-
d*x - c) + a*b*e*arctan(d*x + c) - a*b*e*arctan(-d*x - c) + b^2*e*log((d*x
+ c)^2 + 1))/d
```

Mupad [B] (verification not implemented)

Time = 2.13 (sec) , antiderivative size = 216, normalized size of antiderivative = 2.27

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx$$

$$= \operatorname{atan}(c + dx)^2 \left(\frac{eb^2c^2 + eb^2}{2d} + b^2cex + \frac{b^2dex^2}{2} \right) - x(ae(b - 3ac) + 2a^2ce)$$

$$- d^2 \operatorname{atan}(c + dx) \left(\frac{x(b^2e - 2abce)}{d^2} - \frac{abex^2}{d} \right) + \frac{b^2e \ln(c^2 + 2cdx + d^2x^2 + 1)}{2d}$$

$$+ \frac{a^2dex^2}{2} + \frac{be \operatorname{atan}\left(\frac{bce(ac^2 - bc + a) + bde x(ac^2 - bc + a)}{-eb^2c + aebc^2 + aeb}\right) (ac^2 - bc + a)}{d}$$

input

```
int((c*e + d*e*x)*(a + b*atan(c + d*x))^2,x)
```

output

```
atan(c + d*x)^2*((b^2*e + b^2*c^2*e)/(2*d) + b^2*c*e*x + (b^2*d*e*x^2)/2)
- x*(a*e*(b - 3*a*c) + 2*a^2*c*e) - d^2*atan(c + d*x)*((x*(b^2*e - 2*a*b*c
*e))/d^2 - (a*b*e*x^2)/d) + (b^2*e*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(2*d)
+ (a^2*d*e*x^2)/2 + (b*e*atan((b*c*e*(a - b*c + a*c^2) + b*d*e*x*(a - b*c
+ a*c^2))/(a*b*e - b^2*c*e + a*b*c^2*e))*(a - b*c + a*c^2))/d
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.01

$$\int (ce + dex)(a + b \arctan(c + dx))^2 dx$$

$$= \frac{e(\operatorname{atan}(dx + c)^2 b^2 c^2 + 2\operatorname{atan}(dx + c)^2 b^2 c dx + \operatorname{atan}(dx + c)^2 b^2 d^2 x^2 + \operatorname{atan}(dx + c)^2 b^2 + 2\operatorname{atan}(dx + c)$$

input

```
int((d*e*x+c*e)*(a+b*atan(d*x+c))^2,x)
```

output

```
(e*(atan(c + d*x)**2*b**2*c**2 + 2*atan(c + d*x)**2*b**2*c*d*x + atan(c +
d*x)**2*b**2*d**2*x**2 + atan(c + d*x)**2*b**2 + 2*atan(c + d*x)*a*b*c**2
+ 4*atan(c + d*x)*a*b*c*d*x + 2*atan(c + d*x)*a*b*d**2*x**2 + 2*atan(c + d
*x)*a*b - 2*atan(c + d*x)*b**2*c - 2*atan(c + d*x)*b**2*d*x + log(c**2 + 2
*c*d*x + d**2*x**2 + 1)*b**2 + 2*a**2*c*d*x + a**2*d**2*x**2 - 2*a*b*d*x))
/(2*d)
```

3.18 $\int \frac{(a+b \arctan(c+dx))^2}{ce+dex} dx$

Optimal result	179
Mathematica [B] (verified)	180
Rubi [A] (verified)	180
Maple [C] (warning: unable to verify)	183
Fricas [F]	184
Sympy [F]	184
Maxima [F]	184
Giac [F]	185
Mupad [F(-1)]	185
Reduce [F]	185

Optimal result

Integrand size = 23, antiderivative size = 183

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \frac{2(a + b \arctan(c + dx))^2 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right)}{de} - \frac{ib(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{de} + \frac{ib(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+i(c+dx)}\right)}{de} - \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2de} + \frac{b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1+i(c+dx)}\right)}{2de}$$

output

```
-2*(a+b*arctan(d*x+c))^2*arctanh(-1+2/(1+I*(d*x+c)))/d/e-I*b*(a+b*arctan(d*x+c))*polylog(2,1-2/(1+I*(d*x+c)))/d/e+I*b*(a+b*arctan(d*x+c))*polylog(2,-1+2/(1+I*(d*x+c)))/d/e-1/2*b^2*polylog(3,1-2/(1+I*(d*x+c)))/d/e+1/2*b^2*polylog(3,-1+2/(1+I*(d*x+c)))/d/e
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 381 vs. $2(183) = 366$.

Time = 0.28 (sec) , antiderivative size = 381, normalized size of antiderivative = 2.08

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx$$

$$= \frac{-6iab\pi^2 - ib^2\pi^3 + 24iab\pi \arctan(c + dx) - 48iab \arctan(c + dx)^2 + 16ib^2 \arctan(c + dx)^3 - ab\pi \log(16$$

input `Integrate[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x),x]`

output

```
((-6*I)*a*b*Pi^2 - I*b^2*Pi^3 + (24*I)*a*b*Pi*ArcTan[c + d*x] - (48*I)*a*b*
*ArcTan[c + d*x]^2 + (16*I)*b^2*ArcTan[c + d*x]^3 - a*b*Pi*Log[16777216] +
24*b^2*ArcTan[c + d*x]^2*Log[1 - E^((-2*I)*ArcTan[c + d*x])] + 24*a*b*Pi*
Log[1 + E^((-2*I)*ArcTan[c + d*x])] - 48*a*b*ArcTan[c + d*x]*Log[1 + E^((-
2*I)*ArcTan[c + d*x])] + 48*a*b*ArcTan[c + d*x]*Log[1 - E^((2*I)*ArcTan[c
+ d*x])] - 24*b^2*ArcTan[c + d*x]^2*Log[1 + E^((2*I)*ArcTan[c + d*x])] + 2
4*a^2*Log[c + d*x] + 12*a*b*Pi*Log[1 + c^2 + 2*c*d*x + d^2*x^2] - (24*I)*a
*b*PolyLog[2, -E^((-2*I)*ArcTan[c + d*x])] + (24*I)*b^2*ArcTan[c + d*x]*Po
lyLog[2, E^((-2*I)*ArcTan[c + d*x])] + (24*I)*b^2*ArcTan[c + d*x]*PolyLog[
2, -E^((2*I)*ArcTan[c + d*x])] - (24*I)*a*b*PolyLog[2, E^((2*I)*ArcTan[c +
d*x])] + 12*b^2*PolyLog[3, E^((-2*I)*ArcTan[c + d*x])] - 12*b^2*PolyLog[3
, -E^((2*I)*ArcTan[c + d*x])])/(24*d*e)
```

Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {5566, 27, 5357, 5523, 5529, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx$$

$$\int \frac{(a+b \arctan(c+dx))^2}{e(c+dx)} d(c+dx)$$

5566

$$\int \frac{(a+b \arctan(c+dx))^2}{c+dx} d(c+dx)$$

27

5357

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a+b \arctan(c+dx))^2 - 4b \int \frac{(a+b \arctan(c+dx)) \operatorname{arctanh}\left(1 - \frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx)}{de}$$

5523

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a+b \arctan(c+dx))^2 - 4b \left(\frac{1}{2} \int \frac{(a+b \arctan(c+dx)) \log\left(2 - \frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx) - \frac{1}{2} \int \frac{(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c+dx) \right)}{de}$$

5529

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a+b \arctan(c+dx))^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a+b \arctan(c+dx)) \right) \right)}{de}$$

7164

$$\frac{2 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a+b \arctan(c+dx))^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a+b \arctan(c+dx)) \right) \right)}{de}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x),x]`

output `(2*(a + b*ArcTan[c + d*x])^2*ArcTanh[1 - 2/(1 + I*(c + d*x))] - 4*b*(((I/2)*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] + (b*PolyLog[3, 1 - 2/(1 + I*(c + d*x))])/4)/2 + ((-1/2*I)*(a + b*ArcTan[c + d*x])*PolyLog[2, -1 + 2/(1 + I*(c + d*x))] - (b*PolyLog[3, -1 + 2/(1 + I*(c + d*x))])/4)/2))/(d*e)`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 5357 $\text{Int}[((a_.) + \text{ArcTan}[(c_.)*(x_)]*(b_.))^p/(x_), x_Symbol] \rightarrow \text{Simp}[2*(a + b*\text{ArcTan}[c*x])^p*\text{ArcTanh}[1 - 2/(1 + I*c*x)], x] - \text{Simp}[2*b*c*p \text{ Int}[(a + b*\text{ArcTan}[c*x])^{p-1}*(\text{ArcTanh}[1 - 2/(1 + I*c*x)]/(1 + c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[p, 1]$
- rule 5523 $\text{Int}[(\text{ArcTanh}[u_]*((a_.) + \text{ArcTan}[(c_.)*(x_)]*(b_.))^p)/((d_) + (e_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Int}[\text{Log}[1 + u]*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x] - \text{Simp}[1/2 \text{ Int}[\text{Log}[1 - u]*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2*d] \ \&\& \ \text{EqQ}[u^2 - (1 - 2*(I/(I - c*x)))^2, 0]$
- rule 5529 $\text{Int}[(\text{Log}[u_]*((a_.) + \text{ArcTan}[(c_.)*(x_)]*(b_.))^p)/((d_) + (e_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(-I)*(a + b*\text{ArcTan}[c*x])^p*(\text{PolyLog}[2, 1 - u]/(2*c*d)), x] + \text{Simp}[b*p*(I/2 \text{ Int}[(a + b*\text{ArcTan}[c*x])^{p-1}*(\text{PolyLog}[2, 1 - u]/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2*d] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]$
- rule 5566 $\text{Int}[((a_.) + \text{ArcTan}[(c_) + (d_.)*(x_)]*(b_.))^p*((e_.) + (f_.)*(x_)^m), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$
- rule 7164 $\text{Int}[(u_)*\text{PolyLog}[n_, v_], x_Symbol] \rightarrow \text{With}[\{w = \text{DerivativeDivides}[v, u*v, x]\}, \text{Simp}[w*\text{PolyLog}[n + 1, v], x] /; \! \text{FalseQ}[w]] /; \text{FreeQ}[n, x]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.39 (sec) , antiderivative size = 1154, normalized size of antiderivative = 6.31

method	result	size
derivativeldivides	Expression too large to display	1154
default	Expression too large to display	1154
parts	Expression too large to display	1159

input `int((a+b*arctan(d*x+c))^2/(d*e*x+c*e),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & 1/d*(a^2/e*\ln(d*x+c)+b^2/e*(\ln(d*x+c)*\arctan(d*x+c)^2+I*\arctan(d*x+c)*\text{poly} \\
 & \log(2,-(1+I*(d*x+c))^2/(1+(d*x+c)^2))-1/2*\text{polylog}(3,-(1+I*(d*x+c))^2/(1+(d \\
 & *x+c)^2))-\arctan(d*x+c)^2*\ln((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)+\arctan(d*x+c \\
 &)^2*\ln(1+(1+I*(d*x+c))/(1+(d*x+c)^2)^{(1/2)})-2*I*\arctan(d*x+c)*\text{polylog}(2,-(\\
 & 1+I*(d*x+c))/(1+(d*x+c)^2)^{(1/2)})+2*\text{polylog}(3,-(1+I*(d*x+c))/(1+(d*x+c)^2) \\
 & ^{(1/2)})+\arctan(d*x+c)^2*\ln(1-(1+I*(d*x+c))/(1+(d*x+c)^2)^{(1/2)})-2*I*\arctan \\
 & (d*x+c)*\text{polylog}(2,(1+I*(d*x+c))/(1+(d*x+c)^2)^{(1/2)})+2*\text{polylog}(3,(1+I*(d*x \\
 & +c))/(1+(d*x+c)^2)^{(1/2)})+1/2*I*\text{Pi}*(\text{csgn}(I*((1+I*(d*x+c))^2/(1+(d*x+c) \\
 & ^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*\text{csgn}(((1+I*(d*x+c))^2/(1+(d*x+c) \\
 & ^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-\text{csgn}(((1+I*(d*x+c))^2/(1+(d*x+c) \\
 & ^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2+\text{csgn}(I*((1+I*(d*x+c))^2/(1+(d*x+c) \\
 & ^2)-1))*\text{csgn}(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*\text{csgn}(I*((1+I*(d*x+c))^2/ \\
 & (1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-\text{csgn}(I*((1+I*(d*x+c))^ \\
 & 2/(1+(d*x+c)^2)-1))*\text{csgn}(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+ \\
 & c))^2/(1+(d*x+c)^2)))^2-\text{csgn}(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*\text{csgn}(I*(\\
 & (1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2+\text{csgn} \\
 & (I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^3- \\
 & \text{csgn}(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)) \\
 &)*\text{csgn}(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)) \\
 &)^2+\text{csgn}(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)...
 \end{aligned}$$

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^2}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e),x, algorithm="fricas")`

output `integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)/(d*e*x + c*e), x)`

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \frac{\int \frac{a^2}{c+dx} dx + \int \frac{b^2 \operatorname{atan}^2(c+dx)}{c+dx} dx + \int \frac{2ab \operatorname{atan}(c+dx)}{c+dx} dx}{e}$$

input `integrate((a+b*atan(d*x+c))**2/(d*e*x+c*e),x)`

output `(Integral(a**2/(c + d*x), x) + Integral(b**2*atan(c + d*x)**2/(c + d*x), x) + Integral(2*a*b*atan(c + d*x)/(c + d*x), x))/e`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^2}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e),x, algorithm="maxima")`

output `a^2*log(d*e*x + c*e)/(d*e) + integrate(1/16*(12*b^2*arctan(d*x + c)^2 + b^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 32*a*b*arctan(d*x + c))/(d*e*x + c*e), x)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^2}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^2/(d*e*x + c*e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^2}{ce + dex} dx$$

input `int((a + b*atan(c + d*x))^2/(c*e + d*e*x),x)`

output `int((a + b*atan(c + d*x))^2/(c*e + d*e*x), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^2}{ce + dex} dx \\ &= \frac{2 \left(\int \frac{\operatorname{atan}(dx+c)}{dx+c} dx \right) abd + \left(\int \frac{\operatorname{atan}(dx+c)^2}{dx+c} dx \right) b^2 d + \log(dx + c) a^2}{de} \end{aligned}$$

input `int((a+b*atan(d*x+c))^2/(d*e*x+c*e),x)`

output `(2*int(atan(c + d*x)/(c + d*x),x)*a*b*d + int(atan(c + d*x)**2/(c + d*x),x)*b**2*d + log(c + d*x)*a**2)/(d*e)`

3.19 $\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^2} dx$

Optimal result	186
Mathematica [A] (verified)	187
Rubi [A] (verified)	187
Maple [B] (verified)	189
Fricas [F]	190
Sympy [F]	191
Maxima [F]	191
Giac [F]	192
Mupad [F(-1)]	192
Reduce [F]	192

Optimal result

Integrand size = 23, antiderivative size = 119

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = -\frac{i(a + b \arctan(c + dx))^2}{de^2} - \frac{(a + b \arctan(c + dx))^2}{de^2(c + dx)} + \frac{2b(a + b \arctan(c + dx)) \log\left(2 - \frac{2}{1 - i(c + dx)}\right)}{de^2} - \frac{ib^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - i(c + dx)}\right)}{de^2}$$

output

```
-I*(a+b*arctan(d*x+c))^2/d/e^2-(a+b*arctan(d*x+c))^2/d/e^2/(d*x+c)+2*b*(a+b*arctan(d*x+c))*ln(2-2/(1-I*(d*x+c)))/d/e^2-I*b^2*polylog(2,-1+2/(1-I*(d*x+c)))/d/e^2
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.13

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx$$

$$= \frac{-ib^2(-i + c + dx) \arctan(c + dx)^2 + 2b \arctan(c + dx) (-a + b(c + dx) \log(1 - e^{2i \arctan(c+dx)})) + a(-}{de^2(c + dx)}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^2,x]
```

output

```
((-I)*b^2*(-I + c + d*x)*ArcTan[c + d*x]^2 + 2*b*ArcTan[c + d*x]*(-a + b*(c + d*x)*Log[1 - E^((2*I)*ArcTan[c + d*x])]) + a*(-a + 2*b*(c + d*x)*Log[(c + d*x)/Sqrt[1 + (c + d*x)^2]]) - I*b^2*(c + d*x)*PolyLog[2, E^((2*I)*ArcTan[c + d*x])])/(d*e^2*(c + d*x))
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {5566, 27, 5361, 5459, 5403, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx$$

$$\downarrow \text{5566}$$

$$\int \frac{(a + b \arctan(c + dx))^2}{e^2(c + dx)^2} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^2} d(c + dx)$$

$$\downarrow \text{5361}$$

$$\begin{aligned}
& \frac{2b \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^2}{c+dx}}{de^2} \\
& \quad \downarrow \text{5459} \\
& \frac{-\frac{(a+b \arctan(c+dx))^2}{c+dx} + 2b \left(i \int \frac{a+b \arctan(c+dx)}{(c+dx)(c+dx+i)} d(c+dx) - \frac{i(a+b \arctan(c+dx))^2}{2b} \right)}{de^2} \\
& \quad \downarrow \text{5403} \\
& \frac{-\frac{(a+b \arctan(c+dx))^2}{c+dx} + 2b \left(i \left(i b \int \frac{\log\left(2 - \frac{2}{1-i(c+dx)}\right)}{(c+dx)^2+1} d(c+dx) - i \log\left(2 - \frac{2}{1-i(c+dx)}\right) (a+b \arctan(c+dx)) \right) \right)}{de^2} - \frac{i(a+b \arctan(c+dx))^2}{c+dx} \\
& \quad \downarrow \text{2897} \\
& \frac{-\frac{(a+b \arctan(c+dx))^2}{c+dx} + 2b \left(i \left(-i \log\left(2 - \frac{2}{1-i(c+dx)}\right) (a+b \arctan(c+dx)) - \frac{1}{2} b \text{PolyLog}\left(2, \frac{2}{1-i(c+dx)} - 1\right) \right) \right)}{de^2} - \frac{i(a+b \arctan(c+dx))^2}{c+dx}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^2,x]`

output `((-(a + b*ArcTan[c + d*x])^2/(c + d*x)) + 2*b*(((1/2*I)*(a + b*ArcTan[c + d*x])^2)/b + I*((-I)*(a + b*ArcTan[c + d*x])*Log[2 - 2/(1 - I*(c + d*x))] - (b*PolyLog[2, -1 + 2/(1 - I*(c + d*x))])/2)))/(d*e^2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2897 `Int[Log[u_]*(P_q_)^(m_.), x_Symbol] := With[{C = FullSimplify[P_q^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[P_q, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[P_q, x]]`

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5403

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_
Symbol] := Simp[(a + b*ArcTan[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Si
mp[b*c*(p/d) Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1
+ c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*
d^2 + e^2, 0]
```

rule 5459

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*d*(p + 1))), x] + Si
mp[I/d Int[(a + b*ArcTan[c*x])^p/(x*(I + c*x)), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && GtQ[p, 0]
```

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m
_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 324 vs. $2(115) = 230$.

Time = 0.92 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.73

method	result
derivativedivides	$-\frac{a^2}{e^2(dx+c)} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{dx+c} + 2 \ln(dx+c) \arctan(dx+c) - \arctan(dx+c) \ln(1+(dx+c)^2) - \frac{i \left(\ln(dx+c-i) \ln(1+(dx+c)^2) - \ln(1+(dx+c-i)^2) \right)}{2i} \right)}{e^2(dx+c)}$
default	$-\frac{a^2}{e^2(dx+c)} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{dx+c} + 2 \ln(dx+c) \arctan(dx+c) - \arctan(dx+c) \ln(1+(dx+c)^2) - \frac{i \left(\ln(dx+c-i) \ln(1+(dx+c)^2) - \ln(1+(dx+c-i)^2) \right)}{2i} \right)}{e^2(dx+c)}$
parts	$-\frac{a^2}{e^2(dx+c)d} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{dx+c} + 2 \ln(dx+c) \arctan(dx+c) - \arctan(dx+c) \ln(1+(dx+c)^2) - \frac{i \left(\ln(dx+c-i) \ln(1+(dx+c)^2) - \ln(1+(dx+c-i)^2) \right)}{2i} \right)}{e^2(dx+c)d}$

input `int((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^2,x,method=_RETURNVERBOSE)`

output `1/d*(-a^2/e^2/(d*x+c)+b^2/e^2*(-1/(d*x+c)*arctan(d*x+c)^2+2*ln(d*x+c)*arctan(d*x+c)-arctan(d*x+c)*ln(1+(d*x+c)^2)-1/2*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))+1/2*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))+I*ln(d*x+c)*ln(1+I*(d*x+c))-I*ln(d*x+c)*ln(1-I*(d*x+c))+I*dilog(1+I*(d*x+c))-I*dilog(1-I*(d*x+c)))+2*a*b/e^2*(-1/(d*x+c)*arctan(d*x+c)+ln(d*x+c)-1/2*ln(1+(d*x+c)^2))`

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="fricas")`

output `integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)/(d^2*e^2*x^2 + 2*c*d*e^2*x + c^2*e^2), x)`

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = \frac{\int \frac{a^2}{c^2 + 2cdx + d^2x^2} dx + \int \frac{b^2 \operatorname{atan}^2(c + dx)}{c^2 + 2cdx + d^2x^2} dx + \int \frac{2ab \operatorname{atan}(c + dx)}{c^2 + 2cdx + d^2x^2} dx}{e^2}$$

input `integrate((a+b*atan(d*x+c))**2/(d*e*x+c*e)**2,x)`

output `(Integral(a**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(b**2*atan(c + d*x)**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(2*a*b*atan(c + d*x)/(c**2 + 2*c*d*x + d**2*x**2), x))/e**2`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="maxima")`

output `-(d*(log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2) - 2*log(d*x + c)/(d^2*e^2)) + 2*arctan(d*x + c)/(d^2*e^2*x + c*d*e^2))*a*b - 1/16*(4*arctan(d*x + c)^2 - 16*(d^2*e^2*x + c*d*e^2)*integrate(1/16*(12*(d^2*x^2 + 2*c*d*x + c^2 + 1)*arctan(d*x + c)^2 + (d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 8*(d*x + c)*arctan(d*x + c) - 4*(d^2*x^2 + 2*c*d*x + c^2)*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/(d^4*e^2*x^4 + 4*c*d^3*e^2*x^3 + (6*c^2 + 1)*d^2*e^2*x^2 + 2*(2*c^3 + c)*d*e^2*x + (c^4 + c^2)*e^2), x) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2*b^2/(d^2*e^2*x + c*d*e^2) - a^2/(d^2*e^2*x + c*d*e^2)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^2,x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^2/(d*e*x + c*e)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^2}{(ce + dex)^2} dx$$

input `int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^2,x)`

output `int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^2, x)`

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^2} dx$$

$$= \frac{-\operatorname{atan}(dx + c)^2 b^2 c^3 - \operatorname{atan}(dx + c)^2 b^2 c^2 dx - \operatorname{atan}(dx + c)^2 b^2 c + \operatorname{atan}(dx + c)^2 b^2 dx - 4 \operatorname{atan}(dx + c)}$$

input `int((a+b*atan(d*x+c))^2/(d*e*x+c*e)^2,x)`

output

```
( - atan(c + d*x)**2*b**2*c**3 - atan(c + d*x)**2*b**2*c**2*d*x - atan(c +
d*x)**2*b**2*c + atan(c + d*x)**2*b**2*d*x - 4*atan(c + d*x)*a*b*c - 2*at
an(c + d*x)*b**2*c**2 - 2*int((atan(c + d*x)*x**2)/(c**4 + 4*c**3*d*x + 6*
c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x
)*b**2*c*d**3 - 2*int((atan(c + d*x)*x**2)/(c**4 + 4*c**3*d*x + 6*c**2*d**
2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*b**2*d
**4*x - 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*c**2 - 2*log(c**2 + 2*c*
d*x + d**2*x**2 + 1)*a*b*c*d*x - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*
c**3 - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c**2*d*x + 4*log(c + d*x)*
a*b*c**2 + 4*log(c + d*x)*a*b*c*d*x + 2*log(c + d*x)*b**2*c**3 + 2*log(c +
d*x)*b**2*c**2*d*x + 2*a**2*d*x)/(2*c*d*e**2*(c + d*x))
```

3.20 $\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^3} dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [A] (verified)	198
Fricas [A] (verification not implemented)	199
Sympy [C] (verification not implemented)	199
Maxima [B] (verification not implemented)	200
Giac [F]	201
Mupad [B] (verification not implemented)	201
Reduce [B] (verification not implemented)	202

Optimal result

Integrand size = 23, antiderivative size = 117

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx = -\frac{b(a + b \arctan(c + dx))}{de^3(c + dx)} - \frac{(a + b \arctan(c + dx))^2}{2de^3} - \frac{(a + b \arctan(c + dx))^2}{2de^3(c + dx)^2} + \frac{b^2 \log(c + dx)}{de^3} - \frac{b^2 \log(1 + (c + dx)^2)}{2de^3}$$

output

```
-b*(a+b*arctan(d*x+c))/d/e^3/(d*x+c)-1/2*(a+b*arctan(d*x+c))^2/d/e^3-1/2*(a+b*arctan(d*x+c))^2/d/e^3/(d*x+c)^2+b^2*ln(d*x+c)/d/e^3-1/2*b^2*ln(1+(d*x+c)^2)/d/e^3
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.66

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx = \frac{a^2 + 2abc + 2abdx + 2b(b(c + dx) + a(1 + c^2 + 2cdx + d^2x^2)) \arctan(c + dx) + b^2(1 + c^2 + 2cdx + d^2x^2)}{(ce + dex)^3}$$

input `Integrate[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^3,x]`

output `-1/2*(a^2 + 2*a*b*c + 2*a*b*d*x + 2*b*(b*(c + d*x) + a*(1 + c^2 + 2*c*d*x + d^2*x^2))*ArcTan[c + d*x] + b^2*(1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTan[c + d*x]^2 - 2*b^2*(c + d*x)^2*Log[c + d*x] + b^2*c^2*Log[1 + c^2 + 2*c*d*x + d^2*x^2] + 2*b^2*c*d*x*Log[1 + c^2 + 2*c*d*x + d^2*x^2] + b^2*d^2*x^2*Log[1 + c^2 + 2*c*d*x + d^2*x^2])/(d*e^3*(c + d*x)^2)`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {5566, 27, 5361, 5453, 5361, 243, 47, 14, 16, 5419}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx \\
 & \quad \downarrow \text{5566} \\
 & \int \frac{(a + b \arctan(c + dx))^2}{e^3(c + dx)^3} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^3} d(c + dx) \\
 & \quad \downarrow \text{5361} \\
 & \frac{b \int \frac{a + b \arctan(c + dx)}{(c + dx)^2((c + dx)^2 + 1)} d(c + dx) - \frac{(a + b \arctan(c + dx))^2}{2(c + dx)^2}}{de^3} \\
 & \quad \downarrow \text{5453} \\
 & \frac{b \left(\int \frac{a + b \arctan(c + dx)}{(c + dx)^2} d(c + dx) - \int \frac{a + b \arctan(c + dx)}{(c + dx)^2 + 1} d(c + dx) \right) - \frac{(a + b \arctan(c + dx))^2}{2(c + dx)^2}}{de^3} \\
 & \quad \downarrow \text{5361}
 \end{aligned}$$

$$\frac{b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + b \int \frac{1}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 243

$$\frac{b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c+dx)^2 - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 47

$$\frac{b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b\left(\int \frac{1}{(c+dx)^2} d(c+dx)^2 - \int \frac{1}{(c+dx)^2+1} d(c+dx)^2\right) - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 14

$$\frac{b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b\left(\log((c+dx)^2) - \int \frac{1}{(c+dx)^2+1} d(c+dx)^2\right) - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 16

$$\frac{b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{a+b \arctan(c+dx)}{c+dx} + \frac{1}{2}b(\log((c+dx)^2) - \log((c+dx)^2+1))\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

↓ 5419

$$\frac{b\left(-\frac{(a+b \arctan(c+dx))^2}{2b} - \frac{a+b \arctan(c+dx)}{c+dx} + \frac{1}{2}b(\log((c+dx)^2) - \log((c+dx)^2+1))\right) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}}{de^3}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^3,x]`

output `(-1/2*(a + b*ArcTan[c + d*x])^2/(c + d*x)^2 + b*(-((a + b*ArcTan[c + d*x])/(c + d*x)) - (a + b*ArcTan[c + d*x])^2/(2*b) + (b*(Log[(c + d*x)^2] - Log[1 + (c + d*x)^2]))/2))/(d*e^3)`

Defintions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ /; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ /; FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ /; FreeQ}[b, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ /; FreeQ}[\{a, b, c, d\}, x]$
- rule 243 $\text{Int}[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] \text{ /; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m - 1)/2]$
- rule 5361 $\text{Int}[(a_ + \text{ArcTan}[c_*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[x^(m + 1)*((a + b*\text{ArcTan}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^(m + n)*((a + b*\text{ArcTan}[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] \text{ /; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 5419 $\text{Int}[(a_ + \text{ArcTan}[c_*(x_)]*(b_))^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^(p + 1)/(b*c*d*(p + 1)), x] \text{ /; FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[e, c^2*d] \ \&\& \ \text{NeQ}[p, -1]$
- rule 5453 $\text{Int}[(a_ + \text{ArcTan}[c_*(x_)]*(b_))^(p_)*((f_)*(x_)^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f*x)^m*(a + b*\text{ArcTan}[c*x])^p, x], x] - \text{Simp}[e/(d*f^2) \text{ Int}[(f*x)^(m + 2)*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.10

method	result
derivativdivides	$\frac{-\frac{a^2}{2e^3(dx+c)^2} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{2(dx+c)^2} - \frac{\arctan(dx+c)}{dx+c} - \frac{\arctan(dx+c)^2}{2} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{e^3} + \frac{2ab \left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{2c}{e^3} \right)}{d}$
default	$\frac{-\frac{a^2}{2e^3(dx+c)^2} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{2(dx+c)^2} - \frac{\arctan(dx+c)}{dx+c} - \frac{\arctan(dx+c)^2}{2} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{e^3} + \frac{2ab \left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{2c}{e^3} \right)}{d}$
parts	$-\frac{a^2}{2e^3(dx+c)^2 d} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{2(dx+c)^2} - \frac{\arctan(dx+c)}{dx+c} - \frac{\arctan(dx+c)^2}{2} + \ln(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{e^3 d} + \frac{2ab \left(-\frac{\arctan(dx+c)}{2(dx+c)^2} - \frac{2c}{e^3} \right)}{d}$
parallelrisch	$\frac{ab d^4 x^2 - 3ab c^2 d^2 + 4 \ln(dx+c) b^2 c^3 d^2 - 2 \ln(d^2 x^2 + 2cdx + c^2 + 1) b^2 c^3 d^2 - 4 \arctan(dx+c) b^2 c^2 d^2 - 2 \arctan(dx+c)^2 b^2 c^3 d^2}{e^3 d}$
risch	$\frac{b^2 (d^2 x^2 + 2cdx + c^2 + 1) \ln(1+i(dx+c))^2}{8e^3(dx+c)^2 d} - \frac{b(b d^2 x^2 \ln(1-i(dx+c)) + 2bcdx \ln(1-i(dx+c)) - 2ibdx + \ln(1-i(dx+c))) b c^3}{4e^3(dx+c)^2 d}$

input

```
int((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-1/2*a^2/e^3/(d*x+c)^2+b^2/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)^2-1/(d*x+c)*arctan(d*x+c)-1/2*arctan(d*x+c)^2+ln(d*x+c)-1/2*ln(1+(d*x+c)^2))+2*a*b/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)-1/2/(d*x+c)-1/2*arctan(d*x+c)))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.79

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx =$$

$$\frac{2 abdx + 2 abc + (b^2 d^2 x^2 + 2 b^2 c dx + b^2 c^2 + b^2) \arctan(dx + c)^2 + a^2 + 2 (abd^2 x^2 + abc^2 + b^2 c + (2 ab$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="fricas")`

output `-1/2*(2*a*b*d*x + 2*a*b*c + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2 + b^2)*arctan(d*x + c)^2 + a^2 + 2*(a*b*d^2*x^2 + a*b*c^2 + b^2*c + (2*a*b*c + b^2)*d*x + a*b)*arctan(d*x + c) + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 2*(b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*log(d*x + c))/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 9.22 (sec) , antiderivative size = 1107, normalized size of antiderivative = 9.46

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx = \text{Too large to display}$$

input `integrate((a+b*atan(d*x+c))**2/(d*e*x+c*e)**3,x)`

output

```
Piecewise((-a**2/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - 2*
a*b*c**2*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2
) - 4*a*b*c*d*x*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**
*3*x**2) - 2*a*b*c/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) -
2*a*b*d**2*x**2*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**
*3*x**2) - 2*a*b*d*x/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2)
- 2*a*b*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2)
+ 2*b**2*c**2*log(c/d + x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3
*x**2) - 2*b**2*c**2*log(c/d + x - I/d)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x +
2*d**3*e**3*x**2) - b**2*c**2*atan(c + d*x)**2/(2*c**2*d*e**3 + 4*c*d**2*
e**3*x + 2*d**3*e**3*x**2) + 2*I*b**2*c**2*atan(c + d*x)/(2*c**2*d*e**3 +
4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 4*b**2*c*d*x*log(c/d + x)/(2*c**2*d*
e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - 4*b**2*c*d*x*log(c/d + x - I/
d)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - 2*b**2*c*d*x*ata
n(c + d*x)**2/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**2) + 4*I*b
**2*c*d*x*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3*x**
2) - 2*b**2*c*atan(c + d*x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x + 2*d**3*e**3
*x**2) + 2*b**2*d**2*x**2*log(c/d + x)/(2*c**2*d*e**3 + 4*c*d**2*e**3*x +
2*d**3*e**3*x**2) - 2*b**2*d**2*x**2*log(c/d + x - I/d)/(2*c**2*d*e**3 +
4*c*d**2*e**3*x + 2*d**3*e**3*x**2) - b**2*d**2*x**2*atan(c + d*x)**2/(2...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. $2(111) = 222$.

Time = 0.21 (sec) , antiderivative size = 268, normalized size of antiderivative = 2.29

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx$$

$$= - \left(d \left(\frac{1}{d^3 e^3 x + cd^2 e^3} + \frac{\arctan\left(\frac{d^2 x + cd}{d}\right)}{d^2 e^3} \right) + \frac{\arctan(dx + c)}{d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3} \right) ab$$

$$- \frac{1}{2} \left(2d \left(\frac{1}{d^3 e^3 x + cd^2 e^3} + \frac{\arctan\left(\frac{d^2 x + cd}{d}\right)}{d^2 e^3} \right) \arctan(dx + c) - \frac{\arctan(dx + c)^2 - \log(d^2 x^2 + 2cdx + c^2)}{de^3} \right)$$

$$- \frac{b^2 \arctan(dx + c)^2}{2(d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3)} - \frac{a^2}{2(d^3 e^3 x^2 + 2cd^2 e^3 x + c^2 de^3)}$$

input

```
integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="maxima")
```

output

$$-(d*(1/(d^3*e^3*x + c*d^2*e^3) + \arctan((d^2*x + c*d)/d)/(d^2*e^3)) + \arctan(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3))*a*b - 1/2*(2*d*(1/(d^3*e^3*x + c*d^2*e^3) + \arctan((d^2*x + c*d)/d)/(d^2*e^3))*\arctan(d*x + c) - (\arctan(d*x + c)^2 - \log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 2*\log(d*x + c))/(d*e^3))*b^2 - 1/2*b^2*\arctan(d*x + c)^2/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3) - 1/2*a^2/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)$$
Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^3} dx$$

input

```
integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^3,x, algorithm="giac")
```

output

```
integrate((b*arctan(d*x + c) + a)^2/(d*e*x + c*e)^3, x)
```

Mupad [B] (verification not implemented)

Time = 3.49 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.98

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx = \frac{b^2 \ln(c + dx)}{de^3} - \frac{\frac{a^2 + 2bca}{2d} + abx}{c^2 e^3 + 2cde^3 x + d^2 e^3 x^2} - \frac{\operatorname{atan}(c + dx) \left(\frac{b^2 c}{d^3 e^3} + \frac{b^2 x}{d^2 e^3} + \frac{ab}{d^3 e^3} \right)}{x^2 + \frac{c^2}{d^2} + \frac{2cx}{d}} - \operatorname{atan}(c + dx)^2 \left(\frac{b^2}{2de^3} + \frac{b^2}{2d^3 e^3 \left(x^2 + \frac{c^2}{d^2} + \frac{2cx}{d} \right)} \right) + \frac{\ln(c + dx - i) \left(-\frac{b^2}{2} + \frac{ab \operatorname{li}}{2} \right)}{de^3} - \frac{\ln(c + dx + \operatorname{li}) \left(\frac{b^2}{2} + \frac{\operatorname{li} ab}{2} \right)}{de^3}$$

input

```
int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^3,x)
```

output

$$\begin{aligned} & (b^2 \log(c + dx)) / (d^3 e^3) - ((a^2 + 2ab^2 c) / (2d) + ab^2 x) / (c^2 e^3 + d^2 e^3 x^2 + 2c d e^3 x) - (\operatorname{atan}(c + dx) * ((b^2 c) / (d^3 e^3) + (b^2 x) / (d^2 e^3) + (ab) / (d^3 e^3))) / (x^2 + c^2/d^2 + (2cx)/d) - \operatorname{atan}(c + dx)^2 * \\ & (b^2 / (2d e^3) + b^2 / (2d^3 e^3 * (x^2 + c^2/d^2 + (2cx)/d))) + (\log(c + dx - 1i) * ((ab^2 1i) / 2 - b^2/2)) / (d^3 e^3) - (\log(c + dx + 1i) * ((ab^2 1i) / 2 + b^2/2)) / (d^3 e^3) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.87

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^3} dx$$

$$= \frac{-\operatorname{atan}(dx + c)^2 b^2 c^3 - 2 \operatorname{atan}(dx + c)^2 b^2 c^2 dx - \operatorname{atan}(dx + c)^2 b^2 c d^2 x^2 - \operatorname{atan}(dx + c)^2 b^2 c - 2 \operatorname{atan}(dx + c) a b^2 c^3 - 4 \operatorname{atan}(dx + c) a b^2 c^2 dx - 2 \operatorname{atan}(dx + c) a b^2 c d^2 x^2 - 2 \operatorname{atan}(dx + c) a b^2 c - 2 \operatorname{atan}(dx + c) a b^2 c^2 - 2 \operatorname{atan}(dx + c) b^2 c^2 dx - \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^3 - 2 \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^2 dx - \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^2 dx^2 + 2 \log(c + dx) b^2 c^3 + 4 \log(c + dx) b^2 c^2 dx + 2 \log(c + dx) b^2 c d^2 x^2 - a^2 c - a b^2 c^2 + a b^2 d^2 x^2}{(2c d^3 e^3 (c^2 + 2c dx + d^2 x^2))}$$

input `int((a+b*atan(d*x+c))^2/(d*e*x+c*e)^3,x)`

output

$$\begin{aligned} & (-\operatorname{atan}(c + dx)^2 b^2 c^3 - 2 \operatorname{atan}(c + dx)^2 b^2 c^2 dx - \operatorname{atan}(c + dx)^2 b^2 c^2 d^2 x^2 - \operatorname{atan}(c + dx)^2 b^2 c^2 - 2 \operatorname{atan}(c + dx) a b^2 c^3 - 4 \operatorname{atan}(c + dx) a b^2 c^2 dx - 2 \operatorname{atan}(c + dx) a b^2 c d^2 x^2 - 2 \operatorname{atan}(c + dx) a b^2 c - 2 \operatorname{atan}(c + dx) b^2 c^2 dx - \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^3 - 2 \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^2 dx - \log(c^2 + 2c dx + d^2 x^2 + 1) b^2 c^2 dx^2 + 2 \log(c + dx) b^2 c^3 + 4 \log(c + dx) b^2 c^2 dx + 2 \log(c + dx) b^2 c d^2 x^2 - a^2 c - a b^2 c^2 + a b^2 d^2 x^2) / (2c d^3 e^3 (c^2 + 2c dx + d^2 x^2)) \end{aligned}$$

3.21 $\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^4} dx$

Optimal result	203
Mathematica [A] (verified)	204
Rubi [A] (verified)	204
Maple [B] (verified)	207
Fricas [F]	208
Sympy [F]	209
Maxima [F]	209
Giac [F]	210
Mupad [F(-1)]	210
Reduce [F]	210

Optimal result

Integrand size = 23, antiderivative size = 194

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = -\frac{b^2}{3de^4(c + dx)} - \frac{b^2 \arctan(c + dx)}{3de^4} - \frac{b(a + b \arctan(c + dx))}{3de^4(c + dx)^2} + \frac{i(a + b \arctan(c + dx))^2}{3de^4} - \frac{(a + b \arctan(c + dx))^2}{3de^4(c + dx)^3} - \frac{2b(a + b \arctan(c + dx)) \log\left(2 - \frac{2}{1-i(c+dx)}\right)}{3de^4} + \frac{ib^2 \text{PolyLog}\left(2, -1 + \frac{2}{1-i(c+dx)}\right)}{3de^4}$$

output

```
-1/3*b^2/d/e^4/(d*x+c)-1/3*b^2*arctan(d*x+c)/d/e^4-1/3*b*(a+b*arctan(d*x+c
))/d/e^4/(d*x+c)^2+1/3*I*(a+b*arctan(d*x+c))^2/d/e^4-1/3*(a+b*arctan(d*x+c
))^2/d/e^4/(d*x+c)^3-2/3*b*(a+b*arctan(d*x+c))*ln(2-2/(1-I*(d*x+c)))/d/e^4
+1/3*I*b^2*polylog(2,-1+2/(1-I*(d*x+c)))/d/e^4
```

Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.84

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \frac{ab + \frac{a^2}{(c+dx)^3} + \frac{ab}{(c+dx)^2} + \frac{b^2}{c+dx} + b^2 \left(-i + \frac{1}{(c+dx)^3} \right) \arctan(c + dx)^2 + b \arctan(c + dx) \left(b + \frac{2a}{(c+dx)^3} + \frac{1}{c+dx} \right)}{3de^4}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^4,x]
```

output

```
-1/3*(a*b + a^2/(c + d*x)^3 + (a*b)/(c + d*x)^2 + b^2/(c + d*x) + b^2*(-I + (c + d*x)^(-3))*ArcTan[c + d*x]^2 + b*ArcTan[c + d*x]*(b + (2*a)/(c + d*x)^3 + b/(c + d*x)^2 + 2*b*Log[1 - E^((2*I)*ArcTan[c + d*x])]) + 2*a*b*Log[(c + d*x)/Sqrt[1 + (c + d*x)^2]] - I*b^2*PolyLog[2, E^((2*I)*ArcTan[c + d*x])])/(d*e^4)
```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {5566, 27, 5361, 5453, 5361, 264, 216, 5459, 5403, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx \\ & \quad \downarrow \text{5566} \\ & \int \frac{(a + b \arctan(c + dx))^2}{e^4(c + dx)^4} d(c + dx) \\ & \quad \downarrow \text{27} \\ & \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^4} d(c + dx) \\ & \quad \downarrow \\ & \frac{\int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^4} d(c + dx)}{de^4} \end{aligned}$$

$$\begin{aligned}
& \downarrow 5361 \\
& \frac{\frac{2}{3}b \int \frac{a+b \arctan(c+dx)}{(c+dx)^3((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 5453 \\
& \frac{\frac{2}{3}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^3} d(c+dx) - \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) \right) - \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 5361 \\
& \frac{\frac{2}{3}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) + \frac{1}{2}b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2} \right) - \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 264 \\
& \frac{\frac{2}{3}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) + \frac{1}{2}b \left(- \int \frac{1}{(c+dx)^2+1} d(c+dx) - \frac{1}{c+dx} \right) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2} \right) - \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 216 \\
& \frac{\frac{2}{3}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{2(c+dx)^2} + \frac{1}{2}b \left(- \arctan(c+dx) - \frac{1}{c+dx} \right) \right) - \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3}}{de^4} \\
& \downarrow 5459 \\
& \frac{- \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3} + \frac{2}{3}b \left(-i \int \frac{a+b \arctan(c+dx)}{(c+dx)(c+dx+i)} d(c+dx) + \frac{i(a+b \arctan(c+dx))^2}{2b} - \frac{a+b \arctan(c+dx)}{2(c+dx)^2} + \frac{1}{2}b \left(- \arctan(c+dx) - \frac{1}{c+dx} \right) \right)}{de^4} \\
& \downarrow 5403 \\
& \frac{- \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3} + \frac{2}{3}b \left(-i \left(ib \int \frac{\log \left(2 - \frac{2}{1-i(c+dx)} \right)}{(c+dx)^2+1} d(c+dx) - i \log \left(2 - \frac{2}{1-i(c+dx)} \right) (a+b \arctan(c+dx)) \right) \right) + \frac{i(a+b \arctan(c+dx))^2}{2b}}{de^4} \\
& \downarrow 2897 \\
& \frac{- \frac{(a+b \arctan(c+dx))^2}{3(c+dx)^3} + \frac{2}{3}b \left(-i \left(-i \log \left(2 - \frac{2}{1-i(c+dx)} \right) (a+b \arctan(c+dx)) - \frac{1}{2}b \text{PolyLog} \left(2, \frac{2}{1-i(c+dx)} - 1 \right) \right) \right) + \frac{i(a+b \arctan(c+dx))^2}{2b}}{de^4}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^4,x]`

output `(-1/3*(a + b*ArcTan[c + d*x])^2/(c + d*x)^3 + (2*b*((b*(-(c + d*x)^(-1) - ArcTan[c + d*x]))/2 - (a + b*ArcTan[c + d*x])/(2*(c + d*x)^2) + ((I/2)*(a + b*ArcTan[c + d*x])^2)/b - I*((-I)*(a + b*ArcTan[c + d*x])*Log[2 - 2/(1 - I*(c + d*x))] - (b*PolyLog[2, -1 + 2/(1 - I*(c + d*x))])/2)))/3)/(d*e^4)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2897 `Int[Log[u]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & IntegerQ[m])) && NeQ[m, -1]`

rule 5403

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_))), x_
Symbol] := Simp[(a + b*ArcTan[c*x])^p*(Log[2 - 2/(1 + e*(x/d))]/d), x] - Si
mp[b*c*(p/d) Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2 - 2/(1 + e*(x/d))]/(1
+ c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*
d^2 + e^2, 0]
```

rule 5453

```
Int((((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)*((f_.)*(x_))^(m_))/((d_) + (e
_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTan[c*x])^p, x],
x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*((a + b*ArcTan[c*x])^p/(d + e*x^2)
), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

rule 5459

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*d*(p + 1))), x] + Si
mp[I/d Int[(a + b*ArcTan[c*x])^p/(x*(I + c*x)), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && GtQ[p, 0]
```

rule 5566

```
Int((((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m
_.)), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 367 vs. $2(176) = 352$.

Time = 2.45 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.90

method	result
derivativedivides	$-\frac{a^2}{3e^4(dx+c)^3} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{3(dx+c)^3} - \frac{\arctan(dx+c)}{3(dx+c)^2} - \frac{2 \ln(dx+c) \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{i \left(\ln(dx+c-i) \right)}{3} \right)}{3}$
default	$-\frac{a^2}{3e^4(dx+c)^3} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{3(dx+c)^3} - \frac{\arctan(dx+c)}{3(dx+c)^2} - \frac{2 \ln(dx+c) \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{i \left(\ln(dx+c-i) \right)}{3} \right)}{3}$
parts	$-\frac{a^2}{3e^4(dx+c)^3 d} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{3(dx+c)^3} - \frac{\arctan(dx+c)}{3(dx+c)^2} - \frac{2 \ln(dx+c) \arctan(dx+c)}{3} + \frac{\arctan(dx+c) \ln(1+(dx+c)^2)}{3} + \frac{i \left(\ln(dx+c-i) \right)}{3} \right)}{3}$

input `int((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^4,x,method=_RETURNVERBOSE)`

output `1/d*(-1/3*a^2/e^4/(d*x+c)^3+b^2/e^4*(-1/3/(d*x+c)^3*arctan(d*x+c)^2-1/3/(d*x+c)^2*arctan(d*x+c)-2/3*ln(d*x+c)*arctan(d*x+c)+1/3*arctan(d*x+c)*ln(1+(d*x+c)^2)+1/6*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))-1/6*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))-1/3/(d*x+c)-1/3*arctan(d*x+c)-1/3*I*ln(d*x+c)*ln(1+I*(d*x+c))+1/3*I*ln(d*x+c)*ln(1-I*(d*x+c))-1/3*I*dilog(1+I*(d*x+c))+1/3*I*dilog(1-I*(d*x+c)))+2*a*b/e^4*(-1/3/(d*x+c)^3*arctan(d*x+c)-1/6/(d*x+c)^2-1/3*ln(d*x+c)+1/6*ln(1+(d*x+c)^2))`

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^4} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="fricas")`

output `integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)/(d^4*e^4*x^4 + 4*c*d^3*e^4*x^3 + 6*c^2*d^2*e^4*x^2 + 4*c^3*d*e^4*x + c^4*e^4), x)`

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx$$

$$= \int \frac{a^2}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{b^2 \operatorname{atan}^2(c + dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{2ab \operatorname{atan}(c + dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx$$

input `integrate((a+b*atan(d*x+c))**2/(d*e*x+c*e)**4,x)`

output `(Integral(a**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(b**2*atan(c + d*x)**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(2*a*b*atan(c + d*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x))/e**4`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^4} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="maxima")`

output `-1/3*(d*(1/(d^4*e^4*x^2 + 2*c*d^3*e^4*x + c^2*d^2*e^4) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^4) + 2*log(d*x + c)/(d^2*e^4)) + 2*arctan(d*x + c)/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4))*a*b - 1/48*(4*arctan(d*x + c)^2 - 48*(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4)*integrate(1/48*(36*(d^2*x^2 + 2*c*d*x + c^2 + 1)*arctan(d*x + c)^2 + 3*(d^2*x^2 + 2*c*d*x + c^2 + 1)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 8*(d*x + c)*arctan(d*x + c) - 4*(d^2*x^2 + 2*c*d*x + c^2)*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/(d^6*e^4*x^6 + 6*c*d^5*e^4*x^5 + (15*c^2 + 1)*d^4*e^4*x^4 + 4*(5*c^3 + c)*d^3*e^4*x^3 + 3*(5*c^4 + 2*c^2)*d^2*e^4*x^2 + 2*(3*c^5 + 2*c^3)*d*e^4*x + (c^6 + c^4)*e^4), x) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2)*b^2/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4) - 1/3*a^2/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(dex + ce)^4} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^4,x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^2/(d*e*x + c*e)^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^2}{(ce + dex)^4} dx$$

input `int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^4,x)`

output `int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^4, x)`

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^4} dx = \text{Too large to display}$$

input `int((a+b*atan(d*x+c))^2/(d*e*x+c*e)^4,x)`

output

```
(3*atan(c + d*x)**2*b**2*c**5 + 9*atan(c + d*x)**2*b**2*c**4*d*x + 9*atan(c + d*x)**2*b**2*c**3*d**2*x**2 - 3*atan(c + d*x)**2*b**2*c**3 + 3*atan(c + d*x)**2*b**2*c**2*d**3*x**3 - 9*atan(c + d*x)**2*b**2*c**2*d*x - 9*atan(c + d*x)**2*b**2*c*d**2*x**2 - 6*atan(c + d*x)**2*b**2*c - 3*atan(c + d*x)**2*b**2*d**3*x**3 - 12*atan(c + d*x)*a*b*c + 6*atan(c + d*x)*b**2*c**4 + 12*atan(c + d*x)*b**2*c**3*d*x + 6*atan(c + d*x)*b**2*c**2*d**2*x**2 - 8*atan(c + d*x)*b**2*c**2 - 12*atan(c + d*x)*b**2*c*d*x - 6*atan(c + d*x)*b**2*d**2*x**2 - 6*int((atan(c + d*x)*x**2)/(c**6 + 6*c**5*d*x + 15*c**4*d**2*x**2 + c**4 + 20*c**3*d**3*x**3 + 4*c**3*d*x + 15*c**2*d**4*x**4 + 6*c**2*d**2*x**2 + 6*c*d**5*x**5 + 4*c*d**3*x**3 + d**6*x**6 + d**4*x**4),x)*b**2*c**3*d**3 - 18*int((atan(c + d*x)*x**2)/(c**6 + 6*c**5*d*x + 15*c**4*d**2*x**2 + c**4 + 20*c**3*d**3*x**3 + 4*c**3*d*x + 15*c**2*d**4*x**4 + 6*c**2*d**2*x**2 + 6*c*d**5*x**5 + 4*c*d**3*x**3 + d**6*x**6 + d**4*x**4),x)*b**2*c**2*d**4*x - 18*int((atan(c + d*x)*x**2)/(c**6 + 6*c**5*d*x + 15*c**4*d**2*x**2 + c**4 + 20*c**3*d**3*x**3 + 4*c**3*d*x + 15*c**2*d**4*x**4 + 6*c**2*d**2*x**2 + 6*c*d**5*x**5 + 4*c*d**3*x**3 + d**6*x**6 + d**4*x**4),x)*b**2*c*d**5*x**2 - 6*int((atan(c + d*x)*x**2)/(c**6 + 6*c**5*d*x + 15*c**4*d**2*x**2 + c**4 + 20*c**3*d**3*x**3 + 4*c**3*d*x + 15*c**2*d**4*x**4 + 6*c**2*d**2*x**2 + 6*c*d**5*x**5 + 4*c*d**3*x**3 + d**6*x**6 + d**4*x**4),x)*b**2*d**6*x**3 + 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*c**4 + 18...
```

3.22 $\int \frac{(a+b \arctan(c+dx))^2}{(ce+dex)^5} dx$

Optimal result	212
Mathematica [A] (verified)	213
Rubi [A] (verified)	213
Maple [A] (verified)	217
Fricas [B] (verification not implemented)	218
Sympy [F(-1)]	219
Maxima [B] (verification not implemented)	219
Giac [B] (verification not implemented)	220
Mupad [B] (verification not implemented)	221
Reduce [B] (verification not implemented)	222

Optimal result

Integrand size = 23, antiderivative size = 170

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx = -\frac{b^2}{12de^5(c + dx)^2} - \frac{b(a + b \arctan(c + dx))}{6de^5(c + dx)^3} + \frac{b(a + b \arctan(c + dx))}{2de^5(c + dx)} + \frac{(a + b \arctan(c + dx))^2}{4de^5} - \frac{(a + b \arctan(c + dx))^2}{4de^5(c + dx)^4} - \frac{2b^2 \log(c + dx)}{3de^5} + \frac{b^2 \log(1 + (c + dx)^2)}{3de^5}$$

output

```
-1/12*b^2/d/e^5/(d*x+c)^2-1/6*b*(a+b*arctan(d*x+c))/d/e^5/(d*x+c)^3+1/2*b*(a+b*arctan(d*x+c))/d/e^5/(d*x+c)+1/4*(a+b*arctan(d*x+c))^2/d/e^5-1/4*(a+b*arctan(d*x+c))^2/d/e^5/(d*x+c)^4-2/3*b^2*ln(d*x+c)/d/e^5+1/3*b^2*ln(1+(d*x+c)^2)/d/e^5
```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx =$$

$$3a^2 + 2ab(c + dx) + b^2(c + dx)^2 - 6ab(c + dx)^3 - 2b(b(-c + 3c^3 - dx + 9c^2dx + 9cd^2x^2 + 3d^3x^3) +$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^5,x]
```

output

```
-1/12*(3*a^2 + 2*a*b*(c + d*x) + b^2*(c + d*x)^2 - 6*a*b*(c + d*x)^3 - 2*b
*(b*(-c + 3*c^3 - d*x + 9*c^2*d*x + 9*c*d^2*x^2 + 3*d^3*x^3) + 3*a*(-1 + c
^4 + 4*c^3*d*x + 6*c^2*d^2*x^2 + 4*c*d^3*x^3 + d^4*x^4))*ArcTan[c + d*x] -
3*b^2*(-1 + c^4 + 4*c^3*d*x + 6*c^2*d^2*x^2 + 4*c*d^3*x^3 + d^4*x^4)*ArcT
an[c + d*x]^2 + 8*b^2*(c + d*x)^4*Log[c + d*x] - 4*b^2*(c + d*x)^4*Log[1 +
c^2 + 2*c*d*x + d^2*x^2)]/(d*e^5*(c + d*x)^4)
```

Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.92, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.652$, Rules used = {5566, 27, 5361, 5453, 5361, 243, 54, 2009, 5453, 5361, 243, 47, 14, 16, 5419}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx$$

$$\downarrow \text{5566}$$

$$\int \frac{(a + b \arctan(c + dx))^2}{e^5(c + dx)^5} d(c + dx)$$

$$\frac{d}{d}$$

$$\downarrow \text{27}$$

$$\int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^5} d(c + dx)$$

$$\frac{de^5}{de^5}$$

$$\begin{aligned} & \downarrow 5361 \\ & \frac{\frac{1}{2}b \int \frac{a+b \arctan(c+dx)}{(c+dx)^4((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 5453 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^4} d(c+dx) - \int \frac{a+b \arctan(c+dx)}{(c+dx)^2((c+dx)^2+1)} d(c+dx) \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 5361 \\ & \frac{\frac{1}{2}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2((c+dx)^2+1)} d(c+dx) + \frac{1}{3}b \int \frac{1}{(c+dx)^3((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 243 \\ & \frac{\frac{1}{2}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2((c+dx)^2+1)} d(c+dx) + \frac{1}{6}b \int \frac{1}{(c+dx)^4((c+dx)^2+1)} d(c+dx)^2 - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 54 \\ & \frac{\frac{1}{2}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2((c+dx)^2+1)} d(c+dx) + \frac{1}{6}b \int \left(-\frac{1}{(c+dx)^2} + \frac{1}{(c+dx)^4} + \frac{1}{(c+dx)^2+1} \right) d(c+dx)^2 - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 2009 \\ & \frac{\frac{1}{2}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log((c+dx)^2) + \log((c+dx)^2+1) \right) \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 5453 \\ & \frac{\frac{1}{2}b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2} d(c+dx) + \int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log((c+dx)^2) \right) \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 5361 \\ & \frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - b \int \frac{1}{(c+dx)((c+dx)^2+1)} d(c+dx) + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} - \log((c+dx)^2) \right) \right) - \frac{(a+b \arctan(c+dx))^2}{4(c+dx)^4}}{de^5} \\ & \downarrow 243 \end{aligned}$$

$$\frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c+dx)^2 + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} \right) \right)}{de^5}$$

↓ 47

$$\frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}b \left(\int \frac{1}{(c+dx)^2} d(c+dx)^2 - \int \frac{1}{(c+dx)^2+1} d(c+dx)^2 \right) + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} \right)}{de^5}$$

↓ 14

$$\frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}b \left(\log((c+dx)^2) - \int \frac{1}{(c+dx)^2+1} d(c+dx)^2 \right) + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} \right)}{de^5}$$

↓ 16

$$\frac{\frac{1}{2}b \left(\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} - \frac{1}{2}b \left(\log((c+dx)^2) - \log((c+dx)^2+1) \right) + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} \right) \right)}{de^5}$$

↓ 5419

$$\frac{\frac{1}{2}b \left(\frac{(a+b \arctan(c+dx))^2}{2b} + \frac{a+b \arctan(c+dx)}{c+dx} - \frac{a+b \arctan(c+dx)}{3(c+dx)^3} - \frac{1}{2}b \left(\log((c+dx)^2) - \log((c+dx)^2+1) \right) + \frac{1}{6}b \left(-\frac{1}{(c+dx)^2} \right) \right)}{de^5}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(c*e + d*e*x)^5,x]`

output `(-1/4*(a + b*ArcTan[c + d*x])^2/(c + d*x)^4 + (b*(-1/3*(a + b*ArcTan[c + d*x]))/(c + d*x)^3 + (a + b*ArcTan[c + d*x])/(c + d*x) + (a + b*ArcTan[c + d*x])^2/(2*b) - (b*(Log[(c + d*x)^2] - Log[1 + (c + d*x)^2]))/2 + (b*(-(c + d*x)^(-2) - Log[(c + d*x)^2] + Log[1 + (c + d*x)^2]))/6)/2)/(d*e^5)`

Definitions of rubi rules used

- rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$
- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$
- rule 47 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$
- rule 54 $\text{Int}(((a_) + (b_)*(x_))^{(m_)}*((c_) + (d_)*(x_))^{(n_)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{ILtQ}[m, 0] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ !(\text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m + n + 2, 0])$
- rule 243 $\text{Int}[(x_)^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}], x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] \text{ ; FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$
- rule 5361 $\text{Int}(((a_) + \text{ArcTan}[(c_)*(x_)^{(n_)}])*(b_))^{(p_)}*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTan}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTan}[c*x^n])^{(p-1)/(1+c^2*x^{(2*n)})}), x], x] \text{ ; FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

```
rule 5419 Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)/((d_.) + (e_.)*(x_)^2), x_Symbol]
  :=> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x]
  && EqQ[e, c^2*d] && NeQ[p, -1]

rule 5453 Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_.) + (e_.)*(x_)^2),
  x_Symbol] :=> Simp[1/d Int[(f*x)^m*(a + b*ArcTan[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*
  ((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]

rule 5566 Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m_)),
  x_Symbol] :=> Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /;
  FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.96

method	result
derivativedivides	$-\frac{a^2}{4e^5(dx+c)^4} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{4(dx+c)^4} - \frac{\arctan(dx+c)}{6(dx+c)^3} + \frac{\arctan(dx+c)}{2dx+2c} + \frac{\arctan(dx+c)^2}{4} - \frac{1}{12(dx+c)^2} - \frac{2 \ln(dx+c)}{3} + \frac{\ln(1+(dx+c)^2)}{3} \right)}{e^5}$
default	$-\frac{a^2}{4e^5(dx+c)^4} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{4(dx+c)^4} - \frac{\arctan(dx+c)}{6(dx+c)^3} + \frac{\arctan(dx+c)}{2dx+2c} + \frac{\arctan(dx+c)^2}{4} - \frac{1}{12(dx+c)^2} - \frac{2 \ln(dx+c)}{3} + \frac{\ln(1+(dx+c)^2)}{3} \right)}{e^5 d}$
parts	$-\frac{a^2}{4e^5(dx+c)^4 d} + \frac{b^2 \left(-\frac{\arctan(dx+c)^2}{4(dx+c)^4} - \frac{\arctan(dx+c)}{6(dx+c)^3} + \frac{\arctan(dx+c)}{2dx+2c} + \frac{\arctan(dx+c)^2}{4} - \frac{1}{12(dx+c)^2} - \frac{2 \ln(dx+c)}{3} + \frac{\ln(1+(dx+c)^2)}{3} \right)}{e^5 d}$
parallelrisch	$\frac{18x^2 \arctan(dx+c)b^2 c d^7 + 12c d^8 b^2 \arctan(dx+c)^2 x^3 + 18x^2 abc d^7 + 18x \arctan(dx+c)b^2 c^2 d^6 + 18xab c^2 d^6 + 12x \arctan(dx+c)b^2 c^2 d^6}{e^5 d}$
risch	Expression too large to display

```
input int((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^5,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-1/4*a^2/e^5/(d*x+c)^4+b^2/e^5*(-1/4/(d*x+c)^4*arctan(d*x+c)^2-1/6/(d
*x+c)^3*arctan(d*x+c)+1/2/(d*x+c)*arctan(d*x+c)+1/4*arctan(d*x+c)^2-1/12/(
d*x+c)^2-2/3*ln(d*x+c)+1/3*ln(1+(d*x+c)^2))+2*a*b/e^5*(-1/4/(d*x+c)^4*arct
an(d*x+c)-1/12/(d*x+c)^3+1/4/(d*x+c)+1/4*arctan(d*x+c)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(156) = 312$.

Time = 0.15 (sec) , antiderivative size = 448, normalized size of antiderivative = 2.64

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx$$

$$= \frac{6abd^3x^3 + 6abc^3 + (18abc - b^2)d^2x^2 - b^2c^2 - 2abc + 2(9abc^2 - b^2c - ab)dx + 3(b^2d^4x^4 + 4b^2cd^3x^3 +$$

input

```
integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="fricas")
```

output

```
1/12*(6*a*b*d^3*x^3 + 6*a*b*c^3 + (18*a*b*c - b^2)*d^2*x^2 - b^2*c^2 - 2*a
*b*c + 2*(9*a*b*c^2 - b^2*c - a*b)*d*x + 3*(b^2*d^4*x^4 + 4*b^2*c*d^3*x^3
+ 6*b^2*c^2*d^2*x^2 + 4*b^2*c^3*d*x + b^2*c^4 - b^2)*arctan(d*x + c)^2 - 3
*a^2 + 2*(3*a*b*d^4*x^4 + 3*(4*a*b*c + b^2)*d^3*x^3 + 3*a*b*c^4 + 3*b^2*c^
3 + 9*(2*a*b*c^2 + b^2*c)*d^2*x^2 - b^2*c + (12*a*b*c^3 + 9*b^2*c^2 - b^2)
*d*x - 3*a*b)*arctan(d*x + c) + 4*(b^2*d^4*x^4 + 4*b^2*c*d^3*x^3 + 6*b^2*c
^2*d^2*x^2 + 4*b^2*c^3*d*x + b^2*c^4)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 8
*(b^2*d^4*x^4 + 4*b^2*c*d^3*x^3 + 6*b^2*c^2*d^2*x^2 + 4*b^2*c^3*d*x + b^2*
c^4)*log(d*x + c))/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*
c^3*d^2*e^5*x + c^4*d*e^5)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))**2/(d*e*x+c*e)**5,x)`

output `Timed out`

Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 534 vs. $2(156) = 312$.

Time = 0.20 (sec) , antiderivative size = 534, normalized size of antiderivative = 3.14

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx \\ &= \frac{1}{6} \left(d \left(\frac{3d^2x^2 + 6cdx + 3c^2 - 1}{d^5e^5x^3 + 3cd^4e^5x^2 + 3c^2d^3e^5x + c^3d^2e^5} + \frac{3 \arctan\left(\frac{d^2x+cd}{d}\right)}{d^2e^5} \right) - \frac{3 \arctan(dx + c)}{d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5} \right) \\ &+ \frac{1}{12} \left(2d \left(\frac{3d^2x^2 + 6cdx + 3c^2 - 1}{d^5e^5x^3 + 3cd^4e^5x^2 + 3c^2d^3e^5x + c^3d^2e^5} + \frac{3 \arctan\left(\frac{d^2x+cd}{d}\right)}{d^2e^5} \right) \arctan(dx + c) - \frac{3(d^2x^2 + 6cdx + 3c^2 - 1) \arctan(dx + c)}{4(d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5)} \right) \\ &- \frac{b^2 \arctan(dx + c)^2}{4(d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5)} \\ &- \frac{a^2}{4(d^5e^5x^4 + 4cd^4e^5x^3 + 6c^2d^3e^5x^2 + 4c^3d^2e^5x + c^4de^5)} \end{aligned}$$

input `integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="maxima")`

output

```

1/6*(d*((3*d^2*x^2 + 6*c*d*x + 3*c^2 - 1)/(d^5*e^5*x^3 + 3*c*d^4*e^5*x^2 +
3*c^2*d^3*e^5*x + c^3*d^2*e^5) + 3*arctan((d^2*x + c*d)/d)/(d^2*e^5)) - 3
*arctan(d*x + c)/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^
3*d^2*e^5*x + c^4*d*e^5))*a*b + 1/12*(2*d*((3*d^2*x^2 + 6*c*d*x + 3*c^2 -
1)/(d^5*e^5*x^3 + 3*c*d^4*e^5*x^2 + 3*c^2*d^3*e^5*x + c^3*d^2*e^5) + 3*arc
tan((d^2*x + c*d)/d)/(d^2*e^5))*arctan(d*x + c) - (3*(d^2*x^2 + 2*c*d*x +
c^2)*arctan(d*x + c)^2 - 4*(d^2*x^2 + 2*c*d*x + c^2)*log(d^2*x^2 + 2*c*d*x
+ c^2 + 1) + 8*(d^2*x^2 + 2*c*d*x + c^2)*log(d*x + c) + 1)*d^2/(d^5*e^5*x
^2 + 2*c*d^4*e^5*x + c^2*d^3*e^5))*b^2 - 1/4*b^2*arctan(d*x + c)^2/(d^5*e^
5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*e^5*x + c^4*d*e^5)
- 1/4*a^2/(d^5*e^5*x^4 + 4*c*d^4*e^5*x^3 + 6*c^2*d^3*e^5*x^2 + 4*c^3*d^2*
e^5*x + c^4*d*e^5)

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 836 vs. $2(156) = 312$.

Time = 0.34 (sec) , antiderivative size = 836, normalized size of antiderivative = 4.92

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx = \text{Too large to display}$$

input

```
integrate((a+b*arctan(d*x+c))^2/(d*e*x+c*e)^5,x, algorithm="giac")
```

output

```

-1/192*(3*b^2*arctan((d*e*x + c*e)/e)^2*tan(1/2*arctan((d*e*x + c*e)/e))^8
+ 6*a*b*arctan((d*e*x + c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e))^8 - 12*b
^2*arctan((d*e*x + c*e)/e)^2*tan(1/2*arctan((d*e*x + c*e)/e))^6 - 4*b^2*ar
ctan((d*e*x + c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e))^7 + 3*a^2*tan(1/2*a
rctan((d*e*x + c*e)/e))^8 - 24*a*b*arctan((d*e*x + c*e)/e)*tan(1/2*arctan(
(d*e*x + c*e)/e))^6 - 4*a*b*tan(1/2*arctan((d*e*x + c*e)/e))^7 - 30*b^2*ar
ctan((d*e*x + c*e)/e)^2*tan(1/2*arctan((d*e*x + c*e)/e))^4 + 60*b^2*arctan
((d*e*x + c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e))^5 - 12*a^2*tan(1/2*arct
an((d*e*x + c*e)/e))^6 + 4*b^2*tan(1/2*arctan((d*e*x + c*e)/e))^6 - 60*a*b
*arctan((d*e*x + c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e))^4 + 64*b^2*log(1
6*tan(1/2*arctan((d*e*x + c*e)/e))^2/(tan(1/2*arctan((d*e*x + c*e)/e))^4 +
2*tan(1/2*arctan((d*e*x + c*e)/e))^2 + 1))*tan(1/2*arctan((d*e*x + c*e)/e
))^4 + 60*a*b*tan(1/2*arctan((d*e*x + c*e)/e))^5 - 12*b^2*arctan((d*e*x +
c*e)/e)^2*tan(1/2*arctan((d*e*x + c*e)/e))^2 - 60*b^2*arctan((d*e*x + c*e)
/e)*tan(1/2*arctan((d*e*x + c*e)/e))^3 - 30*a^2*tan(1/2*arctan((d*e*x + c*
e)/e))^4 + 8*b^2*tan(1/2*arctan((d*e*x + c*e)/e))^4 - 24*a*b*arctan((d*e*x
+ c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e))^2 - 60*a*b*tan(1/2*arctan((d*e
*x + c*e)/e))^3 + 3*b^2*arctan((d*e*x + c*e)/e)^2 + 4*b^2*arctan((d*e*x +
c*e)/e)*tan(1/2*arctan((d*e*x + c*e)/e)) - 12*a^2*tan(1/2*arctan((d*e*x +
c*e)/e))^2 + 4*b^2*tan(1/2*arctan((d*e*x + c*e)/e))^2 + 6*a*b*arctan((d...

```

Mupad [B] (verification not implemented)

Time = 4.37 (sec) , antiderivative size = 438, normalized size of antiderivative = 2.58

$$\begin{aligned}
& \int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx \\
&= \operatorname{atan}(c + dx)^2 \left(\frac{b^2}{4de^5} - \frac{b^2}{4d^3e^5 \left(\frac{c^4}{d^2} + 6c^2x^2 + d^2x^4 + \frac{4c^3x}{d} + 4cdx^3 \right)} \right) \\
&\quad - \frac{x^2 \left(\frac{b^2d}{2} - 9abcd \right) + x(b^2c - 9abc^2 + ab) + \frac{3a^2 - 6abc^3 + 2abc + b^2c^2}{2d} - 3abd^2x^3}{6c^4e^5 + 24c^3de^5x + 36c^2d^2e^5x^2 + 24cd^3e^5x^3 + 6d^4e^5x^4} \\
&\quad + \frac{\operatorname{atan}(c + dx) \left(\frac{b^2x^3}{2e^5} - \frac{ab}{2d^3e^5} + \frac{b^2c \left(\frac{c^2-1}{3d^2} + \frac{2c^2}{3d^2} \right)}{2de^5} + \frac{b^2x \left(d \left(\frac{c^2-1}{3d^2} + \frac{2c^2}{3d^2} \right) + \frac{2c^2}{d} \right)}{2de^5} + \frac{3b^2cx^2}{2de^5} \right)}{\frac{c^4}{d^2} + 6c^2x^2 + d^2x^4 + \frac{4c^3x}{d} + 4cdx^3} \\
&\quad - \frac{2b^2 \ln(c + dx)}{3de^5} - \frac{\ln(c + dx - i) \left(-\frac{b^2}{3} + \frac{abli}{4} \right)}{de^5} + \frac{\ln(c + dx + li) \left(\frac{b^2}{3} + \frac{liab}{4} \right)}{de^5}
\end{aligned}$$

input `int((a + b*atan(c + d*x))^2/(c*e + d*e*x)^5,x)`

output `atan(c + d*x)^2*(b^2/(4*d*e^5) - b^2/(4*d^3*e^5*(c^4/d^2 + 6*c^2*x^2 + d^2*x^4 + (4*c^3*x)/d + 4*c*d*x^3))) - (x^2*((b^2*d)/2 - 9*a*b*c*d) + x*(a*b + b^2*c - 9*a*b*c^2) + (3*a^2 + b^2*c^2 + 2*a*b*c - 6*a*b*c^3)/(2*d) - 3*a*b*d^2*x^3)/(6*c^4*e^5 + 6*d^4*e^5*x^4 + 24*c*d^3*e^5*x^3 + 36*c^2*d^2*e^5*x^2 + 24*c^3*d*e^5*x) + (atan(c + d*x)*((b^2*x^3)/(2*e^5) - (a*b)/(2*d^3*e^5) + (b^2*c*((c^2 - 1)/(3*d^2) + (2*c^2)/(3*d^2)))/(2*d*e^5) + (b^2*x*(d*((c^2 - 1)/(3*d^2) + (2*c^2)/(3*d^2)) + (2*c^2)/d))/(2*d*e^5) + (3*b^2*c*x^2)/(2*d*e^5)))/(c^4/d^2 + 6*c^2*x^2 + d^2*x^4 + (4*c^3*x)/d + 4*c*d*x^3) - (2*b^2*log(c + d*x))/(3*d*e^5) - (log(c + d*x - 1i)*((a*b*1i)/4 - b^2/3))/(d*e^5) + (log(c + d*x + 1i)*((a*b*1i)/4 + b^2/3))/(d*e^5)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 679, normalized size of antiderivative = 3.99

$$\int \frac{(a + b \arctan(c + dx))^2}{(ce + dex)^5} dx$$

$$= \frac{-16 \log(dx + c) b^2 c^5 + 12 \operatorname{atan}(dx + c) b^2 c d^3 x^3 + 8 \log(d^2 x^2 + 2cdx + c^2 + 1) b^2 c d^4 x^4 - 16 \log(dx + c) b^2 c d^5}{(ce + dex)^5}$$

input `int((a+b*atan(d*x+c))^2/(d*e*x+c*e)^5,x)`

output

```
(6*atan(c + d*x)**2*b**2*c**5 + 24*atan(c + d*x)**2*b**2*c**4*d*x + 36*atan(c + d*x)**2*b**2*c**3*d**2*x**2 + 24*atan(c + d*x)**2*b**2*c**2*d**3*x**3 + 6*atan(c + d*x)**2*b**2*c*d**4*x**4 - 6*atan(c + d*x)**2*b**2*c + 12*atan(c + d*x)*a*b*c**5 + 48*atan(c + d*x)*a*b*c**4*d*x + 72*atan(c + d*x)*a*b*c**3*d**2*x**2 + 48*atan(c + d*x)*a*b*c**2*d**3*x**3 + 12*atan(c + d*x)*a*b*c*d**4*x**4 - 12*atan(c + d*x)*a*b*c + 12*atan(c + d*x)*b**2*c**4 + 36*atan(c + d*x)*b**2*c**3*d*x + 36*atan(c + d*x)*b**2*c**2*d**2*x**2 - 4*atan(c + d*x)*b**2*c**2 + 12*atan(c + d*x)*b**2*c*d**3*x**3 - 4*atan(c + d*x)*b**2*c*d*x + 8*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c**5 + 32*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c**4*d*x + 48*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c**3*d**2*x**2 + 32*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c**2*d**3*x**3 + 8*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*c*d**4*x**4 - 16*log(c + d*x)*b**2*c**5 - 64*log(c + d*x)*b**2*c**4*d*x - 96*log(c + d*x)*b**2*c**3*d**2*x**2 - 64*log(c + d*x)*b**2*c**2*d**3*x**3 - 16*log(c + d*x)*b**2*c*d**4*x**4 - 6*a**2*c + 9*a*b*c**4 + 24*a*b*c**3*d*x + 18*a*b*c**2*d**2*x**2 - 4*a*b*c**2 - 4*a*b*c*d*x - 3*a*b*d**4*x**4 - 2*b**2*c**3 - 4*b**2*c**2*d*x - 2*b**2*c*d**2*x**2)/(24*c*d**5*(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4))
```


3.23 $\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx$

Optimal result	224
Mathematica [A] (verified)	225
Rubi [A] (verified)	225
Maple [C] (warning: unable to verify)	229
Fricas [F]	230
Sympy [F]	231
Maxima [F]	231
Giac [F]	232
Mupad [F(-1)]	233
Reduce [F]	233

Optimal result

Integrand size = 23, antiderivative size = 271

$$\begin{aligned}
 & \int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx \\
 &= ab^2 e^2 x + \frac{b^3 e^2 (c + dx) \arctan(c + dx)}{d} - \frac{be^2 (a + b \arctan(c + dx))^2}{2d} \\
 & \quad - \frac{be^2 (c + dx)^2 (a + b \arctan(c + dx))^2}{2d} - \frac{ie^2 (a + b \arctan(c + dx))^3}{3d} \\
 & \quad + \frac{e^2 (c + dx)^3 (a + b \arctan(c + dx))^3}{3d} - \frac{be^2 (a + b \arctan(c + dx))^2 \log\left(\frac{2}{1+i(c+dx)}\right)}{d} \\
 & \quad - \frac{b^3 e^2 \log(1 + (c + dx)^2)}{2d} - \frac{ib^2 e^2 (a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d} \\
 & \quad - \frac{b^3 e^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2d}
 \end{aligned}$$

output

```

a*b^2*e^2*x+b^3*e^2*(d*x+c)*arctan(d*x+c)/d-1/2*b*e^2*(a+b*arctan(d*x+c))^
2/d-1/2*b*e^2*(d*x+c)^2*(a+b*arctan(d*x+c))^2/d-1/3*I*e^2*(a+b*arctan(d*x+
c))^3/d+1/3*e^2*(d*x+c)^3*(a+b*arctan(d*x+c))^3/d-b*e^2*(a+b*arctan(d*x+c)
)^2*ln(2/(1+I*(d*x+c)))/d-1/2*b^3*e^2*ln(1+(d*x+c)^2)/d-I*b^2*e^2*(a+b*arc
tan(d*x+c))*polylog(2,1-2/(1+I*(d*x+c)))/d-1/2*b^3*e^2*polylog(3,1-2/(1+I*
(d*x+c)))/d

```

Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.29

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx$$

$$= \frac{e^2 \left(-3a^2b(c + dx)^2 + 2a^3(c + dx)^3 + 6a^2b(c + dx)^3 \arctan(c + dx) + 3a^2b \log(1 + (c + dx)^2) + 6ab^2(c + dx)^3 \arctan^2(c + dx) - 3a^2b \arctan^3(c + dx) + 6ab^2 \log(1 + (c + dx)^2) \arctan(c + dx) + 3ab^2 \log^2(1 + (c + dx)^2) + 6ab^2 \arctan(c + dx) \log(1 + (c + dx)^2) + 3ab^2 \arctan^2(c + dx) \right)}{6d}$$

input

```
Integrate[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x])^3,x]
```

output

```
(e^2*(-3*a^2*b*(c + d*x)^2 + 2*a^3*(c + d*x)^3 + 6*a^2*b*(c + d*x)^3*ArcTan[c + d*x] + 3*a^2*b*Log[1 + (c + d*x)^2] + 6*a*b^2*(c + d*x - ArcTan[c + d*x] - (c + d*x)^2*ArcTan[c + d*x] + I*ArcTan[c + d*x]^2 + (c + d*x)^3*ArcTan[c + d*x]^2 - 2*ArcTan[c + d*x]*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + I*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) + b^3*(6*(c + d*x)*ArcTan[c + d*x] - 3*(1 + (c + d*x)^2)*ArcTan[c + d*x]^2 + (2*I)*ArcTan[c + d*x]^3 - 2*(c + d*x)*ArcTan[c + d*x]^3 + 2*(c + d*x)*(1 + (c + d*x)^2)*ArcTan[c + d*x]^3 - 6*ArcTan[c + d*x]^2*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + 6*Log[1/Sqrt[1 + (c + d*x)^2]] + (6*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) - 3*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])]))/(6*d)
```

Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.85, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$, Rules used = {5566, 27, 5361, 5451, 5361, 5451, 2009, 5419, 5455, 5379, 5529, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx$$

$$\downarrow 5566$$

$$\frac{\int e^2 (c + dx)^2 (a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{e^2 \int (c + dx)^2 (a + b \arctan(c + dx))^3 d(c + dx)}{d} \\ & \downarrow 5361 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \int \frac{(c+dx)^3 (a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c + dx) \right)}{d} \\ & \downarrow 5451 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(\int (c + dx) (a + b \arctan(c + dx))^2 d(c + dx) - \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c + dx) \right) \right)}{d} \\ & \downarrow 5361 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(-b \int \frac{(c+dx)^2 (a+b \arctan(c+dx))}{(c+dx)^2+1} d(c + dx) - \int \frac{(c+dx)(a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c + dx) \right) \right)}{d} \\ & \downarrow 5451 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(-b \left(\int (a + b \arctan(c + dx)) d(c + dx) - \int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c + dx) \right) \right) \right)}{d} \\ & \downarrow 2009 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(- \int \frac{(c+dx)(a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c + dx) - b \left(- \int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c + dx) \right) \right) \right)}{d} \\ & \downarrow 5419 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(- \int \frac{(c+dx)(a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c + dx) + \frac{1}{2} (c + dx)^2 (a + b \arctan(c + dx)) \right) \right)}{d} \\ & \downarrow 5455 \\ & \frac{e^2 \left(\frac{1}{3} (c + dx)^3 (a + b \arctan(c + dx))^3 - b \left(\int \frac{(a+b \arctan(c+dx))^2}{-c-dx+i} d(c + dx) + \frac{i(a+b \arctan(c+dx))^3}{3b} + \frac{1}{2} (c + dx)^2 (a + b \arctan(c + dx)) \right) \right)}{d} \\ & \downarrow 5379 \end{aligned}$$

$$e^2 \left(\frac{1}{3}(c+dx)^3(a+b\arctan(c+dx))^3 - b \left(-2b \int \frac{(a+b\arctan(c+dx)) \log\left(\frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx) + \frac{i(a+b\arctan(c+dx))^3}{3b} + \right. \right.$$

↓ 5529

$$e^2 \left(\frac{1}{3}(c+dx)^3(a+b\arctan(c+dx))^3 - b \left(-2b \left(\frac{1}{2}ib \int \frac{\text{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx) - \frac{1}{2}i \text{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) \right. \right. \right.$$

↓ 7164

$$e^2 \left(\frac{1}{3}(c+dx)^3(a+b\arctan(c+dx))^3 - b \left(-2b \left(-\frac{1}{2}i \text{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a+b\arctan(c+dx)) - \frac{1}{4}b \text{PolyLog}\left(3, 1 - \frac{2}{i(c+dx)+1}\right) \right) \right)$$

input `Int[(c*e + d*e*x)^2*(a + b*ArcTan[c + d*x])^3,x]`

output `(e^2*(((c + d*x)^3*(a + b*ArcTan[c + d*x])^3)/3 - b*(((c + d*x)^2*(a + b*ArcTan[c + d*x])^2)/2 + ((I/3)*(a + b*ArcTan[c + d*x])^3)/b + (a + b*ArcTan[c + d*x])^2*Log[2/(1 + I*(c + d*x))] - b*(a*(c + d*x) + b*(c + d*x)*ArcTan[c + d*x] - (a + b*ArcTan[c + d*x])^2/(2*b) - (b*Log[1 + (c + d*x)^2])/2) - 2*b*((-1/2*I)*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] - (b*PolyLog[3, 1 - 2/(1 + I*(c + d*x))])/4)))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5361 $\text{Int}[(a + \text{ArcTan}[c \cdot x^n] \cdot b)^p \cdot x^m, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot \text{ArcTan}[c \cdot x^n])^p / (m+1), x] - \text{Simp}[b \cdot c \cdot n \cdot (p/(m+1)) \text{Int}[x^{m+n} \cdot (a + b \cdot \text{ArcTan}[c \cdot x^n])^{p-1} / (1 + c^2 \cdot x^{2n})], x], x] /;$ $\text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$

rule 5379 $\text{Int}[(a + \text{ArcTan}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x), x_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTan}[c \cdot x])^p \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/e), x] + \text{Simp}[b \cdot c \cdot (p/e) \text{Int}[(a + b \cdot \text{ArcTan}[c \cdot x])^{p-1} \cdot (\text{Log}[2/(1 + e \cdot (x/d))]/(1 + c^2 \cdot x^2))], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d^2 + e^2, 0]$

rule 5419 $\text{Int}[(a + \text{ArcTan}[c \cdot x] \cdot b)^p / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot \text{ArcTan}[c \cdot x])^{p+1} / (b \cdot c \cdot d \cdot (p+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{NeQ}[p, -1]$

rule 5451 $\text{Int}[(a + \text{ArcTan}[c \cdot x] \cdot b)^p \cdot (f \cdot x)^m / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[f^2/e \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p, x] - \text{Simp}[d \cdot (f^2/e) \text{Int}[(f \cdot x)^{m-2} \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p / (d + e \cdot x^2)], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{GtQ}[m, 1]$

rule 5455 $\text{Int}[(a + \text{ArcTan}[c \cdot x] \cdot b)^p \cdot x / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(-I) \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^{p+1} / (b \cdot e \cdot (p+1)), x] - \text{Simp}[1/(c \cdot d) \text{Int}[(a + b \cdot \text{ArcTan}[c \cdot x])^p / (I - c \cdot x), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{IGtQ}[p, 0]$

rule 5529 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTan}[c \cdot x] \cdot b)^p) / ((d) + (e) \cdot x^2), x_Symbol] \rightarrow \text{Simp}[(-I) \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot p \cdot (I/2) \text{Int}[(a + b \cdot \text{ArcTan}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2 \cdot (I / (I - c \cdot x)))^2, 0]$

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.10 (sec) , antiderivative size = 1277, normalized size of antiderivative = 4.71

method	result	size
derivativedivides	Expression too large to display	1277
default	Expression too large to display	1277
parts	Expression too large to display	1285

input

```
int((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```

1/d*(1/3*e^2*a^3*(d*x+c)^3+e^2*b^3*(1/3*(d*x+c)^3*arctan(d*x+c)^3-1/2*(d*x
+c)^2*arctan(d*x+c)^2+1/2*arctan(d*x+c)^2*ln(1+(d*x+c)^2)-1/4*I*Pi*csgn(I/
(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/
(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^2*arctan(d*x+c)^2-1/2*polylog(3,-(1+I
*(d*x+c))^2/(1+(d*x+c)^2))-arctan(d*x+c)^2*ln((1+I*(d*x+c))/(1+(d*x+c)^2)^
(1/2))+1/4*I*Pi*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+I*(d
*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+
c))^2/(1+(d*x+c)^2))^2)*arctan(d*x+c)^2+1/2*I*Pi*csgn(I*(1+(1+I*(d*x+c))^2
/(1+(d*x+c)^2))^2)*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^2*arctan(d*x+
c)^2-1/4*I*Pi*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+(1+I*(
d*x+c))^2/(1+(d*x+c)^2))^2)*arctan(d*x+c)^2+1/4*I*Pi*csgn(I*(1+I*(d*x+c))^
2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^3*arctan(d*x+c)^2-1/4
*I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1+I*(d*x+c))^2/(1+(d*x
+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^2*arctan(d*x+c)^2-ln(2)*arctan
(d*x+c)^2+ln(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))-1/4*I*Pi*csgn(I*(1+(1+I*(d*x
+c))^2/(1+(d*x+c)^2))^2)^3*arctan(d*x+c)^2+arctan(d*x+c)*(d*x+c-I)-1/2*I*P
i*csgn(I*(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))*csgn(I*(1+I*(d*x+c))^2/(1+(d*x
+c)^2))^2*arctan(d*x+c)^2+1/4*I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))^3
*arctan(d*x+c)^2+1/4*I*Pi*csgn(I*(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))^2*csgn
(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))*arctan(d*x+c)^2+I*arctan(d*x+c)*poly1...

```

Fricas [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^3 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="fricas")
```

output

```

integral(a^3*d^2*e^2*x^2 + 2*a^3*c*d*e^2*x + a^3*c^2*e^2 + (b^3*d^2*e^2*x^
2 + 2*b^3*c*d*e^2*x + b^3*c^2*e^2)*arctan(d*x + c)^3 + 3*(a*b^2*d^2*e^2*x^
2 + 2*a*b^2*c*d*e^2*x + a*b^2*c^2*e^2)*arctan(d*x + c)^2 + 3*(a^2*b*d^2*e^
2*x^2 + 2*a^2*b*c*d*e^2*x + a^2*b*c^2*e^2)*arctan(d*x + c), x)

```

Sympy [F]

$$\begin{aligned}
& \int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx \\
&= e^2 \left(\int a^3 c^2 dx + \int a^3 d^2 x^2 dx + \int b^3 c^2 \operatorname{atan}^3(c + dx) dx \right. \\
&\quad + \int 3ab^2 c^2 \operatorname{atan}^2(c + dx) dx + \int 3a^2 bc^2 \operatorname{atan}(c + dx) dx + \int 2a^3 c dx dx \\
&\quad + \int b^3 d^2 x^2 \operatorname{atan}^3(c + dx) dx + \int 3ab^2 d^2 x^2 \operatorname{atan}^2(c + dx) dx \\
&\quad + \int 3a^2 bd^2 x^2 \operatorname{atan}(c + dx) dx + \int 2b^3 c dx \operatorname{atan}^3(c + dx) dx \\
&\quad \left. + \int 6ab^2 c dx \operatorname{atan}^2(c + dx) dx + \int 6a^2 b c dx \operatorname{atan}(c + dx) dx \right)
\end{aligned}$$

input `integrate((d*e*x+c*e)**2*(a+b*atan(d*x+c))**3,x)`

output `e**2*(Integral(a**3*c**2, x) + Integral(a**3*d**2*x**2, x) + Integral(b**3*c**2*atan(c + d*x)**3, x) + Integral(3*a*b**2*c**2*atan(c + d*x)**2, x) + Integral(3*a**2*b*c**2*atan(c + d*x), x) + Integral(2*a**3*c*d*x, x) + Integral(b**3*d**2*x**2*atan(c + d*x)**3, x) + Integral(3*a*b**2*d**2*x**2*atan(c + d*x)**2, x) + Integral(3*a**2*b*d**2*x**2*atan(c + d*x), x) + Integral(2*b**3*c*d*x*atan(c + d*x)**3, x) + Integral(6*a*b**2*c*d*x*atan(c + d*x)**2, x) + Integral(6*a**2*b*c*d*x*atan(c + d*x), x))`

Maxima [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output

```

7/8*b^3*c^4*e^2*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 3*a*b^2*c^4*
e^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - (3*arctan(d*x + c)*arcta
n((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*a*b^2*c^4*e^2 - 7/32
*(6*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)^2/d - 4*arctan(d*x + c)*arct
an((d^2*x + c*d)/d)^3/d + arctan((d^2*x + c*d)/d)^4/d)*b^3*c^4*e^2 + 1/3*a
^3*d^2*e^2*x^3 + 7/8*b^3*c^2*e^2*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)
/d + 28*b^3*d^4*e^2*integrate(1/32*x^4*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x
+ c^2 + 1), x) + 3*b^3*d^4*e^2*integrate(1/32*x^4*arctan(d*x + c)*log(d^
2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 96*a*b^2*
d^4*e^2*integrate(1/32*x^4*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1)
, x) + 112*b^3*c*d^3*e^2*integrate(1/32*x^3*arctan(d*x + c)^3/(d^2*x^2 + 2
*c*d*x + c^2 + 1), x) + 4*b^3*d^4*e^2*integrate(1/32*x^4*arctan(d*x + c)*l
og(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^3
*c*d^3*e^2*integrate(1/32*x^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2
+ 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 384*a*b^2*c*d^3*e^2*integrate(1
/32*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 168*b^3*c^2*
d^2*e^2*integrate(1/32*x^2*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1)
, x) + 16*b^3*c*d^3*e^2*integrate(1/32*x^3*arctan(d*x + c)*log(d^2*x^2 + 2
*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 18*b^3*c^2*d^2*e^2*i
ntegrate(1/32*x^2*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d...

```

Giac [F]

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx = \int (dex + ce)^2 (b \arctan(dx + c) + a)^3 dx$$

input

```
integrate((d*e*x+c*e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="giac")
```

output

```
sage0*x
```

Mupad [F(-1)]

Timed out.

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx = \int (ce + dex)^2 (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((c*e + d*e*x)^2*(a + b*atan(c + d*x))^3,x)`output `int((c*e + d*e*x)^2*(a + b*atan(c + d*x))^3, x)`**Reduce [F]**

$$\int (ce + dex)^2 (a + b \arctan(c + dx))^3 dx$$

$$= \frac{e^2 \left(6 \operatorname{atan}(dx + c)^3 b^3 c^2 dx + 6 \operatorname{atan}(dx + c)^3 b^3 c d^2 x^2 + 6 \operatorname{atan}(dx + c)^2 a b^2 d^3 x^3 - 6 \operatorname{atan}(dx + c)^2 b^3 c dx - \dots \right)}{\dots}$$

input `int((d*e*x+c*e)^2*(a+b*atan(d*x+c))^3,x)`output `(e**2*(2*atan(c + d*x)**3*b**3*c**3 + 6*atan(c + d*x)**3*b**3*c**2*d*x + 6*atan(c + d*x)**3*b**3*c*d**2*x**2 + 2*atan(c + d*x)**3*b**3*c + 2*atan(c + d*x)**3*b**3*d**3*x**3 + 6*atan(c + d*x)**2*a*b**2*c**3 + 18*atan(c + d*x)**2*a*b**2*c**2*d*x + 18*atan(c + d*x)**2*a*b**2*c*d**2*x**2 + 6*atan(c + d*x)**2*a*b**2*c + 6*atan(c + d*x)**2*a*b**2*d**3*x**3 - 3*atan(c + d*x)**2*b**3*c**2 - 6*atan(c + d*x)**2*b**3*c*d*x - 3*atan(c + d*x)**2*b**3*d*x**2 - 3*atan(c + d*x)**2*b**3 + 6*atan(c + d*x)*a**2*b*c**3 + 18*atan(c + d*x)*a**2*b*c**2*d*x + 18*atan(c + d*x)*a**2*b*c*d**2*x**2 + 6*atan(c + d*x)*a**2*b*d**3*x**3 - 6*atan(c + d*x)*a*b**2*c**2 - 12*atan(c + d*x)*a*b**2*c*d*x - 6*atan(c + d*x)*a*b**2*d**2*x**2 - 6*atan(c + d*x)*a*b**2 + 6*atan(c + d*x)*b**3*c + 6*atan(c + d*x)*b**3*d*x + 12*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*a*b**2*d**2 + 6*int((atan(c + d*x)*2*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**3*d**2 + 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a**2*b - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**3 + 6*a**3*c**2*d*x + 6*a**3*c*d**2*x**2 + 2*a**3*d**3*x**3 - 6*a**2*b*c*d*x - 3*a**2*b*d**2*x**2 + 6*a*b**2*d*x))/(6*d)`

3.24 $\int (ce + dex)(a + b \arctan(c + dx))^3 dx$

Optimal result	234
Mathematica [A] (verified)	235
Rubi [A] (verified)	235
Maple [B] (verified)	239
Fricas [F]	239
Sympy [F]	240
Maxima [F]	241
Giac [F]	241
Mupad [F(-1)]	242
Reduce [F]	242

Optimal result

Integrand size = 21, antiderivative size = 164

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = -\frac{3ibe(a + b \arctan(c + dx))^2}{2d} - \frac{3be(c + dx)(a + b \arctan(c + dx))^2}{2d} + \frac{e(a + b \arctan(c + dx))^3}{2d} + \frac{e(c + dx)^2(a + b \arctan(c + dx))^3}{2d} - \frac{3b^2e(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d} - \frac{3ib^3e \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{2d}$$

output

```
-3/2*I*b*e*(a+b*arctan(d*x+c))^2/d-3/2*b*e*(d*x+c)*(a+b*arctan(d*x+c))^2/d
+1/2*e*(a+b*arctan(d*x+c))^3/d+1/2*e*(d*x+c)^2*(a+b*arctan(d*x+c))^3/d-3*b
^2*e*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/d-3/2*I*b^3*e*polylog(2,1-2/(
1+I*(d*x+c)))/d
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.20

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx$$

$$= \frac{e \left(3b^2(-i + c + dx)(-b + a(i + c + dx)) \arctan(c + dx)^2 + b^3(1 + c^2 + 2cdx + d^2x^2) \arctan(c + dx)^3 + \dots \right)}{d}$$

input

```
Integrate[(c*e + d*e*x)*(a + b*ArcTan[c + d*x])^3,x]
```

output

```
(e*(3*b^2*(-I + c + d*x)*(-b + a*(I + c + d*x))*ArcTan[c + d*x]^2 + b^3*(1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTan[c + d*x]^3 + 3*b*ArcTan[c + d*x]*(a*(-2*b*(c + d*x) + a*(1 + c^2 + 2*c*d*x + d^2*x^2)) - 2*b^2*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + a*(a*(c + d*x)*(-3*b + a*c + a*d*x) - 6*b^2*Log[1/Sqrt[1 + (c + d*x)^2]]) + (3*I)*b^3*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]))/(2*d)
```

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.92, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.476$, Rules used = {5566, 27, 5361, 5451, 5345, 5419, 5455, 5379, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx$$

$$\downarrow 5566$$

$$\frac{\int e(c + dx)(a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow 27$$

$$\frac{e \int (c + dx)(a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

$$\begin{aligned} & \downarrow 5361 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b \int \frac{(c+dx)^2(a+b\arctan(c+dx))^2}{(c+dx)^2+1} d(c+dx)\right)}{d} \\ & \downarrow 5451 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(\int (a+b\arctan(c+dx))^2 d(c+dx) - \int \frac{(a+b\arctan(c+dx))^2}{(c+dx)^2+1} d(c+dx)\right)\right)}{d} \\ & \downarrow 5345 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b \int \frac{(c+dx)(a+b\arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \int \frac{(a+b\arctan(c+dx))^2}{(c+dx)^2+1} d(c+dx)\right)\right)}{d} \\ & \downarrow 5419 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b \int \frac{(c+dx)(a+b\arctan(c+dx))}{(c+dx)^2+1} d(c+dx) - \frac{(a+b\arctan(c+dx))^3}{3b} + (c+dx)\right)\right)}{d} \\ & \downarrow 5455 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b\left(-\int \frac{a+b\arctan(c+dx)}{-c-dx+i} d(c+dx) - \frac{i(a+b\arctan(c+dx))^2}{2b}\right) - \frac{(a+b\arctan(c+dx))^3}{3b}\right)\right)}{d} \\ & \downarrow 5379 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b\left(b \int \frac{\log\left(\frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c+dx) - \frac{i(a+b\arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right)\right)\right)}{d} \\ & \downarrow 2849 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b\left(-ib \int \frac{\log\left(\frac{2}{i(c+dx)+1}\right)}{1-\frac{2}{i(c+dx)+1}} d_{\frac{1}{i(c+dx)+1}} - \frac{i(a+b\arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right)\right)\right)}{d} \\ & \downarrow 2752 \\ & \frac{e\left(\frac{1}{2}(c+dx)^2(a+b\arctan(c+dx))^3 - \frac{3}{2}b\left(-2b\left(-\frac{i(a+b\arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right)\right)(a+b\arctan(c+dx)) - \right)}{d} \end{aligned}$$

input `Int[(c*e + d*e*x)*(a + b*ArcTan[c + d*x])^3,x]`

output `(e*(((c + d*x)^2*(a + b*ArcTan[c + d*x])^3)/2 - (3*b*((c + d*x)*(a + b*ArcTan[c + d*x])^2 - (a + b*ArcTan[c + d*x])^3/(3*b) - 2*b*((-1/2*I)*(a + b*ArcTan[c + d*x])^2)/b - (a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] - (I/2)*b*PolyLog[2, 1 - 2/(1 + I*(c + d*x))])))/2))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] :> Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

rule 5345 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] & IntegerQ[m])) && NeQ[m, -1]`

rule 5379 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
:> Simp[(-a + b*ArcTan[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(
p/e Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2))
, x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0
]`

rule 5419 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
:> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b,
c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]`

rule 5451 `Int[(((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)*((f_.)*(x_)^(m_))/((d_) + (e
.)*(x)^2), x_Symbol] := Simp[f^2/e Int[(f*x)^(m - 2)*(a + b*ArcTan[c*x]
)^p, x], x] - Simp[d*(f^2/e Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/(d
+ e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]`

rule 5455 `Int[(((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Si
mp[1/(c*d) Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]`

rule 5566 `Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m
_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(150) = 300$.

Time = 0.65 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.96

method	result
derivativedivides	$\frac{e a^3 (dx+c)^2}{2} + e b^3 \left(\frac{(dx+c)^2 \arctan(dx+c)^3}{2} + \frac{\arctan(dx+c)^3}{2} - \frac{3(dx+c) \arctan(dx+c)^2}{2} + \frac{3 \arctan(dx+c) \ln(1+(dx+c)^2)}{2} + \frac{3i \left(\ln \right)}{2} \right)$
default	$\frac{e a^3 (dx+c)^2}{2} + e b^3 \left(\frac{(dx+c)^2 \arctan(dx+c)^3}{2} + \frac{\arctan(dx+c)^3}{2} - \frac{3(dx+c) \arctan(dx+c)^2}{2} + \frac{3 \arctan(dx+c) \ln(1+(dx+c)^2)}{2} + \frac{3i \left(\ln \right)}{2} \right)$
parts	$e a^3 \left(\frac{1}{2} dx^2 + cx \right) + \frac{e b^3 \left(\frac{(dx+c)^2 \arctan(dx+c)^3}{2} + \frac{\arctan(dx+c)^3}{2} - \frac{3(dx+c) \arctan(dx+c)^2}{2} + \frac{3 \arctan(dx+c) \ln(1+(dx+c)^2)}{2} \right)}{2}$
risch	Expression too large to display

input

```
int((d*e*x+c*e)*(a+b*arctan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(1/2*e*a^3*(d*x+c)^2+e*b^3*(1/2*(d*x+c)^2*arctan(d*x+c)^3+1/2*arctan(d*x+c)^3-3/2*(d*x+c)*arctan(d*x+c)^2+3/2*arctan(d*x+c)*ln(1+(d*x+c)^2)+3/4*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))-3/4*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I))))+3*e*a*b^2*(1/2*(d*x+c)^2*arctan(d*x+c)^2+1/2*arctan(d*x+c)^2-(d*x+c)*arctan(d*x+c)+1/2*ln(1+(d*x+c)^2))+3*e*a^2*b*(1/2*(d*x+c)^2*arctan(d*x+c)-1/2*d*x-1/2*c+1/2*arctan(d*x+c))
```

Fricas [F]

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = \int (dex + ce)(b \arctan(dx + c) + a)^3 dx$$

input

```
integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^3,x, algorithm="fricas")
```


output

```
integral(a^3*d*e*x + a^3*c*e + (b^3*d*e*x + b^3*c*e)*arctan(d*x + c)^3 + 3
*(a*b^2*d*e*x + a*b^2*c*e)*arctan(d*x + c)^2 + 3*(a^2*b*d*e*x + a^2*b*c*e)
*arctan(d*x + c), x)
```

Sympy [F]

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = e \left(\int a^3 c dx + \int a^3 dx dx \right. \\ \left. + \int b^3 c \operatorname{atan}^3(c + dx) dx \right. \\ \left. + \int 3ab^2 c \operatorname{atan}^2(c + dx) dx \right. \\ \left. + \int 3a^2 bc \operatorname{atan}(c + dx) dx \right. \\ \left. + \int b^3 dx \operatorname{atan}^3(c + dx) dx \right. \\ \left. + \int 3ab^2 dx \operatorname{atan}^2(c + dx) dx \right. \\ \left. + \int 3a^2 b dx \operatorname{atan}(c + dx) dx \right)$$

input

```
integrate((d*e*x+c*e)*(a+b*atan(d*x+c))**3,x)
```

output

```
e*(Integral(a**3*c, x) + Integral(a**3*d*x, x) + Integral(b**3*c*atan(c +
d*x)**3, x) + Integral(3*a*b**2*c*atan(c + d*x)**2, x) + Integral(3*a**2*b
*c*atan(c + d*x), x) + Integral(b**3*d*x*atan(c + d*x)**3, x) + Integral(3
*a*b**2*d*x*atan(c + d*x)**2, x) + Integral(3*a**2*b*d*x*atan(c + d*x), x)
)
```

Maxima [F]

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = \int (dex + ce)(b \arctan(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output

```

1/2*a^3*d*e*x^2 + 3/2*(x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((
d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3))*a^2*b*d*e +
a^3*c*e*x + 3/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*a^2*
b*c*e/d + 1/32*(8*(b^3*d^2*e*x^2 + 2*b^3*c*d*e*x + (b^3*c^2 + b^3)*e)*arct
an(d*x + c)^3 + 12*(a*b^2*d^2*e*x^2 + (2*a*b^2*c - b^3)*d*e*x)*arctan(d*x
+ c)^2 - 3*(a*b^2*d^2*e*x^2 + (2*a*b^2*c - b^3)*d*e*x)*log(d^2*x^2 + 2*c*d
*x + c^2 + 1)^2 + 4*(4*b^3*c^3*e*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)
/d + 18*a*b^2*c^3*e*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 6*(3*arc
tan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*a*
b^2*c^3*e - (6*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)^2/d - 4*arctan(d*
x + c)*arctan((d^2*x + c*d)/d)^3/d + arctan((d^2*x + c*d)/d)^4/d)*b^3*c^3*
e - 3*b^3*c^2*e*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d + 4*b^3*c*e*ar
ctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 128*b^3*d^3*e*integrate(1/32*x
^3*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 576*a*b^2*d^3*e*i
ntegrate(1/32*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 38
4*b^3*c*d^2*e*integrate(1/32*x^2*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^
2 + 1), x) + 48*a*b^2*d^3*e*integrate(1/32*x^3*log(d^2*x^2 + 2*c*d*x + c^2
+ 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 1728*a*b^2*c*d^2*e*integrate(1
/32*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 384*b^3*c^2*
d*e*integrate(1/32*x*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x...

```

Giac [F]

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = \int (dex + ce)(b \arctan(dx + c) + a)^3 dx$$

input `integrate((d*e*x+c*e)*(a+b*arctan(d*x+c))^3,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx = \int (ce + dex) (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((c*e + d*e*x)*(a + b*atan(c + d*x))^3,x)`

output `int((c*e + d*e*x)*(a + b*atan(c + d*x))^3, x)`

Reduce [F]

$$\int (ce + dex)(a + b \arctan(c + dx))^3 dx$$

$$= \frac{e \left(\operatorname{atan}(dx + c)^3 b^3 c^2 + 2 \operatorname{atan}(dx + c)^3 b^3 c dx + \operatorname{atan}(dx + c)^3 b^3 d^2 x^2 + \operatorname{atan}(dx + c)^3 b^3 + 3 \operatorname{atan}(dx + c) \right)}{2d}$$

input `int((d*e*x+c*e)*(a+b*atan(d*x+c))^3,x)`

output `(e*(atan(c + d*x)**3*b**3*c**2 + 2*atan(c + d*x)**3*b**3*c*d*x + atan(c + d*x)**3*b**3*d**2*x**2 + atan(c + d*x)**3*b**3 + 3*atan(c + d*x)**2*a*b**2*c**2 + 6*atan(c + d*x)**2*a*b**2*c*d*x + 3*atan(c + d*x)**2*a*b**2*d**2*x**2 + 3*atan(c + d*x)**2*a*b**2 - 3*atan(c + d*x)**2*b**3*d*x + 3*atan(c + d*x)*a**2*b*c**2 + 6*atan(c + d*x)*a**2*b*c*d*x + 3*atan(c + d*x)*a**2*b*d**2*x**2 + 3*atan(c + d*x)*a**2*b - 6*atan(c + d*x)*a*b**2*c - 6*atan(c + d*x)*a*b**2*d*x + 6*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**3*d**2 + 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b**2 + 2*a**3*c*d*x + a**3*d**2*x**2 - 3*a**2*b*d*x))/(2*d)`

3.25 $\int \frac{(a+b \arctan(c+dx))^3}{ce+dex} dx$

Optimal result	243
Mathematica [B] (verified)	244
Rubi [A] (verified)	245
Maple [C] (warning: unable to verify)	248
Fricas [F]	249
Sympy [F]	249
Maxima [F]	249
Giac [F]	250
Mupad [F(-1)]	250
Reduce [F]	250

Optimal result

Integrand size = 23, antiderivative size = 279

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx \\ &= \frac{2(a + b \arctan(c + dx))^3 \operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right)}{de} \\ & \quad - \frac{3ib(a + b \arctan(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{2de} \\ & \quad + \frac{3ib(a + b \arctan(c + dx))^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1+i(c+dx)}\right)}{2de} \\ & \quad - \frac{3b^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2de} \\ & \quad + \frac{3b^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1+i(c+dx)}\right)}{2de} \\ & \quad + \frac{3ib^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1+i(c+dx)}\right)}{4de} - \frac{3ib^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1+i(c+dx)}\right)}{4de} \end{aligned}$$

output

```
-2*(a+b*arctan(d*x+c))^3*arctanh(-1+2/(1+I*(d*x+c)))/d/e-3/2*I*b*(a+b*arctan(d*x+c))^2*polylog(2,1-2/(1+I*(d*x+c)))/d/e+3/2*I*b*(a+b*arctan(d*x+c))^2*polylog(2,-1+2/(1+I*(d*x+c)))/d/e-3/2*b^2*(a+b*arctan(d*x+c))*polylog(3,1-2/(1+I*(d*x+c)))/d/e+3/2*b^2*(a+b*arctan(d*x+c))*polylog(3,-1+2/(1+I*(d*x+c)))/d/e+3/4*I*b^3*polylog(4,1-2/(1+I*(d*x+c)))/d/e-3/4*I*b^3*polylog(4,-1+2/(1+I*(d*x+c)))/d/e
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 562 vs. $2(279) = 558$.

Time = 0.50 (sec) , antiderivative size = 562, normalized size of antiderivative = 2.01

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx$$

$$= \frac{64a^3 \log(c + dx) - 24ia^2b(\pi^2 - 4\pi \arctan(c + dx) + 8 \arctan(c + dx)^2 - i\pi \log(16) + 4i\pi \log(1 + e^{-2ia}))}{ce + dex}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x),x]
```

output

```
(64*a^3*Log[c + d*x] - (24*I)*a^2*b*(Pi^2 - 4*Pi*ArcTan[c + d*x] + 8*ArcTan[c + d*x]^2 - I*Pi*Log[16] + (4*I)*Pi*Log[1 + E^((-2*I)*ArcTan[c + d*x])]) - (8*I)*ArcTan[c + d*x]*Log[1 + E^((-2*I)*ArcTan[c + d*x])] + (8*I)*ArcTan[c + d*x]*Log[1 - E^((2*I)*ArcTan[c + d*x])] + (2*I)*Pi*Log[1 + c^2 + 2*c*d*x + d^2*x^2] + 4*PolyLog[2, -E^((-2*I)*ArcTan[c + d*x])] + 4*PolyLog[2, E^((2*I)*ArcTan[c + d*x])]) + 8*a*b^2*((-I)*Pi^3 + (16*I)*ArcTan[c + d*x]^3 + 24*ArcTan[c + d*x]^2*Log[1 - E^((-2*I)*ArcTan[c + d*x])] - 24*ArcTan[c + d*x]^2*Log[1 + E^((2*I)*ArcTan[c + d*x])] + (24*I)*ArcTan[c + d*x]*PolyLog[2, E^((-2*I)*ArcTan[c + d*x])] + (24*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])] + 12*PolyLog[3, E^((-2*I)*ArcTan[c + d*x])] - 12*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])]) - I*b^3*(Pi^4 - 32*ArcTan[c + d*x]^4 + (64*I)*ArcTan[c + d*x]^3*Log[1 - E^((-2*I)*ArcTan[c + d*x])] - (64*I)*ArcTan[c + d*x]^3*Log[1 + E^((2*I)*ArcTan[c + d*x])] - 96*ArcTan[c + d*x]^2*PolyLog[2, E^((-2*I)*ArcTan[c + d*x])] - 96*ArcTan[c + d*x]^2*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])] + (96*I)*ArcTan[c + d*x]*PolyLog[3, E^((-2*I)*ArcTan[c + d*x])] - (96*I)*ArcTan[c + d*x]*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])]) + 48*PolyLog[4, E^((-2*I)*ArcTan[c + d*x])] + 48*PolyLog[4, -E^((2*I)*ArcTan[c + d*x])])/(64*d*e)
```

Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {5566, 27, 5357, 5523, 5529, 5533, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx$$

$$\downarrow \text{5566}$$

$$\int \frac{(a + b \arctan(c + dx))^3}{e(c + dx)} d(c + dx)}{d}$$

$$\downarrow \text{27}$$

$$\int \frac{(a + b \arctan(c + dx))^3}{c + dx} d(c + dx)}{de}$$

↓ 5357

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a + b \operatorname{arctan}(c + dx))^3 - 6b \int \frac{(a+b \operatorname{arctan}(c+dx))^2 \operatorname{arctanh}\left(1 - \frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c + dx)}{de}$$

↓ 5523

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a + b \operatorname{arctan}(c + dx))^3 - 6b \left(\frac{1}{2} \int \frac{(a+b \operatorname{arctan}(c+dx))^2 \log\left(2 - \frac{2}{i(c+dx)+1}\right)}{(c+dx)^2+1} d(c + dx) - \frac{1}{2} \int \frac{(a+b \operatorname{arctan}(c+dx))^2}{(c+dx)^2+1} d(c + dx) \right)}{de}$$

↓ 5529

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a + b \operatorname{arctan}(c + dx))^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a + b \operatorname{arctan}(c + dx))^2 \right) \right)}{de}$$

↓ 5533

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a + b \operatorname{arctan}(c + dx))^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a + b \operatorname{arctan}(c + dx))^2 \right) \right)}{de}$$

↓ 7164

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1+i(c+dx)}\right) (a + b \operatorname{arctan}(c + dx))^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a + b \operatorname{arctan}(c + dx))^2 \right) \right)}{de}$$

input

```
Int[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x),x]
```

output

```
(2*(a + b*ArcTan[c + d*x])^3*ArcTanh[1 - 2/(1 + I*(c + d*x))] - 6*b*(((I/2)*
(a + b*ArcTan[c + d*x])^2*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] - I*b*(((I/2)*
(a + b*ArcTan[c + d*x])*PolyLog[3, 1 - 2/(1 + I*(c + d*x))] + (b*PolyLog
[4, 1 - 2/(1 + I*(c + d*x))])/4))/2 + ((-1/2*I)*(a + b*ArcTan[c + d*x])^2
*PolyLog[2, -1 + 2/(1 + I*(c + d*x))] + I*b*(((I/2)*(a + b*ArcTan[c + d*x])
*PolyLog[3, -1 + 2/(1 + I*(c + d*x))] + (b*PolyLog[4, -1 + 2/(1 + I*(c + d
*x))])/4))/2))/(d*e)
```

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 5357 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_/(x_), x_Symbol] := Simp[2*(a + b*ArcTan[c*x])^p*ArcTanh[1 - 2/(1 + I*c*x)], x] - Simp[2*b*c*p Int[(a + b*ArcTan[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 + I*c*x)]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[p, 1]`

rule 5523 `Int[(ArcTanh[u_]*)((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[u^2 - (1 - 2*(I/(I - c*x)))^2, 0]`

rule 5529 `Int[(Log[u_]*)((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*p*(I/2 Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]`

rule 5533 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_*PolyLog[k_, u_] / ((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[I*(a + b*ArcTan[c*x])^p*(PolyLog[k + 1, u]/(2*c*d)), x] - Simp[b*p*(I/2 Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[k + 1, u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[u^2 - (1 - 2*(I/(I - c*x)))^2, 0]`

rule 5566 `Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^p_*((e_.) + (f_.)*(x_))^m_), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.71 (sec) , antiderivative size = 2313, normalized size of antiderivative = 8.29

method	result	size
derivativedivides	Expression too large to display	2313
default	Expression too large to display	2313
parts	Expression too large to display	2321

input

```
int((a+b*arctan(d*x+c))^3/(d*e*x+c*e),x,method=_RETURNVERBOSE)
```

output

```
1/d*(a^3/e*ln(d*x+c)+b^3/e*(ln(d*x+c)*arctan(d*x+c)^3-arctan(d*x+c)^3*ln((
1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)+arctan(d*x+c)^3*ln(1+(1+I*(d*x+c))/(1+(d*x
+c)^2)^(1/2))-3*I*arctan(d*x+c)^2*polylog(2,-(1+I*(d*x+c))/(1+(d*x+c)^2)^(
1/2))+6*arctan(d*x+c)*polylog(3,-(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))+6*I*po
lylog(4,-(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))+arctan(d*x+c)^3*ln(1-(1+I*(d*x
+c))/(1+(d*x+c)^2)^(1/2))-3*I*arctan(d*x+c)^2*polylog(2,(1+I*(d*x+c))/(1+(
d*x+c)^2)^(1/2))+6*arctan(d*x+c)*polylog(3,(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/
2))+6*I*polylog(4,(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))+1/2*I*Pi*(csgn(I*((1+
I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*csgn(((1+
I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-csgn(((1+
I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2+csgn(I*
((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1))*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)
))*csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^
2)))-csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1))*csgn(I*((1+I*(d*x+c))^2/(1+
(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2-csgn(I/(1+(1+I*(d*x+c)
)^2/(1+(d*x+c)^2)))*csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c
))^2/(1+(d*x+c)^2)))^2+csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d
*x+c))^2/(1+(d*x+c)^2)))^3-csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+
I*(d*x+c))^2/(1+(d*x+c)^2)))*csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+
I*(d*x+c))^2/(1+(d*x+c)^2)))^2+csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(...
```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^3}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e),x, algorithm="fricas")`

output `integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3)/(d*e*x + c*e), x)`

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx$$

$$= \frac{\int \frac{a^3}{c+dx} dx + \int \frac{b^3 \operatorname{atan}^3(c+dx)}{c+dx} dx + \int \frac{3ab^2 \operatorname{atan}^2(c+dx)}{c+dx} dx + \int \frac{3a^2b \operatorname{atan}(c+dx)}{c+dx} dx}{e}$$

input `integrate((a+b*atan(d*x+c))**3/(d*e*x+c*e),x)`

output `(Integral(a**3/(c + d*x), x) + Integral(b**3*atan(c + d*x)**3/(c + d*x), x) + Integral(3*a*b**2*atan(c + d*x)**2/(c + d*x), x) + Integral(3*a**2*b*atan(c + d*x)/(c + d*x), x))/e`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^3}{dex + ce} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e),x, algorithm="maxima")`

output

```
a^3*log(d*e*x + c*e)/(d*e) + integrate(1/32*(28*b^3*arctan(d*x + c)^3 + 3*
b^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 96*a*b^2*arctan(d
*x + c)^2 + 96*a^2*b*arctan(d*x + c))/(d*e*x + c*e), x)
```

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx = \int \frac{(b \arctan(dx + c) + a)^3}{dex + ce} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e),x, algorithm="giac")
```

output

```
integrate((b*arctan(d*x + c) + a)^3/(d*e*x + c*e), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{ce + dex} dx$$

input

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x),x)
```

output

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x), x)
```

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{ce + dex} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atan}(dx+c)}{dx+c} dx \right) a^2 b d + \left(\int \frac{\operatorname{atan}(dx+c)^3}{dx+c} dx \right) b^3 d + 3 \left(\int \frac{\operatorname{atan}(dx+c)^2}{dx+c} dx \right) a b^2 d + \log(dx + c) a^3}{de}$$

input `int((a+b*atan(d*x+c))^3/(d*e*x+c*e),x)`

output `(3*int(atan(c + d*x)/(c + d*x),x)*a**2*b*d + int(atan(c + d*x)**3/(c + d*x),x)*b**3*d + 3*int(atan(c + d*x)**2/(c + d*x),x)*a*b**2*d + log(c + d*x)*a**3)/(d*e)`

3.26 $\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^2} dx$

Optimal result	252
Mathematica [A] (verified)	253
Rubi [A] (verified)	253
Maple [C] (warning: unable to verify)	256
Fricas [F]	257
Sympy [F]	257
Maxima [F]	257
Giac [F]	258
Mupad [F(-1)]	258
Reduce [F]	259

Optimal result

Integrand size = 23, antiderivative size = 163

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx \\ &= -\frac{i(a + b \arctan(c + dx))^3}{de^2} - \frac{(a + b \arctan(c + dx))^3}{de^2(c + dx)} \\ & \quad + \frac{3b(a + b \arctan(c + dx))^2 \log\left(2 - \frac{2}{1 - i(c + dx)}\right)}{de^2} \\ & \quad - \frac{3ib^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - i(c + dx)}\right)}{de^2} \\ & \quad + \frac{3b^3 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - i(c + dx)}\right)}{2de^2} \end{aligned}$$

output

```
-I*(a+b*arctan(d*x+c))^3/d/e^2-(a+b*arctan(d*x+c))^3/d/e^2/(d*x+c)+3*b*(a+b*arctan(d*x+c))^2*ln(2-2/(1-I*(d*x+c)))/d/e^2-3*I*b^2*(a+b*arctan(d*x+c))*polylog(2,-1+2/(1-I*(d*x+c)))/d/e^2+3/2*b^3*polylog(3,-1+2/(1-I*(d*x+c)))/d/e^2
```

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.61

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx$$

$$= -\frac{2a^3}{c+dx} - \frac{6a^2b \arctan(c+dx)}{c+dx} + 6a^2b \log(c + dx) - 3a^2b \log(1 + c^2 + 2cdx + d^2x^2) + 6ab^2 (\arctan(c + dx)) ((-$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^2,x]
```

output

```
((-2*a^3)/(c + d*x) - (6*a^2*b*ArcTan[c + d*x])/(c + d*x) + 6*a^2*b*Log[c + d*x] - 3*a^2*b*Log[1 + c^2 + 2*c*d*x + d^2*x^2] + 6*a*b^2*(ArcTan[c + d*x]*((-I - (c + d*x)^(-1))*ArcTan[c + d*x] + 2*Log[1 - E^((2*I)*ArcTan[c + d*x])]) - I*PolyLog[2, E^((2*I)*ArcTan[c + d*x])]) + 2*b^3*((-1/8*I)*Pi^3 + I*ArcTan[c + d*x]^3 - ArcTan[c + d*x]^3/(c + d*x) + 3*ArcTan[c + d*x]^2*Log[1 - E^((-2*I)*ArcTan[c + d*x])]) + (3*I)*ArcTan[c + d*x]*PolyLog[2, E^((-2*I)*ArcTan[c + d*x])]) + (3*PolyLog[3, E^((-2*I)*ArcTan[c + d*x])])]/(2))/ (2*d*e^2)
```

Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {5566, 27, 5361, 5459, 5403, 5527, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx$$

$$\downarrow \text{5566}$$

$$\int \frac{(a + b \arctan(c + dx))^3}{e^2(c + dx)^2} d(c + dx)$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \frac{\int \frac{(a+b \arctan(c+dx))^3}{(c+dx)^2} d(c+dx)}{de^2} \\
& \quad \downarrow \text{5361} \\
& \frac{3b \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^3}{c+dx}}{de^2} \\
& \quad \downarrow \text{5459} \\
& \frac{-\frac{(a+b \arctan(c+dx))^3}{c+dx} + 3b \left(i \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)(c+dx+i)} d(c+dx) - \frac{i(a+b \arctan(c+dx))^3}{3b} \right)}{de^2} \\
& \quad \downarrow \text{5403} \\
& \frac{-\frac{(a+b \arctan(c+dx))^3}{c+dx} + 3b \left(i \left(2ib \int \frac{(a+b \arctan(c+dx)) \log \left(2 - \frac{2}{1-i(c+dx)} \right)}{(c+dx)^2+1} d(c+dx) - i \log \left(2 - \frac{2}{1-i(c+dx)} \right) (a+b \arctan(c+dx)) \right) \right)}{de^2} \\
& \quad \downarrow \text{5527} \\
& \frac{-\frac{(a+b \arctan(c+dx))^3}{c+dx} + 3b \left(i \left(2ib \left(\frac{1}{2} i \operatorname{PolyLog} \left(2, \frac{2}{1-i(c+dx)} \right) - 1 \right) (a+b \arctan(c+dx)) - \frac{1}{2} ib \int \frac{\operatorname{PolyLog} \left(2, \frac{2}{1-i(c+dx)} \right)}{(c+dx)^2+1} \right) \right)}{de^2} \\
& \quad \downarrow \text{7164} \\
& \frac{-\frac{(a+b \arctan(c+dx))^3}{c+dx} + 3b \left(i \left(2ib \left(\frac{1}{2} i \operatorname{PolyLog} \left(2, \frac{2}{1-i(c+dx)} \right) - 1 \right) (a+b \arctan(c+dx)) - \frac{1}{4} b \operatorname{PolyLog} \left(3, \frac{2}{1-i(c+dx)} \right) \right) \right)}{de^2}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^2,x]`

output `((-(a + b*ArcTan[c + d*x])^3/(c + d*x)) + 3*b*(((1/3*I)*(a + b*ArcTan[c + d*x])^3)/b + I*((-I)*(a + b*ArcTan[c + d*x])^2*Log[2 - 2/(1 - I*(c + d*x))] + (2*I)*b*((I/2)*(a + b*ArcTan[c + d*x])*PolyLog[2, -1 + 2/(1 - I*(c + d*x))]) - (b*PolyLog[3, -1 + 2/(1 - I*(c + d*x))])/4)))/(d*e^2)`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 5361 $\text{Int}[((a_.) + \text{ArcTan}[(c_.)(x_)^(n_.)]*(b_.))^(p_.)(x_)^(m_.), x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}*((a + b*\text{ArcTan}[c*x^n])^p/(m+1)), x] - \text{Simp}[b*c*n*(p/(m+1)) \text{ Int}[x^{(m+n)}*((a + b*\text{ArcTan}[c*x^n])^{(p-1)/(1+c^2*x^{2n})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[p, 1] \ || \ (\text{EqQ}[n, 1] \ \&\& \ \text{IntegerQ}[m])) \ \&\& \ \text{NeQ}[m, -1]$
- rule 5403 $\text{Int}[((a_.) + \text{ArcTan}[(c_.)(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)(x_))), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^p*(\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \text{ Int}[(a + b*\text{ArcTan}[c*x])^{(p-1)}*(\text{Log}[2 - 2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2*d^2 + e^2, 0]$
- rule 5459 $\text{Int}[((a_.) + \text{ArcTan}[(c_.)(x_)]*(b_.))^(p_.)/((x_)*((d_) + (e_.)(x_)^2)), x_Symbol] \rightarrow \text{Simp}[(-I)*((a + b*\text{ArcTan}[c*x])^{(p+1)}/(b*d*(p+1))), x] + \text{Simp}[I/d \text{ Int}[(a + b*\text{ArcTan}[c*x])^p/(x*(I + c*x)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[e, c^2*d] \ \&\& \ \text{GtQ}[p, 0]$
- rule 5527 $\text{Int}[(\text{Log}[u_]*((a_.) + \text{ArcTan}[(c_.)(x_)]*(b_.))^(p_.))/((d_) + (e_.)(x_)^2), x_Symbol] \rightarrow \text{Simp}[I*(a + b*\text{ArcTan}[c*x])^p*(\text{PolyLog}[2, 1 - u]/(2*c*d)), x] - \text{Simp}[b*p*(I/2) \text{ Int}[(a + b*\text{ArcTan}[c*x])^{(p-1)}*(\text{PolyLog}[2, 1 - u]/(d + e*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2*d] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2*(I/(I + c*x)))^2, 0]$
- rule 5566 $\text{Int}[((a_.) + \text{ArcTan}[(c_) + (d_.)(x_)]*(b_.))^(p_.)*((e_.) + (f_.)(x_)^m), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e - c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.65 (sec) , antiderivative size = 2104, normalized size of antiderivative = 12.91

method	result	size
derivativedivides	Expression too large to display	2104
default	Expression too large to display	2104
parts	Expression too large to display	2112

input

```
int((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-a^3/e^2/(d*x+c)+b^3/e^2*(-1/(d*x+c)*arctan(d*x+c)^3+3*ln(d*x+c)*arct
an(d*x+c)^2-3/2*arctan(d*x+c)^2*ln(1+(d*x+c)^2)+3*arctan(d*x+c)^2*ln((1+I*
(d*x+c))/(1+(d*x+c)^2)^(1/2))-3*arctan(d*x+c)^2*ln((1+I*(d*x+c))^2/(1+(d*x
+c)^2)-1)-I*arctan(d*x+c)^3+3/4*(2*I*Pi*csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)
^2)-1))*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*csgn(I*((1+I*(d*x+c))^2/
(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-I*Pi*csgn(I/(1+(1+I*(d
*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1
+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)+I*Pi*csgn
(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+
(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)-2*I*Pi*csgn(I*((1+I*(d*x+c))^2/(1+(d*x
+c)^2)-1))*csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+
(d*x+c)^2)))^2-I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^
2/(1+(d*x+c)^2))^2)-2*I*Pi*csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+
I*(d*x+c))^2/(1+(d*x+c)^2)))^2+I*Pi*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2
)))^2*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)-2*I*Pi*csgn(I*(1+(1+I*(d
*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)-I*P
i*csgn(I*(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))^2*csgn(I*(1+I*(d*x+c))^2/(1+(d
*x+c)^2))+2*I*Pi*csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2
/(1+(d*x+c)^2)))^3+I*Pi*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I
*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)-2...
```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="fricas")`

output `integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3)/(d^2*e^2*x^2 + 2*c*d*e^2*x + c^2*e^2), x)`

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx$$

$$= \frac{\int \frac{a^3}{c^2+2cdx+d^2x^2} dx + \int \frac{b^3 \operatorname{atan}^3(c+dx)}{c^2+2cdx+d^2x^2} dx + \int \frac{3ab^2 \operatorname{atan}^2(c+dx)}{c^2+2cdx+d^2x^2} dx + \int \frac{3a^2b \operatorname{atan}(c+dx)}{c^2+2cdx+d^2x^2} dx}{e^2}$$

input `integrate((a+b*atan(d*x+c))**3/(d*e*x+c*e)**2,x)`

output `(Integral(a**3/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(b**3*atan(c + d*x)**3/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(3*a*b**2*atan(c + d*x)**2/(c**2 + 2*c*d*x + d**2*x**2), x) + Integral(3*a**2*b*atan(c + d*x)/(c**2 + 2*c*d*x + d**2*x**2), x))/e**2`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="maxima")`

output

```
-3/2*(d*(log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2) - 2*log(d*x + c)/(d^2*
e^2)) + 2*arctan(d*x + c)/(d^2*e^2*x + c*d*e^2))*a^2*b - a^3/(d^2*e^2*x +
c*d*e^2) - 1/32*(4*b^3*arctan(d*x + c)^3 - 3*b^3*arctan(d*x + c)*log(d^2*x
^2 + 2*c*d*x + c^2 + 1)^2 - 32*(d^2*e^2*x + c*d*e^2)*integrate(1/32*(28*(b
^3*d^2*x^2 + 2*b^3*c*d*x + b^3*c^2 + b^3)*arctan(d*x + c)^3 + 12*(8*a*b^2*
d^2*x^2 + 8*a*b^2*c^2 + b^3*c + 8*a*b^2 + (16*a*b^2*c + b^3)*d*x)*arctan(d
*x + c)^2 - 12*(b^3*d^2*x^2 + 2*b^3*c*d*x + b^3*c^2)*arctan(d*x + c)*log(d
^2*x^2 + 2*c*d*x + c^2 + 1) - 3*(b^3*d*x + b^3*c - (b^3*d^2*x^2 + 2*b^3*c*
d*x + b^3*c^2 + b^3)*arctan(d*x + c))*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2)/
(d^4*e^2*x^4 + 4*c*d^3*e^2*x^3 + (6*c^2 + 1)*d^2*e^2*x^2 + 2*(2*c^3 + c)*d
*e^2*x + (c^4 + c^2)*e^2), x))/(d^2*e^2*x + c*d*e^2)
```

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^2} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^2,x, algorithm="giac")
```

output

```
integrate((b*arctan(d*x + c) + a)^3/(d*e*x + c*e)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{(ce + dex)^2} dx$$

input

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^2,x)
```

output

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^2, x)
```

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^2} dx = \text{Too large to display}$$

input `int((a+b*atan(d*x+c))^3/(d*e*x+c*e)^2,x)`

output `(- atan(c + d*x)**3*b**3*c**3 - atan(c + d*x)**3*b**3*c**2*d*x - atan(c + d*x)**3*b**3*c + atan(c + d*x)**3*b**3*d*x - 6*atan(c + d*x)**2*a*b**2*c**3 - 6*atan(c + d*x)**2*a*b**2*c**2*d*x - 6*atan(c + d*x)**2*a*b**2*c - 3*atan(c + d*x)**2*b**3*c**4 - 3*atan(c + d*x)**2*b**3*c**3*d*x - 3*atan(c + d*x)**2*b**3*c**2 - 6*atan(c + d*x)*a**2*b*c - 12*atan(c + d*x)*a*b**2*c**2 - 6*atan(c + d*x)*b**3*c**3 + 12*int((atan(c + d*x)*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*a*b**2*c**2*d**2 + 12*int((atan(c + d*x)*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*a*b**2*c*d**3*x + 6*int((atan(c + d*x)*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*b**3*c**3*d**2 + 6*int((atan(c + d*x)*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*b**3*c**2*d**3*x - 3*int((atan(c + d*x)**2*x**2)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*b**3*c*d**3 - 3*int((atan(c + d*x)**2*x**2)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + c**2 + 4*c*d**3*x**3 + 2*c*d*x + d**4*x**4 + d**2*x**2),x)*b**3*d**4*x - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a**2*b*c**2 - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a**2*b*c*d*x - 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b**2*c**3 - 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b**2*c**2*d*x - 3*log(c...`

3.27 $\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^3} dx$

Optimal result	260
Mathematica [A] (verified)	261
Rubi [A] (verified)	261
Maple [B] (verified)	264
Fricas [F]	265
Sympy [F]	265
Maxima [F]	266
Giac [F]	267
Mupad [F(-1)]	267
Reduce [F]	267

Optimal result

Integrand size = 23, antiderivative size = 180

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = -\frac{3ib(a + b \arctan(c + dx))^2}{2de^3} - \frac{3b(a + b \arctan(c + dx))^2}{2de^3(c + dx)} - \frac{(a + b \arctan(c + dx))^3}{2de^3} - \frac{(a + b \arctan(c + dx))^3}{2de^3(c + dx)^2} + \frac{3b^2(a + b \arctan(c + dx)) \log\left(2 - \frac{2}{1-i(c+dx)}\right)}{de^3} - \frac{3ib^3 \text{PolyLog}\left(2, -1 + \frac{2}{1-i(c+dx)}\right)}{2de^3}$$

output

```
-3/2*I*b*(a+b*arctan(d*x+c))^2/d/e^3-3/2*b*(a+b*arctan(d*x+c))^2/d/e^3/(d*x+c)-1/2*(a+b*arctan(d*x+c))^3/d/e^3-1/2*(a+b*arctan(d*x+c))^3/d/e^3/(d*x+c)^2+3*b^2*(a+b*arctan(d*x+c))*ln(2-2/(1-I*(d*x+c)))/d/e^3-3/2*I*b^3*polyl
og(2,-1+2/(1-I*(d*x+c)))/d/e^3
```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.25

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \frac{a^3 + b^3(1 + c^2 + 2cdx + d^2x^2) \arctan(c + dx)^3 + 3a^2b(c + dx + (1 + (c + dx)^2) \arctan(c + dx)) + 3ab^2(1 + (c + dx)^2) \arctan(c + dx)^2 + b^3(1 + (c + dx)^2) \arctan(c + dx)}{(ce + dex)^3}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^3,x]
```

output

```
-1/2*(a^3 + b^3*(1 + c^2 + 2*c*d*x + d^2*x^2)*ArcTan[c + d*x]^3 + 3*a^2*b*(c + d*x + (1 + (c + d*x)^2)*ArcTan[c + d*x]) + 3*a*b^2*(2*(c + d*x)*ArcTan[c + d*x] + (1 + (c + d*x)^2)*ArcTan[c + d*x]^2 - 2*(c + d*x)^2*Log[(c + d*x)/Sqrt[1 + (c + d*x)^2]]) + 3*b^3*(c + d*x)*(ArcTan[c + d*x]^2 - 2*(c + d*x)*ArcTan[c + d*x]*Log[1 - E^((2*I)*ArcTan[c + d*x])]) + I*(c + d*x)*(ArcTan[c + d*x]^2 + PolyLog[2, E^((2*I)*ArcTan[c + d*x])]))/(d*e^3*(c + d*x)^2)
```

Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$, Rules used = {5566, 27, 5361, 5453, 5361, 5419, 5459, 5403, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx$$

↓ 5566

$$\int \frac{(a + b \arctan(c + dx))^3}{e^3(c + dx)^3} d(c + dx)$$

↓ 27

$$\frac{\int \frac{(a+b \arctan(c+dx))^3}{(c+dx)^3} d(c+dx)}{de^3}$$

↓ 5361

$$\frac{\frac{3}{2}b \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)^2((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 5453

$$\frac{\frac{3}{2}b \left(\int \frac{(a+b \arctan(c+dx))^2}{(c+dx)^2} d(c+dx) - \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c+dx) \right) - \frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 5361

$$\frac{\frac{3}{2}b \left(2b \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) - \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)^2+1} d(c+dx) - \frac{(a+b \arctan(c+dx))^2}{c+dx} \right) - \frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 5419

$$\frac{\frac{3}{2}b \left(2b \int \frac{a+b \arctan(c+dx)}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^3}{3b} - \frac{(a+b \arctan(c+dx))^2}{c+dx} \right) - \frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2}}{de^3}$$

↓ 5459

$$\frac{-\frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2} + \frac{3}{2}b \left(2b \left(i \int \frac{a+b \arctan(c+dx)}{(c+dx)(c+dx+i)} d(c+dx) - \frac{i(a+b \arctan(c+dx))^2}{2b} \right) - \frac{(a+b \arctan(c+dx))^3}{3b} - \frac{(a+b \arctan(c+dx))^2}{c+dx} \right)}{de^3}$$

↓ 5403

$$\frac{-\frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2} + \frac{3}{2}b \left(2b \left(i \left(ib \int \frac{\log\left(2 - \frac{2}{1-i(c+dx)}\right)}{(c+dx)^2+1} d(c+dx) - i \log\left(2 - \frac{2}{1-i(c+dx)}\right) (a+b \arctan(c+dx)) \right) \right)}{de^3}$$

↓ 2897

$$\frac{-\frac{(a+b \arctan(c+dx))^3}{2(c+dx)^2} + \frac{3}{2}b \left(2b \left(i \left(-i \log\left(2 - \frac{2}{1-i(c+dx)}\right) (a+b \arctan(c+dx)) - \frac{1}{2}b \text{PolyLog}\left(2, \frac{2}{1-i(c+dx)} - 1\right) \right) \right)}{de^3}$$

input

`Int[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^3,x]`

output

$$\begin{aligned} & (-1/2*(a + b*\text{ArcTan}[c + d*x])^3/(c + d*x)^2 + (3*b*(-((a + b*\text{ArcTan}[c + d*x])^2/(c + d*x)) - (a + b*\text{ArcTan}[c + d*x])^3/(3*b) + 2*b*((-1/2*I)*(a + b*\text{ArcTan}[c + d*x])^2)/b + I*(-I)*(a + b*\text{ArcTan}[c + d*x])*Log[2 - 2/(1 - I*(c + d*x))] - (b*\text{PolyLog}[2, -1 + 2/(1 - I*(c + d*x))])/2)))/2)/(d*e^3) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$$

rule 2897

$$\text{Int}[\text{Log}[u_]*(P_q)^{(m_.)}, x_Symbol] \rightarrow \text{With}[\{C = \text{FullSimplify}[P_q^m*((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \&\& \text{PolyQ}[P_q, x] \&\& \text{RationalFunctionQ}[u, x] \&\& \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[P_q, x]]$$

rule 5361

$$\text{Int}[(a_. + \text{ArcTan}[c_*(x_)^{(n_.)}]*(b_.))^{(p_.)}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTan}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \quad \text{Int}[x^{(m + n)}*((a + b*\text{ArcTan}[c*x^n])^{(p - 1)/(1 + c^2*x^{(2*n)})}), x], x] /; \text{FreeQ}[\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \& \& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$$

rule 5403

$$\text{Int}[(a_. + \text{ArcTan}[c_*(x_)]*(b_.))^{(p_.)}/((x_)*((d_) + (e_.)*(x_))), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^p*(\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \quad \text{Int}[(a + b*\text{ArcTan}[c*x])^{(p - 1)}*(\text{Log}[2 - 2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2*d^2 + e^2, 0]$$

rule 5419

$$\text{Int}[(a_. + \text{ArcTan}[c_*(x_)]*(b_.))^{(p_.)}/((d_) + (e_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^{(p + 1)}/(b*c*d*(p + 1)), x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \&\& \text{EqQ}[e, c^2*d] \&\& \text{NeQ}[p, -1]$$


```
rule 5453 Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))^(p_.)*((f_.)*(x_.))^(m_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/d Int[(f*x)^m*(a + b*ArcTan[c*x])^p, x], x] - Simp[e/(d*f^2) Int[(f*x)^(m + 2)*(a + b*ArcTan[c*x])^p/(d + e*x^2), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && LtQ[m, -1]
```

```
rule 5459 Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))^(p_.)/((x_.)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[(-1)*((a + b*ArcTan[c*x])^(p + 1)/(b*d*(p + 1))), x] + Simp[I/d Int[(a + b*ArcTan[c*x])^p/(x*(I + c*x)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && GtQ[p, 0]
```

```
rule 5566 Int[((a_.) + ArcTan[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 422 vs. 2(166) = 332.

Time = 1.52 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.35

method	result
derivativedivides	$-\frac{a^3}{2e^3(dx+c)^2} + \frac{b^3 \left(-\frac{\arctan(dx+c)^3}{2(dx+c)^2} - \frac{3\arctan(dx+c)^2}{2(dx+c)} - \frac{\arctan(dx+c)^3}{2} + 3\ln(dx+c)\arctan(dx+c) - \frac{3\arctan(dx+c)\ln(1+(dx+c))}{2} \right)}{2e^3(dx+c)^2}$
default	$-\frac{a^3}{2e^3(dx+c)^2} + \frac{b^3 \left(-\frac{\arctan(dx+c)^3}{2(dx+c)^2} - \frac{3\arctan(dx+c)^2}{2(dx+c)} - \frac{\arctan(dx+c)^3}{2} + 3\ln(dx+c)\arctan(dx+c) - \frac{3\arctan(dx+c)\ln(1+(dx+c))}{2} \right)}{2e^3(dx+c)^2}$
parts	$-\frac{a^3}{2e^3(dx+c)^2} + \frac{b^3 \left(-\frac{\arctan(dx+c)^3}{2(dx+c)^2} - \frac{3\arctan(dx+c)^2}{2(dx+c)} - \frac{\arctan(dx+c)^3}{2} + 3\ln(dx+c)\arctan(dx+c) - \frac{3\arctan(dx+c)\ln(1+(dx+c))}{2} \right)}{2e^3(dx+c)^2}$

```
input int((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-1/2*a^3/e^3/(d*x+c)^2+b^3/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)^3-3/2/(d*x+c)*arctan(d*x+c)^2-1/2*arctan(d*x+c)^3+3*ln(d*x+c)*arctan(d*x+c)-3/2*arctan(d*x+c)*ln(1+(d*x+c)^2)-3/4*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))+3/4*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))+3/2*I*ln(d*x+c)*ln(1+I*(d*x+c))-3/2*I*ln(d*x+c)*ln(1-I*(d*x+c))+3/2*I*dilog(1+I*(d*x+c))-3/2*I*dilog(1-I*(d*x+c)))+3*a*b^2/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)^2-1/(d*x+c)*arctan(d*x+c)-1/2*arctan(d*x+c)^2+ln(d*x+c)-1/2*ln(1+(d*x+c)^2))+3*a^2*b/e^3*(-1/2/(d*x+c)^2*arctan(d*x+c)-1/2/(d*x+c)-1/2*arctan(d*x+c))
```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^3} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="fricas")
```

output

```
integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3)/(d^3*e^3*x^3 + 3*c*d^2*e^3*x^2 + 3*c^2*d*e^3*x + c^3*e^3), x)
```

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \frac{\int \frac{a^3}{c^3+3c^2dx+3cd^2x^2+d^3x^3} dx + \int \frac{b^3 \operatorname{atan}^3(c+dx)}{c^3+3c^2dx+3cd^2x^2+d^3x^3} dx + \int \frac{3ab^2 \operatorname{atan}^2(c+dx)}{c^3+3c^2dx+3cd^2x^2+d^3x^3} dx + \int \frac{3a^2b \operatorname{atan}(c+dx)}{c^3+3c^2dx+3cd^2x^2+d^3x^3} dx}{e^3}$$

input

```
integrate((a+b*atan(d*x+c))**3/(d*e*x+c*e)**3,x)
```

output

```
(Integral(a**3/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(b**3*atan(c + d*x)**3/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(3*a*b**2*atan(c + d*x)**2/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x) + Integral(3*a**2*b*atan(c + d*x)/(c**3 + 3*c**2*d*x + 3*c*d**2*x**2 + d**3*x**3), x))/e**3
```

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^3} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="maxima")
```

output

```
-3/2*(d*(1/(d^3*e^3*x + c*d^2*e^3) + arctan((d^2*x + c*d)/d)/(d^2*e^3)) + arctan(d*x + c)/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3))*a^2*b - 3/2*(2*d*(1/(d^3*e^3*x + c*d^2*e^3) + arctan((d^2*x + c*d)/d)/(d^2*e^3))*arctan(d*x + c) - (arctan(d*x + c)^2 - log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 2*log(d*x + c))/(d*e^3))*a*b^2 - 3/2*a*b^2*arctan(d*x + c)^2/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3) - 1/32*(8*(d^2*x^2 + 2*c*d*x + c^2 + 1)*arctan(d*x + c)^3 + 12*(d*x + c)*arctan(d*x + c)^2 - 3*(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 - 32*(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)*integrate(1/32*(16*(d^2*x^2 + 2*c*d*x + c^2 + 1)*arctan(d*x + c)^3 + 12*(d^3*x^3 + 3*c*d^2*x^2 + c^3 + (3*c^2 + 1)*d*x + c)*arctan(d*x + c)^2 + 3*(d^3*x^3 + 3*c*d^2*x^2 + c^3 + (3*c^2 + 1)*d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 24*(d^2*x^2 + 2*c*d*x + c^2)*arctan(d*x + c) - 12*(d^3*x^3 + 3*c*d^2*x^2 + 3*c^2*d*x + c^3)*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/(d^5*e^3*x^5 + 5*c*d^4*e^3*x^4 + (10*c^2 + 1)*d^3*e^3*x^3 + (10*c^3 + 3*c)*d^2*e^3*x^2 + (5*c^4 + 3*c^2)*d*e^3*x + (c^5 + c^3)*e^3), x))*b^3/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3) - 1/2*a^3/(d^3*e^3*x^2 + 2*c*d^2*e^3*x + c^2*d*e^3)
```

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^3} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^3,x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^3/(d*e*x + c*e)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{(ce + dex)^3} dx$$

input `int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^3,x)`

output `int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^3, x)`

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^3} dx = \text{Too large to display}$$

input `int((a+b*atan(d*x+c))^3/(d*e*x+c*e)^3,x)`

output

```
( - 2*atan(c + d*x)**3*b**3*c**3 - 4*atan(c + d*x)**3*b**3*c**2*d*x - 2*at
an(c + d*x)**3*b**3*c*d**2*x**2 - 2*atan(c + d*x)**3*b**3*c - 6*atan(c + d
*x)**2*a*b**2*c**3 - 12*atan(c + d*x)**2*a*b**2*c**2*d*x - 6*atan(c + d*x)
**2*a*b**2*c*d**2*x**2 - 6*atan(c + d*x)**2*a*b**2*c + 6*atan(c + d*x)**2*
b**3*c*d*x + 6*atan(c + d*x)**2*b**3*d**2*x**2 - 6*atan(c + d*x)*a**2*b*c*
*3 - 12*atan(c + d*x)*a**2*b*c**2*d*x - 6*atan(c + d*x)*a**2*b*c*d**2*x**2
- 6*atan(c + d*x)*a**2*b*c - 12*atan(c + d*x)*a*b**2*c**2 - 12*atan(c + d
*x)*a*b**2*c*d*x - 6*atan(c + d*x)*b**3*c**3 - 12*atan(c + d*x)*b**3*c**2*
d*x - 6*atan(c + d*x)*b**3*c*d**2*x**2 + 6*atan(c + d*x)*b**3*c + 12*atan(
c + d*x)*b**3*d*x + 12*int((atan(c + d*x)*x)/(c**5 + 5*c**4*d*x + 10*c**3*
d**2*x**2 + c**3 + 10*c**2*d**3*x**3 + 3*c**2*d*x + 5*c*d**4*x**4 + 3*c*d*
*2*x**2 + d**5*x**5 + d**3*x**3),x)*b**3*c**2*d**2 + 24*int((atan(c + d*x)
*x)/(c**5 + 5*c**4*d*x + 10*c**3*d**2*x**2 + c**3 + 10*c**2*d**3*x**3 + 3*
c**2*d*x + 5*c*d**4*x**4 + 3*c*d**2*x**2 + d**5*x**5 + d**3*x**3),x)*b**3*
c*d**3*x + 12*int((atan(c + d*x)*x)/(c**5 + 5*c**4*d*x + 10*c**3*d**2*x**2
+ c**3 + 10*c**2*d**3*x**3 + 3*c**2*d*x + 5*c*d**4*x**4 + 3*c*d**2*x**2 +
d**5*x**5 + d**3*x**3),x)*b**3*d**4*x**2 - 6*log(c**2 + 2*c*d*x + d**2*x*
*2 + 1)*a*b**2*c**3 - 12*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b**2*c**2*d
*x - 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b**2*c*d**2*x**2 + 6*log(c**2
+ 2*c*d*x + d**2*x**2 + 1)*b**3*c**2 + 12*log(c**2 + 2*c*d*x + d**2*x*...
```

3.28 $\int \frac{(a+b \arctan(c+dx))^3}{(ce+dex)^4} dx$

Optimal result	269
Mathematica [A] (verified)	270
Rubi [A] (verified)	271
Maple [C] (warning: unable to verify)	275
Fricas [F]	276
Sympy [F]	277
Maxima [F]	277
Giac [F]	278
Mupad [F(-1)]	278
Reduce [F]	279

Optimal result

Integrand size = 23, antiderivative size = 287

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = -\frac{b^2(a + b \arctan(c + dx))}{de^4(c + dx)} - \frac{b(a + b \arctan(c + dx))^2}{2de^4}$$

$$- \frac{b(a + b \arctan(c + dx))^2}{2de^4(c + dx)^2} + \frac{i(a + b \arctan(c + dx))^3}{3de^4}$$

$$- \frac{(a + b \arctan(c + dx))^3}{3de^4(c + dx)^3}$$

$$+ \frac{b^3 \log(c + dx)}{de^4} - \frac{b^3 \log(1 + (c + dx)^2)}{2de^4}$$

$$- \frac{b(a + b \arctan(c + dx))^2 \log\left(2 - \frac{2}{1 - i(c + dx)}\right)}{de^4}$$

$$+ \frac{ib^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - i(c + dx)}\right)}{de^4}$$

$$- \frac{b^3 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - i(c + dx)}\right)}{2de^4}$$

output

$$-b^2(a+b\arctan(dx+c))/d/e^4/(dx+c)-1/2*b*(a+b\arctan(dx+c))^2/d/e^4-1/2*b*(a+b\arctan(dx+c))^2/d/e^4/(dx+c)^2+1/3*I*(a+b\arctan(dx+c))^3/d/e^4-1/3*(a+b\arctan(dx+c))^3/d/e^4/(dx+c)^3+b^3*\ln(dx+c)/d/e^4-1/2*b^3*\ln(1+(dx+c)^2)/d/e^4-b*(a+b\arctan(dx+c))^2*\ln(2-2/(1-I*(dx+c)))/d/e^4+I*b^2*(a+b\arctan(dx+c))*\text{polylog}(2,-1+2/(1-I*(dx+c)))/d/e^4-1/2*b^3*\text{polylog}(3,-1+2/(1-I*(dx+c)))/d/e^4$$
Mathematica [A] (verified)

Time = 0.96 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.25

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx$$

$$= -\frac{8a^3}{(c+dx)^3} - \frac{12a^2b}{(c+dx)^2} - \frac{24a^2b \arctan(c+dx)}{(c+dx)^3} - 24a^2b \log(c + dx) + 12a^2b \log(1 + c^2 + 2cdx + d^2x^2) + 24ab^2 \left(- \right)$$

input

`Integrate[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^4,x]`

output

$$\left(\frac{(-8a^3)/(c + dx)^3 - (12a^2b)/(c + dx)^2 - (24a^2b*ArcTan[c + dx])/(c + dx)^3 - 24a^2b*Log[c + dx] + 12a^2b*Log[1 + c^2 + 2*c*d*x + d^2*x^2] + 24a*b^2*(-(((c + dx)^2 + ArcTan[c + dx]^2)/(c + dx)^3) + ArcTan[c + dx]*(-1 - (c + dx)^{-2}) + I*ArcTan[c + dx] - 2*Log[1 - E^{(2*I)*ArcTan[c + dx]}]) + I*PolyLog[2, E^{(2*I)*ArcTan[c + dx]}]) + b^3*(I*Pi^3 - (24*ArcTan[c + dx])/(c + dx) - 12*ArcTan[c + dx]^2 - (12*ArcTan[c + dx]^2)/(c + dx)^2 - (8*I)*ArcTan[c + dx]^3 - (8*ArcTan[c + dx]^3)/(c + dx)^3 - 24*ArcTan[c + dx]^2*Log[1 - E^{(-2*I)*ArcTan[c + dx]}] + 24*Log[(c + dx)/Sqrt[1 + (c + dx)^2]] - (24*I)*ArcTan[c + dx]*PolyLog[2, E^{(-2*I)*ArcTan[c + dx]}] - 12*PolyLog[3, E^{(-2*I)*ArcTan[c + dx]}]) \right) / (24*d*e^4)$$

Rubi [A] (verified)

Time = 1.70 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.87, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.696$, Rules used = {5566, 27, 5361, 5453, 5361, 5453, 5361, 243, 47, 14, 16, 5419, 5459, 5403, 5527, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx \\
 & \quad \downarrow \text{5566} \\
 & \int \frac{(a + b \arctan(c + dx))^3}{e^4(c + dx)^4} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(a + b \arctan(c + dx))^3}{(c + dx)^4} d(c + dx) \\
 & \quad \downarrow \text{5361} \\
 & \frac{b \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^3((c + dx)^2 + 1)} d(c + dx) - \frac{(a + b \arctan(c + dx))^3}{3(c + dx)^3}}{de^4} \\
 & \quad \downarrow \text{5453} \\
 & \frac{b \left(\int \frac{(a + b \arctan(c + dx))^2}{(c + dx)^3} d(c + dx) - \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)((c + dx)^2 + 1)} d(c + dx) \right) - \frac{(a + b \arctan(c + dx))^3}{3(c + dx)^3}}{de^4} \\
 & \quad \downarrow \text{5361} \\
 & \frac{b \left(b \int \frac{a + b \arctan(c + dx)}{(c + dx)^2((c + dx)^2 + 1)} d(c + dx) - \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)((c + dx)^2 + 1)} d(c + dx) - \frac{(a + b \arctan(c + dx))^2}{2(c + dx)^2} \right) - \frac{(a + b \arctan(c + dx))^3}{3(c + dx)^3}}{de^4} \\
 & \quad \downarrow \text{5453} \\
 & \frac{b \left(b \left(\int \frac{a + b \arctan(c + dx)}{(c + dx)^2} d(c + dx) - \int \frac{a + b \arctan(c + dx)}{(c + dx)^2 + 1} d(c + dx) \right) - \int \frac{(a + b \arctan(c + dx))^2}{(c + dx)((c + dx)^2 + 1)} d(c + dx) - \frac{(a + b \arctan(c + dx))^2}{2(c + dx)^2} \right) - \frac{(a + b \arctan(c + dx))^3}{3(c + dx)^3}}{de^4} \\
 & \quad \downarrow \text{5361}
 \end{aligned}$$

$$\frac{b\left(b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + b \int \frac{1}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx)\right)}{de^4}$$

↓ 243

$$\frac{b\left(b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b \int \frac{1}{(c+dx)^2((c+dx)^2+1)} d(c+dx)^2 - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx)\right)}{de^4}$$

↓ 47

$$\frac{b\left(b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b\left(\int \frac{1}{(c+dx)^2} d(c+dx)^2 - \int \frac{1}{(c+dx)^2+1} d(c+dx)^2\right) - \frac{a+b \arctan(c+dx)}{c+dx}\right) - \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx)\right)}{de^4}$$

↓ 14

$$\frac{b\left(-\int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx) + b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) + \frac{1}{2}b\left(\log((c+dx)^2) - \int \frac{1}{(c+dx)^2+1} d(c+dx)\right)\right)\right)}{de^4}$$

↓ 16

$$\frac{b\left(-\int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx) + b\left(-\int \frac{a+b \arctan(c+dx)}{(c+dx)^2+1} d(c+dx) - \frac{a+b \arctan(c+dx)}{c+dx} + \frac{1}{2}b(\log((c+dx)^2) - \log((c+dx)^2))\right)\right)}{de^4}$$

↓ 5419

$$\frac{b\left(-\int \frac{(a+b \arctan(c+dx))^2}{(c+dx)((c+dx)^2+1)} d(c+dx) - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2} + b\left(-\frac{(a+b \arctan(c+dx))^2}{2b} - \frac{a+b \arctan(c+dx)}{c+dx} + \frac{1}{2}b(\log((c+dx)^2) - \log((c+dx)^2))\right)\right)}{de^4}$$

↓ 5459

$$\frac{-\frac{(a+b \arctan(c+dx))^3}{3(c+dx)^3} + b\left(-i \int \frac{(a+b \arctan(c+dx))^2}{(c+dx)(c+dx+i)} d(c+dx) + \frac{i(a+b \arctan(c+dx))^3}{3b} - \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2} + b\left(-\frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2} + \frac{(a+b \arctan(c+dx))^2}{2(c+dx)^2}\right)\right)}{de^4}$$

↓ 5403

$$\frac{-\frac{(a+b \arctan(c+dx))^3}{3(c+dx)^3} + b\left(-i\left(2ib \int \frac{(a+b \arctan(c+dx)) \log\left(2 - \frac{2}{1-i(c+dx)}\right)}{(c+dx)^2+1} d(c+dx) - i \log\left(2 - \frac{2}{1-i(c+dx)}\right)\right)(a+b \arctan(c+dx))\right)}{de^4}$$

↓ 5527

$$-\frac{(a+b\arctan(c+dx))^3}{3(c+dx)^3} + b\left(-i\left(2ib\left(\frac{1}{2}i\operatorname{PolyLog}\left(2, \frac{2}{1-i(c+dx)} - 1\right)(a+b\arctan(c+dx)) - \frac{1}{2}ib\int \frac{\operatorname{PolyLog}\left(2, \frac{2}{1-i(c+dx)}\right)}{(c+dx)^2+1}\right.\right.\right.$$

↓ 7164

$$-\frac{(a+b\arctan(c+dx))^3}{3(c+dx)^3} + b\left(-i\left(2ib\left(\frac{1}{2}i\operatorname{PolyLog}\left(2, \frac{2}{1-i(c+dx)} - 1\right)(a+b\arctan(c+dx)) - \frac{1}{4}b\operatorname{PolyLog}\left(3, \frac{2}{1-i(c+dx)}\right)\right.\right.\right.$$

input

```
Int[(a + b*ArcTan[c + d*x])^3/(c*e + d*e*x)^4,x]
```

output

```
(-1/3*(a + b*ArcTan[c + d*x])^3/(c + d*x)^3 + b*(-1/2*(a + b*ArcTan[c + d*x])^2/(c + d*x)^2 + ((I/3)*(a + b*ArcTan[c + d*x])^3)/b + b*(-((a + b*ArcTan[c + d*x])/(c + d*x)) - (a + b*ArcTan[c + d*x])^2/(2*b) + (b*(Log[(c + d*x)^2] - Log[1 + (c + d*x)^2]))/2) - I*((-I)*(a + b*ArcTan[c + d*x])^2*Log[2 - 2/(1 - I*(c + d*x))] + (2*I)*b*((I/2)*(a + b*ArcTan[c + d*x])*PolyLog[2, -1 + 2/(1 - I*(c + d*x))] - (b*PolyLog[3, -1 + 2/(1 - I*(c + d*x))])))/(d*e^4)
```

Defintions of rubi rules used

rule 14

```
Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]
```

rule 16

```
Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 47

```
Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]
```

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m - 1)/2]$

rule 5361 $\text{Int}[(a_. + \text{ArcTan}[c_.*(x_)^{(n_.)}]*(b_.))^{(p_.)}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m + 1)}*((a + b*\text{ArcTan}[c*x^n])^p/(m + 1)), x] - \text{Simp}[b*c*n*(p/(m + 1)) \text{ Int}[x^{(m + n)}*((a + b*\text{ArcTan}[c*x^n])^{(p - 1)/(1 + c^2*x^{2*n})}), x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[p, 1] \|\| (\text{EqQ}[n, 1] \& \& \text{IntegerQ}[m])) \&\& \text{NeQ}[m, -1]$

rule 5403 $\text{Int}[(a_. + \text{ArcTan}[c_.*(x_)]*(b_.))^{(p_.)}/((x_)*((d_) + (e_.)*(x_))), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^p*(\text{Log}[2 - 2/(1 + e*(x/d))]/d), x] - \text{Simp}[b*c*(p/d) \text{ Int}[(a + b*\text{ArcTan}[c*x])^{(p - 1)}*(\text{Log}[2 - 2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{EqQ}[c^2*d^2 + e^2, 0]$

rule 5419 $\text{Int}[(a_. + \text{ArcTan}[c_.*(x_)]*(b_.))^{(p_.)}/((d_) + (e_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(a + b*\text{ArcTan}[c*x])^{(p + 1)}/(b*c*d*(p + 1)), x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x] \&\& \text{EqQ}[e, c^2*d] \&\& \text{NeQ}[p, -1]$

rule 5453 $\text{Int}[(a_. + \text{ArcTan}[c_.*(x_)]*(b_.))^{(p_.)}*((f_.)*(x_)^{(m_.)})/((d_) + (e_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Int}[(f*x)^m*(a + b*\text{ArcTan}[c*x])^p, x], x] - \text{Simp}[e/(d*f^2) \text{ Int}[(f*x)^{(m + 2)}*((a + b*\text{ArcTan}[c*x])^p/(d + e*x^2)), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{GtQ}[p, 0] \&\& \text{LtQ}[m, -1]$

rule 5459 $\text{Int}[(a_. + \text{ArcTan}[c_.*(x_)]*(b_.))^{(p_.)}/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] \rightarrow \text{Simp}[(-I)*((a + b*\text{ArcTan}[c*x])^{(p + 1)}/(b*d*(p + 1))), x] + \text{Simp}[I/d \text{ Int}[(a + b*\text{ArcTan}[c*x])^p/(x*(I + c*x)), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[e, c^2*d] \&\& \text{GtQ}[p, 0]$

rule 5527

```
Int[(Log[u_]*((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2
), x_Symbol] := Simp[I*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x
] - Simp[b*p*(I/2) Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d
+ e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*
d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I + c*x)))^2, 0]
```

rule 5566

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_)^(m
_.), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x],
x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] &&
IGtQ[p, 0]
```

rule 7164

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v,
x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.86 (sec) , antiderivative size = 2446, normalized size of antiderivative = 8.52

method	result	size
derivativedivides	Expression too large to display	2446
default	Expression too large to display	2446
parts	Expression too large to display	2454

input

```
int((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^4,x,method=_RETURNVERBOSE)
```

output

```

1/d*(-1/3*a^3/e^4/(d*x+c)^3+b^3/e^4*(1/4*I*Pi*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*arctan(d*x+c)^2+2*I*arctan(d*x+c)*polylog(2,-(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))+2*I*arctan(d*x+c)*polylog(2,(1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))-1/2/(d*x+c)^2*arctan(d*x+c)^2-1/2*I*Pi*csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^3*arctan(d*x+c)^2-1/2*I*Pi*csgn(I*((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^3*arctan(d*x+c)^2+1/2*I*Pi*csgn(((1+I*(d*x+c))^2/(1+(d*x+c)^2)-1)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2*arctan(d*x+c)^2-1/2*I*Pi*arctan(d*x+c)^2-1/2*arctan(d*x+c)*(I*(d*x+c)+(1+(d*x+c)^2)^(1/2)+1)/(d*x+c)-1/4*I*Pi*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^3*arctan(d*x+c)^2+1/4*I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^3*arctan(d*x+c)^2-1/4*I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2))*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^2*arctan(d*x+c)^2-1/4*I*Pi*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2*csgn(I*(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*arctan(d*x+c)^2-1/3/(d*x+c)^3*arctan(d*x+c)^3-1/4*I*Pi*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)*csgn(I*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))^2)^2*arctan(d*x+c)^2-arctan(d*x+c)^2*ln((1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))-1/2*arctan(d*x+c)^2+1/4*I*Pi*csgn(I*(1+I*(d*x+c))^2/(1+(d*x...

```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^4} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="fricas")
```

output

```

integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3)/(d^4*e^4*x^4 + 4*c*d^3*e^4*x^3 + 6*c^2*d^2*e^4*x^2 + 4*c^3*d*e^4*x + c^4*e^4), x)

```

Sympy [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx$$

$$= \int \frac{a^3}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{b^3 \operatorname{atan}^3(c + dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{3ab^2 \operatorname{atan}^2(c + dx)}{c^4 + 4c^3 dx + 6c^2 d^2 x^2 + 4cd^3 x^3 + d^4 x^4} dx + \int \frac{3ab^2 \operatorname{atan}^2(c + dx)}{e^4} dx + \int \frac{3ab^2 \operatorname{atan}^2(c + dx)}{e^4} dx$$

input `integrate((a+b*atan(d*x+c))**3/(d*e*x+c*e)**4,x)`

output `(Integral(a**3/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(b**3*atan(c + d*x)**3/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(3*a*b**2*atan(c + d*x)**2/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x) + Integral(3*a**2*b*atan(c + d*x)/(c**4 + 4*c**3*d*x + 6*c**2*d**2*x**2 + 4*c*d**3*x**3 + d**4*x**4), x))/e**4`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^4} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="maxima")`

output

```
-1/2*(d*(1/(d^4*e^4*x^2 + 2*c*d^3*e^4*x + c^2*d^2*e^4) - log(d^2*x^2 + 2*c
*d*x + c^2 + 1)/(d^2*e^4) + 2*log(d*x + c)/(d^2*e^4)) + 2*arctan(d*x + c)/
(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4))*a^2*b - 1/3
*a^3/(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4) - 1/96*
(4*b^3*arctan(d*x + c)^3 - 3*b^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c
^2 + 1)^2 - 96*(d^4*e^4*x^3 + 3*c*d^3*e^4*x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^
4)*integrate(1/32*(28*(b^3*d^2*x^2 + 2*b^3*c*d*x + b^3*c^2 + b^3)*arctan(d
*x + c)^3 + 4*(24*a*b^2*d^2*x^2 + 24*a*b^2*c^2 + b^3*c + 24*a*b^2 + (48*a*
b^2*c + b^3)*d*x)*arctan(d*x + c)^2 - 4*(b^3*d^2*x^2 + 2*b^3*c*d*x + b^3*c
^2)*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - (b^3*d*x + b^3*c -
3*(b^3*d^2*x^2 + 2*b^3*c*d*x + b^3*c^2 + b^3)*arctan(d*x + c))*log(d^2*x^2
+ 2*c*d*x + c^2 + 1)^2)/(d^6*e^4*x^6 + 6*c*d^5*e^4*x^5 + (15*c^2 + 1)*d^4
*e^4*x^4 + 4*(5*c^3 + c)*d^3*e^4*x^3 + 3*(5*c^4 + 2*c^2)*d^2*e^4*x^2 + 2*(
3*c^5 + 2*c^3)*d*e^4*x + (c^6 + c^4)*e^4), x))/(d^4*e^4*x^3 + 3*c*d^3*e^4*
x^2 + 3*c^2*d^2*e^4*x + c^3*d*e^4)
```

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(dex + ce)^4} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(d*e*x+c*e)^4,x, algorithm="giac")
```

output

```
integrate((b*arctan(d*x + c) + a)^3/(d*e*x + c*e)^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{(ce + dex)^4} dx$$

input

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^4,x)
```

output

```
int((a + b*atan(c + d*x))^3/(c*e + d*e*x)^4, x)
```

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(ce + dex)^4} dx = \text{too large to display}$$

input `int((a+b*atan(d*x+c))^3/(d*e*x+c*e)^4,x)`

output

```
(3*atan(c + d*x)**3*b**3*c**6 + 9*atan(c + d*x)**3*b**3*c**5*d*x + 9*atan(c + d*x)**3*b**3*c**4*d**2*x**2 - 3*atan(c + d*x)**3*b**3*c**4 + 3*atan(c + d*x)**3*b**3*c**3*d**3*x**3 - 9*atan(c + d*x)**3*b**3*c**3*d*x - 9*atan(c + d*x)**3*b**3*c**2*d**2*x**2 - 6*atan(c + d*x)**3*b**3*c**2 - 3*atan(c + d*x)**3*b**3*c*d**3*x**3 + 18*atan(c + d*x)**2*a*b**2*c**6 + 54*atan(c + d*x)**2*a*b**2*c**5*d*x + 54*atan(c + d*x)**2*a*b**2*c**4*d**2*x**2 + 18*atan(c + d*x)**2*a*b**2*c**3*d**3*x**3 - 18*atan(c + d*x)**2*a*b**2*c**2 + 12*atan(c + d*x)**2*b**3*c**7 + 36*atan(c + d*x)**2*b**3*c**6*d*x + 36*atan(c + d*x)**2*b**3*c**5*d**2*x**2 + 12*atan(c + d*x)**2*b**3*c**4*d**3*x**3 - 9*atan(c + d*x)**2*b**3*c**4*d*x - 18*atan(c + d*x)**2*b**3*c**3*d**2*x**2 - 12*atan(c + d*x)**2*b**3*c**3 - 9*atan(c + d*x)**2*b**3*c**2*d**3*x**3 - 18*atan(c + d*x)**2*b**3*c**2*d*x - 9*atan(c + d*x)**2*b**3*c*d**2*x**2 - 18*atan(c + d*x)*a**2*b*c**2 + 36*atan(c + d*x)*a*b**2*c**5 + 72*atan(c + d*x)*a*b**2*c**4*d*x + 36*atan(c + d*x)*a*b**2*c**3*d**2*x**2 - 12*atan(c + d*x)*a*b**2*c**3 + 33*atan(c + d*x)*b**3*c**6 + 75*atan(c + d*x)*b**3*c**5*d*x + 51*atan(c + d*x)*b**3*c**4*d**2*x**2 - 26*atan(c + d*x)*b**3*c**4 + 9*atan(c + d*x)*b**3*c**3*d**3*x**3 - 54*atan(c + d*x)*b**3*c**3*d*x - 45*atan(c + d*x)*b**3*c**2*d**2*x**2 - 3*atan(c + d*x)*b**3*c**2 - 9*atan(c + d*x)*b**3*c*d**3*x**3 - 9*atan(c + d*x)*b**3*c*d*x + 36*int((atan(c + d*x)*x)/(c**6 + 6*c**5*d*x + 15*c**4*d**2*x**2 + c**4 + 20*c**3*...
```


3.29 $\int \frac{\arctan(1+x)}{2+2x} dx$

Optimal result	280
Mathematica [A] (verified)	280
Rubi [A] (verified)	281
Maple [A] (verified)	282
Fricas [F]	283
Sympy [F]	283
Maxima [B] (verification not implemented)	283
Giac [F]	284
Mupad [B] (verification not implemented)	284
Reduce [F]	284

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{\arctan(1+x)}{2+2x} dx = \frac{1}{4}i \operatorname{PolyLog}(2, -i(1+x)) - \frac{1}{4}i \operatorname{PolyLog}(2, i(1+x))$$

output `1/4*I*polylog(2, -I*(1+x))-1/4*I*polylog(2, I*(1+x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(1+x)}{2+2x} dx = \frac{1}{4}i \operatorname{PolyLog}(2, -i(1+x)) - \frac{1}{4}i \operatorname{PolyLog}(2, i(1+x))$$

input `Integrate[ArcTan[1 + x]/(2 + 2*x), x]`

output `(I/4)*PolyLog[2, (-I)*(1 + x)] - (I/4)*PolyLog[2, I*(1 + x)]`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5566, 27, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(x+1)}{2x+2} dx \\
 & \quad \downarrow \text{5566} \\
 & \int \frac{\arctan(x+1)}{2(x+1)} d(x+1) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{\arctan(x+1)}{x+1} d(x+1) \\
 & \quad \downarrow \text{5355} \\
 & \frac{1}{2} \left(\frac{1}{2} i \int \frac{\log(1-i(x+1))}{x+1} d(x+1) - \frac{1}{2} i \int \frac{\log(i(x+1)+1)}{x+1} d(x+1) \right) \\
 & \quad \downarrow \text{2838} \\
 & \frac{1}{2} \left(\frac{1}{2} i \text{PolyLog}(2, -i(x+1)) - \frac{1}{2} i \text{PolyLog}(2, i(x+1)) \right)
 \end{aligned}$$

input `Int[ArcTan[1 + x]/(2 + 2*x), x]`

output `((I/2)*PolyLog[2, (-I)*(1 + x)] - (I/2)*PolyLog[2, I*(1 + x)])/2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x) /; FreeQ[b, x]]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5355 `Int[((a_) + ArcTan[(c_)*(x_)])*(b_)]/(x_), x_Symbol] := Simp[a*Log[x], x] + (Simp[I*(b/2) Int[Log[1 - I*c*x]/x, x] - Simp[I*(b/2) Int[Log[1 + I*c*x]/x, x], x]) /; FreeQ[{a, b, c}, x]`

rule 5566 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_)^(m_)]/(x_), x_Symbol] := Simp[1/d Subst[Int[(f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

method	result
risch	$-\frac{i \operatorname{dilog}(-ix-i+1)}{4} + \frac{i \operatorname{dilog}(ix+i+1)}{4}$
derivativedivides	$\frac{\ln(1+x) \arctan(1+x)}{2} + \frac{i \ln(1+x) \ln(1+i(1+x))}{4} - \frac{i \ln(1+x) \ln(1-i(1+x))}{4} + \frac{i \operatorname{dilog}(1+i(1+x))}{4} - \frac{i \operatorname{dilog}(1-i(1+x))}{4}$
default	$\frac{\ln(1+x) \arctan(1+x)}{2} + \frac{i \ln(1+x) \ln(1+i(1+x))}{4} - \frac{i \ln(1+x) \ln(1-i(1+x))}{4} + \frac{i \operatorname{dilog}(1+i(1+x))}{4} - \frac{i \operatorname{dilog}(1-i(1+x))}{4}$
parts	$\frac{\ln(1+x) \arctan(1+x)}{2} + \frac{i \ln(1+x) \ln(1+i(1+x))}{4} - \frac{i \ln(1+x) \ln(1-i(1+x))}{4} + \frac{i \operatorname{dilog}(1+i(1+x))}{4} - \frac{i \operatorname{dilog}(1-i(1+x))}{4}$

input `int(arctan(1+x)/(2+2*x), x, method=_RETURNVERBOSE)`

output `-1/4*I*dilog(-I*x+1-I)+1/4*I*dilog(I*x+1+I)`

Fricas [F]

$$\int \frac{\arctan(1+x)}{2+2x} dx = \int \frac{\arctan(x+1)}{2(x+1)} dx$$

input `integrate(arctan(1+x)/(2+2*x),x, algorithm="fricas")`

output `integral(1/2*arctan(x + 1)/(x + 1), x)`

Sympy [F]

$$\int \frac{\arctan(1+x)}{2+2x} dx = \int \frac{\operatorname{atan}(x+1)}{2} dx$$

input `integrate(atan(1+x)/(2+2*x),x)`

output `Integral(atan(x + 1)/(x + 1), x)/2`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(17) = 34$.

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\begin{aligned} \int \frac{\arctan(1+x)}{2+2x} dx = & -\frac{1}{4} \arctan(x+1, 0) \log(x^2 + 2x + 2) \\ & + \frac{1}{2} \arctan(x+1) \log(|x+1|) \\ & - \frac{1}{4} i \operatorname{Li}_2(ix + i + 1) + \frac{1}{4} i \operatorname{Li}_2(-ix - i + 1) \end{aligned}$$

input `integrate(arctan(1+x)/(2+2*x),x, algorithm="maxima")`

output $-1/4*\arctan2(x + 1, 0)*\log(x^2 + 2*x + 2) + 1/2*\arctan(x + 1)*\log(\text{abs}(x + 1)) - 1/4*I*\text{dilog}(I*x + I + 1) + 1/4*I*\text{dilog}(-I*x - I + 1)$

Giac [F]

$$\int \frac{\arctan(1+x)}{2+2x} dx = \int \frac{\arctan(x+1)}{2(x+1)} dx$$

input `integrate(arctan(1+x)/(2+2*x),x, algorithm="giac")`

output `integrate(1/2*arctan(x + 1)/(x + 1), x)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{\arctan(1+x)}{2+2x} dx = -\frac{\text{Li}_2(1-x \text{li} - i) \text{li}}{4} + \frac{\text{Li}_2(x \text{li} + 1 + \text{li}) \text{li}}{4}$$

input `int(atan(x + 1)/(2*x + 2),x)`

output `(dilog(x*1i + (1 + 1i))*1i)/4 - (dilog((1 - 1i) - x*1i)*1i)/4`

Reduce [F]

$$\int \frac{\arctan(1+x)}{2+2x} dx = \frac{\left(\int \frac{\text{atan}(x+1)}{x+1} dx\right)}{2}$$

input `int(atan(1+x)/(2+2*x),x)`

output `int(atan(x + 1)/(x + 1),x)/2`

3.30 $\int \frac{\arctan(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	285
Mathematica [A] (verified)	285
Rubi [A] (verified)	286
Maple [A] (verified)	287
Fricas [F]	288
Sympy [F]	288
Maxima [B] (verification not implemented)	288
Giac [F]	289
Mupad [F(-1)]	289
Reduce [F]	290

Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \frac{i \operatorname{PolyLog}(2, -i(a + bx))}{2d} - \frac{i \operatorname{PolyLog}(2, i(a + bx))}{2d}$$

output `1/2*I*polylog(2,-I*(b*x+a))/d-1/2*I*polylog(2,I*(b*x+a))/d`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \frac{i(\operatorname{PolyLog}(2, -i(a + bx)) - \operatorname{PolyLog}(2, i(a + bx)))}{2d}$$

input `Integrate[ArcTan[a + b*x]/((a*d)/b + d*x),x]`

output `((I/2)*(PolyLog[2, (-I)*(a + b*x)] - PolyLog[2, I*(a + b*x)]))/d`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {5566, 27, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx \\
 & \quad \downarrow \text{5566} \\
 & \int \frac{b \arctan(a+bx)}{d(a+bx)} d(a + bx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{\arctan(a+bx)}{a+bx} d(a + bx) \\
 & \quad \downarrow \text{5355} \\
 & \frac{\frac{1}{2}i \int \frac{\log(1-i(a+bx))}{a+bx} d(a + bx) - \frac{1}{2}i \int \frac{\log(i(a+bx)+1)}{a+bx} d(a + bx)}{d} \\
 & \quad \downarrow \text{2838} \\
 & \frac{\frac{1}{2}i \text{PolyLog}(2, -i(a + bx)) - \frac{1}{2}i \text{PolyLog}(2, i(a + bx))}{d}
 \end{aligned}$$

input `Int[ArcTan[a + b*x]/((a*d)/b + d*x), x]`

output `((I/2)*PolyLog[2, (-I)*(a + b*x)] - (I/2)*PolyLog[2, I*(a + b*x)])/d`

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)(Gx_)] /; \text{FreeQ}[b, x]$

rule 2838 $\text{Int}[\text{Log}[(c_)((d_)+(e_)(x_)^{n_})]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \ \&\& \ \text{EqQ}[c*d, 1]$

rule 5355 $\text{Int}[(a_)+\text{ArcTan}[(c_)(x_)]*(b_)]/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] + (\text{Simp}[I*(b/2) \text{ Int}[\text{Log}[1-I*c*x]/x, x] - \text{Simp}[I*(b/2) \text{ Int}[\text{Log}[1+I*c*x]/x, x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 5566 $\text{Int}[(a_)+\text{ArcTan}[(c_)+(d_)(x_)]*(b_)]^{(p_)}*((e_)+(f_)(x_))^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^{m*}(a+b*\text{ArcTan}[x])^p, x], x, c+d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{EqQ}[d*e-c*f, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

method	result
risch	$-\frac{i \operatorname{dilog}(-ibx-ia+1)}{2d} + \frac{i \operatorname{dilog}(ibx+ia+1)}{2d}$
parts	$\frac{\ln(bx+a) \arctan(bx+a)}{d} - \frac{-\frac{i \ln(bx+a) \ln(1+i(bx+a))}{2} + \frac{i \ln(bx+a) \ln(1-i(bx+a))}{2} - \frac{i \operatorname{dilog}(1+i(bx+a))}{2} + \frac{i \operatorname{dilog}(1-i(bx+a))}{2}}{d}$
derivativedivides	$\frac{\frac{b \ln(bx+a) \arctan(bx+a)}{d} - b \left(-\frac{i \ln(bx+a) \ln(1+i(bx+a))}{2} + \frac{i \ln(bx+a) \ln(1-i(bx+a))}{2} - \frac{i \operatorname{dilog}(1+i(bx+a))}{2} + \frac{i \operatorname{dilog}(1-i(bx+a))}{2} \right)}{b}$
default	$\frac{\frac{b \ln(bx+a) \arctan(bx+a)}{d} - b \left(-\frac{i \ln(bx+a) \ln(1+i(bx+a))}{2} + \frac{i \ln(bx+a) \ln(1-i(bx+a))}{2} - \frac{i \operatorname{dilog}(1+i(bx+a))}{2} + \frac{i \operatorname{dilog}(1-i(bx+a))}{2} \right)}{b}$

input $\text{int}(\arctan(b*x+a)/(a*d/b+d*x), x, \text{method}=_RETURNVERBOSE)$

output $-1/2*I/d*\operatorname{dilog}(1-I*a-I*b*x)+1/2*I/d*\operatorname{dilog}(1+I*a+I*b*x)$

Fricas [F]

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\arctan(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arctan(b*x+a)/(a*d/b+d*x),x, algorithm="fricas")`

output `integral(b*arctan(b*x + a)/(b*d*x + a*d), x)`

Sympy [F]

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{atan}(a+bx)}{a+bx} dx}{d}$$

input `integrate(atan(b*x+a)/(a*d/b+d*x), x)`

output `b*Integral(atan(a + b*x)/(a + b*x), x)/d`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(29) = 58$.

Time = 0.20 (sec) , antiderivative size = 123, normalized size of antiderivative = 3.00

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\arctan(bx + a) \log(dx + \frac{ad}{b})}{d} - \frac{\arctan\left(\frac{b^2x+ab}{b}\right) \log(dx + \frac{ad}{b})}{d} - \frac{\arctan(bx + a, 0) \log(b^2x^2 + 2abx + a^2 + 1) - 2 \arctan(bx + a) \log(|bx + a|) + i \operatorname{Li}_2(ibx + ia + 1)}{2d}$$

input `integrate(arctan(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")`

output

```
arctan(b*x + a)*log(d*x + a*d/b)/d - arctan((b^2*x + a*b)/b)*log(d*x + a*d
/b)/d - 1/2*(arctan2(b*x + a, 0)*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*arct
an(b*x + a)*log(abs(b*x + a)) + I*dilog(I*b*x + I*a + 1) - I*dilog(-I*b*x
- I*a + 1))/d
```

Giac [F]

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\arctan(bx + a)}{dx + \frac{ad}{b}} dx$$

input

```
integrate(arctan(b*x+a)/(a*d/b+d*x),x, algorithm="giac")
```

output

```
integrate(arctan(b*x + a)/(d*x + a*d/b), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{atan}(a + bx)}{dx + \frac{ad}{b}} dx$$

input

```
int(atan(a + b*x)/(d*x + (a*d)/b),x)
```

output

```
int(atan(a + b*x)/(d*x + (a*d)/b), x)
```

Reduce [F]

$$\int \frac{\arctan(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\left(\int \frac{\arctan(bx+a)}{bx+a} dx\right) b}{d}$$

input `int(atan(b*x+a)/(a*d/b+d*x),x)`

output `(int(atan(a + b*x)/(a + b*x),x)*b)/d`

3.31 $\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx$

Optimal result	291
Mathematica [N/A]	291
Rubi [N/A]	292
Maple [N/A]	292
Fricas [F(-2)]	293
Sympy [N/A]	293
Maxima [F(-2)]	293
Giac [N/A]	294
Mupad [N/A]	294
Reduce [N/A]	294

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \text{Int}\left((a + bx)^2 \sqrt{\arctan(a + bx)}, x\right)$$

output `Defer(Int)((b*x+a)^2*arctan(b*x+a)^(1/2), x)`

Mathematica [N/A]

Not integrable

Time = 3.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \int (a + bx)^2 \sqrt{\arctan(a + bx)} dx$$

input `Integrate[(a + b*x)^2*Sqrt[ArcTan[a + b*x]], x]`

output `Integrate[(a + b*x)^2*Sqrt[ArcTan[a + b*x]], x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx$$

$$\downarrow 5572$$

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx$$

input `Int[(a + b*x)^2*Sqrt[ArcTan[a + b*x]],x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int (bx + a)^2 \sqrt{\arctan(bx + a)} dx$$

input `int((b*x+a)^2*arctan(b*x+a)^(1/2),x)`

output `int((b*x+a)^2*arctan(b*x+a)^(1/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \text{Exception raised: TypeError}$$

input `integrate((b*x+a)^2*arctan(b*x+a)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [N/A]

Not integrable

Time = 10.91 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \int (a + bx)^2 \sqrt{\text{atan}(a + bx)} dx$$

input `integrate((b*x+a)**2*atan(b*x+a)**(1/2),x)`

output `Integral((a + b*x)**2*sqrt(atan(a + b*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((b*x+a)^2*arctan(b*x+a)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [N/A]

Not integrable

Time = 121.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.17

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \int (bx + a)^2 \sqrt{\arctan(bx + a)} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)^(1/2),x, algorithm="giac")`

output `sage0*x`

Mupad [N/A]

Not integrable

Time = 1.33 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx = \int \sqrt{\arctan(a + bx)} (a + bx)^2 dx$$

input `int(atan(a + b*x)^(1/2)*(a + b*x)^2,x)`

output `int(atan(a + b*x)^(1/2)*(a + b*x)^2, x)`

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 224, normalized size of antiderivative = 12.44

$$\int (a + bx)^2 \sqrt{\arctan(a + bx)} dx$$

$$= \sqrt{\arctan(bx + a)} a^2 x + \sqrt{\arctan(bx + a)} ab x^2 + \frac{\sqrt{\arctan(bx + a)} b^2 x^3}{3}$$

$$- \frac{\left(\int \frac{\sqrt{\arctan(bx+a)} x^3}{\arctan(bx+a)a^2 + 2\arctan(bx+a)abx + \arctan(bx+a)b^2x^2 + \arctan(bx+a)} dx \right) b^3}{6}$$

$$- \frac{\left(\int \frac{\sqrt{\arctan(bx+a)} x^2}{\arctan(bx+a)a^2 + 2\arctan(bx+a)abx + \arctan(bx+a)b^2x^2 + \arctan(bx+a)} dx \right) a b^2}{2}$$

$$- \frac{\left(\int \frac{\sqrt{\arctan(bx+a)} x}{\arctan(bx+a)a^2 + 2\arctan(bx+a)abx + \arctan(bx+a)b^2x^2 + \arctan(bx+a)} dx \right) a^2 b}{2}$$

input

```
int((b*x+a)^2*atan(b*x+a)^(1/2),x)
```

output

```
(6*sqrt(atan(a + b*x))*a**2*x + 6*sqrt(atan(a + b*x))*a*b*x**2 + 2*sqrt(atan(a + b*x))*b**2*x**3 - int((sqrt(atan(a + b*x))*x**3)/(atan(a + b*x)*a**2 + 2*atan(a + b*x)*a*b*x + atan(a + b*x)*b**2*x**2 + atan(a + b*x)),x)*b**3 - 3*int((sqrt(atan(a + b*x))*x**2)/(atan(a + b*x)*a**2 + 2*atan(a + b*x)*a*b*x + atan(a + b*x)*b**2*x**2 + atan(a + b*x)),x)*a*b**2 - 3*int((sqrt(atan(a + b*x))*x)/(atan(a + b*x)*a**2 + 2*atan(a + b*x)*a*b*x + atan(a + b*x)*b**2*x**2 + atan(a + b*x)),x)*a**2*b)/6
```


3.32 $\int (e + fx)^3 (a + b \arctan(c + dx)) dx$

Optimal result	296
Mathematica [C] (verified)	297
Rubi [A] (verified)	297
Maple [B] (verified)	299
Fricas [A] (verification not implemented)	300
Sympy [C] (verification not implemented)	301
Maxima [A] (verification not implemented)	302
Giac [B] (verification not implemented)	303
Mupad [B] (verification not implemented)	304
Reduce [B] (verification not implemented)	305

Optimal result

Integrand size = 18, antiderivative size = 233

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx$$

$$= -\frac{bf(6d^2e^2 - 12cdef - (1 - 6c^2)f^2)x}{4d^3} - \frac{bf^2(de - cf)(c + dx)^2}{2d^4} - \frac{bf^3(c + dx)^3}{12d^4}$$

$$- \frac{b(d^4e^4 - 4cd^3e^3f - 6(1 - c^2)d^2e^2f^2 + 4c(3 - c^2)def^3 + (1 - 6c^2 + c^4)f^4) \arctan(c + dx)}{4d^4f}$$

$$+ \frac{(e + fx)^4(a + b \arctan(c + dx))}{4f}$$

$$- \frac{b(de - cf)(de + f - cf)(de - (1 + c)f) \log(1 + (c + dx)^2)}{2d^4}$$

output

```
-1/4*b*f*(6*d^2*e^2-12*c*d*e*f-(-6*c^2+1)*f^2)*x/d^3-1/2*b*f^2*(-c*f+d*e)*
(d*x+c)^2/d^4-1/12*b*f^3*(d*x+c)^3/d^4-1/4*b*(d^4*e^4-4*c*d^3*e^3*f-6*(-c^
2+1)*d^2*e^2*f^2+4*c*(-c^2+3)*d*e*f^3+(c^4-6*c^2+1)*f^4)*arctan(d*x+c)/d^4
/f+1/4*(f*x+e)^4*(a+b*arctan(d*x+c))/f-1/2*b*(-c*f+d*e)*(-c*f+d*e+f)*(d*e-
(1+c)*f)*ln(1+(d*x+c)^2)/d^4
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.67

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{(e + fx)^4 (a + b \arctan(c + dx)) - \frac{b(6df^2(6d^2e^2 - 12cdef + (-1 + 6c^2)f^2)x + 12f^3(de - cf)(c + dx)^2 + 2f^4(c + dx)^3 - 3i(de - (-i + c)f)}{6d^4}}{4f}}$$

input `Integrate[(e + f*x)^3*(a + b*ArcTan[c + d*x]),x]`

output $((e + f*x)^4*(a + b*ArcTan[c + d*x]) - (b*(6*d*f^2*(6*d^2*e^2 - 12*c*d*e*f + (-1 + 6*c^2)*f^2)*x + 12*f^3*(d*e - c*f)*(c + d*x)^2 + 2*f^4*(c + d*x)^3 - (3*I)*(d*e - (-I + c)*f)^4*Log[I - c - d*x] + (3*I)*(d*e - (I + c)*f)^4*Log[I + c + d*x]))/(6*d^4))/(4*f)$

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5570, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx$$

$$\downarrow 5570$$

$$\int \frac{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)^3 (a + b \arctan(c + dx))}{d^3} d(c + dx)$$

$$\downarrow 27$$

$$\frac{\int (de - cf + f(c + dx))^3 (a + b \arctan(c + dx)) d(c + dx)}{d^4}$$

$$\begin{aligned} & \downarrow \text{5387} \\ & \frac{(f(c+dx)-cf+de)^4(a+b \arctan(c+dx))}{4f} - \frac{b \int \frac{(de-cf+f(c+dx))^4 d(c+dx)}{(c+dx)^2+1}}{4f} \\ & \downarrow \text{478} \\ & \frac{(f(c+dx)-cf+de)^4(a+b \arctan(c+dx))}{4f} - \frac{b \int \left((c+dx)^2 f^4 + 4(de-cf)(c+dx)f^3 + (6d^2 e^2 - 12cdf e - (1-6c^2)f^2) f^2 + \frac{d^4 e^4 - 4cd^3 f e^3 - 6(1-c^2)d^2 f^2}{4f} \right)}{d^4} \\ & \downarrow \text{2009} \\ & \frac{(f(c+dx)-cf+de)^4(a+b \arctan(c+dx))}{4f} - \frac{b(\arctan(c+dx)(-6(1-c^2)d^2 e^2 f^2 + 4c(3-c^2)de f^3 + (c^4 - 6c^2 + 1)f^4 - 4cd^3 e^3 f + d^4 e^4) + f^2(c+dx)(-))}{d^4} \end{aligned}$$

input `Int[(e + f*x)^3*(a + b*ArcTan[c + d*x]),x]`

output `((d*e - c*f + f*(c + d*x))^4*(a + b*ArcTan[c + d*x]))/(4*f) - (b*(f^2*(6*d^2*e^2 - 12*c*d*e*f - (1 - 6*c^2)*f^2)*(c + d*x) + 2*f^3*(d*e - c*f)*(c + d*x)^2 + (f^4*(c + d*x)^3)/3 + (d^4*e^4 - 4*c*d^3*e^3*f - 6*(1 - c^2)*d^2*e^2*f^2 + 4*c*(3 - c^2)*d*e*f^3 + (1 - 6*c^2 + c^4)*f^4)*ArcTan[c + d*x] + 2*f*(d*e - c*f)*(d*e + f - c*f)*(d*e - (1 + c)*f)*Log[1 + (c + d*x)^2])/(4*f))/d^4`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 478 `Int[((c_) + (d_)*(x_))^(n_)/((a_) + (b_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 5387 Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))*((d_) + (e_.)*(x_.))^(q_.), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(
  c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b
  , c, d, e, q}, x] && NeQ[q, -1]
```

```
rule 5570 Int[((a_.) + ArcTan[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 463 vs. 2(221) = 442.

Time = 0.35 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.99

method	result
parts	$\frac{2f^2bce x}{d^2} - \frac{3f^2bce \arctan(dx+c)}{d^3} + \frac{f^2b c^3 e \arctan(dx+c)}{d^3} - \frac{3fb c^2 e^2 \arctan(dx+c)}{2d^2} + \frac{b f^3 c}{4d^4} - \frac{13b f^3 c^3}{12d^4} - \frac{f^3}{1}$
derivativeldivides	$\frac{a(cf-de-f(dx+c))^4}{4d^3 f} - \frac{b \left(-\frac{f^3 \arctan(dx+c)c^4}{4} + f^2 \arctan(dx+c)c^3 de + f^3 \arctan(dx+c)c^3(dx+c) - \frac{3f \arctan(dx+c)c^2 d^2 e^2}{2} - 3f^2 \right)}{4d^3 f}$
default	$\frac{a(cf-de-f(dx+c))^4}{4d^3 f} - \frac{b \left(-\frac{f^3 \arctan(dx+c)c^4}{4} + f^2 \arctan(dx+c)c^3 de + f^3 \arctan(dx+c)c^3(dx+c) - \frac{3f \arctan(dx+c)c^2 d^2 e^2}{2} - 3f^2 \right)}{4d^3 f}$
parallelsch	$-3 \arctan(dx+c) b c^4 f^3 + 18 \arctan(dx+c) b c^2 f^3 + 3x^4 a d^4 f^3 + 12x a d^4 e^3 + 3x b d f^3 - x^3 b d^3 f^3 + 6 \ln(d^2 x^2 + 2cdx + c^2 + 1)$
risch	$\frac{2f^2bce x}{d^2} - \frac{3f^2bce \arctan(dx+c)}{d^3} + \frac{f^2b c^3 e \arctan(dx+c)}{d^3} - \frac{3fb c^2 e^2 \arctan(dx+c)}{2d^2} - \frac{3f^2b c^2 e \ln(d^2 x^2 + 2cdx + c^2 + 1)}{2d^3}$

```
input int((f*x+e)^3*(a+b*arctan(d*x+c)), x, method=_RETURNVERBOSE)
```

output

```
2/d^2*f^2*b*c*e*x-3/d^3*f^2*b*c*e*arctan(d*x+c)+1/d^3*f^2*b*c^3*e*arctan(d
*x+c)-3/2/d^2*f*b*c^2*e^2*arctan(d*x+c)+1/4*b/d^4*f^3*c-13/12*b/d^4*f^3*c^
3-1/12/d*f^3*b*x^3+1/4/d^3*f^3*b*x-1/4/d^4*f^3*b*arctan(d*x+c)+1/4*a*(f*x+
e)^4/f+1/2*b/d^4*f^3*ln(1+(d*x+c)^2)*c^3-1/2*b/d^4*f^3*ln(1+(d*x+c)^2)*c+1
/2*b/d^3*f^2*ln(1+(d*x+c)^2)*e+b*f^2*arctan(d*x+c)*e*x^3+3/2*b*f*arctan(d*
x+c)*e^2*x^2+5/2*b/d^3*f^2*c^2*e-3/2*b/d^2*f*c*e^2-3/2*b/d^3*f^2*ln(1+(d*x
+c)^2)*c^2*e+3/2*b/d^2*f*ln(1+(d*x+c)^2)*c*e^2-1/2*b/d*ln(1+(d*x+c)^2)*e^3
+1/4*b*f^3*arctan(d*x+c)*x^4+b*arctan(d*x+c)*x*e^3+1/4/d^2*f^3*b*c*x^2-1/2
/d*f^2*b*e*x^2-3/4/d^3*f^3*b*c^2*x-3/2/d*f*b*e^2*x+1/d*b*c*e^3*arctan(d*x+
c)-1/4/d^4*f^3*b*c^4*arctan(d*x+c)+3/2/d^4*f^3*b*c^2*arctan(d*x+c)+3/2/d^2
*f*b*e^2*arctan(d*x+c)
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.36

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{3ad^4f^3x^4 + (12ad^4ef^2 - bd^3f^3)x^3 + 3(6ad^4e^2f - 2bd^3ef^2 + bcd^2f^3)x^2 + 3(4ad^4e^3 - 6bd^3e^2f + 8bcd^2ef^2 - 3b^2c^2f^2 - (3b^2c^2 - b)d^2f^3)x + 3(bd^4f^3x^4 + 4bd^4ef^2x^3 + 6bd^4e^2fx^2 + 4bd^4e^3x + 4b^2cd^3e^3 - 6(b^2c^2 - b)d^2e^2f + 4(b^2c^3 - 3b^2c)d^2ef^2 - (b^2c^4 - 6b^2c^2 + b)f^3) \arctan(dx + c) - 6(bd^3e^3 - 3b^2cd^2e^2f + (3b^2c^2 - b)d^2ef^2 - (b^2c^3 - b^2c)f^3) \log(d^2x^2 + 2cdx + c^2 + 1)}{d^4}$$

input

```
integrate((f*x+e)^3*(a+b*arctan(d*x+c)),x, algorithm="fricas")
```

output

```
1/12*(3*a*d^4*f^3*x^4 + (12*a*d^4*e*f^2 - b*d^3*f^3)*x^3 + 3*(6*a*d^4*e^2*
f - 2*b*d^3*e*f^2 + b*c*d^2*f^3)*x^2 + 3*(4*a*d^4*e^3 - 6*b*d^3*e^2*f + 8*
b*c*d^2*e*f^2 - (3*b*c^2 - b)*d^2*f^3)*x + 3*(b*d^4*f^3*x^4 + 4*b*d^4*e*f^2*
x^3 + 6*b*d^4*e^2*f*x^2 + 4*b*d^4*e^3*x + 4*b*c*d^3*e^3 - 6*(b*c^2 - b)*d^
2*e^2*f + 4*(b*c^3 - 3*b*c)*d^2*e*f^2 - (b*c^4 - 6*b*c^2 + b)*f^3)*arctan(d*
x + c) - 6*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + (3*b*c^2 - b)*d^2*e*f^2 - (b*c^3 -
b*c)*f^3)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^4
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 12.08 (sec) , antiderivative size = 654, normalized size of antiderivative = 2.81

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)**3*(a+b*atan(d*x+c)),x)`

output

```
Piecewise((a*e**3*x + 3*a*e**2*f*x**2/2 + a*e*f**2*x**3 + a*f**3*x**4/4 -
b*c**4*f**3*atan(c + d*x)/(4*d**4) + b*c**3*e*f**2*atan(c + d*x)/d**3 + b*
c**3*f**3*log(c/d + x - I/d)/d**4 - I*b*c**3*f**3*atan(c + d*x)/d**4 - 3*b
*c**2*e**2*f*atan(c + d*x)/(2*d**2) - 3*b*c**2*e*f**2*log(c/d + x - I/d)/d
**3 + 3*I*b*c**2*e*f**2*atan(c + d*x)/d**3 - 3*b*c**2*f**3*x/(4*d**3) + 3*
b*c**2*f**3*atan(c + d*x)/(2*d**4) + b*c*e**3*atan(c + d*x)/d + 3*b*c*e**2
*f*log(c/d + x - I/d)/d**2 - 3*I*b*c*e**2*f*atan(c + d*x)/d**2 + 2*b*c*e*f
**2*x/d**2 + b*c*f**3*x**2/(4*d**2) - 3*b*c*e*f**2*atan(c + d*x)/d**3 - b*
c*f**3*log(c/d + x - I/d)/d**4 + I*b*c*f**3*atan(c + d*x)/d**4 + b*e**3*x*
atan(c + d*x) + 3*b*e**2*f*x**2*atan(c + d*x)/2 + b*e*f**2*x**3*atan(c + d
*x) + b*f**3*x**4*atan(c + d*x)/4 - b*e**3*log(c/d + x - I/d)/d + I*b*e**3
*atan(c + d*x)/d - 3*b*e**2*f*x/(2*d) - b*e*f**2*x**2/(2*d) - b*f**3*x**3/
(12*d) + 3*b*e**2*f*atan(c + d*x)/(2*d**2) + b*e*f**2*log(c/d + x - I/d)/d
**3 - I*b*e*f**2*atan(c + d*x)/d**3 + b*f**3*x/(4*d**3) - b*f**3*atan(c +
d*x)/(4*d**4), Ne(d, 0)), ((a + b*atan(c))*(e**3*x + 3*e**2*f*x**2/2 + e*f
**2*x**3 + f**3*x**4/4), True))
```

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.48

$$\begin{aligned}
\int (e + fx)^3 (a + b \arctan(c + dx)) dx = & \frac{1}{4} af^3 x^4 + aef^2 x^3 + \frac{3}{2} ae^2 fx^2 \\
& + \frac{3}{2} \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^3} - \frac{c \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) be^2 f \\
& + \frac{1}{2} \left(2x^3 \arctan(dx + c) - d \left(\frac{dx^2 - 4cx}{d^3} - \frac{2(c^3 - 3c) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^4} + \frac{(3c^2 - 1) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^4} \right) \right) be^2 f \\
& + \frac{1}{12} \left(3x^4 \arctan(dx + c) - d \left(\frac{d^2 x^3 - 3cdx^2 + 3(3c^2 - 1)x}{d^4} + \frac{3(c^4 - 6c^2 + 1) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^5} - \frac{6(c^3 - c) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^5} \right) \right) be^2 f \\
& + ae^3 x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) be^3}{2d}
\end{aligned}$$

input `integrate((f*x+e)^3*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output

```

1/4*a*f^3*x^4 + a*e*f^2*x^3 + 3/2*a*e^2*f*x^2 + 3/2*(x^2*arctan(d*x + c) -
d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*
x + c^2 + 1)/d^3))*b*e^2*f + 1/2*(2*x^3*arctan(d*x + c) - d*((d*x^2 - 4*c*
x)/d^3 - 2*(c^3 - 3*c)*arctan((d^2*x + c*d)/d)/d^4 + (3*c^2 - 1)*log(d^2*x
^2 + 2*c*d*x + c^2 + 1)/d^4))*b*e*f^2 + 1/12*(3*x^4*arctan(d*x + c) - d*((
d^2*x^3 - 3*c*d*x^2 + 3*(3*c^2 - 1)*x)/d^4 + 3*(c^4 - 6*c^2 + 1)*arctan((d
^2*x + c*d)/d)/d^5 - 6*(c^3 - c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^5))*b*
f^3 + a*e^3*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b
*e^3/d

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 885 vs. $2(216) = 432$.

Time = 1.22 (sec) , antiderivative size = 885, normalized size of antiderivative = 3.80

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)^3*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output

```
1/24*(6*b*d^4*f^3*x^4*arctan(d*x + c) + 6*a*d^4*f^3*x^4 + 24*b*d^4*e*f^2*x
^3*arctan(d*x + c) + 24*a*d^4*e*f^2*x^3 + 36*b*d^4*e^2*f*x^2*arctan(d*x +
c) + 36*a*d^4*e^2*f*x^2 - 2*b*d^3*f^3*x^3 + 24*b*d^4*e^3*x*arctan(d*x + c)
- 12*pi*b*c*d^3*e^3*sgn(d*x + c) + 18*pi*b*c^2*d^2*e^2*f*sgn(d*x + c) - 1
2*pi*b*c^3*d*e*f^2*sgn(d*x + c) + 3*pi*b*c^4*f^3*sgn(d*x + c) + 12*pi*b*c*
d^3*e^3*sgn(-d*x - c) - 18*pi*b*c^2*d^2*e^2*f*sgn(-d*x - c) + 12*pi*b*c^3*
d*e*f^2*sgn(-d*x - c) - 3*pi*b*c^4*f^3*sgn(-d*x - c) + 24*a*d^4*e^3*x - 12
*b*d^3*e*f^2*x^2 + 6*b*c*d^2*f^3*x^2 + 12*b*c*d^3*e^3*arctan(d*x + c) - 18
*b*c^2*d^2*e^2*f*arctan(d*x + c) + 12*b*c^3*d*e*f^2*arctan(d*x + c) - 3*b*
c^4*f^3*arctan(d*x + c) - 12*b*c*d^3*e^3*arctan(-d*x - c) + 18*b*c^2*d^2*e
^2*f*arctan(-d*x - c) - 12*b*c^3*d*e*f^2*arctan(-d*x - c) + 3*b*c^4*f^3*ar
ctan(-d*x - c) - 36*b*d^3*e^2*f*x + 48*b*c*d^2*e*f^2*x - 18*b*c^2*d*f^3*x
- 12*b*d^3*e^3*log((d*x + c)^2 + 1) + 36*b*c*d^2*e^2*f*log((d*x + c)^2 + 1
) - 36*b*c^2*d*e*f^2*log((d*x + c)^2 + 1) + 12*b*c^3*f^3*log((d*x + c)^2 +
1) - 18*pi*b*d^2*e^2*f*sgn(d*x + c) + 36*pi*b*c*d*e*f^2*sgn(d*x + c) - 18
*pi*b*c^2*f^3*sgn(d*x + c) + 18*pi*b*d^2*e^2*f*sgn(-d*x - c) - 36*pi*b*c*d
*e*f^2*sgn(-d*x - c) + 18*pi*b*c^2*f^3*sgn(-d*x - c) + 18*b*d^2*e^2*f*arct
an(d*x + c) - 36*b*c*d*e*f^2*arctan(d*x + c) + 18*b*c^2*f^3*arctan(d*x + c
) - 18*b*d^2*e^2*f*arctan(-d*x - c) + 36*b*c*d*e*f^2*arctan(-d*x - c) - 18
*b*c^2*f^3*arctan(-d*x - c) + 6*b*d*f^3*x + 12*b*d*e*f^2*log((d*x + c)^...
```


Mupad [B] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 787, normalized size of antiderivative = 3.38

$$\begin{aligned}
& \int (e + fx)^3 (a + b \arctan(c + dx)) dx \\
&= \operatorname{atan}(c + dx) \left(b e^3 x + \frac{3 b e^2 f x^2}{2} + b e f^2 x^3 + \frac{b f^3 x^4}{4} \right) \\
&+ x \left(\frac{e (6 a c^2 f^2 + 12 a c d e f + 2 a d^2 e^2 - 3 b d e f + 6 a f^2)}{2 d^2} \right. \\
&\quad \left. - \frac{(4 c^2 + 4) \left(\frac{f^2 (8 a c f - b f + 12 a d e)}{4 d} - \frac{2 a c f^3}{d} \right)}{4 d^2} \right. \\
&\quad \left. + \frac{2 c \left(\frac{2 c \left(\frac{f^2 (8 a c f - b f + 12 a d e)}{4 d} - \frac{2 a c f^3}{d} \right)}{d} - \frac{4 a c^2 f^3 + 24 a c d e f^2 + 12 a d^2 e^2 f - 4 b d e f^2 + 4 a f^3}{4 d^2} + \frac{a f^3 (4 c^2 + 4)}{4 d^2} \right)}{d} \right) \\
&- x^2 \left(\frac{c \left(\frac{f^2 (8 a c f - b f + 12 a d e)}{4 d} - \frac{2 a c f^3}{d} \right)}{d} \right. \\
&\quad \left. - \frac{4 a c^2 f^3 + 24 a c d e f^2 + 12 a d^2 e^2 f - 4 b d e f^2 + 4 a f^3}{8 d^2} + \frac{a f^3 (4 c^2 + 4)}{8 d^2} \right) \\
&+ x^3 \left(\frac{f^2 (8 a c f - b f + 12 a d e)}{12 d} - \frac{2 a c f^3}{3 d} \right) + \frac{a f^3 x^4}{4} \\
&- \frac{\ln(c^2 + 2 c d x + d^2 x^2 + 1) (-64 b c^3 d^4 f^3 + 192 b c^2 d^5 e f^2 - 192 b c d^6 e^2 f + 64 b c d^4 f^3 + 64 b d^7 e^3)}{128 d^8} \\
&- b \operatorname{atan} \left(\frac{4 d^3 \left(\frac{c (c^4 f^3 - 4 c^3 d e f^2 + 6 c^2 d^2 e^2 f - 6 c^2 f^3 - 4 c d^3 e^3 + 12 c d e f^2 - 6 d^2 e^2 f + f^3)}{4 d^3} + \frac{x (c^4 f^3 - 4 c^3 d e f^2 + 6 c^2 d^2 e^2 f - 6 c^2 f^3 - 4 c d^3 e^3 + 12 c d e f^2 - 6 d^2 e^2 f + f^3)}{4 d^2} \right)}{c^4 f^3 - 4 c^3 d e f^2 + 6 c^2 d^2 e^2 f - 6 c^2 f^3 - 4 c d^3 e^3 + 12 c d e f^2 - 6 d^2 e^2 f + f^3} \right)
\end{aligned}$$

4

input `int((e + f*x)^3*(a + b*atan(c + d*x)),x)`

output

```
atan(c + d*x)*((b*f^3*x^4)/4 + b*e^3*x + (3*b*e^2*f*x^2)/2 + b*e*f^2*x^3)
+ x*((e*(6*a*f^2 + 6*a*c^2*f^2 + 2*a*d^2*e^2 - 3*b*d*e*f + 12*a*c*d*e*f))/
(2*d^2) - ((4*c^2 + 4)*(f^2*(8*a*c*f - b*f + 12*a*d*e))/(4*d) - (2*a*c*f^
3)/d))/(4*d^2) + (2*c*((2*c*((f^2*(8*a*c*f - b*f + 12*a*d*e))/(4*d) - (2*a
*c*f^3)/d))/d - (4*a*f^3 + 4*a*c^2*f^3 - 4*b*d*e*f^2 + 12*a*d^2*e^2*f + 24
*a*c*d*e*f^2)/(4*d^2) + (a*f^3*(4*c^2 + 4))/(4*d^2)))/d - x^2*((c*((f^2*(
8*a*c*f - b*f + 12*a*d*e))/(4*d) - (2*a*c*f^3)/d))/d - (4*a*f^3 + 4*a*c^2*
f^3 - 4*b*d*e*f^2 + 12*a*d^2*e^2*f + 24*a*c*d*e*f^2)/(8*d^2) + (a*f^3*(4*c
^2 + 4))/(8*d^2) + x^3*((f^2*(8*a*c*f - b*f + 12*a*d*e))/(12*d) - (2*a*c*
f^3)/(3*d)) + (a*f^3*x^4)/4 - (log(c^2 + d^2*x^2 + 2*c*d*x + 1)*(64*b*d^7*
e^3 - 64*b*c^3*d^4*f^3 + 64*b*c*d^4*f^3 - 64*b*d^5*e*f^2 - 192*b*c*d^6*e^2
*f + 192*b*c^2*d^5*e*f^2))/(128*d^8) - (b*atan((4*d^3*((c*(f^3 - 6*c^2*f^3
+ c^4*f^3 - 4*c*d^3*e^3 - 6*d^2*e^2*f + 6*c^2*d^2*e^2*f + 12*c*d*e*f^2 -
4*c^3*d*e*f^2))/(4*d^3) + (x*(f^3 - 6*c^2*f^3 + c^4*f^3 - 4*c*d^3*e^3 - 6*
d^2*e^2*f + 6*c^2*d^2*e^2*f + 12*c*d*e*f^2 - 4*c^3*d*e*f^2))/(4*d^2)))/(f^
3 - 6*c^2*f^3 + c^4*f^3 - 4*c*d^3*e^3 - 6*d^2*e^2*f + 6*c^2*d^2*e^2*f + 12
*c*d*e*f^2 - 4*c^3*d*e*f^2))*(f^3 - 6*c^2*f^3 + c^4*f^3 - 4*c*d^3*e^3 - 6*
d^2*e^2*f + 6*c^2*d^2*e^2*f + 12*c*d*e*f^2 - 4*c^3*d*e*f^2))/(4*d^4)
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 494, normalized size of antiderivative = 2.12

$$\int (e + fx)^3 (a + b \arctan(c + dx)) dx$$

$$= \frac{-3 \operatorname{atan}(dx + c) b f^3 - b d^3 f^3 x^3 - 3 \operatorname{atan}(dx + c) b c^4 f^3 + 18 \operatorname{atan}(dx + c) b c^2 f^3 + 6 \log(d^2 x^2 + 2cdx + c^2) b c^2 f^3 + 6 \log(d^2 x^2 + 2cdx + c^2) b c^2 f^3}{1}$$

input

```
int((f*x+e)^3*(a+b*atan(d*x+c)),x)
```

output

```
( - 3*atan(c + d*x)*b*c**4*f**3 + 12*atan(c + d*x)*b*c**3*d*e*f**2 - 18*at
an(c + d*x)*b*c**2*d**2*e**2*f + 18*atan(c + d*x)*b*c**2*f**3 + 12*atan(c
+ d*x)*b*c*d**3*e**3 - 36*atan(c + d*x)*b*c*d*e*f**2 + 12*atan(c + d*x)*b*
d**4*e**3*x + 18*atan(c + d*x)*b*d**4*e**2*f*x**2 + 12*atan(c + d*x)*b*d**
4*e*f**2*x**3 + 3*atan(c + d*x)*b*d**4*f**3*x**4 + 18*atan(c + d*x)*b*d**2
*e**2*f - 3*atan(c + d*x)*b*f**3 + 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b
*c**3*f**3 - 18*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c**2*d*e*f**2 + 18*1
og(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d**2*e**2*f - 6*log(c**2 + 2*c*d*x
+ d**2*x**2 + 1)*b*c*f**3 - 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d**3*e
**3 + 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d*e*f**2 + 12*a*d**4*e**3*x
+ 18*a*d**4*e**2*f*x**2 + 12*a*d**4*e*f**2*x**3 + 3*a*d**4*f**3*x**4 - 9*b
*c**2*d*f**3*x + 24*b*c*d**2*e*f**2*x + 3*b*c*d**2*f**3*x**2 - 18*b*d**3*e
**2*f*x - 6*b*d**3*e*f**2*x**2 - b*d**3*f**3*x**3 + 3*b*d*f**3*x)/(12*d**4
)
```

3.33 $\int (e + fx)^2 (a + b \arctan(c + dx)) dx$

Optimal result	307
Mathematica [C] (verified)	308
Rubi [A] (verified)	308
Maple [A] (verified)	310
Fricas [A] (verification not implemented)	311
Sympy [C] (verification not implemented)	311
Maxima [A] (verification not implemented)	312
Giac [B] (verification not implemented)	313
Mupad [B] (verification not implemented)	314
Reduce [B] (verification not implemented)	315

Optimal result

Integrand size = 18, antiderivative size = 155

$$\begin{aligned} & \int (e + fx)^2 (a + b \arctan(c + dx)) dx \\ &= -\frac{bf(de - cf)x}{d^2} - \frac{bf^2(c + dx)^2}{6d^3} \\ & \quad - \frac{b(de - cf)(d^2e^2 - 2cdef - (3 - c^2)f^2) \arctan(c + dx)}{3d^3f} \\ & \quad + \frac{(e + fx)^3(a + b \arctan(c + dx))}{3f} \\ & \quad - \frac{b(3d^2e^2 - 6cdef - (1 - 3c^2)f^2) \log(1 + (c + dx)^2)}{6d^3} \end{aligned}$$

output

```
-b*f*(-c*f+d*e)*x/d^2-1/6*b*f^2*(d*x+c)^2/d^3-1/3*b*(-c*f+d*e)*(d^2*e^2-2*
c*d*e*f-(-c^2+3)*f^2)*arctan(d*x+c)/d^3/f+1/3*(f*x+e)^3*(a+b*arctan(d*x+c)
)/f-1/6*b*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*ln(1+(d*x+c)^2)/d^3
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.76

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{(e + fx)^3 (a + b \arctan(c + dx)) - \frac{b(6df^2(de - cf)x + f^3(c + dx)^2 - i(de - (-i + c)f)^3 \log(i - c - dx) + i(de - (i + c)f)^3 \log(i + c + dx))}{2d^3}}{3f}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcTan[c + d*x]),x]
```

output

```
((e + f*x)^3*(a + b*ArcTan[c + d*x]) - (b*(6*d*f^2*(d*e - c*f)*x + f^3*(c + d*x)^2 - I*(d*e - (-I + c)*f)^3*Log[I - c - d*x] + I*(d*e - (I + c)*f)^3*Log[I + c + d*x]))/(2*d^3))/(3*f)
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5570, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$\downarrow \text{5570}$$

$$\int \frac{\left(\frac{d(e - \frac{cf}{d}) + f(c + dx)}{d^2}\right)^2 (a + b \arctan(c + dx))}{d} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(de - cf + f(c + dx))^2 (a + b \arctan(c + dx))}{d^3} d(c + dx)$$

$$\downarrow \text{5387}$$

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))}{3f} - \frac{b \int \frac{(de-cf+f(c+dx))^3 d(c+dx)}{(c+dx)^2+1}}{3f}}{d^3}$$

↓ 478

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))}{3f} - \frac{b \int \left((c+dx)f^3+3(de-cf)f^2+\frac{(de-cf)(d^2e^2-2cdf e+c^2f^2-3f^2)+f(3d^2e^2-6cdf e-(1-3c^2)f^2)(c+dx)}{(c+dx)^2+1} \right) d(c+dx)}{3f}}{d^3}$$

↓ 2009

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))}{3f} - \frac{b(\arctan(c+dx)(de-cf)(-(3-c^2)f^2-2cdf e+d^2e^2)+\frac{1}{2}f(-(1-3c^2)f^2-6cdf e+3d^2e^2)\log((c+dx)))}{3f}}{d^3}$$

```
input Int[(e + f*x)^2*(a + b*ArcTan[c + d*x]),x]
```

```
output (((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTan[c + d*x]))/(3*f) - (b*(3*f^2*(d*e - c*f)*(c + d*x) + (f^3*(c + d*x)^2)/2 + (d*e - c*f)*(d^2*e^2 - 2*c*d*e*f - (3 - c^2)*f^2)*ArcTan[c + d*x] + (f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*Log[1 + (c + d*x)^2])/2))/(3*f))/d^3
```

Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 478 Int[((c_) + (d_.)*(x_))^(n_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5387 Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))*((d_) + (e_.)*(x_.))^(q_.), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(
  c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b
  , c, d, e, q}, x] && NeQ[q, -1]
```

```
rule 5570 Int[((a_.) + ArcTan[(c_) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.71

method	result
parts	$\frac{a(fx+e)^3}{3f} + \frac{bc e^2 \arctan(dx+c)}{d} - \frac{f^2 b x^2}{6d} + \frac{b f^2 \ln(1+(dx+c)^2)}{6d^3} - \frac{f b c^2 e \arctan(dx+c)}{d^2} + \frac{b f \ln(1+(dx+c)^2)}{d^2}$
derivativedivides	$-\frac{a(cf-de-f(dx+c))^3}{3d^2 f} + \frac{b f^2 \arctan(dx+c)c^2(dx+c)}{d^2} - \frac{2bf \arctan(dx+c)ce(dx+c)}{d} - \frac{b f^2 \arctan(dx+c)c(dx+c)^2}{d^2} + b \arctan(dx+c)$
default	$-\frac{a(cf-de-f(dx+c))^3}{3d^2 f} + \frac{b f^2 \arctan(dx+c)c^2(dx+c)}{d^2} - \frac{2bf \arctan(dx+c)ce(dx+c)}{d} - \frac{b f^2 \arctan(dx+c)c(dx+c)^2}{d^2} + b \arctan(dx+c)$
paralelrisch	$-\frac{\ln(d^2 x^2 + 2cdx + c^2 + 1) b f^2 - f^2 b + 6a c^2 e f d + 6e f a d + 7b c^2 f^2 - 12bcdef + 6 \arctan(dx+c) b c^2 d e f - 6 \ln(d^2 x^2 + 2cdx + c^2 + 1) b c^2 d e f}{6d^3}$
risch	$-\frac{i(fx+e)^3 b \ln(1+i(dx+c))}{6f} + \frac{f b c e \ln(d^2 x^2 + 2cdx + c^2 + 1)}{d^2} - \frac{e^2 b \ln(d^2 x^2 + 2cdx + c^2 + 1)}{2d} + \frac{f^2 b \ln(d^2 x^2 + 2cdx + c^2 + 1)}{6d^3}$

```
input int((f*x+e)^2*(a+b*arctan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/3*a*(f*x+e)^3/f+1/d*b*c*e^2*arctan(d*x+c)-1/6*f^2/d*b*x^2+1/6*b/d^3*f^2*
ln(1+(d*x+c)^2)-f/d^2*b*c^2*e*arctan(d*x+c)+b/d^2*f*ln(1+(d*x+c)^2)*c*e+b*
f*arctan(d*x+c)*e*x^2+1/3*b*f^2*arctan(d*x+c)*x^3+b*arctan(d*x+c)*x*e^2+5/
6*b/d^3*f^2*c^2-b/d^2*f*c*e+1/3*f^2/d^3*b*c^3*arctan(d*x+c)+2/3*f^2/d^2*b*
c*x-f/d*b*e*x-1/2*b/d^3*f^2*ln(1+(d*x+c)^2)*c^2-1/2*b*e^2*ln(1+(d*x+c)^2)/
d-f^2/d^3*b*c*arctan(d*x+c)+f/d^2*b*e*arctan(d*x+c)
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.28

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{2ad^3 f^2 x^3 + (6ad^3 ef - bd^2 f^2)x^2 + 2(3ad^3 e^2 - 3bd^2 ef + 2bcd f^2)x + 2(bd^3 f^2 x^3 + 3bd^3 ef x^2 + 3bd^3 e^2 x + 2ad^3 e^2 + 6ad^3 ef - bd^2 f^2)x + 2(3ad^3 e^2 - 3bd^2 ef + 2bcd f^2)x + 2(bd^3 f^2 x^3 + 3bd^3 ef x^2 + 3bd^3 e^2 x + 2ad^3 e^2 + 6ad^3 ef - bd^2 f^2)}{d^3}$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c)),x, algorithm="fricas")`output `1/6*(2*a*d^3*f^2*x^3 + (6*a*d^3*e*f - b*d^2*f^2)*x^2 + 2*(3*a*d^3*e^2 - 3*b*d^2*e*f + 2*b*c*d*f^2)*x + 2*(b*d^3*f^2*x^3 + 3*b*d^3*e*f*x^2 + 3*b*d^3*e^2*x + 3*b*c*d^2*e^2 - 3*(b*c^2 - b)*d*e*f + (b*c^3 - 3*b*c)*f^2)*arctan(d*x + c) - (3*b*d^2*e^2 - 6*b*c*d*e*f + (3*b*c^2 - b)*f^2)*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/d^3`**Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 4.72 (sec) , antiderivative size = 376, normalized size of antiderivative = 2.43

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$= \left\{ \begin{array}{l} ae^2 x + aefx^2 + \frac{af^2 x^3}{3} + \frac{bc^3 f^2 \operatorname{atan}(c+dx)}{3d^3} - \frac{bc^2 ef \operatorname{atan}(c+dx)}{d^2} - \frac{bc^2 f^2 \log\left(\frac{c}{d} + x - \frac{i}{d}\right)}{d^3} + \frac{ibc^2 f^2 \operatorname{atan}(c+dx)}{d^3} + \frac{bce^2 \operatorname{atan}(c+dx)}{d} \\ (a + b \operatorname{atan}(c)) \left(e^2 x + efx^2 + \frac{f^2 x^3}{3} \right) \end{array} \right.$$

input `integrate((f*x+e)**2*(a+b*atan(d*x+c)),x)`

output

```
Piecewise((a**2*x + a*e*f*x**2 + a*f**2*x**3/3 + b*c**3*f**2*atan(c + d*x)/(3*d**3) - b*c**2*e*f*atan(c + d*x)/d**2 - b*c**2*f**2*log(c/d + x - I/d)/d**3 + I*b*c**2*f**2*atan(c + d*x)/d**3 + b*c*e**2*atan(c + d*x)/d + 2*b*c*e*f*log(c/d + x - I/d)/d**2 - 2*I*b*c*e*f*atan(c + d*x)/d**2 + 2*b*c*f**2*x/(3*d**2) - b*c*f**2*atan(c + d*x)/d**3 + b*e**2*x*atan(c + d*x) + b*e*f*x**2*atan(c + d*x) + b*f**2*x**3*atan(c + d*x)/3 - b*e**2*log(c/d + x - I/d)/d + I*b*e**2*atan(c + d*x)/d - b*e*f*x/d - b*f**2*x**2/(6*d) + b*e*f*atan(c + d*x)/d**2 + b*f**2*log(c/d + x - I/d)/(3*d**3) - I*b*f**2*atan(c + d*x)/(3*d**3), Ne(d, 0)), ((a + b*atan(c))*(e**2*x + e*f*x**2 + f**2*x**3/3), True))
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.42

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx = \frac{1}{3} a f^2 x^3 + a e f x^2 + \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2 x + cd}{d}\right) - \frac{c \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^3}}{d^3} \right) \right) b e f + \frac{1}{6} \left(2 x^3 \arctan(dx + c) - d \left(\frac{dx^2 - 4cx}{d^3} - \frac{2(c^3 - 3c) \arctan\left(\frac{d^2 x + cd}{d}\right) + \frac{(3c^2 - 1) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^4}}{d^4} \right) \right) b f^2 + a e^2 x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) b e^2}{2d}$$

input

```
integrate((f*x+e)^2*(a+b*arctan(d*x+c)),x, algorithm="maxima")
```

output

```
1/3*a*f^2*x^3 + a*e*f*x^2 + (x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d)/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3))*b*e*f + 1/6*(2*x^3*arctan(d*x + c) - d*((d*x^2 - 4*c*x)/d^3 - 2*(c^3 - 3*c)*arctan((d^2*x + c*d)/d)/d^4 + (3*c^2 - 1)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^4))*b*f^2 + a*e^2*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b*e^2/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 512 vs. $2(143) = 286$.

Time = 0.45 (sec) , antiderivative size = 512, normalized size of antiderivative = 3.30

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{2bd^3 f^2 x^3 \arctan(dx + c) + 2ad^3 f^2 x^3 + 6bd^3 e f x^2 \arctan(dx + c) + 6ad^3 e f x^2 + 6bd^3 e^2 x \arctan(dx + c) + \dots}{d^3}$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output

```
1/6*(2*b*d^3*f^2*x^3*arctan(d*x + c) + 2*a*d^3*f^2*x^3 + 6*b*d^3*e*f*x^2*arctan(d*x + c) + 6*a*d^3*e*f*x^2 + 6*b*d^3*e^2*x*arctan(d*x + c) - 3*pi*b*c*d^2*e^2*sgn(d*x + c) + 3*pi*b*c^2*d*e*f*sgn(d*x + c) - pi*b*c^3*f^2*sgn(d*x + c) + 3*pi*b*c*d^2*e^2*sgn(-d*x - c) - 3*pi*b*c^2*d*e*f*sgn(-d*x - c) + pi*b*c^3*f^2*sgn(-d*x - c) + 6*a*d^3*e^2*x - b*d^2*f^2*x^2 + 3*b*c*d^2*e^2*arctan(d*x + c) - 3*b*c^2*d*e*f*arctan(d*x + c) + b*c^3*f^2*arctan(d*x + c) - 3*b*c*d^2*e^2*arctan(-d*x - c) + 3*b*c^2*d*e*f*arctan(-d*x - c) - b*c^3*f^2*arctan(-d*x - c) - 6*b*d^2*e*f*x + 4*b*c*d*f^2*x - 3*b*d^2*e^2*log((d*x + c)^2 + 1) + 6*b*c*d*e*f*log((d*x + c)^2 + 1) - 3*b*c^2*f^2*log((d*x + c)^2 + 1) - 3*pi*b*d*e*f*sgn(d*x + c) + 3*pi*b*c*f^2*sgn(d*x + c) + 3*pi*b*d*e*f*sgn(-d*x - c) - 3*pi*b*c*f^2*sgn(-d*x - c) + 3*b*d*e*f*arctan(d*x + c) - 3*b*c*f^2*arctan(d*x + c) - 3*b*d*e*f*arctan(-d*x - c) + 3*b*c*f^2*arctan(-d*x - c) + b*f^2*log((d*x + c)^2 + 1))/d^3
```

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 411, normalized size of antiderivative = 2.65

$$\begin{aligned}
& \int (e + fx)^2 (a + b \arctan(c + dx)) dx \\
&= x^2 \left(\frac{f(6acf - bf + 6ade)}{6d} - \frac{acf^2}{d} \right) - x \left(\frac{2c \left(\frac{f(6acf - bf + 6ade)}{3d} - \frac{2acf^2}{d} \right)}{d} \right. \\
&\quad \left. - \frac{3ac^2 f^2 + 12acdef + 3ad^2 e^2 - 3bdef + 3af^2}{3d^2} + \frac{af^2(3c^2 + 3)}{3d^2} \right) \\
&+ \operatorname{atan}(c + dx) \left(be^2 x + bef x^2 + \frac{bf^2 x^3}{3} \right) + \frac{af^2 x^3}{3} \\
&- \frac{\ln(c^2 + 2cdx + d^2 x^2 + 1) (36b^2 d^3 f^2 - 72bcd^4 ef + 36bd^5 e^2 - 12bd^3 f^2)}{72d^6} \\
&+ \frac{b \operatorname{atan} \left(\frac{3d^2 \left(\frac{c(c^3 f^2 - 3c^2 def + 3cd^2 e^2 - 3cf^2 + 3def)}{3d^2} + \frac{x(c^3 f^2 - 3c^2 def + 3cd^2 e^2 - 3cf^2 + 3def)}{3d} \right)}{c^3 f^2 - 3c^2 def + 3cd^2 e^2 - 3cf^2 + 3def} \right) (c^3 f^2 - 3c^2 def + 3cd^2 e^2 - 3cf^2 + 3def)}{3d^3}
\end{aligned}$$

input `int((e + f*x)^2*(a + b*atan(c + d*x)),x)`

output

```

x^2*((f*(6*a*c*f - b*f + 6*a*d*e))/(6*d) - (a*c*f^2)/d) - x*((2*c*((f*(6*a
*c*f - b*f + 6*a*d*e))/(3*d) - (2*a*c*f^2)/d))/d - (3*a*f^2 + 3*a*c^2*f^2
+ 3*a*d^2*e^2 - 3*b*d*e*f + 12*a*c*d*e*f)/(3*d^2) + (a*f^2*(3*c^2 + 3))/(3
*d^2)) + atan(c + d*x)*((b*f^2*x^3)/3 + b*e^2*x + b*e*f*x^2) + (a*f^2*x^3)
/3 - (log(c^2 + d^2*x^2 + 2*c*d*x + 1)*(36*b*d^5*e^2 - 12*b*d^3*f^2 + 36*b
*c^2*d^3*f^2 - 72*b*c*d^4*e*f))/(72*d^6) + (b*atan((3*d^2*((c*(c^3*f^2 - 3
*c*f^2 + 3*c*d^2*e^2 + 3*d*e*f - 3*c^2*d*e*f))/(3*d^2) + (x*(c^3*f^2 - 3*c
*f^2 + 3*c*d^2*e^2 + 3*d*e*f - 3*c^2*d*e*f))/(3*d)))/(c^3*f^2 - 3*c*f^2 +
3*c*d^2*e^2 + 3*d*e*f - 3*c^2*d*e*f))*(c^3*f^2 - 3*c*f^2 + 3*c*d^2*e^2 + 3
*d*e*f - 3*c^2*d*e*f))/(3*d^3)

```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.89

$$\int (e + fx)^2 (a + b \arctan(c + dx)) dx$$

$$= \frac{2atan(dx + c)bc^3f^2 - 6atan(dx + c)bc^2def + 6atan(dx + c)bcd^2e^2 - 6atan(dx + c)bcf^2 + 6atan(dx + c)bc^2d^2e^2 - 6atan(dx + c)bc^2d^2e^2 + 6atan(dx + c)bc^2d^2e^2}{6d^3}$$

input `int((f*x+e)^2*(a+b*atan(d*x+c)),x)`output `(2*atan(c + d*x)*b*c**3*f**2 - 6*atan(c + d*x)*b*c**2*d*e*f + 6*atan(c + d*x)*b*c*d**2*e**2 - 6*atan(c + d*x)*b*c*f**2 + 6*atan(c + d*x)*b*d**3*e**2*x + 6*atan(c + d*x)*b*d**3*e*f*x**2 + 2*atan(c + d*x)*b*d**3*f**2*x**3 + 6*atan(c + d*x)*b*d*e*f - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c**2*f**2 + 6*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d*e*f - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d**2*e**2 + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*f**2 + 6*a*d**3*e**2*x + 6*a*d**3*e*f*x**2 + 2*a*d**3*f**2*x**3 + 4*b*c*d*f**2*x - 6*b*d**2*e*f*x - b*d**2*f**2*x**2)/(6*d**3)`

3.34 $\int (e + fx)(a + b \arctan(c + dx)) dx$

Optimal result	316
Mathematica [C] (verified)	316
Rubi [A] (verified)	317
Maple [A] (verified)	319
Fricas [A] (verification not implemented)	319
Sympy [C] (verification not implemented)	320
Maxima [A] (verification not implemented)	320
Giac [B] (verification not implemented)	321
Mupad [B] (verification not implemented)	322
Reduce [B] (verification not implemented)	322

Optimal result

Integrand size = 16, antiderivative size = 94

$$\int (e + fx)(a + b \arctan(c + dx)) dx = -\frac{bf x}{2d} + \frac{b(f^2 - (de - cf)^2) \arctan(c + dx)}{2d^2 f} + \frac{(e + fx)^2(a + b \arctan(c + dx))}{2f} - \frac{b(de - cf) \log(1 + (c + dx)^2)}{2d^2}$$

output

```
-1/2*b*f*x/d+1/2*b*(f^2-(-c*f+d*e)^2)*arctan(d*x+c)/d^2/f+1/2*(f*x+e)^2*(a+b*arctan(d*x+c))/f-1/2*b*(-c*f+d*e)*ln(1+(d*x+c)^2)/d^2
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.73

$$\int (e + fx)(a + b \arctan(c + dx)) dx = aex + \frac{1}{2}afx^2 + bex \arctan(c + dx) + \frac{bf \left(\frac{1}{2}d \left(-\frac{c}{d} + \frac{c+dx}{d} \right)^2 \arctan(c + dx) - \frac{1}{2}d \left(\frac{x}{d} - \frac{i(i-c)^2 \log(i-c-dx)}{2d^2} + \frac{i(i+c)^2 \log(i+c+dx)}{2d^2} \right) \right)}{d} - \frac{be(-2c \arctan(c + dx) + \log(1 + c^2 + 2cdx + d^2x^2))}{2d}$$

input `Integrate[(e + f*x)*(a + b*ArcTan[c + d*x]),x]`

output `a*e*x + (a*f*x^2)/2 + b*e*x*ArcTan[c + d*x] + (b*f*((d*(-(c/d) + (c + d*x)/d)^2*ArcTan[c + d*x])/2 - (d*(x/d - ((I/2)*(I - c)^2*Log[I - c - d*x])/d^2 + ((I/2)*(I + c)^2*Log[I + c + d*x])/d^2))/2))/d - (b*e*(-2*c*ArcTan[c + d*x] + Log[1 + c^2 + 2*c*d*x + d^2*x^2]))/(2*d)`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {5570, 27, 5387, 478, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)(a + b \arctan(c + dx)) dx \\
 & \quad \downarrow \text{5570} \\
 & \int \frac{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)(a + b \arctan(c + dx))}{d} d(c + dx) \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(de - cf + f(c + dx))(a + b \arctan(c + dx))d(c + dx)}{d^2} \\
 & \quad \downarrow \text{5387} \\
 & \frac{\frac{(f(c + dx) - cf + de)^2(a + b \arctan(c + dx))}{2f} - \frac{b \int \frac{(de - cf + f(c + dx))^2 d(c + dx)}{(c + dx)^2 + 1}}{2f}}{d^2} \\
 & \quad \downarrow \text{478} \\
 & \frac{\frac{(f(c + dx) - cf + de)^2(a + b \arctan(c + dx))}{2f} - \frac{b \int \left(f^2 + \frac{(de - cf - f)(de - cf + f) + 2f(de - cf)(c + dx)}{(c + dx)^2 + 1}\right) d(c + dx)}{2f}}{d^2} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{\frac{(f(c+dx)-cf+de)^2(a+b\arctan(c+dx))}{2f} - \frac{b(\arctan(c+dx)(-cf+de+f)(de-(c+1)f)+f(de-cf)\log((c+dx)^2+1)+f^2(c+dx))}{2f}}{d^2}$$

input `Int[(e + f*x)*(a + b*ArcTan[c + d*x]),x]`

output `((((d*e - c*f + f*(c + d*x))^2*(a + b*ArcTan[c + d*x]))/(2*f) - (b*(f^2*(c + d*x) + (d*e + f - c*f)*(d*e - (1 + c)*f)*ArcTan[c + d*x] + f*(d*e - c*f)*Log[1 + (c + d*x)^2]))/(2*f))/d^2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 478 `Int[((c_) + (d_)*(x_))^(n_)/((a_) + (b_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[n, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5387 `Int[((a_) + ArcTan[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 5570 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.20

method	result
parts	$a\left(\frac{1}{2}f x^2 + ex\right) + \frac{b\left(\frac{\arctan(dx+c)(dx+c)^2 f}{2d} - \frac{\arctan(dx+c)cf(dx+c)}{d} + \arctan(dx+c)e(dx+c) - \frac{f(dx+c) + \frac{(-2cf+2de)}{d}}{d}\right)}{d}$
derivativedivides	$\frac{a\left(f c(dx+c) - ed(dx+c) - \frac{f(dx+c)^2}{2}\right)}{d} - \frac{b\left(\arctan(dx+c)fc(dx+c) - \arctan(dx+c)ed(dx+c) - \frac{\arctan(dx+c)f(dx+c)^2}{2} + \frac{f(dx+c)}{2}\right)}{d}$
default	$\frac{a\left(f c(dx+c) - ed(dx+c) - \frac{f(dx+c)^2}{2}\right)}{d} - \frac{b\left(\arctan(dx+c)fc(dx+c) - \arctan(dx+c)ed(dx+c) - \frac{\arctan(dx+c)f(dx+c)^2}{2} + \frac{f(dx+c)}{2}\right)}{d}$
parallelrisch	$\frac{\arctan(dx+c)b d^2 f x^2 + a d^2 f x^2 + 2x \arctan(dx+c)b d^2 e + 2a d^2 ex - \arctan(dx+c)b c^2 f + 2bcde \arctan(dx+c) + bcf \ln(dx+c)}{2d^2}$
risch	$-\frac{ib(f x^2 + 2ex) \ln(1+i(dx+c))}{4} + \frac{ibf x^2 \ln(1-i(dx+c))}{4} + \frac{ibex \ln(1-i(dx+c))}{2} + \frac{af x^2}{2} - \frac{\arctan(dx+c)b c^2 f}{2d^2}$

input `int((f*x+e)*(a+b*arctan(d*x+c)),x,method=_RETURNVERBOSE)`

output `a*(1/2*f*x^2+e*x)+b/d*(1/2/d*arctan(d*x+c)*(d*x+c)^2*f-1/d*arctan(d*x+c)*c*f*(d*x+c)+arctan(d*x+c)*e*(d*x+c)-1/2/d*(f*(d*x+c)+1/2*(-2*c*f+2*d*e)*ln(1+(d*x+c)^2)-f*arctan(d*x+c)))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.10

$$\int (e + fx)(a + b \arctan(c + dx)) dx = \frac{ad^2fx^2 + (2ad^2e - bdf)x + (bd^2fx^2 + 2bd^2ex + 2bcde - (bc^2 - b)f) \arctan(dx + c) - (bde - bcf) \log}{2d^2}$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c)),x, algorithm="fricas")`

output

```
1/2*(a*d^2*f*x^2 + (2*a*d^2*e - b*d*f)*x + (b*d^2*f*x^2 + 2*b*d^2*e*x + 2*
b*c*d*e - (b*c^2 - b)*f)*arctan(d*x + c) - (b*d*e - b*c*f)*log(d^2*x^2 + 2
*c*d*x + c^2 + 1))/d^2
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.94 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.88

$$\int (e + fx)(a + b \arctan(c + dx)) dx$$

$$= \begin{cases} aex + \frac{afx^2}{2} - \frac{bc^2 f \operatorname{atan}(c+dx)}{2d^2} + \frac{bce \operatorname{atan}(c+dx)}{d} + \frac{bcf \log\left(\frac{c}{d} + x - \frac{i}{d}\right)}{d^2} - \frac{ibcf \operatorname{atan}(c+dx)}{d^2} + bex \operatorname{atan}(c + dx) + \frac{bf x^2 \operatorname{atan}(c + dx)}{2d} \\ (a + b \operatorname{atan}(c)) \left(ex + \frac{fx^2}{2} \right) \end{cases}$$

input

```
integrate((f*x+e)*(a+b*atan(d*x+c)),x)
```

output

```
Piecewise((a*e*x + a*f*x**2/2 - b*c**2*f*atan(c + d*x)/(2*d**2) + b*c*e*at
an(c + d*x)/d + b*c*f*log(c/d + x - I/d)/d**2 - I*b*c*f*atan(c + d*x)/d**2
+ b*e*x*atan(c + d*x) + b*f*x**2*atan(c + d*x)/2 - b*e*log(c/d + x - I/d)
/d + I*b*e*atan(c + d*x)/d - b*f*x/(2*d) + b*f*atan(c + d*x)/(2*d**2), Ne(
d, 0)), ((a + b*atan(c))*(e*x + f*x**2/2), True))
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.23

$$\int (e + fx)(a + b \arctan(c + dx)) dx = \frac{1}{2} a f x^2$$

$$+ \frac{1}{2} \left(x^2 \arctan(dx + c) - d \left(\frac{x}{d^2} + \frac{(c^2 - 1) \arctan\left(\frac{d^2 x + cd}{d}\right)}{d^3} - \frac{c \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^3} \right) \right) b f$$

$$+ a e x + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1)) b e}{2d}$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output
$$\frac{1}{2}a*f*x^2 + \frac{1}{2}*(x^2*arctan(d*x + c) - d*(x/d^2 + (c^2 - 1)*arctan((d^2*x + c*d)/d))/d^3 - c*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/d^3)*b*f + a*e*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b*e/d$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 250 vs. $2(86) = 172$.

Time = 0.20 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.66

$$\int (e + fx)(a + b \arctan(c + dx)) dx$$

$$= \frac{2bd^2fx^2 \arctan(dx + c) + 2ad^2fx^2 + 4bd^2ex \arctan(dx + c) - 2\pi bc d \operatorname{sgn}(dx + c) + \pi bc^2 f \operatorname{sgn}(dx + c)}{d^2}$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output
$$\frac{1}{4}*(2*b*d^2*f*x^2*arctan(d*x + c) + 2*a*d^2*f*x^2 + 4*b*d^2*e*x*arctan(d*x + c) - 2*pi*b*c*d*e*sgn(d*x + c) + pi*b*c^2*f*sgn(d*x + c) + 2*pi*b*c*d*e*sgn(-d*x - c) - pi*b*c^2*f*sgn(-d*x - c) + 4*a*d^2*e*x + 2*b*c*d*e*arctan(d*x + c) - b*c^2*f*arctan(d*x + c) - 2*b*c*d*e*arctan(-d*x - c) + b*c^2*f*arctan(-d*x - c) - 2*b*d*f*x - 2*b*d*e*log((d*x + c)^2 + 1) + 2*b*c*f*log((d*x + c)^2 + 1) - pi*b*f*sgn(d*x + c) + pi*b*f*sgn(-d*x - c) + b*f*arctan(d*x + c) - b*f*arctan(-d*x - c))/d^2$$

Mupad [B] (verification not implemented)

Time = 1.61 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.45

$$\int (e + fx)(a + b \arctan(c + dx)) dx = a e x + \frac{a f x^2}{2} - \frac{b e \ln(c^2 + 2 c d x + d^2 x^2 + 1)}{2 d} + \frac{b f \operatorname{atan}(c + dx)}{2 d^2} + \frac{b f x^2 \operatorname{atan}(c + dx)}{2} - \frac{b f x}{2 d} + b e x \operatorname{atan}(c + dx) - \frac{b c^2 f \operatorname{atan}(c + dx)}{2 d^2} + \frac{b c f \ln(c^2 + 2 c d x + d^2 x^2 + 1)}{2 d^2} + \frac{b c e \operatorname{atan}(c + dx)}{d}$$

input `int((e + f*x)*(a + b*atan(c + d*x)),x)`output `a*e*x + (a*f*x^2)/2 - (b*e*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(2*d) + (b*f*atan(c + d*x))/(2*d^2) + (b*f*x^2*atan(c + d*x))/2 - (b*f*x)/(2*d) + b*e*x*atan(c + d*x) - (b*c^2*f*atan(c + d*x))/(2*d^2) + (b*c*f*log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(2*d^2) + (b*c*e*atan(c + d*x))/d`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.46

$$\int (e + fx)(a + b \arctan(c + dx)) dx = \frac{-\operatorname{atan}(dx + c) b c^2 f + 2 \operatorname{atan}(dx + c) b c d e + 2 \operatorname{atan}(dx + c) b d^2 e x + \operatorname{atan}(dx + c) b d^2 f x^2 + \operatorname{atan}(dx + c) b d^2 f x^2}{2 d^2}$$

input `int((f*x+e)*(a+b*atan(d*x+c)),x)`output `(- atan(c + d*x)*b*c**2*f + 2*atan(c + d*x)*b*c*d*e + 2*atan(c + d*x)*b*d**2*e*x + atan(c + d*x)*b*d**2*f*x**2 + atan(c + d*x)*b*f + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*f - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d*e + 2*a*d**2*e*x + a*d**2*f*x**2 - b*d*f*x)/(2*d**2)`

3.35 $\int (a + b \arctan(c + dx)) dx$

Optimal result	323
Mathematica [A] (verified)	323
Rubi [A] (verified)	324
Maple [A] (verified)	325
Fricas [A] (verification not implemented)	325
Sympy [A] (verification not implemented)	326
Maxima [A] (verification not implemented)	326
Giac [A] (verification not implemented)	326
Mupad [B] (verification not implemented)	327
Reduce [B] (verification not implemented)	327

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int (a + b \arctan(c + dx)) dx = ax + \frac{b(c + dx) \arctan(c + dx)}{d} - \frac{b \log(1 + (c + dx)^2)}{2d}$$

output `a*x+b*(d*x+c)*arctan(d*x+c)/d-1/2*b*ln(1+(d*x+c)^2)/d`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int (a + b \arctan(c + dx)) dx = ax + bx \arctan(c + dx) - \frac{b(-2c \arctan(c + dx) + \log(1 + c^2 + 2cdx + d^2x^2))}{2d}$$

input `Integrate[a + b*ArcTan[c + d*x],x]`

output `a*x + b*x*ArcTan[c + d*x] - (b*(-2*c*ArcTan[c + d*x] + Log[1 + c^2 + 2*c*d*x + d^2*x^2]))/(2*d)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \arctan(c + dx)) dx$$

$$\downarrow \text{2009}$$

$$ax + \frac{b(c + dx) \arctan(c + dx)}{d} - \frac{b \log((c + dx)^2 + 1)}{2d}$$

input `Int[a + b*ArcTan[c + d*x],x]`

output `a*x + (b*(c + d*x)*ArcTan[c + d*x])/d - (b*Log[1 + (c + d*x)^2])/(2*d)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

method	result	size
default	$ax + \frac{b \left((dx+c) \arctan(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{d}$	35
parts	$ax + \frac{b \left((dx+c) \arctan(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{d}$	35
derivativedivides	$\frac{(dx+c)a + b \left((dx+c) \arctan(dx+c) - \frac{\ln(1+(dx+c)^2)}{2} \right)}{d}$	40
parallelrisc	$-\frac{b(-2x \arctan(dx+c)d^2 - 2c \arctan(dx+c)d + \ln(d^2x^2 + 2cdx + c^2 + 1)d)}{2d^2} + ax$	54
risc	$ax - \frac{ibx \ln(1+i(dx+c))}{2} + \frac{ibx \ln(1-i(dx+c))}{2} + \frac{bc \arctan(dx+c)}{d} - \frac{b \ln(d^2x^2 + 2cdx + c^2 + 1)}{2d}$	73

input `int(a+b*arctan(d*x+c),x,method=_RETURNVERBOSE)`output `a*x+b/d*((d*x+c)*arctan(d*x+c)-1/2*ln(1+(d*x+c)^2))`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int (a + b \arctan(c + dx)) dx$$

$$= \frac{2adx + 2(bdx + bc) \arctan(dx + c) - b \log(d^2x^2 + 2cdx + c^2 + 1)}{2d}$$

input `integrate(a+b*arctan(d*x+c),x, algorithm="fricas")`output `1/2*(2*a*d*x + 2*(b*d*x + b*c)*arctan(d*x + c) - b*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/d`

Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int (a + b \arctan(c + dx)) dx$$

$$= ax + b \begin{cases} \frac{c \operatorname{atan}(c+dx)}{d} + x \operatorname{atan}(c + dx) - \frac{\log(c^2 + 2cdx + d^2x^2 + 1)}{2d} & \text{for } d \neq 0 \\ x \operatorname{atan}(c) & \text{otherwise} \end{cases}$$

input `integrate(a+b*atan(d*x+c),x)`output `a*x + b*Piecewise((c*atan(c + d*x)/d + x*atan(c + d*x) - log(c**2 + 2*c*d*x + d**2*x**2 + 1)/(2*d), Ne(d, 0)), (x*atan(c), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int (a + b \arctan(c + dx)) dx = ax + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1))b}{2d}$$

input `integrate(a+b*arctan(d*x+c),x, algorithm="maxima")`output `a*x + 1/2*(2*(d*x + c)*arctan(d*x + c) - log((d*x + c)^2 + 1))*b/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int (a + b \arctan(c + dx)) dx = ax + \frac{(2(dx + c) \arctan(dx + c) - \log((dx + c)^2 + 1))b}{2d}$$

input `integrate(a+b*arctan(d*x+c),x, algorithm="giac")`

output $a*x + 1/2*(2*(d*x + c)*\arctan(d*x + c) - \log((d*x + c)^2 + 1))*b/d$

Mupad [B] (verification not implemented)

Time = 1.54 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int (a + b \arctan(c + dx)) dx = ax + bx \operatorname{atan}(c + dx) - \frac{b \ln(c^2 + 2cdx + d^2x^2 + 1)}{2d} + \frac{bc \operatorname{atan}(c + dx)}{d}$$

input $\operatorname{int}(a + b*\operatorname{atan}(c + d*x), x)$

output $a*x + b*x*\operatorname{atan}(c + d*x) - (b*\log(c^2 + d^2*x^2 + 2*c*d*x + 1))/(2*d) + (b*c*\operatorname{atan}(c + d*x))/d$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int (a + b \arctan(c + dx)) dx = \frac{2 \operatorname{atan}(dx + c) bc + 2 \operatorname{atan}(dx + c) bdx - \log(d^2x^2 + 2cdx + c^2 + 1) b + 2adx}{2d}$$

input $\operatorname{int}(a+b*\operatorname{atan}(d*x+c), x)$

output $(2*\operatorname{atan}(c + d*x)*b*c + 2*\operatorname{atan}(c + d*x)*b*d*x - \log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b + 2*a*d*x)/(2*d)$

3.36 $\int \frac{a+b \arctan(c+dx)}{e+fx} dx$

Optimal result	328
Mathematica [A] (verified)	329
Rubi [A] (verified)	329
Maple [A] (verified)	332
Fricas [F]	332
Sympy [F(-1)]	333
Maxima [F]	333
Giac [F]	333
Mupad [F(-1)]	334
Reduce [F]	334

Optimal result

Integrand size = 18, antiderivative size = 162

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = -\frac{(a + b \arctan(c + dx)) \log\left(\frac{2}{1 - i(c + dx)}\right)}{f} + \frac{(a + b \arctan(c + dx)) \log\left(\frac{2d(e + fx)}{(de + (i - c)f)(1 - i(c + dx))}\right)}{f} + \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i(c + dx)}\right)}{2f} - \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2d(e + fx)}{(de + (i - c)f)(1 - i(c + dx))}\right)}{2f}$$

output

```
-(a+b*arctan(d*x+c))*ln(2/(1-I*(d*x+c)))/f+(a+b*arctan(d*x+c))*ln(2*d*(f*x
+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f+1/2*I*b*polylog(2,1-2/(1-I*(d*x+c)))/f-
1/2*I*b*polylog(2,1-2*d*(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.99

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx$$

$$= \frac{2a \log(d(e + fx)) + ib \log\left(\frac{d(e+fx)}{de-(i+c)f}\right) \log(1 - i(c + dx)) - ib \log\left(\frac{d(e+fx)}{de+if-cf}\right) \log(1 + i(c + dx)) - ib \text{PolyLog}[2, (f(-I + c + dx))/(-(d*e) + (-I + c)*f)] + I*b*\text{PolyLog}[2, (f*(I + c + dx))/(-(d*e) + (I + c)*f)]}{2f}$$

input `Integrate[(a + b*ArcTan[c + d*x])/(e + f*x), x]`

output `(2*a*Log[d*(e + f*x)] + I*b*Log[(d*(e + f*x))/(d*e - (I + c)*f])*Log[1 - I*(c + d*x)] - I*b*Log[(d*(e + f*x))/(d*e + I*f - c*f])*Log[1 + I*(c + d*x)] - I*b*PolyLog[2, (f*(-I + c + d*x))/(-(d*e) + (-I + c)*f)] + I*b*PolyLog[2, (f*(I + c + d*x))/(-(d*e) + (I + c)*f)]/(2*f)`

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5570, 27, 5381, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx$$

$$\downarrow \text{5570}$$

$$\int \frac{d(a + b \arctan(c + dx))}{d\left(e - \frac{cf}{d}\right) + f(c + dx)} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{a + b \arctan(c + dx)}{f(c + dx) - cf + de} d(c + dx)$$

$$\downarrow \text{5381}$$

$$\frac{-\frac{b \int \frac{\log\left(\frac{2(de-cf+f(c+dx))}{(de-cf+if)(1-i(c+dx))}\right)}{(c+dx)^2+1} d(c+dx)}{f} + \frac{b \int \frac{\log\left(\frac{2}{1-i(c+dx)}\right)}{(c+dx)^2+1} d(c+dx)}{f} + (a + b \arctan(c + dx)) \log\left(\frac{2(f(c+dx)-cf+de)}{(1-i(c+dx))(-cf+de+if)}\right)}{f} - \frac{\log\left(\frac{2}{1-i(c+dx)}\right) (a + b \arctan(c + dx))}{f}}$$

↓ 2849

$$\frac{-\frac{b \int \frac{\log\left(\frac{2(de-cf+f(c+dx))}{(de-cf+if)(1-i(c+dx))}\right)}{(c+dx)^2+1} d(c+dx)}{f} + \frac{ib \int \frac{\log\left(\frac{2}{1-i(c+dx)}\right)}{1-i(c+dx)} d\frac{1}{1-i(c+dx)}}{f} + (a + b \arctan(c + dx)) \log\left(\frac{2(f(c+dx)-cf+de)}{(1-i(c+dx))(-cf+de+if)}\right)}{f} - \frac{\log\left(\frac{2}{1-i(c+dx)}\right) (a + b \arctan(c + dx))}{f}}$$

↓ 2752

$$\frac{-\frac{b \int \frac{\log\left(\frac{2(de-cf+f(c+dx))}{(de-cf+if)(1-i(c+dx))}\right)}{(c+dx)^2+1} d(c+dx)}{f} + \frac{(a + b \arctan(c + dx)) \log\left(\frac{2(f(c+dx)-cf+de)}{(1-i(c+dx))(-cf+de+if)}\right)}{f} - \frac{\log\left(\frac{2}{1-i(c+dx)}\right) (a + b \arctan(c + dx))}{f} + \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(c+dx)}\right)}{2f}}$$

↓ 2897

$$\frac{(a + b \arctan(c + dx)) \log\left(\frac{2(f(c+dx)-cf+de)}{(1-i(c+dx))(-cf+de+if)}\right)}{f} - \frac{\log\left(\frac{2}{1-i(c+dx)}\right) (a + b \arctan(c + dx))}{f} - \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2(de-cf+f(c+dx))}{(de-cf+if)(1-i(c+dx))}\right)}{2f} + \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(c+dx)}\right)}{2f}$$

input `Int[(a + b*ArcTan[c + d*x])/(e + f*x),x]`

output `-(((a + b*ArcTan[c + d*x])*Log[2/(1 - I*(c + d*x))])/f) + ((a + b*ArcTan[c + d*x])*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/f + ((I/2)*b*PolyLog[2, 1 - 2/(1 - I*(c + d*x))])/f - ((I/2)*b*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/f`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_.)}, x_Symbol] \rightarrow \text{With}\{C = \text{FullSimplify}[Pq^{m*}((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]$
- rule 5381 $\text{Int}[(a_.) + \text{ArcTan}[(c_*)(x_)]*(b_.)/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTan}[c*x])*(\text{Log}[2/(1 - I*c*x)]/e), x] + (\text{Simp}[(a + b*\text{ArcTan}[c*x])*(\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] + \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))]/(1 + c^2*x^2), x], x]) /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[c^2*d^2 + e^2, 0]$
- rule 5570 $\text{Int}[(a_.) + \text{ArcTan}[(c_) + (d_*)(x_)]*(b_.)^{(p_.)*((e_.) + (f_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ \text{IgtQ}[p, 0]$

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.22

method	result
parts	$\frac{a \ln(fx+e)}{f} + \frac{b \left(\frac{d \ln(f(dx+c)-cf+de)}{f} \arctan(dx+c) - d \left(-\frac{i \ln(f(dx+c)-cf+de) \left(\ln\left(\frac{if-f(dx+c)}{-cf+de+if}\right) - \ln\left(\frac{if+f(dx+c)}{cf-de+if}\right) \right)}{2f} \right)}{d}$
derivativedivides	$\frac{ad \ln(cf-de-f(dx+c))}{f} - bd \left(-\frac{\ln(cf-de-f(dx+c))}{f} \arctan(dx+c) + \frac{i \ln(cf-de-f(dx+c)) \left(\ln\left(\frac{if+f(dx+c)}{cf-de+if}\right) - \ln\left(\frac{if-f(dx+c)}{-cf+de+if}\right) \right)}{2f} \right)$
default	$\frac{ad \ln(cf-de-f(dx+c))}{f} - bd \left(-\frac{\ln(cf-de-f(dx+c))}{f} \arctan(dx+c) + \frac{i \ln(cf-de-f(dx+c)) \left(\ln\left(\frac{if+f(dx+c)}{cf-de+if}\right) - \ln\left(\frac{if-f(dx+c)}{-cf+de+if}\right) \right)}{2f} \right)$
risch	$\frac{a \ln(icf-ide+(-idx-ic+1)f-f)}{f} + \frac{ib \operatorname{dilog}\left(\frac{icf-ide+(-idx-ic+1)f-f}{icf-ide-f}\right)}{2f} + \frac{ib \ln(-idx-ic+1) \ln\left(\frac{icf-ide+(-idx-ic+1)f-f}{icf-ide-f}\right)}{2f}$

input `int((a+b*arctan(d*x+c))/(f*x+e),x,method=_RETURNVERBOSE)`

output `a*ln(f*x+e)/f+b/d*(d*ln(f*(d*x+c)-c*f+d*e)/f*arctan(d*x+c)-d*(-1/2*I*ln(f*(d*x+c)-c*f+d*e)*(ln((I*f-f*(d*x+c))/(d*e+I*f-c*f))-ln((I*f+f*(d*x+c))/(c*f-d*e+I*f)))/f-1/2*I*(dilog((I*f-f*(d*x+c))/(d*e+I*f-c*f))-dilog((I*f+f*(d*x+c))/(c*f-d*e+I*f)))/f)`

Fricas [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \int \frac{b \arctan(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e),x, algorithm="fricas")`

output `integral((b*arctan(d*x + c) + a)/(f*x + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))/(f*x+e),x)`output `Timed out`**Maxima [F]**

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \int \frac{b \arctan(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e),x, algorithm="maxima")`output `2*b*integrate(1/2*arctan(d*x + c)/(f*x + e), x) + a*log(f*x + e)/f`**Giac [F]**

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \int \frac{b \arctan(dx + c) + a}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e),x, algorithm="giac")`output `integrate((b*arctan(d*x + c) + a)/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \int \frac{a + b \operatorname{atan}(c + dx)}{e + fx} dx$$

input `int((a + b*atan(c + d*x))/(e + f*x),x)`output `int((a + b*atan(c + d*x))/(e + f*x), x)`**Reduce [F]**

$$\int \frac{a + b \arctan(c + dx)}{e + fx} dx = \frac{\left(\int \frac{\operatorname{atan}(dx+c)}{fx+e} dx \right) bf + \log(fx + e) a}{f}$$

input `int((a+b*atan(d*x+c))/(f*x+e),x)`output `(int(atan(c + d*x)/(e + f*x),x)*b*f + log(e + f*x)*a)/f`

3.37 $\int \frac{a+b \arctan(c+dx)}{(e+fx)^2} dx$

Optimal result	335
Mathematica [C] (verified)	335
Rubi [A] (verified)	336
Maple [A] (verified)	339
Fricas [A] (verification not implemented)	340
Sympy [F(-1)]	340
Maxima [A] (verification not implemented)	341
Giac [B] (verification not implemented)	341
Mupad [B] (verification not implemented)	342
Reduce [B] (verification not implemented)	343

Optimal result

Integrand size = 18, antiderivative size = 151

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \frac{bd(de - cf) \arctan(c + dx)}{f(d^2e^2 - 2cdef + (1 + c^2)f^2)} - \frac{a + b \arctan(c + dx)}{f(e + fx)} + \frac{bd \log(e + fx)}{d^2e^2 - 2cdef + (1 + c^2)f^2} - \frac{bd \log(1 + c^2 + 2cdx + d^2x^2)}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)}$$

output

```
b*d*(-c*f+d*e)*arctan(d*x+c)/f/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-(a+b*arctan
(d*x+c))/f/(f*x+e)+b*d*ln(f*x+e)/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-b*d*ln(d^
2*x^2+2*c*d*x+c^2+1)/(2*d^2*e^2-4*c*d*e*f+2*(c^2+1)*f^2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.80

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \frac{-\frac{a+b \arctan(c+dx)}{e+fx} + \frac{bd(i(-de+(i+c)f) \log(i-c-dx)+i(de+if-cf) \log(i+c+dx)+2f \log(d(e+fx)))}{2(d^2e^2-2cdef+(1+c^2)f^2)}}{f}$$

input `Integrate[(a + b*ArcTan[c + d*x])/(e + f*x)^2,x]`

output `((-(a + b*ArcTan[c + d*x])/(e + f*x)) + (b*d*(I*(-(d*e) + (I + c)*f)*Log[I - c - d*x] + I*(d*e + I*f - c*f)*Log[I + c + d*x] + 2*f*Log[d*(e + f*x)]))/(2*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2))/f`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5568, 2081, 1144, 27, 1142, 27, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx \\
 & \quad \downarrow 5568 \\
 & \frac{bd \int \frac{1}{(e+fx)((c+dx)^2+1)} dx}{f} - \frac{a + b \arctan(c + dx)}{f(e + fx)} \\
 & \quad \downarrow 2081 \\
 & \frac{bd \int \frac{1}{(e+fx)(c^2+2dxc+d^2x^2+1)} dx}{f} - \frac{a + b \arctan(c + dx)}{f(e + fx)} \\
 & \quad \downarrow 1144 \\
 & \frac{bd \left(\frac{\int \frac{d(de-2cf-dfx)}{c^2+2dxc+d^2x^2+1} dx}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{f} - \frac{a + b \arctan(c + dx)}{f(e + fx)} \\
 & \quad \downarrow 27 \\
 & \frac{bd \left(\frac{d \int \frac{de-2cf-dfx}{c^2+2dxc+d^2x^2+1} dx}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{f} - \frac{a + b \arctan(c + dx)}{f(e + fx)} \\
 & \quad \downarrow 1142
 \end{aligned}$$

$$\begin{array}{c}
\frac{bd \left(\frac{d \left((de-cf) \int \frac{1}{c^2+2dxc+d^2x^2+1} dx - \frac{f \int \frac{2d(c+dx)}{c^2+2dxc+d^2x^2+1} dx}{2d} \right)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{\frac{f}{a+b \arctan(c+dx)} \frac{1}{f(e+fx)}} \\
\downarrow 27 \\
\frac{bd \left(\frac{d \left((de-cf) \int \frac{1}{c^2+2dxc+d^2x^2+1} dx - f \int \frac{c+dx}{c^2+2dxc+d^2x^2+1} dx \right)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{\frac{f}{a+b \arctan(c+dx)} \frac{1}{f(e+fx)}} \\
\downarrow 1083 \\
\frac{bd \left(\frac{d \left(-f \int \frac{c+dx}{c^2+2dxc+d^2x^2+1} dx - 2(de-cf) \int \frac{1}{-4d^2-(2xd^2+2cd)^2} d(2xd^2+2cd) \right)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{\frac{f}{a+b \arctan(c+dx)} \frac{1}{f(e+fx)}} \\
\downarrow 217 \\
\frac{bd \left(\frac{d \left(\frac{\arctan\left(\frac{2cd+2d^2x}{2d}\right)(de-cf)}{d} - f \int \frac{c+dx}{c^2+2dxc+d^2x^2+1} dx \right)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{\frac{f}{a+b \arctan(c+dx)} \frac{1}{f(e+fx)}} \\
\downarrow 1103 \\
\frac{bd \left(\frac{d \left(\frac{\arctan\left(\frac{2cd+2d^2x}{2d}\right)(de-cf)}{d} - \frac{f \log(c^2+2cdx+d^2x^2+1)}{2d} \right)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{f \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{\frac{f}{a+b \arctan(c+dx)} \frac{1}{f(e+fx)}}
\end{array}$$

input $\text{Int}[(a + b \cdot \text{ArcTan}[c + d \cdot x]) / (e + f \cdot x)^2, x]$

output
$$-\frac{(a + b \cdot \text{ArcTan}[c + d \cdot x]) / (f \cdot (e + f \cdot x)) + (b \cdot d \cdot ((f \cdot \text{Log}[e + f \cdot x]) / (d^2 \cdot e^2 - 2 \cdot c \cdot d \cdot e \cdot f + (1 + c^2) \cdot f^2)) + (d \cdot ((d \cdot e - c \cdot f) \cdot \text{ArcTan}[(2 \cdot c \cdot d + 2 \cdot d^2 \cdot x) / (2 \cdot d)])) / d - (f \cdot \text{Log}[1 + c^2 + 2 \cdot c \cdot d \cdot x + d^2 \cdot x^2]) / (2 \cdot d)) / (d^2 \cdot e^2 - 2 \cdot c \cdot d \cdot e \cdot f + (1 + c^2) \cdot f^2))}{f}$$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)(F x_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F x, (b_)(G x_)] /; \text{FreeQ}[b, x]$

rule 217 $\text{Int}[((a_)+(b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x / \text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1083 $\text{Int}[((a_)+(b_)(x_)+(c_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[((d_)+(e_)(x_))/((a_)+(b_)(x_)+(c_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142 $\text{Int}[((d_)+(e_)(x_))/((a_)+(b_)(x_)+(c_)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e) / (2 \cdot c) \quad \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e / (2 \cdot c) \quad \text{Int}[(b + 2 \cdot c \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1144 $\text{Int}[1/(((d_)+(e_)(x_)) \cdot ((a_)+(b_)(x_)+(c_)(x_)^2)), x_Symbol] \rightarrow \text{Simp}[e \cdot (\text{Log}[\text{RemoveContent}[d + e \cdot x, x]] / (c \cdot d^2 - b \cdot d \cdot e + a \cdot e^2)), x] + \text{Simp}[1 / (c \cdot d^2 - b \cdot d \cdot e + a \cdot e^2) \quad \text{Int}[(c \cdot d - b \cdot e - c \cdot e \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

```
rule 2081 Int[(u_)^(m_)*(v_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^m*ExpandToSum
[v, x]^p, x] /; FreeQ[{m, p}, x] && LinearQ[u, x] && QuadraticQ[v, x] && !
(LinearMatchQ[u, x] && QuadraticMatchQ[v, x])
```

```
rule 5568 Int[((a_) + ArcTan[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m
_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m +
1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[
c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && IGtQ[p, 0] && ILtQ[m, -1]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.06

method	result
parts	$-\frac{a}{(fx+e)f} + \frac{b \left(-\frac{d^2 \arctan(dx+c)}{(f(dx+c)-cf+de)f} + \frac{d^2 \left(-\frac{f \ln(1+(dx+c)^2)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{(-cf+de) \arctan(dx+c)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{f \ln(f(dx+c)-cf+de)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} \right)}{f} \right)}{d}$
derivativdivides	$\frac{\frac{a d^2}{(cf-de-f(dx+c))f} + b d^2 \left(\frac{\arctan(dx+c)}{(cf-de-f(dx+c))f} - \frac{f \ln(cf-de-f(dx+c))}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{f \ln(1+(dx+c)^2)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{(cf-de) \arctan(dx+c)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} \right)}{d}$
default	$\frac{\frac{a d^2}{(cf-de-f(dx+c))f} + b d^2 \left(\frac{\arctan(dx+c)}{(cf-de-f(dx+c))f} - \frac{f \ln(cf-de-f(dx+c))}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{f \ln(1+(dx+c)^2)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} + \frac{(cf-de) \arctan(dx+c)}{c^2 f^2 - 2cdef + d^2 e^2 + f^2} \right)}{d}$
parallelrisc	$\frac{-2x \arctan(dx+c)bc d^3 f^2 + 2x \arctan(dx+c)b d^4 e f + 2 \ln(fx+e)xb d^3 f^2 - \ln(d^2 x^2 + 2cdx + c^2 + 1)xb d^3 f^2 - 2 \arctan(dx+c)bc d^3 f^2}{2}$
risc	$\frac{ib \ln(1+i(dx+c))}{2f(fx+e)} + \frac{-ib f^2 \ln(1-i(dx+c)) - i \ln((cdf - d^2 e - 3idf)x - 2icf - ide + c^2 f - edc + 3f)bcd f^2 x + i \ln((-cdf + c^2 f - edc + 3f)cd)}{2f(fx+e)}$

```
input int((a+b*arctan(d*x+c))/(f*x+e)^2,x,method=_RETURNVERBOSE)
```

output

```
-a/(f*x+e)/f+b/d*(-d^2/(f*(d*x+c)-c*f+d*e)/f*arctan(d*x+c)+d^2/f*(1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2))*(-1/2*f*ln(1+(d*x+c)^2)+(-c*f+d*e)*arctan(d*x+c))+1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f*ln(f*(d*x+c)-c*f+d*e))
```

Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.26

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \frac{2ad^2e^2 - 4acdef + 2(ac^2 + a)f^2 - 2(bcdef - (bc^2 + b)f^2 + (bd^2ef - bcd^2f^2)x) \arctan(dx + c) + (bd^2e^3f - 2cde^2f^2 + (c^2 + 1)ef^3 + (d^2e^2f^2 - 2cde^2f^2)x)}{2(d^2e^3f - 2cde^2f^2 + (c^2 + 1)ef^3 + (d^2e^2f^2 - 2cde^2f^2)x)}$$

input

```
integrate((a+b*arctan(d*x+c))/(f*x+e)^2,x, algorithm="fricas")
```

output

```
-1/2*(2*a*d^2*e^2 - 4*a*c*d*e*f + 2*(a*c^2 + a)*f^2 - 2*(b*c*d*e*f - (b*c^2 + b)*f^2 + (b*d^2*e*f - b*c*d*f^2)*x)*arctan(d*x + c) + (b*d*f^2*x + b*d*e*f)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 2*(b*d*f^2*x + b*d*e*f)*log(f*x + e))/(d^2*e^3*f - 2*c*d*e^2*f^2 + (c^2 + 1)*e*f^3 + (d^2*e^2*f^2 - 2*c*d*e*f^3 + (c^2 + 1)*f^4)*x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \text{Timed out}$$

input

```
integrate((a+b*atan(d*x+c))/(f*x+e)**2,x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.17

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx$$

$$= \frac{1}{2} \left(d \left(\frac{2(d^2e - cdf) \arctan\left(\frac{d^2x + cd}{d}\right)}{(d^2e^2f - 2cdef^2 + (c^2 + 1)f^3)d} - \frac{\log(d^2x^2 + 2cdx + c^2 + 1)}{d^2e^2 - 2cdef + (c^2 + 1)f^2} + \frac{2 \log(fx + e)}{d^2e^2 - 2cdef + (c^2 + 1)f^2} \right) - \frac{a}{f^2x + ef} \right)$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e)^2,x, algorithm="maxima")`

output `1/2*(d*(2*(d^2*e - c*d*f)*arctan((d^2*x + c*d)/d)/((d^2*e^2*f - 2*c*d*e*f^2 + (c^2 + 1)*f^3)*d) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2) + 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2)) - 2*arctan(d*x + c)/(f^2*x + e*f)*b - a/(f^2*x + e*f)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3974 vs. 2(149) = 298.

Time = 0.35 (sec) , antiderivative size = 3974, normalized size of antiderivative = 26.32

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \text{Too large to display}$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e)^2,x, algorithm="giac")`

output

```

1/2*(d^3*e*log(4*(d^2*e^2*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c
*f/(f*x + e))/f))^4 - 2*c*d*e*f*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x +
e) + c*f/(f*x + e))/f))^4 + c^2*f^2*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*
x + e) + c*f/(f*x + e))/f))^4 - 2*d^2*e^2*tan(1/2*arctan(-(f*x + e)*(d - d
*e/(f*x + e) + c*f/(f*x + e))/f))^2 + 4*c*d*e*f*tan(1/2*arctan(-(f*x + e)*
(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^2 - 2*c^2*f^2*tan(1/2*arctan(-(f*x
+ e)*(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^2 + 4*d*e*f*tan(1/2*arctan(-
(f*x + e)*(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^3 - 4*c*f^2*tan(1/2*arct
an(-(f*x + e)*(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^3 + d^2*e^2 - 2*c*d*
e*f + c^2*f^2 - 4*d*e*f*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c*f
/(f*x + e))/f)) + 4*c*f^2*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c
*f/(f*x + e))/f)) + 4*f^2*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c
*f/(f*x + e))/f))^2)/(tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c*f/(
f*x + e))/f))^4 + 2*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c*f/(f*
x + e))/f))^2 + 1))*tan(1/2*arctan(-(f*x + e)*(d - d*e/(f*x + e) + c*f/(f*
x + e))/f))^2 - c*d^2*f*log(4*(d^2*e^2*tan(1/2*arctan(-(f*x + e)*(d - d*e/
(f*x + e) + c*f/(f*x + e))/f))^4 - 2*c*d*e*f*tan(1/2*arctan(-(f*x + e)*(d
- d*e/(f*x + e) + c*f/(f*x + e))/f))^4 + c^2*f^2*tan(1/2*arctan(-(f*x + e)
*(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^4 - 2*d^2*e^2*tan(1/2*arctan(-(f*
x + e)*(d - d*e/(f*x + e) + c*f/(f*x + e))/f))^2 + 4*c*d*e*f*tan(1/2*ar...

```

Mupad [B] (verification not implemented)

Time = 2.38 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.84

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^2} dx = \frac{bd \ln(e + fx)}{d^2 e^2 - 2cdef + (c^2 + 1) f^2} - \frac{b \operatorname{atan}(c + dx)}{f(e + fx)} - \frac{a}{x f^2 + e f} - \frac{bd \ln(c + dx - i) \operatorname{li}}{2f(de - cf + f \operatorname{li})} - \frac{bd \ln(c + dx + i)}{2f(f - cf \operatorname{li} + de \operatorname{li})}$$

input

```
int((a + b*atan(c + d*x))/(e + f*x)^2,x)
```

output

```

(b*d*log(e + f*x))/(f^2*(c^2 + 1) + d^2*e^2 - 2*c*d*e*f) - (b*atan(c + d*x
))/(f*(e + f*x)) - a/(e*f + f^2*x) - (b*d*log(c + d*x - i)*i)/(2*f*(f*i
- c*f + d*e)) - (b*d*log(c + d*x + i))/(2*f*(f - c*f*i + d*e*i))

```


3.38 $\int \frac{a+b \arctan(c+dx)}{(e+fx)^3} dx$

Optimal result	344
Mathematica [C] (verified)	345
Rubi [A] (verified)	345
Maple [A] (verified)	348
Fricas [B] (verification not implemented)	349
Sympy [F(-1)]	349
Maxima [A] (verification not implemented)	350
Giac [C] (verification not implemented)	350
Mupad [B] (verification not implemented)	352
Reduce [B] (verification not implemented)	353

Optimal result

Integrand size = 18, antiderivative size = 227

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = -\frac{bd}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)(e + fx)} + \frac{bd^2(de + f - cf)(de - (1 + c)f) \arctan(c + dx)}{2f(d^2e^2 - 2cdef + (1 + c^2)f^2)^2} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2} + \frac{bd^2(de - cf) \log(e + fx)}{(d^2e^2 - 2cdef + (1 + c^2)f^2)^2} - \frac{bd^2(de - cf) \log(1 + c^2 + 2cdx + d^2x^2)}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)^2}$$

output

```
-1/2*b*d/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)/(f*x+e)+1/2*b*d^2*(-c*f+d*e+f)*(d
*e-(1+c)*f)*arctan(d*x+c)/f/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)^2-1/2*(a+b*arc
tan(d*x+c))/f/(f*x+e)^2+b*d^2*(-c*f+d*e)*ln(f*x+e)/(d^2*e^2-2*c*d*e*f+(c^2
+1)*f^2)^2-1/2*b*d^2*(-c*f+d*e)*ln(d^2*x^2+2*c*d*x+c^2+1)/(d^2*e^2-2*c*d*e
*f+(c^2+1)*f^2)^2
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.77

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx$$

$$= \frac{-\frac{a+b \arctan(c+dx)}{(e+fx)^2} + \frac{1}{2}bd^2 \left(-\frac{2f}{d(d^2e^2-2cdef+(1+c^2)f^2)(e+fx)} - \frac{i \log(i-c-dx)}{(de-(-i+c)f)^2} + \frac{i \log(i+c+dx)}{(de-(i+c)f)^2} - \frac{4f(-de+cf) \log(d(e+fx))}{(d^2e^2-2cdef+(1+c^2)f^2)^2} \right)}{2f}$$

input `Integrate[(a + b*ArcTan[c + d*x])/(e + f*x)^3,x]`

output

```
(-((a + b*ArcTan[c + d*x])/(e + f*x)^2) + (b*d^2*((-2*f)/(d*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(e + f*x)) - (I*Log[I - c - d*x])/(d*e - (-I + c)*f)^2 + (I*Log[I + c + d*x])/(d*e - (I + c)*f)^2 - (4*f*(-(d*e) + c*f)*Log[d*(e + f*x)])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)^2)/2)/(2*f)
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5568, 2081, 1145, 27, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx$$

$$\downarrow \text{5568}$$

$$\frac{bd \int \frac{1}{(e+fx)^2((c+dx)^2+1)} dx}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2}$$

$$\downarrow \text{2081}$$

$$\frac{bd \int \frac{1}{(e+fx)^2(c^2+2dxc+d^2x^2+1)} dx}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2}$$

$$\begin{array}{c}
 \downarrow 1145 \\
 \frac{bd \left(\frac{\int \frac{d(de-2cf-dfx)}{(e+fx)(c^2+2dxc+d^2x^2+1)} dx}{(c^2+1)f^2-2cdef+d^2e^2} - \frac{f}{(e+fx)((c^2+1)f^2-2cdef+d^2e^2)} \right)}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2} \\
 \downarrow 27 \\
 \frac{bd \left(\frac{d \int \frac{de-2cf-dfx}{(e+fx)(c^2+2dxc+d^2x^2+1)} dx}{(c^2+1)f^2-2cdef+d^2e^2} - \frac{f}{(e+fx)((c^2+1)f^2-2cdef+d^2e^2)} \right)}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2} \\
 \downarrow 1200 \\
 \frac{bd \left(\frac{d \int \left(\frac{2(de-cf)f^2}{(d^2e^2-2cdf e+(c^2+1)f^2)(e+fx)} + \frac{d(d^2e^2-4cdf e-(1-3c^2)f^2-2df(de-cf)x)}{(d^2e^2-2cdf e+(c^2+1)f^2)(c^2+2dxc+d^2x^2+1)} \right) dx}{(c^2+1)f^2-2cdef+d^2e^2} - \frac{f}{(e+fx)((c^2+1)f^2-2cdef+d^2e^2)} \right)}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2} \\
 \downarrow 2009 \\
 \frac{bd \left(\frac{d \left(\frac{\arctan(c+dx)(-cf+de+f)(de-(c+1)f)}{(c^2+1)f^2-2cdef+d^2e^2} - \frac{f(de-cf) \log(c^2+2cdx+d^2x^2+1)}{(c^2+1)f^2-2cdef+d^2e^2} + \frac{2f(de-cf) \log(e+fx)}{(c^2+1)f^2-2cdef+d^2e^2} \right)}{(c^2+1)f^2-2cdef+d^2e^2} - \frac{f}{(e+fx)((c^2+1)f^2-2cdef+d^2e^2)} \right)}{2f} - \frac{a + b \arctan(c + dx)}{2f(e + fx)^2}
 \end{array}$$

input `Int[(a + b*ArcTan[c + d*x])/(e + f*x)^3,x]`

output `-1/2*(a + b*ArcTan[c + d*x])/(f*(e + f*x)^2) + (b*d*(-(f/((d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)*(e + f*x))) + (d*(((d*e + f - c*f)*(d*e - (1 + c)*f)*ArcTan[c + d*x])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (2*f*(d*e - c*f)*Log[e + f*x])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) - (f*(d*e - c*f)*Log[1 + c^2 + 2*c*d*x + d^2*x^2])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2))))/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)))/(2*f)`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 1145 `Int[((d_) + (e_)*(x_)^(m_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[e*((d + e*x)^(m + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))], x] + Simp[1/(c*d^2 - b*d*e + a*e^2) Int[(d + e*x)^(m + 1)*(Simp[c*d - b*e - c*e*x, x]/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[m, -1]`
- rule 1200 `Int((((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_)))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IntegersQ[n]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2081 `Int[(u_)^(m_)*(v_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^m*ExpandToSum[v, x]^p, x] /; FreeQ[{m, p}, x] && LinearQ[u, x] && QuadraticQ[v, x] && !(LinearMatchQ[u, x] && QuadraticMatchQ[v, x])`
- rule 5568 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_)^(m_)), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.08

method	result
parts	$-\frac{a}{2(fx+e)^2 f} + \frac{b}{2(f(dx+c)-cf+de)^2 f} + \frac{d^3}{d} \left(\frac{\frac{(2cf^2-2def)\ln(1+(dx+c)^2)}{2} + (c^2f^2-2cdef+d^2e^2-f^2)\arctan(dx+c)}{(c^2f^2-2cdef+d^2e^2+f^2)^2} \right)$
derivativdivides	$-\frac{a d^3}{2(cf-de-f(dx+c))^2 f} - b d^3 \left(\frac{\arctan(dx+c)}{2(cf-de-f(dx+c))^2 f} - \frac{f}{(c^2f^2-2cdef+d^2e^2+f^2)(cf-de-f(dx+c))} - \frac{2f(cf-de)\ln(cf-de-f(dx+c))}{(c^2f^2-2cdef+d^2e^2+f^2)^2} \right)$
default	$-\frac{a d^3}{2(cf-de-f(dx+c))^2 f} - b d^3 \left(\frac{\arctan(dx+c)}{2(cf-de-f(dx+c))^2 f} - \frac{f}{(c^2f^2-2cdef+d^2e^2+f^2)(cf-de-f(dx+c))} - \frac{2f(cf-de)\ln(cf-de-f(dx+c))}{(c^2f^2-2cdef+d^2e^2+f^2)^2} \right)$
parallelrisch	$-\frac{4 \ln(fx+e) x b c d^4 e f^4 - 2 \ln(d^2 x^2 + 2 c d x + c^2 + 1) x b c d^4 e f^4 + 2 x^2 \arctan(dx+c) b c d^5 e f^4 - 2 x \arctan(dx+c) b c^2 d^4 e f^4}{d}$
risch	Expression too large to display

```
input int((a+b*arctan(d*x+c))/(f*x+e)^3,x,method=_RETURNVERBOSE)
```

```
output -1/2*a/(f*x+e)^2/f+b/d*(-1/2*d^3/(f*(d*x+c)-c*f+d*e)^2/f*arctan(d*x+c)+1/2*d^3/f*(1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)^2*(1/2*(2*c*f^2-2*d*e*f)*ln(1+(d*x+c)^2)+(c^2*f^2-2*c*d*e*f+d^2*e^2-f^2)*arctan(d*x+c))-f/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)/(f*(d*x+c)-c*f+d*e)-2*(c*f-d*e)*f/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)^2*ln(f*(d*x+c)-c*f+d*e)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 682 vs. 2(219) = 438.

Time = 1.22 (sec) , antiderivative size = 682, normalized size of antiderivative = 3.00

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = \frac{ad^4e^4 - (4ac - b)d^3e^3f + 2(3ac^2 - bc + a)d^2e^2f^2 - (4ac^3 - bc^2 + 4ac - b)def^3 + (ac^4 + 2ac^2 + a)}$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e)^3,x, algorithm="fricas")`

output

```
-1/2*(a*d^4*e^4 - (4*a*c - b)*d^3*e^3*f + 2*(3*a*c^2 - b*c + a)*d^2*e^2*f^2 - (4*a*c^3 - b*c^2 + 4*a*c - b)*d*e*f^3 + (a*c^4 + 2*a*c^2 + a)*f^4 + (b*d^3*e^2*f^2 - 2*b*c*d^2*e*f^3 + (b*c^2 + b)*d*f^4)*x - (2*b*c*d^3*e^3*f - (5*b*c^2 + 3*b)*d^2*e^2*f^2 + 4*(b*c^3 + b*c)*d*e*f^3 - (b*c^4 + 2*b*c^2 + b)*f^4 + (b*d^4*e^2*f^2 - 2*b*c*d^3*e*f^3 + (b*c^2 - b)*d^2*f^4)*x^2 + 2*(b*d^4*e^3*f - 2*b*c*d^3*e^2*f^2 + (b*c^2 - b)*d^2*e*f^3)*x)*arctan(d*x + c) + (b*d^3*e^3*f - b*c*d^2*e^2*f^2 + (b*d^3*e*f^3 - b*c*d^2*f^4)*x^2 + 2*(b*d^3*e^2*f^2 - b*c*d^2*e*f^3)*x)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 2*(b*d^3*e^3*f - b*c*d^2*e^2*f^2 + (b*d^3*e*f^3 - b*c*d^2*f^4)*x^2 + 2*(b*d^3*e^2*f^2 - b*c*d^2*e*f^3)*x)*log(f*x + e))/(d^4*e^6*f - 4*c*d^3*e^5*f^2 + 2*(3*c^2 + 1)*d^2*e^4*f^3 - 4*(c^3 + c)*d*e^3*f^4 + (c^4 + 2*c^2 + 1)*e^2*f^5 + (d^4*e^4*f^3 - 4*c*d^3*e^3*f^4 + 2*(3*c^2 + 1)*d^2*e^2*f^5 - 4*(c^3 + c)*d*e*f^6 + (c^4 + 2*c^2 + 1)*f^7)*x^2 + 2*(d^4*e^5*f^2 - 4*c*d^3*e^4*f^3 + 2*(3*c^2 + 1)*d^2*e^3*f^4 - 4*(c^3 + c)*d*e^2*f^5 + (c^4 + 2*c^2 + 1)*e*f^6)*x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))/(f*x+e)**3,x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.80

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx =$$

$$-\frac{1}{2} \left(d \left(\frac{(d^2 e - cdf) \log(d^2 x^2 + 2cdx + c^2 + 1)}{d^4 e^4 - 4cd^3 e^3 f + 2(3c^2 + 1)d^2 e^2 f^2 - 4(c^3 + c)def^3 + (c^4 + 2c^2 + 1)f^4} - \frac{a}{2(f^3 x^2 + 2ef^2 x + e^2 f)} \right) \right)$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e)^3,x, algorithm="maxima")`

output

$$-1/2*(d*((d^2*e - c*d*f)*\log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^4*e^4 - 4*c*d^3*e^3*f + 2*(3*c^2 + 1)*d^2*e^2*f^2 - 4*(c^3 + c)*d*e*f^3 + (c^4 + 2*c^2 + 1)*f^4) - 2*(d^2*e - c*d*f)*\log(f*x + e)/(d^4*e^4 - 4*c*d^3*e^3*f + 2*(3*c^2 + 1)*d^2*e^2*f^2 - 4*(c^3 + c)*d*e*f^3 + (c^4 + 2*c^2 + 1)*f^4) - (d^4*e^2 - 2*c*d^3*e*f + (c^2 - 1)*d^2*f^2)*\arctan((d^2*x + c*d)/d)/((d^4*e^4*f - 4*c*d^3*e^3*f^2 + 2*(3*c^2 + 1)*d^2*e^2*f^3 - 4*(c^3 + c)*d*e*f^4 + (c^4 + 2*c^2 + 1)*f^5)*d) + 1/(d^2*e^3 - 2*c*d*e^2*f + (c^2 + 1)*e*f^2 + (d^2*e^2*f - 2*c*d*e*f^2 + (c^2 + 1)*f^3)*x)) + \arctan(d*x + c)/(f^3*x^2 + 2*e*f^2*x + e^2*f))*b - 1/2*a/(f^3*x^2 + 2*e*f^2*x + e^2*f)$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.59 (sec) , antiderivative size = 1561, normalized size of antiderivative = 6.88

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input `integrate((a+b*arctan(d*x+c))/(f*x+e)^3,x, algorithm="giac")`

output

```

1/4*(I*b*d^4*e^2*f^2*x^2*log(I*d*x + I*c - 1) - 2*I*b*c*d^3*e*f^3*x^2*log(
I*d*x + I*c - 1) + I*b*c^2*d^2*f^4*x^2*log(I*d*x + I*c - 1) - I*b*d^4*e^2*
f^2*x^2*log(-I*d*x - I*c - 1) + 2*I*b*c*d^3*e*f^3*x^2*log(-I*d*x - I*c - 1
) - I*b*c^2*d^2*f^4*x^2*log(-I*d*x - I*c - 1) + 2*I*b*d^4*e^3*f*x*log(I*d*
x + I*c - 1) - 4*I*b*c*d^3*e^2*f^2*x*log(I*d*x + I*c - 1) + 2*I*b*c^2*d^2*
e*f^3*x*log(I*d*x + I*c - 1) - 2*b*d^3*e*f^3*x^2*log(I*d*x + I*c - 1) + 2*
b*c*d^2*f^4*x^2*log(I*d*x + I*c - 1) - 2*I*b*d^4*e^3*f*x*log(-I*d*x - I*c
- 1) + 4*I*b*c*d^3*e^2*f^2*x*log(-I*d*x - I*c - 1) - 2*I*b*c^2*d^2*e*f^3*x
*log(-I*d*x - I*c - 1) - 2*b*d^3*e*f^3*x^2*log(-I*d*x - I*c - 1) + 2*b*c*d
^2*f^4*x^2*log(-I*d*x - I*c - 1) + 4*b*d^3*e*f^3*x^2*log(f*x + e) - 4*b*c*
d^2*f^4*x^2*log(f*x + e) - 2*b*d^4*e^4*arctan(d*x + c) + 8*b*c*d^3*e^3*f*a
rctan(d*x + c) - 12*b*c^2*d^2*e^2*f^2*arctan(d*x + c) + 8*b*c^3*d*e*f^3*ar
ctan(d*x + c) - 2*b*c^4*f^4*arctan(d*x + c) + I*b*d^4*e^4*log(I*d*x + I*c
- 1) - 2*I*b*c*d^3*e^3*f*log(I*d*x + I*c - 1) + I*b*c^2*d^2*e^2*f^2*log(I*
d*x + I*c - 1) - 4*b*d^3*e^2*f^2*x*log(I*d*x + I*c - 1) + 4*b*c*d^2*e*f^3*
x*log(I*d*x + I*c - 1) - I*b*d^2*f^4*x^2*log(I*d*x + I*c - 1) - I*b*d^4*e^
4*log(-I*d*x - I*c - 1) + 2*I*b*c*d^3*e^3*f*log(-I*d*x - I*c - 1) - I*b*c^
2*d^2*e^2*f^2*log(-I*d*x - I*c - 1) - 4*b*d^3*e^2*f^2*x*log(-I*d*x - I*c
- 1) + 4*b*c*d^2*e*f^3*x*log(-I*d*x - I*c - 1) + I*b*d^2*f^4*x^2*log(-I*d*x
- I*c - 1) + 8*b*d^3*e^2*f^2*x*log(f*x + e) - 8*b*c*d^2*e*f^3*x*log(f*...

```


Mupad [B] (verification not implemented)

Time = 8.60 (sec) , antiderivative size = 399, normalized size of antiderivative = 1.76

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = \frac{bd^3 e \ln(e + fx)}{(c^2 f^2 - 2cdef + d^2 e^2 + f^2)^2} - \frac{af}{2(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{bde}{2(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{ac^2 f}{2(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{b \operatorname{atan}(c + dx)}{2f(e + fx)^2} - \frac{bcd^2 f \ln(e + fx)}{(c^2 f^2 - 2cdef + d^2 e^2 + f^2)^2} + \frac{acde}{(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{bdfx}{2(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{ad^2 e^2}{2f(e + fx)^2 (c^2 f^2 - 2cdef + d^2 e^2 + f^2)} - \frac{bd^2 \ln(c + dx - i) \operatorname{li}}{4f(de - cf + fli)^2} + \frac{bd^2 \ln(c + dx + li) \operatorname{li}}{4f(cf - de + fli)^2}$$

input `int((a + b*atan(c + d*x))/(e + f*x)^3,x)`output `(b*d^2*log(c + d*x + 1i)*1i)/(4*f*(f*1i + c*f - d*e)^2) - (a*f)/(2*(e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)) - (b*d*e)/(2*(e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)) - (b*d^2*log(c + d*x - 1i)*1i)/(4*f*(f*1i - c*f + d*e)^2) - (b*atan(c + d*x))/(2*f*(e + f*x)^2) - (a*c^2*f)/(2*(e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)) + (b*d^3*e*log(e + f*x))/(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)^2 - (b*c*d^2*f*log(e + f*x))/(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)^2 + (a*c*d*e)/((e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)) - (b*d*f*x)/(2*(e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f)) - (a*d^2*e^2)/(2*f*(e + f*x)^2*(f^2 + c^2*f^2 + d^2*e^2 - 2*c*d*e*f))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 1110, normalized size of antiderivative = 4.89

$$\int \frac{a + b \arctan(c + dx)}{(e + fx)^3} dx = \text{Too large to display}$$

input `int((a+b*atan(d*x+c))/(f*x+e)^3,x)`

output

```
( - 2*atan(c + d*x)*b*c**4*e*f**4 + 8*atan(c + d*x)*b*c**3*d*e**2*f**3 - 1
0*atan(c + d*x)*b*c**2*d**2*e**3*f**2 + 4*atan(c + d*x)*b*c**2*d**2*e**2*f
**3*x + 2*atan(c + d*x)*b*c**2*d**2*e*f**4*x**2 - 4*atan(c + d*x)*b*c**2*e
*f**4 + 4*atan(c + d*x)*b*c*d**3*e**4*f - 8*atan(c + d*x)*b*c*d**3*e**3*f*
*2*x - 4*atan(c + d*x)*b*c*d**3*e**2*f**3*x**2 + 8*atan(c + d*x)*b*c*d*e**
2*f**3 + 4*atan(c + d*x)*b*d**4*e**4*f*x + 2*atan(c + d*x)*b*d**4*e**3*f**
2*x**2 - 6*atan(c + d*x)*b*d**2*e**3*f**2 - 4*atan(c + d*x)*b*d**2*e**2*f*
*3*x - 2*atan(c + d*x)*b*d**2*e*f**4*x**2 - 2*atan(c + d*x)*b*e*f**4 + 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d**2*e**3*f**2 + 4*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d**2*e**2*f**3*x + 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*c*d**2*e*f**4*x**2 - 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d**3*e**4*f - 4*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d**3*e**3*f**2*x - 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b*d**3*e**2*f**3*x**2 - 4*log(e + f*x)*b*c*d**2*e**3*f**2 - 8*log(e + f*x)*b*c*d**2*e**2*f**3*x - 4*log(e + f*x)*b*c*d**2*e*f**4*x**2 + 4*log(e + f*x)*b*d**3*e**4*f + 8*log(e + f*x)*b*d**3*e**3*f**2*x + 4*log(e + f*x)*b*d**3*e**2*f**3*x**2 - 2*a*c**4*e*f**4 + 8*a*c**3*d*e**2*f**3 - 12*a*c**2*d**2*e**3*f**2 - 4*a*c**2*e*f**4 + 8*a*c*d**3*e**4*f + 8*a*c*d*e**2*f**3 - 2*a*d**4*e**5 - 4*a*d**2*e**3*f**2 - 2*a*e*f**4 - b*c**2*d*e**2*f**3 + b*c**2*d*f**5*x**2 + 2*b*c*d**2*e**3*f**2 - 2*b*c*d**2*e*f**4*x**2 - b*d**3*e**4*f + b*d**3*e**2*f**3*x**2 - b*d*e**2...
```

3.39 $\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx$

Optimal result	354
Mathematica [B] (warning: unable to verify)	355
Rubi [A] (verified)	356
Maple [B] (verified)	358
Fricas [F]	359
Sympy [F]	360
Maxima [F]	360
Giac [F]	361
Mupad [F(-1)]	361
Reduce [F]	361

Optimal result

Integrand size = 20, antiderivative size = 382

$$\begin{aligned}
 & \int (e + fx)^2 (a + b \arctan(c + dx))^2 dx \\
 &= \frac{b^2 f^2 x}{3d^2} - \frac{2abf(de - cf)x}{d^2} - \frac{b^2 f^2 \arctan(c + dx)}{3d^3} \\
 & \quad - \frac{2b^2 f(de - cf)(c + dx) \arctan(c + dx)}{d^3} - \frac{bf^2(c + dx)^2(a + b \arctan(c + dx))}{3d^3} \\
 & \quad + \frac{i(3d^2 e^2 - 6cdef - (1 - 3c^2) f^2)(a + b \arctan(c + dx))^2}{3d^3} \\
 & \quad - \frac{(de - cf)(d^2 e^2 - 2cdef - (3 - c^2) f^2)(a + b \arctan(c + dx))^2}{3d^3 f} \\
 & \quad + \frac{(e + fx)^3 (a + b \arctan(c + dx))^2}{3f} \\
 & \quad + \frac{2b(3d^2 e^2 - 6cdef - (1 - 3c^2) f^2)(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{3d^3} \\
 & \quad + \frac{b^2 f(de - cf) \log(1 + (c + dx)^2)}{d^3} \\
 & \quad + \frac{ib^2(3d^2 e^2 - 6cdef - (1 - 3c^2) f^2) \text{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{3d^3}
 \end{aligned}$$

output

```

1/3*b^2*f^2*x/d^2-2*a*b*f*(-c*f+d*e)*x/d^2-1/3*b^2*f^2*arctan(d*x+c)/d^3-2
*b^2*f*(-c*f+d*e)*(d*x+c)*arctan(d*x+c)/d^3-1/3*b*f^2*(d*x+c)^2*(a+b*arcta
n(d*x+c))/d^3+1/3*I*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*(a+b*arctan(d*x+c
))^2/d^3-1/3*(-c*f+d*e)*(d^2*e^2-2*c*d*e*f-(-c^2+3)*f^2)*(a+b*arctan(d*x+c
))^2/d^3/f+1/3*(f*x+e)^3*(a+b*arctan(d*x+c))^2/f+2/3*b*(3*d^2*e^2-6*c*d*e*
f-(-3*c^2+1)*f^2)*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/d^3+b^2*f*(-c*f+
d*e)*ln(1+(d*x+c)^2)/d^3+1/3*I*b^2*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*po
lylog(2,1-2/(1+I*(d*x+c)))/d^3

```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 801 vs. $2(382) = 764$.

Time = 7.63 (sec) , antiderivative size = 801, normalized size of antiderivative = 2.10

$$\begin{aligned}
& \int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = a^2 e^2 x + a^2 e f x^2 + \frac{1}{3} a^2 f^2 x^3 \\
& + \frac{ab(-dfx(6de - 4cf + dfx) + 2(3def - 3c^2 def + c^3 f^2 + 3c(d^2 e^2 - f^2) + d^3 x(3e^2 + 3efx + f^2 x^2)) \arctan(c + dx) + 3d^3 \log(1 + e^{2i \arctan(c+dx)})) - i \operatorname{PolyLog}(2, -e^{2i \arctan(c+dx)})}{d^3} \\
& + \frac{b^2 e^2 (\arctan(c + dx) ((-i + c + dx) \arctan(c + dx) + 2 \log(1 + e^{2i \arctan(c+dx)})) - i \operatorname{PolyLog}(2, -e^{2i \arctan(c+dx)}))}{d} \\
& + \frac{b^2 e f ((1 + 2ic - c^2 + d^2 x^2) \arctan(c + dx)^2 - 2 \arctan(c + dx) (c + dx + 2c \log(1 + e^{2i \arctan(c+dx)})) + 2 \log(1 + e^{2i \arctan(c+dx)})^2)}{d^2} \\
& + \frac{b^2 f^2 (1 + (c + dx)^2)^{3/2} \left(\frac{c+dx}{\sqrt{1+(c+dx)^2}} + \frac{6c(c+dx) \arctan(c+dx)}{\sqrt{1+(c+dx)^2}} + \frac{3(c+dx) \arctan(c+dx)^2}{\sqrt{1+(c+dx)^2}} + \frac{3c^2(c+dx) \arctan(c+dx)^2}{\sqrt{1+(c+dx)^2}} \right)}{d^3}
\end{aligned}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcTan[c + d*x])^2,x]
```

output

```

a^2*e^2*x + a^2*e*f*x^2 + (a^2*f^2*x^3)/3 + (a*b*(-(d*f*x*(6*d*e - 4*c*f +
d*f*x)) + 2*(3*d*e*f - 3*c^2*d*e*f + c^3*f^2 + 3*c*(d^2*e^2 - f^2) + d^3*
x*(3*e^2 + 3*e*f*x + f^2*x^2))*ArcTan[c + d*x] + (-3*d^2*e^2 + 6*c*d*e*f +
(1 - 3*c^2)*f^2)*Log[1 + (c + d*x)^2])/d + (b^2*e^2*(ArcTan[c + d*
x]*((-I + c + d*x)*ArcTan[c + d*x] + 2*Log[1 + E^((2*I)*ArcTan[c + d*x])])
- I*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]))/d + (b^2*e*f*((1 + (2*I)*c -
c^2 + d^2*x^2)*ArcTan[c + d*x]^2 - 2*ArcTan[c + d*x]*(c + d*x + 2*c*Log[1
+ E^((2*I)*ArcTan[c + d*x])]) + Log[1 + (c + d*x)^2] + (2*I)*c*PolyLog[2,
-E^((2*I)*ArcTan[c + d*x])]))/d^2 + (b^2*f^2*(1 + (c + d*x)^2)^(3/2)*((c
+ d*x)/Sqrt[1 + (c + d*x)^2] + (6*c*(c + d*x)*ArcTan[c + d*x])/Sqrt[1 + (c
+ d*x)^2] + (3*(c + d*x)*ArcTan[c + d*x]^2)/Sqrt[1 + (c + d*x)^2] + (3*c^
2*(c + d*x)*ArcTan[c + d*x]^2)/Sqrt[1 + (c + d*x)^2] + I*ArcTan[c + d*x]^2
*Cos[3*ArcTan[c + d*x]] - (3*I)*c^2*ArcTan[c + d*x]^2*Cos[3*ArcTan[c + d*x
]] - 2*ArcTan[c + d*x]*Cos[3*ArcTan[c + d*x]]*Log[1 + E^((2*I)*ArcTan[c +
d*x])]) + 6*c^2*ArcTan[c + d*x]*Cos[3*ArcTan[c + d*x]]*Log[1 + E^((2*I)*Arc
Tan[c + d*x])]) + 6*c*cos[3*ArcTan[c + d*x]]*Log[1/Sqrt[1 + (c + d*x)^2]] +
((3*I - 12*c - (9*I)*c^2)*ArcTan[c + d*x]^2 + 2*ArcTan[c + d*x]*(-2 + (-3
+ 9*c^2)*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + 18*c*Log[1/Sqrt[1 + (c + d
*x)^2]])/Sqrt[1 + (c + d*x)^2] - ((4*I)*(-1 + 3*c^2)*PolyLog[2, -E^((2*I)*
ArcTan[c + d*x])])/(1 + (c + d*x)^2)^(3/2) + Sin[3*ArcTan[c + d*x]] + 6...

```

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5570, 27, 5389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx$$

$$\downarrow 5570$$

$$\int \frac{\left(\frac{d(e - \frac{ef}{d}) + f(c + dx)}{d^2}\right)^2 (a + b \arctan(c + dx))^2}{d} d(c + dx)$$

$$\downarrow 27$$

$$\frac{\int (de - cf + f(c + dx))^2 (a + b \arctan(c + dx))^2 d(c + dx)}{d^3}$$

↓ 5389

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))^2}{3f} - \frac{2b \int \left((c+dx)(a+b \arctan(c+dx))f^3 + 3(de-cf)(a+b \arctan(c+dx))f^2 + \frac{(de-cf)(d^2e^2 - 2cdf e - (3-c^2)f^2)(a+b \arctan(c+dx))^2}{2b} \right) d(c+dx)}{d^3}}{3f}$$

↓ 2009

$$\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))^2}{3f} - \frac{2b \left(-\frac{if(-1-3c^2)f^2-6cdf+3d^2e^2}{2b}(a+b \arctan(c+dx))^2 + \frac{(de-cf)(-(3-c^2)f^2-2cdf+ d^2e^2)(a+b \arctan(c+dx))^2}{2b} \right)}{d^3}$$

input `Int[(e + f*x)^2*(a + b*ArcTan[c + d*x])^2,x]`

output `((((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTan[c + d*x])^2)/(3*f) - (2*b*(-1/2*(b*f^3*(c + d*x) + 3*a*f^2*(d*e - c*f)*(c + d*x) + (b*f^3*ArcTan[c + d*x])/2 + 3*b*f^2*(d*e - c*f)*(c + d*x)*ArcTan[c + d*x] + (f^3*(c + d*x)^2*(a + b*ArcTan[c + d*x]))/2 - ((I/2)*f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*(a + b*ArcTan[c + d*x])^2)/b + ((d*e - c*f)*(d^2*e^2 - 2*c*d*e*f - (3 - c^2)*f^2)*(a + b*ArcTan[c + d*x])^2)/(2*b) - f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*(a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] - (3*b*f^2*(d*e - c*f)*Log[1 + (c + d*x)^2])/2 - (I/2)*b*f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*PolyLog[2, 1 - 2/(1 + I*(c + d*x))]))/(3*f))/d^3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5389

```
Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))^(p_.)*((d_.) + (e_.)*(x_.))^(q_.), x_Sy
mbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - S
imp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] &&
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

rule 5570

```
Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1017 vs. $2(362) = 724$.

Time = 0.67 (sec) , antiderivative size = 1018, normalized size of antiderivative = 2.66

method	result	size
parts	Expression too large to display	1018
derivativeldivides	Expression too large to display	1042
default	Expression too large to display	1042
risch	Expression too large to display	2416

input

```
int((f*x+e)^2*(a+b*arctan(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```

1/3*a^2*(f*x+e)^3/f+b^2/d*(1/3/d^2*f^2*arctan(d*x+c)^2*(d*x+c)^3-1/d^2*f^2
*arctan(d*x+c)^2*(d*x+c)^2*c+1/d*f*arctan(d*x+c)^2*(d*x+c)^2*e+1/d^2*f^2*a
rctan(d*x+c)^2*(d*x+c)*c^2-2/d*f*arctan(d*x+c)^2*(d*x+c)*c*e+arctan(d*x+c)
^2*(d*x+c)*e^2-1/3/d^2*f^2*arctan(d*x+c)^2*c^3+1/d*f*arctan(d*x+c)^2*c^2*e
-arctan(d*x+c)^2*c*e^2+1/3*d/f*arctan(d*x+c)^2*e^3-2/3/d^2/f*(1/2*arctan(d
*x+c)*f^3*(d*x+c)^2-3*arctan(d*x+c)*c*f^3*(d*x+c)+3*arctan(d*x+c)*d*e*f^2*
(d*x+c)+3/2*arctan(d*x+c)*ln(1+(d*x+c)^2)*c^2*f^3-3*arctan(d*x+c)*ln(1+(d*
x+c)^2)*c*d*e*f^2+3/2*arctan(d*x+c)*ln(1+(d*x+c)^2)*d^2*e^2*f-1/2*arctan(d
*x+c)*ln(1+(d*x+c)^2)*f^3-arctan(d*x+c)^2*c^3*f^3+3*arctan(d*x+c)^2*c^2*d*
e*f^2-3*arctan(d*x+c)^2*c*d^2*e^2*f+arctan(d*x+c)^2*d^3*e^3+3*arctan(d*x+c
)^2*c*f^3-3*arctan(d*x+c)^2*d*e*f^2-1/2*f^2*(f*(d*x+c)+1/2*(-6*c*f+6*d*e)*
ln(1+(d*x+c)^2)-f*arctan(d*x+c))-1/2*f*(3*c^2*f^2-6*c*d*e*f+3*d^2*e^2-f^2)
*(-1/2*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+
c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))+1/2*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)
-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))
)-1/4*(-2*c^3*f^3+6*c^2*d*e*f^2-6*c*d^2*e^2*f+2*d^3*e^3+6*c*f^3-6*d*e*f^2)
*arctan(d*x+c)^2))+1/3*a*b/d^3*f^2*ln(1+(d*x+c)^2)+2*b/d*arctan(d*x+c)*a*c
*e^2-1/3/d*f^2*b*a*x^2+2/3/d^3*c^3*f^2*b*a*arctan(d*x+c)+4/3/d^2*c*f^2*x*b
*a-2/d*e*x*f*b*a-a*b/d^3*f^2*ln(1+(d*x+c)^2)*c^2-a*b/d*ln(1+(d*x+c)^2)*e^2
-2*b/d^3*arctan(d*x+c)*a*c*f^2+2/d^2*e*f*b*a*arctan(d*x+c)-2*b/d^2*arct...

```

Fricas [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")
```

output

```

integral(a^2*f^2*x^2 + 2*a^2*e*f*x + a^2*e^2 + (b^2*f^2*x^2 + 2*b^2*e*f*x
+ b^2*e^2)*arctan(d*x + c)^2 + 2*(a*b*f^2*x^2 + 2*a*b*e*f*x + a*b*e^2)*arc
tan(d*x + c), x)

```


Sympy [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \int (a + b \operatorname{atan}(c + dx))^2 (e + fx)^2 dx$$

input `integrate((f*x+e)**2*(a+b*atan(d*x+c))**2,x)`

output `Integral((a + b*atan(c + d*x))**2*(e + f*x)**2, x)`

Maxima [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output `3/4*b^2*c^2*e^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 1/4*(3*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*b^2*c^2*e^2 + 1/3*a^2*f^2*x^3 + 36*b^2*d^2*f^2*integrate(1/48*x^4*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^2*d^2*f^2*integrate(1/48*x^4*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 72*b^2*d^2*e*f*integrate(1/48*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 72*b^2*c*d*f^2*integrate(1/48*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 4*b^2*d^2*f^2*integrate(1/48*x^4*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b^2*d^2*e*f*integrate(1/48*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 36*b^2*d^2*e^2*integrate(1/48*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 144*b^2*c*d*e*f*integrate(1/48*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 36*b^2*c^2*f^2*integrate(1/48*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*d^2*e*f*integrate(1/48*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 4*b^2*c*d*f^2*integrate(1/48*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^2*d^2*e^2*integrate(1/48*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*c*d*e*f*integrate(1/48*x^2*log(d^2*x^2 + 2...`

Giac [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^2 dx$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c))^2,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \int (e + fx)^2 (a + b \operatorname{atan}(c + dx))^2 dx$$

input `int((e + f*x)^2*(a + b*atan(c + d*x))^2,x)`

output `int((e + f*x)^2*(a + b*atan(c + d*x))^2, x)`

Reduce [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^2 dx = \text{Too large to display}$$

input `int((f*x+e)^2*(a+b*atan(d*x+c))^2,x)`

output

```
( - 2*atan(c + d*x)**2*b**2*c**3*f**2 + 3*atan(c + d*x)**2*b**2*c**2*d*e*f
- 2*atan(c + d*x)**2*b**2*c*f**2 + 3*atan(c + d*x)**2*b**2*d**3*e**2*x +
3*atan(c + d*x)**2*b**2*d**3*e*f*x**2 + atan(c + d*x)**2*b**2*d**3*f**2*x*
*3 + 3*atan(c + d*x)**2*b**2*d*e*f + 2*atan(c + d*x)*a*b*c**3*f**2 - 6*ata
n(c + d*x)*a*b*c**2*d*e*f + 6*atan(c + d*x)*a*b*c*d**2*e**2 - 6*atan(c + d
*x)*a*b*c*f**2 + 6*atan(c + d*x)*a*b*d**3*e**2*x + 6*atan(c + d*x)*a*b*d**
3*e*f*x**2 + 2*atan(c + d*x)*a*b*d**3*f**2*x**3 + 6*atan(c + d*x)*a*b*d*e*
f + 5*atan(c + d*x)*b**2*c**2*f**2 - 6*atan(c + d*x)*b**2*c*d*e*f + 4*atan
(c + d*x)*b**2*c*d*f**2*x - 6*atan(c + d*x)*b**2*d**2*e*f*x - atan(c + d*x
)*b**2*d**2*f**2*x**2 - atan(c + d*x)*b**2*f**2 - 6*int((atan(c + d*x)*x)/
(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*c**2*d**2*f**2 + 12*int((atan(c +
d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*c*d**3*e*f - 6*int((atan
(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*d**4*e**2 + 2*int((a
tan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*d**2*f**2 - 3*log
(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*c**2*f**2 + 6*log(c**2 + 2*c*d*x + d*
**2*x**2 + 1)*a*b*c*d*e*f - 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*d**2*
e**2 + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*f**2 - 3*log(c**2 + 2*c*d*x
+ d**2*x**2 + 1)*b**2*c*f**2 + 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2
*d*e*f + 3*a**2*d**3*e**2*x + 3*a**2*d**3*e*f*x**2 + a**2*d**3*f**2*x**3 +
4*a*b*c*d*f**2*x - 6*a*b*d**2*e*f*x - a*b*d**2*f**2*x**2 + b**2*d*f**2...
```

3.40 $\int (e + fx)(a + b \arctan(c + dx))^2 dx$

Optimal result	363
Mathematica [A] (verified)	364
Rubi [A] (verified)	364
Maple [A] (verified)	366
Fricas [F]	367
Sympy [F]	367
Maxima [F]	368
Giac [F]	368
Mupad [F(-1)]	369
Reduce [F]	369

Optimal result

Integrand size = 18, antiderivative size = 222

$$\begin{aligned} & \int (e + fx)(a + b \arctan(c + dx))^2 dx \\ &= -\frac{abfx}{d} - \frac{b^2 f(c + dx) \arctan(c + dx)}{d^2} + \frac{i(de - cf)(a + b \arctan(c + dx))^2}{d^2} \\ & \quad - \frac{(de + f - cf)(de - (1 + c)f)(a + b \arctan(c + dx))^2}{2d^2 f} \\ & \quad + \frac{(e + fx)^2(a + b \arctan(c + dx))^2}{2f} \\ & \quad + \frac{2b(de - cf)(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d^2} \\ & \quad + \frac{b^2 f \log(1 + (c + dx)^2)}{2d^2} + \frac{ib^2(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d^2} \end{aligned}$$

output

```
-a*b*f*x/d-b^2*f*(d*x+c)*arctan(d*x+c)/d^2+I*(-c*f+d*e)*(a+b*arctan(d*x+c)
)^2/d^2-1/2*(-c*f+d*e+f)*(d*e-(1+c)*f)*(a+b*arctan(d*x+c))^2/d^2/f+1/2*(f*
x+e)^2*(a+b*arctan(d*x+c))^2/f+2*b*(-c*f+d*e)*(a+b*arctan(d*x+c))*ln(2/(1+
I*(d*x+c)))/d^2+1/2*b^2*f*ln(1+(d*x+c)^2)/d^2+I*b^2*(-c*f+d*e)*polylog(2,1
-2/(1+I*(d*x+c)))/d^2
```

Mathematica [A] (verified)

Time = 0.86 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.19

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx$$

$$= \frac{2a^2cde - 2abcf - a^2c^2f + 2a^2d^2ex - 2abdfx + a^2d^2fx^2 + b^2(-i + c + dx)(2de + if - cf + dfx) \arctan(c + dx) + \dots}{d^2}$$

input

```
Integrate[(e + f*x)*(a + b*ArcTan[c + d*x])^2,x]
```

output

```
(2*a^2*c*d*e - 2*a*b*c*f - a^2*c^2*f + 2*a^2*d^2*e*x - 2*a*b*d*f*x + a^2*d^2*f*x^2 + b^2*(-1 + c + d*x)*(2*d*e + 1*f - c*f + d*f*x)*ArcTan[c + d*x]^2 - 2*b*ArcTan[c + d*x]*(b*f*(c + d*x) + a*(-2*c*d*e + c^2*f - 2*d^2*e*x - f*(1 + d^2*x^2))) - 2*b*(d*e - c*f)*Log[1 + E^((2*I)*ArcTan[c + d*x])] + 4*a*b*d*e*Log[1/Sqrt[1 + (c + d*x)^2]] - 2*b^2*f*Log[1/Sqrt[1 + (c + d*x)^2]] - 4*a*b*c*f*Log[1/Sqrt[1 + (c + d*x)^2]] - (2*I)*b^2*(d*e - c*f)*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])])/(2*d^2)
```

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5570, 27, 5389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx$$

$$\downarrow \text{5570}$$

$$\int \frac{(d(e - \frac{cf}{d}) + f(c + dx))(a + b \arctan(c + dx))^2}{d} d(c + dx)$$

$$\downarrow \text{27}$$

$$\int \frac{(de - cf + f(c + dx))(a + b \arctan(c + dx))^2}{d^2} d(c + dx)$$

↓ 5389

$$\frac{(f(c+dx)-cf+de)^2(a+b\arctan(c+dx))^2}{2f} - \frac{b \int \left((a+b\arctan(c+dx))f^2 + \frac{((de-cf+f)(de-(c+1)f)+2f(de-cf)(c+dx))(a+b\arctan(c+dx))}{(c+dx)^2+1} \right) d(c+dx)}{d^2 f}$$

↓ 2009

$$\frac{(f(c+dx)-cf+de)^2(a+b\arctan(c+dx))^2}{2f} - \frac{b \left(-\frac{if(de-cf)(a+b\arctan(c+dx))^2}{b} + \frac{(-cf+de+f)(de-(c+1)f)(a+b\arctan(c+dx))^2}{2b} - 2f(de-cf) \log\left(\frac{1}{1+}\right) \right)}{d^2}$$

input `Int[(e + f*x)*(a + b*ArcTan[c + d*x])^2,x]`

output `((((d*e - c*f + f*(c + d*x))^2*(a + b*ArcTan[c + d*x])^2)/(2*f) - (b*(a*f^2*(c + d*x) + b*f^2*(c + d*x)*ArcTan[c + d*x] - (I*f*(d*e - c*f)*(a + b*ArcTan[c + d*x])^2)/b + ((d*e + f - c*f)*(d*e - (1 + c)*f)*(a + b*ArcTan[c + d*x])^2)/(2*b) - 2*f*(d*e - c*f)*(a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] - (b*f^2*Log[1 + (c + d*x)^2])/2 - I*b*f*(d*e - c*f)*PolyLog[2, 1 - 2/(1 + I*(c + d*x))]))/f)/d^2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5389 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))^ (p_.)*((d_.) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 5570

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^ (p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 413, normalized size of antiderivative = 1.86

method	result
parts	$a^2 \left(\frac{1}{2} f x^2 + e x \right) + \frac{b^2 \left(\frac{\arctan(dx+c)^2 (dx+c)^2 f}{2d} - \frac{\arctan(dx+c)^2 c f (dx+c)}{d} + \arctan(dx+c)^2 e (dx+c) - \frac{-\ln(1+(dx+c))}{d} \right)}{d}$
derivativedivides	$\frac{a^2 \left(f c (dx+c) - e d (dx+c) - \frac{f (dx+c)^2}{2} \right)}{d} - \frac{b^2 \left(\arctan(dx+c)^2 f c (dx+c) - \arctan(dx+c)^2 e d (dx+c) - \frac{\arctan(dx+c)^2 f (dx+c)^2}{2} - \ln(1+(dx+c)) \right)}{d}$
default	$\frac{a^2 \left(f c (dx+c) - e d (dx+c) - \frac{f (dx+c)^2}{2} \right)}{d} - \frac{b^2 \left(\arctan(dx+c)^2 f c (dx+c) - \arctan(dx+c)^2 e d (dx+c) - \frac{\arctan(dx+c)^2 f (dx+c)^2}{2} - \ln(1+(dx+c)) \right)}{d}$
risch	Expression too large to display

input

```
int((f*x+e)*(a+b*arctan(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
a^2*(1/2*f*x^2+e*x)+b^2/d*(1/2/d*arctan(d*x+c)^2*(d*x+c)^2*f-1/d*arctan(d*x+c)^2*c*f*(d*x+c)+arctan(d*x+c)^2*e*(d*x+c)-1/d*(-ln(1+(d*x+c)^2)*arctan(d*x+c)*c*f+ln(1+(d*x+c)^2)*arctan(d*x+c)*d*e-1/2*arctan(d*x+c)^2*f+arctan(d*x+c)*(d*x+c)*f-1/2*f*ln(1+(d*x+c)^2)-1/2*(-2*c*f+2*d*e)*(-1/2*I*(ln(d*x+c-I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(-1/2*I*(d*x+c+I)))+1/2*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-dilog(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))))+2*a*b/d*(1/2/d*arctan(d*x+c)*(d*x+c)^2*f-1/d*arctan(d*x+c)*c*f*(d*x+c)+arctan(d*x+c)*e*(d*x+c)-1/2/d*(f*(d*x+c)+1/2*(-2*c*f+2*d*e)*ln(1+(d*x+c)^2)-f*arctan(d*x+c)))
```

Fricas [F]

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx = \int (fx + e)(b \arctan(dx + c) + a)^2 dx$$

input

```
integrate((f*x+e)*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")
```

output

```
integral(a^2*f*x + a^2*e + (b^2*f*x + b^2*e)*arctan(d*x + c)^2 + 2*(a*b*f*x + a*b*e)*arctan(d*x + c), x)
```

Sympy [F]

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx = \int (a + b \operatorname{atan}(c + dx))^2 (e + fx) dx$$

input

```
integrate((f*x+e)*(a+b*atan(d*x+c))**2,x)
```

output

```
Integral((a + b*atan(c + d*x))**2*(e + f*x), x)
```


Maxima [F]

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx = \int (fx + e)(b \arctan(dx + c) + a)^2 dx$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output `3/4*b^2*c^2*e*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 1/4*(3*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*b^2*c^2*e + 12*b^2*d^2*f*integrate(1/16*x^3*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + b^2*d^2*f*integrate(1/16*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*d^2*e*integrate(1/16*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 24*b^2*c*d*f*integrate(1/16*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 2*b^2*d^2*f*integrate(1/16*x^3*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + b^2*d^2*e*integrate(1/16*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 2*b^2*c*d*f*integrate(1/16*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 24*b^2*c*d*e*integrate(1/16*x*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^2*c^2*f*integrate(1/16*x*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 4*b^2*d^2*e*integrate(1/16*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 2*b^2*c*d*f*integrate(1/16*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 2*b^2*c*d*e*integrate(1/16*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + b^2*c^2*f*integrate(1/16*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 4*b^2*c*d*e*integrate(1/16*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + ...`

Giac [F]

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx = \int (fx + e)(b \arctan(dx + c) + a)^2 dx$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c))^2,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx = \int (e + fx) (a + b \operatorname{atan}(c + dx))^2 dx$$

input `int((e + f*x)*(a + b*atan(c + d*x))^2,x)`output `int((e + f*x)*(a + b*atan(c + d*x))^2, x)`**Reduce [F]**

$$\int (e + fx)(a + b \arctan(c + dx))^2 dx$$

$$= \frac{\operatorname{atan}(dx + c)^2 b^2 c^2 f + 2 \operatorname{atan}(dx + c)^2 b^2 d^2 e x + \operatorname{atan}(dx + c)^2 b^2 d^2 f x^2 + \operatorname{atan}(dx + c)^2 b^2 f - 2 \operatorname{atan}(dx$$

input `int((f*x+e)*(a+b*atan(d*x+c))^2,x)`output `(atan(c + d*x)**2*b**2*c**2*f + 2*atan(c + d*x)**2*b**2*d**2*e*x + atan(c + d*x)**2*b**2*d**2*f*x**2 + atan(c + d*x)**2*b**2*f - 2*atan(c + d*x)*a*b*c**2*f + 4*atan(c + d*x)*a*b*c*d*e + 4*atan(c + d*x)*a*b*d**2*e*x + 2*atan(c + d*x)*a*b*d**2*f*x**2 + 2*atan(c + d*x)*a*b*f - 2*atan(c + d*x)*b**2*c*f - 2*atan(c + d*x)*b**2*d*f*x + 4*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*c*d**2*f - 4*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*d**3*e + 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*c*f - 2*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b*d*e + log(c**2 + 2*c*d*x + d**2*x**2 + 1)*b**2*f + 2*a**2*d**2*e*x + a**2*d**2*f*x**2 - 2*a*b*d*f*x)/(2*d**2)`

3.41 $\int (a + b \arctan(c + dx))^2 dx$

Optimal result	370
Mathematica [A] (verified)	371
Rubi [A] (verified)	371
Maple [A] (verified)	373
Fricas [F]	374
Sympy [F]	374
Maxima [F]	375
Giac [F]	375
Mupad [F(-1)]	376
Reduce [F]	376

Optimal result

Integrand size = 12, antiderivative size = 102

$$\int (a + b \arctan(c + dx))^2 dx = \frac{i(a + b \arctan(c + dx))^2}{d} + \frac{(c + dx)(a + b \arctan(c + dx))^2}{d} + \frac{2b(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d} + \frac{ib^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d}$$

output

```
I*(a+b*arctan(d*x+c))^2/d+(d*x+c)*(a+b*arctan(d*x+c))^2/d+2*b*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/d+I*b^2*polylog(2,1-2/(1+I*(d*x+c)))/d
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.07

$$\int (a + b \arctan(c + dx))^2 dx$$

$$= \frac{b^2(-i + c + dx) \arctan(c + dx)^2 + 2b \arctan(c + dx) (ac + adx + b \log(1 + e^{2i \arctan(c+dx)})) + a(ac + adx + b \log(1 + e^{2i \arctan(c+dx)}))}{d}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^2,x]
```

output

```
(b^2*(-I + c + d*x)*ArcTan[c + d*x]^2 + 2*b*ArcTan[c + d*x]*(a*c + a*d*x +
b*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + a*(a*c + a*d*x + 2*b*Log[1/Sqrt[1
+ (c + d*x)^2]]) - I*b^2*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])])/d
```

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5562, 5345, 5455, 5379, 2849, 2752}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \arctan(c + dx))^2 dx$$

$$\downarrow \text{5562}$$

$$\frac{\int (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow \text{5345}$$

$$\frac{(c + dx)(a + b \arctan(c + dx))^2 - 2b \int \frac{(c+dx)(a+b \arctan(c+dx))}{(c+dx)^2+1} d(c + dx)}{d}$$

$$\downarrow \text{5455}$$

$$\frac{(c + dx)(a + b \arctan(c + dx))^2 - 2b \left(- \int \frac{a+b \arctan(c+dx)}{-c-dx+i} d(c + dx) - \frac{i(a+b \arctan(c+dx))^2}{2b} \right)}{d}$$

↓ 5379

$$\frac{(c + dx)(a + b \arctan(c + dx))^2 - 2b \left(b \int \frac{\log\left(\frac{2}{i(c+dx)+1}\right) d(c + dx) - \frac{i(a+b \arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right) (a + b \arctan(c + dx)) \right)}{d}$$

↓ 2849

$$\frac{(c + dx)(a + b \arctan(c + dx))^2 - 2b \left(-ib \int \frac{\log\left(\frac{2}{1-i(c+dx)+1}\right) d \frac{1}{i(c+dx)+1} - \frac{i(a+b \arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right) (a + b \arctan(c + dx)) \right)}{d}$$

↓ 2752

$$\frac{(c + dx)(a + b \arctan(c + dx))^2 - 2b \left(-\frac{i(a+b \arctan(c+dx))^2}{2b} - \log\left(\frac{2}{1+i(c+dx)}\right) (a + b \arctan(c + dx)) - \frac{1}{2} ib \text{PolyLog}[2, 1 - 2/(1 + I*(c + d*x))] \right)}{d}$$

input `Int[(a + b*ArcTan[c + d*x])^2,x]`

output `((c + d*x)*(a + b*ArcTan[c + d*x])^2 - 2*b*(((1/2*I)*(a + b*ArcTan[c + d*x])^2)/b - (a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] - (I/2)*b*PolyLog[2, 1 - 2/(1 + I*(c + d*x))]))/d`

Defintions of rubi rules used

rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`

rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`

- rule 5345 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`
- rule 5379 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTan[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]`
- rule 5455 `Int[(((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Simp[1/(c*d) Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]`
- rule 5562 `Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[1/d Subst[Int[(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.36

method	result
parts	$a^2x + \frac{b^2 \left(\arctan(dx+c)^2(dx+c+i) + 2 \arctan(dx+c) \ln \left(1 + \frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) - 2i \arctan(dx+c)^2 - i \operatorname{polylog} \left(2, -\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) \right)}{d}$
derivativedivides	$\frac{(dx+c)a^2 - i \arctan(dx+c)^2 b^2 + \arctan(dx+c)^2 b^2 (dx+c) - i \operatorname{polylog} \left(2, -\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) b^2 + 2 \arctan(dx+c) \ln \left(1 + \frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) b^2}{d}$
default	$\frac{(dx+c)a^2 - i \arctan(dx+c)^2 b^2 + \arctan(dx+c)^2 b^2 (dx+c) - i \operatorname{polylog} \left(2, -\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) b^2 + 2 \arctan(dx+c) \ln \left(1 + \frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) b^2}{d}$
risch	$a^2x + \frac{i \ln(-idx-ic+1)abc}{d} - \frac{ab \ln(d^2x^2+2cdx+c^2+1)}{2d} + \frac{a^2c}{d} - \frac{i \ln(-idx-ic+1)^2 b^2}{4d} - \frac{iab \arctan(dx+c)}{d}$

input `int((a+b*arctan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `a^2*x+b^2/d*(arctan(d*x+c)^2*(d*x+c+I)+2*arctan(d*x+c)*ln(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))-2*I*arctan(d*x+c)^2-I*polylog(2,-(1+I*(d*x+c))^2/(1+(d*x+c)^2)))+2*a*b/d*((d*x+c)*arctan(d*x+c)-1/2*ln(1+(d*x+c)^2))`

Fricas [F]

$$\int (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 dx$$

input `integrate((a+b*arctan(d*x+c))^2,x, algorithm="fricas")`

output `integral(b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2, x)`

Sympy [F]

$$\int (a + b \arctan(c + dx))^2 dx = \int (a + b \operatorname{atan}(c + dx))^2 dx$$

input `integrate((a+b*atan(d*x+c))**2,x)`

output `Integral((a + b*atan(c + d*x))**2, x)`

Maxima [F]

$$\int (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 dx$$

input `integrate((a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output

```
1/16*(12*c^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 4*(3*arctan(d*x
+ c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*c^2 + 4*x
*arctan(d*x + c)^2 + 192*d^2*integrate(1/16*x^2*arctan(d*x + c)^2/(d^2*x^2
+ 2*c*d*x + c^2 + 1), x) + 16*d^2*integrate(1/16*x^2*log(d^2*x^2 + 2*c*d*x
+ c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 384*c*d*integrate(1/16*x
*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 64*d^2*integrate(1
/16*x^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x)
+ 32*c*d*integrate(1/16*x*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2
*c*d*x + c^2 + 1), x) + 64*c*d*integrate(1/16*x*log(d^2*x^2 + 2*c*d*x + c^
2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 16*c^2*integrate(1/16*log(d^2*x
^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) - x*log(d^2*x^
2 + 2*c*d*x + c^2 + 1)^2 + 12*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d
- 12*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d + 4*arctan((d^2*x + c*d)/
d)^3/d - 128*d*integrate(1/16*x*arctan(d*x + c)/(d^2*x^2 + 2*c*d*x + c^2 +
1), x) + 16*integrate(1/16*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 +
2*c*d*x + c^2 + 1), x))*b^2 + a^2*x + (2*(d*x + c)*arctan(d*x + c) - log((
d*x + c)^2 + 1))*a*b/d
```

Giac [F]

$$\int (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 dx$$

input `integrate((a+b*arctan(d*x+c))^2,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (a + b \arctan(c + dx))^2 dx = \int (a + b \operatorname{atan}(c + dx))^2 dx$$

input `int((a + b*atan(c + d*x))^2,x)`output `int((a + b*atan(c + d*x))^2, x)`**Reduce [F]**

$$\int (a + b \arctan(c + dx))^2 dx$$

$$= \frac{\operatorname{atan}(dx + c)^2 b^2 dx + 2 \operatorname{atan}(dx + c) abc + 2 \operatorname{atan}(dx + c) abdx - 2 \left(\int \frac{\operatorname{atan}(dx+c)x}{d^2 x^2 + 2cdx + c^2 + 1} dx \right) b^2 d^2 - \log(d^2 x^2 + 2cdx + c^2 + 1) ab^2}{d}$$

input `int((a+b*atan(d*x+c))^2,x)`output `(atan(c + d*x)**2*b**2*d*x + 2*atan(c + d*x)*a*b*c + 2*atan(c + d*x)*a*b*d*x - 2*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**2*d**2 - log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a*b + a**2*d*x)/d`

3.42 $\int \frac{(a+b \arctan(c+dx))^2}{e+fx} dx$

Optimal result	377
Mathematica [F]	378
Rubi [A] (verified)	378
Maple [C] (warning: unable to verify)	380
Fricas [F]	381
Sympy [F(-1)]	381
Maxima [F]	381
Giac [F]	382
Mupad [F(-1)]	382
Reduce [F]	382

Optimal result

Integrand size = 20, antiderivative size = 261

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx$$

$$= -\frac{(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1-i(c+dx)}\right)}{f}$$

$$+ \frac{(a + b \arctan(c + dx))^2 \log\left(\frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{f}$$

$$+ \frac{ib(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(c+dx)}\right)}{f}$$

$$- \frac{ib(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{f}$$

$$- \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1-i(c+dx)}\right)}{2f} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{2f}$$

output

```
-(a+b*arctan(d*x+c))^2*ln(2/(1-I*(d*x+c)))/f+(a+b*arctan(d*x+c))^2*ln(2*d*(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f+I*b*(a+b*arctan(d*x+c))*polylog(2,1-2/(1-I*(d*x+c)))/f-I*b*(a+b*arctan(d*x+c))*polylog(2,1-2*d*(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f-1/2*b^2*polylog(3,1-2/(1-I*(d*x+c)))/f+1/2*b^2*polylog(3,1-2*d*(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f
```

Mathematica [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx$$

input `Integrate[(a + b*ArcTan[c + d*x])^2/(e + f*x), x]`

output `Integrate[(a + b*ArcTan[c + d*x])^2/(e + f*x), x]`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {5570, 27, 5383}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx \\ & \quad \downarrow \text{5570} \\ & \int \frac{d(a + b \arctan(c + dx))^2}{d(e - \frac{cf}{d}) + f(c + dx)} d(c + dx) \\ & \quad \downarrow \text{27} \\ & \int \frac{(a + b \arctan(c + dx))^2}{f(c + dx) - cf + de} d(c + dx) \\ & \quad \downarrow \text{5383} \end{aligned}$$

$$\begin{aligned}
& \frac{ib(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + if)(1 - i(c + dx))}\right)}{f} + \\
& \frac{(a + b \arctan(c + dx))^2 \log\left(\frac{2(f(c + dx) - cf + de)}{(1 - i(c + dx))(-cf + de + if)}\right)}{f} + \\
& \frac{ib \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i(c + dx)}\right) (a + b \arctan(c + dx))}{f} - \\
& \frac{\log\left(\frac{2}{1 - i(c + dx)}\right) (a + b \arctan(c + dx))^2}{f} + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + if)(1 - i(c + dx))}\right)}{2f} - \\
& \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - i(c + dx)}\right)}{2f}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^2/(e + f*x), x]`

output `-(((a + b*ArcTan[c + d*x])^2*Log[2/(1 - I*(c + d*x))])/f) + ((a + b*ArcTan[c + d*x])^2*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/f + (I*b*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 - I*(c + d*x))])/f - (I*b*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x)))]/f - (b^2*PolyLog[3, 1 - 2/(1 - I*(c + d*x))]/(2*f) + (b^2*PolyLog[3, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x)))]/(2*f)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 5383 `Int[((a_) + ArcTan[(c_)*(x_)]*(b_.))^2/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTan[c*x])^2)*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*ArcTan[c*x])^2*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] + Simp[I*b*(a + b*ArcTan[c*x])*PolyLog[2, 1 - 2/(1 - I*c*x)]/e, x] - Simp[I*b*(a + b*ArcTan[c*x])*PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e, x] - Simp[b^2*(PolyLog[3, 1 - 2/(1 - I*c*x)]/(2*e)), x] + Simp[b^2*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]`

rule 5570

```
Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.29 (sec) , antiderivative size = 1877, normalized size of antiderivative = 7.19

method	result	size
derivativeldivides	Expression too large to display	1877
default	Expression too large to display	1877
parts	Expression too large to display	1998

input

```
int((a+b*arctan(d*x+c))^2/(f*x+e),x,method=_RETURNVERBOSE)
```

output

```
1/d*(a^2*d*ln(c*f-d*e-f*(d*x+c))/f-b^2*d*(-ln(c*f-d*e-f*(d*x+c))/f*arctan(
d*x+c)^2+2/f*(1/2*arctan(d*x+c)^2*ln(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f
*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d
*e)-1/4*I*Pi*csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2
/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)/(1+(1+I*(d*x
+c))^2/(1+(d*x+c)^2)))*(csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I
*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e))*
csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(
d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)
^2)-I*f+c*f-d*e)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*csgn(I/(1+(1+I*(d*x+c)
)^2/(1+(d*x+c)^2)))-csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d
*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e))*csgn
(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d
e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)
^2)))+csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d
*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)/(1+(1+I*(d*x+c))^2/
(1+(d*x+c)^2)))^2*arctan(d*x+c)^2-1/2*I*arctan(d*x+c)*polylog(2,-(1+I*(d*
x+c))^2/(1+(d*x+c)^2))+1/4*polylog(3,-(1+I*(d*x+c))^2/(1+(d*x+c)^2))-1/2*f
/(c*f-d*e+I*f)*arctan(d*x+c)*polylog(2,(c*f-d*e+I*f)*(1+I*(d*x+c))^2/(1+(d
*x+c)^2)/(d*e+I*f-c*f))-1/2*I*f/(c*f-d*e+I*f)*arctan(d*x+c)^2*ln(1-(c*f...
```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(f*x+e),x, algorithm="fricas")`

output `integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)/(f*x + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))**2/(f*x+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(f*x+e),x, algorithm="maxima")`

output `a^2*log(f*x + e)/f + integrate(1/16*(12*b^2*arctan(d*x + c)^2 + b^2*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 32*a*b*arctan(d*x + c))/(f*x + e), x)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^2}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(f*x+e),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^2/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^2}{e + fx} dx$$

input `int((a + b*atan(c + d*x))^2/(e + f*x),x)`

output `int((a + b*atan(c + d*x))^2/(e + f*x), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(a + b \arctan(c + dx))^2}{e + fx} dx \\ &= \frac{2 \left(\int \frac{\operatorname{atan}(dx+c)}{fx+e} dx \right) abf + \left(\int \frac{\operatorname{atan}(dx+c)^2}{fx+e} dx \right) b^2 f + \log(fx + e) a^2}{f} \end{aligned}$$

input `int((a+b*atan(d*x+c))^2/(f*x+e),x)`

output `(2*int(atan(c + d*x)/(e + f*x),x)*a*b*f + int(atan(c + d*x)**2/(e + f*x),x)
)*b**2*f + log(e + f*x)*a**2)/f`

3.43 $\int \frac{(a+b \arctan(c+dx))^2}{(e+fx)^2} dx$

Optimal result	383
Mathematica [A] (warning: unable to verify)	384
Rubi [A] (verified)	385
Maple [A] (verified)	387
Fricas [F]	389
Sympy [F(-1)]	389
Maxima [F]	390
Giac [F]	390
Mupad [F(-1)]	391
Reduce [F]	391

Optimal result

Integrand size = 20, antiderivative size = 480

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \frac{id(a + b \arctan(c + dx))^2}{d^2e^2 - 2cdef + (1 + c^2) f^2} + \frac{d(de - cf)(a + b \arctan(c + dx))^2}{f(d^2e^2 - 2cdef + (1 + c^2) f^2)} - \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} - \frac{2bd(a + b \arctan(c + dx)) \log\left(\frac{2}{1-i(c+dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2} + \frac{2bd(a + b \arctan(c + dx)) \log\left(\frac{2d(e+fx)}{(de+if-cf)(1-i(c+dx))}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2} + \frac{2bd(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2} + \frac{ib^2d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(c+dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2} - \frac{ib^2d \operatorname{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+if-cf)(1-i(c+dx))}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2} + \frac{ib^2d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2) f^2}$$

output

```
I*d*(a+b*arctan(d*x+c))^2/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+d*(-c*f+d*e)*(a+b*arctan(d*x+c))^2/f/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-(a+b*arctan(d*x+c))^2/f/(f*x+e)-2*b*d*(a+b*arctan(d*x+c))*ln(2/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+2*b*d*(a+b*arctan(d*x+c))*ln(2*d*(f*x+e)/(d*e+I*f-c*f)/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+2*b*d*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+I*b^2*d*polylog(2,1-2/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-I*b^2*d*polylog(2,1-2*d*(f*x+e)/(d*e+I*f-c*f)/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+I*b^2*d*polylog(2,1-2/(1+I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)
```

Mathematica [A] (warning: unable to verify)

Time = 4.54 (sec) , antiderivative size = 419, normalized size of antiderivative = 0.87

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx$$

$$= \frac{-\frac{a^2}{f} + \frac{2ab \left(-((-cde + f + c^2 f - d^2 ex + cdfx) \arctan(c + dx)) + d(e + fx) \log\left(\frac{d(e + fx)}{\sqrt{1 + (c + dx)^2}}\right) \right)}{d^2 e^2 - 2cde f + (1 + c^2) f^2}}{f} + \frac{b^2 d(e + fx) \left(-\frac{e^{i \arctan\left(\frac{de - cf}{f}\right)} \arctan(c + dx)}{f \sqrt{1 + \frac{(de - cf)^2}{f^2}}} \right)}{f^2}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^2/(e + f*x)^2,x]
```

output

```
(-(a^2/f) + (2*a*b*(-((-c*d*e) + f + c^2*f - d^2*e*x + c*d*f*x)*ArcTan[c + d*x]) + d*(e + f*x)*Log[(d*(e + f*x))/Sqrt[1 + (c + d*x)^2]])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b^2*d*(e + f*x)*(-(E^(I*ArcTan[(d*e - c*f)/f]))*ArcTan[c + d*x]^2)/(f*Sqrt[1 + (d*e - c*f)^2/f^2])) + ((c + d*x)*ArcTan[c + d*x]^2)/(d*(e + f*x)) - ((d*e - c*f)*((-I)*(Pi - 2*ArcTan[(d*e - c*f)/f]))*ArcTan[c + d*x] - Pi*Log[1 + E^((-2*I)*ArcTan[c + d*x])] - 2*(ArcTan[(d*e - c*f)/f] + ArcTan[c + d*x])*Log[1 - E^((2*I)*(ArcTan[(d*e - c*f)/f] + ArcTan[c + d*x]))] + Pi*Log[1/Sqrt[1 + (c + d*x)^2]] + 2*ArcTan[(d*e - c*f)/f]*Log[Sin[ArcTan[(d*e - c*f)/f] + ArcTan[c + d*x]]) + I*PolyLog[2, E^((2*I)*(ArcTan[(d*e - c*f)/f] + ArcTan[c + d*x]))])/(f^2*(1 + (d*e - c*f)^2/f^2)))/(d*e - c*f)/(e + f*x)
```

Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 586, normalized size of antiderivative = 1.22, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5568, 7292, 5580, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx \\
 & \quad \downarrow \text{5568} \\
 & \frac{2bd \int \frac{a+b \arctan(c+dx)}{(e+fx)((c+dx)^2+1)} dx}{f} - \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{7292} \\
 & \frac{2bd \int \frac{a+b \arctan(c+dx)}{(e+fx)(c^2+2dxc+d^2x^2+1)} dx}{f} - \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{5580} \\
 & \frac{2b \int \frac{d(a+b \arctan(c+dx))}{(d(e-\frac{cf}{d})+f(c+dx))((c+dx)^2+1)} d(c + dx)}{f} - \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2bd \int \frac{a+b \arctan(c+dx)}{(de-cf+f(c+dx))((c+dx)^2+1)} d(c + dx)}{f} - \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{7276} \\
 & \frac{2bd \int \left(\frac{a}{(de-cf+f(c+dx))((c+dx)^2+1)} + \frac{b \arctan(c+dx)}{(de-cf+f(c+dx))((c+dx)^2+1)} \right) d(c + dx)}{f} - \\
 & \quad \frac{(a + b \arctan(c + dx))^2}{f(e + fx)} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$2bd \left(\frac{(a + b \arctan(c + dx))^2}{f(e + fx)} + \frac{a \arctan(c+dx)(de-cf)}{(de-cf)^2+f^2} + \frac{af \log(f(c+dx)-cf+de)}{(de-cf)^2+f^2} - \frac{af \log((c+dx)^2+1)}{2((de-cf)^2+f^2)} + \frac{ibf \arctan(c+dx)^2}{2((c^2+1)f^2-2cdef+d^2e^2)} + \frac{b \arctan(c+dx)^2(de-cf)}{2((c^2+1)f^2-2cdef+d^2e^2)} \right)$$

input `Int[(a + b*ArcTan[c + d*x])^2/(e + f*x)^2,x]`

output `-((a + b*ArcTan[c + d*x])^2/(f*(e + f*x))) + (2*b*d*((a*(d*e - c*f)*ArcTan[c + d*x])/(f^2 + (d*e - c*f)^2) + ((I/2)*b*f*ArcTan[c + d*x]^2)/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b*(d*e - c*f)*ArcTan[c + d*x]^2)/(2*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)) - (b*f*ArcTan[c + d*x]*Log[2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b*f*ArcTan[c + d*x]*Log[2/(1 + I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (a*f*Log[d*e - c*f + f*(c + d*x)])/(f^2 + (d*e - c*f)^2) + (b*f*ArcTan[c + d*x]*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) - (a*f*Log[1 + (c + d*x)^2])/(2*(f^2 + (d*e - c*f)^2)) + ((I/2)*b*f*PolyLog[2, 1 - 2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + ((I/2)*b*f*PolyLog[2, 1 - 2/(1 + I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) - ((I/2)*b*f*PolyLog[2, 1 - (2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x)))]/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2))/f`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5568 `Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_.)]*(b_.))^p_*(e_.) + (f_.)*(x_.)^m_], x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]`

rule 5580

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Subst
[Int[(d*e - c*f)/d + f*(x/d)]^m*(C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTan[x])
^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x] &
& EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [A] (verified)

Time = 1.44 (sec) , antiderivative size = 776, normalized size of antiderivative = 1.62

method	result
parts	$-\frac{a^2}{(fx+e)f} + b^2 \left(-\frac{d^2 \arctan(dx+c)^2}{(f(dx+c)-cf+de)f} + 2d^2 \left(-\frac{\arctan(dx+c)f \ln(1+(dx+c)^2)}{2(c^2f^2-2cdef+d^2e^2+f^2)} - \frac{\arctan(dx+c)^2cf}{c^2f^2-2cdef+d^2e^2+f^2} + \frac{\arctan(dx+c)}{c^2f^2-2cdef+d^2e^2+f^2} \right) \right)$
derivativedivides	$\frac{a^2d^2}{(cf-de-f(dx+c))f} + b^2d^2 \left(\frac{\arctan(dx+c)^2}{(cf-de-f(dx+c))f} - 2 \left(-\frac{\arctan(dx+c)f \ln(cf-de-f(dx+c))}{c^2f^2-2cdef+d^2e^2+f^2} + \frac{\arctan(dx+c)f \ln(1+(dx+c)^2)}{2c^2f^2-4cdef+2d^2e^2+2f^2} + \frac{\arctan(dx+c)}{c^2f^2-2cdef+d^2e^2+f^2} \right) \right)$
default	$\frac{a^2d^2}{(cf-de-f(dx+c))f} + b^2d^2 \left(\frac{\arctan(dx+c)^2}{(cf-de-f(dx+c))f} - 2 \left(-\frac{\arctan(dx+c)f \ln(cf-de-f(dx+c))}{c^2f^2-2cdef+d^2e^2+f^2} + \frac{\arctan(dx+c)f \ln(1+(dx+c)^2)}{2c^2f^2-4cdef+2d^2e^2+2f^2} + \frac{\arctan(dx+c)}{c^2f^2-2cdef+d^2e^2+f^2} \right) \right)$

```
input int((a+b*arctan(d*x+c))^2/(f*x+e)^2,x,method=_RETURNVERBOSE)
```

output

```
-a^2/(f*x+e)/f+b^2/d*(-d^2/(f*(d*x+c)-c*f+d*e)/f*arctan(d*x+c)^2+2*d^2/f*(
-1/2*arctan(d*x+c)/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f*ln(1+(d*x+c)^2)-1/(c^
2*f^2-2*c*d*e*f+d^2*e^2+f^2)*arctan(d*x+c)^2*c*f+1/(c^2*f^2-2*c*d*e*f+d^2*
e^2+f^2)*arctan(d*x+c)^2*d*e+arctan(d*x+c)/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)
*f*ln(f*(d*x+c)-c*f+d*e)-1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f^2*(-1/2*I*ln(
f*(d*x+c)-c*f+d*e)*(ln((I*f-f*(d*x+c))/(d*e+I*f-c*f))-ln((I*f+f*(d*x+c))/(
c*f-d*e+I*f)))/f-1/2*I*(dilog((I*f-f*(d*x+c))/(d*e+I*f-c*f))-dilog((I*f+f*
(d*x+c))/(c*f-d*e+I*f)))/f)+1/2/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*(c*f-d*e)*
arctan(d*x+c)^2+1/2/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f*(-1/2*I*(ln(d*x+c-I)
*ln(1+(d*x+c)^2)-1/2*ln(d*x+c-I)^2-dilog(-1/2*I*(d*x+c+I))-ln(d*x+c-I)*ln(
-1/2*I*(d*x+c+I)))+1/2*I*(ln(d*x+c+I)*ln(1+(d*x+c)^2)-1/2*ln(d*x+c+I)^2-di
log(1/2*I*(d*x+c-I))-ln(d*x+c+I)*ln(1/2*I*(d*x+c-I)))))+2*a*b/d*(-d^2/(f*
(d*x+c)-c*f+d*e)/f*arctan(d*x+c)+d^2/f*(1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*
(-1/2*f*ln(1+(d*x+c)^2)+(-c*f+d*e)*arctan(d*x+c))+1/(c^2*f^2-2*c*d*e*f+d^2
*e^2+f^2)*f*ln(f*(d*x+c)-c*f+d*e)))
```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(fx + e)^2} dx$$

input

```
integrate((a+b*arctan(d*x+c))^2/(f*x+e)^2,x, algorithm="fricas")
```

output

```
integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)/(f^2*x^2 +
2*e*f*x + e^2), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \text{Timed out}$$

input

```
integrate((a+b*atan(d*x+c))**2/(f*x+e)**2,x)
```

output Timed out

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(f*x+e)^2,x, algorithm="maxima")`

output `(d*(2*(d^2*e - c*d*f)*arctan((d^2*x + c*d)/d)/((d^2*e^2*f - 2*c*d*e*f^2 + (c^2 + 1)*f^3)*d) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2) + 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2)) - 2*arctan(d*x + c)/(f^2*x + e*f))*a*b - 1/16*(4*arctan(d*x + c)^2 - 16*(f^2*x + e*f)*integrate(1/16*(12*(d^2*f*x^2 + 2*c*d*f*x + (c^2 + 1)*f)*arctan(d*x + c)^2 + (d^2*f*x^2 + 2*c*d*f*x + (c^2 + 1)*f)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 8*(d*f*x + d*e)*arctan(d*x + c) - 4*(d^2*f*x^2 + c*d*e + (d^2*e + c*d*f)*x)*log(d^2*x^2 + 2*c*d*x + c^2 + 1))/(d^2*f^3*x^4 + (c^2 + 1)*e^2*f + 2*(d^2*e*f^2 + c*d*f^3)*x^3 + (d^2*e^2*f + 4*c*d*e*f^2 + (c^2 + 1)*f^3)*x^2 + 2*(c*d*e^2*f + (c^2 + 1)*e*f^2)*x), x) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2)*b^2/(f^2*x + e*f) - a^2/(f^2*x + e*f)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^2}{(fx + e)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^2/(f*x+e)^2,x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^2/(f*x + e)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^2}{(e + fx)^2} dx$$

input `int((a + b*atan(c + d*x))^2/(e + f*x)^2,x)`output `int((a + b*atan(c + d*x))^2/(e + f*x)^2, x)`**Reduce [F]**

$$\int \frac{(a + b \arctan(c + dx))^2}{(e + fx)^2} dx = \text{too large to display}$$

input `int((a+b*atan(d*x+c))^2/(f*x+e)^2,x)`

output

```
( - atan(c + d*x)**2*b**2*c**4*e*f**3 + 2*atan(c + d*x)**2*b**2*c**3*d*e**
2*f**2 - atan(c + d*x)**2*b**2*c**2*d**2*e**3*f - atan(c + d*x)**2*b**2*c*
**2*d**2*e**2*f**2*x - 2*atan(c + d*x)**2*b**2*c**2*e*f**3 + 2*atan(c + d*x
)**2*b**2*c*d**3*e**3*f*x + 2*atan(c + d*x)**2*b**2*c*d*e**2*f**2 - atan(c
+ d*x)**2*b**2*d**4*e**4*x - atan(c + d*x)**2*b**2*d**2*e**3*f - atan(c +
d*x)**2*b**2*d**2*e**2*f**2*x - atan(c + d*x)**2*b**2*e*f**3 - 2*atan(c +
d*x)*a*b*c**4*e*f**3 + 2*atan(c + d*x)*a*b*c**3*d*e**2*f**2 - 2*atan(c +
d*x)*a*b*c**3*d*e*f**3*x + 2*atan(c + d*x)*a*b*c**2*d**2*e**3*f + 2*atan(c
+ d*x)*a*b*c**2*d**2*e**2*f**2*x - 4*atan(c + d*x)*a*b*c**2*e*f**3 - 2*at
an(c + d*x)*a*b*c*d**3*e**4 + 2*atan(c + d*x)*a*b*c*d**3*e**3*f*x + 2*atan
(c + d*x)*a*b*c*d*e**2*f**2 - 2*atan(c + d*x)*a*b*c*d*e*f**3*x - 2*atan(c
+ d*x)*a*b*d**4*e**4*x + 2*atan(c + d*x)*a*b*d**2*e**3*f + 2*atan(c + d*x)
*a*b*d**2*e**2*f**2*x - 2*atan(c + d*x)*a*b*e*f**3 - 2*atan(c + d*x)*b**2*
c**2*d*e**2*f**2 + 2*atan(c + d*x)*b**2*c*d**2*e**3*f - 2*atan(c + d*x)*b
**2*c*d**2*e**2*f**2*x + 2*atan(c + d*x)*b**2*d**3*e**3*f*x - 2*atan(c + d
*x)*b**2*d*e**2*f**2 + 2*int((atan(c + d*x)*x)/(c**4*e**2*f**2 + 2*c**4*e*f
**3*x + c**4*f**4*x**2 + 2*c**3*d*e**2*f**2*x + 4*c**3*d*e*f**3*x**2 + 2*c
**3*d*f**4*x**3 - c**2*d**2*e**4 - 2*c**2*d**2*e**3*f*x + 2*c**2*d**2*e*f*
**3*x**3 + c**2*d**2*f**4*x**4 + 2*c**2*e**2*f**2 + 4*c**2*e*f**3*x + 2*c**
2*f**4*x**2 - 2*c*d**3*e**4*x - 4*c*d**3*e**3*f*x**2 - 2*c*d**3*e**2*f*...
```

3.44 $\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx$

Optimal result	394
Mathematica [B] (warning: unable to verify)	395
Rubi [A] (verified)	396
Maple [C] (warning: unable to verify)	398
Fricas [F]	399
Sympy [F]	399
Maxima [F]	399
Giac [F]	400
Mupad [F(-1)]	401
Reduce [F]	401

Optimal result

Integrand size = 20, antiderivative size = 564

$$\begin{aligned}
\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx &= \frac{ab^2 f^2 x}{d^2} + \frac{b^3 f^2 (c + dx) \arctan(c + dx)}{d^3} \\
&- \frac{bf^2 (a + b \arctan(c + dx))^2}{2d^3} - \frac{3ibf(de - cf)(a + b \arctan(c + dx))^2}{d^3} \\
&- \frac{3bf(de - cf)(c + dx)(a + b \arctan(c + dx))^2}{d^3} - \frac{bf^2(c + dx)^2(a + b \arctan(c + dx))^2}{2d^3} \\
&+ \frac{i(3d^2e^2 - 6cdef - (1 - 3c^2)f^2)(a + b \arctan(c + dx))^3}{3d^3} \\
&- \frac{(de - cf)(d^2e^2 - 2cdef - (3 - c^2)f^2)(a + b \arctan(c + dx))^3}{3d^3 f} \\
&+ \frac{(e + fx)^3 (a + b \arctan(c + dx))^3}{3f} \\
&- \frac{6b^2 f(de - cf)(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d^3} \\
&+ \frac{b(3d^2e^2 - 6cdef - (1 - 3c^2)f^2)(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1+i(c+dx)}\right)}{d^3} \\
&- \frac{b^3 f^2 \log(1 + (c + dx)^2)}{2d^3} - \frac{3ib^3 f(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d^3} \\
&+ \frac{ib^2(3d^2e^2 - 6cdef - (1 - 3c^2)f^2)(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d^3} \\
&+ \frac{b^3(3d^2e^2 - 6cdef - (1 - 3c^2)f^2) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2d^3}
\end{aligned}$$

output

```

a*b^2*f^2*x/d^2+b^3*f^2*(d*x+c)*arctan(d*x+c)/d^3-1/2*b*f^2*(a+b*arctan(d*
x+c))^2/d^3-3*I*b*f*(-c*f+d*e)*(a+b*arctan(d*x+c))^2/d^3-3*b*f*(-c*f+d*e)*
(d*x+c)*(a+b*arctan(d*x+c))^2/d^3-1/2*b*f^2*(d*x+c)^2*(a+b*arctan(d*x+c))^
2/d^3+1/3*I*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*(a+b*arctan(d*x+c))^3/d^3
-1/3*(-c*f+d*e)*(d^2*e^2-2*c*d*e*f-(-c^2+3)*f^2)*(a+b*arctan(d*x+c))^3/d^3
/f+1/3*(f*x+e)^3*(a+b*arctan(d*x+c))^3/f-6*b^2*f*(-c*f+d*e)*(a+b*arctan(d*
x+c))*ln(2/(1+I*(d*x+c)))/d^3+b*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*(a+b*
arctan(d*x+c))^2*ln(2/(1+I*(d*x+c)))/d^3-1/2*b^3*f^2*ln(1+(d*x+c)^2)/d^3-3
*I*b^3*f*(-c*f+d*e)*polylog(2,1-2/(1+I*(d*x+c)))/d^3+I*b^2*(3*d^2*e^2-6*c*
d*e*f-(-3*c^2+1)*f^2)*(a+b*arctan(d*x+c))*polylog(2,1-2/(1+I*(d*x+c)))/d^3
+1/2*b^3*(3*d^2*e^2-6*c*d*e*f-(-3*c^2+1)*f^2)*polylog(3,1-2/(1+I*(d*x+c)))
/d^3

```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1844 vs. $2(564) = 1128$.

Time = 13.52 (sec) , antiderivative size = 1844, normalized size of antiderivative = 3.27

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \text{Too large to display}$$

input

```
Integrate[(e + f*x)^2*(a + b*ArcTan[c + d*x])^3,x]
```

output

```
(a^2*(a*d^2*e^2 - 3*b*d*e*f + 2*b*c*f^2)*x)/d^2 - (a^2*f*(-2*a*d*e + b*f)*
x^2)/(2*d) + (a^3*f^2*x^3)/3 + ((3*a^2*b*c*d^2*e^2 + 3*a^2*b*d*e*f - 3*a^2
*b*c^2*d*e*f - 3*a^2*b*c*f^2 + a^2*b*c^3*f^2)*ArcTan[c + d*x])/d^3 + a^2*b
*x*(3*e^2 + 3*e*f*x + f^2*x^2)*ArcTan[c + d*x] + ((-3*a^2*b*d^2*e^2 + 6*a^
2*b*c*d*e*f + a^2*b*f^2 - 3*a^2*b*c^2*f^2)*Log[1 + c^2 + 2*c*d*x + d^2*x^2
])/d^3 + (3*a*b^2*e^2*((-I)*ArcTan[c + d*x]^2 + (c + d*x)*ArcTan[c + d
*x]^2 + 2*ArcTan[c + d*x]*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - I*PolyLog[2
, -E^((2*I)*ArcTan[c + d*x])])/d + 6*a*b^2*e*f*(-(((c + d*x)*ArcTan[c + d
*x])/d^2) + (I*c*ArcTan[c + d*x]^2)/d^2 - (c*(c + d*x)*ArcTan[c + d*x]^2)/
d^2 + ((1 + (c + d*x)^2)*ArcTan[c + d*x]^2)/(2*d^2) - (2*c*ArcTan[c + d*x]
*Log[1 + E^((2*I)*ArcTan[c + d*x])])/d^2 - Log[1/Sqrt[1 + (c + d*x)^2]]/d^
2 + (I*c*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])])/d^2 + (b^3*e^2*((-I)*Arc
Tan[c + d*x]^3 + (c + d*x)*ArcTan[c + d*x]^3 + 3*ArcTan[c + d*x]^2*Log[1 +
E^((2*I)*ArcTan[c + d*x])]) - (3*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*A
rcTan[c + d*x])]) + (3*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])])/2)/d + (b^3
*e*f*(ArcTan[c + d*x]*((3*I)*ArcTan[c + d*x] + (2*I)*c*ArcTan[c + d*x]^2 +
(1 + (c + d*x)^2)*ArcTan[c + d*x]^2 - (c + d*x)*ArcTan[c + d*x]*(3 + 2*c*
ArcTan[c + d*x]) - 6*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - 6*c*ArcTan[c + d
*x]*Log[1 + E^((2*I)*ArcTan[c + d*x])]) + (3*I)*(1 + 2*c*ArcTan[c + d*x])*
PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) - 3*c*PolyLog[3, -E^((2*I)*ArcTa...
```

Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 554, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5570, 27, 5389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx$$

$$\downarrow 5570$$

$$\int \frac{\left(d\left(e - \frac{ef}{d}\right) + f(c + dx)\right)^2 (a + b \arctan(c + dx))^3}{d^2} d(c + dx)$$

$$\downarrow 27$$

$$\frac{\int (de - cf + f(c + dx))^2 (a + b \arctan(c + dx))^3 d(c + dx)}{d^3}$$

↓ 5389

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))^3}{3f} - \frac{b \int \left((c+dx)(a+b \arctan(c+dx))^2 f^3 + 3(de-cf)(a+b \arctan(c+dx))^2 f^2 + \frac{(de-cf)(d^2e^2-2cdf e-3c^2d^2)}{f} \right) d(c+dx)}{d^3}}{f}$$

↓ 2009

$$\frac{\frac{(f(c+dx)-cf+de)^3(a+b \arctan(c+dx))^3}{3f} - \frac{b \left(-ibf(-1-3c^2)f^2 - 6cdf + 3d^2e^2 \right) \text{PolyLog}\left(2, 1 - \frac{2}{i(c+dx)+1}\right) (a+b \arctan(c+dx)) - \frac{if(-1-3c^2)}{f}}{d^3}}{f}$$

input `Int[(e + f*x)^2*(a + b*ArcTan[c + d*x])^3,x]`

output `((d*e - c*f + f*(c + d*x))^3*(a + b*ArcTan[c + d*x])^3)/(3*f) - (b*(-a*b*f^3*(c + d*x) - b^2*f^3*(c + d*x)*ArcTan[c + d*x] + (f^3*(a + b*ArcTan[c + d*x])^2)/2 + (3*I)*f^2*(d*e - c*f)*(a + b*ArcTan[c + d*x])^2 + 3*f^2*(d*e - c*f)*(c + d*x)*(a + b*ArcTan[c + d*x])^2 + (f^3*(c + d*x)^2*(a + b*ArcTan[c + d*x])^2)/2 - ((I/3)*f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*(a + b*ArcTan[c + d*x])^3)/b + ((d*e - c*f)*(d^2*e^2 - 2*c*d*e*f - (3 - c^2)*f^2)*(a + b*ArcTan[c + d*x])^3)/(3*b) + 6*b*f^2*(d*e - c*f)*(a + b*ArcTan[c + d*x])*Log[2/(1 + I*(c + d*x))] - f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*(a + b*ArcTan[c + d*x])^2*Log[2/(1 + I*(c + d*x))] + (b^2*f^3*Log[1 + (c + d*x)^2])/2 + (3*I)*b^2*f^2*(d*e - c*f)*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] - I*b*f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] - (b^2*f*(3*d^2*e^2 - 6*c*d*e*f - (1 - 3*c^2)*f^2)*PolyLog[3, 1 - 2/(1 + I*(c + d*x))])/2)/f/d^3`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5389 `Int[((a_) + ArcTan[(c_)*(x_)]*(b_))^(p_)*((d_) + (e_)*(x_))^(q_), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Simp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]`

rule 5570 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && IGtQ[p, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 12.27 (sec) , antiderivative size = 5843, normalized size of antiderivative = 10.36

method	result	size
derivativedivides	Expression too large to display	5843
default	Expression too large to display	5843
parts	Expression too large to display	6026

input `int((f*x+e)^2*(a+b*arctan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^3 dx$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="fricas")`

output `integral(a^3*f^2*x^2 + 2*a^3*e*f*x + a^3*e^2 + (b^3*f^2*x^2 + 2*b^3*e*f*x + b^3*e^2)*arctan(d*x + c)^3 + 3*(a*b^2*f^2*x^2 + 2*a*b^2*e*f*x + a*b^2*e^2)*arctan(d*x + c)^2 + 3*(a^2*b*f^2*x^2 + 2*a^2*b*e*f*x + a^2*b*e^2)*arctan(d*x + c), x)`

Sympy [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \int (a + b \operatorname{atan}(c + dx))^3 (e + fx)^2 dx$$

input `integrate((f*x+e)**2*(a+b*atan(d*x+c))**3,x)`

output `Integral((a + b*atan(c + d*x))**3*(e + f*x)**2, x)`

Maxima [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^3 dx$$

input `integrate((f*x+e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output

```

7/8*b^3*c^2*e^2*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 3*a*b^2*c^2*
e^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - (3*arctan(d*x + c)*arcta
n((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*a*b^2*c^2*e^2 - 7/32
*(6*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)^2/d - 4*arctan(d*x + c)*arct
an((d^2*x + c*d)/d)^3/d + arctan((d^2*x + c*d)/d)^4/d)*b^3*c^2*e^2 + 1/3*a
^3*f^2*x^3 + 7/8*b^3*e^2*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 28*
b^3*d^2*f^2*integrate(1/32*x^4*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2
+ 1), x) + 3*b^3*d^2*f^2*integrate(1/32*x^4*arctan(d*x + c)*log(d^2*x^2 +
2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 96*a*b^2*d^2*f^2*
integrate(1/32*x^4*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 5
6*b^3*d^2*e*f*integrate(1/32*x^3*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^
2 + 1), x) + 56*b^3*c*d*f^2*integrate(1/32*x^3*arctan(d*x + c)^3/(d^2*x^2
+ 2*c*d*x + c^2 + 1), x) + 4*b^3*d^2*f^2*integrate(1/32*x^4*arctan(d*x + c
)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b
^3*d^2*e*f*integrate(1/32*x^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2
+ 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b^3*c*d*f^2*integrate(1/32*x^
3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x +
c^2 + 1), x) + 192*a*b^2*d^2*e*f*integrate(1/32*x^3*arctan(d*x + c)^2/(d^2
*x^2 + 2*c*d*x + c^2 + 1), x) + 192*a*b^2*c*d*f^2*integrate(1/32*x^3*arcta
n(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 28*b^3*d^2*e^2*integra...

```

Giac [F]

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \int (fx + e)^2 (b \arctan(dx + c) + a)^3 dx$$

input

```
integrate((f*x+e)^2*(a+b*arctan(d*x+c))^3,x, algorithm="giac")
```

output

```
sage0*x
```

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \int (e + fx)^2 (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((e + f*x)^2*(a + b*atan(c + d*x))^3,x)`output `int((e + f*x)^2*(a + b*atan(c + d*x))^3, x)`**Reduce [F]**

$$\int (e + fx)^2 (a + b \arctan(c + dx))^3 dx = \text{Too large to display}$$

input `int((f*x+e)^2*(a+b*atan(d*x+c))^3,x)`

output

```
( - 4*atan(c + d*x)**3*b**3*c**3*f**2 + 6*atan(c + d*x)**3*b**3*c**2*d*e*f
- 4*atan(c + d*x)**3*b**3*c*f**2 + 6*atan(c + d*x)**3*b**3*d**3*e**2*x +
6*atan(c + d*x)**3*b**3*d**3*e*f*x**2 + 2*atan(c + d*x)**3*b**3*d**3*f**2*
x**3 + 6*atan(c + d*x)**3*b**3*d*e*f - 12*atan(c + d*x)**2*a*b**2*c**3*f**
2 + 18*atan(c + d*x)**2*a*b**2*c**2*d*e*f - 12*atan(c + d*x)**2*a*b**2*c*f
**2 + 18*atan(c + d*x)**2*a*b**2*d**3*e**2*x + 18*atan(c + d*x)**2*a*b**2*
d**3*e*f*x**2 + 6*atan(c + d*x)**2*a*b**2*d**3*f**2*x**3 + 18*atan(c + d*x
)**2*a*b**2*d*e*f - 3*atan(c + d*x)**2*b**3*c**2*f**2 + 12*atan(c + d*x)**
2*b**3*c*d*f**2*x - 18*atan(c + d*x)**2*b**3*d**2*e*f*x - 3*atan(c + d*x)*
*2*b**3*d**2*f**2*x**2 - 3*atan(c + d*x)**2*b**3*f**2 + 6*atan(c + d*x)*a*
*2*b*c**3*f**2 - 18*atan(c + d*x)*a**2*b*c**2*d*e*f + 18*atan(c + d*x)*a**
2*b*c*d**2*e**2 - 18*atan(c + d*x)*a**2*b*c*f**2 + 18*atan(c + d*x)*a**2*b
*d**3*e**2*x + 18*atan(c + d*x)*a**2*b*d**3*e*f*x**2 + 6*atan(c + d*x)*a**
2*b*d**3*f**2*x**3 + 18*atan(c + d*x)*a**2*b*d*e*f + 30*atan(c + d*x)*a*b*
*2*c**2*f**2 - 36*atan(c + d*x)*a*b**2*c*d*e*f + 24*atan(c + d*x)*a*b**2*c
*d*f**2*x - 36*atan(c + d*x)*a*b**2*d**2*e*f*x - 6*atan(c + d*x)*a*b**2*d*
*2*f**2*x**2 - 6*atan(c + d*x)*a*b**2*f**2 + 6*atan(c + d*x)*b**3*c*f**2 +
6*atan(c + d*x)*b**3*d*f**2*x - 36*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x
+ d**2*x**2 + 1),x)*a*b**2*c**2*d**2*f**2 + 72*int((atan(c + d*x)*x)/(c**2
+ 2*c*d*x + d**2*x**2 + 1),x)*a*b**2*c*d**3*e*f - 36*int((atan(c + d*x...
```

3.45 $\int (e + fx)(a + b \arctan(c + dx))^3 dx$

Optimal result	403
Mathematica [A] (verified)	404
Rubi [A] (verified)	405
Maple [C] (warning: unable to verify)	407
Fricas [F]	407
Sympy [F]	407
Maxima [F]	408
Giac [F]	409
Mupad [F(-1)]	409
Reduce [F]	409

Optimal result

Integrand size = 18, antiderivative size = 337

$$\begin{aligned}
 & \int (e + fx)(a + b \arctan(c + dx))^3 dx \\
 &= -\frac{3ibf(a + b \arctan(c + dx))^2}{2d^2} - \frac{3bf(c + dx)(a + b \arctan(c + dx))^2}{2d^2} \\
 &+ \frac{i(de - cf)(a + b \arctan(c + dx))^3}{d^2} \\
 &- \frac{(de + f - cf)(de - (1 + c)f)(a + b \arctan(c + dx))^3}{2d^2 f} \\
 &+ \frac{(e + fx)^2(a + b \arctan(c + dx))^3}{2f} - \frac{3b^2 f(a + b \arctan(c + dx)) \log\left(\frac{2}{1+i(c+dx)}\right)}{d^2} \\
 &+ \frac{3b(de - cf)(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1+i(c+dx)}\right)}{d^2} \\
 &- \frac{3ib^3 f \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{2d^2} \\
 &+ \frac{3ib^2(de - cf)(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d^2} \\
 &+ \frac{3b^3(de - cf) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2d^2}
 \end{aligned}$$

output

```
-3/2*I*b*f*(a+b*arctan(d*x+c))^2/d^2-3/2*b*f*(d*x+c)*(a+b*arctan(d*x+c))^2
/d^2+I*(-c*f+d*e)*(a+b*arctan(d*x+c))^3/d^2-1/2*(-c*f+d*e+f)*(d*e-(1+c)*f)
*(a+b*arctan(d*x+c))^3/d^2/f+1/2*(f*x+e)^2*(a+b*arctan(d*x+c))^3/f-3*b^2*f
*(a+b*arctan(d*x+c))*ln(2/(1+I*(d*x+c)))/d^2+3*b*(-c*f+d*e)*(a+b*arctan(d*
x+c))^2*ln(2/(1+I*(d*x+c)))/d^2-3/2*I*b^3*f*polylog(2,1-2/(1+I*(d*x+c)))/d
^2+3*I*b^2*(-c*f+d*e)*(a+b*arctan(d*x+c))*polylog(2,1-2/(1+I*(d*x+c)))/d^2
+3/2*b^3*(-c*f+d*e)*polylog(3,1-2/(1+I*(d*x+c)))/d^2
```

Mathematica [A] (verified)

Time = 4.56 (sec) , antiderivative size = 592, normalized size of antiderivative = 1.76

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx$$

$$= \frac{a^2(2ade - 3bf - 2acf)(c + dx) + a^3f(c + dx)^2 + 3a^2bf \arctan(c + dx) - 3a^2b(c + dx)(cf - d(2e + fx))}{d^2}$$

input

```
Integrate[(e + f*x)*(a + b*ArcTan[c + d*x])^3,x]
```

output

```
(a^2*(2*a*d*e - 3*b*f - 2*a*c*f)*(c + d*x) + a^3*f*(c + d*x)^2 + 3*a^2*b*f
*ArcTan[c + d*x] - 3*a^2*b*(c + d*x)*(c*f - d*(2*e + f*x))*ArcTan[c + d*x]
+ 6*a*b^2*f*(-((c + d*x)*ArcTan[c + d*x]) + ((1 + (c + d*x)^2)*ArcTan[c +
d*x]^2)/2 - Log[1/Sqrt[1 + (c + d*x)^2]]) - 3*a^2*b*(d*e - c*f)*Log[1 + (
c + d*x)^2] + 6*a*b^2*d*e*(ArcTan[c + d*x]*((-I + c + d*x)*ArcTan[c + d*x]
+ 2*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - I*PolyLog[2, -E^((2*I)*ArcTan[c
+ d*x])]) - 6*a*b^2*c*f*(ArcTan[c + d*x]*((-I + c + d*x)*ArcTan[c + d*x]
+ 2*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - I*PolyLog[2, -E^((2*I)*ArcTan[c
+ d*x])]) + b^3*f*(ArcTan[c + d*x]*((3*I)*ArcTan[c + d*x] - 3*(c + d*x)*A
rcTan[c + d*x] + (1 + (c + d*x)^2)*ArcTan[c + d*x]^2 - 6*Log[1 + E^((2*I)*A
rcTan[c + d*x])]) + (3*I)*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) + 2*b^3*
d*e*(ArcTan[c + d*x]^2*(-I + c + d*x)*ArcTan[c + d*x] + 3*Log[1 + E^((2*I
)*ArcTan[c + d*x])]) - (3*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*ArcTan[c
+ d*x])]) + (3*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])])/2 - 2*b^3*c*f*(Arc
Tan[c + d*x]^2*(-I + c + d*x)*ArcTan[c + d*x] + 3*Log[1 + E^((2*I)*ArcTan
[c + d*x])]) - (3*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*ArcTan[c + d*x]
)] + (3*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])])/2))/(2*d^2)
```

Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5570, 27, 5389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx$$

$$\downarrow 5570$$

$$\int \frac{\left(d\left(e - \frac{cf}{d}\right) + f(c + dx)\right)(a + b \arctan(c + dx))^3}{d} d(c + dx)$$

$$\downarrow 27$$

$$\int \frac{(de - cf + f(c + dx))(a + b \arctan(c + dx))^3}{d^2} d(c + dx)$$

$$\downarrow 5389$$

$$\frac{(f(c + dx) - cf + de)^2 (a + b \arctan(c + dx))^3}{2f} - \frac{3b \int \left(f^2 (a + b \arctan(c + dx))^2 + \frac{(de - cf + f)(de - (c + 1)f) + 2f(de - cf)(c + dx)(a + b \arctan(c + dx))^2}{(c + dx)^2 + 1} \right) d(c + dx)}{2f d^2}$$

$$\downarrow 2009$$

$$\frac{(f(c + dx) - cf + de)^2 (a + b \arctan(c + dx))^3}{2f} - \frac{3b \left(-2ibf(de - cf) \operatorname{PolyLog}\left(2, 1 - \frac{2}{i(c + dx) + 1}\right) (a + b \arctan(c + dx)) - \frac{2if(de - cf)(a + b \arctan(c + dx))^3}{3b} \right)}{2f d^2}$$

input

```
Int[(e + f*x)*(a + b*ArcTan[c + d*x])^3, x]
```

output

```

(((d*e - c*f + f*(c + d*x))^2*(a + b*ArcTan[c + d*x])^3)/(2*f) - (3*b*(I*f
^2*(a + b*ArcTan[c + d*x])^2 + f^2*(c + d*x)*(a + b*ArcTan[c + d*x])^2 - (
((2*I)/3)*f*(d*e - c*f)*(a + b*ArcTan[c + d*x])^3)/b + ((d*e + f - c*f)*(d
*e - (1 + c)*f)*(a + b*ArcTan[c + d*x])^3)/(3*b) + 2*b*f^2*(a + b*ArcTan[c
+ d*x])*Log[2/(1 + I*(c + d*x))] - 2*f*(d*e - c*f)*(a + b*ArcTan[c + d*x]
)^2*Log[2/(1 + I*(c + d*x))] + I*b^2*f^2*PolyLog[2, 1 - 2/(1 + I*(c + d*x)
)] - (2*I)*b*f*(d*e - c*f)*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 + I
*(c + d*x))] - b^2*f*(d*e - c*f)*PolyLog[3, 1 - 2/(1 + I*(c + d*x))]))/(2*f
)/d^2

```

Defintions of rubi rules used

rule 27

```

Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 5389

```

Int[((a_) + ArcTan[(c_)*(x_)]*(b_.))^p_)*((d_) + (e_.)*(x_))^(q_.), x_Sy
mbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - S
imp[b*c*(p/(e*(q + 1))) Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] &&
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]

```

rule 5570

```

Int[((a_) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^p_)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]

```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.86 (sec) , antiderivative size = 8296, normalized size of antiderivative = 24.62

method	result	size
parts	Expression too large to display	8296
derivativeldivides	Expression too large to display	8298
default	Expression too large to display	8298

input `int((f*x+e)*(a+b*arctan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F]

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx = \int (fx + e)(b \arctan(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c))^3,x, algorithm="fricas")`

output `integral(a^3*f*x + a^3*e + (b^3*f*x + b^3*e)*arctan(d*x + c)^3 + 3*(a*b^2*f*x + a*b^2*e)*arctan(d*x + c)^2 + 3*(a^2*b*f*x + a^2*b*e)*arctan(d*x + c), x)`

Sympy [F]

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx = \int (a + b \operatorname{atan}(c + dx))^3 (e + fx) dx$$

input `integrate((f*x+e)*(a+b*atan(d*x+c))**3,x)`

output `Integral((a + b*atan(c + d*x))**3*(e + f*x), x)`

Maxima [F]

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx = \int (fx + e)(b \arctan(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output

```

7/8*b^3*c^2*e*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 3*a*b^2*c^2*e*
arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - (3*arctan(d*x + c)*arctan((d
^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*a*b^2*c^2*e - 7/32*(6*ar
ctan(d*x + c)^2*arctan((d^2*x + c*d)/d)^2/d - 4*arctan(d*x + c)*arctan((d
^2*x + c*d)/d)^3/d + arctan((d^2*x + c*d)/d)^4/d)*b^3*c^2*e + 7/8*b^3*e*arc
tan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 56*b^3*d^2*f*integrate(1/64*x^3
*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b^3*d^2*f*integra
te(1/64*x^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 +
2*c*d*x + c^2 + 1), x) + 192*a*b^2*d^2*f*integrate(1/64*x^3*arctan(d*x + c
)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 56*b^3*d^2*e*integrate(1/64*x^2*ar
ctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 112*b^3*c*d*f*integrat
e(1/64*x^2*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^3*d
^2*f*integrate(1/64*x^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d
^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b^3*d^2*e*integrate(1/64*x^2*arctan(d*
x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x
) + 12*b^3*c*d*f*integrate(1/64*x^2*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x
+ c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 192*a*b^2*d^2*e*integrate
(1/64*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 384*a*b^2*
c*d*f*integrate(1/64*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1),
x) + 112*b^3*c*d*e*integrate(1/64*x*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*...

```

Giac [F]

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx = \int (fx + e)(b \arctan(dx + c) + a)^3 dx$$

input `integrate((f*x+e)*(a+b*arctan(d*x+c))^3,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx = \int (e + fx) (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((e + f*x)*(a + b*atan(c + d*x))^3,x)`

output `int((e + f*x)*(a + b*atan(c + d*x))^3, x)`

Reduce [F]

$$\int (e + fx)(a + b \arctan(c + dx))^3 dx$$

$$= \frac{\operatorname{atan}(dx + c)^3 b^3 d^2 f x^2 + 6 \operatorname{atan}(dx + c)^2 a b^2 d^2 e x + 3 \operatorname{atan}(dx + c)^2 a b^2 d^2 f x^2 + 6 \operatorname{atan}(dx + c) a^2 b c d e + \dots}{\dots}$$

input `int((f*x+e)*(a+b*atan(d*x+c))^3,x)`

output

```
(atan(c + d*x)**3*b**3*c**2*f + 2*atan(c + d*x)**3*b**3*d**2*e*x + atan(c
+ d*x)**3*b**3*d**2*f*x**2 + atan(c + d*x)**3*b**3*f + 3*atan(c + d*x)**2*
a*b**2*c**2*f + 6*atan(c + d*x)**2*a*b**2*d**2*e*x + 3*atan(c + d*x)**2*a*
b**2*d**2*f*x**2 + 3*atan(c + d*x)**2*a*b**2*f - 3*atan(c + d*x)**2*b**3*d
*f*x - 3*atan(c + d*x)*a**2*b*c**2*f + 6*atan(c + d*x)*a**2*b*c*d*e + 6*at
an(c + d*x)*a**2*b*d**2*e*x + 3*atan(c + d*x)*a**2*b*d**2*f*x**2 + 3*atan(
c + d*x)*a**2*b*f - 6*atan(c + d*x)*a*b**2*c*f - 6*atan(c + d*x)*a*b**2*d*
f*x + 12*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*a*b**2*
c*d**2*f - 12*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*a*
b**2*d**3*e + 6*int((atan(c + d*x)*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),x)*
b**3*d**2*f + 6*int((atan(c + d*x)**2*x)/(c**2 + 2*c*d*x + d**2*x**2 + 1),
x)*b**3*c*d**2*f - 6*int((atan(c + d*x)**2*x)/(c**2 + 2*c*d*x + d**2*x**2
+ 1),x)*b**3*d**3*e + 3*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a**2*b*c*f - 3
*log(c**2 + 2*c*d*x + d**2*x**2 + 1)*a**2*b*d*e + 3*log(c**2 + 2*c*d*x + d
**2*x**2 + 1)*a*b**2*f + 2*a**3*d**2*e*x + a**3*d**2*f*x**2 - 3*a**2*b*d*f
*x)/(2*d**2)
```

3.46 $\int (a + b \arctan(c + dx))^3 dx$

Optimal result	411
Mathematica [A] (verified)	412
Rubi [A] (verified)	412
Maple [B] (verified)	415
Fricas [F]	415
Sympy [F]	416
Maxima [F]	416
Giac [F]	417
Mupad [F(-1)]	417
Reduce [F]	417

Optimal result

Integrand size = 12, antiderivative size = 143

$$\begin{aligned}
 \int (a + b \arctan(c + dx))^3 dx = & \frac{i(a + b \arctan(c + dx))^3}{d} \\
 & + \frac{(c + dx)(a + b \arctan(c + dx))^3}{d} \\
 & + \frac{3b(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1+i(c+dx)}\right)}{d} \\
 & + \frac{3ib^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1+i(c+dx)}\right)}{d} \\
 & + \frac{3b^3 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1+i(c+dx)}\right)}{2d}
 \end{aligned}$$

output

```

I*(a+b*arctan(d*x+c))^3/d+(d*x+c)*(a+b*arctan(d*x+c))^3/d+3*b*(a+b*arctan(
d*x+c))^2*ln(2/(1+I*(d*x+c)))/d+3*I*b^2*(a+b*arctan(d*x+c))*polylog(2,1-2/
(1+I*(d*x+c)))/d+3/2*b^3*polylog(3,1-2/(1+I*(d*x+c)))/d

```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.48

$$\int (a + b \arctan(c + dx))^3 dx$$

$$= \frac{2a^3(c + dx) + 6a^2b(c + dx) \arctan(c + dx) - 3a^2b \log(1 + (c + dx)^2) + 6ab^2(\arctan(c + dx) ((-i + c + dx) \arctan(c + dx) + \log[1 + E^{((2I) \arctan(c + dx))}])) - I \text{PolyLog}[2, -E^{((2I) \arctan(c + dx))}] + 2b^3(\arctan(c + dx)^2((-i + c + dx) \arctan(c + dx) + 3 \log[1 + E^{((2I) \arctan(c + dx))}])) - (3I) \arctan(c + dx) \text{PolyLog}[2, -E^{((2I) \arctan(c + dx))}] + (3 \text{PolyLog}[3, -E^{((2I) \arctan(c + dx))}])/2)}{2d}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3,x]
```

output

```
(2*a^3*(c + d*x) + 6*a^2*b*(c + d*x)*ArcTan[c + d*x] - 3*a^2*b*Log[1 + (c + d*x)^2] + 6*a*b^2*(ArcTan[c + d*x]*((-I + c + d*x)*ArcTan[c + d*x] + 2*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - I*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) + 2*b^3*(ArcTan[c + d*x]^2*((-I + c + d*x)*ArcTan[c + d*x] + 3*Log[1 + E^((2*I)*ArcTan[c + d*x])]) - (3*I)*ArcTan[c + d*x]*PolyLog[2, -E^((2*I)*ArcTan[c + d*x])]) + (3*PolyLog[3, -E^((2*I)*ArcTan[c + d*x])])/2))/(2*d)
```

Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5562, 5345, 5455, 5379, 5529, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \arctan(c + dx))^3 dx$$

$$\downarrow \text{5562}$$

$$\frac{\int (a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow \text{5345}$$

$$\frac{(c + dx)(a + b \arctan(c + dx))^3 - 3b \int \frac{(c + dx)(a + b \arctan(c + dx))^2}{(c + dx)^2 + 1} d(c + dx)}{d}$$

$$\downarrow \text{5455}$$

$$\frac{(c + dx)(a + b \arctan(c + dx))^3 - 3b \left(- \int \frac{(a + b \arctan(c + dx))^2}{-c - dx + i} d(c + dx) - \frac{i(a + b \arctan(c + dx))^3}{3b} \right)}{d}$$

↓ 5379

$$\frac{(c + dx)(a + b \arctan(c + dx))^3 - 3b \left(2b \int \frac{(a + b \arctan(c + dx)) \log\left(\frac{2}{i(c + dx) + 1}\right)}{(c + dx)^2 + 1} d(c + dx) - \frac{i(a + b \arctan(c + dx))^3}{3b} - \log\left(\frac{1}{1 + i(c + dx)}\right) \right)}{d}$$

↓ 5529

$$\frac{(c + dx)(a + b \arctan(c + dx))^3 - 3b \left(2b \left(\frac{1}{2} i b \int \frac{\text{PolyLog}\left(2, 1 - \frac{2}{i(c + dx) + 1}\right)}{(c + dx)^2 + 1} d(c + dx) - \frac{1}{2} i \text{PolyLog}\left(2, 1 - \frac{2}{i(c + dx) + 1}\right) \right) \right)}{d}$$

↓ 7164

$$\frac{(c + dx)(a + b \arctan(c + dx))^3 - 3b \left(-\frac{1}{2} i \text{PolyLog}\left(2, 1 - \frac{2}{i(c + dx) + 1}\right) (a + b \arctan(c + dx)) - \frac{1}{4} b \text{PolyLog}\left(3, 1 - \frac{2}{i(c + dx) + 1}\right) \right)}{d}$$

input `Int[(a + b*ArcTan[c + d*x])^3,x]`

output `((c + d*x)*(a + b*ArcTan[c + d*x])^3 - 3*b*(((-1/3*I)*(a + b*ArcTan[c + d*x])^3)/b - (a + b*ArcTan[c + d*x])^2*Log[2/(1 + I*(c + d*x))] + 2*b*(((-1/2*I)*(a + b*ArcTan[c + d*x])*PolyLog[2, 1 - 2/(1 + I*(c + d*x))] - (b*PolyLog[3, 1 - 2/(1 + I*(c + d*x))])/4)))/d`

Definitions of rubi rules used

rule 5345 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 5379 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Simp[b*c*(p/e) Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]`

rule 5455 `Int[(((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Simp[1/(c*d) Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]`

rule 5529 `Int[(Log[u_]*)((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Simp[b*p*(I/2) Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]`

rule 5562 `Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d Subst[Int[(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]`

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 278 vs. $2(136) = 272$.

Time = 0.82 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.95

method	result
derivativedivides	$(dx+c)a^3+b^3 \left(\arctan(dx+c)^3(dx+c+i)-2i \arctan(dx+c)^3+3 \arctan(dx+c)^2 \ln \left(1+\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) -3i \arctan(dx+c) \right)$
default	$(dx+c)a^3+b^3 \left(\arctan(dx+c)^3(dx+c+i)-2i \arctan(dx+c)^3+3 \arctan(dx+c)^2 \ln \left(1+\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) -3i \arctan(dx+c) \right)$
parts	$a^3x + \frac{b^3 \left(\arctan(dx+c)^3(dx+c+i)-2i \arctan(dx+c)^3+3 \arctan(dx+c)^2 \ln \left(1+\frac{(1+i(dx+c))^2}{1+(dx+c)^2} \right) -3i \arctan(dx+c) \right)}{d}$

input `int((a+b*arctan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*((d*x+c)*a^3+b^3*(arctan(d*x+c)^3*(d*x+c+I)-2*I*arctan(d*x+c)^3+3*arctan(d*x+c)^2*ln(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))-3*I*arctan(d*x+c)*polylog(2,-(1+I*(d*x+c))^2/(1+(d*x+c)^2))+3/2*polylog(3,-(1+I*(d*x+c))^2/(1+(d*x+c)^2)))+3*a*b^2*(arctan(d*x+c)^2*(d*x+c+I)+2*arctan(d*x+c)*ln(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2))-2*I*arctan(d*x+c)^2-I*polylog(2,-(1+I*(d*x+c))^2/(1+(d*x+c)^2)))+3*a^2*b*((d*x+c)*arctan(d*x+c)-1/2*ln(1+(d*x+c)^2)))`

Fricas [F]

$$\int (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 dx$$

input `integrate((a+b*arctan(d*x+c))^3,x, algorithm="fricas")`

output `integral(b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3, x)`

Sympy [F]

$$\int (a + b \arctan(c + dx))^3 dx = \int (a + b \operatorname{atan}(c + dx))^3 dx$$

input `integrate((a+b*atan(d*x+c))**3,x)`

output `Integral((a + b*atan(c + d*x))**3, x)`

Maxima [F]

$$\int (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 dx$$

input `integrate((a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output `7/8*b^3*c^2*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 1/8*b^3*x*arctan(d*x + c)^3 + 3*a*b^2*c^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 3/32*b^3*x*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 - (3*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^2/d - arctan((d^2*x + c*d)/d)^3/d)*a*b^2*c^2 - 7/32*(6*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)^2/d - 4*arctan(d*x + c)*arctan((d^2*x + c*d)/d)^3/d + arctan((d^2*x + c*d)/d)^4/d)*b^3*c^2 + 7/8*b^3*arctan(d*x + c)^3*arctan((d^2*x + c*d)/d)/d + 28*b^3*d^2*integrate(1/32*x^2*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^3*d^2*integrate(1/32*x^2*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 96*a*b^2*d^2*integrate(1/32*x^2*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 56*b^3*c*d*integrate(1/32*x*arctan(d*x + c)^3/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^3*d^2*integrate(1/32*x^2*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 6*b^3*c*d*integrate(1/32*x*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 192*a*b^2*c*d*integrate(1/32*x*arctan(d*x + c)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 12*b^3*c*d*integrate(1/32*x*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*b^3*c^2*integrate(1/32*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2/(d^2*x^2 + 2*c*d*x + c^2 + 1), x) + 3*a*b^2*arctan(d*x + c)^2*arctan((d^2*x + c*d)/d)/d - 12*b^3*d*integrate(...`

Giac [F]

$$\int (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 dx$$

input `integrate((a+b*arctan(d*x+c))^3,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (a + b \arctan(c + dx))^3 dx = \int (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((a + b*atan(c + d*x))^3,x)`

output `int((a + b*atan(c + d*x))^3, x)`

Reduce [F]

$$\int (a + b \arctan(c + dx))^3 dx$$

$$= \frac{2 \operatorname{atan}(dx + c)^3 b^3 dx + 6 \operatorname{atan}(dx + c)^2 a b^2 dx + 6 \operatorname{atan}(dx + c) a^2 b c + 6 \operatorname{atan}(dx + c) a^2 b dx - 12 \left(\int \frac{\operatorname{atan}}{d^2 x^2 +} \right)}{2d}$$

input `int((a+b*atan(d*x+c))^3,x)`

output

```
(2*atan(c + d*x)**3*b**3*d*x + 6*atan(c + d*x)**2*a*b**2*d*x + 6*atan(c +
d*x)*a**2*b*c + 6*atan(c + d*x)*a**2*b*d*x - 12*int((atan(c + d*x)*x)/(c**
2 + 2*c*d*x + d**2*x**2 + 1),x)*a*b**2*d**2 - 6*int((atan(c + d*x)**2*x)/(
c**2 + 2*c*d*x + d**2*x**2 + 1),x)*b**3*d**2 - 3*log(c**2 + 2*c*d*x + d**2
*x**2 + 1)*a**2*b + 2*a**3*d*x)/(2*d)
```

$$3.47 \quad \int \frac{(a+b \arctan(c+dx))^3}{e+fx} dx$$

Optimal result	419
Mathematica [F]	420
Rubi [A] (verified)	420
Maple [C] (warning: unable to verify)	422
Fricas [F]	423
Sympy [F(-1)]	424
Maxima [F]	424
Giac [F]	424
Mupad [F(-1)]	425
Reduce [F]	425

Optimal result

Integrand size = 20, antiderivative size = 372

$$\begin{aligned} & \int \frac{(a+b \arctan(c+dx))^3}{e+fx} dx \\ &= -\frac{(a+b \arctan(c+dx))^3 \log\left(\frac{2}{1-i(c+dx)}\right)}{f} \\ & \quad + \frac{(a+b \arctan(c+dx))^3 \log\left(\frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{f} \\ & \quad + \frac{3ib(a+b \arctan(c+dx))^2 \text{PolyLog}\left(2, 1 - \frac{2}{1-i(c+dx)}\right)}{2f} \\ & \quad - \frac{3ib(a+b \arctan(c+dx))^2 \text{PolyLog}\left(2, 1 - \frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{2f} \\ & \quad - \frac{3b^2(a+b \arctan(c+dx)) \text{PolyLog}\left(3, 1 - \frac{2}{1-i(c+dx)}\right)}{2f} \\ & \quad + \frac{3b^2(a+b \arctan(c+dx)) \text{PolyLog}\left(3, 1 - \frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{2f} \\ & \quad - \frac{3ib^3 \text{PolyLog}\left(4, 1 - \frac{2}{1-i(c+dx)}\right)}{4f} + \frac{3ib^3 \text{PolyLog}\left(4, 1 - \frac{2d(e+fx)}{(de+(i-c)f)(1-i(c+dx))}\right)}{4f} \end{aligned}$$

output

```

-(a+b*arctan(d*x+c))^3*ln(2/(1-I*(d*x+c)))/f+(a+b*arctan(d*x+c))^3*ln(2*d*
(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f+3/2*I*b*(a+b*arctan(d*x+c))^2*polylog
og(2,1-2/(1-I*(d*x+c)))/f-3/2*I*b*(a+b*arctan(d*x+c))^2*polylog(2,1-2*d*(f
*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f-3/2*b^2*(a+b*arctan(d*x+c))*polylog(3
,1-2/(1-I*(d*x+c)))/f+3/2*b^2*(a+b*arctan(d*x+c))*polylog(3,1-2*d*(f*x+e)/
(d*e+(I-c)*f)/(1-I*(d*x+c)))/f-3/4*I*b^3*polylog(4,1-2/(1-I*(d*x+c)))/f+3/
4*I*b^3*polylog(4,1-2*d*(f*x+e)/(d*e+(I-c)*f)/(1-I*(d*x+c)))/f

```

Mathematica [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3/(e + f*x), x]
```

output

```
Integrate[(a + b*ArcTan[c + d*x])^3/(e + f*x), x]
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {5570, 27, 5385}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx \\
\downarrow 5570 \\
\int \frac{d(a + b \arctan(c + dx))^3}{d(e - \frac{cf}{d}) + f(c + dx)} d(c + dx) \\
\hline
d \\
\downarrow 27
\end{array}$$

$$\begin{aligned}
& \int \frac{(a + b \arctan(c + dx))^3}{f(c + dx) - cf + de} d(c + dx) \\
& \quad \downarrow \text{5385} \\
& \frac{3b^2(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(3, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + if)(1 - i(c + dx))}\right)}{2f} - \\
& \frac{3b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - i(c + dx)}\right) (a + b \arctan(c + dx))}{2f} - \\
& \frac{3ib(a + b \arctan(c + dx))^2 \operatorname{PolyLog}\left(2, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + if)(1 - i(c + dx))}\right)}{2f} + \\
& \frac{(a + b \arctan(c + dx))^3 \log\left(\frac{2(f(c + dx) - cf + de)}{(1 - i(c + dx))(-cf + de + if)}\right)}{f} + \\
& \frac{3ib \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i(c + dx)}\right) (a + b \arctan(c + dx))^2}{2f} - \\
& \frac{\log\left(\frac{2}{1 - i(c + dx)}\right) (a + b \arctan(c + dx))^3}{f} + \frac{3ib^3 \operatorname{PolyLog}\left(4, 1 - \frac{2(de - cf + f(c + dx))}{(de - cf + if)(1 - i(c + dx))}\right)}{4f} - \\
& \frac{3ib^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - i(c + dx)}\right)}{4f}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^3/(e + f*x),x]`

output

```

-(((a + b*ArcTan[c + d*x])^3*Log[2/(1 - I*(c + d*x))])/f) + ((a + b*ArcTan
[c + d*x])^3*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(
c + d*x))))/f + (((3*I)/2)*b*(a + b*ArcTan[c + d*x])^2*PolyLog[2, 1 - 2/(
1 - I*(c + d*x))])/f - (((3*I)/2)*b*(a + b*ArcTan[c + d*x])^2*PolyLog[2, 1
- (2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))])/f
- (3*b^2*(a + b*ArcTan[c + d*x])*PolyLog[3, 1 - 2/(1 - I*(c + d*x))])/((2*f)
+ (3*b^2*(a + b*ArcTan[c + d*x])*PolyLog[3, 1 - (2*(d*e - c*f + f*(c +
d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x)))]/(2*f) - (((3*I)/4)*b^3*Poly
Log[4, 1 - 2/(1 - I*(c + d*x))])/f + (((3*I)/4)*b^3*PolyLog[4, 1 - (2*(d*e
- c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x)))]/f

```

Definitions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 5385

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^3/((d_.) + (e_.)*(x_)), x_Symbol] :>
Simp[(-(a + b*ArcTan[c*x])^3)*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*ArcTan[c*x])^3*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] + Simp[3*I*b*(a + b*ArcTan[c*x])^2*(PolyLog[2, 1 - 2/(1 - I*c*x)]/(2*e)), x] - Simp[3*I*b*(a + b*ArcTan[c*x])^2*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] - Simp[3*b^2*(a + b*ArcTan[c*x])*(PolyLog[3, 1 - 2/(1 - I*c*x)]/(2*e)), x] + Simp[3*b^2*(a + b*ArcTan[c*x])*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] - Simp[3*I*b^3*(PolyLog[4, 1 - 2/(1 - I*c*x)]/(4*e)), x] + Simp[3*I*b^3*(PolyLog[4, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

rule 5570

```
Int[((a_.) + ArcTan[(c_.) + (d_.)*(x_)]*(b_.))^p*((e_.) + (f_.)*(x_))^m, x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && IGtQ[p, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.72 (sec) , antiderivative size = 3817, normalized size of antiderivative = 10.26

method	result	size
derivativedivides	Expression too large to display	3817
default	Expression too large to display	3817
parts	Expression too large to display	4072

input

```
int((a+b*arctan(d*x+c))^3/(f*x+e),x,method=_RETURNVERBOSE)
```

output

```

1/d*(a^3*d*ln(c*f-d*e-f*(d*x+c))/f-b^3*d*(-ln(c*f-d*e-f*(d*x+c))/f*arctan(
d*x+c)^3+3/f*(1/3*arctan(d*x+c)^3*ln(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f
*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d
*e)-1/3*I*f/(c*f-d*e+I*f)*arctan(d*x+c)^3*ln(1-(c*f-d*e+I*f)*(1+I*(d*x+c))
^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))-1/6*I*Pi*csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d
*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^
2)-I*f+c*f-d*e)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))*(csgn(I*(I*f*(1+I*(d*x+
c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/
(1+(d*x+c)^2)-I*f+c*f-d*e))*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-csgn
(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*
e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)
^2)))*csgn(I/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))-csgn(I*(I*f*(1+I*(d*x+c))^
2/(1+(d*x+c)^2)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(
d*x+c)^2)-I*f+c*f-d*e))*csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I
*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)/(
1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))+csgn(I*(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2
)+c*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+
c*f-d*e)/(1+(1+I*(d*x+c))^2/(1+(d*x+c)^2)))^2)*arctan(d*x+c)^3+1/2*arctan(
d*x+c)*polylog(3,-(1+I*(d*x+c))^2/(1+(d*x+c)^2))+I*d*e*polylog(4,(c*f-d*e+
I*f)*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))/(4*I*f+4*c*f-4*d*e)+1...

```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^3}{fx + e} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(f*x+e),x, algorithm="fricas")
```

output

```
integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arct
an(d*x + c) + a^3)/(f*x + e), x)
```


Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))**3/(f*x+e), x)`output `Timed out`**Maxima [F]**

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^3}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(f*x+e), x, algorithm="maxima")`output `a^3*log(f*x + e)/f + integrate(1/32*(28*b^3*arctan(d*x + c)^3 + 3*b^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 + 96*a*b^2*arctan(d*x + c)^2 + 96*a^2*b*arctan(d*x + c))/(f*x + e), x)`**Giac [F]**

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \int \frac{(b \arctan(dx + c) + a)^3}{fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(f*x+e), x, algorithm="giac")`output `integrate((b*arctan(d*x + c) + a)^3/(f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{e + fx} dx$$

input `int((a + b*atan(c + d*x))^3/(e + f*x),x)`output `int((a + b*atan(c + d*x))^3/(e + f*x), x)`**Reduce [F]**

$$\int \frac{(a + b \arctan(c + dx))^3}{e + fx} dx$$

$$= \frac{3 \left(\int \frac{\operatorname{atan}(dx+c)}{fx+e} dx \right) a^2 b f + \left(\int \frac{\operatorname{atan}(dx+c)^3}{fx+e} dx \right) b^3 f + 3 \left(\int \frac{\operatorname{atan}(dx+c)^2}{fx+e} dx \right) a b^2 f + \log(fx + e) a^3}{f}$$

input `int((a+b*atan(d*x+c))^3/(f*x+e),x)`output `(3*int(atan(c + d*x)/(e + f*x),x)*a**2*b*f + int(atan(c + d*x)**3/(e + f*x),x)*b**3*f + 3*int(atan(c + d*x)**2/(e + f*x),x)*a*b**2*f + log(e + f*x)*a**3)/f`

$$3.48 \quad \int \frac{(a+b \arctan(c+dx))^3}{(e+fx)^2} dx$$

Optimal result	427
Mathematica [F]	428
Rubi [A] (verified)	428
Maple [C] (warning: unable to verify)	431
Fricas [F]	432
Sympy [F(-1)]	433
Maxima [F]	433
Giac [F]	434
Mupad [F(-1)]	434
Reduce [F]	434

Optimal result

Integrand size = 20, antiderivative size = 693

$$\begin{aligned}
& \int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx \\
&= \frac{id(a + b \arctan(c + dx))^3}{d^2e^2 - 2cdef + (1 + c^2)f^2} + \frac{d(de - cf)(a + b \arctan(c + dx))^3}{f(d^2e^2 - 2cdef + (1 + c^2)f^2)} \\
&\quad - \frac{(a + b \arctan(c + dx))^3}{f(e + fx)} - \frac{3bd(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1 - i(c + dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad + \frac{3bd(a + b \arctan(c + dx))^2 \log\left(\frac{2d(e + fx)}{(de + if - cf)(1 - i(c + dx))}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad + \frac{3bd(a + b \arctan(c + dx))^2 \log\left(\frac{2}{1 + i(c + dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad + \frac{3ib^2d(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - i(c + dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad - \frac{3ib^2d(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2d(e + fx)}{(de + if - cf)(1 - i(c + dx))}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad + \frac{3ib^2d(a + b \arctan(c + dx)) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + i(c + dx)}\right)}{d^2e^2 - 2cdef + (1 + c^2)f^2} \\
&\quad - \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - i(c + dx)}\right)}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)} + \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2d(e + fx)}{(de + if - cf)(1 - i(c + dx))}\right)}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)} \\
&\quad + \frac{3b^3d \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 + i(c + dx)}\right)}{2(d^2e^2 - 2cdef + (1 + c^2)f^2)}
\end{aligned}$$

output

```
I*d*(a+b*arctan(d*x+c))^3/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+d*(-c*f+d*e)*(a+
b*arctan(d*x+c))^3/f/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-(a+b*arctan(d*x+c))^3
/f/(f*x+e)-3*b*d*(a+b*arctan(d*x+c))^2*ln(2/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*
e*f+(c^2+1)*f^2)+3*b*d*(a+b*arctan(d*x+c))^2*ln(2*d*(f*x+e)/(d*e+I*f-c*f)/
(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+3*b*d*(a+b*arctan(d*x+c))^2
*ln(2/(1+I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+3*I*b^2*d*(a+b*arctan
(d*x+c))*polylog(2,1-2/(1-I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-3*I*
b^2*d*(a+b*arctan(d*x+c))*polylog(2,1-2*d*(f*x+e)/(d*e+I*f-c*f)/(1-I*(d*x+
c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)+3*I*b^2*d*(a+b*arctan(d*x+c))*polylog
(2,1-2/(1+I*(d*x+c)))/(d^2*e^2-2*c*d*e*f+(c^2+1)*f^2)-3*b^3*d*polylog(3,1-
2/(1-I*(d*x+c)))/(2*d^2*e^2-4*c*d*e*f+2*(c^2+1)*f^2)+3*b^3*d*polylog(3,1-2
*d*(f*x+e)/(d*e+I*f-c*f)/(1-I*(d*x+c)))/(2*d^2*e^2-4*c*d*e*f+2*(c^2+1)*f^2
)+3*b^3*d*polylog(3,1-2/(1+I*(d*x+c)))/(2*d^2*e^2-4*c*d*e*f+2*(c^2+1)*f^2)
```

Mathematica [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx$$

input

```
Integrate[(a + b*ArcTan[c + d*x])^3/(e + f*x)^2,x]
```

output

```
Integrate[(a + b*ArcTan[c + d*x])^3/(e + f*x)^2, x]
```

Rubi [A] (verified)

Time = 2.35 (sec) , antiderivative size = 1265, normalized size of antiderivative = 1.83, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5568, 7292, 5580, 27, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx$$

$$\begin{aligned}
& \downarrow 5568 \\
& \frac{3bd \int \frac{(a+b \arctan(c+dx))^2}{(e+fx)((c+dx)^2+1)} dx}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)} \\
& \downarrow 7292 \\
& \frac{3bd \int \frac{(a+b \arctan(c+dx))^2}{(e+fx)(c^2+2dxc+d^2x^2+1)} dx}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)} \\
& \downarrow 5580 \\
& \frac{3b \int \frac{d(a+b \arctan(c+dx))^2}{\left(d\left(e-\frac{cf}{d}\right)+f(c+dx)\right)((c+dx)^2+1)} d(c+dx)}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)} \\
& \downarrow 27 \\
& \frac{3bd \int \frac{(a+b \arctan(c+dx))^2}{(de-cf+f(c+dx))((c+dx)^2+1)} d(c+dx)}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)} \\
& \downarrow 7276 \\
& \frac{3bd \int \left(\frac{a^2}{(de-cf+f(c+dx))((c+dx)^2+1)} + \frac{2b \arctan(c+dx)a}{(de-cf+f(c+dx))((c+dx)^2+1)} + \frac{b^2 \arctan(c+dx)^2}{(de-cf+f(c+dx))((c+dx)^2+1)} \right) d(c+dx)}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)} \\
& \downarrow 2009 \\
& \frac{3bd \left(\frac{ib^2 f \arctan(c+dx)^3}{3(d^2e^2-2cdf e+(c^2+1)f^2)} + \frac{b^2(de-cf) \arctan(c+dx)^3}{3(d^2e^2-2cdf e+(c^2+1)f^2)} - \frac{b^2 f \log\left(\frac{2}{1-i(c+dx)}\right) \arctan(c+dx)^2}{d^2e^2-2cdf e+(c^2+1)f^2} + \frac{b^2 f \log\left(\frac{2}{i(c+dx)+1}\right) \arctan(c+dx)^2}{d^2e^2-2cdf e+(c^2+1)f^2} \right)}{f} - \frac{(a+b \arctan(c+dx))^3}{f(e+fx)}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])^3/(e + f*x)^2,x]`

output

```

-((a + b*ArcTan[c + d*x])^3/(f*(e + f*x))) + (3*b*d*((a^2*(d*e - c*f)*ArcTan[c + d*x])/(f^2 + (d*e - c*f)^2) + (I*a*b*f*ArcTan[c + d*x]^2)/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (a*b*(d*e - c*f)*ArcTan[c + d*x]^2)/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + ((I/3)*b^2*f*ArcTan[c + d*x]^3)/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b^2*(d*e - c*f)*ArcTan[c + d*x]^3)/(3*(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2)) - (2*a*b*f*ArcTan[c + d*x]*Log[2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) - (b^2*f*ArcTan[c + d*x]^2*Log[2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (2*a*b*f*ArcTan[c + d*x]*Log[2/(1 + I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b^2*f*ArcTan[c + d*x]^2*Log[2/(1 + I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (a^2*f*Log[d*e - c*f + f*(c + d*x)])/(f^2 + (d*e - c*f)^2) + (2*a*b*f*ArcTan[c + d*x]*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (b^2*f*ArcTan[c + d*x]^2*Log[(2*(d*e - c*f + f*(c + d*x))]/((d*e + I*f - c*f)*(1 - I*(c + d*x))))/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) - (a^2*f*Log[1 + (c + d*x)^2])/(2*(f^2 + (d*e - c*f)^2)) + (I*a*b*f*PolyLog[2, 1 - 2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (I*b^2*f*ArcTan[c + d*x]*PolyLog[2, 1 - 2/(1 - I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (I*a*b*f*PolyLog[2, 1 - 2/(1 + I*(c + d*x))])/(d^2*e^2 - 2*c*d*e*f + (1 + c^2)*f^2) + (I*b^2*f*ArcTan[c + d*x]*PolyLog[2, ...

```

Defintions of rubi rules used

rule 27

```

Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]

```

rule 2009

```

Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 5568

```

Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^p_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m + 1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[m, -1]

```

rule 5580

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/d Subs
t[Int[((d*e - c*f)/d + f*(x/d))^m*(C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTan[x])
^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x] &
& EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

rule 7292

```
Int[u_, x_Symbol] := With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v !=
= u]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.97 (sec) , antiderivative size = 3708, normalized size of antiderivative = 5.35

method	result	size
derivativedivides	Expression too large to display	3708
default	Expression too large to display	3708
parts	Expression too large to display	3834

input

```
int((a+b*arctan(d*x+c))^3/(f*x+e)^2,x,method=_RETURNVERBOSE)
```


output

```

1/d*(a^3*d^2/(c*f-d*e-f*(d*x+c))/f+b^3*d^2*(1/(c*f-d*e-f*(d*x+c))/f*arctan
(d*x+c)^3-3/f*(1/2*arctan(d*x+c)^2/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f*ln(1+
(d*x+c)^2)+arctan(d*x+c)^3/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*c*f-arctan(d*x+
c)^3/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*d*e-arctan(d*x+c)^2*f/(c^2*f^2-2*c*d*
e*f+d^2*e^2+f^2)*ln(c*f-d*e-f*(d*x+c))-f/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*a
rctan(d*x+c)^2*ln((1+I*(d*x+c))/(1+(d*x+c)^2)^(1/2))+f/(c^2*f^2-2*c*d*e*f+
d^2*e^2+f^2)*arctan(d*x+c)^2*ln(I*f*(1+I*(d*x+c))^2/(1+(d*x+c)^2)+c*f*(1+I
*(d*x+c))^2/(1+(d*x+c)^2)-d*e*(1+I*(d*x+c))^2/(1+(d*x+c)^2)-I*f+c*f-d*e)-1
/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f^2/(c*f-d*e+I*f)*arctan(d*x+c)*polylog(2
,(c*f-d*e+I*f)*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))-1/2*I/(c^2*f^2
-2*c*d*e*f+d^2*e^2+f^2)*f^2/(c*f-d*e+I*f)*polylog(3,(c*f-d*e+I*f)*(1+I*(d*
x+c))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))+I/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f^2
*c/(c*f-d*e+I*f)*arctan(d*x+c)*polylog(2,(c*f-d*e+I*f)*(1+I*(d*x+c))^2/(1+
(d*x+c)^2)/(d*e+I*f-c*f))+1/3*I*f/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*arctan(d
*x+c)^3-1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f^2*c/(c*f-d*e+I*f)*arctan(d*x+c
)^2*ln(1-(c*f-d*e+I*f)*(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))-1/2/(c
^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f^2*c/(c*f-d*e+I*f)*polylog(3,(c*f-d*e+I*f)*
(1+I*(d*x+c))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))-I/(c^2*f^2-2*c*d*e*f+d^2*e^2+
f^2)*f*d*e/(c*f-d*e+I*f)*arctan(d*x+c)*polylog(2,(c*f-d*e+I*f)*(1+I*(d*x+c
))^2/(1+(d*x+c)^2)/(d*e+I*f-c*f))+1/(c^2*f^2-2*c*d*e*f+d^2*e^2+f^2)*f*d...

```

Fricas [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(fx + e)^2} dx$$

input

```
integrate((a+b*arctan(d*x+c))^3/(f*x+e)^2,x, algorithm="fricas")
```

output

```
integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arct
an(d*x + c) + a^3)/(f^2*x^2 + 2*e*f*x + e^2), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))**3/(f*x+e)**2,x)`

output Timed out

Maxima [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(fx + e)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(f*x+e)^2,x, algorithm="maxima")`

output `3/2*(d*(2*(d^2*e - c*d*f)*arctan((d^2*x + c*d)/d)/((d^2*e^2*f - 2*c*d*e*f^2 + (c^2 + 1)*f^3)*d) - log(d^2*x^2 + 2*c*d*x + c^2 + 1)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2) + 2*log(f*x + e)/(d^2*e^2 - 2*c*d*e*f + (c^2 + 1)*f^2)) - 2*arctan(d*x + c)/(f^2*x + e*f))*a^2*b - a^3/(f^2*x + e*f) - 1/32*(4*b^3*arctan(d*x + c)^3 - 3*b^3*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 - 32*(f^2*x + e*f)*integrate(1/32*(28*(b^3*d^2*f*x^2 + 2*b^3*c*d*f*x + (b^3*c^2 + b^3)*f)*arctan(d*x + c)^3 + 12*(8*a*b^2*d^2*f*x^2 + b^3*d*e + (16*a*b^2*c + b^3)*d*f*x + 8*(a*b^2*c^2 + a*b^2)*f)*arctan(d*x + c)^2 - 12*(b^3*d^2*f*x^2 + b^3*c*d*e + (b^3*d^2*e + b^3*c*d*f)*x)*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 3*(b^3*d*f*x + b^3*d*e - (b^3*d^2*f*x^2 + 2*b^3*c*d*f*x + (b^3*c^2 + b^3)*f)*arctan(d*x + c))*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2)/(d^2*f^3*x^4 + (c^2 + 1)*e^2*f + 2*(d^2*e*f^2 + c*d*f^3)*x^3 + (d^2*e^2*f + 4*c*d*e*f^2 + (c^2 + 1)*f^3)*x^2 + 2*(c*d*e^2*f + (c^2 + 1)*e*f^2)*x), x))/(f^2*x + e*f)`

Giac [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \int \frac{(b \arctan(dx + c) + a)^3}{(fx + e)^2} dx$$

input `integrate((a+b*arctan(d*x+c))^3/(f*x+e)^2,x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)^3/(f*x + e)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \int \frac{(a + b \operatorname{atan}(c + dx))^3}{(e + fx)^2} dx$$

input `int((a + b*atan(c + d*x))^3/(e + f*x)^2,x)`

output `int((a + b*atan(c + d*x))^3/(e + f*x)^2, x)`

Reduce [F]

$$\int \frac{(a + b \arctan(c + dx))^3}{(e + fx)^2} dx = \text{too large to display}$$

input `int((a+b*atan(d*x+c))^3/(f*x+e)^2,x)`

output

```
( - 2*atan(c + d*x)**3*b**3*c**5*e*f**5 + 6*atan(c + d*x)**3*b**3*c**4*d*e
**2*f**4 - 6*atan(c + d*x)**3*b**3*c**3*d**2*e**3*f**3 - 2*atan(c + d*x)**
3*b**3*c**3*d**2*e**2*f**4*x - 4*atan(c + d*x)**3*b**3*c**3*e*f**5 + 2*ata
n(c + d*x)**3*b**3*c**2*d**3*e**4*f**2 + 6*atan(c + d*x)**3*b**3*c**2*d**3
*e**3*f**3*x + 8*atan(c + d*x)**3*b**3*c**2*d*e**2*f**4 - 6*atan(c + d*x)*
*3*b**3*c*d**4*e**4*f**2*x - 6*atan(c + d*x)**3*b**3*c*d**2*e**3*f**3 - 2*
atan(c + d*x)**3*b**3*c*d**2*e**2*f**4*x - 2*atan(c + d*x)**3*b**3*c*e*f**
5 + 2*atan(c + d*x)**3*b**3*d**5*e**5*f*x + 2*atan(c + d*x)**3*b**3*d**3*e
**4*f**2 + 2*atan(c + d*x)**3*b**3*d**3*e**3*f**3*x + 2*atan(c + d*x)**3*b
**3*d*e**2*f**4 - 6*atan(c + d*x)**2*a*b**2*c**5*e*f**5 + 15*atan(c + d*x)
**2*a*b**2*c**4*d*e**2*f**4 - 3*atan(c + d*x)**2*a*b**2*c**4*d*e*f**5*x -
6*atan(c + d*x)**2*a*b**2*c**3*d**2*e**3*f**3 + 6*atan(c + d*x)**2*a*b**2*
c**3*d**2*e**2*f**4*x - 12*atan(c + d*x)**2*a*b**2*c**3*e*f**5 - 12*atan(c
+ d*x)**2*a*b**2*c**2*d**3*e**4*f**2 + 18*atan(c + d*x)**2*a*b**2*c**2*d
e**2*f**4 - 6*atan(c + d*x)**2*a*b**2*c**2*d*e*f**5*x + 12*atan(c + d*x)**
2*a*b**2*c*d**4*e**5*f - 6*atan(c + d*x)**2*a*b**2*c*d**4*e**4*f**2*x - 6*
atan(c + d*x)**2*a*b**2*c*d**2*e**3*f**3 + 6*atan(c + d*x)**2*a*b**2*c*d**
2*e**2*f**4*x - 6*atan(c + d*x)**2*a*b**2*c*e*f**5 - 3*atan(c + d*x)**2*a*
b**2*d**5*e**6 + 3*atan(c + d*x)**2*a*b**2*d**5*e**5*f*x + 3*atan(c + d*x)
**2*a*b**2*d*e**2*f**4 - 3*atan(c + d*x)**2*a*b**2*d*e*f**5*x - 6*atan(...
```

3.49 $\int (e + fx)^m (a + b \arctan(c + dx)) dx$

Optimal result	436
Mathematica [A] (verified)	437
Rubi [A] (verified)	437
Maple [F]	439
Fricas [F]	439
Sympy [F(-1)]	439
Maxima [F]	440
Giac [F]	440
Mupad [F(-1)]	441
Reduce [F]	441

Optimal result

Integrand size = 18, antiderivative size = 177

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx$$

$$= \frac{(e + fx)^{1+m} (a + b \arctan(c + dx))}{f(1 + m)}$$

$$- \frac{ibd(e + fx)^{2+m} \operatorname{Hypergeometric2F1}\left(1, 2 + m, 3 + m, \frac{d(e+fx)}{de+(i-c)f}\right)}{2f(de + (i - c)f)(1 + m)(2 + m)}$$

$$+ \frac{ibd(e + fx)^{2+m} \operatorname{Hypergeometric2F1}\left(1, 2 + m, 3 + m, \frac{d(e+fx)}{de-(i+c)f}\right)}{2f(de - (i + c)f)(1 + m)(2 + m)}$$

output

```
(f*x+e)^(1+m)*(a+b*arctan(d*x+c))/f/(1+m)-1/2*I*b*d*(f*x+e)^(2+m)*hypergeo
m([1, 2+m],[3+m],d*(f*x+e)/(d*e+(I-c)*f))/f/(d*e+(I-c)*f)/(1+m)/(2+m)+1/2*
I*b*d*(f*x+e)^(2+m)*hypergeom([1, 2+m],[3+m],d*(f*x+e)/(d*e-(I+c)*f))/f/(d
*e-(I+c)*f)/(1+m)/(2+m)
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.92

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx$$

$$= \frac{(e + fx)^{1+m} \left(2(a + b \arctan(c + dx)) + \frac{bd(e+fx) \left((de-(i+c)f) \operatorname{Hypergeometric2F1} \left(1, 2+m, 3+m, \frac{d(e+fx)}{de-(-i+c)f} \right) + (-de+(-i+c)f) \operatorname{Hypergeometric2F1} \left(1, 2+m, 3+m, \frac{d(e+fx)}{de-(-i+c)f} \right) \right)}{(ide+f-icf)(de-(-i+c)f)(2+m)} \right)}{2f(1+m)}$$

input

```
Integrate[(e + f*x)^m*(a + b*ArcTan[c + d*x]),x]
```

output

```
((e + f*x)^(1 + m)*(2*(a + b*ArcTan[c + d*x]) + (b*d*(e + f*x)*((d*e - (I + c)*f)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*(e + f*x))/(d*e - (-I + c)*f)]) + (-d*e) + (-I + c)*f)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*(e + f*x))/(d*e - (I + c)*f)])/((I*d*e + f - I*c*f)*(d*e - (-I + c)*f)*(2 + m)))/(2*f*(1 + m))
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.34, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5570, 5387, 485, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx$$

$$\downarrow \text{5570}$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \arctan(c + dx)) d(c + dx)}{d}$$

$$\downarrow \text{5387}$$

$$\frac{\frac{d(a+b \arctan(c+dx)) \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+1}}{f(m+1)} - \frac{bd \int \frac{\left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{(c+dx)^2+1} d(c+dx)}{f(m+1)}}{d}$$

$$\frac{d(a+b \arctan(c+dx)) \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+1}}{f(m+1)} - \frac{bd \int \left(\frac{i \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{2(-c-dx+i)} + \frac{i \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^{m+1}}{2(c+dx+i)} \right) d(c+dx)}{f(m+1)}$$

485

d

2009

$$\frac{d(a+b \arctan(c+dx)) \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+1}}{f(m+1)} - \frac{bd \left(\frac{id \left(\frac{f(c+dx)}{d} - \frac{cf}{d} + e \right)^{m+2} \text{Hypergeometric2F1} \left(1, m+2, m+3, \frac{de-cf+f(c+dx)}{de-cf+if} \right)}{2(m+2)(de+(-c+i)f)} - \frac{id \left(\frac{f(c+dx)}{d} - \frac{cf}{d} \right)}{f(m+1)} \right)}{d}$$

input `Int[(e + f*x)^m*(a + b*ArcTan[c + d*x]),x]`

output `((d*(e - (c*f)/d + (f*(c + d*x))/d)^(1 + m)*(a + b*ArcTan[c + d*x]))/(f*(1 + m)) - (b*d*(((I/2)*d*(e - (c*f)/d + (f*(c + d*x))/d)^(2 + m)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*e - c*f + f*(c + d*x))/(d*e + I*f - c*f]])/((d*e + (I - c)*f)*(2 + m)) - ((I/2)*d*(e - (c*f)/d + (f*(c + d*x))/d)^(2 + m)*Hypergeometric2F1[1, 2 + m, 3 + m, (d*e - c*f + f*(c + d*x))/(d*e - (I + c)*f]])/((d*e - (I + c)*f)*(2 + m))))/(f*(1 + m))/d`

Defintions of rubi rules used

rule 485 `Int[((c_) + (d_.)*(x_))^(n_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Int[Expand Integrand[(c + d*x)^n, 1/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, n}, x] & & !IntegerQ[2*n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5387 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.)*((d_) + (e_.)*(x_))^(q_.), x_Symbol] := Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Simp[b*(c/(e*(q + 1))) Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]`

rule 5570

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^(m*(a + b*A
rcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I
GtQ[p, 0]
```

Maple [F]

$$\int (fx + e)^m (a + b \arctan(dx + c)) dx$$

input

```
int((f*x+e)^m*(a+b*arctan(d*x+c)),x)
```

output

```
int((f*x+e)^m*(a+b*arctan(d*x+c)),x)
```

Fricas [F]

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \int (b \arctan(dx + c) + a)(fx + e)^m dx$$

input

```
integrate((f*x+e)^m*(a+b*arctan(d*x+c)),x, algorithm="fricas")
```

output

```
integral((b*arctan(d*x + c) + a)*(f*x + e)^m, x)
```

Sympy [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \text{Timed out}$$

input

```
integrate((f*x+e)**m*(a+b*atan(d*x+c)),x)
```

output

```
Timed out
```


Maxima [F]

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \int (b \arctan(dx + c) + a)(fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c)),x, algorithm="maxima")`

output `1/2*((3*e*m^2 + 2*e*m + (3*f*m^2 + 2*f*m + f)*x + e)*(f*x + e)^m*arctan(d*x + c) + (e*m + (f*m + f)*x + e)*(f*x + e)^m*log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 2*(f*m^3 + f*m^2 + f*m + f)*integrate(-1/2*(2*((c^2 + 1)*f*m^3 + 2*(c^2 + 1)*f*m^2 + (c^2 + 1)*f*m + (d^2*f*m^3 + 2*d^2*f*m^2 + d^2*f*m)*x^2 + 2*(c*d*f*m^3 + 2*c*d*f*m^2 + c*d*f*m)*x)*(f*x + e)^m*arctan(d*x + c) - ((c^2 + 1)*f*m^3 - (c^2 + 1)*f*m + (d^2*f*m^3 - d^2*f*m)*x^2 + 2*(c*d*f*m^3 - c*d*f*m)*x)*(f*x + e)^m*log(d^2*x^2 + 2*c*d*x + c^2 + 1) - 2*((c - 1)*d*e*m^2 - (c + 1)*d*e*m + (d^2*f*m^2 - d^2*f*m)*x^2 + ((d^2*e + (c - 1)*d*f)*m^2 - (d^2*e + (c + 1)*d*f)*m)*x)*(f*x + e)^m)/((c^2 + 1)*f*m^3 + (c^2 + 1)*f*m^2 + (c^2 + 1)*f*m + (d^2*f*m^3 + d^2*f*m^2 + d^2*f*m + d^2*f)*x^2 + (c^2 + 1)*f + 2*(c*d*f*m^3 + c*d*f*m^2 + c*d*f*m + c*d*f)*x), x))*b/(f*m^3 + f*m^2 + f*m + f) + (f*x + e)^(m + 1)*a/(f*(m + 1))`

Giac [F]

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \int (b \arctan(dx + c) + a)(fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c)),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \int (e + fx)^m (a + b \operatorname{atan}(c + dx)) dx$$

input `int((e + f*x)^m*(a + b*atan(c + d*x)),x)`output `int((e + f*x)^m*(a + b*atan(c + d*x)), x)`**Reduce [F]**

$$\int (e + fx)^m (a + b \arctan(c + dx)) dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atan(d*x+c)),x)`

output

```

((e + f*x)**m*atan(c + d*x)*b*c*e*m + (e + f*x)**m*atan(c + d*x)*b*c*f*m*x
+ (e + f*x)**m*a*c*e*m + (e + f*x)**m*a*c*f*m*x - (e + f*x)**m*b*e + int(
(e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x + 2*
c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 + d*
**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*c**2*e*f*m**2 + in
t((e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x +
2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 +
d**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*c**2*e*f*m - int
((e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x + 2
*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 + d
**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*c*d*e**2*m**2 - i
nt((e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x +
2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 +
d**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*c*d*e**2*m + in
t((e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x +
2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 +
d**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*e*f*m**2 + int((
e + f*x)**m/(c**2*e*m + c**2*e + c**2*f*m*x + c**2*f*x + 2*c*d*e*m*x + 2*c
*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 + d**
2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b*e*f*m - int((e ...

```

3.50 $\int (e + fx)^m (a + b \arctan(c + dx))^2 dx$

Optimal result	443
Mathematica [N/A]	443
Rubi [N/A]	444
Maple [N/A]	444
Fricas [N/A]	445
Sympy [F(-1)]	445
Maxima [N/A]	445
Giac [N/A]	446
Mupad [N/A]	446
Reduce [N/A]	447

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \text{Int}((e + fx)^m (a + b \arctan(c + dx))^2, x)$$

output `Defer(Int)((f*x+e)^m*(a+b*arctan(d*x+c))^2,x)`

Mathematica [N/A]

Not integrable

Time = 3.87 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \int (e + fx)^m (a + b \arctan(c + dx))^2 dx$$

input `Integrate[(e + f*x)^m*(a + b*ArcTan[c + d*x])^2,x]`

output `Integrate[(e + f*x)^m*(a + b*ArcTan[c + d*x])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx$$

$$\downarrow 5570$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

$$\downarrow 5560$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \arctan(c + dx))^2 d(c + dx)}{d}$$

input `Int[(e + f*x)^m*(a + b*ArcTan[c + d*x])^2,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (fx + e)^m (a + b \arctan(dx + c))^2 dx$$

input `int((f*x+e)^m*(a+b*arctan(d*x+c))^2,x)`

output `int((f*x+e)^m*(a+b*arctan(d*x+c))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c))^2,x, algorithm="fricas")`

output `integral((b^2*arctan(d*x + c)^2 + 2*a*b*arctan(d*x + c) + a^2)*(f*x + e)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \text{Timed out}$$

input `integrate((f*x+e)**m*(a+b*atan(d*x+c))**2,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 9.07 (sec) , antiderivative size = 504, normalized size of antiderivative = 25.20

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c))^2,x, algorithm="maxima")`

output

```
(f*x + e)^(m + 1)*a^2/(f*(m + 1)) + 1/16*(4*(b^2*f*x + b^2*e)*(f*x + e)^m*
arctan(d*x + c)^2 - (b^2*f*x + b^2*e)*(f*x + e)^m*log(d^2*x^2 + 2*c*d*x +
c^2 + 1)^2 + 16*(f*m + f)*integrate(1/16*(12*((b^2*c^2 + b^2)*f*m + (b^2*d
^2*f*m + b^2*d^2*f)*x^2 + (b^2*c^2 + b^2)*f + 2*(b^2*c*d*f*m + b^2*c*d*f)*
x)*(f*x + e)^m*arctan(d*x + c)^2 + ((b^2*c^2 + b^2)*f*m + (b^2*d^2*f*m + b
^2*d^2*f)*x^2 + (b^2*c^2 + b^2)*f + 2*(b^2*c*d*f*m + b^2*c*d*f)*x)*(f*x +
e)^m*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2 - 8*(b^2*d*e - 4*(a*b*c^2 + a*b)*f
*m - 4*(a*b*d^2*f*m + a*b*d^2*f)*x^2 - 4*(a*b*c^2 + a*b)*f - (8*a*b*c*d*f*
m + (8*a*b*c - b^2)*d*f)*x)*(f*x + e)^m*arctan(d*x + c) + 4*(b^2*d^2*f*x^2
+ b^2*c*d*e + (b^2*d^2*e + b^2*c*d*f)*x)*(f*x + e)^m*log(d^2*x^2 + 2*c*d*
x + c^2 + 1))/((c^2 + 1)*f*m + (d^2*f*m + d^2*f)*x^2 + (c^2 + 1)*f + 2*(c*
d*f*m + c*d*f)*x), x))/(f*m + f)
```

Giac [N/A]

Not integrable

Time = 108.32 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.15

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \int (b \arctan(dx + c) + a)^2 (fx + e)^m dx$$

input

```
integrate((f*x+e)^m*(a+b*arctan(d*x+c))^2,x, algorithm="giac")
```

output

```
sage0*x
```

Mupad [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \int (e + fx)^m (a + b \operatorname{atan}(c + dx))^2 dx$$

input

```
int((e + f*x)^m*(a + b*atan(c + d*x))^2,x)
```

output `int((e + f*x)^m*(a + b*atan(c + d*x))^2, x)`

Reduce [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 17260, normalized size of antiderivative = 863.00

$$\int (e + fx)^m (a + b \arctan(c + dx))^2 dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atan(d*x+c))^2,x)`

output

```
((e + f*x)**m*atan(c + d*x)**2*b**2*c**4*e*f*m**2 + (e + f*x)**m*atan(c +
d*x)**2*b**2*c**4*f**2*m**2*x + (e + f*x)**m*atan(c + d*x)**2*b**2*c**2*e*
f*m**2 + (e + f*x)**m*atan(c + d*x)**2*b**2*c**2*f**2*m**2*x + 2*(e + f*x)
**m*atan(c + d*x)*a*b*c**4*e*f*m**2 + 2*(e + f*x)**m*atan(c + d*x)*a*b*c**
4*f**2*m**2*x + 2*(e + f*x)**m*atan(c + d*x)*a*b*c**2*e*f*m**2 + 2*(e + f*
x)**m*atan(c + d*x)*a*b*c**2*f**2*m**2*x - 2*(e + f*x)**m*atan(c + d*x)*b*
*2*c**2*d*e**2*m + (e + f*x)**m*a**2*c**4*e*f*m**2 + (e + f*x)**m*a**2*c**
4*f**2*m**2*x + (e + f*x)**m*a**2*c**2*e*f*m**2 + (e + f*x)**m*a**2*c**2*f
**2*m**2*x - 2*(e + f*x)**m*a*b*c**3*e*f*m - 2*(e + f*x)**m*a*b*c*e*f*m +
(e + f*x)**m*b**2*c*d*e**2 + int((e + f*x)**m/(c**4*e*m + c**4*e + c**4*f*
m*x + c**4*f*x + 2*c**3*d*e*m*x + 2*c**3*d*e*x + 2*c**3*d*f*m*x**2 + 2*c**
3*d*f*x**2 + c**2*d**2*e*m*x**2 + c**2*d**2*e*x**2 + c**2*d**2*f*m*x**3 +
c**2*d**2*f*x**3 + 2*c**2*e*m + 2*c**2*e + 2*c**2*f*m*x + 2*c**2*f*x + 2*c
*d*e*m*x + 2*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**
2*e*x**2 + d**2*f*m*x**3 + d**2*f*x**3 + e*m + e + f*m*x + f*x),x)*b**2*c*
*6*e*f**2*m**2 + int((e + f*x)**m/(c**4*e*m + c**4*e + c**4*f*m*x + c**4*f
*x + 2*c**3*d*e*m*x + 2*c**3*d*e*x + 2*c**3*d*f*m*x**2 + 2*c**3*d*f*x**2 +
c**2*d**2*e*m*x**2 + c**2*d**2*e*x**2 + c**2*d**2*f*m*x**3 + c**2*d**2*f*
x**3 + 2*c**2*e*m + 2*c**2*e + 2*c**2*f*m*x + 2*c**2*f*x + 2*c*d*e*m*x + 2
*c*d*e*x + 2*c*d*f*m*x**2 + 2*c*d*f*x**2 + d**2*e*m*x**2 + d**2*e*x**2 ...
```


3.51 $\int (e + fx)^m (a + b \arctan(c + dx))^3 dx$

Optimal result	448
Mathematica [N/A]	448
Rubi [N/A]	449
Maple [N/A]	449
Fricas [N/A]	450
Sympy [F(-1)]	450
Maxima [N/A]	450
Giac [N/A]	451
Mupad [N/A]	452
Reduce [N/A]	452

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \text{Int}((e + fx)^m (a + b \arctan(c + dx))^3, x)$$

output `Defer(Int)((f*x+e)^m*(a+b*arctan(d*x+c))^3,x)`

Mathematica [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \int (e + fx)^m (a + b \arctan(c + dx))^3 dx$$

input `Integrate[(e + f*x)^m*(a + b*ArcTan[c + d*x])^3,x]`

output `Integrate[(e + f*x)^m*(a + b*ArcTan[c + d*x])^3, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx$$

$$\downarrow 5570$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

$$\downarrow 5560$$

$$\frac{\int \left(e - \frac{cf}{d} + \frac{f(c+dx)}{d} \right)^m (a + b \arctan(c + dx))^3 d(c + dx)}{d}$$

input `Int[(e + f*x)^m*(a + b*ArcTan[c + d*x])^3,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (fx + e)^m (a + b \arctan(dx + c))^3 dx$$

input `int((f*x+e)^m*(a+b*arctan(d*x+c))^3,x)`

output `int((f*x+e)^m*(a+b*arctan(d*x+c))^3,x)`

Fricas [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c))^3,x, algorithm="fricas")`

output `integral((b^3*arctan(d*x + c)^3 + 3*a*b^2*arctan(d*x + c)^2 + 3*a^2*b*arctan(d*x + c) + a^3)*(f*x + e)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \text{Timed out}$$

input `integrate((f*x+e)**m*(a+b*atan(d*x+c))**3,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 12.19 (sec) , antiderivative size = 659, normalized size of antiderivative = 32.95

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 (fx + e)^m dx$$

input `integrate((f*x+e)^m*(a+b*arctan(d*x+c))^3,x, algorithm="maxima")`

output

```
(f*x + e)^(m + 1)*a^3/(f*(m + 1)) + 1/32*(4*(b^3*f*x + b^3*e)*(f*x + e)^m*
arctan(d*x + c)^3 - 3*(b^3*f*x + b^3*e)*(f*x + e)^m*arctan(d*x + c)*log(d^
2*x^2 + 2*c*d*x + c^2 + 1)^2 + 32*(f*m + f)*integrate(1/32*(28*((b^3*c^2 +
b^3)*f*m + (b^3*d^2*f*m + b^3*d^2*f)*x^2 + (b^3*c^2 + b^3)*f + 2*(b^3*c*d
*f*m + b^3*c*d*f)*x)*(f*x + e)^m*arctan(d*x + c)^3 - 12*(b^3*d*e - 8*(a*b^
2*c^2 + a*b^2)*f*m - 8*(a*b^2*d^2*f*m + a*b^2*d^2*f)*x^2 - 8*(a*b^2*c^2 +
a*b^2)*f - (16*a*b^2*c*d*f*m + (16*a*b^2*c - b^3)*d*f)*x)*(f*x + e)^m*arct
an(d*x + c)^2 + 12*(b^3*d^2*f*x^2 + b^3*c*d*e + (b^3*d^2*e + b^3*c*d*f)*x)
*(f*x + e)^m*arctan(d*x + c)*log(d^2*x^2 + 2*c*d*x + c^2 + 1) + 96*((a^2*b
*c^2 + a^2*b)*f*m + (a^2*b*d^2*f*m + a^2*b*d^2*f)*x^2 + (a^2*b*c^2 + a^2*b
)*f + 2*(a^2*b*c*d*f*m + a^2*b*c*d*f)*x)*(f*x + e)^m*arctan(d*x + c) + 3*(
((b^3*c^2 + b^3)*f*m + (b^3*d^2*f*m + b^3*d^2*f)*x^2 + (b^3*c^2 + b^3)*f +
2*(b^3*c*d*f*m + b^3*c*d*f)*x)*(f*x + e)^m*arctan(d*x + c) + (b^3*d*f*x +
b^3*d*e)*(f*x + e)^m)*log(d^2*x^2 + 2*c*d*x + c^2 + 1)^2)/((c^2 + 1)*f*m
+ (d^2*f*m + d^2*f)*x^2 + (c^2 + 1)*f + 2*(c*d*f*m + c*d*f)*x), x)/(f*m +
f)
```

Giac [N/A]

Not integrable

Time = 107.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.15

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \int (b \arctan(dx + c) + a)^3 (fx + e)^m dx$$

input

```
integrate((f*x+e)^m*(a+b*arctan(d*x+c))^3,x, algorithm="giac")
```

output

```
sage0*x
```

Mupad [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \int (e + fx)^m (a + b \operatorname{atan}(c + dx))^3 dx$$

input `int((e + f*x)^m*(a + b*atan(c + d*x))^3,x)`output `int((e + f*x)^m*(a + b*atan(c + d*x))^3, x)`**Reduce [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 42351, normalized size of antiderivative = 2117.55

$$\int (e + fx)^m (a + b \arctan(c + dx))^3 dx = \text{Too large to display}$$

input `int((f*x+e)^m*(a+b*atan(d*x+c))^3,x)`

output

```
(2*(e + f*x)**m*atan(c + d*x)**3*b**3*c**5*e*f*m**3 + 2*(e + f*x)**m*atan(
c + d*x)**3*b**3*c**5*f**2*m**3*x + 2*(e + f*x)**m*atan(c + d*x)**3*b**3*c
**3*e*f*m**3 + 2*(e + f*x)**m*atan(c + d*x)**3*b**3*c**3*f**2*m**3*x + 6*(
e + f*x)**m*atan(c + d*x)**2*a*b**2*c**5*e*f*m**3 + 6*(e + f*x)**m*atan(c
+ d*x)**2*a*b**2*c**5*f**2*m**3*x + 6*(e + f*x)**m*atan(c + d*x)**2*a*b**2
*c**3*e*f*m**3 + 6*(e + f*x)**m*atan(c + d*x)**2*a*b**2*c**3*f**2*m**3*x -
6*(e + f*x)**m*atan(c + d*x)**2*b**3*c**3*d*e**2*m**2 + 6*(e + f*x)**m*at
an(c + d*x)*a**2*b*c**5*e*f*m**3 + 6*(e + f*x)**m*atan(c + d*x)*a**2*b*c**
5*f**2*m**3*x + 6*(e + f*x)**m*atan(c + d*x)*a**2*b*c**3*e*f*m**3 + 6*(e +
f*x)**m*atan(c + d*x)*a**2*b*c**3*f**2*m**3*x - 12*(e + f*x)**m*atan(c +
d*x)*a*b**2*c**3*d*e**2*m**2 - 6*(e + f*x)**m*atan(c + d*x)*b**3*c**3*e*f*
m + 18*(e + f*x)**m*atan(c + d*x)*b**3*c**2*d*e**2*m - 6*(e + f*x)**m*atan
(c + d*x)*b**3*c*e*f*m + 2*(e + f*x)**m*a**3*c**5*e*f*m**3 + 2*(e + f*x)**
m*a**3*c**5*f**2*m**3*x + 2*(e + f*x)**m*a**3*c**3*e*f*m**3 + 2*(e + f*x)*
**m*a**3*c**3*f**2*m**3*x - 6*(e + f*x)**m*a**2*b*c**4*e*f*m**2 - 6*(e + f
x)**m*a**2*b*c**2*e*f*m**2 + 6*(e + f*x)**m*a*b**2*c**2*d*e**2*m + 3*(e +
f*x)**m*b**3*c**2*e*f - 9*(e + f*x)**m*b**3*c*d*e**2 + 3*(e + f*x)**m*b**3
*e*f + 6*int((e + f*x)**m/(c**4*e*m + c**4*e + c**4*f*m*x + c**4*f*x + 2*c
**3*d*e*m*x + 2*c**3*d*e*x + 2*c**3*d*f*m*x**2 + 2*c**3*d*f*x**2 + c**2*d
**2*e*m*x**2 + c**2*d**2*e*x**2 + c**2*d**2*f*m*x**3 + c**2*d**2*f*x**3 ...
```

3.52 $\int \frac{\arctan(a+bx)}{c+dx^3} dx$

Optimal result	455
Mathematica [A] (verified)	456
Rubi [A] (verified)	457
Maple [C] (warning: unable to verify)	460
Fricas [F]	461
Sympy [F(-1)]	461
Maxima [F]	462
Giac [F]	462
Mupad [F(-1)]	462
Reduce [F]	463

Optimal result

Integrand size = 16, antiderivative size = 725

$$\begin{aligned}
\int \frac{\arctan(a + bx)}{c + dx^3} dx = & -\frac{\arctan(a + bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{\sqrt[3]{-1} \arctan(a + bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& - \frac{(-1)^{2/3} \arctan(a + bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{\arctan(a + bx) \log\left(\frac{2b\left(\sqrt[3]{c} + \sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} + (i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{(-1)^{2/3} \arctan(a + bx) \log\left(\frac{2b\left(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} - \sqrt[3]{-1}(i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& - \frac{\sqrt[3]{-1} \arctan(a + bx) \log\left(\frac{2b\left(\sqrt[3]{c} + (-1)^{2/3}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} + (-1)^{2/3}(i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{3c^{2/3}\sqrt[3]{d}} \\
& + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{\sqrt[6]{-1} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{(-1)^{5/6} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& - \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2b\left(\sqrt[3]{c} + \sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} + (i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& + \frac{\sqrt[6]{-1} \operatorname{PolyLog}\left(2, 1 - \frac{2b\left(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} - \sqrt[3]{-1}(i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{6c^{2/3}\sqrt[3]{d}} \\
& + \frac{(-1)^{5/6} \operatorname{PolyLog}\left(2, 1 - \frac{2b\left(\sqrt[3]{c} + (-1)^{2/3}\sqrt[3]{dx}\right)}{\left(b\sqrt[3]{c} + (-1)^{2/3}(i-a)\sqrt[3]{d}\right)(1-i(a+bx))}\right)}{6c^{2/3}\sqrt[3]{d}}
\end{aligned}$$

output

```

-1/3*arctan(b*x+a)*ln(2/(1-I*(b*x+a)))/c^(2/3)/d^(1/3)+1/3*(-1)^(1/3)*arct
an(b*x+a)*ln(2/(1-I*(b*x+a)))/c^(2/3)/d^(1/3)-1/3*(-1)^(2/3)*arctan(b*x+a)
*ln(2/(1-I*(b*x+a)))/c^(2/3)/d^(1/3)+1/3*arctan(b*x+a)*ln(2*b*(c^(1/3)+d^(
1/3)*x)/(b*c^(1/3)+(I-a)*d^(1/3))/(1-I*(b*x+a))/c^(2/3)/d^(1/3)+1/3*(-1)^(
2/3)*arctan(b*x+a)*ln(2*b*(c^(1/3)-(-1)^(1/3)*d^(1/3)*x)/(b*c^(1/3)-(-1)^(
1/3)*(I-a)*d^(1/3))/(1-I*(b*x+a))/c^(2/3)/d^(1/3)-1/3*(-1)^(1/3)*arctan(
b*x+a)*ln(2*b*(c^(1/3)+(-1)^(2/3)*d^(1/3)*x)/(b*c^(1/3)+(-1)^(2/3)*(I-a)*d
^(1/3))/(1-I*(b*x+a))/c^(2/3)/d^(1/3)+1/6*I*polylog(2,1-2/(1-I*(b*x+a)))/
c^(2/3)/d^(1/3)-1/6*(-1)^(1/6)*polylog(2,1-2/(1-I*(b*x+a)))/c^(2/3)/d^(1/3
)-1/6*(-1)^(5/6)*polylog(2,1-2/(1-I*(b*x+a)))/c^(2/3)/d^(1/3)-1/6*I*polylo
g(2,1-2*b*(c^(1/3)+d^(1/3)*x)/(b*c^(1/3)+(I-a)*d^(1/3))/(1-I*(b*x+a))/c^(
2/3)/d^(1/3)+1/6*(-1)^(1/6)*polylog(2,1-2*b*(c^(1/3)-(-1)^(1/3)*d^(1/3)*x)
/(b*c^(1/3)-(-1)^(1/3)*(I-a)*d^(1/3))/(1-I*(b*x+a))/c^(2/3)/d^(1/3)+1/6*(
-1)^(5/6)*polylog(2,1-2*b*(c^(1/3)+(-1)^(2/3)*d^(1/3)*x)/(b*c^(1/3)+(-1)^(
2/3)*(I-a)*d^(1/3))/(1-I*(b*x+a))/c^(2/3)/d^(1/3)

```

Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 701, normalized size of antiderivative = 0.97

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx$$

$$= \frac{-i \log(1 + ia + ibx) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (-i+a)\sqrt[3]{d}}\right) + i \log(-i(i + a + bx)) \log\left(\frac{b(\sqrt[3]{c} + \sqrt[3]{dx})}{b\sqrt[3]{c} - (i+a)\sqrt[3]{d}}\right) + \sqrt[6]{-1} \log(1 + \dots)}{\dots}$$

input

```
Integrate[ArcTan[a + b*x]/(c + d*x^3), x]
```

output

```

((-I)*Log[1 + I*a + I*b*x]*Log[(b*(c^(1/3) + d^(1/3)*x))/(b*c^(1/3) - (-I
+ a)*d^(1/3))] + I*Log[(-I)*(I + a + b*x)]*Log[(b*(c^(1/3) + d^(1/3)*x))/(
b*c^(1/3) - (I + a)*d^(1/3))] + (-1)^(1/6)*Log[1 + I*a + I*b*x]*Log[(b*(c^
(1/3) - (-1)^(1/3)*d^(1/3)*x))/(b*c^(1/3) + (-1)^(1/3)*(-I + a)*d^(1/3))]
- (-1)^(1/6)*Log[(-I)*(I + a + b*x)]*Log[(b*(c^(1/3) - (-1)^(1/3)*d^(1/3)*
x))/(b*c^(1/3) + (-1)^(1/3)*(I + a)*d^(1/3))] - (-1)^(5/6)*Log[(-I)*(I + a
+ b*x)]*Log[(b*(c^(1/3) + (-1)^(2/3)*d^(1/3)*x))/(b*c^(1/3) + (-1)^(1/6)*
(1 - I*a)*d^(1/3))] + (-1)^(5/6)*Log[1 + I*a + I*b*x]*Log[(b*(c^(1/3) + (-
1)^(2/3)*d^(1/3)*x))/(b*c^(1/3) - (-1)^(2/3)*(-I + a)*d^(1/3))] - I*PolyLo
g[2, (d^(1/3)*(-I + a + b*x))/(-b*c^(1/3)) + (-I + a)*d^(1/3))] + (-1)^(5
/6)*PolyLog[2, ((-1)^(1/6)*d^(1/3)*(-I + a + b*x))/(I*b*c^(1/3) + (-1)^(1/
6)*(-I + a)*d^(1/3))] + (-1)^(1/6)*PolyLog[2, ((-1)^(1/3)*d^(1/3)*(-I + a
+ b*x))/(b*c^(1/3) + (-1)^(1/3)*(-I + a)*d^(1/3))] + I*PolyLog[2, (d^(1/3)
*(I + a + b*x))/(-b*c^(1/3)) + (I + a)*d^(1/3))] - (-1)^(1/6)*PolyLog[2,
((-1)^(1/3)*d^(1/3)*(I + a + b*x))/(b*c^(1/3) + (-1)^(1/3)*(I + a)*d^(1/3
))] - (-1)^(5/6)*PolyLog[2, ((-1)^(2/3)*d^(1/3)*(I + a + b*x))/(-b*c^(1/3)
) + (-1)^(2/3)*(I + a)*d^(1/3)]]/(6*c^(2/3)*d^(1/3))

```

Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 869, normalized size of antiderivative = 1.20, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(a + bx)}{c + dx^3} dx \\
 & \quad \downarrow \text{5574} \\
 & \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{dx^3 + c} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{dx^3 + c} dx \\
 & \quad \downarrow \text{2856}
 \end{aligned}$$

$$\frac{1}{2}i \int \left(-\frac{\log(-ia - ibx + 1)}{3c^{2/3}(-\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(-ia - ibx + 1)}{3c^{2/3}(\sqrt[3]{-1}\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(-ia - ibx + 1)}{3c^{2/3}(-(-1)^{2/3}\sqrt[3]{dx} - \sqrt[3]{c})} \right) dx -$$

$$\frac{1}{2}i \int \left(-\frac{\log(ia + ibx + 1)}{3c^{2/3}(-\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(ia + ibx + 1)}{3c^{2/3}(\sqrt[3]{-1}\sqrt[3]{dx} - \sqrt[3]{c})} - \frac{\log(ia + ibx + 1)}{3c^{2/3}(-(-1)^{2/3}\sqrt[3]{dx} - \sqrt[3]{c})} \right) dx$$

↓ 2009

$$\frac{1}{2}i \left(\frac{\log(-ia - ibx + 1) \log\left(\frac{b(\sqrt[3]{dx} + \sqrt[3]{c})}{b\sqrt[3]{c} - (a+i)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} + \frac{(-1)^{2/3} \log(-ia - ibx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{\sqrt[3]{-1}\sqrt[3]{d}(a+i) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} - \frac{\sqrt[3]{-1} \log(-ia - ibx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{\sqrt[3]{-1}\sqrt[3]{d}(a+i) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} \right)$$

$$\frac{1}{2}i \left(\frac{\log(ia + ibx + 1) \log\left(\frac{b(\sqrt[3]{dx} + \sqrt[3]{c})}{\sqrt[3]{d}(i-a) + b\sqrt[3]{c}}\right)}{3c^{2/3}\sqrt[3]{d}} + \frac{(-1)^{2/3} \log(ia + ibx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{b\sqrt[3]{c} - \sqrt[3]{-1}(i-a)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} - \frac{\sqrt[3]{-1} \log(ia + ibx + 1) \log\left(\frac{b(\sqrt[3]{c} - \sqrt[3]{-1}\sqrt[3]{dx})}{b\sqrt[3]{c} - \sqrt[3]{-1}(i-a)\sqrt[3]{d}}\right)}{3c^{2/3}\sqrt[3]{d}} \right)$$

input Int[ArcTan[a + b*x]/(c + d*x^3),x]

output

$$\begin{aligned}
& (-1/2*I)*((\text{Log}[1 + I*a + I*b*x]*\text{Log}[(b*(c^{1/3} + d^{1/3}*x))/(b*c^{1/3} + (I - a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) + ((-1)^{2/3}*\text{Log}[1 + I*a + I*b*x] \\
& *\text{Log}[(b*(c^{1/3} - (-1)^{1/3}*d^{1/3}*x))/(b*c^{1/3} - (-1)^{1/3}*(I - a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) - ((-1)^{1/3}*\text{Log}[1 + I*a + I*b*x]*\text{Log}[(b*(c^{1/3} + (-1)^{2/3}*d^{1/3}*x))/(b*c^{1/3} + (-1)^{2/3}*(I - a)*d^{1/3})]) \\
&)/ (3*c^{2/3}*d^{1/3})) + \text{PolyLog}[2, (d^{1/3}*(I - a - b*x))/(b*c^{1/3} + (I - a)*d^{1/3})]/ (3*c^{2/3}*d^{1/3})) - ((-1)^{1/3}*\text{PolyLog}[2, -(((-1)^{1/6})*d^{1/3}*(I - a - b*x))/(I*b*c^{1/3} - (-1)^{1/6}*(I - a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) + ((-1)^{2/3}*\text{PolyLog}[2, -(((-1)^{1/3})*d^{1/3}*(I - a - b*x))/(b*c^{1/3} - (-1)^{1/3}*(I - a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) + (I/2)*((\text{Log}[1 - I*a - I*b*x]*\text{Log}[(b*(c^{1/3} + d^{1/3}*x))/(b*c^{1/3} - (I + a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) + ((-1)^{2/3}*\text{Log}[1 - I*a - I*b*x]*\text{Log}[(b*(c^{1/3} - (-1)^{1/3}*d^{1/3}*x))/(b*c^{1/3} + (-1)^{1/3}*(I + a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) - ((-1)^{1/3}*\text{Log}[1 - I*a - I*b*x]*\text{Log}[(b*(c^{1/3} + (-1)^{2/3}*d^{1/3}*x))/(b*c^{1/3} + (-1)^{1/6}*(1 - I*a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) + \text{PolyLog}[2, -((d^{1/3}*(I + a + b*x))/(b*c^{1/3} - (I + a)*d^{1/3}))]/ (3*c^{2/3}*d^{1/3})) + ((-1)^{2/3}*\text{PolyLog}[2, ((-1)^{1/3})*d^{1/3}*(I + a + b*x))/(b*c^{1/3} + (-1)^{1/3}*(I + a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3})) - ((-1)^{1/3}*\text{PolyLog}[2, -(((-1)^{2/3})*d^{1/3}*(I + a + b*x))/(b*c^{1/3} - (-1)^{2/3}*(I + a)*d^{1/3})])/ (3*c^{2/3}*d^{1/3}))
\end{aligned}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2856 $\text{Int}[(a_.) + \text{Log}[(c_.)*((d_.) + (e_.)*(x_))^{(n_.)}*(b_.)^{(p_.)}*((f_.) + (g_.)*(x_))^{(r_.)}])^{(q_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*\text{Log}[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, n, r\}, x] \&\& I \text{GtQ}[p, 0] \&\& \text{IntegerQ}[q] \&\& (\text{GtQ}[q, 0] \|\| (\text{IntegerQ}[r] \&\& \text{NeQ}[r, 1]))]$

rule 5574 $\text{Int}[\text{ArcTan}[(a_.) + (b_.)*(x_)]/((c_.) + (d_.)*(x_))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[I/2 \text{Int}[\text{Log}[1 - I*a - I*b*x]/(c + d*x^n), x], x] - \text{Simp}[I/2 \text{Int}[\text{Log}[1 + I*a + I*b*x]/(c + d*x^n), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{RationalQ}[n]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.90 (sec) , antiderivative size = 380, normalized size of antiderivative = 0.52

method	result
risch	$ib^2 \left(\frac{\sum_{R1=\text{RootOf}(dZ^3+(3\text{RootOf}(_Z^2+1,\text{index}=1)ad-3d)_Z^2+(-6\text{RootOf}(_Z^2+1,\text{index}=1)ad-3a^2d+3d)_Z-\text{Ro}}$
derivativedivides	$\frac{b^3 \left(\frac{\sum_{R=\text{RootOf}(dZ^3-3adZ^2+3a^2dZ-a^3d+b^3c)} \ln(bx-_R+a)}{-R^2+2_R a-a^2} \arctan(bx+a) \right)}{3d} + \left(b^3 \arctan(bx+a) \left(_R=\text{Ro}$
default	$\frac{b^3 \left(\frac{\sum_{R=\text{RootOf}(dZ^3-3adZ^2+3a^2dZ-a^3d+b^3c)} \ln(bx-_R+a)}{-R^2+2_R a-a^2} \arctan(bx+a) \right)}{3d} + \left(b^3 \arctan(bx+a) \left(_R=\text{Ro}$

input `int(arctan(b*x+a)/(d*x^3+c),x,method=_RETURNVERBOSE)`

output

```
-1/6*I*b^2/d*sum(1/(1+2*I*a*_R1-2*I*a+_R1^2-a^2-2*_R1)*(ln(1-I*a-I*b*x)*ln
((R1+I*b*x+I*a-1)/R1)+dilog((R1+I*b*x+I*a-1)/R1)),_R1=RootOf(d*_Z^3+(3
*RootOf(_Z^2+1,index=1)*a*d-3*d)*_Z^2+(-6*RootOf(_Z^2+1,index=1)*a*d-3*a^2
*d+3*d)*_Z-RootOf(_Z^2+1,index=1)*a^3*d+RootOf(_Z^2+1,index=1)*b^3*c+3*Ro
otOf(_Z^2+1,index=1)*a*d+3*a^2*d-d))+1/6*I*b^2/d*sum(1/(1-2*I*a*_R1+2*I*a+_
R1^2-a^2-2*_R1)*(ln(1+I*a+I*b*x)*ln((R1-I*b*x-I*a-1)/R1)+dilog((R1-I*b*
x-I*a-1)/R1)),_R1=RootOf(d*_Z^3+(-3*RootOf(_Z^2+1,index=1)*a*d-3*d)*_Z^2+
(6*RootOf(_Z^2+1,index=1)*a*d-3*a^2*d+3*d)*_Z+RootOf(_Z^2+1,index=1)*a^3*d
-RootOf(_Z^2+1,index=1)*b^3*c-3*RootOf(_Z^2+1,index=1)*a*d+3*a^2*d-d))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \int \frac{\arctan(bx + a)}{dx^3 + c} dx$$

input

```
integrate(arctan(b*x+a)/(d*x^3+c),x, algorithm="fricas")
```

output

```
integral(arctan(b*x + a)/(d*x^3 + c), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \text{Timed out}$$

input

```
integrate(atan(b*x+a)/(d*x**3+c),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \int \frac{\arctan(bx + a)}{dx^3 + c} dx$$

input `integrate(arctan(b*x+a)/(d*x^3+c),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(d*x^3 + c), x)`

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \int \frac{\arctan(bx + a)}{dx^3 + c} dx$$

input `integrate(arctan(b*x+a)/(d*x^3+c),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(d*x^3 + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{atan}(a + bx)}{dx^3 + c} dx$$

input `int(atan(a + b*x)/(c + d*x^3),x)`

output `int(atan(a + b*x)/(c + d*x^3), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + dx^3} dx = \int \frac{\operatorname{atan}(bx + a)}{dx^3 + c} dx$$

input `int(atan(b*x+a)/(d*x^3+c),x)`

output `int(atan(a + b*x)/(c + d*x**3),x)`

3.53 $\int \frac{\arctan(a+bx)}{c+dx^2} dx$

Optimal result	464
Mathematica [A] (warning: unable to verify)	465
Rubi [A] (verified)	465
Maple [B] (verified)	467
Fricas [F]	468
Sympy [F(-1)]	468
Maxima [B] (verification not implemented)	469
Giac [F]	470
Mupad [F(-1)]	470
Reduce [F]	470

Optimal result

Integrand size = 16, antiderivative size = 319

$$\int \frac{\arctan(a+bx)}{c+dx^2} dx = \frac{\arctan(a+bx) \log\left(\frac{2b(\sqrt{-c}-\sqrt{dx})}{(b\sqrt{-c}-(i-a)\sqrt{d})(1-i(a+bx))}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{\arctan(a+bx) \log\left(\frac{2b(\sqrt{-c}+\sqrt{dx})}{(b\sqrt{-c}+(i-a)\sqrt{d})(1-i(a+bx))}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2b(\sqrt{-c}-\sqrt{dx})}{(b\sqrt{-c}-(i-a)\sqrt{d})(1-i(a+bx))}\right)}{4\sqrt{-c}\sqrt{d}} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2b(\sqrt{-c}+\sqrt{dx})}{(b\sqrt{-c}+(i-a)\sqrt{d})(1-i(a+bx))}\right)}{4\sqrt{-c}\sqrt{d}}$$

output

```
1/2*arctan(b*x+a)*ln(2*b*((-c)^(1/2)-d^(1/2)*x)/(b*(-c)^(1/2)-(I-a)*d^(1/2)))/(1-I*(b*x+a))/(-c)^(1/2)/d^(1/2)-1/2*arctan(b*x+a)*ln(2*b*((-c)^(1/2)+d^(1/2)*x)/(b*(-c)^(1/2)+(I-a)*d^(1/2)))/(1-I*(b*x+a))/(-c)^(1/2)/d^(1/2)-1/4*I*polylog(2,1-2*b*((-c)^(1/2)-d^(1/2)*x)/(b*(-c)^(1/2)-(I-a)*d^(1/2)))/(1-I*(b*x+a))/(-c)^(1/2)/d^(1/2)+1/4*I*polylog(2,1-2*b*((-c)^(1/2)+d^(1/2)*x)/(b*(-c)^(1/2)+(I-a)*d^(1/2)))/(1-I*(b*x+a))/(-c)^(1/2)/d^(1/2)
```

Mathematica [A] (warning: unable to verify)

Time = 0.23 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.28

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx =$$

$$i \left(\log(1 + ia + ibx) \log \left(\frac{b(\sqrt{-c} - \sqrt{dx})}{b\sqrt{-c} + (i+a)\sqrt{d}} \right) - \log(-i(i + a + bx)) \log \left(\frac{b(\sqrt{-c} - \sqrt{dx})}{b\sqrt{-c} + (i+a)\sqrt{d}} \right) - \log(1 + ia + ibx) \right)$$

input `Integrate[ArcTan[a + b*x]/(c + d*x^2),x]`

output

```
((-1/4*I)*(Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (-I + a)*Sqrt[d]]) - Log[(-I)*(I + a + b*x)]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (I + a)*Sqrt[d]]) - Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (-I + a)*Sqrt[d]]) + Log[(-I)*(I + a + b*x)]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (I + a)*Sqrt[d]]) - PolyLog[2, (Sqrt[d]*(-I + a + b*x))/(-b*Sqrt[-c]) + (-I + a)*Sqrt[d]] + PolyLog[2, (Sqrt[d]*(-I + a + b*x))/(b*Sqrt[-c] + (-I + a)*Sqrt[d]] + PolyLog[2, (Sqrt[d]*(I + a + b*x))/(-b*Sqrt[-c]) + (I + a)*Sqrt[d]] - PolyLog[2, (Sqrt[d]*(I + a + b*x))/(b*Sqrt[-c] + (I + a)*Sqrt[d])])/(Sqrt[-c]*Sqrt[d])
```

Rubi [A] (verified)Time = 0.97 (sec) , antiderivative size = 541, normalized size of antiderivative = 1.70, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx$$

$$\downarrow 5574$$

$$\frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{dx^2 + c} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{dx^2 + c} dx$$

$$\begin{aligned}
 & \downarrow 2856 \\
 & \frac{1}{2}i \int \left(\frac{\sqrt{-c} \log(-ia - ibx + 1)}{2c(\sqrt{-c} - \sqrt{dx})} + \frac{\sqrt{-c} \log(-ia - ibx + 1)}{2c(\sqrt{dx} + \sqrt{-c})} \right) dx - \\
 & \frac{1}{2}i \int \left(\frac{\sqrt{-c} \log(ia + ibx + 1)}{2c(\sqrt{-c} - \sqrt{dx})} + \frac{\sqrt{-c} \log(ia + ibx + 1)}{2c(\sqrt{dx} + \sqrt{-c})} \right) dx \\
 & \downarrow 2009 \\
 & \frac{1}{2}i \left(\frac{\text{PolyLog}\left(2, -\frac{\sqrt{d}(a+bx+i)}{b\sqrt{-c}-(a+i)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\text{PolyLog}\left(2, \frac{\sqrt{d}(a+bx+i)}{\sqrt{d}(a+i)+b\sqrt{-c}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\log(-ia - ibx + 1) \log\left(\frac{b(\sqrt{-c}-\sqrt{dx})}{b\sqrt{-c}+(a+i)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} \right) - \\
 & \frac{1}{2}i \left(\frac{\text{PolyLog}\left(2, -\frac{\sqrt{d}(-a-bx+i)}{b\sqrt{-c}-(i-a)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} - \frac{\text{PolyLog}\left(2, \frac{\sqrt{d}(-a-bx+i)}{\sqrt{d}(i-a)+b\sqrt{-c}}\right)}{2\sqrt{-c}\sqrt{d}} + \frac{\log(ia + ibx + 1) \log\left(\frac{b(\sqrt{-c}-\sqrt{dx})}{b\sqrt{-c}-(-a+i)\sqrt{d}}\right)}{2\sqrt{-c}\sqrt{d}} \right) - \dots
 \end{aligned}$$

input `Int[ArcTan[a + b*x]/(c + d*x^2),x]`

output

```

(-1/2*I)*((Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] - (I - a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - (Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] + (I - a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) + PolyLog[2, -((Sqrt[d]*(I - a - b*x))/(b*Sqrt[-c] - (I - a)*Sqrt[d]))]/(2*Sqrt[-c]*Sqrt[d]) - PolyLog[2, (Sqrt[d]*(I - a - b*x))/(b*Sqrt[-c] + (I - a)*Sqrt[d])]/(2*Sqrt[-c]*Sqrt[d])) + (I/2)*((Log[1 - I*a - I*b*x]*Log[(b*(Sqrt[-c] - Sqrt[d]*x))/(b*Sqrt[-c] + (I + a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - (Log[1 - I*a - I*b*x]*Log[(b*(Sqrt[-c] + Sqrt[d]*x))/(b*Sqrt[-c] - (I + a)*Sqrt[d])])/(2*Sqrt[-c]*Sqrt[d]) - PolyLog[2, -((Sqrt[d]*(I + a + b*x))/(b*Sqrt[-c] - (I + a)*Sqrt[d]))]/(2*Sqrt[-c]*Sqrt[d]) + PolyLog[2, (Sqrt[d]*(I + a + b*x))/(b*Sqrt[-c] + (I + a)*Sqrt[d])]/(2*Sqrt[-c]*Sqrt[d]))
    
```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2856 Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.
)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)
^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I
GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))
```

```
rule 5574 Int[ArcTan[(a_) + (b_.)*(x_)]/((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 541 vs. 2(253) = 506.

Time = 1.14 (sec) , antiderivative size = 542, normalized size of antiderivative = 1.70

method	result
risch	$\frac{\ln(-bxi-ai+1) \ln\left(\frac{iad-b\sqrt{cd}+(-bxi-ai+1)d-d}{iad-b\sqrt{cd}-d}\right)\sqrt{cd}}{4cd} - \frac{\ln(-bxi-ai+1) \ln\left(\frac{iad+b\sqrt{cd}+(-bxi-ai+1)d-d}{iad+b\sqrt{cd}-d}\right)\sqrt{cd}}{4cd} - \dots$
derivativedivides	Expression too large to display
default	Expression too large to display

```
input int(arctan(b*x+a)/(d*x^2+c),x,method=_RETURNVERBOSE)
```

output

```
1/4*ln(1-I*a-I*b*x)/c/d*ln((I*a*d-b*(c*d)^(1/2)+(1-I*a-I*b*x)*d-d)/(I*a*d-
b*(c*d)^(1/2)-d))*(c*d)^(1/2)-1/4*ln(1-I*a-I*b*x)/c/d*ln((I*a*d+b*(c*d)^(1
/2)+(1-I*a-I*b*x)*d-d)/(I*a*d+b*(c*d)^(1/2)-d))*(c*d)^(1/2)-1/4/c/d*dilog(
(I*a*d+b*(c*d)^(1/2)+(1-I*a-I*b*x)*d-d)/(I*a*d+b*(c*d)^(1/2)-d))*(c*d)^(1/
2)+1/4/c/d*dilog((I*a*d-b*(c*d)^(1/2)+(1-I*a-I*b*x)*d-d)/(I*a*d-b*(c*d)^(1
/2)-d))*(c*d)^(1/2)+1/4*ln(1+I*a+I*b*x)/c/d*ln((I*a*d+b*(c*d)^(1/2)-(1+I*a
+I*b*x)*d+d)/(I*a*d+b*(c*d)^(1/2)+d))*(c*d)^(1/2)-1/4*ln(1+I*a+I*b*x)/c/d*
ln((I*a*d-b*(c*d)^(1/2)-(1+I*a+I*b*x)*d+d)/(I*a*d-b*(c*d)^(1/2)+d))*(c*d)^(
1/2)+1/4/c/d*dilog((I*a*d+b*(c*d)^(1/2)-(1+I*a+I*b*x)*d+d)/(I*a*d+b*(c*d)
^(1/2)+d))*(c*d)^(1/2)-1/4/c/d*dilog((I*a*d-b*(c*d)^(1/2)-(1+I*a+I*b*x)*d+
d)/(I*a*d-b*(c*d)^(1/2)+d))*(c*d)^(1/2)
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \int \frac{\arctan(bx + a)}{dx^2 + c} dx$$

input

```
integrate(arctan(b*x+a)/(d*x^2+c),x, algorithm="fricas")
```

output

```
integral(arctan(b*x + a)/(d*x^2 + c), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \text{Timed out}$$

input

```
integrate(atan(b*x+a)/(d*x**2+c),x)
```

output

```
Timed out
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 8520 vs. $2(235) = 470$.

Time = 4.49 (sec) , antiderivative size = 8520, normalized size of antiderivative = 26.71

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \text{Too large to display}$$

input `integrate(arctan(b*x+a)/(d*x^2+c),x, algorithm="maxima")`

output

```
1/8*b*(8*arctan(d*x/sqrt(c*d))*arctan((b^2*x + a*b)/b)/b - (4*arctan(sqrt(d)*x/sqrt(c))*arctan2((2*a*b^2*c*d + (a*b^3*c + (a^3 + a)*b*d + (b^4*c + (a^2 + 3)*b^2*d)*x)*sqrt(c)*sqrt(d) + (3*b^3*c*d + (a^2 + 1)*b*d^2)*x)/(b^4*c^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 + 4*(b^3*c + (a^2 + 1)*b*d)*sqrt(c)*sqrt(d)), ((a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 + (2*a*b^2*d*x + b^3*c + 3*(a^2 + 1)*b*d)*sqrt(c)*sqrt(d) + (a*b^3*c*d + (a^3 + a)*b*d^2)*x)/(b^4*c^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 + 4*(b^3*c + (a^2 + 1)*b*d)*sqrt(c)*sqrt(d))) + 4*arctan(sqrt(d)*x/sqrt(c))*arctan2((2*a*b^2*c*d - (a*b^3*c + (a^3 + a)*b*d + (b^4*c + (a^2 + 3)*b^2*d)*x)*sqrt(c)*sqrt(d) + (3*b^3*c*d + (a^2 + 1)*b*d^2)*x)/(b^4*c^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 - 4*(b^3*c + (a^2 + 1)*b*d)*sqrt(c)*sqrt(d)), ((a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 - (2*a*b^2*d*x + b^3*c + 3*(a^2 + 1)*b*d)*sqrt(c)*sqrt(d) + (a*b^3*c*d + (a^3 + a)*b*d^2)*x)/(b^4*c^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*d^2 - 4*(b^3*c + (a^2 + 1)*b*d)*sqrt(c)*sqrt(d))) + log(d*x^2 + c)*log(((a^2 + 1)*b^22*c^11*d + 11*(a^4 + 22*a^2 + 21)*b^20*c^10*d^2 + 55*(a^6 + 39*a^4 + 171*a^2 + 133)*b^18*c^9*d^3 + 33*(5*a^8 + 260*a^6 + 1870*a^4 + 3876*a^2 + 2261)*b^16*c^8*d^4 + 330*(a^10 + 61*a^8 + 570*a^6 + 1802*a^4 + 2261*a^2 + 969)*b^14*c^7*d^5 + 22*(21*a^12 + 1386*a^10 + 15015*a^8 + 60060*a^6 + 109395*a^4 + 92378*a^2 + 29393)*b^12*c^6*d^6 + 22*(21*a^14 + 1407*a^12 + 16401*a^10 + 75075*a^8 + 169...
```

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \int \frac{\arctan(bx + a)}{dx^2 + c} dx$$

input `integrate(arctan(b*x+a)/(d*x^2+c),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(d*x^2 + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{atan}(a + bx)}{dx^2 + c} dx$$

input `int(atan(a + b*x)/(c + d*x^2),x)`

output `int(atan(a + b*x)/(c + d*x^2), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + dx^2} dx = \int \frac{\operatorname{atan}(bx + a)}{dx^2 + c} dx$$

input `int(atan(b*x+a)/(d*x^2+c),x)`

output `int(atan(a + b*x)/(c + d*x**2),x)`

3.54 $\int \frac{\arctan(a+bx)}{c+dx} dx$

Optimal result	471
Mathematica [A] (verified)	472
Rubi [A] (verified)	472
Maple [A] (verified)	475
Fricas [F]	475
Sympy [F(-1)]	476
Maxima [B] (verification not implemented)	476
Giac [F]	477
Mupad [F(-1)]	477
Reduce [F]	477

Optimal result

Integrand size = 14, antiderivative size = 152

$$\int \frac{\arctan(a+bx)}{c+dx} dx = -\frac{\arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{d} + \frac{\arctan(a+bx) \log\left(\frac{2b(c+dx)}{(bc+(i-a)d)(1-i(a+bx))}\right)}{d} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{2d} - \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2b(c+dx)}{(bc+(i-a)d)(1-i(a+bx))}\right)}{2d}$$

output

```
-arctan(b*x+a)*ln(2/(1-I*(b*x+a)))/d+arctan(b*x+a)*ln(2*b*(d*x+c)/(b*c+(I-a)*d)/(1-I*(b*x+a)))/d+1/2*I*polylog(2,1-2/(1-I*(b*x+a)))/d-1/2*I*polylog(2,1-2*b*(d*x+c)/(b*c+(I-a)*d)/(1-I*(b*x+a)))/d
```


Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.52

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \frac{i \log(1 - i(a + bx)) \log\left(-\frac{i\left(\frac{bc-ad}{b} + \frac{d(a+bx)}{b}\right)}{-\frac{d}{b} - \frac{i(bc-ad)}{b}}\right)}{2d} - \frac{i \log(1 + i(a + bx)) \log\left(\frac{i\left(\frac{bc-ad}{b} + \frac{d(a+bx)}{b}\right)}{-\frac{d}{b} + \frac{i(bc-ad)}{b}}\right)}{2d} + \frac{i \operatorname{PolyLog}\left(2, -\frac{id(1-i(a+bx))}{bc-id-ad}\right)}{2d} - \frac{i \operatorname{PolyLog}\left(2, \frac{id(1+i(a+bx))}{bc+id-ad}\right)}{2d}$$

input `Integrate[ArcTan[a + b*x]/(c + d*x), x]`output `((I/2)*Log[1 - I*(a + b*x)]*Log[(-I)*((b*c - a*d)/b + (d*(a + b*x))/b)]/(-d/b - (I*(b*c - a*d))/b)]/d - ((I/2)*Log[1 + I*(a + b*x)]*Log[(I*((b*c - a*d)/b + (d*(a + b*x))/b))/(-d/b + (I*(b*c - a*d))/b)]/d + ((I/2)*PolyLog[2, ((-I)*d*(1 - I*(a + b*x)))/(b*c - I*d - a*d)]/d - ((I/2)*PolyLog[2, (I*d*(1 + I*(a + b*x)))/(b*c + I*d - a*d)]/d`**Rubi [A] (verified)**Time = 0.53 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5570, 27, 5381, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{c + dx} dx$$

↓ 5570

$$\int \frac{b \arctan(a+bx)}{b\left(c - \frac{ad}{b}\right) + d(a+bx)} d(a + bx)$$

b

$$\begin{aligned}
& \downarrow 27 \\
& \int \frac{\arctan(a+bx)}{d(a+bx)-ad+bc} d(a+bx) \\
& \downarrow 5381 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+id)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx)}{d} + \frac{\int \frac{\log\left(\frac{2}{1-i(a+bx)}\right)}{(a+bx)^2+1} d(a+bx)}{d} + \\
& \frac{\arctan(a+bx) \log\left(\frac{d}{(1-i(a+bx))(-ad+bc+id)}\right)}{d} - \frac{\arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{d} \\
& \downarrow 2849 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+id)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx)}{d} + \frac{i \int \frac{\log\left(\frac{2}{1-i(a+bx)}\right)}{1-i(a+bx)} d(a+bx)}{d} + \\
& \frac{\arctan(a+bx) \log\left(\frac{d}{(1-i(a+bx))(-ad+bc+id)}\right)}{d} - \frac{\arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{d} \\
& \downarrow 2752 \\
& -\frac{\int \frac{\log\left(\frac{2(bc-ad+d(a+bx))}{(bc-ad+id)(1-i(a+bx))}\right)}{(a+bx)^2+1} d(a+bx)}{d} + \frac{\arctan(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(1-i(a+bx))(-ad+bc+id)}\right)}{d} - \\
& \frac{\arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{d} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{2d} \\
& \downarrow 2897 \\
& \frac{\arctan(a+bx) \log\left(\frac{2(d(a+bx)-ad+bc)}{(1-i(a+bx))(-ad+bc+id)}\right)}{d} - \frac{\arctan(a+bx) \log\left(\frac{2}{1-i(a+bx)}\right)}{d} - \\
& \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2(bc-ad+d(a+bx))}{(bc-ad+id)(1-i(a+bx))}\right)}{2d} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bx)}\right)}{2d}
\end{aligned}$$

input `Int[ArcTan[a + b*x]/(c + d*x), x]`

output `-((ArcTan[a + b*x]*Log[2/(1 - I*(a + b*x))])/d) + (ArcTan[a + b*x]*Log[(2*(b*c - a*d + d*(a + b*x))]/((b*c + I*d - a*d)*(1 - I*(a + b*x))])/d + ((I/2)*PolyLog[2, 1 - 2/(1 - I*(a + b*x))])/d - ((I/2)*PolyLog[2, 1 - (2*(b*c - a*d + d*(a + b*x))]/((b*c + I*d - a*d)*(1 - I*(a + b*x))])/d`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2752 $\text{Int}[\text{Log}[(c_*)(x_)]/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}\{c, d, e\}, x] \ \&\& \ \text{EqQ}[e + c*d, 0]$
- rule 2849 $\text{Int}[\text{Log}[(c_)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[-e/g \text{ Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}\{c, d, e, f, g\}, x] \ \&\& \ \text{EqQ}[c, 2*d] \ \&\& \ \text{EqQ}[e^2*f + d^2*g, 0]$
- rule 2897 $\text{Int}[\text{Log}[u_]*(Pq_)^{(m_.)}, x_Symbol] \rightarrow \text{With}\{C = \text{FullSimplify}[Pq^{m*}((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{RationalFunctionQ}[u, x] \ \&\& \ \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]$
- rule 5381 $\text{Int}[(a_.) + \text{ArcTan}[(c_*)(x_)]*(b_.)/((d_) + (e_*)(x_)), x_Symbol] \rightarrow \text{Simp}[(-a + b*\text{ArcTan}[c*x])*(\text{Log}[2/(1 - I*c*x)]/e), x] + (\text{Simp}[(a + b*\text{ArcTan}[c*x])*(\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))])/e), x] + \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - \text{Simp}[b*(c/e) \text{ Int}[\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x))]/(1 + c^2*x^2), x], x]) /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[c^2*d^2 + e^2, 0]$
- rule 5570 $\text{Int}[(a_.) + \text{ArcTan}[(c_) + (d_*)(x_)]*(b_.)^{(p_.)*((e_.) + (f_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^m*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ \text{IgtQ}[p, 0]$

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{b \ln(ad-bc-d(bx+a)) \arctan(bx+a)}{d} + b \left(-\frac{i \ln(ad-bc-d(bx+a)) \left(\ln\left(\frac{id+d(bx+a)}{ad-bc+id}\right) - \ln\left(\frac{id-d(bx+a)}{-ad+bc+id}\right) \right)}{2d} - \frac{i \operatorname{dilog}\left(\frac{id+d(bx+a)}{ad-bc+id}\right)}{b} \right)$
default	$\frac{b \ln(ad-bc-d(bx+a)) \arctan(bx+a)}{d} + b \left(-\frac{i \ln(ad-bc-d(bx+a)) \left(\ln\left(\frac{id+d(bx+a)}{ad-bc+id}\right) - \ln\left(\frac{id-d(bx+a)}{-ad+bc+id}\right) \right)}{2d} - \frac{i \operatorname{dilog}\left(\frac{id+d(bx+a)}{ad-bc+id}\right)}{b} \right)$
parts	$\frac{\ln(dx+c) \arctan(bx+a)}{d} - b \left(-\frac{i \ln(dx+c) \left(\ln\left(\frac{id-ad+bc-b(dx+c)}{-ad+bc+id}\right) - \ln\left(\frac{id+ad-bc+b(dx+c)}{ad-bc+id}\right) \right)}{2bd} - \frac{i \operatorname{dilog}\left(\frac{id-ad+bc-b(dx+c)}{-ad+bc+id}\right)}{2bd} \right)$
risch	$\frac{i \operatorname{dilog}\left(\frac{id-ibc+(-bxi-ai+1)d-d}{iad-ibc-d}\right)}{2d} + \frac{i \ln(-bxi-ai+1) \ln\left(\frac{id-ibc+(-bxi-ai+1)d-d}{iad-ibc-d}\right)}{2d} - \frac{i \operatorname{dilog}\left(\frac{-iad+ibc+(bxi+ai+1)d-d}{-iad+ibc-d}\right)}{2d}$

input `int(arctan(b*x+a)/(d*x+c),x,method=_RETURNVERBOSE)`

output `1/b*(b*ln(a*d-b*c-d*(b*x+a))/d*arctan(b*x+a)+b*(-1/2*I*ln(a*d-b*c-d*(b*x+a))*ln((I*d+d*(b*x+a))/(a*d-b*c+I*d))-ln((I*d-d*(b*x+a))/(I*d-a*d+b*c)))/d-1/2*I*(dilog((I*d+d*(b*x+a))/(a*d-b*c+I*d))-dilog((I*d-d*(b*x+a))/(I*d-a*d+b*c)))/d)`

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \int \frac{\arctan(bx + a)}{dx + c} dx$$

input `integrate(arctan(b*x+a)/(d*x+c),x, algorithm="fricas")`

output `integral(arctan(b*x + a)/(d*x + c), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \text{Timed out}$$

input `integrate(atan(b*x+a)/(d*x+c),x)`output `Timed out`**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 284 vs. $2(128) = 256$.

Time = 0.27 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.87

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \frac{\arctan(bx + a) \log(dx + c)}{d} - \frac{\arctan\left(\frac{b^2x+ab}{b}\right) \log(dx + c)}{d} - \frac{\arctan\left(\frac{bd^2x+bcd}{b^2c^2-2abcd+(a^2+1)d^2}, \frac{b^2c^2-abcd+(b^2cd-abd^2)x}{b^2c^2-2abcd+(a^2+1)d^2}\right) \log(b^2x^2 + 2abx + a^2 + 1) - \arctan(bx + a) \log\left(\frac{b^2x^2 + 2abx + a^2 + 1}{b^2}\right)}{2d}$$

input `integrate(arctan(b*x+a)/(d*x+c),x, algorithm="maxima")`

output `arctan(b*x + a)*log(d*x + c)/d - arctan((b^2*x + a*b)/b)*log(d*x + c)/d - 1/2*(arctan2((b*d^2*x + b*c*d)/(b^2*c^2 - 2*a*b*c*d + (a^2 + 1)*d^2), (b^2*c^2 - a*b*c*d + (b^2*c*d - a*b*d^2)*x)/(b^2*c^2 - 2*a*b*c*d + (a^2 + 1)*d^2))*log(b^2*x^2 + 2*a*b*x + a^2 + 1) - arctan(b*x + a)*log((b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)/(b^2*c^2 - 2*a*b*c*d + (a^2 + 1)*d^2)) + I*dilog((I*b*d*x + (I*a + 1)*d)/(-I*b*c + (I*a + 1)*d)) - I*dilog((I*b*d*x + (I*a - 1)*d)/(-I*b*c + (I*a - 1)*d))/d`

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \int \frac{\arctan(bx + a)}{dx + c} dx$$

input `integrate(arctan(b*x+a)/(d*x+c),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \int \frac{\operatorname{atan}(a + bx)}{c + dx} dx$$

input `int(atan(a + b*x)/(c + d*x),x)`

output `int(atan(a + b*x)/(c + d*x), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + dx} dx = \int \frac{\operatorname{atan}(bx + a)}{dx + c} dx$$

input `int(atan(b*x+a)/(d*x+c),x)`

output `int(atan(a + b*x)/(c + d*x),x)`

3.55 $\int \frac{\arctan(a+bx)}{c+\frac{d}{x}} dx$

Optimal result	478
Mathematica [B] (warning: unable to verify)	479
Rubi [A] (verified)	480
Maple [A] (verified)	481
Fricas [F]	482
Sympy [F(-1)]	482
Maxima [A] (verification not implemented)	483
Giac [F]	483
Mupad [F(-1)]	484
Reduce [F]	484

Optimal result

Integrand size = 16, antiderivative size = 244

$$\int \frac{\arctan(a+bx)}{c+\frac{d}{x}} dx = -\frac{(1+ia+ibx)\log(1+ia+ibx)}{2bc} - \frac{(1-ia-ibx)\log(-i(i+a+bx))}{2bc} - \frac{id\log(1-ia-ibx)\log\left(-\frac{b(d+cx)}{(i+a)c-bd}\right)}{2c^2} + \frac{id\log(1+ia+ibx)\log\left(\frac{b(d+cx)}{(i-a)c+bd}\right)}{2c^2} + \frac{id\operatorname{PolyLog}\left(2, \frac{c(i-a-bx)}{ic-ac+bd}\right)}{2c^2} - \frac{id\operatorname{PolyLog}\left(2, \frac{c(i+a+bx)}{(i+a)c-bd}\right)}{2c^2}$$

output

```
-1/2*(1+I*a+I*b*x)*ln(1+I*a+I*b*x)/b/c-1/2*(1-I*a-I*b*x)*ln(-I*(I+a+b*x))/
b/c-1/2*I*d*ln(1-I*a-I*b*x)*ln(-b*(c*x+d)/((I+a)*c-b*d))/c^2+1/2*I*d*ln(1+
I*a+I*b*x)*ln(b*(c*x+d)/((I-a)*c+b*d))/c^2+1/2*I*d*polylog(2,c*(I-a-b*x)/(
I*c-a*c+b*d))/c^2-1/2*I*d*polylog(2,c*(I+a+b*x)/((I+a)*c-b*d))/c^2
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 771 vs. $2(244) = 488$.

Time = 7.96 (sec) , antiderivative size = 771, normalized size of antiderivative = 3.16

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \text{Too large to display}$$

input `Integrate[ArcTan[a + b*x]/(c + d/x), x]`

output

```
(-2*a^2*c^2*ArcTan[a + b*x] + 2*a*b*c*d*ArcTan[a + b*x] + I*a*b*c*d*Pi*ArcTan[a + b*x] - I*b^2*d^2*Pi*ArcTan[a + b*x] - 2*a*b*c^2*x*ArcTan[a + b*x] + 2*b^2*c*d*x*ArcTan[a + b*x] + (2*I)*a*b*c*d*ArcTan[a - (b*d)/c]*ArcTan[a + b*x] - (2*I)*b^2*d^2*ArcTan[a - (b*d)/c]*ArcTan[a + b*x] - b*c*d*ArcTan[a + b*x]^2 + I*a*b*c*d*ArcTan[a + b*x]^2 - I*b^2*d^2*ArcTan[a + b*x]^2 + (b*c*d*Sqrt[1 + a^2 - (2*a*b*d)/c + (b^2*d^2)/c^2]*ArcTan[a + b*x]^2)/E^(I*ArcTan[a - (b*d)/c]) + a*b*c*d*Pi*Log[1 + E^((-2*I)*ArcTan[a + b*x])] - b^2*d^2*Pi*Log[1 + E^((-2*I)*ArcTan[a + b*x])] - 2*a*b*c*d*ArcTan[a + b*x]*Log[1 + E^((2*I)*ArcTan[a + b*x])] + 2*b^2*d^2*ArcTan[a + b*x]*Log[1 + E^((2*I)*ArcTan[a + b*x])] - 2*a*b*c*d*ArcTan[a - (b*d)/c]*Log[1 - E^((2*I)*(-ArcTan[a - (b*d)/c] + ArcTan[a + b*x]))] + 2*b^2*d^2*ArcTan[a - (b*d)/c]*Log[1 - E^((2*I)*(-ArcTan[a - (b*d)/c] + ArcTan[a + b*x]))] + 2*a*b*c*d*ArcTan[a + b*x]*Log[1 - E^((2*I)*(-ArcTan[a - (b*d)/c] + ArcTan[a + b*x]))] - 2*b^2*d^2*ArcTan[a + b*x]*Log[1 - E^((2*I)*(-ArcTan[a - (b*d)/c] + ArcTan[a + b*x]))] - 2*a*c^2*Log[1/Sqrt[1 + (a + b*x)^2]] + 2*b*c*d*Log[1/Sqrt[1 + (a + b*x)^2]] - a*b*c*d*Pi*Log[1/Sqrt[1 + (a + b*x)^2]] + b^2*d^2*Pi*Log[1/Sqrt[1 + (a + b*x)^2]] + 2*a*b*c*d*ArcTan[a - (b*d)/c]*Log[Sin[ArcTan[(-a*c) + b*d]/c] + ArcTan[a + b*x]]] - 2*b^2*d^2*ArcTan[a - (b*d)/c]*Log[Sin[ArcTan[(-a*c) + b*d]/c] + ArcTan[a + b*x]]] + I*b*d*(a*c - b*d)*PolyLog[2, -E^((2*I)*ArcTan[a + b*x])] + I*b*d*(-a*c + b*d)*PolyLog[2, E^...
```


Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx$$

$$\downarrow 5574$$

$$\frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{c + \frac{d}{x}} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{c + \frac{d}{x}} dx$$

$$\downarrow 2856$$

$$\frac{1}{2}i \int \left(\frac{\log(-ia - ibx + 1)}{c} - \frac{d \log(-ia - ibx + 1)}{c(d + cx)} \right) dx -$$

$$\frac{1}{2}i \int \left(\frac{\log(ia + ibx + 1)}{c} - \frac{d \log(ia + ibx + 1)}{c(d + cx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{1}{2}i \left(-\frac{d \operatorname{PolyLog}\left(2, \frac{c(a+bx+i)}{(a+i)c-bd}\right)}{c^2} - \frac{d \log(-ia - ibx + 1) \log\left(-\frac{b(cx+d)}{-bd+(a+i)c}\right)}{c^2} + \frac{i(-ia - ibx + 1) \log(-i(a + bx + i))}{bc} \right)$$

$$\frac{1}{2}i \left(-\frac{d \operatorname{PolyLog}\left(2, \frac{c(-a-bx+i)}{-ac+ic+bd}\right)}{c^2} - \frac{d \log(ia + ibx + 1) \log\left(\frac{b(cx+d)}{bd+(-a+i)c}\right)}{c^2} - \frac{i(ia + ibx + 1) \log(ia + ibx + 1)}{bc} - \frac{x}{c} \right)$$

input `Int[ArcTan[a + b*x]/(c + d/x),x]`

output

```
(-1/2*I)*(-(x/c) - (I*(1 + I*a + I*b*x)*Log[1 + I*a + I*b*x])/(b*c) - (d*Log[1 + I*a + I*b*x]*Log[(b*(d + c*x))/((I - a)*c + b*d)]/c^2 - (d*PolyLog[2, (c*(I - a - b*x))/(I*c - a*c + b*d)]/c^2) + (I/2)*(-(x/c) + (I*(1 - I*a - I*b*x)*Log[(-I)*(I + a + b*x)])/(b*c) - (d*Log[1 - I*a - I*b*x]*Log[-((b*(d + c*x))/((I + a)*c - b*d))]/c^2 - (d*PolyLog[2, (c*(I + a + b*x))/((I + a)*c - b*d)]/c^2)
```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2856 Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]*(b_.))^(p_.)*((f_) + (g_.
)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)
^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && I
GtQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))
```

```
rule 5574 Int[ArcTan[(a_) + (b_.)*(x_)]/((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{\frac{\arctan(bx+a)(bx+a)}{c} - \frac{\arctan(bx+a)d \ln(ac-bd-c(bx+a))}{c^2} + \frac{\ln(a^2c^2-2abcd+d^2b^2-2ac(ac-bd-c(bx+a))+2bd(ac-bd-c(bx+a)))}{2}}{c}$
default	$\frac{\frac{\arctan(bx+a)(bx+a)}{c} - \frac{\arctan(bx+a)d \ln(ac-bd-c(bx+a))}{c^2} + \frac{\ln(a^2c^2-2abcd+d^2b^2-2ac(ac-bd-c(bx+a))+2bd(ac-bd-c(bx+a)))}{2}}{c}$
parts	$\frac{\arctan(bx+a)x}{c} - \frac{\arctan(bx+a)d \ln(cx+d)}{c^2} - \frac{b \left(\frac{\ln(a^2c^2-2abcd+2abc(cx+d)+d^2b^2-2b^2d(cx+d)+b^2(cx+d)^2+c^2)}{2b^2} - \frac{a}{c} \right)}{c}$
risch	$\frac{i \ln(-bxi-ai+1)x}{2c} - \frac{id \operatorname{dilog}\left(\frac{iac-ibd+(-bxi-ai+1)c-c}{iac-ibd-c}\right)}{2c^2} - \frac{i \ln(bxi+ai+1)x}{2c} - \frac{i \ln(bxi+ai+1)a}{2bc} - \frac{\ln(-bxi-ai+1)}{2bc}$

```
input int(arctan(b*x+a)/(c+d/x), x, method=_RETURNVERBOSE)
```

output

```
1/b*(arctan(b*x+a)/c*(b*x+a)-arctan(b*x+a)*d*b/c^2*ln(a*c-b*d-c*(b*x+a))+1
/c*(-1/2*ln(a^2*c^2-2*a*b*c*d+d^2*b^2-2*a*c*(a*c-b*d-c*(b*x+a))+2*b*d*(a*c
-b*d-c*(b*x+a))+c^2+(a*c-b*d-c*(b*x+a))^2)-b*d*(-1/2*I*ln(a*c-b*d-c*(b*x+a
))*(ln((I*c+c*(b*x+a))/(a*c-b*d+I*c))-ln((I*c-c*(b*x+a))/(I*c-a*c+b*d)))/c
-1/2*I*(dilog((I*c+c*(b*x+a))/(a*c-b*d+I*c))-dilog((I*c-c*(b*x+a))/(I*c-a*
c+b*d)))/c))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x}} dx$$

input

```
integrate(arctan(b*x+a)/(c+d/x),x, algorithm="fricas")
```

output

```
integral(x*arctan(b*x + a)/(c*x + d), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \text{Timed out}$$

input

```
integrate(atan(b*x+a)/(c+d/x),x)
```

output

```
Timed out
```

Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.16

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx =$$

$$bd \arctan(bx + a) \log\left(-\frac{b^2c^2x^2 + 2b^2cdx + b^2d^2}{2abcd - b^2d^2 - (a^2 + 1)c^2}\right) + i bd \operatorname{Li}_2\left(-\frac{ibcx + (ia-1)c}{(-ia+1)c + ibd}\right) - i bd \operatorname{Li}_2\left(-\frac{ibcx + (ia+1)c}{(-ia-1)c + ibd}\right) - 2$$

input `integrate(arctan(b*x+a)/(c+d/x),x, algorithm="maxima")`

output

```
-1/2*(b*d*arctan(b*x + a)*log(-(b^2*c^2*x^2 + 2*b^2*c*d*x + b^2*d^2)/(2*a*
b*c*d - b^2*d^2 - (a^2 + 1)*c^2)) + I*b*d*dilog(-(I*b*c*x + (I*a - 1)*c)/
(-I*a + 1)*c + I*b*d)) - I*b*d*dilog(-(I*b*c*x + (I*a + 1)*c)/((-I*a - 1)*
c + I*b*d)) - 2*(b*c*x + a*c)*arctan(b*x + a) - (b*d*arctan2(-(b*c^2*x + b
*c*d)/(2*a*b*c*d - b^2*d^2 - (a^2 + 1)*c^2), (a*b*c*d - b^2*d^2 + (a*b*c^2
- b^2*c*d)*x)/(2*a*b*c*d - b^2*d^2 - (a^2 + 1)*c^2)) - c)*log(b^2*x^2 + 2
*a*b*x + a^2 + 1))/(b*c^2)
```

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x}} dx$$

input `integrate(arctan(b*x+a)/(c+d/x),x, algorithm="giac")`

output

`integrate(arctan(b*x + a)/(c + d/x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{atan}(a + bx)}{c + \frac{d}{x}} dx$$

input `int(atan(a + b*x)/(c + d/x),x)`output `int(atan(a + b*x)/(c + d/x), x)`**Reduce [F]**

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x}} dx = \int \frac{\operatorname{atan}(bx + a)x}{cx + d} dx$$

input `int(atan(b*x+a)/(c+d/x),x)`output `int((atan(a + b*x)*x)/(c*x + d),x)`

$$3.56 \quad \int \frac{\arctan(a+bx)}{c+\frac{d}{x^2}} dx$$

Optimal result	486
Mathematica [A] (verified)	487
Rubi [A] (verified)	488
Maple [A] (verified)	490
Fricas [F]	491
Sympy [F(-1)]	491
Maxima [B] (verification not implemented)	491
Giac [F]	492
Mupad [F(-1)]	493
Reduce [F]	493

Optimal result

Integrand size = 16, antiderivative size = 668

$$\begin{aligned}
 \int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = & -\frac{(1 + ia + ibx) \log(1 + ia + ibx)}{2bc} \\
 & -\frac{(1 - ia - ibx) \log(-i(i + a + bx))}{2bc} \\
 & + \frac{i\sqrt{d} \log(1 + ia + ibx) \log\left(-\frac{b(\sqrt{d} - \sqrt{-cx})}{i\sqrt{-c} - a\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & - \frac{i\sqrt{d} \log(1 - ia - ibx) \log\left(\frac{b(\sqrt{d} - \sqrt{-cx})}{i\sqrt{-c} + a\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & + \frac{i\sqrt{d} \log(1 - ia - ibx) \log\left(-\frac{b(\sqrt{d} + \sqrt{-cx})}{(i+a)\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & - \frac{i\sqrt{d} \log(1 + ia + ibx) \log\left(\frac{b(\sqrt{d} + \sqrt{-cx})}{i\sqrt{-c} - a\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & + \frac{i\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(i-a-bx)}{i\sqrt{-c} - a\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & - \frac{i\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(1+ia+ibx)}{(1+ia)\sqrt{-c} - ib\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & + \frac{i\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(i+a+bx)}{i\sqrt{-c} + a\sqrt{-c} - b\sqrt{d}}\right)}{4(-c)^{3/2}} \\
 & - \frac{i\sqrt{d} \operatorname{PolyLog}\left(2, \frac{\sqrt{-c}(i+a+bx)}{i\sqrt{-c} + a\sqrt{-c} + b\sqrt{d}}\right)}{4(-c)^{3/2}}
 \end{aligned}$$

output

```

-1/2*(1+I*a+I*b*x)*ln(1+I*a+I*b*x)/b/c-1/2*(1-I*a-I*b*x)*ln(-I*(I+a+b*x))/
b/c+1/4*I*d^(1/2)*ln(1+I*a+I*b*x)*ln(-b*(d^(1/2)-(-c)^(1/2)*x)/(I*(-c)^(1/2)
-2)*a*(-c)^(1/2)-b*d^(1/2)))/(-c)^(3/2)-1/4*I*d^(1/2)*ln(1-I*a-I*b*x)*ln(b*
(d^(1/2)-(-c)^(1/2)*x)/(I*(-c)^(1/2)+a*(-c)^(1/2)+b*d^(1/2)))/(-c)^(3/2)+1
/4*I*d^(1/2)*ln(1-I*a-I*b*x)*ln(-b*(d^(1/2)+(-c)^(1/2)*x)/((I+a)*(-c)^(1/2)
-b*d^(1/2)))/(-c)^(3/2)-1/4*I*d^(1/2)*ln(1+I*a+I*b*x)*ln(b*(d^(1/2)+(-c)^(
1/2)*x)/(I*(-c)^(1/2)-a*(-c)^(1/2)+b*d^(1/2)))/(-c)^(3/2)+1/4*I*d^(1/2)*p
olylog(2,(-c)^(1/2)*(I-a-b*x)/(I*(-c)^(1/2)-a*(-c)^(1/2)-b*d^(1/2)))/(-c)^(
3/2)-1/4*I*d^(1/2)*polylog(2,(-c)^(1/2)*(1+I*a+I*b*x)/((1+I*a)*(-c)^(1/2)
-I*b*d^(1/2)))/(-c)^(3/2)+1/4*I*d^(1/2)*polylog(2,(-c)^(1/2)*(I+a+b*x)/(I*
(-c)^(1/2)+a*(-c)^(1/2)-b*d^(1/2)))/(-c)^(3/2)-1/4*I*d^(1/2)*polylog(2,(-c)
^(1/2)*(I+a+b*x)/(I*(-c)^(1/2)+a*(-c)^(1/2)+b*d^(1/2)))/(-c)^(3/2)

```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 660, normalized size of antiderivative = 0.99

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx =$$

$$i \left(2i\sqrt{-c} \log(1 + ia + ibx) - 2a\sqrt{-c} \log(1 + ia + ibx) - 2b\sqrt{-c}x \log(1 + ia + ibx) + 2i\sqrt{-c} \log(-i) \right)$$

input

```
Integrate[ArcTan[a + b*x]/(c + d/x^2),x]
```


output

```

((-1/4*I)*((2*I)*Sqrt[-c]*Log[1 + I*a + I*b*x] - 2*a*Sqrt[-c]*Log[1 + I*a
+ I*b*x] - 2*b*Sqrt[-c]*x*Log[1 + I*a + I*b*x] + (2*I)*Sqrt[-c]*Log[(-I)*(
I + a + b*x)] + 2*a*Sqrt[-c]*Log[(-I)*(I + a + b*x)] + 2*b*Sqrt[-c]*x*Log[
(-I)*(I + a + b*x)] - b*Sqrt[d]*Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[d] - Sqr
t[-c]*x))/((-I)*Sqrt[-c] + a*Sqrt[-c] + b*Sqrt[d])] + b*Sqrt[d]*Log[(-I)*(
I + a + b*x)]*Log[(b*(Sqrt[d] - Sqrt[-c]*x))/(I*Sqrt[-c] + a*Sqrt[-c] + b*
Sqrt[d])] - b*Sqrt[d]*Log[(-I)*(I + a + b*x)]*Log[-((b*(Sqrt[d] + Sqrt[-c]
*x))/((I + a)*Sqrt[-c] - b*Sqrt[d]))] + b*Sqrt[d]*Log[1 + I*a + I*b*x]*Log
[(b*(Sqrt[d] + Sqrt[-c]*x))/(I*Sqrt[-c] - a*Sqrt[-c] + b*Sqrt[d])] + b*Sqr
t[d]*PolyLog[2, (Sqrt[-c]*(-I + a + b*x))/((-I)*Sqrt[-c] + a*Sqrt[-c] - b*
Sqrt[d])] - b*Sqrt[d]*PolyLog[2, (Sqrt[-c]*(-I + a + b*x))/((-I)*Sqrt[-c]
+ a*Sqrt[-c] + b*Sqrt[d])] - b*Sqrt[d]*PolyLog[2, (Sqrt[-c]*(I + a + b*x))
/(I*Sqrt[-c] + a*Sqrt[-c] - b*Sqrt[d])] + b*Sqrt[d]*PolyLog[2, (Sqrt[-c]*(
I + a + b*x))/(I*Sqrt[-c] + a*Sqrt[-c] + b*Sqrt[d])]))/(b*(-c)^(3/2))

```

Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 678, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx \\
 & \quad \downarrow \text{5574} \\
 & \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{c + \frac{d}{x^2}} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{c + \frac{d}{x^2}} dx \\
 & \quad \downarrow \text{2856} \\
 & \frac{1}{2}i \int \left(\frac{\log(-ia - ibx + 1)}{c} - \frac{d \log(-ia - ibx + 1)}{c(cx^2 + d)} \right) dx - \\
 & \quad \frac{1}{2}i \int \left(\frac{\log(ia + ibx + 1)}{c} - \frac{d \log(ia + ibx + 1)}{c(cx^2 + d)} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{1}{2^i} \left(\frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(a+bx+i)}{\sqrt{-ca+i}\sqrt{-c-b\sqrt{d}}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(a+bx+i)}{\sqrt{-ca+i}\sqrt{-c+b\sqrt{d}}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \log(-ia - ibx + 1) \log \left(\frac{b(\sqrt{d}}{a\sqrt{-c}} + \dots \right)}{2(-c)^{3/2}} \right. \\ \left. - \frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(-a-bx+i)}{-\sqrt{-ca+i}\sqrt{-c-b\sqrt{d}}} \right)}{2(-c)^{3/2}} + \frac{\sqrt{d} \operatorname{PolyLog} \left(2, \frac{\sqrt{-c}(ia+ibx+1)}{(ia+1)\sqrt{-c-ib\sqrt{d}}} \right)}{2(-c)^{3/2}} - \frac{\sqrt{d} \log(ia + ibx + 1) \log \left(-\frac{1}{a(-c)} \right)}{2(-c)^{3/2}} \right)$$

input `Int[ArcTan[a + b*x]/(c + d/x^2), x]`

output `(-1/2*I)*(-(x/c) - (I*(1 + I*a + I*b*x)*Log[1 + I*a + I*b*x])/(b*c) - (Sqrt[d]*Log[1 + I*a + I*b*x]*Log[-((b*(Sqrt[d] - Sqrt[-c]*x))/(I*Sqrt[-c] - a*Sqrt[-c] - b*Sqrt[d]))])/(2*(-c)^(3/2)) + (Sqrt[d]*Log[1 + I*a + I*b*x]*Log[(b*(Sqrt[d] + Sqrt[-c]*x))/(I*Sqrt[-c] - a*Sqrt[-c] + b*Sqrt[d]))/(2*(-c)^(3/2)) - (Sqrt[d]*PolyLog[2, (Sqrt[-c]*(I - a - b*x))/(I*Sqrt[-c] - a*Sqrt[-c] - b*Sqrt[d])])/(2*(-c)^(3/2)) + (Sqrt[d]*PolyLog[2, (Sqrt[-c]*(1 + I*a + I*b*x))/((1 + I*a)*Sqrt[-c] - I*b*Sqrt[d])])/(2*(-c)^(3/2)) + (I/2)*(-(x/c) + (I*(1 - I*a - I*b*x)*Log[(-I)*(I + a + b*x)])/(b*c) - (Sqrt[d]*Log[1 - I*a - I*b*x]*Log[(b*(Sqrt[d] - Sqrt[-c]*x))/(I*Sqrt[-c] + a*Sqrt[-c] + b*Sqrt[d])])/(2*(-c)^(3/2)) + (Sqrt[d]*Log[1 - I*a - I*b*x]*Log[-((b*(Sqrt[d] + Sqrt[-c]*x))/((I + a)*Sqrt[-c] - b*Sqrt[d]))])/(2*(-c)^(3/2)) + (Sqrt[d]*PolyLog[2, (Sqrt[-c]*(I + a + b*x))/(I*Sqrt[-c] + a*Sqrt[-c] - b*Sqrt[d])])/(2*(-c)^(3/2)) - (Sqrt[d]*PolyLog[2, (Sqrt[-c]*(I + a + b*x))/(I*Sqrt[-c] + a*Sqrt[-c] + b*Sqrt[d])])/(2*(-c)^(3/2)))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_.) + (e_.)*(x_.))^(n_.)]*(b_.))^(p_.)*((f_.) + (g_.)*(x_.)^(r_.))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && IntegerQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 5574

```
Int[ArcTan[(a_) + (b_)*(x_)]/((c_) + (d_)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 2.14 (sec) , antiderivative size = 647, normalized size of antiderivative = 0.97

method	result
risch	$-\frac{i \ln(bxi+ai+1)a}{2bc} + \frac{i \ln(-bxi-ai+1)a}{2bc} + \frac{i \ln(-bxi-ai+1)x}{2c} - \frac{i \ln(bxi+ai+1)x}{2c} - \frac{\ln(-bxi-ai+1)}{2bc} + \frac{1}{bc} -$
derivativdivides	Expression too large to display
default	Expression too large to display

input

```
int(arctan(b*x+a)/(c+d/x^2),x,method=_RETURNVERBOSE)
```

output

```
-1/2*I/b/c*ln(1+I*a+I*b*x)*a+1/2*I/b/c*ln(1-I*a-I*b*x)*a+1/2*I/c*ln(1-I*a-
I*b*x)*x-1/2*I/c*ln(1+I*a+I*b*x)*x-1/2/b/c*ln(1-I*a-I*b*x)+1/b/c-1/4/c^2*ln
n(1-I*a-I*b*x)*ln((I*a*c-b*(c*d)^(1/2)+(1-I*a-I*b*x)*c-c)/(I*a*c-b*(c*d)^(
1/2)-c))*(c*d)^(1/2)+1/4/c^2*ln(1-I*a-I*b*x)*ln((I*a*c+b*(c*d)^(1/2)+(1-I*
a-I*b*x)*c-c)/(I*a*c+b*(c*d)^(1/2)-c))*(c*d)^(1/2)-1/4/c^2*dilog((I*a*c-b*
(c*d)^(1/2)+(1-I*a-I*b*x)*c-c)/(I*a*c-b*(c*d)^(1/2)-c))*(c*d)^(1/2)+1/4/c^
2*dilog((I*a*c+b*(c*d)^(1/2)+(1-I*a-I*b*x)*c-c)/(I*a*c+b*(c*d)^(1/2)-c))*
(c*d)^(1/2)-1/2/b/c*ln(1+I*a+I*b*x)-1/4/c^2*ln(1+I*a+I*b*x)*(c*d)^(1/2)*ln(
(I*a*c+b*(c*d)^(1/2)-(1+I*a+I*b*x)*c+c)/(I*a*c+b*(c*d)^(1/2)+c))+1/4/c^2*ln
n(1+I*a+I*b*x)*(c*d)^(1/2)*ln((I*a*c-b*(c*d)^(1/2)-(1+I*a+I*b*x)*c+c)/(I*a
*c-b*(c*d)^(1/2)+c))-1/4/c^2*(c*d)^(1/2)*dilog((I*a*c+b*(c*d)^(1/2)-(1+I*a
+I*b*x)*c+c)/(I*a*c+b*(c*d)^(1/2)+c))+1/4/c^2*(c*d)^(1/2)*dilog((I*a*c-b*(
c*d)^(1/2)-(1+I*a+I*b*x)*c+c)/(I*a*c-b*(c*d)^(1/2)+c))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x^2}} dx$$

input `integrate(arctan(b*x+a)/(c+d/x^2),x, algorithm="fricas")`

output `integral(x^2*arctan(b*x + a)/(c*x^2 + d), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \text{Timed out}$$

input `integrate(atan(b*x+a)/(c+d/x**2),x)`

output `Timed out`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 8518 vs. $2(466) = 932$.

Time = 0.81 (sec) , antiderivative size = 8518, normalized size of antiderivative = 12.75

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \text{Too large to display}$$

input `integrate(arctan(b*x+a)/(c+d/x^2),x, algorithm="maxima")`

output

```

-(d*arctan(c*x/sqrt(c*d))/(sqrt(c*d)*c) - x/c)*arctan(b*x + a) + 1/8*(8*a*
c*arctan(b*x + a) + (4*b*arctan(sqrt(c)*x/sqrt(d))*arctan2((2*a*b^2*c*d +
(a*b^3*d + (a^3 + a)*b*c + (b^4*d + (a^2 + 3)*b^2*c)*x)*sqrt(c)*sqrt(d) +
(3*b^3*c*d + (a^2 + 1)*b*c^2)*x)/(b^4*d^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2
*a^2 + 1)*c^2 + 4*(b^3*d + (a^2 + 1)*b*c)*sqrt(c)*sqrt(d)), ((a^2 + 3)*b^2
*c*d + (a^4 + 2*a^2 + 1)*c^2 + (2*a*b^2*c*x + b^3*d + 3*(a^2 + 1)*b*c)*sqr
t(c)*sqrt(d) + (a*b^3*c*d + (a^3 + a)*b*c^2)*x)/(b^4*d^2 + 2*(a^2 + 3)*b^2
*c*d + (a^4 + 2*a^2 + 1)*c^2 + 4*(b^3*d + (a^2 + 1)*b*c)*sqrt(c)*sqrt(d))
+ 4*b*arctan(sqrt(c)*x/sqrt(d))*arctan2((2*a*b^2*c*d - (a*b^3*d + (a^3 +
a)*b*c + (b^4*d + (a^2 + 3)*b^2*c)*x)*sqrt(c)*sqrt(d) + (3*b^3*c*d + (a^2
+ 1)*b*c^2)*x)/(b^4*d^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^2 + 1)*c^2 - 4*
(b^3*d + (a^2 + 1)*b*c)*sqrt(c)*sqrt(d)), ((a^2 + 3)*b^2*c*d + (a^4 + 2*a^
2 + 1)*c^2 - (2*a*b^2*c*x + b^3*d + 3*(a^2 + 1)*b*c)*sqrt(c)*sqrt(d) + (a*
b^3*c*d + (a^3 + a)*b*c^2)*x)/(b^4*d^2 + 2*(a^2 + 3)*b^2*c*d + (a^4 + 2*a^
2 + 1)*c^2 - 4*(b^3*d + (a^2 + 1)*b*c)*sqrt(c)*sqrt(d)) + b*log(c*x^2 + d
)*log(((a^2 + 1)*b^22*c*d^11 + 11*(a^4 + 22*a^2 + 21)*b^20*c^2*d^10 + 55*(
a^6 + 39*a^4 + 171*a^2 + 133)*b^18*c^3*d^9 + 33*(5*a^8 + 260*a^6 + 1870*a^
4 + 3876*a^2 + 2261)*b^16*c^4*d^8 + 330*(a^10 + 61*a^8 + 570*a^6 + 1802*a^
4 + 2261*a^2 + 969)*b^14*c^5*d^7 + 22*(21*a^12 + 1386*a^10 + 15015*a^8 + 6
0060*a^6 + 109395*a^4 + 92378*a^2 + 29393)*b^12*c^6*d^6 + 22*(21*a^14 + ...

```

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x^2}} dx$$

input

```
integrate(arctan(b*x+a)/(c+d/x^2),x, algorithm="giac")
```

output

```
integrate(arctan(b*x + a)/(c + d/x^2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{c + \frac{d}{x^2}} dx$$

input `int(atan(a + b*x)/(c + d/x^2), x)`output `int(atan(a + b*x)/(c + d/x^2), x)`**Reduce [F]**

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^2}} dx = \int \frac{\operatorname{atan}(bx + a) x^2}{c x^2 + d} dx$$

input `int(atan(b*x+a)/(c+d/x^2), x)`output `int((atan(a + b*x)*x**2)/(c*x**2 + d), x)`

$$3.57 \quad \int \frac{\arctan(a+bx)}{c+\frac{d}{x^3}} dx$$

Optimal result	494
Mathematica [A] (verified)	495
Rubi [A] (verified)	496
Maple [C] (warning: unable to verify)	499
Fricas [F]	500
Sympy [F(-1)]	500
Maxima [F]	501
Giac [F]	501
Mupad [F(-1)]	501
Reduce [F]	502

Optimal result

Integrand size = 16, antiderivative size = 933

$$\int \frac{\arctan(a+bx)}{c+\frac{d}{x^3}} dx = \text{Too large to display}$$

output

```

-1/2*(1+I*a+I*b*x)*ln(1+I*a+I*b*x)/b/c-1/2*(1-I*a-I*b*x)*ln(-I*(I+a+b*x))/
b/c-1/6*I*d^(1/3)*ln(1-I*a-I*b*x)*ln(-b*(d^(1/3)+c^(1/3)*x)/((I+a)*c^(1/3)
-b*d^(1/3)))/c^(4/3)+1/6*I*d^(1/3)*ln(1+I*a+I*b*x)*ln(b*(d^(1/3)+c^(1/3)*x)
)/((I-a)*c^(1/3)+b*d^(1/3)))/c^(4/3)-1/6*(-1)^(1/6)*d^(1/3)*ln(1+I*a+I*b*x
)*ln(-b*(d^(1/3)-(-1)^(1/3)*c^(1/3)*x)/((-1)^(1/3)*(I-a)*c^(1/3)-b*d^(1/3)
))/c^(4/3)+1/6*(-1)^(1/6)*d^(1/3)*ln(1-I*a-I*b*x)*ln(b*(d^(1/3)-(-1)^(1/3)
)*c^(1/3)*x)/((-1)^(1/3)*(I+a)*c^(1/3)+b*d^(1/3)))/c^(4/3)-1/6*(-1)^(5/6)*d
^(1/3)*ln(1+I*a+I*b*x)*ln(b*(d^(1/3)+(-1)^(2/3)*c^(1/3)*x)/((-1)^(2/3)*(I-
a)*c^(1/3)+b*d^(1/3)))/c^(4/3)+1/6*(-1)^(5/6)*d^(1/3)*ln(1-I*a-I*b*x)*ln(b
*(d^(1/3)+(-1)^(2/3)*c^(1/3)*x)/((-1)^(1/6)*(1-I*a)*c^(1/3)+b*d^(1/3)))/c^
(4/3)-1/6*(-1)^(1/6)*d^(1/3)*polylog(2,(-1)^(1/3)*c^(1/3)*(I-a-b*x)/((-1)^(
1/3)*(I-a)*c^(1/3)-b*d^(1/3)))/c^(4/3)-1/6*(-1)^(5/6)*d^(1/3)*polylog(2,(
-1)^(1/6)*c^(1/3)*(I-a-b*x)/((-1)^(1/6)*(I-a)*c^(1/3)-I*b*d^(1/3)))/c^(4/3
)+1/6*I*d^(1/3)*polylog(2,c^(1/3)*(I-a-b*x)/((I-a)*c^(1/3)+b*d^(1/3)))/c^(
4/3)-1/6*I*d^(1/3)*polylog(2,c^(1/3)*(I+a+b*x)/((I+a)*c^(1/3)-b*d^(1/3)))/
c^(4/3)+1/6*(-1)^(5/6)*d^(1/3)*polylog(2,(-1)^(2/3)*c^(1/3)*(I+a+b*x)/((-1)
^(2/3)*(I+a)*c^(1/3)-b*d^(1/3)))/c^(4/3)+1/6*(-1)^(1/6)*d^(1/3)*polylog(2
,(-1)^(1/3)*c^(1/3)*(I+a+b*x)/((-1)^(1/3)*(I+a)*c^(1/3)+b*d^(1/3)))/c^(4/3
)

```

Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 896, normalized size of antiderivative = 0.96

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \text{Too large to display}$$

input

```
Integrate[ArcTan[a + b*x]/(c + d/x^3),x]
```


output

```

((I/6)*((3*I)*c^(1/3)*Log[1 + I*a + I*b*x] - 3*a*c^(1/3)*Log[1 + I*a + I*b
*x] - 3*b*c^(1/3)*x*Log[1 + I*a + I*b*x] + (3*I)*c^(1/3)*Log[(-I)*(I + a +
b*x)] + 3*a*c^(1/3)*Log[(-I)*(I + a + b*x)] + 3*b*c^(1/3)*x*Log[(-I)*(I +
a + b*x)] + b*d^(1/3)*Log[1 + I*a + I*b*x]*Log[(b*(d^(1/3) + c^(1/3)*x))/
(-((-I + a)*c^(1/3)) + b*d^(1/3))] - b*d^(1/3)*Log[(-I)*(I + a + b*x)]*Log
[(b*(d^(1/3) + c^(1/3)*x))/(-((I + a)*c^(1/3)) + b*d^(1/3))] + (-1)^(2/3)*
b*d^(1/3)*Log[1 + I*a + I*b*x]*Log[(b*(d^(1/3) - (-1)^(1/3)*c^(1/3)*x))/((-
1)^(1/3)*(-I + a)*c^(1/3) + b*d^(1/3))] - (-1)^(2/3)*b*d^(1/3)*Log[(-I)*(
I + a + b*x)]*Log[(b*(d^(1/3) - (-1)^(1/3)*c^(1/3)*x))/((-1)^(1/3)*(I + a
)*c^(1/3) + b*d^(1/3))] + (-1)^(1/3)*b*d^(1/3)*Log[(-I)*(I + a + b*x)]*Log[
(b*(d^(1/3) + (-1)^(2/3)*c^(1/3)*x))/((-1)^(1/6)*(1 - I*a)*c^(1/3) + b*d^(
1/3))] - (-1)^(1/3)*b*d^(1/3)*Log[1 + I*a + I*b*x]*Log[(b*(d^(1/3) + (-1)^(
2/3)*c^(1/3)*x))/(-((-1)^(2/3)*(-I + a)*c^(1/3)) + b*d^(1/3))] + b*d^(1/3
)*PolyLog[2, (c^(1/3)*(-I + a + b*x))/((-I + a)*c^(1/3) - b*d^(1/3))] - (-
1)^(1/3)*b*d^(1/3)*PolyLog[2, ((-1)^(1/6)*c^(1/3)*(-I + a + b*x))/((-1)^(1
/6)*(-I + a)*c^(1/3) + I*b*d^(1/3))] + (-1)^(2/3)*b*d^(1/3)*PolyLog[2, ((-
1)^(1/3)*c^(1/3)*(-I + a + b*x))/((-1)^(1/3)*(-I + a)*c^(1/3) + b*d^(1/3))
] - b*d^(1/3)*PolyLog[2, (c^(1/3)*(I + a + b*x))/((I + a)*c^(1/3) - b*d^(1
/3))] + (-1)^(1/3)*b*d^(1/3)*PolyLog[2, ((-1)^(2/3)*c^(1/3)*(I + a + b*x))
/((-1)^(2/3)*(I + a)*c^(1/3) - b*d^(1/3))] - (-1)^(2/3)*b*d^(1/3)*PolyL...

```

Rubi [A] (verified)

Time = 1.79 (sec) , antiderivative size = 951, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx \\
 & \quad \downarrow \text{5574} \\
 & \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{c + \frac{d}{x^3}} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{c + \frac{d}{x^3}} dx \\
 & \quad \downarrow \text{2856}
 \end{aligned}$$

$$\frac{1}{2}i \int \left(\frac{\log(-ia - ibx + 1)}{c} - \frac{d \log(-ia - ibx + 1)}{c(cx^3 + d)} \right) dx - \frac{1}{2}i \int \left(\frac{\log(ia + ibx + 1)}{c} - \frac{d \log(ia + ibx + 1)}{c(cx^3 + d)} \right) dx$$

↓ 2009

$$\frac{1}{2}i \left(-\frac{x}{c} + \frac{i(-ia - ibx + 1) \log(-i(a + bx + i))}{bc} - \frac{\sqrt[3]{d} \log(-ia - ibx + 1) \log \left(-\frac{b(\sqrt[3]{cx} + \sqrt[3]{d})}{(a+i)\sqrt[3]{c} - b\sqrt[3]{d}} \right)}{3c^{4/3}} - \frac{(-1)^{2/3} \sqrt[3]{d}}{3c^{4/3}} \right) - \frac{1}{2}i \left(-\frac{x}{c} - \frac{i(ia + ibx + 1) \log(ia + ibx + 1)}{bc} - \frac{\sqrt[3]{d} \log(ia + ibx + 1) \log \left(\frac{b(\sqrt[3]{cx} + \sqrt[3]{d})}{\sqrt[3]{c}(i-a) + b\sqrt[3]{d}} \right)}{3c^{4/3}} - \frac{(-1)^{2/3} \sqrt[3]{d} \log(ia + ibx + 1)}{3c^{4/3}} \right)$$

input Int[ArcTan[a + b*x]/(c + d/x^3),x]

output

$$\begin{aligned}
& (-1/2*I)*(-(x/c) - (I*(1 + I*a + I*b*x)*\text{Log}[1 + I*a + I*b*x])/(b*c) - (d^{1/3}*\text{Log}[1 + I*a + I*b*x]*\text{Log}[(b*(d^{1/3} + c^{1/3}*x))/((I - a)*c^{1/3} + b*d^{1/3})])/(3*c^{4/3}) - ((-1)^{2/3}*d^{1/3}*\text{Log}[1 + I*a + I*b*x]*\text{Log}[-((b*(d^{1/3} - (-1)^{1/3}*c^{1/3}*x))/((-1)^{1/3}*(I - a)*c^{1/3} - b*d^{1/3}))])/(3*c^{4/3}) + ((-1)^{1/3}*d^{1/3}*\text{Log}[1 + I*a + I*b*x]*\text{Log}[(b*(d^{1/3} + (-1)^{2/3}*c^{1/3}*x))/((-1)^{2/3}*(I - a)*c^{1/3} + b*d^{1/3})])/(3*c^{4/3}) - ((-1)^{2/3}*d^{1/3}*\text{PolyLog}[2, ((-1)^{1/3}*c^{1/3}*(I - a - b*x))/((-1)^{1/3}*(I - a)*c^{1/3} - b*d^{1/3})])/(3*c^{4/3}) + ((-1)^{1/3}*d^{1/3}*\text{PolyLog}[2, ((-1)^{1/6}*c^{1/3}*(I - a - b*x))/((-1)^{1/6}*(I - a)*c^{1/3} - I*b*d^{1/3})])/(3*c^{4/3}) - (d^{1/3}*\text{PolyLog}[2, (c^{1/3}*(I - a - b*x))/((I - a)*c^{1/3} + b*d^{1/3})])/(3*c^{4/3})) + (I/2)*(-(x/c) + (I*(1 - I*a - I*b*x)*\text{Log}[(-I)*(I + a + b*x)]/(b*c) - (d^{1/3}*\text{Log}[1 - I*a - I*b*x]*\text{Log}[-(b*(d^{1/3} + c^{1/3}*x))/((I + a)*c^{1/3} - b*d^{1/3})])/(3*c^{4/3}) - ((-1)^{2/3}*d^{1/3}*\text{Log}[1 - I*a - I*b*x]*\text{Log}[(b*(d^{1/3} - (-1)^{1/3}*c^{1/3}*x))/((-1)^{1/3}*(I + a)*c^{1/3} + b*d^{1/3})])/(3*c^{4/3})) + ((-1)^{1/3}*d^{1/3}*\text{Log}[1 - I*a - I*b*x]*\text{Log}[(b*(d^{1/3} + (-1)^{2/3}*c^{1/3}*x))/((-1)^{1/6}*(1 - I*a)*c^{1/3} + b*d^{1/3})])/(3*c^{4/3}) - (d^{1/3}*\text{PolyLog}[2, (c^{1/3}*(I + a + b*x))/((I + a)*c^{1/3} - b*d^{1/3})])/(3*c^{4/3}) + ((-1)^{1/3}*d^{1/3}*\text{PolyLog}[2, ((-1)^{2/3}*c^{1/3}*(I + a + b*x))/((-1)^{2/3}*(I + a)*c^{1/3} - b*d^{1/3})])/(3*c^{4/3}) - ((-1)^{2/3}*\dots
\end{aligned}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2856 $\text{Int}[(a_.) + \text{Log}[(c_.)*((d_.) + (e_.)*(x_.))^{(n_.)}*(b_.)]^{(p_.)*((f_.) + (g_.)*(x_.)^{(r_.))^{(q_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*\text{Log}[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, n, r\}, x] \&\& I \text{GtQ}[p, 0] \&\& \text{IntegerQ}[q] \&\& (\text{GtQ}[q, 0] \|\| (\text{IntegerQ}[r] \&\& \text{NeQ}[r, 1]))$

rule 5574 $\text{Int}[\text{ArcTan}[(a_) + (b_.)*(x_.)]/((c_) + (d_.)*(x_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[I/2 \text{Int}[\text{Log}[1 - I*a - I*b*x]/(c + d*x^n), x], x] - \text{Simp}[I/2 \text{Int}[\text{Log}[1 + I*a + I*b*x]/(c + d*x^n), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{RationalQ}[n]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.09 (sec) , antiderivative size = 511, normalized size of antiderivative = 0.55

method	result
risch	$\frac{i \ln(-bxi-ai+1)x}{2c} + \frac{i \ln(-bxi-ai+1)a}{2bc} - \frac{\ln(-bxi-ai+1)}{2bc} + \frac{1}{bc} + \frac{ib^2 d \left(\sqrt{-R1=RootOf(c_Z^3+(3\text{RootOf}(-Z^2+...))} \right)}{...}$
derivativedivides	$\frac{\arctan(bx+a)(bx+a)}{c} + \frac{\arctan(bx+a) \left(\sqrt{-R=RootOf(c_Z^3-3ac_Z^2+3a^2c_Z-a^3c+b^3d)} - \frac{\ln(bx-_R+a)}{_R^2+2_R a-a^2} \right) d b^3}{3c^2}$
default	$\frac{\arctan(bx+a)(bx+a)}{c} + \frac{\arctan(bx+a) \left(\sqrt{-R=RootOf(c_Z^3-3ac_Z^2+3a^2c_Z-a^3c+b^3d)} - \frac{\ln(bx-_R+a)}{_R^2+2_R a-a^2} \right) d b^3}{3c^2}$

input `int(arctan(b*x+a)/(c+d/x^3),x,method=_RETURNVERBOSE)`

output

```

1/2*I/c*ln(1-I*a-I*b*x)*x+1/2*I/b/c*ln(1-I*a-I*b*x)*a-1/2/b/c*ln(1-I*a-I*b
*x)+1/b/c+1/6*I*b^2*d/c^2*sum(1/(1+2*I*a*_R1-2*I*a+_R1^2-a^2-2*_R1)*(ln(1-
I*a-I*b*x)*ln((_R1+I*b*x+I*a-1)/_R1)+dilog((_R1+I*b*x+I*a-1)/_R1)),_R1=Ro
otOf(c*_Z^3+(3*RootOf(_Z^2+1,index=1)*a*c-3*c)*_Z^2+(-6*RootOf(_Z^2+1,index
=1)*a*c-3*a^2*c+3*c)*_Z-RootOf(_Z^2+1,index=1)*a^3*c+RootOf(_Z^2+1,index=1
)*b^3*d+3*RootOf(_Z^2+1,index=1)*a*c+3*a^2*c-c))-1/2*I/c*ln(1+I*a+I*b*x)*x
-1/2*I/b/c*ln(1+I*a+I*b*x)*a-1/2/b/c*ln(1+I*a+I*b*x)-1/6*I*b^2*d/c^2*sum(1
/(1-2*I*a*_R1+2*I*a+_R1^2-a^2-2*_R1)*(ln(1+I*a+I*b*x)*ln((_R1-I*b*x-I*a-1)
/_R1)+dilog((_R1-I*b*x-I*a-1)/_R1)),_R1=RootOf(c*_Z^3+(-3*RootOf(_Z^2+1,in
dex=1)*a*c-3*c)*_Z^2+(6*RootOf(_Z^2+1,index=1)*a*c-3*a^2*c+3*c)*_Z+RootOf(
_Z^2+1,index=1)*a^3*c-RootOf(_Z^2+1,index=1)*b^3*d-3*RootOf(_Z^2+1,index=1
)*a*c+3*a^2*c-c))

```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x^3}} dx$$

input

```
integrate(arctan(b*x+a)/(c+d/x^3),x, algorithm="fricas")
```

output

```
integral(x^3*arctan(b*x + a)/(c*x^3 + d), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \text{Timed out}$$

input

```
integrate(atan(b*x+a)/(c+d/x**3),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x^3}} dx$$

input `integrate(arctan(b*x+a)/(c+d/x^3),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(c + d/x^3), x)`

Giac [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{x^3}} dx$$

input `integrate(arctan(b*x+a)/(c+d/x^3),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(c + d/x^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{atan}(a + bx)}{c + \frac{d}{x^3}} dx$$

input `int(atan(a + b*x)/(c + d/x^3),x)`

output `int(atan(a + b*x)/(c + d/x^3), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{x^3}} dx = \int \frac{\operatorname{atan}(bx + a) x^3}{c x^3 + d} dx$$

input `int(atan(b*x+a)/(c+d/x^3),x)`

output `int((atan(a + b*x)*x**3)/(c*x**3 + d),x)`

3.58 $\int \frac{\arctan(a+bx)}{1+x^2} dx$

Optimal result	503
Mathematica [A] (verified)	504
Rubi [A] (verified)	504
Maple [A] (verified)	506
Fricas [F]	507
Sympy [F]	507
Maxima [B] (verification not implemented)	507
Giac [F]	508
Mupad [F(-1)]	508
Reduce [F]	509

Optimal result

Integrand size = 14, antiderivative size = 179

$$\int \frac{\arctan(a+bx)}{1+x^2} dx = -\frac{1}{2}i \arctan(a+bx) \log\left(\frac{2b(i-x)}{(a-i(1-b))(1-i(a+bx))}\right) + \frac{1}{2}i \arctan(a+bx) \log\left(-\frac{2b(i+x)}{(a-i(1+b))(1-i(a+bx))}\right) - \frac{1}{4} \text{PolyLog}\left(2, 1 - \frac{2b(i-x)}{(a-i(1-b))(1-i(a+bx))}\right) + \frac{1}{4} \text{PolyLog}\left(2, 1 + \frac{2b(i+x)}{(a-i(1+b))(1-i(a+bx))}\right)$$

output

```
-1/2*I*arctan(b*x+a)*ln(2*b*(I-x)/(a-I*(1-b))/(1-I*(b*x+a)))+1/2*I*arctan(
b*x+a)*ln(-2*b*(I+x)/(a-I*(1+b))/(1-I*(b*x+a)))-1/4*polylog(2,1-2*b*(I-x)/
(a-I*(1-b))/(1-I*(b*x+a)))+1/4*polylog(2,1+2*b*(I+x)/(a-I*(1+b))/(1-I*(b*x
+a)))
```


Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.58

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \frac{1}{4} \log\left(\frac{b(i-x)}{a+i(1+b)}\right) \log(1-ia-ibx) - \frac{1}{4} \log\left(-\frac{b(i+x)}{a+i(1-b)}\right) \log(1-ia-ibx) - \frac{1}{4} \log\left(\frac{b(i-x)}{a-i(1-b)}\right) \log(1+ia+ibx) + \frac{1}{4} \log\left(-\frac{b(i+x)}{a-i(1+b)}\right) \log(1+ia+ibx) - \frac{1}{4} \text{PolyLog}\left(2, \frac{1-ia-ibx}{1-ia-b}\right) + \frac{1}{4} \text{PolyLog}\left(2, \frac{1-ia-ibx}{1-ia+b}\right) - \frac{1}{4} \text{PolyLog}\left(2, \frac{1+ia+ibx}{1+ia-b}\right) + \frac{1}{4} \text{PolyLog}\left(2, \frac{1+ia+ibx}{1+ia+b}\right)$$

input `Integrate[ArcTan[a + b*x]/(1 + x^2), x]`

output `(Log[(b*(I - x))/(a + I*(1 + b))]*Log[1 - I*a - I*b*x])/4 - (Log[-((b*(I + x))/(a + I*(1 - b)))]*Log[1 - I*a - I*b*x])/4 - (Log[(b*(I - x))/(a - I*(1 - b))]*Log[1 + I*a + I*b*x])/4 + (Log[-((b*(I + x))/(a - I*(1 + b)))]*Log[1 + I*a + I*b*x])/4 - PolyLog[2, (1 - I*a - I*b*x)/(1 - I*a - b)]/4 + PolyLog[2, (1 - I*a - I*b*x)/(1 - I*a + b)]/4 - PolyLog[2, (1 + I*a + I*b*x)/(1 + I*a - b)]/4 + PolyLog[2, (1 + I*a + I*b*x)/(1 + I*a + b)]/4`

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.70, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5574, 2856, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{x^2 + 1} dx$$

$$\begin{aligned}
& \downarrow 5574 \\
& \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{x^2 + 1} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{x^2 + 1} dx \\
& \downarrow 2856 \\
& \frac{1}{2}i \int \left(\frac{i \log(-ia - ibx + 1)}{2(i - x)} + \frac{i \log(-ia - ibx + 1)}{2(x + i)} \right) dx - \\
& \frac{1}{2}i \int \left(\frac{i \log(ia + ibx + 1)}{2(i - x)} + \frac{i \log(ia + ibx + 1)}{2(x + i)} \right) dx \\
& \downarrow 2009 \\
& \frac{1}{2}i \left(\frac{1}{2}i \operatorname{PolyLog} \left(2, \frac{a + bx + i}{a - ib + i} \right) - \frac{1}{2}i \operatorname{PolyLog} \left(2, \frac{a + bx + i}{a + i(b + 1)} \right) - \frac{1}{2}i \log \left(\frac{b(-x + i)}{a + i(b + 1)} \right) \log(-ia - ibx + 1) + \right. \\
& \left. \frac{1}{2}i \left(-\frac{1}{2}i \operatorname{PolyLog} \left(2, -\frac{-a - bx + i}{a - i(1 - b)} \right) + \frac{1}{2}i \operatorname{PolyLog} \left(2, -\frac{-a - bx + i}{a - i(b + 1)} \right) - \frac{1}{2}i \log \left(\frac{b(-x + i)}{a - i(1 - b)} \right) \log(ia + ibx + 1) \right)
\end{aligned}$$

input `Int[ArcTan[a + b*x]/(1 + x^2),x]`

output `(-1/2*I)*((-1/2*I)*Log[(b*(I - x))/(a - I*(1 - b))]*Log[1 + I*a + I*b*x] + (I/2)*Log[-((b*(I + x))/(a - I*(1 + b)))]*Log[1 + I*a + I*b*x] - (I/2)*PolyLog[2, -((I - a - b*x)/(a - I*(1 - b)))] + (I/2)*PolyLog[2, -((I - a - b*x)/(a - I*(1 + b)))] + (I/2)*((-1/2*I)*Log[(b*(I - x))/(a + I*(1 + b))]*Log[1 - I*a - I*b*x] + (I/2)*Log[-((b*(I + x))/(a + I*(1 - b)))]*Log[1 - I*a - I*b*x] + (I/2)*PolyLog[2, (I + a + b*x)/(I + a - I*b)] - (I/2)*PolyLog[2, (I + a + b*x)/(a + I*(1 + b))])`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2856 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*((f_) + (g_.)*(x_)^(r_))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, r}, x] && IntegerQ[p, 0] && IntegerQ[q] && (GtQ[q, 0] || (IntegerQ[r] && NeQ[r, 1]))`

rule 5574

```
Int[ArcTan[(a_) + (b_.)*(x_)]/((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [A] (verified)

Time = 0.85 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.26

method	result
risch	$\frac{\ln(-bxi-ai+1) \ln\left(\frac{-bxi-b}{ai-b-1}\right)}{4} + \frac{\operatorname{dilog}\left(\frac{-bxi-b}{ai-b-1}\right)}{4} - \frac{\ln(-bxi-ai+1) \ln\left(\frac{-bxi+b}{ai+b-1}\right)}{4} - \frac{\operatorname{dilog}\left(\frac{-bxi+b}{ai+b-1}\right)}{4} + \frac{\ln(bxi+ai)}{4}$
default	$\arctan(x) \arctan(bx+a) - b \left(\frac{\ln\left(1 - \frac{(-ib+a+i)(ix+1)^2}{(x^2+1)(-ib-a-i)}\right) \arctan(x)}{2(ib+a+i)} - \frac{\ln\left(1 - \frac{(-ib+a+i)(ix+1)^2}{(x^2+1)(-ib-a-i)}\right) \arctan(x)}{2b(ib+a+i)} \right)$
parts	$\arctan(x) \arctan(bx+a) - b \left(\frac{\ln\left(1 - \frac{(-ib+a+i)(ix+1)^2}{(x^2+1)(-ib-a-i)}\right) \arctan(x)}{2(ib+a+i)} - \frac{\ln\left(1 - \frac{(-ib+a+i)(ix+1)^2}{(x^2+1)(-ib-a-i)}\right) \arctan(x)}{2b(ib+a+i)} \right)$
derivativedivides	$b \arctan(x) \arctan(bx+a) - b^2 \left(\frac{\arctan\left(b\left(\frac{bx+a}{b} - \frac{a}{b}\right) + a\right) \arctan\left(-\frac{bx+a}{b} + \frac{a}{b}\right)}{b} - \frac{-\arctan\left(-\frac{bx+a}{b} + \frac{a}{b}\right) \arctan\left(b\left(\frac{bx+a}{b} - \frac{a}{b}\right) + a\right)}{b} \right)$

input

```
int(arctan(b*x+a)/(x^2+1), x, method=_RETURNVERBOSE)
```

output

```
1/4*ln(1-I*a-I*b*x)*ln((-I*b*x-b)/(I*a-b-1))+1/4*dilog((-I*b*x-b)/(I*a-b-1))
-1/4*ln(1-I*a-I*b*x)*ln((-I*b*x+b)/(I*a+b-1))-1/4*dilog((-I*b*x+b)/(I*a+b-1))
+1/4*ln(1+I*a+I*b*x)*ln((I*b*x-b)/(-I*a-b-1))+1/4*dilog((I*b*x-b)/(-I*a-b-1))
-1/4*ln(1+I*a+I*b*x)*ln((I*b*x+b)/(-I*a+b-1))-1/4*dilog((I*b*x+b)/(-I*a+b-1))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \int \frac{\arctan(bx + a)}{x^2 + 1} dx$$

input `integrate(arctan(b*x+a)/(x^2+1),x, algorithm="fricas")`

output `integral(arctan(b*x + a)/(x^2 + 1), x)`

Sympy [F]

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \int \frac{\operatorname{atan}(a + bx)}{x^2 + 1} dx$$

input `integrate(atan(b*x+a)/(x**2+1),x)`

output `Integral(atan(a + b*x)/(x**2 + 1), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 328 vs. $2(129) = 258$.

Time = 0.18 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.83

$$\begin{aligned} & \int \frac{\arctan(a + bx)}{1 + x^2} dx \\ &= \frac{1}{8} b \left(\frac{8 \arctan(x) \arctan\left(\frac{b^2x+ab}{b}\right)}{b} - \frac{4 \arctan(x) \arctan\left(\frac{ab+(b^2+b)x}{a^2+b^2+2b+1}, \frac{abx+a^2+b+1}{a^2+b^2+2b+1}\right)}{b} - 4 \arctan(x) \arctan\left(\frac{b^2x+ab}{b}\right) \right) \end{aligned}$$

input `integrate(arctan(b*x+a)/(x^2+1),x, algorithm="maxima")`

output `1/8*b*(8*arctan(x)*arctan((b^2*x + a*b)/b)/b - (4*arctan(x)*arctan2((a*b + (b^2 + b)*x)/(a^2 + b^2 + 2*b + 1), (a*b*x + a^2 + b + 1)/(a^2 + b^2 + 2*b + 1)) - 4*arctan(x)*arctan2((a*b + (b^2 - b)*x)/(a^2 + b^2 - 2*b + 1), (a*b*x + a^2 - b + 1)/(a^2 + b^2 - 2*b + 1)) + log(x^2 + 1)*log((b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^2 + b^2 + 2*b + 1)) - log(x^2 + 1)*log((b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^2 + b^2 - 2*b + 1)) + 2*dilog(-(I*b*x - b)/(I*a + b + 1)) - 2*dilog(-(I*b*x - b)/(I*a + b - 1)) + 2*dilog((I*b*x + b)/(-I*a + b + 1)) - 2*dilog((I*b*x + b)/(-I*a + b - 1)))/b + arctan(b*x + a)*arctan(x) - arctan(x)*arctan((b^2*x + a*b)/b)`

Giac [F]

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \int \frac{\arctan(bx + a)}{x^2 + 1} dx$$

input `integrate(arctan(b*x+a)/(x^2+1),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(x^2 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \int \frac{\operatorname{atan}(a + bx)}{x^2 + 1} dx$$

input `int(atan(a + b*x)/(x^2 + 1),x)`

output `int(atan(a + b*x)/(x^2 + 1), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{1 + x^2} dx = \int \frac{\operatorname{atan}(bx + a)}{x^2 + 1} dx$$

input `int(atan(b*x+a)/(x^2+1),x)`

output `int(atan(a + b*x)/(x**2 + 1),x)`

3.59
$$\int \frac{a+b \arctan(c+dx)}{e+f\sqrt{x}} dx$$

Optimal result	511
Mathematica [A] (verified)	512
Rubi [F]	513
Maple [C] (verified)	514
Fricas [F]	515
Sympy [F(-1)]	515
Maxima [F]	515
Giac [F]	516
Mupad [F(-1)]	516
Reduce [F]	516

Optimal result

Integrand size = 22, antiderivative size = 706

$$\begin{aligned}
\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = & \frac{2a\sqrt{x}}{f} + \frac{2ib\sqrt{i+c} \arctan\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{i+c}}\right)}{\sqrt{d}f} \\
& - \frac{2ib\sqrt{i-c} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{i-c}}\right)}{\sqrt{d}f} \\
& + \frac{ibe \log\left(\frac{f(\sqrt{-i-c}-\sqrt{d}\sqrt{x})}{\sqrt{de}+\sqrt{-i-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& - \frac{ibe \log\left(\frac{f(\sqrt{i-c}-\sqrt{d}\sqrt{x})}{\sqrt{de}+\sqrt{i-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& + \frac{ibe \log\left(-\frac{f(\sqrt{-i-c}+\sqrt{d}\sqrt{x})}{\sqrt{de}-\sqrt{-i-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& - \frac{ibe \log\left(-\frac{f(\sqrt{i-c}+\sqrt{d}\sqrt{x})}{\sqrt{de}-\sqrt{i-c}f}\right) \log(e + f\sqrt{x})}{f^2} \\
& + \frac{ib\sqrt{x} \log(1 - ic - idx)}{f} \\
& - \frac{e \log(e + f\sqrt{x}) (a + ib \log(1 - ic - idx))}{f^2} \\
& - \frac{ib\sqrt{x} \log(1 + ic + idx)}{f} \\
& - \frac{e \log(e + f\sqrt{x}) (a - ib \log(1 + ic + idx))}{f^2} \\
& + \frac{ibe \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}-\sqrt{-i-c}f}\right)}{f^2} \\
& + \frac{ibe \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}+\sqrt{-i-c}f}\right)}{f^2} \\
& - \frac{ibe \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}-\sqrt{i-c}f}\right)}{f^2} \\
& - \frac{ibe \operatorname{PolyLog}\left(2, \frac{\sqrt{d}(e+f\sqrt{x})}{\sqrt{de}+\sqrt{i-c}f}\right)}{f^2}
\end{aligned}$$

output

```

2*a*x^(1/2)/f+2*I*b*(I+c)^(1/2)*arctan(d^(1/2)*x^(1/2)/(I+c)^(1/2))/d^(1/2)
)/f-2*I*b*(I-c)^(1/2)*arctanh(d^(1/2)*x^(1/2)/(I-c)^(1/2))/d^(1/2)/f+I*b*e
*ln(f*((-I-c)^(1/2)-d^(1/2)*x^(1/2))/(d^(1/2)*e+(-I-c)^(1/2)*f))*ln(e+f*x^
(1/2))/f^2-I*b*e*ln(f*((I-c)^(1/2)-d^(1/2)*x^(1/2))/(d^(1/2)*e+(I-c)^(1/2)
*f))*ln(e+f*x^(1/2))/f^2+I*b*e*ln(-f*((-I-c)^(1/2)+d^(1/2)*x^(1/2))/(d^(1/
2)*e-(-I-c)^(1/2)*f))*ln(e+f*x^(1/2))/f^2-I*b*e*ln(-f*((I-c)^(1/2)+d^(1/2)
*x^(1/2))/(d^(1/2)*e-(I-c)^(1/2)*f))*ln(e+f*x^(1/2))/f^2+I*b*x^(1/2)*ln(1-
I*c-I*d*x)/f-e*ln(e+f*x^(1/2))*(a+I*b*ln(1-I*c-I*d*x))/f^2-I*b*x^(1/2)*ln(
1+I*c+I*d*x)/f-e*ln(e+f*x^(1/2))*(a-I*b*ln(1+I*c+I*d*x))/f^2+I*b*e*polylog
(2,d^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e-(-I-c)^(1/2)*f))/f^2+I*b*e*polylog(2,d
^(1/2)*(e+f*x^(1/2))/(d^(1/2)*e+(-I-c)^(1/2)*f))/f^2-I*b*e*polylog(2,d^(1/
2)*(e+f*x^(1/2))/(d^(1/2)*e-(I-c)^(1/2)*f))/f^2-I*b*e*polylog(2,d^(1/2)*(e
+f*x^(1/2))/(d^(1/2)*e+(I-c)^(1/2)*f))/f^2

```

Mathematica [A] (verified)

Time = 22.83 (sec) , antiderivative size = 631, normalized size of antiderivative = 0.89

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx$$

$$= \frac{2a(f\sqrt{x} - e \log(e + f\sqrt{x})) + ib \left(\frac{2\sqrt{i+c}f \arctan\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{i+c}}\right)}{\sqrt{d}} - \frac{2\sqrt{i-c}f \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{x}}{\sqrt{i-c}}\right)}{\sqrt{d}} + e \log\left(\frac{f(\sqrt{-i-c}-\sqrt{d}\sqrt{x})}{\sqrt{de+\sqrt{-i-c}f}}\right) \right)}{1}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])/(e + f*Sqrt[x]),x]
```

output

```
(2*a*(f*Sqrt[x] - e*Log[e + f*Sqrt[x]]) + I*b*((2*Sqrt[I + c]*f*ArcTan[(Sqrt[d]*Sqrt[x])/Sqrt[I + c]])/Sqrt[d] - (2*Sqrt[I - c]*f*ArcTanh[(Sqrt[d]*Sqrt[x])/Sqrt[I - c]])/Sqrt[d] + e*Log[(f*(Sqrt[-I - c] - Sqrt[d]*Sqrt[x]))/(Sqrt[d]*e + Sqrt[-I - c]*f)]*Log[e + f*Sqrt[x]] - e*Log[(f*(Sqrt[I - c] - Sqrt[d]*Sqrt[x]))/(Sqrt[d]*e + Sqrt[I - c]*f)]*Log[e + f*Sqrt[x]] + e*Log[(f*(Sqrt[-I - c] + Sqrt[d]*Sqrt[x]))/(-(Sqrt[d]*e) + Sqrt[-I - c]*f)]*Log[e + f*Sqrt[x]] - e*Log[(f*(Sqrt[I - c] + Sqrt[d]*Sqrt[x]))/(-(Sqrt[d]*e) + Sqrt[I - c]*f)]*Log[e + f*Sqrt[x]] - f*Sqrt[x]*Log[1 + I*c + I*d*x] + e*Log[e + f*Sqrt[x]]*Log[1 + I*c + I*d*x] + f*Sqrt[x]*Log[(-I)*(I + c + d*x)] - e*Log[e + f*Sqrt[x]]*Log[(-I)*(I + c + d*x)] + e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e - Sqrt[-I - c]*f)] + e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e + Sqrt[-I - c]*f)] - e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e - Sqrt[I - c]*f)] - e*PolyLog[2, (Sqrt[d]*(e + f*Sqrt[x]))/(Sqrt[d]*e + Sqrt[I - c]*f)))/f^2
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{\sqrt{x}(a + b \arctan(c + dx))}{e + f\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{7293} \\
 & 2 \int \left(\frac{\sqrt{x}a}{e + f\sqrt{x}} + \frac{b\sqrt{x} \arctan(c + dx)}{e + f\sqrt{x}} \right) d\sqrt{x} \\
 & \quad \downarrow \text{2009} \\
 & 2 \left(-\frac{be \int \frac{\arctan(c+dx)}{e+f\sqrt{x}} d\sqrt{x}}{f} - \frac{ae \log(e + f\sqrt{x})}{f^2} + \frac{a\sqrt{x}}{f} + \frac{b \arctan\left(\frac{\sqrt{\sqrt{c^2+1}-c-\sqrt{2}\sqrt{d}\sqrt{x}}}{\sqrt{\sqrt{c^2+1}+c}}\right)}{\sqrt{2}\sqrt{\sqrt{c^2+1}+c\sqrt{d}f}} - \frac{b \arctan\left(\frac{\sqrt{\sqrt{c^2+1}-c}}{\sqrt{\sqrt{c^2+1}+c}}\right)}{\sqrt{2}\sqrt{\sqrt{c^2+1}+c}} \right)
 \end{aligned}$$

input

```
Int[(a + b*ArcTan[c + d*x])/(e + f*Sqrt[x]), x]
```

output `$Aborted`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.27 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.53

method	result
derivativedivides	$\frac{2a\sqrt{x}}{f} - \frac{2ae \ln(e+f\sqrt{x})}{f^2} + \frac{2b \arctan(dx+c)\sqrt{x}}{f} - \frac{2b \arctan(dx+c)e \ln(e+f\sqrt{x})}{f^2} - b \left(\dots \right)$
default	$\frac{2a\sqrt{x}}{f} - \frac{2ae \ln(e+f\sqrt{x})}{f^2} + \frac{2b \arctan(dx+c)\sqrt{x}}{f} - \frac{2b \arctan(dx+c)e \ln(e+f\sqrt{x})}{f^2} - b \left(\dots \right)$
parts	$a \left(\frac{2\sqrt{x}}{f} + \frac{e \ln(f\sqrt{x}-e)}{f^2} - \frac{e \ln(e+f\sqrt{x})}{f^2} - \frac{e \ln(f^2x-e^2)}{f^2} \right) + \frac{2b \arctan(dx+c)\sqrt{x}}{f} - \frac{2b \arctan(dx+c)e \ln(e+f\sqrt{x})}{f^2}$

input `int((a+b*arctan(d*x+c))/(e+f*x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*a*x^(1/2)/f-2*a*e/f^2*ln(e+f*x^(1/2))+2*b*arctan(d*x+c)/f*x^(1/2)-2*b*arctan(d*x+c)*e/f^2*ln(e+f*x^(1/2))-b*sum((R^2-2*R*e+e^2)/(R^3*d-3*R^2*d*e+_R*c*f^2+3*_R*d*e^2-c*e*f^2-d*e^3)*ln(f*x^(1/2)-_R+e),_R=RootOf(d^2*_Z^4-4*d^2*e*_Z^3+(2*c*d*f^2+6*d^2*e^2)*_Z^2+(-4*c*d*e*f^2-4*d^2*e^3)*_Z+c^2*f^4+2*c*d*e^2*f^2+d^2*e^4+f^4))+b*e*sum(1/(_R1^2*d-2*_R1*d*e+c*f^2+d*e^2)*(ln(e+f*x^(1/2))*ln((-f*x^(1/2)+_R1-e)/_R1)+dilog((-f*x^(1/2)+_R1-e)/_R1)),_R1=RootOf(d^2*_Z^4-4*d^2*e*_Z^3+(2*c*d*f^2+6*d^2*e^2)*_Z^2+(-4*c*d*e*f^2-4*d^2*e^3)*_Z+c^2*f^4+2*c*d*e^2*f^2+d^2*e^4+f^4))`

Fricas [F]

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \arctan(dx + c) + a}{f\sqrt{x} + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(e+f*x^(1/2)),x, algorithm="fricas")`

output `integral(-(b*e*arctan(d*x + c) + a*e - (b*f*arctan(d*x + c) + a*f)*sqrt(x))/(f^2*x - e^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))/(e+f*x**(1/2)),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \arctan(dx + c) + a}{f\sqrt{x} + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(e+f*x^(1/2)),x, algorithm="maxima")`

output `2*(b*f^2*integrate(1/2*arctan(d*x + c)/(f*sqrt(x) + e), x) - a*e*log(f*sqrt(x) + e) + a*f*sqrt(x))/f^2`

Giac [F]

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \int \frac{b \arctan(dx + c) + a}{f\sqrt{x} + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(e+f*x^(1/2)),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)/(f*sqrt(x) + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \int \frac{a + b \operatorname{atan}(c + dx)}{e + f\sqrt{x}} dx$$

input `int((a + b*atan(c + d*x))/(e + f*x^(1/2)),x)`

output `int((a + b*atan(c + d*x))/(e + f*x^(1/2)), x)`

Reduce [F]

$$\int \frac{a + b \arctan(c + dx)}{e + f\sqrt{x}} dx = \frac{2\sqrt{x}af + \left(\int \frac{\operatorname{atan}(dx+c)}{-f^2x+e^2} dx\right) be f^2 - \left(\int \frac{\sqrt{x} \operatorname{atan}(dx+c)}{-f^2x+e^2} dx\right) b f^3 - 2 \log(\sqrt{x}f + e) ae}{f^2}$$

input `int((a+b*atan(d*x+c))/(e+f*x^(1/2)),x)`

output `(2*sqrt(x)*a*f + int(atan(c + d*x)/(e**2 - f**2*x),x)*b*e*f**2 - int((sqrt(x)*atan(c + d*x))/(e**2 - f**2*x),x)*b*f**3 - 2*log(sqrt(x)*f + e)*a*e)/f**2`

3.60
$$\int \frac{\arctan(a+bx)}{c+d\sqrt{x}} dx$$

Optimal result	518
Mathematica [A] (verified)	519
Rubi [A] (verified)	519
Maple [C] (verified)	522
Fricas [F]	523
Sympy [F(-1)]	523
Maxima [F]	523
Giac [F(-2)]	524
Mupad [F(-1)]	524
Reduce [F]	524

Optimal result

Integrand size = 18, antiderivative size = 673

$$\begin{aligned}
 \int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = & \frac{2i\sqrt{i+a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i+a}}\right)}{\sqrt{bd}} - \frac{2i\sqrt{i-a} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i-a}}\right)}{\sqrt{bd}} \\
 & + \frac{ic \log\left(\frac{d(\sqrt{-i-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{-i-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 & - \frac{ic \log\left(\frac{d(\sqrt{i-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc}+\sqrt{i-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 & + \frac{ic \log\left(-\frac{d(\sqrt{-i-a}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{-i-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 & - \frac{ic \log\left(-\frac{d(\sqrt{i-a}+\sqrt{b}\sqrt{x})}{\sqrt{bc}-\sqrt{i-ad}}\right) \log(c + d\sqrt{x})}{d^2} \\
 & + \frac{i\sqrt{x} \log(1 - ia - ibx)}{d} - \frac{ic \log(c + d\sqrt{x}) \log(1 - ia - ibx)}{d^2} \\
 & - \frac{i\sqrt{x} \log(1 + ia + ibx)}{d} + \frac{ic \log(c + d\sqrt{x}) \log(1 + ia + ibx)}{d^2} \\
 & + \frac{ic \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{-i-ad}}\right)}{d^2} + \frac{ic \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{-i-ad}}\right)}{d^2} \\
 & - \frac{ic \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{i-ad}}\right)}{d^2} - \frac{ic \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{i-ad}}\right)}{d^2}
 \end{aligned}$$

output

```

2*I*(I+a)^(1/2)*arctan(b^(1/2)*x^(1/2)/(I+a)^(1/2))/b^(1/2)/d-2*I*(I-a)^(1/2)*arctanh(b^(1/2)*x^(1/2)/(I-a)^(1/2))/b^(1/2)/d+I*c*ln(d*((-I-a)^(1/2)-b^(1/2)*x^(1/2))/(b^(1/2)*c+(-I-a)^(1/2)*d))*ln(c+d*x^(1/2))/d^2-I*c*ln(d*((I-a)^(1/2)-b^(1/2)*x^(1/2))/(b^(1/2)*c+(I-a)^(1/2)*d))*ln(c+d*x^(1/2))/d^2+I*c*ln(-d*((-I-a)^(1/2)+b^(1/2)*x^(1/2))/(b^(1/2)*c-(-I-a)^(1/2)*d))*ln(c+d*x^(1/2))/d^2-I*c*ln(-d*((I-a)^(1/2)+b^(1/2)*x^(1/2))/(b^(1/2)*c-(I-a)^(1/2)*d))*ln(c+d*x^(1/2))/d^2+I*x^(1/2)*ln(1-I*a-I*b*x)/d-I*c*ln(c+d*x^(1/2))*ln(1-I*a-I*b*x)/d^2-I*x^(1/2)*ln(1+I*a+I*b*x)/d+I*c*ln(c+d*x^(1/2))*ln(1+I*a+I*b*x)/d^2+I*c*polylog(2,b^(1/2)*(c+d*x^(1/2))/(b^(1/2)*c-(-I-a)^(1/2)*d))/d^2+I*c*polylog(2,b^(1/2)*(c+d*x^(1/2))/(b^(1/2)*c+(-I-a)^(1/2)*d))/d^2-I*c*polylog(2,b^(1/2)*(c+d*x^(1/2))/(b^(1/2)*c-(I-a)^(1/2)*d))/d^2-I*c*polylog(2,b^(1/2)*(c+d*x^(1/2))/(b^(1/2)*c+(I-a)^(1/2)*d))/d^2

```

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 604, normalized size of antiderivative = 0.90

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx$$

$$= i \left(\frac{2\sqrt{i+ad} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i+a}}\right)}{\sqrt{b}} - \frac{2\sqrt{i-ad} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i-a}}\right)}{\sqrt{b}} + c \log\left(\frac{d(\sqrt{-i-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc+\sqrt{-i-ad}}}\right) \log(c + d\sqrt{x}) - c \log\left(\frac{d(\sqrt{i-a}-\sqrt{b}\sqrt{x})}{\sqrt{bc+\sqrt{i-a}}}\right) \right)$$

input `Integrate[ArcTan[a + b*x]/(c + d*Sqrt[x]),x]`

output

```
(I*((2*Sqrt[I + a]*d*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[I + a]])/Sqrt[b] - (2*Sqrt[I - a]*d*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[I - a]])/Sqrt[b] + c*Log[(d*(Sqrt[-I - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c + Sqrt[-I - a]*d)]*Log[c + d*Sqrt[x]] - c*Log[(d*(Sqrt[I - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c + Sqrt[I - a]*d)]*Log[c + d*Sqrt[x]] + c*Log[(d*(Sqrt[-I - a] + Sqrt[b]*Sqrt[x]))/(-(Sqrt[b]*c) + Sqrt[-I - a]*d)]*Log[c + d*Sqrt[x]] - c*Log[(d*(Sqrt[I - a] + Sqrt[b]*Sqrt[x]))/(-(Sqrt[b]*c) + Sqrt[I - a]*d)]*Log[c + d*Sqrt[x]] - d*Sqrt[x]*Log[1 + I*a + I*b*x] + c*Log[c + d*Sqrt[x]]*Log[1 + I*a + I*b*x] + d*Sqrt[x]*Log[(-I)*(I + a + b*x)] - c*Log[c + d*Sqrt[x]]*Log[(-I)*(I + a + b*x)] + c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[-I - a]*d)] + c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqrt[-I - a]*d)] - c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[I - a]*d)] - c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqrt[I - a]*d)]))/d^2
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 665, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5574, 2855, 2916, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx$$

$$\begin{aligned}
& \downarrow \text{5574} \\
& \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{c + d\sqrt{x}} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{c + d\sqrt{x}} dx \\
& \downarrow \text{2855} \\
& i \int \frac{\sqrt{x} \log(-ia - ibx + 1)}{c + d\sqrt{x}} d\sqrt{x} - i \int \frac{\sqrt{x} \log(ia + ibx + 1)}{c + d\sqrt{x}} d\sqrt{x} \\
& \downarrow \text{2916} \\
& i \int \left(\frac{\log(-ia - ibx + 1)}{d} - \frac{c \log(-ia - ibx + 1)}{d(c + d\sqrt{x})} \right) d\sqrt{x} - \\
& i \int \left(\frac{\log(ia + ibx + 1)}{d} - \frac{c \log(ia + ibx + 1)}{d(c + d\sqrt{x})} \right) d\sqrt{x} \\
& \downarrow \text{2009} \\
& i \left(\frac{2\sqrt{a+i} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a+i}}\right)}{\sqrt{bd}} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{-a-id}}\right)}{d^2} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{-a-id}}\right)}{d^2} + \frac{c \log(c + d\sqrt{x}) \log}{c} \right) \\
& i \left(\frac{2\sqrt{-a+i} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-a+i}}\right)}{\sqrt{bd}} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}-\sqrt{i-ad}}\right)}{d^2} + \frac{c \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(c+d\sqrt{x})}{\sqrt{bc}+\sqrt{i-ad}}\right)}{d^2} + \frac{c \log(c + d\sqrt{x}) \log}{c} \right)
\end{aligned}$$

input `Int[ArcTan[a + b*x]/(c + d*Sqrt[x]),x]`

output

```

I*((-2*Sqrt[x])/d + (2*Sqrt[I + a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[I + a]])/
(Sqrt[b]*d) + (c*Log[(d*(Sqrt[-I - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c + Sqr
t[-I - a]*d)]*Log[c + d*Sqrt[x]])/d^2 + (c*Log[-((d*(Sqrt[-I - a] + Sqrt[b]
)*Sqrt[x]))/(Sqrt[b]*c - Sqrt[-I - a]*d)]*Log[c + d*Sqrt[x]])/d^2 + (Sqrt
[x]*Log[1 - I*a - I*b*x])/d - (c*Log[c + d*Sqrt[x]]*Log[1 - I*a - I*b*x])/
d^2 + (c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[-I - a]*d
)]/d^2 + (c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqrt[-I - a]
*d)])/d^2) - I*((-2*Sqrt[x])/d + (2*Sqrt[I - a]*ArcTanh[(Sqrt[b]*Sqrt[x])/
Sqrt[I - a]])/(Sqrt[b]*d) + (c*Log[(d*(Sqrt[I - a] - Sqrt[b]*Sqrt[x]))/(Sq
rt[b]*c + Sqrt[I - a]*d)]*Log[c + d*Sqrt[x]])/d^2 + (c*Log[-((d*(Sqrt[I -
a] + Sqrt[b]*Sqrt[x]))/(Sqrt[b]*c - Sqrt[I - a]*d)]*Log[c + d*Sqrt[x]])/d
^2 + (Sqrt[x]*Log[1 + I*a + I*b*x])/d - (c*Log[c + d*Sqrt[x]]*Log[1 + I*a
+ I*b*x])/d^2 + (c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c - Sqrt[
I - a]*d)])/d^2 + (c*PolyLog[2, (Sqrt[b]*(c + d*Sqrt[x]))/(Sqrt[b]*c + Sqr
t[I - a]*d)])/d^2)

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2855

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))* (b_.)]^(p_.))*((f_.) + (g_
.)*(x_)^(r_.))^(q_.), x_Symbol] := With[{k = Denominator[r]}, Simp[k Subst
[Int[x^(k - 1)*(f + g*x^(k*r))^q*(a + b*Log[c*(d + e*x^k)^n])^p, x], x, x^(
1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q}, x] && FractionQ[r] && I
GtQ[p, 0]
```

rule 2916

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))^(p_.)]*(b_.)]^(q_.)*(x_)^(m
_.)*((f_.) + (g_.)*(x_)^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log
[c*(d + e*x^n)^p])^q, x^m*(f + g*x)^r, x], x] /; FreeQ[{a, b, c, d, e, f, g
, n, p, q}, x] && IntegerQ[m] && IntegerQ[r]
```

rule 5574

```
Int[ArcTan[(a_) + (b_.)*(x_)]/((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 364, normalized size of antiderivative = 0.54

method	result
derivativedivides	$\frac{2 \arctan(bx+a)\sqrt{x}}{d} - \frac{2 \arctan(bx+a)c \ln(c+d\sqrt{x})}{d^2} - \frac{d^2 \left(\frac{\sum_{R=\text{RootOf}(b^2 Z^4 - 4b^2 c Z^3 + (2ab d^2 + 6b^2 c^2) Z^2 + \dots)}{d^2} \right)}{4b}$
default	$\frac{2 \arctan(bx+a)\sqrt{x}}{d} - \frac{2 \arctan(bx+a)c \ln(c+d\sqrt{x})}{d^2} - \frac{d^2 \left(\frac{\sum_{R=\text{RootOf}(b^2 Z^4 - 4b^2 c Z^3 + (2ab d^2 + 6b^2 c^2) Z^2 + \dots)}{d^2} \right)}{4b}$

```
input int(arctan(b*x+a)/(c+d*x^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 2*arctan(b*x+a)/d*x^(1/2)-2*arctan(b*x+a)*c/d^2*ln(c+d*x^(1/2))-4*b/d^2*(1/4*d^2/b*sum((_R^2-2*_R*c+c^2)/(_R^3*b-3*_R^2*b*c+_R*a*d^2+3*_R*b*c^2-a*c*d^2-b*c^3)*ln(d*x^(1/2)-_R+c),_R=RootOf(b^2*_Z^4-4*b^2*c*_Z^3+(2*a*b*d^2+6*b^2*c^2)*_Z^2+(-4*a*b*c*d^2-4*b^2*c^3)*_Z+a^2*d^4+2*a*b*c^2*d^2+b^2*c^4+d^4))-1/4*c*d^2/b*sum(1/(_R1^2*b-2*_R1*b*c+a*d^2+b*c^2)*(ln(c+d*x^(1/2))*ln((-d*x^(1/2)+_R1-c)/_R1)+dilog((-d*x^(1/2)+_R1-c)/_R1)),_R1=RootOf(b^2*_Z^4-4*b^2*c*_Z^3+(2*a*b*d^2+6*b^2*c^2)*_Z^2+(-4*a*b*c*d^2-4*b^2*c^3)*_Z+a^2*d^4+2*a*b*c^2*d^2+b^2*c^4+d^4))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\arctan(bx + a)}{d\sqrt{x} + c} dx$$

input `integrate(arctan(b*x+a)/(c+d*x^(1/2)),x, algorithm="fricas")`

output `integral((d*sqrt(x)*arctan(b*x + a) - c*arctan(b*x + a))/(d^2*x - c^2), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \text{Timed out}$$

input `integrate(atan(b*x+a)/(c+d*x**(1/2)),x)`

output `Timed out`

Maxima [F]

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\arctan(bx + a)}{d\sqrt{x} + c} dx$$

input `integrate(arctan(b*x+a)/(c+d*x^(1/2)),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(d*sqrt(x) + c), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \text{Exception raised: TypeError}$$

input `integrate(arctan(b*x+a)/(c+d*x^(1/2)),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \int \frac{\operatorname{atan}(a + bx)}{c + d\sqrt{x}} dx$$

input `int(atan(a + b*x)/(c + d*x^(1/2)),x)`

output `int(atan(a + b*x)/(c + d*x^(1/2)), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + d\sqrt{x}} dx = \left(\int \frac{\operatorname{atan}(bx + a)}{-d^2x + c^2} dx \right) c - \left(\int \frac{\sqrt{x} \operatorname{atan}(bx + a)}{-d^2x + c^2} dx \right) d$$

input `int(atan(b*x+a)/(c+d*x^(1/2)),x)`

output `int(atan(a + b*x)/(c**2 - d**2*x),x)*c - int((sqrt(x)*atan(a + b*x))/(c**2 - d**2*x),x)*d`

$$3.61 \quad \int \frac{\arctan(a+bx)}{c+\frac{d}{\sqrt{x}}} dx$$

Optimal result	526
Mathematica [A] (verified)	527
Rubi [A] (verified)	528
Maple [C] (warning: unable to verify)	531
Fricas [F]	532
Sympy [F(-1)]	532
Maxima [F]	533
Giac [F(-2)]	533
Mupad [F(-1)]	533
Reduce [F]	534

Optimal result

Integrand size = 18, antiderivative size = 770

$$\begin{aligned}
\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = & -\frac{2i\sqrt{i+ad} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i+a}}\right)}{\sqrt{bc^2}} + \frac{2i\sqrt{i-ad} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i-a}}\right)}{\sqrt{bc^2}} \\
& - \frac{id^2 \log\left(\frac{c(\sqrt{-i-a}-\sqrt{b}\sqrt{x})}{\sqrt{-i-ac}+\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& + \frac{id^2 \log\left(\frac{c(\sqrt{i-a}-\sqrt{b}\sqrt{x})}{\sqrt{i-ac}+\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& - \frac{id^2 \log\left(\frac{c(\sqrt{-i-a}+\sqrt{b}\sqrt{x})}{\sqrt{-i-ac}-\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& + \frac{id^2 \log\left(\frac{c(\sqrt{i-a}+\sqrt{b}\sqrt{x})}{\sqrt{i-ac}-\sqrt{bd}}\right) \log(d + c\sqrt{x})}{c^3} \\
& - \frac{id\sqrt{x} \log(1 - ia - ibx)}{c^2} + \frac{id^2 \log(d + c\sqrt{x}) \log(1 - ia - ibx)}{c^3} \\
& + \frac{id\sqrt{x} \log(1 + ia + ibx)}{c^2} - \frac{(1 + ia + ibx) \log(1 + ia + ibx)}{2bc} \\
& - \frac{id^2 \log(d + c\sqrt{x}) \log(1 + ia + ibx)}{c^3} \\
& - \frac{(1 - ia - ibx) \log(-i(i + a + bx))}{2bc} \\
& - \frac{id^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-i-ac}-\sqrt{bd}}\right)}{c^3} \\
& + \frac{id^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{i-ac}-\sqrt{bd}}\right)}{c^3} \\
& - \frac{id^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{-i-ac}+\sqrt{bd}}\right)}{c^3} + \frac{id^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(d+c\sqrt{x})}{\sqrt{i-ac}+\sqrt{bd}}\right)}{c^3}
\end{aligned}$$

output

```
I*d^2*ln(c*((I-a)^(1/2)-b^(1/2)*x^(1/2))/((I-a)^(1/2)*c+b^(1/2)*d))*ln(d+c*x^(1/2))/c^3-2*I*(I+a)^(1/2)*d*arctan(b^(1/2)*x^(1/2)/(I+a)^(1/2))/b^(1/2)/c^2+I*d^2*polylog(2,b^(1/2)*(d+c*x^(1/2))/((I-a)^(1/2)*c+b^(1/2)*d))/c^3+I*d^2*polylog(2,-b^(1/2)*(d+c*x^(1/2))/((I-a)^(1/2)*c-b^(1/2)*d))/c^3-I*d^2*ln(d+c*x^(1/2))*ln(1+I*a+I*b*x)/c^3+I*d*x^(1/2)*ln(1+I*a+I*b*x)/c^2-I*d^2*polylog(2,b^(1/2)*(d+c*x^(1/2))/((-I-a)^(1/2)*c+b^(1/2)*d))/c^3-I*d^2*ln(c*((-I-a)^(1/2)-b^(1/2)*x^(1/2))/((-I-a)^(1/2)*c+b^(1/2)*d))*ln(d+c*x^(1/2))/c^3-I*d*x^(1/2)*ln(1-I*a-I*b*x)/c^2-1/2*(1+I*a+I*b*x)*ln(1+I*a+I*b*x)/b/c+I*d^2*ln(c*((I-a)^(1/2)+b^(1/2)*x^(1/2))/((I-a)^(1/2)*c-b^(1/2)*d))*ln(d+c*x^(1/2))/c^3-1/2*(1-I*a-I*b*x)*ln(-I*(I+a+b*x))/b/c-I*d^2*polylog(2,-b^(1/2)*(d+c*x^(1/2))/((-I-a)^(1/2)*c-b^(1/2)*d))/c^3+2*I*(I-a)^(1/2)*d*arctanh(b^(1/2)*x^(1/2)/(I-a)^(1/2))/b^(1/2)/c^2+I*d^2*ln(d+c*x^(1/2))*ln(1-I*a-I*b*x)/c^3-I*d^2*ln(c*((-I-a)^(1/2)+b^(1/2)*x^(1/2))/((-I-a)^(1/2)*c-b^(1/2)*d))*ln(d+c*x^(1/2))/c^3
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 770, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx =$$

$$i \left(4\sqrt{i+a}\sqrt{bcd} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i+a}}\right) - 4\sqrt{i-a}\sqrt{bcd} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{i-a}}\right) + 2bd^2 \log\left(\frac{c(\sqrt{-i-a}-\sqrt{b}\sqrt{x})}{\sqrt{-i-ac}+\sqrt{bd}}\right) \right) \log(d +$$

input

```
Integrate[ArcTan[a + b*x]/(c + d/Sqrt[x]),x]
```


output

```

((-1/2*I)*(4*Sqrt[I + a]*Sqrt[b]*c*d*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[I + a]]
- 4*Sqrt[I - a]*Sqrt[b]*c*d*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[I - a]] + 2*b*
d^2*Log[(c*(Sqrt[-I - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[-I - a]*c + Sqrt[b]*d)]
*Log[d + c*Sqrt[x]] - 2*b*d^2*Log[(c*(Sqrt[I - a] - Sqrt[b]*Sqrt[x]))/(Sqr
t[I - a]*c + Sqrt[b]*d)]*Log[d + c*Sqrt[x]] + 2*b*d^2*Log[(c*(Sqrt[-I - a]
+ Sqrt[b]*Sqrt[x]))/(Sqrt[-I - a]*c - Sqrt[b]*d)]*Log[d + c*Sqrt[x]] - 2*
b*d^2*Log[(c*(Sqrt[I - a] + Sqrt[b]*Sqrt[x]))/(Sqrt[I - a]*c - Sqrt[b]*d)]
*Log[d + c*Sqrt[x]] - I*c^2*Log[1 + I*a + I*b*x] + a*c^2*Log[1 + I*a + I*b
*x] - 2*b*c*d*Sqrt[x]*Log[1 + I*a + I*b*x] + b*c^2*x*Log[1 + I*a + I*b*x]
+ 2*b*d^2*Log[d + c*Sqrt[x]]*Log[1 + I*a + I*b*x] - I*c^2*Log[(-I)*(I + a
+ b*x)] - a*c^2*Log[(-I)*(I + a + b*x)] + 2*b*c*d*Sqrt[x]*Log[(-I)*(I + a
+ b*x)] - b*c^2*x*Log[(-I)*(I + a + b*x)] - 2*b*d^2*Log[d + c*Sqrt[x]]*Log
[(-I)*(I + a + b*x)] + 2*b*d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]))/(-(Sqr
t[-I - a]*c) + Sqrt[b]*d)] + 2*b*d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]))/
(Sqrt[-I - a]*c + Sqrt[b]*d)] - 2*b*d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]
))/(-(Sqrt[I - a]*c) + Sqrt[b]*d)] - 2*b*d^2*PolyLog[2, (Sqrt[b]*(d + c*Sq
rt[x]))/(Sqrt[I - a]*c + Sqrt[b]*d)))/(b*c^3)

```

Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 792, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5574, 2855, 2005, 2916, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx \\
& \quad \downarrow \text{5574} \\
& \frac{1}{2}i \int \frac{\log(-ia - ibx + 1)}{c + \frac{d}{\sqrt{x}}} dx - \frac{1}{2}i \int \frac{\log(ia + ibx + 1)}{c + \frac{d}{\sqrt{x}}} dx \\
& \quad \downarrow \text{2855} \\
& i \int \frac{\sqrt{x} \log(-ia - ibx + 1)}{c + \frac{d}{\sqrt{x}}} d\sqrt{x} - i \int \frac{\sqrt{x} \log(ia + ibx + 1)}{c + \frac{d}{\sqrt{x}}} d\sqrt{x} \\
& \quad \downarrow \text{2005}
\end{aligned}$$

$$i \int \frac{x \log(-ia - ibx + 1)}{\sqrt{xc + d}} d\sqrt{x} - i \int \frac{x \log(ia + ibx + 1)}{\sqrt{xc + d}} d\sqrt{x}$$

↓ 2916

$$i \int \left(\frac{\log(-ia - ibx + 1)d^2}{c^2(\sqrt{xc + d})} - \frac{\log(-ia - ibx + 1)d}{c^2} + \frac{\sqrt{x} \log(-ia - ibx + 1)}{c} \right) d\sqrt{x} -$$

$$i \int \left(\frac{\log(ia + ibx + 1)d^2}{c^2(\sqrt{xc + d})} - \frac{\log(ia + ibx + 1)d}{c^2} + \frac{\sqrt{x} \log(ia + ibx + 1)}{c} \right) d\sqrt{x}$$

↓ 2009

$$i \left(-\frac{2\sqrt{a + id} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a+i}}\right)}{\sqrt{bc^2}} - \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{-a-ic-\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{-a-ic+\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \log(c\sqrt{x})}{c^3} \right)$$

$$i \left(-\frac{2\sqrt{-a + id} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-a+i}}\right)}{\sqrt{bc^2}} - \frac{d^2 \operatorname{PolyLog}\left(2, -\frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{i-ac-\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \operatorname{PolyLog}\left(2, \frac{\sqrt{b}(\sqrt{xc+d})}{\sqrt{i-ac+\sqrt{bd}}}\right)}{c^3} - \frac{d^2 \log(c\sqrt{x})}{c^3} \right)$$

input

```
Int[ArcTan[a + b*x]/(c + d/Sqrt[x]), x]
```

output

```

I*((2*d*Sqrt[x])/c^2 - x/(2*c) - (2*Sqrt[I + a]*d*ArcTan[(Sqrt[b]*Sqrt[x])
/Sqrt[I + a]])/(Sqrt[b]*c^2) - (d^2*Log[(c*(Sqrt[-I - a] - Sqrt[b]*Sqrt[x]
)))/(Sqrt[-I - a]*c + Sqrt[b]*d)]*Log[d + c*Sqrt[x]])/c^3 - (d^2*Log[(c*(Sq
rt[-I - a] + Sqrt[b]*Sqrt[x])))/(Sqrt[-I - a]*c - Sqrt[b]*d)]*Log[d + c*Sqr
t[x]])/c^3 - (d*Sqrt[x]*Log[1 - I*a - I*b*x])/c^2 + (d^2*Log[d + c*Sqrt[x]
]*Log[1 - I*a - I*b*x])/c^3 + ((I/2)*(1 - I*a - I*b*x)*Log[(-I)*(I + a + b
*x)])/(b*c) - (d^2*PolyLog[2, -(Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[-I - a]*c
- Sqrt[b]*d)))/c^3 - (d^2*PolyLog[2, (Sqrt[b]*(d + c*Sqrt[x]))/(Sqrt[-I -
a]*c + Sqrt[b]*d)])/c^3) - I*((2*d*Sqrt[x])/c^2 - x/(2*c) - (2*Sqrt[I - a
]*d*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[I - a]])/(Sqrt[b]*c^2) - (d^2*Log[(c*(S
qrt[I - a] - Sqrt[b]*Sqrt[x]))/(Sqrt[I - a]*c + Sqrt[b]*d)]*Log[d + c*Sqrt
[x]])/c^3 - (d^2*Log[(c*(Sqrt[I - a] + Sqrt[b]*Sqrt[x]))/(Sqrt[I - a]*c -
Sqrt[b]*d)]*Log[d + c*Sqrt[x]])/c^3 - (d*Sqrt[x]*Log[1 + I*a + I*b*x])/c^2
- ((I/2)*(1 + I*a + I*b*x)*Log[1 + I*a + I*b*x])/(b*c) + (d^2*Log[d + c*S
qrt[x]*Log[1 + I*a + I*b*x])/c^3 - (d^2*PolyLog[2, -(Sqrt[b]*(d + c*Sqrt
[x]))/(Sqrt[I - a]*c - Sqrt[b]*d)])/c^3 - (d^2*PolyLog[2, (Sqrt[b]*(d + c
*Sqrt[x]))/(Sqrt[I - a]*c + Sqrt[b]*d)])/c^3)

```

Defintions of rubi rules used

rule 2005

```

Int[(Fx_)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :=> Int[x^(m
+ n*p)*(b + a/x^n)^p*Fx, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && Neg
Q[n]

```

rule 2009

```

Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]

```

rule 2855

```

Int[((a_) + Log[(c_)*((d_) + (e_)*(x_)^(n_))]*(b_))^(p_)*((f_) + (g_
.)*(x_)^(r_))^(q_), x_Symbol] :=> With[{k = Denominator[r]}, Simp[k Subst
[Int[x^(k - 1)*(f + g*x^(k*r))^q*(a + b*Log[c*(d + e*x^k)^n])^p, x], x, x^(
1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q}, x] && FractionQ[r] && I
GtQ[p, 0]

```

```
rule 2916 Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)]*(b_.))^(q_.)*(x_)^(m
_.)*(f_.) + (g_.)*(x_)^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log
[c*(d + e*x^n)^p])^q, x^m*(f + g*x)^r, x], x] /; FreeQ[{a, b, c, d, e, f, g
, n, p, q}, x] && IntegerQ[m] && IntegerQ[r]
```

```
rule 5574 Int[ArcTan[(a_) + (b_.)*(x_)]/((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[
I/2 Int[Log[1 - I*a - I*b*x]/(c + d*x^n), x], x] - Simp[I/2 Int[Log[1 +
I*a + I*b*x]/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[n]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.17 (sec) , antiderivative size = 388, normalized size of antiderivative = 0.50

method	result
derivativedivides	$\frac{\arctan(bx+a)x}{c} - \frac{2 \arctan(bx+a)d\sqrt{x}}{c^2} + \frac{2 \arctan(bx+a)d^2 \ln(d+c\sqrt{x})}{c^3} - \frac{c}{4b} \left(\frac{c}{-R=\text{RootOf}(b^2 Z^4 - 4b^2 d Z^2 - c^2)} $
default	$\frac{\arctan(bx+a)x}{c} - \frac{2 \arctan(bx+a)d\sqrt{x}}{c^2} + \frac{2 \arctan(bx+a)d^2 \ln(d+c\sqrt{x})}{c^3} - \frac{c}{4b} \left(\frac{c}{-R=\text{RootOf}(b^2 Z^4 - 4b^2 d Z^2 - c^2)} $

```
input int(arctan(b*x+a)/(c+d/x^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
arctan(b*x+a)/c*x-2*arctan(b*x+a)/c^2*d*x^(1/2)+2*arctan(b*x+a)*d^2/c^3*ln
(d+c*x^(1/2))-4*b/c^2*(-1/8*c/b*sum((-_R^3+5*_R^2*d-7*_R*d^2+3*d^3)/(_R^3*
b-3*_R^2*b*d+_R*a*c^2+3*_R*b*d^2-a*c^2*d-b*d^3)*ln(c*x^(1/2)-_R+d),_R=Root
Of(b^2*_Z^4-4*b^2*d*_Z^3+(2*a*b*c^2+6*b^2*d^2)*_Z^2+(-4*a*b*c^2*d-4*b^2*d^
3)*_Z+a^2*c^4+2*a*b*c^2*d^2+b^2*d^4+c^4))+1/4*c*d^2/b*sum(1/(_R1^2*b-2*_R1
*b*d+a*c^2+b*d^2)*(ln(d+c*x^(1/2))*ln((-c*x^(1/2)+_R1-d)/_R1)+dilog((-c*x^
(1/2)+_R1-d)/_R1)),_R1=RootOf(b^2*_Z^4-4*b^2*d*_Z^3+(2*a*b*c^2+6*b^2*d^2)*
_Z^2+(-4*a*b*c^2*d-4*b^2*d^3)*_Z+a^2*c^4+2*a*b*c^2*d^2+b^2*d^4+c^4)))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{\sqrt{x}}} dx$$

input

```
integrate(arctan(b*x+a)/(c+d/x^(1/2)),x, algorithm="fricas")
```

output

```
integral((c*x*arctan(b*x + a) - d*sqrt(x)*arctan(b*x + a))/(c^2*x - d^2),
x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \text{Timed out}$$

input

```
integrate(atan(b*x+a)/(c+d/x**(1/2)),x)
```

output

```
Timed out
```

Maxima [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\arctan(bx + a)}{c + \frac{d}{\sqrt{x}}} dx$$

input `integrate(arctan(b*x+a)/(c+d/x^(1/2)),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(c + d/sqrt(x)), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \text{Exception raised: TypeError}$$

input `integrate(arctan(b*x+a)/(c+d/x^(1/2)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch fo
r the root of a polynomial with parameters. This might be wrong.The choice
was done`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \int \frac{\text{atan}(a + bx)}{c + \frac{d}{\sqrt{x}}} dx$$

input `int(atan(a + b*x)/(c + d/x^(1/2)),x)`

output `int(atan(a + b*x)/(c + d/x^(1/2)), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{c + \frac{d}{\sqrt{x}}} dx = \left(\int \frac{\operatorname{atan}(bx + a) x}{c^2 x - d^2} dx \right) c - \left(\int \frac{\sqrt{x} \operatorname{atan}(bx + a)}{c^2 x - d^2} dx \right) d$$

input `int(atan(b*x+a)/(c+d/x^(1/2)),x)`

output `int((atan(a + b*x)*x)/(c**2*x - d**2),x)*c - int((sqrt(x)*atan(a + b*x))/(c**2*x - d**2),x)*d`

3.62 $\int \frac{a+b \arctan(c+dx)}{e+fx+gx^2} dx$

Optimal result	535
Mathematica [A] (warning: unable to verify)	536
Rubi [A] (verified)	537
Maple [B] (verified)	538
Fricas [F]	539
Sympy [F(-1)]	540
Maxima [F(-2)]	540
Giac [F]	540
Mupad [F(-1)]	541
Reduce [F]	541

Optimal result

Integrand size = 23, antiderivative size = 398

$$\begin{aligned}
 & \int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx \\
 &= \frac{(a + b \arctan(c + dx)) \log \left(-\frac{2(2cg - d(f - \sqrt{f^2 - 4eg}) - 2g(c + dx))}{(df + 2ig - 2cg - d\sqrt{f^2 - 4eg})(1 - i(c + dx))} \right)}{\sqrt{f^2 - 4eg}} \\
 & - \frac{(a + b \arctan(c + dx)) \log \left(-\frac{2(2cg - d(f + \sqrt{f^2 - 4eg}) - 2g(c + dx))}{(2(i - c)g + d(f + \sqrt{f^2 - 4eg}))(1 - i(c + dx))} \right)}{\sqrt{f^2 - 4eg}} \\
 & - \frac{i b \operatorname{PolyLog} \left(2, 1 + \frac{2(2cg - d(f - \sqrt{f^2 - 4eg}) - 2g(c + dx))}{(df + 2ig - 2cg - d\sqrt{f^2 - 4eg})(1 - i(c + dx))} \right)}{2\sqrt{f^2 - 4eg}} \\
 & + \frac{i b \operatorname{PolyLog} \left(2, 1 + \frac{2(2cg - d(f + \sqrt{f^2 - 4eg}) - 2g(c + dx))}{(2(i - c)g + d(f + \sqrt{f^2 - 4eg}))(1 - i(c + dx))} \right)}{2\sqrt{f^2 - 4eg}}
 \end{aligned}$$

output

```
(a+b*arctan(d*x+c))*ln((-4*c*g+2*d*(f-(-4*e*g+f^2)^(1/2))+4*g*(d*x+c))/(d*f+2*I*g-2*c*g-d*(-4*e*g+f^2)^(1/2))/(1-I*(d*x+c)))/(-4*e*g+f^2)^(1/2)-(a+b*arctan(d*x+c))*ln((-4*c*g+2*d*(f+(-4*e*g+f^2)^(1/2))+4*g*(d*x+c))/(2*(I-c)*g+d*(f+(-4*e*g+f^2)^(1/2)))/(1-I*(d*x+c)))/(-4*e*g+f^2)^(1/2)-1/2*I*b*polylog(2,1+2*(2*c*g-d*(f-(-4*e*g+f^2)^(1/2))-2*g*(d*x+c))/(d*f+2*I*g-2*c*g-d*(-4*e*g+f^2)^(1/2))/(1-I*(d*x+c)))/(-4*e*g+f^2)^(1/2)+1/2*I*b*polylog(2,1+2*(2*c*g-d*(f+(-4*e*g+f^2)^(1/2))-2*g*(d*x+c))/(2*(I-c)*g+d*(f+(-4*e*g+f^2)^(1/2)))/(1-I*(d*x+c)))/(-4*e*g+f^2)^(1/2)
```

Mathematica [A] (warning: unable to verify)

Time = 0.54 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.24

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx =$$

$$\frac{4a \operatorname{arctanh}\left(\frac{f+2gx}{\sqrt{f^2-4eg}}\right) - ib \log\left(\frac{d(-f+\sqrt{f^2-4eg}-2gx)}{2(i+c)g+d(-f+\sqrt{f^2-4eg})}\right) \log(1 - i(c + dx)) + ib \log\left(\frac{d(f+\sqrt{f^2-4eg}+2gx)}{-2(i+c)g+d(f+\sqrt{f^2-4eg})}\right) \log(1 + i(c + dx))}{1}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])/(e + f*x + g*x^2),x]
```

output

```
-1/2*(4*a*ArcTanh[(f + 2*g*x)/Sqrt[f^2 - 4*e*g]] - I*b*Log[(d*(-f + Sqrt[f^2 - 4*e*g] - 2*g*x))/(2*(I + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))]*Log[1 - I*(c + d*x)] + I*b*Log[(d*(f + Sqrt[f^2 - 4*e*g] + 2*g*x))/(-2*(I + c)*g + d*(f + Sqrt[f^2 - 4*e*g]))]*Log[1 - I*(c + d*x)] + I*b*Log[(d*(-f + Sqrt[f^2 - 4*e*g] - 2*g*x))/(2*(-I + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))]*Log[1 + I*(c + d*x)] - I*b*Log[(d*(f + Sqrt[f^2 - 4*e*g] + 2*g*x))/(-2*(-I + c)*g + d*(f + Sqrt[f^2 - 4*e*g]))]*Log[1 + I*(c + d*x)] + I*b*PolyLog[2, (2*g*(-I + c + d*x))/(2*(-I + c)*g + d*(-f + Sqrt[f^2 - 4*e*g]))] - I*b*PolyLog[2, (2*g*(-I + c + d*x))/(2*(-I + c)*g - d*(f + Sqrt[f^2 - 4*e*g]))] - I*b*PolyLog[2, (2*g*(I + c + d*x))/(2*(I + c)*g + d*(-f + Sqrt[f^2 - 4*e*g])] + I*b*PolyLog[2, (2*g*(I + c + d*x))/(2*(I + c)*g - d*(f + Sqrt[f^2 - 4*e*g])])]/Sqrt[f^2 - 4*e*g]
```

Rubi [A] (verified)

Time = 1.23 (sec) , antiderivative size = 427, normalized size of antiderivative = 1.07, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx \\
 & \quad \downarrow \text{7279} \\
 & \int \left(\frac{a}{e + fx + gx^2} + \frac{b \arctan(c + dx)}{e + fx + gx^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{2a \operatorname{arctanh}\left(\frac{f+2gx}{\sqrt{f^2-4eg}}\right)}{\sqrt{f^2-4eg}} + \frac{b \arctan(c + dx) \log\left(-\frac{2(-2g(c+dx)+2cg-d(f-\sqrt{f^2-4eg}))}{(1-i(c+dx))(-2cg-d\sqrt{f^2-4eg}+df+2ig)}\right)}{\sqrt{f^2-4eg}} \\
 & \quad - \frac{b \arctan(c + dx) \log\left(-\frac{2(-2g(c+dx)+2cg-d(\sqrt{f^2-4eg}+f))}{(1-i(c+dx))(d(\sqrt{f^2-4eg}+f)+2(-c+i)g)}\right)}{\sqrt{f^2-4eg}} \\
 & \quad + \frac{i b \operatorname{PolyLog}\left(2, \frac{2(2cg-2(c+dx)g-d(f-\sqrt{f^2-4eg}))}{(fd-\sqrt{f^2-4eg}d-2cg+2ig)(1-i(c+dx))} + 1\right)}{2\sqrt{f^2-4eg}} \\
 & \quad + \frac{i b \operatorname{PolyLog}\left(2, \frac{2(2cg-2(c+dx)g-d(f+\sqrt{f^2-4eg}))}{(2(i-c)g+d(f+\sqrt{f^2-4eg}))(1-i(c+dx))} + 1\right)}{2\sqrt{f^2-4eg}}
 \end{aligned}$$

input

```
Int[(a + b*ArcTan[c + d*x])/(e + f*x + g*x^2), x]
```

output

$$\begin{aligned} & (-2*a*ArcTanh[(f + 2*g*x)/Sqrt[f^2 - 4*e*g])/Sqrt[f^2 - 4*e*g] + (b*ArcTan[c + d*x]*Log[(-2*(2*c*g - d*(f - Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((d*f + (2*I)*g - 2*c*g - d*Sqrt[f^2 - 4*e*g])*(1 - I*(c + d*x)))]/Sqrt[f^2 - 4*e*g] - (b*ArcTan[c + d*x]*Log[(-2*(2*c*g - d*(f + Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((2*(I - c)*g + d*(f + Sqrt[f^2 - 4*e*g])*(1 - I*(c + d*x))))]/Sqrt[f^2 - 4*e*g] - ((I/2)*b*PolyLog[2, 1 + (2*(2*c*g - d*(f - Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((d*f + (2*I)*g - 2*c*g - d*Sqrt[f^2 - 4*e*g])*(1 - I*(c + d*x)))]/Sqrt[f^2 - 4*e*g] + ((I/2)*b*PolyLog[2, 1 + (2*(2*c*g - d*(f + Sqrt[f^2 - 4*e*g]) - 2*g*(c + d*x))]/((2*(I - c)*g + d*(f + Sqrt[f^2 - 4*e*g])*(1 - I*(c + d*x)))]/Sqrt[f^2 - 4*e*g] \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp[IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 7279

$$\text{Int}[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] \text{ :> With}[v = \text{RationalFunctionExpand}[u/(a + b*x^n + c*x^(2*n)), x], \text{Int}[v, x] \text{ /; SumQ}[v]] \text{ /; FreeQ}[{a, b, c}, x] \ \&\& \ \text{EqQ}[n2, 2*n] \ \&\& \ \text{IGtQ}[n, 0]$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 966 vs. $2(358) = 716$.

Time = 2.08 (sec) , antiderivative size = 967, normalized size of antiderivative = 2.43

method	result
risch	$\frac{db \ln(-idx-ic+1) \ln\left(\frac{2icg-ifd+2(-idx-ic+1)g-\sqrt{4d^2eg-d^2f^2-2g}}{2icg-ifd-\sqrt{4d^2eg-d^2f^2-2g}}\right)}{2\sqrt{4d^2eg-d^2f^2}} - \frac{db \ln(-idx-ic+1) \ln\left(\frac{2icg-ifd+2(-idx-ic+1)g-\sqrt{4d^2eg-d^2f^2-2g}}{2icg-ifd+\sqrt{4d^2eg-d^2f^2-2g}}\right)}{2\sqrt{4d^2eg-d^2f^2}}$
parts	Expression too large to display
derivativedivides	Expression too large to display
default	Expression too large to display

input

$$\text{int}((a+b*\arctan(d*x+c))/(g*x^2+f*x+e), x, \text{method}=_RETURNVERBOSE)$$

output

```

1/2*d*b*ln(1-I*c-I*d*x)/(4*d^2*e*g-d^2*f^2)^(1/2)*ln((2*I*c*g-I*f*d+2*(1-I
*c-I*d*x)*g-(4*d^2*e*g-d^2*f^2)^(1/2)-2*g)/(2*I*c*g-I*f*d-(4*d^2*e*g-d^2*f
^2)^(1/2)-2*g))-1/2*d*b*ln(1-I*c-I*d*x)/(4*d^2*e*g-d^2*f^2)^(1/2)*ln((2*I*
c*g-I*f*d+2*(1-I*c-I*d*x)*g+(4*d^2*e*g-d^2*f^2)^(1/2)-2*g)/(2*I*c*g-I*f*d+
(4*d^2*e*g-d^2*f^2)^(1/2)-2*g))+1/2*d*b/(4*d^2*e*g-d^2*f^2)^(1/2)*dilog((2
*I*c*g-I*f*d+2*(1-I*c-I*d*x)*g-(4*d^2*e*g-d^2*f^2)^(1/2)-2*g)/(2*I*c*g-I*f
*d-(4*d^2*e*g-d^2*f^2)^(1/2)-2*g))-1/2*d*b/(4*d^2*e*g-d^2*f^2)^(1/2)*dilog
((2*I*c*g-I*f*d+2*(1-I*c-I*d*x)*g+(4*d^2*e*g-d^2*f^2)^(1/2)-2*g)/(2*I*c*g-
I*f*d+(4*d^2*e*g-d^2*f^2)^(1/2)-2*g))-2*I*d*a/(-4*d^2*e*g+d^2*f^2)^(1/2)*a
rctan((2*I*c*g-I*f*d+2*(1-I*c-I*d*x)*g-2*g)/(-4*d^2*e*g+d^2*f^2)^(1/2))+1/
2*b*d*ln(1+I*c+I*d*x)/(4*d^2*e*g-d^2*f^2)^(1/2)*ln((2*I*c*g-I*f*d-2*(1+I*c
+I*d*x)*g+(4*d^2*e*g-d^2*f^2)^(1/2)+2*g)/(2*I*c*g-I*f*d+2*g+(4*d^2*e*g-d^2
*f^2)^(1/2)))-1/2*b*d*ln(1+I*c+I*d*x)/(4*d^2*e*g-d^2*f^2)^(1/2)*ln((2*I*c*
g-I*f*d-2*(1+I*c+I*d*x)*g-(4*d^2*e*g-d^2*f^2)^(1/2)+2*g)/(2*I*c*g-I*f*d+2*
g-(4*d^2*e*g-d^2*f^2)^(1/2)))+1/2*b*d/(4*d^2*e*g-d^2*f^2)^(1/2)*dilog((2*I
*c*g-I*f*d-2*(1+I*c+I*d*x)*g+(4*d^2*e*g-d^2*f^2)^(1/2)+2*g)/(2*I*c*g-I*f*d
+2*g+(4*d^2*e*g-d^2*f^2)^(1/2)))-1/2*b*d/(4*d^2*e*g-d^2*f^2)^(1/2)*dilog((
2*I*c*g-I*f*d-2*(1+I*c+I*d*x)*g-(4*d^2*e*g-d^2*f^2)^(1/2)+2*g)/(2*I*c*g-I*
f*d+2*g-(4*d^2*e*g-d^2*f^2)^(1/2)))

```

Fricas [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \int \frac{b \arctan(dx + c) + a}{gx^2 + fx + e} dx$$

input

```
integrate((a+b*arctan(d*x+c))/(g*x^2+f*x+e),x, algorithm="fricas")
```

output

```
integral((b*arctan(d*x + c) + a)/(g*x^2 + f*x + e), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))/(g*x**2+f*x+e),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arctan(d*x+c))/(g*x^2+f*x+e),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*e*g-f^2>0)', see `assume?` for more deta`

Giac [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \int \frac{b \arctan(dx + c) + a}{gx^2 + fx + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(g*x^2+f*x+e),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)/(g*x^2 + f*x + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \int \frac{a + b \operatorname{atan}(c + dx)}{gx^2 + fx + e} dx$$

input `int((a + b*atan(c + d*x))/(e + f*x + g*x^2),x)`output `int((a + b*atan(c + d*x))/(e + f*x + g*x^2), x)`**Reduce [F]**

$$\int \frac{a + b \arctan(c + dx)}{e + fx + gx^2} dx = \text{Too large to display}$$

input `int((a+b*atan(d*x+c))/(g*x^2+f*x+e),x)`

output

```
(4*atan(c + d*x)**2*b*c*e*g - atan(c + d*x)**2*b*c*f**2 + 2*sqrt(4*e*g - f
**2)*atan((f + 2*g*x)/sqrt(4*e*g - f**2))*a*f + 4*int(atan(c + d*x)/(c**2*
e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**
2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 + e + f*x + g*x**2),x)*b*c**2*e*f*g -
int(atan(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*
x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 + e + f*x +
g*x**2),x)*b*c**2*f**3 - 8*int(atan(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x
**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3
+ d**2*g*x**4 + e + f*x + g*x**2),x)*b*c*d*e**2*g + 2*int(atan(c + d*x)/(c
**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 +
d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 + e + f*x + g*x**2),x)*b*c*d*e*f*
*2 + 4*int(atan(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*
c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 + e +
f*x + g*x**2),x)*b*e*f*g - int(atan(c + d*x)/(c**2*e + c**2*f*x + c**2*g*x
**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3
+ d**2*g*x**4 + e + f*x + g*x**2),x)*b*f**3 - 8*int((atan(c + d*x)*x**2)/(
c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3
+ d**2*e*x**2 + d**2*f*x**3 + d**2*g*x**4 + e + f*x + g*x**2),x)*b*c*d*e*g
**2 + 2*int((atan(c + d*x)*x**2)/(c**2*e + c**2*f*x + c**2*g*x**2 + 2*c*d*
e*x + 2*c*d*f*x**2 + 2*c*d*g*x**3 + d**2*e*x**2 + d**2*f*x**3 + d**2*g*...
```

3.63 $\int \frac{\arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx$

Optimal result	543
Mathematica [A] (verified)	544
Rubi [A] (verified)	544
Maple [A] (verified)	545
Fricas [F]	546
Sympy [F]	546
Maxima [F]	546
Giac [F]	547
Mupad [F(-1)]	547
Reduce [F]	547

Optimal result

Integrand size = 28, antiderivative size = 132

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = -\frac{2i \arctan(a + bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b} + \frac{i \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b} - \frac{i \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b}$$

output

```
-2*I*arctan(b*x+a)*arctan((1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b+I*polylog(2,-I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b-I*polylog(2,I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b
```


Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.51

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \frac{i(2 \arctan(e^{i \arctan(a+bx)}) \arctan(a + bx) - \text{PolyLog}(2, -ie^{i \arctan(a+bx)}) + \text{PolyLog}(2, ie^{i \arctan(a+bx)}))}{b}$$

input `Integrate[ArcTan[a + b*x]/Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2], x]`output `((-I)*(2*ArcTan[E^(I*ArcTan[a + b*x])]*ArcTan[a + b*x] - PolyLog[2, (-I)*E^(I*ArcTan[a + b*x])]) + PolyLog[2, I*E^(I*ArcTan[a + b*x])])/b`**Rubi [A] (verified)**Time = 0.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.96, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {5578, 5421}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx$$

↓ 5578

$$\frac{\int \frac{\arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a + bx)}{b}$$

↓ 5421

$$\frac{-2i \arctan(a + bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) + i \text{PolyLog}\left(2, -\frac{i\sqrt{i(a+bx)+1}}{\sqrt{1-i(a+bx)}}\right) - i \text{PolyLog}\left(2, \frac{i\sqrt{i(a+bx)+1}}{\sqrt{1-i(a+bx)}}\right)}{b}$$

input `Int[ArcTan[a + b*x]/Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2], x]`

output

```
((-2*I)*ArcTan[a + b*x]*ArcTan[Sqrt[1 + I*(a + b*x)]/Sqrt[1 - I*(a + b*x)]
] + I*PolyLog[2, ((-I)*Sqrt[1 + I*(a + b*x)]/Sqrt[1 - I*(a + b*x)]] - I*P
olyLog[2, (I*Sqrt[1 + I*(a + b*x)]/Sqrt[1 - I*(a + b*x)])]/b
```

Defintions of rubi rules used

rule 5421

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol]
:> Simp[-2*I*(a + b*ArcTan[c*x])*(ArcTan[Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x]]/
(c*Sqrt[d])), x] + (Simp[I*b*(PolyLog[2, (-I)*(Sqrt[1 + I*c*x]/Sqrt[1 - I*c
*x]])/(c*Sqrt[d])), x] - Simp[I*b*(PolyLog[2, I*(Sqrt[1 + I*c*x]/Sqrt[1 - I
*c*x]])/(c*Sqrt[d])), x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] &&
GtQ[d, 0]
```

rule 5578

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((A_.) + (B_.)*(x_) + (
C_.)*(x_)^2)^(q_.), x_Symbol] :> Simp[1/d Subst[Int[(C/d^2 + (C/d^2)*x^2)
^q*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, A, B, C,
p, q}, x] && EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.02

method	result
default	$\frac{-\arctan(bx+a) \ln\left(1 + \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right) + \arctan(bx+a) \ln\left(1 - \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right) + i \operatorname{dilog}\left(1 + \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right) - i \operatorname{dilog}\left(1 - \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)}{b}$

input

```
int(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/b*(-arctan(b*x+a)*ln(1+I*(1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))+arctan(b*x+a
)*ln(1-I*(1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))+I*dilog(1+I*(1+I*(b*x+a))/(1+(
b*x+a)^2)^(1/2))-I*dilog(1-I*(1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)))
```

Fricas [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorithm="fricas")`

output `integral(arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1), x)`

Sympy [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx$$

input `integrate(atan(b*x+a)/(b**2*x**2+2*a*b*x+a**2+1)**(1/2),x)`

output `Integral(atan(a + b*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x)`

Maxima [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1), x)`

Giac [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx$$

input `int(atan(a + b*x)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2),x)`

output `int(atan(a + b*x)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `int(atan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x)`

output `int(atan(a + b*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1),x)`

$$3.64 \quad \int \frac{\arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx$$

Optimal result	548
Mathematica [A] (verified)	549
Rubi [A] (verified)	549
Maple [A] (verified)	551
Fricas [F]	551
Sympy [F(-1)]	552
Maxima [F]	552
Giac [F]	552
Mupad [F(-1)]	553
Reduce [F]	553

Optimal result

Integrand size = 33, antiderivative size = 216

$$\begin{aligned} & \int \frac{\arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx \\ &= -\frac{2i\sqrt{1+(a+bx)^2} \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b\sqrt{c+c(a+bx)^2}} \\ & \quad + \frac{i\sqrt{1+(a+bx)^2} \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b\sqrt{c+c(a+bx)^2}} \\ & \quad - \frac{i\sqrt{1+(a+bx)^2} \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b\sqrt{c+c(a+bx)^2}} \end{aligned}$$

output

```
-2*I*(1+(b*x+a)^2)^(1/2)*arctan(b*x+a)*arctan((1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)+I*(1+(b*x+a)^2)^(1/2)*polylog(2,-I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)-I*(1+(b*x+a)^2)^(1/2)*polylog(2,I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.44

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \frac{i\sqrt{1 + (a + bx)^2}(2 \arctan(e^{i \arctan(a+bx)}) \arctan(a + bx) - \text{PolyLog}(2, -ie^{i \arctan(a+bx)}) + \text{PolyLog}(2, ie^{i \arctan(a+bx)}))}{b\sqrt{c(1 + (a + bx)^2)}}$$

input

```
Integrate[ArcTan[a + b*x]/Sqrt[(1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2], x]
```

output

```
((-I)*Sqrt[1 + (a + b*x)^2]*(2*ArcTan[E^(I*ArcTan[a + b*x]])*ArcTan[a + b*x] - PolyLog[2, (-I)*E^(I*ArcTan[a + b*x]]) + PolyLog[2, I*E^(I*ArcTan[a + b*x])])/(b*Sqrt[c*(1 + (a + b*x)^2)])
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.72, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {5578, 5425, 5421}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\arctan(a + bx)}{\sqrt{(a^2 + 1)c + 2abcx + b^2cx^2}} dx \\ & \quad \downarrow \text{5578} \\ & \frac{\int \frac{\arctan(a+bx)}{\sqrt{c(a+bx)^2+c}} d(a + bx)}{b} \\ & \quad \downarrow \text{5425} \\ & \frac{\sqrt{(a + bx)^2 + 1} \int \frac{\arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a + bx)}{b\sqrt{c(a + bx)^2 + c}} \\ & \quad \downarrow \text{5421} \end{aligned}$$

$$\frac{\sqrt{(a+bx)^2+1} \left(-2i \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) + i \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) - i \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) \right)}{b\sqrt{c(a+bx)^2+c}}$$

input `Int[ArcTan[a + b*x]/Sqrt[(1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2],x]`

output `(Sqrt[1 + (a + b*x)^2]*((-2*I)*ArcTan[a + b*x]*ArcTan[Sqrt[1 + I*(a + b*x)]]/Sqrt[1 - I*(a + b*x)]) + I*PolyLog[2, ((-I)*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]] - I*PolyLog[2, (I*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]])/(b*Sqrt[c + c*(a + b*x)^2])`

Defintions of rubi rules used

rule 5421 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol]
:> Simp[-2*I*(a + b*ArcTan[c*x])*(ArcTan[Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x]]/(c*Sqrt[d])), x] + (Simp[I*b*(PolyLog[2, (-I)*(Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x])])/(c*Sqrt[d])), x] - Simp[I*b*(PolyLog[2, I*(Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x])])/(c*Sqrt[d])), x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && GtQ[d, 0]`

rule 5425 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol]
:> Simp[Sqrt[1 + c^2*x^2]/Sqrt[d + e*x^2] Int[(a + b*ArcTan[c*x])^p/Sqrt[1 + c^2*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0] && !GtQ[d, 0]`

rule 5578 `Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^p_*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^q_, x_Symbol]
:> Simp[1/d Subst[Int[(C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, A, B, C, p, q}, x] && EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.81

method	result
default	$-\frac{\left(\arctan(bx+a) \ln\left(1+\frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)-\arctan(bx+a) \ln\left(1-\frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)-i \operatorname{dilog}\left(1+\frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)+i \operatorname{dilog}\left(1-\frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)\right)}{\sqrt{b^2x^2+2abx+a^2+1}bc}$

input `int(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x,method=_RETURNVE
RBOSE)`

output `-(arctan(b*x+a)*ln(1+I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-arctan(b*x+a)*ln
(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-I*dilog(1+I*(1+I*(b*x+a)))/(1+(b*x+
a)^2)^(1/2))+I*dilog(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))*(c*(b*x+a-I)*
(I+a+b*x))^(1/2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)/b/c`

Fricas [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2cx^2 + 2abx + (a^2 + 1)c}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x, algorithm
="fricas")`

output `integral(arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \text{Timed out}$$

input `integrate(atan(b*x+a)/((a**2+1)*c+2*a*b*c*x+b**2*c*x**2)**(1/2),x)`

output Timed out

Maxima [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2cx^2 + 2abcx + (a^2 + 1)c}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Giac [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{\sqrt{b^2cx^2 + 2abcx + (a^2 + 1)c}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{\sqrt{cb^2x^2 + 2acbx + c(a^2 + 1)}} dx$$

input `int(atan(a + b*x)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/2), x)`

output `int(atan(a + b*x)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/2), x)`

Reduce [F]

$$\int \frac{\arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \frac{\int \frac{\operatorname{atan}(bx+a)}{\sqrt{b^2x^2+2abx+a^2+1}} dx}{\sqrt{c}}$$

input `int(atan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2), x)`

output `int(atan(a + b*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x)/sqrt(c)`

3.65
$$\int \frac{\arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx$$

Optimal result	554
Mathematica [B] (warning: unable to verify)	554
Rubi [N/A]	555
Maple [N/A]	556
Fricas [N/A]	556
Sympy [N/A]	556
Maxima [N/A]	557
Giac [N/A]	557
Mupad [N/A]	558
Reduce [N/A]	558

Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx = \text{Int}\left(\frac{\arctan(a+bx)}{\sqrt[3]{1+(a+bx)^2}}, x\right)$$

output `Defer(Int)(arctan(b*x+a)/(1+(b*x+a)^2)^(1/3), x)`

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 163 vs. 2(23) = 46.

Time = 0.29 (sec) , antiderivative size = 163, normalized size of antiderivative = 5.82

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx$$

$$= \frac{6 \Gamma\left(\frac{11}{6}\right) \Gamma\left(\frac{7}{3}\right) \left(15 + 10(a+bx) \arctan(a+bx) + \frac{4(a+bx) \arctan(a+bx) \text{Hypergeometric2F1}\left(1, \frac{4}{3}, \frac{11}{6}, \frac{1}{1+(a+bx)^2}\right)}{1+(a+bx)^2}\right)}{20b\sqrt[3]{1+a^2+2abx+b^2x^2} \Gamma\left(\frac{11}{6}\right) \Gamma\left(\frac{7}{3}\right)}$$

input `Integrate[ArcTan[a + b*x]/(1 + a^2 + 2*a*b*x + b^2*x^2)^(1/3), x]`

output

```
(6*Gamma[11/6]*Gamma[7/3]*(15 + 10*(a + b*x)*ArcTan[a + b*x] + (4*(a + b*x)
)*ArcTan[a + b*x]*Hypergeometric2F1[1, 4/3, 11/6, (1 + (a + b*x)^2)^(-1)])
/(1 + (a + b*x)^2)) + (5*2^(1/3)*Sqrt[Pi]*Gamma[5/3]*HypergeometricPFQ[{1,
4/3, 4/3}, {11/6, 7/3}, (1 + (a + b*x)^2)^(-1)]/(1 + (a + b*x)^2))/(20*b
*(1 + a^2 + 2*a*b*x + b^2*x^2)^(1/3)*Gamma[11/6]*Gamma[7/3])
```

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{a^2+2abx+b^2x^2+1}} dx$$

$$\downarrow \text{5578}$$

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{(a+bx)^2+1}} d(a+bx)$$

$$\frac{\phantom{\int \frac{\arctan(a+bx)}{\sqrt[3]{(a+bx)^2+1}} d(a+bx)}}{b}$$

$$\downarrow \text{5560}$$

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{(a+bx)^2+1}} d(a+bx)$$

$$\frac{\phantom{\int \frac{\arctan(a+bx)}{\sqrt[3]{(a+bx)^2+1}} d(a+bx)}}{b}$$

input

```
Int[ArcTan[a + b*x]/(1 + a^2 + 2*a*b*x + b^2*x^2)^(1/3), x]
```

output

```
$Aborted
```

Maple [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\arctan (bx+a)}{\left(b^2 x^2+2 a b x+a^2+1\right)^{\frac{1}{3}}} dx$$

input `int(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x)`

output `int(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\arctan (a+bx)}{\sqrt[3]{1+a^2+2 a b x+b^2 x^2}} dx = \int \frac{\arctan (bx+a)}{\left(b^2 x^2+2 a b x+a^2+1\right)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x, algorithm="fricas")`

output `integral(arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Sympy [N/A]

Not integrable

Time = 0.92 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{\arctan (a+bx)}{\sqrt[3]{1+a^2+2 a b x+b^2 x^2}} dx = \int \frac{\operatorname{atan}(a+bx)}{\sqrt[3]{a^2+2 a b x+b^2 x^2+1}} dx$$

input `integrate(atan(b*x+a)/(b**2*x**2+2*a*b*x+a**2+1)**(1/3),x)`

output `Integral(atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3), x)`

Maxima [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Mupad [N/A]

Not integrable

Time = 0.86 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{(a^2 + 2abx + b^2x^2 + 1)^{1/3}} dx$$

input `int(atan(a + b*x)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/3), x)`output `int(atan(a + b*x)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/3), x)`**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `int(atan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3), x)`output `int(atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3), x)`

3.66
$$\int \frac{\arctan(a+bx)}{\sqrt[3]{(1+a^2)c+2abcx+b^2cx^2}} dx$$

Optimal result	559
Mathematica [B] (warning: unable to verify)	559
Rubi [N/A]	560
Maple [N/A]	561
Fricas [N/A]	561
Sympy [N/A]	561
Maxima [N/A]	562
Giac [N/A]	562
Mupad [N/A]	563
Reduce [N/A]	563

Optimal result

Integrand size = 33, antiderivative size = 33

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{(1+a^2)c+2abcx+b^2cx^2}} dx = \text{Int}\left(\frac{\arctan(a+bx)}{\sqrt[3]{c+c(a+bx)^2}}, x\right)$$

output `Defer(Int)(arctan(b*x+a)/(c+c*(b*x+a)^2)^(1/3), x)`

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 165 vs. 2(25) = 50.

Time = 0.15 (sec) , antiderivative size = 165, normalized size of antiderivative = 5.00

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{(1+a^2)c+2abcx+b^2cx^2}} dx$$

$$= \frac{6 \Gamma\left(\frac{11}{6}\right) \Gamma\left(\frac{7}{3}\right) \left(15 + 10(a+bx) \arctan(a+bx) + \frac{4(a+bx) \arctan(a+bx) \text{Hypergeometric2F1}\left(1, \frac{4}{3}, \frac{11}{6}, \frac{1}{1+(a+bx)^2}\right)}{1+(a+bx)^2}\right)}{20b \sqrt[3]{c(1+a^2+2abx+b^2x^2)} \Gamma\left(\frac{11}{6}\right) \Gamma\left(\frac{7}{3}\right)}$$

input `Integrate[ArcTan[a + b*x]/((1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2)^(1/3), x]`

output

```
(6*Gamma[11/6]*Gamma[7/3]*(15 + 10*(a + b*x)*ArcTan[a + b*x] + (4*(a + b*x)
)*ArcTan[a + b*x]*Hypergeometric2F1[1, 4/3, 11/6, (1 + (a + b*x)^2)^(-1)])
/(1 + (a + b*x)^2)) + (5*2^(1/3)*Sqrt[Pi]*Gamma[5/3]*HypergeometricPFQ[{1,
4/3, 4/3}, {11/6, 7/3}, (1 + (a + b*x)^2)^(-1)]/(1 + (a + b*x)^2))/(20*b
*(c*(1 + a^2 + 2*a*b*x + b^2*x^2))^(1/3)*Gamma[11/6]*Gamma[7/3])
```

Rubi [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(a^2 + 1)c + 2abcx + b^2cx^2}} dx$$

$$\downarrow \text{5578}$$

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{c(a+bx)^2 + c}} d(a+bx)$$

$$\frac{\phantom{\int \frac{\arctan(a+bx)}{\sqrt[3]{c(a+bx)^2 + c}} d(a+bx)}}{b}$$

$$\downarrow \text{5560}$$

$$\int \frac{\arctan(a+bx)}{\sqrt[3]{c(a+bx)^2 + c}} d(a+bx)$$

$$\frac{\phantom{\int \frac{\arctan(a+bx)}{\sqrt[3]{c(a+bx)^2 + c}} d(a+bx)}}{b}$$

input

```
Int[ArcTan[a + b*x]/((1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2)^(1/3), x]
```

output

```
$Aborted
```

Maple [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{\arctan(bx + a)}{((a^2 + 1)c + 2abcx + b^2cx^2)^{\frac{1}{3}}} dx$$

input `int(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x)`

output `int(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x, algorithm="fricas")`

output `integral(arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c)^(1/3), x)`

Sympy [N/A]

Not integrable

Time = 7.69 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{\sqrt[3]{c(a^2 + 2abx + b^2x^2 + 1)}} dx$$

input `integrate(atan(b*x+a)/((a**2+1)*c+2*a*b*c*x+b**2*c*x**2)**(1/3),x)`

output `Integral(atan(a + b*x)/(c*(a**2 + 2*a*b*x + b**2*x**2 + 1))**(1/3), x)`

Maxima [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x, algorithm="maxima")`

output `integrate(arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c)^(1/3), x)`

Giac [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input `integrate(arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x, algorithm="giac")`

output `integrate(arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c)^(1/3), x)`

Mupad [N/A]

Not integrable

Time = 0.88 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\operatorname{atan}(a + bx)}{(cb^2x^2 + 2acbx + c(a^2 + 1))^{1/3}} dx$$

input `int(atan(a + b*x)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/3), x)`output `int(atan(a + b*x)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/3), x)`**Reduce [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{\arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \frac{\int \frac{\operatorname{atan}(bx+a)}{(b^2x^2+2abx+a^2+1)^{\frac{1}{3}}} dx}{c^{\frac{1}{3}}}$$

input `int(atan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3), x)`output `int(atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3), x)/c**(1/3)`

$$3.67 \quad \int \frac{a+b \arctan(c+dx)}{e+fx^2+gx^4} dx$$

Optimal result	564
Mathematica [A] (warning: unable to verify)	565
Rubi [A] (warning: unable to verify)	566
Maple [C] (warning: unable to verify)	569
Fricas [F]	570
Sympy [F(-1)]	570
Maxima [F]	570
Giac [F]	571
Mupad [F(-1)]	571
Reduce [F]	571

Optimal result

Integrand size = 25, antiderivative size = 1199

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \text{Too large to display}$$

output

```

-1/2*g^(1/2)*(a+b*arctan(d*x+c))*ln(-2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)-g^(
(1/2)*x^2^(1/2)))/(2^(1/2)*(I-c)*g^(1/2)-d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(
1-I*(d*x+c)))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)+1/2
*g^(1/2)*(a+b*arctan(d*x+c))*ln(-2*d*((-f+(-4*e*g+f^2)^(1/2))^(1/2)-g^(1/2
)*x^2^(1/2)))/(2^(1/2)*(I-c)*g^(1/2)-d*(-f+(-4*e*g+f^2)^(1/2))^(1/2))/(1-I*
(d*x+c)))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f+(-4*e*g+f^2)^(1/2))^(1/2)+1/2*g^(
1/2)*(a+b*arctan(d*x+c))*ln(2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)+g^(1/2)*x^2
^(1/2)))/(2^(1/2)*(I-c)*g^(1/2)+d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(1-I*(d*x+
c)))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)-1/2*g^(1/2)*
(a+b*arctan(d*x+c))*ln(2*d*((-f+(-4*e*g+f^2)^(1/2))^(1/2)+g^(1/2)*x^2^(1/2
)))/(2^(1/2)*(I-c)*g^(1/2)+d*(-f+(-4*e*g+f^2)^(1/2))^(1/2))/(1-I*(d*x+c)))*
2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f+(-4*e*g+f^2)^(1/2))^(1/2)+1/4*I*b*g^(1/2)*p
olylog(2,1+2*d*((-f-(-4*e*g+f^2)^(1/2))^(1/2)-g^(1/2)*x^2^(1/2)))/(2^(1/2)*
(I-c)*g^(1/2)-d*(-f-(-4*e*g+f^2)^(1/2))^(1/2))/(1-I*(d*x+c)))*2^(1/2)/(-4*
e*g+f^2)^(1/2)/(-f-(-4*e*g+f^2)^(1/2))^(1/2)-1/4*I*b*g^(1/2)*polylog(2,1+2
*d*((-f+(-4*e*g+f^2)^(1/2))^(1/2)-g^(1/2)*x^2^(1/2)))/(2^(1/2)*(I-c)*g^(1/2
)-d*(-f+(-4*e*g+f^2)^(1/2))^(1/2))/(1-I*(d*x+c)))*2^(1/2)/(-4*e*g+f^2)^(1/
2)/(-f+(-4*e*g+f^2)^(1/2))^(1/2)-1/4*I*b*g^(1/2)*polylog(2,1-2*d*((-f-(-4*
e*g+f^2)^(1/2))^(1/2)+g^(1/2)*x^2^(1/2)))/(2^(1/2)*(I-c)*g^(1/2)+d*(-f-(-4*
e*g+f^2)^(1/2))^(1/2))/(1-I*(d*x+c)))*2^(1/2)/(-4*e*g+f^2)^(1/2)/(-f-(-...

```

Mathematica [A] (warning: unable to verify)

Time = 5.25 (sec) , antiderivative size = 1807, normalized size of antiderivative = 1.51

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \text{Too large to display}$$

input

```
Integrate[(a + b*ArcTan[c + d*x])/(e + f*x^2 + g*x^4),x]
```

output

```
(Sqrt[g]*((4*a*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f - Sqrt[f^2 - 4*e*g]])/Sqrt[f - Sqrt[f^2 - 4*e*g]] - (4*a*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f + Sqrt[f^2 - 4*e*g]])/Sqrt[f + Sqrt[f^2 - 4*e*g]] + (I*b*Log[1 + I*c + I*d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(-I + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])]/Sqrt[-f - Sqrt[f^2 - 4*e*g]] - (I*b*Log[(-I)*(I + c + d*x)]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])]/Sqrt[-f - Sqrt[f^2 - 4*e*g]] - (I*b*Log[1 + I*c + I*d*x]*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(-I + c)*Sqrt[g] + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])]/Sqrt[-f + Sqrt[f^2 - 4*e*g]] + (I*b*Log[(-I)*(I + c + d*x)]*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I + c)*Sqrt[g] + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])]/Sqrt[-f + Sqrt[f^2 - 4*e*g]] - (I*b*Log[1 + I*c + I*d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(-(Sqrt[2]*(-I + c)*Sqrt[g]) + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])]/Sqrt[-f - Sqrt[f^2 - 4*e*g]] + (I*b*Log[(-I)*(I + c + d*x)]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(-(Sqrt[2]*(I + c)*Sqrt[g]) + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])]/Sqrt[-f - Sqrt[f^2 - 4*e*g]] + (I*b*Log[1 + I*c + I*d*x]*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(-(Sqrt[2]*(-I + c)*Sqrt[g]) + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])]/Sqrt[-f + Sqrt[f^2 - 4*e*g]] - (I*b*Log[(-I)*(I + c + ...
```

Rubi [A] (warning: unable to verify)

Time = 4.78 (sec) , antiderivative size = 2244, normalized size of antiderivative = 1.87, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {7279, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx$$

↓ 7279

$$\int \left(\frac{a}{e + fx^2 + gx^4} + \frac{b \arctan(c + dx)}{e + fx^2 + gx^4} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{\sqrt{2a}\sqrt{g} \arctan\left(\frac{\sqrt{2}\sqrt{gx}}{\sqrt{f-\sqrt{f^2-4eg}}}\right)}{\sqrt{f^2-4eg}\sqrt{f-\sqrt{f^2-4eg}}} - \frac{\sqrt{2a}\sqrt{g} \arctan\left(\frac{\sqrt{2}\sqrt{gx}}{\sqrt{f+\sqrt{f^2-4eg}}}\right)}{\sqrt{f^2-4eg}\sqrt{f+\sqrt{f^2-4eg}}} + \\
& \frac{ib\sqrt{g} \log(ic+idx+1) \log\left(-\frac{d\left(\sqrt{-f-\sqrt{f^2-4eg}}-\sqrt{2}\sqrt{gx}\right)}{\sqrt{2}(i-c)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \log(-ic-idx+1) \log\left(\frac{d\left(\sqrt{-f-\sqrt{f^2-4eg}}-\sqrt{2}\sqrt{gx}\right)}{\sqrt{2}\sqrt{g}(c+i)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \log(ic+idx+1) \log\left(-\frac{d\left(\sqrt{f^2-4eg}-f-\sqrt{2}\sqrt{gx}\right)}{\sqrt{2}(i-c)\sqrt{g}-d\sqrt{f^2-4eg}-f}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
& \frac{ib\sqrt{g} \log(-ic-idx+1) \log\left(\frac{d\left(\sqrt{f^2-4eg}-f-\sqrt{2}\sqrt{gx}\right)}{\sqrt{2}\sqrt{g}(c+i)+d\sqrt{f^2-4eg}-f}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
& \frac{ib\sqrt{g} \log(-ic-idx+1) \log\left(-\frac{d\left(\sqrt{2}\sqrt{gx}+\sqrt{-f-\sqrt{f^2-4eg}}\right)}{\sqrt{2}(c+i)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \log(ic+idx+1) \log\left(\frac{d\left(\sqrt{2}\sqrt{gx}+\sqrt{-f-\sqrt{f^2-4eg}}\right)}{\sqrt{2}\sqrt{g}(i-c)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \log(-ic-idx+1) \log\left(-\frac{d\left(\sqrt{2}\sqrt{gx}+\sqrt{\sqrt{f^2-4eg}-f}\right)}{\sqrt{2}(c+i)\sqrt{g}-d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
& \frac{ib\sqrt{g} \log(ic+idx+1) \log\left(\frac{d\left(\sqrt{2}\sqrt{gx}+\sqrt{\sqrt{f^2-4eg}-f}\right)}{\sqrt{2}\sqrt{g}(i-c)+d\sqrt{\sqrt{f^2-4eg}-f}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{\sqrt{f^2-4eg}-f}} + \\
& \frac{ib\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+i)}{\sqrt{2}(i-c)\sqrt{g}-d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+i)}{\sqrt{2}\sqrt{g}(i-c)+d\sqrt{-f-\sqrt{f^2-4eg}}}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} - \\
& \frac{ib\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+i)}{\sqrt{2}\sqrt{g}(-c-dx+i)}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}} + \frac{ib\sqrt{g} \operatorname{PolyLog}\left(2, \frac{\sqrt{2}\sqrt{g}(-c-dx+i)}{\sqrt{2}\sqrt{g}(-c-dx+i)}\right)}{2\sqrt{2}\sqrt{f^2-4eg}\sqrt{-f-\sqrt{f^2-4eg}}}
\end{aligned}$$

input `Int[(a + b*ArcTan[c + d*x])/(e + f*x^2 + g*x^4),x]`

output `(Sqrt[2]*a*Sqrt[g]*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f - Sqrt[f^2 - 4*e*g]])/(Sqrt[f^2 - 4*e*g]*Sqrt[f - Sqrt[f^2 - 4*e*g]]) - (Sqrt[2]*a*Sqrt[g]*ArcTan[(Sqrt[2]*Sqrt[g]*x)/Sqrt[f + Sqrt[f^2 - 4*e*g]])/(Sqrt[f^2 - 4*e*g]*Sqrt[f + Sqrt[f^2 - 4*e*g]]) + ((I/2)*b*Sqrt[g]*Log[1 + I*c + I*d*x]*Log[-(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I - c)*Sqrt[g] - d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])])/(Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - ((I/2)*b*Sqrt[g]*Log[1 - I*c - I*d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I + c)*Sqrt[g] + d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])])/(Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - ((I/2)*b*Sqrt[g]*Log[1 + I*c + I*d*x]*Log[-(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I - c)*Sqrt[g] - d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])])/(Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f + Sqrt[f^2 - 4*e*g]]) + ((I/2)*b*Sqrt[g]*Log[1 - I*c - I*d*x]*Log[(d*(Sqrt[-f + Sqrt[f^2 - 4*e*g]] - Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I + c)*Sqrt[g] + d*Sqrt[-f + Sqrt[f^2 - 4*e*g]])])/(Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f + Sqrt[f^2 - 4*e*g]]) + ((I/2)*b*Sqrt[g]*Log[1 - I*c - I*d*x]*Log[-(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I + c)*Sqrt[g] - d*Sqrt[-f - Sqrt[f^2 - 4*e*g]])])/(Sqrt[2]*Sqrt[f^2 - 4*e*g]*Sqrt[-f - Sqrt[f^2 - 4*e*g]]) - ((I/2)*b*Sqrt[g]*Log[1 + I*c + I*d*x]*Log[(d*(Sqrt[-f - Sqrt[f^2 - 4*e*g]] + Sqrt[2]*Sqrt[g]*x))/(Sqrt[2]*(I - c)*Sqrt[g] + d*Sqrt[...`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7279 `Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.03 (sec) , antiderivative size = 981, normalized size of antiderivative = 0.82

method	result	size
risch	Expression too large to display	981
parts	Expression too large to display	1831
derivativdivides	Expression too large to display	1838
default	Expression too large to display	1838

input `int((a+b*arctan(d*x+c))/(g*x^4+f*x^2+e),x,method=_RETURNVERBOSE)`

output

```
-1/4*d^3*b*sum(1/(6*I*_R1^2*c*g-2*I*c^3*g-I*c*d^2*f-12*I*_R1*c*g+2*_R1^3*g
-6*_R1*c^2*g-_R1*d^2*f+6*I*c*g-6*_R1^2*g+6*c^2*g+d^2*f+6*_R1*g-2*g)*(ln(1-
I*c-I*d*x)*ln((_R1+I*d*x+I*c-1)/_R1)+dilog((_R1+I*d*x+I*c-1)/_R1)),_R1=Ro
otOf(g*_Z^4+(4*RootOf(_Z^2+1,index=1)*c*g-4*g)*_Z^3+(-12*RootOf(_Z^2+1,inde
x=1)*c*g-6*c^2*g-d^2*f+6*g)*_Z^2+(-4*RootOf(_Z^2+1,index=1)*c^3*g-2*RootOf
(_Z^2+1,index=1)*c*d^2*f+12*RootOf(_Z^2+1,index=1)*c*g+12*c^2*g+2*d^2*f-4*
g)*_Z+4*RootOf(_Z^2+1,index=1)*c^3*g+2*RootOf(_Z^2+1,index=1)*c*d^2*f+c^4*
g+c^2*d^2*f+e*d^4-4*RootOf(_Z^2+1,index=1)*c*g-6*c^2*g-d^2*f+g))+1/2*I*d^3
*a*sum(1/(6*I*_R^2*c*g-2*I*c^3*g-I*c*d^2*f-12*I*_R*c*g+2*_R^3*g-6*_R*c^2*g
-_R*d^2*f+6*I*c*g-6*g*_R^2+6*c^2*g+d^2*f+6*g*_R-2*g)*ln(-I*d*x-I*c+1-_R),_
R=RootOf(g*_Z^4+(4*RootOf(_Z^2+1,index=1)*c*g-4*g)*_Z^3+(-12*RootOf(_Z^2+1
,index=1)*c*g-6*c^2*g-d^2*f+6*g)*_Z^2+(-4*RootOf(_Z^2+1,index=1)*c^3*g-2*R
ootOf(_Z^2+1,index=1)*c*d^2*f+12*RootOf(_Z^2+1,index=1)*c*g+12*c^2*g+2*d^2
*f-4*g)*_Z+4*RootOf(_Z^2+1,index=1)*c^3*g+2*RootOf(_Z^2+1,index=1)*c*d^2*f
+c^4*g+c^2*d^2*f+e*d^4-4*RootOf(_Z^2+1,index=1)*c*g-6*c^2*g-d^2*f+g))-1/4*
b*d^3*sum(1/(-6*I*_R1^2*c*g+2*I*c^3*g+I*c*d^2*f+2*_R1^3*g-6*_R1*c^2*g+12*I
*_R1*c*g-_R1*d^2*f-6*_R1^2*g+6*c^2*g-6*I*c*g+d^2*f+6*_R1*g-2*g)*(ln(1+I*c+
I*d*x)*ln((_R1-I*d*x-I*c-1)/_R1)+dilog((_R1-I*d*x-I*c-1)/_R1)),_R1=RootOf(
g*_Z^4+(-4*RootOf(_Z^2+1,index=1)*c*g-4*g)*_Z^3+(-6*c^2*g+12*RootOf(_Z^2+1
,index=1)*c*g-d^2*f+6*g)*_Z^2+(4*RootOf(_Z^2+1,index=1)*c^3*g+2*RootOf(...
```

Fricas [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \arctan(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="fricas")`

output `integral((b*arctan(d*x + c) + a)/(g*x^4 + f*x^2 + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \text{Timed out}$$

input `integrate((a+b*atan(d*x+c))/(g*x**4+f*x**2+e),x)`

output `Timed out`

Maxima [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \arctan(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="maxima")`

output `integrate((b*arctan(d*x + c) + a)/(g*x^4 + f*x^2 + e), x)`

Giac [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{b \arctan(dx + c) + a}{gx^4 + fx^2 + e} dx$$

input `integrate((a+b*arctan(d*x+c))/(g*x^4+f*x^2+e),x, algorithm="giac")`

output `integrate((b*arctan(d*x + c) + a)/(g*x^4 + f*x^2 + e), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx = \int \frac{a + b \operatorname{atan}(c + dx)}{gx^4 + fx^2 + e} dx$$

input `int((a + b*atan(c + d*x))/(e + f*x^2 + g*x^4),x)`

output `int((a + b*atan(c + d*x))/(e + f*x^2 + g*x^4), x)`

Reduce [F]

$$\int \frac{a + b \arctan(c + dx)}{e + fx^2 + gx^4} dx$$

$$= \frac{2\sqrt{e} \sqrt{2\sqrt{g}\sqrt{e+f}} \operatorname{atan}\left(\frac{\sqrt{2\sqrt{g}\sqrt{e-f-2\sqrt{g}x}}}{\sqrt{2\sqrt{g}\sqrt{e+f}}}\right) af - 4\sqrt{g} \sqrt{2\sqrt{g}\sqrt{e+f}} \operatorname{atan}\left(\frac{\sqrt{2\sqrt{g}\sqrt{e-f-2\sqrt{g}x}}}{\sqrt{2\sqrt{g}\sqrt{e+f}}}\right) ae - 2\sqrt{e}}{\dots}$$

input `int((a+b*atan(d*x+c))/(g*x^4+f*x^2+e),x)`

output

```
(2*sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) -
2*sqrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*f - 4*sqrt(g)*sqrt(2*sqrt(g)*
sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) - 2*sqrt(g)*x)/sqrt(2*sqrt(
g)*sqrt(e) + f))*a*e - 2*sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*
sqrt(g)*sqrt(e) - f) + 2*sqrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*f + 4*s
qrt(g)*sqrt(2*sqrt(g)*sqrt(e) + f)*atan((sqrt(2*sqrt(g)*sqrt(e) - f) + 2*s
qrt(g)*x)/sqrt(2*sqrt(g)*sqrt(e) + f))*a*e - sqrt(e)*sqrt(2*sqrt(g)*sqrt(e
) - f)*log( - sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + sqrt(g)*x**2)*a*f
+ sqrt(e)*sqrt(2*sqrt(g)*sqrt(e) - f)*log(sqrt(2*sqrt(g)*sqrt(e) - f)*x +
sqrt(e) + sqrt(g)*x**2)*a*f - 2*sqrt(g)*sqrt(2*sqrt(g)*sqrt(e) - f)*log( -
sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + sqrt(g)*x**2)*a*e + 2*sqrt(g)*s
qrt(2*sqrt(g)*sqrt(e) - f)*log(sqrt(2*sqrt(g)*sqrt(e) - f)*x + sqrt(e) + s
qrt(g)*x**2)*a*e + 16*int(atan(c + d*x)/(e + f*x**2 + g*x**4),x)*b*e**2*g
- 4*int(atan(c + d*x)/(e + f*x**2 + g*x**4),x)*b*e*f**2)/(4*e*(4*e*g - f**
2))
```

$$3.68 \quad \int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx$$

Optimal result	573
Mathematica [A] (verified)	574
Rubi [A] (verified)	574
Maple [A] (verified)	576
Fricas [F]	576
Sympy [F(-1)]	577
Maxima [F]	577
Giac [F]	577
Mupad [F(-1)]	578
Reduce [F]	578

Optimal result

Integrand size = 35, antiderivative size = 187

$$\begin{aligned} \int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx = & -\frac{\sqrt{1+(a+bx)^2}}{2b} \\ & + \frac{(a+bx)\sqrt{1+(a+bx)^2} \arctan(a+bx)}{2b} \\ & + \frac{i \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b} \\ & - \frac{i \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{2b} \\ & + \frac{i \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{2b} \end{aligned}$$

output

```
-1/2*(1+(b*x+a)^2)^(1/2)/b+1/2*(b*x+a)*(1+(b*x+a)^2)^(1/2)*arctan(b*x+a)/b
+I*arctan(b*x+a)*arctan((1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b-1/2*I*
polylog(2,-I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b+1/2*I*polylog(2,I*(
1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b
```

Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.78

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{1+a^2+2abx+b^2x^2}} dx$$

$$= \frac{-\sqrt{1+(a+bx)^2} + (a+bx)\sqrt{1+(a+bx)^2} \arctan(a+bx) - \arctan(a+bx) \log(1 - ie^{i \arctan(a+bx)})}{2b}$$

input

```
Integrate[((a + b*x)^2*ArcTan[a + b*x])/Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2],
x]
```

output

```
(-Sqrt[1 + (a + b*x)^2] + (a + b*x)*Sqrt[1 + (a + b*x)^2]*ArcTan[a + b*x]
- ArcTan[a + b*x]*Log[1 - I*E^(I*ArcTan[a + b*x])] + ArcTan[a + b*x]*Log[1
+ I*E^(I*ArcTan[a + b*x])] - I*PolyLog[2, (-I)*E^(I*ArcTan[a + b*x])] + I
*PolyLog[2, I*E^(I*ArcTan[a + b*x])])/(2*b)
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.95, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.114$, Rules used = {5580, 5487, 241, 5421}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{a^2+2abx+b^2x^2+1}} dx$$

$$\downarrow \text{5580}$$

$$\frac{\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a+bx)}{b}$$

$$\downarrow \text{5487}$$

$$\frac{-\frac{1}{2} \int \frac{\arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a+bx) - \frac{1}{2} \int \frac{a+bx}{\sqrt{(a+bx)^2+1}} d(a+bx) + \frac{1}{2} (a+bx) \sqrt{(a+bx)^2+1} \arctan(a+bx)}{b}$$

$$\frac{-\frac{1}{2} \int \frac{\arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a+bx) + \frac{1}{2}(a+bx)\sqrt{(a+bx)^2+1} \arctan(a+bx) - \frac{1}{2}\sqrt{(a+bx)^2+1}}{b}$$

↓ 241

↓ 5421

$$\frac{\frac{1}{2} \left(2i \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) - i \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) + i \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) \right) + \frac{1}{2}(a+bx)}{b}$$

input `Int[((a + b*x)^2*ArcTan[a + b*x])/Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2], x]`

output `(-1/2*Sqrt[1 + (a + b*x)^2] + ((a + b*x)*Sqrt[1 + (a + b*x)^2]*ArcTan[a + b*x])/2 + ((2*I)*ArcTan[a + b*x]*ArcTan[Sqrt[1 + I*(a + b*x)]/Sqrt[1 - I*(a + b*x)]] - I*PolyLog[2, ((-I)*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]] + I*PolyLog[2, (I*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]])/2)/b`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5421 `Int[((a_) + ArcTan[(c_)*(x_)])*(b_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[-2*I*(a + b*ArcTan[c*x])*(ArcTan[Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x]]/(c*Sqrt[d])), x] + (Simp[I*b*(PolyLog[2, (-I)*(Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x])])/(c*Sqrt[d])), x] - Simp[I*b*(PolyLog[2, I*(Sqrt[1 + I*c*x]/Sqrt[1 - I*c*x])])/(c*Sqrt[d])), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && GtQ[d, 0]`

rule 5487 `Int[(((a_) + ArcTan[(c_)*(x_)])*(b_))^(p_)*((f_)*(x_)^(m_))/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[f*(f*x)^(m - 1)*Sqrt[d + e*x^2]*((a + b*ArcTan[c*x])^p/(c^2*d*m)), x] + (-Simp[b*f*(p/(c*m)) Int[(f*x)^(m - 1)*((a + b*ArcTan[c*x])^(p - 1)/Sqrt[d + e*x^2]), x], x] - Simp[f^2*((m - 1)/(c^2*m)) Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/Sqrt[d + e*x^2]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[e, c^2*d] && GtQ[p, 0] && GtQ[m, 1]`

rule 5580

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.)*((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)^(q_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(C/d^2 + (C/d^2)*x^2)^q*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x] && EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]
```

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.96

method	result
default	$\frac{(\arctan(bx+a)bx+a \arctan(bx+a)-1)\sqrt{b^2x^2+2abx+a^2+1}}{2b} + \frac{\arctan(bx+a) \ln\left(1 + \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right) - \arctan(bx+a) \ln\left(1 - \frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right)}{2b}$

input

```
int((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x,method=_RETURN
VERBOSE)
```

output

```
1/2*(arctan(b*x+a)*b*x+a*arctan(b*x+a)-1)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)/b+
1/2*(arctan(b*x+a)*ln(1+I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-arctan(b*x+a)
*ln(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-I*dilog(1+I*(1+I*(b*x+a)))/(1+(b
*x+a)^2)^(1/2))+I*dilog(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))/b
```

Fricas [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input

```
integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorit
hm="fricas")
```

output

```
integral((b^2*x^2 + 2*a*b*x + a^2)*arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x
+ a^2 + 1), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \text{Timed out}$$

input `integrate((b*x+a)**2*atan(b*x+a)/(b**2*x**2+2*a*b*x+a**2+1)**(1/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*x + a)^2*arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1), x)`

Giac [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*x + a)^2*arctan(b*x + a)/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(a + bx) (a + bx)^2}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx$$

input `int((atan(a + b*x)*(a + b*x)^2)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2), x)`

output `int((atan(a + b*x)*(a + b*x)^2)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2), x)`

Reduce [F]

$$\begin{aligned} \int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{1 + a^2 + 2abx + b^2x^2}} dx &= \left(\int \frac{\operatorname{atan}(bx + a)}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx \right) a^2 \\ &+ \left(\int \frac{\operatorname{atan}(bx + a) x^2}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx \right) b^2 \\ &+ 2 \left(\int \frac{\operatorname{atan}(bx + a) x}{\sqrt{b^2x^2 + 2abx + a^2 + 1}} dx \right) ab \end{aligned}$$

input `int((b*x+a)^2*atan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2), x)`

output `int(atan(a + b*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x)*a**2 + int((atan(a + b*x)*x**2)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x)*b**2 + 2*int((atan(a + b*x)*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x)*a*b`

3.69
$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx$$

Optimal result	579
Mathematica [A] (verified)	580
Rubi [A] (verified)	580
Maple [A] (verified)	583
Fricas [F]	583
Sympy [F(-1)]	584
Maxima [F]	584
Giac [F]	584
Mupad [F(-1)]	585
Reduce [F]	585

Optimal result

Integrand size = 40, antiderivative size = 281

$$\begin{aligned} & \int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx \\ &= -\frac{\sqrt{c+c(a+bx)^2}}{2bc} + \frac{(a+bx)\sqrt{c+c(a+bx)^2} \arctan(a+bx)}{2bc} \\ & \quad + \frac{i\sqrt{1+(a+bx)^2} \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{b\sqrt{c+c(a+bx)^2}} \\ & \quad - \frac{i\sqrt{1+(a+bx)^2} \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{2b\sqrt{c+c(a+bx)^2}} \\ & \quad + \frac{i\sqrt{1+(a+bx)^2} \operatorname{PolyLog}\left(2, \frac{i\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right)}{2b\sqrt{c+c(a+bx)^2}} \end{aligned}$$

output

```
-1/2*(c+c*(b*x+a)^2)^(1/2)/b/c+1/2*(b*x+a)*(c+c*(b*x+a)^2)^(1/2)*arctan(b*x+a)/b/c+I*(1+(b*x+a)^2)^(1/2)*arctan(b*x+a)*arctan((1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)-1/2*I*(1+(b*x+a)^2)^(1/2)*polylog(2,-I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)+1/2*I*(1+(b*x+a)^2)^(1/2)*polylog(2,I*(1+I*(b*x+a))^(1/2)/(1-I*(b*x+a))^(1/2))/b/(c+c*(b*x+a)^2)^(1/2)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.67

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx$$

$$= \frac{\sqrt{1 + a^2 + 2abx + b^2x^2} \left(-\sqrt{1 + (a + bx)^2} + (a + bx)\sqrt{1 + (a + bx)^2} \arctan(a + bx) - \arctan(a + bx) \right)}{2b\sqrt{1 + a^2 + 2abx + b^2x^2}}$$

input

```
Integrate[((a + b*x)^2*ArcTan[a + b*x])/Sqrt[(1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2],x]
```

output

```
(Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]*(-Sqrt[1 + (a + b*x)^2] + (a + b*x)*Sqrt[1 + (a + b*x)^2]*ArcTan[a + b*x] - ArcTan[a + b*x]*Log[1 - I*E^(I*ArcTan[a + b*x])]) + ArcTan[a + b*x]*Log[1 + I*E^(I*ArcTan[a + b*x])] - I*PolyLog[2, (-I)*E^(I*ArcTan[a + b*x])] + I*PolyLog[2, I*E^(I*ArcTan[a + b*x])])/(2*b*Sqrt[c*(1 + a^2 + 2*a*b*x + b^2*x^2)])
```

Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.77, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5580, 5487, 241, 5425, 5421}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(a^2 + 1)c + 2abcx + b^2cx^2}} dx$$

$$\downarrow \text{5580}$$

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{c(a+bx)^2+c}} d(a + bx)$$

$$\frac{\phantom{\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{c(a+bx)^2+c}} d(a + bx)}}{b}$$

$$\downarrow \text{5487}$$

$$\begin{aligned}
 & \frac{-\frac{1}{2} \int \frac{\arctan(a+bx)}{\sqrt{c(a+bx)^2+c}} d(a+bx) - \frac{1}{2} \int \frac{a+bx}{\sqrt{c(a+bx)^2+c}} d(a+bx) + \frac{(a+bx) \arctan(a+bx) \sqrt{c(a+bx)^2+c}}{2c}}{b} \\
 & \quad \downarrow 241 \\
 & \frac{-\frac{1}{2} \int \frac{\arctan(a+bx)}{\sqrt{c(a+bx)^2+c}} d(a+bx) + \frac{(a+bx) \arctan(a+bx) \sqrt{c(a+bx)^2+c}}{2c} - \frac{\sqrt{c(a+bx)^2+c}}{2c}}{b} \\
 & \quad \downarrow 5425 \\
 & \frac{-\frac{\sqrt{(a+bx)^2+1} \int \frac{\arctan(a+bx)}{\sqrt{(a+bx)^2+1}} d(a+bx)}{2\sqrt{c(a+bx)^2+c}} + \frac{(a+bx) \arctan(a+bx) \sqrt{c(a+bx)^2+c}}{2c} - \frac{\sqrt{c(a+bx)^2+c}}{2c}}{b} \\
 & \quad \downarrow 5421 \\
 & \frac{-\frac{\sqrt{(a+bx)^2+1} \left(-2i \arctan(a+bx) \arctan\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}}\right) + i \operatorname{PolyLog}\left(2, -\frac{i\sqrt{i(a+bx)+1}}{\sqrt{1-i(a+bx)}}\right) - i \operatorname{PolyLog}\left(2, \frac{i\sqrt{i(a+bx)+1}}{\sqrt{1-i(a+bx)}}\right) \right)}{2\sqrt{c(a+bx)^2+c}} + \frac{(a+bx) \arctan(a+bx) \sqrt{c(a+bx)^2+c}}{2c}}{b}
 \end{aligned}$$

input

```
Int[((a + b*x)^2*ArcTan[a + b*x])/Sqrt[(1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2],x]
```

output

```
(-1/2*Sqrt[c + c*(a + b*x)^2]/c + ((a + b*x)*Sqrt[c + c*(a + b*x)^2]*ArcTan[a + b*x])/(2*c) - (Sqrt[1 + (a + b*x)^2]*((-2*I)*ArcTan[a + b*x]*ArcTan[Sqrt[1 + I*(a + b*x)]/Sqrt[1 - I*(a + b*x)]] + I*PolyLog[2, ((-I)*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]] - I*PolyLog[2, (I*Sqrt[1 + I*(a + b*x)])/Sqrt[1 - I*(a + b*x)]]))/(2*Sqrt[c + c*(a + b*x)^2])/b
```

Definitions of rubi rules used

rule 241 $\text{Int}[(x_*)*((a_*) + (b_*)*(x_*)^2)^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^2)^{(p + 1)}/(2*b*(p + 1)), x] /;$ FreeQ[{a, b, p}, x] && NeQ[p, -1]

rule 5421 $\text{Int}[(a_*) + \text{ArcTan}[(c_*)*(x_*)]*(b_*)]/\text{Sqrt}[(d_*) + (e_*)*(x_*)^2], x_Symbol] \rightarrow \text{Simp}[-2*I*(a + b*\text{ArcTan}[c*x])*(\text{ArcTan}[\text{Sqrt}[1 + I*c*x]/\text{Sqrt}[1 - I*c*x]]/(c*\text{Sqrt}[d])), x] + (\text{Simp}[I*b*(\text{PolyLog}[2, (-I)*(\text{Sqrt}[1 + I*c*x]/\text{Sqrt}[1 - I*c*x])]/(c*\text{Sqrt}[d])), x] - \text{Simp}[I*b*(\text{PolyLog}[2, I*(\text{Sqrt}[1 + I*c*x]/\text{Sqrt}[1 - I*c*x])]/(c*\text{Sqrt}[d])), x]) /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && GtQ[d, 0]

rule 5425 $\text{Int}[(a_*) + \text{ArcTan}[(c_*)*(x_*)]*(b_*)]^{(p_*)}/\text{Sqrt}[(d_*) + (e_*)*(x_*)^2], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[1 + c^2*x^2]/\text{Sqrt}[d + e*x^2] \text{Int}[(a + b*\text{ArcTan}[c*x])^p/\text{Sqrt}[1 + c^2*x^2], x], x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0] && !GtQ[d, 0]

rule 5487 $\text{Int}[(a_*) + \text{ArcTan}[(c_*)*(x_*)]*(b_*)]^{(p_*)}*((f_*)*(x_*)^m)/\text{Sqrt}[(d_*) + (e_*)*(x_*)^2], x_Symbol] \rightarrow \text{Simp}[f*(f*x)^{(m - 1)}*\text{Sqrt}[d + e*x^2]*((a + b*\text{ArcTan}[c*x])^p/(c^2*d*m)), x] + (-\text{Simp}[b*f*(p/(c*m)) \text{Int}[(f*x)^{(m - 1)}*((a + b*\text{ArcTan}[c*x])^{(p - 1)}/\text{Sqrt}[d + e*x^2]), x], x] - \text{Simp}[f^2*(m - 1)/(c^2*m) \text{Int}[(f*x)^{(m - 2)}*((a + b*\text{ArcTan}[c*x])^p/\text{Sqrt}[d + e*x^2]), x], x]) /;$ FreeQ[{a, b, c, d, e, f}, x] && EqQ[e, c^2*d] && GtQ[p, 0] && GtQ[m, 1]

rule 5580 $\text{Int}[(a_*) + \text{ArcTan}[(c_*) + (d_*)*(x_*)]*(b_*)]^{(p_*)}*((e_*) + (f_*)*(x_*)^m) * ((A_*) + (B_*)*(x_*) + (C_*)*(x_*)^2)^{(q_*)}, x_Symbol] \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(d*e - c*f)/d + f*(x/d)]^m*(C/d^2 + (C/d^2)*x^2)^q*(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, m, p, q}, x] && EqQ[B*(1 + c^2) - 2*A*c*d, 0] && EqQ[2*c*C - B*d, 0]

Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.79

method	result
default	$\frac{(\arctan(bx+a)bx+a \arctan(bx+a)-1)\sqrt{c(bx+a-i)(bx+a+i)}}{2bc} + \frac{(\arctan(bx+a) \ln\left(1+\frac{i(1+i(bx+a))}{\sqrt{1+(bx+a)^2}}\right) - \arctan(bx+a) \ln\left(1-\frac{i(1-i(bx+a))}{\sqrt{1+(bx+a)^2}}\right))}{2bc}$

input `int((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(arctan(b*x+a)*b*x+a*arctan(b*x+a)-1)*(c*(b*x+a-I)*(I+a+b*x))^(1/2)/b/c+1/2*(arctan(b*x+a)*ln(1+I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-arctan(b*x+a)*ln(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))-I*dilog(1+I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))+I*dilog(1-I*(1+I*(b*x+a)))/(1+(b*x+a)^2)^(1/2))*(c*(b*x+a-I)*(I+a+b*x))^(1/2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)/b/c`

Fricas [F]

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt{(1+a^2)c+2abcx+b^2cx^2}} dx = \int \frac{(bx+a)^2 \arctan(bx+a)}{\sqrt{b^2cx^2+2abcx+(a^2+1)c}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x,algorithm="fricas")`

output `integral((b^2*x^2 + 2*a*b*x + a^2)*arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \text{Timed out}$$

input `integrate((b*x+a)**2*atan(b*x+a)/((a**2+1)*c+2*a*b*c*x+b**2*c*x**2)**(1/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{\sqrt{b^2cx^2 + 2abcx + (a^2 + 1)c}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x,algorithm="maxima")`

output `integrate((b*x + a)^2*arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Giac [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{\sqrt{b^2cx^2 + 2abcx + (a^2 + 1)c}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x,algorithm="giac")`

output `integrate((b*x + a)^2*arctan(b*x + a)/sqrt(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\operatorname{atan}(a + bx) (a + bx)^2}{\sqrt{cb^2x^2 + 2acbx + c(a^2 + 1)}} dx$$

input `int((atan(a + b*x)*(a + b*x)^2)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/2),x)`

output `int((atan(a + b*x)*(a + b*x)^2)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/2), x)`

Reduce [F]

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt{(1 + a^2)c + 2abcx + b^2cx^2}} dx$$

$$= \frac{\left(\int \frac{\operatorname{atan}(bx+a)}{\sqrt{b^2x^2+2abx+a^2+1}} dx\right) a^2 + \left(\int \frac{\operatorname{atan}(bx+a)x^2}{\sqrt{b^2x^2+2abx+a^2+1}} dx\right) b^2 + 2\left(\int \frac{\operatorname{atan}(bx+a)x}{\sqrt{b^2x^2+2abx+a^2+1}} dx\right) ab}{\sqrt{c}}$$

input `int((b*x+a)^2*atan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/2),x)`

output `(int(atan(a + b*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1),x)*a**2 + int((atan(a + b*x)*x**2)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1),x)*b**2 + 2*int((atan(a + b*x)*x)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1),x)*a*b)/sqrt(c)`

3.70
$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx$$

Optimal result	586
Mathematica [B] (warning: unable to verify)	586
Rubi [N/A]	587
Maple [N/A]	588
Fricas [N/A]	588
Sympy [N/A]	588
Maxima [N/A]	589
Giac [N/A]	589
Mupad [N/A]	590
Reduce [N/A]	590

Optimal result

Integrand size = 35, antiderivative size = 35

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx = \text{Int} \left(\frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{1+(a+bx)^2}}, x \right)$$

output `Defer(Int)((b*x+a)^2*arctan(b*x+a)/(1+(b*x+a)^2)^(1/3),x)`

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 181 vs. 2(30) = 60.

Time = 4.38 (sec) , antiderivative size = 181, normalized size of antiderivative = 5.17

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{1+a^2+2abx+b^2x^2}} dx =$$

$$3(1+(a+bx)^2)^{2/3} \left(\frac{5\sqrt[3]{2}\sqrt{\pi} \Gamma(\frac{5}{3}) {}_3F_2(1, \frac{4}{3}, \frac{4}{3}; \frac{11}{6}, \frac{7}{3}; \frac{1}{1+(a+bx)^2})}{(1+(a+bx)^2)^2} + \Gamma(\frac{11}{6}) \Gamma(\frac{7}{3}) \left(15 + \frac{90}{1+(a+bx)^2} \right) \right)$$

140b Ga

input `Integrate[((a + b*x)^2*ArcTan[a + b*x])/(1 + a^2 + 2*a*b*x + b^2*x^2)^(1/3),x]`

output

$$\frac{(-3(1 + (a + bx)^2)^{2/3} * ((5 * 2^{1/3}) * \text{Sqrt}[\text{Pi}] * \text{Gamma}[5/3] * \text{HypergeometricPFQ}[\{1, 4/3, 4/3\}, \{11/6, 7/3\}, (1 + (a + bx)^2)^{-1}]) / (1 + (a + bx)^2)^2 + \text{Gamma}[11/6] * \text{Gamma}[7/3] * (15 + 90 / (1 + (a + bx)^2) + (24 * (a + bx) * \text{ArcTan}[a + bx] * \text{Hypergeometric2F1}[1, 4/3, 11/6, (1 + (a + bx)^2)^{-1}]) / (1 + (a + bx)^2)^2 + 5 * \text{ArcTan}[a + bx] * (-4 * (a + bx) + 6 * \text{Sin}[2 * \text{ArcTan}[a + bx]])) / (140 * b * \text{Gamma}[11/6] * \text{Gamma}[7/3])$$
Rubi [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{a^2 + 2abx + b^2x^2 + 1}} dx$$

$$\downarrow \text{5580}$$

$$\int \frac{(a+bx)^2 \arctan(a+bx) d(a + bx)}{\sqrt[3]{(a + bx)^2 + 1}}$$

$$\frac{b}{b}$$

$$\downarrow \text{5560}$$

$$\int \frac{(a+bx)^2 \arctan(a+bx) d(a + bx)}{\sqrt[3]{(a + bx)^2 + 1}}$$

$$\frac{b}{b}$$

input

$$\text{Int}[(a + bx)^2 * \text{ArcTan}[a + bx] / (1 + a^2 + 2 * a * b * x + b^2 * x^2)^{(1/3)}, x]$$

output

\$Aborted

Maple [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `int((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3), x)`

output `int((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3), x)`

Fricas [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3), x, algorithm="fricas")`

output `integral((b^2*x^2 + 2*a*b*x + a^2)*arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Sympy [N/A]

Not integrable

Time = 3.87 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(a + bx)^2 \operatorname{atan}(a + bx)}{\sqrt[3]{a^2 + 2abx + b^2x^2 + 1}} dx$$

input `integrate((b*x+a)**2*atan(b*x+a)/(b**2*x**2+2*a*b*x+a**2+1)**(1/3),x)`

output `Integral((a + b*x)**2*atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3), x)`

Maxima [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x, algorithm="maxima")`

output `integrate((b*x + a)^2*arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Giac [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x, algorithm="giac")`

output `integrate((b*x + a)^2*arctan(b*x + a)/(b^2*x^2 + 2*a*b*x + a^2 + 1)^(1/3), x)`

Mupad [N/A]

Not integrable

Time = 0.93 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \int \frac{\operatorname{atan}(a + bx) (a + bx)^2}{(a^2 + 2abx + b^2x^2 + 1)^{1/3}} dx$$

input `int((atan(a + b*x)*(a + b*x)^2)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/3),x)`

output `int((atan(a + b*x)*(a + b*x)^2)/(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/3), x)`

Reduce [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.89

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{1 + a^2 + 2abx + b^2x^2}} dx = \left(\int \frac{\operatorname{atan}(bx + a)}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx \right) a^2 + \left(\int \frac{\operatorname{atan}(bx + a) x^2}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx \right) b^2 + 2 \left(\int \frac{\operatorname{atan}(bx + a) x}{(b^2x^2 + 2abx + a^2 + 1)^{\frac{1}{3}}} dx \right) ab$$

input `int((b*x+a)^2*atan(b*x+a)/(b^2*x^2+2*a*b*x+a^2+1)^(1/3),x)`

output `int(atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*a**2 + int((atan(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*b**2 + 2*int((atan(a + b*x)*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*a*b`

3.71
$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{(1+a^2)c + 2abcx + b^2cx^2}} dx$$

Optimal result	591
Mathematica [B] (warning: unable to verify)	591
Rubi [N/A]	592
Maple [N/A]	593
Fricas [N/A]	593
Sympy [N/A]	594
Maxima [N/A]	594
Giac [N/A]	595
Mupad [N/A]	595
Reduce [N/A]	596

Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{(1+a^2)c + 2abcx + b^2cx^2}} dx = \text{Int} \left(\frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{c + c(a+bx)^2}}, x \right)$$

output

```
Defer(Int)((b*x+a)^2*arctan(b*x+a)/(c+c*(b*x+a)^2)^(1/3), x)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 225 vs. 2(32) = 64.

Time = 0.69 (sec) , antiderivative size = 225, normalized size of antiderivative = 5.62

$$\int \frac{(a+bx)^2 \arctan(a+bx)}{\sqrt[3]{(1+a^2)c + 2abcx + b^2cx^2}} dx = 3\sqrt[3]{1+a^2+2abx+b^2x^2}(1+(a+bx)^2)^{2/3} \left(\frac{5\sqrt[3]{2}\sqrt{\pi} \Gamma(\frac{5}{3}) {}_3F_2(1, \frac{4}{3}, \frac{4}{3}; \frac{11}{6}, \frac{7}{3}; \frac{1}{1+(a+bx)^2})}{(1+(a+bx)^2)^2} + \Gamma(\frac{11}{6}) \Gamma(\frac{1}{6}) \right)$$

140b³√c(

input

```
Integrate[((a + b*x)^2*ArcTan[a + b*x])/((1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2)^(1/3),x]
```

output

```
(-3*(1 + a^2 + 2*a*b*x + b^2*x^2)^(1/3)*(1 + (a + b*x)^2)^(2/3)*((5*2^(1/3))*Sqrt[Pi]*Gamma[5/3]*HypergeometricPFQ[{1, 4/3, 4/3}, {11/6, 7/3}, (1 + (a + b*x)^2)^(-1)])/(1 + (a + b*x)^2)^2 + Gamma[11/6]*Gamma[7/3]*(15 + 90/(1 + (a + b*x)^2) + (24*(a + b*x)*ArcTan[a + b*x]*Hypergeometric2F1[1, 4/3, 11/6, (1 + (a + b*x)^2)^(-1)])/(1 + (a + b*x)^2)^2 + 5*ArcTan[a + b*x]*(-4*(a + b*x) + 6*Sin[2*ArcTan[a + b*x]])))/(140*b*(c*(1 + a^2 + 2*a*b*x + b^2*x^2))^(1/3)*Gamma[11/6]*Gamma[7/3])
```

Rubi [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(a^2 + 1)c + 2abcx + b^2cx^2}} dx$$

$$\downarrow \text{5580}$$

$$\int \frac{(a+bx)^2 \arctan(a+bx) d(a + bx)}{\sqrt[3]{c(a + bx)^2 + c} b}$$

$$\downarrow \text{5560}$$

$$\int \frac{(a+bx)^2 \arctan(a+bx) d(a + bx)}{\sqrt[3]{c(a + bx)^2 + c} b}$$

input

```
Int[((a + b*x)^2*ArcTan[a + b*x])/((1 + a^2)*c + 2*a*b*c*x + b^2*c*x^2)^(1/3),x]
```

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{(bx + a)^2 \arctan(bx + a)}{((a^2 + 1)c + 2abcx + b^2cx^2)^{\frac{1}{3}}} dx$$

input `int((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x)`

output `int((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x)`

Fricas [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.22

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x,
algorithm="fricas")`

output `integral((b^2*x^2 + 2*a*b*x + a^2)*arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x
+ (a^2 + 1)*c)^(1/3), x)`

Sympy [N/A]

Not integrable

Time = 72.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(a + bx)^2 \operatorname{atan}(a + bx)}{\sqrt[3]{c(a^2 + 2abx + b^2x^2 + 1)}} dx$$

input `integrate((b*x+a)**2*atan(b*x+a)/((a**2+1)*c+2*a*b*c*x+b**2*c*x**2)**(1/3),x)`

output `Integral((a + b*x)**2*atan(a + b*x)/(c*(a**2 + 2*a*b*x + b**2*x**2 + 1))**(1/3), x)`

Maxima [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x, algorithm="maxima")`

output `integrate((b*x + a)^2*arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c)^(1/3), x)`

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{(bx + a)^2 \arctan(bx + a)}{(b^2cx^2 + 2abcx + (a^2 + 1)c)^{\frac{1}{3}}} dx$$

input

```
integrate((b*x+a)^2*arctan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x,
algorithm="giac")
```

output

```
integrate((b*x + a)^2*arctan(b*x + a)/(b^2*c*x^2 + 2*a*b*c*x + (a^2 + 1)*c
)^(1/3), x)
```

Mupad [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx = \int \frac{\operatorname{atan}(a + bx) (a + bx)^2}{(cb^2x^2 + 2acbx + c(a^2 + 1))^{1/3}} dx$$

input

```
int((atan(a + b*x)*(a + b*x)^2)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/3
),x)
```

output

```
int((atan(a + b*x)*(a + b*x)^2)/(c*(a^2 + 1) + b^2*c*x^2 + 2*a*b*c*x)^(1/3
), x)
```

Reduce [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 105, normalized size of antiderivative = 2.62

$$\int \frac{(a + bx)^2 \arctan(a + bx)}{\sqrt[3]{(1 + a^2)c + 2abcx + b^2cx^2}} dx$$

$$= \frac{\left(\int \frac{\operatorname{atan}(bx+a)}{(b^2x^2+2abx+a^2+1)^{\frac{1}{3}}} dx \right) a^2 + \left(\int \frac{\operatorname{atan}(bx+a)x^2}{(b^2x^2+2abx+a^2+1)^{\frac{1}{3}}} dx \right) b^2 + 2 \left(\int \frac{\operatorname{atan}(bx+a)x}{(b^2x^2+2abx+a^2+1)^{\frac{1}{3}}} dx \right) ab}{c^{\frac{1}{3}}}$$

input

```
int((b*x+a)^2*atan(b*x+a)/((a^2+1)*c+2*a*b*c*x+b^2*c*x^2)^(1/3),x)
```

output

```
(int(atan(a + b*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*a**2 + int((
atan(a + b*x)*x**2)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*b**2 + 2*in
t((atan(a + b*x)*x)/(a**2 + 2*a*b*x + b**2*x**2 + 1)**(1/3),x)*a*b)/c**(1/
3)
```

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	597
4.2	Links to plain text integration problems used in this report for each CAS .	615

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
      ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```



```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file