

Computer Algebra Independent Integration Tests

Summer 2024

5-Inverse-trig-functions/5.5-Inverse-secant/286-5.5

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 9:07pm

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42
3	Listing of integrals	44
3.1	$\int \frac{\sec^{-1}(ax^5)}{x} dx$	46
3.2	$\int x^3 \sec^{-1}(\sqrt{x}) dx$	52
3.3	$\int x^2 \sec^{-1}(\sqrt{x}) dx$	58
3.4	$\int x \sec^{-1}(\sqrt{x}) dx$	64
3.5	$\int \sec^{-1}(\sqrt{x}) dx$	70
3.6	$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx$	75

3.7	$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx$	81
3.8	$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx$	87
3.9	$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$	94
3.10	$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx$	101
3.11	$\int x \sec^{-1}\left(\frac{a}{x}\right) dx$	107
3.12	$\int \sec^{-1}\left(\frac{a}{x}\right) dx$	113
3.13	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx$	118
3.14	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx$	124
3.15	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx$	130
3.16	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx$	135
3.17	$\int \frac{\sec^{-1}(ax^n)}{x} dx$	141
3.18	$\int x^4 \sec^{-1}(a + bx) dx$	147
3.19	$\int x^3 \sec^{-1}(a + bx) dx$	155
3.20	$\int x^2 \sec^{-1}(a + bx) dx$	163
3.21	$\int x \sec^{-1}(a + bx) dx$	170
3.22	$\int \sec^{-1}(a + bx) dx$	177
3.23	$\int \frac{\sec^{-1}(a+bx)}{x} dx$	183
3.24	$\int \frac{\sec^{-1}(a+bx)}{x^2} dx$	193
3.25	$\int \frac{\sec^{-1}(a+bx)}{x^3} dx$	201
3.26	$\int \frac{\sec^{-1}(a+bx)}{x^4} dx$	210
3.27	$\int x^3 \sec^{-1}(a + bx)^2 dx$	220
3.28	$\int x^2 \sec^{-1}(a + bx)^2 dx$	228
3.29	$\int x \sec^{-1}(a + bx)^2 dx$	235
3.30	$\int \sec^{-1}(a + bx)^2 dx$	241
3.31	$\int \frac{\sec^{-1}(a+bx)^2}{x} dx$	247
3.32	$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx$	258
3.33	$\int x^2 \sec^{-1}(a + bx)^3 dx$	265
3.34	$\int x \sec^{-1}(a + bx)^3 dx$	274
3.35	$\int \sec^{-1}(a + bx)^3 dx$	281
3.36	$\int \frac{\sec^{-1}(a+bx)^3}{x} dx$	288
3.37	$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx$	300
3.38	$\int x(a + b \sec^{-1}(c + dx^2)) dx$	307
3.39	$\int x^2(a + b \sec^{-1}(c + dx^3)) dx$	313
3.40	$\int x^3(a + b \sec^{-1}(c + dx^4)) dx$	319
3.41	$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$	325
3.42	$\int \sec^{-1}(ce^{a+bx}) dx$	331
3.43	$\int e^{\sec^{-1}(ax)} x^2 dx$	338

3.44	$\int e^{\sec^{-1}(ax)} x dx$	344
3.45	$\int e^{\sec^{-1}(ax)} dx$	349
3.46	$\int \frac{e^{\sec^{-1}(ax)}}{x} dx$	354
3.47	$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$	359
3.48	$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx$	364
3.49	$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$	369
3.50	$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$	374
4	Appendix	381
4.1	Listing of Grading functions	381
4.2	Links to plain text integration problems used in this report for each CAS399	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [50]. This is test number [286].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (50)	0.00 (0)
Mathematica	98.00 (49)	2.00 (1)
Maple	74.00 (37)	26.00 (13)
Fricas	56.00 (28)	44.00 (22)
Giac	54.00 (27)	46.00 (23)
Maxima	36.00 (18)	64.00 (32)
Sympy	26.00 (13)	74.00 (37)
Mupad	20.00 (10)	80.00 (40)
Reduce	4.00 (2)	96.00 (48)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	70.000	14.000	14.000	2.000
Maple	68.000	6.000	0.000	26.000
Fricas	48.000	8.000	0.000	44.000
Giac	32.000	22.000	0.000	46.000
Maxima	32.000	4.000	0.000	64.000
Sympy	6.000	0.000	20.000	74.000
Mupad	0.000	20.000	0.000	80.000
Reduce	0.000	4.000	0.000	96.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	0.00	100.00	0.00
Maple	13	100.00	0.00	0.00
Fricas	22	90.91	0.00	9.09
Giac	23	91.30	0.00	8.70
Maxima	32	100.00	0.00	0.00
Sympy	37	89.19	10.81	0.00
Mupad	40	0.00	100.00	0.00
Reduce	48	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.07
Fricas	0.13
Giac	0.17
Reduce	0.22
Maple	0.42
Rubi	0.47
Mathematica	0.54
Mupad	0.96
Sympy	18.20

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	34.00	0.90	34.00	0.90
Mupad	37.50	0.91	36.50	0.90
Maxima	56.67	1.21	54.50	1.20
Sympy	76.46	1.63	61.00	1.67
Fricas	102.61	1.37	56.50	0.92
Giac	121.11	1.65	82.00	1.72
Rubi	125.98	1.04	74.50	1.00
Maple	174.38	1.36	76.00	1.31
Mathematica	195.20	1.84	107.00	1.00

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {1, 6, 17, 22, 41, 42, 50}

Mathematica {27, 28, 31, 32, 36}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	42

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 18, 19, 20, 21, 23, 27, 28, 29, 30, 33, 34, 35, 43, 44, 45, 46, 47, 48, 49, 50 }

B grade { 14, 31, 32, 36, 40, 41, 42 }

C grade { 17, 22, 24, 25, 26, 38, 39 }

F normal fail { }

F(-1) timedout fail { 37 }

F(-2) exception fail { }

Maple

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 28, 29, 30, 32, 33, 34, 38, 39, 40, 42, 50 }

B grade { 24, 25, 26 }

C grade { }

F normal fail { 1, 31, 35, 36, 37, 41, 43, 44, 45, 46, 47, 48, 49 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 15, 18, 19, 20, 21, 25, 26, 38, 39, 40, 41, 47, 48, 49 }

B grade { 14, 16, 22, 24 }

C grade { }

F normal fail { 1, 6, 13, 23, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 43, 44, 45, 46, 50 }

F(-1) timedout fail { }

F(-2) exception fail { 17, 42 }

Maxima

A grade { 2, 3, 4, 5, 7, 10, 11, 12, 14, 15, 16, 22, 38, 39, 40, 41 }

B grade { 8, 9 }

C grade { }

F normal fail { 1, 6, 13, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 7, 8, 9, 10, 11, 12, 15, 16, 21, 24, 38, 39, 40, 41, 47, 50 }

B grade { 2, 3, 4, 5, 14, 18, 19, 20, 22, 25, 26 }

C grade { }

F normal fail { 1, 13, 17, 23, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 48, 49 }

F(-1) timedout fail { }

F(-2) exception fail { 6, 27 }

Mupad

A grade { }

B grade { 5, 7, 11, 12, 14, 22, 38, 39, 40, 41 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 4, 6, 8, 9, 10, 13, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-2) exception fail { }

Sympy

A grade { 10, 11, 12 }

B grade { }

C grade { 2, 3, 4, 5, 7, 8, 9, 14, 15, 16 }

F normal fail { 1, 6, 13, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { 38, 39, 40, 41 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 11, 12 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	62	72	56	0	0	0	0	0	12	0
N.S.	1	1.16	0.90	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.456	0.024	0.000	0.000	0.000	0.000	0.000	0.252	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	59	40	40	66	32	119	152	9	0
N.S.	1	1.02	0.69	0.69	1.14	0.55	2.05	2.62	0.16	0.00
time (sec)	N/A	0.221	0.020	0.174	0.042	0.106	49.824	0.134	0.277	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	50	35	35	52	27	90	116	9	0
N.S.	1	1.06	0.74	0.74	1.11	0.57	1.91	2.47	0.19	0.00
time (sec)	N/A	0.208	0.016	0.113	0.034	0.111	16.415	0.141	0.257	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	39	28	28	38	20	61	80	7	0
N.S.	1	1.08	0.78	0.78	1.06	0.56	1.69	2.22	0.19	0.00
time (sec)	N/A	0.198	0.014	0.113	0.035	0.106	5.434	0.134	0.217	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	22	21	14	29	41	5	21
N.S.	1	1.00	1.00	1.22	1.17	0.78	1.61	2.28	0.28	1.17
time (sec)	N/A	0.165	0.003	0.105	0.052	0.105	2.102	0.128	0.216	1.281

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	62	54	63	0	0	0	0	9	0
N.S.	1	1.11	0.96	1.12	0.00	0.00	0.00	0.00	0.16	0.00
time (sec)	N/A	0.464	0.018	0.341	0.000	0.000	0.000	0.000	0.251	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	36	32	44	51	19	75	30	9	28
N.S.	1	0.95	0.84	1.16	1.34	0.50	1.97	0.79	0.24	0.74
time (sec)	N/A	0.203	0.016	0.118	0.148	0.119	11.785	0.129	0.220	0.824

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	57	55	57	80	29	144	44	9	0
N.S.	1	1.06	1.02	1.06	1.48	0.54	2.67	0.81	0.17	0.00
time (sec)	N/A	0.208	0.024	0.121	0.116	0.199	38.071	0.134	0.215	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	76	45	67	106	35	180	58	9	0
N.S.	1	1.12	0.66	0.99	1.56	0.51	2.65	0.85	0.13	0.00
time (sec)	N/A	0.218	0.037	0.118	0.124	0.121	108.728	0.390	0.221	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	62	42	56	54	39	51	47	12	0
N.S.	1	1.11	0.75	1.00	0.96	0.70	0.91	0.84	0.21	0.00
time (sec)	N/A	0.266	0.025	0.684	0.126	0.104	0.213	0.148	0.221	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	57	44	67	46	38	41	39	47	38
N.S.	1	1.21	0.94	1.43	0.98	0.81	0.87	0.83	1.00	0.81
time (sec)	N/A	0.231	0.018	0.644	0.134	0.105	0.181	0.124	0.227	0.699

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	28	24	27	22	28	21	24
N.S.	1	1.00	1.00	1.08	0.92	1.04	0.85	1.08	0.81	0.92
time (sec)	N/A	0.201	0.008	0.664	0.032	0.117	0.148	0.123	0.218	0.761

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	67	59	76	0	0	0	0	12	0
N.S.	1	1.14	1.00	1.29	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.368	0.017	0.844	0.000	0.000	0.000	0.000	0.216	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	C	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	93	30	52	107	29	61	12	29
N.S.	1	1.00	3.00	0.97	1.68	3.45	0.94	1.97	0.39	0.94
time (sec)	N/A	0.246	0.089	0.041	0.034	0.124	1.304	0.124	0.208	0.683

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	33	32	39	53	61	12	0
N.S.	1	1.00	0.95	0.87	0.84	1.03	1.39	1.61	0.32	0.00
time (sec)	N/A	0.221	0.016	0.647	0.119	0.122	0.738	0.130	0.214	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	C	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	61	69	54	64	142	100	80	12	0
N.S.	1	1.02	1.15	0.90	1.07	2.37	1.67	1.33	0.20	0.00
time (sec)	N/A	0.245	0.028	0.646	0.110	0.126	1.679	0.126	0.219	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	80	60	93	0	0	0	0	12	0
N.S.	1	1.16	0.87	1.35	0.00	0.00	0.00	0.00	0.17	0.00
time (sec)	N/A	0.492	0.056	0.795	0.000	0.000	0.000	0.000	0.216	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	197	206	173	329	0	152	0	412	12	0
N.S.	1	1.05	0.88	1.67	0.00	0.77	0.00	2.09	0.06	0.00
time (sec)	N/A	0.724	0.115	0.244	0.000	0.125	0.000	0.161	0.226	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	157	150	249	0	130	0	302	12	0
N.S.	1	1.01	0.97	1.61	0.00	0.84	0.00	1.95	0.08	0.00
time (sec)	N/A	0.536	0.178	0.170	0.000	0.122	0.000	0.153	0.223	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	131	190	0	117	0	207	12	0
N.S.	1	1.00	1.13	1.64	0.00	1.01	0.00	1.78	0.10	0.00
time (sec)	N/A	0.399	0.118	0.171	0.000	0.139	0.000	0.162	0.214	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	110	108	0	104	0	136	10	0
N.S.	1	1.00	1.41	1.38	0.00	1.33	0.00	1.74	0.13	0.00
time (sec)	N/A	0.446	0.077	0.155	0.000	0.121	0.000	0.147	0.213	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F	B	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	35	468	45	55	73	0	82	8	35
N.S.	1	0.95	12.65	1.22	1.49	1.97	0.00	2.22	0.22	0.95
time (sec)	N/A	0.228	1.936	0.041	0.030	0.128	0.000	0.140	0.227	1.007

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	200	263	284	374	0	0	0	0	12	0
N.S.	1	1.32	1.42	1.87	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	1.101	0.222	0.904	0.000	0.000	0.000	0.000	0.246	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	73	112	126	0	281	0	94	12	0
N.S.	1	1.04	1.60	1.80	0.00	4.01	0.00	1.34	0.17	0.00
time (sec)	N/A	0.412	0.197	0.348	0.000	0.130	0.000	0.148	0.248	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	148	198	319	0	427	0	216	12	0
N.S.	1	1.18	1.58	2.55	0.00	3.42	0.00	1.73	0.10	0.00
time (sec)	N/A	0.742	0.668	0.334	0.000	0.132	0.000	0.164	0.228	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	181	222	241	528	0	548	0	451	12	0
N.S.	1	1.23	1.33	2.92	0.00	3.03	0.00	2.49	0.07	0.00
time (sec)	N/A	1.062	0.287	0.335	0.000	0.189	0.000	0.186	0.258	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	381	357	667	673	0	0	0	0	14	0
N.S.	1	0.94	1.75	1.77	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.638	8.171	0.854	0.000	0.000	0.000	0.000	0.261	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	288	271	473	498	0	0	0	0	14	0
N.S.	1	0.94	1.64	1.73	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	0.541	4.011	0.760	0.000	0.000	0.000	0.000	0.226	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	146	144	225	0	0	0	0	12	0
N.S.	1	0.95	0.94	1.46	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	0.442	0.094	0.500	0.000	0.000	0.000	0.000	0.229	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	89	111	162	0	0	0	0	10	0
N.S.	1	0.95	1.18	1.72	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.411	0.078	0.266	0.000	0.000	0.000	0.000	0.240	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	310	398	813	0	0	0	0	0	14	0
N.S.	1	1.28	2.62	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	1.579	1.515	0.000	0.000	0.000	0.000	0.000	0.258	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	244	248	686	336	0	0	0	0	14	0
N.S.	1	1.02	2.81	1.38	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.724	1.459	0.772	0.000	0.000	0.000	0.000	0.240	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	494	446	446	716	0	0	0	0	14	0
N.S.	1	0.90	0.90	1.45	0.00	0.00	0.00	0.00	0.03	0.00
time (sec)	N/A	0.728	0.249	0.739	0.000	0.000	0.000	0.000	0.243	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	253	257	379	0	0	0	0	12	0
N.S.	1	0.91	0.92	1.36	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.571	0.095	0.727	0.000	0.000	0.000	0.000	0.249	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	154	149	160	0	0	0	0	0	10	0
N.S.	1	0.97	1.04	0.00	0.00	0.00	0.00	0.00	0.06	0.00
time (sec)	N/A	0.576	0.069	0.000	0.000	0.000	0.000	0.000	0.266	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	430	529	1058	0	0	0	0	0	14	0
N.S.	1	1.23	2.46	0.00	0.00	0.00	0.00	0.00	0.03	0.00
time (sec)	N/A	1.847	2.555	0.000	0.000	0.000	0.000	0.000	0.265	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F(-1)	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	362	365	0	0	0	0	0	0	14	0
N.S.	1	1.01	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00
time (sec)	N/A	0.885	0.000	0.000	0.000	0.000	0.000	0.000	0.258	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	54	516	64	71	96	0	100	21	52
N.S.	1	0.93	8.90	1.10	1.22	1.66	0.00	1.72	0.36	0.90
time (sec)	N/A	0.252	2.014	0.221	0.028	0.113	0.000	0.240	0.231	1.208

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	54	516	64	71	96	0	100	23	52
N.S.	1	0.93	8.90	1.10	1.22	1.66	0.00	1.72	0.40	0.90
time (sec)	N/A	0.273	0.274	0.224	0.040	0.125	0.000	0.250	0.234	0.945

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	54	137	64	71	96	0	100	23	52
N.S.	1	0.93	2.36	1.10	1.22	1.66	0.00	1.72	0.40	0.90
time (sec)	N/A	0.277	0.211	0.242	0.042	0.112	0.000	0.261	0.257	0.909

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	A	A	F(-1)	A	F	B
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	49	45	130	0	66	92	0	75	17	44
N.S.	1	0.92	2.65	0.00	1.35	1.88	0.00	1.53	0.35	0.90
time (sec)	N/A	0.323	0.207	0.000	0.039	0.130	0.000	0.196	0.266	1.317

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	100	280	111	0	0	0	0	12	0
N.S.	1	1.18	3.29	1.31	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.466	0.806	0.817	0.000	0.000	0.000	0.000	0.258	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	99	97	95	0	0	0	0	0	12	0
N.S.	1	0.98	0.96	0.00	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	0.341	0.219	0.000	0.000	0.000	0.000	0.000	0.274	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	91	89	107	0	0	0	0	0	10	0
N.S.	1	0.98	1.18	0.00	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.322	0.137	0.000	0.000	0.000	0.000	0.000	0.300	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	91	89	54	0	0	0	0	0	8	0
N.S.	1	0.98	0.59	0.00	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	0.304	0.048	0.000	0.000	0.000	0.000	0.000	0.270	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	79	0	0	0	0	0	12	0
N.S.	1	1.00	1.76	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.281	0.038	0.000	0.000	0.000	0.000	0.000	0.252	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	39	43	34	0	0	23	0	43	12	0
N.S.	1	1.10	0.87	0.00	0.00	0.59	0.00	1.10	0.31	0.00
time (sec)	N/A	0.230	0.030	0.000	0.000	0.126	0.000	0.138	0.236	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	41	42	30	0	0	30	0	0	12	0
N.S.	1	1.02	0.73	0.00	0.00	0.73	0.00	0.00	0.29	0.00
time (sec)	N/A	0.282	0.031	0.000	0.000	0.116	0.000	0.000	0.251	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	84	79	54	0	0	40	0	0	12	0
N.S.	1	0.94	0.64	0.00	0.00	0.48	0.00	0.00	0.14	0.00
time (sec)	N/A	0.288	0.107	0.000	0.000	0.155	0.000	0.000	0.254	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	A	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	68	59	99	0	0	0	115	21	0
N.S.	1	0.99	0.86	1.43	0.00	0.00	0.00	1.67	0.30	0.00
time (sec)	N/A	0.491	0.050	0.519	0.000	0.000	0.000	0.338	0.237	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [23] had the largest ratio of [1.3999999999999991]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	9	8	1.16	10	0.800
2	A	4	4	1.02	10	0.400
3	A	4	4	1.06	10	0.400
4	A	4	4	1.08	8	0.500
5	A	2	2	1.00	6	0.333
6	A	9	8	1.11	10	0.800
7	A	6	5	0.95	10	0.500
8	A	7	6	1.06	10	0.600
9	A	8	7	1.12	10	0.700
10	A	6	5	1.11	10	0.500
11	A	4	4	1.21	8	0.500
12	A	3	3	1.00	6	0.500
13	A	8	7	1.14	10	0.700
14	A	6	5	1.00	10	0.500
15	A	3	3	1.00	10	0.300
16	A	7	6	1.02	10	0.600
17	A	9	8	1.16	10	0.800
18	A	10	9	1.05	10	0.900
19	A	9	8	1.01	10	0.800
20	A	6	5	1.00	10	0.500
21	A	10	9	1.00	8	1.125

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	6	5	0.95	6	0.833
23	A	15	14	1.32	10	1.400
24	A	8	7	1.04	10	0.700
25	A	13	12	1.18	10	1.200
26	A	14	13	1.23	10	1.300
27	A	7	6	0.94	12	0.500
28	A	6	5	0.94	12	0.417
29	A	7	6	0.95	10	0.600
30	A	8	7	0.95	8	0.875
31	A	16	15	1.28	12	1.250
32	A	6	5	1.02	12	0.417
33	A	6	5	0.90	12	0.417
34	A	7	6	0.91	10	0.600
35	A	9	8	0.97	8	1.000
36	A	16	15	1.23	12	1.250
37	A	6	5	1.01	12	0.417
38	A	3	2	0.93	14	0.143
39	A	3	2	0.93	16	0.125
40	A	3	2	0.93	16	0.125
41	A	7	6	0.92	14	0.429
42	A	9	8	1.18	10	0.800
43	A	5	4	0.98	10	0.400
44	A	5	4	0.98	8	0.500
45	A	4	3	0.98	6	0.500
46	A	5	4	1.00	10	0.400
47	A	4	3	1.10	10	0.300
48	A	6	5	1.02	10	0.500
49	A	5	4	0.94	10	0.400
50	A	10	9	0.99	19	0.474

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\sec^{-1}(ax^5)}{x} dx$	46
3.2	$\int x^3 \sec^{-1}(\sqrt{x}) dx$	52
3.3	$\int x^2 \sec^{-1}(\sqrt{x}) dx$	58
3.4	$\int x \sec^{-1}(\sqrt{x}) dx$	64
3.5	$\int \sec^{-1}(\sqrt{x}) dx$	70
3.6	$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx$	75
3.7	$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx$	81
3.8	$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx$	87
3.9	$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$	94
3.10	$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx$	101
3.11	$\int x \sec^{-1}\left(\frac{a}{x}\right) dx$	107
3.12	$\int \sec^{-1}\left(\frac{a}{x}\right) dx$	113
3.13	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx$	118
3.14	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx$	124
3.15	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx$	130
3.16	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx$	135
3.17	$\int \frac{\sec^{-1}(ax^n)}{x} dx$	141
3.18	$\int x^4 \sec^{-1}(a + bx) dx$	147
3.19	$\int x^3 \sec^{-1}(a + bx) dx$	155
3.20	$\int x^2 \sec^{-1}(a + bx) dx$	163
3.21	$\int x \sec^{-1}(a + bx) dx$	170
3.22	$\int \sec^{-1}(a + bx) dx$	177
3.23	$\int \frac{\sec^{-1}(a+bx)}{x} dx$	183
3.24	$\int \frac{\sec^{-1}(a+bx)}{x^2} dx$	193
3.25	$\int \frac{\sec^{-1}(a+bx)}{x^3} dx$	201

3.26	$\int \frac{\sec^{-1}(a+bx)}{x^4} dx$	210
3.27	$\int x^3 \sec^{-1}(a+bx)^2 dx$	220
3.28	$\int x^2 \sec^{-1}(a+bx)^2 dx$	228
3.29	$\int x \sec^{-1}(a+bx)^2 dx$	235
3.30	$\int \sec^{-1}(a+bx)^2 dx$	241
3.31	$\int \frac{\sec^{-1}(a+bx)^2}{x} dx$	247
3.32	$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx$	258
3.33	$\int x^2 \sec^{-1}(a+bx)^3 dx$	265
3.34	$\int x \sec^{-1}(a+bx)^3 dx$	274
3.35	$\int \sec^{-1}(a+bx)^3 dx$	281
3.36	$\int \frac{\sec^{-1}(a+bx)^3}{x} dx$	288
3.37	$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx$	300
3.38	$\int x(a+b \sec^{-1}(c+dx^2)) dx$	307
3.39	$\int x^2(a+b \sec^{-1}(c+dx^3)) dx$	313
3.40	$\int x^3(a+b \sec^{-1}(c+dx^4)) dx$	319
3.41	$\int x^{-1+n} \sec^{-1}(a+bx^n) dx$	325
3.42	$\int \sec^{-1}(ce^{a+bx}) dx$	331
3.43	$\int e^{\sec^{-1}(ax)} x^2 dx$	338
3.44	$\int e^{\sec^{-1}(ax)} x dx$	344
3.45	$\int e^{\sec^{-1}(ax)} dx$	349
3.46	$\int \frac{e^{\sec^{-1}(ax)}}{x} dx$	354
3.47	$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$	359
3.48	$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx$	364
3.49	$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$	369
3.50	$\int \frac{\sec^{-1}(a+bx)}{\frac{a}{b}+dx} dx$	374

3.1 $\int \frac{\sec^{-1}(ax^5)}{x} dx$

Optimal result	46
Mathematica [A] (verified)	46
Rubi [A] (warning: unable to verify)	47
Maple [F]	49
Fricas [F]	50
Sympy [F]	50
Maxima [F]	50
Giac [F]	51
Mupad [F(-1)]	51
Reduce [F]	51

Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right) + \frac{1}{10}i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right)$$

output

```
1/10*I*arcsec(a*x^5)^2-1/5*arcsec(a*x^5)*ln(1+(1/a/x^5+I*(1-1/a^2/x^10)^(1/2))^(1/2))+1/10*I*polylog(2,-(1/a/x^5+I*(1-1/a^2/x^10)^(1/2))^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.90

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \frac{1}{10}i \left(\sec^{-1}(ax^5) \left(\sec^{-1}(ax^5) + 2i \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right)\right) + \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right) \right)$$

input

```
Integrate[ArcSec[a*x^5]/x,x]
```

output

```
(I/10)*(ArcSec[a*x^5]*(ArcSec[a*x^5] + (2*I)*Log[1 + E^((2*I)*ArcSec[a*x^5]
)]) + PolyLog[2, -E^((2*I)*ArcSec[a*x^5])])
```

Rubi [A] (warning: unable to verify)

Time = 0.46 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.16, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {7282, 5741, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(ax^5)}{x} dx \\
 & \quad \downarrow 7282 \\
 & \frac{1}{5} \int \frac{\sec^{-1}(ax^5)}{x^5} dx^5 \\
 & \quad \downarrow 5741 \\
 & -\frac{1}{5} \int \frac{\arccos\left(\frac{1}{ax^5}\right)}{x^5} d\frac{1}{x^5} \\
 & \quad \downarrow 5137 \\
 & \frac{1}{5} \int a \sqrt{1 - \frac{1}{a^2 x^{10}} x^5} \arccos\left(\frac{1}{ax^5}\right) d\arccos\left(\frac{1}{ax^5}\right) \\
 & \quad \downarrow 3042 \\
 & \frac{1}{5} \int \arccos\left(\frac{1}{ax^5}\right) \tan\left(\arccos\left(\frac{1}{ax^5}\right)\right) d\arccos\left(\frac{1}{ax^5}\right) \\
 & \quad \downarrow 4202 \\
 & \frac{1}{5} \left(\frac{ix^{10}}{2} - 2i \int \frac{e^{2i \arccos\left(\frac{1}{ax^5}\right)} \arccos\left(\frac{1}{ax^5}\right)}{1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)}} d\arccos\left(\frac{1}{ax^5}\right) \right) \\
 & \quad \downarrow 2620 \\
 & \frac{1}{5} \left(\frac{ix^{10}}{2} - 2i \left(\frac{1}{2} i \int \log\left(1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)}\right) d\arccos\left(\frac{1}{ax^5}\right) - \frac{1}{2} i \arccos\left(\frac{1}{ax^5}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)}\right) \right) \right)
 \end{aligned}$$

↓ 2715

$$\frac{1}{5} \left(\frac{ix^{10}}{2} - 2i \left(\frac{1}{4} \int e^{2i \arccos\left(\frac{1}{ax^5}\right)} \log \left(1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)} \right) de^{2i \arccos\left(\frac{1}{ax^5}\right)} - \frac{1}{2} i \arccos \left(\frac{1}{ax^5} \right) \log \left(1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)} \right) \right) \right)$$

↓ 2838

$$\frac{1}{5} \left(\frac{ix^{10}}{2} - 2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \arccos\left(\frac{1}{ax^5}\right)} \right) - \frac{1}{2} i \arccos \left(\frac{1}{ax^5} \right) \log \left(1 + e^{2i \arccos\left(\frac{1}{ax^5}\right)} \right) \right) \right)$$

input `Int[ArcSec[a*x^5]/x, x]`

output `((I/2)*x^10 - (2*I)*((-1/2*I)*ArcCos[1/(a*x^5)]*Log[1 + E^((2*I)*ArcCos[1/(a*x^5)])]) - PolyLog[2, -E^((2*I)*ArcCos[1/(a*x^5)])]/4)/5`

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[
 *((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(
 e + f*x))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGt
 Q[m, 0]`

rule 5137 `Int[((a_.) + ArcCos[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] := -Subst[Int[
 (a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0
]`

rule 5741 `Int[((a_.) + ArcSec[(c_.)*(x_)])*(b_.)/(x_), x_Symbol] := -Subst[Int[(a + b
 *ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7282 `Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1
 /lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (
 lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[
 u] && !RationalFunctionQ[u, x]`

Maple [F]

$$\int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

input `int(arcsec(a*x^5)/x,x)`

output `int(arcsec(a*x^5)/x,x)`

Fricas [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

input `integrate(arcsec(a*x^5)/x,x, algorithm="fricas")`

output `integral(arcsec(a*x^5)/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asec}(ax^5)}{x} dx$$

input `integrate(asec(a*x**5)/x,x)`

output `Integral(asec(a*x**5)/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

input `integrate(arcsec(a*x^5)/x,x, algorithm="maxima")`

output `-5*a^2*integrate(sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1)*log(x)/(a^4*x^11 - a^2*x), x) - 5*I*a^2*integrate(log(x)/(a^4*x^11 - a^2*x), x) + arctan(sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1))*log(x) - 1/2*I*log(a^2*x^10)*log(x) + 1/2*I*log(a*x^5 + 1)*log(x) + 1/2*I*log(-a*x^5 + 1)*log(x) + I*log(a)*log(x) + 5/2*I*log(x)^2 + 1/10*I*dilog(a*x^5) + 1/10*I*dilog(-a*x^5)`

Giac [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

input `integrate(arcsec(a*x^5)/x,x, algorithm="giac")`

output `integrate(arcsec(a*x^5)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\arccos\left(\frac{1}{ax^5}\right)}{x} dx$$

input `int(arccos(1/(a*x^5))/x,x)`

output `int(arccos(1/(a*x^5))/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asec}(ax^5)}{x} dx$$

input `int(asec(a*x^5)/x,x)`

output `int(asec(a*x**5)/x,x)`

3.2 $\int x^3 \sec^{-1}(\sqrt{x}) dx$

Optimal result	52
Mathematica [A] (verified)	52
Rubi [A] (verified)	53
Maple [A] (verified)	54
Fricas [A] (verification not implemented)	55
Sympy [C] (verification not implemented)	55
Maxima [A] (verification not implemented)	56
Giac [B] (verification not implemented)	56
Mupad [F(-1)]	57
Reduce [F]	57

Optimal result

Integrand size = 10, antiderivative size = 58

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{4}\sqrt{-1+x} - \frac{1}{4}(-1+x)^{3/2} - \frac{3}{20}(-1+x)^{5/2} - \frac{1}{28}(-1+x)^{7/2} + \frac{1}{4}x^4 \sec^{-1}(\sqrt{x})$$

output

```
-1/4*(-1+x)^(1/2)-1/4*(-1+x)^(3/2)-3/20*(-1+x)^(5/2)-1/28*(-1+x)^(7/2)+1/4*x^4*arcsec(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{140}\sqrt{-1+x}(16+8x+6x^2+5x^3) + \frac{1}{4}x^4 \sec^{-1}(\sqrt{x})$$

input

```
Integrate[x^3*ArcSec[Sqrt[x]],x]
```

output

```
-1/140*(Sqrt[-1+x]*(16+8*x+6*x^2+5*x^3))+ (x^4*ArcSec[Sqrt[x]])/4
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5793, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \sec^{-1}(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{5793} \\
 & \frac{1}{4} x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \frac{x^3}{2\sqrt{x-1}} \, dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{8} \int \frac{x^3}{\sqrt{x-1}} \, dx \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{4} x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{8} \int \left((x-1)^{5/2} + 3(x-1)^{3/2} + 3\sqrt{x-1} + \frac{1}{\sqrt{x-1}} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{4} x^4 \sec^{-1}(\sqrt{x}) + \frac{1}{8} \left(-\frac{2}{7}(x-1)^{7/2} - \frac{6}{5}(x-1)^{5/2} - 2(x-1)^{3/2} - 2\sqrt{x-1} \right)
 \end{aligned}$$

input `Int [x^3*ArcSec [Sqrt [x]] , x]`

output `(-2*Sqrt [-1 + x] - 2*(-1 + x)^(3/2) - (6*(-1 + x)^(5/2))/5 - (2*(-1 + x)^(7/2))/7)/8 + (x^4*ArcSec [Sqrt [x]])/4`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5793 `Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

method	result	size
parts	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{\sqrt{\frac{-1+x}{x}} \sqrt{x} (5x^3+6x^2+8x+16)}{140}$	40
derivativeldivides	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{(-1+x)(5x^3+6x^2+8x+16)}{140\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	43
default	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{(-1+x)(5x^3+6x^2+8x+16)}{140\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	43

input `int(x^3*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/4*x^4*arcsec(x^(1/2))-1/140*((-1+x)/x)^(1/2)*x^(1/2)*(5*x^3+6*x^2+8*x+16)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.55

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{140} (5x^3 + 6x^2 + 8x + 16) \sqrt{x-1}$$

input `integrate(x^3*arcsec(x^(1/2)),x, algorithm="fricas")`

output `1/4*x^4*arcsec(sqrt(x)) - 1/140*(5*x^3 + 6*x^2 + 8*x + 16)*sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 49.82 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.05

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \frac{x^4 \operatorname{asec}(\sqrt{x})}{4} \begin{cases} \frac{2x^3\sqrt{x-1}}{7} + \frac{12x^2\sqrt{x-1}}{35} + \frac{16x\sqrt{x-1}}{35} + \frac{32\sqrt{x-1}}{35} & \text{for } |x| > 1 \\ \frac{2ix^3\sqrt{1-x}}{7} + \frac{12ix^2\sqrt{1-x}}{35} + \frac{16ix\sqrt{1-x}}{35} + \frac{32i\sqrt{1-x}}{35} & \text{otherwise} \end{cases}$$

8

input `integrate(x**3*asec(x**(1/2)),x)`

output `x**4*asec(sqrt(x))/4 - Piecewise((2*x**3*sqrt(x - 1)/7 + 12*x**2*sqrt(x - 1)/35 + 16*x*sqrt(x - 1)/35 + 32*sqrt(x - 1)/35, Abs(x) > 1), (2*I*x**3*sqrt(1 - x)/7 + 12*I*x**2*sqrt(1 - x)/35 + 16*I*x*sqrt(1 - x)/35 + 32*I*sqrt(1 - x)/35, True))/8`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.14

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{28} x^{\frac{7}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{5}{2}} \\ + \frac{1}{4} x^4 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{4} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

input `integrate(x^3*arcsec(x^(1/2)),x, algorithm="maxima")`

output `-1/28*x^(7/2)*(-1/x + 1)^(7/2) - 3/20*x^(5/2)*(-1/x + 1)^(5/2) + 1/4*x^4*arcsec(sqrt(x)) - 1/4*x^(3/2)*(-1/x + 1)^(3/2) - 1/4*sqrt(x)*sqrt(-1/x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(38) = 76.

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.62

$$\int x^3 \sec^{-1}(\sqrt{x}) dx \\ = -\frac{1}{3584} x^{\frac{7}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^7 - \frac{7}{2560} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^5 \\ + \frac{1}{4} x^4 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{7}{512} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^3 - \frac{35}{512} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1\right) \\ + \frac{1225 x^3 \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^6 + 245 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^4 + 49 x \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^2 + 5}{17920 x^{\frac{7}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^7}$$

input `integrate(x^3*arcsec(x^(1/2)),x, algorithm="giac")`

output

```
-1/3584*x^(7/2)*(sqrt(-1/x + 1) - 1)^7 - 7/2560*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 + 1/4*x^4*arccos(1/sqrt(x)) - 7/512*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 - 35/512*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/17920*(1225*x^3*(sqrt(-1/x + 1) - 1)^6 + 245*x^2*(sqrt(-1/x + 1) - 1)^4 + 49*x*(sqrt(-1/x + 1) - 1)^2 + 5)/(x^(7/2)*(sqrt(-1/x + 1) - 1)^7)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \int x^3 \arccos\left(\frac{1}{\sqrt{x}}\right) dx$$

input

```
int(x^3*acos(1/x^(1/2)),x)
```

output

```
int(x^3*acos(1/x^(1/2)), x)
```

Reduce [F]

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \int asec(\sqrt{x}) x^3 dx$$

input

```
int(x^3*asec(x^(1/2)),x)
```

output

```
int(asec(sqrt(x))*x**3,x)
```

3.3 $\int x^2 \sec^{-1}(\sqrt{x}) dx$

Optimal result	58
Mathematica [A] (verified)	58
Rubi [A] (verified)	59
Maple [A] (verified)	60
Fricas [A] (verification not implemented)	61
Sympy [C] (verification not implemented)	61
Maxima [A] (verification not implemented)	61
Giac [B] (verification not implemented)	62
Mupad [F(-1)]	63
Reduce [F]	63

Optimal result

Integrand size = 10, antiderivative size = 47

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{3}\sqrt{-1+x} - \frac{2}{9}(-1+x)^{3/2} - \frac{1}{15}(-1+x)^{5/2} + \frac{1}{3}x^3 \sec^{-1}(\sqrt{x})$$

output

```
-1/3*(-1+x)^(1/2)-2/9*(-1+x)^(3/2)-1/15*(-1+x)^(5/2)+1/3*x^3*arcsec(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.74

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{45}\sqrt{-1+x}(8+4x+3x^2) + \frac{1}{3}x^3 \sec^{-1}(\sqrt{x})$$

input

```
Integrate[x^2*ArcSec[Sqrt[x]],x]
```

output

```
-1/45*(Sqrt[-1+x]*(8+4*x+3*x^2))+ (x^3*ArcSec[Sqrt[x]])/3
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5793, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sec^{-1}(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{5793} \\
 & \frac{1}{3} x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{3} \int \frac{x^2}{2\sqrt{x-1}} \, dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{6} \int \frac{x^2}{\sqrt{x-1}} \, dx \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{3} x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{6} \int \left((x-1)^{3/2} + 2\sqrt{x-1} + \frac{1}{\sqrt{x-1}} \right) \, dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3} x^3 \sec^{-1}(\sqrt{x}) + \frac{1}{6} \left(-\frac{2}{5} (x-1)^{5/2} - \frac{4}{3} (x-1)^{3/2} - 2\sqrt{x-1} \right)
 \end{aligned}$$

input `Int [x^2*ArcSec[Sqrt [x]] , x]`

output `(-2*Sqrt[-1 + x] - (4*(-1 + x)^(3/2))/3 - (2*(-1 + x)^(5/2))/5)/6 + (x^3*ArcSec[Sqrt[x]])/3`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5793 `Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.74

method	result	size
parts	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{\sqrt{\frac{-1+x}{x}} \sqrt{x} (3x^2+4x+8)}{45}$	35
derivativedivides	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{(-1+x)(3x^2+4x+8)}{45\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	38
default	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{(-1+x)(3x^2+4x+8)}{45\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	38

input `int(x^2*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/3*x^3*arcsec(x^(1/2))-1/45*((-1+x)/x)^(1/2)*x^(1/2)*(3*x^2+4*x+8)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.57

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{45} (3x^2 + 4x + 8)\sqrt{x-1}$$

input `integrate(x^2*arcsec(x^(1/2)),x, algorithm="fricas")`

output `1/3*x^3*arcsec(sqrt(x)) - 1/45*(3*x^2 + 4*x + 8)*sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 16.42 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.91

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \frac{x^3 \operatorname{asec}(\sqrt{x})}{3} - \frac{\begin{cases} \frac{2x^2\sqrt{x-1}}{5} + \frac{8x\sqrt{x-1}}{15} + \frac{16\sqrt{x-1}}{15} & \text{for } |x| > 1 \\ \frac{2ix^2\sqrt{1-x}}{5} + \frac{8ix\sqrt{1-x}}{15} + \frac{16i\sqrt{1-x}}{15} & \text{otherwise} \end{cases}}{6}$$

input `integrate(x**2*asec(x**(1/2)),x)`

output `x**3*asec(sqrt(x))/3 - Piecewise((2*x**2*sqrt(x - 1)/5 + 8*x*sqrt(x - 1)/15 + 16*sqrt(x - 1)/15, Abs(x) > 1), (2*I*x**2*sqrt(1 - x)/5 + 8*I*x*sqrt(1 - x)/15 + 16*I*sqrt(1 - x)/15, True))/6`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{15} x^{\frac{5}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{arcsec}(\sqrt{x}) - \frac{2}{9} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{3} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

input `integrate(x^2*arcsec(x^(1/2)),x, algorithm="maxima")`

output `-1/15*x^(5/2)*(-1/x + 1)^(5/2) + 1/3*x^3*arcsec(sqrt(x)) - 2/9*x^(3/2)*(-1/x + 1)^(3/2) - 1/3*sqrt(x)*sqrt(-1/x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(31) = 62$.

Time = 0.14 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.47

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{480} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^5 - \frac{5}{288} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3 + \frac{1}{3} x^3 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{5}{48} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) + \frac{150 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^4 + 25 x \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^2 + 3}{1440 x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^5}$$

input `integrate(x^2*arcsec(x^(1/2)),x, algorithm="giac")`

output `-1/480*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 - 5/288*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 1/3*x^3*arccos(1/sqrt(x)) - 5/48*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/1440*(150*x^2*(sqrt(-1/x + 1) - 1)^4 + 25*x*(sqrt(-1/x + 1) - 1)^2 + 3)/(x^(5/2)*(sqrt(-1/x + 1) - 1)^5)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \int x^2 \operatorname{acos}\left(\frac{1}{\sqrt{x}}\right) dx$$

input `int(x^2*acos(1/x^(1/2)),x)`output `int(x^2*acos(1/x^(1/2)), x)`**Reduce [F]**

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \int \operatorname{asec}(\sqrt{x}) x^2 dx$$

input `int(x^2*asec(x^(1/2)),x)`output `int(asec(sqrt(x))*x**2,x)`

3.4 $\int x \sec^{-1}(\sqrt{x}) dx$

Optimal result	64
Mathematica [A] (verified)	64
Rubi [A] (verified)	65
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	67
Sympy [C] (verification not implemented)	67
Maxima [A] (verification not implemented)	67
Giac [B] (verification not implemented)	68
Mupad [F(-1)]	68
Reduce [F]	69

Optimal result

Integrand size = 8, antiderivative size = 36

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{2}\sqrt{-1+x} - \frac{1}{6}(-1+x)^{3/2} + \frac{1}{2}x^2 \sec^{-1}(\sqrt{x})$$

output `-1/2*(-1+x)^(1/2)-1/6*(-1+x)^(3/2)+1/2*x^2*arcsec(x^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{6}\sqrt{-1+x}(2+x) + \frac{1}{2}x^2 \sec^{-1}(\sqrt{x})$$

input `Integrate[x*ArcSec[Sqrt[x]],x]`

output `-1/6*(Sqrt[-1 + x]*(2 + x)) + (x^2*ArcSec[Sqrt[x]])/2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5793, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sec^{-1}(\sqrt{x}) dx \\
 & \quad \downarrow \text{5793} \\
 & \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{2} \int \frac{x}{2\sqrt{x-1}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \frac{x}{\sqrt{x-1}} dx \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \left(\sqrt{x-1} + \frac{1}{\sqrt{x-1}} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) + \frac{1}{4} \left(-\frac{2}{3}(x-1)^{3/2} - 2\sqrt{x-1} \right)
 \end{aligned}$$

input `Int[x*ArcSec[Sqrt[x]],x]`

output `(-2*Sqrt[-1 + x] - (2*(-1 + x)^(3/2))/3)/4 + (x^2*ArcSec[Sqrt[x]])/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5793 `Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

method	result	size
parts	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{\sqrt{\frac{-1+x}{x}} \sqrt{x} (x+2)}{6}$	28
derivativedivides	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{(-1+x)(x+2)}{6\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	31
default	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{(-1+x)(x+2)}{6\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	31

input `int(x*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

output `1/2*x^2*arcsec(x^(1/2))-1/6*((-1+x)/x)^(1/2)*x^(1/2)*(x+2)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int x \sec^{-1}(\sqrt{x}) dx = \frac{1}{2} x^2 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{6} (x+2)\sqrt{x-1}$$

input `integrate(x*arcsec(x^(1/2)),x, algorithm="fricas")`

output `1/2*x^2*arcsec(sqrt(x)) - 1/6*(x + 2)*sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 5.43 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.69

$$\int x \sec^{-1}(\sqrt{x}) dx = \frac{x^2 \operatorname{asec}(\sqrt{x})}{2} - \frac{\begin{cases} \frac{2x\sqrt{x-1}}{3} + \frac{4\sqrt{x-1}}{3} & \text{for } |x| > 1 \\ \frac{2ix\sqrt{1-x}}{3} + \frac{4i\sqrt{1-x}}{3} & \text{otherwise} \end{cases}}{4}$$

input `integrate(x*asec(x**(1/2)),x)`

output `x**2*asec(sqrt(x))/2 - Piecewise((2*x*sqrt(x - 1)/3 + 4*sqrt(x - 1)/3, Abs(x) > 1), (2*I*x*sqrt(1 - x)/3 + 4*I*sqrt(1 - x)/3, True))/4`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{6} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

input `integrate(x*arcsec(x^(1/2)),x, algorithm="maxima")`

output

```
-1/6*x^(3/2)*(-1/x + 1)^(3/2) + 1/2*x^2*arcsec(sqrt(x)) - 1/2*sqrt(x)*sqrt(-1/x + 1)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(24) = 48$.

Time = 0.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.22

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{48} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3 + \frac{1}{2} x^2 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{3}{16} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) + \frac{9x \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^2 + 1}{48 x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3}$$

input

```
integrate(x*arcsec(x^(1/2)),x, algorithm="giac")
```

output

```
-1/48*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 1/2*x^2*arccos(1/sqrt(x)) - 3/16*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/48*(9*x*(sqrt(-1/x + 1) - 1)^2 + 1)/(x^(3/2))*(sqrt(-1/x + 1) - 1)^3)
```

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(\sqrt{x}) dx = \int x \arccos\left(\frac{1}{\sqrt{x}}\right) dx$$

input

```
int(x*acos(1/x^(1/2)),x)
```

output

```
int(x*acos(1/x^(1/2)), x)
```

Reduce [F]

$$\int x \sec^{-1}(\sqrt{x}) dx = \int \operatorname{asec}(\sqrt{x}) x dx$$

input `int(x*asec(x^(1/2)),x)`

output `int(asec(sqrt(x))*x,x)`

3.5 $\int \sec^{-1}(\sqrt{x}) dx$

Optimal result	70
Mathematica [A] (verified)	70
Rubi [A] (verified)	71
Maple [A] (verified)	72
Fricas [A] (verification not implemented)	72
Sympy [C] (verification not implemented)	72
Maxima [A] (verification not implemented)	73
Giac [B] (verification not implemented)	73
Mupad [B] (verification not implemented)	74
Reduce [F]	74

Optimal result

Integrand size = 6, antiderivative size = 18

$$\int \sec^{-1}(\sqrt{x}) dx = -\sqrt{-1+x} + x \sec^{-1}(\sqrt{x})$$

output `-(-1+x)^(1/2)+x*arcsec(x^(1/2))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sec^{-1}(\sqrt{x}) dx = -\sqrt{-1+x} + x \sec^{-1}(\sqrt{x})$$

input `Integrate[ArcSec[Sqrt[x]],x]`

output `-Sqrt[-1 + x] + x*ArcSec[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5791, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{-1}(\sqrt{x}) dx$$

$$\downarrow 5791$$

$$x \sec^{-1}(\sqrt{x}) - \int \frac{1}{2\sqrt{x-1}} dx$$

$$\downarrow 17$$

$$x \sec^{-1}(\sqrt{x}) - \sqrt{x-1}$$

input `Int[ArcSec[Sqrt[x]], x]`

output `-Sqrt[-1 + x] + x*ArcSec[Sqrt[x]]`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_)^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 5791 `Int[ArcSec[u_], x_Symbol] :> Simp[x*ArcSec[u], x] - Simp[u/Sqrt[u^2] Int[SimplifyIntegrand[x*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

method	result	size
parts	$x \operatorname{arcsec}(\sqrt{x}) - \sqrt{x} \sqrt{\frac{-1+x}{x}}$	22
derivativedivides	$x \operatorname{arcsec}(\sqrt{x}) - \frac{-1+x}{\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	25
default	$x \operatorname{arcsec}(\sqrt{x}) - \frac{-1+x}{\sqrt{\frac{-1+x}{x}} \sqrt{x}}$	25

input `int(arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

output `x*arcsec(x^(1/2))-x^(1/2)*((-1+x)/x)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{arcsec}(\sqrt{x}) - \sqrt{x-1}$$

input `integrate(arcsec(x^(1/2)),x, algorithm="fricas")`

output `x*arcsec(sqrt(x)) - sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.61

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{asec}(\sqrt{x}) - \frac{\begin{cases} 2\sqrt{x-1} & \text{for } |x| > 1 \\ 2i\sqrt{1-x} & \text{otherwise} \end{cases}}{2}$$

input `integrate(asec(x**(1/2)),x)`

output `x*asec(sqrt(x)) - Piecewise((2*sqrt(x - 1), Abs(x) > 1), (2*I*sqrt(1 - x), True))/2`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.17

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{arcsec}(\sqrt{x}) - \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

input `integrate(arcsec(x^(1/2)),x, algorithm="maxima")`

output `x*arcsec(sqrt(x)) - sqrt(x)*sqrt(-1/x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(14) = 28$.

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.28

$$\int \sec^{-1}(\sqrt{x}) dx = x \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{1}{2} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) + \frac{1}{2 \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)}$$

input `integrate(arcsec(x^(1/2)),x, algorithm="giac")`

output `x*arccos(1/sqrt(x)) - 1/2*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/2/(sqrt(x)*(sqrt(-1/x + 1) - 1))`

Mupad [B] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.17

$$\int \sec^{-1}(\sqrt{x}) dx = x \arccos\left(\frac{1}{\sqrt{x}}\right) - \sqrt{x} \sqrt{1 - \frac{1}{x}}$$

input `int(acos(1/x^(1/2)),x)`

output `x*acos(1/x^(1/2)) - x^(1/2)*(1 - 1/x)^(1/2)`

Reduce [F]

$$\int \sec^{-1}(\sqrt{x}) dx = \int \operatorname{asec}(\sqrt{x}) dx$$

input `int(asec(x^(1/2)),x)`

output `int(asec(sqrt(x)),x)`

3.6 $\int \frac{\sec^{-1}(\sqrt{x})}{x} dx$

Optimal result	75
Mathematica [A] (verified)	75
Rubi [A] (warning: unable to verify)	76
Maple [A] (verified)	78
Fricas [F]	79
Sympy [F]	79
Maxima [F]	79
Giac [F(-2)]	80
Mupad [F(-1)]	80
Reduce [F]	80

Optimal result

Integrand size = 10, antiderivative size = 56

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = i \sec^{-1}(\sqrt{x})^2 - 2 \sec^{-1}(\sqrt{x}) \log\left(1 + e^{2i \sec^{-1}(\sqrt{x})}\right) + i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(\sqrt{x})}\right)$$

output

```
I*arcsec(x^(1/2))^2-2*arcsec(x^(1/2))*ln(1+(1/x^(1/2)+I*(1-1/x)^(1/2))^2)+
I*polylog(2,-(1/x^(1/2)+I*(1-1/x)^(1/2))^2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = i \left(\sec^{-1}(\sqrt{x}) \left(\sec^{-1}(\sqrt{x}) + 2i \log\left(1 + e^{2i \sec^{-1}(\sqrt{x})}\right) \right) + \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(\sqrt{x})}\right) \right)$$

input

```
Integrate[ArcSec[Sqrt[x]]/x,x]
```

output

```
I*(ArcSec[Sqrt[x]]*(ArcSec[Sqrt[x]] + (2*I)*Log[1 + E^((2*I)*ArcSec[Sqrt[x]])])) + PolyLog[2, -E^((2*I)*ArcSec[Sqrt[x]])]
```

Rubi [A] (warning: unable to verify)

Time = 0.46 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {7267, 5741, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(\sqrt{x})}{x} dx \\
 & \quad \downarrow \text{7267} \\
 & 2 \int \frac{\sec^{-1}(\sqrt{x})}{\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{5741} \\
 & -2 \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{\sqrt{x}} d\frac{1}{\sqrt{x}} \\
 & \quad \downarrow \text{5137} \\
 & 2 \int \sqrt{1 - \frac{1}{x}} \sqrt{x} \arccos\left(\frac{1}{\sqrt{x}}\right) d\arccos\left(\frac{1}{\sqrt{x}}\right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \arccos\left(\frac{1}{\sqrt{x}}\right) \tan\left(\arccos\left(\frac{1}{\sqrt{x}}\right)\right) d\arccos\left(\frac{1}{\sqrt{x}}\right) \\
 & \quad \downarrow \text{4202} \\
 & 2 \left(\frac{ix}{2} - 2i \int \frac{e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)} \arccos\left(\frac{1}{\sqrt{x}}\right)}{1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}} d\arccos\left(\frac{1}{\sqrt{x}}\right) \right) \\
 & \quad \downarrow \text{2620}
 \end{aligned}$$

$$\begin{aligned}
& 2\left(\frac{ix}{2} - 2i\left(\frac{1}{2}i \int \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) d\arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{1}{2}i \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right)\right)\right) \\
& \quad \downarrow \text{2715} \\
& 2\left(\frac{ix}{2} - 2i\left(\frac{1}{4} \int e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)} \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) de^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)} - \frac{1}{2}i \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right)\right)\right) \\
& \quad \downarrow \text{2838} \\
& 2\left(\frac{ix}{2} - 2i\left(-\frac{1}{4} \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) - \frac{1}{2}i \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right)\right)\right)
\end{aligned}$$

input `Int[ArcSec[Sqrt[x]]/x, x]`

output `2*((I/2)*x - (2*I)*((-1/2*I)*ArcCos[1/Sqrt[x]]*Log[1 + E^((2*I)*ArcCos[1/Sqrt[x]])] - PolyLog[2, -E^((2*I)*ArcCos[1/Sqrt[x]])]/4))`

Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^((n_)*((c_) + (d_)*(x_))^(m_)))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^((n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^((n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 5137 `Int[((a_.) + ArcCos[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 5741 `Int[((a_.) + ArcSec[(c_.)*(x_)])*(b_.)/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result
derivativedivides	$i \operatorname{arcsec}(\sqrt{x})^2 - 2 \operatorname{arcsec}(\sqrt{x}) \ln \left(1 + \left(\frac{1}{\sqrt{x}} + i \sqrt{1 - \frac{1}{x}} \right)^2 \right) + i \operatorname{polylog} \left(2, -\left(\frac{1}{\sqrt{x}} + i \sqrt{1 - \frac{1}{x}} \right)^2 \right)$
default	$i \operatorname{arcsec}(\sqrt{x})^2 - 2 \operatorname{arcsec}(\sqrt{x}) \ln \left(1 + \left(\frac{1}{\sqrt{x}} + i \sqrt{1 - \frac{1}{x}} \right)^2 \right) + i \operatorname{polylog} \left(2, -\left(\frac{1}{\sqrt{x}} + i \sqrt{1 - \frac{1}{x}} \right)^2 \right)$

input `int(arcsec(x^(1/2))/x,x,method=_RETURNVERBOSE)`

output `I*arcsec(x^(1/2))^2-2*arcsec(x^(1/2))*ln(1+(1/x^(1/2)+I*(1-1/x)^(1/2))^2)+I*polylog(2,-(1/x^(1/2)+I*(1-1/x)^(1/2))^2)`

Fricas [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsec}(\sqrt{x})}{x} dx$$

input `integrate(arcsec(x^(1/2))/x,x, algorithm="fricas")`

output `integral(arcsec(sqrt(x))/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x} dx$$

input `integrate(asec(x**(1/2))/x,x)`

output `Integral(asec(sqrt(x))/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsec}(\sqrt{x})}{x} dx$$

input `integrate(arcsec(x^(1/2))/x,x, algorithm="maxima")`

output `integrate(arcsec(sqrt(x))/x, x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \text{Exception raised: NotImplementedError}$$

input `integrate(arcsec(x^(1/2))/x,x, algorithm="giac")`

output `Exception raised: NotImplementedError >> unable to parse Giac output: Invalid series expansion: non tractable function acos at +infinity`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

input `int(arccos(1/x^(1/2))/x,x)`

output `int(arccos(1/x^(1/2))/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x} dx$$

input `int(asec(x^(1/2))/x,x)`

output `int(asec(sqrt(x))/x,x)`

3.7 $\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	81
Mathematica [A] (verified)	81
Rubi [A] (verified)	82
Maple [A] (verified)	84
Fricas [A] (verification not implemented)	84
Sympy [C] (verification not implemented)	85
Maxima [A] (verification not implemented)	85
Giac [A] (verification not implemented)	86
Mupad [B] (verification not implemented)	86
Reduce [F]	86

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1+x}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{2} \arctan(\sqrt{-1+x})$$

output `1/2*(-1+x)^(1/2)/x-arcsec(x^(1/2))/x+1/2*arctan((-1+x)^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1+x} - 2 \sec^{-1}(\sqrt{x}) - x \arcsin\left(\frac{1}{\sqrt{x}}\right)}{2x}$$

input `Integrate[ArcSec[Sqrt[x]]/x^2,x]`

output `(Sqrt[-1 + x] - 2*ArcSec[Sqrt[x]] - x*ArcSin[1/Sqrt[x]])/(2*x)`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5793, 27, 52, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx \\
 & \quad \downarrow \text{5793} \\
 & \int \frac{1}{2\sqrt{x-1}x^2} dx - \frac{\sec^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{x-1}x^2} dx - \frac{\sec^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\sqrt{x-1}x} dx + \frac{\sqrt{x-1}}{x} \right) - \frac{\sec^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(\int \frac{1}{x} d\sqrt{x-1} + \frac{\sqrt{x-1}}{x} \right) - \frac{\sec^{-1}(\sqrt{x})}{x} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(\arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{x} \right) - \frac{\sec^{-1}(\sqrt{x})}{x}
 \end{aligned}$$

input `Int[ArcSec[Sqrt[x]]/x^2,x]`

output `-(ArcSec[Sqrt[x]]/x) + (Sqrt[-1 + x]/x + ArcTan[Sqrt[-1 + x]])/2`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 52 $\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)}/((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2)/((b*c - a*d)*(m + 1))) \text{ Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{ILTQ}[m, -1] \ \&\& \ \text{FractionQ}[n] \ \&\& \ \text{LtQ}[n, 0]$
- rule 73 $\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 216 $\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$
- rule 5793 $\text{Int}[(a_.) + \text{ArcSec}[u]*(b_.)*((c_.) + (d_.)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{(m + 1)}*((a + b*\text{ArcSec}[u])/((d*(m + 1))), x] - \text{Simp}[b*(u/((d*(m + 1)*\text{Sqrt}[u^2])) \text{ Int}[\text{SimplifyIntegrand}[(c + d*x)^{(m + 1)}*(D[u, x]/(u*\text{Sqrt}[u^2 - 1])), x], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{InverseFunctionFreeQ}[u, x] \ \&\& \ !\text{FunctionOfQ}[(c + d*x)^{(m + 1)}, u, x] \ \&\& \ !\text{FunctionOfExponentialQ}[u, x]$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

method	result	size
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} + \frac{\sqrt{\frac{-1+x}{x}} (\arctan(\sqrt{-1+x})x + \sqrt{-1+x})}{2\sqrt{x}\sqrt{-1+x}}$	44
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} - \frac{\sqrt{-1+x} \left(\arctan\left(\frac{1}{\sqrt{-1+x}}\right)x - \sqrt{-1+x} \right)}{2\sqrt{\frac{-1+x}{x}} x^{\frac{3}{2}}}$	46
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} - \frac{\sqrt{-1+x} \left(\arctan\left(\frac{1}{\sqrt{-1+x}}\right)x - \sqrt{-1+x} \right)}{2\sqrt{\frac{-1+x}{x}} x^{\frac{3}{2}}}$	46

input `int(arcsec(x^(1/2))/x^2,x,method=_RETURNVERBOSE)`output $-\operatorname{arcsec}(x^{1/2})/x + 1/2 * ((-1+x)/x)^{1/2} / x^{1/2} * (\arctan((-1+x)^{1/2}) * x + (-1+x)^{1/2}) / (-1+x)^{1/2}$ **Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.50

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{(x-2) \operatorname{arcsec}(\sqrt{x}) + \sqrt{x-1}}{2x}$$

input `integrate(arcsec(x^(1/2))/x^2,x, algorithm="fricas")`output $1/2 * ((x - 2) * \operatorname{arcsec}(\operatorname{sqrt}(x)) + \operatorname{sqrt}(x - 1)) / x$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 11.79 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.97

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\begin{cases} i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) - \frac{i}{\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{i}{x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right) + \frac{\sqrt{1-\frac{1}{x}}}{\sqrt{x}} & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{asec}(\sqrt{x})}{x}$$

input `integrate(asec(x**(1/2))/x**2,x)`

output `Piecewise((I*acosh(1/sqrt(x)) - I/(sqrt(x)*sqrt(-1 + 1/x)) + I/(x**(3/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-asin(1/sqrt(x)) + sqrt(1 - 1/x)/sqrt(x), True))/2 - asecc(sqrt(x))/x`

Maxima [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = -\frac{\sqrt{x}\sqrt{-\frac{1}{x}+1}}{2\left(x\left(\frac{1}{x}-1\right)-1\right)} - \frac{\operatorname{arcsec}(\sqrt{x})}{x} + \frac{1}{2} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x}+1}\right)$$

input `integrate(arcsec(x^(1/2))/x^2,x, algorithm="maxima")`

output `-1/2*sqrt(x)*sqrt(-1/x + 1)/(x*(1/x - 1) - 1) - arcsec(sqrt(x))/x + 1/2*arctan(sqrt(x)*sqrt(-1/x + 1))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-\frac{1}{x} + 1}}{2\sqrt{x}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x} + \frac{1}{2} \arccos\left(\frac{1}{\sqrt{x}}\right)$$

input `integrate(arcsec(x^(1/2))/x^2,x, algorithm="giac")`output `1/2*sqrt(-1/x + 1)/sqrt(x) - arccos(1/sqrt(x))/x + 1/2*arccos(1/sqrt(x))`**Mupad [B] (verification not implemented)**

Time = 0.82 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{1 - \frac{1}{x}}}{2\sqrt{x}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right) \left(\frac{2}{x} - 1\right)}{2}$$

input `int(acos(1/x^(1/2))/x^2,x)`output `(1 - 1/x)^(1/2)/(2*x^(1/2)) - (acos(1/x^(1/2))*(2/x - 1))/2`**Reduce [F]**

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x^2} dx$$

input `int(asec(x^(1/2))/x^2,x)`output `int(asec(sqrt(x))/x**2,x)`

3.8 $\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	87
Mathematica [A] (verified)	87
Rubi [A] (verified)	88
Maple [A] (verified)	90
Fricas [A] (verification not implemented)	90
Sympy [C] (verification not implemented)	91
Maxima [B] (verification not implemented)	91
Giac [A] (verification not implemented)	92
Mupad [F(-1)]	92
Reduce [F]	93

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{-1+x}}{8x^2} + \frac{3\sqrt{-1+x}}{16x} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \arctan(\sqrt{-1+x})$$

output `1/8*(-1+x)^(1/2)/x^2+3/16*(-1+x)^(1/2)/x-1/2*arcsec(x^(1/2))/x^2+3/16*arctan((-1+x)^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.02

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \left(\frac{1}{8x^{3/2}} + \frac{3}{16\sqrt{x}} \right) \sqrt{\frac{-1+x}{x}} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16} \arcsin\left(\frac{1}{\sqrt{x}}\right)$$

input `Integrate[ArcSec[Sqrt[x]]/x^3,x]`

output `(1/(8*x^(3/2)) + 3/(16*Sqrt[x]))*Sqrt[(-1 + x)/x] - ArcSec[Sqrt[x]]/(2*x^2) - (3*ArcSin[1/Sqrt[x]])/16`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5793, 27, 52, 52, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx \\
 & \quad \downarrow \text{5793} \\
 & \frac{1}{2} \int \frac{1}{2\sqrt{x-1}x^3} dx - \frac{\sec^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} \int \frac{1}{\sqrt{x-1}x^3} dx - \frac{\sec^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{4} \left(\frac{3}{4} \int \frac{1}{\sqrt{x-1}x^2} dx + \frac{\sqrt{x-1}}{2x^2} \right) - \frac{\sec^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{4} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{x-1}} dx + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) - \frac{\sec^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{4} \left(\frac{3}{4} \left(\int \frac{1}{x} d\sqrt{x-1} + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) - \frac{\sec^{-1}(\sqrt{x})}{2x^2} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{4} \left(\frac{3}{4} \left(\arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) - \frac{\sec^{-1}(\sqrt{x})}{2x^2}
 \end{aligned}$$

input `Int[ArcSec[Sqrt[x]]/x^3,x]`

output

```
-1/2*ArcSec[Sqrt[x]]/x^2 + (Sqrt[-1 + x]/(2*x^2) + (3*(Sqrt[-1 + x]/x + ArcTan[Sqrt[-1 + x]]))/4)/4
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 52

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntegerQ[a, b, c, d, m, n, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 5793

```
Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

method	result	size
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} - \frac{\sqrt{-1+x} \left(3 \arctan\left(\frac{1}{\sqrt{-1+x}}\right) x^2 - 3\sqrt{-1+x} x - 2\sqrt{-1+x} \right)}{16\sqrt{\frac{-1+x}{x}} x^{\frac{5}{2}}}$	57
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} - \frac{\sqrt{-1+x} \left(3 \arctan\left(\frac{1}{\sqrt{-1+x}}\right) x^2 - 3\sqrt{-1+x} x - 2\sqrt{-1+x} \right)}{16\sqrt{\frac{-1+x}{x}} x^{\frac{5}{2}}}$	57
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} + \frac{\sqrt{\frac{-1+x}{x}} \left(3 \arctan(\sqrt{-1+x}) x^2 + 3\sqrt{-1+x} x + 2\sqrt{-1+x} \right)}{16x^{\frac{3}{2}} \sqrt{-1+x}}$	57

input `int(arcsec(x^(1/2))/x^3,x,method=_RETURNVERBOSE)`output `-1/2*arcsec(x^(1/2))/x^2-1/16*(-1+x)^(1/2)*(3*arctan(1/(-1+x)^(1/2))*x^2-3*(-1+x)^(1/2)*x-2*(-1+x)^(1/2))/((-1+x)/x)^(1/2)/x^(5/2)`**Fricas [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.54

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{(3x^2 - 8) \operatorname{arcsec}(\sqrt{x}) + (3x + 2)\sqrt{x-1}}{16x^2}$$

input `integrate(arcsec(x^(1/2))/x^3,x, algorithm="fricas")`output `1/16*((3*x^2 - 8)*arcsec(sqrt(x)) + (3*x + 2)*sqrt(x - 1))/x^2`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 38.07 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.67

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{\begin{cases} \frac{3i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{4} - \frac{3i}{4\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{i}{4x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{2x^{\frac{5}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\frac{3 \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{4} + \frac{3}{4\sqrt{x}\sqrt{1-\frac{1}{x}}} - \frac{1}{4x^{\frac{3}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{2x^{\frac{5}{2}}\sqrt{1-\frac{1}{x}}} & \text{otherwise} \end{cases}}{4} - \frac{\operatorname{asec}(\sqrt{x})}{2x^2}$$

input `integrate(asec(x**(1/2))/x**3,x)`

output `Piecewise((3*I*acosh(1/sqrt(x))/4 - 3*I/(4*sqrt(x)*sqrt(-1 + 1/x)) + I/(4*x**(3/2)*sqrt(-1 + 1/x)) + I/(2*x**(5/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-3*asin(1/sqrt(x))/4 + 3/(4*sqrt(x)*sqrt(1 - 1/x)) - 1/(4*x**(3/2)*sqrt(1 - 1/x)) - 1/(2*x**(5/2)*sqrt(1 - 1/x)), True))/4 - asecc(sqrt(x))/(2*x**2)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(38) = 76.

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.48

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{3x^{\frac{3}{2}}\left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + 5\sqrt{x}\sqrt{-\frac{1}{x} + 1}}{16\left(x^2\left(\frac{1}{x} - 1\right)^2 - 2x\left(\frac{1}{x} - 1\right) + 1\right)} - \frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} + \frac{3}{16} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x} + 1}\right)$$

input `integrate(arcsec(x^(1/2))/x^3,x, algorithm="maxima")`

output

```
1/16*(3*x^(3/2)*(-1/x + 1)^(3/2) + 5*sqrt(x)*sqrt(-1/x + 1))/(x^2*(1/x - 1)
)^2 - 2*x*(1/x - 1) + 1) - 1/2*arcsec(sqrt(x))/x^2 + 3/16*arctan(sqrt(x)*s
qrt(-1/x + 1))
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{3\sqrt{-\frac{1}{x}+1}}{16\sqrt{x}} + \frac{\sqrt{-\frac{1}{x}+1}}{8x^{\frac{3}{2}}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{2x^2} + \frac{3}{16}\arccos\left(\frac{1}{\sqrt{x}}\right)$$

input

```
integrate(arcsec(x^(1/2))/x^3,x, algorithm="giac")
```

output

```
3/16*sqrt(-1/x + 1)/sqrt(x) + 1/8*sqrt(-1/x + 1)/x^(3/2) - 1/2*arccos(1/sq
rt(x))/x^2 + 3/16*arccos(1/sqrt(x))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

input

```
int(acos(1/x^(1/2))/x^3,x)
```

output

```
int(acos(1/x^(1/2))/x^3, x)
```

Reduce [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x^3} dx$$

input `int(asec(x^(1/2))/x^3,x)`

output `int(asec(sqrt(x))/x**3,x)`

3.9 $\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	94
Mathematica [A] (verified)	94
Rubi [A] (verified)	95
Maple [A] (verified)	97
Fricas [A] (verification not implemented)	97
Sympy [C] (verification not implemented)	98
Maxima [B] (verification not implemented)	98
Giac [A] (verification not implemented)	99
Mupad [F(-1)]	99
Reduce [F]	100

Optimal result

Integrand size = 10, antiderivative size = 68

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} + \frac{5\sqrt{-1+x}}{48x} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{48} \arctan(\sqrt{-1+x})$$

output `1/18*(-1+x)^(1/2)/x^3+5/72*(-1+x)^(1/2)/x^2+5/48*(-1+x)^(1/2)/x-1/3*arcsec(x^(1/2))/x^3+5/48*arctan((-1+x)^(1/2))`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.66

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1+x}(8+10x+15x^2) - 48\sec^{-1}(\sqrt{x}) - 15x^3 \arcsin\left(\frac{1}{\sqrt{x}}\right)}{144x^3}$$

input `Integrate[ArcSec[Sqrt[x]]/x^4,x]`

output `(Sqrt[-1+x]*(8+10*x+15*x^2)-48*ArcSec[Sqrt[x]]-15*x^3*ArcSin[1/Sqrt[x]])/(144*x^3)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5793, 27, 52, 52, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx \\
 & \quad \downarrow \text{5793} \\
 & \frac{1}{3} \int \frac{1}{2\sqrt{x-1}x^4} dx - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{6} \int \frac{1}{\sqrt{x-1}x^4} dx - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{6} \left(\frac{5}{6} \int \frac{1}{\sqrt{x-1}x^3} dx + \frac{\sqrt{x-1}}{3x^3} \right) - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{6} \left(\frac{5}{6} \left(\frac{3}{4} \int \frac{1}{\sqrt{x-1}x^2} dx + \frac{\sqrt{x-1}}{2x^2} \right) + \frac{\sqrt{x-1}}{3x^3} \right) - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{6} \left(\frac{5}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{1}{\sqrt{x-1}x} dx + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) + \frac{\sqrt{x-1}}{3x^3} \right) - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{6} \left(\frac{5}{6} \left(\frac{3}{4} \left(\int \frac{1}{x} d\sqrt{x-1} + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) + \frac{\sqrt{x-1}}{3x^3} \right) - \frac{\sec^{-1}(\sqrt{x})}{3x^3} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{6} \left(\frac{5}{6} \left(\frac{3}{4} \left(\arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{x} \right) + \frac{\sqrt{x-1}}{2x^2} \right) + \frac{\sqrt{x-1}}{3x^3} \right) - \frac{\sec^{-1}(\sqrt{x})}{3x^3}
 \end{aligned}$$

input `Int[ArcSec[Sqrt[x]]/x^4,x]`

output `-1/3*ArcSec[Sqrt[x]]/x^3 + (Sqrt[-1 + x]/(3*x^3) + (5*(Sqrt[-1 + x]/(2*x^2) + (3*(Sqrt[-1 + x]/x + ArcTan[Sqrt[-1 + x]))/4))/6)/6`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 52 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntegerQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5793 `Int[((a_.) + ArcSec[u]*(b_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Simp[b*(u/(d*(m + 1)*Sqrt[u^2])) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.99

method	result	size
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} - \frac{\sqrt{-1+x} \left(15 \arctan\left(\frac{1}{\sqrt{-1+x}}\right) x^3 - 15\sqrt{-1+x} x^2 - 10\sqrt{-1+x} x - 8\sqrt{-1+x} \right)}{144 \sqrt{\frac{-1+x}{x}} x^{\frac{7}{2}}}$	67
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} - \frac{\sqrt{-1+x} \left(15 \arctan\left(\frac{1}{\sqrt{-1+x}}\right) x^3 - 15\sqrt{-1+x} x^2 - 10\sqrt{-1+x} x - 8\sqrt{-1+x} \right)}{144 \sqrt{\frac{-1+x}{x}} x^{\frac{7}{2}}}$	67
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} + \frac{\sqrt{\frac{-1+x}{x}} \left(15 \arctan(\sqrt{-1+x}) x^3 + 15\sqrt{-1+x} x^2 + 10\sqrt{-1+x} x + 8\sqrt{-1+x} \right)}{144 x^{\frac{5}{2}} \sqrt{-1+x}}$	67

input `int(arcsec(x^(1/2))/x^4,x,method=_RETURNVERBOSE)`

output
$$-1/3*\operatorname{arcsec}(x^{(1/2)})/x^3 - 1/144*(-1+x)^{(1/2)}*(15*\arctan(1/(-1+x)^{(1/2)})*x^3 - 15*(-1+x)^{(1/2)}*x^2 - 10*(-1+x)^{(1/2)}*x - 8*(-1+x)^{(1/2)})/(((-1+x)/x)^{(1/2)}/x^{(7/2)})$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{3(5x^3 - 16) \operatorname{arcsec}(\sqrt{x}) + (15x^2 + 10x + 8)\sqrt{x-1}}{144x^3}$$

input `integrate(arcsec(x^(1/2))/x^4,x, algorithm="fricas")`

output
$$1/144*(3*(5*x^3 - 16)*\operatorname{arcsec}(\operatorname{sqrt}(x)) + (15*x^2 + 10*x + 8)*\operatorname{sqrt}(x - 1))/x^3$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 108.73 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.65

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$$

$$= \frac{\begin{cases} \frac{5i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{8} - \frac{5i}{8\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{5i}{24x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{12x^{\frac{5}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{3x^{\frac{7}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\frac{5 \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{8} + \frac{5}{8\sqrt{x}\sqrt{1-\frac{1}{x}}} - \frac{5}{24x^{\frac{3}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{12x^{\frac{5}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{3x^{\frac{7}{2}}\sqrt{1-\frac{1}{x}}} & \text{otherwise} \end{cases}}{6} - \frac{\operatorname{asec}(\sqrt{x})}{3x^3}$$

input `integrate(asec(x**(1/2))/x**4,x)`

output `Piecewise((5*I*acosh(1/sqrt(x))/8 - 5*I/(8*sqrt(x)*sqrt(-1 + 1/x)) + 5*I/(24*x**(3/2)*sqrt(-1 + 1/x)) + I/(12*x**(5/2)*sqrt(-1 + 1/x)) + I/(3*x**(7/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-5*asin(1/sqrt(x))/8 + 5/(8*sqrt(x)*sqrt(1 - 1/x)) - 5/(24*x**(3/2)*sqrt(1 - 1/x)) - 1/(12*x**(5/2)*sqrt(1 - 1/x)) - 1/(3*x**(7/2)*sqrt(1 - 1/x)), True))/6 - asecc(sqrt(x))/(3*x**3)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(48) = 96.

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.56

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = -\frac{15x^{\frac{5}{2}}\left(-\frac{1}{x}+1\right)^{\frac{5}{2}}+40x^{\frac{3}{2}}\left(-\frac{1}{x}+1\right)^{\frac{3}{2}}+33\sqrt{x}\sqrt{-\frac{1}{x}+1}}{144\left(x^3\left(\frac{1}{x}-1\right)^3-3x^2\left(\frac{1}{x}-1\right)^2+3x\left(\frac{1}{x}-1\right)-1\right)} - \frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} + \frac{5}{48} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x}+1}\right)$$

input `integrate(arcsec(x^(1/2))/x^4,x, algorithm="maxima")`

output

```
-1/144*(15*x^(5/2)*(-1/x + 1)^(5/2) + 40*x^(3/2)*(-1/x + 1)^(3/2) + 33*sqrt(x)*sqrt(-1/x + 1))/(x^3*(1/x - 1)^3 - 3*x^2*(1/x - 1)^2 + 3*x*(1/x - 1) - 1) - 1/3*arcsec(sqrt(x))/x^3 + 5/48*arctan(sqrt(x)*sqrt(-1/x + 1))
```

Giac [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.85

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{5\sqrt{-\frac{1}{x}+1}}{48\sqrt{x}} + \frac{5\sqrt{-\frac{1}{x}+1}}{72x^{\frac{3}{2}}} + \frac{\sqrt{-\frac{1}{x}+1}}{18x^{\frac{5}{2}}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{3x^3} + \frac{5}{48}\arccos\left(\frac{1}{\sqrt{x}}\right)$$

input

```
integrate(arcsec(x^(1/2))/x^4,x, algorithm="giac")
```

output

```
5/48*sqrt(-1/x + 1)/sqrt(x) + 5/72*sqrt(-1/x + 1)/x^(3/2) + 1/18*sqrt(-1/x + 1)/x^(5/2) - 1/3*arccos(1/sqrt(x))/x^3 + 5/48*arccos(1/sqrt(x))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

input

```
int(arccos(1/x^(1/2))/x^4,x)
```

output

```
int(arccos(1/x^(1/2))/x^4, x)
```

Reduce [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x^4} dx$$

input `int(asec(x^(1/2))/x^4,x)`

output `int(asec(sqrt(x))/x**4,x)`

3.10 $\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	101
Mathematica [A] (verified)	101
Rubi [A] (verified)	102
Maple [A] (verified)	103
Fricas [A] (verification not implemented)	104
Sympy [A] (verification not implemented)	104
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [F(-1)]	105
Reduce [F]	106

Optimal result

Integrand size = 10, antiderivative size = 56

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{3}a^3 \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{9}a^3 \left(1 - \frac{x^2}{a^2}\right)^{3/2} + \frac{1}{3}x^3 \arccos \left(\frac{x}{a} \right)$$

output

```
-1/3*a^3*(1-x^2/a^2)^(1/2)+1/9*a^3*(1-x^2/a^2)^(3/2)+1/3*x^3*arccos(x/a)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.75

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{9}a(2a^2 + x^2) \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{3}x^3 \sec^{-1} \left(\frac{a}{x} \right)$$

input

```
Integrate[x^2*ArcSec[a/x],x]
```

output

```
-1/9*(a*(2*a^2 + x^2)*Sqrt[1 - x^2/a^2]) + (x^3*ArcSec[a/x])/3
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5787, 5139, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx \\
 & \quad \downarrow \text{5787} \\
 & \int x^2 \arccos\left(\frac{x}{a}\right) dx \\
 & \quad \downarrow \text{5139} \\
 & \frac{\int \frac{x^3}{\sqrt{1-\frac{x^2}{a^2}}} dx}{3a} + \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) \\
 & \quad \downarrow \text{243} \\
 & \frac{\int \frac{x^2}{\sqrt{1-\frac{x^2}{a^2}}} dx^2}{6a} + \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) \\
 & \quad \downarrow \text{53} \\
 & \frac{\int \left(\frac{a^2}{\sqrt{1-\frac{x^2}{a^2}}} - a^2 \sqrt{1-\frac{x^2}{a^2}} \right) dx^2}{6a} + \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{2}{3}a^4 \left(1 - \frac{x^2}{a^2}\right)^{3/2} - 2a^4 \sqrt{1 - \frac{x^2}{a^2}}}{6a} + \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right)
 \end{aligned}$$

input

```
Int [x^2*ArcSec [a/x] , x]
```

output

```
(-2*a^4*Sqrt [1 - x^2/a^2] + (2*a^4*(1 - x^2/a^2)^(3/2))/3)/(6*a) + (x^3*ArcCos [x/a])/3
```

Definitions of rubi rules used

- rule 53 $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$
- rule 243 $\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 5139 $\text{Int}[(a_.) + \text{ArcCos}[(c_.)(x_)]*(b_.)^{(n_.)}((d_.)(x_)^{(m_.)}), x_Symbol] \rightarrow \text{Simp}[(d*x)^{m+1}*((a + b*\text{ArcCos}[c*x])^n/(d*(m+1))), x] + \text{Simp}[b*c*(n/(d*(m+1))) \ \text{Int}[(d*x)^{m+1}*((a + b*\text{ArcCos}[c*x])^{n-1}/\text{Sqrt}[1 - c^2*x^2]), x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{NeQ}[m, -1]$
- rule 5787 $\text{Int}[\text{ArcSec}[(c_.)/((a_.) + (b_.)(x_)^{(n_.)})]^{(m_.)}(u_.), x_Symbol] \rightarrow \text{Int}[u*\text{ArcCos}[a/c + b*(x^n/c)]^m, x] /; \text{FreeQ}[\{a, b, c, n, m\}, x]$

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

method	result	size
parts	$\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3} + \frac{-\frac{x^2 a^2 \sqrt{1-\frac{x^2}{a^2}}}{3} - \frac{2a^4 \sqrt{1-\frac{x^2}{a^2}}}{3}}{3a}$	56
derivativedivides	$-a^3 \left(-\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3a^3} + \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66
default	$-a^3 \left(-\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3a^3} + \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66

input `int(x^2*arcsec(a/x),x,method=_RETURNVERBOSE)`

output `1/3*x^3*arcsec(a/x)+1/3/a*(-1/3*x^2*a^2*(1-x^2/a^2)^(1/2)-2/3*a^4*(1-x^2/a^2)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.70

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{3} x^3 \operatorname{arcsec} \left(\frac{a}{x} \right) - \frac{1}{9} (2a^2x + x^3) \sqrt{\frac{a^2 - x^2}{x^2}}$$

input `integrate(x^2*arcsec(a/x),x, algorithm="fricas")`

output `1/3*x^3*arcsec(a/x) - 1/9*(2*a^2*x + x^3)*sqrt((a^2 - x^2)/x^2)`

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \begin{cases} -\frac{2a^3 \sqrt{1-\frac{x^2}{a^2}}}{9} - \frac{ax^2 \sqrt{1-\frac{x^2}{a^2}}}{9} + \frac{x^3 \operatorname{asec} \left(\frac{a}{x} \right)}{3} & \text{for } a \neq 0 \\ \tilde{\infty} x^3 & \text{otherwise} \end{cases}$$

input `integrate(x**2*asec(a/x),x)`

output `Piecewise((-2*a**3*sqrt(1 - x**2/a**2)/9 - a*x**2*sqrt(1 - x**2/a**2)/9 + x**3*asec(a/x)/3, Ne(a, 0)), (zoo*x**3, True))`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{3} x^3 \operatorname{arcsec} \left(\frac{a}{x} \right) - \frac{2a^4 \sqrt{-\frac{x^2}{a^2} + 1} + a^2 x^2 \sqrt{-\frac{x^2}{a^2} + 1}}{9a}$$

input `integrate(x^2*arcsec(a/x),x, algorithm="maxima")`output `1/3*x^3*arcsec(a/x) - 1/9*(2*a^4*sqrt(-x^2/a^2 + 1) + a^2*x^2*sqrt(-x^2/a^2 + 1))/a`**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.84

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{3} x^3 \arccos \left(\frac{x}{a} \right) - \frac{2}{9} a^3 \sqrt{-\frac{x^2}{a^2} + 1} - \frac{1}{9} a x^2 \sqrt{-\frac{x^2}{a^2} + 1}$$

input `integrate(x^2*arcsec(a/x),x, algorithm="giac")`output `1/3*x^3*arccos(x/a) - 2/9*a^3*sqrt(-x^2/a^2 + 1) - 1/9*a*x^2*sqrt(-x^2/a^2 + 1)`**Mupad [F(-1)]**

Timed out.

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \begin{cases} \frac{x^3 \operatorname{acos} \left(\frac{x}{a} \right) - \sqrt{a^2 - x^2} (2a^2 + x^2)}{3} & \text{if } 0 < a \\ \int x^2 \operatorname{acos} \left(\frac{x}{a} \right) dx & \text{if } -0 < a \end{cases}$$

input `int(x^2*acos(x/a),x)`

output `piecewise(0 < a, (x^3*acos(x/a))/3 - ((a^2 - x^2)^(1/2)*(2*a^2 + x^2))/9, ~0 < a, int(x^2*acos(x/a), x))`

Reduce [F]

$$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx = \int a \sec\left(\frac{a}{x}\right) x^2 dx$$

input `int(x^2*asec(a/x), x)`

output `int(asec(a/x)*x**2, x)`

3.11 $\int x \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [A] (verified)	109
Fricas [A] (verification not implemented)	110
Sympy [A] (verification not implemented)	111
Maxima [A] (verification not implemented)	111
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	112
Reduce [B] (verification not implemented)	112

Optimal result

Integrand size = 8, antiderivative size = 47

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{4}ax\sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{2}x^2 \arccos \left(\frac{x}{a} \right) + \frac{1}{4}a^2 \arcsin \left(\frac{x}{a} \right)$$

output `-1/4*a*x*(1-x^2/a^2)^(1/2)+1/2*x^2*arccos(x/a)+1/4*a^2*arcsin(x/a)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.94

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{4} \left(-ax\sqrt{1 - \frac{x^2}{a^2}} + 2x^2 \sec^{-1} \left(\frac{a}{x} \right) + a^2 \arcsin \left(\frac{x}{a} \right) \right)$$

input `Integrate[x*ArcSec[a/x],x]`

output `(-(a*x*Sqrt[1 - x^2/a^2]) + 2*x^2*ArcSec[a/x] + a^2*ArcSin[x/a])/4`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5787, 5139, 262, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \sec^{-1}\left(\frac{a}{x}\right) dx \\ & \quad \downarrow \text{5787} \\ & \int x \arccos\left(\frac{x}{a}\right) dx \\ & \quad \downarrow \text{5139} \\ & \frac{\int \frac{x^2}{\sqrt{1-\frac{x^2}{a^2}}} dx}{2a} + \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) \\ & \quad \downarrow \text{262} \\ & \frac{\frac{1}{2}a^2 \int \frac{1}{\sqrt{1-\frac{x^2}{a^2}}} dx - \frac{1}{2}a^2 x \sqrt{1-\frac{x^2}{a^2}}}{2a} + \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) \\ & \quad \downarrow \text{223} \\ & \frac{\frac{1}{2}a^3 \arcsin\left(\frac{x}{a}\right) - \frac{1}{2}a^2 x \sqrt{1-\frac{x^2}{a^2}}}{2a} + \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) \end{aligned}$$

input

```
Int [x*ArcSec [a/x] , x]
```

output

```
(x^2*ArcCos [x/a])/2 + (-1/2*(a^2*x*sqrt [1 - x^2/a^2]) + (a^3*ArcSin [x/a])/2)/(2*a)
```

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 5139 `Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 5787 `Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int[u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]`

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.43

method	result	size
parts	$\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2} + \frac{-\frac{x a^2 \sqrt{1-\frac{x^2}{a^2}}}{2} + \frac{a^2 \arctan\left(\frac{\sqrt{\frac{1}{a^2} x}}{\sqrt{1-\frac{x^2}{a^2}}}\right)}{2\sqrt{\frac{1}{a^2}}}}{2a}$	67
derivativedivides	$-a^2 \left(-\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2a^2} - \frac{\sqrt{\frac{a^2}{x^2}-1} \left(\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2}-1}}\right) a^2}{x^2} - \sqrt{\frac{a^2}{x^2}-1} \right) x^3}{4\sqrt{\frac{(a^2-1)x^2}{a^2}} a^3} \right)$	91
default	$-a^2 \left(-\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2a^2} - \frac{\sqrt{\frac{a^2}{x^2}-1} \left(\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2}-1}}\right) a^2}{x^2} - \sqrt{\frac{a^2}{x^2}-1} \right) x^3}{4\sqrt{\frac{(a^2-1)x^2}{a^2}} a^3} \right)$	91

```
input int(x*arcsec(a/x),x,method=_RETURNVERBOSE)
```

```
output 1/2*x^2*arcsec(a/x)+1/2/a*(-1/2*x*a^2*(1-x^2/a^2)^(1/2)+1/2*a^2/(1/a^2)^(1/2)*arctan((1/a^2)^(1/2)*x/(1-x^2/a^2)^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = -\frac{1}{4} x^2 \sqrt{\frac{a^2-x^2}{x^2}} - \frac{1}{4} (a^2-2x^2) \operatorname{arcsec}\left(\frac{a}{x}\right)$$

```
input integrate(x*arcsec(a/x),x, algorithm="fricas")
```

output `-1/4*x^2*sqrt((a^2 - x^2)/x^2) - 1/4*(a^2 - 2*x^2)*arcsec(a/x)`

Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.87

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \begin{cases} -\frac{a^2 \operatorname{asec} \left(\frac{a}{x} \right)}{4} - \frac{ax \sqrt{1 - \frac{x^2}{a^2}}}{4} + \frac{x^2 \operatorname{asec} \left(\frac{a}{x} \right)}{2} & \text{for } a \neq 0 \\ \tilde{\infty} x^2 & \text{otherwise} \end{cases}$$

input `integrate(x*asec(a/x),x)`

output `Piecewise((-a**2*asec(a/x)/4 - a*x*sqrt(1 - x**2/a**2)/4 + x**2*asec(a/x)/2, Ne(a, 0)), (zoo*x**2, True))`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{2} x^2 \operatorname{arcsec} \left(\frac{a}{x} \right) + \frac{a^3 \arcsin \left(\frac{x}{a} \right) - a^2 x \sqrt{-\frac{x^2}{a^2} + 1}}{4 a}$$

input `integrate(x*arcsec(a/x),x, algorithm="maxima")`

output `1/2*x^2*arcsec(a/x) + 1/4*(a^3*arcsin(x/a) - a^2*x*sqrt(-x^2/a^2 + 1))/a`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = -\frac{1}{4} a^2 \arccos\left(\frac{x}{a}\right) + \frac{1}{2} x^2 \arccos\left(\frac{x}{a}\right) - \frac{1}{4} a x \sqrt{-\frac{x^2}{a^2} + 1}$$

input `integrate(x*arcsec(a/x),x, algorithm="giac")`output `-1/4*a^2*arccos(x/a) + 1/2*x^2*arccos(x/a) - 1/4*a*x*sqrt(-x^2/a^2 + 1)`**Mupad [B] (verification not implemented)**

Time = 0.70 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = \frac{a^2 \arccos\left(\frac{x}{a}\right) \left(\frac{2x^2}{a^2} - 1\right)}{4} - \frac{a x \sqrt{1 - \frac{x^2}{a^2}}}{4}$$

input `int(x*acos(x/a),x)`output `(a^2*acos(x/a)*((2*x^2)/a^2 - 1))/4 - (a*x*(1 - x^2/a^2)^(1/2))/4`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = -\frac{a \sec\left(\frac{a}{x}\right) a^2}{2} + \frac{a \sec\left(\frac{a}{x}\right) x^2}{2} - \frac{a \sin\left(\frac{a}{x}\right) a^2}{4} - \frac{\sqrt{a^2 - x^2} x}{4}$$

input `int(x*asec(a/x),x)`output `(- 2*asec(a/x)*a**2 + 2*asec(a/x)*x**2 - asin(x/a)*a**2 - sqrt(a**2 - x**2)*x)/4`

3.12 $\int \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	113
Mathematica [A] (verified)	113
Rubi [A] (verified)	114
Maple [A] (verified)	115
Fricas [A] (verification not implemented)	115
Sympy [A] (verification not implemented)	116
Maxima [A] (verification not implemented)	116
Giac [A] (verification not implemented)	116
Mupad [B] (verification not implemented)	117
Reduce [B] (verification not implemented)	117

Optimal result

Integrand size = 6, antiderivative size = 26

$$\int \sec^{-1} \left(\frac{a}{x} \right) dx = -a \sqrt{1 - \frac{x^2}{a^2}} + x \arccos \left(\frac{x}{a} \right)$$

output `-a*(1-x^2/a^2)^(1/2)+x*arccos(x/a)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \sec^{-1} \left(\frac{a}{x} \right) dx = -a \sqrt{1 - \frac{x^2}{a^2}} + x \sec^{-1} \left(\frac{a}{x} \right)$$

input `Integrate[ArcSec[a/x], x]`

output `-(a*Sqrt[1 - x^2/a^2]) + x*ArcSec[a/x]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5787, 5131, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sec^{-1}\left(\frac{a}{x}\right) dx \\ & \quad \downarrow \text{5787} \\ & \int \arccos\left(\frac{x}{a}\right) dx \\ & \quad \downarrow \text{5131} \\ & \frac{\int \frac{x}{\sqrt{1-\frac{x^2}{a^2}}} dx}{a} + x \arccos\left(\frac{x}{a}\right) \\ & \quad \downarrow \text{241} \\ & x \arccos\left(\frac{x}{a}\right) - a\sqrt{1-\frac{x^2}{a^2}} \end{aligned}$$

input `Int[ArcSec[a/x], x]`

output `-(a*Sqrt[1 - x^2/a^2]) + x*ArcCos[x/a]`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5131 `Int[((a_.) + ArcCos[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCos[c*x])^n, x] + Simp[b*c*n Int[x*(a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 5787

```
Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

method	result	size
parts	$x \operatorname{arcsec}\left(\frac{a}{x}\right) - a\sqrt{\frac{a^2-x^2}{a^2}}$	28
derivativedivides	$-a\left(-\frac{x \operatorname{arcsec}\left(\frac{a}{x}\right)}{a} + \frac{x^2\left(\frac{a^2}{x^2}-1\right)}{\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}}}\right)$	51
default	$-a\left(-\frac{x \operatorname{arcsec}\left(\frac{a}{x}\right)}{a} + \frac{x^2\left(\frac{a^2}{x^2}-1\right)}{\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}}}\right)$	51

```
input int(arcsec(a/x),x,method=_RETURNVERBOSE)
```

```
output x*arcsec(a/x)-a*((a^2-x^2)/a^2)^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \operatorname{arcsec}\left(\frac{a}{x}\right) - x\sqrt{\frac{a^2-x^2}{x^2}}$$

```
input integrate(arcsec(a/x),x, algorithm="fricas")
```

```
output x*arcsec(a/x) - x*sqrt((a^2 - x^2)/x^2)
```

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = \begin{cases} -a\sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{asec}\left(\frac{a}{x}\right) & \text{for } a \neq 0 \\ \tilde{\infty}x & \text{otherwise} \end{cases}$$

input `integrate(asec(a/x), x)`output `Piecewise((-a*sqrt(1 - x**2/a**2) + x*asec(a/x), Ne(a, 0)), (zoo*x, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \operatorname{arcsec}\left(\frac{a}{x}\right) - a\sqrt{-\frac{x^2}{a^2} + 1}$$

input `integrate(arcsec(a/x), x, algorithm="maxima")`output `x*arcsec(a/x) - a*sqrt(-x^2/a^2 + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = a\left(\frac{x \arccos\left(\frac{x}{a}\right)}{a} - \sqrt{-\frac{x^2}{a^2} + 1}\right)$$

input `integrate(arcsec(a/x), x, algorithm="giac")`output `a*(x*arccos(x/a)/a - sqrt(-x^2/a^2 + 1))`

Mupad [B] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \arccos\left(\frac{x}{a}\right) - a \sqrt{1 - \frac{x^2}{a^2}}$$

input `int(acos(x/a),x)`

output `x*acos(x/a) - a*(1 - x^2/a^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = a \operatorname{asec}\left(\frac{a}{x}\right) x - \sqrt{a^2 - x^2}$$

input `int(asec(a/x),x)`

output `asec(a/x)*x - sqrt(a**2 - x**2)`

3.13 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx$

Optimal result	118
Mathematica [A] (verified)	118
Rubi [A] (verified)	119
Maple [A] (verified)	121
Fricas [F]	122
Sympy [F]	122
Maxima [F]	122
Giac [F]	123
Mupad [F(-1)]	123
Reduce [F]	123

Optimal result

Integrand size = 10, antiderivative size = 59

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x}{a}\right)}\right)$$

output

```
-1/2*I*arccos(x/a)^2+arccos(x/a)*ln(1+(x/a+I*(1-x^2/a^2)^(1/2))^2)-1/2*I*polylog(2,-(x/a+I*(1-x^2/a^2)^(1/2))^2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = -\frac{1}{2}i \sec^{-1}\left(\frac{a}{x}\right)^2 + \sec^{-1}\left(\frac{a}{x}\right) \log\left(1 + e^{2i \sec^{-1}\left(\frac{a}{x}\right)}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}\left(\frac{a}{x}\right)}\right)$$

input

```
Integrate[ArcSec[a/x]/x,x]
```

output

```
(-1/2*I)*ArcSec[a/x]^2 + ArcSec[a/x]*Log[1 + E^((2*I)*ArcSec[a/x])] - (I/2)*PolyLog[2, -E^((2*I)*ArcSec[a/x])]
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5787, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx \\
 & \quad \downarrow \text{5787} \\
 & \int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx \\
 & \quad \downarrow \text{5137} \\
 & - \int \frac{a\sqrt{1-\frac{x^2}{a^2}} \arccos\left(\frac{x}{a}\right)}{x} d\arccos\left(\frac{x}{a}\right) \\
 & \quad \downarrow \text{3042} \\
 & - \int \arccos\left(\frac{x}{a}\right) \tan\left(\arccos\left(\frac{x}{a}\right)\right) d\arccos\left(\frac{x}{a}\right) \\
 & \quad \downarrow \text{4202} \\
 & 2i \int \frac{e^{2i \arccos\left(\frac{x}{a}\right)} \arccos\left(\frac{x}{a}\right)}{1 + e^{2i \arccos\left(\frac{x}{a}\right)}} d\arccos\left(\frac{x}{a}\right) - \frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 \\
 & \quad \downarrow \text{2620} \\
 & 2i \left(\frac{1}{2}i \int \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) d\arccos\left(\frac{x}{a}\right) - \frac{1}{2}i \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) \right) - \\
 & \quad \frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 \\
 & \quad \downarrow \text{2715}
 \end{aligned}$$

$$2i \left(\frac{1}{4} \int e^{-2i \arccos\left(\frac{x}{a}\right)} \log \left(1 + e^{2i \arccos\left(\frac{x}{a}\right)} \right) dx - \frac{1}{2} i \arccos \left(\frac{x}{a} \right) \log \left(1 + e^{2i \arccos\left(\frac{x}{a}\right)} \right) \right) - \frac{1}{2} i \arccos \left(\frac{x}{a} \right)^2$$

↓ 2838

$$2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \arccos\left(\frac{x}{a}\right)} \right) - \frac{1}{2} i \arccos \left(\frac{x}{a} \right) \log \left(1 + e^{2i \arccos\left(\frac{x}{a}\right)} \right) \right) - \frac{1}{2} i \arccos \left(\frac{x}{a} \right)^2$$

input

```
Int[ArcSec[a/x]/x,x]
```

output

```
(-1/2*I)*ArcCos[x/a]^2 + (2*I)*((-1/2*I)*ArcCos[x/a]*Log[1 + E^((2*I)*ArcCos[x/a])] - PolyLog[2, -E^((2*I)*ArcCos[x/a])]/4)
```

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 5137 `Int[((a_.) + ArcCos[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 5787 `Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int[u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]`

Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

method	result	s
derivativedivides	$-\frac{i \operatorname{arcsec}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arcsec}\left(\frac{a}{x}\right) \ln\left(1 + \left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right) - \frac{i \operatorname{polylog}\left(2, -\left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right)}{2}$	7
default	$-\frac{i \operatorname{arcsec}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arcsec}\left(\frac{a}{x}\right) \ln\left(1 + \left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right) - \frac{i \operatorname{polylog}\left(2, -\left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right)}{2}$	7

input `int(arcsec(a/x)/x,x,method=_RETURNVERBOSE)`

output `-1/2*I*arcsec(a/x)^2+arcsec(a/x)*ln(1+(x/a+I*(1-x^2/a^2)^(1/2))^2)-1/2*I*polylog(2,-(x/a+I*(1-x^2/a^2)^(1/2))^2)`

Fricas [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

input `integrate(arcsec(a/x)/x,x, algorithm="fricas")`

output `integral(arcsec(a/x)/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x} dx$$

input `integrate(asec(a/x)/x,x)`

output `Integral(asec(a/x)/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

input `integrate(arcsec(a/x)/x,x, algorithm="maxima")`

output `integrate(arcsec(a/x)/x, x)`

Giac [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

input `integrate(arcsec(a/x)/x,x, algorithm="giac")`

output `integrate(arcsec(a/x)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx$$

input `int(arcos(x/a)/x,x)`

output `int(arcos(x/a)/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x} dx$$

input `int(asec(a/x)/x,x)`

output `int(asec(a/x)/x,x)`

3.14 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx$

Optimal result	124
Mathematica [B] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	127
Fricas [B] (verification not implemented)	127
Sympy [C] (verification not implemented)	128
Maxima [A] (verification not implemented)	128
Giac [B] (verification not implemented)	129
Mupad [B] (verification not implemented)	129
Reduce [F]	129

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\arccos\left(\frac{x}{a}\right)}{x} + \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a}$$

output

```
-arccos(x/a)/x+arctanh((1-x^2/a^2)^(1/2))/a
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. $2(31) = 62$.

Time = 0.09 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.00

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} + \frac{\sqrt{-1 + \frac{a^2}{x^2}} x \left(-\log\left(1 - \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right) + \log\left(1 + \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right) \right)}{2a^2 \sqrt{1 - \frac{x^2}{a^2}}}$$

input

```
Integrate[ArcSec[a/x]/x^2,x]
```

output

$$-(\text{ArcSec}[a/x]/x) + (\text{Sqrt}[-1 + a^2/x^2]*x*(-\text{Log}[1 - a/(\text{Sqrt}[-1 + a^2/x^2]*x)] + \text{Log}[1 + a/(\text{Sqrt}[-1 + a^2/x^2]*x)]))/(2*a^2*\text{Sqrt}[1 - x^2/a^2])$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5787, 5139, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx \\ & \quad \downarrow \text{5787} \\ & \int \frac{\arccos\left(\frac{x}{a}\right)}{x^2} dx \\ & \quad \downarrow \text{5139} \\ & -\frac{\int \frac{1}{x\sqrt{1-\frac{x^2}{a^2}}} dx}{a} - \frac{\arccos\left(\frac{x}{a}\right)}{x} \\ & \quad \downarrow \text{243} \\ & -\frac{\int \frac{1}{x^2\sqrt{1-\frac{x^2}{a^2}}} dx^2}{2a} - \frac{\arccos\left(\frac{x}{a}\right)}{x} \\ & \quad \downarrow \text{73} \\ & a \int \frac{1}{a^2 - a^2 x^4} d\sqrt{1 - \frac{x^2}{a^2}} - \frac{\arccos\left(\frac{x}{a}\right)}{x} \\ & \quad \downarrow \text{221} \\ & \frac{\text{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a} - \frac{\arccos\left(\frac{x}{a}\right)}{x} \end{aligned}$$

input

$$\text{Int}[\text{ArcSec}[a/x]/x^2, x]$$

output $-(\text{ArcCos}[x/a]/x) + \text{ArcTanh}[\text{Sqrt}[1 - x^2/a^2]]/a$

Defintions of rubi rules used

- rule 73 $\text{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_.) + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m+1) - 1)}(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 221 $\text{Int}[(a_. + (b_.)(x_.)^2)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$
- rule 243 $\text{Int}[(x_.)^{(m_.)}((a_.) + (b_.)(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m-1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m-1)/2]$
- rule 5139 $\text{Int}[(a_. + \text{ArcCos}[(c_.)(x_.)]*(b_.))^{(n_.)}((d_.)(x_.))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}((a + b*\text{ArcCos}[c*x])^n/(d*(m+1))), x] + \text{Simp}[b*c*(n/(d*(m+1))) \text{ Int}[(d*x)^{(m+1)}((a + b*\text{ArcCos}[c*x])^{(n-1)}/\text{Sqrt}[1 - c^2*x^2]), x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{NeQ}[m, -1]$
- rule 5787 $\text{Int}[\text{ArcSec}[(c_.)/((a_.) + (b_.)(x_.)^{(n_.)}])^{(m_.)}(u_.), x_Symbol] \rightarrow \text{Int}[u*\text{ArcCos}[a/c + b*(x^n/c)]^m, x] /; \text{FreeQ}[\{a, b, c, n, m\}, x]$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
parts	$-\frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} + \frac{\operatorname{arctanh}\left(\frac{1}{\sqrt{1-\frac{x^2}{a^2}}}\right)}{a}$	30
derivativeldivides	$-\frac{\frac{a}{x} \operatorname{arcsec}\left(\frac{a}{x}\right) - \ln\left(\frac{a}{x} + \frac{a\sqrt{1-\frac{x^2}{a^2}}}{x}\right)}{a}$	44
default	$-\frac{\frac{a}{x} \operatorname{arcsec}\left(\frac{a}{x}\right) - \ln\left(\frac{a}{x} + \frac{a\sqrt{1-\frac{x^2}{a^2}}}{x}\right)}{a}$	44

input `int(arcsec(a/x)/x^2,x,method=_RETURNVERBOSE)`

output `-arcsec(a/x)/x+1/a*arctanh(1/(1-x^2/a^2)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. 2(29) = 58.

Time = 0.12 (sec) , antiderivative size = 107, normalized size of antiderivative = 3.45

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{2ax \arctan\left(-\frac{x^2\sqrt{\frac{a^2-x^2}{x^2}}}{a^2-x^2}\right) - 2(ax-a) \operatorname{arcsec}\left(\frac{a}{x}\right) - x \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} + a\right) + x \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} - a\right)}{2ax}$$

input `integrate(arcsec(a/x)/x^2,x, algorithm="fricas")`

output `-1/2*(2*a*x*arctan(-x^2*sqrt((a^2 - x^2)/x^2)/(a^2 - x^2)) - 2*(a*x - a)*arcsec(a/x) - x*log(x*sqrt((a^2 - x^2)/x^2) + a) + x*log(x*sqrt((a^2 - x^2)/x^2) - a))/(a*x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.30 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x} - \frac{\begin{cases} -\operatorname{acosh}\left(\frac{a}{x}\right) & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ i \operatorname{asin}\left(\frac{a}{x}\right) & \text{otherwise} \end{cases}}{a}$$

input `integrate(asec(a/x)/x**2,x)`

output `-asec(a/x)/x - Piecewise((-acosh(a/x), Abs(a**2/x**2) > 1), (I*asin(a/x), True))/a`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.68

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\frac{2a \operatorname{arcsec}\left(\frac{a}{x}\right)}{x} - \log\left(\sqrt{-\frac{x^2}{a^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{x^2}{a^2} + 1} + 1\right)}{2a}$$

input `integrate(arcsec(a/x)/x^2,x, algorithm="maxima")`

output `-1/2*(2*a*arcsec(a/x)/x - log(sqrt(-x^2/a^2 + 1) + 1) + log(-sqrt(-x^2/a^2 + 1) + 1))/a`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(29) = 58$.

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.97

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{a \left(\frac{\log\left(|a + \sqrt{a^2 - x^2}\right|)}{a} - \frac{\log\left(|-a + \sqrt{a^2 - x^2}\right|)}{a} \right)}{2|a|} - \frac{\arccos\left(\frac{x}{a}\right)}{x}$$

input `integrate(arcsec(a/x)/x^2,x, algorithm="giac")`

output `1/2*a*(log(abs(a + sqrt(a^2 - x^2)))/a - log(abs(-a + sqrt(a^2 - x^2)))/a) /abs(a) - arccos(x/a)/x`

Mupad [B] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{x^2}{a^2}}}\right)}{a} - \frac{\arccos\left(\frac{x}{a}\right)}{x}$$

input `int(acos(x/a)/x^2,x)`

output `atanh(1/(1 - x^2/a^2)^(1/2))/a - acos(x/a)/x`

Reduce [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \int \frac{a \operatorname{sec}\left(\frac{a}{x}\right)}{x^2} dx$$

input `int(asec(a/x)/x^2,x)`

output `int(asec(a/x)/x**2,x)`

3.15 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx$

Optimal result	130
Mathematica [A] (verified)	130
Rubi [A] (verified)	131
Maple [A] (verified)	132
Fricas [A] (verification not implemented)	132
Sympy [C] (verification not implemented)	133
Maxima [A] (verification not implemented)	133
Giac [A] (verification not implemented)	134
Mupad [F(-1)]	134
Reduce [F]	134

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2}$$

output

$$1/2*(1-x^2/a^2)^{(1/2)}/a/x-1/2*\arccos(x/a)/x^2$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{x\sqrt{1 - \frac{x^2}{a^2}} - a \sec^{-1}\left(\frac{a}{x}\right)}{2ax^2}$$

input

```
Integrate[ArcSec[a/x]/x^3,x]
```

output

$$(x*\text{Sqrt}[1 - x^2/a^2] - a*\text{ArcSec}[a/x])/(2*a*x^2)$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5787, 5139, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx \\ & \quad \downarrow 5787 \\ & \int \frac{\arccos\left(\frac{x}{a}\right)}{x^3} dx \\ & \quad \downarrow 5139 \\ & -\frac{\int \frac{1}{x^2 \sqrt{1-\frac{x^2}{a^2}}} dx}{2a} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2} \\ & \quad \downarrow 242 \\ & \frac{\sqrt{1-\frac{x^2}{a^2}}}{2ax} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2} \end{aligned}$$

input `Int[ArcSec[a/x]/x^3,x]`

output `Sqrt[1 - x^2/a^2]/(2*a*x) - ArcCos[x/a]/(2*x^2)`

Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 5139 `Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Simp[b*c*(n
/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2
*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 5787 `Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]`

Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

method	result	size
parts	$-\frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{2x^2} + \frac{\sqrt{1-\frac{x^2}{a^2}}}{2ax}$	33
derivativedivides	$-\frac{\frac{a^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2x^2} - \frac{x\left(\frac{a^2}{x^2}-1\right)}{2\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}}}{a}}{a^2}$	54
default	$-\frac{\frac{a^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2x^2} - \frac{x\left(\frac{a^2}{x^2}-1\right)}{2\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}}}{a}}{a^2}$	54

input `int(arcsec(a/x)/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arcsec(a/x)/x^2+1/2*(1-x^2/a^2)^(1/2)/a/x`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{a^2 \operatorname{arcsec}\left(\frac{a}{x}\right) - x^2 \sqrt{\frac{a^2-x^2}{x^2}}}{2a^2x^2}$$

input `integrate(arcsec(a/x)/x^3,x, algorithm="fricas")`

output `-1/2*(a^2*arcsec(a/x) - x^2*sqrt((a^2 - x^2)/x^2))/(a^2*x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{\operatorname{asec}\left(\frac{a}{x}\right)}{2x^2} - \frac{\begin{cases} -\frac{\sqrt{\frac{a^2}{x^2}-1}}{a} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ -i\frac{\sqrt{-\frac{a^2}{x^2}+1}}{a} & \text{otherwise} \end{cases}}{2a}$$

input `integrate(asec(a/x)/x**3,x)`

output `-asec(a/x)/(2*x**2) - Piecewise((-sqrt(a**2/x**2 - 1)/a, Abs(a**2/x**2) > 1), (-I*sqrt(-a**2/x**2 + 1)/a, True))/(2*a)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{2x^2} + \frac{\sqrt{-\frac{x^2}{a^2}+1}}{2ax}$$

input `integrate(arcsec(a/x)/x^3,x, algorithm="maxima")`

output `-1/2*arcsec(a/x)/x^2 + 1/2*sqrt(-x^2/a^2 + 1)/(a*x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.61

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{a \left(\frac{a + \sqrt{a^2 - x^2}}{a^2 x} - \frac{x}{(a + \sqrt{a^2 - x^2}) a^2} \right)}{4 |a|} - \frac{\arccos\left(\frac{x}{a}\right)}{2 x^2}$$

input `integrate(arcsec(a/x)/x^3,x, algorithm="giac")`

output `1/4*a*((a + sqrt(a^2 - x^2))/(a^2*x) - x/((a + sqrt(a^2 - x^2))*a^2))/abs(a) - 1/2*arccos(x/a)/x^2`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \int \frac{\arccos\left(\frac{x}{a}\right)}{x^3} dx$$

input `int(acos(x/a)/x^3,x)`

output `int(acos(x/a)/x^3, x)`

Reduce [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \int \frac{a \operatorname{asec}\left(\frac{a}{x}\right)}{x^3} dx$$

input `int(asec(a/x)/x^3,x)`

output `int(asec(a/x)/x**3,x)`

3.16 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx$

Optimal result	135
Mathematica [A] (verified)	135
Rubi [A] (verified)	136
Maple [A] (verified)	138
Fricas [B] (verification not implemented)	138
Sympy [C] (verification not implemented)	139
Maxima [A] (verification not implemented)	139
Giac [A] (verification not implemented)	140
Mupad [F(-1)]	140
Reduce [F]	140

Optimal result

Integrand size = 10, antiderivative size = 60

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} + \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3}$$

output $\frac{1}{6}*(1-x^2/a^2)^{(1/2)}/a/x^2-1/3*\arccos(x/a)/x^3+1/6*\operatorname{arctanh}((1-x^2/a^2)^{(1/2)})/a^3$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.15

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{a^2 x \sqrt{1 - \frac{x^2}{a^2}} - 2a^3 \sec^{-1}\left(\frac{a}{x}\right) - x^3 \log(x) + x^3 \log\left(1 + \sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3 x^3}$$

input `Integrate[ArcSec[a/x]/x^4,x]`

output $(a^2*x*\operatorname{Sqrt}[1 - x^2/a^2] - 2*a^3*\operatorname{ArcSec}[a/x] - x^3*\operatorname{Log}[x] + x^3*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - x^2/a^2]])/(6*a^3*x^3)$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5787, 5139, 243, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx \\
 & \quad \downarrow \text{5787} \\
 & \int \frac{\arccos\left(\frac{x}{a}\right)}{x^4} dx \\
 & \quad \downarrow \text{5139} \\
 & -\frac{\int \frac{1}{x^3 \sqrt{1-\frac{x^2}{a^2}}} dx}{3a} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} \\
 & \quad \downarrow \text{243} \\
 & -\frac{\int \frac{1}{x^4 \sqrt{1-\frac{x^2}{a^2}}} dx^2}{6a} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} \\
 & \quad \downarrow \text{52} \\
 & -\frac{\int \frac{1}{x^2 \sqrt{1-\frac{x^2}{a^2}}} dx^2}{6a} - \frac{\sqrt{1-\frac{x^2}{a^2}}}{x^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} \\
 & \quad \downarrow \text{73} \\
 & -\frac{\int \frac{1}{a^2-a^2x^4} d\sqrt{1-\frac{x^2}{a^2}} - \frac{\sqrt{1-\frac{x^2}{a^2}}}{x^2}}{6a} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} \\
 & \quad \downarrow \text{221} \\
 & -\frac{\operatorname{arctanh}\left(\sqrt{1-\frac{x^2}{a^2}}\right)}{a^2} - \frac{\sqrt{1-\frac{x^2}{a^2}}}{x^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3}
 \end{aligned}$$

input

Int [ArcSec [a/x] /x^4, x]

output

$$-1/3 \operatorname{ArcCos}[x/a]/x^3 - (-\sqrt{1 - x^2/a^2}/x^2) - \operatorname{ArcTanh}[\sqrt{1 - x^2/a^2}]/a^2)/(6*a)$$

Defintions of rubi rules used

rule 52

$$\operatorname{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^{(n + 1)}/((b*c - a*d)*(m + 1))), x] - \operatorname{Simp}[d*((m + n + 2)/((b*c - a*d)*(m + 1))) \operatorname{Int}[(a + b*x)^{(m + 1)}(c + d*x)^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x\} \&\& \operatorname{ILtQ}[m, -1] \&\& \operatorname{FractionQ}[n] \&\& \operatorname{LtQ}[n, 0]$$

rule 73

$$\operatorname{Int}[(a_.) + (b_.)(x_)^{(m_.)}((c_.) + (d_.)(x_)^{(n_.)}), x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Simp}[p/b \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}\{a, b, c, d\}, x\} \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 221

$$\operatorname{Int}[(a_.) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a) \operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x\} \&\& \operatorname{NegQ}[a/b]$$

rule 243

$$\operatorname{Int}[(x_)^{(m_.)}((a_.) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \operatorname{FreeQ}\{a, b, m, p\}, x\} \&\& \operatorname{IntegerQ}[(m - 1)/2]$$

rule 5139

$$\operatorname{Int}[(a_.) + \operatorname{ArcCos}[c_.)(x_)](b_.)^{(n_.)}((d_.)(x_)^{(m_.)}), x_Symbol] \rightarrow \operatorname{Simp}[(d*x)^{(m + 1)}((a + b*\operatorname{ArcCos}[c*x])^n/(d*(m + 1))), x] + \operatorname{Simp}[b*c*(n/(d*(m + 1))) \operatorname{Int}[(d*x)^{(m + 1)}((a + b*\operatorname{ArcCos}[c*x])^{(n - 1)}/\sqrt{1 - c^2*x^2}), x], x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x\} \&\& \operatorname{IGtQ}[n, 0] \&\& \operatorname{NeQ}[m, -1]$$

rule 5787

$$\operatorname{Int}[\operatorname{ArcSec}[c_.)/((a_.) + (b_.)(x_)^{(n_.)})]^{(m_.)}(u_.), x_Symbol] \rightarrow \operatorname{Int}[u*\operatorname{ArcCos}[a/c + b*(x^n/c)]^m, x] /; \operatorname{FreeQ}\{a, b, c, n, m\}, x\}$$

Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.90

method	result	size
parts	$\frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\sqrt{1-\frac{x^2}{a^2}}}{2x^2} - \frac{\operatorname{arctanh}\left(\frac{1}{\sqrt{1-\frac{x^2}{a^2}}}\right)}{3a}$	54
derivativeldivides	$\frac{a^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\sqrt{\frac{a^2}{x^2}-1} \left(\frac{a\sqrt{\frac{a^2}{x^2}-1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2}-1}\right) \right) x}{6\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a}$	91
default	$\frac{a^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\sqrt{\frac{a^2}{x^2}-1} \left(\frac{a\sqrt{\frac{a^2}{x^2}-1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2}-1}\right) \right) x}{6\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a}$	91

input `int(arcsec(a/x)/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*arcsec(a/x)/x^3-1/3/a*(-1/2/x^2*(1-x^2/a^2)^(1/2)-1/2/a^2*arctanh(1/(1-x^2/a^2)^(1/2)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. 2(50) = 100.

Time = 0.13 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.37

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{4a^3x^3 \arctan\left(-\frac{x^2\sqrt{\frac{a^2-x^2}{x^2}}}{a^2-x^2}\right) - x^3 \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} + a\right) + x^3 \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} - a\right) - 2ax^2\sqrt{\frac{a^2-x^2}{x^2}} - 4}{12a^3x^3}$$

input `integrate(arcsec(a/x)/x^4,x, algorithm="fricas")`

output

```
-1/12*(4*a^3*x^3*arctan(-x^2*sqrt((a^2 - x^2)/x^2)/(a^2 - x^2)) - x^3*log(
x*sqrt((a^2 - x^2)/x^2) + a) + x^3*log(x*sqrt((a^2 - x^2)/x^2) - a) - 2*a*
x^2*sqrt((a^2 - x^2)/x^2) - 4*(a^3*x^3 - a^3)*arcsec(a/x))/(a^3*x^3)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.68 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.67

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = -\frac{\operatorname{asec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\begin{cases} -\frac{\sqrt{\frac{a^2}{x^2}-1}}{2ax} - \frac{\operatorname{acosh}\left(\frac{a}{x}\right)}{2a^2} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ \frac{ia}{2x^3\sqrt{-\frac{a^2}{x^2}+1}} - \frac{i}{2ax\sqrt{-\frac{a^2}{x^2}+1}} + \frac{i\operatorname{asin}\left(\frac{a}{x}\right)}{2a^2} & \text{otherwise} \end{cases}}{3a}$$

input

```
integrate(asec(a/x)/x**4,x)
```

output

```
-asec(a/x)/(3*x**3) - Piecewise((-sqrt(a**2/x**2 - 1)/(2*a*x) - acosh(a/x)
/(2*a**2), Abs(a**2/x**2) > 1), (I*a/(2*x**3*sqrt(-a**2/x**2 + 1)) - I/(2*
a*x*sqrt(-a**2/x**2 + 1)) + I*asin(a/x)/(2*a**2), True))/(3*a)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.07

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{\log\left(\frac{2\sqrt{-\frac{x^2}{a^2}+1}}{|x|} + \frac{2}{|x|}\right)}{a^2} + \frac{\sqrt{-\frac{x^2}{a^2}+1}}{x^2} - \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{3x^3}$$

input

```
integrate(arcsec(a/x)/x^4,x, algorithm="maxima")
```

output

```
1/6*(log(2*sqrt(-x^2/a^2 + 1)/abs(x) + 2/abs(x))/a^2 + sqrt(-x^2/a^2 + 1)/
x^2)/a - 1/3*arcsec(a/x)/x^3
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.33

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{a \left(\frac{\log\left(|a + \sqrt{a^2 - x^2}\right|)}{a^3} - \frac{\log\left(|-a + \sqrt{a^2 - x^2}\right|)}{a^3} + \frac{2\sqrt{a^2 - x^2}}{a^2 x^2} \right)}{12|a|} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3}$$

input `integrate(arcsec(a/x)/x^4,x, algorithm="giac")`

output `1/12*a*(log(abs(a + sqrt(a^2 - x^2)))/a^3 - log(abs(-a + sqrt(a^2 - x^2)))/a^3 + 2*sqrt(a^2 - x^2)/(a^2*x^2))/abs(a) - 1/3*arccos(x/a)/x^3`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \int \frac{\arccos\left(\frac{x}{a}\right)}{x^4} dx$$

input `int(acos(x/a)/x^4,x)`

output `int(acos(x/a)/x^4, x)`

Reduce [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \int \frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x^4} dx$$

input `int(asec(a/x)/x^4,x)`

output `int(asec(a/x)/x**4,x)`

3.17 $\int \frac{\sec^{-1}(ax^n)}{x} dx$

Optimal result	141
Mathematica [C] (verified)	141
Rubi [A] (warning: unable to verify)	142
Maple [A] (verified)	144
Fricas [F(-2)]	145
Sympy [F]	145
Maxima [F]	145
Giac [F]	146
Mupad [F(-1)]	146
Reduce [F]	146

Optimal result

Integrand size = 10, antiderivative size = 69

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \frac{i \sec^{-1}(ax^n)^2}{2n} - \frac{\sec^{-1}(ax^n) \log\left(1 + e^{2i \sec^{-1}(ax^n)}\right)}{n} + \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^n)}\right)}{2n}$$

output

$1/2*I*\operatorname{arcsec}(a*x^n)^2/n - \operatorname{arcsec}(a*x^n)*\ln(1+(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)/n + 1/2*I*\operatorname{polylog}(2, -(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)/n$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.87

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \frac{x^{-n} {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{2}, \frac{3}{2}, \frac{x^{-2n}}{a^2}\right)}{an} + \left(\sec^{-1}(ax^n) + \arcsin\left(\frac{x^{-n}}{a}\right)\right) \log(x)$$

input

`Integrate[ArcSec[a*x^n]/x, x]`

output

$$\text{HypergeometricPFQ}[\{1/2, 1/2, 1/2\}, \{3/2, 3/2\}, 1/(a^2 x^{2n})]/(a^n x^n) + (\text{ArcSec}[a x^n] + \text{ArcSin}[1/(a x^n)]) \cdot \text{Log}[x]$$
Rubi [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.16, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {7282, 5741, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^{-1}(ax^n)}{x} dx \\ & \quad \downarrow \text{7282} \\ & \frac{\int x^{-n} \sec^{-1}(ax^n) dx^n}{n} \\ & \quad \downarrow \text{5741} \\ & \frac{\int x^{-n} \arccos\left(\frac{x^{-n}}{a}\right) dx^{-n}}{n} \\ & \quad \downarrow \text{5137} \\ & \frac{\int ax^n \sqrt{1 - \frac{x^{-2n}}{a^2}} \arccos\left(\frac{x^{-n}}{a}\right) d \arccos\left(\frac{x^{-n}}{a}\right)}{n} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \arccos\left(\frac{x^{-n}}{a}\right) \tan\left(\arccos\left(\frac{x^{-n}}{a}\right)\right) d \arccos\left(\frac{x^{-n}}{a}\right)}{n} \\ & \quad \downarrow \text{4202} \\ & \frac{\frac{1}{2}ix^{2n} - 2i \int \frac{e^{2i \arccos\left(\frac{x^{-n}}{a}\right)} \arccos\left(\frac{x^{-n}}{a}\right)}{1+e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}} d \arccos\left(\frac{x^{-n}}{a}\right)}{n} \\ & \quad \downarrow \text{2620} \end{aligned}$$

$$\frac{\frac{1}{2}ix^{2n} - 2i\left(\frac{1}{2}i \int \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right) d \arccos\left(\frac{x^{-n}}{a}\right) - \frac{1}{2}i \arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)\right)}{n}$$

↓ 2715

$$\frac{\frac{1}{2}ix^{2n} - 2i\left(\frac{1}{4} \int e^{2i \arccos\left(\frac{x^{-n}}{a}\right)} \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right) de^{2i \arccos\left(\frac{x^{-n}}{a}\right)} - \frac{1}{2}i \arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)\right)}{n}$$

↓ 2838

$$\frac{\frac{1}{2}ix^{2n} - 2i\left(-\frac{1}{4} \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right) - \frac{1}{2}i \arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)\right)}{n}$$

input

```
Int[ArcSec[a*x^n]/x, x]
```

output

```
((I/2)*x^(2*n) - (2*I)*((-1/2*I)*ArcCos[1/(a*x^n)]*Log[1 + E^((2*I)*ArcCos[1/(a*x^n)])] - PolyLog[2, -E^((2*I)*ArcCos[1/(a*x^n)])]/4))/n
```

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_)+(f_)*(x_))))^(n_)*((c_)+(d_)*(x_))^(m_))/((a_)+(b_)*((F_)^((g_)*((e_)+(f_)*(x_))))^(n_)), x_Symbol] := Simp[((c+d*x)^m/(b*f*g*n*Log[F]))*Log[1+b*((F^(g*(e+f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c+d*x)^(m-1)*Log[1+b*((F^(g*(e+f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_)+(b_)*((F_)^((e_)*((c_)+(d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a+b*x]/x, x], x, (F^(e*(c+d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_)+(e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```


rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]`

rule 5137 `Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 5741 `Int[((a_.) + ArcSec[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

rule 7282 `Int[(u_)/(x_), x_Symbol] := With[{lst = PowerVariableExpn[u, 0, x]}, Simp[1/lst[[2]] Subst[Int[NormalizeIntegrand[Simplify[lst[[1]]/x], x], x], x, (lst[[3]]*x)^lst[[2]]], x] /; !FalseQ[lst] && NeQ[lst[[2]], 0] /; NonsumQ[u] && !RationalFunctionQ[u, x]`

Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.35

method	result	size
derivativedivides	$\frac{\frac{i \operatorname{arcsec}(a x^n)^2}{2} - \operatorname{arcsec}(a x^n) \ln \left(1 + \left(\frac{x^{-n}}{a} + i \sqrt{1 - \frac{x^{-2n}}{a^2}} \right)^2 \right)}{n} + \frac{i \operatorname{polylog} \left(2, - \left(\frac{x^{-n}}{a} + i \sqrt{1 - \frac{x^{-2n}}{a^2}} \right)^2 \right)}{2}$	93
default	$\frac{\frac{i \operatorname{arcsec}(a x^n)^2}{2} - \operatorname{arcsec}(a x^n) \ln \left(1 + \left(\frac{x^{-n}}{a} + i \sqrt{1 - \frac{x^{-2n}}{a^2}} \right)^2 \right)}{n} + \frac{i \operatorname{polylog} \left(2, - \left(\frac{x^{-n}}{a} + i \sqrt{1 - \frac{x^{-2n}}{a^2}} \right)^2 \right)}{2}$	93

input `int(arcsec(a*x^n)/x,x,method=_RETURNVERBOSE)`

output $1/n*(1/2*I*arcsec(a*x^n)^2-arcsec(a*x^n)*ln(1+(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)+1/2*I*polylog(2,-(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2))$

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

input `integrate(arcsec(a*x^n)/x,x, algorithm="fricas")`

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asec}(ax^n)}{x} dx$$

input `integrate(asec(a*x**n)/x,x)`

output `Integral(asec(a*x**n)/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsec}(ax^n)}{x} dx$$

input `integrate(arcsec(a*x^n)/x,x, algorithm="maxima")`

output `-a^2*n*integrate(sqrt(a*x^n + 1)*sqrt(a*x^n - 1)*log(x)/(a^4*x*x^(2*n) - a^2*x), x) + arctan(sqrt(a*x^n + 1)*sqrt(a*x^n - 1))*log(x)`

Giac [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsec}(ax^n)}{x} dx$$

input `integrate(arcsec(a*x^n)/x,x, algorithm="giac")`

output `integrate(arcsec(a*x^n)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\arccos\left(\frac{1}{ax^n}\right)}{x} dx$$

input `int(arccos(1/(a*x^n))/x,x)`

output `int(arccos(1/(a*x^n))/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asec}(x^n a)}{x} dx$$

input `int(asec(a*x^n)/x,x)`

output `int(asec(x**n*a)/x,x)`

3.18 $\int x^4 \sec^{-1}(a + bx) dx$

Optimal result	147
Mathematica [A] (verified)	148
Rubi [A] (verified)	148
Maple [A] (verified)	151
Fricas [A] (verification not implemented)	152
Sympy [F]	152
Maxima [F]	153
Giac [B] (verification not implemented)	153
Mupad [F(-1)]	154
Reduce [F]	154

Optimal result

Integrand size = 10, antiderivative size = 197

$$\int x^4 \sec^{-1}(a + bx) dx = \frac{a(20 + 53a^2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{30b^5} + \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{20b^2} - \frac{(9 + 58a^2)(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{120b^5} + \frac{a^5 \sec^{-1}(a + bx)}{5b^5} + \frac{1}{5}x^5 \sec^{-1}(a + bx) - \frac{(3 + 40a^2 + 40a^4) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{40b^5}$$

output

```
1/30*a*(53*a^2+20)*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^5+11/60*a*x^2*(b*x+a)*(
1-1/(b*x+a)^2)^(1/2)/b^3-1/20*x^3*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2-1/120*
(58*a^2+9)*(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)/b^5+1/5*a^5*arcsec(b*x+a)/b^5+1
/5*x^5*arcsec(b*x+a)-1/40*(40*a^4+40*a^2+3)*arctanh((1-1/(b*x+a)^2)^(1/2))
/b^5
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.88

$$\int x^4 \sec^{-1}(a + bx) dx$$

$$= \frac{\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} (a^2(71 + 154a^2) + 2a(31 + 48a^2)bx - 9(1 + 4a^2)b^2x^2 + 16ab^3x^3 - 6b^4x^4) + 24b^5x^5 \sec^{-1}(a + bx)}{120b^5}$$

input

```
Integrate[x^4*ArcSec[a + b*x],x]
```

output

```
(Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^2*(71 + 154*a^2) + 2*a*(31 + 48*a^2)*b*x - 9*(1 + 4*a^2)*b^2*x^2 + 16*a*b^3*x^3 - 6*b^4*x^4) + 24*b^5*x^5*ArcSec[a + b*x] - 24*a^5*ArcSin[(a + b*x)^(-1)] - 3*(3 + 40*a^2 + 40*a^4)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(120*b^5)
```

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {5781, 4926, 3042, 4269, 3042, 4544, 3042, 4536, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \sec^{-1}(a + bx) dx$$

$$\downarrow \text{5781}$$

$$\frac{\int b^4 x^4 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^5}$$

$$\downarrow \text{4926}$$

$$\frac{\frac{1}{5} \int -b^5 x^5 d \sec^{-1}(a + bx) + \frac{1}{5} b^5 x^5 \sec^{-1}(a + bx)}{b^5}$$

$$\downarrow \text{3042}$$

$$\frac{\frac{1}{5} \int (a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}))^5 d \sec^{-1}(a + bx) + \frac{1}{5} b^5 x^5 \sec^{-1}(a + bx)}{b^5}$$

↓ 4269

$$\frac{\frac{1}{5} \left(\frac{1}{4} \int b^2 x^2 (4a^3 + 11(a + bx)^2 a - 3(4a^2 + 1)(a + bx)) d \sec^{-1}(a + bx) - \frac{1}{4} b^3 x^3 (a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) + \frac{1}{5} b^5 x^5}{b^5}$$

↓ 3042

$$\frac{\frac{1}{5} \left(\frac{1}{4} \int (a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}))^2 (4a^3 + 11 \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}))^2 a - 3(4a^2 + 1) \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}) \right)}{b^5}$$

↓ 4544

$$\frac{\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int -bx(12a^4 - (48a^2 + 31)(a + bx)a + (58a^2 + 9)(a + bx)^2) d \sec^{-1}(a + bx) + \frac{11}{3} ab^2 x^2 (a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) \right)}{b^5}$$

↓ 3042

$$\frac{\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \int (a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2})) (12a^4 - (48a^2 + 31) \csc(\sec^{-1}(a + bx) + \frac{\pi}{2})) a + (58a^2 + 9) \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}) \right) \right)}{b^5}$$

↓ 4536

$$\frac{\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \int (24a^5 + 4(53a^2 + 20)(a + bx)^2 a - 3(40a^4 + 40a^2 + 3)(a + bx)) d \sec^{-1}(a + bx) - \frac{1}{2} (58a^2 + 9)(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) \right) \right)}{b^5}$$

↓ 2009

$$\frac{\frac{1}{5} \left(\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} (24a^5 \sec^{-1}(a + bx) + 4(53a^2 + 20) a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} - 3(40a^4 + 40a^2 + 3) \operatorname{arctanh}(\sqrt{1 - \frac{1}{(a+bx)^2}}) \right) \right) \right)}{b^5}$$

input

Int[x^4*ArcSec[a + b*x],x]

output

$$\begin{aligned} & ((b^5 x^5 \operatorname{ArcSec}[a + b x])/5 + (-1/4 (b^3 x^3 (a + b x) \sqrt{1 - (a + b x)^{-2}})) + ((11 a b^2 x^2 (a + b x) \sqrt{1 - (a + b x)^{-2}})/3 + (-1/2 ((9 + 58 a^2) (a + b x)^2 \sqrt{1 - (a + b x)^{-2}}) + (4 a (20 + 53 a^2) (a + b x) \sqrt{1 - (a + b x)^{-2}} + 24 a^5 \operatorname{ArcSec}[a + b x] - 3 (3 + 40 a^2 + 40 a^4) \operatorname{ArcTanh}[\sqrt{1 - (a + b x)^{-2}}]) / (2/3) / 4) / 5) / b^5 \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \;/; \operatorname{SumQ}[u]$$

rule 3042

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \;/; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4269

$$\begin{aligned} \operatorname{Int}[(\operatorname{csc}[c_] + (d_)(x_)](b_) + (a_))^{(n_)}, x_Symbol] \rightarrow & \operatorname{Simp}[(-b^2) \operatorname{Cot}[c + d x] * ((a + b \operatorname{Csc}[c + d x])^{(n-2)} / (d(n-1))), x] + \operatorname{Simp}[1/(n-1) \\ & \operatorname{Int}[(a + b \operatorname{Csc}[c + d x])^{(n-3)} * \operatorname{Simp}[a^{3(n-1)} + (b(b^2(n-2) + 3a^2(n-1))) * \operatorname{Csc}[c + d x] + (a b^2(3n-4)) * \operatorname{Csc}[c + d x]^2, x], x] \;/; \\ & \operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{NeQ}[a^2 - b^2, 0] \ \&\& \operatorname{GtQ}[n, 2] \ \&\& \operatorname{IntegerQ}[2n] \end{aligned}$$

rule 4536

$$\begin{aligned} \operatorname{Int}[(A_) + \operatorname{csc}[e_] + (f_)(x_)](B_) + \operatorname{csc}[e_] + (f_)(x_)]^2(C_), x_Symbol] \rightarrow & \operatorname{Simp}[(-b) * C * \operatorname{Csc}[e + f x] * (\operatorname{Cot}[e + f x] / (2f)), x] + \operatorname{Simp}[1/2 \operatorname{Int}[\operatorname{Simp}[2Aa + (2Ba + b(2A + C)) * \operatorname{Csc}[e + f x] + 2(aC + Bb) * \operatorname{Csc}[e + f x]^2, x], x], x] \;/; \\ & \operatorname{FreeQ}\{a, b, e, f, A, B, C\}, x] \end{aligned}$$

rule 4544

$$\begin{aligned} \operatorname{Int}[(A_) + \operatorname{csc}[e_] + (f_)(x_)](B_) + \operatorname{csc}[e_] + (f_)(x_)]^2(C_), x_Symbol] \rightarrow & \operatorname{Simp}[(-C) * \operatorname{Cot}[e + f x] * ((a + b \operatorname{Csc}[e + f x])^m / (f(m+1))), x] + \operatorname{Simp}[1/(m+1) \operatorname{Int}[(a + b \operatorname{Csc}[e + f x])^{(m-1)} * \operatorname{Simp}[aA(m+1) + ((A*b + a*B)*(m+1) + bC*m) * \operatorname{Csc}[e + f x] + (b*B*(m+1) + aC*m) * \operatorname{Csc}[e + f x]^2, x], x], x] \;/; \\ & \operatorname{FreeQ}\{a, b, e, f, A, B, C\}, x \ \&\& \operatorname{NeQ}[a^2 - b^2, 0] \ \&\& \operatorname{IGtQ}[2m, 0] \end{aligned}$$

rule 4926

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 5781

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)])*(b_.)^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.67

method	result
derivativedivides	$-\frac{\operatorname{arcsec}(bx+a)a^5}{5} + \operatorname{arcsec}(bx+a)a^4(bx+a) - 2 \operatorname{arcsec}(bx+a)a^3(bx+a)^2 + 2 \operatorname{arcsec}(bx+a)a^2(bx+a)^3 - \operatorname{arcsec}(bx+a)a(bx+a)^4$
default	$-\frac{\operatorname{arcsec}(bx+a)a^5}{5} + \operatorname{arcsec}(bx+a)a^4(bx+a) - 2 \operatorname{arcsec}(bx+a)a^3(bx+a)^2 + 2 \operatorname{arcsec}(bx+a)a^2(bx+a)^3 - \operatorname{arcsec}(bx+a)a(bx+a)^4$
parts	$\frac{x^5 \operatorname{arcsec}(bx+a)}{5} + \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(-6x^3\sqrt{b^2x^2+2abx+a^2-1} b^3\sqrt{b^2+22\sqrt{b^2x^2+2abx+a^2-1}} \sqrt{b^2} a b^2x^2 - 24a^2x^3 \right)}{\dots}$

input

```
int(x^4*arcsec(b*x+a), x, method=_RETURNVERBOSE)
```


output

```
1/b^5*(-1/5*arcsec(b*x+a)*a^5+arcsec(b*x+a)*a^4*(b*x+a)-2*arcsec(b*x+a)*a^3*(b*x+a)^2+2*arcsec(b*x+a)*a^2*(b*x+a)^3-arcsec(b*x+a)*a*(b*x+a)^4+1/5*arcsec(b*x+a)*(b*x+a)^5-1/120*((b*x+a)^2-1)^(1/2)*(24*a^5*arctan(1/((b*x+a)^2-1)^(1/2))+120*a^4*ln(b*x+a+((b*x+a)^2-1)^(1/2))-240*a^3*((b*x+a)^2-1)^(1/2)+120*a^2*(b*x+a)*((b*x+a)^2-1)^(1/2)-40*a*(b*x+a)^2*((b*x+a)^2-1)^(1/2)+6*(b*x+a)^3*((b*x+a)^2-1)^(1/2)+120*a^2*ln(b*x+a+((b*x+a)^2-1)^(1/2))-80*a*((b*x+a)^2-1)^(1/2)+9*(b*x+a)*((b*x+a)^2-1)^(1/2)+9*ln(b*x+a+((b*x+a)^2-1)^(1/2)))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.77

$$\int x^4 \sec^{-1}(a + bx) dx$$

$$= \frac{24b^5x^5 \operatorname{arcsec}(bx + a) + 48a^5 \arctan(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) + 3(40a^4 + 40a^2 + 3) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{b^5}$$

input

```
integrate(x^4*arcsec(b*x+a),x, algorithm="fricas")
```

output

```
1/120*(24*b^5*x^5*arcsec(b*x + a) + 48*a^5*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + 3*(40*a^4 + 40*a^2 + 3)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (6*b^3*x^3 - 22*a*b^2*x^2 - 154*a^3 + (58*a^2 + 9)*b*x - 71*a)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/b^5
```

Sympy [F]

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \operatorname{asec}(a + bx) dx$$

input

```
integrate(x**4*asec(b*x+a),x)
```

output

```
Integral(x**4*asec(a + b*x), x)
```

Maxima [F]

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \operatorname{arcsec}(bx + a) dx$$

input `integrate(x^4*arcsec(b*x+a),x, algorithm="maxima")`

output `1/5*x^5*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/5*(b^2*x^6 + a*b*x^5)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x + a - 1)) - 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. $2(173) = 346$.

Time = 0.16 (sec) , antiderivative size = 412, normalized size of antiderivative = 2.09

$$\int x^4 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{960} b \left(\frac{192 (bx + a)^5 \left(\frac{5a}{bx+a} - \frac{10a^2}{(bx+a)^2} + \frac{10a^3}{(bx+a)^3} - \frac{5a^4}{(bx+a)^4} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right) - \frac{3 (bx + a)^6}{b^6} \right)$$

input `integrate(x^4*arcsec(b*x+a),x, algorithm="giac")`

output

```
-1/960*b*(192*(b*x + a)^5*(5*a/(b*x + a) - 10*a^2/(b*x + a)^2 + 10*a^3/(b*x + a)^3 - 5*a^4/(b*x + a)^4 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^6 - (3*(b*x + a)^4*(sqrt(-1/(b*x + a)^2 + 1) - 1)^4 + 40*(b*x + a)^3*a*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 240*(b*x + a)^2*a^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 960*(b*x + a)*a^3*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 24*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 360*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 24*(40*a^4 + 40*a^2 + 3)*log(1/2*abs(b*x + a)*abs(-2*sqrt(-1/(b*x + a)^2 + 1) + 2)) - (120*(8*a^3 + 3*a)*(b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 24*(10*a^2 + 1)*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 40*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 3)/((b*x + a)^4*(sqrt(-1/(b*x + a)^2 + 1) - 1)^4))/b^6)
```

Mupad [F(-1)]

Timed out.

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \operatorname{acos}\left(\frac{1}{a + bx}\right) dx$$

input

```
int(x^4*acos(1/(a + b*x)),x)
```

output

```
int(x^4*acos(1/(a + b*x)), x)
```

Reduce [F]

$$\int x^4 \sec^{-1}(a + bx) dx = \int a \operatorname{sec}(bx + a) x^4 dx$$

input

```
int(x^4*asec(b*x+a),x)
```

output

```
int(asec(a + b*x)*x**4,x)
```

3.19 $\int x^3 \sec^{-1}(a + bx) dx$

Optimal result	155
Mathematica [A] (verified)	156
Rubi [A] (verified)	156
Maple [A] (verified)	159
Fricas [A] (verification not implemented)	159
Sympy [F]	160
Maxima [F]	160
Giac [B] (verification not implemented)	161
Mupad [F(-1)]	161
Reduce [F]	162

Optimal result

Integrand size = 10, antiderivative size = 155

$$\int x^3 \sec^{-1}(a + bx) dx = -\frac{(2 + 17a^2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^4} - \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2}$$

$$+ \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \sec^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4}x^4 \sec^{-1}(a + bx) + \frac{a(1 + 2a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{2b^4}$$

output

```
-1/12*(17*a^2+2)*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4-1/12*x^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2+1/3*a*(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)/b^4-1/4*a^4*arcs
ec(b*x+a)/b^4+1/4*x^4*arcsec(b*x+a)+1/2*a*(2*a^2+1)*arctanh((1-1/(b*x+a)^2)^(1/2))/b^4
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.97

$$\int x^3 \sec^{-1}(a + bx) dx$$

$$= \frac{-\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}(2a + 13a^3 + 2bx + 9a^2bx - 3ab^2x^2 + b^3x^3) + 3b^4x^4 \sec^{-1}(a + bx) + 3a^4 \arcsin\left(\frac{1}{a+bx}\right)}{12b^4}$$

input `Integrate[x^3*ArcSec[a + b*x],x]`

output

```
(-(Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(2*a + 13*a^3 + 2*b*x
+ 9*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3)) + 3*b^4*x^4*ArcSec[a + b*x] + 3*a^4*
ArcSin[(a + b*x)^(-1)] + 6*a*(1 + 2*a^2)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2
+ 2*a*b*x + b^2*x^2)/(a + b*x)^2]])]/(12*b^4)
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5781, 25, 4926, 3042, 4269, 3042, 4536, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \sec^{-1}(a + bx) dx$$

$$\downarrow \text{5781}$$

$$\frac{\int b^3 x^3 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^4}$$

$$\downarrow \text{25}$$

$$\frac{\int -b^3 x^3 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^4}$$

$$\downarrow \text{4926}$$

$$\frac{\frac{1}{4}b^4x^4 \sec^{-1}(a+bx) - \frac{1}{4} \int b^4x^4 d \sec^{-1}(a+bx)}{b^4}$$

↓ 3042

$$\frac{\frac{1}{4}b^4x^4 \sec^{-1}(a+bx) - \frac{1}{4} \int (a - \csc(\sec^{-1}(a+bx) + \frac{\pi}{2}))^4 d \sec^{-1}(a+bx)}{b^4}$$

↓ 4269

$$\frac{\frac{1}{4} \left(-\frac{1}{3} \int -bx(3a^3 + 8(a+bx)^2a - (9a^2 + 2)(a+bx)) d \sec^{-1}(a+bx) - \frac{1}{3}b^2x^2(a+bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \right) + \frac{1}{4}b^4x^4 \sec^{-1}(a+bx)}{b^4}$$

↓ 3042

$$\frac{\frac{1}{4} \left(-\frac{1}{3} \int (a - \csc(\sec^{-1}(a+bx) + \frac{\pi}{2})) \left(3a^3 + 8 \csc(\sec^{-1}(a+bx) + \frac{\pi}{2})^2 a + (-9a^2 - 2) \csc(\sec^{-1}(a+bx) + \frac{\pi}{2}) \right) d \sec^{-1}(a+bx) \right)}{b^4}$$

↓ 4536

$$\frac{\frac{1}{4} \left(\frac{1}{3} \left(4a(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} - \frac{1}{2} \int (6a^4 - 12(2a^2 + 1)(a+bx)a + 2(17a^2 + 2)(a+bx)^2) d \sec^{-1}(a+bx) \right) \right)}{b^4}$$

↓ 2009

$$\frac{\frac{1}{4} \left(\frac{1}{3} \left(\frac{1}{2} \left(-6a^4 \sec^{-1}(a+bx) + 12(2a^2 + 1) \operatorname{arctanh} \left(\sqrt{1 - \frac{1}{(a+bx)^2}} \right) - 2(17a^2 + 2)(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) + 4a^4 \right) \right)}{b^4}$$

input

```
Int[x^3*ArcSec[a + b*x],x]
```

output

```
((b^4*x^4*ArcSec[a + b*x])/4 + (-1/3*(b^2*x^2*(a + b*x)*Sqrt[1 - (a + b*x)
^(-2)]) + (4*a*(a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)] + (-2*(2 + 17*a^2)*(a
+ b*x)*Sqrt[1 - (a + b*x)^(-2)] - 6*a^4*ArcSec[a + b*x] + 12*a*(1 + 2*a^2)
*ArcTanh[Sqrt[1 - (a + b*x)^(-2)]])/2)/3)/4)/b^4
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ /; SumQ}[\text{u}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4269 $\text{Int}[(\text{csc}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.))^{\text{n}_}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b}^2) * \text{Cot}[\text{c} + \text{d} * \text{x}] * ((\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 2} / (\text{d} * (\text{n} - 1))), \text{x}] + \text{Simp}[1 / (\text{n} - 1) \text{ Int}[(\text{a} + \text{b} * \text{Csc}[\text{c} + \text{d} * \text{x}])^{\text{n} - 3} * \text{Simp}[\text{a}^3 * (\text{n} - 1) + (\text{b} * (\text{b}^2 * (\text{n} - 2) + 3 * \text{a}^2 * (\text{n} - 1))) * \text{Csc}[\text{c} + \text{d} * \text{x}] + (\text{a} * \text{b}^2 * (3 * \text{n} - 4)) * \text{Csc}[\text{c} + \text{d} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \&\& \text{GtQ}[\text{n}, 2] \&\& \text{IntegerQ}[2 * \text{n}]$
- rule 4536 $\text{Int}[(\text{A}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{B}_.) + \text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^2 * (\text{C}_.) * (\text{csc}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] * (\text{b}_.) + (\text{a}_.)), \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b}) * \text{C} * \text{Csc}[\text{e} + \text{f} * \text{x}] * (\text{Cot}[\text{e} + \text{f} * \text{x}] / (2 * \text{f})), \text{x}] + \text{Simp}[1 / 2 \quad \text{Int}[\text{Simp}[2 * \text{A} * \text{a} + (2 * \text{B} * \text{a} + \text{b} * (2 * \text{A} + \text{C})) * \text{Csc}[\text{e} + \text{f} * \text{x}] + 2 * (\text{a} * \text{C} + \text{B} * \text{b}) * \text{Csc}[\text{e} + \text{f} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}]$
- rule 4926 $\text{Int}[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{\text{m}_.} * \text{Sec}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * ((\text{a}_.) + (\text{b}_.) * \text{Sec}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)]))^{\text{n}_.} * \text{Tan}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{e} + \text{f} * \text{x})^{\text{m}} * ((\text{a} + \text{b} * \text{Sec}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1} / (\text{b} * \text{d} * (\text{n} + 1))), \text{x}] - \text{Simp}[\text{f} * (\text{m} / (\text{b} * \text{d} * (\text{n} + 1))) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{m} - 1} * (\text{a} + \text{b} * \text{Sec}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1}, \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{NeQ}[\text{n}, -1]$
- rule 5781 $\text{Int}[(\text{a}_.) + \text{ArcSec}[(\text{c}_.) + (\text{d}_.) * (\text{x}_)] * (\text{b}_.)]^{\text{p}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{\text{m}_.}, \text{x_Symbol}] \rightarrow \text{Simp}[1 / \text{d}^{\text{m} + 1} \quad \text{Subst}[\text{Int}[(\text{a} + \text{b} * \text{x})^{\text{p}} * \text{Sec}[\text{x}] * \text{Tan}[\text{x}] * (\text{d} * \text{e} - \text{c} * \text{f} + \text{f} * \text{Sec}[\text{x}])^{\text{m}}, \text{x}], \text{x}, \text{ArcSec}[\text{c} + \text{d} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{IntegerQ}[\text{m}]$

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.61

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a)a^4}{4} - \operatorname{arcsec}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsec}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^4}{4} + \dots$
default	$\frac{\operatorname{arcsec}(bx+a)a^4}{4} - \operatorname{arcsec}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsec}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^4}{4} + \dots$
parts	$\frac{x^4 \operatorname{arcsec}(bx+a)}{4} + \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(3a^4 \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2-x^2} \sqrt{b^2x^2+2abx+a^2-1} b^2 \sqrt{b^2+12} \right)}{\dots}$

```
input int(x^3*arcsec(b*x+a), x, method=_RETURNVERBOSE)
```

```
output 1/b^4*(1/4*arcsec(b*x+a)*a^4-arcsec(b*x+a)*a^3*(b*x+a)+3/2*arcsec(b*x+a)*a^2*(b*x+a)^2-arcsec(b*x+a)*a*(b*x+a)^3+1/4*arcsec(b*x+a)*(b*x+a)^4+1/12*((b*x+a)^2-1)^(1/2)*(3*a^4*arctan(1/((b*x+a)^2-1)^(1/2))+12*a^3*ln(b*x+a+(b*x+a)^2-1)^(1/2))-18*a^2*((b*x+a)^2-1)^(1/2)+6*a*(b*x+a)*((b*x+a)^2-1)^(1/2)-(b*x+a)^2*((b*x+a)^2-1)^(1/2)+6*a*ln(b*x+a+((b*x+a)^2-1)^(1/2))-2*((b*x+a)^2-1)^(1/2))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

$$\int x^3 \sec^{-1}(a + bx) dx = \frac{3b^4x^4 \operatorname{arcsec}(bx+a) - 6a^4 \arctan(-bx-a + \sqrt{b^2x^2+2abx+a^2-1}) - 6(2a^3+a) \log(-bx-a + \sqrt{b^2x^2+2abx+a^2-1})}{12b^4}$$

```
input integrate(x^3*arcsec(b*x+a), x, algorithm="fricas")
```


output

```
1/12*(3*b^4*x^4*arcsec(b*x + a) - 6*a^4*arctan(-b*x - a + sqrt(b^2*x^2 + 2
*a*b*x + a^2 - 1)) - 6*(2*a^3 + a)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x +
a^2 - 1)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(b^2*x^2 - 4*a*b*x + 13*a^2
+ 2))/b^4
```

Sympy [F]

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \operatorname{asec}(a + bx) dx$$

input

```
integrate(x**3*asec(b*x+a),x)
```

output

```
Integral(x**3*asec(a + b*x), x)
```

Maxima [F]

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \operatorname{arcsec}(bx + a) dx$$

input

```
integrate(x^3*arcsec(b*x+a),x, algorithm="maxima")
```

output

```
1/4*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/4*(b^2*x
^5 + a*b*x^4)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2
*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x
+ a - 1)) - 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 302 vs. $2(135) = 270$.

Time = 0.15 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.95

$$\int x^3 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{96}b \left(\frac{24(bx+a)^4 \left(\frac{4a}{bx+a} - \frac{6a^2}{(bx+a)^2} + \frac{4a^3}{(bx+a)^3} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{b^5} + \frac{(bx+a)^3 \left(\sqrt{-\frac{1}{(bx+a)}} \right)}{b^5} \right)$$

input `integrate(x^3*arcsec(b*x+a),x, algorithm="giac")`

output `-1/96*b*(24*(b*x + a)^4*(4*a/(b*x + a) - 6*a^2/(b*x + a)^2 + 4*a^3/(b*x + a)^3 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^5 + ((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 12*(b*x + a)^2*a*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 72*(b*x + a)*a^2*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 9*(b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 48*(2*a^3 + a)*log(1/2*abs(b*x + a)*abs(-2*sqrt(-1/(b*x + a)^2 + 1) + 2)) - (9*(8*a^2 + 1)*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)/((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3))/b^5`

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \arccos \left(\frac{1}{a + bx} \right) dx$$

input `int(x^3*acos(1/(a + b*x)),x)`

output `int(x^3*acos(1/(a + b*x)), x)`

Reduce [F]

$$\int x^3 \sec^{-1}(a + bx) dx = \int a \sec(bx + a) x^3 dx$$

input `int(x^3*asec(b*x+a),x)`

output `int(asec(a + b*x)*x**3,x)`

3.20 $\int x^2 \sec^{-1}(a + bx) dx$

Optimal result	163
Mathematica [A] (verified)	163
Rubi [A] (verified)	164
Maple [A] (verified)	166
Fricas [A] (verification not implemented)	166
Sympy [F]	167
Maxima [F]	167
Giac [B] (verification not implemented)	168
Mupad [F(-1)]	168
Reduce [F]	169

Optimal result

Integrand size = 10, antiderivative size = 116

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{5a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^3} - \frac{x(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \sec^{-1}(a + bx)}{3b^3} + \frac{1}{3} x^3 \sec^{-1}(a + bx) - \frac{(1 + 6a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{6b^3}$$

output

```
5/6*a*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^3-1/6*x*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2+1/3*a^3*arcsec(b*x+a)/b^3+1/3*x^3*arcsec(b*x+a)-1/6*(6*a^2+1)*arctanh((1-1/(b*x+a)^2)^(1/2))/b^3
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.13

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{(5a^2 + 4abx - b^2x^2) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} + 2b^3x^3 \sec^{-1}(a + bx) - 2a^3 \arcsin\left(\frac{1}{a+bx}\right) - (1 + 6a^2) \log\left((a + bx)\right)}{6b^3}$$

input `Integrate[x^2*ArcSec[a + b*x],x]`

output
$$\frac{((5a^2 + 4abx - b^2x^2)\sqrt{(-1 + a^2 + 2abx + b^2x^2)/(a + bx)^2} + 2b^3x^3\text{ArcSec}[a + bx] - 2a^3\text{ArcSin}[(a + bx)^{-1}] - (1 + 6a^2)\text{Log}[(a + bx)(1 + \sqrt{(-1 + a^2 + 2abx + b^2x^2)/(a + bx)^2})])}{(6b^3)}$$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5781, 4926, 3042, 4269, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sec^{-1}(a + bx) dx$$

$$\downarrow 5781$$

$$\frac{\int b^2 x^2 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^3}$$

$$\downarrow 4926$$

$$\frac{\frac{1}{3} \int -b^3 x^3 d \sec^{-1}(a + bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a + bx)}{b^3}$$

$$\downarrow 3042$$

$$\frac{\frac{1}{3} \int (a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}))^3 d \sec^{-1}(a + bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a + bx)}{b^3}$$

$$\downarrow 4269$$

$$\frac{\frac{1}{3} \left(\frac{1}{2} \int (2a^3 + 5(a + bx)^2 a - (6a^2 + 1)(a + bx)) d \sec^{-1}(a + bx) - \frac{1}{2} bx(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) + \frac{1}{3} b^3 x^3 \sec^{-1}(a + bx)}{b^3}$$

$$\downarrow 2009$$

$$\frac{\frac{1}{3} \left(\frac{1}{2} \left(2a^3 \sec^{-1}(a+bx) - (6a^2+1) \operatorname{arctanh} \left(\sqrt{1 - \frac{1}{(a+bx)^2}} \right) + 5a(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) - \frac{1}{2} bx(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{b^3}$$

input `Int[x^2*ArcSec[a + b*x], x]`

output `((b^3*x^3*ArcSec[a + b*x])/3 + (-1/2*(b*x*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]) + (5*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)] + 2*a^3*ArcSec[a + b*x] - (1 + 6*a^2)*ArcTanh[Sqrt[1 - (a + b*x)^(-2)]])/2)/3)/b^3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4269 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^n_, x_Symbol] :=> Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Simp[1/(n - 1) Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]`

rule 4926 `Int[((e_.) + (f_.)*(x_))^m_*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^n_*(e_.) + (f_.)*(x_))^m_*(a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 5781 `Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^p_*((e_.) + (f_.)*(x_))^m_*(a + b*Sec[c + d*x])^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.64

method	result
derivativedivides	$-\frac{\operatorname{arcsec}(bx+a)a^3}{3} + \operatorname{arcsec}(bx+a)a^2(bx+a) - \operatorname{arcsec}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^3}{3} - \frac{\sqrt{(bx+a)^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \right)}{b^3}$
default	$-\frac{\operatorname{arcsec}(bx+a)a^3}{3} + \operatorname{arcsec}(bx+a)a^2(bx+a) - \operatorname{arcsec}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^3}{3} - \frac{\sqrt{(bx+a)^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \right)}{b^3}$
parts	$\frac{x^3 \operatorname{arcsec}(bx+a)}{3} - \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2} + 6 \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2-1}\sqrt{b^2}}{\sqrt{b^2}}\right) \right)}{6b^3 \sqrt{b^2x^2+2abx+a^2-1}}$

```
input int(x^2*arcsec(b*x+a), x, method=_RETURNVERBOSE)
```

```
output 1/b^3*(-1/3*arcsec(b*x+a)*a^3+arcsec(b*x+a)*a^2*(b*x+a)-arcsec(b*x+a)*a*(b*x+a)^2+1/3*arcsec(b*x+a)*(b*x+a)^3-1/6*((b*x+a)^2-1)^(1/2)*(2*a^3*arctan(1/((b*x+a)^2-1)^(1/2))+6*a^2*ln(b*x+a+((b*x+a)^2-1)^(1/2))-6*a*((b*x+a)^2-1)^(1/2)+(b*x+a)*((b*x+a)^2-1)^(1/2)+ln(b*x+a+((b*x+a)^2-1)^(1/2)))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.01

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{2b^3x^3 \operatorname{arcsec}(bx+a) + 4a^3 \arctan(-bx-a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) + (6a^2 + 1) \log(-bx-a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{6b^3}$$

```
input integrate(x^2*arcsec(b*x+a), x, algorithm="fricas")
```

output

```
1/6*(2*b^3*x^3*arcsec(b*x + a) + 4*a^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*
a*b*x + a^2 - 1)) + (6*a^2 + 1)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^
2 - 1)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(b*x - 5*a))/b^3
```

Sympy [F]

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \operatorname{asec}(a + bx) dx$$

input

```
integrate(x**2*asec(b*x+a),x)
```

output

```
Integral(x**2*asec(a + b*x), x)
```

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \operatorname{arcsec}(bx + a) dx$$

input

```
integrate(x^2*arcsec(b*x+a),x, algorithm="maxima")
```

output

```
1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/3*(b^2*x
^4 + a*b*x^3)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2
*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x
+ a - 1)) - 1), x)
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. $2(100) = 200$.

Time = 0.16 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.78

$$\int x^2 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{24} b \left(\frac{8 (bx + a)^3 \left(\frac{3a}{bx+a} - \frac{3a^2}{(bx+a)^2} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right) - (bx + a)^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)}{b^4} \right)$$

input `integrate(x^2*arcsec(b*x+a),x, algorithm="giac")`

output `-1/24*b*(8*(b*x + a)^3*(3*a/(b*x + a) - 3*a^2/(b*x + a)^2 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^4 - ((b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 4*(6*a^2 + 1))*log(1/2*abs(b*x + a)*abs(-2*sqrt(-1/(b*x + a)^2 + 1) + 2)) - (12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)/((b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2))/b^4)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \operatorname{acos} \left(\frac{1}{a + bx} \right) dx$$

input `int(x^2*acos(1/(a + b*x)),x)`

output `int(x^2*acos(1/(a + b*x)), x)`

Reduce [F]

$$\int x^2 \sec^{-1}(a + bx) dx = \int a \sec(bx + a) x^2 dx$$

input `int(x^2*asec(b*x+a),x)`

output `int(asec(a + b*x)*x**2,x)`

3.21 $\int x \sec^{-1}(a + bx) dx$

Optimal result	170
Mathematica [A] (verified)	170
Rubi [A] (verified)	171
Maple [A] (verified)	173
Fricas [A] (verification not implemented)	174
Sympy [F]	175
Maxima [F]	175
Giac [A] (verification not implemented)	175
Mupad [F(-1)]	176
Reduce [F]	176

Optimal result

Integrand size = 8, antiderivative size = 78

$$\int x \sec^{-1}(a + bx) dx = -\frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \sec^{-1}(a + bx)}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx) + \frac{a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^2}$$

output `-1/2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2-1/2*a^2*arcsec(b*x+a)/b^2+1/2*x^2*arcsec(b*x+a)+a*arctanh((1-1/(b*x+a)^2)^(1/2))/b^2`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.41

$$\int x \sec^{-1}(a + bx) dx = \frac{-\left((a + bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right) + b^2x^2 \sec^{-1}(a + bx) + a^2 \arcsin\left(\frac{1}{a+bx}\right) + 2a \log\left((a + bx)\left(1 + \sqrt{\frac{-1+a^2}{(a+bx)^2}}\right)\right)}{2b^2}$$

input `Integrate[x*ArcSec[a + b*x],x]`

output

```
(-((a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]) + b^2*x^2*ArcSec[a + b*x] + a^2*ArcSin[(a + b*x)^(-1)] + 2*a*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]])]/(2*b^2)
```

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$, Rules used = {5781, 25, 4926, 3042, 4260, 3042, 4254, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \sec^{-1}(a + bx) dx$$

$$\downarrow 5781$$

$$\frac{\int bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow 25$$

$$-\frac{\int -bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow 4926$$

$$\frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx) - \frac{1}{2} \int b^2x^2 d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow 3042$$

$$\frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx) - \frac{1}{2} \int (a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2}))^2 d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow 4260$$

$$\frac{\frac{1}{2}(2a \int (a + bx) d \sec^{-1}(a + bx) - \int (a + bx)^2 d \sec^{-1}(a + bx) + a^2(-\sec^{-1}(a + bx))) + \frac{1}{2}b^2x^2 \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow 3042$$

$$\frac{\frac{1}{2} \left(2a \int \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2} \right) d \sec^{-1}(a+bx) - \int \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2} \right)^2 d \sec^{-1}(a+bx) + a^2 \left(-\sec^{-1}(a+bx) \right) \right)}{b^2}$$

↓ 4254

$$\frac{\frac{1}{2} \left(\int 1 d \left(- \left((a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) \right) + 2a \int \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2} \right) d \sec^{-1}(a+bx) + a^2 \left(-\sec^{-1}(a+bx) \right) \right)}{b^2}$$

↓ 24

$$\frac{\frac{1}{2} \left(2a \int \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2} \right) d \sec^{-1}(a+bx) + a^2 \left(-\sec^{-1}(a+bx) \right) - (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) + \frac{1}{2} b^2 x^2 \sec^{-1}(a+bx)}{b^2}$$

↓ 4257

$$\frac{\frac{1}{2} \left(a^2 \left(-\sec^{-1}(a+bx) \right) + 2a \operatorname{arctanh} \left(\sqrt{1 - \frac{1}{(a+bx)^2}} \right) - (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \right) + \frac{1}{2} b^2 x^2 \sec^{-1}(a+bx)}{b^2}$$

input

```
Int[x*ArcSec[a + b*x], x]
```

output

```
((b^2*x^2*ArcSec[a + b*x])/2 + (-((a + b*x)*Sqrt[1 - (a + b*x)^(-2)])) - a^2*ArcSec[a + b*x] + 2*a*ArcTanh[Sqrt[1 - (a + b*x)^(-2)]])/2/b^2
```

Defintions of rubi rules used

rule 24

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[-d^{(-1)} \text{Subst}[\text{Int}[\text{Exp}[\text{andIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4260 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.) + (a_.))^2, x_Symbol] \rightarrow \text{Simp}[a^2*x, x] + (\text{Simp}[2*a*b \text{Int}[\text{Csc}[c + d*x], x], x] + \text{Simp}[b^2 \text{Int}[\text{Csc}[c + d*x]^2, x], x]) /; \text{FreeQ}[\{a, b, c, d\}, x]$

rule 4926 $\text{Int}[(e_.) + (f_.)(x_)]^{(m_.)}*\text{Sec}[(c_.) + (d_.)(x_)]*((a_.) + (b_.)*\text{Sec}[(c_.) + (d_.)(x_)]^{(n_.)}*\text{Tan}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*((a + b*\text{Sec}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] - \text{Simp}[f*(m/(b*d*(n + 1))) \text{Int}[(e + f*x)^{(m - 1)}*(a + b*\text{Sec}[c + d*x])^{(n + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$

rule 5781 $\text{Int}[(a_.) + \text{ArcSec}[(c_.) + (d_.)(x_)]*(b_.)]^{(p_.)}*((e_.) + (f_.)(x_)]^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/d^{(m + 1)} \text{Subst}[\text{Int}[(a + b*x)^p*\text{Sec}[x]*\text{Tan}[x]*(d*e - c*f + f*\text{Sec}[x])^m, x], x, \text{ArcSec}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[m]$

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.38

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsec}(bx+a)(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a) + \frac{\sqrt{(bx+a)^2-1} \left(2a \ln(bx+a + \sqrt{(bx+a)^2-1}) - \sqrt{(bx+a)^2-1} \right)}{2(bx+a)\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}}}{b^2}$
default	$\frac{\frac{\operatorname{arcsec}(bx+a)(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a) + \frac{\sqrt{(bx+a)^2-1} \left(2a \ln(bx+a + \sqrt{(bx+a)^2-1}) - \sqrt{(bx+a)^2-1} \right)}{2(bx+a)\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}}}{b^2}$
parts	$\frac{x^2 \operatorname{arcsec}(bx+a)}{2} + \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(a^2 \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2+2a} \ln\left(\frac{b^2x + \sqrt{b^2x^2+2abx+a^2-1} \sqrt{b^2+2a}}{\sqrt{b^2+2a}}\right) \right)}{2b^2\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}} (bx+a)\sqrt{b^2}}$

input `int(x*arcsec(b*x+a), x, method=_RETURNVERBOSE)`

output
$$\frac{1}{b^2} \left(\frac{1}{2} \operatorname{arcsec}(bx+a) (bx+a)^2 - \operatorname{arcsec}(bx+a) a (bx+a) + \frac{1}{2} (bx+a)^2 - 1 \right)^{1/2} \left(2a \ln(bx+a + \sqrt{(bx+a)^2-1}) - \sqrt{(bx+a)^2-1} \right) - \frac{\sqrt{(bx+a)^2-1}}{(bx+a)^2} \right) / (bx+a)$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.33

$$\int x \sec^{-1}(a + bx) dx$$

$$= \frac{b^2 x^2 \operatorname{arcsec}(bx+a) - 2a^2 \arctan(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) - 2a \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{2b^2}$$

input `integrate(x*arcsec(b*x+a), x, algorithm="fricas")`

output
$$\frac{1}{2} (b^2 x^2 \operatorname{arcsec}(bx+a) - 2a^2 \arctan(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) - 2a \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) - \sqrt{b^2x^2 + 2abx + a^2 - 1}) / b^2$$

Sympy [F]

$$\int x \sec^{-1}(a + bx) dx = \int x \operatorname{asec}(a + bx) dx$$

input `integrate(x*asec(b*x+a),x)`

output `Integral(x*asec(a + b*x), x)`

Maxima [F]

$$\int x \sec^{-1}(a + bx) dx = \int x \operatorname{arcsec}(bx + a) dx$$

input `integrate(x*arcsec(b*x+a),x, algorithm="maxima")`

output `1/2*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/2*(b^2*x^3 + a*b*x^2)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x + a - 1)) - 1), x)`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.74

$$\int x \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{4} b \left(\frac{2 (bx + a)^2 \left(\frac{2a}{bx+a} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{b^3} + \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + 4a \log \left(\frac{1}{2} \right)}{b^3} \right)$$

input `integrate(x*arcsec(b*x+a),x, algorithm="giac")`

output

```
-1/4*b*(2*(b*x + a)^2*(2*a/(b*x + a) - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^3 + ((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 4*a*log(1/2*abs(b*x + a)*abs(-2*sqrt(-1/(b*x + a)^2 + 1) + 2)) - 1/((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1)))/b^3)
```

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx) dx = \int x \arccos\left(\frac{1}{a + bx}\right) dx$$

input

```
int(x*acos(1/(a + b*x)),x)
```

output

```
int(x*acos(1/(a + b*x)), x)
```

Reduce [F]

$$\int x \sec^{-1}(a + bx) dx = \int a \sec(bx + a) x dx$$

input

```
int(x*asec(b*x+a),x)
```

output

```
int(asec(a + b*x)*x,x)
```

3.22 $\int \sec^{-1}(a + bx) dx$

Optimal result	177
Mathematica [C] (verified)	177
Rubi [A] (warning: unable to verify)	178
Maple [A] (verified)	180
Fricas [B] (verification not implemented)	180
Sympy [F]	181
Maxima [A] (verification not implemented)	181
Giac [B] (verification not implemented)	181
Mupad [B] (verification not implemented)	182
Reduce [F]	182

Optimal result

Integrand size = 6, antiderivative size = 37

$$\int \sec^{-1}(a + bx) dx = \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b}$$

output

```
(b*x+a)*arcsec(b*x+a)/b-arctanh((1-1/(b*x+a)^2)^(1/2))/b
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.94 (sec) , antiderivative size = 468, normalized size of antiderivative = 12.65

$$\int \sec^{-1}(a + bx) dx = x \sec^{-1}(a + bx) + \frac{(a + bx) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} \left(\sqrt[4]{-1}(-i + \sqrt{-1 + a^2}) \sqrt{2i - ia^2 + 2\sqrt{-1 + a^2}} \arctan\left(\frac{(-1)^{3/4} \sqrt{2i - ia^2 + 2\sqrt{-1 + a^2}}}{a\sqrt{-1+a^2} - a\sqrt{-1+a^2}}\right) \right)}{b}$$

input

```
Integrate[ArcSec[a + b*x], x]
```

output

```
x*ArcSec[a + b*x] + ((a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + a^2])*Sqrt[2*I - I*a^2 + 2*Sqrt[-1 + a^2]])*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*a^2 + 2*Sqrt[-1 + a^2]]*b*x)/(a*Sqrt[-1 + a^2] - a*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])] + (-1)^(3/4)*(I + Sqrt[-1 + a^2])*Sqrt[-2*I + I*a^2 + 2*Sqrt[-1 + a^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*a^2 + 2*Sqrt[-1 + a^2]]*b*x)/(a*Sqrt[-1 + a^2] - a*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])] + a*(a*ArcTan[(Sqrt[-1 + a^2]*b^2*x^2)/(a^4 + a^3*b*x + b^2*x^2 - a^2*(1 + Sqrt[-1 + a^2])*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])]) - Log[Sqrt[-1 + a^2] - b*x - Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2]] + Log[b^2*(Sqrt[-1 + a^2] + b*x - Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])])]/(a*b*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])
```

Rubi [A] (warning: unable to verify)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5773, 895, 798, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{-1}(a + bx) dx \\
 & \quad \downarrow \text{5773} \\
 & \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \int \frac{1}{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}} dx \\
 & \quad \downarrow \text{895} \\
 & \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\int \frac{1}{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}} d(a + bx)}{b} \\
 & \quad \downarrow \text{798} \\
 & \frac{\int \frac{(a + bx)^2}{\sqrt{-a - bx + 1}} d \frac{1}{(a + bx)^2}}{2b} + \frac{(a + bx) \sec^{-1}(a + bx)}{b} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{(a+bx)\sec^{-1}(a+bx)}{b} - \frac{\int \frac{1}{1-\frac{1}{(a+bx)^4}} d\sqrt{-a-bx+1}}{b}$$

↓ 219

$$\frac{(a+bx)\sec^{-1}(a+bx)}{b} - \frac{\operatorname{arctanh}(\sqrt{-a-bx+1})}{b}$$

input `Int[ArcSec[a + b*x], x]`

output `((a + b*x)*ArcSec[a + b*x])/b - ArcTanh[Sqrt[1 - a - b*x]]/b`

Definitions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 895 `Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_.), x_Symbol] := Simp[u^m/(Coefficient[v, x, 1]*v^m) Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]`

rule 5773 `Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{(bx+a) \operatorname{arcsec}(bx+a) - \ln\left(bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{b}$
default	$\frac{(bx+a) \operatorname{arcsec}(bx+a) - \ln\left(bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{b}$
parts	$x \operatorname{arcsec}(bx+a) - \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(a \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2} + \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2-1}}{\sqrt{b^2}}\right) \right)}{b\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}} (bx+a)\sqrt{b^2}}$

input `int(arcsec(b*x+a),x,method=_RETURNVERBOSE)`output `1/b*((b*x+a)*arcsec(b*x+a)-ln(b*x+a+(b*x+a)*(1-1/(b*x+a)^2)^(1/2)))`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(35) = 70.

Time = 0.13 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.97

$$\int \sec^{-1}(a+bx) dx$$

$$= \frac{bx \operatorname{arcsec}(bx+a) + 2a \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) + \log(-bx-a+\sqrt{b^2x^2+2abx+a^2-1})}{b}$$

input `integrate(arcsec(b*x+a),x, algorithm="fricas")`output `(b*x*arcsec(b*x+a) + 2*a*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)))/b`

Sympy [F]

$$\int \sec^{-1}(a + bx) dx = \int \operatorname{asec}(a + bx) dx$$

input `integrate(asec(b*x+a), x)`

output `Integral(asec(a + b*x), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.49

$$\int \sec^{-1}(a + bx) dx$$

$$= \frac{2(bx + a) \operatorname{arcsec}(bx + a) - \log\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right)}{2b}$$

input `integrate(arcsec(b*x+a), x, algorithm="maxima")`

output `1/2*(2*(b*x + a)*arcsec(b*x + a) - log(sqrt(-1/(b*x + a)^2 + 1) + 1) + log(-sqrt(-1/(b*x + a)^2 + 1) + 1))/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(35) = 70$.

Time = 0.14 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.22

$$\int \sec^{-1}(a + bx) dx$$

$$= \frac{1}{2} b \left(\frac{2(bx + a) \arccos\left(-\frac{1}{(bx+a)\left(\frac{a}{bx+a}-1\right)-a}\right)}{b^2} - \frac{\log\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right)}{b^2} \right)$$

input `integrate(arcsec(b*x+a),x, algorithm="giac")`

output `1/2*b*(2*(b*x + a)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^2 - (log(sqrt(-1/(b*x + a)^2 + 1) + 1) - log(-sqrt(-1/(b*x + a)^2 + 1) + 1))/b^2)`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \sec^{-1}(a + bx) dx = -\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1-\frac{1}{(a+bx)^2}}}\right) - \operatorname{acos}\left(\frac{1}{a+bx}\right) (a + bx)}{b}$$

input `int(acos(1/(a + b*x)),x)`

output `-(atanh(1/(1 - 1/(a + b*x)^2)^(1/2)) - acos(1/(a + b*x))*(a + b*x))/b`

Reduce [F]

$$\int \sec^{-1}(a + bx) dx = \int \operatorname{asec}(bx + a) dx$$

input `int(asec(b*x+a),x)`

output `int(asec(a + b*x),x)`

3.23 $\int \frac{\sec^{-1}(a+bx)}{x} dx$

Optimal result	183
Mathematica [A] (verified)	184
Rubi [A] (verified)	185
Maple [A] (verified)	189
Fricas [F]	190
Sympy [F]	190
Maxima [F]	191
Giac [F]	191
Mupad [F(-1)]	191
Reduce [F]	192

Optimal result

Integrand size = 10, antiderivative size = 200

$$\int \frac{\sec^{-1}(a+bx)}{x} dx = \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) - \sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right) - i \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) - i \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) + \frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)$$

output

```
arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-I*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))-I*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+1/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)
```


Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.42

$$\begin{aligned}
\int \frac{\sec^{-1}(a + bx)}{x} dx = & -4i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \arctan\left(\frac{(1+a) \tan\left(\frac{1}{2} \sec^{-1}(a + bx)\right)}{\sqrt{1-a^2}}\right) \\
& + \left(\sec^{-1}(a + bx) - 2 \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right)\right) \log\left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a}\right) + \left(\sec^{-1}(a + bx) \right. \\
& \left. + 2 \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right)\right) \log\left(1 - \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a}\right) \\
& - \sec^{-1}(a + bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right) \\
& - i \left(\text{PolyLog}\left(2, -\frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a}\right) \right. \\
& \quad \left. + \text{PolyLog}\left(2, \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a}\right)\right) \\
& + \frac{1}{2} i \text{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)
\end{aligned}$$

input `Integrate[ArcSec[a + b*x]/x,x]`output `(-4*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*ArcTan[(((1 + a)*Tan[ArcSec[a + b*x]/2])/Sqrt[1 - a^2]] + (ArcSec[a + b*x] - 2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]])*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + (ArcSec[a + b*x] + 2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]])*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] - ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] - I*(PolyLog[2, -(((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a)] + PolyLog[2, ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a]) + (I/2)*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])]`

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.32, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$, Rules used = {5781, 25, 5062, 5041, 25, 3042, 4202, 2620, 2715, 2838, 5031, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(a+bx)}{x} dx \\
 & \quad \downarrow \text{5781} \\
 & \int \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{bx} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{25} \\
 & - \int - \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{bx} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{5062} \\
 & - \int \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{\frac{a}{a+bx} - 1} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{5041} \\
 & \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx) d \sec^{-1}(a+bx) - \\
 & \quad a \int - \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{25} \\
 & \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx) d \sec^{-1}(a+bx) + \\
 & \quad a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \int \sec^{-1}(a+bx) \tan(\sec^{-1}(a+bx)) d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{4202} \\
& -2i \int \frac{e^{2i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)}{1 + e^{2i \sec^{-1}(a+bx)}} d \sec^{-1}(a+bx) + \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{2} i \sec^{-1}(a+bx)^2 \\
& \quad \downarrow \text{2620} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) - \\
& 2i \left(\frac{1}{2} i \int \log(1 + e^{2i \sec^{-1}(a+bx)}) d \sec^{-1}(a+bx) - \frac{1}{2} i \sec^{-1}(a+bx) \log(1 + e^{2i \sec^{-1}(a+bx)}) \right) + \\
& \quad \frac{1}{2} i \sec^{-1}(a+bx)^2 \\
& \quad \downarrow \text{2715} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) - \\
& 2i \left(\frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \log(1 + e^{2i \sec^{-1}(a+bx)}) d e^{2i \sec^{-1}(a+bx)} - \frac{1}{2} i \sec^{-1}(a+bx) \log(1 + e^{2i \sec^{-1}(a+bx)}) \right) + \\
& \quad \frac{1}{2} i \sec^{-1}(a+bx)^2 \\
& \quad \downarrow \text{2838} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) - \\
& 2i \left(-\frac{1}{4} \text{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) - \frac{1}{2} i \sec^{-1}(a+bx) \log(1 + e^{2i \sec^{-1}(a+bx)}) \right) + \\
& \quad \frac{1}{2} i \sec^{-1}(a+bx)^2 \\
& \quad \downarrow \text{5031}
\end{aligned}$$

$$a \left(-i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)}{-e^{i \sec^{-1}(a+bx)} a - \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)}{-e^{i \sec^{-1}(a+bx)} a + \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - \right. \\ \left. 2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \right) + \right. \\ \left. \frac{1}{2} i \sec^{-1}(a+bx)^2 \right)$$

↓ 2620

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{i \int \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) d \sec^{-1}(a+bx)}{a} \right) - i \left(\frac{i \sec^{-1}(a+bx)}{a} - \frac{1}{a} \int \frac{e^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} d \sec^{-1}(a+bx) \right) \right. \\ \left. 2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \right) + \right. \\ \left. \frac{1}{2} i \sec^{-1}(a+bx)^2 \right)$$

↓ 2715

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{\int e^{-i \sec^{-1}(a+bx)} \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) d e^{i \sec^{-1}(a+bx)}}{a} \right) - i \left(\frac{i \sec^{-1}(a+bx)}{a} - \frac{1}{a} \int \frac{e^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} d \sec^{-1}(a+bx) \right) \right. \\ \left. 2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \right) + \right. \\ \left. \frac{1}{2} i \sec^{-1}(a+bx)^2 \right)$$

↓ 2838

$$a \left(-i \left(\frac{\text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} + \frac{i \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} \right) - i \left(\frac{\text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right)}{a} - \frac{1}{a} \int \frac{e^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} d \sec^{-1}(a+bx) \right) \right. \\ \left. 2i \left(-\frac{1}{4} \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \right) + \right. \\ \left. \frac{1}{2} i \sec^{-1}(a+bx)^2 \right)$$

input `Int[ArcSec[a + b*x]/x,x]`

output

```
(I/2)*ArcSec[a + b*x]^2 + a*(((1/2*I)*ArcSec[a + b*x]^2)/a - I*((I*ArcSec
[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/a + Poly
Log[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])]/a) - I*((I*ArcSec[a
+ b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/a + PolyLog
[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])]/a) - (2*I)*((1/2*I)*A
rcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] - PolyLog[2, -E^((2*I)*A
rcSec[a + b*x])])/4)
```

Definitions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x
)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4202

```
Int[(((c_) + (d_)*(x_)^(m_))*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(
e + f*x))/(1 + E^(2*I*(e + f*x))))], x], x] /; FreeQ[{c, d, e, f}, x] && IGt
Q[m, 0]
```

rule 5031

```
Int[(((e_.) + (f_.)*(x_)^(m_.)*Sin[(c_.) + (d_.)*(x_)])/(Cos[(c_.) + (d_.)
*(x_)*(b_.) + (a_.)), x_Symbol] := Simp[I*((e + f*x)^(m + 1)/(b*f*(m + 1)))
, x] + (-Simp[I Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a - Rt[a^2 - b^2, 2] +
b*E^(I*(c + d*x)))]), x], x] - Simp[I Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a
+ Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x)))]), x], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

rule 5041

```
Int[(((e_.) + (f_.)*(x_)^(m_.)*Tan[(c_.) + (d_.)*(x_)^(n_.)]/(Cos[(c_.) +
(d_.)*(x_)*(b_.) + (a_.)), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Tan[c +
d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Sin[c + d*x]*(Tan[c + d*x]^(n
- 1)/(a + b*Cos[c + d*x]))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[
m, 0] && IGtQ[n, 0]
```

rule 5062

```
Int[(((e_.) + (f_.)*(x_)^(m_.)*(F_)[(c_.) + (d_.)*(x_)^(n_.)*(G_)[(c_.) +
(d_.)*(x_)^(p_.)]/(a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])], x_Symbol] := In
t[(e + f*x)^m*Cos[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cos[c + d*x]))
, x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m
, n, p]
```

rule 5781

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)*(b_.)]^(p_.)*((e_.) + (f_.)*(x_)^(m
_.)), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d
*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e
, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.87

method	result
derivativedivides	$\operatorname{arcsec}(bx + a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right) + \operatorname{arcsec}(bx + a) \ln \left(\frac{a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right)$
default	$\operatorname{arcsec}(bx + a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right) + \operatorname{arcsec}(bx + a) \ln \left(\frac{a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right)$

input `int(arcsec(b*x+a)/x,x,method=_RETURNVERBOSE)`

output `arcsec(b*x+a)*ln((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+arcsec(b*x+a)*ln((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))-arcsec(b*x+a)*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-arcsec(b*x+a)*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-I*dilog((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-I*dilog((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))`

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x} dx$$

input `integrate(arcsec(b*x+a)/x,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asec}(a + bx)}{x} dx$$

input `integrate(asec(b*x+a)/x,x)`

output `Integral(asec(a + b*x)/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x} dx$$

input `integrate(arcsec(b*x+a)/x,x, algorithm="maxima")`

output `integrate(arcsec(b*x + a)/x, x)`

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x} dx$$

input `integrate(arcsec(b*x+a)/x,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x} dx$$

input `int(arccos(1/(a + b*x))/x,x)`

output `int(arccos(1/(a + b*x))/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{asec}(bx + a)}{x} dx$$

input `int(asec(b*x+a)/x,x)`

output `int(asec(a + b*x)/x,x)`

3.24 $\int \frac{\sec^{-1}(a+bx)}{x^2} dx$

Optimal result	193
Mathematica [C] (verified)	194
Rubi [A] (verified)	194
Maple [B] (verified)	197
Fricas [B] (verification not implemented)	197
Sympy [F]	198
Maxima [F]	198
Giac [A] (verification not implemented)	199
Mupad [F(-1)]	199
Reduce [F]	200

Optimal result

Integrand size = 10, antiderivative size = 70

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = -\frac{b \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{x} + \frac{2b \arctan\left(\frac{\sqrt{1+a} \tan\left(\frac{1}{2} \sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}$$

output `-b*arcsec(b*x+a)/a-arcsec(b*x+a)/x+2*b*arctan(((1+a)^(1/2)*tan(1/2*arcsec(b*x+a)))/(1-a)^(1/2))/a/(-a^2+1)^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.60

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx$$

$$= -\frac{\sec^{-1}(a + bx)}{x} + \frac{b \left(\arcsin\left(\frac{1}{a+bx}\right) - \frac{i \log\left(\frac{2\left(\frac{ia(-1+a^2+abx)}{\sqrt{1-a^2}} + a(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{bx}\right)}{\sqrt{1-a^2}} \right)}{a}$$

input `Integrate[ArcSec[a + b*x]/x^2,x]`

output `-(ArcSec[a + b*x]/x) + (b*(ArcSin[(a + b*x)^(-1)] - (I*Log[(2*((I*a*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] + a*(a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/(b*x)])/Sqrt[1 - a^2]))/a`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5781, 4926, 3042, 4270, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx$$

$$\downarrow \text{5781}$$

$$b \int \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^2 x^2} d \sec^{-1}(a + bx)$$

$$\begin{aligned}
& \downarrow 4926 \\
& b \left(- \int - \frac{1}{bx} d \sec^{-1}(a + bx) - \frac{\sec^{-1}(a + bx)}{bx} \right) \\
& \downarrow 3042 \\
& b \left(- \int \frac{1}{a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2})} d \sec^{-1}(a + bx) - \frac{\sec^{-1}(a + bx)}{bx} \right) \\
& \downarrow 4270 \\
& b \left(\frac{\int \frac{1}{1 - \frac{a}{a+bx}} d \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{bx} \right) \\
& \downarrow 3042 \\
& b \left(\frac{\int \frac{1}{1 - a \sin(\sec^{-1}(a+bx) + \frac{\pi}{2})} d \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{bx} \right) \\
& \downarrow 3138 \\
& b \left(\frac{2 \int \frac{1}{(a+1) \tan^2(\frac{1}{2} \sec^{-1}(a+bx)) - a + 1} d \tan(\frac{1}{2} \sec^{-1}(a + bx))}{a} - \frac{\sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{bx} \right) \\
& \downarrow 218 \\
& b \left(\frac{2 \arctan\left(\frac{\sqrt{a+1} \tan(\frac{1}{2} \sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a \sqrt{1-a^2}} - \frac{\sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{bx} \right)
\end{aligned}$$

input `Int[ArcSec[a + b*x]/x^2,x]`

output `b*(-(ArcSec[a + b*x]/a) - ArcSec[a + b*x]/(b*x) + (2*ArcTan[(Sqrt[1 + a]*Tan[ArcSec[a + b*x]/2])/Sqrt[1 - a]])/(a*Sqrt[1 - a^2]))`

Definitions of rubi rules used

- rule 218 $\text{Int}[\{(a_) + (b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3138 $\text{Int}[\{(a_) + (b_)*\sin[\text{Pi}/2 + (c_) + (d_)*(x_)]\}^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x\}, \text{Simp}[2*(e/d) \ \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4270 $\text{Int}[(\text{csc}[(c_) + (d_)*(x_)]*(b_) + (a_))^{-1}, x_Symbol] \rightarrow \text{Simp}[x/a, x] - \text{Simp}[1/a \ \text{Int}[1/(1 + (a/b)*\text{Sin}[c + d*x]), x], x] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 4926 $\text{Int}[\{(e_) + (f_)*(x_)^{(m_)}*\text{Sec}[(c_) + (d_)*(x_)]*\{(a_) + (b_)*\text{Sec}[(c_) + (d_)*(x_)]\}^{(n_)}*\text{Tan}[(c_) + (d_)*(x_)]\}, x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*((a + b*\text{Sec}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] - \text{Simp}[f*(m/(b*d*(n + 1))) \ \text{Int}[(e + f*x)^{(m - 1)}*(a + b*\text{Sec}[c + d*x])^{(n + 1)}, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$
- rule 5781 $\text{Int}[\{(a_) + \text{ArcSec}[(c_) + (d_)*(x_)]*(b_)\}^{(p_)}*\{(e_) + (f_)*(x_)\}^{(m_)}\}, x_Symbol] \rightarrow \text{Simp}[1/d^{(m + 1)} \ \text{Subst}[\text{Int}[(a + b*x)^p*\text{Sec}[x]*\text{Tan}[x]*(d*e - c*f + f*\text{Sec}[x])^m, x], x, \text{ArcSec}[c + d*x]], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[m]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. 2(62) = 124.

Time = 0.35 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.80

method	result
derivativedivides	$b \left(-\frac{\operatorname{arcsec}(bx+a)}{bx} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2\sqrt{a^2-1}\sqrt{(bx+a)^2-1}+2(bx+a)a-2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}} \right)$
default	$b \left(-\frac{\operatorname{arcsec}(bx+a)}{bx} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2\sqrt{a^2-1}\sqrt{(bx+a)^2-1}+2(bx+a)a-2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}} \right)$
parts	$-\frac{\operatorname{arcsec}(bx+a)}{x} + \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left(\arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}\sqrt{b^2x^2+2abx+a^2-1}}{x}\right) \right)}{\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}}$

input `int(arcsec(b*x+a)/x^2,x,method=_RETURNVERBOSE)`

output `b*(-1/b/x*arcsec(b*x+a)+((b*x+a)^2-1)^(1/2)*(arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(1/2)-ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+(b*x+a)*a-1)/b/x))/((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a)/a/(a^2-1)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(62) = 124.

Time = 0.13 (sec) , antiderivative size = 281, normalized size of antiderivative = 4.01

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx$$

$$= \left[\frac{2(a^2-1)bx \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) - \sqrt{a^2-1}bx \log\left(\frac{a^2bx+a^3+\sqrt{b^2x^2+2abx+a^2-1}}{(a^3-a)x}\right)}{2(a^2-1)bx \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) - 2\sqrt{-a^2+1}bx \arctan\left(-\frac{\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}}{a^2-1}\right)}{(a^3-a)x} \right]$$

input `integrate(arcsec(b*x+a)/x^2,x, algorithm="fricas")`

output `[-(2*(a^2 - 1)*b*x*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - sqrt(a^2 - 1)*b*x*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 - sqrt(a^2 - 1)*a - 1) - (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) + (a^3 - a)*arcsec(b*x + a))/((a^3 - a)*x), -(2*(a^2 - 1)*b*x*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - 2*sqrt(-a^2 + 1)*b*x*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) + (a^3 - a)*arcsec(b*x + a))/((a^3 - a)*x)]`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asec}(a + bx)}{x^2} dx$$

input `integrate(asec(b*x+a)/x**2,x)`

output `Integral(asec(a + b*x)/x**2, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x^2} dx$$

input `integrate(arcsec(b*x+a)/x^2,x, algorithm="maxima")`

output `(x*integrate((b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1)) / (b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x + (b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x)* e^(log(b*x + a + 1) + log(b*x + a - 1))), x) - arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))/x`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.34

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx$$

$$= b \left(\frac{2 \arctan \left(\frac{(bx+a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + a}{\sqrt{-a^2 + 1}} \right)}{\sqrt{-a^2 + 1} a} + \frac{\arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{a \left(\frac{a}{bx+a} - 1 \right)} \right)$$

input `integrate(arcsec(b*x+a)/x^2,x, algorithm="giac")`output `b*(2*arctan(((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + a)/sqrt(-a^2 + 1)) / (sqrt(-a^2 + 1)*a) + arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a)) / (a*(a/(b*x + a) - 1)))`**Mupad [F(-1)]**

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x^2} dx$$

input `int(arccos(1/(a + b*x))/x^2,x)`output `int(arccos(1/(a + b*x))/x^2, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{asec}(bx + a)}{x^2} dx$$

input `int(asec(b*x+a)/x^2,x)`

output `int(asec(a + b*x)/x**2,x)`

3.25 $\int \frac{\sec^{-1}(a+bx)}{x^3} dx$

Optimal result	201
Mathematica [C] (verified)	201
Rubi [A] (verified)	202
Maple [B] (verified)	206
Fricas [A] (verification not implemented)	207
Sympy [F]	207
Maxima [F]	208
Giac [B] (verification not implemented)	208
Mupad [F(-1)]	209
Reduce [F]	209

Optimal result

Integrand size = 10, antiderivative size = 125

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \sec^{-1}(a+bx)}{2a^2} - \frac{\sec^{-1}(a+bx)}{2x^2} - \frac{(1-2a^2)b^2 \arctan\left(\frac{\sqrt{1+a}\tan(\frac{1}{2}\sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}}$$

output

```
1/2*b*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a/(-a^2+1)/x+1/2*b^2*arcsec(b*x+a)/a^2
-1/2*arcsec(b*x+a)/x^2-(-2*a^2+1)*b^2*arctan((1+a)^(1/2)*tan(1/2*arcsec(b*
x+a))/(1-a)^(1/2))/a^2/(-a^2+1)^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.67 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.58

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx =$$

$$\frac{bx(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{a(-1+a^2)} + \sec^{-1}(a + bx) + \frac{b^2x^2 \arcsin\left(\frac{1}{a+bx}\right)}{a^2} + \frac{i(-1+2a^2)b^2x^2 \log\left(\frac{4(-1+a)a^2(1+a)\left(-\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}\right)}{(-1+2a^2)}\right)}{2x^2 a^2(1-a^2)^{3/2}}$$

input `Integrate[ArcSec[a + b*x]/x^3,x]`

output `-1/2*((b*x*(a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])/(a*(-1 + a^2)) + ArcSec[a + b*x] + (b^2*x^2*ArcSin[(a + b*x)^(-1)])/a^2 + (I*(-1 + 2*a^2)*b^2*x^2*Log[(4*(-1 + a)*a^2*(1 + a)*((-I)*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] - (a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]])/((-1 + 2*a^2)*b^2*x)]/(a^2*(1 - a^2)^(3/2)))/x^2`

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.18, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {5781, 25, 4926, 3042, 4272, 3042, 4407, 3042, 4318, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx$$

↓ 5781

$$b^2 \int \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3 x^3} dx - d \sec^{-1}(a + bx)$$

↓ 25

$$-b^2 \int -\frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3 x^3} dx - d \sec^{-1}(a + bx)$$

$$\begin{aligned} & \downarrow 4926 \\ & b^2 \left(\frac{1}{2} \int \frac{1}{b^2 x^2} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 3042 \\ & b^2 \left(\frac{1}{2} \int \frac{1}{\left(a - \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^2} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 4272 \\ & b^2 \left(\frac{1}{2} \left(\frac{\int -\frac{-a^2 - (a+bx)a+1}{bx} d \sec^{-1}(a+bx)}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 3042 \\ & b^2 \left(\frac{1}{2} \left(\frac{\int \frac{-a^2 - \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)a+1}{a - \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)} d \sec^{-1}(a+bx)}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 4407 \\ & b^2 \left(\frac{1}{2} \left(\frac{\frac{(1-2a^2) \int -\frac{a+bx}{a} d \sec^{-1}(a+bx)}{a(1-a^2)} + \frac{(1-a^2) \sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 3042 \\ & b^2 \left(\frac{1}{2} \left(\frac{\frac{(1-2a^2) \int \frac{\csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)}{a - \csc \left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)} d \sec^{-1}(a+bx)}{a(1-a^2)} + \frac{(1-a^2) \sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 4318 \\ & b^2 \left(\frac{1}{2} \left(\frac{\frac{(1-a^2) \sec^{-1}(a+bx)}{a} - \frac{(1-2a^2) \int \frac{1}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2 x^2} \right) \\ & \downarrow 3042 \end{aligned}$$

$$b^2 \left(\frac{1}{2} \left(\frac{(1-a^2) \sec^{-1}(a+bx)}{a} - \frac{(1-2a^2) \int \frac{1}{1-a \sin(\sec^{-1}(a+bx) + \frac{\pi}{2})} d \sec^{-1}(a+bx)}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 3138

$$b^2 \left(\frac{1}{2} \left(\frac{(1-a^2) \sec^{-1}(a+bx)}{a} - \frac{2(1-2a^2) \int \frac{1}{(a+1) \tan^2(\frac{1}{2} \sec^{-1}(a+bx)) - a+1} d \tan(\frac{1}{2} \sec^{-1}(a+bx))}{a(1-a^2)} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2x^2} \right)$$

↓ 218

$$b^2 \left(\frac{1}{2} \left(\frac{(1-a^2) \sec^{-1}(a+bx)}{a} - \frac{2(1-2a^2) \arctan\left(\frac{\sqrt{a+1} \tan(\frac{1}{2} \sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}} + \frac{\sqrt{1 - \frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} \right) - \frac{\sec^{-1}(a+bx)}{2b^2x^2} \right)$$

input

```
Int[ArcSec[a + b*x]/x^3,x]
```

output

```
b^2*(-1/2*ArcSec[a + b*x]/(b^2*x^2) + (((a + b*x)*Sqrt[1 - (a + b*x)^(-2)]
)/(a*(1 - a^2)*b*x) + (((1 - a^2)*ArcSec[a + b*x])/a - (2*(1 - 2*a^2)*ArcT
an[(Sqrt[1 + a]*Tan[ArcSec[a + b*x]/2])/Sqrt[1 - a]]/(a*Sqrt[1 - a^2]))/(
a*(1 - a^2)))/2)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3138 $\text{Int}[(a_ + (b_)\sin[\text{Pi}/2 + (c_) + (d_)(x_)])^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4272 $\text{Int}[(\text{csc}[(c_) + (d_)(x_)]*(b_) + (a_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b^2*\text{Cot}[c + d*x]*((a + b*\text{Csc}[c + d*x])^{(n + 1)}/(a*d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/(a*(n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Csc}[c + d*x])^{(n + 1)}*\text{Simp}[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*\text{Csc}[c + d*x] + b^2*(n + 2)*\text{Csc}[c + d*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

rule 4318 $\text{Int}[\text{csc}[(e_) + (f_)(x_)]/(\text{csc}[(e_) + (f_)(x_)]*(b_) + (a_)), x_Symbol] \rightarrow \text{Simp}[1/b \text{ Int}[1/(1 + (a/b)*\text{Sin}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

rule 4407 $\text{Int}[(\text{csc}[(e_) + (f_)(x_)]*(d_) + (c_))/(\text{csc}[(e_) + (f_)(x_)]*(b_) + (a_)), x_Symbol] \rightarrow \text{Simp}[c*(x/a), x] - \text{Simp}[(b*c - a*d)/a \text{ Int}[\text{Csc}[e + f*x]/(a + b*\text{Csc}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$

rule 4926 $\text{Int}[((e_) + (f_)(x_))^{(m_)}*\text{Sec}[(c_) + (d_)(x_)]*((a_) + (b_)*\text{Sec}[(c_) + (d_)(x_)])^{(n_)}*\text{Tan}[(c_) + (d_)(x_)], x_Symbol] \rightarrow \text{Simp}[(e + f*x)^m*((a + b*\text{Sec}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] - \text{Simp}[f*(m/(b*d*(n + 1))) \text{ Int}[(e + f*x)^{(m - 1)}*(a + b*\text{Sec}[c + d*x])^{(n + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[n, -1]$

rule 5781 $\text{Int}[((a_) + \text{ArcSec}[(c_) + (d_)(x_)]*(b_))^{(p_)}*((e_) + (f_)(x_))^{(m_)}, x_Symbol] \rightarrow \text{Simp}[1/d^{(m + 1)} \text{ Subst}[\text{Int}[(a + b*x)^p*\text{Sec}[x]*\text{Tan}[x]*(d*e - c*f + f*\text{Sec}[x])^m, x], x, \text{ArcSec}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{IntegerQ}[m]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 318 vs. 2(109) = 218.

Time = 0.33 (sec) , antiderivative size = 319, normalized size of antiderivative = 2.55

method	result
parts	$-\frac{\operatorname{arcsec}(bx+a)}{2x^2} - \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left((a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) a^2bx - 2 \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}}{x}\right) \right)}{2x^2}$
derivativedivides	$b^2 \left(-\frac{\operatorname{arcsec}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) (a^2-1)^{\frac{3}{2}} a^3 - \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) (a^2-1)^{\frac{3}{2}} a^2 (bx) \right)}{2b^2x^2} \right)$
default	$b^2 \left(-\frac{\operatorname{arcsec}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) (a^2-1)^{\frac{3}{2}} a^3 - \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) (a^2-1)^{\frac{3}{2}} a^2 (bx) \right)}{2b^2x^2} \right)$

input `int(arcsec(b*x+a)/x^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/2*\operatorname{arcsec}(b*x+a)/x^2 - 1/2*b*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}*((a^2-1)^{(3/2)}* \\ & \arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)})*a^2*b*x - 2*\ln(2*(a*b*x+(a^2-1)^{(1/2)} \\ &)*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}+a^2-1)/x)*a^4*b*x - b*\arctan(1/(b^2*x^2+2*a* \\ & b*x+a^2-1)^{(1/2)})*x*(a^2-1)^{(3/2)}+(a^2-1)^{(3/2)}*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)} \\ & *a^3*\ln(2*(a*b*x+(a^2-1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}+a^2-1)/x)* \\ & a^2*b*x - b*\ln(2*(a*b*x+(a^2-1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}+a^2-1)/x \\ &)*x)/((b^2*x^2+2*a*b*x+a^2-1)/(b*x+a)^2)^{(1/2)}/(b*x+a)/a^2/(a^2-1)^{(5/2)}/x \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 427, normalized size of antiderivative = 3.42

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx$$

$$= \frac{\left[(2a^2 - 1)\sqrt{a^2 - 1}b^2x^2 \log\left(\frac{a^2bx + a^3 + \sqrt{b^2x^2 + 2abx + a^2 - 1}(a^2 + \sqrt{a^2 - 1}a - 1) + (abx + a^2 - 1)\sqrt{a^2 - 1} - a}{x}\right) + 2(a^4 - 2a^2 + 1)\sqrt{-a^2 + 1}b^2x^2 \arctan\left(-\frac{\sqrt{-a^2 + 1}bx - \sqrt{b^2x^2 + 2abx + a^2 - 1}\sqrt{-a^2 + 1}}{a^2 - 1}\right) - 2(a^4 - 2a^2 + 1)b^2x^2 \arctan\left(\frac{a^2bx + a^3 + \sqrt{b^2x^2 + 2abx + a^2 - 1}(a^2 + \sqrt{a^2 - 1}a - 1) + (abx + a^2 - 1)\sqrt{a^2 - 1} - a}{x}\right) \right]}{2(a^4 - 2a^2 + 1)b^2x^2}$$

input `integrate(arcsec(b*x+a)/x^3,x, algorithm="fricas")`output

```
[1/2*((2*a^2 - 1)*sqrt(a^2 - 1)*b^2*x^2*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 + sqrt(a^2 - 1)*a - 1) + (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) + 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (a^3 - a)*b^2*x^2 - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x - (a^6 - 2*a^4 + a^2)*arcsec(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2), -1/2*(2*(2*a^2 - 1)*sqrt(-a^2 + 1)*b^2*x^2*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) - 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (a^3 - a)*b^2*x^2 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x + (a^6 - 2*a^4 + a^2)*arcsec(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2)]
```

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asec}(a + bx)}{x^3} dx$$

input `integrate(asec(b*x+a)/x**3,x)`output `Integral(asec(a + b*x)/x**3, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x^3} dx$$

input `integrate(arcsec(b*x+a)/x^3,x, algorithm="maxima")`

output `1/2*(2*x^2*integrate(1/2*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2 + (b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) - arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))/x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. $2(106) = 212$.

Time = 0.16 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.73

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = -\frac{1}{2}b \left(\frac{2(2a^2b - b) \arctan\left(\frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + a}{\sqrt{-a^2 + 1}}\right)}{(a^4 - a^2)\sqrt{-a^2 + 1}} + \frac{2\left((bx+a)ab\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + 2(bx+a)\right)}{\left((bx+a)^2\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + 2(bx+a)\right)^2} \right)$$

input `integrate(arcsec(b*x+a)/x^3,x, algorithm="giac")`

output `-1/2*b*(2*(2*a^2*b - b)*arctan(((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + a)/sqrt(-a^2 + 1))/((a^4 - a^2)*sqrt(-a^2 + 1)) + 2*((b*x + a)*a*b*(sqrt(-1/(b*x + a)^2 + 1) - 1) + b)/(((b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 2*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)*(a^3 - a)) + (2*a*b/(b*x + a) - b)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/(a^2*(a/(b*x + a) - 1)^2))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x^3} dx$$

input `int(acos(1/(a + b*x))/x^3,x)`output `int(acos(1/(a + b*x))/x^3, x)`**Reduce [F]**

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = \int \frac{\operatorname{asec}(bx + a)}{x^3} dx$$

input `int(asec(b*x+a)/x^3,x)`output `int(asec(a + b*x)/x**3,x)`

3.26 $\int \frac{\sec^{-1}(a+bx)}{x^4} dx$

Optimal result	210
Mathematica [C] (verified)	211
Rubi [A] (verified)	212
Maple [B] (verified)	216
Fricas [A] (verification not implemented)	217
Sympy [F]	217
Maxima [F]	218
Giac [B] (verification not implemented)	218
Mupad [F(-1)]	219
Reduce [F]	219

Optimal result

Integrand size = 10, antiderivative size = 181

$$\int \frac{\sec^{-1}(a+bx)}{x^4} dx = \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\sec^{-1}(a+bx)}{3a^3} - \frac{\sec^{-1}(a+bx)}{3x^3} + \frac{(2-5a^2+6a^4)b^3\arctan\left(\frac{\sqrt{1+a}\tan(\frac{1}{2}\sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}$$

output

```
1/6*b*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a/(-a^2+1)/x^2-1/6*(-5*a^2+2)*b^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a^2/(-a^2+1)^2/x-1/3*b^3*arcsec(b*x+a)/a^3-1/3*arcsec(b*x+a)/x^3+1/3*(6*a^4-5*a^2+2)*b^3*arctan((1+a)^(1/2)*tan(1/2*arcsec(b*x+a))/(1-a)^(1/2))/a^3/(-a^2+1)^(5/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.33

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx$$

$$= \frac{1}{6} \left(-\frac{b \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} (a^4 + abx - 4a^3bx + 2b^2x^2 - a^2(1 + 5b^2x^2))}{a^2(-1+a^2)^2 x^2} - \frac{2 \sec^{-1}(a + bx)}{x^3} + \frac{2b^3 \arcsin\left(\frac{1}{a+bx}\right)}{a^3} - \frac{i(2 - 5a^2 + 6a^4) b^3 \log\left(\frac{12a^3(-1+a^2)^2 \left(\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}} + (a+bx) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{(2-5a^2+6a^4)b^3x}\right)}{a^3(1-a^2)^{5/2}} \right)$$

input `Integrate[ArcSec[a + b*x]/x^4,x]`

output `(-((b*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^4 + a*b*x - 4*a^3*b*x + 2*b^2*x^2 - a^2*(1 + 5*b^2*x^2)))/(a^2*(-1 + a^2)^2*x^2)) - (2*ArcSec[a + b*x])/x^3 + (2*b^3*ArcSin[(a + b*x)^(-1)])/a^3 - (I*(2 - 5*a^2 + 6*a^4)*b^3*Log[(12*a^3*(-1 + a^2)^2*((I*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] + (a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]])]/((2 - 5*a^2 + 6*a^4)*b^3*x)))/(a^3*(1 - a^2)^(5/2))/6`

Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.23, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$, Rules used = {5781, 4926, 3042, 4272, 3042, 4548, 3042, 4407, 3042, 4318, 3042, 3138, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(a+bx)}{x^4} dx \\
 & \quad \downarrow \text{5781} \\
 & b^3 \int \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{b^4 x^4} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{4926} \\
 & b^3 \left(-\frac{1}{3} \int -\frac{1}{b^3 x^3} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left(-\frac{1}{3} \int \frac{1}{\left(a - \csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^3} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{4272} \\
 & b^3 \left(\frac{1}{3} \left(\frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2 x^2} - \frac{\int \frac{-(a+bx)^2 - 2a(a+bx) + 2(1-a^2)}{b^2 x^2} d \sec^{-1}(a+bx)}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{3042} \\
 & b^3 \left(\frac{1}{3} \left(\frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2 x^2} - \frac{\int \frac{-\csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)^2 - 2a \csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right) + 2(1-a^2)}{\left(a - \csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^2} d \sec^{-1}(a+bx)}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3 x^3} \right) \\
 & \quad \downarrow \text{4548}
 \end{aligned}$$

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\int -\frac{2(1-a^2)^2 - a(1-4a^2)(a+bx)}{bx} d\sec^{-1}(a+bx) + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\int \frac{2(1-a^2)^2 - a(1-4a^2)\csc(\sec^{-1}(a+bx)+\frac{\pi}{2})}{a-\csc(\sec^{-1}(a+bx)+\frac{\pi}{2})} d\sec^{-1}(a+bx) + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 4407

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\frac{(6a^4-5a^2+2)\int -\frac{a+bx}{bx} d\sec^{-1}(a+bx) + \frac{2(1-a^2)^2\sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\frac{(6a^4-5a^2+2)\int \frac{\csc(\sec^{-1}(a+bx)+\frac{\pi}{2})}{a-\csc(\sec^{-1}(a+bx)+\frac{\pi}{2})} d\sec^{-1}(a+bx) + \frac{2(1-a^2)^2\sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 4318

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\frac{2(1-a^2)^2\sec^{-1}(a+bx)}{a} - \frac{(6a^4-5a^2+2)\int \frac{1}{1-\frac{1}{a+bx}} d\sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3042

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\frac{2(1-a^2)^2\sec^{-1}(a+bx)}{a} - \frac{(6a^4-5a^2+2)\int \frac{1}{1-a\sin(\sec^{-1}(a+bx)+\frac{\pi}{2})} d\sec^{-1}(a+bx)}{a}}{a(1-a^2)} + \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx}}{2a(1-a^2)} \right) - \frac{\sec^{-1}(a+bx)}{3b^3x^3} \right)$$

↓ 3138

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{\frac{2(1-a^2)^2 \sec^{-1}(a+bx)}{a} - \frac{2(6a^4-5a^2+2) \int \frac{1}{(a+1)\tan^2(\frac{1}{2}\sec^{-1}(a+bx))-a+1} d \tan(\frac{1}{2}\sec^{-1}(a+bx))}{a}}{2a(1-a^2)} + \dots \right) \right)$$

↓ 218

$$b^3 \left(\frac{1}{3} \left(\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)b^2x^2} - \frac{(2-5a^2)\sqrt{1-\frac{1}{(a+bx)^2}}(a+bx)}{a(1-a^2)bx} + \frac{\frac{2(1-a^2)^2 \sec^{-1}(a+bx)}{a} - \frac{2(6a^4-5a^2+2) \arctan\left(\frac{\sqrt{a+1}\tan(\frac{1}{2}\sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}}{2a(1-a^2)} \right) \right)$$

input

```
Int[ArcSec[a + b*x]/x^4,x]
```

output

```
b^3*(-1/3*ArcSec[a + b*x]/(b^3*x^3) + (((a + b*x)*Sqrt[1 - (a + b*x)^(-2)]
)/(2*a*(1 - a^2)*b^2*x^2) - (((2 - 5*a^2)*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]
))/((a*(1 - a^2)*b*x) + ((2*(1 - a^2)^2*ArcSec[a + b*x])/a - (2*(2 - 5*a^2
+ 6*a^4)*ArcTan[(Sqrt[1 + a]*Tan[ArcSec[a + b*x]/2])/Sqrt[1 - a]]/(a*Sqr
t[1 - a^2]))/(a*(1 - a^2)))/(2*a*(1 - a^2)))/3
```

Defintions of rubi rules used

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

- rule 3138 $\text{Int}[(a + (b \sin[\pi/2 + (c + d)x])^{-1}), x_{\text{Symbol}}] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + dx)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + b + (a - b)*e^{2*x^2}), x], x, \text{Tan}[(c + dx)/2]/e], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$
- rule 4272 $\text{Int}[(\csc[(c + dx)]*(b + a))^{(n)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[b^2*\text{Cot}[c + dx]*((a + b*\csc[c + dx])^{(n+1)}/(a*d*(n+1)*(a^2 - b^2))), x] + \text{Simp}[1/(a*(n+1)*(a^2 - b^2)) \text{ Int}[(a + b*\csc[c + dx])^{(n+1)}*\text{Simp}[(a^2 - b^2)*(n+1) - a*b*(n+1)*\csc[c + dx] + b^2*(n+2)*\csc[c + dx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$
- rule 4318 $\text{Int}[\csc[(e + f)x]/(\csc[(e + f)x]*(b + a))], x_{\text{Symbol}}] \rightarrow \text{Simp}[1/b \text{ Int}[1/(1 + (a/b)*\sin[e + fx]), x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$
- rule 4407 $\text{Int}[(\csc[(e + f)x]*(d + c))/(\csc[(e + f)x]*(b + a))], x_{\text{Symbol}}] \rightarrow \text{Simp}[c*(x/a), x] - \text{Simp}[(b*c - a*d)/a \text{ Int}[\csc[e + fx]/(a + b*\csc[e + fx]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$
- rule 4548 $\text{Int}[(A + \csc[(e + f)x]*(B + C))^{(m)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(A*b^2 - a*b*B + a^2*C)*\text{Cot}[e + fx]*((a + b*\csc[e + fx])^{(m+1)}/(a*f*(m+1)*(a^2 - b^2))), x] + \text{Simp}[1/(a*(m+1)*(a^2 - b^2)) \text{ Int}[(a + b*\csc[e + fx])^{(m+1)}*\text{Simp}[A*(a^2 - b^2)*(m+1) - a*(A*b - a*B + b*C)*(m+1)*\csc[e + fx] + (A*b^2 - a*b*B + a^2*C)*(m+2)*\csc[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1]$
- rule 4926 $\text{Int}[(e + f*x)^{(m)}*\text{Sec}[(c + d)x]*((a + b*\text{Sec}[(c + d)x])^{(n)}*\text{Tan}[(c + d)x]), x_{\text{Symbol}}] \rightarrow \text{Simp}[(e + f*x)^m*((a + b*\text{Sec}[c + dx])^{(n+1)}/(b*d*(n+1))), x] - \text{Simp}[f*(m/(b*d*(n+1))) \text{ Int}[(e + f*x)^{(m-1)}*(a + b*\text{Sec}[c + dx])^{(n+1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{NeQ}[n, -1]$

rule 5781

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d *e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 527 vs. 2(159) = 318.

Time = 0.34 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.92

method	result
parts	$-\frac{\operatorname{arcsec}(bx+a)}{3x^3} - \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left(-2(a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) a^4b^2x^2 + 6 \ln\left(\frac{2a^2-2+2abx+2\sqrt{b^2x^2+2abx+a^2-1}}{a^2-1}\right) \right)}{3x^3}$
derivatividivides	Expression too large to display
default	Expression too large to display

input

```
int(arcsec(b*x+a)/x^4,x,method=_RETURNVERBOSE)
```

output

```
-1/3*arcsec(b*x+a)/x^3-1/6*b*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(-2*(a^2-1)^(3/2)*arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*a^4*b^2*x^2+6*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*a^6*b^2*x^2+4*(a^2-1)^(3/2)*arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*a^2*b^2*x^2-5*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(3/2)*a^3*b*x-11*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*a^4*b^2*x^2+(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(3/2)*a^4-2*b^2*arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*x^2*(a^2-1)^(3/2)+2*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(3/2)*a*b*x+7*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*a^2*b^2*x^2-(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(3/2)*a^2-2*b^2*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*x^2)/((b^2*x^2+2*a*b*x+a^2-1)/(b*x+a)^2)^(1/2)/(b*x+a)/a^3/(a^2-1)^(7/2)/x^2
```

Fricas [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 548, normalized size of antiderivative = 3.03

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx$$

$$= \left[\frac{(6a^4 - 5a^2 + 2)\sqrt{a^2 - 1}b^3x^3 \log\left(\frac{a^2bx + a^3 + \sqrt{b^2x^2 + 2abx + a^2 - 1}(a^2 - \sqrt{a^2 - 1}a - 1) - (abx + a^2 - 1)\sqrt{a^2 - 1} - a}{x}\right) - 4(a^6 - 3a^4 + 3a^2 - 1)b^3x^3 \arctan\left(\frac{-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}}{\sqrt{a^2 - 1}}\right) + (5a^5 - 7a^3 + 2a)b^3x^3 - 2(a^9 - 3a^7 + 3a^5 - a^3)\operatorname{arcsec}(bx + a) + ((5a^5 - 7a^3 + 2a)b^2x^2 - (a^6 - 2a^4 + a^2)bx)\sqrt{b^2x^2 + 2abx + a^2 - 1}}{(a^9 - 3a^7 + 3a^5 - a^3)x^3}, \frac{1}{6}(2(6a^4 - 5a^2 + 2)\sqrt{-a^2 + 1}b^3x^3 \arctan\left(\frac{-\sqrt{-a^2 + 1}bx - \sqrt{b^2x^2 + 2abx + a^2 - 1}\sqrt{-a^2 + 1}}{a^2 - 1}\right) - 4(a^6 - 3a^4 + 3a^2 - 1)b^3x^3 \arctan\left(\frac{-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}}{\sqrt{a^2 - 1}}\right) + (5a^5 - 7a^3 + 2a)b^3x^3 - 2(a^9 - 3a^7 + 3a^5 - a^3)\operatorname{arcsec}(bx + a) + ((5a^5 - 7a^3 + 2a)b^2x^2 - (a^6 - 2a^4 + a^2)bx)\sqrt{b^2x^2 + 2abx + a^2 - 1}}{(a^9 - 3a^7 + 3a^5 - a^3)x^3} \right]$$

input `integrate(arcsec(b*x+a)/x^4,x, algorithm="fricas")`output

```
[1/6*((6*a^4 - 5*a^2 + 2)*sqrt(a^2 - 1)*b^3*x^3*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 - sqrt(a^2 - 1)*a - 1) - (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) - 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arcsec(b*x + a) + ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3), 1/6*(2*(6*a^4 - 5*a^2 + 2)*sqrt(-a^2 + 1)*b^3*x^3*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))*sqrt(-a^2 + 1))/(a^2 - 1)) - 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arcsec(b*x + a) + ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3)]
```

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asec}(a + bx)}{x^4} dx$$

input `integrate(asec(b*x+a)/x**4,x)`output `Integral(asec(a + b*x)/x**4, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x^4} dx$$

input `integrate(arcsec(b*x+a)/x^4,x, algorithm="maxima")`

output `1/3*(3*x^3*integrate(1/3*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3 + (b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) - arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))/x^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 451 vs. $2(155) = 310$.

Time = 0.19 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.49

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \frac{1}{3} b \left(\frac{(6a^4b^2 - 5a^2b^2 + 2b^2) \arctan\left(\frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + a}{\sqrt{-a^2 + 1}}\right)}{(a^7 - 2a^5 + a^3)\sqrt{-a^2 + 1}} + \frac{4(bx+a)^3 a^3 b^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)}{\dots} \right)$$

input `integrate(arcsec(b*x+a)/x^4,x, algorithm="giac")`

output

```
1/3*b*((6*a^4*b^2 - 5*a^2*b^2 + 2*b^2)*arctan(((b*x + a)*(sqrt(-1/(b*x + a)
)^2 + 1) - 1) + a)/sqrt(-a^2 + 1))/((a^7 - 2*a^5 + a^3)*sqrt(-a^2 + 1)) +
(4*(b*x + a)^3*a^3*b^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 10*(b*x + a)^2*a
^4*b^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 - (b*x + a)^3*a*b^2*(sqrt(-1/(b*x
+ a)^2 + 1) - 1)^3 + (b*x + a)^2*a^2*b^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2
+ 16*(b*x + a)*a^3*b^2*(sqrt(-1/(b*x + a)^2 + 1) - 1) - 2*(b*x + a)^2*b^2*
(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 - 7*(b*x + a)*a*b^2*(sqrt(-1/(b*x + a)^2
+ 1) - 1) + 5*a^2*b^2 - 2*b^2)/((a^6 - 2*a^4 + a^2)*((b*x + a)^2*(sqrt(-1/
(b*x + a)^2 + 1) - 1)^2 + 2*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1
)^2) - (3*a*b^2/(b*x + a) - 3*a^2*b^2/(b*x + a)^2 - b^2)*arccos(-1/((b*x +
a)*(a/(b*x + a) - 1) - a))/(a^3*(a/(b*x + a) - 1)^3))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x^4} dx$$

input

```
int(acos(1/(a + b*x))/x^4,x)
```

output

```
int(acos(1/(a + b*x))/x^4, x)
```

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asec}(bx + a)}{x^4} dx$$

input

```
int(asec(b*x+a)/x^4,x)
```

output

```
int(asec(a + b*x)/x**4,x)
```

3.27 $\int x^3 \sec^{-1}(a + bx)^2 dx$

Optimal result	221
Mathematica [A] (warning: unable to verify)	222
Rubi [A] (verified)	222
Maple [A] (verified)	225
Fricas [F]	226
Sympy [F]	226
Maxima [F]	226
Giac [F(-2)]	227
Mupad [F(-1)]	227
Reduce [F]	227

Optimal result

Integrand size = 12, antiderivative size = 381

$$\begin{aligned}
 \int x^3 \sec^{-1}(a+bx)^2 dx = & -\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} - \frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{3b^4} \\
 & - \frac{3a^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{b^4} \\
 & + \frac{a(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{b^4} \\
 & - \frac{(a+bx)^3\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{6b^4} - \frac{a^4 \sec^{-1}(a+bx)^2}{4b^4} \\
 & + \frac{1}{4}x^4 \sec^{-1}(a+bx)^2 - \frac{2ia \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{4ia^3 \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} + \frac{\log(a+bx)}{3b^4} \\
 & + \frac{3a^2 \log(a+bx)}{b^4} + \frac{ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & + \frac{2ia^3 \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{ia \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{2ia^3 \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4}
 \end{aligned}$$

output

```

-a*x/b^3+1/12*(b*x+a)^2/b^4-1/3*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a
)/b^4-3*a^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a)/b^4+a*(b*x+a)^2*(1
-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a)/b^4-1/6*(b*x+a)^3*(1-1/(b*x+a)^2)^(1/2)*
arcsec(b*x+a)/b^4-1/4*a^4*arcsec(b*x+a)^2/b^4+1/4*x^4*arcsec(b*x+a)^2-2*I*
a*arcsec(b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^4+I*a*polylog(
2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^4+1/3*ln(b*x+a)/b^4+3*a^2*ln(b
*x+a)/b^4+2*I*a^3*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^4-I*
a*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^4-2*I*a^3*polylog(2,I
*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^4-4*I*a^3*arcsec(b*x+a)*arctan(1/(
b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^4

```

Mathematica [A] (warning: unable to verify)

Time = 8.17 (sec) , antiderivative size = 667, normalized size of antiderivative = 1.75

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \text{Too large to display}$$

input `Integrate[x^3*ArcSec[a + b*x]^2,x]`

output

```
((1 - a/(a + b*x))^3*(24*a*(2 + (1 + 2*a^2)*ArcSec[a + b*x]^2) + (2 + (-2
+ 24*a)*ArcSec[a + b*x] + 3*(1 - 4*a + 12*a^2)*ArcSec[a + b*x]^2)/(-1 + Sq
rt[1 - (a + b*x)^(-2)]) + 16*(1 + 9*a^2)*Log[(a + b*x)^(-1)] - 24*a*(1 + 2
*a^2)*((Pi - 2*ArcSec[a + b*x])*(Log[1 - I/E^(I*ArcSec[a + b*x])] - Log[1
+ I/E^(I*ArcSec[a + b*x])]) - Pi*Log[Cot[(Pi + 2*ArcSec[a + b*x])/4]] + (2
*I)*(PolyLog[2, (-I)/E^(I*ArcSec[a + b*x])] - PolyLog[2, I/E^(I*ArcSec[a +
b*x])])) - (3*ArcSec[a + b*x]^2)/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a +
b*x]/2])^4 + (4*ArcSec[a + b*x]*(1 + 6*a*ArcSec[a + b*x])*Sin[ArcSec[a +
b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2])^3 + (8*(2*ArcSe
c[a + b*x] + 18*a^2*ArcSec[a + b*x] + 6*a^3*ArcSec[a + b*x]^2 + 3*a*(2 + A
rcSec[a + b*x]^2))*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[A
rcSec[a + b*x]/2]) - (3*ArcSec[a + b*x]^2)/(Cos[ArcSec[a + b*x]/2] + Sin[A
rcSec[a + b*x]/2])^4 + (4*ArcSec[a + b*x]*(1 - 6*a*ArcSec[a + b*x])*Sin[Arc
Sec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^3 - (2
+ (2 - 24*a)*ArcSec[a + b*x] + 3*(1 - 4*a + 12*a^2)*ArcSec[a + b*x]^2)/(C
os[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^2 - (8*(-2*ArcSec[a + b*x]
- 18*a^2*ArcSec[a + b*x] + 6*a^3*ArcSec[a + b*x]^2 + 3*a*(2 + ArcSec[a +
b*x]^2))*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a +
b*x]/2])))/(48*b^4*(-1 + a/(a + b*x))^3)
```

Rubi [A] (verified)Time = 0.64 (sec) , antiderivative size = 357, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5781, 25, 4926, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x^3 \sec^{-1}(a + bx)^2 dx \\
& \quad \downarrow \text{5781} \\
& \frac{\int b^3 x^3 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b^4} \\
& \quad \downarrow \text{25} \\
& \frac{\int -b^3 x^3 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b^4} \\
& \quad \downarrow \text{4926} \\
& \frac{\frac{1}{4} b^4 x^4 \sec^{-1}(a + bx)^2 - \frac{1}{2} \int b^4 x^4 \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b^4} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{1}{4} b^4 x^4 \sec^{-1}(a + bx)^2 - \frac{1}{2} \int \sec^{-1}(a + bx) \left(a - \csc \left(\sec^{-1}(a + bx) + \frac{\pi}{2} \right) \right)^4 d \sec^{-1}(a + bx)}{b^4} \\
& \quad \downarrow \text{4678} \\
& \frac{\frac{1}{4} b^4 x^4 \sec^{-1}(a + bx)^2 - \frac{1}{2} \int \left(\sec^{-1}(a + bx) a^4 - 4(a + bx) \sec^{-1}(a + bx) a^3 + 6(a + bx)^2 \sec^{-1}(a + bx) a^2 - 4(a + bx)^3 \sec^{-1}(a + bx) a + 4(a + bx)^4 \right) d \sec^{-1}(a + bx)}{b^4} \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{1}{4} b^4 x^4 \sec^{-1}(a + bx)^2 + \frac{1}{2} \left(-\frac{1}{2} a^4 \sec^{-1}(a + bx)^2 - 8ia^3 \sec^{-1}(a + bx) \arctan \left(e^{i \sec^{-1}(a+bx)} \right) + 4ia^3 \text{PolyLog} \left(2, e^{i \sec^{-1}(a+bx)} \right) \right)}{b^4}
\end{aligned}$$

input

Int [x^3*ArcSec[a + b*x]^2,x]

output

$$\begin{aligned} & ((b^4 x^4 \operatorname{ArcSec}[a + b x]^2)/4 + (-2 a (a + b x) + (a + b x)^2/6 - (2 (a + b x) \sqrt{1 - (a + b x)^{-2}}) \operatorname{ArcSec}[a + b x])/3 - 6 a^2 (a + b x) \sqrt{1 - (a + b x)^{-2}} \operatorname{ArcSec}[a + b x] + 2 a (a + b x)^2 \sqrt{1 - (a + b x)^{-2}} \operatorname{ArcSec}[a + b x] - ((a + b x)^3 \sqrt{1 - (a + b x)^{-2}} \operatorname{ArcSec}[a + b x])/3 - (a^4 \operatorname{ArcSec}[a + b x]^2)/2 - (4 I) a \operatorname{ArcSec}[a + b x] \operatorname{ArcTan}[E^{(I \operatorname{ArcSec}[a + b x])}] - (8 I) a^3 \operatorname{ArcSec}[a + b x] \operatorname{ArcTan}[E^{(I \operatorname{ArcSec}[a + b x])}] - (2 \operatorname{Log}[(a + b x)^{-1}])/3 - 6 a^2 \operatorname{Log}[(a + b x)^{-1}] + (2 I) a \operatorname{PolyLog}[2, (-I) E^{(I \operatorname{ArcSec}[a + b x])}] + (4 I) a^3 \operatorname{PolyLog}[2, (-I) E^{(I \operatorname{ArcSec}[a + b x])}] - (2 I) a \operatorname{PolyLog}[2, I E^{(I \operatorname{ArcSec}[a + b x])}] - (4 I) a^3 \operatorname{PolyLog}[2, I E^{(I \operatorname{ArcSec}[a + b x])}]) / (2 b^4) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$$

rule 2009

$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 3042

$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4678

$$\operatorname{Int}[(\operatorname{csc}[e] + (f)(x))(b) + (a)^{(n)}((c) + (d)(x))^{(m)}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(c + d x)^m, (a + b \operatorname{Csc}[e + f x])^n, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, m\}, x \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{IGtQ}[n, 0]$$

rule 4926

$$\operatorname{Int}[(e + (f)(x))^{(m)} \operatorname{Sec}[c + (d)(x)]((a) + (b) \operatorname{Sec}[c + (d)(x)])^{(n)} \operatorname{Tan}[c + (d)(x)], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(e + f x)^m ((a + b \operatorname{Sec}[c + d x])^{(n+1)} / (b d (n+1))), x] - \operatorname{Simp}[f (m / (b d (n+1))) \operatorname{Int}[(e + f x)^{(m-1)} (a + b \operatorname{Sec}[c + d x])^{(n+1)}, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, n\}, x \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{NeQ}[n, -1]$$

rule 5781

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^p_)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d
*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e
, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.85 (sec) , antiderivative size = 673, normalized size of antiderivative = 1.77

method	result
derivativedivides	$-\operatorname{arcsec}(bx+a)^2 a^3 (bx+a) + \frac{3 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a)^2}{2} - \operatorname{arcsec}(bx+a)^2 a (bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^4}{4} - 3 \operatorname{arcsec}(bx+a)$
default	$-\operatorname{arcsec}(bx+a)^2 a^3 (bx+a) + \frac{3 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a)^2}{2} - \operatorname{arcsec}(bx+a)^2 a (bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^4}{4} - 3 \operatorname{arcsec}(bx+a)$

input

```
int(x^3*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/b^4*(-arcsec(b*x+a)^2*a^3*(b*x+a)+3/2*arcsec(b*x+a)^2*a^2*(b*x+a)^2-arcsec(b*x+a)^2*a*(b*x+a)^3+1/4*arcsec(b*x+a)^2*(b*x+a)^4-3*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*a^2*(b*x+a)+arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*a*(b*x+a)^2-1/6*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)^3+I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a-1/3*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)-3*I*a^2*arcsec(b*x+a)-(b*x+a)*a+1/12*(b*x+a)^2+2/3*ln(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))-1/3*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+6*ln(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))*a^2-3*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)*a^2-2*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a^3*arcsec(b*x+a)+2*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a^3*arcsec(b*x+a)+2*I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a^3-I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a-ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a*arcsec(b*x+a)+ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a*arcsec(b*x+a)-1/3*I*arcsec(b*x+a)-2*I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))) *a^3)
```

Fricas [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x^3*arcsec(b*x+a)^2,x, algorithm="fricas")`

output `integral(x^3*arcsec(b*x + a)^2, x)`

Sympy [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{asec}^2(a + bx) dx$$

input `integrate(x**3*asec(b*x+a)**2,x)`

output `Integral(x**3*asec(a + b*x)**2, x)`

Maxima [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x^3*arcsec(b*x+a)^2,x, algorithm="maxima")`

output `1/4*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/16*x^4*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/4*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 4*(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2 - 1)*b*x^4 + (a^3 - a)*x^3)*log(b*x + a)^2 - (b^3*x^6 + 2*a*b^2*x^5 + (a^2 - 1)*b*x^4 + 4*(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2 - 1)*b*x^4 + (a^3 - a)*x^3)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F(-2)]

Exception generated.

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^3*arcsec(b*x+a)^2,x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vector & l) Error: Bad Argument Value`

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \arccos\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(x^3*acos(1/(a + b*x))^2,x)`

output `int(x^3*acos(1/(a + b*x))^2, x)`

Reduce [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int \operatorname{asec}(bx + a)^2 x^3 dx$$

input `int(x^3*asec(b*x+a)^2,x)`

output `int(asec(a + b*x)**2*x**3,x)`

3.28 $\int x^2 \sec^{-1}(a + bx)^2 dx$

Optimal result	228
Mathematica [A] (warning: unable to verify)	229
Rubi [A] (verified)	230
Maple [A] (verified)	232
Fricas [F]	232
Sympy [F]	233
Maxima [F]	233
Giac [F]	233
Mupad [F(-1)]	234
Reduce [F]	234

Optimal result

Integrand size = 12, antiderivative size = 288

$$\begin{aligned}
 \int x^2 \sec^{-1}(a + bx)^2 dx = & \frac{x}{3b^2} + \frac{2a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3} \\
 & - \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{3b^3} + \frac{a^3 \sec^{-1}(a + bx)^2}{3b^3} \\
 & + \frac{1}{3} x^3 \sec^{-1}(a + bx)^2 + \frac{2i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{3b^3} \\
 & + \frac{4ia^2 \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{2a \log(a + bx)}{b^3} - \frac{i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{3b^3} \\
 & - \frac{2ia^2 \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{3b^3} \\
 & + \frac{2ia^2 \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3}
 \end{aligned}$$

output

$$\frac{1}{3}x/b^2 + 2a*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}*\operatorname{arcsec}(b*x+a)/b^3 - 1/3*(b*x+a)^2*(1-1/(b*x+a)^2)^{(1/2)}*\operatorname{arcsec}(b*x+a)/b^3 + 1/3*a^3*\operatorname{arcsec}(b*x+a)^2/b^3 + 1/3*x^3*\operatorname{arcsec}(b*x+a)^2 + 2/3*I*\operatorname{arcsec}(b*x+a)*\arctan(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})/b^3 + 4*I*a^2*\operatorname{arcsec}(b*x+a)*\arctan(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})/b^3 - 2*a*\ln(b*x+a)/b^3 - 1/3*I*\operatorname{polylog}(2, -I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3 - 2*I*a^2*\operatorname{polylog}(2, -I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3 + 1/3*I*\operatorname{polylog}(2, I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3 + 2*I*a^2*\operatorname{polylog}(2, I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3$$
Mathematica [A] (warning: unable to verify)

Time = 4.01 (sec) , antiderivative size = 473, normalized size of antiderivative = 1.64

$$\int x^2 \sec^{-1}(a + bx)^2 dx$$

$$4 + 2(1 + 6a^2) \sec^{-1}(a + bx)^2 + \frac{\sec^{-1}(a+bx)(2+(-1+6a)\sec^{-1}(a+bx))}{-1+\sqrt{1-\frac{1}{(a+bx)^2}}} + 24a \log\left(\frac{1}{a+bx}\right) + 2(-1 - 6a^2) \left((\pi - 2) \right)$$

input

Integrate[x^2*ArcSec[a + b*x]^2,x]

output

$$(4 + 2*(1 + 6*a^2)*\operatorname{ArcSec}[a + b*x]^2 + (\operatorname{ArcSec}[a + b*x]*(2 + (-1 + 6*a)*\operatorname{ArcSec}[a + b*x]))/(-1 + \operatorname{Sqrt}[1 - (a + b*x)^{-2}])) + 24*a*\operatorname{Log}[(a + b*x)^{-1}] + 2*(-1 - 6*a^2)*((\pi - 2*\operatorname{ArcSec}[a + b*x])*(\operatorname{Log}[1 - I/E^{(I*\operatorname{ArcSec}[a + b*x])}] - \operatorname{Log}[1 + I/E^{(I*\operatorname{ArcSec}[a + b*x])}]) - \pi*\operatorname{Log}[\operatorname{Cot}[(\pi + 2*\operatorname{ArcSec}[a + b*x])/4]]) + (2*I)*(PolyLog[2, (-I)/E^{(I*\operatorname{ArcSec}[a + b*x])}] - PolyLog[2, I/E^{(I*\operatorname{ArcSec}[a + b*x])}])) + (2*\operatorname{ArcSec}[a + b*x]^2*\operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])/(\operatorname{Cos}[\operatorname{ArcSec}[a + b*x]/2] - \operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])^3 + (2*(2 + 12*a*\operatorname{ArcSec}[a + b*x] + (1 + 6*a^2)*\operatorname{ArcSec}[a + b*x]^2)*\operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])/(\operatorname{Cos}[\operatorname{ArcSec}[a + b*x]/2] - \operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2]) - (2*\operatorname{ArcSec}[a + b*x]^2*\operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])/(\operatorname{Cos}[\operatorname{ArcSec}[a + b*x]/2] + \operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])^3 + (\operatorname{ArcSec}[a + b*x]*(2 + (1 - 6*a)*\operatorname{ArcSec}[a + b*x]))/(\operatorname{Cos}[\operatorname{ArcSec}[a + b*x]/2] + \operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])^2 - (2*(2 - 12*a*\operatorname{ArcSec}[a + b*x] + (1 + 6*a^2)*\operatorname{ArcSec}[a + b*x]^2)*\operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2])/(\operatorname{Cos}[\operatorname{ArcSec}[a + b*x]/2] + \operatorname{Sin}[\operatorname{ArcSec}[a + b*x]/2]))/(12*b^3)$$

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 271, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5781, 4926, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sec^{-1}(a+bx)^2 dx \\
 & \quad \downarrow \text{5781} \\
 & \frac{\int b^2 x^2 (a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2 d \sec^{-1}(a+bx)}{b^3} \\
 & \quad \downarrow \text{4926} \\
 & \frac{\frac{2}{3} \int -b^3 x^3 \sec^{-1}(a+bx) d \sec^{-1}(a+bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a+bx)^2}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2}{3} \int \sec^{-1}(a+bx) \left(a - \csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^3 d \sec^{-1}(a+bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a+bx)^2}{b^3} \\
 & \quad \downarrow \text{4678} \\
 & \frac{\frac{2}{3} \int (\sec^{-1}(a+bx)a^3 - 3(a+bx) \sec^{-1}(a+bx)a^2 + 3(a+bx)^2 \sec^{-1}(a+bx)a - (a+bx)^3 \sec^{-1}(a+bx)) d \sec^{-1}(a+bx)}{b^3} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{1}{3} b^3 x^3 \sec^{-1}(a+bx)^2 + \frac{2}{3} \left(\frac{1}{2} a^3 \sec^{-1}(a+bx)^2 + 6ia^2 \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right) - 3ia^2 \text{PolyLog}\left(2, -i\right)\right)}{b^3}
 \end{aligned}$$

input `Int [x^2*ArcSec[a + b*x]^2,x]`

output

$$\begin{aligned} & ((b^3x^3\text{ArcSec}[a + bx]^2)/3 + (2*((a + bx)/2 + 3*a*(a + bx)*\text{Sqrt}[1 - \\ & (a + bx)^{-2}])\text{ArcSec}[a + bx] - ((a + bx)^2*\text{Sqrt}[1 - (a + bx)^{-2}])\text{Ar} \\ & \text{cSec}[a + bx])/2 + (a^3\text{ArcSec}[a + bx]^2)/2 + I*\text{ArcSec}[a + bx]*\text{ArcTan}[E^ \\ & (I*\text{ArcSec}[a + bx])] + (6*I)*a^2*\text{ArcSec}[a + bx]*\text{ArcTan}[E^(I*\text{ArcSec}[a + b* \\ & x])] + 3*a*\text{Log}[(a + bx)^{-1}] - (I/2)*\text{PolyLog}[2, (-I)*E^(I*\text{ArcSec}[a + b*x \\ &])] - (3*I)*a^2*\text{PolyLog}[2, (-I)*E^(I*\text{ArcSec}[a + b*x])] + (I/2)*\text{PolyLog}[2, \\ & I*E^(I*\text{ArcSec}[a + b*x])] + (3*I)*a^2*\text{PolyLog}[2, I*E^(I*\text{ArcSec}[a + b*x])]) \\ & /3)/b^3 \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$$

rule 4678

$$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^{(n_.)}*((c_.) + (d_.)*(x_.))^{(m_.)} \\ , x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (a + b*\text{Csc}[e + f*x])^n, x], \\ x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, f, m\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0]$$

rule 4926

$$\text{Int}(((e_.) + (f_.)*(x_.))^{(m_.)}*\text{Sec}[(c_.) + (d_.)*(x_.)]*((a_.) + (b_.)*\text{Sec}[(c \\ _.) + (d_.)*(x_.)])^{(n_.)}*\text{Tan}[(c_.) + (d_.)*(x_.)], x_Symbol] \text{ :> } \text{Simp}[(e + f* \\ x)^m*((a + b*\text{Sec}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] - \text{Simp}[f*(m/(b*d*(n + \\ 1))) \ \text{Int}[(e + f*x)^{(m - 1)}*(a + b*\text{Sec}[c + d*x])^{(n + 1)}, x], x] \text{ /; } \text{FreeQ} \\ \{a, b, c, d, e, f, n\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$$

rule 5781

$$\text{Int}(((a_.) + \text{ArcSec}[(c_.) + (d_.)*(x_.)]*(b_.))^{(p_.)}*((e_.) + (f_.)*(x_.))^{(m \\ _.)}, x_Symbol] \text{ :> } \text{Simp}[1/d^{(m + 1)} \ \text{Subst}[\text{Int}[(a + b*x)^p*\text{Sec}[x]*\text{Tan}[x]*(d \\ *e - c*f + f*\text{Sec}[x])^m, x], x, \text{ArcSec}[c + d*x]], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e \\ , f\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[m]$$

Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 498, normalized size of antiderivative = 1.73

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) - \dots}{\dots}$
default	$\operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) - \dots$

input `int(x^2*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/b^3*(\operatorname{arcsec}(b*x+a)^2*a^2*(b*x+a) - \operatorname{arcsec}(b*x+a)^2*a*(b*x+a)^2 + 1/3*\operatorname{arcsec}(\\ & b*x+a)^2*(b*x+a)^3 + 2*\operatorname{arcsec}(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^{(1/2)}*a*(b*x+ \\ & a) - 1/3*\operatorname{arcsec}(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^{(1/2)}*(b*x+a)^2 + 2*I*\operatorname{dilog}(1 \\ & -I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})))*a^2 + 1/3*b*x + 1/3*a + 1/3*\operatorname{arcsec}(b*x+a) \\ &)*\ln(1+I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})) - 1/3*\operatorname{arcsec}(b*x+a)*\ln(1-I*(1/ \\ & (b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})) + 1/3*I*\operatorname{dilog}(1-I*(1/(b*x+a) + I*(1-1/(b*x+a) \\ &)^2)^{(1/2)})) - 1/3*I*\operatorname{dilog}(1+I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)})) - 4*\ln(1/(\\ & b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))*a + 2*\ln(1+(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2) \\ &))^2)*a - 2*I*\operatorname{dilog}(1+I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2 + 2*I*\operatorname{arcsec}(\\ & b*x+a)*a + 2*\ln(1+I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2*\operatorname{arcsec}(b*x+a) - \\ & 2*\ln(1-I*(1/(b*x+a) + I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2*\operatorname{arcsec}(b*x+a) \end{aligned}$$
Fricas [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x^2*arcsec(b*x+a)^2,x, algorithm="fricas")`

output `integral(x^2*arcsec(b*x + a)^2, x)`

Sympy [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{asec}^2(a + bx) dx$$

input `integrate(x**2*asec(b*x+a)**2,x)`

output `Integral(x**2*asec(a + b*x)**2, x)`

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x^2*arcsec(b*x+a)^2,x, algorithm="maxima")`

output `1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/12*x^3*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/3*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2 - 1)*b*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x^2*arcsec(b*x+a)^2,x, algorithm="giac")`

output `integrate(x^2*arcsec(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(x^2*acos(1/(a + b*x))^2,x)`output `int(x^2*acos(1/(a + b*x))^2, x)`**Reduce [F]**

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int \operatorname{asec}(bx + a)^2 x^2 dx$$

input `int(x^2*asec(b*x+a)^2,x)`output `int(asec(a + b*x)**2*x**2,x)`

3.29 $\int x \sec^{-1}(a + bx)^2 dx$

Optimal result	235
Mathematica [A] (verified)	236
Rubi [A] (verified)	236
Maple [A] (verified)	238
Fricas [F]	239
Sympy [F]	239
Maxima [F]	239
Giac [F]	240
Mupad [F(-1)]	240
Reduce [F]	240

Optimal result

Integrand size = 10, antiderivative size = 154

$$\int x \sec^{-1}(a + bx)^2 dx = -\frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^2} - \frac{a^2 \sec^{-1}(a + bx)^2}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx)^2 - \frac{4ia \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{\log(a + bx)}{b^2} + \frac{2ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{2ia \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2}$$

output

```
- (b*x+a)*(1-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a)/b^2-1/2*a^2*arcsec(b*x+a)^2/b^2+1/2*x^2*arcsec(b*x+a)^2-4*I*a*arcsec(b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^2+ln(b*x+a)/b^2+2*I*a*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2-2*I*a*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.94

$$\int x \sec^{-1}(a + bx)^2 dx$$

$$= \frac{-\left((a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)\right) - a(a + bx) \sec^{-1}(a + bx)^2 + \frac{1}{2}(a + bx)^2 \sec^{-1}(a + bx)^2 - 4ia s}{}$$

input

Integrate[x*ArcSec[a + b*x]^2,x]

output

```
(-((a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]) - a*(a + b*x)*ArcSec[a + b*x]^2 + ((a + b*x)^2*ArcSec[a + b*x]^2)/2 - (4*I)*a*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])] + Log[a + b*x] + (2*I)*a*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - (2*I)*a*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^2
```

Rubi [A] (verified)Time = 0.44 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5781, 25, 4926, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \sec^{-1}(a + bx)^2 dx$$

$$\downarrow \text{5781}$$

$$\frac{\int bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow \text{25}$$

$$\frac{\int -bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b^2}$$

$$\downarrow \text{4926}$$

$$\frac{\frac{1}{2}b^2x^2 \sec^{-1}(a+bx)^2 - \int b^2x^2 \sec^{-1}(a+bx) d \sec^{-1}(a+bx)}{b^2}$$

↓ 3042

$$\frac{\frac{1}{2}b^2x^2 \sec^{-1}(a+bx)^2 - \int \sec^{-1}(a+bx) \left(a - \csc\left(\sec^{-1}(a+bx) + \frac{\pi}{2}\right)\right)^2 d \sec^{-1}(a+bx)}{b^2}$$

↓ 4678

$$\frac{\frac{1}{2}b^2x^2 \sec^{-1}(a+bx)^2 - \int \left(\sec^{-1}(a+bx)a^2 - 2(a+bx) \sec^{-1}(a+bx)a + (a+bx)^2 \sec^{-1}(a+bx)\right) d \sec^{-1}(a+bx)}{b^2}$$

↓ 2009

$$\frac{-\frac{1}{2}a^2 \sec^{-1}(a+bx)^2 - 4ia \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right) + \frac{1}{2}b^2x^2 \sec^{-1}(a+bx)^2 + 2ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2}$$

input `Int[x*ArcSec[a + b*x]^2,x]`

output `((-(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]) - (a^2*ArcSec[a + b*x]^2)/2 + (b^2*x^2*ArcSec[a + b*x]^2)/2 - (4*I)*a*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])] - Log[(a + b*x)^(-1)] + (2*I)*a*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - (2*I)*a*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4678 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 4926 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 5781 `Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.46

method	result
derivativedivides	$\frac{-a \left(\operatorname{arcsec}(bx+a)^2 (bx+a) + 2 \operatorname{arcsec}(bx+a) \ln \left(1 + i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) - 2 \operatorname{arcsec}(bx+a) \ln \left(1 - i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) \right)}{\dots}$
default	$\frac{-a \left(\operatorname{arcsec}(bx+a)^2 (bx+a) + 2 \operatorname{arcsec}(bx+a) \ln \left(1 + i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) - 2 \operatorname{arcsec}(bx+a) \ln \left(1 - i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) \right)}{\dots}$

input `int(x*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/b^2*(-a*(arcsec(b*x+a)^2*(b*x+a)+2*arcsec(b*x+a)*ln(1+I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2)))-2*arcsec(b*x+a)*ln(1-I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2)))-2*I*dilog(1+I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2)))+2*I*dilog(1-I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2)))+1/2*arcsec(b*x+a)^2*(b*x+a)^2-arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)-ln(1/(b*x+a)))`

Fricas [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x*arcsec(b*x+a)^2,x, algorithm="fricas")`

output `integral(x*arcsec(b*x + a)^2, x)`

Sympy [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{asec}^2(a + bx) dx$$

input `integrate(x*asec(b*x+a)**2,x)`

output `Integral(x*asec(a + b*x)**2, x)`

Maxima [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x*arcsec(b*x+a)^2,x, algorithm="maxima")`

output `1/2*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/8*x^2*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/2*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a)^2 - (b^3*x^4 + 2*a*b^2*x^3 + (a^2 - 1)*b*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2)/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(x*arcsec(b*x+a)^2,x, algorithm="giac")`

output `integrate(x*arcsec(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(x*acos(1/(a + b*x))^2,x)`

output `int(x*acos(1/(a + b*x))^2, x)`

Reduce [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int a \sec(bx + a)^2 x dx$$

input `int(x*asec(b*x+a)^2,x)`

output `int(asec(a + b*x)**2*x,x)`

3.30 $\int \sec^{-1}(a + bx)^2 dx$

Optimal result	241
Mathematica [A] (verified)	241
Rubi [A] (verified)	242
Maple [A] (verified)	244
Fricas [F]	245
Sympy [F]	245
Maxima [F]	245
Giac [F]	246
Mupad [F(-1)]	246
Reduce [F]	246

Optimal result

Integrand size = 8, antiderivative size = 94

$$\int \sec^{-1}(a + bx)^2 dx = \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} + \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

output

```
(b*x+a)*arcsec(b*x+a)^2/b+4*I*arcsec(b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b-2*I*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+2*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.18

$$\int \sec^{-1}(a + bx)^2 dx = \frac{\sec^{-1}(a + bx) \left((a + bx) \sec^{-1}(a + bx) - 2 \log\left(1 - ie^{i \sec^{-1}(a + bx)}\right) + 2 \log\left(1 + ie^{i \sec^{-1}(a + bx)}\right) \right) - 2i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right) + 2i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

input

```
Integrate[ArcSec[a + b*x]^2,x]
```

output

$$\frac{(\text{ArcSec}[a + b*x]*((a + b*x)*\text{ArcSec}[a + b*x] - 2*\text{Log}[1 - I*E^(I*\text{ArcSec}[a + b*x])]) + 2*\text{Log}[1 + I*E^(I*\text{ArcSec}[a + b*x])]) - (2*I)*\text{PolyLog}[2, (-I)*E^(I*\text{ArcSec}[a + b*x])]}{b} + (2*I)*\text{PolyLog}[2, I*E^(I*\text{ArcSec}[a + b*x])]/b$$
Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5775, 5739, 4244, 3042, 4669, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{-1}(a + bx)^2 dx$$

$$\downarrow 5775$$

$$\frac{\int \sec^{-1}(a + bx)^2 d(a + bx)}{b}$$

$$\downarrow 5739$$

$$\frac{\int (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b}$$

$$\downarrow 4244$$

$$\frac{(a + bx) \sec^{-1}(a + bx)^2 - 2 \int (a + bx) \sec^{-1}(a + bx) d \sec^{-1}(a + bx)}{b}$$

$$\downarrow 3042$$

$$\frac{(a + bx) \sec^{-1}(a + bx)^2 - 2 \int \sec^{-1}(a + bx) \csc\left(\sec^{-1}(a + bx) + \frac{\pi}{2}\right) d \sec^{-1}(a + bx)}{b}$$

$$\downarrow 4669$$

$$\frac{(a + bx) \sec^{-1}(a + bx)^2 - 2\left(-\int \log\left(1 - ie^{i \sec^{-1}(a+bx)}\right) d \sec^{-1}(a + bx) + \int \log\left(1 + ie^{i \sec^{-1}(a+bx)}\right) d \sec^{-1}(a + bx)\right)}{b}$$

$$\downarrow 2715$$

$$\frac{(a + bx) \sec^{-1}(a + bx)^2 - 2 \left(i \int e^{-i \sec^{-1}(a+bx)} \log \left(1 - i e^{i \sec^{-1}(a+bx)} \right) dx e^{i \sec^{-1}(a+bx)} - i \int e^{-i \sec^{-1}(a+bx)} \log \left(1 + \right. \right.}{b}$$

↓ 2838

$$\frac{(a + bx) \sec^{-1}(a + bx)^2 - 2 \left(-2i \sec^{-1}(a + bx) \arctan \left(e^{i \sec^{-1}(a+bx)} \right) + i \operatorname{PolyLog} \left(2, -i e^{i \sec^{-1}(a+bx)} \right) - i \operatorname{PolyL} \right.}{b}$$

input

```
Int[ArcSec[a + b*x]^2,x]
```

output

```
((a + b*x)*ArcSec[a + b*x]^2 - 2*((-2*I)*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])] + I*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - I*PolyLog[2, I*E^(I*ArcSec[a + b*x])]))/b
```

Defintions of rubi rules used

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4244

```
Int[(x_)^(m_)*Sec[(a_) + (b_)*(x_)^(n_.)]^(p_)*Tan[(a_) + (b_)*(x_)^(n_.)]^(q_), x_Symbol] :> Simp[x^(m - n + 1)*(Sec[a + b*x^n]^p/(b*n*p)), x] - Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sec[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
  := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Simp[
  d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
  x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x)
  )], x], x) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5739

```
Int[((a_.) + ArcSec[(c_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[1/c Subst[
  Int[(a + b*x)^n*Sec[x]*Tan[x], x], x, ArcSec[c*x]], x] /; FreeQ[{a, b, c, n
  }, x] && IGtQ[n, 0]
```

rule 5775

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^p, x_Symbol] := Simp[1/d
  Subst[Int[(a + b*ArcSec[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d},
  x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.72

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a)^2(bx+a)+2 \operatorname{arcsec}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)-2 \operatorname{arcsec}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)}{b}$
default	$\frac{\operatorname{arcsec}(bx+a)^2(bx+a)+2 \operatorname{arcsec}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)-2 \operatorname{arcsec}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)}{b}$

input `int(arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
1/b*(arcsec(b*x+a)^2*(b*x+a)+2*arcsec(b*x+a)*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x
+a)^2)^(1/2)))-2*arcsec(b*x+a)*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))
-2*I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+2*I*dilog(1-I*(1/(b*x+
a)+I*(1-1/(b*x+a)^2)^(1/2))))
```

Fricas [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(arcsec(b*x+a)^2,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^2, x)`

Sympy [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{asec}^2(a + bx) dx$$

input `integrate(asec(b*x+a)**2,x)`

output `Integral(asec(a + b*x)**2, x)`

Maxima [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(arcsec(b*x+a)^2,x, algorithm="maxima")`

output `x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/4*x*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate((2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

input `integrate(arcsec(b*x+a)^2,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

input `int(acos(1/(a + b*x))^2,x)`

output `int(acos(1/(a + b*x))^2, x)`

Reduce [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{asec}(bx + a)^2 dx$$

input `int(asec(b*x+a)^2,x)`

output `int(asec(a + b*x)**2,x)`

3.31 $\int \frac{\sec^{-1}(a+bx)^2}{x} dx$

Optimal result	247
Mathematica [B] (warning: unable to verify)	248
Rubi [A] (verified)	249
Maple [F]	255
Fricas [F]	255
Sympy [F]	255
Maxima [F]	256
Giac [F]	256
Mupad [F(-1)]	256
Reduce [F]	257

Optimal result

Integrand size = 12, antiderivative size = 310

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^2}{x} dx &= \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \sec^{-1}(a+bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 2i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&\quad - 2i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad + i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&\quad - \frac{1}{2} \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right)
\end{aligned}$$

output

```
arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-arcsec(b*x+a)^2*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-2*I*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))-2*I*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+I*arcsec(b*x+a)*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+2*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+2*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-1/2*polylog(3,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 813 vs. $2(310) = 620$.

Time = 1.52 (sec) , antiderivative size = 813, normalized size of antiderivative = 2.62

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \text{Too large to display}$$

input

```
Integrate[ArcSec[a + b*x]^2/x,x]
```

output

```

ArcSec[a + b*x]^2*Log[1 + (a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])]
+ ArcSec[a + b*x]^2*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a
] - 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqrt
[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + ArcSec[a + b*x]^2*Log[1 - (a*E^(I*A
rcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + ArcSec[a + b*x]^2*Log[1 - ((1 + Sq
rt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1
+ a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] -
2*ArcSec[a + b*x]^2*Log[1 + E^((2*I)*ArcSec[a + b*x])] + ArcSec[a + b*x]^
2*Log[(2*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/(a + b*x)] - ArcSe
c[a + b*x]^2*Log[1 + ((-1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a
+ b*x)^(-2)])))/a] + 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Lo
g[1 + ((-1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))
/a] - ArcSec[a + b*x]^2*Log[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*S
qrt[1 - (a + b*x)^(-2)])))/a] - 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/S
qrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x
)^(-2)])))/a] - (2*I)*ArcSec[a + b*x]*PolyLog[2, -((a*E^(I*ArcSec[a + b*x]
)))/(-1 + Sqrt[1 - a^2])] - (2*I)*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec
[a + b*x]))/(1 + Sqrt[1 - a^2])] + I*ArcSec[a + b*x]*PolyLog[2, -E^((2*I)*
ArcSec[a + b*x])] + 2*PolyLog[3, -((a*E^(I*ArcSec[a + b*x])))/(-1 + Sqrt[1
- a^2])] + 2*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])]...

```

Rubi [A] (verified)

Time = 1.58 (sec) , antiderivative size = 398, normalized size of antiderivative = 1.28, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {5781, 25, 5062, 5041, 25, 3042, 4202, 2620, 3011, 2720, 5031, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(a + bx)^2}{x} dx \\
 & \quad \downarrow \text{5781} \\
 & \int \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{bx} d \sec^{-1}(a + bx) \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
& - \int - \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{bx} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{5062} \\
& - \int \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{\frac{a}{a+bx} - 1} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{5041} \\
& \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2 d \sec^{-1}(a+bx) - \\
& \quad a \int - \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{25} \\
& \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2 d \sec^{-1}(a+bx) + \\
& \quad a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{3042} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \int \sec^{-1}(a+ \\
& \quad bx)^2 \tan(\sec^{-1}(a+bx)) d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{4202} \\
& -2i \int \frac{e^{2i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^2}{1 + e^{2i \sec^{-1}(a+bx)}} d \sec^{-1}(a+bx) + \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{3} i \sec^{-1}(a+bx)^3 \\
& \quad \downarrow \text{2620} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) - \\
& 2i \left(i \int \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) - \frac{1}{2} i \sec^{-1}(a+bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \right) + \\
& \quad \frac{1}{3} i \sec^{-1}(a+bx)^3
\end{aligned}$$

↓ 3011

$$-2i \left(i \left(\frac{1}{2} i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \int \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right) - a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{3} i \sec^{-1}(a+bx)^3 \right)$$

↓ 2720

$$-2i \left(i \left(\frac{1}{2} i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) de^{2i \sec^{-1}(a+bx)} \right) - a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{3} i \sec^{-1}(a+bx)^3 \right)$$

↓ 5031

$$a \left(-i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^2}{-e^{i \sec^{-1}(a+bx)} a - \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^2}{-e^{i \sec^{-1}(a+bx)} a + \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - \frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) de^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{3} i \sec^{-1}(a+bx)^3$$

↓ 2620

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{2i \int \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) d \sec^{-1}(a+bx)}{a} \right) - \frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) de^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{3} i \sec^{-1}(a+bx)^3$$

↓ 3011

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{2i \left(i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - i \int \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) de^{i \sec^{-1}(a+bx)} \right)}{a} \right) - \frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) de^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{3} i \sec^{-1}(a+bx)^3$$

↓ 2720

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a} - \frac{2i \left(i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) - \int e^{-i \sec^{-1}(a+bx)} dx\right)}{a} \right. \right. \\ \left. \left. 2i \left(i \left(\frac{1}{2} i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) - \frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) dx \right) \right) \right. \right. \\ \left. \left. \frac{1}{3} i \sec^{-1}(a+bx)^3 \right) \right)$$

↓ 7143

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a} - \frac{2i \left(i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) - \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \right)}{a} \right. \right. \\ \left. \left. 2i \left(i \left(\frac{1}{2} i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) - \frac{1}{4} \operatorname{PolyLog}\left(3, -e^{2i \sec^{-1}(a+bx)}\right) \right) \right) \right. \right. \\ \left. \left. \frac{1}{3} i \sec^{-1}(a+bx)^3 \right) \right)$$

input `Int[ArcSec[a + b*x]^2/x,x]`

output `(I/3)*ArcSec[a + b*x]^3 + a*(((-1/3*I)*ArcSec[a + b*x]^3)/a - I*((I*ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/a - ((2*I)*(I*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])]) - PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])]))/a) - I*((I*ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/a - ((2*I)*(I*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])]) - PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])]))/a) - (2*I)*((-1/2*I)*ArcSec[a + b*x]^2*Log[1 + E^((2*I)*ArcSec[a + b*x])] + I*((I/2)*ArcSec[a + b*x]*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] - PolyLog[3, -E^((2*I)*ArcSec[a + b*x])]/4))`

Definitions of rubi rules used

- rule 25 `Int[-(Fx), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2620 `Int[(((F-)((g-)*(e-) + (f-)*(x-)))(n-)*((c-) + (d-)*(x-)(m-))/
((a-) + (b-)*((F-)((g-)*(e-) + (f-)*(x-)))(n-)), x_Symbol] := Simp
[((c + d*x)m/(b*f*g*n*Log[F]))*Log[1 + b*((F(g*(e + f*x)))n/a], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)(m - 1)*Log[1 + b*((F(g*(e + f*x
)n/a], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2720 `Int[u-, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w-)*((a-)*(v-)(n-))(m-) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E((c-)*(a-) + (b-)*x)
*(F-)[v-] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e-)*((F-)((c-)*(a-) + (b-)*(x-)))(n-)]*((f-) + (g-)
*(x-)(m-)), x_Symbol] := Simp[(-f + g*x)m*(PolyLog[2, (-e)*(F(c*(a +
b*x)))n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)(
m - 1)*PolyLog[2, (-e)*(F(c*(a + b*x)))n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u-, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4202 `Int[(((c-) + (d-)*(x-)(m-))*tan[(e-) + (f-)*(x-)], x_Symbol] := Simp[I
*((c + d*x)(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)m*(E(2*I*(
e + f*x))/(1 + E(2*I*(e + f*x))))), x], x] /; FreeQ[{c, d, e, f}, x] && IGt
Q[m, 0]`

rule 5031

```
Int[(((e_.) + (f_.)*(x_.))^(m_.)*Sin[(c_.) + (d_.)*(x_.)]/(Cos[(c_.) + (d_.)
*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[I*((e + f*x)^(m + 1)/(b*f*(m + 1)))
, x] + (-Simp[I Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a - Rt[a^2 - b^2, 2] +
b*E^(I*(c + d*x)))]), x], x] - Simp[I Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a
+ Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x)))]), x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

rule 5041

```
Int[(((e_.) + (f_.)*(x_.))^(m_.)*Tan[(c_.) + (d_.)*(x_.)]^(n_.))/(Cos[(c_.) +
(d_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] := Simp[1/a Int[(e + f*x)^m*Tan[c +
d*x]^n, x], x] - Simp[b/a Int[(e + f*x)^m*Sin[c + d*x]*(Tan[c + d*x]^(n
- 1)/(a + b*Cos[c + d*x]))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[
m, 0] && IGtQ[n, 0]
```

rule 5062

```
Int[(((e_.) + (f_.)*(x_.))^(m_.)*(F_)[(c_.) + (d_.)*(x_.)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_.)]^(p_.))/(a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_.)]), x_Symbol] := In
t[(e + f*x)^m*Cos[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cos[c + d*x]))
, x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m
, n, p]
```

rule 5781

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d
*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e
, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx+a)^2}{x} dx$$

input `int(arcsec(b*x+a)^2/x,x)`

output `int(arcsec(b*x+a)^2/x,x)`

Fricas [F]

$$\int \frac{\sec^{-1}(a+bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx+a)^2}{x} dx$$

input `integrate(arcsec(b*x+a)^2/x,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^2/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a+bx)^2}{x} dx = \int \frac{\operatorname{asec}^2(a+bx)}{x} dx$$

input `integrate(asec(b*x+a)**2/x,x)`

output `Integral(asec(a + b*x)**2/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

input `integrate(arcsec(b*x+a)^2/x,x, algorithm="maxima")`

output `integrate(arcsec(b*x + a)^2/x, x)`

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

input `integrate(arcsec(b*x+a)^2/x,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{acos}\left(\frac{1}{a+bx}\right)^2}{x} dx$$

input `int(acos(1/(a + b*x))^2/x,x)`

output `int(acos(1/(a + b*x))^2/x, x)`

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{asec}(bx + a)^2}{x} dx$$

input `int(asec(b*x+a)^2/x,x)`

output `int(asec(a + b*x)**2/x,x)`

3.32 $\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx$

Optimal result	258
Mathematica [B] (warning: unable to verify)	259
Rubi [A] (verified)	260
Maple [A] (verified)	262
Fricas [F]	262
Sympy [F]	263
Maxima [F]	263
Giac [F]	263
Mupad [F(-1)]	264
Reduce [F]	264

Optimal result

Integrand size = 12, antiderivative size = 244

$$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx = -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

output

```
-b*arcsec(b*x+a)^2/a-arcsec(b*x+a)^2/x-2*I*b*arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*I*b*arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*b*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 686 vs. $2(244) = 488$.

Time = 1.46 (sec) , antiderivative size = 686, normalized size of antiderivative = 2.81

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \text{Too large to display}$$

input `Integrate[ArcSec[a + b*x]^2/x^2,x]`

output

```

-(((a + b*x)*ArcSec[a + b*x]^2)/x + (2*b*(2*ArcSec[a + b*x]*ArcTanh[(-1
+ a)*Cot[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]] - 2*ArcCos[a^(-1)]*ArcTanh[(((
1 + a)*Tan[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] - (2*I)*A
rcTanh[(-1 + a)*Cot[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]] + (2*I)*ArcTanh[(((
1 + a)*Tan[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[Sqrt[-1 + a^2]/(Sqrt[
2]*Sqrt[a]*E^((I/2)*ArcSec[a + b*x])*Sqrt[-((b*x)/(a + b*x))])] + (ArcCos[
a^(-1)] + (2*I)*(ArcTanh[(-1 + a)*Cot[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]]
- ArcTanh[(((1 + a)*Tan[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]])*Log[(Sqrt[-1
+ a^2]*E^((I/2)*ArcSec[a + b*x]))/(Sqrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x)
))])] - (ArcCos[a^(-1)] - (2*I)*ArcTanh[(((1 + a)*Tan[ArcSec[a + b*x]/2])/Sq
rt[-1 + a^2]])*Log[(-1 + a)*(I + I*a + Sqrt[-1 + a^2])*(-I + Tan[ArcSec[a
+ b*x]/2]))/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] - (ArcC
os[a^(-1)] + (2*I)*ArcTanh[(((1 + a)*Tan[ArcSec[a + b*x]/2])/Sqrt[-1 + a^2]
])*Log[(-1 + a)*(-I - I*a + Sqrt[-1 + a^2])*(I + Tan[ArcSec[a + b*x]/2]))
/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] + I*(-PolyLog[2, ((
1 - I*Sqrt[-1 + a^2])*(1 - a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))/(a*
(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] + PolyLog[2, ((1 + I*Sq
rt[-1 + a^2])*(1 - a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))/(a*(-1 + a
+ Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))])))/Sqrt[-1 + a^2])/a)

```

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5781, 4926, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(a+bx)^2}{x^2} dx \\
 & \quad \downarrow \text{5781} \\
 & b \int \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{b^2 x^2} d \sec^{-1}(a+bx) \\
 & \quad \downarrow \text{4926} \\
 & b \left(-2 \int -\frac{\sec^{-1}(a+bx)}{bx} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^2}{bx} \right) \\
 & \quad \downarrow \text{3042} \\
 & b \left(-2 \int \frac{\sec^{-1}(a+bx)}{a - \csc(\sec^{-1}(a+bx) + \frac{\pi}{2})} d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^2}{bx} \right) \\
 & \quad \downarrow \text{4679} \\
 & b \left(-2 \int \left(\frac{\sec^{-1}(a+bx)}{a} + \frac{\sec^{-1}(a+bx)}{a \left(\frac{a}{a+bx} - 1 \right)} \right) d \sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^2}{bx} \right) \\
 & \quad \downarrow \text{2009} \\
 & b \left(-\frac{\sec^{-1}(a+bx)^2}{bx} - 2 \left(\frac{\text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right)}{a\sqrt{1-a^2}} - \frac{\text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right)}{a\sqrt{1-a^2}} + \frac{i \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right)}{a\sqrt{1-a^2}} \right) \right)
 \end{aligned}$$

input

```
Int[ArcSec[a + b*x]^2/x^2,x]
```

output

```
b*(-(ArcSec[a + b*x]^2/(b*x)) - 2*(ArcSec[a + b*x]^2/(2*a) + (I*ArcSec[a +
b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 -
a^2]) - (I*ArcSec[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 -
a^2])])/(a*Sqrt[1 - a^2]) + PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqr
t[1 - a^2])]/(a*Sqrt[1 - a^2]) - PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 +
Sqrt[1 - a^2])]/(a*Sqrt[1 - a^2]))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4679

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

rule 4926

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Simp[f*m/(b*d*(n +
1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ
[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 5781

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d
*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e
, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.38

method	result
derivativedivides	$b \left(-\frac{(bx+a) \operatorname{arcsec}(bx+a)^2}{abx} - \frac{2i\sqrt{-a^2+1} \operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right)}{a(a^2-1)} \right) + \frac{2i\sqrt{-a^2+1}}{a(a^2-1)}$
default	$b \left(-\frac{(bx+a) \operatorname{arcsec}(bx+a)^2}{abx} - \frac{2i\sqrt{-a^2+1} \operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1 + \sqrt{-a^2+1}} \right)}{a(a^2-1)} \right) + \frac{2i\sqrt{-a^2+1}}{a(a^2-1)}$

input `int(arcsec(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)`

output `b*(-(b*x+a)*arcsec(b*x+a)^2/a/b/x-2*I*(-a^2+1)^(1/2)/a/(a^2-1)*arcsec(b*x+a)*ln((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+2*I*(-a^2+1)^(1/2)/a/(a^2-1)*arcsec(b*x+a)*ln((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))-2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2))))`

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

input `integrate(arcsec(b*x+a)^2/x^2,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^2/x^2, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{asec}^2(a + bx)}{x^2} dx$$

input `integrate(asec(b*x+a)**2/x**2,x)`

output `Integral(asec(a + b*x)**2/x**2, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

input `integrate(arcsec(b*x+a)^2/x^2,x, algorithm="maxima")`

output `-1/4*(4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 4*x*integrate((2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2), x) - log(b^2*x^2 + 2*a*b*x + a^2)^2)/x`

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

input `integrate(arcsec(b*x+a)^2/x^2,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)^2/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)^2}{x^2} dx$$

input `int(acos(1/(a + b*x))^2/x^2,x)`output `int(acos(1/(a + b*x))^2/x^2, x)`**Reduce [F]**

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{asec}(bx + a)^2}{x^2} dx$$

input `int(asec(b*x+a)^2/x^2,x)`output `int(asec(a + b*x)**2/x**2,x)`

3.33 $\int x^2 \sec^{-1}(a + bx)^3 dx$

Optimal result	266
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	270
Fricas [F]	271
Sympy [F]	271
Maxima [F]	272
Giac [F]	272
Mupad [F(-1)]	273
Reduce [F]	273

Optimal result

Integrand size = 12, antiderivative size = 494

$$\begin{aligned}
\int x^2 \sec^{-1}(a+bx)^3 dx = & \frac{(a+bx) \sec^{-1}(a+bx)}{b^3} - \frac{3ia \sec^{-1}(a+bx)^2}{b^3} \\
& + \frac{3a(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{b^3} \\
& - \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{2b^3} + \frac{a^3 \sec^{-1}(a+bx)^3}{3b^3} \\
& + \frac{\frac{1}{3} x^3 \sec^{-1}(a+bx)^3 + \frac{i \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3}}{3} \\
& + \frac{6ia^2 \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^3} \\
& + \frac{6a \sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^3} \\
& - \frac{i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& - \frac{6ia^2 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& + \frac{i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& + \frac{6ia^2 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& - \frac{3ia \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{b^3} \\
& + \frac{\operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& + \frac{6a^2 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
& - \frac{\operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} - \frac{6a^2 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

output

```
(b*x+a)*arcsec(b*x+a)/b^3-3*I*a*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/b^3+3*a*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a)^2/b^3-1/2*(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)*arcsec(b*x+a)^2/b^3+1/3*a^3*arcsec(b*x+a)^3/b^3+1/3*x^3*arcsec(b*x+a)^3+I*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^3-I*arcsec(b*x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3-arctanh((1-1/(b*x+a)^2)^(1/2))/b^3+6*a*arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/b^3+I*arcsec(b*x+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3-6*I*a^2*arcsec(b*x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3+6*I*a^2*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^3-3*I*a*arcsec(b*x+a)^2/b^3+6*I*a^2*arcsec(b*x+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3+polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3+6*a^2*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3-polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3-6*a^2*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.90

$$\int x^2 \sec^{-1}(a + bx)^3 dx$$

$$= \frac{-\coth^{-1}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right) + (a + bx) \sec^{-1}(a + bx) - 3ia \sec^{-1}(a + bx)^2 + 3a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3}$$

input

```
Integrate[x^2*ArcSec[a + b*x]^3,x]
```

output

```
(-ArcCoth[Sqrt[1 - (a + b*x)^(-2)]] + (a + b*x)*ArcSec[a + b*x] - (3*I)*a*
ArcSec[a + b*x]^2 + 3*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]
^2 - ((a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/2 + (a^3*Arc
Sec[a + b*x]^3)/3 + (b^3*x^3*ArcSec[a + b*x]^3)/3 + I*ArcSec[a + b*x]^2*Ar
cTan[E^(I*ArcSec[a + b*x])] + (6*I)*a^2*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcS
ec[a + b*x])] + 6*a*ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] - I
*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - (6*I)*a^2*ArcSec
[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] + I*ArcSec[a + b*x]*PolyL
og[2, I*E^(I*ArcSec[a + b*x])] + (6*I)*a^2*ArcSec[a + b*x]*PolyLog[2, I*E^
(I*ArcSec[a + b*x])] - (3*I)*a*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] + Po
lyLog[3, (-I)*E^(I*ArcSec[a + b*x])] + 6*a^2*PolyLog[3, (-I)*E^(I*ArcSec[a
+ b*x])] - PolyLog[3, I*E^(I*ArcSec[a + b*x])] - 6*a^2*PolyLog[3, I*E^(I*
ArcSec[a + b*x])])/b^3
```

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5781, 4926, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sec^{-1}(a + bx)^3 dx \\
 & \quad \downarrow \text{5781} \\
 & \frac{\int b^2 x^2 (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3 d \sec^{-1}(a + bx)}{b^3} \\
 & \quad \downarrow \text{4926} \\
 & \frac{\int -b^3 x^3 \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a + bx)^3}{b^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sec^{-1}(a + bx)^2 \left(a - \csc \left(\sec^{-1}(a + bx) + \frac{\pi}{2} \right) \right)^3 d \sec^{-1}(a + bx) + \frac{1}{3} b^3 x^3 \sec^{-1}(a + bx)^3}{b^3} \\
 & \quad \downarrow \text{4678}
 \end{aligned}$$

$$\int \frac{(\sec^{-1}(a+bx))^2 a^3 - 3(a+bx)\sec^{-1}(a+bx)^2 a^2 + 3(a+bx)^2 \sec^{-1}(a+bx)^2 a - (a+bx)^3 \sec^{-1}(a+bx)^2}{b^3} dx$$

↓ 2009

$$\frac{1}{3}a^3 \sec^{-1}(a+bx)^3 + 6ia^2 \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right) - 6ia^2 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)$$

input

```
Int[x^2*ArcSec[a + b*x]^3,x]
```

output

```
((a + b*x)*ArcSec[a + b*x] - (3*I)*a*ArcSec[a + b*x]^2 + 3*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2 - ((a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/2 + (a^3*ArcSec[a + b*x]^3)/3 + (b^3*x^3*ArcSec[a + b*x]^3)/3 + I*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] + (6*I)*a^2*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] - ArcTanh[Sqrt[1 - (a + b*x)^(-2)]] + 6*a*ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] - I*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - (6*I)*a^2*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] + I*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])] + (6*I)*a^2*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])] - (3*I)*a*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] + PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])] + 6*a^2*PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])] - PolyLog[3, I*E^(I*ArcSec[a + b*x])] - 6*a^2*PolyLog[3, I*E^(I*ArcSec[a + b*x])])/b^3
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4678

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

rule 4926

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

rule 5781

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)])*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 716, normalized size of antiderivative = 1.45

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a) \left(6 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - 6 \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + 2 \operatorname{arcsec}(bx+a)^2 (bx+a)^3 + 18 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) \right)}{6}$
default	$\frac{\operatorname{arcsec}(bx+a) \left(6 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - 6 \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + 2 \operatorname{arcsec}(bx+a)^2 (bx+a)^3 + 18 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) \right)}{6}$

input

```
int(x^2*arcsec(b*x+a)^3,x,method=_RETURNVERBOSE)
```

output

```

1/b^3*(1/6*arcsec(b*x+a)*(6*arcsec(b*x+a)^2*a^2*(b*x+a)-6*arcsec(b*x+a)^2*
a*(b*x+a)^2+2*arcsec(b*x+a)^2*(b*x+a)^3+18*arcsec(b*x+a)*(((b*x+a)^2-1)/(b
*x+a)^2)^(1/2)*a*(b*x+a)-3*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(
b*x+a)^2+18*I*a*arcsec(b*x+a)+6*b*x+6*a)-I*arcsec(b*x+a)*polylog(2,-I*(1/(
b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+6*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)
^2)^(1/2)))*a^2*arcsec(b*x+a)+3*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))
)*a^2*arcsec(b*x+a)^2+6*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*
a^2-6*I*arcsec(b*x+a)^2*a+I*arcsec(b*x+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b
*x+a)^2)^(1/2)))-3*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a^2*arcsec(
b*x+a)^2-6*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a^2+6*ln(1+(1/
(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)*a*arcsec(b*x+a)+2*I*arctan(1/(b*x+a)+I
*(1-1/(b*x+a)^2)^(1/2))-3*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))
^2)*a+1/2*arcsec(b*x+a)^2*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-6*I*
polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a^2*arcsec(b*x+a)+polylo
g(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-1/2*arcsec(b*x+a)^2*ln(1-I*(1/
(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)
^(1/2))))

```

Fricas [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

input

```
integrate(x^2*arcsec(b*x+a)^3,x, algorithm="fricas")
```

output

```
integral(x^2*arcsec(b*x + a)^3, x)
```

Sympy [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{asec}^3(a + bx) dx$$

input

```
integrate(x**2*asec(b*x+a)**3,x)
```


output `Integral(x**2*asec(a + b*x)**3, x)`

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(x^2*arcsec(b*x+a)^3,x, algorithm="maxima")`

output `1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 1/4*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/4*((4*b*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x^3*log(b^2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) + 4*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2 - 1)*b*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(x^2*arcsec(b*x+a)^3,x, algorithm="giac")`

output `integrate(x^2*arcsec(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{acos}\left(\frac{1}{a + bx}\right)^3 dx$$

input `int(x^2*acos(1/(a + b*x))^3,x)`output `int(x^2*acos(1/(a + b*x))^3, x)`**Reduce [F]**

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int \operatorname{asec}(bx + a)^3 x^2 dx$$

input `int(x^2*asec(b*x+a)^3,x)`output `int(asec(a + b*x)**3*x**2,x)`

3.34 $\int x \sec^{-1}(a + bx)^3 dx$

Optimal result	274
Mathematica [A] (verified)	275
Rubi [A] (verified)	276
Maple [A] (verified)	278
Fricas [F]	278
Sympy [F]	279
Maxima [F]	279
Giac [F]	280
Mupad [F(-1)]	280
Reduce [F]	280

Optimal result

Integrand size = 10, antiderivative size = 278

$$\begin{aligned}
 \int x \sec^{-1}(a + bx)^3 dx = & \frac{3i \sec^{-1}(a + bx)^2}{2b^2} - \frac{3(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{2b^2} \\
 & - \frac{a^2 \sec^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \sec^{-1}(a + bx)^3 \\
 & - \frac{6ia \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{3 \sec^{-1}(a + bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{6ia \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2b^2} \\
 & - \frac{6a \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{6a \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^2}
 \end{aligned}$$

output

$$\frac{3}{2}I \operatorname{arcsec}(bx+a)^2/b^2 - 3/2*(bx+a)*(1-1/(bx+a)^2)^{(1/2)}*\operatorname{arcsec}(bx+a)^2/b^2 - 1/2*a^2*\operatorname{arcsec}(bx+a)^3/b^2 + 1/2*x^2*\operatorname{arcsec}(bx+a)^3 - 6*I*a*\operatorname{arcsec}(bx+a)^2*\arctan(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)})/b^2 - 3*\operatorname{arcsec}(bx+a)*\ln(1+(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)})^2)/b^2 + 6*I*a*\operatorname{arcsec}(bx+a)*\operatorname{polylog}(2,-I*(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)}))/b^2 - 6*I*a*\operatorname{arcsec}(bx+a)*\operatorname{polylog}(2,I*(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)}))/b^2 + 3/2*I*\operatorname{polylog}(2,-(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)})^2)/b^2 - 6*a*\operatorname{polylog}(3,-I*(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)}))/b^2 + 6*a*\operatorname{polylog}(3,I*(1/(bx+a)+I*(1-1/(bx+a)^2)^{(1/2)}))/b^2$$
Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.92

$$\int x \sec^{-1}(a + bx)^3 dx$$

$$= \frac{\frac{3}{2}i \sec^{-1}(a + bx)^2 - \frac{3}{2}(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2 - a(a + bx) \sec^{-1}(a + bx)^3 + \frac{1}{2}(a + bx)^2 \sec^{-1}(a + bx)}{b^2}$$

input

Integrate[x*ArcSec[a + b*x]^3,x]

output

$$\frac{(((3I)/2)*\operatorname{ArcSec}[a + b*x]^2 - (3*(a + b*x)*\operatorname{Sqrt}[1 - (a + b*x)^{-2}])*\operatorname{ArcSec}[a + b*x]^2)/2 - a*(a + b*x)*\operatorname{ArcSec}[a + b*x]^3 + ((a + b*x)^2*\operatorname{ArcSec}[a + b*x]^3)/2 - (6I)*a*\operatorname{ArcSec}[a + b*x]^2*\operatorname{ArcTan}[E^{(I*\operatorname{ArcSec}[a + b*x])}] - 3*\operatorname{ArcSec}[a + b*x]*\operatorname{Log}[1 + E^{((2I)*\operatorname{ArcSec}[a + b*x])}] + (6I)*a*\operatorname{ArcSec}[a + b*x]*\operatorname{PolyLog}[2, (-I)*E^{(I*\operatorname{ArcSec}[a + b*x])}] - (6I)*a*\operatorname{ArcSec}[a + b*x]*\operatorname{PolyLog}[2, I*E^{(I*\operatorname{ArcSec}[a + b*x])}] + ((3I)/2)*\operatorname{PolyLog}[2, -E^{((2I)*\operatorname{ArcSec}[a + b*x])}] - 6*a*\operatorname{PolyLog}[3, (-I)*E^{(I*\operatorname{ArcSec}[a + b*x])}] + 6*a*\operatorname{PolyLog}[3, I*E^{(I*\operatorname{ArcSec}[a + b*x])}])/b^2$$

Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.91, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5781, 25, 4926, 3042, 4678, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sec^{-1}(a + bx)^3 dx \\
 & \quad \downarrow \text{5781} \\
 & \frac{\int bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3 d \sec^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{25} \\
 & - \frac{\int -bx(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3 d \sec^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{4926} \\
 & \frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx)^3 - \frac{3}{2} \int b^2x^2 \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx)^3 - \frac{3}{2} \int \sec^{-1}(a + bx)^2 \left(a - \csc(\sec^{-1}(a + bx) + \frac{\pi}{2})\right)^2 d \sec^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{4678} \\
 & \frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx)^3 - \frac{3}{2} \int (a^2 \sec^{-1}(a + bx)^2 + (a + bx)^2 \sec^{-1}(a + bx)^2 - 2a(a + bx) \sec^{-1}(a + bx)^2) d \sec^{-1}(a + bx)}{b^2} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\frac{1}{2}b^2x^2 \sec^{-1}(a + bx)^3 - \frac{3}{2} \left(\frac{1}{3}a^2 \sec^{-1}(a + bx)^3 + 4ia \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right) - 4ia \sec^{-1}(a + bx) \right)}{b^2}
 \end{aligned}$$

input `Int[x*ArcSec[a + b*x]^3,x]`

output

$$\frac{\left((b^2 x^2 \operatorname{ArcSec}[a + b x]^3) / 2 - (3((-I) \operatorname{ArcSec}[a + b x]^2 + (a + b x) \sqrt{1 - (a + b x)^{-2}}) \operatorname{ArcSec}[a + b x]^2 + (a^2 \operatorname{ArcSec}[a + b x]^3) / 3 + (4I) a \operatorname{ArcSec}[a + b x]^2 \operatorname{ArcTan}[E^{(I \operatorname{ArcSec}[a + b x])}] + 2 \operatorname{ArcSec}[a + b x] \operatorname{Log}[1 + E^{((2I) \operatorname{ArcSec}[a + b x])}] - (4I) a \operatorname{ArcSec}[a + b x] \operatorname{PolyLog}[2, (-I) E^{(I \operatorname{ArcSec}[a + b x])}] + (4I) a \operatorname{ArcSec}[a + b x] \operatorname{PolyLog}[2, I E^{(I \operatorname{ArcSec}[a + b x])}] - I \operatorname{PolyLog}[2, -E^{((2I) \operatorname{ArcSec}[a + b x])}] + 4 a \operatorname{PolyLog}[3, (-I) E^{(I \operatorname{ArcSec}[a + b x])}] - 4 a \operatorname{PolyLog}[3, I E^{(I \operatorname{ArcSec}[a + b x])}]) / 2) \right)}{b^2}$$

Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$$

rule 2009

$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 3042

$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4678

$$\operatorname{Int}[(\operatorname{csc}[e] + (f)(x))(b) + (a)^{(n)}((c) + (d)(x))^{(m)}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(c + d x)^m, (a + b \operatorname{Csc}[e + f x])^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m\}, x] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{IGtQ}[n, 0]$$

rule 4926

$$\operatorname{Int}[(e + (f)(x))^{(m)} \operatorname{Sec}[(c) + (d)(x)]((a) + (b) \operatorname{Sec}[(c) + (d)(x)])^{(n)} \operatorname{Tan}[(c) + (d)(x)], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(e + f x)^m ((a + b \operatorname{Sec}[c + d x])^{(n+1)} / (b d (n+1))), x] - \operatorname{Simp}[f (m / (b d (n+1))) \operatorname{Int}[(e + f x)^{(m-1)} (a + b \operatorname{Sec}[c + d x])^{(n+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n\}, x] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{NeQ}[n, -1]$$

rule 5781

$$\operatorname{Int}[(a) + \operatorname{ArcSec}[(c) + (d)(x)](b)^{(p)}(e + (f)(x))^{(m)}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[1/d^{(m+1)} \operatorname{Subst}[\operatorname{Int}[(a + b x)^p \operatorname{Sec}[x] \operatorname{Tan}[x] (d e - c f + f \operatorname{Sec}[x])^m, x], x, \operatorname{ArcSec}[c + d x]], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \operatorname{IGtQ}[p, 0] \ \&\& \operatorname{IntegerQ}[m]$$

Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.36

method	result
derivativedivides	$-\frac{\operatorname{arcsec}(bx+a)^2 \left(2 \operatorname{arcsec}(bx+a)a(bx+a) - \operatorname{arcsec}(bx+a)(bx+a)^2 + 3 \sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)+3i \right)}{2} + 6i \operatorname{polylog} \left(2, -i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right)$
default	$-\frac{\operatorname{arcsec}(bx+a)^2 \left(2 \operatorname{arcsec}(bx+a)a(bx+a) - \operatorname{arcsec}(bx+a)(bx+a)^2 + 3 \sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)+3i \right)}{2} + 6i \operatorname{polylog} \left(2, -i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right)$

input `int(x*arcsec(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
1/b^2*(-1/2*arcsec(b*x+a)^2*(2*arcsec(b*x+a)*a*(b*x+a)-arcsec(b*x+a)*(b*x+a)^2+3*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)+3*I)+6*I*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a*arcsec(b*x+a)-3*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a*arcsec(b*x+a)^2-6*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a-6*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a*arcsec(b*x+a)+3*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a*arcsec(b*x+a)^2+6*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a+3*I*arcsec(b*x+a)^2-3*arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+3/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2))
```

Fricas [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(x*arcsec(b*x+a)^3,x, algorithm="fricas")`

output `integral(x*arcsec(b*x + a)^3, x)`

Sympy [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{asec}^3(a + bx) dx$$

input `integrate(x*asec(b*x+a)**3,x)`

output `Integral(x*asec(a + b*x)**3, x)`

Maxima [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(x*arcsec(b*x+a)^3,x, algorithm="maxima")`

output `1/2*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 3/8*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(3/8*((4*b*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x^2*log(b^2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) + 4*(2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a)^2 - (b^3*x^4 + 2*a*b^2*x^3 + (a^2 - 1)*b*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(x*arcsec(b*x+a)^3,x, algorithm="giac")`

output `integrate(x*arcsec(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{acos}\left(\frac{1}{a + bx}\right)^3 dx$$

input `int(x*acos(1/(a + b*x))^3,x)`

output `int(x*acos(1/(a + b*x))^3, x)`

Reduce [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int a \sec(bx + a)^3 x dx$$

input `int(x*asec(b*x+a)^3,x)`

output `int(asec(a + b*x)**3*x,x)`

3.35 $\int \sec^{-1}(a + bx)^3 dx$

Optimal result	281
Mathematica [A] (verified)	282
Rubi [A] (verified)	282
Maple [F]	285
Fricas [F]	285
Sympy [F]	286
Maxima [F]	286
Giac [F]	286
Mupad [F(-1)]	287
Reduce [F]	287

Optimal result

Integrand size = 8, antiderivative size = 154

$$\int \sec^{-1}(a + bx)^3 dx = \frac{(a + bx) \sec^{-1}(a + bx)^3}{b} + \frac{6i \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$- \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$+ \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$+ \frac{6 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{6 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

output

```
(b*x+a)*arcsec(b*x+a)^3/b+6*I*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b-6*I*arcsec(b*x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+6*I*arcsec(b*x+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+6*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b-6*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.04

$$\int \sec^{-1}(a + bx)^3 dx$$

$$= \frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \sec^{-1}(a + bx)^2 \left(\log \left(1 - ie^{i \sec^{-1}(a+bx)} \right) - \log \left(1 + ie^{i \sec^{-1}(a+bx)} \right) \right) - 6i \sec^{-1}(a + bx)}{b}$$

input

```
Integrate[ArcSec[a + b*x]^3,x]
```

output

```
((a + b*x)*ArcSec[a + b*x]^3 - 3*ArcSec[a + b*x]^2*(Log[1 - I*E^(I*ArcSec[a + b*x])] - Log[1 + I*E^(I*ArcSec[a + b*x])]) - (6*I)*ArcSec[a + b*x]*(PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - PolyLog[2, I*E^(I*ArcSec[a + b*x])]) + 6*(PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])] - PolyLog[3, I*E^(I*ArcSec[a + b*x])]))/b
```

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5775, 5739, 4244, 3042, 4669, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sec^{-1}(a + bx)^3 dx$$

$$\downarrow 5775$$

$$\frac{\int \sec^{-1}(a + bx)^3 d(a + bx)}{b}$$

$$\downarrow 5739$$

$$\frac{\int (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3 d \sec^{-1}(a + bx)}{b}$$

$$\downarrow 4244$$

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \int (a + bx) \sec^{-1}(a + bx)^2 d \sec^{-1}(a + bx)}{b}$$

↓ 3042

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \int \sec^{-1}(a + bx)^2 \csc \left(\sec^{-1}(a + bx) + \frac{\pi}{2} \right) d \sec^{-1}(a + bx)}{b}$$

↓ 4669

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \left(-2 \int \sec^{-1}(a + bx) \log \left(1 - i e^{i \sec^{-1}(a + bx)} \right) d \sec^{-1}(a + bx) + 2 \int \sec^{-1}(a + bx) \log \right)}{b}$$

↓ 3011

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \left(2 \left(i \sec^{-1}(a + bx) \text{PolyLog} \left(2, -i e^{i \sec^{-1}(a + bx)} \right) - i \int \text{PolyLog} \left(2, -i e^{i \sec^{-1}(a + bx)} \right) d \right) \right)}{b}$$

↓ 2720

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \left(2 \left(i \sec^{-1}(a + bx) \text{PolyLog} \left(2, -i e^{i \sec^{-1}(a + bx)} \right) - \int e^{-i \sec^{-1}(a + bx)} \text{PolyLog} \left(2, -i e^{i \sec^{-1}(a + bx)} \right) d \right) \right)}{b}$$

↓ 7143

$$\frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \left(-2i \sec^{-1}(a + bx)^2 \arctan \left(e^{i \sec^{-1}(a + bx)} \right) + 2 \left(i \sec^{-1}(a + bx) \text{PolyLog} \left(2, -i e^{i \sec^{-1}(a + bx)} \right) \right) \right)}{b}$$

input

```
Int[ArcSec[a + b*x]^3,x]
```

output

```
((a + b*x)*ArcSec[a + b*x]^3 - 3*((-2*I)*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] + 2*(I*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x]]) - PolyLog[3, (-I)*E^(I*ArcSec[a + b*x]])] - 2*(I*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])] - PolyLog[3, I*E^(I*ArcSec[a + b*x])]))/b
```

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4244 `Int[(x_)^(m_.)*Sec[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tan[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[x^(m - n + 1)*(Sec[a + b*x^n]^p/(b*n*p)), x] - Simp[(m - n + 1)/(b*n*p) Int[x^(m - n)*Sec[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]`

rule 4669 `Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5739 `Int[((a_.) + ArcSec[(c_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[1/c Subst[Int[(a + b*x)^n*Sec[x]*Tan[x], x], x, ArcSec[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]`

rule 5775 `Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Simp[1/d
Subst[Int[(a + b*ArcSec[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d},
x] && IGtQ[p, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [F]

$$\int \operatorname{arcsec}(bx + a)^3 dx$$

input `int(arcsec(b*x+a)^3,x)`

output `int(arcsec(b*x+a)^3,x)`

Fricas [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(arcsec(b*x+a)^3,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^3, x)`

Sympy [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{asec}^3(a + bx) dx$$

input `integrate(asec(b*x+a)**3,x)`

output `Integral(asec(a + b*x)**3, x)`

Maxima [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(arcsec(b*x+a)^3,x, algorithm="maxima")`

output `x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 3/4*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(3/4*(4*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x*log(b^2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) + 4*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)`

Giac [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

input `integrate(arcsec(b*x+a)^3,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(a + bx)^3 dx = \int \arccos\left(\frac{1}{a + bx}\right)^3 dx$$

input `int(acos(1/(a + b*x))^3,x)`

output `int(acos(1/(a + b*x))^3, x)`

Reduce [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{asec}(bx + a)^3 dx$$

input `int(asec(b*x+a)^3,x)`

output `int(asec(a + b*x)**3,x)`

3.36 $\int \frac{\sec^{-1}(a+bx)^3}{x} dx$

Optimal result	289
Mathematica [B] (warning: unable to verify)	290
Rubi [A] (verified)	291
Maple [F]	297
Fricas [F]	297
Sympy [F]	298
Maxima [F]	298
Giac [F]	298
Mupad [F(-1)]	299
Reduce [F]	299

Optimal result

Integrand size = 12, antiderivative size = 430

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^3}{x} dx &= \sec^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ \sec^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \sec^{-1}(a+bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&- 3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&- 3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&+ \frac{3}{2} i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&+ 6 \sec^{-1}(a+bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ 6 \sec^{-1}(a+bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \frac{3}{2} \sec^{-1}(a+bx) \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) \\
&+ 6i \operatorname{PolyLog} \left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
&+ 6i \operatorname{PolyLog} \left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
&- \frac{3}{4} i \operatorname{PolyLog} \left(4, -e^{2i \sec^{-1}(a+bx)} \right)
\end{aligned}$$

output

```
arcsec(b*x+a)^3*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x+a)^3*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-arcsec(b*x+a)^3*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-3*I*arcsec(b*x+a)^2*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))-3*I*arcsec(b*x+a)^2*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+3/2*I*arcsec(b*x+a)^2*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+6*arcsec(b*x+a)*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+6*arcsec(b*x+a)*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-3/2*arcsec(b*x+a)*polylog(3,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+6*I*polylog(4,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+6*I*polylog(4,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))-3/4*I*polylog(4,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1058 vs. $2(430) = 860$.

Time = 2.55 (sec) , antiderivative size = 1058, normalized size of antiderivative = 2.46

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \text{Too large to display}$$

input

```
Integrate[ArcSec[a + b*x]^3/x,x]
```

output

```

2*ArcSec[a + b*x]^3*Log[1 + (a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])
] + ArcSec[a + b*x]^3*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))
/a] - 6*ArcSec[a + b*x]^2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 +
Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 2*ArcSec[a + b*x]^3*Log[1 - (a*
E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + ArcSec[a + b*x]^3*Log[1 - ((
1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 6*ArcSec[a + b*x]^2*ArcSin[
Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]
)))/a] - 3*ArcSec[a + b*x]^3*Log[1 + E^((2*I)*ArcSec[a + b*x])] + 2*ArcSec
[a + b*x]^3*Log[(2*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/(a + b*x
)] - ArcSec[a + b*x]^3*Log[1 + (a*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-
2)])))/(-1 + Sqrt[1 - a^2])] - ArcSec[a + b*x]^3*Log[1 + ((-1 + Sqrt[1 - a
^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/a] + 6*ArcSec[a + b*x]
^2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqrt[1 - a^2])*((a + b*
x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/a] - ArcSec[a + b*x]^3*Log[1 - (a*(
(a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/(1 + Sqrt[1 - a^2])] - ArcSe
c[a + b*x]^3*Log[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a
+ b*x)^(-2)]))/a] - 6*ArcSec[a + b*x]^2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*L
og[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))
/a] - (3*I)*ArcSec[a + b*x]^2*PolyLog[2, -(a*E^(I*ArcSec[a + b*x]))/(-1 +
Sqrt[1 - a^2])] - (3*I)*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I*ArcSec[a...

```

Rubi [A] (verified)

Time = 1.85 (sec) , antiderivative size = 529, normalized size of antiderivative = 1.23, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {5781, 25, 5062, 5041, 25, 3042, 4202, 2620, 3011, 5031, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx$$

$$\downarrow \text{5781}$$

$$\int \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3}{bx} d \sec^{-1}(a + bx)$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& - \int - \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{bx} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{5062} \\
& - \int \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{\frac{a}{a+bx} - 1} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{5041} \\
& \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3 d \sec^{-1}(a+bx) - \\
& \quad a \int - \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{25} \\
& \int (a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3 d \sec^{-1}(a+bx) + \\
& \quad a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{3042} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \int \sec^{-1}(a+ \\
& \quad bx)^3 \tan(\sec^{-1}(a+bx)) d \sec^{-1}(a+bx) \\
& \quad \downarrow \text{4202} \\
& -2i \int \frac{e^{2i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^3}{1 + e^{2i \sec^{-1}(a+bx)}} d \sec^{-1}(a+bx) + \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{4} i \sec^{-1}(a+bx)^4 \\
& \quad \downarrow \text{2620} \\
& a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) - \\
& 2i \left(\frac{3}{2} i \int \sec^{-1}(a+bx)^2 \log(1 + e^{2i \sec^{-1}(a+bx)}) d \sec^{-1}(a+bx) - \frac{1}{2} i \sec^{-1}(a+bx)^3 \log(1 + e^{2i \sec^{-1}(a+bx)}) \right) + \\
& \quad \frac{1}{4} i \sec^{-1}(a+bx)^4
\end{aligned}$$

↓ 3011

$$-2i \left(\frac{3}{2}i \left(\frac{1}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \int \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right. \right. \\ \left. \left. a \int \frac{\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^3}{1 - \frac{a}{a+bx}} d \sec^{-1}(a+bx) + \frac{1}{4}i \sec^{-1}(a+bx)^4 \right. \right.$$

↓ 5031

$$a \left(-i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^3}{-e^{i \sec^{-1}(a+bx)} a - \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - i \int \frac{e^{i \sec^{-1}(a+bx)} \sec^{-1}(a+bx)^3}{-e^{i \sec^{-1}(a+bx)} a + \sqrt{1-a^2} + 1} d \sec^{-1}(a+bx) - \right. \\ \left. 2i \left(\frac{3}{2}i \left(\frac{1}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \int \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right. \right. \right. \\ \left. \left. \left. \frac{1}{4}i \sec^{-1}(a+bx)^4 \right. \right. \right.$$

↓ 2620

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{3i \int \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) d \sec^{-1}(a+bx)}{a} \right) \right. \\ \left. 2i \left(\frac{3}{2}i \left(\frac{1}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \int \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right. \right. \right. \\ \left. \left. \left. \frac{1}{4}i \sec^{-1}(a+bx)^4 \right. \right. \right.$$

↓ 3011

$$a \left(-i \left(\frac{i \sec^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{3i \left(i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - 2i \int \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right)}{a} \right) \right. \\ \left. 2i \left(\frac{3}{2}i \left(\frac{1}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \int \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a+bx) \right. \right. \right. \\ \left. \left. \left. \frac{1}{4}i \sec^{-1}(a+bx)^4 \right. \right. \right.$$

↓ 7163

$$a \left(-i \left(\frac{i \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{3i \left(i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - 2i \left(\int \text{PolyLog} \right) \right)}{a} \right) \right. \\ \left. 2i \left(\frac{3}{2} i \left(\frac{1}{2} i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \left(\frac{1}{2} i \int \text{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) d \sec^{-1}(a + bx) - \frac{1}{4} i \sec^{-1}(a + bx)^4 \right) \right) \right) \right.$$

↓ 2720

$$a \left(-i \left(\frac{i \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{3i \left(i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - 2i \left(\int e^{-i \sec^{-1}(a+bx)} \text{PolyLog} \right) \right)}{a} \right) \right. \\ \left. 2i \left(\frac{3}{2} i \left(\frac{1}{2} i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \left(\frac{1}{4} \int e^{-2i \sec^{-1}(a+bx)} \text{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) de^{2i \sec^{-1}(a+bx)} - \frac{1}{4} i \sec^{-1}(a + bx)^4 \right) \right) \right) \right.$$

↓ 7143

$$a \left(-i \left(\frac{i \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a} - \frac{3i \left(i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) - 2i \left(\text{PolyLog} \right) \right)}{a} \right) \right. \\ \left. 2i \left(\frac{3}{2} i \left(\frac{1}{2} i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) - i \left(\frac{1}{4} \text{PolyLog} \left(4, -e^{2i \sec^{-1}(a+bx)} \right) - \frac{1}{2} i \sec^{-1}(a + bx) \right) - \frac{1}{4} i \sec^{-1}(a + bx)^4 \right) \right) \right.$$

input Int[ArcSec[a + b*x]^3/x,x]

output

```
(I/4)*ArcSec[a + b*x]^4 + a*((( -1/4*I)*ArcSec[a + b*x]^4)/a - I*((I*ArcSec
[a + b*x]^3*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/a - ((
3*I)*(I*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1
- a^2]]) - (2*I)*((-I)*ArcSec[a + b*x]*PolyLog[3, (a*E^(I*ArcSec[a + b*x]
)))/(1 - Sqrt[1 - a^2]]) + PolyLog[4, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1
- a^2])])))/a - I*((I*ArcSec[a + b*x]^3*Log[1 - (a*E^(I*ArcSec[a + b*x]
))/(1 + Sqrt[1 - a^2])])/a - ((3*I)*(I*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I
*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2]]) - (2*I)*((-I)*ArcSec[a + b*x]*Poly
Log[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2]]) + PolyLog[4, (a*E^(I
*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])))/a) - (2*I)*((-1/2*I)*ArcSec[a
+ b*x]^3*Log[1 + E^((2*I)*ArcSec[a + b*x])] + ((3*I)/2)*((I/2)*ArcSec[a +
b*x]^2*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] - I*((-1/2*I)*ArcSec[a + b*x
]*PolyLog[3, -E^((2*I)*ArcSec[a + b*x])] + PolyLog[4, -E^((2*I)*ArcSec[a +
b*x]])/4)))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^((n_)*((c_) + (d_)*(x_))^(m_)))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^((n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```


rule 3011 $\text{Int}[\text{Log}[1 + (e_.) * ((F_.)^{((c_.) * (a_.) + (b_.) * (x_)))})^{(n_.)}] * ((f_.) + (g_.) * (x_))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-f + g*x)^m * (\text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n]) / (b*c*n*\text{Log}[F]), x] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \text{Int}[(f + g*x)^{(m - 1)} * \text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4202 $\text{Int}[((c_.) + (d_.) * (x_))^{(m_.)} * \tan[(e_.) + (f_.) * (x_)], x_Symbol] \rightarrow \text{Simp}[I * ((c + d*x)^{(m + 1)} / (d*(m + 1))), x] - \text{Simp}[2*I \text{Int}[(c + d*x)^m * (E^{(2*I*(e + f*x))} / (1 + E^{(2*I*(e + f*x))}))], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0]$

rule 5031 $\text{Int}[(((e_.) + (f_.) * (x_))^{(m_.)} * \sin[(c_.) + (d_.) * (x_)] / (\cos[(c_.) + (d_.) * (x_)] * (b_.) + (a_))), x_Symbol] \rightarrow \text{Simp}[I * ((e + f*x)^{(m + 1)} / (b*f*(m + 1))), x] + (-\text{Simp}[I \text{Int}[(e + f*x)^m * (E^{(I*(c + d*x))} / (a - \text{Rt}[a^2 - b^2, 2] + b * E^{(I*(c + d*x))}))], x], x] - \text{Simp}[I \text{Int}[(e + f*x)^m * (E^{(I*(c + d*x))} / (a + \text{Rt}[a^2 - b^2, 2] + b * E^{(I*(c + d*x))}))], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{PosQ}[a^2 - b^2]$

rule 5041 $\text{Int}[(((e_.) + (f_.) * (x_))^{(m_.)} * \tan[(c_.) + (d_.) * (x_)]^{(n_.)}) / (\cos[(c_.) + (d_.) * (x_)] * (b_.) + (a_))), x_Symbol] \rightarrow \text{Simp}[1/a \text{Int}[(e + f*x)^m * \tan[c + d*x]^n, x], x] - \text{Simp}[b/a \text{Int}[(e + f*x)^m * \sin[c + d*x] * (\tan[c + d*x]^{(n - 1)} / (a + b * \cos[c + d*x]))], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[n, 0]$

rule 5062 $\text{Int}[(((e_.) + (f_.) * (x_))^{(m_.)} * (F_)[(c_.) + (d_.) * (x_)]^{(n_.)} * (G_)[(c_.) + (d_.) * (x_)]^{(p_.)}) / ((a_.) + (b_.) * \sec[(c_.) + (d_.) * (x_)]), x_Symbol] \rightarrow \text{Int}[(e + f*x)^m * \cos[c + d*x] * F[c + d*x]^n * (G[c + d*x]^p / (b + a * \cos[c + d*x])), x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{TrigQ}[F] \&\& \text{TrigQ}[G] \&\& \text{IntegersQ}[m, n, p]$

rule 5781 `Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

input `int(arcsec(b*x+a)^3/x,x)`

output `int(arcsec(b*x+a)^3/x,x)`

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

input `integrate(arcsec(b*x+a)^3/x,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^3/x, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{asec}^3(a + bx)}{x} dx$$

input `integrate(asec(b*x+a)**3/x,x)`

output `Integral(asec(a + b*x)**3/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

input `integrate(arcsec(b*x+a)^3/x,x, algorithm="maxima")`

output `integrate(arcsec(b*x + a)^3/x, x)`

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

input `integrate(arcsec(b*x+a)^3/x,x, algorithm="giac")`

output `integrate(arcsec(b*x + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)^3}{x} dx$$

input `int(acos(1/(a + b*x))^3/x,x)`output `int(acos(1/(a + b*x))^3/x, x)`**Reduce [F]**

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{asec}(bx + a)^3}{x} dx$$

input `int(asec(b*x+a)^3/x,x)`output `int(asec(a + b*x)**3/x,x)`

3.37 $\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx$

Optimal result	300
Mathematica [F(-1)]	301
Rubi [A] (verified)	301
Maple [F]	303
Fricas [F]	304
Sympy [F]	304
Maxima [F]	304
Giac [F]	305
Mupad [F(-1)]	305
Reduce [F]	306

Optimal result

Integrand size = 12, antiderivative size = 362

$$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx = -\frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6b \sec^{-1}(a+bx) \text{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \sec^{-1}(a+bx) \text{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6ib \text{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6ib \text{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

output

```
-b*arcsec(b*x+a)^3/a-arcsec(b*x+a)^3/x-3*I*b*arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+3*I*b*arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*I*b*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*I*b*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Mathematica [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \$Aborted$$

input

```
Integrate[ArcSec[a + b*x]^3/x^2,x]
```

output

```
$Aborted
```

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5781, 4926, 3042, 4679, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx$$

↓ 5781

$$b \int \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^3}{b^2 x^2} dx \sec^{-1}(a + bx)$$

$$\begin{aligned}
& \downarrow 4926 \\
& b \left(-3 \int -\frac{\sec^{-1}(a+bx)^2}{bx} d\sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^3}{bx} \right) \\
& \downarrow 3042 \\
& b \left(-3 \int \frac{\sec^{-1}(a+bx)^2}{a - \csc(\sec^{-1}(a+bx) + \frac{\pi}{2})} d\sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^3}{bx} \right) \\
& \downarrow 4679 \\
& b \left(-3 \int \left(\frac{\sec^{-1}(a+bx)^2}{a} + \frac{\sec^{-1}(a+bx)^2}{a \left(\frac{a}{a+bx} - 1 \right)} \right) d\sec^{-1}(a+bx) - \frac{\sec^{-1}(a+bx)^3}{bx} \right) \\
& \downarrow 2009 \\
& b \left(-\frac{\sec^{-1}(a+bx)^3}{bx} - 3 \left(\frac{2 \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a\sqrt{1-a^2}} - \frac{2 \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right)}{a\sqrt{1-a^2}} \right) \right)
\end{aligned}$$

input

```
Int[ArcSec[a + b*x]^3/x^2,x]
```

output

```
b*(-(ArcSec[a + b*x]^3/(b*x)) - 3*(ArcSec[a + b*x]^3/(3*a) + (I*ArcSec[a +
b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1
- a^2]) - (I*ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt
[1 - a^2])])/(a*Sqrt[1 - a^2]) + (2*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*Arc
Sec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (2*ArcSec[a + b*x
]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a
^2]) + ((2*I)*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(
a*Sqrt[1 - a^2]) - ((2*I)*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1
- a^2])])/(a*Sqrt[1 - a^2])))
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4679 `Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]`

rule 4926 `Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Simp[f*(m/(b*d*(n + 1))) Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

rule 5781 `Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[1/d^(m + 1) Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

input `int(arcsec(b*x+a)^3/x^2,x)`

output `int(arcsec(b*x+a)^3/x^2,x)`

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

input `integrate(arcsec(b*x+a)^3/x^2,x, algorithm="fricas")`

output `integral(arcsec(b*x + a)^3/x^2, x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asec}^3(a + bx)}{x^2} dx$$

input `integrate(asec(b*x+a)**3/x**2,x)`

output `Integral(asec(a + b*x)**3/x**2, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

input `integrate(arcsec(b*x+a)^3/x^2,x, algorithm="maxima")`

output

```
-1/4*(4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 3*arctan(sqrt(b*x
+ a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - 4*x*integrate
(3/4*((4*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x*log(b^2*x
^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) - 4*((b^3*x^3 +
3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 + (b^3*x^3 + 2*a*
b^2*x^2 + (a^2 - 1)*b*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x -
a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*s
qrt(b*x + a - 1)))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*
x^2), x))/x
```

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

input

```
integrate(arcsec(b*x+a)^3/x^2,x, algorithm="giac")
```

output

```
integrate(arcsec(b*x + a)^3/x^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)^3}{x^2} dx$$

input

```
int(arccos(1/(a + b*x))^3/x^2,x)
```

output

```
int(arccos(1/(a + b*x))^3/x^2, x)
```

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asec}(bx + a)^3}{x^2} dx$$

input `int(asec(b*x+a)^3/x^2,x)`

output `int(asec(a + b*x)**3/x**2,x)`

3.38 $\int x(a + b \sec^{-1}(c + dx^2)) dx$

Optimal result	307
Mathematica [C] (verified)	307
Rubi [A] (verified)	308
Maple [A] (verified)	309
Fricas [A] (verification not implemented)	310
Sympy [F(-1)]	310
Maxima [A] (verification not implemented)	310
Giac [A] (verification not implemented)	311
Mupad [B] (verification not implemented)	311
Reduce [F]	312

Optimal result

Integrand size = 14, antiderivative size = 58

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^2)^2}}\right)}{2d}$$

output `1/2*a*x^2+1/2*b*(d*x^2+c)*arcsec(d*x^2+c)/d-1/2*b*arctanh((1-1/(d*x^2+c)^2)^(1/2))/d`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.01 (sec) , antiderivative size = 516, normalized size of antiderivative = 8.90

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2 \sec^{-1}(c + dx^2) + \frac{b(c + dx^2) \sqrt{\frac{-1+c^2+2cdx^2+d^2x^4}{(c+dx^2)^2}} \left(\sqrt[4]{-1}(-i + \sqrt{-1 + c^2}) \sqrt{2i - ic^2 + 2\sqrt{-1 + c^2}} \arctan\left(\frac{(-1)^{3/4} \sqrt{2i - ic^2}}{c\sqrt{-1+c^2} - c\sqrt{-1+c^2}}\right) \right)}{2d}$$

input `Integrate[x*(a + b*ArcSec[c + d*x^2]),x]`

output
$$\begin{aligned} & (a*x^2)/2 + (b*x^2*ArcSec[c + d*x^2])/2 + (b*(c + d*x^2)*Sqrt[(-1 + c^2 + \\ & 2*c*d*x^2 + d^2*x^4)/(c + d*x^2)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + c^2])*Sqrt \\ & [2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*c^2 + 2*S \\ & qrt[-1 + c^2]]*d*x^2)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^ \\ & 2*x^4])) + (-1)^(3/4)*(I + Sqrt[-1 + c^2])*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + \\ & c^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^2)/(c* \\ & Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4])) + c*(c*ArcTan[(S \\ & qrt[-1 + c^2]*d^2*x^4)/(c^4 + c^3*d*x^2 + d^2*x^4 - c^2*(1 + Sqrt[-1 + c^2 \\ &]*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4]))] - Log[Sqrt[-1 + c^2] - d*x^2 - S \\ & qrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4]] + Log[d^2*(Sqrt[-1 + c^2] + d*x^2 - S \\ & qrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4])))])/(2*c*d*Sqrt[-1 + c^2 + 2*c*d*x^2 \\ & + d^2*x^4]) \end{aligned}$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {7266, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(a + b \sec^{-1}(c + dx^2)) dx \\ & \quad \downarrow \text{7266} \\ & \frac{1}{2} \int (a + b \sec^{-1}(dx^2 + c)) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \left(ax^2 - \frac{\text{barctanh}\left(\sqrt{1 - \frac{1}{(c+dx^2)^2}}\right)}{d} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{d} \right) \end{aligned}$$

input `Int[x*(a + b*ArcSec[c + d*x^2]),x]`

output $(a*x^2 + (b*(c + d*x^2)*ArcSec[c + d*x^2])/d - (b*ArcTanh[Sqrt[1 - (c + d*x^2)^{-2}]])/d)/2$

Defintions of rubi rules used

rule 2009 $Int[u_, x_Symbol] \rightarrow Simp[IntSum[u, x], x] /; SumQ[u]$

rule 7266 $Int[(u_)*(x_)^{(m_.)}, x_Symbol] \rightarrow Simp[1/(m + 1) Subst[Int[SubstFor[x^{(m + 1)}, u, x], x], x, x^{(m + 1)}], x] /; FreeQ[m, x] \&\& NeQ[m, -1] \&\& FunctionOfQ[x^{(m + 1)}, u, x]$

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{ax^2}{2} + \frac{b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	64
derivativedivides	$\frac{(dx^2+c)a+b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	68
default	$\frac{(dx^2+c)a+b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	68

input $int(x*(a+b*arcsec(d*x^2+c)),x,method=_RETURNVERBOSE)$

output $1/2*a*x^2+1/2*b/d*((d*x^2+c)*arcsec(d*x^2+c)-ln(d*x^2+c+(d*x^2+c)*(1-1/(d*x^2+c)^2)^(1/2)))$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x(a + b \sec^{-1}(c + dx^2)) dx$$

$$= \frac{bdx^2 \operatorname{arcsec}(dx^2 + c) + adx^2 + 2bc \arctan(-dx^2 - c + \sqrt{d^2x^4 + 2cdx^2 + c^2 - 1}) + b \log(-dx^2 - c + \sqrt{d^2x^4 + 2cdx^2 + c^2 - 1})}{2d}$$

input `integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="fricas")`

output `1/2*(b*d*x^2*arcsec(d*x^2 + c) + a*d*x^2 + 2*b*c*arctan(-d*x^2 - c + sqrt(d^2*x^4 + 2*c*d*x^2 + c^2 - 1)) + b*log(-d*x^2 - c + sqrt(d^2*x^4 + 2*c*d*x^2 + c^2 - 1)))/d`

Sympy [F(-1)]

Timed out.

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \text{Timed out}$$

input `integrate(x*(a+b*asec(d*x**2+c)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{1}{2} ax^2$$

$$+ \frac{\left(2(dx^2 + c) \operatorname{arcsec}(dx^2 + c) - \log\left(\sqrt{-\frac{1}{(dx^2+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^2+c)^2} + 1} + 1\right)\right) b}{4d}$$

input `integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="maxima")`

output

$$\frac{1}{2}ax^2 + \frac{1}{4}b(2\sqrt{dx^2+c}\operatorname{arcsec}(\sqrt{dx^2+c}) - \log(\sqrt{-1/(dx^2+c)^2+1} + 1) + \log(-\sqrt{-1/(dx^2+c)^2+1} + 1))b/d$$

Giac [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{1}{2}ax^2 + \frac{1}{4}bd \left(\frac{2(dx^2+c)\arccos\left(-\frac{1}{(dx^2+c)\left(\frac{c}{dx^2+c}-1\right)-c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^2+c)^2}+1}+1\right) - \log\left(-\sqrt{-\frac{1}{(dx^2+c)^2}+1}\right)}{d^2} \right)$$

input

```
integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="giac")
```

output

$$\frac{1}{2}ax^2 + \frac{1}{4}bd(2\sqrt{dx^2+c}\arccos(-1/((dx^2+c)*(c/(dx^2+c)-1)-c))/d^2 - (\log(\sqrt{-1/(dx^2+c)^2+1} + 1) - \log(-\sqrt{-1/(dx^2+c)^2+1} + 1))/d^2)$$

Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1-\frac{1}{(dx^2+c)^2}}}\right)}{2d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^2+c}\right) (dx^2+c)}{2d}$$

input

```
int(x*(a + b*acos(1/(c + d*x^2))),x)
```

output

$$(a*x^2)/2 - (b*atanh(1/(1 - 1/(c + d*x^2)^2)^(1/2)))/(2*d) + (b*acos(1/(c + d*x^2))*(c + d*x^2))/(2*d)$$

Reduce [F]

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \left(\int a \sec(dx^2 + c) x dx \right) b + \frac{ax^2}{2}$$

input `int(x*(a+b*asec(d*x^2+c)),x)`

output `(2*int(asec(c + d*x**2)*x,x)*b + a*x**2)/2`

3.39 $\int x^2(a + b \sec^{-1}(c + dx^3)) dx$

Optimal result	313
Mathematica [C] (verified)	313
Rubi [A] (verified)	314
Maple [A] (verified)	315
Fricas [A] (verification not implemented)	316
Sympy [F(-1)]	316
Maxima [A] (verification not implemented)	316
Giac [A] (verification not implemented)	317
Mupad [B] (verification not implemented)	317
Reduce [F]	318

Optimal result

Integrand size = 16, antiderivative size = 58

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d} - \frac{b \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^3)^2}}\right)}{3d}$$

output `1/3*a*x^3+1/3*b*(d*x^3+c)*arcsec(d*x^3+c)/d-1/3*b*arctanh((1-1/(d*x^3+c)^2)^(1/2))/d`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 516, normalized size of antiderivative = 8.90

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} + \frac{1}{3}bx^3 \sec^{-1}(c + dx^3) + \frac{b(c + dx^3) \sqrt{\frac{-1+c^2+2cdx^3+d^2x^6}{(c+dx^3)^2}} \left(\sqrt[4]{-1+c^2} (-i + \sqrt{-1+c^2}) \sqrt{2i-ic^2+2\sqrt{-1+c^2}} \arctan\left(\frac{(-1)^{3/4} \sqrt{2i-ic^2}}{c\sqrt{-1+c^2}-c\sqrt{-1+c^2}}\right) \right)}{3d}$$

input `Integrate[x^2*(a + b*ArcSec[c + d*x^3]),x]`

output
$$\begin{aligned} & (a*x^3)/3 + (b*x^3*ArcSec[c + d*x^3])/3 + (b*(c + d*x^3)*Sqrt[(-1 + c^2 + \\ & 2*c*d*x^3 + d^2*x^6)/(c + d*x^3)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + c^2])*Sqrt \\ & [2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*c^2 + 2*S \\ & qrt[-1 + c^2]]*d*x^3)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^ \\ & 2*x^6])) + (-1)^(3/4)*(I + Sqrt[-1 + c^2])*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + \\ & c^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^3)/(c* \\ & Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6])] + c*(c*ArcTan[(S \\ & qrt[-1 + c^2]*d^2*x^6)/(c^4 + c^3*d*x^3 + d^2*x^6 - c^2*(1 + Sqrt[-1 + c^2 \\ &]*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]))] - Log[Sqrt[-1 + c^2] - d*x^3 - S \\ & qrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]] + Log[d^2*(Sqrt[-1 + c^2] + d*x^3 - S \\ & qrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6])))]))/(3*c*d*Sqrt[-1 + c^2 + 2*c*d*x^3 \\ & + d^2*x^6]) \end{aligned}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {7266, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2(a + b \sec^{-1}(c + dx^3)) dx \\ & \quad \downarrow \text{7266} \\ & \frac{1}{3} \int (a + b \sec^{-1}(dx^3 + c)) dx^3 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{3} \left(ax^3 - \frac{\text{barctanh}\left(\sqrt{1 - \frac{1}{(c+dx^3)^2}}\right)}{d} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{d} \right) \end{aligned}$$

input `Int[x^2*(a + b*ArcSec[c + d*x^3]),x]`

output $(a*x^3 + (b*(c + d*x^3)*ArcSec[c + d*x^3])/d - (b*ArcTanh[Sqrt[1 - (c + d*x^3)^{-2}]])/d)/3$

Defintions of rubi rules used

rule 2009 $Int[u_, x_Symbol] \rightarrow Simp[IntSum[u, x], x] /; SumQ[u]$

rule 7266 $Int[(u_)*(x_)^(m_.), x_Symbol] \rightarrow Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] \&\& NeQ[m, -1] \&\& FunctionOfQ[x^(m + 1), u, x]$

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{ax^3}{3} + \frac{b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	64
derivativedivides	$\frac{(dx^3+c)a+b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	68
default	$\frac{(dx^3+c)a+b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	68

input $\text{int}(x^2*(a+b*\operatorname{arcsec}(d*x^3+c)), x, \text{method}=_RETURNVERBOSE)$

output $1/3*a*x^3+1/3*b/d*((d*x^3+c)*\operatorname{arcsec}(d*x^3+c)-\ln(d*x^3+c+(d*x^3+c)*(1-1/(d*x^3+c)^2))^{(1/2)})$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx$$

$$= \frac{bdx^3 \operatorname{arcsec}(dx^3 + c) + adx^3 + 2bc \arctan(-dx^3 - c + \sqrt{d^2x^6 + 2cdx^3 + c^2 - 1}) + b \log(-dx^3 - c + \sqrt{d^2x^6 + 2cdx^3 + c^2 - 1})}{3d}$$

input `integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="fricas")`

output `1/3*(b*d*x^3*arcsec(d*x^3 + c) + a*d*x^3 + 2*b*c*arctan(-d*x^3 - c + sqrt(d^2*x^6 + 2*c*d*x^3 + c^2 - 1)) + b*log(-d*x^3 - c + sqrt(d^2*x^6 + 2*c*d*x^3 + c^2 - 1)))/d`

Sympy [F(-1)]

Timed out.

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \text{Timed out}$$

input `integrate(x**2*(a+b*asec(d*x**3+c)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \frac{1}{3} ax^3$$

$$+ \frac{\left(2(dx^3 + c) \operatorname{arcsec}(dx^3 + c) - \log\left(\sqrt{-\frac{1}{(dx^3+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^3+c)^2} + 1} + 1\right)\right) b}{6d}$$

input `integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="maxima")`

output $\frac{1}{3}ax^3 + \frac{1}{6}(2(dx^3 + c)\operatorname{arcsec}(dx^3 + c) - \log(\sqrt{-1/(dx^3 + c)^2 + 1} + 1) + \log(-\sqrt{-1/(dx^3 + c)^2 + 1} + 1))b/d$

Giac [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{1}{3}ax^3 + \frac{1}{6}bd \left(\frac{2(dx^3 + c) \arccos\left(-\frac{1}{(dx^3 + c)\left(\frac{c}{dx^3 + c} - 1\right) - c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^3 + c)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(dx^3 + c)^2} + 1}\right)}{d^2} \right)$$

input `integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="giac")`

output $\frac{1}{3}ax^3 + \frac{1}{6}bd*(2*(dx^3 + c)*\arccos(-1/((dx^3 + c)*(c/(dx^3 + c) - 1) - c))/d^2 - (\log(\sqrt{-1/(dx^3 + c)^2 + 1} + 1) - \log(-\sqrt{-1/(dx^3 + c)^2 + 1} + 1))/d^2)$

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(dx^3 + c)^2}}}\right)}{3d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^3 + c}\right) (dx^3 + c)}{3d}$$

input `int(x^2*(a + b*acos(1/(c + d*x^3))),x)`

output $(a*x^3)/3 - (b*\operatorname{atanh}(1/(1 - 1/(c + d*x^3)^2)^{(1/2)}))/(3*d) + (b*\operatorname{acos}(1/(c + d*x^3))*(c + d*x^3))/(3*d)$

Reduce [F]

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \left(\int a \sec(dx^3 + c) x^2 dx \right) b + \frac{ax^3}{3}$$

input `int(x^2*(a+b*asec(d*x^3+c)),x)`

output `(3*int(asec(c + d*x**3)*x**2,x)*b + a*x**3)/3`

3.40 $\int x^3(a + b \sec^{-1}(c + dx^4)) dx$

Optimal result	319
Mathematica [B] (verified)	319
Rubi [A] (verified)	320
Maple [A] (verified)	321
Fricas [A] (verification not implemented)	321
Sympy [F(-1)]	322
Maxima [A] (verification not implemented)	322
Giac [A] (verification not implemented)	323
Mupad [B] (verification not implemented)	323
Reduce [F]	324

Optimal result

Integrand size = 16, antiderivative size = 58

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} - \frac{b \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^4)^2}}\right)}{4d}$$

output

`1/4*a*x^4+1/4*b*(d*x^4+c)*arcsec(d*x^4+c)/d-1/4*b*arctanh((1-1/(d*x^4+c)^2)^(1/2))/d`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 137 vs. 2(58) = 116.

Time = 0.21 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.36

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} - \frac{b\sqrt{-1 + (c + dx^4)^2} \left(-\log\left(1 - \frac{c+dx^4}{\sqrt{-1+(c+dx^4)^2}}\right) + \log\left(1 + \frac{c+dx^4}{\sqrt{-1+(c+dx^4)^2}}\right) \right)}{8d(c + dx^4) \sqrt{1 - \frac{1}{(c+dx^4)^2}}}$$

input `Integrate[x^3*(a + b*ArcSec[c + d*x^4]),x]`

output `(a*x^4)/4 + (b*(c + d*x^4)*ArcSec[c + d*x^4])/(4*d) - (b*Sqrt[-1 + (c + d*x^4)^2]*(-Log[1 - (c + d*x^4)/Sqrt[-1 + (c + d*x^4)^2]] + Log[1 + (c + d*x^4)/Sqrt[-1 + (c + d*x^4)^2]]))/(8*d*(c + d*x^4)*Sqrt[1 - (c + d*x^4)^(-2)])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {7266, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx$$

$$\downarrow 7266$$

$$\frac{1}{4} \int (a + b \sec^{-1}(dx^4 + c)) dx^4$$

$$\downarrow 2009$$

$$\frac{1}{4} \left(ax^4 - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^4)^2}}\right)}{d} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{d} \right)$$

input `Int[x^3*(a + b*ArcSec[c + d*x^4]),x]`

output `(a*x^4 + (b*(c + d*x^4)*ArcSec[c + d*x^4])/d - (b*ArcTanh[Sqrt[1 - (c + d*x^4)^(-2)]])/d)/4`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7266 Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{ax^4}{4} + \frac{b \left((dx^4+c) \operatorname{arcsec}(dx^4+c) - \ln \left(dx^4+c+(dx^4+c) \sqrt{1-\frac{1}{(dx^4+c)^2}} \right) \right)}{4d}$	64
derivativedivides	$\frac{(dx^4+c)a+b \left((dx^4+c) \operatorname{arcsec}(dx^4+c) - \ln \left(dx^4+c+(dx^4+c) \sqrt{1-\frac{1}{(dx^4+c)^2}} \right) \right)}{4d}$	68
default	$\frac{(dx^4+c)a+b \left((dx^4+c) \operatorname{arcsec}(dx^4+c) - \ln \left(dx^4+c+(dx^4+c) \sqrt{1-\frac{1}{(dx^4+c)^2}} \right) \right)}{4d}$	68

```
input int(x^3*(a+b*arcsec(d*x^4+c)),x,method=_RETURNVERBOSE)
```

```
output 1/4*a*x^4+1/4*b/d*((d*x^4+c)*arcsec(d*x^4+c)-ln(d*x^4+c+(d*x^4+c)*(1-1/(d*x^4+c)^2)^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{bdx^4 \operatorname{arcsec}(dx^4 + c) + adx^4 + 2bc \arctan(-dx^4 - c + \sqrt{d^2x^8 + 2cdx^4 + c^2 - 1}) + b \log(-dx^4 - c + \sqrt{d^2x^8 + 2cdx^4 + c^2 - 1})}{4d}$$

input `integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="fricas")`

output $\frac{1}{4}*(b*d*x^4*arcsec(d*x^4 + c) + a*d*x^4 + 2*b*c*arctan(-d*x^4 - c + \sqrt{d^2*x^8 + 2*c*d*x^4 + c^2 - 1})) + b*\log(-d*x^4 - c + \sqrt{d^2*x^8 + 2*c*d*x^4 + c^2 - 1}))/d$

Sympy [F(-1)]

Timed out.

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \text{Timed out}$$

input `integrate(x**3*(a+b*asec(d*x**4+c)),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{1}{4} ax^4 + \frac{\left(2(dx^4 + c) \operatorname{arcsec}(dx^4 + c) - \log\left(\sqrt{-\frac{1}{(dx^4+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^4+c)^2} + 1} + 1\right)\right)b}{8d}$$

input `integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="maxima")`

output $\frac{1}{4}*a*x^4 + \frac{1}{8}*(2*(d*x^4 + c)*arcsec(d*x^4 + c) - \log(\sqrt{-1/(d*x^4 + c)^2 + 1} + 1) + \log(-\sqrt{-1/(d*x^4 + c)^2 + 1} + 1))*b/d$

Giac [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{1}{4} ax^4 + \frac{1}{8} bd \left(\frac{2(dx^4 + c) \arccos\left(-\frac{1}{(dx^4+c)\left(\frac{c}{dx^4+c}-1\right)-c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^4+c)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(dx^4+c)^2} + 1}\right)}{d^2} \right)$$

input `integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="giac")`output `1/4*a*x^4 + 1/8*b*d*(2*(d*x^4 + c)*arccos(-1/((d*x^4 + c)*(c/(d*x^4 + c) - 1) - c))/d^2 - (log(sqrt(-1/(d*x^4 + c)^2 + 1) + 1) - log(-sqrt(-1/(d*x^4 + c)^2 + 1) + 1))/d^2)`**Mupad [B] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(dx^4+c)^2}}}\right)}{4d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^4+c}\right) (dx^4 + c)}{4d}$$

input `int(x^3*(a + b*acos(1/(c + d*x^4))),x)`output `(a*x^4)/4 - (b*atanh(1/(1 - 1/(c + d*x^4)^2)^(1/2)))/(4*d) + (b*acos(1/(c + d*x^4))*(c + d*x^4))/(4*d)`

Reduce [F]

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \left(\int a \sec(dx^4 + c) x^3 dx \right) b + \frac{ax^4}{4}$$

input `int(x^3*(a+b*asec(d*x^4+c)),x)`

output `(4*int(asec(c + d*x**4)*x**3,x)*b + a*x**4)/4`

3.41 $\int x^{-1+n} \sec^{-1}(a + bx^n) dx$

Optimal result	325
Mathematica [B] (verified)	325
Rubi [A] (warning: unable to verify)	326
Maple [F]	328
Fricas [A] (verification not implemented)	328
Sympy [F(-1)]	329
Maxima [A] (verification not implemented)	329
Giac [A] (verification not implemented)	329
Mupad [B] (verification not implemented)	330
Reduce [F]	330

Optimal result

Integrand size = 14, antiderivative size = 49

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

output

```
(a+b*x^n)*arcsec(a+b*x^n)/b/n-arctanh((1-1/(a+b*x^n)^2)^(1/2))/b/n
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 130 vs. 2(49) = 98.

Time = 0.21 (sec) , antiderivative size = 130, normalized size of antiderivative = 2.65

$$\begin{aligned} & \int x^{-1+n} \sec^{-1}(a + bx^n) dx \\ &= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} \\ & \quad - \frac{\sqrt{-1 + (a + bx^n)^2} \left(-\log\left(1 - \frac{a+bx^n}{\sqrt{-1+(a+bx^n)^2}}\right) + \log\left(1 + \frac{a+bx^n}{\sqrt{-1+(a+bx^n)^2}}\right) \right)}{2bn(a + bx^n) \sqrt{1 - \frac{1}{(a+bx^n)^2}}} \end{aligned}$$

input `Integrate[x^(-1 + n)*ArcSec[a + b*x^n], x]`

output
$$\frac{((a + b*x^n)*ArcSec[a + b*x^n])/(b*n) - (Sqrt[-1 + (a + b*x^n)^2]*(-Log[1 - (a + b*x^n)/Sqrt[-1 + (a + b*x^n)^2]] + Log[1 + (a + b*x^n)/Sqrt[-1 + (a + b*x^n)^2]]))/(2*b*n*(a + b*x^n)*Sqrt[1 - (a + b*x^n)^(-2)])}$$

Rubi [A] (warning: unable to verify)

Time = 0.32 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.92, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {7266, 5773, 895, 798, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{n-1} \sec^{-1}(a + bx^n) dx \\ & \quad \downarrow \text{7266} \\ & \frac{\int \sec^{-1}(bx^n + a) dx^n}{n} \\ & \quad \downarrow \text{5773} \\ & \frac{\frac{(a+bx^n) \sec^{-1}(a+bx^n)}{b} - \int \frac{1}{(bx^n+a)\sqrt{1-\frac{1}{(bx^n+a)^2}}} dx^n}{n} \\ & \quad \downarrow \text{895} \\ & \frac{\frac{(a+bx^n) \sec^{-1}(a+bx^n)}{b} - \frac{\int \frac{x^{-n}}{\sqrt{1-x^{-2n}}} d(bx^n+a)}{b}}{n} \\ & \quad \downarrow \text{798} \\ & \frac{\frac{\int \frac{x^{-n}}{\sqrt{-bx^n-a+1}} dx^{-2n}}{2b} + \frac{(a+bx^n) \sec^{-1}(a+bx^n)}{b}}{n} \\ & \quad \downarrow \text{73} \\ & \frac{\frac{(a+bx^n) \sec^{-1}(a+bx^n)}{b} - \frac{\int \frac{1}{1-x^{2n}} d\sqrt{-bx^n-a+1}}{b}}{n} \end{aligned}$$

$$\frac{(a+bx^n) \sec^{-1}(a+bx^n)}{b} - \frac{\operatorname{arctanh}(\sqrt{-a-bx^n+1})}{b}$$

↓ 219

$$\frac{\quad}{n}$$

input `Int[x^(-1 + n)*ArcSec[a + b*x^n], x]`

output `((a + b*x^n)*ArcSec[a + b*x^n])/b - ArcTanh[Sqrt[1 - a - b*x^n]]/b/n`

Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
 [Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
 b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 895 `Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_.), x_Symbol] := Simp[u^m/(Coeff
 icient[v, x, 1]*v^m) Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[
 {a, b, m, n, p}, x] && LinearPairQ[u, v, x]`

rule 5773 `Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]
 /d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]`

rule 7266

```
Int[(u_)*(x_)^(m_.), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m
+ 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && Function
OfQ[x^(m + 1), u, x]
```

Maple [F]

$$\int x^{-1+n} \operatorname{arcsec}(a + bx^n) dx$$

input

```
int(x^(-1+n)*arcsec(a+b*x^n),x)
```

output

```
int(x^(-1+n)*arcsec(a+b*x^n),x)
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.88

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{bx^n \operatorname{arcsec}(bx^n + a) + 2a \arctan(-bx^n - a + \sqrt{b^2x^{2n} + 2abx^n + a^2 - 1}) + \log(-bx^n - a + \sqrt{b^2x^{2n} + 2abx^n + a^2 - 1})}{bn}$$

input

```
integrate(x^(-1+n)*arcsec(a+b*x^n),x, algorithm="fricas")
```

output

```
(b*x^n*arcsec(b*x^n + a) + 2*a*arctan(-b*x^n - a + sqrt(b^2*x^(2*n) + 2*a*
b*x^n + a^2 - 1)) + log(-b*x^n - a + sqrt(b^2*x^(2*n) + 2*a*b*x^n + a^2 -
1)))/(b*n)
```

Sympy [F(-1)]

Timed out.

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = \text{Timed out}$$

input `integrate(x**(-1+n)*asec(a+b*x**n),x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.35

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{2(bx^n + a) \operatorname{arcsec}(bx^n + a) - \log\left(\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right)}{2bn}$$

input `integrate(x^(-1+n)*arcsec(a+b*x^n),x, algorithm="maxima")`output `1/2*(2*(b*x^n + a)*arcsec(b*x^n + a) - log(sqrt(-1/(b*x^n + a)^2 + 1) + 1) + log(-sqrt(-1/(b*x^n + a)^2 + 1) + 1))/(b*n)`**Giac [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.53

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{b \left(\frac{2(bx^n+a) \arccos\left(\frac{1}{bx^n+a}\right)}{b^2} - \frac{\log\left(\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right)}{b^2} \right)}{2n}$$

input `integrate(x^(-1+n)*arcsec(a+b*x^n),x, algorithm="giac")`

output $\frac{1}{2} b (2 (b x^n + a) \arccos(1/(b x^n + a)) / b^2 - (\log(\sqrt{-1/(b x^n + a)^2 + 1}) + 1) - \log(-\sqrt{-1/(b x^n + a)^2 + 1}) / b^2) / n$

Mupad [B] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int x^{-1+n} \sec^{-1}(a + b x^n) dx = -\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(a + b x^n)^2}}}\right) - \arccos\left(\frac{1}{a + b x^n}\right) (a + b x^n)}{b n}$$

input `int(x^(n - 1)*acos(1/(a + b*x^n)),x)`

output $-(\operatorname{atanh}(1/(1 - 1/(a + b x^n)^2)^{1/2}) - \arccos(1/(a + b x^n)) * (a + b x^n)) / (b * n)$

Reduce [F]

$$\int x^{-1+n} \sec^{-1}(a + b x^n) dx = \int \frac{x^n \operatorname{asec}(x^n b + a)}{x} dx$$

input `int(x^(-1+n)*asec(a+b*x^n),x)`

output `int((x**n*asec(x**n*b + a))/x,x)`

3.42 $\int \sec^{-1}(ce^{a+bx}) dx$

Optimal result	331
Mathematica [B] (verified)	331
Rubi [A] (warning: unable to verify)	332
Maple [A] (verified)	335
Fricas [F(-2)]	335
Sympy [F]	336
Maxima [F]	336
Giac [F]	337
Mupad [F(-1)]	337
Reduce [F]	337

Optimal result

Integrand size = 10, antiderivative size = 85

$$\int \sec^{-1}(ce^{a+bx}) dx = \frac{i \sec^{-1}(ce^{a+bx})^2}{2b} - \frac{\sec^{-1}(ce^{a+bx}) \log(1 + e^{2i \sec^{-1}(ce^{a+bx})})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i \sec^{-1}(ce^{a+bx})})}{2b}$$

output

```
1/2*I*arcsec(c*exp(b*x+a))^2/b-arcsec(c*exp(b*x+a))*ln(1+(1/c/(exp(1)^(b*x+a)))+I*(1-1/c^2/(exp(1)^(b*x+a))^2)^(1/2))/b+1/2*I*polylog(2,-(1/c/(exp(1)^(b*x+a)))+I*(1-1/c^2/(exp(1)^(b*x+a))^2)^(1/2))/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 280 vs. 2(85) = 170.

Time = 0.81 (sec) , antiderivative size = 280, normalized size of antiderivative = 3.29

$$\int \sec^{-1}(ce^{a+bx}) dx = x \sec^{-1}(ce^{a+bx}) - \frac{e^{-a-bx} \left(4\sqrt{-1 + c^2 e^{2(a+bx)}} \arctan \left(\sqrt{-1 + c^2 e^{2(a+bx)}} \right) (2bx - \log(c^2 e^{2(a+bx)})) + \sqrt{1 - c^2 e^{2(a+bx)}} \left(\log^2 \right) \right)}{\dots}$$

input `Integrate[ArcSec[c*E^(a + b*x)],x]`

output `x*ArcSec[c*E^(a + b*x)] - (E^(-a - b*x)*(4*Sqrt[-1 + c^2*E^(2*(a + b*x))]*ArcTan[Sqrt[-1 + c^2*E^(2*(a + b*x))]]*(2*b*x - Log[c^2*E^(2*(a + b*x))]) + Sqrt[1 - c^2*E^(2*(a + b*x))]*(Log[c^2*E^(2*(a + b*x))]^2 - 4*Log[c^2*E^(2*(a + b*x))]*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2] + 2*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2]^2) - 4*Sqrt[1 - c^2*E^(2*(a + b*x))]*PolyLog[2, (1 - Sqrt[1 - c^2*E^(2*(a + b*x))])/2])/(8*b*c*Sqrt[1 - 1/(c^2*E^(2*(a + b*x))])])`

Rubi [A] (warning: unable to verify)

Time = 0.47 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {2720, 5741, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sec^{-1}(ce^{a+bx}) dx \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int e^{-a-bx} \sec^{-1}(ce^{a+bx}) de^{a+bx}}{b} \\
 & \quad \downarrow \text{5741} \\
 & \frac{\int e^{-a-bx} \arccos\left(\frac{e^{-a-bx}}{c}\right) de^{-a-bx}}{b} \\
 & \quad \downarrow \text{5137} \\
 & \frac{\int ce^{a+bx} \sqrt{1 - \frac{e^{-2a-2bx}}{c^2}} \arccos\left(\frac{e^{-a-bx}}{c}\right) d \arccos\left(\frac{e^{-a-bx}}{c}\right)}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \arccos\left(\frac{e^{-a-bx}}{c}\right) \tan\left(\arccos\left(\frac{e^{-a-bx}}{c}\right)\right) d \arccos\left(\frac{e^{-a-bx}}{c}\right)}{b}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 4202 \\
\frac{\frac{1}{2}ie^{2a+2bx} - 2i \int \frac{e^{a+bx+2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}}{1+e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}} d \arccos\left(\frac{e^{-a-bx}}{c}\right)}{b} \\
\downarrow 2620 \\
\frac{\frac{1}{2}ie^{2a+2bx} - 2i \left(\frac{1}{2}i \int \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) d \arccos\left(\frac{e^{-a-bx}}{c}\right) - \frac{1}{2}i \arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) \right)}{b} \\
\downarrow 2715 \\
\frac{\frac{1}{2}ie^{2a+2bx} - 2i \left(\frac{1}{4} \int e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)} \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) de^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)} - \frac{1}{2}i \arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) \right)}{b} \\
\downarrow 2838 \\
\frac{\frac{1}{2}ie^{2a+2bx} - 2i \left(-\frac{1}{4} \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) - \frac{1}{2}i \arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right) \right)}{b}
\end{array}$$

input `Int[ArcSec[c*E^(a + b*x)], x]`

output `((I/2)*E^(2*a + 2*b*x) - (2*I)*((-1/2*I)*ArcCos[E^(-a - b*x)/c]*Log[1 + E^((2*I)*ArcCos[E^(-a - b*x)/c]]) - PolyLog[2, -E^((2*I)*ArcCos[E^(-a - b*x)/c]])/4)/b`

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
-> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)
*(F_)[v_]) /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4202 `Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] :> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(
e + f*x))/(1 + E^(2*I*(e + f*x))))], x], x] /; FreeQ[{c, d, e, f}, x] && IGt
Q[m, 0]`

rule 5137 `Int[((a_) + ArcCos[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] :> -Subst[Int[
(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0
]`

rule 5741 `Int[((a_) + ArcSec[(c_)*(x_)]*(b_))/(x_), x_Symbol] :> -Subst[Int[(a + b
*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.31

method	result
derivativedivides	$\frac{i \operatorname{arcsec}\left(\frac{e^{bx+a}c}{2}\right)^2 - \operatorname{arcsec}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{b} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)\right)}{2}$
default	$\frac{i \operatorname{arcsec}\left(\frac{e^{bx+a}c}{2}\right)^2 - \operatorname{arcsec}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{b} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)\right)}{2}$

input `int(arcsec(exp(b*x+a)*c),x,method=_RETURNVERBOSE)`

output `1/b*(1/2*I*arcsec(exp(b*x+a)*c)^2-arcsec(exp(b*x+a)*c)*ln(1+(1/exp(b*x+a)/c+I*(1-1/exp(b*x+a)^2/c^2)^(1/2))^2)+1/2*I*polylog(2,-(1/exp(b*x+a)/c+I*(1-1/exp(b*x+a)^2/c^2)^(1/2))^2))`

Fricas [F(-2)]

Exception generated.

$$\int \sec^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

input `integrate(arcsec(c*exp(b*x+a)),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

Sympy [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{asec}(ce^{a+bx}) dx$$

input `integrate(asec(c*exp(b*x+a)),x)`

output `Integral(asec(c*exp(a + b*x)), x)`

Maxima [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsec}(ce^{(bx+a)}) dx$$

input `integrate(arcsec(c*exp(b*x+a)),x, algorithm="maxima")`

output `-1/2*(2*b^2*c^2*integrate(x*e^(2*b*x + 2*a + 1/2*log(c*e^(b*x + a) + 1) + 1/2*log(c*e^(b*x + a) - 1))/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(log(c*e^(b*x + a) + 1) + log(c*e^(b*x + a) - 1)) - 1), x) + 2*I*b^2*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(log(c*e^(b*x + a) + 1) + log(c*e^(b*x + a) - 1)) - 1), x) - I*b^2*x^2 - 2*b*x*arctan(sqrt(c*e^(b*x + a) + 1)*sqrt(c*e^(b*x + a) - 1)) + I*b*x*log(c^2*e^(2*b*x + 2*a)) - I*b*x*log(c*e^(b*x + a) + 1) - I*b*x*log(-c*e^(b*x + a) + 1) - 2*(I*a*b + I*b*log(c))*x - I*dilog(c*e^(b*x + a)) - I*dilog(-c*e^(b*x + a)))/b`

Giac [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsec}(ce^{(bx+a)}) dx$$

input `integrate(arcsec(c*exp(b*x+a)),x, algorithm="giac")`

output `integrate(arcsec(c*e^(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{acos}\left(\frac{e^{-a-bx}}{c}\right) dx$$

input `int(acos(exp(- a - b*x)/c),x)`

output `int(acos(exp(- a - b*x)/c), x)`

Reduce [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{asec}(e^{bx+a}c) dx$$

input `int(asec(c*exp(b*x+a)),x)`

output `int(asec(e**(a + b*x)*c),x)`

3.43 $\int e^{\sec^{-1}(ax)} x^2 dx$

Optimal result	338
Mathematica [A] (verified)	339
Rubi [A] (verified)	339
Maple [F]	341
Fricas [F]	341
Sympy [F]	341
Maxima [F]	342
Giac [F]	342
Mupad [F(-1)]	342
Reduce [F]	343

Optimal result

Integrand size = 10, antiderivative size = 99

$$\int e^{\sec^{-1}(ax)} x^2 dx$$

$$= -\frac{\left(\frac{12}{5} + \frac{4i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 3, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

$$+ \frac{\left(\frac{24}{5} + \frac{8i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

output

```
(-12/5-4/5*I)*exp((1+3*I)*arcsec(a*x))*hypergeom([3, 3/2-1/2*I], [5/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a^3+(24/5+8/5*I)*exp((1+3*I)*arcsec(a*x))*hypergeom([4, 3/2-1/2*I], [5/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a^3
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.96

$$\int e^{\sec^{-1}(ax)} x^2 dx = \frac{e^{\sec^{-1}(ax)} \left((-4 - 4i) \left(-i + a \sqrt{1 - \frac{1}{a^2 x^2}} x \right) \text{Hypergeometric2F1} \left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) + a^4 x^4 (5 + \cos[2 \text{ArcSec}[a*x]] - \sin[2 \text{ArcSec}[a*x]]) \right)}{12a^4 x}$$

input

```
Integrate[E^ArcSec[a*x]*x^2,x]
```

output

```
(E^ArcSec[a*x]*((-4 - 4*I)*(-I + a*Sqrt[1 - 1/(a^2*x^2)]*x)*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])] + a^4*x^4*(5 + Cos[2*ArcSec[a*x]] - Sin[2*ArcSec[a*x]])))/(12*a^4*x)
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5789, 27, 4974, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 e^{\sec^{-1}(ax)} dx \\ & \quad \downarrow \text{5789} \\ & \frac{\int a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} x^4 d \sec^{-1}(ax)}{a} \\ & \quad \downarrow \text{27} \\ & \frac{\int a^4 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} x^4 d \sec^{-1}(ax)}{a^3} \\ & \quad \downarrow \text{4974} \end{aligned}$$

$$\frac{\int \left(\frac{16ie^{(1+3i)\sec^{-1}(ax)}}{(1+e^{2i\sec^{-1}(ax)})^4} - \frac{8ie^{(1+3i)\sec^{-1}(ax)}}{(1+e^{2i\sec^{-1}(ax)})^3} \right) d\sec^{-1}(ax)}{a^3}$$

↓ 2009

$$\frac{\left(\frac{24}{5} + \frac{8i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right) - \left(\frac{12}{5} + \frac{4i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

input `Int [E^ArcSec [a*x]*x^2,x]`

output `((-12/5 - (4*I)/5)*E^((1 + 3*I)*ArcSec[a*x])*Hypergeometric2F1[3/2 - I/2, 3, 5/2 - I/2, -E^((2*I)*ArcSec[a*x])] + (24/5 + (8*I)/5)*E^((1 + 3*I)*ArcSec[a*x])*Hypergeometric2F1[3/2 - I/2, 4, 5/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a^3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4974 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(d_) + (e_)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigToExp[F^(c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]`

rule 5789 `Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] :> Simp[1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple [F]

$$\int e^{\operatorname{arcsec}(ax)} x^2 dx$$

input `int(exp(arcsec(a*x))*x^2,x)`

output `int(exp(arcsec(a*x))*x^2,x)`

Fricas [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x^2,x, algorithm="fricas")`

output `integral(x^2*e^(arcsec(a*x)), x)`

Sympy [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\operatorname{asec}(ax)} dx$$

input `integrate(exp(asec(a*x))*x**2,x)`

output `Integral(x**2*exp(asec(a*x)), x)`

Maxima [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x^2,x, algorithm="maxima")`

output `integrate(x^2*e^(arcsec(a*x)), x)`

Giac [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x^2,x, algorithm="giac")`

output `integrate(x^2*e^(arcsec(a*x)), x)`

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\arccos(\frac{1}{ax})} dx$$

input `int(x^2*exp(acos(1/(a*x))),x)`

output `int(x^2*exp(acos(1/(a*x))), x)`

Reduce [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int e^{a \sec(ax)} x^2 dx$$

input `int(exp(asec(a*x))*x^2,x)`

output `int(e**asec(a*x)*x**2,x)`

3.44 $\int e^{\sec^{-1}(ax)} x dx$

Optimal result	344
Mathematica [A] (verified)	344
Rubi [A] (verified)	345
Maple [F]	346
Fricas [F]	347
Sympy [F]	347
Maxima [F]	347
Giac [F]	348
Mupad [F(-1)]	348
Reduce [F]	348

Optimal result

Integrand size = 8, antiderivative size = 91

$$\int e^{\sec^{-1}(ax)} x dx = -\frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 2, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2} + \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}$$

output

```
(-8/5-4/5*I)*exp((1+2*I)*arcsec(a*x))*hypergeom([2, 1-1/2*I],[2-1/2*I],-(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a^2+(16/5+8/5*I)*exp((1+2*I)*arcsec(a*x))*hypergeom([3, 1-1/2*I],[2-1/2*I],-(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a^2
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.18

$$\int e^{\sec^{-1}(ax)} x dx = \frac{\left(\frac{1}{5} + \frac{i}{10}\right) e^{\sec^{-1}(ax)} \left((-2 + i)ax \left(\sqrt{1 - \frac{1}{a^2x^2}} - ax\right) + (1 + 2i) \operatorname{Hypergeometric2F1}\left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)\right)}{a^2}$$

input `Integrate[E^ArcSec[a*x]*x,x]`

output
$$\frac{((1/5 + I/10)*E^{\text{ArcSec}[a*x]}*((-2 + I)*a*x*\sqrt{1 - 1/(a^2*x^2)} - a*x) + (1 + 2*I)*\text{Hypergeometric2F1}[-1/2*I, 1, 1 - I/2, -E^{((2*I)*\text{ArcSec}[a*x])}] - E^{((2*I)*\text{ArcSec}[a*x])}*\text{Hypergeometric2F1}[1, 1 - I/2, 2 - I/2, -E^{((2*I)*\text{ArcSec}[a*x])}])}{a^2}$$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5789, 27, 4974, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x e^{\sec^{-1}(ax)} dx \\ & \quad \downarrow \text{5789} \\ & \frac{\int a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} x^3 d \sec^{-1}(ax)}{a} \\ & \quad \downarrow \text{27} \\ & \frac{\int a^3 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} x^3 d \sec^{-1}(ax)}{a^2} \\ & \quad \downarrow \text{4974} \\ & \frac{\int \left(\frac{8ie^{(1+2i)\sec^{-1}(ax)}}{(1+e^{2i\sec^{-1}(ax)})^3} - \frac{4ie^{(1+2i)\sec^{-1}(ax)}}{(1+e^{2i\sec^{-1}(ax)})^2} \right) d \sec^{-1}(ax)}{a^2} \\ & \quad \downarrow \text{2009} \\ & \frac{\left(\frac{16}{5} + \frac{8i}{5} \right) e^{(1+2i)\sec^{-1}(ax)} \text{Hypergeometric2F1} \left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)} \right) - \left(\frac{8}{5} + \frac{4i}{5} \right) e^{(1+2i)\sec^{-1}(ax)} \text{Hypergeometric2F1} \left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)} \right)}{a^2} \end{aligned}$$

input `Int [E^ArcSec[a*x]*x,x]`

output `((-8/5 - (4*I)/5)*E^((1 + 2*I)*ArcSec[a*x])*Hypergeometric2F1[1 - I/2, 2, 2 - I/2, -E^((2*I)*ArcSec[a*x])] + (16/5 + (8*I)/5)*E^((1 + 2*I)*ArcSec[a*x])*Hypergeometric2F1[1 - I/2, 3, 2 - I/2, -E^((2*I)*ArcSec[a*x])])/a^2`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4974 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^(c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]`

rule 5789 `Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Simp[1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple **[F]**

$$\int e^{\operatorname{arcsec}(ax)} x dx$$

input `int(exp(arcsec(a*x))*x,x)`

output `int(exp(arcsec(a*x))*x,x)`

Fricas [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x,x, algorithm="fricas")`

output `integral(x*e^(arcsec(a*x)), x)`

Sympy [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{\operatorname{asec}(ax)} dx$$

input `integrate(exp(asec(a*x))*x,x)`

output `Integral(x*exp(asec(a*x)), x)`

Maxima [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x,x, algorithm="maxima")`

output `integrate(x*e^(arcsec(a*x)), x)`

Giac [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x))*x,x, algorithm="giac")`

output `integrate(x*e^(arcsec(a*x)), x)`

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{\operatorname{acos}(\frac{1}{ax})} dx$$

input `int(x*exp(acos(1/(a*x))),x)`

output `int(x*exp(acos(1/(a*x))), x)`

Reduce [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int e^{\operatorname{asec}(ax)} x dx$$

input `int(exp(asec(a*x))*x,x)`

output `int(e**asec(a*x)*x,x)`

3.45 $\int e^{\sec^{-1}(ax)} dx$

Optimal result	349
Mathematica [A] (verified)	349
Rubi [A] (verified)	350
Maple [F]	351
Fricas [F]	351
Sympy [F]	352
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	353
Reduce [F]	353

Optimal result

Integrand size = 6, antiderivative size = 91

$$\int e^{\sec^{-1}(ax)} dx = -\frac{(1+i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 1, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a} + \frac{(2+2i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 2, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

output

```
(-1-I)*exp((1+I)*arcsec(a*x))*hypergeom([1, 1/2-1/2*I], [3/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a+(2+2*I)*exp((1+I)*arcsec(a*x))*hypergeom([2, 1/2-1/2*I], [3/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.59

$$\int e^{\sec^{-1}(ax)} dx = e^{\sec^{-1}(ax)} x - \frac{(1-i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 1, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

input `Integrate[E^ArcSec[a*x], x]`

output `E^ArcSec[a*x]*x - ((1 - I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5789, 4974, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{\sec^{-1}(ax)} dx$$

$$\downarrow 5789$$

$$\frac{\int a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} dx}{a}$$

$$\downarrow 4974$$

$$\frac{\int \left(\frac{4ie^{(1+i)\sec^{-1}(ax)}}{(1+e^{2i\sec^{-1}(ax)})^2} - \frac{2ie^{(1+i)\sec^{-1}(ax)}}{1+e^{2i\sec^{-1}(ax)}} \right) d\sec^{-1}(ax)}{a}$$

$$\downarrow 2009$$

$$\frac{(2 + 2i)e^{(1+i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 2, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right) - (1 + i)e^{(1+i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

input `Int[E^ArcSec[a*x], x]`

output `((-1 - I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])] + (2 + 2*I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 2, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4974 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^(c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]`

rule 5789 `Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_.)), x_Symbol] := Simp[1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple [F]

$$\int e^{\operatorname{arcsec}(ax)} dx$$

input `int(exp(arcsec(a*x)),x)`

output `int(exp(arcsec(a*x)),x)`

Fricas [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x)),x, algorithm="fricas")`

output `integral(e^(arcsec(a*x)), x)`

Sympy [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\operatorname{asec}(ax)} dx$$

input `integrate(exp(asec(a*x)),x)`

output `Integral(exp(asec(a*x)), x)`

Maxima [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x)),x, algorithm="maxima")`

output `integrate(e^(arcsec(a*x)), x)`

Giac [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{(\operatorname{arcsec}(ax))} dx$$

input `integrate(exp(arcsec(a*x)),x, algorithm="giac")`

output `integrate(e^(arcsec(a*x)), x)`

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\arccos(\frac{1}{ax})} dx$$

input `int(exp(acos(1/(a*x))), x)`output `int(exp(acos(1/(a*x))), x)`**Reduce [F]**

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\operatorname{asec}(ax)} dx$$

input `int(exp(asec(a*x)), x)`output `int(e**asec(a*x), x)`

3.46 $\int \frac{e^{\sec^{-1}(ax)}}{x} dx$

Optimal result	354
Mathematica [A] (verified)	354
Rubi [A] (verified)	355
Maple [F]	356
Fricas [F]	356
Sympy [F]	357
Maxima [F]	357
Giac [F]	357
Mupad [F(-1)]	358
Reduce [F]	358

Optimal result

Integrand size = 10, antiderivative size = 45

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = -ie^{\sec^{-1}(ax)} + 2ie^{\sec^{-1}(ax)} \operatorname{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right)$$

output

```
-I*exp(arcsec(a*x))+2*I*exp(arcsec(a*x))*hypergeom([1, -1/2*I],[1-1/2*I],-(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.76

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = -i \left(-e^{\sec^{-1}(ax)} \operatorname{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) + \left(\frac{1}{5} - \frac{2i}{5} \right) e^{(1+2i) \sec^{-1}(ax)} \operatorname{Hypergeometric2F1} \left(1, 1 - \frac{i}{2}, 2 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) \right)$$

input

```
Integrate[E^ArcSec[a*x]/x,x]
```

output

```
(-I)*(-(E^ArcSec[a*x]*Hypergeometric2F1[-1/2*I, 1, 1 - I/2, -E^((2*I)*ArcSec[a*x])]) + (1/5 - (2*I)/5)*E^((1 + 2*I)*ArcSec[a*x])*Hypergeometric2F1[1, 1 - I/2, 2 - I/2, -E^((2*I)*ArcSec[a*x])])
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5789, 27, 4942, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx$$

↓ 5789

$$\frac{\int a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} dx \sec^{-1}(ax)}{a}$$

↓ 27

$$\int ax \sqrt{1 - \frac{1}{a^2 x^2}} e^{\sec^{-1}(ax)} d \sec^{-1}(ax)$$

↓ 4942

$$i \int \left(\frac{2e^{\sec^{-1}(ax)}}{1 + e^{2i \sec^{-1}(ax)}} - e^{\sec^{-1}(ax)} \right) d \sec^{-1}(ax)$$

↓ 2009

$$i \left(-e^{\sec^{-1}(ax)} + 2e^{\sec^{-1}(ax)} \text{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) \right)$$

input

```
Int [E^ArcSec[a*x]/x, x]
```

output

```
I*(-E^ArcSec[a*x] + 2*E^ArcSec[a*x]*Hypergeometric2F1[-1/2*I, 1, 1 - I/2, -E^((2*I)*ArcSec[a*x])])
```

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4942 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*Tan[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Simp[I^n Int[ExpandIntegrand[F^(c*(a + b*x))*((1 - E^(2*I*(d + e*x)))^n/(1 + E^(2*I*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

rule 5789 `Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_.)), x_Symbol] := Simp[1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x} dx$$

input `int(exp(arcsec(a*x))/x,x)`

output `int(exp(arcsec(a*x))/x,x)`

Fricas [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x} dx$$

input `integrate(exp(arcsec(a*x))/x,x, algorithm="fricas")`

output `integral(e^(arcsec(a*x))/x, x)`

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x} dx$$

input `integrate(exp(asec(a*x))/x,x)`

output `Integral(exp(asec(a*x))/x, x)`

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x} dx$$

input `integrate(exp(arcsec(a*x))/x,x, algorithm="maxima")`

output `integrate(e^(arcsec(a*x))/x, x)`

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x} dx$$

input `integrate(exp(arcsec(a*x))/x,x, algorithm="giac")`

output `integrate(e^(arcsec(a*x))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\arccos(\frac{1}{ax})}}{x} dx$$

input `int(exp(acos(1/(a*x)))/x,x)`output `int(exp(acos(1/(a*x)))/x, x)`**Reduce [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x} dx$$

input `int(exp(asec(a*x))/x,x)`output `int(e**asec(a*x)/x,x)`

$$3.47 \quad \int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$$

Optimal result	359
Mathematica [A] (verified)	359
Rubi [A] (verified)	360
Maple [F]	361
Fricas [A] (verification not implemented)	361
Sympy [F]	362
Maxima [F]	362
Giac [A] (verification not implemented)	362
Mupad [F(-1)]	363
Reduce [F]	363

Optimal result

Integrand size = 10, antiderivative size = 39

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} a e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{e^{\sec^{-1}(ax)}}{2x}$$

output `1/2*a*exp(arcsec(a*x))*(1-1/a^2/x^2)^(1/2)-1/2*exp(arcsec(a*x))/x`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} a e^{\sec^{-1}(ax)} \left(\sqrt{1 - \frac{1}{a^2 x^2}} - \frac{1}{ax} \right)$$

input `Integrate[E^ArcSec[a*x]/x^2,x]`

output `(a*E^ArcSec[a*x]*(Sqrt[1 - 1/(a^2*x^2)] - 1/(a*x)))/2`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5789, 27, 4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$$

$$\downarrow \text{5789}$$

$$\frac{\int a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} d \sec^{-1}(ax)}{a}$$

$$\downarrow \text{27}$$

$$a \int e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} d \sec^{-1}(ax)$$

$$\downarrow \text{4932}$$

$$a \left(\frac{1}{2} \sqrt{1 - \frac{1}{a^2 x^2}} e^{\sec^{-1}(ax)} - \frac{e^{\sec^{-1}(ax)}}{2ax} \right)$$

input `Int [E^ArcSec [a*x] /x^2, x]`

output `a*((E^ArcSec [a*x]*Sqrt [1 - 1/(a^2*x^2)])/2 - E^ArcSec [a*x]/(2*a*x))`

Defintions of rubi rules used

rule 27 `Int [(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 4932 `Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :=
 Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
 reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

rule 5789 `Int[(u_.)*(f_)^(ArcSec[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] := Simp[
 1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x,
 ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x^2} dx$$

input `int(exp(arcsec(a*x))/x^2,x)`

output `int(exp(arcsec(a*x))/x^2,x)`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.59

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{(\sqrt{a^2x^2 - 1} - 1)e^{\operatorname{arcsec}(ax)}}{2x}$$

input `integrate(exp(arcsec(a*x))/x^2,x, algorithm="fricas")`

output `1/2*(sqrt(a^2*x^2 - 1) - 1)*e^(arcsec(a*x))/x`

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^2} dx$$

input `integrate(exp(asec(a*x))/x**2,x)`

output `Integral(exp(asec(a*x))/x**2, x)`

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^2} dx$$

input `integrate(exp(arcsec(a*x))/x^2,x, algorithm="maxima")`

output `integrate(e^(arcsec(a*x))/x^2, x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} \left(\sqrt{-\frac{1}{a^2 x^2} + 1} e^{\arccos(\frac{1}{ax})} - \frac{e^{\arccos(\frac{1}{ax})}}{ax} \right) a$$

input `integrate(exp(arcsec(a*x))/x^2,x, algorithm="giac")`

output `1/2*(sqrt(-1/(a^2*x^2) + 1)*e^(arccos(1/(a*x))) - e^(arccos(1/(a*x)))/(a*x)))*a`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\arccos(\frac{1}{ax})}}{x^2} dx$$

input `int(exp(acos(1/(a*x)))/x^2,x)`output `int(exp(acos(1/(a*x)))/x^2, x)`**Reduce [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^2} dx$$

input `int(exp(asec(a*x))/x^2,x)`output `int(e**asec(a*x)/x**2,x)`

$$3.48 \quad \int \frac{e^{\sec^{-1}(ax)}}{x^3} dx$$

Optimal result	364
Mathematica [A] (verified)	364
Rubi [A] (verified)	365
Maple [F]	366
Fricas [A] (verification not implemented)	367
Sympy [F]	367
Maxima [F]	367
Giac [F]	368
Mupad [F(-1)]	368
Reduce [F]	368

Optimal result

Integrand size = 10, antiderivative size = 41

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = -\frac{1}{5}a^2 e^{\sec^{-1}(ax)} \cos(2 \sec^{-1}(ax)) + \frac{1}{10}a^2 e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax))$$

output

```
-1/5*a^2*exp(arcsec(a*x))*cos(2*arcsec(a*x))+1/10*a^2*exp(arcsec(a*x))*sin(2*arcsec(a*x))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \frac{1}{10}a^2 e^{\sec^{-1}(ax)} (-2 \cos(2 \sec^{-1}(ax)) + \sin(2 \sec^{-1}(ax)))$$

input

```
Integrate[E^ArcSec[a*x]/x^3,x]
```

output

```
(a^2*E^ArcSec[a*x]*(-2*Cos[2*ArcSec[a*x]] + Sin[2*ArcSec[a*x]]))/10
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5789, 27, 4972, 27, 4932}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{e^{\sec^{-1}(ax)}}{x^3} dx \\
 & \quad \downarrow \text{5789} \\
 & \int \frac{a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}}}{x} d \sec^{-1}(ax) \\
 & \quad \downarrow \text{27} \\
 & a^2 \int \frac{e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}}}{ax} d \sec^{-1}(ax) \\
 & \quad \downarrow \text{4972} \\
 & a^2 \int \frac{1}{2} e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax)) d \sec^{-1}(ax) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} a^2 \int e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax)) d \sec^{-1}(ax) \\
 & \quad \downarrow \text{4932} \\
 & \frac{1}{2} a^2 \left(\frac{1}{5} e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax)) - \frac{2}{5} e^{\sec^{-1}(ax)} \cos(2 \sec^{-1}(ax)) \right)
 \end{aligned}$$

input `Int [E^ArcSec [a*x] /x^3, x]`

output `(a^2*((-2*E^ArcSec [a*x] *Cos [2*ArcSec [a*x]])/5 + (E^ArcSec [a*x] *Sin [2*ArcSec [a*x]])/5))/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_)] /; FreeQ[b, x]`

rule 4932 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*Sin[(d_) + (e_)*(x_)], x_Symbol] := Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]`

rule 4972 `Int[Cos[(f_) + (g_)*(x_)]^(n_)*(F_)^((c_)*((a_) + (b_)*(x_)))*Sin[(d_) + (e_)*(x_)]^(m_), x_Symbol] := Int[ExpandTrigReduce[F^(c*(a + b*x)), Sin[d + e*x]^m*Cos[f + g*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e, f, g}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

rule 5789 `Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Simp[1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]`

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x^3} dx$$

input `int(exp(arcsec(a*x))/x^3,x)`

output `int(exp(arcsec(a*x))/x^3,x)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \frac{(a^2x^2 + \sqrt{a^2x^2 - 1} - 2)e^{\operatorname{arcsec}(ax)}}{5x^2}$$

input `integrate(exp(arcsec(a*x))/x^3,x, algorithm="fricas")`output `1/5*(a^2*x^2 + sqrt(a^2*x^2 - 1) - 2)*e^(arcsec(a*x))/x^2`**Sympy [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^3} dx$$

input `integrate(exp(asec(a*x))/x**3,x)`output `Integral(exp(asec(a*x))/x**3, x)`**Maxima [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^3} dx$$

input `integrate(exp(arcsec(a*x))/x^3,x, algorithm="maxima")`output `integrate(e^(arcsec(a*x))/x^3, x)`

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^3} dx$$

input `integrate(exp(arcsec(a*x))/x^3,x, algorithm="giac")`

output `integrate(e^(arcsec(a*x))/x^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\arccos(\frac{1}{ax})}}{x^3} dx$$

input `int(exp(acos(1/(a*x)))/x^3,x)`

output `int(exp(acos(1/(a*x)))/x^3, x)`

Reduce [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^3} dx$$

input `int(exp(asec(a*x))/x^3,x)`

output `int(e**asec(a*x)/x**3,x)`

3.49 $\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$

Optimal result	369
Mathematica [A] (verified)	369
Rubi [A] (verified)	370
Maple [F]	371
Fricas [A] (verification not implemented)	372
Sympy [F]	372
Maxima [F]	372
Giac [F]	373
Mupad [F(-1)]	373
Reduce [F]	373

Optimal result

Integrand size = 10, antiderivative size = 84

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{1}{8}a^3 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{a^2 e^{\sec^{-1}(ax)}}{8x} - \frac{3}{40}a^3 e^{\sec^{-1}(ax)} \cos(3 \sec^{-1}(ax)) + \frac{1}{40}a^3 e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax))$$

output

```
1/8*a^3*exp(arcsec(a*x))*(1-1/a^2/x^2)^(1/2)-1/8*a^2*exp(arcsec(a*x))/x-3/40*a^3*exp(arcsec(a*x))*cos(3*arcsec(a*x))+1/40*a^3*exp(arcsec(a*x))*sin(3*arcsec(a*x))
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.64

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{1}{40}a^3 e^{\sec^{-1}(ax)} \left(5\sqrt{1 - \frac{1}{a^2 x^2}} - \frac{5}{ax} - 3 \cos(3 \sec^{-1}(ax)) + \sin(3 \sec^{-1}(ax)) \right)$$

input

```
Integrate[E^ArcSec[a*x]/x^4,x]
```

output

```
(a^3*E^ArcSec[a*x]*(5*Sqrt[1 - 1/(a^2*x^2)] - 5/(a*x) - 3*Cos[3*ArcSec[a*x]] + Sin[3*ArcSec[a*x]]))/40
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.94, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5789, 27, 4972, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$$

$$\downarrow \text{5789}$$

$$\int \frac{a^2 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}}}{x^2} d \sec^{-1}(ax)$$

$$\downarrow \text{27}$$

$$a^3 \int \frac{e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}}}{a^2 x^2} d \sec^{-1}(ax)$$

$$\downarrow \text{4972}$$

$$a^3 \int \left(\frac{1}{4} e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax)) + \frac{1}{4} e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} \right) d \sec^{-1}(ax)$$

$$\downarrow \text{2009}$$

$$a^3 \left(\frac{1}{8} \sqrt{1 - \frac{1}{a^2 x^2}} e^{\sec^{-1}(ax)} - \frac{e^{\sec^{-1}(ax)}}{8ax} - \frac{3}{40} e^{\sec^{-1}(ax)} \cos(3 \sec^{-1}(ax)) + \frac{1}{40} e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax)) \right)$$

input

```
Int[E^ArcSec[a*x]/x^4,x]
```

output

```
a^3*((E^ArcSec[a*x]*Sqrt[1 - 1/(a^2*x^2)])/8 - E^ArcSec[a*x]/(8*a*x) - (3*
E^ArcSec[a*x]*Cos[3*ArcSec[a*x]])/40 + (E^ArcSec[a*x]*Sin[3*ArcSec[a*x]])/
40)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 4972

```
Int[Cos[(f_.) + (g_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_
.) + (e_.)*(x_)]^(m_.), x_Symbol] := Int[ExpandTrigReduce[F^(c*(a + b*x)),
Sin[d + e*x]^m*cos[f + g*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e, f, g}, x]
&& IGtQ[m, 0] && IGtQ[n, 0]
```

rule 5789

```
Int[(u_.)*(f_)^(ArcSec[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] := Simp[
1/b Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x,
ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x^4} dx$$

input

```
int(exp(arcsec(a*x))/x^4,x)
```

output

```
int(exp(arcsec(a*x))/x^4,x)
```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.48

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{(a^2x^2 + (a^2x^2 + 1)\sqrt{a^2x^2 - 1} - 3)e^{\sec^{-1}(ax)}}{10x^3}$$

input `integrate(exp(arcsec(a*x))/x^4,x, algorithm="fricas")`output `1/10*(a^2*x^2 + (a^2*x^2 + 1)*sqrt(a^2*x^2 - 1) - 3)*e^(arcsec(a*x))/x^3`**Sympy [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^4} dx$$

input `integrate(exp(asec(a*x))/x**4,x)`output `Integral(exp(asec(a*x))/x**4, x)`**Maxima [F]**

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^4} dx$$

input `integrate(exp(arcsec(a*x))/x^4,x, algorithm="maxima")`output `integrate(e^(arcsec(a*x))/x^4, x)`

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^4} dx$$

input `integrate(exp(arcsec(a*x))/x^4,x, algorithm="giac")`

output `integrate(e^(arcsec(a*x))/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\arccos(\frac{1}{ax})}}{x^4} dx$$

input `int(exp(acos(1/(a*x)))/x^4,x)`

output `int(exp(acos(1/(a*x)))/x^4, x)`

Reduce [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^4} dx$$

input `int(exp(asec(a*x))/x^4,x)`

output `int(e**asec(a*x)/x**4,x)`

3.50 $\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	374
Mathematica [A] (verified)	374
Rubi [A] (warning: unable to verify)	375
Maple [A] (verified)	378
Fricas [F]	378
Sympy [F]	379
Maxima [F]	379
Giac [A] (verification not implemented)	379
Mupad [F(-1)]	380
Reduce [F]	380

Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{i \sec^{-1}(a+bx)^2}{2d} - \frac{\sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{d} + \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2d}$$

output

```
1/2*I*arcsec(b*x+a)^2/d-arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/d+1/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/d
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.86

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{i \left(\sec^{-1}(a+bx) \left(\sec^{-1}(a+bx) + 2i \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right) \right) + \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) \right)}{2d}$$

input

```
Integrate[ArcSec[a + b*x]/((a*d)/b + d*x), x]
```

output

```
((I/2)*(ArcSec[a + b*x]*(ArcSec[a + b*x] + (2*I)*Log[1 + E^((2*I)*ArcSec[a + b*x])]) + PolyLog[2, -E^((2*I)*ArcSec[a + b*x])]))/d
```

Rubi [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {5779, 27, 5741, 5137, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx \\
 & \quad \downarrow \text{5779} \\
 & \frac{\int \frac{b \sec^{-1}(a+bx)}{d(a+bx)} d(a+bx)}{b} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sec^{-1}(a+bx)}{a+bx} d(a+bx)}{d} \\
 & \quad \downarrow \text{5741} \\
 & \frac{\int (a + bx) \arccos\left(\frac{1}{a+bx}\right) d\frac{1}{a+bx}}{d} \\
 & \quad \downarrow \text{5137} \\
 & \frac{\int (a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \arccos\left(\frac{1}{a+bx}\right) d\arccos\left(\frac{1}{a+bx}\right)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \arccos\left(\frac{1}{a+bx}\right) \tan\left(\arccos\left(\frac{1}{a+bx}\right)\right) d\arccos\left(\frac{1}{a+bx}\right)}{d} \\
 & \quad \downarrow \text{4202}
 \end{aligned}$$

$$\frac{\frac{i}{2(a+bx)^2} - 2i \int \frac{e^{2i \arccos\left(\frac{1}{a+bx}\right)} \arccos\left(\frac{1}{a+bx}\right)}{1+e^{2i \arccos\left(\frac{1}{a+bx}\right)}} d \arccos\left(\frac{1}{a+bx}\right)}{d}$$

↓ 2620

$$\frac{\frac{i}{2(a+bx)^2} - 2i \left(\frac{1}{2} i \int \log \left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)} \right) d \arccos\left(\frac{1}{a+bx}\right) - \frac{1}{2} i \arccos\left(\frac{1}{a+bx}\right) \log \left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)} \right) \right)}{d}$$

↓ 2715

$$\frac{\frac{i}{2(a+bx)^2} - 2i \left(\frac{1}{4} \int (a+bx) \log \left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)} \right) d e^{2i \arccos\left(\frac{1}{a+bx}\right)} - \frac{1}{2} i \arccos\left(\frac{1}{a+bx}\right) \log \left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)} \right) \right)}{d}$$

↓ 2838

$$\frac{\frac{i}{2(a+bx)^2} - 2i \left(-\frac{1}{4} \text{PolyLog}(2, -a - bx) - \frac{1}{2} i \arccos\left(\frac{1}{a+bx}\right) \log \left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)} \right) \right)}{d}$$

input `Int[ArcSec[a + b*x]/((a*d)/b + d*x), x]`

output `((I/2)/(a + b*x)^2 - (2*I)*((-1/2*I)*ArcCos[(a + b*x)^(-1)]*Log[1 + E^((2*I)*ArcCos[(a + b*x)^(-1)])] - PolyLog[2, -a - b*x]/4))/d`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^((n_))], x_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{ Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x] \&\& \text{GtQ}[a, 0]$

rule 2838 $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x] \&\& \text{EqQ}[c*d, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4202 $\text{Int}[(c_) + (d_)*(x_)^(m_)*\tan[(e_) + (f_)*(x_)], x_Symbol] \rightarrow \text{Simp}[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - \text{Simp}[2*I \text{ Int}[(c + d*x)^m*(E^{(2*I*(e + f*x))}/(1 + E^{(2*I*(e + f*x))})), x], x] /; \text{FreeQ}\{c, d, e, f\}, x] \&\& \text{IGtQ}[m, 0]$

rule 5137 $\text{Int}[(a_) + \text{ArcCos}[(c_)*(x_)]*(b_)]^(n_)/(x_), x_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b*x)^n*\text{Tan}[x], x], x, \text{ArcCos}[c*x]] /; \text{FreeQ}\{a, b, c\}, x] \&\& \text{IGtQ}[n, 0]$

rule 5741 $\text{Int}[(a_) + \text{ArcSec}[(c_)*(x_)]*(b_)]/(x_), x_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b*\text{ArcCos}[x/c])/x, x], x, 1/x] /; \text{FreeQ}\{a, b, c\}, x]$

rule 5779 $\text{Int}[(a_) + \text{ArcSec}[(c_) + (d_)*(x_)]*(b_)]^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] \rightarrow \text{Simp}[1/d \text{ Subst}[\text{Int}[(f*(x/d))^m*(a + b*\text{ArcSec}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{EqQ}[d*e - c*f, 0] \&\& \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.43

method	result	size
derivativedivides	$\frac{\frac{ib \operatorname{arcsec}(bx+a)^2}{2d} - \frac{b \operatorname{arcsec}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{d}}{b} + \frac{ib \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{2d}$	99
default	$\frac{\frac{ib \operatorname{arcsec}(bx+a)^2}{2d} - \frac{b \operatorname{arcsec}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{d}}{b} + \frac{ib \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{2d}$	99

input `int(arcsec(b*x+a)/(a*d/b+d*x), x, method=_RETURNVERBOSE)`

output
$$\frac{1}{b} \left(\frac{1}{2} I/d * b * \operatorname{arcsec}(bx+a)^2 - 1/d * b * \operatorname{arcsec}(bx+a) * \ln\left(1 + \left(\frac{1}{bx+a} + I * \left(1 - \frac{1}{(bx+a)^2}\right)^{1/2}\right)^2\right) + 1/2 * I/d * b * \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + I * \left(1 - \frac{1}{(bx+a)^2}\right)^{1/2}\right)^2\right) \right)$$

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsec}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arcsec(b*x+a)/(a*d/b+d*x), x, algorithm="fricas")`

output `integral(b*arcsec(b*x + a)/(b*d*x + a*d), x)`

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{asec}\left(\frac{a+bx}{a+bx}\right) dx}{d}}$$

input `integrate(asec(b*x+a)/(a*d/b+d*x), x)`

output `b*Integral(asec(a + b*x)/(a + b*x), x)/d`

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsec}(bx + a)}{dx + \frac{ad}{b}} dx$$

input `integrate(arcsec(b*x+a)/(a*d/b+d*x), x, algorithm="maxima")`

output `-1/2*(2*b*d*integrate(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*log(b*x + a)/(b^3*d*x^3 + 3*a*b^2*d*x^2 + (3*a^2 - 1)*b*d*x + (a^3 - a)*d), x) + 2*I*b*d*integrate(log(b*x + a)/(b^3*d*x^3 + 3*a*b^2*d*x^2 + (3*a^2 - 1)*b*d*x + (a^3 - a)*d), x) - 2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b*x + a) + I*log(b^2*x^2 + 2*a*b*x + a^2)*log(b*x + a) - I*log(b*x + a + 1)*log(b*x + a) - I*log(b*x + a)^2 - I*log(b*x + a)*log(-b*x - a + 1) - I*dilog(b*x + a) - I*dilog(-b*x - a))/d`

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.67

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = -\frac{1}{4} b^2 \left(\frac{2 (bx + a)^2 \arccos\left(\frac{1}{\left(\frac{bx+a}{\frac{a}{bx+a}-1}\right)-a}\left(\frac{a}{\frac{a}{bx+a}-1}\right)+a\right)}{b^3 d} - \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)}{b^3 d} - \frac{1}{(bx+a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)} \right)$$

input `integrate(arcsec(b*x+a)/(a*d/b+d*x),x, algorithm="giac")`

output
$$-1/4*b^2*(2*(b*x + a)^2*\arccos(1/((b*x + a)*(a/(b*x + a) - 1) - a)*(a/(b*x + a) - 1) + a))/(b^3*d) - ((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) - 1/((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1)))/(b^3*d)$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{dx + \frac{ad}{b}} dx$$

input `int(acos(1/(a + b*x))/(d*x + (a*d)/b),x)`

output `int(acos(1/(a + b*x))/(d*x + (a*d)/b), x)`

Reduce [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{\left(\int \frac{asec(bx+a)}{bx+a} dx\right) b}{d}$$

input `int(asec(b*x+a)/(a*d/b+d*x),x)`

output `(int(asec(a + b*x)/(a + b*x),x)*b)/d`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	381
4.2	Links to plain text integration problems used in this report for each CAS .	399

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```



```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
        If [Head [expn] === RootSum,
            Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
            If [Head [expn] === Integrate || Head [expn] === Int,
                Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
                9]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```



```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file