

# Computer Algebra Independent Integration Tests

Summer 2024

6-Hyperbolic-functions/6.3-Hyperbolic-tangent/307-6.3.2

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 8:34am

# Contents

<b>1</b>	<b>Introduction</b>	<b>10</b>
1.1	Listing of CAS systems tested . . . . .	11
1.2	Results . . . . .	12
1.3	Time and leaf size Performance . . . . .	16
1.4	Performance based on number of rules Rubi used . . . . .	18
1.5	Performance based on number of steps Rubi used . . . . .	19
1.6	Solved integrals histogram based on leaf size of result . . . . .	20
1.7	Solved integrals histogram based on CPU time used . . . . .	21
1.8	Leaf size vs. CPU time used . . . . .	22
1.9	list of integrals with no known antiderivative . . . . .	23
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	23
1.11	list of integrals solved by CAS but failed verification . . . . .	23
1.12	Timing . . . . .	24
1.13	Verification . . . . .	24
1.14	Important notes about some of the results . . . . .	25
1.15	Current tree layout of integration tests . . . . .	28
1.16	Design of the test system . . . . .	29
<b>2</b>	<b>detailed summary tables of results</b>	<b>30</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	31
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	37
2.3	Detailed conclusion table specific for Rubi results . . . . .	102
<b>3</b>	<b>Listing of integrals</b>	<b>111</b>
3.1	$\int \tanh^6(a + bx) dx$ . . . . .	119
3.2	$\int \tanh^5(a + bx) dx$ . . . . .	126
3.3	$\int \tanh^4(a + bx) dx$ . . . . .	133
3.4	$\int \tanh^3(a + bx) dx$ . . . . .	139
3.5	$\int \tanh^2(a + bx) dx$ . . . . .	145
3.6	$\int \tanh(a + bx) dx$ . . . . .	150
3.7	$\int \coth(a + bx) dx$ . . . . .	155

3.8	$\int \coth^2(a + bx) dx$	160
3.9	$\int \coth^3(a + bx) dx$	166
3.10	$\int \coth^4(a + bx) dx$	173
3.11	$\int \coth^5(a + bx) dx$	179
3.12	$\int \coth^6(a + bx) dx$	187
3.13	$\int (b \tanh(c + dx))^{7/2} dx$	194
3.14	$\int (b \tanh(c + dx))^{5/2} dx$	202
3.15	$\int (b \tanh(c + dx))^{3/2} dx$	209
3.16	$\int \sqrt{b \tanh(c + dx)} dx$	216
3.17	$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$	223
3.18	$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx$	230
3.19	$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx$	238
3.20	$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx$	245
3.21	$\int \sqrt[3]{\tanh(8x)} dx$	253
3.22	$\int \tanh^n(a + bx) dx$	261
3.23	$\int (b \tanh(c + dx))^n dx$	266
3.24	$\int (a \tanh^2(x))^{3/2} dx$	271
3.25	$\int \sqrt{a \tanh^2(x)} dx$	278
3.26	$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx$	283
3.27	$\int (-\tanh^2(c + dx))^{5/2} dx$	289
3.28	$\int (-\tanh^2(c + dx))^{3/2} dx$	296
3.29	$\int \sqrt{-\tanh^2(c + dx)} dx$	303
3.30	$\int \frac{1}{\sqrt{-\tanh^2(c + dx)}} dx$	309
3.31	$\int \frac{1}{(-\tanh^2(c + dx))^{3/2}} dx$	315
3.32	$\int \frac{1}{(-\tanh^2(c + dx))^{5/2}} dx$	322
3.33	$\int \sqrt{\tanh^3(x)} dx$	330
3.34	$\int (a \tanh^3(x))^{3/2} dx$	338
3.35	$\int \sqrt{a \tanh^3(x)} dx$	346
3.36	$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$	354
3.37	$\int (a \tanh^4(x))^{3/2} dx$	362
3.38	$\int \sqrt{a \tanh^4(x)} dx$	369
3.39	$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$	375
3.40	$\int (b \tanh^m(c + dx))^n dx$	381

3.41	$\int (a + a \tanh(c + dx))^5 dx$	387
3.42	$\int (a + a \tanh(c + dx))^4 dx$	396
3.43	$\int (a + a \tanh(c + dx))^3 dx$	403
3.44	$\int (a + a \tanh(c + dx))^2 dx$	410
3.45	$\int \frac{1}{a+a \tanh(c+dx)} dx$	416
3.46	$\int \frac{1}{(a+a \tanh(c+dx))^2} dx$	421
3.47	$\int \frac{1}{(a+a \tanh(c+dx))^3} dx$	427
3.48	$\int \frac{1}{(a+a \tanh(c+dx))^4} dx$	433
3.49	$\int \frac{1}{(a+a \tanh(c+dx))^5} dx$	440
3.50	$\int (1 + \tanh(x))^{7/2} dx$	449
3.51	$\int (1 + \tanh(x))^{5/2} dx$	456
3.52	$\int (1 + \tanh(x))^{3/2} dx$	462
3.53	$\int \sqrt{1 + \tanh(x)} dx$	468
3.54	$\int \frac{1}{\sqrt{1+\tanh(x)}} dx$	473
3.55	$\int \frac{1}{(1+\tanh(x))^{3/2}} dx$	479
3.56	$\int \frac{1}{(1+\tanh(x))^{5/2}} dx$	485
3.57	$\int (a + b \tanh(c + dx))^5 dx$	492
3.58	$\int (a + b \tanh(c + dx))^4 dx$	502
3.59	$\int (a + b \tanh(c + dx))^3 dx$	511
3.60	$\int (a + b \tanh(c + dx))^2 dx$	518
3.61	$\int \frac{1}{a+b \tanh(c+dx)} dx$	524
3.62	$\int \frac{1}{(a+b \tanh(c+dx))^2} dx$	530
3.63	$\int \frac{1}{(a+b \tanh(c+dx))^3} dx$	538
3.64	$\int \frac{1}{(a+b \tanh(c+dx))^4} dx$	548
3.65	$\int \frac{1}{4+6 \tanh(c+dx)} dx$	558
3.66	$\int \frac{1}{4-6 \tanh(c+dx)} dx$	563
3.67	$\int \sqrt{a + b \tanh(c + dx)} dx$	568
3.68	$\int \frac{1}{\sqrt{a+b \tanh(c+dx)}} dx$	575
3.69	$\int \frac{\sinh^4(x)}{1+\tanh(x)} dx$	582
3.70	$\int \frac{\sinh^3(x)}{1+\tanh(x)} dx$	588
3.71	$\int \frac{\sinh^2(x)}{1+\tanh(x)} dx$	595
3.72	$\int \frac{\sinh(x)}{1+\tanh(x)} dx$	601
3.73	$\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx$	607
3.74	$\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx$	613
3.75	$\int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx$	619

3.76	$\int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx$	626
3.77	$\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx$	632
3.78	$\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx$	640
3.79	$\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx$	646
3.80	$\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx$	654
3.81	$\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx$	663
3.82	$\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx$	674
3.83	$\int \frac{\sinh(x)}{a+b \tanh(x)} dx$	681
3.84	$\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx$	689
3.85	$\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx$	695
3.86	$\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx$	701
3.87	$\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx$	709
3.88	$\int \frac{\operatorname{csch}^5(x)}{a+b \tanh(x)} dx$	716
3.89	$\int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx$	725
3.90	$\int \frac{\operatorname{csch}(x)}{i+\tanh(x)} dx$	733
3.91	$\int (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n dx$	739
3.92	$\int \frac{\cosh^4(x)}{1+\tanh(x)} dx$	744
3.93	$\int \frac{\cosh^3(x)}{1+\tanh(x)} dx$	750
3.94	$\int \frac{\cosh^2(x)}{1+\tanh(x)} dx$	756
3.95	$\int \frac{\cosh(x)}{1+\tanh(x)} dx$	761
3.96	$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx$	766
3.97	$\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx$	771
3.98	$\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx$	776
3.99	$\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx$	782
3.100	$\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx$	787
3.101	$\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx$	793
3.102	$\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx$	799
3.103	$\int \frac{\operatorname{sech}^8(x)}{a+b \tanh(x)} dx$	806
3.104	$\int \frac{\operatorname{sech}^6(x)}{a+b \tanh(x)} dx$	814

3.105	$\int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx$	822
3.106	$\int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx$	829
3.107	$\int \frac{1}{a+b \tanh(x)} dx$	834
3.108	$\int \frac{\cosh^2(x)}{a+b \tanh(x)} dx$	840
3.109	$\int \frac{\cosh^4(x)}{a+b \tanh(x)} dx$	847
3.110	$\int \frac{\operatorname{sech}^7(x)}{a+b \tanh(x)} dx$	855
3.111	$\int \frac{\operatorname{sech}^5(x)}{a+b \tanh(x)} dx$	867
3.112	$\int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx$	877
3.113	$\int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx$	884
3.114	$\int \frac{\cosh(x)}{a+b \tanh(x)} dx$	889
3.115	$\int \frac{\cosh^3(x)}{a+b \tanh(x)} dx$	896
3.116	$\int \frac{\tanh^5(x)}{1+\tanh(x)} dx$	905
3.117	$\int \frac{\tanh^4(x)}{1+\tanh(x)} dx$	914
3.118	$\int \frac{\tanh^3(x)}{1+\tanh(x)} dx$	922
3.119	$\int \frac{\tanh^2(x)}{1+\tanh(x)} dx$	929
3.120	$\int \frac{\tanh(x)}{1+\tanh(x)} dx$	935
3.121	$\int \frac{1}{1+\tanh(x)} dx$	940
3.122	$\int \frac{\operatorname{coth}(x)}{1+\tanh(x)} dx$	945
3.123	$\int \frac{\operatorname{coth}^2(x)}{1+\tanh(x)} dx$	951
3.124	$\int \frac{\operatorname{coth}^3(x)}{1+\tanh(x)} dx$	958
3.125	$\int \frac{\operatorname{coth}^4(x)}{1+\tanh(x)} dx$	966
3.126	$\int \tanh(x)(1+\tanh(x))^{3/2} dx$	975
3.127	$\int \tanh(x)\sqrt{1+\tanh(x)} dx$	982
3.128	$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx$	988
3.129	$\int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx$	995
3.130	$\int \tanh^2(x)(1+\tanh(x))^{3/2} dx$	1002
3.131	$\int \tanh^2(x)\sqrt{1+\tanh(x)} dx$	1009
3.132	$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx$	1015
3.133	$\int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx$	1021
3.134	$\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx$	1028
3.135	$\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx$	1042
3.136	$\int \frac{\tanh^3(x)}{a+b \tanh(x)} dx$	1052

3.137	$\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx$	1061
3.138	$\int \frac{\tanh(x)}{a+b \tanh(x)} dx$	1068
3.139	$\int \frac{1}{a+b \tanh(x)} dx$	1074
3.140	$\int \frac{\coth(x)}{a+b \tanh(x)} dx$	1080
3.141	$\int \frac{\coth^2(x)}{a+b \tanh(x)} dx$	1087
3.142	$\int \frac{\coth^3(x)}{a+b \tanh(x)} dx$	1095
3.143	$\int \frac{\coth^4(x)}{a+b \tanh(x)} dx$	1105
3.144	$\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx$	1116
3.145	$\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$	1123
3.146	$\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$	1132
3.147	$\int x^5 \tanh(a + 2 \log(x)) dx$	1142
3.148	$\int x^3 \tanh(a + 2 \log(x)) dx$	1148
3.149	$\int x \tanh(a + 2 \log(x)) dx$	1153
3.150	$\int \frac{\tanh(a+2 \log(x))}{x} dx$	1158
3.151	$\int \frac{\tanh(a+2 \log(x))}{x^3} dx$	1163
3.152	$\int \frac{\tanh(a+2 \log(x))}{x^5} dx$	1168
3.153	$\int x^2 \tanh(a + 2 \log(x)) dx$	1174
3.154	$\int \tanh(a + 2 \log(x)) dx$	1182
3.155	$\int \frac{\tanh(a+2 \log(x))}{x^2} dx$	1190
3.156	$\int \frac{\tanh(a+2 \log(x))}{x^4} dx$	1199
3.157	$\int x^5 \tanh^2(a + 2 \log(x)) dx$	1207
3.158	$\int x^3 \tanh^2(a + 2 \log(x)) dx$	1214
3.159	$\int x \tanh^2(a + 2 \log(x)) dx$	1220
3.160	$\int \frac{\tanh^2(a+2 \log(x))}{x} dx$	1226
3.161	$\int \frac{\tanh^2(a+2 \log(x))}{x^3} dx$	1231
3.162	$\int \frac{\tanh^2(a+2 \log(x))}{x^5} dx$	1237
3.163	$\int \frac{\tanh^2(a+2 \log(x))}{x^7} dx$	1243
3.164	$\int \frac{\tanh^2(a+2 \log(x))}{x^9} dx$	1250
3.165	$\int x^4 \tanh^2(a + 2 \log(x)) dx$	1256
3.166	$\int x^2 \tanh^2(a + 2 \log(x)) dx$	1266
3.167	$\int \tanh^2(a + 2 \log(x)) dx$	1275
3.168	$\int \frac{\tanh^2(a+2 \log(x))}{x^2} dx$	1282
3.169	$\int \frac{\tanh^2(a+2 \log(x))}{x^4} dx$	1291
3.170	$\int (ex)^m \tanh(a + 2 \log(x)) dx$	1300
3.171	$\int (ex)^m \tanh^2(a + 2 \log(x)) dx$	1305

3.172	$\int (ex)^m \tanh^3(a + 2 \log(x)) dx$	1311
3.173	$\int \tanh^p(a + b \log(x)) dx$	1318
3.174	$\int (ex)^m \tanh^p(a + b \log(x)) dx$	1323
3.175	$\int \tanh^p\left(a + \frac{\log(x)}{2}\right) dx$	1328
3.176	$\int \tanh^p\left(a + \frac{\log(x)}{4}\right) dx$	1333
3.177	$\int \tanh^p\left(a + \frac{\log(x)}{6}\right) dx$	1338
3.178	$\int \tanh^p\left(a + \frac{\log(x)}{8}\right) dx$	1344
3.179	$\int \tanh^p(a + \log(x)) dx$	1351
3.180	$\int \tanh^p(a + 2 \log(x)) dx$	1356
3.181	$\int \tanh^p(a + 3 \log(x)) dx$	1361
3.182	$\int x^3 \tanh(d(a + b \log(cx^n))) dx$	1366
3.183	$\int x^2 \tanh(d(a + b \log(cx^n))) dx$	1371
3.184	$\int x \tanh(d(a + b \log(cx^n))) dx$	1376
3.185	$\int \tanh(d(a + b \log(cx^n))) dx$	1381
3.186	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx$	1386
3.187	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$	1392
3.188	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^3} dx$	1397
3.189	$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$	1402
3.190	$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx$	1409
3.191	$\int x \tanh^2(d(a + b \log(cx^n))) dx$	1416
3.192	$\int \tanh^2(d(a + b \log(cx^n))) dx$	1423
3.193	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x} dx$	1430
3.194	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^2} dx$	1436
3.195	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^3} dx$	1443
3.196	$\int \frac{\tanh^3(a+b \log(cx^n))}{x} dx$	1450
3.197	$\int \frac{\tanh^4(a+b \log(cx^n))}{x} dx$	1458
3.198	$\int \frac{\tanh^5(a+b \log(cx^n))}{x} dx$	1466
3.199	$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$	1475
3.200	$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx$	1480
3.201	$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx$	1487
3.202	$\int \tanh^p(d(a + b \log(cx^n))) dx$	1496
3.203	$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx$	1502
3.204	$\int \frac{\tanh^{\frac{5}{2}}(a+b \log(cx^n))}{x} dx$	1508
3.205	$\int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx$	1516
3.206	$\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{x} dx$	1524
3.207	$\int \frac{1}{x \sqrt{\tanh(a+b \log(cx^n))}} dx$	1531



3.208	$\int \frac{1}{x \tanh^{\frac{3}{2}}(a+b \log(cx^n))} dx$	1538
3.209	$\int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx$	1546
3.210	$\int \frac{\tanh^5(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1554
3.211	$\int \frac{\tanh^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1562
3.212	$\int \frac{\tanh(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1569
3.213	$\int \frac{\coth(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1576
3.214	$\int \frac{\coth^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1582
3.215	$\int \tanh(x) \sqrt{a+b \tanh^2(x)+c \tanh^4(x)} dx$	1589
3.216	$\int e^{a+bx} \tanh^4(a+bx) dx$	1598
3.217	$\int e^{a+bx} \tanh^3(a+bx) dx$	1604
3.218	$\int e^{a+bx} \tanh^2(a+bx) dx$	1610
3.219	$\int e^{a+bx} \tanh(a+bx) dx$	1616
3.220	$\int e^{a+bx} \coth(a+bx) dx$	1621
3.221	$\int e^{a+bx} \coth^2(a+bx) dx$	1626
3.222	$\int e^{a+bx} \coth^3(a+bx) dx$	1632
3.223	$\int e^{a+bx} \coth^4(a+bx) dx$	1638
3.224	$\int e^x \tanh^2(2x) dx$	1645
3.225	$\int e^x \tanh(2x) dx$	1651
3.226	$\int e^x \coth(2x) dx$	1659
3.227	$\int e^x \coth^2(2x) dx$	1665
3.228	$\int e^x \coth^4(2x) dx$	1671
3.229	$\int e^x \tanh^2(3x) dx$	1678
3.230	$\int e^x \tanh(3x) dx$	1685
3.231	$\int e^x \coth(3x) dx$	1693
3.232	$\int e^x \coth^2(3x) dx$	1701
3.233	$\int e^x \tanh^2(4x) dx$	1708
3.234	$\int e^x \tanh(4x) dx$	1716
3.235	$\int e^x \coth(4x) dx$	1727
3.236	$\int e^x \coth^2(4x) dx$	1737
3.237	$\int \frac{e^x}{a-\tanh(2x)} dx$	1744
3.238	$\int \frac{e^x}{(a-\tanh(2x))^2} dx$	1752
3.239	$\int e^{c(a+bx)} \tanh^3(d+ex) dx$	1761
3.240	$\int e^{c(a+bx)} \tanh^2(d+ex) dx$	1767
3.241	$\int e^{c(a+bx)} \tanh(d+ex) dx$	1773

3.242	$\int e^{c(a+bx)} \coth(d+ex) dx$	1778
3.243	$\int e^{c(a+bx)} \coth^2(d+ex) dx$	1783
3.244	$\int e^{c(a+bx)} \coth^3(d+ex) dx$	1789
3.245	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{5/2} dx$	1795
3.246	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{3/2} dx$	1803
3.247	$\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx$	1810
3.248	$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$	1816
3.249	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$	1822
3.250	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$	1830
3.251	$\int \sin^3(\tanh(a+bx)) dx$	1838
3.252	$\int \sin^2(\tanh(a+bx)) dx$	1845
3.253	$\int \sin(\tanh(a+bx)) dx$	1851
3.254	$\int \csc(\tanh(a+bx)) dx$	1857
3.255	$\int \cos^3(\tanh(a+bx)) dx$	1862
3.256	$\int \cos^2(\tanh(a+bx)) dx$	1869
3.257	$\int \cos(\tanh(a+bx)) dx$	1875
3.258	$\int \sec(\tanh(a+bx)) dx$	1881
<b>4</b>	<b>Appendix</b>	<b>1886</b>
4.1	Listing of Grading functions	1886
4.2	Links to plain text integration problems used in this report for each CAS	1904

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	11
1.2	Results . . . . .	12
1.3	Time and leaf size Performance . . . . .	16
1.4	Performance based on number of rules Rubi used . . . . .	18
1.5	Performance based on number of steps Rubi used . . . . .	19
1.6	Solved integrals histogram based on leaf size of result . . . . .	20
1.7	Solved integrals histogram based on CPU time used . . . . .	21
1.8	Leaf size vs. CPU time used . . . . .	22
1.9	list of integrals with no known antiderivative . . . . .	23
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	23
1.11	list of integrals solved by CAS but failed verification . . . . .	23
1.12	Timing . . . . .	24
1.13	Verification . . . . .	24
1.14	Important notes about some of the results . . . . .	25
1.15	Current tree layout of integration tests . . . . .	28
1.16	Design of the test system . . . . .	29

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 258 ]. This is test number [ 307 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 258 )	0.00 ( 0 )
Mathematica	99.61 ( 257 )	0.39 ( 1 )
Fricas	84.88 ( 219 )	15.12 ( 39 )
Maple	83.33 ( 215 )	16.67 ( 43 )
Giac	73.26 ( 189 )	26.74 ( 69 )
Mupad	71.71 ( 185 )	28.29 ( 73 )
Reduce	64.73 ( 167 )	35.27 ( 91 )
Maxima	62.40 ( 161 )	37.60 ( 97 )
Sympy	27.52 ( 71 )	72.48 ( 187 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Mathematica	79.845	7.364	11.628	1.163
Rubi	79.070	0.000	20.155	0.775
Maple	63.566	4.651	15.116	16.667
Giac	56.977	13.178	2.326	27.519
Maxima	43.798	15.504	2.326	38.372
Fricas	17.054	60.853	6.202	15.891
Sympy	13.566	13.178	0.000	73.256
Mupad	0.000	70.930	0.000	29.070
Reduce	0.000	63.953	0.000	36.047

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.



System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	100.00	0.00	0.00
Fricas	39	100.00	0.00	0.00
Maple	43	95.35	4.65	0.00
Giac	69	72.46	13.04	14.49
Mupad	73	0.00	100.00	0.00
Reduce	91	100.00	0.00	0.00
Maxima	97	84.54	0.00	15.46
Sympy	187	96.79	3.21	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.09
Fricas	0.13
Giac	0.13
Reduce	0.27
Rubi	0.42
Mathematica	0.76
Sympy	1.88
Mupad	1.89
Maple	3.05

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	75.13	1.07	48.00	0.95
Mathematica	83.07	1.18	59.00	1.00
Rubi	83.19	1.11	63.50	1.01
Mupad	84.63	1.36	50.00	1.00
Maxima	87.52	1.57	62.00	1.16
Giac	87.89	1.40	60.00	1.16
Reduce	241.93	3.13	101.00	2.05
Sympy	458.06	4.60	66.00	1.53
Fricas	708.95	8.12	216.00	4.50

Table 1.6: Leaf size performance for each CAS

# 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{254, 258}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {13, 14, 15, 16, 17, 18, 19, 20, 21, 210, 211, 212, 213, 214, 215}

**Mathematica** {173, 174, 178, 179, 180, 181, 202, 203, 222, 249, 250}

**Maple** {245, 246, 248, 249, 250}



**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	31
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	37
2.3	Detailed conclusion table specific for Rubi results . . . . .	102

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	31
Mma . . . . .	32
Maple . . . . .	32
Fricas . . . . .	33
Maxima . . . . .	33
Giac . . . . .	34
Mupad . . . . .	35
Sympy . . . . .	35
Reduce . . . . .	36

### Rubi

**A grade** { 1, 3, 5, 6, 8, 10, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 29, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 74, 76, 78, 80, 82, 85, 87, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 119, 121, 130, 131, 132, 133, 137, 139, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 197, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 255, 256, 257 }

**B grade** { }

**C grade** { 2, 4, 7, 9, 11, 24, 27, 28, 30, 31, 32, 70, 72, 73, 75, 77, 79, 81, 83, 84, 86, 88, 93, 115, 116, 117, 118, 120, 122, 123, 124, 125, 126, 127, 128, 129, 134, 135, 136, 138, 140, 141, 142, 143, 196, 198, 210, 211, 212, 213, 214, 215 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }



**Mma**

**A grade** { 1, 2, 3, 4, 5, 6, 7, 9, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 74, 76, 78, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 126, 127, 128, 130, 131, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 149, 150, 151, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 174, 175, 176, 177, 178, 186, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 223, 225, 226, 228, 237, 239, 240, 243, 244, 245, 246, 247, 248, 250, 251, 252, 253, 255, 256, 257 }

**B grade** { 73, 75, 77, 79, 148, 152, 173, 179, 180, 181, 182, 183, 184, 185, 187, 188, 202, 241, 242 }

**C grade** { 8, 10, 12, 39, 54, 55, 56, 123, 124, 125, 129, 132, 153, 154, 155, 156, 221, 222, 224, 227, 229, 230, 231, 232, 233, 234, 235, 236, 238, 249 }

**F normal fail** { 91 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

**Maple**

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 24, 25, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 90, 92, 93, 94, 95, 96, 97, 99, 101, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 148, 150, 152, 158, 160, 162, 164, 186, 193, 196, 197, 198, 204, 205, 206, 207, 208, 209, 210, 211, 212, 215, 218, 219, 220, 221, 222, 223, 251, 252, 253, 255, 256, 257 }

**B grade** { 26, 75, 89, 98, 100, 102, 103, 104, 105, 110, 145, 146 }

**C grade** { 147, 149, 151, 153, 154, 155, 156, 157, 159, 161, 163, 165, 166, 167, 168, 169, 216, 217, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 245, 246, 247, 248, 249, 250 }

**F normal fail** { 22, 23, 40, 91, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182,

183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 213, 214, 239, 240, 241, 242, 243, 244 }

**F(-1) timedout fail** { 254, 258 }

**F(-2) exception fail** { }

## Fricas

**A grade** { 61, 65, 66, 71, 72, 84, 90, 94, 95, 96, 107, 113, 137, 138, 139, 140, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 161, 162, 163, 164, 165, 166, 167, 168, 169, 219, 225, 230, 235, 247, 248 }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 24, 25, 26, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 67, 68, 69, 70, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 92, 93, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 141, 142, 143, 144, 145, 146, 160, 186, 193, 196, 197, 198, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 220, 221, 222, 223, 224, 226, 227, 228, 229, 231, 232, 236, 245, 246, 249, 250 }

**C grade** { 27, 28, 29, 30, 31, 32, 233, 234, 237, 238, 251, 252, 253, 255, 256, 257 }

**F normal fail** { 22, 23, 40, 91, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 239, 240, 241, 242, 243, 244 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maxima

**A grade** { 5, 6, 7, 8, 24, 25, 37, 38, 39, 44, 45, 46, 47, 48, 49, 50, 56, 59, 60, 61, 62, 65, 66, 69, 70, 71, 72, 73, 74, 80, 82, 90, 92, 93, 94, 95, 96, 97, 106, 107, 108, 109, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 186, 193, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 235, 236, 245, 246, 247, 248, 249, 250 }

**B grade** { 1, 2, 3, 4, 9, 10, 11, 12, 26, 41, 42, 43, 51, 52, 53, 54, 55, 57, 58, 63, 64, 75, 76, 77, 78, 79, 85, 87, 89, 98, 99, 100, 101, 102, 103, 104, 105, 196, 197, 198 }

**C grade** { 27, 28, 29, 30, 31, 32 }

**F normal fail** { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 33, 34, 35, 36, 40, 91, 126, 127, 128, 129, 130, 131, 132, 133, 145, 146, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 233, 234, 239, 240, 241, 242, 243, 244, 251, 252, 253, 255, 256, 257 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { 67, 68, 81, 83, 84, 86, 88, 110, 111, 112, 113, 114, 115, 237, 238 }

## Giac

**A grade** { 1, 2, 3, 4, 5, 8, 9, 10, 11, 12, 24, 26, 33, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 69, 70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 86, 88, 90, 92, 93, 94, 95, 96, 99, 100, 101, 102, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 132, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 193, 197, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 245, 246, 247, 248, 249, 250 }

**B grade** { 6, 7, 21, 25, 34, 35, 50, 51, 55, 56, 75, 85, 87, 89, 97, 98, 103, 104, 105, 106, 110, 126, 127, 128, 129, 130, 131, 133, 144, 186, 196, 198, 237, 238 }

**C grade** { 27, 28, 29, 30, 31, 32 }

**F normal fail** { 22, 23, 40, 91, 145, 146, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 210, 211, 215, 239, 240, 241, 242, 243, 244, 251, 252, 253, 255, 256, 257 }

**F(-1) timedout fail** { 204, 205, 206, 207, 208, 209, 212, 213, 214 }

**F(-2) exception fail** { 13, 14, 15, 16, 17, 18, 19, 20, 67, 68 }

## Mupad

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 26, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 186, 193, 196, 197, 198, 204, 205, 206, 207, 208, 209, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 22, 23, 24, 25, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 91, 145, 146, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 210, 211, 212, 213, 214, 215, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 255, 256, 257 }

**F(-2) exception fail** { }

## Sympy

**A grade** { 1, 2, 3, 4, 5, 21, 41, 42, 43, 44, 57, 58, 59, 60, 65, 66, 96, 126, 127, 128, 129, 130, 131, 132, 133, 150, 160, 186, 196, 197, 198, 205, 206, 207, 208 }

**B grade** { 6, 7, 8, 9, 10, 11, 12, 45, 46, 47, 48, 49, 61, 62, 63, 64, 70, 72, 93, 95, 107, 116, 117, 118, 119, 120, 121, 134, 135, 136, 137, 138, 139, 193 }

**C grade** { }

**F normal fail** { 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 50, 51, 52, 53, 54, 55, 56, 67, 68, 69, 71, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 94, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 108, 109, 110, 111, 112, 113, 114, 115, 122, 123, 124, 125, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 151, 152, 153, 154, 155, 156, 157, 158, 159, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 210, 211, 212, 213, 214, 215, 216, 217,

218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 247, 248, 249, 251, 252, 253, 255, 256, 257 }

**F(-1) timedout fail** { 203, 204, 209, 245, 246, 250 }

**F(-2) exception fail** { }

## Reduce

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 24, 25, 26, 27, 28, 29, 30, 31, 32, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 134, 135, 136, 137, 138, 139, 140, 141, 144, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 186, 193, 196, 197, 198, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 245, 246, 247, 248, 249, 250 }

**C grade** { }

**F normal fail** { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 33, 34, 35, 36, 40, 50, 51, 52, 53, 54, 55, 56, 67, 68, 91, 126, 127, 128, 129, 130, 131, 132, 133, 142, 143, 145, 146, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 192, 194, 195, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 239, 240, 241, 242, 243, 244, 251, 252, 253, 255, 256, 257 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	53	39	115	254	39	74	38	34
N.S.	1	1.00	1.23	0.91	2.67	5.91	0.91	1.72	0.88	0.79
time (sec)	N/A	0.328	0.016	0.152	0.036	0.083	0.131	0.132	0.229	0.098

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	53	37	42	102	968	42	67	252	37
N.S.	1	1.26	0.88	1.00	2.43	23.05	1.00	1.60	6.00	0.88
time (sec)	N/A	0.350	0.036	0.154	0.116	0.103	0.108	0.125	0.214	2.166

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	38	27	71	119	27	52	28	24
N.S.	1	1.00	1.36	0.96	2.54	4.25	0.96	1.86	1.00	0.86
time (sec)	N/A	0.264	0.010	0.136	0.040	0.089	0.092	0.125	0.286	0.062

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	36	25	32	61	339	31	48	138	27
N.S.	1	1.33	0.93	1.19	2.26	12.56	1.15	1.78	5.11	1.00
time (sec)	N/A	0.263	0.010	0.132	0.120	0.099	0.086	0.127	0.300	2.072

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	23	17	25	33	15	24	16	13
N.S.	1	1.00	1.77	1.31	1.92	2.54	1.15	1.85	1.23	1.00
time (sec)	N/A	0.208	0.001	0.126	0.048	0.081	0.075	0.130	0.258	0.065

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	37	17	24	22	16
N.S.	1	1.00	1.00	1.09	1.00	3.36	1.55	2.18	2.00	1.45
time (sec)	N/A	0.189	0.001	0.162	0.028	0.086	0.065	0.117	0.252	2.076

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	15	11	12	11	37	27	25	29	21
N.S.	1	1.36	1.00	1.09	1.00	3.36	2.45	2.27	2.64	1.91
time (sec)	N/A	0.189	0.001	0.280	0.033	0.098	0.189	0.133	0.304	0.040

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	27	17	25	33	65	24	16	13
N.S.	1	1.00	2.08	1.31	1.92	2.54	5.00	1.85	1.23	1.00
time (sec)	N/A	0.198	0.001	0.164	0.036	0.091	0.511	0.133	0.218	0.057

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	40	25	38	79	346	90	49	182	68
N.S.	1	1.48	0.93	1.41	2.93	12.81	3.33	1.81	6.74	2.52
time (sec)	N/A	0.277	0.015	0.253	0.037	0.104	0.700	0.132	0.243	0.048

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	31	29	71	108	78	52	28	24
N.S.	1	1.00	1.11	1.04	2.54	3.86	2.79	1.86	1.00	0.86
time (sec)	N/A	0.282	0.009	0.212	0.041	0.090	1.091	0.123	0.329	0.061

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	57	35	48	122	978	104	70	334	159
N.S.	1	1.36	0.83	1.14	2.90	23.29	2.48	1.67	7.95	3.79
time (sec)	N/A	0.366	0.033	0.277	0.043	0.097	1.605	0.131	0.262	2.085



Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	31	39	115	239	92	74	38	34
N.S.	1	1.00	0.72	0.91	2.67	5.56	2.14	1.72	0.88	0.79
time (sec)	N/A	0.340	0.010	0.239	0.041	0.085	2.454	0.131	0.233	2.112

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	80	80	0	1545	0	0	58	83
N.S.	1	1.00	0.82	0.82	0.00	15.93	0.00	0.00	0.60	0.86
time (sec)	N/A	0.452	0.265	0.444	0.000	0.131	0.000	0.000	0.280	2.631

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	74	66	63	0	969	0	0	24	62
N.S.	1	0.95	0.85	0.81	0.00	12.42	0.00	0.00	0.31	0.79
time (sec)	N/A	0.360	0.108	0.389	0.000	0.129	0.000	0.000	0.254	2.345

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	72	66	62	0	626	0	0	37	61
N.S.	1	0.96	0.88	0.83	0.00	8.35	0.00	0.00	0.49	0.81
time (sec)	N/A	0.346	0.051	0.374	0.000	0.108	0.000	0.000	0.242	2.295

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	51	52	47	0	583	0	0	12	41
N.S.	1	0.88	0.90	0.81	0.00	10.05	0.00	0.00	0.21	0.71
time (sec)	N/A	0.237	0.027	0.480	0.000	0.134	0.000	0.000	0.267	2.215

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	51	49	46	0	587	0	0	24	38
N.S.	1	0.89	0.86	0.81	0.00	10.30	0.00	0.00	0.42	0.67
time (sec)	N/A	0.253	0.025	0.451	0.000	0.102	0.000	0.000	0.249	2.384

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	74	74	65	0	914	0	0	24	64
N.S.	1	0.95	0.95	0.83	0.00	11.72	0.00	0.00	0.31	0.82
time (sec)	N/A	0.329	0.060	0.385	0.000	0.130	0.000	0.000	0.248	2.340

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	76	78	64	0	1425	0	0	24	63
N.S.	1	0.96	0.99	0.81	0.00	18.04	0.00	0.00	0.30	0.80
time (sec)	N/A	0.338	0.075	0.389	0.000	0.129	0.000	0.000	0.227	2.454

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-2)</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	101	90	83	0	2133	0	0	24	80
N.S.	1	1.01	0.90	0.83	0.00	21.33	0.00	0.00	0.24	0.80
time (sec)	N/A	0.447	0.176	0.393	0.000	0.142	0.000	0.000	0.293	2.684

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	70	90	102	0	179	63	110	8	71
N.S.	1	1.01	1.30	1.48	0.00	2.59	0.91	1.59	0.12	1.03
time (sec)	N/A	0.298	0.075	0.160	0.000	0.083	1.046	0.132	0.244	2.598

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	0	0	0	0	0	10	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.226	0.093	0.000	0.000	0.000	0.000	0.000	0.256	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	49	0	0	0	0	0	14	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.237	0.085	0.000	0.000	0.000	0.000	0.000	0.267	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	28	30	42	467	0	52	85	0
N.S.	1	1.00	0.80	0.86	1.20	13.34	0.00	1.49	2.43	0.00
time (sec)	N/A	0.303	0.024	0.288	0.150	0.121	0.000	0.119	0.225	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	26	19	72	0	31	15	0
N.S.	1	1.00	1.00	1.62	1.19	4.50	0.00	1.94	0.94	0.00
time (sec)	N/A	0.240	0.008	0.275	0.127	0.108	0.000	0.129	0.250	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	33	31	76	0	1	22	14
N.S.	1	1.00	1.00	2.06	1.94	4.75	0.00	0.06	1.38	0.88
time (sec)	N/A	0.263	0.008	0.268	0.136	0.103	0.000	0.129	0.237	2.131

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	C	C	<b>F</b>	C	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	73	59	67	113	180	0	142	253	0
N.S.	1	0.83	0.67	0.76	1.28	2.05	0.00	1.61	2.88	0.00
time (sec)	N/A	0.481	0.059	0.395	0.131	0.094	0.000	0.160	0.249	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	C	C	<b>F</b>	C	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	56	47	53	66	100	0	92	138	0
N.S.	1	0.93	0.78	0.88	1.10	1.67	0.00	1.53	2.30	0.00
time (sec)	N/A	0.371	0.063	0.292	0.140	0.086	0.000	0.147	0.246	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	C	<b>F</b>	C	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	45	28	23	0	54	23	0
N.S.	1	1.00	1.00	1.45	0.90	0.74	0.00	1.74	0.74	0.00
time (sec)	N/A	0.239	0.023	0.321	0.127	0.110	0.000	0.129	0.254	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	C	C	<b>F</b>	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	35	31	52	45	23	0	61	33	24
N.S.	1	1.13	1.00	1.68	1.45	0.74	0.00	1.97	1.06	0.77
time (sec)	N/A	0.250	0.042	0.323	0.176	0.117	0.000	0.132	0.302	2.063

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	C	C	<b>F</b>	C	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	47	81	85	100	0	100	183	0
N.S.	1	1.00	0.78	1.35	1.42	1.67	0.00	1.67	3.05	0.00
time (sec)	N/A	0.331	0.080	0.313	0.128	0.098	0.000	0.132	0.252	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	C	C	<b>F</b>	C	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	77	57	91	132	180	0	126	337	0
N.S.	1	0.88	0.65	1.03	1.50	2.05	0.00	1.43	3.83	0.00
time (sec)	N/A	0.414	0.087	0.342	0.136	0.103	0.000	0.147	0.257	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	49	38	43	0	106	0	55	16	0
N.S.	1	0.86	0.67	0.75	0.00	1.86	0.00	0.96	0.28	0.00
time (sec)	N/A	0.319	0.029	0.319	0.000	0.088	0.000	0.132	0.255	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	B	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	64	55	76	0	1267	0	342	14	0
N.S.	1	0.74	0.64	0.88	0.00	14.73	0.00	3.98	0.16	0.00
time (sec)	N/A	0.395	0.048	0.351	0.000	0.122	0.000	0.205	0.248	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	B	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	51	40	62	0	365	0	115	19	0
N.S.	1	0.81	0.63	0.98	0.00	5.79	0.00	1.83	0.30	0.00
time (sec)	N/A	0.312	0.026	0.323	0.000	0.106	0.000	0.161	0.281	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	51	53	65	0	514	0	41	16	0
N.S.	1	0.80	0.83	1.02	0.00	8.03	0.00	0.64	0.25	0.00
time (sec)	N/A	0.327	0.031	0.345	0.000	0.108	0.000	0.143	0.256	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	39	39	46	82	2114	0	45	25	0
N.S.	1	0.57	0.57	0.67	1.19	30.64	0.00	0.65	0.36	0.00
time (sec)	N/A	0.367	0.057	0.308	0.129	0.166	0.000	0.125	0.242	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	22	21	32	19	213	0	16	9	0
N.S.	1	0.71	0.68	1.03	0.61	6.87	0.00	0.52	0.29	0.00
time (sec)	N/A	0.240	0.014	0.282	0.156	0.095	0.000	0.118	0.269	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	22	26	32	23	238	0	19	16	0
N.S.	1	0.71	0.84	1.03	0.74	7.68	0.00	0.61	0.52	0.00
time (sec)	N/A	0.239	0.017	0.275	0.145	0.120	0.000	0.135	0.258	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	57	0	0	0	0	0	16	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.307	0.077	0.000	0.000	0.000	0.000	0.000	0.278	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	108	63	58	302	907	95	85	211	65
N.S.	1	1.08	0.63	0.58	3.02	9.07	0.95	0.85	2.11	0.65
time (sec)	N/A	0.593	0.217	0.329	0.134	0.086	0.131	0.134	0.241	0.148

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	84	51	48	196	562	76	71	162	53
N.S.	1	1.09	0.66	0.62	2.55	7.30	0.99	0.92	2.10	0.69
time (sec)	N/A	0.470	0.123	0.203	0.142	0.085	0.109	0.130	0.312	1.967

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	60	40	38	116	299	61	57	113	43
N.S.	1	1.07	0.71	0.68	2.07	5.34	1.09	1.02	2.02	0.77
time (sec)	N/A	0.358	0.151	0.199	0.127	0.087	0.112	0.133	0.269	1.963



Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	29	28	50	117	44	39	72	33
N.S.	1	1.00	0.81	0.78	1.39	3.25	1.22	1.08	2.00	0.92
time (sec)	N/A	0.270	0.116	0.158	0.040	0.095	0.085	0.126	0.235	0.086

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	33	25	31	50	73	30	36	25
N.S.	1	1.00	1.18	0.89	1.11	1.79	2.61	1.07	1.29	0.89
time (sec)	N/A	0.198	0.065	0.168	0.034	0.097	0.363	0.126	0.218	1.956

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	55	47	42	43	101	223	42	48	41
N.S.	1	1.08	0.92	0.82	0.84	1.98	4.37	0.82	0.94	0.80
time (sec)	N/A	0.281	0.134	0.183	0.044	0.079	0.587	0.130	0.244	1.956

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	82	59	59	56	160	430	53	60	58
N.S.	1	1.12	0.81	0.81	0.77	2.19	5.89	0.73	0.82	0.79
time (sec)	N/A	0.377	0.170	0.211	0.041	0.096	0.738	0.127	0.252	1.978

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	109	96	76	67	220	694	64	72	75
N.S.	1	1.14	1.00	0.79	0.70	2.29	7.23	0.67	0.75	0.78
time (sec)	N/A	0.479	0.168	0.257	0.046	0.096	0.981	0.130	0.225	1.968

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	136	111	91	78	287	1018	75	84	92
N.S.	1	1.12	0.92	0.75	0.64	2.37	8.41	0.62	0.69	0.76
time (sec)	N/A	0.616	0.199	0.302	0.050	0.108	1.313	0.133	0.213	1.994

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	63	47	43	83	331	0	140	46	44
N.S.	1	1.11	0.82	0.75	1.46	5.81	0.00	2.46	0.81	0.77
time (sec)	N/A	0.474	0.378	0.231	0.122	0.098	0.000	0.134	0.272	0.164

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	48	39	35	70	206	0	96	32	54
N.S.	1	1.07	0.87	0.78	1.56	4.58	0.00	2.13	0.71	1.20
time (sec)	N/A	0.358	0.293	0.204	0.168	0.110	0.000	0.128	0.253	1.949

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	27	57	127	0	52	18	26
N.S.	1	1.00	1.00	0.82	1.73	3.85	0.00	1.58	0.55	0.79
time (sec)	N/A	0.273	0.213	0.202	0.116	0.102	0.000	0.131	0.240	0.095

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	17	43	99	0	27	7	16
N.S.	1	1.00	1.00	0.81	2.05	4.71	0.00	1.29	0.33	0.76
time (sec)	N/A	0.202	0.152	0.260	0.110	0.089	0.000	0.141	0.241	0.115

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	26	27	57	141	0	50	26	26
N.S.	1	1.00	0.81	0.84	1.78	4.41	0.00	1.56	0.81	0.81
time (sec)	N/A	0.270	0.175	0.239	0.122	0.105	0.000	0.129	0.236	2.004

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	B	B	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	28	35	69	244	0	95	66	32
N.S.	1	1.00	0.57	0.71	1.41	4.98	0.00	1.94	1.35	0.65
time (sec)	N/A	0.338	0.214	0.226	0.124	0.113	0.000	0.137	0.209	2.021

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	66	28	43	79	338	0	139	125	40
N.S.	1	1.08	0.46	0.70	1.30	5.54	0.00	2.28	2.05	0.66
time (sec)	N/A	0.418	0.238	0.219	0.118	0.096	0.000	0.128	0.223	0.111

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	114	182	310	2739	211	224	1151	153
N.S.	1	1.00	0.80	1.28	2.18	19.29	1.49	1.58	8.11	1.08
time (sec)	N/A	0.835	0.442	0.334	0.147	0.105	0.152	0.135	0.322	2.073

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	91	132	201	1389	144	152	652	113
N.S.	1	1.00	0.90	1.31	1.99	13.75	1.43	1.50	6.46	1.12
time (sec)	N/A	0.592	0.241	0.253	0.121	0.108	0.134	0.134	0.229	2.019

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	67	93	118	646	100	97	390	77
N.S.	1	1.00	0.97	1.35	1.71	9.36	1.45	1.41	5.65	1.12
time (sec)	N/A	0.411	0.231	0.235	0.128	0.095	0.103	0.128	0.223	2.013

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	54	52	49	201	54	56	144	44
N.S.	1	1.00	1.42	1.37	1.29	5.29	1.42	1.47	3.79	1.16
time (sec)	N/A	0.272	0.073	0.192	0.037	0.094	0.085	0.115	0.255	1.985

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	64	55	56	62	224	62	57	60
N.S.	1	1.00	1.28	1.10	1.12	1.24	4.48	1.24	1.14	1.20
time (sec)	N/A	0.340	0.062	0.196	0.036	0.104	1.035	0.129	0.239	0.165

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	98	94	93	127	422	1389	132	428	127
N.S.	1	1.15	1.11	1.09	1.49	4.96	16.34	1.55	5.04	1.49
time (sec)	N/A	0.497	0.712	0.227	0.048	0.094	9.207	0.129	0.226	2.384

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	155	122	130	325	1427	6412	205	1186	304
N.S.	1	1.20	0.95	1.01	2.52	11.06	49.71	1.59	9.19	2.36
time (sec)	N/A	0.733	1.565	0.297	0.081	0.114	12.427	0.139	0.256	2.991

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	208	160	163	525	3693	16643	305	2312	452
N.S.	1	1.23	0.95	0.96	3.11	21.85	98.48	1.80	13.68	2.67
time (sec)	N/A	1.004	2.218	0.348	0.084	0.151	19.447	0.161	0.240	3.976

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	53	28	28	49	42	30	42	34
N.S.	1	1.00	1.71	0.90	0.90	1.58	1.35	0.97	1.35	1.10
time (sec)	N/A	0.291	0.023	0.244	0.038	0.124	0.264	0.127	0.219	0.128

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	53	28	29	48	42	25	37	33
N.S.	1	1.00	1.71	0.90	0.94	1.55	1.35	0.81	1.19	1.06
time (sec)	N/A	0.288	0.023	0.202	0.039	0.080	0.244	0.121	0.219	0.109

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F(-2)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	90	74	63	0	2203	0	0	13	151
N.S.	1	1.22	1.00	0.85	0.00	29.77	0.00	0.00	0.18	2.04
time (sec)	N/A	0.294	0.065	0.602	0.000	0.155	0.000	0.000	0.272	2.300

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F(-2)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	85	74	62	0	2279	0	0	58	240
N.S.	1	1.15	1.00	0.84	0.00	30.80	0.00	0.00	0.78	3.24
time (sec)	N/A	0.261	0.042	0.454	0.000	0.136	0.000	0.000	0.231	2.383

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	62	42	35	36	92	0	42	47	34
N.S.	1	1.03	0.70	0.58	0.60	1.53	0.00	0.70	0.78	0.57
time (sec)	N/A	0.283	0.083	3.898	0.041	0.081	0.000	0.122	0.212	2.342

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	35	34	24	27	56	134	25	32	23
N.S.	1	1.40	1.36	0.96	1.08	2.24	5.36	1.00	1.28	0.92
time (sec)	N/A	0.451	0.078	1.287	0.040	0.084	0.330	0.112	0.241	2.113

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	40	24	23	22	51	0	30	33	22
N.S.	1	1.05	0.63	0.61	0.58	1.34	0.00	0.79	0.87	0.58
time (sec)	N/A	0.296	0.013	0.709	0.029	0.090	0.000	0.123	0.225	2.151

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	25	19	12	11	23	48	11	34	11
N.S.	1	1.47	1.12	0.71	0.65	1.35	2.82	0.65	2.00	0.65
time (sec)	N/A	0.446	0.012	0.345	0.036	0.076	0.205	0.123	0.235	2.015

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	24	49	17	21	38	0	18	29	21
N.S.	1	2.00	4.08	1.42	1.75	3.17	0.00	1.50	2.42	1.75
time (sec)	N/A	0.419	0.103	0.292	0.052	0.091	0.000	0.120	0.269	0.059

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	11	24	29	77	0	29	67	23
N.S.	1	1.00	0.73	1.60	1.93	5.13	0.00	1.93	4.47	1.53
time (sec)	N/A	0.254	0.009	0.349	0.051	0.087	0.000	0.114	0.321	2.040

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	30	59	33	48	209	0	34	97	48
N.S.	1	1.67	3.28	1.83	2.67	11.61	0.00	1.89	5.39	2.67
time (sec)	N/A	0.486	0.201	0.533	0.048	0.093	0.000	0.115	0.241	0.077



Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	20	19	75	84	0	18	36	18
N.S.	1	1.00	1.18	1.12	4.41	4.94	0.00	1.06	2.12	1.06
time (sec)	N/A	0.257	0.172	1.103	0.047	0.074	0.000	0.121	0.235	2.027

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	46	69	48	74	640	0	49	180	117
N.S.	1	1.35	2.03	1.41	2.18	18.82	0.00	1.44	5.29	3.44
time (sec)	N/A	0.521	0.320	2.054	0.054	0.090	0.000	0.119	0.225	2.060

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	27	25	149	185	0	24	57	24
N.S.	1	1.00	0.82	0.76	4.52	5.61	0.00	0.73	1.73	0.73
time (sec)	N/A	0.277	0.227	4.459	0.034	0.084	0.000	0.116	0.211	0.090

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	B	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	58	124	60	98	1260	0	61	260	207
N.S.	1	1.32	2.82	1.36	2.23	28.64	0.00	1.39	5.91	4.70
time (sec)	N/A	0.544	0.356	8.686	0.040	0.097	0.000	0.607	0.242	0.094

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	242	144	183	163	1226	0	214	335	135
N.S.	1	1.65	0.98	1.24	1.11	8.34	0.00	1.46	2.28	0.92
time (sec)	N/A	0.694	0.440	12.167	0.062	0.107	0.000	0.128	0.242	2.508

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	153	180	166	0	1861	0	163	317	261
N.S.	1	1.12	1.31	1.21	0.00	13.58	0.00	1.19	2.31	1.91
time (sec)	N/A	0.920	0.816	4.106	0.000	0.134	0.000	0.121	0.232	2.367

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	142	73	98	83	334	0	101	152	81
N.S.	1	1.69	0.87	1.17	0.99	3.98	0.00	1.20	1.81	0.96
time (sec)	N/A	0.391	0.114	1.228	0.043	0.120	0.000	0.127	0.244	2.333

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	88	79	92	0	427	0	60	171	157
N.S.	1	1.22	1.10	1.28	0.00	5.93	0.00	0.83	2.38	2.18
time (sec)	N/A	0.518	0.146	0.535	0.000	0.134	0.000	0.125	0.247	2.269

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	60	68	53	0	237	0	60	96	177
N.S.	1	1.15	1.31	1.02	0.00	4.56	0.00	1.15	1.85	3.40
time (sec)	N/A	0.392	0.161	0.472	0.000	0.137	0.000	0.128	0.262	2.467

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	34	28	50	65	122	0	78	116	323
N.S.	1	1.17	0.97	1.72	2.24	4.21	0.00	2.69	4.00	11.14
time (sec)	N/A	0.261	0.304	0.556	0.038	0.136	0.000	0.123	0.282	2.444

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F(-2)	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	98	147	110	0	1165	0	125	350	506
N.S.	1	1.20	1.79	1.34	0.00	14.21	0.00	1.52	4.27	6.17
time (sec)	N/A	0.547	0.494	1.209	0.000	0.125	0.000	0.120	0.244	2.478

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	87	70	143	161	912	0	202	567	123
N.S.	1	1.12	0.90	1.83	2.06	11.69	0.00	2.59	7.27	1.58
time (sec)	N/A	0.331	2.502	2.564	0.046	0.099	0.000	0.125	0.294	2.325

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	218	299	335	227	0	5347	0	273	1133	753
N.S.	1	1.37	1.54	1.04	0.00	24.53	0.00	1.25	5.20	3.45
time (sec)	N/A	0.836	0.983	6.108	0.000	0.196	0.000	0.123	0.233	4.338

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	139	119	277	308	2972	0	412	1407	237
N.S.	1	1.07	0.92	2.13	2.37	22.86	0.00	3.17	10.82	1.82
time (sec)	N/A	0.393	5.441	12.853	0.053	0.139	0.000	0.130	0.244	2.391

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	46	29	34	43	0	28	111	61
N.S.	1	1.00	1.39	0.88	1.03	1.30	0.00	0.85	3.36	1.85
time (sec)	N/A	0.346	0.195	0.888	0.124	0.101	0.000	0.114	0.252	0.427

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	153	157	0	0	0	0	0	0	27	0
N.S.	1	1.03	0.00	0.00	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	0.437	0.000	0.000	0.000	0.000	0.000	0.000	0.314	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	62	53	35	36	95	0	42	47	34
N.S.	1	1.03	0.88	0.58	0.60	1.58	0.00	0.70	0.78	0.57
time (sec)	N/A	0.291	0.083	3.799	0.036	0.097	0.000	0.122	0.306	2.329

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	38	36	30	33	60	134	31	39	29
N.S.	1	1.31	1.24	1.03	1.14	2.07	4.62	1.07	1.34	1.00
time (sec)	N/A	0.286	0.090	1.314	0.037	0.082	0.313	0.123	0.272	2.194

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	40	32	23	22	52	0	30	33	22
N.S.	1	1.05	0.84	0.61	0.58	1.37	0.00	0.79	0.87	0.58
time (sec)	N/A	0.272	0.082	0.700	0.040	0.076	0.000	0.123	0.251	2.123

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	23	18	17	25	48	19	25	17
N.S.	1	1.00	1.21	0.95	0.89	1.32	2.53	1.00	1.32	0.89
time (sec)	N/A	0.259	0.066	0.350	0.042	0.088	0.189	0.126	0.216	2.169

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	7	7	6	9	8	6	7	6
N.S.	1	1.00	0.70	0.70	0.60	0.90	0.80	0.60	0.70	0.60
time (sec)	N/A	0.222	0.021	0.722	0.034	0.077	0.158	0.115	0.256	0.036

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	5	5	7	6	5	20	0	13	14	13
N.S.	1	1.00	1.40	1.20	1.00	4.00	0.00	2.60	2.80	2.60
time (sec)	N/A	0.225	0.006	1.903	0.038	0.091	0.000	0.120	0.294	0.061

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	12	21	22	48	0	18	29	18
N.S.	1	1.00	2.00	3.50	3.67	8.00	0.00	3.00	4.83	3.00
time (sec)	N/A	0.269	0.102	4.615	0.127	0.085	0.000	0.119	0.245	2.226

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	10	10	37	53	0	10	18	16
N.S.	1	1.00	0.83	0.83	3.08	4.42	0.00	0.83	1.50	1.33
time (sec)	N/A	0.225	0.029	1.993	0.029	0.081	0.000	0.116	0.234	2.473

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	41	49	288	0	31	85	61
N.S.	1	1.00	1.00	1.71	2.04	12.00	0.00	1.29	3.54	2.54
time (sec)	N/A	0.303	0.104	54.644	0.116	0.079	0.000	0.114	0.241	0.106

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	19	93	140	0	18	43	18
N.S.	1	1.00	0.96	0.76	3.72	5.60	0.00	0.72	1.72	0.72
time (sec)	N/A	0.233	0.038	8.989	0.034	0.076	0.000	0.121	0.233	0.079

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	39	34	67	73	670	0	45	135	137
N.S.	1	1.15	1.00	1.97	2.15	19.71	0.00	1.32	3.97	4.03
time (sec)	N/A	0.394	0.110	0.169	0.115	0.079	0.000	0.123	0.268	2.248

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	139	127	413	386	5275	0	593	1798	301
N.S.	1	0.99	0.91	2.95	2.76	37.68	0.00	4.24	12.84	2.15
time (sec)	N/A	0.374	0.425	0.234	0.145	0.156	0.000	0.132	0.237	2.818

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	82	80	222	204	1827	0	316	893	169
N.S.	1	0.99	0.96	2.67	2.46	22.01	0.00	3.81	10.76	2.04
time (sec)	N/A	0.310	0.215	104.511	0.130	0.112	0.000	0.132	0.255	2.349

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	39	41	99	89	430	0	104	312	88
N.S.	1	0.98	1.02	2.48	2.22	10.75	0.00	2.60	7.80	2.20
time (sec)	N/A	0.267	0.086	21.299	0.116	0.109	0.000	0.125	0.229	2.356

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	12	11	42	0	45	35	50
N.S.	1	1.00	1.00	1.09	1.00	3.82	0.00	4.09	3.18	4.55
time (sec)	N/A	0.217	0.003	3.021	0.031	0.117	0.000	0.126	0.211	0.191

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	49	42	41	42	146	43	42	35
N.S.	1	1.00	1.26	1.08	1.05	1.08	3.74	1.10	1.08	0.90
time (sec)	N/A	0.315	0.067	0.151	0.034	0.081	0.234	0.127	0.252	0.149



Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	117	75	104	86	331	0	111	150	84
N.S.	1	1.18	0.76	1.05	0.87	3.34	0.00	1.12	1.52	0.85
time (sec)	N/A	0.380	0.145	1.228	0.044	0.108	0.000	0.129	0.238	2.514

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	195	206	191	165	1281	0	227	355	143
N.S.	1	1.16	1.23	1.14	0.98	7.62	0.00	1.35	2.11	0.85
time (sec)	N/A	0.514	0.246	11.934	0.038	0.108	0.000	0.132	0.302	2.619

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	165	166	323	0	6509	0	326	1373	447
N.S.	1	1.05	1.06	2.06	0.00	41.46	0.00	2.08	8.75	2.85
time (sec)	N/A	1.531	0.435	199.657	0.000	0.256	0.000	0.134	0.248	8.307

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	103	116	164	0	2043	0	152	564	265
N.S.	1	1.01	1.14	1.61	0.00	20.03	0.00	1.49	5.53	2.60
time (sec)	N/A	0.939	0.266	48.640	0.000	0.138	0.000	0.118	0.218	5.892

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	65	77	0	309	0	63	116	119
N.S.	1	1.00	1.16	1.38	0.00	5.52	0.00	1.12	2.07	2.12
time (sec)	N/A	0.529	0.143	8.367	0.000	0.110	0.000	0.123	0.261	4.679

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	46	39	0	148	0	35	48	35
N.S.	1	1.00	1.24	1.05	0.00	4.00	0.00	0.95	1.30	0.95
time (sec)	N/A	0.248	0.017	0.783	0.000	0.093	0.000	0.125	0.274	0.120

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	64	80	93	0	435	0	61	127	157
N.S.	1	0.88	1.10	1.27	0.00	5.96	0.00	0.84	1.74	2.15
time (sec)	N/A	0.532	0.214	0.591	0.000	0.098	0.000	0.121	0.231	2.271

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F(-2)</b>	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	125	258	172	0	1871	0	162	322	221
N.S.	1	0.95	1.95	1.30	0.00	14.17	0.00	1.23	2.44	1.67
time (sec)	N/A	0.958	0.360	4.119	0.000	0.126	0.000	0.121	0.258	3.127

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	70	52	40	55	571	104	47	62	35
N.S.	1	1.63	1.21	0.93	1.28	13.28	2.42	1.09	1.44	0.81
time (sec)	N/A	0.577	0.124	0.213	0.111	0.092	0.209	0.120	0.223	0.100

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	49	45	30	43	354	85	39	51	29
N.S.	1	1.32	1.22	0.81	1.16	9.57	2.30	1.05	1.38	0.78
time (sec)	N/A	0.501	0.083	0.189	0.109	0.103	0.207	0.120	0.247	0.083

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	44	41	28	29	186	75	35	40	21
N.S.	1	1.42	1.32	0.90	0.94	6.00	2.42	1.13	1.29	0.68
time (sec)	N/A	0.384	0.109	0.168	0.130	0.088	0.203	0.120	0.226	2.025

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	24	34	18	17	73	61	17	31	21
N.S.	1	1.26	1.79	0.95	0.89	3.84	3.21	0.89	1.63	1.11
time (sec)	N/A	0.232	0.062	0.186	0.119	0.089	0.170	0.131	0.255	0.070

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	24	14	11	10	26	27	10	19	10
N.S.	1	1.50	0.88	0.69	0.62	1.62	1.69	0.62	1.19	0.62
time (sec)	N/A	0.210	0.070	0.155	0.028	0.076	0.162	0.117	0.254	0.063

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	11	10	26	27	10	19	10
N.S.	1	1.00	1.12	0.69	0.62	1.62	1.69	0.62	1.19	0.62
time (sec)	N/A	0.184	0.025	0.137	0.029	0.096	0.165	0.123	0.202	0.055

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	32	34	18	24	73	0	18	45	17
N.S.	1	1.68	1.79	0.95	1.26	3.84	0.00	0.95	2.37	0.89
time (sec)	N/A	0.299	0.093	0.398	0.029	0.107	0.000	0.118	0.234	0.064

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	39	47	30	38	196	0	36	95	29
N.S.	1	1.34	1.62	1.03	1.31	6.76	0.00	1.24	3.28	1.00
time (sec)	N/A	0.412	0.197	0.458	0.044	0.091	0.000	0.117	0.237	2.062

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	60	54	30	54	357	0	40	141	35
N.S.	1	1.62	1.46	0.81	1.46	9.65	0.00	1.08	3.81	0.95
time (sec)	N/A	0.534	0.206	0.503	0.034	0.103	0.000	0.129	0.256	2.015

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	67	53	44	64	582	0	48	191	69
N.S.	1	1.56	1.23	1.02	1.49	13.53	0.00	1.12	4.44	1.60
time (sec)	N/A	0.607	0.223	0.536	0.037	0.098	0.000	0.123	0.237	2.081

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	56	39	35	0	206	58	96	23	34
N.S.	1	1.24	0.87	0.78	0.00	4.58	1.29	2.13	0.51	0.76
time (sec)	N/A	0.352	0.493	0.237	0.000	0.087	3.556	0.135	0.235	0.141

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	41	32	26	0	128	48	53	10	25
N.S.	1	1.28	1.00	0.81	0.00	4.00	1.50	1.66	0.31	0.78
time (sec)	N/A	0.264	0.293	0.241	0.000	0.086	0.751	0.130	0.211	2.067

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	41	30	25	0	141	46	52	16	24
N.S.	1	1.37	1.00	0.83	0.00	4.70	1.53	1.73	0.53	0.80
time (sec)	N/A	0.275	0.348	0.250	0.000	0.092	1.146	0.130	0.268	0.129

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	57	36	35	0	240	60	73	22	32
N.S.	1	1.16	0.73	0.71	0.00	4.90	1.22	1.49	0.45	0.65
time (sec)	N/A	0.339	0.389	0.224	0.000	0.091	4.726	0.120	0.275	2.127

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	45	35	0	329	58	140	25	34
N.S.	1	1.00	1.00	0.78	0.00	7.31	1.29	3.11	0.56	0.76
time (sec)	N/A	0.352	0.611	0.233	0.000	0.097	4.874	0.136	0.266	0.146

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	34	26	0	194	49	96	12	25
N.S.	1	1.00	1.00	0.76	0.00	5.71	1.44	2.82	0.35	0.74
time (sec)	N/A	0.264	0.372	0.245	0.000	0.092	0.976	0.144	0.243	0.105

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	34	35	0	161	58	54	18	36
N.S.	1	1.00	0.81	0.83	0.00	3.83	1.38	1.29	0.43	0.86
time (sec)	N/A	0.344	0.478	0.269	0.000	0.116	1.375	0.136	0.320	0.123

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	A	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	48	35	0	244	60	95	24	31
N.S.	1	1.00	0.98	0.71	0.00	4.98	1.22	1.94	0.49	0.63
time (sec)	N/A	0.386	0.781	0.233	0.000	0.108	5.069	0.130	0.281	2.022

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	131	92	96	150	1296	546	142	139	85
N.S.	1	1.39	0.98	1.02	1.60	13.79	5.81	1.51	1.48	0.90
time (sec)	N/A	1.073	0.465	0.228	0.116	0.163	0.472	0.121	0.285	0.222

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	99	77	76	100	644	442	98	115	68
N.S.	1	1.30	1.01	1.00	1.32	8.47	5.82	1.29	1.51	0.89
time (sec)	N/A	0.793	0.316	0.208	0.137	0.125	0.402	0.126	0.261	0.167

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	81	65	67	71	264	330	75	92	59
N.S.	1	1.27	1.02	1.05	1.11	4.12	5.16	1.17	1.44	0.92
time (sec)	N/A	0.565	0.235	0.205	0.121	0.107	0.324	0.124	0.319	2.202

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	61	59	52	56	76	243	58	73	46
N.S.	1	0.97	0.94	0.83	0.89	1.21	3.86	0.92	1.16	0.73
time (sec)	N/A	0.512	0.137	0.167	0.124	0.121	0.276	0.119	0.272	0.118

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	48	50	40	40	43	141	43	43	36
N.S.	1	1.23	1.28	1.03	1.03	1.10	3.62	1.10	1.10	0.92
time (sec)	N/A	0.342	0.137	0.147	0.033	0.088	0.221	0.132	0.266	2.091

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	49	42	41	42	146	43	42	35
N.S.	1	1.00	1.26	1.08	1.05	1.08	3.74	1.10	1.08	0.90
time (sec)	N/A	0.296	0.006	0.121	0.033	0.106	0.218	0.129	0.341	0.002



Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	63	65	56	65	73	0	58	93	58
N.S.	1	1.24	1.27	1.10	1.27	1.43	0.00	1.14	1.82	1.14
time (sec)	N/A	0.413	0.116	0.424	0.043	0.103	0.000	0.123	0.276	2.404

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	77	66	71	86	271	0	75	252	73
N.S.	1	1.28	1.10	1.18	1.43	4.52	0.00	1.25	4.20	1.22
time (sec)	N/A	0.558	0.225	0.442	0.046	0.103	0.000	0.125	3.049	2.391

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	109	76	92	121	641	0	97	15	111
N.S.	1	1.43	1.00	1.21	1.59	8.43	0.00	1.28	0.20	1.46
time (sec)	N/A	0.775	0.231	0.536	0.042	0.115	0.000	0.121	200.039	2.448

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	131	92	114	173	1299	0	142	15	163
N.S.	1	1.35	0.95	1.18	1.78	13.39	0.00	1.46	0.15	1.68
time (sec)	N/A	1.072	0.357	0.575	0.049	0.117	0.000	0.120	200.032	2.663

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	58	49	73	68	182	0	174	2513	69
N.S.	1	1.05	0.89	1.33	1.24	3.31	0.00	3.16	45.69	1.25
time (sec)	N/A	0.407	0.114	8.961	0.221	0.103	0.000	0.128	0.271	2.341

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	231	245	250	953	0	1516	0	0	26	0
N.S.	1	1.06	1.08	4.13	0.00	6.56	0.00	0.00	0.11	0.00
time (sec)	N/A	1.030	1.305	6.418	0.000	0.204	0.000	0.000	0.307	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	351	353	268	1186	0	2110	0	0	28	0
N.S.	1	1.01	0.76	3.38	0.00	6.01	0.00	0.00	0.08	0.00
time (sec)	N/A	1.525	0.705	6.407	0.000	0.175	0.000	0.000	0.234	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	56	50	33	31	0	37	87	33
N.S.	1	1.00	1.75	1.56	1.03	0.97	0.00	1.16	2.72	1.03
time (sec)	N/A	0.236	0.174	0.586	0.118	0.095	0.000	0.118	0.255	2.085

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	27	64	24	23	28	0	23	62	21
N.S.	1	0.93	2.21	0.83	0.79	0.97	0.00	0.79	2.14	0.72
time (sec)	N/A	0.260	0.015	0.367	0.042	0.075	0.000	0.115	0.248	2.065

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	35	41	19	22	0	19	75	25
N.S.	1	1.00	1.52	1.78	0.83	0.96	0.00	0.83	3.26	1.09
time (sec)	N/A	0.209	0.127	0.464	0.115	0.089	0.000	0.118	0.238	2.023

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	18	15	20	48	15
N.S.	1	1.00	1.00	0.92	0.83	1.50	1.25	1.67	4.00	1.25
time (sec)	N/A	0.222	0.019	0.240	0.031	0.089	0.109	0.121	0.264	2.113

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	40	44	19	21	0	16	79	24
N.S.	1	1.00	2.00	2.20	0.95	1.05	0.00	0.80	3.95	1.20
time (sec)	N/A	0.231	0.112	0.296	0.126	0.107	0.000	0.124	0.251	2.030

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	80	32	21	38	0	44	78	29
N.S.	1	1.00	2.11	0.84	0.55	1.00	0.00	1.16	2.05	0.76
time (sec)	N/A	0.257	0.019	0.315	0.046	0.087	0.000	0.119	0.273	2.040

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	109	174	64	37	128	111	0	123	151	47
N.S.	1	1.60	0.59	0.34	1.17	1.02	0.00	1.13	1.39	0.43
time (sec)	N/A	0.472	0.156	0.451	0.132	0.129	0.000	0.119	0.229	2.062

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	158	58	33	124	99	0	119	144	44
N.S.	1	1.52	0.56	0.32	1.19	0.95	0.00	1.14	1.38	0.42
time (sec)	N/A	0.460	0.124	0.233	0.117	0.089	0.000	0.122	0.282	2.126

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	175	59	42	125	107	0	121	141	45
N.S.	1	1.65	0.56	0.40	1.18	1.01	0.00	1.14	1.33	0.42
time (sec)	N/A	0.497	0.125	0.279	0.140	0.122	0.000	0.122	0.255	2.107

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	109	169	61	45	133	116	0	123	150	48
N.S.	1	1.55	0.56	0.41	1.22	1.06	0.00	1.13	1.38	0.44
time (sec)	N/A	0.429	0.125	0.309	0.136	0.104	0.000	0.112	0.276	2.099

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	70	53	77	54	61	0	59	186	54
N.S.	1	1.25	0.95	1.38	0.96	1.09	0.00	1.05	3.32	0.96
time (sec)	N/A	0.314	0.329	0.515	0.120	0.076	0.000	0.126	0.253	2.138

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	86	42	40	58	0	39	144	39
N.S.	1	1.00	1.83	0.89	0.85	1.23	0.00	0.83	3.06	0.83
time (sec)	N/A	0.263	0.071	0.319	0.043	0.094	0.000	0.127	0.253	2.145

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	52	41	57	35	50	0	35	174	41
N.S.	1	1.30	1.02	1.42	0.88	1.25	0.00	0.88	4.35	1.02
time (sec)	N/A	0.261	0.258	0.319	0.127	0.091	0.000	0.121	0.303	2.191

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	24	13	21	28	12	19	12	28
N.S.	1	1.00	1.71	0.93	1.50	2.00	0.86	1.36	0.86	2.00
time (sec)	N/A	0.223	0.030	0.170	0.036	0.105	0.123	0.123	0.303	2.105

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	59	40	66	37	51	0	39	178	47
N.S.	1	1.34	0.91	1.50	0.84	1.16	0.00	0.89	4.05	1.07
time (sec)	N/A	0.280	0.271	0.299	0.120	0.096	0.000	0.128	0.295	2.157

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	60	96	54	35	78	0	54	172	50
N.S.	1	1.09	1.75	0.98	0.64	1.42	0.00	0.98	3.13	0.91
time (sec)	N/A	0.277	0.081	0.330	0.052	0.083	0.000	0.134	0.292	2.124

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	77	57	79	48	62	0	50	192	55
N.S.	1	1.40	1.04	1.44	0.87	1.13	0.00	0.91	3.49	1.00
time (sec)	N/A	0.291	0.272	0.465	0.131	0.100	0.000	0.119	0.250	2.120

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	72	117	63	45	87	0	82	182	59
N.S.	1	1.12	1.83	0.98	0.70	1.36	0.00	1.28	2.84	0.92
time (sec)	N/A	0.284	0.104	0.633	0.042	0.093	0.000	0.121	0.239	2.302

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	214	191	71	165	176	0	159	323	76
N.S.	1	1.47	1.31	0.49	1.13	1.21	0.00	1.09	2.21	0.52
time (sec)	N/A	0.511	0.501	0.464	0.124	0.097	0.000	0.119	0.291	2.302

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	209	174	53	144	169	0	139	317	67
N.S.	1	1.56	1.30	0.40	1.07	1.26	0.00	1.04	2.37	0.50
time (sec)	N/A	0.510	0.498	0.312	0.129	0.097	0.000	0.130	0.251	2.262

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	165	146	47	138	163	0	133	309	61
N.S.	1	1.31	1.16	0.37	1.10	1.29	0.00	1.06	2.45	0.48
time (sec)	N/A	0.360	0.388	0.247	0.154	0.086	0.000	0.126	0.266	2.286

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	213	181	64	146	161	0	143	306	68
N.S.	1	1.54	1.31	0.46	1.06	1.17	0.00	1.04	2.22	0.49
time (sec)	N/A	0.503	0.540	0.290	0.128	0.091	0.000	0.122	0.281	2.296

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	208	189	68	152	184	0	142	316	70
N.S.	1	1.51	1.37	0.49	1.10	1.33	0.00	1.03	2.29	0.51
time (sec)	N/A	0.496	0.495	0.320	0.135	0.107	0.000	0.125	0.306	2.226

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	47	0	0	0	0	0	58	0
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.00	0.00	0.97	0.00
time (sec)	N/A	0.240	0.086	0.000	0.000	0.000	0.000	0.000	0.257	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	97	79	0	0	0	0	0	565	0
N.S.	1	1.23	1.00	0.00	0.00	0.00	0.00	0.00	7.15	0.00
time (sec)	N/A	0.318	0.152	0.000	0.000	0.000	0.000	0.000	0.293	0.000



Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	125	197	111	0	0	0	0	0	0	0
N.S.	1	1.58	0.89	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.477	0.180	0.000	0.000	0.000	0.000	0.000	0.348	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	259	0	0	0	0	0	59	0
N.S.	1	1.00	3.28	0.00	0.00	0.00	0.00	0.00	0.75	0.00
time (sec)	N/A	0.276	0.434	0.000	0.000	0.000	0.000	0.000	0.259	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	126	0	0	0	0	0	77	0
N.S.	1	1.00	1.27	0.00	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	0.312	0.486	0.000	0.000	0.000	0.000	0.000	0.249	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	76	0	0	0	0	0	58	0
N.S.	1	1.00	1.49	0.00	0.00	0.00	0.00	0.00	1.14	0.00
time (sec)	N/A	0.243	0.554	0.000	0.000	0.000	0.000	0.000	0.322	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	109	121	0	0	0	0	0	58	0
N.S.	1	1.03	1.14	0.00	0.00	0.00	0.00	0.00	0.55	0.00
time (sec)	N/A	0.298	0.622	0.000	0.000	0.000	0.000	0.000	0.268	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	158	173	141	0	0	0	0	0	58	0
N.S.	1	1.09	0.89	0.00	0.00	0.00	0.00	0.00	0.37	0.00
time (sec)	N/A	0.355	0.687	0.000	0.000	0.000	0.000	0.000	0.278	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	219	205	228	0	0	0	0	0	58	0
N.S.	1	0.94	1.04	0.00	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.355	0.886	0.000	0.000	0.000	0.000	0.000	0.305	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	47	0
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	0.77	0.00
time (sec)	N/A	0.248	0.390	0.000	0.000	0.000	0.000	0.000	0.249	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	58	0
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	0.95	0.00
time (sec)	N/A	0.270	0.439	0.000	0.000	0.000	0.000	0.000	0.251	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	58	0
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	0.95	0.00
time (sec)	N/A	0.244	0.436	0.000	0.000	0.000	0.000	0.000	0.637	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	95	127	0	0	0	0	0	57	0
N.S.	1	1.61	2.15	0.00	0.00	0.00	0.00	0.00	0.97	0.00
time (sec)	N/A	0.395	5.205	0.000	0.000	0.000	0.000	0.000	0.230	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	99	136	0	0	0	0	0	57	0
N.S.	1	1.57	2.16	0.00	0.00	0.00	0.00	0.00	0.90	0.00
time (sec)	N/A	0.337	4.929	0.000	0.000	0.000	0.000	0.000	0.256	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	55	91	122	0	0	0	0	0	55	0
N.S.	1	1.65	2.22	0.00	0.00	0.00	0.00	0.00	1.00	0.00
time (sec)	N/A	0.338	4.977	0.000	0.000	0.000	0.000	0.000	0.276	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	53	88	126	0	0	0	0	0	52	0
N.S.	1	1.66	2.38	0.00	0.00	0.00	0.00	0.00	0.98	0.00
time (sec)	N/A	0.318	5.648	0.000	0.000	0.000	0.000	0.000	0.233	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	25	24	76	36	88	41	34
N.S.	1	1.00	0.96	1.00	0.96	3.04	1.44	3.52	1.64	1.36
time (sec)	N/A	0.232	0.036	0.427	0.037	0.098	1.390	0.198	0.238	2.135

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	93	126	0	0	0	0	0	60	0
N.S.	1	1.58	2.14	0.00	0.00	0.00	0.00	0.00	1.02	0.00
time (sec)	N/A	0.337	2.056	0.000	0.000	0.000	0.000	0.000	0.304	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	56	92	120	0	0	0	0	0	63	0
N.S.	1	1.64	2.14	0.00	0.00	0.00	0.00	0.00	1.12	0.00
time (sec)	N/A	0.337	2.061	0.000	0.000	0.000	0.000	0.000	0.227	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	133	183	159	0	0	0	0	0	78	0
N.S.	1	1.38	1.20	0.00	0.00	0.00	0.00	0.00	0.59	0.00
time (sec)	N/A	0.508	5.424	0.000	0.000	0.000	0.000	0.000	0.252	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	137	187	169	0	0	0	0	0	78	0
N.S.	1	1.36	1.23	0.00	0.00	0.00	0.00	0.00	0.57	0.00
time (sec)	N/A	0.485	5.174	0.000	0.000	0.000	0.000	0.000	0.266	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	131	181	155	0	0	0	0	0	76	0
N.S.	1	1.38	1.18	0.00	0.00	0.00	0.00	0.00	0.58	0.00
time (sec)	N/A	0.494	5.011	0.000	0.000	0.000	0.000	0.000	0.325	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	127	177	163	0	0	0	0	0	71	0
N.S.	1	1.39	1.28	0.00	0.00	0.00	0.00	0.00	0.56	0.00
time (sec)	N/A	0.474	5.891	0.000	0.000	0.000	0.000	0.000	0.241	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	33	51	33	36	72	70	52	36	34
N.S.	1	1.18	1.82	1.18	1.29	2.57	2.50	1.86	1.29	1.21
time (sec)	N/A	0.245	0.073	0.364	0.132	0.083	2.411	0.209	0.234	2.434

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	135	182	162	0	0	0	0	0	84	0
N.S.	1	1.35	1.20	0.00	0.00	0.00	0.00	0.00	0.62	0.00
time (sec)	N/A	0.497	2.186	0.000	0.000	0.000	0.000	0.000	0.292	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	136	183	159	0	0	0	0	0	87	0
N.S.	1	1.35	1.17	0.00	0.00	0.00	0.00	0.00	0.64	0.00
time (sec)	N/A	0.493	2.264	0.000	0.000	0.000	0.000	0.000	0.243	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	49	38	47	304	566	68	137	211	94
N.S.	1	1.14	0.88	1.09	7.07	13.16	1.58	3.19	4.91	2.19
time (sec)	N/A	0.308	0.109	0.540	0.101	0.102	0.770	0.200	0.250	2.237

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	47	62	42	494	194	65	78	46	162
N.S.	1	1.04	1.38	0.93	10.98	4.31	1.44	1.73	1.02	3.60
time (sec)	N/A	0.333	0.065	0.826	0.129	0.087	1.668	0.185	0.271	2.302

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	71	55	62	829	1568	87	171	385	227
N.S.	1	1.08	0.83	0.94	12.56	23.76	1.32	2.59	5.83	3.44
time (sec)	N/A	0.421	0.119	1.536	0.166	0.112	3.888	0.220	0.280	2.224

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	88	124	160	0	0	0	0	0	80	0
N.S.	1	1.41	1.82	0.00	0.00	0.00	0.00	0.00	0.91	0.00
time (sec)	N/A	0.432	8.852	0.000	0.000	0.000	0.000	0.000	0.235	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	169	213	317	0	0	0	0	0	158	0
N.S.	1	1.26	1.88	0.00	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	0.633	11.968	0.000	0.000	0.000	0.000	0.000	0.260	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	307	362	606	0	0	0	0	0	0	0
N.S.	1	1.18	1.97	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.919	13.103	0.000	0.000	0.000	0.000	0.000	0.477	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	387	0	0	0	0	0	98	0
N.S.	1	1.00	3.37	0.00	0.00	0.00	0.00	0.00	0.85	0.00
time (sec)	N/A	0.438	0.880	0.000	0.000	0.000	0.000	0.000	0.285	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	174	0	0	0	0	0	116	0
N.S.	1	1.00	1.29	0.00	0.00	0.00	0.00	0.00	0.86	0.00
time (sec)	N/A	0.462	0.753	0.000	0.000	0.000	0.000	0.000	0.296	0.000



Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F(-1)</b>	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	62	76	0	625	0	0	31	65
N.S.	1	1.00	0.85	1.04	0.00	8.56	0.00	0.00	0.42	0.89
time (sec)	N/A	0.352	0.203	1.450	0.000	0.117	0.000	0.000	0.298	3.301

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	71	57	74	0	334	87	0	56	51
N.S.	1	1.01	0.81	1.06	0.00	4.77	1.24	0.00	0.80	0.73
time (sec)	N/A	0.341	0.100	1.168	0.000	0.105	17.139	0.000	0.245	2.861

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	49	43	61	0	303	66	0	18	39
N.S.	1	1.02	0.90	1.27	0.00	6.31	1.38	0.00	0.38	0.81
time (sec)	N/A	0.274	0.059	1.191	0.000	0.117	0.997	0.000	0.267	2.406

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	49	47	37	0	305	66	0	31	36
N.S.	1	1.04	1.00	0.79	0.00	6.49	1.40	0.00	0.66	0.77
time (sec)	N/A	0.268	0.085	1.158	0.000	0.108	1.910	0.000	0.310	2.569

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	A	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	97	76	0	625	87	0	31	65
N.S.	1	1.00	1.37	1.07	0.00	8.80	1.23	0.00	0.44	0.92
time (sec)	N/A	0.348	0.112	1.153	0.000	0.108	8.253	0.000	0.228	2.731

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F(-1)</b>	<b>F(-1)</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	73	101	74	0	1110	0	0	31	64
N.S.	1	1.01	1.40	1.03	0.00	15.42	0.00	0.00	0.43	0.89
time (sec)	N/A	0.340	0.176	1.154	0.000	0.154	0.000	0.000	0.321	3.427

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	92	136	149	0	8891	0	0	38	0
N.S.	1	0.68	1.01	1.10	0.00	65.86	0.00	0.00	0.28	0.00
time (sec)	N/A	0.500	0.870	2.351	0.000	1.149	0.000	0.000	0.382	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	53	105	90	0	6663	0	0	38	0
N.S.	1	0.50	1.00	0.86	0.00	63.46	0.00	0.00	0.36	0.00
time (sec)	N/A	0.438	0.051	2.131	0.000	0.842	0.000	0.000	0.309	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	B	F	F(-1)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	29	58	52	0	1748	0	0	36	0
N.S.	1	0.50	1.00	0.90	0.00	30.14	0.00	0.00	0.62	0.00
time (sec)	N/A	0.320	0.024	2.151	0.000	0.560	0.000	0.000	0.349	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	B	F	F(-1)	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	111	109	0	0	6705	0	0	36	0
N.S.	1	1.05	1.03	0.00	0.00	63.25	0.00	0.00	0.34	0.00
time (sec)	N/A	0.458	0.241	0.000	0.000	0.831	0.000	0.000	0.301	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	B	F	F(-1)	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	142	191	142	0	0	9168	0	0	23	0
N.S.	1	1.35	1.00	0.00	0.00	64.56	0.00	0.00	0.16	0.00
time (sec)	N/A	0.580	0.397	0.000	0.000	1.103	0.000	0.000	200.026	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	B	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	85	131	165	0	7896	0	0	502	0
N.S.	1	0.64	0.99	1.25	0.00	59.82	0.00	0.00	3.80	0.00
time (sec)	N/A	0.779	0.211	2.221	0.000	1.353	0.000	0.000	0.422	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	95	76	92	94	604	0	68	159	155
N.S.	1	0.89	0.71	0.86	0.88	5.64	0.00	0.64	1.49	1.45
time (sec)	N/A	0.412	0.098	0.796	0.120	0.104	0.000	0.272	0.245	0.113

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	68	60	80	69	339	0	52	112	93
N.S.	1	0.88	0.78	1.04	0.90	4.40	0.00	0.68	1.45	1.21
time (sec)	N/A	0.362	0.065	0.708	0.127	0.088	0.000	0.116	0.240	2.235

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	45	40	48	47	147	0	41	68	58
N.S.	1	0.88	0.78	0.94	0.92	2.88	0.00	0.80	1.33	1.14
time (sec)	N/A	0.339	0.066	0.390	0.119	0.115	0.000	0.114	0.216	0.082

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	22	22	27	23	38	0	23	22	34
N.S.	1	0.88	0.88	1.08	0.92	1.52	0.00	0.92	0.88	1.36
time (sec)	N/A	0.297	0.013	0.378	0.120	0.086	0.000	0.110	0.238	0.052

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	22	22	27	38	49	0	32	34	38
N.S.	1	0.88	0.88	1.08	1.52	1.96	0.00	1.28	1.36	1.52
time (sec)	N/A	0.296	0.015	0.365	0.039	0.099	0.000	0.113	0.268	0.086

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	47	179	48	62	198	0	56	103	62
N.S.	1	0.89	3.38	0.91	1.17	3.74	0.00	1.06	1.94	1.17
time (sec)	N/A	0.351	1.440	0.388	0.039	0.090	0.000	0.125	0.246	2.092

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	72	286	77	88	459	0	72	177	97
N.S.	1	0.89	3.53	0.95	1.09	5.67	0.00	0.89	2.19	1.20
time (sec)	N/A	0.263	1.792	0.691	0.035	0.097	0.000	0.143	0.271	0.087

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	101	115	88	110	796	0	83	245	160
N.S.	1	0.89	1.02	0.78	0.97	7.04	0.00	0.73	2.17	1.42
time (sec)	N/A	0.258	10.083	0.752	0.038	0.108	0.000	0.127	0.228	0.078

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	113	48	35	89	393	0	89	193	86
N.S.	1	1.28	0.55	0.40	1.01	4.47	0.00	1.01	2.19	0.98
time (sec)	N/A	0.283	0.041	0.603	0.111	0.113	0.000	0.111	0.267	0.248

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	108	95	24	78	94	0	78	79	81
N.S.	1	1.59	1.40	0.35	1.15	1.38	0.00	1.15	1.16	1.19
time (sec)	N/A	0.330	0.026	0.485	0.128	0.097	0.000	0.105	0.244	2.222

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	23	16	36	22	31	0	23	26	26
N.S.	1	1.44	1.00	2.25	1.38	1.94	0.00	1.44	1.62	1.62
time (sec)	N/A	0.192	0.011	0.503	0.121	0.088	0.000	0.107	0.225	0.164

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	113	48	34	230	0	35	81	38
N.S.	1	1.00	3.23	1.37	0.97	6.57	0.00	1.00	2.31	1.09
time (sec)	N/A	0.209	1.125	0.591	0.113	0.098	0.000	0.104	0.277	0.198

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	76	62	61	1097	0	50	186	110
N.S.	1	1.00	1.03	0.84	0.82	14.82	0.00	0.68	2.51	1.49
time (sec)	N/A	0.249	11.059	0.678	0.107	0.108	0.000	0.105	0.251	2.571

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	113	97	59	81	549	0	81	182	86
N.S.	1	1.28	1.10	0.67	0.92	6.24	0.00	0.92	2.07	0.98
time (sec)	N/A	0.409	0.068	0.633	0.114	0.092	0.000	0.107	0.258	0.329

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	104	24	47	69	94	0	69	71	70
N.S.	1	1.49	0.34	0.67	0.99	1.34	0.00	0.99	1.01	1.00
time (sec)	N/A	0.329	0.012	0.546	0.122	0.100	0.000	0.108	0.230	2.347

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	97	22	138	75	113	0	76	82	81
N.S.	1	1.37	0.31	1.94	1.06	1.59	0.00	1.07	1.15	1.14
time (sec)	N/A	0.298	0.015	0.539	0.113	0.095	0.000	0.105	0.260	0.273

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	108	113	150	87	628	0	88	208	93
N.S.	1	1.15	1.20	1.60	0.93	6.68	0.00	0.94	2.21	0.99
time (sec)	N/A	0.498	1.451	0.556	0.122	0.113	0.000	0.108	0.236	2.467

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	C	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	296	382	51	36	0	1303	0	263	513	474
N.S.	1	1.29	0.17	0.12	0.00	4.40	0.00	0.89	1.73	1.60
time (sec)	N/A	1.120	0.047	0.689	0.000	0.143	0.000	0.122	0.285	5.215

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	C	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	373	24	24	0	171	0	251	230	457
N.S.	1	1.33	0.09	0.09	0.00	0.61	0.00	0.90	0.82	1.63
time (sec)	N/A	1.115	0.011	0.540	0.000	0.099	0.000	0.117	0.287	4.844

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	134	22	56	97	120	0	98	101	104
N.S.	1	1.47	0.24	0.62	1.07	1.32	0.00	1.08	1.11	1.14
time (sec)	N/A	0.381	0.014	0.560	0.117	0.120	0.000	0.106	0.227	2.464



Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	109	134	113	68	109	943	0	110	252	122
N.S.	1	1.23	1.04	0.62	1.00	8.65	0.00	1.01	2.31	1.12
time (sec)	N/A	0.295	1.231	0.618	0.118	0.119	0.000	0.113	0.258	2.517

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F(-2)</b>	C	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	109	81	70	0	406	0	328	230	163
N.S.	1	1.02	0.76	0.65	0.00	3.79	0.00	3.07	2.15	1.52
time (sec)	N/A	0.305	0.056	0.944	0.000	0.103	0.000	0.107	0.274	2.437

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	C	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	107	215	0	1604	0	456	5067	280
N.S.	1	1.00	0.69	1.39	0.00	10.35	0.00	2.94	32.69	1.81
time (sec)	N/A	0.386	0.127	1.077	0.000	0.119	0.000	0.113	0.267	24.354

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	205	0	0	0	0	0	0	0
N.S.	1	1.00	1.23	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.464	1.660	0.000	0.000	0.000	0.000	0.000	0.265	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	169	0	0	0	0	0	439	0
N.S.	1	1.00	1.44	0.00	0.00	0.00	0.00	0.00	3.75	0.00
time (sec)	N/A	0.372	0.830	0.000	0.000	0.000	0.000	0.000	0.263	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	141	0	0	0	0	0	46	0
N.S.	1	1.00	2.10	0.00	0.00	0.00	0.00	0.00	0.69	0.00
time (sec)	N/A	0.286	0.176	0.000	0.000	0.000	0.000	0.000	0.244	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	134	0	0	0	0	0	46	0
N.S.	1	1.00	2.06	0.00	0.00	0.00	0.00	0.00	0.71	0.00
time (sec)	N/A	0.290	0.777	0.000	0.000	0.000	0.000	0.000	0.224	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	164	0	0	0	0	0	440	0
N.S.	1	1.00	1.45	0.00	0.00	0.00	0.00	0.00	3.89	0.00
time (sec)	N/A	0.571	0.606	0.000	0.000	0.000	0.000	0.000	0.249	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	161	161	210	0	0	0	0	0	0	0
N.S.	1	1.00	1.30	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.686	1.122	0.000	0.000	0.000	0.000	0.000	0.316	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	311	161	133	195	145	1226	0	193	248	0
N.S.	1	0.52	0.43	0.63	0.47	3.94	0.00	0.62	0.80	0.00
time (sec)	N/A	1.525	0.175	1.982	0.138	0.104	0.000	0.134	0.227	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-1)	A	B	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	193	103	104	131	90	458	0	137	139	0
N.S.	1	0.53	0.54	0.68	0.47	2.37	0.00	0.71	0.72	0.00
time (sec)	N/A	0.486	0.124	1.613	0.141	0.107	0.000	0.134	0.257	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	53	51	218	35	53	0	63	31	0
N.S.	1	0.64	0.61	2.63	0.42	0.64	0.00	0.76	0.37	0.00
time (sec)	N/A	0.356	0.039	1.884	0.129	0.099	0.000	0.118	0.252	0.000

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	83	53	51	48	56	70	0	99	58	0
N.S.	1	0.64	0.61	0.58	0.67	0.84	0.00	1.19	0.70	0.00
time (sec)	N/A	0.390	0.088	1.195	0.139	0.092	0.000	0.128	0.235	0.000

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	197	107	334	131	112	613	0	176	303	0
N.S.	1	0.54	1.70	0.66	0.57	3.11	0.00	0.89	1.54	0.00
time (sec)	N/A	0.888	7.135	1.675	0.159	0.109	0.000	0.134	0.261	0.000

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F(-1)</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	319	169	164	195	167	1617	0	230	830	0
N.S.	1	0.53	0.51	0.61	0.52	5.07	0.00	0.72	2.60	0.00
time (sec)	N/A	1.764	11.478	1.831	0.137	0.112	0.000	0.143	0.282	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	137	124	118	0	697	0	0	11	0
N.S.	1	0.87	0.79	0.75	0.00	4.44	0.00	0.00	0.07	0.00
time (sec)	N/A	1.004	0.382	1.779	0.000	0.126	0.000	0.000	0.243	0.000

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	101	88	88	0	230	0	0	11	0
N.S.	1	0.88	0.77	0.77	0.00	2.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.646	0.327	0.860	0.000	0.115	0.000	0.000	0.233	0.000

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	69	59	58	0	215	0	0	9	0
N.S.	1	0.90	0.77	0.75	0.00	2.79	0.00	0.00	0.12	0.00
time (sec)	N/A	0.353	0.132	0.555	0.000	0.106	0.000	0.000	0.230	0.000

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	9	0	9	9	8	9	9	11
N.S.	1	1.00	1.29	0.00	1.29	1.29	1.14	1.29	1.29	1.57
time (sec)	N/A	0.324	1.907	0.000	0.470	0.100	10.144	0.177	0.273	3.199

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	137	124	118	0	698	0	0	11	0
N.S.	1	0.87	0.79	0.75	0.00	4.45	0.00	0.00	0.07	0.00
time (sec)	N/A	0.608	0.459	1.716	0.000	0.141	0.000	0.000	0.271	0.000

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	101	88	88	0	230	0	0	11	0
N.S.	1	0.88	0.77	0.77	0.00	2.00	0.00	0.00	0.10	0.00
time (sec)	N/A	0.682	0.368	0.839	0.000	0.093	0.000	0.000	0.264	0.000

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	69	62	58	0	216	0	0	9	0
N.S.	1	0.90	0.81	0.75	0.00	2.81	0.00	0.00	0.12	0.00
time (sec)	N/A	0.588	0.130	0.566	0.000	0.102	0.000	0.000	0.290	0.000

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	9	0	9	9	8	9	9	11
N.S.	1	1.00	1.29	0.00	1.29	1.29	1.14	1.29	1.29	1.57
time (sec)	N/A	0.532	3.648	0.000	0.215	0.096	2.013	0.191	0.244	2.852

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [116] had the largest ratio of [1.9090899999999995]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	10	10	1.00	8	1.250
2	C	11	11	1.26	8	1.375
3	A	7	7	1.00	8	0.875
4	C	7	7	1.33	8	0.875
5	A	4	4	1.00	8	0.500
6	A	3	3	1.00	6	0.500
7	C	3	3	1.36	6	0.500
8	A	4	4	1.00	8	0.500
9	C	7	7	1.48	8	0.875
10	A	7	7	1.00	8	0.875
11	C	11	11	1.36	8	1.375
12	A	10	10	1.00	8	1.250
13	A	12	11	1.00	12	0.917
14	A	10	9	0.95	12	0.750
15	A	10	9	0.96	12	0.750
16	A	8	7	0.88	12	0.583
17	A	8	7	0.89	12	0.583
18	A	10	9	0.95	12	0.750
19	A	10	9	0.96	12	0.750
20	A	12	11	1.01	12	0.917
21	A	12	11	1.01	8	1.375

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	1.00	8	0.500
23	A	5	4	1.00	10	0.400
24	C	9	9	1.00	10	0.900
25	A	5	5	1.00	10	0.500
26	A	5	5	1.00	10	0.500
27	C	13	13	0.83	14	0.929
28	C	9	9	0.93	14	0.643
29	A	5	5	1.00	14	0.357
30	C	5	5	1.13	14	0.357
31	C	9	9	1.00	14	0.643
32	C	13	13	0.88	14	0.929
33	A	12	11	0.86	8	1.375
34	A	14	13	0.74	10	1.300
35	A	12	11	0.81	10	1.100
36	A	12	11	0.80	10	1.100
37	A	12	12	0.57	10	1.200
38	A	6	6	0.71	10	0.600
39	A	6	6	0.71	10	0.600
40	A	7	6	1.00	12	0.500
41	A	12	12	1.08	12	1.000
42	A	10	10	1.09	12	0.833
43	A	8	8	1.07	12	0.667
44	A	6	6	1.00	12	0.500
45	A	3	3	1.00	12	0.250
46	A	5	5	1.08	12	0.417
47	A	7	7	1.12	12	0.583
48	A	9	9	1.14	12	0.750
49	A	11	11	1.12	12	0.917
50	A	10	9	1.11	8	1.125
51	A	8	7	1.07	8	0.875
52	A	6	5	1.00	8	0.625
53	A	4	3	1.00	8	0.375

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	6	5	1.00	8	0.625
55	A	8	7	1.00	8	0.875
56	A	10	9	1.08	8	1.125
57	A	12	12	1.00	12	1.000
58	A	10	10	1.00	12	0.833
59	A	8	8	1.00	12	0.667
60	A	6	6	1.00	12	0.500
61	A	5	5	1.00	12	0.417
62	A	7	7	1.15	12	0.583
63	A	9	9	1.20	12	0.750
64	A	11	11	1.23	12	0.917
65	A	5	5	1.00	12	0.417
66	A	5	5	1.00	12	0.417
67	A	8	7	1.22	14	0.500
68	A	7	6	1.15	14	0.429
69	A	7	6	1.03	11	0.545
70	C	12	12	1.40	11	1.091
71	A	7	6	1.05	11	0.545
72	C	11	11	1.47	9	1.222
73	C	12	12	2.00	9	1.333
74	A	6	5	1.00	11	0.455
75	C	11	11	1.67	11	1.000
76	A	7	6	1.00	11	0.545
77	C	12	12	1.35	11	1.091
78	A	7	6	1.00	11	0.545
79	C	11	11	1.32	11	1.000
80	A	10	9	1.65	13	0.692
81	C	24	23	1.12	13	1.769
82	A	8	7	1.69	13	0.538
83	C	15	14	1.22	11	1.273
84	C	8	8	1.15	11	0.727
85	A	6	5	1.17	13	0.385

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	C	8	8	1.20	13	0.615
87	A	6	5	1.12	13	0.385
88	C	8	8	1.37	13	0.615
89	A	6	5	1.07	13	0.385
90	A	9	9	1.00	11	0.818
91	A	4	3	1.03	23	0.130
92	A	5	4	1.03	11	0.364
93	C	6	5	1.31	11	0.455
94	A	5	4	1.05	11	0.364
95	A	4	4	1.00	9	0.444
96	A	2	2	1.00	9	0.222
97	A	4	3	1.00	11	0.273
98	A	4	4	1.00	11	0.364
99	A	4	3	1.00	11	0.273
100	A	6	6	1.00	11	0.545
101	A	5	4	1.00	11	0.364
102	A	8	8	1.15	11	0.727
103	A	6	5	0.99	13	0.385
104	A	6	5	0.99	13	0.385
105	A	6	5	0.98	13	0.385
106	A	4	3	1.00	13	0.231
107	A	5	5	1.00	8	0.625
108	A	6	5	1.18	13	0.385
109	A	6	5	1.16	13	0.385
110	A	21	20	1.05	13	1.538
111	A	15	14	1.01	13	1.077
112	A	9	8	1.00	13	0.615
113	A	4	3	1.00	11	0.273
114	A	9	8	0.88	11	0.727
115	C	15	14	0.95	13	1.077
116	C	21	21	1.63	11	1.909
117	C	15	15	1.32	11	1.364

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	C	11	11	1.42	11	1.000
119	A	4	4	1.26	11	0.364
120	C	4	4	1.50	9	0.444
121	A	3	3	1.00	6	0.500
122	C	9	9	1.68	9	1.000
123	C	13	13	1.34	11	1.182
124	C	18	18	1.62	11	1.636
125	C	20	20	1.56	11	1.818
126	C	9	8	1.24	11	0.727
127	C	7	6	1.28	11	0.545
128	C	7	6	1.37	11	0.545
129	C	9	8	1.16	11	0.727
130	A	10	9	1.00	13	0.692
131	A	8	7	1.00	13	0.538
132	A	10	9	1.00	13	0.692
133	A	9	8	1.00	13	0.615
134	C	21	20	1.39	13	1.538
135	C	16	15	1.30	13	1.154
136	C	13	12	1.27	13	0.923
137	A	11	11	0.97	13	0.846
138	C	6	6	1.23	11	0.545
139	A	5	5	1.00	8	0.625
140	C	8	8	1.24	11	0.727
141	C	12	12	1.28	13	0.923
142	C	16	16	1.43	13	1.231
143	C	19	19	1.35	13	1.462
144	A	6	6	1.05	14	0.429
145	A	9	8	1.06	24	0.333
146	A	10	9	1.01	26	0.346
147	A	6	5	1.00	11	0.455
148	A	6	5	0.93	11	0.455
149	A	5	4	1.00	9	0.444
Continued on next page						

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	5	4	1.00	11	0.364
151	A	5	4	1.00	11	0.364
152	A	6	5	1.00	11	0.455
153	A	11	10	1.60	11	0.909
154	A	11	10	1.52	7	1.429
155	A	11	10	1.65	11	0.909
156	A	11	10	1.55	11	0.909
157	A	8	7	1.25	13	0.538
158	A	5	4	1.00	13	0.308
159	A	7	6	1.30	11	0.545
160	A	6	5	1.00	13	0.385
161	A	7	6	1.34	13	0.462
162	A	5	4	1.09	13	0.308
163	A	8	7	1.40	13	0.538
164	A	5	4	1.12	13	0.308
165	A	14	13	1.47	13	1.000
166	A	13	12	1.56	13	0.923
167	A	3	3	1.31	9	0.333
168	A	13	12	1.54	13	0.923
169	A	13	12	1.51	13	0.923
170	A	3	3	1.00	13	0.231
171	A	5	5	1.23	15	0.333
172	A	7	7	1.58	15	0.467
173	A	3	3	1.00	9	0.333
174	A	3	3	1.00	15	0.200
175	A	2	2	1.00	11	0.182
176	A	5	4	1.03	11	0.364
177	A	6	5	1.09	11	0.455
178	A	7	6	0.94	11	0.545
179	A	3	3	1.00	7	0.429
180	A	3	3	1.00	9	0.333
181	A	3	3	1.00	9	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
182	A	5	4	1.61	17	0.235
183	A	5	4	1.57	17	0.235
184	A	5	4	1.65	15	0.267
185	A	5	4	1.66	13	0.308
186	A	5	4	1.00	17	0.235
187	A	5	4	1.58	17	0.235
188	A	5	4	1.64	17	0.235
189	A	7	6	1.38	19	0.316
190	A	7	6	1.36	19	0.316
191	A	7	6	1.38	17	0.353
192	A	7	6	1.39	15	0.400
193	A	6	5	1.18	19	0.263
194	A	7	6	1.35	19	0.316
195	A	7	6	1.35	19	0.316
196	C	9	8	1.14	17	0.471
197	A	9	8	1.04	17	0.471
198	C	13	12	1.08	17	0.706
199	A	5	4	1.41	19	0.211
200	A	7	6	1.26	21	0.286
201	A	9	8	1.18	21	0.381
202	A	5	4	1.00	15	0.267
203	A	5	4	1.00	21	0.190
204	A	11	10	1.00	19	0.526
205	A	11	10	1.01	19	0.526
206	A	9	8	1.02	19	0.421
207	A	9	8	1.04	19	0.421
208	A	11	10	1.00	19	0.526
209	A	11	10	1.01	19	0.526
210	C	12	11	0.68	23	0.478
211	C	10	9	0.50	23	0.391
212	C	7	6	0.50	21	0.286
213	C	7	6	1.05	21	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
214	C	7	6	1.35	23	0.261
215	C	12	11	0.64	21	0.524
216	A	4	3	0.89	16	0.188
217	A	5	4	0.88	16	0.250
218	A	4	3	0.88	16	0.188
219	A	5	4	0.88	14	0.286
220	A	5	4	0.88	14	0.286
221	A	4	3	0.89	16	0.188
222	A	5	4	0.89	16	0.250
223	A	4	3	0.89	16	0.188
224	A	4	3	1.28	10	0.300
225	A	12	11	1.59	8	1.375
226	A	7	6	1.44	8	0.750
227	A	4	3	1.00	10	0.300
228	A	4	3	1.00	10	0.300
229	A	4	3	1.28	10	0.300
230	A	12	11	1.49	8	1.375
231	A	12	11	1.37	8	1.375
232	A	4	3	1.15	10	0.300
233	A	4	3	1.29	10	0.300
234	A	11	10	1.33	8	1.250
235	A	16	15	1.47	8	1.875
236	A	4	3	1.23	10	0.300
237	A	6	5	1.02	14	0.357
238	A	4	3	1.00	14	0.214
239	A	2	2	1.00	18	0.111
240	A	2	2	1.00	18	0.111
241	A	2	2	1.00	16	0.125
242	A	2	2	1.00	16	0.125
243	A	2	2	1.00	18	0.111
244	A	2	2	1.00	18	0.111
245	A	6	5	0.52	25	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
246	A	6	5	0.53	25	0.200
247	A	6	5	0.64	25	0.200
248	A	6	5	0.64	25	0.200
249	A	6	5	0.54	25	0.200
250	A	6	5	0.53	25	0.200
251	A	4	3	0.87	9	0.333
252	A	4	3	0.88	9	0.333
253	A	4	3	0.90	7	0.429
254	N/A	4	0	1.00	7	0.000
255	A	4	3	0.87	9	0.333
256	A	4	3	0.88	9	0.333
257	A	4	3	0.90	7	0.429
258	N/A	4	0	1.00	7	0.000

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int \tanh^6(a + bx) dx$ . . . . .	119
3.2	$\int \tanh^5(a + bx) dx$ . . . . .	126
3.3	$\int \tanh^4(a + bx) dx$ . . . . .	133
3.4	$\int \tanh^3(a + bx) dx$ . . . . .	139
3.5	$\int \tanh^2(a + bx) dx$ . . . . .	145
3.6	$\int \tanh(a + bx) dx$ . . . . .	150
3.7	$\int \coth(a + bx) dx$ . . . . .	155
3.8	$\int \coth^2(a + bx) dx$ . . . . .	160
3.9	$\int \coth^3(a + bx) dx$ . . . . .	166
3.10	$\int \coth^4(a + bx) dx$ . . . . .	173
3.11	$\int \coth^5(a + bx) dx$ . . . . .	179
3.12	$\int \coth^6(a + bx) dx$ . . . . .	187
3.13	$\int (b \tanh(c + dx))^{7/2} dx$ . . . . .	194
3.14	$\int (b \tanh(c + dx))^{5/2} dx$ . . . . .	202
3.15	$\int (b \tanh(c + dx))^{3/2} dx$ . . . . .	209
3.16	$\int \sqrt{b \tanh(c + dx)} dx$ . . . . .	216
3.17	$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$ . . . . .	223
3.18	$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx$ . . . . .	230
3.19	$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx$ . . . . .	238
3.20	$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx$ . . . . .	245
3.21	$\int \sqrt[3]{\tanh(8x)} dx$ . . . . .	253
3.22	$\int \tanh^n(a + bx) dx$ . . . . .	261
3.23	$\int (b \tanh(c + dx))^n dx$ . . . . .	266
3.24	$\int (a \tanh^2(x))^{3/2} dx$ . . . . .	271
3.25	$\int \sqrt{a \tanh^2(x)} dx$ . . . . .	278



3.26	$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx$	283
3.27	$\int (-\tanh^2(c+dx))^{5/2} dx$	289
3.28	$\int (-\tanh^2(c+dx))^{3/2} dx$	296
3.29	$\int \sqrt{-\tanh^2(c+dx)} dx$	303
3.30	$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx$	309
3.31	$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx$	315
3.32	$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx$	322
3.33	$\int \sqrt{\tanh^3(x)} dx$	330
3.34	$\int (a \tanh^3(x))^{3/2} dx$	338
3.35	$\int \sqrt{a \tanh^3(x)} dx$	346
3.36	$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$	354
3.37	$\int (a \tanh^4(x))^{3/2} dx$	362
3.38	$\int \sqrt{a \tanh^4(x)} dx$	369
3.39	$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$	375
3.40	$\int (b \tanh^m(c+dx))^n dx$	381
3.41	$\int (a + a \tanh(c+dx))^5 dx$	387
3.42	$\int (a + a \tanh(c+dx))^4 dx$	396
3.43	$\int (a + a \tanh(c+dx))^3 dx$	403
3.44	$\int (a + a \tanh(c+dx))^2 dx$	410
3.45	$\int \frac{1}{a+a \tanh(c+dx)} dx$	416
3.46	$\int \frac{1}{(a+a \tanh(c+dx))^2} dx$	421
3.47	$\int \frac{1}{(a+a \tanh(c+dx))^3} dx$	427
3.48	$\int \frac{1}{(a+a \tanh(c+dx))^4} dx$	433
3.49	$\int \frac{1}{(a+a \tanh(c+dx))^5} dx$	440
3.50	$\int (1 + \tanh(x))^{7/2} dx$	449
3.51	$\int (1 + \tanh(x))^{5/2} dx$	456
3.52	$\int (1 + \tanh(x))^{3/2} dx$	462
3.53	$\int \sqrt{1 + \tanh(x)} dx$	468
3.54	$\int \frac{1}{\sqrt{1+\tanh(x)}} dx$	473
3.55	$\int \frac{1}{(1+\tanh(x))^{3/2}} dx$	479
3.56	$\int \frac{1}{(1+\tanh(x))^{5/2}} dx$	485
3.57	$\int (a + b \tanh(c+dx))^5 dx$	492
3.58	$\int (a + b \tanh(c+dx))^4 dx$	502

3.59	$\int (a + b \tanh(c + dx))^3 dx$	511
3.60	$\int (a + b \tanh(c + dx))^2 dx$	518
3.61	$\int \frac{1}{a+b \tanh(c+dx)} dx$	524
3.62	$\int \frac{1}{(a+b \tanh(c+dx))^2} dx$	530
3.63	$\int \frac{1}{(a+b \tanh(c+dx))^3} dx$	538
3.64	$\int \frac{1}{(a+b \tanh(c+dx))^4} dx$	548
3.65	$\int \frac{1}{4+6 \tanh(c+dx)} dx$	558
3.66	$\int \frac{1}{4-6 \tanh(c+dx)} dx$	563
3.67	$\int \sqrt{a + b \tanh(c + dx)} dx$	568
3.68	$\int \frac{1}{\sqrt{a+b \tanh(c+dx)}} dx$	575
3.69	$\int \frac{\sinh^4(x)}{1+\tanh(x)} dx$	582
3.70	$\int \frac{\sinh^3(x)}{1+\tanh(x)} dx$	588
3.71	$\int \frac{\sinh^2(x)}{1+\tanh(x)} dx$	595
3.72	$\int \frac{\sinh(x)}{1+\tanh(x)} dx$	601
3.73	$\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx$	607
3.74	$\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx$	613
3.75	$\int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx$	619
3.76	$\int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx$	626
3.77	$\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx$	632
3.78	$\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx$	640
3.79	$\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx$	646
3.80	$\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx$	654
3.81	$\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx$	663
3.82	$\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx$	674
3.83	$\int \frac{\sinh(x)}{a+b \tanh(x)} dx$	681
3.84	$\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx$	689
3.85	$\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx$	695
3.86	$\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx$	701
3.87	$\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx$	709
3.88	$\int \frac{\operatorname{csch}^5(x)}{a+b \tanh(x)} dx$	716
3.89	$\int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx$	725

3.90	$\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx$	733
3.91	$\int (d\operatorname{sech}(e+fx))^m (a+b\tanh(e+fx))^n dx$	739
3.92	$\int \frac{\cosh^4(x)}{1+\tanh(x)} dx$	744
3.93	$\int \frac{\cosh^3(x)}{1+\tanh(x)} dx$	750
3.94	$\int \frac{\cosh^2(x)}{1+\tanh(x)} dx$	756
3.95	$\int \frac{\cosh(x)}{1+\tanh(x)} dx$	761
3.96	$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx$	766
3.97	$\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx$	771
3.98	$\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx$	776
3.99	$\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx$	782
3.100	$\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx$	787
3.101	$\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx$	793
3.102	$\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx$	799
3.103	$\int \frac{\operatorname{sech}^8(x)}{a+b\tanh(x)} dx$	806
3.104	$\int \frac{\operatorname{sech}^6(x)}{a+b\tanh(x)} dx$	814
3.105	$\int \frac{\operatorname{sech}^4(x)}{a+b\tanh(x)} dx$	822
3.106	$\int \frac{\operatorname{sech}^2(x)}{a+b\tanh(x)} dx$	829
3.107	$\int \frac{1}{a+b\tanh(x)} dx$	834
3.108	$\int \frac{\cosh^2(x)}{a+b\tanh(x)} dx$	840
3.109	$\int \frac{\cosh^4(x)}{a+b\tanh(x)} dx$	847
3.110	$\int \frac{\operatorname{sech}^7(x)}{a+b\tanh(x)} dx$	855
3.111	$\int \frac{\operatorname{sech}^5(x)}{a+b\tanh(x)} dx$	867
3.112	$\int \frac{\operatorname{sech}^3(x)}{a+b\tanh(x)} dx$	877
3.113	$\int \frac{\operatorname{sech}(x)}{a+b\tanh(x)} dx$	884
3.114	$\int \frac{\cosh(x)}{a+b\tanh(x)} dx$	889
3.115	$\int \frac{\cosh^3(x)}{a+b\tanh(x)} dx$	896
3.116	$\int \frac{\tanh^5(x)}{1+\tanh(x)} dx$	905
3.117	$\int \frac{\tanh^4(x)}{1+\tanh(x)} dx$	914
3.118	$\int \frac{\tanh^3(x)}{1+\tanh(x)} dx$	922
3.119	$\int \frac{\tanh^2(x)}{1+\tanh(x)} dx$	929

3.120	$\int \frac{\tanh(x)}{1+\tanh(x)} dx$	935
3.121	$\int \frac{1}{1+\tanh(x)} dx$	940
3.122	$\int \frac{\coth(x)}{1+\tanh(x)} dx$	945
3.123	$\int \frac{\coth^2(x)}{1+\tanh(x)} dx$	951
3.124	$\int \frac{\coth^3(x)}{1+\tanh(x)} dx$	958
3.125	$\int \frac{\coth^4(x)}{1+\tanh(x)} dx$	966
3.126	$\int \tanh(x)(1 + \tanh(x))^{3/2} dx$	975
3.127	$\int \tanh(x)\sqrt{1 + \tanh(x)} dx$	982
3.128	$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx$	988
3.129	$\int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx$	995
3.130	$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx$	1002
3.131	$\int \tanh^2(x)\sqrt{1 + \tanh(x)} dx$	1009
3.132	$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx$	1015
3.133	$\int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx$	1021
3.134	$\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx$	1028
3.135	$\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx$	1042
3.136	$\int \frac{\tanh^3(x)}{a+b \tanh(x)} dx$	1052
3.137	$\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx$	1061
3.138	$\int \frac{\tanh(x)}{a+b \tanh(x)} dx$	1068
3.139	$\int \frac{1}{a+b \tanh(x)} dx$	1074
3.140	$\int \frac{\coth(x)}{a+b \tanh(x)} dx$	1080
3.141	$\int \frac{\coth^2(x)}{a+b \tanh(x)} dx$	1087
3.142	$\int \frac{\coth^3(x)}{a+b \tanh(x)} dx$	1095
3.143	$\int \frac{\coth^4(x)}{a+b \tanh(x)} dx$	1105
3.144	$\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx$	1116
3.145	$\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$	1123
3.146	$\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$	1132
3.147	$\int x^5 \tanh(a + 2 \log(x)) dx$	1142
3.148	$\int x^3 \tanh(a + 2 \log(x)) dx$	1148
3.149	$\int x \tanh(a + 2 \log(x)) dx$	1153
3.150	$\int \frac{\tanh(a+2 \log(x))}{x} dx$	1158
3.151	$\int \frac{\tanh(a+2 \log(x))}{x^3} dx$	1163
3.152	$\int \frac{\tanh(a+2 \log(x))}{x^5} dx$	1168

3.153	$\int x^2 \tanh(a + 2 \log(x)) dx$	1174
3.154	$\int \tanh(a + 2 \log(x)) dx$	1182
3.155	$\int \frac{\tanh(a+2 \log(x))}{x^2} dx$	1190
3.156	$\int \frac{\tanh(a+2 \log(x))}{x^4} dx$	1199
3.157	$\int x^5 \tanh^2(a + 2 \log(x)) dx$	1207
3.158	$\int x^3 \tanh^2(a + 2 \log(x)) dx$	1214
3.159	$\int x \tanh^2(a + 2 \log(x)) dx$	1220
3.160	$\int \frac{\tanh^2(a+2 \log(x))}{x} dx$	1226
3.161	$\int \frac{\tanh^2(a+2 \log(x))}{x^3} dx$	1231
3.162	$\int \frac{\tanh^2(a+2 \log(x))}{x^5} dx$	1237
3.163	$\int \frac{\tanh^2(a+2 \log(x))}{x^7} dx$	1243
3.164	$\int \frac{\tanh^2(a+2 \log(x))}{x^9} dx$	1250
3.165	$\int x^4 \tanh^2(a + 2 \log(x)) dx$	1256
3.166	$\int x^2 \tanh^2(a + 2 \log(x)) dx$	1266
3.167	$\int \tanh^2(a + 2 \log(x)) dx$	1275
3.168	$\int \frac{\tanh^2(a+2 \log(x))}{x^2} dx$	1282
3.169	$\int \frac{\tanh^2(a+2 \log(x))}{x^4} dx$	1291
3.170	$\int (ex)^m \tanh(a + 2 \log(x)) dx$	1300
3.171	$\int (ex)^m \tanh^2(a + 2 \log(x)) dx$	1305
3.172	$\int (ex)^m \tanh^3(a + 2 \log(x)) dx$	1311
3.173	$\int \tanh^p(a + b \log(x)) dx$	1318
3.174	$\int (ex)^m \tanh^p(a + b \log(x)) dx$	1323
3.175	$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx$	1328
3.176	$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx$	1333
3.177	$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$	1338
3.178	$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx$	1344
3.179	$\int \tanh^p(a + \log(x)) dx$	1351
3.180	$\int \tanh^p(a + 2 \log(x)) dx$	1356
3.181	$\int \tanh^p(a + 3 \log(x)) dx$	1361
3.182	$\int x^3 \tanh(d(a + b \log(cx^n))) dx$	1366
3.183	$\int x^2 \tanh(d(a + b \log(cx^n))) dx$	1371
3.184	$\int x \tanh(d(a + b \log(cx^n))) dx$	1376
3.185	$\int \tanh(d(a + b \log(cx^n))) dx$	1381
3.186	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx$	1386
3.187	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$	1392
3.188	$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^3} dx$	1397

3.189	$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$	1402
3.190	$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx$	1409
3.191	$\int x \tanh^2(d(a + b \log(cx^n))) dx$	1416
3.192	$\int \tanh^2(d(a + b \log(cx^n))) dx$	1423
3.193	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x} dx$	1430
3.194	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^2} dx$	1436
3.195	$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^3} dx$	1443
3.196	$\int \frac{\tanh^3(a+b \log(cx^n))}{x} dx$	1450
3.197	$\int \frac{\tanh^4(a+b \log(cx^n))}{x} dx$	1458
3.198	$\int \frac{\tanh^5(a+b \log(cx^n))}{x} dx$	1466
3.199	$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$	1475
3.200	$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx$	1480
3.201	$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx$	1487
3.202	$\int \tanh^p(d(a + b \log(cx^n))) dx$	1496
3.203	$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx$	1502
3.204	$\int \frac{\tanh^{\frac{5}{2}}(a+b \log(cx^n))}{x} dx$	1508
3.205	$\int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx$	1516
3.206	$\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{x} dx$	1524
3.207	$\int \frac{1}{x \sqrt{\tanh(a+b \log(cx^n))}} dx$	1531
3.208	$\int \frac{1}{x \tanh^{\frac{3}{2}}(a+b \log(cx^n))} dx$	1538
3.209	$\int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx$	1546
3.210	$\int \frac{\tanh^5(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1554
3.211	$\int \frac{\tanh^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1562
3.212	$\int \frac{\tanh(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1569
3.213	$\int \frac{\coth(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1576
3.214	$\int \frac{\coth^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$	1582
3.215	$\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$	1589
3.216	$\int e^{a+bx} \tanh^4(a + bx) dx$	1598
3.217	$\int e^{a+bx} \tanh^3(a + bx) dx$	1604
3.218	$\int e^{a+bx} \tanh^2(a + bx) dx$	1610
3.219	$\int e^{a+bx} \tanh(a + bx) dx$	1616
3.220	$\int e^{a+bx} \coth(a + bx) dx$	1621
3.221	$\int e^{a+bx} \coth^2(a + bx) dx$	1626

3.222	$\int e^{a+bx} \coth^3(a+bx) dx$	1632
3.223	$\int e^{a+bx} \coth^4(a+bx) dx$	1638
3.224	$\int e^x \tanh^2(2x) dx$	1645
3.225	$\int e^x \tanh(2x) dx$	1651
3.226	$\int e^x \coth(2x) dx$	1659
3.227	$\int e^x \coth^2(2x) dx$	1665
3.228	$\int e^x \coth^4(2x) dx$	1671
3.229	$\int e^x \tanh^2(3x) dx$	1678
3.230	$\int e^x \tanh(3x) dx$	1685
3.231	$\int e^x \coth(3x) dx$	1693
3.232	$\int e^x \coth^2(3x) dx$	1701
3.233	$\int e^x \tanh^2(4x) dx$	1708
3.234	$\int e^x \tanh(4x) dx$	1716
3.235	$\int e^x \coth(4x) dx$	1727
3.236	$\int e^x \coth^2(4x) dx$	1737
3.237	$\int \frac{e^x}{a-\tanh(2x)} dx$	1744
3.238	$\int \frac{e^x}{(a-\tanh(2x))^2} dx$	1752
3.239	$\int e^{c(a+bx)} \tanh^3(d+ex) dx$	1761
3.240	$\int e^{c(a+bx)} \tanh^2(d+ex) dx$	1767
3.241	$\int e^{c(a+bx)} \tanh(d+ex) dx$	1773
3.242	$\int e^{c(a+bx)} \coth(d+ex) dx$	1778
3.243	$\int e^{c(a+bx)} \coth^2(d+ex) dx$	1783
3.244	$\int e^{c(a+bx)} \coth^3(d+ex) dx$	1789
3.245	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{5/2} dx$	1795
3.246	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{3/2} dx$	1803
3.247	$\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx$	1810
3.248	$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$	1816
3.249	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$	1822
3.250	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$	1830
3.251	$\int \sin^3(\tanh(a+bx)) dx$	1838
3.252	$\int \sin^2(\tanh(a+bx)) dx$	1845
3.253	$\int \sin(\tanh(a+bx)) dx$	1851
3.254	$\int \csc(\tanh(a+bx)) dx$	1857
3.255	$\int \cos^3(\tanh(a+bx)) dx$	1862
3.256	$\int \cos^2(\tanh(a+bx)) dx$	1869
3.257	$\int \cos(\tanh(a+bx)) dx$	1875
3.258	$\int \sec(\tanh(a+bx)) dx$	1881

### 3.1 $\int \tanh^6(a + bx) dx$

Optimal result . . . . .	119
Mathematica [A] (verified) . . . . .	119
Rubi [A] (verified) . . . . .	120
Maple [A] (verified) . . . . .	122
Fricas [B] (verification not implemented) . . . . .	122
Sympy [A] (verification not implemented) . . . . .	123
Maxima [B] (verification not implemented) . . . . .	123
Giac [A] (verification not implemented) . . . . .	124
Mupad [B] (verification not implemented) . . . . .	124
Reduce [B] (verification not implemented) . . . . .	125

#### Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \tanh^6(a + bx) dx = x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b}$$

output

```
x-tanh(b*x+a)/b-1/3*tanh(b*x+a)^3/b-1/5*tanh(b*x+a)^5/b
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

$$\int \tanh^6(a + bx) dx = \frac{\operatorname{arctanh}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b}$$

input

```
Integrate[Tanh[a + b*x]^6,x]
```

output

```
ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b) - Tanh[a + b*x]^5/(5*b)
```



**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$ , Rules used = {3042, 25, 3954, 3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^6(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan(ia + ibx)^6 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan(ia + ibx)^6 dx \\
 & \quad \downarrow \text{3954} \\
 & \int \tanh^4(a + bx) dx - \frac{\tanh^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\tanh^5(a + bx)}{5b} + \int \tan(ia + ibx)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & -\int -\tanh^2(a + bx) dx - \frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25} \\
 & \int \tanh^2(a + bx) dx - \frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan(ia + ibx)^2 dx - \frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
& - \int \tan(ia + ibx)^2 dx - \frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} \\
& \quad \downarrow \text{3954} \\
& \int 1 dx - \frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} \\
& \quad \downarrow \text{24} \\
& -\frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} + x
\end{aligned}$$

input `Int[Tanh[a + b*x]^6, x]`

output `x - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b) - Tanh[a + b*x]^5/(5*b)`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

**Maple [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

method	result	size
parallelrisch	$-\frac{3 \tanh(bx+a)^5 + 5 \tanh(bx+a)^3 - 15bx + 15 \tanh(bx+a)}{15b}$	39
derivativedivides	$\frac{-\frac{\tanh(bx+a)^5}{5} - \frac{\tanh(bx+a)^3}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(1+\tanh(bx+a))}{2}}{b}$	56
default	$\frac{-\frac{\tanh(bx+a)^5}{5} - \frac{\tanh(bx+a)^3}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(1+\tanh(bx+a))}{2}}{b}$	56
risch	$x + \frac{6e^{8bx+8a} + 12e^{6bx+6a} + \frac{56e^{4bx+4a}}{3} + \frac{28e^{2bx+2a}}{3} + \frac{46}{15}}{b(1+e^{2bx+2a})^5}$	67

input `int(tanh(b*x+a)^6,x,method=_RETURNVERBOSE)`output 
$$-1/15*(3*\tanh(b*x+a)^5+5*\tanh(b*x+a)^3-15*b*x+15*\tanh(b*x+a))/b$$
**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(39) = 78.

Time = 0.08 (sec) , antiderivative size = 254, normalized size of antiderivative = 5.91

$$\int \tanh^6(a + bx) dx$$

$$= \frac{(15bx + 23) \cosh(bx + a)^5 + 5(15bx + 23) \cosh(bx + a) \sinh(bx + a)^4 - 23 \sinh(bx + a)^5 + 5(15bx + 23) \cosh(bx + a) \sinh(bx + a)^3 - 23 \sinh(bx + a)^3 \cosh(bx + a) + 5(15bx + 23) \cosh(bx + a) \sinh(bx + a) - 23 \sinh(bx + a) \cosh(bx + a)}{15(b \cosh(bx + a) \sinh(bx + a) + \cosh^2(bx + a) - \sinh^2(bx + a))}$$

input `integrate(tanh(b*x+a)^6,x, algorithm="fricas")`

output

```
1/15*((15*b*x + 23)*cosh(b*x + a)^5 + 5*(15*b*x + 23)*cosh(b*x + a)*sinh(b*x + a)^4 - 23*sinh(b*x + a)^5 + 5*(15*b*x + 23)*cosh(b*x + a)^3 - 5*(46*cosh(b*x + a)^2 + 5)*sinh(b*x + a)^3 + 5*(2*(15*b*x + 23)*cosh(b*x + a)^3 + 3*(15*b*x + 23)*cosh(b*x + a))*sinh(b*x + a)^2 + 10*(15*b*x + 23)*cosh(b*x + a) - 5*(23*cosh(b*x + a)^4 + 15*cosh(b*x + a)^2 + 10)*sinh(b*x + a))/(b*cosh(b*x + a)^5 + 5*b*cosh(b*x + a)*sinh(b*x + a)^4 + 5*b*cosh(b*x + a)^3 + 5*(2*b*cosh(b*x + a)^3 + 3*b*cosh(b*x + a))*sinh(b*x + a)^2 + 10*b*cosh(b*x + a))
```

**Sympy [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

$$\int \tanh^6(a + bx) dx = \begin{cases} x - \frac{\tanh^5(a+bx)}{5b} - \frac{\tanh^3(a+bx)}{3b} - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^6(a) & \text{otherwise} \end{cases}$$

input

```
integrate(tanh(b*x+a)**6,x)
```

output

```
Piecewise((x - tanh(a + b*x)**5/(5*b) - tanh(a + b*x)**3/(3*b) - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**6, True))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 115 vs.  $2(39) = 78$ .

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 2.67

$$\int \tanh^6(a + bx) dx = x + \frac{a}{b} - \frac{2(70e^{(-2bx-2a)} + 140e^{(-4bx-4a)} + 90e^{(-6bx-6a)} + 45e^{(-8bx-8a)} + 23)}{15b(5e^{(-2bx-2a)} + 10e^{(-4bx-4a)} + 10e^{(-6bx-6a)} + 5e^{(-8bx-8a)} + e^{(-10bx-10a)} + 1)}$$

input

```
integrate(tanh(b*x+a)^6,x, algorithm="maxima")
```

output

```
x + a/b - 2/15*(70*e^(-2*b*x - 2*a) + 140*e^(-4*b*x - 4*a) + 90*e^(-6*b*x - 6*a) + 45*e^(-8*b*x - 8*a) + 23)/(b*(5*e^(-2*b*x - 2*a) + 10*e^(-4*b*x - 4*a) + 10*e^(-6*b*x - 6*a) + 5*e^(-8*b*x - 8*a) + e^(-10*b*x - 10*a) + 1) )
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.72

$$\int \tanh^6(a + bx) dx = \frac{15bx + 15a + \frac{2(45e^{(8bx+8a)} + 90e^{(6bx+6a)} + 140e^{(4bx+4a)} + 70e^{(2bx+2a)} + 23)}{(e^{(2bx+2a)} + 1)^5}}{15b}$$

input

```
integrate(tanh(b*x+a)^6,x, algorithm="giac")
```

output

```
1/15*(15*b*x + 15*a + 2*(45*e^(8*b*x + 8*a) + 90*e^(6*b*x + 6*a) + 140*e^(4*b*x + 4*a) + 70*e^(2*b*x + 2*a) + 23)/(e^(2*b*x + 2*a) + 1)^5)/b
```

**Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \tanh^6(a + bx) dx = x - \frac{\tanh(a+bx)^5}{5} + \frac{\tanh(a+bx)^3}{3} + \tanh(a + bx)$$

input

```
int(tanh(a + b*x)^6,x)
```

output

```
x - (tanh(a + b*x) + tanh(a + b*x)^3/3 + tanh(a + b*x)^5/5)/b
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int \tanh^6(a + bx) dx = \frac{-3 \tanh^5(bx + a) - 5 \tanh^3(bx + a) - 15 \tanh(bx + a) + 15bx}{15b}$$

input `int(tanh(b*x+a)^6,x)`

output `( - 3*tanh(a + b*x)**5 - 5*tanh(a + b*x)**3 - 15*tanh(a + b*x) + 15*b*x)/(15*b)`

## 3.2 $\int \tanh^5(a + bx) dx$

Optimal result	126
Mathematica [A] (verified)	126
Rubi [C] (verified)	127
Maple [A] (verified)	129
Fricas [B] (verification not implemented)	129
Sympy [A] (verification not implemented)	130
Maxima [B] (verification not implemented)	131
Giac [A] (verification not implemented)	131
Mupad [B] (verification not implemented)	132
Reduce [B] (verification not implemented)	132

### Optimal result

Integrand size = 8, antiderivative size = 42

$$\int \tanh^5(a + bx) dx = \frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b} - \frac{\tanh^4(a + bx)}{4b}$$

output

```
ln(cosh(b*x+a))/b-1/2*tanh(b*x+a)^2/b-1/4*tanh(b*x+a)^4/b
```

### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int \tanh^5(a + bx) dx = \frac{4 \log(\cosh(a + bx)) + 4 \operatorname{sech}^2(a + bx) - \operatorname{sech}^4(a + bx)}{4b}$$

input

```
Integrate[Tanh[a + b*x]^5,x]
```

output

```
(4*Log[Cosh[a + b*x]] + 4*Sech[a + b*x]^2 - Sech[a + b*x]^4)/(4*b)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.26, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {3042, 26, 3954, 26, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^5(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \tan(ia + ibx)^5 dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \tan(ia + ibx)^5 dx \\
 & \quad \downarrow \text{3954} \\
 & -i \left( - \int -i \tanh^3(a + bx) dx - \frac{i \tanh^4(a + bx)}{4b} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( i \int \tanh^3(a + bx) dx - \frac{i \tanh^4(a + bx)}{4b} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( i \int i \tan(ia + ibx)^3 dx - \frac{i \tanh^4(a + bx)}{4b} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( - \int \tan(ia + ibx)^3 dx - \frac{i \tanh^4(a + bx)}{4b} \right) \\
 & \quad \downarrow \text{3954} \\
 & -i \left( \int i \tanh(a + bx) dx - \frac{i \tanh^4(a + bx)}{4b} - \frac{i \tanh^2(a + bx)}{2b} \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$



$$\begin{aligned}
& -i \left( i \int \tanh(a + bx) dx - \frac{i \tanh^4(a + bx)}{4b} - \frac{i \tanh^2(a + bx)}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& -i \left( i \int -i \tan(ia + ibx) dx - \frac{i \tanh^4(a + bx)}{4b} - \frac{i \tanh^2(a + bx)}{2b} \right) \\
& \quad \downarrow \text{26} \\
& -i \left( \int \tan(ia + ibx) dx - \frac{i \tanh^4(a + bx)}{4b} - \frac{i \tanh^2(a + bx)}{2b} \right) \\
& \quad \downarrow \text{3956} \\
& -i \left( -\frac{i \tanh^4(a + bx)}{4b} - \frac{i \tanh^2(a + bx)}{2b} + \frac{i \log(\cosh(a + bx))}{b} \right)
\end{aligned}$$

input `Int[Tanh[a + b*x]^5, x]`

output `(-I)*((I*Log[Cosh[a + b*x]])/b - ((I/2)*Tanh[a + b*x]^2)/b - ((I/4)*Tanh[a + b*x]^4)/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

**Maple [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

method	result	size
parallelrisch	$-\frac{\tanh(bx+a)^4 + 4bx + 2 \tanh(bx+a)^2 + 4 \ln(1 - \tanh(bx+a))}{4b}$	42
derivativedivides	$-\frac{\frac{\tanh(bx+a)^4}{4} - \frac{\tanh(bx+a)^2}{2} - \frac{\ln(-1 + \tanh(bx+a))}{2} - \frac{\ln(1 + \tanh(bx+a))}{2}}{b}$	48
default	$-\frac{\frac{\tanh(bx+a)^4}{4} - \frac{\tanh(bx+a)^2}{2} - \frac{\ln(-1 + \tanh(bx+a))}{2} - \frac{\ln(1 + \tanh(bx+a))}{2}}{b}$	48
risch	$-x - \frac{2a}{b} + \frac{4e^{2bx+2a}(e^{4bx+4a} + e^{2bx+2a} + 1)}{b(1+e^{2bx+2a})^4} + \frac{\ln(1+e^{2bx+2a})}{b}$	74

input `int(tanh(b*x+a)^5,x,method=_RETURNVERBOSE)`

output `-1/4*(tanh(b*x+a)^4+4*b*x+2*tanh(b*x+a)^2+4*ln(1-tanh(b*x+a)))/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 968 vs. 2(38) = 76.

Time = 0.10 (sec) , antiderivative size = 968, normalized size of antiderivative = 23.05

$$\int \tanh^5(a + bx) dx = \text{Too large to display}$$

input `integrate(tanh(b*x+a)^5,x, algorithm="fricas")`

output

```

-(b*x*cosh(b*x + a)^8 + 8*b*x*cosh(b*x + a)*sinh(b*x + a)^7 + b*x*sinh(b*x
+ a)^8 + 4*(b*x - 1)*cosh(b*x + a)^6 + 4*(7*b*x*cosh(b*x + a)^2 + b*x - 1
)*sinh(b*x + a)^6 + 8*(7*b*x*cosh(b*x + a)^3 + 3*(b*x - 1)*cosh(b*x + a))*
sinh(b*x + a)^5 + 2*(3*b*x - 2)*cosh(b*x + a)^4 + 2*(35*b*x*cosh(b*x + a)^
4 + 30*(b*x - 1)*cosh(b*x + a)^2 + 3*b*x - 2)*sinh(b*x + a)^4 + 8*(7*b*x*c
osh(b*x + a)^5 + 10*(b*x - 1)*cosh(b*x + a)^3 + (3*b*x - 2)*cosh(b*x + a))
*sinh(b*x + a)^3 + 4*(b*x - 1)*cosh(b*x + a)^2 + 4*(7*b*x*cosh(b*x + a)^6
+ 15*(b*x - 1)*cosh(b*x + a)^4 + 3*(3*b*x - 2)*cosh(b*x + a)^2 + b*x - 1)*
sinh(b*x + a)^2 + b*x - (cosh(b*x + a)^8 + 8*cosh(b*x + a)*sinh(b*x + a)^7
+ sinh(b*x + a)^8 + 4*(7*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^6 + 4*cosh(b*
x + a)^6 + 8*(7*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(35
*cosh(b*x + a)^4 + 30*cosh(b*x + a)^2 + 3)*sinh(b*x + a)^4 + 6*cosh(b*x +
a)^4 + 8*(7*cosh(b*x + a)^5 + 10*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b
*x + a)^3 + 4*(7*cosh(b*x + a)^6 + 15*cosh(b*x + a)^4 + 9*cosh(b*x + a)^2
+ 1)*sinh(b*x + a)^2 + 4*cosh(b*x + a)^2 + 8*(cosh(b*x + a)^7 + 3*cosh(b*x
+ a)^5 + 3*cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x + a) + 1)*log(2*cosh
(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))) + 8*(b*x*cosh(b*x + a)^7 + 3*(b
*x - 1)*cosh(b*x + a)^5 + (3*b*x - 2)*cosh(b*x + a)^3 + (b*x - 1)*cosh(b*x
+ a))*sinh(b*x + a))/(b*cosh(b*x + a)^8 + 8*b*cosh(b*x + a)*sinh(b*x + a)
^7 + b*sinh(b*x + a)^8 + 4*b*cosh(b*x + a)^6 + 4*(7*b*cosh(b*x + a)^2 + ...

```

### Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \tanh^5(a + bx) dx = \begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} - \frac{\tanh^4(a+bx)}{4b} - \frac{\tanh^2(a+bx)}{2b} & \text{for } b \neq 0 \\ x \tanh^5(a) & \text{otherwise} \end{cases}$$

input

```
integrate(tanh(b*x+a)**5,x)
```

output

```
Piecewise((x - log(tanh(a + b*x) + 1)/b - tanh(a + b*x)**4/(4*b) - tanh(a
+ b*x)**2/(2*b), Ne(b, 0)), (x*tanh(a)**5, True))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 102 vs.  $2(38) = 76$ .

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.43

$$\int \tanh^5(a + bx) dx = x + \frac{a}{b} + \frac{\log(e^{(-2bx-2a)} + 1)}{b} + \frac{4(e^{(-2bx-2a)} + e^{(-4bx-4a)} + e^{(-6bx-6a)})}{b(4e^{(-2bx-2a)} + 6e^{(-4bx-4a)} + 4e^{(-6bx-6a)} + e^{(-8bx-8a)} + 1)}$$

input `integrate(tanh(b*x+a)^5,x, algorithm="maxima")`

output `x + a/b + log(e^(-2*b*x - 2*a) + 1)/b + 4*(e^(-2*b*x - 2*a) + e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a))/(b*(4*e^(-2*b*x - 2*a) + 6*e^(-4*b*x - 4*a) + 4*e^(-6*b*x - 6*a) + e^(-8*b*x - 8*a) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.60

$$\int \tanh^5(a + bx) dx = -\frac{bx + a - \frac{4(e^{(6bx+6a)} + e^{(4bx+4a)} + e^{(2bx+2a)})}{(e^{(2bx+2a)} + 1)^4} - \log(e^{(2bx+2a)} + 1)}{b}$$

input `integrate(tanh(b*x+a)^5,x, algorithm="giac")`

output `-(b*x + a - 4*(e^(6*b*x + 6*a) + e^(4*b*x + 4*a) + e^(2*b*x + 2*a)))/(e^(2*b*x + 2*a) + 1)^4 - log(e^(2*b*x + 2*a) + 1))/b`

**Mupad [B] (verification not implemented)**

Time = 2.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int \tanh^5(a + bx) dx = x - \frac{\ln(\tanh(a + bx) + 1) + \frac{\tanh(a+bx)^2}{2} + \frac{\tanh(a+bx)^4}{4}}{b}$$

input `int(tanh(a + b*x)^5,x)`output `x - (log(tanh(a + b*x) + 1) + tanh(a + b*x)^2/2 + tanh(a + b*x)^4/4)/b`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 252, normalized size of antiderivative = 6.00

$$\int \tanh^5(a + bx) dx = \frac{e^{8bx+8a} \log(e^{2bx+2a} + 1) - e^{8bx+8a} bx - e^{8bx+8a} + 4e^{6bx+6a} \log(e^{2bx+2a} + 1) - 4e^{6bx+6a} bx + 6e^{4bx+4a} \log(e^{2bx+2a} + 1) - 6e^{4bx+4a} bx - 2e^{4bx+4a} + 4e^{2bx+2a} \log(e^{2bx+2a} + 1) - 4e^{2bx+2a} bx + \log(e^{2bx+2a} + 1) - bx - 1}{b(e^{8bx+8a} + 4e^{6bx+6a} + 6e^{4bx+4a} + 4e^{2bx+2a} + 1)}$$

input `int(tanh(b*x+a)^5,x)`output `(e**(8*a + 8*b*x)*log(e**(2*a + 2*b*x) + 1) - e**(8*a + 8*b*x)*b*x - e**(8*a + 8*b*x) + 4*e**(6*a + 6*b*x)*log(e**(2*a + 2*b*x) + 1) - 4*e**(6*a + 6*b*x)*b*x + 6*e**(4*a + 4*b*x)*log(e**(2*a + 2*b*x) + 1) - 6*e**(4*a + 4*b*x)*b*x - 2*e**(4*a + 4*b*x) + 4*e**(2*a + 2*b*x)*log(e**(2*a + 2*b*x) + 1) - 4*e**(2*a + 2*b*x)*b*x + log(e**(2*a + 2*b*x) + 1) - b*x - 1)/(b*(e**(8*a + 8*b*x) + 4*e**(6*a + 6*b*x) + 6*e**(4*a + 4*b*x) + 4*e**(2*a + 2*b*x) + 1))`

### 3.3 $\int \tanh^4(a + bx) dx$

Optimal result . . . . .	133
Mathematica [A] (verified) . . . . .	133
Rubi [A] (verified) . . . . .	134
Maple [A] (verified) . . . . .	135
Fricas [B] (verification not implemented) . . . . .	136
Sympy [A] (verification not implemented) . . . . .	136
Maxima [B] (verification not implemented) . . . . .	137
Giac [A] (verification not implemented) . . . . .	137
Mupad [B] (verification not implemented) . . . . .	137
Reduce [B] (verification not implemented) . . . . .	138

#### Optimal result

Integrand size = 8, antiderivative size = 28

$$\int \tanh^4(a + bx) dx = x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b}$$

output

```
x-tanh(b*x+a)/b-1/3*tanh(b*x+a)^3/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\int \tanh^4(a + bx) dx = \frac{\operatorname{arctanh}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b}$$

input

```
Integrate[Tanh[a + b*x]^4,x]
```

output

```
ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b)
```

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^4(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(ia + ibx)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int -\tanh^2(a + bx) dx - \frac{\tanh^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25} \\
 & \int \tanh^2(a + bx) dx - \frac{\tanh^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\tanh^3(a + bx)}{3b} + \int -\tan(ia + ibx)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\frac{\tanh^3(a + bx)}{3b} - \int \tan(ia + ibx)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & -\frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} + x
 \end{aligned}$$

input

`Int[Tanh[a + b*x]^4, x]`

output  $x - \tanh[a + b*x]/b - \tanh[a + b*x]^3/(3*b)$

### Defintions of rubi rules used

rule 24  $\text{Int}[a_, x\_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 25  $\text{Int}[-(F x_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F x, x], x]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\text{Int}[(b \cdot \tan(c \cdot x) + d \cdot x)^n, x\_Symbol] \rightarrow \text{Simp}[b \cdot ((b \cdot \tan[c + d \cdot x])^{n-1} / (d \cdot (n-1))), x] - \text{Simp}[b^2 \text{Int}[(b \cdot \tan[c + d \cdot x])^{n-2}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1]$

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

method	result	size
parallelrisch	$-\frac{\tanh(bx+a)^3 - 3bx + 3 \tanh(bx+a)}{3b}$	27
risch	$x + \frac{4e^{4bx+4a} + 4e^{2bx+2a} + \frac{8}{3}}{b(1+e^{2bx+2a})^3}$	45
derivativedivides	$\frac{-\frac{\tanh(bx+a)^3}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(1+\tanh(bx+a))}{2}}{b}$	46
default	$\frac{-\frac{\tanh(bx+a)^3}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(1+\tanh(bx+a))}{2}}{b}$	46

input  $\text{int}(\tanh(b*x+a)^4, x, \text{method}=\_RETURNVERBOSE)$

output  $-1/3*(\tanh(b*x+a)^3 - 3*b*x + 3*\tanh(b*x+a))/b$



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 119 vs.  $2(26) = 52$ .

Time = 0.09 (sec) , antiderivative size = 119, normalized size of antiderivative = 4.25

$$\int \tanh^4(a + bx) dx$$

$$= \frac{(3bx + 4) \cosh(bx + a)^3 + 3(3bx + 4) \cosh(bx + a) \sinh(bx + a)^2 - 12 \cosh(bx + a)^2 \sinh(bx + a) - 4 \sinh(bx + a)^3 + 3(3bx + 4) \cosh(bx + a)}{3(b \cosh(bx + a))^3 + 3b \cosh(bx + a) \sinh(bx + a)^2 + 3b \cosh(bx + a)^2 \sinh(bx + a) + 3b \sinh(bx + a)^3}$$

input `integrate(tanh(b*x+a)^4,x, algorithm="fricas")`

output `1/3*((3*b*x + 4)*cosh(b*x + a)^3 + 3*(3*b*x + 4)*cosh(b*x + a)*sinh(b*x + a)^2 - 12*cosh(b*x + a)^2*sinh(b*x + a) - 4*sinh(b*x + a)^3 + 3*(3*b*x + 4)*cosh(b*x + a))/(b*cosh(b*x + a)^3 + 3*b*cosh(b*x + a)*sinh(b*x + a)^2 + 3*b*cosh(b*x + a)^2*sinh(b*x + a) + 3*b*sinh(b*x + a)^3)`

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \tanh^4(a + bx) dx = \begin{cases} x - \frac{\tanh^3(a+bx)}{3b} - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^4(a) & \text{otherwise} \end{cases}$$

input `integrate(tanh(b*x+a)**4,x)`

output `Piecewise((x - tanh(a + b*x)**3/(3*b) - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**4, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(26) = 52$ .

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.54

$$\int \tanh^4(a + bx) dx = x + \frac{a}{b} - \frac{4(3e^{(-2bx-2a)} + 3e^{(-4bx-4a)} + 2)}{3b(3e^{(-2bx-2a)} + 3e^{(-4bx-4a)} + e^{(-6bx-6a)} + 1)}$$

input `integrate(tanh(b*x+a)^4,x, algorithm="maxima")`

output `x + a/b - 4/3*(3*e^(-2*b*x - 2*a) + 3*e^(-4*b*x - 4*a) + 2)/(b*(3*e^(-2*b*x - 2*a) + 3*e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.86

$$\int \tanh^4(a + bx) dx = \frac{3bx + 3a + \frac{4(3e^{(4bx+4a)} + 3e^{(2bx+2a)} + 2)}{(e^{(2bx+2a)} + 1)^3}}{3b}$$

input `integrate(tanh(b*x+a)^4,x, algorithm="giac")`

output `1/3*(3*b*x + 3*a + 4*(3*e^(4*b*x + 4*a) + 3*e^(2*b*x + 2*a) + 2)/(e^(2*b*x + 2*a) + 1)^3)/b`

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \tanh^4(a + bx) dx = x - \frac{\frac{\tanh(a+bx)^3}{3} + \tanh(a + bx)}{b}$$

input `int(tanh(a + b*x)^4,x)`

output `x - (tanh(a + b*x) + tanh(a + b*x)^3/3)/b`

### Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \tanh^4(a + bx) dx = \frac{-\tanh(bx + a)^3 - 3\tanh(bx + a) + 3bx}{3b}$$

input `int(tanh(b*x+a)^4,x)`

output `( - tanh(a + b*x)**3 - 3*tanh(a + b*x) + 3*b*x)/(3*b)`

### 3.4 $\int \tanh^3(a + bx) dx$

Optimal result . . . . .	139
Mathematica [A] (verified) . . . . .	139
Rubi [C] (verified) . . . . .	140
Maple [A] (verified) . . . . .	141
Fricas [B] (verification not implemented) . . . . .	142
Sympy [A] (verification not implemented) . . . . .	142
Maxima [B] (verification not implemented) . . . . .	143
Giac [A] (verification not implemented) . . . . .	143
Mupad [B] (verification not implemented) . . . . .	144
Reduce [B] (verification not implemented) . . . . .	144

#### Optimal result

Integrand size = 8, antiderivative size = 27

$$\int \tanh^3(a + bx) dx = \frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b}$$

output

```
ln(cosh(b*x+a))/b-1/2*tanh(b*x+a)^2/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \tanh^3(a + bx) dx = \frac{2 \log(\cosh(a + bx)) + \operatorname{sech}^2(a + bx)}{2b}$$

input

```
Integrate[Tanh[a + b*x]^3,x]
```

output

```
(2*Log[Cosh[a + b*x]] + Sech[a + b*x]^2)/(2*b)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.33, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \tan(ia + ibx)^3 dx \\
 & \quad \downarrow \text{26} \\
 & i \int \tan(ia + ibx)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & i \left( \frac{i \tanh^2(a + bx)}{2b} - \int i \tanh(a + bx) dx \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( \frac{i \tanh^2(a + bx)}{2b} - i \int \tanh(a + bx) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & i \left( \frac{i \tanh^2(a + bx)}{2b} - i \int -i \tan(ia + ibx) dx \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( \frac{i \tanh^2(a + bx)}{2b} - \int \tan(ia + ibx) dx \right) \\
 & \quad \downarrow \text{3956} \\
 & i \left( \frac{i \tanh^2(a + bx)}{2b} - \frac{i \log(\cosh(a + bx))}{b} \right)
 \end{aligned}$$

input `Int[Tanh[a + b*x]^3,x]`

output `I*(((I)*Log[Cosh[a + b*x]])/b + ((I/2)*Tanh[a + b*x]^2)/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

### Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19

method	result	size
parallelrisch	$-\frac{2bx + \tanh(bx+a)^2 + 2\ln(1 - \tanh(bx+a))}{2b}$	32
derivativedivides	$\frac{-\frac{\tanh(bx+a)^2}{2} - \frac{\ln(-1 + \tanh(bx+a))}{2} - \frac{\ln(1 + \tanh(bx+a))}{2}}{b}$	38
default	$\frac{-\frac{\tanh(bx+a)^2}{2} - \frac{\ln(-1 + \tanh(bx+a))}{2} - \frac{\ln(1 + \tanh(bx+a))}{2}}{b}$	38
risch	$-x - \frac{2a}{b} + \frac{2e^{2bx+2a}}{b(1+e^{2bx+2a})^2} + \frac{\ln(1+e^{2bx+2a})}{b}$	54

input `int(tanh(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
-1/2*(2*b*x+tanh(b*x+a)^2+2*ln(1-tanh(b*x+a)))/b
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(25) = 50.

Time = 0.10 (sec) , antiderivative size = 339, normalized size of antiderivative = 12.56

$$\int \tanh^3(a + bx) dx =$$

$$\frac{bx \cosh(bx + a)^4 + 4bx \cosh(bx + a) \sinh(bx + a)^3 + bx \sinh(bx + a)^4 + 2(bx - 1) \cosh(bx + a)^2 + \dots}{\dots}$$

input

```
integrate(tanh(b*x+a)^3,x, algorithm="fricas")
```

output

```
-(b*x*cosh(b*x + a)^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 + 2*(b*x - 1)*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 + b*x - 1)*sinh(b*x + a)^2 + b*x - (cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x + a) + 1)*log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))) + 4*(b*x*cosh(b*x + a)^3 + (b*x - 1)*cosh(b*x + a))*sinh(b*x + a))/(b*cosh(b*x + a)^4 + 4*b*cosh(b*x + a)*sinh(b*x + a)^3 + b*sinh(b*x + a)^4 + 2*b*cosh(b*x + a)^2 + 2*(3*b*cosh(b*x + a)^2 + b)*sinh(b*x + a)^2 + 4*(b*cosh(b*x + a)^3 + b*cosh(b*x + a))*sinh(b*x + a) + b)
```

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int \tanh^3(a + bx) dx = \begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} - \frac{\tanh^2(a+bx)}{2b} & \text{for } b \neq 0 \\ x \tanh^3(a) & \text{otherwise} \end{cases}$$

input

```
integrate(tanh(b*x+a)**3,x)
```

output `Piecewise((x - log(tanh(a + b*x) + 1)/b - tanh(a + b*x)**2/(2*b), Ne(b, 0)), (x*tanh(a)**3, True))`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs.  $2(25) = 50$ .

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.26

$$\int \tanh^3(a + bx) dx = x + \frac{a}{b} + \frac{\log(e^{-2bx-2a} + 1)}{b} + \frac{2e^{-2bx-2a}}{b(2e^{-2bx-2a} + e^{-4bx-4a} + 1)}$$

input `integrate(tanh(b*x+a)^3,x, algorithm="maxima")`

output `x + a/b + log(e^(-2*b*x - 2*a) + 1)/b + 2*e^(-2*b*x - 2*a)/(b*(2*e^(-2*b*x - 2*a) + e^(-4*b*x - 4*a) + 1))`

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.78

$$\int \tanh^3(a + bx) dx = -\frac{bx + a - \frac{2e^{(2bx+2a)}}{(e^{(2bx+2a)}+1)^2} - \log(e^{(2bx+2a)} + 1)}{b}$$

input `integrate(tanh(b*x+a)^3,x, algorithm="giac")`

output `-(b*x + a - 2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)^2 - log(e^(2*b*x + 2*a) + 1))/b`



**Mupad [B] (verification not implemented)**

Time = 2.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \tanh^3(a + bx) dx = x - \frac{\ln(\tanh(a + bx) + 1) + \frac{\tanh(a + bx)^2}{2}}{b}$$

input `int(tanh(a + b*x)^3,x)`output `x - (log(tanh(a + b*x) + 1) + tanh(a + b*x)^2/2)/b`**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 138, normalized size of antiderivative = 5.11

$$\int \tanh^3(a + bx) dx = \frac{e^{4bx+4a} \log(e^{2bx+2a} + 1) - e^{4bx+4a} bx - e^{4bx+4a} + 2e^{2bx+2a} \log(e^{2bx+2a} + 1) - 2e^{2bx+2a} bx + \log(e^{2bx+2a} + 1)}{b(e^{4bx+4a} + 2e^{2bx+2a} + 1)}$$

input `int(tanh(b*x+a)^3,x)`output `(e**(4*a + 4*b*x)*log(e**(2*a + 2*b*x) + 1) - e**(4*a + 4*b*x)*b*x - e**(4*a + 4*b*x) + 2*e**(2*a + 2*b*x)*log(e**(2*a + 2*b*x) + 1) - 2*e**(2*a + 2*b*x)*b*x + log(e**(2*a + 2*b*x) + 1) - b*x - 1)/(b*(e**(4*a + 4*b*x) + 2*e**(2*a + 2*b*x) + 1))`

### 3.5 $\int \tanh^2(a + bx) dx$

Optimal result	145
Mathematica [A] (verified)	145
Rubi [A] (verified)	146
Maple [A] (verified)	147
Fricas [B] (verification not implemented)	148
Sympy [A] (verification not implemented)	148
Maxima [A] (verification not implemented)	148
Giac [A] (verification not implemented)	149
Mupad [B] (verification not implemented)	149
Reduce [B] (verification not implemented)	149

#### Optimal result

Integrand size = 8, antiderivative size = 13

$$\int \tanh^2(a + bx) dx = x - \frac{\tanh(a + bx)}{b}$$

output `x-tanh(b*x+a)/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.77

$$\int \tanh^2(a + bx) dx = \frac{\operatorname{arctanh}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b}$$

input `Integrate[Tanh[a + b*x]^2,x]`

output `ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan(ia + ibx)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan(ia + ibx)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{\tanh(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & x - \frac{\tanh(a + bx)}{b}
 \end{aligned}$$

input `Int [Tanh[a + b*x]^2, x]`

output `x - Tanh[a + b*x]/b`

**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

**Maple [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

method	result	size
parallelrisch	$-\frac{-bx + \tanh(bx+a)}{b}$	17
risch	$x + \frac{2}{b(1+e^{2bx+2a})}$	21
derivativedivides	$\frac{-\tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2}}{b} + \frac{\ln(1+\tanh(bx+a))}{2}$	36
default	$\frac{-\tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2}}{b} + \frac{\ln(1+\tanh(bx+a))}{2}$	36

input `int(tanh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `-(-b*x+tanh(b*x+a))/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(13) = 26$ .

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.54

$$\int \tanh^2(a + bx) dx = \frac{(bx + 1) \cosh(bx + a) - \sinh(bx + a)}{b \cosh(bx + a)}$$

input `integrate(tanh(b*x+a)^2,x, algorithm="fricas")`

output `((b*x + 1)*cosh(b*x + a) - sinh(b*x + a))/(b*cosh(b*x + a))`

**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \tanh^2(a + bx) dx = \begin{cases} x - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^2(a) & \text{otherwise} \end{cases}$$

input `integrate(tanh(b*x+a)**2,x)`

output `Piecewise((x - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.92

$$\int \tanh^2(a + bx) dx = x + \frac{a}{b} - \frac{2}{b(e^{-2bx-2a} + 1)}$$

input `integrate(tanh(b*x+a)^2,x, algorithm="maxima")`

output `x + a/b - 2/(b*(e^(-2*b*x - 2*a) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.85

$$\int \tanh^2(a + bx) dx = \frac{bx + a + \frac{2}{e^{(2bx+2a)+1}}}{b}$$

input `integrate(tanh(b*x+a)^2,x, algorithm="giac")`

output `(b*x + a + 2/(e^(2*b*x + 2*a) + 1))/b`

**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \tanh^2(a + bx) dx = x - \frac{\tanh(a + bx)}{b}$$

input `int(tanh(a + b*x)^2,x)`

output `x - tanh(a + b*x)/b`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.23

$$\int \tanh^2(a + bx) dx = \frac{-\tanh(bx + a) + bx}{b}$$

input `int(tanh(b*x+a)^2,x)`

output `( - tanh(a + b*x) + b*x)/b`

### 3.6 $\int \tanh(a + bx) dx$

Optimal result . . . . .	150
Mathematica [A] (verified) . . . . .	150
Rubi [A] (verified) . . . . .	151
Maple [A] (verified) . . . . .	152
Fricas [B] (verification not implemented) . . . . .	152
Sympy [B] (verification not implemented) . . . . .	153
Maxima [A] (verification not implemented) . . . . .	153
Giac [B] (verification not implemented) . . . . .	153
Mupad [B] (verification not implemented) . . . . .	154
Reduce [B] (verification not implemented) . . . . .	154

#### Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \tanh(a + bx) dx = \frac{\log(\cosh(a + bx))}{b}$$

output `ln(cosh(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \tanh(a + bx) dx = \frac{\log(\cosh(a + bx))}{b}$$

input `Integrate[Tanh[a + b*x],x]`

output `Log[Cosh[a + b*x]]/b`

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tanh(a + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \tan(ia + ibx) dx \\ & \quad \downarrow \text{26} \\ & -i \int \tan(ia + ibx) dx \\ & \quad \downarrow \text{3956} \\ & \frac{\log(\cosh(a + bx))}{b} \end{aligned}$$

input `Int[Tanh[a + b*x], x]`

output `Log[Cosh[a + b*x]]/b`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativdivides	$\frac{\ln(\cosh(bx+a))}{b}$	12
default	$\frac{\ln(\cosh(bx+a))}{b}$	12
parallelrisc	$-\frac{bx + \ln(1 - \tanh(bx+a))}{b}$	21
risc	$-x - \frac{2a}{b} + \frac{\ln(1 + e^{2bx+2a})}{b}$	27

input

```
int(tanh(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
ln(cosh(b*x+a))/b
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(11) = 22.

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 3.36

$$\int \tanh(a + bx) dx = -\frac{bx - \log\left(\frac{2 \cosh(bx+a)}{\cosh(bx+a) - \sinh(bx+a)}\right)}{b}$$

input

```
integrate(tanh(b*x+a),x, algorithm="fricas")
```

output

```
-(b*x - log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/b
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 17 vs.  $2(8) = 16$ .

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \tanh(a + bx) dx = \begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} & \text{for } b \neq 0 \\ x \tanh(a) & \text{otherwise} \end{cases}$$

input `integrate(tanh(b*x+a),x)`

output `Piecewise((x - log(tanh(a + b*x) + 1)/b, Ne(b, 0)), (x*tanh(a), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \tanh(a + bx) dx = \frac{\log(\cosh(bx + a))}{b}$$

input `integrate(tanh(b*x+a),x, algorithm="maxima")`

output `log(cosh(b*x + a))/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 24 vs.  $2(11) = 22$ .

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.18

$$\int \tanh(a + bx) dx = -\frac{bx + a - \log(e^{2bx+2a} + 1)}{b}$$

input `integrate(tanh(b*x+a),x, algorithm="giac")`

output  $-(b*x + a - \log(e^{(2*b*x + 2*a) + 1}))/b$

### Mupad [B] (verification not implemented)

Time = 2.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

$$\int \tanh(a + bx) dx = x - \frac{\ln(\tanh(a + bx) + 1)}{b}$$

input `int(tanh(a + b*x),x)`

output  $x - \log(\tanh(a + b*x) + 1)/b$

### Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.00

$$\int \tanh(a + bx) dx = \frac{\log(e^{2bx+2a} + 1) - bx}{b}$$

input `int(tanh(b*x+a),x)`

output  $(\log(e^{(2*a + 2*b*x) + 1}) - b*x)/b$

### 3.7 $\int \coth(a + bx) dx$

Optimal result . . . . .	155
Mathematica [A] (verified) . . . . .	155
Rubi [C] (verified) . . . . .	156
Maple [A] (verified) . . . . .	157
Fricas [B] (verification not implemented) . . . . .	157
Sympy [B] (verification not implemented) . . . . .	158
Maxima [A] (verification not implemented) . . . . .	158
Giac [B] (verification not implemented) . . . . .	158
Mupad [B] (verification not implemented) . . . . .	159
Reduce [B] (verification not implemented) . . . . .	159

#### Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \coth(a + bx) dx = \frac{\log(\sinh(a + bx))}{b}$$

output `ln(sinh(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \coth(a + bx) dx = \frac{\log(\sinh(a + bx))}{b}$$

input `Integrate[Coth[a + b*x],x]`

output `Log[Sinh[a + b*x]]/b`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.36, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \coth(a + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{26} \\ & -i \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right) dx \\ & \quad \downarrow \text{3956} \\ & \frac{\log(-i \sinh(a + bx))}{b} \end{aligned}$$

input `Int[Coth[a + b*x],x]`

output `Log[(-I)*Sinh[a + b*x]]/b`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{\ln(\sinh(bx+a))}{b}$	12
default	$\frac{\ln(\sinh(bx+a))}{b}$	12
risch	$-x - \frac{2a}{b} + \frac{\ln(e^{2bx+2a}-1)}{b}$	27
parallelrisc	$\frac{-bx + \ln(\tanh(bx+a)) - \ln(1 - \tanh(bx+a))}{b}$	30

input

```
int(coth(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
ln(sinh(b*x+a))/b
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 37 vs.  $2(11) = 22$ .

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 3.36

$$\int \coth(a + bx) dx = -\frac{bx - \log\left(\frac{2 \sinh(bx+a)}{\cosh(bx+a) - \sinh(bx+a)}\right)}{b}$$

input

```
integrate(coth(b*x+a), x, algorithm="fricas")
```

output

```
-(b*x - log(2*sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/b
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs.  $2(8) = 16$ .

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.45

$$\int \coth(a + bx) dx = \begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} & \text{for } b \neq 0 \\ x \coth(a) & \text{otherwise} \end{cases}$$

input `integrate(coth(b*x+a), x)`

output `Piecewise((x - log(tanh(a + b*x) + 1)/b + log(tanh(a + b*x))/b, Ne(b, 0)), (x*coth(a), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \coth(a + bx) dx = \frac{\log(\sinh(bx + a))}{b}$$

input `integrate(coth(b*x+a), x, algorithm="maxima")`

output `log(sinh(b*x + a))/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 25 vs.  $2(11) = 22$ .

Time = 0.13 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.27

$$\int \coth(a + bx) dx = -\frac{bx + a - \log(|e^{(2bx+2a)} - 1|)}{b}$$

input `integrate(coth(b*x+a), x, algorithm="giac")`

output  $-(b*x + a - \log(\text{abs}(e^{(2*b*x + 2*a)} - 1)))/b$

### Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.91

$$\int \coth(a + bx) dx = \frac{\ln(e^{2a} e^{2bx} - 1)}{b} - x$$

input `int(coth(a + b*x), x)`

output  $\log(\exp(2*a)*\exp(2*b*x) - 1)/b - x$

### Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.64

$$\int \coth(a + bx) dx = \frac{\log(e^{bx+a} - 1) + \log(e^{bx+a} + 1) - bx}{b}$$

input `int(coth(b*x+a), x)`

output  $(\log(e^{(a + b*x)} - 1) + \log(e^{(a + b*x)} + 1) - b*x)/b$



### 3.8 $\int \coth^2(a + bx) dx$

Optimal result	160
Mathematica [C] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [B] (verification not implemented)	163
Sympy [B] (verification not implemented)	163
Maxima [A] (verification not implemented)	164
Giac [A] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

#### Optimal result

Integrand size = 8, antiderivative size = 13

$$\int \coth^2(a + bx) dx = x - \frac{\coth(a + bx)}{b}$$

output `x-coth(b*x+a)/b`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.08

$$\int \coth^2(a + bx) dx = -\frac{\coth(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \tanh^2(a + bx)\right)}{b}$$

input `Integrate[Coth[a + b*x]^2,x]`

output `-((Coth[a + b*x]*Hypergeometric2F1[-1/2, 1, 1/2, Tanh[a + b*x]^2])/b)`

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^2(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(ia + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \int 1dx - \frac{\coth(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & x - \frac{\coth(a + bx)}{b}
 \end{aligned}$$

input `Int[Coth[a + b*x]^2,x]`

output `x - Coth[a + b*x]/b`

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

## Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

method	result	size
parallelrisch	$\frac{bx - \operatorname{coth}(bx+a)}{b}$	17
risch	$x - \frac{2}{b(e^{2bx+2a}-1)}$	21
derivativedivides	$-\operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}$	36
default	$-\operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}$	36

input `int(coth(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `(b*x-coth(b*x+a))/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(13) = 26$ .

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.54

$$\int \coth^2(a + bx) dx = \frac{(bx + 1) \sinh(bx + a) - \cosh(bx + a)}{b \sinh(bx + a)}$$

input `integrate(coth(b*x+a)^2,x, algorithm="fricas")`

output `((b*x + 1)*sinh(b*x + a) - cosh(b*x + a))/(b*sinh(b*x + a))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(8) = 16$ .

Time = 0.51 (sec) , antiderivative size = 65, normalized size of antiderivative = 5.00

$$\int \coth^2(a + bx) dx = \begin{cases} x \coth^2(a) & \text{for } b = 0 \\ x \coth^2(bx + \log(-e^{-bx})) & \text{for } a = \log(-e^{-bx}) \\ x \coth^2(bx + \log(e^{-bx})) & \text{for } a = \log(e^{-bx}) \\ x - \frac{1}{b \tanh(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(coth(b*x+a)**2,x)`

output `Piecewise((x*coth(a)**2, Eq(b, 0)), (x*coth(b*x + log(-exp(-b*x)))**2, Eq(a, log(-exp(-b*x)))), (x*coth(b*x + log(exp(-b*x)))**2, Eq(a, log(exp(-b*x))))), (x - 1/(b*tanh(a + b*x)), True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.92

$$\int \coth^2(a + bx) dx = x + \frac{a}{b} + \frac{2}{b(e^{(-2bx-2a)} - 1)}$$

input `integrate(coth(b*x+a)^2,x, algorithm="maxima")`output `x + a/b + 2/(b*(e^(-2*b*x - 2*a) - 1))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.85

$$\int \coth^2(a + bx) dx = \frac{bx + a - \frac{2}{e^{(2bx+2a)} - 1}}{b}$$

input `integrate(coth(b*x+a)^2,x, algorithm="giac")`output `(b*x + a - 2/(e^(2*b*x + 2*a) - 1))/b`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \coth^2(a + bx) dx = x - \frac{\coth(a + bx)}{b}$$

input `int(coth(a + b*x)^2,x)`output `x - coth(a + b*x)/b`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.23

$$\int \coth^2(a + bx) dx = \frac{-\coth(bx + a) + bx}{b}$$

input `int(coth(b*x+a)^2,x)`

output `( - coth(a + b*x) + b*x)/b`

### 3.9 $\int \coth^3(a + bx) dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [C] (verified)	167
Maple [A] (verified)	168
Fricas [B] (verification not implemented)	169
Sympy [B] (verification not implemented)	169
Maxima [B] (verification not implemented)	170
Giac [A] (verification not implemented)	171
Mupad [B] (verification not implemented)	171
Reduce [B] (verification not implemented)	171

#### Optimal result

Integrand size = 8, antiderivative size = 27

$$\int \coth^3(a + bx) dx = -\frac{\coth^2(a + bx)}{2b} + \frac{\log(\sinh(a + bx))}{b}$$

output

```
-1/2*coth(b*x+a)^2/b+ln(sinh(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \coth^3(a + bx) dx = -\frac{\operatorname{csch}^2(a + bx) - 2 \log(\sinh(a + bx))}{2b}$$

input

```
Integrate[Coth[a + b*x]^3,x]
```

output

```
-1/2*(Csch[a + b*x]^2 - 2*Log[Sinh[a + b*x]])/b
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.48, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \tan\left(ia + ibx + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{26} \\
 & i \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & i\left(\frac{i \coth^2(a + bx)}{2b} - \int i \coth(a + bx) dx\right) \\
 & \quad \downarrow \text{26} \\
 & i\left(\frac{i \coth^2(a + bx)}{2b} - i \int \coth(a + bx) dx\right) \\
 & \quad \downarrow \text{3042} \\
 & i\left(\frac{i \coth^2(a + bx)}{2b} - i \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right) dx\right) \\
 & \quad \downarrow \text{26} \\
 & i\left(\frac{i \coth^2(a + bx)}{2b} - \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right) dx\right) \\
 & \quad \downarrow \text{3956} \\
 & i\left(\frac{i \coth^2(a + bx)}{2b} - \frac{i \log(-i \sinh(a + bx))}{b}\right)
 \end{aligned}$$



input `Int[Coth[a + b*x]^3,x]`

output `I*(((I/2)*Coth[a + b*x]^2)/b - (I*Log[(-I)*Sinh[a + b*x]])/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

### Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

method	result	size
derivativedivides	$\frac{-\frac{\coth(bx+a)^2}{2} - \frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	38
default	$\frac{-\frac{\coth(bx+a)^2}{2} - \frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	38
parallelrisch	$\frac{-2bx+2\ln(\tanh(bx+a))-2\ln(1-\tanh(bx+a))-\coth(bx+a)^2}{2b}$	43
risch	$-x - \frac{2a}{b} - \frac{2e^{2bx+2a}}{b(e^{2bx+2a}-1)^2} + \frac{\ln(e^{2bx+2a}-1)}{b}$	54

input `int(coth(b*x+a)^3,x,method=_RETURNVERBOSE)`

output  $1/b*(-1/2*\coth(b*x+a)^2-1/2*\ln(\coth(b*x+a)-1)-1/2*\ln(\coth(b*x+a)+1))$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 346 vs.  $2(25) = 50$ .

Time = 0.10 (sec) , antiderivative size = 346, normalized size of antiderivative = 12.81

$$\int \coth^3(a + bx) dx = \frac{bx \cosh(bx + a)^4 + 4bx \cosh(bx + a) \sinh(bx + a)^3 + bx \sinh(bx + a)^4 - 2(bx - 1) \cosh(bx + a)^2 + \dots}{\dots}$$

input `integrate(coth(b*x+a)^3,x, algorithm="fricas")`

output  $-(b*x*\cosh(b*x + a)^4 + 4*b*x*\cosh(b*x + a)*\sinh(b*x + a)^3 + b*x*\sinh(b*x + a)^4 - 2*(b*x - 1)*\cosh(b*x + a)^2 + 2*(3*b*x*\cosh(b*x + a)^2 - b*x + 1)*\sinh(b*x + a)^2 + b*x - (\cosh(b*x + a)^4 + 4*\cosh(b*x + a)*\sinh(b*x + a)^3 + \sinh(b*x + a)^4 + 2*(3*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^2 - 2*\cosh(b*x + a)^2 + 4*(\cosh(b*x + a)^3 - \cosh(b*x + a))*\sinh(b*x + a) + 1)*\log(2*\sinh(b*x + a)/(\cosh(b*x + a) - \sinh(b*x + a))) + 4*(b*x*\cosh(b*x + a)^3 - (b*x - 1)*\cosh(b*x + a))*\sinh(b*x + a))/(b*\cosh(b*x + a)^4 + 4*b*\cosh(b*x + a)*\sinh(b*x + a)^3 + b*\sinh(b*x + a)^4 - 2*b*\cosh(b*x + a)^2 + 2*(3*b*\cosh(b*x + a)^2 - b)*\sinh(b*x + a)^2 + 4*(b*\cosh(b*x + a)^3 - b*\cosh(b*x + a))*\sinh(b*x + a) + b)$

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs.  $2(20) = 40$ .

Time = 0.70 (sec) , antiderivative size = 90, normalized size of antiderivative = 3.33

$$\int \coth^3(a + bx) dx = \begin{cases} x \coth^3(a) & \text{for } b = 0 \\ x \coth^3(bx + \log(-e^{-bx})) & \text{for } a = \log(-e^{-bx}) \\ x \coth^3(bx + \log(e^{-bx})) & \text{for } a = \log(e^{-bx}) \\ x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} - \frac{1}{2b \tanh^2(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(coth(b*x+a)**3,x)`

output `Piecewise((x*coth(a)**3, Eq(b, 0)), (x*coth(b*x + log(-exp(-b*x)))**3, Eq(a, log(-exp(-b*x)))), (x*coth(b*x + log(exp(-b*x)))**3, Eq(a, log(exp(-b*x))))), (x - log(tanh(a + b*x) + 1)/b + log(tanh(a + b*x))/b - 1/(2*b*tanh(a + b*x)**2), True))`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs.  $2(25) = 50$ .

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.93

$$\int \coth^3(a + bx) dx = x + \frac{a}{b} + \frac{\log(e^{-bx-a} + 1)}{b} + \frac{\log(e^{-bx-a} - 1)}{b} + \frac{2e^{-2bx-2a}}{b(2e^{-2bx-2a} - e^{-4bx-4a} - 1)}$$

input `integrate(coth(b*x+a)^3,x, algorithm="maxima")`

output `x + a/b + log(e^(-b*x - a) + 1)/b + log(e^(-b*x - a) - 1)/b + 2*e^(-2*b*x - 2*a)/(b*(2*e^(-2*b*x - 2*a) - e^(-4*b*x - 4*a) - 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.81

$$\int \coth^3(a + bx) dx = -\frac{bx + a + \frac{2e^{(2bx+2a)}}{(e^{(2bx+2a)}-1)^2} - \log(|e^{(2bx+2a)} - 1|)}{b}$$

input `integrate(coth(b*x+a)^3,x, algorithm="giac")`output `-(b*x + a + 2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) - 1)^2 - log(abs(e^(2*b*x + 2*a) - 1)))/b`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.52

$$\int \coth^3(a + bx) dx = \frac{\ln(e^{2a} e^{2bx} - 1)}{b} - x - \frac{2}{b(e^{2a+2bx} - 1)} - \frac{2}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)}$$

input `int(coth(a + b*x)^3,x)`output `log(exp(2*a)*exp(2*b*x) - 1)/b - x - 2/(b*(exp(2*a + 2*b*x) - 1)) - 2/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 182, normalized size of antiderivative = 6.74

$$\int \coth^3(a + bx) dx = \frac{e^{4bx+4a} \log(e^{bx+a} - 1) + e^{4bx+4a} \log(e^{bx+a} + 1) - e^{4bx+4a} bx - e^{4bx+4a} - 2e^{2bx+2a} \log(e^{bx+a} - 1) - 2e^{2bx+2a} \log(e^{bx+a} + 1)}{b(e^{4bx+4a} - 2e^{2bx+2a} + 1)}$$

input `int(coth(b*x+a)^3,x)`

output 
$$\frac{(e^{4a+4bx})\log(e^{a+bx}-1) + e^{4a+4bx}\log(e^{a+bx}+1) - e^{4a+4bx}bx - e^{4a+4bx} - 2e^{2a+2bx}\log(e^{a+bx}-1) - 2e^{2a+2bx}\log(e^{a+bx}+1) + 2e^{2a+2bx}bx + \log(e^{a+bx}-1) + \log(e^{a+bx}+1) - bx - 1}{(e^{4a+4bx} - 2e^{2a+2bx} + 1)}$$

### 3.10 $\int \coth^4(a + bx) dx$

Optimal result . . . . .	173
Mathematica [C] (verified) . . . . .	173
Rubi [A] (verified) . . . . .	174
Maple [A] (verified) . . . . .	175
Fricas [B] (verification not implemented) . . . . .	176
Sympy [B] (verification not implemented) . . . . .	176
Maxima [B] (verification not implemented) . . . . .	177
Giac [A] (verification not implemented) . . . . .	177
Mupad [B] (verification not implemented) . . . . .	177
Reduce [B] (verification not implemented) . . . . .	178

#### Optimal result

Integrand size = 8, antiderivative size = 28

$$\int \coth^4(a + bx) dx = x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b}$$

output

```
x-coth(b*x+a)/b-1/3*coth(b*x+a)^3/b
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

$$\int \coth^4(a + bx) dx = -\frac{\coth^3(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \tanh^2(a + bx)\right)}{3b}$$

input

```
Integrate[Coth[a + b*x]^4,x]
```

output

```
-1/3*(Coth[a + b*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, Tanh[a + b*x]^2])/b
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^4(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(ia + ibx + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int -\coth^2(a + bx) dx - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25} \\
 & \int \coth^2(a + bx) dx - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\coth^3(a + bx)}{3b} + \int -\tan\left(ia + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\frac{\coth^3(a + bx)}{3b} - \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & -\frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} + x
 \end{aligned}$$

input

Int[Coth[a + b\*x]^4, x]

output  $x - \operatorname{Coth}[a + b*x]/b - \operatorname{Coth}[a + b*x]^3/(3*b)$

### Defintions of rubi rules used

rule 24  $\operatorname{Int}[a_, x\_Symbol] \rightarrow \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

rule 25  $\operatorname{Int}[-(F_x_), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 3042  $\operatorname{Int}[u_, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\operatorname{Int}[(b*.)\tan[(c_.) + (d_.)*(x_.)]^{(n_)}, x\_Symbol] \rightarrow \operatorname{Simp}[b*((b*\operatorname{Tan}[c + d*x])^{(n-1)}/(d*(n-1))), x] - \operatorname{Simp}[b^2 \operatorname{Int}[(b*\operatorname{Tan}[c + d*x])^{(n-2)}, x], x] /; \operatorname{FreeQ}[\{b, c, d\}, x] \ \&\& \operatorname{GtQ}[n, 1]$

### Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

method	result	size
parallelrisc	$\frac{-\operatorname{coth}(bx+a)^3 + 3bx - 3\operatorname{coth}(bx+a)}{3b}$	29
risc	$x - \frac{4(3e^{4bx+4a} - 3e^{2bx+2a} + 2)}{3b(e^{2bx+2a} - 1)^3}$	45
derivativedivides	$\frac{-\frac{\operatorname{coth}(bx+a)^3}{3} - \operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	46
default	$\frac{-\frac{\operatorname{coth}(bx+a)^3}{3} - \operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	46

input  $\operatorname{int}(\operatorname{coth}(b*x+a)^4, x, \operatorname{method}=\_RETURNVERBOSE)$

output  $1/3*(-\operatorname{coth}(b*x+a)^3 + 3*b*x - 3*\operatorname{coth}(b*x+a))/b$



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 108 vs.  $2(26) = 52$ .

Time = 0.09 (sec) , antiderivative size = 108, normalized size of antiderivative = 3.86

$$\int \coth^4(a + bx) dx = \frac{(3bx + 4) \sinh(bx + a)^3 - 4 \cosh(bx + a)^3 - 12 \cosh(bx + a) \sinh(bx + a)^2 + 3((3bx + 4) \cosh(bx + a) \sinh(bx + a)^2 + 3(b \sinh(bx + a)^3 + 3(b \cosh(bx + a)^2 - b) \sinh(bx + a))}{3(b \sinh(bx + a)^3 + 3(b \cosh(bx + a)^2 - b) \sinh(bx + a))}$$

input `integrate(coth(b*x+a)^4,x, algorithm="fricas")`

output `1/3*((3*b*x + 4)*sinh(b*x + a)^3 - 4*cosh(b*x + a)^3 - 12*cosh(b*x + a)*sinh(b*x + a)^2 + 3*((3*b*x + 4)*cosh(b*x + a)^2 - 3*b*x - 4)*sinh(b*x + a))/(b*sinh(b*x + a)^3 + 3*(b*cosh(b*x + a)^2 - b)*sinh(b*x + a))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 78 vs.  $2(20) = 40$ .

Time = 1.09 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.79

$$\int \coth^4(a + bx) dx = \begin{cases} x \coth^4(a) & \text{for } b = 0 \\ x \coth^4(bx + \log(-e^{-bx})) & \text{for } a = \log(-e^{-bx}) \\ x \coth^4(bx + \log(e^{-bx})) & \text{for } a = \log(e^{-bx}) \\ x - \frac{1}{b \tanh(a+bx)} - \frac{1}{3b \tanh^3(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(coth(b*x+a)**4,x)`

output `Piecewise((x*coth(a)**4, Eq(b, 0)), (x*coth(b*x + log(-exp(-b*x)))**4, Eq(a, log(-exp(-b*x)))), (x*coth(b*x + log(exp(-b*x)))**4, Eq(a, log(exp(-b*x))))), (x - 1/(b*tanh(a + b*x)) - 1/(3*b*tanh(a + b*x)**3), True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(26) = 52$ .

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.54

$$\int \coth^4(a + bx) dx = x + \frac{a}{b} - \frac{4(3e^{(-2bx-2a)} - 3e^{(-4bx-4a)} - 2)}{3b(3e^{(-2bx-2a)} - 3e^{(-4bx-4a)} + e^{(-6bx-6a)} - 1)}$$

input `integrate(coth(b*x+a)^4,x, algorithm="maxima")`

output `x + a/b - 4/3*(3*e^(-2*b*x - 2*a) - 3*e^(-4*b*x - 4*a) - 2)/(b*(3*e^(-2*b*x - 2*a) - 3*e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a) - 1))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.86

$$\int \coth^4(a + bx) dx = \frac{3bx + 3a - \frac{4(3e^{(4bx+4a)} - 3e^{(2bx+2a)} + 2)}{(e^{(2bx+2a)} - 1)^3}}{3b}$$

input `integrate(coth(b*x+a)^4,x, algorithm="giac")`

output `1/3*(3*b*x + 3*a - 4*(3*e^(4*b*x + 4*a) - 3*e^(2*b*x + 2*a) + 2)/(e^(2*b*x + 2*a) - 1)^3)/b`

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \coth^4(a + bx) dx = x - \frac{\frac{\coth(a+bx)^3}{3} + \coth(a + bx)}{b}$$

input `int(coth(a + b*x)^4,x)`

output  $x - (\coth(a + b*x) + \coth(a + b*x)^3/3)/b$

### Reduce [B] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \coth^4(a + bx) dx = \frac{-\coth(bx + a)^3 - 3\coth(bx + a) + 3bx}{3b}$$

input `int(coth(b*x+a)^4,x)`

output `( - coth(a + b*x)**3 - 3*coth(a + b*x) + 3*b*x)/(3*b)`

### 3.11 $\int \coth^5(a + bx) dx$

Optimal result	179
Mathematica [A] (verified)	179
Rubi [C] (verified)	180
Maple [A] (verified)	182
Fricas [B] (verification not implemented)	182
Sympy [B] (verification not implemented)	183
Maxima [B] (verification not implemented)	184
Giac [A] (verification not implemented)	184
Mupad [B] (verification not implemented)	185
Reduce [B] (verification not implemented)	185

#### Optimal result

Integrand size = 8, antiderivative size = 42

$$\int \coth^5(a + bx) dx = -\frac{\coth^2(a + bx)}{2b} - \frac{\coth^4(a + bx)}{4b} + \frac{\log(\sinh(a + bx))}{b}$$

output

```
-1/2*coth(b*x+a)^2/b-1/4*coth(b*x+a)^4/b+ln(sinh(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.83

$$\int \coth^5(a + bx) dx = -\frac{4\operatorname{csch}^2(a + bx) + \operatorname{csch}^4(a + bx) - 4\log(\sinh(a + bx))}{4b}$$

input

```
Integrate[Coth[a + b*x]^5,x]
```

output

```
-1/4*(4*Csch[a + b*x]^2 + Csch[a + b*x]^4 - 4*Log[Sinh[a + b*x]])/b
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.36, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {3042, 26, 3954, 26, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^5(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^5 dx \\
 & \quad \downarrow \text{3954} \\
 & -i\left(-\int -i \coth^3(a + bx) dx - \frac{i \coth^4(a + bx)}{4b}\right) \\
 & \quad \downarrow \text{26} \\
 & -i\left(i \int \coth^3(a + bx) dx - \frac{i \coth^4(a + bx)}{4b}\right) \\
 & \quad \downarrow \text{3042} \\
 & -i\left(i \int i \tan\left(ia + ibx + \frac{\pi}{2}\right)^3 dx - \frac{i \coth^4(a + bx)}{4b}\right) \\
 & \quad \downarrow \text{26} \\
 & -i\left(-\int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^3 dx - \frac{i \coth^4(a + bx)}{4b}\right) \\
 & \quad \downarrow \text{3954} \\
 & -i\left(\int i \coth(a + bx) dx - \frac{i \coth^4(a + bx)}{4b} - \frac{i \coth^2(a + bx)}{2b}\right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& -i \left( i \int \coth(a + bx) dx - \frac{i \coth^4(a + bx)}{4b} - \frac{i \coth^2(a + bx)}{2b} \right) \\
& \downarrow 3042 \\
& -i \left( i \int -i \tan \left( ia + ibx + \frac{\pi}{2} \right) dx - \frac{i \coth^4(a + bx)}{4b} - \frac{i \coth^2(a + bx)}{2b} \right) \\
& \downarrow 26 \\
& -i \left( \int \tan \left( \frac{1}{2}(2ia + \pi) + ibx \right) dx - \frac{i \coth^4(a + bx)}{4b} - \frac{i \coth^2(a + bx)}{2b} \right) \\
& \downarrow 3956 \\
& -i \left( -\frac{i \coth^4(a + bx)}{4b} - \frac{i \coth^2(a + bx)}{2b} + \frac{i \log(-i \sinh(a + bx))}{b} \right)
\end{aligned}$$

input `Int[Coth[a + b*x]^5,x]`

output `(-I)*((( -1/2*I)*Coth[a + b*x]^2)/b - ((I/4)*Coth[a + b*x]^4)/b + (I*Log[(-I)*Sinh[a + b*x]])/b)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$\frac{-\frac{\operatorname{coth}(bx+a)^4}{4} - \frac{\operatorname{coth}(bx+a)^2}{2} - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} - \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	48
default	$\frac{-\frac{\operatorname{coth}(bx+a)^4}{4} - \frac{\operatorname{coth}(bx+a)^2}{2} - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} - \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	48
parallelrisc	$\frac{-\operatorname{coth}(bx+a)^4 - 2\operatorname{coth}(bx+a)^2 - 4bx + 4\ln(\tanh(bx+a)) - 4\ln(1 - \tanh(bx+a))}{4b}$	53
risc	$-x - \frac{2a}{b} - \frac{4e^{2bx+2a}(e^{4bx+4a} - e^{2bx+2a} + 1)}{b(e^{2bx+2a} - 1)^4} + \frac{\ln(e^{2bx+2a} - 1)}{b}$	76

input

```
int(coth(b*x+a)^5,x,method=_RETURNVERBOSE)
```

output

```
1/b*(-1/4*coth(b*x+a)^4-1/2*coth(b*x+a)^2-1/2*ln(coth(b*x+a)-1)-1/2*ln(cot
h(b*x+a)+1))
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 978 vs.  $2(38) = 76$ .

Time = 0.10 (sec) , antiderivative size = 978, normalized size of antiderivative = 23.29

$$\int \operatorname{coth}^5(a + bx) dx = \text{Too large to display}$$

input

```
integrate(coth(b*x+a)^5,x, algorithm="fricas")
```

output

```

-(b*x*cosh(b*x + a)^8 + 8*b*x*cosh(b*x + a)*sinh(b*x + a)^7 + b*x*sinh(b*x
+ a)^8 - 4*(b*x - 1)*cosh(b*x + a)^6 + 4*(7*b*x*cosh(b*x + a)^2 - b*x + 1
)*sinh(b*x + a)^6 + 8*(7*b*x*cosh(b*x + a)^3 - 3*(b*x - 1)*cosh(b*x + a))*
sinh(b*x + a)^5 + 2*(3*b*x - 2)*cosh(b*x + a)^4 + 2*(35*b*x*cosh(b*x + a)^
4 - 30*(b*x - 1)*cosh(b*x + a)^2 + 3*b*x - 2)*sinh(b*x + a)^4 + 8*(7*b*x*c
osh(b*x + a)^5 - 10*(b*x - 1)*cosh(b*x + a)^3 + (3*b*x - 2)*cosh(b*x + a))
*sinh(b*x + a)^3 - 4*(b*x - 1)*cosh(b*x + a)^2 + 4*(7*b*x*cosh(b*x + a)^6
- 15*(b*x - 1)*cosh(b*x + a)^4 + 3*(3*b*x - 2)*cosh(b*x + a)^2 - b*x + 1)*
sinh(b*x + a)^2 + b*x - (cosh(b*x + a)^8 + 8*cosh(b*x + a)*sinh(b*x + a)^7
+ sinh(b*x + a)^8 + 4*(7*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^6 - 4*cosh(b*
x + a)^6 + 8*(7*cosh(b*x + a)^3 - 3*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(35
*cosh(b*x + a)^4 - 30*cosh(b*x + a)^2 + 3)*sinh(b*x + a)^4 + 6*cosh(b*x +
a)^4 + 8*(7*cosh(b*x + a)^5 - 10*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b
*x + a)^3 + 4*(7*cosh(b*x + a)^6 - 15*cosh(b*x + a)^4 + 9*cosh(b*x + a)^2
- 1)*sinh(b*x + a)^2 - 4*cosh(b*x + a)^2 + 8*(cosh(b*x + a)^7 - 3*cosh(b*x
+ a)^5 + 3*cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*log(2*sinh
(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))) + 8*(b*x*cosh(b*x + a)^7 - 3*(b
*x - 1)*cosh(b*x + a)^5 + (3*b*x - 2)*cosh(b*x + a)^3 - (b*x - 1)*cosh(b*x
+ a))*sinh(b*x + a))/(b*cosh(b*x + a)^8 + 8*b*cosh(b*x + a)*sinh(b*x + a)
^7 + b*sinh(b*x + a)^8 - 4*b*cosh(b*x + a)^6 + 4*(7*b*cosh(b*x + a)^2 - ...

```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs.  $2(32) = 64$ .

Time = 1.61 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.48

$$\int \coth^5(a + bx) dx = \begin{cases} x \coth^5(a) & \text{for } b = 0 \\ x \coth^5(bx + \log(-e^{-bx})) & \text{for } a = \log(-e^{-bx}) \\ x \coth^5(bx + \log(e^{-bx})) & \text{for } a = \log(e^{-bx}) \\ x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} - \frac{1}{2b \tanh^2(a+bx)} - \frac{1}{4b \tanh^4(a+bx)} & \text{otherwise} \end{cases}$$

input

```
integrate(coth(b*x+a)**5,x)
```



output

```
Piecewise((x*coth(a)**5, Eq(b, 0)), (x*coth(b*x + log(-exp(-b*x)))**5, Eq(a, log(-exp(-b*x)))), (x*coth(b*x + log(exp(-b*x)))**5, Eq(a, log(exp(-b*x))))), (x - log(tanh(a + b*x) + 1)/b + log(tanh(a + b*x))/b - 1/(2*b*tanh(a + b*x)**2) - 1/(4*b*tanh(a + b*x)**4), True))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 122 vs.  $2(38) = 76$ .

Time = 0.04 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.90

$$\int \coth^5(a + bx) dx = x + \frac{a}{b} + \frac{\log(e^{-bx-a} + 1)}{b} + \frac{\log(e^{-bx-a} - 1)}{b} + \frac{4(e^{-2bx-2a} - e^{-4bx-4a} + e^{-6bx-6a})}{b(4e^{-2bx-2a} - 6e^{-4bx-4a} + 4e^{-6bx-6a} - e^{-8bx-8a} - 1)}$$

input

```
integrate(coth(b*x+a)^5,x, algorithm="maxima")
```

output

```
x + a/b + log(e^(-b*x - a) + 1)/b + log(e^(-b*x - a) - 1)/b + 4*(e^(-2*b*x - 2*a) - e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a))/(b*(4*e^(-2*b*x - 2*a) - 6*e^(-4*b*x - 4*a) + 4*e^(-6*b*x - 6*a) - e^(-8*b*x - 8*a) - 1))
```

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.67

$$\int \coth^5(a + bx) dx = -\frac{bx + a + \frac{4(e^{6bx+6a} - e^{4bx+4a} + e^{2bx+2a})}{(e^{2bx+2a} - 1)^4} - \log(|e^{2bx+2a} - 1|)}{b}$$

input

```
integrate(coth(b*x+a)^5,x, algorithm="giac")
```

output

```
-(b*x + a + 4*(e^(6*b*x + 6*a) - e^(4*b*x + 4*a) + e^(2*b*x + 2*a)))/(e^(2*b*x + 2*a) - 1)^4 - log(abs(e^(2*b*x + 2*a) - 1))/b
```

**Mupad [B] (verification not implemented)**

Time = 2.09 (sec) , antiderivative size = 159, normalized size of antiderivative = 3.79

$$\int \coth^5(a + bx) dx = \frac{\ln(e^{2a} e^{2bx} - 1)}{b} - x - \frac{4}{b(e^{2a+2bx} - 1)} - \frac{8}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{8}{b(3e^{2a+2bx} - 3e^{4a+4bx} + e^{6a+6bx} - 1)} - \frac{4}{b(6e^{4a+4bx} - 4e^{2a+2bx} - 4e^{6a+6bx} + e^{8a+8bx} + 1)}$$

input `int(coth(a + b*x)^5,x)`output `log(exp(2*a)*exp(2*b*x) - 1)/b - x - 4/(b*(exp(2*a + 2*b*x) - 1)) - 8/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - 8/(b*(3*exp(2*a + 2*b*x) - 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) - 1)) - 4/(b*(6*exp(4*a + 4*b*x) - 4*exp(2*a + 2*b*x) - 4*exp(6*a + 6*b*x) + exp(8*a + 8*b*x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 334, normalized size of antiderivative = 7.95

$$\int \coth^5(a + bx) dx = \frac{e^{8bx+8a} \log(e^{bx+a} - 1) + e^{8bx+8a} \log(e^{bx+a} + 1) - e^{8bx+8a} bx - e^{8bx+8a} - 4e^{6bx+6a} \log(e^{bx+a} - 1) - 4e^{6bx+6a} \log(e^{bx+a} + 1)}{1}$$

input `int(coth(b*x+a)^5,x)`

output

```
(e**(8*a + 8*b*x)*log(e**(a + b*x) - 1) + e**(8*a + 8*b*x)*log(e**(a + b*x) + 1) - e**(8*a + 8*b*x)*b*x - e**(8*a + 8*b*x) - 4*e**(6*a + 6*b*x)*log(e**(a + b*x) - 1) - 4*e**(6*a + 6*b*x)*log(e**(a + b*x) + 1) + 4*e**(6*a + 6*b*x)*b*x + 6*e**(4*a + 4*b*x)*log(e**(a + b*x) - 1) + 6*e**(4*a + 4*b*x)*log(e**(a + b*x) + 1) - 6*e**(4*a + 4*b*x)*b*x - 2*e**(4*a + 4*b*x) - 4*e**(2*a + 2*b*x)*log(e**(a + b*x) - 1) - 4*e**(2*a + 2*b*x)*log(e**(a + b*x) + 1) + 4*e**(2*a + 2*b*x)*b*x + log(e**(a + b*x) - 1) + log(e**(a + b*x) + 1) - b*x - 1)/(b*(e**(8*a + 8*b*x) - 4*e**(6*a + 6*b*x) + 6*e**(4*a + 4*b*x) - 4*e**(2*a + 2*b*x) + 1))
```

## 3.12 $\int \coth^6(a + bx) dx$

Optimal result	187
Mathematica [C] (verified)	187
Rubi [A] (verified)	188
Maple [A] (verified)	190
Fricas [B] (verification not implemented)	190
Sympy [B] (verification not implemented)	191
Maxima [B] (verification not implemented)	191
Giac [A] (verification not implemented)	192
Mupad [B] (verification not implemented)	192
Reduce [B] (verification not implemented)	193

### Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \coth^6(a + bx) dx = x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth^5(a + bx)}{5b}$$

output

```
x-coth(b*x+a)/b-1/3*coth(b*x+a)^3/b-1/5*coth(b*x+a)^5/b
```

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

$$\int \coth^6(a + bx) dx = -\frac{\coth^5(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, 1, -\frac{3}{2}, \tanh^2(a + bx)\right)}{5b}$$

input

```
Integrate[Coth[a + b*x]^6,x]
```

output

```
-1/5*(Coth[a + b*x]^5*Hypergeometric2F1[-5/2, 1, -3/2, Tanh[a + b*x]^2])/b
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$ , Rules used = {3042, 25, 3954, 3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth^6(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(ia + ibx + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^6 dx \\
 & \quad \downarrow \text{3954} \\
 & \int \coth^4(a + bx) dx - \frac{\coth^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\coth^5(a + bx)}{5b} + \int \tan\left(ia + ibx + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & -\int -\coth^2(a + bx) dx - \frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25} \\
 & \int \coth^2(a + bx) dx - \frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(ia + ibx + \frac{\pi}{2}\right)^2 dx - \frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
 & - \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right)^2 dx - \frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & -\frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} + x
 \end{aligned}$$

input `Int[Coth[a + b*x]^6,x]`

output `x - Coth[a + b*x]/b - Coth[a + b*x]^3/(3*b) - Coth[a + b*x]^5/(5*b)`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

method	result	size
parallelrisch	$\frac{-3 \operatorname{coth}(bx+a)^5 - 5 \operatorname{coth}(bx+a)^3 + 15bx - 15 \operatorname{coth}(bx+a)}{15b}$	39
derivativedivides	$\frac{-\frac{\operatorname{coth}(bx+a)^5}{5} - \frac{\operatorname{coth}(bx+a)^3}{3} - \operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	56
default	$\frac{-\frac{\operatorname{coth}(bx+a)^5}{5} - \frac{\operatorname{coth}(bx+a)^3}{3} - \operatorname{coth}(bx+a) - \frac{\ln(\operatorname{coth}(bx+a)-1)}{2} + \frac{\ln(\operatorname{coth}(bx+a)+1)}{2}}{b}$	56
risch	$x - \frac{2(45 e^{8bx+8a} - 90 e^{6bx+6a} + 140 e^{4bx+4a} - 70 e^{2bx+2a} + 23)}{15b(e^{2bx+2a}-1)^5}$	67

input `int(coth(b*x+a)^6,x,method=_RETURNVERBOSE)`output `1/15*(-3*coth(b*x+a)^5-5*coth(b*x+a)^3+15*b*x-15*coth(b*x+a))/b`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(39) = 78.

Time = 0.09 (sec) , antiderivative size = 239, normalized size of antiderivative = 5.56

$$\int \operatorname{coth}^6(a+bx) dx$$

$$= \frac{(15bx+23)\sinh(bx+a)^5 - 23\cosh(bx+a)^5 - 115\cosh(bx+a)\sinh(bx+a)^4 + 5(2(15bx+23)\cosh(bx+a)\sinh(bx+a)^3 - 23\cosh(bx+a)^3 - 115\cosh(bx+a)\sinh(bx+a)^2 + 5(2(15bx+23)\cosh(bx+a)\sinh(bx+a) - 23\cosh(bx+a))\sinh(bx+a) - 50\cosh(bx+a)^2 + 30b\sinh(bx+a) - 50\cosh(bx+a))}{15b(e^{2bx+2a}-1)^5}$$

input `integrate(coth(b*x+a)^6,x, algorithm="fricas")`output `1/15*((15*b*x + 23)*sinh(b*x + a)^5 - 23*cosh(b*x + a)^5 - 115*cosh(b*x + a)*sinh(b*x + a)^4 + 5*(2*(15*b*x + 23)*cosh(b*x + a)^2 - 15*b*x - 23)*sinh(b*x + a)^3 + 25*cosh(b*x + a)^3 - 5*(46*cosh(b*x + a)^3 - 15*cosh(b*x + a))*sinh(b*x + a)^2 + 5*((15*b*x + 23)*cosh(b*x + a)^4 - 3*(15*b*x + 23)*cosh(b*x + a)^2 + 30*b*x + 46)*sinh(b*x + a) - 50*cosh(b*x + a))/(b*sinh(b*x + a)^5 + 5*(2*b*cosh(b*x + a)^2 - b)*sinh(b*x + a)^3 + 5*(b*cosh(b*x + a)^4 - 3*b*cosh(b*x + a)^2 + 2*b)*sinh(b*x + a))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 92 vs.  $2(32) = 64$ .

Time = 2.45 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.14

$$\int \coth^6(a + bx) dx$$

$$= \begin{cases} x \coth^6(a) & \text{for } b = 0 \\ x \coth^6(bx + \log(-e^{-bx})) & \text{for } a = \log(-e^{-bx}) \\ x \coth^6(bx + \log(e^{-bx})) & \text{for } a = \log(e^{-bx}) \\ x - \frac{1}{b \tanh(a+bx)} - \frac{1}{3b \tanh^3(a+bx)} - \frac{1}{5b \tanh^5(a+bx)} & \text{otherwise} \end{cases}$$

input

```
integrate(coth(b*x+a)**6,x)
```

output

```
Piecewise((x*coth(a)**6, Eq(b, 0)), (x*coth(b*x + log(-exp(-b*x)))**6, Eq(a, log(-exp(-b*x)))), (x*coth(b*x + log(exp(-b*x)))**6, Eq(a, log(exp(-b*x)))), (x - 1/(b*tanh(a + b*x)) - 1/(3*b*tanh(a + b*x)**3) - 1/(5*b*tanh(a + b*x)**5), True))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 115 vs.  $2(39) = 78$ .

Time = 0.04 (sec) , antiderivative size = 115, normalized size of antiderivative = 2.67

$$\int \coth^6(a + bx) dx$$

$$= x + \frac{a}{b} - \frac{2(70e^{(-2bx-2a)} - 140e^{(-4bx-4a)} + 90e^{(-6bx-6a)} - 45e^{(-8bx-8a)} - 23)}{15b(5e^{(-2bx-2a)} - 10e^{(-4bx-4a)} + 10e^{(-6bx-6a)} - 5e^{(-8bx-8a)} + e^{(-10bx-10a)} - 1)}$$

input

```
integrate(coth(b*x+a)^6,x, algorithm="maxima")
```



output

```
x + a/b - 2/15*(70*e^(-2*b*x - 2*a) - 140*e^(-4*b*x - 4*a) + 90*e^(-6*b*x - 6*a) - 45*e^(-8*b*x - 8*a) - 23)/(b*(5*e^(-2*b*x - 2*a) - 10*e^(-4*b*x - 4*a) + 10*e^(-6*b*x - 6*a) - 5*e^(-8*b*x - 8*a) + e^(-10*b*x - 10*a) - 1))
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.72

$$\int \coth^6(a + bx) dx = \frac{15bx + 15a - \frac{2(45e^{(8bx+8a)} - 90e^{(6bx+6a)} + 140e^{(4bx+4a)} - 70e^{(2bx+2a)} + 23)}{(e^{(2bx+2a)} - 1)^5}}{15b}$$

input

```
integrate(coth(b*x+a)^6,x, algorithm="giac")
```

output

```
1/15*(15*b*x + 15*a - 2*(45*e^(8*b*x + 8*a) - 90*e^(6*b*x + 6*a) + 140*e^(4*b*x + 4*a) - 70*e^(2*b*x + 2*a) + 23)/(e^(2*b*x + 2*a) - 1)^5)/b
```

**Mupad [B] (verification not implemented)**

Time = 2.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \coth^6(a + bx) dx = x - \frac{\coth(a+bx)^5}{5} + \frac{\coth(a+bx)^3}{3} + \coth(a + bx) \frac{1}{b}$$

input

```
int(coth(a + b*x)^6,x)
```

output

```
x - (coth(a + b*x) + coth(a + b*x)^3/3 + coth(a + b*x)^5/5)/b
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int \coth^6(a + bx) dx = \frac{-3 \coth^5(bx + a) - 5 \coth^3(bx + a) - 15 \coth(bx + a) + 15bx}{15b}$$

input `int(coth(b*x+a)^6,x)`

output `( - 3*coth(a + b*x)**5 - 5*coth(a + b*x)**3 - 15*coth(a + b*x) + 15*b*x)/(15*b)`

### 3.13 $\int (b \tanh(c + dx))^{7/2} dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (warning: unable to verify)	195
Maple [A] (verified)	198
Fricas [B] (verification not implemented)	198
Sympy [F]	199
Maxima [F]	200
Giac [F(-2)]	200
Mupad [B] (verification not implemented)	200
Reduce [F]	201

#### Optimal result

Integrand size = 12, antiderivative size = 97

$$\int (b \tanh(c + dx))^{7/2} dx = \frac{b^{7/2} \arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d}$$

output

```
b^(7/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(7/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d-2*b^3*(b*tanh(d*x+c))^(1/2)/d-2/5*b*(b*tanh(d*x+c))^(5/2)/d
```

#### Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

$$\int (b \tanh(c + dx))^{7/2} dx = \frac{(b \tanh(c + dx))^{7/2} \left( -\arctan\left(\sqrt{\tanh(c + dx)}\right) - \operatorname{arctanh}\left(\sqrt{\tanh(c + dx)}\right) + 2\sqrt{\tanh(c + dx)} + \frac{2}{5} \tanh(c + dx) \right)}{d \tanh^{7/2}(c + dx)}$$

input

```
Integrate[(b*Tanh[c + d*x])^(7/2),x]
```

output

```

-(((b*Tanh[c + d*x])^(7/2)*(-ArcTan[Sqrt[Tanh[c + d*x]]] - ArcTanh[Sqrt[Ta
nh[c + d*x]]] + 2*Sqrt[Tanh[c + d*x]] + (2*Tanh[c + d*x]^(5/2))/5))/(d*Tan
h[c + d*x]^(7/2)))

```

**Rubi [A] (warning: unable to verify)**

Time = 0.45 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3954, 3042, 3954, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (b \tanh(c + dx))^{7/2} dx \\
& \quad \downarrow \text{3042} \\
& \int (-ib \tan(ic + idx))^{7/2} dx \\
& \quad \downarrow \text{3954} \\
& b^2 \int (b \tanh(c + dx))^{3/2} dx - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& -\frac{2b(b \tanh(c + dx))^{5/2}}{5d} + b^2 \int (-ib \tan(ic + idx))^{3/2} dx \\
& \quad \downarrow \text{3954} \\
& b^2 \left( b^2 \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx - \frac{2b\sqrt{b \tanh(c + dx)}}{d} \right) - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& -\frac{2b(b \tanh(c + dx))^{5/2}}{5d} + b^2 \left( -\frac{2b\sqrt{b \tanh(c + dx)}}{d} + b^2 \int \frac{1}{\sqrt{-ib \tan(ic + idx)}} dx \right) \\
& \quad \downarrow \text{3957}
\end{aligned}$$

$$\begin{aligned}
 & b^2 \left( -\frac{b^3 \int -\frac{1}{\sqrt{b \tanh(c+dx)}(b^2-b^2 \tanh^2(c+dx))} d(b \tanh(c+dx))}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \\
 & \qquad \qquad \qquad \frac{2b(b \tanh(c+dx))^{5/2}}{5d} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & b^2 \left( \frac{b^3 \int \frac{1}{\sqrt{b \tanh(c+dx)}(b^2-b^2 \tanh^2(c+dx))} d(b \tanh(c+dx))}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \\
 & \qquad \qquad \qquad \frac{2b(b \tanh(c+dx))^{5/2}}{5d} \\
 & \qquad \qquad \qquad \downarrow \text{266} \\
 & b^2 \left( \frac{2b^3 \int \frac{1}{b^2-b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c+dx)}}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \frac{2b(b \tanh(c+dx))^{5/2}}{5d} \\
 & \qquad \qquad \qquad \downarrow \text{756} \\
 & b^2 \left( \frac{2b^3 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\int \frac{1}{b^2 \tanh^2(c+dx)+b} d\sqrt{b \tanh(c+dx)}}{2b} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \\
 & \qquad \qquad \qquad \frac{2b(b \tanh(c+dx))^{5/2}}{5d} \\
 & \qquad \qquad \qquad \downarrow \text{216} \\
 & b^2 \left( \frac{2b^3 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \\
 & \qquad \qquad \qquad \frac{2b(b \tanh(c+dx))^{5/2}}{5d} \\
 & \qquad \qquad \qquad \downarrow \text{219} \\
 & b^2 \left( \frac{2b^3 \left( \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} + \frac{\operatorname{arctanh}(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \right) - \\
 & \qquad \qquad \qquad \frac{2b(b \tanh(c+dx))^{5/2}}{5d}
 \end{aligned}$$

input `Int[(b*Tanh[c + d*x])^(7/2),x]`

output `(-2*b*(b*Tanh[c + d*x])^(5/2))/(5*d) + b^2*((2*b^3*(ArcTan[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2))) + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)))/d - (2*b*Sqrt[b*Tanh[c + d*x]]/d)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{b^{\frac{7}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b} \tanh(dx+c)}{d} - \frac{2b(b \tanh(dx+c))^{\frac{5}{2}}}{5d}$	80
default	$\frac{b^{\frac{7}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b} \tanh(dx+c)}{d} - \frac{2b(b \tanh(dx+c))^{\frac{5}{2}}}{5d}$	80

input `int((b*tanh(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output `b^(7/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(7/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d-2*b^3*(b*tanh(d*x+c))^(1/2)/d-2/5*b*(b*tanh(d*x+c))^(5/2)/d`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 742 vs.  $2(79) = 158$ .

Time = 0.13 (sec) , antiderivative size = 1545, normalized size of antiderivative = 15.93

$$\int (b \tanh(c + dx))^{7/2} dx = \text{Too large to display}$$

input `integrate((b*tanh(d*x+c))^(7/2),x, algorithm="fricas")`

output

```

[-1/20*(10*(b^3*cosh(d*x + c)^4 + 4*b^3*cosh(d*x + c)*sinh(d*x + c)^3 + b^3*sinh(d*x + c)^4 + 2*b^3*cosh(d*x + c)^2 + b^3 + 2*(3*b^3*cosh(d*x + c)^2 + b^3)*sinh(d*x + c)^2 + 4*(b^3*cosh(d*x + c)^3 + b^3*cosh(d*x + c))*sinh(d*x + c))*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) - 5*(b^3*cosh(d*x + c)^4 + 4*b^3*cosh(d*x + c)*sinh(d*x + c)^3 + b^3*sinh(d*x + c)^4 + 2*b^3*cosh(d*x + c)^2 + b^3 + 2*(3*b^3*cosh(d*x + c)^2 + b^3)*sinh(d*x + c)^2 + 4*(b^3*cosh(d*x + c)^3 + b^3*cosh(d*x + c))*sinh(d*x + c))*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 16*(3*b^3*cosh(d*x + c)^4 + 12*b^3*cosh(d*x + c)*sinh(d*x + c)^3 + 3*b^3*sinh(d*x + c)^4 + 4*b^3*cosh(d*x + c)^2 + 3*b^3 + 2*(9*b^3*cosh(d*x + c)^2 + 2*b^3)*sinh(d*x + c)^2 + 4*(3*b^3*cosh(d*x + c)^3 + 2*b^3*cosh(d*x + c))*sinh(d*x + c))*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(d*cosh(d*x + c)^4 + 4*d*cosh(d*x + c)*sinh(d*x + c)^3 + d*sinh(d*x + c)^4 + 2*d*cosh(d*x + c)^2 + 2*...

```

## Sympy [F]

$$\int (b \tanh(c + dx))^{7/2} dx = \int (b \tanh(c + dx))^{\frac{7}{2}} dx$$

input

```
integrate((b*tanh(d*x+c))**(7/2), x)
```

output

```
Integral((b*tanh(c + d*x))**(7/2), x)
```



**Maxima [F]**

$$\int (b \tanh(c + dx))^{7/2} dx = \int (b \tanh(dx + c))^{\frac{7}{2}} dx$$

input `integrate((b*tanh(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(7/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int (b \tanh(c + dx))^{7/2} dx = \text{Exception raised: TypeError}$$

input `integrate((b*tanh(d*x+c))^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.63 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.86

$$\int (b \tanh(c + dx))^{7/2} dx = \frac{b^{7/2} \operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b \tanh(c+dx)}}{d} - \frac{2b(b \tanh(c+dx))^{5/2}}{5d} - \frac{b^{7/2} \operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)} \operatorname{li}}{\sqrt{b}}\right) \operatorname{li}}{d}$$

input `int((b*tanh(c + d*x))^(7/2),x)`

output

```
(b^(7/2)*atan((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d - (2*b^3*(b*tanh(c + d*x))^(1/2))/d - (2*b*(b*tanh(c + d*x))^(5/2))/(5*d) - (b^(7/2)*atan(((b*tanh(c + d*x))^(1/2)*1i)/b^(1/2))*1i)/d
```

**Reduce [F]**

$$\int (b \tanh(c + dx))^{7/2} dx = \frac{\sqrt{b} b^3 \left( -2\sqrt{\tanh(dx+c)} \tanh(dx+c)^2 - 10\sqrt{\tanh(dx+c)} + 5 \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)} dx \right) d \right)}{5d}$$

input

```
int((b*tanh(d*x+c))^(7/2),x)
```

output

```
(sqrt(b)*b**3*( - 2*sqrt(tanh(c + d*x))*tanh(c + d*x)**2 - 10*sqrt(tanh(c + d*x)) + 5*int(sqrt(tanh(c + d*x))/tanh(c + d*x),x)*d))/(5*d)
```

### 3.14 $\int (b \tanh(c + dx))^{5/2} dx$

Optimal result	202
Mathematica [A] (verified)	202
Rubi [A] (warning: unable to verify)	203
Maple [A] (verified)	205
Fricas [B] (verification not implemented)	206
Sympy [F]	207
Maxima [F]	207
Giac [F(-2)]	207
Mupad [B] (verification not implemented)	208
Reduce [F]	208

#### Optimal result

Integrand size = 12, antiderivative size = 78

$$\int (b \tanh(c + dx))^{5/2} dx = -\frac{b^{5/2} \arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}$$

output

```
-b^(5/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(5/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d-2/3*b*(b*tanh(d*x+c))^(3/2)/d
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.85

$$\int (b \tanh(c + dx))^{5/2} dx = \frac{(b \tanh(c + dx))^{5/2} \left( \arctan\left(\sqrt{\tanh(c + dx)}\right) - \operatorname{arctanh}\left(\sqrt{\tanh(c + dx)}\right) + \frac{2}{3} \tanh^{\frac{3}{2}}(c + dx) \right)}{d \tanh^{\frac{5}{2}}(c + dx)}$$

input

```
Integrate[(b*Tanh[c + d*x])^(5/2),x]
```

output

$$-\left(\left(b \operatorname{Tanh}[c + d*x]\right)^{(5/2)} * \left(\operatorname{ArcTan}\left[\operatorname{Sqrt}\left[\operatorname{Tanh}[c + d*x]\right]\right]\right) - \operatorname{ArcTanh}\left[\operatorname{Sqrt}\left[\operatorname{Tanh}[c + d*x]\right]\right] + \left(2 * \operatorname{Tanh}[c + d*x]^{(3/2)}\right) / 3\right) / \left(d * \operatorname{Tanh}[c + d*x]^{(5/2)}\right)$$
**Rubi [A] (warning: unable to verify)**

Time = 0.36 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3954, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (b \tanh(c + dx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (-ib \tan(ic + idx))^{5/2} dx \\ & \quad \downarrow \text{3954} \\ & b^2 \int \sqrt{b \tanh(c + dx)} dx - \frac{2b(b \tanh(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{3042} \\ & -\frac{2b(b \tanh(c + dx))^{3/2}}{3d} + b^2 \int \sqrt{-ib \tan(ic + idx)} dx \\ & \quad \downarrow \text{3957} \\ & -\frac{b^3 \int -\frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{25} \\ & \frac{b^3 \int \frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{266} \\ & \frac{2b^3 \int \frac{b^2 \tanh^2(c+dx)}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d} \end{aligned}$$

$$\begin{array}{c}
 \downarrow 827 \\
 \frac{2b^3 \left( \frac{1}{2} \int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{1}{2} \int \frac{1}{b^2 \tanh^2(c+dx)+b} d\sqrt{b \tanh(c+dx)} \right)}{\frac{d}{2b(b \tanh(c+dx))^{3/2}} 3d} \\
 \downarrow 216 \\
 \frac{2b^3 \left( \frac{1}{2} \int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} \right)}{d} - \frac{2b(b \tanh(c+dx))^{3/2}}{3d} \\
 \downarrow 219 \\
 \frac{2b^3 \left( \frac{\operatorname{arctanh}(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} \right)}{d} - \frac{2b(b \tanh(c+dx))^{3/2}}{3d}
 \end{array}$$

input `Int[(b*Tanh[c + d*x])^(5/2),x]`

output `(2*b^3*(-1/2*ArcTan[Sqrt[b]*Tanh[c + d*x]]/Sqrt[b] + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*Sqrt[b]))/d - (2*b*(b*Tanh[c + d*x])^(3/2))/(3*d)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{b^{\frac{5}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(dx+c))^{\frac{3}{2}}}{3d}$	63
default	$-\frac{b^{\frac{5}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(dx+c))^{\frac{3}{2}}}{3d}$	63

input `int((b*tanh(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output

```
-b^(5/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(5/2)*arctanh((b*tanh(d
*x+c))^(1/2)/b^(1/2))/d-2/3*b*(b*tanh(d*x+c))^(3/2)/d
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 454 vs.  $2(62) = 124$ .

Time = 0.13 (sec) , antiderivative size = 969, normalized size of antiderivative = 12.42

$$\int (b \tanh(c + dx))^{5/2} dx = \text{Too large to display}$$

input

```
integrate((b*tanh(d*x+c))^(5/2),x, algorithm="fricas")
```

output

```
[-1/12*(6*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*s
inh(d*x + c)^2 + b^2)*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*s
inh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x
+ c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x +
c)^2)) - 3*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2
*sinh(d*x + c)^2 + b^2)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x +
c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x +
c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x +
c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cos
h(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*
cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d
*x + c)^4)) + 8*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) +
b^2*sinh(d*x + c)^2 - b^2)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(d*cosh(d
*x + c)^2 + 2*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x + c)^2 + d), -1/1
2*(6*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh(d
*x + c)^2 + b^2)*sqrt(b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*
x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/sqrt(b))
- 3*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh(d
*x + c)^2 + b^2)*sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*si
nh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*s...
```

**Sympy [F]**

$$\int (b \tanh(c + dx))^{5/2} dx = \int (b \tanh(c + dx))^{5/2} dx$$

input `integrate((b*tanh(d*x+c))**(5/2),x)`

output `Integral((b*tanh(c + d*x))**(5/2), x)`

**Maxima [F]**

$$\int (b \tanh(c + dx))^{5/2} dx = \int (b \tanh(dx + c))^{5/2} dx$$

input `integrate((b*tanh(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(5/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int (b \tanh(c + dx))^{5/2} dx = \text{Exception raised: TypeError}$$

input `integrate((b*tanh(d*x+c))^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`



**Mupad [B] (verification not implemented)**

Time = 2.35 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.79

$$\int (b \tanh(c + dx))^{5/2} dx = \frac{b^{5/2} \operatorname{atanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{b^{5/2} \operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}$$

input `int((b*tanh(c + d*x))^(5/2),x)`output `(b^(5/2)*atanh((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d - (b^(5/2)*atan((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d - (2*b*(b*tanh(c + d*x))^(3/2))/(3*d)`**Reduce [F]**

$$\int (b \tanh(c + dx))^{5/2} dx = \sqrt{b} \left( \int \sqrt{\tanh(dx + c)} \tanh(dx + c)^2 dx \right) b^2$$

input `int((b*tanh(d*x+c))^(5/2),x)`output `sqrt(b)*int(sqrt(tanh(c + d*x))*tanh(c + d*x)**2,x)*b**2`

### 3.15 $\int (b \tanh(c + dx))^{3/2} dx$

Optimal result	209
Mathematica [A] (verified)	209
Rubi [A] (warning: unable to verify)	210
Maple [A] (verified)	212
Fricas [B] (verification not implemented)	213
Sympy [F]	214
Maxima [F]	214
Giac [F(-2)]	214
Mupad [B] (verification not implemented)	215
Reduce [F]	215

#### Optimal result

Integrand size = 12, antiderivative size = 75

$$\int (b \tanh(c + dx))^{3/2} dx = \frac{b^{3/2} \arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

output

$b^{(3/2)}*\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d+b^{(3/2)}*\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d-2*b*(b*\tanh(d*x+c))^{(1/2)}/d$

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.88

$$\int (b \tanh(c + dx))^{3/2} dx = \frac{\left(-\arctan\left(\sqrt{\tanh(c + dx)}\right) - \operatorname{arctanh}\left(\sqrt{\tanh(c + dx)}\right) + 2\sqrt{\tanh(c + dx)}\right) (b \tanh(c + dx))^{3/2}}{d \tanh^{\frac{3}{2}}(c + dx)}$$

input

`Integrate[(b*Tanh[c + d*x])^(3/2),x]`

output

```
-(((ArcTan[Sqrt[Tanh[c + d*x]]] - ArcTanh[Sqrt[Tanh[c + d*x]]] + 2*Sqrt[Tanh[c + d*x]])*(b*Tanh[c + d*x])^(3/2))/(d*Tanh[c + d*x])^(3/2))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.35 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3954, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (b \tanh(c + dx))^{3/2} dx$$

$$\downarrow 3042$$

$$\int (-ib \tan(ic + idx))^{3/2} dx$$

$$\downarrow 3954$$

$$b^2 \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

$$\downarrow 3042$$

$$-\frac{2b\sqrt{b \tanh(c + dx)}}{d} + b^2 \int \frac{1}{\sqrt{-ib \tan(ic + idx)}} dx$$

$$\downarrow 3957$$

$$-\frac{b^3 \int \frac{1}{\sqrt{b \tanh(c + dx)}(b^2 - b^2 \tanh^2(c + dx))} d(b \tanh(c + dx))}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

$$\downarrow 25$$

$$\frac{b^3 \int \frac{1}{\sqrt{b \tanh(c + dx)}(b^2 - b^2 \tanh^2(c + dx))} d(b \tanh(c + dx))}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

$$\downarrow 266$$

$$\frac{2b^3 \int \frac{1}{b^2 - b^4 \tanh^4(c + dx)} d\sqrt{b \tanh(c + dx)}}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

$$\begin{array}{c}
 \downarrow 756 \\
 \frac{2b^3 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\int \frac{1}{b^2 \tanh^2(c+dx)+b} d\sqrt{b \tanh(c+dx)}}{2b} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \\
 \downarrow 216 \\
 \frac{2b^3 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d} \\
 \downarrow 219 \\
 \frac{2b^3 \left( \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} + \frac{\operatorname{arctanh}(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d} - \frac{2b\sqrt{b \tanh(c+dx)}}{d}
 \end{array}$$

input `Int[(b*Tanh[c + d*x])^(3/2),x]`

output `(2*b^3*(ArcTan[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)) + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)))/d - (2*b*Sqrt[b*Tanh[c + d*x]])/d`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{b^{\frac{3}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b} \tanh(dx+c)}{d}$	62
default	$\frac{b^{\frac{3}{2}} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{b^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b} \tanh(dx+c)}{d}$	62

input `int((b*tanh(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output

```
b^(3/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(3/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d-2*b*(b*tanh(d*x+c))^(1/2)/d
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 281 vs.  $2(61) = 122$ .

Time = 0.11 (sec) , antiderivative size = 626, normalized size of antiderivative = 8.35

$$\int (b \tanh(c + dx))^{3/2} dx = \text{Too large to display}$$

input

```
integrate((b*tanh(d*x+c))^(3/2),x, algorithm="fricas")
```

output

```
[-1/4*(2*sqrt(-b)*b*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) - sqrt(-b)*b*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*b*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/d, 1/4*(2*b^(3/2)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/sqrt(b)) + b^(3/2)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b) - 8*b*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/d]
```

**Sympy [F]**

$$\int (b \tanh(c + dx))^{3/2} dx = \int (b \tanh(c + dx))^{\frac{3}{2}} dx$$

input `integrate((b*tanh(d*x+c))**(3/2),x)`

output `Integral((b*tanh(c + d*x))**(3/2), x)`

**Maxima [F]**

$$\int (b \tanh(c + dx))^{3/2} dx = \int (b \tanh(dx + c))^{\frac{3}{2}} dx$$

input `integrate((b*tanh(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(3/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int (b \tanh(c + dx))^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((b*tanh(d*x+c))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.81

$$\int (b \tanh(c + dx))^{3/2} dx = \frac{b^{3/2} \operatorname{atan}\left(\frac{\sqrt{b} \tanh(c+dx)}{\sqrt{b}}\right)}{d} - \frac{2b \sqrt{b \tanh(c+dx)}}{d} + \frac{b^{3/2} \operatorname{atanh}\left(\frac{\sqrt{b} \tanh(c+dx)}{\sqrt{b}}\right)}{d}$$

input `int((b*tanh(c + d*x))^(3/2),x)`output `(b^(3/2)*atan((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d - (2*b*(b*tanh(c + d*x))^(1/2))/d + (b^(3/2)*atanh((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d`**Reduce [F]**

$$\int (b \tanh(c + dx))^{3/2} dx = \frac{\sqrt{b} b \left( -2 \sqrt{\tanh(dx + c)} + \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)} dx \right) d \right)}{d}$$

input `int((b*tanh(d*x+c))^(3/2),x)`output `(sqrt(b)*b*( - 2*sqrt(tanh(c + d*x)) + int(sqrt(tanh(c + d*x))/tanh(c + d*x),x)*d))/d`



### 3.16 $\int \sqrt{b \tanh(c + dx)} dx$

Optimal result	216
Mathematica [A] (verified)	216
Rubi [A] (warning: unable to verify)	217
Maple [A] (verified)	219
Fricas [B] (verification not implemented)	219
Sympy [F]	220
Maxima [F]	220
Giac [F(-2)]	221
Mupad [B] (verification not implemented)	221
Reduce [F]	222

#### Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \sqrt{b \tanh(c + dx)} dx = -\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d} + \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{d}$$

output

```
-b^(1/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(1/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int \sqrt{b \tanh(c + dx)} dx = -\frac{\left(\arctan\left(\sqrt{\tanh(c + dx)}\right) - \operatorname{arctanh}\left(\sqrt{\tanh(c + dx)}\right)\right) \sqrt{b \tanh(c + dx)}}{d \sqrt{\tanh(c + dx)}}$$

input

```
Integrate[Sqrt[b*Tanh[c + d*x]],x]
```

output

```

-(((ArcTan[Sqrt[Tanh[c + d*x]]] - ArcTanh[Sqrt[Tanh[c + d*x]]])*Sqrt[b*Tanh[c + d*x]])/(d*Sqrt[Tanh[c + d*x]]))

```

**Rubi [A] (warning: unable to verify)**

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.88, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \sqrt{b \tanh(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{-ib \tan(ic + idx)} dx \\
& \quad \downarrow \text{3957} \\
& \frac{b \int -\frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\
& \quad \downarrow \text{25} \\
& \frac{b \int \frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\
& \quad \downarrow \text{266} \\
& \frac{2b \int \frac{b^2 \tanh^2(c+dx)}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c + dx)}}{d} \\
& \quad \downarrow \text{827} \\
& \frac{2b \left( \frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c + dx)} - \frac{1}{2} \int \frac{1}{b^2 \tanh^2(c+dx) + b} d\sqrt{b \tanh(c + dx)} \right)}{d} \\
& \quad \downarrow \text{216} \\
& \frac{2b \left( \frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c + dx)} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} \right)}{d}
\end{aligned}$$

$$\frac{2b \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(c+dx)}{2\sqrt{b}}\right)}{2\sqrt{b}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{b} \tanh(c+dx)}{2\sqrt{b}}\right)}{2\sqrt{b}} \right)}{d}$$

input `Int[Sqrt[b*Tanh[c + d*x]],x]`

output `(2*b*(-1/2*ArcTan[Sqrt[b]*Tanh[c + d*x]]/Sqrt[b] + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*Sqrt[b])))/d`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d}$	47
default	$-\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d} + \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{d}$	47

input `int((b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `-b^(1/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d+b^(1/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 261 vs.  $2(46) = 92$ .

Time = 0.13 (sec) , antiderivative size = 583, normalized size of antiderivative = 10.05

$$\int \sqrt{b \tanh(c + dx)} dx = \text{Too large to display}$$

input `integrate((b*tanh(d*x+c))^(1/2),x, algorithm="fricas")`

output

```
[-1/4*(2*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c)
+ sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cos
h(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) - sqr
t(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cos
sh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh
(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*
x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*
x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x +
c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)))/d, -1/4*(2*sqr
t(b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x +
c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/sqrt(b)) - sqrt(b)*log(2*b*c
osh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*
sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4
+ 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 +
(6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x
+ c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(
d*x + c)) - b))/d]
```

**Sympy [F]**

$$\int \sqrt{b \tanh(c + dx)} dx = \int \sqrt{b \tanh(c + dx)} dx$$

input

```
integrate((b*tanh(d*x+c))**(1/2),x)
```

output

```
Integral(sqrt(b*tanh(c + d*x)), x)
```

**Maxima [F]**

$$\int \sqrt{b \tanh(c + dx)} dx = \int \sqrt{b \tanh(dx + c)} dx$$

input

```
integrate((b*tanh(d*x+c))^(1/2),x, algorithm="maxima")
```

output `integrate(sqrt(b*tanh(d*x + c)), x)`

### Giac [F(-2)]

Exception generated.

$$\int \sqrt{b \tanh(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((b*tanh(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

### Mupad [B] (verification not implemented)

Time = 2.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.71

$$\int \sqrt{b \tanh(c + dx)} dx = -\frac{\sqrt{b} \left( \operatorname{atan} \left( \frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}} \right) - \operatorname{atanh} \left( \frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}} \right) \right)}{d}$$

input `int((b*tanh(c + d*x))^(1/2),x)`

output `-(b^(1/2)*(atan((b*tanh(c + d*x))^(1/2)/b^(1/2)) - atanh((b*tanh(c + d*x))  
^(1/2)/b^(1/2))))/d`

**Reduce [F]**

$$\int \sqrt{b \tanh(c + dx)} dx = \sqrt{b} \left( \int \sqrt{\tanh(dx + c)} dx \right)$$

input `int((b*tanh(d*x+c))^(1/2),x)`

output `sqrt(b)*int(sqrt(tanh(c + d*x)),x)`

### 3.17 $\int \frac{1}{\sqrt{b \tanh(c+dx)}} dx$

Optimal result	223
Mathematica [A] (verified)	223
Rubi [A] (warning: unable to verify)	224
Maple [A] (verified)	226
Fricas [B] (verification not implemented)	227
Sympy [F]	227
Maxima [F]	228
Giac [F(-2)]	228
Mupad [B] (verification not implemented)	228
Reduce [F]	229

#### Optimal result

Integrand size = 12, antiderivative size = 57

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \frac{\arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{\sqrt{bd}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{\sqrt{bd}}$$

output

$\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(1/2)}/d+\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(1/2)}/d$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \frac{\left(\arctan\left(\sqrt{\tanh(c + dx)}\right) + \operatorname{arctanh}\left(\sqrt{\tanh(c + dx)}\right)\right) \sqrt{\tanh(c + dx)}}{d\sqrt{b \tanh(c + dx)}}$$

input

`Integrate[1/Sqrt[b*Tanh[c + d*x]],x]`



output

$$\left( \left( \text{ArcTan}[\text{Sqrt}[\text{Tanh}[c + d*x]]] + \text{ArcTanh}[\text{Sqrt}[\text{Tanh}[c + d*x]]] \right) * \text{Sqrt}[\text{Tanh}[c + d*x]] \right) / (d * \text{Sqrt}[b * \text{Tanh}[c + d*x]])$$
**Rubi [A] (warning: unable to verify)**

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.89, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sqrt{-ib \tan(ic + idx)}} dx \\ & \quad \downarrow \text{3957} \\ & \frac{b \int -\frac{1}{\sqrt{b \tanh(c+dx)}(b^2 - b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{d} \\ & \quad \downarrow \text{25} \\ & \frac{b \int \frac{1}{\sqrt{b \tanh(c+dx)}(b^2 - b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{d} \\ & \quad \downarrow \text{266} \\ & \frac{2b \int \frac{1}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c + dx)}}{d} \\ & \quad \downarrow \text{756} \\ & \frac{2b \left( \frac{\int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\int \frac{1}{b^2 \tanh^2(c+dx) + b} d\sqrt{b \tanh(c+dx)}}{2b} \right)}{d} \\ & \quad \downarrow \text{216} \end{aligned}$$

$$\frac{2b \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} + \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d}$$

↓ 219

$$\frac{2b \left( \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} + \frac{\operatorname{arctanh}(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{d}$$

input `Int[1/Sqrt[b*Tanh[c + d*x]],x]`

output `(2*b*(ArcTan[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)) + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)))/d`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{\sqrt{b}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{\sqrt{b}d}$	46
default	$\frac{\arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{\sqrt{b}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{\sqrt{b}d}$	46

input `int(1/(b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(1/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(1/2)/d`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 263 vs.  $2(45) = 90$ .

Time = 0.10 (sec) , antiderivative size = 587, normalized size of antiderivative = 10.30

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \text{Too large to display}$$

input `integrate(1/(b*tanh(d*x+c))^(1/2),x, algorithm="fricas")`

output `[-1/4*(2*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) + sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)))/(b*d), 1/4*(2*sqrt(b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/sqrt(b)) + sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b))/(b*d)]`

**Sympy [F]**

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$$

input `integrate(1/(b*tanh(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(b*tanh(c + d*x)), x)`

**Maxima [F]**

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \int \frac{1}{\sqrt{b \tanh(dx + c)}} dx$$

input `integrate(1/(b*tanh(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*tanh(d*x + c)), x)`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(b*tanh(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Degree mismatch inside factorisation over extensionDegree mismatch inside factorisation over extensionindex.c c index_m`

**Mupad [B] (verification not implemented)**

Time = 2.38 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right) + \operatorname{atanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{\sqrt{b} d}$$

input `int(1/(b*tanh(c + d*x))^(1/2),x)`

output `(atan((b*tanh(c + d*x))^(1/2)/b^(1/2)) + atanh((b*tanh(c + d*x))^(1/2)/b^(1/2)))/(b^(1/2)*d)`

**Reduce [F]**

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx = \frac{\sqrt{b} \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)} dx \right)}{b}$$

input `int(1/(b*tanh(d*x+c))^(1/2),x)`

output `(sqrt(b)*int(sqrt(tanh(c + d*x))/tanh(c + d*x),x))/b`

### 3.18 $\int \frac{1}{(b \tanh(c+dx))^{3/2}} dx$

Optimal result	230
Mathematica [A] (verified)	230
Rubi [A] (warning: unable to verify)	231
Maple [A] (verified)	234
Fricas [B] (verification not implemented)	234
Sympy [F]	235
Maxima [F]	236
Giac [F(-2)]	236
Mupad [B] (verification not implemented)	236
Reduce [F]	237

#### Optimal result

Integrand size = 12, antiderivative size = 78

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = -\frac{\arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} - \frac{2}{bd\sqrt{b \tanh(c + dx)}}$$

output

```
-arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(3/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(3/2)/d-2/b/d/(b*tanh(d*x+c))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \frac{-2 - \arctan\left(\sqrt[4]{\tanh^2(c + dx)}\right) \sqrt[4]{\tanh^2(c + dx)} + \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(c + dx)}\right)}{bd\sqrt{b \tanh(c + dx)}}$$

input

```
Integrate[(b*Tanh[c + d*x])^(-3/2), x]
```

output

```
(-2 - ArcTan[(Tanh[c + d*x]^2)^(1/4)]*(Tanh[c + d*x]^2)^(1/4) + ArcTanh[(Tanh[c + d*x]^2)^(1/4)]*(Tanh[c + d*x]^2)^(1/4))/(b*d*Sqrt[b*Tanh[c + d*x]])
```

**Rubi [A] (warning: unable to verify)**

Time = 0.33 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3955, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \tanh(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-ib \tan(ic + idx))^{3/2}} dx \\
 & \quad \downarrow \text{3955} \\
 & \frac{\int \sqrt{b \tanh(c + dx)} dx}{b^2} - \frac{2}{bd \sqrt{b \tanh(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{bd \sqrt{b \tanh(c + dx)}} + \frac{\int \sqrt{-ib \tan(ic + idx)} dx}{b^2} \\
 & \quad \downarrow \text{3957} \\
 & -\frac{\int -\frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{bd} - \frac{2}{bd \sqrt{b \tanh(c + dx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{bd} - \frac{2}{bd \sqrt{b \tanh(c + dx)}} \\
 & \quad \downarrow \text{266}
 \end{aligned}$$



$$\begin{aligned}
& \frac{2 \int \frac{b^2 \tanh^2(c+dx)}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c+dx)}}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} \\
& \quad \downarrow 827 \\
& \frac{2 \left( \frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{1}{2} \int \frac{1}{b^2 \tanh^2(c+dx) + b} d\sqrt{b \tanh(c+dx)} \right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} \\
& \quad \downarrow 216 \\
& \frac{2 \left( \frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} \right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} \\
& \quad \downarrow 219 \\
& \frac{2 \left( \frac{\operatorname{arctanh}(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} \right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}}
\end{aligned}$$

input `Int[(b*Tanh[c + d*x])^(-3/2),x]`

output `(2*(-1/2*ArcTan[Sqrt[b]*Tanh[c + d*x]]/Sqrt[b] + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*Sqrt[b]))/(b*d) - 2/(b*d*Sqrt[b*Tanh[c + d*x]])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{3}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{3}{2}}d} - \frac{2}{bd\sqrt{b}\tanh(dx+c)}$	65
default	$-\frac{\arctan\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{3}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{3}{2}}d} - \frac{2}{bd\sqrt{b}\tanh(dx+c)}$	65

input `int(1/(b*tanh(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `-arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(3/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(3/2)/d-2/b/d/(b*tanh(d*x+c))^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 427 vs. 2(64) = 128.

Time = 0.13 (sec) , antiderivative size = 914, normalized size of antiderivative = 11.72

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="fricas")`

output

```

[-1/4*(2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^
2 - 1)*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) +
sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/(b*cosh(
d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) + (cosh
(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(-b
)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d
*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x
+ c)^4 + 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x +
c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x +
c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2
+ 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*(cosh(d*x + c)^
2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x +
c)/cosh(d*x + c)))/(b^2*d*cosh(d*x + c)^2 + 2*b^2*d*cosh(d*x + c)*sinh(d*
x + c) + b^2*d*sinh(d*x + c)^2 - b^2*d), -1/4*(2*(cosh(d*x + c)^2 + 2*cosh
(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(b)*arctan((cosh(d*x +
c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*
x + c)/cosh(d*x + c))/sqrt(b)) - (cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d
*x + c) + sinh(d*x + c)^2 - 1)*sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(
d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh
(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + ...

```

### Sympy [F]

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \int \frac{1}{(b \tanh(c + dx))^{\frac{3}{2}}} dx$$

input

```
integrate(1/(b*tanh(d*x+c))**(3/2), x)
```

output

```
Integral((b*tanh(c + d*x))**(-3/2), x)
```

**Maxima [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \int \frac{1}{(b \tanh(dx + c))^{3/2}} dx$$

input `integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(3/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.34 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.82

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2} d} - \frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2} d} - \frac{2}{b d \sqrt{b \tanh(c + dx)}}$$

input `int(1/(b*tanh(c + d*x))^(3/2),x)`

output

```
atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(3/2)*d) - atan((b*tanh(c + d*x)
)^(1/2)/b^(1/2))/(b^(3/2)*d) - 2/(b*d*(b*tanh(c + d*x))^(1/2))
```

**Reduce [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx = \frac{\sqrt{b} \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)^2} dx \right)}{b^2}$$

input

```
int(1/(b*tanh(d*x+c))^(3/2),x)
```

output

```
(sqrt(b)*int(sqrt(tanh(c + d*x))/tanh(c + d*x)**2,x))/b**2
```

### 3.19 $\int \frac{1}{(b \tanh(c+dx))^{5/2}} dx$

Optimal result	238
Mathematica [A] (verified)	238
Rubi [A] (warning: unable to verify)	239
Maple [A] (verified)	241
Fricas [B] (verification not implemented)	242
Sympy [F]	243
Maxima [F]	243
Giac [F(-2)]	243
Mupad [B] (verification not implemented)	244
Reduce [F]	244

#### Optimal result

Integrand size = 12, antiderivative size = 79

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \frac{\arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}}$$

output

```
arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d-2/3/b/d/(b*tanh(d*x+c))^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.99

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \frac{-2 + 3 \arctan\left(\sqrt[4]{\tanh^2(c + dx)}\right) \tanh^2(c + dx)^{3/4} + 3 \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(c + dx)}\right)}{3bd(b \tanh(c + dx))^{3/2}}$$

input

```
Integrate[(b*Tanh[c + d*x])^(-5/2),x]
```

output

```
(-2 + 3*ArcTan[(Tanh[c + d*x]^2)^(1/4)]*(Tanh[c + d*x]^2)^(3/4) + 3*ArcTan
h[(Tanh[c + d*x]^2)^(1/4)]*(Tanh[c + d*x]^2)^(3/4))/(3*b*d*(b*Tanh[c + d*x
])^(3/2))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.34 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3955, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(b \tanh(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-ib \tan(ic + idx))^{5/2}} dx \\
 & \quad \downarrow \text{3955} \\
 & \frac{\int \frac{1}{\sqrt{b \tanh(c+dx)}} dx}{b^2} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{3bd(b \tanh(c + dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{-ib \tan(ic+idx)}} dx}{b^2} \\
 & \quad \downarrow \text{3957} \\
 & -\frac{\int -\frac{1}{\sqrt{b \tanh(c+dx)}(b^2 - b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{bd} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{1}{\sqrt{b \tanh(c+dx)}(b^2 - b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{bd} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}} \\
 & \quad \downarrow \text{266}
 \end{aligned}$$



$$\begin{aligned}
& \frac{2 \int \frac{1}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c+dx)}}{bd} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}} \\
& \quad \downarrow \text{756} \\
& \frac{2 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\int \frac{1}{b^2 \tanh^2(c+dx)+b} d\sqrt{b \tanh(c+dx)}}{2b} \right)}{bd} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}} \\
& \quad \downarrow \text{216} \\
& \frac{2 \left( \frac{\int \frac{1}{b-b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)}}{2b} + \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{bd} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{2 \left( \frac{\arctan(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} + \frac{\operatorname{arctanh}(\sqrt{b \tanh(c+dx)})}{2b^{3/2}} \right)}{bd} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}}
\end{aligned}$$

input `Int[(b*Tanh[c + d*x])^(-5/2),x]`

output `(2*(ArcTan[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)) + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*b^(3/2)))/(b*d) - 2/(3*b*d*(b*Tanh[c + d*x])^(3/2))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}d} - \frac{2}{3bd(b \tanh(dx+c))^{\frac{3}{2}}}$	64
default	$\frac{\arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}d} - \frac{2}{3bd(b \tanh(dx+c))^{\frac{3}{2}}}$	64

input `int(1/(b*tanh(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output

```
arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d-2/3/b/d/(b*tanh(d*x+c))^(3/2)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 682 vs.  $2(63) = 126$ .

Time = 0.13 (sec) , antiderivative size = 1425, normalized size of antiderivative = 18.04

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \text{Too large to display}$$

input

```
integrate(1/(b*tanh(d*x+c))^(5/2),x, algorithm="fricas")
```

output

```
[-1/12*(6*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^2 - 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 1)*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) + 3*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^2 - 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 1)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c) + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^3*d*cosh(d*x + c)^4 + 4*b^3*d*cosh(d*x + c)*sinh(d*x + c)^3 + b^3*d*sinh(d*x + c)^4 - 2*b^3*d*cosh(d*x + c)^2 + b^3*d + 2*(3*b^3*d*cosh(d*x + c)^2 - b^3*d)*sinh(d*x + c)^2 + 4*(b^3*d*cosh(d*x + c)^3 - b^3*d*co...
```

**Sympy [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \int \frac{1}{(b \tanh(c + dx))^{\frac{5}{2}}} dx$$

input `integrate(1/(b*tanh(d*x+c))**(5/2), x)`

output `Integral((b*tanh(c + d*x))**(-5/2), x)`

**Maxima [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \int \frac{1}{(b \tanh(dx + c))^{\frac{5}{2}}} dx$$

input `integrate(1/(b*tanh(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(5/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(b*tanh(d*x+c))^(5/2), x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.45 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.80

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2} d} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}} + \frac{\operatorname{atanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2} d}$$

input `int(1/(b*tanh(c + d*x))^(5/2),x)`output `atan((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(5/2)*d) - 2/(3*b*d*(b*tanh(c + d*x))^(3/2)) + atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(5/2)*d)`**Reduce [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx = \frac{\sqrt{b} \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)^3} dx \right)}{b^3}$$

input `int(1/(b*tanh(d*x+c))^(5/2),x)`output `(sqrt(b)*int(sqrt(tanh(c + d*x))/tanh(c + d*x)**3,x))/b**3`

### 3.20 $\int \frac{1}{(b \tanh(c+dx))^{7/2}} dx$

Optimal result	245
Mathematica [A] (verified)	245
Rubi [A] (warning: unable to verify)	246
Maple [A] (verified)	249
Fricas [B] (verification not implemented)	249
Sympy [F]	250
Maxima [F]	251
Giac [F(-2)]	251
Mupad [B] (verification not implemented)	251
Reduce [F]	252

#### Optimal result

Integrand size = 12, antiderivative size = 100

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = -\frac{\arctan\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} - \frac{2}{5bd(b \tanh(c + dx))^{5/2}} - \frac{2}{b^3d\sqrt{b \tanh(c + dx)}}$$

output

```
-arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(7/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(7/2)/d-2/5/b/d/(b*tanh(d*x+c))^(5/2)-2/b^3/d/(b*tanh(d*x+c))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.90

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \frac{-2 \operatorname{coth}^2(c + dx) + 5 \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(c + dx)}\right) \sqrt[4]{\tanh^2(c + dx)} - 5 \left(2 + \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(c + dx)}\right)\right)}{5b^3d\sqrt{b \tanh(c + dx)}}$$

input

```
Integrate[(b*Tanh[c + d*x])^(-7/2),x]
```

output

$$\frac{(-2*\text{Coth}[c + d*x]^2 + 5*\text{ArcTanh}[(\text{Tanh}[c + d*x]^2)^{1/4}]*(\text{Tanh}[c + d*x]^2)^{1/4} - 5*(2 + \text{ArcTan}[(\text{Tanh}[c + d*x]^2)^{1/4}]*(\text{Tanh}[c + d*x]^2)^{1/4}))}{(5*b^3*d*\text{Sqrt}[b*\text{Tanh}[c + d*x]])}$$
**Rubi [A] (warning: unable to verify)**

Time = 0.45 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3955, 3042, 3955, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(b \tanh(c + dx))^{7/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(-ib \tan(ic + idx))^{7/2}} dx \\ & \quad \downarrow \text{3955} \\ & \frac{\int \frac{1}{(b \tanh(c+dx))^{3/2}} dx}{b^2} - \frac{2}{5bd(b \tanh(c + dx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} + \frac{\int \frac{1}{(-ib \tan(ic+idx))^{3/2}} dx}{b^2} \\ & \quad \downarrow \text{3955} \\ & \frac{\int \frac{\sqrt{b \tanh(c+dx)} dx}{b^2} - \frac{2}{bd\sqrt{b \tanh(c+dx)}}}{b^2} - \frac{2}{5bd(b \tanh(c + dx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} + \frac{-\frac{2}{bd\sqrt{b \tanh(c+dx)}} + \frac{\int \sqrt{-ib \tan(ic+idx)} dx}{b^2}}{b^2} \\ & \quad \downarrow \text{3957} \end{aligned}$$

$$\begin{aligned}
 & \frac{\int -\frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c+dx))}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sqrt{b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c+dx))}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2 \int \frac{b^2 \tanh^2(c+dx)}{b^2 - b^4 \tanh^4(c+dx)} d\sqrt{b \tanh(c+dx)}}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}} \\
 & \quad \downarrow \text{827} \\
 & \frac{2\left(\frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{1}{2} \int \frac{1}{b^2 \tanh^2(c+dx) + b} d\sqrt{b \tanh(c+dx)}\right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \\
 & \quad \frac{b^2}{2} \\
 & \quad \frac{2}{5bd(b \tanh(c+dx))^{5/2}} \\
 & \quad \downarrow \text{216} \\
 & \frac{2\left(\frac{1}{2} \int \frac{1}{b - b^2 \tanh^2(c+dx)} d\sqrt{b \tanh(c+dx)} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}}\right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \\
 & \quad \frac{b^2}{2} \\
 & \quad \frac{2}{5bd(b \tanh(c+dx))^{5/2}} \\
 & \quad \downarrow \text{219} \\
 & \frac{2\left(\frac{\operatorname{arctanh}(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}} - \frac{\arctan(\sqrt{b} \tanh(c+dx))}{2\sqrt{b}}\right)}{bd} - \frac{2}{bd\sqrt{b \tanh(c+dx)}} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}}
 \end{aligned}$$

input `Int[(b*Tanh[c + d*x])^(-7/2),x]`

output `-2/(5*b*d*(b*Tanh[c + d*x])^(5/2)) + ((2*(-1/2*ArcTan[Sqrt[b]*Tanh[c + d*x]]/Sqrt[b] + ArcTanh[Sqrt[b]*Tanh[c + d*x]]/(2*Sqrt[b])))/(b*d) - 2/(b*d*Sqrt[b*Tanh[c + d*x]]))/b^2`



## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_-), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216  $\text{Int}[(\text{a}_- + (\text{b}_-)(\text{x}_-)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[\text{b}, 2]))*\text{ArcTan}[\text{Rt}[\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 219  $\text{Int}[(\text{a}_- + (\text{b}_-)(\text{x}_-)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[-\text{b}, 2]))*\text{ArcTanh}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 266  $\text{Int}[(\text{c}_-)(\text{x}_-)^{\text{m}_-}(\text{a}_- + (\text{b}_-)(\text{x}_-)^2)^{\text{p}_-}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[\text{x}^{\text{k}(\text{m} + 1) - 1}(\text{a} + \text{b}(\text{x}^{2\text{k}}/\text{c}^2))^{\text{p}}, \text{x}], \text{x}, (\text{c}\text{x})^{(1/\text{k})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \&\& \text{FractionQ}[\text{m}] \&\& \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 827  $\text{Int}[(\text{x}_-)^2/(\text{a}_- + (\text{b}_-)(\text{x}_-)^4), \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{s}/(2*\text{b}) \quad \text{Int}[1/(\text{r} + \text{s}\text{x}^2), \text{x}], \text{x}] - \text{Simp}[\text{s}/(2*\text{b}) \quad \text{Int}[1/(\text{r} - \text{s}\text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{!GtQ}[\text{a}/\text{b}, 0]$
- rule 3042  $\text{Int}[\text{u}_-, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3955  $\text{Int}[(\text{b}_-)\text{tan}[(\text{c}_- + (\text{d}_-)(\text{x}_-))], \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b}*\text{Tan}[\text{c} + \text{d}\text{x}])^{(\text{n} + 1)}/(\text{b}\text{d}(\text{n} + 1)), \text{x}] - \text{Simp}[1/\text{b}^2 \quad \text{Int}[(\text{b}*\text{Tan}[\text{c} + \text{d}\text{x}])^{(\text{n} + 2)}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[\text{n}, -1]$
- rule 3957  $\text{Int}[(\text{b}_-)\text{tan}[(\text{c}_- + (\text{d}_-)(\text{x}_-))], \text{x\_Symbol}] \rightarrow \text{Simp}[\text{b}/\text{d} \quad \text{Subst}[\text{Int}[\text{x}^{\text{n}}/(\text{b}^2 + \text{x}^2), \text{x}], \text{x}, \text{b}*\text{Tan}[\text{c} + \text{d}\text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}, \text{n}\}, \text{x}] \&\& \text{!IntegerQ}[\text{n}]$

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{7}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{7}{2}}d} - \frac{2}{5bd(b\tanh(dx+c))^{\frac{5}{2}}} - \frac{2}{b^3d\sqrt{b\tanh(dx+c)}}$	83
default	$-\frac{\arctan\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{7}{2}}d} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(dx+c)}{\sqrt{b}}\right)}{b^{\frac{7}{2}}d} - \frac{2}{5bd(b\tanh(dx+c))^{\frac{5}{2}}} - \frac{2}{b^3d\sqrt{b\tanh(dx+c)}}$	83

input `int(1/(b*tanh(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output `-arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(7/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(7/2)/d-2/5/b/d/(b*tanh(d*x+c))^(5/2)-2/b^3/d/(b*tanh(d*x+c))^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1036 vs. 2(82) = 164.

Time = 0.14 (sec) , antiderivative size = 2133, normalized size of antiderivative = 21.33

$$\int \frac{1}{(b\tanh(c+dx))^{7/2}} dx = \text{Too large to display}$$

input `integrate(1/(b*tanh(d*x+c))^(7/2),x, algorithm="fricas")`

output

```

[-1/20*(10*(cosh(d*x + c)^6 + 6*cosh(d*x + c)*sinh(d*x + c)^5 + sinh(d*x +
c)^6 + 3*(5*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^4 - 3*cosh(d*x + c)^4 + 4*
(5*cosh(d*x + c)^3 - 3*cosh(d*x + c))*sinh(d*x + c)^3 + 3*(5*cosh(d*x + c)
^4 - 6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + 3*cosh(d*x + c)^2 + 6*(cosh(
d*x + c)^5 - 2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c) - 1)*sqrt(-b
)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^
2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/(b*cosh(d*x + c)^2 + 2
*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2)) + 5*(cosh(d*x + c)^6
+ 6*cosh(d*x + c)*sinh(d*x + c)^5 + sinh(d*x + c)^6 + 3*(5*cosh(d*x + c)^2
- 1)*sinh(d*x + c)^4 - 3*cosh(d*x + c)^4 + 4*(5*cosh(d*x + c)^3 - 3*cosh(
d*x + c))*sinh(d*x + c)^3 + 3*(5*cosh(d*x + c)^4 - 6*cosh(d*x + c)^2 + 1)*
sinh(d*x + c)^2 + 3*cosh(d*x + c)^2 + 6*(cosh(d*x + c)^5 - 2*cosh(d*x + c)
^3 + cosh(d*x + c))*sinh(d*x + c) - 1)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 +
4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 +
4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^2
+ 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*si
nh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*si
nh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x
+ c)^3 + sinh(d*x + c)^4)) + 16*(3*cosh(d*x + c)^6 + 18*cosh(d*x + c)*sinh
(d*x + c)^5 + 3*sinh(d*x + c)^6 + (45*cosh(d*x + c)^2 - 1)*sinh(d*x + c...

```

### Sympy [F]

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \int \frac{1}{(b \tanh(c + dx))^{7/2}} dx$$

input

```
integrate(1/(b*tanh(d*x+c))**(7/2),x)
```

output

```
Integral((b*tanh(c + d*x))**(-7/2), x)
```

**Maxima [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \int \frac{1}{(b \tanh(dx + c))^{7/2}} dx$$

input `integrate(1/(b*tanh(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^(7/2), x)`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(b*tanh(d*x+c))^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m i_lex_is_greater E  
rror: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.68 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.80

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2} d} - \frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2} d} - \frac{\frac{2}{5b} + \frac{2 \tanh(c+dx)^2}{b}}{d (b \tanh(c + dx))^{5/2}}$$

input `int(1/(b*tanh(c + d*x))^(7/2),x)`

output

```
atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(7/2)*d) - atan((b*tanh(c + d*x)
)^(1/2)/b^(1/2))/(b^(7/2)*d) - (2/(5*b) + (2*tanh(c + d*x)^2)/b)/(d*(b*tan
h(c + d*x))^(5/2))
```

**Reduce [F]**

$$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx = \frac{\sqrt{b} \left( \int \frac{\sqrt{\tanh(dx+c)}}{\tanh(dx+c)^4} dx \right)}{b^4}$$

input

```
int(1/(b*tanh(d*x+c))^(7/2),x)
```

output

```
(sqrt(b)*int(sqrt(tanh(c + d*x))/tanh(c + d*x)**4,x))/b**4
```

### 3.21 $\int \sqrt[3]{\tanh(8x)} dx$

Optimal result	253
Mathematica [A] (verified)	253
Rubi [A] (warning: unable to verify)	254
Maple [A] (verified)	257
Fricas [B] (verification not implemented)	257
Sympy [A] (verification not implemented)	258
Maxima [F]	258
Giac [B] (verification not implemented)	259
Mupad [B] (verification not implemented)	259
Reduce [F]	260

#### Optimal result

Integrand size = 8, antiderivative size = 69

$$\int \sqrt[3]{\tanh(8x)} dx = -\frac{1}{16}\sqrt{3} \arctan\left(\frac{1 + 2 \tanh^{\frac{2}{3}}(8x)}{\sqrt{3}}\right) - \frac{1}{16} \log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32} \log\left(1 + \tanh^{\frac{2}{3}}(8x) + \tanh^{\frac{4}{3}}(8x)\right)$$

output

```
-1/16*3^(1/2)*arctan(1/3*(1+2*tanh(8*x)^(2/3))*3^(1/2))-1/16*ln(1-tanh(8*x)^(2/3))+1/32*ln(1+tanh(8*x)^(2/3)+tanh(8*x)^(4/3))
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.30

$$\int \sqrt[3]{\tanh(8x)} dx = \frac{\left(\log\left(1 - \sqrt[3]{\tanh^2(8x)}\right) - \sqrt[3]{-1} \log\left(1 + \sqrt[3]{-1} \sqrt[3]{\tanh^2(8x)}\right) + (-1)^{2/3} \log\left(1 - (-1)^{2/3} \sqrt[3]{\tanh^2(8x)}\right)\right)}{16 \tanh^2(8x)^{2/3}}$$

input

```
Integrate[Tanh[8*x]^(1/3), x]
```

output

$$-1/16*((\text{Log}[1 - (\text{Tanh}[8*x]^2)^{(1/3)}] - (-1)^{(1/3)}*\text{Log}[1 + (-1)^{(1/3)}*(\text{Tanh}[8*x]^2)^{(1/3)}] + (-1)^{(2/3)}*\text{Log}[1 - (-1)^{(2/3)}*(\text{Tanh}[8*x]^2)^{(1/3)}])*(\text{Tanh}[8*x]^{(4/3)})/(\text{Tanh}[8*x]^2)^{(2/3)}$$
**Rubi [A] (warning: unable to verify)**

Time = 0.30 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {3042, 3957, 25, 266, 807, 821, 16, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt[3]{\tanh(8x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt[3]{-i \tan(8ix)} dx \\ & \quad \downarrow \text{3957} \\ & -\frac{1}{8} \int -\frac{\sqrt[3]{\tanh(8x)}}{1 - \tanh^2(8x)} d \tanh(8x) \\ & \quad \downarrow \text{25} \\ & \frac{1}{8} \int \frac{\sqrt[3]{\tanh(8x)}}{1 - \tanh^2(8x)} d \tanh(8x) \\ & \quad \downarrow \text{266} \\ & \frac{3}{8} \int \frac{\tanh(8x)}{1 - \tanh^2(8x)} d \sqrt[3]{\tanh(8x)} \\ & \quad \downarrow \text{807} \\ & \frac{3}{16} \int \frac{\tanh^{\frac{2}{3}}(8x)}{1 - \tanh(8x)} d \tanh^{\frac{2}{3}}(8x) \\ & \quad \downarrow \text{821} \\ & \frac{3}{16} \left( \frac{1}{3} \int \frac{1}{1 - \tanh^{\frac{2}{3}}(8x)} d \tanh^{\frac{2}{3}}(8x) - \frac{1}{3} \int \frac{1 - \tanh^{\frac{2}{3}}(8x)}{2 \tanh^{\frac{2}{3}}(8x) + 1} d \tanh^{\frac{2}{3}}(8x) \right) \end{aligned}$$

$$\begin{aligned}
& \downarrow 16 \\
& \frac{3}{16} \left( -\frac{1}{3} \int \frac{1 - \tanh^{\frac{2}{3}}(8x)}{2 \tanh^{\frac{2}{3}}(8x) + 1} d \tanh^{\frac{2}{3}}(8x) - \frac{1}{3} \log \left( 1 - \tanh^{\frac{2}{3}}(8x) \right) \right) \\
& \downarrow 1142 \\
& \frac{3}{16} \left( \frac{1}{3} \left( \frac{1}{2} \int 1 d \tanh^{\frac{2}{3}}(8x) - \frac{3}{2} \int \frac{1}{2 \tanh^{\frac{2}{3}}(8x) + 1} d \tanh^{\frac{2}{3}}(8x) \right) - \frac{1}{3} \log \left( 1 - \tanh^{\frac{2}{3}}(8x) \right) \right) \\
& \downarrow 1083 \\
& \frac{3}{16} \left( \frac{1}{3} \left( \frac{1}{2} \int 1 d \tanh^{\frac{2}{3}}(8x) + 3 \int \frac{1}{-2 \tanh^{\frac{2}{3}}(8x) - 4} d \left( 2 \tanh^{\frac{2}{3}}(8x) + 1 \right) \right) - \frac{1}{3} \log \left( 1 - \tanh^{\frac{2}{3}}(8x) \right) \right) \\
& \downarrow 217 \\
& \frac{3}{16} \left( \frac{1}{3} \left( \frac{1}{2} \int 1 d \tanh^{\frac{2}{3}}(8x) - \sqrt{3} \arctan \left( \frac{2 \tanh^{\frac{2}{3}}(8x) + 1}{\sqrt{3}} \right) \right) - \frac{1}{3} \log \left( 1 - \tanh^{\frac{2}{3}}(8x) \right) \right) \\
& \downarrow 1103 \\
& \frac{3}{16} \left( \frac{1}{3} \left( \frac{1}{2} \log \left( 2 \tanh^{\frac{2}{3}}(8x) + 1 \right) - \sqrt{3} \arctan \left( \frac{2 \tanh^{\frac{2}{3}}(8x) + 1}{\sqrt{3}} \right) \right) - \frac{1}{3} \log \left( 1 - \tanh^{\frac{2}{3}}(8x) \right) \right)
\end{aligned}$$

input `Int [Tanh [8*x]^(1/3), x]`

output `(3*(-1/3*Log[1 - Tanh[8*x]^(2/3)] + (-Sqrt[3]*ArcTan[(1 + 2*Tanh[8*x]^(2/3))/Sqrt[3]]) + Log[1 + 2*Tanh[8*x]^(2/3)]/2)/3)/16`

### Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`



rule 217  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^2)^p, x\_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot (x^{2k}/c^2))^p, x], x, (c \cdot x)^{1/k}], x] /;$   $\text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 807  $\text{Int}[(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^n)^p, x\_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m+1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{(m+1)/k-1} \cdot (a + b \cdot x^{n/k})^p, x], x, x^k], x] /;$   $k \neq 1 /;$   $\text{FreeQ}\{a, b, p, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 821  $\text{Int}[(x_ ) / ((a_ + (b_ \cdot)(x_ )^3)), x\_Symbol] \rightarrow \text{Simp}[-(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3])^{(-1)} \ \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1/(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3]) \ \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x) / (\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /;$   $\text{FreeQ}\{a, b, x\}$

rule 1083  $\text{Int}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /;$   $\text{FreeQ}\{a, b, c, x\}$

rule 1103  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)), x\_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e) / (2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e / (2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\}$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.48

method	result
derivativedivides	$-\frac{\ln(\tanh(8x)^{\frac{1}{3}}-1)}{16} + \frac{\ln(\tanh(8x)^{\frac{2}{3}}+\tanh(8x)^{\frac{1}{3}}+1)}{32} + \frac{\sqrt{3} \arctan\left(\frac{(2 \tanh(8x)^{\frac{1}{3}}+1)\sqrt{3}}{3}\right)}{16} - \frac{\ln(\tanh(8x)^{\frac{1}{3}}+1)}{16}$
default	$-\frac{\ln(\tanh(8x)^{\frac{1}{3}}-1)}{16} + \frac{\ln(\tanh(8x)^{\frac{2}{3}}+\tanh(8x)^{\frac{1}{3}}+1)}{32} + \frac{\sqrt{3} \arctan\left(\frac{(2 \tanh(8x)^{\frac{1}{3}}+1)\sqrt{3}}{3}\right)}{16} - \frac{\ln(\tanh(8x)^{\frac{1}{3}}+1)}{16}$

input

```
int(tanh(8*x)^(1/3), x, method=_RETURNVERBOSE)
```

output

```
-1/16*ln(tanh(8*x)^(1/3)-1)+1/32*ln(tanh(8*x)^(2/3)+tanh(8*x)^(1/3)+1)+1/16*3^(1/2)*arctan(1/3*(2*tanh(8*x)^(1/3)+1)*3^(1/2))-1/16*ln(tanh(8*x)^(1/3)+1)+1/32*ln(tanh(8*x)^(2/3)-tanh(8*x)^(1/3)+1)-1/16*3^(1/2)*arctan(1/3*(2*tanh(8*x)^(1/3)-1)*3^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(52) = 104.

Time = 0.08 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.59

$$\int \sqrt[3]{\tanh(8x)} dx$$

$$= -\frac{1}{16} \sqrt{3} \arctan\left(\frac{2}{3} \sqrt{3} \left(\frac{\sinh(8x)}{\cosh(8x)}\right)^{\frac{2}{3}} + \frac{1}{3} \sqrt{3}\right) - \frac{1}{16} \log\left(\left(\frac{\sinh(8x)}{\cosh(8x)}\right)^{\frac{2}{3}} - 1\right)$$

$$+ \frac{1}{32} \log\left(\frac{\cosh(8x)^2 + 2 \cosh(8x) \sinh(8x) + \sinh(8x)^2 + (\cosh(8x)^2 + 2 \cosh(8x) \sinh(8x) + \sinh(8x)^2)}{\cosh(8x)^2 + 2 \cosh(8x) \sinh(8x) + \sinh(8x)^2}\right)$$

input `integrate(tanh(8*x)^(1/3),x, algorithm="fricas")`

output `-1/16*sqrt(3)*arctan(2/3*sqrt(3)*(sinh(8*x)/cosh(8*x))^(2/3) + 1/3*sqrt(3)  
) - 1/16*log((sinh(8*x)/cosh(8*x))^(2/3) - 1) + 1/32*log((cosh(8*x)^2 + 2*  
cosh(8*x)*sinh(8*x) + sinh(8*x)^2 + (cosh(8*x)^2 + 2*cosh(8*x)*sinh(8*x) +  
sinh(8*x)^2 + 1)*(sinh(8*x)/cosh(8*x))^(2/3) + (cosh(8*x)^2 + 2*cosh(8*x)  
*sinh(8*x) + sinh(8*x)^2 - 1)*(sinh(8*x)/cosh(8*x))^(1/3) + 1)/(cosh(8*x)^  
2 + 2*cosh(8*x)*sinh(8*x) + sinh(8*x)^2 + 1))`

### Sympy [A] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.91

$$\int \sqrt[3]{\tanh(8x)} dx = -\frac{\log\left(\tanh^{\frac{2}{3}}(8x) - 1\right)}{16} + \frac{\log\left(\tanh^{\frac{4}{3}}(8x) + \tanh^{\frac{2}{3}}(8x) + 1\right)}{32} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}\left(\tanh^{\frac{2}{3}}(8x) + \frac{1}{2}\right)}{3}\right)}{16}$$

input `integrate(tanh(8*x)**(1/3),x)`

output `-log(tanh(8*x)**(2/3) - 1)/16 + log(tanh(8*x)**(4/3) + tanh(8*x)**(2/3) +  
1)/32 - sqrt(3)*atan(2*sqrt(3)*(tanh(8*x)**(2/3) + 1/2)/3)/16`

### Maxima [F]

$$\int \sqrt[3]{\tanh(8x)} dx = \int \tanh(8x)^{\frac{1}{3}} dx$$

input `integrate(tanh(8*x)^(1/3),x, algorithm="maxima")`

output `integrate(tanh(8*x)^(1/3), x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 110 vs.  $2(52) = 104$ .

Time = 0.13 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.59

$$\int \sqrt[3]{\tanh(8x)} dx = -\frac{1}{16} \sqrt{3} \arctan \left( \frac{1}{3} \sqrt{3} \left( 2 \left( \frac{e^{16x} - 1}{e^{16x} + 1} \right)^{\frac{2}{3}} + 1 \right) \right) \\ + \frac{1}{32} \log \left( \left( \frac{e^{16x} - 1}{e^{16x} + 1} \right)^{\frac{2}{3}} + \frac{\left( \frac{e^{16x} - 1}{e^{16x} + 1} \right)^{\frac{1}{3}} (e^{16x} - 1)}{e^{16x} + 1} + 1 \right) \\ - \frac{1}{16} \log \left( \left| \left( \frac{e^{16x} - 1}{e^{16x} + 1} \right)^{\frac{2}{3}} - 1 \right| \right)$$

input `integrate(tanh(8*x)^(1/3),x, algorithm="giac")`

output `-1/16*sqrt(3)*arctan(1/3*sqrt(3)*(2*((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) + 1)) + 1/32*log(((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) + ((e^(16*x) - 1)/(e^(16*x) + 1))^(1/3)*(e^(16*x) - 1)/(e^(16*x) + 1) + 1) - 1/16*log(abs(((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) - 1))`

**Mupad [B] (verification not implemented)**

Time = 2.60 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.03

$$\int \sqrt[3]{\tanh(8x)} dx = -\frac{\ln \left( 81 \tanh(8x)^{2/3} - 81 \right)}{16} \\ - \ln \left( 162 \tanh(8x)^{2/3} \left( -\frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right) - 81 \right) \left( -\frac{1}{32} + \frac{\sqrt{3} \text{li}}{32} \right) \\ + \ln \left( -162 \tanh(8x)^{2/3} \left( \frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right) - 81 \right) \left( \frac{1}{32} + \frac{\sqrt{3} \text{li}}{32} \right)$$

input `int(tanh(8*x)^(1/3),x)`

output

```
log(- 162*tanh(8*x)^(2/3)*((3^(1/2)*1i)/4 + 1/4) - 81)*((3^(1/2)*1i)/32 +
1/32) - log(162*tanh(8*x)^(2/3)*((3^(1/2)*1i)/4 - 1/4) - 81)*((3^(1/2)*1i)
/32 - 1/32) - log(81*tanh(8*x)^(2/3) - 81)/16
```

**Reduce [F]**

$$\int \sqrt[3]{\tanh(8x)} dx = \int \tanh(8x)^{\frac{1}{3}} dx$$

input

```
int(tanh(8*x)^(1/3), x)
```

output

```
int(tanh(8*x)**(1/3), x)
```

## 3.22 $\int \tanh^n(a + bx) dx$

Optimal result	261
Mathematica [A] (verified)	261
Rubi [A] (verified)	262
Maple [F]	263
Fricas [F]	263
Sympy [F]	264
Maxima [F]	264
Giac [F]	264
Mupad [F(-1)]	265
Reduce [F]	265

### Optimal result

Integrand size = 8, antiderivative size = 43

$$\int \tanh^n(a + bx) dx = \frac{\text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \tanh^2(a + bx)\right) \tanh^{1+n}(a + bx)}{b(1 + n)}$$

output

```
hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], tanh(b*x+a)^2)*tanh(b*x+a)^(1+n)/b/(1+n)
```

### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \tanh^n(a + bx) dx = \frac{\text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \tanh^2(a + bx)\right) \tanh^{1+n}(a + bx)}{b(1 + n)}$$

input

```
Integrate[Tanh[a + b*x]^n,x]
```

output

```
(Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[a + b*x]^2]*Tanh[a + b*x]^(1 + n))/(b*(1 + n))
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3957, 25, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^n(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (-i \tan(ia + ibx))^n dx \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int -\frac{\tanh^n(a+bx)}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\tanh^n(a+bx)}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b} \\
 & \quad \downarrow \text{278} \\
 & \frac{\tanh^{n+1}(a + bx) \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, \tanh^2(a + bx)\right)}{b(n+1)}
 \end{aligned}$$

input `Int [Tanh[a + b*x]^n, x]`

output `(Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[a + b*x]^2]*Tanh[a + b*x]^(1 + n))/(b*(1 + n))`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))]*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**Maple [F]**

$$\int \tanh (bx + a)^n dx$$

input `int(tanh(b*x+a)^n,x)`

output `int(tanh(b*x+a)^n,x)`

**Fricas [F]**

$$\int \tanh^n(a + bx) dx = \int \tanh (bx + a)^n dx$$

input `integrate(tanh(b*x+a)^n,x, algorithm="fricas")`



output `integral(tanh(b*x + a)^n, x)`

### Sympy [F]

$$\int \tanh^n(a + bx) dx = \int \tanh^n(a + bx) dx$$

input `integrate(tanh(b*x+a)**n,x)`

output `Integral(tanh(a + b*x)**n, x)`

### Maxima [F]

$$\int \tanh^n(a + bx) dx = \int \tanh(bx + a)^n dx$$

input `integrate(tanh(b*x+a)^n,x, algorithm="maxima")`

output `integrate(tanh(b*x + a)^n, x)`

### Giac [F]

$$\int \tanh^n(a + bx) dx = \int \tanh(bx + a)^n dx$$

input `integrate(tanh(b*x+a)^n,x, algorithm="giac")`

output `integrate(tanh(b*x + a)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^n(a + bx) dx = \int \tanh(a + bx)^n dx$$

input `int(tanh(a + b*x)^n, x)`output `int(tanh(a + b*x)^n, x)`**Reduce [F]**

$$\int \tanh^n(a + bx) dx = \int \tanh(bx + a)^n dx$$

input `int(tanh(b*x+a)^n, x)`output `int(tanh(a + b*x)**n, x)`

### 3.23 $\int (b \tanh(c + dx))^n dx$

Optimal result	266
Mathematica [A] (verified)	266
Rubi [A] (verified)	267
Maple [F]	268
Fricas [F]	268
Sympy [F]	269
Maxima [F]	269
Giac [F]	269
Mupad [F(-1)]	270
Reduce [F]	270

#### Optimal result

Integrand size = 10, antiderivative size = 48

$$\int (b \tanh(c + dx))^n dx = \frac{\text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \tanh^2(c + dx)\right) (b \tanh(c + dx))^{1+n}}{bd(1+n)}$$

output

```
hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], tanh(d*x+c)^2)*(b*tanh(d*x+c))^(1+n)/
b/d/(1+n)
```

#### Mathematica [A] (verified)

Time = 0.09 (sec), antiderivative size = 49, normalized size of antiderivative = 1.02

$$\int (b \tanh(c + dx))^n dx = \frac{\text{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh(c + dx))^n}{d(1+n)}$$

input

```
Integrate[(b*Tanh[c + d*x])^n,x]
```

output

```
(Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[c + d*x]^2]*Tanh[c + d*x]
*(b*Tanh[c + d*x])^n)/(d*(1 + n))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 3957, 25, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \tanh(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (-ib \tan(ic + idx))^n dx \\
 & \quad \downarrow \text{3957} \\
 & \frac{b \int -\frac{(b \tanh(c+dx))^n}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & \frac{b \int \frac{(b \tanh(c+dx))^n}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\
 & \quad \downarrow \text{278} \\
 & \frac{(b \tanh(c + dx))^{n+1} \text{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, \tanh^2(c + dx)\right)}{bd(n + 1)}
 \end{aligned}$$

input

```
Int[(b*Tanh[c + d*x])^n,x]
```

output

```
(Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[c + d*x]^2]*(b*Tanh[c + d
*x])^(1 + n))/(b*d*(1 + n))
```

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))]*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**Maple [F]**

$$\int (b \tanh(dx + c))^n dx$$

input `int((b*tanh(d*x+c))^n,x)`

output `int((b*tanh(d*x+c))^n,x)`

**Fricas [F]**

$$\int (b \tanh(c + dx))^n dx = \int (b \tanh(dx + c))^n dx$$

input `integrate((b*tanh(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*tanh(d*x + c))^n, x)`

### Sympy [F]

$$\int (b \tanh(c + dx))^n dx = \int (b \tanh(c + dx))^n dx$$

input `integrate((b*tanh(d*x+c))^n,x)`

output `Integral((b*tanh(c + d*x))^n, x)`

### Maxima [F]

$$\int (b \tanh(c + dx))^n dx = \int (b \tanh(dx + c))^n dx$$

input `integrate((b*tanh(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c))^n, x)`

### Giac [F]

$$\int (b \tanh(c + dx))^n dx = \int (b \tanh(dx + c))^n dx$$

input `integrate((b*tanh(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*tanh(d*x + c))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \tanh(c + dx))^n dx = \int (b \tanh(c + dx))^n dx$$

input `int((b*tanh(c + d*x))^n,x)`output `int((b*tanh(c + d*x))^n, x)`**Reduce [F]**

$$\int (b \tanh(c + dx))^n dx = b^n \left( \int \tanh(dx + c)^n dx \right)$$

input `int((b*tanh(d*x+c))^n,x)`output `b**n*int(tanh(c + d*x)**n,x)`

## 3.24 $\int (a \tanh^2(x))^{3/2} dx$

Optimal result	271
Mathematica [A] (verified)	271
Rubi [C] (verified)	272
Maple [A] (verified)	274
Fricas [B] (verification not implemented)	274
Sympy [F]	275
Maxima [A] (verification not implemented)	275
Giac [A] (verification not implemented)	276
Mupad [F(-1)]	276
Reduce [B] (verification not implemented)	276

### Optimal result

Integrand size = 10, antiderivative size = 35

$$\int (a \tanh^2(x))^{3/2} dx = a \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)} - \frac{1}{2} a \tanh(x) \sqrt{a \tanh^2(x)}$$

output

```
a*coth(x)*ln(cosh(x))*(a*tanh(x)^2)^(1/2)-1/2*a*tanh(x)*(a*tanh(x)^2)^(1/2)
```

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

$$\int (a \tanh^2(x))^{3/2} dx = \frac{1}{2} a (2 \coth(x) \log(\cosh(x)) + \operatorname{csch}(x) \operatorname{sech}(x)) \sqrt{a \tanh^2(x)}$$

input

```
Integrate[(a*Tanh[x]^2)^(3/2),x]
```

output

```
(a*(2*Coth[x]*Log[Cosh[x]] + Csch[x]*Sech[x])*Sqrt[a*Tanh[x]^2])/2
```



**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4141, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (-a \tan(ix)^2)^{3/2} dx \\
 & \quad \downarrow \text{4141} \\
 & a \coth(x) \sqrt{a \tanh^2(x)} \int \tanh^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a \coth(x) \sqrt{a \tanh^2(x)} \int i \tan(ix)^3 dx \\
 & \quad \downarrow \text{26} \\
 & ia \coth(x) \sqrt{a \tanh^2(x)} \int \tan(ix)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & ia \coth(x) \sqrt{a \tanh^2(x)} \left( \frac{1}{2} i \tanh^2(x) - \int i \tanh(x) dx \right) \\
 & \quad \downarrow \text{26} \\
 & ia \coth(x) \sqrt{a \tanh^2(x)} \left( \frac{1}{2} i \tanh^2(x) - i \int \tanh(x) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & ia \coth(x) \sqrt{a \tanh^2(x)} \left( \frac{1}{2} i \tanh^2(x) - i \int -i \tan(ix) dx \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$ia \operatorname{coth}(x) \sqrt{a \tanh^2(x)} \left( \frac{1}{2} i \tanh^2(x) - \int \tan(ix) dx \right)$$

↓ 3956

$$ia \operatorname{coth}(x) \sqrt{a \tanh^2(x)} \left( \frac{1}{2} i \tanh^2(x) - i \log(\cosh(x)) \right)$$

input `Int[(a*Tanh[x]^2)^(3/2),x]`

output `I*a*Coth[x]*Sqrt[a*Tanh[x]^2]*((-I)*Log[Cosh[x]] + (I/2)*Tanh[x]^2)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x, x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])`

**Maple [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$-\frac{(\tanh(x)^2 a)^{\frac{3}{2}} (\tanh(x)^2 + \ln(\tanh(x)-1) + \ln(1+\tanh(x)))}{2 \tanh(x)^3}$	30
default	$-\frac{(\tanh(x)^2 a)^{\frac{3}{2}} (\tanh(x)^2 + \ln(\tanh(x)-1) + \ln(1+\tanh(x)))}{2 \tanh(x)^3}$	30
risch	$a \sqrt{\frac{(e^{2x}-1)^2 a}{(e^{2x}+1)^2}} \frac{(e^{4x} \ln(e^{2x}+1) - e^{4x} x + 2 e^{2x} \ln(e^{2x}+1) - 2 e^{2x} x + 2 e^{2x} + \ln(e^{2x}+1) - x)}{(e^{2x}-1)(e^{2x}+1)}$	95

input `int((tanh(x)^2*a)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/2*(tanh(x)^2*a)^(3/2)*(tanh(x)^2+ln(tanh(x)-1)+ln(1+tanh(x)))/tanh(x)^3`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 467 vs. 2(29) = 58.

Time = 0.12 (sec) , antiderivative size = 467, normalized size of antiderivative = 13.34

$$\int (a \tanh^2(x))^{3/2} dx = \text{Too large to display}$$

input `integrate((a*tanh(x)^2)^(3/2),x, algorithm="fricas")`

output

```

-(a*x*cosh(x)^4 + (a*x*e^(2*x) + a*x)*sinh(x)^4 + 4*(a*x*cosh(x)*e^(2*x) +
a*x*cosh(x))*sinh(x)^3 + 2*(a*x - a)*cosh(x)^2 + 2*(3*a*x*cosh(x)^2 + a*x
+ (3*a*x*cosh(x)^2 + a*x - a)*e^(2*x) - a)*sinh(x)^2 + a*x + (a*x*cosh(x)
^4 + 2*(a*x - a)*cosh(x)^2 + a*x)*e^(2*x) - (a*cosh(x)^4 + (a*e^(2*x) + a)
*sinh(x)^4 + 4*(a*cosh(x)*e^(2*x) + a*cosh(x))*sinh(x)^3 + 2*a*cosh(x)^2 +
2*(3*a*cosh(x)^2 + (3*a*cosh(x)^2 + a)*e^(2*x) + a)*sinh(x)^2 + (a*cosh(x)
)^4 + 2*a*cosh(x)^2 + a)*e^(2*x) + 4*(a*cosh(x)^3 + a*cosh(x) + (a*cosh(x)
^3 + a*cosh(x))*e^(2*x))*sinh(x) + a)*log(2*cosh(x)/(cosh(x) - sinh(x))) +
4*(a*x*cosh(x)^3 + (a*x - a)*cosh(x) + (a*x*cosh(x)^3 + (a*x - a)*cosh(x)
)*e^(2*x))*sinh(x))*sqrt((a*e^(4*x) - 2*a*e^(2*x) + a)/(e^(4*x) + 2*e^(2*x)
+ 1))/((e^(2*x) - 1)*sinh(x)^4 - cosh(x)^4 + 4*(cosh(x)*e^(2*x) - cosh(x)
))*sinh(x)^3 - 2*(3*cosh(x)^2 - (3*cosh(x)^2 + 1)*e^(2*x) + 1)*sinh(x)^2 -
2*cosh(x)^2 + (cosh(x)^4 + 2*cosh(x)^2 + 1)*e^(2*x) - 4*(cosh(x)^3 - (cos
h(x)^3 + cosh(x))*e^(2*x) + cosh(x))*sinh(x) - 1)

```

**Sympy [F]**

$$\int (a \tanh^2(x))^{3/2} dx = \int (a \tanh^2(x))^{\frac{3}{2}} dx$$

input

```
integrate((a*tanh(x)**2)**(3/2),x)
```

output

```
Integral((a*tanh(x)**2)**(3/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int (a \tanh^2(x))^{3/2} dx = -a^{\frac{3}{2}}x - a^{\frac{3}{2}} \log(e^{(-2x)} + 1) - \frac{2a^{\frac{3}{2}}e^{(-2x)}}{2e^{(-2x)} + e^{(-4x)} + 1}$$

input

```
integrate((a*tanh(x)^2)^(3/2),x, algorithm="maxima")
```

output

```

-a^(3/2)*x - a^(3/2)*log(e^(-2*x) + 1) - 2*a^(3/2)*e^(-2*x)/(2*e^(-2*x) +
e^(-4*x) + 1)

```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.49

$$\int (a \tanh^2(x))^{3/2} dx = -\left(x \operatorname{sgn}(e^{4x} - 1) - \log(e^{2x} + 1) \operatorname{sgn}(e^{4x} - 1) - \frac{2e^{2x} \operatorname{sgn}(e^{4x} - 1)}{(e^{2x} + 1)^2}\right) a^{3/2}$$

input `integrate((a*tanh(x)^2)^(3/2),x, algorithm="giac")`output `-(x*sgn(e^(4*x) - 1) - log(e^(2*x) + 1)*sgn(e^(4*x) - 1) - 2*e^(2*x)*sgn(e^(4*x) - 1)/(e^(2*x) + 1)^2)*a^(3/2)`**Mupad [F(-1)]**

Timed out.

$$\int (a \tanh^2(x))^{3/2} dx = \int (a \tanh(x)^2)^{3/2} dx$$

input `int((a*tanh(x)^2)^(3/2),x)`output `int((a*tanh(x)^2)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.43

$$\int (a \tanh^2(x))^{3/2} dx = \frac{\sqrt{a} a (e^{4x} \log(e^{2x} + 1) - e^{4x} x - e^{4x} + 2e^{2x} \log(e^{2x} + 1) - 2e^{2x} x + \log(e^{2x} + 1) - x)}{e^{4x} + 2e^{2x} + 1}$$

input `int((a*tanh(x)^2)^(3/2),x)`

output

```
(sqrt(a)*a*(e**(4*x)*log(e**(2*x) + 1) - e**(4*x)*x - e**(4*x) + 2*e**(2*x)
)*log(e**(2*x) + 1) - 2*e**(2*x)*x + log(e**(2*x) + 1) - x - 1))/(e**(4*x)
+ 2*e**(2*x) + 1)
```

### 3.25 $\int \sqrt{a \tanh^2(x)} dx$

Optimal result	278
Mathematica [A] (verified)	278
Rubi [A] (verified)	279
Maple [A] (verified)	280
Fricas [B] (verification not implemented)	281
Sympy [F]	281
Maxima [A] (verification not implemented)	281
Giac [B] (verification not implemented)	282
Mupad [F(-1)]	282
Reduce [B] (verification not implemented)	282

#### Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \sqrt{a \tanh^2(x)} dx = \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)}$$

output

```
coth(x)*ln(cosh(x))*(a*tanh(x)^2)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sqrt{a \tanh^2(x)} dx = \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)}$$

input

```
Integrate[Sqrt[a*Tanh[x]^2],x]
```

output

```
Coth[x]*Log[Cosh[x]]*Sqrt[a*Tanh[x]^2]
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4141, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \tanh^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-a \tan(ix)^2} dx \\
 & \quad \downarrow \text{4141} \\
 & \coth(x) \sqrt{a \tanh^2(x)} \int \tanh(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \coth(x) \sqrt{a \tanh^2(x)} \int -i \tan(ix) dx \\
 & \quad \downarrow \text{26} \\
 & -i \coth(x) \sqrt{a \tanh^2(x)} \int \tan(ix) dx \\
 & \quad \downarrow \text{3956} \\
 & \coth(x) \sqrt{a \tanh^2(x)} \log(\cosh(x))
 \end{aligned}$$

input `Int[Sqrt[a*Tanh[x]^2], x]`

output `Coth[x]*Log[Cosh[x]]*Sqrt[a*Tanh[x]^2]`



## Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)^(n_)])^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

method	result	size
derivativedivides	$-\frac{\sqrt{\tanh(x)^2 a} (\ln(\tanh(x)-1) + \ln(1+\tanh(x)))}{2 \tanh(x)}$	26
default	$-\frac{\sqrt{\tanh(x)^2 a} (\ln(\tanh(x)-1) + \ln(1+\tanh(x)))}{2 \tanh(x)}$	26
risch	$-\sqrt{\frac{(e^{2x}-1)^2 a}{(e^{2x}+1)^2}} (e^{2x}+1)x + \sqrt{\frac{(e^{2x}-1)^2 a}{(e^{2x}+1)^2}} (e^{2x}+1) \ln(e^{2x}+1)$	81

input `int((tanh(x)^2*a)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(tanh(x)^2*a)^(1/2)*(ln(tanh(x)-1)+ln(1+tanh(x)))/tanh(x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 72 vs.  $2(14) = 28$ .

Time = 0.11 (sec) , antiderivative size = 72, normalized size of antiderivative = 4.50

$$\int \sqrt{a \tanh^2(x)} dx = -\frac{\left(xe^{(2x)} - (e^{(2x)} + 1) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right) + x\right) \sqrt{\frac{ae^{(4x)} - 2ae^{(2x)} + a}{e^{(4x)} + 2e^{(2x)} + 1}}}{e^{(2x)} - 1}$$

input `integrate((a*tanh(x)^2)^(1/2),x, algorithm="fricas")`

output `-(x*e^(2*x) - (e^(2*x) + 1)*log(2*cosh(x)/(cosh(x) - sinh(x))) + x)*sqrt((a*e^(4*x) - 2*a*e^(2*x) + a)/(e^(4*x) + 2*e^(2*x) + 1))/(e^(2*x) - 1)`

**Sympy [F]**

$$\int \sqrt{a \tanh^2(x)} dx = \int \sqrt{a \tanh^2(x)} dx$$

input `integrate((a*tanh(x)**2)**(1/2),x)`

output `Integral(sqrt(a*tanh(x)**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \sqrt{a \tanh^2(x)} dx = -\sqrt{a}x - \sqrt{a} \log(e^{(-2x)} + 1)$$

input `integrate((a*tanh(x)^2)^(1/2),x, algorithm="maxima")`

output `-sqrt(a)*x - sqrt(a)*log(e^(-2*x) + 1)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs.  $2(14) = 28$ .

Time = 0.13 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \sqrt{a \tanh^2(x)} dx = -(x \operatorname{sgn}(e^{4x} - 1) - \log(e^{2x} + 1) \operatorname{sgn}(e^{4x} - 1)) \sqrt{a}$$

input `integrate((a*tanh(x)^2)^(1/2),x, algorithm="giac")`

output `-(x*sgn(e^(4*x) - 1) - log(e^(2*x) + 1)*sgn(e^(4*x) - 1))*sqrt(a)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \tanh^2(x)} dx = \int \sqrt{a \tanh(x)^2} dx$$

input `int((a*tanh(x)^2)^(1/2),x)`

output `int((a*tanh(x)^2)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \sqrt{a \tanh^2(x)} dx = \sqrt{a} (\log(e^{2x} + 1) - x)$$

input `int((a*tanh(x)^2)^(1/2),x)`

output `sqrt(a)*(log(e**(2*x) + 1) - x)`

$$3.26 \quad \int \frac{1}{\sqrt{a \tanh^2(x)}} dx$$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [B] (verified)	285
Fricas [B] (verification not implemented)	286
Sympy [F]	286
Maxima [B] (verification not implemented)	287
Giac [A] (verification not implemented)	287
Mupad [B] (verification not implemented)	287
Reduce [B] (verification not implemented)	288

### Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = \frac{\log(\sinh(x)) \tanh(x)}{\sqrt{a \tanh^2(x)}}$$

output `ln(sinh(x))*tanh(x)/(a*tanh(x)^2)^(1/2)`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = \frac{\log(\sinh(x)) \tanh(x)}{\sqrt{a \tanh^2(x)}}$$

input `Integrate[1/Sqrt[a*Tanh[x]^2],x]`

output `(Log[Sinh[x]]*Tanh[x])/Sqrt[a*Tanh[x]^2]`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4141, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \tanh^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-a \tan(ix)^2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\tanh(x) \int \coth(x) dx}{\sqrt{a \tanh^2(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh(x) \int -i \tan(ix + \frac{\pi}{2}) dx}{\sqrt{a \tanh^2(x)}} \\
 & \quad \downarrow \text{26} \\
 & -\frac{i \tanh(x) \int \tan(ix + \frac{\pi}{2}) dx}{\sqrt{a \tanh^2(x)}} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\tanh(x) \log(\sinh(x))}{\sqrt{a \tanh^2(x)}}
 \end{aligned}$$

input

 $\text{Int}[1/\text{Sqrt}[a*\text{Tanh}[x]^2], x]$ 

output

 $(\text{Log}[\text{Sinh}[x]]*\text{Tanh}[x])/\text{Sqrt}[a*\text{Tanh}[x]^2]$

## Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)^(n_)])^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*(b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs.  $2(14) = 28$ .

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

method	result	size
derivativedivides	$\frac{\tanh(x)(2 \ln(\tanh(x)) - \ln(\tanh(x)-1) - \ln(1+\tanh(x)))}{2\sqrt{\tanh(x)^2 a}}$	33
default	$\frac{\tanh(x)(2 \ln(\tanh(x)) - \ln(\tanh(x)-1) - \ln(1+\tanh(x)))}{2\sqrt{\tanh(x)^2 a}}$	33
risch	$-\frac{(e^{2x}-1)x}{\sqrt{\frac{(e^{2x}-1)^2 a}{(e^{2x}+1)^2}}(e^{2x}+1)} + \frac{(e^{2x}-1) \ln(e^{2x}-1)}{\sqrt{\frac{(e^{2x}-1)^2 a}{(e^{2x}+1)^2}}(e^{2x}+1)}$	81

input `int(1/(tanh(x)^2*a)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*tanh(x)*(2*ln(tanh(x))-ln(tanh(x)-1)-ln(1+tanh(x)))/(tanh(x)^2*a)^(1/2)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs.  $2(14) = 28$ .

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 4.75

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = -\frac{\left(xe^{(2x)} - (e^{(2x)} + 1) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) + x\right) \sqrt{\frac{ae^{(4x)} - 2ae^{(2x)} + a}{e^{(4x)} + 2e^{(2x)} + 1}}}{ae^{(2x)} - a}$$

input `integrate(1/(a*tanh(x)^2)^(1/2),x, algorithm="fricas")`

output `-(x*e^(2*x) - (e^(2*x) + 1)*log(2*sinh(x)/(cosh(x) - sinh(x))) + x)*sqrt((a*e^(4*x) - 2*a*e^(2*x) + a)/(e^(4*x) + 2*e^(2*x) + 1))/(a*e^(2*x) - a)`

### Sympy [F]

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = \int \frac{1}{\sqrt{a \tanh^2(x)}} dx$$

input `integrate(1/(a*tanh(x)**2)**(1/2),x)`

output `Integral(1/sqrt(a*tanh(x)**2), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs.  $2(14) = 28$ .

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = -\frac{x}{\sqrt{a}} - \frac{\log(e^{-x} + 1)}{\sqrt{a}} - \frac{\log(e^{-x} - 1)}{\sqrt{a}}$$

input `integrate(1/(a*tanh(x)^2)^(1/2),x, algorithm="maxima")`

output `-x/sqrt(a) - log(e^(-x) + 1)/sqrt(a) - log(e^(-x) - 1)/sqrt(a)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 1, normalized size of antiderivative = 0.06

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = 0$$

input `integrate(1/(a*tanh(x)^2)^(1/2),x, algorithm="giac")`

output `0`

**Mupad [B] (verification not implemented)**

Time = 2.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\tanh(x)}{\sqrt{\tanh(x)^2}}\right)}{\sqrt{a}}$$

input `int(1/(a*tanh(x)^2)^(1/2),x)`



output `atanh(tanh(x)/(tanh(x)^2)^(1/2))/a^(1/2)`

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx = \frac{\sqrt{a} (\log(e^x - 1) + \log(e^x + 1) - x)}{a}$$

input `int(1/(a*tanh(x)^2)^(1/2),x)`

output `(sqrt(a)*(log(e**x - 1) + log(e**x + 1) - x))/a`

### 3.27 $\int (-\tanh^2(c + dx))^{5/2} dx$

Optimal result	289
Mathematica [A] (verified)	289
Rubi [C] (verified)	290
Maple [A] (verified)	292
Fricas [C] (verification not implemented)	293
Sympy [F]	293
Maxima [C] (verification not implemented)	294
Giac [C] (verification not implemented)	294
Mupad [F(-1)]	295
Reduce [B] (verification not implemented)	295

#### Optimal result

Integrand size = 14, antiderivative size = 88

$$\int (-\tanh^2(c + dx))^{5/2} dx = \frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d} - \frac{\tanh(c + dx) \sqrt{-\tanh^2(c + dx)}}{2d} - \frac{\tanh^3(c + dx) \sqrt{-\tanh^2(c + dx)}}{4d}$$

output

```
coth(d*x+c)*ln(cosh(d*x+c))*(-tanh(d*x+c)^2)^(1/2)/d-1/2*tanh(d*x+c)*(-tanh(d*x+c)^2)^(1/2)/d-1/4*tanh(d*x+c)^3*(-tanh(d*x+c)^2)^(1/2)/d
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.67

$$\int (-\tanh^2(c + dx))^{5/2} dx = \frac{\coth^5(c + dx) (4 \log(\cosh(c + dx)) + 4 \operatorname{sech}^2(c + dx) - \operatorname{sech}^4(c + dx)) (-\tanh^2(c + dx))^{5/2}}{4d}$$

input

```
Integrate[(-Tanh[c + d*x]^2)^(5/2), x]
```

output

$$\frac{(\operatorname{Coth}[c + dx]^5(4\operatorname{Log}[\operatorname{Cosh}[c + dx]] + 4\operatorname{Sech}[c + dx]^2 - \operatorname{Sech}[c + dx]^4)(-\operatorname{Tanh}[c + dx]^2)^{5/2})}{(4d)}$$
**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.83, number of steps used = 13, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$ , Rules used = {3042, 4141, 3042, 26, 3954, 26, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (-\tanh^2(c + dx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (\tan(ic + idx)^2)^{5/2} dx \\ & \quad \downarrow \text{4141} \\ & \sqrt{-\tanh^2(c + dx) \operatorname{coth}(c + dx)} \int \tanh^5(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{-\tanh^2(c + dx) \operatorname{coth}(c + dx)} \int -i \tan(ic + idx)^5 dx \\ & \quad \downarrow \text{26} \\ & -i \sqrt{-\tanh^2(c + dx) \operatorname{coth}(c + dx)} \int \tan(ic + idx)^5 dx \\ & \quad \downarrow \text{3954} \\ & -i \sqrt{-\tanh^2(c + dx) \operatorname{coth}(c + dx)} \left( - \int -i \tanh^3(c + dx) dx - \frac{i \tanh^4(c + dx)}{4d} \right) \\ & \quad \downarrow \text{26} \\ & -i \sqrt{-\tanh^2(c + dx) \operatorname{coth}(c + dx)} \left( i \int \tanh^3(c + dx) dx - \frac{i \tanh^4(c + dx)}{4d} \right) \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( i \int i \tan(ic+idx)^3 dx - \frac{i \tanh^4(c+dx)}{4d} \right) \\
& \downarrow 26 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( - \int \tan(ic+idx)^3 dx - \frac{i \tanh^4(c+dx)}{4d} \right) \\
& \downarrow 3954 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( \int i \tanh(c+dx) dx - \frac{i \tanh^4(c+dx)}{4d} - \frac{i \tanh^2(c+dx)}{2d} \right) \\
& \downarrow 26 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( i \int \tanh(c+dx) dx - \frac{i \tanh^4(c+dx)}{4d} - \frac{i \tanh^2(c+dx)}{2d} \right) \\
& \downarrow 3042 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( i \int -i \tan(ic+idx) dx - \frac{i \tanh^4(c+dx)}{4d} - \frac{i \tanh^2(c+dx)}{2d} \right) \\
& \downarrow 26 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( \int \tan(ic+idx) dx - \frac{i \tanh^4(c+dx)}{4d} - \frac{i \tanh^2(c+dx)}{2d} \right) \\
& \downarrow 3956 \\
& -i\sqrt{-\tanh^2(c+dx)} \operatorname{coth}(c+dx) \left( -\frac{i \tanh^4(c+dx)}{4d} - \frac{i \tanh^2(c+dx)}{2d} + \frac{i \log(\cosh(c+dx))}{d} \right)
\end{aligned}$$

input `Int[(-Tanh[c + d*x]^2)^(5/2), x]`

output `(-I)*Coth[c + d*x]*Sqrt[-Tanh[c + d*x]^2]*((I*Log[Cosh[c + d*x]])/d - ((I/2)*Tanh[c + d*x]^2)/d - ((I/4)*Tanh[c + d*x]^4)/d)`

Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.76

method	result
derivativedivides	$-\frac{\left(-\tanh(dx+c)^2\right)^{\frac{5}{2}}\left(\tanh(dx+c)^4+2\tanh(dx+c)^2+2\ln(\tanh(dx+c)-1)+2\ln(\tanh(dx+c)+1)\right)}{4d\tanh(dx+c)^5}$
default	$-\frac{\left(-\tanh(dx+c)^2\right)^{\frac{5}{2}}\left(\tanh(dx+c)^4+2\tanh(dx+c)^2+2\ln(\tanh(dx+c)-1)+2\ln(\tanh(dx+c)+1)\right)}{4d\tanh(dx+c)^5}$
risch	$\frac{(e^{2dx+2c}+1)\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}x}{e^{2dx+2c}-1} - \frac{2(e^{2dx+2c}+1)\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}(dx+c)}{(e^{2dx+2c}-1)d} + \frac{4\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}e^{2dx+2c}(e^{4dx+4c}-1)}{(e^{2dx+2c}-1)(e^{2dx+2c}+1)}$

input `int((-tanh(d*x+c)^2)^(5/2),x,method=_RETURNVERBOSE)`

output `-1/4/d*(-tanh(d*x+c)^2)^(5/2)*(tanh(d*x+c)^4+2*tanh(d*x+c)^2+2*ln(tanh(d*x+c)-1)+2*ln(tanh(d*x+c)+1))/tanh(d*x+c)^5`

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.05

$$\int (-\tanh^2(c + dx))^{5/2} dx = \frac{-i dx e^{(8 dx + 8 c)} - i dx - 4 (i dx - i) e^{(6 dx + 6 c)} - 2 (3i dx - 2i) e^{(4 dx + 4 c)} - 4 (i dx - i) e^{(2 dx + 2 c)}}{de^{(8 dx + 8 c)} + 4 de^{(6 dx + 6 c)} + 6 de^{(4 dx + 4 c)} + 4 de^{(2 dx + 2 c)} + d}$$

input `integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="fricas")`

output `(-I*d*x*e^(8*d*x + 8*c) - I*d*x - 4*(I*d*x - I)*e^(6*d*x + 6*c) - 2*(3*I*d*x - 2*I)*e^(4*d*x + 4*c) - 4*(I*d*x - I)*e^(2*d*x + 2*c) + (I*e^(8*d*x + 8*c) + 4*I*e^(6*d*x + 6*c) + 6*I*e^(4*d*x + 4*c) + 4*I*e^(2*d*x + 2*c) + I)*log(e^(2*d*x + 2*c) + 1)/(d*e^(8*d*x + 8*c) + 4*d*e^(6*d*x + 6*c) + 6*d*e^(4*d*x + 4*c) + 4*d*e^(2*d*x + 2*c) + d)`

### Sympy [F]

$$\int (-\tanh^2(c + dx))^{5/2} dx = \int (-\tanh^2(c + dx))^{\frac{5}{2}} dx$$

input `integrate((-tanh(d*x+c)**2)**(5/2),x)`

output `Integral((-tanh(c + d*x)**2)**(5/2), x)`

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.28

$$\int (-\tanh^2(c+dx))^{5/2} dx = -\frac{i(dx+c)}{d} - \frac{i \log(e^{(-2dx-2c)}+1)}{d} + \frac{4(-ie^{(-2dx-2c)} - ie^{(-4dx-4c)} - ie^{(-6dx-6c)})}{d(4e^{(-2dx-2c)} + 6e^{(-4dx-4c)} + 4e^{(-6dx-6c)} + e^{(-8dx-8c)} + 1)}$$

input `integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="maxima")`

output `-I*(d*x + c)/d - I*log(e^(-2*d*x - 2*c) + 1)/d + 4*(-I*e^(-2*d*x - 2*c) - I*e^(-4*d*x - 4*c) - I*e^(-6*d*x - 6*c))/(d*(4*e^(-2*d*x - 2*c) + 6*e^(-4*d*x - 4*c) + 4*e^(-6*d*x - 6*c) + e^(-8*d*x - 8*c) + 1))`

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.61

$$\int (-\tanh^2(c+dx))^{5/2} dx = \frac{i(dx+c)\operatorname{sgn}(-e^{(4dx+4c)}+1) - i \log(e^{(2dx+2c)}+1)\operatorname{sgn}(-e^{(4dx+4c)}+1) - \frac{4i(e^{(6dx+6c)}\operatorname{sgn}(-e^{(4dx+4c)}+1))}{d}}{d}$$

input `integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="giac")`

output `(I*(d*x + c)*sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) + 1)*sgn(-e^(4*d*x + 4*c) + 1) - 4*I*(e^(6*d*x + 6*c))*sgn(-e^(4*d*x + 4*c) + 1) + e^(4*d*x + 4*c)*sgn(-e^(4*d*x + 4*c) + 1) + e^(2*d*x + 2*c)*sgn(-e^(4*d*x + 4*c) + 1))/(e^(2*d*x + 2*c) + 1)^4/d`

**Mupad [F(-1)]**

Timed out.

$$\int (-\tanh^2(c + dx))^{5/2} dx = \int (-\tanh(c + dx)^2)^{5/2} dx$$

input `int((-tanh(c + d*x)^2)^(5/2), x)`output `int((-tanh(c + d*x)^2)^(5/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.88

$$\int (-\tanh^2(c + dx))^{5/2} dx = \frac{i(e^{8dx+8c}\log(e^{2dx+2c} + 1) - e^{8dx+8c}dx - e^{8dx+8c} + 4e^{6dx+6c}\log(e^{2dx+2c} + 1) - 4e^{6dx+6c}dx + 6e^{4dx+4c}\log(e^{2dx+2c} + 1) - 6e^{4dx+4c}dx + 4e^{2dx+2c}\log(e^{2dx+2c} + 1) - 4e^{2dx+2c}dx + \log(e^{2dx+2c} + 1) - dx - 1)}{d(e^{8dx+8c} + 4e^{6dx+6c} + 6e^{4dx+4c} + 4e^{2dx+2c} + 1)}$$

input `int((-tanh(d*x+c)^2)^(5/2), x)`output `(i*(e**(8*c + 8*d*x)*log(e**(2*c + 2*d*x) + 1) - e**(8*c + 8*d*x)*d*x - e**(8*c + 8*d*x) + 4*e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x) + 1) - 4*e**(6*c + 6*d*x)*d*x + 6*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1) - 6*e**(4*c + 4*d*x)*d*x - 2*e**(4*c + 4*d*x) + 4*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1) - 4*e**(2*c + 2*d*x)*d*x + log(e**(2*c + 2*d*x) + 1) - d*x - 1))/(d*(e**(8*c + 8*d*x) + 4*e**(6*c + 6*d*x) + 6*e**(4*c + 4*d*x) + 4*e**(2*c + 2*d*x) + 1))`



### 3.28 $\int (-\tanh^2(c + dx))^{3/2} dx$

Optimal result	296
Mathematica [A] (verified)	296
Rubi [C] (verified)	297
Maple [A] (verified)	299
Fricas [C] (verification not implemented)	299
Sympy [F]	300
Maxima [C] (verification not implemented)	300
Giac [C] (verification not implemented)	301
Mupad [F(-1)]	301
Reduce [B] (verification not implemented)	302

#### Optimal result

Integrand size = 14, antiderivative size = 60

$$\int (-\tanh^2(c + dx))^{3/2} dx = -\frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d} + \frac{\tanh(c + dx) \sqrt{-\tanh^2(c + dx)}}{2d}$$

output

`-coth(d*x+c)*ln(cosh(d*x+c))*(-tanh(d*x+c)^2)^(1/2)/d+1/2*tanh(d*x+c)*(-tanh(d*x+c)^2)^(1/2)/d`

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int (-\tanh^2(c + dx))^{3/2} dx = \frac{\coth^3(c + dx) (2 \log(\cosh(c + dx)) + \operatorname{sech}^2(c + dx)) (-\tanh^2(c + dx))^{3/2}}{2d}$$

input

`Integrate[(-Tanh[c + d*x]^2)^(3/2), x]`

output

```
(Coth[c + d*x]^3*(2*Log[Cosh[c + d*x]] + Sech[c + d*x]^2)*(-Tanh[c + d*x]^2)^(3/2))/(2*d)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$ , Rules used = {3042, 4141, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (-\tanh^2(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (\tan(ic + idx)^2)^{3/2} dx \\
 & \quad \downarrow \text{4141} \\
 & \sqrt{-\tanh^2(c + dx)}(-\coth(c + dx)) \int \tanh^3(c + dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{-\tanh^2(c + dx)}(-\coth(c + dx)) \int i \tan(ic + idx)^3 dx \\
 & \quad \downarrow \text{26} \\
 & -i\sqrt{-\tanh^2(c + dx)} \coth(c + dx) \int \tan(ic + idx)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & -i\sqrt{-\tanh^2(c + dx)} \coth(c + dx) \left( \frac{i \tanh^2(c + dx)}{2d} - \int i \tanh(c + dx) dx \right) \\
 & \quad \downarrow \text{26} \\
 & -i\sqrt{-\tanh^2(c + dx)} \coth(c + dx) \left( \frac{i \tanh^2(c + dx)}{2d} - i \int \tanh(c + dx) dx \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& -i\sqrt{-\tanh^2(c+dx)} \coth(c+dx) \left( \frac{i \tanh^2(c+dx)}{2d} - i \int -i \tan(ic+idx) dx \right) \\
& \downarrow 26 \\
& -i\sqrt{-\tanh^2(c+dx)} \coth(c+dx) \left( \frac{i \tanh^2(c+dx)}{2d} - \int \tan(ic+idx) dx \right) \\
& \downarrow 3956 \\
& -i\sqrt{-\tanh^2(c+dx)} \coth(c+dx) \left( \frac{i \tanh^2(c+dx)}{2d} - \frac{i \log(\cosh(c+dx))}{d} \right)
\end{aligned}$$

input `Int[(-Tanh[c + d*x]^2)^(3/2), x]`

output `(-I)*Coth[c + d*x]*Sqrt[-Tanh[c + d*x]^2]*(((I)*Log[Cosh[c + d*x]])/d + (I/2)*Tanh[c + d*x]^2)/d)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n-1)/(d*(n-1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

### Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.88

method	result
derivativedivides	$-\frac{\left(-\tanh(dx+c)^2\right)^{\frac{3}{2}}\left(\tanh(dx+c)^2+\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1)\right)}{2d \tanh(dx+c)^3}$
default	$-\frac{\left(-\tanh(dx+c)^2\right)^{\frac{3}{2}}\left(\tanh(dx+c)^2+\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1)\right)}{2d \tanh(dx+c)^3}$
risch	$-\frac{\sqrt{\frac{\left(e^{2dx+2c}-1\right)^2}{\left(e^{2dx+2c}+1\right)^2}}\left(-e^{4dx+4c}dx+e^{4dx+4c}\ln\left(e^{2dx+2c}+1\right)-2e^{4dx+4c}c-2e^{2dx+2c}dx+2e^{2dx+2c}\ln\left(e^{2dx+2c}+1\right)-4\right)}{\left(e^{2dx+2c}-1\right)\left(e^{2dx+2c}+1\right)d}$

input

```
int((-tanh(d*x+c)^2)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
-1/2/d*(-tanh(d*x+c)^2)^(3/2)*(tanh(d*x+c)^2+ln(tanh(d*x+c)-1)+ln(tanh(d*x+c)+1))/tanh(d*x+c)^3
```

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.67

$$\int (-\tanh^2(dx))^{3/2} dx = \frac{i dx e^{(4 dx+4 c)} + i dx - 2(-i dx + i) e^{(2 dx+2 c)} + (-i e^{(4 dx+4 c)} - 2i e^{(2 dx+2 c)} - i) \log(e^{(2 dx+2 c)} + 1)}{d e^{(4 dx+4 c)} + 2 d e^{(2 dx+2 c)} + d}$$

input `integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="fricas")`

output `(I*d*x*e^(4*d*x + 4*c) + I*d*x - 2*(-I*d*x + I)*e^(2*d*x + 2*c) + (-I*e^(4*d*x + 4*c) - 2*I*e^(2*d*x + 2*c) - I)*log(e^(2*d*x + 2*c) + 1))/(d*e^(4*d*x + 4*c) + 2*d*e^(2*d*x + 2*c) + d)`

## Sympy [F]

$$\int (-\tanh^2(c + dx))^{3/2} dx = \int (-\tanh^2(c + dx))^{\frac{3}{2}} dx$$

input `integrate((-tanh(d*x+c)**2)**(3/2),x)`

output `Integral((-tanh(c + d*x)**2)**(3/2), x)`

## Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

$$\int (-\tanh^2(c + dx))^{3/2} dx = \frac{i(dx + c)}{d} + \frac{i \log(e^{-2dx-2c} + 1)}{d} + \frac{2i e^{-2dx-2c}}{d(2e^{-2dx-2c} + e^{-4dx-4c} + 1)}$$

input `integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="maxima")`

output `I*(d*x + c)/d + I*log(e^(-2*d*x - 2*c) + 1)/d + 2*I*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c) + 1))`

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.53

$$\int (-\tanh^2(c + dx))^{3/2} dx = \frac{-i(dx + c)\operatorname{sgn}(-e^{(4dx+4c)} + 1) + i \log(e^{(2dx+2c)} + 1) \operatorname{sgn}(-e^{(4dx+4c)} + 1) + \frac{2i e^{(2dx+2c)} \operatorname{sgn}(-e^{(4dx+4c)} + 1)}{(e^{(2dx+2c)} + 1)^2}}{d}$$

input `integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="giac")`

output `(-I*(d*x + c)*sgn(-e^(4*d*x + 4*c) + 1) + I*log(e^(2*d*x + 2*c) + 1)*sgn(-e^(4*d*x + 4*c) + 1) + 2*I*e^(2*d*x + 2*c)*sgn(-e^(4*d*x + 4*c) + 1)/(e^(2*d*x + 2*c) + 1)^2)/d`

**Mupad [F(-1)]**

Timed out.

$$\int (-\tanh^2(c + dx))^{3/2} dx = \int (-\tanh(c + dx)^2)^{3/2} dx$$

input `int((-tanh(c + d*x)^2)^(3/2),x)`

output `int((-tanh(c + d*x)^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.30

$$\int (-\tanh^2(c + dx))^{3/2} dx = \frac{i(-e^{4dx+4c}\log(e^{2dx+2c} + 1) + e^{4dx+4c}dx + e^{4dx+4c} - 2e^{2dx+2c}\log(e^{2dx+2c} + 1) + 2e^{2dx+2c}dx - \log(e^{4dx+4c} + 2e^{2dx+2c} + 1))}{d(e^{4dx+4c} + 2e^{2dx+2c} + 1)}$$

input `int((-tanh(d*x+c)^2)^(3/2),x)`output `(i*(- e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1) + e**(4*c + 4*d*x)*d*x + e**(4*c + 4*d*x) - 2*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1) + 2*e**(2*c + 2*d*x)*d*x - log(e**(2*c + 2*d*x) + 1) + d*x + 1))/(d*(e**(4*c + 4*d*x) + 2*e**(2*c + 2*d*x) + 1))`

### 3.29 $\int \sqrt{-\tanh^2(c + dx)} dx$

Optimal result	303
Mathematica [A] (verified)	303
Rubi [A] (verified)	304
Maple [A] (verified)	305
Fricas [C] (verification not implemented)	306
Sympy [F]	306
Maxima [C] (verification not implemented)	307
Giac [C] (verification not implemented)	307
Mupad [F(-1)]	308
Reduce [B] (verification not implemented)	308

#### Optimal result

Integrand size = 14, antiderivative size = 31

$$\int \sqrt{-\tanh^2(c + dx)} dx = \frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d}$$

output `coth(d*x+c)*ln(cosh(d*x+c))*(-tanh(d*x+c)^2)^(1/2)/d`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \sqrt{-\tanh^2(c + dx)} dx = \frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d}$$

input `Integrate[Sqrt[-Tanh[c + d*x]^2],x]`

output `(Coth[c + d*x]*Log[Cosh[c + d*x]]*Sqrt[-Tanh[c + d*x]^2])/d`



**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3042, 4141, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{-\tanh^2(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(ic+idx)^2} dx \\
 & \quad \downarrow \text{4141} \\
 & \sqrt{-\tanh^2(c+dx)} \coth(c+dx) \int \tanh(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{-\tanh^2(c+dx)} \coth(c+dx) \int -i \tan(ic+idx) dx \\
 & \quad \downarrow \text{26} \\
 & -i \sqrt{-\tanh^2(c+dx)} \coth(c+dx) \int \tan(ic+idx) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\sqrt{-\tanh^2(c+dx)} \coth(c+dx) \log(\cosh(c+dx))}{d}
 \end{aligned}$$

input `Int[Sqrt[-Tanh[c + d*x]^2],x]`

output `(Coth[c + d*x]*Log[Cosh[c + d*x]]*Sqrt[-Tanh[c + d*x]^2])/d`

## Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)^(n_)])^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.45

method	result
derivativedivides	$-\frac{\sqrt{-\tanh(dx+c)^2 (\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1))}}{2d \tanh(dx+c)}$
default	$-\frac{\sqrt{-\tanh(dx+c)^2 (\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1))}}{2d \tanh(dx+c)}$
risch	$\frac{(e^{2dx+2c}+1)\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}}{e^{2dx+2c}-1} - \frac{2(e^{2dx+2c}+1)\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}}{(e^{2dx+2c}-1)d} + \frac{(e^{2dx+2c}+1)\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}}{(e^{2dx+2c}-1)d}$

input `int((-tanh(d*x+c)^2)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/2/d*(-\tanh(d*x+c)^2)^{(1/2)}*(\ln(\tanh(d*x+c)-1)+\ln(\tanh(d*x+c)+1))/\tanh(d*x+c)$$

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \sqrt{-\tanh^2(c+dx)} dx = \frac{-i dx + i \log(e^{(2dx+2c)} + 1)}{d}$$

input `integrate((-tanh(d*x+c)^2)^(1/2),x, algorithm="fricas")`

output 
$$(-I*d*x + I*\log(e^{(2*d*x + 2*c)} + 1))/d$$

### Sympy [F]

$$\int \sqrt{-\tanh^2(c+dx)} dx = \int \sqrt{-\tanh^2(c+dx)} dx$$

input `integrate((-tanh(d*x+c)**2)**(1/2),x)`

output `Integral(sqrt(-tanh(c + d*x)**2), x)`

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \sqrt{-\tanh^2(c + dx)} dx = -\frac{i(dx + c)}{d} - \frac{i \log(e^{(-2dx-2c)} + 1)}{d}$$

input `integrate((-tanh(d*x+c)^2)^(1/2),x, algorithm="maxima")`

output `-I*(d*x + c)/d - I*log(e^(-2*d*x - 2*c) + 1)/d`

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.74

$$\int \sqrt{-\tanh^2(c + dx)} dx = \frac{i(dx + c)\operatorname{sgn}(-e^{(4dx+4c)} + 1) - i \log(e^{(2dx+2c)} + 1)\operatorname{sgn}(-e^{(4dx+4c)} + 1)}{d}$$

input `integrate((-tanh(d*x+c)^2)^(1/2),x, algorithm="giac")`

output `(I*(d*x + c)*sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) + 1)*sgn(-e^(4*d*x + 4*c) + 1))/d`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{-\tanh^2(c + dx)} dx = \int \sqrt{-\tanh(c + dx)^2} dx$$

input `int((-tanh(c + d*x)^2)^(1/2),x)`output `int((-tanh(c + d*x)^2)^(1/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \sqrt{-\tanh^2(c + dx)} dx = \frac{i(\log(e^{2dx+2c} + 1) - dx)}{d}$$

input `int((-tanh(d*x+c)^2)^(1/2),x)`output `(i*(log(e**(2*c + 2*d*x) + 1) - d*x))/d`

### 3.30 $\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx$

Optimal result . . . . .	309
Mathematica [A] (verified) . . . . .	309
Rubi [C] (verified) . . . . .	310
Maple [A] (verified) . . . . .	311
Fricas [C] (verification not implemented) . . . . .	312
Sympy [F] . . . . .	312
Maxima [C] (verification not implemented) . . . . .	313
Giac [C] (verification not implemented) . . . . .	313
Mupad [B] (verification not implemented) . . . . .	314
Reduce [B] (verification not implemented) . . . . .	314

#### Optimal result

Integrand size = 14, antiderivative size = 31

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

output `ln(sinh(d*x+c))*tanh(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

input `Integrate[1/Sqrt[-Tanh[c + d*x]^2],x]`

output `(Log[Sinh[c + d*x]]*Tanh[c + d*x])/(d*Sqrt[-Tanh[c + d*x]^2])`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3042, 4141, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{\tan(ic+idx)^2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\tanh(c+dx) \int \coth(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh(c+dx) \int -i \tan(ic+idx + \frac{\pi}{2}) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{26} \\
 & -\frac{i \tanh(c+dx) \int \tan(\frac{1}{2}(2ic+\pi) + idx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\tanh(c+dx) \log(-i \sinh(c+dx))}{d\sqrt{-\tanh^2(c+dx)}}
 \end{aligned}$$

input `Int[1/Sqrt[-Tanh[c + d*x]^2],x]`

output `(Log[(-I)*Sinh[c + d*x]]*Tanh[c + d*x])/(d*Sqrt[-Tanh[c + d*x]^2])`

Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)^(n_)])^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.68

method	result	size
derivativedivides	$-\frac{\tanh(dx+c)(\ln(\tanh(dx+c)+1)-2\ln(\tanh(dx+c))+\ln(\tanh(dx+c)-1))}{2d\sqrt{-\tanh(dx+c)^2}}$	52
default	$-\frac{\tanh(dx+c)(\ln(\tanh(dx+c)+1)-2\ln(\tanh(dx+c))+\ln(\tanh(dx+c)-1))}{2d\sqrt{-\tanh(dx+c)^2}}$	52
risch	$\frac{(e^{2dx+2c-1})x}{\sqrt{-\frac{(e^{2dx+2c-1})^2}{(e^{2dx+2c+1})^2}(e^{2dx+2c+1})}} - \frac{2(e^{2dx+2c-1})(dx+c)}{\sqrt{-\frac{(e^{2dx+2c-1})^2}{(e^{2dx+2c+1})^2}(e^{2dx+2c+1})}d} + \frac{(e^{2dx+2c-1})\ln(e^{2dx+2c-1})}{\sqrt{-\frac{(e^{2dx+2c-1})^2}{(e^{2dx+2c+1})^2}(e^{2dx+2c+1})}d}$	192

input `int(1/(-tanh(d*x+c)^2)^(1/2),x,method=_RETURNVERBOSE)`



output 
$$\frac{-1/2/d*\tanh(d*x+c)*(ln(\tanh(d*x+c)+1)-2*ln(\tanh(d*x+c))+ln(\tanh(d*x+c)-1))}{(-\tanh(d*x+c)^2)^{(1/2)}}$$

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{i dx - i \log(e^{(2dx+2c)} - 1)}{d}$$

input `integrate(1/(-tanh(d*x+c)**2)**(1/2),x, algorithm="fricas")`

output `(I*d*x - I*log(e^(2*d*x + 2*c) - 1))/d`

### Sympy [F]

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx$$

input `integrate(1/(-tanh(d*x+c)**2)**(1/2),x)`

output `Integral(1/sqrt(-tanh(c + d*x)**2), x)`

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.45

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{i(dx+c)}{d} + \frac{i \log(e^{(-dx-c)} + 1)}{d} + \frac{i \log(e^{(-dx-c)} - 1)}{d}$$

input `integrate(1/(-tanh(d*x+c)^2)^(1/2),x, algorithm="maxima")`

output `I*(d*x + c)/d + I*log(e^(-d*x - c) + 1)/d + I*log(e^(-d*x - c) - 1)/d`

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.97

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = -\frac{\frac{i dx + i c}{\operatorname{sgn}(-e^{(4 dx + 4 c)} + 1)} - \frac{i \log(e^{(2 dx + 2 c)} - 1)}{\operatorname{sgn}(-e^{(4 dx + 4 c)} + 1)}}{d}$$

input `integrate(1/(-tanh(d*x+c)^2)^(1/2),x, algorithm="giac")`

output `-((I*d*x + I*c)/sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) - 1)/sgn(-e^(4*d*x + 4*c) + 1))/d`

**Mupad [B] (verification not implemented)**

Time = 2.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.77

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{\operatorname{atan}\left(\frac{\tanh(c+dx)}{\sqrt{-\tanh^2(c+dx)^2}}\right)}{d}$$

input `int(1/(-tanh(c + d*x)^2)^(1/2),x)`output `atan(tanh(c + d*x)/(-tanh(c + d*x)^2)^(1/2))/d`**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx = \frac{i(-\log(e^{dx+c} - 1) - \log(e^{dx+c} + 1) + dx)}{d}$$

input `int(1/(-tanh(d*x+c)^2)^(1/2),x)`output `(i*( - log(e**(c + d*x) - 1) - log(e**(c + d*x) + 1) + d*x))/d`

**3.31**  $\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx$

Optimal result	315
Mathematica [A] (verified)	315
Rubi [C] (verified)	316
Maple [A] (verified)	318
Fricas [C] (verification not implemented)	319
Sympy [F]	319
Maxima [C] (verification not implemented)	319
Giac [C] (verification not implemented)	320
Mupad [F(-1)]	320
Reduce [B] (verification not implemented)	321

**Optimal result**

Integrand size = 14, antiderivative size = 60

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = \frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

output `1/2*coth(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)-ln(sinh(d*x+c))*tanh(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)`

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = -\frac{(\operatorname{csch}^2(c+dx) - 2 \log(\sinh(c+dx))) \tanh^3(c+dx)}{2d(-\tanh^2(c+dx))^{3/2}}$$

input `Integrate[(-Tanh[c + d*x]^2)^(-3/2), x]`

output `-1/2*((Csch[c + d*x]^2 - 2*Log[Sinh[c + d*x]])*Tanh[c + d*x]^3)/(d*(-Tanh[c + d*x]^2)^(3/2))`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$ , Rules used = {3042, 4141, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\tan(ic+idx))^{3/2}} dx \\
 & \quad \downarrow \text{4141} \\
 & -\frac{\tanh(c+dx) \int \coth^3(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\tanh(c+dx) \int i \tan(ic+idx + \frac{\pi}{2})^3 dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{26} \\
 & -\frac{i \tanh(c+dx) \int \tan(\frac{1}{2}(2ic+\pi) + idx)^3 dx}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{3954} \\
 & -\frac{i \tanh(c+dx) \left( \frac{i \coth^2(c+dx)}{2d} - \int i \coth(c+dx) dx \right)}{\sqrt{-\tanh^2(c+dx)}} \\
 & \quad \downarrow \text{26} \\
 & -\frac{i \tanh(c+dx) \left( \frac{i \coth^2(c+dx)}{2d} - i \int \coth(c+dx) dx \right)}{\sqrt{-\tanh^2(c+dx)}}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 \frac{i \tanh(c + dx) \left( \frac{i \coth^2(c+dx)}{2d} - i \int -i \tan \left( ic + idx + \frac{\pi}{2} \right) dx \right)}{\sqrt{-\tanh^2(c + dx)}} \\
 \downarrow \text{26} \\
 \frac{i \tanh(c + dx) \left( \frac{i \coth^2(c+dx)}{2d} - \int \tan \left( \frac{1}{2}(2ic + \pi) + idx \right) dx \right)}{\sqrt{-\tanh^2(c + dx)}} \\
 \downarrow \text{3956} \\
 \frac{i \tanh(c + dx) \left( \frac{i \coth^2(c+dx)}{2d} - \frac{i \log(-i \sinh(c+dx))}{d} \right)}{\sqrt{-\tanh^2(c + dx)}}
 \end{array}$$

input `Int[(-Tanh[c + d*x]^2)^(-3/2),x]`

output `((-I)*(((I/2)*Coth[c + d*x]^2)/d - (I*Log[(-I)*Sinh[c + d*x]])/d)*Tanh[c + d*x])/Sqrt[-Tanh[c + d*x]^2]`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

### Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.35

method	result
derivativedivides	$\frac{\tanh(dx+c) \left( 2 \ln(\tanh(dx+c)) \tanh(dx+c)^2 - \ln(\tanh(dx+c)-1) \tanh(dx+c)^2 - \ln(\tanh(dx+c)+1) \tanh(dx+c)^2 - 1 \right)}{2d \left( -\tanh(dx+c)^2 \right)^{\frac{3}{2}}}$
default	$\frac{\tanh(dx+c) \left( 2 \ln(\tanh(dx+c)) \tanh(dx+c)^2 - \ln(\tanh(dx+c)-1) \tanh(dx+c)^2 - \ln(\tanh(dx+c)+1) \tanh(dx+c)^2 - 1 \right)}{2d \left( -\tanh(dx+c)^2 \right)^{\frac{3}{2}}}$
risch	$-\frac{e^{4dx+4c} dx + e^{4dx+4c} \ln(e^{2dx+2c}-1) - 2e^{4dx+4c} c + 2e^{2dx+2c} dx - 2e^{2dx+2c} \ln(e^{2dx+2c}-1) + 4e^{2dx+2c} c - dx - 2e^{2dx+2c}}{(e^{2dx+2c}-1)(e^{2dx+2c}+1) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2} d}}$

input `int(1/(-tanh(d*x+c)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2/d*tanh(d*x+c)*(2*ln(tanh(d*x+c))*tanh(d*x+c)^2-ln(tanh(d*x+c)-1)*tanh(d*x+c)^2-ln(tanh(d*x+c)+1)*tanh(d*x+c)^2-1)/(-tanh(d*x+c)^2)^(3/2)`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.67

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = \frac{-i dx e^{(4dx+4c)} - i dx - 2(-i dx + i)e^{(2dx+2c)} + (i e^{(4dx+4c)} - 2i e^{(2dx+2c)} + d)}{d e^{(4dx+4c)} - 2 d e^{(2dx+2c)} + d}$$

input `integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="fricas")`

output `(-I*d*x*e^(4*d*x + 4*c) - I*d*x - 2*(-I*d*x + I)*e^(2*d*x + 2*c) + (I*e^(4*d*x + 4*c) - 2*I*e^(2*d*x + 2*c) + I)*log(e^(2*d*x + 2*c) - 1))/(d*e^(4*d*x + 4*c) - 2*d*e^(2*d*x + 2*c) + d)`

**Sympy [F]**

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = \int \frac{1}{(-\tanh^2(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(1/(-tanh(d*x+c)**2)**(3/2),x)`

output `Integral((-tanh(c + d*x)**2)**(-3/2), x)`

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.42

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = -\frac{i(dx+c)}{d} - \frac{i \log(e^{(-dx-c)} + 1)}{d} - \frac{i \log(e^{(-dx-c)} - 1)}{d} - \frac{2i e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} - e^{(-4dx-4c)} - 1)}$$



input `integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="maxima")`

output `-I*(d*x + c)/d - I*log(e^(-d*x - c) + 1)/d - I*log(e^(-d*x - c) - 1)/d - 2*I*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) - e^(-4*d*x - 4*c) - 1))`

### Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.67

$$\int \frac{1}{(-\tanh^2(c + dx))^{3/2}} dx = \frac{\frac{i dx + i c}{\operatorname{sgn}(-e^{(4 dx + 4 c) + 1})} - \frac{i \log(e^{(2 dx + 2 c) - 1})}{\operatorname{sgn}(-e^{(4 dx + 4 c) + 1})}}{d} + \frac{2i e^{(2 dx + 2 c)}}{(e^{(2 dx + 2 c) - 1})^2 \operatorname{sgn}(-e^{(4 dx + 4 c) + 1})}$$

input `integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="giac")`

output `((I*d*x + I*c)/sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) - 1)/sgn(-e^(4*d*x + 4*c) + 1) + 2*I*e^(2*d*x + 2*c)/((e^(2*d*x + 2*c) - 1)^2*sgn(-e^(4*d*x + 4*c) + 1)))/d`

### Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(-\tanh^2(c + dx))^{3/2}} dx = \int \frac{1}{(-\tanh(c + dx)^2)^{3/2}} dx$$

input `int(1/(-tanh(c + d*x)^2)^(3/2),x)`

output `int(1/(-tanh(c + d*x)^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 183, normalized size of antiderivative = 3.05

$$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx = \frac{i(e^{4dx+4c}\log(e^{dx+c}-1) + e^{4dx+4c}\log(e^{dx+c}+1) - e^{4dx+4c}dx - e^{4dx+4c} - 2e^{2c+2dx}dx + \log(e^{dx+c}-1) + \log(e^{dx+c}+1) - dx - 1)}{(d(e^{4c+4dx} - 2e^{2c+2dx} + 1))}$$

input `int(1/(-tanh(d*x+c)^2)^(3/2),x)`output `(i*(e**(4*c + 4*d*x))*log(e**(c + d*x) - 1) + e**(4*c + 4*d*x))*log(e**(c + d*x) + 1) - e**(4*c + 4*d*x)*d*x - e**(4*c + 4*d*x) - 2*e**(2*c + 2*d*x)*log(e**(c + d*x) - 1) - 2*e**(2*c + 2*d*x)*log(e**(c + d*x) + 1) + 2*e**(2*c + 2*d*x)*d*x + log(e**(c + d*x) - 1) + log(e**(c + d*x) + 1) - d*x - 1)) / (d*(e**(4*c + 4*d*x) - 2*e**(2*c + 2*d*x) + 1))`

### 3.32 $\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx$

Optimal result	322
Mathematica [A] (verified)	322
Rubi [C] (verified)	323
Maple [A] (verified)	326
Fricas [C] (verification not implemented)	326
Sympy [F]	327
Maxima [C] (verification not implemented)	327
Giac [C] (verification not implemented)	328
Mupad [F(-1)]	328
Reduce [B] (verification not implemented)	329

#### Optimal result

Integrand size = 14, antiderivative size = 88

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx = -\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} + \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

output

`-1/2*coth(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)-1/4*coth(d*x+c)^3/d/(-tanh(d*x+c)^2)^(1/2)+ln(sinh(d*x+c))*tanh(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.65

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx = \frac{(4\operatorname{csch}^2(c+dx) + \operatorname{csch}^4(c+dx) - 4\log(\sinh(c+dx))) \tanh^5(c+dx)}{4d(-\tanh^2(c+dx))^{5/2}}$$

input `Integrate[(-Tanh[c + d*x]^2)^(-5/2), x]`

output `-1/4*((4*Csch[c + d*x]^2 + Csch[c + d*x]^4 - 4*Log[Sinh[c + d*x]])*Tanh[c + d*x]^5)/(d*(-Tanh[c + d*x]^2)^(5/2))`

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88, number of steps used = 13, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$ , Rules used = {3042, 4141, 3042, 26, 3954, 26, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-\tanh^2(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(\tan(ic + idx)^2)^{5/2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\tanh(c + dx) \int \coth^5(c + dx) dx}{\sqrt{-\tanh^2(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh(c + dx) \int -i \tan\left(ic + idx + \frac{\pi}{2}\right)^5 dx}{\sqrt{-\tanh^2(c + dx)}} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \tanh(c + dx) \int \tan\left(\frac{1}{2}(2ic + \pi) + idx\right)^5 dx}{\sqrt{-\tanh^2(c + dx)}} \\
 & \quad \downarrow \text{3954}
 \end{aligned}$$

$$\begin{aligned}
& \frac{i \tanh(c+dx) \left( -\int -i \coth^3(c+dx) dx - \frac{i \coth^4(c+dx)}{4d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 26 \\
& \frac{i \tanh(c+dx) \left( i \int \coth^3(c+dx) dx - \frac{i \coth^4(c+dx)}{4d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{i \tanh(c+dx) \left( i \int i \tan \left( ic + idx + \frac{\pi}{2} \right)^3 dx - \frac{i \coth^4(c+dx)}{4d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 26 \\
& \frac{i \tanh(c+dx) \left( -\int \tan \left( \frac{1}{2}(2ic + \pi) + idx \right)^3 dx - \frac{i \coth^4(c+dx)}{4d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 3954 \\
& \frac{i \tanh(c+dx) \left( \int i \coth(c+dx) dx - \frac{i \coth^4(c+dx)}{4d} - \frac{i \coth^2(c+dx)}{2d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 26 \\
& \frac{i \tanh(c+dx) \left( i \int \coth(c+dx) dx - \frac{i \coth^4(c+dx)}{4d} - \frac{i \coth^2(c+dx)}{2d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{i \tanh(c+dx) \left( i \int -i \tan \left( ic + idx + \frac{\pi}{2} \right) dx - \frac{i \coth^4(c+dx)}{4d} - \frac{i \coth^2(c+dx)}{2d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 26 \\
& \frac{i \tanh(c+dx) \left( \int \tan \left( \frac{1}{2}(2ic + \pi) + idx \right) dx - \frac{i \coth^4(c+dx)}{4d} - \frac{i \coth^2(c+dx)}{2d} \right)}{\sqrt{-\tanh^2(c+dx)}} \\
& \quad \downarrow 3956
\end{aligned}$$

$$\frac{i \tanh(c + dx) \left( -\frac{i \coth^4(c+dx)}{4d} - \frac{i \coth^2(c+dx)}{2d} + \frac{i \log(-i \sinh(c+dx))}{d} \right)}{\sqrt{-\tanh^2(c + dx)}}$$

input `Int[(-Tanh[c + d*x]^2)^(-5/2),x]`

output `((-I)*(((1/2*I)*Coth[c + d*x]^2)/d - ((I/4)*Coth[c + d*x]^4)/d + (I*Log[(-I)*Sinh[c + d*x]])/d)*Tanh[c + d*x])/Sqrt[-Tanh[c + d*x]^2]`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.03

method	result
derivativedivides	$-\frac{\tanh(dx+c)\left(2\ln(\tanh(dx+c)+1)\tanh(dx+c)^4-4\ln(\tanh(dx+c))\tanh(dx+c)^4+2\ln(\tanh(dx+c)-1)\tanh(dx+c)\right)}{4d\left(-\tanh(dx+c)^2\right)^{\frac{5}{2}}}$
default	$-\frac{\tanh(dx+c)\left(2\ln(\tanh(dx+c)+1)\tanh(dx+c)^4-4\ln(\tanh(dx+c))\tanh(dx+c)^4+2\ln(\tanh(dx+c)-1)\tanh(dx+c)\right)}{4d\left(-\tanh(dx+c)^2\right)^{\frac{5}{2}}}$
risch	$\frac{(e^{2dx+2c}-1)x}{\sqrt{\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}(e^{2dx+2c}+1)}} - \frac{2(e^{2dx+2c}-1)(dx+c)}{\sqrt{\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}(e^{2dx+2c}+1)d}} - \frac{4e^{2dx+2c}(e^{4dx+4c}-e^{2dx+2c}+1)}{(e^{2dx+2c}-1)^3(e^{2dx+2c}+1)}\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(e^{2dx+2c}+1)^2}}$

input `int(1/(-tanh(d*x+c)^2)^(5/2),x,method=_RETURNVERBOSE)`

output `-1/4/d*tanh(d*x+c)*(2*ln(tanh(d*x+c)+1)*tanh(d*x+c)^4-4*ln(tanh(d*x+c))*tanh(d*x+c)^4+2*ln(tanh(d*x+c)-1)*tanh(d*x+c)^4+2*tanh(d*x+c)^2+1)/(-tanh(d*x+c)^2)^(5/2)`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.05

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx = \frac{i dx e^{(8dx+8c)} + i dx - 4(i dx - i)e^{(6dx+6c)} - 2(-3i dx + 2i)e^{(4dx+4c)} - 4(i dx - i)e^{(2dx+2c)}}{de^{(8dx+8c)} - 4de^{(6dx+6c)} + 6de^{(4dx+4c)} - 4de^{(2dx+2c)} + d}$$

input `integrate(1/(-tanh(d*x+c)^2)^(5/2),x, algorithm="fricas")`

output `(I*d*x*e^(8*d*x + 8*c) + I*d*x - 4*(I*d*x - I)*e^(6*d*x + 6*c) - 2*(-3*I*d*x + 2*I)*e^(4*d*x + 4*c) - 4*(I*d*x - I)*e^(2*d*x + 2*c) + (-I*e^(8*d*x + 8*c) + 4*I*e^(6*d*x + 6*c) - 6*I*e^(4*d*x + 4*c) + 4*I*e^(2*d*x + 2*c) - I)*log(e^(2*d*x + 2*c) - 1)/(d*e^(8*d*x + 8*c) - 4*d*e^(6*d*x + 6*c) + 6*d*e^(4*d*x + 4*c) - 4*d*e^(2*d*x + 2*c) + d)`

**Sympy [F]**

$$\int \frac{1}{(-\tanh^2(c + dx))^{5/2}} dx = \int \frac{1}{(-\tanh^2(c + dx))^{\frac{5}{2}}} dx$$

input `integrate(1/(-tanh(d*x+c)**2)**(5/2), x)`

output `Integral((-tanh(c + d*x)**2)**(-5/2), x)`

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.50

$$\int \frac{1}{(-\tanh^2(c + dx))^{5/2}} dx = \frac{i(dx + c)}{d} + \frac{i \log(e^{(-dx-c)} + 1)}{d} + \frac{i \log(e^{(-dx-c)} - 1)}{d} - \frac{4(-i e^{(-2dx-2c)} + i e^{(-4dx-4c)} - i e^{(-6dx-6c)})}{d(4e^{(-2dx-2c)} - 6e^{(-4dx-4c)} + 4e^{(-6dx-6c)} - e^{(-8dx-8c)} - 1)}$$

input `integrate(1/(-tanh(d*x+c)^2)^(5/2), x, algorithm="maxima")`

output `I*(d*x + c)/d + I*log(e^(-d*x - c) + 1)/d + I*log(e^(-d*x - c) - 1)/d - 4*(-I*e^(-2*d*x - 2*c) + I*e^(-4*d*x - 4*c) - I*e^(-6*d*x - 6*c))/(d*(4*e^(-2*d*x - 2*c) - 6*e^(-4*d*x - 4*c) + 4*e^(-6*d*x - 6*c) - e^(-8*d*x - 8*c) - 1))`



**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.43

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx =$$

$$\frac{\frac{i dx + ic}{\operatorname{sgn}(-e^{(4dx+4c)+1})} - \frac{i \log(e^{(2dx+2c)-1})}{\operatorname{sgn}(-e^{(4dx+4c)+1})} + \frac{4(i e^{(6dx+6c)} - i e^{(4dx+4c)} + i e^{(2dx+2c)})}{(e^{(2dx+2c)-1})^4 \operatorname{sgn}(-e^{(4dx+4c)+1})}}{d}$$

input `integrate(1/(-tanh(d*x+c)^2)^(5/2),x, algorithm="giac")`

output `-((I*d*x + I*c)/sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) - 1)/sgn(-e^(4*d*x + 4*c) + 1) + 4*(I*e^(6*d*x + 6*c) - I*e^(4*d*x + 4*c) + I*e^(2*d*x + 2*c))/(e^(2*d*x + 2*c) - 1)^4*sgn(-e^(4*d*x + 4*c) + 1))/d`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx = \int \frac{1}{(-\tanh(c+dx)^2)^{5/2}} dx$$

input `int(1/(-tanh(c + d*x)^2)^(5/2),x)`

output `int(1/(-tanh(c + d*x)^2)^(5/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 337, normalized size of antiderivative = 3.83

$$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx = \frac{i(-e^{8dx+8c}\log(e^{dx+c}-1) - e^{8dx+8c}\log(e^{dx+c}+1) + e^{8dx+8c}dx + e^{8dx+8c} + 4$$

input `int(1/(-tanh(d*x+c)^2)^(5/2),x)`

output

```
(i*( - e**(8*c + 8*d*x)*log(e**(c + d*x) - 1) - e**(8*c + 8*d*x)*log(e**(c + d*x) + 1) + e**(8*c + 8*d*x)*d*x + e**(8*c + 8*d*x) + 4*e**(6*c + 6*d*x)*log(e**(c + d*x) - 1) + 4*e**(6*c + 6*d*x)*log(e**(c + d*x) + 1) - 4*e**(6*c + 6*d*x)*d*x - 6*e**(4*c + 4*d*x)*log(e**(c + d*x) - 1) - 6*e**(4*c + 4*d*x)*log(e**(c + d*x) + 1) + 6*e**(4*c + 4*d*x)*d*x + 2*e**(4*c + 4*d*x) + 4*e**(2*c + 2*d*x)*log(e**(c + d*x) - 1) + 4*e**(2*c + 2*d*x)*log(e**(c + d*x) + 1) - 4*e**(2*c + 2*d*x)*d*x - log(e**(c + d*x) - 1) - log(e**(c + d*x) + 1) + d*x + 1))/(d*(e**(8*c + 8*d*x) - 4*e**(6*c + 6*d*x) + 6*e**(4*c + 4*d*x) - 4*e**(2*c + 2*d*x) + 1))
```

### 3.33 $\int \sqrt{\tanh^3(x)} dx$

Optimal result . . . . .	330
Mathematica [A] (verified) . . . . .	330
Rubi [A] (verified) . . . . .	331
Maple [A] (verified) . . . . .	334
Fricas [B] (verification not implemented) . . . . .	334
Sympy [F] . . . . .	335
Maxima [F] . . . . .	336
Giac [A] (verification not implemented) . . . . .	336
Mupad [F(-1)] . . . . .	336
Reduce [F] . . . . .	337

#### Optimal result

Integrand size = 8, antiderivative size = 57

$$\int \sqrt{\tanh^3(x)} dx = -2 \coth(x) \sqrt{\tanh^3(x)} + \frac{\arctan\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

output

```
-2*coth(x)*(tanh(x)^3)^(1/2)+arctan(tanh(x)^(1/2))*(tanh(x)^3)^(1/2)/tanh(x)^(3/2)+arctanh(tanh(x)^(1/2))*(tanh(x)^3)^(1/2)/tanh(x)^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.67

$$\int \sqrt{\tanh^3(x)} dx = \frac{\left(\arctan\left(\sqrt{\tanh(x)}\right) + \operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) - 2\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

input

```
Integrate[Sqrt[Tanh[x]^3],x]
```

output

$$\left( \left( \text{ArcTan}[\text{Sqrt}[\text{Tanh}[x]]] + \text{ArcTanh}[\text{Sqrt}[\text{Tanh}[x]]] - 2*\text{Sqrt}[\text{Tanh}[x]] \right) * \text{Sqrt}[\text{Tanh}[x]^3] \right) / \text{Tanh}[x]^{(3/2)}$$
**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.86, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\tanh^3(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{i \tan(ix)^3} dx \\ & \quad \downarrow \text{4141} \\ & \frac{\sqrt{\tanh^3(x)} \int \tanh^{\frac{3}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\ & \quad \downarrow \text{3042} \\ & \frac{\sqrt{\tanh^3(x)} \int (-i \tan(ix))^{3/2} dx}{\tanh^{\frac{3}{2}}(x)} \\ & \quad \downarrow \text{3954} \\ & \frac{\sqrt{\tanh^3(x)} \left( \int \frac{1}{\sqrt{\tanh(x)}} dx - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\ & \quad \downarrow \text{3042} \\ & \frac{\sqrt{\tanh^3(x)} \left( -2\sqrt{\tanh(x)} + \int \frac{1}{\sqrt{-i \tan(ix)}} dx \right)}{\tanh^{\frac{3}{2}}(x)} \\ & \quad \downarrow \text{3957} \end{aligned}$$

$$\frac{\sqrt{\tanh^3(x)} \left( -\int -\frac{1}{\sqrt{\tanh(x)(1-\tanh^2(x))}} d\tanh(x) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 25

$$\frac{\sqrt{\tanh^3(x)} \left( \int \frac{1}{\sqrt{\tanh(x)(1-\tanh^2(x))}} d\tanh(x) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 266

$$\frac{\sqrt{\tanh^3(x)} \left( 2 \int \frac{1}{1-\tanh^2(x)} d\sqrt{\tanh(x)} - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 756

$$\frac{\sqrt{\tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} + \frac{1}{2} \int \frac{1}{\tanh(x)+1} d\sqrt{\tanh(x)} \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 216

$$\frac{\sqrt{\tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} + \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 219

$$\frac{\sqrt{\tanh^3(x)} \left( 2 \left( \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) + \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(x)} \right) \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}$$

input `Int [Sqrt [Tanh [x] ^3] , x]`

output `((2*(ArcTan [Sqrt [Tanh [x]]] /2 + ArcTanh [Sqrt [Tanh [x]]] /2) - 2*Sqrt [Tanh [x]] ) *Sqrt [Tanh [x] ^3] ) /Tanh [x] ^ (3/2)`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 219  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 266  $\text{Int}[(\text{c}_) * (\text{x}_)]^{(\text{m}_)} * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{k} * (\text{m} + 1) - 1)} * (\text{a} + \text{b} * (\text{x}^{(2 * \text{k})}/\text{c}^2))^{\text{p}}, \text{x}], \text{x}, (\text{c} * \text{x})^{(1/\text{k})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \&\& \text{FractionQ}[\text{m}] \&\& \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 756  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^4)^{-1}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{r}/(2 * \text{a}) \quad \text{Int}[1/(\text{r} - \text{s} * \text{x}^2), \text{x}], \text{x}] + \text{Simp}[\text{r}/(2 * \text{a}) \quad \text{Int}[1/(\text{r} + \text{s} * \text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{!GtQ}[\text{a}/\text{b}, 0]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3954  $\text{Int}[(\text{b}_) * \tan[(\text{c}_) + (\text{d}_) * (\text{x}_)]]^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{b} * ((\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} - 1)}/(\text{d} * (\text{n} - 1))), \text{x}] - \text{Simp}[\text{b}^2 \quad \text{Int}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} - 2)}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{GtQ}[\text{n}, 1]$
- rule 3957  $\text{Int}[(\text{b}_) * \tan[(\text{c}_) + (\text{d}_) * (\text{x}_)]]^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{b}/\text{d} \quad \text{Subst}[\text{Int}[\text{x}^{\text{n}}/(\text{b}^2 + \text{x}^2), \text{x}], \text{x}, \text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}, \text{n}\}, \text{x}] \&\& \text{!IntegerQ}[\text{n}]$

rule 4141

```

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :> With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

```

**Maple [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$-\frac{\sqrt{\tanh(x)^3} \left( 4\sqrt{\tanh(x)} + \ln(\sqrt{\tanh(x)} - 1) - \ln(\sqrt{\tanh(x)} + 1) - 2 \arctan(\sqrt{\tanh(x)}) \right)}{2 \tanh(x)^{\frac{3}{2}}}$	43
default	$-\frac{\sqrt{\tanh(x)^3} \left( 4\sqrt{\tanh(x)} + \ln(\sqrt{\tanh(x)} - 1) - \ln(\sqrt{\tanh(x)} + 1) - 2 \arctan(\sqrt{\tanh(x)}) \right)}{2 \tanh(x)^{\frac{3}{2}}}$	43

input

```
int((tanh(x)^3)^(1/2), x, method=_RETURNVERBOSE)
```

output

```

-1/2*(tanh(x)^3)^(1/2)*(4*tanh(x)^(1/2)+ln(tanh(x)^(1/2)-1)-ln(tanh(x)^(1/
2)+1)-2*arctan(tanh(x)^(1/2)))/tanh(x)^(3/2)

```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 106 vs.  $2(43) = 86$ .

Time = 0.09 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.86

$$\int \sqrt{\tanh^3(x)} dx = -2 \sqrt{\frac{\sinh(x)}{\cosh(x)}} + \arctan \left( -\cosh(x)^2 - 2 \cosh(x) \sinh(x) - \sinh(x)^2 \right. \\ \left. + (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{\frac{\sinh(x)}{\cosh(x)}} \right) \\ - \frac{1}{2} \log \left( -\cosh(x)^2 - 2 \cosh(x) \sinh(x) - \sinh(x)^2 \right. \\ \left. + (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{\frac{\sinh(x)}{\cosh(x)}} \right)$$

input `integrate((tanh(x)^3)^(1/2),x, algorithm="fricas")`

output `-2*sqrt(sinh(x)/cosh(x)) + arctan(-cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(sinh(x)/cosh(x))) - 1/2*log(-cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(sinh(x)/cosh(x)))`

## Sympy [F]

$$\int \sqrt{\tanh^3(x)} dx = \int \sqrt{\tanh^3(x)} dx$$

input `integrate((tanh(x)**3)**(1/2),x)`

output `Integral(sqrt(tanh(x)**3), x)`



**Maxima [F]**

$$\int \sqrt{\tanh^3(x)} dx = \int \sqrt{\tanh(x)^3} dx$$

input `integrate((tanh(x)^3)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(tanh(x)^3), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.96

$$\int \sqrt{\tanh^3(x)} dx = \frac{4}{\sqrt{e^{(4x)} - 1} - e^{(2x)} - 1} + \arctan\left(\sqrt{e^{(4x)} - 1} - e^{(2x)}\right) - \frac{1}{2} \log\left(-\sqrt{e^{(4x)} - 1} + e^{(2x)}\right)$$

input `integrate((tanh(x)^3)^(1/2),x, algorithm="giac")`

output `4/(sqrt(e^(4*x) - 1) - e^(2*x) - 1) + arctan(sqrt(e^(4*x) - 1) - e^(2*x)) - 1/2*log(-sqrt(e^(4*x) - 1) + e^(2*x))`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{\tanh^3(x)} dx = \int \sqrt{\tanh(x)^3} dx$$

input `int((tanh(x)^3)^(1/2),x)`

output `int((tanh(x)^3)^(1/2), x)`

**Reduce [F]**

$$\int \sqrt{\tanh^3(x)} dx = -2\sqrt{\tanh(x)} + \int \frac{\sqrt{\tanh(x)}}{\tanh(x)} dx$$

input `int((tanh(x)^3)^(1/2),x)`

output `- 2*sqrt(tanh(x)) + int(sqrt(tanh(x))/tanh(x),x)`

### 3.34 $\int (a \tanh^3(x))^{3/2} dx$

Optimal result	338
Mathematica [A] (verified)	338
Rubi [A] (verified)	339
Maple [A] (verified)	342
Fricas [B] (verification not implemented)	343
Sympy [F]	344
Maxima [F]	344
Giac [B] (verification not implemented)	344
Mupad [F(-1)]	345
Reduce [F]	345

#### Optimal result

Integrand size = 10, antiderivative size = 86

$$\int (a \tanh^3(x))^{3/2} dx = -\frac{2}{3}a\sqrt{a \tanh^3(x)} - \frac{a \arctan\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{a \operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} - \frac{2}{7}a \tanh^2(x) \sqrt{a \tanh^3(x)}$$

output

```
-2/3*a*(a*tanh(x)^3)^(1/2)-a*arctan(tanh(x)^(1/2))*(a*tanh(x)^3)^(1/2)/tanh(x)^(3/2)+a*arctanh(tanh(x)^(1/2))*(a*tanh(x)^3)^(1/2)/tanh(x)^(3/2)-2/7*a*tanh(x)^2*(a*tanh(x)^3)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.64

$$\int (a \tanh^3(x))^{3/2} dx = \frac{(a \tanh^3(x))^{3/2} \left( 21 \arctan\left(\sqrt{\tanh(x)}\right) - 21 \operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) + 14 \tanh^{\frac{3}{2}}(x) + 6 \tanh^{\frac{7}{2}}(x) \right)}{21 \tanh^{\frac{9}{2}}(x)}$$

input `Integrate[(a*Tanh[x]^3)^(3/2),x]`

output `-1/21*((a*Tanh[x]^3)^(3/2)*(21*ArcTan[Sqrt[Tanh[x]]] - 21*ArcTanh[Sqrt[Tanh[x]]] + 14*Tanh[x]^(3/2) + 6*Tanh[x]^(7/2)))/Tanh[x]^(9/2)`

### Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.74, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3954, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh^3(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (ia \tan(ix)^3)^{3/2} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{a \sqrt{a \tanh^3(x)} \int \tanh^{\frac{9}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt{a \tanh^3(x)} \int (-i \tan(ix))^{9/2} dx}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3954} \\
 & \frac{a \sqrt{a \tanh^3(x)} \left( \int \tanh^{\frac{5}{2}}(x) dx - \frac{2}{7} \tanh^{\frac{7}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt{a \tanh^3(x)} \left( -\frac{2}{7} \tanh^{\frac{7}{2}}(x) + \int (-i \tan(ix))^{5/2} dx \right)}{\tanh^{\frac{3}{2}}(x)}
 \end{aligned}$$

↓ 3954

$$\frac{a\sqrt{a \tanh^3(x)} \left( \int \sqrt{\tanh(x)} dx - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 3042

$$\frac{a\sqrt{a \tanh^3(x)} \left( \int \sqrt{-i \tan(ix)} dx - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 3957

$$\frac{a\sqrt{a \tanh^3(x)} \left( -\int -\frac{\sqrt{\tanh(x)}}{1-\tanh^2(x)} d \tanh(x) - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 25

$$\frac{a\sqrt{a \tanh^3(x)} \left( \int \frac{\sqrt{\tanh(x)}}{1-\tanh^2(x)} d \tanh(x) - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 266

$$\frac{a\sqrt{a \tanh^3(x)} \left( 2 \int \frac{\tanh(x)}{1-\tanh^2(x)} d\sqrt{\tanh(x)} - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 827

$$\frac{a\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} - \frac{1}{2} \int \frac{1}{\tanh(x)+1} d\sqrt{\tanh(x)} \right) - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 216

$$\frac{a\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} - \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

↓ 219

$$\frac{a\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(x)} \right) - \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - \frac{2}{7} \tanh^{\frac{7}{2}}(x) - \frac{2}{3} \tanh^{\frac{3}{2}}(x) \right)}{\tanh^{\frac{3}{2}}(x)}$$

input `Int[(a*Tanh[x]^3)^(3/2),x]`

output `(a*Sqrt[a*Tanh[x]^3]*(2*(-1/2*ArcTan[Sqrt[Tanh[x]]] + ArcTanh[Sqrt[Tanh[x]]])/2) - (2*Tanh[x]^(3/2))/3 - (2*Tanh[x]^(7/2)/7))/Tanh[x]^(3/2)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

### Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.88

method	result	si
derivativedivides	$\frac{(a \tanh(x)^3)^{\frac{3}{2}} \left( 21a^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) - 21a^{\frac{7}{2}} \operatorname{arctan}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) - 6(a \tanh(x))^{\frac{7}{2}} - 14a^2(a \tanh(x))^{\frac{3}{2}} \right)}{21 \tanh(x)^3 (a \tanh(x))^{\frac{3}{2}} a^2}$	7
default	$\frac{(a \tanh(x)^3)^{\frac{3}{2}} \left( 21a^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) - 21a^{\frac{7}{2}} \operatorname{arctan}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) - 6(a \tanh(x))^{\frac{7}{2}} - 14a^2(a \tanh(x))^{\frac{3}{2}} \right)}{21 \tanh(x)^3 (a \tanh(x))^{\frac{3}{2}} a^2}$	7

input `int((a*tanh(x)^3)^(3/2),x,method=_RETURNVERBOSE)`

output `1/21*(a*tanh(x)^3)^(3/2)*(21*a^(7/2)*arctanh((a*tanh(x))^(1/2)/a^(1/2))-21*a^(7/2)*arctan((a*tanh(x))^(1/2)/a^(1/2))-6*(a*tanh(x))^(7/2)-14*a^2*(a*tanh(x))^(3/2))/tanh(x)^3/(a*tanh(x))^(3/2)/a^2`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 615 vs.  $2(66) = 132$ .

Time = 0.12 (sec) , antiderivative size = 1267, normalized size of antiderivative = 14.73

$$\int (a \tanh^3(x))^{3/2} dx = \text{Too large to display}$$

input `integrate((a*tanh(x)^3)^(3/2),x, algorithm="fricas")`

output

```
[-1/84*(42*(a*cosh(x)^6 + 6*a*cosh(x)*sinh(x)^5 + a*sinh(x)^6 + 3*a*cosh(x)^4 + 3*(5*a*cosh(x)^2 + a)*sinh(x)^4 + 4*(5*a*cosh(x)^3 + 3*a*cosh(x))*sinh(x)^3 + 3*a*cosh(x)^2 + 3*(5*a*cosh(x)^4 + 6*a*cosh(x)^2 + a)*sinh(x)^2 + 6*(a*cosh(x)^5 + 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) + a)*sqrt(-a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2)) - 21*(a*cosh(x)^6 + 6*a*cosh(x)*sinh(x)^5 + a*sinh(x)^6 + 3*a*cosh(x)^4 + 3*(5*a*cosh(x)^2 + a)*sinh(x)^4 + 4*(5*a*cosh(x)^3 + 3*a*cosh(x))*sinh(x)^3 + 3*a*cosh(x)^2 + 3*(5*a*cosh(x)^4 + 6*a*cosh(x)^2 + a)*sinh(x)^2 + 6*(a*cosh(x)^5 + 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) + a)*sqrt(-a)*log(-(a*cosh(x)^4 + 4*a*cosh(x)^3*sinh(x) + 6*a*cosh(x)^2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 - 2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x)) - 2*a)/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + 16*(5*a*cosh(x)^6 + 30*a*cosh(x)*sinh(x)^5 + 5*a*sinh(x)^6 - a*cosh(x)^4 + (75*a*cosh(x)^2 - a)*sinh(x)^4 + 4*(25*a*cosh(x)^3 - a*cosh(x))*sinh(x)^3 + a*cosh(x)^2 + (75*a*cosh(x)^4 - 6*a*cosh(x)^2 + a)*sinh(x)^2 + 2*(15*a*cosh(x)^5 - 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) - 5*a)*sqrt(a*sinh(x)/cosh(x)))/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(5*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 + 6*cosh(x)^2 + 1)*sin...
```



**Sympy [F]**

$$\int (a \tanh^3(x))^{3/2} dx = \int (a \tanh^3(x))^{\frac{3}{2}} dx$$

input `integrate((a*tanh(x)**3)**(3/2),x)`

output `Integral((a*tanh(x)**3)**(3/2), x)`

**Maxima [F]**

$$\int (a \tanh^3(x))^{3/2} dx = \int (a \tanh^3(x))^{\frac{3}{2}} dx$$

input `integrate((a*tanh(x)^3)^(3/2),x, algorithm="maxima")`

output `integrate((a*tanh(x)^3)^(3/2), x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 342 vs.  $2(66) = 132$ .

Time = 0.20 (sec) , antiderivative size = 342, normalized size of antiderivative = 3.98

$$\int (a \tanh^3(x))^{3/2} dx =$$

$$-\frac{1}{42} \left( 42 \sqrt{a} \arctan \left( -\frac{\sqrt{a}e^{(2x)} - \sqrt{ae^{(4x)} - a}}{\sqrt{a}} \right) \operatorname{sgn}(e^{(4x)} - 1) + 21 \sqrt{a} \log \left( \left| -\sqrt{a}e^{(2x)} + \sqrt{ae^{(4x)} - a} \right| \right) \right)$$

input `integrate((a*tanh(x)^3)^(3/2),x, algorithm="giac")`

output

```
-1/42*(42*sqrt(a)*arctan(-(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))/sqrt(a))
*sgn(e^(4*x) - 1) + 21*sqrt(a)*log(abs(-sqrt(a)*e^(2*x) + sqrt(a*e^(4*x) -
a)))*sgn(e^(4*x) - 1) + 16*(21*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))^6*
a*sgn(e^(4*x) - 1) + 42*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))^5*a^(3/2)*
sgn(e^(4*x) - 1) + 119*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))^4*a^2*sgn(e
^(4*x) - 1) + 56*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))^3*a^(5/2)*sgn(e^(
4*x) - 1) + 63*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))^2*a^3*sgn(e^(4*x) -
1) + 14*(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))*a^(7/2)*sgn(e^(4*x) - 1)
+ 5*a^4*sgn(e^(4*x) - 1))/(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a) + sqrt(a)
)^7)*a
```

**Mupad [F(-1)]**

Timed out.

$$\int (a \tanh^3(x))^{3/2} dx = \int (a \tanh(x)^3)^{3/2} dx$$

input

```
int((a*tanh(x)^3)^(3/2),x)
```

output

```
int((a*tanh(x)^3)^(3/2), x)
```

**Reduce [F]**

$$\int (a \tanh^3(x))^{3/2} dx = \sqrt{a} \left( \int \sqrt{\tanh(x)} \tanh(x)^4 dx \right) a$$

input

```
int((a*tanh(x)^3)^(3/2),x)
```

output

```
sqrt(a)*int(sqrt(tanh(x))*tanh(x)**4,x)*a
```

### 3.35 $\int \sqrt{a \tanh^3(x)} dx$

Optimal result	346
Mathematica [A] (verified)	346
Rubi [A] (verified)	347
Maple [A] (verified)	350
Fricas [B] (verification not implemented)	350
Sympy [F]	351
Maxima [F]	351
Giac [B] (verification not implemented)	352
Mupad [F(-1)]	352
Reduce [F]	353

#### Optimal result

Integrand size = 10, antiderivative size = 63

$$\int \sqrt{a \tanh^3(x)} dx = -2 \operatorname{coth}(x) \sqrt{a \tanh^3(x)} + \frac{\arctan\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

output

$-2*\operatorname{coth}(x)*(a*\tanh(x)^3)^{(1/2)}+\arctan(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}+\operatorname{arctanh}(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.63

$$\int \sqrt{a \tanh^3(x)} dx = \frac{\left(\arctan\left(\sqrt{\tanh(x)}\right) + \operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) - 2\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

input

`Integrate[Sqrt[a*Tanh[x]^3], x]`

output

```
((ArcTan[Sqrt[Tanh[x]]] + ArcTanh[Sqrt[Tanh[x]]] - 2*Sqrt[Tanh[x]])*Sqrt[a
*Tanh[x]^3])/Tanh[x]^(3/2)
```

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.81, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \tanh^3(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{ia \tan(ix)^3} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\sqrt{a \tanh^3(x)} \int \tanh^{\frac{3}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a \tanh^3(x)} \int (-i \tan(ix))^{3/2} dx}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3954} \\
 & \frac{\sqrt{a \tanh^3(x)} \left( \int \frac{1}{\sqrt{\tanh(x)}} dx - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a \tanh^3(x)} \left( -2\sqrt{\tanh(x)} + \int \frac{1}{\sqrt{-i \tan(ix)}} dx \right)}{\tanh^{\frac{3}{2}}(x)} \\
 & \quad \downarrow \text{3957}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{a \tanh^3(x)} \left( -\int -\frac{1}{\sqrt{\tanh(x)(1-\tanh^2(x))}} d \tanh(x) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{25} \\
& \frac{\sqrt{a \tanh^3(x)} \left( \int \frac{1}{\sqrt{\tanh(x)(1-\tanh^2(x))}} d \tanh(x) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{266} \\
& \frac{\sqrt{a \tanh^3(x)} \left( 2 \int \frac{1}{1-\tanh^2(x)} d\sqrt{\tanh(x)} - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{756} \\
& \frac{\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} + \frac{1}{2} \int \frac{1}{\tanh(x)+1} d\sqrt{\tanh(x)} \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{216} \\
& \frac{\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)} + \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{219} \\
& \frac{\sqrt{a \tanh^3(x)} \left( 2 \left( \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) + \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(x)} \right) \right) - 2\sqrt{\tanh(x)} \right)}{\tanh^{\frac{3}{2}}(x)}
\end{aligned}$$

input `Int [Sqrt [a*Tanh [x] ^3] , x]`

output `((2*(ArcTan [Sqrt [Tanh [x]]]/2 + ArcTanh [Sqrt [Tanh [x]]]/2) - 2*Sqrt [Tanh [x]])*Sqrt [a*Tanh [x] ^3])/Tanh [x]^(3/2)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`
- rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4141

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

**Maple [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.98

method	result	size
derivativedivides	$-\frac{\sqrt{a \tanh(x)^3} \left( 2\sqrt{a \tanh(x)} - \sqrt{a} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) - \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) \right)}{\tanh(x) \sqrt{a \tanh(x)}}$	62
default	$-\frac{\sqrt{a \tanh(x)^3} \left( 2\sqrt{a \tanh(x)} - \sqrt{a} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) - \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) \right)}{\tanh(x) \sqrt{a \tanh(x)}}$	62

input

```
int((a*tanh(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-(a*tanh(x)^3)^(1/2)*(2*(a*tanh(x))^(1/2)-a^(1/2)*arctan((a*tanh(x))^(1/2)
/a^(1/2))-a^(1/2)*arctanh((a*tanh(x))^(1/2)/a^(1/2)))/tanh(x)/(a*tanh(x))^(
1/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(49) = 98.

Time = 0.11 (sec) , antiderivative size = 365, normalized size of antiderivative = 5.79

$$\int \sqrt{a \tanh^3(x)} dx = \text{Too large to display}$$

input

```
integrate((a*tanh(x)^3)^(1/2),x, algorithm="fricas")
```

output

```
[-1/2*sqrt(-a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2)) + 1/4*sqrt(-a)*log(-(a*cosh(x)^4 + 4*a*cosh(x)^3*sinh(x) + 6*a*cosh(x)^2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 + 2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x)) - 2*a)/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) - 2*sqrt(a*sinh(x)/cosh(x)), 1/2*sqrt(a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a*sinh(x)/cosh(x))/sqrt(a)) + 1/4*sqrt(a)*log(2*a*cosh(x)^4 + 8*a*cosh(x)^3*sinh(x) + 12*a*cosh(x)^2*sinh(x)^2 + 8*a*cosh(x)*sinh(x)^3 + 2*a*sinh(x)^4 + 2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))*sqrt(a)*sqrt(a*sinh(x)/cosh(x)) - a) - 2*sqrt(a*sinh(x)/cosh(x))]
```

**Sympy [F]**

$$\int \sqrt{a \tanh^3(x)} dx = \int \sqrt{a \tanh^3(x)} dx$$

input

```
integrate((a*tanh(x)**3)**(1/2),x)
```

output

```
Integral(sqrt(a*tanh(x)**3), x)
```

**Maxima [F]**

$$\int \sqrt{a \tanh^3(x)} dx = \int \sqrt{a \tanh^3(x)} dx$$

input

```
integrate((a*tanh(x)^3)^(1/2),x, algorithm="maxima")
```

output

```
integrate(sqrt(a*tanh(x)^3), x)
```



**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 115 vs.  $2(49) = 98$ .

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.83

$$\int \sqrt{a \tanh^3(x)} dx = \sqrt{a} \arctan \left( -\frac{\sqrt{a}e^{(2x)} - \sqrt{ae^{(4x)} - a}}{\sqrt{a}} \right) \operatorname{sgn}(e^{(4x)} - 1) \\ - \frac{1}{2} \sqrt{a} \log \left( \left| -\sqrt{a}e^{(2x)} + \sqrt{ae^{(4x)} - a} \right| \right) \operatorname{sgn}(e^{(4x)} - 1) \\ - \frac{4a \operatorname{sgn}(e^{(4x)} - 1)}{\sqrt{a}e^{(2x)} - \sqrt{ae^{(4x)} - a} + \sqrt{a}}$$

input `integrate((a*tanh(x)^3)^(1/2),x, algorithm="giac")`

output `sqrt(a)*arctan(-(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))/sqrt(a))*sgn(e^(4*x) - 1) - 1/2*sqrt(a)*log(abs(-sqrt(a)*e^(2*x) + sqrt(a*e^(4*x) - a)))*sgn(e^(4*x) - 1) - 4*a*sgn(e^(4*x) - 1)/(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a) + sqrt(a))`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \tanh^3(x)} dx = \int \sqrt{a \tanh(x)^3} dx$$

input `int((a*tanh(x)^3)^(1/2),x)`

output `int((a*tanh(x)^3)^(1/2), x)`

**Reduce [F]**

$$\int \sqrt{a \tanh^3(x)} dx = \sqrt{a} \left( -2\sqrt{\tanh(x)} + \int \frac{\sqrt{\tanh(x)}}{\tanh(x)} dx \right)$$

input `int((a*tanh(x)^3)^(1/2),x)`

output `sqrt(a)*(-2*sqrt(tanh(x)) + int(sqrt(tanh(x))/tanh(x),x))`

### 3.36 $\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$

Optimal result . . . . .	354
Mathematica [A] (verified) . . . . .	354
Rubi [A] (verified) . . . . .	355
Maple [A] (verified) . . . . .	358
Fricas [B] (verification not implemented) . . . . .	358
Sympy [F] . . . . .	359
Maxima [F] . . . . .	360
Giac [A] (verification not implemented) . . . . .	360
Mupad [F(-1)] . . . . .	360
Reduce [F] . . . . .	361

#### Optimal result

Integrand size = 10, antiderivative size = 64

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} - \frac{\arctan\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}}$$

output

```
-2*tanh(x)/(a*tanh(x)^3)^(1/2)-arctan(tanh(x)^(1/2))*tanh(x)^(3/2)/(a*tanh(x)^3)^(1/2)+arctanh(tanh(x)^(1/2))*tanh(x)^(3/2)/(a*tanh(x)^3)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \frac{\tanh(x) \left( 2 + \arctan\left(\sqrt[4]{\tanh^2(x)}\right) \sqrt[4]{\tanh^2(x)} - \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(x)}\right) \sqrt[4]{\tanh^2(x)} \right)}{\sqrt{a \tanh^3(x)}}$$

input `Integrate[1/Sqrt[a*Tanh[x]^3],x]`

output `-((Tanh[x]*(2 + ArcTan[(Tanh[x]^2)^(1/4)]*(Tanh[x]^2)^(1/4) - ArcTanh[(Tanh[x]^2)^(1/4)]*(Tanh[x]^2)^(1/4)))/Sqrt[a*Tanh[x]^3])`

### Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.80, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$ , Rules used = {3042, 4141, 3042, 3955, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \tanh^3(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{ia \tan(ix)^3}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\tanh^{\frac{3}{2}}(x) \int \frac{1}{\tanh^{\frac{3}{2}}(x)} dx}{\sqrt{a \tanh^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh^{\frac{3}{2}}(x) \int \frac{1}{(-i \tan(ix))^{3/2}} dx}{\sqrt{a \tanh^3(x)}} \\
 & \quad \downarrow \text{3955} \\
 & \frac{\tanh^{\frac{3}{2}}(x) \left( \int \sqrt{\tanh(x)} dx - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\tanh^{\frac{3}{2}}(x) \left( -\frac{2}{\sqrt{\tanh(x)}} + \int \sqrt{-i \tan(ix)} dx \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{3957} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( -\int -\frac{\sqrt{\tanh(x)}}{1-\tanh^2(x)} d \tanh(x) - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{25} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( \int \frac{\sqrt{\tanh(x)}}{1-\tanh^2(x)} d \tanh(x) - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{266} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( 2 \int \frac{\tanh(x)}{1-\tanh^2(x)} d \sqrt{\tanh(x)} - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{827} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d \sqrt{\tanh(x)} - \frac{1}{2} \int \frac{1}{\tanh(x)+1} d \sqrt{\tanh(x)} \right) - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{216} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(x)} d \sqrt{\tanh(x)} - \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}} \\
& \quad \downarrow \text{219} \\
& \frac{\tanh^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(x)} \right) - \frac{1}{2} \arctan \left( \sqrt{\tanh(x)} \right) \right) - \frac{2}{\sqrt{\tanh(x)}} \right)}{\sqrt{a \tanh^3(x)}}
\end{aligned}$$

input `Int [1/Sqrt [a*Tanh [x]^3] ,x]`

output `((2*(-1/2*ArcTan[Sqrt[Tanh[x]]] + ArcTanh[Sqrt[Tanh[x]]])/2) - 2/Sqrt[Tanh[x]])*Tanh[x]^(3/2))/Sqrt[a*Tanh[x]^3]`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 219  $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 266  $\text{Int}[(\text{c}_) * (\text{x}_)]^{(\text{m}_)} * ((\text{a}_) + (\text{b}_) * (\text{x}_)^2)^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{k} * (\text{m} + 1) - 1)} * (\text{a} + \text{b} * (\text{x}^{(2 * \text{k})}/\text{c}^2))^{\text{p}}, \text{x}], \text{x}, (\text{c} * \text{x})^{(1/\text{k})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \&\& \text{FractionQ}[\text{m}] \&\& \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 827  $\text{Int}[(\text{x}_)^2 / ((\text{a}_) + (\text{b}_) * (\text{x}_)^4), \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{s}/(2 * \text{b}) \quad \text{Int}[1/(\text{r} + \text{s} * \text{x}^2), \text{x}], \text{x}] - \text{Simp}[\text{s}/(2 * \text{b}) \quad \text{Int}[1/(\text{r} - \text{s} * \text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{!GtQ}[\text{a}/\text{b}, 0]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3955  $\text{Int}[(\text{b}_) * \tan[(\text{c}_) + (\text{d}_) * (\text{x}_)]]^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} + 1)} / (\text{b} * \text{d} * (\text{n} + 1)), \text{x}] - \text{Simp}[1/\text{b}^2 \quad \text{Int}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} + 2)}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[\text{n}, -1]$
- rule 3957  $\text{Int}[(\text{b}_) * \tan[(\text{c}_) + (\text{d}_) * (\text{x}_)]]^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{b}/\text{d} \quad \text{Subst}[\text{Int}[\text{x}^{\text{n}} / (\text{b}^2 + \text{x}^2), \text{x}], \text{x}, \text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}, \text{n}\}, \text{x}] \&\& \text{!IntegerQ}[\text{n}]$

rule 4141

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$-\frac{\tanh(x) \left( 2a^{\frac{5}{2}} + \arctan\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} - \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} \right)}{\sqrt{a \tanh(x)^3} a^{\frac{5}{2}}}$	65
default	$-\frac{\tanh(x) \left( 2a^{\frac{5}{2}} + \arctan\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} - \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} \right)}{\sqrt{a \tanh(x)^3} a^{\frac{5}{2}}}$	65

input

```
int(1/(a*tanh(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-tanh(x)*(2*a^(5/2)+arctan((a*tanh(x))^(1/2)/a^(1/2))*a^2*(a*tanh(x))^(1/2)
)-arctanh((a*tanh(x))^(1/2)/a^(1/2))*a^2*(a*tanh(x))^(1/2)/(a*tanh(x)^3)^(
(1/2)/a^(5/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(50) = 100.

Time = 0.11 (sec) , antiderivative size = 514, normalized size of antiderivative = 8.03

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \text{Too large to display}$$

input

```
integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="fricas")
```

output

```
[-1/4*(2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*arctan((
cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cos
h(x))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2)) + (cosh(x)^2 + 2*
cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*log(-(a*cosh(x)^4 + 4*a*cosh(x)^
3*sinh(x) + 6*a*cosh(x)^2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4
+ 2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)
)/cosh(x)) - 2*a)/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2
+ 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + 8*(cosh(x)^2 + 2*cosh(x)*sinh(x) +
sinh(x)^2 + 1)*sqrt(a*sinh(x)/cosh(x)))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x)
+ a*sinh(x)^2 - a), -1/4*(2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 -
1)*sqrt(a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a*si
nh(x)/cosh(x))/sqrt(a)) - (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)
*sqrt(a)*log(2*a*cosh(x)^4 + 8*a*cosh(x)^3*sinh(x) + 12*a*cosh(x)^2*sinh(x)
)^2 + 8*a*cosh(x)*sinh(x)^3 + 2*a*sinh(x)^4 + 2*(cosh(x)^4 + 4*cosh(x)*sin
h(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)
)^3 + cosh(x))*sinh(x))*sqrt(a)*sqrt(a*sinh(x)/cosh(x)) - a) + 8*(cosh(x)^
2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a*sinh(x)/cosh(x)))/(a*cosh(x)
^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a)]
```

## Sympy [F]

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \int \frac{1}{\sqrt{a \tanh^3(x)}} dx$$

input

```
integrate(1/(a*tanh(x)**3)**(1/2), x)
```

output

```
Integral(1/sqrt(a*tanh(x)**3), x)
```



**Maxima [F]**

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \int \frac{1}{\sqrt{a \tanh(x)^3}} dx$$

input `integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(a*tanh(x)^3), x)`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.64

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \frac{4}{(\sqrt{a}e^{(2x)} - \sqrt{a}e^{(4x)} - a - \sqrt{a}) \operatorname{sgn}(e^{(4x)} - 1)}$$

input `integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="giac")`

output `4/((sqrt(a)*e^(2*x) - sqrt(a)*e^(4*x) - a) - sqrt(a))*sgn(e^(4*x) - 1)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \int \frac{1}{\sqrt{a \tanh(x)^3}} dx$$

input `int(1/(a*tanh(x)^3)^(1/2),x)`

output `int(1/(a*tanh(x)^3)^(1/2), x)`

Reduce [F]

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx = \frac{\sqrt{a} \left( \int \frac{\sqrt{\tanh(x)}}{\tanh(x)^2} dx \right)}{a}$$

input `int(1/(a*tanh(x)^3)^(1/2),x)`

output `(sqrt(a)*int(sqrt(tanh(x))/tanh(x)**2,x))/a`

### 3.37 $\int (a \tanh^4(x))^{3/2} dx$

Optimal result	362
Mathematica [A] (verified)	362
Rubi [A] (verified)	363
Maple [A] (verified)	365
Fricas [B] (verification not implemented)	366
Sympy [F]	367
Maxima [A] (verification not implemented)	367
Giac [A] (verification not implemented)	367
Mupad [F(-1)]	368
Reduce [B] (verification not implemented)	368

#### Optimal result

Integrand size = 10, antiderivative size = 69

$$\int (a \tanh^4(x))^{3/2} dx = -a \coth(x) \sqrt{a \tanh^4(x)} + ax \coth^2(x) \sqrt{a \tanh^4(x)} - \frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)} - \frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)}$$

output

`-a*coth(x)*(a*tanh(x)^4)^(1/2)+a*x*coth(x)^2*(a*tanh(x)^4)^(1/2)-1/3*a*tanh(x)*(a*tanh(x)^4)^(1/2)-1/5*a*tanh(x)^3*(a*tanh(x)^4)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.57

$$\int (a \tanh^4(x))^{3/2} dx = \frac{1}{15} \coth(x) (-3 - 5 \coth^2(x) - 15 \coth^4(x) + 15 \arctanh(\tanh(x)) \coth^5(x)) (a \tanh^4(x))^{3/2}$$

input

`Integrate[(a*Tanh[x]^4)^(3/2),x]`

output

```
(Coth[x]*(-3 - 5*Coth[x]^2 - 15*Coth[x]^4 + 15*ArcTanh[Tanh[x]]*Coth[x]^5)
*(a*Tanh[x]^4)^(3/2))/15
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.57, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$ , Rules used = {3042, 4141, 3042, 25, 3954, 3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh^4(x))^{3/2} dx \\
 & \quad \downarrow 3042 \\
 & \int (a \tan(ix)^4)^{3/2} dx \\
 & \quad \downarrow 4141 \\
 & a \coth^2(x) \sqrt{a \tanh^4(x)} \int \tanh^6(x) dx \\
 & \quad \downarrow 3042 \\
 & a \coth^2(x) \sqrt{a \tanh^4(x)} \int -\tan(ix)^6 dx \\
 & \quad \downarrow 25 \\
 & -a \coth^2(x) \sqrt{a \tanh^4(x)} \int \tan(ix)^6 dx \\
 & \quad \downarrow 3954 \\
 & -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( \frac{\tanh^5(x)}{5} - \int \tanh^4(x) dx \right) \\
 & \quad \downarrow 3042 \\
 & -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( \frac{\tanh^5(x)}{5} - \int \tan(ix)^4 dx \right) \\
 & \quad \downarrow 3954
 \end{aligned}$$

$$\begin{aligned}
& -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( \int -\tanh^2(x) dx + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} \right) \\
& \quad \downarrow 25 \\
& -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( -\int \tanh^2(x) dx + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} \right) \\
& \quad \downarrow 3042 \\
& -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( -\int -\tan(ix)^2 dx + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} \right) \\
& \quad \downarrow 25 \\
& -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( \int \tan(ix)^2 dx + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} \right) \\
& \quad \downarrow 3954 \\
& -a \coth^2(x) \sqrt{a \tanh^4(x)} \left( -\int 1 dx + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} + \tanh(x) \right) \\
& \quad \downarrow 24 \\
& -a \left( -x + \frac{\tanh^5(x)}{5} + \frac{\tanh^3(x)}{3} + \tanh(x) \right) \coth^2(x) \sqrt{a \tanh^4(x)}
\end{aligned}$$

input `Int[(a*Tanh[x]^4)^(3/2),x]`

output `-(a*Coth[x]^2*sqrt[a*Tanh[x]^4]*(-x + Tanh[x] + Tanh[x]^3/3 + Tanh[x]^5/5))`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\frac{(a \tanh(x)^4)^{\frac{3}{2}} (6 \tanh(x)^5 + 10 \tanh(x)^3 + 15 \ln(\tanh(x) - 1) - 15 \ln(1 + \tanh(x)) + 30 \tanh(x))}{30 \tanh(x)^6}$	46
default	$-\frac{(a \tanh(x)^4)^{\frac{3}{2}} (6 \tanh(x)^5 + 10 \tanh(x)^3 + 15 \ln(\tanh(x) - 1) - 15 \ln(1 + \tanh(x)) + 30 \tanh(x))}{30 \tanh(x)^6}$	46
risch	$\frac{a(e^{2x}+1)^2 \sqrt{\frac{a(e^{2x}-1)^4}{(e^{2x}+1)^4}} x}{(e^{2x}-1)^2} + \frac{2a \sqrt{\frac{a(e^{2x}-1)^4}{(e^{2x}+1)^4}} (45 e^{8x} + 90 e^{6x} + 140 e^{4x} + 70 e^{2x} + 23)}{15(e^{2x}-1)^2 (e^{2x}+1)^3}$	106

input `int((a*tanh(x)^4)^(3/2), x, method=_RETURNVERBOSE)`

output `-1/30*(a*tanh(x)^4)^(3/2)*(6*tanh(x)^5+10*tanh(x)^3+15*ln(tanh(x)-1)-15*ln(1+tanh(x))+30*tanh(x))/tanh(x)^6`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2114 vs.  $2(57) = 114$ .

Time = 0.17 (sec) , antiderivative size = 2114, normalized size of antiderivative = 30.64

$$\int (a \tanh^4(x))^{3/2} dx = \text{Too large to display}$$

input `integrate((a*tanh(x)^4)^(3/2),x, algorithm="fricas")`

output

```
1/15*(15*a*x*cosh(x)^10 + 15*(a*x*e^(4*x) + 2*a*x*e^(2*x) + a*x)*sinh(x)^10 + 150*(a*x*cosh(x)*e^(4*x) + 2*a*x*cosh(x)*e^(2*x) + a*x*cosh(x))*sinh(x)^9 + 15*(5*a*x + 6*a)*cosh(x)^8 + 15*(45*a*x*cosh(x)^2 + 5*a*x + (45*a*x*cosh(x)^2 + 5*a*x + 6*a)*e^(4*x) + 2*(45*a*x*cosh(x)^2 + 5*a*x + 6*a)*e^(2*x) + 6*a)*sinh(x)^8 + 120*(15*a*x*cosh(x)^3 + (5*a*x + 6*a)*cosh(x) + (15*a*x*cosh(x)^3 + (5*a*x + 6*a)*cosh(x))*e^(4*x) + 2*(15*a*x*cosh(x)^3 + (5*a*x + 6*a)*cosh(x))*e^(2*x))*sinh(x)^7 + 30*(5*a*x + 6*a)*cosh(x)^6 + 30*(105*a*x*cosh(x)^4 + 14*(5*a*x + 6*a)*cosh(x)^2 + 5*a*x + (105*a*x*cosh(x)^4 + 14*(5*a*x + 6*a)*cosh(x)^2 + 5*a*x + 6*a)*e^(4*x) + 2*(105*a*x*cosh(x)^4 + 14*(5*a*x + 6*a)*cosh(x)^2 + 5*a*x + 6*a)*e^(2*x) + 6*a)*sinh(x)^6 + 60*(63*a*x*cosh(x)^5 + 14*(5*a*x + 6*a)*cosh(x)^3 + 3*(5*a*x + 6*a)*cosh(x) + (63*a*x*cosh(x)^5 + 14*(5*a*x + 6*a)*cosh(x)^3 + 3*(5*a*x + 6*a)*cosh(x))*e^(4*x) + 2*(63*a*x*cosh(x)^5 + 14*(5*a*x + 6*a)*cosh(x)^3 + 3*(5*a*x + 6*a)*cosh(x))*e^(2*x))*sinh(x)^5 + 10*(15*a*x + 28*a)*cosh(x)^4 + 10*(315*a*x*cosh(x)^6 + 105*(5*a*x + 6*a)*cosh(x)^4 + 45*(5*a*x + 6*a)*cosh(x)^2 + 15*a*x + (315*a*x*cosh(x)^6 + 105*(5*a*x + 6*a)*cosh(x)^4 + 45*(5*a*x + 6*a)*cosh(x)^2 + 15*a*x + 28*a)*e^(4*x) + 2*(315*a*x*cosh(x)^6 + 105*(5*a*x + 6*a)*cosh(x)^4 + 45*(5*a*x + 6*a)*cosh(x)^2 + 15*a*x + 28*a)*e^(2*x) + 28*a)*sinh(x)^4 + 40*(45*a*x*cosh(x)^7 + 21*(5*a*x + 6*a)*cosh(x)^5 + 15*(5*a*x + 6*a)*cosh(x)^3 + (15*a*x + 28*a)*cosh(x) + (45*a*x*cosh(x)^7...
```

**Sympy [F]**

$$\int (a \tanh^4(x))^{3/2} dx = \int (a \tanh^4(x))^{\frac{3}{2}} dx$$

input `integrate((a*tanh(x)**4)**(3/2),x)`

output `Integral((a*tanh(x)**4)**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.19

$$\int (a \tanh^4(x))^{3/2} dx = a^{\frac{3}{2}} x - \frac{2 \left( 70 a^{\frac{3}{2}} e^{-2x} + 140 a^{\frac{3}{2}} e^{-4x} + 90 a^{\frac{3}{2}} e^{-6x} + 45 a^{\frac{3}{2}} e^{-8x} + 23 a^{\frac{3}{2}} \right)}{15 (5 e^{-2x} + 10 e^{-4x} + 10 e^{-6x} + 5 e^{-8x} + e^{-10x} + 1)}$$

input `integrate((a*tanh(x)^4)^(3/2),x, algorithm="maxima")`

output `a^(3/2)*x - 2/15*(70*a^(3/2)*e^(-2*x) + 140*a^(3/2)*e^(-4*x) + 90*a^(3/2)*e^(-6*x) + 45*a^(3/2)*e^(-8*x) + 23*a^(3/2))/(5*e^(-2*x) + 10*e^(-4*x) + 10*e^(-6*x) + 5*e^(-8*x) + e^(-10*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int (a \tanh^4(x))^{3/2} dx = \frac{1}{15} a^{\frac{3}{2}} \left( 15x + \frac{2(45e^{8x} + 90e^{6x} + 140e^{4x} + 70e^{2x} + 23)}{(e^{2x} + 1)^5} \right)$$

input `integrate((a*tanh(x)^4)^(3/2),x, algorithm="giac")`



output

```
1/15*a^(3/2)*(15*x + 2*(45*e^(8*x) + 90*e^(6*x) + 140*e^(4*x) + 70*e^(2*x)
+ 23)/(e^(2*x) + 1)^5)
```

**Mupad [F(-1)]**

Timed out.

$$\int (a \tanh^4(x))^{3/2} dx = \int (a \tanh(x)^4)^{3/2} dx$$

input

```
int((a*tanh(x)^4)^(3/2),x)
```

output

```
int((a*tanh(x)^4)^(3/2), x)
```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int (a \tanh^4(x))^{3/2} dx = \frac{\sqrt{a} a (-3 \tanh(x)^5 - 5 \tanh(x)^3 - 15 \tanh(x) + 15x)}{15}$$

input

```
int((a*tanh(x)^4)^(3/2),x)
```

output

```
(sqrt(a)*a*( - 3*tanh(x)**5 - 5*tanh(x)**3 - 15*tanh(x) + 15*x))/15
```

### 3.38 $\int \sqrt{a \tanh^4(x)} dx$

Optimal result	369
Mathematica [A] (verified)	369
Rubi [A] (verified)	370
Maple [A] (verified)	371
Fricas [B] (verification not implemented)	372
Sympy [F]	372
Maxima [A] (verification not implemented)	373
Giac [A] (verification not implemented)	373
Mupad [F(-1)]	373
Reduce [B] (verification not implemented)	374

#### Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \sqrt{a \tanh^4(x)} dx = -\coth(x)\sqrt{a \tanh^4(x)} + x \coth^2(x)\sqrt{a \tanh^4(x)}$$

output

```
-coth(x)*(a*tanh(x)^4)^(1/2)+x*coth(x)^2*(a*tanh(x)^4)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \sqrt{a \tanh^4(x)} dx = \coth(x)(-1 + \operatorname{arctanh}(\tanh(x)) \coth(x))\sqrt{a \tanh^4(x)}$$

input

```
Integrate[Sqrt[a*Tanh[x]^4],x]
```

output

```
Coth[x]*(-1 + ArcTanh[Tanh[x]]*Coth[x])*Sqrt[a*Tanh[x]^4]
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4141, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \tanh^4(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \tan(ix)^4} dx \\
 & \quad \downarrow \text{4141} \\
 & \coth^2(x) \sqrt{a \tanh^4(x)} \int \tanh^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \coth^2(x) \sqrt{a \tanh^4(x)} \int -\tan(ix)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\coth^2(x) \sqrt{a \tanh^4(x)} \int \tan(ix)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & -\coth^2(x) \sqrt{a \tanh^4(x)} (\tanh(x) - \int 1 dx) \\
 & \quad \downarrow \text{24} \\
 & (\tanh(x) - x) (-\coth^2(x)) \sqrt{a \tanh^4(x)}
 \end{aligned}$$

input `Int [Sqrt [a*Tanh [x]^4] , x]`

output `-(Coth [x]^2*Sqrt [a*Tanh [x]^4]*(-x + Tanh [x]))`

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$-\frac{\sqrt{a \tanh(x)^4} (2 \tanh(x) + \ln(\tanh(x) - 1) - \ln(1 + \tanh(x)))}{2 \tanh(x)^2}$	32
default	$-\frac{\sqrt{a \tanh(x)^4} (2 \tanh(x) + \ln(\tanh(x) - 1) - \ln(1 + \tanh(x)))}{2 \tanh(x)^2}$	32
risch	$\frac{\sqrt{\frac{a(e^{2x}-1)^4}{(e^{2x}+1)^4}} (e^{2x}+1)^2 x}{(e^{2x}-1)^2} + \frac{2\sqrt{\frac{a(e^{2x}-1)^4}{(e^{2x}+1)^4}} (e^{2x}+1)}{(e^{2x}-1)^2}$	76

input `int((a*tanh(x)^4)^(1/2), x, method=_RETURNVERBOSE)`

output  $-1/2*(a*\tanh(x)^4)^{(1/2)*(2*\tanh(x)+\ln(\tanh(x)-1)-\ln(1+\tanh(x)))/\tanh(x)^2$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(27) = 54.

Time = 0.09 (sec) , antiderivative size = 213, normalized size of antiderivative = 6.87

$$\int \sqrt{a \tanh^4(x)} dx$$

$$= \frac{(x \cosh(x)^2 + (x e^{4x} + 2x e^{2x} + x) \sinh(x)^2 + (x \cosh(x)^2 + x + 2) e^{4x} + 2(x \cosh(x)^2 + x + 2) e^{2x})}{(e^{4x} - 2e^{2x} + 1) \sinh(x)^2 + \cosh(x)^2 + (\cosh(x)^2 + 1) e^{4x} - 2(\cosh(x)^2 + 1) e^{2x}}$$

input `integrate((a*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output  $(x*\cosh(x)^2 + (x*e^{4*x} + 2*x*e^{2*x} + x)*\sinh(x)^2 + (x*\cosh(x)^2 + x + 2)*e^{4*x} + 2*(x*\cosh(x)^2 + x + 2)*e^{2*x} + 2*(x*\cosh(x)*e^{4*x} + 2*x*\cosh(x)*e^{2*x} + x*\cosh(x))*\sinh(x) + x + 2)*\text{sqrt}((a*e^{8*x} - 4*a*e^{6*x} + 6*a*e^{4*x} - 4*a*e^{2*x} + a)/(e^{8*x} + 4*e^{6*x} + 6*e^{4*x} + 4*e^{2*x} + 1))/((e^{4*x} - 2*e^{2*x} + 1)*\sinh(x)^2 + \cosh(x)^2 + (\cosh(x)^2 + 1)*e^{4*x} - 2*(\cosh(x)^2 + 1)*e^{2*x} + 2*(\cosh(x)*e^{4*x} - 2*\cosh(x)*e^{2*x} + \cosh(x))*\sinh(x) + 1)$

### Sympy [F]

$$\int \sqrt{a \tanh^4(x)} dx = \int \sqrt{a \tanh^4(x)} dx$$

input `integrate((a*tanh(x)**4)**(1/2),x)`

output `Integral(sqrt(a*tanh(x)**4), x)`

**Maxima [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \sqrt{a \tanh^4(x)} dx = \sqrt{a}x - \frac{2\sqrt{a}}{e^{(-2x)} + 1}$$

input `integrate((a*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `sqrt(a)*x - 2*sqrt(a)/(e^(-2*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.52

$$\int \sqrt{a \tanh^4(x)} dx = \sqrt{a} \left( x + \frac{2}{e^{(2x)} + 1} \right)$$

input `integrate((a*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `sqrt(a)*(x + 2/(e^(2*x) + 1))`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \tanh^4(x)} dx = \int \sqrt{a \tanh(x)^4} dx$$

input `int((a*tanh(x)^4)^(1/2),x)`

output `int((a*tanh(x)^4)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.29

$$\int \sqrt{a \tanh^4(x)} dx = \sqrt{a} (-\tanh(x) + x)$$

input `int((a*tanh(x)^4)^(1/2),x)`

output `sqrt(a)*(-tanh(x) + x)`

### 3.39 $\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$

Optimal result	375
Mathematica [C] (verified)	375
Rubi [A] (verified)	376
Maple [A] (verified)	378
Fricas [B] (verification not implemented)	378
Sympy [F]	379
Maxima [A] (verification not implemented)	379
Giac [A] (verification not implemented)	380
Mupad [F(-1)]	380
Reduce [B] (verification not implemented)	380

#### Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = -\frac{\tanh(x)}{\sqrt{a \tanh^4(x)}} + \frac{x \tanh^2(x)}{\sqrt{a \tanh^4(x)}}$$

output `-tanh(x)/(a*tanh(x)^4)^(1/2)+x*tanh(x)^2/(a*tanh(x)^4)^(1/2)`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = -\frac{\text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \tanh^2(x)\right) \tanh(x)}{\sqrt{a \tanh^4(x)}}$$

input `Integrate[1/Sqrt[a*Tanh[x]^4],x]`

output `-((Hypergeometric2F1[-1/2, 1, 1/2, Tanh[x]^2]*Tanh[x])/Sqrt[a*Tanh[x]^4])`



**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4141, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \tanh^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a \tan(ix)^4}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\tanh^2(x) \int \coth^2(x) dx}{\sqrt{a \tanh^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh^2(x) \int -\tan\left(ix + \frac{\pi}{2}\right)^2 dx}{\sqrt{a \tanh^4(x)}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\tanh^2(x) \int \tan\left(ix + \frac{\pi}{2}\right)^2 dx}{\sqrt{a \tanh^4(x)}} \\
 & \quad \downarrow \text{3954} \\
 & -\frac{\tanh^2(x)(\coth(x) - \int 1 dx)}{\sqrt{a \tanh^4(x)}} \\
 & \quad \downarrow \text{24} \\
 & -\frac{\tanh^2(x)(\coth(x) - x)}{\sqrt{a \tanh^4(x)}}
 \end{aligned}$$

input `Int[1/Sqrt[a*Tanh[x]^4],x]`

output `-((( -x + Coth[x])*Tanh[x]^2)/Sqrt[a*Tanh[x]^4])`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$-\frac{\tanh(x)(\ln(\tanh(x)-1)\tanh(x)-\ln(1+\tanh(x))\tanh(x)+2)}{2\sqrt{a}\tanh(x)^4}$	32
default	$-\frac{\tanh(x)(\ln(\tanh(x)-1)\tanh(x)-\ln(1+\tanh(x))\tanh(x)+2)}{2\sqrt{a}\tanh(x)^4}$	32
risch	$\frac{e^{4x}x-2e^{2x}x-2e^{2x}+x+2}{\sqrt{\frac{a(e^{2x}-1)^4}{(e^{2x}+1)^4}}(e^{2x}+1)^2}$	52

input `int(1/(a*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)`output 
$$-1/2*\tanh(x)*(\ln(\tanh(x)-1)*\tanh(x)-\ln(1+\tanh(x))*\tanh(x)+2)/(a*\tanh(x)^4)^{(1/2)}$$
**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 238 vs.  $2(27) = 54$ .

Time = 0.12 (sec) , antiderivative size = 238, normalized size of antiderivative = 7.68

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$$

$$= \frac{(x \cosh(x)^2 + (xe^{4x} + 2xe^{2x} + x) \sinh(x)^2 + (x \cosh(x)^2 - x - 2)e^{4x} + 2(x \cosh(x)^2 - x - 2)e^{2x})}{a \cosh(x)^2 + (ae^{4x} - 2ae^{2x} + a) \sinh(x)^2 + (a \cosh(x)^2 - a)e^{4x} - 2(a \cosh(x)^2 - a)e^{2x}}$$

input `integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output

```
(x*cosh(x)^2 + (x*e^(4*x) + 2*x*e^(2*x) + x)*sinh(x)^2 + (x*cosh(x)^2 - x
- 2)*e^(4*x) + 2*(x*cosh(x)^2 - x - 2)*e^(2*x) + 2*(x*cosh(x)*e^(4*x) + 2*
x*cosh(x)*e^(2*x) + x*cosh(x))*sinh(x) - x - 2)*sqrt((a*e^(8*x) - 4*a*e^(6
*x) + 6*a*e^(4*x) - 4*a*e^(2*x) + a)/(e^(8*x) + 4*e^(6*x) + 6*e^(4*x) + 4*
e^(2*x) + 1))/(a*cosh(x)^2 + (a*e^(4*x) - 2*a*e^(2*x) + a)*sinh(x)^2 + (a*
cosh(x)^2 - a)*e^(4*x) - 2*(a*cosh(x)^2 - a)*e^(2*x) + 2*(a*cosh(x)*e^(4*x
) - 2*a*cosh(x)*e^(2*x) + a*cosh(x))*sinh(x) - a)
```

**Sympy [F]**

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = \int \frac{1}{\sqrt{a \tanh^4(x)}} dx$$

input

```
integrate(1/(a*tanh(x)**4)**(1/2),x)
```

output

```
Integral(1/sqrt(a*tanh(x)**4), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = \frac{x}{\sqrt{a}} + \frac{2\sqrt{a}}{ae^{(-2x)} - a}$$

input

```
integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="maxima")
```

output

```
x/sqrt(a) + 2*sqrt(a)/(a*e^(-2*x) - a)
```

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = \frac{x}{\sqrt{a}} - \frac{2}{\sqrt{a}(e^{2x} - 1)}$$

input `integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `x/sqrt(a) - 2/(sqrt(a)*(e^(2*x) - 1))`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = \int \frac{1}{\sqrt{a \tanh(x)^4}} dx$$

input `int(1/(a*tanh(x)^4)^(1/2),x)`

output `int(1/(a*tanh(x)^4)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx = \frac{\sqrt{a}(\tanh(x)x - 1)}{\tanh(x)a}$$

input `int(1/(a*tanh(x)^4)^(1/2),x)`

output `(sqrt(a)*(tanh(x)*x - 1))/(tanh(x)*a)`

### 3.40 $\int (b \tanh^m(c + dx))^n dx$

Optimal result	381
Mathematica [A] (verified)	381
Rubi [A] (verified)	382
Maple [F]	384
Fricas [F]	384
Sympy [F]	384
Maxima [F]	385
Giac [F]	385
Mupad [F(-1)]	385
Reduce [F]	386

#### Optimal result

Integrand size = 12, antiderivative size = 57

$$\int (b \tanh^m(c + dx))^n dx$$

$$= \frac{\text{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + mn), \frac{1}{2}(3 + mn), \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh^m(c + dx))^n}{d(1 + mn)}$$

output

```
hypergeom([1, 1/2*m*n+1/2], [1/2*m*n+3/2], tanh(d*x+c)^2)*tanh(d*x+c)*(b*tanh(d*x+c)^m)^n/d/(m*n+1)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec), antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int (b \tanh^m(c + dx))^n dx$$

$$= \frac{\text{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + mn), \frac{1}{2}(3 + mn), \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh^m(c + dx))^n}{d(1 + mn)}$$

input

```
Integrate[(b*Tanh[c + d*x]^m)^n,x]
```

output

$$\text{(Hypergeometric2F1[1, (1 + m*n)/2, (3 + m*n)/2, \text{Tanh}[c + d*x]^2]*\text{Tanh}[c + d*x]*(b*\text{Tanh}[c + d*x]^m)^n)/(d*(1 + m*n))}$$
**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4142, 3042, 3957, 25, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (b \tanh^m(c + dx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int (b(-i \tan(ic + idx))^m)^n dx \\ & \quad \downarrow \text{4142} \\ & \tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n \int \tanh^{mn}(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n \int (-i \tan(ic + idx))^{mn} dx \\ & \quad \downarrow \text{3957} \\ & \frac{\tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n \int -\frac{\tanh^{mn}(c+dx)}{1-\tanh^2(c+dx)} d \tanh(c + dx)}{d} \\ & \quad \downarrow \text{25} \\ & \frac{\tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n \int \frac{\tanh^{mn}(c+dx)}{1-\tanh^2(c+dx)} d \tanh(c + dx)}{d} \\ & \quad \downarrow \text{278} \\ & \frac{\tanh(c + dx) (b \tanh^m(c + dx))^n \text{Hypergeometric2F1}\left(1, \frac{1}{2}(mn + 1), \frac{1}{2}(mn + 3), \tanh^2(c + dx)\right)}{d(mn + 1)} \end{aligned}$$

input `Int[(b*Tanh[c + d*x]^m)^n,x]`

output `(Hypergeometric2F1[1, (1 + m*n)/2, (3 + m*n)/2, Tanh[c + d*x]^2]*Tanh[c + d*x]*(b*Tanh[c + d*x]^m)^n)/(d*(1 + m*n))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4142 `Int[(u_)*((b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := Simp[b^IntPart[p]*((b*(c*Tan[e + f*x])^n)^FracPart[p]/(c*Tan[e + f*x])^(n*FracPart[p])) Int[ActivateTrig[u]*(c*Tan[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_)*(trig_)[e + f*x])^(m_) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])`



**Maple [F]**

$$\int (b \tanh(dx + c)^m)^n dx$$

input `int((b*tanh(d*x+c)^m)^n,x)`

output `int((b*tanh(d*x+c)^m)^n,x)`

**Fricas [F]**

$$\int (b \tanh^m(c + dx))^n dx = \int (b \tanh(dx + c)^m)^n dx$$

input `integrate((b*tanh(d*x+c)^m)^n,x, algorithm="fricas")`

output `integral((b*tanh(d*x + c)^m)^n, x)`

**Sympy [F]**

$$\int (b \tanh^m(c + dx))^n dx = \int (b \tanh^m(c + dx))^n dx$$

input `integrate((b*tanh(d*x+c)**m)**n,x)`

output `Integral((b*tanh(c + d*x)**m)**n, x)`

**Maxima [F]**

$$\int (b \tanh^m(c + dx))^n dx = \int (b \tanh(dx + c)^m)^n dx$$

input `integrate((b*tanh(d*x+c)^m)^n,x, algorithm="maxima")`

output `integrate((b*tanh(d*x + c)^m)^n, x)`

**Giac [F]**

$$\int (b \tanh^m(c + dx))^n dx = \int (b \tanh(dx + c)^m)^n dx$$

input `integrate((b*tanh(d*x+c)^m)^n,x, algorithm="giac")`

output `integrate((b*tanh(d*x + c)^m)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \tanh^m(c + dx))^n dx = \int (b \tanh(c + dx)^m)^n dx$$

input `int((b*tanh(c + d*x)^m)^n,x)`

output `int((b*tanh(c + d*x)^m)^n, x)`

**Reduce [F]**

$$\int (b \tanh^m(c + dx))^n dx = b^n \left( \int \tanh(dx + c)^{mn} dx \right)$$

input `int((b*tanh(d*x+c)^m)^n,x)`

output `b**n*int(tanh(c + d*x)**(m*n),x)`

### 3.41 $\int (a + a \tanh(c + dx))^5 dx$

Optimal result . . . . .	387
Mathematica [A] (verified) . . . . .	387
Rubi [A] (verified) . . . . .	388
Maple [A] (verified) . . . . .	391
Fricas [B] (verification not implemented) . . . . .	391
Sympy [A] (verification not implemented) . . . . .	392
Maxima [B] (verification not implemented) . . . . .	393
Giac [A] (verification not implemented) . . . . .	393
Mupad [B] (verification not implemented) . . . . .	394
Reduce [B] (verification not implemented) . . . . .	394

#### Optimal result

Integrand size = 12, antiderivative size = 100

$$\int (a + a \tanh(c + dx))^5 dx = 16a^5 x + \frac{16a^5 \log(\cosh(c + dx))}{d} - \frac{8a^5 \tanh(c + dx)}{d} - \frac{2a^2(a + a \tanh(c + dx))^3}{3d} - \frac{a(a + a \tanh(c + dx))^4}{4d} - \frac{2a(a^2 + a^2 \tanh(c + dx))^2}{d}$$

output

```
16*a^5*x+16*a^5*ln(cosh(d*x+c))/d-8*a^5*tanh(d*x+c)/d-2/3*a^2*(a+a*tanh(d*x+c))^3/d-1/4*a*(a+a*tanh(d*x+c))^4/d-2*a*(a^2+a^2*tanh(d*x+c))^2/d
```

#### Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.63

$$\int (a + a \tanh(c + dx))^5 dx = \frac{a^5 (35 + 192 \log(1 - \tanh(c + dx)) + 180 \tanh(c + dx) + 66 \tanh^2(c + dx) + 20 \tanh^3(c + dx) + 3 \tanh^4(c + dx))}{12d}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^5,x]
```

output

```
-1/12*(a^5*(35 + 192*Log[1 - Tanh[c + d*x]] + 180*Tanh[c + d*x] + 66*Tanh[
c + d*x]^2 + 20*Tanh[c + d*x]^3 + 3*Tanh[c + d*x]^4))/d
```

**Rubi [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3959, 3042, 3959, 3042, 3959, 3042, 3958, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh(c + dx) + a)^5 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ia \tan(ic + idx))^5 dx \\
 & \quad \downarrow \text{3959} \\
 & 2a \int (\tanh(c + dx)a + a)^4 dx - \frac{a(a \tanh(c + dx) + a)^4}{4d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a(a \tanh(c + dx) + a)^4}{4d} + 2a \int (a - ia \tan(ic + idx))^4 dx \\
 & \quad \downarrow \text{3959} \\
 & 2a \left( 2a \int (\tanh(c + dx)a + a)^3 dx - \frac{a(a \tanh(c + dx) + a)^3}{3d} \right) - \frac{a(a \tanh(c + dx) + a)^4}{4d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a(a \tanh(c + dx) + a)^4}{4d} + 2a \left( -\frac{a(a \tanh(c + dx) + a)^3}{3d} + 2a \int (a - ia \tan(ic + idx))^3 dx \right) \\
 & \quad \downarrow \text{3959} \\
 & 2a \left( 2a \left( 2a \int (\tanh(c + dx)a + a)^2 dx - \frac{a(a \tanh(c + dx) + a)^2}{2d} \right) - \frac{a(a \tanh(c + dx) + a)^3}{3d} \right) - \\
 & \quad \frac{a(a \tanh(c + dx) + a)^4}{4d}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & 2a \left( -\frac{a(a \tanh(c+dx) + a)^3}{3d} + 2a \left( -\frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \int (a - ia \tan(ic + idx))^2 dx \right) \right) \\
 & \downarrow 3958 \\
 & 2a \left( -\frac{a(a \tanh(c+dx) + a)^3}{3d} + 2a \left( -\frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( -2ia^2 \int i \tanh(c+dx) dx - \frac{a^2 \tanh(c+dx)}{d} \right) \right) \right) \\
 & \downarrow 26 \\
 & 2a \left( 2a \left( 2a \left( 2a^2 \int \tanh(c+dx) dx - \frac{a^2 \tanh(c+dx)}{d} + 2a^2 x \right) - \frac{a(a \tanh(c+dx) + a)^2}{2d} \right) - \frac{a(a \tanh(c+dx) + a)^4}{4d} - \frac{a(a \tanh(c+dx) + a)^3}{3d} \right) \\
 & \downarrow 3042 \\
 & 2a \left( -\frac{a(a \tanh(c+dx) + a)^3}{3d} + 2a \left( -\frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( 2a^2 \int -i \tan(ic + idx) dx - \frac{a^2 \tanh(c+dx)}{d} \right) \right) \right) \\
 & \downarrow 26 \\
 & 2a \left( -\frac{a(a \tanh(c+dx) + a)^3}{3d} + 2a \left( -\frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( -2ia^2 \int \tan(ic + idx) dx - \frac{a^2 \tanh(c+dx)}{d} \right) \right) \right) \\
 & \downarrow 3956 \\
 & 2a \left( 2a \left( 2a \left( -\frac{a^2 \tanh(c+dx)}{d} + \frac{2a^2 \log(\cosh(c+dx))}{d} + 2a^2 x \right) - \frac{a(a \tanh(c+dx) + a)^2}{2d} \right) - \frac{a(a \tanh(c+dx) + a)^4}{4d} - \frac{a(a \tanh(c+dx) + a)^3}{3d} \right)
 \end{aligned}$$

input `Int[(a + a*Tanh[c + d*x])^5,x]`

output

```
-1/4*(a*(a + a*Tanh[c + d*x])^4)/d + 2*a*(-1/3*(a*(a + a*Tanh[c + d*x])^3)
/d + 2*a*(-1/2*(a*(a + a*Tanh[c + d*x])^2)/d + 2*a*(2*a^2*x + (2*a^2*Log[C
osh[c + d*x]])/d - (a^2*Tanh[c + d*x])/d))
```

**Defintions of rubi rules used**

rule 26

```
Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 3958

```
Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)
*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x]
, x]) /; FreeQ[{a, b, c, d}, x]
```

rule 3959

```
Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a +
b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*
x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n
, 1]
```

**Maple [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.58

method	result
derivativedivides	$\frac{a^5 \left( -\frac{\tanh(dx+c)^4}{4} - \frac{5 \tanh(dx+c)^3}{3} - \frac{11 \tanh(dx+c)^2}{2} - 15 \tanh(dx+c) - 16 \ln(\tanh(dx+c)-1) \right)}{d}$
default	$\frac{a^5 \left( -\frac{\tanh(dx+c)^4}{4} - \frac{5 \tanh(dx+c)^3}{3} - \frac{11 \tanh(dx+c)^2}{2} - 15 \tanh(dx+c) - 16 \ln(\tanh(dx+c)-1) \right)}{d}$
parallelrisc	$-\frac{3 \tanh(dx+c)^4 a^5 + 20 \tanh(dx+c)^3 a^5 + 66 \tanh(dx+c)^2 a^5 + 192 \ln(1 - \tanh(dx+c)) a^5 + 180 a^5 \tanh(dx+c)}{12d}$
risc	$-\frac{32a^5 c}{d} + \frac{4a^5 (48 e^{6dx+6c} + 108 e^{4dx+4c} + 88 e^{2dx+2c} + 25)}{3d(e^{2dx+2c} + 1)^4} + \frac{16a^5 \ln(e^{2dx+2c} + 1)}{d}$
parts	$a^5 x + \frac{a^5 \left( -\frac{\tanh(dx+c)^4}{4} - \frac{\tanh(dx+c)^2}{2} - \frac{\ln(\tanh(dx+c)-1)}{2} - \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{5a^5 \ln(\cosh(dx+c))}{d} + \frac{10a^5}{d}$

input `int((a+a*tanh(d*x+c))^5,x,method=_RETURNVERBOSE)`output `1/d*a^5*(-1/4*tanh(d*x+c)^4-5/3*tanh(d*x+c)^3-11/2*tanh(d*x+c)^2-15*tanh(d*x+c)-16*ln(tanh(d*x+c)-1))`**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 907 vs.  $2(96) = 192$ .

Time = 0.09 (sec) , antiderivative size = 907, normalized size of antiderivative = 9.07

$$\int (a + a \tanh(c + dx))^5 dx = \text{Too large to display}$$

input `integrate((a+a*tanh(d*x+c))^5,x, algorithm="fricas")`



output

```

4/3*(48*a^5*cosh(d*x + c)^6 + 288*a^5*cosh(d*x + c)*sinh(d*x + c)^5 + 48*a
^5*sinh(d*x + c)^6 + 108*a^5*cosh(d*x + c)^4 + 88*a^5*cosh(d*x + c)^2 + 25
*a^5 + 36*(20*a^5*cosh(d*x + c)^2 + 3*a^5)*sinh(d*x + c)^4 + 48*(20*a^5*co
sh(d*x + c)^3 + 9*a^5*cosh(d*x + c))*sinh(d*x + c)^3 + 8*(90*a^5*cosh(d*x
+ c)^4 + 81*a^5*cosh(d*x + c)^2 + 11*a^5)*sinh(d*x + c)^2 + 12*(a^5*cosh(d
*x + c)^8 + 8*a^5*cosh(d*x + c)*sinh(d*x + c)^7 + a^5*sinh(d*x + c)^8 + 4*
a^5*cosh(d*x + c)^6 + 6*a^5*cosh(d*x + c)^4 + 4*a^5*cosh(d*x + c)^2 + 4*(7
*a^5*cosh(d*x + c)^2 + a^5)*sinh(d*x + c)^6 + 8*(7*a^5*cosh(d*x + c)^3 + 3
*a^5*cosh(d*x + c))*sinh(d*x + c)^5 + a^5 + 2*(35*a^5*cosh(d*x + c)^4 + 30
*a^5*cosh(d*x + c)^2 + 3*a^5)*sinh(d*x + c)^4 + 8*(7*a^5*cosh(d*x + c)^5 +
10*a^5*cosh(d*x + c)^3 + 3*a^5*cosh(d*x + c))*sinh(d*x + c)^3 + 4*(7*a^5*
cosh(d*x + c)^6 + 15*a^5*cosh(d*x + c)^4 + 9*a^5*cosh(d*x + c)^2 + a^5)*si
nh(d*x + c)^2 + 8*(a^5*cosh(d*x + c)^7 + 3*a^5*cosh(d*x + c)^5 + 3*a^5*cos
h(d*x + c)^3 + a^5*cosh(d*x + c))*sinh(d*x + c))*log(2*cosh(d*x + c)/(cosh
(d*x + c) - sinh(d*x + c))) + 16*(18*a^5*cosh(d*x + c)^5 + 27*a^5*cosh(d*x
+ c)^3 + 11*a^5*cosh(d*x + c))*sinh(d*x + c))/(d*cosh(d*x + c)^8 + 8*d*co
sh(d*x + c)*sinh(d*x + c)^7 + d*sinh(d*x + c)^8 + 4*d*cosh(d*x + c)^6 + 4*
(7*d*cosh(d*x + c)^2 + d)*sinh(d*x + c)^6 + 8*(7*d*cosh(d*x + c)^3 + 3*d*c
osh(d*x + c))*sinh(d*x + c)^5 + 6*d*cosh(d*x + c)^4 + 2*(35*d*cosh(d*x + c
)^4 + 30*d*cosh(d*x + c)^2 + 3*d)*sinh(d*x + c)^4 + 8*(7*d*cosh(d*x + c...

```

### Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.95

$$\int (a + a \tanh(c + dx))^5 dx$$

$$= \begin{cases} 32a^5x - \frac{16a^5 \log(\tanh(c+dx)+1)}{d} - \frac{a^5 \tanh^4(c+dx)}{4d} - \frac{5a^5 \tanh^3(c+dx)}{3d} - \frac{11a^5 \tanh^2(c+dx)}{2d} - \frac{15a^5 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^5 & \text{otherwise} \end{cases}$$

input

```
integrate((a+a*tanh(d*x+c))**5,x)
```

output

```

Piecewise(((32*a**5*x - 16*a**5*log(tanh(c + d*x) + 1)/d - a**5*tanh(c + d*
x)**4/(4*d) - 5*a**5*tanh(c + d*x)**3/(3*d) - 11*a**5*tanh(c + d*x)**2/(2*
d) - 15*a**5*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**5, True))

```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 302 vs.  $2(96) = 192$ .

Time = 0.13 (sec) , antiderivative size = 302, normalized size of antiderivative = 3.02

$$\int (a + a \tanh(c + dx))^5 dx$$

$$= \frac{5}{3} a^5 \left( 3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)} \right)$$

$$+ a^5 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{4(e^{(-2dx-2c)} + e^{(-4dx-4c)} + e^{(-6dx-6c)})}{d(4e^{(-2dx-2c)} + 6e^{(-4dx-4c)} + 4e^{(-6dx-6c)} + e^{(-8dx-8c)} + 1)} \right)$$

$$+ 10a^5 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 10a^5 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^5 x + \frac{5a^5 \log(\cosh(dx + c))}{d}$$

input `integrate((a+a*tanh(d*x+c))^5,x, algorithm="maxima")`

output

```
5/3*a^5*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + 2)/(d*(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) + 1))) + a^5*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 4*(e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c))/(d*(4*e^(-2*d*x - 2*c) + 6*e^(-4*d*x - 4*c) + 4*e^(-6*d*x - 6*c) + e^(-8*d*x - 8*c) + 1))) + 10*a^5*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 2*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c) + 1))) + 10*a^5*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^5*x + 5*a^5*log(cosh(d*x + c))/d
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.85

$$\int (a + a \tanh(c + dx))^5 dx$$

$$= \frac{4 \left( 12 a^5 \log(e^{(2dx+2c)} + 1) + \frac{48 a^5 e^{(6dx+6c)} + 108 a^5 e^{(4dx+4c)} + 88 a^5 e^{(2dx+2c)} + 25 a^5}{(e^{(2dx+2c)} + 1)^4} \right)}{3d}$$

input `integrate((a+a*tanh(d*x+c))^5,x, algorithm="giac")`

output 
$$\frac{4}{3} * (12 * a^5 * \log(e^{(2 * d * x + 2 * c)} + 1) + (48 * a^5 * e^{(6 * d * x + 6 * c)} + 108 * a^5 * e^{(4 * d * x + 4 * c)} + 88 * a^5 * e^{(2 * d * x + 2 * c)} + 25 * a^5) / (e^{(2 * d * x + 2 * c)} + 1)^4) / d$$

### Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.65

$$\int (a + a \tanh(c + dx))^5 dx = 32 a^5 x - \frac{a^5 (192 \ln(\tanh(c + dx) + 1) + 180 \tanh(c + dx) + 66 \tanh(c + dx)^2 + 20 \tanh(c + dx)^3 + 3 \tanh(c + dx)^4)}{12 d}$$

input `int((a + a*tanh(c + d*x))^5,x)`

output 
$$32 * a^5 * x - (a^5 * (192 * \log(\tanh(c + d * x) + 1) + 180 * \tanh(c + d * x) + 66 * \tanh(c + d * x)^2 + 20 * \tanh(c + d * x)^3 + 3 * \tanh(c + d * x)^4)) / (12 * d)$$

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.11

$$\int (a + a \tanh(c + dx))^5 dx = \frac{4a^5 (12e^{8dx+8c} \log(e^{2dx+2c} + 1) - 12e^{8dx+8c} + 48e^{6dx+6c} \log(e^{2dx+2c} + 1) + 72e^{4dx+4c} \log(e^{2dx+2c} + 1) + 36e^{4dx+4c} - 36e^{8dx+8c} + 36e^{6dx+6c} + 36e^{4dx+4c})}{3d(e^{8dx+8c} + 4e^{6dx+6c} + 6e^{4dx+4c} + 4e^{2dx+2c} + 1)}$$

input `int((a+a*tanh(d*x+c))^5,x)`

output

```
(4*a**5*(12*e**(8*c + 8*d*x)*log(e**(2*c + 2*d*x) + 1) - 12*e**(8*c + 8*d*x) + 48*e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x) + 1) + 72*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1) + 36*e**(4*c + 4*d*x) + 48*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1) + 40*e**(2*c + 2*d*x) + 12*log(e**(2*c + 2*d*x) + 1) + 13))/(3*d*(e**(8*c + 8*d*x) + 4*e**(6*c + 6*d*x) + 6*e**(4*c + 4*d*x) + 4*e**(2*c + 2*d*x) + 1))
```

### 3.42 $\int (a + a \tanh(c + dx))^4 dx$

Optimal result . . . . .	396
Mathematica [A] (verified) . . . . .	396
Rubi [A] (verified) . . . . .	397
Maple [A] (verified) . . . . .	399
Fricas [B] (verification not implemented) . . . . .	400
Sympy [A] (verification not implemented) . . . . .	400
Maxima [B] (verification not implemented) . . . . .	401
Giac [A] (verification not implemented) . . . . .	401
Mupad [B] (verification not implemented) . . . . .	402
Reduce [B] (verification not implemented) . . . . .	402

#### Optimal result

Integrand size = 12, antiderivative size = 77

$$\int (a + a \tanh(c + dx))^4 dx = 8a^4x + \frac{8a^4 \log(\cosh(c + dx))}{d} - \frac{4a^4 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^3}{3d} - \frac{(a^2 + a^2 \tanh(c + dx))^2}{d}$$

output

```
8*a^4*x+8*a^4*ln(cosh(d*x+c))/d-4*a^4*tanh(d*x+c)/d-1/3*a*(a+a*tanh(d*x+c))^3/d-(a^2+a^2*tanh(d*x+c))^2/d
```

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.66

$$\int (a + a \tanh(c + dx))^4 dx = \frac{a^4(4 + 24 \log(1 - \tanh(c + dx)) + 21 \tanh(c + dx) + 6 \tanh^2(c + dx) + \tanh^3(c + dx))}{3d}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^4,x]
```

output

```
-1/3*(a^4*(4 + 24*Log[1 - Tanh[c + d*x]] + 21*Tanh[c + d*x] + 6*Tanh[c + d
*x]^2 + Tanh[c + d*x]^3))/d
```

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$ , Rules used = {3042, 3959, 3042, 3959, 3042, 3958, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh(c + dx) + a)^4 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ia \tan(ic + idx))^4 dx \\
 & \quad \downarrow \text{3959} \\
 & 2a \int (\tanh(c + dx)a + a)^3 dx - \frac{a(a \tanh(c + dx) + a)^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a(a \tanh(c + dx) + a)^3}{3d} + 2a \int (a - ia \tan(ic + idx))^3 dx \\
 & \quad \downarrow \text{3959} \\
 & 2a \left( 2a \int (\tanh(c + dx)a + a)^2 dx - \frac{a(a \tanh(c + dx) + a)^2}{2d} \right) - \frac{a(a \tanh(c + dx) + a)^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{a(a \tanh(c + dx) + a)^3}{3d} + 2a \left( -\frac{a(a \tanh(c + dx) + a)^2}{2d} + 2a \int (a - ia \tan(ic + idx))^2 dx \right) \\
 & \quad \downarrow \text{3958} \\
 & -\frac{a(a \tanh(c + dx) + a)^3}{3d} + \\
 & 2a \left( -\frac{a(a \tanh(c + dx) + a)^2}{2d} + 2a \left( -2ia^2 \int i \tanh(c + dx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \right) \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& 2a \left( 2a \left( 2a^2 \int \tanh(c+dx) dx - \frac{a^2 \tanh(c+dx)}{d} + 2a^2 x \right) - \frac{a(a \tanh(c+dx) + a)^2}{2d} \right) - \\
& \quad \frac{a(a \tanh(c+dx) + a)^3}{3d} \\
& \downarrow 3042 \\
& - \frac{a(a \tanh(c+dx) + a)^3}{3d} + \\
& 2a \left( - \frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( 2a^2 \int -i \tan(ic+idx) dx - \frac{a^2 \tanh(c+dx)}{d} + 2a^2 x \right) \right) \\
& \downarrow 26 \\
& - \frac{a(a \tanh(c+dx) + a)^3}{3d} + \\
& 2a \left( - \frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( -2ia^2 \int \tan(ic+idx) dx - \frac{a^2 \tanh(c+dx)}{d} + 2a^2 x \right) \right) \\
& \downarrow 3956 \\
& 2a \left( 2a \left( - \frac{a^2 \tanh(c+dx)}{d} + \frac{2a^2 \log(\cosh(c+dx))}{d} + 2a^2 x \right) - \frac{a(a \tanh(c+dx) + a)^2}{2d} \right) - \\
& \quad \frac{a(a \tanh(c+dx) + a)^3}{3d}
\end{aligned}$$

input `Int[(a + a*Tanh[c + d*x])^4,x]`

output `-1/3*(a*(a + a*Tanh[c + d*x])^3)/d + 2*a*(-1/2*(a*(a + a*Tanh[c + d*x])^2)/d + 2*a*(2*a^2*x + (2*a^2*Log[Cosh[c + d*x]]))/d - (a^2*Tanh[c + d*x])/d)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.62

method	result
derivativedivides	$\frac{a^4 \left( -\frac{\tanh(dx+c)^3}{3} - 2 \tanh(dx+c)^2 - 7 \tanh(dx+c) - 8 \ln(\tanh(dx+c)-1) \right)}{d}$
default	$\frac{a^4 \left( -\frac{\tanh(dx+c)^3}{3} - 2 \tanh(dx+c)^2 - 7 \tanh(dx+c) - 8 \ln(\tanh(dx+c)-1) \right)}{d}$
parallelrisch	$-\frac{\tanh(dx+c)^3 a^4 + 6 \tanh(dx+c)^2 a^4 + 24 \ln(1 - \tanh(dx+c)) a^4 + 21 a^4 \tanh(dx+c)}{3d}$
risch	$-\frac{16a^4c}{d} + \frac{4a^4(18e^{4dx+4c} + 27e^{2dx+2c} + 11)}{3d(e^{2dx+2c} + 1)^3} + \frac{8a^4 \ln(e^{2dx+2c} + 1)}{d}$
parts	$a^4 x + \frac{a^4 \left( -\frac{\tanh(dx+c)^3}{3} - \tanh(dx+c) - \frac{\ln(\tanh(dx+c)-1)}{2} + \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{4a^4 \ln(\cosh(dx+c))}{d} + \frac{6a^4 (-\tanh(dx+c))}{d}$

input `int((a+a*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*a^4*(-1/3*tanh(d*x+c)^3-2*tanh(d*x+c)^2-7*tanh(d*x+c)-8*ln(tanh(d*x+c)-1))`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 562 vs.  $2(75) = 150$ .

Time = 0.08 (sec) , antiderivative size = 562, normalized size of antiderivative = 7.30

$$\int (a + a \tanh(c + dx))^4 dx = \text{Too large to display}$$

input `integrate((a+a*tanh(d*x+c))^4,x, algorithm="fricas")`

output

$$\begin{aligned} & \frac{4}{3}(18a^4 \cosh(dx+c)^4 + 72a^4 \cosh(dx+c) \sinh(dx+c)^3 + 18a^4 \sinh(dx+c)^4 \\ & + 27a^4 \cosh(dx+c)^2 + 11a^4 + 27(4a^4 \cosh(dx+c)^2 + a^4) \sinh(dx+c)^2 + 6(a^4 \cosh(dx+c)^6 + 6a^4 \cosh(dx+c) \\ & ) \sinh(dx+c)^5 + a^4 \sinh(dx+c)^6 + 3a^4 \cosh(dx+c)^4 + 3a^4 \cosh(dx+c)^2 + 3(5a^4 \cosh(dx+c)^2 + a^4) \sinh(dx+c)^4 + a^4 + 4( \\ & 5a^4 \cosh(dx+c)^3 + 3a^4 \cosh(dx+c)) \sinh(dx+c)^3 + 3(5a^4 \cosh(dx+c)^4 + 6a^4 \cosh(dx+c)^2 + a^4) \sinh(dx+c)^2 + 6(a^4 \cosh \\ & (dx+c)^5 + 2a^4 \cosh(dx+c)^3 + a^4 \cosh(dx+c)) \sinh(dx+c) \log\left(\frac{2 \cosh(dx+c)}{\cosh(dx+c) - \sinh(dx+c)}\right) + 18(4a^4 \cosh(dx+c)^3 \\ & + 3a^4 \cosh(dx+c)) \sinh(dx+c) / (d \cosh(dx+c)^6 + 6d \cosh(dx+c) \sinh(dx+c)^5 + d \sinh(dx+c)^6 + 3d \cosh(dx+c)^4 + 3(5d \cosh(dx+c)^2 + d) \sinh(dx+c)^4 + 4(5d \cosh(dx+c)^3 + 3d \cosh(dx+c)) \sinh(dx+c)^3 + 3d \cosh(dx+c)^2 + 3(5d \cosh(dx+c)^4 + 6d \cosh(dx+c)^2 + d) \sinh(dx+c)^2 + 6(d \cosh(dx+c)^5 + 2d \cosh(dx+c) \sinh(dx+c)^3 + d \cosh(dx+c)) \sinh(dx+c) + d) \end{aligned}$$
**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.99

$$\begin{aligned} & \int (a + a \tanh(c + dx))^4 dx \\ & = \begin{cases} 16a^4 x - \frac{8a^4 \log(\tanh(c+dx)+1)}{d} - \frac{a^4 \tanh^3(c+dx)}{3d} - \frac{2a^4 \tanh^2(c+dx)}{d} - \frac{7a^4 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^4 & \text{otherwise} \end{cases} \end{aligned}$$

input `integrate((a+a*tanh(d*x+c))**4,x)`

output

```
Piecewise((16*a**4*x - 8*a**4*log(tanh(c + d*x) + 1)/d - a**4*tanh(c + d*x)
)**3/(3*d) - 2*a**4*tanh(c + d*x)**2/d - 7*a**4*tanh(c + d*x)/d, Ne(d, 0))
, (x*(a*tanh(c) + a)**4, True))
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs.  $2(75) = 150$ .

Time = 0.14 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.55

$$\int (a + a \tanh(c + dx))^4 dx$$

$$= \frac{1}{3} a^4 \left( 3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)} \right)$$

$$+ 4a^4 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 6a^4 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^4 x + \frac{4a^4 \log(\cosh(dx + c))}{d}$$

input

```
integrate((a+a*tanh(d*x+c))^4,x, algorithm="maxima")
```

output

```
1/3*a^4*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + 2)/(d*
(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) + 1))) + 4*a^4
*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 2*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d
*x - 2*c) + e^(-4*d*x - 4*c) + 1))) + 6*a^4*(x + c/d - 2/(d*(e^(-2*d*x - 2
*c) + 1))) + a^4*x + 4*a^4*log(cosh(d*x + c))/d
```

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.92

$$\int (a + a \tanh(c + dx))^4 dx = \frac{4 \left( 6a^4 \log(e^{(2dx+2c)} + 1) + \frac{18a^4 e^{(4dx+4c)} + 27a^4 e^{(2dx+2c)} + 11a^4}{(e^{(2dx+2c)} + 1)^3} \right)}{3d}$$

input

```
integrate((a+a*tanh(d*x+c))^4,x, algorithm="giac")
```

output  $\frac{4}{3}*(6*a^4*\log(e^{(2*d*x + 2*c)} + 1) + (18*a^4*e^{(4*d*x + 4*c)} + 27*a^4*e^{(2*d*x + 2*c)} + 11*a^4)/(e^{(2*d*x + 2*c)} + 1)^3)/d$

### Mupad [B] (verification not implemented)

Time = 1.97 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.69

$$\int (a + a \tanh(c + dx))^4 dx = 16 a^4 x - \frac{a^4 (24 \ln(\tanh(c + dx) + 1) + 21 \tanh(c + dx) + 6 \tanh(c + dx)^2 + \tanh(c + dx)^3)}{3d}$$

input `int((a + a*tanh(c + d*x))^4,x)`

output  $16*a^4*x - (a^4*(24*\log(\tanh(c + d*x) + 1) + 21*\tanh(c + d*x) + 6*\tanh(c + d*x)^2 + \tanh(c + d*x)^3))/(3*d)$

### Reduce [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.10

$$\int (a + a \tanh(c + dx))^4 dx = \frac{4a^4(6e^{6dx+6c}\log(e^{2dx+2c} + 1) - 6e^{6dx+6c} + 18e^{4dx+4c}\log(e^{2dx+2c} + 1) + 18e^{2dx+2c}\log(e^{2dx+2c} + 1) + 9e^{2dx+2c})}{3d(e^{6dx+6c} + 3e^{4dx+4c} + 3e^{2dx+2c} + 1)}$$

input `int((a+a*tanh(d*x+c))^4,x)`

output  $(4*a**4*(6*e**(6*c + 6*d*x)*\log(e**(2*c + 2*d*x) + 1) - 6*e**(6*c + 6*d*x) + 18*e**(4*c + 4*d*x)*\log(e**(2*c + 2*d*x) + 1) + 18*e**(2*c + 2*d*x)*\log(e**(2*c + 2*d*x) + 1) + 9*e**(2*c + 2*d*x) + 6*\log(e**(2*c + 2*d*x) + 1) + 5))/(3*d*(e**(6*c + 6*d*x) + 3*e**(4*c + 4*d*x) + 3*e**(2*c + 2*d*x) + 1))$

### 3.43 $\int (a + a \tanh(c + dx))^3 dx$

Optimal result . . . . .	403
Mathematica [A] (verified) . . . . .	403
Rubi [A] (verified) . . . . .	404
Maple [A] (verified) . . . . .	406
Fricas [B] (verification not implemented) . . . . .	406
Sympy [A] (verification not implemented) . . . . .	407
Maxima [B] (verification not implemented) . . . . .	407
Giac [A] (verification not implemented) . . . . .	408
Mupad [B] (verification not implemented) . . . . .	408
Reduce [B] (verification not implemented) . . . . .	409

#### Optimal result

Integrand size = 12, antiderivative size = 56

$$\int (a + a \tanh(c + dx))^3 dx = 4a^3x + \frac{4a^3 \log(\cosh(c + dx))}{d} - \frac{2a^3 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^2}{2d}$$

output

```
4*a^3*x+4*a^3*ln(cosh(d*x+c))/d-2*a^3*tanh(d*x+c)/d-1/2*a*(a+a*tanh(d*x+c))^2/d
```

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.71

$$\int (a + a \tanh(c + dx))^3 dx = -\frac{a^3(8 \log(1 - \tanh(c + dx)) + 6 \tanh(c + dx) + \tanh^2(c + dx))}{2d}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^3,x]
```

output

$$\frac{-1/2*(a^3*(8*\text{Log}[1 - \text{Tanh}[c + d*x]] + 6*\text{Tanh}[c + d*x] + \text{Tanh}[c + d*x]^2))/d}{d}$$
**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3959, 3042, 3958, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \tanh(c + dx) + a)^3 dx \\ & \quad \downarrow \text{3042} \\ & \int (a - ia \tan(ic + id x))^3 dx \\ & \quad \downarrow \text{3959} \\ & 2a \int (\tanh(c + dx)a + a)^2 dx - \frac{a(a \tanh(c + dx) + a)^2}{2d} \\ & \quad \downarrow \text{3042} \\ & -\frac{a(a \tanh(c + dx) + a)^2}{2d} + 2a \int (a - ia \tan(ic + id x))^2 dx \\ & \quad \downarrow \text{3958} \\ & -\frac{a(a \tanh(c + dx) + a)^2}{2d} + 2a \left( -2ia^2 \int i \tanh(c + dx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \right) \\ & \quad \downarrow \text{26} \\ & 2a \left( 2a^2 \int \tanh(c + dx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \right) - \frac{a(a \tanh(c + dx) + a)^2}{2d} \\ & \quad \downarrow \text{3042} \\ & -\frac{a(a \tanh(c + dx) + a)^2}{2d} + 2a \left( 2a^2 \int -i \tan(ic + id x) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \right) \\ & \quad \downarrow \text{26} \end{aligned}$$

$$-\frac{a(a \tanh(c+dx) + a)^2}{2d} + 2a \left( -2ia^2 \int \tan(ic + idx) dx - \frac{a^2 \tanh(c+dx)}{d} + 2a^2 x \right)$$

↓ 3956

$$2a \left( -\frac{a^2 \tanh(c+dx)}{d} + \frac{2a^2 \log(\cosh(c+dx))}{d} + 2a^2 x \right) - \frac{a(a \tanh(c+dx) + a)^2}{2d}$$

input `Int[(a + a*Tanh[c + d*x])^3,x]`

output `-1/2*(a*(a + a*Tanh[c + d*x])^2)/d + 2*a*(2*a^2*x + (2*a^2*Log[Cosh[c + d*x]])/d - (a^2*Tanh[c + d*x])/d)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

### Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.68

method	result
derivativedivides	$\frac{a^3 \left( -\frac{\tanh(dx+c)^2}{2} - 3 \tanh(dx+c) - 4 \ln(\tanh(dx+c)-1) \right)}{d}$
default	$\frac{a^3 \left( -\frac{\tanh(dx+c)^2}{2} - 3 \tanh(dx+c) - 4 \ln(\tanh(dx+c)-1) \right)}{d}$
parallelrisch	$-\frac{\tanh(dx+c)^2 a^3 + 8 \ln(1-\tanh(dx+c)) a^3 + 6 a^3 \tanh(dx+c)}{2d}$
risch	$-\frac{8a^3 c}{d} + \frac{2a^3 (4e^{2dx+2c} + 3)}{d(e^{2dx+2c} + 1)^2} + \frac{4a^3 \ln(e^{2dx+2c} + 1)}{d}$
parts	$a^3 x + \frac{a^3 \left( -\frac{\tanh(dx+c)^2}{2} - \frac{\ln(\tanh(dx+c)-1)}{2} - \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{3a^3 \ln(\cosh(dx+c))}{d} + \frac{3a^3 (-\tanh(dx+c)-1)}{d}$

input `int((a+a*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-1/2*tanh(d*x+c)^2-3*tanh(d*x+c)-4*ln(tanh(d*x+c)-1))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(54) = 108.

Time = 0.09 (sec) , antiderivative size = 299, normalized size of antiderivative = 5.34

$$\int (a + a \tanh(c + dx))^3 dx$$

$$= \frac{2 \left( 4 a^3 \cosh(dx + c)^2 + 8 a^3 \cosh(dx + c) \sinh(dx + c) + 4 a^3 \sinh(dx + c)^2 + 3 a^3 + 2 (a^3 \cosh(dx + c) \right)}{d \cosh(dx + c)^4 + 4 d \cosh(dx + c) \sinh(dx + c)}$$

input `integrate((a+a*tanh(d*x+c))^3,x, algorithm="fricas")`

output

```
2*(4*a^3*cosh(d*x + c)^2 + 8*a^3*cosh(d*x + c)*sinh(d*x + c) + 4*a^3*sinh(
d*x + c)^2 + 3*a^3 + 2*(a^3*cosh(d*x + c)^4 + 4*a^3*cosh(d*x + c)*sinh(d*x
+ c)^3 + a^3*sinh(d*x + c)^4 + 2*a^3*cosh(d*x + c)^2 + a^3 + 2*(3*a^3*cos
h(d*x + c)^2 + a^3)*sinh(d*x + c)^2 + 4*(a^3*cosh(d*x + c)^3 + a^3*cosh(d*
x + c))*sinh(d*x + c))*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c))
))/(d*cosh(d*x + c)^4 + 4*d*cosh(d*x + c)*sinh(d*x + c)^3 + d*sinh(d*x + c
)^4 + 2*d*cosh(d*x + c)^2 + 2*(3*d*cosh(d*x + c)^2 + d)*sinh(d*x + c)^2 +
4*(d*cosh(d*x + c)^3 + d*cosh(d*x + c))*sinh(d*x + c) + d)
```

**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09

$$\int (a + a \tanh(c + dx))^3 dx$$

$$= \begin{cases} 8a^3x - \frac{4a^3 \log(\tanh(c+dx)+1)}{d} - \frac{a^3 \tanh^2(c+dx)}{2d} - \frac{3a^3 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^3 & \text{otherwise} \end{cases}$$

input

```
integrate((a+a*tanh(d*x+c))**3,x)
```

output

```
Piecewise((8*a**3*x - 4*a**3*log(tanh(c + d*x) + 1)/d - a**3*tanh(c + d*x)
**2/(2*d) - 3*a**3*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**3, True
))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 116 vs.  $2(54) = 108$ .

Time = 0.13 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.07

$$\int (a + a \tanh(c + dx))^3 dx$$

$$= a^3 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 3a^3 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^3x + \frac{3a^3 \log(\cosh(dx + c))}{d}$$



input `integrate((a+a*tanh(d*x+c))^3,x, algorithm="maxima")`

output `a^3*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 2*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c) + 1))) + 3*a^3*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^3*x + 3*a^3*log(cosh(d*x + c))/d`

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.02

$$\int (a + a \tanh(c + dx))^3 dx = \frac{2 \left( 2 a^3 \log(e^{2dx+2c} + 1) + \frac{4 a^3 e^{2dx+2c} + 3 a^3}{(e^{2dx+2c} + 1)^2} \right)}{d}$$

input `integrate((a+a*tanh(d*x+c))^3,x, algorithm="giac")`

output `2*(2*a^3*log(e^(2*d*x + 2*c) + 1) + (4*a^3*e^(2*d*x + 2*c) + 3*a^3)/(e^(2*d*x + 2*c) + 1)^2)/d`

### Mupad [B] (verification not implemented)

Time = 1.96 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.77

$$\begin{aligned} \int (a + a \tanh(c + dx))^3 dx \\ = 8 a^3 x - \frac{a^3 (8 \ln(\tanh(c + dx) + 1) + 6 \tanh(c + dx) + \tanh(c + dx)^2)}{2d} \end{aligned}$$

input `int((a + a*tanh(c + d*x))^3,x)`

output `8*a^3*x - (a^3*(8*log(tanh(c + d*x) + 1) + 6*tanh(c + d*x) + tanh(c + d*x)^2))/(2*d)`

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.02

$$\int (a + a \tanh(c + dx))^3 dx$$

$$= \frac{2a^3 (2e^{4dx+4c} \log(e^{2dx+2c} + 1) - 2e^{4dx+4c} + 4e^{2dx+2c} \log(e^{2dx+2c} + 1) + 2 \log(e^{2dx+2c} + 1) + 1)}{d(e^{4dx+4c} + 2e^{2dx+2c} + 1)}$$

input `int((a+a*tanh(d*x+c))^3,x)`output `(2*a**3*(2*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1) - 2*e**(4*c + 4*d*x) + 4*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1) + 2*log(e**(2*c + 2*d*x) + 1) + 1))/(d*(e**(4*c + 4*d*x) + 2*e**(2*c + 2*d*x) + 1))`

### 3.44 $\int (a + a \tanh(c + dx))^2 dx$

Optimal result	410
Mathematica [A] (verified)	410
Rubi [A] (verified)	411
Maple [A] (verified)	412
Fricas [B] (verification not implemented)	413
Sympy [A] (verification not implemented)	413
Maxima [A] (verification not implemented)	414
Giac [A] (verification not implemented)	414
Mupad [B] (verification not implemented)	414
Reduce [B] (verification not implemented)	415

#### Optimal result

Integrand size = 12, antiderivative size = 36

$$\int (a + a \tanh(c + dx))^2 dx = 2a^2 x + \frac{2a^2 \log(\cosh(c + dx))}{d} - \frac{a^2 \tanh(c + dx)}{d}$$

output

```
2*a^2*x+2*a^2*ln(cosh(d*x+c))/d-a^2*tanh(d*x+c)/d
```

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81

$$\int (a + a \tanh(c + dx))^2 dx = \frac{a(-2a \log(1 - \tanh(c + dx)) - a \tanh(c + dx))}{d}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^2,x]
```

output

```
(a*(-2*a*Log[1 - Tanh[c + d*x]] - a*Tanh[c + d*x]))/d
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3958, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \tanh(c + dx) + a)^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ia \tan(ic + idx))^2 dx \\
 & \quad \downarrow \text{3958} \\
 & -2ia^2 \int i \tanh(c + dx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \\
 & \quad \downarrow \text{26} \\
 & 2a^2 \int \tanh(c + dx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \\
 & \quad \downarrow \text{3042} \\
 & 2a^2 \int -i \tan(ic + idx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \\
 & \quad \downarrow \text{26} \\
 & -2ia^2 \int \tan(ic + idx) dx - \frac{a^2 \tanh(c + dx)}{d} + 2a^2 x \\
 & \quad \downarrow \text{3956} \\
 & -\frac{a^2 \tanh(c + dx)}{d} + \frac{2a^2 \log(\cosh(c + dx))}{d} + 2a^2 x
 \end{aligned}$$

input `Int[(a + a*Tanh[c + d*x])^2,x]`

output `2*a^2*x + (2*a^2*Log[Cosh[c + d*x]])/d - (a^2*Tanh[c + d*x])/d`

## Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

## Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

method	result	size
derivativdivides	$\frac{a^2(-\tanh(dx+c)-2\ln(\tanh(dx+c)-1))}{d}$	28
default	$\frac{a^2(-\tanh(dx+c)-2\ln(\tanh(dx+c)-1))}{d}$	28
parallelrisch	$-\frac{2\ln(1-\tanh(dx+c))a^2+a^2\tanh(dx+c)}{d}$	33
risch	$-\frac{4a^2c}{d} + \frac{2a^2}{d(e^{2dx+2c}+1)} + \frac{2a^2\ln(e^{2dx+2c}+1)}{d}$	52
parts	$a^2x + \frac{a^2\left(-\tanh(dx+c) - \frac{\ln(\tanh(\frac{dx+c}{2})-1)}{2} + \frac{\ln(\tanh(\frac{dx+c}{2})+1)}{2}\right)}{d} + \frac{2a^2\ln(\cosh(dx+c))}{d}$	60

input `int((a+a*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*a^2*(-tanh(d*x+c)-2*ln(tanh(d*x+c)-1))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 117 vs.  $2(36) = 72$ .

Time = 0.09 (sec) , antiderivative size = 117, normalized size of antiderivative = 3.25

$$\int (a + a \tanh(c + dx))^2 dx$$

$$= \frac{2 \left( a^2 + (a^2 \cosh(dx + c))^2 + 2 a^2 \cosh(dx + c) \sinh(dx + c) + a^2 \sinh(dx + c)^2 + a^2 \right) \log \left( \frac{2 \cosh(dx + c) - \sinh(dx + c)}{\cosh(dx + c) - \sinh(dx + c)} \right)}{d \cosh(dx + c)^2 + 2 d \cosh(dx + c) \sinh(dx + c) + d \sinh(dx + c)^2 + d}$$

input `integrate((a+a*tanh(d*x+c))^2,x, algorithm="fricas")`

output `2*(a^2 + (a^2*cosh(d*x + c)^2 + 2*a^2*cosh(d*x + c)*sinh(d*x + c) + a^2*sinh(d*x + c)^2 + a^2)*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c))))/(d*cosh(d*x + c)^2 + 2*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x + c)^2 + d)`

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.22

$$\int (a + a \tanh(c + dx))^2 dx = \begin{cases} 4a^2x - \frac{2a^2 \log(\tanh(c+dx)+1)}{d} - \frac{a^2 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^2 & \text{otherwise} \end{cases}$$

input `integrate((a+a*tanh(d*x+c))**2,x)`

output `Piecewise((4*a**2*x - 2*a**2*log(tanh(c + d*x) + 1)/d - a**2*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.39

$$\int (a + a \tanh(c + dx))^2 dx = a^2 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^2 x + \frac{2a^2 \log(\cosh(dx + c))}{d}$$

input `integrate((a+a*tanh(d*x+c))^2,x, algorithm="maxima")`output `a^2*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^2*x + 2*a^2*log(cosh(d*x + c))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

$$\int (a + a \tanh(c + dx))^2 dx = \frac{2 \left( a^2 \log(e^{(2dx+2c)} + 1) + \frac{a^2}{e^{(2dx+2c)} + 1} \right)}{d}$$

input `integrate((a+a*tanh(d*x+c))^2,x, algorithm="giac")`output `2*(a^2*log(e^(2*d*x + 2*c) + 1) + a^2/(e^(2*d*x + 2*c) + 1))/d`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int (a + a \tanh(c + dx))^2 dx = 4a^2 x - \frac{a^2 (2 \ln(\tanh(c + dx) + 1) + \tanh(c + dx))}{d}$$

input `int((a + a*tanh(c + d*x))^2,x)`output `4*a^2*x - (a^2*(2*log(tanh(c + d*x) + 1) + tanh(c + d*x)))/d`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.00

$$\int (a + a \tanh(c + dx))^2 dx = \frac{2a^2 (e^{2dx+2c} \log(e^{2dx+2c} + 1) - e^{2dx+2c} + \log(e^{2dx+2c} + 1))}{d(e^{2dx+2c} + 1)}$$

input `int((a+a*tanh(d*x+c))^2,x)`

output `(2*a**2*(e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1) - e**(2*c + 2*d*x) + 1  
og(e**(2*c + 2*d*x) + 1)))/(d*(e**(2*c + 2*d*x) + 1))`



### 3.45 $\int \frac{1}{a+a \tanh(c+dx)} dx$

Optimal result	416
Mathematica [A] (verified)	416
Rubi [A] (verified)	417
Maple [A] (verified)	418
Fricas [B] (verification not implemented)	418
Sympy [B] (verification not implemented)	419
Maxima [A] (verification not implemented)	419
Giac [A] (verification not implemented)	420
Mupad [B] (verification not implemented)	420
Reduce [B] (verification not implemented)	420

#### Optimal result

Integrand size = 12, antiderivative size = 28

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{x}{2a} - \frac{1}{2d(a + a \tanh(c + dx))}$$

output `1/2*x/a-1/2/d/(a+a*tanh(d*x+c))`

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{\operatorname{arctanh}(\tanh(c+dx))}{a} - \frac{1}{a+a \tanh(c+dx)} \frac{1}{2d}$$

input `Integrate[(a + a*Tanh[c + d*x])^(-1),x]`

output `(ArcTanh[Tanh[c + d*x]]/a - (a + a*Tanh[c + d*x])^(-1))/(2*d)`

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a \tanh(c + dx) + a} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{a - ia \tan(ic + idx)} dx$$

$$\downarrow \text{3960}$$

$$\frac{\int 1 dx}{2a} - \frac{1}{2d(a \tanh(c + dx) + a)}$$

$$\downarrow \text{24}$$

$$\frac{x}{2a} - \frac{1}{2d(a \tanh(c + dx) + a)}$$

input `Int[(a + a*Tanh[c + d*x])^(-1),x]`

output `x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x]))`

**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960

```
Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a +
b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^
(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

method	result	size
risch	$\frac{x}{2a} - \frac{e^{-2dx-2c}}{4ad}$	25
parallelrisch	$\frac{-1+\tanh(dx+c)xd+dx}{2da(\tanh(dx+c)+1)}$	33
derivativedivides	$-\frac{1}{2(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{4} - \frac{\ln(\tanh(dx+c)-1)}{4}$	43
default	$-\frac{1}{2(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{4} - \frac{\ln(\tanh(dx+c)-1)}{4}$	43

input

```
int(1/(a+a*tanh(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/2*x/a-1/4/a/d*exp(-2*d*x-2*c)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(24) = 48.

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.79

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{(2dx - 1) \cosh(dx + c) + (2dx + 1) \sinh(dx + c)}{4(ad \cosh(dx + c) + ad \sinh(dx + c))}$$

input

```
integrate(1/(a+a*tanh(d*x+c)),x, algorithm="fricas")
```

output

```
1/4*((2*d*x - 1)*cosh(d*x + c) + (2*d*x + 1)*sinh(d*x + c))/(a*d*cosh(d*x
+ c) + a*d*sinh(d*x + c))
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 73 vs.  $2(19) = 38$ .

Time = 0.36 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.61

$$\int \frac{1}{a + a \tanh(c + dx)} dx$$

$$= \begin{cases} \frac{dx \tanh(c+dx)}{2ad \tanh(c+dx)+2ad} + \frac{dx}{2ad \tanh(c+dx)+2ad} - \frac{1}{2ad \tanh(c+dx)+2ad} & \text{for } d \neq 0 \\ \frac{x}{a \tanh(c)+a} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+a*tanh(d*x+c)),x)`

output `Piecewise((d*x*tanh(c + d*x)/(2*a*d*tanh(c + d*x) + 2*a*d) + d*x/(2*a*d*tanh(c + d*x) + 2*a*d) - 1/(2*a*d*tanh(c + d*x) + 2*a*d), Ne(d, 0)), (x/(a*tanh(c) + a), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{dx + c}{2ad} - \frac{e^{(-2dx-2c)}}{4ad}$$

input `integrate(1/(a+a*tanh(d*x+c)),x, algorithm="maxima")`

output `1/2*(d*x + c)/(a*d) - 1/4*e^(-2*d*x - 2*c)/(a*d)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{2(dx+c)}{a} - \frac{e^{(-2dx-2c)}}{a} \frac{1}{4d}$$

input `integrate(1/(a+a*tanh(d*x+c)),x, algorithm="giac")`

output `1/4*(2*(d*x + c)/a - e^(-2*d*x - 2*c)/a)/d`

**Mupad [B] (verification not implemented)**

Time = 1.96 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{x}{2a} - \frac{1}{2ad (\tanh(c + dx) + 1)}$$

input `int(1/(a + a*tanh(c + d*x)),x)`

output `x/(2*a) - 1/(2*a*d*(tanh(c + d*x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.29

$$\int \frac{1}{a + a \tanh(c + dx)} dx = \frac{2e^{2dx+2c}dx - 1}{4e^{2dx+2c}ad}$$

input `int(1/(a+a*tanh(d*x+c)),x)`

output `(2*e**(2*c + 2*d*x)*d*x - 1)/(4*e**(2*c + 2*d*x)*a*d)`

### 3.46 $\int \frac{1}{(a+a \tanh(c+dx))^2} dx$

Optimal result	421
Mathematica [A] (verified)	421
Rubi [A] (verified)	422
Maple [A] (verified)	423
Fricas [B] (verification not implemented)	424
Sympy [B] (verification not implemented)	424
Maxima [A] (verification not implemented)	425
Giac [A] (verification not implemented)	425
Mupad [B] (verification not implemented)	425
Reduce [B] (verification not implemented)	426

#### Optimal result

Integrand size = 12, antiderivative size = 51

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = \frac{x}{4a^2} - \frac{1}{4d(a + a \tanh(c + dx))^2} - \frac{1}{4d(a^2 + a^2 \tanh(c + dx))}$$

output

$$1/4*x/a^2-1/4/d/(a+a*\tanh(d*x+c))^2-1/4/d/(a^2+a^2*\tanh(d*x+c))$$

#### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.92

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = -\frac{2 + \tanh(c + dx) - \operatorname{arctanh}(\tanh(c + dx))(1 + \tanh(c + dx))^2}{4a^2d(1 + \tanh(c + dx))^2}$$

input

$$\text{Integrate}[(a + a*\text{Tanh}[c + d*x])^{-2}, x]$$

output

$$-1/4*(2 + \text{Tanh}[c + d*x] - \text{ArcTanh}[\text{Tanh}[c + d*x]]*(1 + \text{Tanh}[c + d*x])^2)/(a^2*d*(1 + \text{Tanh}[c + d*x])^2)$$

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \tanh(c + dx) + a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - ia \tan(ic + idx))^2} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{\int \frac{1}{\tanh(c+dx)a+a} dx}{2a} - \frac{1}{4d(a \tanh(c + dx) + a)^2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{4d(a \tanh(c + dx) + a)^2} + \frac{\int \frac{1}{a-ia \tan(ic+idx)} dx}{2a} \\
 & \quad \downarrow \text{3960} \\
 & \frac{\int \frac{1 dx}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c + dx) + a)^2} \\
 & \quad \downarrow \text{24} \\
 & \frac{\frac{x}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c + dx) + a)^2}
 \end{aligned}$$

input `Int[(a + a*Tanh[c + d*x])^(-2),x]`

output `-1/4*1/(d*(a + a*Tanh[c + d*x])^2) + (x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x])))/(2*a)`

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

## Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

method	result	size
risch	$\frac{x}{4a^2} - \frac{e^{-2dx-2c}}{4da^2} - \frac{e^{-4dx-4c}}{16da^2}$	42
parallelrisch	$\frac{-2 + \tanh(dx+c)^2 x d + 2 \tanh(dx+c) x d + dx - \tanh(dx+c)}{4da^2(\tanh(dx+c)+1)^2}$	53
derivativedivides	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{8} - \frac{1}{4(\tanh(dx+c)+1)^2} - \frac{1}{4(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{8}}{da^2}$	55
default	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{8} - \frac{1}{4(\tanh(dx+c)+1)^2} - \frac{1}{4(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{8}}{da^2}$	55

input `int(1/(a+a*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/4*x/a^2-1/4/d/a^2*exp(-2*d*x-2*c)-1/16/d/a^2*exp(-4*d*x-4*c)`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 101 vs.  $2(45) = 90$ .

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.98

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx$$

$$= \frac{(4 dx - 1) \cosh(dx + c)^2 + 2(4 dx + 1) \cosh(dx + c) \sinh(dx + c) + (4 dx - 1) \sinh(dx + c)^2 - 4}{16(a^2 d \cosh(dx + c)^2 + 2 a^2 d \cosh(dx + c) \sinh(dx + c) + a^2 d \sinh(dx + c)^2)}$$

input `integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="fricas")`

output `1/16*((4*d*x - 1)*cosh(d*x + c)^2 + 2*(4*d*x + 1)*cosh(d*x + c)*sinh(d*x + c) + (4*d*x - 1)*sinh(d*x + c)^2 - 4)/(a^2*d*cosh(d*x + c)^2 + 2*a^2*d*cosh(d*x + c)*sinh(d*x + c) + a^2*d*sinh(d*x + c)^2)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 223 vs.  $2(41) = 82$ .

Time = 0.59 (sec) , antiderivative size = 223, normalized size of antiderivative = 4.37

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx$$

$$= \begin{cases} \frac{dx \tanh^2(c+dx)}{4a^2 d \tanh^2(c+dx)+8a^2 d \tanh(c+dx)+4a^2 d} + \frac{2dx \tanh(c+dx)}{4a^2 d \tanh^2(c+dx)+8a^2 d \tanh(c+dx)+4a^2 d} + \frac{dx}{4a^2 d \tanh^2(c+dx)+8a^2 d \tanh(c+dx)+4a^2 d} \\ \frac{x}{(a \tanh(c)+a)^2} \end{cases}$$

input `integrate(1/(a+a*tanh(d*x+c))**2,x)`

output `Piecewise((d*x*tanh(c + d*x)**2/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) + 2*d*x*tanh(c + d*x)/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) + d*x/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) - tanh(c + d*x)/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) - 2/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d), Ne(d, 0)), (x/(a*tanh(c) + a)**2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = \frac{dx + c}{4 a^2 d} - \frac{4 e^{(-2 dx - 2c)} + e^{(-4 dx - 4c)}}{16 a^2 d}$$

input `integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="maxima")`output `1/4*(d*x + c)/(a^2*d) - 1/16*(4*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c))/(a^2*d)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = -\frac{(4 e^{(2 dx + 2c)} + 1) e^{(-4 dx - 4c)}}{a^2} - \frac{4(dx+c)}{a^2} \frac{1}{16 d}$$

input `integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="giac")`output `-1/16*((4*e^(2*d*x + 2*c) + 1)*e^(-4*d*x - 4*c)/a^2 - 4*(d*x + c)/a^2)/d`**Mupad [B] (verification not implemented)**

Time = 1.96 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.80

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = \frac{x}{4 a^2} - \frac{e^{-2c-2dx}}{4 a^2 d} - \frac{e^{-4c-4dx}}{16 a^2 d}$$

input `int(1/(a + a*tanh(c + d*x))^2,x)`output `x/(4*a^2) - exp(- 2*c - 2*d*x)/(4*a^2*d) - exp(- 4*c - 4*d*x)/(16*a^2*d)`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + a \tanh(c + dx))^2} dx = \frac{4e^{4dx+4c} dx - 4e^{2dx+2c} - 1}{16e^{4dx+4c} a^2 d}$$

input `int(1/(a+a*tanh(d*x+c))^2,x)`

output `(4*e**(4*c + 4*d*x)*d*x - 4*e**(2*c + 2*d*x) - 1)/(16*e**(4*c + 4*d*x)*a**2*d)`

### 3.47 $\int \frac{1}{(a+a \tanh(c+dx))^3} dx$

Optimal result . . . . .	427
Mathematica [A] (verified) . . . . .	427
Rubi [A] (verified) . . . . .	428
Maple [A] (verified) . . . . .	429
Fricas [B] (verification not implemented) . . . . .	430
Sympy [B] (verification not implemented) . . . . .	430
Maxima [A] (verification not implemented) . . . . .	431
Giac [A] (verification not implemented) . . . . .	431
Mupad [B] (verification not implemented) . . . . .	432
Reduce [B] (verification not implemented) . . . . .	432

#### Optimal result

Integrand size = 12, antiderivative size = 73

$$\int \frac{1}{(a+a \tanh(c+dx))^3} dx = \frac{x}{8a^3} - \frac{1}{6d(a+a \tanh(c+dx))^3} - \frac{1}{8ad(a+a \tanh(c+dx))^2} - \frac{1}{8d(a^3+a^3 \tanh(c+dx))}$$

output

```
1/8*x/a^3-1/6/d/(a+a*tanh(d*x+c))^3-1/8/a/d/(a+a*tanh(d*x+c))^2-1/8/d/(a^3+a^3*tanh(d*x+c))
```

#### Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.81

$$\int \frac{1}{(a+a \tanh(c+dx))^3} dx = \frac{10+9 \tanh(c+dx)+3 \tanh^2(c+dx)-3 \arctanh(\tanh(c+dx))(1+\tanh(c+dx))^3}{24a^3d(1+\tanh(c+dx))^3}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^(-3),x]
```

output

```
-1/24*(10 + 9*Tanh[c + d*x] + 3*Tanh[c + d*x]^2 - 3*ArcTanh[Tanh[c + d*x]]
*(1 + Tanh[c + d*x])^3)/(a^3*d*(1 + Tanh[c + d*x])^3)
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3960, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \tanh(c + dx) + a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - ia \tan(ic + idx))^3} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{\int \frac{1}{(\tanh(c+dx)a+a)^2} dx}{2a} - \frac{1}{6d(a \tanh(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{6d(a \tanh(c + dx) + a)^3} + \frac{\int \frac{1}{(a-ia \tan(ic+idx))^2} dx}{2a} \\
 & \quad \downarrow \text{3960} \\
 & \frac{\int \frac{1}{\tanh(c+dx)a+a} dx}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{6d(a \tanh(c + dx) + a)^3} + \frac{-\frac{1}{4d(a \tanh(c+dx)+a)^2} + \frac{\int \frac{1}{a-ia \tan(ic+idx)} dx}{2a}}{2a} \\
 & \quad \downarrow \text{3960} \\
 & \frac{\int \frac{1 dx}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c + dx) + a)^3}
 \end{aligned}$$

$$\frac{\frac{x}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3}$$

input `Int[(a + a*Tanh[c + d*x])^(-3),x]`

output `-1/6*1/(d*(a + a*Tanh[c + d*x])^3) + (-1/4*1/(d*(a + a*Tanh[c + d*x])^2) + (x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x]))) / (2*a)) / (2*a)`

**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.81

method	result	size
risch	$\frac{x}{8a^3} - \frac{3e^{-2dx-2c}}{16a^3d} - \frac{3e^{-4dx-4c}}{32a^3d} - \frac{e^{-6dx-6c}}{48a^3d}$	59
derivativedivides	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{16} - \frac{1}{6(\tanh(dx+c)+1)^3} - \frac{1}{8(\tanh(dx+c)+1)^2} - \frac{1}{8(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{16}}{da^3}$	67
default	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{16} - \frac{1}{6(\tanh(dx+c)+1)^3} - \frac{1}{8(\tanh(dx+c)+1)^2} - \frac{1}{8(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{16}}{da^3}$	67
parallelrisch	$\frac{-10-3 \tanh(dx+c)^2-9 \tanh(dx+c)+9 \tanh(dx+c)^2xd+3dx+9 \tanh(dx+c)xd+3 \tanh(dx+c)^3xd}{24da^3(\tanh(dx+c)+1)^3}$	77

input `int(1/(a+a*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/8*x/a^3-3/16/a^3/d*exp(-2*d*x-2*c)-3/32/a^3/d*exp(-4*d*x-4*c)-1/48/a^3/d*exp(-6*d*x-6*c)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(65) = 130.

Time = 0.10 (sec) , antiderivative size = 160, normalized size of antiderivative = 2.19

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx$$

$$= \frac{2(6dx - 1) \cosh(dx + c)^3 + 6(6dx - 1) \cosh(dx + c) \sinh(dx + c)^2 + 2(6dx + 1) \sinh(dx + c)^3 + 3(6dx + 1) \cosh(dx + c) \sinh(dx + c)^2}{96(a^3 d \cosh(dx + c))^3 + 3a^3 d \cosh(dx + c)^2 \sinh(dx + c) + 3a^3 d \cosh(dx + c) \sinh(dx + c)^2 + a^3 d \sinh(dx + c)^3}$$

input `integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="fricas")`

output `1/96*(2*(6*d*x - 1)*cosh(d*x + c)^3 + 6*(6*d*x - 1)*cosh(d*x + c)*sinh(d*x + c)^2 + 2*(6*d*x + 1)*sinh(d*x + c)^3 + 3*(2*(6*d*x + 1)*cosh(d*x + c)^2 - 3)*sinh(d*x + c) - 27*cosh(d*x + c))/(a^3*d*cosh(d*x + c)^3 + 3*a^3*d*cosh(d*x + c)^2*sinh(d*x + c) + 3*a^3*d*cosh(d*x + c)*sinh(d*x + c)^2 + a^3*d*sinh(d*x + c)^3)`

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 430 vs. 2(60) = 120.

Time = 0.74 (sec) , antiderivative size = 430, normalized size of antiderivative = 5.89

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx$$

$$= \left\{ \begin{array}{l} \frac{3dx \tanh^3(c+dx)}{24a^3 d \tanh^3(c+dx) + 72a^3 d \tanh^2(c+dx) + 72a^3 d \tanh(c+dx) + 24a^3 d} + \frac{9dx \tanh^2(c+dx)}{24a^3 d \tanh^3(c+dx) + 72a^3 d \tanh^2(c+dx) + 72a^3 d \tanh(c+dx) + 24a^3 d} \\ \frac{x}{(a \tanh(c) + a)^3} \end{array} \right.$$

input `integrate(1/(a+a*tanh(d*x+c))**3,x)`

output `Piecewise((3*d*x*tanh(c + d*x)**3/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) + 9*d*x*tanh(c + d*x)**2/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) + 9*d*x*tanh(c + d*x)/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 3*d*x/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 3*tanh(c + d*x)**2/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 9*tanh(c + d*x)/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 10/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d), Ne(d, 0)), (x/(a*tanh(c) + a)**3, True))`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.77

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx = \frac{dx + c}{8a^3d} - \frac{18e^{(-2dx-2c)} + 9e^{(-4dx-4c)} + 2e^{(-6dx-6c)}}{96a^3d}$$

input `integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="maxima")`

output `1/8*(d*x + c)/(a^3*d) - 1/96*(18*e^(-2*d*x - 2*c) + 9*e^(-4*d*x - 4*c) + 2*e^(-6*d*x - 6*c))/(a^3*d)`

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx = -\frac{(18e^{(4dx+4c)}+9e^{(2dx+2c)}+2)e^{(-6dx-6c)}}{a^3} - \frac{12(dx+c)}{a^3}$$

input `integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="giac")`



output

$$-1/96*((18*e^(4*d*x + 4*c) + 9*e^(2*d*x + 2*c) + 2)*e^(-6*d*x - 6*c)/a^3 - 12*(d*x + c)/a^3)/d$$

**Mupad [B] (verification not implemented)**

Time = 1.98 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx = \frac{x}{8a^3} - \frac{3e^{-2c-2dx}}{16a^3d} - \frac{3e^{-4c-4dx}}{32a^3d} - \frac{e^{-6c-6dx}}{48a^3d}$$

input

$$\text{int}(1/(a + a*\tanh(c + d*x))^3,x)$$

output

$$x/(8*a^3) - (3*\exp(- 2*c - 2*d*x))/(16*a^3*d) - (3*\exp(- 4*c - 4*d*x))/(32*a^3*d) - \exp(- 6*c - 6*d*x)/(48*a^3*d)$$

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a + a \tanh(c + dx))^3} dx = \frac{12e^{6dx+6c}dx - 18e^{4dx+4c} - 9e^{2dx+2c} - 2}{96e^{6dx+6c}a^3d}$$

input

$$\text{int}(1/(a+a*\tanh(d*x+c))^3,x)$$

output

$$(12*e**(6*c + 6*d*x)*d*x - 18*e**(4*c + 4*d*x) - 9*e**(2*c + 2*d*x) - 2)/(96*e**(6*c + 6*d*x)*a**3*d)$$

**3.48**       $\int \frac{1}{(a+a \tanh(c+dx))^4} dx$

Optimal result . . . . .	433
Mathematica [A] (verified) . . . . .	434
Rubi [A] (verified) . . . . .	434
Maple [A] (verified) . . . . .	436
Fricas [B] (verification not implemented) . . . . .	437
Sympy [B] (verification not implemented) . . . . .	437
Maxima [A] (verification not implemented) . . . . .	438
Giac [A] (verification not implemented) . . . . .	439
Mupad [B] (verification not implemented) . . . . .	439
Reduce [B] (verification not implemented) . . . . .	439

**Optimal result**

Integrand size = 12, antiderivative size = 96

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx = \frac{x}{16a^4} - \frac{1}{8d(a + a \tanh(c + dx))^4} - \frac{1}{12ad(a + a \tanh(c + dx))^3} - \frac{1}{16d(a^2 + a^2 \tanh(c + dx))^2} - \frac{1}{16d(a^4 + a^4 \tanh(c + dx))}$$

```
output 1/16*x/a^4-1/8/d/(a+a*tanh(d*x+c))^4-1/12/a/d/(a+a*tanh(d*x+c))^3-1/16/d/(a^2+a^2*tanh(d*x+c))^2-1/16/d/(a^4+a^4*tanh(d*x+c))
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx$$

$$= \frac{a \left( \frac{\operatorname{arctanh}(\tanh(c+dx))}{16a^5} - \frac{1}{8a(a+a \tanh(c+dx))^4} - \frac{1}{12a^2(a+a \tanh(c+dx))^3} - \frac{1}{16a^3(a+a \tanh(c+dx))^2} - \frac{1}{16a^4(a+a \tanh(c+dx))} \right)}{d}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^(-4), x]
```

output

```
(a*(ArcTanh[Tanh[c + d*x]]/(16*a^5) - 1/(8*a*(a + a*Tanh[c + d*x])^4) - 1/(12*a^2*(a + a*Tanh[c + d*x])^3) - 1/(16*a^3*(a + a*Tanh[c + d*x])^2) - 1/(16*a^4*(a + a*Tanh[c + d*x]))) / d
```

**Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3960, 3042, 3960, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \tanh(c + dx) + a)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(a - ia \tan(ic + idx))^4} dx$$

$$\downarrow \text{3960}$$

$$\frac{\int \frac{1}{(\tanh(c+dx)a+a)^3} dx}{2a} - \frac{1}{8d(a \tanh(c + dx) + a)^4}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
 & -\frac{1}{8d(a \tanh(c + dx) + a)^4} + \frac{\int \frac{1}{(a - ia \tan(ic + idx))^3} dx}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\int \frac{1}{(\tanh(c+dx)a+a)^2} dx}{2a} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c + dx) + a)^4} \\
 & \quad \downarrow 3042 \\
 & -\frac{1}{8d(a \tanh(c + dx) + a)^4} + \frac{-\frac{1}{6d(a \tanh(c+dx)+a)^3} + \frac{\int \frac{1}{(a - ia \tan(ic + idx))^2} dx}{2a}}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\int \frac{1}{\tanh(c+dx)a+a} dx}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c + dx) + a)^4} \\
 & \quad \downarrow 3042 \\
 & -\frac{1}{8d(a \tanh(c + dx) + a)^4} + \frac{-\frac{1}{6d(a \tanh(c+dx)+a)^3} + \frac{-\frac{1}{4d(a \tanh(c+dx)+a)^2} + \frac{\int \frac{1}{a - ia \tan(ic + idx)} dx}{2a}}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\int \frac{1 dx}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c + dx) + a)^4} \\
 & \quad \downarrow 24 \\
 & \frac{\frac{x}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)}}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c + dx) + a)^4}
 \end{aligned}$$

input

```
Int[(a + a*Tanh[c + d*x])^(-4), x]
```

output

```
-1/8*1/(d*(a + a*Tanh[c + d*x])^4) + (-1/6*1/(d*(a + a*Tanh[c + d*x])^3) + (-1/4*1/(d*(a + a*Tanh[c + d*x])^2) + (x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x])))/(2*a))/(2*a)/(2*a)
```

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

## Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.79

method	result
risch	$\frac{x}{16a^4} - \frac{e^{-2dx-2c}}{8a^4d} - \frac{3e^{-4dx-4c}}{32a^4d} - \frac{e^{-6dx-6c}}{24a^4d} - \frac{e^{-8dx-8c}}{128a^4d}$
derivativdivides	$-\frac{1}{8(\tanh(dx+c)+1)^4} - \frac{1}{12(\tanh(dx+c)+1)^3} - \frac{1}{16(\tanh(dx+c)+1)^2} - \frac{1}{16(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{32} - \frac{\ln(\tanh(dx+c)-1)}{32}$ $d a^4$
default	$-\frac{1}{8(\tanh(dx+c)+1)^4} - \frac{1}{12(\tanh(dx+c)+1)^3} - \frac{1}{16(\tanh(dx+c)+1)^2} - \frac{1}{16(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{32} - \frac{\ln(\tanh(dx+c)-1)}{32}$ $d a^4$
parallelrisch	$-\frac{16-3 \tanh(dx+c)^3+3 \tanh(dx+c)^4 x d-12 \tanh(dx+c)^2-19 \tanh(dx+c)+18 \tanh(dx+c)^2 x d+3 d x+12 \tanh(dx+c)}{48 d a^4 (\tanh(dx+c)+1)^4}$

input `int(1/(a+a*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/16*x/a^4-1/8/a^4/d*exp(-2*d*x-2*c)-3/32/a^4/d*exp(-4*d*x-4*c)-1/24/a^4/d*exp(-6*d*x-6*c)-1/128/a^4/d*exp(-8*d*x-8*c)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 220 vs.  $2(86) = 172$ .

Time = 0.10 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.29

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx$$

$$= \frac{3(8dx - 1) \cosh(dx + c)^4 + 12(8dx + 1) \cosh(dx + c) \sinh(dx + c)^3 + 3(8dx - 1) \sinh(dx + c)^4 + 2}{384(a^4 d \cosh(dx + c)^4 + 4a^4 d \cosh(dx + c)^3 \sinh(dx + c)}$$

input `integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="fricas")`

output `1/384*(3*(8*d*x - 1)*cosh(d*x + c)^4 + 12*(8*d*x + 1)*cosh(d*x + c)*sinh(d*x + c)^3 + 3*(8*d*x - 1)*sinh(d*x + c)^4 + 2*(9*(8*d*x - 1)*cosh(d*x + c)^2 - 32)*sinh(d*x + c)^2 - 64*cosh(d*x + c)^2 + 4*(3*(8*d*x + 1)*cosh(d*x + c)^3 - 16*cosh(d*x + c))*sinh(d*x + c) - 36)/(a^4*d*cosh(d*x + c)^4 + 4*a^4*d*cosh(d*x + c)^3*sinh(d*x + c) + 6*a^4*d*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*a^4*d*cosh(d*x + c)*sinh(d*x + c)^3 + a^4*d*sinh(d*x + c)^4)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 694 vs.  $2(80) = 160$ .

Time = 0.98 (sec) , antiderivative size = 694, normalized size of antiderivative = 7.23

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(a+a*tanh(d*x+c))**4,x)`

output

```
Piecewise((3*d*x*tanh(c + d*x)**4/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d
*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x)
+ 48*a**4*d) + 12*d*x*tanh(c + d*x)**3/(48*a**4*d*tanh(c + d*x)**4 + 192*a
**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c
+ d*x) + 48*a**4*d) + 18*d*x*tanh(c + d*x)**2/(48*a**4*d*tanh(c + d*x)**4
+ 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*t
anh(c + d*x) + 48*a**4*d) + 12*d*x*tanh(c + d*x)/(48*a**4*d*tanh(c + d*x)*
**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*
d*tanh(c + d*x) + 48*a**4*d) + 3*d*x/(48*a**4*d*tanh(c + d*x)**4 + 192*a**
4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d
*x) + 48*a**4*d) - 3*tanh(c + d*x)**3/(48*a**4*d*tanh(c + d*x)**4 + 192*a**
4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c +
d*x) + 48*a**4*d) - 12*tanh(c + d*x)**2/(48*a**4*d*tanh(c + d*x)**4 + 192*a
**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c
+ d*x) + 48*a**4*d) - 19*tanh(c + d*x)/(48*a**4*d*tanh(c + d*x)**4 + 192*a
**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c +
d*x) + 48*a**4*d) - 16/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c +
d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4
*d), Ne(d, 0)), (x/(a*tanh(c) + a)**4, True))
```

### Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.70

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx$$

$$= \frac{dx + c}{16 a^4 d} - \frac{48 e^{(-2 dx - 2c)} + 36 e^{(-4 dx - 4c)} + 16 e^{(-6 dx - 6c)} + 3 e^{(-8 dx - 8c)}}{384 a^4 d}$$

input

```
integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="maxima")
```

output

```
1/16*(d*x + c)/(a^4*d) - 1/384*(48*e^(-2*d*x - 2*c) + 36*e^(-4*d*x - 4*c)
+ 16*e^(-6*d*x - 6*c) + 3*e^(-8*d*x - 8*c))/(a^4*d)
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.67

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx = -\frac{(48 e^{(6 dx + 6 c)} + 36 e^{(4 dx + 4 c)} + 16 e^{(2 dx + 2 c)} + 3) e^{(-8 dx - 8 c)} - \frac{24 (dx + c)}{a^4}}{384 d}$$

input `integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="giac")`output `-1/384*((48*e^(6*d*x + 6*c) + 36*e^(4*d*x + 4*c) + 16*e^(2*d*x + 2*c) + 3) *e^(-8*d*x - 8*c)/a^4 - 24*(d*x + c)/a^4)/d`**Mupad [B] (verification not implemented)**

Time = 1.97 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx = \frac{x}{16 a^4} - \frac{e^{-2c-2dx}}{8 a^4 d} - \frac{3 e^{-4c-4dx}}{32 a^4 d} - \frac{e^{-6c-6dx}}{24 a^4 d} - \frac{e^{-8c-8dx}}{128 a^4 d}$$

input `int(1/(a + a*tanh(c + d*x))^4,x)`output `x/(16*a^4) - exp(- 2*c - 2*d*x)/(8*a^4*d) - (3*exp(- 4*c - 4*d*x))/(32*a^4 *d) - exp(- 6*c - 6*d*x)/(24*a^4*d) - exp(- 8*c - 8*d*x)/(128*a^4*d)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.75

$$\int \frac{1}{(a + a \tanh(c + dx))^4} dx = \frac{24 e^{8dx+8c} dx - 48 e^{6dx+6c} - 36 e^{4dx+4c} - 16 e^{2dx+2c} - 3}{384 e^{8dx+8c} a^4 d}$$

input `int(1/(a+a*tanh(d*x+c))^4,x)`output `(24*e**(8*c + 8*d*x)*d*x - 48*e**(6*c + 6*d*x) - 36*e**(4*c + 4*d*x) - 16*e**(2*c + 2*d*x) - 3)/(384*e**(8*c + 8*d*x)*a**4*d)`



### 3.49 $\int \frac{1}{(a+a \tanh(c+dx))^5} dx$

Optimal result . . . . .	440
Mathematica [A] (verified) . . . . .	441
Rubi [A] (verified) . . . . .	441
Maple [A] (verified) . . . . .	444
Fricas [B] (verification not implemented) . . . . .	444
Sympy [B] (verification not implemented) . . . . .	445
Maxima [A] (verification not implemented) . . . . .	446
Giac [A] (verification not implemented) . . . . .	447
Mupad [B] (verification not implemented) . . . . .	447
Reduce [B] (verification not implemented) . . . . .	448

#### Optimal result

Integrand size = 12, antiderivative size = 121

$$\int \frac{1}{(a+a \tanh(c+dx))^5} dx = \frac{x}{32a^5} - \frac{1}{10d(a+a \tanh(c+dx))^5} - \frac{1}{16ad(a+a \tanh(c+dx))^4} - \frac{1}{24a^2d(a+a \tanh(c+dx))^3} - \frac{1}{32ad(a^2+a^2 \tanh(c+dx))^2} - \frac{1}{32d(a^5+a^5 \tanh(c+dx))}$$

output

```
1/32*x/a^5-1/10/d/(a+a*tanh(d*x+c))^5-1/16/a/d/(a+a*tanh(d*x+c))^4-1/24/a^
2/d/(a+a*tanh(d*x+c))^3-1/32/a/d/(a^2+a^2*tanh(d*x+c))^2-1/32/d/(a^5+a^5*t
anh(d*x+c))
```

**Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.92

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx$$

$$= \frac{\operatorname{sech}^5(c + dx)(-500 \cosh(c + dx) - 375 \cosh(3(c + dx)) - 149 \cosh(5(c + dx)) - 100 \sinh(c + dx) - 22 \sinh(3(c + dx)))}{3840a^5d(1 + \tanh(c + dx))^5}$$

input

```
Integrate[(a + a*Tanh[c + d*x])^(-5),x]
```

output

```
(Sech[c + d*x]^5*(-500*Cosh[c + d*x] - 375*Cosh[3*(c + d*x)] - 149*Cosh[5*(c + d*x)] - 100*Sinh[c + d*x] - 225*Sinh[3*(c + d*x)] - 125*Sinh[5*(c + d*x)] + 120*ArcTanh[Tanh[c + d*x]]*(Cosh[5*(c + d*x)] + Sinh[5*(c + d*x)]))/ (3840*a^5*d*(1 + Tanh[c + d*x])^5)
```

**Rubi [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.12, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3960, 3042, 3960, 3042, 3960, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \tanh(c + dx) + a)^5} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(a - ia \tan(ic + idx))^5} dx$$

$$\downarrow \text{3960}$$

$$\frac{\int \frac{1}{(\tanh(c+dx)a+a)^4} dx}{2a} - \frac{1}{10d(a \tanh(c + dx) + a)^5}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
 & -\frac{1}{10d(a \tanh(c + dx) + a)^5} + \frac{\int \frac{1}{(a - ia \tan(ic + idx))^4} dx}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\frac{\int \frac{1}{(\tanh(c+dx)a+a)^3} dx}{2a} - \frac{1}{8d(a \tanh(c+dx)+a)^4}}{2a} - \frac{1}{10d(a \tanh(c + dx) + a)^5} \\
 & \quad \downarrow 3042 \\
 & -\frac{1}{10d(a \tanh(c + dx) + a)^5} + \frac{-\frac{1}{8d(a \tanh(c+dx)+a)^4} + \frac{\int \frac{1}{(a - ia \tan(ic + idx))^3} dx}{2a}}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\frac{\int \frac{1}{(\tanh(c+dx)a+a)^2} dx}{2a} - \frac{1}{6d(a \tanh(c+dx)+a)^3}}{2a} - \frac{1}{8d(a \tanh(c+dx)+a)^4} - \frac{1}{10d(a \tanh(c + dx) + a)^5} \\
 & \quad \downarrow 3042 \\
 & -\frac{1}{10d(a \tanh(c + dx) + a)^5} + \frac{-\frac{1}{8d(a \tanh(c+dx)+a)^4} + \frac{-\frac{1}{6d(a \tanh(c+dx)+a)^3} + \frac{\int \frac{1}{(a - ia \tan(ic + idx))^2} dx}{2a}}{2a}}{2a} \\
 & \quad \downarrow 3960 \\
 & \frac{\frac{\int \frac{1}{\tanh(c+dx)a+a} dx}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2}}{2a} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c+dx)+a)^4} - \\
 & \quad \frac{2a}{1} \\
 & \quad \frac{1}{10d(a \tanh(c + dx) + a)^5} \\
 & \quad \downarrow 3042 \\
 & -\frac{1}{10d(a \tanh(c + dx) + a)^5} + \\
 & \quad \frac{-\frac{1}{6d(a \tanh(c+dx)+a)^3} + \frac{-\frac{1}{4d(a \tanh(c+dx)+a)^2} + \frac{\int \frac{1}{a - ia \tan(ic + idx)} dx}{2a}}{2a}}{2a} \\
 & \quad \downarrow 3960
 \end{aligned}$$

$$\frac{\frac{\int \frac{1 dx}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c+dx)+a)^4}}{2a}}{2a} - \frac{1}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c+dx)+a)^4} - \frac{1}{10d(a \tanh(c+dx)+a)^5}$$

↓ 24

$$\frac{\frac{\frac{x}{2a} - \frac{1}{2d(a \tanh(c+dx)+a)} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c+dx)+a)^4}}{2a}}{2a} - \frac{1}{2a} - \frac{1}{4d(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3} - \frac{1}{8d(a \tanh(c+dx)+a)^4} - \frac{1}{10d(a \tanh(c+dx)+a)^5}$$

input `Int[(a + a*Tanh[c + d*x])^(-5), x]`

output `-1/10*1/(d*(a + a*Tanh[c + d*x])^5) + (-1/8*1/(d*(a + a*Tanh[c + d*x])^4) + (-1/6*1/(d*(a + a*Tanh[c + d*x])^3) + (-1/4*1/(d*(a + a*Tanh[c + d*x])^2) + (x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x])))/(2*a))/(2*a))/(2*a))/(2*a)`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

### Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.75

method	result
derivativedivides	$-\frac{1}{10(\tanh(dx+c)+1)^5} - \frac{1}{16(\tanh(dx+c)+1)^4} - \frac{1}{24(\tanh(dx+c)+1)^3} - \frac{1}{32(\tanh(dx+c)+1)^2} - \frac{1}{32(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{64} + \frac{1}{d a^5}$
default	$-\frac{1}{10(\tanh(dx+c)+1)^5} - \frac{1}{16(\tanh(dx+c)+1)^4} - \frac{1}{24(\tanh(dx+c)+1)^3} - \frac{1}{32(\tanh(dx+c)+1)^2} - \frac{1}{32(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{64} + \frac{1}{d a^5}$
risch	$\frac{x}{32a^5} - \frac{5e^{-2dx-2c}}{64a^5d} - \frac{5e^{-4dx-4c}}{64a^5d} - \frac{5e^{-6dx-6c}}{96a^5d} - \frac{5e^{-8dx-8c}}{256a^5d} - \frac{e^{-10dx-10c}}{320a^5d}$
parallelrisch	$\frac{-128-75 \tanh(dx+c)^3+75 \tanh(dx+c)^4xd-155 \tanh(dx+c)^2-175 \tanh(dx+c)+150 \tanh(dx+c)^2xd+15dx+75 \tanh(dx+c)}{480d a^5 (\tanh(dx+c)+1)^5}$

input `int(1/(a+a*tanh(d*x+c))^5,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{d/a^5} \left( -\frac{1}{10}(\tanh(dx+c)+1)^{-5} - \frac{1}{16}(\tanh(dx+c)+1)^{-4} - \frac{1}{24}(\tanh(dx+c)+1)^{-3} - \frac{1}{32}(\tanh(dx+c)+1)^{-2} - \frac{1}{32}(\tanh(dx+c)+1)^{-1} + \frac{1}{64} \ln(\tanh(dx+c)+1) - \frac{1}{64} \ln(\tanh(dx+c)-1) \right)$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. 2(109) = 218.

Time = 0.11 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.37

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx$$

$$= \frac{12(10dx - 1) \cosh(dx + c)^5 + 60(10dx - 1) \cosh(dx + c) \sinh(dx + c)^4 + 12(10dx + 1) \sinh(dx + c)^3 + 12(10dx + 1) \sinh(dx + c) \cosh(dx + c)^2 + 12(10dx + 1) \cosh(dx + c)^2 + 12(10dx + 1)}{3840 a^5 d \cosh(dx + c)^5 + \dots}$$

input `integrate(1/(a+a*tanh(d*x+c))^5,x, algorithm="fricas")`

output

```
1/3840*(12*(10*d*x - 1)*cosh(d*x + c)^5 + 60*(10*d*x - 1)*cosh(d*x + c)*sinh(d*x + c)^4 + 12*(10*d*x + 1)*sinh(d*x + c)^5 + 15*(8*(10*d*x + 1)*cosh(d*x + c)^2 - 15)*sinh(d*x + c)^3 - 375*cosh(d*x + c)^3 + 15*(8*(10*d*x - 1)*cosh(d*x + c)^3 - 75*cosh(d*x + c))*sinh(d*x + c)^2 + 5*(12*(10*d*x + 1)*cosh(d*x + c)^4 - 135*cosh(d*x + c)^2 - 20)*sinh(d*x + c) - 500*cosh(d*x + c))/(a^5*d*cosh(d*x + c)^5 + 5*a^5*d*cosh(d*x + c)^4*sinh(d*x + c) + 10*a^5*d*cosh(d*x + c)^3*sinh(d*x + c)^2 + 10*a^5*d*cosh(d*x + c)^2*sinh(d*x + c)^3 + 5*a^5*d*cosh(d*x + c)*sinh(d*x + c)^4 + a^5*d*sinh(d*x + c)^5)
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1018 vs.  $2(102) = 204$ .

Time = 1.31 (sec) , antiderivative size = 1018, normalized size of antiderivative = 8.41

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx = \text{Too large to display}$$

input

```
integrate(1/(a+a*tanh(d*x+c))**5,x)
```



**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.62

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx$$

$$= -\frac{(300 e^{(8 dx+8 c)}+300 e^{(6 dx+6 c)}+200 e^{(4 dx+4 c)}+75 e^{(2 dx+2 c)}+12) e^{(-10 dx-10 c)}}{a^5} - \frac{120 (dx+c)}{a^5}$$

$$3840 d$$

input `integrate(1/(a+a*tanh(d*x+c))^5,x, algorithm="giac")`output `-1/3840*((300*e^(8*d*x + 8*c) + 300*e^(6*d*x + 6*c) + 200*e^(4*d*x + 4*c) + 75*e^(2*d*x + 2*c) + 12)*e^(-10*d*x - 10*c)/a^5 - 120*(d*x + c)/a^5)/d`**Mupad [B] (verification not implemented)**

Time = 1.99 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx = \frac{x}{32 a^5} - \frac{5 e^{-2c-2dx}}{64 a^5 d} - \frac{5 e^{-4c-4dx}}{64 a^5 d}$$

$$- \frac{5 e^{-6c-6dx}}{96 a^5 d} - \frac{5 e^{-8c-8dx}}{256 a^5 d} - \frac{e^{-10c-10dx}}{320 a^5 d}$$

input `int(1/(a + a*tanh(c + d*x))^5,x)`output `x/(32*a^5) - (5*exp(- 2*c - 2*d*x))/(64*a^5*d) - (5*exp(- 4*c - 4*d*x))/(64*a^5*d) - (5*exp(- 6*c - 6*d*x))/(96*a^5*d) - (5*exp(- 8*c - 8*d*x))/(256*a^5*d) - exp(- 10*c - 10*d*x)/(320*a^5*d)`



**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a + a \tanh(c + dx))^5} dx$$

$$= \frac{120e^{10dx+10c}dx - 300e^{8dx+8c} - 300e^{6dx+6c} - 200e^{4dx+4c} - 75e^{2dx+2c} - 12}{3840e^{10dx+10c}a^5d}$$

input `int(1/(a+a*tanh(d*x+c))^5,x)`output `(120*e**(10*c + 10*d*x)*d*x - 300*e**(8*c + 8*d*x) - 300*e**(6*c + 6*d*x) - 200*e**(4*c + 4*d*x) - 75*e**(2*c + 2*d*x) - 12)/(3840*e**(10*c + 10*d*x)*a**5*d)`

### 3.50 $\int (1 + \tanh(x))^{7/2} dx$

Optimal result	449
Mathematica [A] (verified)	449
Rubi [A] (verified)	450
Maple [A] (verified)	452
Fricas [B] (verification not implemented)	452
Sympy [F]	453
Maxima [A] (verification not implemented)	453
Giac [B] (verification not implemented)	454
Mupad [B] (verification not implemented)	454
Reduce [F]	455

#### Optimal result

Integrand size = 8, antiderivative size = 57

$$\int (1 + \tanh(x))^{7/2} dx = 8\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 8\sqrt{1 + \tanh(x)} - \frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2}$$

output

$8*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-8*(1+\tanh(x))^{(1/2)}-4/3*(1+\tanh(x))^{(3/2)}-2/5*(1+\tanh(x))^{(5/2)}$

#### Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.82

$$\int (1 + \tanh(x))^{7/2} dx = 8\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2}{15}\sqrt{1 + \tanh(x)}(73 + 16 \tanh(x) + 3 \tanh^2(x))$$

input

`Integrate[(1 + Tanh[x])^(7/2), x]`

output

$$8\sqrt{2}\operatorname{ArcTanh}[\sqrt{1 + \operatorname{Tanh}[x]}/\sqrt{2}] - (2\sqrt{1 + \operatorname{Tanh}[x]}*(73 + 16*\operatorname{Tanh}[x] + 3*\operatorname{Tanh}[x]^2))/15$$
**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.11, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$ , Rules used = {3042, 3959, 3042, 3959, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\tanh(x) + 1)^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (1 - i \tan(ix))^{7/2} dx \\
 & \quad \downarrow \text{3959} \\
 & 2 \int (\tanh(x) + 1)^{5/2} dx - \frac{2}{5} (\tanh(x) + 1)^{5/2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{5} (\tanh(x) + 1)^{5/2} + 2 \int (1 - i \tan(ix))^{5/2} dx \\
 & \quad \downarrow \text{3959} \\
 & 2 \left( 2 \int (\tanh(x) + 1)^{3/2} dx - \frac{2}{3} (\tanh(x) + 1)^{3/2} \right) - \frac{2}{5} (\tanh(x) + 1)^{5/2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{5} (\tanh(x) + 1)^{5/2} + 2 \left( -\frac{2}{3} (\tanh(x) + 1)^{3/2} + 2 \int (1 - i \tan(ix))^{3/2} dx \right) \\
 & \quad \downarrow \text{3959} \\
 & 2 \left( 2 \left( 2 \int \sqrt{\tanh(x) + 1} dx - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2} \right) - \frac{2}{5} (\tanh(x) + 1)^{5/2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{2}{5}(\tanh(x) + 1)^{5/2} + \\
& 2\left(-\frac{2}{3}(\tanh(x) + 1)^{3/2} + 2\left(-2\sqrt{\tanh(x) + 1} + 2 \int \sqrt{1 - i \tan(ix)} dx\right)\right) \\
& \quad \downarrow \text{3961} \\
& 2\left(2\left(4 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2\sqrt{\tanh(x) + 1}\right) - \frac{2}{3}(\tanh(x) + 1)^{3/2}\right) - \\
& \quad \frac{2}{5}(\tanh(x) + 1)^{5/2} \\
& \quad \downarrow \text{219} \\
& 2\left(2\left(2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right) - 2\sqrt{\tanh(x) + 1}\right) - \frac{2}{3}(\tanh(x) + 1)^{3/2}\right) - \\
& \quad \frac{2}{5}(\tanh(x) + 1)^{5/2}
\end{aligned}$$

input `Int[(1 + Tanh[x])^(7/2), x]`

output `(-2*(1 + Tanh[x])^(5/2))/5 + 2*((-2*(1 + Tanh[x])^(3/2))/3 + 2*(2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]))`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961

```
Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

**Maple [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$8\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 8\sqrt{1+\tanh(x)} - \frac{4(1+\tanh(x))^{\frac{3}{2}}}{3} - \frac{2(1+\tanh(x))^{\frac{5}{2}}}{5}$	43
default	$8\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 8\sqrt{1+\tanh(x)} - \frac{4(1+\tanh(x))^{\frac{3}{2}}}{3} - \frac{2(1+\tanh(x))^{\frac{5}{2}}}{5}$	43

input

```
int((1+tanh(x))^(7/2),x,method=_RETURNVERBOSE)
```

output

```
8*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-8*(1+tanh(x))^(1/2)-4/3*(
1+tanh(x))^(3/2)-2/5*(1+tanh(x))^(5/2)
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 331 vs.  $2(42) = 84$ .

Time = 0.10 (sec) , antiderivative size = 331, normalized size of antiderivative = 5.81

$$\int (1 + \tanh(x))^{7/2} dx = \frac{4 \left( 15 (\sqrt{2} \cosh(x))^4 + 4 \sqrt{2} \cosh(x) \sinh(x)^3 + \sqrt{2} \sinh(x)^4 + 2 (3 \sqrt{2} \cosh(x)^2 + \sqrt{2}) \right)}{\dots}$$

input

```
integrate((1+tanh(x))^(7/2),x, algorithm="fricas")
```

output

```
4/15*(15*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(23*cosh(x)^5 + 115*cosh(x)*sinh(x)^4 + 23*sinh(x)^5 + 5*(46*cosh(x)^2 + 7)*sinh(x)^3 + 35*cosh(x)^3 + 5*(46*cosh(x)^3 + 21*cosh(x))*sinh(x)^2 + 5*(23*cosh(x)^4 + 21*cosh(x)^2 + 3)*sinh(x) + 15*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)
```

### Sympy [F]

$$\int (1 + \tanh(x))^{7/2} dx = \int (\tanh(x) + 1)^{\frac{7}{2}} dx$$

input

```
integrate((1+tanh(x))**(7/2),x)
```

output

```
Integral((tanh(x) + 1)**(7/2), x)
```

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.46

$$\int (1 + \tanh(x))^{7/2} dx = -4\sqrt{2} \log\left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}\right) - \frac{8\sqrt{2}}{\sqrt{e^{(-2x)}+1}} - \frac{8\sqrt{2}}{3(e^{(-2x)}+1)^{\frac{3}{2}}} - \frac{8\sqrt{2}}{5(e^{(-2x)}+1)^{\frac{5}{2}}}$$

input

```
integrate((1+tanh(x))^(7/2),x, algorithm="maxima")
```

output

```
-4*sqrt(2)*log(-sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/
sqrt(e^(-2*x) + 1)) - 8*sqrt(2)/sqrt(e^(-2*x) + 1) - 8/3*sqrt(2)/(e^(-2*x)
+ 1)^(3/2) - 8/5*sqrt(2)/(e^(-2*x) + 1)^(5/2)
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 140 vs.  $2(42) = 84$ .

Time = 0.13 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.46

$$\int (1 + \tanh(x))^{7/2} dx = \frac{4}{15} \sqrt{2} \left( \frac{2 \left( 45 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^4 - 135 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3 + 170 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 100 \sqrt{e^{4x} + e^{2x}} + 100 e^{2x} + 23 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^5 - 15 \log(-2 \sqrt{e^{4x} + e^{2x}} + 2 e^{2x} + 1)} \right)$$

input

```
integrate((1+tanh(x))^(7/2),x, algorithm="giac")
```

output

```
4/15*sqrt(2)*(2*(45*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^4 - 135*(sqrt(e^(4
*x) + e^(2*x)) - e^(2*x))^3 + 170*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 -
100*sqrt(e^(4*x) + e^(2*x)) + 100*e^(2*x) + 23)/(sqrt(e^(4*x) + e^(2*x)) -
e^(2*x) - 1)^5 - 15*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))
```

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

$$\int (1 + \tanh(x))^{7/2} dx = -8 \sqrt{\tanh(x) + 1} - \frac{4(\tanh(x) + 1)^{3/2}}{3} - \frac{2(\tanh(x) + 1)^{5/2}}{5} - \sqrt{2} \operatorname{atan} \left( \frac{\sqrt{2} \sqrt{\tanh(x) + 1} \operatorname{li}}{2} \right) \operatorname{Si}$$

input

```
int((tanh(x) + 1)^(7/2),x)
```

output

```
- 2^(1/2)*atan((2^(1/2)*(tanh(x) + 1)^(1/2)*1i)/2)*8i - 8*(tanh(x) + 1)^(1/2) - (4*(tanh(x) + 1)^(3/2))/3 - (2*(tanh(x) + 1)^(5/2))/5
```

**Reduce [F]**

$$\int (1 + \tanh(x))^{7/2} dx = \int \sqrt{\tanh(x) + 1} dx + \int \sqrt{\tanh(x) + 1} \tanh(x)^3 dx + 3 \left( \int \sqrt{\tanh(x) + 1} \tanh(x)^2 dx \right) + 3 \left( \int \sqrt{\tanh(x) + 1} \tanh(x) dx \right)$$

input

```
int((1+tanh(x))^(7/2),x)
```

output

```
int(sqrt(tanh(x) + 1),x) + int(sqrt(tanh(x) + 1)*tanh(x)**3,x) + 3*int(sqrt(tanh(x) + 1)*tanh(x)**2,x) + 3*int(sqrt(tanh(x) + 1)*tanh(x),x)
```



### 3.51 $\int (1 + \tanh(x))^{5/2} dx$

Optimal result	456
Mathematica [A] (verified)	456
Rubi [A] (verified)	457
Maple [A] (verified)	458
Fricas [B] (verification not implemented)	459
Sympy [F]	459
Maxima [B] (verification not implemented)	460
Giac [B] (verification not implemented)	460
Mupad [B] (verification not implemented)	461
Reduce [F]	461

#### Optimal result

Integrand size = 8, antiderivative size = 45

$$\int (1 + \tanh(x))^{5/2} dx = 4\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 4\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}$$

output `4*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-4*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)`

#### Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int (1 + \tanh(x))^{5/2} dx = 4\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2}{3}\sqrt{1 + \tanh(x)}(7 + \tanh(x))$$

input `Integrate[(1 + Tanh[x])^(5/2), x]`

output `4*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*Sqrt[1 + Tanh[x]]*(7 + Tanh[x]))/3`

**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 3959, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\tanh(x) + 1)^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (1 - i \tan(ix))^{5/2} dx \\
 & \quad \downarrow \text{3959} \\
 & 2 \int (\tanh(x) + 1)^{3/2} dx - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{3} (\tanh(x) + 1)^{3/2} + 2 \int (1 - i \tan(ix))^{3/2} dx \\
 & \quad \downarrow \text{3959} \\
 & 2 \left( 2 \int \sqrt{\tanh(x) + 1} dx - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{3} (\tanh(x) + 1)^{3/2} + 2 \left( -2\sqrt{\tanh(x) + 1} + 2 \int \sqrt{1 - i \tan(ix)} dx \right) \\
 & \quad \downarrow \text{3961} \\
 & 2 \left( 4 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{219} \\
 & 2 \left( 2\sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2}
 \end{aligned}$$

input

Int[(1 + Tanh[x])^(5/2), x]

output  $(-2*(1 + \tanh[x])^{(3/2)})/3 + 2*(2*\text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[1 + \tanh[x]]/\text{Sqrt}[2]] - 2*\text{Sqrt}[1 + \tanh[x]])$

### Defintions of rubi rules used

rule 219  $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3959  $\text{Int}[(a_ + (b_)*\tan[(c_.) + (d_)*(x_)])^{(n_)}, x\_Symbol] \rightarrow \text{Simp}[b*((a + b*\tan[c + d*x])^{(n - 1)}/(d*(n - 1))), x] + \text{Simp}[2*a \ \text{Int}[(a + b*\tan[c + d*x])^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[n, 1]$

rule 3961  $\text{Int}[\text{Sqrt}[(a_ + (b_)*\tan[(c_.) + (d_)*(x_)]]], x\_Symbol] \rightarrow \text{Simp}[-2*(b/d) \ \text{Subst}[\text{Int}[1/(2*a - x^2), x], x, \text{Sqrt}[a + b*\tan[c + d*x]]], x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$

### Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$4\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 4\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{3/2}}{3}$	35
default	$4\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 4\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{3/2}}{3}$	35

input  $\text{int}((1+\tanh(x))^{(5/2)}, x, \text{method}=\_RETURNVERBOSE)$

output  $4*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-4*(1+\tanh(x))^{(1/2)}-2/3*(1+\tanh(x))^{(3/2)}$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 206 vs.  $2(34) = 68$ .

Time = 0.11 (sec) , antiderivative size = 206, normalized size of antiderivative = 4.58

$$\int (1 + \tanh(x))^{5/2} dx = \frac{2 \left( 3 \left( \sqrt{2} \cosh(x)^2 + 2 \sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2} \right) \log \left( -2 \cosh(x)^2 - \right) \right)}{\dots}$$

input `integrate((1+tanh(x))^(5/2),x, algorithm="fricas")`

output  $2/3*(3*(\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 + \sqrt{2})*\log(-2*\cosh(x)^2 - 4*\cosh(x)*\sinh(x) - 2*\sinh(x)^2 - \sqrt{2}*(\sqrt{2}*\cosh(x)^3 + 3*\sqrt{2}*\cosh(x)*\sinh(x)^2 + \sqrt{2}*\sinh(x)^3 + (3*\sqrt{2}*\cosh(x)^2 + \sqrt{2})*\sinh(x) + \sqrt{2}*\cosh(x))/\sqrt{\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1) - 1) - 2*\sqrt{2}*(4*\cosh(x)^3 + 12*\cosh(x)*\sinh(x)^2 + 4*\sinh(x)^3 + 3*(4*\cosh(x)^2 + 1)*\sinh(x) + 3*\cosh(x))/\sqrt{\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1})/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)$

### Sympy [F]

$$\int (1 + \tanh(x))^{5/2} dx = \int (\tanh(x) + 1)^{5/2} dx$$

input `integrate((1+tanh(x))**(5/2),x)`

output `Integral((tanh(x) + 1)**(5/2), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 70 vs.  $2(34) = 68$ .

Time = 0.17 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.56

$$\int (1 + \tanh(x))^{5/2} dx = -2\sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}} \right) - \frac{4\sqrt{2}}{\sqrt{e^{(-2x)}+1}} - \frac{4\sqrt{2}}{3(e^{(-2x)}+1)^{3/2}}$$

input `integrate((1+tanh(x))^(5/2),x, algorithm="maxima")`

output `-2*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1))) - 4*sqrt(2)/sqrt(e^(-2*x) + 1) - 4/3*sqrt(2)/(e^(-2*x) + 1)^(3/2)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 96 vs.  $2(34) = 68$ .

Time = 0.13 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.13

$$\int (1 + \tanh(x))^{5/2} dx = \frac{2}{3}\sqrt{2} \left( \frac{2 \left( 6 \left( \sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^2 - 9 \sqrt{e^{(4x)} + e^{(2x)}} + 9e^{(2x)} + 4 \right)}{\left( \sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} - 1 \right)^3} - 3 \log \left( -2 \sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} - 1 \right) \right)$$

input `integrate((1+tanh(x))^(5/2),x, algorithm="giac")`

output `2/3*sqrt(2)*(2*(6*(sqrt(e^(4*x) + e^(2*x))) - e^(2*x))^2 - 9*sqrt(e^(4*x) + e^(2*x)) + 9*e^(2*x) + 4)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1)^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 1.95 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.20

$$\int (1 + \tanh(x))^{5/2} dx = \sqrt{8} \ln \left( -2\sqrt{8} \sqrt{\tanh(x) + 1} - 8 \right) - \frac{2(\tanh(x) + 1)^{3/2}}{3} - 2\sqrt{2} \ln \left( 4\sqrt{2} \sqrt{\tanh(x) + 1} - 8 \right) - 4\sqrt{\tanh(x) + 1}$$

input `int((tanh(x) + 1)^(5/2),x)`output `8^(1/2)*log(- 2*8^(1/2)*(tanh(x) + 1)^(1/2) - 8) - (2*(tanh(x) + 1)^(3/2))/3 - 2*2^(1/2)*log(4*2^(1/2)*(tanh(x) + 1)^(1/2) - 8) - 4*(tanh(x) + 1)^(1/2)`**Reduce [F]**

$$\int (1 + \tanh(x))^{5/2} dx = \int \sqrt{\tanh(x) + 1} dx + \int \sqrt{\tanh(x) + 1} \tanh(x)^2 dx + 2 \left( \int \sqrt{\tanh(x) + 1} \tanh(x) dx \right)$$

input `int((1+tanh(x))^(5/2),x)`output `int(sqrt(tanh(x) + 1),x) + int(sqrt(tanh(x) + 1)*tanh(x)**2,x) + 2*int(sqrt(tanh(x) + 1)*tanh(x),x)`

### 3.52 $\int (1 + \tanh(x))^{3/2} dx$

Optimal result	462
Mathematica [A] (verified)	462
Rubi [A] (verified)	463
Maple [A] (verified)	464
Fricas [B] (verification not implemented)	465
Sympy [F]	465
Maxima [B] (verification not implemented)	466
Giac [A] (verification not implemented)	466
Mupad [B] (verification not implemented)	467
Reduce [F]	467

#### Optimal result

Integrand size = 8, antiderivative size = 33

$$\int (1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)}$$

output

```
2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int (1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)}$$

input

```
Integrate[(1 + Tanh[x])^(3/2), x]
```

output

```
2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (\tanh(x) + 1)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (1 - i \tan(ix))^{3/2} dx \\
 & \quad \downarrow \text{3959} \\
 & 2 \int \sqrt{\tanh(x) + 1} dx - 2\sqrt{\tanh(x) + 1} \\
 & \quad \downarrow \text{3042} \\
 & -2\sqrt{\tanh(x) + 1} + 2 \int \sqrt{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3961} \\
 & 4 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2\sqrt{\tanh(x) + 1} \\
 & \quad \downarrow \text{219} \\
 & 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right) - 2\sqrt{\tanh(x) + 1}
 \end{aligned}$$

input `Int[(1 + Tanh[x])^(3/2), x]`

output `2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]`



## Definitions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)}$	27
default	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)}$	27

input `int((1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)`

output `2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 127 vs.  $2(26) = 52$ .

Time = 0.10 (sec) , antiderivative size = 127, normalized size of antiderivative = 3.85

$$\int (1 + \tanh(x))^{3/2} dx = \sqrt{2} \log \left( \frac{-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 + \sqrt{2}(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (3\sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x) + \sqrt{2} \cosh(x) - 1)}{\sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}} \right) - \frac{2\sqrt{2}(\cosh(x) + \sinh(x))}{\sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}}$$

input `integrate((1+tanh(x))^(3/2),x, algorithm="fricas")`

output `sqrt(2)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(cosh(x) + sinh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)`

**Sympy [F]**

$$\int (1 + \tanh(x))^{3/2} dx = \int (\tanh(x) + 1)^{\frac{3}{2}} dx$$

input `integrate((1+tanh(x))**(3/2),x)`

output `Integral((tanh(x) + 1)**(3/2), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 57 vs.  $2(26) = 52$ .

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.73

$$\int (1 + \tanh(x))^{3/2} dx = -\sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}} \right) - \frac{2\sqrt{2}}{\sqrt{e^{(-2x)}+1}}$$

input `integrate((1+tanh(x))^(3/2),x, algorithm="maxima")`

output `-sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1))) - 2*sqrt(2)/sqrt(e^(-2*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.58

$$\int (1 + \tanh(x))^{3/2} dx = \sqrt{2} \left( \frac{2}{\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} - 1} - \log \left( -2 \sqrt{e^{(4x)} + e^{(2x)}} + 2e^{(2x)} + 1 \right) \right)$$

input `integrate((1+tanh(x))^(3/2),x, algorithm="giac")`

output `sqrt(2)*(2/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1) - log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int (1 + \tanh(x))^{3/2} dx = 2\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right) - 2\sqrt{\tanh(x)+1}$$

input `int((tanh(x) + 1)^(3/2),x)`

output `2*2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2)`

**Reduce [F]**

$$\int (1 + \tanh(x))^{3/2} dx = \int \sqrt{\tanh(x)+1} dx + \int \sqrt{\tanh(x)+1} \tanh(x) dx$$

input `int((1+tanh(x))^(3/2),x)`

output `int(sqrt(tanh(x) + 1),x) + int(sqrt(tanh(x) + 1)*tanh(x),x)`

### 3.53 $\int \sqrt{1 + \tanh(x)} dx$

Optimal result	468
Mathematica [A] (verified)	468
Rubi [A] (verified)	469
Maple [A] (verified)	470
Fricas [B] (verification not implemented)	470
Sympy [F]	471
Maxima [B] (verification not implemented)	471
Giac [A] (verification not implemented)	472
Mupad [B] (verification not implemented)	472
Reduce [F]	472

#### Optimal result

Integrand size = 8, antiderivative size = 21

$$\int \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)$$

output

```
2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)$$

input

```
Integrate[Sqrt[1 + Tanh[x]], x]
```

output

```
Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]
```

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\tanh(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{1 - i \tan(ix)} dx \\ & \quad \downarrow \text{3961} \\ & 2 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} \\ & \quad \downarrow \text{219} \\ & \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) \end{aligned}$$

input `Int[Sqrt[1 + Tanh[x]], x]`

output `Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]`

**Defintions of rubi rules used**

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961

```
Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
  , b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)$	17
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)$	17

input

```
int((1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 99 vs.  $2(16) = 32$ .

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 4.71

$$\int \sqrt{1 + \tanh(x)} dx = \frac{1}{2} \sqrt{2} \log \left( \frac{-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - \sqrt{2}(\sqrt{2} \cosh(x)^3 + 3 \sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (3 \sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x) + \sqrt{2} \cosh(x) \sinh(x) + \sqrt{2})}{\sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}} - 1 \right)$$

input

```
integrate((1+tanh(x))^(1/2),x, algorithm="fricas")
```

output `1/2*sqrt(2)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1)`

## Sympy [F]

$$\int \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} dx$$

input `integrate((1+tanh(x))**(1/2),x)`

output `Integral(sqrt(tanh(x) + 1), x)`

## Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(16) = 32.

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.05

$$\int \sqrt{1 + \tanh(x)} dx = -\frac{1}{2} \sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}} \right)$$

input `integrate((1+tanh(x))^(1/2),x, algorithm="maxima")`

output `-1/2*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1)))`



**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \sqrt{1 + \tanh(x)} dx = -\frac{1}{2} \sqrt{2} \log \left( -2 \sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1 \right)$$

input `integrate((1+tanh(x))^(1/2),x, algorithm="giac")`

output `-1/2*sqrt(2)*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1)`

**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{atanh} \left( \frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right)$$

input `int((tanh(x) + 1)^(1/2),x)`

output `2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2)`

**Reduce [F]**

$$\int \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} dx$$

input `int((1+tanh(x))^(1/2),x)`

output `int(sqrt(tanh(x) + 1),x)`

### 3.54 $\int \frac{1}{\sqrt{1+\tanh(x)}} dx$

Optimal result	473
Mathematica [C] (verified)	473
Rubi [A] (verified)	474
Maple [A] (verified)	475
Fricas [B] (verification not implemented)	476
Sympy [F]	476
Maxima [B] (verification not implemented)	477
Giac [A] (verification not implemented)	477
Mupad [B] (verification not implemented)	478
Reduce [F]	478

#### Optimal result

Integrand size = 8, antiderivative size = 32

$$\int \frac{1}{\sqrt{1+\tanh(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1+\tanh(x)}}$$

output `1/2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/(1+tanh(x))^(1/2)`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{1+\tanh(x)}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1+\tanh(x))\right)}{\sqrt{1+\tanh(x)}}$$

input `Integrate[1/Sqrt[1 + Tanh[x]], x]`

output `-(Hypergeometric2F1[-1/2, 1, 1/2, (1 + Tanh[x])/2]/Sqrt[1 + Tanh[x]])`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{\tanh(x)+1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{1-i \tan(ix)}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \int \sqrt{\tanh(x)+1} dx - \frac{1}{\sqrt{\tanh(x)+1}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{\sqrt{\tanh(x)+1}} + \frac{1}{2} \int \sqrt{1-i \tan(ix)} dx \\
 & \quad \downarrow \text{3961} \\
 & \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)+1} - \frac{1}{\sqrt{\tanh(x)+1}} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{\tanh(x)+1}}
 \end{aligned}$$

input `Int[1/Sqrt[1 + Tanh[x]], x]`

output `ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]]`

## Definitions of rubi rules used

rule 219  $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$   $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 3042  $\text{Int}[u_+, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3960  $\text{Int}[(a_+) + (b_+)*\tan[(c_+) + (d_+)(x_+)]^{(n_+)}, x\_Symbol] \rightarrow \text{Simp}[a*((a + b*\text{Tan}[c + d*x])^n/(2*b*d*n)), x] + \text{Simp}[1/(2*a) \ \text{Int}[(a + b*\text{Tan}[c + d*x])^{(n + 1)}, x], x] /;$   $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[n, 0]$

rule 3961  $\text{Int}[\text{Sqrt}[(a_+) + (b_+)*\tan[(c_+) + (d_+)(x_+)]], x\_Symbol] \rightarrow \text{Simp}[-2*(b/d) \ \text{Subst}[\text{Int}[1/(2*a - x^2), x], x, \text{Sqrt}[a + b*\text{Tan}[c + d*x]], x] /;$   $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} - \frac{1}{\sqrt{1+\tanh(x)}}$	27
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} - \frac{1}{\sqrt{1+\tanh(x)}}$	27

input `int(1/(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/(1+tanh(x))^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 141 vs.  $2(26) = 52$ .

Time = 0.11 (sec) , antiderivative size = 141, normalized size of antiderivative = 4.41

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx$$

$$= \frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - \frac{\sqrt{2}(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x) + 3\sqrt{2} \sinh(x)^3)}{4(\cosh(x) + \sinh(x))}\right)}{4(\cosh(x) + \sinh(x))}$$

input `integrate(1/(1+tanh(x))^(1/2),x, algorithm="fricas")`

output `1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x) + 3*sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x) + sinh(x))`

**Sympy [F]**

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx = \int \frac{1}{\sqrt{\tanh(x) + 1}} dx$$

input `integrate(1/(1+tanh(x))**(1/2),x)`

output `Integral(1/sqrt(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 57 vs.  $2(26) = 52$ .

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.78

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx = -\frac{1}{4} \sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}} \right) - \frac{1}{2} \sqrt{2} \sqrt{e^{(-2x)} + 1}$$

input `integrate(1/(1+tanh(x))^(1/2),x, algorithm="maxima")`

output `-1/4*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1))) - 1/2*sqrt(2)*sqrt(e^(-2*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.56

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx = -\frac{1}{4} \sqrt{2} \left( \frac{2}{\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)}} + \log \left( -2 \sqrt{e^{(4x)} + e^{(2x)}} + 2e^{(2x)} + 1 \right) \right)$$

input `integrate(1/(1+tanh(x))^(1/2),x, algorithm="giac")`

output `-1/4*sqrt(2)*(2/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x)) + log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 2.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx = \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} - \frac{1}{\sqrt{\tanh(x) + 1}}$$

input `int(1/(tanh(x) + 1)^(1/2),x)`output `(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/2 - 1/(tanh(x) + 1)^(1/2)`**Reduce [F]**

$$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx = 2\sqrt{\tanh(x) + 1} + \int \frac{\sqrt{\tanh(x) + 1} \tanh(x)^2}{\tanh(x) + 1} dx$$

input `int(1/(1+tanh(x))^(1/2),x)`output `2*sqrt(tanh(x) + 1) + int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x) + 1),x)`

### 3.55 $\int \frac{1}{(1+\tanh(x))^{3/2}} dx$

Optimal result	479
Mathematica [C] (verified)	479
Rubi [A] (verified)	480
Maple [A] (verified)	481
Fricas [B] (verification not implemented)	482
Sympy [F]	482
Maxima [B] (verification not implemented)	483
Giac [B] (verification not implemented)	483
Mupad [B] (verification not implemented)	484
Reduce [F]	484

#### Optimal result

Integrand size = 8, antiderivative size = 49

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}}$$

output `1/4*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/3/(1+tanh(x))^(3/2)-1/2/(1+tanh(x))^(1/2)`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.21 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.57

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{1}{2}(1 + \tanh(x))\right)}{3(1 + \tanh(x))^{3/2}}$$

input `Integrate[(1 + Tanh[x])^(-3/2), x]`

output `-1/3*Hypergeometric2F1[-3/2, 1, -1/2, (1 + Tanh[x])/2]/(1 + Tanh[x])^(3/2)`



**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 3960, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\tanh(x) + 1)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{\tanh(x) + 1}} dx - \frac{1}{3(\tanh(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{3(\tanh(x) + 1)^{3/2}} + \frac{1}{2} \int \frac{1}{\sqrt{1 - i \tan(ix)}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \left( \frac{1}{2} \int \sqrt{\tanh(x) + 1} dx - \frac{1}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{3(\tanh(x) + 1)^{3/2}} + \frac{1}{2} \left( -\frac{1}{\sqrt{\tanh(x) + 1}} + \frac{1}{2} \int \sqrt{1 - i \tan(ix)} dx \right) \\
 & \quad \downarrow \text{3961} \\
 & \frac{1}{2} \left( \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}}
 \end{aligned}$$

input `Int[(1 + Tanh[x])^(-3/2), x]`

output `-1/3*1/(1 + Tanh[x])^(3/2) + (ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]])/2`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))* ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

### Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} - \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{2\sqrt{1+\tanh(x)}}$	35
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} - \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{2\sqrt{1+\tanh(x)}}$	35

input `int(1/(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)`

output  $\frac{1}{4} \cdot 2^{1/2} \cdot \operatorname{arctanh}\left(\frac{1}{2} \cdot (1 + \tanh(x))^{1/2} \cdot 2^{1/2}\right) - \frac{1}{3} \cdot (1 + \tanh(x))^{3/2} - \frac{1}{2} \cdot (1 + \tanh(x))^{1/2}$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs.  $2(34) = 68$ .

Time = 0.11 (sec) , antiderivative size = 244, normalized size of antiderivative = 4.98

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = \frac{3(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x)^2 \sinh(x) + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3)}{(1 + \tanh(x))^{3/2}}$$

input `integrate(1/(1+tanh(x))^(3/2),x, algorithm="fricas")`

output  $\frac{1}{24} \cdot (3 \cdot (\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x)^2 \sinh(x) + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3) \cdot \log(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - \sqrt{2} (\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (3\sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x) + \sqrt{2} \cosh(x))) / \sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1} - 1) - 2 \sqrt{2} \cdot (4 \cosh(x)^4 + 16 \cosh(x) \sinh(x)^3 + 4 \sinh(x)^4 + (24 \cosh(x)^2 + 5) \sinh(x)^2 + 5 \cosh(x)^2 + 2 \cdot (8 \cosh(x)^3 + 5 \cosh(x)) \sinh(x) + 1) / \sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}) / (\cosh(x)^3 + 3 \cosh(x)^2 \sinh(x) + 3 \cosh(x) \sinh(x)^2 + \sinh(x)^3)$

### Sympy [F]

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = \int \frac{1}{(\tanh(x) + 1)^{3/2}} dx$$

input `integrate(1/(1+tanh(x))**(3/2),x)`

output `Integral((tanh(x) + 1)**(-3/2), x)`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs.  $2(34) = 68$ .

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.41

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = -\frac{1}{12} \sqrt{2} \left( \frac{3}{e^{(-2x)} + 1} + 1 \right) (e^{(-2x)} + 1)^{\frac{3}{2}} - \frac{1}{8} \sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}} \right)$$

input `integrate(1/(1+tanh(x))^(3/2),x, algorithm="maxima")`

output `-1/12*sqrt(2)*(3/(e^(-2*x) + 1) + 1)*(e^(-2*x) + 1)^(3/2) - 1/8*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1)))`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs.  $2(34) = 68$ .

Time = 0.14 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.94

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = -\frac{1}{24} \sqrt{2} \left( \frac{2 \left( 6 \left( \sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^2 - 3 \sqrt{e^{(4x)} + e^{(2x)}} + 3 e^{(2x)} + 1 \right)}{\left( \sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^3} + 3 \log \left( -2 \sqrt{e^{(4x)} + e^{(2x)}} + \dots \right) \right)$$

input `integrate(1/(1+tanh(x))^(3/2),x, algorithm="giac")`

output

```
-1/24*sqrt(2)*(2*(6*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 3*sqrt(e^(4*x)
+ e^(2*x)) + 3*e^(2*x) + 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 3*log
(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))
```

**Mupad [B] (verification not implemented)**

Time = 2.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.65

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x)+1}}{2}\right)}{4} - \frac{\frac{\tanh(x)}{2} + \frac{5}{6}}{(\tanh(x) + 1)^{3/2}}$$

input

```
int(1/(tanh(x) + 1)^(3/2),x)
```

output

```
(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 - (tanh(x)/2 + 5/6)/(ta
nh(x) + 1)^(3/2)
```

**Reduce [F]**

$$\int \frac{1}{(1 + \tanh(x))^{3/2}} dx = \frac{-2\sqrt{\tanh(x) + 1} + \left(\int \frac{\sqrt{\tanh(x)+1} \tanh(x)^2}{\tanh(x)^2 + 2\tanh(x)+1} dx\right) \tanh(x) + \int \frac{\sqrt{\tanh(x)+1} \tanh(x)^2}{\tanh(x)^2 + 2\tanh(x)+1} dx}{\tanh(x) + 1}$$

input

```
int(1/(1+tanh(x))^(3/2),x)
```

output

```
( - 2*sqrt(tanh(x) + 1) + int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**2 +
2*tanh(x) + 1),x)*tanh(x) + int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**
2 + 2*tanh(x) + 1),x))/(tanh(x) + 1)
```

### 3.56 $\int \frac{1}{(1+\tanh(x))^{5/2}} dx$

Optimal result	485
Mathematica [C] (verified)	485
Rubi [A] (verified)	486
Maple [A] (verified)	488
Fricas [B] (verification not implemented)	488
Sympy [F]	489
Maxima [A] (verification not implemented)	489
Giac [B] (verification not implemented)	490
Mupad [B] (verification not implemented)	490
Reduce [F]	491

#### Optimal result

Integrand size = 8, antiderivative size = 61

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}} - \frac{1}{4\sqrt{1 + \tanh(x)}}$$

output

$1/8*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-1/5/(1+\tanh(x))^{(5/2)}-1/6/(1+\tanh(x))^{(3/2)}-1/4/(1+\tanh(x))^{(1/2)}$

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.46

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = -\frac{\operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, 1, -\frac{3}{2}, \frac{1}{2}(1 + \tanh(x))\right)}{5(1 + \tanh(x))^{5/2}}$$

input

`Integrate[(1 + Tanh[x])^(-5/2), x]`

output

$$-1/5*\text{Hypergeometric2F1}[-5/2, 1, -3/2, (1 + \text{Tanh}[x])/2]/(1 + \text{Tanh}[x])^{5/2}$$
**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$ , Rules used = {3042, 3960, 3042, 3960, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(\tanh(x) + 1)^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix))^{5/2}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \int \frac{1}{(\tanh(x) + 1)^{3/2}} dx - \frac{1}{5(\tanh(x) + 1)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{5(\tanh(x) + 1)^{5/2}} + \frac{1}{2} \int \frac{1}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\sqrt{\tanh(x) + 1}} dx - \frac{1}{3(\tanh(x) + 1)^{3/2}} \right) - \frac{1}{5(\tanh(x) + 1)^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{5(\tanh(x) + 1)^{5/2}} + \frac{1}{2} \left( -\frac{1}{3(\tanh(x) + 1)^{3/2}} + \frac{1}{2} \int \frac{1}{\sqrt{1 - i \tan(ix)}} dx \right) \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \left( \frac{1}{2} \left( \frac{1}{2} \int \sqrt{\tanh(x) + 1} dx - \frac{1}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}} \right) - \frac{1}{5(\tanh(x) + 1)^{5/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{5(\tanh(x)+1)^{5/2}} + \\
& \frac{1}{2} \left( -\frac{1}{3(\tanh(x)+1)^{3/2}} + \frac{1}{2} \left( -\frac{1}{\sqrt{\tanh(x)+1}} + \frac{1}{2} \int \sqrt{1-i \tan(ix)} dx \right) \right) \\
& \quad \downarrow \text{3961} \\
& \frac{1}{2} \left( \frac{1}{2} \left( \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)+1} - \frac{1}{\sqrt{\tanh(x)+1}} \right) - \frac{1}{3(\tanh(x)+1)^{3/2}} \right) - \\
& \quad \frac{1}{5(\tanh(x)+1)^{5/2}} \\
& \quad \downarrow \text{219} \\
& \frac{1}{2} \left( \frac{1}{2} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{\tanh(x)+1}} \right) - \frac{1}{3(\tanh(x)+1)^{3/2}} \right) - \frac{1}{5(\tanh(x)+1)^{5/2}}
\end{aligned}$$

input `Int[(1 + Tanh[x])^(-5/2), x]`

output `-1/5*1/(1 + Tanh[x])^(5/2) + (-1/3*1/(1 + Tanh[x])^(3/2) + (ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]])/2)/2`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`



rule 3961

```
Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.70

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{8} - \frac{1}{5(1+\tanh(x))^{\frac{5}{2}}} - \frac{1}{6(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{4\sqrt{1+\tanh(x)}}$	43
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{8} - \frac{1}{5(1+\tanh(x))^{\frac{5}{2}}} - \frac{1}{6(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{4\sqrt{1+\tanh(x)}}$	43

input

```
int(1/(1+tanh(x))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
1/8*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/5/(1+tanh(x))^(5/2)-1
/6/(1+tanh(x))^(3/2)-1/4/(1+tanh(x))^(1/2)
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 338 vs.  $2(42) = 84$ .

Time = 0.10 (sec) , antiderivative size = 338, normalized size of antiderivative = 5.54

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = \frac{15 (\sqrt{2} \cosh(x))^5 + 5 \sqrt{2} \cosh(x)^4 \sinh(x) + 10 \sqrt{2} \cosh(x)^3 \sinh(x)^2 + 10 \sqrt{2} \cosh(x)^2 \sinh(x)^3 + 5 \sqrt{2} \cosh(x) \sinh(x)^4 + 5 \sinh(x)^5}{(1 + \tanh(x))^{5/2}}$$

input

```
integrate(1/(1+tanh(x))^(5/2),x, algorithm="fricas")
```

output

```
1/240*(15*(sqrt(2)*cosh(x)^5 + 5*sqrt(2)*cosh(x)^4*sinh(x) + 10*sqrt(2)*cosh(x)^3*sinh(x)^2 + 10*sqrt(2)*cosh(x)^2*sinh(x)^3 + 5*sqrt(2)*cosh(x)*sinh(x)^4 + sqrt(2)*sinh(x)^5)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(23*cosh(x)^6 + 138*cosh(x)*sinh(x)^5 + 23*sinh(x)^6 + (345*cosh(x)^2 + 34)*sinh(x)^4 + 34*cosh(x)^4 + 4*(115*cosh(x)^3 + 34*cosh(x))*sinh(x)^3 + (345*cosh(x)^4 + 204*cosh(x)^2 + 14)*sinh(x)^2 + 14*cosh(x)^2 + 2*(69*cosh(x)^5 + 68*cosh(x)^3 + 14*cosh(x))*sinh(x) + 3)/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x)^5 + 5*cosh(x)^4*sinh(x) + 10*cosh(x)^3*sinh(x)^2 + 10*cosh(x)^2*sinh(x)^3 + 5*cosh(x)*sinh(x)^4 + sinh(x)^5)
```

**Sympy [F]**

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = \int \frac{1}{(\tanh(x) + 1)^{5/2}} dx$$

input

```
integrate(1/(1+tanh(x))**(5/2),x)
```

output

```
Integral((tanh(x) + 1)**(-5/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.30

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = -\frac{1}{120} \sqrt{2} \left( \frac{5}{e^{(-2x)} + 1} + \frac{15}{(e^{(-2x)} + 1)^2} + 3 \right) (e^{(-2x)} + 1)^{\frac{5}{2}} - \frac{1}{16} \sqrt{2} \log \left( -\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}} \right)$$

input

```
integrate(1/(1+tanh(x))^(5/2),x, algorithm="maxima")
```

output

```
-1/120*sqrt(2)*(5/(e^(-2*x) + 1) + 15/(e^(-2*x) + 1)^2 + 3)*(e^(-2*x) + 1)
^(5/2) - 1/16*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2)
+ sqrt(2)/sqrt(e^(-2*x) + 1)))
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 139 vs.  $2(42) = 84$ .

Time = 0.13 (sec) , antiderivative size = 139, normalized size of antiderivative = 2.28

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx =$$

$$-\frac{1}{240} \sqrt{2} \left( \frac{2 \left( 45 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^4 - 45 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3 + 35 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^5} \right)$$

input

```
integrate(1/(1+tanh(x))^(5/2),x, algorithm="giac")
```

output

```
-1/240*sqrt(2)*(2*(45*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^4 - 45*(sqrt(e^(
4*x) + e^(2*x)) - e^(2*x))^3 + 35*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 -
15*sqrt(e^(4*x) + e^(2*x)) + 15*e^(2*x) + 3)/(sqrt(e^(4*x) + e^(2*x)) - e^
(2*x))^5 + 15*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))
```

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.66

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x)+1}}{2}\right)}{8} - \frac{\frac{\tanh(x)}{6} + \frac{(\tanh(x)+1)^2}{4} + \frac{11}{30}}{(\tanh(x) + 1)^{5/2}}$$

input

```
int(1/(tanh(x) + 1)^(5/2),x)
```

output

```
(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/8 - (tanh(x)/6 + (tanh(x)
+ 1)^2/4 + 11/30)/(tanh(x) + 1)^(5/2)
```

**Reduce [F]**

$$\int \frac{1}{(1 + \tanh(x))^{5/2}} dx = \frac{-2\sqrt{\tanh(x) + 1} + 3 \left( \int \frac{\sqrt{\tanh(x)+1} \tanh(x)^2}{\tanh(x)^3 + 3 \tanh(x)^2 + 3 \tanh(x) + 1} dx \right) \tanh(x)^2 + 6 \left( \int \frac{\sqrt{\tanh(x)+1}}{\tanh(x)^3 + 3 \tanh(x)^2 + 3 \tanh(x) + 1} dx \right) \tanh(x)}{3 \tanh(x)^2 + 6 \tanh(x) + 3}$$

input `int(1/(1+tanh(x))^(5/2),x)`

output `( - 2*sqrt(tanh(x) + 1) + 3*int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**3 + 3*tanh(x)**2 + 3*tanh(x) + 1),x)*tanh(x)**2 + 6*int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**3 + 3*tanh(x)**2 + 3*tanh(x) + 1),x)*tanh(x) + 3*int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**3 + 3*tanh(x)**2 + 3*tanh(x) + 1),x))/(3*(tanh(x)**2 + 2*tanh(x) + 1))`

### 3.57 $\int (a + b \tanh(c + dx))^5 dx$

Optimal result	492
Mathematica [A] (verified)	493
Rubi [A] (verified)	493
Maple [A] (verified)	496
Fricas [B] (verification not implemented)	497
Sympy [A] (verification not implemented)	498
Maxima [B] (verification not implemented)	498
Giac [A] (verification not implemented)	499
Mupad [B] (verification not implemented)	500
Reduce [B] (verification not implemented)	500

#### Optimal result

Integrand size = 12, antiderivative size = 142

$$\int (a + b \tanh(c + dx))^5 dx = a(a^4 + 10a^2b^2 + 5b^4)x + \frac{b(5a^4 + 10a^2b^2 + b^4) \log(\cosh(c + dx))}{d} - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} - \frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} - \frac{b(a + b \tanh(c + dx))^4}{4d}$$

output

```
a*(a^4+10*a^2*b^2+5*b^4)*x+b*(5*a^4+10*a^2*b^2+b^4)*ln(cosh(d*x+c))/d-4*a*b^2*(a^2+b^2)*tanh(d*x+c)/d-1/2*b*(3*a^2+b^2)*(a+b*tanh(d*x+c))^2/d-2/3*a*b*(a+b*tanh(d*x+c))^3/d-1/4*b*(a+b*tanh(d*x+c))^4/d
```

**Mathematica [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.80

$$\int (a + b \tanh(c + dx))^5 dx = \frac{6(a + b)^5 \log(1 - \tanh(c + dx)) - 6(a - b)^5 \log(1 + \tanh(c + dx)) + 60ab^2(2a^2 + b^2) \tanh(c + dx) + 12d}{12d}$$

input `Integrate[(a + b*Tanh[c + d*x])^5,x]`

output `-1/12*(6*(a + b)^5*Log[1 - Tanh[c + d*x]] - 6*(a - b)^5*Log[1 + Tanh[c + d*x]] + 60*a*b^2*(2*a^2 + b^2)*Tanh[c + d*x] + 6*b^3*(10*a^2 + b^2)*Tanh[c + d*x]^2 + 20*a*b^4*Tanh[c + d*x]^3 + 3*b^5*Tanh[c + d*x]^4)/d`

**Rubi [A] (verified)**

Time = 0.83 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 3963, 3042, 4011, 3042, 4011, 3042, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tanh(c + dx))^5 dx \\ & \quad \downarrow \text{3042} \\ & \int (a - ib \tan(ic + idx))^5 dx \\ & \quad \downarrow \text{3963} \\ & \int (a + b \tanh(c + dx))^3 (a^2 + 2b \tanh(c + dx)a + b^2) dx - \frac{b(a + b \tanh(c + dx))^4}{4d} \\ & \quad \downarrow \text{3042} \\ & -\frac{b(a + b \tanh(c + dx))^4}{4d} + \int (a - ib \tan(ic + idx))^3 (a^2 - 2ib \tan(ic + idx)a + b^2) dx \end{aligned}$$

$$\begin{aligned}
& \downarrow 4011 \\
& \int (a + b \tanh(c + dx))^2 (a(a^2 + 3b^2) + b(3a^2 + b^2) \tanh(c + dx)) dx - \\
& \quad \frac{b(a + b \tanh(c + dx))^4}{4d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 3042 \\
& \int (a - ib \tan(ic + idx))^2 (a(a^2 + 3b^2) - ib(3a^2 + b^2) \tan(ic + idx)) dx - \\
& \quad \frac{b(a + b \tanh(c + dx))^4}{4d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 4011 \\
& \int (a + b \tanh(c + dx)) (a^4 + 6b^2a^2 + 4b(a^2 + b^2) \tanh(c + dx)a + b^4) dx - \\
& \quad \frac{b(3a^2 + b^2) (a + b \tanh(c + dx))^2}{2d} - \frac{b(a + b \tanh(c + dx))^4}{4d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 3042 \\
& \int (a - ib \tan(ic + idx)) (a^4 + 6b^2a^2 - 4ib(a^2 + b^2) \tan(ic + idx)a + b^4) dx - \\
& \quad \frac{b(3a^2 + b^2) (a + b \tanh(c + dx))^2}{2d} - \frac{b(a + b \tanh(c + dx))^4}{4d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 4008 \\
& -ib(5a^4 + 10a^2b^2 + b^4) \int i \tanh(c + dx) dx - \frac{b(3a^2 + b^2) (a + b \tanh(c + dx))^2}{2d} - \\
& \quad \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \\
& \quad \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 26 \\
& b(5a^4 + 10a^2b^2 + b^4) \int \tanh(c + dx) dx - \frac{b(3a^2 + b^2) (a + b \tanh(c + dx))^2}{2d} - \\
& \quad \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \\
& \quad \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& b(5a^4 + 10a^2b^2 + b^4) \int -i \tan(ic + idx) dx - \frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \\
& \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \\
& \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \quad \downarrow \text{26} \\
& -ib(5a^4 + 10a^2b^2 + b^4) \int \tan(ic + idx) dx - \frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \\
& \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \\
& \frac{2ab(a + b \tanh(c + dx))^3}{3d} \\
& \quad \downarrow \text{3956} \\
& -\frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + \\
& \frac{b(5a^4 + 10a^2b^2 + b^4) \log(\cosh(c + dx))}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \\
& \frac{2ab(a + b \tanh(c + dx))^3}{3d}
\end{aligned}$$

input `Int[(a + b*Tanh[c + d*x])^5,x]`

output `a*(a^4 + 10*a^2*b^2 + 5*b^4)*x + (b*(5*a^4 + 10*a^2*b^2 + b^4)*Log[Cosh[c + d*x]])/d - (4*a*b^2*(a^2 + b^2)*Tanh[c + d*x])/d - (b*(3*a^2 + b^2)*(a + b*Tanh[c + d*x])^2)/(2*d) - (2*a*b*(a + b*Tanh[c + d*x])^3)/(3*d) - (b*(a + b*Tanh[c + d*x])^4)/(4*d)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 3956  $\text{Int}[\tan[(c_.) + (d_.)(x_.)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 3963  $\text{Int}[(a_. + (b_.)\tan[(c_.) + (d_.)(x_.)]^{(n_.)}, x\_Symbol] \rightarrow \text{Simp}[b*((a + b*\text{Tan}[c + d*x])^{(n - 1)}/(d*(n - 1))), x] + \text{Int}[(a^2 - b^2 + 2*a*b*\text{Tan}[c + d*x])*(a + b*\text{Tan}[c + d*x])^{(n - 2)}, x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[n, 1]$

rule 4008  $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]*((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x\_Symbol] \rightarrow \text{Simp}[(a*c - b*d)*x, x] + (\text{Simp}[b*d*(\text{Tan}[e + f*x]/f), x] + \text{Simp}[(b*c + a*d) \text{Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[b*c + a*d, 0]$

rule 4011  $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}*((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x\_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m - 1)}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$

## Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.28

method	result
derivativdivides	$\frac{(a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5) \ln(\tanh(dx+c)+1) - b^5 \tanh(dx+c)^4 - 5ab^4 \tanh(dx+c)^3 - 5a^2b^3 \tanh(dx+c)^2 - 5ab^4}{d}$
default	$\frac{(a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5) \ln(\tanh(dx+c)+1) - b^5 \tanh(dx+c)^4 - 5ab^4 \tanh(dx+c)^3 - 5a^2b^3 \tanh(dx+c)^2 - 5ab^4}{d}$
parallelrisch	$- \frac{3b^5 \tanh(dx+c)^4 + 20ab^4 \tanh(dx+c)^3 - 12a^5 dx + 60a^4 b dx - 120a^3 b^2 dx + 120a^2 b^3 dx - 60ab^4 dx + 12b^5 dx + 60a^2 b^3 \tanh(dx+c)}{d}$
parts	$a^5 x + \frac{b^5 \left( -\frac{\tanh(dx+c)^4}{4} - \frac{\tanh(dx+c)^2}{2} - \frac{\ln(\tanh(dx+c)-1)}{2} - \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{5ab^4 \left( -\frac{\tanh(dx+c)^3}{3} - \tanh(dx+c) \right)}{d}$
risch	$a^5 x - 5ba^4 x + 10a^3 b^2 x - 10b^3 a^2 x + 5ab^4 x - b^5 x - \frac{10ba^4 c}{d} - \frac{20b^3 a^2 c}{d} - \frac{2b^5 c}{d} + \frac{4b^2 (15a^3 e^6)}{d}$

input  $\text{int}((a+b*\tanh(d*x+c))^5, x, \text{method}=\_RETURNVERBOSE)$

output

```
1/d*(1/2*(a^5-5*a^4*b+10*a^3*b^2-10*a^2*b^3+5*a*b^4-b^5)*ln(tanh(d*x+c)+1)
-1/4*b^5*tanh(d*x+c)^4-5/3*a*b^4*tanh(d*x+c)^3-5*a^2*b^3*tanh(d*x+c)^2-5*a
*b^4*tanh(d*x+c)-1/2*(a^5+5*a^4*b+10*a^3*b^2+10*a^2*b^3+5*a*b^4+b^5)*ln(ta
nh(d*x+c)-1)-1/2*b^5*tanh(d*x+c)^2-10*a^3*b^2*tanh(d*x+c))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2739 vs.  $2(136) = 272$ .

Time = 0.10 (sec) , antiderivative size = 2739, normalized size of antiderivative = 19.29

$$\int (a + b \tanh(c + dx))^5 dx = \text{Too large to display}$$

input

```
integrate((a+b*tanh(d*x+c))^5,x, algorithm="fricas")
```

output

```
1/3*(3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(
d*x + c)^8 + 24*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*
d*x*cosh(d*x + c)*sinh(d*x + c)^7 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2
*b^3 + 5*a*b^4 - b^5)*d*x*sinh(d*x + c)^8 + 12*(5*a^3*b^2 + 5*a^2*b^3 + 5*
a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*
x)*cosh(d*x + c)^6 + 12*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + 7*(a^5 -
5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x + c)^2 + (
a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*sinh(d*x + c
)^6 + 24*(7*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*
cosh(d*x + c)^3 + 3*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*
b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c))*sinh(d*x
+ c)^5 + 60*a^3*b^2 + 40*a*b^4 + 6*(30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2
*b^5 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*co
sh(d*x + c)^4 + 6*(35*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 -
b^5)*d*x*cosh(d*x + c)^4 + 30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2*b^5 + 3
*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x + 30*(5*a^3
*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^
3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^4 + 24*(7*(a^5 - 5*
a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x + c)^5 + 10*
(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - ...
```

**Sympy [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.49

$$\int (a + b \tanh(c + dx))^5 dx$$

$$= \begin{cases} a^5 x + 5a^4 b x - \frac{5a^4 b \log(\tanh(c+dx)+1)}{d} + 10a^3 b^2 x - \frac{10a^3 b^2 \tanh(c+dx)}{d} + 10a^2 b^3 x - \frac{10a^2 b^3 \log(\tanh(c+dx)+1)}{d} - 5a b^4 x + \frac{5a b^4 \tanh(c+dx)}{d} + 5a^2 b^2 x - \frac{5a^2 b^2 \log(\tanh(c+dx)+1)}{d} + 5a^3 b x - \frac{5a^3 b \tanh(c+dx)}{d} + 5a^4 x - \frac{5a^4 \log(\tanh(c+dx)+1)}{d} \\ x(a + b \tanh(c))^5 \end{cases}$$

input `integrate((a+b*tanh(d*x+c))**5,x)`

output

```
Piecewise((a**5*x + 5*a**4*b*x - 5*a**4*b*log(tanh(c + d*x) + 1)/d + 10*a**3*b**2*x - 10*a**3*b**2*tanh(c + d*x)/d + 10*a**2*b**3*x - 10*a**2*b**3*log(tanh(c + d*x) + 1)/d - 5*a**2*b**3*tanh(c + d*x)**2/d + 5*a*b**4*x - 5*a*b**4*tanh(c + d*x)**3/(3*d) - 5*a*b**4*tanh(c + d*x)/d + b**5*x - b**5*log(tanh(c + d*x) + 1)/d - b**5*tanh(c + d*x)**4/(4*d) - b**5*tanh(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tanh(c))**5, True))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 310 vs. 2(136) = 272.

Time = 0.15 (sec) , antiderivative size = 310, normalized size of antiderivative = 2.18

$$\int (a + b \tanh(c + dx))^5 dx$$

$$= \frac{5}{3} ab^4 \left( 3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)} \right)$$

$$+ b^5 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{4(e^{(-2dx-2c)} + e^{(-4dx-4c)} + e^{(-6dx-6c)})}{d(4e^{(-2dx-2c)} + 6e^{(-4dx-4c)} + 4e^{(-6dx-6c)} + e^{(-8dx-8c)} + 1)} \right)$$

$$+ 10a^2 b^3 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 10a^3 b^2 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^5 x + \frac{5a^4 b \log(\cosh(dx + c))}{d}$$

input `integrate((a+b*tanh(d*x+c))^5,x, algorithm="maxima")`

output

```
5/3*a*b^4*(3*x + 3*c/d - 4*(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + 2)/(
d*(3*e^(-2*d*x - 2*c) + 3*e^(-4*d*x - 4*c) + e^(-6*d*x - 6*c) + 1))) + b^5
*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 4*(e^(-2*d*x - 2*c) + e^(-4*d*x
- 4*c) + e^(-6*d*x - 6*c))/(d*(4*e^(-2*d*x - 2*c) + 6*e^(-4*d*x - 4*c) + 4
*e^(-6*d*x - 6*c) + e^(-8*d*x - 8*c) + 1))) + 10*a^2*b^3*(x + c/d + log(e^
(-2*d*x - 2*c) + 1)/d + 2*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) + e^(-4*
d*x - 4*c) + 1))) + 10*a^3*b^2*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) +
a^5*x + 5*a^4*b*log(cosh(d*x + c))/d
```

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.58

$$\int (a + b \tanh(c + dx))^5 dx$$

$$= \frac{3(a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5)(dx + c) + 3(5a^4b + 10a^2b^3 + b^5) \log(e^{(2dx+2c)} + 1) + 4(a^5x + 5a^4b \log(\cosh(dx + c)))}{d}$$

input

```
integrate((a+b*tanh(d*x+c))^5,x, algorithm="giac")
```

output

```
1/3*(3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*(d*x + c)
+ 3*(5*a^4*b + 10*a^2*b^3 + b^5)*log(e^(2*d*x + 2*c) + 1) + 4*(15*a^3*b^2
+ 10*a*b^4 + 3*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5)*e^(6*d*x + 6*c) +
3*(15*a^3*b^2 + 10*a^2*b^3 + 10*a*b^4 + b^5)*e^(4*d*x + 4*c) + (45*a^3*b^2
+ 15*a^2*b^3 + 25*a*b^4 + 3*b^5)*e^(2*d*x + 2*c))/(e^(2*d*x + 2*c) + 1)^4
)/d
```

**Mupad [B] (verification not implemented)**

Time = 2.07 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.08

$$\int (a + b \tanh(c + dx))^5 dx = x (a^5 + 5 a^4 b + 10 a^3 b^2 + 10 a^2 b^3 + 5 a b^4 + b^5) - \frac{5 \tanh(c + dx) (2 a^3 b^2 + a b^4)}{d} - \frac{b^5 \tanh(c + dx)^4}{4 d} - \frac{\ln(\tanh(c + dx) + 1) (5 a^4 b + 10 a^2 b^3 + b^5)}{d} - \frac{\tanh(c + dx)^2 (10 a^2 b^3 + b^5)}{2 d} - \frac{5 a b^4 \tanh(c + dx)^3}{3 d}$$

input `int((a + b*tanh(c + d*x))^5,x)`output `x*(5*a*b^4 + 5*a^4*b + a^5 + b^5 + 10*a^2*b^3 + 10*a^3*b^2) - (5*tanh(c + d*x)*(a*b^4 + 2*a^3*b^2))/d - (b^5*tanh(c + d*x)^4)/(4*d) - (log(tanh(c + d*x) + 1)*(5*a^4*b + b^5 + 10*a^2*b^3))/d - (tanh(c + d*x)^2*(b^5 + 10*a^2*b^3))/(2*d) - (5*a*b^4*tanh(c + d*x)^3)/(3*d)`**Reduce [B] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 1151, normalized size of antiderivative = 8.11

$$\int (a + b \tanh(c + dx))^5 dx = \text{Too large to display}$$

input `int((a+b*tanh(d*x+c))^5,x)`

output

```
(15***e**(8*c + 8*d*x)*log(e**(2*c + 2*d*x) + 1)*a**4*b + 30***e**(8*c + 8*d*x)
)*log(e**(2*c + 2*d*x) + 1)*a**2*b**3 + 3***e**(8*c + 8*d*x)*log(e**(2*c + 2
*d*x) + 1)*b**5 + 3***e**(8*c + 8*d*x)*a**5*d*x - 15***e**(8*c + 8*d*x)*a**4*b
*d*x + 30***e**(8*c + 8*d*x)*a**3*b**2*d*x - 15***e**(8*c + 8*d*x)*a**3*b**2 -
30***e**(8*c + 8*d*x)*a**2*b**3*d*x - 15***e**(8*c + 8*d*x)*a**2*b**3 + 15***e
*(8*c + 8*d*x)*a*b**4*d*x - 15***e**(8*c + 8*d*x)*a*b**4 - 3***e**(8*c + 8*d*x
)*b**5*d*x - 3***e**(8*c + 8*d*x)*b**5 + 60***e**(6*c + 6*d*x)*log(e**(2*c + 2
*d*x) + 1)*a**4*b + 120***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x) + 1)*a**2*b*
*3 + 12***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x) + 1)*b**5 + 12***e**(6*c + 6*d
*x)*a**5*d*x - 60***e**(6*c + 6*d*x)*a**4*b*d*x + 120***e**(6*c + 6*d*x)*a**3*
b**2*d*x - 120***e**(6*c + 6*d*x)*a**2*b**3*d*x + 60***e**(6*c + 6*d*x)*a*b**4
*d*x - 12***e**(6*c + 6*d*x)*b**5*d*x + 90***e**(4*c + 4*d*x)*log(e**(2*c + 2*
d*x) + 1)*a**4*b + 180***e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1)*a**2*b**
3 + 18***e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1)*b**5 + 18***e**(4*c + 4*d*
x)*a**5*d*x - 90***e**(4*c + 4*d*x)*a**4*b*d*x + 180***e**(4*c + 4*d*x)*a**3*b
**2*d*x + 90***e**(4*c + 4*d*x)*a**3*b**2 - 180***e**(4*c + 4*d*x)*a**2*b**3*d
*x + 30***e**(4*c + 4*d*x)*a**2*b**3 + 90***e**(4*c + 4*d*x)*a*b**4*d*x + 30***e
**(4*c + 4*d*x)*a*b**4 - 18***e**(4*c + 4*d*x)*b**5*d*x - 6***e**(4*c + 4*d*x)
*b**5 + 60***e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1)*a**4*b + 120***e**(2*c
+ 2*d*x)*log(e**(2*c + 2*d*x) + 1)*a**2*b**3 + 12***e**(2*c + 2*d*x)*log...
```

### 3.58 $\int (a + b \tanh(c + dx))^4 dx$

Optimal result . . . . .	502
Mathematica [A] (verified) . . . . .	502
Rubi [A] (verified) . . . . .	503
Maple [A] (verified) . . . . .	506
Fricas [B] (verification not implemented) . . . . .	506
Sympy [A] (verification not implemented) . . . . .	507
Maxima [B] (verification not implemented) . . . . .	508
Giac [A] (verification not implemented) . . . . .	508
Mupad [B] (verification not implemented) . . . . .	509
Reduce [B] (verification not implemented) . . . . .	509

#### Optimal result

Integrand size = 12, antiderivative size = 101

$$\int (a + b \tanh(c + dx))^4 dx = (a^4 + 6a^2b^2 + b^4) x + \frac{4ab(a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} - \frac{ab(a + b \tanh(c + dx))^2}{d} - \frac{b(a + b \tanh(c + dx))^3}{3d}$$

output

```
(a^4+6*a^2*b^2+b^4)*x+4*a*b*(a^2+b^2)*ln(cosh(d*x+c))/d-b^2*(3*a^2+b^2)*tanh(d*x+c)/d-a*b*(a+b*tanh(d*x+c))^2/d-1/3*b*(a+b*tanh(d*x+c))^3/d
```

#### Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.90

$$\int (a + b \tanh(c + dx))^4 dx = \frac{3(a + b)^4 \log(1 - \tanh(c + dx)) - 3(a - b)^4 \log(1 + \tanh(c + dx)) + 6b^2(6a^2 + b^2) \tanh(c + dx) + 12a^2b \tanh^2(c + dx) + 4b^3 \tanh^3(c + dx)}{6d}$$

input

```
Integrate[(a + b*Tanh[c + d*x])^4,x]
```

output

```
-1/6*(3*(a + b)^4*Log[1 - Tanh[c + d*x]] - 3*(a - b)^4*Log[1 + Tanh[c + d*
x]] + 6*b^2*(6*a^2 + b^2)*Tanh[c + d*x] + 12*a*b^3*Tanh[c + d*x]^2 + 2*b^4
*Tanh[c + d*x]^3)/d
```

**Rubi [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$ , Rules used = {3042, 3963, 3042, 4011, 3042, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tanh(c + dx))^4 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ib \tan(ic + idx))^4 dx \\
 & \quad \downarrow \text{3963} \\
 & \int (a + b \tanh(c + dx))^2 (a^2 + 2b \tanh(c + dx)a + b^2) dx - \frac{b(a + b \tanh(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b(a + b \tanh(c + dx))^3}{3d} + \int (a - ib \tan(ic + idx))^2 (a^2 - 2ib \tan(ic + idx)a + b^2) dx \\
 & \quad \downarrow \text{4011} \\
 & \int (a + b \tanh(c + dx)) (a(a^2 + 3b^2) + b(3a^2 + b^2) \tanh(c + dx)) dx - \\
 & \quad \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ib \tan(ic + idx)) (a(a^2 + 3b^2) - ib(3a^2 + b^2) \tan(ic + idx)) dx - \\
 & \quad \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d} \\
 & \quad \downarrow \text{4008}
 \end{aligned}$$



$$-4iab(a^2 + b^2) \int i \tanh(c + dx) dx - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

↓ 26

$$4ab(a^2 + b^2) \int \tanh(c + dx) dx - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

↓ 3042

$$4ab(a^2 + b^2) \int -i \tan(ic + idx) dx - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

↓ 26

$$-4iab(a^2 + b^2) \int \tan(ic + idx) dx - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

↓ 3956

$$-\frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + \frac{4ab(a^2 + b^2) \log(\cosh(c + dx))}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

input

```
Int[(a + b*Tanh[c + d*x])^4,x]
```

output

```
(a^4 + 6*a^2*b^2 + b^4)*x + (4*a*b*(a^2 + b^2)*Log[Cosh[c + d*x]])/d - (b^2*(3*a^2 + b^2)*Tanh[c + d*x])/d - (a*b*(a + b*Tanh[c + d*x])^2)/d - (b*(a + b*Tanh[c + d*x])^3)/(3*d)
```

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956  $\text{Int}[\tan[(c.) + (d.)*(x)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 3963  $\text{Int}[(a.) + (b.)*\tan[(c.) + (d.)*(x)]^{(n)}, x\_Symbol] \rightarrow \text{Simp}[b*((a + b*\text{Tan}[c + d*x])^{(n-1)}/(d*(n-1))), x] + \text{Int}[(a^2 - b^2 + 2*a*b*\text{Tan}[c + d*x])*(a + b*\text{Tan}[c + d*x])^{(n-2)}, x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[n, 1]$
- rule 4008  $\text{Int}[(a.) + (b.)*\tan[(e.) + (f.)*(x)]*((c.) + (d.)*\tan[(e.) + (f.)*(x)]), x\_Symbol] \rightarrow \text{Simp}[(a*c - b*d)*x, x] + (\text{Simp}[b*d*(\text{Tan}[e + f*x]/f), x] + \text{Simp}[(b*c + a*d) \text{Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[b*c + a*d, 0]$
- rule 4011  $\text{Int}[(a.) + (b.)*\tan[(e.) + (f.)*(x)]^{(m)}*((c.) + (d.)*\tan[(e.) + (f.)*(x)]), x\_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

### Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.31

method	result
parallelsch	$\frac{-b^4 \tanh(dx+c)^3 - 3a^4 dx + 12a^3 b dx - 18a^2 b^2 dx + 12a b^3 dx - 3b^4 dx + 6a b^3 \tanh(dx+c)^2 + 12 \ln(1 - \tanh(dx+c)) a^3 b + 12 \ln(1 + \tanh(dx+c)) a^3 b}{3d}$
derivativedivides	$\frac{-\frac{b^4 \tanh(dx+c)^3}{3} - 2a b^3 \tanh(dx+c)^2 - 6a^2 b^2 \tanh(dx+c) - b^4 \tanh(dx+c) - \frac{(a^4 + 4a^3 b + 6a^2 b^2 + 4a b^3 + b^4) \ln(\tanh(dx+c))}{2}}{d}$
default	$\frac{-\frac{b^4 \tanh(dx+c)^3}{3} - 2a b^3 \tanh(dx+c)^2 - 6a^2 b^2 \tanh(dx+c) - b^4 \tanh(dx+c) - \frac{(a^4 + 4a^3 b + 6a^2 b^2 + 4a b^3 + b^4) \ln(\tanh(dx+c))}{2}}{d}$
parts	$a^4 x + \frac{b^4 \left( -\frac{\tanh(dx+c)^3}{3} - \tanh(dx+c) - \frac{\ln(\tanh(dx+c)-1)}{2} + \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{4a b^3 \left( -\frac{\tanh(dx+c)^2}{2} - \frac{\ln(\tanh(dx+c))}{2} \right)}{d}$
risch	$a^4 x - 4a^3 b x + 6a^2 b^2 x - 4a b^3 x + b^4 x - \frac{8a^3 b c}{d} - \frac{8a b^3 c}{d} + \frac{4b^2 (9e^{4dx+4c} a^2 + 6e^{4dx+4c} ab + 3e^{4dx+4c})}{d}$

input `int((a+b*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `-1/3*(b^4*tanh(d*x+c)^3-3*a^4*d*x+12*a^3*b*d*x-18*a^2*b^2*d*x+12*a*b^3*d*x-3*b^4*d*x+6*a*b^3*tanh(d*x+c)^2+12*ln(1-tanh(d*x+c))*a^3*b+12*ln(1+tanh(d*x+c))*a*b^3+18*a^2*b^2*tanh(d*x+c)+3*b^4*tanh(d*x+c))/d`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1389 vs. 2(99) = 198.

Time = 0.11 (sec) , antiderivative size = 1389, normalized size of antiderivative = 13.75

$$\int (a + b \tanh(c + dx))^4 dx = \text{Too large to display}$$

input `integrate((a+b*tanh(d*x+c))^4,x, algorithm="fricas")`

output

```

1/3*(3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*cosh(d*x + c)^6 + 1
8*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*cosh(d*x + c)*sinh(d*x +
c)^5 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*sinh(d*x + c)^6
+ 3*(12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3
+ b^4)*d*x)*cosh(d*x + c)^4 + 3*(15*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3
+ b^4)*d*x*cosh(d*x + c)^2 + 12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3
*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*sinh(d*x + c)^4 + 36*a^2*b^2 + 8*b^4
+ 12*(5*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*cosh(d*x + c)^3 +
(12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b
^4)*d*x)*cosh(d*x + c))*sinh(d*x + c)^3 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4
*a*b^3 + b^4)*d*x + 3*(24*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6
*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*cosh(d*x + c)^2 + 3*(15*(a^4 - 4*a^3*b + 6*
a^2*b^2 - 4*a*b^3 + b^4)*d*x*cosh(d*x + c)^4 + 24*a^2*b^2 + 8*a*b^3 + 4*b^
4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x + 6*(12*a^2*b^2 + 8*
a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*cosh(d*
x + c)^2)*sinh(d*x + c)^2 + 12*((a^3*b + a*b^3)*cosh(d*x + c)^6 + 6*(a^3*b
+ a*b^3)*cosh(d*x + c)*sinh(d*x + c)^5 + (a^3*b + a*b^3)*sinh(d*x + c)^6
+ 3*(a^3*b + a*b^3)*cosh(d*x + c)^4 + 3*(a^3*b + a*b^3 + 5*(a^3*b + a*b^3)
*cosh(d*x + c)^2)*sinh(d*x + c)^4 + a^3*b + a*b^3 + 4*(5*(a^3*b + a*b^3)*c
osh(d*x + c)^3 + 3*(a^3*b + a*b^3)*cosh(d*x + c))*sinh(d*x + c)^3 + 3*(...

```

### Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.43

$$\int (a + b \tanh(c + dx))^4 dx$$

$$= \begin{cases} a^4 x + 4a^3 b x - \frac{4a^3 b \log(\tanh(c+dx)+1)}{d} + 6a^2 b^2 x - \frac{6a^2 b^2 \tanh(c+dx)}{d} + 4ab^3 x - \frac{4ab^3 \log(\tanh(c+dx)+1)}{d} - \frac{2ab^3 \tanh(c+dx)}{d} \\ x(a + b \tanh(c))^4 \end{cases}$$

input

```
integrate((a+b*tanh(d*x+c))**4,x)
```

output

```

Piecewise((a**4*x + 4*a**3*b*x - 4*a**3*b*log(tanh(c + d*x) + 1)/d + 6*a**
2*b**2*x - 6*a**2*b**2*tanh(c + d*x)/d + 4*a*b**3*x - 4*a*b**3*log(tanh(c
+ d*x) + 1)/d - 2*a*b**3*tanh(c + d*x)**2/d + b**4*x - b**4*tanh(c + d*x)*
*3/(3*d) - b**4*tanh(c + d*x)/d, Ne(d, 0)), (x*(a + b*tanh(c))**4, True))

```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 201 vs.  $2(99) = 198$ .

Time = 0.12 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.99

$$\int (a + b \tanh(c + dx))^4 dx$$

$$= \frac{1}{3} b^4 \left( 3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)} \right)$$

$$+ 4ab^3 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 6a^2b^2 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^4x + \frac{4a^3b \log(\cosh(dx + c))}{d}$$

input `integrate((a+b*tanh(d*x+c))^4,x, algorithm="maxima")`

output  $\frac{1}{3}b^4(3x + 3c/d - 4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)/(d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1))) + 4a^3b^3(x + c/d + \log(e^{(-2dx-2c)} + 1)/d + 2e^{(-2dx-2c)}/(d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1))) + 6a^2b^2(x + c/d - 2/(d(e^{(-2dx-2c)} + 1))) + a^4x + 4a^3b \log(\cosh(dx + c))/d$

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.50

$$\int (a + b \tanh(c + dx))^4 dx$$

$$= \frac{3(a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4)(dx + c) + 12(a^3b + ab^3) \log(e^{(2dx+2c)} + 1) + \frac{4(9a^2b^2 + 2b^4 + 3a^2b^2 + 2ab^3)}{3d}}{3d}$$

input `integrate((a+b*tanh(d*x+c))^4,x, algorithm="giac")`



output

```
(12***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x) + 1)*a**3*b + 12***e**(6*c + 6*d*x)
)*log(e**(2*c + 2*d*x) + 1)*a*b**3 + 3***e**(6*c + 6*d*x)*a**4*d*x - 12***e**(
6*c + 6*d*x)*a**3*b*d*x + 18***e**(6*c + 6*d*x)*a**2*b**2*d*x - 12***e**(6*c +
6*d*x)*a**2*b**2 - 12***e**(6*c + 6*d*x)*a*b**3*d*x - 8***e**(6*c + 6*d*x)*a*
b**3 + 3***e**(6*c + 6*d*x)*b**4*d*x - 4***e**(6*c + 6*d*x)*b**4 + 36***e**(4*c
+ 4*d*x)*log(e**(2*c + 2*d*x) + 1)*a**3*b + 36***e**(4*c + 4*d*x)*log(e**(2*
c + 2*d*x) + 1)*a*b**3 + 9***e**(4*c + 4*d*x)*a**4*d*x - 36***e**(4*c + 4*d*x)
*a**3*b*d*x + 54***e**(4*c + 4*d*x)*a**2*b**2*d*x - 36***e**(4*c + 4*d*x)*a*b*
**3*d*x + 9***e**(4*c + 4*d*x)*b**4*d*x + 36***e**(2*c + 2*d*x)*log(e**(2*c + 2
*d*x) + 1)*a**3*b + 36***e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1)*a*b**3 +
9***e**(2*c + 2*d*x)*a**4*d*x - 36***e**(2*c + 2*d*x)*a**3*b*d*x + 54***e**(2*c
+ 2*d*x)*a**2*b**2*d*x + 36***e**(2*c + 2*d*x)*a**2*b**2 - 36***e**(2*c + 2*d
*x)*a*b**3*d*x + 9***e**(2*c + 2*d*x)*b**4*d*x + 12*log(e**(2*c + 2*d*x) + 1
)*a**3*b + 12*log(e**(2*c + 2*d*x) + 1)*a*b**3 + 3*a**4*d*x - 12*a**3*b*d*
x + 18*a**2*b**2*d*x + 24*a**2*b**2 - 12*a*b**3*d*x - 8*a*b**3 + 3*b**4*d*
x + 4*b**4)/(3*d*(e**(6*c + 6*d*x) + 3***e**(4*c + 4*d*x) + 3***e**(2*c + 2*d*
x) + 1))
```

### 3.59 $\int (a + b \tanh(c + dx))^3 dx$

Optimal result . . . . .	511
Mathematica [A] (verified) . . . . .	511
Rubi [A] (verified) . . . . .	512
Maple [A] (verified) . . . . .	514
Fricas [B] (verification not implemented) . . . . .	514
Sympy [A] (verification not implemented) . . . . .	515
Maxima [A] (verification not implemented) . . . . .	516
Giac [A] (verification not implemented) . . . . .	516
Mupad [B] (verification not implemented) . . . . .	517
Reduce [B] (verification not implemented) . . . . .	517

#### Optimal result

Integrand size = 12, antiderivative size = 69

$$\int (a + b \tanh(c + dx))^3 dx = a(a^2 + 3b^2)x + \frac{b(3a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d}$$

output a\*(a^2+3\*b^2)\*x+b\*(3\*a^2+b^2)\*ln(cosh(d\*x+c))/d-2\*a\*b^2\*tanh(d\*x+c)/d-1/2\*b\*(a+b\*tanh(d\*x+c))^2/d

#### Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.97

$$\int (a + b \tanh(c + dx))^3 dx = \frac{(a + b)^3 \log(1 - \tanh(c + dx)) - (a - b)^3 \log(1 + \tanh(c + dx)) + 6ab^2 \tanh(c + dx) + b^3 \tanh^2(c + dx)}{2d}$$

input Integrate[(a + b\*Tanh[c + d\*x])^3,x]



output

$$-1/2*((a + b)^3 \text{Log}[1 - \text{Tanh}[c + d*x]] - (a - b)^3 \text{Log}[1 + \text{Tanh}[c + d*x]] + 6*a*b^2*\text{Tanh}[c + d*x] + b^3*\text{Tanh}[c + d*x]^2)/d$$

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {3042, 3963, 3042, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tanh(c + dx))^3 dx \\ & \quad \downarrow 3042 \\ & \int (a - ib \tan(ic + idx))^3 dx \\ & \quad \downarrow 3963 \\ & \int (a + b \tanh(c + dx)) (a^2 + 2b \tanh(c + dx)a + b^2) dx - \frac{b(a + b \tanh(c + dx))^2}{2d} \\ & \quad \downarrow 3042 \\ & -\frac{b(a + b \tanh(c + dx))^2}{2d} + \int (a - ib \tan(ic + idx)) (a^2 - 2ib \tan(ic + idx)a + b^2) dx \\ & \quad \downarrow 4008 \\ & -ib(3a^2 + b^2) \int i \tanh(c + dx) dx + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d} \\ & \quad \downarrow 26 \\ & b(3a^2 + b^2) \int \tanh(c + dx) dx + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d} \\ & \quad \downarrow 3042 \\ & b(3a^2 + b^2) \int -i \tan(ic + idx) dx + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d} \\ & \quad \downarrow 26 \end{aligned}$$

$$-ib(3a^2 + b^2) \int \tan(ic + idx)dx + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d}$$

↓ 3956

$$\frac{b(3a^2 + b^2) \log(\cosh(c + dx))}{d} + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d}$$

input `Int[(a + b*Tanh[c + d*x])^3,x]`

output `a*(a^2 + 3*b^2)*x + (b*(3*a^2 + b^2)*Log[Cosh[c + d*x]])/d - (2*a*b^2*Tanh[c + d*x])/d - (b*(a + b*Tanh[c + d*x])^2)/(2*d)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3963 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Int[(a^2 - b^2 + 2*a*b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n - 2), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.35

method	result
derivativedivides	$\frac{-\frac{b^3 \tanh(dx+c)^2}{2} - 3b^2 a \tanh(dx+c) - \frac{(a^3 + 3a^2 b + 3b^2 a + b^3) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^3 - 3a^2 b + 3b^2 a - b^3) \ln(\tanh(dx+c)+1)}{2}}{d}$
default	$\frac{-\frac{b^3 \tanh(dx+c)^2}{2} - 3b^2 a \tanh(dx+c) - \frac{(a^3 + 3a^2 b + 3b^2 a + b^3) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^3 - 3a^2 b + 3b^2 a - b^3) \ln(\tanh(dx+c)+1)}{2}}{d}$
parallelrisch	$-\frac{-2a^3 dx + 6a^2 b dx - 6a b^2 dx + 2b^3 dx + b^3 \tanh(dx+c)^2 + 6 \ln(1 - \tanh(dx+c)) a^2 b + 2 \ln(1 - \tanh(dx+c)) b^3 + 6b^2 a \tanh(dx+c)}{2d}$
parts	$a^3 x + \frac{b^3 \left( -\frac{\tanh(dx+c)^2}{2} - \frac{\ln(\tanh(dx+c)-1)}{2} - \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{3a^2 b \ln(\cosh(dx+c))}{d} + \frac{3b^2 a (-\tanh(dx+c))}{d}$
risch	$a^3 x - 3a^2 b x + 3a b^2 x - b^3 x - \frac{6b a^2 c}{d} - \frac{2b^3 c}{d} + \frac{2b^2 (3e^{2dx+2c} a + e^{2dx+2c} b + 3a)}{d(e^{2dx+2c} + 1)^2} + \frac{3b \ln(e^{2dx+2c} + 1) a}{d}$

input `int((a+b*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(-1/2*b^3*tanh(d*x+c)^2-3*b^2*a*tanh(d*x+c)-1/2*(a^3+3*a^2*b+3*a*b^2+b^3)*ln(tanh(d*x+c)-1)+1/2*(a^3-3*a^2*b+3*a*b^2-b^3)*ln(tanh(d*x+c)+1))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 646 vs. 2(67) = 134.

Time = 0.09 (sec) , antiderivative size = 646, normalized size of antiderivative = 9.36

$$\int (a + b \tanh(c + dx))^3 dx = \text{Too large to display}$$

input `integrate((a+b*tanh(d*x+c))^3,x, algorithm="fricas")`

output

```
((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*cosh(d*x + c)^4 + 4*(a^3 - 3*a^2*b +
3*a*b^2 - b^3)*d*x*cosh(d*x + c)*sinh(d*x + c)^3 + (a^3 - 3*a^2*b + 3*a*b^
2 - b^3)*d*x*sinh(d*x + c)^4 + 6*a*b^2 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d
*x + 2*(3*a*b^2 + b^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*cosh(d*x + c)
^2 + 2*(3*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*cosh(d*x + c)^2 + 3*a*b^2 +
b^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*sinh(d*x + c)^2 + ((3*a^2*b + b
^3)*cosh(d*x + c)^4 + 4*(3*a^2*b + b^3)*cosh(d*x + c)*sinh(d*x + c)^3 + (3
*a^2*b + b^3)*sinh(d*x + c)^4 + 3*a^2*b + b^3 + 2*(3*a^2*b + b^3)*cosh(d*x
+ c)^2 + 2*(3*a^2*b + b^3 + 3*(3*a^2*b + b^3)*cosh(d*x + c)^2)*sinh(d*x +
c)^2 + 4*((3*a^2*b + b^3)*cosh(d*x + c)^3 + (3*a^2*b + b^3)*cosh(d*x + c)
)*sinh(d*x + c))*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c))) + 4*
((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*cosh(d*x + c)^3 + (3*a*b^2 + b^3 + (a
^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*cosh(d*x + c))*sinh(d*x + c))/(d*cosh(d
*x + c)^4 + 4*d*cosh(d*x + c)*sinh(d*x + c)^3 + d*sinh(d*x + c)^4 + 2*d*co
sh(d*x + c)^2 + 2*(3*d*cosh(d*x + c)^2 + d)*sinh(d*x + c)^2 + 4*(d*cosh(d*
x + c)^3 + d*cosh(d*x + c))*sinh(d*x + c) + d)
```

### Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.45

$$\int (a + b \tanh(c + dx))^3 dx$$

$$= \begin{cases} a^3 x + 3a^2 b x - \frac{3a^2 b \log(\tanh(c+dx)+1)}{d} + 3ab^2 x - \frac{3ab^2 \tanh(c+dx)}{d} + b^3 x - \frac{b^3 \log(\tanh(c+dx)+1)}{d} - \frac{b^3 \tanh^2(c+dx)}{2d} \\ x(a + b \tanh(c))^3 \end{cases}$$

input

```
integrate((a+b*tanh(d*x+c))**3,x)
```

output

```
Piecewise((a**3*x + 3*a**2*b*x - 3*a**2*b*log(tanh(c + d*x) + 1)/d + 3*a*b
**2*x - 3*a*b**2*tanh(c + d*x)/d + b**3*x - b**3*log(tanh(c + d*x) + 1)/d
- b**3*tanh(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tanh(c))**3, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.71

$$\int (a + b \tanh(c + dx))^3 dx$$

$$= b^3 \left( x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right)$$

$$+ 3ab^2 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^3x + \frac{3a^2b \log(\cosh(dx + c))}{d}$$

input `integrate((a+b*tanh(d*x+c))^3,x, algorithm="maxima")`output `b^3*(x + c/d + log(e^(-2*d*x - 2*c) + 1)/d + 2*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c) + 1))) + 3*a*b^2*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^3*x + 3*a^2*b*log(cosh(d*x + c))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.41

$$\int (a + b \tanh(c + dx))^3 dx$$

$$= \frac{(a^3 - 3a^2b + 3ab^2 - b^3)(dx + c) + (3a^2b + b^3) \log(e^{(2dx+2c)} + 1) + \frac{2(3ab^2 + (3ab^2 + b^3)e^{(2dx+2c)})}{(e^{(2dx+2c)} + 1)^2}}{d}$$

input `integrate((a+b*tanh(d*x+c))^3,x, algorithm="giac")`output `((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*(d*x + c) + (3*a^2*b + b^3)*log(e^(2*d*x + 2*c) + 1) + 2*(3*a*b^2 + (3*a*b^2 + b^3)*e^(2*d*x + 2*c))/(e^(2*d*x + 2*c) + 1)^2)/d`

**Mupad [B] (verification not implemented)**

Time = 2.01 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.12

$$\int (a + b \tanh(c + dx))^3 dx = x (a^3 + 3 a^2 b + 3 a b^2 + b^3) - \frac{\ln(\tanh(c + dx) + 1) (3 a^2 b + b^3)}{d} - \frac{b^3 \tanh(c + dx)^2}{2 d} - \frac{3 a b^2 \tanh(c + dx)}{d}$$

input `int((a + b*tanh(c + d*x))^3,x)`output `x*(3*a*b^2 + 3*a^2*b + a^3 + b^3) - (log(tanh(c + d*x) + 1)*(3*a^2*b + b^3))/d - (b^3*tanh(c + d*x)^2)/(2*d) - (3*a*b^2*tanh(c + d*x))/d`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 390, normalized size of antiderivative = 5.65

$$\int (a + b \tanh(c + dx))^3 dx = \frac{3e^{4dx+4c} \log(e^{2dx+2c} + 1) a^2 b + e^{4dx+4c} \log(e^{2dx+2c} + 1) b^3 + e^{4dx+4c} a^3 dx - 3e^{4dx+4c} a^2 b dx + 3e^{4dx+4c} a b^2 dx}{1}$$

input `int((a+b*tanh(d*x+c))^3,x)`output `(3*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1)*a**2*b + e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x) + 1)*b**3 + e**(4*c + 4*d*x)*a**3*d*x - 3*e**(4*c + 4*d*x)*a**2*b*d*x + 3*e**(4*c + 4*d*x)*a*b**2*d*x - 3*e**(4*c + 4*d*x)*a*b**2 - e**(4*c + 4*d*x)*b**3*d*x - e**(4*c + 4*d*x)*b**3 + 6*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1)*a**2*b + 2*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1)*b**3 + 2*e**(2*c + 2*d*x)*a**3*d*x - 6*e**(2*c + 2*d*x)*a**2*b*d*x + 6*e**(2*c + 2*d*x)*a*b**2*d*x - 2*e**(2*c + 2*d*x)*b**3*d*x + 3*log(e**(2*c + 2*d*x) + 1)*a**2*b + log(e**(2*c + 2*d*x) + 1)*b**3 + a**3*d*x - 3*a**2*b*d*x + 3*a*b**2*d*x + 3*a*b**2 - b**3*d*x - b**3)/(d*(e**(4*c + 4*d*x) + 2*e**(2*c + 2*d*x) + 1))`

### 3.60 $\int (a + b \tanh(c + dx))^2 dx$

Optimal result	518
Mathematica [A] (verified)	518
Rubi [A] (verified)	519
Maple [A] (verified)	520
Fricas [B] (verification not implemented)	521
Sympy [A] (verification not implemented)	521
Maxima [A] (verification not implemented)	522
Giac [A] (verification not implemented)	522
Mupad [B] (verification not implemented)	523
Reduce [B] (verification not implemented)	523

#### Optimal result

Integrand size = 12, antiderivative size = 38

$$\int (a + b \tanh(c + dx))^2 dx = (a^2 + b^2) x + \frac{2ab \log(\cosh(c + dx))}{d} - \frac{b^2 \tanh(c + dx)}{d}$$

output

```
(a^2+b^2)*x+2*a*b*ln(cosh(d*x+c))/d-b^2*tanh(d*x+c)/d
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

$$\int (a + b \tanh(c + dx))^2 dx = \frac{-(a + b)^2 \log(1 - \tanh(c + dx)) + (a - b)^2 \log(1 + \tanh(c + dx)) - 2b^2 \tanh(c + dx)}{2d}$$

input

```
Integrate[(a + b*Tanh[c + d*x])^2,x]
```

output

```
(-((a + b)^2*Log[1 - Tanh[c + d*x]]) + (a - b)^2*Log[1 + Tanh[c + d*x]] - 2*b^2*Tanh[c + d*x])/(2*d)
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3958, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tanh(c + dx))^2 dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a - ib \tan(ic + idx))^2 dx \\
 & \quad \downarrow \text{3958} \\
 & -2iab \int i \tanh(c + dx) dx + x(a^2 + b^2) - \frac{b^2 \tanh(c + dx)}{d} \\
 & \quad \downarrow \text{26} \\
 & 2ab \int \tanh(c + dx) dx + x(a^2 + b^2) - \frac{b^2 \tanh(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & 2ab \int -i \tan(ic + idx) dx + x(a^2 + b^2) - \frac{b^2 \tanh(c + dx)}{d} \\
 & \quad \downarrow \text{26} \\
 & -2iab \int \tan(ic + idx) dx + x(a^2 + b^2) - \frac{b^2 \tanh(c + dx)}{d} \\
 & \quad \downarrow \text{3956} \\
 & x(a^2 + b^2) + \frac{2ab \log(\cosh(c + dx))}{d} - \frac{b^2 \tanh(c + dx)}{d}
 \end{aligned}$$

input `Int[(a + b*Tanh[c + d*x])^2,x]`

output `(a^2 + b^2)*x + (2*a*b*Log[Cosh[c + d*x]])/d - (b^2*Tanh[c + d*x])/d`



## Definitions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_{x_}, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956  $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /;$   $\text{FreeQ}\{c, d\}, x]$

rule 3958  $\text{Int}[(a_ + (b_)*\tan[(c_.) + (d_.)*(x_)]^2, x\_Symbol] \rightarrow \text{Simp}[(a^2 - b^2)*x, x] + (\text{Simp}[b^2*(\text{Tan}[c + d*x]/d), x] + \text{Simp}[2*a*b \text{Int}[\text{Tan}[c + d*x], x], x]) /;$   $\text{FreeQ}\{a, b, c, d\}, x]$

## Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.37

method	result	size
parallelrisc	$-\frac{-a^2 dx + 2abdx - b^2 dx + 2 \ln(1 - \tanh(dx+c))ab + b^2 \tanh(dx+c)}{d}$	52
parts	$a^2 x + \frac{b^2 \left( -\tanh(dx+c) - \frac{\ln(\tanh(dx+c)-1)}{2} + \frac{\ln(\tanh(dx+c)+1)}{2} \right)}{d} + \frac{2ab \ln(\cosh(dx+c))}{d}$	59
derivativedivides	$\frac{-b^2 \tanh(dx+c) - \frac{(a^2 + 2ab + b^2) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^2 - 2ab + b^2) \ln(\tanh(dx+c)+1)}{2}}{d}$	61
default	$\frac{-b^2 \tanh(dx+c) - \frac{(a^2 + 2ab + b^2) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^2 - 2ab + b^2) \ln(\tanh(dx+c)+1)}{2}}{d}$	61
risc	$a^2 x - 2abx + b^2 x - \frac{4abc}{d} + \frac{2b^2}{d(e^{2dx+2c}+1)} + \frac{2ab \ln(e^{2dx+2c}+1)}{d}$	65

input  $\text{int}((a+b*\tanh(d*x+c))^2, x, \text{method}=\_RETURNVERBOSE)$

output  $-(-a^2*d*x+2*a*b*d*x-b^2*d*x+2*\ln(1-\tanh(d*x+c))*a*b+b^2*\tanh(d*x+c))/d$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 201 vs.  $2(38) = 76$ .

Time = 0.09 (sec) , antiderivative size = 201, normalized size of antiderivative = 5.29

$$\int (a + b \tanh(c + dx))^2 dx$$

$$= \frac{(a^2 - 2ab + b^2)dx \cosh(dx + c)^2 + 2(a^2 - 2ab + b^2)dx \cosh(dx + c) \sinh(dx + c) + (a^2 - 2ab + b^2)dx}{d \cosh(dx + c)}$$

input `integrate((a+b*tanh(d*x+c))^2,x, algorithm="fricas")`

output `((a^2 - 2*a*b + b^2)*d*x*cosh(d*x + c)^2 + 2*(a^2 - 2*a*b + b^2)*d*x*cosh(d*x + c)*sinh(d*x + c) + (a^2 - 2*a*b + b^2)*d*x*sinh(d*x + c)^2 + (a^2 - 2*a*b + b^2)*d*x + 2*b^2 + 2*(a*b*cosh(d*x + c)^2 + 2*a*b*cosh(d*x + c)*sinh(d*x + c) + a*b*sinh(d*x + c)^2 + a*b)*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c)))/(d*cosh(d*x + c)^2 + 2*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x + c)^2 + d)`

**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

$$\int (a + b \tanh(c + dx))^2 dx$$

$$= \begin{cases} a^2 x + 2abx - \frac{2ab \log(\tanh(c+dx)+1)}{d} + b^2 x - \frac{b^2 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a + b \tanh(c))^2 & \text{otherwise} \end{cases}$$

input `integrate((a+b*tanh(d*x+c))**2,x)`

output `Piecewise((a**2*x + 2*a*b*x - 2*a*b*log(tanh(c + d*x) + 1)/d + b**2*x - b**2*tanh(c + d*x)/d, Ne(d, 0)), (x*(a + b*tanh(c))**2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int (a + b \tanh(c + dx))^2 dx = b^2 \left( x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^2 x + \frac{2ab \log(\cosh(dx + c))}{d}$$

input `integrate((a+b*tanh(d*x+c))^2,x, algorithm="maxima")`output `b^2*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^2*x + 2*a*b*log(cosh(d*x + c))/d`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.47

$$\int (a + b \tanh(c + dx))^2 dx = \frac{2ab \log(e^{(2dx+2c)} + 1) + (a^2 - 2ab + b^2)(dx + c) + \frac{2b^2}{e^{(2dx+2c)} + 1}}{d}$$

input `integrate((a+b*tanh(d*x+c))^2,x, algorithm="giac")`output `(2*a*b*log(e^(2*d*x + 2*c) + 1) + (a^2 - 2*a*b + b^2)*(d*x + c) + 2*b^2/(e^(2*d*x + 2*c) + 1))/d`

**Mupad [B] (verification not implemented)**

Time = 1.98 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int (a + b \tanh(c + dx))^2 dx = x (a^2 + 2ab + b^2) - \frac{b^2 \tanh(c + dx)}{d} - \frac{2ab \ln(\tanh(c + dx) + 1)}{d}$$

input `int((a + b*tanh(c + d*x))^2,x)`output `x*(2*a*b + a^2 + b^2) - (b^2*tanh(c + d*x))/d - (2*a*b*log(tanh(c + d*x) + 1))/d`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.79

$$\int (a + b \tanh(c + dx))^2 dx = \frac{2e^{2dx+2c} \log(e^{2dx+2c} + 1) ab + e^{2dx+2c} a^2 dx - 2e^{2dx+2c} ab dx + e^{2dx+2c} b^2 dx - 2e^{2dx+2c} b^2 + 2 \log(e^{2dx+2c} + 1)}{d(e^{2dx+2c} + 1)}$$

input `int((a+b*tanh(d*x+c))^2,x)`output `(2*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x) + 1)*a*b + e**(2*c + 2*d*x)*a**2*d*x - 2*e**(2*c + 2*d*x)*a*b*d*x + e**(2*c + 2*d*x)*b**2*d*x - 2*e**(2*c + 2*d*x)*b**2 + 2*log(e**(2*c + 2*d*x) + 1)*a*b + a**2*d*x - 2*a*b*d*x + b**2*d*x)/(d*(e**(2*c + 2*d*x) + 1))`

### 3.61 $\int \frac{1}{a+b \tanh(c+dx)} dx$

Optimal result . . . . .	524
Mathematica [A] (verified) . . . . .	524
Rubi [A] (verified) . . . . .	525
Maple [A] (verified) . . . . .	526
Fricas [A] (verification not implemented) . . . . .	527
Sympy [B] (verification not implemented) . . . . .	527
Maxima [A] (verification not implemented) . . . . .	528
Giac [A] (verification not implemented) . . . . .	528
Mupad [B] (verification not implemented) . . . . .	529
Reduce [B] (verification not implemented) . . . . .	529

#### Optimal result

Integrand size = 12, antiderivative size = 50

$$\int \frac{1}{a+b \tanh(c+dx)} dx = \frac{ax}{a^2-b^2} - \frac{b \log(a \cosh(c+dx) + b \sinh(c+dx))}{(a^2-b^2)d}$$

output

```
a*x/(a^2-b^2)-b*ln(a*cosh(d*x+c)+b*sinh(d*x+c))/(a^2-b^2)/d
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.28

$$\int \frac{1}{a+b \tanh(c+dx)} dx = \frac{(-a+b) \log(1-\tanh(c+dx)) + (a+b) \log(1+\tanh(c+dx)) - 2b \log(a+b \tanh(c+dx))}{2(a-b)(a+b)d}$$

input

```
Integrate[(a + b*Tanh[c + d*x])^(-1), x]
```

output

```
((-a + b)*Log[1 - Tanh[c + d*x]] + (a + b)*Log[1 + Tanh[c + d*x]] - 2*b*Log[a + b*Tanh[c + d*x]])/(2*(a - b)*(a + b)*d)
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \tanh(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a - ib \tan(ic + idx)} dx \\
 & \quad \downarrow \text{3965} \\
 & \frac{ax}{a^2 - b^2} - \frac{ib \int -\frac{i(b+a \tanh(c+dx))}{a+b \tanh(c+dx)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b+a \tanh(c+dx)}{a+b \tanh(c+dx)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b-ia \tan(ic+idx)}{a-ib \tan(ic+idx)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{4013} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)}
 \end{aligned}$$

input `Int[(a + b*Tanh[c + d*x])^(-1),x]`

output `(a*x)/(a^2 - b^2) - (b*Log[a*Cosh[c + d*x] + b*Sinh[c + d*x]])/((a^2 - b^2)*d)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3965  $\text{Int}[(a + (b \cdot \tan[c + d \cdot x])^{-1}), x\_Symbol] \rightarrow \text{Simp}[a \cdot (x / (a^2 + b^2)), x] + \text{Simp}[b / (a^2 + b^2) \text{Int}[(b - a \cdot \tan[c + d \cdot x]) / (a + b \cdot \tan[c + d \cdot x]), x], x] /;$   $\text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4013  $\text{Int}[(c + (d \cdot \tan[e + f \cdot x]) / ((a + (b \cdot \tan[e + f \cdot x]) * (x))), x\_Symbol] \rightarrow \text{Simp}[(c / (b \cdot f)) * \text{Log}[\text{RemoveContent}[a \cdot \text{Cos}[e + f \cdot x] + b \cdot \text{Sin}[e + f \cdot x], x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a \cdot c + b \cdot d, 0]$

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.10

method	result	size
parallelrisc	$-\frac{-dxa-dxb-\ln(1-\tanh(dx+c))b+b\ln(a+b\tanh(dx+c))}{d(a^2-b^2)}$	55
derivativedivides	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{2b+2a} - \frac{b\ln(a+b\tanh(dx+c))}{(a-b)(a+b)} + \frac{\ln(\tanh(dx+c)+1)}{2a-2b}}{d}$	71
default	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{2b+2a} - \frac{b\ln(a+b\tanh(dx+c))}{(a-b)(a+b)} + \frac{\ln(\tanh(dx+c)+1)}{2a-2b}}{d}$	71
risc	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} + \frac{2bc}{d(a^2-b^2)} - \frac{b\ln\left(e^{2dx+2c} + \frac{a-b}{a+b}\right)}{d(a^2-b^2)}$	81

input `int(1/(a+b*tanh(d*x+c)),x,method=_RETURNVERBOSE)`output `-(-d*x*a-d*x*b-ln(1-tanh(d*x+c))*b+b*ln(a+b*tanh(d*x+c)))/d/(a^2-b^2)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.24

$$\int \frac{1}{a + b \tanh(c + dx)} dx = \frac{(a + b)dx - b \log\left(\frac{2(a \cosh(dx+c) + b \sinh(dx+c))}{\cosh(dx+c) - \sinh(dx+c)}\right)}{(a^2 - b^2)d}$$

input `integrate(1/(a+b*tanh(d*x+c)),x, algorithm="fricas")`

output `((a + b)*d*x - b*log(2*(a*cosh(d*x + c) + b*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))))/((a^2 - b^2)*d)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(37) = 74.

Time = 1.03 (sec) , antiderivative size = 224, normalized size of antiderivative = 4.48

$$\int \frac{1}{a + b \tanh(c + dx)} dx = \begin{cases} \frac{\infty x}{\tanh(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ -\frac{dx \tanh(c+dx)}{2bd \tanh(c+dx) - 2bd} + \frac{dx}{2bd \tanh(c+dx) - 2bd} + \frac{1}{2bd \tanh(c+dx) - 2bd} & \text{for } a = -b \\ \frac{dx \tanh(c+dx)}{2bd \tanh(c+dx) + 2bd} + \frac{dx}{2bd \tanh(c+dx) + 2bd} - \frac{1}{2bd \tanh(c+dx) + 2bd} & \text{for } a = b \\ \frac{x}{a+b \tanh(c)} & \text{for } d = 0 \\ \frac{adx}{a^2d - b^2d} - \frac{bdx}{a^2d - b^2d} - \frac{b \log\left(\frac{a}{b} + \tanh(c+dx)\right)}{a^2d - b^2d} + \frac{b \log(\tanh(c+dx)+1)}{a^2d - b^2d} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+b*tanh(d*x+c)),x)`



output

```
Piecewise((zoo*x/tanh(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a, Eq(b, 0))
, (-d*x*tanh(c + d*x)/(2*b*d*tanh(c + d*x) - 2*b*d) + d*x/(2*b*d*tanh(c +
d*x) - 2*b*d) + 1/(2*b*d*tanh(c + d*x) - 2*b*d), Eq(a, -b)), (d*x*tanh(c +
d*x)/(2*b*d*tanh(c + d*x) + 2*b*d) + d*x/(2*b*d*tanh(c + d*x) + 2*b*d) -
1/(2*b*d*tanh(c + d*x) + 2*b*d), Eq(a, b)), (x/(a + b*tanh(c)), Eq(d, 0)),
(a*d*x/(a**2*d - b**2*d) - b*d*x/(a**2*d - b**2*d) - b*log(a/b + tanh(c +
d*x))/(a**2*d - b**2*d) + b*log(tanh(c + d*x) + 1)/(a**2*d - b**2*d), Tru
e))
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12

$$\int \frac{1}{a + b \tanh(c + dx)} dx = -\frac{b \log(-(a - b)e^{(-2dx - 2c)} - a - b)}{(a^2 - b^2)d} + \frac{dx + c}{(a + b)d}$$

input

```
integrate(1/(a+b*tanh(d*x+c)),x, algorithm="maxima")
```

output

```
-b*log(-(a - b)*e^(-2*d*x - 2*c) - a - b)/((a^2 - b^2)*d) + (d*x + c)/((a
+ b)*d)
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.24

$$\int \frac{1}{a + b \tanh(c + dx)} dx = -\frac{b \log(|ae^{(2dx+2c)} + be^{(2dx+2c)} + a - b|)}{a^2 - b^2} - \frac{dx + c}{a - b}$$

input

```
integrate(1/(a+b*tanh(d*x+c)),x, algorithm="giac")
```

output

```
-(b*log(abs(a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) + a - b))/(a^2 - b^2) -
(d*x + c)/(a - b))/d
```

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.20

$$\int \frac{1}{a + b \tanh(c + dx)} dx$$

$$= \frac{ax - bx}{a^2 - b^2} + \frac{b(\ln(\tanh(c + dx) + 1) - \ln(a + b \tanh(c + dx)))}{d(a^2 - b^2)}$$

input `int(1/(a + b*tanh(c + d*x)),x)`output `(a*x - b*x)/(a^2 - b^2) + (b*(log(tanh(c + d*x) + 1) - log(a + b*tanh(c + d*x))))/(d*(a^2 - b^2))`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.14

$$\int \frac{1}{a + b \tanh(c + dx)} dx = \frac{-\log(e^{2dx+2c}a + e^{2dx+2c}b + a - b) b + adx + bdx}{d(a^2 - b^2)}$$

input `int(1/(a+b*tanh(d*x+c)),x)`output `( - log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*b + a*d*x + b*d*x )/(d*(a**2 - b**2))`

### 3.62 $\int \frac{1}{(a+b \tanh(c+dx))^2} dx$

Optimal result . . . . .	530
Mathematica [A] (verified) . . . . .	530
Rubi [A] (verified) . . . . .	531
Maple [A] (verified) . . . . .	533
Fricas [B] (verification not implemented) . . . . .	533
Sympy [B] (verification not implemented) . . . . .	534
Maxima [A] (verification not implemented) . . . . .	535
Giac [A] (verification not implemented) . . . . .	536
Mupad [B] (verification not implemented) . . . . .	536
Reduce [B] (verification not implemented) . . . . .	537

#### Optimal result

Integrand size = 12, antiderivative size = 85

$$\int \frac{1}{(a+b \tanh(c+dx))^2} dx = \frac{(a^2+b^2)x}{(a^2-b^2)^2} - \frac{2ab \log(a \cosh(c+dx) + b \sinh(c+dx))}{(a^2-b^2)^2 d} + \frac{b}{(a^2-b^2)d(a+b \tanh(c+dx))}$$

output

```
(a^2+b^2)*x/(a^2-b^2)^2-2*a*b*ln(a*cosh(d*x+c)+b*sinh(d*x+c))/(a^2-b^2)^2/d+b/(a^2-b^2)/d/(a+b*tanh(d*x+c))
```

#### Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.11

$$\int \frac{1}{(a+b \tanh(c+dx))^2} dx = \frac{-\frac{\log(1-\tanh(c+dx))}{(a+b)^2} + \frac{\log(1+\tanh(c+dx))}{(a-b)^2} + \frac{2b(-2a \log(a+b \tanh(c+dx)) + \frac{a^2-b^2}{a+b \tanh(c+dx)})}{(a^2-b^2)^2}}{2d}$$

input

```
Integrate[(a + b*Tanh[c + d*x])^(-2), x]
```

output

$$\frac{(-\text{Log}[1 - \text{Tanh}[c + d*x]]/(a + b)^2) + \text{Log}[1 + \text{Tanh}[c + d*x]]/(a - b)^2 + (2*b*(-2*a*\text{Log}[a + b*\text{Tanh}[c + d*x]] + (a^2 - b^2)/(a + b*\text{Tanh}[c + d*x])))/(a^2 - b^2)^2)/(2*d)}$$
**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {3042, 3964, 3042, 4014, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + b \tanh(c + dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(a - ib \tan(ic + idx))^2} dx \\ & \quad \downarrow \text{3964} \\ & \frac{\int \frac{a - b \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2} + \frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} \\ & \quad \downarrow \text{3042} \\ & \frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\int \frac{a + ib \tan(ic + idx)}{a - ib \tan(ic + idx)} dx}{a^2 - b^2} \\ & \quad \downarrow \text{4014} \\ & \frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\frac{x(a^2 + b^2)}{a^2 - b^2} - \frac{2iab \int -\frac{i(b + a \tanh(c + dx))}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} \\ & \quad \downarrow \text{26} \\ & \frac{\frac{x(a^2 + b^2)}{a^2 - b^2} - \frac{2ab \int \frac{b + a \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} + \frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\frac{x(a^2 + b^2)}{a^2 - b^2} - \frac{2ab \int \frac{b - ia \tan(ic + idx)}{a - ib \tan(ic + idx)} dx}{a^2 - b^2}}{a^2 - b^2}$$

↓ 4013

$$\frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\frac{x(a^2 + b^2)}{a^2 - b^2} - \frac{2ab \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)}}{a^2 - b^2}$$

input `Int[(a + b*Tanh[c + d*x])^(-2), x]`

output `((a^2 + b^2)*x)/(a^2 - b^2) - (2*a*b*Log[a*Cosh[c + d*x] + b*Sinh[c + d*x]])/((a^2 - b^2)*d)/(a^2 - b^2) + b/((a^2 - b^2)*d*(a + b*Tanh[c + d*x]))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3964 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] :> Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

**Maple [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.09

method	result
derivativedivides	$-\frac{\ln(\tanh(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^2} + \frac{b}{(a-b)(a+b)(a+b \tanh(dx+c))} - \frac{2ab \ln(a+b \tanh(dx+c))}{(a+b)^2(a-b)^2}$
default	$-\frac{\ln(\tanh(dx+c)-1)}{2(a+b)^2} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^2} + \frac{b}{(a-b)(a+b)(a+b \tanh(dx+c))} - \frac{2ab \ln(a+b \tanh(dx+c))}{(a+b)^2(a-b)^2}$
risch	$\frac{x}{a^2+2ab+b^2} + \frac{4abx}{a^4-2a^2b^2+b^4} + \frac{4abc}{d(a^4-2a^2b^2+b^4)} + \frac{2b^2}{(a-b)d(a^2+2ab+b^2)(e^{2dx+2c}a+e^{2dx+2c}b+a-b)} - \frac{2ab \ln(a+b \tanh(dx+c))}{(a+b)^2(a-b)^2}$
parallelrisc	$-\frac{2 \ln(1-\tanh(dx+c)) \tanh(dx+c) a^2 b^2 + 2 \ln(a+b \tanh(dx+c)) \tanh(dx+c) a^2 b^2 - b^4 \tanh(dx+c) - a^2 b^2 dx + a^2 b^2 \tanh(dx+c)}{(a^4 - 2a^2b^2 + b^4)d}$

input

```
int(1/(a+b*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-1/2/(a+b)^2*ln(tanh(d*x+c)-1)+1/2/(a-b)^2*ln(tanh(d*x+c)+1)+b/(a-b)/
(a+b)/(a+b*tanh(d*x+c))-2*a*b/(a+b)^2/(a-b)^2*ln(a+b*tanh(d*x+c)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 422 vs. 2(85) = 170.

Time = 0.09 (sec) , antiderivative size = 422, normalized size of antiderivative = 4.96

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx$$

$$= \frac{(a^3 + 3a^2b + 3ab^2 + b^3)dx \cosh(dx + c)^2 + 2(a^3 + 3a^2b + 3ab^2 + b^3)dx \cosh(dx + c) \sinh(dx + c) + (a^5 + a^4b - 2a^3b^2 - 2a^2b^3 + ab^4 + b^5)d \cosh(dx + c) \sinh(dx + c)}{(a^5 + a^4b - 2a^3b^2 - 2a^2b^3 + ab^4 + b^5)d \cosh(dx + c) \sinh(dx + c)}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="fricas")
```

output

```
((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x*cosh(d*x + c)^2 + 2*(a^3 + 3*a^2*b +
3*a*b^2 + b^3)*d*x*cosh(d*x + c)*sinh(d*x + c) + (a^3 + 3*a^2*b + 3*a*b^2
+ b^3)*d*x*sinh(d*x + c)^2 + 2*a*b^2 - 2*b^3 + (a^3 + a^2*b - a*b^2 - b^3)
*d*x - 2*(a^2*b - a*b^2 + (a^2*b + a*b^2)*cosh(d*x + c)^2 + 2*(a^2*b + a*b
^2)*cosh(d*x + c)*sinh(d*x + c) + (a^2*b + a*b^2)*sinh(d*x + c)^2)*log(2*(
a*cosh(d*x + c) + b*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c)))/((a^5
+ a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*cosh(d*x + c)^2 + 2*(a^5
+ a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*cosh(d*x + c)*sinh(d*x +
c) + (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*sinh(d*x + c)^
2 + (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*d)
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1389 vs.  $2(70) = 140$ .

Time = 9.21 (sec) , antiderivative size = 1389, normalized size of antiderivative = 16.34

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*tanh(d*x+c))**2,x)
```

output

```
Piecewise((zoo*x/tanh(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**2, Eq(b, 0)), (d*x*tanh(c + d*x)**2/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - 2*d*x*tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + d*x/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + 2/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d), Eq(a, -b)), (d*x*tanh(c + d*x)**2/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + 2*d*x*tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + d*x/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - 2/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d), Eq(a, b)), (x/(a + b*tanh(c))**2, Eq(d, 0)), (a**3*d*x/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*a**3*b**2*d - 2*a**2*b**3*d*tanh(c + d*x) + a*b**4*d + b**5*d*tanh(c + d*x)) + a**2*b*d*x*tanh(c + d*x)/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*a**3*b**2*d - 2*a**2*b**3*d*tanh(c + d*x) + a*b**4*d + b**5*d*tanh(c + d*x)) - 2*a**2*b*d*x/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*a**3*b**2*d - 2*a**2*b**3*d*tanh(c + d*x) + a*b**4*d + b**5*d*tanh(c + d*x)) - 2*a**2*b*log(a/b + tanh(c + d*x))/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*a**3*b**2*d - 2*a**2*b**3*d*tanh(c + d*x) + a*b**4*d + b**5*d*tanh(c + d*x)) + 2*a**2*b*1...
```

### Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.49

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx = -\frac{2ab \log(-(a-b)e^{(-2dx-2c)} - a - b)}{(a^4 - 2a^2b^2 + b^4)d} - \frac{2b^2}{(a^4 - 2a^2b^2 + b^4 + (a^4 - 2a^3b + 2ab^3 - b^4)e^{(-2dx-2c)})d} + \frac{dx + c}{(a^2 + 2ab + b^2)d}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="maxima")
```

output

```
-2*a*b*log(-(a - b)*e^(-2*d*x - 2*c) - a - b)/((a^4 - 2*a^2*b^2 + b^4)*d) - 2*b^2/((a^4 - 2*a^2*b^2 + b^4 + (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^(-2*d*x - 2*c))*d) + (d*x + c)/((a^2 + 2*a*b + b^2)*d)
```



**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.55

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx$$

$$= -\frac{\frac{2ab \log(|-ae^{(2dx+2c)} - be^{(2dx+2c)} - a + b|)}{a^4 - 2a^2b^2 + b^4} - \frac{dx+c}{a^2 - 2ab + b^2} - \frac{2(ab^2 - b^3)}{(ae^{(2dx+2c)} + be^{(2dx+2c)} + a - b)(a+b)^2(a-b)^2}}{d}$$

input `integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="giac")`

output

```
-(2*a*b*log(abs(-a*e^(2*d*x + 2*c) - b*e^(2*d*x + 2*c) - a + b))/(a^4 - 2*
a^2*b^2 + b^4) - (d*x + c)/(a^2 - 2*a*b + b^2) - 2*(a*b^2 - b^3)/((a*e^(2*
d*x + 2*c) + b*e^(2*d*x + 2*c) + a - b)*(a + b)^2*(a - b)^2))/d
```

**Mupad [B] (verification not implemented)**

Time = 2.38 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.49

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx = \frac{\frac{ax}{(a+b)^2} + \frac{bx \tanh(c+dx)}{(a+b)^2} - \frac{b^2 \tanh(c+dx)}{ad(a^2-b^2)}}{a + b \tanh(c + dx)}$$

$$- \frac{2ab \ln(a + b \tanh(c + dx))}{d(a^4 - 2a^2b^2 + b^4)}$$

$$+ \frac{2ab \ln(\tanh(c + dx) + 1)}{d(a^2 - b^2)^2}$$

input `int(1/(a + b*tanh(c + d*x))^2,x)`

output

```
((a*x)/(a + b)^2 + (b*x*tanh(c + d*x))/(a + b)^2 - (b^2*tanh(c + d*x))/(a*
d*(a^2 - b^2)))/(a + b*tanh(c + d*x)) - (2*a*b*log(a + b*tanh(c + d*x)))/(
d*(a^4 + b^4 - 2*a^2*b^2)) + (2*a*b*log(tanh(c + d*x) + 1))/(d*(a^2 - b^2)
^2)
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 428, normalized size of antiderivative = 5.04

$$\int \frac{1}{(a + b \tanh(c + dx))^2} dx$$

$$= \frac{-2e^{2dx+2c} \log(e^{2dx+2c}a + e^{2dx+2c}b + a - b) a^2 b - 2e^{2dx+2c} \log(e^{2dx+2c}a + e^{2dx+2c}b + a - b) a b^2 + e^{2dx+2c} a^3 + e^{2dx+2c} a^2 b + e^{2dx+2c} a b^2 + e^{2dx+2c} b^3}{d(e^{2dx+2c} a^5 + e^{2dx+2c} a^4 b + e^{2dx+2c} a^3 b^2 + e^{2dx+2c} a^2 b^3 + e^{2dx+2c} a b^4 + e^{2dx+2c} b^5)}$$

input `int(1/(a+b*tanh(d*x+c))^2,x)`

output

```
( - 2*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)
*a**2*b - 2*e**(2*c + 2*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b
+ a - b)*a*b**2 + e**(2*c + 2*d*x)*a**3*d*x + 3*e**(2*c + 2*d*x)*a**2*b*d*
x + 3*e**(2*c + 2*d*x)*a*b**2*d*x - 2*e**(2*c + 2*d*x)*a*b**2 + e**(2*c +
2*d*x)*b**3*d*x - 2*e**(2*c + 2*d*x)*b**3 - 2*log(e**(2*c + 2*d*x)*a + e**
(2*c + 2*d*x)*b + a - b)*a**2*b + 2*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d
*x)*b + a - b)*a*b**2 + a**3*d*x + a**2*b*d*x - a*b**2*d*x - b**3*d*x)/(d*
(e**(2*c + 2*d*x)*a**5 + e**(2*c + 2*d*x)*a**4*b - 2*e**(2*c + 2*d*x)*a**3
*b**2 - 2*e**(2*c + 2*d*x)*a**2*b**3 + e**(2*c + 2*d*x)*a*b**4 + e**(2*c +
2*d*x)*b**5 + a**5 - a**4*b - 2*a**3*b**2 + 2*a**2*b**3 + a*b**4 - b**5))
```

### 3.63 $\int \frac{1}{(a+b \tanh(c+dx))^3} dx$

Optimal result . . . . .	538
Mathematica [A] (verified) . . . . .	539
Rubi [A] (verified) . . . . .	539
Maple [A] (verified) . . . . .	542
Fricas [B] (verification not implemented) . . . . .	542
Sympy [B] (verification not implemented) . . . . .	543
Maxima [B] (verification not implemented) . . . . .	544
Giac [A] (verification not implemented) . . . . .	545
Mupad [B] (verification not implemented) . . . . .	545
Reduce [B] (verification not implemented) . . . . .	546

#### Optimal result

Integrand size = 12, antiderivative size = 129

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = \frac{a(a^2 + 3b^2)x}{(a^2 - b^2)^3} - \frac{b(3a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^3 d} + \frac{b}{2(a^2 - b^2)d(a + b \tanh(c + dx))^2} + \frac{2ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))}$$

output

```
a*(a^2+3*b^2)*x/(a^2-b^2)^3-b*(3*a^2+b^2)*ln(a*cosh(d*x+c)+b*sinh(d*x+c))/(a^2-b^2)^3/d+1/2*b/(a^2-b^2)/d/(a+b*tanh(d*x+c))^2+2*a*b/(a^2-b^2)^2/d/(a+b*tanh(d*x+c))
```

**Mathematica [A] (verified)**

Time = 1.56 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx$$

$$= \frac{-\frac{\log(1-\tanh(c+dx))}{(a+b)^3} + \frac{\log(1+\tanh(c+dx))}{(a-b)^3} + \frac{b \left( -2(3a^2+b^2) \log(a+b \tanh(c+dx)) + \frac{(a^2-b^2)(5a^2-b^2+4ab \tanh(c+dx))}{(a+b \tanh(c+dx))^2} \right)}{(a^2-b^2)^3}}{2d}$$

input `Integrate[(a + b*Tanh[c + d*x])^(-3), x]`

output `(-(Log[1 - Tanh[c + d*x]]/(a + b)^3) + Log[1 + Tanh[c + d*x]]/(a - b)^3 + (b*(-2*(3*a^2 + b^2)*Log[a + b*Tanh[c + d*x]] + ((a^2 - b^2)*(5*a^2 - b^2 + 4*a*b*Tanh[c + d*x]))/(a + b*Tanh[c + d*x])^2))/(a^2 - b^2)^3)/(2*d)`

**Rubi [A] (verified)**

Time = 0.73 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.20, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {3042, 3964, 3042, 4012, 3042, 4014, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(a - ib \tan(ic + idx))^3} dx$$

$$\downarrow \text{3964}$$

$$\frac{\int \frac{a - b \tanh(c + dx)}{(a + b \tanh(c + dx))^2} dx}{a^2 - b^2} + \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{\int \frac{a + ib \tan(ic + idx)}{(a - ib \tan(ic + idx))^2} dx}{a^2 - b^2} \\
& \quad \downarrow 4012 \\
& \frac{\int \frac{a^2 - 2b \tanh(c + dx)a + b^2}{a + b \tanh(c + dx)} dx}{a^2 - b^2} + \frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\int \frac{a^2 + 2ib \tan(ic + idx)a + b^2}{a - ib \tan(ic + idx)} dx}{a^2 - b^2} \\
& \quad \downarrow 4014 \\
& \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \\
& \quad \frac{\frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{ax(a^2 + 3b^2)}{a^2 - b^2} - \frac{ib(3a^2 + b^2) \int \frac{b + a \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} \\
& \quad \downarrow 26 \\
& \frac{\frac{ax(a^2 + 3b^2)}{a^2 - b^2} - \frac{b(3a^2 + b^2) \int \frac{b + a \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} + \frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \\
& \quad \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \\
& \quad \frac{\frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{ax(a^2 + 3b^2)}{a^2 - b^2} - \frac{b(3a^2 + b^2) \int \frac{b - ia \tan(ic + idx)}{a - ib \tan(ic + idx)} dx}{a^2 - b^2}}{a^2 - b^2} \\
& \quad \downarrow 4013 \\
& \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \\
& \quad \frac{\frac{2ab}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{ax(a^2 + 3b^2)}{a^2 - b^2} - \frac{b(3a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)}}{a^2 - b^2}
\end{aligned}$$

input `Int[(a + b*Tanh[c + d*x])^(-3),x]`

output `b/(2*(a^2 - b^2)*d*(a + b*Tanh[c + d*x])^2) + (((a*(a^2 + 3*b^2)*x)/(a^2 - b^2) - (b*(3*a^2 + b^2)*Log[a*Cosh[c + d*x] + b*Sinh[c + d*x]])/(a^2 - b^2)*d)/(a^2 - b^2) + (2*a*b)/((a^2 - b^2)*d*(a + b*Tanh[c + d*x]))/(a^2 - b^2)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3964 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

**Maple [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{b}{2(a-b)(a+b)(a+b \tanh(dx+c))^2} + \frac{2ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))} - \frac{b(3a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^3(a-b)^3} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^3} - \frac{\ln(\tanh(dx+c)-1)}{2(a-b)^3}$
default	$\frac{b}{2(a-b)(a+b)(a+b \tanh(dx+c))^2} + \frac{2ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))} - \frac{b(3a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^3(a-b)^3} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^3} - \frac{\ln(\tanh(dx+c)-1)}{2(a-b)^3}$
risch	$\frac{x}{a^3+3a^2b+3b^2a+b^3} + \frac{6b a^2 x}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{2b^3 x}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{6b a^2 c}{d(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{2}{d(a^6-3a^4b^2+3a^2b^4-b^6)}$
parallelrisc	$-\frac{2a^5 b^2 dx - 6a^4 b^3 dx - 2x \tanh(dx+c)^2 b^7 d - 6 \ln(1 - \tanh(dx+c)) \tanh(dx+c)^2 a^2 b^5 + 6 \ln(a+b \tanh(dx+c)) \tanh(dx+c)^2 a^2 b^5}{d(a^6-3a^4b^2+3a^2b^4-b^6)}$

input

```
int(1/(a+b*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(1/2*b/(a-b)/(a+b)/(a+b*tanh(d*x+c))^2+2*a*b/(a+b)^2/(a-b)^2/(a+b*tanh
(d*x+c))-b*(3*a^2+b^2)/(a+b)^3/(a-b)^3*ln(a+b*tanh(d*x+c))+1/2/(a-b)^3*ln(
tanh(d*x+c)+1)-1/2/(a+b)^3*ln(tanh(d*x+c)-1))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1427 vs. 2(127) = 254.

Time = 0.11 (sec) , antiderivative size = 1427, normalized size of antiderivative = 11.06

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="fricas")
```

output

```
((a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x*cosh(d*x +
c)^4 + 4*(a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x*cos
h(d*x + c)*sinh(d*x + c)^3 + (a^5 + 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*
a*b^4 + b^5)*d*x*sinh(d*x + c)^4 + 6*a^3*b^2 - 12*a^2*b^3 + 6*a*b^4 + (a^5
+ a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*x + 2*(3*a^3*b^2 - a^2*b
^3 - 3*a*b^4 + b^5 + (a^5 + 3*a^4*b + 2*a^3*b^2 - 2*a^2*b^3 - 3*a*b^4 - b^
5)*d*x)*cosh(d*x + c)^2 + 2*(3*a^3*b^2 - a^2*b^3 - 3*a*b^4 + b^5 + 3*(a^5
+ 5*a^4*b + 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x*cosh(d*x + c)^2 +
(a^5 + 3*a^4*b + 2*a^3*b^2 - 2*a^2*b^3 - 3*a*b^4 - b^5)*d*x)*sinh(d*x + c
)^2 - (3*a^4*b - 6*a^3*b^2 + 4*a^2*b^3 - 2*a*b^4 + b^5 + (3*a^4*b + 6*a^3*
b^2 + 4*a^2*b^3 + 2*a*b^4 + b^5)*cosh(d*x + c)^4 + 4*(3*a^4*b + 6*a^3*b^2
+ 4*a^2*b^3 + 2*a*b^4 + b^5)*cosh(d*x + c)*sinh(d*x + c)^3 + (3*a^4*b + 6*
a^3*b^2 + 4*a^2*b^3 + 2*a*b^4 + b^5)*sinh(d*x + c)^4 + 2*(3*a^4*b - 2*a^2*
b^3 - b^5)*cosh(d*x + c)^2 + 2*(3*a^4*b - 2*a^2*b^3 - b^5 + 3*(3*a^4*b + 6
*a^3*b^2 + 4*a^2*b^3 + 2*a*b^4 + b^5)*cosh(d*x + c)^2)*sinh(d*x + c)^2 + 4
*((3*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + 2*a*b^4 + b^5)*cosh(d*x + c)^3 + (3*a
^4*b - 2*a^2*b^3 - b^5)*cosh(d*x + c))*sinh(d*x + c))*log(2*(a*cosh(d*x +
c) + b*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))) + 4*((a^5 + 5*a^4*b
+ 10*a^3*b^2 + 10*a^2*b^3 + 5*a*b^4 + b^5)*d*x*cosh(d*x + c)^3 + (3*a^3*b
^2 - a^2*b^3 - 3*a*b^4 + b^5 + (a^5 + 3*a^4*b + 2*a^3*b^2 - 2*a^2*b^3 - ...
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6412 vs.  $2(107) = 214$ .

Time = 12.43 (sec) , antiderivative size = 6412, normalized size of antiderivative = 49.71

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*tanh(d*x+c))**3,x)
```



output

```
Piecewise((zoo*x/tanh(c)**3, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**3, Eq(
b, 0)), (-3*d*x*tanh(c + d*x)**3/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*t
anh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) + 9*d*x*tanh(c + d*
x)**2/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d
*tanh(c + d*x) - 24*b**3*d) - 9*d*x*tanh(c + d*x)/(24*b**3*d*tanh(c + d*x)
**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) +
3*d*x/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d
*tanh(c + d*x) - 24*b**3*d) + 3*tanh(c + d*x)**2/(24*b**3*d*tanh(c + d*x)*
**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) - 9
*tanh(c + d*x)/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 +
72*b**3*d*tanh(c + d*x) - 24*b**3*d) + 10/(24*b**3*d*tanh(c + d*x)**3 - 72
*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d), Eq(a, -b)
), (3*d*x*tanh(c + d*x)**3/(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c
+ d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) + 9*d*x*tanh(c + d*x)**2/
(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(
c + d*x) + 24*b**3*d) + 9*d*x*tanh(c + d*x)/(24*b**3*d*tanh(c + d*x)**3 +
72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) + 3*d*x/
(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(
c + d*x) + 24*b**3*d) - 3*tanh(c + d*x)**2/(24*b**3*d*tanh(c + d*x)**3 + 7
2*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) - 9*ta...
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 325 vs.  $2(127) = 254$ .

Time = 0.08 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.52

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = -\frac{(3a^2b + b^3) \log(-(a - b)e^{(-2dx - 2c)} - a - b)}{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)d}$$

$$-\frac{2(3a^2b^2 + 3ab^3 + (3a^2b^2 - a^7 + a^6b - 3a^5b^2 - 3a^4b^3 + 3a^3b^4 + 3a^2b^5 - ab^6 - b^7 + 2(a^7 - a^6b - 3a^5b^2 + 3a^4b^3 + 3a^3b^4 - 3a^2b^5 - ab^6 - b^7)dx + c)}{(a^3 + 3a^2b + 3ab^2 + b^3)d}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="maxima")
```

output

$$\begin{aligned} & -(3a^2b + b^3) \log(-(a - b)e^{-2dx - 2c} - a - b) / ((a^6 - 3a^4b^2 + 3a^2b^4 - b^6) * d) - 2 * (3a^2b^2 + 3ab^3 + (3a^2b^2 - 2ab^3 - b^4) * e^{-2dx - 2c}) / ((a^7 + a^6b - 3a^5b^2 - 3a^4b^3 + 3a^3b^4 + 3a^2b^5 - ab^6 - b^7 + 2(a^7 - a^6b - 3a^5b^2 + 3a^4b^3 + 3a^3b^4 - 3a^2b^5 - ab^6 + b^7) * e^{-2dx - 2c} + (a^7 - 3a^6b + a^5b^2 + 5a^4b^3 - 5a^3b^4 - a^2b^5 + 3ab^6 - b^7) * e^{-4dx - 4c}) * d) + (dx + c) / ((a^3 + 3a^2b + 3ab^2 + b^3) * d) \end{aligned}$$

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.59

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = \frac{\frac{(3a^2b + b^3) \log(|-ae^{(2dx+2c)} - be^{(2dx+2c)} - a + b|)}{a^6 - 3a^4b^2 + 3a^2b^4 - b^6} - \frac{dx + c}{a^3 - 3a^2b + 3ab^2 - b^3} - \frac{2 \left( (3a^2b^2 - 4ab^3 + b^4) e^{(2dx+2c)} + \frac{3(a^3b^2 - 2a^2b^3 + ab^4)}{a+b} \right)}{(ae^{(2dx+2c)} + be^{(2dx+2c)} + a - b)^2 (a+b)^2 (a-b)^3}}{d}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="giac")
```

output

$$\begin{aligned} & -((3a^2b + b^3) \log(\text{abs}(-a * e^{(2dx + 2c)} - b * e^{(2dx + 2c)} - a + b)) / (a^6 - 3a^4b^2 + 3a^2b^4 - b^6) - (dx + c) / (a^3 - 3a^2b + 3ab^2 - b^3) - 2 * ((3a^2b^2 - 4ab^3 + b^4) * e^{(2dx + 2c)} + 3 * (a^3b^2 - 2a^2b^3 + ab^4) / (a + b)) / ((a * e^{(2dx + 2c)} + b * e^{(2dx + 2c)} + a - b)^2 * (a + b)^2 * (a - b)^3)) / d \end{aligned}$$

**Mupad [B] (verification not implemented)**

Time = 2.99 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.36

$$\begin{aligned} & \int \frac{1}{(a + b \tanh(c + dx))^3} dx \\ & = \frac{\tanh(c + dx) \left( \frac{1}{ad} - \frac{a^4 + a^2b^2}{ad(a^4 - 2a^2b^2 + b^4)} \right) + \frac{a^2x}{(a+b)(a^2 + 2ab + b^2)} + \frac{b^2x \tanh(c+dx)^2}{a^3 + 3a^2b + 3ab^2 + b^3} + \frac{\tanh(c+dx)^2 \left( \frac{b^5}{2} - \frac{5a^2b^3}{2} \right)}{a^2d(a^4 - 2a^2b^2 + b^4)} + \frac{2c}{a^3} - \frac{\ln(a + b \tanh(c + dx)) (3a^2b + b^3)}{d(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)} + \frac{\ln(\tanh(c + dx) + 1) (3a^2b + b^3)}{d(a^2 - b^2)^3} \end{aligned}$$

input `int(1/(a + b*tanh(c + d*x))^3,x)`

output 
$$\begin{aligned} & (\tanh(c + d*x)*(1/(a*d) - (a^4 + a^2*b^2)/(a*d*(a^4 + b^4 - 2*a^2*b^2))) + \\ & (a^2*x)/((a + b)*(2*a*b + a^2 + b^2)) + (b^2*x*\tanh(c + d*x)^2)/(3*a*b^2 \\ & + 3*a^2*b + a^3 + b^3) + (\tanh(c + d*x)^2*(b^5/2 - (5*a^2*b^3)/2))/(a^2*d* \\ & (a^4 + b^4 - 2*a^2*b^2)) + (2*a*b*x*\tanh(c + d*x))/(3*a*b^2 + 3*a^2*b + a^ \\ & 3 + b^3))/(a^2 + b^2*\tanh(c + d*x)^2 + 2*a*b*\tanh(c + d*x)) - (\log(a + b*t \\ & \tanh(c + d*x))*(3*a^2*b + b^3))/(d*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)) + ( \\ & \log(\tanh(c + d*x) + 1)*(3*a^2*b + b^3))/(d*(a^2 - b^2)^3) \end{aligned}$$

### Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 1186, normalized size of antiderivative = 9.19

$$\int \frac{1}{(a + b \tanh(c + dx))^3} dx = \text{Too large to display}$$

input `int(1/(a+b*tanh(d*x+c))^3,x)`

output

```
( - 3*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)
*a**4*b - 6*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b
+ a - b)*a**3*b**2 - 4*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c +
2*d*x)*b + a - b)*a**2*b**3 - 2*e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)*a +
e**(2*c + 2*d*x)*b + a - b)*a*b**4 - e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)
)*a + e**(2*c + 2*d*x)*b + a - b)*b**5 + e**(4*c + 4*d*x)*a**5*d*x + 5*e**
(4*c + 4*d*x)*a**4*b*d*x + 10*e**(4*c + 4*d*x)*a**3*b**2*d*x - 3*e**(4*c +
4*d*x)*a**3*b**2 + 10*e**(4*c + 4*d*x)*a**2*b**3*d*x - 5*e**(4*c + 4*d*x)
*a**2*b**3 + 5*e**(4*c + 4*d*x)*a*b**4*d*x - e**(4*c + 4*d*x)*a*b**4 + e**
(4*c + 4*d*x)*b**5*d*x + e**(4*c + 4*d*x)*b**5 - 6*e**(2*c + 2*d*x)*log(e
*(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**4*b + 4*e**(2*c + 2*d*x)
*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**2*b**3 + 2*e**(2*
c + 2*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*b**5 + 2*e
**(2*c + 2*d*x)*a**5*d*x + 6*e**(2*c + 2*d*x)*a**4*b*d*x + 4*e**(2*c + 2*d
*x)*a**3*b**2*d*x - 4*e**(2*c + 2*d*x)*a**2*b**3*d*x - 6*e**(2*c + 2*d*x)*
a*b**4*d*x - 2*e**(2*c + 2*d*x)*b**5*d*x - 3*log(e**(2*c + 2*d*x)*a + e**
(2*c + 2*d*x)*b + a - b)*a**4*b + 6*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*
x)*b + a - b)*a**3*b**2 - 4*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b +
a - b)*a**2*b**3 + 2*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*
a*b**4 - log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*b**5 + a...
```

### 3.64 $\int \frac{1}{(a+b \tanh(c+dx))^4} dx$

Optimal result . . . . .	548
Mathematica [A] (verified) . . . . .	549
Rubi [A] (verified) . . . . .	549
Maple [A] (verified) . . . . .	552
Fricas [B] (verification not implemented) . . . . .	553
Sympy [B] (verification not implemented) . . . . .	553
Maxima [B] (verification not implemented) . . . . .	554
Giac [A] (verification not implemented) . . . . .	555
Mupad [B] (verification not implemented) . . . . .	556
Reduce [B] (verification not implemented) . . . . .	556

#### Optimal result

Integrand size = 12, antiderivative size = 169

$$\int \frac{1}{(a+b \tanh(c+dx))^4} dx = \frac{(a^4 + 6a^2b^2 + b^4)x}{(a^2 - b^2)^4} - \frac{4ab(a^2 + b^2) \log(a \cosh(c+dx) + b \sinh(c+dx))}{(a^2 - b^2)^4 d} + \frac{b}{3(a^2 - b^2)d(a+b \tanh(c+dx))^3} + \frac{ab}{(a^2 - b^2)^2 d(a+b \tanh(c+dx))^2} + \frac{b(3a^2 + b^2)}{(a^2 - b^2)^3 d(a+b \tanh(c+dx))}$$

output

```
(a^4+6*a^2*b^2+b^4)*x/(a^2-b^2)^4-4*a*b*(a^2+b^2)*ln(a*cosh(d*x+c)+b*sinh(d*x+c))/(a^2-b^2)^4/d+1/3*b/(a^2-b^2)/d/(a+b*tanh(d*x+c))^3+a*b/(a^2-b^2)^2/d/(a+b*tanh(d*x+c))^2+b*(3*a^2+b^2)/(a^2-b^2)^3/d/(a+b*tanh(d*x+c))
```

### Mathematica [A] (verified)

Time = 2.22 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.95

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx$$

$$= \frac{-\frac{3 \log(1 - \tanh(c + dx))}{(a + b)^4} + \frac{3 \log(1 + \tanh(c + dx))}{(a - b)^4} + \frac{2b \left( -12a(a^2 + b^2) \log(a + b \tanh(c + dx)) + \frac{(a^2 - b^2)(13a^4 - 2a^2b^2 + b^4 + 3ab(7a^2 + b^2) \tanh(c + dx))}{(a + b \tanh(c + dx))^2} \right)}{(a^2 - b^2)^4}}{6d}$$

input

```
Integrate[(a + b*Tanh[c + d*x])^(-4), x]
```

output

```
((-3*Log[1 - Tanh[c + d*x]])/(a + b)^4 + (3*Log[1 + Tanh[c + d*x]])/(a - b)^4 + (2*b*(-12*a*(a^2 + b^2)*Log[a + b*Tanh[c + d*x]] + ((a^2 - b^2)*(13*a^4 - 2*a^2*b^2 + b^4 + 3*a*b*(7*a^2 + b^2)*Tanh[c + d*x] + 3*b^2*(3*a^2 + b^2)*Tanh[c + d*x]^2))/(a + b*Tanh[c + d*x]^3))/(a^2 - b^2)^4)/(6*d)
```

### Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.23, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3964, 3042, 4012, 3042, 4012, 3042, 4014, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(a - ib \tan(ic + idx))^4} dx$$

$$\downarrow \text{3964}$$

$$\frac{\int \frac{a - b \tanh(c + dx)}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2} + \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \frac{\int \frac{a + ib \tan(ic + idx)}{(a - ib \tan(ic + idx))^3} dx}{a^2 - b^2} \\
 & \quad \downarrow 4012 \\
 & \frac{\int \frac{a^2 - 2b \tanh(c + dx)a + b^2}{(a + b \tanh(c + dx))^2} dx}{a^2 - b^2} + \frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} \\
 & \quad \downarrow 3042 \\
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{\int \frac{a^2 + 2ib \tan(ic + idx)a + b^2}{(a - ib \tan(ic + idx))^2} dx}{a^2 - b^2} \\
 & \quad \downarrow 4012 \\
 & \frac{\int \frac{a(a^2 + 3b^2) - b(3a^2 + b^2) \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \\
 & \quad \frac{a^2 - b^2}{b} \\
 & \quad \frac{3d(a^2 - b^2)(a + b \tanh(c + dx))^3}{b} \\
 & \quad \downarrow 3042 \\
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \\
 & \frac{\frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{\int \frac{a(a^2 + 3b^2) + ib(3a^2 + b^2) \tan(ic + idx)}{a - ib \tan(ic + idx)} dx}{a^2 - b^2}}{a^2 - b^2} \\
 & \quad \downarrow 4014 \\
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \\
 & \frac{\frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{x(a^4 + 6a^2b^2 + b^4)}{a^2 - b^2} - \frac{4iab(a^2 + b^2) \int \frac{i(b + a \tanh(c + dx))}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} \\
 & \quad \downarrow 26 \\
 & \frac{\frac{x(a^4 + 6a^2b^2 + b^4)}{a^2 - b^2} - \frac{4iab(a^2 + b^2) \int \frac{b + a \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2}}{a^2 - b^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \\
 & \quad \frac{a^2 - b^2}{b} \\
 & \quad \frac{3d(a^2 - b^2)(a + b \tanh(c + dx))^3}{b}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \\
 & \frac{\frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{x(a^4 + 6a^2b^2 + b^4)}{a^2 - b^2} - \frac{4ab(a^2 + b^2) \int \frac{b - ia \tan(ic + idx)}{a - ib \tan(ic + idx)} dx}{a^2 - b^2}}{a^2 - b^2} \\
 & \downarrow 4013 \\
 & \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} + \\
 & \frac{\frac{ab}{d(a^2 - b^2)(a + b \tanh(c + dx))^2} + \frac{b(3a^2 + b^2)}{d(a^2 - b^2)(a + b \tanh(c + dx))} + \frac{x(a^4 + 6a^2b^2 + b^4)}{a^2 - b^2} - \frac{4ab(a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{a^2 - b^2}}{a^2 - b^2}
 \end{aligned}$$

input `Int[(a + b*Tanh[c + d*x])^(-4), x]`

output `b/(3*(a^2 - b^2)*d*(a + b*Tanh[c + d*x])^3) + ((a*b)/((a^2 - b^2)*d*(a + b*Tanh[c + d*x])^2) + (((a^4 + 6*a^2*b^2 + b^4)*x)/(a^2 - b^2) - (4*a*b*(a^2 + b^2)*Log[a*Cosh[c + d*x] + b*Sinh[c + d*x]])/((a^2 - b^2)*d))/(a^2 - b^2) + (b*(3*a^2 + b^2))/((a^2 - b^2)*d*(a + b*Tanh[c + d*x]))/(a^2 - b^2)))/(a^2 - b^2)`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



```
rule 3964 Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((a +
b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2)
Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]
```

```
rule 4012 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.96

method	result
derivativedivides	$\frac{b}{3(a-b)(a+b)(a+b \tanh(dx+c))^3} + \frac{ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))^2} + \frac{b(3a^2+b^2)}{(a+b)^3(a-b)^3(a+b \tanh(dx+c))} - \frac{4ba(a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^4(a-b)^4}$
default	$\frac{b}{3(a-b)(a+b)(a+b \tanh(dx+c))^3} + \frac{ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))^2} + \frac{b(3a^2+b^2)}{(a+b)^3(a-b)^3(a+b \tanh(dx+c))} - \frac{4ba(a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^4(a-b)^4}$
risch	$\frac{x}{a^4+4a^3b+6a^2b^2+4ab^3+b^4} + \frac{8ba^3x}{a^8-4a^6b^2+6a^4b^4-4a^2b^6+b^8} + \frac{8b^3ax}{a^8-4a^6b^2+6a^4b^4-4a^2b^6+b^8} + \frac{8ba^5}{d(a^8-4a^6b^2+6a^4b^4-4a^2b^6+b^8)}$
parallelrisc	$- \frac{-3a^9bdx-12a^8b^2dx-18a^7b^3dx-12a^6b^4dx-3a^5b^5dx+9a^6b^4-4a^4b^6+a^2b^8-6a^8b^2-3x \tanh(dx+c)^3a^6b^4d-12x \tanh(dx+c)^2a^6b^4d}{d(a^8-4a^6b^2+6a^4b^4-4a^2b^6+b^8)}$

input `int(1/(a+b*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(1/3*b/(a-b)/(a+b)/(a+b*tanh(d*x+c))^3+a*b/(a+b)^2/(a-b)^2/(a+b*tanh(d*x+c))^2+b*(3*a^2+b^2)/(a+b)^3/(a-b)^3/(a+b*tanh(d*x+c))-4*b*a*(a^2+b^2)/(a+b)^4/(a-b)^4*ln(a+b*tanh(d*x+c))+1/2/(a-b)^4*ln(tanh(d*x+c)+1)-1/2/(a+b)^4*ln(tanh(d*x+c)-1))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3693 vs.  $2(167) = 334$ .

Time = 0.15 (sec) , antiderivative size = 3693, normalized size of antiderivative = 21.85

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(a+b*tanh(d*x+c))^4,x, algorithm="fricas")`

output Too large to include

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16643 vs.  $2(144) = 288$ .

Time = 19.45 (sec) , antiderivative size = 16643, normalized size of antiderivative = 98.48

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(1/(a+b*tanh(d*x+c))**4,x)`

output

```
Piecewise((zoo*x/tanh(c)**4, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**4, Eq(
b, 0)), (3*d*x*tanh(c + d*x)**4/(48*b**4*d*tanh(c + d*x)**4 - 192*b**4*d*t
anh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c + d*x) +
48*b**4*d) - 12*d*x*tanh(c + d*x)**3/(48*b**4*d*tanh(c + d*x)**4 - 192*b*
**4*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c +
d*x) + 48*b**4*d) + 18*d*x*tanh(c + d*x)**2/(48*b**4*d*tanh(c + d*x)**4 -
192*b**4*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tan
h(c + d*x) + 48*b**4*d) - 12*d*x*tanh(c + d*x)/(48*b**4*d*tanh(c + d*x)**4
- 192*b**4*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*
tanh(c + d*x) + 48*b**4*d) + 3*d*x/(48*b**4*d*tanh(c + d*x)**4 - 192*b**4*
d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c + d*x
) + 48*b**4*d) - 3*tanh(c + d*x)**3/(48*b**4*d*tanh(c + d*x)**4 - 192*b**4
*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c + d*
x) + 48*b**4*d) + 12*tanh(c + d*x)**2/(48*b**4*d*tanh(c + d*x)**4 - 192*b*
**4*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c +
d*x) + 48*b**4*d) - 19*tanh(c + d*x)/(48*b**4*d*tanh(c + d*x)**4 - 192*b**
4*d*tanh(c + d*x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c + d
*x) + 48*b**4*d) + 16/(48*b**4*d*tanh(c + d*x)**4 - 192*b**4*d*tanh(c + d*
x)**3 + 288*b**4*d*tanh(c + d*x)**2 - 192*b**4*d*tanh(c + d*x) + 48*b**4*d
), Eq(a, -b)), (3*d*x*tanh(c + d*x)**4/(48*b**4*d*tanh(c + d*x)**4 + 19...
```

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 525 vs.  $2(167) = 334$ .

Time = 0.08 (sec) , antiderivative size = 525, normalized size of antiderivative = 3.11

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = -\frac{4(a^3b + ab^3) \log(-(a-b)e^{(-2dx-2c)} - a - b)}{(a^8 - 4a^6b^2 + 6a^4b^4 - 4a^2b^6 + b^8)d}$$

$$-\frac{3(a^{10} + 2a^9b - 3a^8b^2 - 8a^7b^3 + 2a^6b^4 + 12a^5b^5 + 2a^4b^6 - 8a^3b^7 - 3a^2b^8 + 2ab^9 + b^{10} + 3(a^{10} - 5$$

$$+ \frac{dx + c}{(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)d}$$

input

```
integrate(1/(a+b*tanh(d*x+c))^4,x, algorithm="maxima")
```

output

```

-4*(a^3*b + a*b^3)*log(-(a - b)*e^(-2*d*x - 2*c) - a - b)/((a^8 - 4*a^6*b^
2 + 6*a^4*b^4 - 4*a^2*b^6 + b^8)*d) - 4/3*(9*a^4*b^2 + 18*a^3*b^3 + 11*a^2
*b^4 + 4*a*b^5 + 2*b^6 + 3*(6*a^4*b^2 + 2*a^3*b^3 - 5*a^2*b^4 - 2*a*b^5 -
b^6)*e^(-2*d*x - 2*c) + 3*(3*a^4*b^2 - 4*a^3*b^3 + b^6)*e^(-4*d*x - 4*c))/
((a^10 + 2*a^9*b - 3*a^8*b^2 - 8*a^7*b^3 + 2*a^6*b^4 + 12*a^5*b^5 + 2*a^4*
b^6 - 8*a^3*b^7 - 3*a^2*b^8 + 2*a*b^9 + b^10 + 3*(a^10 - 5*a^8*b^2 + 10*a^
6*b^4 - 10*a^4*b^6 + 5*a^2*b^8 - b^10)*e^(-2*d*x - 2*c) + 3*(a^10 - 2*a^9*
b - 3*a^8*b^2 + 8*a^7*b^3 + 2*a^6*b^4 - 12*a^5*b^5 + 2*a^4*b^6 + 8*a^3*b^7
- 3*a^2*b^8 - 2*a*b^9 + b^10)*e^(-4*d*x - 4*c) + (a^10 - 4*a^9*b + 3*a^8*
b^2 + 8*a^7*b^3 - 14*a^6*b^4 + 14*a^4*b^6 - 8*a^3*b^7 - 3*a^2*b^8 + 4*a*b^
9 - b^10)*e^(-6*d*x - 6*c))*d) + (d*x + c)/((a^4 + 4*a^3*b + 6*a^2*b^2 + 4
*a*b^3 + b^4)*d)

```

**Giac [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.80

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = \frac{12 \frac{(a^3 b + ab^3) \log(|-ae^{(2dx+2c)} - be^{(2dx+2c)} - a + b|)}{a^8 - 4a^6 b^2 + 6a^4 b^4 - 4a^2 b^6 + b^8}}{a^4 - 4a^3 b + 6a^2 b^2 - 4ab^3 + b^4} - \frac{3(dx+c)}{a^4 - 4a^3 b + 6a^2 b^2 - 4ab^3 + b^4} - \frac{4 \left( 3a^4 b^2 - 2a^3 b^3 - 2a^2 b^4 + 2ab^5 - b^6 \right) e^{(4dx+4c)}}{a^4 - 4a^3 b + 6a^2 b^2 - 4ab^3 + b^4}$$

3d

input

```
integrate(1/(a+b*tanh(d*x+c))^4,x, algorithm="giac")
```

output

```

-1/3*(12*(a^3*b + a*b^3)*log(abs(-a*e^(2*d*x + 2*c) - b*e^(2*d*x + 2*c) -
a + b))/(a^8 - 4*a^6*b^2 + 6*a^4*b^4 - 4*a^2*b^6 + b^8) - 3*(d*x + c)/(a^4
- 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4) - 4*(3*(3*a^4*b^2 - 2*a^3*b^3 - 2*
a^2*b^4 + 2*a*b^5 - b^6)*e^(4*d*x + 4*c) + 3*(6*a^4*b^2 - 14*a^3*b^3 + 11*
a^2*b^4 - 4*a*b^5 + b^6)*e^(2*d*x + 2*c) + (9*a^5*b^2 - 27*a^4*b^3 + 29*a^
3*b^4 - 15*a^2*b^5 + 6*a*b^6 - 2*b^7)/(a + b))/((a*e^(2*d*x + 2*c) + b*e^(
2*d*x + 2*c) + a - b)^3*(a + b)^3*(a - b)^4)/d

```

**Mupad [B] (verification not implemented)**

Time = 3.98 (sec) , antiderivative size = 452, normalized size of antiderivative = 2.67

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = \frac{\ln(\tanh(c + dx) + 1)}{2da^4 - 8da^3b + 12da^2b^2 - 8dab^3 + 2db^4}$$

$$- \frac{\frac{\tanh(c+dx)(6a^4b^2-3a^2b^4+b^6)}{ad(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{\tanh(c+dx)^2(10a^4b^3-3a^2b^5+b^7)}{a^2d(a^3-3a^2b+3ab^2-b^3)(a^3+3a^2b+3ab^2+b^3)} + \frac{\tanh(c+dx)^3\left(\frac{13a^4b^4}{3} - \frac{2a^2b^6}{3} + \frac{b^8}{3}\right)}{a^3d(a^3-3a^2b+3ab^2-b^3)(a^3+3a^2b+3ab^2+b^3)}}{a^3 + 3a^2b \tanh(c + dx) + 3ab^2 \tanh(c + dx)^2 + b^3 \tanh(c + dx)^3}$$

$$- \frac{\ln(1 - \tanh(c + dx))}{2da^4 + 8da^3b + 12da^2b^2 + 8dab^3 + 2db^4}$$

$$- \frac{4 \ln(a + b \tanh(c + dx)) (a^3b + ab^3)}{d(a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4)(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)}$$

input `int(1/(a + b*tanh(c + d*x))^4,x)`

output

```
log(tanh(c + d*x) + 1)/(2*a^4*d + 2*b^4*d + 12*a^2*b^2*d - 8*a*b^3*d - 8*a^3*b*d) - ((tanh(c + d*x)*(b^6 - 3*a^2*b^4 + 6*a^4*b^2))/(a*d*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)) + (tanh(c + d*x)^2*(b^7 - 3*a^2*b^5 + 10*a^4*b^3))/(a^2*d*(3*a*b^2 - 3*a^2*b + a^3 - b^3)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)) + (tanh(c + d*x)^3*(b^8/3 - (2*a^2*b^6)/3 + (13*a^4*b^4)/3))/(a^3*d*(3*a*b^2 - 3*a^2*b + a^3 - b^3)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)))/(a^3 + b^3*tanh(c + d*x)^3 + 3*a*b^2*tanh(c + d*x)^2 + 3*a^2*b*tanh(c + d*x)) - log(1 - tanh(c + d*x))/(2*a^4*d + 2*b^4*d + 12*a^2*b^2*d + 8*a*b^3*d + 8*a^3*b*d) - (4*log(a + b*tanh(c + d*x))*(a*b^3 + a^3*b))/(d*(a^4 - 4*a^3*b - 4*a*b^3 + b^4 + 6*a^2*b^2)*(4*a*b^3 + 4*a^3*b + a^4 + b^4 + 6*a^2*b^2))
```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 2312, normalized size of antiderivative = 13.68

$$\int \frac{1}{(a + b \tanh(c + dx))^4} dx = \text{Too large to display}$$

input `int(1/(a+b*tanh(d*x+c))^4,x)`

output

```
( - 12***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a -
b)*a**6*b - 36***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*
b + a - b)*a**5*b**2 - 48***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*
c + 2*d*x)*b + a - b)*a**4*b**3 - 48***e**(6*c + 6*d*x)*log(e**(2*c + 2*d*x)
*a + e**(2*c + 2*d*x)*b + a - b)*a**3*b**4 - 36***e**(6*c + 6*d*x)*log(e**(2
*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**2*b**5 - 12***e**(6*c + 6*d*x
)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a*b**6 + 3***e**(6*c
+ 6*d*x)*a**7*d*x + 21***e**(6*c + 6*d*x)*a**6*b*d*x + 63***e**(6*c + 6*d*x)*a
**5*b**2*d*x - 12***e**(6*c + 6*d*x)*a**5*b**2 + 105***e**(6*c + 6*d*x)*a**4*b
**3*d*x - 28***e**(6*c + 6*d*x)*a**4*b**3 + 105***e**(6*c + 6*d*x)*a**3*b**4*d
*x - 16***e**(6*c + 6*d*x)*a**3*b**4 + 63***e**(6*c + 6*d*x)*a**2*b**5*d*x + 2
1***e**(6*c + 6*d*x)*a*b**6*d*x - 4***e**(6*c + 6*d*x)*a*b**6 + 3***e**(6*c + 6
*d*x)*b**7*d*x - 4***e**(6*c + 6*d*x)*b**7 - 36***e**(4*c + 4*d*x)*log(e**(2*c
+ 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**6*b - 36***e**(4*c + 4*d*x)*log(
e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**5*b**2 + 36***e**(4*c +
4*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a**2*b**5 + 36
***e**(4*c + 4*d*x)*log(e**(2*c + 2*d*x)*a + e**(2*c + 2*d*x)*b + a - b)*a*b
**6 + 9***e**(4*c + 4*d*x)*a**7*d*x + 45***e**(4*c + 4*d*x)*a**6*b*d*x + 81***e
*(4*c + 4*d*x)*a**5*b**2*d*x + 45***e**(4*c + 4*d*x)*a**4*b**3*d*x - 45***e**
(4*c + 4*d*x)*a**3*b**4*d*x - 81***e**(4*c + 4*d*x)*a**2*b**5*d*x - 45***e**...
```

### 3.65 $\int \frac{1}{4+6 \tanh(c+dx)} dx$

Optimal result . . . . .	558
Mathematica [A] (verified) . . . . .	558
Rubi [A] (verified) . . . . .	559
Maple [A] (verified) . . . . .	560
Fricas [A] (verification not implemented) . . . . .	561
Sympy [A] (verification not implemented) . . . . .	561
Maxima [A] (verification not implemented) . . . . .	561
Giac [A] (verification not implemented) . . . . .	562
Mupad [B] (verification not implemented) . . . . .	562
Reduce [B] (verification not implemented) . . . . .	562

#### Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{4+6 \tanh(c+dx)} dx = -\frac{x}{5} + \frac{3 \log(2 \cosh(c+dx) + 3 \sinh(c+dx))}{10d}$$

output `-1/5*x+3/10*ln(2*cosh(d*x+c)+3*sinh(d*x+c))/d`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

$$\int \frac{1}{4+6 \tanh(c+dx)} dx = -\frac{\log(1-\tanh(c+dx))}{20d} - \frac{\log(1+\tanh(c+dx))}{4d} + \frac{3 \log(2+3 \tanh(c+dx))}{10d}$$

input `Integrate[(4 + 6*Tanh[c + d*x])^(-1),x]`

output `-1/20*Log[1 - Tanh[c + d*x]]/d - Log[1 + Tanh[c + d*x]]/(4*d) + (3*Log[2 + 3*Tanh[c + d*x]])/(10*d)`

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{6 \tanh(c + dx) + 4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{4 - 6i \tan(ic + idx)} dx \\
 & \quad \downarrow \text{3965} \\
 & -\frac{x}{5} + \frac{3}{10} i \int -\frac{i(2 \tanh(c + dx) + 3)}{3 \tanh(c + dx) + 2} dx \\
 & \quad \downarrow \text{26} \\
 & \frac{3}{10} \int \frac{2 \tanh(c + dx) + 3}{3 \tanh(c + dx) + 2} dx - \frac{x}{5} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{x}{5} + \frac{3}{10} \int \frac{3 - 2i \tan(ic + idx)}{2 - 3i \tan(ic + idx)} dx \\
 & \quad \downarrow \text{4013} \\
 & \frac{3 \log(3 \sinh(c + dx) + 2 \cosh(c + dx))}{10d} - \frac{x}{5}
 \end{aligned}$$

input `Int[(4 + 6*Tanh[c + d*x])^(-1),x]`

output `-1/5*x + (3*Log[2*Cosh[c + d*x] + 3*Sinh[c + d*x]])/(10*d)`



## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_{x_}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u_ , x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u_ , x], x] /; \text{FunctionOfTrigOfLinearQ}[u_ , x]$
- rule 3965  $\text{Int}[((a_ ) + (b_ )*\tan[(c_ ) + (d_ )*(x_)])^{-1}, x\_Symbol] \rightarrow \text{Simp}[a*(x/(a^2 + b^2)), x] + \text{Simp}[b/(a^2 + b^2) \text{Int}[(b - a*\tan[c + d*x])/(a + b*\tan[c + d*x]), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4013  $\text{Int}[((c_ ) + (d_ )*\tan[(e_ ) + (f_ )*(x_)])/((a_ ) + (b_ )*\tan[(e_ ) + (f_ )*(x_)]), x\_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
risch	$-\frac{x}{2} - \frac{3c}{5d} + \frac{3 \ln(e^{2dx+2c} - \frac{1}{5})}{10d}$	28
parallelrisc	$-\frac{5dx+3 \ln(1-\tanh(dx+c))-3 \ln(\frac{2}{3}+\tanh(dx+c))}{10d}$	35
derivativedivides	$\frac{\frac{3 \ln(2+3 \tanh(dx+c))}{5} - \frac{\ln(\tanh(dx+c)+1)}{2} - \frac{\ln(\tanh(dx+c)-1)}{10}}{2d}$	42
default	$\frac{\frac{3 \ln(2+3 \tanh(dx+c))}{5} - \frac{\ln(\tanh(dx+c)+1)}{2} - \frac{\ln(\tanh(dx+c)-1)}{10}}{2d}$	42

input `int(1/(4+6*tanh(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/2*x-3/5*c/d+3/10/d*ln(exp(2*d*x+2*c)-1/5)`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = -\frac{5 dx - 3 \log\left(\frac{2(2 \cosh(dx+c)+3 \sinh(dx+c))}{\cosh(dx+c)-\sinh(dx+c)}\right)}{10 d}$$

input `integrate(1/(4+6*tanh(d*x+c)),x, algorithm="fricas")`output `-1/10*(5*d*x - 3*log(2*(2*cosh(d*x + c) + 3*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))))/d`**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = \begin{cases} \frac{x}{10} - \frac{3 \log(\tanh(c+dx)+1)}{10d} + \frac{3 \log(3 \tanh(c+dx)+2)}{10d} & \text{for } d \neq 0 \\ \frac{x}{6 \tanh(c)+4} & \text{otherwise} \end{cases}$$

input `integrate(1/(4+6*tanh(d*x+c)),x)`output `Piecewise((x/10 - 3*log(tanh(c + d*x) + 1)/(10*d) + 3*log(3*tanh(c + d*x) + 2)/(10*d), Ne(d, 0)), (x/(6*tanh(c) + 4), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = \frac{dx + c}{10 d} + \frac{3 \log(e^{(-2 dx - 2c)} - 5)}{10 d}$$

input `integrate(1/(4+6*tanh(d*x+c)),x, algorithm="maxima")`output `1/10*(d*x + c)/d + 3/10*log(e^(-2*d*x - 2*c) - 5)/d`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = -\frac{5 dx + 5 c - 3 \log(|5 e^{(2 dx + 2 c)} - 1|)}{10 d}$$

input `integrate(1/(4+6*tanh(d*x+c)),x, algorithm="giac")`output `-1/10*(5*d*x + 5*c - 3*log(abs(5*e^(2*d*x + 2*c) - 1)))/d`**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = \frac{x}{10} - \frac{\frac{3 \ln(\tanh(c+dx)+1)}{10} - \frac{3 \ln(3 \tanh(c+dx)+2)}{10}}{d}$$

input `int(1/(6*tanh(c + d*x) + 4),x)`output `x/10 - ((3*log(tanh(c + d*x) + 1))/10 - (3*log(3*tanh(c + d*x) + 2))/10)/d`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{1}{4 + 6 \tanh(c + dx)} dx = \frac{3 \log(5 e^{dx+c} - \sqrt{5}) + 3 \log(5 e^{dx+c} + \sqrt{5}) - 5 dx}{10 d}$$

input `int(1/(4+6*tanh(d*x+c)),x)`output `(3*log(5*e**(c + d*x) - sqrt(5)) + 3*log(5*e**(c + d*x) + sqrt(5)) - 5*d*x)/(10*d)`

### 3.66 $\int \frac{1}{4-6 \tanh(c+dx)} dx$

Optimal result . . . . .	563
Mathematica [A] (verified) . . . . .	563
Rubi [A] (verified) . . . . .	564
Maple [A] (verified) . . . . .	565
Fricas [A] (verification not implemented) . . . . .	566
Sympy [A] (verification not implemented) . . . . .	566
Maxima [A] (verification not implemented) . . . . .	566
Giac [A] (verification not implemented) . . . . .	567
Mupad [B] (verification not implemented) . . . . .	567
Reduce [B] (verification not implemented) . . . . .	567

#### Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{4-6 \tanh(c+dx)} dx = -\frac{x}{5} - \frac{3 \log(2 \cosh(c+dx) - 3 \sinh(c+dx))}{10d}$$

output

```
-1/5*x-3/10*ln(2*cosh(d*x+c)-3*sinh(d*x+c))/d
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

$$\int \frac{1}{4-6 \tanh(c+dx)} dx = -\frac{3 \log(2-3 \tanh(c+dx))}{10d} + \frac{\log(1-\tanh(c+dx))}{4d} + \frac{\log(1+\tanh(c+dx))}{20d}$$

input

```
Integrate[(4 - 6*Tanh[c + d*x])^(-1),x]
```

output

```
(-3*Log[2 - 3*Tanh[c + d*x]])/(10*d) + Log[1 - Tanh[c + d*x]]/(4*d) + Log[1 + Tanh[c + d*x]]/(20*d)
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{4 - 6 \tanh(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{4 + 6i \tan(ic + idx)} dx \\
 & \quad \downarrow \text{3965} \\
 & -\frac{x}{5} - \frac{3}{10} i \int \frac{i(3 - 2 \tanh(c + dx))}{2 - 3 \tanh(c + dx)} dx \\
 & \quad \downarrow \text{26} \\
 & \frac{3}{10} \int \frac{3 - 2 \tanh(c + dx)}{2 - 3 \tanh(c + dx)} dx - \frac{x}{5} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{x}{5} + \frac{3}{10} \int \frac{2i \tan(ic + idx) + 3}{3i \tan(ic + idx) + 2} dx \\
 & \quad \downarrow \text{4013} \\
 & -\frac{3 \log(2 \cosh(c + dx) - 3 \sinh(c + dx))}{10d} - \frac{x}{5}
 \end{aligned}$$

input `Int[(4 - 6*Tanh[c + d*x])^(-1),x]`

output `-1/5*x - (3*Log[2*Cosh[c + d*x] - 3*Sinh[c + d*x]])/(10*d)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3965  $\text{Int}[(a + (b \cdot \tan[c + (d \cdot x)])^{-1}), x\_Symbol] \rightarrow \text{Simp}[a \cdot (x / (a^2 + b^2)), x] + \text{Simp}[b / (a^2 + b^2) \text{Int}[(b - a \cdot \tan[c + d \cdot x]) / (a + b \cdot \tan[c + d \cdot x]), x], x] /;$   $\text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4013  $\text{Int}[(c + (d \cdot \tan[e + (f \cdot x)]) / ((a + (b \cdot \tan[e + (f \cdot x)])) \cdot \sin[e + f \cdot x]), x\_Symbol] \rightarrow \text{Simp}[(c / (b \cdot f)) \cdot \text{Log}[\text{RemoveContent}[a \cdot \cos[e + f \cdot x] + b \cdot \sin[e + f \cdot x], x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a \cdot c + b \cdot d, 0]$

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
risch	$\frac{x}{10} + \frac{3c}{5d} - \frac{3 \ln(e^{2dx+2c}-5)}{10d}$	28
parallelrisch	$-\frac{-dx-3 \ln(1-\tanh(dx+c))+3 \ln(-\frac{2}{3}+\tanh(dx+c))}{10d}$	35
derivativedivides	$\frac{\frac{\ln(\tanh(dx+c)-1)}{2} + \frac{\ln(\tanh(dx+c)+1)}{10} - \frac{3 \ln(-2+3 \tanh(dx+c))}{5}}{2d}$	42
default	$\frac{\frac{\ln(\tanh(dx+c)-1)}{2} + \frac{\ln(\tanh(dx+c)+1)}{10} - \frac{3 \ln(-2+3 \tanh(dx+c))}{5}}{2d}$	42

input `int(1/(4-6*tanh(d*x+c)),x,method=_RETURNVERBOSE)`output `1/10*x+3/5*c/d-3/10/d*ln(exp(2*d*x+2*c)-5)`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.55

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = \frac{dx - 3 \log \left( \frac{-2(2 \cosh(dx+c) - 3 \sinh(dx+c))}{\cosh(dx+c) - \sinh(dx+c)} \right)}{10d}$$

input `integrate(1/(4-6*tanh(d*x+c)),x, algorithm="fricas")`output `1/10*(d*x - 3*log(-2*(2*cosh(d*x + c) - 3*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))))/d`**Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = \begin{cases} -\frac{x}{2} + \frac{3 \log(\tanh(c+dx)+1)}{10d} - \frac{3 \log(3 \tanh(c+dx)-2)}{10d} & \text{for } d \neq 0 \\ \frac{x}{4-6 \tanh(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/(4-6*tanh(d*x+c)),x)`output `Piecewise((-x/2 + 3*log(tanh(c + d*x) + 1)/(10*d) - 3*log(3*tanh(c + d*x) - 2)/(10*d), Ne(d, 0)), (x/(4 - 6*tanh(c)), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = -\frac{1}{2}x - \frac{c}{2d} - \frac{3 \log(5e^{(-2dx-2c)} - 1)}{10d}$$

input `integrate(1/(4-6*tanh(d*x+c)),x, algorithm="maxima")`output `-1/2*x - 1/2*c/d - 3/10*log(5*e^(-2*d*x - 2*c) - 1)/d`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.81

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = \frac{dx + c - 3 \log(|e^{(2dx+2c)} - 5|)}{10d}$$

input `integrate(1/(4-6*tanh(d*x+c)),x, algorithm="giac")`output `1/10*(d*x + c - 3*log(abs(e^(2*d*x + 2*c) - 5)))/d`**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = \frac{\frac{3 \ln(\tanh(c+dx)+1)}{10} - \frac{3 \ln(3 \tanh(c+dx)-2)}{10}}{d} - \frac{x}{2}$$

input `int(-1/(6*tanh(c + d*x) - 4),x)`output `((3*log(tanh(c + d*x) + 1))/10 - (3*log(3*tanh(c + d*x) - 2))/10)/d - x/2`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \frac{1}{4 - 6 \tanh(c + dx)} dx = \frac{-3 \log(e^{dx+c} - \sqrt{5}) - 3 \log(e^{dx+c} + \sqrt{5}) + dx}{10d}$$

input `int(1/(4-6*tanh(d*x+c)),x)`output `( - 3*log(e**(c + d*x) - sqrt(5)) - 3*log(e**(c + d*x) + sqrt(5)) + d*x)/(10*d)`



### 3.67 $\int \sqrt{a + b \tanh(c + dx)} dx$

Optimal result	568
Mathematica [A] (verified)	568
Rubi [A] (verified)	569
Maple [A] (verified)	571
Fricas [B] (verification not implemented)	571
Sympy [F]	572
Maxima [F(-2)]	573
Giac [F(-2)]	573
Mupad [B] (verification not implemented)	574
Reduce [F]	574

#### Optimal result

Integrand size = 14, antiderivative size = 74

$$\int \sqrt{a + b \tanh(c + dx)} dx = -\frac{\sqrt{a - b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{d} + \frac{\sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{d}$$

output

$-(a-b)^{(1/2)}*\operatorname{arctanh}((a+b*\tanh(d*x+c))^{(1/2)/(a-b)^{(1/2)})/d+(a+b)^{(1/2)}*\operatorname{arctanh}((a+b*\tanh(d*x+c))^{(1/2)/(a+b)^{(1/2)})/d$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \tanh(c + dx)} dx = -\frac{\sqrt{a - b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{d} + \frac{\sqrt{a + b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{d}$$

input

`Integrate[Sqrt[a + b*Tanh[c + d*x]],x]`

output

$$-\left(\frac{\sqrt{a-b} \operatorname{ArcTanh}\left[\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right]}{d}\right) + \left(\frac{\sqrt{a+b} \operatorname{ArcTanh}\left[\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right]}{d}\right)$$
**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.22, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3966, 25, 483, 25, 1450, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a + b \tanh(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{a - ib \tan(ic + idx)} dx \\ & \quad \downarrow \text{3966} \\ & \frac{b \int -\frac{\sqrt{a+b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\ & \quad \downarrow \text{25} \\ & \frac{b \int \frac{\sqrt{a+b \tanh(c+dx)}}{b^2 - b^2 \tanh^2(c+dx)} d(b \tanh(c + dx))}{d} \\ & \quad \downarrow \text{483} \\ & \frac{2b \int -\frac{b^2 \tanh^2(c+dx)}{b^4 \tanh^4(c+dx) - 2ab^2 \tanh^2(c+dx) + a^2 - b^2} d\sqrt{a + b \tanh(c + dx)}}{d} \\ & \quad \downarrow \text{25} \\ & \frac{2b \int \frac{b^2 \tanh^2(c+dx)}{b^4 \tanh^4(c+dx) - 2ab^2 \tanh^2(c+dx) + a^2 - b^2} d\sqrt{a + b \tanh(c + dx)}}{d} \\ & \quad \downarrow \text{1450} \end{aligned}$$

$$\frac{2b \left( -\frac{(a+b) \int \frac{1}{b^2 \tanh^2(c+dx) - a - b} d\sqrt{a+b \tanh(c+dx)}}{2b} - \frac{1}{2} \left(1 - \frac{a}{b}\right) \int \frac{1}{b^2 \tanh^2(c+dx) - a + b} d\sqrt{a + b \tanh(c + dx)} \right)}{d}$$

↓ 220

$$\frac{2b \left( \frac{\left(1 - \frac{a}{b}\right) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{2\sqrt{a-b}} + \frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{2b} \right)}{d}$$

input `Int[Sqrt[a + b*Tanh[c + d*x]],x]`

output `(2*b*((1 - a/b)*ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a - b]]/(2*Sqrt[a - b]) + (Sqrt[a + b]*ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a + b]])/(2*b)))/d`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 483 `Int[Sqrt[(c_) + (d_.)*(x_)]/((a_) + (b_.)*(x_)^2), x_Symbol] :> Simp[2*d Subst[Int[x^2/(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x]`

rule 1450 `Int[((d_.)*(x_))^(m_)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(d^2/2)*(b/q + 1) Int[(d*x)^(m - 2)/(b/2 + q/2 + c*x^2), x], x] - Simp[(d^2/2)*(b/q - 1) Int[(d*x)^(m - 2)/(b/2 - q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - 4*a*c, 0] && GeQ[m, 2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3966 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 + b^2, 0]`

### Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$\frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \tanh(dx+c)}{\sqrt{a+b}}\right)}{d} - \frac{\sqrt{-a+b} \operatorname{arctan}\left(\frac{\sqrt{a+b} \tanh(dx+c)}{\sqrt{-a+b}}\right)}{d}$	63
default	$\frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b} \tanh(dx+c)}{\sqrt{a+b}}\right)}{d} - \frac{\sqrt{-a+b} \operatorname{arctan}\left(\frac{\sqrt{a+b} \tanh(dx+c)}{\sqrt{-a+b}}\right)}{d}$	63

input `int((a+b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `(a+b)^(1/2)*arctanh((a+b*tanh(d*x+c))^(1/2)/(a+b)^(1/2))/d-1/d*(-a+b)^(1/2)*arctan((a+b*tanh(d*x+c))^(1/2)/(-a+b)^(1/2))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs.  $2(62) = 124$ .

Time = 0.15 (sec) , antiderivative size = 2203, normalized size of antiderivative = 29.77

$$\int \sqrt{a + b \tanh(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="fricas")`

output

```
[1/4*(sqrt(a + b)*log(2*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 8*(a^2 + 2*a
*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + 2*(a^2 + 2*a*b + b^2)*sinh(d*x +
c)^4 + 4*(a^2 + a*b)*cosh(d*x + c)^2 + 4*(3*(a^2 + 2*a*b + b^2)*cosh(d*x
+ c)^2 + a^2 + a*b)*sinh(d*x + c)^2 + 2*a^2 - b^2 + 2*((a + b)*cosh(d*x +
c)^4 + 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 +
(2*a + b)*cosh(d*x + c)^2 + (6*(a + b)*cosh(d*x + c)^2 + 2*a + b)*sinh(d*
x + c)^2 + 2*(2*(a + b)*cosh(d*x + c)^3 + (2*a + b)*cosh(d*x + c))*sinh(d*
x + c) + a)*sqrt(a + b)*sqrt((a*cosh(d*x + c) + b*sinh(d*x + c))/cosh(d*x
+ c)) + 8*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 + (a^2 + a*b)*cosh(d*x + c)
)*sinh(d*x + c)) + sqrt(a - b)*log(((2*a^2 - b^2)*cosh(d*x + c)^4 + 4*(2*a
^2 - b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (2*a^2 - b^2)*sinh(d*x + c)^4 +
4*(a^2 - a*b)*cosh(d*x + c)^2 + 2*(3*(2*a^2 - b^2)*cosh(d*x + c)^2 + 2*a^2
- 2*a*b)*sinh(d*x + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - 2*(a*cosh(d*x + c)^4 +
4*a*cosh(d*x + c)*sinh(d*x + c)^3 + a*sinh(d*x + c)^4 + (2*a - b)*cosh(d*
x + c)^2 + (6*a*cosh(d*x + c)^2 + 2*a - b)*sinh(d*x + c)^2 + 2*(2*a*cosh(d
*x + c)^3 + (2*a - b)*cosh(d*x + c))*sinh(d*x + c) + a - b)*sqrt(a - b)*sq
rt((a*cosh(d*x + c) + b*sinh(d*x + c))/cosh(d*x + c)) + 4*((2*a^2 - b^2)*c
osh(d*x + c)^3 + 2*(a^2 - a*b)*cosh(d*x + c))*sinh(d*x + c))/(cosh(d*x + c
)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2
+ 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)))/d, -1/4*(2*sqrt(...
```

### Sympy [F]

$$\int \sqrt{a + b \tanh(c + dx)} dx = \int \sqrt{a + b \tanh(c + dx)} dx$$

input

```
integrate((a+b*tanh(d*x+c))**(1/2), x)
```

output

```
Integral(sqrt(a + b*tanh(c + d*x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{a + b \tanh(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see `assume?` for more detail`

**Giac [F(-2)]**

Exception generated.

$$\int \sqrt{a + b \tanh(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Degree mismatch inside factorisation over extensionDegree mismatch inside factorisation over extensionindex.c c index_m`

**Mupad [B] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.04

$$\int \sqrt{a + b \tanh(c + dx)} dx$$

$$= \frac{\operatorname{atan}\left(\frac{b^2 \sqrt{a+b} \sqrt{a+b \tanh(c+dx)} \operatorname{li} - a b \sqrt{a+b} \sqrt{a+b \tanh(c+dx)} \operatorname{li}}{a^2 b - b^3}\right) \sqrt{a+b} \operatorname{li}}{d} + \frac{\operatorname{atan}\left(\frac{b^2 \sqrt{a-b} \sqrt{a+b \tanh(c+dx)} \operatorname{li} + a b \sqrt{a-b} \sqrt{a+b \tanh(c+dx)} \operatorname{li}}{a^2 b - b^3}\right) \sqrt{a-b} \operatorname{li}}{d}$$

input `int((a + b*tanh(c + d*x))^(1/2),x)`output `(atan((b^2*(a + b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i - a*b*(a + b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i)/(a^2*b - b^3))*(a + b)^(1/2)*1i)/d + (atan((b^2*(a - b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i + a*b*(a - b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i)/(a^2*b - b^3))*(a - b)^(1/2)*1i)/d`**Reduce [F]**

$$\int \sqrt{a + b \tanh(c + dx)} dx = \int \sqrt{\tanh(dx + c) b + a} dx$$

input `int((a+b*tanh(d*x+c))^(1/2),x)`output `int(sqrt(tanh(c + d*x)*b + a),x)`

### 3.68 $\int \frac{1}{\sqrt{a+b \tanh(c+dx)}} dx$

Optimal result	575
Mathematica [A] (verified)	575
Rubi [A] (verified)	576
Maple [A] (verified)	578
Fricas [B] (verification not implemented)	578
Sympy [F]	579
Maxima [F(-2)]	580
Giac [F(-2)]	580
Mupad [B] (verification not implemented)	581
Reduce [F]	581

#### Optimal result

Integrand size = 14, antiderivative size = 74

$$\int \frac{1}{\sqrt{a+b \tanh(c+dx)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{\sqrt{a-bd}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{\sqrt{a+bd}}$$

output

$-\operatorname{arctanh}\left(\frac{(a+b*\tanh(d*x+c))^{1/2}}{(a-b)^{1/2}}\right)/(a-b)^{1/2}/d+\operatorname{arctanh}\left(\frac{(a+b*\tanh(d*x+c))^{1/2}}{(a+b)^{1/2}}\right)/(a+b)^{1/2}/d$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a+b \tanh(c+dx)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{\sqrt{a-bd}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{\sqrt{a+bd}}$$

input

`Integrate[1/Sqrt[a + b*Tanh[c + d*x]],x]`

output

$-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[a - b]]/(\operatorname{Sqrt}[a - b]*d)) + \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[a + b]]/(\operatorname{Sqrt}[a + b]*d)$



**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {3042, 3966, 25, 484, 1406, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a - ib \tan(ic + idx)}} dx \\
 & \quad \downarrow \text{3966} \\
 & \frac{b \int -\frac{1}{\sqrt{a+b \tanh(c+dx)}(b^2-b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & \frac{b \int \frac{1}{\sqrt{a+b \tanh(c+dx)}(b^2-b^2 \tanh^2(c+dx))} d(b \tanh(c + dx))}{d} \\
 & \quad \downarrow \text{484} \\
 & \frac{2b \int \frac{1}{-b^4 \tanh^4(c+dx)+2ab^2 \tanh^2(c+dx)-a^2+b^2} d\sqrt{a + b \tanh(c + dx)}}{d} \\
 & \quad \downarrow \text{1406} \\
 & \frac{2b \left( \frac{\int \frac{1}{-b^2 \tanh^2(c+dx)+a+b} d\sqrt{a+b \tanh(c+dx)}}{2b} - \frac{\int \frac{1}{-b^2 \tanh^2(c+dx)+a-b} d\sqrt{a+b \tanh(c+dx)}}{2b} \right)}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{2b \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{2b\sqrt{a+b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{2b\sqrt{a-b}} \right)}{d}
 \end{aligned}$$

input

```
Int[1/Sqrt[a + b*Tanh[c + d*x]], x]
```

output  $(2*b*(-1/2*ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a - b]]/(Sqrt[a - b]*b) + ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a + b]]/(2*b*Sqrt[a + b]))/d$

### Defintions of rubi rules used

rule 25  $Int[-(Fx_), x\_Symbol] \rightarrow Simp[Identity[-1] Int[Fx, x], x]$

rule 219  $Int[((a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[\{a, b\}, x] \&\& NegQ[a/b] \&\& (GtQ[a, 0] || LtQ[b, 0])$

rule 484  $Int[1/(Sqrt[(c_) + (d_)*(x_)]*((a_) + (b_)*(x_)^2)), x\_Symbol] \rightarrow Simp[2*d Subst[Int[1/(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4), x], x, Sqrt[c + d*x]], x] /; FreeQ[\{a, b, c, d\}, x]$

rule 1406  $Int[((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^{-1}, x\_Symbol] \rightarrow With[\{q = Rt[b^2 - 4*a*c, 2]\}, Simp[c/q Int[1/(b/2 - q/2 + c*x^2), x], x] - Simp[c/q Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[\{a, b, c\}, x] \&\& NeQ[b^2 - 4*a*c, 0] \&\& PosQ[b^2 - 4*a*c]$

rule 3042  $Int[u_, x\_Symbol] \rightarrow Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]$

rule 3966  $Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^{(n_)}, x\_Symbol] \rightarrow Simp[b/d Subst[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[\{a, b, c, d, n\}, x] \&\& NeQ[a^2 + b^2, 0]$

**Maple [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b}\tanh(dx+c)}{\sqrt{a+b}}\right)}{\sqrt{a+b}d} + \frac{\operatorname{arctan}\left(\frac{\sqrt{a+b}\tanh(dx+c)}{\sqrt{-a+b}}\right)}{d\sqrt{-a+b}}$	62
default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b}\tanh(dx+c)}{\sqrt{a+b}}\right)}{\sqrt{a+b}d} + \frac{\operatorname{arctan}\left(\frac{\sqrt{a+b}\tanh(dx+c)}{\sqrt{-a+b}}\right)}{d\sqrt{-a+b}}$	62

input `int(1/(a+b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `arctanh((a+b*tanh(d*x+c))^(1/2)/(a+b)^(1/2))/(a+b)^(1/2)/d+1/d/(-a+b)^(1/2)*arctan((a+b*tanh(d*x+c))^(1/2)/(-a+b)^(1/2))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 351 vs. 2(62) = 124.

Time = 0.14 (sec) , antiderivative size = 2279, normalized size of antiderivative = 30.80

$$\int \frac{1}{\sqrt{a+b\tanh(c+dx)}} dx = \text{Too large to display}$$

input `integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="fricas")`

output

```
[1/4*(sqrt(a + b)*(a - b)*log(2*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 8*(a
^2 + 2*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + 2*(a^2 + 2*a*b + b^2)*si
nh(d*x + c)^4 + 4*(a^2 + a*b)*cosh(d*x + c)^2 + 4*(3*(a^2 + 2*a*b + b^2)*c
osh(d*x + c)^2 + a^2 + a*b)*sinh(d*x + c)^2 + 2*a^2 - b^2 + 2*((a + b)*cos
h(d*x + c)^4 + 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x
+ c)^4 + (2*a + b)*cosh(d*x + c)^2 + (6*(a + b)*cosh(d*x + c)^2 + 2*a + b)
*sinh(d*x + c)^2 + 2*(2*(a + b)*cosh(d*x + c)^3 + (2*a + b)*cosh(d*x + c))
*sinh(d*x + c) + a)*sqrt(a + b)*sqrt((a*cosh(d*x + c) + b*sinh(d*x + c))/c
osh(d*x + c)) + 8*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 + (a^2 + a*b)*cosh(
d*x + c))*sinh(d*x + c)) + (a + b)*sqrt(a - b)*log(((2*a^2 - b^2)*cosh(d*x
+ c)^4 + 4*(2*a^2 - b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + (2*a^2 - b^2)*si
nh(d*x + c)^4 + 4*(a^2 - a*b)*cosh(d*x + c)^2 + 2*(3*(2*a^2 - b^2)*cosh(d*
x + c)^2 + 2*a^2 - 2*a*b)*sinh(d*x + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - 2*(a*c
osh(d*x + c)^4 + 4*a*cosh(d*x + c)*sinh(d*x + c)^3 + a*sinh(d*x + c)^4 + (
2*a - b)*cosh(d*x + c)^2 + (6*a*cosh(d*x + c)^2 + 2*a - b)*sinh(d*x + c)^2
+ 2*(2*a*cosh(d*x + c)^3 + (2*a - b)*cosh(d*x + c))*sinh(d*x + c) + a - b
)*sqrt(a - b)*sqrt((a*cosh(d*x + c) + b*sinh(d*x + c))/cosh(d*x + c)) + 4*
((2*a^2 - b^2)*cosh(d*x + c)^3 + 2*(a^2 - a*b)*cosh(d*x + c))*sinh(d*x + c
)))/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*
sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4))/...
```

SymPy [F]

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx = \int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$$

input

```
integrate(1/(a+b*tanh(d*x+c))**(1/2),x)
```

output

```
Integral(1/sqrt(a + b*tanh(c + d*x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx = \text{Exception raised: ValueError}$$

input `integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see `assume?` for more detail`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m operator + Error: Bad Argument Value`

**Mupad [B] (verification not implemented)**

Time = 2.38 (sec) , antiderivative size = 240, normalized size of antiderivative = 3.24

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$$

$$= \frac{\operatorname{atanh}\left(\frac{(a d^3 + b d^3) \sqrt{a + b \tanh(c + dx)}}{b d^3 \sqrt{a + b}} - \frac{16 a b^2 \sqrt{a + b \tanh(c + dx)}}{\left(\frac{16 b^4 d^3}{a d^3 + b d^3} + \frac{16 a b^3 d^3}{a d^3 + b d^3}\right) \sqrt{a + b}}\right)}{d \sqrt{a + b}}$$

$$+ \frac{\operatorname{atanh}\left(\frac{16 a b^2 \sqrt{a + b \tanh(c + dx)}}{\left(\frac{16 b^4 d^3}{a d^3 - b d^3} - \frac{16 a b^3 d^3}{a d^3 - b d^3}\right) \sqrt{a - b}} + \frac{(a d^3 - b d^3) \sqrt{a + b \tanh(c + dx)}}{b d^3 \sqrt{a - b}}\right)}{d \sqrt{a - b}}$$

input `int(1/(a + b*tanh(c + d*x))^(1/2),x)`output `atanh(((a*d^3 + b*d^3)*(a + b*tanh(c + d*x))^(1/2))/(b*d^3*(a + b)^(1/2)) - (16*a*b^2*(a + b*tanh(c + d*x))^(1/2))/(((16*b^4*d^3)/(a*d^3 + b*d^3) + (16*a*b^3*d^3)/(a*d^3 + b*d^3))*(a + b)^(1/2)))/(d*(a + b)^(1/2)) + atanh((16*a*b^2*(a + b*tanh(c + d*x))^(1/2))/(((16*b^4*d^3)/(a*d^3 - b*d^3) - (16*a*b^3*d^3)/(a*d^3 - b*d^3))*(a - b)^(1/2)) + ((a*d^3 - b*d^3)*(a + b*tanh(c + d*x))^(1/2))/(b*d^3*(a - b)^(1/2)))/(d*(a - b)^(1/2))`**Reduce [F]**

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$$

$$= \frac{2\sqrt{\tanh(dx + c)b + a} + \left(\int \frac{\sqrt{\tanh(dx+c)b+a} \tanh(dx+c)^2 dx}{\tanh(dx+c)b+a}\right) bd}{bd}$$

input `int(1/(a+b*tanh(d*x+c))^(1/2),x)`output `(2*sqrt(tanh(c + d*x)*b + a) + int((sqrt(tanh(c + d*x)*b + a)*tanh(c + d*x)**2)/(tanh(c + d*x)*b + a),x)*b*d)/(b*d)`

### 3.69 $\int \frac{\sinh^4(x)}{1+\tanh(x)} dx$

Optimal result	582
Mathematica [A] (verified)	582
Rubi [A] (verified)	583
Maple [A] (verified)	585
Fricas [B] (verification not implemented)	585
Sympy [F]	586
Maxima [A] (verification not implemented)	586
Giac [A] (verification not implemented)	586
Mupad [B] (verification not implemented)	587
Reduce [B] (verification not implemented)	587

#### Optimal result

Integrand size = 11, antiderivative size = 60

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \frac{x}{16} + \frac{1}{32(1 - \tanh(x))^2} - \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} + \frac{5}{32(1 + \tanh(x))^2} - \frac{3}{16(1 + \tanh(x))}$$

output `1/16*x+1/32/(1-tanh(x))^2-1/(8-8*tanh(x))-1/24/(1+tanh(x))^3+5/32/(1+tanh(x))^2-3/(16+16*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.70

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \frac{1}{192}(12x - 15 \cosh(2x) + 6 \cosh(4x) - \cosh(6x) - 3 \sinh(2x) - 3 \sinh(4x) + \sinh(6x))$$

input `Integrate[Sinh[x]^4/(1 + Tanh[x]),x]`

output

```
(12*x - 15*Cosh[2*x] + 6*Cosh[4*x] - Cosh[6*x] - 3*Sinh[2*x] - 3*Sinh[4*x]
+ Sinh[6*x])/192
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 3999, 25, 516, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sinh^4(x)}{\tanh(x) + 1} dx$$

$$\downarrow 3042$$

$$\int \frac{\sin(ix)^4}{1 - i \tan(ix)} dx$$

$$\downarrow 3999$$

$$- \int -\frac{\tanh^4(x)}{(\tanh(x) + 1)(1 - \tanh^2(x))^3} d \tanh(x)$$

$$\downarrow 25$$

$$\int \frac{\tanh^4(x)}{(\tanh(x) + 1)(1 - \tanh^2(x))^3} d \tanh(x)$$

$$\downarrow 516$$

$$\int \frac{\tanh^4(x)}{(1 - \tanh(x))^3(\tanh(x) + 1)^4} d \tanh(x)$$

$$\downarrow 99$$

$$\int \left( -\frac{1}{16(\tanh^2(x) - 1)} - \frac{1}{8(\tanh(x) - 1)^2} + \frac{3}{16(\tanh(x) + 1)^2} - \frac{1}{16(\tanh(x) - 1)^3} - \frac{5}{16(\tanh(x) + 1)^3} + \frac{1}{8(\tanh(x) - 1)^4} \right) dx$$

$$\downarrow 2009$$



$$\frac{1}{16} \operatorname{arctanh}(\tanh(x)) - \frac{1}{8(1 - \tanh(x))} - \frac{3}{16(\tanh(x) + 1)} + \frac{1}{32(1 - \tanh(x))^2} + \frac{1}{32(\tanh(x) + 1)^2} - \frac{1}{24(\tanh(x) + 1)^3}$$

input `Int[Sinh[x]^4/(1 + Tanh[x]),x]`

output `ArcTanh[Tanh[x]]/16 + 1/(32*(1 - Tanh[x])^2) - 1/(8*(1 - Tanh[x])) - 1/(24*(1 + Tanh[x])^3) + 5/(32*(1 + Tanh[x])^2) - 3/(16*(1 + Tanh[x]))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))`

rule 516 `Int[((e_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[(e*x)^m*(c + d*x)^(n + p)*(a/c + (b/d)*x)^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && EqQ[b*c^2 + a*d^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0] && !IntegerQ[n]))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3999 `Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

**Maple [A] (verified)**

Time = 3.90 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.58

method	result
risch	$\frac{x}{16} + \frac{e^{4x}}{128} - \frac{3e^{2x}}{64} - \frac{e^{-2x}}{32} + \frac{3e^{-4x}}{128} - \frac{e^{-6x}}{192}$
parallelrisc	$\frac{(-12 \cosh(x) - 12 \sinh(x)) \ln(1 - \tanh(x)) + (12 \cosh(x) + 12 \sinh(x)) \ln(1 + \tanh(x)) - 88 \cosh(x) - 64 \sinh(x) - 9 \cosh(3x) + \cosh(5x)}{384 \cosh(x) + 384 \sinh(x)}$
default	$\frac{1}{8(\tanh(\frac{x}{2}) - 1)^4} + \frac{1}{4(\tanh(\frac{x}{2}) - 1)^3} - \frac{1}{8(\tanh(\frac{x}{2}) - 1)} - \frac{\ln(\tanh(\frac{x}{2}) - 1)}{16} - \frac{1}{3(1 + \tanh(\frac{x}{2}))^6} + \frac{1}{(1 + \tanh(\frac{x}{2}))^5} - \frac{1}{8(1 + \tanh(\frac{x}{2}))^4}$

input `int(sinh(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`output `1/16*x+1/128*exp(4*x)-3/64*exp(2*x)-1/32*exp(-2*x)+3/128*exp(-4*x)-1/192*exp(-6*x)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(44) = 88.

Time = 0.08 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.53

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \frac{\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + 5 \sinh(x)^5 + (50 \cosh(x)^2 - 27) \sinh(x)^3 - 9 \cosh(x)^3 + (10 \cosh(x) - 27) \sinh(x)}{384 (\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="fricas")`output `1/384*(cosh(x)^5 + 5*cosh(x)*sinh(x)^4 + 5*sinh(x)^5 + (50*cosh(x)^2 - 27)*sinh(x)^3 - 9*cosh(x)^3 + (10*cosh(x)^3 - 27*cosh(x))*sinh(x)^2 + 12*(2*x - 1)*cosh(x) + (25*cosh(x)^4 - 81*cosh(x)^2 + 24*x + 12)*sinh(x))/(cosh(x) + sinh(x))`

**Sympy [F]**

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \int \frac{\sinh^4(x)}{\tanh(x) + 1} dx$$

input `integrate(sinh(x)**4/(1+tanh(x)),x)`

output `Integral(sinh(x)**4/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.60

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = -\frac{1}{128} (6e^{(-2x)} - 1)e^{(4x)} + \frac{1}{16}x - \frac{1}{32}e^{(-2x)} + \frac{3}{128}e^{(-4x)} - \frac{1}{192}e^{(-6x)}$$

input `integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="maxima")`

output `-1/128*(6*e^(-2*x) - 1)*e^(4*x) + 1/16*x - 1/32*e^(-2*x) + 3/128*e^(-4*x)  
- 1/192*e^(-6*x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.70

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = -\frac{1}{384} (22e^{(6x)} + 12e^{(4x)} - 9e^{(2x)} + 2)e^{(-6x)} + \frac{1}{16}x + \frac{1}{128}e^{(4x)} - \frac{3}{64}e^{(2x)}$$

input `integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="giac")`

output `-1/384*(22*e^(6*x) + 12*e^(4*x) - 9*e^(2*x) + 2)*e^(-6*x) + 1/16*x + 1/128  
*e^(4*x) - 3/64*e^(2*x)`

**Mupad [B] (verification not implemented)**

Time = 2.34 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.57

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \frac{x}{16} - \frac{e^{-2x}}{32} - \frac{3e^{2x}}{64} + \frac{3e^{-4x}}{128} + \frac{e^{4x}}{128} - \frac{e^{-6x}}{192}$$

input `int(sinh(x)^4/(tanh(x) + 1),x)`output `x/16 - exp(-2*x)/32 - (3*exp(2*x))/64 + (3*exp(-4*x))/128 + exp(4*x)/128 - exp(-6*x)/192`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx = \frac{3e^{10x} - 18e^{8x} + 24e^{6x}x - 12e^{4x} + 9e^{2x} - 2}{384e^{6x}}$$

input `int(sinh(x)^4/(1+tanh(x)),x)`output `(3*e**(10*x) - 18*e**(8*x) + 24*e**(6*x)*x - 12*e**(4*x) + 9*e**(2*x) - 2)/(384*e**(6*x))`

### 3.70 $\int \frac{\sinh^3(x)}{1+\tanh(x)} dx$

Optimal result	588
Mathematica [A] (verified)	588
Rubi [C] (verified)	589
Maple [A] (verified)	591
Fricas [B] (verification not implemented)	592
Sympy [B] (verification not implemented)	592
Maxima [A] (verification not implemented)	593
Giac [A] (verification not implemented)	593
Mupad [B] (verification not implemented)	594
Reduce [B] (verification not implemented)	594

#### Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = -\frac{1}{3} \cosh^3(x) + \frac{\cosh^5(x)}{5} - \frac{\sinh^5(x)}{5}$$

output `-1/3*cosh(x)^3+1/5*cosh(x)^5-1/5*sinh(x)^5`

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = \frac{1}{120}(\cosh(x) - \sinh(x))(-20 \cosh(2x) + 4 \cosh(4x) - 10 \sinh(2x) + \sinh(4x))$$

input `Integrate[Sinh[x]^3/(1 + Tanh[x]),x]`

output `((Cosh[x] - Sinh[x])*(-20*Cosh[2*x] + 4*Cosh[4*x] - 10*Sinh[2*x] + Sinh[4*x]))/120`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.40, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.091$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 25, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh^3(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \sin(ix)^3}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\sin(ix)^3}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4001} \\
 & i \int -\frac{i \cosh(x) \sinh^3(x)}{\cosh(x) + \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh^3(x) \cosh(x)}{\sinh(x) + \cosh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \sin(ix)^3 \cos(ix)}{\cos(ix) - i \sin(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\cos(ix) \sin(ix)^3}{\cos(ix) - i \sin(ix)} dx \\
 & \quad \downarrow \text{3587} \\
 & - \int -\cosh(x)(\cosh(x) - \sinh(x)) \sinh^3(x) dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 25 \\
& \int \sinh^3(x) \cosh(x) (\cosh(x) - \sinh(x)) dx \\
& \downarrow 3042 \\
& \int i \sin(ix)^3 \cos(ix) (i \sin(ix) + \cos(ix)) dx \\
& \downarrow 26 \\
& i \int \cos(ix) (\cos(ix) + i \sin(ix)) \sin(ix)^3 dx \\
& \downarrow 3586 \\
& i \int (i \cosh(x) \sinh^4(x) - i \cosh^2(x) \sinh^3(x)) dx \\
& \downarrow 2009 \\
& i \left( \frac{1}{5} i \sinh^5(x) - \frac{1}{5} i \cosh^5(x) + \frac{1}{3} i \cosh^3(x) \right)
\end{aligned}$$

input

```
Int[Sinh[x]^3/(1 + Tanh[x]),x]
```

output

```
I*((I/3)*Cosh[x]^3 - (I/5)*Cosh[x]^5 + (I/5)*Sinh[x]^5)
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 26

```
Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`

rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

## Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

method	result
risch	$\frac{e^{3x}}{48} - \frac{e^x}{8} - \frac{e^{-3x}}{24} + \frac{e^{-5x}}{80}$
parallelrisch	$\frac{4 \cosh(4x) - 16 \sinh(x) + \sinh(4x) - 10 \sinh(2x) - 20 \cosh(2x) - 16 \cosh(x)}{120 \sinh(x) + 120 \cosh(x)}$
default	$\frac{2}{5(1+\tanh(\frac{x}{2}))^5} - \frac{1}{(1+\tanh(\frac{x}{2}))^4} + \frac{2}{3(1+\tanh(\frac{x}{2}))^3} - \frac{1}{8(1+\tanh(\frac{x}{2}))} - \frac{1}{6(\tanh(\frac{x}{2})-1)^3} - \frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{8}$

input `int(sinh(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `1/48*exp(3*x)-1/8*exp(x)-1/24*exp(-3*x)+1/80*exp(-5*x)`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 56 vs.  $2(19) = 38$ .

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.24

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{\cosh(x)^4 + \cosh(x) \sinh(x)^3 + \sinh(x)^4 + (6 \cosh(x)^2 - 5) \sinh(x)^2 - 5 \cosh(x)^2 + (\cosh(x)^3 - 5 \cosh(x)) \sinh(x)}{30 (\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output `1/30*(cosh(x)^4 + cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 5)*sinh(x)^2 - 5*cosh(x)^2 + (cosh(x)^3 - 5*cosh(x))*sinh(x))/(cosh(x) + sinh(x))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 134 vs.  $2(19) = 38$ .

Time = 0.33 (sec) , antiderivative size = 134, normalized size of antiderivative = 5.36

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = \frac{3 \sinh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{3 \sinh^3(x)}{15 \tanh(x) + 15}$$

$$+ \frac{6 \sinh^2(x) \cosh(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{9 \sinh^2(x) \cosh(x)}{15 \tanh(x) + 15}$$

$$- \frac{6 \sinh(x) \cosh^2(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{6 \sinh(x) \cosh^2(x)}{15 \tanh(x) + 15}$$

$$- \frac{8 \cosh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{2 \cosh^3(x)}{15 \tanh(x) + 15}$$

input `integrate(sinh(x)**3/(1+tanh(x)),x)`

output

```
3*sinh(x)**3*tanh(x)/(15*tanh(x) + 15) - 3*sinh(x)**3/(15*tanh(x) + 15) +
6*sinh(x)**2*cosh(x)*tanh(x)/(15*tanh(x) + 15) + 9*sinh(x)**2*cosh(x)/(15*
tanh(x) + 15) - 6*sinh(x)*cosh(x)**2*tanh(x)/(15*tanh(x) + 15) + 6*sinh(x)
*cosh(x)**2/(15*tanh(x) + 15) - 8*cosh(x)**3*tanh(x)/(15*tanh(x) + 15) - 2
*cosh(x)**3/(15*tanh(x) + 15)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = -\frac{1}{48} (6e^{(-2x)} - 1)e^{(3x)} - \frac{1}{24} e^{(-3x)} + \frac{1}{80} e^{(-5x)}$$

input

```
integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="maxima")
```

output

```
-1/48*(6*e^(-2*x) - 1)*e^(3*x) - 1/24*e^(-3*x) + 1/80*e^(-5*x)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = -\frac{1}{240} (10e^{(2x)} - 3)e^{(-5x)} + \frac{1}{48} e^{(3x)} - \frac{1}{8} e^x$$

input

```
integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="giac")
```

output

```
-1/240*(10*e^(2*x) - 3)*e^(-5*x) + 1/48*e^(3*x) - 1/8*e^x
```

**Mupad [B] (verification not implemented)**

Time = 2.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = \frac{e^{3x}}{48} - \frac{e^{-3x}}{24} + \frac{e^{-5x}}{80} - \frac{e^x}{8}$$

input `int(sinh(x)^3/(tanh(x) + 1),x)`output `exp(3*x)/48 - exp(-3*x)/24 + exp(-5*x)/80 - exp(x)/8`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.28

$$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx = \frac{5e^{8x} - 30e^{6x} - 10e^{2x} + 3}{240e^{5x}}$$

input `int(sinh(x)^3/(1+tanh(x)),x)`output `(5*e**(8*x) - 30*e**(6*x) - 10*e**(2*x) + 3)/(240*e**(5*x))`

### 3.71 $\int \frac{\sinh^2(x)}{1+\tanh(x)} dx$

Optimal result	595
Mathematica [A] (verified)	595
Rubi [A] (verified)	596
Maple [A] (verified)	598
Fricas [A] (verification not implemented)	598
Sympy [F]	599
Maxima [A] (verification not implemented)	599
Giac [A] (verification not implemented)	599
Mupad [B] (verification not implemented)	600
Reduce [B] (verification not implemented)	600

#### Optimal result

Integrand size = 11, antiderivative size = 38

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = -\frac{x}{8} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} + \frac{1}{4(1 + \tanh(x))}$$

output `-1/8*x+1/(8-8*tanh(x))-1/8/(1+tanh(x))^2+1/(4+4*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.63

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = \frac{1}{32}(-4x + 4 \cosh(2x) - \cosh(4x) + \sinh(4x))$$

input `Integrate[Sinh[x]^2/(1 + Tanh[x]),x]`

output `(-4*x + 4*Cosh[2*x] - Cosh[4*x] + Sinh[4*x])/32`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 25, 3999, 516, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh^2(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sin(ix)^2}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\sin(ix)^2}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3999} \\
 & \int \frac{\tanh^2(x)}{(\tanh(x) + 1)(1 - \tanh^2(x))^2} d \tanh(x) \\
 & \quad \downarrow \text{516} \\
 & \int \frac{\tanh^2(x)}{(1 - \tanh(x))^2(\tanh(x) + 1)^3} d \tanh(x) \\
 & \quad \downarrow \text{99} \\
 & \int \left( \frac{1}{8(\tanh^2(x) - 1)} + \frac{1}{8(\tanh(x) - 1)^2} - \frac{1}{4(\tanh(x) + 1)^2} + \frac{1}{4(\tanh(x) + 1)^3} \right) d \tanh(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{8} \operatorname{arctanh}(\tanh(x)) + \frac{1}{8(1 - \tanh(x))} + \frac{1}{4(\tanh(x) + 1)} - \frac{1}{8(\tanh(x) + 1)^2}
 \end{aligned}$$

input

```
Int[Sinh[x]^2/(1 + Tanh[x]),x]
```

output 
$$-1/8*\text{ArcTanh}[\text{Tanh}[x]] + 1/(8*(1 - \text{Tanh}[x])) - 1/(8*(1 + \text{Tanh}[x])^2) + 1/(4*(1 + \text{Tanh}[x]))$$

### Defintions of rubi rules used

rule 25 
$$\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \text{:>} \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 99 
$$\text{Int}[((\text{a}_.) + (\text{b}_.)*(\text{x}_))^{\text{(m}_.)}*((\text{c}_.) + (\text{d}_.)*(\text{x}_))^{\text{(n}_.)}*((\text{e}_.) + (\text{f}_.)*(\text{x}_))^{\text{(p}_.)}], \text{x}_] \text{:>} \text{Int}[\text{ExpandIntegrand}[(\text{a} + \text{b}*x)^{\text{m}}*(\text{c} + \text{d}*x)^{\text{n}}*(\text{e} + \text{f}*x)^{\text{p}}, \text{x}], \text{x}] \text{;/}; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{p}\}, \text{x}] \ \&\& \ \text{IntegersQ}[\text{m}, \text{n}] \ \&\& \ (\text{IntegerQ}[\text{p}] \ | \ | \ (\text{GtQ}[\text{m}, 0] \ \&\& \ \text{GeQ}[\text{n}, -1]))$$

rule 516 
$$\text{Int}[(\text{e}_.)*(\text{x}_))^{\text{(m}_.)}*((\text{c}_.) + (\text{d}_.)*(\text{x}_))^{\text{(n}_.)}*((\text{a}_.) + (\text{b}_.)*(\text{x}_)^2)^{\text{(p}_.)}], \text{x\_Symbol}] \text{:>} \text{Int}[(\text{e}*x)^{\text{m}}*(\text{c} + \text{d}*x)^{\text{n} + \text{p}}*(\text{a}/\text{c} + (\text{b}/\text{d})*x)^{\text{p}}, \text{x}] \text{;/}; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{m}, \text{n}, \text{p}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{b}*c^2 + \text{a}*d^2, 0] \ \&\& \ (\text{IntegerQ}[\text{p}] \ || \ | \ (\text{GtQ}[\text{a}, 0] \ \&\& \ \text{GtQ}[\text{c}, 0] \ \&\& \ !\text{IntegerQ}[\text{n}]))$$

rule 2009 
$$\text{Int}[\text{u}_, \text{x\_Symbol}] \text{:>} \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{;/}; \text{SumQ}[\text{u}]$$

rule 3042 
$$\text{Int}[\text{u}_, \text{x\_Symbol}] \text{:>} \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{;/}; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$$

rule 3999 
$$\text{Int}[\sin[(\text{e}_.) + (\text{f}_.)*(\text{x}_))]^{\text{(m}_.)}*((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])^{\text{(n}_.)}], \text{x\_Symbol}] \text{:>} \text{Simp}[\text{b}/\text{f} \quad \text{Subst}[\text{Int}[\text{x}^{\text{m}}*((\text{a} + \text{x})^{\text{n}}/(\text{b}^2 + \text{x}^2)^{\text{m}/2 + 1})], \text{x}], \text{x}, \text{b}*\text{Tan}[\text{e} + \text{f}*x]], \text{x}] \text{;/}; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{n}\}, \text{x}] \ \&\& \ \text{IntegerQ}[\text{m}/2]$$

**Maple [A] (verified)**

Time = 0.71 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.61

method	result
risch	$-\frac{x}{8} + \frac{e^{2x}}{16} + \frac{e^{-2x}}{16} - \frac{e^{-4x}}{32}$
parallelrisch	$\frac{\cosh(3x)+3\sinh(3x)+(-4x-1)\cosh(x)+(-4x-5)\sinh(x)}{32\sinh(x)+32\cosh(x)}$
default	$-\frac{1}{2(1+\tanh(\frac{x}{2}))^4} + \frac{1}{(1+\tanh(\frac{x}{2}))^3} - \frac{1}{2(1+\tanh(\frac{x}{2}))^2} - \frac{\ln(1+\tanh(\frac{x}{2}))}{8} + \frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{4\tanh(\frac{x}{2})-4} + \dots$

input `int(sinh(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)`output `-1/8*x+1/16*exp(2*x)+1/16*exp(-2*x)-1/32*exp(-4*x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + 3 \sinh(x)^3 - 2(2x - 1) \cosh(x) + (9 \cosh(x)^2 - 4x - 2) \sinh(x)}{32(\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="fricas")`output `1/32*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + 3*sinh(x)^3 - 2*(2*x - 1)*cosh(x) + (9*cosh(x)^2 - 4*x - 2)*sinh(x))/(cosh(x) + sinh(x))`

**Sympy [F]**

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = \int \frac{\sinh^2(x)}{\tanh(x) + 1} dx$$

input `integrate(sinh(x)**2/(1+tanh(x)),x)`

output `Integral(sinh(x)**2/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = -\frac{1}{8}x + \frac{1}{16}e^{(2x)} + \frac{1}{16}e^{(-2x)} - \frac{1}{32}e^{(-4x)}$$

input `integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="maxima")`

output `-1/8*x + 1/16*e^(2*x) + 1/16*e^(-2*x) - 1/32*e^(-4*x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = \frac{1}{32} (3e^{(4x)} + 2e^{(2x)} - 1)e^{(-4x)} - \frac{1}{8}x + \frac{1}{16}e^{(2x)}$$

input `integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="giac")`

output `1/32*(3*e^(4*x) + 2*e^(2*x) - 1)*e^(-4*x) - 1/8*x + 1/16*e^(2*x)`



**Mupad [B] (verification not implemented)**

Time = 2.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = \frac{e^{-2x}}{16} - \frac{x}{8} + \frac{e^{2x}}{16} - \frac{e^{-4x}}{32}$$

input `int(sinh(x)^2/(tanh(x) + 1),x)`output `exp(-2*x)/16 - x/8 + exp(2*x)/16 - exp(-4*x)/32`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx = \frac{2e^{6x} - 4e^{4x}x + 2e^{2x} - 1}{32e^{4x}}$$

input `int(sinh(x)^2/(1+tanh(x)),x)`output `(2*e**(6*x) - 4*e**(4*x)*x + 2*e**(2*x) - 1)/(32*e**(4*x))`

### 3.72 $\int \frac{\sinh(x)}{1+\tanh(x)} dx$

Optimal result . . . . .	601
Mathematica [A] (verified) . . . . .	601
Rubi [C] (verified) . . . . .	602
Maple [A] (verified) . . . . .	604
Fricas [A] (verification not implemented) . . . . .	604
Sympy [B] (verification not implemented) . . . . .	605
Maxima [A] (verification not implemented) . . . . .	605
Giac [A] (verification not implemented) . . . . .	606
Mupad [B] (verification not implemented) . . . . .	606
Reduce [B] (verification not implemented) . . . . .	606

#### Optimal result

Integrand size = 9, antiderivative size = 17

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{\cosh^3(x)}{3} - \frac{\sinh^3(x)}{3}$$

output `1/3*cosh(x)^3-1/3*sinh(x)^3`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{1}{12} (3 \cosh(x) + \cosh(3x) - 4 \sinh^3(x))$$

input `Integrate[Sinh[x]/(1 + Tanh[x]),x]`

output `(3*Cosh[x] + Cosh[3*x] - 4*Sinh[x]^3)/12`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.222$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4001} \\
 & -i \int \frac{i \cosh(x) \sinh(x)}{\cosh(x) + \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh(x) \cosh(x)}{\sinh(x) + \cosh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix) \cos(ix)}{\cos(ix) - i \sin(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\cos(ix) \sin(ix)}{\cos(ix) - i \sin(ix)} dx \\
 & \quad \downarrow \text{3587} \\
 & \int \sinh(x) \cosh(x) (\cosh(x) - \sinh(x)) dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int -i \sin(ix) \cos(ix) (i \sin(ix) + \cos(ix)) dx \\
& \quad \downarrow \text{26} \\
& -i \int \cos(ix) (\cos(ix) + i \sin(ix)) \sin(ix) dx \\
& \quad \downarrow \text{3586} \\
& -i \int (i \cosh^2(x) \sinh(x) - i \cosh(x) \sinh^2(x)) dx \\
& \quad \downarrow \text{2009} \\
& -i \left( \frac{1}{3} i \cosh^3(x) - \frac{1}{3} i \sinh^3(x) \right)
\end{aligned}$$

input `Int[Sinh[x]/(1 + Tanh[x]),x]`

output `(-I)*((I/3)*Cosh[x]^3 - (I/3)*Sinh[x]^3)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] :> Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`

rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

### Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
risch	$\frac{e^x}{4} + \frac{e^{-3x}}{12}$	12
parallelrisc	$\frac{4 \sinh(x) + \sinh(2x) + 2 \cosh(2x) + 4 \cosh(x)}{6 \cosh(x) + 6 \sinh(x)}$	32
default	$\frac{2}{3(1+\tanh(\frac{x}{2}))^3} - \frac{1}{(1+\tanh(\frac{x}{2}))^2} + \frac{1}{2 \tanh(\frac{x}{2})+2} - \frac{1}{2(\tanh(\frac{x}{2})-1)}$	42

input `int(sinh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `1/4*exp(x)+1/12*exp(-3*x)`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{\cosh(x)^2 + \cosh(x) \sinh(x) + \sinh(x)^2}{3(\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)/(1+tanh(x)),x, algorithm="fricas")`

output `1/3*(cosh(x)^2 + cosh(x)*sinh(x) + sinh(x)^2)/(cosh(x) + sinh(x))`

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs.  $2(12) = 24$ .

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.82

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{\sinh(x) \tanh(x)}{3 \tanh(x) + 3} - \frac{\sinh(x)}{3 \tanh(x) + 3} + \frac{2 \cosh(x) \tanh(x)}{3 \tanh(x) + 3} + \frac{\cosh(x)}{3 \tanh(x) + 3}$$

input `integrate(sinh(x)/(1+tanh(x)),x)`

output `sinh(x)*tanh(x)/(3*tanh(x) + 3) - sinh(x)/(3*tanh(x) + 3) + 2*cosh(x)*tanh(x)/(3*tanh(x) + 3) + cosh(x)/(3*tanh(x) + 3)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{1}{12} e^{(-3x)} + \frac{1}{4} e^x$$

input `integrate(sinh(x)/(1+tanh(x)),x, algorithm="maxima")`

output `1/12*e^(-3*x) + 1/4*e^x`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{1}{12} e^{(-3x)} + \frac{1}{4} e^x$$

input `integrate(sinh(x)/(1+tanh(x)),x, algorithm="giac")`

output `1/12*e^(-3*x) + 1/4*e^x`

**Mupad [B] (verification not implemented)**

Time = 2.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{e^{-3x}}{12} + \frac{e^x}{4}$$

input `int(sinh(x)/(tanh(x) + 1),x)`

output `exp(-3*x)/12 + exp(x)/4`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

$$\int \frac{\sinh(x)}{1 + \tanh(x)} dx = \frac{12e^{3x} \cosh(x) - 3e^{4x} - 6e^{2x} + 1}{12e^{3x}}$$

input `int(sinh(x)/(1+tanh(x)),x)`

output `(12*e**(3*x)*cosh(x) - 3*e**(4*x) - 6*e**(2*x) + 1)/(12*e**(3*x))`

### 3.73 $\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx$

Optimal result	607
Mathematica [B] (verified)	607
Rubi [C] (verified)	608
Maple [A] (verified)	610
Fricas [B] (verification not implemented)	611
Sympy [F]	611
Maxima [A] (verification not implemented)	611
Giac [A] (verification not implemented)	612
Mupad [B] (verification not implemented)	612
Reduce [B] (verification not implemented)	612

#### Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx = -\operatorname{arctanh}(\cosh(x)) + \cosh(x) - \sinh(x)$$

output

```
-arctanh(cosh(x))+cosh(x)-sinh(x)
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 49 vs.  $2(12) = 24$ .

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 4.08

$$\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx = \frac{\cosh(x) - \log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) - \left(\log\left(\cosh\left(\frac{x}{2}\right)\right) - \log\left(\sinh\left(\frac{x}{2}\right)\right) + \sinh(x)\right) \tanh(x)}{1+\tanh(x)}$$

input

```
Integrate[Csch[x]/(1 + Tanh[x]), x]
```



output

```
(Cosh[x] - Log[Cosh[x/2]] + Log[Sinh[x/2]] - (Log[Cosh[x/2]] - Log[Sinh[x/2]] + Sinh[x])*Tanh[x])/(1 + Tanh[x])
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.00, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.333$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 25, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\sin(ix)(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\sin(ix)(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & i \int -\frac{i \operatorname{coth}(x)}{\cosh(x) + \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\operatorname{coth}(x)}{\sinh(x) + \cosh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \cos(ix)}{\sin(ix)(\cos(ix) - i \sin(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\cos(ix)}{(\cos(ix) - i \sin(ix)) \sin(ix)} dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3587} \\
& - \int -\coth(x)(\cosh(x) - \sinh(x))dx \\
& \downarrow \text{25} \\
& \int \coth(x)(\cosh(x) - \sinh(x))dx \\
& \downarrow \text{3042} \\
& \int \frac{i \cos(ix)(i \sin(ix) + \cos(ix))}{\sin(ix)} dx \\
& \downarrow \text{26} \\
& i \int \frac{\cos(ix)(\cos(ix) + i \sin(ix))}{\sin(ix)} dx \\
& \downarrow \text{3586} \\
& i \int (i \cosh(x) - i \cosh(x) \coth(x)) dx \\
& \downarrow \text{2009} \\
& i(\operatorname{arctanh}(\cosh(x)) + i \sinh(x) - i \cosh(x))
\end{aligned}$$

input `Int [Csch[x]/(1 + Tanh[x]), x]`

output `I*(I*ArcTanh[Cosh[x]] - I*Cosh[x] + I*Sinh[x])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`

rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

## Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

method	result	size
default	$\ln\left(\tanh\left(\frac{x}{2}\right)\right) + \frac{2}{1+\tanh\left(\frac{x}{2}\right)}$	17
risch	$e^{-x} + \ln(e^x - 1) - \ln(e^x + 1)$	18

input `int(csch(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `ln(tanh(1/2*x))+2/(1+tanh(1/2*x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 38 vs.  $2(12) = 24$ .

Time = 0.09 (sec) , antiderivative size = 38, normalized size of antiderivative = 3.17

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = \frac{-(\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) + 1) - (\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) - 1)}{\cosh(x) + \sinh(x)}$$

input `integrate(csch(x)/(1+tanh(x)),x, algorithm="fricas")`

output `-((cosh(x) + sinh(x))*log(cosh(x) + sinh(x) + 1) - (cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) - 1)/(cosh(x) + sinh(x))`

**Sympy [F]**

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}(x)}{\tanh(x) + 1} dx$$

input `integrate(csch(x)/(1+tanh(x)),x)`

output `Integral(csch(x)/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = e^{(-x)} - \log(e^{(-x)} + 1) + \log(e^{(-x)} - 1)$$

input `integrate(csch(x)/(1+tanh(x)),x, algorithm="maxima")`

output `e^(-x) - log(e^(-x) + 1) + log(e^(-x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = e^{-x} - \log(e^x + 1) + \log(|e^x - 1|)$$

input `integrate(csch(x)/(1+tanh(x)),x, algorithm="giac")`

output `e^(-x) - log(e^x + 1) + log(abs(e^x - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = \ln(2 - 2e^x) - \ln(-2e^x - 2) + e^{-x}$$

input `int(1/(sinh(x)*(tanh(x) + 1)),x)`

output `log(2 - 2*exp(x)) - log(- 2*exp(x) - 2) + exp(-x)`

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.42

$$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx = \frac{e^x \log(e^x - 1) - e^x \log(e^x + 1) + 1}{e^x}$$

input `int(csch(x)/(1+tanh(x)),x)`

output `(e**x*log(e**x - 1) - e**x*log(e**x + 1) + 1)/e**x`

### 3.74 $\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx$

Optimal result	613
Mathematica [A] (verified)	613
Rubi [A] (verified)	614
Maple [A] (verified)	615
Fricas [B] (verification not implemented)	616
Sympy [F]	616
Maxima [A] (verification not implemented)	617
Giac [A] (verification not implemented)	617
Mupad [B] (verification not implemented)	617
Reduce [B] (verification not implemented)	618

#### Optimal result

Integrand size = 11, antiderivative size = 15

$$\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx = -\operatorname{coth}(x) - \log(\tanh(x)) + \log(1+\tanh(x))$$

output `-coth(x)-ln(tanh(x))+ln(1+tanh(x))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx = x - \operatorname{coth}(x) - \log(\sinh(x))$$

input `Integrate[Csch[x]^2/(1 + Tanh[x]), x]`

output `x - Coth[x] - Log[Sinh[x]]`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {3042, 25, 3999, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^2(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\sin(ix)^2(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\sin(ix)^2(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{3999} \\
 & \int \frac{\operatorname{coth}^2(x)}{\tanh(x) + 1} d \tanh(x) \\
 & \quad \downarrow \text{54} \\
 & \int \left( \frac{1}{\tanh(x) + 1} + \operatorname{coth}^2(x) - \operatorname{coth}(x) \right) d \tanh(x) \\
 & \quad \downarrow \text{2009} \\
 & -\operatorname{coth}(x) - \log(\tanh(x)) + \log(\tanh(x) + 1)
 \end{aligned}$$

input

```
Int [Csch[x]^2/(1 + Tanh[x]), x]
```

output

```
-Coth[x] - Log[Tanh[x]] + Log[1 + Tanh[x]]
```

## Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 54 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3999 `Int[sin[(e_) + (f_)*(x_)^(m_))*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

method	result	size
risch	$2x - \frac{2}{e^{2x}-1} - \ln(e^{2x}-1)$	24
default	$-\frac{\tanh(\frac{x}{2})}{2} + 2 \ln(1 + \tanh(\frac{x}{2})) - \frac{1}{2 \tanh(\frac{x}{2})} - \ln(\tanh(\frac{x}{2}))$	32

input `int(csch(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `2*x-2/(exp(2*x)-1)-ln(exp(2*x)-1)`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 77 vs.  $2(15) = 30$ .

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 5.13

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{2x \cosh(x)^2 + 4x \cosh(x) \sinh(x) + 2x \sinh(x)^2 - (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \log(2 \sinh(x) / (\cosh(x) - \sinh(x))) - 2x - 2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1}$$

input `integrate(csch(x)^2/(1+tanh(x)),x, algorithm="fricas")`

output `(2*x*cosh(x)^2 + 4*x*cosh(x)*sinh(x) + 2*x*sinh(x)^2 - (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*log(2*sinh(x)/(cosh(x) - sinh(x))) - 2*x - 2)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)`

**Sympy [F]**

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}^2(x)}{\tanh(x) + 1} dx$$

input `integrate(csch(x)**2/(1+tanh(x)),x)`

output `Integral(csch(x)**2/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx = \frac{2}{e^{(-2x)} - 1} - \log(e^{(-x)} + 1) - \log(e^{(-x)} - 1)$$

input `integrate(csch(x)^2/(1+tanh(x)),x, algorithm="maxima")`output `2/(e^(-2*x) - 1) - log(e^(-x) + 1) - log(e^(-x) - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx = 2x + \frac{e^{(2x)} - 3}{e^{(2x)} - 1} - \log(|e^{(2x)} - 1|)$$

input `integrate(csch(x)^2/(1+tanh(x)),x, algorithm="giac")`output `2*x + (e^(2*x) - 3)/(e^(2*x) - 1) - log(abs(e^(2*x) - 1))`**Mupad [B] (verification not implemented)**

Time = 2.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx = 2x - \ln(e^{2x} - 1) - \frac{2}{e^{2x} - 1}$$

input `int(1/(sinh(x)^2*(tanh(x) + 1)),x)`output `2*x - log(exp(2*x) - 1) - 2/(exp(2*x) - 1)`

**Reduce [B] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 67, normalized size of antiderivative = 4.47

$$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{-e^{2x} \log(e^x - 1) - e^{2x} \log(e^x + 1) + 2e^{2x}x - 2e^{2x} + \log(e^x - 1) + \log(e^x + 1) - 2x}{e^{2x} - 1}$$

input `int(csch(x)^2/(1+tanh(x)),x)`output `( - e**(2*x)*log(e**x - 1) - e**(2*x)*log(e**x + 1) + 2*e**(2*x)*x - 2*e**(2*x) + log(e**x - 1) + log(e**x + 1) - 2*x)/(e**(2*x) - 1)`

### 3.75 $\int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx$

Optimal result	619
Mathematica [B] (verified)	619
Rubi [C] (verified)	620
Maple [B] (verified)	622
Fricas [B] (verification not implemented)	623
Sympy [F]	623
Maxima [B] (verification not implemented)	624
Giac [B] (verification not implemented)	624
Mupad [B] (verification not implemented)	624
Reduce [B] (verification not implemented)	625

#### Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx = -\frac{1}{2}\operatorname{arctanh}(\cosh(x)) + \operatorname{csch}(x) - \frac{1}{2}\operatorname{coth}(x)\operatorname{csch}(x)$$

output `-1/2*arctanh(cosh(x))+csch(x)-1/2*coth(x)*csch(x)`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 59 vs.  $2(18) = 36$ .

Time = 0.20 (sec) , antiderivative size = 59, normalized size of antiderivative = 3.28

$$\int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx = \frac{1}{8} \left( 4 \operatorname{coth} \left( \frac{x}{2} \right) - \operatorname{csch}^2 \left( \frac{x}{2} \right) - 4 \log \left( \cosh \left( \frac{x}{2} \right) \right) + 4 \log \left( \sinh \left( \frac{x}{2} \right) \right) \right. \\ \left. - \operatorname{sech}^2 \left( \frac{x}{2} \right) - 4 \tanh \left( \frac{x}{2} \right) \right)$$

input `Integrate[Csch[x]^3/(1 + Tanh[x]),x]`

output

$$(4*\text{Coth}[x/2] - \text{Csch}[x/2]^2 - 4*\text{Log}[\text{Cosh}[x/2]] + 4*\text{Log}[\text{Sinh}[x/2]] - \text{Sech}[x/2]^2 - 4*\text{Tanh}[x/2])/8$$
**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.67, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\text{csch}^3(x)}{\tanh(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{i}{\sin(ix)^3(1 - i \tan(ix))} dx \\ & \quad \downarrow \text{26} \\ & -i \int \frac{1}{\sin(ix)^3(1 - i \tan(ix))} dx \\ & \quad \downarrow \text{4001} \\ & -i \int \frac{i \coth(x) \text{csch}^2(x)}{\cosh(x) + \sinh(x)} dx \\ & \quad \downarrow \text{26} \\ & \int \frac{\coth(x) \text{csch}^2(x)}{\sinh(x) + \cosh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{i \cos(ix)}{\sin(ix)^3(\cos(ix) - i \sin(ix))} dx \\ & \quad \downarrow \text{26} \\ & -i \int \frac{\cos(ix)}{(\cos(ix) - i \sin(ix)) \sin(ix)^3} dx \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3587} \\
& \int \coth(x) \operatorname{csch}^2(x) (\cosh(x) - \sinh(x)) dx \\
& \downarrow \text{3042} \\
& \int -\frac{i \cos(ix) (i \sin(ix) + \cos(ix))}{\sin(ix)^3} dx \\
& \downarrow \text{26} \\
& -i \int \frac{\cos(ix) (\cos(ix) + i \sin(ix))}{\sin(ix)^3} dx \\
& \downarrow \text{3586} \\
& -i \int (i \coth^2(x) \operatorname{csch}(x) - i \coth(x) \operatorname{csch}(x)) dx \\
& \downarrow \text{2009} \\
& -i \left( -\frac{1}{2} i \operatorname{arctanh}(\cosh(x)) + i \operatorname{csch}(x) - \frac{1}{2} i \coth(x) \operatorname{csch}(x) \right)
\end{aligned}$$

input `Int[Csch[x]^3/(1 + Tanh[x]), x]`

output `(-I)*((-1/2*I)*ArcTanh[Cosh[x]] + I*Csch[x] - (I/2)*Coth[x]*Csch[x])`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`

rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs.  $2(14) = 28$ .

Time = 0.53 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.83

method	result	size
risch	$\frac{e^x(e^{2x}-3)}{(e^{2x}-1)^2} + \frac{\ln(e^x-1)}{2} - \frac{\ln(e^x+1)}{2}$	33
default	$\frac{\tanh(\frac{x}{2})^2}{8} - \frac{\tanh(\frac{x}{2})}{2} - \frac{1}{8\tanh(\frac{x}{2})^2} + \frac{1}{2\tanh(\frac{x}{2})} + \frac{\ln(\tanh(\frac{x}{2}))}{2}$	39

input `int(csch(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `exp(x)*(exp(2*x)-3)/(exp(2*x)-1)^2+1/2*ln(exp(x)-1)-1/2*ln(exp(x)+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 209 vs.  $2(14) = 28$ .

Time = 0.09 (sec) , antiderivative size = 209, normalized size of antiderivative = 11.61

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{2 \cosh(x)^3 + 6 \cosh(x) \sinh(x)^2 + 2 \sinh(x)^3 - (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^3 - \cosh(x)) \sinh(x) + 1) \log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^3 - \cosh(x)) \sinh(x) + 1) \log(\cosh(x) + \sinh(x) - 1) + 6(\cosh(x)^2 - 1) \sinh(x) - 6 \cosh(x)}{\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^3 - \cosh(x)) \sinh(x) + 1}$$

input `integrate(csch(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output

$$\frac{1/2*(2*\cosh(x)^3 + 6*\cosh(x)*\sinh(x)^2 + 2*\sinh(x)^3 - (\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 - 1)*\sinh(x)^2 - 2*\cosh(x)^2 + 4*(\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)*\log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 - 1)*\sinh(x)^2 - 2*\cosh(x)^2 + 4*(\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)*\log(\cosh(x) + \sinh(x) - 1) + 6*(\cosh(x)^2 - 1)*\sinh(x) - 6*\cosh(x))}{(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 - 1)*\sinh(x)^2 - 2*\cosh(x)^2 + 4*(\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)}$$
**Sympy [F]**

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}^3(x)}{\tanh(x) + 1} dx$$

input `integrate(csch(x)**3/(1+tanh(x)),x)`

output `Integral(csch(x)**3/(tanh(x) + 1), x)`



**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 48 vs.  $2(14) = 28$ .

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.67

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx = -\frac{e^{(-x)} - 3e^{(-3x)}}{2e^{(-2x)} - e^{(-4x)} - 1} - \frac{1}{2} \log(e^{(-x)} + 1) + \frac{1}{2} \log(e^{(-x)} - 1)$$

input `integrate(csch(x)^3/(1+tanh(x)),x, algorithm="maxima")`

output `-(e^(-x) - 3*e^(-3*x))/(2*e^(-2*x) - e^(-4*x) - 1) - 1/2*log(e^(-x) + 1) + 1/2*log(e^(-x) - 1)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 34 vs.  $2(14) = 28$ .

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.89

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx = \frac{e^{(3x)} - 3e^x}{(e^{(2x)} - 1)^2} - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(csch(x)^3/(1+tanh(x)),x, algorithm="giac")`

output `(e^(3*x) - 3*e^x)/(e^(2*x) - 1)^2 - 1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.67

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx = \frac{\ln(1 - e^x)}{2} - \frac{\ln(-e^x - 1)}{2} + \frac{e^x}{e^{2x} - 1} - \frac{2e^x}{e^{4x} - 2e^{2x} + 1}$$

input `int(1/(sinh(x)^3*(tanh(x) + 1)),x)`

output  $\frac{\log(1 - \exp(x))/2 - \log(-\exp(x) - 1)/2 + \exp(x)/(\exp(2x) - 1) - (2\exp(x))/(\exp(4x) - 2\exp(2x) + 1)}$

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 97, normalized size of antiderivative = 5.39

$$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{e^{4x} \log(e^x - 1) - e^{4x} \log(e^x + 1) + 2e^{3x} - 2e^{2x} \log(e^x - 1) + 2e^{2x} \log(e^x + 1) - 6e^x + \log(e^x - 1) - \log(e^x + 1)}{2e^{4x} - 4e^{2x} + 2}$$

input `int(csch(x)^3/(1+tanh(x)),x)`

output  $(e^{4x} \log(e^x - 1) - e^{4x} \log(e^x + 1) + 2e^{3x} - 2e^{2x} \log(e^x - 1) + 2e^{2x} \log(e^x + 1) - 6e^x + \log(e^x - 1) - \log(e^x + 1))/(2e^{4x} - 2e^{2x} + 1)$

### 3.76 $\int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx$

Optimal result	626
Mathematica [A] (verified)	626
Rubi [A] (verified)	627
Maple [A] (verified)	628
Fricas [B] (verification not implemented)	629
Sympy [F]	629
Maxima [B] (verification not implemented)	630
Giac [A] (verification not implemented)	630
Mupad [B] (verification not implemented)	631
Reduce [B] (verification not implemented)	631

#### Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx = \frac{\operatorname{coth}^2(x)}{2} - \frac{\operatorname{coth}^3(x)}{3}$$

output `1/2*coth(x)^2-1/3*coth(x)^3`

#### Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx = -\frac{1}{6}\operatorname{csch}(x)(2\cosh(x) + (-3 + 2\operatorname{coth}(x))\operatorname{csch}(x))$$

input `Integrate[Csch[x]^4/(1 + Tanh[x]), x]`

output `-1/6*(Csch[x]*(2*Cosh[x] + (-3 + 2*Coth[x])*Csch[x]))`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 3999, 25, 516, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^4(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(ix)^4(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{3999} \\
 & - \int -\frac{\operatorname{coth}^4(x)(1 - \tanh^2(x))}{\tanh(x) + 1} d \tanh(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{(1 - \tanh^2(x)) \operatorname{coth}^4(x)}{\tanh(x) + 1} d \tanh(x) \\
 & \quad \downarrow \text{516} \\
 & \int (1 - \tanh(x)) \operatorname{coth}^4(x) d \tanh(x) \\
 & \quad \downarrow \text{53} \\
 & \int (\operatorname{coth}^4(x) - \operatorname{coth}^3(x)) d \tanh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{\operatorname{coth}^2(x)}{2} - \frac{\operatorname{coth}^3(x)}{3}
 \end{aligned}$$

input `Int [Csch[x]^4/(1 + Tanh[x]), x]`

output `Coth[x]^2/2 - Coth[x]^3/3`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 53 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`
- rule 516 `Int[((e_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[(e*x)^m*(c + d*x)^(n + p)*(a/c + (b/d)*x)^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && EqQ[b*c^2 + a*d^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0] && !IntegerQ[n]))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3999 `Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

method	result	size
risch	$-\frac{2(3e^{2x}+1)}{3(e^{2x}-1)^3}$	19
parallelrisc	$\frac{\operatorname{sech}(\frac{x}{2})^3 \operatorname{csch}(\frac{x}{2})^3 (-4 \cosh(3x) - 12 \cosh(x) + 3 \sinh(x) + 7 \sinh(3x))}{384}$	36
default	$-\frac{\tanh(\frac{x}{2})^3}{24} + \frac{\tanh(\frac{x}{2})^2}{8} - \frac{\tanh(\frac{x}{2})}{8} - \frac{1}{24 \tanh(\frac{x}{2})^3} - \frac{1}{8 \tanh(\frac{x}{2})} + \frac{1}{8 \tanh(\frac{x}{2})^2}$	48



**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs.  $2(13) = 26$ .

Time = 0.05 (sec) , antiderivative size = 75, normalized size of antiderivative = 4.41

$$\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx = -\frac{2e^{(-2x)}}{3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1} + \frac{4e^{(-4x)}}{3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1} + \frac{2}{3(3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1)}$$

input `integrate(csch(x)^4/(1+tanh(x)),x, algorithm="maxima")`

output `-2*e^(-2*x)/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1) + 4*e^(-4*x)/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1) + 2/3/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx = -\frac{2(3e^{(2x)} + 1)}{3(e^{(2x)} - 1)^3}$$

input `integrate(csch(x)^4/(1+tanh(x)),x, algorithm="giac")`

output `-2/3*(3*e^(2*x) + 1)/(e^(2*x) - 1)^3`

**Mupad [B] (verification not implemented)**

Time = 2.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx = -\frac{2(3e^{2x} + 1)}{3(e^{2x} - 1)^3}$$

input `int(1/(sinh(x)^4*(tanh(x) + 1)),x)`output `-(2*(3*exp(2*x) + 1))/(3*(exp(2*x) - 1)^3)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.12

$$\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx = \frac{-6e^{2x} - 2}{3e^{6x} - 9e^{4x} + 9e^{2x} - 3}$$

input `int(csch(x)^4/(1+tanh(x)),x)`output `(2*( - 3*e**(2*x) - 1))/(3*(e**(6*x) - 3*e**(4*x) + 3*e**(2*x) - 1))`



### 3.77 $\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx$

Optimal result	632
Mathematica [B] (verified)	632
Rubi [C] (verified)	633
Maple [A] (verified)	636
Fricas [B] (verification not implemented)	636
Sympy [F]	637
Maxima [B] (verification not implemented)	638
Giac [A] (verification not implemented)	638
Mupad [B] (verification not implemented)	639
Reduce [B] (verification not implemented)	639

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx = \frac{1}{8} \operatorname{arctanh}(\cosh(x)) - \frac{1}{8} \operatorname{coth}(x) \operatorname{csch}(x) + \frac{\operatorname{csch}^3(x)}{3} - \frac{1}{4} \operatorname{coth}(x) \operatorname{csch}^3(x)$$

output

```
1/8*arctanh(cosh(x))-1/8*coth(x)*csch(x)+1/3*csch(x)^3-1/4*coth(x)*csch(x)^3
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 69 vs.  $2(34) = 68$ .

Time = 0.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.03

$$\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx = \frac{1}{192} \operatorname{csch}^4(x) \left( -42 \cosh(x) - 6 \cosh(3x) + 2 \sinh(x) \left( 32 - 9 \left( \log \left( \cosh \left( \frac{x}{2} \right) \right) - \log \left( \sinh \left( \frac{x}{2} \right) \right) \right) \sinh(x) + 3 \left( \log \left( \cosh \left( \frac{x}{2} \right) \right) - \log \left( \sinh \left( \frac{x}{2} \right) \right) \right) \sinh(3x) \right)$$

input `Integrate[Csch[x]^5/(1 + Tanh[x]),x]`

output `(Csch[x]^4*(-42*Cosh[x] - 6*Cosh[3*x] + 2*Sinh[x]*(32 - 9*(Log[Cosh[x/2]] - Log[Sinh[x/2]])*Sinh[x] + 3*(Log[Cosh[x/2]] - Log[Sinh[x/2]])*Sinh[3*x]))/192`

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.091$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 25, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^5(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\sin(ix)^5(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\sin(ix)^5(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & i \int -\frac{i \coth(x) \operatorname{csch}^4(x)}{\cosh(x) + \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x) \operatorname{csch}^4(x)}{\sinh(x) + \cosh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \cos(ix)}{\sin(ix)^5(\cos(ix) - i \sin(ix))} dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& i \int \frac{\cos(ix)}{(\cos(ix) - i \sin(ix)) \sin(ix)^5} dx \\
& \downarrow 3587 \\
& - \int -\coth(x) \operatorname{csch}^4(x) (\cosh(x) - \sinh(x)) dx \\
& \downarrow 25 \\
& \int \coth(x) \operatorname{csch}^4(x) (\cosh(x) - \sinh(x)) dx \\
& \downarrow 3042 \\
& \int \frac{i \cos(ix) (i \sin(ix) + \cos(ix))}{\sin(ix)^5} dx \\
& \downarrow 26 \\
& i \int \frac{\cos(ix) (\cos(ix) + i \sin(ix))}{\sin(ix)^5} dx \\
& \downarrow 3586 \\
& i \int (i \coth(x) \operatorname{csch}^3(x) - i \coth^2(x) \operatorname{csch}^3(x)) dx \\
& \downarrow 2009 \\
& i \left( -\frac{1}{8} i \operatorname{arctanh}(\cosh(x)) - \frac{1}{3} i \operatorname{csch}^3(x) + \frac{1}{4} i \coth(x) \operatorname{csch}^3(x) + \frac{1}{8} i \coth(x) \operatorname{csch}(x) \right)
\end{aligned}$$

input `Int [Csch[x]^5/(1 + Tanh[x]), x]`

output `I*((-1/8*I)*ArcTanh[Cosh[x]] + (I/8)*Coth[x]*Csch[x] - (I/3)*Csch[x]^3 + (I/4)*Coth[x]*Csch[x]^3)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`
- rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`
- rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

**Maple [A] (verified)**

Time = 2.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.41

method	result	size
risch	$-\frac{e^x(3e^{6x}-11e^{4x}+53e^{2x}+3)}{12(e^{2x}-1)^4} - \frac{\ln(e^x-1)}{8} + \frac{\ln(e^x+1)}{8}$	48
default	$\frac{\tanh(\frac{x}{2})^4}{64} - \frac{\tanh(\frac{x}{2})^3}{24} + \frac{\tanh(\frac{x}{2})}{8} - \frac{1}{64 \tanh(\frac{x}{2})^4} + \frac{1}{24 \tanh(\frac{x}{2})^3} - \frac{1}{8 \tanh(\frac{x}{2})} - \frac{\ln(\tanh(\frac{x}{2}))}{8}$	55

input `int(csch(x)^5/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-1/12*exp(x)*(3*exp(6*x)-11*exp(4*x)+53*exp(2*x)+3)/(exp(2*x)-1)^4-1/8*ln(exp(x)-1)+1/8*ln(exp(x)+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 640 vs.  $2(26) = 52$ .

Time = 0.09 (sec) , antiderivative size = 640, normalized size of antiderivative = 18.82

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(csch(x)^5/(1+tanh(x)),x, algorithm="fricas")`

output

```

-1/24*(6*cosh(x)^7 + 42*cosh(x)*sinh(x)^6 + 6*sinh(x)^7 + 2*(63*cosh(x)^2
- 11)*sinh(x)^5 - 22*cosh(x)^5 + 10*(21*cosh(x)^3 - 11*cosh(x))*sinh(x)^4
+ 2*(105*cosh(x)^4 - 110*cosh(x)^2 + 53)*sinh(x)^3 + 106*cosh(x)^3 + 2*(63
*cosh(x)^5 - 110*cosh(x)^3 + 159*cosh(x))*sinh(x)^2 - 3*(cosh(x)^8 + 8*cos
h(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 - 1)*sinh(x)^6 - 4*cosh(x)^6 +
8*(7*cosh(x)^3 - 3*cosh(x))*sinh(x)^5 + 2*(35*cosh(x)^4 - 30*cosh(x)^2 +
3)*sinh(x)^4 + 6*cosh(x)^4 + 8*(7*cosh(x)^5 - 10*cosh(x)^3 + 3*cosh(x))*si
nh(x)^3 + 4*(7*cosh(x)^6 - 15*cosh(x)^4 + 9*cosh(x)^2 - 1)*sinh(x)^2 - 4*c
osh(x)^2 + 8*(cosh(x)^7 - 3*cosh(x)^5 + 3*cosh(x)^3 - cosh(x))*sinh(x) + 1
)*log(cosh(x) + sinh(x) + 1) + 3*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)
)^8 + 4*(7*cosh(x)^2 - 1)*sinh(x)^6 - 4*cosh(x)^6 + 8*(7*cosh(x)^3 - 3*cos
h(x))*sinh(x)^5 + 2*(35*cosh(x)^4 - 30*cosh(x)^2 + 3)*sinh(x)^4 + 6*cosh(x)
)^4 + 8*(7*cosh(x)^5 - 10*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^
6 - 15*cosh(x)^4 + 9*cosh(x)^2 - 1)*sinh(x)^2 - 4*cosh(x)^2 + 8*(cosh(x)^7
- 3*cosh(x)^5 + 3*cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x)
- 1) + 2*(21*cosh(x)^6 - 55*cosh(x)^4 + 159*cosh(x)^2 + 3)*sinh(x) + 6*c
osh(x))/(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 - 1)*
sinh(x)^6 - 4*cosh(x)^6 + 8*(7*cosh(x)^3 - 3*cosh(x))*sinh(x)^5 + 2*(35*c
osh(x)^4 - 30*cosh(x)^2 + 3)*sinh(x)^4 + 6*cosh(x)^4 + 8*(7*cosh(x)^5 - 10*
cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^6 - 15*cosh(x)^4 + 9*co...

```

## Sympy [F]

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}^5(x)}{\tanh(x) + 1} dx$$

input

```
integrate(csch(x)**5/(1+tanh(x)),x)
```

output

```
Integral(csch(x)**5/(tanh(x) + 1), x)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 74 vs.  $2(26) = 52$ .

Time = 0.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.18

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = \frac{3e^{(-x)} - 11e^{(-3x)} + 53e^{(-5x)} + 3e^{(-7x)}}{12(4e^{(-2x)} - 6e^{(-4x)} + 4e^{(-6x)} - e^{(-8x)} - 1)} + \frac{1}{8} \log(e^{(-x)} + 1) - \frac{1}{8} \log(e^{(-x)} - 1)$$

input `integrate(csch(x)^5/(1+tanh(x)),x, algorithm="maxima")`

output `1/12*(3*e^(-x) - 11*e^(-3*x) + 53*e^(-5*x) + 3*e^(-7*x))/(4*e^(-2*x) - 6*e^(-4*x) + 4*e^(-6*x) - e^(-8*x) - 1) + 1/8*log(e^(-x) + 1) - 1/8*log(e^(-x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = -\frac{3e^{(7x)} - 11e^{(5x)} + 53e^{(3x)} + 3e^x}{12(e^{(2x)} - 1)^4} + \frac{1}{8} \log(e^x + 1) - \frac{1}{8} \log(|e^x - 1|)$$

input `integrate(csch(x)^5/(1+tanh(x)),x, algorithm="giac")`

output `-1/12*(3*e^(7*x) - 11*e^(5*x) + 53*e^(3*x) + 3*e^x)/(e^(2*x) - 1)^4 + 1/8*log(e^x + 1) - 1/8*log(abs(e^x - 1))`

**Mupad [B] (verification not implemented)**

Time = 2.06 (sec) , antiderivative size = 117, normalized size of antiderivative = 3.44

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = \frac{\ln\left(\frac{e^x}{4} + \frac{1}{4}\right)}{8} - \frac{\ln\left(\frac{e^x}{4} - \frac{1}{4}\right)}{8} - \frac{e^x}{4(e^{2x} - 1)} - \frac{2e^{3x} + 2e^x}{6e^{4x} - 4e^{2x} - 4e^{6x} + e^{8x} + 1} - \frac{4e^x}{3(3e^{2x} - 3e^{4x} + e^{6x} - 1)} + \frac{e^x}{6(e^{4x} - 2e^{2x} + 1)}$$

input `int(1/(sinh(x)^5*(tanh(x) + 1)),x)`output `log(exp(x)/4 + 1/4)/8 - log(exp(x)/4 - 1/4)/8 - exp(x)/(4*(exp(2*x) - 1)) - (2*exp(3*x) + 2*exp(x))/(6*exp(4*x) - 4*exp(2*x) - 4*exp(6*x) + exp(8*x) + 1) - (4*exp(x))/(3*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) + exp(x)/(6*(exp(4*x) - 2*exp(2*x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 180, normalized size of antiderivative = 5.29

$$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx = \frac{-3e^{8x}\log(e^x - 1) + 3e^{8x}\log(e^x + 1) - 6e^{7x} + 12e^{6x}\log(e^x - 1) - 12e^{6x}\log(e^x + 1) + 22e^{5x} - 18e^{4x}\log(e^x - 1) + 18e^{4x}\log(e^x + 1) - 106e^{3x} + 12e^{2x}\log(e^x - 1) - 12e^{2x}\log(e^x + 1) - 6e^{2x} - 3\log(e^x - 1) + 3\log(e^x + 1)}{24e^{8x} - 96e^{6x}}$$

input `int(csch(x)^5/(1+tanh(x)),x)`output `( - 3*e**(8*x)*log(e**x - 1) + 3*e**(8*x)*log(e**x + 1) - 6*e**(7*x) + 12*e**(6*x)*log(e**x - 1) - 12*e**(6*x)*log(e**x + 1) + 22*e**(5*x) - 18*e**(4*x)*log(e**x - 1) + 18*e**(4*x)*log(e**x + 1) - 106*e**(3*x) + 12*e**(2*x)*log(e**x - 1) - 12*e**(2*x)*log(e**x + 1) - 6*e**x - 3*log(e**x - 1) + 3*log(e**x + 1))/(24*(e**(8*x) - 4*e**(6*x) + 6*e**(4*x) - 4*e**(2*x) + 1))`



### 3.78 $\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx$

Optimal result	640
Mathematica [A] (verified)	640
Rubi [A] (verified)	641
Maple [A] (verified)	642
Fricas [B] (verification not implemented)	643
Sympy [F]	643
Maxima [B] (verification not implemented)	644
Giac [A] (verification not implemented)	644
Mupad [B] (verification not implemented)	645
Reduce [B] (verification not implemented)	645

#### Optimal result

Integrand size = 11, antiderivative size = 33

$$\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx = -\frac{1}{2} \operatorname{coth}^2(x) + \frac{\operatorname{coth}^3(x)}{3} + \frac{\operatorname{coth}^4(x)}{4} - \frac{\operatorname{coth}^5(x)}{5}$$

output `-1/2*coth(x)^2+1/3*coth(x)^3+1/4*coth(x)^4-1/5*coth(x)^5`

#### Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx = \frac{1}{120} \operatorname{csch}^5(x) (-20 \cosh(x) - 5 \cosh(3x) + \cosh(5x) + 30 \sinh(x))$$

input `Integrate[Csch[x]^6/(1 + Tanh[x]), x]`

output `(Csch[x]^5*(-20*Cosh[x] - 5*Cosh[3*x] + Cosh[5*x] + 30*Sinh[x]))/120`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 25, 3999, 516, 84, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^6(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\sin(ix)^6(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\sin(ix)^6(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{3999} \\
 & \int \frac{(1 - \tanh^2(x))^2 \operatorname{coth}^6(x)}{\tanh(x) + 1} d \tanh(x) \\
 & \quad \downarrow \text{516} \\
 & \int (1 - \tanh(x))^2 (\tanh(x) + 1) \operatorname{coth}^6(x) d \tanh(x) \\
 & \quad \downarrow \text{84} \\
 & \int (\operatorname{coth}^6(x) - \operatorname{coth}^5(x) - \operatorname{coth}^4(x) + \operatorname{coth}^3(x)) d \tanh(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{5} \operatorname{coth}^5(x) + \frac{\operatorname{coth}^4(x)}{4} + \frac{\operatorname{coth}^3(x)}{3} - \frac{\operatorname{coth}^2(x)}{2}
 \end{aligned}$$

input `Int [Csch[x]^6/(1 + Tanh[x]), x]`

output `-1/2*Coth[x]^2 + Coth[x]^3/3 + Coth[x]^4/4 - Coth[x]^5/5`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 84 `Int[((d_)*(x_))^(n_)*((a_) + (b_)*(x_))*((e_) + (f_)*(x_))^(p_), x_] :> Int[ExpandIntegrand[(a + b*x)*(d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, d, e, f, n}, x] && IGtQ[p, 0] && EqQ[b*e + a*f, 0] && !(ILtQ[n + p + 2, 0] && GtQ[n + 2*p, 0])`
- rule 516 `Int[((e_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[(e*x)^m*(c + d*x)^(n + p)*(a/c + (b/d)*x)^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && EqQ[b*c^2 + a*d^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0] && !IntegerQ[n]))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3999 `Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 4.46 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

method	result
risch	$-\frac{4(20e^{4x} + 5e^{2x} - 1)}{15(e^{2x} - 1)^5}$
parallelrisc	$\frac{\operatorname{sech}\left(\frac{x}{2}\right)^5 \operatorname{csch}\left(\frac{x}{2}\right)^5 (64 \cosh(5x) - 320 \cosh(3x) - 1280 \cosh(x) + 330 \sinh(x) - 159 \sinh(5x) + 795 \sinh(3x))}{245760}$
default	$-\frac{\tanh\left(\frac{x}{2}\right)^5}{160} + \frac{\tanh\left(\frac{x}{2}\right)^4}{64} + \frac{\tanh\left(\frac{x}{2}\right)^3}{96} - \frac{\tanh\left(\frac{x}{2}\right)^2}{16} + \frac{\tanh\left(\frac{x}{2}\right)}{16} + \frac{1}{64 \tanh\left(\frac{x}{2}\right)^4} - \frac{1}{160 \tanh\left(\frac{x}{2}\right)^5} + \frac{1}{96 \tanh\left(\frac{x}{2}\right)^3}$

input `int(csch(x)^6/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-4/15*(20*exp(4*x)+5*exp(2*x)-1)/(exp(2*x)-1)^5`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs.  $2(25) = 50$ .

Time = 0.08 (sec) , antiderivative size = 185, normalized size of antiderivative = 5.61

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx =$$

$$-\frac{15 (\cosh(x)^8 + 8 \cosh(x) \sinh(x)^7 + \sinh(x)^8 + (28 \cosh(x)^2 - 5) \sinh(x)^6 - 5 \cosh(x)^6 + 2 (28 \cosh(x)^4 - 15 \cosh(x)^2) \sinh(x)^4 + 10 \cosh(x)^4 + 4 (14 \cosh(x)^5 - 25 \cosh(x)^3 + 10 \cosh(x) \sinh(x))^2 + 2 (28 \cosh(x)^6 - 75 \cosh(x)^4 + 60 \cosh(x)^2 - 11) \sinh(x)^2 - 11 \cosh(x)^2 + 2 (4 \cosh(x)^7 - 15 \cosh(x)^5 + 20 \cosh(x)^3 - 9 \cosh(x) \sinh(x))^2 + 5)}{15 (\cosh(x)^8 + 8 \cosh(x) \sinh(x)^7 + \sinh(x)^8 + (28 \cosh(x)^2 - 5) \sinh(x)^6 - 5 \cosh(x)^6 + 2 (28 \cosh(x)^4 - 15 \cosh(x)^2) \sinh(x)^4 + 10 \cosh(x)^4 + 4 (14 \cosh(x)^5 - 25 \cosh(x)^3 + 10 \cosh(x) \sinh(x))^2 + 2 (28 \cosh(x)^6 - 75 \cosh(x)^4 + 60 \cosh(x)^2 - 11) \sinh(x)^2 - 11 \cosh(x)^2 + 2 (4 \cosh(x)^7 - 15 \cosh(x)^5 + 20 \cosh(x)^3 - 9 \cosh(x) \sinh(x))^2 + 5)}$$

input `integrate(csch(x)^6/(1+tanh(x)),x, algorithm="fricas")`

output `-4/15*(19*cosh(x)^2 + 42*cosh(x)*sinh(x) + 19*sinh(x)^2 + 5)/(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 - 5)*sinh(x)^6 - 5*cosh(x)^6 + 2*(28*cosh(x)^3 - 15*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 - 15*cosh(x)^2 + 2)*sinh(x)^4 + 10*cosh(x)^4 + 4*(14*cosh(x)^5 - 25*cosh(x)^3 + 10*cosh(x)*sinh(x))^2 + 2*(28*cosh(x)^6 - 75*cosh(x)^4 + 60*cosh(x)^2 - 11)*sinh(x)^2 - 11*cosh(x)^2 + 2*(4*cosh(x)^7 - 15*cosh(x)^5 + 20*cosh(x)^3 - 9*cosh(x)*sinh(x))^2 + 5)`

### Sympy [F]

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}^6(x)}{\tanh(x) + 1} dx$$

input `integrate(csch(x)**6/(1+tanh(x)),x)`

output `Integral(csch(x)**6/(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 149 vs.  $2(25) = 50$ .

Time = 0.03 (sec) , antiderivative size = 149, normalized size of antiderivative = 4.52

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx = \frac{4e^{-2x}}{3(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1)} - \frac{8e^{-4x}}{3(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1)} + \frac{8e^{-6x}}{5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1} - \frac{4}{15(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1)}$$

input `integrate(csch(x)^6/(1+tanh(x)),x, algorithm="maxima")`

output `4/3*e^(-2*x)/(5*e^(-2*x) - 10*e^(-4*x) + 10*e^(-6*x) - 5*e^(-8*x) + e^(-10*x) - 1) - 8/3*e^(-4*x)/(5*e^(-2*x) - 10*e^(-4*x) + 10*e^(-6*x) - 5*e^(-8*x) + e^(-10*x) - 1) + 8*e^(-6*x)/(5*e^(-2*x) - 10*e^(-4*x) + 10*e^(-6*x) - 5*e^(-8*x) + e^(-10*x) - 1) - 4/15/(5*e^(-2*x) - 10*e^(-4*x) + 10*e^(-6*x) - 5*e^(-8*x) + e^(-10*x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx = -\frac{4(20e^{4x} + 5e^{2x} - 1)}{15(e^{2x} - 1)^5}$$

input `integrate(csch(x)^6/(1+tanh(x)),x, algorithm="giac")`

output `-4/15*(20*e^(4*x) + 5*e^(2*x) - 1)/(e^(2*x) - 1)^5`

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx = -\frac{4(5e^{2x} + 20e^{4x} - 1)}{15(e^{2x} - 1)^5}$$

input `int(1/(sinh(x)^6*(tanh(x) + 1)),x)`output `-(4*(5*exp(2*x) + 20*exp(4*x) - 1))/(15*(exp(2*x) - 1)^5)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.73

$$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx = \frac{-80e^{4x} - 20e^{2x} + 4}{15e^{10x} - 75e^{8x} + 150e^{6x} - 150e^{4x} + 75e^{2x} - 15}$$

input `int(csch(x)^6/(1+tanh(x)),x)`output `(4*( - 20*e**(4*x) - 5*e**(2*x) + 1))/(15*(e**(10*x) - 5*e**(8*x) + 10*e**(6*x) - 10*e**(4*x) + 5*e**(2*x) - 1))`

### 3.79 $\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx$

Optimal result	646
Mathematica [B] (verified)	646
Rubi [C] (verified)	647
Maple [A] (verified)	649
Fricas [B] (verification not implemented)	650
Sympy [F]	651
Maxima [B] (verification not implemented)	651
Giac [A] (verification not implemented)	652
Mupad [B] (verification not implemented)	652
Reduce [B] (verification not implemented)	653

#### Optimal result

Integrand size = 11, antiderivative size = 44

$$\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx = -\frac{1}{16}\operatorname{arctanh}(\cosh(x)) + \frac{1}{16}\operatorname{coth}(x)\operatorname{csch}(x) - \frac{1}{24}\operatorname{coth}(x)\operatorname{csch}^3(x) + \frac{\operatorname{csch}^5(x)}{5} - \frac{1}{6}\operatorname{coth}(x)\operatorname{csch}^5(x)$$

output

```
-1/16*arctanh(cosh(x))+1/16*coth(x)*csch(x)-1/24*coth(x)*csch(x)^3+1/5*csc
h(x)^5-1/6*coth(x)*csch(x)^5
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 124 vs. 2(44) = 88.

Time = 0.36 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.82

$$\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx = \frac{72 \operatorname{coth}\left(\frac{x}{2}\right) + 30 \operatorname{csch}^2\left(\frac{x}{2}\right) - 120 \log\left(\cosh\left(\frac{x}{2}\right)\right) + 120 \log\left(\sinh\left(\frac{x}{2}\right)\right) + 30 \operatorname{sech}^2\left(\frac{x}{2}\right) - 5 \operatorname{sech}^6\left(\frac{x}{2}\right) - 288 \operatorname{csch}^2\left(\frac{x}{2}\right)}{1}$$

input

```
Integrate[Csch[x]^7/(1 + Tanh[x]), x]
```

output

```
(72*Coth[x/2] + 30*Csch[x/2]^2 - 120*Log[Cosh[x/2]] + 120*Log[Sinh[x/2]] +
 30*Sech[x/2]^2 - 5*Sech[x/2]^6 - 288*Csch[x]^3*Sinh[x/2]^4 - 384*Csch[x]^
5*Sinh[x/2]^6 - 18*Csch[x/2]^4*Sinh[x] + Csch[x/2]^6*(-5 + 6*Sinh[x]) - 72
*Tanh[x/2])/1920
```

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.32, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3587, 3042, 26, 3586, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^7(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i}{\sin(ix)^7(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{1}{\sin(ix)^7(1 - i \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & -i \int \frac{i \coth(x) \operatorname{csch}^6(x)}{\cosh(x) + \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x) \operatorname{csch}^6(x)}{\sinh(x) + \cosh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \cos(ix)}{\sin(ix)^7(\cos(ix) - i \sin(ix))} dx \\
 & \quad \downarrow \text{26}
 \end{aligned}$$



$$\begin{aligned}
& -i \int \frac{\cos(ix)}{(\cos(ix) - i \sin(ix)) \sin(ix)^7} dx \\
& \quad \downarrow \text{3587} \\
& \int \coth(x) \operatorname{csch}^6(x) (\cosh(x) - \sinh(x)) dx \\
& \quad \downarrow \text{3042} \\
& \int -\frac{i \cos(ix) (i \sin(ix) + \cos(ix))}{\sin(ix)^7} dx \\
& \quad \downarrow \text{26} \\
& -i \int \frac{\cos(ix) (\cos(ix) + i \sin(ix))}{\sin(ix)^7} dx \\
& \quad \downarrow \text{3586} \\
& -i \int (i \coth^2(x) \operatorname{csch}^5(x) - i \coth(x) \operatorname{csch}^5(x)) dx \\
& \quad \downarrow \text{2009} \\
& -i \left( -\frac{1}{16} i \operatorname{arctanh}(\cosh(x)) + \frac{1}{5} i \operatorname{csch}^5(x) - \frac{1}{6} i \coth(x) \operatorname{csch}^5(x) - \frac{1}{24} i \coth(x) \operatorname{csch}^3(x) + \frac{1}{16} i \coth(x) \operatorname{csch}(x) \right)
\end{aligned}$$

input `Int[Csch[x]^7/(1 + Tanh[x]), x]`

output `(-I)*((-1/16*I)*ArcTanh[Cosh[x]] + (I/16)*Coth[x]*Csch[x] - (I/24)*Coth[x]*Csch[x]^3 + (I/5)*Csch[x]^5 - (I/6)*Coth[x]*Csch[x]^5)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3586 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]`

rule 3587 `Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(p_), x_Symbol] := Simp[a^p*b^p Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

## Maple [A] (verified)

Time = 8.69 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.36

method	result
risch	$\frac{e^x(15e^{10x} - 85e^{8x} + 198e^{6x} - 1338e^{4x} - 85e^{2x} + 15)}{120(e^{2x} - 1)^6} - \frac{\ln(e^x + 1)}{16} + \frac{\ln(e^x - 1)}{16}$
default	$\frac{\tanh(\frac{x}{2})^6}{384} - \frac{\tanh(\frac{x}{2})^5}{160} - \frac{\tanh(\frac{x}{2})^4}{128} + \frac{\tanh(\frac{x}{2})^3}{32} - \frac{\tanh(\frac{x}{2})^2}{128} - \frac{\tanh(\frac{x}{2})}{16} - \frac{1}{384 \tanh(\frac{x}{2})^6} + \frac{1}{16 \tanh(\frac{x}{2})} + \frac{1}{160 \tanh(\frac{x}{2})}$

input `int(csch(x)^7/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{120} \exp(x) (15 \exp(10x) - 85 \exp(8x) + 198 \exp(6x) - 1338 \exp(4x) - 85 \exp(2x) + 15) / (\exp(2x) - 1)^6 - 1/16 * \ln(\exp(x) + 1) + 1/16 * \ln(\exp(x) - 1)$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1260 vs.  $2(34) = 68$ .

Time = 0.10 (sec) , antiderivative size = 1260, normalized size of antiderivative = 28.64

$$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(csch(x)^7/(1+tanh(x)),x, algorithm="fricas")`

output

```
1/240*(30*cosh(x)^11 + 330*cosh(x)*sinh(x)^10 + 30*sinh(x)^11 + 10*(165*cosh(x)^2 - 17)*sinh(x)^9 - 170*cosh(x)^9 + 90*(55*cosh(x)^3 - 17*cosh(x))*sinh(x)^8 + 36*(275*cosh(x)^4 - 170*cosh(x)^2 + 11)*sinh(x)^7 + 396*cosh(x)^7 + 84*(165*cosh(x)^5 - 170*cosh(x)^3 + 33*cosh(x))*sinh(x)^6 + 12*(1155*cosh(x)^6 - 1785*cosh(x)^4 + 693*cosh(x)^2 - 223)*sinh(x)^5 - 2676*cosh(x)^5 + 60*(165*cosh(x)^7 - 357*cosh(x)^5 + 231*cosh(x)^3 - 223*cosh(x))*sinh(x)^4 + 10*(495*cosh(x)^8 - 1428*cosh(x)^6 + 1386*cosh(x)^4 - 2676*cosh(x)^2 - 17)*sinh(x)^3 - 170*cosh(x)^3 + 6*(275*cosh(x)^9 - 1020*cosh(x)^7 + 1386*cosh(x)^5 - 4460*cosh(x)^3 - 85*cosh(x))*sinh(x)^2 - 15*(cosh(x)^12 + 12*cosh(x)*sinh(x)^11 + sinh(x)^12 + 6*(11*cosh(x)^2 - 1)*sinh(x)^10 - 6*cosh(x)^10 + 20*(11*cosh(x)^3 - 3*cosh(x))*sinh(x)^9 + 15*(33*cosh(x)^4 - 18*cosh(x)^2 + 1)*sinh(x)^8 + 15*cosh(x)^8 + 24*(33*cosh(x)^5 - 30*cosh(x)^3 + 5*cosh(x))*sinh(x)^7 + 4*(231*cosh(x)^6 - 315*cosh(x)^4 + 105*cosh(x)^2 - 5)*sinh(x)^6 - 20*cosh(x)^6 + 24*(33*cosh(x)^7 - 63*cosh(x)^5 + 35*cosh(x)^3 - 5*cosh(x))*sinh(x)^5 + 15*(33*cosh(x)^8 - 84*cosh(x)^6 + 70*cosh(x)^4 - 20*cosh(x)^2 + 1)*sinh(x)^4 + 15*cosh(x)^4 + 20*(11*cosh(x)^9 - 36*cosh(x)^7 + 42*cosh(x)^5 - 20*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 6*(11*cosh(x)^10 - 45*cosh(x)^8 + 70*cosh(x)^6 - 50*cosh(x)^4 + 15*cosh(x)^2 - 1)*sinh(x)^2 - 6*cosh(x)^2 + 12*(cosh(x)^11 - 5*cosh(x)^9 + 10*cosh(x)^7 - 10*cosh(x)^5 + 5*cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) + ...
```

**Sympy [F]**

$$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{csch}^7(x)}{\tanh(x) + 1} dx$$

input `integrate(csch(x)**7/(1+tanh(x)),x)`

output `Integral(csch(x)**7/(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 98 vs.  $2(34) = 68$ .

Time = 0.04 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.23

$$\begin{aligned} & \int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx \\ &= -\frac{15e^{-x} - 85e^{-3x} + 198e^{-5x} - 1338e^{-7x} - 85e^{-9x} + 15e^{-11x}}{120(6e^{-2x} - 15e^{-4x} + 20e^{-6x} - 15e^{-8x} + 6e^{-10x} - e^{-12x} - 1)} \\ & \quad - \frac{1}{16} \log(e^{-x} + 1) + \frac{1}{16} \log(e^{-x} - 1) \end{aligned}$$

input `integrate(csch(x)^7/(1+tanh(x)),x, algorithm="maxima")`

output `-1/120*(15*e^(-x) - 85*e^(-3*x) + 198*e^(-5*x) - 1338*e^(-7*x) - 85*e^(-9*x) + 15*e^(-11*x))/(6*e^(-2*x) - 15*e^(-4*x) + 20*e^(-6*x) - 15*e^(-8*x) + 6*e^(-10*x) - e^(-12*x) - 1) - 1/16*log(e^(-x) + 1) + 1/16*log(e^(-x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.61 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.39

$$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx = \frac{15e^{(11x)} - 85e^{(9x)} + 198e^{(7x)} - 1338e^{(5x)} - 85e^{(3x)} + 15e^x}{120(e^{(2x)} - 1)^6} - \frac{1}{16} \log(e^x + 1) + \frac{1}{16} \log(|e^x - 1|)$$

input `integrate(csch(x)^7/(1+tanh(x)),x, algorithm="giac")`output `1/120*(15*e^(11*x) - 85*e^(9*x) + 198*e^(7*x) - 1338*e^(5*x) - 85*e^(3*x) + 15*e^x)/(e^(2*x) - 1)^6 - 1/16*log(e^x + 1) + 1/16*log(abs(e^x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 207, normalized size of antiderivative = 4.70

$$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx = \frac{\ln\left(\frac{1}{8} - \frac{e^x}{8}\right)}{16} - \frac{\ln\left(-\frac{e^x}{8} - \frac{1}{8}\right)}{16} - \frac{\frac{16e^{3x}}{3} + \frac{16e^{5x}}{3}}{15e^{4x} - 6e^{2x} - 20e^{6x} + 15e^{8x} - 6e^{10x} + e^{12x} + 1} - \frac{\frac{8e^{3x}}{3} + \frac{8e^x}{5}}{5e^{2x} - 10e^{4x} + 10e^{6x} - 5e^{8x} + e^{10x} - 1} - \frac{e^x}{6e^x} - \frac{e^x}{5(6e^{4x} - 4e^{2x} - 4e^{6x} + e^{8x} + 1)} + \frac{e^x}{8(e^{2x} - 1)} + \frac{e^x}{15(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{e^x}{12(e^{4x} - 2e^{2x} + 1)}$$

input `int(1/(sinh(x)^7*(tanh(x) + 1)),x)`

output

```
log(1/8 - exp(x)/8)/16 - log(- exp(x)/8 - 1/8)/16 - ((16*exp(3*x))/3 + (16
*exp(5*x))/3)/(15*exp(4*x) - 6*exp(2*x) - 20*exp(6*x) + 15*exp(8*x) - 6*ex
p(10*x) + exp(12*x) + 1) - ((8*exp(3*x))/3 + (8*exp(x))/5)/(5*exp(2*x) - 1
0*exp(4*x) + 10*exp(6*x) - 5*exp(8*x) + exp(10*x) - 1) - (6*exp(x))/(5*(6*
exp(4*x) - 4*exp(2*x) - 4*exp(6*x) + exp(8*x) + 1)) + exp(x)/(8*(exp(2*x)
- 1)) + exp(x)/(15*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) - exp(x)/(12*
(exp(4*x) - 2*exp(2*x) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 260, normalized size of antiderivative = 5.91

$$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx$$

$$= \frac{15e^{12x}\log(e^x - 1) - 15e^{12x}\log(e^x + 1) + 30e^{11x} - 90e^{10x}\log(e^x - 1) + 90e^{10x}\log(e^x + 1) - 170e^{9x} + 225e^{8x}\log(e^x - 1) - 225e^{8x}\log(e^x + 1) + 396e^{7x} - 300e^{6x}\log(e^x - 1) + 300e^{6x}\log(e^x + 1) - 2676e^{5x} + 225e^{4x}\log(e^x - 1) - 225e^{4x}\log(e^x + 1) - 170e^{3x} - 90e^{2x}\log(e^x - 1) + 90e^{2x}\log(e^x + 1) + 30e^{2x} + 15\log(e^x - 1) - 15\log(e^x + 1)}{(240*(e^{12x} - 6e^{10x} + 15e^{8x} - 20e^{6x} + 15e^{4x} - 6e^{2x} + 1))}$$

input

```
int(csch(x)^7/(1+tanh(x)),x)
```

output

```
(15*e**(12*x)*log(e**x - 1) - 15*e**(12*x)*log(e**x + 1) + 30*e**(11*x) -
90*e**(10*x)*log(e**x - 1) + 90*e**(10*x)*log(e**x + 1) - 170*e**(9*x) + 2
25*e**(8*x)*log(e**x - 1) - 225*e**(8*x)*log(e**x + 1) + 396*e**(7*x) - 30
0*e**(6*x)*log(e**x - 1) + 300*e**(6*x)*log(e**x + 1) - 2676*e**(5*x) + 22
5*e**(4*x)*log(e**x - 1) - 225*e**(4*x)*log(e**x + 1) - 170*e**(3*x) - 90*
e**(2*x)*log(e**x - 1) + 90*e**(2*x)*log(e**x + 1) + 30*e**x + 15*log(e**x
- 1) - 15*log(e**x + 1))/(240*(e**(12*x) - 6*e**(10*x) + 15*e**(8*x) - 20
*e**(6*x) + 15*e**(4*x) - 6*e**(2*x) + 1))
```

### 3.80 $\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx$

Optimal result	654
Mathematica [A] (verified)	654
Rubi [A] (verified)	655
Maple [A] (verified)	658
Fricas [B] (verification not implemented)	659
Sympy [F]	660
Maxima [A] (verification not implemented)	660
Giac [A] (verification not implemented)	661
Mupad [B] (verification not implemented)	661
Reduce [B] (verification not implemented)	662

#### Optimal result

Integrand size = 13, antiderivative size = 147

$$\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx = -\frac{a(3a+b) \log(1-\tanh(x))}{16(a+b)^3} + \frac{a(3a-b) \log(1+\tanh(x))}{16(a-b)^3} - \frac{a^4 b \log(a+b \tanh(x))}{(a^2-b^2)^3} - \frac{\cosh^4(x)(b-a \tanh(x))}{4(a^2-b^2)} + \frac{\cosh^2(x)(4b(2a^2-b^2)-a(5a^2-b^2)\tanh(x))}{8(a^2-b^2)^2}$$

output 
$$-1/16*a*(3*a+b)*\ln(1-\tanh(x))/(a+b)^3+1/16*a*(3*a-b)*\ln(1+\tanh(x))/(a-b)^3 -a^4*b*\ln(a+b*\tanh(x))/(a^2-b^2)^3-\cosh(x)^4*(b-a*\tanh(x))/(4*a^2-4*b^2)+1/8*\cosh(x)^2*(4*b*(2*a^2-b^2)-a*(5*a^2-b^2)*\tanh(x))/(a^2-b^2)^2$$

#### Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.98

$$\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx = \frac{12a^5x + 24a^3b^2x - 4ab^4x + 4b(3a^4 - 4a^2b^2 + b^4) \cosh(2x) - b(a^2 - b^2)^2 \cosh(4x) - 32a^4b \log(a \cosh(x))}{32(a-b)^3(a+b)^3}$$

input `Integrate[Sinh[x]^4/(a + b*Tanh[x]),x]`

output  $(12a^5x + 24a^3b^2x - 4ab^4x + 4b(3a^4 - 4a^2b^2 + b^4)\text{Cosh}[2x] - b(a^2 - b^2)^2\text{Cosh}[4x] - 32a^4b\text{Log}[a\text{Cosh}[x] + b\text{Sinh}[x]] - 8a^3(a^2 - b^2)\text{Sinh}[2x] + a^5\text{Sinh}[4x] - 2a^3b^2\text{Sinh}[4x] + ab^4\text{Sinh}[4x]) / (32(a - b)^3(a + b)^3)$

### Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.65, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$ , Rules used = {3042, 3999, 25, 601, 2178, 25, 27, 657, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sinh^4(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\sin(ix)^4}{a - ib \tan(ix)} dx \\ & \quad \downarrow 3999 \\ & -b \int -\frac{b^4 \tanh^4(x)}{(a + b \tanh(x)) (b^2 - b^2 \tanh^2(x))^3} d(b \tanh(x)) \\ & \quad \downarrow 25 \\ & b \int \frac{b^4 \tanh^4(x)}{(a + b \tanh(x)) (b^2 - b^2 \tanh^2(x))^3} d(b \tanh(x)) \\ & \quad \downarrow 601 \\ & -b \left( \frac{\int \frac{-\frac{3a \tanh(x)b^5}{a^2 - b^2} + 4 \tanh^2(x)b^4 + \frac{a^2 b^4}{a^2 - b^2}}{(a + b \tanh(x)) (b^2 - b^2 \tanh^2(x))^2} d(b \tanh(x))}{4b^2} + \frac{b^2 \left( \frac{b^2}{a^2 - b^2} - \frac{ab \tanh(x)}{a^2 - b^2} \right)}{4 (b^2 - b^2 \tanh^2(x))^2} \right) \\ & \quad \downarrow 2178 \end{aligned}$$



$$-b \left( \frac{\int -\frac{ab^4(a(3a^2+b^2)-b(5a^2-b^2)\tanh(x))}{(a^2-b^2)^2(a+b\tanh(x))(b^2-b^2\tanh^2(x))}d(b\tanh(x))}{2b^2} - \frac{b^2(4b^2(2a^2-b^2)-ab(5a^2-b^2)\tanh(x))}{2(a^2-b^2)^2(b^2-b^2\tanh^2(x))} + \frac{b^2\left(\frac{b^2}{a^2-b^2} - \frac{ab\tanh(x)}{a^2-b^2}\right)}{4(b^2-b^2\tanh^2(x))^2} \right)$$

↓ 25

$$-b \left( \frac{\int -\frac{ab^4(a(3a^2+b^2)-b(5a^2-b^2)\tanh(x))}{(a^2-b^2)^2(a+b\tanh(x))(b^2-b^2\tanh^2(x))}d(b\tanh(x))}{2b^2} - \frac{b^2(4b^2(2a^2-b^2)-ab(5a^2-b^2)\tanh(x))}{2(a^2-b^2)^2(b^2-b^2\tanh^2(x))} + \frac{b^2\left(\frac{b^2}{a^2-b^2} - \frac{ab\tanh(x)}{a^2-b^2}\right)}{4(b^2-b^2\tanh^2(x))^2} \right)$$

↓ 27

$$-b \left( \frac{ab^2 \int \frac{a(3a^2+b^2)-b(5a^2-b^2)\tanh(x)}{(a+b\tanh(x))(b^2-b^2\tanh^2(x))}d(b\tanh(x))}{2(a^2-b^2)^2} - \frac{b^2(4b^2(2a^2-b^2)-ab(5a^2-b^2)\tanh(x))}{2(a^2-b^2)^2(b^2-b^2\tanh^2(x))} + \frac{b^2\left(\frac{b^2}{a^2-b^2} - \frac{ab\tanh(x)}{a^2-b^2}\right)}{4(b^2-b^2\tanh^2(x))^2} \right)$$

↓ 657

$$-b \left( \frac{ab^2 \int \left( -\frac{8a^3}{(a-b)(a+b)(a+b\tanh(x))} + \frac{(a-b)^2(3a+b)}{2b(a+b)(b-b\tanh(x))} + \frac{(3a-b)(a+b)^2}{2(a-b)b(\tanh(x)b+b)} \right) d(b\tanh(x))}{2(a^2-b^2)^2} - \frac{b^2(4b^2(2a^2-b^2)-ab(5a^2-b^2)\tanh(x))}{2(a^2-b^2)^2(b^2-b^2\tanh^2(x))} \right)$$

↓ 2009

$$-b \left( \frac{b^2\left(\frac{b^2}{a^2-b^2} - \frac{ab\tanh(x)}{a^2-b^2}\right)}{4(b^2-b^2\tanh^2(x))^2} + \frac{-\frac{b^2(4b^2(2a^2-b^2)-ab(5a^2-b^2)\tanh(x))}{2(a^2-b^2)^2(b^2-b^2\tanh^2(x))} - \frac{ab^2\left(-\frac{8a^3\log(a+b\tanh(x))}{a^2-b^2} - \frac{(a-b)^2(3a+b)\log(b-b\tanh(x))}{2b(a+b)}\right)}{2(a^2-b^2)^2}}{4b^2} \right)$$

input `Int[Sinh[x]^4/(a + b*Tanh[x]),x]`

output

```
-(b*((b^2*(b^2/(a^2 - b^2) - (a*b*Tanh[x])/(a^2 - b^2)))/(4*(b^2 - b^2*Tanh[x]^2)^2) + (-1/2*(a*b^2*(-1/2*((a - b)^2*(3*a + b)*Log[b - b*Tanh[x]])/(b*(a + b)) - (8*a^3*Log[a + b*Tanh[x]])/(a^2 - b^2) + ((3*a - b)*(a + b)^2*Log[b + b*Tanh[x]])/(2*(a - b)*b)))/(a^2 - b^2)^2 - (b^2*(4*b^2*(2*a^2 - b^2) - a*b*(5*a^2 - b^2)*Tanh[x]))/(2*(a^2 - b^2)^2*(b^2 - b^2*Tanh[x]^2)))/(4*b^2))
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 601

```
Int[(x_)^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[x^m*(c + d*x)^n, a + b*x^2, x], e = Coeff[PolynomialRemainder[x^m*(c + d*x)^n, a + b*x^2, x], x, 0], f = Coeff[PolynomialRemainder[x^m*(c + d*x)^n, a + b*x^2, x], x, 1]}, Simp[(a*f - b*e*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1)) Int[(c + d*x)^n*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*(p + 1)*Qx)/(c + d*x)^n + (e*(2*p + 3))/(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 1] && LtQ[p, -1] && ILtQ[n, 0] && NeQ[b*c^2 + a*d^2, 0]
```

rule 657

```
Int[(((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_))/((a_) + (c_)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g, m}, x] && IntegersQ[n]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2178

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :
> With[{Qx = PolynomialQuotient[(d + e*x)^m*Pq, a + b*x^2, x], R = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 0], S = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x^2, x], x, 1]}, Simp[(a*S - b*R*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*b*(p + 1)) Int[(d + e*x)^m*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*b*(p + 1)*Qx)/(d + e*x)^m + (b*R*(2*p + 3))/(d + e*x)^m, x], x], x] /; FreeQ[{a, b, d, e}, x] && PolyQ[Pq, x] && NeQ[b*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3999

```
Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :=> Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]
```

## Maple [A] (verified)

Time = 12.17 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.24

method	result
risch	$\frac{3a^2x}{8(a+b)^3} + \frac{axb}{8(a+b)^3} + \frac{e^{4x}}{64a+64b} - \frac{e^{2x}a}{8(a+b)^2} - \frac{e^{2x}b}{16(a+b)^2} + \frac{e^{-2x}a}{8(a-b)^2} - \frac{e^{-2x}b}{16(a-b)^2} - \frac{e^{-4x}}{64(a-b)} + \frac{2a^4bx}{a^6-3a^4b^2+3a^2b^4-b^6}$
default	$-\frac{8}{(32a-32b)(1+\tanh(\frac{x}{2}))^4} + \frac{32}{(64a-64b)(1+\tanh(\frac{x}{2}))^3} - \frac{-a-b}{8(a-b)^2(1+\tanh(\frac{x}{2}))^2} - \frac{3a-b}{8(a-b)^2(1+\tanh(\frac{x}{2}))} + \frac{a(3a-b)\ln(\dots)}{8(a-b)}$

input

```
int(sinh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
3/8*a^2*x/(a+b)^3+1/8*a*x/(a+b)^3*b+1/64/(a+b)*exp(4*x)-1/8/(a+b)^2*exp(2*x)*a-1/16/(a+b)^2*exp(2*x)*b+1/8/(a-b)^2*exp(-2*x)*a-1/16/(a-b)^2*exp(-2*x)*b-1/64/(a-b)*exp(-4*x)+2*a^4*b/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*x-a^4*b/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*ln(exp(2*x)+(a-b)/(a+b))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1226 vs.  $2(139) = 278$ .

Time = 0.11 (sec) , antiderivative size = 1226, normalized size of antiderivative = 8.34

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

output

```
1/64*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^8 + 8*(a
^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^7 + (a^5
- a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^8 - 4*(2*a^5 - 3*a
^4*b - 2*a^3*b^2 + 4*a^2*b^3 - b^5)*cosh(x)^6 - 4*(2*a^5 - 3*a^4*b - 2*a^3
*b^2 + 4*a^2*b^3 - b^5 - 7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 -
b^5)*cosh(x)^2)*sinh(x)^6 + 8*(3*a^5 + 8*a^4*b + 6*a^3*b^2 - a*b^4)*x*cosh
(x)^4 + 8*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^3
- 3*(2*a^5 - 3*a^4*b - 2*a^3*b^2 + 4*a^2*b^3 - b^5)*cosh(x))*sinh(x)^5 -
a^5 - a^4*b + 2*a^3*b^2 + 2*a^2*b^3 - a*b^4 - b^5 + 2*(35*(a^5 - a^4*b - 2
*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 - 30*(2*a^5 - 3*a^4*b - 2*a^
3*b^2 + 4*a^2*b^3 - b^5)*cosh(x)^2 + 4*(3*a^5 + 8*a^4*b + 6*a^3*b^2 - a*b^
4)*x)*sinh(x)^4 + 8*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)
*cosh(x)^5 - 10*(2*a^5 - 3*a^4*b - 2*a^3*b^2 + 4*a^2*b^3 - b^5)*cosh(x)^3
+ 4*(3*a^5 + 8*a^4*b + 6*a^3*b^2 - a*b^4)*x*cosh(x))*sinh(x)^3 + 4*(2*a^5
+ 3*a^4*b - 2*a^3*b^2 - 4*a^2*b^3 + b^5)*cosh(x)^2 + 4*(7*(a^5 - a^4*b - 2
*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 + 2*a^5 + 3*a^4*b - 2*a^3*b^
2 - 4*a^2*b^3 + b^5 - 15*(2*a^5 - 3*a^4*b - 2*a^3*b^2 + 4*a^2*b^3 - b^5)*c
osh(x)^4 + 12*(3*a^5 + 8*a^4*b + 6*a^3*b^2 - a*b^4)*x*cosh(x)^2)*sinh(x)^2
- 64*(a^4*b*cosh(x)^4 + 4*a^4*b*cosh(x)^3*sinh(x) + 6*a^4*b*cosh(x)^2*si
nh(x)^2 + 4*a^4*b*cosh(x)*sinh(x)^3 + a^4*b*sinh(x)^4)*log(2*(a*cosh(x) ...
```

**Sympy [F]**

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx = \int \frac{\sinh^4(x)}{a + b \tanh(x)} dx$$

input `integrate(sinh(x)**4/(a+b*tanh(x)),x)`

output `Integral(sinh(x)**4/(a + b*tanh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.11

$$\begin{aligned} \int \frac{\sinh^4(x)}{a + b \tanh(x)} dx = & -\frac{a^4 b \log(-(a-b)e^{-2x} - a - b)}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} \\ & + \frac{(3a^2 + ab)x}{8(a^3 + 3a^2 b + 3ab^2 + b^3)} - \frac{(4(2a+b)e^{-2x} - a - b)e^{4x}}{64(a^2 + 2ab + b^2)} \\ & + \frac{4(2a-b)e^{-2x} - (a-b)e^{-4x}}{64(a^2 - 2ab + b^2)} \end{aligned}$$

input `integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

output `-a^4*b*log(-(a - b)*e^(-2*x) - a - b)/(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) + 1/8*(3*a^2 + a*b)*x/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 1/64*(4*(2*a + b)*e^(-2*x) - a - b)*e^(4*x)/(a^2 + 2*a*b + b^2) + 1/64*(4*(2*a - b)*e^(-2*x) - (a - b)*e^(-4*x))/(a^2 - 2*a*b + b^2)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.46

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx$$

$$= -\frac{a^4 b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{(3a^2 - ab)x}{8(a^3 - 3a^2 b + 3ab^2 - b^3)}$$

$$- \frac{(18a^2 e^{(4x)} - 6abe^{(4x)} - 8a^2 e^{(2x)} + 12abe^{(2x)} - 4b^2 e^{(2x)} + a^2 - 2ab + b^2)e^{(-4x)}}{64(a^3 - 3a^2 b + 3ab^2 - b^3)}$$

$$+ \frac{ae^{(4x)} + be^{(4x)} - 8ae^{(2x)} - 4be^{(2x)}}{64(a^2 + 2ab + b^2)}$$

input `integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="giac")`output `-a^4*b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) + 1/8*(3*a^2 - a*b)*x/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) - 1/64*(18*a^2*e^(4*x) - 6*a*b*e^(4*x) - 8*a^2*e^(2*x) + 12*a*b*e^(2*x) - 4*b^2*e^(2*x) + a^2 - 2*a*b + b^2)*e^(-4*x)/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) + 1/64*(a*e^(4*x) + b*e^(4*x) - 8*a*e^(2*x) - 4*b*e^(2*x))/(a^2 + 2*a*b + b^2)`**Mupad [B] (verification not implemented)**

Time = 2.51 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.92

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx = \frac{e^{4x}}{64a + 64b} - \frac{e^{-4x}}{64a - 64b} + \frac{e^{-2x}(2a - b)}{16(a - b)^2} - \frac{e^{2x}(2a + b)}{16(a + b)^2}$$

$$- \frac{a^4 b \ln(a - b + ae^{2x} + be^{2x})}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{ax(3a - b)}{8(a - b)^3}$$

input `int(sinh(x)^4/(a + b*tanh(x)),x)`output `exp(4*x)/(64*a + 64*b) - exp(-4*x)/(64*a - 64*b) + (exp(-2*x)*(2*a - b))/(16*(a - b)^2) - (exp(2*x)*(2*a + b))/(16*(a + b)^2) - (a^4*b*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2) + (a*x*(3*a - b))/(8*(a - b)^3)`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.28

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx$$

$$= \frac{2e^{8x}a^2b^3 - e^{8x}b^5 + 12e^{2x}a^4b - 16e^{2x}a^2b^3 - e^{8x}a^4b - 64e^{4x}\log(e^{2x}a + e^{2x}b + a - b)a^4b + 64e^{4x}a^4bx + 48e^{4x}a^4b^2 - 48e^{4x}a^4b^3 - 48e^{4x}a^4b^4 + 48e^{4x}a^4b^5}{(a + b \tanh(x))^4}$$

input `int(sinh(x)^4/(a+b*tanh(x)),x)`output

```
(e**(8*x)*a**5 - e**(8*x)*a**4*b - 2*e**(8*x)*a**3*b**2 + 2*e**(8*x)*a**2*
b**3 + e**(8*x)*a*b**4 - e**(8*x)*b**5 - 8*e**(6*x)*a**5 + 12*e**(6*x)*a**
4*b + 8*e**(6*x)*a**3*b**2 - 16*e**(6*x)*a**2*b**3 + 4*e**(6*x)*b**5 - 64*
e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**4*b + 24*e**(4*x)*a**5*x
+ 64*e**(4*x)*a**4*b*x + 48*e**(4*x)*a**3*b**2*x - 8*e**(4*x)*a*b**4*x + 8
*e**(2*x)*a**5 + 12*e**(2*x)*a**4*b - 8*e**(2*x)*a**3*b**2 - 16*e**(2*x)*a
**2*b**3 + 4*e**(2*x)*b**5 - a**5 - a**4*b + 2*a**3*b**2 + 2*a**2*b**3 - a
*b**4 - b**5)/(64*e**(4*x)*(a**6 - 3*a**4*b**2 + 3*a**2*b**4 - b**6))
```

### 3.81 $\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx$

Optimal result	663
Mathematica [A] (verified)	663
Rubi [C] (verified)	664
Maple [A] (verified)	669
Fricas [B] (verification not implemented)	670
Sympy [F]	671
Maxima [F(-2)]	671
Giac [A] (verification not implemented)	671
Mupad [B] (verification not implemented)	672
Reduce [B] (verification not implemented)	672

#### Optimal result

Integrand size = 13, antiderivative size = 137

$$\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx = -\frac{a^3 b \arctan\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{5/2}} - \frac{ab^2 \cosh(x)}{(a^2-b^2)^2} - \frac{a \cosh(x)}{a^2-b^2} + \frac{a \cosh^3(x)}{3(a^2-b^2)} + \frac{a^2 b \sinh(x)}{(a^2-b^2)^2} - \frac{b \sinh^3(x)}{3(a^2-b^2)}$$

output `-a^3*b*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))/(a^2-b^2)^(5/2)-a*b^2*cosh(x)/(a^2-b^2)^2-a*cosh(x)/(a^2-b^2)+a*cosh(x)^3/(3*a^2-3*b^2)+a^2*b*sinh(x)/(a^2-b^2)^2-b*sinh(x)^3/(3*a^2-3*b^2)`

#### Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.31

$$\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx = \frac{-3a\sqrt{a-b}\sqrt{a+b}(3a^2+b^2) \cosh(x) + a\sqrt{a-b}\sqrt{a+b}(a^2-b^2) \cosh(3x) + b\left(-24a^3 \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}}\right)\right)}{12(a-b)^{5/2}(a+b)^{5/2}}$$



input `Integrate[Sinh[x]^3/(a + b*Tanh[x]),x]`

output `(-3*a*Sqrt[a - b]*Sqrt[a + b]*(3*a^2 + b^2)*Cosh[x] + a*Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*Cosh[3*x] + b*(-24*a^3*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])]) + 3*Sqrt[a - b]*Sqrt[a + b]*(5*a^2 - b^2)*Sinh[x] - Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*Sinh[3*x])/(12*(a - b)^(5/2)*(a + b)^(5/2))`

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.92 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.12, number of steps used = 24, number of rules used = 23,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.769$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3588, 25, 26, 3042, 25, 26, 3044, 15, 3113, 2009, 3578, 26, 3042, 26, 3118, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh^3(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \sin(ix)^3}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\sin(ix)^3}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{4001} \\
 & i \int -\frac{i \cosh(x) \sinh^3(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh^3(x) \cosh(x)}{a \cosh(x) + b \sinh(x)} dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{i \sin(ix)^3 \cos(ix)}{a \cos(ix) - ib \sin(ix)} dx \\
& \downarrow 26 \\
& i \int \frac{\cos(ix) \sin(ix)^3}{a \cos(ix) - ib \sin(ix)} dx \\
& \downarrow 3588 \\
& i \left( \frac{a \int -i \sinh^3(x) dx}{a^2 - b^2} - \frac{ib \int -\cosh(x) \sinh^2(x) dx}{a^2 - b^2} + \frac{iab \int -\frac{\sinh^2(x)}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \right) \\
& \downarrow 25 \\
& i \left( \frac{a \int -i \sinh^3(x) dx}{a^2 - b^2} + \frac{ib \int \cosh(x) \sinh^2(x) dx}{a^2 - b^2} - \frac{iab \int \frac{\sinh^2(x)}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \right) \\
& \downarrow 26 \\
& i \left( -\frac{ia \int \sinh^3(x) dx}{a^2 - b^2} + \frac{ib \int \cosh(x) \sinh^2(x) dx}{a^2 - b^2} - \frac{iab \int \frac{\sinh^2(x)}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \right) \\
& \downarrow 3042 \\
& i \left( -\frac{ia \int i \sin(ix)^3 dx}{a^2 - b^2} + \frac{ib \int -\cos(ix) \sin(ix)^2 dx}{a^2 - b^2} - \frac{iab \int -\frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right) \\
& \downarrow 25 \\
& i \left( -\frac{ia \int i \sin(ix)^3 dx}{a^2 - b^2} - \frac{ib \int \cos(ix) \sin(ix)^2 dx}{a^2 - b^2} + \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right) \\
& \downarrow 26 \\
& i \left( \frac{a \int \sin(ix)^3 dx}{a^2 - b^2} - \frac{ib \int \cos(ix) \sin(ix)^2 dx}{a^2 - b^2} + \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right) \\
& \downarrow 3044 \\
& i \left( \frac{a \int \sin(ix)^3 dx}{a^2 - b^2} - \frac{b \int -\sinh^2(x) d(i \sinh(x))}{a^2 - b^2} + \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right)
\end{aligned}$$

$$\begin{aligned}
& \downarrow 15 \\
& i \left( \frac{a \int \sin(ix)^3 dx}{a^2 - b^2} + \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} \right) \\
& \downarrow 3113 \\
& i \left( \frac{ia \int (1 - \cosh^2(x)) d \cosh(x)}{a^2 - b^2} + \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} \right) \\
& \downarrow 2009 \\
& i \left( \frac{iab \int \frac{\sin(ix)^2}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right) \\
& \downarrow 3578 \\
& i \left( \frac{iab \left( -\frac{ib \int i \sinh(x) dx}{a^2 - b^2} + \frac{a^2 \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right) \\
& \downarrow 26 \\
& i \left( \frac{iab \left( \frac{b \int \sinh(x) dx}{a^2 - b^2} + \frac{a^2 \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right) \\
& \downarrow 3042 \\
& i \left( \frac{iab \left( \frac{b \int -i \sin(ix) dx}{a^2 - b^2} + \frac{a^2 \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right) \\
& \downarrow 26 \\
& i \left( \frac{iab \left( -\frac{ib \int \sin(ix) dx}{a^2 - b^2} + \frac{a^2 \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right)
\end{aligned}$$

$$\downarrow \text{3118}$$

$$i \left( \frac{iab \left( \frac{a^2 \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} + \frac{b \cosh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right)$$

$$\downarrow \text{3553}$$

$$i \left( \frac{iab \left( \frac{ia^2 \int \frac{1}{a^2 - b^2 - (-ib \cosh(x) - ia \sinh(x))^2} d(-ib \cosh(x) - ia \sinh(x))}{a^2 - b^2} - \frac{a \sinh(x)}{a^2 - b^2} + \frac{b \cosh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right)$$

$$\downarrow \text{219}$$

$$i \left( \frac{iab \left( \frac{ia^2 \operatorname{arctanh} \left( \frac{-ia \sinh(x) - ib \cosh(x)}{\sqrt{a^2 - b^2}} \right)}{(a^2 - b^2)^{3/2}} - \frac{a \sinh(x)}{a^2 - b^2} + \frac{b \cosh(x)}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{ib \sinh^3(x)}{3(a^2 - b^2)} + \frac{ia \left( \cosh(x) - \frac{\cosh^3(x)}{3} \right)}{a^2 - b^2} \right)$$

input `Int[Sinh[x]^3/(a + b*Tanh[x]),x]`

output `I*((I*a*(Cosh[x] - Cosh[x]^3/3))/(a^2 - b^2) + ((I/3)*b*Sinh[x]^3)/(a^2 - b^2) + (I*a*b*((I*a^2*ArcTanh[(-I)*b*Cosh[x] - I*a*Sinh[x])/Sqrt[a^2 - b^2]])/(a^2 - b^2)^(3/2) + (b*Cosh[x])/(a^2 - b^2) - (a*Sinh[x])/(a^2 - b^2)))/(a^2 - b^2)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 219  $\text{Int}[(a\_ + (b\_)*(x\_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 2009  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3044  $\text{Int}[\cos[(e\_ + (f\_)*(x\_)]^{(n\_)}*((a\_)*\sin[(e\_ + (f\_)*(x\_)]^{(m\_)}), x\_Symbol] \rightarrow \text{Simp}[1/(a*f) \ \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{(n-1)/2}], x], x, a*\sin[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{LtQ}[0, m, n])$
- rule 3113  $\text{Int}[\sin[(c\_ + (d\_)*(x\_)]^{(n\_)}), x\_Symbol] \rightarrow \text{Simp}[-d^{-1} \ \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{(n-1)/2}], x], x], x, \text{Cos}[c + d*x]], x] /; \text{FreeQ}\{c, d\}, x] \ \&\& \ \text{IGtQ}[(n-1)/2, 0]$
- rule 3118  $\text{Int}[\sin[(c\_ + (d\_)*(x\_)]), x\_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /; \text{FreeQ}\{c, d\}, x]$
- rule 3553  $\text{Int}[(\cos[(c\_ + (d\_)*(x\_)]*(a\_ + (b\_)*\sin[(c\_ + (d\_)*(x\_)]^{-1}), x\_Symbol] \rightarrow \text{Simp}[-d^{-1} \ \text{Subst}[\text{Int}[1/(a^2 + b^2 - x^2)], x], x, b*\text{Cos}[c + d*x] - a*\text{Sin}[c + d*x]], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

rule 3578

```
Int[sin[(c_.) + (d_.)*(x_)]^(m_)/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin
[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[(-a)*(Sin[c + d*x]^(m - 1)/(d*(a^2
+ b^2)*(m - 1))), x] + (Simp[a^2/(a^2 + b^2) Int[Sin[c + d*x]^(m - 2)/(a
*Cos[c + d*x] + b*Sin[c + d*x]), x], x] + Simp[b/(a^2 + b^2) Int[Sin[c +
d*x]^(m - 1), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ
[m, 1]
```

rule 3588

```
Int[(cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.))/(cos[(c_.
) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[b
/(a^2 + b^2) Int[Cos[c + d*x]^m*Sin[c + d*x]^(n - 1), x], x] + (Simp[a/(a
^2 + b^2) Int[Cos[c + d*x]^(m - 1)*Sin[c + d*x]^n, x], x] - Simp[a*(b/(a^
2 + b^2)) Int[Cos[c + d*x]^(m - 1)*(Sin[c + d*x]^(n - 1)/(a*Cos[c + d*x]
+ b*Sin[c + d*x])), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
&& IGtQ[m, 0] && IGtQ[n, 0]
```

rule 4001

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

## Maple [A] (verified)

Time = 4.11 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.21

method	result
default	$-\frac{8}{(16a-16b)(1+\tanh(\frac{x}{2}))^2} + \frac{16}{3(1+\tanh(\frac{x}{2}))^3(16a-16b)} - \frac{a}{2(a-b)^2(1+\tanh(\frac{x}{2}))} - \frac{16}{3(\tanh(\frac{x}{2})-1)^3(16a+16b)} - \frac{1}{(16a+16b)}$
risch	$\frac{e^{3x}}{24a+24b} - \frac{3e^x a}{8(a+b)^2} - \frac{e^x b}{8(a+b)^2} - \frac{3e^{-x} a}{8(a-b)^2} + \frac{e^{-x} b}{8(a-b)^2} + \frac{e^{-3x}}{24a-24b} - \frac{b a^3 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} (a+b)^2 (a-b)^2} + \frac{b a^3 \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} (a+b)^2 (a-b)^2}$

input

```
int(sinh(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-8/(16*a-16*b)/(1+tanh(1/2*x))^2+16/3/(1+tanh(1/2*x))^3/(16*a-16*b)-1/2*a/
(a-b)^2/(1+tanh(1/2*x))-16/3/(tanh(1/2*x)-1)^3/(16*a+16*b)-8/(16*a+16*b)/(
tanh(1/2*x)-1)^2+1/2*a/(a+b)^2/(tanh(1/2*x)-1)-2*a^3*b/(a-b)^2/(a+b)^2/(a^
2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 903 vs.  $2(129) = 258$ .

Time = 0.13 (sec) , antiderivative size = 1861, normalized size of antiderivative = 13.58

$$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
[1/24*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 + 6*(
a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^5 + (a^
5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^6 + a^5 + a^4*b -
2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*
a^2*b^3 - a*b^4 - b^5)*cosh(x)^4 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*
b^3 - a*b^4 - b^5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*
cosh(x)^2)*sinh(x)^4 + 4*(5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 -
b^5)*cosh(x)^3 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5
)*cosh(x))*sinh(x)^3 - 3*(3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4
+ b^5)*cosh(x)^2 - 3*(3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^
5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 + 6*(3
*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*cosh(x)^2)*sinh(x)^2
- 24*(a^3*b*cosh(x)^3 + 3*a^3*b*cosh(x)^2*sinh(x) + 3*a^3*b*cosh(x)*sinh(
x)^2 + a^3*b*sinh(x)^3)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b
)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + 2*sqrt(-a^2 + b^2)*(cosh(x) + sin
h(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sin
h(x)^2 + a - b)) + 6*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*
cosh(x)^5 - 2*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*cosh
(x)^3 - (3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^5)*cosh(x))*s
inh(x))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^3 + 3*(a^6 - 3*a^4...
```

**Sympy [F]**

$$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx = \int \frac{\sinh^3(x)}{a + b \tanh(x)} dx$$

input `integrate(sinh(x)**3/(a+b*tanh(x)),x)`

output `Integral(sinh(x)**3/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

$$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx = -\frac{2a^3b \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}} - \frac{(9ae^{2x} - 3be^{2x} - a + b)e^{-3x}}{24(a^2 - 2ab + b^2)} + \frac{a^2e^{3x} + 2abe^{3x} + b^2e^{3x} - 9a^2e^x - 12abe^x - 3b^2e^x}{24(a^3 + 3a^2b + 3ab^2 + b^3)}$$

input `integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="giac")`



output

$$\begin{aligned} & -2a^3b \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right) / \left((a^4 - 2a^2b^2 + b^4) \sqrt{a^2 - b^2}\right) - 1/24(9ae^{2x} - 3be^{2x} - a + b)e^{-3x} / (a^2 - 2ab + b^2) \\ & + 1/24(a^2e^{3x} + 2ab^2e^{3x} + b^2e^{3x} - 9a^2e^x - 12ab^2e^x - 3b^2e^x) / (a^3 + 3a^2b + 3ab^2 + b^3) \end{aligned}$$

**Mupad [B] (verification not implemented)**

Time = 2.37 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.91

$$\begin{aligned} & \int \frac{\sinh^3(x)}{a + b \tanh(x)} dx \\ & = \frac{e^{-3x}}{24a - 24b} + \frac{e^{3x}}{24a + 24b} - \frac{e^x(3a + b)}{8(a + b)^2} - \frac{e^{-x}(3a - b)}{8(a - b)^2} \\ & \quad - \frac{2 \operatorname{atan}\left(\frac{a^3 b e^x \sqrt{a^{10} - 5a^8 b^2 + 10a^6 b^4 - 10a^4 b^6 + 5a^2 b^8 - b^{10}}}{a^5 \sqrt{a^6 b^2 - b^5} \sqrt{a^6 b^2 + 2a^2 b^3} \sqrt{a^6 b^2 - 2a^3 b^2} \sqrt{a^6 b^2 + a b^4} \sqrt{a^6 b^2 - a^4 b} \sqrt{a^6 b^2}}\right) \sqrt{a^6 b^2}}{\sqrt{a^{10} - 5a^8 b^2 + 10a^6 b^4 - 10a^4 b^6 + 5a^2 b^8 - b^{10}}} \end{aligned}$$

input

int(sinh(x)^3/(a + b\*tanh(x)),x)

output

$$\begin{aligned} & \exp(-3x)/(24a - 24b) + \exp(3x)/(24a + 24b) - (\exp(x)(3a + b))/(8(a + b)^2) - (\exp(-x)(3a - b))/(8(a - b)^2) \\ & - (2 \operatorname{atan}\left(\frac{a^3 b \exp(x)(a^{10} - b^{10} + 5a^2 b^8 - 10a^4 b^6 + 10a^6 b^4 - 5a^8 b^2)^{1/2}}{a^5 (a^6 b^2)^{1/2} - b^5 (a^6 b^2)^{1/2} + 2a^2 b^3 (a^6 b^2)^{1/2} - 2a^3 b^2 (a^6 b^2)^{1/2} + a b^4 (a^6 b^2)^{1/2} - a^4 b (a^6 b^2)^{1/2}}\right) (a^6 b^2)^{1/2}) / (a^{10} - b^{10} + 5a^2 b^8 - 10a^4 b^6 + 10a^6 b^4 - 5a^8 b^2)^{1/2} \end{aligned}$$

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.31

$$\begin{aligned} & \int \frac{\sinh^3(x)}{a + b \tanh(x)} dx \\ & = \frac{15e^{4x} a^4 b - 18e^{4x} a^2 b^3 - 15e^{2x} a^4 b + 18e^{2x} a^2 b^3 + 2e^{6x} a^2 b^3 + a^5 - 2a^3 b^2 + 6e^{4x} a^3 b^2 + 3e^{4x} a b^4 - 2e^{6x} a^3 b^2}{a^2 + b^2} \end{aligned}$$

input `int(sinh(x)^3/(a+b*tanh(x)),x)`

output 
$$\begin{aligned} & (-48e^{3x}\sqrt{a^2-b^2}\operatorname{atan}\left(\frac{e^{2x}a+e^{2x}b}{\sqrt{a^2-b^2}}\right) \\ & + e^{6x}a^3b - e^{6x}a^5 - e^{6x}a^4b - 2e^{6x}a^3b^2 + 2e^{6x} \\ & + e^{6x}a^2b^3 + e^{6x}ab^4 - e^{6x}b^5 - 9e^{4x}a^5 + 15e^{4x} \\ & + e^{4x}a^4b + 6e^{4x}a^3b^2 - 18e^{4x}a^2b^3 + 3e^{4x}a \\ & + e^{4x}ab^4 + 3e^{4x}b^5 - 9e^{2x}a^5 - 15e^{2x}a^4b + 6e^{2x} \\ & + e^{2x}a^3b^2 + 18e^{2x}a^2b^3 + 3e^{2x}ab^4 - 3e^{2x}b^5 \\ & + a^5 + a^4b - 2a^3b^2 - 2a^2b^3 + ab^4 + b^5) / (24e^{3x} \\ & + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6)) \end{aligned}$$

### 3.82 $\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx$

Optimal result	674
Mathematica [A] (verified)	674
Rubi [A] (verified)	675
Maple [A] (verified)	677
Fricas [B] (verification not implemented)	678
Sympy [F]	678
Maxima [A] (verification not implemented)	679
Giac [A] (verification not implemented)	679
Mupad [B] (verification not implemented)	680
Reduce [B] (verification not implemented)	680

#### Optimal result

Integrand size = 13, antiderivative size = 84

$$\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx = \frac{a \log(1 - \tanh(x))}{4(a+b)^2} - \frac{a \log(1 + \tanh(x))}{4(a-b)^2} + \frac{a^2 b \log(a+b \tanh(x))}{(a^2-b^2)^2} - \frac{\cosh^2(x)(b-a \tanh(x))}{2(a^2-b^2)}$$

output

```
1/4*a*ln(1-tanh(x))/(a+b)^2-1/4*a*ln(1+tanh(x))/(a-b)^2+a^2*b*ln(a+b*tanh(x))/(a^2-b^2)^2-cosh(x)^2*(b-a*tanh(x))/(2*a^2-2*b^2)
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.87

$$\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx = \frac{(-a^2 b + b^3) \cosh(2x) + a(-2(a^2 + b^2)x + 4ab \log(a \cosh(x) + b \sinh(x)) + (a^2 - b^2) \sinh(2x))}{4(a-b)^2(a+b)^2}$$

input

```
Integrate[Sinh[x]^2/(a + b*Tanh[x]),x]
```

output

$$\frac{((-a^2b) + b^3)\text{Cosh}[2x] + a(-2(a^2 + b^2)x + 4ab\text{Log}[a\text{Cosh}[x] + b\text{Sinh}[x]] + (a^2 - b^2)\text{Sinh}[2x])}{4(a - b)^2(a + b)^2}$$

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.69, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {3042, 25, 3999, 601, 27, 657, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sinh^2(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{\sin(ix)^2}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{25} \\ & -\int \frac{\sin(ix)^2}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{3999} \\ & b \int \frac{b^2 \tanh^2(x)}{(a + b \tanh(x)) (b^2 - b^2 \tanh^2(x))^2} d(b \tanh(x)) \\ & \quad \downarrow \text{601} \\ & b \left( -\frac{\int \frac{ab^2(a - b \tanh(x))}{(a^2 - b^2)(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))} d(b \tanh(x))}{2b^2} - \frac{\frac{b^2}{a^2 - b^2} - \frac{ab \tanh(x)}{a^2 - b^2}}{2(b^2 - b^2 \tanh^2(x))} \right) \\ & \quad \downarrow \text{27} \\ & b \left( -\frac{a \int \frac{a - b \tanh(x)}{(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))} d(b \tanh(x))}{2(a^2 - b^2)} - \frac{\frac{b^2}{a^2 - b^2} - \frac{ab \tanh(x)}{a^2 - b^2}}{2(b^2 - b^2 \tanh^2(x))} \right) \\ & \quad \downarrow \text{657} \end{aligned}$$

$$b \left( -\frac{a \int \left( -\frac{2a}{(a-b)(a+b)(a+b \tanh(x))} + \frac{a-b}{2b(a+b)(b-b \tanh(x))} + \frac{a+b}{2(a-b)b(\tanh(x)b+b)} \right) d(b \tanh(x))}{2(a^2 - b^2)} - \frac{\frac{b^2}{a^2 - b^2} - \frac{ab \tanh(x)}{a^2 - b^2}}{2(b^2 - b^2 \tanh^2(x))} \right)$$

↓ 2009

$$b \left( -\frac{\frac{b^2}{a^2 - b^2} - \frac{ab \tanh(x)}{a^2 - b^2}}{2(b^2 - b^2 \tanh^2(x))} - \frac{a \left( -\frac{2a \log(a+b \tanh(x))}{a^2 - b^2} - \frac{(a-b) \log(b-b \tanh(x))}{2b(a+b)} + \frac{(a+b) \log(b \tanh(x)+b)}{2b(a-b)} \right)}{2(a^2 - b^2)} \right)$$

input `Int[Sinh[x]^2/(a + b*Tanh[x]),x]`

output `b*(-1/2*(a*(-1/2*((a - b)*Log[b - b*Tanh[x]])/(b*(a + b)) - (2*a*Log[a + b*Tanh[x]])/(a^2 - b^2) + ((a + b)*Log[b + b*Tanh[x]]/(2*(a - b)*b)))/(a^2 - b^2) - (b^2/(a^2 - b^2) - (a*b*Tanh[x]/(a^2 - b^2))/(2*(b^2 - b^2*Tanh[x]^2)))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 601 `Int[(x_)^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[x^m*(c + d*x)^n, a + b*x^2, x], e = Coeff[PolynomialRemainder[x^m*(c + d*x)^n, a + b*x^2, x], x, 0], f = Coeff[PolynomialRemainder[x^m*(c + d*x)^n, a + b*x^2, x], x, 1]}, Simp[(a*f - b*e*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Simp[1/(2*a*(p + 1) Int[(c + d*x)^n*(a + b*x^2)^(p + 1)*ExpandToSum[(2*a*(p + 1)*Qx)/(c + d*x)^n + (e*(2*p + 3))/(c + d*x)^n, x], x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 1] && LtQ[p, -1] && ILtQ[n, 0] && NeQ[b*c^2 + a*d^2, 0]`

rule 657 `Int[(((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.)))/((a_.) + (c_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g, m}, x] && IntegersQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3999 `Int[sin[(e_.) + (f_.)*(x_)^(m_.)]*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.17

method	result
risch	$-\frac{ax}{2(a+b)^2} + \frac{e^{2x}}{8a+8b} - \frac{e^{-2x}}{8(a-b)} - \frac{2a^2bx}{a^4-2a^2b^2+b^4} + \frac{a^2b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^4-2a^2b^2+b^4}$
default	$\frac{4}{(8a+8b)\left(\tanh\left(\frac{x}{2}\right)-1\right)^2} + \frac{8}{(16a+16b)\left(\tanh\left(\frac{x}{2}\right)-1\right)} + \frac{a \ln\left(\tanh\left(\frac{x}{2}\right)-1\right)}{2(a+b)^2} + \frac{a^2b \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{(a-b)^2(a+b)^2} - \frac{1}{(8a-8b)\left(\tanh\left(\frac{x}{2}\right)+1\right)}$

input `int(sinh(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output 
$$-1/2*a*x/(a+b)^2+1/8/(a+b)*\exp(2*x)-1/8/(a-b)*\exp(-2*x)-2*a^2*b/(a^4-2*a^2*b^2+b^4)*x+a^2*b/(a^4-2*a^2*b^2+b^4)*\ln(\exp(2*x)+(a-b)/(a+b))$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 334 vs.  $2(79) = 158$ .

Time = 0.12 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.98

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{(a^3 - a^2b - ab^2 + b^3) \cosh(x)^4 + 4(a^3 - a^2b - ab^2 + b^3) \cosh(x) \sinh(x)^3 + (a^3 - a^2b - ab^2 + b^3) \sinh(x)^4}{(a^4 - 2a^2b^2 + b^4) \cosh(x)^2 + 2(a^4 - 2a^2b^2 + b^4) \cosh(x) \sinh(x) + (a^4 - 2a^2b^2 + b^4) \sinh(x)^2} + \frac{2(a^3 - a^2b - ab^2 + b^3) \cosh(x)^2 \sinh(x) + 2(a^3 - a^2b - ab^2 + b^3) \cosh(x) \sinh(x)^2 + 2(a^3 - a^2b - ab^2 + b^3) \sinh(x)^3}{(a^4 - 2a^2b^2 + b^4) \cosh(x)^2 + 2(a^4 - 2a^2b^2 + b^4) \cosh(x) \sinh(x) + (a^4 - 2a^2b^2 + b^4) \sinh(x)^2}$$

input `integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`

output

```
1/8*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 - a^2*b - a*b^2 + b^3)
*cosh(x)*sinh(x)^3 + (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^4 - 4*(a^3 + 2*a^
2*b + a*b^2)*x*cosh(x)^2 - a^3 - a^2*b + a*b^2 + b^3 + 2*(3*(a^3 - a^2*b -
a*b^2 + b^3)*cosh(x)^2 - 2*(a^3 + 2*a^2*b + a*b^2)*x)*sinh(x)^2 + 8*(a^2*
b*cosh(x)^2 + 2*a^2*b*cosh(x)*sinh(x) + a^2*b*sinh(x)^2)*log(2*(a*cosh(x)
+ b*sinh(x))/(cosh(x) - sinh(x))) + 4*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)
^3 - 2*(a^3 + 2*a^2*b + a*b^2)*x*cosh(x))*sinh(x))/((a^4 - 2*a^2*b^2 + b^4
)*cosh(x)^2 + 2*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x) + (a^4 - 2*a^2*b^2
+ b^4)*sinh(x)^2)
```

**Sympy [F]**

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx = \int \frac{\sinh^2(x)}{a + b \tanh(x)} dx$$

input `integrate(sinh(x)**2/(a+b*tanh(x)),x)`

output `Integral(sinh(x)**2/(a + b*tanh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.99

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx = \frac{a^2 b \log(-(a-b)e^{(-2x)} - a - b)}{a^4 - 2a^2 b^2 + b^4} - \frac{ax}{2(a^2 + 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)} - \frac{e^{(-2x)}}{8(a-b)}$$

input `integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")`output `a^2*b*log(-(a - b)*e^(-2*x) - a - b)/(a^4 - 2*a^2*b^2 + b^4) - 1/2*a*x/(a^2 + 2*a*b + b^2) + 1/8*e^(2*x)/(a + b) - 1/8*e^(-2*x)/(a - b)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.20

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx = \frac{a^2 b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^4 - 2a^2 b^2 + b^4} - \frac{ax}{2(a^2 - 2ab + b^2)} + \frac{(2ae^{(2x)} - a + b)e^{(-2x)}}{8(a^2 - 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)}$$

input `integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="giac")`output `a^2*b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^4 - 2*a^2*b^2 + b^4) - 1/2*a*x/(a^2 - 2*a*b + b^2) + 1/8*(2*a*e^(2*x) - a + b)*e^(-2*x)/(a^2 - 2*a*b + b^2) + 1/8*e^(2*x)/(a + b)`



**Mupad [B] (verification not implemented)**

Time = 2.33 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.96

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx = \frac{e^{2x}}{8a + 8b} - \frac{e^{-2x}}{8a - 8b} - \frac{ax}{2(a - b)^2} + \frac{a^2 b \ln(a - b + a e^{2x} + b e^{2x})}{a^4 - 2a^2 b^2 + b^4}$$

input `int(sinh(x)^2/(a + b*tanh(x)),x)`output `exp(2*x)/(8*a + 8*b) - exp(-2*x)/(8*a - 8*b) - (a*x)/(2*(a - b)^2) + (a^2*b*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^4 + b^4 - 2*a^2*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.81

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx = \frac{e^{4x} a^3 - e^{4x} a^2 b - e^{4x} a b^2 + e^{4x} b^3 + 8e^{2x} \log(e^{2x} a + e^{2x} b + a - b) a^2 b - 4e^{2x} a^3 x - 8e^{2x} a^2 b x - 4e^{2x} a b^2 x - 8e^{2x} (a^4 - 2a^2 b^2 + b^4)}{8e^{2x} (a^4 - 2a^2 b^2 + b^4)}$$

input `int(sinh(x)^2/(a+b*tanh(x)),x)`output `(e**(4*x)*a**3 - e**(4*x)*a**2*b - e**(4*x)*a*b**2 + e**(4*x)*b**3 + 8*e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*b - 4*e**(2*x)*a**3*x - 8*e**(2*x)*a**2*b*x - 4*e**(2*x)*a*b**2*x - a**3 - a**2*b + a*b**2 + b**3)/(8*e**(2*x)*(a**4 - 2*a**2*b**2 + b**4))`

### 3.83 $\int \frac{\sinh(x)}{a+b \tanh(x)} dx$

Optimal result . . . . .	681
Mathematica [A] (verified) . . . . .	681
Rubi [C] (verified) . . . . .	682
Maple [A] (verified) . . . . .	685
Fricas [B] (verification not implemented) . . . . .	685
Sympy [F] . . . . .	686
Maxima [F(-2)] . . . . .	686
Giac [A] (verification not implemented) . . . . .	687
Mupad [B] (verification not implemented) . . . . .	687
Reduce [B] (verification not implemented) . . . . .	688

#### Optimal result

Integrand size = 11, antiderivative size = 72

$$\int \frac{\sinh(x)}{a+b \tanh(x)} dx = \frac{ab \arctan\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} + \frac{a \cosh(x)}{a^2-b^2} - \frac{b \sinh(x)}{a^2-b^2}$$

output `a*b*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))/(a^2-b^2)^(3/2)+a*cosh(x)/(a^2-b^2)-b*sinh(x)/(a^2-b^2)`

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.10

$$\int \frac{\sinh(x)}{a+b \tanh(x)} dx = \frac{2ab \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}}\right)}{(a-b)^{3/2}(a+b)^{3/2}} + \frac{a \cosh(x)}{a^2-b^2} + \frac{b \sinh(x)}{-a^2+b^2}$$

input `Integrate[Sinh[x]/(a + b*Tanh[x]),x]`

output `(2*a*b*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])])/((a - b)^(3/2)* (a + b)^(3/2)) + (a*Cosh[x])/(a^2 - b^2) + (b*Sinh[x])/(-a^2 + b^2)`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.22, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.273$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3588, 26, 3042, 26, 3117, 3118, 3553, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sinh(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{4001} \\
 & -i \int \frac{i \cosh(x) \sinh(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\sinh(x) \cosh(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix) \cos(ix)}{a \cos(ix) - ib \sin(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\cos(ix) \sin(ix)}{a \cos(ix) - ib \sin(ix)} dx \\
 & \quad \downarrow \text{3588} \\
 & -i \left( \frac{a \int i \sinh(x) dx}{a^2 - b^2} - \frac{ib \int \cosh(x) dx}{a^2 - b^2} + \frac{iab \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& -i \left( \frac{ia \int \sinh(x) dx}{a^2 - b^2} - \frac{ib \int \cosh(x) dx}{a^2 - b^2} + \frac{iab \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \right) \\
& \downarrow 3042 \\
& -i \left( -\frac{ib \int \sin(ix + \frac{\pi}{2}) dx}{a^2 - b^2} + \frac{ia \int -i \sin(ix) dx}{a^2 - b^2} + \frac{iab \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right) \\
& \downarrow 26 \\
& -i \left( -\frac{ib \int \sin(ix + \frac{\pi}{2}) dx}{a^2 - b^2} + \frac{a \int \sin(ix) dx}{a^2 - b^2} + \frac{iab \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} \right) \\
& \downarrow 3117 \\
& -i \left( \frac{a \int \sin(ix) dx}{a^2 - b^2} + \frac{iab \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} - \frac{ib \sinh(x)}{a^2 - b^2} \right) \\
& \downarrow 3118 \\
& -i \left( \frac{iab \int \frac{1}{a \cos(ix) - ib \sin(ix)} dx}{a^2 - b^2} - \frac{ib \sinh(x)}{a^2 - b^2} + \frac{ia \cosh(x)}{a^2 - b^2} \right) \\
& \downarrow 3553 \\
& -i \left( -\frac{ab \int \frac{1}{a^2 - b^2 - (-ib \cosh(x) - ia \sinh(x))^2} d(-ib \cosh(x) - ia \sinh(x))}{a^2 - b^2} - \frac{ib \sinh(x)}{a^2 - b^2} + \frac{ia \cosh(x)}{a^2 - b^2} \right) \\
& \downarrow 219 \\
& -i \left( -\frac{ab \operatorname{arctanh} \left( \frac{-ia \sinh(x) - ib \cosh(x)}{\sqrt{a^2 - b^2}} \right)}{(a^2 - b^2)^{3/2}} - \frac{ib \sinh(x)}{a^2 - b^2} + \frac{ia \cosh(x)}{a^2 - b^2} \right)
\end{aligned}$$

input `Int [Sinh [x] / (a + b*Tanh [x]), x]`

output `(-I)*(-((a*b*ArcTanh[((-I)*b*Cosh[x] - I*a*Sinh[x])/Sqrt[a^2 - b^2]])/(a^2 - b^2)^(3/2)) + (I*a*Cosh[x])/(a^2 - b^2) - (I*b*Sinh[x])/(a^2 - b^2))`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 219  $\text{Int}[(a + (b \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) * \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3117  $\text{Int}[\sin[\pi/2 + (c \cdot x) + (d \cdot x)], x\_Symbol] \rightarrow \text{Simp}[\sin[c + d \cdot x]/d, x] /; \text{FreeQ}\{c, d\}, x]$
- rule 3118  $\text{Int}[\sin[(c \cdot x) + (d \cdot x)], x\_Symbol] \rightarrow \text{Simp}[-\cos[c + d \cdot x]/d, x] /; \text{FreeQ}\{c, d\}, x]$
- rule 3553  $\text{Int}[(\cos[(c \cdot x) + (d \cdot x)] \cdot (a \cdot x) + (b \cdot x) \cdot \sin[(c \cdot x) + (d \cdot x)])^{-1}, x\_Symbol] \rightarrow \text{Simp}[-d^{-1} \text{Subst}[\text{Int}[1/(a^2 + b^2 - x^2), x], x, b \cdot \cos[c + d \cdot x] - a \cdot \sin[c + d \cdot x]], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 3588  $\text{Int}[(\cos[(c \cdot x) + (d \cdot x)]^m \cdot \sin[(c \cdot x) + (d \cdot x)]^n) / (\cos[(c \cdot x) + (d \cdot x)] \cdot (a \cdot x) + (b \cdot x) \cdot \sin[(c \cdot x) + (d \cdot x)]), x\_Symbol] \rightarrow \text{Simp}[b / (a^2 + b^2) \text{Int}[\cos[c + d \cdot x]^m \cdot \sin[c + d \cdot x]^{n-1}, x], x] + (\text{Simp}[a / (a^2 + b^2) \text{Int}[\cos[c + d \cdot x]^{m-1} \cdot \sin[c + d \cdot x]^n, x], x] - \text{Simp}[a \cdot (b / (a^2 + b^2)) \text{Int}[\cos[c + d \cdot x]^{m-1} \cdot (\sin[c + d \cdot x]^{n-1} / (a \cdot \cos[c + d \cdot x] + b \cdot \sin[c + d \cdot x])), x], x]) /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0]$

rule 4001

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

**Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.28

method	result	size
default	$\frac{2ab \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{(a+b)(a-b)\sqrt{a^2 - b^2}} + \frac{4}{(4a-4b)(1+\tanh\left(\frac{x}{2}\right))} - \frac{4}{(4a+4b)(\tanh\left(\frac{x}{2}\right)-1)}$	92
risch	$\frac{e^x}{2b+2a} + \frac{e^{-x}}{2a-2b} - \frac{ba \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)(a-b)} + \frac{ba \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)(a-b)}$	120

input

```
int(sinh(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
2*a*b/(a+b)/(a-b)/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))+4/(4*a-4*b)/(1+tanh(1/2*x))-4/(4*a+4*b)/(tanh(1/2*x)-1)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(68) = 136.

Time = 0.13 (sec) , antiderivative size = 427, normalized size of antiderivative = 5.93

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx$$

$$= \frac{a^3 + a^2b - ab^2 - b^3 + (a^3 - a^2b - ab^2 + b^3) \cosh(x)^2 + 2(a^3 - a^2b - ab^2 + b^3) \cosh(x) \sinh(x) + (a^3 - a^2b - ab^2 + b^3) \sinh(x)^2}{2((a^4 - b^4) \cosh(x)^2 + (a^4 - b^4) \sinh(x)^2)}$$

input

```
integrate(sinh(x)/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
[1/2*(a^3 + a^2*b - a*b^2 - b^3 + (a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^2 +
2*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh(x) + (a^3 - a^2*b - a*b^2 + b^3
)*sinh(x)^2 + 2*(a*b*cosh(x) + a*b*sinh(x))*sqrt(-a^2 + b^2)*log(((a + b)*
cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + 2*sqrt(-a^2 +
b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*s
inh(x) + (a + b)*sinh(x)^2 + a - b)))/((a^4 - 2*a^2*b^2 + b^4)*cosh(x) + (
a^4 - 2*a^2*b^2 + b^4)*sinh(x)), 1/2*(a^3 + a^2*b - a*b^2 - b^3 + (a^3 - a
^2*b - a*b^2 + b^3)*cosh(x)^2 + 2*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh
(x) + (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^2 - 4*(a*b*cosh(x) + a*b*sinh(x)
)*sqrt(a^2 - b^2)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x
)))))/((a^4 - 2*a^2*b^2 + b^4)*cosh(x) + (a^4 - 2*a^2*b^2 + b^4)*sinh(x))]
```

**Sympy [F]**

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx = \int \frac{\sinh(x)}{a + b \tanh(x)} dx$$

input

```
integrate(sinh(x)/(a+b*tanh(x)),x)
```

output

```
Integral(sinh(x)/(a + b*tanh(x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input

```
integrate(sinh(x)/(a+b*tanh(x)),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.83

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx = \frac{2ab \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{\frac{3}{2}}} + \frac{e^{-x}}{2(a - b)} + \frac{e^x}{2(a + b)}$$

input `integrate(sinh(x)/(a+b*tanh(x)),x, algorithm="giac")`output `2*a*b*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(a^2 - b^2)^(3/2) + 1/2*e^(-x)/(a - b) + 1/2*e^x/(a + b)`**Mupad [B] (verification not implemented)**

Time = 2.27 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.18

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx = \frac{e^x}{2a + 2b} + \frac{e^{-x}}{2a - 2b} + \frac{2 \operatorname{atan}\left(\frac{ab e^x \sqrt{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6}}{a^3 \sqrt{a^2 b^2 + b^3} \sqrt{a^2 b^2 - a b^2} \sqrt{a^2 b^2 - a^2 b} \sqrt{a^2 b^2}}\right) \sqrt{a^2 b^2}}{\sqrt{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6}}$$

input `int(sinh(x)/(a + b*tanh(x)),x)`output `exp(x)/(2*a + 2*b) + exp(-x)/(2*a - 2*b) + (2*atan((a*b*exp(x))*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2)))/(a^3*(a^2*b^2)^(1/2) + b^3*(a^2*b^2)^(1/2) - a*b^2*(a^2*b^2)^(1/2) - a^2*b*(a^2*b^2)^(1/2))*(a^2*b^2)^(1/2))/(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2)`



**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.38

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx$$

$$= \frac{4e^x \sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) a^2 b + 2e^x \cosh(x) a^4 - 4e^x \cosh(x) a^2 b^2 + 2e^x \cosh(x) b^4 - e^{2x} a^3 b + e^{2x} a^2 b^2}{2e^x a (a^4 - 2a^2 b^2 + b^4)}$$

input `int(sinh(x)/(a+b*tanh(x)),x)`output `(4***x*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**2*b + 2***x*cosh(x)*a**4 - 4***x*cosh(x)*a**2*b**2 + 2***x*cosh(x)*b**4 - e**(2*x)*a**3*b + e**(2*x)*a**2*b**2 + e**(2*x)*a*b**3 - e**(2*x)*b**4 + a**3*b + a**2*b**2 - a*b**3 - b**4)/(2***x*a*(a**4 - 2*a**2*b**2 + b**4))`

### 3.84 $\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx$

Optimal result	689
Mathematica [A] (verified)	689
Rubi [C] (verified)	690
Maple [A] (verified)	692
Fricas [A] (verification not implemented)	692
Sympy [F]	693
Maxima [F(-2)]	693
Giac [A] (verification not implemented)	693
Mupad [B] (verification not implemented)	694
Reduce [B] (verification not implemented)	694

#### Optimal result

Integrand size = 11, antiderivative size = 52

$$\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx = -\frac{b \arctan\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{a \sqrt{a^2-b^2}} - \frac{\operatorname{arctanh}(\cosh(x))}{a}$$

output `-b*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))/a/(a^2-b^2)^(1/2)-arctanh(cosh(x))/a`

#### Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.31

$$\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx = \frac{2b \arctan\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right)}{\sqrt{a-b}\sqrt{a+b}} - \frac{\log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right)}{a}$$

input `Integrate[Csch[x]/(a + b*Tanh[x]),x]`

output `((-2*b*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])])/(Sqrt[a - b]*Sqrt[a + b]) - Log[Cosh[x/2]] + Log[Sinh[x/2]])/a`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3589, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\sin(ix)(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\sin(ix)(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & i \int -\frac{i \coth(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \cos(ix)}{\sin(ix)(a \cos(ix) - ib \sin(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\cos(ix)}{\sin(ix)(a \cos(ix) - ib \sin(ix))} dx \\
 & \quad \downarrow \text{3589} \\
 & i \int \left( \frac{ib}{a(a \cosh(x) + b \sinh(x))} - \frac{i \operatorname{csch}(x)}{a} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$i \left( \frac{ib \arctan\left(\frac{a \sinh(x) + b \cosh(x)}{\sqrt{a^2 - b^2}}\right)}{a\sqrt{a^2 - b^2}} + \frac{i \operatorname{arctanh}(\cosh(x))}{a} \right)$$

input `Int[Csch[x]/(a + b*Tanh[x]),x]`

output `I*((I*b*ArcTan[(b*Cosh[x] + a*Sinh[x])/Sqrt[a^2 - b^2]])/(a*Sqrt[a^2 - b^2]) + (I*ArcTanh[Cosh[x]])/a)`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3589 `Int[(cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.))/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]`

rule 4001 `Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

**Maple [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.02

method	result	size
default	$\frac{\ln(\tanh(\frac{x}{2}))}{a} - \frac{2b \arctan\left(\frac{2a \tanh(\frac{x}{2}) + 2b}{2\sqrt{a^2 - b^2}}\right)}{a\sqrt{a^2 - b^2}}$	53
risch	$\frac{\ln(e^x - 1)}{a} - \frac{\ln(e^x + 1)}{a} - \frac{b \ln\left(e^x + \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2} a} + \frac{b \ln\left(e^x - \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2} a}$	97

input `int(csch(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`output `1/a*ln(tanh(1/2*x))-2*b/a/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 237, normalized size of antiderivative = 4.56

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx$$

$$= \left[ -\frac{\sqrt{-a^2 + b^2} b \log\left(\frac{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + 2\sqrt{-a^2 + b^2}(\cosh(x) + \sinh(x)) - a + b}{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a - b}\right) + (a^2 - b^2)}{a^3 - ab^2} \right]$$

input `integrate(csch(x)/(a+b*tanh(x)),x, algorithm="fricas")`output `[-(sqrt(-a^2 + b^2)*b*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b)) + (a^2 - b^2)*log(cosh(x) + sinh(x) + 1) - (a^2 - b^2)*log(cosh(x) + sinh(x) - 1))/(a^3 - a*b^2), (2*sqrt(a^2 - b^2)*b*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x))) - (a^2 - b^2)*log(cosh(x) + sinh(x) + 1) + (a^2 - b^2)*log(cosh(x) + sinh(x) - 1))/(a^3 - a*b^2)]`

**Sympy [F]**

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx$$

input `integrate(csch(x)/(a+b*tanh(x)),x)`

output `Integral(csch(x)/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(csch(x)/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.15

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx = -\frac{2b \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}a} - \frac{\log(e^x + 1)}{a} + \frac{\log(|e^x - 1|)}{a}$$

input `integrate(csch(x)/(a+b*tanh(x)),x, algorithm="giac")`

output `-2*b*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*a) - log(e^x + 1)/a + log(abs(e^x - 1))/a`

**Mupad [B] (verification not implemented)**

Time = 2.47 (sec) , antiderivative size = 177, normalized size of antiderivative = 3.40

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx = \frac{\ln(32ab - 32a^2 + 32a^2 e^x - 32ab e^x)}{a} - \frac{\ln(32ab - 32a^2 - 32a^2 e^x + 32ab e^x)}{a} - \frac{b \ln(32ab^2 e^x + 32a^2 b e^x - 32ab \sqrt{b^2 - a^2}) \sqrt{b^2 - a^2}}{ab^2 - a^3} + \frac{b \ln(32ab^2 e^x + 32a^2 b e^x + 32ab \sqrt{b^2 - a^2}) \sqrt{b^2 - a^2}}{ab^2 - a^3}$$

input `int(1/(sinh(x)*(a + b*tanh(x))),x)`output `log(32*a*b - 32*a^2 + 32*a^2*exp(x) - 32*a*b*exp(x))/a - log(32*a*b - 32*a^2 - 32*a^2*exp(x) + 32*a*b*exp(x))/a - (b*log(32*a*b^2*exp(x) + 32*a^2*b*exp(x) - 32*a*b*(b^2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/(a*b^2 - a^3) + (b*log(32*a*b^2*exp(x) + 32*a^2*b*exp(x) + 32*a*b*(b^2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/(a*b^2 - a^3)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.85

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx = \frac{-2\sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b + \log(e^x - 1) a^2 - \log(e^x - 1) b^2 - \log(e^x + 1) a^2 + \log(e^x + 1) b^2}{a(a^2 - b^2)}$$

input `int(csch(x)/(a+b*tanh(x)),x)`output `( - 2*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b + log(e**x - 1)*a**2 - log(e**x - 1)*b**2 - log(e**x + 1)*a**2 + log(e**x + 1)*b**2)/(a*(a**2 - b**2))`

### 3.85 $\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx$

Optimal result	695
Mathematica [A] (verified)	695
Rubi [A] (verified)	696
Maple [A] (verified)	697
Fricas [B] (verification not implemented)	698
Sympy [F]	698
Maxima [B] (verification not implemented)	699
Giac [B] (verification not implemented)	699
Mupad [B] (verification not implemented)	700
Reduce [B] (verification not implemented)	700

#### Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx = -\frac{\operatorname{coth}(x)}{a} - \frac{b \log(\tanh(x))}{a^2} + \frac{b \log(a+b \tanh(x))}{a^2}$$

output `-coth(x)/a-b*ln(tanh(x))/a^2+b*ln(a+b*tanh(x))/a^2`

#### Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx = -\frac{a \operatorname{coth}(x) + b \log(\sinh(x)) - b \log(a \cosh(x) + b \sinh(x))}{a^2}$$

input `Integrate[Csch[x]^2/(a + b*Tanh[x]),x]`

output `-((a*Coth[x] + b*Log[Sinh[x]] - b*Log[a*Cosh[x] + b*Sinh[x]))/a^2)`



**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 25, 3999, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\sin(ix)^2(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\sin(ix)^2(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{3999} \\
 & b \int \frac{\operatorname{coth}^2(x)}{b^2(a + b \tanh(x))} d(b \tanh(x)) \\
 & \quad \downarrow \text{54} \\
 & b \int \left( \frac{\operatorname{coth}^2(x)}{ab^2} - \frac{\operatorname{coth}(x)}{a^2b} + \frac{1}{a^2(a + b \tanh(x))} \right) d(b \tanh(x)) \\
 & \quad \downarrow \text{2009} \\
 & b \left( -\frac{\log(b \tanh(x))}{a^2} + \frac{\log(a + b \tanh(x))}{a^2} - \frac{\operatorname{coth}(x)}{ab} \right)
 \end{aligned}$$

input

```
Int [Csch[x]^2/(a + b*Tanh[x]), x]
```

output

```
b*(-(Coth[x]/(a*b)) - Log[b*Tanh[x]]/a^2 + Log[a + b*Tanh[x]]/a^2)
```

## Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 54 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3999 `Int[sin[(e_) + (f_)*(x_)^(m_))*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.72

method	result	size
risch	$-\frac{2}{a(e^{2x}-1)} + \frac{b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2} - \frac{b \ln(e^{2x}-1)}{a^2}$	50
default	$-\frac{\tanh\left(\frac{x}{2}\right)}{2a} + \frac{b \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{a^2} - \frac{1}{2a \tanh\left(\frac{x}{2}\right)} - \frac{b \ln\left(\tanh\left(\frac{x}{2}\right)\right)}{a^2}$	56

input `int(csch(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output `-2/a/(exp(2*x)-1)+1/a^2*b*ln(exp(2*x)+(a-b)/(a+b))-1/a^2*b*ln(exp(2*x)-1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 122 vs.  $2(29) = 58$ .

Time = 0.14 (sec) , antiderivative size = 122, normalized size of antiderivative = 4.21

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 - b) \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 - b) \log\left(\frac{2(a \cosh(x) - b \sinh(x))}{\cosh(x) + \sinh(x)}\right) - 2a}{a^2 \cosh(x)^2 + 2a^2 \cosh(x) \sinh(x) + a^2 \sinh(x)^2 - a^2}$$

input `integrate(csch(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`

output `((b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - (b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*log(2*sinh(x)/(cosh(x) + sinh(x))) - 2*a)/(a^2*cosh(x)^2 + 2*a^2*cosh(x)*sinh(x) + a^2*sinh(x)^2 - a^2)`

**Sympy [F]**

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$$

input `integrate(csch(x)**2/(a+b*tanh(x)),x)`

output `Integral(csch(x)**2/(a + b*tanh(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(29) = 58$ .

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.24

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx = \frac{b \log(-(a-b)e^{(-2x)} - a - b)}{a^2} - \frac{b \log(e^{(-x)} + 1)}{a^2} - \frac{b \log(e^{(-x)} - 1)}{a^2} + \frac{2}{ae^{(-2x)} - a}$$

input `integrate(csch(x)^2/(a+b*tanh(x)),x, algorithm="maxima")`

output `b*log(-(a - b)*e^(-2*x) - a - b)/a^2 - b*log(e^(-x) + 1)/a^2 - b*log(e^(-x) - 1)/a^2 + 2/(a*e^(-2*x) - a)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 78 vs.  $2(29) = 58$ .

Time = 0.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.69

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx = \frac{(ab + b^2) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^3 + a^2b} - \frac{b \log(|e^{(2x)} - 1|)}{a^2} + \frac{be^{(2x)} - 2a - b}{a^2(e^{(2x)} - 1)}$$

input `integrate(csch(x)^2/(a+b*tanh(x)),x, algorithm="giac")`

output `(a*b + b^2)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^3 + a^2*b) - b*log(abs(e^(2*x) - 1))/a^2 + (b*e^(2*x) - 2*a - b)/(a^2*(e^(2*x) - 1))`

**Mupad [B] (verification not implemented)**

Time = 2.44 (sec) , antiderivative size = 323, normalized size of antiderivative = 11.14

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{2 \operatorname{atan}\left(\frac{b(a^4(b^2)^{3/2} - a^6\sqrt{b^2})}{(ab^5\sqrt{-a^4-b^6}\sqrt{-a^4+a^2}b^4\sqrt{-a^4-a^3}b^3\sqrt{-a^4+b^6}e^{2x}\sqrt{-a^4-2a^2}b^4e^{2x}\sqrt{-a^4+a^4}b^2e^{2x}\sqrt{-a^4})+b^2}\right)}{-a^{12}b^4+3a^{10}b^6-3a^8b^8+a^6b^{10}} + \frac{2}{a(e^{2x}-1)}$$

input `int(1/(sinh(x)^2*(a + b*tanh(x))),x)`

output

```
(2*atan((b*(a^4*(b^2)^(3/2) - a^6*(b^2)^(1/2))*(a*b^5*(-a^4)^(1/2) - b^6*(-a^4)^(1/2) + a^2*b^4*(-a^4)^(1/2) - a^3*b^3*(-a^4)^(1/2) + b^6*exp(2*x)*(-a^4)^(1/2) - 2*a^2*b^4*exp(2*x)*(-a^4)^(1/2) + a^4*b^2*exp(2*x)*(-a^4)^(1/2)) + b^2*(a^3*(b^2)^(3/2) - a^5*(b^2)^(1/2))*(a*b^5*(-a^4)^(1/2) - b^6*(-a^4)^(1/2) + a^2*b^4*(-a^4)^(1/2) - a^3*b^3*(-a^4)^(1/2) + b^6*exp(2*x)*(-a^4)^(1/2) - 2*a^2*b^4*exp(2*x)*(-a^4)^(1/2) + a^4*b^2*exp(2*x)*(-a^4)^(1/2)))/(a^6*b^10 - 3*a^8*b^8 + 3*a^10*b^6 - a^12*b^4))*(b^2)^(1/2)/(-a^4)^(1/2) - 2/(a*(exp(2*x) - 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 116, normalized size of antiderivative = 4.00

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{-e^{2x}\log(e^x - 1)b - e^{2x}\log(e^x + 1)b + e^{2x}\log(e^{2x}a + e^{2x}b + a - b)b - 2e^{2x}a + \log(e^x - 1)b + \log(e^x + 1)b}{a^2(e^{2x} - 1)}$$

input `int(csch(x)^2/(a+b*tanh(x)),x)`

output

```
( - e**(2*x)*log(e**x - 1)*b - e**(2*x)*log(e**x + 1)*b + e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b - 2*e**(2*x)*a + log(e**x - 1)*b + log(e**x + 1)*b - log(e**(2*x)*a + e**(2*x)*b + a - b)*b)/(a**2*(e**(2*x) - 1))
```

### 3.86 $\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx$

Optimal result	701
Mathematica [A] (verified)	701
Rubi [C] (verified)	702
Maple [A] (verified)	704
Fricas [B] (verification not implemented)	704
Sympy [F]	705
Maxima [F(-2)]	706
Giac [A] (verification not implemented)	706
Mupad [B] (verification not implemented)	707
Reduce [B] (verification not implemented)	708

#### Optimal result

Integrand size = 13, antiderivative size = 82

$$\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx = \frac{b\sqrt{a^2-b^2} \arctan\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{a^3} + \frac{\operatorname{arctanh}(\cosh(x))}{2a} - \frac{b^2 \operatorname{arctanh}(\cosh(x))}{a^3} + \frac{b \operatorname{csch}(x)}{a^2} - \frac{\operatorname{coth}(x) \operatorname{csch}(x)}{2a}$$

output

```
b*(a^2-b^2)^(1/2)*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))/a^3+1/2*arctanh(cosh(x))/a-b^2*arctanh(cosh(x))/a^3+b*csch(x)/a^2-1/2*coth(x)*csch(x)/a
```

#### Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.79

$$\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx = \frac{-16\sqrt{a-b}\sqrt{a+b} \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}}\right) - 4ab \operatorname{coth}\left(\frac{x}{2}\right) + a^2 \operatorname{csch}^2\left(\frac{x}{2}\right) - 4a^2 \log\left(\cosh\left(\frac{x}{2}\right)\right) + 8b^2 \log\left(\cosh\left(\frac{x}{2}\right)\right)}{8a^3}$$

input

```
Integrate[Csch[x]^3/(a + b*Tanh[x]), x]
```

output

```
-1/8*(-16*sqrt[a - b]*b*sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(sqrt[a - b]*
sqrt[a + b])] - 4*a*b*Coth[x/2] + a^2*Csch[x/2]^2 - 4*a^2*Log[Cosh[x/2]] +
8*b^2*Log[Cosh[x/2]] + 4*a^2*Log[Sinh[x/2]] - 8*b^2*Log[Sinh[x/2]] + a^2*
Sech[x/2]^2 + 4*a*b*Tanh[x/2])/a^3
```

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.55 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.20, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3589, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i}{\sin(ix)^3(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{1}{\sin(ix)^3(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & -i \int \frac{i \coth(x) \operatorname{csch}^2(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{\coth(x) \operatorname{csch}^2(x)}{a \cosh(x) + b \sinh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \cos(ix)}{\sin(ix)^3(a \cos(ix) - ib \sin(ix))} dx \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\begin{aligned}
 & -i \int \frac{\cos(ix)}{\sin(ix)^3 (a \cos(ix) - ib \sin(ix))} dx \\
 & \quad \downarrow \text{3589} \\
 & -i \int \left( -\frac{i \operatorname{sech}^2(x) b^3}{a^3 (a \cosh(x) + b \sinh(x))} + \frac{i \operatorname{csch}(x) \operatorname{sech}^2(x) b^2}{a^3} - \frac{i \operatorname{csch}^2(x) \operatorname{sech}(x) b}{a^2} + \frac{i \operatorname{csch}^3(x)}{a} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -i \left( -\frac{i b^2 \operatorname{arctanh}(\cosh(x))}{a^3} + \frac{i b \operatorname{csch}(x)}{a^2} + \frac{i b \sqrt{a^2 - b^2} \operatorname{arctan}\left(\frac{a \sinh(x) + b \cosh(x)}{\sqrt{a^2 - b^2}}\right)}{a^3} + \frac{i \operatorname{arctanh}(\cosh(x))}{2a} - \frac{i \operatorname{coth}(x)}{2a} \right)
 \end{aligned}$$

input `Int [Csch[x]^3/(a + b*Tanh[x]), x]`

output `(-I)*((I*b*Sqrt[a^2 - b^2]*ArcTan[(b*Cosh[x] + a*Sinh[x])/Sqrt[a^2 - b^2]])/a^3 + ((I/2)*ArcTanh[Cosh[x]])/a - (I*b^2*ArcTanh[Cosh[x]])/a^3 + (I*b*Csch[x])/a^2 - ((I/2)*Coth[x]*Csch[x])/a`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3589 `Int[(cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.))/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]`



rule 4001

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

**Maple [A] (verified)**

Time = 1.21 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.34

method	result
default	$\frac{\frac{\tanh\left(\frac{x}{2}\right)^2 a - 2b \tanh\left(\frac{x}{2}\right)}{4a^2} - \frac{1}{8a \tanh\left(\frac{x}{2}\right)^2} + \frac{(-2a^2 + 4b^2) \ln\left(\tanh\left(\frac{x}{2}\right)\right)}{4a^3} + \frac{b}{2a^2 \tanh\left(\frac{x}{2}\right)} + \frac{2b\sqrt{a^2 - b^2} \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{a^3}$
risch	$-\frac{e^x (e^{2x} a - 2e^{2x} b + a + 2b)}{(e^{2x} - 1)^2 a^2} - \frac{\ln(e^x - 1)}{2a} + \frac{\ln(e^x - 1)b^2}{a^3} + \frac{\ln(e^x + 1)}{2a} - \frac{\ln(e^x + 1)b^2}{a^3} + \frac{\sqrt{-a^2 + b^2} b \ln\left(e^x + \frac{\sqrt{-a^2 + b^2}}{a + b}\right)}{a^3} - \sqrt{-a^2 + b^2}$

input

```
int(csch(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
1/4/a^2*(1/2*tanh(1/2*x)^2*a-2*b*tanh(1/2*x))-1/8/a/tanh(1/2*x)^2+1/4/a^3*(-2*a^2+4*b^2)*ln(tanh(1/2*x))+1/2/a^2*b/tanh(1/2*x)+2*b*(a^2-b^2)^(1/2)/a^3*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 555 vs. 2(74) = 148.

Time = 0.13 (sec) , antiderivative size = 1165, normalized size of antiderivative = 14.21

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```

[-1/2*(2*(a^2 - 2*a*b)*cosh(x)^3 + 6*(a^2 - 2*a*b)*cosh(x)*sinh(x)^2 + 2*(
a^2 - 2*a*b)*sinh(x)^3 - 2*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)
)^4 - 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - b
*cosh(x))*sinh(x) + b)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)
*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(
x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh
(x)^2 + a - b)) + 2*(a^2 + 2*a*b)*cosh(x) - ((a^2 - 2*b^2)*cosh(x)^4 + 4*(
a^2 - 2*b^2)*cosh(x)*sinh(x)^3 + (a^2 - 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)
*cosh(x)^2 + 2*(3*(a^2 - 2*b^2)*cosh(x)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 -
2*b^2 + 4*((a^2 - 2*b^2)*cosh(x)^3 - (a^2 - 2*b^2)*cosh(x))*sinh(x))*log(
cosh(x) + sinh(x) + 1) + ((a^2 - 2*b^2)*cosh(x)^4 + 4*(a^2 - 2*b^2)*cosh(x)
)*sinh(x)^3 + (a^2 - 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)*cosh(x)^2 + 2*(3*(
a^2 - 2*b^2)*cosh(x)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 - 2*b^2 + 4*((a^2 -
2*b^2)*cosh(x)^3 - (a^2 - 2*b^2)*cosh(x))*sinh(x))*log(cosh(x) + sinh(x) -
1) + 2*(3*(a^2 - 2*a*b)*cosh(x)^2 + a^2 + 2*a*b)*sinh(x))/(a^3*cosh(x)^4
+ 4*a^3*cosh(x)*sinh(x)^3 + a^3*sinh(x)^4 - 2*a^3*cosh(x)^2 + a^3 + 2*(3*a
^3*cosh(x)^2 - a^3)*sinh(x)^2 + 4*(a^3*cosh(x)^3 - a^3*cosh(x))*sinh(x)),
-1/2*(2*(a^2 - 2*a*b)*cosh(x)^3 + 6*(a^2 - 2*a*b)*cosh(x)*sinh(x)^2 + 2*(a
^2 - 2*a*b)*sinh(x)^3 + 4*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)
)^4 - 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - ...

```

## Sympy [F]

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx$$

input

```
integrate(csch(x)**3/(a+b*tanh(x)), x)
```

output

```
Integral(csch(x)**3/(a + b*tanh(x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.52

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx = \frac{(a^2 - 2b^2) \log(e^x + 1)}{2a^3} - \frac{(a^2 - 2b^2) \log(|e^x - 1|)}{2a^3} + \frac{2(a^2b - b^3) \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}a^3} - \frac{ae^{3x} - 2be^{3x} + ae^x + 2be^x}{a^2(e^{2x} - 1)^2}$$

input `integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="giac")`

output `1/2*(a^2 - 2*b^2)*log(e^x + 1)/a^3 - 1/2*(a^2 - 2*b^2)*log(abs(e^x - 1))/a^3 + 2*(a^2*b - b^3)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*a^3) - (a*e^(3*x) - 2*b*e^(3*x) + a*e^x + 2*b*e^x)/(a^2*(e^(2*x) - 1)^2)`

**Mupad [B] (verification not implemented)**

Time = 2.48 (sec) , antiderivative size = 506, normalized size of antiderivative = 6.17

$$\begin{aligned}
& \int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx \\
&= \frac{\ln(8a^3b - 16ab^3 - 4a^4 + 8b^4 + 4a^2b^2 - 4a^4e^x + 8b^4e^x - 16ab^3e^x + 8a^3be^x + 4a^2b^2e^x)}{2a} \\
&\quad - \frac{2e^x}{a - 2ae^{2x} + ae^{4x}} \\
&\quad - \frac{\ln(16ab^3 - 8a^3b + 4a^4 - 8b^4 - 4a^2b^2 - 4a^4e^x + 8b^4e^x - 16ab^3e^x + 8a^3be^x + 4a^2b^2e^x)}{2a} \\
&\quad - \frac{b^2 \ln(8a^3b - 16ab^3 - 4a^4 + 8b^4 + 4a^2b^2 - 4a^4e^x + 8b^4e^x - 16ab^3e^x + 8a^3be^x + 4a^2b^2e^x)}{a^3} \\
&\quad + \frac{b^2 \ln(16ab^3 - 8a^3b + 4a^4 - 8b^4 - 4a^2b^2 - 4a^4e^x + 8b^4e^x - 16ab^3e^x + 8a^3be^x + 4a^2b^2e^x)}{a^3} \\
&\quad - \frac{ae^x}{a^2e^{2x} - a^2} + \frac{2be^x}{a^2e^{2x} - a^2} \\
&\quad - \frac{b \ln(8b^2\sqrt{b^2 - a^2} - 8b^3e^x + 8a^2be^x - 8ab\sqrt{b^2 - a^2})\sqrt{b^2 - a^2}}{a^3} \\
&\quad + \frac{b \ln(8b^2\sqrt{b^2 - a^2} + 8b^3e^x - 8a^2be^x - 8ab\sqrt{b^2 - a^2})\sqrt{b^2 - a^2}}{a^3}
\end{aligned}$$

input `int(1/(sinh(x)^3*(a + b*tanh(x))),x)`

output

```

log(8*a^3*b - 16*a*b^3 - 4*a^4 + 8*b^4 + 4*a^2*b^2 - 4*a^4*exp(x) + 8*b^4*
exp(x) - 16*a*b^3*exp(x) + 8*a^3*b*exp(x) + 4*a^2*b^2*exp(x))/(2*a) - (2*
exp(x))/(a - 2*a*exp(2*x) + a*exp(4*x)) - log(16*a*b^3 - 8*a^3*b + 4*a^4 -
8*b^4 - 4*a^2*b^2 - 4*a^4*exp(x) + 8*b^4*exp(x) - 16*a*b^3*exp(x) + 8*a^3*
b*exp(x) + 4*a^2*b^2*exp(x))/(2*a) - (b^2*log(8*a^3*b - 16*a*b^3 - 4*a^4 +
8*b^4 + 4*a^2*b^2 - 4*a^4*exp(x) + 8*b^4*exp(x) - 16*a*b^3*exp(x) + 8*a^3
*b*exp(x) + 4*a^2*b^2*exp(x)))/a^3 + (b^2*log(16*a*b^3 - 8*a^3*b + 4*a^4 -
8*b^4 - 4*a^2*b^2 - 4*a^4*exp(x) + 8*b^4*exp(x) - 16*a*b^3*exp(x) + 8*a^3
*b*exp(x) + 4*a^2*b^2*exp(x)))/a^3 - (a*exp(x))/(a^2*exp(2*x) - a^2) + (2*
b*exp(x))/(a^2*exp(2*x) - a^2) - (b*log(8*b^2*(b^2 - a^2)^(1/2) - 8*b^3*ex
p(x) + 8*a^2*b*exp(x) - 8*a*b*(b^2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/a^3 +
(b*log(8*b^2*(b^2 - a^2)^(1/2) + 8*b^3*exp(x) - 8*a^2*b*exp(x) - 8*a*b*(b^
2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/a^3

```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 350, normalized size of antiderivative = 4.27

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx$$

$$= \frac{4e^{4x} \sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b - 8e^{2x} \sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b + 4\sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b - e^{4x} \log(e^x a + e^x b)}{1}$$

input `int(csch(x)^3/(a+b*tanh(x)),x)`output

```
(4***e**(4*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b
- 8***e**(2*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b
+ 4*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b - e**(4
*x)*log(e**x - 1)*a**2 + 2*e**(4*x)*log(e**x - 1)*b**2 + e**(4*x)*log(e**x
+ 1)*a**2 - 2*e**(4*x)*log(e**x + 1)*b**2 - 2*e**(3*x)*a**2 + 4*e**(3*x)*
a*b + 2*e**(2*x)*log(e**x - 1)*a**2 - 4*e**(2*x)*log(e**x - 1)*b**2 - 2*e
*(2*x)*log(e**x + 1)*a**2 + 4*e**(2*x)*log(e**x + 1)*b**2 - 2*e**x*a**2 -
4*e**x*a*b - log(e**x - 1)*a**2 + 2*log(e**x - 1)*b**2 + log(e**x + 1)*a**
2 - 2*log(e**x + 1)*b**2)/(2*a**3*(e**(4*x) - 2*e**(2*x) + 1))
```

### 3.87 $\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx$

Optimal result	709
Mathematica [A] (verified)	709
Rubi [A] (verified)	710
Maple [A] (verified)	711
Fricas [B] (verification not implemented)	712
Sympy [F]	713
Maxima [B] (verification not implemented)	713
Giac [B] (verification not implemented)	714
Mupad [B] (verification not implemented)	714
Reduce [B] (verification not implemented)	715

#### Optimal result

Integrand size = 13, antiderivative size = 78

$$\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx = \frac{(a^2 - b^2) \operatorname{coth}(x)}{a^3} + \frac{b \operatorname{coth}^2(x)}{2a^2} - \frac{\operatorname{coth}^3(x)}{3a} + \frac{b(a^2 - b^2) \log(\tanh(x))}{a^4} - \frac{b(a^2 - b^2) \log(a + b \tanh(x))}{a^4}$$

output

$$(a^2-b^2)*\operatorname{coth}(x)/a^3+1/2*b*\operatorname{coth}(x)^2/a^2-1/3*\operatorname{coth}(x)^3/a+b*(a^2-b^2)*\ln(\tanh(x))/a^4-b*(a^2-b^2)*\ln(a+b*\tanh(x))/a^4$$

#### Mathematica [A] (verified)

Time = 2.50 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx = \frac{3a^2b \operatorname{csch}^2(x) - 2 \operatorname{coth}(x) (-2a^3 + 3ab^2 + a^3 \operatorname{csch}^2(x)) + 6b(a^2 - b^2) (\log(\sinh(x)) - \log(a \cosh(x) + b \sinh(x)))}{6a^4}$$

input

```
Integrate[Csch[x]^4/(a + b*Tanh[x]), x]
```

output

$$(3a^2b\operatorname{Csch}[x]^2 - 2\operatorname{Coth}[x]*(-2a^3 + 3a*b^2 + a^3\operatorname{Csch}[x]^2) + 6b*(a^2 - b^2)*(Log[\operatorname{Sinh}[x]] - Log[a*\operatorname{Cosh}[x] + b*\operatorname{Sinh}[x]]))/(6a^4)$$
**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3999, 25, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{1}{\sin(ix)^4(a - ib \tan(ix))} dx \\ & \quad \downarrow 3999 \\ & -b \int -\frac{\operatorname{coth}^4(x) (b^2 - b^2 \tanh^2(x))}{b^4(a + b \tanh(x))} d(b \tanh(x)) \\ & \quad \downarrow 25 \\ & b \int \frac{\operatorname{coth}^4(x) (b^2 - b^2 \tanh^2(x))}{b^4(a + b \tanh(x))} d(b \tanh(x)) \\ & \quad \downarrow 522 \\ & b \int \left( \frac{\operatorname{coth}^4(x)}{ab^2} - \frac{\operatorname{coth}^3(x)}{a^2b} + \frac{(b^2 - a^2) \operatorname{coth}^2(x)}{a^3b^2} + \frac{(a^2 - b^2) \operatorname{coth}(x)}{a^4b} + \frac{b^2 - a^2}{a^4(a + b \tanh(x))} \right) d(b \tanh(x)) \\ & \quad \downarrow 2009 \\ & -b \left( -\frac{\operatorname{coth}^2(x)}{2a^2} - \frac{(a^2 - b^2) \log(b \tanh(x))}{a^4} + \frac{(a^2 - b^2) \log(a + b \tanh(x))}{a^4} - \frac{(a^2 - b^2) \operatorname{coth}(x)}{a^3b} + \frac{\operatorname{coth}^3(x)}{3ab} \right) \end{aligned}$$

input

$$\operatorname{Int}[\operatorname{Csch}[x]^4/(a + b*\operatorname{Tanh}[x]), x]$$

output

$$-(b*(-((a^2 - b^2)*\text{Coth}[x])/(a^3*b)) - \text{Coth}[x]^2/(2*a^2) + \text{Coth}[x]^3/(3*a*b) - ((a^2 - b^2)*\text{Log}[b*\text{Tanh}[x]])/a^4 + ((a^2 - b^2)*\text{Log}[a + b*\text{Tanh}[x]])/a^4))$$
**Defintions of rubi rules used**

rule 25

$$\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$$

rule 522

$$\text{Int}[(e_*)*(x_)^{(m_*)}*((c_*) + (d_*)*(x_)^{(n_*)}*((a_*) + (b_*)*(x_)^2)^{(p_*)}), x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0]$$

rule 2009

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 3042

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3999

$$\text{Int}[\sin[(e_*) + (f_*)*(x_)]^{(m_*)}*((a_*) + (b_*)*\tan[(e_*) + (f_*)*(x_)]^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[b/f \text{ Subst}[\text{Int}[x^m*((a + x)^n/(b^2 + x^2)^{(m/2 + 1))}, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \ \&\& \ \text{IntegerQ}[m/2]$$
**Maple [A] (verified)**

Time = 2.56 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.83

method	result
risch	$-\frac{2(-3e^{4x}ab+3b^2e^{4x}+6e^{2x}a^2+3e^{2x}ab-6b^2e^{2x}-2a^2+3b^2)}{3a^3(e^{2x}-1)^3} - \frac{b \ln(e^{2x} + \frac{a-b}{a+b})}{a^2} + \frac{b^3 \ln(e^{2x} + \frac{a-b}{a+b})}{a^4} + \frac{b \ln(e^{2x}-1)}{a^2} - \frac{b^3 \ln(e^{2x}-1)}{a^4}$
default	$-\frac{\tanh(\frac{x}{2})^3 a^2}{3} - a \tanh(\frac{x}{2})^2 b - 3 \tanh(\frac{x}{2}) a^2 + 4b^2 \tanh(\frac{x}{2}) - \frac{1}{24a \tanh(\frac{x}{2})^3} - \frac{-3a^2+4b^2}{8a^3 \tanh(\frac{x}{2})} + \frac{b}{8a^2 \tanh(\frac{x}{2})^2} + \frac{b(a^2-b^2) \ln(\frac{x}{2})}{a^4}$

input

$$\text{int}(\text{csch}(x)^4/(a+b*\text{tanh}(x)), x, \text{method}=\_RETURNVERBOSE)$$



output

```
-2/3*(-3*exp(4*x)*a*b+3*b^2*exp(4*x)+6*exp(2*x)*a^2+3*exp(2*x)*a*b-6*b^2*exp(2*x)-2*a^2+3*b^2)/a^3/(exp(2*x)-1)^3-1/a^2*b*ln(exp(2*x)+(a-b)/(a+b))+1/a^4*b^3*ln(exp(2*x)+(a-b)/(a+b))+1/a^2*b*ln(exp(2*x)-1)-1/a^4*b^3*ln(exp(2*x)-1)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 912 vs.  $2(74) = 148$ .

Time = 0.10 (sec) , antiderivative size = 912, normalized size of antiderivative = 11.69

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
1/3*(6*(a^2*b - a*b^2)*cosh(x)^4 + 24*(a^2*b - a*b^2)*cosh(x)*sinh(x)^3 + 6*(a^2*b - a*b^2)*sinh(x)^4 + 4*a^3 - 6*a*b^2 - 6*(2*a^3 + a^2*b - 2*a*b^2)*cosh(x)^2 - 6*(2*a^3 + a^2*b - 2*a*b^2 - 6*(a^2*b - a*b^2)*cosh(x)^2)*sinh(x)^2 - 3*((a^2*b - b^3)*cosh(x)^6 + 6*(a^2*b - b^3)*cosh(x)*sinh(x)^5 + (a^2*b - b^3)*sinh(x)^6 - 3*(a^2*b - b^3)*cosh(x)^4 - 3*(a^2*b - b^3 - 5*(a^2*b - b^3)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^2*b - b^3)*cosh(x)^3 - 3*(a^2*b - b^3)*cosh(x))*sinh(x)^3 - a^2*b + b^3 + 3*(a^2*b - b^3)*cosh(x)^2 + 3*(5*(a^2*b - b^3)*cosh(x)^4 + a^2*b - b^3 - 6*(a^2*b - b^3)*cosh(x)^2)*sinh(x)^2 + 6*((a^2*b - b^3)*cosh(x)^5 - 2*(a^2*b - b^3)*cosh(x)^3 + (a^2*b - b^3)*cosh(x))*sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + 3*((a^2*b - b^3)*cosh(x)^6 + 6*(a^2*b - b^3)*cosh(x)*sinh(x)^5 + (a^2*b - b^3)*sinh(x)^6 - 3*(a^2*b - b^3)*cosh(x)^4 - 3*(a^2*b - b^3 - 5*(a^2*b - b^3)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^2*b - b^3)*cosh(x)^3 - 3*(a^2*b - b^3)*cosh(x))*sinh(x)^3 - a^2*b + b^3 + 3*(a^2*b - b^3)*cosh(x)^2 + 3*(5*(a^2*b - b^3)*cosh(x)^4 + a^2*b - b^3 - 6*(a^2*b - b^3)*cosh(x)^2)*sinh(x)^2 + 6*((a^2*b - b^3)*cosh(x)^5 - 2*(a^2*b - b^3)*cosh(x)^3 + (a^2*b - b^3)*cosh(x))*sinh(x))*log(2*sinh(x)/(cosh(x) - sinh(x))) + 12*(2*(a^2*b - a*b^2)*cosh(x)^3 - (2*a^3 + a^2*b - 2*a*b^2)*cosh(x))*sinh(x))/(a^4*cosh(x)^6 + 6*a^4*cosh(x)*sinh(x)^5 + a^4*sinh(x)^6 - 3*a^4*cosh(x)^4 + 3*a^4*cosh(x)^2 + 3*(5*a^4*cosh(x)^2 - a^4)*sinh(x)^4 - a^4 + 4*(5*a^4*cosh(x)^3 - 3...
```

**Sympy [F]**

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx$$

input `integrate(csch(x)**4/(a+b*tanh(x)),x)`

output `Integral(csch(x)**4/(a + b*tanh(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 161 vs.  $2(74) = 148$ .

Time = 0.05 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.06

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx = -\frac{2(2a^2 - 3b^2 - 3(2a^2 - ab - 2b^2)e^{-2x}) - 3(ab + b^2)e^{-4x}}{3(3a^3e^{-2x} - 3a^3e^{-4x} + a^3e^{-6x} - a^3)} - \frac{(a^2b - b^3) \log(-(a - b)e^{-2x} - a - b)}{a^4} + \frac{(a^2b - b^3) \log(e^{-x} + 1)}{a^4} + \frac{(a^2b - b^3) \log(e^{-x} - 1)}{a^4}$$

input `integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

output `-2/3*(2*a^2 - 3*b^2 - 3*(2*a^2 - a*b - 2*b^2)*e^(-2*x) - 3*(a*b + b^2)*e^(-4*x))/(3*a^3*e^(-2*x) - 3*a^3*e^(-4*x) + a^3*e^(-6*x) - a^3) - (a^2*b - b^3)*log(-(a - b)*e^(-2*x) - a - b)/a^4 + (a^2*b - b^3)*log(e^(-x) + 1)/a^4 + (a^2*b - b^3)*log(e^(-x) - 1)/a^4`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(74) = 148.

Time = 0.13 (sec) , antiderivative size = 202, normalized size of antiderivative = 2.59

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx$$

$$= -\frac{(a^3b + a^2b^2 - ab^3 - b^4) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^5 + a^4b} + \frac{(a^2b - b^3) \log(|e^{(2x)} - 1|)}{a^4}$$

$$- \frac{11a^2be^{(6x)} - 11b^3e^{(6x)} - 45a^2be^{(4x)} + 12ab^2e^{(4x)} + 33b^3e^{(4x)} + 24a^3e^{(2x)} + 45a^2be^{(2x)} - 24ab^2e^{(2x)}}{6a^4(e^{(2x)} - 1)^3}$$

input `integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="giac")`

output 
$$\begin{aligned} & -(a^3b + a^2b^2 - ab^3 - b^4) \cdot \log(\operatorname{abs}(a \cdot e^{(2x)} + b \cdot e^{(2x)} + a - b)) / ( \\ & a^5 + a^4b) + (a^2b - b^3) \cdot \log(\operatorname{abs}(e^{(2x)} - 1)) / a^4 - 1/6 \cdot (11a^2b \cdot e^{(6x)} - \\ & 11b^3 \cdot e^{(6x)} - 45a^2b \cdot e^{(4x)} + 12a \cdot b^2 \cdot e^{(4x)} + 33b^3 \cdot e^{(4x)} \\ & + 24a^3 \cdot e^{(2x)} + 45a^2b \cdot e^{(2x)} - 24a \cdot b^2 \cdot e^{(2x)} - 33b^3 \cdot e^{(2x)} \\ & - 8a^3 - 11a^2b + 12ab^2 + 11b^3) / (a^4 \cdot (e^{(2x)} - 1)^3) \end{aligned}$$

**Mupad [B] (verification not implemented)**

Time = 2.32 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.58

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx = \frac{2b(a-b)}{a^3(e^{2x}-1)} - \frac{2(2a-b)}{a^2(e^{4x}-2e^{2x}+1)}$$

$$- \frac{3a(3e^{2x}-3e^{4x}+e^{6x}-1)}{8}$$

$$- \frac{b \ln(a-b+ae^{2x}+be^{2x})(a+b)(a-b)}{a^4}$$

$$+ \frac{b \ln(e^{2x}-1)(a+b)(a-b)}{a^4}$$

input `int(1/(sinh(x)^4*(a + b*tanh(x))),x)`

output

$$\frac{(2*b*(a - b))/(a^3*(\exp(2*x) - 1)) - (2*(2*a - b))/(a^2*(\exp(4*x) - 2*\exp(2*x) + 1)) - 8/(3*a*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (b*\log(a - b + a*\exp(2*x) + b*\exp(2*x))*(a + b)*(a - b))/a^4 + (b*\log(\exp(2*x) - 1)*(a + b)*(a - b))/a^4}$$
**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 567, normalized size of antiderivative = 7.27

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx$$

$$= \frac{-9e^{2x} \log(e^{2x}a + e^{2x}b + a - b) a^2 b - 4a b^2 - 2e^{6x} a b^2 - 9e^{4x} \log(e^x - 1) a^2 b - 9e^{4x} \log(e^x + 1) a^2 b + 9e^{2x} \log(e^x - 1) a^2 b + 9e^{2x} \log(e^x + 1) a^2 b + 9e^{2x} \log(e^x - 1) a^2 b + 9e^{2x} \log(e^x + 1) a^2 b}{(3*a*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (2*(2*a - b))/(a^2*(\exp(4*x) - 2*\exp(2*x) + 1)) - 8/(3*a*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (b*\log(a - b + a*\exp(2*x) + b*\exp(2*x))*(a + b)*(a - b))/a^4 + (b*\log(\exp(2*x) - 1)*(a + b)*(a - b))/a^4}$$

input

`int(csch(x)^4/(a+b*tanh(x)),x)`

output

$$\frac{(3*e^{6*x}*\log(e^{2*x} - 1)*a^{**2}*b - 3*e^{6*x}*\log(e^{2*x} - 1)*b^{**3} + 3*e^{6*x}*\log(e^{2*x} + 1)*a^{**2}*b - 3*e^{6*x}*\log(e^{2*x} + 1)*b^{**3} - 3*e^{6*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*a^{**2}*b + 3*e^{6*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*b^{**3} + 2*e^{6*x}*a^{**2}*b - 2*e^{6*x}*a*b^{**2} - 9*e^{4*x}*\log(e^{2*x} - 1)*a^{**2}*b + 9*e^{4*x}*\log(e^{2*x} - 1)*b^{**3} - 9*e^{4*x}*\log(e^{2*x} + 1)*a^{**2}*b + 9*e^{4*x}*\log(e^{2*x} + 1)*b^{**3} + 9*e^{4*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*a^{**2}*b - 9*e^{4*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*b^{**3} + 9*e^{2*x}*\log(e^{2*x} - 1)*a^{**2}*b - 9*e^{2*x}*\log(e^{2*x} - 1)*b^{**3} + 9*e^{2*x}*\log(e^{2*x} + 1)*a^{**2}*b - 9*e^{2*x}*\log(e^{2*x} + 1)*b^{**3} - 9*e^{2*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*a^{**2}*b + 9*e^{2*x}*\log(e^{2*x})*a + e^{2*x}*b + a - b)*b^{**3} - 12*e^{2*x}*a^{**3} + 6*e^{2*x}*a*b^{**2} - 3*\log(e^{2*x} - 1)*a^{**2}*b + 3*\log(e^{2*x} - 1)*b^{**3} - 3*\log(e^{2*x} + 1)*a^{**2}*b + 3*\log(e^{2*x} + 1)*b^{**3} + 3*\log(e^{2*x})*a + e^{2*x}*b + a - b)*a^{**2}*b - 3*\log(e^{2*x})*a + e^{2*x}*b + a - b)*b^{**3} + 4*a^{**3} - 2*a^{**2}*b - 4*a*b^{**2})/(3*a^{**4}*(e^{6*x} - 3*e^{4*x} + 3*e^{2*x} - 1))$$

### 3.88 $\int \frac{\operatorname{csch}^5(x)}{a+b \tanh(x)} dx$

Optimal result	716
Mathematica [A] (verified)	717
Rubi [C] (verified)	717
Maple [A] (verified)	720
Fricas [B] (verification not implemented)	720
Sympy [F]	721
Maxima [F(-2)]	721
Giac [A] (verification not implemented)	721
Mupad [B] (verification not implemented)	722
Reduce [B] (verification not implemented)	723

#### Optimal result

Integrand size = 13, antiderivative size = 218

$$\int \frac{\operatorname{csch}^5(x)}{a+b \tanh(x)} dx = -\frac{b \arctan(\sinh(x))}{a^2} + \frac{b^3 \arctan(\sinh(x))}{a^4} + \frac{b(a^2 - b^2) \arctan(\sinh(x))}{a^4} - \frac{b(a^2 - b^2)^{3/2} \arctan\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{a^5} - \frac{3 \operatorname{arctanh}(\cosh(x))}{a^5} + \frac{3b^2 \operatorname{arctanh}(\cosh(x))}{2a^3} - \frac{b^4 \operatorname{arctanh}(\cosh(x))}{a^5} - \frac{8a}{a^2} \operatorname{csch}(x) + \frac{b^3 \operatorname{csch}(x)}{a^4} + \frac{3 \operatorname{coth}(x) \operatorname{csch}(x)}{8a} - \frac{b^2 \operatorname{coth}(x) \operatorname{csch}(x)}{2a^3} + \frac{b \operatorname{csch}^3(x)}{3a^2} - \frac{\operatorname{coth}(x) \operatorname{csch}^3(x)}{4a} - \frac{b^2 \operatorname{sech}(x)}{a^3} + \frac{b^4 \operatorname{sech}(x)}{a^5} + \frac{b^2(a^2 - b^2) \operatorname{sech}(x)}{a^5}$$

output

```
-b*arctan(sinh(x))/a^2+b^3*arctan(sinh(x))/a^4+b*(a^2-b^2)*arctan(sinh(x))
/a^4-b*(a^2-b^2)^(3/2)*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))/a^5-3
/8*arctanh(cosh(x))/a+3/2*b^2*arctanh(cosh(x))/a^3-b^4*arctanh(cosh(x))/a^
5-b*csch(x)/a^2+b^3*csch(x)/a^4+3/8*coth(x)*csch(x)/a-1/2*b^2*coth(x)*csch
(x)/a^3+1/3*b*csch(x)^3/a^2-1/4*coth(x)*csch(x)^3/a-b^2*sech(x)/a^3+b^4*se
ch(x)/a^5+b^2*(a^2-b^2)*sech(x)/a^5
```

**Mathematica [A] (verified)**

Time = 0.98 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.54

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx$$

$$= \frac{-384a^2\sqrt{a-b}b\sqrt{a+b} \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}}\right) + 384\sqrt{a-b}b^3\sqrt{a+b} \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}}\right) - 16ab(7a^2 - 6b^2) \operatorname{Coth}\left[\frac{x}{2}\right] + 6a^2(3a^2 - 4b^2) \operatorname{Csch}\left[\frac{x}{2}\right]^2 - 72a^4 \operatorname{Log}\left[\operatorname{Cosh}\left[\frac{x}{2}\right]\right] + 288a^2b^2 \operatorname{Log}\left[\operatorname{Cosh}\left[\frac{x}{2}\right]\right] - 192b^4 \operatorname{Log}\left[\operatorname{Cosh}\left[\frac{x}{2}\right]\right] + 72a^4 \operatorname{Log}\left[\operatorname{Sinh}\left[\frac{x}{2}\right]\right] - 288a^2b^2 \operatorname{Log}\left[\operatorname{Sinh}\left[\frac{x}{2}\right]\right] + 192b^4 \operatorname{Log}\left[\operatorname{Sinh}\left[\frac{x}{2}\right]\right] + 18a^4 \operatorname{Sech}\left[\frac{x}{2}\right]^2 - 24a^2b^2 \operatorname{Sech}\left[\frac{x}{2}\right]^2 + 3a^4 \operatorname{Sech}\left[\frac{x}{2}\right]^4 + 64a^3b \operatorname{Csch}\left[\frac{x}{2}\right]^3 \operatorname{Sinh}\left[\frac{x}{2}\right]^4 + a^3 \operatorname{Csch}\left[\frac{x}{2}\right]^4 (-3a + 4b \operatorname{Sinh}\left[\frac{x}{2}\right]) + 112a^3b \operatorname{Tanh}\left[\frac{x}{2}\right] - 96a^2b^3 \operatorname{Tanh}\left[\frac{x}{2}\right]}{(192a^5)}$$

input

```
Integrate[Csch[x]^5/(a + b*Tanh[x]), x]
```

output

```
(-384*a^2*Sqrt[a - b]*b*Sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])] + 384*Sqrt[a - b]*b^3*Sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])] - 16*a*b*(7*a^2 - 6*b^2)*Coth[x/2] + 6*a^2*(3*a^2 - 4*b^2)*Csch[x/2]^2 - 72*a^4*Log[Cosh[x/2]] + 288*a^2*b^2*Log[Cosh[x/2]] - 192*b^4*Log[Cosh[x/2]] + 72*a^4*Log[Sinh[x/2]] - 288*a^2*b^2*Log[Sinh[x/2]] + 192*b^4*Log[Sinh[x/2]] + 18*a^4*Sech[x/2]^2 - 24*a^2*b^2*Sech[x/2]^2 + 3*a^4*Sech[x/2]^4 + 64*a^3*b*Csch[x]^3*Sinh[x/2]^4 + a^3*Csch[x/2]^4*(-3*a + 4*b*Sinh[x]) + 112*a^3*b*Tanh[x/2] - 96*a^2*b^3*Tanh[x/2])/(192*a^5)
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.84 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.37, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$ , Rules used = {3042, 26, 4001, 26, 3042, 26, 3589, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{i}{\sin(ix)^5(a - ib \tan(ix))} dx$$

$$\begin{aligned}
& \downarrow 26 \\
& i \int \frac{1}{\sin(ix)^5(a - ib \tan(ix))} dx \\
& \downarrow 4001 \\
& i \int -\frac{i \coth(x) \operatorname{csch}^4(x)}{a \cosh(x) + b \sinh(x)} dx \\
& \downarrow 26 \\
& \int \frac{\coth(x) \operatorname{csch}^4(x)}{a \cosh(x) + b \sinh(x)} dx \\
& \downarrow 3042 \\
& \int \frac{i \cos(ix)}{\sin(ix)^5(a \cos(ix) - ib \sin(ix))} dx \\
& \downarrow 26 \\
& i \int \frac{\cos(ix)}{\sin(ix)^5(a \cos(ix) - ib \sin(ix))} dx \\
& \downarrow 3589 \\
& i \int \left( \frac{i \operatorname{sech}^4(x) b^5}{a^5(a \cosh(x) + b \sinh(x))} - \frac{i \operatorname{csch}(x) \operatorname{sech}^4(x) b^4}{a^5} + \frac{i \operatorname{csch}^2(x) \operatorname{sech}^3(x) b^3}{a^4} - \frac{i \operatorname{csch}^3(x) \operatorname{sech}^2(x) b^2}{a^3} + \frac{i \operatorname{csch}^4(x)}{a^2} \right) dx \\
& \downarrow 2009 \\
& i \left( \frac{ib^4 \operatorname{arctanh}(\cosh(x))}{a^5} - \frac{ib^4 \operatorname{sech}(x)}{a^5} - \frac{ib^3 \operatorname{arctan}(\sinh(x))}{a^4} - \frac{3ib^3 \operatorname{csch}(x)}{2a^4} + \frac{ib^3 \operatorname{csch}(x) \operatorname{sech}^2(x)}{2a^4} + \frac{ib^3 \tanh(x) \operatorname{sech}(x)}{2a^4} \right)
\end{aligned}$$

input

```
Int [Csch[x]^5/(a + b*Tanh[x]), x]
```

output

$$I*((I*b*ArcTan[Sinh[x]])/a^2 - (I*b^3*ArcTan[Sinh[x]])/a^4 - (I*b*(a^2 - b^2)*ArcTan[Sinh[x]])/a^4 + (I*b*(a^2 - b^2)^{(3/2)}*ArcTan[(b*Cosh[x] + a*Sinh[x])/Sqrt[a^2 - b^2]])/a^5 + (((3*I)/8)*ArcTanh[Cosh[x]])/a - (((3*I)/2)*b^2*ArcTanh[Cosh[x]])/a^3 + (I*b^4*ArcTanh[Cosh[x]])/a^5 + (I*b*Csch[x])/a^2 - (((3*I)/2)*b^3*Csch[x])/a^4 - (((3*I)/8)*Coth[x]*Csch[x])/a - ((I/3)*b*Csch[x]^3)/a^2 + ((I/4)*Coth[x]*Csch[x]^3)/a + (((3*I)/2)*b^2*Sech[x])/a^3 - (I*b^4*Sech[x])/a^5 - (I*b^2*(a^2 - b^2)*Sech[x])/a^5 + ((I/2)*b^2*Csch[x]^2*Sech[x])/a^3 + ((I/2)*b^3*Csch[x]*Sech[x]^2)/a^4 + ((I/2)*b^3*Sech[x]*Tanh[x])/a^4)$$

### Defintions of rubi rules used

rule 26

$$\text{Int}[(\text{Complex}[0, a_])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$$

rule 2009

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 3042

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3589

$$\text{Int}[(\cos[(c_.) + (d_.)*(x_)]^{(m_.)} \sin[(c_.) + (d_.)*(x_)]^{(n_.)}) / (\cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.) \sin[(c_.) + (d_.)*(x_)]), x\_Symbol] \rightarrow \text{Int}[\text{ExpandTrig}[\cos[c + d*x]^m (\sin[c + d*x]^n / (a \cos[c + d*x] + b \sin[c + d*x])), x], x] \text{ ; FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{IntegersQ}[m, n]$$

rule 4001

$$\text{Int}[\sin[(e_.) + (f_.)*(x_)]^{(m_.)} ((a_.) + (b_.) \tan[(e_.) + (f_.)*(x_)]^{(n_.)}), x\_Symbol] \rightarrow \text{Int}[\text{Sin}[e + f*x]^m ((a \cos[e + f*x] + b \sin[e + f*x])^n / \cos[e + f*x]^n), x] \text{ ; FreeQ}\{a, b, e, f\}, x\} \ \&\& \ \text{IntegerQ}[(m - 1)/2] \ \&\& \ \text{ILtQ}[n, 0] \ \&\& \ ((\text{LtQ}[m, 5] \ \&\& \ \text{GtQ}[n, -4]) \ || \ (\text{EqQ}[m, 5] \ \&\& \ \text{EqQ}[n, -1]))$$



**Maple [A] (verified)**

Time = 6.11 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.04

method	result
default	$\frac{a^3 \tanh\left(\frac{x}{2}\right)^4}{4} - \frac{2b \tanh\left(\frac{x}{2}\right)^3 a^2}{3} - \frac{2a^3 \tanh\left(\frac{x}{2}\right)^2 + 2a b^2 \tanh\left(\frac{x}{2}\right)^2 + 10a^2 b \tanh\left(\frac{x}{2}\right) - 8b^3 \tanh\left(\frac{x}{2}\right)}{16a^4} - \frac{2b(a^4 - 2a^2 b^2 + b^4) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + b}{a^2 - b^2}\right)}{a^5 \sqrt{a^2 - b^2}}$
risch	$\frac{e^x (9a^3 e^{6x} - 24e^{6x} a^2 b - 12a b^2 e^{6x} + 24b^3 e^{6x} - 33a^3 e^{4x} + 104a^2 b e^{4x} + 12e^{4x} b^2 a - 72b^3 e^{4x} - 33a^3 e^{2x} - 104e^{2x} a^2 b + 12e^{2x} a b^2 + 72b^3 e^{2x})}{12a^4 (e^{2x} - 1)^4}$

input `int(csch(x)^5/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 1/16/a^4*(1/4*a^3*tanh(1/2*x)^4-2/3*b*tanh(1/2*x)^3*a^2-2*a^3*tanh(1/2*x)^2 \\ & +2*a*b^2*tanh(1/2*x)^2+10*a^2*b*tanh(1/2*x)-8*b^3*tanh(1/2*x))-2*b*(a^4-2 \\ & *a^2*b^2+b^4)/a^5/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2) \\ & ^{(1/2)})-1/64/a/tanh(1/2*x)^4-1/32*(-4*a^2+4*b^2)/a^3/tanh(1/2*x)^2+1/16/ \\ & a^5*(6*a^4-24*a^2*b^2+16*b^4)*ln(tanh(1/2*x))+1/24/a^2*b/tanh(1/2*x)^3-1/8 \\ & *b*(5*a^2-4*b^2)/a^4/tanh(1/2*x) \end{aligned}$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2646 vs. 2(202) = 404.

Time = 0.20 (sec) , antiderivative size = 5347, normalized size of antiderivative = 24.53

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="fricas")`

output Too large to include

**Sympy [F]**

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx$$

input `integrate(csch(x)**5/(a+b*tanh(x)),x)`

output `Integral(csch(x)**5/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.25

$$\begin{aligned} \int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = & -\frac{(3a^4 - 12a^2b^2 + 8b^4) \log(e^x + 1)}{8a^5} \\ & + \frac{(3a^4 - 12a^2b^2 + 8b^4) \log(|e^x - 1|)}{8a^5} - \frac{2(a^4b - 2a^2b^3 + b^5) \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}a^5} \\ & + \frac{9a^3e^{(7x)} - 24a^2be^{(7x)} - 12ab^2e^{(7x)} + 24b^3e^{(7x)} - 33a^3e^{(5x)} + 104a^2be^{(5x)} + 12ab^2e^{(5x)} - 72b^3e^{(5x)}}{12a^4(e^{(2x)} - } \end{aligned}$$

input `integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="giac")`

output `-1/8*(3*a^4 - 12*a^2*b^2 + 8*b^4)*log(e^x + 1)/a^5 + 1/8*(3*a^4 - 12*a^2*b^2 + 8*b^4)*log(abs(e^x - 1))/a^5 - 2*(a^4*b - 2*a^2*b^3 + b^5)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*a^5) + 1/12*(9*a^3*e^(7*x) - 24*a^2*b*e^(7*x) - 12*a*b^2*e^(7*x) + 24*b^3*e^(7*x) - 33*a^3*e^(5*x) + 104*a^2*b*e^(5*x) + 12*a*b^2*e^(5*x) - 72*b^3*e^(5*x) - 33*a^3*e^(3*x) - 104*a^2*b*e^(3*x) + 12*a*b^2*e^(3*x) + 72*b^3*e^(3*x) + 9*a^3*e^x + 24*a^2*b*e^x - 12*a*b^2*e^x - 24*b^3*e^x)/(a^4*(e^(2*x) - 1)^4)`

### Mupad [B] (verification not implemented)

Time = 4.34 (sec) , antiderivative size = 753, normalized size of antiderivative = 3.45

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `int(1/(sinh(x)^5*(a + b*tanh(x))),x)`

output

```
(log(exp(x) - 1)*(3*a^4 + 8*b^4 - 12*a^2*b^2))/(8*a^5) - (log(exp(x) + 1)*
(3*a^4 + 8*b^4 - 12*a^2*b^2))/(8*a^5) - (4*exp(x))/(a*(6*exp(4*x) - 4*exp(
2*x) - 4*exp(6*x) + exp(8*x) + 1)) - (2*exp(x)*(9*a - 4*b))/(3*a^2*(3*exp(
2*x) - 3*exp(4*x) + exp(6*x) - 1)) - (exp(x)*(3*a^2 - 16*a*b + 12*b^2))/(6
*a^3*(exp(4*x) - 2*exp(2*x) + 1)) - (exp(x)*(4*a*b^2 + 8*a^2*b - 3*a^3 - 8
*b^3))/(4*a^4*(exp(2*x) - 1)) + (b*log((b*exp(x)*(a - b)^2*(288*a*b^6 - 24
*a^6*b - 9*a^7 - 192*b^7 + 224*a^2*b^5 - 456*a^3*b^4 + 24*a^4*b^3 + 144*a^
5*b^2)))/(2*a^12*(a + b)) - (b*(a - b)*(-(a + b)^3*(a - b)^3)^(1/2)*(8*a^5*
b^3 - 9*a^8 - 9*a^7*b + 8*a^6*b^2 + 192*b^5*exp(x)*(-(a^2 - b^2)^3)^(1/2)
- 224*a^2*b^3*exp(x)*(-(a^2 - b^2)^3)^(1/2) - 88*a^3*b^2*exp(x)*(-(a^2 - b
^2)^3)^(1/2) + 96*a*b^4*exp(x)*(-(a^2 - b^2)^3)^(1/2) + 24*a^4*b*exp(x)*(-
(a^2 - b^2)^3)^(1/2)))/(2*a^12*(a + b)^4)*(-(a + b)^3*(a - b)^3)^(1/2))/a
^5 - (b*log((b*exp(x)*(a - b)^2*(288*a*b^6 - 24*a^6*b - 9*a^7 - 192*b^7 +
224*a^2*b^5 - 456*a^3*b^4 + 24*a^4*b^3 + 144*a^5*b^2)))/(2*a^12*(a + b)) -
(b*(a - b)*(-(a + b)^3*(a - b)^3)^(1/2)*(9*a^7*b + 9*a^8 - 8*a^5*b^3 - 8*a
^6*b^2 + 192*b^5*exp(x)*(-(a^2 - b^2)^3)^(1/2) - 224*a^2*b^3*exp(x)*(-(a^2
- b^2)^3)^(1/2) - 88*a^3*b^2*exp(x)*(-(a^2 - b^2)^3)^(1/2) + 96*a*b^4*exp
(x)*(-(a^2 - b^2)^3)^(1/2) + 24*a^4*b*exp(x)*(-(a^2 - b^2)^3)^(1/2)))/(2*a
^12*(a + b)^4)*(-(a + b)^3*(a - b)^3)^(1/2))/a^5
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 1133, normalized size of antiderivative = 5.20

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
int(csch(x)^5/(a+b*tanh(x)),x)
```

output

```
( - 48***8*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2)
)*a**2*b + 48***8*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2
- b**2))*b**3 + 192***6*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt
(a**2 - b**2))*a**2*b - 192***6*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x
*b)/sqrt(a**2 - b**2))*b**3 - 288***4*x)*sqrt(a**2 - b**2)*atan((e**x*a
+ e**x*b)/sqrt(a**2 - b**2))*a**2*b + 288***4*x)*sqrt(a**2 - b**2)*atan(
(e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**3 + 192***2*x)*sqrt(a**2 - b**2)
*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**2*b - 192***2*x)*sqrt(a**2
- b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**3 - 48*sqrt(a**2 - b
**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**2*b + 48*sqrt(a**2 - b**
2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**3 + 9***8*x)*log(e**x -
1)*a**4 - 36***8*x)*log(e**x - 1)*a**2*b**2 + 24***8*x)*log(e**x - 1)*
b**4 - 9***8*x)*log(e**x + 1)*a**4 + 36***8*x)*log(e**x + 1)*a**2*b**2
- 24***8*x)*log(e**x + 1)*b**4 + 18***7*x)*a**4 - 48***7*x)*a**3*b
- 24***7*x)*a**2*b**2 + 48***7*x)*a*b**3 - 36***6*x)*log(e**x - 1)*a
**4 + 144***6*x)*log(e**x - 1)*a**2*b**2 - 96***6*x)*log(e**x - 1)*b**
4 + 36***6*x)*log(e**x + 1)*a**4 - 144***6*x)*log(e**x + 1)*a**2*b**2
+ 96***6*x)*log(e**x + 1)*b**4 - 66***5*x)*a**4 + 208***5*x)*a**3*b
+ 24***5*x)*a**2*b**2 - 144***5*x)*a*b**3 + 54***4*x)*log(e**x - 1)*
a**4 - 216***4*x)*log(e**x - 1)*a**2*b**2 + 144***4*x)*log(e**x - 1...
```

### 3.89 $\int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx$

Optimal result . . . . .	725
Mathematica [A] (verified) . . . . .	725
Rubi [A] (verified) . . . . .	726
Maple [B] (verified) . . . . .	728
Fricas [B] (verification not implemented) . . . . .	728
Sympy [F] . . . . .	729
Maxima [B] (verification not implemented) . . . . .	730
Giac [B] (verification not implemented) . . . . .	730
Mupad [B] (verification not implemented) . . . . .	731
Reduce [B] (verification not implemented) . . . . .	732

#### Optimal result

Integrand size = 13, antiderivative size = 130

$$\int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx = -\frac{(a^2-b^2)^2 \operatorname{coth}(x)}{a^5} - \frac{b(2a^2-b^2) \operatorname{coth}^2(x)}{2a^4} + \frac{(2a^2-b^2) \operatorname{coth}^3(x)}{3a^3} + \frac{b \operatorname{coth}^4(x)}{4a^2} - \frac{\operatorname{coth}^5(x)}{5a} - \frac{b(a^2-b^2)^2 \log(\tanh(x))}{a^6} + \frac{b(a^2-b^2)^2 \log(a+b \tanh(x))}{a^6}$$

output

```
-(a^2-b^2)^2*coth(x)/a^5-1/2*b*(2*a^2-b^2)*coth(x)^2/a^4+1/3*(2*a^2-b^2)*c
oth(x)^3/a^3+1/4*b*coth(x)^4/a^2-1/5*coth(x)^5/a-b*(a^2-b^2)^2*ln(tanh(x))
/a^6+b*(a^2-b^2)^2*ln(a+b*tanh(x))/a^6
```

#### Mathematica [A] (verified)

Time = 5.44 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx = \frac{-4 \operatorname{coth}(x) (8a^5 - 25a^3b^2 + 15ab^4 + (-4a^5 + 5a^3b^2) \operatorname{csch}^2(x) + 3a^5 \operatorname{csch}^4(x)) + 15b(-2a^2(a^2 - b^2) \operatorname{csch}^2(x) + 3a^2 \operatorname{csch}^4(x))}{60a^6}$$

input `Integrate[Csch[x]^6/(a + b*Tanh[x]), x]`

output  $(-4*\text{Coth}[x]*(8*a^5 - 25*a^3*b^2 + 15*a*b^4 + (-4*a^5 + 5*a^3*b^2)*\text{Csch}[x]^2 + 3*a^5*\text{Csch}[x]^4) + 15*b*(-2*a^2*(a^2 - b^2)*\text{Csch}[x]^2 + a^4*\text{Csch}[x]^4 - 4*(a^2 - b^2)^2*(\text{Log}[\text{Sinh}[x]] - \text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])))/(60*a^6)$

### Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 25, 3999, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\text{csch}^6(x)}{a + b \tanh(x)} dx$$

$$\downarrow 3042$$

$$\int -\frac{1}{\sin(ix)^6(a - ib \tan(ix))} dx$$

$$\downarrow 25$$

$$-\int \frac{1}{\sin(ix)^6(a - ib \tan(ix))} dx$$

$$\downarrow 3999$$

$$b \int \frac{\text{coth}^6(x) (b^2 - b^2 \tanh^2(x))^2}{b^6(a + b \tanh(x))} d(b \tanh(x))$$

$$\downarrow 522$$

$$b \int \left( \frac{\text{coth}^6(x)}{ab^2} - \frac{\text{coth}^5(x)}{a^2b} + \frac{(b^4 - 2a^2b^2) \text{coth}^4(x)}{a^3b^4} + \frac{(2a^2b^2 - b^4) \text{coth}^3(x)}{a^4b^3} + \frac{(a^2 - b^2)^2 \text{coth}^2(x)}{a^5b^2} - \frac{(a^2 - b^2)}{a^6} \right) dx$$

$$\downarrow 2009$$

$$b \left( \frac{\coth^4(x)}{4a^2} - \frac{(a^2 - b^2)^2 \log(b \tanh(x))}{a^6} + \frac{(a^2 - b^2)^2 \log(a + b \tanh(x))}{a^6} - \frac{(a^2 - b^2)^2 \coth(x)}{a^5 b} - \frac{(2a^2 - b^2) \coth(x)}{2a^4} \right)$$

input `Int[Csch[x]^6/(a + b*Tanh[x]),x]`

output `b*(-((a^2 - b^2)^2*Coth[x])/(a^5*b)) - ((2*a^2 - b^2)*Coth[x]^2)/(2*a^4) + ((2*a^2 - b^2)*Coth[x]^3)/(3*a^3*b) + Coth[x]^4/(4*a^2) - Coth[x]^5/(5*a*b) - ((a^2 - b^2)^2*Log[b*Tanh[x]])/a^6 + ((a^2 - b^2)^2*Log[a + b*Tanh[x]])/a^6)`

### Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 522 `Int[((e_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3999 `Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b/f Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`



**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 276 vs.  $2(122) = 244$ .

Time = 12.85 (sec) , antiderivative size = 277, normalized size of antiderivative = 2.13

method	result
default	$-\frac{a^4 \tanh\left(\frac{x}{2}\right)^5}{5} - \frac{b \tanh\left(\frac{x}{2}\right)^4 a^3}{2} - \frac{5 \tanh\left(\frac{x}{2}\right)^3 a^4}{3} + \frac{4a^2 b^2 \tanh\left(\frac{x}{2}\right)^3}{3} + \frac{6a^3 b \tanh\left(\frac{x}{2}\right)^2 - 4b^3 \tanh\left(\frac{x}{2}\right)^2 a + 10a^4 \tanh\left(\frac{x}{2}\right) - 28a^2 b^2 \tanh\left(\frac{x}{2}\right)}{32a^5}$
risch	$-\frac{2(15a^3 b e^{8x} - 15a^2 b^2 e^{8x} - 15a b^3 e^{8x} + 15b^4 e^{8x} - 75b a^3 e^{6x} + 90a^2 b^2 e^{6x} + 45a b^3 e^{6x} - 60b^4 e^{6x} + 80e^{4x} a^4 + 75e^{4x} a^3 b - 160e^{4x} a^2 b^2 - 15a^5 (e^{2x} - 1)^5)}{15a^5 (e^{2x} - 1)^5}$

input `int(csch(x)^6/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output

```
-1/32/a^5*(1/5*a^4*tanh(1/2*x)^5-1/2*b*tanh(1/2*x)^4*a^3-5/3*tanh(1/2*x)^3
*a^4+4/3*a^2*b^2*tanh(1/2*x)^3+6*a^3*b*tanh(1/2*x)^2-4*b^3*tanh(1/2*x)^2*a
+10*a^4*tanh(1/2*x)-28*a^2*b^2*tanh(1/2*x)+16*b^4*tanh(1/2*x))-1/160/a/tan
h(1/2*x)^5-1/96*(-5*a^2+4*b^2)/a^3/tanh(1/2*x)^3-1/32/a^5*(10*a^4-28*a^2*b
^2+16*b^4)/tanh(1/2*x)+1/64/a^2*b/tanh(1/2*x)^4-1/16/a^4*b*(3*a^2-2*b^2)/t
anh(1/2*x)^2-1/a^6*b*(a^4-2*a^2*b^2+b^4)*ln(tanh(1/2*x))+2/a^6*b*(1/2*a^4-
a^2*b^2+1/2*b^4)*ln(tanh(1/2*x)^2*a+2*b*tanh(1/2*x)+a)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2972 vs.  $2(122) = 244$ .

Time = 0.14 (sec) , antiderivative size = 2972, normalized size of antiderivative = 22.86

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(csch(x)^6/(a+b*tanh(x)),x, algorithm="fricas")`

output

```

-1/15*(30*(a^4*b - a^3*b^2 - a^2*b^3 + a*b^4)*cosh(x)^8 + 240*(a^4*b - a^3
*b^2 - a^2*b^3 + a*b^4)*cosh(x)*sinh(x)^7 + 30*(a^4*b - a^3*b^2 - a^2*b^3
+ a*b^4)*sinh(x)^8 - 30*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4)*cosh(x
)^6 - 30*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4 - 28*(a^4*b - a^3*b^2
- a^2*b^3 + a*b^4)*cosh(x)^2)*sinh(x)^6 + 60*(28*(a^4*b - a^3*b^2 - a^2*b^
3 + a*b^4)*cosh(x)^3 - 3*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4)*cosh(
x))*sinh(x)^5 + 16*a^5 - 50*a^3*b^2 + 30*a*b^4 + 10*(16*a^5 + 15*a^4*b - 3
2*a^3*b^2 - 9*a^2*b^3 + 18*a*b^4)*cosh(x)^4 + 10*(16*a^5 + 15*a^4*b - 32*a
^3*b^2 - 9*a^2*b^3 + 18*a*b^4 + 210*(a^4*b - a^3*b^2 - a^2*b^3 + a*b^4)*co
sh(x)^4 - 45*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4)*cosh(x)^2)*sinh(x
)^4 + 40*(42*(a^4*b - a^3*b^2 - a^2*b^3 + a*b^4)*cosh(x)^5 - 15*(5*a^4*b -
6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4)*cosh(x)^3 + (16*a^5 + 15*a^4*b - 32*a^3*
b^2 - 9*a^2*b^3 + 18*a*b^4)*cosh(x))*sinh(x)^3 - 10*(8*a^5 + 3*a^4*b - 22*
a^3*b^2 - 3*a^2*b^3 + 12*a*b^4)*cosh(x)^2 + 10*(84*(a^4*b - a^3*b^2 - a^2*
b^3 + a*b^4)*cosh(x)^6 - 8*a^5 - 3*a^4*b + 22*a^3*b^2 + 3*a^2*b^3 - 12*a*b
^4 - 45*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a*b^4)*cosh(x)^4 + 6*(16*a^5
+ 15*a^4*b - 32*a^3*b^2 - 9*a^2*b^3 + 18*a*b^4)*cosh(x)^2)*sinh(x)^2 - 15*
((a^4*b - 2*a^2*b^3 + b^5)*cosh(x)^10 + 10*(a^4*b - 2*a^2*b^3 + b^5)*cosh(
x))*sinh(x)^9 + (a^4*b - 2*a^2*b^3 + b^5)*sinh(x)^10 - 5*(a^4*b - 2*a^2*b^3
+ b^5)*cosh(x)^8 - 5*(a^4*b - 2*a^2*b^3 + b^5 - 9*(a^4*b - 2*a^2*b^3 +...

```

## Sympy [F]

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx$$

input

```
integrate(csch(x)**6/(a+b*tanh(x)), x)
```

output

```
Integral(csch(x)**6/(a + b*tanh(x)), x)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 308 vs.  $2(122) = 244$ .

Time = 0.05 (sec) , antiderivative size = 308, normalized size of antiderivative = 2.37

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx$$

$$= \frac{2(8a^4 - 25a^2b^2 + 15b^4 - 5(8a^4 - 3a^3b - 22a^2b^2 + 3ab^3 + 12b^4)e^{-2x}) + 5(16a^4 - 15a^3b - 32a^2b^2 + 9ab^3 + 18b^4)e^{-4x} + 15(5a^5e^{-2x} - 10a^5e^{-4x} + 10a^5e^{-6x} - 5a^5e^{-8x})}{15(5a^5e^{-2x} - 10a^5e^{-4x} + 10a^5e^{-6x} - 5a^5e^{-8x})} + \frac{(a^4b - 2a^2b^3 + b^5) \log(-(a-b)e^{-2x} - a - b)}{a^6} - \frac{(a^4b - 2a^2b^3 + b^5) \log(e^{-x} + 1)}{a^6} - \frac{(a^4b - 2a^2b^3 + b^5) \log(e^{-x} - 1)}{a^6}$$

input `integrate(csch(x)^6/(a+b*tanh(x)),x, algorithm="maxima")`

output `2/15*(8*a^4 - 25*a^2*b^2 + 15*b^4 - 5*(8*a^4 - 3*a^3*b - 22*a^2*b^2 + 3*a*b^3 + 12*b^4)*e^(-2*x) + 5*(16*a^4 - 15*a^3*b - 32*a^2*b^2 + 9*a*b^3 + 18*b^4)*e^(-4*x) + 15*(5*a^3*b + 6*a^2*b^2 - 3*a*b^3 - 4*b^4)*e^(-6*x) - 15*(a^3*b + a^2*b^2 - a*b^3 - b^4)*e^(-8*x))/(5*a^5*e^(-2*x) - 10*a^5*e^(-4*x) + 10*a^5*e^(-6*x) - 5*a^5*e^(-8*x) + a^5*e^(-10*x) - a^5) + (a^4*b - 2*a^2*b^3 + b^5)*log(-(a - b)*e^(-2*x) - a - b)/a^6 - (a^4*b - 2*a^2*b^3 + b^5)*log(e^(-x) + 1)/a^6 - (a^4*b - 2*a^2*b^3 + b^5)*log(e^(-x) - 1)/a^6`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 412 vs.  $2(122) = 244$ .

Time = 0.13 (sec) , antiderivative size = 412, normalized size of antiderivative = 3.17

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx$$

$$= \frac{(a^5b + a^4b^2 - 2a^3b^3 - 2a^2b^4 + ab^5 + b^6) \log(|ae^{2x} + be^{2x} + a - b|)}{a^7 + a^6b} - \frac{(a^4b - 2a^2b^3 + b^5) \log(|e^{2x} - 1|)}{a^6} + \frac{137a^4be^{10x} - 274a^2b^3e^{10x} + 137b^5e^{10x} - 805a^4be^{8x} + 120a^3b^2e^{8x} + 1490a^2b^3e^{8x} - 120ab^4e^{8x} - 137a^4be^{6x} + 274a^2b^3e^{6x} - 137b^5e^{6x} - 805a^4be^{4x} + 120a^3b^2e^{4x} + 1490a^2b^3e^{4x} - 120ab^4e^{4x} - 137a^4be^{2x} + 274a^2b^3e^{2x} - 137b^5e^{2x} - 805a^4be^{0x} + 120a^3b^2e^{0x} + 1490a^2b^3e^{0x} - 120ab^4e^{0x}}{a^7 + a^6b}$$

input `integrate(csch(x)^6/(a+b*tanh(x)),x, algorithm="giac")`

output 
$$\begin{aligned} & (a^5b + a^4b^2 - 2a^3b^3 - 2a^2b^4 + ab^5 + b^6) \log(\text{abs}(a e^{2x} \\ & + b e^{2x} + a - b)) / (a^7 + a^6b) - (a^4b - 2a^2b^3 + b^5) \log(\text{abs}(e^{2x} - 1)) / a^6 \\ & + 1/60 * (137a^4b e^{10x} - 274a^2b^3 e^{10x} + 137b^5 e^{10x} - 805a^4b e^{8x} + 120a^3b^2 e^{8x} \\ & + 1490a^2b^3 e^{8x} - 120ab^4 e^{8x} - 685b^5 e^{8x} + 1970a^4b e^{6x} - 720a^3b^2 e^{6x} \\ & - 3100a^2b^3 e^{6x} + 480ab^4 e^{6x} + 1370b^5 e^{6x} - 640a^5 e^{4x} - 1970a^4b e^{4x} \\ & + 1280a^3b^2 e^{4x} + 3100a^2b^3 e^{4x} - 720ab^4 e^{4x} - 1370b^5 e^{4x} + 320a^5 e^{2x} \\ & + 805a^4b e^{2x} - 880a^3b^2 e^{2x} - 1490a^2b^3 e^{2x} + 480ab^4 e^{2x} + 685b^5 e^{2x} \\ & - 64a^5 - 137a^4b + 200a^3b^2 + 274a^2b^3 - 120ab^4 - 137b^5) / (a^6(e^{2x} - 1)^5) \end{aligned}$$

### Mupad [B] (verification not implemented)

Time = 2.39 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.82

$$\begin{aligned} \int \frac{\text{csch}^6(x)}{a + b \tanh(x)} dx &= \frac{2(a-b)(ab-b^2)}{a^4(e^{4x}-2e^{2x}+1)} - \frac{8(4a^2-3ab+b^2)}{3a^3(3e^{2x}-3e^{4x}+e^{6x}-1)} \\ &\quad - \frac{4(4a-b)}{a^2(6e^{4x}-4e^{2x}-4e^{6x}+e^{8x}+1)} \\ &\quad - \frac{5a(5e^{2x}-10e^{4x}+10e^{6x}-5e^{8x}+e^{10x}-1)}{32} \\ &\quad - \frac{2(a+b)(a-b)(ab-b^2)}{a^5(e^{2x}-1)} \\ &\quad + \frac{b \ln(a-b+ae^{2x}+be^{2x})(a+b)^2(a-b)^2}{a^6} \\ &\quad - \frac{b \ln(e^{2x}-1)(a+b)^2(a-b)^2}{a^6} \end{aligned}$$

input `int(1/(sinh(x)^6*(a + b*tanh(x))),x)`

output

$$\begin{aligned} & (2*(a - b)*(a*b - b^2))/(a^4*(\exp(4*x) - 2*\exp(2*x) + 1)) - (8*(4*a^2 - 3* \\ & a*b + b^2))/(3*a^3*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (4*(4*a - b \\ & ))/(a^2*(6*\exp(4*x) - 4*\exp(2*x) - 4*\exp(6*x) + \exp(8*x) + 1)) - 32/(5*a*( \\ & 5*\exp(2*x) - 10*\exp(4*x) + 10*\exp(6*x) - 5*\exp(8*x) + \exp(10*x) - 1)) - (2 \\ & *(a + b)*(a - b)*(a*b - b^2))/(a^5*(\exp(2*x) - 1)) + (b*\log(a - b + a*\exp( \\ & 2*x) + b*\exp(2*x))*(a + b)^2*(a - b)^2)/a^6 - (b*\log(\exp(2*x) - 1)*(a + b) \\ & ^2*(a - b)^2)/a^6 \end{aligned}$$
**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 1407, normalized size of antiderivative = 10.82

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
int(csch(x)^6/(a+b*tanh(x)),x)
```

output

```
( - 15***e**(10*x)*log(e***x - 1)*a**4*b + 30***e**(10*x)*log(e***x - 1)*a**2*b*
*3 - 15***e**(10*x)*log(e***x - 1)*b**5 - 15***e**(10*x)*log(e***x + 1)*a**4*b +
30***e**(10*x)*log(e***x + 1)*a**2*b**3 - 15***e**(10*x)*log(e***x + 1)*b**5 +
15***e**(10*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**4*b - 30***e**(10*x)*lo
g(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*b**3 + 15***e**(10*x)*log(e**(2*x)*a
+ e**(2*x)*b + a - b)*b**5 - 6***e**(10*x)*a**4*b + 6***e**(10*x)*a**3*b**2 +
6***e**(10*x)*a**2*b**3 - 6***e**(10*x)*a*b**4 + 75***e**(8*x)*log(e***x - 1)*a*
*4*b - 150***e**(8*x)*log(e***x - 1)*a**2*b**3 + 75***e**(8*x)*log(e***x - 1)*b*
*5 + 75***e**(8*x)*log(e***x + 1)*a**4*b - 150***e**(8*x)*log(e***x + 1)*a**2*b*
*3 + 75***e**(8*x)*log(e***x + 1)*b**5 - 75***e**(8*x)*log(e**(2*x)*a + e**(2*x)
)*b + a - b)*a**4*b + 150***e**(8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a*
*2*b**3 - 75***e**(8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**5 - 150***e**
(6*x)*log(e***x - 1)*a**4*b + 300***e**(6*x)*log(e***x - 1)*a**2*b**3 - 150***e**
(6*x)*log(e***x - 1)*b**5 - 150***e**(6*x)*log(e***x + 1)*a**4*b + 300***e**(6*x)
*log(e***x + 1)*a**2*b**3 - 150***e**(6*x)*log(e***x + 1)*b**5 + 150***e**(6*x)
*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**4*b - 300***e**(6*x)*log(e**(2*x)*a
+ e**(2*x)*b + a - b)*a**2*b**3 + 150***e**(6*x)*log(e**(2*x)*a + e**(2*x)*
b + a - b)*b**5 + 90***e**(6*x)*a**4*b - 120***e**(6*x)*a**3*b**2 - 30***e**(6*x)
)*a**2*b**3 + 60***e**(6*x)*a*b**4 + 150***e**(4*x)*log(e***x - 1)*a**4*b - 300
***e**(4*x)*log(e***x - 1)*a**2*b**3 + 150***e**(4*x)*log(e***x - 1)*b**5 + 1...
```

### 3.90 $\int \frac{\operatorname{csch}(x)}{i+\tanh(x)} dx$

Optimal result	733
Mathematica [A] (verified)	733
Rubi [A] (verified)	734
Maple [A] (verified)	736
Fricas [A] (verification not implemented)	736
Sympy [F]	737
Maxima [A] (verification not implemented)	737
Giac [A] (verification not implemented)	737
Mupad [B] (verification not implemented)	738
Reduce [B] (verification not implemented)	738

#### Optimal result

Integrand size = 11, antiderivative size = 33

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = i \operatorname{arctanh}(\cosh(x)) - \frac{i \operatorname{arctanh}\left(\frac{\cosh(x) + i \sinh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

output `I*arctanh(cosh(x))-1/2*I*arctanh(1/2*(cosh(x)+I*sinh(x))*2^(1/2))*2^(1/2)`

#### Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = -i \left( \sqrt{2} \operatorname{arctanh}\left(\frac{1 + i \tanh\left(\frac{x}{2}\right)}{\sqrt{2}}\right) - \log\left(\cosh\left(\frac{x}{2}\right)\right) + \log\left(\sinh\left(\frac{x}{2}\right)\right) \right)$$

input `Integrate[Csch[x]/(I + Tanh[x]), x]`

output

```
(-I)*(Sqrt[2]*ArcTanh[(1 + I*Tanh[x/2])/Sqrt[2]] - Log[Cosh[x/2]] + Log[Sinh[x/2]])
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$ , Rules used = {3042, 26, 26, 4001, 26, 3042, 26, 3589, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{csch}(x)}{\tanh(x) + i} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\sin(ix)(i - i \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int -\frac{i}{\sin(ix)(1 - \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & \int \frac{1}{\sin(ix)(1 - \tan(ix))} dx \\
 & \quad \downarrow \text{4001} \\
 & \int -\frac{i \coth(x)}{\cosh(x) - i \sinh(x)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\coth(x)}{\cosh(x) - i \sinh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & -i \int \frac{i \cos(ix)}{(\cos(ix) - \sin(ix)) \sin(ix)} dx \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\int \frac{\cos(ix)}{\sin(ix)(\cos(ix) - \sin(ix))} dx$$

↓ 3589

$$\int \left( \frac{1}{\cosh(x) - i \sinh(x)} - i \operatorname{csch}(x) \right) dx$$

↓ 2009

$$i \operatorname{arctanh}(\cosh(x)) - \frac{i \operatorname{arctanh}\left(\frac{\cosh(x) + i \sinh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

input `Int [Csch[x]/(I + Tanh[x]), x]`

output `I*ArcTanh[Cosh[x]] - (I*ArcTanh[(Cosh[x] + I*Sinh[x])/Sqrt[2]])/Sqrt[2]`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3589 `Int[(cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.))/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[ExpandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]`



rule 4001

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

**Maple [A] (verified)**

Time = 0.89 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

method	result	size
default	$-i \ln\left(\tanh\left(\frac{x}{2}\right)\right) + \sqrt{2} \arctan\left(\frac{(2 \tanh(\frac{x}{2}) - 2i)\sqrt{2}}{4}\right)$	29
risch	$-i \ln(e^x - 1) + \frac{i\sqrt{2} \ln(e^x - \frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2})}{2} - \frac{i\sqrt{2} \ln(e^x + \frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2})}{2} + i \ln(e^x + 1)$	60

input

```
int(csch(x)/(I+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-I*ln(tanh(1/2*x))+2^(1/2)*arctan(1/4*(2*tanh(1/2*x)-2*I)*2^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.30

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = -\frac{1}{2}i\sqrt{2} \log\left(-\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2} + e^x\right) + \frac{1}{2}i\sqrt{2} \log\left(\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2} + e^x\right) + i \log(e^x + 1) - i \log(e^x - 1)$$

input

```
integrate(csch(x)/(I+tanh(x)),x, algorithm="fricas")
```

output

```
-1/2*I*sqrt(2)*log(-(1/2*I - 1/2)*sqrt(2) + e^x) + 1/2*I*sqrt(2)*log((1/2*I - 1/2)*sqrt(2) + e^x) + I*log(e^x + 1) - I*log(e^x - 1)
```

**Sympy [F]**

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = \int \frac{\operatorname{csch}(x)}{\tanh(x) + i} dx$$

input `integrate(csch(x)/(I+tanh(x)),x)`

output `Integral(csch(x)/(tanh(x) + I), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = -\sqrt{2} \arctan \left( \left( \frac{1}{2}i + \frac{1}{2} \right) \sqrt{2}e^{(-x)} \right) + i \log(e^{(-x)} + 1) - i \log(e^{(-x)} - 1)$$

input `integrate(csch(x)/(I+tanh(x)),x, algorithm="maxima")`

output `-sqrt(2)*arctan((1/2*I + 1/2)*sqrt(2)*e^(-x)) + I*log(e^(-x) + 1) - I*log(e^(-x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = \sqrt{2} \arctan \left( -\left( \frac{1}{2}i - \frac{1}{2} \right) \sqrt{2}e^x \right) + i \log(e^x + 1) - i \log(|e^x - 1|)$$

input `integrate(csch(x)/(I+tanh(x)),x, algorithm="giac")`

output `sqrt(2)*arctan(-(1/2*I - 1/2)*sqrt(2)*e^x) + I*log(e^x + 1) - I*log(abs(e^x - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.43 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.85

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = \ln(-8e^x - 8) i - \ln(8 - 8e^x) i - \frac{\sqrt{2} \ln(e^x(4 - 4i) - \sqrt{2}4i) i}{2} + \frac{\sqrt{2} \ln(e^x(4 - 4i) + \sqrt{2}4i) i}{2}$$

input `int(1/(sinh(x)*(tanh(x) + 1i)),x)`output `log(- 8*exp(x) - 8)*1i - log(8 - 8*exp(x))*1i - (2^(1/2)*log(exp(x)*(4 - 4i) - 2^(1/2)*4i)*1i)/2 + (2^(1/2)*log(exp(x)*(4 - 4i) + 2^(1/2)*4i)*1i)/2`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 111, normalized size of antiderivative = 3.36

$$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx = \frac{-\sqrt{2} \log(e^x i + e^x - \sqrt{2}) i - \sqrt{2} \log(e^x i + e^x - \sqrt{2}) + \sqrt{2} \log(e^x i + e^x + \sqrt{2}) i + \sqrt{2} \log(e^x i + e^x + \sqrt{2})}{2i - 2}$$

input `int(csch(x)/(I+tanh(x)),x)`output `( - sqrt(2)*log(e**x*i + e**x - sqrt(2))*i - sqrt(2)*log(e**x*i + e**x - sqrt(2)) + sqrt(2)*log(e**x*i + e**x + sqrt(2))*i + sqrt(2)*log(e**x*i + e**x + sqrt(2)) + 2*log(e**x - 1)*i + 2*log(e**x - 1) - 2*log(e**x + 1)*i - 2*log(e**x + 1))/(2*(i - 1))`

### 3.91 $\int (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n dx$

Optimal result	739
Mathematica [F]	739
Rubi [A] (verified)	740
Maple [F]	741
Fricas [F]	742
Sympy [F]	742
Maxima [F]	742
Giac [F]	743
Mupad [F(-1)]	743
Reduce [F]	743

#### Optimal result

Integrand size = 23, antiderivative size = 153

$$\int (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n dx = \frac{b \operatorname{AppellF1}\left(1+n, 1-\frac{m}{2}, 1-\frac{m}{2}, 2+n, \frac{a+b \tanh(e+fx)}{a-b}, \frac{a+b \tanh(e+fx)}{a+b}\right) (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n}{(a^2-b^2) f(1+n)}$$

output

```
-b*AppellF1(1+n,1-1/2*m,1-1/2*m,2+n,(a+b*tanh(f*x+e))/(a-b),(a+b*tanh(f*x+e))/(a+b))*(d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^(1+n)/(a^2-b^2)/f/(1+n)/((1-(a+b*tanh(f*x+e))/(a-b))^(1/2*m))/((1-(a+b*tanh(f*x+e))/(a+b))^(1/2*m))
```

#### Mathematica [F]

$$\int (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n dx = \int (d\operatorname{sech}(e+fx))^m (a+b \tanh(e+fx))^n dx$$

input

```
Integrate[(d*Sech[e + f*x])^m*(a + b*Tanh[e + f*x])^n,x]
```

output

```
Integrate[(d*Sech[e + f*x])^m*(a + b*Tanh[e + f*x])^n, x]
```

**Rubi [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {3042, 3995, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx$$

$$\downarrow 3042$$

$$\int (d \sec(ie + ifx))^m (a - ib \tan(ie + ifx))^n dx$$

$$\downarrow 3995$$

$$\frac{b(d \operatorname{sech}(e + fx))^m \left(1 - \frac{a+b \tanh(e+fx)}{a-b}\right)^{-m/2} \left(1 - \frac{a+b \tanh(e+fx)}{a+b}\right)^{-m/2} \int (a + b \tanh(e + fx))^n \left(1 - \frac{a+b \tanh(e+fx)}{a-b}\right)^{-m/2} dx}{f(a^2 - b^2)}$$

$$\downarrow 150$$

$$\frac{b(d \operatorname{sech}(e + fx))^m \left(1 - \frac{a+b \tanh(e+fx)}{a-b}\right)^{-m/2} \left(1 - \frac{a+b \tanh(e+fx)}{a+b}\right)^{-m/2} (a + b \tanh(e + fx))^{n+1} \operatorname{AppellF1}\left(n + \frac{1}{2}, \frac{2-m}{2}, \frac{2-m}{2}, 2+n, \frac{a+b \tanh(e+fx)}{a-b}, \frac{a+b \tanh(e+fx)}{a+b}\right)}{f(n+1)(a^2 - b^2)}$$

input

```
Int[(d*Sech[e + f*x])^m*(a + b*Tanh[e + f*x])^n,x]
```

output

```
-((b*AppellF1[1 + n, (2 - m)/2, (2 - m)/2, 2 + n, (a + b*Tanh[e + f*x])/(a - b), (a + b*Tanh[e + f*x])/(a + b)]*(d*Sech[e + f*x])^m*(a + b*Tanh[e + f*x])^(1 + n))/((a^2 - b^2)*f*(1 + n)*(1 - (a + b*Tanh[e + f*x])/(a - b))^(m/2)*(1 - (a + b*Tanh[e + f*x])/(a + b))^(m/2)))
```

## Definitions of rubi rules used

rule 150

```
Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_]
  := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2,
  (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m]
  && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3995

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
  := Simp[d^(2*IntPart[m/2])*(a^2 + b^2)^(IntPart[m/2] - 1)*((d*Sec[e + f*x])^(2*FracPart[m/2])/(f*b^(2*IntPart[m/2] - 1)*(1 - (a + b*Tan[e + f*x])/(a - Rt[-b^2, 2]))^FracPart[m/2]*(1 - (a + b*Tan[e + f*x])/(a + Rt[-b^2, 2]))^FracPart[m/2])) Subst[Int[x^n*(1 - x/(a - Rt[-b^2, 2]))^(m/2 - 1)*(1 - x/(a + Rt[-b^2, 2]))^(m/2 - 1), x], x, a + b*Tan[e + f*x]], x] /; FreeQ[{a, b, d, e, f, m, n}, x] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n]
```

## Maple [F]

$$\int (d \operatorname{sech}(fx + e))^m (a + b \tanh(fx + e))^n dx$$

input

```
int((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x)
```

output

```
int((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x)
```

**Fricas [F]**

$$\int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx$$

$$= \int (d \operatorname{sech}(fx + e))^m (b \tanh(fx + e) + a)^n dx$$

input `integrate((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x, algorithm="fricas")`

output `integral((d*sech(f*x + e))^m*(b*tanh(f*x + e) + a)^n, x)`

**Sympy [F]**

$$\int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx$$

$$= \int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx$$

input `integrate((d*sech(f*x+e))**m*(a+b*tanh(f*x+e))**n,x)`

output `Integral((d*sech(e + f*x))**m*(a + b*tanh(e + f*x))**n, x)`

**Maxima [F]**

$$\int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx$$

$$= \int (d \operatorname{sech}(fx + e))^m (b \tanh(fx + e) + a)^n dx$$

input `integrate((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x, algorithm="maxima")`

output `integrate((d*sech(f*x + e))^m*(b*tanh(f*x + e) + a)^n, x)`

**Giac [F]**

$$\begin{aligned} & \int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx \\ &= \int (d \operatorname{sech}(fx + e))^m (b \tanh(fx + e) + a)^n dx \end{aligned}$$

input `integrate((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x, algorithm="giac")`

output `integrate((d*sech(f*x + e))^m*(b*tanh(f*x + e) + a)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\begin{aligned} & \int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx \\ &= \int \left( \frac{d}{\cosh(e + fx)} \right)^m (a + b \tanh(e + fx))^n dx \end{aligned}$$

input `int((d/cosh(e + f*x))^m*(a + b*tanh(e + f*x))^n,x)`

output `int((d/cosh(e + f*x))^m*(a + b*tanh(e + f*x))^n, x)`

**Reduce [F]**

$$\begin{aligned} & \int (d \operatorname{sech}(e + fx))^m (a + b \tanh(e + fx))^n dx \\ &= d^m \left( \int \operatorname{sech}(fx + e)^m (\tanh(fx + e) b + a)^n dx \right) \end{aligned}$$

input `int((d*sech(f*x+e))^m*(a+b*tanh(f*x+e))^n,x)`

output `d**m*int(sech(e + f*x)**m*(tanh(e + f*x)*b + a)**n,x)`



### 3.92 $\int \frac{\cosh^4(x)}{1+\tanh(x)} dx$

Optimal result	744
Mathematica [A] (verified)	744
Rubi [A] (verified)	745
Maple [A] (verified)	746
Fricas [B] (verification not implemented)	747
Sympy [F]	747
Maxima [A] (verification not implemented)	748
Giac [A] (verification not implemented)	748
Mupad [B] (verification not implemented)	748
Reduce [B] (verification not implemented)	749

#### Optimal result

Integrand size = 11, antiderivative size = 60

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \frac{5x}{16} + \frac{1}{32(1 - \tanh(x))^2} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} - \frac{3}{32(1 + \tanh(x))^2} - \frac{3}{16(1 + \tanh(x))}$$

output

```
5/16*x+1/32/(1-tanh(x))^2+1/(8-8*tanh(x))-1/24/(1+tanh(x))^3-3/32/(1+tanh(x))^2-3/(16+16*tanh(x))
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.88

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \frac{\operatorname{sech}(x)(-80 \cosh(x) + 15 \cosh(3x) + \cosh(5x) + 40 \sinh(x) + 120 \operatorname{arctanh}(\tanh(x))(\cosh(x) + \sinh(x)))}{384(1 + \tanh(x))}$$

input

```
Integrate[Cosh[x]^4/(1 + Tanh[x]), x]
```

output

```
(Sech[x]*(-80*Cosh[x] + 15*Cosh[3*x] + Cosh[5*x] + 40*Sinh[x] + 120*ArcTan
h[Tanh[x]]*(Cosh[x] + Sinh[x]) + 45*Sinh[3*x] + 5*Sinh[5*x]))/(384*(1 + Ta
nh[x]))
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3042, 3968, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cosh^4(x)}{\tanh(x) + 1} dx$$

$$\downarrow 3042$$

$$\int \frac{1}{(1 - i \tan(ix)) \sec(ix)^4} dx$$

$$\downarrow 3968$$

$$\int \frac{1}{(1 - \tanh(x))^3 (\tanh(x) + 1)^4} d \tanh(x)$$

$$\downarrow 54$$

$$\int \left( -\frac{5}{16 (\tanh^2(x) - 1)} + \frac{1}{8 (\tanh(x) - 1)^2} + \frac{3}{16 (\tanh(x) + 1)^2} - \frac{1}{16 (\tanh(x) - 1)^3} + \frac{3}{16 (\tanh(x) + 1)^3} + \frac{1}{8 (\tanh(x) - 1)^4} - \frac{3}{8 (\tanh(x) + 1)^4} \right) dx$$

$$\downarrow 2009$$

$$\frac{5}{16} \operatorname{arctanh}(\tanh(x)) + \frac{1}{8(1 - \tanh(x))} - \frac{3}{16(\tanh(x) + 1)} + \frac{1}{32(1 - \tanh(x))^2} - \frac{3}{32(\tanh(x) + 1)^2} - \frac{1}{24(\tanh(x) + 1)^3}$$

input

```
Int[Cosh[x]^4/(1 + Tanh[x]),x]
```

output  $(5 \operatorname{ArcTanh}[\operatorname{Tanh}[x]])/16 + 1/(32(1 - \operatorname{Tanh}[x])^2) + 1/(8(1 - \operatorname{Tanh}[x])) - 1/(24(1 + \operatorname{Tanh}[x])^3) - 3/(32(1 + \operatorname{Tanh}[x])^2) - 3/(16(1 + \operatorname{Tanh}[x]))$

### Defintions of rubi rules used

rule 54  $\operatorname{Int}[(a + (b \cdot x)^m) \cdot (c + (d \cdot x)^n), x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x \&\& \operatorname{ILtQ}[m, 0] \&\& \operatorname{IntegerQ}[n] \&\& \operatorname{!(IGtQ}[n, 0] \&\& \operatorname{LtQ}[m + n + 2, 0])$

rule 2009  $\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$

rule 3042  $\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3968  $\operatorname{Int}[\sec[(e + (f \cdot x)^m) \cdot (a + (b \cdot \tan[(e + (f \cdot x)^m])^n)], x\_Symbol] \rightarrow \operatorname{Simp}[1/(a^{m-2} \cdot b \cdot f) \operatorname{Subst}[\operatorname{Int}[(a - x)^{m/2-1} \cdot (a + x)^{n+m/2-1}, x], x, b \cdot \tan[e + f \cdot x]], x] /; \operatorname{FreeQ}\{a, b, e, f, n\}, x \&\& \operatorname{EqQ}[a^2 + b^2, 0] \&\& \operatorname{IntegerQ}[m/2]$

### Maple [A] (verified)

Time = 3.80 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.58

method	result
risch	$\frac{5x}{16} + \frac{e^{4x}}{128} + \frac{5e^{2x}}{64} - \frac{5e^{-2x}}{32} - \frac{5e^{-4x}}{128} - \frac{e^{-6x}}{192}$
parallelrisch	$\frac{(-12 \cosh(x) - 12 \sinh(x)) \ln(1 - \tanh(x)) + (12 \cosh(x) + 12 \sinh(x)) \ln(1 + \tanh(x)) + 15 \cosh(3x) + \cosh(5x) + 45 \sinh(3x) + 5 \sinh(5x)}{384 \cosh(x) + 384 \sinh(x)}$
default	$\frac{1}{8(\tanh(\frac{x}{2}) - 1)^4} + \frac{1}{4(\tanh(\frac{x}{2}) - 1)^3} + \frac{1}{2(\tanh(\frac{x}{2}) - 1)^2} + \frac{3}{8(\tanh(\frac{x}{2}) - 1)} - \frac{5 \ln(\tanh(\frac{x}{2}) - 1)}{16} - \frac{1}{3(1 + \tanh(\frac{x}{2}))^6} + \dots$

input  $\operatorname{int}(\cosh(x)^4/(1 + \tanh(x)), x, \operatorname{method} = \_RETURNVERBOSE)$

output  $5/16*x+1/128*\exp(4*x)+5/64*\exp(2*x)-5/32*\exp(-2*x)-5/128*\exp(-4*x)-1/192*\exp(-6*x)$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs.  $2(44) = 88$ .

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.58

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx$$

$$= \frac{\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + 5 \sinh(x)^5 + 5(10 \cosh(x)^2 + 9) \sinh(x)^3 + 15 \cosh(x)^3 + 5(2 \cosh(x)^2 + 3) \sinh(x)}{384(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="fricas")`

output `1/384*(cosh(x)^5 + 5*cosh(x)*sinh(x)^4 + 5*sinh(x)^5 + 5*(10*cosh(x)^2 + 9)*sinh(x)^3 + 15*cosh(x)^3 + 5*(2*cosh(x)^3 + 9*cosh(x))*sinh(x)^2 + 60*(2*x - 1)*cosh(x) + 5*(5*cosh(x)^4 + 27*cosh(x)^2 + 24*x + 12)*sinh(x))/(cosh(x) + sinh(x))`

### Sympy [F]

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \int \frac{\cosh^4(x)}{\tanh(x) + 1} dx$$

input `integrate(cosh(x)**4/(1+tanh(x)),x)`

output `Integral(cosh(x)**4/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.60

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \frac{1}{128} (10 e^{(-2x)} + 1) e^{(4x)} + \frac{5}{16} x - \frac{5}{32} e^{(-2x)} - \frac{5}{128} e^{(-4x)} - \frac{1}{192} e^{(-6x)}$$

input `integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="maxima")`output `1/128*(10*e^(-2*x) + 1)*e^(4*x) + 5/16*x - 5/32*e^(-2*x) - 5/128*e^(-4*x) - 1/192*e^(-6*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.70

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = -\frac{1}{384} (110 e^{(6x)} + 60 e^{(4x)} + 15 e^{(2x)} + 2) e^{(-6x)} + \frac{5}{16} x + \frac{1}{128} e^{(4x)} + \frac{5}{64} e^{(2x)}$$

input `integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="giac")`output `-1/384*(110*e^(6*x) + 60*e^(4*x) + 15*e^(2*x) + 2)*e^(-6*x) + 5/16*x + 1/128*e^(4*x) + 5/64*e^(2*x)`**Mupad [B] (verification not implemented)**

Time = 2.33 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.57

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \frac{5x}{16} - \frac{5e^{-2x}}{32} + \frac{5e^{2x}}{64} - \frac{5e^{-4x}}{128} + \frac{e^{4x}}{128} - \frac{e^{-6x}}{192}$$

input `int(cosh(x)^4/(tanh(x) + 1),x)`

output  $(5x)/16 - (5\exp(-2x))/32 + (5\exp(2x))/64 - (5\exp(-4x))/128 + \exp(4x)/128 - \exp(-6x)/192$

### Reduce [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx = \frac{3e^{10x} + 30e^{8x} + 120e^{6x}x - 60e^{4x} - 15e^{2x} - 2}{384e^{6x}}$$

input `int(cosh(x)^4/(1+tanh(x)),x)`

output  $(3e^{10x} + 30e^{8x} + 120e^{6x}x - 60e^{4x} - 15e^{2x} - 2)/(384e^{6x})$

### 3.93 $\int \frac{\cosh^3(x)}{1+\tanh(x)} dx$

Optimal result	750
Mathematica [A] (verified)	750
Rubi [C] (verified)	751
Maple [A] (verified)	752
Fricas [B] (verification not implemented)	753
Sympy [B] (verification not implemented)	753
Maxima [A] (verification not implemented)	754
Giac [A] (verification not implemented)	754
Mupad [B] (verification not implemented)	755
Reduce [B] (verification not implemented)	755

#### Optimal result

Integrand size = 11, antiderivative size = 29

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = \frac{4 \sinh(x)}{5} + \frac{4 \sinh^3(x)}{15} - \frac{\cosh^3(x)}{5(1 + \tanh(x))}$$

output `4/5*sinh(x)+4/15*sinh(x)^3-cosh(x)^3/(5+5*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.24

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = \frac{\operatorname{sech}(x)(-45 + 20 \cosh(2x) + \cosh(4x) + 40 \sinh(2x) + 4 \sinh(4x))}{120(1 + \tanh(x))}$$

input `Integrate[Cosh[x]^3/(1 + Tanh[x]),x]`

output `(Sech[x]*(-45 + 20*Cosh[2*x] + Cosh[4*x] + 40*Sinh[2*x] + 4*Sinh[4*x]))/(120*(1 + Tanh[x]))`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.31, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {3042, 3983, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh^3(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix)) \sec(ix)^3} dx \\
 & \quad \downarrow \text{3983} \\
 & \frac{4}{5} \int \cosh^3(x) dx - \frac{\cosh^3(x)}{5(\tanh(x) + 1)} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\cosh^3(x)}{5(\tanh(x) + 1)} + \frac{4}{5} \int \sin\left(ix + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{3113} \\
 & -\frac{\cosh^3(x)}{5(\tanh(x) + 1)} + \frac{4}{5} i \int (\sinh^2(x) + 1) d(-i \sinh(x)) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\cosh^3(x)}{5(\tanh(x) + 1)} + \frac{4}{5} i \left( -\frac{1}{3} i \sinh^3(x) - i \sinh(x) \right)
 \end{aligned}$$

input `Int[Cosh[x]^3/(1 + Tanh[x]),x]`

output `((4*I)/5)*((-I)*Sinh[x] - (I/3)*Sinh[x]^3) - Cosh[x]^3/(5*(1 + Tanh[x]))`



## Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3983 `Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[a*(d*Sec[e + f*x])^m*((a + b*Tan[e + f*x])^n/(b*f*(m + 2*n))), x] + Simp[Simplify[m + n]/(a*(m + 2*n)) Int[(d*Sec[e + f*x])^m*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && NeQ[m + 2*n, 0] && IntegersQ[2*m, 2*n]`

## Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

method	result
risch	$\frac{e^{3x}}{48} + \frac{e^x}{4} - \frac{3e^{-x}}{8} - \frac{e^{-3x}}{12} - \frac{e^{-5x}}{80}$
parallelrisch	$\frac{-45 + \cosh(4x) + 20 \cosh(2x) + 4 \sinh(4x) + 40 \sinh(2x) - 24 \sinh(x) - 24 \cosh(x)}{120 \sinh(x) + 120 \cosh(x)}$
default	$-\frac{1}{6(\tanh(\frac{x}{2})-1)^3} - \frac{1}{4(\tanh(\frac{x}{2})-1)^2} - \frac{5}{8(\tanh(\frac{x}{2})-1)} - \frac{2}{5(1+\tanh(\frac{x}{2}))^5} + \frac{1}{(1+\tanh(\frac{x}{2}))^4} - \frac{5}{3(1+\tanh(\frac{x}{2}))^3} + \dots$

input `int(cosh(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `1/48*exp(3*x)+1/4*exp(x)-3/8*exp(-x)-1/12*exp(-3*x)-1/80*exp(-5*x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 60 vs.  $2(23) = 46$ .

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.07

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{\cosh(x)^4 + 16 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 10) \sinh(x)^2 + 20 \cosh(x)^2 + 16(\cosh(x) + \sinh(x))}{120(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output `1/120*(cosh(x)^4 + 16*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 10)*sinh(x)^2 + 20*cosh(x)^2 + 16*(cosh(x)^3 + 5*cosh(x))*sinh(x) - 45)/(cosh(x) + sinh(x))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 134 vs.  $2(26) = 52$ .

Time = 0.31 (sec) , antiderivative size = 134, normalized size of antiderivative = 4.62

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = -\frac{8 \sinh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{2 \sinh^3(x)}{15 \tanh(x) + 15}$$

$$- \frac{6 \sinh^2(x) \cosh(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{6 \sinh^2(x) \cosh(x)}{15 \tanh(x) + 15}$$

$$+ \frac{6 \sinh(x) \cosh^2(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{9 \sinh(x) \cosh^2(x)}{15 \tanh(x) + 15}$$

$$+ \frac{3 \cosh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{3 \cosh^3(x)}{15 \tanh(x) + 15}$$

input `integrate(cosh(x)**3/(1+tanh(x)),x)`

output

```
-8*sinh(x)**3*tanh(x)/(15*tanh(x) + 15) - 2*sinh(x)**3/(15*tanh(x) + 15) -
6*sinh(x)**2*cosh(x)*tanh(x)/(15*tanh(x) + 15) + 6*sinh(x)**2*cosh(x)/(15
*tanh(x) + 15) + 6*sinh(x)*cosh(x)**2*tanh(x)/(15*tanh(x) + 15) + 9*sinh(x
)*cosh(x)**2/(15*tanh(x) + 15) + 3*cosh(x)**3*tanh(x)/(15*tanh(x) + 15) -
3*cosh(x)**3/(15*tanh(x) + 15)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = \frac{1}{48} (12 e^{(-2x)} + 1) e^{(3x)} - \frac{3}{8} e^{(-x)} - \frac{1}{12} e^{(-3x)} - \frac{1}{80} e^{(-5x)}$$

input

```
integrate(cosh(x)^3/(1+tanh(x)),x, algorithm="maxima")
```

output

```
1/48*(12*e^(-2*x) + 1)*e^(3*x) - 3/8*e^(-x) - 1/12*e^(-3*x) - 1/80*e^(-5*x)
)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = -\frac{1}{240} (90 e^{(4x)} + 20 e^{(2x)} + 3) e^{(-5x)} + \frac{1}{48} e^{(3x)} + \frac{1}{4} e^x$$

input

```
integrate(cosh(x)^3/(1+tanh(x)),x, algorithm="giac")
```

output

```
-1/240*(90*e^(4*x) + 20*e^(2*x) + 3)*e^(-5*x) + 1/48*e^(3*x) + 1/4*e^x
```

**Mupad [B] (verification not implemented)**

Time = 2.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = \frac{e^{3x}}{48} - \frac{e^{-3x}}{12} - \frac{3e^{-x}}{8} - \frac{e^{-5x}}{80} + \frac{e^x}{4}$$

input `int(cosh(x)^3/(tanh(x) + 1),x)`output `exp(3*x)/48 - exp(-3*x)/12 - (3*exp(-x))/8 - exp(-5*x)/80 + exp(x)/4`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.34

$$\int \frac{\cosh^3(x)}{1 + \tanh(x)} dx = \frac{5e^{8x} + 60e^{6x} - 90e^{4x} - 20e^{2x} - 3}{240e^{5x}}$$

input `int(cosh(x)^3/(1+tanh(x)),x)`output `(5*e**(8*x) + 60*e**(6*x) - 90*e**(4*x) - 20*e**(2*x) - 3)/(240*e**(5*x))`

### 3.94 $\int \frac{\cosh^2(x)}{1+\tanh(x)} dx$

Optimal result	756
Mathematica [A] (verified)	756
Rubi [A] (verified)	757
Maple [A] (verified)	758
Fricas [A] (verification not implemented)	759
Sympy [F]	759
Maxima [A] (verification not implemented)	759
Giac [A] (verification not implemented)	760
Mupad [B] (verification not implemented)	760
Reduce [B] (verification not implemented)	760

#### Optimal result

Integrand size = 11, antiderivative size = 38

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \frac{3x}{8} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} - \frac{1}{4(1 + \tanh(x))}$$

output `3/8*x+1/(8-8*tanh(x))-1/8/(1+tanh(x))^2-1/(4+4*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \frac{1}{16} (6 \operatorname{arctanh}(\tanh(x)) + \cosh(x)(\cosh(x) - \sinh(x))(-5 + \cosh(2x) + 3 \sinh(2x)))$$

input `Integrate[Cosh[x]^2/(1 + Tanh[x]),x]`

output `(6*ArcTanh[Tanh[x]] + Cosh[x]*(Cosh[x] - Sinh[x])*(-5 + Cosh[2*x] + 3*Sinh[2*x]))/16`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3042, 3968, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh^2(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix)) \sec(ix)^2} dx \\
 & \quad \downarrow \text{3968} \\
 & \int \frac{1}{(1 - \tanh(x))^2 (\tanh(x) + 1)^3} d \tanh(x) \\
 & \quad \downarrow \text{54} \\
 & \int \left( -\frac{3}{8(\tanh^2(x) - 1)} + \frac{1}{8(\tanh(x) - 1)^2} + \frac{1}{4(\tanh(x) + 1)^2} + \frac{1}{4(\tanh(x) + 1)^3} \right) d \tanh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{3}{8} \operatorname{arctanh}(\tanh(x)) + \frac{1}{8(1 - \tanh(x))} - \frac{1}{4(\tanh(x) + 1)} - \frac{1}{8(\tanh(x) + 1)^2}
 \end{aligned}$$

input `Int [Cosh[x]^2/(1 + Tanh[x]), x]`

output `(3*ArcTanh[Tanh[x]])/8 + 1/(8*(1 - Tanh[x])) - 1/(8*(1 + Tanh[x])^2) - 1/(4*(1 + Tanh[x]))`

## Definitions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3968 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/(a^(m - 2)*b*f) Subst[Int[(a - x)^(m/2 - 1)*(a + x)^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && EqQ[a^2 + b^2, 0] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.61

method	result
risch	$\frac{3x}{8} + \frac{e^{2x}}{16} - \frac{3e^{-2x}}{16} - \frac{e^{-4x}}{32}$
parallelrisc	$\frac{\cosh(3x)+3\sinh(3x)+(12x-1)\cosh(x)+(12x+11)\sinh(x)}{32\sinh(x)+32\cosh(x)}$
default	$\frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{4\tanh(\frac{x}{2})-4} - \frac{3\ln(\tanh(\frac{x}{2})-1)}{8} - \frac{1}{2(1+\tanh(\frac{x}{2}))^4} + \frac{1}{(1+\tanh(\frac{x}{2}))^3} - \frac{3}{2(1+\tanh(\frac{x}{2}))^2} + \frac{1}{1+\tanh(\frac{x}{2})}$

input `int(cosh(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `3/8*x+1/16*exp(2*x)-3/16*exp(-2*x)-1/32*exp(-4*x)`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.37

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + 3 \sinh(x)^3 + 6(2x - 1) \cosh(x) + 3(3 \cosh(x)^2 + 4x + 2) \sinh(x)}{32(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="fricas")`

output `1/32*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + 3*sinh(x)^3 + 6*(2*x - 1)*cosh(x) + 3*(3*cosh(x)^2 + 4*x + 2)*sinh(x))/(cosh(x) + sinh(x))`

**Sympy [F]**

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \int \frac{\cosh^2(x)}{\tanh(x) + 1} dx$$

input `integrate(cosh(x)**2/(1+tanh(x)),x)`

output `Integral(cosh(x)**2/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \frac{3}{8}x + \frac{1}{16}e^{(2x)} - \frac{3}{16}e^{(-2x)} - \frac{1}{32}e^{(-4x)}$$

input `integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="maxima")`

output `3/8*x + 1/16*e^(2*x) - 3/16*e^(-2*x) - 1/32*e^(-4*x)`



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = -\frac{1}{32} (9e^{(4x)} + 6e^{(2x)} + 1)e^{(-4x)} + \frac{3}{8}x + \frac{1}{16}e^{(2x)}$$

input `integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="giac")`

output `-1/32*(9*e^(4*x) + 6*e^(2*x) + 1)*e^(-4*x) + 3/8*x + 1/16*e^(2*x)`

**Mupad [B] (verification not implemented)**

Time = 2.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \frac{3x}{8} - \frac{3e^{-2x}}{16} + \frac{e^{2x}}{16} - \frac{e^{-4x}}{32}$$

input `int(cosh(x)^2/(tanh(x) + 1),x)`

output `(3*x)/8 - (3*exp(-2*x))/16 + exp(2*x)/16 - exp(-4*x)/32`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx = \frac{2e^{6x} + 12e^{4x}x - 6e^{2x} - 1}{32e^{4x}}$$

input `int(cosh(x)^2/(1+tanh(x)),x)`

output `(2*e**(6*x) + 12*e**(4*x)*x - 6*e**(2*x) - 1)/(32*e**(4*x))`

### 3.95 $\int \frac{\cosh(x)}{1+\tanh(x)} dx$

Optimal result . . . . .	761
Mathematica [A] (verified) . . . . .	761
Rubi [A] (verified) . . . . .	762
Maple [A] (verified) . . . . .	763
Fricas [A] (verification not implemented) . . . . .	764
Sympy [B] (verification not implemented) . . . . .	764
Maxima [A] (verification not implemented) . . . . .	764
Giac [A] (verification not implemented) . . . . .	765
Mupad [B] (verification not implemented) . . . . .	765
Reduce [B] (verification not implemented) . . . . .	765

#### Optimal result

Integrand size = 9, antiderivative size = 19

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{2 \sinh(x)}{3} - \frac{\cosh(x)}{3(1 + \tanh(x))}$$

output `2/3*sinh(x)-cosh(x)/(3+3*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{1}{12}(-3 \cosh(x) - \cosh(3x) + 9 \sinh(x) + \sinh(3x))$$

input `Integrate[Cosh[x]/(1 + Tanh[x]),x]`

output `(-3*Cosh[x] - Cosh[3*x] + 9*Sinh[x] + Sinh[3*x])/12`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {3042, 3983, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix)) \sec(ix)} dx \\
 & \quad \downarrow \text{3983} \\
 & \frac{2 \int \cosh(x) dx}{3} - \frac{\cosh(x)}{3(\tanh(x) + 1)} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\cosh(x)}{3(\tanh(x) + 1)} + \frac{2}{3} \int \sin\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3117} \\
 & \frac{2 \sinh(x)}{3} - \frac{\cosh(x)}{3(\tanh(x) + 1)}
 \end{aligned}$$

input `Int [Cosh[x]/(1 + Tanh[x]), x]`

output `(2*Sinh[x])/3 - Cosh[x]/(3*(1 + Tanh[x]))`

## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3983 `Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(d*Sec[e + f*x])^m*((a + b*Tan[e + f*x])^n/(b*f*(m + 2*n))), x] + Simp[Simplify[m + n]/(a*(m + 2*n)) Int[(d*Sec[e + f*x])^m*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && NeQ[m + 2*n, 0] && IntegersQ[2*m, 2*n]`

## Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
risch	$\frac{e^x}{4} - \frac{e^{-x}}{2} - \frac{e^{-3x}}{12}$	18
parallelrisc	$-\frac{\cosh(3x)}{12} + \frac{1}{3} - \frac{\cosh(x)}{4} + \frac{3\sinh(x)}{4} + \frac{\sinh(3x)}{12}$	23
default	$-\frac{1}{2(\tanh(\frac{x}{2})-1)} - \frac{2}{3(1+\tanh(\frac{x}{2}))^3} + \frac{1}{(1+\tanh(\frac{x}{2}))^2} - \frac{3}{2(1+\tanh(\frac{x}{2}))}$	40

input `int(cosh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `1/4*exp(x)-1/2*exp(-x)-1/12*exp(-3*x)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{\cosh(x)^2 + 4 \cosh(x) \sinh(x) + \sinh(x)^2 - 3}{6(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)/(1+tanh(x)),x, algorithm="fricas")`

output `1/6*(cosh(x)^2 + 4*cosh(x)*sinh(x) + sinh(x)^2 - 3)/(cosh(x) + sinh(x))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. 2(15) = 30.

Time = 0.19 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.53

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{2 \sinh(x) \tanh(x)}{3 \tanh(x) + 3} + \frac{\sinh(x)}{3 \tanh(x) + 3} + \frac{\cosh(x) \tanh(x)}{3 \tanh(x) + 3} - \frac{\cosh(x)}{3 \tanh(x) + 3}$$

input `integrate(cosh(x)/(1+tanh(x)),x)`

output `2*sinh(x)*tanh(x)/(3*tanh(x) + 3) + sinh(x)/(3*tanh(x) + 3) + cosh(x)*tanh(x)/(3*tanh(x) + 3) - cosh(x)/(3*tanh(x) + 3)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = -\frac{1}{2} e^{(-x)} - \frac{1}{12} e^{(-3x)} + \frac{1}{4} e^x$$

input `integrate(cosh(x)/(1+tanh(x)),x, algorithm="maxima")`

output  $-1/2*e^{-x} - 1/12*e^{-3*x} + 1/4*e^x$

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = -\frac{1}{12} (6e^{2x} + 1)e^{-3x} + \frac{1}{4} e^x$$

input `integrate(cosh(x)/(1+tanh(x)),x, algorithm="giac")`

output  $-1/12*(6*e^{2*x} + 1)*e^{-3*x} + 1/4*e^x$

### Mupad [B] (verification not implemented)

Time = 2.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{e^x}{4} - \frac{e^{-3x}}{12} - \frac{e^{-x}}{2}$$

input `int(cosh(x)/(tanh(x) + 1),x)`

output  $\exp(x)/4 - \exp(-3*x)/12 - \exp(-x)/2$

### Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{\cosh(x)}{1 + \tanh(x)} dx = \frac{3e^{4x} - 6e^{2x} - 1}{12e^{3x}}$$

input `int(cosh(x)/(1+tanh(x)),x)`

output  $(3*e^{4*x} - 6*e^{2*x} - 1)/(12*e^{3*x})$

### 3.96 $\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx$

Optimal result	766
Mathematica [A] (verified)	766
Rubi [A] (verified)	767
Maple [A] (verified)	768
Fricas [A] (verification not implemented)	768
Sympy [A] (verification not implemented)	768
Maxima [A] (verification not implemented)	769
Giac [A] (verification not implemented)	769
Mupad [B] (verification not implemented)	769
Reduce [B] (verification not implemented)	770

#### Optimal result

Integrand size = 9, antiderivative size = 10

$$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx = -\frac{\operatorname{sech}(x)}{1+\tanh(x)}$$

output `-sech(x)/(1+tanh(x))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx = -\cosh(x) + \sinh(x)$$

input `Integrate[Sech[x]/(1 + Tanh[x]), x]`

output `-Cosh[x] + Sinh[x]`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {3042, 3969}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}(x)}{\tanh(x) + 1} dx$$

↓ 3042

$$\int \frac{\sec(ix)}{1 - i \tan(ix)} dx$$

↓ 3969

$$-\frac{\operatorname{sech}(x)}{\tanh(x) + 1}$$

input `Int [Sech [x] / (1 + Tanh [x]), x]`

output `-(Sech [x] / (1 + Tanh [x]))`

**Defintions of rubi rules used**

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 3969 `Int [((d_)*sec [(e_) + (f_)*(x_)] )^(m_)*((a_) + (b_)*tan [(e_) + (f_)*(x_)] )^(n_), x_Symbol] :> Simp [b*(d*Sec [e + f*x])^m*((a + b*Tan [e + f*x])^n / (a*f*m)), x] /; FreeQ [ {a, b, d, e, f, m, n}, x] && EqQ [a^2 + b^2, 0] && EqQ [Simplify [m + n], 0]`



**Maple [A] (verified)**

Time = 0.72 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

method	result	size
risch	$-e^{-x}$	7
gosper	$-\frac{\operatorname{sech}(x)}{1+\tanh(x)}$	11
default	$-\frac{2}{1+\tanh(\frac{x}{2})}$	11
orering	$-\frac{\operatorname{sech}(x)}{1+\tanh(x)}$	11

input `int(sech(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)`output `-exp(-x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx = -\frac{1}{\cosh(x) + \sinh(x)}$$

input `integrate(sech(x)/(1+tanh(x)),x, algorithm="fricas")`output `-1/(cosh(x) + sinh(x))`**Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx = -\frac{\operatorname{sech}(x)}{\tanh(x) + 1}$$

input `integrate(sech(x)/(1+tanh(x)),x)`

output `-sech(x)/(tanh(x) + 1)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{sech}(x)}{1 + \tanh(x)} dx = -e^{(-x)}$$

input `integrate(sech(x)/(1+tanh(x)),x, algorithm="maxima")`

output `-e^(-x)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{sech}(x)}{1 + \tanh(x)} dx = -e^{(-x)}$$

input `integrate(sech(x)/(1+tanh(x)),x, algorithm="giac")`

output `-e^(-x)`

### Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{sech}(x)}{1 + \tanh(x)} dx = -e^{-x}$$

input `int(1/(cosh(x)*(tanh(x) + 1)),x)`

output `-exp(-x)`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{sech}(x)}{1 + \tanh(x)} dx = -\frac{1}{e^x}$$

input `int(sech(x)/(1+tanh(x)),x)`

output `( - 1)/e**x`

### 3.97 $\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx$

Optimal result	771
Mathematica [A] (verified)	771
Rubi [A] (verified)	772
Maple [A] (verified)	773
Fricas [B] (verification not implemented)	773
Sympy [F]	774
Maxima [A] (verification not implemented)	774
Giac [B] (verification not implemented)	774
Mupad [B] (verification not implemented)	775
Reduce [B] (verification not implemented)	775

#### Optimal result

Integrand size = 11, antiderivative size = 5

$$\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx = \log(1+\tanh(x))$$

output `ln(1+tanh(x))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.40

$$\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx = x - \log(\cosh(x))$$

input `Integrate[Sech[x]^2/(1 + Tanh[x]), x]`

output `x - Log[Cosh[x]]`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {3042, 3968, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{sech}^2(x)}{\tanh(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sec(ix)^2}{1 - i \tan(ix)} dx \\ & \quad \downarrow \text{3968} \\ & \int \frac{1}{\tanh(x) + 1} d \tanh(x) \\ & \quad \downarrow \text{16} \\ & \log(\tanh(x) + 1) \end{aligned}$$

input `Int [Sech [x]^2/(1 + Tanh [x]), x]`

output `Log [1 + Tanh [x]]`

**Defintions of rubi rules used**

rule 16 `Int [(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp [c*(Log [RemoveContent [a + b*x, x]]/b), x] /; FreeQ [{a, b, c}, x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 3968

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_
), x_Symbol] :> Simp[1/(a^(m - 2)*b*f) Subst[Int[(a - x)^(m/2 - 1)*(a + x
)^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] &&
EqQ[a^2 + b^2, 0] && IntegerQ[m/2]
```

**Maple [A] (verified)**

Time = 1.90 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\ln(1 + \tanh(x))$	6
default	$\ln(1 + \tanh(x))$	6
risch	$2x - \ln(e^{2x} + 1)$	14

input

```
int(sech(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
ln(1+tanh(x))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 20 vs.  $2(5) = 10$ .

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 4.00

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = 2x - \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)$$

input

```
integrate(sech(x)^2/(1+tanh(x)),x, algorithm="fricas")
```

output

```
2*x - log(2*cosh(x)/(cosh(x) - sinh(x)))
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^2(x)}{\tanh(x) + 1} dx$$

input `integrate(sech(x)**2/(1+tanh(x)),x)`

output `Integral(sech(x)**2/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = \log(\tanh(x) + 1)$$

input `integrate(sech(x)^2/(1+tanh(x)),x, algorithm="maxima")`

output `log(tanh(x) + 1)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 13 vs.  $2(5) = 10$ .

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 2.60

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = 2x - \log(e^{(2x)} + 1)$$

input `integrate(sech(x)^2/(1+tanh(x)),x, algorithm="giac")`

output `2*x - log(e^(2*x) + 1)`

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 2.60

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = 2x - \ln(e^{2x} + 1)$$

input `int(1/(cosh(x)^2*(tanh(x) + 1)),x)`

output `2*x - log(exp(2*x) + 1)`

**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 2.80

$$\int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx = -\log(e^{2x} + 1) + 2x$$

input `int(sech(x)^2/(1+tanh(x)),x)`

output `- log(e**(2*x) + 1) + 2*x`



### 3.98 $\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx$

Optimal result	776
Mathematica [A] (verified)	776
Rubi [A] (verified)	777
Maple [B] (verified)	778
Fricas [B] (verification not implemented)	779
Sympy [F]	779
Maxima [B] (verification not implemented)	779
Giac [B] (verification not implemented)	780
Mupad [B] (verification not implemented)	780
Reduce [B] (verification not implemented)	781

#### Optimal result

Integrand size = 11, antiderivative size = 6

$$\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx = \arctan(\sinh(x)) + \operatorname{sech}(x)$$

output `arctan(sinh(x))+sech(x)`

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx = 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + \operatorname{sech}(x)$$

input `Integrate[Sech[x]^3/(1 + Tanh[x]), x]`

output `2*ArcTan[Tanh[x/2]] + Sech[x]`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3042, 3982, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^3(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^3}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3982} \\
 & \int \operatorname{sech}(x) dx + \operatorname{sech}(x) \\
 & \quad \downarrow \text{3042} \\
 & \operatorname{sech}(x) + \int \csc\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4257} \\
 & \arctan(\sinh(x)) + \operatorname{sech}(x)
 \end{aligned}$$

input `Int [Sech[x]^3/(1 + Tanh[x]), x]`

output `ArcTan[Sinh[x]] + Sech[x]`

## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3982 `Int[((d_)*sec[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Simp[d^2*((m - 2)/(a*(m + n - 1)))*Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !IntegerQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs.  $2(6) = 12$ .

Time = 4.62 (sec) , antiderivative size = 21, normalized size of antiderivative = 3.50

method	result	size
default	$\frac{2}{\tanh(\frac{x}{2})^2 + 1} + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\right)$	21
risch	$\frac{2e^x}{e^{2x} + 1} + i \ln(e^x + i) - i \ln(e^x - i)$	32

input `int(sech(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `2/(tanh(1/2*x)^2+1)+2*arctan(tanh(1/2*x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 48 vs.  $2(6) = 12$ .

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 8.00

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{2 \left( (\cosh(x))^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1 \right) \arctan(\cosh(x) + \sinh(x)) + \cosh(x) + \sinh(x)}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}$$

input `integrate(sech(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output `2*((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*arctan(cosh(x) + sinh(x)) + cosh(x) + sinh(x))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^3(x)}{\tanh(x) + 1} dx$$

input `integrate(sech(x)**3/(1+tanh(x)),x)`

output `Integral(sech(x)**3/(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 22 vs.  $2(6) = 12$ .

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 3.67

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx = \frac{2 e^{(-x)}}{e^{(-2x)} + 1} - 2 \arctan(e^{(-x)})$$

input `integrate(sech(x)^3/(1+tanh(x)),x, algorithm="maxima")`

output `2*e^(-x)/(e^(-2*x) + 1) - 2*arctan(e^(-x))`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. 2(6) = 12.

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 3.00

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx = \frac{2e^x}{e^{(2x)} + 1} + 2 \arctan(e^x)$$

input `integrate(sech(x)^3/(1+tanh(x)),x, algorithm="giac")`

output `2*e^x/(e^(2*x) + 1) + 2*arctan(e^x)`

### Mupad [B] (verification not implemented)

Time = 2.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 3.00

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx = 2 \operatorname{atan}(e^x) + \frac{2e^x}{e^{2x} + 1}$$

input `int(1/(cosh(x)^3*(tanh(x) + 1)),x)`

output `2*atan(exp(x)) + (2*exp(x))/(exp(2*x) + 1)`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 4.83

$$\int \frac{\operatorname{sech}^3(x)}{1 + \tanh(x)} dx = \frac{2e^{2x} \operatorname{atan}(e^x) + 2\operatorname{atan}(e^x) + 2e^x}{e^{2x} + 1}$$

input `int(sech(x)^3/(1+tanh(x)),x)`

output `(2*(e**(2*x))*atan(e**x) + atan(e**x) + e**x)/(e**(2*x) + 1)`

### 3.99 $\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx$

Optimal result	782
Mathematica [A] (verified)	782
Rubi [A] (verified)	783
Maple [A] (verified)	784
Fricas [B] (verification not implemented)	784
Sympy [F]	785
Maxima [B] (verification not implemented)	785
Giac [A] (verification not implemented)	786
Mupad [B] (verification not implemented)	786
Reduce [B] (verification not implemented)	786

#### Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx = -\frac{1}{2}(1-\tanh(x))^2$$

output `-1/2*(1-tanh(x))^2`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx = -\frac{1}{2}(-2+\tanh(x))\tanh(x)$$

input `Integrate[Sech[x]^4/(1+Tanh[x]),x]`

output `-1/2*((-2+Tanh[x])*Tanh[x])`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {3042, 3968, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^4(x)}{\tanh(x) + 1} dx$$

↓ 3042

$$\int \frac{\sec(ix)^4}{1 - i \tan(ix)} dx$$

↓ 3968

$$\int (1 - \tanh(x)) d \tanh(x)$$

↓ 17

$$-\frac{1}{2}(1 - \tanh(x))^2$$

input `Int [Sech [x]^4/(1 + Tanh [x]), x]`

output `-1/2*(1 - Tanh [x])^2`

**Defintions of rubi rules used**

rule 17 `Int [(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp [c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 3042 `Int [u_, x_Symbol] := Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinearQ [u, x]`



rule 3968

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_
), x_Symbol] :> Simp[1/(a^(m - 2)*b*f) Subst[Int[(a - x)^(m/2 - 1)*(a + x
)^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] &&
EqQ[a^2 + b^2, 0] && IntegerQ[m/2]
```

**Maple [A] (verified)**

Time = 1.99 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\tanh(x) - \frac{\tanh(x)^2}{2}$	10
default	$\tanh(x) - \frac{\tanh(x)^2}{2}$	10
risch	$-\frac{2}{(e^{2x}+1)^2}$	11
parallelrisch	$\frac{7+18 \sinh(2x)-11 \cosh(2x)}{18+18 \cosh(2x)}$	26

input

```
int(sech(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
tanh(x)-1/2*tanh(x)^2
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(8) = 16$ .

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 4.42

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx =$$

$$\frac{2}{\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)$$

input

```
integrate(sech(x)^4/(1+tanh(x)),x, algorithm="fricas")
```

output 
$$\frac{-2/(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 + 1)*\sinh(x)^2 + 2*\cosh(x)^2 + 4*(\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)}{1}$$

### Sympy [F]

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^4(x)}{\tanh(x) + 1} dx$$

input `integrate(sech(x)**4/(1+tanh(x)), x)`

output `Integral(sech(x)**4/(tanh(x) + 1), x)`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs.  $2(8) = 16$ .

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 3.08

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx = \frac{4e^{(-2x)}}{2e^{(-2x)} + e^{(-4x)} + 1} + \frac{2}{2e^{(-2x)} + e^{(-4x)} + 1}$$

input `integrate(sech(x)^4/(1+tanh(x)), x, algorithm="maxima")`

output `4*e^(-2*x)/(2*e^(-2*x) + e^(-4*x) + 1) + 2/(2*e^(-2*x) + e^(-4*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx = -\frac{2}{(e^{2x} + 1)^2}$$

input `integrate(sech(x)^4/(1+tanh(x)),x, algorithm="giac")`output `-2/(e^(2*x) + 1)^2`**Mupad [B] (verification not implemented)**

Time = 2.47 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.33

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx = -\frac{2}{2e^{2x} + e^{4x} + 1}$$

input `int(1/(cosh(x)^4*(tanh(x) + 1)),x)`output `-2/(2*exp(2*x) + exp(4*x) + 1)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{\operatorname{sech}^4(x)}{1 + \tanh(x)} dx = -\frac{2}{e^{4x} + 2e^{2x} + 1}$$

input `int(sech(x)^4/(1+tanh(x)),x)`output `( - 2)/(e**(4*x) + 2*e**(2*x) + 1)`

### 3.100 $\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx$

Optimal result	787
Mathematica [A] (verified)	787
Rubi [A] (verified)	788
Maple [B] (verified)	789
Fricas [B] (verification not implemented)	790
Sympy [F]	790
Maxima [B] (verification not implemented)	791
Giac [A] (verification not implemented)	791
Mupad [B] (verification not implemented)	791
Reduce [B] (verification not implemented)	792

#### Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx = \frac{1}{2} \arctan(\sinh(x)) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \operatorname{sech}(x) \tanh(x)$$

output `1/2*arctan(sinh(x))+1/3*sech(x)^3+1/2*sech(x)*tanh(x)`

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx = \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \operatorname{sech}(x) \tanh(x)$$

input `Integrate[Sech[x]^5/(1 + Tanh[x]), x]`

output `ArcTan[Tanh[x/2]] + Sech[x]^3/3 + (Sech[x]*Tanh[x])/2`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 3982, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^5(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^5}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3982} \\
 & \int \operatorname{sech}^3(x) dx + \frac{\operatorname{sech}^3(x)}{3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\operatorname{sech}^3(x)}{3} + \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{\int \operatorname{sech}(x) dx}{2} + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \tanh(x) \operatorname{sech}(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \tanh(x) \operatorname{sech}(x) \\
 & \quad \downarrow \text{4257} \\
 & \frac{1}{2} \arctan(\sinh(x)) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \tanh(x) \operatorname{sech}(x)
 \end{aligned}$$

input `Int [Sech [x]^5/(1 + Tanh [x]), x]`

output `ArcTan [Sinh [x]]/2 + Sech [x]^3/3 + (Sech [x]*Tanh [x])/2`

## Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3982 `Int[((d_)*sec[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Simp[d^2*((m - 2)/(a*(m + n - 1))) Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !IntegerQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 4255 `Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs.  $2(18) = 36$ .

Time = 54.64 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.71

method	result	size
default	$\frac{-\tanh\left(\frac{x}{2}\right)^5 + 2\tanh\left(\frac{x}{2}\right)^4 + \tanh\left(\frac{x}{2}\right) + \frac{2}{3}}{\left(\tanh\left(\frac{x}{2}\right)^2 + 1\right)^3} + \arctan\left(\tanh\left(\frac{x}{2}\right)\right)$	41
risch	$\frac{e^x(3e^{4x} + 8e^{2x} - 3)}{3(e^{2x} + 1)^3} + \frac{i \ln(e^x + i)}{2} - \frac{i \ln(e^x - i)}{2}$	46

input `int(sech(x)^5/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output  $2*(-1/2*\tanh(1/2*x)^5+\tanh(1/2*x)^4+1/2*\tanh(1/2*x)+1/3)/(\tanh(1/2*x)^2+1)^3+\arctan(\tanh(1/2*x))$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 288 vs. 2(18) = 36.

Time = 0.08 (sec) , antiderivative size = 288, normalized size of antiderivative = 12.00

$$\int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx$$

$$= \frac{3 \cosh(x)^5 + 15 \cosh(x) \sinh(x)^4 + 3 \sinh(x)^5 + 2(15 \cosh(x)^2 + 4) \sinh(x)^3 + 8 \cosh(x)^3 + 6(5 \cosh(x)^2 + 1) \sinh(x) + \arctan(\cosh(x) + \sinh(x))}{\cosh(x)^6 + 6 \cosh(x) \sinh(x)^5 + \sinh(x)^6 + 3(5 \cosh(x)^2 + 1) \sinh(x)^4 + 3 \cosh(x)^4 + 4(5 \cosh(x)^3 + 3 \cosh(x)) \sinh(x)^3 + 3(5 \cosh(x)^4 + 6 \cosh(x)^2 + 1) \sinh(x)^2 + 3 \cosh(x)^2 + 6(\cosh(x)^5 + 2 \cosh(x)^3 + \cosh(x)) \sinh(x) + 1}$$

input `integrate(sech(x)^5/(1+tanh(x)),x, algorithm="fricas")`

output  $1/3*(3*\cosh(x)^5 + 15*\cosh(x)*\sinh(x)^4 + 3*\sinh(x)^5 + 2*(15*\cosh(x)^2 + 4)*\sinh(x)^3 + 8*\cosh(x)^3 + 6*(5*\cosh(x)^3 + 4*\cosh(x))*\sinh(x)^2 + 3*(\cosh(x)^6 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6 + 3*(5*\cosh(x)^2 + 1)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(5*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 3*(5*\cosh(x)^4 + 6*\cosh(x)^2 + 1)*\sinh(x)^2 + 3*\cosh(x)^2 + 6*(\cosh(x)^5 + 2*\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)*\arctan(\cosh(x) + \sinh(x)) + 3*(5*\cosh(x)^4 + 8*\cosh(x)^2 - 1)*\sinh(x) - 3*\cosh(x))/(\cosh(x)^6 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6 + 3*(5*\cosh(x)^2 + 1)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(5*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 3*(5*\cosh(x)^4 + 6*\cosh(x)^2 + 1)*\sinh(x)^2 + 3*\cosh(x)^2 + 6*(\cosh(x)^5 + 2*\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)$

### Sympy [F]

$$\int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^5(x)}{\tanh(x) + 1} dx$$

input `integrate(sech(x)**5/(1+tanh(x)),x)`

output `Integral(sech(x)**5/(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 49 vs.  $2(18) = 36$ .

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.04

$$\int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx = \frac{3e^{(-x)} + 8e^{(-3x)} - 3e^{(-5x)}}{3(3e^{(-2x)} + 3e^{(-4x)} + e^{(-6x)} + 1)} - \arctan(e^{(-x)})$$

input `integrate(sech(x)^5/(1+tanh(x)),x, algorithm="maxima")`

output `1/3*(3*e^(-x) + 8*e^(-3*x) - 3*e^(-5*x))/(3*e^(-2*x) + 3*e^(-4*x) + e^(-6*x) + 1) - arctan(e^(-x))`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.29

$$\int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx = \frac{3e^{(5x)} + 8e^{(3x)} - 3e^x}{3(e^{(2x)} + 1)^3} + \arctan(e^x)$$

input `integrate(sech(x)^5/(1+tanh(x)),x, algorithm="giac")`

output `1/3*(3*e^(5*x) + 8*e^(3*x) - 3*e^x)/(e^(2*x) + 1)^3 + arctan(e^x)`

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.54

$$\begin{aligned} & \int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx \\ &= \operatorname{atan}(e^x) + \frac{e^x}{e^{2x} + 1} - \frac{8e^x}{3(3e^{2x} + 3e^{4x} + e^{6x} + 1)} + \frac{2e^x}{3(2e^{2x} + e^{4x} + 1)} \end{aligned}$$

input `int(1/(cosh(x)^5*(tanh(x) + 1)),x)`



output `atan(exp(x)) + exp(x)/(exp(2*x) + 1) - (8*exp(x))/(3*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) + (2*exp(x))/(3*(2*exp(2*x) + exp(4*x) + 1))`

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 85, normalized size of antiderivative = 3.54

$$\int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx$$

$$= \frac{3e^{6x} \operatorname{atan}(e^x) + 9e^{4x} \operatorname{atan}(e^x) + 9e^{2x} \operatorname{atan}(e^x) + 3 \operatorname{atan}(e^x) + 3e^{5x} + 8e^{3x} - 3e^x}{3e^{6x} + 9e^{4x} + 9e^{2x} + 3}$$

input `int(sech(x)^5/(1+tanh(x)),x)`

output `(3*e**(6*x)*atan(e**x) + 9*e**(4*x)*atan(e**x) + 9*e**(2*x)*atan(e**x) + 3*atan(e**x) + 3*e**(5*x) + 8*e**(3*x) - 3*e**x)/(3*(e**(6*x) + 3*e**(4*x) + 3*e**(2*x) + 1))`

### 3.101 $\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx$

Optimal result	793
Mathematica [A] (verified)	793
Rubi [A] (verified)	794
Maple [A] (verified)	795
Fricas [B] (verification not implemented)	796
Sympy [F]	796
Maxima [B] (verification not implemented)	797
Giac [A] (verification not implemented)	797
Mupad [B] (verification not implemented)	798
Reduce [B] (verification not implemented)	798

#### Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx = -\frac{2}{3}(1-\tanh(x))^3 + \frac{1}{4}(1-\tanh(x))^4$$

output `-2/3*(1-tanh(x))^3+1/4*(1-tanh(x))^4`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx = \frac{1}{12} \tanh(x) (12 - 6 \tanh(x) - 4 \tanh^2(x) + 3 \tanh^3(x))$$

input `Integrate[Sech[x]^6/(1 + Tanh[x]), x]`

output `(Tanh[x]*(12 - 6*Tanh[x] - 4*Tanh[x]^2 + 3*Tanh[x]^3))/12`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3042, 3968, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{sech}^6(x)}{\tanh(x) + 1} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sec(ix)^6}{1 - i \tan(ix)} dx \\ & \quad \downarrow \text{3968} \\ & \int (1 - \tanh(x))^2 (\tanh(x) + 1) d \tanh(x) \\ & \quad \downarrow \text{49} \\ & \int (2(1 - \tanh(x))^2 - (1 - \tanh(x))^3) d \tanh(x) \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4}(1 - \tanh(x))^4 - \frac{2}{3}(1 - \tanh(x))^3 \end{aligned}$$

input `Int [Sech[x]^6/(1 + Tanh[x]), x]`

output `(-2*(1 - Tanh[x])^3)/3 + (1 - Tanh[x])^4/4`

## Definitions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int  
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]  
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3968 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)  
, x_Symbol] := Simp[1/(a^(m - 2)*b*f) Subst[Int[(a - x)^(m/2 - 1)*(a + x)  
]^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] &&  
EqQ[a^2 + b^2, 0] && IntegerQ[m/2]`

## Maple [A] (verified)

Time = 8.99 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
risch	$-\frac{4(4e^{2x}+1)}{3(e^{2x}+1)^4}$	19
derivativedivides	$\frac{\tanh(x)^4}{4} - \frac{\tanh(x)^3}{3} - \frac{\tanh(x)^2}{2} + \tanh(x)$	22
default	$\frac{\tanh(x)^4}{4} - \frac{\tanh(x)^3}{3} - \frac{\tanh(x)^2}{2} + \tanh(x)$	22
parallelrisch	$\frac{800 \sinh(2x)+249+200 \sinh(4x)-117 \cosh(4x)-468 \cosh(2x)}{300 \cosh(4x)+1200 \cosh(2x)+900}$	44

input `int(sech(x)^6/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-4/3*(4*exp(2*x)+1)/(exp(2*x)+1)^4`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 140 vs.  $2(17) = 34$ .

Time = 0.08 (sec) , antiderivative size = 140, normalized size of antiderivative = 5.60

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx =$$

$$\frac{-4/3 (5 \cosh(x) + 3 \sinh(x))}{3 (\cosh(x)^7 + 7 \cosh(x) \sinh(x)^6 + \sinh(x)^7 + (21 \cosh(x)^2 + 4) \sinh(x)^5 + 4 \cosh(x)^5 + 5 (7 \cosh(x)^3 + 4 \cosh(x)) \sinh(x)^4 + (35 \cosh(x)^4 + 40 \cosh(x)^2 + 6) \sinh(x)^3 + 6 \cosh(x)^3 + (21 \cosh(x)^5 + 40 \cosh(x)^3 + 18 \cosh(x)) \sinh(x)^2 + (7 \cosh(x)^6 + 20 \cosh(x)^4 + 18 \cosh(x)^2 + 3) \sinh(x) + 5 \cosh(x)}$$

input `integrate(sech(x)^6/(1+tanh(x)),x, algorithm="fricas")`

output `-4/3*(5*cosh(x) + 3*sinh(x))/(cosh(x)^7 + 7*cosh(x)*sinh(x)^6 + sinh(x)^7 + (21*cosh(x)^2 + 4)*sinh(x)^5 + 4*cosh(x)^5 + 5*(7*cosh(x)^3 + 4*cosh(x))*sinh(x)^4 + (35*cosh(x)^4 + 40*cosh(x)^2 + 6)*sinh(x)^3 + 6*cosh(x)^3 + (21*cosh(x)^5 + 40*cosh(x)^3 + 18*cosh(x))*sinh(x)^2 + (7*cosh(x)^6 + 20*cosh(x)^4 + 18*cosh(x)^2 + 3)*sinh(x) + 5*cosh(x))`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^6(x)}{\tanh(x) + 1} dx$$

input `integrate(sech(x)**6/(1+tanh(x)),x)`

output `Integral(sech(x)**6/(tanh(x) + 1), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 93 vs.  $2(17) = 34$ .

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.72

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx = \frac{16 e^{(-2x)}}{3(4 e^{(-2x)} + 6 e^{(-4x)} + 4 e^{(-6x)} + e^{(-8x)} + 1)} + \frac{8 e^{(-4x)}}{4 e^{(-2x)} + 6 e^{(-4x)} + 4 e^{(-6x)} + e^{(-8x)} + 1} + \frac{4}{3(4 e^{(-2x)} + 6 e^{(-4x)} + 4 e^{(-6x)} + e^{(-8x)} + 1)}$$

input `integrate(sech(x)^6/(1+tanh(x)),x, algorithm="maxima")`

output `16/3*e^(-2*x)/(4*e^(-2*x) + 6*e^(-4*x) + 4*e^(-6*x) + e^(-8*x) + 1) + 8*e^(-4*x)/(4*e^(-2*x) + 6*e^(-4*x) + 4*e^(-6*x) + e^(-8*x) + 1) + 4/3/(4*e^(-2*x) + 6*e^(-4*x) + 4*e^(-6*x) + e^(-8*x) + 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx = -\frac{4(4e^{(2x)} + 1)}{3(e^{(2x)} + 1)^4}$$

input `integrate(sech(x)^6/(1+tanh(x)),x, algorithm="giac")`

output `-4/3*(4*e^(2*x) + 1)/(e^(2*x) + 1)^4`

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx = -\frac{4(4e^{2x} + 1)}{3(e^{2x} + 1)^4}$$

input `int(1/(cosh(x)^6*(tanh(x) + 1)),x)`output `-(4*(4*exp(2*x) + 1))/(3*(exp(2*x) + 1)^4)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\int \frac{\operatorname{sech}^6(x)}{1 + \tanh(x)} dx = \frac{-16e^{2x} - 4}{3e^{8x} + 12e^{6x} + 18e^{4x} + 12e^{2x} + 3}$$

input `int(sech(x)^6/(1+tanh(x)),x)`output `(4*( - 4*e**(2*x) - 1))/(3*(e**(8*x) + 4*e**(6*x) + 6*e**(4*x) + 4*e**(2*x) + 1))`

### 3.102 $\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx$

Optimal result	799
Mathematica [A] (verified)	799
Rubi [A] (verified)	800
Maple [B] (verified)	802
Fricas [B] (verification not implemented)	802
Sympy [F]	803
Maxima [B] (verification not implemented)	804
Giac [A] (verification not implemented)	804
Mupad [B] (verification not implemented)	805
Reduce [B] (verification not implemented)	805

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx = \frac{3}{8} \arctan(\sinh(x)) + \frac{\operatorname{sech}^5(x)}{5} + \frac{3}{8} \operatorname{sech}(x) \tanh(x) + \frac{1}{4} \operatorname{sech}^3(x) \tanh(x)$$

output

`3/8*arctan(sinh(x))+1/5*sech(x)^5+3/8*sech(x)*tanh(x)+1/4*sech(x)^3*tanh(x)`

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx = \frac{1}{40} \left( 30 \arctan \left( \tanh \left( \frac{x}{2} \right) \right) + 8 \operatorname{sech}^5(x) + 15 \operatorname{sech}(x) \tanh(x) + 10 \operatorname{sech}^3(x) \tanh(x) \right)$$

input

`Integrate[Sech[x]^7/(1 + Tanh[x]), x]`



output

```
(30*ArcTan[Tanh[x/2]] + 8*Sech[x]^5 + 15*Sech[x]*Tanh[x] + 10*Sech[x]^3*Tanh[x])/40
```

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.15, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 3982, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^7(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^7}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3982} \\
 & \int \operatorname{sech}^5(x) dx + \frac{\operatorname{sech}^5(x)}{5} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\operatorname{sech}^5(x)}{5} + \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \int \operatorname{sech}^3(x) dx + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x) \\
 & \quad \downarrow \text{3042} \\
 & \frac{3}{4} \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x) \\
 & \quad \downarrow \text{4255} \\
 & \frac{3}{4} \left( \frac{\int \operatorname{sech}(x) dx}{2} + \frac{1}{2} \tanh(x) \operatorname{sech}(x) \right) + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{3}{4} \left( \frac{1}{2} \tanh(x) \operatorname{sech}(x) + \frac{1}{2} \int \csc \left( ix + \frac{\pi}{2} \right) dx \right) + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x)$$

↓ 4257

$$\frac{3}{4} \left( \frac{1}{2} \arctan(\sinh(x)) + \frac{1}{2} \tanh(x) \operatorname{sech}(x) \right) + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x)$$

input `Int[Sech[x]^7/(1 + Tanh[x]),x]`

output `Sech[x]^5/5 + (Sech[x]^3*Tanh[x])/4 + (3*(ArcTan[Sinh[x]]/2 + (Sech[x]*Tanh[x])/2))/4`

### Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3982 `Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Simp[d^2*((m - 2)/(a*(m + n - 1)) Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !IntegerQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 66 vs.  $2(26) = 52$ .

Time = 0.17 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.97

$$\frac{-\frac{5 \tanh\left(\frac{x}{2}\right)^9}{4} + 2 \tanh\left(\frac{x}{2}\right)^8 - \frac{\tanh\left(\frac{x}{2}\right)^7}{2} + 4 \tanh\left(\frac{x}{2}\right)^4 + \frac{\tanh\left(\frac{x}{2}\right)^3}{2} + \frac{5 \tanh\left(\frac{x}{2}\right)}{4} + \frac{2}{5} + \frac{3 \arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{4}}{\left(\tanh\left(\frac{x}{2}\right)^2 + 1\right)^5}$$

input `int(sech(x)^7/(1+tanh(x)),x)`

output `2*(-5/8*tanh(1/2*x)^9+tanh(1/2*x)^8-1/4*tanh(1/2*x)^7+2*tanh(1/2*x)^4+1/4*tanh(1/2*x)^3+5/8*tanh(1/2*x)+1/5)/(tanh(1/2*x)^2+1)^5+3/4*arctan(tanh(1/2*x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 670 vs.  $2(26) = 52$ .

Time = 0.08 (sec) , antiderivative size = 670, normalized size of antiderivative = 19.71

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sech(x)^7/(1+tanh(x)),x, algorithm="fricas")`

output

```

1/20*(15*cosh(x)^9 + 135*cosh(x)*sinh(x)^8 + 15*sinh(x)^9 + 10*(54*cosh(x)
^2 + 7)*sinh(x)^7 + 70*cosh(x)^7 + 70*(18*cosh(x)^3 + 7*cosh(x))*sinh(x)^6
+ 2*(945*cosh(x)^4 + 735*cosh(x)^2 + 64)*sinh(x)^5 + 128*cosh(x)^5 + 10*(
189*cosh(x)^5 + 245*cosh(x)^3 + 64*cosh(x))*sinh(x)^4 + 10*(126*cosh(x)^6
+ 245*cosh(x)^4 + 128*cosh(x)^2 - 7)*sinh(x)^3 - 70*cosh(x)^3 + 10*(54*cos
h(x)^7 + 147*cosh(x)^5 + 128*cosh(x)^3 - 21*cosh(x))*sinh(x)^2 + 15*(cosh(
x)^10 + 10*cosh(x)*sinh(x)^9 + sinh(x)^10 + 5*(9*cosh(x)^2 + 1)*sinh(x)^8
+ 5*cosh(x)^8 + 40*(3*cosh(x)^3 + cosh(x))*sinh(x)^7 + 10*(21*cosh(x)^4 +
14*cosh(x)^2 + 1)*sinh(x)^6 + 10*cosh(x)^6 + 4*(63*cosh(x)^5 + 70*cosh(x)^
3 + 15*cosh(x))*sinh(x)^5 + 10*(21*cosh(x)^6 + 35*cosh(x)^4 + 15*cosh(x)^2
+ 1)*sinh(x)^4 + 10*cosh(x)^4 + 40*(3*cosh(x)^7 + 7*cosh(x)^5 + 5*cosh(x)
^3 + cosh(x))*sinh(x)^3 + 5*(9*cosh(x)^8 + 28*cosh(x)^6 + 30*cosh(x)^4 + 1
2*cosh(x)^2 + 1)*sinh(x)^2 + 5*cosh(x)^2 + 10*(cosh(x)^9 + 4*cosh(x)^7 + 6
*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*sinh(x) + 1)*arctan(cosh(x) + sinh(x))
+ 5*(27*cosh(x)^8 + 98*cosh(x)^6 + 128*cosh(x)^4 - 42*cosh(x)^2 - 3)*sinh
(x) - 15*cosh(x))/(cosh(x)^10 + 10*cosh(x)*sinh(x)^9 + sinh(x)^10 + 5*(9*c
osh(x)^2 + 1)*sinh(x)^8 + 5*cosh(x)^8 + 40*(3*cosh(x)^3 + cosh(x))*sinh(x)
^7 + 10*(21*cosh(x)^4 + 14*cosh(x)^2 + 1)*sinh(x)^6 + 10*cosh(x)^6 + 4*(63
*cosh(x)^5 + 70*cosh(x)^3 + 15*cosh(x))*sinh(x)^5 + 10*(21*cosh(x)^6 + 35*
cosh(x)^4 + 15*cosh(x)^2 + 1)*sinh(x)^4 + 10*cosh(x)^4 + 40*(3*cosh(x)^...

```

## Sympy [F]

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \int \frac{\operatorname{sech}^7(x)}{\tanh(x) + 1} dx$$

input

```
integrate(sech(x)**7/(1+tanh(x)),x)
```

output

```
Integral(sech(x)**7/(tanh(x) + 1), x)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 73 vs.  $2(26) = 52$ .

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.15

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \frac{15e^{-x} + 70e^{-3x} + 128e^{-5x} - 70e^{-7x} - 15e^{-9x}}{20(5e^{-2x} + 10e^{-4x} + 10e^{-6x} + 5e^{-8x} + e^{-10x} + 1)} - \frac{3}{4} \arctan(e^{-x})$$

input `integrate(sech(x)^7/(1+tanh(x)),x, algorithm="maxima")`

output `1/20*(15*e^(-x) + 70*e^(-3*x) + 128*e^(-5*x) - 70*e^(-7*x) - 15*e^(-9*x))/  
(5*e^(-2*x) + 10*e^(-4*x) + 10*e^(-6*x) + 5*e^(-8*x) + e^(-10*x) + 1) - 3/  
4*arctan(e^(-x))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.32

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \frac{15e^{9x} + 70e^{7x} + 128e^{5x} - 70e^{3x} - 15e^x}{20(e^{2x} + 1)^5} + \frac{3}{4} \arctan(e^x)$$

input `integrate(sech(x)^7/(1+tanh(x)),x, algorithm="giac")`

output `1/20*(15*e^(9*x) + 70*e^(7*x) + 128*e^(5*x) - 70*e^(3*x) - 15*e^x)/(e^(2*x)  
) + 1)^5 + 3/4*arctan(e^x)`

**Mupad [B] (verification not implemented)**

Time = 2.25 (sec) , antiderivative size = 137, normalized size of antiderivative = 4.03

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \frac{3 \operatorname{atan}(e^x)}{4} - \frac{32 e^{3x}}{5 (5 e^{2x} + 10 e^{4x} + 10 e^{6x} + 5 e^{8x} + e^{10x} + 1)} - \frac{12 e^x}{5 (4 e^{2x} + 6 e^{4x} + 4 e^{6x} + e^{8x} + 1)} + \frac{3 e^x}{4 (e^{2x} + 1)} + \frac{2 e^x}{5 (3 e^{2x} + 3 e^{4x} + e^{6x} + 1)} + \frac{e^x}{2 (2 e^{2x} + e^{4x} + 1)}$$

input `int(1/(cosh(x)^7*(tanh(x) + 1)),x)`output `(3*atan(exp(x)))/4 - (32*exp(3*x))/(5*(5*exp(2*x) + 10*exp(4*x) + 10*exp(6*x) + 5*exp(8*x) + exp(10*x) + 1)) - (12*exp(x))/(5*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) + (3*exp(x))/(4*(exp(2*x) + 1)) + (2*exp(x))/(5*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) + exp(x)/(2*(2*exp(2*x) + exp(4*x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 135, normalized size of antiderivative = 3.97

$$\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx = \frac{15e^{10x} \operatorname{atan}(e^x) + 75e^{8x} \operatorname{atan}(e^x) + 150e^{6x} \operatorname{atan}(e^x) + 150e^{4x} \operatorname{atan}(e^x) + 75e^{2x} \operatorname{atan}(e^x) + 15 \operatorname{atan}(e^x) + 15}{20e^{10x} + 100e^{8x} + 200e^{6x} + 200e^{4x} + 100e^{2x} + 20}$$

input `int(sech(x)^7/(1+tanh(x)),x)`output `(15*e**(10*x)*atan(e**x) + 75*e**(8*x)*atan(e**x) + 150*e**(6*x)*atan(e**x) + 150*e**(4*x)*atan(e**x) + 75*e**(2*x)*atan(e**x) + 15*atan(e**x) + 15*e**(9*x) + 70*e**(7*x) + 128*e**(5*x) - 70*e**(3*x) - 15*e**x)/(20*(e**(10*x) + 5*e**(8*x) + 10*e**(6*x) + 10*e**(4*x) + 5*e**(2*x) + 1))`

### 3.103 $\int \frac{\operatorname{sech}^8(x)}{a+b \tanh(x)} dx$

Optimal result . . . . .	806
Mathematica [A] (verified) . . . . .	806
Rubi [A] (verified) . . . . .	807
Maple [B] (verified) . . . . .	809
Fricas [B] (verification not implemented) . . . . .	809
Sympy [F] . . . . .	810
Maxima [B] (verification not implemented) . . . . .	810
Giac [B] (verification not implemented) . . . . .	811
Mupad [B] (verification not implemented) . . . . .	812
Reduce [B] (verification not implemented) . . . . .	813

#### Optimal result

Integrand size = 13, antiderivative size = 140

$$\int \frac{\operatorname{sech}^8(x)}{a+b \tanh(x)} dx = -\frac{(a^2 - b^2)^3 \log(a + b \tanh(x))}{b^7} + \frac{a(a^4 - 3a^2b^2 + 3b^4) \tanh(x)}{b^6} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^2(x)}{2b^5} + \frac{a(a^2 - 3b^2) \tanh^3(x)}{3b^4} - \frac{(a^2 - 3b^2) \tanh^4(x)}{4b^3} + \frac{a \tanh^5(x)}{5b^2} - \frac{\tanh^6(x)}{6b}$$

output

```
-(a^2-b^2)^3*ln(a+b*tanh(x))/b^7+a*(a^4-3*a^2*b^2+3*b^4)*tanh(x)/b^6-1/2*(a^4-3*a^2*b^2+3*b^4)*tanh(x)^2/b^5+1/3*a*(a^2-3*b^2)*tanh(x)^3/b^4-1/4*(a^2-3*b^2)*tanh(x)^4/b^3+1/5*a*tanh(x)^5/b^2-1/6*tanh(x)^6/b
```

#### Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91

$$\int \frac{\operatorname{sech}^8(x)}{a+b \tanh(x)} dx = \frac{-60(a^2 - b^2)^3 \log(a + b \tanh(x)) + 15b^4(-a^2 + b^2) \operatorname{sech}^4(x) + 10b^6 \operatorname{sech}^6(x) + 60ab(a^4 - 3a^2b^2 + 3b^4) \tanh(x)}{60b^7}$$

input `Integrate[Sech[x]^8/(a + b*Tanh[x]), x]`

output  $(-60*(a^2 - b^2)^3*\text{Log}[a + b*\text{Tanh}[x]] + 15*b^4*(-a^2 + b^2)*\text{Sech}[x]^4 + 10*b^6*\text{Sech}[x]^6 + 60*a*b*(a^4 - 3*a^2*b^2 + 3*b^4)*\text{Tanh}[x] - 30*b^2*(a^2 - b^2)^2*\text{Tanh}[x]^2 + 20*a*b^3*(a^2 - 3*b^2)*\text{Tanh}[x]^3 + 12*a*b^5*\text{Tanh}[x]^5)/(60*b^7)$

### Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3987, 27, 476, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\text{sech}^8(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^8}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{3987} \\
 & \frac{\int \frac{(b^2 - b^2 \tanh^2(x))^3}{b^6(a + b \tanh(x))} d(b \tanh(x))}{b} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(b^2 - b^2 \tanh^2(x))^3}{a + b \tanh(x)} d(b \tanh(x))}{b^7} \\
 & \quad \downarrow \text{476} \\
 & \frac{\int \left( \left( \frac{3(b^2 - a^2)b^2}{a^4} + 1 \right) a^5 + b^4 \tanh^4(x)a + b^2(a^2 - 3b^2) \tanh^2(x)a - b^5 \tanh^5(x) - b^3(a^2 - 3b^2) \tanh^3(x) - b(a^4 - \right.}{b^7} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$



$$\frac{-(a^2 - b^2)^3 \log(a + b \tanh(x)) - \frac{1}{4}b^4(a^2 - 3b^2) \tanh^4(x) + \frac{1}{3}ab^3(a^2 - 3b^2) \tanh^3(x) - \frac{1}{2}b^2(a^4 - 3a^2b^2 + 3b^4) \tanh^2(x) + \frac{1}{5}ab^5 \tanh^5(x) - \frac{1}{6}b^6 \tanh^6(x)}{b^7}$$

input `Int[Sech[x]^8/(a + b*Tanh[x]),x]`

output `((-(a^2 - b^2)^3*Log[a + b*Tanh[x]]) + a*b*(a^4 - 3*a^2*b^2 + 3*b^4)*Tanh[x] - (b^2*(a^4 - 3*a^2*b^2 + 3*b^4)*Tanh[x]^2)/2 + (a*b^3*(a^2 - 3*b^2)*Tanh[x]^3)/3 - (b^4*(a^2 - 3*b^2)*Tanh[x]^4)/4 + (a*b^5*Tanh[x]^5)/5 - (b^6*Tanh[x]^6)/6)/b^7`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 476 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3987 `Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 412 vs.  $2(130) = 260$ .

Time = 0.23 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.95

$$\frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{b^7} + \frac{2\left((a^5b - 3a^3b^3 + 3ab^5) \tanh\left(\frac{x}{2}\right)^{11} + (-a^4b^2 + 3a^2b^4 - 3b^6)\right)}{b^7}$$

input `int(sech(x)^8/(a+b*tanh(x)),x)`

output 
$$\begin{aligned} & -(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)/b^7 * \ln(\tanh(1/2*x)^2*a + 2*b*\tanh(1/2*x) + a) + \\ & /b^7 * (((a^5*b - 3*a^3*b^3 + 3*a*b^5)*\tanh(1/2*x)^{11} + (-a^4*b^2 + 3*a^2*b^4 - 3*b^6) \\ & *\tanh(1/2*x)^{10} + (5*a^5*b - 41/3*a^3*b^3 + 11*a*b^5)*\tanh(1/2*x)^9 + (-4*a^4*b^2 + \\ & 10*a^2*b^4 - 6*b^6)*\tanh(1/2*x)^8 + (10*a^5*b - 26*a^3*b^3 + 106/5*a*b^5)*\tanh(1/2 \\ & *x)^7 + (-6*a^4*b^2 + 14*a^2*b^4 - 34/3*b^6)*\tanh(1/2*x)^6 + (10*a^5*b - 26*a^3*b^3 + \\ & 106/5*a*b^5)*\tanh(1/2*x)^5 + (-4*a^4*b^2 + 10*a^2*b^4 - 6*b^6)*\tanh(1/2*x)^4 + (5* \\ & a^5*b - 41/3*a^3*b^3 + 11*a*b^5)*\tanh(1/2*x)^3 + (-a^4*b^2 + 3*a^2*b^4 - 3*b^6)*\tanh \\ & (1/2*x)^2 + (a^5*b - 3*a^3*b^3 + 3*a*b^5)*\tanh(1/2*x)) / (\tanh(1/2*x)^2 + 1)^6 + 1/2 * ( \\ & a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) * \ln(\tanh(1/2*x)^2 + 1) \end{aligned}$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 5275 vs.  $2(130) = 260$ .

Time = 0.16 (sec) , antiderivative size = 5275, normalized size of antiderivative = 37.68

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="fricas")`

output `Too large to include`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx$$

input `integrate(sech(x)**8/(a+b*tanh(x)),x)`

output `Integral(sech(x)**8/(a + b*tanh(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 386 vs.  $2(130) = 260$ .

Time = 0.14 (sec) , antiderivative size = 386, normalized size of antiderivative = 2.76

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx$$

$$= \frac{2(15a^5 - 40a^3b^2 + 33ab^4 + 3(25a^5 + 5a^4b - 70a^3b^2 - 10a^2b^3 + 61ab^4 + 5b^5)e^{(-2x)} + 30(5a^5 + 2a^4b - 10a^3b^2 + 33a^2b^3 - 24a^2b^3 + 33a^2b^3 + 23b^5)e^{(-6x)} + 15(5a^5 + 4a^4b - 12a^3b^2 - 10a^2b^3 + 7a^2b^3 + 6b^5)e^{(-8x)} + 15(a^5 + a^4b - 2a^3b^2 - 2a^2b^3 + ab^4 + b^5)e^{(-10x)})}{(6b^6e^{(-2x)} + 15b^6e^{(-4x)} + 20b^6e^{(-6x)} + 15b^6e^{(-8x)} + 6b^6e^{(-10x)} + b^6e^{(-12x)} + b^6) - (a^6 - 3a^4b^2 + 3a^2b^4 - b^6)*\log(-(a - b)*e^{(-2x)} - a - b) / b^7 + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6)*\log(e^{(-2x)} + 1) / b^7}$$

input `integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="maxima")`

output `2/15*(15*a^5 - 40*a^3*b^2 + 33*a*b^4 + 3*(25*a^5 + 5*a^4*b - 70*a^3*b^2 - 10*a^2*b^3 + 61*a*b^4 + 5*b^5)*e^(-2*x) + 30*(5*a^5 + 2*a^4*b - 14*a^3*b^2 - 5*a^2*b^3 + 13*a*b^4 + 3*b^5)*e^(-4*x) + 10*(15*a^5 + 9*a^4*b - 40*a^3*b^2 - 24*a^2*b^3 + 33*a*b^4 + 23*b^5)*e^(-6*x) + 15*(5*a^5 + 4*a^4*b - 12*a^3*b^2 - 10*a^2*b^3 + 7*a*b^4 + 6*b^5)*e^(-8*x) + 15*(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*e^(-10*x))/(6*b^6*e^(-2*x) + 15*b^6*e^(-4*x) + 20*b^6*e^(-6*x) + 15*b^6*e^(-8*x) + 6*b^6*e^(-10*x) + b^6*e^(-12*x) + b^6) - (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*log(-(a - b)*e^(-2*x) - a - b) / b^7 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*log(e^(-2*x) + 1) / b^7`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 593 vs.  $2(130) = 260$ .

Time = 0.13 (sec) , antiderivative size = 593, normalized size of antiderivative = 4.24

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="giac")`

output

```

-(a^7 + a^6*b - 3*a^5*b^2 - 3*a^4*b^3 + 3*a^3*b^4 + 3*a^2*b^5 - a*b^6 - b^7)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a*b^7 + b^8) + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*log(e^(2*x) + 1)/b^7 - 1/60*(147*a^6*e^(12*x) - 441*a^4*b^2*e^(12*x) + 441*a^2*b^4*e^(12*x) - 147*b^6*e^(12*x) + 882*a^6*e^(10*x) + 120*a^5*b*e^(10*x) - 2766*a^4*b^2*e^(10*x) - 240*a^3*b^3*e^(10*x) + 2886*a^2*b^4*e^(10*x) + 120*a*b^5*e^(10*x) - 1002*b^6*e^(10*x) + 2205*a^6*e^(8*x) + 600*a^5*b*e^(8*x) - 7095*a^4*b^2*e^(8*x) - 1440*a^3*b^3*e^(8*x) + 7815*a^2*b^4*e^(8*x) + 840*a*b^5*e^(8*x) - 2925*b^6*e^(8*x) + 2940*a^6*e^(6*x) + 1200*a^5*b*e^(6*x) - 9540*a^4*b^2*e^(6*x) - 3200*a^3*b^3*e^(6*x) + 10740*a^2*b^4*e^(6*x) + 2640*a*b^5*e^(6*x) - 4780*b^6*e^(6*x) + 2205*a^6*e^(4*x) + 1200*a^5*b*e^(4*x) - 7095*a^4*b^2*e^(4*x) - 3360*a^3*b^3*e^(4*x) + 7815*a^2*b^4*e^(4*x) + 3120*a*b^5*e^(4*x) - 2925*b^6*e^(4*x) + 882*a^6*e^(2*x) + 600*a^5*b*e^(2*x) - 2766*a^4*b^2*e^(2*x) - 1680*a^3*b^3*e^(2*x) + 2886*a^2*b^4*e^(2*x) + 1464*a*b^5*e^(2*x) - 1002*b^6*e^(2*x) + 147*a^6 + 120*a^5*b - 441*a^4*b^2 - 320*a^3*b^3 + 441*a^2*b^4 + 264*a*b^5 - 147*b^6)/(b^7*(e^(2*x) + 1)^6)

```

**Mupad [B] (verification not implemented)**

Time = 2.82 (sec) , antiderivative size = 301, normalized size of antiderivative = 2.15

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx = \frac{\ln(e^{2x} + 1) (a + b)^3 (a - b)^3}{b^7} - \frac{32(a - 5b)}{5b^2 (5e^{2x} + 10e^{4x} + 10e^{6x} + 5e^{8x} + e^{10x} + 1)} - \frac{4(a^2 - 4ab + 7b^2)}{b^3 (4e^{2x} + 6e^{4x} + 4e^{6x} + e^{8x} + 1)} - \frac{\ln(a - b + ae^{2x} + be^{2x}) (a + b)^3 (a - b)^3}{b^7} - \frac{32}{3b (6e^{2x} + 15e^{4x} + 20e^{6x} + 15e^{8x} + 6e^{10x} + e^{12x} + 1)} - \frac{8(a - b)(a^2 - 2ab + b^2)}{3b^4 (3e^{2x} + 3e^{4x} + e^{6x} + 1)} - \frac{2(a + b)^2 (a - b)(a^2 - 2ab + b^2)}{b^6 (e^{2x} + 1)} - \frac{2(a + b)(a - b)(a^2 - 2ab + b^2)}{b^5 (2e^{2x} + e^{4x} + 1)}$$

input `int(1/(cosh(x)^8*(a + b*tanh(x))),x)`output `(log(exp(2*x) + 1)*(a + b)^3*(a - b)^3)/b^7 - (32*(a - 5*b))/(5*b^2*(5*exp(2*x) + 10*exp(4*x) + 10*exp(6*x) + 5*exp(8*x) + exp(10*x) + 1)) - (4*(a^2 - 4*a*b + 7*b^2))/(b^3*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) - (log(a - b + a*exp(2*x) + b*exp(2*x))*(a + b)^3*(a - b)^3)/b^7 - 32/(3*b*(6*exp(2*x) + 15*exp(4*x) + 20*exp(6*x) + 15*exp(8*x) + 6*exp(10*x) + exp(12*x) + 1)) - (8*(a - b)*(a^2 - 2*a*b + b^2))/(3*b^4*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) - (2*(a + b)^2*(a - b)*(a^2 - 2*a*b + b^2))/(b^6*(exp(2*x) + 1)) - (2*(a + b)*(a - b)*(a^2 - 2*a*b + b^2))/(b^5*(2*exp(2*x) + exp(4*x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 1798, normalized size of antiderivative = 12.84

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `int(sech(x)^8/(a+b*tanh(x)),x)`

output

```
(15***e**(12*x)*log(e**(2*x) + 1)*a**6 - 45***e**(12*x)*log(e**(2*x) + 1)*a**4
*b**2 + 45***e**(12*x)*log(e**(2*x) + 1)*a**2*b**4 - 15***e**(12*x)*log(e**(2*
x) + 1)*b**6 - 15***e**(12*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**6 + 45
***e**(12*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**4*b**2 - 45***e**(12*x)*l
og(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*b**4 + 15***e**(12*x)*log(e**(2*x)*
a + e**(2*x)*b + a - b)*b**6 + 5***e**(12*x)*a**5*b - 5***e**(12*x)*a**4*b**2
- 10***e**(12*x)*a**3*b**3 + 10***e**(12*x)*a**2*b**4 + 5***e**(12*x)*a*b**5 - 5
***e**(12*x)*b**6 + 90***e**(10*x)*log(e**(2*x) + 1)*a**6 - 270***e**(10*x)*log(
e**(2*x) + 1)*a**4*b**2 + 270***e**(10*x)*log(e**(2*x) + 1)*a**2*b**4 - 90***e
**(10*x)*log(e**(2*x) + 1)*b**6 - 90***e**(10*x)*log(e**(2*x)*a + e**(2*x)*b
+ a - b)*a**6 + 270***e**(10*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**4*b
**2 - 270***e**(10*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*b**4 + 90***e
**(10*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**6 + 225***e**(8*x)*log(e**(2
*x) + 1)*a**6 - 675***e**(8*x)*log(e**(2*x) + 1)*a**4*b**2 + 675***e**(8*x)*lo
g(e**(2*x) + 1)*a**2*b**4 - 225***e**(8*x)*log(e**(2*x) + 1)*b**6 - 225***e**(
8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**6 + 675***e**(8*x)*log(e**(2*x)
*a + e**(2*x)*b + a - b)*a**4*b**2 - 675***e**(8*x)*log(e**(2*x)*a + e**(2*x)
)*b + a - b)*a**2*b**4 + 225***e**(8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)
*b**6 - 75***e**(8*x)*a**5*b + 45***e**(8*x)*a**4*b**2 + 210***e**(8*x)*a**3*b**
3 - 150***e**(8*x)*a**2*b**4 - 135***e**(8*x)*a*b**5 + 105***e**(8*x)*b**6 + ...
```

### 3.104 $\int \frac{\operatorname{sech}^6(x)}{a+b \tanh(x)} dx$

Optimal result	814
Mathematica [A] (verified)	814
Rubi [A] (verified)	815
Maple [B] (verified)	816
Fricas [B] (verification not implemented)	817
Sympy [F]	818
Maxima [B] (verification not implemented)	819
Giac [B] (verification not implemented)	819
Mupad [B] (verification not implemented)	820
Reduce [B] (verification not implemented)	821

#### Optimal result

Integrand size = 13, antiderivative size = 83

$$\int \frac{\operatorname{sech}^6(x)}{a+b \tanh(x)} dx = \frac{(a^2 - b^2)^2 \log(a + b \tanh(x))}{b^5} - \frac{a(a^2 - 2b^2) \tanh(x)}{b^4} + \frac{(a^2 - 2b^2) \tanh^2(x)}{2b^3} - \frac{a \tanh^3(x)}{3b^2} + \frac{\tanh^4(x)}{4b}$$

output

$(a^2 - b^2)^2 \ln(a + b \tanh(x)) / b^5 - a(a^2 - 2b^2) \tanh(x) / b^4 + 1/2(a^2 - 2b^2) \tanh(x)^2 / b^3 - 1/3 a \tanh(x)^3 / b^2 + 1/4 \tanh(x)^4 / b$

#### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{sech}^6(x)}{a+b \tanh(x)} dx = \frac{12(a^2 - b^2)^2 \log(a + b \tanh(x)) + 3b^4 \operatorname{sech}^4(x) - 12ab(a^2 - 2b^2) \tanh(x) + 6b^2(a^2 - b^2) \tanh^2(x) - 4ab^3}{12b^5}$$

input

`Integrate[Sech[x]^6/(a + b*Tanh[x]), x]`

output

$$(12*(a^2 - b^2)^2*\text{Log}[a + b*\text{Tanh}[x]] + 3*b^4*\text{Sech}[x]^4 - 12*a*b*(a^2 - 2*b^2)*\text{Tanh}[x] + 6*b^2*(a^2 - b^2)*\text{Tanh}[x]^2 - 4*a*b^3*\text{Tanh}[x]^3)/(12*b^5)$$

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.99, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3987, 27, 476, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\text{sech}^6(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sec(ix)^6}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{3987} \\ & \frac{\int \frac{(b^2 - b^2 \tanh^2(x))^2}{b^4(a + b \tanh(x))} d(b \tanh(x))}{b} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{(b^2 - b^2 \tanh^2(x))^2}{a + b \tanh(x)} d(b \tanh(x))}{b^5} \\ & \quad \downarrow \text{476} \\ & \frac{\int \left( -\left( \left( 1 - \frac{2b^2}{a^2} \right) a^3 \right) - b^2 \tanh^2(x)a + b^3 \tanh^3(x) + b(a^2 - 2b^2) \tanh(x) + \frac{(a^2 - b^2)^2}{a + b \tanh(x)} \right) d(b \tanh(x))}{b^5} \\ & \quad \downarrow \text{2009} \\ & \frac{\frac{1}{2}b^2(a^2 - 2b^2) \tanh^2(x) - ab(a^2 - 2b^2) \tanh(x) + (a^2 - b^2)^2 \log(a + b \tanh(x)) - \frac{1}{3}ab^3 \tanh^3(x) + \frac{1}{4}b^4 \tanh^4(x)}{b^5} \end{aligned}$$



input `Int[Sech[x]^6/(a + b*Tanh[x]),x]`

output `((a^2 - b^2)^2*Log[a + b*Tanh[x]] - a*b*(a^2 - 2*b^2)*Tanh[x] + (b^2*(a^2 - 2*b^2)*Tanh[x]^2)/2 - (a*b^3*Tanh[x]^3)/3 + (b^4*Tanh[x]^4)/4)/b^5`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 476 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3987 `Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. 2(77) = 154.

Time = 104.51 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.67

method	result
default	$2 \frac{\left( (a^3 b - 2 a b^3) \tanh\left(\frac{x}{2}\right)^7 + (-a^2 b^2 + 2 b^4) \tanh\left(\frac{x}{2}\right)^6 + (3 a^3 b - \frac{14}{3} a b^3) \tanh\left(\frac{x}{2}\right)^5 + (-2 a^2 b^2 + 2 b^4) \tanh\left(\frac{x}{2}\right)^4 + (3 a^3 b - \frac{14}{3} a b^3) \tanh\left(\frac{x}{2}\right)^3 + \dots \right)}{\left( \tanh\left(\frac{x}{2}\right)^2 + 1 \right)^4}$
risch	$\frac{2 a^3 e^{6 x} - 2 e^{6 x} a^2 b - 2 a b^2 e^{6 x} + 2 b^3 e^{6 x} + 6 a^3 e^{4 x} - 4 a^2 b e^{4 x} - 10 e^{4 x} b^2 a + 8 b^3 e^{4 x} + 6 a^3 e^{2 x} - 2 e^{2 x} a^2 b - \frac{34 e^{2 x} a b^2}{3} + 2 b^3 e^{2 x} + 2 a^3 - \frac{10 b^2 a}{3}}{b^4 (e^{2 x} + 1)^4}$

input `int(sech(x)^6/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2/b^5 * ((a^3*b - 2*a*b^3) * \tanh(1/2*x)^7 + (-a^2*b^2 + 2*b^4) * \tanh(1/2*x)^6 + (3*a^3*b - 14/3*a*b^3) * \tanh(1/2*x)^5 + (-2*a^2*b^2 + 2*b^4) * \tanh(1/2*x)^4 + (3*a^3*b - 14/3*a*b^3) * \tanh(1/2*x)^3 + (-a^2*b^2 + 2*b^4) * \tanh(1/2*x)^2 + (a^3*b - 2*a*b^3) * \tanh(1/2*x)) / (\tanh(1/2*x)^2 + 1)^4 + 1/2 * (a^4 - 2*a^2*b^2 + b^4) * \ln(\tanh(1/2*x)^2 + 1) \\ & + (a^4 - 2*a^2*b^2 + b^4) / b^5 * \ln(\tanh(1/2*x)^2 * a + 2*b * \tanh(1/2*x) + a) \end{aligned}$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1827 vs.  $2(77) = 154$ .

Time = 0.11 (sec) , antiderivative size = 1827, normalized size of antiderivative = 22.01

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="fricas")`

output

```

1/3*(6*(a^3*b - a^2*b^2 - a*b^3 + b^4)*cosh(x)^6 + 36*(a^3*b - a^2*b^2 - a
*b^3 + b^4)*cosh(x)*sinh(x)^5 + 6*(a^3*b - a^2*b^2 - a*b^3 + b^4)*sinh(x)^
6 + 6*(3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*cosh(x)^4 + 6*(3*a^3*b - 2*a
^2*b^2 - 5*a*b^3 + 4*b^4 + 15*(a^3*b - a^2*b^2 - a*b^3 + b^4)*cosh(x)^2)*s
inh(x)^4 + 6*a^3*b - 10*a*b^3 + 24*(5*(a^3*b - a^2*b^2 - a*b^3 + b^4)*cosh
(x)^3 + (3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*cosh(x))*sinh(x)^3 + 2*(9*
a^3*b - 3*a^2*b^2 - 17*a*b^3 + 3*b^4)*cosh(x)^2 + 2*(45*(a^3*b - a^2*b^2 -
a*b^3 + b^4)*cosh(x)^4 + 9*a^3*b - 3*a^2*b^2 - 17*a*b^3 + 3*b^4 + 18*(3*a
^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*cosh(x)^2)*sinh(x)^2 + 3*((a^4 - 2*a^2
*b^2 + b^4)*cosh(x)^8 + 8*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x)^7 + (a^4
- 2*a^2*b^2 + b^4)*sinh(x)^8 + 4*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^6 + 4*(a
^4 - 2*a^2*b^2 + b^4 + 7*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^6 + 8*
(7*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x))*
sinh(x)^5 + 6*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + 2*(35*(a^4 - 2*a^2*b^2 +
b^4)*cosh(x)^4 + 3*a^4 - 6*a^2*b^2 + 3*b^4 + 30*(a^4 - 2*a^2*b^2 + b^4)*c
osh(x)^2)*sinh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 8*(7*(a^4 - 2*a^2*b^2 + b^4)
*cosh(x)^5 + 10*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b
^4)*cosh(x))*sinh(x)^3 + 4*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2 + 4*(7*(a^4 -
2*a^2*b^2 + b^4)*cosh(x)^6 + 15*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + a^4 -
2*a^2*b^2 + b^4 + 9*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^2 + 8*(...

```

## Sympy [F]

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx$$

input

```
integrate(sech(x)**6/(a+b*tanh(x)), x)
```

output

```
Integral(sech(x)**6/(a + b*tanh(x)), x)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 204 vs.  $2(77) = 154$ .

Time = 0.13 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.46

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx =$$

$$-\frac{2(3a^3 - 5ab^2 + (9a^3 + 3a^2b - 17ab^2 - 3b^3)e^{-2x}) + 3(3a^3 + 2a^2b - 5ab^2 - 4b^3)e^{-4x} + 3(a^3 + a^2b - 5ab^2 - 4b^3)e^{-6x} + 3(a^3 + a^2b - 5ab^2 - 4b^3)e^{-8x} + 3(a^3 + a^2b - 5ab^2 - 4b^3)}{3(4b^4e^{-2x} + 6b^4e^{-4x} + 4b^4e^{-6x} + b^4e^{-8x} + b^4)}$$

$$+ \frac{(a^4 - 2a^2b^2 + b^4) \log(-(a-b)e^{-2x} - a - b)}{b^5} - \frac{(a^4 - 2a^2b^2 + b^4) \log(e^{-2x} + 1)}{b^5}$$

input `integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="maxima")`

output 
$$-2/3*(3*a^3 - 5*a*b^2 + (9*a^3 + 3*a^2*b - 17*a*b^2 - 3*b^3)*e^{-2*x}) + 3*(3*a^3 + 2*a^2*b - 5*a*b^2 - 4*b^3)*e^{-4*x} + 3*(a^3 + a^2*b - a*b^2 - b^3)*e^{-6*x})/(4*b^4*e^{-2*x} + 6*b^4*e^{-4*x} + 4*b^4*e^{-6*x} + b^4*e^{-8*x} + b^4) + (a^4 - 2*a^2*b^2 + b^4)*\log(-(a - b)*e^{-2*x} - a - b)/b^5 - (a^4 - 2*a^2*b^2 + b^4)*\log(e^{-2*x} + 1)/b^5$$

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 316 vs.  $2(77) = 154$ .

Time = 0.13 (sec) , antiderivative size = 316, normalized size of antiderivative = 3.81

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx$$

$$= \frac{(a^5 + a^4b - 2a^3b^2 - 2a^2b^3 + ab^4 + b^5) \log(|ae^{2x} + be^{2x} + a - b|)}{ab^5 + b^6}$$

$$- \frac{(a^4 - 2a^2b^2 + b^4) \log(e^{2x} + 1)}{b^5}$$

$$+ \frac{25a^4e^{8x} - 50a^2b^2e^{8x} + 25b^4e^{8x} + 100a^4e^{6x} + 24a^3be^{6x} - 224a^2b^2e^{6x} - 24ab^3e^{6x} + 124b^4e^{6x} - 25a^4e^{4x} + 50a^2b^2e^{4x} - 25b^4e^{4x} + 100a^4e^{2x} + 24a^3be^{2x} - 224a^2b^2e^{2x} - 24ab^3e^{2x} + 124b^4e^{2x} - 25a^4 + 50a^2b^2 - 25b^4}{b^5}$$

input `integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="giac")`

output

```
(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*log(abs(a*e^(2*x) + b*
e^(2*x) + a - b))/(a*b^5 + b^6) - (a^4 - 2*a^2*b^2 + b^4)*log(e^(2*x) + 1)
/b^5 + 1/12*(25*a^4*e^(8*x) - 50*a^2*b^2*e^(8*x) + 25*b^4*e^(8*x) + 100*a^
4*e^(6*x) + 24*a^3*b*e^(6*x) - 224*a^2*b^2*e^(6*x) - 24*a*b^3*e^(6*x) + 12
4*b^4*e^(6*x) + 150*a^4*e^(4*x) + 72*a^3*b*e^(4*x) - 348*a^2*b^2*e^(4*x) -
120*a*b^3*e^(4*x) + 246*b^4*e^(4*x) + 100*a^4*e^(2*x) + 72*a^3*b*e^(2*x)
- 224*a^2*b^2*e^(2*x) - 136*a*b^3*e^(2*x) + 124*b^4*e^(2*x) + 25*a^4 + 24*
a^3*b - 50*a^2*b^2 - 40*a*b^3 + 25*b^4)/(b^5*(e^(2*x) + 1)^4)
```

**Mupad [B] (verification not implemented)**

Time = 2.35 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.04

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx = \frac{4}{b(4e^{2x} + 6e^{4x} + 4e^{6x} + e^{8x} + 1)} + \frac{2(a-b)^2}{b^3(2e^{2x} + e^{4x} + 1)}$$

$$+ \frac{8(a-3b)}{3b^2(3e^{2x} + 3e^{4x} + e^{6x} + 1)} + \frac{2(a+b)(a-b)^2}{b^4(e^{2x} + 1)}$$

$$+ \frac{\ln(a-b + ae^{2x} + be^{2x})(a+b)^2(a-b)^2}{b^5}$$

$$- \frac{\ln(e^{2x} + 1)(a+b)^2(a-b)^2}{b^5}$$

input

```
int(1/(cosh(x)^6*(a + b*tanh(x))),x)
```

output

```
4/(b*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) + (2*(a - b)^2
)/(b^3*(2*exp(2*x) + exp(4*x) + 1)) + (8*(a - 3*b))/(3*b^2*(3*exp(2*x) + 3
*exp(4*x) + exp(6*x) + 1)) + (2*(a + b)*(a - b)^2)/(b^4*(exp(2*x) + 1)) +
(log(a - b + a*exp(2*x) + b*exp(2*x))*(a + b)^2*(a - b)^2)/b^5 - (log(exp(
2*x) + 1)*(a + b)^2*(a - b)^2)/b^5
```

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 893, normalized size of antiderivative = 10.76

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `int(sech(x)^6/(a+b*tanh(x)),x)`

output

```
( - 6**e**(8*x)*log(e**(2*x) + 1)*a**4 + 12**e**(8*x)*log(e**(2*x) + 1)*a**2
*b**2 - 6**e**(8*x)*log(e**(2*x) + 1)*b**4 + 6**e**(8*x)*log(e**(2*x)*a + e
*(2*x)*b + a - b)*a**4 - 12**e**(8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*
a**2*b**2 + 6**e**(8*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**4 - 3**e**(8
*x)*a**3*b + 3**e**(8*x)*a**2*b**2 + 3**e**(8*x)*a*b**3 - 3**e**(8*x)*b**4 -
24**e**(6*x)*log(e**(2*x) + 1)*a**4 + 48**e**(6*x)*log(e**(2*x) + 1)*a**2*b**
*2 - 24**e**(6*x)*log(e**(2*x) + 1)*b**4 + 24**e**(6*x)*log(e**(2*x)*a + e**
(2*x)*b + a - b)*a**4 - 48**e**(6*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a
**2*b**2 + 24**e**(6*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**4 - 36**e**(
4*x)*log(e**(2*x) + 1)*a**4 + 72**e**(4*x)*log(e**(2*x) + 1)*a**2*b**2 - 36
**e**(4*x)*log(e**(2*x) + 1)*b**4 + 36**e**(4*x)*log(e**(2*x)*a + e**(2*x)*b
+ a - b)*a**4 - 72**e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*b**
2 + 36**e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**4 + 18**e**(4*x)*a*
*3*b - 6**e**(4*x)*a**2*b**2 - 42**e**(4*x)*a*b**3 + 30**e**(4*x)*b**4 - 24**e
**(2*x)*log(e**(2*x) + 1)*a**4 + 48**e**(2*x)*log(e**(2*x) + 1)*a**2*b**2 -
24**e**(2*x)*log(e**(2*x) + 1)*b**4 + 24**e**(2*x)*log(e**(2*x)*a + e**(2*x)
)*b + a - b)*a**4 - 48**e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2*
b**2 + 24**e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**4 + 24**e**(2*x)
*a**3*b - 56**e**(2*x)*a*b**3 - 6*log(e**(2*x) + 1)*a**4 + 12*log(e**(2*x)
+ 1)*a**2*b**2 - 6*log(e**(2*x) + 1)*b**4 + 6*log(e**(2*x)*a + e**(2*x)...
```

### 3.105 $\int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx$

Optimal result	822
Mathematica [A] (verified)	822
Rubi [A] (verified)	823
Maple [B] (verified)	824
Fricas [B] (verification not implemented)	825
Sympy [F]	826
Maxima [B] (verification not implemented)	826
Giac [B] (verification not implemented)	827
Mupad [B] (verification not implemented)	827
Reduce [B] (verification not implemented)	828

#### Optimal result

Integrand size = 13, antiderivative size = 40

$$\int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx = -\frac{(a^2 - b^2) \log(a + b \tanh(x))}{b^3} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

output  $-(a^2-b^2)*\ln(a+b*\tanh(x))/b^3+a*\tanh(x)/b^2-1/2*\tanh(x)^2/b$

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx = \frac{2(-a^2 + b^2) \log(a + b \tanh(x)) + 2ab \tanh(x) - b^2 \tanh^2(x)}{2b^3}$$

input `Integrate[Sech[x]^4/(a + b*Tanh[x]), x]`

output  $(2*(-a^2 + b^2)*\operatorname{Log}[a + b*\operatorname{Tanh}[x]] + 2*a*b*\operatorname{Tanh}[x] - b^2*\operatorname{Tanh}[x]^2)/(2*b^3)$

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3987, 27, 476, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^4}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{3987} \\
 & \frac{\int \frac{b^2 - b^2 \tanh^2(x)}{b^2(a + b \tanh(x))} d(b \tanh(x))}{b} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{b^2 - b^2 \tanh^2(x)}{a + b \tanh(x)} d(b \tanh(x))}{b^3} \\
 & \quad \downarrow \text{476} \\
 & \frac{\int \left( a - b \tanh(x) + \frac{b^2 - a^2}{a + b \tanh(x)} \right) d(b \tanh(x))}{b^3} \\
 & \quad \downarrow \text{2009} \\
 & \frac{-(a^2 - b^2) \log(a + b \tanh(x)) + ab \tanh(x) - \frac{1}{2} b^2 \tanh^2(x)}{b^3}
 \end{aligned}$$

input `Int [Sech [x]^4/(a + b*Tanh [x]), x]`

output `((-(a^2 - b^2)*Log[a + b*Tanh[x]]) + a*b*Tanh[x] - (b^2*Tanh[x]^2)/2)/b^3`



**Defintions of rubi rules used**

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_) /; FreeQ[b, x]]`

rule 476 `Int[((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, n, x}] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3987 `Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. 2(38) = 76.

Time = 21.30 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.48

method	result	size
default	$-\frac{(a^2-b^2) \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{b^3} + \frac{2\left(\tanh\left(\frac{x}{2}\right)^3 ab - b^2 \tanh\left(\frac{x}{2}\right)^2 + \tanh\left(\frac{x}{2}\right) ab\right)}{\left(\tanh\left(\frac{x}{2}\right)^2 + 1\right)^2} + (a^2 - b^2) \ln\left(\tanh\left(\frac{x}{2}\right)^2 + 1\right)}$	99
risch	$-\frac{2(e^{2x}a - e^{2x}b + a)}{(e^{2x} + 1)^2 b^2} + \frac{\ln(e^{2x} + 1)a^2}{b^3} - \frac{\ln(e^{2x} + 1)}{b} - \frac{\ln\left(e^{2x} + \frac{a-b}{a+b}\right)a^2}{b^3} + \frac{\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b}$	102

input `int(sech(x)^4/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

output

```
-(a^2-b^2)/b^3*ln(tanh(1/2*x)^2*a+2*b*tanh(1/2*x)+a)+2/b^3*((tanh(1/2*x)^3
*a*b-b^2*tanh(1/2*x)^2+tanh(1/2*x)*a*b)/(tanh(1/2*x)^2+1)^2+1/2*(a^2-b^2)*
ln(tanh(1/2*x)^2+1))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 430 vs.  $2(38) = 76$ .

Time = 0.11 (sec) , antiderivative size = 430, normalized size of antiderivative = 10.75

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx =$$

$$\frac{2(ab - b^2) \cosh(x)^2 + 4(ab - b^2) \cosh(x) \sinh(x) + 2(ab - b^2) \sinh(x)^2 + 2ab + ((a^2 - b^2) \cosh(x))}{-}$$

input

```
integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
-(2*(a*b - b^2)*cosh(x)^2 + 4*(a*b - b^2)*cosh(x)*sinh(x) + 2*(a*b - b^2)*
sinh(x)^2 + 2*a*b + ((a^2 - b^2)*cosh(x)^4 + 4*(a^2 - b^2)*cosh(x)*sinh(x)
^3 + (a^2 - b^2)*sinh(x)^4 + 2*(a^2 - b^2)*cosh(x)^2 + 2*(3*(a^2 - b^2)*co
sh(x)^2 + a^2 - b^2)*sinh(x)^2 + a^2 - b^2 + 4*((a^2 - b^2)*cosh(x)^3 + (a
^2 - b^2)*cosh(x))*sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(
x))) - ((a^2 - b^2)*cosh(x)^4 + 4*(a^2 - b^2)*cosh(x)*sinh(x)^3 + (a^2 - b
^2)*sinh(x)^4 + 2*(a^2 - b^2)*cosh(x)^2 + 2*(3*(a^2 - b^2)*cosh(x)^2 + a^2
- b^2)*sinh(x)^2 + a^2 - b^2 + 4*((a^2 - b^2)*cosh(x)^3 + (a^2 - b^2)*cos
h(x))*sinh(x))*log(2*cosh(x)/(cosh(x) - sinh(x))))/(b^3*cosh(x)^4 + 4*b^3*
cosh(x)*sinh(x)^3 + b^3*sinh(x)^4 + 2*b^3*cosh(x)^2 + b^3 + 2*(3*b^3*cosh(
x)^2 + b^3)*sinh(x)^2 + 4*(b^3*cosh(x)^3 + b^3*cosh(x))*sinh(x))
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx$$

input `integrate(sech(x)**4/(a+b*tanh(x)),x)`

output `Integral(sech(x)**4/(a + b*tanh(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 89 vs.  $2(38) = 76$ .

Time = 0.12 (sec) , antiderivative size = 89, normalized size of antiderivative = 2.22

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx = \frac{2((a+b)e^{-2x} + a)}{2b^2e^{-2x} + b^2e^{-4x} + b^2} - \frac{(a^2 - b^2) \log(-(a-b)e^{-2x} - a - b)}{b^3} + \frac{(a^2 - b^2) \log(e^{-2x} + 1)}{b^3}$$

input `integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

output `2*((a + b)*e^(-2*x) + a)/(2*b^2*e^(-2*x) + b^2*e^(-4*x) + b^2) - (a^2 - b^2)*log(-(a - b)*e^(-2*x) - a - b)/b^3 + (a^2 - b^2)*log(e^(-2*x) + 1)/b^3`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 104 vs.  $2(38) = 76$ .

Time = 0.13 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.60

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx = -\frac{(a^3 + a^2b - ab^2 - b^3) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{ab^3 + b^4} + \frac{(a^2 - b^2) \log(e^{(2x)} + 1)}{b^3} - \frac{2(ab + (ab - b^2)e^{(2x)})}{b^3(e^{(2x)} + 1)^2}$$

input `integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="giac")`

output `-(a^3 + a^2*b - a*b^2 - b^3)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a*b^3 + b^4) + (a^2 - b^2)*log(e^(2*x) + 1)/b^3 - 2*(a*b + (a*b - b^2)*e^(2*x))/(b^3*(e^(2*x) + 1)^2)`

**Mupad [B] (verification not implemented)**

Time = 2.36 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.20

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx = \frac{\ln(e^{2x} + 1) (a + b) (a - b)}{b^3} - \frac{2(a - b)}{b^2 (e^{2x} + 1)} - \frac{\ln(a - b + ae^{2x} + be^{2x}) (a + b) (a - b)}{b^3} - \frac{2}{b (2e^{2x} + e^{4x} + 1)}$$

input `int(1/(cosh(x)^4*(a + b*tanh(x))),x)`

output `(log(exp(2*x) + 1)*(a + b)*(a - b))/b^3 - (2*(a - b))/(b^2*(exp(2*x) + 1)) - (log(a - b + a*exp(2*x) + b*exp(2*x))*(a + b)*(a - b))/b^3 - 2/(b*(2*exp(2*x) + exp(4*x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 312, normalized size of antiderivative = 7.80

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx$$

$$= \frac{e^{4x} \log(e^{2x} + 1) a^2 - e^{4x} \log(e^{2x} + 1) b^2 - e^{4x} \log(e^{2x} a + e^{2x} b + a - b) a^2 + e^{4x} \log(e^{2x} a + e^{2x} b + a - b) b^2}{(b^2 - a^2) \cosh(2x)}$$

input `int(sech(x)^4/(a+b*tanh(x)),x)`output `(e**(4*x)*log(e**(2*x) + 1)*a**2 - e**(4*x)*log(e**(2*x) + 1)*b**2 - e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2 + e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**2 + e**(4*x)*a*b - e**(4*x)*b**2 + 2*e**(2*x)*log(e**(2*x) + 1)*a**2 - 2*e**(2*x)*log(e**(2*x) + 1)*b**2 - 2*e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2 + 2*e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**2 + log(e**(2*x) + 1)*a**2 - log(e**(2*x) + 1)*b**2 - log(e**(2*x)*a + e**(2*x)*b + a - b)*a**2 + log(e**(2*x)*a + e**(2*x)*b + a - b)*b**2 - a*b - b**2)/(b**3*(e**(4*x) + 2*e**(2*x) + 1))`

### 3.106 $\int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx$

Optimal result	829
Mathematica [A] (verified)	829
Rubi [A] (verified)	830
Maple [A] (verified)	831
Fricas [B] (verification not implemented)	831
Sympy [F]	832
Maxima [A] (verification not implemented)	832
Giac [B] (verification not implemented)	832
Mupad [B] (verification not implemented)	833
Reduce [B] (verification not implemented)	833

#### Optimal result

Integrand size = 13, antiderivative size = 11

$$\int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx = \frac{\log(a+b \tanh(x))}{b}$$

output `ln(a+b*tanh(x))/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx = \frac{\log(a+b \tanh(x))}{b}$$

input `Integrate[Sech[x]^2/(a + b*Tanh[x]), x]`

output `Log[a + b*Tanh[x]]/b`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {3042, 3987, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sec(ix)^2}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{3987} \\ & \frac{\int \frac{1}{a + b \tanh(x)} d(b \tanh(x))}{b} \\ & \quad \downarrow \text{16} \\ & \frac{\log(a + b \tanh(x))}{b} \end{aligned}$$

input `Int[Sech[x]^2/(a + b*Tanh[x]),x]`

output `Log[a + b*Tanh[x]]/b`

**Defintions of rubi rules used**

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3987

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_
), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1),
x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2,
0] && IntegerQ[m/2]
```

**Maple [A] (verified)**

Time = 3.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{\ln(a+b \tanh(x))}{b}$	12
default	$\frac{\ln(a+b \tanh(x))}{b}$	12
risch	$-\frac{\ln(e^{2x}+1)}{b} + \frac{\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b}$	35

input

```
int(sech(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
ln(a+b*tanh(x))/b
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(11) = 22.

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.82

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = \frac{\log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)}{b}$$

input

```
integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
(log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - log(2*cosh(x)/(cosh(
x) - sinh(x))))/b
```



**Sympy [F]**

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx$$

input `integrate(sech(x)**2/(a+b*tanh(x)),x)`

output `Integral(sech(x)**2/(a + b*tanh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = \frac{\log(b \tanh(x) + a)}{b}$$

input `integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="maxima")`

output `log(b*tanh(x) + a)/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(11) = 22.

Time = 0.13 (sec) , antiderivative size = 45, normalized size of antiderivative = 4.09

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = \frac{(a + b) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{ab + b^2} - \frac{\log(e^{(2x)} + 1)}{b}$$

input `integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="giac")`

output `(a + b)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a*b + b^2) - log(e^(2*x) + 1)/b`

**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 50, normalized size of antiderivative = 4.55

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = -\frac{2 \operatorname{atan}\left(\frac{a\sqrt{-b^2} + a e^{2x}\sqrt{-b^2} + b e^{2x}\sqrt{-b^2}}{b^2}\right)}{\sqrt{-b^2}}$$

input `int(1/(cosh(x)^2*(a + b*tanh(x))),x)`output `-(2*atan((a*(-b^2)^(1/2) + a*exp(2*x)*(-b^2)^(1/2) + b*exp(2*x)*(-b^2)^(1/2))/b^2))/(-b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 3.18

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx = \frac{-\log(e^{2x} + 1) + \log(e^{2x}a + e^{2x}b + a - b)}{b}$$

input `int(sech(x)^2/(a+b*tanh(x)),x)`output `( - log(e**(2*x) + 1) + log(e**(2*x)*a + e**(2*x)*b + a - b))/b`

### 3.107 $\int \frac{1}{a+b \tanh(x)} dx$

Optimal result	834
Mathematica [A] (verified)	834
Rubi [A] (verified)	835
Maple [A] (verified)	836
Fricas [A] (verification not implemented)	837
Sympy [B] (verification not implemented)	837
Maxima [A] (verification not implemented)	838
Giac [A] (verification not implemented)	838
Mupad [B] (verification not implemented)	838
Reduce [B] (verification not implemented)	839

#### Optimal result

Integrand size = 8, antiderivative size = 39

$$\int \frac{1}{a+b \tanh(x)} dx = \frac{ax}{a^2-b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2-b^2}$$

output

```
a*x/(a^2-b^2)-b*ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.26

$$\int \frac{1}{a+b \tanh(x)} dx = \frac{(-a+b) \log(1-\tanh(x)) + (a+b) \log(1+\tanh(x)) - 2b \log(a+b \tanh(x))}{2(a-b)(a+b)}$$

input

```
Integrate[(a + b*Tanh[x])^(-1), x]
```

output

```
((-a + b)*Log[1 - Tanh[x]] + (a + b)*Log[1 + Tanh[x]] - 2*b*Log[a + b*Tanh[x]])/(2*(a - b)*(a + b))
```

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{3965} \\
 & \frac{ax}{a^2 - b^2} - \frac{ib \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{4013} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}
 \end{aligned}$$

input `Int[(a + b*Tanh[x])^(-1),x]`

output `(a*x)/(a^2 - b^2) - (b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3965  $\text{Int}[(a + (b \cdot \tan[c + d \cdot x])^{-1}), x\_Symbol] \rightarrow \text{Simp}[a \cdot (x / (a^2 + b^2)), x] + \text{Simp}[b / (a^2 + b^2) \text{Int}[(b - a \cdot \tan[c + d \cdot x]) / (a + b \cdot \tan[c + d \cdot x]), x], x] /;$   $\text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4013  $\text{Int}[(c + (d \cdot \tan[e + f \cdot x]) / (a + (b \cdot \tan[e + f \cdot x]) \cdot (x))), x\_Symbol] \rightarrow \text{Simp}[(c / (b \cdot f)) \cdot \text{Log}[\text{RemoveContent}[a \cdot \text{Cos}[e + f \cdot x] + b \cdot \text{Sin}[e + f \cdot x], x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a \cdot c + b \cdot d, 0]$

## Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

method	result	size
parallelrisch	$-\frac{-\ln(1-\tanh(x))b+b\ln(a+b\tanh(x))-ax-bx}{a^2-b^2}$	42
derivativedivides	$-\frac{b\ln(a+b\tanh(x))}{(a-b)(a+b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
default	$-\frac{b\ln(a+b\tanh(x))}{(a-b)(a+b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
risch	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} - \frac{b\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2-b^2}$	55

input `int(1/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`output 
$$-(-\ln(1-\tanh(x))*b+b*\ln(a+b*\tanh(x))-a*x-b*x)/(a^2-b^2)$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{(a + b)x - b \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^2 - b^2}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="fricas")`

output `((a + b)*x - b*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))))/(a^2 - b^2)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. 2(29) = 58.

Time = 0.23 (sec) , antiderivative size = 146, normalized size of antiderivative = 3.74

$$\int \frac{1}{a + b \tanh(x)} dx = \begin{cases} \tilde{\infty}(x - \log(\tanh(x) + 1) + \log(\tanh(x))) & \text{for } a = 0 \wedge b = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ -\frac{x \tanh(x)}{2b \tanh(x) - 2b} + \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} - \frac{bx}{a^2 - b^2} - \frac{b \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} + \frac{b \log(\tanh(x) + 1)}{a^2 - b^2} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+b*tanh(x)),x)`

output `Piecewise((zoo*(x - log(tanh(x) + 1) + log(tanh(x))), Eq(a, 0) & Eq(b, 0)), (x/a, Eq(b, 0)), (-x*tanh(x)/(2*b*tanh(x) - 2*b) + x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) - b*x/(a**2 - b**2) - b*log(a/b + tanh(x))/(a**2 - b**2) + b*log(tanh(x) + 1)/(a**2 - b**2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{1}{a + b \tanh(x)} dx = -\frac{b \log(-(a-b)e^{(-2x)} - a - b)}{a^2 - b^2} + \frac{x}{a + b}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="maxima")`output `-b*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{1}{a + b \tanh(x)} dx = -\frac{b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} + \frac{x}{a - b}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="giac")`output `-b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2 - b^2) + x/(a - b)`**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{ax - b(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

input `int(1/(a + b*tanh(x)),x)`output `(a*x - b*(x - log(tanh(x) + 1) + log(a + b*tanh(x))))/(a^2 - b^2)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{-\log(e^{2x}a + e^{2x}b + a - b)b + ax + bx}{a^2 - b^2}$$

input `int(1/(a+b*tanh(x)),x)`

output `( - log(e**(2*x)*a + e**(2*x)*b + a - b)*b + a*x + b*x)/(a**2 - b**2)`



### 3.108 $\int \frac{\cosh^2(x)}{a+b \tanh(x)} dx$

Optimal result	840
Mathematica [A] (verified)	840
Rubi [A] (verified)	841
Maple [A] (verified)	843
Fricas [B] (verification not implemented)	843
Sympy [F]	844
Maxima [A] (verification not implemented)	844
Giac [A] (verification not implemented)	845
Mupad [B] (verification not implemented)	845
Reduce [B] (verification not implemented)	846

#### Optimal result

Integrand size = 13, antiderivative size = 99

$$\int \frac{\cosh^2(x)}{a+b \tanh(x)} dx = -\frac{(a+2b) \log(1-\tanh(x))}{4(a+b)^2} + \frac{(a-2b) \log(1+\tanh(x))}{4(a-b)^2} + \frac{b^3 \log(a+b \tanh(x))}{(a^2-b^2)^2} + \frac{1}{4(a+b)(1-\tanh(x))} - \frac{1}{4(a-b)(1+\tanh(x))}$$

output

$$-1/4*(a+2*b)*\ln(1-\tanh(x))/(a+b)^2+1/4*(a-2*b)*\ln(1+\tanh(x))/(a-b)^2+b^3*\ln(a+b*\tanh(x))/(a^2-b^2)^2+1/4/(a+b)/(1-\tanh(x))-1/4/(a-b)/(1+\tanh(x))$$

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.76

$$\int \frac{\cosh^2(x)}{a+b \tanh(x)} dx = \frac{2a^3x - 6ab^2x + (-a^2b + b^3) \cosh(2x) + 4b^3 \log(a \cosh(x) + b \sinh(x)) + a(a^2 - b^2) \sinh(2x)}{4(a-b)^2(a+b)^2}$$

input `Integrate[Cosh[x]^2/(a + b*Tanh[x]), x]`

output  $(2a^3x - 6ab^2x + (-(a^2b) + b^3)*\text{Cosh}[2x] + 4b^3*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]] + a*(a^2 - b^2)*\text{Sinh}[2x])/(4*(a - b)^2*(a + b)^2)$

### Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3987, 27, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx$$

↓ 3042

$$\int \frac{1}{\sec(ix)^2(a - ib \tan(ix))} dx$$

↓ 3987

$$\int \frac{b^4}{(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))^2} d(b \tanh(x))$$

↓ 27

$$b^3 \int \frac{1}{(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))^2} d(b \tanh(x))$$

↓ 477

$$\int \left( \frac{b^4}{(a^2 - b^2)^2(a + b \tanh(x))} + \frac{b^2}{4(a+b)(b-b \tanh(x))^2} + \frac{b^2}{4(a-b)(\tanh(x)b+b)^2} + \frac{(a+2b)b}{4(a+b)^2(b-b \tanh(x))} + \frac{(a-2b)b}{4(a-b)^2(\tanh(x)b+b)} \right) d(b \tanh(x))$$

↓ 2009

$$\frac{b^4 \log(a + b \tanh(x))}{(a^2 - b^2)^2} + \frac{b^2}{4(a+b)(b-b \tanh(x))} - \frac{b^2}{4(a-b)(b \tanh(x) + b)} - \frac{b(a+2b) \log(b-b \tanh(x))}{4(a+b)^2} + \frac{b(a-2b) \log(b \tanh(x) + b)}{4(a-b)^2}$$

input `Int[Cosh[x]^2/(a + b*Tanh[x]),x]`

output `(-1/4*(b*(a + 2*b)*Log[b - b*Tanh[x]])/(a + b)^2 + (b^4*Log[a + b*Tanh[x]]  
)/(a^2 - b^2)^2 + ((a - 2*b)*b*Log[b + b*Tanh[x]])/(4*(a - b)^2) + b^2/(4*  
(a + b)*(b - b*Tanh[x])) - b^2/(4*(a - b)*(b + b*Tanh[x]))) / b`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma  
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 477 `Int[((c_) + (d_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[  
a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2  
]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] &  
& NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3987 `Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_  
) , x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1),  
x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2,  
0] && IntegerQ[m/2]`

**Maple [A] (verified)**

Time = 1.23 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.05

method	result
risch	$\frac{ax}{2(a+b)^2} + \frac{xb}{(a+b)^2} + \frac{e^{2x}}{8a+8b} - \frac{e^{-2x}}{8(a-b)} - \frac{2b^3x}{a^4-2a^2b^2+b^4} + \frac{b^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^4-2a^2b^2+b^4}$
default	$\frac{b^3 \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{(a-b)^2(a+b)^2} + \frac{1}{(2b+2a)\left(\tanh\left(\frac{x}{2}\right)-1\right)^2} + \frac{2}{(4a+4b)\left(\tanh\left(\frac{x}{2}\right)-1\right)} + \frac{(-a-2b) \ln\left(\tanh\left(\frac{x}{2}\right)-1\right)}{2(a+b)^2} - \frac{1}{(2a-b)}$

input `int(cosh(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`output  $\frac{1}{2}ax/(a+b)^2 + x/(a+b)^2 + b + 1/8/(a+b) \exp(2x) - 1/8/(a-b) \exp(-2x) - 2b^3/(a^4 - 2a^2b^2 + b^4) * x + b^3/(a^4 - 2a^2b^2 + b^4) * \ln(\exp(2x) + (a-b)/(a+b))$ **Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 331 vs. 2(89) = 178.

Time = 0.11 (sec) , antiderivative size = 331, normalized size of antiderivative = 3.34

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{(a^3 - a^2b - ab^2 + b^3) \cosh(x)^4 + 4(a^3 - a^2b - ab^2 + b^3) \cosh(x) \sinh(x)^3 + (a^3 - a^2b - ab^2 + b^3) \sinh(x)^4}{(a^3 - a^2b - ab^2 + b^3)}$$

input `integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`

output

```
1/8*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 - a^2*b - a*b^2 + b^3)
*cosh(x)*sinh(x)^3 + (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^4 + 4*(a^3 - 3*a*
b^2 - 2*b^3)*x*cosh(x)^2 - a^3 - a^2*b + a*b^2 + b^3 + 2*(3*(a^3 - a^2*b -
a*b^2 + b^3)*cosh(x)^2 + 2*(a^3 - 3*a*b^2 - 2*b^3)*x)*sinh(x)^2 + 8*(b^3*
cosh(x)^2 + 2*b^3*cosh(x)*sinh(x) + b^3*sinh(x)^2)*log(2*(a*cosh(x) + b*si
nh(x))/(cosh(x) - sinh(x))) + 4*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^3 + 2
*(a^3 - 3*a*b^2 - 2*b^3)*x*cosh(x))*sinh(x)/((a^4 - 2*a^2*b^2 + b^4)*cosh
(x)^2 + 2*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x) + (a^4 - 2*a^2*b^2 + b^4
)*sinh(x)^2)
```

**Sympy [F]**

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx = \int \frac{\cosh^2(x)}{a + b \tanh(x)} dx$$

input

```
integrate(cosh(x)**2/(a+b*tanh(x)),x)
```

output

```
Integral(cosh(x)**2/(a + b*tanh(x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.87

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx = \frac{b^3 \log(-(a-b)e^{(-2x)} - a - b)}{a^4 - 2a^2b^2 + b^4} + \frac{(a+2b)x}{2(a^2 + 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)} - \frac{e^{(-2x)}}{8(a-b)}$$

input

```
integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")
```

output

```
b^3*log(-(a - b)*e^(-2*x) - a - b)/(a^4 - 2*a^2*b^2 + b^4) + 1/2*(a + 2*b)
*x/(a^2 + 2*a*b + b^2) + 1/8*e^(2*x)/(a + b) - 1/8*e^(-2*x)/(a - b)
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.12

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx = \frac{b^3 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^4 - 2a^2b^2 + b^4} + \frac{(a - 2b)x}{2(a^2 - 2ab + b^2)} - \frac{(2ae^{(2x)} - 4be^{(2x)} + a - b)e^{(-2x)}}{8(a^2 - 2ab + b^2)} + \frac{e^{(2x)}}{8(a + b)}$$

input `integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="giac")`

output `b^3*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^4 - 2*a^2*b^2 + b^4) + 1/2*(a - 2*b)*x/(a^2 - 2*a*b + b^2) - 1/8*(2*a*e^(2*x) - 4*b*e^(2*x) + a - b)*e^(-2*x)/(a^2 - 2*a*b + b^2) + 1/8*e^(2*x)/(a + b)`

**Mupad [B] (verification not implemented)**

Time = 2.51 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.85

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx = \frac{e^{2x}}{8a + 8b} - \frac{e^{-2x}}{8a - 8b} + \frac{b^3 \ln(a - b + ae^{2x} + be^{2x})}{a^4 - 2a^2b^2 + b^4} + \frac{x(a - 2b)}{2(a - b)^2}$$

input `int(cosh(x)^2/(a + b*tanh(x)),x)`

output `exp(2*x)/(8*a + 8*b) - exp(-2*x)/(8*a - 8*b) + (b^3*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^4 + b^4 - 2*a^2*b^2) + (x*(a - 2*b))/(2*(a - b)^2)`

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.52

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx$$

$$= \frac{e^{4x}a^3 - e^{4x}a^2b - e^{4x}ab^2 + e^{4x}b^3 + 8e^{2x}\log(e^{2x}a + e^{2x}b + a - b)b^3 + 4e^{2x}a^3x - 12e^{2x}ab^2x - 8e^{2x}b^3x - a^3 - a^2b + ab^2 + b^3}{8e^{2x}(a^4 - 2a^2b^2 + b^4)}$$

input `int(cosh(x)^2/(a+b*tanh(x)),x)`output `(e**(4*x)*a**3 - e**(4*x)*a**2*b - e**(4*x)*a*b**2 + e**(4*x)*b**3 + 8*e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**3 + 4*e**(2*x)*a**3*x - 12*e**(2*x)*a*b**2*x - 8*e**(2*x)*b**3*x - a**3 - a**2*b + a*b**2 + b**3)/(8*e**(2*x)*(a**4 - 2*a**2*b**2 + b**4))`

### 3.109 $\int \frac{\cosh^4(x)}{a+b \tanh(x)} dx$

Optimal result	847
Mathematica [A] (verified)	848
Rubi [A] (verified)	848
Maple [A] (verified)	850
Fricas [B] (verification not implemented)	850
Sympy [F]	851
Maxima [A] (verification not implemented)	852
Giac [A] (verification not implemented)	852
Mupad [B] (verification not implemented)	853
Reduce [B] (verification not implemented)	853

#### Optimal result

Integrand size = 13, antiderivative size = 168

$$\int \frac{\cosh^4(x)}{a+b \tanh(x)} dx = -\frac{(3a^2 + 9ab + 8b^2) \log(1 - \tanh(x))}{16(a+b)^3} + \frac{(3a^2 - 9ab + 8b^2) \log(1 + \tanh(x))}{16(a-b)^3} - \frac{b^5 \log(a + b \tanh(x))}{(a^2 - b^2)^3} + \frac{1}{16(a+b)(1 - \tanh(x))^2} + \frac{3a + 5b}{16(a+b)^2(1 - \tanh(x))} - \frac{1}{16(a-b)(1 + \tanh(x))^2} - \frac{3a - 5b}{16(a-b)^2(1 + \tanh(x))}$$

output

```
-1/16*(3*a^2+9*a*b+8*b^2)*ln(1-tanh(x))/(a+b)^3+1/16*(3*a^2-9*a*b+8*b^2)*ln(1+tanh(x))/(a-b)^3-b^5*ln(a+b*tanh(x))/(a^2-b^2)^3+1/16/(a+b)/(1-tanh(x))^2+1/16*(3*a+5*b)/(a+b)^2/(1-tanh(x))-1/16/(a-b)/(1+tanh(x))^2-1/16*(3*a-5*b)/(a-b)^2/(1+tanh(x))
```



**Mathematica [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.23

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx$$

$$= \frac{8b^3(a^2 - b^2) \cosh^2(x) - 4b(a^2 - b^2)^2 \cosh^4(x) - 3a^5 \log(1 - \tanh(x)) + 10a^3b^2 \log(1 - \tanh(x)) - 15ab^4 \log(1 + \tanh(x)) + 8b^5 \log(1 + \tanh(x)) - 3a^5 \log(1 + \tanh(x)) - 10a^3b^2 \log(1 + \tanh(x)) + 15a^2b^4 \log(1 + \tanh(x)) + 8b^5 \log(1 + \tanh(x)) - 16b^5 \log(a + b \tanh(x)) + 4a(a^2 - b^2)^2 \cosh(x)^3 \sinh(x) + a(3a^4 - 10a^2b^2 + 7b^4) \sinh(2x)}{(16(a - b)^3(a + b)^3)}$$

input

```
Integrate[Cosh[x]^4/(a + b*Tanh[x]), x]
```

output

```
(8*b^3*(a^2 - b^2)*Cosh[x]^2 - 4*b*(a^2 - b^2)^2*Cosh[x]^4 - 3*a^5*Log[1 - Tanh[x]] + 10*a^3*b^2*Log[1 - Tanh[x]] - 15*a*b^4*Log[1 - Tanh[x]] + 8*b^5*Log[1 - Tanh[x]] + 3*a^5*Log[1 + Tanh[x]] - 10*a^3*b^2*Log[1 + Tanh[x]] + 15*a*b^4*Log[1 + Tanh[x]] + 8*b^5*Log[1 + Tanh[x]] - 16*b^5*Log[a + b*Tanh[x]] + 4*a*(a^2 - b^2)^2*Cosh[x]^3*Sinh[x] + a*(3*a^4 - 10*a^2*b^2 + 7*b^4)*Sinh[2*x])/(16*(a - b)^3*(a + b)^3)
```

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3042, 3987, 27, 477, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{\sec(ix)^4(a - ib \tan(ix))} dx$$

$$\downarrow \text{3987}$$

$$\int \frac{b^6}{(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))^3} d(b \tanh(x))$$

$$\frac{\int \frac{b^6}{(a + b \tanh(x))(b^2 - b^2 \tanh^2(x))^3} d(b \tanh(x))}{b}$$

$$\begin{array}{c}
 \downarrow 27 \\
 b^5 \int \frac{1}{(a + b \tanh(x)) (b^2 - b^2 \tanh^2(x))^3} d(b \tanh(x)) \\
 \downarrow 477 \\
 \int \left( -\frac{b^6}{(a^2 - b^2)^3 (a + b \tanh(x))} + \frac{b^3}{8(a+b)(b-b \tanh(x))^3} + \frac{b^3}{8(a-b)(\tanh(x)b+b)^3} + \frac{(3a+5b)b^2}{16(a+b)^2(b-b \tanh(x))^2} + \frac{(3a-5b)b^2}{16(a-b)^2(\tanh(x)b+b)^2} + \right. \\
 \left. \frac{b^3}{16(a+b)(b-b \tanh(x))^2} - \frac{b^3}{16(a-b)(b+b \tanh(x))^2} \right) dx \\
 \downarrow 2009 \\
 -\frac{b(3a^2+9ab+8b^2) \log(b-b \tanh(x))}{16(a+b)^3} + \frac{b(3a^2-9ab+8b^2) \log(b \tanh(x)+b)}{16(a-b)^3} - \frac{b^6 \log(a+b \tanh(x))}{(a^2-b^2)^3} + \frac{b^3}{16(a+b)(b-b \tanh(x))^2} - \frac{b^3}{16(a-b)(b+b \tanh(x))^2}
 \end{array}$$

input `Int[Cosh[x]^4/(a + b*Tanh[x]), x]`

output 
$$\begin{aligned}
 & (-1/16*(b*(3*a^2 + 9*a*b + 8*b^2)*\text{Log}[b - b*\text{Tanh}[x]])/(a + b)^3 - (b^6*\text{Log}[a + b*\text{Tanh}[x]])/(a^2 - b^2)^3 + (b*(3*a^2 - 9*a*b + 8*b^2)*\text{Log}[b + b*\text{Tanh}[x]])/(16*(a - b)^3) + b^3/(16*(a + b)*(b - b*\text{Tanh}[x])^2) + (b^2*(3*a + 5*b))/(16*(a + b)^2*(b - b*\text{Tanh}[x])) - b^3/(16*(a - b)*(b + b*\text{Tanh}[x])^2) - ((3*a - 5*b)*b^2)/(16*(a - b)^2*(b + b*\text{Tanh}[x])))/b
 \end{aligned}$$

### Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 477 `Int[((c_) + (d_.)*(x_))^(n_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p Int[ExpandIntegrand[(c + d*x)^n*(1 - Rt[-b/a, 2]*x)^p*(1 + Rt[-b/a, 2]*x)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IntegerQ[n] && NiceSqrtQ[-b/a] && !FractionalPowerFactorQ[Rt[-b/a, 2]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3987 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/(b*f) Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

### Maple [A] (verified)

Time = 11.93 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.14

method	result
risch	$\frac{3a^2x}{8(a+b)^3} + \frac{9axb}{8(a+b)^3} + \frac{xb^2}{(a+b)^3} + \frac{e^{4x}}{64a+64b} + \frac{e^{2x}a}{8(a+b)^2} + \frac{3e^{2x}b}{16(a+b)^2} - \frac{e^{-2x}a}{8(a-b)^2} + \frac{3e^{-2x}b}{16(a-b)^2} - \frac{e^{-4x}}{64(a-b)} + \frac{2b^5}{a^6-3a^4b^2+}$
default	$-\frac{b^5 \ln\left(\tanh\left(\frac{x}{2}\right)^2 a + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{(a+b)^3(a-b)^3} + \frac{1}{2(2b+2a)(\tanh\left(\frac{x}{2}\right)-1)^4} + \frac{2}{(4a+4b)(\tanh\left(\frac{x}{2}\right)-1)^3} - \frac{-7a-9b}{8(a+b)^2(\tanh\left(\frac{x}{2}\right)-1)^2} - \frac{1}{8(a+b)}$

input `int(cosh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output  $\frac{3}{8}a^2x/(a+b)^3 + 9/8ax/(a+b)^3 + bx/(a+b)^3 + 1/64/(a+b) \exp(4x) + 1/8/(a+b)^2 \exp(2x)a + 3/16/(a+b)^2 \exp(2x)b - 1/8/(a-b)^2 \exp(-2x)a + 3/16/(a-b)^2 \exp(-2x)b - 1/64/(a-b) \exp(-4x) + 2b^5/(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) * x - b^5/(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) * \ln(\exp(2x) + (a-b)/(a+b))$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1281 vs.  $2(152) = 304$ .

Time = 0.11 (sec) , antiderivative size = 1281, normalized size of antiderivative = 7.62

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

output

```

1/64*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^8 + 8*(a
^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^7 + (a^5
- a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^8 + 4*(2*a^5 - a^4
*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^6 + 4*(2*a^5 - a^4*b
- 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5 + 7*(a^5 - a^4*b - 2*a^3*b^2 +
2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^2)*sinh(x)^6 + 8*(3*a^5 - 10*a^3*b^2 + 15
*a*b^4 + 8*b^5)*x*cosh(x)^4 + 8*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 +
a*b^4 - b^5)*cosh(x)^3 + 3*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^
4 - 3*b^5)*cosh(x))*sinh(x)^5 - a^5 - a^4*b + 2*a^3*b^2 + 2*a^2*b^3 - a*b^
4 - b^5 + 2*(35*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)
)^4 + 30*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)
^2 + 4*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x)*sinh(x)^4 + 8*(7*(a^5 -
a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^5 + 10*(2*a^5 - a^4*b
- 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^3 + 4*(3*a^5 - 10*a^3*
b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x))*sinh(x)^3 - 4*(2*a^5 + a^4*b - 6*a^3*b^
2 - 4*a^2*b^3 + 4*a*b^4 + 3*b^5)*cosh(x)^2 + 4*(7*(a^5 - a^4*b - 2*a^3*b^2
+ 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 - 2*a^5 - a^4*b + 6*a^3*b^2 + 4*a^2*
b^3 - 4*a*b^4 - 3*b^5 + 15*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^
4 - 3*b^5)*cosh(x)^4 + 12*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x)
)^2)*sinh(x)^2 - 64*(b^5*cosh(x)^4 + 4*b^5*cosh(x)^3*sinh(x) + 6*b^5*co...

```

## Sympy [F]

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = \int \frac{\cosh^4(x)}{a + b \tanh(x)} dx$$

input

```
integrate(cosh(x)**4/(a+b*tanh(x)), x)
```

output

```
Integral(cosh(x)**4/(a + b*tanh(x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.98

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = -\frac{b^5 \log(-(a-b)e^{-2x} - a - b)}{a^6 - 3a^4b^2 + 3a^2b^4 - b^6} + \frac{(3a^2 + 9ab + 8b^2)x}{8(a^3 + 3a^2b + 3ab^2 + b^3)} + \frac{(4(2a + 3b)e^{-2x} + a + b)e^{4x}}{64(a^2 + 2ab + b^2)} - \frac{4(2a - 3b)e^{-2x} + (a - b)e^{-4x}}{64(a^2 - 2ab + b^2)}$$

input `integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

output `-b^5*log(-(a - b)*e^(-2*x) - a - b)/(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) + 1/8*(3*a^2 + 9*a*b + 8*b^2)*x/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) + 1/64*(4*(2*a + 3*b)*e^(-2*x) + a + b)*e^(4*x)/(a^2 + 2*a*b + b^2) - 1/64*(4*(2*a - 3*b)*e^(-2*x) + (a - b)*e^(-4*x))/(a^2 - 2*a*b + b^2)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.35

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = -\frac{b^5 \log(|ae^{2x} + be^{2x} + a - b|)}{a^6 - 3a^4b^2 + 3a^2b^4 - b^6} + \frac{(3a^2 - 9ab + 8b^2)x}{8(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(18a^2e^{4x} - 54abe^{4x} + 48b^2e^{4x} + 8a^2e^{2x} - 20abe^{2x} + 12b^2e^{2x} + a^2 - 2ab + b^2)e^{-4x}}{64(a^3 - 3a^2b + 3ab^2 - b^3)} + \frac{ae^{4x} + be^{4x} + 8ae^{2x} + 12be^{2x}}{64(a^2 + 2ab + b^2)}$$

input `integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="giac")`

output

```
-b^5*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) + 1/8*(3*a^2 - 9*a*b + 8*b^2)*x/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) - 1/64*(18*a^2*e^(4*x) - 54*a*b*e^(4*x) + 48*b^2*e^(4*x) + 8*a^2*e^(2*x) - 20*a*b*e^(2*x) + 12*b^2*e^(2*x) + a^2 - 2*a*b + b^2)*e^(-4*x)/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) + 1/64*(a*e^(4*x) + b*e^(4*x) + 8*a*e^(2*x) + 12*b*e^(2*x))/(a^2 + 2*a*b + b^2)
```

**Mupad [B] (verification not implemented)**

Time = 2.62 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.85

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = \frac{e^{4x}}{64a + 64b} - \frac{e^{-4x}}{64a - 64b} - \frac{e^{-2x}(2a - 3b)}{16(a - b)^2} - \frac{b^5 \ln(a - b + a e^{2x} + b e^{2x})}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{x(3a^2 - 9ab + 8b^2)}{8(a - b)^3} + \frac{e^{2x}(2a + 3b)}{16(a + b)^2}$$

input

```
int(cosh(x)^4/(a + b*tanh(x)),x)
```

output

```
exp(4*x)/(64*a + 64*b) - exp(-4*x)/(64*a - 64*b) - (exp(-2*x)*(2*a - 3*b))/(16*(a - b)^2) - (b^5*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2) + (x*(3*a^2 - 9*a*b + 8*b^2))/(8*(a - b)^3) + (exp(2*x)*(2*a + 3*b))/(16*(a + b)^2)
```

**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 355, normalized size of antiderivative = 2.11

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx = \frac{2e^{8x}a^2b^3 - 64e^{4x}\log(e^{2x}a + e^{2x}b + a - b)b^5 - e^{8x}b^5 - 4e^{2x}a^4b + 16e^{2x}a^2b^3 - e^{8x}a^4b - 80e^{4x}a^3b^2x + 120}{16(a + b)^2}$$

input

```
int(cosh(x)^4/(a+b*tanh(x)),x)
```

output

```
(e**(8*x)*a**5 - e**(8*x)*a**4*b - 2*e**(8*x)*a**3*b**2 + 2*e**(8*x)*a**2*
b**3 + e**(8*x)*a*b**4 - e**(8*x)*b**5 + 8*e**(6*x)*a**5 - 4*e**(6*x)*a**4
*b - 24*e**(6*x)*a**3*b**2 + 16*e**(6*x)*a**2*b**3 + 16*e**(6*x)*a*b**4 -
12*e**(6*x)*b**5 - 64*e**(4*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**5 +
24*e**(4*x)*a**5*x - 80*e**(4*x)*a**3*b**2*x + 120*e**(4*x)*a*b**4*x + 64
*e**(4*x)*b**5*x - 8*e**(2*x)*a**5 - 4*e**(2*x)*a**4*b + 24*e**(2*x)*a**3*
b**2 + 16*e**(2*x)*a**2*b**3 - 16*e**(2*x)*a*b**4 - 12*e**(2*x)*b**5 - a**
5 - a**4*b + 2*a**3*b**2 + 2*a**2*b**3 - a*b**4 - b**5)/(64*e**(4*x)*(a**6
- 3*a**4*b**2 + 3*a**2*b**4 - b**6))
```

### 3.110 $\int \frac{\operatorname{sech}^7(x)}{a+b \tanh(x)} dx$

Optimal result	855
Mathematica [A] (verified)	856
Rubi [A] (verified)	856
Maple [B] (verified)	862
Fricas [B] (verification not implemented)	862
Sympy [F]	863
Maxima [F(-2)]	863
Giac [B] (verification not implemented)	863
Mupad [B] (verification not implemented)	864
Reduce [B] (verification not implemented)	865

#### Optimal result

Integrand size = 13, antiderivative size = 157

$$\int \frac{\operatorname{sech}^7(x)}{a+b \tanh(x)} dx = \frac{a(8a^4 - 20a^2b^2 + 15b^4) \arctan(\sinh(x))}{8b^6} - \frac{(a^2 - b^2)^{5/2} \arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{b^6} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)}{b^5} - \frac{(a^2 - b^2) \operatorname{sech}^3(x)}{3b^3} + \frac{\operatorname{sech}^5(x)}{5b} - \frac{a(4a^2 - 7b^2) \operatorname{sech}(x) \tanh(x)}{8b^4} + \frac{a \operatorname{sech}^3(x) \tanh(x)}{4b^2}$$

output

```
1/8*a*(8*a^4-20*a^2*b^2+15*b^4)*arctan(sinh(x))/b^6-(a^2-b^2)^(5/2)*arctan
(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/b^6+(a^2-b^2)^2*sech(x)/b^5-1/3*(a
^2-b^2)*sech(x)^3/b^3+1/5*sech(x)^5/b-1/8*a*(4*a^2-7*b^2)*sech(x)*tanh(x)/
b^4+1/4*a*sech(x)^3*tanh(x)/b^2
```



**Mathematica [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx$$

$$= \frac{30 \left( a(8a^4 - 20a^2b^2 + 15b^4) \arctan \left( \tanh \left( \frac{x}{2} \right) \right) - 8\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)^2 \arctan \left( \frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b}\sqrt{a+b}} \right) \right) + 24b^5 \operatorname{sech}^5(x) + 10b^3 \operatorname{sech}^3(x) (-4a^2 + 4b^2 + 3ab \tanh(x)) + 15b \operatorname{sech}(x) (8(a^2 - b^2)^2 + (-4a^3b + 7ab^3) \tanh(x))}{120b^6}$$

input

Integrate[Sech[x]^7/(a + b\*Tanh[x]), x]

output

```
(30*(a*(8*a^4 - 20*a^2*b^2 + 15*b^4)*ArcTan[Tanh[x/2]] - 8*Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)^2*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])]) + 24*b^5*Sech[x]^5 + 10*b^3*Sech[x]^3*(-4*a^2 + 4*b^2 + 3*a*b*Tanh[x]) + 15*b*Sech[x]*(8*(a^2 - b^2)^2 + (-4*a^3*b + 7*a*b^3)*Tanh[x]))/(120*b^6)
```

**Rubi [A] (verified)**Time = 1.53 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 20,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.538$ , Rules used = {3042, 3989, 3042, 3967, 3042, 3989, 3042, 3967, 3042, 3989, 3042, 3967, 3042, 3988, 219, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sec(ix)^7}{a - ib \tan(ix)} dx$$

$$\downarrow \text{3989}$$

$$\frac{\int \operatorname{sech}^5(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx}{b^2}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{\int \sec(ix)^5(a + ib \tan(ix))dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^5}{a-ib \tan(ix)} dx}{b^2} \\
 & \downarrow 3967 \\
 & \frac{a \int \operatorname{sech}^5(x)dx + \frac{1}{5}b \operatorname{sech}^5(x)}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^5}{a-ib \tan(ix)} dx}{b^2} \\
 & \downarrow 3042 \\
 & \frac{\frac{1}{5}b \operatorname{sech}^5(x) + a \int \csc(ix + \frac{\pi}{2})^5 dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^5}{a-ib \tan(ix)} dx}{b^2} \\
 & \downarrow 3989 \\
 & - \frac{(a^2 - b^2) \left( \frac{\int \operatorname{sech}^3(x)(a-b \tanh(x))dx}{b^2} - \frac{(a^2-b^2) \int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx}{b^2} \right)}{b^2} + \\
 & \frac{\frac{1}{5}b \operatorname{sech}^5(x) + a \int \csc(ix + \frac{\pi}{2})^5 dx}{b^2} \\
 & \downarrow 3042 \\
 & \frac{\frac{1}{5}b \operatorname{sech}^5(x) + a \int \csc(ix + \frac{\pi}{2})^5 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{\int \sec(ix)^3(a+ib \tan(ix))dx}{b^2} - \frac{(a^2-b^2) \int \frac{\sec(ix)^3}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
 & \downarrow 3967 \\
 & \frac{\frac{1}{5}b \operatorname{sech}^5(x) + a \int \csc(ix + \frac{\pi}{2})^5 dx}{b^2} - \\
 & \frac{(a^2 - b^2) \left( \frac{a \int \operatorname{sech}^3(x)dx + \frac{1}{3}b \operatorname{sech}^3(x)}{b^2} - \frac{(a^2-b^2) \int \frac{\sec(ix)^3}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
 & \downarrow 3042 \\
 & \frac{\frac{1}{5}b \operatorname{sech}^5(x) + a \int \csc(ix + \frac{\pi}{2})^5 dx}{b^2} - \\
 & \frac{(a^2 - b^2) \left( \frac{\frac{1}{3}b \operatorname{sech}^3(x) + a \int \csc(ix + \frac{\pi}{2})^3 dx}{b^2} - \frac{(a^2-b^2) \int \frac{\sec(ix)^3}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
 & \downarrow 3989
 \end{aligned}$$

$$\frac{\frac{1}{5}b\operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - (a^2 - b^2) \left( \frac{(a^2 - b^2) \left( \frac{\int \operatorname{sech}(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{b^2} \right)}{b^2} + \frac{\frac{1}{3}b\operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} \right)$$

---

$b^2$

↓ 3042

$$\frac{\frac{1}{5}b\operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - (a^2 - b^2) \left( \frac{\frac{1}{3}b\operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{\int \sec(ix)(a + ib \tan(ix)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2} \right)}{b^2} \right)$$

---

$b^2$

↓ 3967

$$\frac{\frac{1}{5}b\operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - (a^2 - b^2) \left( \frac{\frac{1}{3}b\operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{a \int \operatorname{sech}(x) dx + b \operatorname{sech}(x)}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2} \right)}{b^2} \right)$$

---

$b^2$

↓ 3042

$$\frac{\frac{1}{5}b\operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - (a^2 - b^2) \left( \frac{\frac{1}{3}b\operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{b\operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2} \right)}{b^2} \right)$$

---

$b^2$

↓ 3988

$$(a^2 - b^2) \left( \frac{\frac{1}{5} b \operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} - \frac{i(a^2 - b^2) \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b + a \tanh(x))^2} d(-i \cosh(x))}{b^2} \right)}{b^2} \right)$$


---

↓ 219

$$(a^2 - b^2) \left( \frac{\frac{1}{5} b \operatorname{sech}^5(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^5 dx}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2} \right)$$


---

↓ 4255

$$(a^2 - b^2) \left( \frac{a \left( \frac{3}{4} \int \operatorname{sech}^3(x) dx + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x) \right) + \frac{1}{5} b \operatorname{sech}^5(x)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2} \right)$$


---

↓ 3042

$$(a^2 - b^2) \left( \frac{\frac{1}{5} b \operatorname{sech}^5(x) + a \left( \frac{1}{4} \tanh(x) \operatorname{sech}^3(x) + \frac{3}{4} \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx \right)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2} \right)$$


---

↓ 4255

$$\begin{aligned}
 & \frac{a\left(\frac{3}{4}\left(\frac{\int \operatorname{sech}(x) dx}{2} + \frac{1}{2} \tanh(x) \operatorname{sech}(x)\right) + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x)\right) + \frac{1}{5} b \operatorname{sech}^5(x)}{b^2} \\
 (a^2 - b^2) & \left( \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a\left(\frac{1}{2} \tanh(x) \operatorname{sech}(x) + \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx\right)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc}{b^2} \right)}{b^2} \right) \\
 & \hspace{10em} \downarrow \text{3042} \\
 & \frac{\frac{1}{5} b \operatorname{sech}^5(x) + a\left(\frac{1}{4} \tanh(x) \operatorname{sech}^3(x) + \frac{3}{4}\left(\frac{1}{2} \tanh(x) \operatorname{sech}(x) + \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx\right)\right)}{b^2} \\
 (a^2 - b^2) & \left( \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a\left(\frac{1}{2} \tanh(x) \operatorname{sech}(x) + \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx\right)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc}{b^2} \right)}{b^2} \right) \\
 & \hspace{10em} \downarrow \text{4257} \\
 & \frac{a\left(\frac{3}{4}\left(\frac{1}{2} \arctan(\sinh(x)) + \frac{1}{2} \tanh(x) \operatorname{sech}(x)\right) + \frac{1}{4} \tanh(x) \operatorname{sech}^3(x)\right) + \frac{1}{5} b \operatorname{sech}^5(x)}{b^2} \\
 (a^2 - b^2) & \left( \frac{a\left(\frac{1}{2} \arctan(\sinh(x)) + \frac{1}{2} \tanh(x) \operatorname{sech}(x)\right) + \frac{1}{3} b \operatorname{sech}^3(x)}{b^2} - \frac{(a^2 - b^2) \left( \frac{a \arctan(\sinh(x)) + b \operatorname{sech}(x)}{b^2} - \frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} \right)}{b^2} \right)
 \end{aligned}$$

input `Int [Sech [x]^7/(a + b*Tanh [x]), x]`

output `((b*Sech [x]^5)/5 + a*((Sech [x]^3*Tanh [x])/4 + (3*(ArcTan [Sinh [x]]/2 + (Sech [x]*Tanh [x])/2))/4))/b^2 - ((a^2 - b^2)*(-((a^2 - b^2)*(-(Sqrt [a^2 - b^2]*ArcTan [(Cosh [x]*(b + a*Tanh [x]))/Sqrt [a^2 - b^2]]))/b^2) + (a*ArcTan [Sinh [x]] + b*Sech [x])/b^2))/b^2 + ((b*Sech [x]^3)/3 + a*(ArcTan [Sinh [x]]/2 + (Sech [x]*Tanh [x])/2))/b^2))/b^2`

## Defintions of rubi rules used

- rule 219  $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3967  $\text{Int}[(d_ \cdot \sec[(e_ + (f_ \cdot x)])^{(m_)} \cdot ((a_ + (b_ \cdot \tan[(e_ + (f_ \cdot x)])) \cdot x))], x\_Symbol] \rightarrow \text{Simp}[b \cdot ((d \cdot \text{Sec}[e + f \cdot x])^m / (f \cdot m)), x] + \text{Simp}[a \ \text{Int}[(d \cdot \text{Sec}[e + f \cdot x])^m, x], x] /; \text{FreeQ}\{a, b, d, e, f, m\}, x \ \&\& \ (\text{IntegerQ}[2 \cdot m] \ || \ \text{NeQ}[a^2 + b^2, 0])$
- rule 3988  $\text{Int}[\sec[(e_ + (f_ \cdot x))] / ((a_ + (b_ \cdot \tan[(e_ + (f_ \cdot x)])) \cdot x)], x\_Symbol] \rightarrow \text{Simp}[-f^{(-1)} \ \text{Subst}[\text{Int}[1/(a^2 + b^2 - x^2), x], x, (b - a \cdot \tan[e + f \cdot x]) / \text{Sec}[e + f \cdot x]], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 3989  $\text{Int}[\sec[(e_ + (f_ \cdot x))]^{(m_)} / ((a_ + (b_ \cdot \tan[(e_ + (f_ \cdot x)])) \cdot x)], x\_Symbol] \rightarrow \text{Simp}[-(b^2)^{-1} \ \text{Int}[\text{Sec}[e + f \cdot x]^{(m - 2)} \cdot (a - b \cdot \tan[e + f \cdot x]), x], x] + \text{Simp}[(a^2 + b^2) / b^2 \ \text{Int}[\text{Sec}[e + f \cdot x]^{(m - 2)} / (a + b \cdot \tan[e + f \cdot x]), x], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{IGtQ}[(m - 1) / 2, 0]$
- rule 4255  $\text{Int}[(\csc[(c_ + (d_ \cdot x)] \cdot (b_ \cdot x))^n], x\_Symbol] \rightarrow \text{Simp}[(-b) \cdot \text{Cos}[c + d \cdot x] \cdot ((b \cdot \csc[c + d \cdot x])^{(n - 1)} / (d \cdot (n - 1))), x] + \text{Simp}[b^2 \cdot ((n - 2) / (n - 1)) \ \text{Int}[(b \cdot \csc[c + d \cdot x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2 \cdot n]$
- rule 4257  $\text{Int}[\csc[(c_ + (d_ \cdot x)]), x\_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d \cdot x]] / d, x] /; \text{FreeQ}\{c, d\}, x]$

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 322 vs.  $2(143) = 286$ .

Time = 199.66 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.06

method	result
default	$\frac{2(-a^6+3a^4b^2-3a^2b^4+b^6) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right)+2b}{2\sqrt{a^2-b^2}}\right)}{b^6\sqrt{a^2-b^2}} + \frac{2\left(\left(\frac{1}{2}a^3b^2-\frac{9}{8}ab^4\right) \tanh\left(\frac{x}{2}\right)^9 + (a^4b-3a^2b^3+3b^5) \tanh\left(\frac{x}{2}\right)^8 + (a^3b^2-\frac{5}{4}ab^4) \tanh\left(\frac{x}{2}\right)^7 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right)^6 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right)^5 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right)^4 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right)^3 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right)^2 + (a^2b^3-\frac{3}{2}ab^5) \tanh\left(\frac{x}{2}\right) + (a^2b^3-\frac{3}{2}ab^5)\right)}{60b^5}$
risch	$\frac{e^x(120a^4e^{8x}-60a^3be^{8x}-240a^2b^2e^{8x}+105ab^3e^{8x}+120b^4e^{8x}+480a^4e^{6x}-120ba^3e^{6x}-1120a^2b^2e^{6x}+330ab^3e^{6x}+640b^4e^{6x}+720e^{4x}-120a^4e^{2x}-60a^3be^{2x}-240a^2b^2e^{2x}+105ab^3e^{2x}+120b^4e^{2x}+480a^4e^{0x}-120ba^3e^{0x}-1120a^2b^2e^{0x}+330ab^3e^{0x}+640b^4e^{0x}+720e^{0x})}{60b^5}$

input

```
int(sech(x)^7/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
2*(-a^6+3*a^4*b^2-3*a^2*b^4+b^6)/b^6/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))+2/b^6*(((1/2*a^3*b^2-9/8*a*b^4)*tanh(1/2*x)^9+(a^4*b-3*a^2*b^3+3*b^5)*tanh(1/2*x)^8+(a^3*b^2-5/4*a*b^4)*tanh(1/2*x)^7+(4*a^4*b-10*a^2*b^3+6*b^5)*tanh(1/2*x)^6+(6*a^4*b-40/3*a^2*b^3+28/3*b^5)*tanh(1/2*x)^4+(-a^3*b^2+5/4*a*b^4)*tanh(1/2*x)^3+(4*a^4*b-26/3*a^2*b^3+14/3*b^5)*tanh(1/2*x)^2+(-1/2*a^3*b^2+9/8*a*b^4)*tanh(1/2*x)+a^4*b-7/3*a^2*b^3+23/15*b^5)/(tanh(1/2*x)^2+1)^5+1/8*a*(8*a^4-20*a^2*b^2+15*b^4)*arctan(tanh(1/2*x)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 3227 vs.  $2(143) = 286$ .

Time = 0.26 (sec) , antiderivative size = 6509, normalized size of antiderivative = 41.46

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
Too large to include
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx$$

input `integrate(sech(x)**7/(a+b*tanh(x)), x)`

output `Integral(sech(x)**7/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(143) = 286.

Time = 0.13 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.08

$$\begin{aligned} & \int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx \\ &= \frac{(8a^5 - 20a^3b^2 + 15ab^4) \arctan(e^x)}{4b^6} - \frac{2(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}b^6} \\ &+ \frac{120a^4e^{(9x)} - 60a^3be^{(9x)} - 240a^2b^2e^{(9x)} + 105ab^3e^{(9x)} + 120b^4e^{(9x)} + 480a^4e^{(7x)} - 120a^3be^{(7x)} - 11}{\dots} \end{aligned}$$



input `integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="giac")`

output 
$$\begin{aligned} & \frac{1}{4}(8a^5 - 20a^3b^2 + 15ab^4) \arctan(e^x)/b^6 - 2(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \arctan((a e^x + b e^x)/\sqrt{a^2 - b^2})/(\sqrt{a^2 - b^2} \\ & * b^6) + 1/60(120a^4e^{9x} - 60a^3b e^{9x} - 240a^2b^2e^{9x} + 105ab^3e^{9x} + 120b^4e^{9x} + 480a^4e^{7x} - 120a^3b e^{7x} - \\ & 1120a^2b^2e^{7x} + 330ab^3e^{7x} + 640b^4e^{7x} + 720a^4e^{5x} - 1760a^2b^2e^{5x} + 1424b^4e^{5x} + 480a^4e^{3x} + 120a^3b \\ & b e^{3x} - 1120a^2b^2e^{3x} - 330ab^3e^{3x} + 640b^4e^{3x} + 120a^4e^x + 60a^3b e^x - 240a^2b^2e^x - 105ab^3e^x + 120b^4e^x) \\ & / (b^5(e^{2x} + 1)^5) \end{aligned}$$

### Mupad [B] (verification not implemented)

Time = 8.31 (sec) , antiderivative size = 447, normalized size of antiderivative = 2.85

$$\begin{aligned} \int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx &= \frac{32 e^x}{5 b (5 e^{2x} + 10 e^{4x} + 10 e^{6x} + 5 e^{8x} + e^{10x} + 1)} \\ & - \frac{\ln \left( \sqrt{-(a+b)^5 (a-b)^5} + a^5 e^x + b^5 e^x + a b^4 e^x + a^4 b e^x - 2 a^2 b^3 e^x - 2 a^3 b^2 e^x \right) \sqrt{-(a+b)^5 (a-b)^5}}{b^6} \\ & + \frac{\ln \left( a^5 e^x - \sqrt{-(a+b)^5 (a-b)^5} + b^5 e^x + a b^4 e^x + a^4 b e^x - 2 a^2 b^3 e^x - 2 a^3 b^2 e^x \right) \sqrt{-(a+b)^5 (a-b)^5}}{b^6} \\ & - \frac{e^x (-12 a^3 + 16 a^2 b + 9 a b^2 - 16 b^3)}{6 b^4 (2 e^{2x} + e^{4x} + 1)} + \frac{e^x (8 a^4 - 4 a^3 b - 16 a^2 b^2 + 7 a b^3 + 8 b^4)}{4 b^5 (e^{2x} + 1)} \\ & + \frac{4 e^x (5 a - 16 b)}{5 b^2 (4 e^{2x} + 6 e^{4x} + 4 e^{6x} + e^{8x} + 1)} + \frac{2 e^x (20 a^2 - 45 a b + 28 b^2)}{15 b^3 (3 e^{2x} + 3 e^{4x} + e^{6x} + 1)} \\ & - \frac{a \ln(e^x - i) (8 a^4 - 20 a^2 b^2 + 15 b^4) \operatorname{li}}{8 b^6} + \frac{a \ln(e^x + i) (8 a^4 - 20 a^2 b^2 + 15 b^4) \operatorname{li}}{8 b^6} \end{aligned}$$

input `int(1/(cosh(x)^7*(a + b*tanh(x))),x)`

output

```
(32*exp(x))/(5*b*(5*exp(2*x) + 10*exp(4*x) + 10*exp(6*x) + 5*exp(8*x) + exp(10*x) + 1)) - (log((-a + b)^5*(a - b)^5)^(1/2) + a^5*exp(x) + b^5*exp(x) + a*b^4*exp(x) + a^4*b*exp(x) - 2*a^2*b^3*exp(x) - 2*a^3*b^2*exp(x))*(-(a + b)^5*(a - b)^5)^(1/2))/b^6 + (log(a^5*exp(x) - (-a + b)^5*(a - b)^5)^(1/2) + b^5*exp(x) + a*b^4*exp(x) + a^4*b*exp(x) - 2*a^2*b^3*exp(x) - 2*a^3*b^2*exp(x))*(-(a + b)^5*(a - b)^5)^(1/2))/b^6 - (exp(x)*(9*a*b^2 + 16*a^2*b - 12*a^3 - 16*b^3))/(6*b^4*(2*exp(2*x) + exp(4*x) + 1)) + (exp(x)*(7*a*b^3 - 4*a^3*b + 8*a^4 + 8*b^4 - 16*a^2*b^2))/(4*b^5*(exp(2*x) + 1)) - (a*log(exp(x) - 1i)*(8*a^4 + 15*b^4 - 20*a^2*b^2)*1i)/(8*b^6) + (a*log(exp(x) + 1i)*(8*a^4 + 15*b^4 - 20*a^2*b^2)*1i)/(8*b^6) + (4*exp(x)*(5*a - 16*b))/(5*b^2*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) + (2*exp(x)*(20*a^2 - 45*a*b + 28*b^2))/(15*b^3*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 1373, normalized size of antiderivative = 8.75

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
int(sech(x)^7/(a+b*tanh(x)),x)
```

output

```
(120***e**(10*x)*atan(e**x)*a**5 - 300***e**(10*x)*atan(e**x)*a**3*b**2 + 225*
e**(10*x)*atan(e**x)*a*b**4 + 600***e**(8*x)*atan(e**x)*a**5 - 1500***e**(8*x)
*atan(e**x)*a**3*b**2 + 1125***e**(8*x)*atan(e**x)*a*b**4 + 1200***e**(6*x)*at
an(e**x)*a**5 - 3000***e**(6*x)*atan(e**x)*a**3*b**2 + 2250***e**(6*x)*atan(e*
*x)*a*b**4 + 1200***e**(4*x)*atan(e**x)*a**5 - 3000***e**(4*x)*atan(e**x)*a**3
*b**2 + 2250***e**(4*x)*atan(e**x)*a*b**4 + 600***e**(2*x)*atan(e**x)*a**5 - 1
500***e**(2*x)*atan(e**x)*a**3*b**2 + 1125***e**(2*x)*atan(e**x)*a*b**4 + 120*
atan(e**x)*a**5 - 300*atan(e**x)*a**3*b**2 + 225*atan(e**x)*a*b**4 - 120*e
**(10*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**4
+ 240***e**(10*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2)
)*a**2*b**2 - 120***e**(10*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(
a**2 - b**2))*b**4 - 600***e**(8*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)
/sqrt(a**2 - b**2))*a**4 + 1200***e**(8*x)*sqrt(a**2 - b**2)*atan((e**x*a +
e**x*b)/sqrt(a**2 - b**2))*a**2*b**2 - 600***e**(8*x)*sqrt(a**2 - b**2)*atan
((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**4 - 1200***e**(6*x)*sqrt(a**2 - b**
2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**4 + 2400***e**(6*x)*sqrt(a**
2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**2*b**2 - 1200***e**(6
*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**4 - 120
0***e**(4*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*a**
4 + 2400***e**(4*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - ...
```

### 3.111 $\int \frac{\operatorname{sech}^5(x)}{a+b \tanh(x)} dx$

Optimal result	867
Mathematica [A] (verified)	867
Rubi [A] (verified)	868
Maple [A] (verified)	871
Fricas [B] (verification not implemented)	872
Sympy [F]	873
Maxima [F(-2)]	874
Giac [A] (verification not implemented)	874
Mupad [B] (verification not implemented)	875
Reduce [B] (verification not implemented)	875

#### Optimal result

Integrand size = 13, antiderivative size = 102

$$\int \frac{\operatorname{sech}^5(x)}{a+b \tanh(x)} dx = -\frac{a(2a^2 - 3b^2) \arctan(\sinh(x))}{2b^4} + \frac{(a^2 - b^2)^{3/2} \arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{b^4} - \frac{(a^2 - b^2) \operatorname{sech}(x)}{b^3} + \frac{\operatorname{sech}^3(x)}{3b} + \frac{a \operatorname{sech}(x) \tanh(x)}{2b^2}$$

output

```
-1/2*a*(2*a^2-3*b^2)*arctan(sinh(x))/b^4+(a^2-b^2)^(3/2)*arctan(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/b^4-(a^2-b^2)*sech(x)/b^3+1/3*sech(x)^3/b+1/2*a*sech(x)*tanh(x)/b^2
```

#### Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.14

$$\int \frac{\operatorname{sech}^5(x)}{a+b \tanh(x)} dx = \frac{-6\left(a(2a^2 - 3b^2) \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + 2\sqrt{a-b}\sqrt{a+b}(-a^2 + b^2) \arctan\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right)\right) + 2b^3 \operatorname{sech}^3(x) + a \operatorname{sech}(x) \tanh(x)}{6b^4}$$

input `Integrate[Sech[x]^5/(a + b*Tanh[x]), x]`

output `(-6*(a*(2*a^2 - 3*b^2)*ArcTan[Tanh[x/2]] + 2*Sqrt[a - b]*Sqrt[a + b]*(-a^2 + b^2)*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])]) + 2*b^3*Sech[x]^3 + 3*b*Sech[x]*(-2*a^2 + 2*b^2 + a*b*Tanh[x]))/(6*b^4)`

### Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.077$ , Rules used = {3042, 3989, 3042, 3967, 3042, 3989, 3042, 3967, 3042, 3988, 219, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sec(ix)^5}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{3989} \\
 & \frac{\int \operatorname{sech}^3(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sec(ix)^3(a + ib \tan(ix)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^3}{a - ib \tan(ix)} dx}{b^2} \\
 & \quad \downarrow \text{3967} \\
 & \frac{a \int \operatorname{sech}^3(x) dx + \frac{1}{3} b \operatorname{sech}^3(x)}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^3}{a - ib \tan(ix)} dx}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)^3}{a - ib \tan(ix)} dx}{b^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3989} \\
& \frac{(a^2 - b^2) \left( \frac{\int \operatorname{sech}(x)(a-b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx}{b^2} \right)}{b^2} + \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} \\
& \downarrow \text{3042} \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{\int \sec(ix)(a+ib \tan(ix)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
& \downarrow \text{3967} \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \frac{(a^2 - b^2) \left( \frac{a \int \operatorname{sech}(x) dx + b \operatorname{sech}(x)}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
& \downarrow \text{3042} \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \\
& \frac{(a^2 - b^2) \left( \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{b^2} \right)}{b^2} \\
& \downarrow \text{3988} \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \\
& \frac{(a^2 - b^2) \left( \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} - \frac{i(a^2 - b^2) \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b+a \tanh(x))^2} d(-i \cosh(x)(b+a \tanh(x)))}{b^2} \right)}{b^2} \\
& \downarrow \text{219} \\
& \frac{\frac{1}{3} b \operatorname{sech}^3(x) + a \int \csc\left(ix + \frac{\pi}{2}\right)^3 dx}{b^2} - \\
& \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2} \\
& \downarrow \text{4255}
\end{aligned}$$

$$\frac{a\left(\frac{\int \operatorname{sech}(x) dx}{2} + \frac{1}{2} \tanh(x) \operatorname{sech}(x)\right) + \frac{1}{3} b \operatorname{sech}^3(x)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2}$$

↓ 3042

$$\frac{\frac{1}{3} b \operatorname{sech}^3(x) + a\left(\frac{1}{2} \tanh(x) \operatorname{sech}(x) + \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx\right)}{b^2} - \frac{(a^2 - b^2) \left( -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \right)}{b^2}$$

↓ 4257

$$\frac{a\left(\frac{1}{2} \arctan(\sinh(x)) + \frac{1}{2} \tanh(x) \operatorname{sech}(x)\right) + \frac{1}{3} b \operatorname{sech}^3(x)}{b^2} - \frac{(a^2 - b^2) \left( \frac{a \arctan(\sinh(x)) + b \operatorname{sech}(x)}{b^2} - \frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} \right)}{b^2}$$

input `Int [Sech [x]^5/(a + b*Tanh [x]), x]`

output `-(((a^2 - b^2)*(-(Sqrt[a^2 - b^2]*ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]])/b^2) + (a*ArcTan[Sinh[x]] + b*Sech[x])/b^2))/b^2 + ((b*Sech[x]^3)/3 + a*(ArcTan[Sinh[x]]/2 + (Sech[x]*Tanh[x])/2))/b^2`

**Defintions of rubi rules used**

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3967 Int[((d.)*sec[(e.) + (f.)*(x.)])^(m.)*((a.) + (b.)*tan[(e.) + (f.)*(x.)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Simp[a Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] || NeQ[a^2 + b^2, 0])
```

```
rule 3988 Int[sec[(e.) + (f.)*(x.)]/((a.) + (b.)*tan[(e.) + (f.)*(x.)]), x_Symbol] := Simp[-f^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]
```

```
rule 3989 Int[sec[(e.) + (f.)*(x.)]^(m.)/((a.) + (b.)*tan[(e.) + (f.)*(x.)]), x_Symbol] := Simp[-(b^2)^(-1) Int[Sec[e + f*x]^(m - 2)*(a - b*Tan[e + f*x]), x], x] + Simp[(a^2 + b^2)/b^2 Int[Sec[e + f*x]^(m - 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[(m - 1)/2, 0]
```

```
rule 4255 Int[(csc[(c.) + (d.)*(x.)]*(b.))^(n.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

```
rule 4257 Int[csc[(c.) + (d.)*(x.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

**Maple [A] (verified)**

Time = 48.64 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.61

method	result
default	$\frac{2(a^4 - 2a^2b^2 + b^4) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{b^4\sqrt{a^2 - b^2}} - \frac{2\left(\frac{b^2 a \tanh\left(\frac{x}{2}\right)^5}{2} + (a^2 b - 2b^3) \tanh\left(\frac{x}{2}\right)^4 + (2a^2 b - 2b^3) \tanh\left(\frac{x}{2}\right)^2 - \frac{b^2 a \tanh\left(\frac{x}{2}\right)}{2} + a^2 b - b^3\right)}{b^4 \left(\tanh\left(\frac{x}{2}\right)^2 + 1\right)^3}$
risch	$-\frac{e^x(6a^2e^{4x} - 3e^{4x}ab - 6b^2e^{4x} + 12e^{2x}a^2 - 20b^2e^{2x} + 6a^2 + 3ab - 6b^2)}{3b^3(e^{2x} + 1)^3} + \frac{\sqrt{-a^2 + b^2} \ln\left(e^x + \frac{\sqrt{-a^2 + b^2}}{a + b}\right)a^2}{b^4} - \frac{\sqrt{-a^2 + b^2} \ln\left(e^x + \frac{\sqrt{-a^2 + b^2}}{a - b}\right)a^2}{b^2}$



input `int(sech(x)^5/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output `2*(a^4-2*a^2*b^2+b^4)/b^4/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))-2/b^4*((1/2*b^2*a*tanh(1/2*x)^5+(a^2*b-2*b^3)*tanh(1/2*x)^4+(2*a^2*b-2*b^3)*tanh(1/2*x)^2-1/2*b^2*a*tanh(1/2*x)+a^2*b-4/3*b^3)/(tanh(1/2*x)^2+1)^3+1/2*a*(2*a^2-3*b^2)*arctan(tanh(1/2*x))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 994 vs.  $2(92) = 184$ .

Time = 0.14 (sec) , antiderivative size = 2043, normalized size of antiderivative = 20.03

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="fricas")`

output

```

[-1/3*(3*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^5 + 15*(2*a^2*b - a*b^2 - 2*b^3)
)*cosh(x)*sinh(x)^4 + 3*(2*a^2*b - a*b^2 - 2*b^3)*sinh(x)^5 + 4*(3*a^2*b -
5*b^3)*cosh(x)^3 + 2*(6*a^2*b - 10*b^3 + 15*(2*a^2*b - a*b^2 - 2*b^3)*cos
h(x)^2)*sinh(x)^3 + 6*(5*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^3 + 2*(3*a^2*b
- 5*b^3)*cosh(x))*sinh(x)^2 + 3*((a^2 - b^2)*cosh(x)^6 + 6*(a^2 - b^2)*cos
h(x)*sinh(x)^5 + (a^2 - b^2)*sinh(x)^6 + 3*(a^2 - b^2)*cosh(x)^4 + 3*(5*(a
^2 - b^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^4 + 4*(5*(a^2 - b^2)*cosh(x)^3 +
3*(a^2 - b^2)*cosh(x))*sinh(x)^3 + 3*(a^2 - b^2)*cosh(x)^2 + 3*(5*(a^2 - b
^2)*cosh(x)^4 + 6*(a^2 - b^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^2 + a^2 - b^2
+ 6*((a^2 - b^2)*cosh(x)^5 + 2*(a^2 - b^2)*cosh(x)^3 + (a^2 - b^2)*cosh(x
))*sinh(x))*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*si
nh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b
)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a -
b)) + 3*((2*a^3 - 3*a*b^2)*cosh(x)^6 + 6*(2*a^3 - 3*a*b^2)*cosh(x)*sinh(x)
)^5 + (2*a^3 - 3*a*b^2)*sinh(x)^6 + 3*(2*a^3 - 3*a*b^2)*cosh(x)^4 + 3*(2*a
^3 - 3*a*b^2 + 5*(2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^4 + 4*(5*(2*a^3 - 3*
a*b^2)*cosh(x)^3 + 3*(2*a^3 - 3*a*b^2)*cosh(x))*sinh(x)^3 + 2*a^3 - 3*a*b^
2 + 3*(2*a^3 - 3*a*b^2)*cosh(x)^2 + 3*(5*(2*a^3 - 3*a*b^2)*cosh(x)^4 + 2*a
^3 - 3*a*b^2 + 6*(2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^2 + 6*((2*a^3 - 3*a*
b^2)*cosh(x)^5 + 2*(2*a^3 - 3*a*b^2)*cosh(x)^3 + (2*a^3 - 3*a*b^2)*cosh...

```

## Sympy [F]

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx$$

input

```
integrate(sech(x)**5/(a+b*tanh(x)),x)
```

output

```
Integral(sech(x)**5/(a + b*tanh(x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.49

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx$$

$$= -\frac{(2a^3 - 3ab^2) \arctan(e^x)}{b^4} + \frac{2(a^4 - 2a^2b^2 + b^4) \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2} b^4}$$

$$-\frac{6a^2e^{5x} - 3abe^{5x} - 6b^2e^{5x} + 12a^2e^{3x} - 20b^2e^{3x} + 6a^2e^x + 3abe^x - 6b^2e^x}{3b^3(e^{2x} + 1)^3}$$

input `integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="giac")`

output `-(2*a^3 - 3*a*b^2)*arctan(e^x)/b^4 + 2*(a^4 - 2*a^2*b^2 + b^4)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*b^4) - 1/3*(6*a^2*e^(5*x) - 3*a*b*e^(5*x) - 6*b^2*e^(5*x) + 12*a^2*e^(3*x) - 20*b^2*e^(3*x) + 6*a^2*e^x + 3*a*b*e^x - 6*b^2*e^x)/(b^3*(e^(2*x) + 1)^3)`

### Mupad [B] (verification not implemented)

Time = 5.89 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.60

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx$$

$$= \frac{\ln \left( \sqrt{-(a+b)^3(a-b)^3} + a^3 e^x - b^3 e^x - a b^2 e^x + a^2 b e^x \right) \sqrt{-(a+b)^3(a-b)^3}}{b^4}$$

$$- \frac{8 e^x}{3 b (3 e^{2x} + 3 e^{4x} + e^{6x} + 1)}$$

$$- \frac{\ln \left( \sqrt{-(a+b)^3(a-b)^3} - a^3 e^x + b^3 e^x + a b^2 e^x - a^2 b e^x \right) \sqrt{-(a+b)^3(a-b)^3}}{b^4}$$

$$- \frac{2 e^x (3 a - 4 b)}{3 b^2 (2 e^{2x} + e^{4x} + 1)} + \frac{e^x (-2 a^2 + a b + 2 b^2)}{b^3 (e^{2x} + 1)}$$

$$+ \frac{a \ln(e^x - i) (2 a^2 - 3 b^2) i}{2 b^4} - \frac{a \ln(e^x + i) (2 a^2 - 3 b^2) i}{2 b^4}$$

input

```
int(1/(cosh(x)^5*(a + b*tanh(x))),x)
```

output

```
(log((-a + b)^3*(a - b)^3)^(1/2) + a^3*exp(x) - b^3*exp(x) - a*b^2*exp(x)
+ a^2*b*exp(x))*(-(a + b)^3*(a - b)^3)^(1/2)/b^4 - (8*exp(x))/(3*b*(3*exp
p(2*x) + 3*exp(4*x) + exp(6*x) + 1)) - (log((-a + b)^3*(a - b)^3)^(1/2) -
a^3*exp(x) + b^3*exp(x) + a*b^2*exp(x) - a^2*b*exp(x))*(-(a + b)^3*(a - b
)^3)^(1/2)/b^4 - (2*exp(x)*(3*a - 4*b))/(3*b^2*(2*exp(2*x) + exp(4*x) + 1
)) + (exp(x)*(a*b - 2*a^2 + 2*b^2))/(b^3*(exp(2*x) + 1)) + (a*log(exp(x) -
1i)*(2*a^2 - 3*b^2)*1i)/(2*b^4) - (a*log(exp(x) + 1i)*(2*a^2 - 3*b^2)*1i)
/(2*b^4)
```

### Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 564, normalized size of antiderivative = 5.53

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx$$

$$= \frac{-6e^{6x} \operatorname{atan}(e^x) a^3 + 9e^{6x} \operatorname{atan}(e^x) a b^2 - 18e^{4x} \operatorname{atan}(e^x) a^3 + 27e^{4x} \operatorname{atan}(e^x) a b^2 - 18e^{2x} \operatorname{atan}(e^x) a^3 + 27e^{2x} \operatorname{atan}(e^x) a b^2}{\dots}$$

input `int(sech(x)^5/(a+b*tanh(x)),x)`

output

$$\begin{aligned} & (-6e^{6x} \operatorname{atan}(e^x) a^3 + 9e^{6x} \operatorname{atan}(e^x) a b^2 - 18e^{4x} \operatorname{atan}(e^x) a^3 + 27e^{4x} \operatorname{atan}(e^x) a b^2 - 18e^{2x} \operatorname{atan}(e^x) a^3 \\ & + 27e^{2x} \operatorname{atan}(e^x) a b^2 - 6 \operatorname{atan}(e^x) a^3 + 9 \operatorname{atan}(e^x) a b^2 + 6e^{6x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) \\ & a^2 - 6e^{6x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) b^2 + 18e^{4x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) \\ & a^2 - 18e^{4x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) b^2 + 18e^{2x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) \\ & a^2 - 18e^{2x} \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) b^2 + 6 \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) \\ & a^2 - 6 \sqrt{a^2 - b^2} \operatorname{atan}((e^x a + e^x b) / \sqrt{a^2 - b^2}) b^2 - 6e^{5x} a^2 b + 3e^{5x} a b^2 + 6e^{5x} b^3 \\ & - 12e^{3x} a^2 b + 20e^{3x} b^3 - 6e^{5x} a^2 b - 3e^{5x} a b^2 + 6e^{5x} b^3) / (3b^4 (e^{6x} + 3e^{4x} + 3e^{2x} + 1)) \end{aligned}$$

### 3.112 $\int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx$

Optimal result . . . . .	877
Mathematica [A] (verified) . . . . .	877
Rubi [A] (verified) . . . . .	878
Maple [A] (verified) . . . . .	880
Fricas [B] (verification not implemented) . . . . .	881
Sympy [F] . . . . .	881
Maxima [F(-2)] . . . . .	882
Giac [A] (verification not implemented) . . . . .	882
Mupad [B] (verification not implemented) . . . . .	882
Reduce [B] (verification not implemented) . . . . .	883

#### Optimal result

Integrand size = 13, antiderivative size = 56

$$\int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx = \frac{a \arctan(\sinh(x))}{b^2} - \frac{\sqrt{a^2-b^2} \arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{b^2} + \frac{\operatorname{sech}(x)}{b}$$

output

`a*arctan(sinh(x))/b^2-(a^2-b^2)^(1/2)*arctan(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/b^2+sech(x)/b`

#### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx = \frac{2a \arctan\left(\tanh\left(\frac{x}{2}\right)\right) - 2\sqrt{a-b}\sqrt{a+b} \arctan\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right) + b \operatorname{sech}(x)}{b^2}$$

input

`Integrate[Sech[x]^3/(a + b*Tanh[x]), x]`

output

```
(2*a*ArcTan[Tanh[x/2]] - 2*Sqrt[a - b]*Sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])] + b*Sech[x])/b^2
```

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$ , Rules used = {3042, 3989, 3042, 3967, 3042, 3988, 219, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sec(ix)^3}{a - ib \tan(ix)} dx$$

$$\downarrow \text{3989}$$

$$\frac{\int \operatorname{sech}(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{b^2}$$

$$\downarrow \text{3042}$$

$$\frac{\int \sec(ix)(a + ib \tan(ix)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2}$$

$$\downarrow \text{3967}$$

$$\frac{a \int \operatorname{sech}(x) dx + b \operatorname{sech}(x)}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2}$$

$$\downarrow \text{3042}$$

$$\frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{b^2}$$

$$\downarrow \text{3988}$$

$$\begin{aligned}
 & \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \\
 & \frac{i(a^2 - b^2) \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b + a \tanh(x))^2} d(-i \cosh(x)(b + a \tanh(x)))}{b^2} \\
 & \quad \downarrow 219 \\
 & -\frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{b \operatorname{sech}(x) + a \int \csc\left(ix + \frac{\pi}{2}\right) dx}{b^2} \\
 & \quad \downarrow 4257 \\
 & \frac{a \arctan(\sinh(x)) + b \operatorname{sech}(x)}{b^2} - \frac{\sqrt{a^2 - b^2} \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{b^2}
 \end{aligned}$$

input `Int [Sech[x]^3/(a + b*Tanh[x]), x]`

output `-((Sqrt[a^2 - b^2]*ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]])/b^2) + (a*ArcTan[Sinh[x]] + b*Sech[x])/b^2`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3967 `Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Simp[a Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] || NeQ[a^2 + b^2, 0])`



rule 3988

```
Int[sec[(e_.) + (f_.)*(x_)]/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
:= Simp[-f^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x]
/; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]
```

rule 3989

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
:= Simp[-(b^2)^(-1) Int[Sec[e + f*x]^(m - 2)*(a - b*Tan[e + f*x]), x], x]
+ Simp[(a^2 + b^2)/b^2 Int[Sec[e + f*x]^(m - 2)/(a + b*Tan[e + f*x]), x], x]
/; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[(m - 1)/2, 0]
```

rule 4257

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

### Maple [A] (verified)

Time = 8.37 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.38

method	result	size
default	$\frac{\frac{2b}{\tanh\left(\frac{x}{2}\right)^2+1} + 2a \arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{b^2} + \frac{2(-a^2+b^2) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right)+2b}{2\sqrt{a^2-b^2}}\right)}{b^2\sqrt{a^2-b^2}}$	77
risch	$\frac{2e^x}{b(e^{2x}+1)} + \frac{ia \ln(e^x+i)}{b^2} - \frac{ia \ln(e^x-i)}{b^2} + \frac{\sqrt{-a^2+b^2} \ln\left(e^x - \frac{\sqrt{-a^2+b^2}}{a+b}\right)}{b^2} - \frac{\sqrt{-a^2+b^2} \ln\left(e^x + \frac{\sqrt{-a^2+b^2}}{a+b}\right)}{b^2}$	117

input

```
int(sech(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
2/b^2*(b/(tanh(1/2*x)^2+1)+a*arctan(tanh(1/2*x)))+2*(-a^2+b^2)/b^2/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 126 vs.  $2(52) = 104$ .

Time = 0.11 (sec) , antiderivative size = 309, normalized size of antiderivative = 5.52

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx$$

$$= \left[ \frac{\sqrt{-a^2 + b^2} (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \log \left( \frac{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2}{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2} \right) + \dots}{b^2 \cosh(x)^2 + 2b^2 \cosh(x) \sinh(x) + b^2 \sinh(x)^2 + b^2} \right]$$

input `integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

output `[(sqrt(-a^2 + b^2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b)) + 2*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*arctan(cosh(x) + sinh(x)) + 2*b*cosh(x) + 2*b*sinh(x))/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2 + b^2), 2*(sqrt(a^2 - b^2)*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x))) + (a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 + a)*arctan(cosh(x) + sinh(x)) + b*cosh(x) + b*sinh(x))/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2 + b^2)]`

**Sympy [F]**

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx$$

input `integrate(sech(x)**3/(a+b*tanh(x)),x)`

output `Integral(sech(x)**3/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(4\*b^2-4\*a^2>0)', see `assume?` f or more de

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx = \frac{2a \arctan(e^x)}{b^2} - \frac{2\sqrt{a^2 - b^2} \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{2e^x}{b(e^{2x} + 1)}$$

input `integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="giac")`

output `2*a*arctan(e^x)/b^2 - 2*sqrt(a^2 - b^2)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/b^2 + 2*e^x/(b*(e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 4.68 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.12

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx = \frac{\ln(ae^x + be^x - \sqrt{b^2 - a^2}) \sqrt{-(a+b)(a-b)}}{b^2} - \frac{\ln(ae^x + be^x + \sqrt{b^2 - a^2}) \sqrt{-(a+b)(a-b)}}{b^2} + \frac{2e^x}{b(e^{2x} + 1)} - \frac{a \ln(e^x - i) \operatorname{li}}{b^2} + \frac{a \ln(e^x + 1) \operatorname{li}}{b^2}$$

input `int(1/(cosh(x)^3*(a + b*tanh(x))),x)`

output `(a*log(exp(x) + 1i)*1i)/b^2 - (a*log(exp(x) - 1i)*1i)/b^2 - (log(a*exp(x) + b*exp(x) + (b^2 - a^2)^(1/2)))*(-(a + b)*(a - b))^(1/2))/b^2 + (log(a*exp(x) + b*exp(x) - (b^2 - a^2)^(1/2)))*(-(a + b)*(a - b))^(1/2))/b^2 + (2*exp(x))/(b*(exp(2*x) + 1))`

### Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.07

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx$$

$$= \frac{2e^{2x} \operatorname{atan}(e^x) a + 2 \operatorname{atan}(e^x) a - 2e^{2x} \sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) - 2\sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) + 2e^x b}{b^2 (e^{2x} + 1)}$$

input `int(sech(x)^3/(a+b*tanh(x)),x)`

output `(2*(e**(2*x))*atan(e**x)*a + atan(e**x)*a - e**(2*x)*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2)) - sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2)) + e**x*b)/(b**2*(e**(2*x) + 1))`

### 3.113 $\int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx$

Optimal result	884
Mathematica [A] (verified)	884
Rubi [A] (verified)	885
Maple [A] (verified)	886
Fricas [A] (verification not implemented)	886
Sympy [F]	887
Maxima [F(-2)]	887
Giac [A] (verification not implemented)	888
Mupad [B] (verification not implemented)	888
Reduce [B] (verification not implemented)	888

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx = \frac{\arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

output `arctan(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/(a^2-b^2)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24

$$\int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx = \frac{2 \arctan\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right)}{\sqrt{a-b}\sqrt{a+b}}$$

input `Integrate[Sech[x]/(a + b*Tanh[x]), x]`

output `(2*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])])/(Sqrt[a - b]*Sqrt[a + b])`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {3042, 3988, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx$$

↓ 3042

$$\int \frac{\sec(ix)}{a - ib \tan(ix)} dx$$

↓ 3988

$$i \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b + a \tanh(x))^2} d(-i \cosh(x)(b + a \tanh(x)))$$

↓ 219

$$\frac{\arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}}$$

input `Int[Sech[x]/(a + b*Tanh[x]), x]`

output `ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]]/Sqrt[a^2 - b^2]`

**Defintions of rubi rules used**

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3988

```
Int[sec[(e_.) + (f_.)*(x_)]/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
  := Simp[-f^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f
*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]
```

**Maple [A] (verified)**

Time = 0.78 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05

method	result	size
default	$\frac{2 \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}}$	39
risch	$-\frac{\ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + \frac{\ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$	70

input

```
int(sech(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
2/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 148, normalized size of antiderivative = 4.00

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx$$

$$= \left[ -\frac{\sqrt{-a^2 + b^2} \log\left(\frac{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 - 2\sqrt{-a^2+b^2}(\cosh(x)+\sinh(x))-a+b}{(a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a-b}\right)}{a^2 - b^2}, \right. \\ \left. -\frac{2 \arctan\left(\frac{\sqrt{a^2-b^2}}{(a+b) \cosh(x) + (a+b) \sinh(x)}\right)}{\sqrt{a^2 - b^2}} \right]$$

input

```
integrate(sech(x)/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
[-sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b))/(a^2 - b^2), -2*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x)))/sqrt(a^2 - b^2)]
```

**Sympy [F]**

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx = \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx$$

input

```
integrate(sech(x)/(a+b*tanh(x)),x)
```

output

```
Integral(sech(x)/(a + b*tanh(x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input

```
integrate(sech(x)/(a+b*tanh(x)),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` for more de
```



**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx = \frac{2 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}}$$

input `integrate(sech(x)/(a+b*tanh(x)),x, algorithm="giac")`output `2*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/sqrt(a^2 - b^2)`**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx = \frac{2 \operatorname{atan}\left(\frac{e^x \sqrt{a^2 - b^2}}{a - b}\right)}{\sqrt{a^2 - b^2}}$$

input `int(1/(cosh(x)*(a + b*tanh(x))),x)`output `(2*atan((exp(x)*(a^2 - b^2)^(1/2))/(a - b)))/(a^2 - b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.30

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx = \frac{2\sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right)}{a^2 - b^2}$$

input `int(sech(x)/(a+b*tanh(x)),x)`output `(2*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2)))/(a**2 - b**2)`

### 3.114 $\int \frac{\cosh(x)}{a+b \tanh(x)} dx$

Optimal result . . . . .	889
Mathematica [A] (verified) . . . . .	889
Rubi [A] (verified) . . . . .	890
Maple [A] (verified) . . . . .	892
Fricas [B] (verification not implemented) . . . . .	892
Sympy [F] . . . . .	893
Maxima [F(-2)] . . . . .	893
Giac [A] (verification not implemented) . . . . .	894
Mupad [B] (verification not implemented) . . . . .	894
Reduce [B] (verification not implemented) . . . . .	895

#### Optimal result

Integrand size = 11, antiderivative size = 73

$$\int \frac{\cosh(x)}{a+b \tanh(x)} dx = -\frac{b^2 \arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} - \frac{b \cosh(x)}{a^2-b^2} + \frac{a \sinh(x)}{a^2-b^2}$$

output

$-b^2 \arctan(\cosh(x) * (b+a \tanh(x)) / (a^2-b^2)^{(1/2)}) / (a^2-b^2)^{(3/2)} - b \cosh(x) / (a^2-b^2) + a \sinh(x) / (a^2-b^2)$

#### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.10

$$\int \frac{\cosh(x)}{a+b \tanh(x)} dx = -\frac{2b^2 \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b} \sqrt{a+b}}\right)}{(a-b)^{3/2} (a+b)^{3/2}} + \frac{b \cosh(x)}{-a^2+b^2} + \frac{a \sinh(x)}{a^2-b^2}$$

input

`Integrate[Cosh[x]/(a + b*Tanh[x]),x]`

output

$(-2*b^2 \text{ArcTan}[(b + a \text{Tanh}[x/2]) / (\text{Sqrt}[a - b] \text{Sqrt}[a + b])]) / ((a - b)^{(3/2)} * (a + b)^{(3/2)}) + (b \text{Cosh}[x]) / (-a^2 + b^2) + (a \text{Sinh}[x]) / (a^2 - b^2)$

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.88, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 3990, 3042, 3967, 3042, 3117, 3988, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cosh(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sec(ix)(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{3990} \\
 & \frac{\int \cosh(x)(a - b \tanh(x)) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a + ib \tan(ix)}{\sec(ix)} dx}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3967} \\
 & \frac{a \int \cosh(x) dx - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-b \cosh(x) + a \int \sin(ix + \frac{\pi}{2}) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3117} \\
 & \frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a - ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3988} \\
 & \frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{ib^2 \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b + a \tanh(x))^2} d(-i \cosh(x)(b + a \tanh(x)))}{a^2 - b^2} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

$$\frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{3/2}}$$

input `Int[Cosh[x]/(a + b*Tanh[x]),x]`

output `-((b^2*ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]])/(a^2 - b^2)^(3/2)) + (-b*Cosh[x] + a*Sinh[x])/(a^2 - b^2)`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_) + (d_)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3967 `Int[((d_)*sec[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Simp[a Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] || NeQ[a^2 + b^2, 0])`

rule 3988 `Int[sec[(e_) + (f_)*(x_)]/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[-f^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]`

rule 3990

```

Int[sec[(e_.) + (f_.)*(x_)]^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_
Symbol] := Simp[1/(a^2 + b^2)  Int[Sec[e + f*x]^m*(a - b*Tan[e + f*x]), x]
, x] + Simp[b^2/(a^2 + b^2)  Int[Sec[e + f*x]^(m + 2)/(a + b*Tan[e + f*x])
, x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[(m - 1)/2,
0]

```

**Maple [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.27

method	result	size
default	$-\frac{2b^2 \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{(a-b)(a+b)\sqrt{a^2 - b^2}} - \frac{2}{(2b+2a)(\tanh\left(\frac{x}{2}\right) - 1)} - \frac{2}{(2a-2b)(1+\tanh\left(\frac{x}{2}\right))}$	93
risch	$\frac{e^x}{2b+2a} - \frac{e^{-x}}{2(a-b)} - \frac{b^2 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)(a-b)} + \frac{b^2 \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)(a-b)}$	122

input

```
int(cosh(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```

-2*b^2/(a-b)/(a+b)/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b
^2)^(1/2))-2/(2*b+2*a)/(tanh(1/2*x)-1)-2/(2*a-2*b)/(1+tanh(1/2*x))

```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 190 vs. 2(69) = 138.

Time = 0.10 (sec) , antiderivative size = 435, normalized size of antiderivative = 5.96

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx$$

$$= \left[ \frac{a^3 + a^2b - ab^2 - b^3 - (a^3 - a^2b - ab^2 + b^3) \cosh(x)^2 - 2(a^3 - a^2b - ab^2 + b^3) \cosh(x) \sinh(x) - a^4}{2((a^4 - 2a^2b^2 + b^4) \cosh(x) + \dots)} \right]$$

input `integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="fricas")`

output `[-1/2*(a^3 + a^2*b - a*b^2 - b^3 - (a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^2 - 2*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh(x) - (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^2 - 2*(b^2*cosh(x) + b^2*sinh(x))*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b)))/((a^4 - 2*a^2*b^2 + b^4)*cosh(x) + (a^4 - 2*a^2*b^2 + b^4)*sinh(x)), -1/2*(a^3 + a^2*b - a*b^2 - b^3 - (a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^2 - 2*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh(x) - (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^2 - 4*(b^2*cosh(x) + b^2*sinh(x))*sqrt(a^2 - b^2)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x)))))/((a^4 - 2*a^2*b^2 + b^4)*cosh(x) + (a^4 - 2*a^2*b^2 + b^4)*sinh(x))]`

## Sympy [F]

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx = \int \frac{\cosh(x)}{a + b \tanh(x)} dx$$

input `integrate(cosh(x)/(a+b*tanh(x)),x)`

output `Integral(cosh(x)/(a + b*tanh(x)), x)`

## Maxima [F(-2)]

Exception generated.

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f
or more de
```

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.84

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx = -\frac{2b^2 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{\frac{3}{2}}} - \frac{e^{-x}}{2(a - b)} + \frac{e^x}{2(a + b)}$$

input

```
integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="giac")
```

output

```
-2*b^2*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(a^2 - b^2)^(3/2) - 1/2*e^(-
-x)/(a - b) + 1/2*e^x/(a + b)
```

### Mupad [B] (verification not implemented)

Time = 2.27 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.15

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx = \frac{e^x}{2a + 2b} - \frac{e^{-x}}{2a - 2b} - \frac{b^2 \ln\left(-\frac{2b^2}{(a+b)^{5/2} \sqrt{b-a}} - \frac{2b^2 e^x}{-a^3 - a^2 b + a b^2 + b^3}\right)}{(a + b)^{3/2} (b - a)^{3/2}} + \frac{b^2 \ln\left(\frac{2b^2}{(a+b)^{5/2} \sqrt{b-a}} - \frac{2b^2 e^x}{-a^3 - a^2 b + a b^2 + b^3}\right)}{(a + b)^{3/2} (b - a)^{3/2}}$$

input

```
int(cosh(x)/(a + b*tanh(x)),x)
```

output

```
exp(x)/(2*a + 2*b) - exp(-x)/(2*a - 2*b) - (b^2*log(- (2*b^2)/((a + b)^(5/
2)*(b - a)^(1/2)) - (2*b^2*exp(x))/(a*b^2 - a^2*b - a^3 + b^3)))/((a + b)^(
3/2)*(b - a)^(3/2)) + (b^2*log((2*b^2)/((a + b)^(5/2)*(b - a)^(1/2)) - (2
*b^2*exp(x))/(a*b^2 - a^2*b - a^3 + b^3)))/((a + b)^(3/2)*(b - a)^(3/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.74

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx$$

$$= \frac{-4e^x \sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b^2 + e^{2x} a^3 - e^{2x} a^2 b - e^{2x} a b^2 + e^{2x} b^3 - a^3 - a^2 b + a b^2 + b^3}{2e^x (a^4 - 2a^2 b^2 + b^4)}$$

input `int(cosh(x)/(a+b*tanh(x)),x)`output `( - 4*e**x*sqrt(a**2 - b**2)*atan((e**x*a + e**x*b)/sqrt(a**2 - b**2))*b**2 + e**(2*x)*a**3 - e**(2*x)*a**2*b - e**(2*x)*a*b**2 + e**(2*x)*b**3 - a**3 - a**2*b + a*b**2 + b**3)/(2*e**x*(a**4 - 2*a**2*b**2 + b**4))`



### 3.115 $\int \frac{\cosh^3(x)}{a+b \tanh(x)} dx$

Optimal result	896
Mathematica [A] (verified)	896
Rubi [C] (verified)	897
Maple [A] (verified)	900
Fricas [B] (verification not implemented)	901
Sympy [F]	902
Maxima [F(-2)]	902
Giac [A] (verification not implemented)	902
Mupad [B] (verification not implemented)	903
Reduce [B] (verification not implemented)	903

#### Optimal result

Integrand size = 13, antiderivative size = 132

$$\int \frac{\cosh^3(x)}{a+b \tanh(x)} dx = \frac{b^4 \arctan\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{5/2}} + \frac{b^3 \cosh(x)}{(a^2-b^2)^2} - \frac{b \cosh^3(x)}{3(a^2-b^2)} - \frac{ab^2 \sinh(x)}{(a^2-b^2)^2} + \frac{a \sinh(x)}{a^2-b^2} + \frac{a \sinh^3(x)}{3(a^2-b^2)}$$

output `b^4*arctan(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/(a^2-b^2)^(5/2)+b^3*cosh(x)/(a^2-b^2)^2-b*cosh(x)^3/(3*a^2-3*b^2)-a*b^2*sinh(x)/(a^2-b^2)^2+a*sinh(x)/(a^2-b^2)+a*sinh(x)^3/(3*a^2-3*b^2)`

#### Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.95

$$\int \frac{\cosh^3(x)}{a+b \tanh(x)} dx = \frac{24b^4 \sqrt{a+b} \arctan\left(\frac{b+a \tanh(\frac{x}{2})}{\sqrt{a-b} \sqrt{a+b}}\right) - 3\sqrt{a-b} b(a^3 + a^2b - 5ab^2 - 5b^3) \cosh(x) - (a-b)^{3/2} b(a+b)^2 \cosh^3(x)}{\dots}$$

input `Integrate[Cosh[x]^3/(a + b*Tanh[x]),x]`

output  $(24*b^4*\text{Sqrt}[a + b]*\text{ArcTan}[(b + a*\text{Tanh}[x/2])/(\text{Sqrt}[a - b]*\text{Sqrt}[a + b])] - 3*\text{Sqrt}[a - b]*b*(a^3 + a^2*b - 5*a*b^2 - 5*b^3)*\text{Cosh}[x] - (a - b)^{(3/2)}*b*(a + b)^2*\text{Cosh}[3*x] + 9*a^4*\text{Sqrt}[a - b]*\text{Sinh}[x] + 9*a^3*\text{Sqrt}[a - b]*b*\text{Sinh}[x] - 21*a^2*\text{Sqrt}[a - b]*b^2*\text{Sinh}[x] - 21*a*\text{Sqrt}[a - b]*b^3*\text{Sinh}[x] + a^4*\text{Sqrt}[a - b]*\text{Sinh}[3*x] + a^3*\text{Sqrt}[a - b]*b*\text{Sinh}[3*x] - a^2*\text{Sqrt}[a - b]*b^2*\text{Sinh}[3*x] - a*\text{Sqrt}[a - b]*b^3*\text{Sinh}[3*x])/(12*(a - b)^{(5/2)}*(a + b)^3)$

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.95, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.077$ , Rules used = {3042, 3990, 3042, 3967, 3042, 3113, 2009, 3990, 3042, 3967, 3042, 3117, 3988, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cosh^3(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sec(ix)^3(a - ib \tan(ix))} dx \\ & \quad \downarrow \text{3990} \\ & \frac{\int \cosh^3(x)(a - b \tanh(x)) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\cosh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{a + ib \tan(ix)}{\sec(ix)^3} dx}{a^2 - b^2} - \frac{b^2 \int \frac{1}{\sec(ix)(a - ib \tan(ix))} dx}{a^2 - b^2} \\ & \quad \downarrow \text{3967} \end{aligned}$$

$$\begin{aligned}
& \frac{a \int \cosh^3(x) dx - \frac{1}{3} b \cosh^3(x)}{a^2 - b^2} - \frac{b^2 \int \frac{1}{\sec(ix)(a-ib \tan(ix))} dx}{a^2 - b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + a \int \sin(ix + \frac{\pi}{2})^3 dx}{a^2 - b^2} - \frac{b^2 \int \frac{1}{\sec(ix)(a-ib \tan(ix))} dx}{a^2 - b^2} \\
& \quad \downarrow \text{3113} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia \int (\sinh^2(x) + 1) d(-i \sinh(x))}{a^2 - b^2} - \frac{b^2 \int \frac{1}{\sec(ix)(a-ib \tan(ix))} dx}{a^2 - b^2} \\
& \quad \downarrow \text{2009} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} - \frac{b^2 \int \frac{1}{\sec(ix)(a-ib \tan(ix))} dx}{a^2 - b^2} \\
& \quad \downarrow \text{3990} \\
& - \frac{b^2 \left( \frac{\int \cosh(x)(a-b \tanh(x)) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx}{a^2 - b^2} \right)}{a^2 - b^2} + \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} - \frac{b^2 \left( \frac{\int \frac{a+ib \tan(ix)}{\sec(ix)} dx}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \right)}{a^2 - b^2} \\
& \quad \downarrow \text{3967} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} - \frac{b^2 \left( \frac{a \int \cosh(x) dx - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \right)}{a^2 - b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} - \frac{b^2 \left( \frac{-b \cosh(x) + a \int \sin(ix + \frac{\pi}{2}) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \right)}{a^2 - b^2} \\
& \quad \downarrow \text{3117} \\
& \frac{-\frac{1}{3} b \cosh^3(x) + ia(-\frac{1}{3} i \sinh^3(x) - i \sinh(x))}{a^2 - b^2} - \frac{b^2 \left( \frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \int \frac{\sec(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \right)}{a^2 - b^2}
\end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3988} \\
& \frac{-\frac{1}{3}b \cosh^3(x) + ia\left(-\frac{1}{3}i \sinh^3(x) - i \sinh(x)\right)}{a^2 - b^2} - \\
& \frac{b^2 \left( \frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{ib^2 \int \frac{1}{a^2 - b^2 + \cosh^2(x)(b + a \tanh(x))^2} d(-i \cosh(x)(b + a \tanh(x)))}{a^2 - b^2} \right)}{a^2 - b^2} \\
& \downarrow \text{219} \\
& - \frac{b^2 \left( \frac{a \sinh(x) - b \cosh(x)}{a^2 - b^2} - \frac{b^2 \arctan\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{3/2}} \right)}{a^2 - b^2} + \\
& \frac{-\frac{1}{3}b \cosh^3(x) + ia\left(-\frac{1}{3}i \sinh^3(x) - i \sinh(x)\right)}{a^2 - b^2}
\end{aligned}$$

input `Int[Cosh[x]^3/(a + b*Tanh[x]),x]`

output `-((b^2*(-((b^2*ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]])/(a^2 - b^2)^(3/2)) + (-b*Cosh[x]) + a*Sinh[x])/(a^2 - b^2)))/(a^2 - b^2) + (-1/3*(b*Cosh[x]^3) + I*a*((-I)*Sinh[x] - (I/3)*Sinh[x]^3))/(a^2 - b^2)`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3967 `Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Simp[a Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] || NeQ[a^2 + b^2, 0])`

rule 3988 `Int[sec[(e_.) + (f_.)*(x_)]/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[-f^(-1) Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]`

rule 3990 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[Sec[e + f*x]^m*(a - b*Tan[e + f*x]), x], x] + Simp[b^2/(a^2 + b^2) Int[Sec[e + f*x]^(m + 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[(m - 1)/2, 0]`

## Maple [A] (verified)

Time = 4.12 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.30

method	result
risch	$\frac{e^{3x}}{24a+24b} + \frac{3e^x a}{8(a+b)^2} + \frac{5e^x b}{8(a+b)^2} - \frac{3e^{-x} a}{8(a-b)^2} + \frac{5e^{-x} b}{8(a-b)^2} - \frac{e^{-3x}}{24(a-b)} - \frac{b^4 \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)^2(a-b)^2} + \frac{b^4 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)^2(a-b)^2}$
default	$-\frac{2}{3(1+\tanh(\frac{x}{2}))^3(2a-2b)} + \frac{1}{(2a-2b)(1+\tanh(\frac{x}{2}))^2} - \frac{2a-3b}{2(a-b)^2(1+\tanh(\frac{x}{2}))} - \frac{2}{3(\tanh(\frac{x}{2})-1)^3(2b+2a)} - \frac{1}{(2b+2a)(\tanh(\frac{x}{2}))}$

input `int(cosh(x)^3/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

output

```
1/24/(a+b)*exp(x)^3+3/8/(a+b)^2*exp(x)*a+5/8/(a+b)^2*exp(x)*b-3/8/(a-b)^2/
exp(x)*a+5/8/(a-b)^2/exp(x)*b-1/24/(a-b)/exp(x)^3-1/(-a^2+b^2)^(1/2)*b^4/(
a+b)^2/(a-b)^2*ln(exp(x)-(a-b)/(-a^2+b^2)^(1/2))+1/(-a^2+b^2)^(1/2)*b^4/(a
+b)^2/(a-b)^2*ln(exp(x)+(a-b)/(-a^2+b^2)^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 908 vs.  $2(124) = 248$ .

Time = 0.13 (sec) , antiderivative size = 1871, normalized size of antiderivative = 14.17

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
[1/24*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 + 6*(
a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^5 + (a^
5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^6 - a^5 - a^4*b +
2*a^3*b^2 + 2*a^2*b^3 - a*b^4 - b^5 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^
2*b^3 + 7*a*b^4 - 5*b^5)*cosh(x)^4 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^
2*b^3 + 7*a*b^4 - 5*b^5 + 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 -
b^5)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a
*b^4 - b^5)*cosh(x)^3 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^
4 - 5*b^5)*cosh(x))*sinh(x)^3 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3
+ 7*a*b^4 + 5*b^5)*cosh(x)^2 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 +
7*a*b^4 + 5*b^5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*c
osh(x)^4 - 6*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*co
sh(x)^2)*sinh(x)^2 - 24*(b^4*cosh(x)^3 + 3*b^4*cosh(x)^2*sinh(x) + 3*b^4*c
osh(x)*sinh(x)^2 + b^4*sinh(x)^3)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2
+ 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh
(x) + sinh(x)) - a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (
a + b)*sinh(x)^2 + a - b)) + 6*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b
^4 - b^5)*cosh(x)^5 + 2*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4
- 5*b^5)*cosh(x)^3 - (3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 + 5
*b^5)*cosh(x))*sinh(x))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^3 ...
```

**Sympy [F]**

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \int \frac{\cosh^3(x)}{a + b \tanh(x)} dx$$

input `integrate(cosh(x)**3/(a+b*tanh(x)),x)`

output `Integral(cosh(x)**3/(a + b*tanh(x)), x)`

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see `assume?` f or more de`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.23

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \frac{2b^4 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}} - \frac{(9ae^{2x} - 15be^{2x} + a - b)e^{-3x}}{24(a^2 - 2ab + b^2)} + \frac{a^2e^{3x} + 2abe^{3x} + b^2e^{3x} + 9a^2e^x + 24abe^x + 15b^2e^x}{24(a^3 + 3a^2b + 3ab^2 + b^3)}$$

input `integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="giac")`

output

```
2*b^4*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/((a^4 - 2*a^2*b^2 + b^4)*sqrt(a^2 - b^2)) - 1/24*(9*a*e^(2*x) - 15*b*e^(2*x) + a - b)*e^(-3*x)/(a^2 - 2*a*b + b^2) + 1/24*(a^2*e^(3*x) + 2*a*b*e^(3*x) + b^2*e^(3*x) + 9*a^2*e^x + 24*a*b*e^x + 15*b^2*e^x)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3)
```

**Mupad [B] (verification not implemented)**

Time = 3.13 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.67

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \frac{e^{3x}}{24a + 24b} - \frac{e^{-3x}}{24a - 24b} - \frac{e^{-x}(3a - 5b)}{8(a - b)^2} + \frac{e^x(3a + 5b)}{8(a + b)^2} - \frac{b^4 \ln\left(-\frac{2b^4 e^x}{a^5 + a^4 b - 2a^3 b^2 - 2a^2 b^3 + a b^4 + b^5} - \frac{2b^4}{(a+b)^{7/2}(b-a)^{3/2}}\right)}{(a+b)^{5/2}(b-a)^{5/2}} + \frac{b^4 \ln\left(\frac{2b^4}{(a+b)^{7/2}(b-a)^{3/2}} - \frac{2b^4 e^x}{a^5 + a^4 b - 2a^3 b^2 - 2a^2 b^3 + a b^4 + b^5}\right)}{(a+b)^{5/2}(b-a)^{5/2}}$$

input

```
int(cosh(x)^3/(a + b*tanh(x)), x)
```

output

```
exp(3*x)/(24*a + 24*b) - exp(-3*x)/(24*a - 24*b) - (exp(-x)*(3*a - 5*b))/(8*(a - b)^2) + (exp(x)*(3*a + 5*b))/(8*(a + b)^2) - (b^4*log(-(2*b^4*exp(x))/(a*b^4 + a^4*b + a^5 + b^5 - 2*a^2*b^3 - 2*a^3*b^2) - (2*b^4)/((a + b)^(7/2)*(b - a)^(3/2))))/((a + b)^(5/2)*(b - a)^(5/2)) + (b^4*log((2*b^4)/((a + b)^(7/2)*(b - a)^(3/2)) - (2*b^4*exp(x))/(a*b^4 + a^4*b + a^5 + b^5 - 2*a^2*b^3 - 2*a^3*b^2)))/((a + b)^(5/2)*(b - a)^(5/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.44

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx = \frac{48e^{3x}\sqrt{a^2 - b^2} \operatorname{atan}\left(\frac{e^x a + e^x b}{\sqrt{a^2 - b^2}}\right) b^4 - 3e^{4x} a^4 b + 18e^{4x} a^2 b^3 - 3e^{2x} a^4 b + 18e^{2x} a^2 b^3 + 2e^{6x} a^2 b^3 - a^5 + 2a^3 b^2}{\dots}$$



input `int(cosh(x)^3/(a+b*tanh(x)),x)`

output `(48*exp(3*x)*sqrt(a**2 - b**2)*atan((exp(x)*a + exp(x)*b)/sqrt(a**2 - b**2))*b**4 + exp(6*x)*a**5 - exp(6*x)*a**4*b - 2*exp(6*x)*a**3*b**2 + 2*exp(6*x)*a**2*b**3 + exp(6*x)*a*b**4 - exp(6*x)*b**5 + 9*exp(4*x)*a**5 - 3*exp(4*x)*a**4*b - 30*exp(4*x)*a**3*b**2 + 18*exp(4*x)*a**2*b**3 + 21*exp(4*x)*a*b**4 - 15*exp(4*x)*b**5 - 9*exp(2*x)*a**5 - 3*exp(2*x)*a**4*b + 30*exp(2*x)*a**3*b**2 + 18*exp(2*x)*a**2*b**3 - 21*exp(2*x)*a*b**4 - 15*exp(2*x)*b**5 - a**5 - a**4*b + 2*a**3*b**2 + 2*a**2*b**3 - a*b**4 - b**5)/(24*exp(3*x)*(a**6 - 3*a**4*b**2 + 3*a**2*b**4 - b**6))`

### 3.116 $\int \frac{\tanh^5(x)}{1+\tanh(x)} dx$

Optimal result . . . . .	905
Mathematica [A] (verified) . . . . .	905
Rubi [C] (verified) . . . . .	906
Maple [A] (verified) . . . . .	910
Fricas [B] (verification not implemented) . . . . .	910
Sympy [B] (verification not implemented) . . . . .	911
Maxima [A] (verification not implemented) . . . . .	912
Giac [A] (verification not implemented) . . . . .	912
Mupad [B] (verification not implemented) . . . . .	913
Reduce [B] (verification not implemented) . . . . .	913

#### Optimal result

Integrand size = 11, antiderivative size = 43

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{5x}{2} - 2 \log(\cosh(x)) - \frac{5 \tanh(x)}{2} + \tanh^2(x) - \frac{5 \tanh^3(x)}{6} + \frac{\tanh^4(x)}{2(1 + \tanh(x))}$$

output

`5/2*x-2*ln(cosh(x))-5/2*tanh(x)+tanh(x)^2-5/6*tanh(x)^3+tanh(x)^4/(2+2*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.21

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{-12 \log(\cosh(x)) - 3(5 + 4 \log(\cosh(x))) \tanh(x) - 9 \tanh^2(x) + \tanh^3(x) - 2 \tanh^4(x) + 15 \operatorname{arctanh}(\tanh(x))}{6(1 + \tanh(x))}$$

input

`Integrate[Tanh[x]^5/(1 + Tanh[x]), x]`

output

```
(-12*Log[Cosh[x]] - 3*(5 + 4*Log[Cosh[x]])*Tanh[x] - 9*Tanh[x]^2 + Tanh[x]^3 - 2*Tanh[x]^4 + 15*ArcTanh[Tanh[x]]*(1 + Tanh[x]))/(6*(1 + Tanh[x]))
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.63, number of steps used = 21, number of rules used = 21,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.909$ , Rules used = {3042, 26, 4033, 26, 3042, 26, 4011, 25, 26, 3042, 25, 4011, 26, 26, 3042, 26, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^5(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)^5}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)^5}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4033} \\
 & -i \left( \frac{1}{2} \int -i(4 - 5 \tanh(x)) \tanh^3(x) dx + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} - \frac{1}{2} i \int (4 - 5 \tanh(x)) \tanh^3(x) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} - \frac{1}{2} i \int i(5i \tan(ix) + 4) \tan(ix)^3 dx \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\begin{aligned}
& -i \left( \frac{1}{2} \int (5i \tan(ix) + 4) \tan(ix)^3 dx + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 4011 \\
& -i \left( \frac{1}{2} \left( \int -((4i \tanh(x) - 5i) \tanh^2(x)) dx - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 25 \\
& -i \left( \frac{1}{2} \left( - \int -i(5 - 4 \tanh(x)) \tanh^2(x) dx - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( i \int (5 - 4 \tanh(x)) \tanh^2(x) dx - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 3042 \\
& -i \left( \frac{1}{2} \left( i \int -((4i \tan(ix) + 5) \tan(ix)^2) dx - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 25 \\
& -i \left( \frac{1}{2} \left( -i \int (4i \tan(ix) + 5) \tan(ix)^2 dx - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 4011 \\
& -i \left( \frac{1}{2} \left( -i \left( -2 \tanh^2(x) + \int i(5i \tanh(x) - 4i) \tanh(x) dx \right) - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( -i \left( -2 \tanh^2(x) + i \int -i(4 - 5 \tanh(x)) \tanh(x) dx \right) - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( -i \left( \int (4 - 5 \tanh(x)) \tanh(x) dx - 2 \tanh^2(x) \right) - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 3042 \\
& -i \left( \frac{1}{2} \left( -i \left( -2 \tanh^2(x) + \int -i(5i \tan(ix) + 4) \tan(ix) dx \right) - \frac{5}{3} i \tanh^3(x) \right) + \frac{i \tanh^4(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26
\end{aligned}$$

$$-i\left(\frac{1}{2}\left(-i\left(-2\tanh^2(x) - i\int(5i\tan(ix) + 4)\tan(ix)dx\right) - \frac{5}{3}i\tanh^3(x)\right) + \frac{i\tanh^4(x)}{2(\tanh(x) + 1)}\right)$$

↓ 4008

$$-i\left(\frac{1}{2}\left(-i\left(-2\tanh^2(x) - i(4\int i\tanh(x)dx - 5ix + 5i\tanh(x))\right) - \frac{5}{3}i\tanh^3(x)\right) + \frac{i\tanh^4(x)}{2(\tanh(x) + 1)}\right)$$

↓ 26

$$-i\left(\frac{1}{2}\left(-i\left(-2\tanh^2(x) - i(4i\int \tanh(x)dx - 5ix + 5i\tanh(x))\right) - \frac{5}{3}i\tanh^3(x)\right) + \frac{i\tanh^4(x)}{2(\tanh(x) + 1)}\right)$$

↓ 3042

$$-i\left(\frac{1}{2}\left(-i\left(-2\tanh^2(x) - i(4i\int -i\tan(ix)dx - 5ix + 5i\tanh(x))\right) - \frac{5}{3}i\tanh^3(x)\right) + \frac{i\tanh^4(x)}{2(\tanh(x) + 1)}\right)$$

↓ 26

$$-i\left(\frac{1}{2}\left(-i\left(-2\tanh^2(x) - i(4\int \tan(ix)dx - 5ix + 5i\tanh(x))\right) - \frac{5}{3}i\tanh^3(x)\right) + \frac{i\tanh^4(x)}{2(\tanh(x) + 1)}\right)$$

↓ 3956

$$-i\left(\frac{i\tanh^4(x)}{2(\tanh(x) + 1)} + \frac{1}{2}\left(-\frac{5}{3}i\tanh^3(x) - i(-2\tanh^2(x) - i(-5ix + 5i\tanh(x) + 4i\log(\cosh(x))))\right)\right)$$

input `Int [Tanh[x]^5/(1 + Tanh[x]), x]`

output `(-I)*(((I/2)*Tanh[x]^4)/(1 + Tanh[x]) + (((-5*I)/3)*Tanh[x]^3 - I*((-I)*((-5*I)*x + (4*I)*Log[Cosh[x]] + (5*I)*Tanh[x]) - 2*Tanh[x]^2))/2)`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 26  $\text{Int}[(\text{Complex}[0, \text{a}_])*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], \text{a}]) \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2, 1]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3956  $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x\_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d}*x], \text{x}]]/\text{d}, \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4008  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]*(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{a}*c - \text{b}*d)*x, \text{x}] + (\text{Simp}[\text{b}*d*(\text{Tan}[\text{e} + \text{f}*x]/\text{f}), \text{x}] + \text{Simp}[(\text{b}*c + \text{a}*d) \quad \text{Int}[\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{b}*c + \text{a}*d, 0]$
- rule 4011  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_.}*(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{d}*(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}/(\text{f}*\text{m}), \text{x}] + \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} - 1}*\text{Simp}[\text{a}*c - \text{b}*d + (\text{b}*c + \text{a}*d)*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{GtQ}[\text{m}, 0]$
- rule 4033  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{n}_.}/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b}*c - \text{a}*d)*((\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} - 1}/(2*\text{a}*f*(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])), \text{x}] + \text{Simp}[1/(2*\text{a}^2) \quad \text{Int}[(\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} - 2}*\text{Simp}[\text{a}*c^2 + \text{a}*d^2*(\text{n} - 1) - \text{b}*c*d*\text{n} - \text{d}*(\text{a}*c*(\text{n} - 2) + \text{b}*d*\text{n})*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{EqQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{GtQ}[\text{n}, 1]$

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$-\frac{\tanh(x)^3}{3} + \frac{\tanh(x)^2}{2} - 2 \tanh(x) + \frac{1}{2+2 \tanh(x)} + \frac{9 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	40
default	$-\frac{\tanh(x)^3}{3} + \frac{\tanh(x)^2}{2} - 2 \tanh(x) + \frac{1}{2+2 \tanh(x)} + \frac{9 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	40
risch	$\frac{9x}{2} + \frac{e^{-2x}}{4} + \frac{4 e^{4x} + 6 e^{2x} + \frac{14}{3}}{(e^{2x}+1)^3} - 2 \ln(e^{2x} + 1)$	44
parallelrisch	$-\frac{2 \tanh(x)^4 - 15 - \tanh(x)^3 - 12 \ln(1 - \tanh(x)) \tanh(x) - 27 \tanh(x)x + 9 \tanh(x)^2 - 12 \ln(1 - \tanh(x)) - 27x}{6(1 + \tanh(x))}$	57

input `int(tanh(x)^5/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-1/3*tanh(x)^3+1/2*tanh(x)^2-2*tanh(x)+1/2/(1+tanh(x))+9/4*ln(1+tanh(x))-1/4*ln(tanh(x)-1)`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 571 vs.  $2(35) = 70$ .

Time = 0.09 (sec) , antiderivative size = 571, normalized size of antiderivative = 13.28

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="fricas")`

output

```

1/12*(54*x*cosh(x)^8 + 432*x*cosh(x)*sinh(x)^7 + 54*x*sinh(x)^8 + 3*(54*x
+ 17)*cosh(x)^6 + 3*(504*x*cosh(x)^2 + 54*x + 17)*sinh(x)^6 + 18*(168*x*co
sh(x)^3 + (54*x + 17)*cosh(x))*sinh(x)^5 + 81*(2*x + 1)*cosh(x)^4 + 9*(420
*x*cosh(x)^4 + 5*(54*x + 17)*cosh(x)^2 + 18*x + 9)*sinh(x)^4 + 12*(252*x*c
osh(x)^5 + 5*(54*x + 17)*cosh(x)^3 + 27*(2*x + 1)*cosh(x))*sinh(x)^3 + (54
*x + 65)*cosh(x)^2 + (1512*x*cosh(x)^6 + 45*(54*x + 17)*cosh(x)^4 + 486*(2
*x + 1)*cosh(x)^2 + 54*x + 65)*sinh(x)^2 - 24*(cosh(x)^8 + 8*cosh(x)*sinh(
x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 + 2*(28*cosh
(x)^3 + 9*cosh(x))*sinh(x)^5 + (70*cosh(x)^4 + 45*cosh(x)^2 + 3)*sinh(x)^4
+ 3*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + (
28*cosh(x)^6 + 45*cosh(x)^4 + 18*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*
(4*cosh(x)^7 + 9*cosh(x)^5 + 6*cosh(x)^3 + cosh(x))*sinh(x))*log(2*cosh(x)
/(cosh(x) - sinh(x))) + 2*(216*x*cosh(x)^7 + 9*(54*x + 17)*cosh(x)^5 + 162
*(2*x + 1)*cosh(x)^3 + (54*x + 65)*cosh(x))*sinh(x) + 3)/(cosh(x)^8 + 8*co
sh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 + 3)*sinh(x)^6 + 3*cosh(x)^6 +
2*(28*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + (70*cosh(x)^4 + 45*cosh(x)^2 + 3
)*sinh(x)^4 + 3*cosh(x)^4 + 4*(14*cosh(x)^5 + 15*cosh(x)^3 + 3*cosh(x))*si
nh(x)^3 + (28*cosh(x)^6 + 45*cosh(x)^4 + 18*cosh(x)^2 + 1)*sinh(x)^2 + cos
h(x)^2 + 2*(4*cosh(x)^7 + 9*cosh(x)^5 + 6*cosh(x)^3 + cosh(x))*sinh(x))

```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs.  $2(42) = 84$ .

Time = 0.21 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.42

$$\begin{aligned}
 \int \frac{\tanh^5(x)}{1 + \tanh(x)} dx &= \frac{3x \tanh(x)}{6 \tanh(x) + 6} + \frac{3x}{6 \tanh(x) + 6} + \frac{12 \log(\tanh(x) + 1) \tanh(x)}{6 \tanh(x) + 6} \\
 &+ \frac{12 \log(\tanh(x) + 1)}{6 \tanh(x) + 6} - \frac{2 \tanh^4(x)}{6 \tanh(x) + 6} \\
 &+ \frac{\tanh^3(x)}{6 \tanh(x) + 6} - \frac{9 \tanh^2(x)}{6 \tanh(x) + 6} + \frac{15}{6 \tanh(x) + 6}
 \end{aligned}$$

input

```
integrate(tanh(x)**5/(1+tanh(x)), x)
```



output

```
3*x*tanh(x)/(6*tanh(x) + 6) + 3*x/(6*tanh(x) + 6) + 12*log(tanh(x) + 1)*tanh(x)/(6*tanh(x) + 6) + 12*log(tanh(x) + 1)/(6*tanh(x) + 6) - 2*tanh(x)**4/(6*tanh(x) + 6) + tanh(x)**3/(6*tanh(x) + 6) - 9*tanh(x)**2/(6*tanh(x) + 6) + 15/(6*tanh(x) + 6)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.28

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{1}{2}x - \frac{2(15e^{-2x} + 12e^{-4x} + 7)}{3(3e^{-2x} + 3e^{-4x} + e^{-6x} + 1)} + \frac{1}{4}e^{-2x} - 2 \log(e^{-2x} + 1)$$

input

```
integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="maxima")
```

output

```
1/2*x - 2/3*(15*e^(-2*x) + 12*e^(-4*x) + 7)/(3*e^(-2*x) + 3*e^(-4*x) + e^(-6*x) + 1) + 1/4*e^(-2*x) - 2*log(e^(-2*x) + 1)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{9}{2}x + \frac{(51e^{6x} + 81e^{4x} + 65e^{2x} + 3)e^{-2x}}{12(e^{2x} + 1)^3} - 2 \log(e^{2x} + 1)$$

input

```
integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="giac")
```

output

```
9/2*x + 1/12*(51*e^(6*x) + 81*e^(4*x) + 65*e^(2*x) + 3)*e^(-2*x)/(e^(2*x) + 1)^3 - 2*log(e^(2*x) + 1)
```

**Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{x}{2} + 2 \ln(\tanh(x) + 1) - 2 \tanh(x) + \frac{\tanh(x)^2}{2} - \frac{\tanh(x)^3}{3} + \frac{1}{2(\tanh(x) + 1)}$$

input `int(tanh(x)^5/(tanh(x) + 1),x)`output `x/2 + 2*log(tanh(x) + 1) - 2*tanh(x) + tanh(x)^2/2 - tanh(x)^3/3 + 1/(2*(tanh(x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx = \frac{24e^{2x} \log(\tanh(x) + 1) - 4e^{2x} \tanh(x)^3 + 6e^{2x} \tanh(x)^2 - 24e^{2x} \tanh(x) + 6e^{2x} x + 3}{12e^{2x}}$$

input `int(tanh(x)^5/(1+tanh(x)),x)`output `(24*e**(2*x)*log(tanh(x) + 1) - 4*e**(2*x)*tanh(x)**3 + 6*e**(2*x)*tanh(x)**2 - 24*e**(2*x)*tanh(x) + 6*e**(2*x)*x + 3)/(12*e**(2*x))`

### 3.117 $\int \frac{\tanh^4(x)}{1+\tanh(x)} dx$

Optimal result	914
Mathematica [A] (verified)	914
Rubi [C] (verified)	915
Maple [A] (verified)	918
Fricas [B] (verification not implemented)	918
Sympy [B] (verification not implemented)	919
Maxima [A] (verification not implemented)	920
Giac [A] (verification not implemented)	920
Mupad [B] (verification not implemented)	920
Reduce [B] (verification not implemented)	921

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = -\frac{3x}{2} + 2 \log(\cosh(x)) + \frac{3 \tanh(x)}{2} - \tanh^2(x) + \frac{\tanh^3(x)}{2(1 + \tanh(x))}$$

output

```
-3/2*x+2*ln(cosh(x))+3/2*tanh(x)-tanh(x)^2+tanh(x)^3/(2+2*tanh(x))
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = \frac{4 \log(\cosh(x)) + (3 + 4 \log(\cosh(x))) \tanh(x) + \tanh^2(x) - \tanh^3(x) - 3 \operatorname{arctanh}(\tanh(x))(1 + \tanh(x))}{2(1 + \tanh(x))}$$

input

```
Integrate[Tanh[x]^4/(1 + Tanh[x]),x]
```

output

```
(4*Log[Cosh[x]] + (3 + 4*Log[Cosh[x]])*Tanh[x] + Tanh[x]^2 - Tanh[x]^3 - 3 *ArcTanh[Tanh[x]]*(1 + Tanh[x]))/(2*(1 + Tanh[x]))
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.32, number of steps used = 15, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.364$ , Rules used = {3042, 4033, 25, 3042, 25, 4011, 26, 26, 3042, 26, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^4(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(ix)^4}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4033} \\
 & \frac{1}{2} \int -((3 - 4 \tanh(x)) \tanh^2(x)) dx + \frac{\tanh^3(x)}{2(\tanh(x) + 1)} \\
 & \quad \downarrow \text{25} \\
 & \frac{\tanh^3(x)}{2(\tanh(x) + 1)} - \frac{1}{2} \int (3 - 4 \tanh(x)) \tanh^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh^3(x)}{2(\tanh(x) + 1)} - \frac{1}{2} \int -((4i \tan(ix) + 3) \tan(ix)^2) dx \\
 & \quad \downarrow \text{25} \\
 & \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \int (4i \tan(ix) + 3) \tan(ix)^2 dx \\
 & \quad \downarrow \text{4011} \\
 & \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) + \int i(3i \tanh(x) - 4i) \tanh(x) dx \right) \\
 & \quad \downarrow \text{26} \\
 & \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) + i \int -i(4 - 3 \tanh(x)) \tanh(x) dx \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& \frac{1}{2} \left( \int (4 - 3 \tanh(x)) \tanh(x) dx - 2 \tanh^2(x) \right) + \frac{\tanh^3(x)}{2(\tanh(x) + 1)} \\
& \downarrow 3042 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) + \int -i(3i \tan(ix) + 4) \tan(ix) dx \right) \\
& \downarrow 26 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) - i \int (3i \tan(ix) + 4) \tan(ix) dx \right) \\
& \downarrow 4008 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) - i(4 \int i \tanh(x) dx - 3ix + 3i \tanh(x)) \right) \\
& \downarrow 26 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) - i(4i \int \tanh(x) dx - 3ix + 3i \tanh(x)) \right) \\
& \downarrow 3042 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) - i(4i \int -i \tan(ix) dx - 3ix + 3i \tanh(x)) \right) \\
& \downarrow 26 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( -2 \tanh^2(x) - i(4 \int \tan(ix) dx - 3ix + 3i \tanh(x)) \right) \\
& \downarrow 3956 \\
& \frac{\tanh^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} (-2 \tanh^2(x) - i(-3ix + 3i \tanh(x) + 4i \log(\cosh(x))))
\end{aligned}$$

input `Int [Tanh[x]^4/(1 + Tanh[x]), x]`

output `Tanh[x]^3/(2*(1 + Tanh[x])) + ((-I)*((-3*I)*x + (4*I)*Log[Cosh[x]] + (3*I)*Tanh[x]) - 2*Tanh[x]^2)/2`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 26  $\text{Int}[(\text{Complex}[0, \text{a}_])*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], \text{a}]) \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2, 1]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinear} \text{Q}[\text{u}, \text{x}]$
- rule 3956  $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x\_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d}*x], \text{x}]]/\text{d}, \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4008  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]*(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{a}*c - \text{b}*d)*x, \text{x}] + (\text{Simp}[\text{b}*d*(\text{Tan}[\text{e} + \text{f}*x]/\text{f}), \text{x}] + \text{Simp}[(\text{b}*c + \text{a}*d) \quad \text{Int}[\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{b}*c + \text{a}*d, 0]$
- rule 4011  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_.}*(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{d}*((\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}/(\text{f}*\text{m})), \text{x}] + \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} - 1}*\text{Simp}[\text{a}*c - \text{b}*d + (\text{b}*c + \text{a}*d)*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{GtQ}[\text{m}, 0]$
- rule 4033  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{n}_.}/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b}*c - \text{a}*d)*((\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} - 1}/(2*\text{a}*f*(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])), \text{x}] + \text{Simp}[1/(2*\text{a}^2) \quad \text{Int}[(\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} - 2}*\text{Simp}[\text{a}*c^2 + \text{a}*d^2*(\text{n} - 1) - \text{b}*c*d*\text{n} - \text{d}*(\text{a}*c*(\text{n} - 2) + \text{b}*d*\text{n})*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{EqQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{GtQ}[\text{n}, 1]$

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

method	result	size
risch	$-\frac{7x}{2} - \frac{e^{-2x}}{4} - \frac{2}{(e^{2x}+1)^2} + 2 \ln(e^{2x} + 1)$	30
derivativedivides	$-\frac{\tanh(x)^2}{2} + \tanh(x) - \frac{1}{2(1+\tanh(x))} - \frac{7 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	32
default	$-\frac{\tanh(x)^2}{2} + \tanh(x) - \frac{1}{2(1+\tanh(x))} - \frac{7 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	32
parallelrisch	$-\frac{3+\tanh(x)^3+4 \ln(1-\tanh(x)) \tanh(x)+7 \tanh(x)x-\tanh(x)^2+4 \ln(1-\tanh(x))+7x}{2(1+\tanh(x))}$	49

input `int(tanh(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-7/2*x-1/4*exp(-2*x)-2/(exp(2*x)+1)^2+2*ln(exp(2*x)+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 354 vs. 2(31) = 62.

Time = 0.10 (sec) , antiderivative size = 354, normalized size of antiderivative = 9.57

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="fricas")`

output

```
-1/4*(14*x*cosh(x)^6 + 84*x*cosh(x)*sinh(x)^5 + 14*x*sinh(x)^6 + (28*x + 1)
)*cosh(x)^4 + (210*x*cosh(x)^2 + 28*x + 1)*sinh(x)^4 + 4*(70*x*cosh(x)^3 +
(28*x + 1)*cosh(x))*sinh(x)^3 + 2*(7*x + 5)*cosh(x)^2 + 2*(105*x*cosh(x)^
4 + 3*(28*x + 1)*cosh(x)^2 + 7*x + 5)*sinh(x)^2 - 8*(cosh(x)^6 + 6*cosh(x)
*sinh(x)^5 + sinh(x)^6 + (15*cosh(x)^2 + 2)*sinh(x)^4 + 2*cosh(x)^4 + 4*(5
*cosh(x)^3 + 2*cosh(x))*sinh(x)^3 + (15*cosh(x)^4 + 12*cosh(x)^2 + 1)*sinh
(x)^2 + cosh(x)^2 + 2*(3*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*sinh(x))*log(2
*cosh(x)/(cosh(x) - sinh(x))) + 4*(21*x*cosh(x)^5 + (28*x + 1)*cosh(x)^3 +
(7*x + 5)*cosh(x))*sinh(x) + 1)/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)
)^6 + (15*cosh(x)^2 + 2)*sinh(x)^4 + 2*cosh(x)^4 + 4*(5*cosh(x)^3 + 2*cosh
(x))*sinh(x)^3 + (15*cosh(x)^4 + 12*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 +
2*(3*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*sinh(x))
```

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs.  $2(34) = 68$ .

Time = 0.21 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.30

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = \frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2} - \frac{4 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{\tanh^3(x)}{2 \tanh(x) + 2} + \frac{\tanh^2(x)}{2 \tanh(x) + 2} - \frac{3}{2 \tanh(x) + 2}$$

input

```
integrate(tanh(x)**4/(1+tanh(x)),x)
```

output

```
x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) - 4*log(tanh(x) + 1)*tanh(x)
/(2*tanh(x) + 2) - 4*log(tanh(x) + 1)/(2*tanh(x) + 2) - tanh(x)**3/(2*tanh
(x) + 2) + tanh(x)**2/(2*tanh(x) + 2) - 3/(2*tanh(x) + 2)
```



**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = \frac{1}{2}x + \frac{2(2e^{(-2x)} + 1)}{2e^{(-2x)} + e^{(-4x)} + 1} - \frac{1}{4}e^{(-2x)} + 2 \log(e^{(-2x)} + 1)$$

input `integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="maxima")`output `1/2*x + 2*(2*e^(-2*x) + 1)/(2*e^(-2*x) + e^(-4*x) + 1) - 1/4*e^(-2*x) + 2*log(e^(-2*x) + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = -\frac{7}{2}x - \frac{(e^{4x} + 10e^{2x} + 1)e^{(-2x)}}{4(e^{2x} + 1)^2} + 2 \log(e^{2x} + 1)$$

input `integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="giac")`output `-7/2*x - 1/4*(e^(4*x) + 10*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) + 1)^2 + 2*log(e^(2*x) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx = \frac{x}{2} - 2 \ln(\tanh(x) + 1) + \tanh(x) - \frac{\tanh(x)^2}{2} - \frac{1}{2(\tanh(x) + 1)}$$

input `int(tanh(x)^4/(tanh(x) + 1),x)`output `x/2 - 2*log(tanh(x) + 1) + tanh(x) - tanh(x)^2/2 - 1/(2*(tanh(x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.38

$$\int \frac{\tanh^4(x)}{1 + \tanh(x)} dx$$
$$= \frac{-8e^{2x} \log(\tanh(x) + 1) - 2e^{2x} \tanh(x)^2 + 4e^{2x} \tanh(x) + 2e^{2x} x - 1}{4e^{2x}}$$

input `int(tanh(x)^4/(1+tanh(x)),x)`output `( - 8*e**(2*x)*log(tanh(x) + 1) - 2*e**(2*x)*tanh(x)**2 + 4*e**(2*x)*tanh(x) + 2*e**(2*x)*x - 1)/(4*e**(2*x))`

### 3.118 $\int \frac{\tanh^3(x)}{1+\tanh(x)} dx$

Optimal result	922
Mathematica [A] (verified)	922
Rubi [C] (verified)	923
Maple [A] (verified)	925
Fricas [B] (verification not implemented)	926
Sympy [B] (verification not implemented)	926
Maxima [A] (verification not implemented)	927
Giac [A] (verification not implemented)	927
Mupad [B] (verification not implemented)	927
Reduce [B] (verification not implemented)	928

#### Optimal result

Integrand size = 11, antiderivative size = 31

$$\int \frac{\tanh^3(x)}{1+\tanh(x)} dx = \frac{3x}{2} - \log(\cosh(x)) - \frac{3 \tanh(x)}{2} + \frac{\tanh^2(x)}{2(1+\tanh(x))}$$

output

```
3/2*x-ln(cosh(x))-3/2*tanh(x)+tanh(x)^2/(2+2*tanh(x))
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int \frac{\tanh^3(x)}{1+\tanh(x)} dx = \frac{2 \log(\cosh(x)) + (3 + 2 \log(\cosh(x))) \tanh(x) + 2 \tanh^2(x) - 3 \operatorname{arctanh}(\tanh(x))(1 + \tanh(x))}{2(1 + \tanh(x))}$$

input

```
Integrate[Tanh[x]^3/(1 + Tanh[x]), x]
```

output

```
-1/2*(2*Log[Cosh[x]] + (3 + 2*Log[Cosh[x]])*Tanh[x] + 2*Tanh[x]^2 - 3*ArcTanh[Tanh[x]]*(1 + Tanh[x]))/(1 + Tanh[x])
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 26, 4033, 26, 3042, 26, 4008, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^3(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \tan(ix)^3}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\tan(ix)^3}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4033} \\
 & i \left( \frac{1}{2} \int i(2 - 3 \tanh(x)) \tanh(x) dx - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( \frac{1}{2} i \int (2 - 3 \tanh(x)) \tanh(x) dx - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{3042} \\
 & i \left( \frac{1}{2} i \int -i(3i \tan(ix) + 2) \tan(ix) dx - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( \frac{1}{2} \int (3i \tan(ix) + 2) \tan(ix) dx - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{4008} \\
 & i \left( \frac{1}{2} (2 \int i \tanh(x) dx - 3ix + 3i \tanh(x)) - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 26 \\
& i \left( \frac{1}{2} (2i \int \tanh(x) dx - 3ix + 3i \tanh(x)) - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
& \downarrow 3042 \\
& i \left( \frac{1}{2} (2i \int -i \tan(ix) dx - 3ix + 3i \tanh(x)) - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
& \downarrow 26 \\
& i \left( \frac{1}{2} (2 \int \tan(ix) dx - 3ix + 3i \tanh(x)) - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right) \\
& \downarrow 3956 \\
& i \left( \frac{1}{2} (-3ix + 3i \tanh(x) + 2i \log(\cosh(x))) - \frac{i \tanh^2(x)}{2(\tanh(x) + 1)} \right)
\end{aligned}$$

input `Int [Tanh[x]^3/(1 + Tanh[x]), x]`

output `I*((( -3*I)*x + (2*I)*Log[Cosh[x]] + (3*I)*Tanh[x])/2 - ((I/2)*Tanh[x]^2)/(1 + Tanh[x]))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

rule 4033

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)/((a_) + (b_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((c + d*Tan[e + f*x])^(n - 1)/(
2*a*f*(a + b*Tan[e + f*x]))), x] + Simp[1/(2*a^2) Int[(c + d*Tan[e + f*x]
)^(n - 2)*Simp[a*c^2 + a*d^2*(n - 1) - b*c*d*n - d*(a*c*(n - 2) + b*d*n)*Ta
n[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[n, 1]
```

## Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$-\tanh(x) + \frac{1}{2+2\tanh(x)} + \frac{5\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	28
default	$-\tanh(x) + \frac{1}{2+2\tanh(x)} + \frac{5\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	28
risch	$\frac{5x}{2} + \frac{e^{-2x}}{4} + \frac{2}{e^{2x}+1} - \ln(e^{2x} + 1)$	30
parallelrisc	$\frac{3+2\ln(1-\tanh(x))\tanh(x)+5\tanh(x)x-2\tanh(x)^2+2\ln(1-\tanh(x))+5x}{2+2\tanh(x)}$	45

input

```
int(tanh(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-tanh(x)+1/2/(1+tanh(x))+5/4*ln(1+tanh(x))-1/4*ln(tanh(x)-1)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 186 vs.  $2(25) = 50$ .

Time = 0.09 (sec) , antiderivative size = 186, normalized size of antiderivative = 6.00

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{10x \cosh(x)^4 + 40x \cosh(x) \sinh(x)^3 + 10x \sinh(x)^4 + (10x + 9) \cosh(x)^2 + (60x \cosh(x)^2 + 10x + 9) \sinh(x)^2 - 4(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + (6 \cosh(x)^2 + 1) \sinh(x)^2 + \cosh(x)^2 + 2(2 \cosh(x)^3 + \cosh(x)) \sinh(x)) \log(2 \cosh(x) / (\cosh(x) - \sinh(x))) + 2(20x \cosh(x)^3 + (10x + 9) \cosh(x)) \sinh(x) + 1}{4(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + (6 \cosh(x)^2 + 1) \sinh(x)^2 + \cosh(x)^2 + 2(2 \cosh(x)^3 + \cosh(x)) \sinh(x))}$$

input `integrate(tanh(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output `1/4*(10*x*cosh(x)^4 + 40*x*cosh(x)*sinh(x)^3 + 10*x*sinh(x)^4 + (10*x + 9)*cosh(x)^2 + (60*x*cosh(x)^2 + 10*x + 9)*sinh(x)^2 - 4*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))*log(2*cosh(x)/(cosh(x) - sinh(x))) + 2*(20*x*cosh(x)^3 + (10*x + 9)*cosh(x))*sinh(x) + 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs.  $2(27) = 54$ .

Time = 0.20 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.42

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx = \frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} + \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2}$$

$$+ \frac{2 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{2 \tanh^2(x)}{2 \tanh(x) + 2} + \frac{3}{2 \tanh(x) + 2}$$

input `integrate(tanh(x)**3/(1+tanh(x)),x)`

output `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) + 2*log(tanh(x) + 1)*tanh(x)/(2*tanh(x) + 2) + 2*log(tanh(x) + 1)/(2*tanh(x) + 2) - 2*tanh(x)**2/(2*tanh(x) + 2) + 3/(2*tanh(x) + 2)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx = \frac{1}{2}x - \frac{2}{e^{(-2x)} + 1} + \frac{1}{4}e^{(-2x)} - \log(e^{(-2x)} + 1)$$

input `integrate(tanh(x)^3/(1+tanh(x)),x, algorithm="maxima")`output `1/2*x - 2/(e^(-2*x) + 1) + 1/4*e^(-2*x) - log(e^(-2*x) + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx = \frac{5}{2}x + \frac{(9e^{(2x)} + 1)e^{(-2x)}}{4(e^{(2x)} + 1)} - \log(e^{(2x)} + 1)$$

input `integrate(tanh(x)^3/(1+tanh(x)),x, algorithm="giac")`output `5/2*x + 1/4*(9*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) + 1) - log(e^(2*x) + 1)`**Mupad [B] (verification not implemented)**

Time = 2.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx = \frac{x}{2} + \ln(\tanh(x) + 1) - \tanh(x) + \frac{1}{2(\tanh(x) + 1)}$$

input `int(tanh(x)^3/(tanh(x) + 1),x)`output `x/2 + log(tanh(x) + 1) - tanh(x) + 1/(2*(tanh(x) + 1))`



**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx = \frac{4e^{2x} \log(\tanh(x) + 1) - 4e^{2x} \tanh(x) + 2e^{2x} x + 1}{4e^{2x}}$$

input `int(tanh(x)^3/(1+tanh(x)),x)`

output `(4*e**(2*x)*log(tanh(x) + 1) - 4*e**(2*x)*tanh(x) + 2*e**(2*x)*x + 1)/(4*e**(2*x))`

### 3.119 $\int \frac{\tanh^2(x)}{1+\tanh(x)} dx$

Optimal result	929
Mathematica [A] (verified)	929
Rubi [A] (verified)	930
Maple [A] (verified)	931
Fricas [B] (verification not implemented)	932
Sympy [B] (verification not implemented)	932
Maxima [A] (verification not implemented)	933
Giac [A] (verification not implemented)	933
Mupad [B] (verification not implemented)	933
Reduce [B] (verification not implemented)	934

#### Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = -\frac{x}{2} + \log(\cosh(x)) - \frac{1}{2(1 + \tanh(x))}$$

output `-1/2*x+ln(cosh(x))-1/(2+2*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{2 \log(\cosh(x)) + \tanh(x) + 2 \log(\cosh(x)) \tanh(x) - \operatorname{arctanh}(\tanh(x))(1 + \tanh(x))}{2(1 + \tanh(x))}$$

input `Integrate[Tanh[x]^2/(1 + Tanh[x]),x]`

output `(2*Log[Cosh[x]] + Tanh[x] + 2*Log[Cosh[x]]*Tanh[x] - ArcTanh[Tanh[x]]*(1 + Tanh[x]))/(2*(1 + Tanh[x]))`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.26, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3042, 25, 4023, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan(ix)^2}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan(ix)^2}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4023} \\
 & -\frac{1}{2} \int (1 - 2 \tanh(x)) dx - \frac{1}{2(\tanh(x) + 1)} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} (2 \log(\cosh(x)) - x) - \frac{1}{2(\tanh(x) + 1)}
 \end{aligned}$$

input `Int [Tanh[x]^2/(1 + Tanh[x]), x]`

output `(-x + 2*Log[Cosh[x]])/2 - 1/(2*(1 + Tanh[x]))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4023 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-b)*(a*c + b*d)^2*((a + b*Tan[e + f*x])^m/(2*a^3*f*m)), x] + Simp[1/(2*a^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c^2 - 2*b*c*d + a*d^2 - 2*b*d^2*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LeQ[m, -1] && EqQ[a^2 + b^2, 0]`

### Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
risch	$-\frac{3x}{2} - \frac{e^{-2x}}{4} + \ln(e^{2x} + 1)$	18
derivativedivides	$-\frac{\ln(\tanh(x)-1)}{4} - \frac{1}{2(1+\tanh(x))} - \frac{3\ln(1+\tanh(x))}{4}$	24
default	$-\frac{\ln(\tanh(x)-1)}{4} - \frac{1}{2(1+\tanh(x))} - \frac{3\ln(1+\tanh(x))}{4}$	24
parallelrisch	$-\frac{1+2\ln(1-\tanh(x))\tanh(x)+3\tanh(x)x+2\ln(1-\tanh(x))+3x}{2(1+\tanh(x))}$	39

input `int(tanh(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `-3/2*x-1/4*exp(-2*x)+ln(exp(2*x)+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 73 vs.  $2(15) = 30$ .

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 3.84

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{6x \cosh(x)^2 + 12x \cosh(x) \sinh(x) + 6x \sinh(x)^2 - 4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2) \log(2 \cosh(x) / (\cosh(x) - \sinh(x))) + 1}{4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)}$$

input `integrate(tanh(x)^2/(1+tanh(x)),x, algorithm="fricas")`

output `-1/4*(6*x*cosh(x)^2 + 12*x*cosh(x)*sinh(x) + 6*x*sinh(x)^2 - 4*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)*log(2*cosh(x)/(cosh(x) - sinh(x))) + 1)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 61 vs.  $2(15) = 30$ .

Time = 0.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 3.21

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2} - \frac{2 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{1}{2 \tanh(x) + 2}$$

input `integrate(tanh(x)**2/(1+tanh(x)),x)`

output `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) - 2*log(tanh(x) + 1)*tanh(x)/(2*tanh(x) + 2) - 2*log(tanh(x) + 1)/(2*tanh(x) + 2) - 1/(2*tanh(x) + 2)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{1}{2}x - \frac{1}{4}e^{(-2x)} + \log(e^{(-2x)} + 1)$$

input `integrate(tanh(x)^2/(1+tanh(x)),x, algorithm="maxima")`output `1/2*x - 1/4*e^(-2*x) + log(e^(-2*x) + 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = -\frac{3}{2}x - \frac{1}{4}e^{(-2x)} + \log(e^{(2x)} + 1)$$

input `integrate(tanh(x)^2/(1+tanh(x)),x, algorithm="giac")`output `-3/2*x - 1/4*e^(-2*x) + log(e^(2*x) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{x}{2} - \ln(\tanh(x) + 1) - \frac{1}{2(\tanh(x) + 1)}$$

input `int(tanh(x)^2/(tanh(x) + 1),x)`output `x/2 - log(tanh(x) + 1) - 1/(2*(tanh(x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.63

$$\int \frac{\tanh^2(x)}{1 + \tanh(x)} dx = \frac{-4e^{2x}\log(\tanh(x) + 1) + 2e^{2x}x - 1}{4e^{2x}}$$

input `int(tanh(x)^2/(1+tanh(x)),x)`

output `( - 4*e**(2*x)*log(tanh(x) + 1) + 2*e**(2*x)*x - 1)/(4*e**(2*x))`

### 3.120 $\int \frac{\tanh(x)}{1+\tanh(x)} dx$

Optimal result . . . . .	935
Mathematica [A] (verified) . . . . .	935
Rubi [C] (verified) . . . . .	936
Maple [A] (verified) . . . . .	937
Fricas [B] (verification not implemented) . . . . .	938
Sympy [B] (verification not implemented) . . . . .	938
Maxima [A] (verification not implemented) . . . . .	938
Giac [A] (verification not implemented) . . . . .	939
Mupad [B] (verification not implemented) . . . . .	939
Reduce [B] (verification not implemented) . . . . .	939

#### Optimal result

Integrand size = 9, antiderivative size = 16

$$\int \frac{\tanh(x)}{1+\tanh(x)} dx = \frac{x}{2} + \frac{1}{2(1+\tanh(x))}$$

output `1/2*x+1/(2+2*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\tanh(x)}{1+\tanh(x)} dx = \frac{1}{2} \left( \operatorname{arctanh}(\tanh(x)) + \frac{1}{1+\tanh(x)} \right)$$

input `Integrate[Tanh[x]/(1 + Tanh[x]), x]`

output `(ArcTanh[Tanh[x]] + (1 + Tanh[x])^(-1))/2`



**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {3042, 26, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)}{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{4009} \\
 & -i \left( \frac{i \int 1 dx}{2} + \frac{i}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{24} \\
 & -i \left( \frac{ix}{2} + \frac{i}{2(\tanh(x) + 1)} \right)
 \end{aligned}$$

input `Int [Tanh[x]/(1 + Tanh[x]),x]`

output `(-I)*((I/2)*x + (I/2)/(1 + Tanh[x]))`

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-*(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

## Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

method	result	size
risch	$\frac{x}{2} + \frac{e^{-2x}}{4}$	11
parallelrisch	$-\frac{-1 - \tanh(x)x - x}{2(1 + \tanh(x))}$	19
derivativedivides	$\frac{1}{2+2 \tanh(x)} + \frac{\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24
default	$\frac{1}{2+2 \tanh(x)} + \frac{\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24

input `int(tanh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)`

output `1/2*x+1/4*exp(-2*x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs.  $2(12) = 24$ .

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{(2x + 1) \cosh(x) + (2x - 1) \sinh(x)}{4(\cosh(x) + \sinh(x))}$$

input `integrate(tanh(x)/(1+tanh(x)),x, algorithm="fricas")`

output `1/4*((2*x + 1)*cosh(x) + (2*x - 1)*sinh(x))/(cosh(x) + sinh(x))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs.  $2(10) = 20$ .

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} + \frac{1}{2 \tanh(x) + 2}$$

input `integrate(tanh(x)/(1+tanh(x)),x)`

output `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) + 1/(2*tanh(x) + 2)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{1}{2} x + \frac{1}{4} e^{(-2x)}$$

input `integrate(tanh(x)/(1+tanh(x)),x, algorithm="maxima")`

output `1/2*x + 1/4*e^(-2*x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{1}{2}x + \frac{1}{4}e^{(-2x)}$$

input `integrate(tanh(x)/(1+tanh(x)),x, algorithm="giac")`output `1/2*x + 1/4*e^(-2*x)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{x}{2} + \frac{e^{-2x}}{4}$$

input `int(tanh(x)/(tanh(x) + 1),x)`output `x/2 + exp(-2*x)/4`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{\tanh(x)}{1 + \tanh(x)} dx = \frac{2e^{2x}x + 1}{4e^{2x}}$$

input `int(tanh(x)/(1+tanh(x)),x)`output `(2*e**(2*x)*x + 1)/(4*e**(2*x))`

### 3.121 $\int \frac{1}{1+\tanh(x)} dx$

Optimal result	940
Mathematica [A] (verified)	940
Rubi [A] (verified)	941
Maple [A] (verified)	942
Fricas [B] (verification not implemented)	942
Sympy [B] (verification not implemented)	943
Maxima [A] (verification not implemented)	943
Giac [A] (verification not implemented)	943
Mupad [B] (verification not implemented)	944
Reduce [B] (verification not implemented)	944

#### Optimal result

Integrand size = 6, antiderivative size = 16

$$\int \frac{1}{1+\tanh(x)} dx = \frac{x}{2} - \frac{1}{2(1+\tanh(x))}$$

output `1/2*x-1/(2+2*tanh(x))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{1}{1+\tanh(x)} dx = \frac{1}{2} \operatorname{arctanh}(\tanh(x)) - \frac{1}{2(1+\tanh(x))}$$

input `Integrate[(1 + Tanh[x])^(-1), x]`

output `ArcTanh[Tanh[x]]/2 - 1/(2*(1 + Tanh[x]))`

**Rubi [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\tanh(x) + 1} dx$$

↓ 3042

$$\int \frac{1}{1 - i \tan(ix)} dx$$

↓ 3960

$$\frac{\int 1 dx}{2} - \frac{1}{2(\tanh(x) + 1)}$$

↓ 24

$$\frac{x}{2} - \frac{1}{2(\tanh(x) + 1)}$$

input `Int[(1 + Tanh[x])^(-1), x]`

output `x/2 - 1/(2*(1 + Tanh[x]))`

**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960

```
Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a +
b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^
(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

method	result	size
risch	$\frac{x}{2} - \frac{e^{-2x}}{4}$	11
parallelsch	$-\frac{1 - \tanh(x)x - x}{2(1 + \tanh(x))}$	19
derivativdivides	$-\frac{1}{2(1 + \tanh(x))} + \frac{\ln(1 + \tanh(x))}{4} - \frac{\ln(\tanh(x) - 1)}{4}$	24
default	$-\frac{1}{2(1 + \tanh(x))} + \frac{\ln(1 + \tanh(x))}{4} - \frac{\ln(\tanh(x) - 1)}{4}$	24

input

```
int(1/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
1/2*x-1/4*exp(-2*x)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 26 vs.  $2(12) = 24$ .

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{(2x - 1) \cosh(x) + (2x + 1) \sinh(x)}{4(\cosh(x) + \sinh(x))}$$

input

```
integrate(1/(1+tanh(x)),x, algorithm="fricas")
```

output

```
1/4*((2*x - 1)*cosh(x) + (2*x + 1)*sinh(x))/(cosh(x) + sinh(x))
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs.  $2(10) = 20$ .

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{1}{2 \tanh(x) + 2}$$

input `integrate(1/(1+tanh(x)),x)`

output `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) - 1/(2*tanh(x) + 2)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{1}{2} x - \frac{1}{4} e^{(-2x)}$$

input `integrate(1/(1+tanh(x)),x, algorithm="maxima")`

output `1/2*x - 1/4*e^(-2*x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{1}{2} x - \frac{1}{4} e^{(-2x)}$$

input `integrate(1/(1+tanh(x)),x, algorithm="giac")`

output `1/2*x - 1/4*e^(-2*x)`



**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{x}{2} - \frac{e^{-2x}}{4}$$

input `int(1/(tanh(x) + 1),x)`

output `x/2 - exp(-2*x)/4`

**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{1 + \tanh(x)} dx = \frac{2e^{2x}x - 1}{4e^{2x}}$$

input `int(1/(1+tanh(x)),x)`

output `(2*e**(2*x)*x - 1)/(4*e**(2*x))`

### 3.122 $\int \frac{\coth(x)}{1+\tanh(x)} dx$

Optimal result	945
Mathematica [A] (verified)	945
Rubi [C] (verified)	946
Maple [A] (verified)	948
Fricas [B] (verification not implemented)	948
Sympy [F]	949
Maxima [A] (verification not implemented)	949
Giac [A] (verification not implemented)	949
Mupad [B] (verification not implemented)	950
Reduce [B] (verification not implemented)	950

#### Optimal result

Integrand size = 9, antiderivative size = 19

$$\int \frac{\coth(x)}{1+\tanh(x)} dx = -\frac{x}{2} + \log(\sinh(x)) + \frac{1}{2(1+\tanh(x))}$$

output

```
-1/2*x+ln(sinh(x))+1/(2+2*tanh(x))
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{\coth(x)}{1+\tanh(x)} dx = -\frac{1}{4} \log(1-\tanh(x)) + \log(\tanh(x)) - \frac{3}{4} \log(1+\tanh(x)) + \frac{1}{2(1+\tanh(x))}$$

input

```
Integrate[Coth[x]/(1 + Tanh[x]), x]
```

output

```
-1/4*Log[1 - Tanh[x]] + Log[Tanh[x]] - (3*Log[1 + Tanh[x]])/4 + 1/(2*(1 + Tanh[x]))
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.68, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {3042, 26, 4034, 26, 3042, 26, 3956, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{(1 - i \tan(ix)) \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{(1 - i \tan(ix)) \tan(ix)} dx \\
 & \quad \downarrow \text{4034} \\
 & i \left( i \int \frac{1}{\tanh(x) + 1} dx + \int -i \coth(x) dx \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( i \int \frac{1}{\tanh(x) + 1} dx - i \int \coth(x) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & i \left( i \int \frac{1}{1 - i \tan(ix)} dx - i \int -i \tan \left( ix + \frac{\pi}{2} \right) dx \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( i \int \frac{1}{1 - i \tan(ix)} dx - \int \tan \left( ix + \frac{\pi}{2} \right) dx \right) \\
 & \quad \downarrow \text{3956} \\
 & i \left( i \int \frac{1}{1 - i \tan(ix)} dx - i \log(\sinh(x)) \right) \\
 & \quad \downarrow \text{3960}
 \end{aligned}$$

$$i \left( i \left( \frac{\int 1 dx}{2} - \frac{1}{2(\tanh(x) + 1)} \right) - i \log(\sinh(x)) \right)$$

↓ 24

$$i \left( i \left( \frac{x}{2} - \frac{1}{2(\tanh(x) + 1)} \right) - i \log(\sinh(x)) \right)$$

input `Int[Coth[x]/(1 + Tanh[x]),x]`

output `I*((-I)*Log[Sinh[x]] + I*(x/2 - 1/(2*(1 + Tanh[x])))`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4034

```
Int[1/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (
f_.)*(x_)])), x_Symbol] :> Simp[b/(b*c - a*d) Int[1/(a + b*Tan[e + f*x]),
x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*Tan[e + f*x]), x], x] /; FreeQ[
{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2
+ d^2, 0]
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
risch	$-\frac{3x}{2} + \frac{e^{-2x}}{4} + \ln(e^{2x} - 1)$	18
default	$-\frac{\ln(\tanh(\frac{x}{2})-1)}{2} + \ln(\tanh(\frac{x}{2})) - \frac{1}{1+\tanh(\frac{x}{2})} + \frac{1}{(1+\tanh(\frac{x}{2}))^2} - \frac{3\ln(1+\tanh(\frac{x}{2}))}{2}$	43
paralelrisch	$\frac{(-2-2\tanh(x))\ln(1-\tanh(x))+(2+2\tanh(x))\ln(\tanh(x))-3\tanh(x)x-3x+1}{2+2\tanh(x)}$	44

input

```
int(coth(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-3/2*x+1/4*exp(-2*x)+ln(exp(2*x)-1)
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 73 vs.  $2(15) = 30$ .

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 3.84

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx =$$

$$\frac{6x \cosh(x)^2 + 12x \cosh(x) \sinh(x) + 6x \sinh(x)^2 - 4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2) \ln(\cosh(x) + \sinh(x))}{4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)}$$

input

```
integrate(coth(x)/(1+tanh(x)),x, algorithm="fricas")
```

output

```
-1/4*(6*x*cosh(x)^2 + 12*x*cosh(x)*sinh(x) + 6*x*sinh(x)^2 - 4*(cosh(x)^2
+ 2*cosh(x)*sinh(x) + sinh(x)^2)*log(2*sinh(x)/(cosh(x) - sinh(x))) - 1)/(
cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)
```

**Sympy [F]**

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx = \int \frac{\coth(x)}{\tanh(x) + 1} dx$$

input

```
integrate(coth(x)/(1+tanh(x)),x)
```

output

```
Integral(coth(x)/(tanh(x) + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.26

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx = \frac{1}{2}x + \frac{1}{4}e^{(-2x)} + \log(e^{(-x)} + 1) + \log(e^{(-x)} - 1)$$

input

```
integrate(coth(x)/(1+tanh(x)),x, algorithm="maxima")
```

output

```
1/2*x + 1/4*e^(-2*x) + log(e^(-x) + 1) + log(e^(-x) - 1)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx = -\frac{3}{2}x + \frac{1}{4}e^{(-2x)} + \log(|e^{(2x)} - 1|)$$

input

```
integrate(coth(x)/(1+tanh(x)),x, algorithm="giac")
```

output `-3/2*x + 1/4*e^(-2*x) + log(abs(e^(2*x) - 1))`

### Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx = \ln(e^{2x} - 1) - \frac{3x}{2} + \frac{e^{-2x}}{4}$$

input `int(coth(x)/(tanh(x) + 1),x)`

output `log(exp(2*x) - 1) - (3*x)/2 + exp(-2*x)/4`

### Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.37

$$\int \frac{\coth(x)}{1 + \tanh(x)} dx = \frac{4e^{2x}\log(e^x - 1) + 4e^{2x}\log(e^x + 1) - 6e^{2x}x + 1}{4e^{2x}}$$

input `int(coth(x)/(1+tanh(x)),x)`

output `(4*e**(2*x)*log(e**x - 1) + 4*e**(2*x)*log(e**x + 1) - 6*e**(2*x)*x + 1)/(4*e**(2*x))`

### 3.123 $\int \frac{\coth^2(x)}{1+\tanh(x)} dx$

Optimal result	951
Mathematica [C] (verified)	951
Rubi [C] (verified)	952
Maple [A] (verified)	955
Fricas [B] (verification not implemented)	955
Sympy [F]	956
Maxima [A] (verification not implemented)	956
Giac [A] (verification not implemented)	956
Mupad [B] (verification not implemented)	957
Reduce [B] (verification not implemented)	957

#### Optimal result

Integrand size = 11, antiderivative size = 29

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \frac{3x}{2} - \frac{3 \coth(x)}{2} - \log(\sinh(x)) + \frac{\coth(x)}{2(1 + \tanh(x))}$$

output `3/2*x-3/2*coth(x)-ln(sinh(x))+coth(x)/(2+2*tanh(x))`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.20 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.62

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \frac{1}{2} \left( \coth^2(x) + \frac{\coth^4(x)}{1 + \coth(x)} - \coth^3(x) \operatorname{Hypergeometric2F1} \left( -\frac{3}{2}, 1, -\frac{1}{2}, \tanh^2(x) \right) - 2(\log(\cosh(x)) + \log(\tanh(x))) \right)$$

input `Integrate[Coth[x]^2/(1 + Tanh[x]),x]`



output

```
(Coth[x]^2 + Coth[x]^4/(1 + Coth[x]) - Coth[x]^3*Hypergeometric2F1[-3/2, 1, -1/2, Tanh[x]^2] - 2*(Log[Cosh[x]] + Log[Tanh[x]]))/2
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.34, number of steps used = 13, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.182$ , Rules used = {3042, 25, 4035, 3042, 25, 4012, 3042, 26, 4014, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^2(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{(1 - i \tan(ix)) \tan(ix)^2} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{(1 - i \tan(ix)) \tan(ix)^2} dx \\
 & \quad \downarrow \text{4035} \\
 & \frac{1}{2} \int \coth^2(x)(3 - 2 \tanh(x)) dx + \frac{\coth(x)}{2(\tanh(x) + 1)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\coth(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \int -\frac{2i \tan(ix) + 3}{\tan(ix)^2} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{\coth(x)}{2(\tanh(x) + 1)} - \frac{1}{2} \int \frac{2i \tan(ix) + 3}{\tan(ix)^2} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{1}{2} \left( -\int \coth(x)(2 - 3 \tanh(x)) dx - 3 \coth(x) \right) + \frac{\coth(x)}{2(\tanh(x) + 1)}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - \int \frac{i(3i \tan(ix) + 2)}{\tan(ix)} dx \right) \\
& \downarrow 26 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i \int \frac{3i \tan(ix) + 2}{\tan(ix)} dx \right) \\
& \downarrow 4014 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i \left( 2 \int -i \coth(x) dx + 3ix \right) \right) \\
& \downarrow 26 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i(3ix - 2i \int \coth(x) dx) \right) \\
& \downarrow 3042 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i \left( 3ix - 2i \int -i \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) \\
& \downarrow 26 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i \left( 3ix - 2 \int \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) \\
& \downarrow 3956 \\
& \frac{\coth(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -3 \coth(x) - i(3ix - 2i \log(\sinh(x))) \right)
\end{aligned}$$

input `Int [Coth[x]^2/(1 + Tanh[x]), x]`

output `(-3*Coth[x] - I*((3*I)*x - (2*I)*Log[Sinh[x]]))/2 + Coth[x]/(2*(1 + Tanh[x]))`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 26  $\text{Int}[(\text{Complex}[0, \text{a}_])*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], \text{a}]) \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2, 1]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3956  $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x\_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d}*x], \text{x}]]/\text{d}, \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4012  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b}*c - \text{a}*d)*((\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1})/(\text{f}*(\text{m} + 1)*(a^2 + b^2)), \text{x}] + \text{Simp}[1/(a^2 + b^2) \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}*\text{Simp}[\text{a}*c + \text{b}*d - (\text{b}*c - \text{a}*d)*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1]$
- rule 4014  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{a}*c + \text{b}*d)*(x/(a^2 + b^2)), \text{x}] + \text{Simp}[(\text{b}*c - \text{a}*d)/(a^2 + b^2) \quad \text{Int}[(\text{b} - \text{a}*\text{Tan}[\text{e} + \text{f}*x])/(a + \text{b}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{a}*c + \text{b}*d, 0]$
- rule 4035  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{n}_.}/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{a})*((\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} + 1})/(2*\text{f}*(\text{b}*c - \text{a}*d)*(a + \text{b}*\text{Tan}[\text{e} + \text{f}*x])), \text{x}] + \text{Simp}[1/(2*\text{a}*(\text{b}*c - \text{a}*d)) \quad \text{Int}[(\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n}}*\text{Simp}[\text{b}*c + \text{a}*d*(\text{n} - 1) - \text{b}*d*\text{n}*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{EqQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!GtQ}[\text{n}, 0]$

**Maple [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

method	result
risch	$\frac{5x}{2} - \frac{e^{-2x}}{4} - \frac{2}{e^{2x}-1} - \ln(e^{2x}-1)$
parallelrisc	$\frac{(2+2 \tanh(x)) \ln(1-\tanh(x))+(-2-2 \tanh(x)) \ln(\tanh(x))+5 \tanh(x)x+5x-2 \coth(x)-3}{2+2 \tanh(x)}$
default	$-\frac{\tanh(\frac{x}{2})}{2} - \frac{\ln(\tanh(\frac{x}{2})-1)}{2} - \frac{1}{2 \tanh(\frac{x}{2})} - \ln\left(\tanh\left(\frac{x}{2}\right)\right) - \frac{1}{(1+\tanh(\frac{x}{2}))^2} + \frac{1}{1+\tanh(\frac{x}{2})} + \frac{5 \ln(1+\tanh(\frac{x}{2}))}{2}$

input `int(coth(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)`output `5/2*x-1/4*exp(-2*x)-2/(exp(2*x)-1)-ln(exp(2*x)-1)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(23) = 46.

Time = 0.09 (sec) , antiderivative size = 196, normalized size of antiderivative = 6.76

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{10x \cosh(x)^4 + 40x \cosh(x) \sinh(x)^3 + 10x \sinh(x)^4 - (10x + 9) \cosh(x)^2 + (60x \cosh(x))^2 - 10x - 4(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + (6 \cosh(x)^2 - 1) \sinh(x)^2 - \cosh(x)^2 + 2(2 \cosh(x)^3 - \cosh(x)) \sinh(x)) \log(2 \sinh(x) / (\cosh(x) - \sinh(x))) + 2(20x \cosh(x)^3 - (10x + 9) \cosh(x)) \sinh(x) + 1}{(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + (6 \cosh(x)^2 - 1) \sinh(x)^2 - \cosh(x)^2 + 2(2 \cosh(x)^3 - \cosh(x)) \sinh(x))}$$

input `integrate(coth(x)^2/(1+tanh(x)),x, algorithm="fricas")`output `1/4*(10*x*cosh(x)^4 + 40*x*cosh(x)*sinh(x)^3 + 10*x*sinh(x)^4 - (10*x + 9)*cosh(x)^2 + (60*x*cosh(x)^2 - 10*x - 9)*sinh(x)^2 - 4*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x))*log(2*sinh(x)/(cosh(x) - sinh(x))) + 2*(20*x*cosh(x)^3 - (10*x + 9)*cosh(x))*sinh(x) + 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x))`

**Sympy [F]**

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \int \frac{\coth^2(x)}{\tanh(x) + 1} dx$$

input `integrate(coth(x)**2/(1+tanh(x)),x)`

output `Integral(coth(x)**2/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.31

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \frac{1}{2}x + \frac{2}{e^{(-2x)} - 1} - \frac{1}{4}e^{(-2x)} - \log(e^{(-x)} + 1) - \log(e^{(-x)} - 1)$$

input `integrate(coth(x)^2/(1+tanh(x)),x, algorithm="maxima")`

output `1/2*x + 2/(e^(-2*x) - 1) - 1/4*e^(-2*x) - log(e^(-x) + 1) - log(e^(-x) - 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.24

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \frac{5}{2}x - \frac{(9e^{(2x)} - 1)e^{(-2x)}}{4(e^{(2x)} - 1)} - \log(|e^{(2x)} - 1|)$$

input `integrate(coth(x)^2/(1+tanh(x)),x, algorithm="giac")`

output `5/2*x - 1/4*(9*e^(2*x) - 1)*e^(-2*x)/(e^(2*x) - 1) - log(abs(e^(2*x) - 1))`

**Mupad [B] (verification not implemented)**

Time = 2.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx = \frac{5x}{2} - \ln(e^{2x} - 1) - \frac{e^{-2x}}{4} - \frac{2}{e^{2x} - 1}$$

input `int(coth(x)^2/(tanh(x) + 1),x)`output `(5*x)/2 - log(exp(2*x) - 1) - exp(-2*x)/4 - 2/(exp(2*x) - 1)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 95, normalized size of antiderivative = 3.28

$$\int \frac{\coth^2(x)}{1 + \tanh(x)} dx$$

$$= \frac{-4e^{4x}\log(e^x - 1) - 4e^{4x}\log(e^x + 1) + 10e^{4x}x - 9e^{4x} + 4e^{2x}\log(e^x - 1) + 4e^{2x}\log(e^x + 1) - 10e^{2x}x + 1}{4e^{2x}(e^{2x} - 1)}$$

input `int(coth(x)^2/(1+tanh(x)),x)`output `( - 4*e**(4*x)*log(e**x - 1) - 4*e**(4*x)*log(e**x + 1) + 10*e**(4*x)*x - 9*e**(4*x) + 4*e**(2*x)*log(e**x - 1) + 4*e**(2*x)*log(e**x + 1) - 10*e**(2*x)*x + 1)/(4*e**(2*x)*(e**(2*x) - 1))`

### 3.124 $\int \frac{\coth^3(x)}{1+\tanh(x)} dx$

Optimal result	958
Mathematica [C] (verified)	958
Rubi [C] (verified)	959
Maple [A] (verified)	962
Fricas [B] (verification not implemented)	963
Sympy [F]	963
Maxima [A] (verification not implemented)	964
Giac [A] (verification not implemented)	964
Mupad [B] (verification not implemented)	964
Reduce [B] (verification not implemented)	965

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = -\frac{3x}{2} + \frac{3 \coth(x)}{2} - \coth^2(x) + 2 \log(\sinh(x)) + \frac{\coth^2(x)}{2(1 + \tanh(x))}$$

output `-3/2*x+3/2*coth(x)-coth(x)^2+2*ln(sinh(x))+coth(x)^2/(2+2*tanh(x))`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.21 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.46

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = \frac{1}{2} \left( -2 \coth^2(x) - \coth^4(x) + \frac{\coth^5(x)}{1 + \coth(x)} + \coth^3(x) \operatorname{Hypergeometric2F1} \left( -\frac{3}{2}, 1, -\frac{1}{2}, \tanh^2(x) \right) + 4(\log(\cosh(x)) + \log(\tanh(x))) \right)$$

input `Integrate[Coth[x]^3/(1 + Tanh[x]), x]`

output

```
(-2*Coth[x]^2 - Coth[x]^4 + Coth[x]^5/(1 + Coth[x]) + Coth[x]^3*Hypergeometric2F1[-3/2, 1, -1/2, Tanh[x]^2] + 4*(Log[Cosh[x]] + Log[Tanh[x]]))/2
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.62, number of steps used = 18, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.636$ , Rules used = {3042, 26, 4035, 26, 3042, 26, 4012, 26, 3042, 25, 4012, 3042, 26, 4014, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^3(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i}{(1 - i \tan(ix)) \tan(ix)^3} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{1}{(1 - i \tan(ix)) \tan(ix)^3} dx \\
 & \quad \downarrow \text{4035} \\
 & -i \left( \frac{i \coth^2(x)}{2(\tanh(x) + 1)} - \frac{1}{2} \int -i \coth^3(x)(4 - 3 \tanh(x)) dx \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( \frac{1}{2} i \int \coth^3(x)(4 - 3 \tanh(x)) dx + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{1}{2} i \int -\frac{i(3i \tan(ix) + 4)}{\tan(ix)^3} dx + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
 & \quad \downarrow \text{26}
 \end{aligned}$$



$$\begin{aligned}
& -i \left( \frac{1}{2} \int \frac{3i \tan(ix) + 4}{\tan(ix)^3} dx + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 4012 \\
& -i \left( \frac{1}{2} \left( \int -i \coth^2(x)(3 - 4 \tanh(x)) dx - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( -i \int \coth^2(x)(3 - 4 \tanh(x)) dx - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 3042 \\
& -i \left( \frac{1}{2} \left( -i \int -\frac{4i \tan(ix) + 3}{\tan(ix)^2} dx - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 25 \\
& -i \left( \frac{1}{2} \left( i \int \frac{4i \tan(ix) + 3}{\tan(ix)^2} dx - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 4012 \\
& -i \left( \frac{1}{2} \left( i \left( \int \coth(x)(4 - 3 \tanh(x)) dx + 3 \coth(x) \right) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 3042 \\
& -i \left( \frac{1}{2} \left( i \left( 3 \coth(x) + \int \frac{i(3i \tan(ix) + 4)}{\tan(ix)} dx \right) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( i \left( 3 \coth(x) + i \int \frac{3i \tan(ix) + 4}{\tan(ix)} dx \right) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 4014 \\
& -i \left( \frac{1}{2} \left( i(3 \coth(x) + i(4 \int -i \coth(x) dx + 3ix)) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 26 \\
& -i \left( \frac{1}{2} \left( i(3 \coth(x) + i(3ix - 4i \int \coth(x) dx)) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& -i \left( \frac{1}{2} \left( i \left( 3 \coth(x) + i \left( 3ix - 4i \int -i \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow \text{26} \\
& -i \left( \frac{1}{2} \left( i \left( 3 \coth(x) + i \left( 3ix - 4 \int \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) - 2i \coth^2(x) \right) + \frac{i \coth^2(x)}{2(\tanh(x) + 1)} \right) \\
& \quad \downarrow \text{3956} \\
& -i \left( \frac{i \coth^2(x)}{2(\tanh(x) + 1)} + \frac{1}{2} (i(3 \coth(x) + i(3ix - 4i \log(\sinh(x)))) - 2i \coth^2(x)) \right)
\end{aligned}$$

input `Int[Coth[x]^3/(1 + Tanh[x]),x]`

output `(-I)*((( -2*I)*Coth[x]^2 + I*(3*Coth[x] + I*((3*I)*x - (4*I)*Log[Sinh[x]])))/2 + ((I/2)*Coth[x]^2)/(1 + Tanh[x]))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

rule 4035

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_.) + (b_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-a)*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*
c - a*d)*(a + b*Tan[e + f*x]))), x] + Simp[1/(2*a*(b*c - a*d)) Int[(c + d
*Tan[e + f*x])^n*Simp[b*c + a*d*(n - 1) - b*d*n*Tan[e + f*x], x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& NeQ[c^2 + d^2, 0] && !GtQ[n, 0]
```

## Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

method	result
risch	$-\frac{7x}{2} + \frac{e^{-2x}}{4} - \frac{2}{(e^{2x}-1)^2} + 2 \ln(e^{2x} - 1)$
parallelrisch	$\frac{(-4 \tanh(x)-4) \ln(1-\tanh(x))+(4+4 \tanh(x)) \ln(\tanh(x))-7 \tanh(x)x-\coth(x)^2-7x+\coth(x)+3}{2+2 \tanh(x)}$
default	$-\frac{\tanh(\frac{x}{2})^2}{8} + \frac{\tanh(\frac{x}{2})}{2} - \frac{\ln(\tanh(\frac{x}{2})-1)}{2} - \frac{1}{8 \tanh(\frac{x}{2})^2} + \frac{1}{2 \tanh(\frac{x}{2})} + 2 \ln\left(\tanh\left(\frac{x}{2}\right)\right) + \frac{1}{(1+\tanh(\frac{x}{2}))^2}$

input

```
int(coth(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-7/2*x+1/4*exp(-2*x)-2/(exp(2*x)-1)^2+2*ln(exp(2*x)-1)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 357 vs.  $2(31) = 62$ .

Time = 0.10 (sec) , antiderivative size = 357, normalized size of antiderivative = 9.65

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="fricas")`

output

```
-1/4*(14*x*cosh(x)^6 + 84*x*cosh(x)*sinh(x)^5 + 14*x*sinh(x)^6 - (28*x + 1)
)*cosh(x)^4 + (210*x*cosh(x)^2 - 28*x - 1)*sinh(x)^4 + 4*(70*x*cosh(x)^3 -
(28*x + 1)*cosh(x))*sinh(x)^3 + 2*(7*x + 5)*cosh(x)^2 + 2*(105*x*cosh(x)^
4 - 3*(28*x + 1)*cosh(x)^2 + 7*x + 5)*sinh(x)^2 - 8*(cosh(x)^6 + 6*cosh(x)
*sinh(x)^5 + sinh(x)^6 + (15*cosh(x)^2 - 2)*sinh(x)^4 - 2*cosh(x)^4 + 4*(5
*cosh(x)^3 - 2*cosh(x))*sinh(x)^3 + (15*cosh(x)^4 - 12*cosh(x)^2 + 1)*sinh
(x)^2 + cosh(x)^2 + 2*(3*cosh(x)^5 - 4*cosh(x)^3 + cosh(x))*sinh(x))*log(2
*sinh(x)/(cosh(x) - sinh(x))) + 4*(21*x*cosh(x)^5 - (28*x + 1)*cosh(x)^3 +
(7*x + 5)*cosh(x))*sinh(x) - 1)/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)
)^6 + (15*cosh(x)^2 - 2)*sinh(x)^4 - 2*cosh(x)^4 + 4*(5*cosh(x)^3 - 2*cosh
(x))*sinh(x)^3 + (15*cosh(x)^4 - 12*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 +
2*(3*cosh(x)^5 - 4*cosh(x)^3 + cosh(x))*sinh(x))
```

**Sympy [F]**

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = \int \frac{\coth^3(x)}{\tanh(x) + 1} dx$$

input `integrate(coth(x)**3/(1+tanh(x)),x)`

output `Integral(coth(x)**3/(tanh(x) + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.46

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = \frac{1}{2}x + \frac{2(2e^{-2x} - 1)}{2e^{-2x} - e^{-4x} - 1} + \frac{1}{4}e^{-2x} + 2 \log(e^{-x} + 1) + 2 \log(e^{-x} - 1)$$

input `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="maxima")`output `1/2*x + 2*(2*e^(-2*x) - 1)/(2*e^(-2*x) - e^(-4*x) - 1) + 1/4*e^(-2*x) + 2*log(e^(-x) + 1) + 2*log(e^(-x) - 1)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.08

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = -\frac{7}{2}x + \frac{(e^{4x} - 10e^{2x} + 1)e^{-2x}}{4(e^{2x} - 1)^2} + 2 \log(|e^{2x} - 1|)$$

input `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="giac")`output `-7/2*x + 1/4*(e^(4*x) - 10*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) - 1)^2 + 2*log(abs(e^(2*x) - 1))`**Mupad [B] (verification not implemented)**

Time = 2.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx = 2 \ln(e^{2x} - 1) - \frac{7x}{2} + \frac{e^{-2x}}{4} - \frac{2}{e^{4x} - 2e^{2x} + 1}$$

input `int(coth(x)^3/(tanh(x) + 1),x)`

output `2*log(exp(2*x) - 1) - (7*x)/2 + exp(-2*x)/4 - 2/(exp(4*x) - 2*exp(2*x) + 1)`

### Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 3.81

$$\int \frac{\coth^3(x)}{1 + \tanh(x)} dx$$

$$= \frac{16e^{6x}\log(e^x - 1) + 16e^{6x}\log(e^x + 1) - 28e^{6x}x + e^{6x} - 32e^{4x}\log(e^x - 1) - 32e^{4x}\log(e^x + 1) + 56e^{4x}x + 1}{8e^{2x}(e^{4x} - 2e^{2x} + 1)}$$

input `int(coth(x)^3/(1+tanh(x)),x)`

output `(16*e**(6*x)*log(e**x - 1) + 16*e**(6*x)*log(e**x + 1) - 28*e**(6*x)*x + e**  
 (6*x) - 32*e**(4*x)*log(e**x - 1) - 32*e**(4*x)*log(e**x + 1) + 56*e**(4  
 *x)*x + 16*e**(2*x)*log(e**x - 1) + 16*e**(2*x)*log(e**x + 1) - 28*e**(2*x  
 )*x - 19*e**(2*x) + 2)/(8*e**(2*x)*(e**(4*x) - 2*e**(2*x) + 1))`

### 3.125 $\int \frac{\coth^4(x)}{1+\tanh(x)} dx$

Optimal result	966
Mathematica [C] (verified)	966
Rubi [C] (verified)	967
Maple [A] (verified)	971
Fricas [B] (verification not implemented)	971
Sympy [F]	972
Maxima [A] (verification not implemented)	973
Giac [A] (verification not implemented)	973
Mupad [B] (verification not implemented)	973
Reduce [B] (verification not implemented)	974

#### Optimal result

Integrand size = 11, antiderivative size = 43

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \frac{5x}{2} - \frac{5 \coth(x)}{2} + \coth^2(x) - \frac{5 \coth^3(x)}{6} - 2 \log(\sinh(x)) + \frac{\coth^3(x)}{2(1 + \tanh(x))}$$

output

`5/2*x-5/2*coth(x)+coth(x)^2-5/6*coth(x)^3-2*ln(sinh(x))+coth(x)^3/(2+2*tanh(x))`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \frac{1}{2} \left( 2 \coth^2(x) + \coth^4(x) + \frac{\coth^6(x)}{1 + \coth(x)} - \coth^5(x) \operatorname{Hypergeometric2F1} \left( -\frac{5}{2}, 1, -\frac{3}{2}, \tanh^2(x) \right) - 4(\log(\cosh(x)) + \log(\tanh(x))) \right)$$

input `Integrate[Coth[x]^4/(1 + Tanh[x]),x]`

output `(2*Coth[x]^2 + Coth[x]^4 + Coth[x]^6/(1 + Coth[x]) - Coth[x]^5*Hypergeometric2F1[-5/2, 1, -3/2, Tanh[x]^2] - 4*(Log[Cosh[x]] + Log[Tanh[x]]))/2`

### Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.56, number of steps used = 20, number of rules used = 20,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.818$ , Rules used = {3042, 4035, 25, 3042, 4012, 25, 3042, 26, 4012, 26, 3042, 25, 4012, 3042, 26, 4014, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^4(x)}{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(1 - i \tan(ix)) \tan(ix)^4} dx \\
 & \quad \downarrow \text{4035} \\
 & \frac{\coth^3(x)}{2(\tanh(x) + 1)} - \frac{1}{2} \int -\coth^4(x)(5 - 4 \tanh(x)) dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int \coth^4(x)(5 - 4 \tanh(x)) dx + \frac{\coth^3(x)}{2(\tanh(x) + 1)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \int \frac{4i \tan(ix) + 5}{\tan(ix)^4} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{1}{2} \left( \int -\coth^3(x)(4 - 5 \tanh(x)) dx - \frac{5 \coth^3(x)}{3} \right) + \frac{\coth^3(x)}{2(\tanh(x) + 1)}
 \end{aligned}$$



$$\begin{aligned}
& \downarrow 25 \\
& \frac{1}{2} \left( - \int \coth^3(x)(4 - 5 \tanh(x)) dx - \frac{5}{3} \coth^3(x) \right) + \frac{\coth^3(x)}{2(\tanh(x) + 1)} \\
& \downarrow 3042 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5}{3} \coth^3(x) - \int - \frac{i(5i \tan(ix) + 4)}{\tan(ix)^3} dx \right) \\
& \downarrow 26 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \int \frac{5i \tan(ix) + 4}{\tan(ix)^3} dx \right) \\
& \downarrow 4012 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( \int -i \coth^2(x)(5 - 4 \tanh(x)) dx - 2i \coth^2(x) \right) \right) \\
& \downarrow 26 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( -i \int \coth^2(x)(5 - 4 \tanh(x)) dx - 2i \coth^2(x) \right) \right) \\
& \downarrow 3042 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( -i \int - \frac{4i \tan(ix) + 5}{\tan(ix)^2} dx - 2i \coth^2(x) \right) \right) \\
& \downarrow 25 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( i \int \frac{4i \tan(ix) + 5}{\tan(ix)^2} dx - 2i \coth^2(x) \right) \right) \\
& \downarrow 4012 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \\
& \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( i \left( \int \coth(x)(4 - 5 \tanh(x)) dx + 5 \coth(x) \right) - 2i \coth^2(x) \right) \right) \\
& \downarrow 3042 \\
& \frac{\coth^3(x)}{2(\tanh(x) + 1)} + \\
& \frac{1}{2} \left( - \frac{5 \coth^3(x)}{3} + i \left( i \left( 5 \coth(x) + \int \frac{i(5i \tan(ix) + 4)}{\tan(ix)} dx \right) - 2i \coth^2(x) \right) \right) \\
& \downarrow 26
\end{aligned}$$

$$\begin{aligned}
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i \left( i \left( 5 \operatorname{coth}(x) + i \int \frac{5i \tan(ix) + 4}{\tan(ix)} dx \right) - 2i \operatorname{coth}^2(x) \right) \right) \\
& \quad \downarrow 4014 \\
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \\
& \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i \left( i(5 \operatorname{coth}(x) + i(4 \int -i \operatorname{coth}(x) dx + 5ix)) - 2i \operatorname{coth}^2(x) \right) \right) \\
& \quad \downarrow 26 \\
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \\
& \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i \left( i(5 \operatorname{coth}(x) + i(5ix - 4i \int \operatorname{coth}(x) dx)) - 2i \operatorname{coth}^2(x) \right) \right) \\
& \quad \downarrow 3042 \\
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \\
& \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i \left( i \left( 5 \operatorname{coth}(x) + i \left( 5ix - 4i \int -i \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) - 2i \operatorname{coth}^2(x) \right) \right) \\
& \quad \downarrow 26 \\
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \\
& \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i \left( i \left( 5 \operatorname{coth}(x) + i \left( 5ix - 4 \int \tan \left( ix + \frac{\pi}{2} \right) dx \right) \right) - 2i \operatorname{coth}^2(x) \right) \right) \\
& \quad \downarrow 3956 \\
& \frac{\operatorname{coth}^3(x)}{2(\tanh(x)+1)} + \frac{1}{2} \left( -\frac{5 \operatorname{coth}^3(x)}{3} + i(i(5 \operatorname{coth}(x) + i(5ix - 4i \log(\sinh(x)))) - 2i \operatorname{coth}^2(x)) \right)
\end{aligned}$$

input `Int [Coth[x]^4/(1 + Tanh[x]), x]`

output `((-5*Coth[x]^3)/3 + I*((-2*I)*Coth[x]^2 + I*(5*Coth[x] + I*((5*I)*x - (4*I)*Log[Sinh[x]]))))/2 + Coth[x]^3/(2*(1 + Tanh[x]))`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 26  $\text{Int}[(\text{Complex}[0, \text{a}_])*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], \text{a}]) \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2, 1]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3956  $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x\_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d}*x], \text{x}]]/\text{d}, \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4012  $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b}*c - \text{a}*d)*((\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}/(\text{f}*(\text{m} + 1)*(a^2 + b^2))), \text{x}] + \text{Simp}[1/(a^2 + b^2) \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}*\text{Simp}[\text{a}*c + \text{b}*d - (\text{b}*c - \text{a}*d)*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1]$
- rule 4014  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{a}*c + \text{b}*d)*(x/(a^2 + b^2)), \text{x}] + \text{Simp}[(\text{b}*c - \text{a}*d)/(a^2 + b^2) \quad \text{Int}[(\text{b} - \text{a}*\text{Tan}[\text{e} + \text{f}*x])/(a + \text{b}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{a}*c + \text{b}*d, 0]$
- rule 4035  $\text{Int}[(\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{n}_.}/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{a})*((\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} + 1}/(2*\text{f}*(\text{b}*c - \text{a}*d)*(a + \text{b}*\text{Tan}[\text{e} + \text{f}*x])), \text{x}] + \text{Simp}[1/(2*\text{a}*(\text{b}*c - \text{a}*d)) \quad \text{Int}[(\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{\text{n}}*\text{Simp}[\text{b}*c + \text{a}*d*(\text{n} - 1) - \text{b}*d*\text{n}*\text{Tan}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{EqQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!GtQ}[\text{n}, 0]$

**Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.02

method	result
risch	$\frac{9x}{2} - \frac{e^{-2x}}{4} - \frac{2(6e^{4x} - 9e^{2x} + 7)}{3(e^{2x} - 1)^3} - 2 \ln(e^{2x} - 1)$
parallelrisch	$\frac{(12 \tanh(x) + 12) \ln(1 - \tanh(x)) + (-12 \tanh(x) - 12) \ln(\tanh(x)) - 2 \coth(x)^3 + 27 \tanh(x)x + \coth(x)^2 + 27x - 9 \coth(x) - 15}{6 + 6 \tanh(x)}$
default	$-\frac{\tanh(\frac{x}{2})^3}{24} + \frac{\tanh(\frac{x}{2})^2}{8} - \frac{9 \tanh(\frac{x}{2})}{8} - \frac{\ln(\tanh(\frac{x}{2}) - 1)}{2} - \frac{1}{24 \tanh(\frac{x}{2})^3} + \frac{1}{8 \tanh(\frac{x}{2})^2} - \frac{9}{8 \tanh(\frac{x}{2})} - 2 \ln(t$

input `int(coth(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`output `9/2*x-1/4*exp(-2*x)-2/3*(6*exp(4*x)-9*exp(2*x)+7)/(exp(2*x)-1)^3-2*ln(exp(2*x)-1)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 582 vs. 2(35) = 70.

Time = 0.10 (sec) , antiderivative size = 582, normalized size of antiderivative = 13.53

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \text{Too large to display}$$

input `integrate(coth(x)^4/(1+tanh(x)),x, algorithm="fricas")`

output

```

1/12*(54*x*cosh(x)^8 + 432*x*cosh(x)*sinh(x)^7 + 54*x*sinh(x)^8 - 3*(54*x
+ 17)*cosh(x)^6 + 3*(504*x*cosh(x)^2 - 54*x - 17)*sinh(x)^6 + 18*(168*x*c
osh(x)^3 - (54*x + 17)*cosh(x))*sinh(x)^5 + 81*(2*x + 1)*cosh(x)^4 + 9*(420
*x*cosh(x)^4 - 5*(54*x + 17)*cosh(x)^2 + 18*x + 9)*sinh(x)^4 + 12*(252*x*c
osh(x)^5 - 5*(54*x + 17)*cosh(x)^3 + 27*(2*x + 1)*cosh(x))*sinh(x)^3 - (54
*x + 65)*cosh(x)^2 + (1512*x*cosh(x)^6 - 45*(54*x + 17)*cosh(x)^4 + 486*(2
*x + 1)*cosh(x)^2 - 54*x - 65)*sinh(x)^2 - 24*(cosh(x)^8 + 8*cosh(x)*sinh(
x)^7 + sinh(x)^8 + (28*cosh(x)^2 - 3)*sinh(x)^6 - 3*cosh(x)^6 + 2*(28*cosh
(x)^3 - 9*cosh(x))*sinh(x)^5 + (70*cosh(x)^4 - 45*cosh(x)^2 + 3)*sinh(x)^4
+ 3*cosh(x)^4 + 4*(14*cosh(x)^5 - 15*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + (
28*cosh(x)^6 - 45*cosh(x)^4 + 18*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*
(4*cosh(x)^7 - 9*cosh(x)^5 + 6*cosh(x)^3 - cosh(x))*sinh(x))*log(2*sinh(x)
/(cosh(x) - sinh(x))) + 2*(216*x*cosh(x)^7 - 9*(54*x + 17)*cosh(x)^5 + 162
*(2*x + 1)*cosh(x)^3 - (54*x + 65)*cosh(x))*sinh(x) + 3)/(cosh(x)^8 + 8*c
osh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 - 3)*sinh(x)^6 - 3*cosh(x)^6 +
2*(28*cosh(x)^3 - 9*cosh(x))*sinh(x)^5 + (70*cosh(x)^4 - 45*cosh(x)^2 + 3
)*sinh(x)^4 + 3*cosh(x)^4 + 4*(14*cosh(x)^5 - 15*cosh(x)^3 + 3*cosh(x))*si
nh(x)^3 + (28*cosh(x)^6 - 45*cosh(x)^4 + 18*cosh(x)^2 - 1)*sinh(x)^2 - cos
h(x)^2 + 2*(4*cosh(x)^7 - 9*cosh(x)^5 + 6*cosh(x)^3 - cosh(x))*sinh(x))

```

## Sympy [F]

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \int \frac{\coth^4(x)}{\tanh(x) + 1} dx$$

input

```
integrate(coth(x)**4/(1+tanh(x)),x)
```

output

```
Integral(coth(x)**4/(tanh(x) + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.49

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \frac{1}{2}x - \frac{2(15e^{-2x} - 12e^{-4x} - 7)}{3(3e^{-2x} - 3e^{-4x} + e^{-6x} - 1)} - \frac{1}{4}e^{-2x} - 2 \log(e^{-x} + 1) - 2 \log(e^{-x} - 1)$$

input `integrate(coth(x)^4/(1+tanh(x)),x, algorithm="maxima")`output `1/2*x - 2/3*(15*e^(-2*x) - 12*e^(-4*x) - 7)/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1) - 1/4*e^(-2*x) - 2*log(e^(-x) + 1) - 2*log(e^(-x) - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.12

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \frac{9}{2}x - \frac{(51e^{6x} - 81e^{4x} + 65e^{2x} - 3)e^{-2x}}{12(e^{2x} - 1)^3} - 2 \log(|e^{2x} - 1|)$$

input `integrate(coth(x)^4/(1+tanh(x)),x, algorithm="giac")`output `9/2*x - 1/12*(51*e^(6*x) - 81*e^(4*x) + 65*e^(2*x) - 3)*e^(-2*x)/(e^(2*x) - 1)^3 - 2*log(abs(e^(2*x) - 1))`**Mupad [B] (verification not implemented)**

Time = 2.08 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.60

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx = \frac{9x}{2} - 2 \ln(e^{2x} - 1) - \frac{e^{-2x}}{4} - \frac{8}{3(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{2}{e^{4x} - 2e^{2x} + 1} - \frac{4}{e^{2x} - 1}$$

input `int(coth(x)^4/(tanh(x) + 1),x)`

output  $(9x)/2 - 2\log(\exp(2x) - 1) - \exp(-2x)/4 - 8/(3(3\exp(2x) - 3\exp(4x) + \exp(6x) - 1)) - 2/(\exp(4x) - 2\exp(2x) + 1) - 4/(\exp(2x) - 1)$

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 191, normalized size of antiderivative = 4.44

$$\int \frac{\coth^4(x)}{1 + \tanh(x)} dx$$

$$= \frac{-24e^{8x}\log(e^x - 1) - 24e^{8x}\log(e^x + 1) + 54e^{8x}x - 17e^{8x} + 72e^{6x}\log(e^x - 1) + 72e^{6x}\log(e^x + 1) - 162e^{6x}}{12e^{2x}}$$

input `int(coth(x)^4/(1+tanh(x)),x)`

output  $(-24e^{8x}\log(e^{**x} - 1) - 24e^{8x}\log(e^{**x} + 1) + 54e^{8x}x - 17e^{8x} + 72e^{6x}\log(e^{**x} - 1) + 72e^{6x}\log(e^{**x} + 1) - 162e^{6x}x - 72e^{4x}\log(e^{**x} - 1) - 72e^{4x}\log(e^{**x} + 1) + 162e^{4x}x + 30e^{4x} + 24e^{2x}\log(e^{**x} - 1) + 24e^{2x}\log(e^{**x} + 1) - 54e^{2x}x - 48e^{2x} + 3)/(12e^{2x}(e^{**6x} - 3e^{**4x} + 3e^{**2x} - 1))$

### 3.126 $\int \tanh(x)(1 + \tanh(x))^{3/2} dx$

Optimal result	975
Mathematica [A] (verified)	975
Rubi [C] (verified)	976
Maple [A] (verified)	978
Fricas [B] (verification not implemented)	978
Sympy [A] (verification not implemented)	979
Maxima [F]	979
Giac [B] (verification not implemented)	980
Mupad [B] (verification not implemented)	980
Reduce [F]	981

#### Optimal result

Integrand size = 11, antiderivative size = 45

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}$$

output

```
2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2}{3}\sqrt{1 + \tanh(x)}(4 + \tanh(x))$$

input

```
Integrate[Tanh[x]*(1 + Tanh[x])^(3/2),x]
```



output

```
2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*Sqrt[1 + Tanh[x]]*(4 + T
anh[x]))/3
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.24, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 26, 4010, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(x)(\tanh(x) + 1)^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i(1 - i \tan(ix))^{3/2} \tan(ix) dx \\
 & \quad \downarrow \text{26} \\
 & -i \int (1 - i \tan(ix))^{3/2} \tan(ix) dx \\
 & \quad \downarrow \text{4010} \\
 & -i \left( i \int (\tanh(x) + 1)^{3/2} dx - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( i \int (1 - i \tan(ix))^{3/2} dx - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right) \\
 & \quad \downarrow \text{3959} \\
 & -i \left( i \left( 2 \int \sqrt{\tanh(x) + 1} dx - 2 \sqrt{\tanh(x) + 1} \right) - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( i \left( -2 \sqrt{\tanh(x) + 1} + 2 \int \sqrt{1 - i \tan(ix)} dx \right) - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3961} \\
 & -i \left( i \left( 4 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right) \\
 & \downarrow \text{219} \\
 & -i \left( i \left( 2\sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - 2\sqrt{\tanh(x) + 1} \right) - \frac{2}{3} i (\tanh(x) + 1)^{3/2} \right)
 \end{aligned}$$

input `Int[Tanh[x]*(1 + Tanh[x])^(3/2),x]`

output `(-I)*((( -2*I)/3)*(1 + Tanh[x])^(3/2) + I*(2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961

```
Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

rule 4010

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{\frac{3}{2}}}{3}$	35
default	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{\frac{3}{2}}}{3}$	35

input

```
int(tanh(x)*(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)-2/3*(
1+tanh(x))^(3/2)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 206 vs.  $2(34) = 68$ .

Time = 0.09 (sec) , antiderivative size = 206, normalized size of antiderivative = 4.58

$$\int \tanh(x) \left( 1 + \frac{3(\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}) \log\left(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2\right)}{2 \cosh(x)^2 + 2 \cosh(x) \sinh(x) + 2 \sinh(x)^2} \right)^{3/2} dx =$$

input `integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="fricas")`

output `1/3*(3*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(5*cosh(x)^3 + 15*cosh(x)*sinh(x)^2 + 5*sinh(x)^3 + 3*(5*cosh(x)^2 + 1)*sinh(x) + 3*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)`

### Sympy [A] (verification not implemented)

Time = 3.56 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx =$$

$$-\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)$$

$$- \frac{2(\tanh(x) + 1)^{3/2}}{3} - 2\sqrt{\tanh(x) + 1}$$

input `integrate(tanh(x)*(1+tanh(x))**(3/2),x)`

output `-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2))) - 2*(tanh(x) + 1)**(3/2)/3 - 2*sqrt(tanh(x) + 1)`

### Maxima [F]

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = \int (\tanh(x) + 1)^{3/2} \tanh(x) dx$$

input `integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="maxima")`

output `-2/3*sqrt(2)/(e^(-2*x) + 1)^(3/2) + integrate(2*sqrt(2)*e^(-x)/((e^(-x) + e^(-3*x))*(e^(-2*x) + 1)^(3/2)), x)`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs.  $2(34) = 68$ .

Time = 0.13 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.13

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = \frac{1}{3} \sqrt{2} \left( \frac{2 \left( 9 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 12 \sqrt{e^{4x} + e^{2x}} + 12 e^{2x} + 5 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} - 1 \right)^3} - 3 \log \left( -2 \sqrt{e^{4x} + e^{2x}} - e^{2x} + 1 \right) \right)$$

input `integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="giac")`

output `1/3*sqrt(2)*(2*(9*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 12*sqrt(e^(4*x) + e^(2*x)) + 12*e^(2*x) + 5)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1)^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

### Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2} \operatorname{atanh} \left( \frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right) - 2\sqrt{\tanh(x) + 1} - \frac{2(\tanh(x) + 1)^{3/2}}{3}$$

input `int(tanh(x)*(tanh(x) + 1)^(3/2),x)`

output `2*2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2) - (2*(tanh(x) + 1)^(3/2))/3`

**Reduce [F]**

$$\int \tanh(x)(1 + \tanh(x))^{3/2} dx = \int \sqrt{\tanh(x) + 1} \tanh(x)^2 dx \\ + \int \sqrt{\tanh(x) + 1} \tanh(x) dx$$

input `int(tanh(x)*(1+tanh(x))^(3/2),x)`

output `int(sqrt(tanh(x) + 1)*tanh(x)**2,x) + int(sqrt(tanh(x) + 1)*tanh(x),x)`

### 3.127 $\int \tanh(x) \sqrt{1 + \tanh(x)} dx$

Optimal result	982
Mathematica [A] (verified)	982
Rubi [C] (verified)	983
Maple [A] (verified)	984
Fricas [B] (verification not implemented)	985
Sympy [A] (verification not implemented)	986
Maxima [F]	986
Giac [B] (verification not implemented)	986
Mupad [B] (verification not implemented)	987
Reduce [F]	987

#### Optimal result

Integrand size = 11, antiderivative size = 32

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}$$

output

```
2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}$$

input

```
Integrate[Tanh[x]*Sqrt[1 + Tanh[x]], x]
```

output

```
Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.28, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 26, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(x) \sqrt{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \sqrt{1 - i \tan(ix)} \tan(ix) dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \sqrt{1 - i \tan(ix)} \tan(ix) dx \\
 & \quad \downarrow \text{4010} \\
 & -i \left( i \int \sqrt{\tanh(x) + 1} dx - 2i \sqrt{\tanh(x) + 1} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( i \int \sqrt{1 - i \tan(ix)} dx - 2i \sqrt{\tanh(x) + 1} \right) \\
 & \quad \downarrow \text{3961} \\
 & -i \left( 2i \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2i \sqrt{\tanh(x) + 1} \right) \\
 & \quad \downarrow \text{219} \\
 & -i \left( i\sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - 2i \sqrt{\tanh(x) + 1} \right)
 \end{aligned}$$

input

```
Int [Tanh [x] * Sqrt [1 + Tanh [x]] , x]
```



output  $(-I)*(I*\text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[1 + \text{Tanh}[x]]/\text{Sqrt}[2]] - (2*I)*\text{Sqrt}[1 + \text{Tanh}[x]])$

### Defintions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F x_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 219  $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3961  $\text{Int}[\text{Sqrt}[(a_) + (b_)*\tan[(c_) + (d_)*(x_)]], x\_Symbol] \rightarrow \text{Simp}[-2*(b/d) \text{Subst}[\text{Int}[1/(2*a - x^2), x], x, \text{Sqrt}[a + b*\tan[c + d*x]], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$

rule 4010  $\text{Int}[(a_) + (b_)*\tan[(e_) + (f_)*(x_)])^m * ((c_) + (d_)*\tan[(e_) + (f_)*(x_)]), x\_Symbol] \rightarrow \text{Simp}[d*((a + b*\tan[e + f*x])^m/(f*m)), x] + \text{Simp}[(b*c + a*d)/b \text{Int}[(a + b*\tan[e + f*x])^m, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ !\text{LtQ}[m, 0]$

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)}$	26
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)}$	26

input `int(tanh(x)*(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)`

output `2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs.  $2(25) = 50$ .

Time = 0.09 (sec) , antiderivative size = 128, normalized size of antiderivative = 4.00

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \frac{1}{2} \sqrt{2} \log \left( \frac{-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 + \sqrt{2}(\sqrt{2} \cosh(x)^3 + 3 \sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (3 \sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x) + \sqrt{2} \cosh(x))}{\sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}} - 1 \right) - \frac{2 \sqrt{2}(\cosh(x) + \sinh(x))}{\sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}}$$

input `integrate(tanh(x)*(1+tanh(x))^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(2)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(cosh(x) + sinh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)`

**Sympy [A] (verification not implemented)**

Time = 0.75 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.50

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx$$

$$= -\frac{\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)}{2} - 2\sqrt{\tanh(x) + 1}$$

input `integrate(tanh(x)*(1+tanh(x))**(1/2),x)`

output `-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/2 - 2*sqrt(tanh(x) + 1)`

**Maxima [F]**

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} \tanh(x) dx$$

input `integrate(tanh(x)*(1+tanh(x))^(1/2),x, algorithm="maxima")`

output `-sqrt(2)/sqrt(e^(-2*x) + 1) + integrate(sqrt(2)*e^(-x)/((e^(-x) + e^(-3*x))*sqrt(e^(-2*x) + 1)), x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(25) = 50.

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.66

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx$$

$$= \frac{1}{2} \sqrt{2} \left( \frac{4}{\sqrt{e^{(4x)} + e^{(2x)} - e^{(2x)} - 1}} - \log \left( -2 \sqrt{e^{(4x)} + e^{(2x)}} + 2e^{(2x)} + 1 \right) \right)$$

input `integrate(tanh(x)*(1+tanh(x))^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*(4/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1) - log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

### Mupad [B] (verification not implemented)

Time = 2.07 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{atanh} \left( \frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right) - 2 \sqrt{\tanh(x) + 1}$$

input `int(tanh(x)*(tanh(x) + 1)^(1/2),x)`

output `2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2)`

### Reduce [F]

$$\int \tanh(x) \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} \tanh(x) dx$$

input `int(tanh(x)*(1+tanh(x))^(1/2),x)`

output `int(sqrt(tanh(x) + 1)*tanh(x),x)`

### 3.128 $\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx$

Optimal result	988
Mathematica [A] (verified)	988
Rubi [C] (verified)	989
Maple [A] (verified)	991
Fricas [B] (verification not implemented)	991
Sympy [A] (verification not implemented)	992
Maxima [F]	992
Giac [B] (verification not implemented)	993
Mupad [B] (verification not implemented)	993
Reduce [F]	994

#### Optimal result

Integrand size = 11, antiderivative size = 30

$$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{\sqrt{1+\tanh(x)}}$$

output

```
1/2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))+1/(1+tanh(x))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{\sqrt{1+\tanh(x)}}$$

input

```
Integrate[Tanh[x]/Sqrt[1 + Tanh[x]], x]
```

output

```
ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] + 1/Sqrt[1 + Tanh[x]]
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 26, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{\sqrt{\tanh(x)+1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)}{\sqrt{1-i \tan(ix)}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)}{\sqrt{1-i \tan(ix)}} dx \\
 & \quad \downarrow \text{4009} \\
 & -i \left( \frac{1}{2} i \int \sqrt{\tanh(x)+1} dx + \frac{i}{\sqrt{\tanh(x)+1}} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{1}{2} i \int \sqrt{1-i \tan(ix)} dx + \frac{i}{\sqrt{\tanh(x)+1}} \right) \\
 & \quad \downarrow \text{3961} \\
 & -i \left( i \int \frac{1}{1-\tanh(x)} d\sqrt{\tanh(x)+1} + \frac{i}{\sqrt{\tanh(x)+1}} \right) \\
 & \quad \downarrow \text{219} \\
 & -i \left( \frac{i \operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{i}{\sqrt{\tanh(x)+1}} \right)
 \end{aligned}$$

input `Int [Tanh[x]/Sqrt[1 + Tanh[x]],x]`

output `(-I)*((I*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]])/Sqrt[2] + I/Sqrt[1 + Tanh[x]])`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

**Maple [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} + \frac{1}{\sqrt{1+\tanh(x)}}$	25
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} + \frac{1}{\sqrt{1+\tanh(x)}}$	25

input `int(tanh(x)/(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)`output `1/2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))+1/(1+tanh(x))^(1/2)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(24) = 48.

Time = 0.09 (sec) , antiderivative size = 141, normalized size of antiderivative = 4.70

$$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx$$

$$= \frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - \frac{\sqrt{2}(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x) + 3\sqrt{2} \sinh(x)^3)}{4(\cosh(x) + \sinh(x))}\right)}{4(\cosh(x) + \sinh(x))}$$

input `integrate(tanh(x)/(1+tanh(x))^(1/2),x, algorithm="fricas")`output `1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) + 2*sqrt(2)*sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x) + sinh(x))`



**Sympy [A] (verification not implemented)**

Time = 1.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.53

$$\int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx = \frac{\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)}{4} + \frac{1}{\sqrt{\tanh(x) + 1}}$$

input `integrate(tanh(x)/(1+tanh(x))**(1/2), x)`output `-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/4 + 1/sqrt(tanh(x) + 1)`**Maxima [F]**

$$\int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx = \int \frac{\tanh(x)}{\sqrt{\tanh(x) + 1}} dx$$

input `integrate(tanh(x)/(1+tanh(x))^(1/2), x, algorithm="maxima")`output `1/2*sqrt(2)*sqrt(e^(-2*x) + 1) + integrate(e^(-x)/(sqrt(2)*e^(-x)/sqrt(e^(-2*x) + 1) + sqrt(2)*e^(-3*x)/sqrt(e^(-2*x) + 1)), x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 52 vs.  $2(24) = 48$ .

Time = 0.13 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.73

$$\int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx$$

$$= \frac{1}{4} \sqrt{2} \left( \frac{2}{\sqrt{e^{4x} + e^{2x}} - e^{2x}} - \log \left( -2 \sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1 \right) \right)$$

input `integrate(tanh(x)/(1+tanh(x))^(1/2),x, algorithm="giac")`

output `1/4*sqrt(2)*(2/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x)) - log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx = \frac{\sqrt{2} \operatorname{atanh} \left( \frac{\sqrt{2} \sqrt{\tanh(x)+1}}{2} \right)}{2} + \frac{1}{\sqrt{\tanh(x) + 1}}$$

input `int(tanh(x)/(tanh(x) + 1)^(1/2),x)`

output `(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/2 + 1/(tanh(x) + 1)^(1/2)`

**Reduce [F]**

$$\int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx = \int \frac{\sqrt{\tanh(x) + 1} \tanh(x)}{\tanh(x) + 1} dx$$

input `int(tanh(x)/(1+tanh(x))^(1/2),x)`

output `int((sqrt(tanh(x) + 1)*tanh(x))/(tanh(x) + 1),x)`

**3.129**       $\int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx$

Optimal result	995
Mathematica [C] (verified)	995
Rubi [C] (verified)	996
Maple [A] (verified)	998
Fricas [B] (verification not implemented)	998
Sympy [A] (verification not implemented)	999
Maxima [F]	999
Giac [B] (verification not implemented)	1000
Mupad [B] (verification not implemented)	1000
Reduce [F]	1001

**Optimal result**

Integrand size = 11, antiderivative size = 49

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}}$$

output `1/4*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))+1/3/(1+tanh(x))^(3/2)-1/2/(1+tanh(x))^(1/2)`

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.39 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.73

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{2 - 3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1 + \tanh(x))\right) (1 + \tanh(x))}{6(1 + \tanh(x))^{3/2}}$$

input `Integrate[Tanh[x]/(1 + Tanh[x])^(3/2), x]`

output `(2 - 3*Hypergeometric2F1[-1/2, 1, 1/2, (1 + Tanh[x])/2]*(1 + Tanh[x]))/(6*(1 + Tanh[x])^(3/2))`

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.16, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 26, 4009, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{(\tanh(x) + 1)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{4009} \\
 & -i \left( \frac{1}{2} i \int \frac{1}{\sqrt{\tanh(x) + 1}} dx + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{1}{2} i \int \frac{1}{\sqrt{1 - i \tan(ix)}} dx + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right) \\
 & \quad \downarrow \text{3960} \\
 & -i \left( \frac{1}{2} i \left( \frac{1}{2} \int \sqrt{\tanh(x) + 1} dx - \frac{1}{\sqrt{\tanh(x) + 1}} \right) + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{1}{2} i \left( -\frac{1}{\sqrt{\tanh(x) + 1}} + \frac{1}{2} \int \sqrt{1 - i \tan(ix)} dx \right) + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right) \\
 & \quad \downarrow \text{3961}
 \end{aligned}$$

$$\begin{aligned}
 & -i \left( \frac{1}{2} i \left( \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}} \right) + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right) \\
 & \quad \downarrow \text{219} \\
 & -i \left( \frac{1}{2} i \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{\tanh(x) + 1}} \right) + \frac{i}{3(\tanh(x) + 1)^{3/2}} \right)
 \end{aligned}$$

input `Int[Tanh[x]/(1 + Tanh[x])^(3/2), x]`

output `(-I)*((I/3)/(1 + Tanh[x])^(3/2) + (I/2)*(ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]]))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} + \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{2\sqrt{1+\tanh(x)}}$	35
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} + \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} - \frac{1}{2\sqrt{1+\tanh(x)}}$	35

input

```
int(tanh(x)/(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
1/4*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))+1/3/(1+tanh(x))^(3/2)-1
/2/(1+tanh(x))^(1/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(34) = 68.

Time = 0.09 (sec) , antiderivative size = 240, normalized size of antiderivative = 4.90

$$\int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx = \frac{3(\sqrt{2} \cosh(x))^3 + 3\sqrt{2} \cosh(x)^2 \sinh(x) + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)}{...}$$

input

```
integrate(tanh(x)/(1+tanh(x))^(3/2),x, algorithm="fricas")
```

output

```
1/24*(3*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)^2*sinh(x) + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x)))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(2*cosh(x)^4 + 8*cosh(x)*sinh(x)^3 + 2*sinh(x)^4 + (12*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(4*cosh(x)^3 + cosh(x))*sinh(x) - 1)/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x)^3 + 3*cosh(x)^2*sinh(x) + 3*cosh(x)*sinh(x)^2 + sinh(x)^3)
```

**Sympy [A] (verification not implemented)**

Time = 4.73 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.22

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx =$$

$$\frac{\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)}{8}$$

$$- \frac{1}{2\sqrt{\tanh(x) + 1}} + \frac{1}{3(\tanh(x) + 1)^{3/2}}$$

input

```
integrate(tanh(x)/(1+tanh(x))**(3/2), x)
```

output

```
-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/8 - 1/(2*sqrt(tanh(x) + 1)) + 1/(3*(tanh(x) + 1)**(3/2))
```

**Maxima [F]**

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = \int \frac{\tanh(x)}{(\tanh(x) + 1)^{3/2}} dx$$

input

```
integrate(tanh(x)/(1+tanh(x))^(3/2), x, algorithm="maxima")
```



output `1/12*sqrt(2)*(e^(-2*x) + 1)^(3/2) + integrate(1/2*e^(-x)/(sqrt(2)*e^(-x)/(e^(-2*x) + 1)^(3/2) + sqrt(2)*e^(-3*x)/(e^(-2*x) + 1)^(3/2)), x)`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs.  $2(34) = 68$ .

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.49

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = -\frac{1}{24} \sqrt{2} \left( \frac{2 \left( 3 \sqrt{e^{4x} + e^{2x}} - 3e^{2x} - 1 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3} + 3 \log \left( -2 \sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1 \right) \right)$$

input `integrate(tanh(x)/(1+tanh(x))^(3/2),x, algorithm="giac")`

output `-1/24*sqrt(2)*(2*(3*sqrt(e^(4*x) + e^(2*x)) - 3*e^(2*x) - 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

### Mupad [B] (verification not implemented)

Time = 2.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.65

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x)+1}}{2}\right)}{4} - \frac{\frac{\tanh(x)}{2} + \frac{1}{6}}{(\tanh(x) + 1)^{3/2}}$$

input `int(tanh(x)/(tanh(x) + 1)^(3/2),x)`

output `(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 - (tanh(x)/2 + 1/6)/(tanh(x) + 1)^(3/2)`

**Reduce [F]**

$$\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx = \int \frac{\sqrt{\tanh(x) + 1} \tanh(x)}{\tanh(x)^2 + 2 \tanh(x) + 1} dx$$

input `int(tanh(x)/(1+tanh(x))^(3/2),x)`

output `int((sqrt(tanh(x) + 1)*tanh(x))/(tanh(x)**2 + 2*tanh(x) + 1),x)`

### 3.130 $\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx$

Optimal result	1002
Mathematica [A] (verified)	1002
Rubi [A] (verified)	1003
Maple [A] (verified)	1005
Fricas [B] (verification not implemented)	1005
Sympy [A] (verification not implemented)	1006
Maxima [F]	1007
Giac [B] (verification not implemented)	1007
Mupad [B] (verification not implemented)	1008
Reduce [F]	1008

#### Optimal result

Integrand size = 13, antiderivative size = 45

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2}$$

output

```
2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2*(1+tanh(x))^(1/2)-2/5*(1+tanh(x))^(5/2)
```

#### Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2}$$

input

```
Integrate[Tanh[x]^2*(1 + Tanh[x])^(3/2), x]
```

output

$$2\sqrt{2}\operatorname{ArcTanh}\left[\frac{\sqrt{1+\operatorname{Tanh}[x]}}{\sqrt{2}}\right] - 2\sqrt{1+\operatorname{Tanh}[x]} - \frac{2(1+\operatorname{Tanh}[x])^{5/2}}{5}$$
**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$ , Rules used = {3042, 25, 4026, 25, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^2(x)(\tanh(x) + 1)^{3/2} dx \\
 & \quad \downarrow 3042 \\
 & \int -(1 - i \tan(ix))^{3/2} \tan(ix)^2 dx \\
 & \quad \downarrow 25 \\
 & - \int (1 - i \tan(ix))^{3/2} \tan(ix)^2 dx \\
 & \quad \downarrow 4026 \\
 & - \int -(\tanh(x) + 1)^{3/2} dx - \frac{2}{5}(\tanh(x) + 1)^{5/2} \\
 & \quad \downarrow 25 \\
 & \int (\tanh(x) + 1)^{3/2} dx - \frac{2}{5}(\tanh(x) + 1)^{5/2} \\
 & \quad \downarrow 3042 \\
 & -\frac{2}{5}(\tanh(x) + 1)^{5/2} + \int (1 - i \tan(ix))^{3/2} dx \\
 & \quad \downarrow 3959 \\
 & 2 \int \sqrt{\tanh(x) + 1} dx - \frac{2}{5}(\tanh(x) + 1)^{5/2} - 2\sqrt{\tanh(x) + 1} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\begin{aligned}
& 2 \int \sqrt{1 - i \tan(ix)} dx - \frac{2}{5} (\tanh(x) + 1)^{5/2} - 2\sqrt{\tanh(x) + 1} \\
& \quad \downarrow \text{3961} \\
& 4 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - \frac{2}{5} (\tanh(x) + 1)^{5/2} - 2\sqrt{\tanh(x) + 1} \\
& \quad \downarrow \text{219} \\
& 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right) - \frac{2}{5} (\tanh(x) + 1)^{5/2} - 2\sqrt{\tanh(x) + 1}
\end{aligned}$$

input `Int[Tanh[x]^2*(1 + Tanh[x])^(3/2), x]`

output `2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]] - (2*(1 + Tanh[x])^(5/2))/5`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4026 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[d^2*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[c^2 - d^2 + 2*c*d*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1] && !(EqQ[m, 2] && EqQ[a, 0])`

### Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{5/2}}{5}$	35
default	$2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - 2\sqrt{1+\tanh(x)} - \frac{2(1+\tanh(x))^{5/2}}{5}$	35

input `int(tanh(x)^2*(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)`

output  $2*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-2*(1+\tanh(x))^{(1/2)}-2/5*(1+\tanh(x))^{(5/2)}$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 329 vs.  $2(34) = 68$ .

Time = 0.10 (sec) , antiderivative size = 329, normalized size of antiderivative = 7.31

$$\int \tanh^2(x) (1 + \tanh(x))^{3/2} dx = \frac{5(\sqrt{2} \cosh(x))^4 + 4\sqrt{2} \cosh(x) \sinh(x)^3 + \sqrt{2} \sinh(x)^4 + 2(3\sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x)}{\dots}$$

input `integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="fricas")`

output

```
1/5*(5*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(9*cosh(x)^5 + 45*cosh(x)*sinh(x)^4 + 9*sinh(x)^5 + 10*(9*cosh(x)^2 + 1)*sinh(x)^3 + 10*cosh(x)^3 + 30*(3*cosh(x)^3 + cosh(x))*sinh(x)^2 + 5*(9*cosh(x)^4 + 6*cosh(x)^2 + 1)*sinh(x) + 5*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)
```

### Sympy [A] (verification not implemented)

Time = 4.87 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx =$$

$$-\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)$$

$$- \frac{2(\tanh(x) + 1)^{5/2}}{5} - 2\sqrt{\tanh(x) + 1}$$

input `integrate(tanh(x)**2*(1+tanh(x))**(3/2),x)`

output

```
-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2))) - 2*(tanh(x) + 1)**(5/2)/5 - 2*sqrt(tanh(x) + 1)
```

**Maxima [F]**

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = \int (\tanh(x) + 1)^{\frac{3}{2}} \tanh(x)^2 dx$$

input `integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="maxima")`

output `integrate((tanh(x) + 1)^(3/2)*tanh(x)^2, x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 140 vs.  $2(34) = 68$ .

Time = 0.14 (sec) , antiderivative size = 140, normalized size of antiderivative = 3.11

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = \frac{1}{5} \sqrt{2} \left( \frac{2 \left( 25 \left( \sqrt{e^{4x}} + e^{2x} \right) - e^{2x} \right)^4 - 60 \left( \sqrt{e^{4x}} + e^{2x} \right) - e^{2x} \right)^3 + 70 \left( \sqrt{e^{4x}} + e^{2x} \right) - e^{2x} \right)^2 - 40 \sqrt{e^{4x}} + e^{2x} + 9}{\left( \sqrt{e^{4x}} + e^{2x} - 1 \right)^5} - 5 \log(-2 \sqrt{e^{4x}} + e^{2x}) + 2 e^{2x} + 1 \right)$$

input `integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="giac")`

output `1/5*sqrt(2)*(2*(25*(sqrt(e^(4*x)) + e^(2*x)) - e^(2*x))^4 - 60*(sqrt(e^(4*x)) + e^(2*x)) - e^(2*x))^3 + 70*(sqrt(e^(4*x)) + e^(2*x)) - e^(2*x))^2 - 40*sqrt(e^(4*x)) + e^(2*x) + 9)/(sqrt(e^(4*x)) + e^(2*x)) - e^(2*x) - 1)^5 - 5*log(-2*sqrt(e^(4*x)) + e^(2*x)) + 2*e^(2*x) + 1)`



**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = 2\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right) - 2\sqrt{\tanh(x)+1} - \frac{2(\tanh(x)+1)^{5/2}}{5}$$

input `int(tanh(x)^2*(tanh(x) + 1)^(3/2),x)`output `2*2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2) - (2*(tanh(x) + 1)^(5/2))/5`**Reduce [F]**

$$\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx = \int \sqrt{\tanh(x)+1} \tanh(x)^3 dx + \int \sqrt{\tanh(x)+1} \tanh(x)^2 dx$$

input `int(tanh(x)^2*(1+tanh(x))^(3/2),x)`output `int(sqrt(tanh(x) + 1)*tanh(x)**3,x) + int(sqrt(tanh(x) + 1)*tanh(x)**2,x)`

### 3.131 $\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx$

Optimal result	1009
Mathematica [A] (verified)	1009
Rubi [A] (verified)	1010
Maple [A] (verified)	1011
Fricas [B] (verification not implemented)	1012
Sympy [A] (verification not implemented)	1012
Maxima [F]	1013
Giac [B] (verification not implemented)	1013
Mupad [B] (verification not implemented)	1014
Reduce [F]	1014

#### Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2}{3} (1 + \tanh(x))^{3/2}$$

output `2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-2/3*(1+tanh(x))^(3/2)`

#### Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2}{3} (1 + \tanh(x))^{3/2}$$

input `Integrate[Tanh[x]^2*Sqrt[1 + Tanh[x]],x]`

output `Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*(1 + Tanh[x])^(3/2))/3`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {3042, 25, 4026, 25, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^2(x) \sqrt{\tanh(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sqrt{1 - i \tan(ix)} \tan(ix)^2 dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sqrt{1 - i \tan(ix)} \tan(ix)^2 dx \\
 & \quad \downarrow \text{4026} \\
 & - \int -\sqrt{\tanh(x) + 1} dx - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{25} \\
 & \int \sqrt{\tanh(x) + 1} dx - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{3} (\tanh(x) + 1)^{3/2} + \int \sqrt{1 - i \tan(ix)} dx \\
 & \quad \downarrow \text{3961} \\
 & 2 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - \frac{2}{3} (\tanh(x) + 1)^{3/2} \\
 & \quad \downarrow \text{219} \\
 & \sqrt{2} \operatorname{arctanh} \left( \frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2}
 \end{aligned}$$

input

```
Int [Tanh [x] ^2* Sqrt [1 + Tanh [x]] , x]
```

output  $\text{Sqrt}[2] \cdot \text{ArcTanh}[\text{Sqrt}[1 + \text{Tanh}[x]]/\text{Sqrt}[2]] - (2 \cdot (1 + \text{Tanh}[x])^{3/2})/3$

### Defintions of rubi rules used

rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$

rule 219  $\text{Int}[(\text{a}_) + (\text{b}_) \cdot (\text{x}_)^2]^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] \cdot \text{Rt}[-\text{b}, 2])) \cdot \text{ArcTanh}[\text{Rt}[-\text{b}, 2] \cdot (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}] \ \&\& \ (\text{GtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$

rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3961  $\text{Int}[\text{Sqrt}[(\text{a}_) + (\text{b}_) \cdot \tan[(\text{c}_) + (\text{d}_) \cdot (\text{x}_)]], \text{x\_Symbol}] \rightarrow \text{Simp}[-2 \cdot (\text{b}/\text{d}) \text{ Subst}[\text{Int}[1/(2 \cdot \text{a} - \text{x}^2), \text{x}], \text{x}, \text{Sqrt}[\text{a} + \text{b} \cdot \tan[\text{c} + \text{d} \cdot \text{x}]]], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{EqQ}[\text{a}^2 + \text{b}^2, 0]$

rule 4026  $\text{Int}[(\text{a}_) + (\text{b}_) \cdot \tan[(\text{e}_) + (\text{f}_) \cdot (\text{x}_)]^{\text{m}_} \cdot ((\text{c}_) + (\text{d}_) \cdot \tan[(\text{e}_) + (\text{f}_) \cdot (\text{x}_)]^2, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{d}^2 \cdot ((\text{a} + \text{b} \cdot \tan[\text{e} + \text{f} \cdot \text{x}])^{\text{m} + 1} / (\text{b} \cdot \text{f} \cdot (\text{m} + 1))), \text{x}] + \text{Int}[(\text{a} + \text{b} \cdot \tan[\text{e} + \text{f} \cdot \text{x}])^{\text{m}} \cdot \text{Simp}[\text{c}^2 - \text{d}^2 + 2 \cdot \text{c} \cdot \text{d} \cdot \tan[\text{e} + \text{f} \cdot \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b} \cdot \text{c} - \text{a} \cdot \text{d}, 0] \ \&\& \ !\text{LeQ}[\text{m}, -1] \ \&\& \ !(\text{EqQ}[\text{m}, 2] \ \&\& \ \text{EqQ}[\text{a}, 0])$

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

method	result	size
derivativedivides	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - \frac{2(1+\tanh(x))^{3/2}}{3}$	26
default	$\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right) - \frac{2(1+\tanh(x))^{3/2}}{3}$	26

input `int(tanh(x)^2*(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)`

output  $2^{1/2}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{1/2})*2^{1/2})-2/3*(1+\tanh(x))^{3/2}$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs.  $2(25) = 50$ .

Time = 0.09 (sec) , antiderivative size = 194, normalized size of antiderivative = 5.71

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx$$

$$= \frac{3(\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}) \log\left(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - \sinh(x)^2\right) - 6 \cosh(x) \sinh(x) \sqrt{2}}{6 \cosh(x) \sinh(x) \sqrt{2}}$$

input `integrate(tanh(x)^2*(1+tanh(x))^(1/2),x, algorithm="fricas")`

output 
$$\frac{1}{6} \frac{3(\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}) \log(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - \sinh(x)^2 - \sqrt{2} (\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (3\sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x) + \sqrt{2} \cosh(x)) / \sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}) - 1) - 8\sqrt{2} (\cosh(x)^3 + 3 \cosh(x)^2 \sinh(x) + 3 \cosh(x) \sinh(x)^2 + \sinh(x)^3) / \sqrt{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}}{6 \cosh(x) \sinh(x) \sqrt{2}}$$

### Sympy [A] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.44

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx$$

$$= -\frac{\sqrt{2} \left( \log\left(\sqrt{\tanh(x)+1} - \sqrt{2}\right) - \log\left(\sqrt{\tanh(x)+1} + \sqrt{2}\right) \right)}{2} - \frac{2(\tanh(x)+1)^{\frac{3}{2}}}{3}$$

input `integrate(tanh(x)**2*(1+tanh(x))**(1/2),x)`

output `-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/2 - 2*(tanh(x) + 1)**(3/2)/3`

### Maxima [F]

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} \tanh(x)^2 dx$$

input `integrate(tanh(x)^2*(1+tanh(x))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(tanh(x) + 1)*tanh(x)^2, x)`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs.  $2(25) = 50$ .

Time = 0.14 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.82

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx$$

$$= \frac{1}{6} \sqrt{2} \left( \frac{8 \left( 3 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 3 \sqrt{e^{4x} + e^{2x}} + 3 e^{2x} + 1 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} - 1 \right)^3} - 3 \log \left( -2 \sqrt{e^{4x} + e^{2x}} + 2 \right) \right)$$

input `integrate(tanh(x)^2*(1+tanh(x))^(1/2),x, algorithm="giac")`

output `1/6*sqrt(2)*(8*(3*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 3*sqrt(e^(4*x) + e^(2*x)) + 3*e^(2*x) + 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1)^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx = \sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right) - \frac{2(\tanh(x) + 1)^{3/2}}{3}$$

input `int(tanh(x)^2*(tanh(x) + 1)^(1/2),x)`

output `2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - (2*(tanh(x) + 1)^(3/2))/3`

**Reduce [F]**

$$\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx = \int \sqrt{\tanh(x) + 1} \tanh(x)^2 dx$$

input `int(tanh(x)^2*(1+tanh(x))^(1/2),x)`

output `int(sqrt(tanh(x) + 1)*tanh(x)**2,x)`

**3.132**  $\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx$

Optimal result	1015
Mathematica [C] (verified)	1015
Rubi [A] (verified)	1016
Maple [A] (verified)	1018
Fricas [B] (verification not implemented)	1018
Sympy [A] (verification not implemented)	1019
Maxima [F]	1019
Giac [A] (verification not implemented)	1020
Mupad [B] (verification not implemented)	1020
Reduce [F]	1020

**Optimal result**

Integrand size = 13, antiderivative size = 42

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1+\tanh(x)}} - 2\sqrt{1+\tanh(x)}$$

output

$1/2*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-1/(1+\tanh(x))^{(1/2)}-2*(1+\tanh(x))^{(1/2)}$

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.48 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx = \frac{-\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1+\tanh(x))\right) - 2(1+\tanh(x))}{\sqrt{1+\tanh(x)}}$$

input

`Integrate[Tanh[x]^2/Sqrt[1 + Tanh[x]], x]`



output

```
(-Hypergeometric2F1[-1/2, 1, 1/2, (1 + Tanh[x])/2] - 2*(1 + Tanh[x]))/Sqrt
[1 + Tanh[x]]
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$ , Rules used = {3042, 25, 4026, 25, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{\sqrt{\tanh(x)+1}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan(ix)^2}{\sqrt{1-i\tan(ix)}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan(ix)^2}{\sqrt{1-i\tan(ix)}} dx \\
 & \quad \downarrow \text{4026} \\
 & -\int -\frac{1}{\sqrt{\tanh(x)+1}} dx - 2\sqrt{\tanh(x)+1} \\
 & \quad \downarrow \text{25} \\
 & \int \frac{1}{\sqrt{\tanh(x)+1}} dx - 2\sqrt{\tanh(x)+1} \\
 & \quad \downarrow \text{3042} \\
 & -2\sqrt{\tanh(x)+1} + \int \frac{1}{\sqrt{1-i\tan(ix)}} dx \\
 & \quad \downarrow \text{3960} \\
 & \frac{1}{2} \int \sqrt{\tanh(x)+1} dx - 2\sqrt{\tanh(x)+1} - \frac{1}{\sqrt{\tanh(x)+1}}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 \frac{1}{2} \int \sqrt{1 - i \tanh(ix)} dx - 2\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}} \\
 \downarrow \text{3961} \\
 \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} - 2\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}} \\
 \downarrow \text{219} \\
 \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - 2\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}}
 \end{array}$$

input `Int [Tanh[x]^2/Sqrt[1 + Tanh[x]], x]`

output `ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]] - 2*Sqrt[1 + Tanh[x]]`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

```
rule 3961 Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
  , b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4026 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
  (f_)*(x_)])^2, x_Symbol] := Simp[d^2*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(
  m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[c^2 - d^2 + 2*c*d*Tan[e + f*
  x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && !LeQ
  [m, -1] && !(EqQ[m, 2] && EqQ[a, 0])
```

**Maple [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} - \frac{1}{\sqrt{1+\tanh(x)}} - 2\sqrt{1+\tanh(x)}$	35
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{2} - \frac{1}{\sqrt{1+\tanh(x)}} - 2\sqrt{1+\tanh(x)}$	35

```
input int(tanh(x)^2/(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/(1+tanh(x))^(1/2)-2*(
  1+tanh(x))^(1/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 161 vs. 2(34) = 68.

Time = 0.12 (sec) , antiderivative size = 161, normalized size of antiderivative = 3.83

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx$$

$$= \frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(-2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - \frac{\sqrt{2}(\sqrt{2} \cosh(x)^3 + 3 \sqrt{2} \cosh(x) \sinh(x) + 3 \sqrt{2} \sinh(x)^3)}{4(\cosh(x) - \sinh(x))}\right)}{4(\cosh(x) - \sinh(x))} + C$$

input `integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="fricas")`

output `1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) - 2*sqrt(2)*(5*cosh(x)^2 + 10*cosh(x)*sinh(x) + 5*sinh(x)^2 + 1)/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x) + sinh(x))`

### Sympy [A] (verification not implemented)

Time = 1.38 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.38

$$\int \frac{\tanh^2(x)}{\sqrt{1 + \tanh(x)}} dx = \frac{\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)}{-2\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}}}$$

input `integrate(tanh(x)**2/(1+tanh(x))**(1/2),x)`

output `-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/4 - 2*sqrt(tanh(x) + 1) - 1/sqrt(tanh(x) + 1)`

### Maxima [F]

$$\int \frac{\tanh^2(x)}{\sqrt{1 + \tanh(x)}} dx = \int \frac{\tanh(x)^2}{\sqrt{\tanh(x) + 1}} dx$$

input `integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)^2/sqrt(tanh(x) + 1), x)`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.29

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx = -\frac{1}{4}\sqrt{2}\log\left(-4\sqrt{e^{4x}+e^{2x}}+4e^{2x}+2\right) - \frac{5\sqrt{2}e^{2x}+\sqrt{2}}{2\sqrt{e^{4x}+e^{2x}}}$$

input `integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="giac")`

output `-1/4*sqrt(2)*log(-4*sqrt(e^(4*x) + e^(2*x)) + 4*e^(2*x) + 2) - 1/2*(5*sqrt(2)*e^(2*x) + sqrt(2))/sqrt(e^(4*x) + e^(2*x))`

**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx = \frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} - \frac{3}{\sqrt{\tanh(x)+1}} - \frac{2\tanh(x)}{\sqrt{\tanh(x)+1}}$$

input `int(tanh(x)^2/(tanh(x) + 1)^(1/2),x)`

output `(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/2 - 3/(tanh(x) + 1)^(1/2) - (2*tanh(x))/(tanh(x) + 1)^(1/2)`

**Reduce [F]**

$$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx = \int \frac{\sqrt{\tanh(x)+1}\tanh(x)^2}{\tanh(x)+1} dx$$

input `int(tanh(x)^2/(1+tanh(x))^(1/2),x)`

output `int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x) + 1),x)`

### 3.133 $\int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx$

Optimal result	1021
Mathematica [A] (verified)	1021
Rubi [A] (verified)	1022
Maple [A] (verified)	1024
Fricas [B] (verification not implemented)	1024
Sympy [A] (verification not implemented)	1025
Maxima [F]	1025
Giac [B] (verification not implemented)	1026
Mupad [B] (verification not implemented)	1026
Reduce [F]	1027

#### Optimal result

Integrand size = 13, antiderivative size = 49

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{3}{2\sqrt{1 + \tanh(x)}}$$

output

$1/4*2^{(1/2)}*\operatorname{arctanh}(1/2*(1+\tanh(x))^{(1/2)}*2^{(1/2)})-1/3/(1+\tanh(x))^{(3/2)}+3/2/(1+\tanh(x))^{(1/2)}$

#### Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.98

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{14 + 18 \tanh(x) + 3\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{12(1 + \tanh(x))^{3/2}} (1 + \tanh(x))^{3/2}$$

input

`Integrate[Tanh[x]^2/(1 + Tanh[x])^(3/2), x]`

output

$(14 + 18*\operatorname{Tanh}[x] + 3*\operatorname{Sqrt}[2]*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 + \operatorname{Tanh}[x]]/\operatorname{Sqrt}[2]]*(1 + \operatorname{Tanh}[x])^{(3/2)})/(12*(1 + \operatorname{Tanh}[x])^{(3/2)})$

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$ , Rules used = {3042, 25, 4023, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{(\tanh(x) + 1)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan(ix)^2}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\tan(ix)^2}{(1 - i \tan(ix))^{3/2}} dx \\
 & \quad \downarrow \text{4023} \\
 & -\frac{1}{2} \int \frac{1 - 2 \tanh(x)}{\sqrt{\tanh(x) + 1}} dx - \frac{1}{3(\tanh(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{3(\tanh(x) + 1)^{3/2}} - \frac{1}{2} \int \frac{2i \tan(ix) + 1}{\sqrt{1 - i \tan(ix)}} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{1}{2} \left( \frac{1}{2} \int \sqrt{\tanh(x) + 1} dx + \frac{3}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{3(\tanh(x) + 1)^{3/2}} + \frac{1}{2} \left( \frac{3}{\sqrt{\tanh(x) + 1}} + \frac{1}{2} \int \sqrt{1 - i \tan(ix)} dx \right) \\
 & \quad \downarrow \text{3961} \\
 & \frac{1}{2} \left( \int \frac{1}{1 - \tanh(x)} d\sqrt{\tanh(x) + 1} + \frac{3}{\sqrt{\tanh(x) + 1}} \right) - \frac{1}{3(\tanh(x) + 1)^{3/2}}
 \end{aligned}$$

$$\frac{1}{2} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{3}{\sqrt{\tanh(x)+1}} \right) - \frac{1}{3(\tanh(x)+1)^{3/2}}$$

input `Int [Tanh[x]^2/(1 + Tanh[x])^(3/2), x]`

output `-1/3*1/(1 + Tanh[x])^(3/2) + (ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] + 3/Sqrt[1 + Tanh[x]])/2`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))* ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`



rule 4023

```
Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_) +
(f_.)*(x_)])^2, x_Symbol] := Simp[(-b)*(a*c + b*d)^(2*(a + b*Tan[e + f*x])^
m/(2*a^3*f*m)), x] + Simp[1/(2*a^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp
[a*c^2 - 2*b*c*d + a*d^2 - 2*b*d^2*Tan[e + f*x], x], x], x] /; FreeQ[{a, b,
c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LeQ[m, -1] && EqQ[a^2 + b^2, 0]
```

**Maple [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.71

method	result	size
derivativedivides	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} - \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} + \frac{3}{2\sqrt{1+\tanh(x)}}$	35
default	$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)}{4} - \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}} + \frac{3}{2\sqrt{1+\tanh(x)}}$	35

input

```
int(tanh(x)^2/(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
1/4*2^(1/2)*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))-1/3/(1+tanh(x))^(3/2)+3
/2/(1+tanh(x))^(1/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. 2(34) = 68.

Time = 0.11 (sec) , antiderivative size = 244, normalized size of antiderivative = 4.98

$$\int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx = \frac{3(\sqrt{2} \cosh(x))^3 + 3\sqrt{2} \cosh(x)^2 \sinh(x) + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)}{...}$$

input

```
integrate(tanh(x)^2/(1+tanh(x))^(3/2),x, algorithm="fricas")
```

output

```
1/24*(3*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)^2*sinh(x) + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3)*log(-2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + sqrt(2)*cosh(x)))/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1) - 1) + 2*sqrt(2)*(8*cosh(x)^4 + 32*cosh(x)*sinh(x)^3 + 8*sinh(x)^4 + (48*cosh(x)^2 + 7)*sinh(x)^2 + 7*cosh(x)^2 + 2*(16*cosh(x)^3 + 7*cosh(x))*sinh(x) - 1)/sqrt(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))/(cosh(x)^3 + 3*cosh(x)^2*sinh(x) + 3*cosh(x)*sinh(x)^2 + sinh(x)^3)
```

### Sympy [A] (verification not implemented)

Time = 5.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.22

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{\sqrt{2} \left( \log \left( \sqrt{\tanh(x) + 1} - \sqrt{2} \right) - \log \left( \sqrt{\tanh(x) + 1} + \sqrt{2} \right) \right)}{8} + \frac{3}{2\sqrt{\tanh(x) + 1}} - \frac{1}{3(\tanh(x) + 1)^{3/2}}$$

input

```
integrate(tanh(x)**2/(1+tanh(x))**(3/2), x)
```

output

```
-sqrt(2)*(log(sqrt(tanh(x) + 1) - sqrt(2)) - log(sqrt(tanh(x) + 1) + sqrt(2)))/8 + 3/(2*sqrt(tanh(x) + 1)) - 1/(3*(tanh(x) + 1)**(3/2))
```

### Maxima [F]

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \int \frac{\tanh(x)^2}{(\tanh(x) + 1)^{3/2}} dx$$

input

```
integrate(tanh(x)^2/(1+tanh(x))^(3/2), x, algorithm="maxima")
```

output

```
integrate(tanh(x)^2/(tanh(x) + 1)^(3/2), x)
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 95 vs.  $2(34) = 68$ .

Time = 0.13 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.94

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{1}{24} \sqrt{2} \left( \frac{2 \left( 6 \left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 + 3 \sqrt{e^{4x} + e^{2x}} - 3e^{2x} - 1 \right)}{\left( \sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3} - 3 \log \right)$$

input `integrate(tanh(x)^2/(1+tanh(x))^(3/2),x, algorithm="giac")`

output `1/24*sqrt(2)*(2*(6*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 + 3*sqrt(e^(4*x) + e^(2*x)) - 3*e^(2*x) - 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))`

**Mupad [B] (verification not implemented)**

Time = 2.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.63

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x)+1}}{2}\right)}{4} + \frac{\frac{3 \tanh(x)}{2} + \frac{7}{6}}{(\tanh(x) + 1)^{3/2}}$$

input `int(tanh(x)^2/(tanh(x) + 1)^(3/2),x)`

output `(2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 + ((3*tanh(x))/2 + 7/6)/(tanh(x) + 1)^(3/2)`

**Reduce [F]**

$$\int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx = \int \frac{\sqrt{\tanh(x) + 1} \tanh(x)^2}{\tanh(x)^2 + 2 \tanh(x) + 1} dx$$

input `int(tanh(x)^2/(1+tanh(x))^(3/2),x)`

output `int((sqrt(tanh(x) + 1)*tanh(x)**2)/(tanh(x)**2 + 2*tanh(x) + 1),x)`

### 3.134 $\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx$

Optimal result	1028
Mathematica [A] (verified)	1028
Rubi [C] (verified)	1029
Maple [A] (verified)	1037
Fricas [B] (verification not implemented)	1038
Sympy [B] (verification not implemented)	1039
Maxima [A] (verification not implemented)	1039
Giac [A] (verification not implemented)	1040
Mupad [B] (verification not implemented)	1040
Reduce [B] (verification not implemented)	1041

#### Optimal result

Integrand size = 13, antiderivative size = 94

$$\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx = -\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^5 \log(a+b \tanh(x))}{b^4(a^2-b^2)} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b}$$

output

`-b*x/(a^2-b^2)+a*ln(cosh(x))/(a^2-b^2)+a^5*ln(a+b*tanh(x))/b^4/(a^2-b^2)-(a^2+b^2)*tanh(x)/b^3+1/2*a*tanh(x)^2/b^2-1/3*tanh(x)^3/b`

#### Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.98

$$\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx = \frac{1}{6} \left( -\frac{3 \log(1-\tanh(x))}{a+b} - \frac{3 \log(1+\tanh(x))}{a-b} + \frac{6a^5 \log(a+b \tanh(x))}{b^4(a^2-b^2)} - \frac{6(a^2+b^2) \tanh(x)}{b^3} + \frac{3a \tanh^2(x)}{b^2} - \frac{2 \tanh^3(x)}{b} \right)$$

input `Integrate[Tanh[x]^5/(a + b*Tanh[x]),x]`

output  $((-3*\text{Log}[1 - \text{Tanh}[x]])/(a + b) - (3*\text{Log}[1 + \text{Tanh}[x]])/(a - b) + (6*a^5*\text{Log}[a + b*\text{Tanh}[x]])/(b^4*(a^2 - b^2)) - (6*(a^2 + b^2)*\text{Tanh}[x])/b^3 + (3*a*\text{Tanh}[x]^2)/b^2 - (2*\text{Tanh}[x]^3)/b)/6$

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.07 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.39, number of steps used = 21, number of rules used = 20,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.538$ , Rules used = {3042, 26, 4049, 27, 3042, 25, 4130, 27, 3042, 26, 4131, 25, 3042, 4109, 26, 3042, 26, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^5(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{i \tan(ix)^5}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{26} \\ & -i \int \frac{\tan(ix)^5}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{4049} \\ & -i \left( \frac{i \int \frac{3 \tanh^2(x)(-a \tanh^2(x) + b \tanh(x) + a)}{a + b \tanh(x)} dx}{3b} - \frac{i \tanh^3(x)}{3b} \right) \\ & \quad \downarrow \text{27} \\ & -i \left( \frac{i \int \frac{\tanh^2(x)(-a \tanh^2(x) + b \tanh(x) + a)}{a + b \tanh(x)} dx}{b} - \frac{i \tanh^3(x)}{3b} \right) \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& -i \left( \frac{i \int -\frac{\tan(ix)^2 (a \tan(ix)^2 - ib \tan(ix) + a)}{a - ib \tan(ix)} dx}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 25 \\
& -i \left( -\frac{i \int \frac{\tan(ix)^2 (a \tan(ix)^2 - ib \tan(ix) + a)}{a - ib \tan(ix)} dx}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 4130 \\
& -i \left( -\frac{i \left( -\frac{a \tanh^2(x)}{2b} + \frac{i \int -\frac{2i \tanh(x) (a^2 - (a^2 + b^2) \tanh^2(x))}{a + b \tanh(x)} dx}{2b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 27 \\
& -i \left( -\frac{i \left( \frac{\int \frac{\tanh(x) (a^2 - (a^2 + b^2) \tanh^2(x))}{a + b \tanh(x)} dx}{b} - \frac{a \tanh^2(x)}{2b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 3042 \\
& -i \left( -\frac{i \left( -\frac{a \tanh^2(x)}{2b} + \frac{\int -\frac{i \tan(ix) (a^2 + (a^2 + b^2) \tan(ix)^2)}{a - ib \tan(ix)} dx}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 26 \\
& -i \left( -\frac{i \left( -\frac{a \tanh^2(x)}{2b} - \frac{i \int \frac{\tan(ix) (a^2 + (a^2 + b^2) \tan(ix)^2)}{a - ib \tan(ix)} dx}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right) \\
& \downarrow 4131
\end{aligned}$$

$$-i \left( \frac{i \left( -\frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{\int -\tanh(x)b^3 - a(a^2+b^2)\tanh^2(x) + a(a^2+b^2) dx}{a+b \tanh(x)} + \frac{i(a^2+b^2)\tanh(x)}{b} \right)}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right)$$

↓ 25

$$-i \left( \frac{i \left( -\frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{i(a^2+b^2)\tanh(x)}{b} - \frac{\int \tanh(x)b^3 - a(a^2+b^2)\tanh^2(x) + a(a^2+b^2) dx}{a+b \tanh(x)} \right)}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right)$$

↓ 3042

$$-i \left( \frac{i \left( -\frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{i(a^2+b^2)\tanh(x)}{b} - \frac{\int -i \tan(ix)b^3 + a(a^2+b^2)\tan(ix)^2 + a(a^2+b^2) dx}{a-ib \tan(ix)} \right)}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right)$$

↓ 4109



$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( -\frac{iab^3 \int i \tanh(x) dx}{a^2-b^2} + \frac{a^5 \int \frac{1-\tanh^2(x)}{a+b \tanh(x)} dx - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \frac{\phantom{i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( -\frac{iab^3 \int i \tanh(x) dx}{a^2-b^2} + \frac{a^5 \int \frac{1-\tanh^2(x)}{a+b \tanh(x)} dx - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right)}}{b} \end{array} \right) - \frac{i \tanh^3(x)}{3b}$$

↓ 26

$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{ab^3 \int \tanh(x) dx}{a^2-b^2} + \frac{a^5 \int \frac{1-\tanh^2(x)}{a+b \tanh(x)} dx - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \frac{\phantom{i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{ab^3 \int \tanh(x) dx}{a^2-b^2} + \frac{a^5 \int \frac{1-\tanh^2(x)}{a+b \tanh(x)} dx - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right)}}{b} \end{array} \right) - \frac{i \tanh^3(x)}{3b}$$

↓ 3042

$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{ab^3 \int -i \tan(ix) dx}{a^2-b^2} + \frac{a^5 \int \frac{\tan(ix)^2+1}{a-ib \tan(ix)} dx}{a^2-b^2} - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \end{array} \right) \frac{i \tanh^3(x)}{3b}$$

↓ 26

$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( -\frac{iab^3 \int \tan(ix) dx}{a^2-b^2} + \frac{a^5 \int \frac{\tan(ix)^2+1}{a-ib \tan(ix)} dx}{a^2-b^2} - \frac{b^4 x}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \end{array} \right) \frac{i \tanh^3(x)}{3b}$$

↓ 3956

$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{a^5 \int \frac{\tan(ix)^2+1}{a-ib \tan(ix)} dx}{a^2-b^2} - \frac{b^4 x}{a^2-b^2} + \frac{ab^3 \log(\cosh(x))}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \frac{\phantom{i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{a^5 \int \frac{\tan(ix)^2+1}{a-ib \tan(ix)} dx}{a^2-b^2} - \frac{b^4 x}{a^2-b^2} + \frac{ab^3 \log(\cosh(x))}{a^2-b^2} \right)}{b} \right)}{b} \right)}{b} \end{array} \right) - \frac{i \tanh^3(x)}{3b}$$

↓ 4100

$$\left( \begin{array}{c} i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{a^5 \int \frac{1}{a+b \tanh(x)} d(b \tanh(x))}{b(a^2-b^2)} - \frac{b^4 x}{a^2-b^2} + \frac{ab^3 \log(\cosh(x))}{a^2-b^2} \right)}{b} \right)}{b} \right) \\ -i \frac{\phantom{i \left( \frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{(a^2+b^2) \tanh(x)}{b} - \frac{i \left( \frac{a^5 \int \frac{1}{a+b \tanh(x)} d(b \tanh(x))}{b(a^2-b^2)} - \frac{b^4 x}{a^2-b^2} + \frac{ab^3 \log(\cosh(x))}{a^2-b^2} \right)}{b} \right)}{b} \right)}{b} \end{array} \right) - \frac{i \tanh^3(x)}{3b}$$

↓ 16

$$-i \left( \frac{i \left( -\frac{a \tanh^2(x)}{2b} - \frac{i \left( \frac{i(a^2+b^2) \tanh(x)}{b} - \frac{i \left( -\frac{b^4 x}{a^2-b^2} + \frac{ab^3 \log(\cosh(x))}{a^2-b^2} + \frac{a^5 \log(a+b \tanh(x))}{b(a^2-b^2)} \right)}{b} \right)}{b} \right)}{b} - \frac{i \tanh^3(x)}{3b} \right)$$

input `Int [Tanh[x]^5/(a + b*Tanh[x]),x]`

output `(-I)*((( (-1/3*I)*Tanh[x]^3)/b - (I*(-1/2*(a*Tanh[x]^2)/b - (I*((( -I)*(-(b^4*x)/(a^2 - b^2)) + (a*b^3*Log[Cosh[x]])/(a^2 - b^2) + (a^5*Log[a + b*Tanh[x]])/(b*(a^2 - b^2)))))/b + (I*(a^2 + b^2)*Tanh[x]/b)/b)/b)`

**Defintions of rubi rules used**

rule 16 `Int[(c.)/((a.) + (b.)*(x.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a])* (Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4049 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m - 2)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n - 1))), x] + Simp[1/(d*(m + n - 1)) Int[(a + b*Tan[e + f*x])^(m - 3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m + n - 1) - b^2*(b*c*(m - 2) + a*d*(1 + n)) + b*d*(m + n - 1)*(3*a^2 - b^2)*Tan[e + f*x] - b^2*(b*c*(m - 2) - a*d*(3*m + 2*n - 4))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || IntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && ( !IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4131

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n +
1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d
*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b
- b*C)*(m + n + 1)*Tan[e + f*x] - C*m*(b*c - a*d)*Tan[e + f*x]^2, x], x
] /; FreeQ[{a, b, c, d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2
+ b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && ( !IntegerQ
[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

## Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{\tanh(x)^3}{3b} + \frac{a \tanh(x)^2}{2b^2} - \frac{a^2 \tanh(x)}{b^3} - \frac{\tanh(x)}{b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{a^5 \ln(a+b \tanh(x))}{b^4(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b}$
default	$-\frac{\tanh(x)^3}{3b} + \frac{a \tanh(x)^2}{2b^2} - \frac{a^2 \tanh(x)}{b^3} - \frac{\tanh(x)}{b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{a^5 \ln(a+b \tanh(x))}{b^4(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b}$
parallelrisch	$-\frac{2 \tanh(x)^3 a^2 b^3 + 2 \tanh(x)^3 b^5 + 3 \tanh(x)^2 b^2 a^3 - 3 \tanh(x)^2 a b^4 + 6 a^5 \ln(a+b \tanh(x)) - 6 \ln(1-\tanh(x)) a b^4 - 6 a b^4 x}{6 b^4 (a^2 - b^2)}$
risch	$\frac{x}{a+b} + \frac{2x a^3}{b^4} + \frac{2ax}{b^2} - \frac{2x a^5}{b^4(a^2-b^2)} + \frac{2a^2 e^{4x} - 2e^{4x} ab + 4b^2 e^{4x} + 4e^{2x} a^2 - 2e^{2x} ab + 4b^2 e^{2x} + 2a^2 + \frac{8b^2}{3}}{b^3(e^{2x}+1)^3} - \frac{a^3 \ln(e^{2x})}{b^4}$

input

```
int(tanh(x)^5/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-1/3*tanh(x)^3/b+1/2*a*tanh(x)^2/b^2-1/b^3*a^2*tanh(x)-tanh(x)/b-1/(2*b+2*
a)*ln(tanh(x)-1)+1/b^4*a^5/(a+b)/(a-b)*ln(a+b*tanh(x))-1/(2*a-2*b)*ln(1+ta
nh(x))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1296 vs.  $2(90) = 180$ .

Time = 0.16 (sec) , antiderivative size = 1296, normalized size of antiderivative = 13.79

$$\int \frac{\tanh^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```
-1/3*(3*(a*b^4 + b^5)*x*cosh(x)^6 + 18*(a*b^4 + b^5)*x*cosh(x)*sinh(x)^5 +
3*(a*b^4 + b^5)*x*sinh(x)^6 - 6*a^4*b - 2*a^2*b^3 + 8*b^5 - 3*(2*a^4*b -
2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*cosh(x)^4 - 3
*(2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 15*(a*b^4 + b^5)*x*c
osh(x)^2 - 3*(a*b^4 + b^5)*x)*sinh(x)^4 + 12*(5*(a*b^4 + b^5)*x*cosh(x)^3
- (2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*
cosh(x))*sinh(x)^3 - 3*(4*a^4*b - 2*a^3*b^2 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 +
b^5)*x)*cosh(x)^2 + 3*(15*(a*b^4 + b^5)*x*cosh(x)^4 - 4*a^4*b + 2*a^3*b^2
- 2*a*b^4 + 4*b^5 - 6*(2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5
- 3*(a*b^4 + b^5)*x)*cosh(x)^2 + 3*(a*b^4 + b^5)*x)*sinh(x)^2 + 3*(a*b^4 +
b^5)*x - 3*(a^5*cosh(x)^6 + 6*a^5*cosh(x)*sinh(x)^5 + a^5*sinh(x)^6 + 3*a
^5*cosh(x)^4 + 3*a^5*cosh(x)^2 + a^5 + 3*(5*a^5*cosh(x)^2 + a^5)*sinh(x)^4
+ 4*(5*a^5*cosh(x)^3 + 3*a^5*cosh(x))*sinh(x)^3 + 3*(5*a^5*cosh(x)^4 + 6*
a^5*cosh(x)^2 + a^5)*sinh(x)^2 + 6*(a^5*cosh(x)^5 + 2*a^5*cosh(x)^3 + a^5*
cosh(x))*sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + 3*(
(a^5 - a*b^4)*cosh(x)^6 + 6*(a^5 - a*b^4)*cosh(x)*sinh(x)^5 + (a^5 - a*b^4
)*sinh(x)^6 + a^5 - a*b^4 + 3*(a^5 - a*b^4)*cosh(x)^4 + 3*(a^5 - a*b^4 + 5
*(a^5 - a*b^4)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a^5 - a*b^4)*cosh(x)^3 + 3*(a^
5 - a*b^4)*cosh(x))*sinh(x)^3 + 3*(a^5 - a*b^4)*cosh(x)^2 + 3*(a^5 - a*b^4
+ 5*(a^5 - a*b^4)*cosh(x)^4 + 6*(a^5 - a*b^4)*cosh(x)^2)*sinh(x)^2 + 6...
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 546 vs.  $2(78) = 156$ .

Time = 0.47 (sec) , antiderivative size = 546, normalized size of antiderivative = 5.81

$$\int \frac{\tanh^5(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(tanh(x)**5/(a+b*tanh(x)),x)`

output

```
Piecewise((zoo*(x - tanh(x)**3/3 - tanh(x)), Eq(a, 0) & Eq(b, 0)), ((x - 1
og(tanh(x) + 1) - tanh(x)**4/4 - tanh(x)**2/2)/a, Eq(b, 0)), (27*x*tanh(x)
/(6*b*tanh(x) - 6*b) - 27*x/(6*b*tanh(x) - 6*b) - 12*log(tanh(x) + 1)*tanh
(x)/(6*b*tanh(x) - 6*b) + 12*log(tanh(x) + 1)/(6*b*tanh(x) - 6*b) - 2*tanh
(x)**4/(6*b*tanh(x) - 6*b) - tanh(x)**3/(6*b*tanh(x) - 6*b) - 9*tanh(x)**2
/(6*b*tanh(x) - 6*b) + 15/(6*b*tanh(x) - 6*b), Eq(a, -b)), (3*x*tanh(x)/(6
*b*tanh(x) + 6*b) + 3*x/(6*b*tanh(x) + 6*b) + 12*log(tanh(x) + 1)*tanh(x)/
(6*b*tanh(x) + 6*b) + 12*log(tanh(x) + 1)/(6*b*tanh(x) + 6*b) - 2*tanh(x)*
*4/(6*b*tanh(x) + 6*b) + tanh(x)**3/(6*b*tanh(x) + 6*b) - 9*tanh(x)**2/(6*
b*tanh(x) + 6*b) + 15/(6*b*tanh(x) + 6*b), Eq(a, b)), (6*a**5*log(a/b + ta
nh(x))/(6*a**2*b**4 - 6*b**6) - 6*a**4*b*tanh(x)/(6*a**2*b**4 - 6*b**6) +
3*a**3*b**2*tanh(x)**2/(6*a**2*b**4 - 6*b**6) - 2*a**2*b**3*tanh(x)**3/(6*
a**2*b**4 - 6*b**6) + 6*a*b**4*x/(6*a**2*b**4 - 6*b**6) - 6*a*b**4*log(tan
h(x) + 1)/(6*a**2*b**4 - 6*b**6) - 3*a*b**4*tanh(x)**2/(6*a**2*b**4 - 6*b*
*6) - 6*b**5*x/(6*a**2*b**4 - 6*b**6) + 2*b**5*tanh(x)**3/(6*a**2*b**4 - 6
*b**6) + 6*b**5*tanh(x)/(6*a**2*b**4 - 6*b**6), True))
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.60

$$\begin{aligned} & \int \frac{\tanh^5(x)}{a + b \tanh(x)} dx \\ &= \frac{a^5 \log(-(a-b)e^{-2x} - a - b)}{a^2 b^4 - b^6} \\ & \quad - \frac{2(3a^2 + 4b^2 + 3(2a^2 + ab + 2b^2)e^{-2x} + 3(a^2 + ab + 2b^2)e^{-4x})}{3(3b^3e^{-2x} + 3b^3e^{-4x} + b^3e^{-6x} + b^3)} \\ & \quad + \frac{x}{a+b} - \frac{(a^3 + ab^2) \log(e^{-2x} + 1)}{b^4} \end{aligned}$$



input `integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="maxima")`

output  $a^5 \log(-(a-b)e^{-2x} - a - b)/(a^2 b^4 - b^6) - 2/3(3a^2 + 4b^2 + 3(2a^2 + ab + 2b^2)e^{-2x} + 3(a^2 + ab + 2b^2)e^{-4x})/(3b^3 e^{-2x} + 3b^3 e^{-4x} + b^3 e^{-6x} + b^3) + x/(a+b) - (a^3 + ab^2) \log(e^{-2x} + 1)/b^4$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.51

$$\int \frac{\tanh^5(x)}{a + b \tanh(x)} dx = \frac{a^5 \log(|ae^{2x} + be^{2x} + a - b|)}{a^2 b^4 - b^6} - \frac{x}{a - b} - \frac{(a^3 + ab^2) \log(e^{2x} + 1)}{b^4} + \frac{2(3a^2 b + 4b^3 + 3(a^2 b - ab^2 + 2b^3)e^{4x}) + 3(2a^2 b - ab^2 + 2b^3)e^{2x}}{3b^4(e^{2x} + 1)^3}$$

input `integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="giac")`

output  $a^5 \log(\text{abs}(a e^{2x} + b e^{2x} + a - b))/(a^2 b^4 - b^6) - x/(a - b) - (a^3 + ab^2) \log(e^{2x} + 1)/b^4 + 2/3(3a^2 b + 4b^3 + 3(a^2 b - ab^2 + 2b^3)e^{4x} + 3(2a^2 b - ab^2 + 2b^3)e^{2x})/(b^4(e^{2x} + 1)^3)$

### Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.90

$$\int \frac{\tanh^5(x)}{a + b \tanh(x)} dx = \frac{x}{a + b} - \frac{\tanh(x)^3}{3b} - \frac{a \ln(\tanh(x) + 1)}{a^2 - b^2} + \frac{a \tanh(x)^2}{2b^2} - \frac{\tanh(x)(a^2 + b^2)}{b^3} + \frac{a^5 \ln(a + b \tanh(x))}{b^4(a^2 - b^2)}$$

input `int(tanh(x)^5/(a + b*tanh(x)),x)`

output

$$\frac{x}{a+b} - \frac{\tanh(x)^3}{3b} - \frac{(a \log(\tanh(x) + 1))}{(a^2 - b^2)} + \frac{(a \tanh(x))^2}{(2b^2)} - \frac{(\tanh(x)(a^2 + b^2))}{b^3} + \frac{(a^5 \log(a + b \tanh(x)))}{(b^4(a^2 - b^2))}$$
**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.48

$$\int \frac{\tanh^5(x)}{a + b \tanh(x)} dx$$

$$= \frac{6 \log(e^{2x} a + e^{2x} b + a - b) a b^4 + 6 \log(\tanh(x) b + a) a^5 - 6 \log(\tanh(x) b + a) a b^4 - 2 \tanh(x)^3 a^2 b^3 + \dots}{6b^4(a^2 - b^2)}$$

input

`int(tanh(x)^5/(a+b*tanh(x)),x)`

output

$$\frac{(6 \log(e^{2x} a + e^{2x} b + a - b) a b^4 + 6 \log(\tanh(x) b + a) a^5 - 6 \log(\tanh(x) b + a) a b^4 - 2 \tanh(x)^3 a^2 b^3 + 2 \tanh(x)^3 b^5 + 3 \tanh(x)^2 a^3 b^2 - 3 \tanh(x)^2 a b^4 - 6 \tanh(x) a^4 b + 6 \tanh(x) b^5 - 6 a b^4 x - 6 b^5 x)}{(6 b^4 (a^2 - b^2))}$$

### 3.135 $\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx$

Optimal result	1042
Mathematica [A] (verified)	1042
Rubi [C] (verified)	1043
Maple [A] (verified)	1047
Fricas [B] (verification not implemented)	1048
Sympy [B] (verification not implemented)	1049
Maxima [A] (verification not implemented)	1050
Giac [A] (verification not implemented)	1050
Mupad [B] (verification not implemented)	1051
Reduce [B] (verification not implemented)	1051

#### Optimal result

Integrand size = 13, antiderivative size = 76

$$\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx = \frac{ax}{a^2-b^2} - \frac{b \log(\cosh(x))}{a^2-b^2} - \frac{a^4 \log(a+b \tanh(x))}{b^3(a^2-b^2)} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

output

```
a*x/(a^2-b^2)-b*ln(cosh(x))/(a^2-b^2)-a^4*ln(a+b*tanh(x))/b^3/(a^2-b^2)+a*tanh(x)/b^2-1/2*tanh(x)^2/b
```

#### Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.01

$$\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx = -\frac{\log(1-\tanh(x))}{2(a+b)} + \frac{\log(1+\tanh(x))}{2(a-b)} - \frac{a^4 \log(a+b \tanh(x))}{b^3(a^2-b^2)} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

input

```
Integrate[Tanh[x]^4/(a + b*Tanh[x]),x]
```

output

$$-1/2*\text{Log}[1 - \text{Tanh}[x]]/(a + b) + \text{Log}[1 + \text{Tanh}[x]]/(2*(a - b)) - (a^4*\text{Log}[a + b*\text{Tanh}[x]])/(b^3*(a^2 - b^2)) + (a*\text{Tanh}[x])/b^2 - \text{Tanh}[x]^2/(2*b)$$

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.79 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.30, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.154$ , Rules used = {3042, 4049, 27, 3042, 26, 4130, 25, 3042, 4110, 26, 3042, 26, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^4(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(ix)^4}{a - ib \tan(ix)} dx \\ & \quad \downarrow \text{4049} \\ & -\frac{\tanh^2(x)}{2b} + \frac{i \int -\frac{2i \tanh(x)(-a \tanh^2(x) + b \tanh(x) + a)}{a + b \tanh(x)} dx}{2b} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\tanh(x)(-a \tanh^2(x) + b \tanh(x) + a)}{a + b \tanh(x)} dx}{b} - \frac{\tanh^2(x)}{2b} \\ & \quad \downarrow \text{3042} \\ & -\frac{\tanh^2(x)}{2b} + \frac{\int -\frac{i \tan(ix)(a \tan(ix)^2 - ib \tan(ix) + a)}{a - ib \tan(ix)} dx}{b} \\ & \quad \downarrow \text{26} \\ & -\frac{\tanh^2(x)}{2b} - \frac{i \int \frac{\tan(ix)(a \tan(ix)^2 - ib \tan(ix) + a)}{a - ib \tan(ix)} dx}{b} \\ & \quad \downarrow \text{4130} \end{aligned}$$

$$\begin{aligned}
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{i \int -\frac{a^2 - (a^2 + b^2) \tanh^2(x)}{a + b \tanh(x)} dx}{b} + \frac{ia \tanh(x)}{b} \right)}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \int \frac{a^2 - (a^2 + b^2) \tanh^2(x)}{a + b \tanh(x)} dx}{b} \right)}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \int \frac{a^2 + (a^2 + b^2) \tan(ix)^2}{a - ib \tan(ix)} dx}{b} \right)}{b} \\
 & \quad \downarrow \text{4110} \\
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( -\frac{ib^3 \int i \tanh(x) dx}{a^2 - b^2} + \frac{a^4 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{a^2 - b^2} - \frac{ab^2 x}{a^2 - b^2} \right)}{b} \right)}{b} \\
 & \quad \downarrow \text{26} \\
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( \frac{b^3 \int \tanh(x) dx}{a^2 - b^2} + \frac{a^4 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{a^2 - b^2} - \frac{ab^2 x}{a^2 - b^2} \right)}{b} \right)}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( \frac{b^3 \int -i \tan(ix) dx}{a^2 - b^2} + \frac{a^4 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx}{a^2 - b^2} - \frac{ab^2 x}{a^2 - b^2} \right)}{b} \right)}{b} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( -\frac{ib^3 \int \tan(ix) dx}{a^2 - b^2} + \frac{a^4 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx - \frac{ab^2 x}{a^2 - b^2} \right)}{b} \right)}{b}$$

↓ 3956

$$\frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( \frac{a^4 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx - \frac{ab^2 x}{a^2 - b^2} + \frac{b^3 \log(\cosh(x))}{a^2 - b^2} \right)}{b} \right)}{b}$$

↓ 4100

$$\frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( \frac{a^4 \int \frac{1}{a + b \tanh(x)} d(b \tanh(x))}{b(a^2 - b^2)} - \frac{ab^2 x}{a^2 - b^2} + \frac{b^3 \log(\cosh(x))}{a^2 - b^2} \right)}{b} \right)}{b}$$

↓ 16

$$\frac{\tanh^2(x)}{2b} - \frac{i \left( \frac{ia \tanh(x)}{b} - \frac{i \left( -\frac{ab^2 x}{a^2 - b^2} + \frac{b^3 \log(\cosh(x))}{a^2 - b^2} + \frac{a^4 \log(a + b \tanh(x))}{b(a^2 - b^2)} \right)}{b} \right)}{b}$$

input `Int [Tanh [x]^4/(a + b*Tanh [x]), x]`

output `-1/2*Tanh[x]^2/b - (I*((( -I)*(-(a*b^2*x)/(a^2 - b^2)) + (b^3*Log[Cosh[x]])/(a^2 - b^2) + (a^4*Log[a + b*Tanh[x]]/(b*(a^2 - b^2)))))/b + (I*a*Tanh[x])/b)/b`

## Definitions of rubi rules used

- rule 16  $\text{Int}[(c\_)/((a\_)+(b\_)*(x\_)), x\_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25  $\text{Int}[-(Fx\_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 26  $\text{Int}[(\text{Complex}[0, a\_])*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 27  $\text{Int}[(a\_)*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b\_)*(Gx\_)] /; \text{FreeQ}[b, x]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956  $\text{Int}[\tan[(c\_)+(d\_)*(x\_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4049  $\text{Int}[(a\_)+(b\_)*\tan[(e\_)+(f\_)*(x\_)]^{(m\_)}*((c\_)+(d\_)*\tan[(e\_)+(f\_)*(x\_)]^{(n\_)}), x\_Symbol] \rightarrow \text{Simp}[b^2*(a + b*\tan[e + f*x])^{(m-2)}*((c + d*\tan[e + f*x])^{(n+1)}/(d*f*(m+n-1))), x] + \text{Simp}[1/(d*(m+n-1)) \text{ Int}[(a + b*\tan[e + f*x])^{(m-3)}*(c + d*\tan[e + f*x])^n*\text{Simp}[a^3*d*(m+n-1) - b^2*(b*c*(m-2) + a*d*(1+n)) + b*d*(m+n-1)*(3*a^2 - b^2)*\tan[e + f*x] - b^2*(b*c*(m-2) - a*d*(3*m+2*n-4))*\tan[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{IntegerQ}[2*m] \ \&\& \ \text{GtQ}[m, 2] \ \&\& \ (\text{GeQ}[n, -1] \ || \ \text{IntegerQ}[m]) \ \&\& \ !(\text{IGtQ}[n, 2] \ \&\& \ (!\text{IntegerQ}[m] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{NeQ}[a, 0])))$
- rule 4100  $\text{Int}[(a\_)+(b\_)*\tan[(e\_)+(f\_)*(x\_)]^{(m\_)}*((A\_)+(C\_)*\tan[(e\_)+(f\_)*(x\_)]^2), x\_Symbol] \rightarrow \text{Simp}[A/(b*f) \ \text{Subst}[\text{Int}[(a + x)^m, x], x, b*\tan[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$

rule 4110

```
Int[((A_) + (C_)*tan[(e_) + (f_)*(x_)]^2)/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[a*(A - C)*(x/(a^2 + b^2)), x] + (Simp[(a^2*C +
A*b^2)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x]
- Simp[b*((A - C)/(a^2 + b^2)) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b,
e, f, A, C}, x] && NeQ[a^2*C + A*b^2, 0] && NeQ[a^2 + b^2, 0] && NeQ[A, C]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*(c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1)), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

### Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00

method	result
derivativedivides	$-\frac{\tanh(x)^2}{2b} + \frac{a \tanh(x)}{b^2} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{a^4 \ln(a+b \tanh(x))}{b^3(a+b)(a-b)} + \frac{\ln(1+\tanh(x))}{2a-2b}$
default	$-\frac{\tanh(x)^2}{2b} + \frac{a \tanh(x)}{b^2} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{a^4 \ln(a+b \tanh(x))}{b^3(a+b)(a-b)} + \frac{\ln(1+\tanh(x))}{2a-2b}$
parallelrisc	$-\frac{\tanh(x)^2 a^2 b^2 - \tanh(x)^2 b^4 - 2 \ln(1 - \tanh(x)) b^4 + 2 a^4 \ln(a + b \tanh(x)) - 2 a b^3 x - 2 b^4 x - 2 \tanh(x) a^3 b + 2 \tanh(x) a b^3}{2 b^3 (a^2 - b^2)}$
risc	$\frac{x}{a+b} - \frac{2x a^2}{b^3} - \frac{2x}{b} + \frac{2x a^4}{b^3(a^2-b^2)} - \frac{2(e^{2x} a - e^{2x} b + a)}{(e^{2x} + 1)^2 b^2} + \frac{\ln(e^{2x} + 1) a^2}{b^3} + \frac{\ln(e^{2x} + 1)}{b} - \frac{a^4 \ln(e^{2x} + \frac{a-b}{a+b})}{b^3(a^2-b^2)}$

input

```
int(tanh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*tanh(x)^2/b+a*tanh(x)/b^2-1/(2*b+2*a)*ln(tanh(x)-1)-1/b^3*a^4/(a+b)/(
a-b)*ln(a+b*tanh(x))+1/(2*a-2*b)*ln(1+tanh(x))
```



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 644 vs.  $2(74) = 148$ .

Time = 0.13 (sec) , antiderivative size = 644, normalized size of antiderivative = 8.47

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

output

```
((a*b^3 + b^4)*x*cosh(x)^4 + 4*(a*b^3 + b^4)*x*cosh(x)*sinh(x)^3 + (a*b^3
+ b^4)*x*sinh(x)^4 - 2*a^3*b + 2*a*b^3 - 2*(a^3*b - a^2*b^2 - a*b^3 + b^4
- (a*b^3 + b^4)*x)*cosh(x)^2 - 2*(a^3*b - a^2*b^2 - a*b^3 + b^4 - 3*(a*b^3
+ b^4)*x*cosh(x)^2 - (a*b^3 + b^4)*x)*sinh(x)^2 + (a*b^3 + b^4)*x - (a^4*
cosh(x)^4 + 4*a^4*cosh(x)*sinh(x)^3 + a^4*sinh(x)^4 + 2*a^4*cosh(x)^2 + a^
4 + 2*(3*a^4*cosh(x)^2 + a^4)*sinh(x)^2 + 4*(a^4*cosh(x)^3 + a^4*cosh(x))*
sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + ((a^4 - b^4)
*cosh(x)^4 + 4*(a^4 - b^4)*cosh(x)*sinh(x)^3 + (a^4 - b^4)*sinh(x)^4 + a^4
- b^4 + 2*(a^4 - b^4)*cosh(x)^2 + 2*(a^4 - b^4 + 3*(a^4 - b^4)*cosh(x)^2)
*sinh(x)^2 + 4*((a^4 - b^4)*cosh(x)^3 + (a^4 - b^4)*cosh(x))*sinh(x))*log(
2*cosh(x)/(cosh(x) - sinh(x))) + 4*((a*b^3 + b^4)*x*cosh(x)^3 - (a^3*b - a
^2*b^2 - a*b^3 + b^4 - (a*b^3 + b^4)*x)*cosh(x))*sinh(x))/(a^2*b^3 - b^5 +
(a^2*b^3 - b^5)*cosh(x)^4 + 4*(a^2*b^3 - b^5)*cosh(x)*sinh(x)^3 + (a^2*b^
3 - b^5)*sinh(x)^4 + 2*(a^2*b^3 - b^5)*cosh(x)^2 + 2*(a^2*b^3 - b^5 + 3*(a
^2*b^3 - b^5)*cosh(x)^2)*sinh(x)^2 + 4*((a^2*b^3 - b^5)*cosh(x)^3 + (a^2*b
^3 - b^5)*cosh(x))*sinh(x))
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 442 vs.  $2(61) = 122$ .

Time = 0.40 (sec) , antiderivative size = 442, normalized size of antiderivative = 5.82

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx$$

$$= \begin{cases} \tilde{\infty} \left( x - \log(\tanh(x) + 1) - \frac{\tanh^2(x)}{2} \right) \\ \frac{x - \frac{\tanh^3(x)}{3} - \tanh(x)}{a} \\ \frac{7x \tanh(x)}{2b \tanh(x) - 2b} - \frac{7x}{2b \tanh(x) - 2b} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) - 2b} + \frac{4 \log(\tanh(x) + 1)}{2b \tanh(x) - 2b} - \frac{\tanh^3(x)}{2b \tanh(x) - 2b} - \frac{\tanh^2(x)}{2b \tanh(x) - 2b} + \frac{\tanh(x)}{2b \tanh(x) - 2b} \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) + 2b} - \frac{4 \log(\tanh(x) + 1)}{2b \tanh(x) + 2b} - \frac{\tanh^3(x)}{2b \tanh(x) + 2b} + \frac{\tanh^2(x)}{2b \tanh(x) + 2b} - \frac{\tanh(x)}{2b \tanh(x) + 2b} \\ - \frac{2a^4 \log\left(\frac{a}{b} + \tanh(x)\right)}{2a^2 b^3 - 2b^5} + \frac{2a^3 b \tanh(x)}{2a^2 b^3 - 2b^5} - \frac{a^2 b^2 \tanh^2(x)}{2a^2 b^3 - 2b^5} + \frac{2ab^3 x}{2a^2 b^3 - 2b^5} - \frac{2ab^3 \tanh(x)}{2a^2 b^3 - 2b^5} - \frac{2b^4 x}{2a^2 b^3 - 2b^5} + \frac{2b^4 \log(\tanh(x) + 1)}{2a^2 b^3 - 2b^5} + \end{cases}$$

input `integrate(tanh(x)**4/(a+b*tanh(x)), x)`

output

```
Piecewise((zoo*(x - log(tanh(x) + 1) - tanh(x)**2/2), Eq(a, 0) & Eq(b, 0)),
((x - tanh(x)**3/3 - tanh(x))/a, Eq(b, 0)), (7*x*tanh(x)/(2*b*tanh(x) -
2*b) - 7*x/(2*b*tanh(x) - 2*b) - 4*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) -
2*b) + 4*log(tanh(x) + 1)/(2*b*tanh(x) - 2*b) - tanh(x)**3/(2*b*tanh(x) -
2*b) - tanh(x)**2/(2*b*tanh(x) - 2*b) + 3/(2*b*tanh(x) - 2*b), Eq(a, -b)),
(x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 4*log(tanh(x) +
1)*tanh(x)/(2*b*tanh(x) + 2*b) - 4*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) -
tanh(x)**3/(2*b*tanh(x) + 2*b) + tanh(x)**2/(2*b*tanh(x) + 2*b) - 3/(2*b*
tanh(x) + 2*b), Eq(a, b)), (-2*a**4*log(a/b + tanh(x))/(2*a**2*b**3 - 2*b*
**5) + 2*a**3*b*tanh(x)/(2*a**2*b**3 - 2*b**5) - a**2*b**2*tanh(x)**2/(2*a*
**2*b**3 - 2*b**5) + 2*a*b**3*x/(2*a**2*b**3 - 2*b**5) - 2*a*b**3*tanh(x)/(
2*a**2*b**3 - 2*b**5) - 2*b**4*x/(2*a**2*b**3 - 2*b**5) + 2*b**4*log(tanh(
x) + 1)/(2*a**2*b**3 - 2*b**5) + b**4*tanh(x)**2/(2*a**2*b**3 - 2*b**5), T
rue))
```

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.32

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx = -\frac{a^4 \log(-(a-b)e^{-2x} - a - b)}{a^2 b^3 - b^5} + \frac{2((a+b)e^{-2x} + a)}{2b^2 e^{-2x} + b^2 e^{-4x} + b^2} + \frac{x}{a+b} + \frac{(a^2 + b^2) \log(e^{-2x} + 1)}{b^3}$$

input `integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

output `-a^4*log(-(a - b)*e^(-2*x) - a - b)/(a^2*b^3 - b^5) + 2*((a + b)*e^(-2*x) + a)/(2*b^2*e^(-2*x) + b^2*e^(-4*x) + b^2) + x/(a + b) + (a^2 + b^2)*log(e^(-2*x) + 1)/b^3`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.29

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx = -\frac{a^4 \log(|ae^{2x} + be^{2x} + a - b|)}{a^2 b^3 - b^5} + \frac{x}{a - b} + \frac{(a^2 + b^2) \log(e^{2x} + 1)}{b^3} - \frac{2(ab + (ab - b^2)e^{2x})}{b^3(e^{2x} + 1)^2}$$

input `integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="giac")`

output `-a^4*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2*b^3 - b^5) + x/(a - b) + (a^2 + b^2)*log(e^(2*x) + 1)/b^3 - 2*(a*b + (a*b - b^2)*e^(2*x))/(b^3*(e^(2*x) + 1)^2)`

**Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx = \frac{x}{a + b} - \frac{\tanh(x)^2}{2b} + \frac{b \ln(\tanh(x) + 1)}{a^2 - b^2} + \frac{a \tanh(x)}{b^2} - \frac{a^4 \ln(a + b \tanh(x))}{b^3 (a^2 - b^2)}$$

input `int(tanh(x)^4/(a + b*tanh(x)),x)`output `x/(a + b) - tanh(x)^2/(2*b) + (b*log(tanh(x) + 1))/(a^2 - b^2) + (a*tanh(x))/b^2 - (a^4*log(a + b*tanh(x)))/(b^3*(a^2 - b^2))`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.51

$$\int \frac{\tanh^4(x)}{a + b \tanh(x)} dx = \frac{-2 \log(e^{2x}a + e^{2x}b + a - b) b^4 - 2 \log(\tanh(x) b + a) a^4 + 2 \log(\tanh(x) b + a) b^4 - \tanh(x)^2 a^2 b^2 + \tanh(x)^2 a^2 b^2}{2b^3 (a^2 - b^2)}$$

input `int(tanh(x)^4/(a+b*tanh(x)),x)`output `( - 2*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**4 - 2*log(tanh(x)*b + a)*a**4 + 2*log(tanh(x)*b + a)*b**4 - tanh(x)**2*a**2*b**2 + tanh(x)**2*b**4 + 2*tanh(x)*a**3*b - 2*tanh(x)*a*b**3 + 2*a*b**3*x + 2*b**4*x)/(2*b**3*(a**2 - b**2))`

### 3.136 $\int \frac{\tanh^3(x)}{a+b \tanh(x)} dx$

Optimal result	1052
Mathematica [A] (verified)	1052
Rubi [C] (verified)	1053
Maple [A] (verified)	1056
Fricas [B] (verification not implemented)	1057
Sympy [B] (verification not implemented)	1057
Maxima [A] (verification not implemented)	1058
Giac [A] (verification not implemented)	1059
Mupad [B] (verification not implemented)	1059
Reduce [B] (verification not implemented)	1059

#### Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{\tanh^3(x)}{a+b \tanh(x)} dx = -\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^3 \log(a+b \tanh(x))}{b^2(a^2-b^2)} - \frac{\tanh(x)}{b}$$

output

$$-\frac{b*x}{(a^2-b^2)} + \frac{a*\ln(\cosh(x))}{(a^2-b^2)} + \frac{a^3*\ln(a+b*\tanh(x))}{b^2*(a^2-b^2)} - \frac{\tanh(x)}{b}$$

#### Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

$$\int \frac{\tanh^3(x)}{a+b \tanh(x)} dx = -\frac{\log(1-\tanh(x))}{2(a+b)} - \frac{\log(1+\tanh(x))}{2(a-b)} + \frac{a^3 \log(a+b \tanh(x))}{b^2(a^2-b^2)} - \frac{\tanh(x)}{b}$$

input

`Integrate[Tanh[x]^3/(a + b*Tanh[x]), x]`

output

$$-\frac{1}{2}*\text{Log}[1 - \text{Tanh}[x]]/(a + b) - \text{Log}[1 + \text{Tanh}[x]]/(2*(a - b)) + (a^3*\text{Log}[a + b*\text{Tanh}[x]])/(b^2*(a^2 - b^2)) - \text{Tanh}[x]/b$$

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.27, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.923$ , Rules used = {3042, 26, 4049, 25, 3042, 4109, 26, 3042, 26, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^3(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \tan(ix)^3}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\tan(ix)^3}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{4049} \\
 & i \left( \frac{i \int \frac{-a \tanh^2(x) + b \tanh(x) + a}{a + b \tanh(x)} dx}{b} + \frac{i \tanh(x)}{b} \right) \\
 & \quad \downarrow \text{25} \\
 & i \left( \frac{i \tanh(x)}{b} - \frac{i \int \frac{-a \tanh^2(x) + b \tanh(x) + a}{a + b \tanh(x)} dx}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & i \left( \frac{i \tanh(x)}{b} - \frac{i \int \frac{a \tan(ix)^2 - ib \tan(ix) + a}{a - ib \tan(ix)} dx}{b} \right) \\
 & \quad \downarrow \text{4109} \\
 & i \left( \frac{i \tanh(x)}{b} - \frac{i \left( -\frac{iab \int i \tanh(x) dx}{a^2 - b^2} + \frac{a^3 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{a^2 - b^2} - \frac{b^2 x}{a^2 - b^2} \right)}{b} \right)
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 26 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( \frac{ab \int \tanh(x) dx}{a^2 - b^2} + \frac{a^3 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{a^2 - b^2} - \frac{b^2 x}{a^2 - b^2} \right)}{b} \right) \\
\downarrow 3042 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( \frac{ab \int -i \tan(ix) dx}{a^2 - b^2} + \frac{a^3 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx}{a^2 - b^2} - \frac{b^2 x}{a^2 - b^2} \right)}{b} \right) \\
\downarrow 26 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( -\frac{iab \int \tan(ix) dx}{a^2 - b^2} + \frac{a^3 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx}{a^2 - b^2} - \frac{b^2 x}{a^2 - b^2} \right)}{b} \right) \\
\downarrow 3956 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( \frac{a^3 \int \frac{\tan(ix)^2 + 1}{a - ib \tan(ix)} dx}{a^2 - b^2} - \frac{b^2 x}{a^2 - b^2} + \frac{ab \log(\cosh(x))}{a^2 - b^2} \right)}{b} \right) \\
\downarrow 4100 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( \frac{a^3 \int \frac{1}{a + b \tanh(x)} d(b \tanh(x))}{b(a^2 - b^2)} - \frac{b^2 x}{a^2 - b^2} + \frac{ab \log(\cosh(x))}{a^2 - b^2} \right)}{b} \right) \\
\downarrow 16 \\
i \left( \frac{i \tanh(x)}{b} - \frac{i \left( -\frac{b^2 x}{a^2 - b^2} + \frac{ab \log(\cosh(x))}{a^2 - b^2} + \frac{a^3 \log(a + b \tanh(x))}{b(a^2 - b^2)} \right)}{b} \right)
\end{array}$$

input `Int [Tanh[x]^3/(a + b*Tanh[x]), x]`

output  $I * (((-I) * (-((b^2 * x) / (a^2 - b^2)) + (a * b * \text{Log}[\text{Cosh}[x]]) / (a^2 - b^2) + (a^3 * \text{Log}[a + b * \text{Tanh}[x]]) / (b * (a^2 - b^2)))) / b + (I * \text{Tanh}[x]) / b)$

### Defintions of rubi rules used

rule 16  $\text{Int}[(c\_)/((a\_)+(b\_)*(x\_)), x\_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 25  $\text{Int}[-(Fx\_), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[Fx, x], x]$

rule 26  $\text{Int}[(\text{Complex}[0, a\_])*(Fx\_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956  $\text{Int}[\tan[(c\_)+(d\_)*(x\_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4049  $\text{Int}[(a\_)+(b\_)*\tan[(e\_)+(f\_)*(x\_)]^{(m\_)}*((c\_)+(d\_)*\tan[(e\_)+(f\_)*(x\_)]^{(n\_)}), x\_Symbol] \rightarrow \text{Simp}[b^2*(a + b*\text{Tan}[e + f*x])^{(m-2)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(d*f*(m+n-1))), x] + \text{Simp}[1/(d*(m+n-1)) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-3)}*(c + d*\text{Tan}[e + f*x])^n * \text{Simp}[a^3*d*(m+n-1) - b^2*(b*c*(m-2) + a*d*(1+n)) + b*d*(m+n-1)*(3*a^2 - b^2)*\text{Tan}[e + f*x] - b^2*(b*c*(m-2) - a*d*(3*m+2*n-4))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{IntegerQ}[2*m] \ \&\& \ \text{GtQ}[m, 2] \ \&\& \ (\text{GeQ}[n, -1] \ || \ \text{IntegerQ}[m]) \ \&\& \ !( \ \text{IGtQ}[n, 2] \ \&\& \ ( \ !\text{IntegerQ}[m] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{NeQ}[a, 0]) \ )$



rule 4100

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

rule 4109

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$-\frac{\tanh(x)}{b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{a^3 \ln(a+b \tanh(x))}{b^2(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b}$	67
default	$-\frac{\tanh(x)}{b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{a^3 \ln(a+b \tanh(x))}{b^2(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b}$	67
parallelrisch	$-\frac{\ln(1-\tanh(x))a b^2 - a^3 \ln(a+b \tanh(x)) + a b^2 x + b^3 x + \tanh(x) a^2 b - \tanh(x) b^3}{b^2(a^2 - b^2)}$	67
risch	$\frac{x}{a+b} + \frac{2ax}{b^2} - \frac{2a^3 x}{b^2(a^2 - b^2)} + \frac{2}{b(e^{2x}+1)} - \frac{a \ln(e^{2x}+1)}{b^2} + \frac{a^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b^2(a^2 - b^2)}$	97

input

```
int(tanh(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
-tanh(x)/b-1/(2*b+2*a)*ln(tanh(x)-1)+1/b^2*a^3/(a+b)/(a-b)*ln(a+b*tanh(x))
-1/(2*a-2*b)*ln(1+tanh(x))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 264 vs.  $2(64) = 128$ .

Time = 0.11 (sec) , antiderivative size = 264, normalized size of antiderivative = 4.12

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \frac{(ab^2 + b^3)x \cosh(x)^2 + 2(ab^2 + b^3)x \cosh(x) \sinh(x) + (ab^2 + b^3)x \sinh(x)^2 - 2a^2b + 2b^3 + (ab^2 + b^3)}{\dots}$$

input `integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

output 
$$-\left(\frac{(a^2b^2 + b^3)x \cosh(x)^2 + 2(a^2b^2 + b^3)x \cosh(x) \sinh(x) + (a^2b^2 + b^3)x \sinh(x)^2 - 2a^2b + 2b^3 + (a^2b^2 + b^3)x - (a^3 \cosh(x)^2 + 2a^3 \cosh(x) \sinh(x) + a^3 \sinh(x)^2 + a^3) \log(2(a \cosh(x) + b \sinh(x)) / (\cosh(x) - \sinh(x))) + (a^3 - a^2b^2 + (a^3 - a^2b^2) \cosh(x)^2 + 2(a^3 - a^2b^2) \cosh(x) \sinh(x) + (a^3 - a^2b^2) \sinh(x)^2) \log(2 \cosh(x) / (\cosh(x) - \sinh(x)))}{(a^2b^2 - b^4 + (a^2b^2 - b^4) \cosh(x)^2 + 2(a^2b^2 - b^4) \cosh(x) \sinh(x) + (a^2b^2 - b^4) \sinh(x)^2)}\right)$$

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 330 vs.  $2(49) = 98$ .

Time = 0.32 (sec) , antiderivative size = 330, normalized size of antiderivative = 5.16

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \begin{cases} \tilde{\infty}(x - \tanh(x)) & \text{for } a = \\ \frac{x - \log(\tanh(x)+1) - \frac{\tanh^2(x)}{2}}{a} & \text{for } b = \\ \frac{5x \tanh(x)}{2b \tanh(x) - 2b} - \frac{5x}{2b \tanh(x) - 2b} - \frac{2 \log(\tanh(x)+1) \tanh(x)}{2b \tanh(x) - 2b} + \frac{2 \log(\tanh(x)+1)}{2b \tanh(x) - 2b} - \frac{2 \tanh^2(x)}{2b \tanh(x) - 2b} + \frac{3}{2b \tanh(x) - 2b} & \text{for } a = \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} + \frac{2 \log(\tanh(x)+1) \tanh(x)}{2b \tanh(x) + 2b} + \frac{2 \log(\tanh(x)+1)}{2b \tanh(x) + 2b} - \frac{2 \tanh^2(x)}{2b \tanh(x) + 2b} + \frac{3}{2b \tanh(x) + 2b} & \text{for } a = \\ \frac{a^3 \log(\frac{a}{b} + \tanh(x))}{a^2b^2 - b^4} - \frac{a^2b \tanh(x)}{a^2b^2 - b^4} + \frac{ab^2x}{a^2b^2 - b^4} - \frac{ab^2 \log(\tanh(x)+1)}{a^2b^2 - b^4} - \frac{b^3x}{a^2b^2 - b^4} + \frac{b^3 \tanh(x)}{a^2b^2 - b^4} & \text{otherwise} \end{cases}$$

input `integrate(tanh(x)**3/(a+b*tanh(x)),x)`

output `Piecewise((zoo*(x - tanh(x)), Eq(a, 0) & Eq(b, 0)), ((x - log(tanh(x) + 1) - tanh(x)**2/2)/a, Eq(b, 0)), (5*x*tanh(x)/(2*b*tanh(x) - 2*b) - 5*x/(2*b*tanh(x) - 2*b) - 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) - 2*b) + 2*log(tanh(x) + 1)/(2*b*tanh(x) - 2*b) - 2*tanh(x)**2/(2*b*tanh(x) - 2*b) + 3/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) + 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) + 2*b) + 2*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) - 2*tanh(x)**2/(2*b*tanh(x) + 2*b) + 3/(2*b*tanh(x) + 2*b), Eq(a, b)), (a**3*log(a/b + tanh(x))/(a**2*b**2 - b**4) - a**2*b*tanh(x)/(a**2*b**2 - b**4) + a*b**2*x/(a**2*b**2 - b**4) - a*b**2*log(tanh(x) + 1)/(a**2*b**2 - b**4) - b**3*x/(a**2*b**2 - b**4) + b**3*tanh(x)/(a**2*b**2 - b**4), True))`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.11

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \frac{a^3 \log(-(a-b)e^{-2x} - a - b)}{a^2 b^2 - b^4} + \frac{x}{a+b} - \frac{a \log(e^{-2x} + 1)}{b^2} - \frac{2}{be^{-2x} + b}$$

input `integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output `a^3*log(-(a - b)*e^(-2*x) - a - b)/(a^2*b^2 - b^4) + x/(a + b) - a*log(e^(-2*x) + 1)/b^2 - 2/(b*e^(-2*x) + b)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.17

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \frac{a^3 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 b^2 - b^4} - \frac{x}{a - b} - \frac{a \log(e^{(2x)} + 1)}{b^2} + \frac{2}{b(e^{(2x)} + 1)}$$

input `integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="giac")`output `a^3*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2*b^2 - b^4) - x/(a - b) - a*log(e^(2*x) + 1)/b^2 + 2/(b*(e^(2*x) + 1))`**Mupad [B] (verification not implemented)**

Time = 2.20 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.92

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \frac{x}{a + b} - \frac{\tanh(x)}{b} - \frac{a \ln(\tanh(x) + 1)}{a^2 - b^2} + \frac{a^3 \ln(a + b \tanh(x))}{b^2 (a^2 - b^2)}$$

input `int(tanh(x)^3/(a + b*tanh(x)),x)`output `x/(a + b) - tanh(x)/b - (a*log(tanh(x) + 1))/(a^2 - b^2) + (a^3*log(a + b*tanh(x)))/(b^2*(a^2 - b^2))`**Reduce [B] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.44

$$\int \frac{\tanh^3(x)}{a + b \tanh(x)} dx = \frac{\log(e^{2x}a + e^{2x}b + a - b) a b^2 + \log(\tanh(x) b + a) a^3 - \log(\tanh(x) b + a) a b^2 - \tanh(x) a^2 b + \tanh(x) a^2 b}{b^2 (a^2 - b^2)}$$

input `int(tanh(x)^3/(a+b*tanh(x)),x)`

output `(log(e**(2*x)*a + e**(2*x)*b + a - b)*a*b**2 + log(tanh(x)*b + a)*a**3 - 1  
og(tanh(x)*b + a)*a*b**2 - tanh(x)*a**2*b + tanh(x)*b**3 - a*b**2*x - b**3  
*x)/(b**2*(a**2 - b**2))`

### 3.137 $\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx$

Optimal result	1061
Mathematica [A] (verified)	1061
Rubi [A] (verified)	1062
Maple [A] (verified)	1064
Fricas [A] (verification not implemented)	1065
Sympy [B] (verification not implemented)	1065
Maxima [A] (verification not implemented)	1066
Giac [A] (verification not implemented)	1066
Mupad [B] (verification not implemented)	1067
Reduce [B] (verification not implemented)	1067

#### Optimal result

Integrand size = 13, antiderivative size = 63

$$\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx = -\frac{ax}{b^2} + \frac{a^3x}{b^2(a^2-b^2)} + \frac{\log(\cosh(x))}{b} - \frac{a^2 \log(a \cosh(x) + b \sinh(x))}{b(a^2-b^2)}$$

output

```
-a*x/b^2+a^3*x/b^2/(a^2-b^2)+ln(cosh(x))/b-a^2*ln(a*cosh(x)+b*sinh(x))/b/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.94

$$\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx = -\frac{\log(1-\tanh(x))}{2(a+b)} + \frac{\log(1+\tanh(x))}{2(a-b)} - \frac{a^2 \log(a+b \tanh(x))}{b(a^2-b^2)}$$

input

```
Integrate[Tanh[x]^2/(a + b*Tanh[x]),x]
```

output

```
-1/2*Log[1 - Tanh[x]]/(a + b) + Log[1 + Tanh[x]]/(2*(a - b)) - (a^2*Log[a + b*Tanh[x]])/(b*(a^2 - b^2))
```

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.97, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.846$ , Rules used = {3042, 25, 4024, 26, 3042, 26, 3956, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan(ix)^2}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\tan(ix)^2}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{4024} \\
 & \frac{a^2 \int \frac{1}{a+b \tanh(x)} dx}{b^2} - \frac{i \int i \tanh(x) dx}{b} - \frac{ax}{b^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{a^2 \int \frac{1}{a+b \tanh(x)} dx}{b^2} + \frac{\int \tanh(x) dx}{b} - \frac{ax}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a^2 \int \frac{1}{a-ib \tan(ix)} dx}{b^2} + \frac{\int -i \tan(ix) dx}{b} - \frac{ax}{b^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{a^2 \int \frac{1}{a-ib \tan(ix)} dx}{b^2} - \frac{i \int \tan(ix) dx}{b} - \frac{ax}{b^2} \\
 & \quad \downarrow \text{3956} \\
 & \frac{a^2 \int \frac{1}{a-ib \tan(ix)} dx}{b^2} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b} \\
 & \quad \downarrow \text{3965}
 \end{aligned}$$

$$\begin{aligned}
& \frac{a^2 \left( \frac{ax}{a^2-b^2} - \frac{ib \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a^2-b^2} \right)}{b^2} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b} \\
& \quad \downarrow \text{26} \\
& \frac{a^2 \left( \frac{ax}{a^2-b^2} - \frac{b \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a^2-b^2} \right)}{b^2} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2 \left( \frac{ax}{a^2-b^2} - \frac{b \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a^2-b^2} \right)}{b^2} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b} \\
& \quad \downarrow \text{4013} \\
& \frac{a^2 \left( \frac{ax}{a^2-b^2} - \frac{b \log(a \cosh(x)+b \sinh(x))}{a^2-b^2} \right)}{b^2} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b}
\end{aligned}$$

input `Int [Tanh[x]^2/(a + b*Tanh[x]), x]`

output `-((a*x)/b^2) + Log[Cosh[x]]/b + (a^2*((a*x)/(a^2 - b^2) - (b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)))/b^2`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`



rule 3965  $\text{Int}[\left((a_{\_}) + (b_{\_})\tan[(c_{\_}) + (d_{\_})(x_{\_})]\right)^{-1}, x\_Symbol] \rightarrow \text{Simp}[a*(x/(a^2 + b^2)), x] + \text{Simp}[b/(a^2 + b^2) \text{Int}[(b - a*\text{Tan}[c + d*x])/(a + b*\text{Tan}[c + d*x]), x], x] /;$   $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

rule 4013  $\text{Int}[\left((c_{\_}) + (d_{\_})\tan[(e_{\_}) + (f_{\_})(x_{\_})]\right)/\left((a_{\_}) + (b_{\_})\tan[(e_{\_}) + (f_{\_})(x_{\_})]\right), x\_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

rule 4024  $\text{Int}[\left((c_{\_}) + (d_{\_})\tan[(e_{\_}) + (f_{\_})(x_{\_})]\right)^2/\left((a_{\_}) + (b_{\_})\tan[(e_{\_}) + (f_{\_})(x_{\_})]\right), x\_Symbol] \rightarrow \text{Simp}[d*(2*b*c - a*d)*(x/b^2), x] + (\text{Simp}[d^2/b \text{Int}[\text{Tan}[e + f*x], x], x] + \text{Simp}[(b*c - a*d)^2/b^2 \text{Int}[1/(a + b*\text{Tan}[e + f*x]), x], x]) /;$   $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

## Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

method	result	size
parallelrisch	$-\frac{-\ln(1-\tanh(x))b^2+a^2\ln(a+b\tanh(x))-abx-b^2x}{b(a^2-b^2)}$	52
derivativedivides	$-\frac{a^2\ln(a+b\tanh(x))}{(a+b)(a-b)b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	60
default	$-\frac{a^2\ln(a+b\tanh(x))}{(a+b)(a-b)b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	60
risch	$\frac{x}{a+b} + \frac{2xa^2}{b(a^2-b^2)} - \frac{2x}{b} - \frac{a^2\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b(a^2-b^2)} + \frac{\ln(e^{2x}+1)}{b}$	82

input `int(tanh(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output 
$$-(-\ln(1-\tanh(x))*b^2+a^2*\ln(a+b*\tanh(x))-a*b*x-b^2*x)/b/(a^2-b^2)$$

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.21

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx$$

$$= -\frac{a^2 \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (ab + b^2)x - (a^2 - b^2) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)}{a^2 b - b^3}$$

input `integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`output `-(a^2*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - (a*b + b^2)*x - (a^2 - b^2)*log(2*cosh(x)/(cosh(x) - sinh(x))))/(a^2*b - b^3)`**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. 2(51) = 102.

Time = 0.28 (sec) , antiderivative size = 243, normalized size of antiderivative = 3.86

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx$$

$$= \begin{cases} \tilde{\infty}(x - \log(\tanh(x) + 1)) & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \tanh(x)}{a} & \text{for } b = 0 \\ \frac{3x \tanh(x)}{2b \tanh(x) - 2b} - \frac{3x}{2b \tanh(x) - 2b} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) - 2b} + \frac{2 \log(\tanh(x) + 1)}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) + 2b} - \frac{2 \log(\tanh(x) + 1)}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ -\frac{a^2 \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 b - b^3} + \frac{abx}{a^2 b - b^3} - \frac{b^2 x}{a^2 b - b^3} + \frac{b^2 \log(\tanh(x) + 1)}{a^2 b - b^3} & \text{otherwise} \end{cases}$$

input `integrate(tanh(x)**2/(a+b*tanh(x)),x)`

output

```
Piecewise((zoo*(x - log(tanh(x) + 1)), Eq(a, 0) & Eq(b, 0)), ((x - tanh(x)
)/a, Eq(b, 0)), (3*x*tanh(x)/(2*b*tanh(x) - 2*b) - 3*x/(2*b*tanh(x) - 2*b)
- 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) - 2*b) + 2*log(tanh(x) + 1)/(2*
b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh
(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(
x) + 2*b) - 2*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b)
, Eq(a, b)), (-a**2*log(a/b + tanh(x))/(a**2*b - b**3) + a*b*x/(a**2*b - b
**3) - b**2*x/(a**2*b - b**3) + b**2*log(tanh(x) + 1)/(a**2*b - b**3), Tru
e))
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx = -\frac{a^2 \log(-(a-b)e^{-2x} - a - b)}{a^2b - b^3} + \frac{x}{a+b} + \frac{\log(e^{-2x} + 1)}{b}$$

input

```
integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")
```

output

```
-a^2*log(-(a - b)*e^(-2*x) - a - b)/(a^2*b - b^3) + x/(a + b) + log(e^(-2*
x) + 1)/b
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.92

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx = -\frac{a^2 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2b - b^3} + \frac{x}{a-b} + \frac{\log(e^{(2x)} + 1)}{b}$$

input

```
integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="giac")
```

output

```
-a^2*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2*b - b^3) + x/(a - b) + 1
og(e^(2*x) + 1)/b
```

**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx = -\frac{b^2(x - \ln(\tanh(x) + 1)) + a^2 \ln(a + b \tanh(x)) - a b x}{b(a^2 - b^2)}$$

input `int(tanh(x)^2/(a + b*tanh(x)),x)`output `-(b^2*(x - log(tanh(x) + 1)) + a^2*log(a + b*tanh(x)) - a*b*x)/(b*(a^2 - b^2))`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.16

$$\int \frac{\tanh^2(x)}{a + b \tanh(x)} dx = \frac{-\log(e^{2x}a + e^{2x}b + a - b)b^2 - \log(\tanh(x)b + a)a^2 + \log(\tanh(x)b + a)b^2 + abx + b^2x}{b(a^2 - b^2)}$$

input `int(tanh(x)^2/(a+b*tanh(x)),x)`output `( - log(e**(2*x)*a + e**(2*x)*b + a - b)*b**2 - log(tanh(x)*b + a)*a**2 + log(tanh(x)*b + a)*b**2 + a*b*x + b**2*x)/(b*(a**2 - b**2))`

### 3.138 $\int \frac{\tanh(x)}{a+b \tanh(x)} dx$

Optimal result	1068
Mathematica [A] (verified)	1068
Rubi [C] (verified)	1069
Maple [A] (verified)	1071
Fricas [A] (verification not implemented)	1071
Sympy [B] (verification not implemented)	1072
Maxima [A] (verification not implemented)	1072
Giac [A] (verification not implemented)	1073
Mupad [B] (verification not implemented)	1073
Reduce [B] (verification not implemented)	1073

#### Optimal result

Integrand size = 11, antiderivative size = 39

$$\int \frac{\tanh(x)}{a+b \tanh(x)} dx = -\frac{bx}{a^2-b^2} + \frac{a \log(a \cosh(x) + b \sinh(x))}{a^2-b^2}$$

output

```
-b*x/(a^2-b^2)+a*ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28

$$\begin{aligned} & \int \frac{\tanh(x)}{a+b \tanh(x)} dx \\ &= \frac{(-a+b) \log(1-\tanh(x)) - (a+b) \log(1+\tanh(x)) + 2a \log(a+b \tanh(x))}{2(a-b)(a+b)} \end{aligned}$$

input

```
Integrate[Tanh[x]/(a + b*Tanh[x]),x]
```

output

```
((-a + b)*Log[1 - Tanh[x]] - (a + b)*Log[1 + Tanh[x]] + 2*a*Log[a + b*Tanh[x]])/(2*(a - b)*(a + b))
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.23, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {3042, 26, 4014, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{4014} \\
 & -i \left( -\frac{a \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a^2 - b^2} - \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( \frac{ia \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a^2 - b^2} - \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{ia \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} - \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{4013} \\
 & -i \left( \frac{ia \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} - \frac{ibx}{a^2 - b^2} \right)
 \end{aligned}$$

input `Int[Tanh[x]/(a + b*Tanh[x]),x]`

output `(-I)*(((I)*b*x)/(a^2 - b^2) + (I*a*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)])*(x_)), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

**Maple [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

method	result	size
parallelrisc	$-\frac{a \ln(1-\tanh(x))-a \ln(a+b \tanh(x))+ax+bx}{a^2-b^2}$	40
risc	$\frac{x}{a+b} - \frac{2ax}{a^2-b^2} + \frac{a \ln\left(\frac{e^{2x} + \frac{a-b}{a+b}}{a^2-b^2}\right)}{a^2-b^2}$	54
derivativedivides	$\frac{a \ln(a+b \tanh(x))}{(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	55
default	$\frac{a \ln(a+b \tanh(x))}{(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	55

input `int(tanh(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

output `-(a*ln(1-tanh(x))-a*ln(a+b*tanh(x))+a*x+b*x)/(a^2-b^2)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{\tanh(x)}{a+b \tanh(x)} dx = -\frac{(a+b)x - a \log\left(\frac{2(a \cosh(x)+b \sinh(x))}{\cosh(x)-\sinh(x)}\right)}{a^2-b^2}$$

input `integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="fricas")`

output `-((a + b)*x - a*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))))/(a^2 - b^2)`



**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 141 vs.  $2(29) = 58$ .

Time = 0.22 (sec) , antiderivative size = 141, normalized size of antiderivative = 3.62

$$\int \frac{\tanh(x)}{a + b \tanh(x)} dx$$

$$= \begin{cases} \tilde{\infty}x & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \log(\tanh(x) + 1)}{a} & \text{for } b = 0 \\ \frac{x \tanh(x)}{2b \tanh(x) - 2b} - \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} + \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} + \frac{a \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} - \frac{a \log(\tanh(x) + 1)}{a^2 - b^2} - \frac{bx}{a^2 - b^2} & \text{otherwise} \end{cases}$$

input `integrate(tanh(x)/(a+b*tanh(x)),x)`

output `Piecewise((zoo*x, Eq(a, 0) & Eq(b, 0)), ((x - log(tanh(x) + 1))/a, Eq(b, 0)), (x*tanh(x)/(2*b*tanh(x) - 2*b) - x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) + 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) + a*log(a/b + tanh(x))/(a**2 - b**2) - a*log(tanh(x) + 1)/(a**2 - b**2) - b*x/(a**2 - b**2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

$$\int \frac{\tanh(x)}{a + b \tanh(x)} dx = \frac{a \log(-(a - b)e^{(-2x)} - a - b)}{a^2 - b^2} + \frac{x}{a + b}$$

input `integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="maxima")`

output `a*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{\tanh(x)}{a + b \tanh(x)} dx = \frac{a \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} - \frac{x}{a - b}$$

input `integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="giac")`output `a*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2 - b^2) - x/(a - b)`**Mupad [B] (verification not implemented)**

Time = 2.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int \frac{\tanh(x)}{a + b \tanh(x)} dx = -\frac{bx - a(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

input `int(tanh(x)/(a + b*tanh(x)),x)`output `-(b*x - a*(x - log(tanh(x) + 1) + log(a + b*tanh(x))))/(a^2 - b^2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{\tanh(x)}{a + b \tanh(x)} dx = \frac{\log(e^{2x}a + e^{2x}b + a - b)a - ax - bx}{a^2 - b^2}$$

input `int(tanh(x)/(a+b*tanh(x)),x)`output `(log(e**(2*x)*a + e**(2*x)*b + a - b)*a - a*x - b*x)/(a**2 - b**2)`

### 3.139 $\int \frac{1}{a+b \tanh(x)} dx$

Optimal result	1074
Mathematica [A] (verified)	1074
Rubi [A] (verified)	1075
Maple [A] (verified)	1076
Fricas [A] (verification not implemented)	1077
Sympy [B] (verification not implemented)	1077
Maxima [A] (verification not implemented)	1078
Giac [A] (verification not implemented)	1078
Mupad [B] (verification not implemented)	1078
Reduce [B] (verification not implemented)	1079

#### Optimal result

Integrand size = 8, antiderivative size = 39

$$\int \frac{1}{a+b \tanh(x)} dx = \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

output

```
a*x/(a^2-b^2)-b*ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.26

$$\int \frac{1}{a+b \tanh(x)} dx = \frac{(-a+b) \log(1 - \tanh(x)) + (a+b) \log(1 + \tanh(x)) - 2b \log(a + b \tanh(x))}{2(a-b)(a+b)}$$

input

```
Integrate[(a + b*Tanh[x])^(-1), x]
```

output

```
((-a + b)*Log[1 - Tanh[x]] + (a + b)*Log[1 + Tanh[x]] - 2*b*Log[a + b*Tanh[x]])/(2*(a - b)*(a + b))
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a - ib \tan(ix)} dx \\
 & \quad \downarrow \text{3965} \\
 & \frac{ax}{a^2 - b^2} - \frac{ib \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{4013} \\
 & \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}
 \end{aligned}$$

input `Int[(a + b*Tanh[x])^(-1),x]`

output `(a*x)/(a^2 - b^2) - (b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3965  $\text{Int}[(a + (b \cdot \tan[c + d \cdot x])^{-1}), x\_Symbol] \rightarrow \text{Simp}[a \cdot (x / (a^2 + b^2)), x] + \text{Simp}[b / (a^2 + b^2) \text{Int}[(b - a \cdot \tan[c + d \cdot x]) / (a + b \cdot \tan[c + d \cdot x]), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4013  $\text{Int}[(c + (d \cdot \tan[e + f \cdot x]) / (a + (b \cdot \tan[e + f \cdot x]) \cdot (x))), x\_Symbol] \rightarrow \text{Simp}[(c / (b \cdot f)) \cdot \text{Log}[\text{RemoveContent}[a \cdot \text{Cos}[e + f \cdot x] + b \cdot \text{Sin}[e + f \cdot x], x]], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a \cdot c + b \cdot d, 0]$

## Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

method	result	size
parallelrisch	$-\frac{-\ln(1-\tanh(x))b+b\ln(a+b\tanh(x))-ax-bx}{a^2-b^2}$	42
derivativedivides	$-\frac{b\ln(a+b\tanh(x))}{(a-b)(a+b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
default	$-\frac{b\ln(a+b\tanh(x))}{(a-b)(a+b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
risch	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} - \frac{b\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2-b^2}$	55

input `int(1/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`output 
$$-(-\ln(1-\tanh(x))*b+b*\ln(a+b*\tanh(x))-a*x-b*x)/(a^2-b^2)$$

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{(a + b)x - b \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^2 - b^2}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="fricas")`

output `((a + b)*x - b*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))))/(a^2 - b^2)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. 2(29) = 58.

Time = 0.22 (sec) , antiderivative size = 146, normalized size of antiderivative = 3.74

$$\int \frac{1}{a + b \tanh(x)} dx = \begin{cases} \tilde{\infty}(x - \log(\tanh(x) + 1) + \log(\tanh(x))) & \text{for } a = 0 \wedge b = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ -\frac{x \tanh(x)}{2b \tanh(x) - 2b} + \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} - \frac{bx}{a^2 - b^2} - \frac{b \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} + \frac{b \log(\tanh(x) + 1)}{a^2 - b^2} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+b*tanh(x)),x)`

output `Piecewise((zoo*(x - log(tanh(x) + 1) + log(tanh(x))), Eq(a, 0) & Eq(b, 0)), (x/a, Eq(b, 0)), (-x*tanh(x)/(2*b*tanh(x) - 2*b) + x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) - b*x/(a**2 - b**2) - b*log(a/b + tanh(x))/(a**2 - b**2) + b*log(tanh(x) + 1)/(a**2 - b**2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{1}{a + b \tanh(x)} dx = -\frac{b \log(-(a - b)e^{(-2x)} - a - b)}{a^2 - b^2} + \frac{x}{a + b}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="maxima")`output `-b*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{1}{a + b \tanh(x)} dx = -\frac{b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} + \frac{x}{a - b}$$

input `integrate(1/(a+b*tanh(x)),x, algorithm="giac")`output `-b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2 - b^2) + x/(a - b)`**Mupad [B] (verification not implemented)**

Time = 0.00 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{ax - b(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

input `int(1/(a + b*tanh(x)),x)`output `(a*x - b*(x - log(tanh(x) + 1) + log(a + b*tanh(x))))/(a^2 - b^2)`

**Reduce [B] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

$$\int \frac{1}{a + b \tanh(x)} dx = \frac{-\log(e^{2x}a + e^{2x}b + a - b)b + ax + bx}{a^2 - b^2}$$

input `int(1/(a+b*tanh(x)),x)`

output `( - log(e**(2*x)*a + e**(2*x)*b + a - b)*b + a*x + b*x)/(a**2 - b**2)`



### 3.140 $\int \frac{\coth(x)}{a+b \tanh(x)} dx$

Optimal result	1080
Mathematica [A] (verified)	1080
Rubi [C] (verified)	1081
Maple [A] (verified)	1083
Fricas [A] (verification not implemented)	1083
Sympy [F]	1084
Maxima [A] (verification not implemented)	1084
Giac [A] (verification not implemented)	1085
Mupad [B] (verification not implemented)	1085
Reduce [B] (verification not implemented)	1085

#### Optimal result

Integrand size = 11, antiderivative size = 51

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = -\frac{bx}{a^2 - b^2} + \frac{\log(\sinh(x))}{a} + \frac{b^2 \log(a \cosh(x) + b \sinh(x))}{a(a^2 - b^2)}$$

output

```
-b*x/(a^2-b^2)+ln(sinh(x))/a+b^2*ln(a*cosh(x)+b*sinh(x))/a/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.27

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = -\frac{\log(1 - \tanh(x))}{2(a + b)} + \frac{\log(\tanh(x))}{a} - \frac{\log(1 + \tanh(x))}{2(a - b)} + \frac{b^2 \log(a + b \tanh(x))}{a(a^2 - b^2)}$$

input

```
Integrate[Coth[x]/(a + b*Tanh[x]),x]
```

output

```
-1/2*Log[1 - Tanh[x]]/(a + b) + Log[Tanh[x]]/a - Log[1 + Tanh[x]]/(2*(a - b)) + (b^2*Log[a + b*Tanh[x]])/(a*(a^2 - b^2))
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.24, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$ , Rules used = {3042, 26, 4054, 26, 3042, 26, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i}{\tan(ix)(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{1}{\tan(ix)(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{4054} \\
 & i \left( \frac{b^2 \int \frac{-i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a(a^2 - b^2)} + \frac{\int -i \coth(x) dx}{a} + \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( -\frac{ib^2 \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a(a^2 - b^2)} - \frac{i \int \coth(x) dx}{a} + \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & i \left( -\frac{ib^2 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2 - b^2)} - \frac{i \int -i \tan \left( ix + \frac{\pi}{2} \right) dx}{a} + \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{26} \\
 & i \left( -\frac{ib^2 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2 - b^2)} - \frac{\int \tan \left( ix + \frac{\pi}{2} \right) dx}{a} + \frac{ibx}{a^2 - b^2} \right) \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

$$i \left( -\frac{ib^2 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} + \frac{ibx}{a^2-b^2} - \frac{i \log(\sinh(x))}{a} \right)$$

↓ 4013

$$i \left( \frac{ibx}{a^2-b^2} - \frac{ib^2 \log(a \cosh(x) + b \sinh(x))}{a(a^2-b^2)} - \frac{i \log(\sinh(x))}{a} \right)$$

input `Int[Coth[x]/(a + b*Tanh[x]),x]`

output `I*((I*b*x)/(a^2 - b^2) - (I*Log[Sinh[x]])/a - (I*b^2*Log[a*Cosh[x] + b*Sinh[x]])/(a*(a^2 - b^2)))`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sinh[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4054

```
Int[1/(((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(a*c - b*d)*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[b^2/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[d^2/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.10

method	result	size
parallelrisch	$\frac{b^2 \ln(a+b \tanh(x)) - a^2 \ln(1 - \tanh(x)) - (a+b)((-a+b) \ln(\tanh(x)) + ax)}{a^3 - b^2 a}$	56
derivativedivides	$-\frac{\ln(1+\tanh(x))}{2a-2b} + \frac{\ln(\tanh(x))}{a} + \frac{b^2 \ln(a+b \tanh(x))}{a(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	67
default	$-\frac{\ln(1+\tanh(x))}{2a-2b} + \frac{\ln(\tanh(x))}{a} + \frac{b^2 \ln(a+b \tanh(x))}{a(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	67
risch	$\frac{x}{a+b} - \frac{2x}{a} - \frac{2xb^2}{a(a^2-b^2)} + \frac{\ln(e^{2x}-1)}{a} + \frac{b^2 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a(a^2-b^2)}$	81

input

```
int(coth(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
(b^2*ln(a+b*tanh(x))-a^2*ln(1-tanh(x))-(a+b)*((-a+b)*ln(tanh(x))+a*x))/(a^3-a*b^2)
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.43

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx$$

$$= \frac{b^2 \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (a^2 + ab)x + (a^2 - b^2) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right)}{a^3 - ab^2}$$

input

```
integrate(coth(x)/(a+b*tanh(x)),x, algorithm="fricas")
```

output  $(b^2 \log(2(a \cosh(x) + b \sinh(x)) / (\cosh(x) - \sinh(x))) - (a^2 + a b)x + (a^2 - b^2) \log(2 \sinh(x) / (\cosh(x) - \sinh(x)))) / (a^3 - a b^2)$

### Sympy [F]

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = \int \frac{\coth(x)}{a + b \tanh(x)} dx$$

input `integrate(coth(x)/(a+b*tanh(x)),x)`

output `Integral(coth(x)/(a + b*tanh(x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.27

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = \frac{b^2 \log(-(a - b)e^{-2x} - a - b)}{a^3 - ab^2} + \frac{x}{a + b} + \frac{\log(e^{-x} + 1)}{a} + \frac{\log(e^{-x} - 1)}{a}$$

input `integrate(coth(x)/(a+b*tanh(x)),x, algorithm="maxima")`

output  $b^2 \log(-(a - b)e^{-2x} - a - b) / (a^3 - a b^2) + x / (a + b) + \log(e^{-x} + 1) / a + \log(e^{-x} - 1) / a$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = \frac{b^2 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^3 - ab^2} - \frac{x}{a - b} + \frac{\log(|e^{(2x)} - 1|)}{a}$$

input `integrate(coth(x)/(a+b*tanh(x)),x, algorithm="giac")`output `b^2*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^3 - a*b^2) - x/(a - b) + log(abs(e^(2*x) - 1))/a`**Mupad [B] (verification not implemented)**

Time = 2.40 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = \frac{\ln(e^{2x} - 1)}{a} - \frac{x}{a - b} - \frac{b^2 \ln(a - b + ae^{2x} + be^{2x})}{ab^2 - a^3}$$

input `int(coth(x)/(a + b*tanh(x)),x)`output `log(exp(2*x) - 1)/a - x/(a - b) - (b^2*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a*b^2 - a^3)`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.82

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = \frac{\log(e^x - 1) a^2 - \log(e^x - 1) b^2 + \log(e^x + 1) a^2 - \log(e^x + 1) b^2 + \log(e^{2x} a + e^{2x} b + a - b) b^2 - a^2 x - a}{a(a^2 - b^2)}$$

input `int(coth(x)/(a+b*tanh(x)),x)`

output

```
(log(e**x - 1)*a**2 - log(e**x - 1)*b**2 + log(e**x + 1)*a**2 - log(e**x + 1)*b**2 + log(e**(2*x)*a + e**(2*x)*b + a - b)*b**2 - a**2*x - a*b*x)/(a*(a**2 - b**2))
```

### 3.141 $\int \frac{\coth^2(x)}{a+b \tanh(x)} dx$

Optimal result	1087
Mathematica [A] (verified)	1087
Rubi [C] (verified)	1088
Maple [A] (verified)	1091
Fricas [B] (verification not implemented)	1091
Sympy [F]	1092
Maxima [A] (verification not implemented)	1092
Giac [A] (verification not implemented)	1093
Mupad [B] (verification not implemented)	1093
Reduce [B] (verification not implemented)	1093

#### Optimal result

Integrand size = 13, antiderivative size = 60

$$\int \frac{\coth^2(x)}{a+b \tanh(x)} dx = \frac{ax}{a^2-b^2} - \frac{\coth(x)}{a} - \frac{b \log(\sinh(x))}{a^2} - \frac{b^3 \log(a \cosh(x) + b \sinh(x))}{a^2(a^2-b^2)}$$

output

$a*x/(a^2-b^2)-\coth(x)/a-b*\ln(\sinh(x))/a^2-b^3*\ln(a*\cosh(x)+b*\sinh(x))/a^2/(a^2-b^2)$

#### Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

$$\int \frac{\coth^2(x)}{a+b \tanh(x)} dx = -\frac{\coth(x)}{a} - \frac{\log(1-\coth(x))}{2(a+b)} + \frac{\log(1+\coth(x))}{2(a-b)} - \frac{b^3 \log(b+a \coth(x))}{a^2(a^2-b^2)}$$

input

`Integrate[Coth[x]^2/(a + b*Tanh[x]), x]`

output

$-(\text{Coth}[x]/a) - \text{Log}[1 - \text{Coth}[x]]/(2*(a + b)) + \text{Log}[1 + \text{Coth}[x]]/(2*(a - b)) - (b^3*\text{Log}[b + a*\text{Coth}[x]])/(a^2*(a^2 - b^2))$



**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.28, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.923$ , Rules used = {3042, 25, 4052, 25, 3042, 26, 4134, 26, 3042, 26, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\coth^2(x)}{a + b \tanh(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{\tan(ix)^2(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1}{\tan(ix)^2(a - ib \tan(ix))} dx \\
 & \quad \downarrow \text{4052} \\
 & \frac{\int -\frac{\coth(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{a} - \frac{\coth(x)}{a} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\int \frac{\coth(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{a} - \frac{\coth(x)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\coth(x)}{a} - \frac{\int \frac{i(b \tan(ix)^2 + ia \tan(ix) + b)}{\tan(ix)(a - ib \tan(ix))} dx}{a} \\
 & \quad \downarrow \text{26} \\
 & \frac{\coth(x)}{a} - \frac{i \int \frac{b \tan(ix)^2 + ia \tan(ix) + b}{\tan(ix)(a - ib \tan(ix))} dx}{a} \\
 & \quad \downarrow \text{4134}
 \end{aligned}$$

$$\frac{\coth(x)}{a} - \frac{i \left( \frac{b^3 \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx}{a(a^2-b^2)} + \frac{b \int -i \coth(x) dx}{a} + \frac{ia^2 x}{a^2-b^2} \right)}{a}$$

↓ 26

$$\frac{\coth(x)}{a} - \frac{i \left( -\frac{ib^3 \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx}{a(a^2-b^2)} - \frac{ib \int \coth(x) dx}{a} + \frac{ia^2 x}{a^2-b^2} \right)}{a}$$

↓ 3042

$$\frac{\coth(x)}{a} - \frac{i \left( -\frac{ib^3 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} - \frac{ib \int -i \tan(ix+\frac{\pi}{2}) dx}{a} + \frac{ia^2 x}{a^2-b^2} \right)}{a}$$

↓ 26

$$\frac{\coth(x)}{a} - \frac{i \left( -\frac{ib^3 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} - \frac{b \int \tan(ix+\frac{\pi}{2}) dx}{a} + \frac{ia^2 x}{a^2-b^2} \right)}{a}$$

↓ 3956

$$\frac{\coth(x)}{a} - \frac{i \left( -\frac{ib^3 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} + \frac{ia^2 x}{a^2-b^2} - \frac{ib \log(\sinh(x))}{a} \right)}{a}$$

↓ 4013

$$\frac{\coth(x)}{a} - \frac{i \left( \frac{ia^2 x}{a^2-b^2} - \frac{ib^3 \log(a \cosh(x)+b \sinh(x))}{a(a^2-b^2)} - \frac{ib \log(\sinh(x))}{a} \right)}{a}$$

input `Int [Coth[x]^2/(a + b*Tanh[x]), x]`

output `-(Coth[x]/a) - (I*((I*a^2*x)/(a^2 - b^2) - (I*b*Log[Sinh[x]])/a - (I*b^3*Log[a*Cosh[x] + b*Sinh[x]])/(a*(a^2 - b^2))))/a`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sinn[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`
- rule 4052 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] :> Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

### Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.18

method	result	size
parallelrisch	$\frac{-b^3 \ln(a+b \tanh(x)) + \ln(1 - \tanh(x)) a^2 b + (a+b)(-b(a-b) \ln(\tanh(x)) + a((-a+b) \coth(x) + ax))}{a^4 - a^2 b^2}$	71
derivativedivides	$\frac{\ln(1 + \tanh(x))}{2a - 2b} - \frac{\ln(\tanh(x) - 1)}{2b + 2a} - \frac{b \ln(\tanh(x))}{a^2} - \frac{1}{a \tanh(x)} - \frac{b^3 \ln(a + b \tanh(x))}{(a - b)(a + b)a^2}$	78
default	$\frac{\ln(1 + \tanh(x))}{2a - 2b} - \frac{\ln(\tanh(x) - 1)}{2b + 2a} - \frac{b \ln(\tanh(x))}{a^2} - \frac{1}{a \tanh(x)} - \frac{b^3 \ln(a + b \tanh(x))}{(a - b)(a + b)a^2}$	78
risch	$\frac{x}{a+b} + \frac{2xb}{a^2} + \frac{2xb^3}{a^2(a^2-b^2)} - \frac{2}{a(e^{2x}-1)} - \frac{b \ln(e^{2x}-1)}{a^2} - \frac{b^3 \ln(e^{2x} + \frac{a-b}{a+b})}{a^2(a^2-b^2)}$	98

input

```
int(coth(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
(-b^3*ln(a+b*tanh(x))+ln(1-tanh(x))*a^2*b+(a+b)*(-b*(a-b)*ln(tanh(x))+a*((-a+b)*coth(x)+a*x)))/(a^4-a^2*b^2)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(60) = 120.

Time = 0.10 (sec) , antiderivative size = 271, normalized size of antiderivative = 4.52

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx =$$

$$\frac{(a^3 + a^2b)x \cosh(x)^2 + 2(a^3 + a^2b)x \cosh(x) \sinh(x) + (a^3 + a^2b)x \sinh(x)^2 - 2a^3 + 2ab^2 - (a^3 + a^2b)}{\dots}$$

input `integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`

output 
$$\begin{aligned} & -((a^3 + a^2*b)*x*\cosh(x)^2 + 2*(a^3 + a^2*b)*x*\cosh(x)*\sinh(x) + (a^3 + a^2*b)*x*\sinh(x)^2 - 2*a^3 + 2*a*b^2 - (a^3 + a^2*b)*x - (b^3*\cosh(x)^2 + 2*b^3*\cosh(x)*\sinh(x) + b^3*\sinh(x)^2 - b^3)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + (a^2*b - b^3 - (a^2*b - b^3)*\cosh(x)^2 - 2*(a^2*b - b^3)*\cosh(x)*\sinh(x) - (a^2*b - b^3)*\sinh(x)^2)*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))))/(a^4 - a^2*b^2 - (a^4 - a^2*b^2)*\cosh(x)^2 - 2*(a^4 - a^2*b^2)*\cosh(x)*\sinh(x) - (a^4 - a^2*b^2)*\sinh(x)^2) \end{aligned}$$

### Sympy [F]

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx = \int \frac{\coth^2(x)}{a + b \tanh(x)} dx$$

input `integrate(coth(x)**2/(a+b*tanh(x)),x)`

output `Integral(coth(x)**2/(a + b*tanh(x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.43

$$\begin{aligned} \int \frac{\coth^2(x)}{a + b \tanh(x)} dx = & -\frac{b^3 \log(-(a-b)e^{(-2x)} - a - b)}{a^4 - a^2 b^2} + \frac{x}{a + b} \\ & - \frac{b \log(e^{(-x)} + 1)}{a^2} - \frac{b \log(e^{(-x)} - 1)}{a^2} + \frac{2}{ae^{(-2x)} - a} \end{aligned}$$

input `integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="maxima")`

output 
$$-b^3*\log(-(a - b)*e^{(-2*x)} - a - b)/(a^4 - a^2*b^2) + x/(a + b) - b*\log(e^{(-x)} + 1)/a^2 - b*\log(e^{(-x)} - 1)/a^2 + 2/(a*e^{(-2*x)} - a)$$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.25

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx = -\frac{b^3 \log(|ae^{2x} + be^{2x} + a - b|)}{a^4 - a^2 b^2} + \frac{x}{a - b} - \frac{b \log(|e^{2x} - 1|)}{a^2} - \frac{2}{a(e^{2x} - 1)}$$

input `integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="giac")`output `-b^3*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^4 - a^2*b^2) + x/(a - b) - b*log(abs(e^(2*x) - 1))/a^2 - 2/(a*(e^(2*x) - 1))`**Mupad [B] (verification not implemented)**

Time = 2.39 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.22

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx = \frac{x}{a - b} - \frac{2}{a(e^{2x} - 1)} - \frac{b^3 \ln(a - b + a e^{2x} + b e^{2x})}{a^4 - a^2 b^2} - \frac{b \ln(e^{2x} - 1)}{a^2}$$

input `int(coth(x)^2/(a + b*tanh(x)),x)`output `x/(a - b) - 2/(a*(exp(2*x) - 1)) - (b^3*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^4 - a^2*b^2) - (b*log(exp(2*x) - 1))/a^2`**Reduce [B] (verification not implemented)**

Time = 3.05 (sec) , antiderivative size = 252, normalized size of antiderivative = 4.20

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx = \frac{-e^{2x} \log(e^x - 1) a^2 b + e^{2x} \log(e^x - 1) b^3 - e^{2x} \log(e^x + 1) a^2 b + e^{2x} \log(e^x + 1) b^3 - e^{2x} \log(e^{2x} a + e^{2x} b + \dots)}{\dots}$$

input `int(coth(x)^2/(a+b*tanh(x)),x)`

output `( - e**(2*x)*log(e**x - 1)*a**2*b + e**(2*x)*log(e**x - 1)*b**3 - e**(2*x)*log(e**x + 1)*a**2*b + e**(2*x)*log(e**x + 1)*b**3 - e**(2*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*b**3 + e**(2*x)*a**3*x - 2*e**(2*x)*a**3 + e**(2*x)*a**2*b*x + 2*e**(2*x)*a*b**2 + log(e**x - 1)*a**2*b - log(e**x - 1)*b**3 + log(e**x + 1)*a**2*b - log(e**x + 1)*b**3 + log(e**(2*x)*a + e**(2*x)*b + a - b)*b**3 - a**3*x - a**2*b*x)/(a**2*(e**(2*x)*a**2 - e**(2*x)*b**2 - a**2 + b**2))`

### 3.142 $\int \frac{\coth^3(x)}{a+b \tanh(x)} dx$

Optimal result	1095
Mathematica [A] (verified)	1095
Rubi [C] (verified)	1096
Maple [A] (verified)	1101
Fricas [B] (verification not implemented)	1101
Sympy [F]	1102
Maxima [A] (verification not implemented)	1102
Giac [A] (verification not implemented)	1103
Mupad [B] (verification not implemented)	1103
Reduce [F]	1104

#### Optimal result

Integrand size = 13, antiderivative size = 76

$$\int \frac{\coth^3(x)}{a+b \tanh(x)} dx = -\frac{bx}{a^2-b^2} + \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} + \frac{(a^2+b^2) \log(\sinh(x))}{a^3} + \frac{b^4 \log(a \cosh(x) + b \sinh(x))}{a^3(a^2-b^2)}$$

output

```
-b*x/(a^2-b^2)+b*coth(x)/a^2-1/2*coth(x)^2/a+(a^2+b^2)*ln(sinh(x))/a^3+b^4
*ln(a*cosh(x)+b*sinh(x))/a^3/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{a+b \tanh(x)} dx = \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} - \frac{\log(1-\coth(x))}{2(a+b)} - \frac{\log(1+\coth(x))}{2(a-b)} + \frac{b^4 \log(b+a \coth(x))}{a^3(a^2-b^2)}$$

input

```
Integrate[Coth[x]^3/(a + b*Tanh[x]), x]
```



output

$$(b \operatorname{Coth}[x])/a^2 - \operatorname{Coth}[x]^2/(2a) - \operatorname{Log}[1 - \operatorname{Coth}[x]]/(2(a + b)) - \operatorname{Log}[1 + \operatorname{Coth}[x]]/(2(a - b)) + (b^4 \operatorname{Log}[b + a \operatorname{Coth}[x]])/(a^3(a^2 - b^2))$$

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.43, number of steps used = 16, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.231$ , Rules used = {3042, 26, 4052, 27, 3042, 25, 4132, 25, 3042, 26, 4135, 26, 3042, 26, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{coth}^3(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow 3042 \\ & \int -\frac{i}{\tan(ix)^3(a - ib \tan(ix))} dx \\ & \quad \downarrow 26 \\ & -i \int \frac{1}{\tan(ix)^3(a - ib \tan(ix))} dx \\ & \quad \downarrow 4052 \\ & -i \left( -\frac{\int \frac{2i \operatorname{coth}^2(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{2a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\ & \quad \downarrow 27 \\ & -i \left( -\frac{i \int \frac{\operatorname{coth}^2(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\ & \quad \downarrow 3042 \\ & -i \left( -\frac{i \int -\frac{b \tan(ix)^2 + ia \tan(ix) + b}{\tan(ix)^2(a - ib \tan(ix))} dx}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\ & \quad \downarrow 25 \end{aligned}$$

$$\begin{aligned}
 & -i \left( \frac{i \int \frac{b \tan(ix)^2 + ia \tan(ix) + b}{\tan(ix)^2(a - ib \tan(ix))} dx}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\
 & \quad \downarrow \text{4132} \\
 & -i \left( \frac{i \left( \frac{b \operatorname{coth}(x)}{a} - \frac{\int -\frac{\operatorname{coth}(x)(a^2 + b^2 - b^2 \tanh^2(x))}{a + b \tanh(x)} dx}{a} \right)}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\
 & \quad \downarrow \text{25} \\
 & -i \left( \frac{i \left( \frac{\int \frac{\operatorname{coth}(x)(a^2 + b^2 - b^2 \tanh^2(x))}{a + b \tanh(x)} dx}{a} + \frac{b \operatorname{coth}(x)}{a} \right)}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\
 & \quad \downarrow \text{3042} \\
 & -i \left( \frac{i \left( \frac{b \operatorname{coth}(x)}{a} + \frac{\int \frac{i(a^2 + b^2 + b^2 \tan(ix)^2)}{\tan(ix)(a - ib \tan(ix))} dx}{a} \right)}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\
 & \quad \downarrow \text{26} \\
 & -i \left( \frac{i \left( \frac{b \operatorname{coth}(x)}{a} + \frac{i \int \frac{a^2 + b^2 + b^2 \tan(ix)^2}{\tan(ix)(a - ib \tan(ix))} dx}{a} \right)}{a} - \frac{i \operatorname{coth}^2(x)}{2a} \right) \\
 & \quad \downarrow \text{4135}
 \end{aligned}$$

$$-i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( \frac{(a^2+b^2) \int -i \coth(x) dx}{a} + \frac{b^4 \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx + \frac{ia^2 bx}{a^2-b^2} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right)$$

↓ 26

$$-i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( -\frac{i(a^2+b^2) \int \coth(x) dx}{a} - \frac{ib^4 \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx + \frac{ia^2 bx}{a^2-b^2} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right)$$

↓ 3042

$$-i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( -\frac{i(a^2+b^2) \int -i \tan(ix + \frac{\pi}{2}) dx}{a} - \frac{ib^4 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx + \frac{ia^2 bx}{a^2-b^2} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right)$$

↓ 26

$$-i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( -\frac{(a^2+b^2) \int \tan(ix + \frac{\pi}{2}) dx}{a} - \frac{ib^4 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx + \frac{ia^2 bx}{a^2-b^2} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right)$$

↓ 3956

$$\begin{aligned}
 & -i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( -\frac{ib^4 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx + \frac{ia^2 bx}{a^2-b^2} - \frac{i(a^2+b^2) \log(\sinh(x))}{a} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right) \\
 & \quad \downarrow \text{4013} \\
 & -i \left( \frac{i \left( \frac{b \coth(x)}{a} + \frac{i \left( \frac{ia^2 bx}{a^2-b^2} - \frac{i(a^2+b^2) \log(\sinh(x))}{a} - \frac{ib^4 \log(a \cosh(x) + b \sinh(x))}{a(a^2-b^2)} \right)}{a} \right)}{a} - \frac{i \coth^2(x)}{2a} \right)
 \end{aligned}$$

input `Int[Coth[x]^3/(a + b*Tanh[x]),x]`

output `(-I)*((( (-1/2*I)*Coth[x]^2)/a + (I*((b*Coth[x])/a + (I*((I*a^2*b*x)/(a^2 - b^2) - (I*(a^2 + b^2)*Log[Sinh[x]))/a - (I*b^4*Log[a*Cosh[x] + b*Sinh[x]])/(a*(a^2 - b^2)))))/a)/a)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4052 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4135

```
Int[((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] := Simp[(a*(A*c - c*C) - b*(A*d - C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

**Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

method	result
parallelrisch	$\frac{2 \ln(a+b \tanh(x))b^4 - 2 \ln(1-\tanh(x))a^4 + (2a^4 - 2b^4) \ln(\tanh(x)) - 2(a+b) \left( \frac{\coth(x)^2 a(a-b)}{2} - b \coth(x)(a-b) + a^2 x \right) a}{2a^5 - 2a^3 b^2}$
derivativedivides	$\frac{b^4 \ln(a+b \tanh(x))}{a^3(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{b}{a^2 \tanh(x)} - \frac{(-a^2-b^2) \ln(\tanh(x))}{a^3} - \frac{1}{2a \tanh(x)^2} - \frac{\ln(1+\tanh(x))}{2a-2b}$
default	$\frac{b^4 \ln(a+b \tanh(x))}{a^3(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{b}{a^2 \tanh(x)} - \frac{(-a^2-b^2) \ln(\tanh(x))}{a^3} - \frac{1}{2a \tanh(x)^2} - \frac{\ln(1+\tanh(x))}{2a-2b}$
risch	$\frac{x}{a+b} - \frac{2x}{a} - \frac{2xb^2}{a^3} - \frac{2xb^4}{a^3(a^2-b^2)} - \frac{2(e^{2x}a - e^{2x}b + b)}{(e^{2x}-1)^2 a^2} + \frac{\ln(e^{2x}-1)}{a} + \frac{\ln(e^{2x}-1)b^2}{a^3} + \frac{b^4 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^3(a^2-b^2)}$

input

```
int(coth(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
(2*ln(a+b*tanh(x))*b^4-2*ln(1-tanh(x))*a^4+(2*a^4-2*b^4)*ln(tanh(x))-2*(a+b)*(1/2*coth(x)^2*a*(a-b)-b*coth(x)*(a-b)+a^2*x)*a)/(2*a^5-2*a^3*b^2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 641 vs. 2(74) = 148.

Time = 0.12 (sec) , antiderivative size = 641, normalized size of antiderivative = 8.43

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input

```
integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="fricas")
```

output

```

-((a^4 + a^3*b)*x*cosh(x)^4 + 4*(a^4 + a^3*b)*x*cosh(x)*sinh(x)^3 + (a^4 +
a^3*b)*x*sinh(x)^4 + 2*a^3*b - 2*a*b^3 + 2*(a^4 - a^3*b - a^2*b^2 + a*b^3
- (a^4 + a^3*b)*x)*cosh(x)^2 + 2*(a^4 - a^3*b - a^2*b^2 + a*b^3 + 3*(a^4
+ a^3*b)*x*cosh(x)^2 - (a^4 + a^3*b)*x)*sinh(x)^2 + (a^4 + a^3*b)*x - (b^4
*cosh(x)^4 + 4*b^4*cosh(x)*sinh(x)^3 + b^4*sinh(x)^4 - 2*b^4*cosh(x)^2 + b
^4 + 2*(3*b^4*cosh(x)^2 - b^4)*sinh(x)^2 + 4*(b^4*cosh(x)^3 - b^4*cosh(x))
*sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - ((a^4 - b^4
)*cosh(x)^4 + 4*(a^4 - b^4)*cosh(x)*sinh(x)^3 + (a^4 - b^4)*sinh(x)^4 + a^
4 - b^4 - 2*(a^4 - b^4)*cosh(x)^2 - 2*(a^4 - b^4 - 3*(a^4 - b^4)*cosh(x)^2
)*sinh(x)^2 + 4*((a^4 - b^4)*cosh(x)^3 - (a^4 - b^4)*cosh(x))*sinh(x))*log
(2*sinh(x)/(cosh(x) - sinh(x))) + 4*((a^4 + a^3*b)*x*cosh(x)^3 + (a^4 - a^
3*b - a^2*b^2 + a*b^3 - (a^4 + a^3*b)*x)*cosh(x))*sinh(x))/(a^5 - a^3*b^2
+ (a^5 - a^3*b^2)*cosh(x)^4 + 4*(a^5 - a^3*b^2)*cosh(x)*sinh(x)^3 + (a^5 -
a^3*b^2)*sinh(x)^4 - 2*(a^5 - a^3*b^2)*cosh(x)^2 - 2*(a^5 - a^3*b^2 - 3*(
a^5 - a^3*b^2)*cosh(x)^2)*sinh(x)^2 + 4*((a^5 - a^3*b^2)*cosh(x)^3 - (a^5
- a^3*b^2)*cosh(x))*sinh(x))

```

### Sympy [F]

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \int \frac{\coth^3(x)}{a + b \tanh(x)} dx$$

input

```
integrate(coth(x)**3/(a+b*tanh(x)),x)
```

output

```
Integral(coth(x)**3/(a + b*tanh(x)), x)
```

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.59

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \frac{b^4 \log(-(a-b)e^{(-2x)} - a - b)}{a^5 - a^3 b^2} + \frac{2((a+b)e^{(-2x)} - b)}{2a^2 e^{(-2x)} - a^2 e^{(-4x)} - a^2} + \frac{x}{a+b} + \frac{(a^2 + b^2) \log(e^{(-x)} + 1)}{a^3} + \frac{(a^2 + b^2) \log(e^{(-x)} - 1)}{a^3}$$

input `integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

output 
$$b^4 \log(-(a-b)e^{-2x} - a - b)/(a^5 - a^3 b^2) + 2*((a+b)e^{-2x} - b)/(2a^2 e^{-2x} - a^2 e^{-4x} - a^2) + x/(a+b) + (a^2 + b^2) \log(e^{-x} + 1)/a^3 + (a^2 + b^2) \log(e^{-x} - 1)/a^3$$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.28

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \frac{b^4 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^5 - a^3 b^2} - \frac{x}{a - b} + \frac{(a^2 + b^2) \log(|e^{(2x)} - 1|)}{a^3} - \frac{2(ab + (a^2 - ab)e^{(2x)})}{a^3(e^{(2x)} - 1)^2}$$

input `integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="giac")`

output 
$$b^4 \log(\text{abs}(a e^{(2x)} + b e^{(2x)} + a - b))/(a^5 - a^3 b^2) - x/(a - b) + (a^2 + b^2) \log(\text{abs}(e^{(2x)} - 1))/a^3 - 2*(a*b + (a^2 - a*b)*e^{(2x)})/(a^3 * (e^{(2x)} - 1)^2)$$

### Mupad [B] (verification not implemented)

Time = 2.45 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.46

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \frac{\ln(e^{2x} - 1)(a^2 + b^2)}{a^3} - \frac{x}{a - b} - \frac{2}{a(e^{4x} - 2e^{2x} + 1)} + \frac{b^4 \ln(a - b + a e^{2x} + b e^{2x})}{a^5 - a^3 b^2} - \frac{2(a^2 - b^2)}{a^2(a + b)(e^{2x} - 1)}$$

input `int(coth(x)^3/(a + b*tanh(x)),x)`



output  $(\log(\exp(2*x) - 1)*(a^2 + b^2))/a^3 - x/(a - b) - 2/(a*(\exp(4*x) - 2*\exp(2*x) + 1)) + (b^4*\log(a - b + a*\exp(2*x) + b*\exp(2*x)))/(a^5 - a^3*b^2) - (2*(a^2 - b^2))/(a^2*(a + b)*(exp(2*x) - 1))$

### Reduce [F]

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx = \int \frac{\coth(x)^3}{\tanh(x) b + a} dx$$

input `int(coth(x)^3/(a+b*tanh(x)),x)`

output `int(coth(x)^3/(a+b*tanh(x)),x)`

### 3.143 $\int \frac{\coth^4(x)}{a+b \tanh(x)} dx$

Optimal result	1105
Mathematica [A] (verified)	1105
Rubi [C] (verified)	1106
Maple [A] (verified)	1112
Fricas [B] (verification not implemented)	1112
Sympy [F]	1113
Maxima [A] (verification not implemented)	1114
Giac [A] (verification not implemented)	1114
Mupad [B] (verification not implemented)	1115
Reduce [F]	1115

#### Optimal result

Integrand size = 13, antiderivative size = 97

$$\int \frac{\coth^4(x)}{a+b \tanh(x)} dx = \frac{ax}{a^2-b^2} - \frac{(a^2+b^2)\coth(x)}{a^3} + \frac{b\coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} - \frac{b(a^2+b^2)\log(\sinh(x))}{a^4} - \frac{b^5\log(a\cosh(x)+b\sinh(x))}{a^4(a^2-b^2)}$$

output

```
a*x/(a^2-b^2)-(a^2+b^2)*coth(x)/a^3+1/2*b*coth(x)^2/a^2-1/3*coth(x)^3/a-b*(a^2+b^2)*ln(sinh(x))/a^4-b^5*ln(a*cosh(x)+b*sinh(x))/a^4/(a^2-b^2)
```

#### Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.95

$$\int \frac{\coth^4(x)}{a+b \tanh(x)} dx = \frac{1}{6} \left( -\frac{6(a^2+b^2)\coth(x)}{a^3} + \frac{3b\coth^2(x)}{a^2} - \frac{2\coth^3(x)}{a} - \frac{3\log(1-\coth(x))}{a+b} + \frac{3\log(1+\coth(x))}{a-b} + \frac{6b^5\log(b+a\coth(x))}{a^4(-a^2+b^2)} \right)$$

input

```
Integrate[Coth[x]^4/(a + b*Tanh[x]), x]
```

output

$$\frac{((-6*(a^2 + b^2)*\text{Coth}[x])/a^3 + (3*b*\text{Coth}[x]^2)/a^2 - (2*\text{Coth}[x]^3)/a - (3*\text{Log}[1 - \text{Coth}[x]])/(a + b) + (3*\text{Log}[1 + \text{Coth}[x]])/(a - b) + (6*b^5*\text{Log}[b + a*\text{Coth}[x]])/(a^4*(-a^2 + b^2)))/6}$$
**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.07 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.35, number of steps used = 19, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.462$ , Rules used = {3042, 4052, 27, 3042, 26, 4132, 27, 3042, 25, 4133, 25, 3042, 26, 4134, 26, 3042, 26, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\coth^4(x)}{a + b \tanh(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\tan(ix)^4(a - ib \tan(ix))} dx \\ & \quad \downarrow \text{4052} \\ & -\frac{\int \frac{3 \coth^3(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{3a} - \frac{\coth^3(x)}{3a} \\ & \quad \downarrow \text{27} \\ & -\frac{\int \frac{\coth^3(x)(-b \tanh^2(x) - a \tanh(x) + b)}{a + b \tanh(x)} dx}{a} - \frac{\coth^3(x)}{3a} \\ & \quad \downarrow \text{3042} \\ & -\frac{\coth^3(x)}{3a} - \frac{\int -\frac{i(b \tan(ix)^2 + ia \tan(ix) + b)}{\tan(ix)^3(a - ib \tan(ix))} dx}{a} \\ & \quad \downarrow \text{26} \\ & -\frac{\coth^3(x)}{3a} + \frac{i \int \frac{b \tan(ix)^2 + ia \tan(ix) + b}{\tan(ix)^3(a - ib \tan(ix))} dx}{a} \end{aligned}$$

$$\begin{aligned}
 & \downarrow 4132 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( -\int \frac{2i \operatorname{coth}^2(x)(a^2+b^2-b^2 \tanh^2(x))}{a+b \tanh(x)} dx - \frac{ib \operatorname{coth}^2(x)}{2a} \right)}{a} \\
 & \downarrow 27 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( -i \int \frac{\operatorname{coth}^2(x)(a^2+b^2-b^2 \tanh^2(x))}{a+b \tanh(x)} dx - \frac{ib \operatorname{coth}^2(x)}{2a} \right)}{a} \\
 & \downarrow 3042 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( -\int \frac{a^2+b^2+b^2 \tan(ix)^2}{\tan(ix)^2(a-ib \tan(ix))} dx - \frac{ib \operatorname{coth}^2(x)}{2a} \right)}{a} \\
 & \downarrow 25 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( \int \frac{a^2+b^2+b^2 \tan(ix)^2}{\tan(ix)^2(a-ib \tan(ix))} dx - \frac{ib \operatorname{coth}^2(x)}{2a} \right)}{a} \\
 & \downarrow 4133 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( \frac{(a^2+b^2) \operatorname{coth}(x)}{a} - \int \frac{\operatorname{coth}(x)(-\tanh(x)a^3-b(a^2+b^2) \tanh^2(x)+b(a^2+b^2))}{a+b \tanh(x)} dx \right)}{a} - \frac{ib \operatorname{coth}^2(x)}{2a} \\
 & \downarrow 25 \\
 & -\frac{\operatorname{coth}^3(x)}{3a} + \frac{i \left( \frac{\int \frac{\operatorname{coth}(x)(-\tanh(x)a^3-b(a^2+b^2) \tanh^2(x)+b(a^2+b^2))}{a+b \tanh(x)} dx + \frac{(a^2+b^2) \operatorname{coth}(x)}{a}}{a} - \frac{ib \operatorname{coth}^2(x)}{2a} \right)}{a} \\
 & \downarrow 3042
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{\coth^3(x)}{3a} + \frac{i \left( \frac{\left( \frac{(a^2+b^2)\coth(x)}{a} + \frac{\int \frac{i \tan(ix)a^3+b(a^2+b^2)\tan(ix)^2+b(a^2+b^2)}{\tan(ix)(a-ib \tan(ix))} dx}{a} \right)}{a} - \frac{ib \coth^2(x)}{2a} \right)}{a} \\
 & \quad \downarrow 26 \\
 & -\frac{\coth^3(x)}{3a} + \frac{i \left( \frac{\left( \frac{(a^2+b^2)\coth(x)}{a} + i \int \frac{i \tan(ix)a^3+b(a^2+b^2)\tan(ix)^2+b(a^2+b^2)}{\tan(ix)(a-ib \tan(ix))} dx \right)}{a} - \frac{ib \coth^2(x)}{2a} \right)}{a} \\
 & \quad \downarrow 4134 \\
 & -\frac{\coth^3(x)}{3a} + \frac{i \left( \frac{\left( \frac{(a^2+b^2)\coth(x)}{a} + i \left( \frac{b(a^2+b^2) \int -i \coth(x) dx}{a} + \frac{b^5 \int -\frac{i(b+a \tanh(x))}{a+b \tanh(x)} dx + \frac{ia^4 x}{a^2-b^2} \right) \right)}{a} - \frac{ib \coth^2(x)}{2a} \right)}{a} \\
 & \quad \downarrow 26 \\
 & -\frac{\coth^3(x)}{3a} + \frac{i \left( \frac{\left( \frac{(a^2+b^2)\coth(x)}{a} + i \left( -\frac{ib(a^2+b^2) \int \coth(x) dx}{a} - \frac{ib^5 \int \frac{b+a \tanh(x)}{a+b \tanh(x)} dx + \frac{ia^4 x}{a^2-b^2} \right) \right)}{a} - \frac{ib \coth^2(x)}{2a} \right)}{a} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\begin{array}{c}
 \frac{\coth^3(x)}{3a} + \\
 i \left( \frac{i \left( \frac{(a^2+b^2) \coth(x)}{a} + \frac{i \left( -\frac{ib(a^2+b^2) \int -i \tan(ix+\frac{\pi}{2}) dx}{a} - \frac{ib^5 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} + \frac{ia^4 x}{a^2-b^2} \right)}{a} \right)}{a} - \frac{ib \coth^2(x)}{2a} \right) \\
 \hline
 a \\
 \downarrow \text{26} \\
 \frac{\coth^3(x)}{3a} + \\
 i \left( \frac{i \left( \frac{(a^2+b^2) \coth(x)}{a} + \frac{i \left( -\frac{b(a^2+b^2) \int \tan(ix+\frac{\pi}{2}) dx}{a} - \frac{ib^5 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} + \frac{ia^4 x}{a^2-b^2} \right)}{a} \right)}{a} - \frac{ib \coth^2(x)}{2a} \right) \\
 \hline
 a \\
 \downarrow \text{3956} \\
 \frac{\coth^3(x)}{3a} + \\
 i \left( \frac{i \left( \frac{(a^2+b^2) \coth(x)}{a} + \frac{i \left( -\frac{ib^5 \int \frac{b-ia \tan(ix)}{a-ib \tan(ix)} dx}{a(a^2-b^2)} - \frac{ib(a^2+b^2) \log(\sinh(x))}{a} + \frac{ia^4 x}{a^2-b^2} \right)}{a} \right)}{a} - \frac{ib \coth^2(x)}{2a} \right) \\
 \hline
 a \\
 \downarrow \text{4013} \\
 \frac{\coth^3(x)}{3a} + \\
 i \left( \frac{i \left( \frac{(a^2+b^2) \coth(x)}{a} + \frac{i \left( -\frac{ib(a^2+b^2) \log(\sinh(x))}{a} - \frac{ib^5 \log(a \cosh(x)+b \sinh(x))}{a(a^2-b^2)} + \frac{ia^4 x}{a^2-b^2} \right)}{a} \right)}{a} - \frac{ib \coth^2(x)}{2a} \right) \\
 \hline
 a
 \end{array}$$

input `Int[Coth[x]^4/(a + b*Tanh[x]),x]`

output `-1/3*Coth[x]^3/a + (I*((( -1/2*I)*b*Coth[x]^2)/a + (I*(((a^2 + b^2)*Coth[x])/a + (I*((I*a^4*x)/(a^2 - b^2) - (I*b*(a^2 + b^2)*Log[Sinh[x]))/a - (I*b^5*Log[a*Cosh[x] + b*Sinh[x]])/(a*(a^2 - b^2))))/a))/a)/a`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sinh[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4052

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c
+ d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1
/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c +
d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c -
a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0]
&& NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || Integ
erQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4132

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4133

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[(A*b^2 + a^2*C)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d
)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Sim
p[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - a*C*(b*c*(m + 1) + a*d*(n
+ 1)) - (m + 1)*(b*c - a*d)*(A*b - b*C)*Tan[e + f*x] - d*(A*b^2 + a^2*C)*(
m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, n},
x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m
, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```



rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] :> Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

### Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.18

method	result
derivativedivides	$\frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{b}{2a^2 \tanh(x)^2} + \frac{-a^2-b^2}{a^3 \tanh(x)} - \frac{(a^2+b^2)b \ln(\tanh(x))}{a^4} - \frac{1}{3a \tanh(x)^3} - \frac{b^5}{3a \tanh(x)^3}$
default	$\frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{b}{2a^2 \tanh(x)^2} + \frac{-a^2-b^2}{a^3 \tanh(x)} - \frac{(a^2+b^2)b \ln(\tanh(x))}{a^4} - \frac{1}{3a \tanh(x)^3} - \frac{b^5}{3a \tanh(x)^3}$
parallelrisch	$\frac{-6b^5 \ln(a+b \tanh(x))+6 \ln(1-\tanh(x))a^4b+(-6a^4b+6b^5) \ln(\tanh(x))+(-2a^5+2a^3b^2) \coth(x)^3+(3a^4b-3a^2b^3) \coth(x)}{6a^6-6a^4b^2}$
risch	$\frac{x}{a+b} + \frac{2xb}{a^2} + \frac{2xb^3}{a^4} + \frac{2xb^5}{a^4(a^2-b^2)} - \frac{2(6a^2e^{4x}-3e^{4x}ab+3b^2e^{4x}-6e^{2x}a^2+3e^{2x}ab-6b^2e^{2x}+4a^2+3b^2)}{3a^3(e^{2x}-1)^3} - \frac{b \ln(e^{2x}-1)}{3a^3(e^{2x}-1)^3}$

input

```
int(coth(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

output

```
1/(2*a-2*b)*ln(1+tanh(x))-1/(2*b+2*a)*ln(tanh(x)-1)+1/2*b/a^2/tanh(x)^2+(-a^2-b^2)/a^3/tanh(x)-(a^2+b^2)/a^4*b*ln(tanh(x))-1/3/a/tanh(x)^3-b^5/(a-b)/(a+b)/a^4*ln(a+b*tanh(x))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1299 vs. 2(93) = 186.

Time = 0.12 (sec) , antiderivative size = 1299, normalized size of antiderivative = 13.39

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx = \text{Too large to display}$$

input `integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

output

$$\begin{aligned} & 1/3*(3*(a^5 + a^4*b)*x*\cosh(x)^6 + 18*(a^5 + a^4*b)*x*\cosh(x)*\sinh(x)^5 + \\ & 3*(a^5 + a^4*b)*x*\sinh(x)^6 - 8*a^5 + 2*a^3*b^2 + 6*a*b^4 - 3*(4*a^5 - 2*a^4*b - \\ & 2*a^3*b^2 + 2*a^2*b^3 - 2*a*b^4 + 3*(a^5 + a^4*b)*x)*\cosh(x)^4 - 3*(4*a^5 - \\ & 2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 - 2*a*b^4 - 15*(a^5 + a^4*b)*x*\cosh(x)^2 + \\ & 3*(a^5 + a^4*b)*x)*\sinh(x)^4 + 12*(5*(a^5 + a^4*b)*x*\cosh(x)^3 - (4*a^5 - \\ & 2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 - 2*a*b^4 + 3*(a^5 + a^4*b)*x)*\cosh(x))*\sinh(x)^3 + \\ & 3*(4*a^5 - 2*a^4*b + 2*a^2*b^3 - 4*a*b^4 + 3*(a^5 + a^4*b)*x)*\cosh(x)^2 + 3*(15*(a^5 + a^4*b)*x*\cosh(x)^4 + \\ & 4*a^5 - 2*a^4*b + 2*a^2*b^3 - 4*a*b^4 - 6*(4*a^5 - 2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 - 2*a*b^4 + \\ & 3*(a^5 + a^4*b)*x)*\cosh(x)^2 + 3*(a^5 + a^4*b)*x)*\sinh(x)^2 - 3*(a^5 + a^4*b)*x - \\ & 3*(b^5*\cosh(x)^6 + 6*b^5*\cosh(x)*\sinh(x)^5 + b^5*\sinh(x)^6 - 3*b^5*\cosh(x)^4 + \\ & 3*b^5*\cosh(x)^2 - b^5 + 3*(5*b^5*\cosh(x)^2 - b^5)*\sinh(x)^4 + 4*(5*b^5*\cosh(x)^3 - \\ & 3*b^5*\cosh(x))*\sinh(x)^3 + 3*(5*b^5*\cosh(x)^4 - 6*b^5*\cosh(x)^2 + b^5)*\sinh(x)^2 + \\ & 6*(b^5*\cosh(x)^5 - 2*b^5*\cosh(x)^3 + b^5*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) - \\ & 3*((a^4*b - b^5)*\cosh(x)^6 + 6*(a^4*b - b^5)*\cosh(x)*\sinh(x)^5 + (a^4*b - b^5)*\sinh(x)^6 - \\ & a^4*b + b^5 - 3*(a^4*b - b^5)*\cosh(x)^4 - 3*(a^4*b - b^5 - 5*(a^4*b - b^5)*\cosh(x)^2)*\sinh(x)^4 + \\ & 4*(5*(a^4*b - b^5)*\cosh(x)^3 - 3*(a^4*b - b^5)*\cosh(x))*\sinh(x)^3 + 3*(a^4*b - b^5)*\cosh(x)^2 + \\ & 3*(a^4*b - b^5 + 5*(a^4*b - b^5)*\cosh(x)^4 - 6*(a^4*b - b^5)*\cosh(x)^2)*\sinh(x)^2 + 6*... \end{aligned}$$

## Sympy [F]

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx = \int \frac{\coth^4(x)}{a + b \tanh(x)} dx$$

input `integrate(coth(x)**4/(a+b*tanh(x)),x)`

output `Integral(coth(x)**4/(a + b*tanh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.78

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx$$

$$= -\frac{b^5 \log(-(a-b)e^{-2x} - a - b)}{a^6 - a^4 b^2}$$

$$+ \frac{2(4a^2 + 3b^2 - 3(2a^2 + ab + 2b^2)e^{-2x}) + 3(2a^2 + ab + b^2)e^{-4x}}{3(3a^3 e^{-2x} - 3a^3 e^{-4x} + a^3 e^{-6x} - a^3)}$$

$$+ \frac{x}{a+b} - \frac{(a^2 b + b^3) \log(e^{-x} + 1)}{a^4} - \frac{(a^2 b + b^3) \log(e^{-x} - 1)}{a^4}$$

input `integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`output `-b^5*log(-(a - b)*e^(-2*x) - a - b)/(a^6 - a^4*b^2) + 2/3*(4*a^2 + 3*b^2 - 3*(2*a^2 + a*b + 2*b^2)*e^(-2*x) + 3*(2*a^2 + a*b + b^2)*e^(-4*x))/(3*a^3 *e^(-2*x) - 3*a^3*e^(-4*x) + a^3*e^(-6*x) - a^3) + x/(a + b) - (a^2*b + b^3)*log(e^(-x) + 1)/a^4 - (a^2*b + b^3)*log(e^(-x) - 1)/a^4`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.46

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx$$

$$= -\frac{b^5 \log(|ae^{2x} + be^{2x} + a - b|)}{a^6 - a^4 b^2} + \frac{x}{a-b} - \frac{(a^2 b + b^3) \log(|e^{2x} - 1|)}{a^4}$$

$$- \frac{2(4a^3 + 3ab^2 + 3(2a^3 - a^2 b + ab^2)e^{4x}) - 3(2a^3 - a^2 b + 2ab^2)e^{2x}}{3a^4(e^{2x} - 1)^3}$$

input `integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="giac")`

output

```
-b^5*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^6 - a^4*b^2) + x/(a - b) -
(a^2*b + b^3)*log(abs(e^(2*x) - 1))/a^4 - 2/3*(4*a^3 + 3*a*b^2 + 3*(2*a^3
- a^2*b + a*b^2)*e^(4*x) - 3*(2*a^3 - a^2*b + 2*a*b^2)*e^(2*x))/(a^4*(e^(
2*x) - 1)^3)
```

**Mupad [B] (verification not implemented)**

Time = 2.66 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.68

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx = \frac{x}{a - b} - \frac{8}{3a(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{b^5 \ln(a - b + ae^{2x} + be^{2x})}{a^6 - a^4 b^2} - \frac{\ln(e^{2x} - 1)(a^2 b + b^3)}{a^4} - \frac{2(2a^3 + a^2 b + b^3)}{a^3(a + b)(e^{2x} - 1)} - \frac{2(2a^2 + ab - b^2)}{a^2(a + b)(e^{4x} - 2e^{2x} + 1)}$$

input

```
int(coth(x)^4/(a + b*tanh(x)),x)
```

output

```
x/(a - b) - 8/(3*a*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) - (b^5*log(a
- b + a*exp(2*x) + b*exp(2*x)))/(a^6 - a^4*b^2) - (log(exp(2*x) - 1)*(a^2*
b + b^3))/a^4 - (2*(a^2*b + 2*a^3 + b^3))/(a^3*(a + b)*(exp(2*x) - 1)) - (
2*(a*b + 2*a^2 - b^2))/(a^2*(a + b)*(exp(4*x) - 2*exp(2*x) + 1))
```

**Reduce [F]**

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx = \int \frac{\coth(x)^4}{\tanh(x)b + a} dx$$

input

```
int(coth(x)^4/(a+b*tanh(x)),x)
```

output

```
int(coth(x)^4/(a+b*tanh(x)),x)
```

### 3.144 $\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx$

Optimal result	1116
Mathematica [A] (verified)	1116
Rubi [A] (verified)	1117
Maple [A] (verified)	1119
Fricas [B] (verification not implemented)	1119
Sympy [F]	1120
Maxima [A] (verification not implemented)	1120
Giac [B] (verification not implemented)	1120
Mupad [B] (verification not implemented)	1121
Reduce [B] (verification not implemented)	1121

#### Optimal result

Integrand size = 14, antiderivative size = 55

$$\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx = \frac{ax}{b(a^2-b^2)} - \frac{\log(a \cosh(x) + b \sinh(x))}{a^2-b^2} - \frac{x}{b(a+b \tanh(x))}$$

output

```
a*x/b/(a^2-b^2)-ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)-x/b/(a+b*tanh(x))
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

$$\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx = \frac{bx - a \log(a \cosh(x) + b \sinh(x))}{a^3 - ab^2} + \frac{x \sinh(x)}{a^2 \cosh(x) + ab \sinh(x)}$$

input

```
Integrate[(x*Sech[x]^2)/(a + b*Tanh[x])^2,x]
```

output

```
(b*x - a*Log[a*Cosh[x] + b*Sinh[x]])/(a^3 - a*b^2) + (x*Sinh[x])/(a^2*Cosh[x] + a*b*Sinh[x])
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {5989, 3042, 3965, 26, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx \\
 & \quad \downarrow \text{5989} \\
 & \frac{\int \frac{1}{a + b \tanh(x)} dx}{b} - \frac{x}{b(a + b \tanh(x))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{x}{b(a + b \tanh(x))} + \frac{\int \frac{1}{a - ib \tan(ix)} dx}{b} \\
 & \quad \downarrow \text{3965} \\
 & -\frac{x}{b(a + b \tanh(x))} + \frac{\frac{ax}{a^2 - b^2} - \frac{ib \int \frac{i(b+a \tanh(x))}{a + b \tanh(x)} dx}{b}}{b} \\
 & \quad \downarrow \text{26} \\
 & \frac{\frac{ax}{a^2 - b^2} - \frac{b \int \frac{b+a \tanh(x)}{a + b \tanh(x)} dx}{a^2 - b^2}}{b} - \frac{x}{b(a + b \tanh(x))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{x}{b(a + b \tanh(x))} + \frac{\frac{ax}{a^2 - b^2} - \frac{b \int \frac{b-ia \tan(ix)}{a - ib \tan(ix)} dx}{a^2 - b^2}}{b} \\
 & \quad \downarrow \text{4013} \\
 & \frac{\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}}{b} - \frac{x}{b(a + b \tanh(x))}
 \end{aligned}$$

input `Int [(x*Sech[x]^2)/(a + b*Tanh[x])^2, x]`

output  $((a*x)/(a^2 - b^2) - (b*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])/(a^2 - b^2))/b - x/(b*(a + b*\text{Tanh}[x]))$

### Defintions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3965  $\text{Int}[(a + (b*\text{tan}[c + (d*x)])^{-1}), x\_Symbol] \rightarrow \text{Simp}[a*(x/(a^2 + b^2)), x] + \text{Simp}[b/(a^2 + b^2) \text{Int}[(b - a*\text{Tan}[c + d*x])/(a + b*\text{Tan}[c + d*x]), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

rule 4013  $\text{Int}[(c + (d*\text{tan}[e + (f*x)])/(a + (b*\text{tan}[e + (f*x)])), x\_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

rule 5989  $\text{Int}[(e + (f*x))^m * \text{Sech}[c + (d*x)]^2 * (a + (b*\text{Tanh}[c + (d*x)]))^n], x\_Symbol] \rightarrow \text{Simp}[(e + f*x)^m * ((a + b*\text{Tanh}[c + d*x])^{n+1} / (b*d*(n+1))), x] - \text{Simp}[f*(m/(b*d*(n+1))) \text{Int}[(e + f*x)^{m-1} * (a + b*\text{Tanh}[c + d*x])^{n+1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{NeQ}[n, -1]$

**Maple [A] (verified)**

Time = 8.96 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.33

method	result	size
risch	$\frac{2x}{a^2-b^2} - \frac{2x}{(e^{2x}a+e^{2x}b+a-b)(a+b)} - \frac{\ln\left(\frac{e^{2x}+a-b}{a+b}\right)}{a^2-b^2}$	73

input `int(x*sech(x)^2/(a+b*tanh(x))^2,x,method=_RETURNVERBOSE)`

output  $\frac{2}{(a^2-b^2)*x-2*x/(\exp(2*x)*a+\exp(2*x)*b+a-b)/(a+b)-1/(a^2-b^2)*\ln(\exp(2*x)+(a-b)/(a+b))}$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 182 vs.  $2(55) = 110$ .

Time = 0.10 (sec) , antiderivative size = 182, normalized size of antiderivative = 3.31

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx$$

$$= \frac{2(a+b)x \cosh(x)^2 + 4(a+b)x \cosh(x) \sinh(x) + 2(a+b)x \sinh(x)^2 - ((a+b) \cosh(x))^2 + 2(a+b) \cosh(x) \sinh(x) - (a+b) \sinh(x)^2}{a^3 - a^2b - ab^2 + b^3 + (a^3 + a^2b - ab^2 - b^3) \cosh(x)^2 + 2(a^3 + a^2b - ab^2 - b^3) \cosh(x) \sinh(x) + (a^3 + a^2b - ab^2 - b^3) \sinh(x)^2}$$

input `integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="fricas")`

output  $(2*(a + b)*x*\cosh(x)^2 + 4*(a + b)*x*\cosh(x)*\sinh(x) + 2*(a + b)*x*\sinh(x)^2 - ((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x)))/(a^3 - a^2*b - a*b^2 + b^3 + (a^3 + a^2*b - a*b^2 - b^3)*\cosh(x)^2 + 2*(a^3 + a^2*b - a*b^2 - b^3)*\cosh(x)*\sinh(x) + (a^3 + a^2*b - a*b^2 - b^3)*\sinh(x)^2)$



**Sympy [F]**

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx = \int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx$$

input `integrate(x*sech(x)**2/(a+b*tanh(x))**2,x)`

output `Integral(x*sech(x)**2/(a + b*tanh(x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.24

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx = \frac{2 x e^{(2x)}}{a^2 - 2 a b + b^2 + (a^2 - b^2) e^{(2x)}} - \frac{\log\left(\frac{(a+b)e^{(2x)}+a-b}{a+b}\right)}{a^2 - b^2}$$

input `integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="maxima")`

output `2*x*e^(2*x)/(a^2 - 2*a*b + b^2 + (a^2 - b^2)*e^(2*x)) - log(((a + b)*e^(2*x) + a - b)/(a + b))/(a^2 - b^2)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 174 vs. 2(55) = 110.

Time = 0.13 (sec) , antiderivative size = 174, normalized size of antiderivative = 3.16

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx = \frac{2 a x e^{(2x)} + 2 b x e^{(2x)} - a e^{(2x)} \log(-a e^{(2x)} - b e^{(2x)} - a + b) - b e^{(2x)} \log(-a e^{(2x)} - b e^{(2x)} - a + b) - a \log(-a e^{(2x)} - b e^{(2x)} - a + b)}{a^3 e^{(2x)} + a^2 b e^{(2x)} - a b^2 e^{(2x)} - b^3 e^{(2x)} + a^3 - a^2 b}$$

input `integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="giac")`

output

$$\frac{(2ax^2e^{2x} + 2bx^2e^{2x} - ae^{2x}\log(-ae^{2x} - be^{2x} - a + b) - be^{2x}\log(-ae^{2x} - be^{2x} - a + b) - a\log(-ae^{2x} - be^{2x} - a + b) + b\log(-ae^{2x} - be^{2x} - a + b))/(a^3e^{2x} + a^2be^{2x} - ab^2e^{2x} - b^3e^{2x} + a^3 - a^2b - ab^2 + b^3)}$$
**Mupad [B] (verification not implemented)**

Time = 2.34 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.25

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx = \frac{2x}{a^2 - b^2} - \frac{\ln(a - b + ae^{2x} + be^{2x})}{a^2 - b^2} - \frac{2x}{(a + b)(a - b + e^{2x}(a + b))}$$

input

`int(x/(cosh(x)^2*(a + b*tanh(x))^2),x)`

output

$$\frac{(2x)/(a^2 - b^2) - \log(a - b + a\exp(2x) + b\exp(2x))/(a^2 - b^2) - (2x)/((a + b)*(a - b + \exp(2x)*(a + b)))}$$
**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 2513, normalized size of antiderivative = 45.69

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx = \text{Too large to display}$$

input

`int(x*sech(x)^2/(a+b*tanh(x))^2,x)`

output

```
( - 18***(6*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*tanh(x)*a**4*b**2 - 2
4***(6*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*tanh(x)*a**3*b**3 - 6***(
6*x)*log(e**(2*x)*a + e**(2*x)*b + a - b)*tanh(x)*a**2*b**4 - 18***(6*x)*
log(e**(2*x)*a + e**(2*x)*b + a - b)*a**5*b - 24***(6*x)*log(e**(2*x)*a +
e**(2*x)*b + a - b)*a**4*b**2 - 6***(6*x)*log(e**(2*x)*a + e**(2*x)*b +
a - b)*a**3*b**3 + 3***(6*x)*sech(x)**2*tanh(x)*a**4*b**2 + 4***(6*x)*se
ch(x)**2*tanh(x)*a**3*b**3 - 2***(6*x)*sech(x)**2*tanh(x)*a**2*b**4 - 4***
(6*x)*sech(x)**2*tanh(x)*a*b**5 - e**(6*x)*sech(x)**2*tanh(x)*b**6 - 18*
e**(6*x)*sech(x)**2*a**6*x - 24***(6*x)*sech(x)**2*a**5*b*x - 9***(6*x)*
sech(x)**2*a**5*b + 12***(6*x)*sech(x)**2*a**4*b**2*x - 12***(6*x)*sech(
x)**2*a**4*b**2 + 24***(6*x)*sech(x)**2*a**3*b**3*x + 6***(6*x)*sech(x)*
**2*a**3*b**3 + 6***(6*x)*sech(x)**2*a**2*b**4*x + 12***(6*x)*sech(x)**2*
a**2*b**4 + 3***(6*x)*sech(x)**2*a*b**5 + 36***(6*x)*tanh(x)*a**4*b**2*x
- 12***(6*x)*tanh(x)*a**4*b**2 + 48***(6*x)*tanh(x)*a**3*b**3*x - 8***
(6*x)*tanh(x)*a**3*b**3 + 12***(6*x)*tanh(x)*a**2*b**4*x + 16***(6*x)*ta
nh(x)*a**2*b**4 + 8***(6*x)*tanh(x)*a*b**5 - 4***(6*x)*tanh(x)*b**6 + 36
***(6*x)*a**5*b*x - 12***(6*x)*a**5*b + 48***(6*x)*a**4*b**2*x - 8***(
6*x)*a**4*b**2 + 12***(6*x)*a**3*b**3*x + 16***(6*x)*a**3*b**3 + 8***(6
*x)*a**2*b**4 - 4***(6*x)*a*b**5 - 54***(4*x)*log(e**(2*x)*a + e**(2*x)*
b + a - b)*tanh(x)*a**4*b**2 - 36***(4*x)*log(e**(2*x)*a + e**(2*x)*b ...
```

**3.145**  $\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$

Optimal result	1123
Mathematica [A] (verified)	1124
Rubi [A] (verified)	1124
Maple [B] (verified)	1127
Fricas [B] (verification not implemented)	1128
Sympy [F]	1129
Maxima [F]	1130
Giac [F]	1130
Mupad [F(-1)]	1130
Reduce [F]	1131

**Optimal result**

Integrand size = 24, antiderivative size = 231

$$\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx = \frac{x \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd}} - \frac{x \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd}} + \frac{\operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a}\sqrt{bd}^2} - \frac{\operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a}\sqrt{bd}^2}$$

output

```
1/2*x*ln(1+(a+b)*exp(2*d*x+2*c)/(a-2*(-a)^(1/2)*b^(1/2)-b))/(-a)^(1/2)/b^(1/2)/d-1/2*x*ln(1+(a+b)*exp(2*d*x+2*c)/(a+2*(-a)^(1/2)*b^(1/2)-b))/(-a)^(1/2)/b^(1/2)/d+1/4*polylog(2,-(a+b)*exp(2*d*x+2*c)/(a-2*(-a)^(1/2)*b^(1/2)-b))/(-a)^(1/2)/b^(1/2)/d^2-1/4*polylog(2,-(a+b)*exp(2*d*x+2*c)/(a+2*(-a)^(1/2)*b^(1/2)-b))/(-a)^(1/2)/b^(1/2)/d^2
```

**Mathematica [A] (verified)**

Time = 1.31 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.08

$$\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$$

$$= \frac{-4\sqrt{-ac} \arctan\left(\frac{a-b+(a+b)e^{2(c+dx)}}{2\sqrt{a}\sqrt{b}}\right) + 2\sqrt{a}(c+dx) \log\left(1 + \frac{(a+b)e^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b-b}}\right) - 2\sqrt{a}(c+dx) \log\left(1 + \frac{(a+b)e^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a^2}\sqrt{bd^2}}$$

input `Integrate[(x*Sech[c + d*x]^2)/(a + b*Tanh[c + d*x]^2),x]`

output `(-4*Sqrt[-a]*c*ArcTan[(a - b + (a + b)*E^(2*(c + d*x)))/(2*Sqrt[a]*Sqrt[b])] + 2*Sqrt[a]*(c + d*x)*Log[1 + ((a + b)*E^(2*(c + d*x)))/(a - 2*Sqrt[-a]*Sqrt[b] - b)] - 2*Sqrt[a]*(c + d*x)*Log[1 + ((a + b)*E^(2*(c + d*x)))/(a + 2*Sqrt[-a]*Sqrt[b] - b)] + Sqrt[a]*PolyLog[2, -((a + b)*E^(2*(c + d*x)))/(a - 2*Sqrt[-a]*Sqrt[b] - b))] - Sqrt[a]*PolyLog[2, -((a + b)*E^(2*(c + d*x)))/(a + 2*Sqrt[-a]*Sqrt[b] - b))]/(4*Sqrt[-a^2]*Sqrt[b]*d^2)`

**Rubi [A] (verified)**

Time = 1.03 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6166, 3042, 3801, 2694, 27, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$$

$$\downarrow \text{6166}$$

$$2 \int \frac{x}{a-b+(a+b) \cosh(2c+2dx)} dx$$

$$\downarrow \text{3042}$$

$$2 \int \frac{x}{a-b+(a+b) \sin(2ic+2idx+\frac{\pi}{2})} dx$$

$$\begin{aligned}
& \downarrow \text{3801} \\
& 4 \int \frac{e^{2c+2dx} x}{a + (a+b)e^{4(c+dx)} + 2(a-b)e^{2c+2dx} + b} dx \\
& \downarrow \text{2694} \\
& 4 \left( \frac{(a+b) \int \frac{e^{2c+2dx} x}{2(a+(a+b)e^{2c+2dx}-b-2\sqrt{-a}\sqrt{b})} dx}{2\sqrt{-a}\sqrt{b}} - \frac{(a+b) \int \frac{e^{2c+2dx} x}{2(a+(a+b)e^{2c+2dx}-b+2\sqrt{-a}\sqrt{b})} dx}{2\sqrt{-a}\sqrt{b}} \right) \\
& \downarrow \text{27} \\
& 4 \left( \frac{(a+b) \int \frac{e^{2c+2dx} x}{a+(a+b)e^{2c+2dx}-b-2\sqrt{-a}\sqrt{b}} dx}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \int \frac{e^{2c+2dx} x}{a+(a+b)e^{2c+2dx}-b+2\sqrt{-a}\sqrt{b}} dx}{4\sqrt{-a}\sqrt{b}} \right) \\
& \downarrow \text{2620} \\
& 4 \left( \frac{(a+b) \left( \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int \log\left(\frac{e^{2c+2dx}(a+b)}{a-b-2\sqrt{-a}\sqrt{b}}+1\right) dx}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \left( \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int \log\left(\frac{e^{2c+2dx}(a+b)}{a-b+2\sqrt{-a}\sqrt{b}}+1\right) dx}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) \\
& \downarrow \text{2715} \\
& 4 \left( \frac{(a+b) \left( \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int e^{-2c-2dx} \log\left(\frac{e^{2c+2dx}(a+b)}{a-b-2\sqrt{-a}\sqrt{b}}+1\right) de^{2c+2dx}}{4d^2(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \left( \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int e^{-2c-2dx} \log\left(\frac{e^{2c+2dx}(a+b)}{a-b+2\sqrt{-a}\sqrt{b}}+1\right) de^{2c+2dx}}{4d^2(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) \\
& \downarrow \text{2838} \\
& 4 \left( \frac{(a+b) \left( \frac{\text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{4d^2(a+b)} + \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \left( \frac{\text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b+2\sqrt{-a}\sqrt{b}}\right)}{4d^2(a+b)} + \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right)
\end{aligned}$$

input `Int[(x*Sech[c + d*x]^2)/(a + b*Tanh[c + d*x]^2),x]`

output `4*(((a + b)*((x*Log[1 + ((a + b)*E^(2*c + 2*d*x)]/(a - 2*Sqrt[-a]*Sqrt[b] - b)))/(2*(a + b)*d) + PolyLog[2, -(((a + b)*E^(2*c + 2*d*x)]/(a - 2*Sqrt[-a]*Sqrt[b] - b)))/(4*(a + b)*d^2)))/(4*Sqrt[-a]*Sqrt[b]) - ((a + b)*((x*Log[1 + ((a + b)*E^(2*c + 2*d*x)]/(a + 2*Sqrt[-a]*Sqrt[b] - b)))/(2*(a + b)*d) + PolyLog[2, -(((a + b)*E^(2*c + 2*d*x)]/(a + 2*Sqrt[-a]*Sqrt[b] - b)))/(4*(a + b)*d^2)))/(4*Sqrt[-a]*Sqrt[b]))`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2694 `Int[((F_)^(u_)*((f_) + (g_)*(x_))^(m_))/((a_) + (b_)*(F_)^(u_) + (c_)*((F_)^(v_))), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3801 `Int[((c_) + (d_)*(x_)^(m_))/((a_) + (b_)*sin[(e_) + Pi*(k_) + (Complex[0, fz_]*(f_)*(x_)])], x_Symbol] := Simp[2 Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]`

rule 6166 `Int[((f_) + (g_)*(x_)^(m_))*Sech[(d_) + (e_)*(x_)]^2/((b_) + (c_)*Tanh[(d_) + (e_)*(x_)]^2), x_Symbol] := Simp[2 Int[(f + g*x)^m/(b - c + (b + c)*Cosh[2*d + 2*e*x]), x], x] /; FreeQ[{b, c, d, e, f, g}, x] && IGtQ[m, 0]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 952 vs.  $2(187) = 374$ .

Time = 6.42 (sec) , antiderivative size = 953, normalized size of antiderivative = 4.13

method	result
risch	$-\frac{acx}{d\sqrt{-ab}(-2\sqrt{-ab}-a+b)} + \frac{bcx}{d\sqrt{-ab}(-2\sqrt{-ab}-a+b)} + \frac{\ln\left(1 - \frac{(a+b)e^{2dx+2c}}{-2\sqrt{-ab}-a+b}\right)ac}{2d^2\sqrt{-ab}(-2\sqrt{-ab}-a+b)} - \frac{\ln\left(1 - \frac{(a+b)e^{2dx+2c}}{-2\sqrt{-ab}-a+b}\right)bc}{2d^2\sqrt{-ab}(-2\sqrt{-ab}-a+b)} - \frac{2c}{-2\sqrt{-ab}}$

input `int(x*sech(d*x+c)^2/(a+tanh(d*x+c)^2*b),x,method=_RETURNVERBOSE)`



output

```

-1/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*c*x+1/d/(-a*b)^(1/2)/(-2*(-a*b)^(
(1/2)-a+b)*b*c*x+1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp
(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a*c-1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/
2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*b*c-1/(-2*(-a*b)^(
(1/2)-a+b)*x^2-1/2/(-a*b)^(1/2)*x^2+1/2/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+
b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a*x-1/2/d/(-a*b)^(1/2)
/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*b*
x-1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*c^2+1/2/d^2/(-a*b)^(1/2)/(-
2*(-a*b)^(1/2)-a+b)*b*c^2+1/4/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*polyl
og(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a-1/4/d^2/(-a*b)^(1/2)/(-
2*(-a*b)^(1/2)-a+b)*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*
b-1/2/d^2/(-a*b)^(1/2)*c^2+1/4/d^2/(-a*b)^(1/2)*polylog(2,(a+b)*exp(2*d*x+
2*c)/(2*(-a*b)^(1/2)-a+b))-1/d^2/(-2*(-a*b)^(1/2)-a+b)*c^2+1/2/d^2/(-2*(-a
*b)^(1/2)-a+b)*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))-1/2/(
-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*x^2+1/2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-
a+b)*b*x^2+1/d^2/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b
)^(1/2)-a+b))*c+1/d/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-
a*b)^(1/2)-a+b))*x-2/d/(-2*(-a*b)^(1/2)-a+b)*c*x+1/2/d^2/(-a*b)^(1/2)*ln(1
-(a+b)*exp(2*d*x+2*c)/(2*(-a*b)^(1/2)-a+b))*c-1/d^2*c/(a*b)^(1/2)*arctan(1
/4*(2*(a+b)*exp(2*d*x+2*c)+2*a-2*b)/(a*b)^(1/2))+1/2/d/(-a*b)^(1/2)*ln(...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1516 vs.  $2(185) = 370$ .

Time = 0.20 (sec) , antiderivative size = 1516, normalized size of antiderivative = 6.56

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="fricas")
```

output

```

-1/2*((a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(2*sqrt(-(2*(a + b)*sqrt
(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*
x + c)) + (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(-2*sqrt(-(2*(a + b)
*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*si
nh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(2*sqrt((2*(a +
b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + 2*cosh(d*x + c) + 2
*sinh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(-2*sqrt((2*
(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + 2*cosh(d*x + c)
+ 2*sinh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog(-(((a -
b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a
+ b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(
-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + a + b)/(a + b) + 1) - (a + b
)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog((((a - b)*cosh(d*x + c) + (a - b)*s
inh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b
/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a
- b)/(a + b)) - a - b)/(a + b) + 1) + (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^
2))*dilog(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*co
sh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(
(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + a + b)/(a +
b) + 1) + (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog((((a - b)*cosh(d...

```

### Sympy [F]

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx$$

input

```
integrate(x*sech(d*x+c)**2/(a+b*tanh(d*x+c)**2),x)
```

output

```
Integral(x*sech(c + d*x)**2/(a + b*tanh(c + d*x)**2), x)
```

**Maxima [F]**

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x \operatorname{sech}(dx + c)^2}{b \tanh(dx + c)^2 + a} dx$$

input `integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="maxima")`

output `integrate(x*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)`

**Giac [F]**

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x \operatorname{sech}(dx + c)^2}{b \tanh(dx + c)^2 + a} dx$$

input `integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="giac")`

output `integrate(x*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x}{\cosh(c + dx)^2 (b \tanh(c + dx)^2 + a)} dx$$

input `int(x/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)),x)`

output `int(x/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)), x)`

**Reduce [F]**

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{\operatorname{sech}(dx + c)^2 x}{\tanh(dx + c)^2 b + a} dx$$

input `int(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x)`

output `int((sech(c + d*x)**2*x)/(tanh(c + d*x)**2*b + a),x)`

**3.146**  $\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$

Optimal result	1132
Mathematica [A] (verified)	1133
Rubi [A] (verified)	1133
Maple [B] (verified)	1137
Fricas [B] (verification not implemented)	1138
Sympy [F]	1139
Maxima [F]	1140
Giac [F]	1140
Mupad [F(-1)]	1140
Reduce [F]	1141

**Optimal result**

Integrand size = 26, antiderivative size = 351

$$\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx = \frac{x^2 \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd}} - \frac{x^2 \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd}}$$

$$+ \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd^2}}$$

$$- \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{2\sqrt{-a}\sqrt{bd^2}}$$

$$- \frac{\operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a}\sqrt{bd^3}}$$

$$+ \frac{\operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a}\sqrt{bd^3}}$$

output

$$\frac{1/2*x^2*\ln(1+(a+b)*\exp(2*d*x+2*c)/(a-2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d-1/2*x^2*\ln(1+(a+b)*\exp(2*d*x+2*c)/(a+2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d+1/2*x*\text{polylog}(2,-(a+b)*\exp(2*d*x+2*c)/(a-2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d^2-1/2*x*\text{polylog}(2,-(a+b)*\exp(2*d*x+2*c)/(a+2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d^2-1/4*\text{polylog}(3,-(a+b)*\exp(2*d*x+2*c)/(a-2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d^3+1/4*\text{polylog}(3,-(a+b)*\exp(2*d*x+2*c)/(a+2*(-a)^{(1/2)*b^{(1/2)}-b)))/(-a)^{(1/2)}/b^{(1/2)}/d^3}$$

**Mathematica [A] (verified)**

Time = 0.70 (sec) , antiderivative size = 268, normalized size of antiderivative = 0.76

$$\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$$

$$= \frac{2d^2 x^2 \log\left(1 + \frac{(a+b)e^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b-b}}\right) - 2d^2 x^2 \log\left(1 + \frac{(a+b)e^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b-b}}\right) + 2dx \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b-b}}\right) - 2dx \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b-b}}\right)}{4\sqrt{-a}\sqrt{bd^3}}$$

input

Integrate[(x^2\*Sech[c + d\*x]^2)/(a + b\*Tanh[c + d\*x]^2),x]

output

$$\frac{(2*d^2*x^2*\text{Log}[1 + ((a + b)*E^(2*(c + d*x)))/(a - 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b)] - 2*d^2*x^2*\text{Log}[1 + ((a + b)*E^(2*(c + d*x)))/(a + 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b)] + 2*d*x*\text{PolyLog}[2, -(((a + b)*E^(2*(c + d*x)))/(a - 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b))] - 2*d*x*\text{PolyLog}[2, -(((a + b)*E^(2*(c + d*x)))/(a + 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b))] - \text{PolyLog}[3, -(((a + b)*E^(2*(c + d*x)))/(a - 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b))] + \text{PolyLog}[3, -(((a + b)*E^(2*(c + d*x)))/(a + 2*\text{Sqrt}[-a]*\text{Sqrt}[b] - b))])/(4*\text{Sqrt}[-a]*\text{Sqrt}[b]*d^3)}$$

**Rubi [A] (verified)**Time = 1.52 (sec) , antiderivative size = 353, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$ , Rules used = {6166, 3042, 3801, 2694, 27, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx \\
& \quad \downarrow \text{6166} \\
& 2 \int \frac{x^2}{a-b+(a+b) \cosh(2c+2dx)} dx \\
& \quad \downarrow \text{3042} \\
& 2 \int \frac{x^2}{a-b+(a+b) \sin(2ic+2idx+\frac{\pi}{2})} dx \\
& \quad \downarrow \text{3801} \\
& 4 \int \frac{e^{2c+2dx} x^2}{a+(a+b)e^{4(c+dx)}+2(a-b)e^{2c+2dx}+b} dx \\
& \quad \downarrow \text{2694} \\
& 4 \left( \frac{(a+b) \int \frac{e^{2c+2dx} x^2}{2(a+(a+b)e^{2c+2dx}-b-2\sqrt{-a}\sqrt{b})} dx}{2\sqrt{-a}\sqrt{b}} - \frac{(a+b) \int \frac{e^{2c+2dx} x^2}{2(a+(a+b)e^{2c+2dx}-b+2\sqrt{-a}\sqrt{b})} dx}{2\sqrt{-a}\sqrt{b}} \right) \\
& \quad \downarrow \text{27} \\
& 4 \left( \frac{(a+b) \int \frac{e^{2c+2dx} x^2}{a+(a+b)e^{2c+2dx}-b-2\sqrt{-a}\sqrt{b}} dx}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \int \frac{e^{2c+2dx} x^2}{a+(a+b)e^{2c+2dx}-b+2\sqrt{-a}\sqrt{b}} dx}{4\sqrt{-a}\sqrt{b}} \right) \\
& \quad \downarrow \text{2620} \\
& 4 \left( \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int x \log\left(\frac{e^{2c+2dx}(a+b)}{a-b-2\sqrt{-a}\sqrt{b}}+1\right) dx}{d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} - \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b}+1\right)}{2d(a+b)} - \frac{\int x \log\left(\frac{e^{2c+2dx}}{a-b+2\sqrt{-a}\sqrt{b}}+1\right) dx}{d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) \\
& \quad \downarrow \text{3011}
\end{aligned}$$

$$4 \left( \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b+a-b}}+1\right)}{2d(a+b)} - \frac{\int \text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right) dx}{2d} - x \text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) - \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c}}{2\sqrt{-a}\sqrt{b}}\right)}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}}$$

↓ 2720

$$4 \left( \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b+a-b}}+1\right)}{2d(a+b)} - \frac{\int e^{-2c-2dx} \text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right) de^{2c+2dx}}{4d^2} - x \text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) - \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c}}{2\sqrt{-a}\sqrt{b}}\right)}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}}$$

↓ 7143

$$4 \left( \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b+a-b}}+1\right)}{2d(a+b)} - \frac{\text{PolyLog}\left(3, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{4d^2} - \frac{x \text{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}} \right) - \frac{(a+b) \left( \frac{x^2 \log\left(\frac{(a+b)e^{2c}}{2\sqrt{-a}\sqrt{b}}\right)}{2d(a+b)} \right)}{4\sqrt{-a}\sqrt{b}}$$

input `Int[(x^2*Sech[c + d*x]^2)/(a + b*Tanh[c + d*x]^2),x]`

output `4*(((a + b)*((x^2*Log[1 + ((a + b)*E^(2*c + 2*d*x))]/(a - 2*Sqrt[-a]*Sqrt[b] - b)))/(2*(a + b)*d) - (-1/2*(x*PolyLog[2, -(((a + b)*E^(2*c + 2*d*x))/(a - 2*Sqrt[-a]*Sqrt[b] - b))])/d + PolyLog[3, -(((a + b)*E^(2*c + 2*d*x))/(a - 2*Sqrt[-a]*Sqrt[b] - b))]/(4*d^2)))/((a + b)*d))/(4*Sqrt[-a]*Sqrt[b]) - (((a + b)*((x^2*Log[1 + ((a + b)*E^(2*c + 2*d*x))]/(a + 2*Sqrt[-a]*Sqrt[b] - b)))/(2*(a + b)*d) - (-1/2*(x*PolyLog[2, -(((a + b)*E^(2*c + 2*d*x))/(a + 2*Sqrt[-a]*Sqrt[b] - b))])/d + PolyLog[3, -(((a + b)*E^(2*c + 2*d*x))/(a + 2*Sqrt[-a]*Sqrt[b] - b))]/(4*d^2)))/((a + b)*d))/(4*Sqrt[-a]*Sqrt[b])`



## Definitions of rubi rules used

- rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_) /; FreeQ[b, x]]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2694 `Int[((F_)^(u_)*((f_) + (g_)*(x_))^(m_))/((a_) + (b_)*(F_)^(u_) + (c_)*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Simp[2*(c/q) Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3801

```
Int[((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Simp[2 Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]
```

rule 6166

```
Int[(((f_.) + (g_.)*(x_)^(m_.)*Sech[(d_.) + (e_.)*(x_)]^2)/((b_) + (c_.)*Tanh[(d_.) + (e_.)*(x_)]^2), x_Symbol] := Simp[2 Int[(f + g*x)^m/(b - c + (b + c)*Cosh[2*d + 2*e*x]), x], x] /; FreeQ[{b, c, d, e, f, g}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.))/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1185 vs.  $2(285) = 570$ .

Time = 6.41 (sec) , antiderivative size = 1186, normalized size of antiderivative = 3.38

method	result	size
risch	Expression too large to display	1186

input

```
int(x^2*sech(d*x+c)^2/(a+tanh(d*x+c)^2*b),x,method=_RETURNVERBOSE)
```

output

```

2/3/d^3/(-a*b)^(1/2)*c^3-1/4/d^3/(-a*b)^(1/2)*polylog(3,(a+b)*exp(2*d*x+2*
c)/(2*(-a*b)^(1/2)-a+b))-1/2/d^3/(-2*(-a*b)^(1/2)-a+b)*polylog(3,(a+b)*exp
(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))+4/3/d^3/(-2*(-a*b)^(1/2)-a+b)*c^3+1/2/d
/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)
^(1/2)-a+b))*x^2+1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*polylog(2,(a
+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*x-1/2/d/(-a*b)^(1/2)/(-2*(-a*b)
^(1/2)-a+b)*b*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*x^2-1/2/d^2/
(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-
a*b)^(1/2)-a+b))*x-1/2/d^3/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*ln(1-(a+b)
*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*c^2+1/2/d^3/(-a*b)^(1/2)/(-2*(-a*b)
^(1/2)-a+b)*b*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*c^2-1/d^2/(
-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*c^2*x+1/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1
/2)-a+b)*a*c^2*x-2/3/d^3/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*c^3+2/3/d^3/
(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*c^3-1/4/d^3/(-a*b)^(1/2)/(-2*(-a*b)^(
1/2)-a+b)*a*polylog(3,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))+1/4/d^3/
(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*polylog(3,(a+b)*exp(2*d*x+2*c)/(-2*(-
a*b)^(1/2)-a+b))+1/3/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*x^3-1/3/(-a*b)^(
1/2)/(-2*(-a*b)^(1/2)-a+b)*a*x^3+1/d^3*c^2/(a*b)^(1/2)*arctan(1/4*(2*(a+b)
*exp(2*d*x+2*c)+2*a-2*b)/(a*b)^(1/2))-1/d^3/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+
b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*c^2+1/d^2/(-a*b)^(1/2)*c^2*x+1...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2110 vs.  $2(283) = 566$ .

Time = 0.17 (sec) , antiderivative size = 2110, normalized size of antiderivative = 6.01

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="fricas")
```

output

```

1/2*(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*d*x*dilog(-((a - b)*cosh(d*
x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d
*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 +
2*a*b + b^2)) + a - b)/(a + b)) + a + b)/(a + b) + 1) + 2*(a + b)*sqrt(-a
*b/(a^2 + 2*a*b + b^2))*d*x*dilog(((a - b)*cosh(d*x + c) + (a - b)*sinh(d
*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2
+ 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)
/(a + b)) - a - b)/(a + b) + 1) - 2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))
*d*x*dilog(-((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*c
osh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt
((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + a + b)/(a +
b) + 1) - 2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*d*x*dilog((((a - b)*co
sh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cosh(d*x + c) + (a + b)*s
inh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*(a + b)*sqrt(-a*b/(a
^2 + 2*a*b + b^2)) - a + b)/(a + b)) - a - b)/(a + b) + 1) + (a + b)*sqrt(
-a*b/(a^2 + 2*a*b + b^2))*c^2*log(2*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*
b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) + (a + b)
*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c^2*log(-2*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2
+ 2*a*b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) -
(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c^2*log(2*sqrt((2*(a + b)*sqrt(-...

```

## Sympy [F]

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx$$

input

```
integrate(x**2*sech(d*x+c)**2/(a+b*tanh(d*x+c)**2),x)
```

output

```
Integral(x**2*sech(c + d*x)**2/(a + b*tanh(c + d*x)**2), x)
```

**Maxima [F]**

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x^2 \operatorname{sech}(dx + c)^2}{b \tanh(dx + c)^2 + a} dx$$

input `integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="maxima")`

output `integrate(x^2*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)`

**Giac [F]**

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x^2 \operatorname{sech}(dx + c)^2}{b \tanh(dx + c)^2 + a} dx$$

input `integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="giac")`

output `integrate(x^2*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{x^2}{\cosh(c + dx)^2 (b \tanh(c + dx)^2 + a)} dx$$

input `int(x^2/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)),x)`

output `int(x^2/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)), x)`

**Reduce [F]**

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx = \int \frac{\operatorname{sech}(dx + c)^2 x^2}{\tanh(dx + c)^2 b + a} dx$$

input `int(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x)`

output `int((sech(c + d*x)**2*x**2)/(tanh(c + d*x)**2*b + a),x)`

### 3.147 $\int x^5 \tanh(a + 2 \log(x)) dx$

Optimal result	1142
Mathematica [A] (verified)	1142
Rubi [A] (verified)	1143
Maple [C] (verified)	1144
Fricas [A] (verification not implemented)	1145
Sympy [F]	1145
Maxima [A] (verification not implemented)	1146
Giac [A] (verification not implemented)	1146
Mupad [B] (verification not implemented)	1146
Reduce [B] (verification not implemented)	1147

#### Optimal result

Integrand size = 11, antiderivative size = 32

$$\int x^5 \tanh(a + 2 \log(x)) dx = -e^{-2a} x^2 + \frac{x^6}{6} + e^{-3a} \arctan(e^a x^2)$$

output `-x^2/exp(2*a)+1/6*x^6+arctan(exp(a)*x^2)/exp(3*a)`

#### Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.75

$$\int x^5 \tanh(a + 2 \log(x)) dx = \frac{x^6}{6} - x^2 \cosh(2a) + \arctan(x^2(\cosh(a) + \sinh(a))) \cosh(3a) + x^2 \sinh(2a) - \arctan(x^2(\cosh(a) + \sinh(a))) \sinh(3a)$$

input `Integrate[x^5*Tanh[a + 2*Log[x]],x]`

output `x^6/6 - x^2*Cosh[2*a] + ArcTan[x^2*(Cosh[a] + Sinh[a])]*Cosh[3*a] + x^2*Sinh[2*a] - ArcTan[x^2*(Cosh[a] + Sinh[a])]*Sinh[3*a]`

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {6071, 959, 807, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \tanh(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{x^5 (e^{2a} x^4 - 1)}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{x^6}{6} - 2 \int \frac{x^5}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{x^6}{6} - \int \frac{x^4}{e^{2a} x^4 + 1} dx^2 \\
 & \quad \downarrow \text{262} \\
 & e^{-2a} \int \frac{1}{e^{2a} x^4 + 1} dx^2 - e^{-2a} x^2 + \frac{x^6}{6} \\
 & \quad \downarrow \text{216} \\
 & e^{-3a} \arctan(e^a x^2) - e^{-2a} x^2 + \frac{x^6}{6}
 \end{aligned}$$

input `Int[x^5*Tanh[a + 2*Log[x]],x]`

output `-(x^2/E^(2*a)) + x^6/6 + ArcTan[E^a*x^2]/E^(3*a)`



## Definitions of rubi rules used

rule 216  $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*ArcTan[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /;$  FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

rule 262  $\text{Int}[(c_)*(x_))^{(m_)}*((a_ + (b_)*(x_)^2)^{(p_)}], x\_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a + b*x^2)^{(p+1)}/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \text{Int}[(c*x)^{(m-2)}*(a + b*x^2)^p, x], x] /;$  FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2\*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 807  $\text{Int}[(x_)^{(m_)}*((a_ + (b_)*(x_)^n)^{(p_)}], x\_Symbol] \rightarrow \text{With}[\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \text{Subst}[\text{Int}[x^{(m+1)/k - 1}*(a + b*x^{(n/k)})^p, x], x, x^k], x] /;$  k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

rule 959  $\text{Int}[(e_)*(x_))^{(m_)}*((a_ + (b_)*(x_)^n)^{(p_)}*((c_ + (d_)*(x_)^n)), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /;$  FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b\*c - a\*d, 0] && NeQ[m + n\*(p+1) + 1, 0]

rule 6071  $\text{Int}[(e_)*(x_))^{(m_)}*\text{Tanh}[(a_ + \text{Log}[x_]*(b_))*(d_)]^{(p_)}], x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)})^p/(1 + E^{(2*a*d)*x^{(2*b*d)})^p}), x] /;$  FreeQ[{a, b, d, e, m, p}, x]

## Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.56

method	result	size
risch	$\frac{x^6}{6} - e^{-2a}x^2 + \frac{ie^{-3a} \ln(e^a x^2 + i)}{2} - \frac{ie^{-3a} \ln(e^a x^2 - i)}{2}$	50

input `int(x^5*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)`

output `1/6*x^6-exp(-2*a)*x^2+1/2*I*exp(-3*a)*ln(exp(a)*x^2+I)-1/2*I*exp(-3*a)*ln(exp(a)*x^2-I)`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

$$\int x^5 \tanh(a + 2 \log(x)) dx = \frac{1}{6} (x^6 e^{(3a)} - 6 x^2 e^a + 6 \arctan(x^2 e^a)) e^{(-3a)}$$

input `integrate(x^5*tanh(a+2*log(x)),x, algorithm="fricas")`

output `1/6*(x^6*e^(3*a) - 6*x^2*e^a + 6*arctan(x^2*e^a))*e^(-3*a)`

### Sympy [F]

$$\int x^5 \tanh(a + 2 \log(x)) dx = \int x^5 \tanh(a + 2 \log(x)) dx$$

input `integrate(x**5*tanh(a+2*ln(x)),x)`

output `Integral(x**5*tanh(a + 2*log(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

$$\int x^5 \tanh(a + 2 \log(x)) dx = \frac{1}{6} (x^6 e^{(2a)} - 6x^2) e^{(-2a)} + \arctan(x^2 e^a) e^{(-3a)}$$

input `integrate(x^5*tanh(a+2*log(x)),x, algorithm="maxima")`output `1/6*(x^6*e^(2*a) - 6*x^2)*e^(-2*a) + arctan(x^2*e^a)*e^(-3*a)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16

$$\int x^5 \tanh(a + 2 \log(x)) dx = \arctan(x^2 e^a) e^{(-3a)} + \frac{1}{6} (x^6 e^{(6a)} - 6x^2 e^{(4a)}) e^{(-6a)}$$

input `integrate(x^5*tanh(a+2*log(x)),x, algorithm="giac")`output `arctan(x^2*e^a)*e^(-3*a) + 1/6*(x^6*e^(6*a) - 6*x^2*e^(4*a))*e^(-6*a)`**Mupad [B] (verification not implemented)**

Time = 2.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.03

$$\int x^5 \tanh(a + 2 \log(x)) dx = \frac{\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{(e^{2a})^{3/2}} - x^2 e^{-2a} + \frac{x^6}{6}$$

input `int(x^5*tanh(a + 2*log(x)),x)`output `atan(x^2*exp(2*a)^(1/2))/exp(2*a)^(3/2) - x^2*exp(-2*a) + x^6/6`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.72

$$\int x^5 \tanh(a + 2 \log(x)) dx = \frac{-6 \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) - 6 \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) + e^{3a}x^6 - 6e^ax^2}{6e^{3a}}$$

input `int(x^5*tanh(a+2*log(x)),x)`output `( - 6*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) - 6*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) + e**(3*a)*x**6 - 6*e**a*x**2)/(6*e**(3*a))`

### 3.148 $\int x^3 \tanh(a + 2 \log(x)) dx$

Optimal result	1148
Mathematica [B] (verified)	1148
Rubi [A] (verified)	1149
Maple [A] (verified)	1150
Fricas [A] (verification not implemented)	1151
Sympy [F]	1151
Maxima [A] (verification not implemented)	1151
Giac [A] (verification not implemented)	1152
Mupad [B] (verification not implemented)	1152
Reduce [B] (verification not implemented)	1152

#### Optimal result

Integrand size = 11, antiderivative size = 29

$$\int x^3 \tanh(a + 2 \log(x)) dx = \frac{x^4}{4} - \frac{1}{2} e^{-2a} \log(1 + e^{2a} x^4)$$

output

```
1/4*x^4-1/2*ln(1+exp(2*a)*x^4)/exp(2*a)
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 64 vs. 2(29) = 58.

Time = 0.02 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.21

$$\begin{aligned} \int x^3 \tanh(a + 2 \log(x)) dx = & \frac{x^4}{4} - \frac{1}{2} \cosh(2a) \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) \\ & + x^4 \sinh(a)) + \frac{1}{2} \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) \\ & + x^4 \sinh(a)) \sinh(2a) \end{aligned}$$

input

```
Integrate[x^3*Tanh[a + 2*Log[x]],x]
```

output

$$x^4/4 - (\text{Cosh}[2*a]*\text{Log}[\text{Cosh}[a] + x^4*\text{Cosh}[a] - \text{Sinh}[a] + x^4*\text{Sinh}[a]])/2 + (\text{Log}[\text{Cosh}[a] + x^4*\text{Cosh}[a] - \text{Sinh}[a] + x^4*\text{Sinh}[a]]*\text{Sinh}[2*a])/2$$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {6071, 946, 25, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \tanh(a + 2 \log(x)) dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{x^3 (e^{2a} x^4 - 1)}{e^{2a} x^4 + 1} dx \\ & \quad \downarrow \text{946} \\ & \frac{1}{4} \int -\frac{1 - e^{2a} x^4}{e^{2a} x^4 + 1} dx^4 \\ & \quad \downarrow \text{25} \\ & -\frac{1}{4} \int \frac{1 - e^{2a} x^4}{e^{2a} x^4 + 1} dx^4 \\ & \quad \downarrow \text{49} \\ & -\frac{1}{4} \int \left( \frac{2}{e^{2a} x^4 + 1} - 1 \right) dx^4 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} (x^4 - 2e^{-2a} \log(e^{2a} x^4 + 1)) \end{aligned}$$

input

$$\text{Int}[x^3*\text{Tanh}[a + 2*\text{Log}[x]],x]$$

output

$$(x^4 - (2*\text{Log}[1 + E^(2*a)*x^4])/E^(2*a))/4$$

## Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6071 `Int[((e_.)*(x_)^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p], x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
risch	$\frac{x^4}{4} - \frac{e^{-2a} \ln(1+e^{2a}x^4)}{2}$	24

input `int(x^3*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)`

output `1/4*x^4-1/2*exp(-2*a)*ln(1+exp(2*a)*x^4)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int x^3 \tanh(a + 2 \log(x)) dx = \frac{1}{4} (x^4 e^{(2a)} - 2 \log(x^4 e^{(2a)} + 1)) e^{(-2a)}$$

input `integrate(x^3*tanh(a+2*log(x)),x, algorithm="fricas")`

output `1/4*(x^4*e^(2*a) - 2*log(x^4*e^(2*a) + 1))*e^(-2*a)`

**Sympy [F]**

$$\int x^3 \tanh(a + 2 \log(x)) dx = \int x^3 \tanh(a + 2 \log(x)) dx$$

input `integrate(x**3*tanh(a+2*ln(x)),x)`

output `Integral(x**3*tanh(a + 2*log(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int x^3 \tanh(a + 2 \log(x)) dx = \frac{1}{4} x^4 - \frac{1}{2} e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

input `integrate(x^3*tanh(a+2*log(x)),x, algorithm="maxima")`

output `1/4*x^4 - 1/2*e^(-2*a)*log(x^4*e^(2*a) + 1)`



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int x^3 \tanh(a + 2 \log(x)) dx = \frac{1}{4} x^4 - \frac{1}{2} e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

input `integrate(x^3*tanh(a+2*log(x)),x, algorithm="giac")`

output `1/4*x^4 - 1/2*e^(-2*a)*log(x^4*e^(2*a) + 1)`

**Mupad [B] (verification not implemented)**

Time = 2.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int x^3 \tanh(a + 2 \log(x)) dx = \frac{x^4}{4} - \frac{e^{-2a} \ln(x^4 + e^{-2a})}{2}$$

input `int(x^3*tanh(a + 2*log(x)),x)`

output `x^4/4 - (exp(-2*a)*log(exp(-2*a) + x^4))/2`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.14

$$\begin{aligned} & \int x^3 \tanh(a + 2 \log(x)) dx \\ &= \frac{e^{2a} x^4 - 2 \log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) - 2 \log(e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1)}{4e^{2a}} \end{aligned}$$

input `int(x^3*tanh(a+2*log(x)),x)`

output `(e**(2*a)*x**4 - 2*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) - 2*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1))/(4*e**(2*a))`

### 3.149 $\int x \tanh(a + 2 \log(x)) dx$

Optimal result	1153
Mathematica [A] (verified)	1153
Rubi [A] (verified)	1154
Maple [C] (verified)	1155
Fricas [A] (verification not implemented)	1156
Sympy [F]	1156
Maxima [A] (verification not implemented)	1156
Giac [A] (verification not implemented)	1157
Mupad [B] (verification not implemented)	1157
Reduce [B] (verification not implemented)	1157

#### Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x \tanh(a + 2 \log(x)) dx = \frac{x^2}{2} - e^{-a} \arctan(e^a x^2)$$

output `1/2*x^2-arctan(exp(a)*x^2)/exp(a)`

#### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.52

$$\int x \tanh(a + 2 \log(x)) dx = \frac{x^2}{2} - \arctan(x^2(\cosh(a) + \sinh(a))) \cosh(a) + \arctan(x^2(\cosh(a) + \sinh(a))) \sinh(a)$$

input `Integrate[x*Tanh[a + 2*Log[x]],x]`

output `x^2/2 - ArcTan[x^2*(Cosh[a] + Sinh[a])]*Cosh[a] + ArcTan[x^2*(Cosh[a] + Sinh[a])]*Sinh[a]`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {6071, 959, 807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tanh(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{x(e^{2a}x^4 - 1)}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{x^2}{2} - 2 \int \frac{x}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{x^2}{2} - \int \frac{1}{e^{2a}x^4 + 1} dx^2 \\
 & \quad \downarrow \text{216} \\
 & \frac{x^2}{2} - e^{-a} \arctan(e^a x^2)
 \end{aligned}$$

input `Int[x*Tanh[a + 2*Log[x]],x]`

output `x^2/2 - ArcTan[E^a*x^2]/E^a`

## Definitions of rubi rules used

rule 216

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 807

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

rule 959

```
Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

rule 6071

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]
```

## Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.78

method	result	size
risch	$\frac{x^2}{2} + \frac{ie^{-a} \ln(e^a x^2 - i)}{2} - \frac{ie^{-a} \ln(e^a x^2 + i)}{2}$	41

input

```
int(x*tanh(a+2*ln(x)), x, method=_RETURNVERBOSE)
```

output

```
1/2*x^2+1/2*I/exp(a)*ln(exp(a)*x^2-I)-1/2*I/exp(a)*ln(exp(a)*x^2+I)
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int x \tanh(a + 2 \log(x)) dx = \frac{1}{2} (x^2 e^a - 2 \arctan(x^2 e^a)) e^{-a}$$

input `integrate(x*tanh(a+2*log(x)),x, algorithm="fricas")`output `1/2*(x^2*e^a - 2*arctan(x^2*e^a))*e^(-a)`**Sympy [F]**

$$\int x \tanh(a + 2 \log(x)) dx = \int x \tanh(a + 2 \log(x)) dx$$

input `integrate(x*tanh(a+2*ln(x)),x)`output `Integral(x*tanh(a + 2*log(x)), x)`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \tanh(a + 2 \log(x)) dx = \frac{1}{2} x^2 - \arctan(x^2 e^a) e^{-a}$$

input `integrate(x*tanh(a+2*log(x)),x, algorithm="maxima")`output `1/2*x^2 - arctan(x^2*e^a)*e^(-a)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \tanh(a + 2 \log(x)) dx = \frac{1}{2} x^2 - \arctan(x^2 e^a) e^{-a}$$

input `integrate(x*tanh(a+2*log(x)),x, algorithm="giac")`output `1/2*x^2 - arctan(x^2*e^a)*e^(-a)`**Mupad [B] (verification not implemented)**

Time = 2.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x \tanh(a + 2 \log(x)) dx = \frac{x^2}{2} - \frac{\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{\sqrt{e^{2a}}}$$

input `int(x*tanh(a + 2*log(x)),x)`output `x^2/2 - atan(x^2*exp(2*a)^(1/2))/exp(2*a)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 3.26

$$\int x \tanh(a + 2 \log(x)) dx = \frac{2 \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) + 2 \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) + e^a x^2}{2e^a}$$

input `int(x*tanh(a+2*log(x)),x)`output `(2*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) + 2*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) + e**a*x**2)/(2*e**a)`

$$3.150 \quad \int \frac{\tanh(a+2 \log(x))}{x} dx$$

Optimal result	1158
Mathematica [A] (verified)	1158
Rubi [A] (verified)	1159
Maple [A] (verified)	1160
Fricas [A] (verification not implemented)	1161
Sympy [A] (verification not implemented)	1161
Maxima [A] (verification not implemented)	1161
Giac [A] (verification not implemented)	1162
Mupad [B] (verification not implemented)	1162
Reduce [B] (verification not implemented)	1162

### Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

output `1/2*ln(cosh(a+2*ln(x)))`

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

input `Integrate[Tanh[a + 2*Log[x]]/x,x]`

output `Log[Cosh[a + 2*Log[x]]]/2`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {3039, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(a + 2 \log(x))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \tanh(a + 2 \log(x)) d \log(x) \\
 & \quad \downarrow \text{3042} \\
 & \int -i \tan(ia + 2i \log(x)) d \log(x) \\
 & \quad \downarrow \text{26} \\
 & -i \int \tan(ia + 2i \log(x)) d \log(x) \\
 & \quad \downarrow \text{3956} \\
 & \frac{1}{2} \log(\cosh(a + 2 \log(x)))
 \end{aligned}$$

input `Int [Tanh[a + 2*Log[x]]/x,x]`

output `Log[Cosh[a + 2*Log[x]]]/2`



## Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst [[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d *x], x]]/d, x] /; FreeQ[{c, d}, x]`

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$\frac{\ln(\cosh(a+2\ln(x)))}{2}$	11
default	$\frac{\ln(\cosh(a+2\ln(x)))}{2}$	11
risch	$-\ln(x) + \frac{\ln(-e^{2a}x^4-1)}{2}$	20
parallelrisc	$-\ln(x) - \frac{\ln(1-\tanh(a+2\ln(x)))}{2}$	20

input `int(tanh(a+2*ln(x))/x,x,method=_RETURNVERBOSE)`

output `1/2*ln(cosh(a+2*ln(x)))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.50

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{1}{2} \log(x^4 e^{2a} + 1) - \log(x)$$

input `integrate(tanh(a+2*log(x))/x,x, algorithm="fricas")`output `1/2*log(x^4*e^(2*a) + 1) - log(x)`**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \log(x) - \frac{\log(\tanh(a + 2 \log(x)) + 1)}{2}$$

input `integrate(tanh(a+2*ln(x))/x,x)`output `log(x) - log(tanh(a + 2*log(x)) + 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

input `integrate(tanh(a+2*log(x))/x,x, algorithm="maxima")`output `1/2*log(cosh(a + 2*log(x)))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{1}{2} \log(x^4 e^{(2a)} + 1) - \frac{1}{4} \log(x^4)$$

input `integrate(tanh(a+2*log(x))/x,x, algorithm="giac")`

output `1/2*log(x^4*e^(2*a) + 1) - 1/4*log(x^4)`

**Mupad [B] (verification not implemented)**

Time = 2.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \ln(x) - \frac{\ln(\tanh(a + 2 \ln(x)) + 1)}{2}$$

input `int(tanh(a + 2*log(x))/x,x)`

output `log(x) - log(tanh(a + 2*log(x)) + 1)/2`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 4.00

$$\int \frac{\tanh(a + 2 \log(x))}{x} dx = \frac{\log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1)}{2} + \frac{\log(e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1)}{2} - \log(x)$$

input `int(tanh(a+2*log(x))/x,x)`

output `(log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) - 2*log(x))/2`

### 3.151 $\int \frac{\tanh(a+2 \log(x))}{x^3} dx$

Optimal result	1163
Mathematica [A] (verified)	1163
Rubi [A] (verified)	1164
Maple [C] (verified)	1165
Fricas [A] (verification not implemented)	1166
Sympy [F]	1166
Maxima [A] (verification not implemented)	1166
Giac [A] (verification not implemented)	1167
Mupad [B] (verification not implemented)	1167
Reduce [B] (verification not implemented)	1167

#### Optimal result

Integrand size = 11, antiderivative size = 20

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \frac{1}{2x^2} + e^a \arctan(e^a x^2)$$

output `1/2/x^2+exp(a)*arctan(exp(a)*x^2)`

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.00

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \frac{1}{2x^2} - \arctan\left(\frac{\cosh(a) - \sinh(a)}{x^2}\right) \cosh(a) - \arctan\left(\frac{\cosh(a) - \sinh(a)}{x^2}\right) \sinh(a)$$

input `Integrate[Tanh[a + 2*Log[x]]/x^3,x]`

output `1/(2*x^2) - ArcTan[(Cosh[a] - Sinh[a])/x^2]*Cosh[a] - ArcTan[(Cosh[a] - Sinh[a])/x^2]*Sinh[a]`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6071, 955, 807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh(a + 2 \log(x))}{x^3} dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{e^{2a}x^4 - 1}{x^3 (e^{2a}x^4 + 1)} dx \\ & \quad \downarrow \text{955} \\ & 2e^{2a} \int \frac{x}{e^{2a}x^4 + 1} dx + \frac{1}{2x^2} \\ & \quad \downarrow \text{807} \\ & e^{2a} \int \frac{1}{e^{2a}x^4 + 1} dx^2 + \frac{1}{2x^2} \\ & \quad \downarrow \text{216} \\ & e^a \arctan(e^a x^2) + \frac{1}{2x^2} \end{aligned}$$

input `Int[Tanh[a + 2*Log[x]]/x^3,x]`

output `1/(2*x^2) + E^a*ArcTan[E^a*x^2]`

## Definitions of rubi rules used

rule 216

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 807

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

rule 955

```
Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1)), x] + Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*e^n*(m + 1)) Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]
```

rule 6071

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]
```

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.30 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.20

method	result	size
risch	$\frac{1}{2x^2} + \frac{\sum_{R=\text{RootOf}(e^{2a}+z^2)} -R \ln((4e^{2a}+5-R^2)x^2-R)}{2}$	44

input

```
int(tanh(a+2*ln(x))/x^3,x,method=_RETURNVERBOSE)
```

output `1/2/x^2+1/2*sum(_R*ln((4*exp(2*a)+5*_R^2)*x^2-_R),_R=RootOf(exp(2*a)+_Z^2))`

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \frac{2 x^2 \arctan(x^2 e^a) e^a + 1}{2 x^2}$$

input `integrate(tanh(a+2*log(x))/x^3,x, algorithm="fricas")`

output `1/2*(2*x^2*arctan(x^2*e^a)*e^a + 1)/x^2`

### Sympy [F]

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \int \frac{\tanh(a + 2 \log(x))}{x^3} dx$$

input `integrate(tanh(a+2*ln(x))/x**3,x)`

output `Integral(tanh(a + 2*log(x))/x**3, x)`

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = -\arctan\left(\frac{e^{-a}}{x^2}\right) e^a + \frac{1}{2x^2}$$

input `integrate(tanh(a+2*log(x))/x^3,x, algorithm="maxima")`

output `-arctan(e^(-a)/x^2)*e^a + 1/2/x^2`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \arctan(x^2 e^a) e^a + \frac{1}{2x^2}$$

input `integrate(tanh(a+2*log(x))/x^3,x, algorithm="giac")`output `arctan(x^2*e^a)*e^a + 1/2/x^2`**Mupad [B] (verification not implemented)**

Time = 2.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \operatorname{atan}(x^2 \sqrt{e^{2a}}) \sqrt{e^{2a}} + \frac{1}{2x^2}$$

input `int(tanh(a + 2*log(x))/x^3,x)`output `atan(x^2*exp(2*a)^(1/2))*exp(2*a)^(1/2) + 1/(2*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.95

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \frac{-2e^a \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^2 - 2e^a \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^2 + 1}{2x^2}$$

input `int(tanh(a+2*log(x))/x^3,x)`output `( - 2*e**a*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**2 - 2*e**a*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**2 + 1)/(2*x**2)`



### 3.152 $\int \frac{\tanh(a+2 \log(x))}{x^5} dx$

Optimal result	1168
Mathematica [B] (verified)	1168
Rubi [A] (verified)	1169
Maple [A] (verified)	1170
Fricas [A] (verification not implemented)	1171
Sympy [F]	1171
Maxima [A] (verification not implemented)	1171
Giac [A] (verification not implemented)	1172
Mupad [B] (verification not implemented)	1172
Reduce [B] (verification not implemented)	1172

#### Optimal result

Integrand size = 11, antiderivative size = 38

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = \frac{1}{4x^4} + 2e^{2a} \log(x) - \frac{1}{2}e^{2a} \log(1 + e^{2a}x^4)$$

output

$1/4/x^4+2*\exp(2*a)*\ln(x)-1/2*\exp(2*a)*\ln(1+\exp(2*a)*x^4)$

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 80 vs.  $2(38) = 76$ .

Time = 0.02 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.11

$$\begin{aligned} \int \frac{\tanh(a + 2 \log(x))}{x^5} dx &= \frac{1}{4x^4} + 2 \cosh(2a) \log(x) \\ &\quad - \frac{1}{2} \cosh(2a) \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) + x^4 \sinh(a)) \\ &\quad + 2 \log(x) \sinh(2a) \\ &\quad - \frac{1}{2} \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) + x^4 \sinh(a)) \sinh(2a) \end{aligned}$$

input

`Integrate[Tanh[a + 2*Log[x]]/x^5,x]`

output

$$\frac{1}{4x^4} + 2\operatorname{Cosh}[2a]\operatorname{Log}[x] - (\operatorname{Cosh}[2a]\operatorname{Log}[\operatorname{Cosh}[a] + x^4\operatorname{Cosh}[a] - \operatorname{Sinh}[a] + x^4\operatorname{Sinh}[a]])/2 + 2\operatorname{Log}[x]\operatorname{Sinh}[2a] - (\operatorname{Log}[\operatorname{Cosh}[a] + x^4\operatorname{Cosh}[a] - \operatorname{Sinh}[a] + x^4\operatorname{Sinh}[a]]\operatorname{Sinh}[2a])/2$$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {6071, 948, 25, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh(a + 2 \log(x))}{x^5} dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{e^{2a}x^4 - 1}{x^5 (e^{2a}x^4 + 1)} dx \\ & \quad \downarrow \text{948} \\ & \frac{1}{4} \int -\frac{1 - e^{2a}x^4}{x^8 (e^{2a}x^4 + 1)} dx^4 \\ & \quad \downarrow \text{25} \\ & -\frac{1}{4} \int \frac{1 - e^{2a}x^4}{x^8 (e^{2a}x^4 + 1)} dx^4 \\ & \quad \downarrow \text{86} \\ & -\frac{1}{4} \int \left( -\frac{2e^{2a}}{x^4} + \frac{1}{x^8} + \frac{2e^{4a}}{e^{2a}x^4 + 1} \right) dx^4 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \left( 2e^{2a} \log(x^4) - 2e^{2a} \log(e^{2a}x^4 + 1) + \frac{1}{x^4} \right) \end{aligned}$$

input

$$\operatorname{Int}[\operatorname{Tanh}[a + 2\operatorname{Log}[x]]/x^5, x]$$

output  $(x^{-4} + 2e^{2a}\log[x^4] - 2e^{2a}\log[1 + e^{2a}x^4])/4$

### Defintions of rubi rules used

rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 86  $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{(\text{n}_.)}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{(\text{p}_.)}), \text{x}_.] \rightarrow \text{Int}[\text{ExpandIntegrand}[(\text{a} + \text{b} * \text{x}) * (\text{c} + \text{d} * \text{x})^{\text{n}} * (\text{e} + \text{f} * \text{x})^{\text{p}}, \text{x}], \text{x}] /;$   
 $\text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{n}\}, \text{x}] \ \&\& \ ((\text{ILtQ}[\text{n}, 0] \ \&\& \ \text{ILtQ}[\text{p}, 0]) \ || \ \text{EqQ}[\text{p}, 1]) \ || \ (\text{IGtQ}[\text{p}, 0] \ \&\& \ (!\text{IntegerQ}[\text{n}] \ || \ \text{LeQ}[9 * \text{p} + 5 * (\text{n} + 2), 0]) \ || \ \text{GeQ}[\text{n} + \text{p} + 1, 0]) \ || \ (\text{GeQ}[\text{n} + \text{p} + 2, 0] \ \&\& \ \text{RationalQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}])))$

rule 948  $\text{Int}[(\text{x}_.)^{(\text{m}_.)} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{(\text{n}_.)})^{(\text{p}_.)} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{(\text{n}_.)})^{(\text{q}_.)}), \text{x\_Symbol}] \rightarrow \text{Simp}[1/\text{n} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{Simplify}[(\text{m} + 1)/\text{n}] - 1) * (\text{a} + \text{b} * \text{x})^{\text{p}} * (\text{c} + \text{d} * \text{x})^{\text{q}}, \text{x}, \text{x}^{\text{n}}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{p}, \text{q}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(\text{m} + 1)/\text{n}]]$

rule 2009  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] /; \text{SumQ}[\text{u}]$

rule 6071  $\text{Int}[(\text{e}_.) * (\text{x}_.)^{(\text{m}_.)} * \text{Tanh}[(\text{a}_.) + \text{Log}[\text{x}_.] * (\text{b}_.) * (\text{d}_.)]^{(\text{p}_.)}), \text{x\_Symbol}] \rightarrow \text{Int}[(\text{e} * \text{x})^{\text{m}} * (-1 + \text{E}^{(2 * \text{a} * \text{d}) * \text{x}^{(2 * \text{b} * \text{d})}})^{\text{p}} / (1 + \text{E}^{(2 * \text{a} * \text{d}) * \text{x}^{(2 * \text{b} * \text{d})}})^{\text{p}}), \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{d}, \text{e}, \text{m}, \text{p}\}, \text{x}]$

### Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

method	result	size
risch	$\frac{1}{4x^4} + 2e^{2a} \ln(x) - \frac{e^{2a} \ln(1+e^{2a}x^4)}{2}$	32

input  $\text{int}(\tanh(\text{a}+2*\ln(\text{x}))/\text{x}^5, \text{x}, \text{method}=\_RETURNVERBOSE)$

output  $1/4/x^4+2*\exp(2*a)*\ln(x)-1/2*\exp(2*a)*\ln(1+\exp(2*a)*x^4)$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = -\frac{2x^4 e^{(2a)} \log(x^4 e^{(2a)} + 1) - 8x^4 e^{(2a)} \log(x) - 1}{4x^4}$$

input `integrate(tanh(a+2*log(x))/x^5,x, algorithm="fricas")`

output  $-1/4*(2*x^4*e^{(2*a)}*\log(x^4*e^{(2*a)} + 1) - 8*x^4*e^{(2*a)}*\log(x) - 1)/x^4$

### Sympy [F]

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = \int \frac{\tanh(a + 2 \log(x))}{x^5} dx$$

input `integrate(tanh(a+2*ln(x))/x**5,x)`

output `Integral(tanh(a + 2*log(x))/x**5, x)`

### Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.55

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = -\frac{1}{2} e^{(2a)} \log\left(\frac{1}{x^4} + e^{(2a)}\right) + \frac{1}{4x^4}$$

input `integrate(tanh(a+2*log(x))/x^5,x, algorithm="maxima")`

output  $-1/2*e^{(2*a)}*\log(1/x^4 + e^{(2*a)}) + 1/4/x^4$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = -\frac{1}{2} e^{(2a)} \log(x^4 e^{(2a)} + 1) + \frac{1}{2} e^{(2a)} \log(x^4) - \frac{2x^4 e^{(2a)} - 1}{4x^4}$$

input `integrate(tanh(a+2*log(x))/x^5,x, algorithm="giac")`output `-1/2*e^(2*a)*log(x^4*e^(2*a) + 1) + 1/2*e^(2*a)*log(x^4) - 1/4*(2*x^4*e^(2*a) - 1)/x^4`**Mupad [B] (verification not implemented)**

Time = 2.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = 2e^{2a} \ln(x) - \frac{e^{2a} \ln(x^4 + e^{-2a})}{2} + \frac{1}{4x^4}$$

input `int(tanh(a + 2*log(x))/x^5,x)`output `2*exp(2*a)*log(x) - (exp(2*a)*log(exp(-2*a) + x^4))/2 + 1/(4*x^4)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.05

$$\int \frac{\tanh(a + 2 \log(x))}{x^5} dx = \frac{-2e^{2a} \log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^4 - 2e^{2a} \log(e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^4 + 8e^{2a} \log(x) x^4 + 1}{4x^4}$$

input `int(tanh(a+2*log(x))/x^5,x)`

output

```
( - 2*e**(2*a)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - 2*e**(2*a)
)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + 8*e**(2*a)*log(x)*x**4 +
1)/(4*x**4)
```

### 3.153 $\int x^2 \tanh(a + 2 \log(x)) dx$

Optimal result	1174
Mathematica [C] (verified)	1174
Rubi [A] (verified)	1175
Maple [C] (verified)	1178
Fricas [A] (verification not implemented)	1179
Sympy [F]	1179
Maxima [A] (verification not implemented)	1180
Giac [A] (verification not implemented)	1180
Mupad [B] (verification not implemented)	1181
Reduce [B] (verification not implemented)	1181

#### Optimal result

Integrand size = 11, antiderivative size = 109

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{x^3}{3} + \frac{e^{-3a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-3a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{\sqrt{2}} + \frac{e^{-3a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^{ax^2}}\right)}{\sqrt{2}}$$

output

```
1/3*x^3-1/2*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(3/2*a)-1/2*arctan(
1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(3/2*a)+1/2*arctanh(2^(1/2)*exp(1/2*a)*
x/(1+exp(a)*x^2))*2^(1/2)/exp(3/2*a)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.16 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.59

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{1}{6} \left( 2x^3 + 3 \operatorname{RootSum} \left[ \cosh(a) - \sinh(a) + \cosh(a)\#1^4 + \sinh(a)\#1^4 \&, \frac{\log(x) - \log(x - \#1)}{\#1} \& \right] (\cosh(2a) - \sinh(2a)) \right)$$

input `Integrate[x^2*Tanh[a + 2*Log[x]],x]`

output `(2*x^3 + 3*RootSum[Cosh[a] - Sinh[a] + Cosh[a]**#1^4 + Sinh[a]**#1^4 & , (Log[x] - Log[x - #1])/#1 & ]*(Cosh[2*a] - Sinh[2*a]))/6`

### Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.60, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.909$ , Rules used = {6071, 959, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tanh(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{x^2 (e^{2a} x^4 - 1)}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{x^3}{3} - 2 \int \frac{x^2}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{826} \\
 & \frac{x^3}{3} - 2 \left( \frac{1}{2} e^{-a} \int \frac{e^a x^2 + 1}{e^{2a} x^4 + 1} dx - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) \\
 & \quad \downarrow \text{1476} \\
 & \frac{x^3}{3} - 2 \left( \frac{1}{2} e^{-a} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx \right) - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$



$$\begin{aligned}
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{1}{-(1-\sqrt{2}e^{a/2}x)^2-1} d(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x+1)^2-1} d(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1}{e^{2ax}} dx \right) \\
& \quad \downarrow \text{217} \\
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1-e^ax^2}{e^{2ax^4}+1} dx \right) \\
& \quad \downarrow \text{1479} \\
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( -\frac{e^{-a/2} \int -\frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} \right) \right) \\
& \quad \downarrow \text{25} \\
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} + \dots \right) \right) \\
& \quad \downarrow \text{27} \\
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} + \dots \right) \right) \\
& \quad \downarrow \text{1103} \\
& 2 \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \log(e^ax^2 + \sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} \right) \right)
\end{aligned}$$

input `Int[x^2*Tanh[a + 2*Log[x]],x]`

output

$$x^3/3 - 2*((-\text{ArcTan}[1 - \text{Sqrt}[2]*E^{(a/2)*x}/(\text{Sqrt}[2]*E^{(a/2)})] + \text{ArcTan}[1 + \text{Sqrt}[2]*E^{(a/2)*x}/(\text{Sqrt}[2]*E^{(a/2)})])/(2*E^a) - (-1/2*\text{Log}[1 - \text{Sqrt}[2]*E^{(a/2)*x} + E^a*x^2]/(\text{Sqrt}[2]*E^{(a/2)}) + \text{Log}[1 + \text{Sqrt}[2]*E^{(a/2)*x} + E^a*x^2]/(2*\text{Sqrt}[2]*E^{(a/2)}))/(2*E^a)$$

## Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \text{:>} \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x\_Symbol}] \text{:>} \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{;/;} \text{FreeQ}[a, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) \text{;/;} \text{FreeQ}[b, \text{x}]]$$

rule 217

$$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, \text{x\_Symbol}] \text{:>} \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] \text{;/;} \text{FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 826

$$\text{Int}[(x_)^2/((a_) + (b_)*(x_)^4), \text{x\_Symbol}] \text{:>} \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}] - \text{Simp}[1/(2*s) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}]] \text{;/;} \text{FreeQ}[\{a, b\}, \text{x}] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 959

$$\text{Int}[(e_)*(x_)^{(m_)*((a_) + (b_)*(x_)^{(n_)})^{(p_)*((c_) + (d_)*(x_)^{(n_))}}, \text{x\_Symbol}] \text{:>} \text{Simp}[d*(e*x)^{(m+1)*((a+b*x^n)^{(p+1)/(b*e*(m+n*(p+1)+1))}, \text{x}] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \quad \text{Int}[(e*x)^m*(a+b*x^n)^p, \text{x}], \text{x}] \text{;/;} \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, \text{x}] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$$

rule 1082

$$\text{Int}[((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, \text{x\_Symbol}] \text{:>} \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), \text{x}], \text{x}, 1 + 2*c*(x/b)], \text{x}] \text{;/;} \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4*a*c])] \text{;/;} \text{FreeQ}[\{a, b, c\}, \text{x}]$$

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6071 `Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.45 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.34

method	result	size
risch	$\frac{x^3}{3} - \frac{e^{-2a} \left( \sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-R)}{-R} \right)}{2}$	37

input `int(x^2*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)`

output `1/3*x^3-1/2*exp(-2*a)*sum(1/_R*ln(x-_R),_R=RootOf(exp(2*a)*_Z^4+1))`

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.02

$$\int x^2 \tanh(a + 2 \log(x)) dx$$

$$= \frac{1}{12} \left( 4x^3 e^a - 6\sqrt{2} \arctan\left(\sqrt{2}x e^{\frac{1}{2}a} + 1\right) e^{(-\frac{1}{2}a)} - 6\sqrt{2} \arctan\left(\sqrt{2}x e^{\frac{1}{2}a} - 1\right) e^{(-\frac{1}{2}a)} + 3\sqrt{2} e^{(-\frac{1}{2}a)} \log(x^2 e^a + \sqrt{2}x e^{\frac{1}{2}a} + 1) - 3\sqrt{2} e^{(-\frac{1}{2}a)} \log(x^2 e^a - \sqrt{2}x e^{\frac{1}{2}a} + 1) \right) e^{-a}$$

input `integrate(x^2*tanh(a+2*log(x)),x, algorithm="fricas")`

output `1/12*(4*x^3*e^a - 6*sqrt(2)*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(-1/2*a) - 6*sqrt(2)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(-1/2*a) + 3*sqrt(2)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 3*sqrt(2)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1))*e^(-a)`

**Sympy [F]**

$$\int x^2 \tanh(a + 2 \log(x)) dx = \int x^2 \tanh(a + 2 \log(x)) dx$$

input `integrate(x**2*tanh(a+2*ln(x)),x)`

output `Integral(x**2*tanh(a + 2*log(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.17

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{1}{3} x^3 - \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2 x e^a + \sqrt{2} e^{\frac{1}{2} a}) e^{(-\frac{1}{2} a)} \right) e^{(-\frac{3}{2} a)} - \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2 x e^a - \sqrt{2} e^{\frac{1}{2} a}) e^{(-\frac{1}{2} a)} \right) e^{(-\frac{3}{2} a)} + \frac{1}{4} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( x^2 e^a + \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) - \frac{1}{4} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( x^2 e^a - \sqrt{2} x e^{\frac{1}{2} a} + 1 \right)$$

input `integrate(x^2*tanh(a+2*log(x)),x, algorithm="maxima")`output `1/3*x^3 - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) + 1/4*sqrt(2)*e^(-3/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 1/4*sqrt(2)*e^(-3/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.13

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{1}{3} x^3 - \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2} e^{(-\frac{1}{2} a)} + 2 x) e^{\frac{1}{2} a} \right) e^{(-\frac{3}{2} a)} - \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2} e^{(-\frac{1}{2} a)} - 2 x) e^{\frac{1}{2} a} \right) e^{(-\frac{3}{2} a)} + \frac{1}{4} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( \sqrt{2} x e^{(-\frac{1}{2} a)} + x^2 + e^{(-a)} \right) - \frac{1}{4} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( -\sqrt{2} x e^{(-\frac{1}{2} a)} + x^2 + e^{(-a)} \right)$$

input `integrate(x^2*tanh(a+2*log(x)),x, algorithm="giac")`

output

```
1/3*x^3 - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2
*a))*e^(-3/2*a) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*
x)*e^(1/2*a))*e^(-3/2*a) + 1/4*sqrt(2)*e^(-3/2*a)*log(sqrt(2)*x*e^(-1/2*a)
+ x^2 + e^(-a)) - 1/4*sqrt(2)*e^(-3/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2
+ e^(-a))
```

**Mupad [B] (verification not implemented)**

Time = 2.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.43

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{3/4}} - \frac{\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{3/4}} + \frac{x^3}{3}$$

input

```
int(x^2*tanh(a + 2*log(x)),x)
```

output

```
atan(x*(-exp(2*a))^(1/4))/(-exp(2*a))^(3/4) - atanh(x*(-exp(2*a))^(1/4))/(-
exp(2*a))^(3/4) + x^3/3
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.39

$$\int x^2 \tanh(a + 2 \log(x)) dx = \frac{6e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) - 6e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) - 3e^{\frac{a}{2}}\sqrt{2} \log(-e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1) + 3e^{\frac{a}{2}}\sqrt{2} \log(e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1)}{12e^{2a}}$$

input

```
int(x^2*tanh(a+2*log(x)),x)
```

output

```
(6***(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2***a*x)/(e**(a/2)*sqrt(2)))
- 6***(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2***a*x)/(e**(a/2)*sqrt(2)
)) - 3***(a/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + ***a*x**2 + 1) + 3***
(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + ***a*x**2 + 1) + 4***(2*a)*x**3)/(
12***(2*a))
```

### 3.154 $\int \tanh(a + 2 \log(x)) dx$

Optimal result	1182
Mathematica [C] (verified)	1182
Rubi [A] (verified)	1183
Maple [C] (verified)	1186
Fricas [A] (verification not implemented)	1187
Sympy [F]	1187
Maxima [A] (verification not implemented)	1188
Giac [A] (verification not implemented)	1188
Mupad [B] (verification not implemented)	1189
Reduce [B] (verification not implemented)	1189

#### Optimal result

Integrand size = 7, antiderivative size = 104

$$\int \tanh(a + 2 \log(x)) dx = x + \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{\sqrt{2}}$$

output

```
x-1/2*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(1/2*a)-1/2*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(1/2*a)-1/2*arctanh(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)/exp(1/2*a)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.56

$$\int \tanh(a + 2 \log(x)) dx = x + \frac{1}{2} \operatorname{RootSum} \left[ \cosh(a) - \sinh(a) + \cosh(a) \#1^4 + \sinh(a) \#1^4 \&, \frac{\log(x) - \log(x - \#1)}{\#1^3} \& \right] (\cosh(2a) - \sinh(2a))$$

input `Integrate[Tanh[a + 2*Log[x]],x]`

output `x + (RootSum[Cosh[a] - Sinh[a] + Cosh[a]**#1^4 + Sinh[a]**#1^4 & , (Log[x] - Log[x - #1])/#1^3 & ]*(Cosh[2*a] - Sinh[2*a]))/2`

### Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.52, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.429$ , Rules used = {6067, 913, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6067} \\
 & \int \frac{e^{2a}x^4 - 1}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{913} \\
 & x - 2 \int \frac{1}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{755} \\
 & x - 2 \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \int \frac{e^ax^2 + 1}{e^{2a}x^4 + 1} dx \right) \\
 & \quad \downarrow \text{1476} \\
 & 2 \left( \frac{1}{2} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) + \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) \\
 & \quad \downarrow \text{1082} \\
 & 2 \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{1}{-(1 - \sqrt{2}e^{a/2}x)^2 - 1} d(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x + 1)^2 - 1} d(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} \right) \right)
 \end{aligned}$$



$$\begin{aligned}
& \downarrow 217 \\
x - 2 \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) \right) \\
& \downarrow 1479 \\
2 \left( \frac{1}{2} \left( - \frac{e^{-a/2} \int - \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} - \frac{e^{-a/2} \int - \frac{\sqrt{2}(\sqrt{2}x + e^{-a/2})}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) \right) \\
& \downarrow 25 \\
2 \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} + \frac{e^{-a/2} \int \frac{\sqrt{2}(\sqrt{2}x + e^{-a/2})}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) \right) \\
& \downarrow 27 \\
2 \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} + \frac{1}{2} e^{-a/2} \int \frac{\sqrt{2}x + e^{-a/2}}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) \right) \\
& \downarrow 1103 \\
2 \left( \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \log(e^a x^2 + \sqrt{2} e^{a/2} x + 1)}{2\sqrt{2}} - \frac{e^{-a/2} \log(e^a x^2 - \sqrt{2} e^{a/2} x + 1)}{2\sqrt{2}} \right) \right)
\end{aligned}$$

input `Int[Tanh[a + 2*Log[x]], x]`

output `x - 2*((-(ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))/2 + (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/2`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 913 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6067 `Int[Tanh[((a_) + Log[x]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.23 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.32

method	result	size
risch	$x - \frac{e^{-2a} \left( \sum_{R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-R)}{-R^3} \right)}{2}$	33

input `int(tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)`

output `x-1/2*exp(-2*a)*sum(1/_R^3*ln(x-_R),_R=RootOf(exp(2*a)*_Z^4+1))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.95

$$\int \tanh(a + 2 \log(x)) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) e^{-\frac{1}{2} a} \\ - \frac{1}{2} \sqrt{2} \arctan \left( \sqrt{2} x e^{\frac{1}{2} a} - 1 \right) e^{-\frac{1}{2} a} \\ - \frac{1}{4} \sqrt{2} e^{-\frac{1}{2} a} \log \left( x^2 e^a + \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) \\ + \frac{1}{4} \sqrt{2} e^{-\frac{1}{2} a} \log \left( x^2 e^a - \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) + x$$

input `integrate(tanh(a+2*log(x)),x, algorithm="fricas")`output `-1/2*sqrt(2)*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(-1/2*a) - 1/2*sqrt(2)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(-1/2*a) - 1/4*sqrt(2)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) + 1/4*sqrt(2)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + x`**Sympy [F]**

$$\int \tanh(a + 2 \log(x)) dx = \int \tanh(a + 2 \log(x)) dx$$

input `integrate(tanh(a+2*ln(x)),x)`output `Integral(tanh(a + 2*log(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.19

$$\int \tanh(a + 2 \log(x)) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a + \sqrt{2}e^{\frac{1}{2}a}) e^{(-\frac{1}{2}a)} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a - \sqrt{2}e^{\frac{1}{2}a}) e^{(-\frac{1}{2}a)} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{4} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( x^2 e^a + \sqrt{2} x e^{\frac{1}{2}a} + 1 \right)$$

$$+ \frac{1}{4} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( x^2 e^a - \sqrt{2} x e^{\frac{1}{2}a} + 1 \right) + x$$

input `integrate(tanh(a+2*log(x)),x, algorithm="maxima")`

output

```
-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*
e^(-1/2*a) - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*
e^(-1/2*a))*e^(-1/2*a) - 1/4*sqrt(2)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^
(1/2*a) + 1) + 1/4*sqrt(2)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) +
1) + x
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.14

$$\int \tanh(a + 2 \log(x)) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} + 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} - 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{4} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( \sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right)$$

$$+ \frac{1}{4} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( -\sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right) + x$$

input `integrate(tanh(a+2*log(x)),x, algorithm="giac")`

output

```
-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-
1/2*a) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2
*a))*e^(-1/2*a) - 1/4*sqrt(2)*e^(-1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 +
e^(-a)) + 1/4*sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a))
+ x
```

**Mupad [B] (verification not implemented)**

Time = 2.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.42

$$\int \tanh(a + 2 \log(x)) dx = x - \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{1/4}} - \frac{\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{1/4}}$$

input

```
int(tanh(a + 2*log(x)),x)
```

output

```
x - atan(x*(-exp(2*a))^(1/4))/(-exp(2*a))^(1/4) - atanh(x*(-exp(2*a))^(1/4
))/(-exp(2*a))^(1/4)
```

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.38

$$\int \tanh(a + 2 \log(x)) dx = \frac{2e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) - 2e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) + e^{\frac{a}{2}}\sqrt{2} \log(-e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1) - e^{\frac{a}{2}}\sqrt{2} \log(e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1)}{4e^a}$$

input

```
int(tanh(a+2*log(x)),x)
```

output

```
(2***(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2***a*x)/(e**(a/2)*sqrt(2)))
- 2***(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2***a*x)/(e**(a/2)*sqrt(2)
)) + e**(a/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) - e**(a/2
)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 4***a*x)/(4***a)
```

### 3.155 $\int \frac{\tanh(a+2 \log(x))}{x^2} dx$

Optimal result	1190
Mathematica [C] (verified)	1190
Rubi [A] (verified)	1191
Maple [C] (verified)	1195
Fricas [A] (verification not implemented)	1195
Sympy [F]	1196
Maxima [A] (verification not implemented)	1196
Giac [A] (verification not implemented)	1197
Mupad [B] (verification not implemented)	1197
Reduce [B] (verification not implemented)	1198

#### Optimal result

Integrand size = 11, antiderivative size = 106

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \frac{1}{x} - \frac{e^{a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} + \frac{e^{a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{\sqrt{2}}$$

output

```
1/x+1/2*exp(1/2*a)*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)+1/2*exp(1/2*a)*
arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)-1/2*exp(1/2*a)*arctanh(2^(1/2)*exp(
1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.56

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \frac{2 - x \operatorname{RootSum}\left[\cosh(a) + \sinh(a) + \cosh(a)\#1^4 - \sinh(a)\#1^4 \&, \frac{\log(x) + \log\left(\frac{1}{x} - \#1\right)}{\#1^3} \&\right] (\cosh(a) + \sinh(a))}{2x}$$

input `Integrate[Tanh[a + 2*Log[x]]/x^2,x]`

output `(2 - x*RootSum[Cosh[a] + Sinh[a] + Cosh[a]**#1^4 - Sinh[a]**#1^4 & , (Log[x] + Log[x^(-1) - #1])/#1^3 & ]*(Cosh[a] + Sinh[a])^2)/(2*x)`

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.65, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.909$ , Rules used = {6071, 955, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(a + 2 \log(x))}{x^2} dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{e^{2a}x^4 - 1}{x^2 (e^{2a}x^4 + 1)} dx \\
 & \quad \downarrow \text{955} \\
 & 2e^{2a} \int \frac{x^2}{e^{2a}x^4 + 1} dx + \frac{1}{x} \\
 & \quad \downarrow \text{826} \\
 & 2e^{2a} \left( \frac{1}{2} e^{-a} \int \frac{e^a x^2 + 1}{e^{2a}x^4 + 1} dx - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a}x^4 + 1} dx \right) + \frac{1}{x} \\
 & \quad \downarrow \text{1476} \\
 & 2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a}x^4 + 1} dx \right) + \frac{1}{x} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$



$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{1}{-(1-\sqrt{2}e^{a/2}x)^2-1} d(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x+1)^2-1} d(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1}{e^{2ax^4}+1} dx \right)$$

$\frac{1}{x}$   
↓ 217

$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1-e^ax^2}{e^{2ax^4}+1} dx \right) +$$

$\frac{1}{x}$   
↓ 1479

$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( -\frac{e^{-a/2} \int -\frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} \right) \right)$$

$\frac{1}{x}$   
↓ 25

$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} \right) \right)$$

$\frac{1}{x}$   
↓ 27

$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} \right) \right)$$

$\frac{1}{x}$   
↓ 1103

$$2e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \log(e^a x^2 + \sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} \right) \right) - \frac{1}{x}$$

input `Int[Tanh[a + 2*Log[x]]/x^2,x]`

output `x^(-1) + 2*E^(2*a)*((-ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))/(2*E^a) - (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/(2*E^a)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 955 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))), x] + Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*e^n*(m + 1)) Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_) + Log[x]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.40

method	result	size
risch	$\frac{1}{x} + \frac{\sum_{R=\text{RootOf}(\_Z^4+e^{2a})} -R \ln((5\_R^4+4e^{2a})x - \_R^3)}{2}$	42

input `int(tanh(a+2*ln(x))/x^2,x,method=_RETURNVERBOSE)`

output `1/x+1/2*sum(_R*ln((5*_R^4+4*exp(2*a))*x-_R^3),_R=RootOf(_Z^4+exp(2*a)))`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.01

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx$$

$$= \frac{2 \sqrt{2} x \arctan\left(\sqrt{2} x e^{\frac{1}{2} a} + 1\right) e^{\frac{1}{2} a} + 2 \sqrt{2} x \arctan\left(\sqrt{2} x e^{\frac{1}{2} a} - 1\right) e^{\frac{1}{2} a} - \sqrt{2} x e^{\frac{1}{2} a} \log\left(x^2 e^a + \sqrt{2} x\right)}{4 x}$$

input `integrate(tanh(a+2*log(x))/x^2,x, algorithm="fricas")`

output `1/4*(2*sqrt(2)*x*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(1/2*a) + 2*sqrt(2)*x*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(1/2*a) - sqrt(2)*x*e^(1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) + sqrt(2)*x*e^(1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + 4)/x`

**Sympy [F]**

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \int \frac{\tanh(a + 2 \log(x))}{x^2} dx$$

input `integrate(tanh(a+2*ln(x))/x**2,x)`

output `Integral(tanh(a + 2*log(x))/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.18

$$\begin{aligned} \int \frac{\tanh(a + 2 \log(x))}{x^2} dx = & -\frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2} a} + \frac{2}{x} \right) e^{-\frac{1}{2} a} \right) e^{\frac{1}{2} a} \\ & - \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2} a} - \frac{2}{x} \right) e^{-\frac{1}{2} a} \right) e^{\frac{1}{2} a} \\ & - \frac{1}{4} \sqrt{2} e^{\frac{1}{2} a} \log \left( \frac{\sqrt{2} e^{\frac{1}{2} a}}{x} + \frac{1}{x^2} + e^a \right) \\ & + \frac{1}{4} \sqrt{2} e^{\frac{1}{2} a} \log \left( -\frac{\sqrt{2} e^{\frac{1}{2} a}}{x} + \frac{1}{x^2} + e^a \right) + \frac{1}{x} \end{aligned}$$

input `integrate(tanh(a+2*log(x))/x^2,x, algorithm="maxima")`

output `-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) + 2/x)*e^(-1/2*a))*e^(1/2*a) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) - 2/x)*e^(-1/2*a))*e^(1/2*a) - 1/4*sqrt(2)*e^(1/2*a)*log(sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + 1/4*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + 1/x`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{(-\frac{1}{2} a)} + 2x \right) e^{\frac{1}{2} a} \right) e^{\frac{1}{2} a}$$

$$+ \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{(-\frac{1}{2} a)} - 2x \right) e^{\frac{1}{2} a} \right) e^{\frac{1}{2} a}$$

$$- \frac{1}{4} \sqrt{2} e^{\frac{1}{2} a} \log \left( \sqrt{2} x e^{(-\frac{1}{2} a)} + x^2 + e^{(-a)} \right)$$

$$+ \frac{1}{4} \sqrt{2} e^{\frac{1}{2} a} \log \left( -\sqrt{2} x e^{(-\frac{1}{2} a)} + x^2 + e^{(-a)} \right) + \frac{1}{x}$$

input `integrate(tanh(a+2*log(x))/x^2,x, algorithm="giac")`

output

```
1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(1/2*a) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(1/2*a) - 1/4*sqrt(2)*e^(1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/4*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/x
```

**Mupad [B] (verification not implemented)**

Time = 2.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.42

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \operatorname{atan} \left( x (-e^{2a})^{1/4} \right) (-e^{2a})^{1/4}$$

$$- \operatorname{atanh} \left( x (-e^{2a})^{1/4} \right) (-e^{2a})^{1/4} + \frac{1}{x}$$

input `int(tanh(a + 2*log(x))/x^2,x)`

output

```
atan(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4) - atanh(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4) + 1/x
```

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.33

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx$$

$$= \frac{-2e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x + 2e^{\frac{a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x + e^{\frac{a}{2}}\sqrt{2} \log(-e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1) x - e^{\frac{a}{2}}\sqrt{2} \log(e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1) x}{4x}$$

input

```
int(tanh(a+2*log(x))/x^2,x)
```

output

```
( - 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))
*x + 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))
*x + e**(a/2)*sqrt(2)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x
- e**(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x + 4)/(4*x)
```

### 3.156 $\int \frac{\tanh(a+2 \log(x))}{x^4} dx$

Optimal result	1199
Mathematica [C] (verified)	1199
Rubi [A] (verified)	1200
Maple [C] (verified)	1203
Fricas [A] (verification not implemented)	1204
Sympy [F]	1204
Maxima [A] (verification not implemented)	1205
Giac [A] (verification not implemented)	1205
Mupad [B] (verification not implemented)	1206
Reduce [B] (verification not implemented)	1206

#### Optimal result

Integrand size = 11, antiderivative size = 109

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \frac{1}{3x^3} - \frac{e^{3a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} + \frac{e^{3a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{\sqrt{2}} + \frac{e^{3a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{\sqrt{2}}$$

output

```
1/3/x^3+1/2*exp(3/2*a)*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)+1/2*exp(3/2*a)*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)+1/2*exp(3/2*a)*arctanh(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.56

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \frac{2 - 3x^3 \operatorname{RootSum}\left[\cosh(a) + \sinh(a) + \cosh(a)\#1^4 - \sinh(a)\#1^4 \&, \frac{\log(x) + \log\left(\frac{1}{x} - \#1\right)}{\#1}\right] \&}{6x^3} (\cosh(a) + \sinh(a))$$



input `Integrate[Tanh[a + 2*Log[x]]/x^4,x]`

output `(2 - 3*x^3*RootSum[Cosh[a] + Sinh[a] + Cosh[a]**#1^4 - Sinh[a]**#1^4 & , (Log[x] + Log[x^(-1) - #1])/#1 & ]*(Cosh[a] + Sinh[a])^2)/(6*x^3)`

### Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.55, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.909$ , Rules used = {6071, 955, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(a + 2 \log(x))}{x^4} dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{e^{2a}x^4 - 1}{x^4 (e^{2a}x^4 + 1)} dx \\
 & \quad \downarrow \text{955} \\
 & 2e^{2a} \int \frac{1}{e^{2a}x^4 + 1} dx + \frac{1}{3x^3} \\
 & \quad \downarrow \text{755} \\
 & 2e^{2a} \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \int \frac{e^ax^2 + 1}{e^{2a}x^4 + 1} dx \right) + \frac{1}{3x^3} \\
 & \quad \downarrow \text{1476} \\
 & 2e^{2a} \left( \frac{1}{2} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) + \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) + \frac{1}{3x^3} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

$$2e^{2a} \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{1}{-(1 - \sqrt{2}e^{a/2}x)^2 - 1} d(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x + 1)^2 - 1} d(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} \right) \right)$$

$$\frac{1}{3x^3}$$

↓ 217

$$2e^{2a} \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) +$$

$$\frac{1}{3x^3}$$

↓ 1479

$$2e^{2a} \left( \frac{1}{2} \left( -\frac{e^{-a/2} \int -\frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} - \frac{e^{-a/2} \int -\frac{\sqrt{2}(\sqrt{2}x + e^{-a/2})}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right)$$

$$\frac{1}{3x^3}$$

↓ 25

$$2e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} + \frac{e^{-a/2} \int \frac{\sqrt{2}(\sqrt{2}x + e^{-a/2})}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right)$$

$$\frac{1}{3x^3}$$

↓ 27

$$2e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} + \frac{1}{2} e^{-a/2} \int \frac{\sqrt{2}x + e^{-a/2}}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right)$$

$$\frac{1}{3x^3}$$

↓ 1103

$$2e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \log(e^a x^2 + \sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} - \frac{e^{-a/2} \log(e^a x^2 - \sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} \right) \right)$$

$$\frac{1}{3x^3}$$

input `Int[Tanh[a + 2*Log[x]]/x^4,x]`

output 
$$\frac{1}{3x^3} + 2E^{2a} \left( \frac{-\operatorname{ArcTan}\left[\frac{1 - \sqrt{2}E^{a/2}x}{\sqrt{2}E^{a/2}}\right]}{\sqrt{2}E^{a/2}} + \frac{\operatorname{ArcTan}\left[\frac{1 + \sqrt{2}E^{a/2}x}{\sqrt{2}E^{a/2}}\right]}{\sqrt{2}E^{a/2}} \right) / 2 + \frac{-1/2 \operatorname{Log}\left[\frac{1 - \sqrt{2}E^{a/2}x + E^a x^2}{\sqrt{2}E^{a/2}}\right] + \operatorname{Log}\left[\frac{1 + \sqrt{2}E^{a/2}x + E^a x^2}{2\sqrt{2}E^{a/2}}\right]}{2}$$

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 955 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))), x] + Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*e^n*(m + 1)) Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6071 `Int[((e_)*(x_)^(m_)*Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.31 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

method	result	size
risch	$\frac{1}{3x^3} + \frac{\sum_{-R=\text{RootOf}(e^{6a}+_Z^4)} -R \ln((-4e^{6a}-5_R^4)x+_R e^{4a})}{2}$	45

input `int(tanh(a+2*ln(x))/x^4,x,method=_RETURNVERBOSE)`

output `1/3/x^3+1/2*sum(_R*ln((-4*exp(6*a)-5*_R^4)*x+_R*exp(4*a)),_R=RootOf(exp(6*a)+_Z^4))`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.06

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx$$

$$= \frac{6 \sqrt{2} x^3 \arctan\left(\sqrt{2} x e^{\frac{1}{2} a} + 1\right) e^{\frac{3}{2} a} + 6 \sqrt{2} x^3 \arctan\left(\sqrt{2} x e^{\frac{1}{2} a} - 1\right) e^{\frac{3}{2} a} + 3 \sqrt{2} x^3 e^{\frac{3}{2} a} \log\left(x^2 e^a + \sqrt{2} x e^{\frac{1}{2} a} + 1\right) - 3 \sqrt{2} x^3 e^{\frac{3}{2} a} \log\left(x^2 e^a - \sqrt{2} x e^{\frac{1}{2} a} + 1\right) + 4}{12 x^3}$$

input `integrate(tanh(a+2*log(x))/x^4,x, algorithm="fricas")`

output `1/12*(6*sqrt(2)*x^3*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(3/2*a) + 6*sqrt(2)*x^3*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(3/2*a) + 3*sqrt(2)*x^3*e^(3/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 3*sqrt(2)*x^3*e^(3/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + 4)/x^3`

### Sympy [F]

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \int \frac{\tanh(a + 2 \log(x))}{x^4} dx$$

input `integrate(tanh(a+2*ln(x))/x**4,x)`

output `Integral(tanh(a + 2*log(x))/x**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.22

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx =$$

$$-\frac{1}{4} \left( 2\sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} + \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{-\frac{1}{2}a} + 2\sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} - \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{-\frac{1}{2}a} \right)$$

$$+ \frac{1}{3x^3}$$

input `integrate(tanh(a+2*log(x))/x^4,x, algorithm="maxima")`output `-1/4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) + 2/x)*e^(-1/2*a))*e^(-1/2*a) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) - 2/x)*e^(-1/2*a))*e^(-1/2*a) - sqrt(2)*e^(-1/2*a)*log(sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a))*e^(2*a) + 1/3/x^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.13

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{-\frac{1}{2}a} + 2x \right) e^{\frac{1}{2}a} \right) e^{\frac{3}{2}a}$$

$$+ \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{-\frac{1}{2}a} - 2x \right) e^{\frac{1}{2}a} \right) e^{\frac{3}{2}a}$$

$$+ \frac{1}{4} \sqrt{2} e^{\frac{3}{2}a} \log \left( \sqrt{2} x e^{-\frac{1}{2}a} + x^2 + e^{-a} \right)$$

$$- \frac{1}{4} \sqrt{2} e^{\frac{3}{2}a} \log \left( -\sqrt{2} x e^{-\frac{1}{2}a} + x^2 + e^{-a} \right) + \frac{1}{3x^3}$$

input `integrate(tanh(a+2*log(x))/x^4,x, algorithm="giac")`

output

$$\begin{aligned} & 1/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} + 2*x)*e^{(1/2*a)})*e^{(3/2*a)} \\ & + 1/2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} - 2*x)*e^{(1/2*a)})*e^{(3/2*a)} \\ & + 1/4*\sqrt{2}*e^{(3/2*a)}*\log(\sqrt{2}*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) - 1/4*\sqrt{2}*e^{(3/2*a)}*\log(-\sqrt{2}*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + 1/3/x^3 \end{aligned}$$
**Mupad [B] (verification not implemented)**

Time = 2.10 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.44

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \frac{1}{3x^3} - \operatorname{atanh}\left(x(-e^{2a})^{1/4}\right) (-e^{2a})^{3/4} - \operatorname{atan}\left(x(-e^{2a})^{1/4}\right) (-e^{2a})^{3/4}$$

input

`int(tanh(a + 2*log(x))/x^4,x)`

output

$$1/(3*x^3) - \operatorname{atanh}(x*(-\exp(2*a))^{(1/4)})*(-\exp(2*a))^{(3/4)} - \operatorname{atan}(x*(-\exp(2*a))^{(1/4)})*(-\exp(2*a))^{(3/4)}$$
**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.38

$$\int \frac{\tanh(a + 2 \log(x))}{x^4} dx = \frac{-6e^{\frac{3a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^3 + 6e^{\frac{3a}{2}}\sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^3 - 3e^{\frac{3a}{2}}\sqrt{2} \log(-e^{\frac{a}{2}}\sqrt{2}x + e^ax^2 + 1) x^3 + \dots}{12x^3}$$

input

`int(tanh(a+2*log(x))/x^4,x)`

output

$$\begin{aligned} & (-6*e^{((3*a)/2)}*\sqrt{2}*\operatorname{atan}((e^{(a/2)}*\sqrt{2} - 2*e^{a*x})/(e^{(a/2)}*\sqrt{2}))*x^{**3} + 6*e^{((3*a)/2)}*\sqrt{2}*\operatorname{atan}((e^{(a/2)}*\sqrt{2} + 2*e^{a*x})/(e^{(a/2)}*\sqrt{2}))*x^{**3} \\ & - 3*e^{((3*a)/2)}*\sqrt{2}*\log(-e^{(a/2)}*\sqrt{2}*x + e^{a*x} + 1)*x^{**3} + 3*e^{((3*a)/2)}*\sqrt{2}*\log(e^{(a/2)}*\sqrt{2}*x + e^{a*x} + 1)*x^{**3} + 4)/(12*x^{**3}) \end{aligned}$$

### 3.157 $\int x^5 \tanh^2(a + 2 \log(x)) dx$

Optimal result	1207
Mathematica [A] (verified)	1207
Rubi [A] (verified)	1208
Maple [C] (verified)	1210
Fricas [A] (verification not implemented)	1210
Sympy [F]	1211
Maxima [A] (verification not implemented)	1211
Giac [A] (verification not implemented)	1212
Mupad [B] (verification not implemented)	1212
Reduce [B] (verification not implemented)	1212

#### Optimal result

Integrand size = 13, antiderivative size = 56

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = -2e^{-2a}x^2 + \frac{x^6}{6} - \frac{e^{-2a}x^2}{1 + e^{2a}x^4} + 3e^{-3a} \arctan(e^a x^2)$$

output

`-2*x^2/exp(2*a)+1/6*x^6-x^2/exp(2*a)/(1+exp(2*a)*x^4)+3*arctan(exp(a)*x^2)/exp(3*a)`

#### Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = \frac{1}{6}x^2 \left( -12e^{-2a} + x^4 - \frac{6}{e^{2a} + e^{4a}x^4} \right) + 3e^{-3a} \arctan(e^a x^2)$$

input

`Integrate[x^5*Tanh[a + 2*Log[x]]^2,x]`

output

`(x^2*(-12/E^(2*a) + x^4 - 6/(E^(2*a) + E^(4*a)*x^4)))/6 + (3*ArcTan[E^a*x^2])/E^(3*a)`



**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.25, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {6071, 963, 27, 959, 807, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \tanh^2(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{x^5 (e^{2a} x^4 - 1)^2}{(e^{2a} x^4 + 1)^2} dx \\
 & \quad \downarrow \text{963} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - \frac{1}{4} e^{-4a} \int \frac{4x^5 (5e^{4a} - e^{6a} x^4)}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - e^{-4a} \int \frac{x^5 (5e^{4a} - e^{6a} x^4)}{e^{2a} x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - e^{-4a} \left( 6e^{4a} \int \frac{x^5}{e^{2a} x^4 + 1} dx - \frac{1}{6} e^{4a} x^6 \right) \\
 & \quad \downarrow \text{807} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - e^{-4a} \left( 3e^{4a} \int \frac{x^4}{e^{2a} x^4 + 1} dx^2 - \frac{1}{6} e^{4a} x^6 \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - e^{-4a} \left( 3e^{4a} \left( e^{-2a} x^2 - e^{-2a} \int \frac{1}{e^{2a} x^4 + 1} dx^2 \right) - \frac{1}{6} e^{4a} x^6 \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{x^6}{e^{2a} x^4 + 1} - e^{-4a} \left( 3e^{4a} \left( e^{-2a} x^2 - e^{-3a} \arctan(e^a x^2) \right) - \frac{1}{6} e^{4a} x^6 \right)
 \end{aligned}$$

input `Int[x^5*Tanh[a + 2*Log[x]]^2,x]`

output `x^6/(1 + E^(2*a)*x^4) - (-1/6*(E^(4*a)*x^6) + 3*E^(4*a)*(x^2/E^(2*a) - ArcTan[E^a*x^2]/E^(3*a)))/E^(4*a)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(b*(m+2*p+1))), x] - Simp[a*c^2*((m-1)/(b*(m+2*p+1)) Int[(c*x)^(m-2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2-1] && NeQ[m+2*p+1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m+1, n]}, Simp[1/k Subst[Int[x^((m+1)/k-1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 959 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m+1)*((a + b*x^n)^(p+1)/(b*e*(m+n*(p+1)+1))), x] - Simp[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m+n*(p+1)+1, 0]`

rule 963

```
Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^(n._))^(p._)*((c._) + (d._)*(x._)^(n._))
^2, x_Symbol] := Simp[(-b*c - a*d)^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)
/(a*b^2*e*n*(p + 1))), x] + Simp[1/(a*b^2*n*(p + 1)) Int[(e*x)^m*(a + b*x
^n)^(p + 1)*Simp[(b*c - a*d)^2*(m + 1) + b^2*c^2*n*(p + 1) + a*b*d^2*n*(p +
1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0]
&& IGtQ[n, 0] && LtQ[p, -1]
```

rule 6071

```
Int[((e._)*(x._))^(m._)*Tanh[((a._) + Log[x_]*(b._))*(d._)]^(p._), x_Symbol]
:= Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

### Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.38

method	result	size
risch	$\frac{x^6}{6} - 2e^{-2a}x^2 + \frac{8e^{-3a}}{3} - \frac{x^2e^{-2a}}{1+e^{2a}x^4} + \frac{3ie^{-3a}\ln(e^ax^2+i)}{2} - \frac{3ie^{-3a}\ln(e^ax^2-i)}{2}$	77

input

```
int(x^5*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/6*x^6-2/exp(a)^2*x^2+8/3/exp(a)^3-x^2/exp(a)^2/(exp(a)^2*x^4+1)+3/2*I/ex
p(a)^3*ln(exp(a)*x^2+I)-3/2*I/exp(a)^3*ln(exp(a)*x^2-I)
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09

$$\int x^5 \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{x^{10}e^{(5a)} - 11x^6e^{(3a)} - 18x^2e^a + 18(x^4e^{(2a)} + 1) \arctan(x^2e^a)}{6(x^4e^{(5a)} + e^{(3a)})}$$

input

```
integrate(x^5*tanh(a+2*log(x))^2,x, algorithm="fricas")
```

output  $1/6*(x^{10}*e^{(5*a)} - 11*x^6*e^{(3*a)} - 18*x^2*e^a + 18*(x^4*e^{(2*a)} + 1)*\arctan(x^2*e^a))/(x^4*e^{(5*a)} + e^{(3*a)})$

### Sympy [F]

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = \int x^5 \tanh^2(a + 2 \log(x)) dx$$

input `integrate(x**5*tanh(a+2*ln(x))**2,x)`

output `Integral(x**5*tanh(a + 2*log(x))**2, x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = \frac{1}{6} (x^6 e^{(2a)} - 12 x^2) e^{(-2a)} + 3 \arctan(x^2 e^a) e^{(-3a)} - \frac{x^2}{x^4 e^{(4a)} + e^{(2a)}}$$

input `integrate(x^5*tanh(a+2*log(x))^2,x, algorithm="maxima")`

output  $1/6*(x^6*e^{(2*a)} - 12*x^2)*e^{(-2*a)} + 3*\arctan(x^2*e^a)*e^{(-3*a)} - x^2/(x^4*e^{(4*a)} + e^{(2*a)})$

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = -\frac{x^2 e^{(-2a)}}{x^4 e^{(2a)} + 1} + 3 \arctan(x^2 e^a) e^{(-3a)} + \frac{1}{6} (x^6 e^{(12a)} - 12 x^2 e^{(10a)}) e^{(-12a)}$$

input `integrate(x^5*tanh(a+2*log(x))^2,x, algorithm="giac")`output `-x^2*e^(-2*a)/(x^4*e^(2*a) + 1) + 3*arctan(x^2*e^a)*e^(-3*a) + 1/6*(x^6*e^(12*a) - 12*x^2*e^(10*a))*e^(-12*a)`**Mupad [B] (verification not implemented)**

Time = 2.14 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = \frac{3 \operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{(e^{2a})^{3/2}} - 2x^2 e^{-2a} - \frac{x^2}{e^{4a} x^4 + e^{2a}} + \frac{x^6}{6}$$

input `int(x^5*tanh(a + 2*log(x))^2,x)`output `(3*atan(x^2*exp(2*a)^(1/2)))/exp(2*a)^(3/2) - 2*x^2*exp(-2*a) - x^2/(exp(2*a) + x^4*exp(4*a)) + x^6/6`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 186, normalized size of antiderivative = 3.32

$$\int x^5 \tanh^2(a + 2 \log(x)) dx = \frac{-18e^{2a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 - 18 \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) - 18e^{2a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 - 18 \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) + 6e^{3a} (e^{2a} x^4 + 1)}{6e^{3a} (e^{2a} x^4 + 1)}$$

input `int(x^5*tanh(a+2*log(x))^2,x)`

output  $(-18e^{2a}\operatorname{atan}\left(\frac{e^{a/2}\sqrt{2}-2e^{ax}}{e^{a/2}\sqrt{2}}\right)x^4 - 18\operatorname{atan}\left(\frac{e^{a/2}\sqrt{2}-2e^{ax}}{e^{a/2}\sqrt{2}}\right) - 18e^{2a}\operatorname{atan}\left(\frac{e^{a/2}\sqrt{2}+2e^{ax}}{e^{a/2}\sqrt{2}}\right)x^4 - 18\operatorname{atan}\left(\frac{e^{a/2}\sqrt{2}+2e^{ax}}{e^{a/2}\sqrt{2}}\right) + e^{5a}x^{10} - 11e^{3a}x^6 - 18e^{ax^2})/(6e^{3a}(e^{2a}x^4 + 1))$

### 3.158 $\int x^3 \tanh^2(a + 2 \log(x)) dx$

Optimal result	1214
Mathematica [A] (verified)	1214
Rubi [A] (verified)	1215
Maple [A] (verified)	1216
Fricas [A] (verification not implemented)	1217
Sympy [F]	1217
Maxima [A] (verification not implemented)	1217
Giac [A] (verification not implemented)	1218
Mupad [B] (verification not implemented)	1218
Reduce [B] (verification not implemented)	1218

#### Optimal result

Integrand size = 13, antiderivative size = 47

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{x^4}{4} - \frac{e^{-2a}}{1 + e^{2a}x^4} - e^{-2a} \log(1 + e^{2a}x^4)$$

output `1/4*x^4-1/exp(2*a)/(1+exp(2*a)*x^4)-ln(1+exp(2*a)*x^4)/exp(2*a)`

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.83

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{x^4}{4} - \cosh(2a) \log((1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)) + \log((1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)) \sinh(2a) + \frac{-\cosh(3a) + \sinh(3a)}{(1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)}$$

input `Integrate[x^3*Tanh[a + 2*Log[x]]^2,x]`

output

$$x^4/4 - \text{Cosh}[2*a]*\text{Log}[(1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]] + \text{Log}[(1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]]*\text{Sinh}[2*a] + (-\text{Cosh}[3*a] + \text{Sinh}[3*a])/((1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a])$$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6071, 946, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \tanh^2(a + 2 \log(x)) dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{x^3 (e^{2a} x^4 - 1)^2}{(e^{2a} x^4 + 1)^2} dx \\ & \quad \downarrow \text{946} \\ & \frac{1}{4} \int \frac{(1 - e^{2a} x^4)^2}{(e^{2a} x^4 + 1)^2} dx^4 \\ & \quad \downarrow \text{49} \\ & \frac{1}{4} \int \left( 1 - \frac{4}{e^{2a} x^4 + 1} + \frac{4}{(e^{2a} x^4 + 1)^2} \right) dx^4 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \left( -\frac{4e^{-2a}}{e^{2a} x^4 + 1} - 4e^{-2a} \log(e^{2a} x^4 + 1) + x^4 \right) \end{aligned}$$

input

$$\text{Int}[x^3*\text{Tanh}[a + 2*\text{Log}[x]]^2,x]$$

output

$$(x^4 - 4/(E^(2*a)*(1 + E^(2*a)*x^4)) - (4*\text{Log}[1 + E^(2*a)*x^4])/E^(2*a))/4$$



## Definitions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

method	result	size
risch	$\frac{x^4}{4} - \frac{e^{-2a}}{1+e^{2a}x^4} - e^{-2a} \ln(1 + e^{2a}x^4)$	42

input `int(x^3*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)`

output `1/4*x^4-exp(-2*a)/(1+exp(2*a)*x^4)-exp(-2*a)*ln(1+exp(2*a)*x^4)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.23

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{x^8 e^{(4a)} + x^4 e^{(2a)} - 4(x^4 e^{(2a)} + 1) \log(x^4 e^{(2a)} + 1) - 4}{4(x^4 e^{(4a)} + e^{(2a)})}$$

input `integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="fricas")`

output `1/4*(x^8*e^(4*a) + x^4*e^(2*a) - 4*(x^4*e^(2*a) + 1)*log(x^4*e^(2*a) + 1) - 4)/(x^4*e^(4*a) + e^(2*a))`

**Sympy [F]**

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \int x^3 \tanh^2(a + 2 \log(x)) dx$$

input `integrate(x**3*tanh(a+2*ln(x))**2,x)`

output `Integral(x**3*tanh(a + 2*log(x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.85

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{1}{4} x^4 - e^{(-2a)} \log(x^4 e^{(2a)} + 1) - \frac{1}{x^4 e^{(4a)} + e^{(2a)}}$$

input `integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="maxima")`

output `1/4*x^4 - e^(-2*a)*log(x^4*e^(2*a) + 1) - 1/(x^4*e^(4*a) + e^(2*a))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{1}{4} x^4 + \frac{x^4}{x^4 e^{(2a)} + 1} - e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

input `integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="giac")`output `1/4*x^4 + x^4/(x^4*e^(2*a) + 1) - e^(-2*a)*log(x^4*e^(2*a) + 1)`**Mupad [B] (verification not implemented)**

Time = 2.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{x^4}{4} - \frac{e^{-2a}}{e^{2a} x^4 + 1} - e^{-2a} \ln(x^4 + e^{-2a})$$

input `int(x^3*tanh(a + 2*log(x))^2,x)`output `x^4/4 - exp(-2*a)/(x^4*exp(2*a) + 1) - exp(-2*a)*log(exp(-2*a) + x^4)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.06

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \frac{e^{4a} x^8 - 4e^{2a} \log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^4 - 4e^{2a} \log(e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^4 + 5e^{2a} x^4 - 4 \log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^4}{4e^{2a} (e^{2a} x^4 + 1)}$$

input `int(x^3*tanh(a+2*log(x))^2,x)`

output

```
(e**(4*a)*x**8 - 4*e**(2*a)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - 4*e**(2*a)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + 5*e**(2*a)*x**4 - 4*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) - 4*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1))/(4*e**(2*a)*(e**(2*a)*x**4 + 1))
```

### 3.159 $\int x \tanh^2(a + 2 \log(x)) dx$

Optimal result	1220
Mathematica [A] (verified)	1220
Rubi [A] (verified)	1221
Maple [C] (verified)	1223
Fricas [A] (verification not implemented)	1223
Sympy [F]	1223
Maxima [A] (verification not implemented)	1224
Giac [A] (verification not implemented)	1224
Mupad [B] (verification not implemented)	1224
Reduce [B] (verification not implemented)	1225

#### Optimal result

Integrand size = 11, antiderivative size = 40

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{x^2}{2} + \frac{x^2}{1 + e^{2a}x^4} - e^{-a} \arctan(e^a x^2)$$

output  $1/2*x^2+x^2/(1+\exp(2*a)*x^4)-\arctan(\exp(a)*x^2)/\exp(a)$

#### Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{x^2}{2} + \frac{x^2}{1 + e^{2(a+2 \log(x))}} - e^{-a} \arctan(e^a x^2)$$

input `Integrate[x*Tanh[a + 2*Log[x]]^2,x]`

output  $x^2/2 + x^2/(1 + E^{(2*(a + 2*Log[x]))}) - \text{ArcTan}[E^a*x^2]/E^a$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.30, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {6071, 963, 27, 959, 807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tanh^2(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{x(e^{2a}x^4 - 1)^2}{(e^{2a}x^4 + 1)^2} dx \\
 & \quad \downarrow \text{963} \\
 & \frac{x^2}{e^{2a}x^4 + 1} - \frac{1}{4}e^{-4a} \int \frac{4x(e^{4a} - e^{6a}x^4)}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{x^2}{e^{2a}x^4 + 1} - e^{-4a} \int \frac{x(e^{4a} - e^{6a}x^4)}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{x^2}{e^{2a}x^4 + 1} - e^{-4a} \left( 2e^{4a} \int \frac{x}{e^{2a}x^4 + 1} dx - \frac{1}{2}e^{4a}x^2 \right) \\
 & \quad \downarrow \text{807} \\
 & \frac{x^2}{e^{2a}x^4 + 1} - e^{-4a} \left( e^{4a} \int \frac{1}{e^{2a}x^4 + 1} dx^2 - \frac{1}{2}e^{4a}x^2 \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{x^2}{e^{2a}x^4 + 1} - e^{-4a} \left( e^{3a} \arctan(e^a x^2) - \frac{1}{2}e^{4a}x^2 \right)
 \end{aligned}$$

input

```
Int[x*Tanh[a + 2*Log[x]]^2,x]
```

output  $x^2/(1 + E^{(2*a)*x^4}) - (-1/2*(E^{(4*a)*x^2}) + E^{(3*a)*ArcTan[E^a*x^2]})/E^{(4*a)}$

### Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$

rule 216  $\text{Int}[(a_*) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 807  $\text{Int}[(x_)^{(m_)*((a_*) + (b_)*(x_)^{(n_)})^{(p_)}}, x\_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{((m + 1)/k - 1)*(a + b*x^{(n/k)})^p}, x], x, x^k], x] /; k \neq 1] /; \text{FreeQ}\{a, b, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 959  $\text{Int}[(e_)*(x_)^{(m_)*((a_*) + (b_)*(x_)^{(n_)})^{(p_)*((c_*) + (d_)*(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m + 1)*((a + b*x^n)^{(p + 1))/(b*e*(m + n*(p + 1) + 1))}, x] - \text{Simp}[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) \ \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p + 1) + 1, 0]$

rule 963  $\text{Int}[(e_)*(x_)^{(m_)*((a_*) + (b_)*(x_)^{(n_)})^{(p_)*((c_*) + (d_)*(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)^2*(e*x)^{(m + 1)*((a + b*x^n)^{(p + 1))/(a*b^2*e*n*(p + 1))}, x] + \text{Simp}[1/(a*b^2*n*(p + 1)) \ \text{Int}[(e*x)^m*(a + b*x^n)^{(p + 1)*\text{Simp}[(b*c - a*d)^2*(m + 1) + b^2*c^2*n*(p + 1) + a*b*d^2*n*(p + 1)*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1]$

rule 6071  $\text{Int}[(e_)*(x_)^{(m_)*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_)]*(d_)]^{(p_)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)})^p/(1 + E^{(2*a*d)*x^{(2*b*d)})^p), x] /; \text{FreeQ}\{a, b, d, e, m, p\}, x]$

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.42

method	result	size
risch	$\frac{x^2}{2} + \frac{x^2}{1+e^{2a}x^4} + \frac{ie^{-a} \ln(e^a x^2 - i)}{2} - \frac{ie^{-a} \ln(e^a x^2 + i)}{2}$	57

input `int(x*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}x^2 + x^2 / (\exp(a)^2 x^4 + 1) + 1/2 * I / \exp(a) * \ln(\exp(a) * x^2 - I) - 1/2 * I / \exp(a) * \ln(\exp(a) * x^2 + I)$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.25

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{x^6 e^{(3a)} + 3 x^2 e^a - 2 (x^4 e^{(2a)} + 1) \arctan(x^2 e^a)}{2 (x^4 e^{(3a)} + e^a)}$$

input `integrate(x*tanh(a+2*log(x))^2,x, algorithm="fricas")`

output  $\frac{1}{2} * (x^6 * e^{(3*a)} + 3 * x^2 * e^a - 2 * (x^4 * e^{(2*a)} + 1) * \arctan(x^2 * e^a)) / (x^4 * e^{(3*a)} + e^a)$

**Sympy [F]**

$$\int x \tanh^2(a + 2 \log(x)) dx = \int x \tanh^2(a + 2 \log(x)) dx$$

input `integrate(x*tanh(a+2*ln(x))**2,x)`

output `Integral(x*tanh(a + 2*log(x))**2, x)`



**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{1}{2} x^2 - \arctan(x^2 e^a) e^{-a} + \frac{x^2}{x^4 e^{2a} + 1}$$

input `integrate(x*tanh(a+2*log(x))^2,x, algorithm="maxima")`output `1/2*x^2 - arctan(x^2*e^a)*e^(-a) + x^2/(x^4*e^(2*a) + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{1}{2} x^2 - \arctan(x^2 e^a) e^{-a} + \frac{x^2}{x^4 e^{2a} + 1}$$

input `integrate(x*tanh(a+2*log(x))^2,x, algorithm="giac")`output `1/2*x^2 - arctan(x^2*e^a)*e^(-a) + x^2/(x^4*e^(2*a) + 1)`**Mupad [B] (verification not implemented)**

Time = 2.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int x \tanh^2(a + 2 \log(x)) dx = \frac{x^2}{e^{2a} x^4 + 1} - \frac{\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{\sqrt{e^{2a}}} + \frac{x^2}{2}$$

input `int(x*tanh(a + 2*log(x))^2,x)`output `x^2/(x^4*exp(2*a) + 1) - atan(x^2*exp(2*a)^(1/2))/exp(2*a)^(1/2) + x^2/2`

**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 174, normalized size of antiderivative = 4.35

$$\int x \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{2e^{2a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^4 + 2\operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) + 2e^{2a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^4 + 2\operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) + e^{3a}x^6}{2e^a(e^{2a}x^4 + 1)}$$

input `int(x*tanh(a+2*log(x))^2,x)`

output

```
(2*e**(2*a)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 +
2*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) + 2*e**(2*a)*atan
((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 + 2*atan((e**(a/2)
*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) + e**(3*a)*x**6 + 3*e**a*x**2)/(2
*e**a*(e**(2*a)*x**4 + 1))
```

### 3.160 $\int \frac{\tanh^2(a+2\log(x))}{x} dx$

Optimal result	1226
Mathematica [A] (verified)	1226
Rubi [A] (verified)	1227
Maple [A] (verified)	1228
Fricas [B] (verification not implemented)	1229
Sympy [A] (verification not implemented)	1229
Maxima [A] (verification not implemented)	1229
Giac [A] (verification not implemented)	1230
Mupad [B] (verification not implemented)	1230
Reduce [B] (verification not implemented)	1230

#### Optimal result

Integrand size = 13, antiderivative size = 14

$$\int \frac{\tanh^2(a + 2\log(x))}{x} dx = \log(x) - \frac{1}{2} \tanh(a + 2\log(x))$$

output `ln(x)-1/2*tanh(a+2*ln(x))`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.71

$$\int \frac{\tanh^2(a + 2\log(x))}{x} dx = \frac{1}{2} \operatorname{arctanh}(\tanh(a + 2\log(x))) - \frac{1}{2} \tanh(a + 2\log(x))$$

input `Integrate[Tanh[a + 2*Log[x]]^2/x,x]`

output `ArcTanh[Tanh[a + 2*Log[x]]]/2 - Tanh[a + 2*Log[x]]/2`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {3039, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(a + 2 \log(x))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \tanh^2(a + 2 \log(x)) d \log(x) \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan(ia + 2i \log(x))^2 d \log(x) \\
 & \quad \downarrow \text{25} \\
 & - \int \tan(ia + 2i \log(x))^2 d \log(x) \\
 & \quad \downarrow \text{3954} \\
 & \int 1 d \log(x) - \frac{1}{2} \tanh(a + 2 \log(x)) \\
 & \quad \downarrow \text{24} \\
 & \log(x) - \frac{1}{2} \tanh(a + 2 \log(x))
 \end{aligned}$$

input

```
Int[Tanh[a + 2*Log[x]]^2/x,x]
```

output

```
Log[x] - Tanh[a + 2*Log[x]]/2
```

## Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst  
[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /;  
NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d  
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]  
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

## Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
parallelrisc	$\ln(x) - \frac{\tanh(a+2\ln(x))}{2}$	13
risc	$\frac{1}{1+e^{2ax^4}} + \ln(x)$	16
derivativedivides	$-\frac{\tanh(a+2\ln(x))}{2} - \frac{\ln(\tanh(a+2\ln(x))-1)}{4} + \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	35
default	$-\frac{\tanh(a+2\ln(x))}{2} - \frac{\ln(\tanh(a+2\ln(x))-1)}{4} + \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	35

input `int(tanh(a+2*ln(x))^2/x,x,method=_RETURNVERBOSE)`

output `ln(x)-1/2*tanh(a+2*ln(x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 28 vs.  $2(12) = 24$ .

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \frac{(x^4 e^{(2a)} + 1) \log(x) + 1}{x^4 e^{(2a)} + 1}$$

input `integrate(tanh(a+2*log(x))^2/x,x, algorithm="fricas")`

output `((x^4*e^(2*a) + 1)*log(x) + 1)/(x^4*e^(2*a) + 1)`

**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \log(x) - \frac{\tanh(a + 2 \log(x))}{2}$$

input `integrate(tanh(a+2*ln(x))**2/x,x)`

output `log(x) - tanh(a + 2*log(x))/2`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.50

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \frac{1}{2} a - \frac{1}{e^{(-2a - 4 \log(x))} + 1} + \log(x)$$

input `integrate(tanh(a+2*log(x))^2/x,x, algorithm="maxima")`

output `1/2*a - 1/(e^(-2*a - 4*log(x)) + 1) + log(x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \frac{1}{x^4 e^{(2a)} + 1} + \frac{1}{4} \log(x^4)$$

input `integrate(tanh(a+2*log(x))^2/x,x, algorithm="giac")`

output `1/(x^4*e^(2*a) + 1) + 1/4*log(x^4)`

**Mupad [B] (verification not implemented)**

Time = 2.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \ln(x) - \frac{x^4 e^{2a} - 1}{2(e^{2a} x^4 + 1)}$$

input `int(tanh(a + 2*log(x))^2/x,x)`

output `log(x) - (x^4*exp(2*a) - 1)/(2*(x^4*exp(2*a) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{\tanh^2(a + 2 \log(x))}{x} dx = \log(x) - \frac{\tanh(2 \log(x) + a)}{2}$$

input `int(tanh(a+2*log(x))^2/x,x)`

output `(2*log(x) - tanh(2*log(x) + a))/2`

### 3.161 $\int \frac{\tanh^2(a+2\log(x))}{x^3} dx$

Optimal result	1231
Mathematica [A] (verified)	1231
Rubi [A] (verified)	1232
Maple [C] (verified)	1234
Fricas [A] (verification not implemented)	1234
Sympy [F]	1235
Maxima [A] (verification not implemented)	1235
Giac [A] (verification not implemented)	1235
Mupad [B] (verification not implemented)	1236
Reduce [B] (verification not implemented)	1236

#### Optimal result

Integrand size = 13, antiderivative size = 44

$$\int \frac{\tanh^2(a + 2\log(x))}{x^3} dx = -\frac{1}{2x^2} - \frac{e^{2a}x^2}{1 + e^{2a}x^4} - e^a \arctan(e^a x^2)$$

output

```
-1/2/x^2-exp(2*a)*x^2/(1+exp(2*a)*x^4)-exp(a)*arctan(exp(a)*x^2)
```

#### Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.91

$$\int \frac{\tanh^2(a + 2\log(x))}{x^3} dx = \frac{-1 - \frac{2}{1+e^{-2(a+2\log(x))}}}{2x^2} + e^a \arctan\left(\frac{e^{-a}}{x^2}\right)$$

input

```
Integrate[Tanh[a + 2*Log[x]]^2/x^3,x]
```

output

```
(-1 - 2/(1 + E^(-2*(a + 2*Log[x]))) )/(2*x^2) + E^a*ArcTan[1/(E^a*x^2)]
```



**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.34, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {6071, 962, 27, 957, 807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{(e^{2a}x^4 - 1)^2}{x^3 (e^{2a}x^4 + 1)^2} dx \\
 & \quad \downarrow \text{962} \\
 & \frac{1}{2} \int -\frac{2x(5e^{2a} - e^{4a}x^4)}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{2x^2 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{27} \\
 & - \int \frac{x(5e^{2a} - e^{4a}x^4)}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{2x^2 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{957} \\
 & -2e^{2a} \int \frac{x}{e^{2a}x^4 + 1} dx - \frac{3e^{2a}x^2}{2(e^{2a}x^4 + 1)} - \frac{1}{2x^2 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{807} \\
 & -e^{2a} \int \frac{1}{e^{2a}x^4 + 1} dx^2 - \frac{3e^{2a}x^2}{2(e^{2a}x^4 + 1)} - \frac{1}{2x^2 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{216} \\
 & -e^a \arctan(e^a x^2) - \frac{3e^{2a}x^2}{2(e^{2a}x^4 + 1)} - \frac{1}{2x^2 (e^{2a}x^4 + 1)}
 \end{aligned}$$

input

```
Int[Tanh[a + 2*Log[x]]^2/x^3,x]
```

output 
$$-1/2*1/(x^2*(1 + E^{(2*a)*x^4})) - (3*E^{(2*a)*x^2})/(2*(1 + E^{(2*a)*x^4})) - E^{a*ArcTan[E^a*x^2]}$$

### Defintions of rubi rules used

rule 27 
$$\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$$

rule 216 
$$\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$$

rule 807 
$$\text{Int}[(x_)^{(m_*)}*((a_*) + (b_*)(x_)^{(n_)})^{(p_)}, x\_Symbol] \rightarrow \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Simp}[1/k \ \text{Subst}[\text{Int}[x^{((m + 1)/k - 1)*(a + b*x^{(n/k)})^p}], x, x^k], x] /; k \neq 1 /; \text{FreeQ}\{a, b, p\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$$

rule 957 
$$\text{Int}[(e_*)(x_)^{(m_*)}*((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}*((c_*) + (d_*)(x_)^{(n_)})], x\_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*(e*x)^{(m + 1)}*((a + b*x^n)^{(p + 1)})/(a*b*e*n*(p + 1)), x] - \text{Simp}[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*b*n*(p + 1)) \ \text{Int}[(e*x)^m*(a + b*x^n)^{(p + 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ ((\ !\text{IntegerQ}[p + 1/2] \ \&\& \ \text{NeQ}[p, -5/4]) \ || \ \ !\text{RationalQ}[m] \ || \ (\text{IGtQ}[n, 0] \ \&\& \ \text{ILtQ}[p + 1/2, 0] \ \&\& \ \text{LeQ}[-1, m, (-n)*(p + 1)]))$$

rule 962 
$$\text{Int}[(e_*)(x_)^{(m_*)}*((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}*((c_*) + (d_*)(x_)^{(n_)})^2], x\_Symbol] \rightarrow \text{Simp}[c^2*(e*x)^{(m + 1)}*((a + b*x^n)^{(p + 1)})/(a*e*(m + 1)), x] - \text{Simp}[1/(a*e^n*(m + 1)) \ \text{Int}[(e*x)^{(m + n)}*(a + b*x^n)^p * \text{Simp}[b*c^2*n*(p + 1) + c*(b*c - 2*a*d)*(m + 1) - a*(m + 1)*d^2*x^n, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{GtQ}[n, 0]$$

rule 6071

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.30 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.50

method	result	size
risch	$\frac{-\frac{3}{2}e^{2a}x^4 - \frac{1}{2}}{x^2(1+e^{2a}x^4)} + \frac{\sum_{R=\text{RootOf}(e^{2a}+_Z^2)} -R \ln((-4e^{2a}-5_R^2)x^2 - R)}{2}$	66

input

```
int(tanh(a+2*ln(x))^2/x^3,x,method=_RETURNVERBOSE)
```

output

```
(-3/2*exp(2*a)*x^4-1/2)/x^2/(1+exp(2*a)*x^4)+1/2*sum(_R*ln((-4*exp(2*a)-5*_R^2)*x^2-_R),_R=RootOf(exp(2*a)+_Z^2))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.16

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = -\frac{3x^4e^{(2a)} + 2(x^6e^{(3a)} + x^2e^a) \arctan(x^2e^a) + 1}{2(x^6e^{(2a)} + x^2)}$$

input

```
integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="fricas")
```

output

```
-1/2*(3*x^4*e^(2*a) + 2*(x^6*e^(3*a) + x^2*e^a)*arctan(x^2*e^a) + 1)/(x^6*
e^(2*a) + x^2)
```

**Sympy [F]**

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx$$

input `integrate(tanh(a+2*ln(x))**2/x**3,x)`

output `Integral(tanh(a + 2*log(x))**2/x**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = \arctan\left(\frac{e^{(-a)}}{x^2}\right) e^a - \frac{1}{2x^2} - \frac{e^{(2a)}}{x^2\left(\frac{1}{x^4} + e^{(2a)}\right)}$$

input `integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="maxima")`

output `arctan(e^(-a)/x^2)*e^a - 1/2/x^2 - e^(2*a)/(x^2*(1/x^4 + e^(2*a)))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = -\arctan(x^2 e^a) e^a - \frac{3x^4 e^{(2a)} + 1}{2(x^6 e^{(2a)} + x^2)}$$

input `integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="giac")`

output `-arctan(x^2*e^a)*e^a - 1/2*(3*x^4*e^(2*a) + 1)/(x^6*e^(2*a) + x^2)`

**Mupad [B] (verification not implemented)**

Time = 2.16 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.07

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = -\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right) \sqrt{e^{2a}} - \frac{3e^{2a}x^4}{2} + \frac{1}{2} \frac{1}{e^{2a}x^6 + x^2}$$

input `int(tanh(a + 2*log(x))^2/x^3,x)`output `- atan(x^2*exp(2*a)^(1/2))*exp(2*a)^(1/2) - ((3*x^4*exp(2*a))/2 + 1/2)/(x^6*exp(2*a) + x^2)`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 178, normalized size of antiderivative = 4.05

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = \frac{2e^{3a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^6 + 2e^a \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}-2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^2 + 2e^{3a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^6 + 2e^a \operatorname{atan}\left(\frac{e^{\frac{a}{2}}\sqrt{2}+2e^ax}{e^{\frac{a}{2}}\sqrt{2}}\right) x^2}{2x^2(e^{2a}x^4 + 1)}$$

input `int(tanh(a+2*log(x))^2/x^3,x)`output `(2***3*a)*atan((**(a/2)*sqrt(2) - 2***a*x)/(**(a/2)*sqrt(2)))*x**6 + 2***a*atan((**(a/2)*sqrt(2) - 2***a*x)/(**(a/2)*sqrt(2)))*x**2 + 2***3*a)*atan((**(a/2)*sqrt(2) + 2***a*x)/(**(a/2)*sqrt(2)))*x**6 + 2***a*atan((**(a/2)*sqrt(2) + 2***a*x)/(**(a/2)*sqrt(2)))*x**2 - 3***2*a)*x**4 - 1)/(2*x**2*(e**(2*a)*x**4 + 1))`

### 3.162 $\int \frac{\tanh^2(a+2\log(x))}{x^5} dx$

Optimal result	1237
Mathematica [A] (verified)	1237
Rubi [A] (verified)	1238
Maple [A] (verified)	1239
Fricas [A] (verification not implemented)	1240
Sympy [F]	1240
Maxima [A] (verification not implemented)	1240
Giac [A] (verification not implemented)	1241
Mupad [B] (verification not implemented)	1241
Reduce [B] (verification not implemented)	1241

#### Optimal result

Integrand size = 13, antiderivative size = 55

$$\int \frac{\tanh^2(a + 2\log(x))}{x^5} dx = -\frac{1}{4x^4} - \frac{e^{2a}}{1 + e^{2a}x^4} - 4e^{2a}\log(x) + e^{2a}\log(1 + e^{2a}x^4)$$

output `-1/4/x^4-exp(2*a)/(1+exp(2*a)*x^4)-4*exp(2*a)*ln(x)+exp(2*a)*ln(1+exp(2*a)*x^4)`

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.75

$$\begin{aligned} \int \frac{\tanh^2(a + 2\log(x))}{x^5} dx &= -\frac{1}{4x^4} - 4\cosh(2a)\log(x) \\ &+ \cosh(2a)\log((1 + x^4)\cosh(a) + (-1 + x^4)\sinh(a)) \\ &- \frac{\cosh(a) + \sinh(a)}{(1 + x^4)\cosh(a) + (-1 + x^4)\sinh(a)} - 4\log(x)\sinh(2a) \\ &+ \log((1 + x^4)\cosh(a) + (-1 + x^4)\sinh(a))\sinh(2a) \end{aligned}$$

input `Integrate[Tanh[a + 2*Log[x]]^2/x^5,x]`

output

$$-1/4*1/x^4 - 4*\text{Cosh}[2*a]*\text{Log}[x] + \text{Cosh}[2*a]*\text{Log}[(1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]] - (\text{Cosh}[a] + \text{Sinh}[a])/((1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]) - 4*\text{Log}[x]*\text{Sinh}[2*a] + \text{Log}[(1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]]*\text{Sinh}[2*a]$$
**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6071, 948, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{(e^{2a}x^4 - 1)^2}{x^5 (e^{2a}x^4 + 1)^2} dx \\ & \quad \downarrow \text{948} \\ & \frac{1}{4} \int \frac{(1 - e^{2a}x^4)^2}{x^8 (e^{2a}x^4 + 1)^2} dx^4 \\ & \quad \downarrow \text{99} \\ & \frac{1}{4} \int \left( -\frac{4e^{2a}}{x^4} + \frac{1}{x^8} + \frac{4e^{4a}}{e^{2a}x^4 + 1} + \frac{4e^{4a}}{(e^{2a}x^4 + 1)^2} \right) dx^4 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \left( -\frac{4e^{2a}}{e^{2a}x^4 + 1} - 4e^{2a} \log(x^4) + 4e^{2a} \log(e^{2a}x^4 + 1) - \frac{1}{x^4} \right) \end{aligned}$$

input

$$\text{Int}[\text{Tanh}[a + 2*\text{Log}[x]]^2/x^5, x]$$

output

$$(-x^{(-4)} - (4*E^{(2*a)})/(1 + E^{(2*a)}*x^4) - 4*E^{(2*a)}*\text{Log}[x^4] + 4*E^{(2*a)}*\text{Log}[1 + E^{(2*a)}*x^4])/4$$

## Definitions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))`

rule 948 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.98

method	result	size
risch	$\frac{-5e^{2a}x^4 - 1}{x^4(1+e^{2a}x^4)} - 4e^{2a} \ln(x) + e^{2a} \ln(-e^{2a}x^4 - 1)$	54

input `int(tanh(a+2*ln(x))^2/x^5,x,method=_RETURNVERBOSE)`

output `(-5/4*exp(2*a)*x^4-1/4)/x^4/(1+exp(2*a)*x^4)-4*exp(2*a)*ln(x)+exp(2*a)*ln(-exp(2*a)*x^4-1)`



**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.42

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = \frac{5x^4 e^{(2a)} - 4(x^8 e^{(4a)} + x^4 e^{(2a)}) \log(x^4 e^{(2a)} + 1) + 16(x^8 e^{(4a)} + x^4 e^{(2a)}) \log(x) + 1}{4(x^8 e^{(2a)} + x^4)}$$

input `integrate(tanh(a+2*log(x))^2/x^5,x, algorithm="fricas")`

output `-1/4*(5*x^4*e^(2*a) - 4*(x^8*e^(4*a) + x^4*e^(2*a))*log(x^4*e^(2*a) + 1) + 16*(x^8*e^(4*a) + x^4*e^(2*a))*log(x) + 1)/(x^8*e^(2*a) + x^4)`

**Sympy [F]**

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx$$

input `integrate(tanh(a+2*ln(x))**2/x**5,x)`

output `Integral(tanh(a + 2*log(x))**2/x**5, x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.64

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = e^{(2a)} \log\left(\frac{1}{x^4} + e^{(2a)}\right) + \frac{e^{(4a)}}{\frac{1}{x^4} + e^{(2a)}} - \frac{1}{4x^4}$$

input `integrate(tanh(a+2*log(x))^2/x^5,x, algorithm="maxima")`

output `e^(2*a)*log(1/x^4 + e^(2*a)) + e^(4*a)/(1/x^4 + e^(2*a)) - 1/4/x^4`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.98

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = e^{(2a)} \log(x^4 e^{(2a)} + 1) - e^{(2a)} \log(x^4) - \frac{5x^4 e^{(2a)} + 1}{4(x^8 e^{(2a)} + x^4)}$$

input `integrate(tanh(a+2*log(x))^2/x^5,x, algorithm="giac")`output `e^(2*a)*log(x^4*e^(2*a) + 1) - e^(2*a)*log(x^4) - 1/4*(5*x^4*e^(2*a) + 1)/  
(x^8*e^(2*a) + x^4)`**Mupad [B] (verification not implemented)**

Time = 2.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.91

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = e^{2a} \ln(x^4 + e^{-2a}) - \frac{\frac{5e^{2a}x^4}{4} + \frac{1}{4}}{e^{2a}x^8 + x^4} - 4e^{2a} \ln(x)$$

input `int(tanh(a + 2*log(x))^2/x^5,x)`output `exp(2*a)*log(exp(-2*a) + x^4) - ((5*x^4*exp(2*a))/4 + 1/4)/(x^8*exp(2*a) +  
x^4) - 4*exp(2*a)*log(x)`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 172, normalized size of antiderivative = 3.13

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^5} dx = \frac{4e^{4a} \log(-e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^8 + 4e^{4a} \log(e^{\frac{a}{2}} \sqrt{2} x + e^a x^2 + 1) x^8 - 16e^{4a} \log(x) x^8 + 5e^{4a} x^8 + 4e^{2a} \log(x)}{4x^4 (e^{2a} x^4 + 1)}$$

input `int(tanh(a+2*log(x))^2/x^5,x)`

output

```
(4*e**(4*a)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**8 + 4*e**(4*a)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**8 - 16*e**(4*a)*log(x)*x**8 + 5*e**(4*a)*x**8 + 4*e**(2*a)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + 4*e**(2*a)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - 16*e**(2*a)*log(x)*x**4 - 1)/(4*x**4*(e**(2*a)*x**4 + 1))
```

### 3.163 $\int \frac{\tanh^2(a+2\log(x))}{x^7} dx$

Optimal result	1243
Mathematica [A] (verified)	1243
Rubi [A] (verified)	1244
Maple [C] (verified)	1246
Fricas [A] (verification not implemented)	1246
Sympy [F]	1247
Maxima [A] (verification not implemented)	1247
Giac [A] (verification not implemented)	1248
Mupad [B] (verification not implemented)	1248
Reduce [B] (verification not implemented)	1248

#### Optimal result

Integrand size = 13, antiderivative size = 55

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = -\frac{1}{6x^6} + \frac{2e^{2a}}{x^2} + \frac{e^{4a}x^2}{1 + e^{2a}x^4} + 3e^{3a} \arctan(e^a x^2)$$

output

`-1/6/x^6+2*exp(2*a)/x^2+exp(4*a)*x^2/(1+exp(2*a)*x^4)+3*exp(3*a)*arctan(exp(a)*x^2)`

#### Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.04

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = -\frac{1}{6x^6} + \frac{2e^{2a}}{x^2} + \frac{e^{4a}x^2}{1 + e^{2a}x^4} - 3e^{3a} \arctan\left(\frac{e^{-a}}{x^2}\right)$$

input

`Integrate[Tanh[a + 2*Log[x]]^2/x^7,x]`

output

`-1/6*1/x^6 + (2*E^(2*a))/x^2 + (E^(4*a)*x^2)/(1 + E^(2*a)*x^4) - 3*E^(3*a)*ArcTan[1/(E^a*x^2)]`

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.40, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {6071, 962, 27, 957, 807, 264, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{(e^{2a}x^4 - 1)^2}{x^7 (e^{2a}x^4 + 1)^2} dx \\
 & \quad \downarrow \text{962} \\
 & \frac{1}{6} \int -\frac{2(11e^{2a} - 3e^{4a}x^4)}{x^3 (e^{2a}x^4 + 1)^2} dx - \frac{1}{6x^6 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{3} \int \frac{11e^{2a} - 3e^{4a}x^4}{x^3 (e^{2a}x^4 + 1)^2} dx - \frac{1}{6x^6 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{957} \\
 & \frac{1}{3} \left( -18e^{2a} \int \frac{1}{x^3 (e^{2a}x^4 + 1)} dx - \frac{7e^{2a}}{2x^2 (e^{2a}x^4 + 1)} \right) - \frac{1}{6x^6 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{3} \left( -9e^{2a} \int \frac{1}{x^4 (e^{2a}x^4 + 1)} dx^2 - \frac{7e^{2a}}{2x^2 (e^{2a}x^4 + 1)} \right) - \frac{1}{6x^6 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{264} \\
 & \frac{1}{3} \left( -9e^{2a} \left( -e^{2a} \int \frac{1}{e^{2a}x^4 + 1} dx^2 - \frac{1}{x^2} \right) - \frac{7e^{2a}}{2x^2 (e^{2a}x^4 + 1)} \right) - \frac{1}{6x^6 (e^{2a}x^4 + 1)} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{3} \left( -9e^{2a} \left( -e^a \arctan(e^a x^2) - \frac{1}{x^2} \right) - \frac{7e^{2a}}{2x^2 (e^{2a}x^4 + 1)} \right) - \frac{1}{6x^6 (e^{2a}x^4 + 1)}
 \end{aligned}$$

input `Int[Tanh[a + 2*Log[x]]^2/x^7,x]`

output `-1/6*1/(x^6*(1 + E^(2*a)*x^4)) + ((-7*E^(2*a))/(2*x^2*(1 + E^(2*a)*x^4)) - 9*E^(2*a)*(-x^(-2) - E^a*ArcTan[E^a*x^2]))/3`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 264 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 807 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 957 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(-b*c - a*d)*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*b*e*n*(p + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*b*n*(p + 1)) Int[(e*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (( !IntegerQ[p + 1/2] && NeQ[p, -5/4]) || !RationalQ[m] || (IGtQ[n, 0] && ILtQ[p + 1/2, 0] && LeQ[-1, m, (-n)*(p + 1)]))`

rule 962

```
Int[((e._)*(x_))^(m_)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))
^2, x_Symbol] :> Simp[c^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))
), x] - Simp[1/(a*e^n*(m + 1)) Int[(e*x)^(m + n)*(a + b*x^n)^p*Simp[b*c^2
*n*(p + 1) + c*(b*c - 2*a*d)*(m + 1) - a*(m + 1)*d^2*x^n, x], x] /; Fre
eQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1]
&& GtQ[n, 0]
```

rule 6071

```
Int[((e._)*(x_))^(m_)*Tanh[((a_) + Log[x]*(b._))*(d._)]^(p_), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p,
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.44

method	result	size
risch	$\frac{3e^{4a}x^8 + \frac{11e^{2a}x^4}{6} - \frac{1}{6}}{x^6(1+e^{2a}x^4)} + \frac{3 \left( \sum_{-R=\text{RootOf}(e^{6a}+Z^2)} -R \ln((4e^{6a}+5-R^2)x^2 - e^{2a}-R) \right)}{2}$	79

input

```
int(tanh(a+2*ln(x))^2/x^7,x,method=_RETURNVERBOSE)
```

output

```
(3*exp(4*a)*x^8+11/6*exp(2*a)*x^4-1/6)/x^6/(1+exp(2*a)*x^4)+3/2*sum(_R*ln(
(4*exp(6*a)+5*_R^2)*x^2-exp(2*a)*_R),_R=RootOf(exp(6*a)+_Z^2))
```

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.13

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx$$

$$= \frac{18x^8e^{(4a)} + 11x^4e^{(2a)} + 18(x^{10}e^{(5a)} + x^6e^{(3a)}) \arctan(x^2e^a) - 1}{6(x^{10}e^{(2a)} + x^6)}$$

input `integrate(tanh(a+2*log(x))^2/x^7,x, algorithm="fricas")`

output  $\frac{1}{6} \cdot (18x^8e^{4a} + 11x^4e^{2a} + 18(x^{10}e^{5a} + x^6e^{3a})) \cdot \arctan\left(\frac{x^2e^a - 1}{x^{10}e^{2a} + x^6}\right)$

## Sympy [F]

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx$$

input `integrate(tanh(a+2*ln(x))**2/x**7,x)`

output `Integral(tanh(a + 2*log(x))**2/x**7, x)`

## Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.87

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = -3 \arctan\left(\frac{e^{-a}}{x^2}\right) e^{3a} + \frac{2e^{2a}}{x^2} + \frac{e^{4a}}{x^2\left(\frac{1}{x^4} + e^{2a}\right)} - \frac{1}{6x^6}$$

input `integrate(tanh(a+2*log(x))^2/x^7,x, algorithm="maxima")`

output  $-3 \cdot \arctan\left(\frac{e^{-a}}{x^2}\right) \cdot e^{3a} + 2 \cdot e^{2a} / x^2 + e^{4a} / (x^2 \cdot (1/x^4 + e^{2a})) - 1/6/x^6$



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.91

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = \frac{x^2 e^{(4a)}}{x^4 e^{(2a)} + 1} + 3 \arctan(x^2 e^a) e^{(3a)} + \frac{12 x^4 e^{(2a)} - 1}{6 x^6}$$

input `integrate(tanh(a+2*log(x))^2/x^7,x, algorithm="giac")`output `x^2*e^(4*a)/(x^4*e^(2*a) + 1) + 3*arctan(x^2*e^a)*e^(3*a) + 1/6*(12*x^4*e^(2*a) - 1)/x^6`**Mupad [B] (verification not implemented)**

Time = 2.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = 3 \operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right) (e^{2a})^{3/2} + \frac{3 e^{4a} x^8 + \frac{11 e^{2a} x^4}{6} - \frac{1}{6}}{e^{2a} x^{10} + x^6}$$

input `int(tanh(a + 2*log(x))^2/x^7,x)`output `3*atan(x^2*exp(2*a)^(1/2))*exp(2*a)^(3/2) + ((11*x^4*exp(2*a))/6 + 3*x^8*exp(4*a) - 1/6)/(x^10*exp(2*a) + x^6)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 192, normalized size of antiderivative = 3.49

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^7} dx = \frac{-18e^{5a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2-2e^a x}}{e^{\frac{a}{2}} \sqrt{2}}\right) x^{10} - 18e^{3a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2-2e^a x}}{e^{\frac{a}{2}} \sqrt{2}}\right) x^6 - 18e^{5a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2+2e^a x}}{e^{\frac{a}{2}} \sqrt{2}}\right) x^{10} - 18e^{3a} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2+2e^a x}}{e^{\frac{a}{2}} \sqrt{2}}\right) x^6}{6x^6 (e^{2a} x^4 + 1)}$$

input `int(tanh(a+2*log(x))^2/x^7,x)`

output

```
( - 18*e**(5*a)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**
10 - 18*e**(3*a)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**
*6 - 18*e**(5*a)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**
*10 - 18*e**(3*a)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**
**6 + 18*e**(4*a)*x**8 + 11*e**(2*a)*x**4 - 1)/(6*x**6*(e**(2*a)*x**4 + 1)
)
```

### 3.164 $\int \frac{\tanh^2(a+2 \log(x))}{x^9} dx$

Optimal result . . . . .	1250
Mathematica [A] (verified) . . . . .	1250
Rubi [A] (verified) . . . . .	1251
Maple [A] (verified) . . . . .	1252
Fricas [A] (verification not implemented) . . . . .	1253
Sympy [F] . . . . .	1253
Maxima [A] (verification not implemented) . . . . .	1253
Giac [A] (verification not implemented) . . . . .	1254
Mupad [B] (verification not implemented) . . . . .	1254
Reduce [B] (verification not implemented) . . . . .	1255

#### Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = -\frac{1}{8x^8} + \frac{e^{2a}}{x^4} + \frac{e^{4a}}{1 + e^{2a}x^4} + 8e^{4a} \log(x) - 2e^{4a} \log(1 + e^{2a}x^4)$$

output

```
-1/8/x^8+exp(2*a)/x^4+exp(4*a)/(1+exp(2*a)*x^4)+8*exp(4*a)*ln(x)-2*exp(4*a)*ln(1+exp(2*a)*x^4)
```

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.83

$$\begin{aligned} \int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = & -\frac{1}{8x^8} + \frac{\cosh(2a)}{x^4} + 8 \cosh(4a) \log(x) \\ & - 2 \cosh(4a) \log((1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)) \\ & + \frac{\sinh(2a)}{x^4} + \frac{\cosh(3a) + \sinh(3a)}{(1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)} \\ & + 8 \log(x) \sinh(4a) \\ & - 2 \log((1 + x^4) \cosh(a) + (-1 + x^4) \sinh(a)) \sinh(4a) \end{aligned}$$

input

```
Integrate[Tanh[a + 2*Log[x]]^2/x^9,x]
```

output

$$\begin{aligned} & -1/8*1/x^8 + \text{Cosh}[2*a]/x^4 + 8*\text{Cosh}[4*a]*\text{Log}[x] - 2*\text{Cosh}[4*a]*\text{Log}[(1 + x^4) \\ & )*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]] + \text{Sinh}[2*a]/x^4 + (\text{Cosh}[3*a] + \text{Sinh}[3*a])/ \\ & ((1 + x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]) + 8*\text{Log}[x]*\text{Sinh}[4*a] - 2*\text{Log}[(1 + \\ & x^4)*\text{Cosh}[a] + (-1 + x^4)*\text{Sinh}[a]]*\text{Sinh}[4*a] \end{aligned}$$

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.12, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6071, 948, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{(e^{2a}x^4 - 1)^2}{x^9 (e^{2a}x^4 + 1)^2} dx \\ & \quad \downarrow \text{948} \\ & \frac{1}{4} \int \frac{(1 - e^{2a}x^4)^2}{x^{12} (e^{2a}x^4 + 1)^2} dx^4 \\ & \quad \downarrow \text{99} \\ & \frac{1}{4} \int \left( \frac{8e^{4a}}{x^4} - \frac{4e^{2a}}{x^8} + \frac{1}{x^{12}} - \frac{8e^{6a}}{e^{2a}x^4 + 1} - \frac{4e^{6a}}{(e^{2a}x^4 + 1)^2} \right) dx^4 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \left( \frac{4e^{4a}}{e^{2a}x^4 + 1} + \frac{4e^{2a}}{x^4} + 8e^{4a} \log(x^4) - 8e^{4a} \log(e^{2a}x^4 + 1) - \frac{1}{2x^8} \right) \end{aligned}$$

input

$$\text{Int}[\text{Tanh}[a + 2*\text{Log}[x]]^2/x^9, x]$$

output

$$\begin{aligned} & (-1/2*1/x^8 + (4*E^{(2*a)})/x^4 + (4*E^{(4*a)})/(1 + E^{(2*a)}*x^4) + 8*E^{(4*a)}* \\ & \text{Log}[x^4] - 8*E^{(4*a)}*\text{Log}[1 + E^{(2*a)}*x^4])/4 \end{aligned}$$

## Definitions of rubi rules used

rule 99 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] | (GtQ[m, 0] && GeQ[n, -1]))`

rule 948 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.98

method	result	size
risch	$\frac{2e^{4a}x^8 + \frac{7e^{2a}x^4}{8} - \frac{1}{8}}{x^8(1+e^{2a}x^4)} + 8e^{4a} \ln(x) - 2e^{4a} \ln(1 + e^{2a}x^4)$	63

input `int(tanh(a+2*ln(x))^2/x^9,x,method=_RETURNVERBOSE)`

output `(2*exp(4*a)*x^8+7/8*exp(2*a)*x^4-1/8)/x^8/(1+exp(2*a)*x^4)+8*exp(4*a)*ln(x)-2*exp(4*a)*ln(1+exp(2*a)*x^4)`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.36

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx$$

$$= \frac{16 x^8 e^{(4a)} + 7 x^4 e^{(2a)} - 16 (x^{12} e^{(6a)} + x^8 e^{(4a)}) \log(x^4 e^{(2a)} + 1) + 64 (x^{12} e^{(6a)} + x^8 e^{(4a)}) \log(x) - 1}{8 (x^{12} e^{(2a)} + x^8)}$$

input `integrate(tanh(a+2*log(x))^2/x^9,x, algorithm="fricas")`

output `1/8*(16*x^8*e^(4*a) + 7*x^4*e^(2*a) - 16*(x^12*e^(6*a) + x^8*e^(4*a))*log(x^4*e^(2*a) + 1) + 64*(x^12*e^(6*a) + x^8*e^(4*a))*log(x) - 1)/(x^12*e^(2*a) + x^8)`

**Sympy [F]**

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx$$

input `integrate(tanh(a+2*ln(x))**2/x**9,x)`

output `Integral(tanh(a + 2*log(x))**2/x**9, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.70

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = -2 e^{(4a)} \log\left(\frac{1}{x^4} + e^{(2a)}\right) - \frac{e^{(6a)}}{\frac{1}{x^4} + e^{(2a)}} + \frac{e^{(2a)}}{x^4} - \frac{1}{8 x^8}$$

input `integrate(tanh(a+2*log(x))^2/x^9,x, algorithm="maxima")`

output  $-2e^{(4a)} \log(1/x^4 + e^{(2a)}) - e^{(6a)}/(1/x^4 + e^{(2a)}) + e^{(2a)}/x^4 - 1/8/x^8$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.28

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = -2e^{(4a)} \log(x^4 e^{(2a)} + 1) + 2e^{(4a)} \log(x^4) + \frac{2x^4 e^{(6a)} + 3e^{(4a)}}{x^4 e^{(2a)} + 1} - \frac{24x^8 e^{(4a)} - 8x^4 e^{(2a)} + 1}{8x^8}$$

input `integrate(tanh(a+2*log(x))^2/x^9,x, algorithm="giac")`

output  $-2e^{(4a)} \log(x^4 e^{(2a)} + 1) + 2e^{(4a)} \log(x^4) + (2x^4 e^{(6a)} + 3e^{(4a)})/(x^4 e^{(2a)} + 1) - 1/8*(24x^8 e^{(4a)} - 8x^4 e^{(2a)} + 1)/x^8$

### Mupad [B] (verification not implemented)

Time = 2.30 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.92

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx = 8e^{4a} \ln(x) - 2e^{4a} \ln(x^4 + e^{-2a}) + \frac{2e^{4a} x^8 + \frac{7e^{2a} x^4}{8} - \frac{1}{8}}{e^{2a} x^{12} + x^8}$$

input `int(tanh(a + 2*log(x))^2/x^9,x)`

output  $8*\exp(4*a)*\log(x) - 2*\exp(4*a)*\log(\exp(-2*a) + x^4) + ((7*x^4*\exp(2*a))/8 + 2*x^8*\exp(4*a) - 1/8)/(x^{12}*\exp(2*a) + x^8)$

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.84

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^9} dx$$

$$= \frac{-16e^{6a} \log(-e^{\frac{a}{2}} \sqrt{2}x + e^ax^2 + 1) x^{12} - 16e^{6a} \log(e^{\frac{a}{2}} \sqrt{2}x + e^ax^2 + 1) x^{12} + 64e^{6a} \log(x) x^{12} - 16e^{6a} x^{12}}{8x^8 (e^{2a}x^2 + 1)}$$

input `int(tanh(a+2*log(x))^2/x^9,x)`output `( - 16*e**(6*a)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**12 - 16*e**(6*a)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**12 + 64*e**(6*a)*log(x)*x**12 - 16*e**(6*a)*x**12 - 16*e**(4*a)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**8 - 16*e**(4*a)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**8 + 64*e**(4*a)*log(x)*x**8 + 7*e**(2*a)*x**4 - 1)/(8*x**8*(e**(2*a)*x**2 + 1))`



### 3.165 $\int x^4 \tanh^2(a + 2 \log(x)) dx$

Optimal result	1256
Mathematica [A] (verified)	1257
Rubi [A] (verified)	1257
Maple [C] (verified)	1261
Fricas [A] (verification not implemented)	1262
Sympy [F]	1262
Maxima [A] (verification not implemented)	1263
Giac [A] (verification not implemented)	1263
Mupad [B] (verification not implemented)	1264
Reduce [B] (verification not implemented)	1264

#### Optimal result

Integrand size = 13, antiderivative size = 146

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = -4e^{-2a}x + \frac{x^5}{5} - \frac{e^{-2a}x}{1 + e^{2a}x^4} - \frac{5e^{-5a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} + \frac{5e^{-5a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} + \frac{5e^{-5a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{2\sqrt{2}}$$

output

```
-4*x/exp(2*a)+1/5*x^5-x/exp(2*a)/(1+exp(2*a)*x^4)+5/4*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(5/2*a)+5/4*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(5/2*a)+5/4*arctanh(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)/exp(5/2*a)
```

**Mathematica [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.31

$$\int x^4 \tanh^2(a + 2 \log(x)) dx$$

$$= -4e^{-2a}x + \frac{x^5}{5} - \frac{e^{-2a}x}{1 + e^{2a}x^4} - \frac{5}{4}\sqrt[4]{-1}e^{-5a/2} \log(\sqrt[4]{-1}e^{-5a/2} - e^{-2a}x)$$

$$- \frac{5}{4}(-1)^{3/4}e^{-5a/2} \log((-1)^{3/4}e^{-5a/2} - e^{-2a}x) + \frac{5}{4}\sqrt[4]{-1}e^{-5a/2} \log(\sqrt[4]{-1}e^{-5a/2} + e^{-2a}x) + \frac{5}{4}(-1)^{3/4}e^{-5a/2} \log((-1)^{3/4}e^{-5a/2} + e^{-2a}x)$$

input `Integrate[x^4*Tanh[a + 2*Log[x]]^2,x]`

output 
$$\frac{(-4*x)}{E^{(2*a)}} + \frac{x^5}{5} - \frac{x}{(E^{(2*a)}*(1 + E^{(2*a)}*x^4))} - (5*(-1)^{(1/4)}*\text{Log}[(-1)^{(1/4)}/E^{((5*a)/2)} - x/E^{(2*a)}])/(4*E^{((5*a)/2)}) - (5*(-1)^{(3/4)}*\text{Log}[(-1)^{(3/4)}/E^{((5*a)/2)} - x/E^{(2*a)}])/(4*E^{((5*a)/2)}) + (5*(-1)^{(1/4)}*\text{Log}[(-1)^{(1/4)}/E^{((5*a)/2)} + x/E^{(2*a)}])/(4*E^{((5*a)/2)}) + (5*(-1)^{(3/4)}*\text{Log}[(-1)^{(3/4)}/E^{((5*a)/2)} + x/E^{(2*a)}])/(4*E^{((5*a)/2)})$$

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.47, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6071, 963, 27, 959, 843, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \tanh^2(a + 2 \log(x)) dx$$

$$\downarrow 6071$$

$$\int \frac{x^4(e^{2a}x^4 - 1)^2}{(e^{2a}x^4 + 1)^2} dx$$

$$\downarrow 963$$

$$\frac{x^5}{e^{2a}x^4 + 1} - \frac{1}{4}e^{-4a} \int \frac{4x^4(4e^{4a} - e^{6a}x^4)}{e^{2a}x^4 + 1} dx$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \int \frac{x^4(4e^{4a} - e^{6a}x^4)}{e^{2a}x^4 + 1} dx \\
& \downarrow 959 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \int \frac{x^4}{e^{2a}x^4 + 1} dx - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 843 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \int \frac{1}{e^{2a}x^4 + 1} dx \right) - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 755 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \int \frac{e^ax^2 + 1}{e^{2a}x^4 + 1} dx \right) \right) - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 1476 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) \right) + \frac{1}{2} \int \frac{1}{e^{2a}x^4 + 1} dx \right) - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 1082 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{1}{-(1 - \sqrt{2}e^{a/2}x)^2 - 1}} d(1 - \sqrt{2}e^{a/2}x) - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x)^2 - 1}} d(\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 217 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) - \frac{1}{5} e^{4a} x^5 \right) \\
& \downarrow 1479 \\
& \frac{x^5}{e^{2a}x^4 + 1} - e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int -\frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx - \frac{e^{-a/2} \int -\frac{\sqrt{2}(\sqrt{2}x + e^{-a/2})}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) - \frac{1}{5} e^{4a} x^5 \right)
\end{aligned}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{x^5}{e^{2a}x^4 + 1} - \\
 e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} + \frac{e^{-a/2} \int \frac{\sqrt{2}(\sqrt{2}x+e^{-a/2})}{x^2+\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\dots)}{\sqrt{\dots}} \right) \right) \right) \\
 & \downarrow 27 \\
 & \frac{x^5}{e^{2a}x^4 + 1} - \\
 e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} + \frac{1}{2} e^{-a/2} \int \frac{\sqrt{2}x + e^{-a/2}}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\dots)}{\sqrt{\dots}} \right) \right) \right) \\
 & \downarrow 1103 \\
 & \frac{x^5}{e^{2a}x^4 + 1} - \\
 e^{-4a} \left( 5e^{4a} \left( e^{-2a}x - e^{-2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{e^{-a/2} \log(e^a x)}{\sqrt{\dots}} \right) \right) \right)
 \end{aligned}$$

input `Int[x^4*Tanh[a + 2*Log[x]]^2,x]`

output `x^5/(1 + E^(2*a)*x^4) - (-1/5*(E^(4*a)*x^5) + 5*E^(4*a)*(x/E^(2*a) - ((-ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))/2 + (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/2)/E^(4*a)`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1}] \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 755  $\text{Int}[(a_ + (b_ \cdot)(x_ )^4)^{-1}, x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2 \cdot r) \ \text{Int}[(r - s \cdot x^2)/(a + b \cdot x^4), x], x] + \text{Simp}[1/(2 \cdot r) \ \text{Int}[(r + s \cdot x^2)/(a + b \cdot x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 843  $\text{Int}[(c_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^n)^p, x\_Symbol] \rightarrow \text{Simp}[c^{n-1} \cdot (c \cdot x)^{m-n+1} \cdot ((a + b \cdot x^n)^{p+1} / (b \cdot (m+n \cdot p+1))), x] - \text{Simp}[a \cdot c^{n-1} \cdot (m-n+1) / (b \cdot (m+n \cdot p+1)) \ \text{Int}[(c \cdot x)^{m-n} \cdot (a + b \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n \cdot p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 959  $\text{Int}[(e_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^n)^p \cdot ((c_ ) + (d_ \cdot)(x_ )^n), x\_Symbol] \rightarrow \text{Simp}[d \cdot (e \cdot x)^{m+1} \cdot ((a + b \cdot x^n)^{p+1} / (b \cdot e \cdot (m+n \cdot (p+1)+1))), x] - \text{Simp}[(a \cdot d \cdot (m+1) - b \cdot c \cdot (m+n \cdot (p+1)+1)) / (b \cdot (m+n \cdot (p+1)+1)) \ \text{Int}[(e \cdot x)^m \cdot (a + b \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[m+n \cdot (p+1)+1, 0]$

rule 963  $\text{Int}[(e_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^n)^p \cdot ((c_ ) + (d_ \cdot)(x_ )^n)^2, x\_Symbol] \rightarrow \text{Simp}[(-b \cdot c - a \cdot d)^2 \cdot (e \cdot x)^{m+1} \cdot ((a + b \cdot x^n)^{p+1} / (a \cdot b^2 \cdot e \cdot n \cdot (p+1))), x] + \text{Simp}[1/(a \cdot b^2 \cdot n \cdot (p+1)) \ \text{Int}[(e \cdot x)^m \cdot (a + b \cdot x^n)^{p+1} \cdot \text{Simp}[(b \cdot c - a \cdot d)^2 \cdot (m+1) + b^2 \cdot c^2 \cdot n \cdot (p+1) + a \cdot b \cdot d^2 \cdot n \cdot (p+1) \cdot x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1]$

rule 1082  $\text{Int}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot S \ \text{imply}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4 \cdot a \cdot c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S  
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,  
e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[  
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[  
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]  
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[  
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],  
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F  
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 6071 `Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol]  
:= Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p,  
x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.49

method	result	size
risch	$\frac{e^{2a}e^{-2a}x^5}{5} - 4e^{-2a}x - \frac{xe^{-2a}}{1+e^{2a}x^4} + \frac{5e^{-4a}}{4} \left( \sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-\frac{R}{e^{2a}})}{-R^3} \right)$	71

input `int(x^4*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)`

output `1/5*exp(2*a)*exp(-2*a)*x^5-4*exp(-2*a)*x-x*exp(-2*a)/(1+exp(2*a)*x^4)+5/4*  
exp(-4*a)*sum(1/_R^3*ln(x-_R),_R=RootOf(exp(2*a)*_Z^4+1))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.21

$$\int x^4 \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{8 x^9 e^{(4a)} - 152 x^5 e^{(2a)} + 50 \sqrt{2} (x^4 e^{(2a)} + 1) \arctan \left( \sqrt{2} x e^{(\frac{1}{2} a)} + 1 \right) e^{(-\frac{1}{2} a)} + 50 \sqrt{2} (x^4 e^{(2a)} + 1) \arctan \left( \sqrt{2} x e^{(\frac{1}{2} a)} - 1 \right) e^{(-\frac{1}{2} a)} + 25 \sqrt{2} (x^4 e^{(2a)} + 1) e^{(-\frac{1}{2} a)} \log(x^2 e^a + \sqrt{2} x e^{(1/2 a)} + 1) - 25 \sqrt{2} (x^4 e^{(2a)} + 1) e^{(-\frac{1}{2} a)} \log(x^2 e^a - \sqrt{2} x e^{(1/2 a)} + 1) - 200 x}{x^4 e^{(4a)} + e^{(2a)}}$$

input `integrate(x^4*tanh(a+2*log(x))^2,x, algorithm="fricas")`

output

```
1/40*(8*x^9*e^(4*a) - 152*x^5*e^(2*a) + 50*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(-1/2*a) + 50*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(-1/2*a) + 25*sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 25*sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) - 200*x)/(x^4*e^(4*a) + e^(2*a))
```

**Sympy [F]**

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = \int x^4 \tanh^2(a + 2 \log(x)) dx$$

input `integrate(x**4*tanh(a+2*ln(x))**2,x)`

output

```
Integral(x**4*tanh(a + 2*log(x))**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.13

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = \frac{1}{5} (x^5 e^{(2a)} - 20x) e^{(-2a)} + \frac{5}{8} \left( 2\sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a + \sqrt{2}e^{\frac{1}{2}a}) \right) e^{(-\frac{1}{2}a)} + 2\sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a - \sqrt{2}e^{\frac{1}{2}a}) \right) e^{(-\frac{1}{2}a)} - \frac{x}{x^4 e^{(4a)} + e^{(2a)}} \right)$$

input `integrate(x^4*tanh(a+2*log(x))^2,x, algorithm="maxima")`

output

```
1/5*(x^5*e^(2*a) - 20*x)*e^(-2*a) + 5/8*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x
*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-1/2*a) + 2*sqrt(2)*arctan(1/2*sq
rt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*e^(-1/2*a)) + sqrt(2)*e^(-1
/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - sqrt(2)*e^(-1/2*a)*log(x^2*
e^a - sqrt(2)*x*e^(1/2*a) + 1))*e^(-2*a) - x/(x^4*e^(4*a) + e^(2*a))
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.09

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = \frac{5}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} + 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{5}{2}a)} + \frac{5}{4} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} - 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{5}{2}a)} + \frac{5}{8} \sqrt{2} e^{(-\frac{5}{2}a)} \log \left( \sqrt{2}x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right) - \frac{5}{8} \sqrt{2} e^{(-\frac{5}{2}a)} \log \left( -\sqrt{2}x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right) + \frac{1}{5} (x^5 e^{(20a)} - 20x e^{(18a)}) e^{(-20a)} - \frac{x e^{(-2a)}}{x^4 e^{(2a)} + 1}$$

input `integrate(x^4*tanh(a+2*log(x))^2,x, algorithm="giac")`



output

```
5/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-5/2*a) + 5/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(-5/2*a) + 5/8*sqrt(2)*e^(-5/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - 5/8*sqrt(2)*e^(-5/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/5*(x^5*e^(20*a) - 20*x*e^(18*a))*e^(-20*a) - x*e^(-2*a)/(x^4*e^(2*a) + 1)
```

**Mupad [B] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.52

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = \frac{x^5}{5} - \frac{5 \operatorname{atan}\left(x (-e^{2a})^{1/4}\right)}{2 (-e^{2a})^{5/4}} - \frac{x}{e^{4a} x^4 + e^{2a}} - 4 x e^{-2a} + \frac{\operatorname{atan}\left(x (-e^{2a})^{1/4} \operatorname{li}\right) 5i}{2 (-e^{2a})^{5/4}}$$

input

```
int(x^4*tanh(a + 2*log(x))^2,x)
```

output

```
(atan(x*(-exp(2*a))^(1/4)*1i)*5i)/(2*(-exp(2*a))^(5/4)) - (5*atan(x*(-exp(2*a))^(1/4)))/(2*(-exp(2*a))^(5/4)) - x/(exp(2*a) + x^4*exp(4*a)) + x^5/5 - 4*x*exp(-2*a)
```

**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.21

$$\int x^4 \tanh^2(a + 2 \log(x)) dx = \frac{-50e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 - 50e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) + 50e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 + 50e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right)}{2}$$

input

```
int(x^4*tanh(a+2*log(x))^2,x)
```

output

```
( - 50*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 - 50*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) + 50*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 + 50*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) - 25*e**((5*a)/2)*sqrt(2)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + 25*e**((5*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - 25*e**(a/2)*sqrt(2)*log( - e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 25*e**(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 8*e**(5*a)*x**9 - 152*e**(3*a)*x**5 - 200*e**a*x)/(40*e**(3*a)*(e**(2*a)*x**4 + 1))
```

### 3.166 $\int x^2 \tanh^2(a + 2 \log(x)) dx$

Optimal result	1266
Mathematica [A] (verified)	1266
Rubi [A] (verified)	1267
Maple [C] (verified)	1271
Fricas [A] (verification not implemented)	1271
Sympy [F]	1272
Maxima [A] (verification not implemented)	1272
Giac [A] (verification not implemented)	1273
Mupad [B] (verification not implemented)	1274
Reduce [B] (verification not implemented)	1274

#### Optimal result

Integrand size = 13, antiderivative size = 134

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \frac{x^3}{3} + \frac{x^3}{1 + e^{2a}x^4} + \frac{3e^{-3a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{-3a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} + \frac{3e^{-3a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{2\sqrt{2}}$$

output

```
1/3*x^3+x^3/(1+exp(2*a)*x^4)-3/4*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/e
xp(3/2*a)-3/4*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(3/2*a)+3/4*arctan
h(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)/exp(3/2*a)
```

#### Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.30

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \frac{1}{12} \left( 4x^3 + \frac{12x^3}{1 + e^{2a}x^4} + 9(-1)^{3/4} e^{-3a/2} \log(\sqrt[4]{-1} e^{-3a/2} - e^{-a}x) + 9\sqrt[4]{-1} e^{-3a/2} \log((-1)^{3/4} e^{-3a/2} - e^{-a}x) - 9(-1)^{3/4} e^{-3a/2} \log(\dots) \right)$$

input `Integrate[x^2*Tanh[a + 2*Log[x]]^2,x]`

output 
$$\frac{(4x^3 + (12x^3)/(1 + E^{(2a)x^4}) + (9(-1)^{3/4} \text{Log}[-1^{1/4}/E^{(3a)/2} - x/E^a])/E^{(3a)/2} + (9(-1)^{1/4} \text{Log}[-1^{3/4}/E^{(3a)/2} - x/E^a])/E^{(3a)/2} - (9(-1)^{3/4} \text{Log}[-1^{1/4}/E^{(3a)/2} + x/E^a])/E^{(3a)/2} - (9(-1)^{1/4} \text{Log}[-1^{3/4}/E^{(3a)/2} + x/E^a])/E^{(3a)/2})}{12}$$

### Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.56, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.923$ , Rules used = {6071, 963, 27, 959, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \tanh^2(a + 2 \log(x)) dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{x^2 (e^{2a} x^4 - 1)^2}{(e^{2a} x^4 + 1)^2} dx \\ & \quad \downarrow \text{963} \\ & \frac{x^3}{e^{2a} x^4 + 1} - \frac{1}{4} e^{-4a} \int \frac{4x^2 (2e^{4a} - e^{6a} x^4)}{e^{2a} x^4 + 1} dx \\ & \quad \downarrow \text{27} \\ & \frac{x^3}{e^{2a} x^4 + 1} - e^{-4a} \int \frac{x^2 (2e^{4a} - e^{6a} x^4)}{e^{2a} x^4 + 1} dx \\ & \quad \downarrow \text{959} \\ & \frac{x^3}{e^{2a} x^4 + 1} - e^{-4a} \left( 3e^{4a} \int \frac{x^2}{e^{2a} x^4 + 1} dx - \frac{1}{3} e^{4a} x^3 \right) \\ & \quad \downarrow \text{826} \end{aligned}$$

$$\frac{x^3}{e^{2a}x^4 + 1} - e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \int \frac{e^ax^2 + 1}{e^{2a}x^4 + 1} dx - \frac{1}{2}e^{-a} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 1476

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \left( \frac{1}{2}e^{-a} \int \frac{1}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx + \frac{1}{2}e^{-a} \int \frac{1}{x^2 + \sqrt{2}e^{-a/2}x + e^{-a}} dx \right) - \frac{1}{2}e^{-a} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 1082

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \left( \frac{e^{-a/2} \int \frac{1}{-(1-\sqrt{2}e^{a/2}x)^2 - 1} d(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2}e^{a/2}x + 1)^2 - 1} d(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} \right) - \frac{1}{2}e^{-a} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 217

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2}e^{-a} \int \frac{1 - e^ax^2}{e^{2a}x^4 + 1} dx \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 1479

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2}e^{-a} \left( -\frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 25

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2}e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2}e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) \right) - \frac{1}{3}e^{4a}x^3 \right)$$

↓ 27

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2a}{x^2 - \sqrt{2}e^{-a/2}x + 1} dx}{2\sqrt{2}} \right) \right) \right)$$

↓ 1103

$$e^{-4a} \left( 3e^{4a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \log(e^ax^2 + \sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} \right) \right) \right)$$

input `Int[x^2*Tanh[a + 2*Log[x]]^2,x]`

output `x^3/(1 + E^(2*a)*x^4) - (-1/3*(E^(4*a)*x^3) + 3*E^(4*a)*((-ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))/(2*E^a) - (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/(2*E^a))/E^(4*a)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 826  $\text{Int}[(x\_)^2/((a\_)+(b\_)*(x\_)^4), x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 959  $\text{Int}[(e_)*(x_)^{m_}*((a_)+(b_)*(x_)^{n_})^{p_}*((c_)+(d_)*(x_)^{n_})], x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{m+1}*((a + b*x^n)^{p+1}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 963  $\text{Int}[(e_)*(x_)^{m_}*((a_)+(b_)*(x_)^{n_})^{p_}*((c_)+(d_)*(x_)^{n_})^2], x\_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)^2*(e*x)^{m+1}*((a + b*x^n)^{p+1}/(a*b^2*e*n*(p+1))), x] + \text{Simp}[1/(a*b^2*n*(p+1)) \text{Int}[(e*x)^m*(a + b*x^n)^{p+1}* \text{Simp}[(b*c - a*d)^2*(m+1) + b^2*c^2*n*(p+1) + a*b*d^2*n*(p+1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[p, -1]$

rule 1082  $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2]^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476  $\text{Int}[(d_)+(e_)*(x_)^2/((a_)+(c_)*(x_)^4), x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6071

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.31 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.40

method	result	size
risch	$\frac{x^3}{3} + \frac{x^3}{1+e^{2a}x^4} - \frac{3e^{-2a} \left( \sum_{R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-R)}{-R} \right)}{4}$	53

input

```
int(x^2*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*x^3+x^3/(1+exp(2*a)*x^4)-3/4*exp(-2*a)*sum(1/_R*ln(x-_R),_R=RootOf(exp(2*a)*_Z^4+1))
```

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.26

$$\int x^2 \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{8x^7e^{(3a)} + 32x^3e^a - 18\sqrt{2}(x^4e^{(2a)} + 1) \arctan\left(\sqrt{2}xe^{(\frac{1}{2}a)} + 1\right) e^{(-\frac{1}{2}a)} - 18\sqrt{2}(x^4e^{(2a)} + 1) \arctan\left(\sqrt{2}xe^{(\frac{1}{2}a)} - 1\right) e^{(-\frac{1}{2}a)}}{4}$$

input

```
integrate(x^2*tanh(a+2*log(x))^2,x, algorithm="fricas")
```



output

```
1/24*(8*x^7*e^(3*a) + 32*x^3*e^a - 18*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(-1/2*a) - 18*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(-1/2*a) + 9*sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 9*sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1))/(x^4*e^(3*a) + e^a)
```

**Sympy [F]**

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \int x^2 \tanh^2(a + 2 \log(x)) dx$$

input

```
integrate(x**2*tanh(a+2*ln(x))**2,x)
```

output

```
Integral(x**2*tanh(a + 2*log(x))**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.07

$$\begin{aligned} \int x^2 \tanh^2(a + 2 \log(x)) dx = & \frac{1}{3} x^3 \\ & - \frac{3}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2 x e^a + \sqrt{2} e^{\frac{1}{2} a}) e^{(-\frac{1}{2} a)} \right) e^{(-\frac{3}{2} a)} \\ & - \frac{3}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2 x e^a - \sqrt{2} e^{\frac{1}{2} a}) e^{(-\frac{1}{2} a)} \right) e^{(-\frac{3}{2} a)} \\ & + \frac{3}{8} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( x^2 e^a + \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) \\ & - \frac{3}{8} \sqrt{2} e^{(-\frac{3}{2} a)} \log \left( x^2 e^a - \sqrt{2} x e^{\frac{1}{2} a} + 1 \right) + \frac{x^3}{x^4 e^{(2a)} + 1} \end{aligned}$$

input

```
integrate(x^2*tanh(a+2*log(x))^2,x, algorithm="maxima")
```

output

```
1/3*x^3 - 3/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) - 3/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) + 3/8*sqrt(2)*e^(-3/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 3/8*sqrt(2)*e^(-3/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + x^3/(x^4*e^(2*a) + 1)
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.04

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \frac{1}{3} x^3 - \frac{3}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} e^{(-\frac{1}{2}a)} + 2x) e^{\frac{1}{2}a}\right) e^{(-\frac{3}{2}a)} - \frac{3}{4} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2} (\sqrt{2} e^{(-\frac{1}{2}a)} - 2x) e^{\frac{1}{2}a}\right) e^{(-\frac{3}{2}a)} + \frac{3}{8} \sqrt{2} e^{(-\frac{3}{2}a)} \log\left(\sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}\right) - \frac{3}{8} \sqrt{2} e^{(-\frac{3}{2}a)} \log\left(-\sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}\right) + \frac{x^3}{x^4 e^{(2a)} + 1}$$

input

```
integrate(x^2*tanh(a+2*log(x))^2,x, algorithm="giac")
```

output

```
1/3*x^3 - 3/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-3/2*a) - 3/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(-3/2*a) + 3/8*sqrt(2)*e^(-3/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - 3/8*sqrt(2)*e^(-3/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + x^3/(x^4*e^(2*a) + 1)
```

**Mupad [B] (verification not implemented)**

Time = 2.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.50

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \frac{x^3}{e^{2a} x^4 + 1} + \frac{3 \operatorname{atan}\left(x (-e^{2a})^{1/4}\right)}{2 (-e^{2a})^{3/4}} + \frac{x^3}{3} + \frac{\operatorname{atan}\left(x (-e^{2a})^{1/4} \operatorname{li}\right) 3i}{2 (-e^{2a})^{3/4}}$$

input `int(x^2*tanh(a + 2*log(x))^2,x)`output `x^3/(x^4*exp(2*a) + 1) + (3*atan(x*(-exp(2*a))^(1/4)))/(2*(-exp(2*a))^(3/4)) + (atan(x*(-exp(2*a))^(1/4)*1i)*3i)/(2*(-exp(2*a))^(3/4)) + x^3/3`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.37

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \frac{18e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 + 18e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) - 18e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 - 18e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right)}{2}$$

input `int(x^2*tanh(a+2*log(x))^2,x)`output `(18*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 + 18*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) - 18*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 - 18*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) - 9*e**((5*a)/2)*sqrt(2)*log(-e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + 9*e**((5*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - 9*e**(a/2)*sqrt(2)*log(-e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 9*e**(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 8*e**(4*a)*x**7 + 32*e**(2*a)*x**3)/(24*e**(2*a)*(e**(2*a)*x**4 + 1))`

### 3.167 $\int \tanh^2(a + 2 \log(x)) dx$

Optimal result	1275
Mathematica [A] (verified)	1275
Rubi [A] (verified)	1276
Maple [C] (verified)	1277
Fricas [A] (verification not implemented)	1278
Sympy [F]	1278
Maxima [A] (verification not implemented)	1279
Giac [A] (verification not implemented)	1279
Mupad [B] (verification not implemented)	1280
Reduce [B] (verification not implemented)	1280

#### Optimal result

Integrand size = 9, antiderivative size = 126

$$\int \tanh^2(a + 2 \log(x)) dx = x + \frac{x}{1 + e^{2a}x^4} + \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{e^{-a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{e^{-a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{2\sqrt{2}}$$

output

```
x+x/(1+exp(2*a)*x^4)-1/4*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(1/2*a)
)-1/4*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)/exp(1/2*a)-1/4*arctanh(2^(1/2)
)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)/exp(1/2*a)
```

#### Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.16

$$\int \tanh^2(a + 2 \log(x)) dx = \frac{1}{4} \left( 4x + \frac{4x}{1 + e^{2a}x^4} + \sqrt[4]{-1}e^{-a/2} \log(\sqrt[4]{-1}e^{-a/2} - x) + (-1)^{3/4}e^{-a/2} \log((-1)^{3/4}e^{-a/2} - x) - \sqrt[4]{-1}e^{-a/2} \log(\sqrt[4]{-1}e^{-a/2} + x) - (-1)^{3/4}e^{-a/2} \log((-1)^{3/4}e^{-a/2} + x) \right)$$

input

```
Integrate[Tanh[a + 2*Log[x]]^2,x]
```

output

$$\frac{(4x + (4x)/(1 + E^{(2a)}x^4) + ((-1)^{(1/4)}\text{Log}[(-1)^{(1/4)}/E^{(a/2)} - x])/E^{(a/2)} + ((-1)^{(3/4)}\text{Log}[(-1)^{(3/4)}/E^{(a/2)} - x])/E^{(a/2)} - ((-1)^{(1/4)}\text{Log}[(-1)^{(1/4)}/E^{(a/2)} + x])/E^{(a/2)} - ((-1)^{(3/4)}\text{Log}[(-1)^{(3/4)}/E^{(a/2)} + x])/E^{(a/2)})/4$$

**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.31, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6067, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tanh^2(a + 2 \log(x)) dx \\ & \quad \downarrow \text{6067} \\ & \int \frac{(e^{2a}x^4 - 1)^2}{(e^{2a}x^4 + 1)^2} dx \\ & \quad \downarrow \text{915} \\ & \int \left( 1 - \frac{4e^{2a}x^4}{(e^{2a}x^4 + 1)^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{2\sqrt{2}} + \frac{x}{e^{2a}x^4 + 1} + \\ & \frac{e^{-a/2} \log(e^ax^2 - \sqrt{2}e^{a/2}x + 1)}{4\sqrt{2}} - \frac{e^{-a/2} \log(e^ax^2 + \sqrt{2}e^{a/2}x + 1)}{4\sqrt{2}} + x \end{aligned}$$

input

```
Int[Tanh[a + 2*Log[x]]^2,x]
```

output

$$x + x/(1 + E^{(2a)}x^4) + \text{ArcTan}[1 - \text{Sqrt}[2]*E^{(a/2)}x]/(2*\text{Sqrt}[2]*E^{(a/2)}) - \text{ArcTan}[1 + \text{Sqrt}[2]*E^{(a/2)}x]/(2*\text{Sqrt}[2]*E^{(a/2)}) + \text{Log}[1 - \text{Sqrt}[2]*E^{(a/2)}x + E^ax^2]/(4*\text{Sqrt}[2]*E^{(a/2)}) - \text{Log}[1 + \text{Sqrt}[2]*E^{(a/2)}x + E^ax^2]/(4*\text{Sqrt}[2]*E^{(a/2)})$$

## Definitions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.37

method	result	size
risch	$x + \frac{x}{1+e^{2a}x^4} - \frac{e^{-2a} \left( \sum_{R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-R)}{-R^3} \right)}{4}$	47

input `int(tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)`

output `x+x/(1+exp(2*a)*x^4)-1/4*exp(-2*a)*sum(1/_R^3*ln(x-_R),_R=RootOf(exp(2*a)*
_Z^4+1))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.29

$$\int \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{8x^5e^{2a} - 2\sqrt{2}(x^4e^{2a} + 1) \arctan\left(\sqrt{2}xe^{\frac{1}{2}a} + 1\right) e^{(-\frac{1}{2}a)} - 2\sqrt{2}(x^4e^{2a} + 1) \arctan\left(\sqrt{2}xe^{\frac{1}{2}a} - 1\right) e^{(-\frac{1}{2}a)} + 16x}{(x^4e^{2a} + 1)^2}$$

input `integrate(tanh(a+2*log(x))^2,x, algorithm="fricas")`

output

```
1/8*(8*x^5*e^(2*a) - 2*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a)
+ 1)*e^(-1/2*a) - 2*sqrt(2)*(x^4*e^(2*a) + 1)*arctan(sqrt(2)*x*e^(1/2*a)
- 1)*e^(-1/2*a) - sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a + sqrt
(2)*x*e^(1/2*a) + 1) + sqrt(2)*(x^4*e^(2*a) + 1)*e^(-1/2*a)*log(x^2*e^a -
sqrt(2)*x*e^(1/2*a) + 1) + 16*x)/(x^4*e^(2*a) + 1)
```

**Sympy [F]**

$$\int \tanh^2(a + 2 \log(x)) dx = \int \tanh^2(a + 2 \log(x)) dx$$

input `integrate(tanh(a+2*ln(x))**2,x)`

output

`Integral(tanh(a + 2*log(x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.10

$$\int \tanh^2(a + 2 \log(x)) dx = -\frac{1}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a + \sqrt{2}e^{\frac{1}{2}a}) e^{(-\frac{1}{2}a)} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (2xe^a - \sqrt{2}e^{\frac{1}{2}a}) e^{(-\frac{1}{2}a)} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{8} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( x^2 e^a + \sqrt{2} x e^{\frac{1}{2}a} + 1 \right)$$

$$+ \frac{1}{8} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( x^2 e^a - \sqrt{2} x e^{\frac{1}{2}a} + 1 \right) + x + \frac{x}{x^4 e^{(2a)} + 1}$$

input `integrate(tanh(a+2*log(x))^2,x, algorithm="maxima")`

output

```
-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*
e^(-1/2*a) - 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*
e^(-1/2*a))*e^(-1/2*a) - 1/8*sqrt(2)*e^(-1/2*a)*log(x^2*e^a + sqrt(2)*x*e^
(1/2*a) + 1) + 1/8*sqrt(2)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) +
1) + x + x/(x^4*e^(2*a) + 1)
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.06

$$\int \tanh^2(a + 2 \log(x)) dx = -\frac{1}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} + 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{4} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2}e^{(-\frac{1}{2}a)} - 2x) e^{\frac{1}{2}a} \right) e^{(-\frac{1}{2}a)}$$

$$- \frac{1}{8} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( \sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right)$$

$$+ \frac{1}{8} \sqrt{2} e^{(-\frac{1}{2}a)} \log \left( -\sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right)$$

$$+ x + \frac{x}{x^4 e^{(2a)} + 1}$$

input `integrate(tanh(a+2*log(x))^2,x, algorithm="giac")`



output

```
-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/8*sqrt(2)*e^(-1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/8*sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + x + x/(x^4*e^(2*a) + 1)
```

**Mupad [B] (verification not implemented)**

Time = 2.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.48

$$\int \tanh^2(a + 2 \log(x)) dx = x - \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{2(-e^{2a})^{1/4}} + \frac{x}{e^{2a}x^4 + 1} + \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4} \operatorname{li}\right)}{2(-e^{2a})^{1/4}}$$

input

```
int(tanh(a + 2*log(x))^2,x)
```

output

```
x - atan(x*(-exp(2*a))^(1/4))/(2*(-exp(2*a))^(1/4)) + (atan(x*(-exp(2*a))^(1/4)*1i)*1i)/(2*(-exp(2*a))^(1/4)) + x/(x^4*exp(2*a) + 1)
```

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 309, normalized size of antiderivative = 2.45

$$\int \tanh^2(a + 2 \log(x)) dx$$

$$= \frac{2e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 + 2e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) - 2e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^4 - 2e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right)}{4}$$

input

```
int(tanh(a+2*log(x))^2,x)
```

output

```
(2*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 + 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2))) - 2*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**4 - 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2))) + e**((5*a)/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 - e**((5*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**4 + e**(a/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) - e**(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1) + 8*e**(3*a)*x**5 + 16*e**a*x)/(8*e**a*(e**(2*a)*x**4 + 1))
```

### 3.168 $\int \frac{\tanh^2(a+2\log(x))}{x^2} dx$

Optimal result	1282
Mathematica [A] (verified)	1282
Rubi [A] (verified)	1283
Maple [C] (verified)	1287
Fricas [A] (verification not implemented)	1287
Sympy [F]	1288
Maxima [A] (verification not implemented)	1288
Giac [A] (verification not implemented)	1289
Mupad [B] (verification not implemented)	1289
Reduce [B] (verification not implemented)	1290

#### Optimal result

Integrand size = 13, antiderivative size = 138

$$\int \frac{\tanh^2(a + 2\log(x))}{x^2} dx = -\frac{1}{x} - \frac{e^{2a}x^3}{1 + e^{2a}x^4} + \frac{e^{a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{e^{a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} + \frac{e^{a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{2\sqrt{2}}$$

output

```
-1/x-exp(2*a)*x^3/(1+exp(2*a)*x^4)-1/4*exp(1/2*a)*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)-1/4*exp(1/2*a)*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)+1/4*exp(1/2*a)*arctanh(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.31

$$\int \frac{\tanh^2(a + 2\log(x))}{x^2} dx = \frac{1}{4} \left( -\frac{4}{x} - \frac{4}{\frac{e^{-2a}}{x^3} + x} + (-1)^{3/4} e^{a/2} \log\left(\frac{e^{-2a}(\sqrt[4]{-1} - e^{a/2}x)}{x^4}\right) + \sqrt[4]{-1} e^{a/2} \log\left(\frac{e^{-2a}((-1)^{3/4} - e^{a/2}x)}{x^4}\right) - (-1)^{3/4} e^{a/2} \log\left(\frac{e^{-2a}(\sqrt[4]{-1} + e^{a/2}x)}{x^4}\right) - \sqrt[4]{-1} e^{a/2} \log\left(\frac{e^{-2a}((-1)^{3/4} + e^{a/2}x)}{x^4}\right) \right)$$

input `Integrate[Tanh[a + 2*Log[x]]^2/x^2,x]`

output  $(-4/x - 4/(1/(E^{(2*a)*x^3}) + x) + (-1)^{(3/4)}*E^{(a/2)}*Log[((-1)^{(1/4)} - E^{(a/2)*x})/(E^{(2*a)*x^4})] + (-1)^{(1/4)}*E^{(a/2)}*Log[((-1)^{(3/4)} - E^{(a/2)*x})/(E^{(2*a)*x^4})] - (-1)^{(3/4)}*E^{(a/2)}*Log[((-1)^{(1/4)} + E^{(a/2)*x})/(E^{(2*a)*x^4})] - (-1)^{(1/4)}*E^{(a/2)}*Log[((-1)^{(3/4)} + E^{(a/2)*x})/(E^{(2*a)*x^4})])/4$

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.54, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.923$ , Rules used = {6071, 962, 25, 957, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx \\
 & \quad \downarrow 6071 \\
 & \int \frac{(e^{2a}x^4 - 1)^2}{x^2 (e^{2a}x^4 + 1)^2} dx \\
 & \quad \downarrow 962 \\
 & \int -\frac{x^2(7e^{2a} - e^{4a}x^4)}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{x(e^{2a}x^4 + 1)} \\
 & \quad \downarrow 25 \\
 & -\int \frac{x^2(7e^{2a} - e^{4a}x^4)}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{x(e^{2a}x^4 + 1)} \\
 & \quad \downarrow 957 \\
 & -e^{2a} \int \frac{x^2}{e^{2a}x^4 + 1} dx - \frac{1}{x(e^{2a}x^4 + 1)} - \frac{2e^{2a}x^3}{e^{2a}x^4 + 1} \\
 & \quad \downarrow 826 \\
 & -e^{2a} \left( \frac{1}{2} e^{-a} \int \frac{e^a x^2 + 1}{e^{2a}x^4 + 1} dx - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a}x^4 + 1} dx \right) - \frac{1}{x(e^{2a}x^4 + 1)} - \frac{2e^{2a}x^3}{e^{2a}x^4 + 1}
 \end{aligned}$$

↓ 1476

$$-e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx \right) - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) - \frac{1}{x(e^{2a} x^4 + 1)} - \frac{2e^{2a} x^3}{e^{2a} x^4 + 1}$$

↓ 1082

$$-e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{1}{-(1 - \sqrt{2} e^{a/2} x)^2 - 1} d(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2} e^{a/2} x + 1)^2 - 1} d(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) - \frac{1}{x(e^{2a} x^4 + 1)} - \frac{2e^{2a} x^3}{e^{2a} x^4 + 1}$$

↓ 217

$$-e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) - \frac{1}{x(e^{2a} x^4 + 1)} - \frac{2e^{2a} x^3}{e^{2a} x^4 + 1}$$

↓ 1479

$$-e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( -\frac{e^{-a/2} \int -\frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) \right) - \frac{1}{x(e^{2a} x^4 + 1)} - \frac{2e^{2a} x^3}{e^{2a} x^4 + 1}$$

↓ 25

$$-e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) \right) - \frac{1}{x(e^{2a} x^4 + 1)} - \frac{2e^{2a} x^3}{e^{2a} x^4 + 1}$$

↓ 27

$$\begin{aligned}
& -e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2} - 2x}{x^2 - \sqrt{2}e^{-a/2}x + e^{-a}} dx}{2\sqrt{2}} \right) \right. \\
& \quad \left. \frac{1}{x(e^{2a}x^4 + 1)} - \frac{2e^{2a}x^3}{e^{2a}x^4 + 1} \right) \\
& \quad \downarrow \text{1103} \\
& -e^{2a} \left( \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) - \frac{1}{2} e^{-a} \left( \frac{e^{-a/2} \log(e^ax^2 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} \right) \right. \\
& \quad \left. \frac{1}{x(e^{2a}x^4 + 1)} - \frac{2e^{2a}x^3}{e^{2a}x^4 + 1} \right)
\end{aligned}$$

input `Int[Tanh[a + 2*Log[x]]^2/x^2,x]`

output `-(1/(x*(1 + E^(2*a)*x^4))) - (2*E^(2*a)*x^3)/(1 + E^(2*a)*x^4) - E^(2*a)*(-ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))/(2*E^a) - (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/(2*E^a)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 826  $\text{Int}[(x\_)^2/((a\_)+(b\_)*(x\_)^4), x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \text{ Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \text{ Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 957  $\text{Int}[(e\_)*(x\_)^{(m\_)}*((a\_)+(b\_)*(x\_)^{(n\_)})^{(p\_)}*((c\_)+(d\_)*(x\_)^{(n\_)}), x\_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1))/(a*b*e*n*(p+1))}, x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[p, -1] \&\& ((! \text{IntegerQ}[p + 1/2] \&\& \text{NeQ}[p, -5/4]) \parallel ! \text{RationalQ}[m] \parallel (\text{IGtQ}[n, 0] \&\& \text{ILtQ}[p + 1/2, 0] \&\& \text{LeQ}[-1, m, (-n)*(p+1)]))$

rule 962  $\text{Int}[(e\_)*(x\_)^{(m\_)}*((a\_)+(b\_)*(x\_)^{(n\_)})^{(p\_)}*((c\_)+(d\_)*(x\_)^{(n\_)})^2, x\_Symbol] \rightarrow \text{Simp}[c^2*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1))/(a*e*(m+1))}, x] - \text{Simp}[1/(a*e^n*(m+1)) \text{ Int}[(e*x)^{(m+n)}*(a + b*x^n)^p * \text{Simp}[b*c^2*n*(p+1) + c*(b*c - 2*a*d)*(m+1) - a*(m+1)*d^2*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[m, -1] \&\& \text{GtQ}[n, 0]$

rule 1082  $\text{Int}[(a\_)+(b\_)*(x\_)+(c\_)*(x\_)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel ! \text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d\_)+(e\_)*(x\_)/((a\_)+(b\_)*(x\_)+(c\_)*(x\_)^2), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1476  $\text{Int}[(d\_)+(e\_)*(x\_)^2/((a\_)+(c\_)*(x\_)^4), x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{ Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6071

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.29 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.46

method	result	size
risch	$\frac{-2e^{2a}x^4 - 1}{x(1 + e^{2a}x^4)} + \frac{\sum_{R=\text{RootOf}(\_Z^4 + e^{2a})} -R \ln((5\_R^4 + 4e^{2a})x + \_R^3)}{4}$	64

input

```
int(tanh(a+2*ln(x))^2/x^2,x,method=_RETURNVERBOSE)
```

output

```
(-2*exp(2*a)*x^4-1)/x/(1+exp(2*a)*x^4)+1/4*sum(_R*ln((5*_R^4+4*exp(2*a))*x+_R^3),_R=RootOf(_Z^4+exp(2*a)))
```

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.17

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx = \frac{16x^4 e^{(2a)} + 2\sqrt{2}(x^5 e^{(2a)} + x) \arctan\left(\sqrt{2}x e^{(\frac{1}{2}a)} + 1\right) e^{(\frac{1}{2}a)} + 2\sqrt{2}(x^5 e^{(2a)} + x) \arctan\left(\sqrt{2}x e^{(\frac{1}{2}a)} - 1\right) e^{(\frac{1}{2}a)}}{x^2}$$

input

```
integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="fricas")
```



output

```
-1/8*(16*x^4*e^(2*a) + 2*sqrt(2)*(x^5*e^(2*a) + x)*arctan(sqrt(2)*x*e^(1/2*a) + 1)*e^(1/2*a) + 2*sqrt(2)*(x^5*e^(2*a) + x)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(1/2*a) - sqrt(2)*(x^5*e^(2*a) + x)*e^(1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) + sqrt(2)*(x^5*e^(2*a) + x)*e^(1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + 8)/(x^5*e^(2*a) + x)
```

**Sympy [F]**

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx$$

input

```
integrate(tanh(a+2*ln(x))**2/x**2,x)
```

output

```
Integral(tanh(a + 2*log(x))**2/x**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.06

$$\begin{aligned} \int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx &= \frac{1}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} + \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{\frac{1}{2}a} \\ &+ \frac{1}{4} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} - \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{\frac{1}{2}a} \\ &+ \frac{1}{8} \sqrt{2} e^{\frac{1}{2}a} \log \left( \frac{\sqrt{2} e^{\frac{1}{2}a}}{x} + \frac{1}{x^2} + e^a \right) \\ &- \frac{1}{8} \sqrt{2} e^{\frac{1}{2}a} \log \left( -\frac{\sqrt{2} e^{\frac{1}{2}a}}{x} + \frac{1}{x^2} + e^a \right) \\ &- \frac{1}{x} - \frac{e^{2a}}{x \left( \frac{1}{x^4} + e^{2a} \right)} \end{aligned}$$

input

```
integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="maxima")
```

output

```
1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) + 2/x)*e^(-1/2*a))*e^(1/2*a) + 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) - 2/x)*e^(-1/2*a))*e^(1/2*a) + 1/8*sqrt(2)*e^(1/2*a)*log(sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) - 1/8*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) - 1/x - e^(2*a)/(x*(1/x^4 + e^(2*a)))
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.04

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx = -\frac{1}{4} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{(-\frac{1}{2}a)} + 2x \right) e^{\frac{1}{2}a} \right) e^{\frac{1}{2}a} - \frac{1}{4} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{(-\frac{1}{2}a)} - 2x \right) e^{\frac{1}{2}a} \right) e^{\frac{1}{2}a} + \frac{1}{8} \sqrt{2} e^{\frac{1}{2}a} \log \left( \sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right) - \frac{1}{8} \sqrt{2} e^{\frac{1}{2}a} \log \left( -\sqrt{2} x e^{(-\frac{1}{2}a)} + x^2 + e^{(-a)} \right) - \frac{2x^4 e^{(2a)} + 1}{x^5 e^{(2a)} + x}$$

input

```
integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="giac")
```

output

```
-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(1/2*a) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(1/2*a) + 1/8*sqrt(2)*e^(1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - 1/8*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - (2*x^4*e^(2*a) + 1)/(x^5*e^(2*a) + x)
```

**Mupad [B] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.49

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx = \frac{\operatorname{atanh} \left( x (-e^{2a})^{1/4} \right) (-e^{2a})^{1/4}}{2} - \frac{\operatorname{atan} \left( x (-e^{2a})^{1/4} \right) (-e^{2a})^{1/4}}{2} - \frac{2e^{2a} x^4 + 1}{e^{2a} x^5 + x}$$

input `int(tanh(a + 2*log(x))^2/x^2,x)`

output `(atanh(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4))/2 - (atan(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4))/2 - (2*x^4*exp(2*a) + 1)/(x + x^5*exp(2*a))`

### Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.22

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx$$

$$= \frac{2e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^5 + 2e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x - 2e^{\frac{5a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x^5 - 2e^{\frac{a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^a x}{e^{\frac{a}{2}} \sqrt{2}}\right) x}{(x + x^5 \exp(2a))}$$

input `int(tanh(a+2*log(x))^2/x^2,x)`

output `(2*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**5 + 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x - 2*e**((5*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**5 - 2*e**(a/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x - e**((5*a)/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**5 + e**((5*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**5 - e**(a/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x + e**(a/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x - 16*e**(-2*a)*x**4 - 8)/(8*x*(e**(2*a)*x**4 + 1))`

### 3.169 $\int \frac{\tanh^2(a+2 \log(x))}{x^4} dx$

Optimal result	1291
Mathematica [A] (verified)	1291
Rubi [A] (verified)	1292
Maple [C] (verified)	1296
Fricas [A] (verification not implemented)	1296
Sympy [F]	1297
Maxima [A] (verification not implemented)	1297
Giac [A] (verification not implemented)	1298
Mupad [B] (verification not implemented)	1298
Reduce [B] (verification not implemented)	1299

#### Optimal result

Integrand size = 13, antiderivative size = 138

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx = -\frac{1}{3x^3} - \frac{e^{2a}x}{1 + e^{2a}x^4} + \frac{3e^{3a/2} \arctan(1 - \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{3a/2} \arctan(1 + \sqrt{2}e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{3a/2} \operatorname{arctanh}\left(\frac{\sqrt{2}e^{a/2}x}{1+e^ax^2}\right)}{2\sqrt{2}}$$

output

```
-1/3/x^3-exp(2*a)*x/(1+exp(2*a)*x^4)-3/4*exp(3/2*a)*arctan(-1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)-3/4*exp(3/2*a)*arctan(1+2^(1/2)*exp(1/2*a)*x)*2^(1/2)-3/4*exp(3/2*a)*arctanh(2^(1/2)*exp(1/2*a)*x/(1+exp(a)*x^2))*2^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.37

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx = \frac{1}{12} \left( -\frac{4}{x^3} - \frac{12e^{2a}x}{1 + e^{2a}x^4} + 9\sqrt[4]{-1}e^{3a/2} \log\left(\frac{e^{-2a}(\sqrt[4]{-1} - e^{a/2}x)}{x^4}\right) + 9(-1)^{3/4}e^{3a/2} \log\left(\frac{e^{-2a}((-1)^{3/4} - e^{a/2}x)}{x^4}\right) - 9\sqrt[4]{-1}e^{3a/2} \log\left(\frac{e^{-2a}((-1)^{3/4} - e^{a/2}x)}{x^4}\right) \right)$$

input `Integrate[Tanh[a + 2*Log[x]]^2/x^4,x]`

output 
$$\frac{(-4/x^3 - (12E^{(2a)x})/(1 + E^{(2a)x^4}) + 9(-1)^{1/4}E^{(3a)/2} \text{Log}[(-1)^{1/4} - E^{(a/2)x}]/(E^{(2a)x^4}) + 9(-1)^{3/4}E^{(3a)/2} \text{Log}[(-1)^{3/4} - E^{(a/2)x}]/(E^{(2a)x^4}) - 9(-1)^{1/4}E^{(3a)/2} \text{Log}[(-1)^{1/4} + E^{(a/2)x}]/(E^{(2a)x^4}) - 9(-1)^{3/4}E^{(3a)/2} \text{Log}[(-1)^{3/4} + E^{(a/2)x}]/(E^{(2a)x^4})]/12$$

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.51, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.923$ , Rules used = {6071, 962, 25, 910, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx \\ & \quad \downarrow \text{6071} \\ & \int \frac{(e^{2a}x^4 - 1)^2}{x^4 (e^{2a}x^4 + 1)^2} dx \\ & \quad \downarrow \text{962} \\ & \frac{1}{3} \int -\frac{13e^{2a} - 3e^{4a}x^4}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{3x^3 (e^{2a}x^4 + 1)} \\ & \quad \downarrow \text{25} \\ & -\frac{1}{3} \int \frac{13e^{2a} - 3e^{4a}x^4}{(e^{2a}x^4 + 1)^2} dx - \frac{1}{3x^3 (e^{2a}x^4 + 1)} \\ & \quad \downarrow \text{910} \\ & \frac{1}{3} \left( -9e^{2a} \int \frac{1}{e^{2a}x^4 + 1} dx - \frac{4e^{2a}x}{e^{2a}x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a}x^4 + 1)} \\ & \quad \downarrow \text{755} \end{aligned}$$

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \int \frac{e^a x^2 + 1}{e^{2a} x^4 + 1} dx \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 1476

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \left( \frac{1}{2} e^{-a} \int \frac{1}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx + \frac{1}{2} e^{-a} \int \frac{1}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx \right) + \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 1082

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{1}{-(1 - \sqrt{2} e^{a/2} x)^2 - 1} d(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} - \frac{e^{-a/2} \int \frac{1}{-(\sqrt{2} e^{a/2} x + 1)^2 - 1} d(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} \right) \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 217

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \int \frac{1 - e^a x^2}{e^{2a} x^4 + 1} dx + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1 - \sqrt{2} e^{a/2} x)}{\sqrt{2}} \right) \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 1479

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \left( -\frac{e^{-a/2} \int \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} - \frac{e^{-a/2} \int \frac{\sqrt{2}(\sqrt{2} x + e^{-a/2})}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} \right) \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 25

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2} e^{-a/2} - 2x}{x^2 - \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} + \frac{e^{-a/2} \int \frac{\sqrt{2}(\sqrt{2} x + e^{-a/2})}{x^2 + \sqrt{2} e^{-a/2} x + e^{-a}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2} e^{a/2} x + 1)}{\sqrt{2}} \right) \right) - \frac{4e^{2a} x}{e^{2a} x^4 + 1} \right) - \frac{1}{3x^3 (e^{2a} x^4 + 1)}$$

↓ 27

$$\frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \int \frac{\sqrt{2}e^{-a/2}-2x}{x^2-\sqrt{2}e^{-a/2}x+e^{-a}} dx}{2\sqrt{2}} + \frac{1}{2} e^{-a/2} \int \frac{\sqrt{2}x+e^{-a/2}}{x^2+\sqrt{2}e^{-a/2}x+e^{-a}} dx \right) + \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) \right) \right. \\ \left. \frac{1}{3x^3(e^{2a}x^4+1)} \right) \\ \downarrow 1103 \\ \frac{1}{3} \left( -9e^{2a} \left( \frac{1}{2} \left( \frac{e^{-a/2} \arctan(\sqrt{2}e^{a/2}x+1)}{\sqrt{2}} - \frac{e^{-a/2} \arctan(1-\sqrt{2}e^{a/2}x)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{e^{-a/2} \log(e^ax^2+\sqrt{2}e^{a/2}x+1)}{2\sqrt{2}} \right) \right) \right) \\ \left. \frac{1}{3x^3(e^{2a}x^4+1)} \right)$$

input `Int[Tanh[a + 2*Log[x]]^2/x^4,x]`

output `-1/3*1/(x^3*(1 + E^(2*a)*x^4)) + ((-4*E^(2*a)*x)/(1 + E^(2*a)*x^4) - 9*E^(2*a)*((-ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2))) + ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^(a/2)))/2 + (-1/2*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(Sqrt[2]*E^(a/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^(a/2)))/2)/3`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 910 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(- (b*c - a*d))*x*((a + b*x^n)^(p + 1)/(a*b*n*(p + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)) Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])`

rule 962 `Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^2, x_Symbol] := Simp[c^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))), x] - Simp[1/(a*e^n*(m + 1)) Int[(e*x)^(m + n)*(a + b*x^n)^p*Simp[b*c^2*n*(p + 1) + c*(b*c - 2*a*d)*(m + 1) - a*(m + 1)*d^2*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1] && GtQ[n, 0]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`



rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 6071

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p], x] /; FreeQ[{a, b, d, e, m, p}, x]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.32 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.49

method	result	size
risch	$\frac{-\frac{4}{3}e^{2a}x^4 - \frac{1}{3}}{x^3(1+e^{2a}x^4)} + \frac{3 \left( \sum_{R=\text{RootOf}(e^{6a}+_Z^4)} -R \ln((-4e^{6a}-5_R^4)x -_R e^{4a}) \right)}{4}$	68

input

```
int(tanh(a+2*ln(x))^2/x^4,x,method=_RETURNVERBOSE)
```

output

```
(-4/3*exp(2*a)*x^4-1/3)/x^3/(1+exp(2*a)*x^4)+3/4*sum(_R*ln((-4*exp(6*a)-5*_R^4)*x-_R*exp(4*a)),_R=RootOf(exp(6*a)+_Z^4))
```

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.33

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx =$$

$$\frac{32 x^4 e^{(2a)} + 18 \sqrt{2} (x^7 e^{(3a)} + x^3 e^a) \arctan \left( \sqrt{2} x e^{(\frac{1}{2} a)} + 1 \right) e^{(\frac{1}{2} a)} + 18 \sqrt{2} (x^7 e^{(3a)} + x^3 e^a) \arctan \left( \sqrt{2} x e^{(\frac{1}{2} a)} + 1 \right) e^{(\frac{1}{2} a)}}{x^4}$$

input

```
integrate(tanh(a+2*log(x))^2/x^4,x, algorithm="fricas")
```

output

```
-1/24*(32*x^4*e^(2*a) + 18*sqrt(2)*(x^7*e^(3*a) + x^3*e^a)*arctan(sqrt(2)*
x*e^(1/2*a) + 1)*e^(1/2*a) + 18*sqrt(2)*(x^7*e^(3*a) + x^3*e^a)*arctan(sqrt(2)*x*e^(1/2*a) - 1)*e^(1/2*a) + 9*sqrt(2)*(x^7*e^(3*a) + x^3*e^a)*e^(1/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 9*sqrt(2)*(x^7*e^(3*a) + x^3*e^a)*e^(1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + 8)/(x^7*e^(2*a) + x^3)
```

**Sympy [F]**

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx$$

input

```
integrate(tanh(a+2*ln(x))**2/x**4,x)
```

output

```
Integral(tanh(a + 2*log(x))**2/x**4, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.10

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx$$

$$= \frac{3}{8} \left( 2\sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} + \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{-\frac{1}{2}a} + 2\sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} \left( \sqrt{2} e^{\frac{1}{2}a} - \frac{2}{x} \right) e^{-\frac{1}{2}a} \right) e^{-\frac{1}{2}a} \right) - \frac{1}{3x^3} - \frac{e^{(2a)}}{x^3 \left( \frac{1}{x^4} + e^{(2a)} \right)}$$

input

```
integrate(tanh(a+2*log(x))^2/x^4,x, algorithm="maxima")
```

output

```
3/8*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) + 2/x)*e^(-1/2*a))*e^(-1/2*a) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) - 2/x)*e^(-1/2*a))*e^(-1/2*a) - sqrt(2)*e^(-1/2*a)*log(sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a))*e^(2*a) - 1/3/x^3 - e^(2*a)/(x^3*(1/x^4 + e^(2*a)))
```

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.03

$$\int \frac{\tanh^2(a + 2\log(x))}{x^4} dx = -\frac{3}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{(-\frac{1}{2}a)} + 2x\right)e^{(\frac{1}{2}a)}\right)e^{(\frac{3}{2}a)}$$

$$-\frac{3}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{(-\frac{1}{2}a)} - 2x\right)e^{(\frac{1}{2}a)}\right)e^{(\frac{3}{2}a)}$$

$$-\frac{3}{8}\sqrt{2}e^{(\frac{3}{2}a)}\log\left(\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}\right)$$

$$+\frac{3}{8}\sqrt{2}e^{(\frac{3}{2}a)}\log\left(-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}\right)$$

$$-\frac{xe^{(2a)}}{x^4e^{(2a)} + 1} - \frac{1}{3x^3}$$

input `integrate(tanh(a+2*log(x))^2/x^4,x, algorithm="giac")`output `-3/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(3/2*a) - 3/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(3/2*a) - 3/8*sqrt(2)*e^(3/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 3/8*sqrt(2)*e^(3/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - x*e^(2*a)/(x^4*e^(2*a) + 1) - 1/3/x^3`**Mupad [B] (verification not implemented)**

Time = 2.23 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.51

$$\int \frac{\tanh^2(a + 2\log(x))}{x^4} dx = \frac{3\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{3/4}}{2} - \frac{\frac{4e^{2a}x^4}{3} + \frac{1}{3}}{e^{2a}x^7 + x^3}$$

$$+ \frac{3\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{3/4}}{2}$$

input `int(tanh(a + 2*log(x))^2/x^4,x)`output `(3*atan(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(3/4))/2 - ((4*x^4*exp(2*a))/3 + 1/3)/(x^7*exp(2*a) + x^3) + (3*atanh(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(3/4))/2`

**Reduce [B] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 316, normalized size of antiderivative = 2.29

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^4} dx$$

$$= \frac{18e^{\frac{7a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^ax}{e^{\frac{a}{2}} \sqrt{2}}\right) x^7 + 18e^{\frac{3a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} - 2e^ax}{e^{\frac{a}{2}} \sqrt{2}}\right) x^3 - 18e^{\frac{7a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^ax}{e^{\frac{a}{2}} \sqrt{2}}\right) x^7 - 18e^{\frac{3a}{2}} \sqrt{2} \operatorname{atan}\left(\frac{e^{\frac{a}{2}} \sqrt{2} + 2e^ax}{e^{\frac{a}{2}} \sqrt{2}}\right) x^3}{x^4}$$

input `int(tanh(a+2*log(x))^2/x^4,x)`output

```
(18*e**((7*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**7 + 18*e**((3*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) - 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**3 - 18*e**((7*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**7 - 18*e**((3*a)/2)*sqrt(2)*atan((e**(a/2)*sqrt(2) + 2*e**a*x)/(e**(a/2)*sqrt(2)))*x**3 + 9*e**((7*a)/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**7 - 9*e**((7*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**7 + 9*e**((3*a)/2)*sqrt(2)*log(- e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**3 - 9*e**((3*a)/2)*sqrt(2)*log(e**(a/2)*sqrt(2)*x + e**a*x**2 + 1)*x**3 - 32*e**(2*a)*x**4 - 8)/(24*x**3*(e**(2*a)*x**4 + 1))
```

### 3.170 $\int (ex)^m \tanh(a + 2 \log(x)) dx$

Optimal result	1300
Mathematica [A] (verified)	1300
Rubi [A] (verified)	1301
Maple [F]	1302
Fricas [F]	1302
Sympy [F]	1303
Maxima [F]	1303
Giac [F]	1303
Mupad [F(-1)]	1304
Reduce [F]	1304

#### Optimal result

Integrand size = 13, antiderivative size = 60

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \frac{(ex)^{1+m}}{e(1+m)} - \frac{2(ex)^{1+m} \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -e^{2a}x^4\right)}{e(1+m)}$$

output

```
(e*x)^(1+m)/e/(1+m)-2*(e*x)^(1+m)*hypergeom([1, 1/4+1/4*m],[5/4+1/4*m],-exp(2*a)*x^4)/e/(1+m)
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.78

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \frac{x(ex)^m (-1 + 2 \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -x^4(\cosh(2a) + \sinh(2a))\right))}{1+m}$$

input

```
Integrate[(e*x)^m*Tanh[a + 2*Log[x]],x]
```

output

```

-((x*(e*x)^m*(-1 + 2*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(x^4*(Cos
h[2*a] + Sinh[2*a]))]))/(1 + m))

```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ex)^m \tanh(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{(e^{2a}x^4 - 1)(ex)^m}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{(ex)^{m+1}}{e(m+1)} - 2 \int \frac{(ex)^m}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{888} \\
 & \frac{(ex)^{m+1}}{e(m+1)} - \frac{2(ex)^{m+1} \text{Hypergeometric2F1}\left(1, \frac{m+1}{4}, \frac{m+5}{4}, -e^{2a}x^4\right)}{e(m+1)}
 \end{aligned}$$

input

```

Int[(e*x)^m*Tanh[a + 2*Log[x]],x]

```

output

```

(e*x)^(1 + m)/(e*(1 + m)) - (2*(e*x)^(1 + m)*Hypergeometric2F1[1, (1 + m)/
4, (5 + m)/4, -(E^(2*a)*x^4)]/(e*(1 + m))

```

## Definitions of rubi rules used

rule 888 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 959 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6071 `Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]`

## Maple [F]

$$\int (ex)^m \tanh(a + 2 \ln(x)) dx$$

input `int((e*x)^m*tanh(a+2*ln(x)),x)`

output `int((e*x)^m*tanh(a+2*ln(x)),x)`

## Fricas [F]

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x)) dx$$

input `integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="fricas")`

output `integral((e*x)^m*tanh(a + 2*log(x)), x)`

### Sympy [F]

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x)) dx$$

input `integrate((e*x)**m*tanh(a+2*ln(x)),x)`

output `Integral((e*x)**m*tanh(a + 2*log(x)), x)`

### Maxima [F]

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x)) dx$$

input `integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="maxima")`

output `integrate((e*x)^m*tanh(a + 2*log(x)), x)`

### Giac [F]

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x)) dx$$

input `integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="giac")`

output `integrate((e*x)^m*tanh(a + 2*log(x)), x)`



**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int \tanh(a + 2 \ln(x)) (ex)^m dx$$

input `int(tanh(a + 2*log(x))*(e*x)^m,x)`output `int(tanh(a + 2*log(x))*(e*x)^m, x)`**Reduce [F]**

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \frac{e^m (x^m x - 2(\int \frac{x^m}{e^{2a}x^4+1} dx) m - 2(\int \frac{x^m}{e^{2a}x^4+1} dx))}{m + 1}$$

input `int((e*x)^m*tanh(a+2*log(x)),x)`output `(e**m*(x**m*x - 2*int(x**m/(e**(2*a)*x**4 + 1),x)*m - 2*int(x**m/(e**(2*a)*x**4 + 1),x)))/(m + 1)`

### 3.171 $\int (ex)^m \tanh^2(a + 2 \log(x)) dx$

Optimal result	1305
Mathematica [A] (verified)	1305
Rubi [A] (verified)	1306
Maple [F]	1308
Fricas [F]	1308
Sympy [F]	1308
Maxima [F]	1309
Giac [F]	1309
Mupad [F(-1)]	1309
Reduce [F]	1310

#### Optimal result

Integrand size = 15, antiderivative size = 79

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \frac{(ex)^{1+m}}{e(1+m)} + \frac{(ex)^{1+m}}{e(1+e^{2a}x^4)} - \frac{(ex)^{1+m} \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -e^{2a}x^4\right)}{e}$$

output

```
(e*x)^(1+m)/e/(1+m)+(e*x)^(1+m)/e/(1+exp(2*a)*x^4)-(e*x)^(1+m)*hypergeom([
1, 1/4+1/4*m],[5/4+1/4*m],[-exp(2*a)*x^4)/e
```

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \frac{x(ex)^m (-1 + 4 \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -x^4(\cosh(2a) + \sinh(2a))\right) - 4 \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -x^4\right)}{1+m}$$

input

```
Integrate[(e*x)^m*Tanh[a + 2*Log[x]]^2,x]
```

output

```

-((x*(e*x)^m*(-1 + 4*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(x^4*(Cos
h[2*a] + Sinh[2*a]))] - 4*Hypergeometric2F1[2, (1 + m)/4, (5 + m)/4, -(x^4
*(Cosh[2*a] + Sinh[2*a]))]))/(1 + m)

```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.23, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6071, 963, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ex)^m \tanh^2(a + 2 \log(x)) dx \\
 & \quad \downarrow \text{6071} \\
 & \int \frac{(e^{2a}x^4 - 1)^2 (ex)^m}{(e^{2a}x^4 + 1)^2} dx \\
 & \quad \downarrow \text{963} \\
 & \frac{(ex)^{m+1}}{e(e^{2a}x^4 + 1)} - \frac{1}{4}e^{-4a} \int \frac{4(ex)^m (e^{4a}m - e^{6a}x^4)}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{(ex)^{m+1}}{e(e^{2a}x^4 + 1)} - e^{-4a} \int \frac{(ex)^m (e^{4a}m - e^{6a}x^4)}{e^{2a}x^4 + 1} dx \\
 & \quad \downarrow \text{959} \\
 & \frac{(ex)^{m+1}}{e(e^{2a}x^4 + 1)} - e^{-4a} \left( e^{4a}(m+1) \int \frac{(ex)^m}{e^{2a}x^4 + 1} dx - \frac{e^{4a}(ex)^{m+1}}{e(m+1)} \right) \\
 & \quad \downarrow \text{888} \\
 & \frac{(ex)^{m+1}}{e(e^{2a}x^4 + 1)} - e^{-4a} \left( \frac{e^{4a}(ex)^{m+1} \text{Hypergeometric2F1}\left(1, \frac{m+1}{4}, \frac{m+5}{4}, -e^{2a}x^4\right)}{e} - \frac{e^{4a}(ex)^{m+1}}{e(m+1)} \right)
 \end{aligned}$$

input

```

Int[(e*x)^m*Tanh[a + 2*Log[x]]^2,x]

```

output

$$\frac{(e^x)^{(1+m)} / (e \cdot (1 + E^{(2a) \cdot x^4})) - (-(E^{(4a)} \cdot (e^x)^{(1+m)}) / (e \cdot (1+m))) + (E^{(4a)} \cdot (e^x)^{(1+m)} \cdot \text{Hypergeometric2F1}[1, (1+m)/4, (5+m)/4, -(E^{(2a)} \cdot x^4)]) / e}{E^{(4a)}}$$
**Defintions of rubi rules used**

rule 27

$$\text{Int}[(a_*) \cdot (F x_*), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F x, x], x] \text{ ; FreeQ}[a, x] \&\& \text{ !MatchQ}[F x, (b_*) \cdot (G x_*) \text{ ; FreeQ}[b, x]$$

rule 888

$$\text{Int}[((c_*) \cdot (x_*))^{(m_*)} \cdot ((a_*) + (b_*) \cdot (x_*)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p \cdot ((c \cdot x)^{(m+1)} / (c \cdot (m+1))) \cdot \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b) \cdot (x^n/a)], x] \text{ ; FreeQ}[\{a, b, c, m, n, p\}, x] \&\& \text{ !IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] \text{ || GtQ}[a, 0])$$

rule 959

$$\text{Int}[((e_*) \cdot (x_*))^{(m_*)} \cdot ((a_*) + (b_*) \cdot (x_*)^{(n_*)})^{(p_*)} \cdot ((c_*) + (d_*) \cdot (x_*)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d \cdot (e^x)^{(m+1)} \cdot ((a + b \cdot x^n)^{(p+1)} / (b \cdot e \cdot (m + n \cdot (p+1) + 1))), x] - \text{Simp}[(a \cdot d \cdot (m+1) - b \cdot c \cdot (m + n \cdot (p+1) + 1)) / (b \cdot (m + n \cdot (p+1) + 1)) \quad \text{Int}[(e^x)^m \cdot (a + b \cdot x^n)^p, x], x] \text{ ; FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \&\& \text{ NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{ NeQ}[m + n \cdot (p+1) + 1, 0]$$

rule 963

$$\text{Int}[((e_*) \cdot (x_*))^{(m_*)} \cdot ((a_*) + (b_*) \cdot (x_*)^{(n_*)})^{(p_*)} \cdot ((c_*) + (d_*) \cdot (x_*)^{(n_*)})^2, x\_Symbol] \rightarrow \text{Simp}[(-b \cdot c - a \cdot d)^2 \cdot (e^x)^{(m+1)} \cdot ((a + b \cdot x^n)^{(p+1)} / (a \cdot b^2 \cdot e \cdot n \cdot (p+1))), x] + \text{Simp}[1 / (a \cdot b^2 \cdot n \cdot (p+1)) \quad \text{Int}[(e^x)^m \cdot (a + b \cdot x^n)^{(p+1)} \cdot \text{Simp}[(b \cdot c - a \cdot d)^2 \cdot (m+1) + b^2 \cdot c^2 \cdot n \cdot (p+1) + a \cdot b \cdot d^2 \cdot n \cdot (p+1) \cdot x^n, x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, m, n\}, x] \&\& \text{ NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{ IGtQ}[n, 0] \&\& \text{ LtQ}[p, -1]$$

rule 6071

$$\text{Int}[((e_*) \cdot (x_*))^{(m_*)} \cdot \text{Tanh}[(a_*) + \text{Log}[x_*] \cdot (b_*)] \cdot (d_*)^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e^x)^m \cdot ((-1 + E^{(2 \cdot a \cdot d)} \cdot x^{(2 \cdot b \cdot d)})^p / (1 + E^{(2 \cdot a \cdot d)} \cdot x^{(2 \cdot b \cdot d)}))^p, x] \text{ ; FreeQ}[\{a, b, d, e, m, p\}, x]$$

**Maple [F]**

$$\int (ex)^m \tanh(a + 2 \ln(x))^2 dx$$

input `int((e*x)^m*tanh(a+2*ln(x))^2,x)`

output `int((e*x)^m*tanh(a+2*ln(x))^2,x)`

**Fricas [F]**

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^2 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="fricas")`

output `integral((e*x)^m*tanh(a + 2*log(x))^2, x)`

**Sympy [F]**

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int (ex)^m \tanh^2(a + 2 \log(x)) dx$$

input `integrate((e*x)**m*tanh(a+2*ln(x))**2,x)`

output `Integral((e*x)**m*tanh(a + 2*log(x))**2, x)`

**Maxima [F]**

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^2 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="maxima")`

output `integrate((e*x)^m*tanh(a + 2*log(x))^2, x)`

**Giac [F]**

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^2 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="giac")`

output `integrate((e*x)^m*tanh(a + 2*log(x))^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int \tanh(a + 2 \ln(x))^2 (e x)^m dx$$

input `int(tanh(a + 2*log(x))^2*(e*x)^m,x)`

output `int(tanh(a + 2*log(x))^2*(e*x)^m, x)`



### 3.172 $\int (ex)^m \tanh^3(a + 2 \log(x)) dx$

Optimal result	1311
Mathematica [A] (verified)	1311
Rubi [A] (verified)	1312
Maple [F]	1315
Fricas [F]	1315
Sympy [F]	1315
Maxima [F]	1316
Giac [F]	1316
Mupad [F(-1)]	1316
Reduce [F]	1317

#### Optimal result

Integrand size = 15, antiderivative size = 125

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \frac{(ex)^{1+m}}{e(1+m)} - \frac{(ex)^{1+m}}{e(1+e^{2a}x^4)^2} + \frac{(5+m)(ex)^{1+m}}{4e(1+e^{2a}x^4)} - \frac{(9+2m+m^2)(ex)^{1+m} \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -e^{2a}x^4\right)}{4e(1+m)}$$

output

```
(e*x)^(1+m)/e/(1+m)-(e*x)^(1+m)/e/(1+exp(2*a)*x^4)^2+1/4*(5+m)*(e*x)^(1+m)/e/(1+exp(2*a)*x^4)-1/4*(m^2+2*m+9)*(e*x)^(1+m)*hypergeom([1, 1/4+1/4*m],[5/4+1/4*m],-exp(2*a)*x^4)/e/(1+m)
```

#### Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.89

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \frac{x(ex)^m (-1 + 6 \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -x^4(\cosh(2a) + \sinh(2a))\right) - 12 \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{4}, \frac{5+m}{4}, -x^4\right)}{4e(1+m)}$$



input `Integrate[(e*x)^m*Tanh[a + 2*Log[x]]^3,x]`

output 
$$-\left(\frac{x(e^x)^m(-1 + 6\text{Hypergeometric2F1}[1, (1 + m)/4, (5 + m)/4, -(x^4(\text{Cosh}[2a] + \text{Sinh}[2a]))]) - 12\text{Hypergeometric2F1}[2, (1 + m)/4, (5 + m)/4, -(x^4(\text{Cosh}[2a] + \text{Sinh}[2a]))]) + 8\text{Hypergeometric2F1}[3, (1 + m)/4, (5 + m)/4, -(x^4(\text{Cosh}[2a] + \text{Sinh}[2a]))])}{(1 + m)}\right)$$

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.58, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {6071, 968, 27, 1047, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (ex)^m \tanh^3(a + 2 \log(x)) dx \\ & \quad \downarrow 6071 \\ & \int \frac{(e^{2a}x^4 - 1)^3 (ex)^m}{(e^{2a}x^4 + 1)^3} dx \\ & \quad \downarrow 968 \\ & -\frac{1}{8}e^{-2a} \int \frac{2(ex)^m (1 - e^{2a}x^4) (e^{4a}(m + 5)x^4 + e^{2a}(3 - m))}{(e^{2a}x^4 + 1)^2} dx - \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e (e^{2a}x^4 + 1)^2} \\ & \quad \downarrow 27 \\ & -\frac{1}{4}e^{-2a} \int \frac{(ex)^m (1 - e^{2a}x^4) (e^{4a}(m + 5)x^4 + e^{2a}(3 - m))}{(e^{2a}x^4 + 1)^2} dx - \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e (e^{2a}x^4 + 1)^2} \\ & \quad \downarrow 1047 \\ & -\frac{1}{4}e^{-2a} \left( \frac{(e^{4a}(m + 5)x^4 + e^{2a}(3 - m)) (ex)^{m+1}}{2e (e^{2a}x^4 + 1)} - \frac{1}{4}e^{-2a} \int -\frac{2(ex)^m (e^{4a}(1 - m)(3 - m) - e^{6a}(m + 3)(m + 5))}{e^{2a}x^4 + 1} \right. \\ & \quad \left. \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e (e^{2a}x^4 + 1)^2} \right) \end{aligned}$$

↓ 27

$$-\frac{1}{4}e^{-2a} \left( \frac{1}{2}e^{-2a} \int \frac{(ex)^m (e^{4a}(1-m)(3-m) - e^{6a}(m+3)(m+5)x^4)}{e^{2a}x^4 + 1} dx + \frac{(e^{4a}(m+5)x^4 + e^{2a}(3-m))(ex)^m}{2e(e^{2a}x^4 + 1)} \right. \\ \left. \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e(e^{2a}x^4 + 1)^2} \right)$$

↓ 959

$$-\frac{1}{4}e^{-2a} \left( \frac{1}{2}e^{-2a} \left( 2e^{4a}(m^2 + 2m + 9) \int \frac{(ex)^m}{e^{2a}x^4 + 1} dx - \frac{e^{4a}(m+3)(m+5)(ex)^{m+1}}{e(m+1)} \right) + \frac{(e^{4a}(m+5)x^4 + e^{2a}(3-m))(ex)^m}{2e(e^{2a}x^4 + 1)} \right. \\ \left. \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e(e^{2a}x^4 + 1)^2} \right)$$

↓ 888

$$-\frac{1}{4}e^{-2a} \left( \frac{1}{2}e^{-2a} \left( \frac{2e^{4a}(m^2 + 2m + 9)(ex)^{m+1} \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{4}, \frac{m+5}{4}, -e^{2a}x^4\right)}{e(m+1)} - \frac{e^{4a}(m+3)(m+5)(ex)^m}{e(m+1)} \right) \right. \\ \left. \frac{(1 - e^{2a}x^4)^2 (ex)^{m+1}}{4e(e^{2a}x^4 + 1)^2} \right)$$

input `Int[(e*x)^m*Tanh[a + 2*Log[x]]^3,x]`

output `-1/4*((e*x)^(1+m)*(1 - E^(2*a)*x^4)^2)/(e*(1 + E^(2*a)*x^4)^2) - (((e*x)^(1+m)*(E^(2*a)*(3 - m) + E^(4*a)*(5 + m)*x^4))/(2*e*(1 + E^(2*a)*x^4)) + (-((E^(4*a)*(3 + m)*(5 + m)*(e*x)^(1+m))/(e*(1 + m))) + (2*E^(4*a)*(9 + 2*m + m^2)*(e*x)^(1+m)*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(E^(2*a)*x^4)]/(e*(1 + m)))/(2*E^(2*a)))/(4*E^(2*a))`

## Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (!\text{LtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 968  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_}))^{(q_)}), x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(a*b*e*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 1047  $\text{Int}[((g_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_}))^{(q_*)}((e_*) + (f_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(-(b*e - a*f))*(g*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^q/(a*b*g*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(g*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-1)}*\text{Simp}[c*(b*e*n*(p+1) + (b*e - a*f)*(m+1)) + d*(b*e*n*(p+1) + (b*e - a*f)*(m+n*q+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 0] \ \&\& \ !(EqQ[q, 1] \ \&\& \ \text{SimplerQ}[b*c - a*d, b*e - a*f])$

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]  
 :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),  
 x] /; FreeQ[{a, b, d, e, m, p}, x]`

### Maple [F]

$$\int (ex)^m \tanh(a + 2 \ln(x))^3 dx$$

input `int((e*x)^m*tanh(a+2*ln(x))^3,x)`

output `int((e*x)^m*tanh(a+2*ln(x))^3,x)`

### Fricas [F]

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^3 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="fricas")`

output `integral((e*x)^m*tanh(a + 2*log(x))^3, x)`

### Sympy [F]

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int (ex)^m \tanh^3(a + 2 \log(x)) dx$$

input `integrate((e*x)**m*tanh(a+2*ln(x))**3,x)`

output `Integral((e*x)**m*tanh(a + 2*log(x))**3, x)`

**Maxima [F]**

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^3 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="maxima")`

output `integrate((e*x)^m*tanh(a + 2*log(x))^3, x)`

**Giac [F]**

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x))^3 dx$$

input `integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="giac")`

output `integrate((e*x)^m*tanh(a + 2*log(x))^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int \tanh(a + 2 \ln(x))^3 (ex)^m dx$$

input `int(tanh(a + 2*log(x))^3*(e*x)^m,x)`

output `int(tanh(a + 2*log(x))^3*(e*x)^m, x)`

**Reduce [F]**

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \text{too large to display}$$

input `int((e*x)^m*tanh(a+2*log(x))^3,x)`

output

```
(e**m*(x**m*e**(4*a)*m**2*x**9 - 10*x**m*e**(4*a)*m*x**9 + 21*x**m*e**(4*a)
)*x**9 - 8*e**(4*a)*int(x**m/(e**(6*a)*m**2*x**12 - 10*e**(6*a)*m*x**12 +
21*e**(6*a)*x**12 + 3*e**(4*a)*m**2*x**8 - 30*e**(4*a)*m*x**8 + 63*e**(4*a)
)*x**8 + 3*e**(2*a)*m**2*x**4 - 30*e**(2*a)*m*x**4 + 63*e**(2*a)*x**4 + m*
*2 - 10*m + 21),x)*m**5*x**8 + 56*e**(4*a)*int(x**m/(e**(6*a)*m**2*x**12 -
10*e**(6*a)*m*x**12 + 21*e**(6*a)*x**12 + 3*e**(4*a)*m**2*x**8 - 30*e**(4
*a)*m*x**8 + 63*e**(4*a)*x**8 + 3*e**(2*a)*m**2*x**4 - 30*e**(2*a)*m*x**4
+ 63*e**(2*a)*x**4 + m**2 - 10*m + 21),x)*m**4*x**8 - 16*e**(4*a)*int(x**m
/(e**(6*a)*m**2*x**12 - 10*e**(6*a)*m*x**12 + 21*e**(6*a)*x**12 + 3*e**(4
a)*m**2*x**8 - 30*e**(4*a)*m*x**8 + 63*e**(4*a)*x**8 + 3*e**(2*a)*m**2*x**
4 - 30*e**(2*a)*m*x**4 + 63*e**(2*a)*x**4 + m**2 - 10*m + 21),x)*m**3*x**8
+ 304*e**(4*a)*int(x**m/(e**(6*a)*m**2*x**12 - 10*e**(6*a)*m*x**12 + 21*e
**(6*a)*x**12 + 3*e**(4*a)*m**2*x**8 - 30*e**(4*a)*m*x**8 + 63*e**(4*a)*x*
*8 + 3*e**(2*a)*m**2*x**4 - 30*e**(2*a)*m*x**4 + 63*e**(2*a)*x**4 + m**2 -
10*m + 21),x)*m**2*x**8 - 1128*e**(4*a)*int(x**m/(e**(6*a)*m**2*x**12 - 1
0*e**(6*a)*m*x**12 + 21*e**(6*a)*x**12 + 3*e**(4*a)*m**2*x**8 - 30*e**(4*a)
)*m*x**8 + 63*e**(4*a)*x**8 + 3*e**(2*a)*m**2*x**4 - 30*e**(2*a)*m*x**4 +
63*e**(2*a)*x**4 + m**2 - 10*m + 21),x)*m*x**8 - 1512*e**(4*a)*int(x**m/(e
**(6*a)*m**2*x**12 - 10*e**(6*a)*m*x**12 + 21*e**(6*a)*x**12 + 3*e**(4*a)*
m**2*x**8 - 30*e**(4*a)*m*x**8 + 63*e**(4*a)*x**8 + 3*e**(2*a)*m**2*x**...
```

### 3.173 $\int \tanh^p(a + b \log(x)) dx$

Optimal result	1318
Mathematica [B] (warning: unable to verify)	1318
Rubi [A] (verified)	1319
Maple [F]	1320
Fricas [F]	1320
Sympy [F]	1321
Maxima [F]	1321
Giac [F]	1321
Mupad [F(-1)]	1322
Reduce [F]	1322

#### Optimal result

Integrand size = 9, antiderivative size = 79

$$\int \tanh^p(a + b \log(x)) dx = x(1 - e^{2a}x^{2b})^{-p} (-1 + e^{2a}x^{2b})^p \operatorname{AppellF1}\left(\frac{1}{2b}, -p, p, \frac{1}{2}\left(2 + \frac{1}{b}\right), e^{2a}x^{2b}, -e^{2a}x^{2b}\right)$$

output

```
x*(-1+exp(2*a)*x^(2*b))^p*AppellF1(1/2/b,-p,p,1+1/2/b,exp(2*a)*x^(2*b),-exp(2*a)*x^(2*b))/((1-exp(2*a)*x^(2*b))^p)
```

#### Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 259 vs. 2(79) = 158.

Time = 0.43 (sec) , antiderivative size = 259, normalized size of antiderivative = 3.28

$$\int \tanh^p(a + b \log(x)) dx = \frac{(1 + 2b)x \left(\frac{-1 + e^{2a}x^{2b}}{1 + e^{2a}x^{2b}}\right)^p \operatorname{AppellF1}\left(\frac{1}{2b}, -p, p, 1 + \frac{1}{2b}, -p, p, 1 + \frac{1}{2b}\right) - 2be^{2a}px^{2b} \operatorname{AppellF1}\left(1 + \frac{1}{2b}, 1 - p, p, 2 + \frac{1}{2b}, e^{2a}x^{2b}, -e^{2a}x^{2b}\right) - 2be^{2a}px^{2b} \operatorname{AppellF1}\left(1 + \frac{1}{2b}, -p, 1 + p, 1 + \frac{1}{2b}, -p, 1 + p, 1 + \frac{1}{2b}\right)}{-2be^{2a}px^{2b} \operatorname{AppellF1}\left(1 + \frac{1}{2b}, 1 - p, p, 2 + \frac{1}{2b}, e^{2a}x^{2b}, -e^{2a}x^{2b}\right) - 2be^{2a}px^{2b} \operatorname{AppellF1}\left(1 + \frac{1}{2b}, -p, 1 + p, 1 + \frac{1}{2b}, -p, 1 + p, 1 + \frac{1}{2b}\right)}$$

input

```
Integrate[Tanh[a + b*Log[x]]^p,x]
```

output

$$\begin{aligned} & ((1 + 2*b)*x*((-1 + E^{(2*a)*x^{(2*b)}})/(1 + E^{(2*a)*x^{(2*b)}}))^p * \text{AppellF1}[1/(2*b), -p, p, 1 + 1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] / (-2*b * E^{(2*a)*x^{(2*b)}} * \text{AppellF1}[1 + 1/(2*b), 1 - p, p, 2 + 1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] - 2*b * E^{(2*a)*x^{(2*b)}} * \text{AppellF1}[1 + 1/(2*b), -p, 1 + p, 2 + 1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] + (1 + 2*b) * \text{AppellF1}[1/(2*b), -p, p, 1 + 1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})]) \end{aligned}$$
**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6067, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tanh^p(a + b \log(x)) dx \\ & \quad \downarrow 6067 \\ & \int (e^{2a}x^{2b} - 1)^p (e^{2a}x^{2b} + 1)^{-p} dx \\ & \quad \downarrow 937 \\ & (1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p \int (1 - e^{2a}x^{2b})^p (e^{2a}x^{2b} + 1)^{-p} dx \\ & \quad \downarrow 936 \\ & x(1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p \text{AppellF1}\left(\frac{1}{2b}, -p, p, \frac{1}{2}\left(2 + \frac{1}{b}\right), e^{2a}x^{2b}, -e^{2a}x^{2b}\right) \end{aligned}$$

input

$$\text{Int}[\text{Tanh}[a + b*\text{Log}[x]]^p, x]$$

output

$$(x*(-1 + E^{(2*a)*x^{(2*b)}}))^p * \text{AppellF1}[1/(2*b), -p, p, (2 + b^{(-1)})/2, E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] / (1 - E^{(2*a)*x^{(2*b)}})^p$$



## Definitions of rubi rules used

rule 936 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)
], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 937 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p])
Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}
, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [F]

$$\int \tanh(a + b \ln(x))^p dx$$

input `int(tanh(a+b*ln(x))^p,x)`

output `int(tanh(a+b*ln(x))^p,x)`

## Fricas [F]

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh(b \log(x) + a)^p dx$$

input `integrate(tanh(a+b*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(b*log(x) + a)^p, x)`

**Sympy [F]**

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh^p(a + b \log(x)) dx$$

input `integrate(tanh(a+b*ln(x))**p,x)`

output `Integral(tanh(a + b*log(x))**p, x)`

**Maxima [F]**

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh(b \log(x) + a)^p dx$$

input `integrate(tanh(a+b*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(b*log(x) + a)^p, x)`

**Giac [F]**

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh(b \log(x) + a)^p dx$$

input `integrate(tanh(a+b*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(b*log(x) + a)^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh(a + b \ln(x))^p dx$$

input `int(tanh(a + b*log(x))^p,x)`output `int(tanh(a + b*log(x))^p, x)`**Reduce [F]**

$$\int \tanh^p(a + b \log(x)) dx = \tanh(\log(x) b + a)^p x - \left( \int \frac{\tanh(\log(x) b + a)^p}{\tanh(\log(x) b + a)} dx \right) b p$$

$$+ \left( \int \tanh(\log(x) b + a)^p \tanh(\log(x) b + a) dx \right) b p$$

input `int(tanh(a+b*log(x))^p,x)`output `tanh(log(x)*b + a)**p*x - int(tanh(log(x)*b + a)**p/tanh(log(x)*b + a),x)*  
b*p + int(tanh(log(x)*b + a)**p*tanh(log(x)*b + a),x)*b*p`

### 3.174 $\int (ex)^m \tanh^p(a + b \log(x)) dx$

Optimal result	1323
Mathematica [A] (warning: unable to verify)	1323
Rubi [A] (verified)	1324
Maple [F]	1325
Fricas [F]	1325
Sympy [F]	1326
Maxima [F]	1326
Giac [F]	1326
Mupad [F(-1)]	1327
Reduce [F]	1327

#### Optimal result

Integrand size = 15, antiderivative size = 99

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \frac{(ex)^{1+m} (1 - e^{2a}x^{2b})^{-p} (-1 + e^{2a}x^{2b})^p \operatorname{AppellF1}\left(\frac{1+m}{2b}, -p, p, 1 + \frac{1+m}{2b}, e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{e(1+m)}$$

output  $(e*x)^{(1+m)}*(-1+\exp(2*a)*x^{(2*b)})^p*\operatorname{AppellF1}(1/2*(1+m)/b, -p, p, 1+1/2*(1+m)/b, \exp(2*a)*x^{(2*b)}, -\exp(2*a)*x^{(2*b)})/e/(1+m)/((1-\exp(2*a)*x^{(2*b)})^p)$

#### Mathematica [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.27

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \frac{x(ex)^m (1 - e^{2a}x^{2b})^{-p} \left(\frac{-1+e^{2a}x^{2b}}{1+e^{2a}x^{2b}}\right)^p (1 + e^{2a}x^{2b})^p \operatorname{AppellF1}\left(\frac{1+m}{2b}, -p, p, 1 + \frac{1+m}{2b}, e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{1+m}$$

input  $\operatorname{Integrate}[(e*x)^m*\operatorname{Tanh}[a + b*\operatorname{Log}[x]]^p, x]$

output

$$\frac{(x*(e*x)^m*((-1 + E^{(2*a)*x^{(2*b)}})/(1 + E^{(2*a)*x^{(2*b)}}))^p*(1 + E^{(2*a)*x^{(2*b)}})^p*AppellF1[(1 + m)/(2*b), -p, p, 1 + (1 + m)/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})])}{((1 + m)*(1 - E^{(2*a)*x^{(2*b)}}))^p}$$
**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6071, 1013, 1012}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (ex)^m \tanh^p(a + b \log(x)) dx \\ & \quad \downarrow \text{6071} \\ & \int (ex)^m (e^{2a}x^{2b} - 1)^p (e^{2a}x^{2b} + 1)^{-p} dx \\ & \quad \downarrow \text{1013} \\ & (1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p \int (ex)^m (1 - e^{2a}x^{2b})^p (e^{2a}x^{2b} + 1)^{-p} dx \\ & \quad \downarrow \text{1012} \\ & \frac{(ex)^{m+1} (1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p \text{AppellF1}\left(\frac{m+1}{2b}, -p, p, \frac{m+1}{2b} + 1, e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{e(m+1)} \end{aligned}$$

input

$$\text{Int}[(e*x)^m*\text{Tanh}[a + b*\text{Log}[x]]^p,x]$$

output

$$\frac{((e*x)^{(1 + m)*(-1 + E^{(2*a)*x^{(2*b)}}))^p*AppellF1[(1 + m)/(2*b), -p, p, 1 + (1 + m)/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})])}{(e*(1 + m)*(1 - E^{(2*a)*x^{(2*b)}}))^p}$$

## Definitions of rubi rules used

rule 1012

```
Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x]
&& NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])
```

rule 1013

```
Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a)^FracPart[p]) Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;
FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])
```

rule 6071

```
Int[((e._)*(x_))^(m._)*Tanh[((a._) + Log[x_]*(b._))*(d._)]^(p._), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]
```

## Maple [F]

$$\int (ex)^m \tanh(a + b \ln(x))^p dx$$

input

```
int((e*x)^m*tanh(a+b*ln(x))^p,x)
```

output

```
int((e*x)^m*tanh(a+b*ln(x))^p,x)
```

## Fricas [F]

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int (ex)^m \tanh(b \log(x) + a)^p dx$$

input

```
integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="fricas")
```

output `integral((e*x)^m*tanh(b*log(x) + a)^p, x)`

### Sympy [F]

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int (ex)^m \tanh^p(a + b \log(x)) dx$$

input `integrate((e*x)**m*tanh(a+b*ln(x))**p,x)`

output `Integral((e*x)**m*tanh(a + b*log(x))**p, x)`

### Maxima [F]

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int (ex)^m \tanh(b \log(x) + a)^p dx$$

input `integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="maxima")`

output `integrate((e*x)^m*tanh(b*log(x) + a)^p, x)`

### Giac [F]

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int (ex)^m \tanh(b \log(x) + a)^p dx$$

input `integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="giac")`

output `integrate((e*x)^m*tanh(b*log(x) + a)^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int \tanh(a + b \ln(x))^p (ex)^m dx$$

input `int(tanh(a + b*log(x))^p*(e*x)^m,x)`output `int(tanh(a + b*log(x))^p*(e*x)^m, x)`**Reduce [F]**

$$\int (ex)^m \tanh^p(a + b \log(x)) dx$$

$$= \frac{e^m \left( x^m \tanh(\log(x) b + a)^p x - \left( \int \frac{x^m \tanh(\log(x) b + a)^p}{\tanh(\log(x) b + a)} dx \right) b p + \left( \int x^m \tanh(\log(x) b + a)^p \tanh(\log(x) b + a) dx \right) \right)}{m + 1}$$

input `int((e*x)^m*tanh(a+b*log(x))^p,x)`output `(e**m*(x**m*tanh(log(x)*b + a)**p*x - int((x**m*tanh(log(x)*b + a)**p)/tanh(log(x)*b + a),x)*b*p + int(x**m*tanh(log(x)*b + a)**p*tanh(log(x)*b + a),x)*b*p))/(m + 1)`



### 3.175 $\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx$

Optimal result	1328
Mathematica [A] (verified)	1328
Rubi [A] (verified)	1329
Maple [F]	1330
Fricas [F]	1330
Sympy [F]	1331
Maxima [F]	1331
Giac [F]	1331
Mupad [F(-1)]	1332
Reduce [F]	1332

#### Optimal result

Integrand size = 11, antiderivative size = 51

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \frac{2^{-p} e^{-2a} (-1 + e^{2ax})^{1+p} \operatorname{Hypergeometric2F1} \left( p, 1 + p, 2 + p, \frac{1}{2}(1 - e^{2ax}) \right)}{1 + p}$$

output `(-1+exp(2*a)*x)^(p+1)*hypergeom([p, p+1], [2+p], 1/2-1/2*exp(2*a)*x)/(2^p)/exp(2*a)/(p+1)`

#### Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.49

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \frac{2^{-p} e^{-2a} \left( \frac{-1+e^{2ax}}{1+e^{2ax}} \right)^{1+p} (1 + e^{2ax})^{1+p} \operatorname{Hypergeometric2F1} \left( p, 1 + p, 2 + p, \frac{1}{2} - \frac{1}{2}e^{2ax} \right)}{1 + p}$$

input `Integrate[Tanh[a + Log[x]/2]^p,x]`

output

```
(((-1 + E^(2*a)*x)/(1 + E^(2*a)*x))^(1 + p)*(1 + E^(2*a)*x)^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 + p, 1/2 - (E^(2*a)*x)/2])/(2^p*E^(2*a)*(1 + p))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {6067, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx$$

↓ 6067

$$\int (e^{2a}x - 1)^p (e^{2a}x + 1)^{-p} dx$$

↓ 79

$$\frac{e^{-2a}2^{-p}(e^{2a}x - 1)^{p+1} \text{Hypergeometric2F1} \left( p, p + 1, p + 2, \frac{1}{2}(1 - e^{2a}x) \right)}{p + 1}$$

input

```
Int[Tanh[a + Log[x]/2]^p,x]
```

output

```
((-1 + E^(2*a)*x)^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 + p, (1 - E^(2*a)*x)/2])/(2^p*E^(2*a)*(1 + p))
```

## Definitions of rubi rules used

rule 79 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 6067 `Int[Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [F]

$$\int \tanh \left( a + \frac{\ln(x)}{2} \right)^p dx$$

input `int(tanh(a+1/2*ln(x))^p,x)`

output `int(tanh(a+1/2*ln(x))^p,x)`

## Fricas [F]

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \int \tanh \left( a + \frac{1}{2} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/2*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 1/2*log(x))^p, x)`

**Sympy [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx$$

input `integrate(tanh(a+1/2*ln(x))**p,x)`

output `Integral(tanh(a + log(x)/2)**p, x)`

**Maxima [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \int \tanh \left( a + \frac{1}{2} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/2*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 1/2*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \int \tanh \left( a + \frac{1}{2} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/2*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 1/2*log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \int \tanh \left( a + \frac{\ln(x)}{2} \right)^p dx$$

input `int(tanh(a + log(x)/2)^p,x)`output `int(tanh(a + log(x)/2)^p, x)`**Reduce [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{2} \right) dx = \tanh \left( \frac{\log(x)}{2} + a \right)^p x - \frac{\left( \int \frac{\tanh \left( \frac{\log(x)}{2} + a \right)^p}{\tanh \left( \frac{\log(x)}{2} + a \right)} dx \right) p}{2} + \frac{\left( \int \tanh \left( \frac{\log(x)}{2} + a \right)^p \tanh \left( \frac{\log(x)}{2} + a \right) dx \right) p}{2}$$

input `int(tanh(a+1/2*log(x))^p,x)`output `(2*tanh((log(x) + 2*a)/2)**p*x - int(tanh((log(x) + 2*a)/2)**p/tanh((log(x) + 2*a)/2),x)*p + int(tanh((log(x) + 2*a)/2)**p*tanh((log(x) + 2*a)/2),x)*p)/2`

### 3.176 $\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx$

Optimal result	1333
Mathematica [A] (verified)	1333
Rubi [A] (verified)	1334
Maple [F]	1335
Fricas [F]	1336
Sympy [F]	1336
Maxima [F]	1336
Giac [F]	1337
Mupad [F(-1)]	1337
Reduce [F]	1337

#### Optimal result

Integrand size = 11, antiderivative size = 106

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = e^{-4a} (-1 + e^{2a} \sqrt{x})^{1+p} (1 + e^{2a} \sqrt{x})^{1-p} - \frac{2^{1-p} e^{-4a} p (-1 + e^{2a} \sqrt{x})^{1+p} \text{Hypergeometric2F1} \left( p, 1 + p, 2 + p, \frac{1}{2} (1 - e^{2a} \sqrt{x}) \right)}{1 + p}$$

output

```
(-1+exp(2*a)*x^(1/2))^(p+1)*(1+exp(2*a)*x^(1/2))^(1-p)/exp(4*a)-2^(1-p)*p*
(-1+exp(2*a)*x^(1/2))^(p+1)*hypergeom([p, p+1],[2+p],1/2-1/2*exp(2*a)*x^(1
/2))/exp(4*a)/(p+1)
```

#### Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \frac{e^{-4a} (-1 + e^{2a} \sqrt{x}) \left( \frac{-1 + e^{2a} \sqrt{x}}{2 + 2e^{2a} \sqrt{x}} \right)^p (2^p (1 + p) (1 + e^{2a} \sqrt{x}) - 2p (1 + e^{2a} \sqrt{x})^p \text{Hypergeometric2F1} (p, 1 + p, 2 + p, \frac{1}{2} (1 - e^{2a} \sqrt{x})))}{1 + p}$$

input `Integrate[Tanh[a + Log[x]/4]^p, x]`

output `((-1 + E^(2*a)*Sqrt[x])*((-1 + E^(2*a)*Sqrt[x])/(2 + 2*E^(2*a)*Sqrt[x]))^p * (2^p*(1 + p)*(1 + E^(2*a)*Sqrt[x]) - 2*p*(1 + E^(2*a)*Sqrt[x])^p*Hypergeometric2F1[p, 1 + p, 2 + p, 1/2 - (E^(2*a)*Sqrt[x])/2])/ (E^(4*a)*(1 + p))`

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6067, 900, 90, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx \\
 & \quad \downarrow \text{6067} \\
 & \int (e^{2a}\sqrt{x} - 1)^p (e^{2a}\sqrt{x} + 1)^{-p} dx \\
 & \quad \downarrow \text{900} \\
 & 2 \int (e^{2a}\sqrt{x} - 1)^p (e^{2a}\sqrt{x} + 1)^{-p} \sqrt{x} d\sqrt{x} \\
 & \quad \downarrow \text{90} \\
 & 2 \left( \frac{1}{2} e^{-4a} (e^{2a}\sqrt{x} - 1)^{p+1} (e^{2a}\sqrt{x} + 1)^{1-p} - e^{-2a} p \int (e^{2a}\sqrt{x} - 1)^p (e^{2a}\sqrt{x} + 1)^{-p} d\sqrt{x} \right) \\
 & \quad \downarrow \text{79} \\
 & 2 \left( \frac{1}{2} e^{-4a} (e^{2a}\sqrt{x} - 1)^{p+1} (e^{2a}\sqrt{x} + 1)^{1-p} - \frac{e^{-4a} 2^{-p} p (e^{2a}\sqrt{x} - 1)^{p+1} \text{Hypergeometric2F1} \left( p, p + 1, p + 2, \frac{1}{2} (1 - \right)}{p + 1} \right)
 \end{aligned}$$

input `Int [Tanh[a + Log[x]/4]^p, x]`

output

```
2*((( -1 + E^(2*a)*Sqrt[x])^(1 + p)*(1 + E^(2*a)*Sqrt[x])^(1 - p))/(2*E^(4*a)) - (p*(-1 + E^(2*a)*Sqrt[x])^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 + p, (1 - E^(2*a)*Sqrt[x])/2])/(2^p*E^(4*a)*(1 + p)))
```

**Defintions of rubi rules used**

rule 79

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

rule 90

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
```

rule 900

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := With[{g = Denominator[n]}, Simp[g Subst[Int[x^(g - 1)*(a + b*x^(g*n))^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x]] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]
```

rule 6067

```
Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d))*x^(2*b*d)]^p/(1 + E^(2*a*d))*x^(2*b*d)]^p, x] /; FreeQ[{a, b, d, p}, x]
```

**Maple [F]**

$$\int \tanh\left(a + \frac{\ln(x)}{4}\right)^p dx$$

input

```
int(tanh(a+1/4*ln(x))^p,x)
```



output `int(tanh(a+1/4*ln(x))^p,x)`

### Fricas [F]

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \int \tanh \left( a + \frac{1}{4} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/4*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 1/4*log(x))^p, x)`

### Sympy [F]

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx$$

input `integrate(tanh(a+1/4*ln(x))**p,x)`

output `Integral(tanh(a + log(x)/4)**p, x)`

### Maxima [F]

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \int \tanh \left( a + \frac{1}{4} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/4*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 1/4*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \int \tanh \left( a + \frac{1}{4} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/4*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 1/4*log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \int \tanh \left( a + \frac{\ln(x)}{4} \right)^p dx$$

input `int(tanh(a + log(x)/4)^p,x)`

output `int(tanh(a + log(x)/4)^p, x)`

**Reduce [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{4} \right) dx = \tanh \left( \frac{\log(x)}{4} + a \right)^p x - \frac{\left( \int \frac{\tanh \left( \frac{\log(x)}{4} + a \right)^p}{\tanh \left( \frac{\log(x)}{4} + a \right)} dx \right) p}{4} + \frac{\left( \int \tanh \left( \frac{\log(x)}{4} + a \right)^p \tanh \left( \frac{\log(x)}{4} + a \right) dx \right) p}{4}$$

input `int(tanh(a+1/4*log(x))^p,x)`

output `(4*tanh((log(x) + 4*a)/4)**p*x - int(tanh((log(x) + 4*a)/4)**p/tanh((log(x) + 4*a)/4),x)*p + int(tanh((log(x) + 4*a)/4)**p*tanh((log(x) + 4*a)/4),x)*p)/4`

### 3.177 $\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$

Optimal result	1338
Mathematica [A] (verified)	1338
Rubi [A] (verified)	1339
Maple [F]	1341
Fricas [F]	1341
Sympy [F]	1342
Maxima [F]	1342
Giac [F]	1342
Mupad [F(-1)]	1343
Reduce [F]	1343

#### Optimal result

Integrand size = 11, antiderivative size = 158

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$$

$$= -e^{-6a} p (-1 + e^{2a \sqrt[3]{x}})^{1+p} (1 + e^{2a \sqrt[3]{x}})^{1-p} + e^{-4a} (-1 + e^{2a \sqrt[3]{x}})^{1+p} (1 + e^{2a \sqrt[3]{x}})^{1-p} \sqrt[3]{x}$$

$$+ \frac{2^{-p} e^{-6a} (1 + 2p^2) (-1 + e^{2a \sqrt[3]{x}})^{1+p} \text{Hypergeometric2F1} \left( p, 1 + p, 2 + p, \frac{1}{2} (1 - e^{2a \sqrt[3]{x}}) \right)}{1 + p}$$

output

```
-p*(-1+exp(2*a)*x^(1/3))^(p+1)*(1+exp(2*a)*x^(1/3))^(1-p)/exp(6*a)+(-1+exp
(2*a)*x^(1/3))^(p+1)*(1+exp(2*a)*x^(1/3))^(1-p)*x^(1/3)/exp(4*a)+(2*p^2+1)
*(-1+exp(2*a)*x^(1/3))^(p+1)*hypergeom([p, p+1],[2+p],1/2-1/2*exp(2*a)*x^(
1/3))/(2^p)/exp(6*a)/(p+1)
```

#### Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.89

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$$

$$= \frac{e^{-6a} (-1 + e^{2a \sqrt[3]{x}}) \left( \frac{-1 + e^{2a \sqrt[3]{x}}}{2 + 2e^{2a \sqrt[3]{x}}} \right)^p (2^p (1 + p) (1 + e^{2a \sqrt[3]{x}}) (-p + e^{2a \sqrt[3]{x}}) + (1 + 2p^2) (1 + e^{2a \sqrt[3]{x}})^p \text{Hyp}}{1 + p}$$

input `Integrate[Tanh[a + Log[x]/6]^p,x]`

output  $((-1 + E^{(2*a)*x^{(1/3)}})*((-1 + E^{(2*a)*x^{(1/3)}})/(2 + 2*E^{(2*a)*x^{(1/3)}}))^p * (2^p*(1 + p)*(1 + E^{(2*a)*x^{(1/3)}})*(-p + E^{(2*a)*x^{(1/3)}}) + (1 + 2*p^2)*(1 + E^{(2*a)*x^{(1/3)}})^p*Hypergeometric2F1[p, 1 + p, 2 + p, 1/2 - (E^{(2*a)*x^{(1/3)}}/2)])/(E^{(6*a)*x^{(1/3)}}*(1 + p))$

### Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {6067, 900, 101, 90, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$$

↓ 6067

$$\int (e^{2a \sqrt[3]{x}} - 1)^p (e^{2a \sqrt[3]{x}} + 1)^{-p} dx$$

↓ 900

$$3 \int (e^{2a \sqrt[3]{x}} - 1)^p (e^{2a \sqrt[3]{x}} + 1)^{-p} x^{2/3} d\sqrt[3]{x}$$

↓ 101

$$3 \left( \frac{1}{3} e^{-4a} \int (e^{2a \sqrt[3]{x}} - 1)^p (e^{2a \sqrt[3]{x}} + 1)^{-p} (1 - 2e^{2a} p \sqrt[3]{x}) d\sqrt[3]{x} + \frac{1}{3} e^{-4a \sqrt[3]{x}} (e^{2a \sqrt[3]{x}} - 1)^{p+1} (e^{2a \sqrt[3]{x}} + 1)^{1-p} \right)$$

↓ 90

$$3 \left( \frac{1}{3} e^{-4a} \left( (2p^2 + 1) \int (e^{2a \sqrt[3]{x}} - 1)^p (e^{2a \sqrt[3]{x}} + 1)^{-p} d\sqrt[3]{x} - e^{-2a} p (e^{2a \sqrt[3]{x}} - 1)^{p+1} (e^{2a \sqrt[3]{x}} + 1)^{1-p} \right) + \frac{1}{3} e^{-4a \sqrt[3]{x}} \right)$$

↓ 79

$$3 \left( \frac{1}{3} e^{-4a} \left( \frac{e^{-2a} 2^{-p} (2p^2 + 1) (e^{2a} \sqrt[3]{x} - 1)^{p+1} \operatorname{Hypergeometric2F1} \left( p, p + 1, p + 2, \frac{1}{2} (1 - e^{2a} \sqrt[3]{x}) \right)}{p + 1} \right) - e^{-2a} p (e^{2a} \sqrt[3]{x}) \right)$$

input `Int[Tanh[a + Log[x]/6]^p,x]`

output `3*((( -1 + E^(2*a)*x^(1/3))^(1 + p)*(1 + E^(2*a)*x^(1/3))^(1 - p)*x^(1/3))/(3*E^(4*a)) + (-((p*(-1 + E^(2*a)*x^(1/3))^(1 + p)*(1 + E^(2*a)*x^(1/3))^(1 - p))/E^(2*a)) + ((1 + 2*p^2)*(-1 + E^(2*a)*x^(1/3))^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 + p, (1 - E^(2*a)*x^(1/3))/2])/(2^p*E^(2*a)*(1 + p)))/(3*E^(4*a))`

### Defintions of rubi rules used

rule 79 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 101 `Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 900 `Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := With[{g = Denominator[n]}, Simp[g Subst[Int[x^(g - 1)*(a + b*x^(g*n))^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x]] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]`

rule 6067 `Int[Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

### Maple [F]

$$\int \tanh \left( a + \frac{\ln(x)}{6} \right)^p dx$$

input `int(tanh(a+1/6*ln(x))^p,x)`

output `int(tanh(a+1/6*ln(x))^p,x)`

### Fricas [F]

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \int \tanh \left( a + \frac{1}{6} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/6*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 1/6*log(x))^p, x)`

**Sympy [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx$$

input `integrate(tanh(a+1/6*ln(x))**p,x)`

output `Integral(tanh(a + log(x)/6)**p, x)`

**Maxima [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \int \tanh \left( a + \frac{1}{6} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/6*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 1/6*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \int \tanh \left( a + \frac{1}{6} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/6*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 1/6*log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \int \tanh \left( a + \frac{\ln(x)}{6} \right)^p dx$$

input `int(tanh(a + log(x)/6)^p,x)`output `int(tanh(a + log(x)/6)^p, x)`**Reduce [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{6} \right) dx = \tanh \left( \frac{\log(x)}{6} + a \right)^p x - \frac{\left( \int \frac{\tanh \left( \frac{\log(x)}{6} + a \right)^p}{\tanh \left( \frac{\log(x)}{6} + a \right)} dx \right) p}{6} + \frac{\left( \int \tanh \left( \frac{\log(x)}{6} + a \right)^p \tanh \left( \frac{\log(x)}{6} + a \right) dx \right) p}{6}$$

input `int(tanh(a+1/6*log(x))^p,x)`output `(6*tanh((log(x) + 6*a)/6)**p*x - int(tanh((log(x) + 6*a)/6)**p/tanh((log(x) + 6*a)/6),x)*p + int(tanh((log(x) + 6*a)/6)**p*tanh((log(x) + 6*a)/6),x)*p)/6`



### 3.178 $\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx$

Optimal result	1344
Mathematica [A] (warning: unable to verify)	1345
Rubi [A] (verified)	1345
Maple [F]	1348
Fricas [F]	1348
Sympy [F]	1348
Maxima [F]	1349
Giac [F]	1349
Mupad [F(-1)]	1349
Reduce [F]	1350

#### Optimal result

Integrand size = 11, antiderivative size = 219

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx = \frac{1}{3} e^{-8a} (3 - 2p + 2p^2) (-1 + e^{2a} \sqrt[4]{x})^{1+p} (1 + e^{2a} \sqrt[4]{x})^{1-p} - \frac{2}{3} e^{-8a} p (-1 + e^{2a} \sqrt[4]{x})^{2+p} (1 + e^{2a} \sqrt[4]{x})^{1-p} + e^{-4a} (-1 + e^{2a} \sqrt[4]{x})^{1+p} (1 + e^{2a} \sqrt[4]{x})^{1-p} \sqrt{x} - \frac{2^{2-p} e^{-8a} p (2 + p^2) (-1 + e^{2a} \sqrt[4]{x})^{1+p} \text{Hypergeometric2F1} \left( p, 1 + p, 2 + p, \frac{1}{2} (1 - e^{2a} \sqrt[4]{x}) \right)}{3(1 + p)}$$

output

```
1/3*(2*p^2-2*p+3)*(-1+exp(2*a)*x^(1/4))^(p+1)*(1+exp(2*a)*x^(1/4))^(1-p)/exp(8*a)-2/3*p*(-1+exp(2*a)*x^(1/4))^(2+p)*(1+exp(2*a)*x^(1/4))^(1-p)/exp(8*a)+(-1+exp(2*a)*x^(1/4))^(p+1)*(1+exp(2*a)*x^(1/4))^(1-p)*x^(1/2)/exp(4*a)-1/3*2^(2-p)*p*(p^2+2)*(-1+exp(2*a)*x^(1/4))^(p+1)*hypergeom([p, p+1], [2+p], 1/2-1/2*exp(2*a)*x^(1/4))/exp(8*a)/(p+1)
```

**Mathematica [A] (warning: unable to verify)**

Time = 0.89 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.04

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx$$

$$= \frac{e^{-8a} (-1 + e^{2a} \sqrt[4]{x}) \left( \frac{-1 + e^{2a} \sqrt[4]{x}}{2 + 2e^{2a} \sqrt[4]{x}} \right)^p (-8p(1 + e^{2a} \sqrt[4]{x})^p \text{Hypergeometric2F1}(-2 + p, 1 + p, 2 + p, \frac{1}{2} - \frac{1}{2}e^{2a} \sqrt[4]{x}))}{(1 + e^{2a} \sqrt[4]{x})^{2p+1}}$$

input `Integrate[Tanh[a + Log[x]/8]^p,x]`output
$$\frac{((-1 + E^{(2*a)*x^{(1/4)}})*((-1 + E^{(2*a)*x^{(1/4)}})/(2 + 2*E^{(2*a)*x^{(1/4)}}))^p * (-8*p*(1 + E^{(2*a)*x^{(1/4)}})^p * \text{Hypergeometric2F1}[-2 + p, 1 + p, 2 + p, 1/2 - (E^{(2*a)*x^{(1/4)}})/2] + 4*(1 + 2*p)*(1 + E^{(2*a)*x^{(1/4)}})^p * \text{Hypergeometric2F1}[-1 + p, 1 + p, 2 + p, 1/2 - (E^{(2*a)*x^{(1/4)}})/2] + (1 + p)*(2^p * E^{(4*a)}*(1 + E^{(2*a)*x^{(1/4)}})*\text{Sqrt}[x] - 2*(1 + E^{(2*a)*x^{(1/4)}})^p * \text{Hypergeometric2F1}[p, 1 + p, 2 + p, 1/2 - (E^{(2*a)*x^{(1/4)}})/2])}}{(E^{(8*a)}*(1 + p))}$$
**Rubi [A] (verified)**Time = 0.35 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$ , Rules used = {6067, 900, 111, 27, 164, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx$$

$$\downarrow \text{6067}$$

$$\int (e^{2a} \sqrt[4]{x} - 1)^p (e^{2a} \sqrt[4]{x} + 1)^{-p} dx$$

$$\downarrow \text{900}$$

$$4 \int (e^{2a} \sqrt[4]{x} - 1)^p (e^{2a} \sqrt[4]{x} + 1)^{-p} x^{3/4} d\sqrt[4]{x}$$

↓ 111

$$4 \left( \frac{1}{4} e^{-4a} \int 2(e^{2a} \sqrt[4]{x} - 1)^p (e^{2a} \sqrt[4]{x} + 1)^{-p} (1 - e^{2a} p \sqrt[4]{x}) \sqrt[4]{x} d\sqrt[4]{x} + \frac{1}{4} e^{-4a} \sqrt{x} (e^{2a} \sqrt[4]{x} - 1)^{p+1} (e^{2a} \sqrt[4]{x} + 1)^{1-p} \right)$$

↓ 27

$$4 \left( \frac{1}{2} e^{-4a} \int (e^{2a} \sqrt[4]{x} - 1)^p (e^{2a} \sqrt[4]{x} + 1)^{-p} (1 - e^{2a} p \sqrt[4]{x}) \sqrt[4]{x} d\sqrt[4]{x} + \frac{1}{4} e^{-4a} \sqrt{x} (e^{2a} \sqrt[4]{x} - 1)^{p+1} (e^{2a} \sqrt[4]{x} + 1)^{1-p} \right)$$

↓ 164

$$4 \left( \frac{1}{2} e^{-4a} \left( \frac{1}{6} e^{-8a} (e^{2a} \sqrt[4]{x} - 1)^{p+1} (e^{2a} \sqrt[4]{x} + 1)^{1-p} (e^{4a} (2p^2 + 3) - 2e^{6a} p \sqrt[4]{x}) - \frac{2}{3} e^{-2a} p (p^2 + 2) \int (e^{2a} \sqrt[4]{x} - 1)^p \right. \right.$$

↓ 79

$$\left. 4 \left( \frac{1}{2} e^{-4a} \left( \frac{1}{6} e^{-8a} (e^{2a} \sqrt[4]{x} - 1)^{p+1} (e^{2a} \sqrt[4]{x} + 1)^{1-p} (e^{4a} (2p^2 + 3) - 2e^{6a} p \sqrt[4]{x}) - \frac{e^{-4a} 2^{1-p} p (p^2 + 2) (e^{2a} \sqrt[4]{x} - 1)^2}{\dots} \right) \right)$$

input

```
Int [Tanh[a + Log[x]/8]^p, x]
```

output

```
4*((( -1 + E^(2*a)*x^(1/4))^(1 + p)*(1 + E^(2*a)*x^(1/4))^(1 - p)*Sqrt[x])/
(4*E^(4*a)) + ((( -1 + E^(2*a)*x^(1/4))^(1 + p)*(1 + E^(2*a)*x^(1/4))^(1 -
p)*(E^(4*a)*(3 + 2*p^2) - 2*E^(6*a)*p*x^(1/4)))/(6*E^(8*a)) - (2^(1 - p)*
p*(2 + p^2)*(-1 + E^(2*a)*x^(1/4))^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 +
p, (1 - E^(2*a)*x^(1/4))/2])/(3*E^(4*a)*(1 + p)))/(2*E^(4*a))
```

## Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 79 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 164 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x))*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Simp[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)) Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]`
- rule 900 `Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := With[{g = Denominator[n]}, Simp[g Subst[Int[x^(g - 1)*(a + b*x^(g*n))^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x]] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

### Maple [F]

$$\int \tanh\left(a + \frac{\ln(x)}{8}\right)^p dx$$

input `int(tanh(a+1/8*ln(x))^p,x)`

output `int(tanh(a+1/8*ln(x))^p,x)`

### Fricas [F]

$$\int \tanh^p\left(a + \frac{\log(x)}{8}\right) dx = \int \tanh\left(a + \frac{1}{8} \log(x)\right)^p dx$$

input `integrate(tanh(a+1/8*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 1/8*log(x))^p, x)`

### Sympy [F]

$$\int \tanh^p\left(a + \frac{\log(x)}{8}\right) dx = \int \tanh^p\left(a + \frac{\log(x)}{8}\right) dx$$

input `integrate(tanh(a+1/8*ln(x))**p,x)`

output `Integral(tanh(a + log(x)/8)**p, x)`

**Maxima [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx = \int \tanh \left( a + \frac{1}{8} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/8*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 1/8*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx = \int \tanh \left( a + \frac{1}{8} \log(x) \right)^p dx$$

input `integrate(tanh(a+1/8*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 1/8*log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx = \int \tanh \left( a + \frac{\ln(x)}{8} \right)^p dx$$

input `int(tanh(a + log(x)/8)^p,x)`

output `int(tanh(a + log(x)/8)^p, x)`

**Reduce [F]**

$$\int \tanh^p \left( a + \frac{\log(x)}{8} \right) dx = \tanh \left( \frac{\log(x)}{8} + a \right)^p x - \frac{\left( \int \frac{\tanh \left( \frac{\log(x)}{8} + a \right)^p}{\tanh \left( \frac{\log(x)}{8} + a \right)} dx \right) p}{8} + \frac{\left( \int \tanh \left( \frac{\log(x)}{8} + a \right)^p \tanh \left( \frac{\log(x)}{8} + a \right) dx \right) p}{8}$$

input `int(tanh(a+1/8*log(x))^p,x)`

output `(8*tanh((log(x) + 8*a)/8)**p*x - int(tanh((log(x) + 8*a)/8)**p/tanh((log(x) + 8*a)/8),x)*p + int(tanh((log(x) + 8*a)/8)**p*tanh((log(x) + 8*a)/8),x)*p)/8`

### 3.179 $\int \tanh^p(a + \log(x)) dx$

Optimal result	1351
Mathematica [B] (warning: unable to verify)	1351
Rubi [A] (verified)	1352
Maple [F]	1353
Fricas [F]	1353
Sympy [F]	1354
Maxima [F]	1354
Giac [F]	1354
Mupad [F(-1)]	1355
Reduce [F]	1355

#### Optimal result

Integrand size = 7, antiderivative size = 61

$$\int \tanh^p(a + \log(x)) dx = x(1 - e^{2a}x^2)^{-p} (-1 + e^{2a}x^2)^p \operatorname{AppellF1}\left(\frac{1}{2}, -p, p, \frac{3}{2}, e^{2a}x^2, -e^{2a}x^2\right)$$

output

```
x*(-1+exp(2*a)*x^2)^p*AppellF1(1/2,-p,p,3/2,exp(2*a)*x^2,-exp(2*a)*x^2)/((1-exp(2*a)*x^2)^p)
```

#### Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 171 vs. 2(61) = 122.

Time = 0.39 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.80

$$\int \tanh^p(a + \log(x)) dx$$

$$= \frac{3x \left(\frac{-1+e^{2a}x^2}{1+e^{2a}x^2}\right)^p \operatorname{AppellF1}\left(\frac{1}{2}, -p, p, \frac{3}{2}, e^{2a}x^2, -e^{2a}x^2\right)}{3 \operatorname{AppellF1}\left(\frac{1}{2}, -p, p, \frac{3}{2}, e^{2a}x^2, -e^{2a}x^2\right) - 2e^{2a}px^2 \left(\operatorname{AppellF1}\left(\frac{3}{2}, 1-p, p, \frac{5}{2}, e^{2a}x^2, -e^{2a}x^2\right) + \operatorname{AppellF1}\left(\frac{3}{2}, 1-p, p, \frac{5}{2}, e^{2a}x^2, -e^{2a}x^2\right)\right)}$$

input

```
Integrate[Tanh[a + Log[x]]^p,x]
```



output

```
(3*x*((-1 + E^(2*a)*x^2)/(1 + E^(2*a)*x^2))^p*AppellF1[1/2, -p, p, 3/2, E^(2*a)*x^2, -(E^(2*a)*x^2)]/(3*AppellF1[1/2, -p, p, 3/2, E^(2*a)*x^2, -(E^(2*a)*x^2)] - 2*E^(2*a)*p*x^2*(AppellF1[3/2, 1 - p, p, 5/2, E^(2*a)*x^2, -(E^(2*a)*x^2)] + AppellF1[3/2, -p, 1 + p, 5/2, E^(2*a)*x^2, -(E^(2*a)*x^2)]))
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6067, 334, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p(a + \log(x)) dx$$

$$\downarrow \text{6067}$$

$$\int (e^{2a}x^2 - 1)^p (e^{2a}x^2 + 1)^{-p} dx$$

$$\downarrow \text{334}$$

$$(1 - e^{2a}x^2)^{-p} (e^{2a}x^2 - 1)^p \int (1 - e^{2a}x^2)^p (e^{2a}x^2 + 1)^{-p} dx$$

$$\downarrow \text{333}$$

$$x(1 - e^{2a}x^2)^{-p} (e^{2a}x^2 - 1)^p \text{AppellF1}\left(\frac{1}{2}, -p, p, \frac{3}{2}, e^{2a}x^2, -e^{2a}x^2\right)$$

input

```
Int[Tanh[a + Log[x]]^p,x]
```

output

```
(x*(-1 + E^(2*a)*x^2)^p*AppellF1[1/2, -p, p, 3/2, E^(2*a)*x^2, -(E^(2*a)*x^2)]/(1 - E^(2*a)*x^2)^p
```

## Definitions of rubi rules used

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d))*x^(2*b*d)]^p/(1 + E^(2*a*d))*x^(2*b*d)]^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [F]

$$\int \tanh(a + \ln(x))^p dx$$

input `int(tanh(a+ln(x))^p,x)`

output `int(tanh(a+ln(x))^p,x)`

## Fricas [F]

$$\int \tanh^p(a + \log(x)) dx = \int \tanh(a + \log(x))^p dx$$

input `integrate(tanh(a+log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + log(x))^p, x)`

**Sympy [F]**

$$\int \tanh^p(a + \log(x)) dx = \int \tanh^p(a + \log(x)) dx$$

input `integrate(tanh(a+ln(x))**p,x)`

output `Integral(tanh(a + log(x))**p, x)`

**Maxima [F]**

$$\int \tanh^p(a + \log(x)) dx = \int \tanh(a + \log(x))^p dx$$

input `integrate(tanh(a+log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p(a + \log(x)) dx = \int \tanh(a + \log(x))^p dx$$

input `integrate(tanh(a+log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p(a + \log(x)) dx = \int \tanh(a + \ln(x))^p dx$$

input `int(tanh(a + log(x))^p, x)`output `int(tanh(a + log(x))^p, x)`**Reduce [F]**

$$\int \tanh^p(a + \log(x)) dx = \tanh(\log(x) + a)^p x - \left( \int \frac{\tanh(\log(x) + a)^p}{\tanh(\log(x) + a)} dx \right) p + \left( \int \tanh(\log(x) + a)^p \tanh(\log(x) + a) dx \right) p$$

input `int(tanh(a+log(x))^p, x)`output `tanh(log(x) + a)**p*x - int(tanh(log(x) + a)**p/tanh(log(x) + a), x)*p + int(tanh(log(x) + a)**p*tanh(log(x) + a), x)*p`

### 3.180 $\int \tanh^p(a + 2 \log(x)) dx$

Optimal result	1356
Mathematica [B] (warning: unable to verify)	1356
Rubi [A] (verified)	1357
Maple [F]	1358
Fricas [F]	1358
Sympy [F]	1359
Maxima [F]	1359
Giac [F]	1359
Mupad [F(-1)]	1360
Reduce [F]	1360

#### Optimal result

Integrand size = 9, antiderivative size = 61

$$\int \tanh^p(a + 2 \log(x)) dx = x(1 - e^{2a}x^4)^{-p} (-1 + e^{2a}x^4)^p \operatorname{AppellF1}\left(\frac{1}{4}, -p, p, \frac{5}{4}, e^{2a}x^4, -e^{2a}x^4\right)$$

output

```
x*(-1+exp(2*a)*x^4)^p*AppellF1(1/4,-p,p,5/4,exp(2*a)*x^4,-exp(2*a)*x^4)/((1-exp(2*a)*x^4)^p)
```

#### Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 171 vs. 2(61) = 122.

Time = 0.44 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.80

$$\int \tanh^p(a + 2 \log(x)) dx$$

$$= \frac{5x \left(\frac{-1+e^{2a}x^4}{1+e^{2a}x^4}\right)^p \operatorname{AppellF1}\left(\frac{1}{4}, -p, p, \frac{5}{4}, e^{2a}x^4, -e^{2a}x^4\right)}{5 \operatorname{AppellF1}\left(\frac{1}{4}, -p, p, \frac{5}{4}, e^{2a}x^4, -e^{2a}x^4\right) - 4e^{2a}px^4 \left(\operatorname{AppellF1}\left(\frac{5}{4}, 1 - p, p, \frac{9}{4}, e^{2a}x^4, -e^{2a}x^4\right) + \operatorname{AppellF1}\left(\frac{5}{4}, 1 - p, p, \frac{9}{4}, e^{2a}x^4, -e^{2a}x^4\right)\right)}$$

input

```
Integrate[Tanh[a + 2*Log[x]]^p,x]
```

output

```
(5*x*((-1 + E^(2*a)*x^4)/(1 + E^(2*a)*x^4))^p*AppellF1[1/4, -p, p, 5/4, E^(2*a)*x^4, -(E^(2*a)*x^4)]/(5*AppellF1[1/4, -p, p, 5/4, E^(2*a)*x^4, -(E^(2*a)*x^4)] - 4*E^(2*a)*p*x^4*(AppellF1[5/4, 1 - p, p, 9/4, E^(2*a)*x^4, -(E^(2*a)*x^4)] + AppellF1[5/4, -p, 1 + p, 9/4, E^(2*a)*x^4, -(E^(2*a)*x^4)]))
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6067, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p(a + 2 \log(x)) dx$$

$$\downarrow 6067$$

$$\int (e^{2a}x^4 - 1)^p (e^{2a}x^4 + 1)^{-p} dx$$

$$\downarrow 937$$

$$(1 - e^{2a}x^4)^{-p} (e^{2a}x^4 - 1)^p \int (1 - e^{2a}x^4)^p (e^{2a}x^4 + 1)^{-p} dx$$

$$\downarrow 936$$

$$x(1 - e^{2a}x^4)^{-p} (e^{2a}x^4 - 1)^p \text{AppellF1}\left(\frac{1}{4}, -p, p, \frac{5}{4}, e^{2a}x^4, -e^{2a}x^4\right)$$

input

```
Int[Tanh[a + 2*Log[x]]^p,x]
```

output

```
(x*(-1 + E^(2*a)*x^4))^p*AppellF1[1/4, -p, p, 5/4, E^(2*a)*x^4, -(E^(2*a)*x^4)]/(1 - E^(2*a)*x^4)^p
```

## Definitions of rubi rules used

rule 936 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)
], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 937 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p])
Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}
, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [F]

$$\int \tanh(a + 2 \ln(x))^p dx$$

input `int(tanh(a+2*ln(x))^p,x)`

output `int(tanh(a+2*ln(x))^p,x)`

## Fricas [F]

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh(a + 2 \log(x))^p dx$$

input `integrate(tanh(a+2*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 2*log(x))^p, x)`

**Sympy [F]**

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh^p(a + 2 \log(x)) dx$$

input `integrate(tanh(a+2*ln(x))**p,x)`

output `Integral(tanh(a + 2*log(x))**p, x)`

**Maxima [F]**

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh(a + 2 \log(x))^p dx$$

input `integrate(tanh(a+2*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 2*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh(a + 2 \log(x))^p dx$$

input `integrate(tanh(a+2*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 2*log(x))^p, x)`



**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh(a + 2 \ln(x))^p dx$$

input `int(tanh(a + 2*log(x))^p,x)`output `int(tanh(a + 2*log(x))^p, x)`**Reduce [F]**

$$\int \tanh^p(a + 2 \log(x)) dx = \tanh(2 \log(x) + a)^p x - 2 \left( \int \frac{\tanh(2 \log(x) + a)^p}{\tanh(2 \log(x) + a)} dx \right)^p$$

$$+ 2 \left( \int \tanh(2 \log(x) + a)^p \tanh(2 \log(x) + a) dx \right)^p$$

input `int(tanh(a+2*log(x))^p,x)`output `tanh(2*log(x) + a)**p*x - 2*int(tanh(2*log(x) + a)**p/tanh(2*log(x) + a),x)*p + 2*int(tanh(2*log(x) + a)**p*tanh(2*log(x) + a),x)*p`

### 3.181 $\int \tanh^p(a + 3 \log(x)) dx$

Optimal result	1361
Mathematica [B] (warning: unable to verify)	1361
Rubi [A] (verified)	1362
Maple [F]	1363
Fricas [F]	1363
Sympy [F]	1364
Maxima [F]	1364
Giac [F]	1364
Mupad [F(-1)]	1365
Reduce [F]	1365

#### Optimal result

Integrand size = 9, antiderivative size = 61

$$\int \tanh^p(a + 3 \log(x)) dx = x(1 - e^{2a}x^6)^{-p} (-1 + e^{2a}x^6)^p \operatorname{AppellF1}\left(\frac{1}{6}, -p, p, \frac{7}{6}, e^{2a}x^6, -e^{2a}x^6\right)$$

output

```
x*(-1+exp(2*a)*x^6)^p*AppellF1(1/6,-p,p,7/6,exp(2*a)*x^6,-exp(2*a)*x^6)/((1-exp(2*a)*x^6)^p)
```

#### Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 171 vs. 2(61) = 122.

Time = 0.44 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.80

$$\int \tanh^p(a + 3 \log(x)) dx$$

$$= \frac{7x \left(\frac{-1+e^{2a}x^6}{1+e^{2a}x^6}\right)^p \operatorname{AppellF1}\left(\frac{1}{6}, -p, p, \frac{7}{6}, e^{2a}x^6, -e^{2a}x^6\right)}{7 \operatorname{AppellF1}\left(\frac{1}{6}, -p, p, \frac{7}{6}, e^{2a}x^6, -e^{2a}x^6\right) - 6e^{2a}px^6 \left(\operatorname{AppellF1}\left(\frac{7}{6}, 1 - p, p, \frac{13}{6}, e^{2a}x^6, -e^{2a}x^6\right) + \operatorname{AppellF1}\left(\frac{7}{6}, 1 - p, p, \frac{13}{6}, e^{2a}x^6, -e^{2a}x^6\right)\right)}$$

input

```
Integrate[Tanh[a + 3*Log[x]]^p,x]
```

output

```
(7*x*((-1 + E^(2*a)*x^6)/(1 + E^(2*a)*x^6))^p*AppellF1[1/6, -p, p, 7/6, E^(2*a)*x^6, -(E^(2*a)*x^6)]/(7*AppellF1[1/6, -p, p, 7/6, E^(2*a)*x^6, -(E^(2*a)*x^6)] - 6*E^(2*a)*p*x^6*(AppellF1[7/6, 1 - p, p, 13/6, E^(2*a)*x^6, -(E^(2*a)*x^6)] + AppellF1[7/6, -p, 1 + p, 13/6, E^(2*a)*x^6, -(E^(2*a)*x^6)]))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6067, 937, 936}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p(a + 3 \log(x)) dx$$

$$\downarrow 6067$$

$$\int (e^{2a}x^6 - 1)^p (e^{2a}x^6 + 1)^{-p} dx$$

$$\downarrow 937$$

$$(1 - e^{2a}x^6)^{-p} (e^{2a}x^6 - 1)^p \int (1 - e^{2a}x^6)^p (e^{2a}x^6 + 1)^{-p} dx$$

$$\downarrow 936$$

$$x(1 - e^{2a}x^6)^{-p} (e^{2a}x^6 - 1)^p \text{AppellF1}\left(\frac{1}{6}, -p, p, \frac{7}{6}, e^{2a}x^6, -e^{2a}x^6\right)$$

input

```
Int[Tanh[a + 3*Log[x]]^p,x]
```

output

```
(x*(-1 + E^(2*a)*x^6))^p*AppellF1[1/6, -p, p, 7/6, E^(2*a)*x^6, -(E^(2*a)*x^6)]/(1 - E^(2*a)*x^6)^p
```

## Definitions of rubi rules used

rule 936 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)
], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 937 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p])
Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}
, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 6067 `Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]`

## Maple [F]

$$\int \tanh(a + 3 \ln(x))^p dx$$

input `int(tanh(a+3*ln(x))^p,x)`

output `int(tanh(a+3*ln(x))^p,x)`

## Fricas [F]

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh(a + 3 \log(x))^p dx$$

input `integrate(tanh(a+3*log(x))^p,x, algorithm="fricas")`

output `integral(tanh(a + 3*log(x))^p, x)`

**Sympy [F]**

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh^p(a + 3 \log(x)) dx$$

input `integrate(tanh(a+3*ln(x))**p,x)`

output `Integral(tanh(a + 3*log(x))**p, x)`

**Maxima [F]**

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh(a + 3 \log(x))^p dx$$

input `integrate(tanh(a+3*log(x))^p,x, algorithm="maxima")`

output `integrate(tanh(a + 3*log(x))^p, x)`

**Giac [F]**

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh(a + 3 \log(x))^p dx$$

input `integrate(tanh(a+3*log(x))^p,x, algorithm="giac")`

output `integrate(tanh(a + 3*log(x))^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh(a + 3 \ln(x))^p dx$$

input `int(tanh(a + 3*log(x))^p,x)`output `int(tanh(a + 3*log(x))^p, x)`**Reduce [F]**

$$\int \tanh^p(a + 3 \log(x)) dx = \tanh(3 \log(x) + a)^p x - 3 \left( \int \frac{\tanh(3 \log(x) + a)^p}{\tanh(3 \log(x) + a)} dx \right)^p + 3 \left( \int \tanh(3 \log(x) + a)^p \tanh(3 \log(x) + a) dx \right)^p$$

input `int(tanh(a+3*log(x))^p,x)`output `tanh(3*log(x) + a)**p*x - 3*int(tanh(3*log(x) + a)**p/tanh(3*log(x) + a),x)*p + 3*int(tanh(3*log(x) + a)**p*tanh(3*log(x) + a),x)*p`

### 3.182 $\int x^3 \tanh(d(a + b \log(cx^n))) dx$

Optimal result	1366
Mathematica [B] (verified)	1366
Rubi [A] (verified)	1367
Maple [F]	1368
Fricas [F]	1369
Sympy [F]	1369
Maxima [F]	1369
Giac [F]	1370
Mupad [F(-1)]	1370
Reduce [F]	1370

#### Optimal result

Integrand size = 17, antiderivative size = 59

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \frac{x^4}{4} - \frac{1}{2}x^4 \operatorname{Hypergeometric2F1}\left(1, \frac{2}{bdn}, 1 + \frac{2}{bdn}, -e^{2ad}(cx^n)^{2bd}\right)$$

output

```
1/4*x^4-1/2*x^4*hypergeom([1, 2/b/d/n], [1+2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 127 vs. 2(59) = 118.

Time = 5.20 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.15

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \frac{x^4 \left( 2e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{2}{bdn}, 2 + \frac{2}{bdn}, -e^{2d(a+b \log(cx^n))}\right) - (2 + bdn) \operatorname{Hypergeometric2F1}\left(1, \frac{2}{bdn}, 1 + \frac{2}{bdn}, -e^{2d(a+b \log(cx^n))}\right) \right)}{8 + 4bdn}$$

input

```
Integrate[x^3*Tanh[d*(a + b*Log[c*x^n])], x]
```

output

$$\frac{(x^4(2E^{(2d(a + b\log(cx^n))})\text{Hypergeometric2F1}[1, 1 + 2/(b*d*n), 2 + 2/(b*d*n), -E^{(2d(a + b\log(cx^n))})] - (2 + b*d*n)\text{Hypergeometric2F1}[1, 2/(b*d*n), 1 + 2/(b*d*n), -E^{(2d(a + b\log(cx^n))})]))/(8 + 4*b*d*n)}$$
**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.61, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^4(cx^n)^{-4/n} \int (cx^n)^{\frac{4}{n}-1} \tanh(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^4(cx^n)^{-4/n} \int \frac{(cx^n)^{\frac{4}{n}-1} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{n}$$

$$\downarrow 959$$

$$\frac{x^4(cx^n)^{-4/n} \left( \frac{1}{4}n(cx^n)^{4/n} - 2 \int \frac{(cx^n)^{\frac{4}{n}-1}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) \right)}{n}$$

$$\downarrow 888$$

$$\frac{x^4(cx^n)^{-4/n} \left( \frac{1}{4}n(cx^n)^{4/n} - \frac{1}{2}n(cx^n)^{4/n} \text{Hypergeometric2F1} \left( 1, \frac{2}{bdn}, 1 + \frac{2}{bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{n}$$

input

$$\text{Int}[x^3 \text{Tanh}[d*(a + b*\text{Log}[c*x^n])], x]$$

output

$$(x^4*((n*(cx^n)^{(4/n)})/4 - (n*(cx^n)^{(4/n})*\text{Hypergeometric2F1}[1, 2/(b*d*n), 1 + 2/(b*d*n), -(E^{(2*a*d)}*(cx^n)^{(2*b*d)})]/2)))/(n*(cx^n)^{(4/n)})$$



## Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 959 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

rule 6073 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] := Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])`

## Maple [F]

$$\int x^3 \tanh(d(a + b \ln(cx^n))) dx$$

input `int(x^3*tanh(d*(a+b*ln(c*x^n))),x)`

output `int(x^3*tanh(d*(a+b*ln(c*x^n))),x)`

**Fricas [F]**

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

output `integral(x^3*tanh(b*d*log(c*x^n) + a*d), x)`

**Sympy [F]**

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh(ad + bd \log(cx^n)) dx$$

input `integrate(x**3*tanh(d*(a+b*ln(c*x**n))),x)`

output `Integral(x**3*tanh(a*d + b*d*log(c*x**n)), x)`

**Maxima [F]**

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

output `1/4*x^4 - 2*integrate(x^3/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)`

**Giac [F]**

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

output `integrate(x^3*tanh((b*log(c*x^n) + a)*d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh(d(a + b \ln(cx^n))) dx$$

input `int(x^3*tanh(d*(a + b*log(c*x^n))),x)`

output `int(x^3*tanh(d*(a + b*log(c*x^n))), x)`

**Reduce [F]**

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = 2e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}x^3}{x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) - \frac{x^4}{4}$$

input `int(x^3*tanh(d*(a+b*log(c*x^n))),x)`

output `(8*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x**3)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) - x**4)/4`

### 3.183 $\int x^2 \tanh(d(a + b \log(cx^n))) dx$

Optimal result	1371
Mathematica [B] (verified)	1371
Rubi [A] (verified)	1372
Maple [F]	1373
Fricas [F]	1374
Sympy [F]	1374
Maxima [F]	1374
Giac [F]	1375
Mupad [F(-1)]	1375
Reduce [F]	1375

#### Optimal result

Integrand size = 17, antiderivative size = 63

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \frac{x^3}{3} - \frac{2}{3}x^3 \operatorname{Hypergeometric2F1}\left(1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)$$

output

`1/3*x^3-2/3*x^3*hypergeom([1, 3/2/b/d/n], [1+3/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 136 vs. 2(63) = 126.

Time = 4.93 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.16

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \frac{x^3 \left( 3e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{3}{2bdn}, 2 + \frac{3}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) - (3 + 2bdn) \operatorname{Hypergeometric2F1}\left(1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) \right)}{9 + 6bdn}$$

input

`Integrate[x^2*Tanh[d*(a + b*Log[c*x^n])],x]`

output

```
(x^3*(3*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 3/(2*b*d*n), 2 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - (3 + 2*b*d*n)*Hypergeometric2F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])))/(9 + 6*b*d*n)
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.57, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^3 (cx^n)^{-3/n} \int (cx^n)^{\frac{3}{n}-1} \tanh(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^3 (cx^n)^{-3/n} \int \frac{(cx^n)^{\frac{3}{n}-1} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{n}$$

$$\downarrow 959$$

$$\frac{x^3 (cx^n)^{-3/n} \left( \frac{1}{3} n (cx^n)^{3/n} - 2 \int \frac{(cx^n)^{\frac{3}{n}-1}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) \right)}{n}$$

$$\downarrow 888$$

$$\frac{x^3 (cx^n)^{-3/n} \left( \frac{1}{3} n (cx^n)^{3/n} - \frac{2}{3} n (cx^n)^{3/n} \text{Hypergeometric2F1} \left( 1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{n}$$

input

```
Int[x^2*Tanh[d*(a + b*Log[c*x^n])],x]
```

output

```
(x^3*((n*(c*x^n)^(3/n))/3 - (2*n*(c*x^n)^(3/n)*Hypergeometric2F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/3))/3)/(n*(c*x^n)^(3/n))
```

### Defintions of rubi rules used

rule 888

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 959

```
Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

rule 6071

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]
```

rule 6073

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] := Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

### Maple [F]

$$\int x^2 \tanh(d(a + b \ln(cx^n))) dx$$

input

```
int(x^2*tanh(d*(a+b*ln(c*x^n))),x)
```

output `int(x^2*tanh(d*(a+b*ln(c*x^n))),x)`

### Fricas [F]

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

output `integral(x^2*tanh(b*d*log(c*x^n) + a*d), x)`

### Sympy [F]

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh(ad + bd \log(cx^n)) dx$$

input `integrate(x**2*tanh(d*(a+b*ln(c*x**n))),x)`

output `Integral(x**2*tanh(a*d + b*d*log(c*x**n)), x)`

### Maxima [F]

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

output `1/3*x^3 - 2*integrate(x^2/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)`

**Giac [F]**

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

output `integrate(x^2*tanh((b*log(c*x^n) + a)*d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh(d(a + b \ln(cx^n))) dx$$

input `int(x^2*tanh(d*(a + b*log(c*x^n))),x)`

output `int(x^2*tanh(d*(a + b*log(c*x^n))), x)`

**Reduce [F]**

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = 2e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}x^2}{x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) - \frac{x^3}{3}$$

input `int(x^2*tanh(d*(a+b*log(c*x^n))),x)`

output `(6*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x**2)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) - x**3)/3`



### 3.184 $\int x \tanh (d(a + b \log (c x^n))) dx$

Optimal result	1376
Mathematica [B] (verified)	1376
Rubi [A] (verified)	1377
Maple [F]	1378
Fricas [F]	1379
Sympy [F]	1379
Maxima [F]	1379
Giac [F]	1380
Mupad [F(-1)]	1380
Reduce [F]	1380

#### Optimal result

Integrand size = 15, antiderivative size = 55

$$\int x \tanh (d(a + b \log (c x^n))) dx = \frac{x^2}{2} - x^2 \operatorname{Hypergeometric2F1} \left( 1, \frac{1}{bdn}, 1 + \frac{1}{bdn}, -e^{2ad}(c x^n)^{2bd} \right)$$

output

```
1/2*x^2-x^2*hypergeom([1, 1/b/d/n],[1+1/b/d/n],-exp(2*a*d)*(c*x^n)^(2*b*d))
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 122 vs. 2(55) = 110.

Time = 4.98 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.22

$$\int x \tanh (d(a + b \log (c x^n))) dx = \frac{x^2 \left( e^{2d(a+b \log (c x^n))} \operatorname{Hypergeometric2F1} \left( 1, 1 + \frac{1}{bdn}, 2 + \frac{1}{bdn}, -e^{2d(a+b \log (c x^n))} \right) - (1 + bdn) \operatorname{Hypergeometric2F1} \left( 1, \frac{1}{bdn}, 1 + \frac{1}{bdn}, -e^{2d(a+b \log (c x^n))} \right) \right)}{2 + 2bdn}$$

input

```
Integrate[x*Tanh[d*(a + b*Log[c*x^n])],x]
```

output

$$\frac{(x^2(E^{2d(a+b\log(cx^n))})\text{Hypergeometric2F1}[1, 1+1/(bdn), 2+1/(bdn), -E^{2d(a+b\log(cx^n))}] - (1+bdn)\text{Hypergeometric2F1}[1, 1/(bdn), 1+1/(bdn), -E^{2d(a+b\log(cx^n))}]))/2+2bdn}$$
**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.65, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \tanh(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^2(cx^n)^{-2/n} \int (cx^n)^{\frac{2}{n}-1} \tanh(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^2(cx^n)^{-2/n} \int \frac{(cx^n)^{\frac{2}{n}-1} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{n}$$

$$\downarrow 959$$

$$\frac{x^2(cx^n)^{-2/n} \left( \frac{1}{2}n(cx^n)^{2/n} - 2 \int \frac{(cx^n)^{\frac{2}{n}-1}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) \right)}{n}$$

$$\downarrow 888$$

$$\frac{x^2(cx^n)^{-2/n} \left( \frac{1}{2}n(cx^n)^{2/n} - n(cx^n)^{2/n} \text{Hypergeometric2F1} \left( 1, \frac{1}{bdn}, 1 + \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{n}$$

input

$$\text{Int}[x*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])], x]$$

output

$$\frac{(x^2*((n*(cx^n)^{(2/n)})/2 - n*(cx^n)^{(2/n)}*\text{Hypergeometric2F1}[1, 1/(bdn), 1 + 1/(bdn), -(E^{2ad})*(cx^n)^{(2bd)}]))/(n*(cx^n)^{(2/n)})}$$

## Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 959 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]`

rule 6073 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] := Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])`

## Maple [F]

$$\int x \tanh(d(a + b \ln(cx^n))) dx$$

input `int(x*tanh(d*(a+b*ln(c*x^n))),x)`

output `int(x*tanh(d*(a+b*ln(c*x^n))),x)`

**Fricas [F]**

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

output `integral(x*tanh(b*d*log(c*x^n) + a*d), x)`

**Sympy [F]**

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh(ad + bd \log(cx^n)) dx$$

input `integrate(x*tanh(d*(a+b*ln(c*x**n))),x)`

output `Integral(x*tanh(a*d + b*d*log(c*x**n)), x)`

**Maxima [F]**

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

output `1/2*x^2 - 2*integrate(x/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)`

**Giac [F]**

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

output `integrate(x*tanh((b*log(c*x^n) + a)*d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh(d(a + b \ln(cx^n))) dx$$

input `int(x*tanh(d*(a + b*log(c*x^n))),x)`

output `int(x*tanh(d*(a + b*log(c*x^n))), x)`

**Reduce [F]**

$$\int x \tanh(d(a + b \log(cx^n))) dx = 2e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn} x}{x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) - \frac{x^2}{2}$$

input `int(x*tanh(d*(a+b*log(c*x^n))),x)`

output `(4*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) - x**2)/2`

### 3.185 $\int \tanh (d(a + b \log (c x^n))) dx$

Optimal result	1381
Mathematica [B] (verified)	1381
Rubi [A] (verified)	1382
Maple [F]	1383
Fricas [F]	1384
Sympy [F]	1384
Maxima [F]	1384
Giac [F]	1385
Mupad [F(-1)]	1385
Reduce [F]	1385

#### Optimal result

Integrand size = 13, antiderivative size = 53

$$\int \tanh (d(a + b \log (c x^n))) dx = x - 2x \operatorname{Hypergeometric2F1} \left( 1, \frac{1}{2bdn}, 1 + \frac{1}{2bdn}, -e^{2ad}(c x^n)^{2bd} \right)$$

output

`x-2*x*hypergeom([1, 1/2/b/d/n], [1+1/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 126 vs. 2(53) = 106.

Time = 5.65 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.38

$$\begin{aligned} & \int \tanh (d(a + b \log (c x^n))) dx \\ &= \frac{e^{2d(a+b \log (c x^n))} x \operatorname{Hypergeometric2F1} \left( 1, 1 + \frac{1}{2bdn}, 2 + \frac{1}{2bdn}, -e^{2d(a+b \log (c x^n))} \right)}{1 + 2bdn} \\ & \quad - x \operatorname{Hypergeometric2F1} \left( 1, \frac{1}{2bdn}, 1 + \frac{1}{2bdn}, -e^{2d(a+b \log (c x^n))} \right) \end{aligned}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])],x]`

output  $(E^{(2*d*(a + b*Log[c*x^n]))}*x*Hypergeometric2F1[1, 1 + 1/(2*b*d*n), 2 + 1/(2*b*d*n), -E^{(2*d*(a + b*Log[c*x^n]))}]/(1 + 2*b*d*n) - x*Hypergeometric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -E^{(2*d*(a + b*Log[c*x^n]))}])$

### Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.66, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6069, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(d(a + b \log(cx^n))) dx \\
 & \quad \downarrow \text{6069} \\
 & \frac{x(cx^n)^{-1/n} \int (cx^n)^{\frac{1}{n}-1} \tanh(d(a + b \log(cx^n))) d(cx^n)}{n} \\
 & \quad \downarrow \text{6071} \\
 & \frac{x(cx^n)^{-1/n} \int \frac{(cx^n)^{\frac{1}{n}-1} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{n} \\
 & \quad \downarrow \text{959} \\
 & \frac{x(cx^n)^{-1/n} \left( n(cx^n)^{\frac{1}{n}} - 2 \int \frac{(cx^n)^{\frac{1}{n}-1}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) \right)}{n} \\
 & \quad \downarrow \text{888} \\
 & \frac{x(cx^n)^{-1/n} \left( n(cx^n)^{\frac{1}{n}} - 2n(cx^n)^{\frac{1}{n}} \text{Hypergeometric2F1} \left( 1, \frac{1}{2bdn}, 1 + \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{n}
 \end{aligned}$$

input `Int [Tanh[d*(a + b*Log[c*x^n])],x]`

output  $(x*(n*(c*x^n)^n^{(-1)} - 2*n*(c*x^n)^n^{(-1)}*Hypergeometric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^{(2*b*d)}]))/(n*(c*x^n)^n^{(-1)})$

### Defintions of rubi rules used

rule 888  $\text{Int}[(c_*)(x_*)^{(m_*)}*((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /;$   $\text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[(e_*)(x_*)^{(m_*)}*((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)(x_*)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m + n*(p+1) + 1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m + n*(p+1) + 1))/(b*(m + n*(p+1) + 1)) \ \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 6069  $\text{Int}[\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_*)^{(n_*)}*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[x/(n*(c*x^n)^{(1/n)}) \ \text{Subst}[\text{Int}[x^{(1/n - 1)}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /;$   $\text{FreeQ}\{a, b, c, d, n, p\}, x \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

rule 6071  $\text{Int}[(e_*)(x_*)^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]* (b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p], x] /;$   $\text{FreeQ}\{a, b, d, e, m, p\}, x]$

### Maple [F]

$$\int \tanh(d(a + b \ln(cx^n))) dx$$

input  $\text{int}(\tanh(d*(a+b*\ln(c*x^n))),x)$

output  $\text{int}(\tanh(d*(a+b*\ln(c*x^n))),x)$



**Fricas [F]**

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d), x)`

**Sympy [F]**

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \log(cx^n))) dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n))),x)`

output `Integral(tanh(d*(a + b*log(c*x**n))), x)`

**Maxima [F]**

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

output `x - 2*integrate(1/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)`

**Giac [F]**

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d) dx$$

input `integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n))) dx$$

input `int(tanh(d*(a + b*log(c*x^n))),x)`

output `int(tanh(d*(a + b*log(c*x^n))), x)`

**Reduce [F]**

$$\int \tanh(d(a + b \log(cx^n))) dx = 2e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) - x$$

input `int(tanh(d*(a+b*log(c*x^n))),x)`

output `2*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) - x`

$$3.186 \quad \int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx$$

Optimal result	1386
Mathematica [A] (verified)	1386
Rubi [A] (verified)	1387
Maple [A] (verified)	1388
Fricas [B] (verification not implemented)	1389
Sympy [A] (verification not implemented)	1389
Maxima [A] (verification not implemented)	1390
Giac [B] (verification not implemented)	1390
Mupad [B] (verification not implemented)	1391
Reduce [B] (verification not implemented)	1391

### Optimal result

Integrand size = 17, antiderivative size = 25

$$\int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx = \frac{\log(\cosh(ad + bd \log(cx^n)))}{bdn}$$

output

```
ln(cosh(a*d+b*d*ln(c*x^n)))/b/d/n
```

### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx = \frac{\log(\cosh(d(a+b \log(cx^n))))}{bdn}$$

input

```
Integrate[Tanh[d*(a + b*Log[c*x^n])]/x,x]
```

output

```
Log[Cosh[d*(a + b*Log[c*x^n])]]/(b*d*n)
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {3039, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx \\
 \downarrow \text{3039} \\
 \int \frac{\tanh(d(a + b \log(cx^n))) d \log(cx^n)}{n} \\
 \downarrow \text{3042} \\
 \int \frac{-i \tan(iad + ib \log(cx^n) d) d \log(cx^n)}{n} \\
 \downarrow \text{26} \\
 \frac{i \int \tan(iad + ib \log(cx^n) d) d \log(cx^n)}{n} \\
 \downarrow \text{3956} \\
 \frac{\log(\cosh(ad + bd \log(cx^n)))}{bdn}
 \end{array}$$

input `Int[Tanh[d*(a + b*Log[c*x^n])]/x,x]`

output `Log[Cosh[a*d + b*d*Log[c*x^n]]]/(b*d*n)`

## Definitions of rubi rules used

- rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F x_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3039  $\text{Int}[u_, x\_Symbol] \rightarrow \text{With}[\{\text{lst} = \text{FunctionOfLog}[\text{Cancel}[x*u], x]\}, \text{Simp}[1/\text{lst}[\text{[3]}] \text{Subst}[\text{Int}[\text{lst}[\text{[1]}], x], x, \text{Log}[\text{lst}[\text{[2]}]]], x] /; \text{!FalseQ}[\text{lst}] /; \text{NonsumQ}[u]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956  $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

## Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

method	result
derivativdivides	$\frac{\ln(\cosh(d(a+b \ln(cx^n))))}{nbd}$
default	$\frac{\ln(\cosh(d(a+b \ln(cx^n))))}{nbd}$
parallelrisch	$-\frac{\ln(x)dbn + \ln(1 - \tanh(d(a+b \ln(cx^n))))}{dbn}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2 \ln(c)}{n} - \frac{2 \ln(x^n)}{n} - \frac{i\pi \text{csgn}(ix^n) \text{csgn}(icx^n)^2}{n} + \frac{i\pi \text{csgn}(ix^n) \text{csgn}(icx^n) \text{csgn}(ic)}{n} + \frac{i\pi \text{csgn}(ic)}{n}$

input  $\text{int}(\tanh(d*(a+b*\ln(c*x^n)))/x,x,\text{method}=\_RETURNVERBOSE)$

output  $1/n/b/d*\ln(\cosh(d*(a+b*\ln(c*x^n))))$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs.  $2(25) = 50$ .

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 3.04

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx$$

$$= -\frac{bdn \log(x) - \log\left(\frac{2 \cosh(bdn \log(x) + bd \log(c) + ad)}{\cosh(bdn \log(x) + bd \log(c) + ad) - \sinh(bdn \log(x) + bd \log(c) + ad)}\right)}{bdn}$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="fricas")`

output `-(b*d*n*log(x) - log(2*cosh(b*d*n*log(x) + b*d*log(c) + a*d)/(cosh(b*d*n*log(x) + b*d*log(c) + a*d) - sinh(b*d*n*log(x) + b*d*log(c) + a*d)))/(b*d*n)`

**Sympy [A] (verification not implemented)**

Time = 1.39 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx = -\frac{\log(bdn \tanh^2(ad + bd \log(cx^n)) - bdn)}{2bdn}$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))/x,x)`

output `-log(b*d*n*tanh(a*d + b*d*log(c*x**n))**2 - b*d*n)/(2*b*d*n)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx = \frac{\log(\cosh((b \log(cx^n) + a)d))}{bdn}$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="maxima")`

output `log(cosh((b*log(c*x^n) + a)*d))/(b*d*n)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 88 vs. 2(25) = 50.

Time = 0.20 (sec) , antiderivative size = 88, normalized size of antiderivative = 3.52

$$\begin{aligned} & \int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx \\ &= -\frac{\log(x^{bdn})}{bdn} \\ & \quad + \frac{\log\left(\sqrt{2x^{2bdn}|c|^{2bd} \cos(\pi b d \operatorname{sgn}(c) - \pi b d) e^{(2ad)} + x^{4bdn}|c|^{4bd} e^{(4ad)} + 1}\right)}{bdn} \end{aligned}$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="giac")`

output `-log(x^(b*d*n))/(b*d*n) + log(sqrt(2*x^(2*b*d*n)*abs(c)^(2*b*d)*cos(pi*b*d*sgn(c) - pi*b*d)*e^(2*a*d) + x^(4*b*d*n)*abs(c)^(4*b*d)*e^(4*a*d) + 1))/(b*d*n)`

**Mupad [B] (verification not implemented)**

Time = 2.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx = \frac{\ln(e^{2ad}(cx^n)^{2bd} + 1)}{bdn} - \ln(x)$$

input `int(tanh(d*(a + b*log(c*x^n)))/x,x)`output `log(exp(2*a*d)*(c*x^n)^(2*b*d) + 1)/(b*d*n) - log(x)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx = \frac{\log(x^{2bdn}e^{2ad}c^{2bd} + 1) - \log(x) bdn}{bdn}$$

input `int(tanh(d*(a+b*log(c*x^n)))/x,x)`output `(log(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1) - log(x)*b*d*n)/(b*d*n)`



**3.187**  $\int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$

Optimal result	1392
Mathematica [B] (verified)	1392
Rubi [A] (verified)	1393
Maple [F]	1394
Fricas [F]	1395
Sympy [F]	1395
Maxima [F]	1395
Giac [F]	1396
Mupad [F(-1)]	1396
Reduce [F]	1396

**Optimal result**

Integrand size = 17, antiderivative size = 59

$$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$$

$$= -\frac{1}{x} + \frac{2 \operatorname{Hypergeometric2F1}\left(1, -\frac{1}{2bdn}, 1 - \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{x}$$

output

```
-1/x+2*hypergeom([1, -1/2/b/d/n], [1-1/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/x
```

**Mathematica [B] (verified)**

Leaf count is larger than twice the leaf count of optimal. 126 vs. 2(59) = 118.

Time = 2.06 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.14

$$\int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$$

$$= \frac{e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 - \frac{1}{2bdn}, 2 - \frac{1}{2bdn}, -e^{2d(a+b \log(cx^n))}\right)}{-1+2bdn} + \operatorname{Hypergeometric2F1}\left(1, -\frac{1}{2bdn}, 1 - \frac{1}{2bdn}, -e^{2d(a+b \log(cx^n))}\right)$$

input

```
Integrate[Tanh[d*(a + b*Log[c*x^n])]/x^2,x]
```

output

```
((E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(2*b*d*n), 2 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))])/(-1 + 2*b*d*n) + Hypergeometric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))])/x
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.58, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx$$

$$\downarrow 6073$$

$$\frac{(cx^n)^{\frac{1}{n}} \int (cx^n)^{-1-\frac{1}{n}} \tanh(d(a + b \log(cx^n))) d(cx^n)}{nx}$$

$$\downarrow 6071$$

$$\frac{(cx^n)^{\frac{1}{n}} \int \frac{(cx^n)^{-1-\frac{1}{n}} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{nx}$$

$$\downarrow 959$$

$$\frac{(cx^n)^{\frac{1}{n}} \left( -2 \int \frac{(cx^n)^{-1-\frac{1}{n}}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) - n(cx^n)^{-1/n} \right)}{nx}$$

$$\downarrow 888$$

$$\frac{(cx^n)^{\frac{1}{n}} \left( 2n(cx^n)^{-1/n} \text{Hypergeometric2F1} \left( 1, -\frac{1}{2bdn}, 1 - \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) - n(cx^n)^{-1/n} \right)}{nx}$$

input

```
Int [Tanh [d*(a + b*Log[c*x^n])] / x^2, x]
```

output 
$$\frac{((c*x^n)^n)^{-1}*(-(n/(c*x^n)^n)^{-1}) + (2*n*Hypergeometric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})])/(c*x^n)^n)^{-1}}{(n*x)}$$

### Defintions of rubi rules used

rule 888 
$$\text{Int}[\frac{((c\_)*(x\_))^{(m\_)}*((a\_)+(b\_)*(x\_)^{(n\_))^{(p\_)}}, x\_Symbol]}{((c*x)^{(m+1)}(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)]}, x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}\{p, 0\} \ \&\& \ (\text{ILtQ}\{p, 0\} \ || \ \text{GtQ}\{a, 0\})$$

rule 959 
$$\text{Int}[\frac{((e\_)*(x\_))^{(m\_)}*((a\_)+(b\_)*(x\_)^{(n\_))^{(p\_)}*((c\_)+(d\_)*(x\_)^{(n\_)}), x\_Symbol]}{d*(e*x)^{(m+1)}*((a+b*x^n)^{(p+1)}(b*e*(m+n*(p+1)+1)))}, x] - \text{Simp}[\frac{a*d*(m+1) - b*c*(m+n*(p+1)+1)}{b*(m+n*(p+1)+1)} \text{Int}[(e*x)^m*(a+b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}\{b*c - a*d, 0\} \ \&\& \ \text{NeQ}\{m+n*(p+1)+1, 0\}$$

rule 6071 
$$\text{Int}[\frac{((e\_)*(x\_))^{(m\_)}*\text{Tanh}[\frac{(a\_)+\text{Log}[x]*(b\_)]*(d\_)]^{(p\_)}}, x\_Symbol]}{(e*x)^m*((-1+E^{(2*a*d)}*x^{(2*b*d)})^p/(1+E^{(2*a*d)}*x^{(2*b*d)})^p)}, x] /; \text{FreeQ}\{a, b, d, e, m, p\}, x]$$

rule 6073 
$$\text{Int}[\frac{((e\_)*(x\_))^{(m\_)}*\text{Tanh}[\frac{(a\_)+\text{Log}[(c\_)*(x\_)^{(n\_)}]*(b\_)]*(d\_)]^{(p\_)}}, x\_Symbol]}{(e*x)^{(m+1)}(e*n*(c*x^n)^{(m+1)/n})} \text{Subst}[\text{Int}[x^{(m+1)/n-1}*\text{Tanh}[d*(a+b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ (\text{NeQ}\{c, 1\} \ || \ \text{NeQ}\{n, 1\})$$

### Maple [F]

$$\int \frac{\tanh(d(a+b \ln(cx^n)))}{x^2} dx$$

input 
$$\text{int}(\tanh(d*(a+b*\ln(c*x^n)))/x^2, x)$$

output `int(tanh(d*(a+b*ln(c*x^n)))/x^2,x)`

### Fricas [F]

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)/x^2, x)`

### Sympy [F]

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh(ad + bd \log(cx^n))}{x^2} dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))/x**2,x)`

output `Integral(tanh(a*d + b*d*log(c*x**n))/x**2, x)`

### Maxima [F]

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="maxima")`

output `-1/x - 2*integrate(1/(c^(2*b*d)*x^2*e^(2*b*d*log(x^n) + 2*a*d) + x^2), x)`

**Giac [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)/x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh(d(a + b \ln(cx^n)))}{x^2} dx$$

input `int(tanh(d*(a + b*log(c*x^n)))/x^2,x)`

output `int(tanh(d*(a + b*log(c*x^n)))/x^2, x)`

**Reduce [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \frac{2e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{2bdn}e^{2ad}c^{2bd}x^2+x^2} dx \right) x + 1}{x}$$

input `int(tanh(d*(a+b*log(c*x^n)))/x^2,x)`

output `(2*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*x**2 + x**2),x)*x + 1)/x`

### 3.188 $\int \frac{\tanh(d(a+b \log(cx^n)))}{x^3} dx$

Optimal result	1397
Mathematica [B] (verified)	1397
Rubi [A] (verified)	1398
Maple [F]	1399
Fricas [F]	1400
Sympy [F]	1400
Maxima [F]	1400
Giac [F]	1401
Mupad [F(-1)]	1401
Reduce [F]	1401

#### Optimal result

Integrand size = 17, antiderivative size = 56

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx$$

$$= -\frac{1}{2x^2} + \frac{\text{Hypergeometric2F1}\left(1, -\frac{1}{bdn}, 1 - \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{x^2}$$

output

```
-1/2/x^2+hypergeom([1, -1/b/d/n], [1-1/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/x^2
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 2.06 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.14

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx$$

$$= \frac{e^{2d(a+b \log(cx^n))} \text{Hypergeometric2F1}\left(1, 1 - \frac{1}{bdn}, 2 - \frac{1}{bdn}, -e^{2d(a+b \log(cx^n))}\right)}{-1+bdn} + \frac{\text{Hypergeometric2F1}\left(1, -\frac{1}{bdn}, 1 - \frac{1}{bdn}, -e^{2d(a+b \log(cx^n))}\right)}{2x^2}$$

input

```
Integrate[Tanh[d*(a + b*Log[c*x^n])]/x^3,x]
```

output

$$\frac{((E^{(2*d*(a + b*\text{Log}[c*x^n]))})*\text{Hypergeometric2F1}[1, 1 - 1/(b*d*n), 2 - 1/(b*d*n), -E^{(2*d*(a + b*\text{Log}[c*x^n]))}]))/(-1 + b*d*n) + \text{Hypergeometric2F1}[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -E^{(2*d*(a + b*\text{Log}[c*x^n]))}])/(2*x^2)}$$
**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.64, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx \\ & \quad \downarrow \text{6073} \\ & \frac{(cx^n)^{2/n} \int (cx^n)^{-1-\frac{2}{n}} \tanh(d(a + b \log(cx^n))) d(cx^n)}{nx^2} \\ & \quad \downarrow \text{6071} \\ & \frac{(cx^n)^{2/n} \int \frac{(cx^n)^{-1-\frac{2}{n}} (e^{2ad}(cx^n)^{2bd} - 1)}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{nx^2} \\ & \quad \downarrow \text{959} \\ & \frac{(cx^n)^{2/n} \left( -2 \int \frac{(cx^n)^{-1-\frac{2}{n}}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n) - \frac{1}{2} n (cx^n)^{-2/n} \right)}{nx^2} \\ & \quad \downarrow \text{888} \\ & \frac{(cx^n)^{2/n} \left( n (cx^n)^{-2/n} \text{Hypergeometric2F1} \left( 1, -\frac{1}{bdn}, 1 - \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd} \right) - \frac{1}{2} n (cx^n)^{-2/n} \right)}{nx^2} \end{aligned}$$

input

$$\text{Int}[\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]/x^3, x]$$

output

$$\frac{((c*x^n)^{(2/n)}*(-1/2*n/(c*x^n)^{(2/n)} + (n*\text{Hypergeometric2F1}[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})])/(c*x^n)^{(2/n)})))/(n*x^2)}$$

## Definitions of rubi rules used

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 959 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]`

rule 6073 `Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] := Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])`

## Maple [F]

$$\int \frac{\tanh(d(a + b \ln(cx^n)))}{x^3} dx$$

input `int(tanh(d*(a+b*ln(c*x^n)))/x^3,x)`

output `int(tanh(d*(a+b*ln(c*x^n)))/x^3,x)`



**Fricas [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)/x^3, x)`

**Sympy [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh(ad + bd \log(cx^n))}{x^3} dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))/x**3,x)`

output `Integral(tanh(a*d + b*d*log(c*x**n))/x**3, x)`

**Maxima [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="maxima")`

output `-1/2/x^2 - 2*integrate(1/(c^(2*b*d)*x^3*e^(2*b*d*log(x^n) + 2*a*d) + x^3), x)`

**Giac [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh(d(a + b \ln(cx^n)))}{x^3} dx$$

input `int(tanh(d*(a + b*log(c*x^n)))/x^3,x)`

output `int(tanh(d*(a + b*log(c*x^n)))/x^3, x)`

**Reduce [F]**

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \frac{4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{2bdn}e^{2ad}c^{2bd}x^3+x^3} dx \right) x^2 + 1}{2x^2}$$

input `int(tanh(d*(a+b*log(c*x^n)))/x^3,x)`

output `(4*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*x**3 + x**3),x)*x**2 + 1)/(2*x**2)`

### 3.189 $\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$

Optimal result	1402
Mathematica [A] (verified)	1403
Rubi [A] (verified)	1403
Maple [F]	1406
Fricas [F]	1406
Sympy [F]	1406
Maxima [F]	1407
Giac [F]	1407
Mupad [F(-1)]	1407
Reduce [F]	1408

#### Optimal result

Integrand size = 19, antiderivative size = 133

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{1}{4} \left(1 + \frac{4}{bdn}\right) x^4 + \frac{x^4 (1 - e^{2ad}(cx^n)^{2bd})}{bdn (1 + e^{2ad}(cx^n)^{2bd})}$$

$$- \frac{2x^4 \operatorname{Hypergeometric2F1}\left(1, \frac{2}{bdn}, 1 + \frac{2}{bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

output

```
1/4*(1+4/b/d/n)*x^4+x^4*(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/(1+exp(2*a*d)
*(c*x^n)^(2*b*d))-2*x^4*hypergeom([1, 2/b/d/n],[1+2/b/d/n],-exp(2*a*d)*(c*
x^n)^(2*b*d))/b/d/n
```

**Mathematica [A] (verified)**

Time = 5.42 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.20

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{x^4 (8e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{2}{bdn}, 2 + \frac{2}{bdn}, -e^{2d(a+b \log(cx^n))}\right) + (2 + bdn)(bdn - 4 \operatorname{Hypergeometric2F1}\left[1, \frac{2}{(b*d*n)}, 1 + \frac{2}{(b*d*n)}, -E^{(2*d*(a + b*Log[c*x^n])}\right] - 4 \operatorname{Tanh}[d*(a + b*Log[c*x^n])])\right)}{4bdn(2 + bdn)}$$

input

```
Integrate[x^3*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

output

```
(x^4*(8*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 2/(b*d*n), 2 + 2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (2 + b*d*n)*(b*d*n - 4*Hypergeometric2F1[1, 2/(b*d*n), 1 + 2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - 4*Tanh[d*(a + b*Log[c*x^n])]))/(4*b*d*n*(2 + b*d*n))
```

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.38, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^4 (cx^n)^{-4/n} \int (cx^n)^{\frac{4}{n}-1} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^4 (cx^n)^{-4/n} \int \frac{(cx^n)^{\frac{4}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)^2}{(e^{2ad}(cx^n)^{2bd}+1)^2} d(cx^n)}{n}$$

$$\downarrow 1004$$

$$x^4 (cx^n)^{-4/n} \left( \frac{(cx^n)^{4/n} (1 - e^{2ad} (cx^n)^{2bd})}{bd (e^{2ad} (cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{4}{n}-1} \left( \frac{e^{2ad} (4-bdn)}{n} - \frac{e^{4ad} (bdn+4) (cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad} (cx^n)^{2bd} + 1}}{2bd} \right)$$

$n$   
↓ 27

$$x^4 (cx^n)^{-4/n} \left( \frac{(cx^n)^{4/n} (1 - e^{2ad} (cx^n)^{2bd})}{bd (e^{2ad} (cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{4}{n}-1} \left( \frac{e^{2ad} (4-bdn)}{n} - \frac{e^{4ad} (bdn+4) (cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad} (cx^n)^{2bd} + 1}}{bd} \right)$$

$n$   
↓ 959

$$x^4 (cx^n)^{-4/n} \left( \frac{(cx^n)^{4/n} (1 - e^{2ad} (cx^n)^{2bd})}{bd (e^{2ad} (cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( \frac{8e^{2ad} \int \frac{(cx^n)^{\frac{4}{n}-1} d(cx^n)}{e^{2ad} (cx^n)^{2bd} + 1}}{n} - \frac{1}{4} e^{2ad} (bdn+4) (cx^n)^{4/n} \right)}{bd} \right)$$

$n$   
↓ 888

$$x^4 (cx^n)^{-4/n} \left( \frac{(cx^n)^{4/n} (1 - e^{2ad} (cx^n)^{2bd})}{bd (e^{2ad} (cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( 2e^{2ad} (cx^n)^{4/n} \text{Hypergeometric2F1} \left( 1, \frac{2}{bdn}, 1 + \frac{2}{bdn}, -e^{2ad} (cx^n)^{2bd} \right) - \frac{1}{4} e^{2ad} (bdn+4) (cx^n)^{4/n} \right)}{bd} \right)$$

$n$

input `Int [x^3*Tanh [d*(a + b*Log [c*x^n])]^2,x]`

output `(x^4*(((c*x^n)^(4/n)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d)))/(b*d*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) - (-1/4*(E^(2*a*d)*(4 + b*d*n)*(c*x^n)^(4/n)) + 2*E^(2*a*d)*(c*x^n)^(4/n)*Hypergeometric2F1[1, 2/(b*d*n), 1 + 2/(b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/(b*d*E^(2*a*d))))/(n*(c*x^n)^(4/n))`

## Definitions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(a*b*e*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_*)}*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{(m+1)/n}) \text{ Subst}[\text{Int}[x^{(m+1)/n-1}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

**Maple [F]**

$$\int x^3 \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x^3*tanh(d*(a+b*ln(c*x^n)))^2,x)`

output `int(x^3*tanh(d*(a+b*ln(c*x^n)))^2,x)`

**Fricas [F]**

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

output `integral(x^3*tanh(b*d*log(c*x^n) + a*d)^2, x)`

**Sympy [F]**

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh^2(ad + bd \log(cx^n)) dx$$

input `integrate(x**3*tanh(d*(a+b*ln(c*x**n)))**2,x)`

output `Integral(x**3*tanh(a*d + b*d*log(c*x**n))**2, x)`

**Maxima [F]**

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

output `1/4*(b*c^(2*b*d)*d*n*x^4*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 8)*x^4)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 8*integrate(x^3/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

**Giac [F]**

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

output `integrate(x^3*tanh((b*log(c*x^n) + a)*d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x^3*tanh(d*(a + b*log(c*x^n)))^2,x)`

output `int(x^3*tanh(d*(a + b*log(c*x^n)))^2, x)`



**Reduce [F]**

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = -4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}x^3}{x^{4bdn}e^{4ad}c^{4bd} + 2x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) + \frac{x^4}{4}$$

input `int(x^3*tanh(d*(a+b*log(c*x^n)))^2,x)`

output `( - 16*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x**3)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) + x**4)/4`

### 3.190 $\int x^2 \tanh^2 (d(a + b \log (cx^n))) dx$

Optimal result	1409
Mathematica [A] (verified)	1410
Rubi [A] (verified)	1410
Maple [F]	1413
Fricas [F]	1413
Sympy [F]	1413
Maxima [F]	1414
Giac [F]	1414
Mupad [F(-1)]	1414
Reduce [F]	1415

#### Optimal result

Integrand size = 19, antiderivative size = 137

$$\int x^2 \tanh^2 (d(a + b \log (cx^n))) dx$$

$$= \frac{1}{3} \left( 1 + \frac{3}{bdn} \right) x^3 + \frac{x^3 \left( 1 - e^{2ad} (cx^n)^{2bd} \right)}{bdn \left( 1 + e^{2ad} (cx^n)^{2bd} \right)}$$

$$- \frac{2x^3 \operatorname{Hypergeometric2F1} \left( 1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2ad} (cx^n)^{2bd} \right)}{bdn}$$

output

```
1/3*(1+3/b/d/n)*x^3+x^3*(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/(1+exp(2*a*d)
*(c*x^n)^(2*b*d))-2*x^3*hypergeom([1, 3/2/b/d/n],[1+3/2/b/d/n],-exp(2*a*d)
*(c*x^n)^(2*b*d))/b/d/n
```

**Mathematica [A] (verified)**

Time = 5.17 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.23

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{x^3 (9e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{3}{2bdn}, 2 + \frac{3}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) + (3 + 2bdn)(bdn - 3 \operatorname{Hypergeometric2F1}\left(1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2d(a+b \log(cx^n))}\right)) - 3 \operatorname{Tanh}[d(a + b \log(cx^n))])}{3bdn(3 + 2bdn)}$$

input

```
Integrate[x^2*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

output

```
(x^3*(9*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 3/(2*b*d*n), 2 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (3 + 2*b*d*n)*(b*d*n - 3*Hypergeometric2F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - 3*Tanh[d*(a + b*Log[c*x^n])]))/(3*b*d*n*(3 + 2*b*d*n))
```

**Rubi [A] (verified)**Time = 0.48 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.36, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^3 (cx^n)^{-3/n} \int (cx^n)^{\frac{3}{n}-1} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^3 (cx^n)^{-3/n} \int \frac{(cx^n)^{\frac{3}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)^2}{(e^{2ad}(cx^n)^{2bd}+1)^2} d(cx^n)}{n}$$

$$\downarrow 1004$$

$$x^3(cx^n)^{-3/n} \left( \frac{(cx^n)^{3/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{3}{n}-1} \left( \frac{e^{2ad}(3-bdn)}{n} - \frac{e^{4ad}(bdn+3)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{2bd} \right)$$

$n$   
↓ 27

$$x^3(cx^n)^{-3/n} \left( \frac{(cx^n)^{3/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{3}{n}-1} \left( \frac{e^{2ad}(3-bdn)}{n} - \frac{e^{4ad}(bdn+3)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{bd} \right)$$

$n$   
↓ 959

$$x^3(cx^n)^{-3/n} \left( \frac{(cx^n)^{3/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( \frac{6e^{2ad} \int \frac{(cx^n)^{\frac{3}{n}-1} d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{n} - \frac{1}{3} e^{2ad}(bdn+3)(cx^n)^{3/n} \right)}{bd} \right)$$

$n$   
↓ 888

$$x^3(cx^n)^{-3/n} \left( \frac{(cx^n)^{3/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( 2e^{2ad}(cx^n)^{3/n} \text{Hypergeometric2F1} \left( 1, \frac{3}{2bdn}, 1 + \frac{3}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) - \frac{1}{3} e^{2ad}(bdn+3) \right)}{bd} \right)$$

$n$

input `Int [x^2*Tanh [d*(a + b*Log [c*x^n])]^2, x]`

output  $(x^3 * (((cx^n)^{(3/n)} * (1 - E^{(2*a*d)} * (cx^n)^{(2*b*d)})) / (b*d * (1 + E^{(2*a*d)} * (cx^n)^{(2*b*d)})) - (-1/3 * (E^{(2*a*d)} * (3 + b*d*n) * (cx^n)^{(3/n)} + 2 * E^{(2*a*d)} * (cx^n)^{(3/n)} * \text{Hypergeometric2F1}[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -(E^{(2*a*d)} * (cx^n)^{(2*b*d)}))]) / (b*d * E^{(2*a*d)}))) / (n * (cx^n)^{(3/n)})$

## Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)/(c*(m+1))}) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)/(b*e*(m+n*(p+1)+1))}), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)/(a*b*e*n*(p+1))}), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_)}]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{((m+1)/n)}) \text{ Subst}[\text{Int}[x^{(m+1)/n - 1}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

**Maple [F]**

$$\int x^2 \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x^2*tanh(d*(a+b*ln(c*x^n)))^2,x)`

output `int(x^2*tanh(d*(a+b*ln(c*x^n)))^2,x)`

**Fricas [F]**

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

output `integral(x^2*tanh(b*d*log(c*x^n) + a*d)^2, x)`

**Sympy [F]**

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh^2(ad + bd \log(cx^n)) dx$$

input `integrate(x**2*tanh(d*(a+b*ln(c*x**n)))**2,x)`

output `Integral(x**2*tanh(a*d + b*d*log(c*x**n))**2, x)`

**Maxima [F]**

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

output `1/3*(b*c^(2*b*d)*d*n*x^3*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 6)*x^3)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 6*integrate(x^2/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

**Giac [F]**

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

output `integrate(x^2*tanh((b*log(c*x^n) + a)*d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x^2*tanh(d*(a + b*log(c*x^n)))^2,x)`

output `int(x^2*tanh(d*(a + b*log(c*x^n)))^2, x)`

**Reduce [F]**

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = -4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}x^2}{x^{4bdn}e^{4ad}c^{4bd} + 2x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) + \frac{x^3}{3}$$

input `int(x^2*tanh(d*(a+b*log(c*x^n)))^2,x)`

output `( - 12*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x**2)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) + x**3)/3`



### 3.191 $\int x \tanh^2(d(a + b \log(cx^n))) dx$

Optimal result	1416
Mathematica [A] (verified)	1417
Rubi [A] (verified)	1417
Maple [F]	1420
Fricas [F]	1420
Sympy [F]	1420
Maxima [F]	1421
Giac [F]	1421
Mupad [F(-1)]	1421
Reduce [F]	1422

#### Optimal result

Integrand size = 17, antiderivative size = 131

$$\int x \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{1}{2} \left(1 + \frac{2}{bdn}\right) x^2 + \frac{x^2 (1 - e^{2ad}(cx^n)^{2bd})}{bdn (1 + e^{2ad}(cx^n)^{2bd})}$$

$$- \frac{2x^2 \operatorname{Hypergeometric2F1}\left(1, \frac{1}{bdn}, 1 + \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

output

```
1/2*(1+2/b/d/n)*x^2+x^2*(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/(1+exp(2*a*d)
*(c*x^n)^(2*b*d))-2*x^2*hypergeom([1, 1/b/d/n],[1+1/b/d/n],-exp(2*a*d)*(c*
x^n)^(2*b*d))/b/d/n
```

**Mathematica [A] (verified)**

Time = 5.01 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.18

$$\int x \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{x^2(2e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{1}{bdn}, 2 + \frac{1}{bdn}, -e^{2d(a+b \log(cx^n))}\right) + (1 + bdn)(bdn - 2 \operatorname{Hypergeometric2F1}\left[1, 1/(b*d*n), 1 + 1/(b*d*n), -E^{(2*d*(a + b*Log[c*x^n])}\right] - 2*\operatorname{Tanh}[d*(a + b*Log[c*x^n])]\right))}{2bdn(1 + bdn)}$$

input

```
Integrate[x*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

output

```
(x^2*(2*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 1/(b*d*n), 2 + 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (1 + b*d*n)*(b*d*n - 2*Hypergeometric2F1[1, 1/(b*d*n), 1 + 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - 2*Tanh[d*(a + b*Log[c*x^n])]))/(2*b*d*n*(1 + b*d*n))
```

**Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.38, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \tanh^2(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{x^2(cx^n)^{-2/n} \int (cx^n)^{\frac{2}{n}-1} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x^2(cx^n)^{-2/n} \int \frac{(cx^n)^{\frac{2}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)^2}{(e^{2ad}(cx^n)^{2bd}+1)^2} d(cx^n)}{n}$$

$$\downarrow 1004$$

$$x^2(cx^n)^{-2/n} \left( \frac{(cx^n)^{2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{2}{n}-1} \left( \frac{e^{2ad}(2-bdn) - e^{4ad}(bdn+2)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{2bd} \right)$$

$n$   
↓ 27

$$x^2(cx^n)^{-2/n} \left( \frac{(cx^n)^{2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{2}{n}-1} \left( \frac{e^{2ad}(2-bdn) - e^{4ad}(bdn+2)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{bd} \right)$$

$n$   
↓ 959

$$x^2(cx^n)^{-2/n} \left( \frac{(cx^n)^{2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( \frac{4e^{2ad} \int \frac{(cx^n)^{\frac{2}{n}-1} d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{n} - \frac{1}{2} e^{2ad}(bdn+2)(cx^n)^{2/n} \right)}{bd} \right)$$

$n$   
↓ 888

$$x^2(cx^n)^{-2/n} \left( \frac{(cx^n)^{2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( 2e^{2ad}(cx^n)^{2/n} \text{Hypergeometric2F1} \left( 1, \frac{1}{bdn}, 1 + \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd} \right) - \frac{1}{2} e^{2ad}(bdn+2)(cx^n)^{2/n} \right)}{bd} \right)$$

$n$

input `Int [x*Tanh [d*(a + b*Log [c*x^n])]^2, x]`

output `(x^2*(((c*x^n)^(2/n)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d)))/(b*d*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) - (-1/2*(E^(2*a*d)*(2 + b*d*n)*(c*x^n)^(2/n)) + 2*E^(2*a*d)*(c*x^n)^(2/n)*Hypergeometric2F1[1, 1/(b*d*n), 1 + 1/(b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/(b*d*E^(2*a*d))))/(n*(c*x^n)^(2/n))`

## Definitions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)/(c*(m+1))}) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)/(b*e*(m+n*(p+1)+1))}), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)/(a*b*e*n*(p+1))}), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_*)}*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{(m+1)/n}) \text{ Subst}[\text{Int}[x^{(m+1)/n - 1}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

**Maple [F]**

$$\int x \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x*tanh(d*(a+b*ln(c*x^n)))^2,x)`

output `int(x*tanh(d*(a+b*ln(c*x^n)))^2,x)`

**Fricas [F]**

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

output `integral(x*tanh(b*d*log(c*x^n) + a*d)^2, x)`

**Sympy [F]**

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh^2(ad + bd \log(cx^n)) dx$$

input `integrate(x*tanh(d*(a+b*ln(c*x**n)))**2,x)`

output `Integral(x*tanh(a*d + b*d*log(c*x**n))**2, x)`

**Maxima [F]**

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

output `1/2*(b*c^(2*b*d)*d*n*x^2*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 4)*x^2)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 4*integrate(x/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

**Giac [F]**

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

output `integrate(x*tanh((b*log(c*x^n) + a)*d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(x*tanh(d*(a + b*log(c*x^n)))^2,x)`

output `int(x*tanh(d*(a + b*log(c*x^n)))^2, x)`

**Reduce [F]**

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = -4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn} x}{x^{4bdn} e^{4ad} c^{4bd} + 2x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) + \frac{x^2}{2}$$

input `int(x*tanh(d*(a+b*log(c*x^n)))^2,x)`

output `( - 8*e**(2*a*d)*c**(2*b*d)*int((x**(2*b*d*n)*x)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) + x**2)/2`

### 3.192 $\int \tanh^2(d(a + b \log(cx^n))) dx$

Optimal result	1423
Mathematica [A] (verified)	1424
Rubi [A] (verified)	1424
Maple [F]	1427
Fricas [F]	1427
Sympy [F]	1427
Maxima [F]	1428
Giac [F]	1428
Mupad [F(-1)]	1428
Reduce [F]	1429

#### Optimal result

Integrand size = 15, antiderivative size = 127

$$\int \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \left(1 + \frac{1}{bdn}\right) x + \frac{x \left(1 - e^{2ad}(cx^n)^{2bd}\right)}{bdn \left(1 + e^{2ad}(cx^n)^{2bd}\right)}$$

$$- \frac{2x \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2bdn}, 1 + \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

output

```
(1+1/b/d/n)*x+x*(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/(1+exp(2*a*d)*(c*x^n)^(2*b*d))-2*x*hypergeom([1, 1/2/b/d/n], [1+1/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n
```



**Mathematica [A] (verified)**

Time = 5.89 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.28

$$\int \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{x(e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{1}{2bdn}, 2 + \frac{1}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) + (1 + 2bdn)(bdn - \operatorname{Hypergeometric2F1}\left[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -E^{(2*d*(a + b*Log[c*x^n])}\right] - \operatorname{Tanh}[d*(a + b*Log[c*x^n])]\right))}{bdn(1 + 2bdn)}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])]^2,x]`

output `(x*(E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 1/(2*b*d*n), 2 + 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (1 + 2*b*d*n)*(b*d*n - Hypergeometric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - Tanh[d*(a + b*Log[c*x^n])]))/(b*d*n*(1 + 2*b*d*n))`

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.39, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6069, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^2(d(a + b \log(cx^n))) dx$$

$$\downarrow \text{6069}$$

$$\frac{x(cx^n)^{-1/n} \int (cx^n)^{\frac{1}{n}-1} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow \text{6071}$$

$$\frac{x(cx^n)^{-1/n} \int \frac{(cx^n)^{\frac{1}{n}-1} (e^{2ad(cx^n)2bd}-1)^2}{(e^{2ad(cx^n)2bd}+1)^2} d(cx^n)}{n}$$

$$\downarrow \text{1004}$$

$$x(cx^n)^{-1/n} \left( \frac{(cx^n)^{\frac{1}{n}} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{1}{n}-1} \left( \frac{e^{2ad}(1-bdn)}{n} - \frac{e^{4ad}(bdn+1)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{2bd} \right)$$

$n$   
↓ 27

$$x(cx^n)^{-1/n} \left( \frac{(cx^n)^{\frac{1}{n}} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{1}{n}-1} \left( \frac{e^{2ad}(1-bdn)}{n} - \frac{e^{4ad}(bdn+1)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{bd} \right)$$

$n$   
↓ 959

$$x(cx^n)^{-1/n} \left( \frac{(cx^n)^{\frac{1}{n}} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( \frac{2e^{2ad} \int \frac{(cx^n)^{\frac{1}{n}-1}}{e^{2ad}(cx^n)^{2bd} + 1} d(cx^n)}{n} - e^{2ad}(bdn+1)(cx^n)^{\frac{1}{n}} \right)}{bd} \right)$$

$n$   
↓ 888

$$x(cx^n)^{-1/n} \left( \frac{(cx^n)^{\frac{1}{n}} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \left( 2e^{2ad}(cx^n)^{\frac{1}{n}} \text{Hypergeometric2F1} \left( 1, \frac{1}{2bdn}, 1 + \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) - e^{2ad}(bdn+1)(cx^n)^{\frac{1}{n}} \right)}{bd} \right)$$

$n$

input `Int [Tanh [d*(a + b*Log [c*x^n])]^2, x]`

output `(x*(((c*x^n)^n^(-1)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d)))/(b*d*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) - (- (E^(2*a*d)*(1 + b*d*n)*(c*x^n)^n^(-1)) + 2*E^(2*a*d)*(c*x^n)^n^(-1)*Hypergeometric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/(b*d*E^(2*a*d))))/(n*(c*x^n)^n^(-1))`

## Definitions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_}))^{(p_*)}((c_*) + (d_*)(x_)^{(n_)}))^{(q_)}, x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(a*b*e*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6069  $\text{Int}[\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_)}]*(b_*)*(d_*)]^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[x/(n*(c*x^n)^{(1/n)}) \text{ Subst}[\text{Int}[x^{(1/n-1)}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)}*x^{(2*b*d)})^p/(1 + E^{(2*a*d)}*x^{(2*b*d)})^p], x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

**Maple [F]**

$$\int \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(tanh(d*(a+b*ln(c*x^n)))^2,x)`

output `int(tanh(d*(a+b*ln(c*x^n)))^2,x)`

**Fricas [F]**

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)^2, x)`

**Sympy [F]**

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh^2(d(a + b \log(cx^n))) dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))**2,x)`

output `Integral(tanh(d*(a + b*log(c*x**n)))**2, x)`

**Maxima [F]**

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

output `(b*c^(2*b*d)*d*n*x*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 2)*x)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 2*integrate(1/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

**Giac [F]**

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n)))^2 dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^2,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^2, x)`

**Reduce [F]**

$$\int \tanh^2(d(a+b \log(cx^n))) dx = -4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{4bdn}e^{4ad}c^{4bd} + 2x^{2bdn}e^{2ad}c^{2bd} + 1} dx \right) + x$$

input `int(tanh(d*(a+b*log(c*x^n)))^2,x)`

output `- 4*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) + x`

### 3.193 $\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x} dx$

Optimal result	1430
Mathematica [A] (verified)	1430
Rubi [A] (verified)	1431
Maple [A] (verified)	1432
Fricas [B] (verification not implemented)	1433
Sympy [B] (verification not implemented)	1433
Maxima [A] (verification not implemented)	1434
Giac [A] (verification not implemented)	1434
Mupad [B] (verification not implemented)	1434
Reduce [B] (verification not implemented)	1435

#### Optimal result

Integrand size = 19, antiderivative size = 28

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \log(x) - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

output `ln(x)-tanh(a*d+b*d*ln(c*x^n))/b/d/n`

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.82

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \frac{\operatorname{arctanh}(\tanh(ad + bd \log(cx^n)))}{bdn} - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x,x]`

output `ArcTanh[Tanh[a*d + b*d*Log[c*x^n]]]/(b*d*n) - Tanh[a*d + b*d*Log[c*x^n]]/(b*d*n)`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {3039, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{\tanh^2(d(a + b \log(cx^n))) d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{-\tan(iad + ib \log(cx^n) d)^2 d \log(cx^n)}{n} \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\tan(iad + ib \log(cx^n) d)^2 d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3954} \\
 & \int \frac{1 d \log(cx^n) - \frac{\tanh(ad + bd \log(cx^n))}{bd}}{n} \\
 & \quad \downarrow \text{24} \\
 & \frac{\log(cx^n) - \frac{\tanh(ad + bd \log(cx^n))}{bd}}{n}
 \end{aligned}$$

input `Int[Tanh[d*(a + b*Log[c*x^n])]^2/x,x]`

output `(Log[c*x^n] - Tanh[a*d + b*d*Log[c*x^n]]/(b*d))/n`



**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst  
[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /;  
NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d  
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]  
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

**Maple [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18

method	result
parallelrisch	$-\frac{\ln(x)dbn+\tanh(d(a+b\ln(cx^n)))}{dbn}$
derivativedivides	$-\tanh(d(a+b\ln(cx^n)))-\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$ $\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$ $-\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$ $\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$
default	$-\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$ $\frac{\ln(\tanh(d(a+b\ln(cx^n)))-1)}{2}+\frac{\ln(\tanh(d(a+b\ln(cx^n)))+1)}{2}$
risch	$\ln(x) + \frac{2}{dbn \left( c^{2bd} (x^n)^{2bd} e^{d \left( ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2 - ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n) \operatorname{csgn}(ic) - ib\pi \operatorname{csgn}(icx^n)^3 + ib\pi \operatorname{csgn}(ic) \right)} \right)}$

input `int(tanh(d*(a+b*ln(c*x^n)))^2/x,x,method=_RETURNVERBOSE)`

output `-(-ln(x)*d*b*n+tanh(d*(a+b*ln(c*x^n))))/d/b/n`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 72 vs.  $2(28) = 56$ .

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.57

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx$$

$$= \frac{(bdn \log(x) + 1) \cosh(bdn \log(x) + bd \log(c) + ad) - \sinh(bdn \log(x) + bd \log(c) + ad)}{bdn \cosh(bdn \log(x) + bd \log(c) + ad)}$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="fricas")`

output `((b*d*n*log(x) + 1)*cosh(b*d*n*log(x) + b*d*log(c) + a*d) - sinh(b*d*n*log(x) + b*d*log(c) + a*d))/(b*d*n*cosh(b*d*n*log(x) + b*d*log(c) + a*d))`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 70 vs.  $2(22) = 44$ .

Time = 2.41 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.50

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = -\frac{\log(\tanh(ad + bd \log(cx^n)) - 1)}{2bdn}$$

$$+ \frac{\log(\tanh(ad + bd \log(cx^n)) + 1)}{2bdn}$$

$$- \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))**2/x,x)`

output `-log(tanh(a*d + b*d*log(c*x**n)) - 1)/(2*b*d*n) + log(tanh(a*d + b*d*log(c*x**n)) + 1)/(2*b*d*n) - tanh(a*d + b*d*log(c*x**n))/(b*d*n)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.29

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \frac{2}{bc^{2bd} d n e^{(2bd \log(x^n) + 2ad)} + bdn} + \log(x)$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="maxima")`output `2/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) + log(x)`**Giac [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.86

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \frac{\log(x^{bdn})}{bdn} + \frac{2}{(c^{2bd} x^{2bdn} e^{(2ad)} + 1) bdn}$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="giac")`output `log(x^(b*d*n))/(b*d*n) + 2/((c^(2*b*d)*x^(2*b*d*n)*e^(2*a*d) + 1)*b*d*n)`**Mupad [B] (verification not implemented)**

Time = 2.43 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \ln(x) + \frac{2}{bdn (e^{2ad} (cx^n)^{2bd} + 1)}$$

input `int(tanh(d*(a + b*log(c*x^n)))^2/x,x)`output `log(x) + 2/(b*d*n*(exp(2*a*d)*(c*x^n)^(2*b*d) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.29

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x} dx = \frac{\log(x^n c) bd - \tanh(\log(x^n c) bd + ad)}{bdn}$$

input `int(tanh(d*(a+b*log(c*x^n)))^2/x,x)`output `(log(x**n*c)*b*d - tanh(log(x**n*c)*b*d + a*d))/(b*d*n)`

### 3.194 $\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^2} dx$

Optimal result	1436
Mathematica [A] (verified)	1437
Rubi [A] (verified)	1437
Maple [F]	1440
Fricas [F]	1440
Sympy [F]	1440
Maxima [F]	1441
Giac [F]	1441
Mupad [F(-1)]	1441
Reduce [F]	1442

#### Optimal result

Integrand size = 19, antiderivative size = 135

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx$$

$$= -\frac{1 - \frac{1}{bdn}}{x} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx(1 + e^{2ad}(cx^n)^{2bd})}$$

$$= \frac{2 \operatorname{Hypergeometric2F1}\left(1, -\frac{1}{2bdn}, 1 - \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{bdnx}$$

output

```
-(1-1/b/d/n)/x+(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/x/(1+exp(2*a*d)*(c*x^n)^(2*b*d))-2*hypergeom([1, -1/2/b/d/n], [1-1/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/x
```

**Mathematica [A] (verified)**

Time = 2.19 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.20

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \frac{e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 - \frac{1}{2bdn}, 2 - \frac{1}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) + (-1 + 2bdn)(bdn + \operatorname{Hypergeometric2F1}\left[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -E^{(2*d*(a + b*Log[c*x^n])}\right])\right]}{bdn(-1 + 2bdn)x}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x^2,x]`output `-((E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(2*b*d*n), 2 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (-1 + 2*b*d*n)*(b*d*n + Hypergeometric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])]) + Tanh[d*(a + b*Log[c*x^n])])/(b*d*n*(-1 + 2*b*d*n)*x)`**Rubi [A] (verified)**Time = 0.50 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.35, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx \\ & \quad \downarrow \text{6073} \\ & \frac{(cx^n)^{\frac{1}{n}} \int (cx^n)^{-1-\frac{1}{n}} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{nx} \\ & \quad \downarrow \text{6071} \\ & \frac{(cx^n)^{\frac{1}{n}} \int \frac{(cx^n)^{-1-\frac{1}{n}} (e^{2ad}(cx^n)^{2bd} - 1)^2}{(e^{2ad}(cx^n)^{2bd} + 1)^2} d(cx^n)}{nx} \\ & \quad \downarrow \text{1004} \end{aligned}$$

$$(cx^n)^{\frac{1}{n}} \left( \frac{(cx^n)^{-1/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{-1-\frac{1}{n}} \left( \frac{e^{2ad}(bdn+1)}{n} - \frac{e^{4ad}(1-bdn)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{2bd} \right)$$

$nx$   
↓ 27

$$(cx^n)^{\frac{1}{n}} \left( \frac{e^{-2ad} \int \frac{(cx^n)^{-1-\frac{1}{n}} \left( \frac{e^{2ad}(bdn+1)}{n} - \frac{e^{4ad}(1-bdn)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{bd} + \frac{(cx^n)^{-1/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)$$

$nx$   
↓ 959

$$(cx^n)^{\frac{1}{n}} \left( \frac{e^{-2ad} \left( \frac{2e^{2ad} \int \frac{(cx^n)^{-1-\frac{1}{n}} d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{n} + e^{2ad}(1-bdn)(cx^n)^{-1/n} \right)}{bd} + \frac{(cx^n)^{-1/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)$$

$nx$   
↓ 888

$$(cx^n)^{\frac{1}{n}} \left( \frac{e^{-2ad} \left( e^{2ad}(1-bdn)(cx^n)^{-1/n} - 2e^{2ad}(cx^n)^{-1/n} \operatorname{Hypergeometric2F1} \left( 1, -\frac{1}{2bdn}, 1 - \frac{1}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{bd} + \frac{(cx^n)^{-1/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)$$

$nx$

input `Int [Tanh [d*(a + b*Log [c*x^n])]^2/x^2, x]`

output `((c*x^n)^n^(-1))*((1 - E^(2*a*d)*(c*x^n)^(2*b*d))/(b*d*(c*x^n)^n^(-1)*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) + ((E^(2*a*d)*(1 - b*d*n))/(c*x^n)^n^(-1) - (2 *E^(2*a*d)*Hypergeometric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -(E^(2*a*d)*d)*(c*x^n)^(2*b*d)]))/(c*x^n)^n^(-1))/(b*d*E^(2*a*d)))/(n*x)`

## Definitions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(a*b*e*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_*)}*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{(m+1)/n}) \text{ Subst}[\text{Int}[x^{(m+1)/n-1}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$



**Maple [F]**

$$\int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^2} dx$$

input `int(tanh(d*(a+b*ln(c*x^n)))^2/x^2,x)`

output `int(tanh(d*(a+b*ln(c*x^n)))^2/x^2,x)`

**Fricas [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)^2/x^2, x)`

**Sympy [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh^2(ad + bd \log(cx^n))}{x^2} dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))**2/x**2,x)`

output `Integral(tanh(a*d + b*d*log(c*x**n))**2/x**2, x)`

**Maxima [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="maxima")`

output `-(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n - 2)/(b*c^(2*b*d)*d*n*x*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x) + 2*integrate(1/(b*c^(2*b*d)*d*n*x^2*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^2), x)`

**Giac [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^2} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)^2/x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^2} dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^2/x^2,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^2/x^2, x)`

**Reduce [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \frac{-4e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{4bdn}e^{4ad}c^{4bdx^2} + 2x^{2bdn}e^{2ad}c^{2bdx^2} + x^2} dx \right) x - 1}{x}$$

input `int(tanh(d*(a+b*log(c*x^n)))^2/x^2,x)`

output `( - 4*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*x**2 + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*x**2 + x**2),x)*x - 1)/x`

### 3.195 $\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^3} dx$

Optimal result	1443
Mathematica [A] (verified)	1444
Rubi [A] (verified)	1444
Maple [F]	1447
Fricas [F]	1447
Sympy [F]	1447
Maxima [F]	1448
Giac [F]	1448
Mupad [F(-1)]	1448
Reduce [F]	1449

#### Optimal result

Integrand size = 19, antiderivative size = 136

$$\int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^3} dx$$

$$= \frac{2 - bdn}{2bdnx^2} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx^2(1 + e^{2ad}(cx^n)^{2bd})}$$

$$- \frac{2 \operatorname{Hypergeometric2F1}\left(1, -\frac{1}{bdn}, 1 - \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{bdnx^2}$$

output

```
1/2*(-b*d*n+2)/b/d/n/x^2+(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/n/x^2/(1+exp(2
*a*d)*(c*x^n)^(2*b*d))-2*hypergeom([1, -1/b/d/n],[1-1/b/d/n],-exp(2*a*d)*(
c*x^n)^(2*b*d))/b/d/n/x^2
```

**Mathematica [A] (verified)**

Time = 2.26 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.17

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \frac{2e^{2d(a+b \log(cx^n))} \operatorname{Hypergeometric2F1}\left(1, 1 - \frac{1}{bdn}, 2 - \frac{1}{bdn}, -e^{2d(a+b \log(cx^n))}\right) + (-1 + bdn)(bdn + 2 \operatorname{Hypergeometric2F1}\left[1, -\frac{1}{(b*d*n)}, 1 - \frac{1}{(b*d*n)}, -E^{(2*d*(a + b*Log[c*x^n])}\right] + 2*\operatorname{Tanh}[d*(a + b*Log[c*x^n])])}{2bdn(-1 + bdn)x^2}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x^3,x]`

output `-1/2*(2*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(b*d*n), 2 - 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (-1 + b*d*n)*(b*d*n + 2*Hypergeometric2F1[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + 2*Tanh[d*(a + b*Log[c*x^n])])/(b*d*n*(-1 + b*d*n)*x^2)`

**Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.35, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx \\ & \quad \downarrow 6073 \\ & \frac{(cx^n)^{2/n} \int (cx^n)^{-1-\frac{2}{n}} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{nx^2} \\ & \quad \downarrow 6071 \\ & \frac{(cx^n)^{2/n} \int \frac{(cx^n)^{-1-\frac{2}{n}} (e^{2ad}(cx^n)^{2bd}-1)^2}{(e^{2ad}(cx^n)^{2bd}+1)^2} d(cx^n)}{nx^2} \\ & \quad \downarrow 1004 \end{aligned}$$

$$\frac{(cx^n)^{2/n} \left( \frac{(cx^n)^{-2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{-1 - \frac{2}{n}} \left( \frac{e^{2ad}(bdn+2)}{n} - \frac{e^{4ad}(2-bdn)(cx^n)^{2bd}}{n} \right) d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{2bd} \right)}{nx^2}$$

$nx^2$   
↓ 27

$$\frac{(cx^n)^{2/n} \left( \frac{e^{-2ad} \int \frac{(cx^n)^{-1 - \frac{2}{n}} \left( \frac{e^{2ad}(bdn+2)}{n} - \frac{e^{4ad}(2-bdn)(cx^n)^{2bd}}{n} \right) d(cx^n)}{bd} + \frac{(cx^n)^{-2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)}{nx^2}$$

$nx^2$   
↓ 959

$$\frac{(cx^n)^{2/n} \left( \frac{e^{-2ad} \left( \frac{4e^{2ad} \int \frac{(cx^n)^{-1 - \frac{2}{n}} d(cx^n)}{e^{2ad}(cx^n)^{2bd} + 1}}{n} + \frac{1}{2} e^{2ad}(2-bdn)(cx^n)^{-2/n} \right)}{bd} + \frac{(cx^n)^{-2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)}{nx^2}$$

$nx^2$   
↓ 888

$$\frac{(cx^n)^{2/n} \left( \frac{e^{-2ad} \left( \frac{1}{2} e^{2ad}(2-bdn)(cx^n)^{-2/n} - 2e^{2ad}(cx^n)^{-2/n} \operatorname{Hypergeometric2F1} \left( 1, -\frac{1}{bdn}, 1 - \frac{1}{bdn}, -e^{2ad}(cx^n)^{2bd} \right) \right)}{bd} + \frac{(cx^n)^{-2/n} (1 - e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd} + 1)} \right)}{nx^2}$$

input `Int [Tanh [d*(a + b*Log [c*x^n])]^2/x^3, x]`

output `((c*x^n)^(2/n)*((1 - E^(2*a*d)*(c*x^n)^(2*b*d))/(b*d*(c*x^n)^(2/n)*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) + ((E^(2*a*d)*(2 - b*d*n))/(2*(c*x^n)^(2/n)) - (2*E^(2*a*d)*Hypergeometric2F1[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/(c*x^n)^(2/n))/(b*d*E^(2*a*d))))/(n*x^2)`

## Defintions of rubi rules used

rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$

rule 888  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 1004  $\text{Int}[((e_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}((c_*) + (d_*)(x_)^{(n_*)})^{(q_*)}, x\_Symbol] \rightarrow \text{Simp}[(-(c*b - a*d))*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}*((c + d*x^n)^{(q-1)}/(a*b*e*n*(p+1))), x] + \text{Simp}[1/(a*b*n*(p+1)) \text{ Int}[(e*x)^m*(a + b*x^n)^{(p+1)}*(c + d*x^n)^{(q-2)}*\text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m+n*(q-1)+1))*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

rule 6071  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_]*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)(x_))^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[(c_*)(x_)^{(n_*)}*(b_*)*(d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{(m+1)/n}) \text{ Subst}[\text{Int}[x^{(m+1)/n-1}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

**Maple [F]**

$$\int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^3} dx$$

input `int(tanh(d*(a+b*ln(c*x^n)))^2/x^3,x)`

output `int(tanh(d*(a+b*ln(c*x^n)))^2/x^3,x)`

**Fricas [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)^2/x^3, x)`

**Sympy [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh^2(ad + bd \log(cx^n))}{x^3} dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))**2/x**3,x)`

output `Integral(tanh(a*d + b*d*log(c*x**n))**2/x**3, x)`



**Maxima [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="maxima")`

output `-1/2*(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n - 4)/(b*c^(2*b*d)*d*n*x^2*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^2) + 4*integrate(1/(b*c^(2*b*d)*d*n*x^3*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^3), x)`

**Giac [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh((b \log(cx^n) + a)d)^2}{x^3} dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)^2/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^3} dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^2/x^3,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^2/x^3, x)`

**Reduce [F]**

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx$$

$$= \frac{-8e^{2ad}c^{2bd} \left( \int \frac{x^{2bdn}}{x^{4bdn}e^{4ad}c^{4bd}x^3 + 2x^{2bdn}e^{2ad}c^{2bd}x^3 + x^3} dx \right) x^2 - 1}{2x^2}$$

input `int(tanh(d*(a+b*log(c*x^n)))^2/x^3,x)`

output `( - 8*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*x**3 + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*x**3 + x**3),x)*x**2 - 1)/(2*x**2)`

### 3.196 $\int \frac{\tanh^3(a+b \log(cx^n))}{x} dx$

Optimal result	1450
Mathematica [A] (verified)	1450
Rubi [C] (verified)	1451
Maple [A] (verified)	1453
Fricas [B] (verification not implemented)	1453
Sympy [A] (verification not implemented)	1454
Maxima [B] (verification not implemented)	1455
Giac [B] (verification not implemented)	1455
Mupad [B] (verification not implemented)	1456
Reduce [B] (verification not implemented)	1457

#### Optimal result

Integrand size = 17, antiderivative size = 43

$$\int \frac{\tanh^3(a+b \log(cx^n))}{x} dx = \frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn}$$

output  $\ln(\cosh(a+b*\ln(c*x^n)))/b/n-1/2*\tanh(a+b*\ln(c*x^n))^2/b/n$

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.88

$$\int \frac{\tanh^3(a+b \log(cx^n))}{x} dx = \frac{2 \log(\cosh(a+b \log(cx^n))) + \operatorname{sech}^2(a+b \log(cx^n))}{2bn}$$

input  $\text{Integrate}[\text{Tanh}[a + b*\text{Log}[c*x^n]]^3/x, x]$

output  $(2*\text{Log}[\text{Cosh}[a + b*\text{Log}[c*x^n]]] + \text{Sech}[a + b*\text{Log}[c*x^n]]^2)/(2*b*n)$

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$ , Rules used = {3039, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\tanh^3(a + b \log(cx^n))}{x} dx \\
 \downarrow \text{3039} \\
 \frac{\int \tanh^3(a + b \log(cx^n)) d \log(cx^n)}{n} \\
 \downarrow \text{3042} \\
 \frac{\int i \tan(ia + ib \log(cx^n))^3 d \log(cx^n)}{n} \\
 \downarrow \text{26} \\
 \frac{i \int \tan(ia + ib \log(cx^n))^3 d \log(cx^n)}{n} \\
 \downarrow \text{3954} \\
 \frac{i \left( \frac{i \tanh^2(a + b \log(cx^n))}{2b} - \int i \tanh(a + b \log(cx^n)) d \log(cx^n) \right)}{n} \\
 \downarrow \text{26} \\
 \frac{i \left( \frac{i \tanh^2(a + b \log(cx^n))}{2b} - i \int \tanh(a + b \log(cx^n)) d \log(cx^n) \right)}{n} \\
 \downarrow \text{3042} \\
 \frac{i \left( \frac{i \tanh^2(a + b \log(cx^n))}{2b} - i \int -i \tan(ia + ib \log(cx^n)) d \log(cx^n) \right)}{n} \\
 \downarrow \text{26} \\
 \frac{i \left( \frac{i \tanh^2(a + b \log(cx^n))}{2b} - \int \tan(ia + ib \log(cx^n)) d \log(cx^n) \right)}{n}
 \end{array}$$

$$\frac{i \left( \frac{i \tanh^2(a+b \log(cx^n))}{2b} - \frac{i \log(\cosh(a+b \log(cx^n)))}{b} \right)}{n}$$

↓ 3956

input `Int[Tanh[a + b*Log[c*x^n]]^3/x,x]`

output `(I*(((-I)*Log[Cosh[a + b*Log[c*x^n]]])/b + ((I/2)*Tanh[a + b*Log[c*x^n]]^2)/b))/n`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst [[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst]] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

**Maple [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09

method	result
parallelrisch	$-\frac{2 \ln(x)bn + \tanh(a+b \ln(cx^n))^2 + 2 \ln(1 - \tanh(a+b \ln(cx^n)))}{2nb}$
derivativedivides	$-\frac{\tanh(a+b \ln(cx^n))^2}{2} - \frac{\ln(\tanh(a+b \ln(cx^n)) - 1)}{2nb} - \frac{\ln(\tanh(a+b \ln(cx^n)) + 1)}{2}$
default	$-\frac{\tanh(a+b \ln(cx^n))^2}{2} - \frac{\ln(\tanh(a+b \ln(cx^n)) - 1)}{2nb} - \frac{\ln(\tanh(a+b \ln(cx^n)) + 1)}{2}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2 \ln(c)}{n} - \frac{2 \ln(x^n)}{n} + \frac{i\pi \operatorname{csgn}(icx^n)^3}{n} - \frac{i\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)}{n} - \frac{i\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2}{n}$

input `int(tanh(a+b*ln(c*x^n))^3/x,x,method=_RETURNVERBOSE)`output `-1/2*(2*ln(x)*b*n+tanh(a+b*ln(c*x^n))^2+2*ln(1-tanh(a+b*ln(c*x^n))))/n/b`**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 566 vs.  $2(41) = 82$ .

Time = 0.10 (sec) , antiderivative size = 566, normalized size of antiderivative = 13.16

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx = \text{Too large to display}$$

input `integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="fricas")`

output

```

-(b*n*cosh(b*n*log(x) + b*log(c) + a)^4*log(x) + 4*b*n*cosh(b*n*log(x) + b
*log(c) + a)*log(x)*sinh(b*n*log(x) + b*log(c) + a)^3 + b*n*log(x)*sinh(b*
n*log(x) + b*log(c) + a)^4 + 2*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c)
+ a)^2 + b*n*log(x) + 2*(3*b*n*cosh(b*n*log(x) + b*log(c) + a)^2*log(x) +
b*n*log(x) - 1)*sinh(b*n*log(x) + b*log(c) + a)^2 - (cosh(b*n*log(x) + b*
log(c) + a)^4 + 4*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(
c) + a)^3 + sinh(b*n*log(x) + b*log(c) + a)^4 + 2*(3*cosh(b*n*log(x) + b*l
og(c) + a)^2 + 1)*sinh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) +
b*log(c) + a)^2 + 4*(cosh(b*n*log(x) + b*log(c) + a)^3 + cosh(b*n*log(x) +
b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a) + 1)*log(2*cosh(b*n*log(x)
+ b*log(c) + a)/(cosh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*lo
g(c) + a))) + 4*(b*n*cosh(b*n*log(x) + b*log(c) + a)^3*log(x) + (b*n*log(x)
- 1)*cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a))/(
b*n*cosh(b*n*log(x) + b*log(c) + a)^4 + 4*b*n*cosh(b*n*log(x) + b*log(c) +
a)*sinh(b*n*log(x) + b*log(c) + a)^3 + b*n*sinh(b*n*log(x) + b*log(c) + a
)^4 + 2*b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + 2*(3*b*n*cosh(b*n*log(x) +
b*log(c) + a)^2 + b*n)*sinh(b*n*log(x) + b*log(c) + a)^2 + b*n + 4*(b*n*c
osh(b*n*log(x) + b*log(c) + a)^3 + b*n*cosh(b*n*log(x) + b*log(c) + a))*si
nh(b*n*log(x) + b*log(c) + a))

```

### Sympy [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.58

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx$$

$$= \begin{cases} \log(x) \tanh^3(a) & \text{for } b = 0 \wedge (b = 0 \vee n = 0) \\ \log(x) \tanh^3(a + b \log(c)) & \text{for } n = 0 \\ \frac{\log(cx^n)}{n} - \frac{\log(\tanh(a + b \log(cx^n)) + 1)}{bn} - \frac{\tanh^2(a + b \log(cx^n))}{2bn} & \text{otherwise} \end{cases}$$

input

```
integrate(tanh(a+b*ln(c*x**n))**3/x,x)
```

output

```

Piecewise((log(x)*tanh(a)**3, Eq(b, 0) & (Eq(b, 0) | Eq(n, 0))), (log(x)*t
anh(a + b*log(c))**3, Eq(n, 0)), (log(c*x**n)/n - log(tanh(a + b*log(c*x**
n)) + 1)/(b*n) - tanh(a + b*log(c*x**n))**2/(2*b*n), True))

```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 304 vs.  $2(41) = 82$ .

Time = 0.10 (sec) , antiderivative size = 304, normalized size of antiderivative = 7.07

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx = \frac{4c^{2b}e^{(2b \log(x^n)+2a)} + 3}{4(bc^{4b}ne^{(4b \log(x^n)+4a)} + 2bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)} - \frac{2c^{2b}e^{(2b \log(x^n)+2a)} + 3}{4(bc^{4b}ne^{(4b \log(x^n)+4a)} + 2bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)} + \frac{3(2c^{2b}e^{(2b \log(x^n)+2a)} + 1)}{4(bc^{4b}ne^{(4b \log(x^n)+4a)} + 2bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)} - \frac{3}{4(bc^{4b}ne^{(4b \log(x^n)+4a)} + 2bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)} + \frac{\log\left(\frac{(c^{2b}e^{(2b \log(x^n)+2a)} + 1)e^{(-2a)}}{c^{2b}}\right)}{bn} - \log(x)$$

input `integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="maxima")`

output `1/4*(4*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 3)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 1/4*(2*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 3)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 3/4*(2*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 3/4/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + log((c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)*e^(-2*a)/c^(2*b))/(b*n) - log(x)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 137 vs.  $2(41) = 82$ .



Time = 0.20 (sec) , antiderivative size = 137, normalized size of antiderivative = 3.19

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx$$

$$= -\frac{\log(x^{bn})}{bn} + \frac{\log\left(\sqrt{2x^{2bn}|c|^{2b} \cos(\pi b \operatorname{sgn}(c) - \pi b) e^{(2a)} + x^{4bn}|c|^{4b} e^{(4a)} + 1}\right)}{bn}$$

$$- \frac{3c^{4b}x^{4bn}e^{(4a)} + 2c^{2b}x^{2bn}e^{(2a)} + 3}{2(c^{2b}x^{2bn}e^{(2a)} + 1)^2bn}$$

input `integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="giac")`

output `-log(x^(b*n))/(b*n) + log(sqrt(2*x^(2*b*n)*abs(c)^(2*b)*cos(pi*b*sgn(c) - pi*b)*e^(2*a) + x^(4*b*n)*abs(c)^(4*b)*e^(4*a) + 1))/(b*n) - 1/2*(3*c^(4*b)*x^(4*b*n)*e^(4*a) + 2*c^(2*b)*x^(2*b*n)*e^(2*a) + 3)/((c^(2*b)*x^(2*b*n)*e^(2*a) + 1)^2*b*n)`

### Mupad [B] (verification not implemented)

Time = 2.24 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.19

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx = \frac{2}{bn + bn e^{2a} (cx^n)^{2b}} - \ln(x)$$

$$- \frac{2}{bn + 2bn e^{2a} (cx^n)^{2b} + bn e^{4a} (cx^n)^{4b}}$$

$$+ \frac{\ln(e^{2a} (cx^n)^{2b} + 1)}{bn}$$

input `int(tanh(a + b*log(c*x^n))^3/x,x)`

output `2/(b*n + b*n*exp(2*a)*(c*x^n)^(2*b)) - log(x) - 2/(b*n + 2*b*n*exp(2*a)*(c*x^n)^(2*b) + b*n*exp(4*a)*(c*x^n)^(4*b)) + log(exp(2*a)*(c*x^n)^(2*b) + 1)/(b*n)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 211, normalized size of antiderivative = 4.91

$$\int \frac{\tanh^3(a + b \log(cx^n))}{x} dx$$

$$= \frac{x^{4bn} e^{4a} c^{4b} \log(x^{2bn} e^{2a} c^{2b} + 1) - x^{4bn} e^{4a} c^{4b} \log(x) bn - x^{4bn} e^{4a} c^{4b} + 2x^{2bn} e^{2a} c^{2b} \log(x^{2bn} e^{2a} c^{2b} + 1) - 2x^{2bn} e^{2a} c^{2b} \log(x) bn - 1}{bn (x^{4bn} e^{4a} c^{4b} + 2x^{2bn} e^{2a} c^{2b} + 1)}$$

input `int(tanh(a+b*log(c*x^n))^3/x,x)`output `(x**(4*b*n)*e**(4*a)*c**(4*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - x**(4*b*n)*e**(4*a)*c**(4*b)*log(x)*b*n - x**(4*b*n)*e**(4*a)*c**(4*b) + 2*x**(2*b*n)*e**(2*a)*c**(2*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - 2*x**(2*b*n)*e**(2*a)*c**(2*b)*log(x)*b*n + log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - log(x)*b*n - 1)/(b*n*(x**(4*b*n)*e**(4*a)*c**(4*b) + 2*x**(2*b*n)*e**(2*a)*c**(2*b) + 1))`

### 3.197 $\int \frac{\tanh^4(a+b \log(cx^n))}{x} dx$

Optimal result	1458
Mathematica [A] (verified)	1458
Rubi [A] (verified)	1459
Maple [A] (verified)	1461
Fricas [B] (verification not implemented)	1461
Sympy [A] (verification not implemented)	1462
Maxima [B] (verification not implemented)	1463
Giac [A] (verification not implemented)	1464
Mupad [B] (verification not implemented)	1464
Reduce [B] (verification not implemented)	1465

#### Optimal result

Integrand size = 17, antiderivative size = 45

$$\int \frac{\tanh^4(a+b \log(cx^n))}{x} dx = \log(x) - \frac{\tanh(a+b \log(cx^n))}{bn} - \frac{\tanh^3(a+b \log(cx^n))}{3bn}$$

output `ln(x)-tanh(a+b*ln(c*x^n))/b/n-1/3*tanh(a+b*ln(c*x^n))^3/b/n`

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.38

$$\int \frac{\tanh^4(a+b \log(cx^n))}{x} dx = \frac{\operatorname{arctanh}(\tanh(a+b \log(cx^n)))}{bn} - \frac{\tanh(a+b \log(cx^n))}{bn} - \frac{\tanh^3(a+b \log(cx^n))}{3bn}$$

input `Integrate[Tanh[a + b*Log[c*x^n]]^4/x,x]`

output `ArcTanh[Tanh[a + b*Log[c*x^n]]]/(b*n) - Tanh[a + b*Log[c*x^n]]/(b*n) - Tanh[a + b*Log[c*x^n]]^3/(3*b*n)`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$ , Rules used = {3039, 3042, 3954, 25, 3042, 25, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\tanh^4(a + b \log(cx^n))}{x} dx \\
 \downarrow 3039 \\
 \int \frac{\tanh^4(a + b \log(cx^n)) d \log(cx^n)}{n} \\
 \downarrow 3042 \\
 \int \frac{\tan(ia + ib \log(cx^n))^4 d \log(cx^n)}{n} \\
 \downarrow 3954 \\
 \frac{-\int -\tanh^2(a + b \log(cx^n)) d \log(cx^n) - \frac{\tanh^3(a+b \log(cx^n))}{3b}}{n} \\
 \downarrow 25 \\
 \frac{\int \tanh^2(a + b \log(cx^n)) d \log(cx^n) - \frac{\tanh^3(a+b \log(cx^n))}{3b}}{n} \\
 \downarrow 3042 \\
 \frac{-\frac{\tanh^3(a+b \log(cx^n))}{3b} + \int -\tan(ia + ib \log(cx^n))^2 d \log(cx^n)}{n} \\
 \downarrow 25 \\
 \frac{-\frac{\tanh^3(a+b \log(cx^n))}{3b} - \int \tan(ia + ib \log(cx^n))^2 d \log(cx^n)}{n} \\
 \downarrow 3954 \\
 \frac{\int 1 d \log(cx^n) - \frac{\tanh^3(a+b \log(cx^n))}{3b} - \frac{\tanh(a+b \log(cx^n))}{b}}{n} \\
 \downarrow 24
 \end{array}$$

$$\frac{-\frac{\tanh^3(a+b\log(cx^n))}{3b} - \frac{\tanh(a+b\log(cx^n))}{b} + \log(cx^n)}{n}$$

input `Int[Tanh[a + b*Log[c*x^n]]^4/x,x]`

output `(Log[c*x^n] - Tanh[a + b*Log[c*x^n]]/b - Tanh[a + b*Log[c*x^n]]^3/(3*b))/n`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst  
[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /;  
NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d  
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]  
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

### Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

method	result
parallelsch	$-\frac{-3 \ln(x)bn + \tanh(a+b \ln(cx^n))^3 + 3 \tanh(a+b \ln(cx^n))}{3nb}$
derivativdivides	$\frac{-\frac{\tanh(a+b \ln(cx^n))^3}{3} - \tanh(a+b \ln(cx^n)) - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2} + \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2}}{nb}$
default	$\frac{-\frac{\tanh(a+b \ln(cx^n))^3}{3} - \tanh(a+b \ln(cx^n)) - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2} + \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2}}{nb}$
risch	$\ln(x) + \frac{4(x^n)^{4b} c^{4b} e^{4a} e^{-2ib\pi \operatorname{csgn}(icx^n)^3} e^{2ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)} e^{2ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2} e^{-2ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2} e^{-ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2}}{bn \left( (x^n)^{2b} c^{2b} e^{2a} e^{ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2} e^{-ib\pi \operatorname{csgn}(ix^n) \operatorname{csgn}(icx^n)^2} \right)}$

input `int(tanh(a+b*ln(c*x^n))^4/x,x,method=_RETURNVERBOSE)`

output `-1/3*(-3*ln(x)*b*n+tanh(a+b*ln(c*x^n))^3+3*tanh(a+b*ln(c*x^n)))/n/b`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(43) = 86.

Time = 0.09 (sec) , antiderivative size = 194, normalized size of antiderivative = 4.31

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx$$

$$= \frac{(3bn \log(x) + 4) \cosh(bn \log(x) + b \log(c) + a)^3 + 3(3bn \log(x) + 4) \cosh(bn \log(x) + b \log(c) + a) \sinh(bn \log(x) + b \log(c) + a)^2}{3(bn \cosh(bn \log(x) + b \log(c) + a) + \sinh(bn \log(x) + b \log(c) + a)^2)}$$

input `integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="fricas")`

output

```
1/3*((3*b*n*log(x) + 4)*cosh(b*n*log(x) + b*log(c) + a)^3 + 3*(3*b*n*log(x)
) + 4)*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a)^2 -
12*cosh(b*n*log(x) + b*log(c) + a)^2*sinh(b*n*log(x) + b*log(c) + a) - 4*
sinh(b*n*log(x) + b*log(c) + a)^3 + 3*(3*b*n*log(x) + 4)*cosh(b*n*log(x) +
b*log(c) + a))/(b*n*cosh(b*n*log(x) + b*log(c) + a)^3 + 3*b*n*cosh(b*n*lo
g(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a)^2 + 3*b*n*cosh(b*n*lo
g(x) + b*log(c) + a))
```

**Sympy [A] (verification not implemented)**

Time = 1.67 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.44

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx$$

$$= \begin{cases} \log(x) \tanh^4(a) & \text{for } b = 0 \wedge (b = 0 \vee n = 0) \\ \log(x) \tanh^4(a + b \log(c)) & \text{for } n = 0 \\ \frac{\log(cx^n)}{n} - \frac{\tanh^3(a + b \log(cx^n))}{3bn} - \frac{\tanh(a + b \log(cx^n))}{bn} & \text{otherwise} \end{cases}$$

input

```
integrate(tanh(a+b*ln(c*x**n))**4/x,x)
```

output

```
Piecewise((log(x)*tanh(a)**4, Eq(b, 0) & (Eq(b, 0) | Eq(n, 0))), (log(x)*t
anh(a + b*log(c))**4, Eq(n, 0)), (log(c*x**n)/n - tanh(a + b*log(c*x**n))*
*3/(3*b*n) - tanh(a + b*log(c*x**n))/(b*n), True))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 494 vs.  $2(43) = 86$ .

Time = 0.13 (sec) , antiderivative size = 494, normalized size of antiderivative = 10.98

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx$$

$$= \frac{18c^{4b}e^{(4b \log(x^n)+4a)} + 27c^{2b}e^{(2b \log(x^n)+2a)} + 11}{12(bc^{6b}ne^{(6b \log(x^n)+6a)} + 3bc^4bne^{(4b \log(x^n)+4a)} + 3bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)}$$

$$+ \frac{6c^{4b}e^{(4b \log(x^n)+4a)} + 15c^{2b}e^{(2b \log(x^n)+2a)} + 11}{12(bc^{6b}ne^{(6b \log(x^n)+6a)} + 3bc^4bne^{(4b \log(x^n)+4a)} + 3bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)}$$

$$+ \frac{2(3c^{4b}e^{(4b \log(x^n)+4a)} + 3c^{2b}e^{(2b \log(x^n)+2a)} + 1)}{3(bc^{6b}ne^{(6b \log(x^n)+6a)} + 3bc^4bne^{(4b \log(x^n)+4a)} + 3bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)}$$

$$- \frac{3c^{2b}e^{(2b \log(x^n)+2a)} + 1}{2(bc^{6b}ne^{(6b \log(x^n)+6a)} + 3bc^4bne^{(4b \log(x^n)+4a)} + 3bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)}$$

$$+ \frac{3(bc^{6b}ne^{(6b \log(x^n)+6a)} + 3bc^4bne^{(4b \log(x^n)+4a)} + 3bc^{2b}ne^{(2b \log(x^n)+2a)} + bn)}{2}$$

$$+ \log(x)$$

input `integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="maxima")`

output

```
1/12*(18*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 27*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 11)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 1/12*(6*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 15*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 11)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 2/3*(3*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 3*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 1/2*(3*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 2/3/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + log(x)
```



**Giac [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.73

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx = \frac{\log(x^{bn})}{bn} + \frac{4(3c^{4b}x^{4bn}e^{(4a)} + 3c^{2b}x^{2bn}e^{(2a)} + 2)}{3(c^{2b}x^{2bn}e^{(2a)} + 1)^3bn}$$

input `integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="giac")`output `log(x^(b*n))/(b*n) + 4/3*(3*c^(4*b)*x^(4*b*n)*e^(4*a) + 3*c^(2*b)*x^(2*b*n)*e^(2*a) + 2)/((c^(2*b)*x^(2*b*n)*e^(2*a) + 1)^3*b*n)`**Mupad [B] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 162, normalized size of antiderivative = 3.60

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx = \ln(x) + \frac{\frac{4}{3bn} + \frac{4e^{4a}(cx^n)^{4b}}{3bn}}{3e^{2a}(cx^n)^{2b} + 3e^{4a}(cx^n)^{4b} + e^{6a}(cx^n)^{6b} + 1} + \frac{4}{3bn(e^{2a}(cx^n)^{2b} + 1)} + \frac{4e^{2a}(cx^n)^{2b}}{3bn(2e^{2a}(cx^n)^{2b} + e^{4a}(cx^n)^{4b} + 1)}$$

input `int(tanh(a + b*log(c*x^n))^4/x,x)`output `log(x) + (4/(3*b*n) + (4*exp(4*a)*(c*x^n)^(4*b))/(3*b*n))/(3*exp(2*a)*(c*x^n)^(2*b) + 3*exp(4*a)*(c*x^n)^(4*b) + exp(6*a)*(c*x^n)^(6*b) + 1) + 4/(3*b*n*(exp(2*a)*(c*x^n)^(2*b) + 1)) + (4*exp(2*a)*(c*x^n)^(2*b))/(3*b*n*(2*exp(2*a)*(c*x^n)^(2*b) + exp(4*a)*(c*x^n)^(4*b) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int \frac{\tanh^4(a + b \log(cx^n))}{x} dx$$
$$= \frac{3 \log(x^n c) b - \tanh(\log(x^n c) b + a)^3 - 3 \tanh(\log(x^n c) b + a)}{3bn}$$

input `int(tanh(a+b*log(c*x^n))^4/x,x)`output `(3*log(x**n*c)*b - tanh(log(x**n*c)*b + a)**3 - 3*tanh(log(x**n*c)*b + a)) / (3*b*n)`

### 3.198 $\int \frac{\tanh^5(a+b \log(cx^n))}{x} dx$

Optimal result . . . . .	1466
Mathematica [A] (verified) . . . . .	1466
Rubi [C] (verified) . . . . .	1467
Maple [A] (verified) . . . . .	1469
Fricas [B] (verification not implemented) . . . . .	1470
Sympy [A] (verification not implemented) . . . . .	1471
Maxima [B] (verification not implemented) . . . . .	1471
Giac [B] (verification not implemented) . . . . .	1472
Mupad [B] (verification not implemented) . . . . .	1473
Reduce [B] (verification not implemented) . . . . .	1474

#### Optimal result

Integrand size = 17, antiderivative size = 66

$$\int \frac{\tanh^5(a+b \log(cx^n))}{x} dx = \frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} - \frac{\tanh^4(a+b \log(cx^n))}{4bn}$$

output

```
ln(cosh(a+b*ln(c*x^n)))/b/n-1/2*tanh(a+b*ln(c*x^n))^2/b/n-1/4*tanh(a+b*ln(c*x^n))^4/b/n
```

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.83

$$\int \frac{\tanh^5(a+b \log(cx^n))}{x} dx = \frac{4 \log(\cosh(a+b \log(cx^n))) + 4\operatorname{sech}^2(a+b \log(cx^n)) - \operatorname{sech}^4(a+b \log(cx^n))}{4bn}$$

input

```
Integrate[Tanh[a + b*Log[c*x^n]]^5/x,x]
```

output

$$(4*\text{Log}[\text{Cosh}[a + b*\text{Log}[c*x^n]]] + 4*\text{Sech}[a + b*\text{Log}[c*x^n]]^2 - \text{Sech}[a + b*\text{Log}[c*x^n]]^4)/(4*b*n)$$
**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.706$ , Rules used = {3039, 3042, 26, 3954, 26, 3042, 26, 3954, 26, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^5(a + b \log(cx^n))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \frac{\int \tanh^5(a + b \log(cx^n)) d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -i \tan(ia + ib \log(cx^n))^5 d \log(cx^n)}{n} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int \tan(ia + ib \log(cx^n))^5 d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3954} \\
 & \frac{i \left( - \int -i \tanh^3(a + b \log(cx^n)) d \log(cx^n) - \frac{i \tanh^4(a + b \log(cx^n))}{4b} \right)}{n} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \left( i \int \tanh^3(a + b \log(cx^n)) d \log(cx^n) - \frac{i \tanh^4(a + b \log(cx^n))}{4b} \right)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{i \left( i \int i \tan(ia + ib \log(cx^n))^3 d \log(cx^n) - \frac{i \tanh^4(a + b \log(cx^n))}{4b} \right)}{n}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 26 \\
 & \frac{i \left( - \int \tan (ia + ib \log (cx^n))^3 d \log (cx^n) - \frac{i \tanh^4(a+b \log(cx^n))}{4b} \right)}{n} \\
 & \downarrow 3954 \\
 & \frac{i \left( \int i \tanh (a + b \log (cx^n)) d \log (cx^n) - \frac{i \tanh^4(a+b \log(cx^n))}{4b} - \frac{i \tanh^2(a+b \log(cx^n))}{2b} \right)}{n} \\
 & \downarrow 26 \\
 & \frac{i \left( i \int \tanh (a + b \log (cx^n)) d \log (cx^n) - \frac{i \tanh^4(a+b \log(cx^n))}{4b} - \frac{i \tanh^2(a+b \log(cx^n))}{2b} \right)}{n} \\
 & \downarrow 3042 \\
 & \frac{i \left( i \int -i \tan (ia + ib \log (cx^n)) d \log (cx^n) - \frac{i \tanh^4(a+b \log(cx^n))}{4b} - \frac{i \tanh^2(a+b \log(cx^n))}{2b} \right)}{n} \\
 & \downarrow 26 \\
 & \frac{i \left( \int \tan (ia + ib \log (cx^n)) d \log (cx^n) - \frac{i \tanh^4(a+b \log(cx^n))}{4b} - \frac{i \tanh^2(a+b \log(cx^n))}{2b} \right)}{n} \\
 & \downarrow 3956 \\
 & \frac{i \left( - \frac{i \tanh^4(a+b \log(cx^n))}{4b} - \frac{i \tanh^2(a+b \log(cx^n))}{2b} + \frac{i \log(\cosh(a+b \log(cx^n)))}{b} \right)}{n}
 \end{aligned}$$

input `Int [Tanh[a + b*Log[c*x^n]]^5/x,x]`

output `((-1)*((I*Log[Cosh[a + b*Log[c*x^n]]])/b - ((I/2)*Tanh[a + b*Log[c*x^n]]^2)/b - ((I/4)*Tanh[a + b*Log[c*x^n]]^4)/b))/n`

**Defintions of rubi rules used**

rule 26 `Int[(Complex[0, a_])*(Fx), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst [[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d *x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d *x], x]]/d, x] /; FreeQ[{c, d}, x]`

**Maple [A] (verified)**

Time = 1.54 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.94

method	result
parallelrisch	$\frac{-\tanh(a+b \ln(c x^n))^4+4 \ln(x) b n+2 \tanh(a+b \ln(c x^n))^2+4 \ln(1-\tanh(a+b \ln(c x^n)))}{4 n b}$
derivativedivides	$\frac{-\frac{\tanh(a+b \ln(c x^n))^4}{4}-\frac{\tanh(a+b \ln(c x^n))^2}{2}-\frac{\ln(\tanh(a+b \ln(c x^n))-1)}{2}-\frac{\ln(\tanh(a+b \ln(c x^n))+1)}{2}}{n b}$
default	$\frac{-\frac{\tanh(a+b \ln(c x^n))^4}{4}-\frac{\tanh(a+b \ln(c x^n))^2}{2}-\frac{\ln(\tanh(a+b \ln(c x^n))-1)}{2}-\frac{\ln(\tanh(a+b \ln(c x^n))+1)}{2}}{n b}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2 \ln(c)}{n} - \frac{2 \ln(x^n)}{n} + \frac{i \pi \operatorname{csgn}(i x^n) \operatorname{csgn}(i c x^n) \operatorname{csgn}(i c)}{n} - \frac{i \pi \operatorname{csgn}(i c x^n)^2 \operatorname{csgn}(i c)}{n} - \frac{i \pi \operatorname{csgn}(i c)}{n}$

input `int(tanh(a+b*ln(c*x^n))^5/x,x,method=_RETURNVERBOSE)`

output

```
-1/4*(tanh(a+b*ln(c*x^n))^4+4*ln(x)*b*n+2*tanh(a+b*ln(c*x^n))^2+4*ln(1-tanh(a+b*ln(c*x^n))))/n/b
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1568 vs.  $2(62) = 124$ .

Time = 0.11 (sec) , antiderivative size = 1568, normalized size of antiderivative = 23.76

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx = \text{Too large to display}$$

input

```
integrate(tanh(a+b*log(c*x^n))^5/x,x, algorithm="fricas")
```

output

```
-(b*n*cosh(b*n*log(x) + b*log(c) + a)^8*log(x) + 8*b*n*cosh(b*n*log(x) + b*log(c) + a)*log(x)*sinh(b*n*log(x) + b*log(c) + a)^7 + b*n*log(x)*sinh(b*n*log(x) + b*log(c) + a)^8 + 4*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^6 + 4*(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^2*log(x) + b*n*log(x) - 1)*sinh(b*n*log(x) + b*log(c) + a)^6 + 8*(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^3*log(x) + 3*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a)^5 + 2*(3*b*n*log(x) - 2)*cosh(b*n*log(x) + b*log(c) + a)^4 + 2*(35*b*n*cosh(b*n*log(x) + b*log(c) + a)^4*log(x) + 30*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^2 + 3*b*n*log(x) - 2)*sinh(b*n*log(x) + b*log(c) + a)^4 + 8*(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^5*log(x) + 10*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^3 + (3*b*n*log(x) - 2)*cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a)^3 + 4*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^2 + b*n*log(x) + 4*(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^6*log(x) + 15*(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^4 + 3*(3*b*n*log(x) - 2)*cosh(b*n*log(x) + b*log(c) + a)^2 + b*n*log(x) - 1)*sinh(b*n*log(x) + b*log(c) + a)^2 - (cosh(b*n*log(x) + b*log(c) + a)^8 + 8*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a)^7 + sinh(b*n*log(x) + b*log(c) + a)^8 + 4*(7*cosh(b*n*log(x) + b*log(c) + a)^2 + 1)*sinh(b*n*log(x) + b*log(c) + a)^6 + 4*cosh(b*n*log(x) + b*log(c) + a)^6 + 8*(7*cosh(b*n*log(x) + b*log(c) + a)^3 ...
```

**Sympy [A] (verification not implemented)**

Time = 3.89 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.32

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx$$

$$= \begin{cases} \log(x) \tanh^5(a) & \text{for } b = 0 \wedge (b = 0 \vee n = 0) \\ \log(x) \tanh^5(a + b \log(c)) & \text{for } n = 0 \\ \frac{\log(cx^n)}{n} - \frac{\log(\tanh(a + b \log(cx^n)) + 1)}{bn} - \frac{\tanh^4(a + b \log(cx^n))}{4bn} - \frac{\tanh^2(a + b \log(cx^n))}{2bn} & \text{otherwise} \end{cases}$$

input `integrate(tanh(a+b*ln(c*x**n))**5/x,x)`

output `Piecewise((log(x)*tanh(a)**5, Eq(b, 0) & (Eq(b, 0) | Eq(n, 0))), (log(x)*tanh(a + b*log(c))**5, Eq(n, 0)), (log(c*x**n)/n - log(tanh(a + b*log(c*x**n)) + 1)/(b*n) - tanh(a + b*log(c*x**n))**4/(4*b*n) - tanh(a + b*log(c*x**n))**2/(2*b*n), True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 829 vs. 2(62) = 124.

Time = 0.17 (sec) , antiderivative size = 829, normalized size of antiderivative = 12.56

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx = \text{Too large to display}$$

input `integrate(tanh(a+b*log(c*x^n))^5/x,x, algorithm="maxima")`



output

```

1/24*(48*c^(6*b)*e^(6*b*log(x^n) + 6*a) + 108*c^(4*b)*e^(4*b*log(x^n) + 4*
a) + 88*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 25)/(b*c^(8*b)*n*e^(8*b*log(x^n)
+ 8*a) + 4*b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x
^n) + 4*a) + 4*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 1/24*(12*c^(6*b
)*e^(6*b*log(x^n) + 6*a) + 42*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 52*c^(2*b)*
e^(2*b*log(x^n) + 2*a) + 25)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(
6*b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b
*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 5/8*(4*c^(6*b)*e^(6*b*log(x^n)
+ 6*a) + 6*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 4*c^(2*b)*e^(2*b*log(x^n) + 2*
a) + 1)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(6*b)*n*e^(6*b*log(x^n)
) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b*c^(2*b)*n*e^(2*b*log
(x^n) + 2*a) + b*n) - 5/12*(6*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 4*c^(2*b)*e
^(2*b*log(x^n) + 2*a) + 1)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(6*
b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b*c
^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 5/12*(4*c^(2*b)*e^(2*b*log(x^n) +
2*a) + 1)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(6*b)*n*e^(6*b*log(
x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b*c^(2*b)*n*e^(2*b*
log(x^n) + 2*a) + b*n) - 5/8/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(
6*b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b
*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + log((c^(2*b)*e^(2*b*log(x^n)...

```

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 171 vs.  $2(62) = 124$ .

Time = 0.22 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.59

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx$$

$$= -\frac{\log(x^{bn})}{bn} + \frac{\log\left(\sqrt{2x^{2bn}|c|^{2b}} \cos(\pi b \operatorname{sgn}(c) - \pi b) e^{(2a)} + x^{4bn}|c|^{4b} e^{(4a)} + 1\right)}{12(c^{2b}x^{2bn}e^{(2a)} + 1)^4bn}$$

input

```
integrate(tanh(a+b*log(c*x^n))^5/x,x, algorithm="giac")
```

output

```
-log(x^(b*n))/(b*n) + log(sqrt(2*x^(2*b*n)*abs(c)^(2*b)*cos(pi*b*sgn(c) -
pi*b)*e^(2*a) + x^(4*b*n)*abs(c)^(4*b)*e^(4*a) + 1))/(b*n) - 1/12*(25*c^(8
*b)*x^(8*b*n)*e^(8*a) + 52*c^(6*b)*x^(6*b*n)*e^(6*a) + 102*c^(4*b)*x^(4*b*
n)*e^(4*a) + 52*c^(2*b)*x^(2*b*n)*e^(2*a) + 25)/((c^(2*b)*x^(2*b*n)*e^(2*a
) + 1)^4*b*n)
```

### Mupad [B] (verification not implemented)

Time = 2.22 (sec) , antiderivative size = 227, normalized size of antiderivative = 3.44

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx$$

$$= \frac{8}{bn + 3bne^{2a}(cx^n)^{2b} + 3bne^{4a}(cx^n)^{4b} + bne^{6a}(cx^n)^{6b}} - \ln(x) + \frac{4}{bn + bne^{2a}(cx^n)^{2b}} - \frac{4}{bn + 4bne^{2a}(cx^n)^{2b} + 6bne^{4a}(cx^n)^{4b} + 4bne^{6a}(cx^n)^{6b} + bne^{8a}(cx^n)^{8b}} - \frac{8}{bn + 2bne^{2a}(cx^n)^{2b} + bne^{4a}(cx^n)^{4b}} + \frac{\ln(e^{2a}(cx^n)^{2b} + 1)}{bn}$$

input

```
int(tanh(a + b*log(c*x^n))^5/x,x)
```

output

```
8/(b*n + 3*b*n*exp(2*a)*(c*x^n)^(2*b) + 3*b*n*exp(4*a)*(c*x^n)^(4*b) + b*n
*exp(6*a)*(c*x^n)^(6*b)) - log(x) + 4/(b*n + b*n*exp(2*a)*(c*x^n)^(2*b)) -
4/(b*n + 4*b*n*exp(2*a)*(c*x^n)^(2*b) + 6*b*n*exp(4*a)*(c*x^n)^(4*b) + 4*
b*n*exp(6*a)*(c*x^n)^(6*b) + b*n*exp(8*a)*(c*x^n)^(8*b)) - 8/(b*n + 2*b*n*
exp(2*a)*(c*x^n)^(2*b) + b*n*exp(4*a)*(c*x^n)^(4*b)) + log(exp(2*a)*(c*x^n
)^(2*b) + 1)/(b*n)
```

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 385, normalized size of antiderivative = 5.83

$$\int \frac{\tanh^5(a + b \log(cx^n))}{x} dx$$

$$= \frac{x^{8bn} e^{8a} c^{8b} \log(x^{2bn} e^{2a} c^{2b} + 1) - x^{8bn} e^{8a} c^{8b} \log(x) bn - x^{8bn} e^{8a} c^{8b} + 4x^{6bn} e^{6a} c^{6b} \log(x^{2bn} e^{2a} c^{2b} + 1) - 4x^{6bn} e^{6a} c^{6b} \log(x) b + 4x^{4bn} e^{4a} c^{4b} \log(x^{2bn} e^{2a} c^{2b} + 1) - 4x^{4bn} e^{4a} c^{4b} \log(x) b + 4x^{2bn} e^{2a} c^{2b} \log(x^{2bn} e^{2a} c^{2b} + 1) - 4x^{2bn} e^{2a} c^{2b} \log(x) b + 4}{(b^n (x^{8bn} e^{8a} c^{8b} + 4x^{6bn} e^{6a} c^{6b} + 4x^{4bn} e^{4a} c^{4b} + 4x^{2bn} e^{2a} c^{2b} + 1))}$$

input `int(tanh(a+b*log(c*x^n))^5/x,x)`

output

```
(x**(8*b*n)*e**(8*a)*c**(8*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - x**(8*b*n)*e**(8*a)*c**(8*b)*log(x)*b*n - x**(8*b*n)*e**(8*a)*c**(8*b) + 4*x**(6*b*n)*e**(6*a)*c**(6*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - 4*x**(6*b*n)*e**(6*a)*c**(6*b)*log(x)*b*n + 6*x**(4*b*n)*e**(4*a)*c**(4*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - 6*x**(4*b*n)*e**(4*a)*c**(4*b)*log(x)*b*n - 2*x**(4*b*n)*e**(4*a)*c**(4*b) + 4*x**(2*b*n)*e**(2*a)*c**(2*b)*log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - 4*x**(2*b*n)*e**(2*a)*c**(2*b)*log(x)*b*n + log(x**(2*b*n)*e**(2*a)*c**(2*b) + 1) - log(x)*b*n - 1)/(b*n*(x**(8*b*n)*e**(8*a)*c**(8*b) + 4*x**(6*b*n)*e**(6*a)*c**(6*b) + 6*x**(4*b*n)*e**(4*a)*c**(4*b) + 4*x**(2*b*n)*e**(2*a)*c**(2*b) + 1))
```

### 3.199 $\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$

Optimal result	1475
Mathematica [A] (verified)	1475
Rubi [A] (verified)	1476
Maple [F]	1477
Fricas [F]	1478
Sympy [F]	1478
Maxima [F]	1478
Giac [F]	1479
Mupad [F(-1)]	1479
Reduce [F]	1479

#### Optimal result

Integrand size = 19, antiderivative size = 88

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$$

$$= \frac{(ex)^{1+m}}{e(1+m)} - \frac{2(ex)^{1+m} \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{e(1+m)}$$

output

```
(e*x)^(1+m)/e/(1+m)-2*(e*x)^(1+m)*hypergeom([1, 1/2*(1+m)/b/d/n], [1+1/2*(1+m)/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/e/(1+m)
```

#### Mathematica [A] (verified)

Time = 8.85 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.82

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$$

$$= \frac{x(ex)^m \left( -\operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn}, -e^{2d(a+b \log(cx^n))}\right) + \frac{e^{2ad(1+m)}(cx^n)^{2bd} \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn}, -e^{2ad(1+m)}(cx^n)^{2bd}\right)}{1+m+2bd} \right)}{1+m}$$

input

```
Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])],x]
```

output

$$\frac{(x*(e*x)^m*(-\text{Hypergeometric2F1}[1, (1+m)/(2*b*d*n), 1+(1+m)/(2*b*d*n), -E^{2*d*(a+b*\text{Log}[c*x^n])}] + (E^{2*a*d}*(1+m)*(c*x^n)^{2*b*d}*\text{Hypergeometric2F1}[1, (1+m+2*b*d*n)/(2*b*d*n), (1+m+4*b*d*n)/(2*b*d*n), -E^{2*a*d}*(c*x^n)^{2*b*d}]))/(1+m+2*b*d*n)))/(1+m)}$$
**Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.41, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6073, 6071, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int (cx^n)^{\frac{m+1}{n}-1} \tanh(d(a + b \log(cx^n))) d(cx^n)}{en}$$

$$\downarrow 6071$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int \frac{(cx^n)^{\frac{m+1}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)}{e^{2ad}(cx^n)^{2bd}+1} d(cx^n)}{en}$$

$$\downarrow 959$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{n(cx^n)^{\frac{m+1}{n}}}{m+1} - 2 \int \frac{(cx^n)^{\frac{m+1}{n}-1}}{e^{2ad}(cx^n)^{2bd}+1} d(cx^n) \right)}{en}$$

$$\downarrow 888$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{n(cx^n)^{\frac{m+1}{n}}}{m+1} - \frac{2n(cx^n)^{\frac{m+1}{n}} \text{Hypergeometric2F1}\left(1, \frac{m+1}{2bdn}, \frac{m+1}{2bdn}+1, -e^{2ad}(cx^n)^{2bd}\right)}{m+1} \right)}{en}$$

input

$$\text{Int}[(e*x)^m*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])], x]$$

output  $((e*x)^{(1+m)}*((n*(c*x^n)^{((1+m)/n)})/(1+m) - (2*n*(c*x^n)^{((1+m)/n)} * \text{Hypergeometric2F1}[1, (1+m)/(2*b*d*n), 1 + (1+m)/(2*b*d*n), -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})]/(1+m)))/(e*n*(c*x^n)^{((1+m)/n)})$

### Defintions of rubi rules used

rule 888  $\text{Int}[((c_*)*(x_))^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_))^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /;$   $\text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 959  $\text{Int}[((e_*)*(x_))^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_))^{(p_*)}*((c_*) + (d_*)*(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)) \ \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

rule 6071  $\text{Int}[((e_*)*(x_))^{(m_*)}*\text{Tanh}[((a_*) + \text{Log}[x_]*(b_))* (d_)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)}*x^{(2*b*d)})^p/(1 + E^{(2*a*d)}*x^{(2*b*d)})^p), x] /;$   $\text{FreeQ}\{a, b, d, e, m, p\}, x]$

rule 6073  $\text{Int}[((e_*)*(x_))^{(m_*)}*\text{Tanh}[((a_*) + \text{Log}[(c_*)*(x_)^{(n_)}]*(b_))* (d_)]^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{((m+1)/n)}) \ \text{Subst}[\text{Int}[x^{((m+1)/n - 1)}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /;$   $\text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ (\text{NeQ}[c, 1] \ || \ \text{NeQ}[n, 1])$

### Maple [F]

$$\int (ex)^m \tanh(d(a + b \ln(cx^n))) dx$$

input  $\text{int}((e*x)^m*\tanh(d*(a+b*\ln(c*x^n))),x)$

output `int((e*x)^m*tanh(d*(a+b*ln(c*x^n))),x)`

### Fricas [F]

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d) dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

output `integral((e*x)^m*tanh(b*d*log(c*x^n) + a*d), x)`

### Sympy [F]

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh(ad + bd \log(cx^n)) dx$$

input `integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))),x)`

output `Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n)), x)`

### Maxima [F]

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d) dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")`

output `e^m*x^m/(m + 1) - 2*e^m*integrate(x^m/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)`

**Giac [F]**

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d) dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

output `integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n))) (ex)^m dx$$

input `int(tanh(d*(a + b*log(c*x^n)))*(e*x)^m,x)`

output `int(tanh(d*(a + b*log(c*x^n)))*(e*x)^m, x)`

**Reduce [F]**

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$$

$$= \frac{e^m \left( x^m x - 2 \left( \int \frac{x^m}{x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) m - 2 \left( \int \frac{x^m}{x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) \right)}{m + 1}$$

input `int((e*x)^m*tanh(d*(a+b*log(c*x^n))),x)`

output `(e**m*(x**m*x - 2*int(x**m/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x)*m - 2*int(x**m/(x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x)))/(m + 1)`



### 3.200 $\int (ex)^m \tanh^2 (d(a + b \log (cx^n))) dx$

Optimal result	1480
Mathematica [A] (verified)	1481
Rubi [A] (verified)	1481
Maple [F]	1484
Fricas [F]	1484
Sympy [F]	1484
Maxima [F]	1485
Giac [F]	1485
Mupad [F(-1)]	1485
Reduce [F]	1486

#### Optimal result

Integrand size = 21, antiderivative size = 169

$$\int (ex)^m \tanh^2 (d(a + b \log (cx^n))) dx$$

$$= \frac{(1 + m + bdn)(ex)^{1+m}}{bde(1 + m)n} + \frac{(ex)^{1+m} (1 - e^{2ad}(cx^n)^{2bd})}{bden (1 + e^{2ad}(cx^n)^{2bd})}$$

$$- \frac{2(ex)^{1+m} \text{Hypergeometric2F1} \left( 1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right)}{bden}$$

output

```
(b*d*n+m+1)*(e*x)^(1+m)/b/d/e/(1+m)/n+(e*x)^(1+m)*(1-exp(2*a*d)*(c*x^n)^(2
*b*d))/b/d/e/n/(1+exp(2*a*d)*(c*x^n)^(2*b*d))-2*(e*x)^(1+m)*hypergeom([1,
1/2*(1+m)/b/d/n],[1+1/2*(1+m)/b/d/n],[-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/e/n
```

**Mathematica [A] (verified)**

Time = 11.97 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.88

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = (ex)^m \left( \frac{x}{1+m} \right. \\ \left. e^{-\frac{(1+2m)(a-bn \log(x)+b \log(cx^n))}{bn}} x^{-2m} \left( e^{\frac{(1+2m)(a+b \log(cx^n))}{bn}} (1+m+2bdn) \operatorname{Hypergeometric2F1} \left( 1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn} \right) \right) \right)$$

input

```
Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

output

```
(e*x)^m*(x/(1+m) - (E^(((1+2*m)*(a + b*Log[c*x^n]))/(b*n))*(1+m+2*
b*d*n)*Hypergeometric2F1[1, (1+m)/(2*b*d*n), 1+(1+m)/(2*b*d*n), -E^(
2*d*(a + b*Log[c*x^n]))] - E^(((1+2*m+2*b*d*n)*(a - b*n*Log[x] + b*Log
[c*x^n]))/(b*n))*(1+m)*x^(1+2*m+2*b*d*n)*Hypergeometric2F1[1, (1+m
+2*b*d*n)/(2*b*d*n), (1+m+4*b*d*n)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x
^n]))] + E^(((1+2*m)*(a + b*Log[c*x^n]))/(b*n))*(1+m+2*b*d*n)*Tanh[d
*(a + b*Log[c*x^n]))]/(b*d*E^(((1+2*m)*(a - b*n*Log[x] + b*Log[c*x^n]))/
(b*n))*n*(1+m+2*b*d*n)*x^(2*m)))
```

**Rubi [A] (verified)**

Time = 0.63 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.26, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6073, 6071, 1004, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx \\ \downarrow 6073 \\ \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int (cx^n)^{\frac{m+1}{n}-1} \tanh^2(d(a + b \log(cx^n))) d(cx^n)}{en}$$

$$\begin{array}{c}
 \downarrow 6071 \\
 \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int \frac{(cx^n)^{\frac{m+1}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)^2}{(e^{2ad}(cx^n)^{2bd}+1)^2} d(cx^n)}{en} \\
 \downarrow 1004 \\
 \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{(cx^n)^{\frac{m+1}{n}} (1-e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd}+1)} - \frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{m+1}{n}-1} \left( \frac{e^{2ad}(m-bdn+1)}{n} - \frac{e^{4ad}(m+bdn+1)(cx^n)^{2bd}}{n} \right)}{e^{2ad}(cx^n)^{2bd}+1} d(cx^n)}{2bd} \right)}{en} \\
 \downarrow 27 \\
 \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{(cx^n)^{\frac{m+1}{n}} (1-e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd}+1)} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{m+1}{n}-1} \left( \frac{e^{2ad}(m-bdn+1)}{n} - \frac{e^{4ad}(m+bdn+1)(cx^n)^{2bd}}{n} \right)}{e^{2ad}(cx^n)^{2bd}+1} d(cx^n)}{bd} \right)}{en} \\
 \downarrow 959 \\
 \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{(cx^n)^{\frac{m+1}{n}} (1-e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd}+1)} - \frac{e^{-2ad} \left( \frac{2(m+1)e^{2ad} \int \frac{(cx^n)^{\frac{m+1}{n}-1}}{e^{2ad}(cx^n)^{2bd}+1} d(cx^n)}{n} - \frac{e^{2ad}(bdn+m+1)(cx^n)^{\frac{m+1}{n}}}{m+1} \right)}{bd} \right)}{en} \\
 \downarrow 888 \\
 \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{(cx^n)^{\frac{m+1}{n}} (1-e^{2ad}(cx^n)^{2bd})}{bd(e^{2ad}(cx^n)^{2bd}+1)} - \frac{e^{-2ad} \left( 2e^{2ad}(cx^n)^{\frac{m+1}{n}} \text{Hypergeometric2F1} \left( 1, \frac{m+1}{2bdn}, \frac{m+1}{2bdn}+1, -e^{2ad}(cx^n)^{2bd} \right) \right)}{bd} \right)}{en}
 \end{array}$$

input

`Int[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^2,x]`

output 
$$\frac{((e*x)^{(1+m)} * (((c*x^n)^{((1+m)/n)} * (1 - E^{(2*a*d)} * (c*x^n)^{(2*b*d)})) / (b*d * (1 + E^{(2*a*d)} * (c*x^n)^{(2*b*d)})) - ((E^{(2*a*d)} * (1+m + b*d*n) * (c*x^n)^{((1+m)/n)}) / (1+m)) + 2 * E^{(2*a*d)} * (c*x^n)^{((1+m)/n)} * \text{Hypergeometric2F1}[1, (1+m)/(2*b*d*n), 1 + (1+m)/(2*b*d*n), -(E^{(2*a*d)} * (c*x^n)^{(2*b*d)})] / (b*d * E^{(2*a*d)})) / (e*n * (c*x^n)^{((1+m)/n)})$$

### Defintions of rubi rules used

rule 27 
$$\text{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$$

rule 888 
$$\text{Int}[(c_*)(x_*)^{(m_*)} * ((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)} / (c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}[\{a, b, c, m, n, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{LtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$$

rule 959 
$$\text{Int}[(e_*)(x_*)^{(m_*)} * ((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)} * ((c_*) + (d_*)(x_*)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d * (e*x)^{(m+1)} * ((a + b*x^n)^{(p+1)} / (b * e * (m + n * (p + 1) + 1))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m + n*(p+1) + 1)) / (b*(m + n*(p + 1) + 1)) \text{ Int}[(e*x)^m * (a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p + 1) + 1, 0]$$

rule 1004 
$$\text{Int}[(e_*)(x_*)^{(m_*)} * ((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)} * ((c_*) + (d_*)(x_*)^{(n_*)})^{(q_*)}, x\_Symbol] \rightarrow \text{Simp}[(-c*b - a*d) * (e*x)^{(m+1)} * (a + b*x^n)^{(p+1)} * ((c + d*x^n)^{(q-1)} / (a*b*e*n*(p+1))), x] + \text{Simp}[1 / (a*b*n*(p+1)) \text{ Int}[(e*x)^m * (a + b*x^n)^{(p+1)} * (c + d*x^n)^{(q-2)} * \text{Simp}[c*(c*b*n*(p+1) + (c*b - a*d)*(m+1)) + d*(c*b*n*(p+1) + (c*b - a*d)*(m + n*(q-1) + 1)) * x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[q, 1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$$

rule 6071 
$$\text{Int}[(e_*)(x_*)^{(m_*)} * \text{Tanh}[(a_*) + \text{Log}[x_]* (b_*) * (d_*)]^{(p_*)}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m * ((-1 + E^{(2*a*d)} * x^{(2*b*d)})^p / (1 + E^{(2*a*d)} * x^{(2*b*d)}))^p, x] /; \text{FreeQ}[\{a, b, d, e, m, p\}, x]$$

rule 6073

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p
_.), x_Symbol] :> Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[
x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a,
b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

**Maple [F]**

$$\int (ex)^m \tanh(d(a + b \ln(cx^n)))^2 dx$$

input

```
int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^2,x)
```

output

```
int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^2,x)
```

**Fricas [F]**

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^2 dx$$

input

```
integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")
```

output

```
integral((e*x)^m*tanh(b*d*log(c*x^n) + a*d)^2, x)
```

**Sympy [F]**

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh^2(ad + bd \log(cx^n)) dx$$

input

```
integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n)))**2,x)
```

output

```
Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n))**2, x)
```

**Maxima [F]**

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

output `-2*e^m*(m + 1)*integrate(x^m/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x) + (b*c^(2*b*d)*d*e^m*n*x*e^(2*b*d*log(x^n) + 2*a*d + m*log(x)) + (b*d*e^m*n + 2*e^m*(m + 1))*x*x^m)/((m*n + n)*b*c^(2*b*d)*d*e^(2*b*d*log(x^n) + 2*a*d) + (m*n + n)*b*d)`

**Giac [F]**

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^2 dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

output `integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n)))^2 (ex)^m dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^2*(e*x)^m,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^2*(e*x)^m, x)`

**Reduce [F]**

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx$$

$$= \frac{e^m \left( -4e^{2ad} c^{2bd} \left( \int \frac{x^{2bdn+m}}{x^{4bdn} e^{4ad} c^{4bd} + 2x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) m - 4e^{2ad} c^{2bd} \left( \int \frac{x^{2bdn+m}}{x^{4bdn} e^{4ad} c^{4bd} + 2x^{2bdn} e^{2ad} c^{2bd} + 1} dx \right) + x^m x \right)}{m + 1}$$

input `int((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x)`

output `(e**m*( - 4*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n + m)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x)*m - 4*e**(2*a*d)*c**(2*b*d)*int(x**(2*b*d*n + m)/(x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d) + 2*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 1),x) + x**m*x)/(m + 1)`

### 3.201 $\int (ex)^m \tanh^3 (d(a + b \log (cx^n))) dx$

Optimal result	1487
Mathematica [A] (verified)	1488
Rubi [A] (verified)	1489
Maple [F]	1493
Fricas [F]	1493
Sympy [F]	1493
Maxima [F]	1494
Giac [F]	1494
Mupad [F(-1)]	1494
Reduce [F]	1495

#### Optimal result

Integrand size = 21, antiderivative size = 307

$$\int (ex)^m \tanh^3 (d(a + b \log (cx^n))) dx = \frac{(1 + m + bdn)(1 + m + 2bdn)(ex)^{1+m}}{2b^2d^2e(1 + m)n^2} - \frac{(ex)^{1+m} (1 - e^{2ad}(cx^n)^{2bd})^2}{2bden (1 + e^{2ad} (cx^n)^{2bd})^2} + \frac{e^{-2ad}(ex)^{1+m} \left( \frac{e^{2ad}(1+m-2bdn)}{n} - \frac{e^{4ad}(1+m+2bdn)(cx^n)^{2bd}}{n} \right)}{2b^2d^2en (1 + e^{2ad} (cx^n)^{2bd})} - \frac{(1 + 2m + m^2 + 2b^2d^2n^2) (ex)^{1+m} \text{Hypergeometric2F1} \left( 1, \frac{1+m}{2bdn}, 1 + \frac{1+m}{2bdn}, -e^{2ad}(cx^n)^{2bd} \right)}{b^2d^2e(1 + m)n^2}$$

output

```
1/2*(b*d*n+m+1)*(2*b*d*n+m+1)*(e*x)^(1+m)/b^2/d^2/e/(1+m)/n^2-1/2*(e*x)^(1+m)*(1-exp(2*a*d)*(c*x^n)^(2*b*d))^2/b/d/e/n/(1+exp(2*a*d)*(c*x^n)^(2*b*d))^2+1/2*(e*x)^(1+m)*(exp(2*a*d)*(-2*b*d*n+m+1)/n-exp(4*a*d)*(2*b*d*n+m+1)*(c*x^n)^(2*b*d)/n)/b^2/d^2/e/exp(2*a*d)/n/(1+exp(2*a*d)*(c*x^n)^(2*b*d))- (2*b^2*d^2*n^2+m^2+2*m+1)*(e*x)^(1+m)*hypergeom([1, 1/2*(1+m)/b/d/n],[1+1/2*(1+m)/b/d/n],-exp(2*a*d)*(c*x^n)^(2*b*d))/b^2/d^2/e/(1+m)/n^2
```



**Mathematica [A] (verified)**

Time = 13.10 (sec) , antiderivative size = 606, normalized size of antiderivative = 1.97

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx$$

$$= \frac{x(ex)^m \operatorname{sech}^2(bdn \log(x) + d(a + b(-n \log(x) + \log(cx^n))))}{2bdn} - \frac{(1+m)x(ex)^m \operatorname{sech}(d(a + b(-n \log(x) + \log(cx^n)))) \operatorname{sech}(bdn \log(x) + d(a + b(-n \log(x) + \log(cx^n))))}{2b^2d^2n^2}$$

$$+ \frac{(1+2m+m^2+2b^2d^2n^2)x^{-m}(ex)^m \operatorname{sech}(d(a + b(-n \log(x) + \log(cx^n))))}{1+m} \left( \frac{x^{1+m} \operatorname{sech}(d(a+b \log(cx^n))) \sinh(d(a+b \log(cx^n)))}{1+m} \right)$$

$$+ \frac{x(ex)^m \tanh(d(a + b(-n \log(x) + \log(cx^n))))}{1+m}$$

input `Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^3,x]`

output

```
(x*(e*x)^m*Sech[b*d*n*Log[x] + d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]^2)/(2
*b*d*n) - ((1 + m)*x*(e*x)^m*Sech[d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*Se
ch[b*d*n*Log[x] + d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*Sinh[b*d*n*Log[x]]
)/(2*b^2*d^2*n^2) + ((1 + 2*m + m^2 + 2*b^2*d^2*n^2)*(e*x)^m*Sech[d*(a + b
*(-(n*Log[x]) + Log[c*x^n]))]*(x^(1 + m)*Sech[d*(a + b*Log[c*x^n]])*Sinh[
b*d*n*Log[x]])/(1 + m) - (Cosh[d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*(E^((
a + 2*a*m + b*(1 + m)*n*Log[x] + b*(1 + 2*m)*(-(n*Log[x]) + Log[c*x^n])))/(
b*n))*(1 + m + 2*b*d*n)*Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m
)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - E^((a*(1 + 2*m + 2*b*d*n))/(b*
n) + (1 + m + 2*b*d*n)*Log[x] + ((1 + 2*m + 2*b*d*n)*(-(n*Log[x]) + Log[c*
x^n]))/n)*(1 + m)*Hypergeometric2F1[1, (1 + m + 2*b*d*n)/(2*b*d*n), (1 + m
+ 4*b*d*n)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + E^((a + 2*a*m + b*(1
+ m)*n*Log[x] + b*(1 + 2*m)*(-(n*Log[x]) + Log[c*x^n]))/(b*n))*(1 + m + 2
*b*d*n)*Tanh[d*(a + b*Log[c*x^n])])/(E^(((1 + 2*m)*(a + b*(-(n*Log[x]) +
Log[c*x^n])))/(b*n))*(1 + m)*(1 + m + 2*b*d*n)))/(2*b^2*d^2*n^2*x^m) + (x
*(e*x)^m*Tanh[d*(a + b*(-(n*Log[x]) + Log[c*x^n]))])/(1 + m)
```

**Rubi [A] (verified)**

Time = 0.92 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$ , Rules used = {6073, 6071, 1004, 27, 1064, 27, 959, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx \\
 & \quad \downarrow \text{6073} \\
 & \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int (cx^n)^{\frac{m+1}{n}-1} \tanh^3(d(a + b \log(cx^n))) d(cx^n)}{en} \\
 & \quad \downarrow \text{6071} \\
 & \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int \frac{(cx^n)^{\frac{m+1}{n}-1} (e^{2ad}(cx^n)^{2bd}-1)^3}{(e^{2ad}(cx^n)^{2bd}+1)^3} d(cx^n)}{en} \\
 & \quad \downarrow \text{1004} \\
 & \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( -\frac{e^{-2ad} \int \frac{2(cx^n)^{\frac{m+1}{n}-1} (1-e^{2ad}(cx^n)^{2bd}) \left( \frac{e^{2ad}(m-2bdn+1)}{n} - \frac{e^{4ad}(m+2bdn+1)(cx^n)^{2bd}}{n} \right) d(cx^n)}{(e^{2ad}(cx^n)^{2bd}+1)^2}}{4bd} - \frac{(cx^n)^{\frac{m+1}{n}}}{2bd(e^{2ad}(cx^n)^{2bd}+1)} \right)}{en} \\
 & \quad \downarrow \text{27} \\
 & \frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{m+1}{n}-1} (1-e^{2ad}(cx^n)^{2bd}) \left( \frac{e^{2ad}(m-2bdn+1)}{n} - \frac{e^{4ad}(m+2bdn+1)(cx^n)^{2bd}}{n} \right) d(cx^n)}{(e^{2ad}(cx^n)^{2bd}+1)^2}}{2bd} - \frac{(cx^n)^{\frac{m+1}{n}} (1-e^{2ad}(cx^n)^{2bd})}{2bd(e^{2ad}(cx^n)^{2bd}+1)} \right)}{en} \\
 & \quad \downarrow \text{1064}
 \end{aligned}$$

$$(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{e^{-2ad} \left( (cx^n)^{\frac{m+1}{n}} \left( \frac{e^{2ad(-2bdn+m+1)}}{n} - \frac{e^{4ad(2bdn+m+1)}(cx^n)2bd}{n} \right) \right)}{bd(e^{2ad}(cx^n)^{2bd+1})} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{m+1}{n}-1} \left( \frac{e^{4ad(m-2bdn+1)}(m-bd)}{n^2} \right)}{e^{2ad}}}{2bd} \right)$$

en

↓ 27

$$(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{e^{-2ad} \left( (cx^n)^{\frac{m+1}{n}} \left( \frac{e^{2ad(-2bdn+m+1)}}{n} - \frac{e^{4ad(2bdn+m+1)}(cx^n)2bd}{n} \right) \right)}{bd(e^{2ad}(cx^n)^{2bd+1})} - \frac{e^{-2ad} \int \frac{(cx^n)^{\frac{m+1}{n}-1} \left( \frac{e^{4ad(m-2bdn+1)}(m-bd)}{n^2} \right)}{e^{2ad}}}{2bd} \right)$$

en

↓ 959

$$(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{e^{-2ad} \left( (cx^n)^{\frac{m+1}{n}} \left( \frac{e^{2ad(-2bdn+m+1)}}{n} - \frac{e^{4ad(2bdn+m+1)}(cx^n)2bd}{n} \right) \right)}{bd(e^{2ad}(cx^n)^{2bd+1})} - \frac{e^{-2ad} \left( \frac{2e^{4ad}(2b^2d^2n^2+m^2+2m+1) \int \frac{(cx^n)^{\frac{m}{n}}}{e^{2ad}(cx^n)^{\frac{m}{n}}}}{n^2} \right)}{2bd} \right)$$

en

↓ 888

$$(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \left( \frac{e^{-2ad} \left( \frac{(cx^n)^{\frac{m+1}{n}} \left( \frac{e^{2ad}(-2bdn+m+1)}{n} - \frac{e^{4ad}(2bdn+m+1)(cx^n)^{2bd}}{n} \right)}{bd(e^{2ad}(cx^n)^{2bd+1})} \right) - e^{-2ad} \left( \frac{2e^{4ad}(2b^2d^2n^2+m^2+2m+1)(cx^n)^{\frac{m+1}{n}}}{2bd} \right)}{2bd} \right)$$

en

input `Int[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^3,x]`

output `((e*x)^(1 + m)*(-1/2*((c*x^n)^((1 + m)/n)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d))^2)/(b*d*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))^2) + (((c*x^n)^((1 + m)/n)*(E^(2*a*d)*(1 + m - 2*b*d*n))/n - (E^(4*a*d)*(1 + m + 2*b*d*n)*(c*x^n)^(2*b*d))/n))/(b*d*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) - (-((E^(4*a*d)*(1 + m + b*d*n)*(1 + m + 2*b*d*n)*(c*x^n)^((1 + m)/n))/((1 + m)*n)) + (2*E^(4*a*d)*(1 + 2*m + m^2 + 2*b^2*d^2*n^2)*(c*x^n)^((1 + m)/n)*Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m)/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/((1 + m)*n))/(b*d*E^(2*a*d))/(2*b*d*E^(2*a*d)))/(e*n*(c*x^n)^((1 + m)/n))`

**Defintions of rubi rules used**

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 888 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 959

```
Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p._)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Simp[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

rule 1004

```
Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p._)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-(c*b - a*d)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Simp[1/(a*b*n*(p + 1)) Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

rule 1064

```
Int[((g._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p._)*((c_) + (d._)*(x_)^(n_))^(q_)*(e_) + (f._)*(x_)^(n_)), x_Symbol] := Simp[(-(b*e - a*f))*(g*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(a*b*g*n*(p + 1))), x] + Simp[1/(a*b*n*(p + 1)) Int[(g*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + (b*e - a*f)*(m + 1)) + d*(b*e*n*(p + 1) + (b*e - a*f)*(m + n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && LtQ[p, -1] && GtQ[q, 0] && !(EqQ[q, 1] && SimplerQ[b*c - a*d, b*e - a*f])
```

rule 6071

```
Int[((e._)*(x_))^(m._)*Tanh[((a_) + Log[x]*(b._))*(d._)]^(p._), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]
```

rule 6073

```
Int[((e._)*(x_))^(m._)*Tanh[((a_) + Log[(c._)*(x_)^(n_.)]*(b._))*(d._)]^(p._), x_Symbol] := Simp[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)) Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

**Maple [F]**

$$\int (ex)^m \tanh(d(a + b \ln(cx^n)))^3 dx$$

input `int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^3,x)`

output `int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^3,x)`

**Fricas [F]**

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^3 dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="fricas")`

output `integral((e*x)^m*tanh(b*d*log(c*x^n) + a*d)^3, x)`

**Sympy [F]**

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh^3(ad + bd \log(cx^n)) dx$$

input `integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))))**3,x)`

output `Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n))**3, x)`

**Maxima [F]**

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^3 dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="maxima")`

output `-(2*b^2*d^2*e^m*n^2 + (m^2 + 2*m + 1)*e^m)*integrate(x^m/(b^2*c^(2*b*d)*d^2*n^2*e^(2*b*d*log(x^n) + 2*a*d) + b^2*d^2*n^2), x) + (b^2*c^(4*b*d)*d^2*e^m*n^2*x*e^(4*b*d*log(x^n) + 4*a*d + m*log(x)) + (b^2*d^2*e^m*n^2 + (m^2 + 2*m + 1)*e^m)*x*x^m + (2*b^2*c^(2*b*d)*d^2*e^m*n^2*e^(2*a*d) + 2*(m*n + n)*b*c^(2*b*d)*d*e^m*e^(2*a*d) + (m^2 + 2*m + 1)*c^(2*b*d)*e^m*e^(2*a*d))*x*e^(2*b*d*log(x^n) + m*log(x)))/((m*n^2 + n^2)*b^2*c^(4*b*d)*d^2*e^(4*b*d*log(x^n) + 4*a*d) + 2*(m*n^2 + n^2)*b^2*c^(2*b*d)*d^2*e^(2*b*d*log(x^n) + 2*a*d) + (m*n^2 + n^2)*b^2*d^2)`

**Giac [F]**

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^3 dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="giac")`

output `integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n)))^3 (ex)^m dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^3*(e*x)^m,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^3*(e*x)^m, x)`

## Reduce [F]

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \text{too large to display}$$

input `int((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x)`

output

```
(e**m*(8*x**(4*b*d*n + m)*e**(4*a*d)*c**(4*b*d)*b**2*d**2*n**2*x - 6*x**(4
*b*d*n + m)*e**(4*a*d)*c**(4*b*d)*b*d*m*n*x - 6*x**(4*b*d*n + m)*e**(4*a*d
)*c**(4*b*d)*b*d*n*x + x**(4*b*d*n + m)*e**(4*a*d)*c**(4*b*d)*m**2*x + 2*x
**(4*b*d*n + m)*e**(4*a*d)*c**(4*b*d)*m*x + x**(4*b*d*n + m)*e**(4*a*d)*c*
*(4*b*d)*x - 128*x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*int(x**m/(8*x**(6*b*d*
n)*e**(6*a*d)*c**(6*b*d)*b**2*d**2*n**2 - 6*x**(6*b*d*n)*e**(6*a*d)*c**(6*
b*d)*b*d*m*n - 6*x**(6*b*d*n)*e**(6*a*d)*c**(6*b*d)*b*d*n + x**(6*b*d*n)*e
**(6*a*d)*c**(6*b*d)*m**2 + 2*x**(6*b*d*n)*e**(6*a*d)*c**(6*b*d)*m + x**(6
*b*d*n)*e**(6*a*d)*c**(6*b*d) + 24*x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*b**2
*d**2*n**2 - 18*x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*b*d*m*n - 18*x**(4*b*d*
n)*e**(4*a*d)*c**(4*b*d)*b*d*n + 3*x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*m**2
+ 6*x**(4*b*d*n)*e**(4*a*d)*c**(4*b*d)*m + 3*x**(4*b*d*n)*e**(4*a*d)*c**(
4*b*d) + 24*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*b**2*d**2*n**2 - 18*x**(2*b
*d*n)*e**(2*a*d)*c**(2*b*d)*b*d*m*n - 18*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d
)*b*d*n + 3*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d)*m**2 + 6*x**(2*b*d*n)*e**(2
*a*d)*c**(2*b*d)*m + 3*x**(2*b*d*n)*e**(2*a*d)*c**(2*b*d) + 8*b**2*d**2*n*
*2 - 6*b*d*m*n - 6*b*d*n + m**2 + 2*m + 1),x)*b**4*d**4*m*n**4 - 128*x**(4
*b*d*n)*e**(4*a*d)*c**(4*b*d)*int(x**m/(8*x**(6*b*d*n)*e**(6*a*d)*c**(6*b*
d)*b**2*d**2*n**2 - 6*x**(6*b*d*n)*e**(6*a*d)*c**(6*b*d)*b*d*m*n - 6*x**(6
*b*d*n)*e**(6*a*d)*c**(6*b*d)*b*d*n + x**(6*b*d*n)*e**(6*a*d)*c**(6*b*d...
```



### 3.202 $\int \tanh^p (d(a + b \log (cx^n))) dx$

Optimal result	1496
Mathematica [B] (warning: unable to verify)	1496
Rubi [A] (verified)	1497
Maple [F]	1499
Fricas [F]	1499
Sympy [F]	1499
Maxima [F]	1500
Giac [F]	1500
Mupad [F(-1)]	1500
Reduce [F]	1501

#### Optimal result

Integrand size = 15, antiderivative size = 115

$$\int \tanh^p (d(a + b \log (cx^n))) dx = x \left( 1 - e^{2ad} (cx^n)^{2bd} \right)^{-p} \left( -1 + e^{2ad} (cx^n)^{2bd} \right)^p \operatorname{AppellF1} \left( \frac{1}{2bdn}, -p, p, 1 + \frac{1}{2bdn}, \exp(2ad) (cx^n)^{2bd}, -\exp(2ad) (cx^n)^{2bd} \right)$$

output

```
x*(-1+exp(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1(1/2/b/d/n,-p,p,1+1/2/b/d/n,exp(2*a*d)*(c*x^n)^(2*b*d),-exp(2*a*d)*(c*x^n)^(2*b*d))/((1-exp(2*a*d)*(c*x^n)^(2*b*d))^p)
```

#### Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 387 vs. 2(115) = 230.

Time = 0.88 (sec) , antiderivative size = 387, normalized size of antiderivative = 3.37

$$\int \tanh^p (d(a + b \log (cx^n))) dx = \frac{(1 + 2bdn)x \left( \frac{-1 + e^{2ad}(cx^n)^{2bd}}{1 + e^{2ad}(cx^n)^{2bd}} \right)^p}{-2bde^{2ad}np (cx^n)^{2bd} \operatorname{AppellF1} \left( 1 + \frac{1}{2bdn}, 1 - p, p, 2 + \frac{1}{2bdn}, e^{2ad} (cx^n)^{2bd}, -e^{2ad} (cx^n)^{2bd} \right) - 2bde^{2ad}np (cx^n)^{2bd}}$$

input `Integrate[Tanh[d*(a + b*Log[c*x^n])]^p,x]`

output 
$$\left( (1 + 2*b*d*n)*x*((-1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)})}/(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}))\right)^p \text{AppellF1}\left[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^{(2*a*d)*(c*x^n)^{(2*b*d)}}, -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right] / (-2*b*d*E^{(2*a*d)*n*p*(c*x^n)^{(2*b*d)}}*\text{AppellF1}\left[1 + 1/(2*b*d*n), 1 - p, p, 2 + 1/(2*b*d*n), E^{(2*a*d)*(c*x^n)^{(2*b*d)}}, -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right] - 2*b*d*E^{(2*a*d)*n*p*(c*x^n)^{(2*b*d)}}*\text{AppellF1}\left[1 + 1/(2*b*d*n), -p, 1 + p, 2 + 1/(2*b*d*n), E^{(2*a*d)*(c*x^n)^{(2*b*d)}}, -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right] + (1 + 2*b*d*n)*\text{AppellF1}\left[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^{(2*a*d)*(c*x^n)^{(2*b*d)}}, -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right]$$

### Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {6069, 6071, 1013, 1012}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tanh^p(d(a + b \log(cx^n))) dx$$

$$\downarrow 6069$$

$$\frac{x(cx^n)^{-1/n} \int (cx^n)^{\frac{1}{n}-1} \tanh^p(d(a + b \log(cx^n))) d(cx^n)}{n}$$

$$\downarrow 6071$$

$$\frac{x(cx^n)^{-1/n} \int (cx^n)^{\frac{1}{n}-1} \left(e^{2ad}(cx^n)^{2bd} - 1\right)^p \left(e^{2ad}(cx^n)^{2bd} + 1\right)^{-p} d(cx^n)}{n}$$

$$\downarrow 1013$$

$$\frac{x(cx^n)^{-1/n} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(e^{2ad}(cx^n)^{2bd} - 1\right)^p \int (cx^n)^{\frac{1}{n}-1} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^p \left(e^{2ad}(cx^n)^{2bd} + 1\right)^{-p} d(cx^n)}{n}$$

$$\downarrow 1012$$

$$x \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(e^{2ad}(cx^n)^{2bd} - 1\right)^p \operatorname{AppellF1} \left(\frac{1}{2bdn}, -p, p, 1 + \frac{1}{2bdn}, e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)$$

input `Int [Tanh [d*(a + b*Log [c*x^n])]^p, x]`

output `(x*(-1 + E^(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/(1 - E^(2*a*d)*(c*x^n)^(2*b*d))^p`

### Defintions of rubi rules used

rule 1012 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 1013 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a)^FracPart[p]) Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 6069 `Int [Tanh [(a_.) + Log [(c_.)*(x_)^(n_.)]*(b_.)]*(d_.)]^(p_.), x_Symbol] := Simp[x/(n*(c*x^n)^(1/n)) Subst [Int [x^(1/n - 1)*Tanh [d*(a + b*Log [x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])`

rule 6071 `Int[((e_.)*(x_))^(m_.)*Tanh[(a_.) + Log[x]*(b_.)]*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]`

**Maple [F]**

$$\int \tanh(d(a + b \ln(cx^n)))^p dx$$

input `int(tanh(d*(a+b*ln(c*x^n)))^p,x)`

output `int(tanh(d*(a+b*ln(c*x^n)))^p,x)`

**Fricas [F]**

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^p dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="fricas")`

output `integral(tanh(b*d*log(c*x^n) + a*d)^p, x)`

**Sympy [F]**

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh^p(d(a + b \log(cx^n))) dx$$

input `integrate(tanh(d*(a+b*ln(c*x**n)))**p,x)`

output `Integral(tanh(d*(a + b*log(c*x**n)))**p, x)`

**Maxima [F]**

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^p dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="maxima")`

output `integrate(tanh((b*log(c*x^n) + a)*d)^p, x)`

**Giac [F]**

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh((b \log(cx^n) + a)d)^p dx$$

input `integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="giac")`

output `integrate(tanh((b*log(c*x^n) + a)*d)^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \ln(cx^n)))^p dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^p,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^p, x)`

**Reduce [F]**

$$\int \tanh^p (d(a + b \log (cx^n))) dx = \tanh (\log (x^n c) bd + ad)^p x$$

$$- \left( \int \frac{\tanh (\log (x^n c) bd + ad)^p}{\tanh (\log (x^n c) bd + ad)} dx \right) bdn p$$

$$+ \left( \int \tanh (\log (x^n c) bd + ad)^p \tanh (\log (x^n c) bd + ad) dx \right) bdn p$$

input `int(tanh(d*(a+b*log(c*x^n)))^p,x)`

output `tanh(log(x**n*c)*b*d + a*d)**p*x - int(tanh(log(x**n*c)*b*d + a*d)**p/tanh(log(x**n*c)*b*d + a*d),x)*b*d*n*p + int(tanh(log(x**n*c)*b*d + a*d)**p*tanh(log(x**n*c)*b*d + a*d),x)*b*d*n*p`

### 3.203 $\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx$

Optimal result	1502
Mathematica [A] (warning: unable to verify)	1502
Rubi [A] (verified)	1503
Maple [F]	1505
Fricas [F]	1505
Sympy [F(-1)]	1505
Maxima [F]	1506
Giac [F]	1506
Mupad [F(-1)]	1506
Reduce [F]	1507

#### Optimal result

Integrand size = 21, antiderivative size = 135

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx$$

$$= \frac{(ex)^{1+m} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(-1 + e^{2ad}(cx^n)^{2bd}\right)^p \operatorname{AppellF1}\left(\frac{1+m}{2bdn}, -p, p, 1 + \frac{1+m}{2bdn}, e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{e(1+m)}$$

output

$$\frac{(e*x)^{(1+m)}*(-1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})^p*\operatorname{AppellF1}\left(\frac{1}{2}*(1+m)/b/d/n,-p,p,1+1/2*(1+m)/b/d/n,\exp(2*a*d)*(c*x^n)^{(2*b*d)},-\exp(2*a*d)*(c*x^n)^{(2*b*d)}\right)}{e/(1+m)/((1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})^p)}$$

#### Mathematica [A] (warning: unable to verify)

Time = 0.75 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.29

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx$$

$$= \frac{x(ex)^m \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(\frac{-1+e^{2ad}(cx^n)^{2bd}}{1+e^{2ad}(cx^n)^{2bd}}\right)^p \left(1 + e^{2ad}(cx^n)^{2bd}\right)^p \operatorname{AppellF1}\left(\frac{1+m}{2bdn}, -p, p, 1 + \frac{1+m}{2bdn}, e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{1+m}$$

input

$$\operatorname{Integrate}[(e*x)^m*\operatorname{Tanh}[d*(a + b*\operatorname{Log}[c*x^n])]^p,x]$$

output

```
(x*(e*x)^m*((-1 + E^(2*a*d)*(c*x^n)^(2*b*d))/(1 + E^(2*a*d)*(c*x^n)^(2*b*d)))^p*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1[(1 + m)/(2*b*d*n), -p, p, 1 + (1 + m)/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/((1 + m)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d))^p)
```

**Rubi [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6073, 6071, 1013, 1012}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx$$

$$\downarrow 6073$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int (cx^n)^{\frac{m+1}{n}-1} \tanh^p(d(a + b \log(cx^n))) d(cx^n)}{en}$$

$$\downarrow 6071$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} \int (cx^n)^{\frac{m+1}{n}-1} (e^{2ad}(cx^n)^{2bd} - 1)^p (e^{2ad}(cx^n)^{2bd} + 1)^{-p} d(cx^n)}{en}$$

$$\downarrow 1013$$

$$\frac{(ex)^{m+1} (cx^n)^{-\frac{m+1}{n}} (1 - e^{2ad}(cx^n)^{2bd})^{-p} (e^{2ad}(cx^n)^{2bd} - 1)^p \int (cx^n)^{\frac{m+1}{n}-1} (1 - e^{2ad}(cx^n)^{2bd})^p (e^{2ad}(cx^n)^{2bd} - 1)^{-p} d(cx^n)}{en}$$

$$\downarrow 1012$$

$$\frac{(ex)^{m+1} (1 - e^{2ad}(cx^n)^{2bd})^{-p} (e^{2ad}(cx^n)^{2bd} - 1)^p \text{AppellF1}\left(\frac{m+1}{2bdn}, -p, p, \frac{m+1}{2bdn} + 1, e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{e(m+1)}$$

input

```
Int[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^p,x]
```



output

$$\frac{((e*x)^{(1+m)}*(-1 + E^{(2*a*d)}*(c*x^n)^{(2*b*d)})^p * \text{AppellF1}[(1+m)/(2*b*d*n), -p, p, 1 + (1+m)/(2*b*d*n), E^{(2*a*d)}*(c*x^n)^{(2*b*d)}, -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})])}{(e*(1+m)*(1 - E^{(2*a*d)}*(c*x^n)^{(2*b*d)})^p)}$$
**Defintions of rubi rules used**

rule 1012

$$\text{Int}[(e_{.})*(x_{.})^{(m_{.})}*((a_{.}) + (b_{.})*(x_{.})^{(n_{.})})^{(p_{.})}*((c_{.}) + (d_{.})*(x_{.})^{(n_{.})})^{(q_{.})}, x\_Symbol] \rightarrow \text{Simp}[a^p*c^q*((e*x)^{(m+1)}/(e*(m+1))) * \text{AppellF1}[(m+1)/n, -p, -q, 1 + (m+1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$$

FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b\*c - a\*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

rule 1013

$$\text{Int}[(e_{.})*(x_{.})^{(m_{.})}*((a_{.}) + (b_{.})*(x_{.})^{(n_{.})})^{(p_{.})}*((c_{.}) + (d_{.})*(x_{.})^{(n_{.})})^{(q_{.})}, x\_Symbol] \rightarrow \text{Simp}[a^p * \text{IntPart}[p] * ((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a)^{\text{FracPart}[p]})) \text{Int}[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$$

FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b\*c - a\*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])

rule 6071

$$\text{Int}[(e_{.})*(x_{.})^{(m_{.})} * \text{Tanh}[(a_{.}) + \text{Log}[x_{.}] * (b_{.}) * (d_{.})]^{(p_{.})}, x\_Symbol] \rightarrow \text{Int}[(e*x)^m * ((-1 + E^{(2*a*d)}*x^{(2*b*d)})^p / (1 + E^{(2*a*d)}*x^{(2*b*d)})^p), x] /;$$

FreeQ[{a, b, d, e, m, p}, x]

rule 6073

$$\text{Int}[(e_{.})*(x_{.})^{(m_{.})} * \text{Tanh}[(a_{.}) + \text{Log}[(c_{.})*(x_{.})^{(n_{.})}] * (b_{.}) * (d_{.})]^{(p_{.})}, x\_Symbol] \rightarrow \text{Simp}[(e*x)^{(m+1)}/(e*n*(c*x^n)^{((m+1)/n)}) \text{Subst}[\text{Int}[x^{((m+1)/n - 1)} * \text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /;$$

FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

**Maple [F]**

$$\int (ex)^m \tanh(d(a + b \ln(cx^n)))^p dx$$

input `int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^p,x)`

output `int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^p,x)`

**Fricas [F]**

$$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh((b \log(cx^n) + a)d)^p dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="fricas")`

output `integral((e*x)^m*tanh(b*d*log(c*x^n) + a*d)^p, x)`

**Sympy [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx = \text{Timed out}$$

input `integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n)))**p,x)`

output `Timed out`

**Maxima [F]**

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx = \int (ex)^m \tanh ((b \log (cx^n) + a)d)^p dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="maxima")`

output `integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^p, x)`

**Giac [F]**

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx = \int (ex)^m \tanh ((b \log (cx^n) + a)d)^p dx$$

input `integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="giac")`

output `integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^p, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx = \int \tanh(d(a + b \ln (cx^n)))^p (ex)^m dx$$

input `int(tanh(d*(a + b*log(c*x^n)))^p*(e*x)^m,x)`

output `int(tanh(d*(a + b*log(c*x^n)))^p*(e*x)^m, x)`

**Reduce [F]**

$$\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx$$

$$= \frac{e^m \left( x^m \tanh (\log (x^n c) b d + a d)^p x - \left( \int \frac{x^m \tanh (\log (x^n c) b d + a d)^p}{\tanh (\log (x^n c) b d + a d)} dx \right) b d n p + \left( \int x^m \tanh (\log (x^n c) b d + a d)^p \tanh (\log (x^n c) b d + a d) dx \right) \right)}{m + 1}$$

input `int((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x)`

output `(e**m*(x**m*tanh(log(x**n*c)*b*d + a*d)**p*x - int((x**m*tanh(log(x**n*c)*b*d + a*d)**p)/tanh(log(x**n*c)*b*d + a*d),x)*b*d*n*p + int(x**m*tanh(log(x**n*c)*b*d + a*d)**p*tanh(log(x**n*c)*b*d + a*d),x)*b*d*n*p))/(m + 1)`

**3.204**  $\int \frac{\tanh^{\frac{5}{2}}(a+b \log (c x^n))}{x} d x$

Optimal result	1508
Mathematica [A] (verified)	1508
Rubi [A] (verified)	1509
Maple [A] (verified)	1512
Fricas [B] (verification not implemented)	1512
Sympy [F(-1)]	1513
Maxima [F]	1514
Giac [F(-1)]	1514
Mupad [B] (verification not implemented)	1514
Reduce [F]	1515

**Optimal result**

Integrand size = 19, antiderivative size = 73

$$\int \frac{\tanh^{\frac{5}{2}}(a+b \log (c x^n))}{x} d x = -\frac{\arctan \left(\sqrt{\tanh (a+b \log (c x^n))}\right)}{b n} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh (a+b \log (c x^n))}\right)}{b n} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log (c x^n))}{3 b n}$$

output -arctan(tanh(a+b\*ln(c\*x^n))^(1/2))/b/n+arctanh(tanh(a+b\*ln(c\*x^n))^(1/2))/b/n-2/3\*tanh(a+b\*ln(c\*x^n))^(3/2)/b/n

**Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.85

$$\int \frac{\tanh^{\frac{5}{2}}(a+b \log (c x^n))}{x} d x = \frac{\arctan \left(\sqrt{\tanh (a+b \log (c x^n))}\right) - \operatorname{arctanh}\left(\sqrt{\tanh (a+b \log (c x^n))}\right) + \frac{2}{3} \tanh^{\frac{3}{2}}(a+b \log (c x^n))}{b n}$$

input `Integrate[Tanh[a + b*Log[c*x^n]]^(5/2)/x,x]`

output `-((ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]] - ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]) + (2*Tanh[a + b*Log[c*x^n]]^(3/2))/3)/(b*n))`

### Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$ , Rules used = {3039, 3042, 3954, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n)) d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-i \tan(ia + ib \log(cx^n)))^{5/2} d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3954} \\
 & \int \frac{\sqrt{\tanh(a + b \log(cx^n))} d \log(cx^n) - \frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3b}}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3b} + \int \sqrt{-i \tan(ia + ib \log(cx^n))} d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3957} \\
 & \frac{-\frac{\int -\frac{\sqrt{\tanh(a + b \log(cx^n))}}{1 - \tanh^2(a + b \log(cx^n))} d \tanh(a + b \log(cx^n))}{b} - \frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3b}}{n} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{array}{c}
 \frac{\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{1-\tanh^2(a+b \log(cx^n))} d \tanh(a+b \log(cx^n))}{b} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3b} \\
 \hline
 n \\
 \downarrow 266 \\
 \frac{2 \int \frac{\tanh(a+b \log(cx^n))}{1-\tanh^2(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))}}{b} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3b} \\
 \hline
 n \\
 \downarrow 827 \\
 \frac{2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} - \frac{1}{2} \int \frac{1}{\tanh(a+b \log(cx^n))+1} d \sqrt{\tanh(a+b \log(cx^n))} \right)}{b} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3b} \\
 \hline
 n \\
 \downarrow 216 \\
 \frac{2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} - \frac{1}{2} \arctan \left( \sqrt{\tanh(a+b \log(cx^n))} \right) \right)}{b} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3b} \\
 \hline
 n \\
 \downarrow 219 \\
 \frac{2 \left( \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(a+b \log(cx^n))} \right) - \frac{1}{2} \arctan \left( \sqrt{\tanh(a+b \log(cx^n))} \right) \right)}{b} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3b} \\
 \hline
 n
 \end{array}$$

input `Int[Tanh[a + b*Log[c*x^n]]^(5/2)/x,x]`

output `((2*(-1/2*ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]])/2)/b - (2*Tanh[a + b*Log[c*x^n]]^(3/2))/(3*b))/n`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^m)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`



**Maple [A] (verified)**

Time = 1.45 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$\frac{-\frac{2\tanh(a+b\ln(cx^n))^{\frac{3}{2}}}{3} - \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))})}{nb}$	76
default	$\frac{-\frac{2\tanh(a+b\ln(cx^n))^{\frac{3}{2}}}{3} - \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))})}{nb}$	76

input `int(tanh(a+b*ln(c*x^n))^(5/2)/x,x,method=_RETURNVERBOSE)`

output `1/n/b*(-2/3*tanh(a+b*ln(c*x^n))^(3/2)-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)-arctan(tanh(a+b*ln(c*x^n))^(1/2)))`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 625 vs.  $2(65) = 130$ .

Time = 0.12 (sec) , antiderivative size = 625, normalized size of antiderivative = 8.56

$$\int \frac{\tanh^{\frac{5}{2}}(a+b\log(cx^n))}{x} dx = \text{Too large to display}$$

input `integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="fricas")`

output

```
-1/6*(6*(cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 +
1)*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^
2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)
*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*
sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + 4
*cosh(b*n*log(x) + b*log(c) + a)^2 + 3*(cosh(b*n*log(x) + b*log(c) + a)^2
+ 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh
(b*n*log(x) + b*log(c) + a)^2 + 1)*log(-cosh(b*n*log(x) + b*log(c) + a)^2
- 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh
(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cos
h(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*lo
g(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*
log(x) + b*log(c) + a))) + 8*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(
x) + b*log(c) + a) + 4*sinh(b*n*log(x) + b*log(c) + a)^2 + 4*(cosh(b*n*log
(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x)
+ b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 - 1)*sqrt(sinh(b*n*log
(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) + 4)/(b*n*cosh(b*n*lo
g(x) + b*log(c) + a)^2 + 2*b*n*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n...
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx = \text{Timed out}$$

input

```
integrate(tanh(a+b*ln(c*x**n))**(5/2)/x,x)
```

output

```
Timed out
```

**Maxima [F]**

$$\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx = \int \frac{\tanh(b \log(cx^n) + a)^{\frac{5}{2}}}{x} dx$$

input `integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="maxima")`

output `integrate(tanh(b*log(c*x^n) + a)^(5/2)/x, x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx = \text{Timed out}$$

input `integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="giac")`

output `Timed out`

**Mupad [B] (verification not implemented)**

Time = 3.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.89

$$\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx = \frac{\operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{2 \tanh(a + b \ln(cx^n))^{3/2}}{3bn}$$

input `int(tanh(a + b*log(c*x^n))^(5/2)/x,x)`

output `atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - (2*tanh(a + b*log(c*x^n))^(3/2))/(3*b*n)`

### Reduce [F]

$$\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx = \int \frac{\sqrt{\tanh(\log(x^n c) b + a)} \tanh(\log(x^n c) b + a)^2}{x} dx$$

input `int(tanh(a+b*log(c*x^n))^(5/2)/x,x)`

output `int((sqrt(tanh(log(x**n*c)*b + a))*tanh(log(x**n*c)*b + a)**2)/x,x)`

### 3.205 $\int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx$

Optimal result	1516
Mathematica [A] (verified)	1516
Rubi [A] (verified)	1517
Maple [A] (verified)	1520
Fricas [B] (verification not implemented)	1520
Sympy [A] (verification not implemented)	1521
Maxima [F]	1522
Giac [F(-1)]	1522
Mupad [B] (verification not implemented)	1522
Reduce [F]	1523

#### Optimal result

Integrand size = 19, antiderivative size = 70

$$\int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx = \frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{bn}$$

output

$\arctan(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n+\operatorname{arctanh}(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n-2*\tanh(a+b*\ln(c*x^n))^{(1/2)}/b/n$

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.81

$$\int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx = \frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right) + \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right) - 2\sqrt{\tanh(a+b \log(cx^n))}}{bn}$$

input `Integrate[Tanh[a + b*Log[c*x^n]]^(3/2)/x,x]`

output `(ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]] + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]] - 2*Sqrt[Tanh[a + b*Log[c*x^n]]])/(b*n)`

### Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$ , Rules used = {3039, 3042, 3954, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n)) d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-i \tan(ia + ib \log(cx^n)))^{3/2} d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3954} \\
 & \int \frac{\frac{1}{\sqrt{\tanh(a+b \log(cx^n))}} d \log(cx^n) - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b}}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} + \int \frac{1}{\sqrt{-i \tan(ia+ib \log(cx^n))}} d \log(cx^n)}{n} \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int -\frac{1}{\sqrt{\tanh(a+b \log(cx^n))(1-\tanh^2(a+b \log(cx^n))}} d \tanh(a+b \log(cx^n))}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b}}{n}
 \end{aligned}$$

$$\begin{array}{c}
 \int \frac{\frac{1}{\sqrt{\tanh(a+b \log(cx^n))(1-\tanh^2(a+b \log(cx^n)))}} d \tanh(a+b \log(cx^n))}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n \\
 \downarrow 25 \\
 \frac{2 \int \frac{1}{1-\tanh^2(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))}}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n \\
 \downarrow 266 \\
 \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \int \frac{1}{\tanh(a+b \log(cx^n))+1} d \sqrt{\tanh(a+b \log(cx^n))}\right)}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n \\
 \downarrow 756 \\
 \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n \\
 \downarrow 216 \\
 \frac{2\left(\frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right) + \frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n \\
 \downarrow 219 \\
 \frac{2\left(\frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right) + \frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{b} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{b} \\
 \hline
 n
 \end{array}$$

input `Int [Tanh [a + b*Log [c*x^n]]^(3/2)/x, x]`

output `((2*(ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2 + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2))/b - (2*Sqrt[Tanh[a + b*Log[c*x^n]]])/b)/n`

**Defintions of rubi rules used**

rule 25 `Int [-(Fx_), x_Symbol] := Simp [Identity [-1] Int [Fx, x], x]`

rule 216 `Int [((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp [(1/(Rt [a, 2]*Rt [b, 2]))*ArcTan [Rt [b, 2]*(x/Rt [a, 2])], x] /; FreeQ [{a, b}, x] && PosQ [a/b] && (GtQ [a, 0] || GtQ [b, 0])`

rule 219  $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])] \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_ \cdot x)^m \cdot (a_ + (b_ \cdot x)^2)^{p_}, x\_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot x^{2k}/c^2)]^p, x], x, (c \cdot x)^{1/k}], x] /;$   $\text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 756  $\text{Int}[(a_ + (b_ \cdot x)^4)^{-1}, x\_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2 \cdot a) \ \text{Int}[1/(r - s \cdot x^2), x], x] + \text{Simp}[r/(2 \cdot a) \ \text{Int}[1/(r + s \cdot x^2), x], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ !\text{GtQ}[a/b, 0]$

rule 3039  $\text{Int}[u_, x\_Symbol] \rightarrow \text{With}\{lst = \text{FunctionOfLog}[\text{Cancel}[x \cdot u], x]\}, \text{Simp}[1/lst \ \text{Subst}[\text{Int}[lst[[1]], x], x, \text{Log}[lst[[2]]]], x] /;$   $!\text{FalseQ}[lst] /;$   $\text{NonsumQ}[u]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\text{Int}[(b_ \cdot \tan[(c_ \cdot x) + (d_ \cdot x)])^n, x\_Symbol] \rightarrow \text{Simp}[b \cdot ((b \cdot \tan[c + d \cdot x])^{n-1} / (d \cdot (n-1))), x] - \text{Simp}[b^2 \ \text{Int}[(b \cdot \tan[c + d \cdot x])^{n-2}, x], x] /;$   $\text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1]$

rule 3957  $\text{Int}[(b_ \cdot \tan[(c_ \cdot x) + (d_ \cdot x)])^n, x\_Symbol] \rightarrow \text{Simp}[b/d \ \text{Subst}[\text{Int}[x^n / (b^2 + x^2), x], x, b \cdot \tan[c + d \cdot x]], x] /;$   $\text{FreeQ}\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$



**Maple [A] (verified)**

Time = 1.17 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.06

method	result
derivativedivides	$\frac{-2\sqrt{\tanh(a+b\ln(cx^n))} - \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}+1)}{2} + \arctan(\sqrt{\tanh(a+b\ln(cx^n))})}{nb}$
default	$\frac{-2\sqrt{\tanh(a+b\ln(cx^n))} - \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}+1)}{2} + \arctan(\sqrt{\tanh(a+b\ln(cx^n))})}{nb}$

input `int(tanh(a+b*ln(c*x^n))^(3/2)/x,x,method=_RETURNVERBOSE)`output `1/n/b*(-2*tanh(a+b*ln(c*x^n))^(1/2)-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)+arctan(tanh(a+b*ln(c*x^n))^(1/2)))`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(64) = 128.

Time = 0.10 (sec) , antiderivative size = 334, normalized size of antiderivative = 4.77

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx =$$

$$\frac{4 \sqrt{\frac{\sinh(bn \log(x) + b \log(c) + a)}{\cosh(bn \log(x) + b \log(c) + a)}} - 2 \arctan\left(-\cosh(bn \log(x) + b \log(c) + a)^2 - 2 \cosh(bn \log(x) + b \log(c) + a)\right)}{nb}$$

input `integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="fricas")`

output

```
-1/2*(4*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) +
a)) - 2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*
log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) +
a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 +
1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)))
+ log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) +
a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 +
(cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sin
h(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt
(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))))/(b*n)
```

**Sympy [A] (verification not implemented)**

Time = 17.14 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.24

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx = -\frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} - 1\right)}{2bn} + \frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} + 1\right)}{2bn} - \frac{2\sqrt{\tanh(a + b \log(cx^n))}}{bn} + \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

input

```
integrate(tanh(a+b*ln(c*x**n))**(3/2)/x,x)
```

output

```
-log(sqrt(tanh(a + b*log(c*x**n))) - 1)/(2*b*n) + log(sqrt(tanh(a + b*log(
c*x**n))) + 1)/(2*b*n) - 2*sqrt(tanh(a + b*log(c*x**n)))/(b*n) + atan(sqrt
(tanh(a + b*log(c*x**n))))/(b*n)
```

**Maxima [F]**

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx = \int \frac{\tanh(b \log(cx^n) + a)^{\frac{3}{2}}}{x} dx$$

input `integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="maxima")`

output `integrate(tanh(b*log(c*x^n) + a)^(3/2)/x, x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx = \text{Timed out}$$

input `integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="giac")`

output `Timed out`

**Mupad [B] (verification not implemented)**

Time = 2.86 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.73

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx$$

$$= \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) + \operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) - 2\sqrt{\tanh(a + b \ln(cx^n))}}{bn}$$

input `int(tanh(a + b*log(c*x^n))^(3/2)/x,x)`

output `(atan(tanh(a + b*log(c*x^n))^(1/2)) + atanh(tanh(a + b*log(c*x^n))^(1/2)) - 2*tanh(a + b*log(c*x^n))^(1/2))/(b*n)`

**Reduce [F]**

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx = \frac{-2\sqrt{\tanh(\log(x^n c) b + a)} + \left( \int \frac{\sqrt{\tanh(\log(x^n c) b + a)}}{\tanh(\log(x^n c) b + a) x} dx \right) b n}{b n}$$

input `int(tanh(a+b*log(c*x^n))^(3/2)/x,x)`

output `( - 2*sqrt(tanh(log(x**n*c)*b + a)) + int(sqrt(tanh(log(x**n*c)*b + a))/(tanh(log(x**n*c)*b + a)*x),x)*b*n)/(b*n)`

**3.206**  $\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{x} dx$

Optimal result . . . . .	1524
Mathematica [A] (verified) . . . . .	1524
Rubi [A] (verified) . . . . .	1525
Maple [A] (verified) . . . . .	1527
Fricas [B] (verification not implemented) . . . . .	1528
Sympy [A] (verification not implemented) . . . . .	1528
Maxima [F] . . . . .	1529
Giac [F(-1)] . . . . .	1529
Mupad [B] (verification not implemented) . . . . .	1529
Reduce [F] . . . . .	1530

**Optimal result**

Integrand size = 19, antiderivative size = 48

$$\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{x} dx = -\frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn}$$

output `-arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n`

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{x} dx = -\frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right) - \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn}$$

input `Integrate[Sqrt[Tanh[a + b*Log[c*x^n]]]/x,x]`

output

$$-\left(\left(\text{ArcTan}\left[\text{Sqrt}\left[\text{Tanh}\left[a + b \cdot \text{Log}\left[cx^n\right]\right]\right]\right] - \text{ArcTanh}\left[\text{Sqrt}\left[\text{Tanh}\left[a + b \cdot \text{Log}\left[cx^n\right]\right]\right]\right]\right)\right)/(b \cdot n)$$
**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {3039, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx \\ & \quad \downarrow \text{3039} \\ & \int \frac{\sqrt{\tanh(a + b \log(cx^n))} d \log(cx^n)}{n} \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{-i \tan(ia + ib \log(cx^n))} d \log(cx^n)}{n} \\ & \quad \downarrow \text{3957} \\ & - \frac{\int - \frac{\sqrt{\tanh(a + b \log(cx^n))}}{1 - \tanh^2(a + b \log(cx^n))} d \tanh(a + b \log(cx^n))}{bn} \\ & \quad \downarrow \text{25} \\ & \frac{\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{1 - \tanh^2(a + b \log(cx^n))} d \tanh(a + b \log(cx^n))}{bn} \\ & \quad \downarrow \text{266} \\ & \frac{2 \int \frac{\tanh(a + b \log(cx^n))}{1 - \tanh^2(a + b \log(cx^n))} d \sqrt{\tanh(a + b \log(cx^n))}}{bn} \\ & \quad \downarrow \text{827} \\ & \frac{2 \left( \frac{1}{2} \int \frac{1}{1 - \tanh(a + b \log(cx^n))} d \sqrt{\tanh(a + b \log(cx^n))} - \frac{1}{2} \int \frac{1}{\tanh(a + b \log(cx^n)) + 1} d \sqrt{\tanh(a + b \log(cx^n))} \right)}{bn} \end{aligned}$$

$$\frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d\sqrt{\tanh(a+b \log(cx^n))} - \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{bn}$$

$$\frac{2\left(\frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right) - \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{bn}$$

input `Int[Sqrt[Tanh[a + b*Log[c*x^n]]]/x,x]`

output `(2*(-1/2*ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]] + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2))/(b*n)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst [[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst]] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 1.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

method	result	size
derivativedivides	$\frac{-\frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))-1})}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2}}{nb} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))})$	61
default	$\frac{-\frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))-1})}{2} + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2}}{nb} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))})$	61

input `int(tanh(a+b*ln(c*x^n))^(1/2)/x,x,method=_RETURNVERBOSE)`

output `1/n/b*(-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)-arctan(tanh(a+b*ln(c*x^n))^(1/2)))`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 303 vs.  $2(44) = 88$ .

Time = 0.12 (sec) , antiderivative size = 303, normalized size of antiderivative = 6.31

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx = \frac{2 \arctan\left(-\cosh(bn \log(x) + b \log(c) + a)^2 - 2 \cosh(bn \log(x) + b \log(c) + a) \sinh(bn \log(x) + b \log(c) + a)\right)}{bn}$$

input `integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="fricas")`

output `-1/2*(2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)))/(b*n)`

**Sympy [A] (verification not implemented)**

Time = 1.00 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.38

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx = -\frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} - 1\right)}{2bn} + \frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} + 1\right)}{2bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

input `integrate(tanh(a+b*ln(c*x**n))**(1/2)/x,x)`

output

```
-log(sqrt(tanh(a + b*log(c*x**n))) - 1)/(2*b*n) + log(sqrt(tanh(a + b*log(
c*x**n))) + 1)/(2*b*n) - atan(sqrt(tanh(a + b*log(c*x**n))))/(b*n)
```

**Maxima [F]**

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx = \int \frac{\sqrt{\tanh(b \log(cx^n) + a)}}{x} dx$$

input

```
integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="maxima")
```

output

```
integrate(sqrt(tanh(b*log(c*x^n) + a))/x, x)
```

**Giac [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx = \text{Timed out}$$

input

```
integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="giac")
```

output

```
Timed out
```

**Mupad [B] (verification not implemented)**

Time = 2.41 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.81

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx$$

$$= -\frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) - \operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn}$$

input

```
int(tanh(a + b*log(c*x^n))^(1/2)/x,x)
```

output `-(atan(tanh(a + b*log(c*x^n))^(1/2)) - atanh(tanh(a + b*log(c*x^n))^(1/2)))/(b*n)`

### Reduce [F]

$$\int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx = \int \frac{\sqrt{\tanh(\log(x^n c) b + a)}}{x} dx$$

input `int(tanh(a+b*log(c*x^n))^(1/2)/x,x)`

output `int(sqrt(tanh(log(x**n*c)*b + a))/x,x)`

**3.207**  $\int \frac{1}{x\sqrt{\tanh(a+b\log(cx^n))}} dx$

Optimal result	1531
Mathematica [A] (verified)	1531
Rubi [A] (verified)	1532
Maple [A] (verified)	1534
Fricas [B] (verification not implemented)	1535
Sympy [A] (verification not implemented)	1535
Maxima [F]	1536
Giac [F(-1)]	1536
Mupad [B] (verification not implemented)	1536
Reduce [F]	1537

**Optimal result**

Integrand size = 19, antiderivative size = 47

$$\int \frac{1}{x\sqrt{\tanh(a+b\log(cx^n))}} dx = \frac{\arctan\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn}$$

output

```
arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{\tanh(a+b\log(cx^n))}} dx = \frac{\arctan\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn}$$

input `Integrate[1/(x*Sqrt[Tanh[a + b*Log[c*x^n]]]),x]`

output `ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]/(b*n) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]/(b*n)]`

### Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {3039, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{1}{\sqrt{\tanh(a + b \log(cx^n))}} d \log(cx^n) \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-i \tan(ia + ib \log(cx^n))}} d \log(cx^n) \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int -\frac{1}{\sqrt{\tanh(a + b \log(cx^n))(1 - \tanh^2(a + b \log(cx^n)))}} d \tanh(a + b \log(cx^n))}{bn} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{1}{\sqrt{\tanh(a + b \log(cx^n))(1 - \tanh^2(a + b \log(cx^n)))}} d \tanh(a + b \log(cx^n))}{bn} \\
 & \quad \downarrow \text{266} \\
 & \frac{2 \int \frac{1}{1 - \tanh^2(a + b \log(cx^n))} d \sqrt{\tanh(a + b \log(cx^n))}}{bn} \\
 & \quad \downarrow \text{756}
 \end{aligned}$$

$$\begin{aligned}
& \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d\sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \int \frac{1}{\tanh(a+b \log(cx^n))+1} d\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} \\
& \quad \downarrow \text{216} \\
& \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d\sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{bn} \\
& \quad \downarrow \text{219} \\
& \frac{2\left(\frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right) + \frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{bn}
\end{aligned}$$

input `Int[1/(x*Sqrt[Tanh[a + b*Log[c*x^n]]]),x]`

output `(2*(ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]/2 + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2))/(b*n)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 3039 `Int[u_, x_Symbol] := With[{lst = FunctionOfLog[Cancel[x*u], x]}, Simp[1/lst [[3]] Subst[Int[lst[[1]], x], x, Log[lst[[2]]]], x] /; !FalseQ[lst]] /; NonsumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$	37
default	$\frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$	37

input `int(1/x/tanh(a+b*ln(c*x^n))^(1/2),x,method=_RETURNVERBOSE)`

output `1/n/b*(arctanh(tanh(a+b*ln(c*x^n))^(1/2))+arctan(tanh(a+b*ln(c*x^n))^(1/2)))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 305 vs.  $2(43) = 86$ .

Time = 0.11 (sec) , antiderivative size = 305, normalized size of antiderivative = 6.49

$$\int \frac{1}{x\sqrt{\tanh(a + b \log(cx^n))}} dx$$

$$= \frac{2 \arctan\left(-\cosh(bn \log(x) + b \log(c) + a)^2 - 2 \cosh(bn \log(x) + b \log(c) + a) \sinh(bn \log(x) + b \log(c) + a)\right)}{bn}$$

input `integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="fricas")`

output

```
1/2*(2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) - log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)))/(b*n)
```

**Sympy [A] (verification not implemented)**

Time = 1.91 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.40

$$\int \frac{1}{x\sqrt{\tanh(a + b \log(cx^n))}} dx = -\frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} - 1\right)}{2bn}$$

$$+ \frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} + 1\right)}{2bn}$$

$$+ \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

input `integrate(1/x/tanh(a+b*ln(c*x**n))**(1/2),x)`



output  $-\log(\sqrt{\tanh(a + b \log(cx^n))}) - 1)/(2bn) + \log(\sqrt{\tanh(a + b \log(cx^n))}) + 1)/(2bn) + \operatorname{atan}(\sqrt{\tanh(a + b \log(cx^n))})/(bn)$

### Maxima [F]

$$\int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx = \int \frac{1}{x \sqrt{\tanh(b \log(cx^n) + a)}} dx$$

input `integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="maxima")`

output `integrate(1/(x*sqrt(tanh(b*log(c*x^n) + a))), x)`

### Giac [F(-1)]

Timed out.

$$\int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx = \text{Timed out}$$

input `integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="giac")`

output `Timed out`

### Mupad [B] (verification not implemented)

Time = 2.57 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.77

$$\begin{aligned} & \int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx \\ &= \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) + \operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} \end{aligned}$$

input `int(1/(x*tanh(a + b*log(c*x^n))^(1/2)),x)`

output `(atan(tanh(a + b*log(c*x^n))^(1/2)) + atanh(tanh(a + b*log(c*x^n))^(1/2)))/(b*n)`

### Reduce [F]

$$\int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx = \int \frac{\sqrt{\tanh(\log(x^n c) b + a)}}{\tanh(\log(x^n c) b + a) x} dx$$

input `int(1/x/tanh(a+b*log(c*x^n))^(1/2),x)`

output `int(sqrt(tanh(log(x**n*c)*b + a))/(tanh(log(x**n*c)*b + a)*x),x)`

**3.208**  $\int \frac{1}{x \tanh^{\frac{3}{2}}(a+b \log(cx^n))} dx$

Optimal result	1538
Mathematica [A] (verified)	1538
Rubi [A] (verified)	1539
Maple [A] (verified)	1542
Fricas [B] (verification not implemented)	1542
Sympy [A] (verification not implemented)	1543
Maxima [F]	1544
Giac [F(-1)]	1544
Mupad [B] (verification not implemented)	1544
Reduce [F]	1545

**Optimal result**

Integrand size = 19, antiderivative size = 71

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a+b \log(cx^n))} dx = -\frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{2bn} - \frac{1}{bn\sqrt{\tanh(a+b \log(cx^n))}}$$

output

$$-\arctan(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n+\operatorname{arctanh}(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n-2/b/n/\tanh(a+b*\ln(c*x^n))^{(1/2)}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.37

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a+b \log(cx^n))} dx = \frac{-2 - \arctan\left(\sqrt[4]{\tanh^2(a+b \log(cx^n))}\right) \sqrt[4]{\tanh^2(a+b \log(cx^n))} + \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(a+b \log(cx^n))}\right)}{bn\sqrt{\tanh(a+b \log(cx^n))}}$$

input `Integrate[1/(x*Tanh[a + b*Log[c*x^n]]^(3/2)),x]`

output  $(-2 - \text{ArcTan}[(\text{Tanh}[a + b \cdot \text{Log}[c \cdot x^n]]^2)^{1/4}] \cdot (\text{Tanh}[a + b \cdot \text{Log}[c \cdot x^n]]^2)^{1/4} + \text{ArcTanh}[(\text{Tanh}[a + b \cdot \text{Log}[c \cdot x^n]]^2)^{1/4}] \cdot (\text{Tanh}[a + b \cdot \text{Log}[c \cdot x^n]]^2)^{1/4}) / (b \cdot n \cdot \text{Sqrt}[\text{Tanh}[a + b \cdot \text{Log}[c \cdot x^n]])]$

### Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$ , Rules used = {3039, 3042, 3955, 3042, 3957, 25, 266, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{1}{\tanh^{\frac{3}{2}}(a + b \log(cx^n))} d \log(cx^n) \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-i \tan(ia + ib \log(cx^n)))^{3/2}} d \log(cx^n) \\
 & \quad \downarrow \text{3955} \\
 & \int \sqrt{\tanh(a + b \log(cx^n))} d \log(cx^n) - \frac{2}{b \sqrt{\tanh(a + b \log(cx^n))}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{b \sqrt{\tanh(a + b \log(cx^n))}} + \int \sqrt{-i \tan(ia + ib \log(cx^n))} d \log(cx^n) \\
 & \quad \downarrow \text{3957} \\
 & -\frac{\int -\frac{\sqrt{\tanh(a + b \log(cx^n))}}{1 - \tanh^2(a + b \log(cx^n))} d \tanh(a + b \log(cx^n))}{b} - \frac{2}{b \sqrt{\tanh(a + b \log(cx^n))}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 25 \\
 \frac{\int \frac{\sqrt{\tanh(a+b \log(cx^n))}}{1-\tanh^2(a+b \log(cx^n))} d \tanh(a+b \log(cx^n))}{b} - \frac{2}{b\sqrt{\tanh(a+b \log(cx^n))}} \\
 \hline
 n \\
 \downarrow 266 \\
 \frac{2 \int \frac{\tanh(a+b \log(cx^n))}{1-\tanh^2(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))}}{b} - \frac{2}{b\sqrt{\tanh(a+b \log(cx^n))}} \\
 \hline
 n \\
 \downarrow 827 \\
 \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} - \frac{1}{2} \int \frac{1}{\tanh(a+b \log(cx^n))+1} d \sqrt{\tanh(a+b \log(cx^n))}\right)}{b} - \frac{2}{b\sqrt{\tanh(a+b \log(cx^n))}} \\
 \hline
 n \\
 \downarrow 216 \\
 \frac{2\left(\frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} - \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{b} - \frac{2}{b\sqrt{\tanh(a+b \log(cx^n))}} \\
 \hline
 n \\
 \downarrow 219 \\
 \frac{2\left(\frac{1}{2} \operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right) - \frac{1}{2} \arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)\right)}{b} - \frac{2}{b\sqrt{\tanh(a+b \log(cx^n))}} \\
 \hline
 n
 \end{array}$$

input `Int [1/(x*Tanh[a + b*Log[c*x^n]]^(3/2)), x]`

output `((2*(-1/2*ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]])/2)/b - 2/(b*Sqrt[Tanh[a + b*Log[c*x^n]]])/n`

## Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216  $\text{Int}[(\text{a}_) + (\text{b}_.) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 219  $\text{Int}[(\text{a}_) + (\text{b}_.) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 266  $\text{Int}[(\text{c}_.) * (\text{x}_)]^{(\text{m}_)} * ((\text{a}_) + (\text{b}_.) * (\text{x}_)^2)^{(\text{p}_)}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{k} * (\text{m} + 1) - 1)} * (\text{a} + \text{b} * (\text{x}^{(2 * \text{k})}/\text{c}^2))^{\text{p}}, \text{x}], \text{x}, (\text{c} * \text{x})^{(1/\text{k})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \&\& \text{FractionQ}[\text{m}] \&\& \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 827  $\text{Int}[(\text{x}_)^2 / ((\text{a}_) + (\text{b}_.) * (\text{x}_)^4), \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{s}/(2 * \text{b}) \quad \text{Int}[1/(\text{r} + \text{s} * \text{x}^2), \text{x}], \text{x}] - \text{Simp}[\text{s}/(2 * \text{b}) \quad \text{Int}[1/(\text{r} - \text{s} * \text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{!GtQ}[\text{a}/\text{b}, 0]$
- rule 3039  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{lst} = \text{FunctionOfLog}[\text{Cancel}[\text{x} * \text{u}], \text{x}]\}, \text{Simp}[1/\text{lst}[[3]] \quad \text{Subst}[\text{Int}[\text{lst}[[1]], \text{x}], \text{x}, \text{Log}[\text{lst}[[2]]]], \text{x}] /; \text{!FalseQ}[\text{lst}] /; \text{NonsumQ}[\text{u}]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3955  $\text{Int}[(\text{b}_.) * \tan[(\text{c}_.) + (\text{d}_.) * (\text{x}_)]]^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} + 1)} / (\text{b} * \text{d} * (\text{n} + 1)), \text{x}] - \text{Simp}[1/\text{b}^2 \quad \text{Int}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} + 2)}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[\text{n}, -1]$

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

**Maple [A] (verified)**

Time = 1.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$\frac{-\frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} - \frac{2}{\sqrt{\tanh(a+b\ln(cx^n))}} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))}) + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2}}{nb}$	76
default	$\frac{-\frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))}-1)}{2} - \frac{2}{\sqrt{\tanh(a+b\ln(cx^n))}} - \arctan(\sqrt{\tanh(a+b\ln(cx^n))}) + \frac{\ln(\sqrt{\tanh(a+b\ln(cx^n))+1})}{2}}{nb}$	76

input

```
int(1/x/tanh(a+b*ln(c*x^n))^(3/2), x, method=_RETURNVERBOSE)
```

output

```
1/n/b*(-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)-2/tanh(a+b*ln(c*x^n))^(1/2)-ar
ctan(tanh(a+b*ln(c*x^n))^(1/2))+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 625 vs. 2(65) = 130.

Time = 0.11 (sec) , antiderivative size = 625, normalized size of antiderivative = 8.80

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = \text{Too large to display}$$

input

```
integrate(1/x/tanh(a+b*log(c*x^n))^(3/2), x, algorithm="fricas")
```

output

```

-1/2*(2*(cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 -
1)*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^
2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)
*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*
sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + 4
*cosh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 +
2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b
*n*log(x) + b*log(c) + a)^2 - 1)*log(-cosh(b*n*log(x) + b*log(c) + a)^2 -
2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b
*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(
b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(
x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*lo
g(x) + b*log(c) + a))) + 8*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x)
+ b*log(c) + a) + 4*sinh(b*n*log(x) + b*log(c) + a)^2 + 4*(cosh(b*n*log(x)
) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) +
b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x)
) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) - 4)/(b*n*cosh(b*n*log(
x) + b*log(c) + a)^2 + 2*b*n*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*1...

```

### Sympy [A] (verification not implemented)

Time = 8.25 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.23

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = -\frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} - 1\right)}{2bn}$$

$$+ \frac{\log\left(\sqrt{\tanh(a + b \log(cx^n))} + 1\right)}{2bn}$$

$$- \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

$$- \frac{2}{bn\sqrt{\tanh(a + b \log(cx^n))}}$$

input

```
integrate(1/x/tanh(a+b*ln(c*x**n))**(3/2), x)
```



output

```
-log(sqrt(tanh(a + b*log(c*x**n))) - 1)/(2*b*n) + log(sqrt(tanh(a + b*log(
c*x**n))) + 1)/(2*b*n) - atan(sqrt(tanh(a + b*log(c*x**n))))/(b*n) - 2/(b*
n*sqrt(tanh(a + b*log(c*x**n))))
```

**Maxima [F]**

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = \int \frac{1}{x \tanh(b \log(cx^n) + a)^{\frac{3}{2}}} dx$$

input

```
integrate(1/x/tanh(a+b*log(c*x^n))^(3/2),x, algorithm="maxima")
```

output

```
integrate(1/(x*tanh(b*log(c*x^n) + a)^(3/2)), x)
```

**Giac [F(-1)]**

Timed out.

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = \text{Timed out}$$

input

```
integrate(1/x/tanh(a+b*log(c*x^n))^(3/2),x, algorithm="giac")
```

output

```
Timed out
```

**Mupad [B] (verification not implemented)**

Time = 2.73 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.92

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = \frac{\operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{\frac{bn}{2}} - \frac{1}{bn \sqrt{\tanh(a + b \ln(cx^n))}}$$

input `int(1/(x*tanh(a + b*log(c*x^n))^(3/2)),x)`

output `atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - 2/(b*n*tanh(a + b*log(c*x^n))^(1/2))`

### Reduce [F]

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx = \int \frac{\sqrt{\tanh(\log(x^n c) b + a)}}{\tanh(\log(x^n c) b + a)^2 x} dx$$

input `int(1/x/tanh(a+b*log(c*x^n))^(3/2),x)`

output `int(sqrt(tanh(log(x**n*c)*b + a))/(tanh(log(x**n*c)*b + a)**2*x),x)`

**3.209**  $\int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx$

Optimal result	1546
Mathematica [A] (verified)	1546
Rubi [A] (verified)	1547
Maple [A] (verified)	1550
Fricas [B] (verification not implemented)	1550
Sympy [F(-1)]	1551
Maxima [F]	1552
Giac [F(-1)]	1552
Mupad [B] (verification not implemented)	1552
Reduce [F]	1553

**Optimal result**

Integrand size = 19, antiderivative size = 72

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx = \frac{\arctan\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{2bn} - \frac{1}{3bn \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

output

```
arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n-2/3/b/n/tanh(a+b*ln(c*x^n))^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx = \frac{-2 + 3 \arctan\left(\sqrt[4]{\tanh^2(a+b \log(cx^n))}\right) \tanh^2(a+b \log(cx^n))^{3/4} + 3 \operatorname{arctanh}\left(\sqrt[4]{\tanh^2(a+b \log(cx^n))}\right)}{3bn \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

input `Integrate[1/(x*Tanh[a + b*Log[c*x^n]]^(5/2)),x]`

output `(-2 + 3*ArcTan[(Tanh[a + b*Log[c*x^n]]^2)^(1/4)]*(Tanh[a + b*Log[c*x^n]]^2)^(3/4) + 3*ArcTanh[(Tanh[a + b*Log[c*x^n]]^2)^(1/4)]*(Tanh[a + b*Log[c*x^n]]^2)^(3/4))/(3*b*n*Tanh[a + b*Log[c*x^n]]^(3/2))`

### Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.01, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$ , Rules used = {3039, 3042, 3955, 3042, 3957, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx \\
 & \quad \downarrow \text{3039} \\
 & \int \frac{1}{\tanh^{\frac{5}{2}}(a + b \log(cx^n))} d \log(cx^n) \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-i \tan(ia + ib \log(cx^n)))^{5/2}} d \log(cx^n) \\
 & \quad \downarrow \text{3955} \\
 & \int \frac{1}{\sqrt{\tanh(a + b \log(cx^n))}} d \log(cx^n) - \frac{2}{3b \tanh^{\frac{3}{2}}(a + b \log(cx^n))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2}{3b \tanh^{\frac{3}{2}}(a + b \log(cx^n))} + \int \frac{1}{\sqrt{-i \tan(ia + ib \log(cx^n))}} d \log(cx^n) \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int -\frac{1}{\sqrt{\tanh(a + b \log(cx^n))(1 - \tanh^2(a + b \log(cx^n)))}} d \tanh(a + b \log(cx^n))}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a + b \log(cx^n))} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\frac{\int \frac{1}{\sqrt{\tanh(a+b \log(cx^n))(1-\tanh^2(a+b \log(cx^n)))}} d \tanh(a+b \log(cx^n))}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

25

$$\frac{2 \int \frac{1}{1-\tanh^2(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))}}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

266

$$\frac{2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \int \frac{1}{\tanh(a+b \log(cx^n))+1} d \sqrt{\tanh(a+b \log(cx^n))} \right)}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

756

$$\frac{2 \left( \frac{1}{2} \int \frac{1}{1-\tanh(a+b \log(cx^n))} d \sqrt{\tanh(a+b \log(cx^n))} + \frac{1}{2} \arctan \left( \sqrt{\tanh(a+b \log(cx^n))} \right) \right)}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

216

$$\frac{2 \left( \frac{1}{2} \arctan \left( \sqrt{\tanh(a+b \log(cx^n))} \right) + \frac{1}{2} \operatorname{arctanh} \left( \sqrt{\tanh(a+b \log(cx^n))} \right) \right)}{b} - \frac{2}{3b \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

219

input `Int [1/(x*Tanh[a + b*Log[c*x^n]]^(5/2)), x]`

output `((2*(ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2 + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/2))/b - 2/(3*b*Tanh[a + b*Log[c*x^n]]^(3/2)))/n`

## Definitions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 216  $\text{Int}[\text{((a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[\text{b}, 2])) * \text{ArcTan}[\text{Rt}[\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{PosQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{GtQ}[\text{b}, 0])$
- rule 219  $\text{Int}[\text{((a}_) + (\text{b}_) * (\text{x}_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2] * \text{Rt}[-\text{b}, 2])) * \text{ArcTanh}[\text{Rt}[-\text{b}, 2] * (\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}] \&\& (\text{GtQ}[\text{a}, 0] \parallel \text{LtQ}[\text{b}, 0])$
- rule 266  $\text{Int}[\text{((c}_) * (\text{x}_))^{\text{m}} * \text{((a}_) + (\text{b}_) * (\text{x}_)^2)^{\text{p}}], \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[\text{x}^{\text{k} * (\text{m} + 1) - 1} * (\text{a} + \text{b} * (\text{x}^{2 * \text{k}}/\text{c}^2))^{\text{p}}, \text{x}], \text{x}, (\text{c} * \text{x})^{(1/\text{k})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \&\& \text{FractionQ}[\text{m}] \&\& \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 756  $\text{Int}[\text{((a}_) + (\text{b}_) * (\text{x}_)^4)^{-1}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[-\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[-\text{a}/\text{b}, 2]]\}, \text{Simp}[\text{r}/(2 * \text{a}) \quad \text{Int}[1/(\text{r} - \text{s} * \text{x}^2), \text{x}], \text{x}] + \text{Simp}[\text{r}/(2 * \text{a}) \quad \text{Int}[1/(\text{r} + \text{s} * \text{x}^2), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{!GtQ}[\text{a}/\text{b}, 0]$
- rule 3039  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{lst} = \text{FunctionOfLog}[\text{Cancel}[\text{x} * \text{u}], \text{x}]\}, \text{Simp}[1/\text{lst}[[3]] \quad \text{Subst}[\text{Int}[\text{lst}[[1]], \text{x}], \text{x}, \text{Log}[\text{lst}[[2]]]], \text{x}] /; \text{!FalseQ}[\text{lst}] /; \text{NonsumQ}[\text{u}]$
- rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3955  $\text{Int}[\text{((b}_) * \text{tan}[(\text{c}_) + (\text{d}_) * (\text{x}_)])^{\text{n}}], \text{x\_Symbol}] \rightarrow \text{Simp}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{\text{n} + 1} / (\text{b} * \text{d} * (\text{n} + 1)), \text{x}] - \text{Simp}[1/\text{b}^2 \quad \text{Int}[(\text{b} * \text{Tan}[\text{c} + \text{d} * \text{x}])^{\text{n} + 2}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[\text{n}, -1]$

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

**Maple [A] (verified)**

Time = 1.15 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{\frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) - \frac{2}{3\tanh(a+b\ln(cx^n))^{\frac{3}{2}}} - \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2}}{nb}$	74
default	$\frac{\frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) - \frac{2}{3\tanh(a+b\ln(cx^n))^{\frac{3}{2}}} - \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2}}{nb}$	74

input

```
int(1/x/tanh(a+b*ln(c*x^n))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
1/n/b*(1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)+arctan(tanh(a+b*ln(c*x^n))^(1/2
))-2/3/tanh(a+b*ln(c*x^n))^(3/2)-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1110 vs. 2(64) = 128.

Time = 0.15 (sec) , antiderivative size = 1110, normalized size of antiderivative = 15.42

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \text{Too large to display}$$

input

```
integrate(1/x/tanh(a+b*log(c*x^n))^(5/2), x, algorithm="fricas")
```

output

```
-1/6*(4*cosh(b*n*log(x) + b*log(c) + a)^4 + 16*cosh(b*n*log(x) + b*log(c)
+ a)*sinh(b*n*log(x) + b*log(c) + a)^3 + 4*sinh(b*n*log(x) + b*log(c) + a)
^4 + 8*(3*cosh(b*n*log(x) + b*log(c) + a)^2 - 1)*sinh(b*n*log(x) + b*log(c)
) + a)^2 - 6*(cosh(b*n*log(x) + b*log(c) + a)^4 + 4*cosh(b*n*log(x) + b*lo
g(c) + a)*sinh(b*n*log(x) + b*log(c) + a)^3 + sinh(b*n*log(x) + b*log(c) +
a)^4 + 2*(3*cosh(b*n*log(x) + b*log(c) + a)^2 - 1)*sinh(b*n*log(x) + b*lo
g(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)^2 + 4*(cosh(b*n*log(x) + b
*log(c) + a)^3 - cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(
c) + a) + 1)*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x)
+ b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log
(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*lo
g(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a
)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) +
a))) - 8*cosh(b*n*log(x) + b*log(c) + a)^2 + 3*(cosh(b*n*log(x) + b*log(c)
) + a)^4 + 4*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) +
a)^3 + sinh(b*n*log(x) + b*log(c) + a)^4 + 2*(3*cosh(b*n*log(x) + b*log(c)
+ a)^2 - 1)*sinh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log
(c) + a)^2 + 4*(cosh(b*n*log(x) + b*log(c) + a)^3 - cosh(b*n*log(x) + b*lo
g(c) + a))*sinh(b*n*log(x) + b*log(c) + a) + 1)*log(-cosh(b*n*log(x) + b*l
og(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*lo...
```

### Sympy [F(-1)]

Timed out.

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \text{Timed out}$$

input

```
integrate(1/x/tanh(a+b*ln(c*x**n))**(5/2), x)
```

output

```
Timed out
```



**Maxima [F]**

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \int \frac{1}{x \tanh(b \log(cx^n) + a)^{\frac{5}{2}}} dx$$

input `integrate(1/x/tanh(a+b*log(c*x^n))^(5/2),x, algorithm="maxima")`

output `integrate(1/(x*tanh(b*log(c*x^n) + a)^(5/2)), x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \text{Timed out}$$

input `integrate(1/x/tanh(a+b*log(c*x^n))^(5/2),x, algorithm="giac")`

output `Timed out`

**Mupad [B] (verification not implemented)**

Time = 3.43 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.89

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} + \frac{\operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{\frac{bn}{2}} - \frac{1}{3bn \tanh(a + b \ln(cx^n))^{3/2}}$$

input `int(1/(x*tanh(a + b*log(c*x^n))^(5/2)),x)`

output `atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) + atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - 2/(3*b*n*tanh(a + b*log(c*x^n))^(3/2))`

### Reduce [F]

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx = \int \frac{\sqrt{\tanh(\log(x^n c) b + a)}}{\tanh(\log(x^n c) b + a)^3 x} dx$$

input `int(1/x/tanh(a+b*log(c*x^n))^(5/2),x)`

output `int(sqrt(tanh(log(x**n*c)*b + a))/(tanh(log(x**n*c)*b + a)**3*x),x)`

**3.210**  $\int \frac{\tanh^5(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$

Optimal result	1554
Mathematica [A] (verified)	1555
Rubi [C] (warning: unable to verify)	1555
Maple [A] (verified)	1559
Fricas [B] (verification not implemented)	1559
Sympy [F]	1560
Maxima [F]	1560
Giac [F]	1560
Mupad [F(-1)]	1561
Reduce [F]	1561

**Optimal result**

Integrand size = 23, antiderivative size = 135

$$\int \frac{\tanh^5(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = \frac{(b-2c)\operatorname{arctanh}\left(\frac{b+2c \tanh^2(x)}{2\sqrt{c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{4c^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}{2c}$$

output

```
1/4*(b-2*c)*arctanh(1/2*(b+2*c*tanh(x)^2)/c^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(3/2)+1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)-1/2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)/c
```

**Mathematica [A] (verified)**

Time = 0.87 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.01

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \frac{1}{4} \left( \frac{(-b + 2c) \operatorname{arctanh} \left( \frac{-b - 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{c^{3/2}} \right. \\ \left. + \frac{2 \operatorname{arctanh} \left( \frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{\sqrt{a + b + c}} \right. \\ \left. - \frac{2\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{c} \right)$$

input `Integrate[Tanh[x]^5/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

output `(((-b + 2*c)*ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/c^(3/2) + (2*ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])])/Sqrt[a + b + c] - (2*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])/c)/4`

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.68, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {3042, 26, 4183, 1578, 1267, 27, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int -\frac{i \tan(ix)^5}{\sqrt{a - b \tan(ix)^2 + c \tan(ix)^4}} dx \\
& \downarrow 26 \\
& -i \int \frac{\tan(ix)^5}{\sqrt{c \tan(ix)^4 - b \tan(ix)^2 + a}} dx \\
& \downarrow 4183 \\
& - \int \frac{i \tanh^5(x)}{(1 - \tanh^2(x)) \sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} d(i \tanh(x)) \\
& \downarrow 1578 \\
& -\frac{1}{2} \int -\frac{\tanh^2(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \\
& \downarrow 1267 \\
& \frac{1}{2} \left( -\frac{\int \frac{b - (b - 2c) \tanh^2(x)}{2(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x))}{c} - \frac{\sqrt{a - ib \tanh(x) - c \tanh^2(x)}}{c} \right) \\
& \downarrow 27 \\
& \frac{1}{2} \left( -\frac{\int \frac{b - (b - 2c) \tanh^2(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x))}{2c} - \frac{\sqrt{a - ib \tanh(x) - c \tanh^2(x)}}{c} \right) \\
& \downarrow 1269 \\
& \frac{1}{2} \left( -\frac{(b - 2c) \int \frac{1}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) + 2c \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x))}{2c} \right) \\
& \downarrow 1092 \\
& \frac{1}{2} \left( -\frac{2(b - 2c) \int \frac{1}{\tanh^2(x) + 4c} d\left(-\frac{b - 2ic \tanh(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}}\right) + 2c \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x))}{2c} \right)
\end{aligned}$$

↓ 219

$$\frac{1}{2} \left( \frac{2c \int \frac{1}{(1-\tanh^2(x))\sqrt{-c \tanh^2(x)-ib \tanh(x)+a}} d(-\tanh^2(x)) + \frac{i(b-2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}}}{2c} - \frac{\sqrt{a-ib \tanh(x)-c \tanh^2(x)}}{c} \right)$$

↓ 1154

$$\frac{1}{2} \left( \frac{\frac{i(b-2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} - 4c \int \frac{1}{\tanh^2(x)+4(a+b+c)} d \frac{2a+b-i(b+2c) \tanh(x)}{\sqrt{-c \tanh^2(x)-ib \tanh(x)+a}}}{2c} - \frac{\sqrt{a-ib \tanh(x)-c \tanh^2(x)}}{c} \right)$$

↓ 219

$$\frac{1}{2} \left( \frac{\frac{i(b-2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} - \frac{2ic \arctan\left(\frac{\tanh(x)}{2\sqrt{a+b+c}}\right)}{\sqrt{a+b+c}}}{2c} - \frac{\sqrt{a-ib \tanh(x)-c \tanh^2(x)}}{c} \right)$$

input

```
Int [Tanh[x]^5/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]
```

output

```
(-1/2*((I*(b - 2*c)*ArcTan[Tanh[x]/(2*Sqrt[c])])/Sqrt[c] - ((2*I)*c*ArcTan
[Tanh[x]/(2*Sqrt[a + b + c])])/Sqrt[a + b + c])/c - Sqrt[a - I*b*Tanh[x] -
c*Tanh[x]^2]/c)/2
```

### Defintions of rubi rules used

rule 26

```
Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

- rule 219  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1092  $\text{Int}[1/\text{Sqrt}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)], x\_Symbol] \rightarrow \text{Simp}[2 \ \text{Subst}[\text{Int}[1/(4 \cdot c - x^2), x], x, (b + 2 \cdot c \cdot x)/\text{Sqrt}[a + b \cdot x + c \cdot x^2]], x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 1154  $\text{Int}[1/(((d_ ) + (e_ \cdot)(x_ )) \cdot \text{Sqrt}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)]), x\_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/(4 \cdot c \cdot d^2 - 4 \cdot b \cdot d \cdot e + 4 \cdot a \cdot e^2 - x^2), x], x, (2 \cdot a \cdot e - b \cdot d - (2 \cdot c \cdot d - b \cdot e) \cdot x)/\text{Sqrt}[a + b \cdot x + c \cdot x^2]], x] /; \text{FreeQ}\{a, b, c, d, e\}, x]$
- rule 1267  $\text{Int}(((d_ ) + (e_ \cdot)(x_ ))^{(m_ )} \cdot ((f_ ) + (g_ \cdot)(x_ ))^{(n_ )} \cdot ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{(p_ )}), x\_Symbol] \rightarrow \text{Simp}[g^n \cdot (d + e \cdot x)^{(m + n - 1)} \cdot ((a + b \cdot x + c \cdot x^2)^{(p + 1)} / (c \cdot e^{(n - 1)} \cdot (m + n + 2 \cdot p + 1))), x] + \text{Simp}[1 / (c \cdot e^{(n \cdot (m + n + 2 \cdot p + 1))}) \ \text{Int}[(d + e \cdot x)^m \cdot (a + b \cdot x + c \cdot x^2)^p \cdot \text{ExpandToSum}[c \cdot e^{(n \cdot (m + n + 2 \cdot p + 1))} \cdot (f + g \cdot x)^n - c \cdot g^n \cdot (m + n + 2 \cdot p + 1) \cdot (d + e \cdot x)^n - g^n \cdot (d + e \cdot x)^{(n - 2)} \cdot (b \cdot d \cdot e \cdot (p + 1) + a \cdot e^2 \cdot (m + n - 1) - c \cdot d^2 \cdot (m + n + 2 \cdot p + 1) - e \cdot (2 \cdot c \cdot d - b \cdot e) \cdot (m + n + p) \cdot x)], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, m, p\}, x \ \&\& \ \text{IGtQ}[n, 1] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{NeQ}[m + n + 2 \cdot p + 1, 0]$
- rule 1269  $\text{Int}(((d_ ) + (e_ \cdot)(x_ ))^{(m_ )} \cdot ((f_ ) + (g_ \cdot)(x_ )) \cdot ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{(p_ )}), x\_Symbol] \rightarrow \text{Simp}[g/e \ \text{Int}[(d + e \cdot x)^{(m + 1)} \cdot (a + b \cdot x + c \cdot x^2)^p, x], x] + \text{Simp}[(e \cdot f - d \cdot g)/e \ \text{Int}[(d + e \cdot x)^m \cdot (a + b \cdot x + c \cdot x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, m, p\}, x \ \&\& \ !\text{IGtQ}[m, 0]$
- rule 1578  $\text{Int}[(x_ )^{(m_ )} \cdot ((d_ ) + (e_ \cdot)(x_ )^2)^{(q_ )} \cdot ((a_ + (b_ \cdot)(x_ )^2 + (c_ \cdot)(x_ )^4)^{(p_ )}), x\_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{((m - 1)/2)} \cdot (d + e \cdot x)^q \cdot (a + b \cdot x + c \cdot x^2)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, c, d, e, p, q\}, x \ \&\& \ \text{IntegerQ}[(m - 1)/2]$
- rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4183

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n2_.))^(p_)), x_Symbol]
:> Simp[f/e Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

**Maple [A] (verified)**

Time = 2.35 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.10

method	result
derivativedivides	$-\frac{\ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{2\sqrt{c}} - \frac{\sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{4c^{\frac{3}{2}}}$
default	$-\frac{\ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{2\sqrt{c}} - \frac{\sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{4c^{\frac{3}{2}}}$

input

```
int(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-1/2*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)-1/2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)/c+1/4*b/c^(3/2)*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))+1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 2075 vs. 2(111) = 222.

Time = 1.15 (sec) , antiderivative size = 8891, normalized size of antiderivative = 65.86

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Too large to display}$$

input

```
integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")
```



output Too large to include

### Sympy [F]

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

input `integrate(tanh(x)**5/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2), x)`

output `Integral(tanh(x)**5/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)`

### Maxima [F]

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^5}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, algorithm="maxima")`

output `integrate(tanh(x)^5/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

### Giac [F]

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^5}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, algorithm="giac")`

output `integrate(tanh(x)^5/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^5}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `int(tanh(x)^5/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`

output `int(tanh(x)^5/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`

### Reduce [F]

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\sqrt{\tanh(x)^4 c + \tanh(x)^2 b + a} \tanh(x)^5}{\tanh(x)^4 c + \tanh(x)^2 b + a} dx$$

input `int(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x)**5)/(tanh(x)**4*c + tanh(x)**2*b + a),x)`

**3.211**  $\int \frac{\tanh^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$

Optimal result	1562
Mathematica [A] (verified)	1563
Rubi [C] (warning: unable to verify)	1563
Maple [A] (verified)	1566
Fricas [B] (verification not implemented)	1566
Sympy [F]	1567
Maxima [F]	1567
Giac [F]	1568
Mupad [F(-1)]	1568
Reduce [F]	1568

**Optimal result**

Integrand size = 23, antiderivative size = 105

$$\int \frac{\tanh^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{b+2c \tanh^2(x)}{2\sqrt{c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

```
output -1/2*arctanh(1/2*(b+2*c*tanh(x)^2)/c^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)+1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \frac{1}{2} \left( \frac{\operatorname{arctanh}\left(\frac{-b - 2c \tanh^2(x)}{2\sqrt{c}\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c}\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{\sqrt{a + b + c}} \right)$$

input `Integrate[Tanh[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

output `(ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/Sqrt[c] + ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/Sqrt[a + b + c])/2`

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.50, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$ , Rules used = {3042, 26, 4183, 1578, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

↓ 3042

$$\int \frac{i \tan(ix)^3}{\sqrt{a - b \tan(ix)^2 + c \tan(ix)^4}} dx$$

$$\begin{aligned}
& \downarrow 26 \\
& i \int \frac{\tan(ix)^3}{\sqrt{c \tan(ix)^4 - b \tan(ix)^2 + a}} dx \\
& \downarrow 4183 \\
& \int -\frac{i \tanh^3(x)}{(1 - \tanh^2(x)) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} d(i \tanh(x)) \\
& \downarrow 1578 \\
& \frac{1}{2} \int -\frac{\tanh^2(x)}{(i \tanh(x) + 1) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \\
& \downarrow 1269 \\
& \frac{1}{2} \left( \int \frac{1}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) - \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \\
& \downarrow 1092 \\
& \frac{1}{2} \left( 2 \int \frac{1}{\tanh^2(x) + 4c} d\left(-\frac{b - 2ic \tanh(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}}\right) - \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \\
& \downarrow 219 \\
& \frac{1}{2} \left( \frac{i \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} - \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \\
& \downarrow 1154 \\
& \frac{1}{2} \left( 2 \int \frac{1}{\tanh^2(x) + 4(a + b + c)} d\frac{2a + b - i(b + 2c) \tanh(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} + \frac{i \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} \right) \\
& \downarrow 219 \\
& \frac{1}{2} \left( \frac{i \arctan\left(\frac{\tanh(x)}{2\sqrt{a+b+c}}\right)}{\sqrt{a+b+c}} + \frac{i \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} \right)
\end{aligned}$$

input `Int [Tanh[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]`

output `((I*ArcTan[Tanh[x]/(2*Sqrt[c])))/Sqrt[c] + (I*ArcTan[Tanh[x]/(2*Sqrt[a + b + c])))/Sqrt[a + b + c])/2`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092 `Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Simp[2 Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x]`

rule 1154 `Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1269 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[g/e Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Simp[(e*f - d*g)/e Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && !IGtQ[m, 0]`

rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4183 `Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_.))^p, x_Symbol] := Simp[f/e Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n 2, 2*n] && NeQ[b^2 - 4*a*c, 0]`

### Maple [A] (verified)

Time = 2.13 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$-\frac{\ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{b \tanh(x)^2 + 2c \tanh(x)^2 + 2a + b}{2\sqrt{a+b+c} \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}}\right)}{2\sqrt{a+b+c}}$	90
default	$-\frac{\ln\left(\frac{\frac{b}{2} + c \tanh(x)^2}{\sqrt{c}} + \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}\right)}{2\sqrt{c}} + \frac{\operatorname{arctanh}\left(\frac{b \tanh(x)^2 + 2c \tanh(x)^2 + 2a + b}{2\sqrt{a+b+c} \sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}}\right)}{2\sqrt{a+b+c}}$	90

input `int(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)+1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1518 vs. 2(85) = 170.

Time = 0.84 (sec) , antiderivative size = 6663, normalized size of antiderivative = 63.46

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Too large to display}$$

input `integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output Too large to include

### Sympy [F]

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

input `integrate(tanh(x)**3/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)`

output `Integral(tanh(x)**3/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)`

### Maxima [F]

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)^3/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`



**Giac [F]**

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `integrate(tanh(x)^3/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `int(tanh(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`

output `int(tanh(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`

**Reduce [F]**

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\sqrt{\tanh(x)^4 c + \tanh(x)^2 b + a} \tanh(x)^3}{\tanh(x)^4 c + \tanh(x)^2 b + a} dx$$

input `int(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x)**3)/(tanh(x)**4*c + tanh(x)**2*b + a),x)`

**3.212** 
$$\int \frac{\tanh(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$$

Optimal result	1569
Mathematica [A] (verified)	1569
Rubi [C] (warning: unable to verify)	1570
Maple [A] (verified)	1572
Fricas [B] (verification not implemented)	1572
Sympy [F]	1573
Maxima [F]	1574
Giac [F(-1)]	1574
Mupad [F(-1)]	1574
Reduce [F]	1575

**Optimal result**

Integrand size = 21, antiderivative size = 58

$$\int \frac{\tanh(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

output

```
1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

input

```
Integrate[Tanh[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]
```

output

```
ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh
[x]^2 + c*Tanh[x]^4])]/(2*Sqrt[a + b + c])
```

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.50, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 26, 4183, 1576, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(ix)}{\sqrt{a - b \tan^2(ix) + c \tan^4(ix)}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(ix)}{\sqrt{c \tan^4(ix) - b \tan^2(ix) + a}} dx \\
 & \quad \downarrow \text{4183} \\
 & -\int \frac{i \tanh(x)}{(1 - \tanh^2(x)) \sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} d(i \tanh(x)) \\
 & \quad \downarrow \text{1576} \\
 & -\frac{1}{2} \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \\
 & \quad \downarrow \text{1154} \\
 & \int \frac{1}{4(a + b + c) + \tanh^2(x)} d \frac{2a - i(b + 2c) \tanh(x) + b}{\sqrt{a - ib \tanh(x) - c \tanh^2(x)}} \\
 & \quad \downarrow \text{219}
 \end{aligned}$$

$$\frac{i \arctan\left(\frac{\tanh(x)}{2\sqrt{a+b+c}}\right)}{2\sqrt{a+b+c}}$$

input `Int [Tanh[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]`

output `((I/2)*ArcTan[Tanh[x]/(2*Sqrt[a + b + c]))/Sqrt[a + b + c]`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1154 `Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Simp[-2 Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1576 `Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4183

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n2_.))^(p_), x_Symbol]
:> Simp[f/e Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

**Maple [A] (verified)**

Time = 2.15 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{b \tanh(x)^2 + 2c \tanh(x)^2 + 2a + b}{2\sqrt{a+b+c} \sqrt{a+b \tanh(x)^2 + c \tanh(x)^4}}\right)}{2\sqrt{a+b+c}}$	52
default	$\frac{\operatorname{arctanh}\left(\frac{b \tanh(x)^2 + 2c \tanh(x)^2 + 2a + b}{2\sqrt{a+b+c} \sqrt{a+b \tanh(x)^2 + c \tanh(x)^4}}\right)}{2\sqrt{a+b+c}}$	52

input

```
int(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2))/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 882 vs.  $2(48) = 96$ .

Time = 0.56 (sec) , antiderivative size = 1748, normalized size of antiderivative = 30.14

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Too large to display}$$

input

```
integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")
```

output

```
[1/4*log(((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^8 + 8*(a^2 + 2*a
*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(
a + b)*c + c^2)*sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*cosh(x)^6 + 4*(7*(a^
2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^2 + a^2 + a*b - b*c - c^2)*si
nh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^3 + 3*(a^2
+ a*b - b*c - c^2)*cosh(x))*sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + b)*c + 3
*c^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^4
+ 30*(a^2 + a*b - b*c - c^2)*cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a + b)*c + 3*c
^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^5 + 1
0*(a^2 + a*b - b*c - c^2)*cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2
)*cosh(x))*sinh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*cosh(x)^2 + 4*(7*(a^2 + 2
*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*cos
h(x)^4 + 3*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*cosh(x)^2 + a^2 + a*b - b
*c - c^2)*sinh(x)^2 + sqrt(2)*((a + b + c)*cosh(x)^4 + 4*(a + b + c)*cosh(
x)*sinh(x)^3 + (a + b + c)*sinh(x)^4 + 2*(a - c)*cosh(x)^2 + 2*(3*(a + b +
c)*cosh(x)^2 + a - c)*sinh(x)^2 + 4*((a + b + c)*cosh(x)^3 + (a - c)*cosh
(x))*sinh(x) + a + b + c)*sqrt(a + b + c)*sqrt(((a + b + c)*cosh(x)^4 + (a
+ b + c)*sinh(x)^4 + 4*(a - c)*cosh(x)^2 + 2*(3*(a + b + c)*cosh(x)^2 + 2
*a - 2*c)*sinh(x)^2 + 3*a - b + 3*c)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*
cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + a^2 + 2*a*b + ...
```

## Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

input

```
integrate(tanh(x)/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)
```

output

```
Integral(tanh(x)/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)
```

**Maxima [F]**

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)}{\sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} dx$$

input `integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(tanh(x)/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Timed out}$$

input `integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\tanh(x)}{\sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} dx$$

input `int(tanh(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`

output `int(tanh(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`

**Reduce [F]**

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\sqrt{\tanh(x)^4 c + \tanh(x)^2 b + a} \tanh(x)}{\tanh(x)^4 c + \tanh(x)^2 b + a} dx$$

input `int(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*c + tanh(x)**2*b + a),x)`



**3.213** 
$$\int \frac{\coth(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$$

Optimal result	1576
Mathematica [A] (verified)	1577
Rubi [C] (warning: unable to verify)	1577
Maple [F]	1579
Fricas [B] (verification not implemented)	1580
Sympy [F]	1580
Maxima [F]	1580
Giac [F(-1)]	1581
Mupad [F(-1)]	1581
Reduce [F]	1581

**Optimal result**

Integrand size = 21, antiderivative size = 106

$$\int \frac{\coth(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

output

`-1/2*arctanh(1/2*(2*a+b*tanh(x)^2)/a^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/a^(1/2)+1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)`

**Mathematica [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.03

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a}} - \frac{\operatorname{arctanh}\left(\frac{-2a-b-(b+2c) \tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

input `Integrate[Coth[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

output `-1/2*ArcTanh[(2*a + b*Tanh[x]^2)/(2*Sqrt[a]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])/Sqrt[a] - ArcTanh[(-2*a - b - (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/(2*Sqrt[a + b + c])]`

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 26, 4183, 1578, 1289, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

↓ 3042

$$\int \frac{i}{\tan(ix)\sqrt{a - b \tan^2(ix) + c \tan^4(ix)}} dx$$

↓ 26

$$\begin{aligned}
& i \int \frac{1}{\tan(ix) \sqrt{c \tan(ix)^4 - b \tan(ix)^2 + a}} dx \\
& \quad \downarrow \text{4183} \\
& \int -\frac{i \coth(x)}{(1 - \tanh^2(x)) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} d(i \tanh(x)) \\
& \quad \downarrow \text{1578} \\
& \frac{1}{2} \int -\frac{i \coth(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \\
& \quad \downarrow \text{1289} \\
& \frac{1}{2} \int \left( \frac{1}{(-i \tanh(x) - 1) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} - \frac{i \coth(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} \right) d(-\tanh^2(x)) \\
& \quad \downarrow \text{2009} \\
& \frac{1}{2} \left( \frac{\operatorname{arctanh}\left(\frac{2a - i(b+2c) \tanh(x) + b}{2\sqrt{a+b+c} \sqrt{a - ib \tanh(x) - c \tanh^2(x)}}\right)}{\sqrt{a+b+c}} - \frac{\operatorname{arctanh}\left(\frac{2a - ib \tanh(x)}{2\sqrt{a} \sqrt{a - ib \tanh(x) - c \tanh^2(x)}}\right)}{\sqrt{a}} \right)
\end{aligned}$$

input `Int[Coth[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]`

output `(-(ArcTanh[(2*a - I*b*Tanh[x])/(2*Sqrt[a]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2))]/Sqrt[a]) + ArcTanh[(2*a + b - I*(b + 2*c)*Tanh[x])/(2*Sqrt[a + b + c]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])]/Sqrt[a + b + c])/2`

### Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 1289 `Int[((d_.) + (e_.)*(x_)^(m_.))*((f_.) + (g_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && (IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0]))`

rule 1578 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.))*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4183 `Int[tan[(d_.) + (e_.)*(x_)]^(m_.))*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_.))^(p_), x_Symbol] := Simp[f/e Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]`

## Maple **[F]**

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}} dx$$

input `int(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1520 vs.  $2(86) = 172$ .

Time = 0.83 (sec) , antiderivative size = 6705, normalized size of antiderivative = 63.25

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Too large to display}$$

input `integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output Too large to include

**Sympy [F]**

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

input `integrate(coth(x)/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)`

output `Integral(coth(x)/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)`

**Maxima [F]**

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Timed out}$$

input `integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")`

output Timed out

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)}{\sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} dx$$

input `int(coth(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`

output `int(coth(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`

**Reduce [F]**

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\sqrt{\tanh^4(x) c + \tanh^2(x) b + a} \coth(x)}{\tanh^4(x) c + \tanh^2(x) b + a} dx$$

input `int(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*coth(x))/(tanh(x)**4*c + tanh(x)**2*b + a),x)`

**3.214** 
$$\int \frac{\coth^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx$$

Optimal result	1582
Mathematica [A] (verified)	1583
Rubi [C] (warning: unable to verify)	1583
Maple [F]	1585
Fricas [B] (verification not implemented)	1586
Sympy [F]	1586
Maxima [F]	1587
Giac [F(-1)]	1587
Mupad [F(-1)]	1587
Reduce [F]	1588

**Optimal result**

Integrand size = 23, antiderivative size = 142

$$\int \frac{\coth^3(x)}{\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} dx = -\frac{(2a-b)\operatorname{arctanh}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{4a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\coth^2(x)\sqrt{a+b \tanh^2(x)+c \tanh^4(x)}}{2a}$$

output

```
-1/4*(2*a-b)*arctanh(1/2*(2*a+b*tanh(x)^2)/a^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/a^(3/2)+1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)-1/2*coth(x)^2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)/a
```

**Mathematica [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = -\frac{(2a - b) \operatorname{arctanh}\left(\frac{2a + b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{4a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a + b + c}} - \frac{\coth^2(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2a}$$

input `Integrate[Coth[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

output `-1/4*((2*a - b)*ArcTanh[(2*a + b*Tanh[x]^2)/(2*Sqrt[a]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/a^(3/2) + ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/(2*Sqrt[a + b + c]) - (Coth[x]^2*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4))/(2*a)`

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.35, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {3042, 26, 4183, 1578, 1289, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

↓ 3042



$$\begin{aligned}
 & \int -\frac{i}{\tan(ix)^3 \sqrt{a - b \tan(ix)^2 + c \tan(ix)^4}} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{1}{\tan(ix)^3 \sqrt{c \tan(ix)^4 - b \tan(ix)^2 + a}} dx \\
 & \quad \downarrow \text{4183} \\
 & - \int \frac{i \coth^3(x)}{(1 - \tanh^2(x)) \sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} d(i \tanh(x)) \\
 & \quad \downarrow \text{1578} \\
 & -\frac{1}{2} \int -\frac{\coth^2(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \\
 & \quad \downarrow \text{1289} \\
 & -\frac{1}{2} \int \left( -\frac{\coth^2(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} + \frac{i \coth(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} + \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left( \frac{\operatorname{barctanh}\left(\frac{2a - ib \tanh(x)}{2\sqrt{a} \sqrt{a - ib \tanh(x) - c \tanh^2(x)}}\right)}{2a^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{2a - ib \tanh(x)}{2\sqrt{a} \sqrt{a - ib \tanh(x) - c \tanh^2(x)}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{2a - i(b+2c) \tanh(x)}{2\sqrt{a+b+c} \sqrt{a - ib \tanh(x) - c \tanh^2(x)}}\right)}{\sqrt{a+b+c}} \right)
 \end{aligned}$$

input `Int[Coth[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]`

output `(-(ArcTanh[(2*a - I*b*Tanh[x])/(2*Sqrt[a]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])]/Sqrt[a]) + (b*ArcTanh[(2*a - I*b*Tanh[x])/(2*Sqrt[a]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])])/(2*a^(3/2)) + ArcTanh[(2*a + b - I*(b + 2*c)*Tanh[x])/(2*Sqrt[a + b + c]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])]/Sqrt[a + b + c] - (I*Coth[x]*Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])/a)/2`

## Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 1289 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && (IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0]))`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_))*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4183 `Int[tan[(d_) + (e_)*(x_)]^(m_))*((a_) + (b_))*((f_)*tan[(d_) + (e_)*(x_)]^(n_) + (c_))*((f_)*tan[(d_) + (e_)*(x_)]^(n2_))^(p_), x_Symbol] := Simp[f/e Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]`

## Maple [F]

$$\int \frac{\coth(x)^3}{\sqrt{a + b \tanh(x)^2 + c \tanh(x)^4}} dx$$

input `int(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2136 vs. 2(118) = 236.

Time = 1.10 (sec) , antiderivative size = 9168, normalized size of antiderivative = 64.56

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Too large to display}$$

input `integrate(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output `Too large to include`

### Sympy [F]

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

input `integrate(coth(x)**3/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)`

output `Integral(coth(x)**3/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)`

**Maxima [F]**

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `integrate(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(coth(x)^3/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \text{Timed out}$$

input `integrate(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

input `int(coth(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`

output `int(coth(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`

**Reduce [F]**

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx = \int \frac{\coth(x)^3}{\sqrt{\tanh(x)^4 c + \tanh(x)^2 b + a}} dx$$

input `int(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output `int(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

### 3.215 $\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$

Optimal result	1589
Mathematica [A] (verified)	1590
Rubi [C] (warning: unable to verify)	1590
Maple [A] (verified)	1594
Fricas [B] (verification not implemented)	1594
Sympy [F]	1595
Maxima [F]	1595
Giac [F]	1595
Mupad [F(-1)]	1596
Reduce [F]	1596

#### Optimal result

Integrand size = 21, antiderivative size = 132

$$\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$$

$$= -\frac{(b + 2c) \operatorname{arctanh}\left(\frac{b + 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{4\sqrt{c}}$$

$$+ \frac{1}{2} \sqrt{a + b + c} \operatorname{arctanh}\left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)$$

$$- \frac{1}{2} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}$$

output

```
-1/4*(b+2*c)*arctanh(1/2*(b+2*c*tanh(x)^2)/c^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)+1/2*(a+b+c)^(1/2)*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))-1/2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.99

$$\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$$

$$= \frac{1}{4} \left( \frac{(b + 2c) \operatorname{arctanh} \left( \frac{-b - 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{\sqrt{c}} \right. \\ \left. + 2\sqrt{a + b + c} \operatorname{arctanh} \left( \frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right) \right. \\ \left. - 2\sqrt{a + b \tanh^2(x) + c \tanh^4(x)} \right)$$

input

```
Integrate[Tanh[x]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]
```

output

```
((b + 2*c)*ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/Sqrt[c] + 2*Sqrt[a + b + c]*ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])] - 2*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])/4
```

**Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.64, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.524$ , Rules used = {3042, 26, 4183, 1576, 1162, 25, 1269, 1092, 219, 1154, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\
& \quad \downarrow \text{3042} \\
& \int -i \tan(ix) \sqrt{a - b \tan^2(ix) + c \tan^4(ix)} dx \\
& \quad \downarrow \text{26} \\
& -i \int \tan(ix) \sqrt{c \tan^4(ix) - b \tan^2(ix) + a} dx \\
& \quad \downarrow \text{4183} \\
& - \int \frac{i \tanh(x) \sqrt{c \tanh^4(x) + b \tanh^2(x) + a}}{1 - \tanh^2(x)} d(i \tanh(x)) \\
& \quad \downarrow \text{1576} \\
& -\frac{1}{2} \int \frac{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}}{1 - \tanh^2(x)} d(-\tanh^2(x)) \\
& \quad \downarrow \text{1162} \\
& \frac{1}{2} \left( \frac{1}{2} \int -\frac{2a + b - i(b + 2c) \tanh(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) - \sqrt{a - ib \tanh(x) - c \tanh^2(x)} \right) \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left( -\frac{1}{2} \int \frac{2a + b - i(b + 2c) \tanh(x)}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) - \sqrt{a - ib \tanh(x) - c \tanh^2(x)} \right) \\
& \quad \downarrow \text{1269} \\
& \frac{1}{2} \left( \frac{1}{2} \left( (b + 2c) \int \frac{1}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) - 2(a + b + c) \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \right. \\
& \quad \downarrow \text{1092} \\
& \left. \frac{1}{2} \left( \frac{1}{2} \left( 2(b + 2c) \int \frac{1}{\tanh^2(x) + 4c} d\left( -\frac{b - 2ic \tanh(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} \right) - 2(a + b + c) \int \frac{1}{(1 - \tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \right) \right)
\end{aligned}$$



↓ 219

$$\frac{1}{2} \left( \frac{1}{2} \left( \frac{i(b+2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} - 2(a+b+c) \int \frac{1}{(1-\tanh^2(x)) \sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} d(-\tanh^2(x)) \right) \right)$$

↓ 1154

$$\frac{1}{2} \left( \frac{1}{2} \left( 4(a+b+c) \int \frac{1}{\tanh^2(x) + 4(a+b+c)} d \frac{2a+b-i(b+2c)\tanh(x)}{\sqrt{-c \tanh^2(x) - ib \tanh(x) + a}} + \frac{i(b+2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} \right) \right)$$

↓ 219

$$\frac{1}{2} \left( \frac{1}{2} \left( 2i\sqrt{a+b+c} \arctan\left(\frac{\tanh(x)}{2\sqrt{a+b+c}}\right) + \frac{i(b+2c) \arctan\left(\frac{\tanh(x)}{2\sqrt{c}}\right)}{\sqrt{c}} \right) - \sqrt{a - ib \tanh(x) - c \tanh^2(x)} \right)$$

input `Int [Tanh[x]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4],x]`

output `((((I*(b + 2*c)*ArcTan[Tanh[x]/(2*Sqrt[c])))/Sqrt[c] + (2*I)*Sqrt[a + b + c]*ArcTan[Tanh[x]/(2*Sqrt[a + b + c]))]/2 - Sqrt[a - I*b*Tanh[x] - c*Tanh[x]^2])/2`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1092  $\text{Int}[1/\text{Sqrt}[(a\_)+(b\_)(x\_)+(c\_)(x\_)^2], x\_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1154  $\text{Int}[1/(((d\_)+(e\_)(x\_))\text{Sqrt}[(a\_)+(b\_)(x\_)+(c\_)(x\_)^2]), x\_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1162  $\text{Int}(((d\_)+(e\_)(x\_))^{(m\_)}*((a\_)+(b\_)(x\_)+(c\_)(x\_)^2)^{(p\_)}, x\_Symbol] \rightarrow \text{Simp}[(d + e*x)^{(m+1)}*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - \text{Simp}[p/(e*(m + 2*p + 1)) \text{ Int}[(d + e*x)^m*\text{Simp}[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^{(p-1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&\& \text{GtQ}[p, 0] \&\& \text{NeQ}[m + 2*p + 1, 0] \&\& (!\text{RationalQ}[m] || \text{LtQ}[m, 1]) \&\& !\text{LtQ}[m + 2*p, 0] \&\& \text{IntQuadraticQ}[a, b, c, d, e, m, p, x]$

rule 1269  $\text{Int}(((d\_)+(e\_)(x\_))^{(m\_)}*((f\_)+(g\_)(x\_))*((a\_)+(b\_)(x\_)+(c\_)(x\_)^2)^{(p\_)}, x\_Symbol] \rightarrow \text{Simp}[g/e \text{ Int}[(d + e*x)^{(m+1)}*(a + b*x + c*x^2)^p, x], x] + \text{Simp}[(e*f - d*g)/e \text{ Int}[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m, p\}, x] \&\& !\text{IGtQ}[m, 0]$

rule 1576  $\text{Int}[(x\_)*((d\_)+(e\_)(x\_)^2)^{(q\_)}*((a\_)+(b\_)(x\_)^2+(c\_)(x\_)^4)^{(p\_)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, e, p, q\}, x]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4183  $\text{Int}[\tan[(d\_)+(e\_)(x\_)]^{(m\_)}*((a\_)+(b\_)((f\_)\tan[(d\_)+(e\_)(x\_)]^{(n\_)}+(c\_)((f\_)\tan[(d\_)+(e\_)(x\_)]^{(n2\_)}))^{(p\_)}, x\_Symbol] \rightarrow \text{Simp}[f/e \text{ Subst}[\text{Int}[(x/f)^m*((a + b*x^n + c*x^{(2*n)})^p/(f^2 + x^2)), x], x, f*\text{Tan}[d + e*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{EqQ}[n^2, 2*n] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

**Maple [A] (verified)**

Time = 2.22 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.25

method	result
derivativedivides	$-\frac{\sqrt{(\tanh(x)^2-1)^2 c+(b+2c)(\tanh(x)^2-1)+a+b+c}}{2} - \frac{(b+2c) \ln\left(\frac{\frac{b}{2}+c+c(\tanh(x)^2-1)}{\sqrt{c}} + \sqrt{(\tanh(x)^2-1)^2 c+(b+2c)(\tanh(x)^2-1)+a+b+c}\right)}{4\sqrt{c}}$
default	$-\frac{\sqrt{(\tanh(x)^2-1)^2 c+(b+2c)(\tanh(x)^2-1)+a+b+c}}{2} - \frac{(b+2c) \ln\left(\frac{\frac{b}{2}+c+c(\tanh(x)^2-1)}{\sqrt{c}} + \sqrt{(\tanh(x)^2-1)^2 c+(b+2c)(\tanh(x)^2-1)+a+b+c}\right)}{4\sqrt{c}}$

input `int(tanh(x)*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/2*((\tanh(x)^2-1)^2*c+(b+2*c)*(\tanh(x)^2-1)+a+b+c)^(1/2)-1/4*(b+2*c)*\ln\left(\frac{(1/2*b+c+c*(\tanh(x)^2-1))/c^(1/2)+((\tanh(x)^2-1)^2*c+(b+2*c)*(\tanh(x)^2-1)+a+b+c)^(1/2))/c^(1/2)+1/2*(a+b+c)^(1/2)*\ln((2*a+2*b+2*c+(b+2*c)*(\tanh(x)^2-1)+2*(a+b+c)^(1/2)*((\tanh(x)^2-1)^2*c+(b+2*c)*(\tanh(x)^2-1)+a+b+c)^(1/2))/(\tanh(x)^2-1))\right)$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1827 vs. 2(108) = 216.

Time = 1.35 (sec) , antiderivative size = 7896, normalized size of antiderivative = 59.82

$$\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx = \text{Too large to display}$$

input `integrate(tanh(x)*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")`

output Too large to include

**Sympy [F]**

$$\begin{aligned} & \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\ &= \int \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} \tanh(x) dx \end{aligned}$$

input `integrate(tanh(x)*(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)`

output `Integral(sqrt(a + b*tanh(x)**2 + c*tanh(x)**4)*tanh(x), x)`

**Maxima [F]**

$$\begin{aligned} & \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\ &= \int \sqrt{c \tanh^4(x) + b \tanh^2(x) + a} \tanh(x) dx \end{aligned}$$

input `integrate(tanh(x)*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a)*tanh(x), x)`

**Giac [F]**

$$\begin{aligned} & \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\ &= \int \sqrt{c \tanh^4(x) + b \tanh^2(x) + a} \tanh(x) dx \end{aligned}$$

input `integrate(tanh(x)*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a)*tanh(x), x)`

**Mupad [F(-1)]**

Timed out.

$$\begin{aligned} & \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\ &= \int \tanh(x) \sqrt{c \tanh^4(x) + b \tanh^2(x) + a} dx \end{aligned}$$

input `int(tanh(x)*(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)`output `int(tanh(x)*(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)`**Reduce [F]**

$$\begin{aligned} & \int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx \\ &= \frac{-\sqrt{\tanh^4(x) c + \tanh^2(x) b + a} b - \left( \int \frac{\sqrt{\tanh^4(x) c + \tanh^2(x) b + a} \tanh^5(x)}{\tanh^4(x) b c - 2 \tanh^4(x) c^2 + \tanh^2(x) b^2 - 2 \tanh^2(x) b c + a b - 2 a c} dx \right) b^2 c + 4 \left( \int \frac{\tanh^5(x)}{\tanh^4(x) b c - 2 \tanh^4(x) c^2 + \tanh^2(x) b^2 - 2 \tanh^2(x) b c + a b - 2 a c} dx \right) b^2 c}{\dots} \end{aligned}$$

input `int(tanh(x)*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x)`

output

```
( - sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*b - int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x)**5)/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*b**2*c + 4*int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x)**5)/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*c**3 + int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*a*b**2 - 4*int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*a*b*c + 4*int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*a*c**2 + int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*b**3 - 2*int((sqrt(tanh(x)**4*c + tanh(x)**2*b + a)*tanh(x))/(tanh(x)**4*b*c - 2*tanh(x)**4*c**2 + tanh(x)**2*b**2 - 2*tanh(x)**2*b*c + a*b - 2*a*c),x)*b**2*c)/(b - 2*c)
```

### 3.216 $\int e^{a+bx} \tanh^4(a + bx) dx$

Optimal result	1598
Mathematica [A] (verified)	1598
Rubi [A] (verified)	1599
Maple [C] (verified)	1600
Fricas [B] (verification not implemented)	1601
Sympy [F]	1601
Maxima [A] (verification not implemented)	1602
Giac [A] (verification not implemented)	1602
Mupad [B] (verification not implemented)	1603
Reduce [B] (verification not implemented)	1603

#### Optimal result

Integrand size = 16, antiderivative size = 107

$$\int e^{a+bx} \tanh^4(a + bx) dx = \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1 + e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1 + e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1 + e^{2a+2bx})} - \frac{3 \arctan(e^{a+bx})}{b}$$

output

$\exp(b*x+a)/b+8/3*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))^3-14/3*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))^2+5*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))-3*\arctan(\exp(b*x+a))/b$

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.71

$$\int e^{a+bx} \tanh^4(a + bx) dx = \frac{e^{a+bx}(12 + 25e^{2(a+bx)} + 24e^{4(a+bx)} + 3e^{6(a+bx)})}{3b(1 + e^{2(a+bx)})^3} - \frac{3 \arctan(e^{a+bx})}{b}$$

input

$\text{Integrate}[E^{(a + b*x)}*\text{Tanh}[a + b*x]^4,x]$

output

$$\frac{(E^{(a + b*x)}*(12 + 25*E^{(2*(a + b*x))} + 24*E^{(4*(a + b*x))} + 3*E^{(6*(a + b*x))}))/((3*b*(1 + E^{(2*(a + b*x)))})^3) - (3*ArcTan[E^{(a + b*x)}])/b}$$

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {2720, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{a+bx} \tanh^4(a+bx) dx \\ & \quad \downarrow 2720 \\ & \frac{\int \frac{(1-e^{2a+2bx})^4}{(1+e^{2a+2bx})^4} de^{a+bx}}{b} \\ & \quad \downarrow 300 \\ & \frac{\int \left( 1 - \frac{8e^{2a+2bx}(1+e^{4a+4bx})}{(1+e^{2a+2bx})^4} \right) de^{a+bx}}{b} \\ & \quad \downarrow 2009 \\ & \frac{-3 \arctan(e^{a+bx}) + e^{a+bx} + \frac{5e^{a+bx}}{e^{2a+2bx}+1} - \frac{14e^{a+bx}}{3(e^{2a+2bx}+1)^2} + \frac{8e^{a+bx}}{3(e^{2a+2bx}+1)^3}}{b} \end{aligned}$$

input

$$\text{Int}[E^{(a + b*x)}*Tanh[a + b*x]^4,x]$$

output

$$\frac{(E^{(a + b*x)} + (8*E^{(a + b*x)})/(3*(1 + E^{(2*a + 2*b*x))})^3) - (14*E^{(a + b*x)})/(3*(1 + E^{(2*a + 2*b*x))})^2 + (5*E^{(a + b*x)})/(1 + E^{(2*a + 2*b*x)}) - 3*ArcTan[E^{(a + b*x)}])/b}$$



**Defintions of rubi rules used**

```
rule 300 Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int
[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c
, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.86

method	result
risch	$\frac{e^{bx+a}}{b} + \frac{e^{bx+a}(15e^{4bx+4a}+16e^{2bx+2a}+9)}{3b(1+e^{2bx+2a})^3} + \frac{3i \ln(e^{bx+a-i})}{2b} - \frac{3i \ln(e^{bx+a+i})}{2b}$
derivativedivides	$\frac{\frac{\sinh(bx+a)^4}{\cosh(bx+a)^3} + \frac{4 \sinh(bx+a)^2}{\cosh(bx+a)^3} + \frac{8}{3 \cosh(bx+a)^3} + \frac{\sinh(bx+a)^3}{\cosh(bx+a)^2} + \frac{3 \sinh(bx+a)}{\cosh(bx+a)^2} - \frac{3 \operatorname{sech}(bx+a) \tanh(bx+a)}{2} - 3 \arctan(e^{bx+a})}{b}$
default	$\frac{\sinh(bx+a)^4}{\cosh(bx+a)^3} + \frac{4 \sinh(bx+a)^2}{\cosh(bx+a)^3} + \frac{8}{3 \cosh(bx+a)^3} + \frac{\sinh(bx+a)^3}{\cosh(bx+a)^2} + \frac{3 \sinh(bx+a)}{\cosh(bx+a)^2} - \frac{3 \operatorname{sech}(bx+a) \tanh(bx+a)}{2} - 3 \arctan(e^{bx+a})}{b}$

```
input int(exp(b*x+a)*tanh(b*x+a)^4,x,method=_RETURNVERBOSE)
```

```
output exp(b*x+a)/b+1/3*exp(b*x+a)*(15*exp(4*b*x+4*a)+16*exp(2*b*x+2*a)+9)/b/(1+e
xp(2*b*x+2*a))^3+3/2*I/b*ln(exp(b*x+a)-I)-3/2*I/b*ln(exp(b*x+a)+I)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 604 vs.  $2(95) = 190$ .

Time = 0.10 (sec) , antiderivative size = 604, normalized size of antiderivative = 5.64

$$\int e^{a+bx} \tanh^4(a+bx) dx = \text{Too large to display}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="fricas")`

output

$$\begin{aligned} & 1/3*(3*\cosh(b*x + a)^7 + 21*\cosh(b*x + a)*\sinh(b*x + a)^6 + 3*\sinh(b*x + a) \\ & )^7 + 3*(21*\cosh(b*x + a)^2 + 8)*\sinh(b*x + a)^5 + 24*\cosh(b*x + a)^5 + 15 \\ & *(7*\cosh(b*x + a)^3 + 8*\cosh(b*x + a))*\sinh(b*x + a)^4 + 5*(21*\cosh(b*x + \\ & a)^4 + 48*\cosh(b*x + a)^2 + 5)*\sinh(b*x + a)^3 + 25*\cosh(b*x + a)^3 + 3*(2 \\ & 1*\cosh(b*x + a)^5 + 80*\cosh(b*x + a)^3 + 25*\cosh(b*x + a))*\sinh(b*x + a)^2 \\ & - 9*(\cosh(b*x + a)^6 + 6*\cosh(b*x + a)*\sinh(b*x + a)^5 + \sinh(b*x + a)^6 \\ & + 3*(5*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^4 + 3*\cosh(b*x + a)^4 + 4*(5*\cos \\ & h(b*x + a)^3 + 3*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*(5*\cosh(b*x + a)^4 + 6 \\ & *\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^2 + 3*\cosh(b*x + a)^2 + 6*(\cosh(b*x + \\ & a)^5 + 2*\cosh(b*x + a)^3 + \cosh(b*x + a))*\sinh(b*x + a) + 1)*\arctan(\cosh(b \\ & *x + a) + \sinh(b*x + a)) + 3*(7*\cosh(b*x + a)^6 + 40*\cosh(b*x + a)^4 + 25* \\ & \cosh(b*x + a)^2 + 4)*\sinh(b*x + a) + 12*\cosh(b*x + a))/(b*\cosh(b*x + a)^6 \\ & + 6*b*\cosh(b*x + a)*\sinh(b*x + a)^5 + b*\sinh(b*x + a)^6 + 3*b*\cosh(b*x + a) \\ & )^4 + 3*(5*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^4 + 4*(5*b*\cosh(b*x + a)^3 \\ & + 3*b*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*b*\cosh(b*x + a)^2 + 3*(5*b*\cosh( \\ & b*x + a)^4 + 6*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^2 + 6*(b*\cosh(b*x + a) \\ & ^5 + 2*b*\cosh(b*x + a)^3 + b*\cosh(b*x + a))*\sinh(b*x + a) + b) \end{aligned}$$
**Sympy [F]**

$$\int e^{a+bx} \tanh^4(a+bx) dx = e^a \int e^{bx} \tanh^4(a+bx) dx$$

input `integrate(exp(b*x+a)*tanh(b*x+a)**4,x)`

output `exp(a)*Integral(exp(b*x)*tanh(a + b*x)**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.88

$$\int e^{a+bx} \tanh^4(a+bx) dx = -\frac{3 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b} + \frac{15 e^{(5bx+5a)} + 16 e^{(3bx+3a)} + 9 e^{(bx+a)}}{3b(e^{(6bx+6a)} + 3e^{(4bx+4a)} + 3e^{(2bx+2a)} + 1)}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="maxima")`

output `-3*arctan(e^(b*x + a))/b + e^(b*x + a)/b + 1/3*(15*e^(5*b*x + 5*a) + 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(b*(e^(6*b*x + 6*a) + 3*e^(4*b*x + 4*a) + 3*e^(2*b*x + 2*a) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.64

$$\int e^{a+bx} \tanh^4(a+bx) dx = \frac{15 e^{(5bx+5a)} + 16 e^{(3bx+3a)} + 9 e^{(bx+a)}}{(e^{(2bx+2a)} + 1)^3} - \frac{9 \arctan(e^{(bx+a)}) + 3 e^{(bx+a)}}{3b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="giac")`

output `1/3*((15*e^(5*b*x + 5*a) + 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(e^(2*b*x + 2*a) + 1)^3 - 9*arctan(e^(b*x + a)) + 3*e^(b*x + a))/b`

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.45

$$\int e^{a+bx} \tanh^4(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} + \frac{\frac{4e^{a+bx}}{3b} + \frac{4e^{5a+5bx}}{3b}}{3e^{2a+2bx} + 3e^{4a+4bx} + e^{6a+6bx} + 1} - \frac{2e^{a+bx}}{b(2e^{2a+2bx} + e^{4a+4bx} + 1)} + \frac{11e^{a+bx}}{3b(e^{2a+2bx} + 1)}$$

input `int(exp(a + b*x)*tanh(a + b*x)^4,x)`output `exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) + (4*exp(a + b*x))/(3*b) + (4*exp(5*a + 5*b*x))/(3*b)/(3*exp(2*a + 2*b*x) + 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) + 1) - (2*exp(a + b*x))/(b*(2*exp(2*a + 2*b*x) + exp(4*a + 4*b*x) + 1)) + (11*exp(a + b*x))/(3*b*(exp(2*a + 2*b*x) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.49

$$\int e^{a+bx} \tanh^4(a+bx) dx = \frac{-9e^{6bx+6a} \operatorname{atan}(e^{bx+a}) - 27e^{4bx+4a} \operatorname{atan}(e^{bx+a}) - 27e^{2bx+2a} \operatorname{atan}(e^{bx+a}) - 9 \operatorname{atan}(e^{bx+a}) + 3e^{7bx+7a} + 24e^{a+bx}}{3b(e^{6bx+6a} + 3e^{4bx+4a} + 3e^{2bx+2a} + 1)}$$

input `int(exp(b*x+a)*tanh(b*x+a)^4,x)`output `( - 9e**(6*a + 6*b*x)*atan(e**(a + b*x)) - 27e**(4*a + 4*b*x)*atan(e**(a + b*x)) - 27e**(2*a + 2*b*x)*atan(e**(a + b*x)) - 9*atan(e**(a + b*x)) + 3e**(7*a + 7*b*x) + 24e**(5*a + 5*b*x) + 25e**(3*a + 3*b*x) + 12e**(a + b*x))/(3*b*(e**(6*a + 6*b*x) + 3e**(4*a + 4*b*x) + 3e**(2*a + 2*b*x) + 1))`

### 3.217 $\int e^{a+bx} \tanh^3(a+bx) dx$

Optimal result	1604
Mathematica [A] (verified)	1604
Rubi [A] (verified)	1605
Maple [C] (verified)	1606
Fricas [B] (verification not implemented)	1607
Sympy [F]	1607
Maxima [A] (verification not implemented)	1608
Giac [A] (verification not implemented)	1608
Mupad [B] (verification not implemented)	1608
Reduce [B] (verification not implemented)	1609

#### Optimal result

Integrand size = 16, antiderivative size = 77

$$\int e^{a+bx} \tanh^3(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3 \arctan(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b-2*exp(b*x+a)/b/(1+exp(2*b*x+2*a))^2+3*exp(b*x+a)/b/(1+exp(2*b*x+2*a))-3*arctan(exp(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.78

$$\int e^{a+bx} \tanh^3(a+bx) dx = \frac{e^{a+bx} (2 + 5e^{2(a+bx)} + e^{4(a+bx)})}{b(1+e^{2(a+bx)})^2} - \frac{3 \arctan(e^{a+bx})}{b}$$

input

```
Integrate[E^(a + b*x)*Tanh[a + b*x]^3,x]
```

output

```
(E^(a + b*x)*(2 + 5*E^(2*(a + b*x)) + E^(4*(a + b*x))))/(b*(1 + E^(2*(a + b*x)))^2) - (3*ArcTan[E^(a + b*x)])/b
```

**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.88, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{a+bx} \tanh^3(a+bx) dx \\
 \downarrow 2720 \\
 \frac{\int -\frac{(1-e^{2a+2bx})^3}{(1+e^{2a+2bx})^3} de^{a+bx}}{b} \\
 \downarrow 25 \\
 \frac{\int \frac{(1-e^{2a+2bx})^3}{(1+e^{2a+2bx})^3} de^{a+bx}}{b} \\
 \downarrow 300 \\
 \frac{\int \left( \frac{2(1+3e^{4a+4bx})}{(1+e^{2a+2bx})^3} - 1 \right) de^{a+bx}}{b} \\
 \downarrow 2009 \\
 \frac{-3 \arctan(e^{a+bx}) + e^{a+bx} + \frac{3e^{a+bx}}{e^{2a+2bx}+1} - \frac{2e^{a+bx}}{(e^{2a+2bx}+1)^2}}{b}
 \end{array}$$

input

```
Int [E^(a + b*x)*Tanh[a + b*x]^3,x]
```

output

```
(E^(a + b*x) - (2*E^(a + b*x))/(1 + E^(2*a + 2*b*x))^2 + (3*E^(a + b*x))/(1 + E^(2*a + 2*b*x)) - 3*ArcTan[E^(a + b*x)])/b
```

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int [PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.04

method	result	size
risch	$\frac{e^{bx+a}}{b} + \frac{e^{bx+a}(3e^{2bx+2a}+1)}{b(1+e^{2bx+2a})^2} + \frac{3i \ln(e^{bx+a}-i)}{2b} - \frac{3i \ln(e^{bx+a}+i)}{2b}$	80
derivativedivides	$\frac{\sinh(bx+a)^3 + \frac{3 \sinh(bx+a)}{\cosh(bx+a)^2} - \frac{3 \operatorname{sech}(bx+a) \tanh(bx+a)}{2} - 3 \arctan(e^{bx+a}) + \frac{\sinh(bx+a)^2}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)}}{b}$	89
default	$\frac{\sinh(bx+a)^3 + \frac{3 \sinh(bx+a)}{\cosh(bx+a)^2} - \frac{3 \operatorname{sech}(bx+a) \tanh(bx+a)}{2} - 3 \arctan(e^{bx+a}) + \frac{\sinh(bx+a)^2}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)}}{b}$	89

input `int(exp(b*x+a)*tanh(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `exp(b*x+a)/b+exp(b*x+a)*(3*exp(2*b*x+2*a)+1)/b/(1+exp(2*b*x+2*a))^2+3/2*I/b*ln(exp(b*x+a)-I)-3/2*I/b*ln(exp(b*x+a)+I)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 339 vs.  $2(71) = 142$ .

Time = 0.09 (sec) , antiderivative size = 339, normalized size of antiderivative = 4.40

$$\int e^{a+bx} \tanh^3(a+bx) dx$$

$$= \frac{\cosh(bx+a)^5 + 5 \cosh(bx+a) \sinh(bx+a)^4 + \sinh(bx+a)^5 + 5(2 \cosh(bx+a)^2 + 1) \sinh(bx+a)}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^3,x, algorithm="fricas")`

output `(cosh(b*x + a)^5 + 5*cosh(b*x + a)*sinh(b*x + a)^4 + sinh(b*x + a)^5 + 5*(2*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^3 + 5*cosh(b*x + a)^3 + 5*(2*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^2 - 3*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x + a) + 1)*arctan(cosh(b*x + a) + sinh(b*x + a)) + (5*cosh(b*x + a)^4 + 15*cosh(b*x + a)^2 + 2)*sinh(b*x + a) + 2*cosh(b*x + a))/(b*cosh(b*x + a)^4 + 4*b*cosh(b*x + a)*sinh(b*x + a)^3 + b*sinh(b*x + a)^4 + 2*b*cosh(b*x + a)^2 + 2*(3*b*cosh(b*x + a)^2 + b)*sinh(b*x + a)^2 + 4*(b*cosh(b*x + a)^3 + b*cosh(b*x + a))*sinh(b*x + a) + b)`

**Sympy [F]**

$$\int e^{a+bx} \tanh^3(a+bx) dx = e^a \int e^{bx} \tanh^3(a+bx) dx$$

input `integrate(exp(b*x+a)*tanh(b*x+a)**3,x)`

output `exp(a)*Integral(exp(b*x)*tanh(a + b*x)**3, x)`



**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.90

$$\int e^{a+bx} \tanh^3(a+bx) dx = -\frac{3 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b} + \frac{3e^{(3bx+3a)} + e^{(bx+a)}}{b(e^{(4bx+4a)} + 2e^{(2bx+2a)} + 1)}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^3,x, algorithm="maxima")`output `-3*arctan(e^(b*x + a))/b + e^(b*x + a)/b + (3*e^(3*b*x + 3*a) + e^(b*x + a))/(b*(e^(4*b*x + 4*a) + 2*e^(2*b*x + 2*a) + 1))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.68

$$\int e^{a+bx} \tanh^3(a+bx) dx = \frac{\frac{3e^{(3bx+3a)} + e^{(bx+a)}}{(e^{(2bx+2a)} + 1)^2} - 3 \arctan(e^{(bx+a)}) + e^{(bx+a)}}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^3,x, algorithm="giac")`output `((3*e^(3*b*x + 3*a) + e^(b*x + a))/(e^(2*b*x + 2*a) + 1)^2 - 3*arctan(e^(b*x + a)) + e^(b*x + a))/b`**Mupad [B] (verification not implemented)**

Time = 2.23 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.21

$$\int e^{a+bx} \tanh^3(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} - \frac{2e^{a+bx}}{b(2e^{2a+2bx} + e^{4a+4bx} + 1)} + \frac{3e^{a+bx}}{b(e^{2a+2bx} + 1)}$$

input `int(exp(a + b*x)*tanh(a + b*x)^3,x)`

output

```
exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) - (
2*exp(a + b*x))/(b*(2*exp(2*a + 2*b*x) + exp(4*a + 4*b*x) + 1)) + (3*exp(a
+ b*x))/(b*(exp(2*a + 2*b*x) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.45

$$\int e^{a+bx} \tanh^3(a+bx) dx$$

$$= \frac{-3e^{4bx+4a} \operatorname{atan}(e^{bx+a}) - 6e^{2bx+2a} \operatorname{atan}(e^{bx+a}) - 3\operatorname{atan}(e^{bx+a}) + e^{5bx+5a} + 5e^{3bx+3a} + 2e^{bx+a}}{b(e^{4bx+4a} + 2e^{2bx+2a} + 1)}$$

input

```
int(exp(b*x+a)*tanh(b*x+a)^3,x)
```

output

```
( - 3*e**(4*a + 4*b*x)*atan(e**(a + b*x)) - 6*e**(2*a + 2*b*x)*atan(e**(a
+ b*x)) - 3*atan(e**(a + b*x)) + e**(5*a + 5*b*x) + 5*e**(3*a + 3*b*x) + 2
*e**(a + b*x))/(b*(e**(4*a + 4*b*x) + 2*e**(2*a + 2*b*x) + 1))
```

### 3.218 $\int e^{a+bx} \tanh^2(a+bx) dx$

Optimal result	1610
Mathematica [A] (verified)	1610
Rubi [A] (verified)	1611
Maple [A] (verified)	1612
Fricas [B] (verification not implemented)	1613
Sympy [F]	1613
Maxima [A] (verification not implemented)	1614
Giac [A] (verification not implemented)	1614
Mupad [B] (verification not implemented)	1614
Reduce [B] (verification not implemented)	1615

#### Optimal result

Integrand size = 16, antiderivative size = 51

$$\int e^{a+bx} \tanh^2(a+bx) dx = \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{2 \arctan(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b+2*exp(b*x+a)/b/(1+exp(2*b*x+2*a))-2*arctan(exp(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.78

$$\int e^{a+bx} \tanh^2(a+bx) dx = \frac{e^{a+bx} \left(1 + \frac{2}{1+e^{2(a+bx)}}\right) - 2 \arctan(e^{a+bx})}{b}$$

input

```
Integrate[E^(a + b*x)*Tanh[a + b*x]^2,x]
```

output

```
(E^(a + b*x)*(1 + 2/(1 + E^(2*(a + b*x)))) - 2*ArcTan[E^(a + b*x)])/b
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {2720, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{a+bx} \tanh^2(a+bx) dx \\
 \downarrow 2720 \\
 \frac{\int \frac{(1-e^{2a+2bx})^2}{(1+e^{2a+2bx})^2} de^{a+bx}}{b} \\
 \downarrow 300 \\
 \frac{\int \left(1 - \frac{4e^{2a+2bx}}{(1+e^{2a+2bx})^2}\right) de^{a+bx}}{b} \\
 \downarrow 2009 \\
 \frac{-2 \arctan(e^{a+bx}) + e^{a+bx} + \frac{2e^{a+bx}}{e^{2a+2bx}+1}}{b}
 \end{array}$$

input `Int [E^(a + b*x)*Tanh[a + b*x]^2,x]`

output `(E^(a + b*x) + (2*E^(a + b*x))/(1 + E^(2*a + 2*b*x)) - 2*ArcTan[E^(a + b*x)])/b`

## Definitions of rubi rules used

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{\frac{\sinh(bx+a)^2}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)} + \sinh(bx+a) - 2 \arctan(e^{bx+a})}{b}$	48
default	$\frac{\frac{\sinh(bx+a)^2}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)} + \sinh(bx+a) - 2 \arctan(e^{bx+a})}{b}$	48
risch	$\frac{e^{bx+a}}{b} + \frac{2e^{bx+a}}{b(1+e^{2bx+2a})} + \frac{i \ln(e^{bx+a}-i)}{b} - \frac{i \ln(e^{bx+a}+i)}{b}$	68

input `int(exp(b*x+a)*tanh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/b*(sinh(b*x+a)^2/cosh(b*x+a)+2/cosh(b*x+a)+sinh(b*x+a)-2*arctan(exp(b*x+a)))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 147 vs.  $2(47) = 94$ .

Time = 0.12 (sec) , antiderivative size = 147, normalized size of antiderivative = 2.88

$$\int e^{a+bx} \tanh^2(a+bx) dx$$

$$= \frac{\cosh(bx+a)^3 + 3 \cosh(bx+a) \sinh(bx+a)^2 + \sinh(bx+a)^3 - 2(\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 + 1) \operatorname{arctan}(\cosh(bx+a) + \sinh(bx+a)) + 3(\cosh(bx+a)^2 + 1) \sinh(bx+a) + 3 \cosh(bx+a)}{b \cosh(bx+a)^2 + 2 b \cosh(bx+a) \sinh(bx+a) + b \sinh(bx+a)^2 + b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="fricas")`

output `(cosh(b*x + a)^3 + 3*cosh(b*x + a)*sinh(b*x + a)^2 + sinh(b*x + a)^3 - 2*(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)*arc tan(cosh(b*x + a) + sinh(b*x + a)) + 3*(cosh(b*x + a)^2 + 1)*sinh(b*x + a) + 3*cosh(b*x + a))/(b*cosh(b*x + a)^2 + 2*b*cosh(b*x + a)*sinh(b*x + a) + b*sinh(b*x + a)^2 + b)`

**Sympy [F]**

$$\int e^{a+bx} \tanh^2(a+bx) dx = e^a \int e^{bx} \tanh^2(a+bx) dx$$

input `integrate(exp(b*x+a)*tanh(b*x+a)**2,x)`

output `exp(a)*Integral(exp(b*x)*tanh(a + b*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.92

$$\int e^{a+bx} \tanh^2(a+bx) dx = -\frac{2 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b} + \frac{2e^{(bx+a)}}{b(e^{(2bx+2a)} + 1)}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="maxima")`output `-2*arctan(e^(b*x + a))/b + e^(b*x + a)/b + 2*e^(b*x + a)/(b*(e^(2*b*x + 2*a) + 1))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.80

$$\int e^{a+bx} \tanh^2(a+bx) dx = \frac{\frac{2e^{(bx+a)}}{e^{(2bx+2a)}+1} - 2 \arctan(e^{(bx+a)}) + e^{(bx+a)}}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="giac")`output `(2*e^(b*x + a)/(e^(2*b*x + 2*a) + 1) - 2*arctan(e^(b*x + a)) + e^(b*x + a))/b`**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int e^{a+bx} \tanh^2(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} + \frac{2e^{a+bx}}{b(e^{2a+2bx} + 1)}$$

input `int(exp(a + b*x)*tanh(a + b*x)^2,x)`output `exp(a + b*x)/b - (2*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) + (2*exp(a + b*x))/(b*(exp(2*a + 2*b*x) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.33

$$\int e^{a+bx} \tanh^2(a+bx) dx = \frac{-2e^{2bx+2a} \operatorname{atan}(e^{bx+a}) - 2\operatorname{atan}(e^{bx+a}) + e^{3bx+3a} + 3e^{bx+a}}{b(e^{2bx+2a} + 1)}$$

input `int(exp(b*x+a)*tanh(b*x+a)^2,x)`output `( - 2*e**(2*a + 2*b*x)*atan(e**(a + b*x)) - 2*atan(e**(a + b*x)) + e**(3*a + 3*b*x) + 3*e**(a + b*x))/(b*(e**(2*a + 2*b*x) + 1))`



### 3.219 $\int e^{a+bx} \tanh(a + bx) dx$

Optimal result	1616
Mathematica [A] (verified)	1616
Rubi [A] (verified)	1617
Maple [A] (verified)	1618
Fricas [A] (verification not implemented)	1619
Sympy [F]	1619
Maxima [A] (verification not implemented)	1619
Giac [A] (verification not implemented)	1620
Mupad [B] (verification not implemented)	1620
Reduce [B] (verification not implemented)	1620

#### Optimal result

Integrand size = 14, antiderivative size = 25

$$\int e^{a+bx} \tanh(a + bx) dx = \frac{e^{a+bx}}{b} - \frac{2 \arctan(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b-2*arctan(exp(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int e^{a+bx} \tanh(a + bx) dx = \frac{e^{a+bx} - 2 \arctan(e^{a+bx})}{b}$$

input

```
Integrate[E^(a + b*x)*Tanh[a + b*x], x]
```

output

```
(E^(a + b*x) - 2*ArcTan[E^(a + b*x)])/b
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {2720, 25, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{a+bx} \tanh(a+bx) dx \\
 \downarrow 2720 \\
 \frac{\int -\frac{1-e^{2a+2bx}}{1+e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 25 \\
 \frac{\int \frac{1-e^{2a+2bx}}{1+e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 299 \\
 \frac{e^{a+bx} - 2 \int \frac{1}{1+e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 216 \\
 \frac{e^{a+bx} - 2 \arctan(e^{a+bx})}{b}
 \end{array}$$

input `Int [E^(a + b*x)*Tanh[a + b*x], x]`

output `(E^(a + b*x) - 2*ArcTan[E^(a + b*x)])/b`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{\sinh(bx+a) - 2 \arctan(e^{bx+a}) + \cosh(bx+a)}{b}$	27
default	$\frac{\sinh(bx+a) - 2 \arctan(e^{bx+a}) + \cosh(bx+a)}{b}$	27
risch	$\frac{e^{bx+a}}{b} + \frac{i \ln(e^{bx+a} - i)}{b} - \frac{i \ln(e^{bx+a} + i)}{b}$	44

input `int(exp(b*x+a)*tanh(b*x+a), x, method=_RETURNVERBOSE)`

output `1/b*(sinh(b*x+a)-2*arctan(exp(b*x+a))+cosh(b*x+a))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.52

$$\int e^{a+bx} \tanh(a+bx) dx = -\frac{2 \arctan(\cosh(bx+a) + \sinh(bx+a)) - \cosh(bx+a) - \sinh(bx+a)}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="fricas")`

output `-(2*arctan(cosh(b*x + a) + sinh(b*x + a)) - cosh(b*x + a) - sinh(b*x + a)) / b`

**Sympy [F]**

$$\int e^{a+bx} \tanh(a+bx) dx = e^a \int e^{bx} \tanh(a+bx) dx$$

input `integrate(exp(b*x+a)*tanh(b*x+a),x)`

output `exp(a)*Integral(exp(b*x)*tanh(a + b*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int e^{a+bx} \tanh(a+bx) dx = -\frac{2 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="maxima")`

output `-2*arctan(e^(b*x + a))/b + e^(b*x + a)/b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int e^{a+bx} \tanh(a+bx) dx = -\frac{2 \arctan(e^{(bx+a)}) - e^{(bx+a)}}{b}$$

input `integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="giac")`output `-(2*arctan(e^(b*x + a)) - e^(b*x + a))/b`**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int e^{a+bx} \tanh(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}}$$

input `int(exp(a + b*x)*tanh(a + b*x),x)`output `exp(a + b*x)/b - (2*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int e^{a+bx} \tanh(a+bx) dx = \frac{-2 \operatorname{atan}(e^{bx+a}) + e^{bx+a}}{b}$$

input `int(exp(b*x+a)*tanh(b*x+a),x)`output `( - 2*atan(e**(a + b*x)) + e**(a + b*x))/b`

### 3.220 $\int e^{a+bx} \coth(a + bx) dx$

Optimal result	1621
Mathematica [A] (verified)	1621
Rubi [A] (verified)	1622
Maple [A] (verified)	1623
Fricas [B] (verification not implemented)	1624
Sympy [F]	1624
Maxima [A] (verification not implemented)	1624
Giac [A] (verification not implemented)	1625
Mupad [B] (verification not implemented)	1625
Reduce [B] (verification not implemented)	1625

#### Optimal result

Integrand size = 14, antiderivative size = 25

$$\int e^{a+bx} \coth(a + bx) dx = \frac{e^{a+bx}}{b} - \frac{2\operatorname{arctanh}(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b-2*arctanh(exp(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int e^{a+bx} \coth(a + bx) dx = \frac{e^{a+bx} - 2\operatorname{arctanh}(e^{a+bx})}{b}$$

input

```
Integrate[E^(a + b*x)*Coth[a + b*x],x]
```

output

```
(E^(a + b*x) - 2*ArcTanh[E^(a + b*x)])/b
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {2720, 25, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{a+bx} \coth(a+bx) dx \\
 \downarrow 2720 \\
 \frac{\int -\frac{1+e^{2a+2bx}}{1-e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 25 \\
 \frac{\int \frac{1+e^{2a+2bx}}{1-e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 299 \\
 \frac{e^{a+bx} - 2 \int \frac{1}{1-e^{2a+2bx}} de^{a+bx}}{b} \\
 \downarrow 219 \\
 \frac{e^{a+bx} - 2\operatorname{arctanh}(e^{a+bx})}{b}
 \end{array}$$

input `Int [E^(a + b*x)*Coth[a + b*x], x]`

output `(E^(a + b*x) - 2*ArcTanh[E^(a + b*x)])/b`

## Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{\sinh(bx+a)+\cosh(bx+a)-2 \operatorname{arctanh}(e^{bx+a})}{b}$	27
default	$\frac{\sinh(bx+a)+\cosh(bx+a)-2 \operatorname{arctanh}(e^{bx+a})}{b}$	27
risch	$\frac{e^{bx+a}}{b} + \frac{\ln(e^{bx+a}-1)}{b} - \frac{\ln(e^{bx+a}+1)}{b}$	39

input `int(exp(b*x+a)*coth(b*x+a),x,method=_RETURNVERBOSE)`

output `1/b*(sinh(b*x+a)+cosh(b*x+a)-2*arctanh(exp(b*x+a)))`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 49 vs.  $2(23) = 46$ .

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.96

$$\int e^{a+bx} \coth(a+bx) dx = \frac{\cosh(bx+a) - \log(\cosh(bx+a) + \sinh(bx+a) + 1) + \log(\cosh(bx+a) + \sinh(bx+a) - 1) + \sinh(bx+a)}{b}$$

input `integrate(exp(b*x+a)*coth(b*x+a),x, algorithm="fricas")`

output `(cosh(b*x + a) - log(cosh(b*x + a) + sinh(b*x + a) + 1) + log(cosh(b*x + a) + sinh(b*x + a) - 1) + sinh(b*x + a))/b`

**Sympy [F]**

$$\int e^{a+bx} \coth(a+bx) dx = e^a \int e^{bx} \coth(a+bx) dx$$

input `integrate(exp(b*x+a)*coth(b*x+a),x)`

output `exp(a)*Integral(exp(b*x)*coth(a + b*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.52

$$\int e^{a+bx} \coth(a+bx) dx = \frac{e^{(bx+a)}}{b} - \frac{\log(e^{(bx+a)} + 1)}{b} + \frac{\log(e^{(bx+a)} - 1)}{b}$$

input `integrate(exp(b*x+a)*coth(b*x+a),x, algorithm="maxima")`

output `e^(b*x + a)/b - log(e^(b*x + a) + 1)/b + log(e^(b*x + a) - 1)/b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.28

$$\int e^{a+bx} \coth(a+bx) dx = \frac{e^{(bx+a)} - \log(e^{(bx+a)} + 1) + \log(|e^{(bx+a)} - 1|)}{b}$$

input `integrate(exp(b*x+a)*coth(b*x+a),x, algorithm="giac")`output `(e^(b*x + a) - log(e^(b*x + a) + 1) + log(abs(e^(b*x + a) - 1)))/b`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.52

$$\int e^{a+bx} \coth(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}}$$

input `int(coth(a + b*x)*exp(a + b*x),x)`output `exp(a + b*x)/b - (2*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

$$\int e^{a+bx} \coth(a+bx) dx = \frac{e^{bx+a} + \log(e^{bx+a} - 1) - \log(e^{bx+a} + 1)}{b}$$

input `int(exp(b*x+a)*coth(b*x+a),x)`output `(e**(a + b*x) + log(e**(a + b*x) - 1) - log(e**(a + b*x) + 1))/b`

### 3.221 $\int e^{a+bx} \coth^2(a + bx) dx$

Optimal result	1626
Mathematica [C] (verified)	1626
Rubi [A] (verified)	1627
Maple [A] (verified)	1628
Fricas [B] (verification not implemented)	1629
Sympy [F]	1629
Maxima [A] (verification not implemented)	1630
Giac [A] (verification not implemented)	1630
Mupad [B] (verification not implemented)	1630
Reduce [B] (verification not implemented)	1631

#### Optimal result

Integrand size = 16, antiderivative size = 53

$$\int e^{a+bx} \coth^2(a + bx) dx = \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1 - e^{2a+2bx})} - \frac{2\operatorname{arctanh}(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b+2*exp(b*x+a)/b/(1-exp(2*b*x+2*a))-2*arctanh(exp(b*x+a))/b
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.44 (sec) , antiderivative size = 179, normalized size of antiderivative = 3.38

$$\int e^{a+bx} \coth^2(a + bx) dx = \frac{e^{a+bx} \left( \frac{1}{48} e^{-4(a+bx)} \left( -375 - 713e^{2(a+bx)} - 181e^{4(a+bx)} + 61e^{6(a+bx)} \right) + \frac{3(125+196e^{2(a+bx)} - 14e^{4(a+bx)} - 52e^{6(a+bx)} + e^8)}{\sqrt{e^{2(a+bx)}}} \right)}{b}$$

input

```
Integrate[E^(a + b*x)*Coth[a + b*x]^2,x]
```

output

$$\frac{(E^{(a + b*x)}*((-375 - 713*E^{(2*(a + b*x))} - 181*E^{(4*(a + b*x))} + 61*E^{(6*(a + b*x))} + (3*(125 + 196*E^{(2*(a + b*x))} - 14*E^{(4*(a + b*x))} - 52*E^{(6*(a + b*x))} + E^{(8*(a + b*x))))*ArcTanh[Sqrt[E^{(2*(a + b*x))}]]/Sqrt[E^{(2*(a + b*x))}])/(48*E^{(4*(a + b*x))} + (4*(E^{(a + b*x)} + E^{(3*(a + b*x)))^2*HypergeometricPFQ[{3/2, 2, 2, 2}, {1, 1, 9/2}, E^{(2*(a + b*x))}])/105))/b$$
**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {2720, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{a+bx} \coth^2(a+bx) dx$$

$$\downarrow \text{2720}$$

$$\frac{\int \frac{(1+e^{2a+2bx})^2}{(1-e^{2a+2bx})^2} de^{a+bx}}{b}$$

$$\downarrow \text{300}$$

$$\frac{\int \left(1 + \frac{4e^{2a+2bx}}{(1-e^{2a+2bx})^2}\right) de^{a+bx}}{b}$$

$$\downarrow \text{2009}$$

$$\frac{-2\operatorname{arctanh}(e^{a+bx}) + e^{a+bx} + \frac{2e^{a+bx}}{1-e^{2a+2bx}}}{b}$$

input

$$\text{Int}[E^{(a + b*x)}*\text{Coth}[a + b*x]^2, x]$$

output

$$(E^{(a + b*x)} + (2*E^{(a + b*x)})/(1 - E^{(2*a + 2*b*x)}) - 2*ArcTanh[E^{(a + b*x)}])/b$$

## Definitions of rubi rules used

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{\cosh(bx+a) - 2 \operatorname{arctanh}(e^{bx+a}) + \frac{\cosh(bx+a)^2}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)}}{b}$	48
default	$\frac{\cosh(bx+a) - 2 \operatorname{arctanh}(e^{bx+a}) + \frac{\cosh(bx+a)^2}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)}}{b}$	48
risch	$\frac{e^{bx+a}}{b} - \frac{2e^{bx+a}}{b(-1+e^{2bx+2a})} + \frac{\ln(e^{bx+a}-1)}{b} - \frac{\ln(e^{bx+a}+1)}{b}$	63

input `int(exp(b*x+a)*coth(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/b*(cosh(b*x+a)-2*arctanh(exp(b*x+a))+cosh(b*x+a)^2/sinh(b*x+a)-2/sinh(b*x+a))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 198 vs.  $2(47) = 94$ .

Time = 0.09 (sec) , antiderivative size = 198, normalized size of antiderivative = 3.74

$$\int e^{a+bx} \coth^2(a+bx) dx$$

$$= \frac{\cosh(bx+a)^3 + 3 \cosh(bx+a) \sinh(bx+a)^2 + \sinh(bx+a)^3 - (\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 - 1) \log(\cosh(bx+a) + \sinh(bx+a) + 1) + (\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 - 1) \log(\cosh(bx+a) + \sinh(bx+a) - 1) + 3(\cosh(bx+a)^2 - 1) \sinh(bx+a) - 3 \cosh(bx+a)}{b \cosh(bx+a)^2 + 2b \cosh(bx+a) \sinh(bx+a) + b \sinh(bx+a)^2 - b}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="fricas")`

output `(cosh(b*x + a)^3 + 3*cosh(b*x + a)*sinh(b*x + a)^2 + sinh(b*x + a)^3 - (cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + (cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + 3*(cosh(b*x + a)^2 - 1)*sinh(b*x + a) - 3*cosh(b*x + a))/(b*cosh(b*x + a)^2 + 2*b*cosh(b*x + a)*sinh(b*x + a) + b*sinh(b*x + a)^2 - b)`

**Sympy [F]**

$$\int e^{a+bx} \coth^2(a+bx) dx = e^a \int e^{bx} \coth^2(a+bx) dx$$

input `integrate(exp(b*x+a)*coth(b*x+a)**2,x)`

output `exp(a)*Integral(exp(b*x)*coth(a + b*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.17

$$\int e^{a+bx} \coth^2(a+bx) dx = \frac{e^{(bx+a)}}{b} - \frac{\log(e^{(bx+a)} + 1)}{b} + \frac{\log(e^{(bx+a)} - 1)}{b} - \frac{2e^{(bx+a)}}{b(e^{2bx+2a} - 1)}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="maxima")`output `e^(b*x + a)/b - log(e^(b*x + a) + 1)/b + log(e^(b*x + a) - 1)/b - 2*e^(b*x + a)/(b*(e^(2*b*x + 2*a) - 1))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.06

$$\int e^{a+bx} \coth^2(a+bx) dx = -\frac{\frac{2e^{(bx+a)}}{e^{(2bx+2a)} - 1} - e^{(bx+a)} + \log(e^{(bx+a)} + 1) - \log(|e^{(bx+a)} - 1|)}{b}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="giac")`output `-(2*e^(b*x + a)/(e^(2*b*x + 2*a) - 1) - e^(b*x + a) + log(e^(b*x + a) + 1) - log(abs(e^(b*x + a) - 1)))/b`**Mupad [B] (verification not implemented)**

Time = 2.09 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.17

$$\int e^{a+bx} \coth^2(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{2e^{a+bx}}{b(e^{2a+2bx} - 1)}$$

input `int(coth(a + b*x)^2*exp(a + b*x),x)`

output

$$\frac{\exp(a + b*x)/b - (2*\operatorname{atan}((\exp(b*x)*\exp(a)*(-b^2)^{(1/2)})/b))/(-b^2)^{(1/2)} - (2*\exp(a + b*x))/(b*(\exp(2*a + 2*b*x) - 1))}{b(e^{2bx+2a} - 1)}$$

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.94

$$\int e^{a+bx} \coth^2(a + bx) dx$$

$$= \frac{e^{3bx+3a} + e^{2bx+2a} \log(e^{bx+a} - 1) - e^{2bx+2a} \log(e^{bx+a} + 1) - 3e^{bx+a} - \log(e^{bx+a} - 1) + \log(e^{bx+a} + 1)}{b(e^{2bx+2a} - 1)}$$

input

```
int(exp(b*x+a)*coth(b*x+a)^2,x)
```

output

```
(e**(3*a + 3*b*x) + e**(2*a + 2*b*x)*log(e**(a + b*x) - 1) - e**(2*a + 2*b*x)*log(e**(a + b*x) + 1) - 3*e**(a + b*x) - log(e**(a + b*x) - 1) + log(e**(a + b*x) + 1))/(b*(e**(2*a + 2*b*x) - 1))
```



### 3.222 $\int e^{a+bx} \coth^3(a + bx) dx$

Optimal result . . . . .	1632
Mathematica [C] (warning: unable to verify) . . . . .	1632
Rubi [A] (verified) . . . . .	1633
Maple [A] (verified) . . . . .	1634
Fricas [B] (verification not implemented) . . . . .	1635
Sympy [F] . . . . .	1636
Maxima [A] (verification not implemented) . . . . .	1636
Giac [A] (verification not implemented) . . . . .	1636
Mupad [B] (verification not implemented) . . . . .	1637
Reduce [B] (verification not implemented) . . . . .	1637

#### Optimal result

Integrand size = 16, antiderivative size = 81

$$\int e^{a+bx} \coth^3(a + bx) dx = \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1 - e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1 - e^{2a+2bx})} - \frac{3\operatorname{arctanh}(e^{a+bx})}{b}$$

output

```
exp(b*x+a)/b-2*exp(b*x+a)/b/(1-exp(2*b*x+2*a))^2+3*exp(b*x+a)/b/(1-exp(2*b*x+2*a))-3*arctanh(exp(b*x+a))/b
```

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.79 (sec) , antiderivative size = 286, normalized size of antiderivative = 3.53

$$\int e^{a+bx} \coth^3(a + bx) dx = \frac{e^{-5(a+bx)} \left( -21(252105 + 507305e^{2(a+bx)} + 173916e^{4(a+bx)} - 154296e^{6(a+bx)} - 73885e^{8(a+bx)} + 4887e^{10(a+bx)}) \right)}{\dots}$$

input

```
Integrate[E^(a + b*x)*Coth[a + b*x]^3,x]
```

output

```

-1/60480*(-21*(252105 + 507305*E^(2*(a + b*x)) + 173916*E^(4*(a + b*x)) -
154296*E^(6*(a + b*x)) - 73885*E^(8*(a + b*x)) + 4887*E^(10*(a + b*x))) -
(315*(-16807 - 28218*E^(2*(a + b*x)) + 1173*E^(4*(a + b*x)) + 17748*E^(6*(
a + b*x)) + 4299*E^(8*(a + b*x)) - 1434*E^(10*(a + b*x)) + 7*E^(12*(a + b*
x)))*ArcTanh[Sqrt[E^(2*(a + b*x))]]/Sqrt[E^(2*(a + b*x))] + 384*E^(8*(a +
b*x))*(1 + E^(2*(a + b*x)))^2*(7 + 5*E^(2*(a + b*x)))*HypergeometricPFQ[{
3/2, 2, 2, 2, 2}, {1, 1, 1, 11/2}, E^(2*(a + b*x))] + 256*E^(8*(a + b*x))*
(1 + E^(2*(a + b*x)))^3*HypergeometricPFQ[{3/2, 2, 2, 2, 2, 2}, {1, 1, 1,
1, 11/2}, E^(2*(a + b*x))]/(b*E^(5*(a + b*x)))

```

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.89, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{a+bx} \coth^3(a+bx) dx \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int -\frac{(1+e^{2a+2bx})^3}{(1-e^{2a+2bx})^3} de^{a+bx}}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{(1+e^{2a+2bx})^3}{(1-e^{2a+2bx})^3} de^{a+bx}}{b} \\
 & \quad \downarrow \text{300} \\
 & \frac{\int \left( \frac{2(1+3e^{4a+4bx})}{(1-e^{2a+2bx})^3} - 1 \right) de^{a+bx}}{b} \\
 & \quad \downarrow \text{2009} \\
 & \frac{-3\arctanh(e^{a+bx}) + e^{a+bx} + \frac{3e^{a+bx}}{1-e^{2a+2bx}} - \frac{2e^{a+bx}}{(1-e^{2a+2bx})^2}}{b}
 \end{aligned}$$

input `Int[E^(a + b*x)*Coth[a + b*x]^3,x]`

output 
$$\frac{(E^{(a + b*x)} - (2*E^{(a + b*x)})/(1 - E^{(2*a + 2*b*x)})^2 + (3*E^{(a + b*x)})/(1 - E^{(2*a + 2*b*x)}) - 3*ArcTanh[E^{(a + b*x)}])/b}$$

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 300 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

### Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.95

method	result	size
risch	$\frac{e^{bx+a}}{b} - \frac{e^{bx+a}(3e^{2bx+2a}-1)}{b(-1+e^{2bx+2a})^2} + \frac{3\ln(e^{bx+a}-1)}{2b} - \frac{3\ln(e^{bx+a}+1)}{2b}$	77
derivativedivides	$\frac{\frac{\cosh(bx+a)^2}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)} + \frac{\cosh(bx+a)^3}{\sinh(bx+a)^2} - \frac{3\cosh(bx+a)}{\sinh(bx+a)^2} + \frac{3\operatorname{csch}(bx+a)\operatorname{coth}(bx+a)}{2} - 3\operatorname{arctanh}(e^{bx+a})}{b}$	89
default	$\frac{\frac{\cosh(bx+a)^2}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)} + \frac{\cosh(bx+a)^3}{\sinh(bx+a)^2} - \frac{3\cosh(bx+a)}{\sinh(bx+a)^2} + \frac{3\operatorname{csch}(bx+a)\operatorname{coth}(bx+a)}{2} - 3\operatorname{arctanh}(e^{bx+a})}{b}$	89

input `int(exp(b*x+a)*coth(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `exp(b*x+a)/b-exp(b*x+a)*(3*exp(2*b*x+2*a)-1)/b/(-1+exp(2*b*x+2*a))^2+3/2/b  
*ln(exp(b*x+a)-1)-3/2/b*ln(exp(b*x+a)+1)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 459 vs.  $2(71) = 142$ .

Time = 0.10 (sec) , antiderivative size = 459, normalized size of antiderivative = 5.67

$$\int e^{a+bx} \coth^3(a+bx) dx$$

$$= \frac{2 \cosh(bx+a)^5 + 10 \cosh(bx+a) \sinh(bx+a)^4 + 2 \sinh(bx+a)^5 + 10 (2 \cosh(bx+a)^2 - 1) \sinh(bx+a)}{b}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="fricas")`

output `1/2*(2*cosh(b*x + a)^5 + 10*cosh(b*x + a)*sinh(b*x + a)^4 + 2*sinh(b*x + a)  
)^5 + 10*(2*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^3 - 10*cosh(b*x + a)^3 + 10  
*(2*cosh(b*x + a)^3 - 3*cosh(b*x + a))*sinh(b*x + a)^2 - 3*(cosh(b*x + a)^  
4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)  
^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*  
x + a))*sinh(b*x + a) + 1)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 3*(cos  
h(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*co  
sh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^  
3 - cosh(b*x + a))*sinh(b*x + a) + 1)*log(cosh(b*x + a) + sinh(b*x + a) -  
1) + 2*(5*cosh(b*x + a)^4 - 15*cosh(b*x + a)^2 + 2)*sinh(b*x + a) + 4*cosh  
(b*x + a))/(b*cosh(b*x + a)^4 + 4*b*cosh(b*x + a)*sinh(b*x + a)^3 + b*sinh  
(b*x + a)^4 - 2*b*cosh(b*x + a)^2 + 2*(3*b*cosh(b*x + a)^2 - b)*sinh(b*x +  
a)^2 + 4*(b*cosh(b*x + a)^3 - b*cosh(b*x + a))*sinh(b*x + a) + b)`

**Sympy [F]**

$$\int e^{a+bx} \coth^3(a+bx) dx = e^a \int e^{bx} \coth^3(a+bx) dx$$

input `integrate(exp(b*x+a)*coth(b*x+a)**3,x)`

output `exp(a)*Integral(exp(b*x)*coth(a + b*x)**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.09

$$\int e^{a+bx} \coth^3(a+bx) dx = \frac{e^{(bx+a)}}{b} - \frac{3 \log(e^{(bx+a)} + 1)}{2b} + \frac{3 \log(e^{(bx+a)} - 1)}{2b} - \frac{3e^{(3bx+3a)} - e^{(bx+a)}}{b(e^{(4bx+4a)} - 2e^{(2bx+2a)} + 1)}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="maxima")`

output `e^(b*x + a)/b - 3/2*log(e^(b*x + a) + 1)/b + 3/2*log(e^(b*x + a) - 1)/b - (3*e^(3*b*x + 3*a) - e^(b*x + a))/(b*(e^(4*b*x + 4*a) - 2*e^(2*b*x + 2*a) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.89

$$\int e^{a+bx} \coth^3(a+bx) dx = \frac{2(3e^{(3bx+3a)} - e^{(bx+a)})}{(e^{(2bx+2a)} - 1)^2} - 2e^{(bx+a)} + 3 \log(e^{(bx+a)} + 1) - 3 \log(|e^{(bx+a)} - 1|)$$

$$= - \frac{\hspace{15em}}{2b}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="giac")`

output

```
-1/2*(2*(3*e^(3*b*x + 3*a) - e^(b*x + a))/(e^(2*b*x + 2*a) - 1)^2 - 2*e^(b*x + a) + 3*log(e^(b*x + a) + 1) - 3*log(abs(e^(b*x + a) - 1)))/b
```

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.20

$$\int e^{a+bx} \coth^3(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{2e^{a+bx}}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{3e^{a+bx}}{b(e^{2a+2bx} - 1)}$$

input

```
int(coth(a + b*x)^3*exp(a + b*x),x)
```

output

```
exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2) - (2*exp(a + b*x))/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - (3*exp(a + b*x))/(b*(exp(2*a + 2*b*x) - 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.19

$$\int e^{a+bx} \coth^3(a+bx) dx = \frac{2e^{5bx+5a} + 3e^{4bx+4a} \log(e^{bx+a} - 1) - 3e^{4bx+4a} \log(e^{bx+a} + 1) - 10e^{3bx+3a} - 6e^{2bx+2a} \log(e^{bx+a} - 1) + 6e^{2bx+2a} \log(e^{bx+a} + 1)}{2b(e^{4bx+4a} - 2e^{2bx+2a} + 1)}$$

input

```
int(exp(b*x+a)*coth(b*x+a)^3,x)
```

output

```
(2***e**(5*a + 5*b*x) + 3***e**(4*a + 4*b*x)*log(e**(a + b*x) - 1) - 3***e**(4*a + 4*b*x)*log(e**(a + b*x) + 1) - 10***e**(3*a + 3*b*x) - 6***e**(2*a + 2*b*x)*log(e**(a + b*x) - 1) + 6***e**(2*a + 2*b*x)*log(e**(a + b*x) + 1) + 4***e**(a + b*x) + 3*log(e**(a + b*x) - 1) - 3*log(e**(a + b*x) + 1))/(2*b*(e**(4*a + 4*b*x) - 2*e**(2*a + 2*b*x) + 1))
```

### 3.223 $\int e^{a+bx} \coth^4(a + bx) dx$

Optimal result	1638
Mathematica [A] (verified)	1638
Rubi [A] (verified)	1639
Maple [A] (verified)	1640
Fricas [B] (verification not implemented)	1641
Sympy [F]	1642
Maxima [A] (verification not implemented)	1642
Giac [A] (verification not implemented)	1642
Mupad [B] (verification not implemented)	1643
Reduce [B] (verification not implemented)	1643

#### Optimal result

Integrand size = 16, antiderivative size = 113

$$\int e^{a+bx} \coth^4(a + bx) dx = \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1 - e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1 - e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1 - e^{2a+2bx})} - \frac{3\operatorname{arctanh}(e^{a+bx})}{b}$$

output

$\exp(b*x+a)/b+8/3*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))^3-14/3*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))^2+5*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))-3*\operatorname{arctanh}(\exp(b*x+a))/b$

#### Mathematica [A] (verified)

Time = 10.08 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02

$$\int e^{a+bx} \coth^4(a + bx) dx = \frac{-24e^{a+bx} + 50e^{3(a+bx)} - 48e^{5(a+bx)} + 6e^{7(a+bx)} + 9(-1 + e^{2(a+bx)})^3 \log(1 - e^{a+bx}) - 9(-1 + e^{2(a+bx)})^3}{6b(-1 + e^{2(a+bx)})^3}$$

input

$\operatorname{Integrate}[E^{(a + b*x)}*\operatorname{Coth}[a + b*x]^4,x]$

output

$$\frac{(-24E^{(a + bx)} + 50E^{(3(a + bx))} - 48E^{(5(a + bx))} + 6E^{(7(a + bx))} + 9(-1 + E^{(2(a + bx))})^3 \text{Log}[1 - E^{(a + bx)}] - 9(-1 + E^{(2(a + bx))})^3 \text{Log}[1 + E^{(a + bx)}])}{(6b(-1 + E^{(2(a + bx))})^3)}$$

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$ , Rules used = {2720, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{a+bx} \coth^4(a+bx) dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{(1+e^{2a+2bx})^4}{(1-e^{2a+2bx})^4} de^{a+bx} \\ & \quad \downarrow \text{300} \\ & \int \left( \frac{8e^{2a+2bx}(1+e^{4a+4bx})}{(1-e^{2a+2bx})^4} + 1 \right) de^{a+bx} \\ & \quad \downarrow \text{2009} \\ & \frac{-3\text{arctanh}(e^{a+bx}) + e^{a+bx} + \frac{5e^{a+bx}}{1-e^{2a+2bx}} - \frac{14e^{a+bx}}{3(1-e^{2a+2bx})^2} + \frac{8e^{a+bx}}{3(1-e^{2a+2bx})^3}}{b} \end{aligned}$$

input

```
Int[E^(a + b*x)*Coth[a + b*x]^4,x]
```

output

```
(E^(a + b*x) + (8*E^(a + b*x))/(3*(1 - E^(2*a + 2*b*x))^3) - (14*E^(a + b*x))/(3*(1 - E^(2*a + 2*b*x))^2) + (5*E^(a + b*x))/(1 - E^(2*a + 2*b*x)) - 3*ArcTanh[E^(a + b*x)])/b
```



## Definitions of rubi rules used

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.78

method	result
risch	$\frac{e^{bx+a}}{b} - \frac{e^{bx+a} (15e^{4bx+4a} - 16e^{2bx+2a} + 9)}{3b(-1+e^{2bx+2a})^3} + \frac{3\ln(e^{bx+a}-1)}{2b} - \frac{3\ln(e^{bx+a}+1)}{2b}$
derivativedivides	$\frac{\frac{\cosh(bx+a)^3}{\sinh(bx+a)^2} - \frac{3\cosh(bx+a)}{\sinh(bx+a)^2} + \frac{3\operatorname{csch}(bx+a)\coth(bx+a)}{2} - 3\operatorname{arctanh}(e^{bx+a}) + \frac{\cosh(bx+a)^4}{\sinh(bx+a)^3} - \frac{4\cosh(bx+a)^2}{\sinh(bx+a)^3} + \frac{8}{3\sinh(bx+a)^3}}{b}$
default	$\frac{\frac{\cosh(bx+a)^3}{\sinh(bx+a)^2} - \frac{3\cosh(bx+a)}{\sinh(bx+a)^2} + \frac{3\operatorname{csch}(bx+a)\coth(bx+a)}{2} - 3\operatorname{arctanh}(e^{bx+a}) + \frac{\cosh(bx+a)^4}{\sinh(bx+a)^3} - \frac{4\cosh(bx+a)^2}{\sinh(bx+a)^3} + \frac{8}{3\sinh(bx+a)^3}}{b}$

input `int(exp(b*x+a)*coth(b*x+a)^4,x,method=_RETURNVERBOSE)`

output `exp(b*x+a)/b-1/3*exp(b*x+a)*(15*exp(4*b*x+4*a)-16*exp(2*b*x+2*a)+9)/b/(-1+exp(2*b*x+2*a))^3+3/2/b*ln(exp(b*x+a)-1)-3/2/b*ln(exp(b*x+a)+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 796 vs.  $2(95) = 190$ .

Time = 0.11 (sec) , antiderivative size = 796, normalized size of antiderivative = 7.04

$$\int e^{a+bx} \coth^4(a+bx) dx = \text{Too large to display}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="fricas")`

output

```

1/6*(6*cosh(b*x + a)^7 + 42*cosh(b*x + a)*sinh(b*x + a)^6 + 6*sinh(b*x + a)
)^7 + 6*(21*cosh(b*x + a)^2 - 8)*sinh(b*x + a)^5 - 48*cosh(b*x + a)^5 + 30
*(7*cosh(b*x + a)^3 - 8*cosh(b*x + a))*sinh(b*x + a)^4 + 10*(21*cosh(b*x +
a)^4 - 48*cosh(b*x + a)^2 + 5)*sinh(b*x + a)^3 + 50*cosh(b*x + a)^3 + 6*(
21*cosh(b*x + a)^5 - 80*cosh(b*x + a)^3 + 25*cosh(b*x + a))*sinh(b*x + a)^
2 - 9*(cosh(b*x + a)^6 + 6*cosh(b*x + a)*sinh(b*x + a)^5 + sinh(b*x + a)^6
+ 3*(5*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^4 - 3*cosh(b*x + a)^4 + 4*(5*co
sh(b*x + a)^3 - 3*cosh(b*x + a))*sinh(b*x + a)^3 + 3*(5*cosh(b*x + a)^4 -
6*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 3*cosh(b*x + a)^2 + 6*(cosh(b*x +
a)^5 - 2*cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x + a) - 1)*log(cosh(b*x
+ a) + sinh(b*x + a) + 1) + 9*(cosh(b*x + a)^6 + 6*cosh(b*x + a)*sinh(b*x
+ a)^5 + sinh(b*x + a)^6 + 3*(5*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^4 - 3*
cosh(b*x + a)^4 + 4*(5*cosh(b*x + a)^3 - 3*cosh(b*x + a))*sinh(b*x + a)^3
+ 3*(5*cosh(b*x + a)^4 - 6*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 3*cosh(b
*x + a)^2 + 6*(cosh(b*x + a)^5 - 2*cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b
*x + a) - 1)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + 6*(7*cosh(b*x + a)^6
- 40*cosh(b*x + a)^4 + 25*cosh(b*x + a)^2 - 4)*sinh(b*x + a) - 24*cosh(b*
x + a))/(b*cosh(b*x + a)^6 + 6*b*cosh(b*x + a)*sinh(b*x + a)^5 + b*sinh(b*
x + a)^6 - 3*b*cosh(b*x + a)^4 + 3*(5*b*cosh(b*x + a)^2 - b)*sinh(b*x + a)
^4 + 4*(5*b*cosh(b*x + a)^3 - 3*b*cosh(b*x + a))*sinh(b*x + a)^3 + 3*b*...
```

**Sympy [F]**

$$\int e^{a+bx} \coth^4(a+bx) dx = e^a \int e^{bx} \coth^4(a+bx) dx$$

input `integrate(exp(b*x+a)*coth(b*x+a)**4,x)`

output `exp(a)*Integral(exp(b*x)*coth(a + b*x)**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.97

$$\int e^{a+bx} \coth^4(a+bx) dx = \frac{e^{(bx+a)}}{b} - \frac{3 \log(e^{(bx+a)} + 1)}{2b} + \frac{3 \log(e^{(bx+a)} - 1)}{2b} - \frac{15e^{(5bx+5a)} - 16e^{(3bx+3a)} + 9e^{(bx+a)}}{3b(e^{(6bx+6a)} - 3e^{(4bx+4a)} + 3e^{(2bx+2a)} - 1)}$$

input `integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="maxima")`

output `e^(b*x + a)/b - 3/2*log(e^(b*x + a) + 1)/b + 3/2*log(e^(b*x + a) - 1)/b - 1/3*(15*e^(5*b*x + 5*a) - 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(b*(e^(6*b*x + 6*a) - 3*e^(4*b*x + 4*a) + 3*e^(2*b*x + 2*a) - 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.73

$$\int e^{a+bx} \coth^4(a+bx) dx = \frac{2(15e^{(5bx+5a)} - 16e^{(3bx+3a)} + 9e^{(bx+a)})}{(e^{(2bx+2a)} - 1)^3} - 6e^{(bx+a)} + 9 \log(e^{(bx+a)} + 1) - 9 \log(|e^{(bx+a)} - 1|)$$

$6b$

input `integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="giac")`

output

```
-1/6*(2*(15*e^(5*b*x + 5*a) - 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(e^(2*b*x + 2*a) - 1)^3 - 6*e^(b*x + a) + 9*log(e^(b*x + a) + 1) - 9*log(abs(e^(b*x + a) - 1)))/b
```

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.42

$$\int e^{a+bx} \coth^4(a+bx) dx = \frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{\frac{4e^{a+bx}}{3b} + \frac{4e^{5a+5bx}}{3b}}{3e^{2a+2bx} - 3e^{4a+4bx} + e^{6a+6bx} - 1} - \frac{2e^{a+bx}}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{11e^{a+bx}}{3b(e^{2a+2bx} - 1)}$$

input

```
int(coth(a + b*x)^4*exp(a + b*x),x)
```

output

```
exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2) - ((4*exp(a + b*x))/(3*b) + (4*exp(5*a + 5*b*x))/(3*b))/(3*exp(2*a + 2*b*x) - 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) - 1) - (2*exp(a + b*x))/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - (11*exp(a + b*x))/(3*b*(exp(2*a + 2*b*x) - 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 245, normalized size of antiderivative = 2.17

$$\int e^{a+bx} \coth^4(a+bx) dx = \frac{6e^{7bx+7a} + 9e^{6bx+6a} \log(e^{bx+a} - 1) - 9e^{6bx+6a} \log(e^{bx+a} + 1) - 48e^{5bx+5a} - 27e^{4bx+4a} \log(e^{bx+a} - 1) + 27e^{4bx+4a} \log(e^{bx+a} + 1)}{6b}$$

input

```
int(exp(b*x+a)*coth(b*x+a)^4,x)
```

output

```
(6***(7*a + 7*b*x) + 9***(6*a + 6*b*x)*log(e**(a + b*x) - 1) - 9***(6*a
+ 6*b*x)*log(e**(a + b*x) + 1) - 48***(5*a + 5*b*x) - 27***(4*a + 4*b*x
)*log(e**(a + b*x) - 1) + 27***(4*a + 4*b*x)*log(e**(a + b*x) + 1) + 50*e
**(3*a + 3*b*x) + 27***(2*a + 2*b*x)*log(e**(a + b*x) - 1) - 27***(2*a +
2*b*x)*log(e**(a + b*x) + 1) - 24***(a + b*x) - 9*log(e**(a + b*x) - 1)
+ 9*log(e**(a + b*x) + 1))/(6*b*(e**(6*a + 6*b*x) - 3***e**(4*a + 4*b*x) + 3
***e**(2*a + 2*b*x) - 1))
```

### 3.224 $\int e^x \tanh^2(2x) dx$

Optimal result	1645
Mathematica [C] (verified)	1645
Rubi [A] (verified)	1646
Maple [C] (verified)	1647
Fricas [B] (verification not implemented)	1648
Sympy [F]	1648
Maxima [A] (verification not implemented)	1649
Giac [A] (verification not implemented)	1649
Mupad [B] (verification not implemented)	1650
Reduce [B] (verification not implemented)	1650

#### Optimal result

Integrand size = 10, antiderivative size = 88

$$\int e^x \tanh^2(2x) dx = e^x + \frac{e^x}{1 + e^{4x}} + \frac{\arctan(1 - \sqrt{2}e^x)}{2\sqrt{2}} - \frac{\arctan(1 + \sqrt{2}e^x)}{2\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^x}{1+e^{2x}}\right)}{2\sqrt{2}}$$

output

```
exp(x)+exp(x)/(1+exp(4*x))-1/4*arctan(-1+2^(1/2)*exp(x))*2^(1/2)-1/4*arctan(1+2^(1/2)*exp(x))*2^(1/2)-1/4*arctanh(2^(1/2)*exp(x)/(1+exp(2*x)))*2^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.55

$$\int e^x \tanh^2(2x) dx = e^x + \frac{e^x}{1 + e^{4x}} + \frac{1}{4} \operatorname{RootSum}\left[1 + \#1^4 \&, \frac{x - \log(e^x - \#1)}{\#1^3} \&\right]$$

input

```
Integrate[E^x*Tanh[2*x]^2,x]
```

output  $E^x + E^x/(1 + E^{(4*x)}) + \text{RootSum}[1 + \#1^4 \& , (x - \text{Log}[E^x - \#1])/ \#1^3 \& ]/4$

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.28, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x \tanh^2(2x) dx \\ & \quad \downarrow 2720 \\ & \int \frac{(1 - e^{4x})^2}{(e^{4x} + 1)^2} de^x \\ & \quad \downarrow 915 \\ & \int \left( 1 - \frac{4e^{4x}}{(e^{4x} + 1)^2} \right) de^x \\ & \quad \downarrow 2009 \\ & \frac{\arctan(1 - \sqrt{2}e^x)}{2\sqrt{2}} - \frac{\arctan(\sqrt{2}e^x + 1)}{2\sqrt{2}} + e^x + \frac{e^x}{e^{4x} + 1} + \frac{\log(-\sqrt{2}e^x + e^{2x} + 1)}{4\sqrt{2}} - \\ & \quad \frac{\log(\sqrt{2}e^x + e^{2x} + 1)}{4\sqrt{2}} \end{aligned}$$

input  $\text{Int}[E^x * \text{Tanh}[2*x]^2, x]$

output  $E^x + E^x/(1 + E^{(4*x)}) + \text{ArcTan}[1 - \text{Sqrt}[2]*E^x]/(2*\text{Sqrt}[2]) - \text{ArcTan}[1 + \text{Sqrt}[2]*E^x]/(2*\text{Sqrt}[2]) + \text{Log}[1 - \text{Sqrt}[2]*E^x + E^{(2*x)}]/(4*\text{Sqrt}[2]) - \text{Log}[1 + \text{Sqrt}[2]*E^x + E^{(2*x)}]/(4*\text{Sqrt}[2])$

## Definitions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.60 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.40

method	result	size
risch	$e^x + \frac{e^x}{1+e^{4x}} + \left( \sum_{_R=\text{RootOf}(256\_Z^4+1)} \_R \ln(e^x - 4\_R) \right)$	35

input `int(exp(x)*tanh(2*x)^2,x,method=_RETURNVERBOSE)`

output `exp(x)+exp(x)/(1+exp(4*x))+sum(_R*ln(exp(x)-4*_R),_R=RootOf(256*_Z^4+1))`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 393 vs.  $2(62) = 124$ .

Time = 0.11 (sec) , antiderivative size = 393, normalized size of antiderivative = 4.47

$$\int e^x \tanh^2(2x) dx = \text{Too large to display}$$

input `integrate(exp(x)*tanh(2*x)^2,x, algorithm="fricas")`

output

```
1/8*(8*cosh(x)^5 + 80*cosh(x)^3*sinh(x)^2 + 80*cosh(x)^2*sinh(x)^3 + 40*cosh(x)*sinh(x)^4 + 8*sinh(x)^5 - 2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)^3*sinh(x) + 6*sqrt(2)*cosh(x)^2*sinh(x)^2 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + sqrt(2))*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) + 1) - 2*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)^3*sinh(x) + 6*sqrt(2)*cosh(x)^2*sinh(x)^2 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + sqrt(2))*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - 1) - (sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)^3*sinh(x) + 6*sqrt(2)*cosh(x)^2*sinh(x)^2 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + sqrt(2))*log((sqrt(2) + 2*cosh(x))/(cosh(x) - sinh(x))) + (sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)^3*sinh(x) + 6*sqrt(2)*cosh(x)^2*sinh(x)^2 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + sqrt(2))*log(-(sqrt(2) - 2*cosh(x))/(cosh(x) - sinh(x))) + 8*(5*cosh(x)^4 + 2)*sinh(x) + 16*cosh(x))/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 1)
```

**Sympy [F]**

$$\int e^x \tanh^2(2x) dx = \int e^x \tanh^2(2x) dx$$

input `integrate(exp(x)*tanh(2*x)**2,x)`

output `Integral(exp(x)*tanh(2*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

$$\int e^x \tanh^2(2x) dx = -\frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) - \frac{1}{4} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) - \frac{1}{8} \sqrt{2} \log\left(\sqrt{2}e^x + e^{(2x)} + 1\right) + \frac{1}{8} \sqrt{2} \log\left(-\sqrt{2}e^x + e^{(2x)} + 1\right) + \frac{e^x}{e^{(4x)} + 1} + e^x$$

input `integrate(exp(x)*tanh(2*x)^2,x, algorithm="maxima")`output `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x/(e^(4*x) + 1) + e^x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

$$\int e^x \tanh^2(2x) dx = -\frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) - \frac{1}{4} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) - \frac{1}{8} \sqrt{2} \log\left(\sqrt{2}e^x + e^{(2x)} + 1\right) + \frac{1}{8} \sqrt{2} \log\left(-\sqrt{2}e^x + e^{(2x)} + 1\right) + \frac{e^x}{e^{(4x)} + 1} + e^x$$

input `integrate(exp(x)*tanh(2*x)^2,x, algorithm="giac")`output `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x/(e^(4*x) + 1) + e^x`

**Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.98

$$\int e^x \tanh^2(2x) dx = e^x + \frac{e^x}{e^{4x} + 1} - \frac{\sqrt{2} \operatorname{atan}\left(\sqrt{2}\left(e^x - \frac{\sqrt{2}}{2}\right)\right)}{4} - \frac{\sqrt{2} \operatorname{atan}\left(\sqrt{2}\left(e^x + \frac{\sqrt{2}}{2}\right)\right)}{4} + \frac{\sqrt{2} \ln\left(\left(e^x - \frac{\sqrt{2}}{2}\right)^2 + \frac{1}{2}\right)}{8} - \frac{\sqrt{2} \ln\left(\left(e^x + \frac{\sqrt{2}}{2}\right)^2 + \frac{1}{2}\right)}{8}$$

input `int(tanh(2*x)^2*exp(x),x)`output `exp(x) + exp(x)/(exp(4*x) + 1) - (2^(1/2)*atan(2^(1/2)*(exp(x) - 2^(1/2)/2)))/4 - (2^(1/2)*atan(2^(1/2)*(exp(x) + 2^(1/2)/2)))/4 + (2^(1/2)*log((exp(x) - 2^(1/2)/2)^2 + 1/2))/8 - (2^(1/2)*log((exp(x) + 2^(1/2)/2)^2 + 1/2))/8`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 193, normalized size of antiderivative = 2.19

$$\int e^x \tanh^2(2x) dx = \frac{-2e^{4x}\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - 2\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - 2e^{4x}\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) - 2\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) + 8e^{5x} + e^x}{8}$$

input `int(exp(x)*tanh(2*x)^2,x)`output `( - 2*e**(4*x)*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - 2*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - 2*e**(4*x)*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) - 2*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) + 8*e**(5*x) + e**(4*x))*sqrt(2)*log(e**(2*x) - e**x*sqrt(2) + 1) - e**(4*x)*sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1) + 16*e**x + sqrt(2)*log(e**(2*x) - e**x*sqrt(2) + 1) - sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1))/(8*(e**(4*x) + 1))`

### 3.225 $\int e^x \tanh(2x) dx$

Optimal result	1651
Mathematica [A] (verified)	1651
Rubi [A] (verified)	1652
Maple [C] (verified)	1655
Fricas [A] (verification not implemented)	1656
Sympy [F]	1656
Maxima [A] (verification not implemented)	1657
Giac [A] (verification not implemented)	1657
Mupad [B] (verification not implemented)	1658
Reduce [B] (verification not implemented)	1658

#### Optimal result

Integrand size = 8, antiderivative size = 68

$$\int e^x \tanh(2x) dx = e^x + \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} - \frac{\arctan(1 + \sqrt{2}e^x)}{\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^x}{1+e^{2x}}\right)}{\sqrt{2}}$$

output

```
exp(x)-1/2*arctan(-1+2^(1/2)*exp(x))*2^(1/2)-1/2*arctan(1+2^(1/2)*exp(x))*
2^(1/2)-1/2*arctanh(2^(1/2)*exp(x)/(1+exp(2*x)))*2^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.40

$$\int e^x \tanh(2x) dx = e^x + \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} - \frac{\arctan(1 + \sqrt{2}e^x)}{\sqrt{2}} + \frac{\log(1 - \sqrt{2}e^x + e^{2x})}{2\sqrt{2}} - \frac{\log(1 + \sqrt{2}e^x + e^{2x})}{2\sqrt{2}}$$

input

```
Integrate[E^x*Tanh[2*x],x]
```

output

$$E^x + \text{ArcTan}[1 - \text{Sqrt}[2]*E^x]/\text{Sqrt}[2] - \text{ArcTan}[1 + \text{Sqrt}[2]*E^x]/\text{Sqrt}[2] + \text{Log}[1 - \text{Sqrt}[2]*E^x + E^{(2*x)}]/(2*\text{Sqrt}[2]) - \text{Log}[1 + \text{Sqrt}[2]*E^x + E^{(2*x)}]/(2*\text{Sqrt}[2])$$
**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.59, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {2720, 25, 913, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \tanh(2x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int -\frac{1 - e^{4x}}{e^{4x} + 1} de^x \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1 - e^{4x}}{1 + e^{4x}} de^x \\
 & \quad \downarrow \text{913} \\
 & e^x - 2 \int \frac{1}{1 + e^{4x}} de^x \\
 & \quad \downarrow \text{755} \\
 & e^x - 2 \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \int \frac{1 + e^{2x}}{1 + e^{4x}} de^x \right) \\
 & \quad \downarrow \text{1476} \\
 & e^x - 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{1 - \sqrt{2}e^x + e^{2x}} de^x + \frac{1}{2} \int \frac{1}{1 + \sqrt{2}e^x + e^{2x}} de^x \right) + \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x \right) \\
 & \quad \downarrow \text{1082} \\
 & e^x - 2 \left( \frac{1}{2} \left( \frac{\int \frac{1}{-1 - e^{2x}} d(1 - \sqrt{2}e^x)}{\sqrt{2}} - \frac{\int \frac{1}{-1 - e^{2x}} d(1 + \sqrt{2}e^x)}{\sqrt{2}} \right) + \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 217 \\
& e^x - 2 \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} dx + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) \\
& \downarrow 1479 \\
& 2 \left( \frac{1}{2} \left( -\frac{\int -\frac{\sqrt{2}-2e^x}{1-\sqrt{2}e^x+e^{2x}} dx}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(1+\sqrt{2}e^x)}{1+\sqrt{2}e^x+e^{2x}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) \\
& \downarrow 25 \\
& 2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}-2e^x}{1-\sqrt{2}e^x+e^{2x}} dx}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(1+\sqrt{2}e^x)}{1+\sqrt{2}e^x+e^{2x}} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) \\
& \downarrow 27 \\
& 2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}-2e^x}{1-\sqrt{2}e^x+e^{2x}} dx}{2\sqrt{2}} + \frac{1}{2} \int \frac{1 + \sqrt{2}e^x}{1 + \sqrt{2}e^x + e^{2x}} dx \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) \\
& \downarrow 1103 \\
& 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\sqrt{2}e^x + e^{2x} + 1)}{2\sqrt{2}} - \frac{\log(-\sqrt{2}e^x + e^{2x} + 1)}{2\sqrt{2}} \right) \right)
\end{aligned}$$

input

Int [E<sup>x</sup>\*Tanh [2\*x] , x]

output

$$E^x - 2 * ((- (\text{ArcTan}[1 - \text{Sqrt}[2] * E^x] / \text{Sqrt}[2]) + \text{ArcTan}[1 + \text{Sqrt}[2] * E^x] / \text{Sqrt}[2])) / 2 + (-1/2 * \text{Log}[1 - \text{Sqrt}[2] * E^x + E^{(2*x)}] / \text{Sqrt}[2] + \text{Log}[1 + \text{Sqrt}[2] * E^x + E^{(2*x)}] / (2 * \text{Sqrt}[2])) / 2$$

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 913 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.48 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.35

method	result	size
risch	$e^x + \left( \sum_{_R=\text{RootOf}(16\_Z^4+1)} \_R \ln(e^x - 2\_R) \right)$	24

input `int(exp(x)*tanh(2*x), x, method=_RETURNVERBOSE)`

output `exp(x)+sum(_R*ln(exp(x)-2*_R), _R=RootOf(16*_Z^4+1))`



**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.38

$$\int e^x \tanh(2x) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) + 1 \right) - \frac{1}{2} \sqrt{2} \arctan \left( \sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) - 1 \right) - \frac{1}{4} \sqrt{2} \log \left( \frac{\sqrt{2} + 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) + \frac{1}{4} \sqrt{2} \log \left( -\frac{\sqrt{2} - 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) + \cosh(x) + \sinh(x)$$

input `integrate(exp(x)*tanh(2*x),x, algorithm="fricas")`output `-1/2*sqrt(2)*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) + 1) - 1/2*sqrt(2)*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - 1) - 1/4*sqrt(2)*log((sqrt(2) + 2*cosh(x))/(cosh(x) - sinh(x))) + 1/4*sqrt(2)*log(-(sqrt(2) - 2*cosh(x))/(cosh(x) - sinh(x))) + cosh(x) + sinh(x)`**Sympy [F]**

$$\int e^x \tanh(2x) dx = \int e^x \tanh(2x) dx$$

input `integrate(exp(x)*tanh(2*x),x)`output `Integral(exp(x)*tanh(2*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.15

$$\int e^x \tanh(2x) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2} + 2e^x) \right) \\ - \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2} - 2e^x) \right) \\ - \frac{1}{4} \sqrt{2} \log \left( \sqrt{2}e^x + e^{(2x)} + 1 \right) + \frac{1}{4} \sqrt{2} \log \left( -\sqrt{2}e^x + e^{(2x)} + 1 \right) + e^x$$

input `integrate(exp(x)*tanh(2*x),x, algorithm="maxima")`output `-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/4*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/4*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x`**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.15

$$\int e^x \tanh(2x) dx = -\frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2} + 2e^x) \right) \\ - \frac{1}{2} \sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2} - 2e^x) \right) \\ - \frac{1}{4} \sqrt{2} \log \left( \sqrt{2}e^x + e^{(2x)} + 1 \right) + \frac{1}{4} \sqrt{2} \log \left( -\sqrt{2}e^x + e^{(2x)} + 1 \right) + e^x$$

input `integrate(exp(x)*tanh(2*x),x, algorithm="giac")`output `-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/4*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/4*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x`

**Mupad [B] (verification not implemented)**

Time = 2.22 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.19

$$\int e^x \tanh(2x) dx = e^x - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x - \sqrt{2})}{2}\right)}{2} + \frac{\sqrt{2} \ln\left((2e^x - \sqrt{2})^2 + 2\right)}{4}$$

$$- \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x + \sqrt{2})}{2}\right)}{2} - \frac{\sqrt{2} \ln\left((2e^x + \sqrt{2})^2 + 2\right)}{4}$$

input `int(tanh(2*x)*exp(x), x)`output `exp(x) - (2^(1/2)*atan((2^(1/2)*(2*exp(x) - 2^(1/2)))/2))/2 + (2^(1/2)*log((2*exp(x) - 2^(1/2))^2 + 2))/4 - (2^(1/2)*atan((2^(1/2)*(2*exp(x) + 2^(1/2)))/2))/2 - (2^(1/2)*log((2*exp(x) + 2^(1/2))^2 + 2))/4`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.16

$$\int e^x \tanh(2x) dx = -\frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right)}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{2} + e^x$$

$$+ \frac{\sqrt{2} \log(e^{2x} - e^x \sqrt{2} + 1)}{4} - \frac{\sqrt{2} \log(e^{2x} + e^x \sqrt{2} + 1)}{4}$$

input `int(exp(x)*tanh(2*x), x)`output `( - 2*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - 2*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) + 4*e**x + sqrt(2)*log(e**(2*x) - e**x*sqrt(2) + 1) - sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1))/4`

### 3.226 $\int e^x \coth(2x) dx$

Optimal result	1659
Mathematica [A] (verified)	1659
Rubi [A] (verified)	1660
Maple [C] (verified)	1662
Fricas [B] (verification not implemented)	1662
Sympy [F]	1663
Maxima [A] (verification not implemented)	1663
Giac [A] (verification not implemented)	1663
Mupad [B] (verification not implemented)	1664
Reduce [B] (verification not implemented)	1664

#### Optimal result

Integrand size = 8, antiderivative size = 16

$$\int e^x \coth(2x) dx = e^x - \arctan(e^x) - \operatorname{arctanh}(e^x)$$

output `exp(x)-arctan(exp(x))-arctanh(exp(x))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int e^x \coth(2x) dx = e^x - \arctan(e^x) - \operatorname{arctanh}(e^x)$$

input `Integrate[E^x*Coth[2*x],x]`

output `E^x - ArcTan[E^x] - ArcTanh[E^x]`

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {2720, 25, 913, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \coth(2x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int -\frac{e^{4x} + 1}{1 - e^{4x}} de^x \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1 + e^{4x}}{1 - e^{4x}} de^x \\
 & \quad \downarrow \text{913} \\
 & e^x - 2 \int \frac{1}{1 - e^{4x}} de^x \\
 & \quad \downarrow \text{756} \\
 & e^x - 2 \left( \frac{1}{2} \int \frac{1}{1 - e^{2x}} de^x + \frac{1}{2} \int \frac{1}{1 + e^{2x}} de^x \right) \\
 & \quad \downarrow \text{216} \\
 & e^x - 2 \left( \frac{1}{2} \int \frac{1}{1 - e^{2x}} de^x + \frac{\arctan(e^x)}{2} \right) \\
 & \quad \downarrow \text{219} \\
 & e^x - 2 \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right)
 \end{aligned}$$

input `Int [E^x*Coth [2*x] , x]`

output `E^x - 2*(ArcTan [E^x]/2 + ArcTanh [E^x]/2)`

## Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`
- rule 913 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.25

method	result	size
risch	$e^x - \frac{\ln(e^x+1)}{2} + \frac{i \ln(e^x-i)}{2} - \frac{i \ln(e^x+i)}{2} + \frac{\ln(e^x-1)}{2}$	36

input `int(exp(x)*coth(2*x),x,method=_RETURNVERBOSE)`

output `exp(x)-1/2*ln(exp(x)+1)+1/2*I*ln(exp(x)-I)-1/2*I*ln(exp(x)+I)+1/2*ln(exp(x)-1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(13) = 26.

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int e^x \coth(2x) dx = -\arctan(\cosh(x) + \sinh(x)) + \cosh(x) - \frac{1}{2} \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{2} \log(\cosh(x) + \sinh(x) - 1) + \sinh(x)$$

input `integrate(exp(x)*coth(2*x),x, algorithm="fricas")`

output `-arctan(cosh(x) + sinh(x)) + cosh(x) - 1/2*log(cosh(x) + sinh(x) + 1) + 1/2*log(cosh(x) + sinh(x) - 1) + sinh(x)`

**Sympy [F]**

$$\int e^x \coth(2x) dx = \int e^x \coth(2x) dx$$

input `integrate(exp(x)*coth(2*x),x)`

output `Integral(exp(x)*coth(2*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int e^x \coth(2x) dx = -\arctan(e^x) + e^x - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

input `integrate(exp(x)*coth(2*x),x, algorithm="maxima")`

output `-arctan(e^x) + e^x - 1/2*log(e^x + 1) + 1/2*log(e^x - 1)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int e^x \coth(2x) dx = -\arctan(e^x) + e^x - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(2*x),x, algorithm="giac")`

output `-arctan(e^x) + e^x - 1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`



**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int e^x \coth(2x) dx = \frac{\ln(2 - 2e^x)}{2} - \frac{\ln(-2e^x - 2)}{2} - \operatorname{atan}(e^x) + e^x$$

input `int(coth(2*x)*exp(x), x)`output `log(2 - 2*exp(x))/2 - log(- 2*exp(x) - 2)/2 - atan(exp(x)) + exp(x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int e^x \coth(2x) dx = -\operatorname{atan}(e^x) + e^x + \frac{\log(e^x - 1)}{2} - \frac{\log(e^x + 1)}{2}$$

input `int(exp(x)*coth(2*x), x)`output `( - 2*atan(e**x) + 2*e**x + log(e**x - 1) - log(e**x + 1))/2`

### 3.227 $\int e^x \coth^2(2x) dx$

Optimal result	1665
Mathematica [C] (verified)	1665
Rubi [A] (verified)	1666
Maple [C] (verified)	1667
Fricas [B] (verification not implemented)	1668
Sympy [F]	1668
Maxima [A] (verification not implemented)	1669
Giac [A] (verification not implemented)	1669
Mupad [B] (verification not implemented)	1669
Reduce [B] (verification not implemented)	1670

#### Optimal result

Integrand size = 10, antiderivative size = 35

$$\int e^x \coth^2(2x) dx = e^x + \frac{e^x}{1 - e^{4x}} - \frac{\arctan(e^x)}{2} - \frac{\operatorname{arctanh}(e^x)}{2}$$

output `exp(x)+exp(x)/(1-exp(4*x))-1/2*arctan(exp(x))-1/2*arctanh(exp(x))`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.12 (sec) , antiderivative size = 113, normalized size of antiderivative = 3.23

$$\int e^x \coth^2(2x) dx = \frac{1}{640} e^{-7x} \left( -3645 - 6769e^{4x} - 1483e^{8x} + 681e^{12x} + 5(729 + 1208e^{4x} + 102e^{8x} - 248e^{12x} + e^{16x}) \operatorname{Hypergeometric2F1} \left( \frac{1}{4}, 1, \frac{5}{4}, e^{4x} \right) \right) + \frac{16}{585} e^{5x} (1 + e^{4x})^2 {}_4F_3 \left( \frac{5}{4}, 2, 2, 2; 1, 1, \frac{17}{4}; e^{4x} \right)$$

input `Integrate[E^x*Coth[2*x]^2,x]`

output

$$\begin{aligned} & (-3645 - 6769E^{(4x)} - 1483E^{(8x)} + 681E^{(12x)} + 5(729 + 1208E^{(4x)} \\ & + 102E^{(8x)} - 248E^{(12x)} + E^{(16x)}) \text{Hypergeometric2F1}[1/4, 1, 5/4, \\ & E^{(4x)}] / (640E^{(7x)}) + (16E^{(5x)}(1 + E^{(4x)})^2 \text{HypergeometricPFQ}[\{5 \\ & /4, 2, 2, 2\}, \{1, 1, 17/4\}, E^{(4x)}]) / 585 \end{aligned}$$
**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x \coth^2(2x) dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{(e^{4x} + 1)^2}{(1 - e^{4x})^2} de^x \\ & \quad \downarrow \text{915} \\ & \int \left( \frac{4e^{4x}}{(1 - e^{4x})^2} + 1 \right) de^x \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{2} \arctan(e^x) - \frac{\operatorname{arctanh}(e^x)}{2} + e^x + \frac{e^x}{1 - e^{4x}} \end{aligned}$$

input

$$\text{Int}[E^x \text{Coth}[2x]^2, x]$$

output

$$E^x + E^x / (1 - E^{(4x)}) - \text{ArcTan}[E^x] / 2 - \text{ArcTanh}[E^x] / 2$$

## Definitions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.37

method	result	size
risch	$e^x - \frac{e^x}{e^{4x}-1} + \frac{i \ln(e^x-i)}{4} - \frac{i \ln(e^x+i)}{4} - \frac{\ln(e^x+1)}{4} + \frac{\ln(e^x-1)}{4}$	48

input `int(exp(x)*coth(2*x)^2,x,method=_RETURNVERBOSE)`

output `exp(x)-exp(x)/(exp(4*x)-1)+1/4*I*ln(exp(x)-I)-1/4*I*ln(exp(x)+I)-1/4*ln(ex
p(x)+1)+1/4*ln(exp(x)-1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 230 vs.  $2(25) = 50$ .

Time = 0.10 (sec) , antiderivative size = 230, normalized size of antiderivative = 6.57

$$\int e^x \coth^2(2x) dx$$

$$= \frac{4 \cosh(x)^5 + 40 \cosh(x)^3 \sinh(x)^2 + 40 \cosh(x)^2 \sinh(x)^3 + 20 \cosh(x) \sinh(x)^4 + 4 \sinh(x)^5 - 2(\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 - 1) \arctan(\cosh(x) + \sinh(x)) - (\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 - 1) \log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 - 1) \log(\cosh(x) + \sinh(x) - 1) + 4(5 \cosh(x)^4 - 2) \sinh(x) - 8 \cosh(x)}{\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 - 1}$$

input `integrate(exp(x)*coth(2*x)^2,x, algorithm="fricas")`

output `1/4*(4*cosh(x)^5 + 40*cosh(x)^3*sinh(x)^2 + 40*cosh(x)^2*sinh(x)^3 + 20*cosh(x)*sinh(x)^4 + 4*sinh(x)^5 - 2*(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 - 1)*arctan(cosh(x) + sinh(x)) - (cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 - 1)*log(cosh(x) + sinh(x) + 1) + (cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 - 1)*log(cosh(x) + sinh(x) - 1) + 4*(5*cosh(x)^4 - 2)*sinh(x) - 8*cosh(x))/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 - 1)`

**Sympy [F]**

$$\int e^x \coth^2(2x) dx = \int e^x \coth^2(2x) dx$$

input `integrate(exp(x)*coth(2*x)**2,x)`

output `Integral(exp(x)*coth(2*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int e^x \coth^2(2x) dx = -\frac{e^x}{e^{4x} - 1} - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x + 1) + \frac{1}{4} \log(e^x - 1)$$

input `integrate(exp(x)*coth(2*x)^2,x, algorithm="maxima")`output `-e^x/(e^(4*x) - 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(e^x - 1)`**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int e^x \coth^2(2x) dx = -\frac{e^x}{e^{4x} - 1} - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x + 1) + \frac{1}{4} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(2*x)^2,x, algorithm="giac")`output `-e^x/(e^(4*x) - 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(abs(e^x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int e^x \coth^2(2x) dx = \frac{\ln(1 - e^x)}{4} - \frac{\ln(-e^x - 1)}{4} - \frac{\operatorname{atan}(e^x)}{2} + e^x - \frac{e^x}{e^{4x} - 1}$$

input `int(coth(2*x)^2*exp(x), x)`output `log(1 - exp(x))/4 - log(- exp(x) - 1)/4 - atan(exp(x))/2 + exp(x) - exp(x)/(exp(4*x) - 1)`

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 81, normalized size of antiderivative = 2.31

$$\int e^x \coth^2(2x) dx$$

$$= \frac{-2e^{4x} \operatorname{atan}(e^x) + 2\operatorname{atan}(e^x) + 4e^{5x} + e^{4x} \log(e^x - 1) - e^{4x} \log(e^x + 1) - 8e^x - \log(e^x - 1) + \log(e^x + 1)}{4e^{4x} - 4}$$

input `int(exp(x)*coth(2*x)^2,x)`output `( - 2*e**(4*x)*atan(e**x) + 2*atan(e**x) + 4*e**(5*x) + e**(4*x)*log(e**x - 1) - e**(4*x)*log(e**x + 1) - 8*e**x - log(e**x - 1) + log(e**x + 1))/(4 * (e**(4*x) - 1))`

### 3.228 $\int e^x \coth^4(2x) dx$

Optimal result . . . . .	1671
Mathematica [A] (verified) . . . . .	1671
Rubi [A] (verified) . . . . .	1672
Maple [C] (verified) . . . . .	1673
Fricas [B] (verification not implemented) . . . . .	1674
Sympy [F] . . . . .	1675
Maxima [A] (verification not implemented) . . . . .	1675
Giac [A] (verification not implemented) . . . . .	1675
Mupad [B] (verification not implemented) . . . . .	1676
Reduce [B] (verification not implemented) . . . . .	1676

#### Optimal result

Integrand size = 10, antiderivative size = 74

$$\int e^x \coth^4(2x) dx = e^x + \frac{4e^x}{3(1 - e^{4x})^3} - \frac{13e^x}{6(1 - e^{4x})^2} + \frac{53e^x}{24(1 - e^{4x})} - \frac{11 \arctan(e^x)}{16} - \frac{11 \operatorname{arctanh}(e^x)}{16}$$

output

```
exp(x)+4/3*exp(x)/(1-exp(4*x))^3-13/6*exp(x)/(1-exp(4*x))^2+53*exp(x)/(24-24*exp(4*x))-11/16*arctan(exp(x))-11/16*arctanh(exp(x))
```

#### Mathematica [A] (verified)

Time = 11.06 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.03

$$\int e^x \coth^4(2x) dx = \frac{1}{96} \left( 96e^x - \frac{128e^x}{(-1 + e^{4x})^3} - \frac{208e^x}{(-1 + e^{4x})^2} - \frac{212e^x}{-1 + e^{4x}} - 66 \arctan(e^x) + 33 \log(1 - e^x) - 33 \log(1 + e^x) \right)$$

input

```
Integrate[E^x*Coth[2*x]^4,x]
```



output

$$\frac{(96E^x - (128E^x)/(-1 + E^{(4x)})^3 - (208E^x)/(-1 + E^{(4x)})^2 - (212E^x)/(-1 + E^{(4x)}) - 66\text{ArcTan}[E^x] + 33\text{Log}[1 - E^x] - 33\text{Log}[1 + E^x])/9}{6}$$

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^x \coth^4(2x) dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{(e^{4x} + 1)^4}{(1 - e^{4x})^4} de^x \\ & \quad \downarrow \text{915} \\ & \int \left( \frac{8e^{4x}(e^{8x} + 1)}{(1 - e^{4x})^4} + 1 \right) de^x \\ & \quad \downarrow \text{2009} \\ & -\frac{11}{16} \arctan(e^x) - \frac{11 \operatorname{arctanh}(e^x)}{16} + e^x + \frac{53e^x}{24(1 - e^{4x})} - \frac{13e^x}{6(1 - e^{4x})^2} + \frac{4e^x}{3(1 - e^{4x})^3} \end{aligned}$$

input

Int [E^x\*Coth[2\*x]^4, x]

output

$$\frac{E^x + (4E^x)/(3*(1 - E^{(4x)})^3) - (13E^x)/(6*(1 - E^{(4x)})^2) + (53E^x)/(24*(1 - E^{(4x)})) - (11*\text{ArcTan}[E^x])/16 - (11*\text{ArcTanh}[E^x])/16}{1}$$

## Definitions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

method	result	size
risch	$e^x - \frac{e^x(53e^{8x} - 54e^{4x} + 33)}{24(e^{4x} - 1)^3} + \frac{11i \ln(e^x - i)}{32} - \frac{11i \ln(e^x + i)}{32} - \frac{11 \ln(e^x + 1)}{32} + \frac{11 \ln(e^x - 1)}{32}$	62

input `int(exp(x)*coth(2*x)^4,x,method=_RETURNVERBOSE)`

output `exp(x)-1/24*exp(x)*(53*exp(8*x)-54*exp(4*x)+33)/(exp(4*x)-1)^3+11/32*I*ln(
exp(x)-I)-11/32*I*ln(exp(x)+I)-11/32*ln(exp(x)+1)+11/32*ln(exp(x)-1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1097 vs.  $2(49) = 98$ .

Time = 0.11 (sec) , antiderivative size = 1097, normalized size of antiderivative = 14.82

$$\int e^x \coth^4(2x) dx = \text{Too large to display}$$

input `integrate(exp(x)*coth(2*x)^4,x, algorithm="fricas")`

output

```

1/96*(96*cosh(x)^13 + 27456*cosh(x)^3*sinh(x)^10 + 7488*cosh(x)^2*sinh(x)^
11 + 1248*cosh(x)*sinh(x)^12 + 96*sinh(x)^13 + 20*(3432*cosh(x)^4 - 25)*si
nh(x)^9 - 500*cosh(x)^9 + 36*(3432*cosh(x)^5 - 125*cosh(x))*sinh(x)^8 + 14
4*(1144*cosh(x)^6 - 125*cosh(x)^2)*sinh(x)^7 + 48*(3432*cosh(x)^7 - 875*co
sh(x)^3)*sinh(x)^6 + 72*(1716*cosh(x)^8 - 875*cosh(x)^4 + 7)*sinh(x)^5 + 5
04*cosh(x)^5 + 120*(572*cosh(x)^9 - 525*cosh(x)^5 + 21*cosh(x))*sinh(x)^4
+ 48*(572*cosh(x)^10 - 875*cosh(x)^6 + 105*cosh(x)^2)*sinh(x)^3 + 144*(52*
cosh(x)^11 - 125*cosh(x)^7 + 35*cosh(x)^3)*sinh(x)^2 - 66*(cosh(x)^12 + 22
0*cosh(x)^3*sinh(x)^9 + 66*cosh(x)^2*sinh(x)^10 + 12*cosh(x)*sinh(x)^11 +
sinh(x)^12 + 3*(165*cosh(x)^4 - 1)*sinh(x)^8 - 3*cosh(x)^8 + 24*(33*cosh(x)
)^5 - cosh(x))*sinh(x)^7 + 84*(11*cosh(x)^6 - cosh(x)^2)*sinh(x)^6 + 24*(3
3*cosh(x)^7 - 7*cosh(x)^3)*sinh(x)^5 + 3*(165*cosh(x)^8 - 70*cosh(x)^4 + 1
)*sinh(x)^4 + 3*cosh(x)^4 + 4*(55*cosh(x)^9 - 42*cosh(x)^5 + 3*cosh(x))*si
nh(x)^3 + 6*(11*cosh(x)^10 - 14*cosh(x)^6 + 3*cosh(x)^2)*sinh(x)^2 + 12*(c
osh(x)^11 - 2*cosh(x)^7 + cosh(x)^3)*sinh(x) - 1)*arctan(cosh(x) + sinh(x)
) - 33*(cosh(x)^12 + 220*cosh(x)^3*sinh(x)^9 + 66*cosh(x)^2*sinh(x)^10 + 1
2*cosh(x)*sinh(x)^11 + sinh(x)^12 + 3*(165*cosh(x)^4 - 1)*sinh(x)^8 - 3*co
sh(x)^8 + 24*(33*cosh(x)^5 - cosh(x))*sinh(x)^7 + 84*(11*cosh(x)^6 - cosh(x)
)^2)*sinh(x)^6 + 24*(33*cosh(x)^7 - 7*cosh(x)^3)*sinh(x)^5 + 3*(165*cosh(
x)^8 - 70*cosh(x)^4 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(55*cosh(x)^9 - 42...
```

**Sympy [F]**

$$\int e^x \coth^4(2x) dx = \int e^x \coth^4(2x) dx$$

input `integrate(exp(x)*coth(2*x)**4,x)`

output `Integral(exp(x)*coth(2*x)**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.82

$$\int e^x \coth^4(2x) dx = -\frac{53 e^{(9x)} - 54 e^{(5x)} + 33 e^x}{24 (e^{(12x)} - 3 e^{(8x)} + 3 e^{(4x)} - 1)} - \frac{11}{16} \arctan(e^x) + e^x - \frac{11}{32} \log(e^x + 1) + \frac{11}{32} \log(e^x - 1)$$

input `integrate(exp(x)*coth(2*x)^4,x, algorithm="maxima")`

output `-1/24*(53*e^(9*x) - 54*e^(5*x) + 33*e^x)/(e^(12*x) - 3*e^(8*x) + 3*e^(4*x) - 1) - 11/16*arctan(e^x) + e^x - 11/32*log(e^x + 1) + 11/32*log(e^x - 1)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.68

$$\int e^x \coth^4(2x) dx = -\frac{53 e^{(9x)} - 54 e^{(5x)} + 33 e^x}{24 (e^{(4x)} - 1)^3} - \frac{11}{16} \arctan(e^x) + e^x - \frac{11}{32} \log(e^x + 1) + \frac{11}{32} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(2*x)^4,x, algorithm="giac")`

output

```
-1/24*(53*e^(9*x) - 54*e^(5*x) + 33*e^x)/(e^(4*x) - 1)^3 - 11/16*arctan(e^x) + e^x - 11/32*log(e^x + 1) + 11/32*log(abs(e^x - 1))
```

**Mupad [B] (verification not implemented)**

Time = 2.57 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.49

$$\int e^x \coth^4(2x) dx = \frac{11 \ln\left(\frac{11}{8} - \frac{11e^x}{8}\right)}{32} - \frac{11 \ln\left(-\frac{11e^x}{8} - \frac{11}{8}\right)}{32} + e^x - \frac{37e^x}{24(e^{4x} - 1)} - \frac{\frac{2e^{9x}}{3} + \frac{2e^x}{3}}{3e^{4x} - 3e^{8x} + e^{12x} - 1} - \frac{5e^x}{6(e^{8x} - 2e^{4x} + 1)} - \frac{\ln\left(-\frac{11e^x}{8} - \frac{11i}{8}\right) 11i}{32} + \frac{\ln\left(-\frac{11e^x}{8} + \frac{11i}{8}\right) 11i}{32}$$

input

```
int(coth(2*x)^4*exp(x), x)
```

output

```
(11*log(11/8 - (11*exp(x))/8))/32 - (11*log(- (11*exp(x))/8 - 11/8))/32 - (log(- (11*exp(x))/8 - 11i/8)*11i)/32 + (log(11i/8 - (11*exp(x))/8)*11i)/32 + exp(x) - (37*exp(x))/(24*(exp(4*x) - 1)) - ((2*exp(9*x))/3 + (2*exp(x))/3)/(3*exp(4*x) - 3*exp(8*x) + exp(12*x) - 1) - (5*exp(x))/(6*(exp(8*x) - 2*exp(4*x) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.51

$$\int e^x \coth^4(2x) dx = \frac{-66e^{12x} \operatorname{atan}(e^x) + 198e^{8x} \operatorname{atan}(e^x) - 198e^{4x} \operatorname{atan}(e^x) + 66 \operatorname{atan}(e^x) + 96e^{13x} + 33e^{12x} \log(e^x - 1) - 33e^{11x}}{24(e^{4x} - 1)}$$

input

```
int(exp(x)*coth(2*x)^4, x)
```

output

```
( - 66*e**(12*x)*atan(e**x) + 198*e**(8*x)*atan(e**x) - 198*e**(4*x)*atan(e**x) + 66*atan(e**x) + 96*e**(13*x) + 33*e**(12*x)*log(e**x - 1) - 33*e**(12*x)*log(e**x + 1) - 500*e**(9*x) - 99*e**(8*x)*log(e**x - 1) + 99*e**(8*x)*log(e**x + 1) + 504*e**(5*x) + 99*e**(4*x)*log(e**x - 1) - 99*e**(4*x)*log(e**x + 1) - 228*e**x - 33*log(e**x - 1) + 33*log(e**x + 1))/(96*(e**(12*x) - 3*e**(8*x) + 3*e**(4*x) - 1))
```

### 3.229 $\int e^x \tanh^2(3x) dx$

Optimal result	1678
Mathematica [C] (verified)	1678
Rubi [A] (verified)	1679
Maple [C] (verified)	1680
Fricas [B] (verification not implemented)	1681
Sympy [F]	1682
Maxima [A] (verification not implemented)	1682
Giac [A] (verification not implemented)	1682
Mupad [B] (verification not implemented)	1683
Reduce [B] (verification not implemented)	1683

#### Optimal result

Integrand size = 10, antiderivative size = 88

$$\int e^x \tanh^2(3x) dx = e^x + \frac{2e^x}{3(1 + e^{6x})} - \frac{2 \arctan(e^x)}{9} + \frac{1}{9} \arctan(\sqrt{3} - 2e^x) - \frac{1}{9} \arctan(\sqrt{3} + 2e^x) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}e^x}{1+e^{2x}}\right)}{3\sqrt{3}}$$

output

```
exp(x)+2*exp(x)/(3+3*exp(6*x))-2/9*arctan(exp(x))-1/9*arctan(-3^(1/2)+2*exp(x))-1/9*arctan(3^(1/2)+2*exp(x))-1/9*arctanh(3^(1/2)*exp(x)/(1+exp(2*x)))*3^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.10

$$\int e^x \tanh^2(3x) dx = e^x + \frac{2e^x}{3(1 + e^{6x})} - \frac{2 \arctan(e^x)}{9} - \frac{1}{9} \operatorname{RootSum}\left[1 - \#1^2 + \#1^4 \&, \frac{-2x + 2 \log(e^x - \#1) + x\#1^2 - \log(e^x - \#1)\#1^2}{-\#1 + 2\#1^3} \&\right]$$

input `Integrate[E^x*Tanh[3*x]^2,x]`

output `E^x + (2*E^x)/(3*(1 + E^(6*x))) - (2*ArcTan[E^x])/9 - RootSum[1 - #1^2 + #1^4 & , (-2*x + 2*Log[E^x - #1] + x*#1^2 - Log[E^x - #1]*#1^2)/(-#1 + 2*#1^3) & ]/9`

### Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.28, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \tanh^2(3x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int \frac{(1 - e^{6x})^2}{(e^{6x} + 1)^2} de^x \\
 & \quad \downarrow \text{915} \\
 & \int \left( 1 - \frac{4e^{6x}}{(e^{6x} + 1)^2} \right) de^x \\
 & \quad \downarrow \text{2009} \\
 & -\frac{2}{9} \arctan(e^x) + \frac{1}{9} \arctan(\sqrt{3} - 2e^x) - \frac{1}{9} \arctan(2e^x + \sqrt{3}) + e^x + \frac{2e^x}{3(e^{6x} + 1)} + \\
 & \quad \frac{\log(-\sqrt{3}e^x + e^{2x} + 1)}{6\sqrt{3}} - \frac{\log(\sqrt{3}e^x + e^{2x} + 1)}{6\sqrt{3}}
 \end{aligned}$$

input `Int[E^x*Tanh[3*x]^2,x]`



output

```
E^x + (2*E^x)/(3*(1 + E^(6*x))) - (2*ArcTan[E^x])/9 + ArcTan[Sqrt[3] - 2*E^x]/9 - ArcTan[Sqrt[3] + 2*E^x]/9 + Log[1 - Sqrt[3]*E^x + E^(2*x)]/(6*Sqrt[3]) - Log[1 + Sqrt[3]*E^x + E^(2*x)]/(6*Sqrt[3])
```

**Defintions of rubi rules used**

rule 915

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2720

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.63 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.67

method	result	size
risch	$e^x + \frac{2e^x}{3(e^{6x}+1)} + \left( \sum_{_R=\text{RootOf}(6561_Z^4-81_Z^2+1)} -R \ln(e^x - 9_R) \right) + \frac{i \ln(e^x - i)}{9} - \frac{i \ln(e^x + i)}{9}$	59

input

```
int(exp(x)*tanh(3*x)^2,x,method=_RETURNVERBOSE)
```

output

```
exp(x)+2/3*exp(x)/(exp(6*x)+1)+sum(_R*ln(exp(x)-9*_R),_R=RootOf(6561*_Z^4-81*_Z^2+1))+1/9*I*ln(exp(x)-I)-1/9*I*ln(exp(x)+I)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 549 vs.  $2(64) = 128$ .

Time = 0.09 (sec) , antiderivative size = 549, normalized size of antiderivative = 6.24

$$\int e^x \tanh^2(3x) dx = \text{Too large to display}$$

input `integrate(exp(x)*tanh(3*x)^2,x, algorithm="fricas")`

output

```
1/18*(18*cosh(x)^7 + 378*cosh(x)^5*sinh(x)^2 + 630*cosh(x)^4*sinh(x)^3 + 6
30*cosh(x)^3*sinh(x)^4 + 378*cosh(x)^2*sinh(x)^5 + 126*cosh(x)*sinh(x)^6 +
18*sinh(x)^7 - 2*(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^
2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5
+ sinh(x)^6 + 1)*arctan(sqrt(3) + 2*cosh(x) + 2*sinh(x)) - 2*(cosh(x)^6 +
6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15
*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 1)*arctan(-sqrt(3)
) + 2*cosh(x) + 2*sinh(x)) - 4*(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(
x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(
x)*sinh(x)^5 + sinh(x)^6 + 1)*arctan(cosh(x) + sinh(x)) - (sqrt(3)*cosh(x)
^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*sqrt(3)*cosh(x)^4*sinh(x)^2 + 20*sq
rt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*cosh(x)^2*sinh(x)^4 + 6*sqrt(3)*co
sh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 + sqrt(3))*log((sqrt(3) + 2*cosh(x))/(co
sh(x) - sinh(x))) + (sqrt(3)*cosh(x)^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*
sqrt(3)*cosh(x)^4*sinh(x)^2 + 20*sqrt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*
cosh(x)^2*sinh(x)^4 + 6*sqrt(3)*cosh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 + sq
rt(3))*log(-(sqrt(3) - 2*cosh(x))/(cosh(x) - sinh(x))) + 6*(21*cosh(x)^6 +
5)*sinh(x) + 30*cosh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*
sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*si
nh(x)^5 + sinh(x)^6 + 1)
```

**Sympy [F]**

$$\int e^x \tanh^2(3x) dx = \int e^x \tanh^2(3x) dx$$

input `integrate(exp(x)*tanh(3*x)**2,x)`

output `Integral(exp(x)*tanh(3*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

$$\begin{aligned} \int e^x \tanh^2(3x) dx = & -\frac{1}{18} \sqrt{3} \log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{18} \sqrt{3} \log(-\sqrt{3}e^x + e^{2x} + 1) \\ & + \frac{2e^x}{3(e^{6x} + 1)} - \frac{1}{9} \arctan(\sqrt{3} + 2e^x) \\ & - \frac{1}{9} \arctan(-\sqrt{3} + 2e^x) - \frac{2}{9} \arctan(e^x) + e^x \end{aligned}$$

input `integrate(exp(x)*tanh(3*x)^2,x, algorithm="maxima")`

output `-1/18*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/18*sqrt(3)*log(-sqrt(3)*e^x + e^(2*x) + 1) + 2/3*e^x/(e^(6*x) + 1) - 1/9*arctan(sqrt(3) + 2*e^x) - 1/9*arctan(-sqrt(3) + 2*e^x) - 2/9*arctan(e^x) + e^x`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

$$\begin{aligned} \int e^x \tanh^2(3x) dx = & -\frac{1}{18} \sqrt{3} \log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{18} \sqrt{3} \log(-\sqrt{3}e^x + e^{2x} + 1) \\ & + \frac{2e^x}{3(e^{6x} + 1)} - \frac{1}{9} \arctan(\sqrt{3} + 2e^x) \\ & - \frac{1}{9} \arctan(-\sqrt{3} + 2e^x) - \frac{2}{9} \arctan(e^x) + e^x \end{aligned}$$

input `integrate(exp(x)*tanh(3*x)^2,x, algorithm="giac")`

output 
$$-1/18*\sqrt{3}*\log(\sqrt{3}*e^x + e^{2*x} + 1) + 1/18*\sqrt{3}*\log(-\sqrt{3}*e^x + e^{2*x} + 1) + 2/3*e^x/(e^{6*x} + 1) - 1/9*\arctan(\sqrt{3} + 2*e^x) - 1/9*\arctan(-\sqrt{3} + 2*e^x) - 2/9*\arctan(e^x) + e^x$$

### Mupad [B] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.98

$$\int e^x \tanh^2(3x) dx = e^x - \frac{\operatorname{atan}(2e^x + \sqrt{3})}{9} - \frac{\operatorname{atan}(2e^x - \sqrt{3})}{9} - \frac{2\operatorname{atan}(e^x)}{9} + \frac{2e^x}{3(e^{6x} + 1)} + \frac{\sqrt{3} \ln\left(\left(\frac{2e^x}{3} - \frac{\sqrt{3}}{3}\right)^2 + \frac{1}{9}\right)}{18} - \frac{\sqrt{3} \ln\left(\left(\frac{2e^x}{3} + \frac{\sqrt{3}}{3}\right)^2 + \frac{1}{9}\right)}{18}$$

input `int(tanh(3*x)^2*exp(x), x)`

output 
$$\frac{\exp(x) - \operatorname{atan}(2*\exp(x) + 3^{(1/2)})}{9} - \frac{\operatorname{atan}(2*\exp(x) - 3^{(1/2)})}{9} - \frac{(2*\operatorname{atan}(\exp(x)))}{9} + \frac{(2*\exp(x))}{(3*(\exp(6*x) + 1))} + \frac{(3^{(1/2)}*\log(((2*\exp(x))/3 - 3^{(1/2)}/3)^2 + 1/9))}{18} - \frac{(3^{(1/2)}*\log(((2*\exp(x))/3 + 3^{(1/2)}/3)^2 + 1/9))}{18}$$

### Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.07

$$\int e^x \tanh^2(3x) dx = \frac{-4e^{6x} \operatorname{atan}(e^x) - 4\operatorname{atan}(e^x) - 2e^{6x} \operatorname{atan}(2e^x - \sqrt{3}) - 2\operatorname{atan}(2e^x - \sqrt{3}) - 2e^{6x} \operatorname{atan}(2e^x + \sqrt{3}) - 2\operatorname{atan}(2e^x + \sqrt{3})}{18}$$

input `int(exp(x)*tanh(3*x)^2,x)`

output

```
( - 4*e**(6*x)*atan(e**x) - 4*atan(e**x) - 2*e**(6*x)*atan(2*e**x - sqrt(3)) - 2*atan(2*e**x - sqrt(3)) - 2*e**(6*x)*atan(2*e**x + sqrt(3)) - 2*atan(2*e**x + sqrt(3)) + 18*e**(7*x) + e**(6*x)*sqrt(3)*log(e**(2*x) - e**x*sqrt(3) + 1) - e**(6*x)*sqrt(3)*log(e**(2*x) + e**x*sqrt(3) + 1) + 30*e**x + sqrt(3)*log(e**(2*x) - e**x*sqrt(3) + 1) - sqrt(3)*log(e**(2*x) + e**x*sqrt(3) + 1))/(18*(e**(6*x) + 1))
```

### 3.230 $\int e^x \tanh(3x) dx$

Optimal result	1685
Mathematica [C] (verified)	1685
Rubi [A] (verified)	1686
Maple [C] (verified)	1689
Fricas [A] (verification not implemented)	1690
Sympy [F]	1690
Maxima [A] (verification not implemented)	1691
Giac [A] (verification not implemented)	1691
Mupad [B] (verification not implemented)	1692
Reduce [B] (verification not implemented)	1692

#### Optimal result

Integrand size = 8, antiderivative size = 70

$$\int e^x \tanh(3x) dx = e^x - \frac{2 \arctan(e^x)}{3} + \frac{1}{3} \arctan(\sqrt{3} - 2e^x) - \frac{1}{3} \arctan(\sqrt{3} + 2e^x) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}e^x}{1+e^{2x}}\right)}{\sqrt{3}}$$

output

$\exp(x) - 2/3 \arctan(\exp(x)) - 1/3 \arctan(-3^{1/2} + 2 \exp(x)) - 1/3 \arctan(3^{1/2} + 2 \exp(x)) - 1/3 \operatorname{arctanh}(3^{1/2} \exp(x) / (1 + \exp(2x))) * 3^{1/2}$

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.34

$$\int e^x \tanh(3x) dx = e^x - 2e^x \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, 1, \frac{7}{6}, -e^{6x}\right)$$

input

`Integrate[E^x*Tanh[3*x], x]`

output  $E^x - 2E^x \text{Hypergeometric2F1}[1/6, 1, 7/6, -E^{(6*x)}]$

### Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.49, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {2720, 25, 913, 753, 27, 216, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \tanh(3x) dx \\
 & \quad \downarrow 2720 \\
 & \int -\frac{1 - e^{6x}}{e^{6x} + 1} de^x \\
 & \quad \downarrow 25 \\
 & -\int \frac{1 - e^{6x}}{1 + e^{6x}} de^x \\
 & \quad \downarrow 913 \\
 & e^x - 2 \int \frac{1}{1 + e^{6x}} de^x \\
 & \quad \downarrow 753 \\
 & e^x - 2 \left( \frac{1}{3} \int \frac{1}{1 + e^{2x}} de^x + \frac{1}{3} \int \frac{2 - \sqrt{3}e^x}{2(1 - \sqrt{3}e^x + e^{2x})} de^x + \frac{1}{3} \int \frac{2 + \sqrt{3}e^x}{2(1 + \sqrt{3}e^x + e^{2x})} de^x \right) \\
 & \quad \downarrow 27 \\
 & e^x - 2 \left( \frac{1}{3} \int \frac{1}{1 + e^{2x}} de^x + \frac{1}{6} \int \frac{2 - \sqrt{3}e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x + \frac{1}{6} \int \frac{2 + \sqrt{3}e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x \right) \\
 & \quad \downarrow 216 \\
 & e^x - 2 \left( \frac{1}{6} \int \frac{2 - \sqrt{3}e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x + \frac{1}{6} \int \frac{2 + \sqrt{3}e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x + \frac{\arctan(e^x)}{3} \right) \\
 & \quad \downarrow 1142
 \end{aligned}$$

$$\begin{aligned}
& 2 \left( \frac{1}{6} \left( \frac{1}{2} \int \frac{1}{1 - \sqrt{3}e^x + e^{2x}} de^x - \frac{1}{2} \sqrt{3} \int -\frac{e^x - \sqrt{3} - 2e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x \right) + \frac{1}{6} \left( \frac{1}{2} \int \frac{1}{1 + \sqrt{3}e^x + e^{2x}} de^x + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} - 2e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x \right) \right) \\
& \quad \downarrow \text{25} \\
& 2 \left( \frac{1}{6} \left( \frac{1}{2} \int \frac{1}{1 - \sqrt{3}e^x + e^{2x}} de^x + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} - 2e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x \right) + \frac{1}{6} \left( \frac{1}{2} \int \frac{1}{1 + \sqrt{3}e^x + e^{2x}} de^x + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} - 2e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x \right) \right) \\
& \quad \downarrow \text{1083} \\
& 2 \left( \frac{1}{6} \left( \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} - 2e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x - \int \frac{1}{-1 - e^{2x}} d(-\sqrt{3} + 2e^x) \right) + \frac{1}{6} \left( \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} + 2e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x - \int \frac{1}{-1 - e^{2x}} d(\sqrt{3} + 2e^x) \right) \right) \\
& \quad \downarrow \text{217} \\
& 2 \left( \frac{1}{6} \left( \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} - 2e^x}{1 - \sqrt{3}e^x + e^{2x}} de^x - \arctan(\sqrt{3} - 2e^x) \right) + \frac{1}{6} \left( \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} + 2e^x}{1 + \sqrt{3}e^x + e^{2x}} de^x + \arctan(2e^x + \sqrt{3}) \right) \right) \\
& \quad \downarrow \text{1103} \\
& 2 \left( \frac{\arctan(e^x)}{3} + \frac{1}{6} \left( -\arctan(\sqrt{3} - 2e^x) - \frac{1}{2} \sqrt{3} \log(-\sqrt{3}e^x + e^{2x} + 1) \right) + \frac{1}{6} \left( \arctan(2e^x + \sqrt{3}) + \frac{1}{2} \sqrt{3} \log(\sqrt{3}e^x + e^{2x} + 1) \right) \right)
\end{aligned}$$

input `Int [E^x*Tanh[3*x], x]`

output `E^x - 2*(ArcTan[E^x]/3 + (-ArcTan[Sqrt[3] - 2*E^x] - (Sqrt[3]*Log[1 - Sqrt[3]*E^x + E^(2*x)]))/2)/6 + (ArcTan[Sqrt[3] + 2*E^x] + (Sqrt[3]*Log[1 + Sqrt[3]*E^x + E^(2*x)]))/2)/6)`



## Defintions of rubi rules used

- rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27  $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 216  $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[\text{b}, 2]))*\text{ArcTan}[\text{Rt}[\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{GtQ}[\text{a}, 0] \ || \ \text{GtQ}[\text{b}, 0])$
- rule 217  $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$
- rule 753  $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^{(n_)} )^{-1}, \text{x\_Symbol}] \rightarrow \text{Module}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, \text{n}]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, \text{n}]], \text{k}, \text{u}, \text{v}\}, \text{Simp}[\text{u} = \text{Int}[(\text{r} - \text{s}*\text{Cos}[(2*\text{k} - 1)*(Pi/\text{n}])*x]/(\text{r}^2 - 2*\text{r}*\text{s}*\text{Cos}[(2*\text{k} - 1)*(Pi/\text{n}])*x + \text{s}^2*x^2), \text{x}] + \text{Int}[(\text{r} + \text{s}*\text{Cos}[(2*\text{k} - 1)*(Pi/\text{n}])*x]/(\text{r}^2 + 2*\text{r}*\text{s}*\text{Cos}[(2*\text{k} - 1)*(Pi/\text{n}])*x + \text{s}^2*x^2), \text{x}]; 2*(\text{r}^2/(\text{a}*\text{n})) \quad \text{Int}[1/(\text{r}^2 + \text{s}^2*x^2), \text{x}] + 2*(\text{r}/(\text{a}*\text{n})) \quad \text{Sum}[\text{u}, \{\text{k}, 1, (\text{n} - 2)/4\}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{IGtQ}[(\text{n} - 2)/4, 0] \ \&\& \ \text{PosQ}[\text{a}/\text{b}]$
- rule 913  $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_)^{(n_)} )^{(p_)}*((\text{c}_) + (\text{d}_.)*(x_)^{(n_)}), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{d}*x*((\text{a} + \text{b}*x^n)^{(p+1})/(\text{b}*(\text{n}*(p+1) + 1))), \text{x}] - \text{Simp}[(\text{a}*\text{d} - \text{b}*\text{c}*(\text{n}*(p+1) + 1))/(\text{b}*(\text{n}*(p+1) + 1)) \quad \text{Int}[(\text{a} + \text{b}*x^n)^p, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{n}, \text{p}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{n}*(\text{p} + 1) + 1, 0]$
- rule 1083  $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[\text{b}^2 - 4*\text{a}*\text{c} - \text{x}^2, \text{x}], \text{x}], \text{x}, \text{b} + 2*\text{c}*x], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

## Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.55 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.67

method	result	size
risch	$e^x + \left( \sum_{R=\text{RootOf}(81Z^4-9Z^2+1)} -R \ln(e^x - 3R) \right) + \frac{i \ln(e^x - i)}{3} - \frac{i \ln(e^x + i)}{3}$	47

input `int(exp(x)*tanh(3*x), x, method=_RETURNVERBOSE)`

output `exp(x)+sum(_R*ln(exp(x)-3*_R), _R=RootOf(81*_Z^4-9*_Z^2+1))+1/3*I*ln(exp(x)-I)-1/3*I*ln(exp(x)+I)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.34

$$\int e^x \tanh(3x) dx = -\frac{1}{6} \sqrt{3} \log \left( \frac{\sqrt{3} + 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) + \frac{1}{6} \sqrt{3} \log \left( -\frac{\sqrt{3} - 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) - \frac{1}{3} \arctan \left( \sqrt{3} + 2 \cosh(x) + 2 \sinh(x) \right) - \frac{1}{3} \arctan \left( -\sqrt{3} + 2 \cosh(x) + 2 \sinh(x) \right) - \frac{2}{3} \arctan(\cosh(x) + \sinh(x)) + \cosh(x) + \sinh(x)$$

input `integrate(exp(x)*tanh(3*x),x, algorithm="fricas")`output `-1/6*sqrt(3)*log((sqrt(3) + 2*cosh(x))/(cosh(x) - sinh(x))) + 1/6*sqrt(3)*log(-(sqrt(3) - 2*cosh(x))/(cosh(x) - sinh(x))) - 1/3*arctan(sqrt(3) + 2*cosh(x) + 2*sinh(x)) - 1/3*arctan(-sqrt(3) + 2*cosh(x) + 2*sinh(x)) - 2/3*arctan(cosh(x) + sinh(x)) + cosh(x) + sinh(x)`**Sympy [F]**

$$\int e^x \tanh(3x) dx = \int e^x \tanh(3x) dx$$

input `integrate(exp(x)*tanh(3*x),x)`output `Integral(exp(x)*tanh(3*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.99

$$\int e^x \tanh(3x) dx = -\frac{1}{6} \sqrt{3} \log(\sqrt{3}e^x + e^{(2x)} + 1) + \frac{1}{6} \sqrt{3} \log(-\sqrt{3}e^x + e^{(2x)} + 1) - \frac{1}{3} \arctan(\sqrt{3} + 2e^x) - \frac{1}{3} \arctan(-\sqrt{3} + 2e^x) - \frac{2}{3} \arctan(e^x) + e^x$$

input `integrate(exp(x)*tanh(3*x),x, algorithm="maxima")`output `-1/6*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/6*sqrt(3)*log(-sqrt(3)*e^x + e^(2*x) + 1) - 1/3*arctan(sqrt(3) + 2*e^x) - 1/3*arctan(-sqrt(3) + 2*e^x) - 2/3*arctan(e^x) + e^x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.99

$$\int e^x \tanh(3x) dx = -\frac{1}{6} \sqrt{3} \log(\sqrt{3}e^x + e^{(2x)} + 1) + \frac{1}{6} \sqrt{3} \log(-\sqrt{3}e^x + e^{(2x)} + 1) - \frac{1}{3} \arctan(\sqrt{3} + 2e^x) - \frac{1}{3} \arctan(-\sqrt{3} + 2e^x) - \frac{2}{3} \arctan(e^x) + e^x$$

input `integrate(exp(x)*tanh(3*x),x, algorithm="giac")`output `-1/6*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/6*sqrt(3)*log(-sqrt(3)*e^x + e^(2*x) + 1) - 1/3*arctan(sqrt(3) + 2*e^x) - 1/3*arctan(-sqrt(3) + 2*e^x) - 2/3*arctan(e^x) + e^x`

**Mupad [B] (verification not implemented)**

Time = 2.35 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00

$$\int e^x \tanh(3x) dx = e^x - \frac{\operatorname{atan}(2e^x + \sqrt{3})}{3} - \frac{\operatorname{atan}(2e^x - \sqrt{3})}{3} - \frac{2\operatorname{atan}(e^x)}{3} + \frac{\sqrt{3} \ln\left((2e^x - \sqrt{3})^2 + 1\right)}{6} - \frac{\sqrt{3} \ln\left((2e^x + \sqrt{3})^2 + 1\right)}{6}$$

input `int(tanh(3*x)*exp(x),x)`output `exp(x) - atan(2*exp(x) + 3^(1/2))/3 - atan(2*exp(x) - 3^(1/2))/3 - (2*atan(exp(x)))/3 + (3^(1/2)*log((2*exp(x) - 3^(1/2))^2 + 1))/6 - (3^(1/2)*log((2*exp(x) + 3^(1/2))^2 + 1))/6`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.01

$$\int e^x \tanh(3x) dx = -\frac{2\operatorname{atan}(e^x)}{3} - \frac{\operatorname{atan}(2e^x - \sqrt{3})}{3} - \frac{\operatorname{atan}(2e^x + \sqrt{3})}{3} + e^x + \frac{\sqrt{3} \log(e^{2x} - e^x \sqrt{3} + 1)}{6} - \frac{\sqrt{3} \log(e^{2x} + e^x \sqrt{3} + 1)}{6}$$

input `int(exp(x)*tanh(3*x),x)`output `( - 4*atan(e**x) - 2*atan(2*e**x - sqrt(3)) - 2*atan(2*e**x + sqrt(3)) + 6*e**x + sqrt(3)*log(e**(2*x) - e**x*sqrt(3) + 1) - sqrt(3)*log(e**(2*x) + e**x*sqrt(3) + 1))/6`

### 3.231 $\int e^x \coth(3x) dx$

Optimal result	1693
Mathematica [C] (verified)	1693
Rubi [A] (verified)	1694
Maple [C] (verified)	1697
Fricas [B] (verification not implemented)	1697
Sympy [F]	1698
Maxima [A] (verification not implemented)	1699
Giac [A] (verification not implemented)	1699
Mupad [B] (verification not implemented)	1700
Reduce [B] (verification not implemented)	1700

#### Optimal result

Integrand size = 8, antiderivative size = 71

$$\int e^x \coth(3x) dx = e^x + \frac{\arctan\left(\frac{1-2e^x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1+2e^x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{2\operatorname{arctanh}(e^x)}{3} - \frac{1}{3}\operatorname{arctanh}\left(\frac{e^x}{1+e^{2x}}\right)$$

output

```
exp(x)+1/3*arctan(1/3*(1-2*exp(x))*3^(1/2))*3^(1/2)-1/3*arctan(1/3*(1+2*exp(x))*3^(1/2))*3^(1/2)-2/3*arctanh(exp(x))-1/3*arctanh(exp(x)/(1+exp(2*x)))
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.31

$$\int e^x \coth(3x) dx = e^x - 2e^x \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, 1, \frac{7}{6}, e^{6x}\right)$$

input

```
Integrate[E^x*Coth[3*x],x]
```

output

$$E^x - 2E^x \text{Hypergeometric2F1}[1/6, 1, 7/6, E^{(6*x)}]$$
**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.37, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {2720, 25, 913, 754, 27, 219, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \coth(3x) dx \\
 & \quad \downarrow 2720 \\
 & \int -\frac{e^{6x} + 1}{1 - e^{6x}} de^x \\
 & \quad \downarrow 25 \\
 & -\int \frac{1 + e^{6x}}{1 - e^{6x}} de^x \\
 & \quad \downarrow 913 \\
 & e^x - 2 \int \frac{1}{1 - e^{6x}} de^x \\
 & \quad \downarrow 754 \\
 & e^x - 2 \left( \frac{1}{3} \int \frac{1}{1 - e^{2x}} de^x + \frac{1}{3} \int \frac{2 - e^x}{2(1 - e^x + e^{2x})} de^x + \frac{1}{3} \int \frac{2 + e^x}{2(1 + e^x + e^{2x})} de^x \right) \\
 & \quad \downarrow 27 \\
 & e^x - 2 \left( \frac{1}{3} \int \frac{1}{1 - e^{2x}} de^x + \frac{1}{6} \int \frac{2 - e^x}{1 - e^x + e^{2x}} de^x + \frac{1}{6} \int \frac{2 + e^x}{1 + e^x + e^{2x}} de^x \right) \\
 & \quad \downarrow 219 \\
 & e^x - 2 \left( \frac{1}{3} \int \frac{2 - e^x}{1 - e^x + e^{2x}} de^x + \frac{1}{6} \int \frac{2 + e^x}{1 + e^x + e^{2x}} de^x + \frac{\operatorname{arctanh}(e^x)}{3} \right) \\
 & \quad \downarrow 1142
 \end{aligned}$$

$$\begin{aligned}
& 2\left(\frac{1}{6}\left(\frac{3}{2}\int\frac{1}{1-e^x+e^{2x}}de^x-\frac{1}{2}\int-\frac{1-2e^x}{1-e^x+e^{2x}}de^x\right)+\frac{1}{6}\left(\frac{3}{2}\int\frac{1}{1+e^x+e^{2x}}de^x+\frac{1}{2}\int\frac{1+2e^x}{1+e^x+e^{2x}}de^x\right)+\arctan\left(\frac{e^x-1}{\sqrt{3}}\right)\right) \\
& \quad \downarrow 25 \\
& 2\left(\frac{1}{6}\left(\frac{3}{2}\int\frac{1}{1-e^x+e^{2x}}de^x+\frac{1}{2}\int\frac{1-2e^x}{1-e^x+e^{2x}}de^x\right)+\frac{1}{6}\left(\frac{3}{2}\int\frac{1}{1+e^x+e^{2x}}de^x+\frac{1}{2}\int\frac{1+2e^x}{1+e^x+e^{2x}}de^x\right)+\arctan\left(\frac{e^x-1}{\sqrt{3}}\right)\right) \\
& \quad \downarrow 1083 \\
& 2\left(\frac{1}{6}\left(\frac{1}{2}\int\frac{1-2e^x}{1-e^x+e^{2x}}de^x-3\int\frac{1}{-3-e^{2x}}d(-1+2e^x)\right)+\frac{1}{6}\left(\frac{1}{2}\int\frac{1+2e^x}{1+e^x+e^{2x}}de^x-3\int\frac{1}{-3-e^{2x}}d(1+2e^x)\right)+\arctan\left(\frac{e^x-1}{\sqrt{3}}\right)\right) \\
& \quad \downarrow 217 \\
& 2\left(\frac{1}{6}\left(\frac{1}{2}\int\frac{1-2e^x}{1-e^x+e^{2x}}de^x+\sqrt{3}\arctan\left(\frac{2e^x-1}{\sqrt{3}}\right)\right)+\frac{1}{6}\left(\frac{1}{2}\int\frac{1+2e^x}{1+e^x+e^{2x}}de^x+\sqrt{3}\arctan\left(\frac{2e^x+1}{\sqrt{3}}\right)\right)+\arctan\left(\frac{e^x-1}{\sqrt{3}}\right)\right) \\
& \quad \downarrow 1103 \\
& 2\left(\frac{1}{6}\left(\sqrt{3}\arctan\left(\frac{2e^x-1}{\sqrt{3}}\right)-\frac{1}{2}\log(-e^x+e^{2x}+1)\right)+\frac{1}{6}\left(\sqrt{3}\arctan\left(\frac{2e^x+1}{\sqrt{3}}\right)+\frac{1}{2}\log(e^x+e^{2x}+1)\right)+\arctan\left(\frac{e^x-1}{\sqrt{3}}\right)\right)
\end{aligned}$$

input `Int[E^x*Coth[3*x], x]`

output `E^x - 2*(ArcTanh[E^x]/3 + (Sqrt[3]*ArcTan[(-1 + 2*E^x)/Sqrt[3]] - Log[1 - E^x + E^(2*x)]/2)/6 + (Sqrt[3]*ArcTan[(1 + 2*E^x)/Sqrt[3]] + Log[1 + E^x + E^(2*x)]/2)/6)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`



rule 217  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1}] \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 219  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 754  $\text{Int}[(a_ + (b_ \cdot)(x_ )^{(n_ )})^{-1}, x\_Symbol] \rightarrow \text{Module}[\{r = \text{Numerator}[\text{Rt}[-a/b, n]], s = \text{Denominator}[\text{Rt}[-a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r - s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x)/(r^2 - 2 \cdot r \cdot s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x + s^2 \cdot x^2), x] + \text{Int}[(r + s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x)/(r^2 + 2 \cdot r \cdot s \cdot \text{Cos}[(2 \cdot k \cdot \text{Pi})/n] \cdot x + s^2 \cdot x^2), x]; 2 \cdot (r^2/(a \cdot n)) \ \text{Int}[1/(r^2 - s^2 \cdot x^2), x] + 2 \cdot (r/(a \cdot n)) \ \text{Sum}[u, \{k, 1, (n - 2)/4\}], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[(n - 2)/4, 0] \ \&\& \ \text{NegQ}[a/b]$

rule 913  $\text{Int}[(a_ + (b_ \cdot)(x_ )^{(n_ )})^{(p_ )} \cdot ((c_ ) + (d_ \cdot)(x_ )^{(n_ )}), x\_Symbol] \rightarrow \text{Simp}[d \cdot x \cdot ((a + b \cdot x^n)^{(p + 1})/(b \cdot (n \cdot (p + 1) + 1))), x] - \text{Simp}[(a \cdot d - b \cdot c \cdot (n \cdot (p + 1) + 1))/(b \cdot (n \cdot (p + 1) + 1)) \ \text{Int}[(a + b \cdot x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, n, p\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[n \cdot (p + 1) + 1, 0]$

rule 1083  $\text{Int}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d_ ) + (e_ \cdot)(x_ )]/((a_ \cdot) + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142  $\text{Int}[(d_ \cdot) + (e_ \cdot)(x_ )]/((a_ ) + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 2720

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.94

method	result
risch	$e^x - \frac{\ln\left(e^x + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}{6} + \frac{i \ln\left(e^x + \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)\sqrt{3}}{6} - \frac{\ln\left(e^x + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{6} - \frac{i \ln\left(e^x + \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)\sqrt{3}}{6} - \frac{\ln(e^x + 1)}{3} + \frac{\ln\left(e^x - \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}{6}$

input

```
int(exp(x)*coth(3*x), x, method=_RETURNVERBOSE)
```

output

```
exp(x)-1/6*ln(exp(x)+1/2-1/2*I*3^(1/2))+1/6*I*ln(exp(x)+1/2-1/2*I*3^(1/2))
*3^(1/2)-1/6*ln(exp(x)+1/2+1/2*I*3^(1/2))-1/6*I*ln(exp(x)+1/2+1/2*I*3^(1/2))
)*3^(1/2)-1/3*ln(exp(x)+1)+1/6*ln(exp(x)-1/2-1/2*I*3^(1/2))+1/6*I*ln(exp(x)-1/2-1/2*I*3^(1/2))
)*3^(1/2)+1/6*ln(exp(x)-1/2+1/2*I*3^(1/2))-1/6*I*ln(exp(x)-1/2+1/2*I*3^(1/2))
)*3^(1/2)+1/3*ln(exp(x)-1)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(56) = 112.

Time = 0.09 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.59

$$\begin{aligned} \int e^x \coth(3x) dx = & -\frac{1}{3} \sqrt{3} \arctan \left( \frac{2}{3} \sqrt{3} \cosh(x) + \frac{2}{3} \sqrt{3} \sinh(x) + \frac{1}{3} \sqrt{3} \right) \\ & - \frac{1}{3} \sqrt{3} \arctan \left( \frac{2}{3} \sqrt{3} \cosh(x) + \frac{2}{3} \sqrt{3} \sinh(x) - \frac{1}{3} \sqrt{3} \right) \\ & + \cosh(x) - \frac{1}{6} \log \left( \frac{2 \cosh(x) + 1}{\cosh(x) - \sinh(x)} \right) \\ & + \frac{1}{6} \log \left( \frac{2 \cosh(x) - 1}{\cosh(x) - \sinh(x)} \right) - \frac{1}{3} \log(\cosh(x) + \sinh(x) + 1) \\ & + \frac{1}{3} \log(\cosh(x) + \sinh(x) - 1) + \sinh(x) \end{aligned}$$

input `integrate(exp(x)*coth(3*x),x, algorithm="fricas")`

output `-1/3*sqrt(3)*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) + 1/3*sqrt(3)) - 1/3*sqrt(3)*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) - 1/3*sqrt(3)) + cosh(x) - 1/6*log((2*cosh(x) + 1)/(cosh(x) - sinh(x))) + 1/6*log((2*cosh(x) - 1)/(cosh(x) - sinh(x))) - 1/3*log(cosh(x) + sinh(x) + 1) + 1/3*log(cosh(x) + sinh(x) - 1) + sinh(x)`

## Sympy [F]

$$\int e^x \coth(3x) dx = \int e^x \coth(3x) dx$$

input `integrate(exp(x)*coth(3*x),x)`

output `Integral(exp(x)*coth(3*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.06

$$\int e^x \coth(3x) dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x + 1)\right) - \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x - 1)\right) \\ + e^x - \frac{1}{6} \log(e^{(2x)} + e^x + 1) + \frac{1}{6} \log(e^{(2x)} - e^x + 1) \\ - \frac{1}{3} \log(e^x + 1) + \frac{1}{3} \log(e^x - 1)$$

input `integrate(exp(x)*coth(3*x),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) + e^x - 1/6*log(e^(2*x) + e^x + 1) + 1/6*log(e^(2*x) - e^x + 1) - 1/3*log(e^x + 1) + 1/3*log(e^x - 1)`**Giac [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.07

$$\int e^x \coth(3x) dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x + 1)\right) - \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x - 1)\right) \\ + e^x - \frac{1}{6} \log(e^{(2x)} + e^x + 1) + \frac{1}{6} \log(e^{(2x)} - e^x + 1) \\ - \frac{1}{3} \log(e^x + 1) + \frac{1}{3} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(3*x),x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) + e^x - 1/6*log(e^(2*x) + e^x + 1) + 1/6*log(e^(2*x) - e^x + 1) - 1/3*log(e^x + 1) + 1/3*log(abs(e^x - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.14

$$\int e^x \coth(3x) dx = \frac{\ln(2 - 2e^x)}{3} - \frac{\ln(-2e^x - 2)}{3} + \frac{\ln((2e^x - 1)^2 + 3)}{6} - \frac{\ln((2e^x + 1)^2 + 3)}{6} + e^x - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2e^x - 1)}{3}\right)}{3} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2e^x + 1)}{3}\right)}{3}$$

input `int(coth(3*x)*exp(x),x)`output `log(2 - 2*exp(x))/3 - log(- 2*exp(x) - 2)/3 + log((2*exp(x) - 1)^2 + 3)/6 - log((2*exp(x) + 1)^2 + 3)/6 + exp(x) - (3^(1/2)*atan((3^(1/2)*(2*exp(x) - 1))/3))/3 - (3^(1/2)*atan((3^(1/2)*(2*exp(x) + 1))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.15

$$\int e^x \coth(3x) dx = -\frac{\sqrt{3} \operatorname{atan}\left(\frac{2e^x - 1}{\sqrt{3}}\right)}{3} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2e^x + 1}{\sqrt{3}}\right)}{3} + e^x - \frac{\log(e^{2x} + e^x + 1)}{6} + \frac{\log(e^{2x} - e^x + 1)}{6} + \frac{\log(e^x - 1)}{3} - \frac{\log(e^x + 1)}{3}$$

input `int(exp(x)*coth(3*x),x)`output `( - 2*sqrt(3)*atan((2*e**x - 1)/sqrt(3)) - 2*sqrt(3)*atan((2*e**x + 1)/sqrt(3)) + 6*e**x - log(e**(2*x) + e**x + 1) + log(e**(2*x) - e**x + 1) + 2*log(e**x - 1) - 2*log(e**x + 1))/6`

### 3.232 $\int e^x \coth^2(3x) dx$

Optimal result	1701
Mathematica [C] (verified)	1701
Rubi [A] (verified)	1702
Maple [C] (verified)	1703
Fricas [B] (verification not implemented)	1704
Sympy [F]	1705
Maxima [A] (verification not implemented)	1705
Giac [A] (verification not implemented)	1705
Mupad [B] (verification not implemented)	1706
Reduce [B] (verification not implemented)	1706

#### Optimal result

Integrand size = 10, antiderivative size = 94

$$\int e^x \coth^2(3x) dx = e^x + \frac{2e^x}{3(1 - e^{6x})} + \frac{\arctan\left(\frac{1-2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{\arctan\left(\frac{1+2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{2\operatorname{arctanh}(e^x)}{9} - \frac{1}{9}\operatorname{arctanh}\left(\frac{e^x}{1 + e^{2x}}\right)$$

output

```
exp(x)+2*exp(x)/(3-3*exp(6*x))+1/9*arctan(1/3*(1-2*exp(x))*3^(1/2))*3^(1/2)
)-1/9*arctan(1/3*(1+2*exp(x))*3^(1/2))*3^(1/2)-2/9*arctanh(exp(x))-1/9*arc
tanh(exp(x)/(1+exp(2*x)))
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.45 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.20

$$\int e^x \coth^2(3x) dx = \frac{e^{-11x}(-15379 - 28153e^{6x} - 5633e^{12x} + 3109e^{18x} + 7(2197 + 3708e^{6x} + 538e^{12x} - 684e^{18x} + e^{24x}) \operatorname{Hypergeometric2F1}\left(\frac{7}{6}, 2, 2, 2; 1, 1, \frac{25}{6}; e^{6x}\right)}{3024} + \frac{36e^{7x}(1 + e^{6x})^2 {}_4F_3\left(\frac{7}{6}, 2, 2, 2; 1, 1, \frac{25}{6}; e^{6x}\right)}{1729}$$

input `Integrate[E^x*Coth[3*x]^2,x]`

output  $(-15379 - 28153E^{(6*x)} - 5633E^{(12*x)} + 3109E^{(18*x)} + 7*(2197 + 3708E^{(6*x)} + 538E^{(12*x)} - 684E^{(18*x)} + E^{(24*x)})\text{Hypergeometric2F1}[1/6, 1, 7/6, E^{(6*x)}]) / (3024E^{(11*x)}) + (36E^{(7*x)}*(1 + E^{(6*x)})^2\text{HypergeometricPFQ}[\{7/6, 2, 2, 2\}, \{1, 1, 25/6\}, E^{(6*x)}]) / 1729$

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.15, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \coth^2(3x) dx$$

$$\downarrow 2720$$

$$\int \frac{(e^{6x} + 1)^2}{(1 - e^{6x})^2} de^x$$

$$\downarrow 915$$

$$\int \left( \frac{4e^{6x}}{(1 - e^{6x})^2} + 1 \right) de^x$$

$$\downarrow 2009$$

$$\frac{\arctan\left(\frac{1-2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{\arctan\left(\frac{2e^x+1}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{2\operatorname{arctanh}(e^x)}{9} + e^x + \frac{2e^x}{3(1 - e^{6x})} + \frac{1}{18} \log(-e^x + e^{2x} + 1) - \frac{1}{18} \log(e^x + e^{2x} + 1)$$

input `Int[E^x*Coth[3*x]^2,x]`

output

$$E^x + (2E^x)/(3(1 - E^{6x})) + \text{ArcTan}[(1 - 2E^x)/\text{Sqrt}[3]]/(3\text{Sqrt}[3]) - \text{ArcTan}[(1 + 2E^x)/\text{Sqrt}[3]]/(3\text{Sqrt}[3]) - (2\text{ArcTanh}[E^x])/9 + \text{Log}[1 - E^x + E^{2x}]/18 - \text{Log}[1 + E^x + E^{2x}]/18$$
**Defintions of rubi rules used**

rule 915

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2720

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.60

method	result
risch	$e^x - \frac{2e^x}{3(e^{6x}-1)} + \frac{\ln(e^x-1)}{9} - \frac{\ln(e^x+1)}{9} + \frac{\ln\left(e^x - \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}{18} + \frac{i \ln\left(e^x - \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)\sqrt{3}}{18} + \frac{\ln\left(e^x - \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{18} - \frac{i \ln\left(e^x - \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)\sqrt{3}}{18}$

input

```
int(exp(x)*coth(3*x)^2,x,method=_RETURNVERBOSE)
```



output

```
exp(x)-2/3*exp(x)/(exp(6*x)-1)+1/9*ln(exp(x)-1)-1/9*ln(exp(x)+1)+1/18*ln(e
xp(x)-1/2-1/2*I*3^(1/2))+1/18*I*3^(1/2)*ln(exp(x)-1/2-1/2*I*3^(1/2))+1/18*
ln(exp(x)-1/2+1/2*I*3^(1/2))-1/18*I*3^(1/2)*ln(exp(x)-1/2+1/2*I*3^(1/2))-1
/18*ln(exp(x)+1/2-1/2*I*3^(1/2))+1/18*I*3^(1/2)*ln(exp(x)+1/2-1/2*I*3^(1/2
))-1/18*ln(exp(x)+1/2+1/2*I*3^(1/2))-1/18*I*3^(1/2)*ln(exp(x)+1/2+1/2*I*3^
(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 628 vs.  $2(68) = 136$ .

Time = 0.11 (sec) , antiderivative size = 628, normalized size of antiderivative = 6.68

$$\int e^x \coth^2(3x) dx = \text{Too large to display}$$

input

```
integrate(exp(x)*coth(3*x)^2,x, algorithm="fricas")
```

output

```
1/18*(18*cosh(x)^7 + 378*cosh(x)^5*sinh(x)^2 + 630*cosh(x)^4*sinh(x)^3 + 6
30*cosh(x)^3*sinh(x)^4 + 378*cosh(x)^2*sinh(x)^5 + 126*cosh(x)*sinh(x)^6 +
  18*sinh(x)^7 - 2*(sqrt(3)*cosh(x)^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*sq
rt(3)*cosh(x)^4*sinh(x)^2 + 20*sqrt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*co
sh(x)^2*sinh(x)^4 + 6*sqrt(3)*cosh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 - sqrt
(3))*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) + 1/3*sqrt(3)) - 2*(
sqrt(3)*cosh(x)^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*sqrt(3)*cosh(x)^4*si
nh(x)^2 + 20*sqrt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*cosh(x)^2*sinh(x)^4 +
  6*sqrt(3)*cosh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 - sqrt(3))*arctan(2/3*sq
rt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) - 1/3*sqrt(3)) - (cosh(x)^6 + 6*cosh(x)
^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^
2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log((2*cosh(x) + 1)/(co
sh(x) - sinh(x))) + (cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)
)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^
5 + sinh(x)^6 - 1)*log((2*cosh(x) - 1)/(cosh(x) - sinh(x))) - 2*(cosh(x)^6
+ 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 +
  15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log(cosh(x)
+ sinh(x) + 1) + 2*(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)
)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^
5 + sinh(x)^6 - 1)*log(cosh(x) + sinh(x) - 1) + 6*(21*cosh(x)^6 - 5)*si...
```

**Sympy [F]**

$$\int e^x \coth^2(3x) dx = \int e^x \coth^2(3x) dx$$

input `integrate(exp(x)*coth(3*x)**2,x)`

output `Integral(exp(x)*coth(3*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.93

$$\begin{aligned} \int e^x \coth^2(3x) dx = & -\frac{1}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x + 1)\right) - \frac{1}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x - 1)\right) \\ & - \frac{2e^x}{3(e^{6x} - 1)} + e^x - \frac{1}{18} \log(e^{2x} + e^x + 1) \\ & + \frac{1}{18} \log(e^{2x} - e^x + 1) - \frac{1}{9} \log(e^x + 1) + \frac{1}{9} \log(e^x - 1) \end{aligned}$$

input `integrate(exp(x)*coth(3*x)^2,x, algorithm="maxima")`

output `-1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) - 2/3*e^x/(e^(6*x) - 1) + e^x - 1/18*log(e^(2*x) + e^x + 1) + 1/18*log(e^(2*x) - e^x + 1) - 1/9*log(e^x + 1) + 1/9*log(e^x - 1)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.94

$$\begin{aligned} \int e^x \coth^2(3x) dx = & -\frac{1}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x + 1)\right) - \frac{1}{9} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2e^x - 1)\right) \\ & - \frac{2e^x}{3(e^{6x} - 1)} + e^x - \frac{1}{18} \log(e^{2x} + e^x + 1) \\ & + \frac{1}{18} \log(e^{2x} - e^x + 1) - \frac{1}{9} \log(e^x + 1) + \frac{1}{9} \log(|e^x - 1|) \end{aligned}$$

input `integrate(exp(x)*coth(3*x)^2,x, algorithm="giac")`

output `-1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) - 2/3*e^x/(e^(6*x) - 1) + e^x - 1/18*log(e^(2*x) + e^x + 1) + 1/18*log(e^(2*x) - e^x + 1) - 1/9*log(e^x + 1) + 1/9*log(abs(e^x - 1))`

### Mupad [B] (verification not implemented)

Time = 2.47 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.99

$$\int e^x \coth^2(3x) dx = \frac{\ln\left(\frac{2}{3} - \frac{2e^x}{3}\right)}{9} - \frac{\ln\left(-\frac{2e^x}{3} - \frac{2}{3}\right)}{9} + \frac{\ln\left(\left(\frac{2e^x}{3} - \frac{1}{3}\right)^2 + \frac{1}{3}\right)}{18} - \frac{\ln\left(\left(\frac{2e^x}{3} + \frac{1}{3}\right)^2 + \frac{1}{3}\right)}{18} + e^x - \frac{2e^x}{3(e^{6x} - 1)} - \frac{\sqrt{3} \operatorname{atan}\left(\sqrt{3}\left(\frac{2e^x}{3} - \frac{1}{3}\right)\right)}{9} - \frac{\sqrt{3} \operatorname{atan}\left(\sqrt{3}\left(\frac{2e^x}{3} + \frac{1}{3}\right)\right)}{9}$$

input `int(coth(3*x)^2*exp(x),x)`

output `log(2/3 - (2*exp(x))/3)/9 - log(-(2*exp(x))/3 - 2/3)/9 + log(((2*exp(x))/3 - 1/3)^2 + 1/3)/18 - log(((2*exp(x))/3 + 1/3)^2 + 1/3)/18 + exp(x) - (2*exp(x))/(3*(exp(6*x) - 1)) - (3^(1/2)*atan(3^(1/2)*((2*exp(x))/3 - 1/3)))/9 - (3^(1/2)*atan(3^(1/2)*((2*exp(x))/3 + 1/3)))/9`

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.21

$$\int e^x \coth^2(3x) dx = \frac{-2e^{6x}\sqrt{3} \operatorname{atan}\left(\frac{2e^x-1}{\sqrt{3}}\right) + 2\sqrt{3} \operatorname{atan}\left(\frac{2e^x-1}{\sqrt{3}}\right) - 2e^{6x}\sqrt{3} \operatorname{atan}\left(\frac{2e^x+1}{\sqrt{3}}\right) + 2\sqrt{3} \operatorname{atan}\left(\frac{2e^x+1}{\sqrt{3}}\right) + 18e^{7x} - e^{6x}\log}{=}$$

input `int(exp(x)*coth(3*x)^2,x)`

output  $(-2e^{6x}\sqrt{3}\operatorname{atan}((2e^{3x}-1)/\sqrt{3}) + 2\sqrt{3}\operatorname{atan}((2e^{3x}-1)/\sqrt{3}) - 2e^{6x}\sqrt{3}\operatorname{atan}((2e^{3x}+1)/\sqrt{3}) + 2\sqrt{3}\operatorname{atan}((2e^{3x}+1)/\sqrt{3}) + 18e^{7x} - e^{6x}\log(e^{2x} + e^{3x} + 1) + e^{6x}\log(e^{2x} - e^{3x} + 1) + 2e^{6x}\log(e^{3x} - 1) - 2e^{6x}\log(e^{3x} + 1) - 30e^{3x} + \log(e^{2x} + e^{3x} + 1) - \log(e^{2x} - e^{3x} + 1) - 2\log(e^{3x} - 1) + 2\log(e^{3x} + 1))/(18(e^{6x} - 1))$

### 3.233 $\int e^x \tanh^2(4x) dx$

Optimal result	1708
Mathematica [C] (verified)	1709
Rubi [A] (verified)	1709
Maple [C] (verified)	1711
Fricas [C] (verification not implemented)	1711
Sympy [F]	1712
Maxima [F]	1713
Giac [A] (verification not implemented)	1713
Mupad [B] (verification not implemented)	1714
Reduce [B] (verification not implemented)	1715

#### Optimal result

Integrand size = 10, antiderivative size = 296

$$\begin{aligned}
 \int e^x \tanh^2(4x) dx = & e^x + \frac{e^x}{2(1+e^{8x})} + \frac{1}{16} \sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}}-2e^x}{\sqrt{2+\sqrt{2}}}\right) \\
 & + \frac{1}{16} \sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}}-2e^x}{\sqrt{2-\sqrt{2}}}\right) \\
 & - \frac{1}{16} \sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}}+2e^x}{\sqrt{2+\sqrt{2}}}\right) \\
 & - \frac{1}{16} \sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}}+2e^x}{\sqrt{2-\sqrt{2}}}\right) \\
 & - \frac{1}{16} \sqrt{2-\sqrt{2}} \operatorname{arctanh}\left(\frac{\sqrt{2-\sqrt{2}}e^x}{1+e^{2x}}\right) \\
 & - \frac{1}{16} \sqrt{2+\sqrt{2}} \operatorname{arctanh}\left(\frac{\sqrt{2+\sqrt{2}}e^x}{1+e^{2x}}\right)
 \end{aligned}$$

output

```
exp(x)+exp(x)/(2+2*exp(8*x))+1/16*(2+2^(1/2))^(1/2)*arctan(((2-2^(1/2))^(1/2)-2*exp(x))/(2+2^(1/2))^(1/2))+1/16*(2-2^(1/2))^(1/2)*arctan(((2+2^(1/2))^(1/2)-2*exp(x))/(2-2^(1/2))^(1/2))-1/16*(2+2^(1/2))^(1/2)*arctan(((2-2^(1/2))^(1/2)+2*exp(x))/(2+2^(1/2))^(1/2))-1/16*(2-2^(1/2))^(1/2)*arctan(((2+2^(1/2))^(1/2)+2*exp(x))/(2-2^(1/2))^(1/2))-1/16*(2-2^(1/2))^(1/2)*arctanh(((2-2^(1/2))^(1/2)*exp(x)/(1+exp(2*x)))-1/16*(2+2^(1/2))^(1/2)*arctanh((2+2^(1/2))^(1/2)*exp(x)/(1+exp(2*x)))
```

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.17

$$\int e^x \tanh^2(4x) dx = e^x + \frac{e^x}{2(1 + e^{8x})} + \frac{1}{16} \text{RootSum} \left[ 1 + \#1^8 \&, \frac{x - \log(e^x - \#1)}{\#1^7} \& \right]$$

input

```
Integrate[E^x*Tanh[4*x]^2,x]
```

output

```
E^x + E^x/(2*(1 + E^(8*x))) + RootSum[1 + #1^8 & , (x - Log[E^x - #1])/#1^7 & ]/16
```

### Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.29, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \tanh^2(4x) dx$$

$$\downarrow 2720$$

$$\int \frac{(1 - e^{8x})^2}{(e^{8x} + 1)^2} de^x$$

$$\begin{aligned}
 & \int \left( 1 - \frac{4e^{8x}}{(e^{8x} + 1)^2} \right) dx \\
 & \quad \downarrow \text{915} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\arctan\left(\frac{\sqrt{2-\sqrt{2}-2e^x}}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2}(2-\sqrt{2})} + \frac{\arctan\left(\frac{\sqrt{2+\sqrt{2}-2e^x}}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2}(2+\sqrt{2})} - \frac{\arctan\left(\frac{2e^x+\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2}(2-\sqrt{2})} - \\
 & \frac{\arctan\left(\frac{2e^x+\sqrt{2+\sqrt{2}}}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2}(2+\sqrt{2})} + e^x + \frac{e^x}{2(e^{8x}+1)} + \frac{1}{32}\sqrt{2-\sqrt{2}}\log\left(-\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1\right) - \\
 & \frac{1}{32}\sqrt{2-\sqrt{2}}\log\left(\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1\right) + \frac{1}{32}\sqrt{2+\sqrt{2}}\log\left(-\sqrt{2+\sqrt{2}}e^x + e^{2x} + 1\right) - \\
 & \frac{1}{32}\sqrt{2+\sqrt{2}}\log\left(\sqrt{2+\sqrt{2}}e^x + e^{2x} + 1\right)
 \end{aligned}$$

input `Int [E^x*Tanh [4*x]^2, x]`

output `E^x + E^x/(2*(1 + E^(8*x))) + ArcTan[(Sqrt[2 - Sqrt[2]] - 2*E^x)/Sqrt[2 + Sqrt[2]]]/(8*Sqrt[2*(2 - Sqrt[2])]) + ArcTan[(Sqrt[2 + Sqrt[2]] - 2*E^x)/Sqrt[2 - Sqrt[2]]]/(8*Sqrt[2*(2 + Sqrt[2])]) - ArcTan[(Sqrt[2 - Sqrt[2]] + 2*E^x)/Sqrt[2 + Sqrt[2]]]/(8*Sqrt[2*(2 - Sqrt[2])]) - ArcTan[(Sqrt[2 + Sqrt[2]] + 2*E^x)/Sqrt[2 - Sqrt[2]]]/(8*Sqrt[2*(2 + Sqrt[2])]) + (Sqrt[2 - Sqrt[2]]*Log[1 - Sqrt[2 - Sqrt[2]]*E^x + E^(2*x)])/32 - (Sqrt[2 - Sqrt[2]]*Log[1 + Sqrt[2 - Sqrt[2]]*E^x + E^(2*x)])/32 + (Sqrt[2 + Sqrt[2]]*Log[1 - Sqrt[2 + Sqrt[2]]*E^x + E^(2*x)])/32 - (Sqrt[2 + Sqrt[2]]*Log[1 + Sqrt[2 + Sqrt[2]]*E^x + E^(2*x)])/32`

### Defintions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.69 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.12

method	result	size
risch	$e^x + \frac{e^x}{2+2e^{8x}} + \left( \sum_{_R=\text{RootOf}(4294967296\_Z^8+1)} \_R \ln(e^x - 16\_R) \right)$	36

input

```
int(exp(x)*tanh(4*x)^2,x,method=_RETURNVERBOSE)
```

output

```
exp(x)+1/2*exp(x)/(1+exp(8*x))+sum(_R*ln(exp(x)-16*_R),_R=RootOf(4294967296*_Z^8+1))
```

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 1303, normalized size of antiderivative = 4.40

$$\int e^x \tanh^2(4x) dx = \text{Too large to display}$$

input

```
integrate(exp(x)*tanh(4*x)^2,x, algorithm="fricas")
```



output

```

1/32*(32*cosh(x)^9 + 1152*cosh(x)^7*sinh(x)^2 + 2688*cosh(x)^6*sinh(x)^3 +
4032*cosh(x)^5*sinh(x)^4 + 4032*cosh(x)^4*sinh(x)^5 + 2688*cosh(x)^3*sinh
(x)^6 + 1152*cosh(x)^2*sinh(x)^7 + 288*cosh(x)*sinh(x)^8 + 32*sinh(x)^9 +
(-(I + 1)*sqrt(2)*(-1)^(1/8)*cosh(x)^8 - (8*I + 8)*sqrt(2)*(-1)^(1/8)*cosh
(x)^7*sinh(x) - (28*I + 28)*sqrt(2)*(-1)^(1/8)*cosh(x)^6*sinh(x)^2 - (56*I
+ 56)*sqrt(2)*(-1)^(1/8)*cosh(x)^5*sinh(x)^3 - (70*I + 70)*sqrt(2)*(-1)^(
1/8)*cosh(x)^4*sinh(x)^4 - (56*I + 56)*sqrt(2)*(-1)^(1/8)*cosh(x)^3*sinh(x
)^5 - (28*I + 28)*sqrt(2)*(-1)^(1/8)*cosh(x)^2*sinh(x)^6 - (8*I + 8)*sqrt(
2)*(-1)^(1/8)*cosh(x)*sinh(x)^7 - (I + 1)*sqrt(2)*(-1)^(1/8)*sinh(x)^8 - (
I + 1)*sqrt(2)*(-1)^(1/8))*log((I + 1)*sqrt(2)*(-1)^(1/8) + 2*cosh(x) + 2*
sinh(x)) + ((I - 1)*sqrt(2)*(-1)^(1/8)*cosh(x)^8 + (8*I - 8)*sqrt(2)*(-1)^(
1/8)*cosh(x)^7*sinh(x) + (28*I - 28)*sqrt(2)*(-1)^(1/8)*cosh(x)^6*sinh(x)
^2 + (56*I - 56)*sqrt(2)*(-1)^(1/8)*cosh(x)^5*sinh(x)^3 + (70*I - 70)*sqrt
(2)*(-1)^(1/8)*cosh(x)^4*sinh(x)^4 + (56*I - 56)*sqrt(2)*(-1)^(1/8)*cosh(x
)^3*sinh(x)^5 + (28*I - 28)*sqrt(2)*(-1)^(1/8)*cosh(x)^2*sinh(x)^6 + (8*I
- 8)*sqrt(2)*(-1)^(1/8)*cosh(x)*sinh(x)^7 + (I - 1)*sqrt(2)*(-1)^(1/8)*sin
h(x)^8 + (I - 1)*sqrt(2)*(-1)^(1/8))*log(-(I - 1)*sqrt(2)*(-1)^(1/8) + 2*c
osh(x) + 2*sinh(x)) + (-(I - 1)*sqrt(2)*(-1)^(1/8)*cosh(x)^8 - (8*I - 8)*s
qrt(2)*(-1)^(1/8)*cosh(x)^7*sinh(x) - (28*I - 28)*sqrt(2)*(-1)^(1/8)*cosh(
x)^6*sinh(x)^2 - (56*I - 56)*sqrt(2)*(-1)^(1/8)*cosh(x)^5*sinh(x)^3 - (...

```

## Sympy [F]

$$\int e^x \tanh^2(4x) dx = \int e^x \tanh^2(4x) dx$$

input

```
integrate(exp(x)*tanh(4*x)**2,x)
```

output

```
Integral(exp(x)*tanh(4*x)**2, x)
```

**Maxima [F]**

$$\int e^x \tanh^2(4x) dx = \int e^x \tanh(4x)^2 dx$$

input `integrate(exp(x)*tanh(4*x)^2,x, algorithm="maxima")`

output `1/2*(2*e^(9*x) + 3*e^x)/(e^(8*x) + 1) - integrate(1/2*e^x/(e^(8*x) + 1), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.89

$$\begin{aligned} \int e^x \tanh^2(4x) dx = & -\frac{1}{16} \sqrt{-\sqrt{2} + 2} \arctan\left(\frac{\sqrt{\sqrt{2} + 2} + 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\ & -\frac{1}{16} \sqrt{-\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{\sqrt{2} + 2} - 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\ & -\frac{1}{16} \sqrt{\sqrt{2} + 2} \arctan\left(\frac{\sqrt{-\sqrt{2} + 2} + 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \\ & -\frac{1}{16} \sqrt{\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{-\sqrt{2} + 2} - 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \\ & -\frac{1}{32} \sqrt{\sqrt{2} + 2} \log\left(\sqrt{\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\ & +\frac{1}{32} \sqrt{\sqrt{2} + 2} \log\left(-\sqrt{\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\ & -\frac{1}{32} \sqrt{-\sqrt{2} + 2} \log\left(\sqrt{-\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\ & +\frac{1}{32} \sqrt{-\sqrt{2} + 2} \log\left(-\sqrt{-\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\ & +\frac{e^x}{2(e^{(8x)} + 1)} + e^x \end{aligned}$$

input `integrate(exp(x)*tanh(4*x)^2,x, algorithm="giac")`

output

```
-1/16*sqrt(-sqrt(2) + 2)*arctan((sqrt(sqrt(2) + 2) + 2*e^x)/sqrt(-sqrt(2)
+ 2)) - 1/16*sqrt(-sqrt(2) + 2)*arctan(-(sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-
sqrt(2) + 2)) - 1/16*sqrt(sqrt(2) + 2)*arctan((sqrt(-sqrt(2) + 2) + 2*e^x)
/sqrt(sqrt(2) + 2)) - 1/16*sqrt(sqrt(2) + 2)*arctan(-(sqrt(-sqrt(2) + 2) -
2*e^x)/sqrt(sqrt(2) + 2)) - 1/32*sqrt(sqrt(2) + 2)*log(sqrt(sqrt(2) + 2)*
e^x + e^(2*x) + 1) + 1/32*sqrt(sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2)*e^x + e
^(2*x) + 1) - 1/32*sqrt(-sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x)
+ 1) + 1/32*sqrt(-sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1)
+ 1/2*e^x/(e^(8*x) + 1) + e^x
```

### Mupad [B] (verification not implemented)

Time = 5.21 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.60

$$\int e^x \tanh^2(4x) dx = \text{Too large to display}$$

input

```
int(tanh(4*x)^2*exp(x), x)
```

output

```
exp(x) + exp(x)/(2*(exp(8*x) + 1)) + log(exp(x)/2 - (2^(1/2) + 2)^(1/2)/4
- ((2 - 2^(1/2))^(1/2)*1i)/4)*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)
*1i)/32) - log(exp(x)/2 + (2^(1/2) + 2)^(1/2)/4 + ((2 - 2^(1/2))^(1/2)*1
i)/4)*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32) + log(exp(x)/
2 - ((2^(1/2) + 2)^(1/2)*1i)/4 + (2 - 2^(1/2))^(1/2)/4)*(((2^(1/2) + 2)^(1
/2)*1i)/32 - (2 - 2^(1/2))^(1/2)/32) - log(exp(x)/2 + ((2^(1/2) + 2)^(1/2)
*1i)/4 - (2 - 2^(1/2))^(1/2)/4)*(((2^(1/2) + 2)^(1/2)*1i)/32 - (2 - 2^(1/2)
))^(1/2)/32) + 2^(1/2)*log(exp(x)/2 - 2^(1/2)*((2^(1/2) + 2)^(1/2)/32 + ((
2 - 2^(1/2))^(1/2)*1i)/32)*(4 + 4i))*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/
2))^(1/2)*1i)/32)*(1/2 + 1i/2) + 2^(1/2)*log(exp(x)/2 - 2^(1/2)*((2^(1/2)
+ 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32)*(4 - 4i))*((2^(1/2) + 2)^(1/2)
)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32)*(1/2 - 1i/2) - 2^(1/2)*log(exp(x)/2 +
2^(1/2)*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32)*(4 - 4i))*
(2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32)*(1/2 - 1i/2) - 2^(1/
2)*log(exp(x)/2 + 2^(1/2)*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1
i)/32)*(4 + 4i))*((2^(1/2) + 2)^(1/2)/32 + ((2 - 2^(1/2))^(1/2)*1i)/32)*(1
/2 + 1i/2)
```

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 513, normalized size of antiderivative = 1.73

$$\int e^x \tanh^2(4x) dx$$

$$= \frac{2e^{8x} \sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right) + 2\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right) - 2e^{8x} \sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}+2e^x}{\sqrt{\sqrt{2}+2}}\right)}{1}$$

input

```
int(exp(x)*tanh(4*x)^2,x)
```

output

```
(2*e**(8*x)*sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) - 2*e**x)/sqrt(sqrt(2) + 2)) + 2*sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) - 2*e**x)/sqrt(sqrt(2) + 2)) - 2*e**(8*x)*sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) + 2*e**x)/sqrt(sqrt(2) + 2)) - 2*sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) + 2*e**x)/sqrt(sqrt(2) + 2)) + 2*e**(8*x)*sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) - 2*e**x)/sqrt(-sqrt(2) + 2)) + 2*sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) - 2*e**x)/sqrt(-sqrt(2) + 2)) - 2*e**(8*x)*sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) + 2*e**x)/sqrt(-sqrt(2) + 2)) - 2*sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) + 2*e**x)/sqrt(-sqrt(2) + 2)) + e**(8*x)*sqrt(-sqrt(2) + 2)*log(-e**x*sqrt(-sqrt(2) + 2) + e**(2*x) + 1) - e**(8*x)*sqrt(-sqrt(2) + 2)*log(e**x*sqrt(-sqrt(2) + 2) + e**(2*x) + 1) + sqrt(-sqrt(2) + 2)*log(-e**x*sqrt(-sqrt(2) + 2) + e**(2*x) + 1) - sqrt(-sqrt(2) + 2)*log(e**x*sqrt(-sqrt(2) + 2) + e**(2*x) + 1) + e**(8*x)*sqrt(sqrt(2) + 2)*log(-e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) - e**(8*x)*sqrt(sqrt(2) + 2)*log(e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) + sqrt(sqrt(2) + 2)*log(-e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) - sqrt(sqrt(2) + 2)*log(e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) + 32*e**(9*x) + 48*e**x)/(32*(e**(8*x) + 1))
```

### 3.234 $\int e^x \tanh(4x) dx$

Optimal result	1716
Mathematica [C] (verified)	1717
Rubi [A] (verified)	1717
Maple [C] (verified)	1721
Fricas [C] (verification not implemented)	1721
Sympy [F]	1722
Maxima [F]	1722
Giac [A] (verification not implemented)	1723
Mupad [B] (verification not implemented)	1724
Reduce [B] (verification not implemented)	1725

#### Optimal result

Integrand size = 8, antiderivative size = 280

$$\begin{aligned}
 \int e^x \tanh(4x) dx = & e^x + \frac{1}{4} \sqrt{2 + \sqrt{2}} \arctan \left( \frac{\sqrt{2 - \sqrt{2}} - 2e^x}{\sqrt{2 + \sqrt{2}}} \right) \\
 & + \frac{1}{4} \sqrt{2 - \sqrt{2}} \arctan \left( \frac{\sqrt{2 + \sqrt{2}} - 2e^x}{\sqrt{2 - \sqrt{2}}} \right) \\
 & - \frac{1}{4} \sqrt{2 + \sqrt{2}} \arctan \left( \frac{\sqrt{2 - \sqrt{2}} + 2e^x}{\sqrt{2 + \sqrt{2}}} \right) \\
 & - \frac{1}{4} \sqrt{2 - \sqrt{2}} \arctan \left( \frac{\sqrt{2 + \sqrt{2}} + 2e^x}{\sqrt{2 - \sqrt{2}}} \right) \\
 & - \frac{1}{4} \sqrt{2 - \sqrt{2}} \operatorname{arctanh} \left( \frac{\sqrt{2 - \sqrt{2}} e^x}{1 + e^{2x}} \right) \\
 & - \frac{1}{4} \sqrt{2 + \sqrt{2}} \operatorname{arctanh} \left( \frac{\sqrt{2 + \sqrt{2}} e^x}{1 + e^{2x}} \right)
 \end{aligned}$$

output

```
exp(x)+1/4*(2+2^(1/2))^(1/2)*arctan(((2-2^(1/2))^(1/2)-2*exp(x))/(2+2^(1/2))^(1/2))+1/4*(2-2^(1/2))^(1/2)*arctan(((2+2^(1/2))^(1/2)-2*exp(x))/(2-2^(1/2))^(1/2))-1/4*(2+2^(1/2))^(1/2)*arctan(((2-2^(1/2))^(1/2)+2*exp(x))/(2+2^(1/2))^(1/2))-1/4*(2-2^(1/2))^(1/2)*arctan(((2+2^(1/2))^(1/2)+2*exp(x))/(2-2^(1/2))^(1/2))-1/4*(2-2^(1/2))^(1/2)*arctanh((2-2^(1/2))^(1/2)*exp(x)/(1+exp(2*x)))-1/4*(2+2^(1/2))^(1/2)*arctanh((2+2^(1/2))^(1/2)*exp(x)/(1+exp(2*x)))
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.09

$$\int e^x \tanh(4x) dx = e^x - 2e^x \operatorname{Hypergeometric2F1}\left(\frac{1}{8}, 1, \frac{9}{8}, -e^{8x}\right)$$

input

```
Integrate[E^x*Tanh[4*x],x]
```

output

```
E^x - 2*E^x*Hypergeometric2F1[1/8, 1, 9/8, -E^(8*x)]
```

**Rubi [A] (verified)**

Time = 1.11 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.33, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$ , Rules used = {2720, 25, 913, 757, 1483, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x \tanh(4x) dx$$

↓ 2720

$$\int -\frac{1 - e^{8x}}{e^{8x} + 1} de^x$$

$$\begin{aligned}
& \downarrow 25 \\
& - \int \frac{1 - e^{8x}}{1 + e^{8x}} dx \\
& \downarrow 913 \\
& e^x - 2 \int \frac{1}{1 + e^{8x}} dx \\
& \downarrow 757 \\
& e^x - 2 \left( \frac{\int \frac{\sqrt{2} - e^{2x}}{1 - \sqrt{2}e^{2x} + e^{4x}} dx}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2} + e^{2x}}{1 + \sqrt{2}e^{2x} + e^{4x}} dx}{2\sqrt{2}} \right) \\
& \downarrow 1483 \\
& 2 \left( \frac{e^x - \frac{\int \frac{\sqrt{2(2-\sqrt{2}) + (1-\sqrt{2})e^x}}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\int \frac{\sqrt{2(2-\sqrt{2}) - (1-\sqrt{2})e^x}}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} + \frac{\frac{\int \frac{\sqrt{2(2+\sqrt{2}) - (1+\sqrt{2})e^x}}{1 - \sqrt{2} + \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2+\sqrt{2}}} + \frac{\int \frac{\sqrt{2(2+\sqrt{2}) + (1+\sqrt{2})e^x}}{1 + \sqrt{2} + \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2+\sqrt{2}}}}{2\sqrt{2}} \right) \\
& \downarrow 1142 \\
& 2 \left( \frac{e^x - \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx + \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}-2e^x}}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}+2e^x}}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} \right) \\
& \downarrow 25 \\
& 2 \left( \frac{e^x - \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}-2e^x}}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}} + \frac{\frac{1}{2}\sqrt{2+\sqrt{2}} \int \frac{1}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}+2e^x}}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} \right) \\
& \downarrow 1083 \\
& 2 \left( \frac{e^x - \frac{-\sqrt{2+\sqrt{2}} \int \frac{1}{-2 - \sqrt{2} - e^{2x}} d(-\sqrt{2-\sqrt{2}+2e^x}) - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}-2e^x}}{1 - \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}} + \frac{-\sqrt{2+\sqrt{2}} \int \frac{1}{-2 - \sqrt{2} - e^{2x}} d(\sqrt{2-\sqrt{2}+2e^x}) - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}+2e^x}}{1 + \sqrt{2} - \sqrt{2}e^x + e^{2x}} dx}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} \right)
\end{aligned}$$

$$\begin{aligned}
 & \downarrow 217 \\
 & 2 \left( \frac{\frac{\arctan\left(\frac{2e^x - \sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}} - 2e^x}{1 - \sqrt{2-\sqrt{2}}e^x + e^{2x}} de^x}{2\sqrt{2-\sqrt{2}}} + \frac{\arctan\left(\frac{2e^x + \sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \int \frac{\sqrt{2-\sqrt{2}} + 2e^x}{1 + \sqrt{2-\sqrt{2}}e^x + e^{2x}} de^x}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} + \frac{\frac{1}{2}(1+\sqrt{2}) \int \dots}{\dots} \right) \\
 & \downarrow 1103 \\
 & 2 \left( \frac{\frac{\arctan\left(\frac{2e^x - \sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) + \frac{1}{2}(1-\sqrt{2}) \log\left(-\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1\right)}{2\sqrt{2-\sqrt{2}}} + \frac{\arctan\left(\frac{2e^x + \sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right) - \frac{1}{2}(1-\sqrt{2}) \log\left(\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1\right)}{2\sqrt{2-\sqrt{2}}}}{2\sqrt{2}} + \frac{\arctan\left(\dots\right)}{\dots} \right)
 \end{aligned}$$

```
input Int [E^x*Tanh [4*x] , x]
```

```
output E^x - 2*(((ArcTan[(-Sqrt[2 - Sqrt[2]] + 2*E^x)/Sqrt[2 + Sqrt[2]]] + ((1 - Sqrt[2])*Log[1 - Sqrt[2 - Sqrt[2]]*E^x + E^(2*x)]))/2)/(2*Sqrt[2 - Sqrt[2]]) + (ArcTan[(Sqrt[2 - Sqrt[2]] + 2*E^x)/Sqrt[2 + Sqrt[2]]] - ((1 - Sqrt[2])*Log[1 + Sqrt[2 - Sqrt[2]]*E^x + E^(2*x)]))/2)/(2*Sqrt[2 - Sqrt[2]]))/(2*Sqrt[2]) + ((ArcTan[(-Sqrt[2 + Sqrt[2]] + 2*E^x)/Sqrt[2 - Sqrt[2]]] - ((1 + Sqrt[2])*Log[1 - Sqrt[2 + Sqrt[2]]*E^x + E^(2*x)]))/2)/(2*Sqrt[2 + Sqrt[2]]) + (ArcTan[(Sqrt[2 + Sqrt[2]] + 2*E^x)/Sqrt[2 - Sqrt[2]]] + ((1 + Sqrt[2])*Log[1 + Sqrt[2 + Sqrt[2]]*E^x + E^(2*x)]))/2)/(2*Sqrt[2 + Sqrt[2]]))/(2*Sqrt[2]))
```

**Defintions of rubi rules used**

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(- (Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```



rule 757  $\text{Int}[(a_ + (b_)(x_)^{(n_)} )^{-1}, x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 4]], s = \text{Denominator}[\text{Rt}[a/b, 4]]\}, \text{Simp}[r/(2*\text{Sqrt}[2]*a) \text{Int}[(\text{Sqrt}[2]*r - s*x^{(n/4)})/(r^2 - \text{Sqrt}[2]*r*s*x^{(n/4)} + s^2*x^{(n/2)}), x], x] + \text{Simp}[r/(2*\text{Sqrt}[2]*a) \text{Int}[(\text{Sqrt}[2]*r + s*x^{(n/4)})/(r^2 + \text{Sqrt}[2]*r*s*x^{(n/4)} + s^2*x^{(n/2)}), x], x]] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[n/4, 1] \ \&\& \ \text{GtQ}[a/b, 0]$

rule 913  $\text{Int}[(a_ + (b_)(x_)^{(n_)} )^{(p_)} ((c_ + (d_)(x_)^{(n_)})), x\_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^n)^{(p + 1)})/(b*(n*(p + 1) + 1)), x] - \text{Simp}[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n*(p + 1) + 1, 0]$

rule 1083  $\text{Int}[(a_ + (b_)(x_ + (c_)(x_)^2))^{-1}, x\_Symbol] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d_ + (e_)(x_))/((a_ + (b_)(x_ + (c_)(x_)^2)), x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142  $\text{Int}[(d_ + (e_)(x_))/((a_ + (b_)(x_ + (c_)(x_)^2)), x\_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x]$

rule 1483  $\text{Int}[(d_ + (e_)(x_)^2)/((a_ + (b_)(x_)^2 + (c_)(x_)^4), x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a/c, 2]\}, \text{With}[\{r = \text{Rt}[2*q - b/c, 2]\}, \text{Simp}[1/(2*c*q*r) \text{Int}[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + \text{Simp}[1/(2*c*q*r) \text{Int}[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NegQ}[b^2 - 4*a*c]$

rule 2720  $\text{Int}[u_, x\_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (w_)*((a_)(v_)^{(n_)} )^{(m_)} /; \text{FreeQ}\{a, m, n\}, x \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ !\text{MatchQ}[u, E^((c_)*((a_ + (b_)*x))*(F_)[v_]) /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{InverseFunctionQ}[F[x]]]$

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.54 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.09

method	result	size
risch	$e^x + \left( \sum_{_R=\text{RootOf}(65536\_Z^8+1)} \_R \ln(e^x - 4\_R) \right)$	24

input `int(exp(x)*tanh(4*x), x, method=_RETURNVERBOSE)`

output `exp(x)+sum(_R*ln(exp(x)-4*_R), _R=RootOf(65536*_Z^8+1))`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.61

$$\begin{aligned}
 \int e^x \tanh(4x) dx = & -\left(\frac{1}{8}i + \frac{1}{8}\right) \sqrt{2}(-1)^{\frac{1}{8}} \log\left(\left(i + 1\right) \sqrt{2}(-1)^{\frac{1}{8}} + 2 \cosh(x) \right. \\
 & \left. + 2 \sinh(x)\right) + \left(\frac{1}{8}i - \frac{1}{8}\right) \sqrt{2}(-1)^{\frac{1}{8}} \log\left(-\left(i - 1\right) \sqrt{2}(-1)^{\frac{1}{8}} \right. \\
 & \left. + 2 \cosh(x) + 2 \sinh(x)\right) - \left(\frac{1}{8}i \right. \\
 & \left. - \frac{1}{8}\right) \sqrt{2}(-1)^{\frac{1}{8}} \log\left(\left(i - 1\right) \sqrt{2}(-1)^{\frac{1}{8}} + 2 \cosh(x) + 2 \sinh(x)\right) \\
 & + \left(\frac{1}{8}i + \frac{1}{8}\right) \sqrt{2}(-1)^{\frac{1}{8}} \log\left(-\left(i + 1\right) \sqrt{2}(-1)^{\frac{1}{8}} + 2 \cosh(x) \right. \\
 & \left. + 2 \sinh(x)\right) - \frac{1}{4}(-1)^{\frac{1}{8}} \log\left(\left(-1\right)^{\frac{1}{8}} + \cosh(x) + \sinh(x)\right) \\
 & - \frac{1}{4}i(-1)^{\frac{1}{8}} \log\left(i(-1)^{\frac{1}{8}} + \cosh(x) + \sinh(x)\right) \\
 & + \frac{1}{4}i(-1)^{\frac{1}{8}} \log\left(-i(-1)^{\frac{1}{8}} + \cosh(x) + \sinh(x)\right) \\
 & + \frac{1}{4}(-1)^{\frac{1}{8}} \log\left(-\left(-1\right)^{\frac{1}{8}} + \cosh(x) + \sinh(x)\right) + \cosh(x) + \sinh(x)
 \end{aligned}$$

input `integrate(exp(x)*tanh(4*x),x, algorithm="fricas")`

output `-(1/8*I + 1/8)*sqrt(2)*(-1)^(1/8)*log((I + 1)*sqrt(2)*(-1)^(1/8) + 2*cosh(x) + 2*sinh(x)) + (1/8*I - 1/8)*sqrt(2)*(-1)^(1/8)*log(-(I - 1)*sqrt(2)*(-1)^(1/8) + 2*cosh(x) + 2*sinh(x)) - (1/8*I - 1/8)*sqrt(2)*(-1)^(1/8)*log((I - 1)*sqrt(2)*(-1)^(1/8) + 2*cosh(x) + 2*sinh(x)) + (1/8*I + 1/8)*sqrt(2)*(-1)^(1/8)*log(-(I + 1)*sqrt(2)*(-1)^(1/8) + 2*cosh(x) + 2*sinh(x)) - 1/4*(-1)^(1/8)*log((-1)^(1/8) + cosh(x) + sinh(x)) - 1/4*I*(-1)^(1/8)*log(I*(-1)^(1/8) + cosh(x) + sinh(x)) + 1/4*I*(-1)^(1/8)*log(-I*(-1)^(1/8) + cosh(x) + sinh(x)) + 1/4*(-1)^(1/8)*log(-(-1)^(1/8) + cosh(x) + sinh(x)) + cosh(x) + sinh(x)`

### Sympy [F]

$$\int e^x \tanh(4x) dx = \int e^x \tanh(4x) dx$$

input `integrate(exp(x)*tanh(4*x),x)`

output `Integral(exp(x)*tanh(4*x), x)`

### Maxima [F]

$$\int e^x \tanh(4x) dx = \int e^x \tanh(4x) dx$$

input `integrate(exp(x)*tanh(4*x),x, algorithm="maxima")`

output `e^x - 2*integrate(e^x/(e^(8*x) + 1), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.90

$$\begin{aligned}
\int e^x \tanh(4x) dx = & -\frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan\left(\frac{\sqrt{\sqrt{2} + 2} + 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\
& -\frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{\sqrt{2} + 2} - 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\
& -\frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan\left(\frac{\sqrt{-\sqrt{2} + 2} + 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \\
& -\frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{-\sqrt{2} + 2} - 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \\
& -\frac{1}{8} \sqrt{\sqrt{2} + 2} \log\left(\sqrt{\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\
& +\frac{1}{8} \sqrt{\sqrt{2} + 2} \log\left(-\sqrt{\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\
& -\frac{1}{8} \sqrt{-\sqrt{2} + 2} \log\left(\sqrt{-\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) \\
& +\frac{1}{8} \sqrt{-\sqrt{2} + 2} \log\left(-\sqrt{-\sqrt{2} + 2}e^x + e^{(2x)} + 1\right) + e^x
\end{aligned}$$

input `integrate(exp(x)*tanh(4*x),x, algorithm="giac")`

output `-1/4*sqrt(-sqrt(2) + 2)*arctan((sqrt(sqrt(2) + 2) + 2*e^x)/sqrt(-sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(-(sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqrt(2) + 2)) - 1/4*sqrt(sqrt(2) + 2)*arctan((sqrt(-sqrt(2) + 2) + 2*e^x)/sqrt(sqrt(2) + 2)) - 1/4*sqrt(sqrt(2) + 2)*arctan(-(sqrt(-sqrt(2) + 2) - 2*e^x)/sqrt(sqrt(2) + 2)) - 1/8*sqrt(sqrt(2) + 2)*log(sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/8*sqrt(sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) - 1/8*sqrt(-sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/8*sqrt(-sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + e^x`

**Mupad [B] (verification not implemented)**

Time = 4.84 (sec) , antiderivative size = 457, normalized size of antiderivative = 1.63

$$\begin{aligned}
& \int e^x \tanh(4x) dx = e^x \\
& -\ln\left(2e^x + \sqrt{\sqrt{2}+2} + \sqrt{2-\sqrt{2}}1i\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \\
& +\ln\left(2e^x + \sqrt{2-\sqrt{2}} - \sqrt{\sqrt{2}+2}1i\right) \left(-\frac{\sqrt{2-\sqrt{2}}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{\sqrt{2}+2}1i}{8}\right) \\
& +\ln\left(2e^x - \sqrt{\sqrt{2}+2} - \sqrt{2-\sqrt{2}}1i\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \\
& -\ln\left(2e^x - \sqrt{2-\sqrt{2}} + \sqrt{\sqrt{2}+2}1i\right) \left(-\frac{\sqrt{2-\sqrt{2}}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{\sqrt{2}+2}1i}{8}\right) + \sqrt{2} \ln\left(2e^x \right. \\
& \qquad \qquad \qquad \left. + \sqrt{2} \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) (-4-4i)\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \left(\frac{1}{2} + \frac{1}{2}i\right) + \sqrt{2} \ln\left(2e^x \right. \\
& \qquad \qquad \qquad \left. + \sqrt{2} \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) (-4+4i)\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \left(\frac{1}{2} - \frac{1}{2}i\right) + \sqrt{2} \ln\left(2e^x \right. \\
& \qquad \qquad \qquad \left. + \sqrt{2} \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) (4-4i)\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \left(-\frac{1}{2} + \frac{1}{2}i\right) + \sqrt{2} \ln\left(2e^x \right. \\
& \qquad \qquad \qquad \left. + \sqrt{2} \left(\frac{\sqrt{\sqrt{2}+2}}{8} + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) (4+4i)\right) \left(\frac{\sqrt{\sqrt{2}+2}}{8} \right. \\
& \qquad \qquad \qquad \left. + \frac{\sqrt{2-\sqrt{2}}1i}{8}\right) \left(-\frac{1}{2} - \frac{1}{2}i\right)
\end{aligned}$$

input `int(tanh(4*x)*exp(x),x)`

output

$$\begin{aligned} & \exp(x) - \log(2*\exp(x) + (2^{(1/2)} + 2)^{(1/2)} + (2 - 2^{(1/2)})^{(1/2)}*1i)*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8) + \log(2*\exp(x) - (2^{(1/2)} + 2)^{(1/2)}*1i + (2 - 2^{(1/2)})^{(1/2)})*((2^{(1/2)} + 2)^{(1/2)}*1i)/8 - (2 - 2^{(1/2)})^{(1/2)}/8) + \log(2*\exp(x) - (2^{(1/2)} + 2)^{(1/2)} - (2 - 2^{(1/2)})^{(1/2)}*1i)*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8) - \log(2*\exp(x) + (2^{(1/2)} + 2)^{(1/2)}*1i - (2 - 2^{(1/2)})^{(1/2)})*((2^{(1/2)} + 2)^{(1/2)}*1i)/8 - (2 - 2^{(1/2)})^{(1/2)}/8) + 2^{(1/2)}*\log(2*\exp(x) - 2^{(1/2)}*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(4 + 4i))*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(1/2 + 1i/2) + 2^{(1/2)}*\log(2*\exp(x) - 2^{(1/2)}*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(4 - 4i))*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(1/2 - 1i/2) - 2^{(1/2)}*\log(2*\exp(x) + 2^{(1/2)}*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(4 - 4i))*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(1/2 - 1i/2) - 2^{(1/2)}*\log(2*\exp(x) + 2^{(1/2)}*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(4 + 4i))*((2^{(1/2)} + 2)^{(1/2)}/8 + ((2 - 2^{(1/2)})^{(1/2)}*1i)/8)*(1/2 + 1i/2) \end{aligned}$$

### Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.82

$$\begin{aligned} \int e^x \tanh(4x) dx = & \frac{\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2-2e^x}}{\sqrt{\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2+2e^x}}{\sqrt{\sqrt{2}+2}}\right)}{4} \\ & + \frac{\sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2-2e^x}}{\sqrt{-\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2+2e^x}}{\sqrt{-\sqrt{2}+2}}\right)}{4} \\ & + \frac{\sqrt{-\sqrt{2} + 2} \log\left(-e^x \sqrt{-\sqrt{2} + 2} + e^{2x} + 1\right)}{8} \\ & - \frac{\sqrt{-\sqrt{2} + 2} \log\left(e^x \sqrt{-\sqrt{2} + 2} + e^{2x} + 1\right)}{8} \\ & + \frac{\sqrt{\sqrt{2} + 2} \log\left(-e^x \sqrt{\sqrt{2} + 2} + e^{2x} + 1\right)}{8} \\ & - \frac{\sqrt{\sqrt{2} + 2} \log\left(e^x \sqrt{\sqrt{2} + 2} + e^{2x} + 1\right)}{8} + e^x \end{aligned}$$

input `int(exp(x)*tanh(4*x),x)`

output 
$$\begin{aligned} & (2\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2} + 2} - 2e^{2x}}{\sqrt{\sqrt{2} + 2}}\right) - 2\sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2} + 2} + 2e^{2x}}{\sqrt{\sqrt{2} + 2}}\right) + 2\sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2} + 2} - 2e^{2x}}{\sqrt{-\sqrt{2} + 2}}\right) - 2\sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2} + 2} + 2e^{2x}}{\sqrt{-\sqrt{2} + 2}}\right) + \sqrt{-\sqrt{2} + 2} \log(-e^{2x}\sqrt{-\sqrt{2} + 2} + e^{4x} + 1) - \sqrt{-\sqrt{2} + 2} \log(e^{2x}\sqrt{-\sqrt{2} + 2} + e^{4x} + 1) + \sqrt{\sqrt{2} + 2} \log(-e^{2x}\sqrt{\sqrt{2} + 2} + e^{4x} + 1) - \sqrt{\sqrt{2} + 2} \log(e^{2x}\sqrt{\sqrt{2} + 2} + e^{4x} + 1) + 8e^{2x})/8 \end{aligned}$$

### 3.235 $\int e^x \coth(4x) dx$

Optimal result	1727
Mathematica [C] (verified)	1727
Rubi [A] (verified)	1728
Maple [C] (verified)	1732
Fricas [A] (verification not implemented)	1733
Sympy [F]	1733
Maxima [A] (verification not implemented)	1734
Giac [A] (verification not implemented)	1734
Mupad [B] (verification not implemented)	1735
Reduce [B] (verification not implemented)	1735

#### Optimal result

Integrand size = 8, antiderivative size = 91

$$\int e^x \coth(4x) dx = e^x - \frac{\arctan(e^x)}{2} + \frac{\arctan(1 - \sqrt{2}e^x)}{2\sqrt{2}} - \frac{\arctan(1 + \sqrt{2}e^x)}{2\sqrt{2}} - \frac{\operatorname{arctanh}(e^x)}{2} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^x}{1+e^{2x}}\right)}{2\sqrt{2}}$$

output

```
exp(x)-1/2*arctan(exp(x))-1/4*arctan(-1+2^(1/2)*exp(x))*2^(1/2)-1/4*arctan
(1+2^(1/2)*exp(x))*2^(1/2)-1/2*arctanh(exp(x))-1/4*arctanh(2^(1/2)*exp(x)/
(1+exp(2*x)))*2^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.24

$$\int e^x \coth(4x) dx = e^x - 2e^x \operatorname{Hypergeometric2F1}\left(\frac{1}{8}, 1, \frac{9}{8}, e^{8x}\right)$$

input

```
Integrate[E^x*Coth[4*x],x]
```



output

$$E^x - 2E^x \text{Hypergeometric2F1}[1/8, 1, 9/8, E^{(8*x)}]$$
**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.47, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.875$ , Rules used = {2720, 25, 913, 758, 755, 756, 216, 219, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \coth(4x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int -\frac{e^{8x} + 1}{1 - e^{8x}} de^x \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{1 + e^{8x}}{1 - e^{8x}} de^x \\
 & \quad \downarrow \text{913} \\
 & e^x - 2 \int \frac{1}{1 - e^{8x}} de^x \\
 & \quad \downarrow \text{758} \\
 & e^x - 2 \left( \frac{1}{2} \int \frac{1}{1 - e^{4x}} de^x + \frac{1}{2} \int \frac{1}{1 + e^{4x}} de^x \right) \\
 & \quad \downarrow \text{755} \\
 & e^x - 2 \left( \frac{1}{2} \int \frac{1}{1 - e^{4x}} de^x + \frac{1}{2} \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \int \frac{1 + e^{2x}}{1 + e^{4x}} de^x \right) \right) \\
 & \quad \downarrow \text{756} \\
 & e^x - 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{1 - e^{2x}} de^x + \frac{1}{2} \int \frac{1}{1 + e^{2x}} de^x \right) + \frac{1}{2} \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \int \frac{1 + e^{2x}}{1 + e^{4x}} de^x \right) \right) \\
 & \quad \downarrow \text{216}
 \end{aligned}$$

$$\begin{aligned}
& e^x - 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{1 - e^{2x}} de^x + \frac{\arctan(e^x)}{2} \right) + \frac{1}{2} \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \int \frac{1 + e^{2x}}{1 + e^{4x}} de^x \right) \right) \\
& \quad \downarrow \text{219} \\
& e^x - 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \int \frac{1 + e^{2x}}{1 + e^{4x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{1476} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{1 - \sqrt{2}e^x + e^{2x}} de^x + \frac{1}{2} \int \frac{1}{1 + \sqrt{2}e^x + e^{2x}} de^x \right) + \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{1082} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \left( \int \frac{1}{-1 - e^{2x}} d(1 - \sqrt{2}e^x) - \int \frac{1}{-1 - e^{2x}} d(1 + \sqrt{2}e^x) \right) + \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{217} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1 - e^{2x}}{1 + e^{4x}} de^x + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{1479} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \left( - \int \frac{\sqrt{2} - 2e^x}{1 - \sqrt{2}e^x + e^{2x}} de^x - \int \frac{\sqrt{2}(1 + \sqrt{2}e^x)}{1 + \sqrt{2}e^x + e^{2x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{25} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \left( \int \frac{\sqrt{2} - 2e^x}{1 - \sqrt{2}e^x + e^{2x}} de^x + \int \frac{\sqrt{2}(1 + \sqrt{2}e^x)}{1 + \sqrt{2}e^x + e^{2x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{27} \\
& 2 \left( \frac{1}{2} \left( \frac{1}{2} \left( \int \frac{\sqrt{2} - 2e^x}{1 - \sqrt{2}e^x + e^{2x}} de^x + \frac{1}{2} \int \frac{1 + \sqrt{2}e^x}{1 + \sqrt{2}e^x + e^{2x}} de^x \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) \right) \\
& \quad \downarrow \text{1103}
\end{aligned}$$

$$2 \left( \frac{1}{2} \left( \frac{\arctan(e^x)}{2} + \frac{\operatorname{arctanh}(e^x)}{2} \right) + \frac{1}{2} \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}e^x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^x)}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\sqrt{2}e^x + e^{2x} + 1)}{2\sqrt{2}} + \frac{\log(\sqrt{2}e^x - e^{2x} + 1)}{2\sqrt{2}} \right) \right) \right)$$

input `Int[E^x*Coth[4*x],x]`

output `E^x - 2*((ArcTan[E^x]/2 + ArcTanh[E^x]/2)/2 + ((-(ArcTan[1 - Sqrt[2]*E^x]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*E^x]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*E^x + E^(2*x)]/Sqrt[2] + Log[1 + Sqrt[2]*E^x + E^(2*x)]/(2*Sqrt[2]))/2)/2)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 758 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^(n/2)), x], x] + Simp[r/(2*a) Int[1/(r + s*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && !GtQ[a/b, 0]`

rule 913 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x)]*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.56 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.62

method	result	size
risch	$e^x + \left( \sum_{_R=\text{RootOf}(256\_Z^4+1)} \_R \ln(e^x - 4\_R) \right) + \frac{\ln(e^x-1)}{4} - \frac{\ln(e^x+1)}{4} + \frac{i \ln(e^x-i)}{4} - \frac{i \ln(e^x+i)}{4}$	56

input

```
int(exp(x)*coth(4*x), x, method=_RETURNVERBOSE)
```

output

```
exp(x)+sum(_R*ln(exp(x)-4*_R), _R=RootOf(256*_Z^4+1))+1/4*ln(exp(x)-1)-1/4*ln(exp(x)+1)+1/4*I*ln(exp(x)-I)-1/4*I*ln(exp(x)+I)
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.32

$$\begin{aligned}
\int e^x \coth(4x) dx = & -\frac{1}{4} \sqrt{2} \arctan \left( \sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) + 1 \right) \\
& -\frac{1}{4} \sqrt{2} \arctan \left( \sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) - 1 \right) \\
& -\frac{1}{8} \sqrt{2} \log \left( \frac{\sqrt{2} + 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) \\
& +\frac{1}{8} \sqrt{2} \log \left( -\frac{\sqrt{2} - 2 \cosh(x)}{\cosh(x) - \sinh(x)} \right) -\frac{1}{2} \arctan(\cosh(x) + \sinh(x)) \\
& + \cosh(x) - \frac{1}{4} \log(\cosh(x) + \sinh(x) + 1) \\
& + \frac{1}{4} \log(\cosh(x) + \sinh(x) - 1) + \sinh(x)
\end{aligned}$$

input `integrate(exp(x)*coth(4*x),x, algorithm="fricas")`output `-1/4*sqrt(2)*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) + 1) - 1/4*sqrt(2)*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - 1) - 1/8*sqrt(2)*log((sqrt(2) + 2*cosh(x))/(cosh(x) - sinh(x))) + 1/8*sqrt(2)*log(-(sqrt(2) - 2*cosh(x))/(cosh(x) - sinh(x))) - 1/2*arctan(cosh(x) + sinh(x)) + cosh(x) - 1/4*log(cosh(x) + sinh(x) + 1) + 1/4*log(cosh(x) + sinh(x) - 1) + sinh(x)`**Sympy [F]**

$$\int e^x \coth(4x) dx = \int e^x \coth(4x) dx$$

input `integrate(exp(x)*coth(4*x),x)`output `Integral(exp(x)*coth(4*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.07

$$\int e^x \coth(4x) dx = -\frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) - \frac{1}{4} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) - \frac{1}{8} \sqrt{2} \log(\sqrt{2}e^x + e^{(2x)} + 1) + \frac{1}{8} \sqrt{2} \log(-\sqrt{2}e^x + e^{(2x)} + 1) - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x + 1) + \frac{1}{4} \log(e^x - 1)$$

input `integrate(exp(x)*coth(4*x),x, algorithm="maxima")`

output

```
-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(e^x - 1)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.08

$$\int e^x \coth(4x) dx = -\frac{1}{4} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) - \frac{1}{4} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) - \frac{1}{8} \sqrt{2} \log(\sqrt{2}e^x + e^{(2x)} + 1) + \frac{1}{8} \sqrt{2} \log(-\sqrt{2}e^x + e^{(2x)} + 1) - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x + 1) + \frac{1}{4} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(4*x),x, algorithm="giac")`

output

```
-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(abs(e^x - 1))
```

**Mupad [B] (verification not implemented)**

Time = 2.46 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.14

$$\int e^x \coth(4x) dx = \frac{\ln(2 - 2e^x)}{4} - \frac{\ln(-2e^x - 2)}{4} - \frac{\operatorname{atan}(e^x)}{2} + e^x - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x - \sqrt{2})}{2}\right)}{4} + \frac{\sqrt{2} \ln\left((2e^x - \sqrt{2})^2 + 2\right)}{8} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x + \sqrt{2})}{2}\right)}{4} - \frac{\sqrt{2} \ln\left((2e^x + \sqrt{2})^2 + 2\right)}{8}$$

input

```
int(coth(4*x)*exp(x), x)
```

output

```
log(2 - 2*exp(x))/4 - log(- 2*exp(x) - 2)/4 - atan(exp(x))/2 + exp(x) - (2^(1/2)*atan((2^(1/2)*(2*exp(x) - 2^(1/2)))/2))/4 + (2^(1/2)*log((2*exp(x) - 2^(1/2))^2 + 2))/8 - (2^(1/2)*atan((2^(1/2)*(2*exp(x) + 2^(1/2)))/2))/4 - (2^(1/2)*log((2*exp(x) + 2^(1/2))^2 + 2))/8
```

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.11

$$\int e^x \coth(4x) dx = -\frac{\operatorname{atan}(e^x)}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right)}{4} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{4} + e^x + \frac{\sqrt{2} \log(e^{2x} - e^x \sqrt{2} + 1)}{8} - \frac{\sqrt{2} \log(e^{2x} + e^x \sqrt{2} + 1)}{8} + \frac{\log(e^x - 1)}{4} - \frac{\log(e^x + 1)}{4}$$



input `int(exp(x)*coth(4*x),x)`

output  $(-4*\operatorname{atan}(e^{**x}) - 2*\sqrt{2}*\operatorname{atan}((2*e^{**x} - \sqrt{2})/\sqrt{2}) - 2*\sqrt{2}*\operatorname{atan}((2*e^{**x} + \sqrt{2})/\sqrt{2}) + 8*e^{**x} + \sqrt{2}*\log(e^{**}(2*x) - e^{**x}*\sqrt{2} + 1) - \sqrt{2}*\log(e^{**}(2*x) + e^{**x}*\sqrt{2} + 1) + 2*\log(e^{**x} - 1) - 2*\log(e^{**x} + 1))/8$

### 3.236 $\int e^x \coth^2(4x) dx$

Optimal result	1737
Mathematica [C] (verified)	1737
Rubi [A] (verified)	1738
Maple [C] (verified)	1739
Fricas [B] (verification not implemented)	1740
Sympy [F]	1741
Maxima [A] (verification not implemented)	1741
Giac [A] (verification not implemented)	1742
Mupad [B] (verification not implemented)	1742
Reduce [B] (verification not implemented)	1743

#### Optimal result

Integrand size = 10, antiderivative size = 109

$$\int e^x \coth^2(4x) dx = e^x + \frac{e^x}{2(1 - e^{8x})} - \frac{\arctan(e^x)}{8} + \frac{\arctan(1 - \sqrt{2}e^x)}{8\sqrt{2}} - \frac{\arctan(1 + \sqrt{2}e^x)}{8\sqrt{2}} - \frac{\operatorname{arctanh}(e^x)}{8} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^x}{1+e^{2x}}\right)}{8\sqrt{2}}$$

output

```
exp(x)+exp(x)/(2-2*exp(8*x))-1/8*arctan(exp(x))-1/16*arctan(-1+2^(1/2)*exp(x))*2^(1/2)-1/16*arctan(1+2^(1/2)*exp(x))*2^(1/2)-1/8*arctanh(exp(x))-1/16*arctanh(2^(1/2)*exp(x)/(1+exp(2*x)))*2^(1/2)
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.23 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.04

$$\int e^x \coth^2(4x) dx = \frac{e^{-15x}(-44217 - 80225e^{8x} - 15127e^{16x} + 9361e^{24x} + 9(4913 + 8368e^{8x} + 1486e^{16x} - 1456e^{24x} + e^{32x})H_5(x))}{9216} + \frac{64e^{9x}(1 + e^{8x})^2 {}_4F_3\left(\frac{9}{8}, 2, 2, 2; 1, 1, \frac{33}{8}; e^{8x}\right)}{3825}$$

input `Integrate[E^x*Coth[4*x]^2,x]`

output  $(-44217 - 80225E^{(8*x)} - 15127E^{(16*x)} + 9361E^{(24*x)} + 9*(4913 + 8368*E^{(8*x)} + 1486E^{(16*x)} - 1456E^{(24*x)} + E^{(32*x)})*Hypergeometric2F1[1/8, 1, 9/8, E^{(8*x)}])/(9216E^{(15*x)}) + (64E^{(9*x)}*(1 + E^{(8*x)})^2*HypergeometricPFQ[{9/8, 2, 2, 2}, {1, 1, 33/8}, E^{(8*x)}])/3825$

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.23, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^x \coth^2(4x) dx \\
 & \quad \downarrow \text{2720} \\
 & \int \frac{(e^{8x} + 1)^2}{(1 - e^{8x})^2} de^x \\
 & \quad \downarrow \text{915} \\
 & \int \left( \frac{4e^{8x}}{(1 - e^{8x})^2} + 1 \right) de^x \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{8} \arctan(e^x) + \frac{\arctan(1 - \sqrt{2}e^x)}{8\sqrt{2}} - \frac{\arctan(\sqrt{2}e^x + 1)}{8\sqrt{2}} - \frac{\operatorname{arctanh}(e^x)}{8} + e^x + \frac{e^x}{2(1 - e^{8x})} + \\
 & \quad \frac{\log(-\sqrt{2}e^x + e^{2x} + 1)}{16\sqrt{2}} - \frac{\log(\sqrt{2}e^x + e^{2x} + 1)}{16\sqrt{2}}
 \end{aligned}$$

input `Int[E^x*Coth[4*x]^2,x]`

output

```
E^x + E^x/(2*(1 - E^(8*x))) - ArcTan[E^x]/8 + ArcTan[1 - Sqrt[2]*E^x]/(8*Sqrt[2]) - ArcTan[1 + Sqrt[2]*E^x]/(8*Sqrt[2]) - ArcTanh[E^x]/8 + Log[1 - Sqrt[2]*E^x + E^(2*x)]/(16*Sqrt[2]) - Log[1 + Sqrt[2]*E^x + E^(2*x)]/(16*Sqrt[2])
```

### Defintions of rubi rules used

rule 915

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2720

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

### Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.62 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.62

method	result
risch	$e^x - \frac{e^x}{2(e^{8x}-1)} + \frac{i \ln(e^x-i)}{16} - \frac{i \ln(e^x+i)}{16} + \left( \sum_{R=\text{RootOf}(65536_Z^4+1)} R \ln(e^x - 16_R) \right) + \frac{\ln(e^x-1)}{16} - \frac{\ln(e^x+1)}{16}$

input

```
int(exp(x)*coth(4*x)^2,x,method=_RETURNVERBOSE)
```

output

```
exp(x)-1/2*exp(x)/(exp(8*x)-1)+1/16*I*ln(exp(x)-I)-1/16*I*ln(exp(x)+I)+sum
(_R*ln(exp(x)-16*_R),_R=RootOf(65536*_Z^4+1))+1/16*ln(exp(x)-1)-1/16*ln(ex
p(x)+1)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 943 vs.  $2(73) = 146$ .

Time = 0.12 (sec) , antiderivative size = 943, normalized size of antiderivative = 8.65

$$\int e^x \coth^2(4x) dx = \text{Too large to display}$$

input

```
integrate(exp(x)*coth(4*x)^2,x, algorithm="fricas")
```

output

```
1/32*(32*cosh(x)^9 + 1152*cosh(x)^7*sinh(x)^2 + 2688*cosh(x)^6*sinh(x)^3 +
4032*cosh(x)^5*sinh(x)^4 + 4032*cosh(x)^4*sinh(x)^5 + 2688*cosh(x)^3*sinh
(x)^6 + 1152*cosh(x)^2*sinh(x)^7 + 288*cosh(x)*sinh(x)^8 + 32*sinh(x)^9 -
2*(sqrt(2)*cosh(x)^8 + 8*sqrt(2)*cosh(x)^7*sinh(x) + 28*sqrt(2)*cosh(x)^6*
sinh(x)^2 + 56*sqrt(2)*cosh(x)^5*sinh(x)^3 + 70*sqrt(2)*cosh(x)^4*sinh(x)^
4 + 56*sqrt(2)*cosh(x)^3*sinh(x)^5 + 28*sqrt(2)*cosh(x)^2*sinh(x)^6 + 8*sq
rt(2)*cosh(x)*sinh(x)^7 + sqrt(2)*sinh(x)^8 - sqrt(2))*arctan(sqrt(2)*cosh
(x) + sqrt(2)*sinh(x) + 1) - 2*(sqrt(2)*cosh(x)^8 + 8*sqrt(2)*cosh(x)^7*si
nh(x) + 28*sqrt(2)*cosh(x)^6*sinh(x)^2 + 56*sqrt(2)*cosh(x)^5*sinh(x)^3 +
70*sqrt(2)*cosh(x)^4*sinh(x)^4 + 56*sqrt(2)*cosh(x)^3*sinh(x)^5 + 28*sqrt(
2)*cosh(x)^2*sinh(x)^6 + 8*sqrt(2)*cosh(x)*sinh(x)^7 + sqrt(2)*sinh(x)^8 -
sqrt(2))*arctan(sqrt(2)*cosh(x) + sqrt(2)*sinh(x) - 1) - 4*(cosh(x)^8 + 8
*cosh(x)^7*sinh(x) + 28*cosh(x)^6*sinh(x)^2 + 56*cosh(x)^5*sinh(x)^3 + 70*
cosh(x)^4*sinh(x)^4 + 56*cosh(x)^3*sinh(x)^5 + 28*cosh(x)^2*sinh(x)^6 + 8*
cosh(x)*sinh(x)^7 + sinh(x)^8 - 1)*arctan(cosh(x) + sinh(x)) - (sqrt(2)*co
sh(x)^8 + 8*sqrt(2)*cosh(x)^7*sinh(x) + 28*sqrt(2)*cosh(x)^6*sinh(x)^2 + 5
6*sqrt(2)*cosh(x)^5*sinh(x)^3 + 70*sqrt(2)*cosh(x)^4*sinh(x)^4 + 56*sqrt(2)
)*cosh(x)^3*sinh(x)^5 + 28*sqrt(2)*cosh(x)^2*sinh(x)^6 + 8*sqrt(2)*cosh(x)
*sinh(x)^7 + sqrt(2)*sinh(x)^8 - sqrt(2))*log((sqrt(2) + 2*cosh(x))/(cosh(
x) - sinh(x))) + (sqrt(2)*cosh(x)^8 + 8*sqrt(2)*cosh(x)^7*sinh(x) + 28*...
```

**Sympy [F]**

$$\int e^x \coth^2(4x) dx = \int e^x \coth^2(4x) dx$$

input `integrate(exp(x)*coth(4*x)**2,x)`

output `Integral(exp(x)*coth(4*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00

$$\begin{aligned} \int e^x \coth^2(4x) dx = & -\frac{1}{16} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) \\ & - \frac{1}{16} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) \\ & - \frac{1}{32} \sqrt{2} \log\left(\sqrt{2}e^x + e^{(2x)} + 1\right) \\ & + \frac{1}{32} \sqrt{2} \log\left(-\sqrt{2}e^x + e^{(2x)} + 1\right) - \frac{e^x}{2(e^{(8x)} - 1)} \\ & - \frac{1}{8} \arctan(e^x) + e^x - \frac{1}{16} \log(e^x + 1) + \frac{1}{16} \log(e^x - 1) \end{aligned}$$

input `integrate(exp(x)*coth(4*x)^2,x, algorithm="maxima")`

output `-1/16*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/16*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/32*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/32*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*e^x/(e^(8*x) - 1) - 1/8*arctan(e^x) + e^x - 1/16*log(e^x + 1) + 1/16*log(e^x - 1)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.01

$$\int e^x \coth^2(4x) dx = -\frac{1}{16} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2e^x)\right) - \frac{1}{16} \sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2e^x)\right) - \frac{1}{32} \sqrt{2} \log(\sqrt{2}e^x + e^{2x} + 1) + \frac{1}{32} \sqrt{2} \log(-\sqrt{2}e^x + e^{2x} + 1) - \frac{e^x}{2(e^{8x} - 1)} - \frac{1}{8} \arctan(e^x) + e^x - \frac{1}{16} \log(e^x + 1) + \frac{1}{16} \log(|e^x - 1|)$$

input `integrate(exp(x)*coth(4*x)^2,x, algorithm="giac")`output `-1/16*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/16*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/32*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/32*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*e^x/(e^(8*x) - 1) - 1/8*arctan(e^x) + e^x - 1/16*log(e^x + 1) + 1/16*log(abs(e^x - 1))`**Mupad [B] (verification not implemented)**

Time = 2.52 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.12

$$\int e^x \coth^2(4x) dx = \frac{\ln\left(\frac{1}{2} - \frac{e^x}{2}\right)}{16} - \frac{\ln\left(-\frac{e^x}{2} - \frac{1}{2}\right)}{16} - \frac{\operatorname{atan}(e^x)}{8} + e^x - \frac{e^x}{2(e^{8x} - 1)} - \frac{\sqrt{2} \operatorname{atan}\left(2\sqrt{2}\left(\frac{e^x}{2} - \frac{\sqrt{2}}{4}\right)\right)}{16} - \frac{\sqrt{2} \operatorname{atan}\left(2\sqrt{2}\left(\frac{e^x}{2} + \frac{\sqrt{2}}{4}\right)\right)}{16} + \frac{\sqrt{2} \ln\left(\left(\frac{e^x}{2} - \frac{\sqrt{2}}{4}\right)^2 + \frac{1}{8}\right)}{32} - \frac{\sqrt{2} \ln\left(\left(\frac{e^x}{2} + \frac{\sqrt{2}}{4}\right)^2 + \frac{1}{8}\right)}{32}$$

input `int(coth(4*x)^2*exp(x), x)`

output

```
log(1/2 - exp(x)/2)/16 - log(- exp(x)/2 - 1/2)/16 - atan(exp(x))/8 + exp(x)
) - exp(x)/(2*(exp(8*x) - 1)) - (2^(1/2)*atan(2*2^(1/2)*(exp(x)/2 - 2^(1/2)
)/4))/16 - (2^(1/2)*atan(2*2^(1/2)*(exp(x)/2 + 2^(1/2)/4)))/16 + (2^(1/2)
*log((exp(x)/2 - 2^(1/2)/4)^2 + 1/8))/32 - (2^(1/2)*log((exp(x)/2 + 2^(1/2)
)/4)^2 + 1/8))/32
```

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.31

$$\int e^x \coth^2(4x) dx$$

$$= \frac{-4e^{8x} \operatorname{atan}(e^x) + 4\operatorname{atan}(e^x) - 2e^{8x}\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) + 2\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - 2e^{8x}\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) + 2\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{32}$$

input

```
int(exp(x)*coth(4*x)^2,x)
```

output

```
( - 4*e**(8*x)*atan(e**x) + 4*atan(e**x) - 2*e**(8*x)*sqrt(2)*atan((2*e**x
- sqrt(2))/sqrt(2)) + 2*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - 2*e**
(8*x)*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) + 2*sqrt(2)*atan((2*e**x + s
qrt(2))/sqrt(2)) + 32*e**(9*x) + e**(8*x)*sqrt(2)*log(e**(2*x) - e**x*sqrt
(2) + 1) - e**(8*x)*sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1) + 2*e**(8*x)*
log(e**x - 1) - 2*e**(8*x)*log(e**x + 1) - 48*e**x - sqrt(2)*log(e**(2*x)
- e**x*sqrt(2) + 1) + sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1) - 2*log(e**
x - 1) + 2*log(e**x + 1))/(32*(e**(8*x) - 1))
```



### 3.237 $\int \frac{e^x}{a - \tanh(2x)} dx$

Optimal result	1744
Mathematica [A] (verified)	1744
Rubi [A] (verified)	1745
Maple [C] (verified)	1747
Fricas [C] (verification not implemented)	1747
Sympy [F]	1748
Maxima [F(-2)]	1748
Giac [B] (verification not implemented)	1749
Mupad [B] (verification not implemented)	1750
Reduce [B] (verification not implemented)	1750

#### Optimal result

Integrand size = 14, antiderivative size = 107

$$\int \frac{e^x}{a - \tanh(2x)} dx = -\frac{e^x}{1-a} + \frac{\arctan\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{1+a}}\right)}{(1-a)\sqrt{1+a}\sqrt[4]{1-a^2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{1+a}}\right)}{(1-a)\sqrt{1+a}\sqrt[4]{1-a^2}}$$

output

```
-exp(x)/(1-a)+arctan((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)/(1+a)^(1/2)/(-a
^2+1)^(1/4)+arctanh((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)/(1+a)^(1/2)/(-a
^2+1)^(1/4)
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.76

$$\int \frac{e^x}{a - \tanh(2x)} dx = \frac{-\sqrt[4]{1-a}(1+a)^{3/4}e^x + \arctan\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{1+a}}\right) + \operatorname{arctanh}\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{1+a}}\right)}{(1-a)^{5/4}(1+a)^{3/4}}$$

input

```
Integrate[E^x/(a - Tanh[2*x]), x]
```

output

$$\left( -((1-a)^{1/4}*(1+a)^{3/4}*E^x) + \text{ArcTan}[\frac{(1-a)^{1/4}*E^x}{(1+a)^{1/4}}] + \text{ArcTanh}[\frac{(1-a)^{1/4}*E^x}{(1+a)^{1/4}}] \right) / ((1-a)^{5/4}*(1+a)^{3/4})$$
**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {2720, 913, 756, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{e^x}{a - \tanh(2x)} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{4x} + 1}{(1-a)(-e^{4x}) + a + 1} de^x \\ & \quad \downarrow \text{913} \\ & \frac{2 \int \frac{1}{-e^{4x}(1-a)+a+1} de^x}{1-a} - \frac{e^x}{1-a} \\ & \quad \downarrow \text{756} \\ & \frac{2 \left( \frac{\int \frac{1}{\sqrt{a+1}-\sqrt{1-ae^{2x}}} de^x}{2\sqrt{a+1}} + \frac{\int \frac{1}{e^{2x}\sqrt{1-a+\sqrt{a+1}}} de^x}{2\sqrt{a+1}} \right)}{1-a} - \frac{e^x}{1-a} \\ & \quad \downarrow \text{218} \\ & \frac{2 \left( \frac{\int \frac{1}{\sqrt{a+1}-\sqrt{1-ae^{2x}}} de^x}{2\sqrt{a+1}} + \frac{\arctan\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}} \right)}{1-a} - \frac{e^x}{1-a} \\ & \quad \downarrow \text{221} \end{aligned}$$

$$\frac{2 \left( \frac{\arctan\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{a+1}}\right)}{2\sqrt{a+1}\sqrt[4]{1-a^2}} \right)}{1-a} - \frac{e^x}{1-a}$$

input `Int[E^x/(a - Tanh[2*x]),x]`

output `-(E^x/(1 - a)) + (2*(ArcTan[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)]/(2*sqrt[1 + a]*(1 - a^2)^(1/4)) + ArcTanh[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)]/(2*sqrt[1 + a]*(1 - a^2)^(1/4))))/(1 - a)`

### Definitions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 913 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Simp[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)) Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.94 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.65

method	result	s
risch	$\frac{e^x}{a-1} + \left( \sum_{_R=\text{RootOf}(1+(16a^8-32a^7-32a^6+96a^5-96a^3+32a^2+32a-16)_Z^4)} \_R \ln(e^x + (-2a^2 + 2)_R) \right)$	7

input

```
int(exp(x)/(a-tanh(2*x)),x,method=_RETURNVERBOSE)
```

output

```
exp(x)/(a-1)+sum(_R*ln(exp(x)+(-2*a^2+2)*_R),_R=RootOf(1+(16*a^8-32*a^7-32
*a^6+96*a^5-96*a^3+32*a^2+32*a-16)*_Z^4))
```

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 406, normalized size of antiderivative = 3.79

$$\int \frac{e^x}{a - \tanh(2x)} dx = \frac{(a-1) \left( -\frac{1}{a^8-2a^7-2a^6+6a^5-6a^3+2a^2+2a-1} \right)^{\frac{1}{4}} \log \left( (a^2-1) \left( -\frac{1}{a^8-2a^7-2a^6+6a^5-6a^3+2a^2+2a-1} \right)^{\frac{1}{4}} + \cosh(x) \right)}{1}$$

input

```
integrate(exp(x)/(a-tanh(2*x)),x, algorithm="fricas")
```

output

```
-1/2*((a - 1)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*a - 1))
^(1/4)*log((a^2 - 1)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*
a - 1))^(1/4) + cosh(x) + sinh(x)) - (a - 1)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*
a^5 - 6*a^3 + 2*a^2 + 2*a - 1))^(1/4)*log(-(a^2 - 1)*(-1/(a^8 - 2*a^7 - 2*
a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*a - 1))^(1/4) + cosh(x) + sinh(x)) - (I*a
- I)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*a - 1))^(1/4)*lo
g(-(I*a^2 - I)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*a - 1)
)^(1/4) + cosh(x) + sinh(x)) - (-I*a + I)*(-1/(a^8 - 2*a^7 - 2*a^6 + 6*a^5
- 6*a^3 + 2*a^2 + 2*a - 1))^(1/4)*log(-(-I*a^2 + I)*(-1/(a^8 - 2*a^7 - 2*
a^6 + 6*a^5 - 6*a^3 + 2*a^2 + 2*a - 1))^(1/4) + cosh(x) + sinh(x)) - 2*cos
h(x) - 2*sinh(x))/(a - 1)
```

**Sympy [F]**

$$\int \frac{e^x}{a - \tanh(2x)} dx = \int \frac{e^x}{a - \tanh(2x)} dx$$

input

```
integrate(exp(x)/(a-tanh(2*x)),x)
```

output

```
Integral(exp(x)/(a - tanh(2*x)), x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{e^x}{a - \tanh(2x)} dx = \text{Exception raised: ValueError}$$

input

```
integrate(exp(x)/(a-tanh(2*x)),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a-1>0)', see `assume?` for more
details)Is
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 328 vs. 2(83) = 166.

Time = 0.11 (sec) , antiderivative size = 328, normalized size of antiderivative = 3.07

$$\int \frac{e^x}{a - \tanh(2x)} dx = -\frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} + 2e^x\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2}} - \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} - 2e^x\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2}} - \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \log\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{2(\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2})} + \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}} \log\left(-\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{2(\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2})} + \frac{e^x}{a - 1}$$

input `integrate(exp(x)/(a-tanh(2*x)),x, algorithm="giac")`

output `-(a^4 - 2*a^3 + 2*a - 1)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*((a + 1)/(a - 1)))^(1/4) + 2*e^x)/((a + 1)/(a - 1))^(1/4)/(sqrt(2)*a^3 - sqrt(2)*a^2 - sqrt(2)*a + sqrt(2)) - (a^4 - 2*a^3 + 2*a - 1)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*((a + 1)/(a - 1))^(1/4) - 2*e^x)/((a + 1)/(a - 1))^(1/4))/(sqrt(2)*a^3 - sqrt(2)*a^2 - sqrt(2)*a + sqrt(2)) - 1/2*(a^4 - 2*a^3 + 2*a - 1)^(1/4)*log(sqrt(2)*((a + 1)/(a - 1))^(1/4)*e^x + sqrt((a + 1)/(a - 1)) + e^(2*x))/(sqrt(2)*a^3 - sqrt(2)*a^2 - sqrt(2)*a + sqrt(2)) + 1/2*(a^4 - 2*a^3 + 2*a - 1)^(1/4)*log(-sqrt(2)*((a + 1)/(a - 1))^(1/4)*e^x + sqrt((a + 1)/(a - 1)) + e^(2*x))/(sqrt(2)*a^3 - sqrt(2)*a^2 - sqrt(2)*a + sqrt(2)) + e^x/(a - 1)`

**Mupad [B] (verification not implemented)**

Time = 2.44 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.52

$$\int \frac{e^x}{a - \tanh(2x)} dx$$

$$= \frac{\ln \left( 8 a (-a - 1)^{1/4} + 8 e^x (a - 1)^{5/4} - 8 (-a - 1)^{1/4} \right) - \ln \left( 8 e^x (a - 1)^{5/4} - 8 a (-a - 1)^{1/4} + 8 (-a - 1)^{1/4} \right)}{2}$$

input `int(exp(x)/(a - tanh(2*x)),x)`output `(log(8*a*(- a - 1)^(1/4) + 8*exp(x)*(a - 1)^(5/4) - 8*(- a - 1)^(1/4)) - log(8*exp(x)*(a - 1)^(5/4) - 8*a*(- a - 1)^(1/4) + 8*(- a - 1)^(1/4)) - log(8*exp(x)*(a - 1)^(5/4) - a*(- a - 1)^(1/4)*8i + (- a - 1)^(1/4)*8i)*1i + log(a*(- a - 1)^(1/4)*8i + 8*exp(x)*(a - 1)^(5/4) - (- a - 1)^(1/4)*8i)*1i + 2*exp(x)*(a - 1)^(1/4)*(- a - 1)^(3/4))/(2*(a - 1)^(5/4)*(- a - 1)^(3/4))`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.15

$$\int \frac{e^x}{a - \tanh(2x)} dx$$

$$= \frac{2(a + 1)^{1/4} (a - 1)^{3/4} \sqrt{2} \operatorname{atan} \left( \frac{(a+1)^{1/4} (a-1)^{1/4} \sqrt{2} - 2e^x \sqrt{a-1}}{(a+1)^{1/4} (a-1)^{1/4} \sqrt{2}} \right) - 2(a + 1)^{1/4} (a - 1)^{3/4} \sqrt{2} \operatorname{atan} \left( \frac{(a+1)^{1/4} (a-1)^{1/4} \sqrt{2} + 2e^x}{(a+1)^{1/4} (a-1)^{1/4} \sqrt{2}} \right)}{2}$$

input `int(exp(x)/(a-tanh(2*x)),x)`

output

```
(2*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2))) - 2*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) + 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2))) + (a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*log(- e**x*(a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) + e**(2*x)*sqrt(a - 1) + sqrt(a + 1)) - (a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*log(e**x*(a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) + e**(2*x)*sqrt(a - 1) + sqrt(a + 1)) + 4*e**x*a**2 - 4*e**x)/(4*(a**3 - a**2 - a + 1))
```



### 3.238 $\int \frac{e^x}{(a - \tanh(2x))^2} dx$

Optimal result	1752
Mathematica [C] (verified)	1753
Rubi [A] (verified)	1753
Maple [C] (verified)	1755
Fricas [C] (verification not implemented)	1755
Sympy [F]	1756
Maxima [F(-2)]	1757
Giac [B] (verification not implemented)	1757
Mupad [B] (verification not implemented)	1758
Reduce [B] (verification not implemented)	1759

#### Optimal result

Integrand size = 14, antiderivative size = 155

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx = \frac{e^x}{(1 - a)^2} + \frac{e^x}{(1 - a)^2(1 + a)(1 + a - (1 - a)e^{4x})} - \frac{(1 + 4a) \arctan\left(\frac{\sqrt[4]{1 - ae^x}}{\sqrt[4]{1 + a}}\right)}{2(1 - a)^2(1 + a)^{3/2}\sqrt[4]{1 - a^2}} - \frac{(1 + 4a) \operatorname{arctanh}\left(\frac{\sqrt[4]{1 - ae^x}}{\sqrt[4]{1 + a}}\right)}{2(1 - a)^2(1 + a)^{3/2}\sqrt[4]{1 - a^2}}$$

output

```
exp(x)/(1-a)^2+exp(x)/(1-a)^2/(1+a)/(1+a-(1-a)*exp(4*x))-1/2*(1+4*a)*arctan((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)^2/(1+a)^(3/2)/(-a^2+1)^(1/4)-1/2*(1+4*a)*arctanh((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)^2/(1+a)^(3/2)/(-a^2+1)^(1/4)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.69

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx$$

$$= \frac{\frac{4(-1+a)e^x(2+2a-e^{4x}+a^2(1+e^{4x}))}{1+a-e^{4x}+ae^{4x}} + (1+4a)\text{RootSum}\left[1+a-\#1^4+a\#1^4\&, \frac{x-\log(e^x-\#1)}{\#1^3}\&\right]}{4(-1+a)^3(1+a)}$$

input `Integrate[E^x/(a - Tanh[2*x])^2,x]`

output `((4*(-1 + a)*E^x*(2 + 2*a - E^(4*x)) + a^2*(1 + E^(4*x)))/(1 + a - E^(4*x) + a*E^(4*x)) + (1 + 4*a)*RootSum[1 + a - #1^4 + a*#1^4 & , (x - Log[E^x - #1])/#1^3 & ])/(4*(-1 + a)^3*(1 + a))`

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {2720, 915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx$$

$$\downarrow \text{2720}$$

$$\int \frac{(e^{4x} + 1)^2}{((1 - a)(-e^{4x}) + a + 1)^2} de^x$$

$$\downarrow \text{915}$$

$$\int \left( \frac{1}{(a - 1)^2} - \frac{4(a - (1 - a)e^{4x})}{(a - 1)^2((a - 1)e^{4x} + a + 1)^2} \right) de^x$$

$$\begin{aligned} & \downarrow 2009 \\ & -\frac{(4a+1)\arctan\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{a+1}}\right)}{2(1-a)^2(a+1)^{3/2}\sqrt[4]{1-a^2}} - \frac{(4a+1)\operatorname{arctanh}\left(\frac{\sqrt[4]{1-ae^x}}{\sqrt[4]{a+1}}\right)}{2(1-a)^2(a+1)^{3/2}\sqrt[4]{1-a^2}} + \frac{e^x}{(1-a)^2} + \\ & \frac{e^x}{(1-a)^2(a+1)((1-a)(-e^{4x})+a+1)} \end{aligned}$$

input `Int[E^x/(a - Tanh[2*x])^2,x]`

output `E^x/(1 - a)^2 + E^x/((1 - a)^2*(1 + a)*(1 + a - (1 - a)*E^(4*x))) - ((1 + 4*a)*ArcTan[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)]/(2*(1 - a)^2*(1 + a)^(3/2)*(1 - a^2)^(1/4)) - ((1 + 4*a)*ArcTanh[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)]/(2*(1 - a)^2*(1 + a)^(3/2)*(1 - a^2)^(1/4)))`

### Defintions of rubi rules used

rule 915 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

**Maple [C] (verified)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.08 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.39

method	result
risch	$\frac{e^x}{a^2-2a+1} + \frac{e^x}{(a+1)(a^2-2a+1)(e^{4x}a-e^{4x}+a+1)} + \left( \text{--}_R=\text{RootOf}((256a^{16}-512a^{15}-1536a^{14}+3584a^{13}+3584a^{12}-10752a^{11}-536a^2+512a-256)*_Z^4+256a^4+256a^3+96a^2+16a+1) \right)$

input `int(exp(x)/(a-tanh(2*x))^2,x,method=_RETURNVERBOSE)`

output `exp(x)/(a^2-2*a+1)+1/(a+1)*exp(x)/(a^2-2*a+1)/(a*exp(x)^4-exp(x)^4+a+1)+sum(_R*ln(exp(x)+(-4/(1+4*a))*a^4+8/(1+4*a))*a^2-4/(1+4*a))/(4/(1+4*a)*a+1/(1+4*a))*_R),_R=RootOf((256*a^16-512*a^15-1536*a^14+3584*a^13+3584*a^12-10752*a^11-3584*a^10+17920*a^9-17920*a^7+3584*a^6+10752*a^5-3584*a^4-3584*a^3+1536*a^2+512*a-256)*_Z^4+256*a^4+256*a^3+96*a^2+16*a+1))`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 1604, normalized size of antiderivative = 10.35

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx = \text{Too large to display}$$

input `integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="fricas")`

output

```

1/4*(4*(a^2 - 1)*cosh(x)^5 + 40*(a^2 - 1)*cosh(x)^3*sinh(x)^2 + 40*(a^2 -
1)*cosh(x)^2*sinh(x)^3 + 20*(a^2 - 1)*cosh(x)*sinh(x)^4 + 4*(a^2 - 1)*sinh
(x)^5 - ((a^4 - 2*a^3 + 2*a - 1)*cosh(x)^4 + 4*(a^4 - 2*a^3 + 2*a - 1)*cos
h(x)^3*sinh(x) + 6*(a^4 - 2*a^3 + 2*a - 1)*cosh(x)^2*sinh(x)^2 + 4*(a^4 -
2*a^3 + 2*a - 1)*cosh(x)*sinh(x)^3 + (a^4 - 2*a^3 + 2*a - 1)*sinh(x)^4 + a
^4 - 2*a^2 + 1)*(-(256*a^4 + 256*a^3 + 96*a^2 + 16*a + 1)/(a^16 - 2*a^15 -
6*a^14 + 14*a^13 + 14*a^12 - 42*a^11 - 14*a^10 + 70*a^9 - 70*a^7 + 14*a^6
+ 42*a^5 - 14*a^4 - 14*a^3 + 6*a^2 + 2*a - 1))^(1/4)*log((4*a + 1)*cosh(x)
) + (4*a + 1)*sinh(x) + (a^4 - 2*a^2 + 1)*(-(256*a^4 + 256*a^3 + 96*a^2 +
16*a + 1)/(a^16 - 2*a^15 - 6*a^14 + 14*a^13 + 14*a^12 - 42*a^11 - 14*a^10
+ 70*a^9 - 70*a^7 + 14*a^6 + 42*a^5 - 14*a^4 - 14*a^3 + 6*a^2 + 2*a - 1))^(
1/4)) + ((a^4 - 2*a^3 + 2*a - 1)*cosh(x)^4 + 4*(a^4 - 2*a^3 + 2*a - 1)*co
sh(x)^3*sinh(x) + 6*(a^4 - 2*a^3 + 2*a - 1)*cosh(x)^2*sinh(x)^2 + 4*(a^4 -
2*a^3 + 2*a - 1)*cosh(x)*sinh(x)^3 + (a^4 - 2*a^3 + 2*a - 1)*sinh(x)^4 +
a^4 - 2*a^2 + 1)*(-(256*a^4 + 256*a^3 + 96*a^2 + 16*a + 1)/(a^16 - 2*a^15
- 6*a^14 + 14*a^13 + 14*a^12 - 42*a^11 - 14*a^10 + 70*a^9 - 70*a^7 + 14*a^
6 + 42*a^5 - 14*a^4 - 14*a^3 + 6*a^2 + 2*a - 1))^(1/4)*log((4*a + 1)*cosh(
x) + (4*a + 1)*sinh(x) - (a^4 - 2*a^2 + 1)*(-(256*a^4 + 256*a^3 + 96*a^2 +
16*a + 1)/(a^16 - 2*a^15 - 6*a^14 + 14*a^13 + 14*a^12 - 42*a^11 - 14*a^10
+ 70*a^9 - 70*a^7 + 14*a^6 + 42*a^5 - 14*a^4 - 14*a^3 + 6*a^2 + 2*a - ...

```

SymPy [F]

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx = \int \frac{e^x}{(a - \tanh(2x))^2} dx$$

input

```
integrate(exp(x)/(a-tanh(2*x))**2,x)
```

output

```
Integral(exp(x)/(a - tanh(2*x))**2, x)
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx = \text{Exception raised: ValueError}$$

input `integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(a-1>0)', see `assume?` for more details)Is

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 456 vs. 2(119) = 238.

Time = 0.11 (sec) , antiderivative size = 456, normalized size of antiderivative = 2.94

$$\begin{aligned} & \int \frac{e^x}{(a - \tanh(2x))^2} dx \\ &= \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}}(4a + 1) \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} + 2e^x\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{2(\sqrt{2}a^5 - \sqrt{2}a^4 - 2\sqrt{2}a^3 + 2\sqrt{2}a^2 + \sqrt{2}a - \sqrt{2})} \\ & \quad - \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}}(4a + 1) \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} - 2e^x\right)}{2\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}}}\right)}{2(\sqrt{2}a^5 - \sqrt{2}a^4 - 2\sqrt{2}a^3 + 2\sqrt{2}a^2 + \sqrt{2}a - \sqrt{2})} \\ & \quad - \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}}(4a + 1) \log\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{4(\sqrt{2}a^5 - \sqrt{2}a^4 - 2\sqrt{2}a^3 + 2\sqrt{2}a^2 + \sqrt{2}a - \sqrt{2})} \\ & \quad + \frac{(a^4 - 2a^3 + 2a - 1)^{\frac{1}{4}}(4a + 1) \log\left(-\sqrt{2}\left(\frac{a+1}{a-1}\right)^{\frac{1}{4}} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{4(\sqrt{2}a^5 - \sqrt{2}a^4 - 2\sqrt{2}a^3 + 2\sqrt{2}a^2 + \sqrt{2}a - \sqrt{2})} \\ & \quad + \frac{e^x}{a^2 - 2a + 1} + \frac{e^x}{(a^3 - a^2 - a + 1)(ae^{(4x)} + a - e^{(4x)} + 1)} \end{aligned}$$

input `integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="giac")`

output

$$\begin{aligned}
 & -1/2*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\arctan(1/2*\sqrt{2}*(\sqrt{2})* \\
 & (a + 1)/(a - 1))^{(1/4)} + 2*e^x/((a + 1)/(a - 1))^{(1/4)}/(\sqrt{2}*a^5 - \sqrt{2} \\
 & *a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) - 1/2*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\arctan(-1/2*\sqrt{2}*(\sqrt{2})* \\
 & (a + 1)/(a - 1))^{(1/4)} - 2*e^x/((a + 1)/(a - 1))^{(1/4)}/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) - 1/4*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\log(\sqrt{2}*(a + 1)/(a - 1))^{(1/4)}*e^x + \sqrt{2}*\sqrt{(a + 1)/(a - 1)} + e^{(2*x)})/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) + 1/4*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\log(-\sqrt{2}*(a + 1)/(a - 1))^{(1/4)}*e^x + \sqrt{2}*\sqrt{(a + 1)/(a - 1)} + e^{(2*x)})/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) + e^x/(a^2 - 2*a + 1) + e^x/((a^3 - a^2 - a + 1)*(a*e^{(4*x)} + a - e^{(4*x)} + 1))
 \end{aligned}$$

### Mupad [B] (verification not implemented)

Time = 24.35 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.81

$$\begin{aligned}
 \int \frac{e^x}{(a - \tanh(2x))^2} dx &= \frac{e^x}{(a - 1)^2} + \frac{\ln\left(\frac{4a+1}{(a-1)^{13/4}(-a-1)^{3/4}} + \frac{e^x(4a+1)}{a^4-2a^3+2a-1}\right)(4a+1)}{4(a-1)^{9/4}(-a-1)^{7/4}} \\
 &\quad - \frac{\ln\left(\frac{e^x(4a+1)}{a^4-2a^3+2a-1} - \frac{4a+1}{(a-1)^{13/4}(-a-1)^{3/4}}\right)(4a+1)}{4(a-1)^{9/4}(-a-1)^{7/4}} \\
 &\quad + \frac{e^x}{(a-1)^2(a+1)(a+e^{4x}(a-1)+1)} \\
 &\quad - \frac{\ln\left(\frac{e^x(4a+1)}{(a-1)^3(a+1)} - \frac{(4a+1)\operatorname{li}}{(a-1)^{13/4}(-a-1)^{3/4}}\right)(4a+1)\operatorname{li}}{4(a-1)^{9/4}(-a-1)^{7/4}} \\
 &\quad + \frac{\ln\left(\frac{e^x(4a+1)}{(a-1)^3(a+1)} + \frac{(4a+1)\operatorname{li}}{(a-1)^{13/4}(-a-1)^{3/4}}\right)(4a+1)\operatorname{li}}{4(a-1)^{9/4}(-a-1)^{7/4}}
 \end{aligned}$$

input `int(exp(x)/(a - tanh(2*x))^2,x)`

output

```
exp(x)/(a - 1)^2 - (log((exp(x)*(4*a + 1))/((a - 1)^3*(a + 1)) - ((4*a + 1)
)*1i)/((a - 1)^(13/4)*(- a - 1)^(3/4)))*(4*a + 1)*1i)/(4*(a - 1)^(9/4)*(-
a - 1)^(7/4)) + (log(((4*a + 1)*1i)/((a - 1)^(13/4)*(- a - 1)^(3/4)) + (ex
p(x)*(4*a + 1))/((a - 1)^3*(a + 1)))*(4*a + 1)*1i)/(4*(a - 1)^(9/4)*(- a -
1)^(7/4)) + (log((4*a + 1)/((a - 1)^(13/4)*(- a - 1)^(3/4)) + (exp(x)*(4*
a + 1))/(2*a - 2*a^3 + a^4 - 1))*(4*a + 1))/(4*(a - 1)^(9/4)*(- a - 1)^(7/
4)) - (log((exp(x)*(4*a + 1))/(2*a - 2*a^3 + a^4 - 1) - (4*a + 1)/((a - 1)
^(13/4)*(- a - 1)^(3/4)))*(4*a + 1))/(4*(a - 1)^(9/4)*(- a - 1)^(7/4)) + e
xp(x)/((a - 1)^2*(a + 1)*(a + exp(4*x)*(a - 1) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 5067, normalized size of antiderivative = 32.69

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx = \text{Too large to display}$$

input

```
int(exp(x)/(a-tanh(2*x))^2,x)
```



output

```
(16*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a
- 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*
sqrt(2)))*tanh(2*x)*a**4 - 4*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)
)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a +
1)**(1/4)*(a - 1)**(1/4)*sqrt(2)))*tanh(2*x)*a**3 + 6*e**(4*x)*(a + 1)**(1
/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2
*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2)))*tanh(2*x)*a**2
- 14*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*
(a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)
)*sqrt(2)))*tanh(2*x)*a - 4*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)
)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)
)**(1/4)*(a - 1)**(1/4)*sqrt(2)))*tanh(2*x) - 16*e**(4*x)*(a + 1)**(1/4)*(
a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*e**x
*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2)))*a**5 + 4*e**(4*x)*(
a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(a - 1)**(1/4)*s
qrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2)))*a**4
- 6*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan(((a + 1)**(1/4)*(
a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(1/4)*(a - 1)**(1/4)
)*sqrt(2)))*a**3 + 14*e**(4*x)*(a + 1)**(1/4)*(a - 1)**(3/4)*sqrt(2)*atan((
(a + 1)**(1/4)*(a - 1)**(1/4)*sqrt(2) - 2*e**x*sqrt(a - 1))/((a + 1)**(...
```

### 3.239 $\int e^{c(a+bx)} \tanh^3(d+ex) dx$

Optimal result	1761
Mathematica [A] (verified)	1762
Rubi [A] (verified)	1762
Maple [F]	1763
Fricas [F]	1764
Sympy [F]	1764
Maxima [F]	1764
Giac [F]	1765
Mupad [F(-1)]	1765
Reduce [F]	1766

#### Optimal result

Integrand size = 18, antiderivative size = 167

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(3, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc}$$

output

```
exp(c*(b*x+a))/b/c-6*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c+12*exp(c*(b*x+a))*hypergeom([2, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c-8*exp(c*(b*x+a))*hypergeom([3, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c
```

**Mathematica [A] (verified)**

Time = 1.66 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.23

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx$$

$$= \frac{1}{2} e^{ac} \left( \frac{2(b^2 c^2 + 2e^2) e^{2d} \left( \frac{e^{(bc+2e)x} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc+2e} - \frac{e^{bcx} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc} \right)}{e^2 (1 + e^{2d})} + \frac{e^{bcx} \operatorname{sech}^2(d+ex)}{e} - \frac{bce^{bcx} \operatorname{sech}(d) \operatorname{sech}(d+ex) \sinh(ex)}{e^2} + \frac{2e^{bcx} \tanh(d)}{bc} \right)$$

input

```
Integrate[E^(c*(a + b*x))*Tanh[d + e*x]^3,x]
```

output

```
(E^(a*c)*((2*(b^2*c^2 + 2*e^2)*E^(2*d))*((E^((b*c + 2*e)*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c + 2*e) - (E^(b*c*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c))/(e^2*(1 + E^(2*d))) + (E^(b*c*x))*Sech[d + e*x]^2/e - (b*c*E^(b*c*x))*Sech[d]*Sech[d + e*x]*Sinh[e*x])/e^2 + (2*E^(b*c*x))*Tanh[d])/(b*c))/2
```

**Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6007, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx$$

↓ 6007

$$\int \left( -\frac{6e^{c(a+bx)}}{e^{2(d+ex)} + 1} + \frac{12e^{c(a+bx)}}{(e^{2(d+ex)} + 1)^2} - \frac{8e^{c(a+bx)}}{(e^{2(d+ex)} + 1)^3} + e^{c(a+bx)} \right) dx$$

↓ 2009

$$-\frac{6e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc} +$$

$$\frac{12e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc} -$$

$$\frac{8e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(3, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

input `Int[E^(c*(a + b*x))*Tanh[d + e*x]^3,x]`

output `E^(c*(a + b*x))/(b*c) - (6*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c) + (12*E^(c*(a + b*x))*Hypergeometric2F1[2, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c) - (8*E^(c*(a + b*x))*Hypergeometric2F1[3, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c)`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6007 `Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Tanh[(d_.) + (e_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((-1 + E^(2*(d + e*x)))^n/(1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

### Maple [F]

$$\int e^{c(bx+a)} \tanh(ex+d)^3 dx$$

input `int(exp(c*(b*x+a))*tanh(e*x+d)^3,x)`

output `int(exp(c*(b*x+a))*tanh(e*x+d)^3,x)`

### Fricas [F]

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = \int e^{((bx+a)c)} \tanh^3(ex+d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="fricas")`

output `integral(e^(b*c*x + a*c)*tanh(e*x + d)^3, x)`

### Sympy [F]

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = e^{ac} \int e^{bcx} \tanh^3(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)**3,x)`

output `exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x)**3, x)`

### Maxima [F]

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = \int e^{((bx+a)c)} \tanh^3(ex+d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="maxima")`

output

```

48*(b^2*c^2*e^e^(a*c) + 2*e^3*e^(a*c))*integrate(e^(b*c*x)/(b^3*c^3 - 12*b
^2*c^2*e + 44*b*c*e^2 - 48*e^3 + (b^3*c^3*e^(8*d) - 12*b^2*c^2*e*e^(8*d) +
44*b*c*e^2*e^(8*d) - 48*e^3*e^(8*d))*e^(8*e*x) + 4*(b^3*c^3*e^(6*d) - 12*
b^2*c^2*e*e^(6*d) + 44*b*c*e^2*e^(6*d) - 48*e^3*e^(6*d))*e^(6*e*x) + 6*(b^
3*c^3*e^(4*d) - 12*b^2*c^2*e*e^(4*d) + 44*b*c*e^2*e^(4*d) - 48*e^3*e^(4*d)
)*e^(4*e*x) + 4*(b^3*c^3*e^(2*d) - 12*b^2*c^2*e*e^(2*d) + 44*b*c*e^2*e^(2*
d) - 48*e^3*e^(2*d))*e^(2*e*x)), x) - (b^3*c^3*e^(a*c) + 36*b^2*c^2*e*e^(a
*c) + 44*b*c*e^2*e^(a*c) + 48*e^3*e^(a*c) - (b^3*c^3*e^(a*c + 6*d) - 12*b^
2*c^2*e*e^(a*c + 6*d) + 44*b*c*e^2*e^(a*c + 6*d) - 48*e^3*e^(a*c + 6*d))*e
^(6*e*x) + 3*(b^3*c^3*e^(a*c + 4*d) - 8*b^2*c^2*e*e^(a*c + 4*d) + 4*b*c*e^
2*e^(a*c + 4*d) + 48*e^3*e^(a*c + 4*d))*e^(4*e*x) - 3*(b^3*c^3*e^(a*c + 2*
d) - 28*b*c*e^2*e^(a*c + 2*d) - 48*e^3*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)
/(b^4*c^4 - 12*b^3*c^3*e + 44*b^2*c^2*e^2 - 48*b*c*e^3 + (b^4*c^4*e^(6*d)
- 12*b^3*c^3*e*e^(6*d) + 44*b^2*c^2*e^2*e^(6*d) - 48*b*c*e^3*e^(6*d))*e^(6
*e*x) + 3*(b^4*c^4*e^(4*d) - 12*b^3*c^3*e*e^(4*d) + 44*b^2*c^2*e^2*e^(4*d)
- 48*b*c*e^3*e^(4*d))*e^(4*e*x) + 3*(b^4*c^4*e^(2*d) - 12*b^3*c^3*e*e^(2*
d) + 44*b^2*c^2*e^2*e^(2*d) - 48*b*c*e^3*e^(2*d))*e^(2*e*x))

```

**Giac [F]**

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = \int e^{(bx+a)c} \tanh(ex+d)^3 dx$$

input

```
integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="giac")
```

output

```
integrate(e^((b*x + a)*c)*tanh(e*x + d)^3, x)
```

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^3(d+ex) dx = \int e^{c(a+bx)} \tanh(d+ex)^3 dx$$

input

```
int(exp(c*(a + b*x))*tanh(d + e*x)^3,x)
```

output `int(exp(c*(a + b*x))*tanh(d + e*x)^3, x)`

## Reduce [F]

$$\int e^{c(a+bx)} \tanh^3(d + ex) dx = \text{too large to display}$$

input `int(exp(c*(b*x+a))*tanh(e*x+d)^3,x)`

output

```
(e**(a*c)*(e**(b*c*x + 4*d + 4*e*x)*b**2*c**2 - 6*e**(b*c*x + 4*d + 4*e*x)
*b*c*e + 8*e**(b*c*x + 4*d + 4*e*x)*e**2 - 4*e**(b*c*x + 2*d + 2*e*x)*b**2
*c**2 + 12*e**(b*c*x + 2*d + 2*e*x)*b*c*e + 16*e**(b*c*x + 2*d + 2*e*x)*e
**2 + 7*e**(b*c*x)*b**2*c**2 + 6*e**(b*c*x)*b*c*e + 8*e**(b*c*x)*e**2 - 8*e
**(4*d + 4*e*x)*int(e**(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6
*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e**2 + 3*e**(4*d + 4*e*x)*b**2*c**2 - 18*
e**(4*d + 4*e*x)*b*c*e + 24*e**(4*d + 4*e*x)*e**2 + 3*e**(2*d + 2*e*x)*b**
2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e + 24*e**(2*d + 2*e*x)*e**2 + b**2*c**2
- 6*b*c*e + 8*e**2),x)*b**5*c**5 + 48*e**(4*d + 4*e*x)*int(e**(b*c*x)/(e**
(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e
**2 + 3*e**(4*d + 4*e*x)*b**2*c**2 - 18*e**(4*d + 4*e*x)*b*c*e + 24*e**(4*d
+ 4*e*x)*e**2 + 3*e**(2*d + 2*e*x)*b**2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e
+ 24*e**(2*d + 2*e*x)*e**2 + b**2*c**2 - 6*b*c*e + 8*e**2),x)*b**4*c**4*e
- 80*e**(4*d + 4*e*x)*int(e**(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6
*d + 6*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e**2 + 3*e**(4*d + 4*e*x)*b**2*c**2
- 18*e**(4*d + 4*e*x)*b*c*e + 24*e**(4*d + 4*e*x)*e**2 + 3*e**(2*d + 2*e*
x)*b**2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e + 24*e**(2*d + 2*e*x)*e**2 + b**2
*c**2 - 6*b*c*e + 8*e**2),x)*b**3*c**3*e**2 + 96*e**(4*d + 4*e*x)*int(e**
(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6*e*x)*b*c*e + 8*e**(6*d
+ 6*e*x)*e**2 + 3*e**(4*d + 4*e*x)*b**2*c**2 - 18*e**(4*d + 4*e*x)*b*c...
```

### 3.240 $\int e^{c(a+bx)} \tanh^2(d+ex) dx$

Optimal result	1767
Mathematica [A] (verified)	1767
Rubi [A] (verified)	1768
Maple [F]	1769
Fricas [F]	1769
Sympy [F]	1770
Maxima [F]	1770
Giac [F]	1771
Mupad [F(-1)]	1771
Reduce [F]	1771

#### Optimal result

Integrand size = 18, antiderivative size = 117

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc}$$

output

```
exp(c*(b*x+a))/b/c-4*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c+4*exp(c*(b*x+a))*hypergeom([2, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c
```

#### Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.44

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \frac{e^{c(a+bx)} \left( 2b^2c^2e^{2(d+ex)} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, -e^{2(d+ex)}\right) - (bc + 2e) \left( 2bce^{2d} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right) - (bc + 2e) \right) \right)}{bce(bc + 2e)(1 + e^{2d})}$$



input `Integrate[E^(c*(a + b*x))*Tanh[d + e*x]^2,x]`

output `(E^(c*(a + b*x))*(2*b^2*c^2*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))] - (b*c + 2*e)*(2*b*c*E^(2*d)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))] - (1 + E^(2*d))*(e - b*c*Sech[d]*Sech[d + e*x]*Sinh[e*x]))) / (b*c*e*(b*c + 2*e)*(1 + E^(2*d)))`

### Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6007, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx$$

$$\downarrow 6007$$

$$\int \left( -\frac{4e^{c(a+bx)}}{e^{2(d+ex)} + 1} + \frac{4e^{c(a+bx)}}{(e^{2(d+ex)} + 1)^2} + e^{c(a+bx)} \right) dx$$

$$\downarrow 2009$$

$$-\frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

input `Int[E^(c*(a + b*x))*Tanh[d + e*x]^2,x]`

output `E^(c*(a + b*x))/(b*c) - (4*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c) + (4*E^(c*(a + b*x))*Hypergeometric2F1[2, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c)`

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6007 `Int[(F_)^((c_)*((a_) + (b_)*(x_)))*Tanh[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((-1 + E^(2*(d + e*x)))^n/(1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

**Maple [F]**

$$\int e^{c(bx+a)} \tanh(ex+d)^2 dx$$

input `int(exp(c*(b*x+a))*tanh(e*x+d)^2,x)`

output `int(exp(c*(b*x+a))*tanh(e*x+d)^2,x)`

**Fricas [F]**

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \int e^{((bx+a)c)} \tanh(ex+d)^2 dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="fricas")`

output `integral(e^(b*c*x + a*c)*tanh(e*x + d)^2, x)`

**Sympy [F]**

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = e^{ac} \int e^{bcx} \tanh^2(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)**2,x)`

output `exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x)**2, x)`

**Maxima [F]**

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \int e^{(bx+a)c} \tanh^2(ex+d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="maxima")`

output `-16*b*c*e*integrate(e^(b*c*x + a*c)/(b^2*c^2 - 6*b*c*e + 8*e^2 + (b^2*c^2*e^(6*d) - 6*b*c*e*e^(6*d) + 8*e^2*e^(6*d))*e^(6*e*x) + 3*(b^2*c^2*e^(4*d) - 6*b*c*e*e^(4*d) + 8*e^2*e^(4*d))*e^(4*e*x) + 3*(b^2*c^2*e^(2*d) - 6*b*c*e*e^(2*d) + 8*e^2*e^(2*d))*e^(2*e*x)), x) + (b^2*c^2*e^(a*c) + 10*b*c*e*e^(a*c) + 8*e^2*e^(a*c) + (b^2*c^2*e^(a*c + 4*d) - 6*b*c*e*e^(a*c + 4*d) + 8*e^2*e^(a*c + 4*d))*e^(4*e*x) - 2*(b^2*c^2*e^(a*c + 2*d) - 2*b*c*e*e^(a*c + 2*d) - 8*e^2*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)/(b^3*c^3 - 6*b^2*c^2*e + 8*b*c*e^2 + (b^3*c^3*e^(4*d) - 6*b^2*c^2*e*e^(4*d) + 8*b*c*e^2*e^(4*d))*e^(4*e*x) + 2*(b^3*c^3*e^(2*d) - 6*b^2*c^2*e*e^(2*d) + 8*b*c*e^2*e^(2*d))*e^(2*e*x))`

**Giac [F]**

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \int e^{((bx+a)c)} \tanh(ex+d)^2 dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="giac")`

output `integrate(e^((b*x + a)*c))*tanh(e*x + d)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx = \int e^{c(a+bx)} \tanh(d+ex)^2 dx$$

input `int(exp(c*(a + b*x))*tanh(d + e*x)^2,x)`

output `int(exp(c*(a + b*x))*tanh(d + e*x)^2, x)`

**Reduce [F]**

$$\int e^{c(a+bx)} \tanh^2(d+ex) dx$$

$$= \frac{e^{ac} \left( e^{bcx+2ex+2d} bc - 2e^{bcx+2ex+2d} e - 3e^{bcx} bc - 2e^{bcx} e + 4e^{2ex+2d} \left( \int \frac{e^{bcx}}{e^{4ex+4d} bc - 2e^{4ex+4d} e + 2e^{2ex+2d} bc - 4e^{2ex+2d} e + bc} dx \right) \right)}{e^{4ex+4d} bc - 2e^{4ex+4d} e + 2e^{2ex+2d} bc - 4e^{2ex+2d} e + bc}$$

input `int(exp(c*(b*x+a))*tanh(e*x+d)^2,x)`

output

```
(e**(a*c)*(e**(b*c*x + 2*d + 2*e*x)*b*c - 2*e**(b*c*x + 2*d + 2*e*x)*e - 3
*e**(b*c*x)*b*c - 2*e**(b*c*x)*e + 4*e**(2*d + 2*e*x)*int(e**(b*c*x)/(e**(
4*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e + 2*e**(2*d + 2*e*x)*b*c - 4*e**(2
*d + 2*e*x)*e + b*c - 2*e),x)*b**3*c**3 - 8*e**(2*d + 2*e*x)*int(e**(b*c*x
)/(e**(4*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e + 2*e**(2*d + 2*e*x)*b*c -
4*e**(2*d + 2*e*x)*e + b*c - 2*e),x)*b**2*c**2*e + 4*int(e**(b*c*x)/(e**(4
*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e + 2*e**(2*d + 2*e*x)*b*c - 4*e**(2*
d + 2*e*x)*e + b*c - 2*e),x)*b**3*c**3 - 8*int(e**(b*c*x)/(e**(4*d + 4*e*x
)*b*c - 2*e**(4*d + 4*e*x)*e + 2*e**(2*d + 2*e*x)*b*c - 4*e**(2*d + 2*e*x)
*e + b*c - 2*e),x)*b**2*c**2*e))/(b*c*(e**(2*d + 2*e*x)*b*c - 2*e**(2*d +
2*e*x)*e + b*c - 2*e))
```

### 3.241 $\int e^{c(a+bx)} \tanh(d + ex) dx$

Optimal result	1773
Mathematica [B] (verified)	1773
Rubi [A] (verified)	1774
Maple [F]	1775
Fricas [F]	1775
Sympy [F]	1776
Maxima [F]	1776
Giac [F]	1776
Mupad [F(-1)]	1777
Reduce [F]	1777

#### Optimal result

Integrand size = 16, antiderivative size = 67

$$\int e^{c(a+bx)} \tanh(d + ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, -e^{2(d+ex)}\right)}{bc}$$

output

```
exp(c*(b*x+a))/b/c-2*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], -exp(2*e*x+2*d))/b/c
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 141 vs. 2(67) = 134.

Time = 0.18 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.10

$$\int e^{c(a+bx)} \tanh(d + ex) dx = \frac{e^{c(a+bx)} (2bce^{2(d+ex)} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, -e^{2(d+ex)}\right) - (bc + 2e) (1 - e^{2d} + 2e^{2d} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, -e^{2(d+ex)}\right)))}{bc(bc + 2e) (1 + e^{2d})}$$

input

```
Integrate[E^(c*(a + b*x))*Tanh[d + e*x], x]
```

output

```
(E^(c*(a + b*x))*(2*b*c*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*
e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))] - (b*c + 2*e)*(1 - E^(2*d) + 2*E^(2
*d))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])
)/(b*c*(b*c + 2*e)*(1 + E^(2*d)))
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {6007, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \tanh(d+ex) dx$$

$$\downarrow \text{6007}$$

$$\int \left( e^{c(a+bx)} - \frac{2e^{c(a+bx)}}{e^{2(d+ex)} + 1} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} \text{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, -e^{2(d+ex)}\right)}{bc}$$

input

```
Int[E^(c*(a + b*x))*Tanh[d + e*x],x]
```

output

```
E^(c*(a + b*x))/(b*c) - (2*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e
), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c)
```

**Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6007 `Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Tanh[(d_.) + (e_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((-1 + E^(2*(d + e*x)))^n/(1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

**Maple [F]**

$$\int e^{c(bx+a)} \tanh(ex + d) dx$$

input `int(exp(c*(b*x+a))*tanh(e*x+d),x)`

output `int(exp(c*(b*x+a))*tanh(e*x+d),x)`

**Fricas [F]**

$$\int e^{c(a+bx)} \tanh(d + ex) dx = \int e^{((bx+a)c)} \tanh(ex + d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d),x, algorithm="fricas")`

output `integral(e^(b*c*x + a*c)*tanh(e*x + d), x)`



**Sympy [F]**

$$\int e^{c(a+bx)} \tanh(d+ex) dx = e^{ac} \int e^{bcx} \tanh(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d), x)`

output `exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x), x)`

**Maxima [F]**

$$\int e^{c(a+bx)} \tanh(d+ex) dx = \int e^{(bx+a)c} \tanh(ex+d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d), x, algorithm="maxima")`

output `4*e*integrate(e^(b*c*x + a*c)/(b*c + (b*c*e^(4*d) - 2*e*e^(4*d))*e^(4*e*x) + 2*(b*c*e^(2*d) - 2*e*e^(2*d))*e^(2*e*x) - 2*e), x) - (b*c*e^(a*c) + 2*e*e^(a*c) - (b*c*e^(a*c + 2*d) - 2*e*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)/(b^2*c^2 - 2*b*c*e + (b^2*c^2*e^(2*d) - 2*b*c*e*e^(2*d))*e^(2*e*x))`

**Giac [F]**

$$\int e^{c(a+bx)} \tanh(d+ex) dx = \int e^{(bx+a)c} \tanh(ex+d) dx$$

input `integrate(exp(c*(b*x+a))*tanh(e*x+d), x, algorithm="giac")`

output `integrate(e^((b*x + a)*c)*tanh(e*x + d), x)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh(d+ex) dx = \int e^{c(a+bx)} \tanh(d+ex) dx$$

input `int(exp(c*(a + b*x))*tanh(d + e*x), x)`

output `int(exp(c*(a + b*x))*tanh(d + e*x), x)`

**Reduce [F]**

$$\int e^{c(a+bx)} \tanh(d+ex) dx = \frac{e^{ac} \left( e^{bcx} - 2 \left( \int \frac{e^{bcx}}{e^{2ex+2d}+1} dx \right) bc \right)}{bc}$$

input `int(exp(c*(b*x+a))*tanh(e*x+d), x)`

output `(e**(a*c)*(e**(b*c*x) - 2*int(e**(b*c*x)/(e**(2*d + 2*e*x) + 1), x)*b*c))/(b*c)`

### 3.242 $\int e^{c(a+bx)} \coth(d+ex) dx$

Optimal result	1778
Mathematica [B] (verified)	1778
Rubi [A] (verified)	1779
Maple [F]	1780
Fricas [F]	1780
Sympy [F]	1781
Maxima [F]	1781
Giac [F]	1781
Mupad [F(-1)]	1782
Reduce [F]	1782

#### Optimal result

Integrand size = 16, antiderivative size = 65

$$\int e^{c(a+bx)} \coth(d+ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc}$$

output `exp(c*(b*x+a))/b/c-2*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 134 vs. 2(65) = 130.

Time = 0.78 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.06

$$\int e^{c(a+bx)} \coth(d+ex) dx = \frac{e^{c(a+bx)} (2bce^{2(d+ex)} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, e^{2(d+ex)}\right) + (bc+2e)(1+e^{2d} - 2e^{2d} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, e^{2(d+ex)}\right)))}{bc(bc+2e)(-1+e^{2d})}$$

input `Integrate[E^(c*(a + b*x))*Coth[d + e*x], x]`

output

$$\frac{(E^{c(a+bx)})*(2*b*c*E^{2*(d+e*x)}*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^{2*(d+e*x)}]) + (b*c + 2*e)*(1 + E^{2*d}) - 2*E^{2*d}*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d+e*x)}])}{(b*c*(b*c + 2*e)*(-1 + E^{2*d}))}$$
**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {6008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \coth(d+ex) dx$$

$$\downarrow \text{6008}$$

$$\int \left( \frac{2e^{c(a+bx)}}{e^{2(d+ex)} - 1} + e^{c(a+bx)} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} \text{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc}$$

input

$$\text{Int}[E^{c*(a + b*x)}*Coth[d + e*x], x]$$

output

$$\frac{E^{c*(a + b*x)}}{(b*c)} - \frac{(2*E^{c*(a + b*x)}*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])}{(b*c)}$$

## Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6008 `Int[Coth[(d_.) + (e_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((1 + E^(2*(d + e*x)))^n/(-1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

## Maple [F]

$$\int e^{c(bx+a)} \coth(ex + d) dx$$

input `int(exp(c*(b*x+a))*coth(e*x+d), x)`

output `int(exp(c*(b*x+a))*coth(e*x+d), x)`

## Fricas [F]

$$\int e^{c(a+bx)} \coth(d + ex) dx = \int \coth(ex + d) e^{((bx+a)c)} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d), x, algorithm="fricas")`

output `integral(coth(e*x + d)*e^(b*c*x + a*c), x)`

**Sympy [F]**

$$\int e^{c(a+bx)} \coth(d+ex) dx = e^{ac} \int e^{bcx} \coth(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d), x)`

output `exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x), x)`

**Maxima [F]**

$$\int e^{c(a+bx)} \coth(d+ex) dx = \int \coth(ex+d) e^{(bx+a)c} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d), x, algorithm="maxima")`

output `4*e*integrate(e^(b*c*x + a*c)/(b*c + (b*c*e^(4*d) - 2*e*e^(4*d))*e^(4*e*x) - 2*(b*c*e^(2*d) - 2*e*e^(2*d))*e^(2*e*x) - 2*e), x) - (b*c*e^(a*c) + 2*e*e^(a*c) + (b*c*e^(a*c + 2*d) - 2*e*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)/(b^2*c^2 - 2*b*c*e - (b^2*c^2*e^(2*d) - 2*b*c*e*e^(2*d))*e^(2*e*x))`

**Giac [F]**

$$\int e^{c(a+bx)} \coth(d+ex) dx = \int \coth(ex+d) e^{(bx+a)c} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d), x, algorithm="giac")`

output `integrate(coth(e*x + d)*e^((b*x + a)*c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \coth(d+ex) dx = \int \coth(d+ex) e^{c(a+bx)} dx$$

input `int(coth(d + e*x)*exp(c*(a + b*x)), x)`output `int(coth(d + e*x)*exp(c*(a + b*x)), x)`**Reduce [F]**

$$\int e^{c(a+bx)} \coth(d+ex) dx = \frac{e^{ac} \left( e^{bcx} + 2 \left( \int \frac{e^{bcx}}{e^{2ex+2d}-1} dx \right) bc \right)}{bc}$$

input `int(exp(c*(b*x+a))*coth(e*x+d), x)`output `(e**(a*c)*(e**(b*c*x) + 2*int(e**(b*c*x)/(e**(2*d + 2*e*x) - 1), x)*b*c))/(b*c)`

### 3.243 $\int e^{c(a+bx)} \coth^2(d+ex) dx$

Optimal result	1783
Mathematica [A] (verified)	1783
Rubi [A] (verified)	1784
Maple [F]	1785
Fricas [F]	1785
Sympy [F]	1786
Maxima [F]	1786
Giac [F]	1787
Mupad [F(-1)]	1787
Reduce [F]	1787

#### Optimal result

Integrand size = 18, antiderivative size = 113

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc}$$

output

```
exp(c*(b*x+a))/b/c-4*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c+4*exp(c*(b*x+a))*hypergeom([2, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c
```

#### Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.45

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \frac{e^{c(a+bx)} \left( 2b^2c^2e^{2(d+ex)} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, e^{2(d+ex)}\right) - (bc+2e) \left( 2bce^{2d} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right) - bce^{2d} \right) \right)}{bce(bc+2e)(-1+e^{2d})}$$



input `Integrate[E^(c*(a + b*x))*Coth[d + e*x]^2,x]`

output `(E^(c*(a + b*x))*(2*b^2*c^2*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^(2*(d + e*x))] - (b*c + 2*e)*(2*b*c*E^(2*d)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))] - (-1 + E^(2*d))*(e + b*c*Csch[d]*Csch[d + e*x]*Sinh[e*x]))) / (b*c*e*(b*c + 2*e)*(-1 + E^(2*d)))`

### Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \coth^2(d+ex) dx$$

$$\downarrow 6008$$

$$\int \left( \frac{4e^{c(a+bx)}}{e^{2(d+ex)} - 1} + \frac{4e^{c(a+bx)}}{(e^{2(d+ex)} - 1)^2} + e^{c(a+bx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

input `Int[E^(c*(a + b*x))*Coth[d + e*x]^2,x]`

output `E^(c*(a + b*x))/(b*c) - (4*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c) + (4*E^(c*(a + b*x))*Hypergeometric2F1[2, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c)`

## Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6008 `Int[Coth[(d_.) + (e_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((1 + E^(2*(d + e*x)))^n/(-1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

## Maple [F]

$$\int e^{c(bx+a)} \coth(ex+d)^2 dx$$

input `int(exp(c*(b*x+a))*coth(e*x+d)^2,x)`

output `int(exp(c*(b*x+a))*coth(e*x+d)^2,x)`

## Fricas [F]

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \int \coth(ex+d)^2 e^{((bx+a)c)} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="fricas")`

output `integral(coth(e*x + d)^2*e^(b*c*x + a*c), x)`

**Sympy [F]**

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = e^{ac} \int e^{bcx} \coth^2(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)**2,x)`

output `exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x)**2, x)`

**Maxima [F]**

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \int \coth(ex+d)^2 e^{(bx+a)c} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="maxima")`

output `16*b*c*e*integrate(-e^(b*c*x + a*c)/(b^2*c^2 - 6*b*c*e + 8*e^2 - (b^2*c^2*e^(6*d) - 6*b*c*e*e^(6*d) + 8*e^2*e^(6*d))*e^(6*e*x) + 3*(b^2*c^2*e^(4*d) - 6*b*c*e*e^(4*d) + 8*e^2*e^(4*d))*e^(4*e*x) - 3*(b^2*c^2*e^(2*d) - 6*b*c*e*e^(2*d) + 8*e^2*e^(2*d))*e^(2*e*x)), x) + (b^2*c^2*e^(a*c) + 10*b*c*e*e^(a*c) + 8*e^2*e^(a*c) + (b^2*c^2*e^(a*c + 4*d) - 6*b*c*e*e^(a*c + 4*d) + 8*e^2*e^(a*c + 4*d))*e^(4*e*x) + 2*(b^2*c^2*e^(a*c + 2*d) - 2*b*c*e*e^(a*c + 2*d) - 8*e^2*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)/(b^3*c^3 - 6*b^2*c^2*e + 8*b*c*e^2 + (b^3*c^3*e^(4*d) - 6*b^2*c^2*e*e^(4*d) + 8*b*c*e^2*e^(4*d))*e^(4*e*x) - 2*(b^3*c^3*e^(2*d) - 6*b^2*c^2*e*e^(2*d) + 8*b*c*e^2*e^(2*d))*e^(2*e*x))`

**Giac [F]**

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \int \coth(ex+d)^2 e^{((bx+a)c)} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="giac")`

output `integrate(coth(e*x + d)^2*e^((b*x + a)*c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \coth^2(d+ex) dx = \int \coth(d+ex)^2 e^{c(a+bx)} dx$$

input `int(coth(d + e*x)^2*exp(c*(a + b*x)),x)`

output `int(coth(d + e*x)^2*exp(c*(a + b*x)), x)`

**Reduce [F]**

$$\int e^{c(a+bx)} \coth^2(d+ex) dx$$

$$= \frac{e^{ac} \left( e^{bcx+2ex+2d} bc - 2e^{bcx+2ex+2d} e + 3e^{bcx} bc + 2e^{bcx} e + 4e^{2ex+2d} \left( \int \frac{e^{bcx}}{e^{4ex+4d} bc - 2e^{4ex+4d} e - 2e^{2ex+2d} bc + 4e^{2ex+2d} e + bc} dx \right) \right)}{e^{4ex+4d} bc - 2e^{4ex+4d} e - 2e^{2ex+2d} bc + 4e^{2ex+2d} e + bc}$$

input `int(exp(c*(b*x+a))*coth(e*x+d)^2,x)`

output

```
(e**(a*c)*(e**(b*c*x + 2*d + 2*e*x)*b*c - 2*e**(b*c*x + 2*d + 2*e*x)*e + 3
*e**(b*c*x)*b*c + 2*e**(b*c*x)*e + 4*e**(2*d + 2*e*x)*int(e**(b*c*x)/(e**(
4*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e - 2*e**(2*d + 2*e*x)*b*c + 4*e**(2
*d + 2*e*x)*e + b*c - 2*e),x)*b**3*c**3 - 8*e**(2*d + 2*e*x)*int(e**(b*c*x
)/(e**(4*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e - 2*e**(2*d + 2*e*x)*b*c +
4*e**(2*d + 2*e*x)*e + b*c - 2*e),x)*b**2*c**2*e - 4*int(e**(b*c*x)/(e**(4
*d + 4*e*x)*b*c - 2*e**(4*d + 4*e*x)*e - 2*e**(2*d + 2*e*x)*b*c + 4*e**(2*
d + 2*e*x)*e + b*c - 2*e),x)*b**3*c**3 + 8*int(e**(b*c*x)/(e**(4*d + 4*e*x
)*b*c - 2*e**(4*d + 4*e*x)*e - 2*e**(2*d + 2*e*x)*b*c + 4*e**(2*d + 2*e*x)
*e + b*c - 2*e),x)*b**2*c**2*e))/(b*c*(e**(2*d + 2*e*x)*b*c - 2*e**(2*d +
2*e*x)*e - b*c + 2*e))
```

### 3.244 $\int e^{c(a+bx)} \coth^3(d+ex) dx$

Optimal result	1789
Mathematica [A] (verified)	1790
Rubi [A] (verified)	1790
Maple [F]	1791
Fricas [F]	1792
Sympy [F]	1792
Maxima [F]	1792
Giac [F]	1793
Mupad [F(-1)]	1793
Reduce [F]	1794

#### Optimal result

Integrand size = 18, antiderivative size = 161

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(3, \frac{bc}{2e}, 1 + \frac{bc}{2e}, e^{2(d+ex)}\right)}{bc}$$

output

```
exp(c*(b*x+a))/b/c-6*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c+12*exp(c*(b*x+a))*hypergeom([2, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c-8*exp(c*(b*x+a))*hypergeom([3, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c
```

**Mathematica [A] (verified)**

Time = 1.12 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.30

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \frac{e^{c(a+bx)} \coth(d)}{bc} - \frac{e^{c(a+bx)} \operatorname{csch}^2(d+ex)}{2e} + \frac{(b^2c^2 + 2e^2) e^{ac+2d+bcx} (bce^{2ex} \operatorname{Hypergeometric2F1}(1, 1 + \frac{bc}{2e}, 2 + \frac{bc}{2e}, e^{2(d+ex)}) - (bc + 2e) \operatorname{Hypergeometric2F1}(1, \frac{bc}{2e}, 1 + \frac{bc}{2e}, E^{2(d+ex)}))}{bce^2(bc + 2e)(-1 + e^{2d})} + \frac{bce^{c(a+bx)} \operatorname{csch}(d) \operatorname{csch}(d+ex) \sinh(ex)}{2e^2}$$

input `Integrate[E^(c*(a + b*x))*Coth[d + e*x]^3,x]`

output

```
(E^(c*(a + b*x))*Coth[d])/(b*c) - (E^(c*(a + b*x))*Csch[d + e*x]^2)/(2*e) + ((b^2*c^2 + 2*e^2)*E^(a*c + 2*d + b*c*x)*(b*c*E^(2*e*x)*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^(2*(d + e*x))] - (b*c + 2*e)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c*e^2*(b*c + 2*e)*(-1 + E^(2*d))) + (b*c*E^(c*(a + b*x))*Csch[d]*Csch[d + e*x]*Sinh[e*x])/(2*e^2)
```

**Rubi [A] (verified)**

Time = 0.69 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6008, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \coth^3(d+ex) dx$$

$$\downarrow 6008$$

$$\int \left( \frac{6e^{c(a+bx)}}{e^{2(d+ex)} - 1} + \frac{12e^{c(a+bx)}}{(e^{2(d+ex)} - 1)^2} + \frac{8e^{c(a+bx)}}{(e^{2(d+ex)} - 1)^3} + e^{c(a+bx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{6e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} \operatorname{Hypergeometric2F1}\left(3, \frac{bc}{2e}, \frac{bc}{2e} + 1, e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

input `Int[E^(c*(a + b*x))*Coth[d + e*x]^3,x]`

output `E^(c*(a + b*x))/(b*c) - (6*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c) + (12*E^(c*(a + b*x))*Hypergeometric2F1[2, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c) - (8*E^(c*(a + b*x))*Hypergeometric2F1[3, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/(b*c)`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6008 `Int[Coth[(d_.) + (e_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((1 + E^(2*(d + e*x)))^n/(-1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]`

### Maple [F]

$$\int e^{c(bx+a)} \coth(ex+d)^3 dx$$

input `int(exp(c*(b*x+a))*coth(e*x+d)^3,x)`

output `int(exp(c*(b*x+a))*coth(e*x+d)^3,x)`



**Fricas [F]**

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \int \coth(ex+d)^3 e^{((bx+a)c)} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="fricas")`

output `integral(coth(e*x + d)^3*e^(b*c*x + a*c), x)`

**Sympy [F]**

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = e^{ac} \int e^{bcx} \coth^3(d+ex) dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)**3,x)`

output `exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x)**3, x)`

**Maxima [F]**

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \int \coth(ex+d)^3 e^{((bx+a)c)} dx$$

input `integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="maxima")`

output

```

48*(b^2*c^2*e*e^(a*c) + 2*e^3*e^(a*c))*integrate(e^(b*c*x)/(b^3*c^3 - 12*b
^2*c^2*e + 44*b*c*e^2 - 48*e^3 + (b^3*c^3*e^(8*d) - 12*b^2*c^2*e*e^(8*d) +
44*b*c*e^2*e^(8*d) - 48*e^3*e^(8*d))*e^(8*e*x) - 4*(b^3*c^3*e^(6*d) - 12*
b^2*c^2*e*e^(6*d) + 44*b*c*e^2*e^(6*d) - 48*e^3*e^(6*d))*e^(6*e*x) + 6*(b^
3*c^3*e^(4*d) - 12*b^2*c^2*e*e^(4*d) + 44*b*c*e^2*e^(4*d) - 48*e^3*e^(4*d)
)*e^(4*e*x) - 4*(b^3*c^3*e^(2*d) - 12*b^2*c^2*e*e^(2*d) + 44*b*c*e^2*e^(2*
d) - 48*e^3*e^(2*d))*e^(2*e*x)), x) - (b^3*c^3*e^(a*c) + 36*b^2*c^2*e*e^(a
*c) + 44*b*c*e^2*e^(a*c) + 48*e^3*e^(a*c) + (b^3*c^3*e^(a*c + 6*d) - 12*b^
2*c^2*e*e^(a*c + 6*d) + 44*b*c*e^2*e^(a*c + 6*d) - 48*e^3*e^(a*c + 6*d))*e
^(6*e*x) + 3*(b^3*c^3*e^(a*c + 4*d) - 8*b^2*c^2*e*e^(a*c + 4*d) + 4*b*c*e^
2*e^(a*c + 4*d) + 48*e^3*e^(a*c + 4*d))*e^(4*e*x) + 3*(b^3*c^3*e^(a*c + 2*
d) - 28*b*c*e^2*e^(a*c + 2*d) - 48*e^3*e^(a*c + 2*d))*e^(2*e*x))*e^(b*c*x)
/(b^4*c^4 - 12*b^3*c^3*e + 44*b^2*c^2*e^2 - 48*b*c*e^3 - (b^4*c^4*e^(6*d)
- 12*b^3*c^3*e*e^(6*d) + 44*b^2*c^2*e^2*e^(6*d) - 48*b*c*e^3*e^(6*d))*e^(6
*e*x) + 3*(b^4*c^4*e^(4*d) - 12*b^3*c^3*e*e^(4*d) + 44*b^2*c^2*e^2*e^(4*d)
- 48*b*c*e^3*e^(4*d))*e^(4*e*x) - 3*(b^4*c^4*e^(2*d) - 12*b^3*c^3*e*e^(2*
d) + 44*b^2*c^2*e^2*e^(2*d) - 48*b*c*e^3*e^(2*d))*e^(2*e*x))

```

**Giac [F]**

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \int \coth(ex+d)^3 e^{((bx+a)c)} dx$$

input

```
integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="giac")
```

output

```
integrate(coth(e*x + d)^3*e^((b*x + a)*c), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \int \coth(d+ex)^3 e^{c(a+bx)} dx$$

input

```
int(coth(d + e*x)^3*exp(c*(a + b*x)),x)
```

output `int(coth(d + e*x)^3*exp(c*(a + b*x)), x)`

## Reduce [F]

$$\int e^{c(a+bx)} \coth^3(d+ex) dx = \text{too large to display}$$

input `int(exp(c*(b*x+a))*coth(e*x+d)^3,x)`

output

```
(e**(a*c)*(e**(b*c*x + 4*d + 4*e*x)*b**2*c**2 - 6*e**(b*c*x + 4*d + 4*e*x)
*b*c*e + 8*e**(b*c*x + 4*d + 4*e*x)*e**2 + 4*e**(b*c*x + 2*d + 2*e*x)*b**2
*c**2 - 12*e**(b*c*x + 2*d + 2*e*x)*b*c*e - 16*e**(b*c*x + 2*d + 2*e*x)*e*
*2 + 7*e**(b*c*x)*b**2*c**2 + 6*e**(b*c*x)*b*c*e + 8*e**(b*c*x)*e**2 + 8*e
**(4*d + 4*e*x)*int(e**(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6
*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e**2 - 3*e**(4*d + 4*e*x)*b**2*c**2 + 18*
e**(4*d + 4*e*x)*b*c*e - 24*e**(4*d + 4*e*x)*e**2 + 3*e**(2*d + 2*e*x)*b**
2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e + 24*e**(2*d + 2*e*x)*e**2 - b**2*c**2
+ 6*b*c*e - 8*e**2),x)*b**5*c**5 - 48*e**(4*d + 4*e*x)*int(e**(b*c*x)/(e**
(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e
*2 - 3*e**(4*d + 4*e*x)*b**2*c**2 + 18*e**(4*d + 4*e*x)*b*c*e - 24*e**(4*d
+ 4*e*x)*e**2 + 3*e**(2*d + 2*e*x)*b**2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e
+ 24*e**(2*d + 2*e*x)*e**2 - b**2*c**2 + 6*b*c*e - 8*e**2),x)*b**4*c**4*e
+ 80*e**(4*d + 4*e*x)*int(e**(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6
*d + 6*e*x)*b*c*e + 8*e**(6*d + 6*e*x)*e**2 - 3*e**(4*d + 4*e*x)*b**2*c**2
+ 18*e**(4*d + 4*e*x)*b*c*e - 24*e**(4*d + 4*e*x)*e**2 + 3*e**(2*d + 2*e*
x)*b**2*c**2 - 18*e**(2*d + 2*e*x)*b*c*e + 24*e**(2*d + 2*e*x)*e**2 - b**2
*c**2 + 6*b*c*e - 8*e**2),x)*b**3*c**3*e**2 - 96*e**(4*d + 4*e*x)*int(e**
(b*c*x)/(e**(6*d + 6*e*x)*b**2*c**2 - 6*e**(6*d + 6*e*x)*b*c*e + 8*e**(6*d
+ 6*e*x)*e**2 - 3*e**(4*d + 4*e*x)*b**2*c**2 + 18*e**(4*d + 4*e*x)*b*c*...
```

### 3.245 $\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx$

Optimal result	1795
Mathematica [A] (verified)	1796
Rubi [A] (verified)	1796
Maple [C] (warning: unable to verify)	1798
Fricas [B] (verification not implemented)	1799
Sympy [F(-1)]	1800
Maxima [A] (verification not implemented)	1801
Giac [A] (verification not implemented)	1801
Mupad [F(-1)]	1802
Reduce [B] (verification not implemented)	1802

#### Optimal result

Integrand size = 25, antiderivative size = 311

$$\begin{aligned}
 \int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx &= \frac{e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc} \\
 &- \frac{4e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})^4} \\
 &+ \frac{26e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{3bc(1 + e^{2c(a+bx)})^3} \\
 &- \frac{55e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{6bc(1 + e^{2c(a+bx)})^2} \\
 &+ \frac{25e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{4bc(1 + e^{2c(a+bx)})} \\
 &- \frac{15 \arctan(e^{c(a+bx)}) \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{4bc}
 \end{aligned}$$

output

```
exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c-4*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^4+26/3*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^3-55/6*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^2+25/4*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))-15/4*arctan(exp(c*(b*x+a)))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.43

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = \frac{\left( e^{c(a+bx)} (33 + 157e^{2c(a+bx)} + 187e^{4c(a+bx)} + 123e^{6c(a+bx)} + 12e^{8c(a+bx)}) - 45(1 + e^{2c(a+bx)})^4 \right) a}{12bc(1 + e^{2c(a+bx)})^4}$$

input

```
Integrate[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(5/2), x]
```

output

```
((E^(c*(a + b*x))*(33 + 157*E^(2*c*(a + b*x)) + 187*E^(4*c*(a + b*x)) + 123*E^(6*c*(a + b*x)) + 12*E^(8*c*(a + b*x))) - 45*(1 + E^(2*c*(a + b*x)))^4 *ArcTan[E^(c*(a + b*x))])*Coth[c*(a + b*x)]*Sqrt[Tanh[c*(a + b*x)]^2]/(12*b*c*(1 + E^(2*c*(a + b*x)))^4)
```

**Rubi [A] (verified)**Time = 1.52 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.52, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx$$

$$\begin{aligned}
 & \downarrow 7271 \\
 & \sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int e^{c(a+bx)} \tanh^5(ac + bcx) dx \\
 & \downarrow 2720 \\
 & \frac{\sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int -\frac{(1-e^{2c(a+bx)})^5}{(1+e^{2c(a+bx)})^5} de^{c(a+bx)}}{bc} \\
 & \downarrow 25 \\
 & -\frac{\sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int \frac{(1-e^{2c(a+bx)})^5}{(1+e^{2c(a+bx)})^5} de^{c(a+bx)}}{bc} \\
 & \downarrow 300 \\
 & -\frac{\sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int \left( \frac{2(1+10e^{4c(a+bx)}+5e^{8c(a+bx)})}{(1+e^{2c(a+bx)})^5} - 1 \right) de^{c(a+bx)}}{bc} \\
 & \downarrow 2009 \\
 & \frac{\left( -\frac{15}{4} \arctan(e^{c(a+bx)}) + e^{c(a+bx)} + \frac{25e^{c(a+bx)}}{4(e^{2c(a+bx)}+1)} - \frac{55e^{c(a+bx)}}{6(e^{2c(a+bx)}+1)^2} + \frac{26e^{c(a+bx)}}{3(e^{2c(a+bx)}+1)^3} - \frac{4e^{c(a+bx)}}{(e^{2c(a+bx)}+1)^4} \right) \sqrt{\tanh^2(ac + bcx)}}{bc}
 \end{aligned}$$

input

Int [E^(c\*(a + b\*x))\*(Tanh[a\*c + b\*c\*x]^2)^(5/2), x]

output

((E^(c\*(a + b\*x)) - (4\*E^(c\*(a + b\*x))))/(1 + E^(2\*c\*(a + b\*x)))^4 + (26\*E^(c\*(a + b\*x)))/(3\*(1 + E^(2\*c\*(a + b\*x)))^3) - (55\*E^(c\*(a + b\*x)))/(6\*(1 + E^(2\*c\*(a + b\*x)))^2) + (25\*E^(c\*(a + b\*x)))/(4\*(1 + E^(2\*c\*(a + b\*x)))) - (15\*ArcTan[E^(c\*(a + b\*x))])/4)\*Coth[a\*c + b\*c\*x]\*Sqrt[Tanh[a\*c + b\*c\*x]^2]/(b\*c)

**Defintions of rubi rules used**

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
  
- rule 300 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int [PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`
  
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
  
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
  
- rule 7271 `Int[(u_)*((a_)*(v_)^(m_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ[v, x] && EqQ[m, 1])`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.98 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.63

method	result
default	$\frac{\text{csgn}(\tanh(c(bx+a))) \left( \frac{\sinh(bcx+ac)^5}{\cosh(bcx+ac)^4} + \frac{5 \sinh(bcx+ac)^3}{\cosh(bcx+ac)^4} + \frac{5 \sinh(bcx+ac)}{\cosh(bcx+ac)^4} - 5 \left( \frac{\text{sech}(bcx+ac)^3}{4} + \frac{3 \text{sech}(bcx+ac)}{8} \right) \tanh(bcx+ac) - \frac{15 \arctan(\tanh(bcx+ac))}{cb} \right)}{cb}$
risch	$\frac{(1+e^{2c(bx+a)}) \sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}} e^{c(bx+a)}}{(e^{2c(bx+a)}-1)bc} + \frac{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}} e^{c(bx+a)} (75 e^{6c(bx+a)} + 115 e^{4c(bx+a)} + 109 e^{2c(bx+a)} + 21)}{12(e^{2c(bx+a)}-1)(1+e^{2c(bx+a)})^3 cb} + \dots$

input `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x,method=_RETURNVERBOSE)`

output `csgn(tanh(c*(b*x+a)))/c/b*(sinh(b*c*x+a*c)^5/cosh(b*c*x+a*c)^4+5*sinh(b*c*x+a*c)^3/cosh(b*c*x+a*c)^4+5*sinh(b*c*x+a*c)/cosh(b*c*x+a*c)^4-5*(1/4*sech(b*c*x+a*c)^3+3/8*sech(b*c*x+a*c))*tanh(b*c*x+a*c)-15/4*arctan(exp(b*c*x+a*c))+sinh(b*c*x+a*c)^4/cosh(b*c*x+a*c)^3+4*sinh(b*c*x+a*c)^2/cosh(b*c*x+a*c)^3+8/3/cosh(b*c*x+a*c)^3)`

### **Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1226 vs.  $2(281) = 562$ .

Time = 0.10 (sec) , antiderivative size = 1226, normalized size of antiderivative = 3.94

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = \text{Too large to display}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="fricas")`



output

```

1/12*(12*cosh(b*c*x + a*c)^9 + 108*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^8 +
12*sinh(b*c*x + a*c)^9 + 3*(144*cosh(b*c*x + a*c)^2 + 41)*sinh(b*c*x + a*
c)^7 + 123*cosh(b*c*x + a*c)^7 + 21*(48*cosh(b*c*x + a*c)^3 + 41*cosh(b*c*
x + a*c))*sinh(b*c*x + a*c)^6 + (1512*cosh(b*c*x + a*c)^4 + 2583*cosh(b*c*
x + a*c)^2 + 187)*sinh(b*c*x + a*c)^5 + 187*cosh(b*c*x + a*c)^5 + (1512*co
sh(b*c*x + a*c)^5 + 4305*cosh(b*c*x + a*c)^3 + 935*cosh(b*c*x + a*c))*sinh
(b*c*x + a*c)^4 + (1008*cosh(b*c*x + a*c)^6 + 4305*cosh(b*c*x + a*c)^4 + 1
870*cosh(b*c*x + a*c)^2 + 157)*sinh(b*c*x + a*c)^3 + 157*cosh(b*c*x + a*c)
^3 + (432*cosh(b*c*x + a*c)^7 + 2583*cosh(b*c*x + a*c)^5 + 1870*cosh(b*c*x
+ a*c)^3 + 471*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^2 - 45*(cosh(b*c*x +
a*c)^8 + 8*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^7 + sinh(b*c*x + a*c)^8 + 4
*(7*cosh(b*c*x + a*c)^2 + 1)*sinh(b*c*x + a*c)^6 + 4*cosh(b*c*x + a*c)^6 +
8*(7*cosh(b*c*x + a*c)^3 + 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^5 + 2*(
35*cosh(b*c*x + a*c)^4 + 30*cosh(b*c*x + a*c)^2 + 3)*sinh(b*c*x + a*c)^4 +
6*cosh(b*c*x + a*c)^4 + 8*(7*cosh(b*c*x + a*c)^5 + 10*cosh(b*c*x + a*c)^3
+ 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^3 + 4*(7*cosh(b*c*x + a*c)^6 + 1
5*cosh(b*c*x + a*c)^4 + 9*cosh(b*c*x + a*c)^2 + 1)*sinh(b*c*x + a*c)^2 + 4
*cosh(b*c*x + a*c)^2 + 8*(cosh(b*c*x + a*c)^7 + 3*cosh(b*c*x + a*c)^5 + 3*
cosh(b*c*x + a*c)^3 + cosh(b*c*x + a*c))*sinh(b*c*x + a*c) + 1)*arctan(cos
h(b*c*x + a*c) + sinh(b*c*x + a*c)) + (108*cosh(b*c*x + a*c)^8 + 861*co...

```

**Sympy [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = \text{Timed out}$$

input

```
integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(5/2),x)
```

output

Timed out

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.47

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = -\frac{15 \arctan(e^{(bcx+ac)})}{4bc} + \frac{12e^{(9bcx+9ac)} + 123e^{(7bcx+7ac)} + 187e^{(5bcx+5ac)} + 157e^{(3bcx+3ac)} + 33e^{(bcx+ac)}}{12bc(e^{(8bcx+8ac)} + 4e^{(6bcx+6ac)} + 6e^{(4bcx+4ac)} + 4e^{(2bcx+2ac)} + 1)}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="maxima")`

output `-15/4*arctan(e^(b*c*x + a*c))/(b*c) + 1/12*(12*e^(9*b*c*x + 9*a*c) + 123*e^(7*b*c*x + 7*a*c) + 187*e^(5*b*c*x + 5*a*c) + 157*e^(3*b*c*x + 3*a*c) + 33*e^(b*c*x + a*c))/(b*c*(e^(8*b*c*x + 8*a*c) + 4*e^(6*b*c*x + 6*a*c) + 6*e^(4*b*c*x + 4*a*c) + 4*e^(2*b*c*x + 2*a*c) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.62

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = -\frac{15 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{4bc} + \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} + \frac{75e^{(7bcx+7ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 115e^{(5bcx+5ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 109e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{12bc(e^{(2bcx+2ac)} + 1)^4}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="giac")`

output `-15/4*arctan(e^(b*c*x + a*c))*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + 1/12*(75*e^(7*b*c*x + 7*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 115*e^(5*b*c*x + 5*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 109*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 21*e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(b*c*(e^(2*b*c*x + 2*a*c) + 1)^4)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = \int e^{c(a+bx)} (\tanh(ac + bcx)^2)^{5/2} dx$$

input `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(5/2), x)`

output `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(5/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.80

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx = \frac{-45e^{8bcx+8ac} \operatorname{atan}(e^{bcx+ac}) - 180e^{6bcx+6ac} \operatorname{atan}(e^{bcx+ac}) - 270e^{4bcx+4ac} \operatorname{atan}(e^{bcx+ac}) - 180e^{2bcx+2ac} \operatorname{atan}(e^{bcx+ac})}{12bc(e^{8bcx+8ac} + 4e^{6bcx+6ac} + 4e^{4bcx+4ac} + 4e^{2bcx+2ac} + 1)}$$

input `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2), x)`

output `( - 45*e**(8*a*c + 8*b*c*x)*atan(e**(a*c + b*c*x)) - 180*e**(6*a*c + 6*b*c*x)*atan(e**(a*c + b*c*x)) - 270*e**(4*a*c + 4*b*c*x)*atan(e**(a*c + b*c*x)) - 180*e**(2*a*c + 2*b*c*x)*atan(e**(a*c + b*c*x)) - 45*atan(e**(a*c + b*c*x)) + 12*e**(9*a*c + 9*b*c*x) + 123*e**(7*a*c + 7*b*c*x) + 187*e**(5*a*c + 5*b*c*x) + 157*e**(3*a*c + 3*b*c*x) + 33*e**(a*c + b*c*x))/(12*b*c*(e**(8*a*c + 8*b*c*x) + 4*e**(6*a*c + 6*b*c*x) + 4*e**(4*a*c + 4*b*c*x) + 4*e**(2*a*c + 2*b*c*x) + 1))`

### 3.246 $\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx$

Optimal result	1803
Mathematica [A] (verified)	1804
Rubi [A] (verified)	1804
Maple [C] (warning: unable to verify)	1806
Fricas [B] (verification not implemented)	1807
Sympy [F(-1)]	1807
Maxima [A] (verification not implemented)	1808
Giac [A] (verification not implemented)	1808
Mupad [F(-1)]	1809
Reduce [B] (verification not implemented)	1809

#### Optimal result

Integrand size = 25, antiderivative size = 193

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx = \frac{e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})^2} + \frac{3e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})} - \frac{3 \arctan(e^{c(a+bx)}) \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc}$$

```
output exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c-2*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^2+3*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))-3*arctan(exp(c*(b*x+a)))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c
```

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.54

$$\int e^{c(a+bx)} \tanh^2(ac + bxc)^{3/2} dx = \frac{\left( e^{c(a+bx)} (2 + 5e^{2c(a+bx)} + e^{4c(a+bx)}) - 3(1 + e^{2c(a+bx)})^2 \arctan(e^{c(a+bx)}) \right) \coth(c(a+bx)) \sqrt{\tanh^2(ac + bxc)}}{bc(1 + e^{2c(a+bx)})^2}$$

input `Integrate[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(3/2),x]`

output `((E^(c*(a + b*x))*(2 + 5E^(2*c*(a + b*x)) + E^(4*c*(a + b*x))) - 3*(1 + E^(2*c*(a + b*x)))^2*ArcTan[E^(c*(a + b*x))])*Coth[c*(a + b*x)]*Sqrt[Tanh[c*(a + b*x)]^2])/(b*c*(1 + E^(2*c*(a + b*x)))^2)`

**Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.53, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{c(a+bx)} \tanh^2(ac + bxc)^{3/2} dx \\ & \quad \downarrow \text{7271} \\ & \sqrt{\tanh^2(ac + bxc) \coth(ac + bxc)} \int e^{c(a+bx)} \tanh^3(ac + bxc) dx \\ & \quad \downarrow \text{2720} \\ & \frac{\sqrt{\tanh^2(ac + bxc) \coth(ac + bxc)} \int -\frac{(1-e^{2c(a+bx)})^3}{(1+e^{2c(a+bx)})^3} de^{c(a+bx)}}{bc} \\ & \quad \downarrow \text{25} \end{aligned}$$

$$\begin{array}{c}
 \frac{\sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int \frac{(1 - e^{2c(a+bx)})^3}{(1 + e^{2c(a+bx)})^3} de^{c(a+bx)}}{bc} \\
 \downarrow 300 \\
 \frac{\sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)} \int \left( \frac{2(1 + 3e^{4c(a+bx)})}{(1 + e^{2c(a+bx)})^3} - 1 \right) de^{c(a+bx)}}{bc} \\
 \downarrow 2009 \\
 \frac{\left( -3 \arctan(e^{c(a+bx)}) + e^{c(a+bx)} + \frac{3e^{c(a+bx)}}{e^{2c(a+bx)} + 1} - \frac{2e^{c(a+bx)}}{(e^{2c(a+bx)} + 1)^2} \right) \sqrt{\tanh^2(ac + bcx) \coth(ac + bcx)}}{bc}
 \end{array}$$

input `Int[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(3/2), x]`

output `((E^(c*(a + b*x)) - (2*E^(c*(a + b*x))))/(1 + E^(2*c*(a + b*x)))^2 + (3*E^(c*(a + b*x)))/(1 + E^(2*c*(a + b*x))) - 3*ArcTan[E^(c*(a + b*x))]*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 300 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 7271

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p},
x] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(Eq
Q[v, x] && EqQ[m, 1])
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.61 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.68

method	result
default	$\frac{\operatorname{csgn}(\tanh(c(bx+a))) \left( \frac{\sinh(bcx+ac)^3}{\cosh(bcx+ac)^2} + \frac{3 \sinh(bcx+ac)}{\cosh(bcx+ac)^2} - \frac{3 \operatorname{sech}(bcx+ac) \tanh(bcx+ac)}{2} - 3 \arctan(e^{bcx+ac}) + \frac{\sinh(bcx+ac)^2}{\cosh(bcx+ac)} + \frac{2}{\cosh(bcx+ac)} \right)}{cb}$
risch	$\frac{\sqrt{\frac{(e^{2c(bx+a)} - 1)^2}{(1 + e^{2c(bx+a)})^2}} (3i \ln(e^{c(bx+a)} - i) e^{4c(bx+a)} - 3i \ln(e^{c(bx+a)} + i) e^{4c(bx+a)} + 2e^{5c(bx+a)} + 6i \ln(e^{c(bx+a)} - i) e^{2c(bx+a)} - 6i \ln(e^{c(bx+a)} + i) e^{2c(bx+a)})}{2(e^{2c(bx+a)} - 1)(1 + e^{2c(bx+a)})cb}$

input

```
int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2), x, method=_RETURNVERBOSE)
```

output

```
csgn(tanh(c*(b*x+a)))/c/b*(sinh(b*c*x+a*c)^3/cosh(b*c*x+a*c)^2+3*sinh(b*c*
x+a*c)/cosh(b*c*x+a*c)^2-3/2*sech(b*c*x+a*c)*tanh(b*c*x+a*c)-3*arctan(exp(
b*c*x+a*c))+sinh(b*c*x+a*c)^2/cosh(b*c*x+a*c)+2/cosh(b*c*x+a*c))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 458 vs.  $2(179) = 358$ .

Time = 0.11 (sec) , antiderivative size = 458, normalized size of antiderivative = 2.37

$$\int e^{c(a+bx)} \tanh^2(ac + b cx)^{3/2} dx = \frac{\cosh(bc x + ac)^5 + 5 \cosh(bc x + ac) \sinh(bc x + ac)^4 + \sinh(bc x + ac)^5 + 5(2 \cosh(bc x + ac) \sinh(bc x + ac)^3 + \cosh(bc x + ac)^2 \sinh(bc x + ac)^2 + \sinh(bc x + ac)^4 + 2(3 \cosh(bc x + ac)^2 + 1) \sinh(bc x + ac)^2 + 2 \cosh(bc x + ac)^2 + 4(\cosh(bc x + ac)^3 + \cosh(bc x + ac)) \sinh(bc x + ac) + 1) \arctan(\cosh(bc x + ac) + \sinh(bc x + ac)) + (5 \cosh(bc x + ac)^4 + 15 \cosh(bc x + ac)^2 + 2) \sinh(bc x + ac) + 2 \cosh(bc x + ac)}{(b^2 c^2 \cosh(bc x + ac)^4 + 4 b^2 c \cosh(bc x + ac) \sinh(bc x + ac)^3 + b^2 c \sinh(bc x + ac)^4 + 2 b^2 c \cosh(bc x + ac)^2 + 2(3 b^2 c \cosh(bc x + ac)^2 + b^2 c) \sinh(bc x + ac)^2 + b^2 c + 4(b^2 c \cosh(bc x + ac)^3 + b^2 c \cosh(bc x + ac)) \sinh(bc x + ac)}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="fricas")`

output `(cosh(b*c*x + a*c)^5 + 5*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^4 + sinh(b*c*x + a*c)^5 + 5*(2*cosh(b*c*x + a*c)^2 + 1)*sinh(b*c*x + a*c)^3 + 5*cosh(b*c*x + a*c)^3 + 5*(2*cosh(b*c*x + a*c)^3 + 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^2 - 3*(cosh(b*c*x + a*c)^4 + 4*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 + sinh(b*c*x + a*c)^4 + 2*(3*cosh(b*c*x + a*c)^2 + 1)*sinh(b*c*x + a*c)^2 + 2*cosh(b*c*x + a*c)^2 + 4*(cosh(b*c*x + a*c)^3 + cosh(b*c*x + a*c))*sinh(b*c*x + a*c) + 1)*arctan(cosh(b*c*x + a*c) + sinh(b*c*x + a*c)) + (5*cosh(b*c*x + a*c)^4 + 15*cosh(b*c*x + a*c)^2 + 2)*sinh(b*c*x + a*c) + 2*cosh(b*c*x + a*c))/(b*c*cosh(b*c*x + a*c)^4 + 4*b*c*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 + b*c*sinh(b*c*x + a*c)^4 + 2*b*c*cosh(b*c*x + a*c)^2 + 2*(3*b*c*cosh(b*c*x + a*c)^2 + b*c)*sinh(b*c*x + a*c)^2 + b*c + 4*(b*c*cosh(b*c*x + a*c)^3 + b*c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c))`

**Sympy [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^2(ac + b cx)^{3/2} dx = \text{Timed out}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(3/2),x)`

output `Timed out`



**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.47

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx = -\frac{3 \arctan(e^{(bcx+ac)})}{bc} + \frac{e^{(5bcx+5ac)} + 5e^{(3bcx+3ac)} + 2e^{(bcx+ac)}}{bc(e^{(4bcx+4ac)} + 2e^{(2bcx+2ac)} + 1)}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="maxima")`

output `-3*arctan(e^(b*c*x + a*c))/(b*c) + (e^(5*b*c*x + 5*a*c) + 5*e^(3*b*c*x + 3*a*c) + 2*e^(b*c*x + a*c))/(b*c*(e^(4*b*c*x + 4*a*c) + 2*e^(2*b*c*x + 2*a*c) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.71

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx = -\frac{3 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} + \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} + \frac{3e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc(e^{(2bcx+2ac)} + 1)^2}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="giac")`

output `-3*arctan(e^(b*c*x + a*c))*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + (3*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(b*c*(e^(2*b*c*x + 2*a*c) + 1)^2)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx = \int e^{c(a+bx)} (\tanh(ac + bcx)^2)^{3/2} dx$$

input `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(3/2),x)`

output `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.72

$$\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx = \frac{-3e^{4bcx+4ac} \operatorname{atan}(e^{bcx+ac}) - 6e^{2bcx+2ac} \operatorname{atan}(e^{bcx+ac}) - 3\operatorname{atan}(e^{bcx+ac}) + e^{5bcx+5ac} + 5e^{3bcx+3ac}}{bc(e^{4bcx+4ac} + 2e^{2bcx+2ac} + 1)}$$

input `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x)`

output `( - 3*e**(4*a*c + 4*b*c*x)*atan(e**(a*c + b*c*x)) - 6*e**(2*a*c + 2*b*c*x)*atan(e**(a*c + b*c*x)) - 3*atan(e**(a*c + b*c*x)) + e**(5*a*c + 5*b*c*x) + 5*e**(3*a*c + 3*b*c*x) + 2*e**(a*c + b*c*x))/(b*c*(e**(4*a*c + 4*b*c*x) + 2*e**(2*a*c + 2*b*c*x) + 1))`

### 3.247 $\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx$

Optimal result	1810
Mathematica [A] (verified)	1810
Rubi [A] (verified)	1811
Maple [C] (verified)	1813
Fricas [A] (verification not implemented)	1813
Sympy [F]	1814
Maxima [A] (verification not implemented)	1814
Giac [A] (verification not implemented)	1814
Mupad [F(-1)]	1815
Reduce [B] (verification not implemented)	1815

#### Optimal result

Integrand size = 25, antiderivative size = 83

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx = \frac{e^{c(a+bx)} \operatorname{coth}(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc} - \frac{2 \arctan(e^{c(a+bx)}) \operatorname{coth}(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc}$$

output `exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c-2*arctan(exp(c*(b*x+a)))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.61

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx = \frac{(e^{c(a+bx)} - 2 \arctan(e^{c(a+bx)})) \operatorname{coth}(c(a + bx)) \sqrt{\tanh^2(c(a + bx))}}{bc}$$

input `Integrate[E^(c*(a + b*x))*Sqrt[Tanh[a*c + b*c*x]^2],x]`

output

$$\frac{((E^{c(a+bx)}) - 2*ArcTan[E^{c(a+bx)}])*Coth[c(a+bx)]*Sqrt[Tanh[c(a+bx)]^2])}{(b*c)}$$
**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.64, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx \\ & \quad \downarrow \text{7271} \\ & \frac{\sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx) \int e^{c(a+bx)} \tanh(ac+bcx) dx}{bc} \\ & \quad \downarrow \text{2720} \\ & \frac{\sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx) \int -\frac{1-e^{2c(a+bx)}}{1+e^{2c(a+bx)}} de^{c(a+bx)}}{bc} \\ & \quad \downarrow \text{25} \\ & -\frac{\sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx) \int \frac{1-e^{2c(a+bx)}}{1+e^{2c(a+bx)}} de^{c(a+bx)}}{bc} \\ & \quad \downarrow \text{299} \\ & \frac{\sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx) \left( e^{c(a+bx)} - 2 \int \frac{1}{1+e^{2c(a+bx)}} de^{c(a+bx)} \right)}{bc} \\ & \quad \downarrow \text{216} \\ & \frac{(e^{c(a+bx)} - 2 \arctan(e^{c(a+bx)})) \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{bc} \end{aligned}$$

input

$$\text{Int}[E^{c(a+bx)}*Sqrt[Tanh[a*c + b*c*x]^2], x]$$

output  $((E^{(c*(a + b*x))} - 2*ArcTan[E^{(c*(a + b*x))}])*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c)$

### Defintions of rubi rules used

rule 25  $Int[-(Fx_), x\_Symbol] \rightarrow Simp[Identity[-1] Int[Fx, x], x]$

rule 216  $Int[((a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& PosQ[a/b] \&\& (GtQ[a, 0] || GtQ[b, 0])$

rule 299  $Int[((a_) + (b_)*(x_)^2)^{(p_)}*((c_) + (d_)*(x_)^2), x\_Symbol] \rightarrow Simp[d*x*((a + b*x^2)^{(p + 1})/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] \&\& NeQ[b*c - a*d, 0] \&\& NeQ[2*p + 3, 0]$

rule 2720  $Int[u_, x\_Symbol] \rightarrow With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] \&\& !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] \&\& IntegerQ[m*n] \&\& !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] \&\& InverseFunctionQ[F[x]]$

rule 7271  $Int[(u_)*((a_)*(v_)^(m_))^(p_), x\_Symbol] \rightarrow Simp[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x] \&\& !IntegerQ[p] \&\& !FreeQ[v, x] \&\& !(EqQ[a, 1] \&\& EqQ[m, 1]) \&\& !(EqQ[v, x] \&\& EqQ[m, 1])$

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 1.88 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.63

method	result
risch	$\frac{(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}e^{c(bx+a)}}{(e^{2c(bx+a)}-1)bc} + \frac{i(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}\ln(e^{c(bx+a)}-i)}{(e^{2c(bx+a)}-1)cb} - \frac{i(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}}{(e^{2c(bx+a)}-1)cb}$

input

```
int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/(exp(2*c*(b*x+a))-1)*(1+exp(2*c*(b*x+a)))*((exp(2*c*(b*x+a))-1)^2/(1+exp(2*c*(b*x+a)))^2)^(1/2)*exp(c*(b*x+a))/b/c+I*((exp(2*c*(b*x+a))-1)^2/(1+exp(2*c*(b*x+a)))^2)^(1/2)/(exp(2*c*(b*x+a))-1)*(1+exp(2*c*(b*x+a)))/c/b*ln(exp(c*(b*x+a))-I)-I*((exp(2*c*(b*x+a))-1)^2/(1+exp(2*c*(b*x+a)))^2)^(1/2)/(exp(2*c*(b*x+a))-1)*(1+exp(2*c*(b*x+a)))/c/b*ln(exp(c*(b*x+a))+I)
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.64

$$\int e^{c(a+bx)}\sqrt{\tanh^2(ac+bcx)}dx$$

$$= -\frac{2\arctan(\cosh(bcx+ac)+\sinh(bcx+ac))-\cosh(bcx+ac)-\sinh(bcx+ac)}{bc}$$

input

```
integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="fricas")
```

output

```
-(2*arctan(cosh(b*c*x + a*c) + sinh(b*c*x + a*c)) - cosh(b*c*x + a*c) - sinh(b*c*x + a*c))/(b*c)
```

**Sympy [F]**

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx = e^{ac} \int \sqrt{\tanh^2(ac + bcx)} e^{bcx} dx$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(1/2),x)`

output `exp(a*c)*Integral(sqrt(tanh(a*c + b*c*x)**2)*exp(b*c*x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.42

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx = -\frac{2 \arctan(e^{(bcx+ac)})}{bc} + \frac{e^{(bcx+ac)}}{bc}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="maxima")`

output `-2*arctan(e^(b*c*x + a*c))/(b*c) + e^(b*c*x + a*c)/(b*c)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.76

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} dx = -\frac{2 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} + \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc}$$

input `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="giac")`

output `-2*arctan(e^(b*c*x + a*c))*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c)`

**Mupad [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx = \int e^{c(a+bx)} \sqrt{\tanh(ac+bcx)^2} dx$$

input `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(1/2),x)`

output `int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.37

$$\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx = \frac{-2\operatorname{atan}(e^{bcx+ac}) + e^{bcx+ac}}{bc}$$

input `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x)`

output `( - 2*atan(e**(a*c + b*c*x)) + e**(a*c + b*c*x))/(b*c)`



**3.248** 
$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$$

Optimal result	1816
Mathematica [A] (verified)	1816
Rubi [A] (verified)	1817
Maple [C] (warning: unable to verify)	1819
Fricas [A] (verification not implemented)	1819
Sympy [F]	1820
Maxima [A] (verification not implemented)	1820
Giac [A] (verification not implemented)	1820
Mupad [F(-1)]	1821
Reduce [B] (verification not implemented)	1821

**Optimal result**

Integrand size = 25, antiderivative size = 83

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx = \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2\arctanh(e^{c(a+bx)}) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}}$$

output `exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-2*arctanh(exp(c*(b*x+a)))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)`

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.61

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx = \frac{(e^{c(a+bx)} - 2\arctanh(e^{c(a+bx)})) \tanh(c(a+bx))}{bc\sqrt{\tanh^2(c(a+bx))}}$$

input `Integrate[E^(c*(a + b*x))/Sqrt[Tanh[a*c + b*c*x]^2], x]`

output `((E^(c*(a + b*x)) - 2*ArcTanh[E^(c*(a + b*x))])*Tanh[c*(a + b*x)]/(b*c*Sqrt[Tanh[c*(a + b*x)]^2])`

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.64, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx \\
 & \quad \downarrow \text{7271} \\
 & \frac{\tanh(ac+bcx) \int e^{c(a+bx)} \coth(ac+bcx) dx}{\sqrt{\tanh^2(ac+bcx)}} \\
 & \quad \downarrow \text{2720} \\
 & \frac{\tanh(ac+bcx) \int -\frac{1+e^{2c(a+bx)}}{1-e^{2c(a+bx)}} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ac+bcx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\tanh(ac+bcx) \int \frac{1+e^{2c(a+bx)}}{1-e^{2c(a+bx)}} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ac+bcx)}} \\
 & \quad \downarrow \text{299} \\
 & \frac{\tanh(ac+bcx) \left( e^{c(a+bx)} - 2 \int \frac{1}{1-e^{2c(a+bx)}} de^{c(a+bx)} \right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
 & \quad \downarrow \text{219} \\
 & \frac{(e^{c(a+bx)} - 2\operatorname{arctanh}(e^{c(a+bx)})) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}}
 \end{aligned}$$

input

```
Int[E^(c*(a + b*x))/Sqrt[Tanh[a*c + b*c*x]^2], x]
```

output  $((E^{(c*(a + b*x))} - 2*ArcTanh[E^{(c*(a + b*x))}])*Tanh[a*c + b*c*x]/(b*c*Sqrt[Tanh[a*c + b*c*x]^2])$

### Defintions of rubi rules used

rule 25  $Int[-(Fx_), x\_Symbol] \rightarrow Simp[Identity[-1] Int[Fx, x], x]$

rule 219  $Int[((a_) + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] \&\& NegQ[a/b] \&\& (GtQ[a, 0] || LtQ[b, 0])$

rule 299  $Int[((a_) + (b_)*(x_)^2)^{(p_)}*((c_) + (d_)*(x_)^2), x\_Symbol] \rightarrow Simp[d*x*((a + b*x^2)^{(p + 1})/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] \&\& NeQ[b*c - a*d, 0] \&\& NeQ[2*p + 3, 0]$

rule 2720  $Int[u_, x\_Symbol] \rightarrow With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] \&\& !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] \&\& IntegerQ[m*n] \&\& !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] \&\& InverseFunctionQ[F[x]]$

rule 7271  $Int[(u_)*((a_)*(v_)^(m_))^(p_), x\_Symbol] \rightarrow Simp[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x] \&\& !IntegerQ[p] \&\& !FreeQ[v, x] \&\& !(EqQ[a, 1] \&\& EqQ[m, 1]) \&\& !(EqQ[v, x] \&\& EqQ[m, 1])$

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.58

method	result	size
default	$\frac{\text{csgn}(\tanh(c(bx+a)))(\sinh(bcx+ac)+\cosh(bcx+ac)-2 \operatorname{arctanh}(e^{bcx+ac}))}{cb}$	48
risch	$\frac{(e^{2c(bx+a)}-1)e^{c(bx+a)}}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc} + \frac{(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)}-1)}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})cb} - \frac{(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)}+1)}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})cb}$	213

input `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `csgn(tanh(c*(b*x+a)))/c/b*(sinh(b*c*x+a*c)+cosh(b*c*x+a*c)-2*arctanh(exp(b*c*x+a*c)))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$$

$$= \frac{\cosh(bcx+ac) - \log(\cosh(bcx+ac) + \sinh(bcx+ac) + 1) + \log(\cosh(bcx+ac) + \sinh(bcx+ac) - 1) + \sinh(bcx+ac)}{bc}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="fricas")`

output `(cosh(b*c*x + a*c) - log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) + 1) + log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) - 1) + sinh(b*c*x + a*c))/(b*c)`

**Sympy [F]**

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx = e^{ac} \int \frac{e^{bcx}}{\sqrt{\tanh^2(ac+bcx)}} dx$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(1/2),x)`

output `exp(a*c)*Integral(exp(b*c*x)/sqrt(tanh(a*c + b*c*x)**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.67

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx = \frac{e^{(bcx+ac)}}{bc} - \frac{\log(e^{(bcx+ac)} + 1)}{bc} + \frac{\log(e^{(bcx+ac)} - 1)}{bc}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="maxima")`

output `e^(b*c*x + a*c)/(b*c) - log(e^(b*c*x + a*c) + 1)/(b*c) + log(e^(b*c*x + a*c) - 1)/(b*c)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.19

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx = \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} - \frac{\log(e^{(bcx+ac)} + 1) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} + \frac{\log(|e^{(bcx+ac)} - 1|) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="giac")`

output  $e^{(b*c*x + a*c)} * \text{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) / (b*c) - \log(e^{(b*c*x + a*c)} + 1) * \text{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) / (b*c) + \log(\text{abs}(e^{(b*c*x + a*c)} - 1)) * \text{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) / (b*c)$

### Mupad [F(-1)]

Timed out.

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac + bcx)}} dx = \int \frac{e^{c(a+bx)}}{\sqrt{\tanh(ac + bcx)^2}} dx$$

input `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(1/2),x)`

output `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(1/2), x)`

### Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.70

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac + bcx)}} dx = \frac{e^{bcx+ac} - \log(e^{bcx+2ac} + e^{ac}) + \log(e^{bcx+2ac} - e^{ac})}{bc}$$

input `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x)`

output  $(e^{(a*c + b*c*x)} - \log(e^{(2*a*c + b*c*x)} + e^{(a*c)}) + \log(e^{(2*a*c + b*c*x)} - e^{(a*c)})) / (b*c)$

**3.249** 
$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$$

Optimal result	1822
Mathematica [C] (warning: unable to verify)	1823
Rubi [A] (verified)	1823
Maple [C] (warning: unable to verify)	1825
Fricas [B] (verification not implemented)	1826
Sympy [F]	1827
Maxima [A] (verification not implemented)	1827
Giac [A] (verification not implemented)	1828
Mupad [F(-1)]	1828
Reduce [B] (verification not implemented)	1829

**Optimal result**

Integrand size = 25, antiderivative size = 197

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{3e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)}) \sqrt{\tanh^2(ac+bcx)}} - \frac{3\arctanh(e^{c(a+bx)}) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}}$$

output

```
exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-2*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^2/(tanh(b*c*x+a*c)^2)^(1/2)+3*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))/(tanh(b*c*x+a*c)^2)^(1/2)-3*arctanh(exp(c*(b*x+a)))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)
```

**Mathematica [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.14 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.70

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = e^{-5c(a+bx)} \left( -21(252105 + 507305e^{2c(a+bx)} + 173916e^{4c(a+bx)} - 154296e^{6c(a+bx)} - 73885e^{8c(a+bx)} + 4887e^{10c(a+bx)}) \right)$$

input `Integrate[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(3/2),x]`

output `-1/60480*((-21*(252105 + 507305*E^(2*c*(a + b*x)) + 173916*E^(4*c*(a + b*x)) - 154296*E^(6*c*(a + b*x)) - 73885*E^(8*c*(a + b*x)) + 4887*E^(10*c*(a + b*x))) - (315*(-16807 - 28218*E^(2*c*(a + b*x)) + 1173*E^(4*c*(a + b*x)) + 17748*E^(6*c*(a + b*x)) + 4299*E^(8*c*(a + b*x)) - 1434*E^(10*c*(a + b*x)) + 7*E^(12*c*(a + b*x)))*ArcTanh[Sqrt[E^(2*c*(a + b*x))]]/Sqrt[E^(2*c*(a + b*x))] + 384*E^(8*c*(a + b*x))*(1 + E^(2*c*(a + b*x)))^2*(7 + 5*E^(2*c*(a + b*x)))*HypergeometricPFQ[{3/2, 2, 2, 2, 2}, {1, 1, 1, 11/2}, E^(2*c*(a + b*x))] + 256*E^(8*c*(a + b*x))*(1 + E^(2*c*(a + b*x)))^3*HypergeometricPFQ[{3/2, 2, 2, 2, 2, 2}, {1, 1, 1, 1, 11/2}, E^(2*c*(a + b*x))]*Tanh[c*(a + b*x)]^3)/(b*c*E^(5*c*(a + b*x))*(Tanh[c*(a + b*x)]^2)^(3/2))`

**Rubi [A] (verified)**

Time = 0.89 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.54, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$$

↓ 7271



$$\begin{aligned}
& \frac{\tanh(ax + bx) \int e^{c(a+bx)} \coth^3(ax + bxc) dx}{\sqrt{\tanh^2(ax + bcx)}} \\
& \quad \downarrow \text{2720} \\
& \frac{\tanh(ax + bcx) \int -\frac{(1+e^{2c(a+bx)})^3}{(1-e^{2c(a+bx)})^3} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ax + bcx)}} \\
& \quad \downarrow \text{25} \\
& \frac{\tanh(ax + bcx) \int \frac{(1+e^{2c(a+bx)})^3}{(1-e^{2c(a+bx)})^3} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ax + bcx)}} \\
& \quad \downarrow \text{300} \\
& \frac{\tanh(ax + bcx) \int \left( \frac{2(1+3e^{4c(a+bx)})}{(1-e^{2c(a+bx)})^3} - 1 \right) de^{c(a+bx)}}{bc\sqrt{\tanh^2(ax + bcx)}} \\
& \quad \downarrow \text{2009} \\
& \frac{\left( -3\operatorname{arctanh}(e^{c(a+bx)}) + e^{c(a+bx)} + \frac{3e^{c(a+bx)}}{1-e^{2c(a+bx)}} - \frac{2e^{c(a+bx)}}{(1-e^{2c(a+bx)})^2} \right) \tanh(ax + bcx)}{bc\sqrt{\tanh^2(ax + bcx)}}
\end{aligned}$$

input

```
Int [E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(3/2), x]
```

output

```
((E^(c*(a + b*x)) - (2*E^(c*(a + b*x))))/(1 - E^(2*c*(a + b*x)))^2 + (3*E^(c*(a + b*x)))/(1 - E^(2*c*(a + b*x))) - 3*ArcTanh[E^(c*(a + b*x))]*Tanh[a*c + b*c*x]/(b*c*Sqrt[Tanh[a*c + b*c*x]^2])
```

## Definitions of rubi rules used

rule 25	<code>Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]</code>
rule 300	<code>Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] &amp;&amp; NeQ[b*c - a*d, 0] &amp;&amp; IGtQ[p, 0] &amp;&amp; ILtQ[q, 0] &amp;&amp; GeQ[p, -q]</code>
rule 2009	<code>Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]</code>
rule 2720	<code>Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] &amp;&amp; !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] &amp;&amp; IntegerQ[m*n] &amp;&amp; !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] &amp;&amp; InverseFunctionQ[F[x]]]</code>
rule 7271	<code>Int[(u_)*((a_)*(v_)^(m_))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x] &amp;&amp; !IntegerQ[p] &amp;&amp; !FreeQ[v, x] &amp;&amp; !(EqQ[a, 1] &amp;&amp; EqQ[m, 1]) &amp;&amp; !(EqQ[v, x] &amp;&amp; EqQ[m, 1])</code>

## Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.68 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.66

method	result
default	$\text{csgn}(\tanh(c(bx+a))) \left( \frac{\cosh(bc x+ac)^2}{\sinh(bc x+ac)} - \frac{2}{\sinh(bc x+ac)} + \frac{\cosh(bc x+ac)^3}{\sinh(bc x+ac)^2} - \frac{3 \cosh(bc x+ac)}{\sinh(bc x+ac)^2} + \frac{3 \operatorname{csch}(bc x+ac) \operatorname{coth}(bc x+ac)}{2} - 3 \operatorname{arctanh}(e^{bc x+ac}) \right)$
risch	$\frac{2e^{5c(bx+a)} + 3 \ln(e^{c(bx+a)} - 1) e^{4c(bx+a)} - 3 \ln(e^{c(bx+a)} + 1) e^{4c(bx+a)} - 10e^{3c(bx+a)} - 6 \ln(e^{c(bx+a)} - 1) e^{2c(bx+a)} + 6 \ln(e^{c(bx+a)} + 1) e^{2c(bx+a)}}{2(e^{2c(bx+a)} - 1)(1 + e^{2c(bx+a)})} \sqrt{\frac{(e^{2c(bx+a)} - 1)^2}{(1 + e^{2c(bx+a)})^2}} cb$

input `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2), x, method=_RETURNVERBOSE)`

output

```
csgn(tanh(c*(b*x+a)))/c/b*(1/sinh(b*c*x+a*c)*cosh(b*c*x+a*c)^2-2/sinh(b*c*x+a*c)+cosh(b*c*x+a*c)^3/sinh(b*c*x+a*c)^2-3/sinh(b*c*x+a*c)^2*cosh(b*c*x+a*c)+3/2*csch(b*c*x+a*c)*coth(b*c*x+a*c)-3*arctanh(exp(b*c*x+a*c)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 613 vs.  $2(179) = 358$ .

Time = 0.11 (sec) , antiderivative size = 613, normalized size of antiderivative = 3.11

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ax+bcx)^{3/2}} dx = \text{Too large to display}$$

input

```
integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="fricas")
```

output

```
1/2*(2*cosh(b*c*x + a*c)^5 + 10*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^4 + 2*
sinh(b*c*x + a*c)^5 + 10*(2*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^3 -
10*cosh(b*c*x + a*c)^3 + 10*(2*cosh(b*c*x + a*c)^3 - 3*cosh(b*c*x + a*c))
*sinh(b*c*x + a*c)^2 - 3*(cosh(b*c*x + a*c)^4 + 4*cosh(b*c*x + a*c)*sinh(b
*c*x + a*c)^3 + sinh(b*c*x + a*c)^4 + 2*(3*cosh(b*c*x + a*c)^2 - 1)*sinh(b
*c*x + a*c)^2 - 2*cosh(b*c*x + a*c)^2 + 4*(cosh(b*c*x + a*c)^3 - cosh(b*c*
x + a*c))*sinh(b*c*x + a*c) + 1)*log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c)
+ 1) + 3*(cosh(b*c*x + a*c)^4 + 4*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 +
sinh(b*c*x + a*c)^4 + 2*(3*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^2 -
2*cosh(b*c*x + a*c)^2 + 4*(cosh(b*c*x + a*c)^3 - cosh(b*c*x + a*c))*sinh(
b*c*x + a*c) + 1)*log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) - 1) + 2*(5*co
sh(b*c*x + a*c)^4 - 15*cosh(b*c*x + a*c)^2 + 2)*sinh(b*c*x + a*c) + 4*cosh
(b*c*x + a*c))/(b*c*cosh(b*c*x + a*c)^4 + 4*b*c*cosh(b*c*x + a*c)*sinh(b*c
*x + a*c)^3 + b*c*sinh(b*c*x + a*c)^4 - 2*b*c*cosh(b*c*x + a*c)^2 + 2*(3*b
*c*cosh(b*c*x + a*c)^2 - b*c)*sinh(b*c*x + a*c)^2 + b*c + 4*(b*c*cosh(b*c*
x + a*c)^3 - b*c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c))
```

**Sympy [F]**

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = e^{ac} \int \frac{e^{bcx}}{(\tanh^2(ac+bcx))^{3/2}} dx$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(3/2),x)`

output `exp(a*c)*Integral(exp(b*c*x)/(tanh(a*c + b*c*x)**2)**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.57

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = -\frac{3 \log(e^{(bcx+ac)} + 1)}{2bc} + \frac{3 \log(e^{(bcx+ac)} - 1)}{2bc} + \frac{e^{(5bcx+5ac)} - 5e^{(3bcx+3ac)} + 2e^{(bcx+ac)}}{bc(e^{(4bcx+4ac)} - 2e^{(2bcx+2ac)} + 1)}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="maxima")`

output `-3/2*log(e^(b*c*x + a*c) + 1)/(b*c) + 3/2*log(e^(b*c*x + a*c) - 1)/(b*c) + (e^(5*b*c*x + 5*a*c) - 5*e^(3*b*c*x + 3*a*c) + 2*e^(b*c*x + a*c))/(b*c*(e^(4*b*c*x + 4*a*c) - 2*e^(2*b*c*x + 2*a*c) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.89

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} - \frac{3 \log(e^{(bcx+ac)} + 1) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{2bc} + \frac{3 \log(|e^{(bcx+ac)} - 1|) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{2bc} - \frac{3e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc(e^{(2bcx+2ac)} - 1)^2}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="giac")`

output `e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) - 3/2*log(e^(b*c*x + a*c) + 1)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + 3/2*log(abs(e^(b*c*x + a*c) - 1))*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) - (3*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(b*c*(e^(2*b*c*x + 2*a*c) - 1)^2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = \int \frac{e^{c(a+bx)}}{(\tanh(ac+bcx))^2)^{3/2}} dx$$

input `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(3/2),x)`

output `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.54

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx = \frac{e^{3bcx+3ac} \tanh(bcx+ac)^2 + 2e^{3bcx+3ac} \tanh(bcx+ac) - e^{3bcx+3ac} - 3e^{2bcx+2ac}}{\dots}$$

input `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x)`output `(e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x)**2 + 2*e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x) - e**(3*a*c + 3*b*c*x) - 3*e**(2*a*c + 2*b*c*x)*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**2 + 3*e**(2*a*c + 2*b*c*x)*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**2 - 7*e**(a*c + b*c*x)*tanh(a*c + b*c*x)**2 - 2*e**(a*c + b*c*x)*tanh(a*c + b*c*x) + e**(a*c + b*c*x) + 3*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**2 - 3*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**2)/(2*tanh(a*c + b*c*x)**2*b*c*(e**(2*a*c + 2*b*c*x) - 1))`

**3.250**  $\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$

Optimal result	1830
Mathematica [A] (warning: unable to verify)	1831
Rubi [A] (verified)	1831
Maple [C] (warning: unable to verify)	1833
Fricas [B] (verification not implemented)	1834
Sympy [F(-1)]	1835
Maxima [A] (verification not implemented)	1835
Giac [A] (verification not implemented)	1836
Mupad [F(-1)]	1836
Reduce [B] (verification not implemented)	1837

**Optimal result**

Integrand size = 25, antiderivative size = 319

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)} \tanh(ac+bcx)}{3bc(1-e^{2c(a+bx)})^3 \sqrt{\tanh^2(ac+bcx)}} - \frac{55e^{c(a+bx)} \tanh(ac+bcx)}{6bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{25e^{c(a+bx)} \tanh(ac+bcx)}{4bc(1-e^{2c(a+bx)}) \sqrt{\tanh^2(ac+bcx)}} - \frac{15\operatorname{arctanh}(e^{c(a+bx)}) \tanh(ac+bcx)}{4bc\sqrt{\tanh^2(ac+bcx)}}$$

output

```
exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-4*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^4/(tanh(b*c*x+a*c)^2)^(1/2)+26/3*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^3/(tanh(b*c*x+a*c)^2)^(1/2)-55/6*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^2/(tanh(b*c*x+a*c)^2)^(1/2)+25/4*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))/(tanh(b*c*x+a*c)^2)^(1/2)-15/4*arctanh(exp(c*(b*x+a)))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)
```

**Mathematica [A] (warning: unable to verify)**

Time = 11.48 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.51

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \frac{(66e^{c(a+bx)} - 314e^{3c(a+bx)} + 374e^{5c(a+bx)} - 246e^{7c(a+bx)} + 24e^{9c(a+bx)} + 45(-1 + e^{2c(a+bx)})) \operatorname{Log}[1 - E^{c(a+bx)}] - 45(-1 + E^{2c(a+bx)})^4 \operatorname{Log}[1 + E^{c(a+bx)}] + 45(-1 + E^{2c(a+bx)})^4 \operatorname{Sqrt}[\operatorname{Tanh}[c(a+bx)]]}{24bc(-1 + e^{2c(a+bx)})}$$

input `Integrate[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(5/2),x]`output `((66*E^(c*(a + b*x)) - 314*E^(3*c*(a + b*x)) + 374*E^(5*c*(a + b*x)) - 246*E^(7*c*(a + b*x)) + 24*E^(9*c*(a + b*x)) + 45*(-1 + E^(2*c*(a + b*x)))^4*Log[1 - E^(c*(a + b*x))] - 45*(-1 + E^(2*c*(a + b*x)))^4*Log[1 + E^(c*(a + b*x))])*Tanh[c*(a + b*x)]/(24*b*c*(-1 + E^(2*c*(a + b*x)))^4*Sqrt[Tanh[c*(a + b*x)]^2])`**Rubi [A] (verified)**Time = 1.76 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.53, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {7271, 2720, 25, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$$

$$\downarrow 7271$$

$$\frac{\tanh(ac+bcx) \int e^{c(a+bx)} \coth^5(ac+bcx) dx}{\sqrt{\tanh^2(ac+bcx)}}$$

$$\downarrow 2720$$

$$\frac{\tanh(ac+bcx) \int -\frac{(1+e^{2c(a+bx)})^5}{(1-e^{2c(a+bx)})^5} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ac+bcx)}}$$



$$\begin{aligned}
& \downarrow 25 \\
& \frac{\tanh(ac + bcx) \int \frac{(1+e^{2c(a+bx)})^5}{(1-e^{2c(a+bx)})^5} de^{c(a+bx)}}{bc\sqrt{\tanh^2(ac + bcx)}} \\
& \downarrow 300 \\
& \frac{\tanh(ac + bcx) \int \left( \frac{2(1+10e^{4c(a+bx)}+5e^{8c(a+bx)})}{(1-e^{2c(a+bx)})^5} - 1 \right) de^{c(a+bx)}}{bc\sqrt{\tanh^2(ac + bcx)}} \\
& \downarrow 2009 \\
& \frac{\left( -\frac{15}{4}\operatorname{arctanh}(e^{c(a+bx)}) + e^{c(a+bx)} + \frac{25e^{c(a+bx)}}{4(1-e^{2c(a+bx)})} - \frac{55e^{c(a+bx)}}{6(1-e^{2c(a+bx)})^2} + \frac{26e^{c(a+bx)}}{3(1-e^{2c(a+bx)})^3} - \frac{4e^{c(a+bx)}}{(1-e^{2c(a+bx)})^4} \right) \tanh(ac + bcx)}{bc\sqrt{\tanh^2(ac + bcx)}}
\end{aligned}$$

input `Int[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(5/2), x]`

output `((E^(c*(a + b*x)) - (4*E^(c*(a + b*x))))/(1 - E^(2*c*(a + b*x)))^4 + (26*E^(c*(a + b*x)))/(3*(1 - E^(2*c*(a + b*x)))^3) - (55*E^(c*(a + b*x)))/(6*(1 - E^(2*c*(a + b*x)))^2) + (25*E^(c*(a + b*x)))/(4*(1 - E^(2*c*(a + b*x)))) - (15*ArcTanh[E^(c*(a + b*x))])/4)*Tanh[a*c + b*c*x]/(b*c*Sqrt[Tanh[a*c + b*c*x]^2])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 300 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 7271

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] := Simp[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])) Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p},
x] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(Eq
Q[v, x] && EqQ[m, 1])
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.83 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.61

method	result
default	$\text{csgn}(\tanh(c(bx+a))) \left( \frac{\cosh(bx+ac)^4}{\sinh(bx+ac)^3} - \frac{4 \cosh(bx+ac)^2}{\sinh(bx+ac)^3} + \frac{8}{3 \sinh(bx+ac)^3} + \frac{\cosh(bx+ac)^5}{\sinh(bx+ac)^4} - \frac{5 \cosh(bx+ac)^3}{\sinh(bx+ac)^4} + \frac{5 \cosh(bx+ac)}{\sinh(bx+ac)^4} + 5 \left( -\frac{\text{csch}(bx+ac)}{\sinh(bx+ac)^4} \right) \right)$
risch	$\frac{(e^{2c(bx+a)} - 1)e^{c(bx+a)}}{\sqrt{\frac{(e^{2c(bx+a)} - 1)^2}{(1+e^{2c(bx+a)})^2}} (1+e^{2c(bx+a)})} bc - \frac{e^{c(bx+a)} (75 e^{6c(bx+a)} - 115 e^{4c(bx+a)} + 109 e^{2c(bx+a)} - 21)}{12 (e^{2c(bx+a)} - 1)^3 (1+e^{2c(bx+a)})} \sqrt{\frac{(e^{2c(bx+a)} - 1)^2}{(1+e^{2c(bx+a)})^2}} cb - \frac{15 (e^{2c(bx+a)} - 1) \ln(e^{c(bx+a)})}{8 \sqrt{\frac{(e^{2c(bx+a)} - 1)^2}{(1+e^{2c(bx+a)})^2}} (1+e^{2c(bx+a)})} cb$

input

```
int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2), x, method=_RETURNVERBOSE)
```

output

```
csgn(tanh(c*(b*x+a)))/c/b*(1/sinh(b*c*x+a*c)^3*cosh(b*c*x+a*c)^4-4/sinh(b*
c*x+a*c)^3*cosh(b*c*x+a*c)^2+8/3/sinh(b*c*x+a*c)^3+cosh(b*c*x+a*c)^5/sinh(
b*c*x+a*c)^4-5/sinh(b*c*x+a*c)^4*cosh(b*c*x+a*c)^3+5/sinh(b*c*x+a*c)^4*cos
h(b*c*x+a*c)+5*(-1/4*csch(b*c*x+a*c)^3+3/8*csch(b*c*x+a*c))*coth(b*c*x+a*c
)-15/4*arctanh(exp(b*c*x+a*c)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 1617 vs.  $2(281) = 562$ .

Time = 0.11 (sec) , antiderivative size = 1617, normalized size of antiderivative = 5.07

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \text{Too large to display}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="fricas")`

output

```
1/24*(24*cosh(b*c*x + a*c)^9 + 216*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^8 +
 24*sinh(b*c*x + a*c)^9 + 6*(144*cosh(b*c*x + a*c)^2 - 41)*sinh(b*c*x + a*
c)^7 - 246*cosh(b*c*x + a*c)^7 + 42*(48*cosh(b*c*x + a*c)^3 - 41*cosh(b*c*
x + a*c))*sinh(b*c*x + a*c)^6 + 2*(1512*cosh(b*c*x + a*c)^4 - 2583*cosh(b*
c*x + a*c)^2 + 187)*sinh(b*c*x + a*c)^5 + 374*cosh(b*c*x + a*c)^5 + 2*(151
2*cosh(b*c*x + a*c)^5 - 4305*cosh(b*c*x + a*c)^3 + 935*cosh(b*c*x + a*c))*
sinh(b*c*x + a*c)^4 + 2*(1008*cosh(b*c*x + a*c)^6 - 4305*cosh(b*c*x + a*c)
^4 + 1870*cosh(b*c*x + a*c)^2 - 157)*sinh(b*c*x + a*c)^3 - 314*cosh(b*c*x
+ a*c)^3 + 2*(432*cosh(b*c*x + a*c)^7 - 2583*cosh(b*c*x + a*c)^5 + 1870*co
sh(b*c*x + a*c)^3 - 471*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^2 - 45*(cosh(
b*c*x + a*c)^8 + 8*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^7 + sinh(b*c*x + a*
c)^8 + 4*(7*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^6 - 4*cosh(b*c*x +
a*c)^6 + 8*(7*cosh(b*c*x + a*c)^3 - 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)
^5 + 2*(35*cosh(b*c*x + a*c)^4 - 30*cosh(b*c*x + a*c)^2 + 3)*sinh(b*c*x +
a*c)^4 + 6*cosh(b*c*x + a*c)^4 + 8*(7*cosh(b*c*x + a*c)^5 - 10*cosh(b*c*x
+ a*c)^3 + 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^3 + 4*(7*cosh(b*c*x + a*
c)^6 - 15*cosh(b*c*x + a*c)^4 + 9*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*
c)^2 - 4*cosh(b*c*x + a*c)^2 + 8*(cosh(b*c*x + a*c)^7 - 3*cosh(b*c*x + a*c)
)^5 + 3*cosh(b*c*x + a*c)^3 - cosh(b*c*x + a*c))*sinh(b*c*x + a*c) + 1)*lo
g(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) + 1) + 45*(cosh(b*c*x + a*c)^8 ...
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \text{Timed out}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(5/2),x)`

output Timed out

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.52

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = -\frac{15 \log(e^{(bcx+ac)} + 1)}{8bc} + \frac{15 \log(e^{(bcx+ac)} - 1)}{8bc} + \frac{12e^{(9bcx+9ac)} - 123e^{(7bcx+7ac)} + 187e^{(5bcx+5ac)} - 157e^{(3bcx+3ac)} + 33e^{(bcx+ac)}}{12bc(e^{(8bcx+8ac)} - 4e^{(6bcx+6ac)} + 6e^{(4bcx+4ac)} - 4e^{(2bcx+2ac)} + 1)}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="maxima")`

output `-15/8*log(e^(b*c*x + a*c) + 1)/(b*c) + 15/8*log(e^(b*c*x + a*c) - 1)/(b*c) + 1/12*(12*e^(9*b*c*x + 9*a*c) - 123*e^(7*b*c*x + 7*a*c) + 187*e^(5*b*c*x + 5*a*c) - 157*e^(3*b*c*x + 3*a*c) + 33*e^(b*c*x + a*c))/(b*c*(e^(8*b*c*x + 8*a*c) - 4*e^(6*b*c*x + 6*a*c) + 6*e^(4*b*c*x + 4*a*c) - 4*e^(2*b*c*x + 2*a*c) + 1))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.72

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc} - \frac{15 \log(e^{(bcx+ac)} + 1) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{8bc} + \frac{15 \log(|e^{(bcx+ac)} - 1|) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{8bc} - \frac{75 e^{(7bcx+7ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 115 e^{(5bcx+5ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 109 e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 21 e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{12bc(e^{(2bcx+2ac)} - 1)^4}$$

input `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="giac")`

output `e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) - 15/8*log(e^(b*c*x + a*c) + 1)*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) + 15/8*log(abs(e^(b*c*x + a*c) - 1))*sgn(e^(2*b*c*x + 2*a*c) - 1)/(b*c) - 1/12*(75*e^(7*b*c*x + 7*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 115*e^(5*b*c*x + 5*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 109*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 21*e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(b*c*(e^(2*b*c*x + 2*a*c) - 1)^4)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \int \frac{e^{c(a+bx)}}{(\tanh(ac+bcx))^2)^{5/2}} dx$$

input `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(5/2),x)`

output `int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(5/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 830, normalized size of antiderivative = 2.60

$$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx = \frac{-45e^{6bcx+6ac}\log(e^{bcx+2ac} + e^{ac})\tanh(bcx+ac)^4 + 45e^{6bcx+6ac}\log(e^{bcx+2ac} - e^{ac})\tanh(bcx+ac)^4}{(24\tanh(ac+bcx)^4bc(e^{6ac+6bcx} + e^{6ac}))}$$

input

```
int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x)
```

output

```
(14***e**(7*a*c + 7*b*c*x)*tanh(a*c + b*c*x)**4 + 16***e**(7*a*c + 7*b*c*x)*tanh(a*c + b*c*x)**3 - 8***e**(7*a*c + 7*b*c*x)*tanh(a*c + b*c*x)**2 + 8***e**(7*a*c + 7*b*c*x)*tanh(a*c + b*c*x) - 6***e**(7*a*c + 7*b*c*x) - 45***e**(6*a*c + 6*b*c*x)*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**4 + 45***e**(6*a*c + 6*b*c*x)*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**4 - 192***e**(5*a*c + 5*b*c*x)*tanh(a*c + b*c*x)**4 - 48***e**(5*a*c + 5*b*c*x)*tanh(a*c + b*c*x)**3 + 24***e**(5*a*c + 5*b*c*x)*tanh(a*c + b*c*x)**2 - 24***e**(5*a*c + 5*b*c*x)*tanh(a*c + b*c*x) + 18***e**(5*a*c + 5*b*c*x) + 135***e**(4*a*c + 4*b*c*x)*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**4 - 135***e**(4*a*c + 4*b*c*x)*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**4 + 202***e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x)**4 + 48***e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x)**3 - 24***e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x)**2 + 24***e**(3*a*c + 3*b*c*x)*tanh(a*c + b*c*x) - 18***e**(3*a*c + 3*b*c*x) - 135***e**(2*a*c + 2*b*c*x)*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**4 + 135***e**(2*a*c + 2*b*c*x)*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**4 - 104***e**(a*c + b*c*x)*tanh(a*c + b*c*x)**4 - 16***e**(a*c + b*c*x)*tanh(a*c + b*c*x)**3 + 8***e**(a*c + b*c*x)*tanh(a*c + b*c*x)**2 - 8***e**(a*c + b*c*x)*tanh(a*c + b*c*x) + 6***e**(a*c + b*c*x) + 45*log(e**(2*a*c + b*c*x) + e**(a*c))*tanh(a*c + b*c*x)**4 - 45*log(e**(2*a*c + b*c*x) - e**(a*c))*tanh(a*c + b*c*x)**4)/(24*tanh(a*c + b*c*x)**4*b*c*(e**(6*a*c + 6*b*c*x) + e**(6*a*c)))
```

### 3.251 $\int \sin^3(\tanh(a + bx)) dx$

Optimal result	1838
Mathematica [A] (verified)	1839
Rubi [A] (verified)	1839
Maple [A] (verified)	1840
Fricas [C] (verification not implemented)	1841
Sympy [F]	1842
Maxima [F]	1843
Giac [F]	1843
Mupad [F(-1)]	1843
Reduce [F]	1844

#### Optimal result

Integrand size = 9, antiderivative size = 157

$$\int \sin^3(\tanh(a + bx)) dx = -\frac{3 \operatorname{CosIntegral}(1 - \tanh(a + bx)) \sin(1)}{8b} - \frac{3 \operatorname{CosIntegral}(1 + \tanh(a + bx)) \sin(1)}{8b} + \frac{\operatorname{CosIntegral}(3 - 3 \tanh(a + bx)) \sin(3)}{8b} + \frac{\operatorname{CosIntegral}(3 + 3 \tanh(a + bx)) \sin(3)}{8b} - \frac{\cos(3) \operatorname{Si}(3 - 3 \tanh(a + bx))}{8b} + \frac{3 \cos(1) \operatorname{Si}(1 - \tanh(a + bx))}{8b} + \frac{3 \cos(1) \operatorname{Si}(1 + \tanh(a + bx))}{8b} - \frac{\cos(3) \operatorname{Si}(3 + 3 \tanh(a + bx))}{8b}$$

output

```
-3/8*Ci(1-tanh(b*x+a))*sin(1)/b-3/8*Ci(1+tanh(b*x+a))*sin(1)/b+1/8*Ci(3-3*
tanh(b*x+a))*sin(3)/b+1/8*Ci(3+3*tanh(b*x+a))*sin(3)/b+1/8*cos(3)*Si(-3+3*
tanh(b*x+a))/b-3/8*cos(1)*Si(-1+tanh(b*x+a))/b+3/8*cos(1)*Si(1+tanh(b*x+a)
)/b-1/8*cos(3)*Si(3+3*tanh(b*x+a))/b
```

**Mathematica [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.79

$$\int \sin^3(\tanh(a + bx)) dx$$


---


$$= \frac{-6 \operatorname{CosIntegral}(1 - \tanh(a + bx)) \sin(1) - 6 \operatorname{CosIntegral}(1 + \tanh(a + bx)) \sin(1) + 2 \operatorname{CosIntegral}(3 -$$

input `Integrate[Sin[Tanh[a + b*x]]^3,x]`

output `(-6*CosIntegral[1 - Tanh[a + b*x]]*Sin[1] - 6*CosIntegral[1 + Tanh[a + b*x]]*Sin[1] + 2*CosIntegral[3 - 3*Tanh[a + b*x]]*Sin[3] + 2*CosIntegral[3 + 3*Tanh[a + b*x]]*Sin[3] - 2*Cos[3]*SinIntegral[3 - 3*Tanh[a + b*x]] + 6*Cos[1]*SinIntegral[1 - Tanh[a + b*x]] + 6*Cos[1]*SinIntegral[1 + Tanh[a + b*x]] - 2*Cos[3]*SinIntegral[3 + 3*Tanh[a + b*x]])/(16*b)`

**Rubi [A] (verified)**

Time = 1.00 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.87, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {4853, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin^3(\tanh(a + bx)) dx$$

$$\downarrow \text{4853}$$

$$\frac{\int \frac{\sin^3(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b}$$

$$\downarrow \text{7276}$$

$$\frac{\int \left( \frac{\sin^3(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\sin^3(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)}{b}$$

$$\downarrow \text{2009}$$



$$\frac{1}{8} \sin(3) \operatorname{CosIntegral}(3 - 3 \tanh(a + bx)) + \frac{1}{8} \sin(3) \operatorname{CosIntegral}(3 \tanh(a + bx) + 3) - \frac{3}{8} \sin(1) \operatorname{CosIntegral}(1 - \tanh(a + bx))$$

input `Int[Sin[Tanh[a + b*x]]^3,x]`

output `((-3*CosIntegral[1 - Tanh[a + b*x]]*Sin[1])/8 - (3*CosIntegral[1 + Tanh[a + b*x]]*Sin[1])/8 + (CosIntegral[3 - 3*Tanh[a + b*x]]*Sin[3])/8 + (CosIntegral[3 + 3*Tanh[a + b*x]]*Sin[3])/8 - (Cos[3]*SinIntegral[3 - 3*Tanh[a + b*x]])/8 + (3*Cos[1]*SinIntegral[1 - Tanh[a + b*x]])/8 + (3*Cos[1]*SinIntegral[1 + Tanh[a + b*x]])/8 - (Cos[3]*SinIntegral[3 + 3*Tanh[a + b*x]])/8)/b`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4853 `Int[u_, x_Symbol] :=> With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x]]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :=> With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

### Maple [A] (verified)

Time = 1.78 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{\frac{\operatorname{Si}(-3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(-3+3 \tanh(bx+a)) \sin(3) - \frac{\operatorname{Si}(3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(3+3 \tanh(bx+a)) \sin(3) - 3 \operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1) + 3 \operatorname{Si}(1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{8} + \frac{\operatorname{Si}(-3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(-3+3 \tanh(bx+a)) \sin(3) - \frac{\operatorname{Si}(3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(3+3 \tanh(bx+a)) \sin(3) - 3 \operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1) + 3 \operatorname{Si}(1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{8}}{b}}$
default	$\frac{\frac{\operatorname{Si}(-3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(-3+3 \tanh(bx+a)) \sin(3) - \frac{\operatorname{Si}(3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(3+3 \tanh(bx+a)) \sin(3) - 3 \operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1) + 3 \operatorname{Si}(1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{8} + \frac{\operatorname{Si}(-3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(-3+3 \tanh(bx+a)) \sin(3) - \frac{\operatorname{Si}(3+3 \tanh(bx+a)) \cos(3) + \operatorname{Ci}(3+3 \tanh(bx+a)) \sin(3) - 3 \operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1) + 3 \operatorname{Si}(1+\tanh(bx+a)) \cos(1) - 3 \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{8}}{b}}$
risch	$-\frac{ie^{-3i} \operatorname{expIntegral}_1\left(-\frac{6i}{1+e^{2bx+2a}}\right)}{16b} + \frac{ie^{3i} \operatorname{expIntegral}_1\left(-\frac{6i}{1+e^{2bx+2a}}+6i\right)}{16b} - \frac{3ie^i \operatorname{expIntegral}_1\left(\frac{2i}{1+e^{2bx+2a}}\right)}{16b} + \dots$

input `int(sin(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `1/b*(1/8*Si(-3+3*tanh(b*x+a))*cos(3)+1/8*Ci(-3+3*tanh(b*x+a))*sin(3)-1/8*Si(3+3*tanh(b*x+a))*cos(3)+1/8*Ci(3+3*tanh(b*x+a))*sin(3)-3/8*Si(-1+tanh(b*x+a))*cos(1)-3/8*Ci(-1+tanh(b*x+a))*sin(1)+3/8*Si(1+tanh(b*x+a))*cos(1)-3/8*Ci(1+tanh(b*x+a))*sin(1))`

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 697, normalized size of antiderivative = 4.44

$$\int \sin^3(\tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(sin(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```

1/16*((-I*cos(3)^2*cos(1) - (-I*cos(1) + sin(1))*sin(3)^2 - 2*I*(I*cos(3)*
cos(1) - cos(3)*sin(1))*sin(3) + I*(-I*cos(3)^2 + I)*sin(1) + I*cos(1))*co
s_integral(3*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) - 3*(2*cos(3)*
cos(1)*sin(1) + I*cos(3)*sin(1)^2 + (-I*cos(1)^2 + I)*cos(3) + I*(-I*cos(1)
)^2 + 2*cos(1)*sin(1) + I*sin(1)^2 + I)*sin(3))*cos_integral((cosh(b*x + a)
+ sinh(b*x + a))/cosh(b*x + a)) + (-I*cos(3)^2*cos(1) - (-I*cos(1) + sin
(1))*sin(3)^2 - 2*I*(I*cos(3)*cos(1) - cos(3)*sin(1))*sin(3) + I*(-I*cos(3)
)^2 + I)*sin(1) + I*cos(1))*cos_integral(6/(cosh(b*x + a)^2 + 2*cosh(b*x +
a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) - 3*(2*cos(3)*cos(1)*sin(1) + I*
cos(3)*sin(1)^2 + (-I*cos(1)^2 + I)*cos(3) + I*(-I*cos(1)^2 + 2*cos(1)*sin
(1) + I*sin(1)^2 + I)*sin(3))*cos_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x
+ a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) - (cos(3)^2*cos(1) - (cos(1) +
I*sin(1))*sin(3)^2 + 2*I*(cos(3)*cos(1) + I*cos(3)*sin(1))*sin(3) + I*(co
s(3)^2 + 1)*sin(1) + cos(1))*sin_integral(3*(cosh(b*x + a) + sinh(b*x + a)
)/cosh(b*x + a)) - 3*(-2*I*cos(3)*cos(1)*sin(1) + cos(3)*sin(1)^2 - (cos(1)
)^2 + 1)*cos(3) - I*(cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*sin(3))*
sin_integral((cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) - (cos(3)^2*co
s(1) - (cos(1) + I*sin(1))*sin(3)^2 + 2*I*(cos(3)*cos(1) + I*cos(3)*sin(1)
)*sin(3) + I*(cos(3)^2 + 1)*sin(1) + cos(1))*sin_integral(6/(cosh(b*x + a)
^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) - 3*(-2*I*co...

```

## Sympy [F]

$$\int \sin^3(\tanh(a + bx)) dx = \int \sin^3(\tanh(a + bx)) dx$$

input

```
integrate(sin(tanh(b*x+a))**3,x)
```

output

```
Integral(sin(tanh(a + b*x))**3, x)
```

**Maxima [F]**

$$\int \sin^3(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^3 dx$$

input `integrate(sin(tanh(b*x+a))^3,x, algorithm="maxima")`

output `integrate(sin(tanh(b*x + a))^3, x)`

**Giac [F]**

$$\int \sin^3(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^3 dx$$

input `integrate(sin(tanh(b*x+a))^3,x, algorithm="giac")`

output `integrate(sin(tanh(b*x + a))^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sin^3(\tanh(a + bx)) dx = \int \sin(\tanh(a + bx))^3 dx$$

input `int(sin(tanh(a + b*x))^3,x)`

output `int(sin(tanh(a + b*x))^3, x)`

**Reduce [F]**

$$\int \sin^3(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^3 dx$$

input `int(sin(tanh(b*x+a))^3,x)`

output `int(sin(tanh(a + b*x))**3,x)`

### 3.252 $\int \sin^2(\tanh(a + bx)) dx$

Optimal result	1845
Mathematica [A] (verified)	1846
Rubi [A] (verified)	1846
Maple [A] (verified)	1848
Fricas [C] (verification not implemented)	1848
Sympy [F]	1849
Maxima [F]	1849
Giac [F]	1850
Mupad [F(-1)]	1850
Reduce [F]	1850

#### Optimal result

Integrand size = 9, antiderivative size = 115

$$\int \sin^2(\tanh(a + bx)) dx = \frac{\cos(2) \operatorname{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} - \frac{\cos(2) \operatorname{CosIntegral}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} + \frac{\sin(2) \operatorname{Si}(2 - 2 \tanh(a + bx))}{4b} - \frac{\sin(2) \operatorname{Si}(2 + 2 \tanh(a + bx))}{4b}$$

output

```
1/4*cos(2)*Ci(2-2*tanh(b*x+a))/b-1/4*cos(2)*Ci(2+2*tanh(b*x+a))/b-1/4*ln(1-tanh(b*x+a))/b+1/4*ln(1+tanh(b*x+a))/b-1/4*sin(2)*Si(-2+2*tanh(b*x+a))/b-1/4*sin(2)*Si(2+2*tanh(b*x+a))/b
```

**Mathematica [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

$$\int \sin^2(\tanh(a + bx)) dx$$

$$= \frac{\cos(2) \operatorname{CosIntegral}(2 - 2 \tanh(a + bx)) - \cos(2) \operatorname{CosIntegral}(2(1 + \tanh(a + bx))) - \log(1 - \tanh(a + bx)) + \log(1 + \tanh(a + bx)) + \operatorname{SinIntegral}(2 - 2 \tanh(a + bx)) - \operatorname{SinIntegral}(2(1 + \tanh(a + bx)))}{4b}$$

input `Integrate[Sin[Tanh[a + b*x]]^2,x]`

output `(Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]] - Cos[2]*CosIntegral[2*(1 + Tanh[a + b*x])] - Log[1 - Tanh[a + b*x]] + Log[1 + Tanh[a + b*x]] + Sin[2]*SinIntegral[2 - 2*Tanh[a + b*x]] - Sin[2]*SinIntegral[2*(1 + Tanh[a + b*x])])/(4*b)`

**Rubi [A] (verified)**

Time = 0.65 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {4853, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin^2(\tanh(a + bx)) dx$$

$$\downarrow 4853$$

$$\int \frac{\sin^2(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)$$

$$\downarrow 7276$$

$$\int \left( \frac{\sin^2(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\sin^2(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)$$

$$\downarrow 2009$$





**Maple [A] (verified)**

Time = 0.86 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{4} + \frac{\ln(1+\tanh(bx+a))}{4} - \frac{\text{Si}(2+2\tanh(bx+a))\sin(2)}{4} - \frac{\text{Ci}(2+2\tanh(bx+a))\cos(2)}{4} - \frac{\text{Si}(-2+2\tanh(bx+a))\sin(2)}{4}}{b}$
default	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{4} + \frac{\ln(1+\tanh(bx+a))}{4} - \frac{\text{Si}(2+2\tanh(bx+a))\sin(2)}{4} - \frac{\text{Ci}(2+2\tanh(bx+a))\cos(2)}{4} - \frac{\text{Si}(-2+2\tanh(bx+a))\sin(2)}{4}}{b}$
risch	$-\frac{e^{-2i} \exp\text{Integral}_1\left(-\frac{4i}{1+e^{2bx+2a}}\right)}{8b} + \frac{e^{-2i} \exp\text{Integral}_1\left(\frac{4i}{1+e^{2bx+2a}}-4i\right)}{8b} + \frac{e^{2i} \exp\text{Integral}_1\left(-\frac{4i}{1+e^{2bx+2a}}+4i\right)}{8b}$

input `int(sin(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`output `1/b*(-1/4*ln(-1+tanh(b*x+a))+1/4*ln(1+tanh(b*x+a))-1/4*Si(2+2*tanh(b*x+a))*sin(2)-1/4*Ci(2+2*tanh(b*x+a))*cos(2)-1/4*Si(-2+2*tanh(b*x+a))*sin(2)+1/4*Ci(-2+2*tanh(b*x+a))*cos(2))`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.00

$$\int \sin^2(\tanh(a + bx)) dx$$

$$= \frac{4bx \cos(2) + 4i bx \sin(2) - (\cos(2)^2 + 2i \cos(2) \sin(2) - \sin(2)^2 + 1) \text{Ci}\left(\frac{2(\cosh(bx+a) + \sinh(bx+a))}{\cosh(bx+a)}\right) + (c$$

input `integrate(sin(tanh(b*x+a))^2,x, algorithm="fricas")`

output

```
1/8*(4*b*x*cos(2) + 4*I*b*x*sin(2) - (cos(2)^2 + 2*I*cos(2)*sin(2) - sin(2)^2 + 1)*cos_integral(2*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (cos(2)^2 + 2*I*cos(2)*sin(2) - sin(2)^2 + 1)*cos_integral(4/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) + (I*cos(2)^2 - 2*cos(2)*sin(2) - I*sin(2)^2 - I)*sin_integral(2*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (-I*cos(2)^2 + 2*cos(2)*sin(2) + I*sin(2)^2 + I)*sin_integral(4/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)))/(b*cos(2) + I*b*sin(2))
```

**Sympy [F]**

$$\int \sin^2(\tanh(a + bx)) dx = \int \sin^2(\tanh(a + bx)) dx$$

input

```
integrate(sin(tanh(b*x+a))**2,x)
```

output

```
Integral(sin(tanh(a + b*x))**2, x)
```

**Maxima [F]**

$$\int \sin^2(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^2 dx$$

input

```
integrate(sin(tanh(b*x+a))^2,x, algorithm="maxima")
```

output

```
1/2*x - 1/2*integrate(cos(2*(e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)), x)
```

**Giac [F]**

$$\int \sin^2(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^2 dx$$

input `integrate(sin(tanh(b*x+a))^2,x, algorithm="giac")`

output `integrate(sin(tanh(b*x + a))^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sin^2(\tanh(a + bx)) dx = \int \sin(\tanh(a + bx))^2 dx$$

input `int(sin(tanh(a + b*x))^2,x)`

output `int(sin(tanh(a + b*x))^2, x)`

**Reduce [F]**

$$\int \sin^2(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a))^2 dx$$

input `int(sin(tanh(b*x+a))^2,x)`

output `int(sin(tanh(a + b*x))**2,x)`

### 3.253 $\int \sin(\tanh(a + bx)) dx$

Optimal result	1851
Mathematica [A] (verified)	1851
Rubi [A] (verified)	1852
Maple [A] (verified)	1853
Fricas [C] (verification not implemented)	1854
Sympy [F]	1854
Maxima [F]	1855
Giac [F]	1855
Mupad [F(-1)]	1855
Reduce [F]	1856

#### Optimal result

Integrand size = 7, antiderivative size = 77

$$\int \sin(\tanh(a + bx)) dx = -\frac{\text{CosIntegral}(1 - \tanh(a + bx)) \sin(1)}{2b} - \frac{\text{CosIntegral}(1 + \tanh(a + bx)) \sin(1)}{2b} + \frac{\cos(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{Si}(1 + \tanh(a + bx))}{2b}$$

output

$$-1/2*\text{Ci}(1-\tanh(b*x+a))*\sin(1)/b-1/2*\text{Ci}(1+\tanh(b*x+a))*\sin(1)/b-1/2*\cos(1)*\text{Si}(-1+\tanh(b*x+a))/b+1/2*\cos(1)*\text{Si}(1+\tanh(b*x+a))/b$$

#### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.77

$$\int \sin(\tanh(a + bx)) dx = \frac{\text{CosIntegral}(1 - \tanh(a + bx)) \sin(1) + \text{CosIntegral}(1 + \tanh(a + bx)) \sin(1) - \cos(1)(\text{Si}(1 - \tanh(a + bx)) + \text{Si}(1 + \tanh(a + bx)))}{2b}$$

input

`Integrate[Sin[Tanh[a + b*x]],x]`

output

```
-1/2*(CosIntegral[1 - Tanh[a + b*x]]*Sin[1] + CosIntegral[1 + Tanh[a + b*x]]*Sin[1] - Cos[1]*(SinIntegral[1 - Tanh[a + b*x]] + SinIntegral[1 + Tanh[a + b*x]]))/b
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.90, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {4853, 3814, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(\tanh(a + bx)) dx$$

$$\downarrow 4853$$

$$\frac{\int \frac{\sin(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b}$$

$$\downarrow 3814$$

$$\frac{\int \left( \frac{\sin(\tanh(a+bx))}{2(1-\tanh(a+bx))} + \frac{\sin(\tanh(a+bx))}{2(\tanh(a+bx)+1)} \right) d \tanh(a + bx)}{b}$$

$$\downarrow 2009$$

$$\frac{-\frac{1}{2} \sin(1) \text{CosIntegral}(1 - \tanh(a + bx)) - \frac{1}{2} \sin(1) \text{CosIntegral}(\tanh(a + bx) + 1) + \frac{1}{2} \cos(1) \text{Si}(1 - \tanh(a + bx))}{b}$$

input

```
Int[Sin[Tanh[a + b*x]],x]
```

output

```
(-1/2*(CosIntegral[1 - Tanh[a + b*x]]*Sin[1]) - (CosIntegral[1 + Tanh[a + b*x]]*Sin[1])/2 + (Cos[1]*SinIntegral[1 - Tanh[a + b*x]])/2 + (Cos[1]*SinIntegral[1 + Tanh[a + b*x]])/2)/b
```

## Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3814 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*Sin[(c_.) + (d_.)*(x_)], x_Symbol] := Int[ExpandIntegrand[Sin[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])`

rule 4853 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x]]`

## Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{\frac{\operatorname{Si}(1+\tanh(bx+a)) \cos(1) - \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{2} - \frac{\operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1)}{2}}{b}$
default	$\frac{\frac{\operatorname{Si}(1+\tanh(bx+a)) \cos(1) - \operatorname{Ci}(1+\tanh(bx+a)) \sin(1)}{2} - \frac{\operatorname{Si}(-1+\tanh(bx+a)) \cos(1) - \operatorname{Ci}(-1+\tanh(bx+a)) \sin(1)}{2}}{b}$
risch	$\frac{ie^{-i} \operatorname{expIntegral}_1\left(-\frac{2i}{1+e^{2bx+2a}}\right)}{4b} + \frac{ie^{-i} \operatorname{expIntegral}_1\left(\frac{2i}{1+e^{2bx+2a}} - 2i\right)}{4b} - \frac{ie^i \operatorname{expIntegral}_1\left(-\frac{2i}{1+e^{2bx+2a}} + 2i\right)}{4b} -$

input `int(sin(tanh(b*x+a)), x, method=_RETURNVERBOSE)`

output `1/b*(1/2*Si(1+tanh(b*x+a))*cos(1)-1/2*Ci(1+tanh(b*x+a))*sin(1)-1/2*Si(-1+tanh(b*x+a))*cos(1)-1/2*Ci(-1+tanh(b*x+a))*sin(1))`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.79

$$\int \sin(\tanh(a + bx)) dx$$

$$= \frac{(i \cos(1)^2 - 2 \cos(1) \sin(1) - i \sin(1)^2 - i) \operatorname{Ci}\left(\frac{\cosh(bx+a)+\sinh(bx+a)}{\cosh(bx+a)}\right) + (i \cos(1)^2 - 2 \cos(1) \sin(1) - i \sin(1)^2 - i) \operatorname{Si}\left(\frac{\cosh(bx+a)+\sinh(bx+a)}{\cosh(bx+a)}\right) + (i \cos(1)^2 - 2 \cos(1) \sin(1) - i \sin(1)^2 - i) \operatorname{Ci}\left(\frac{\cosh(bx+a)-\sinh(bx+a)}{\cosh(bx+a)}\right) + (i \cos(1)^2 - 2 \cos(1) \sin(1) - i \sin(1)^2 - i) \operatorname{Si}\left(\frac{\cosh(bx+a)-\sinh(bx+a)}{\cosh(bx+a)}\right)}{b}$$

input `integrate(sin(tanh(b*x+a)),x, algorithm="fricas")`

output `1/4*((I*cos(1)^2 - 2*cos(1)*sin(1) - I*sin(1)^2 - I)*cos_integral((cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (I*cos(1)^2 - 2*cos(1)*sin(1) - I*sin(1)^2 - I)*cos_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) + (cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*sin_integral((cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*sin_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)))/(b*cos(1) + I*b*sin(1))`

**Sympy [F]**

$$\int \sin(\tanh(a + bx)) dx = \int \sin(\tanh(a + bx)) dx$$

input `integrate(sin(tanh(b*x+a)),x)`

output `Integral(sin(tanh(a + b*x)), x)`

**Maxima [F]**

$$\int \sin(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a)) dx$$

input `integrate(sin(tanh(b*x+a)),x, algorithm="maxima")`

output `integrate(sin(tanh(b*x + a)), x)`

**Giac [F]**

$$\int \sin(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a)) dx$$

input `integrate(sin(tanh(b*x+a)),x, algorithm="giac")`

output `integrate(sin(tanh(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sin(\tanh(a + bx)) dx = \int \sin(\tanh(a + bx)) dx$$

input `int(sin(tanh(a + b*x)),x)`

output `int(sin(tanh(a + b*x)), x)`



**Reduce [F]**

$$\int \sin(\tanh(a + bx)) dx = \int \sin(\tanh(bx + a)) dx$$

input `int(sin(tanh(b*x+a)),x)`

output `int(sin(tanh(a + b*x)),x)`

### 3.254 $\int \csc(\tanh(a + bx)) dx$

Optimal result	1857
Mathematica [N/A]	1857
Rubi [N/A]	1858
Maple [F(-1)]	1859
Fricas [N/A]	1859
Sympy [N/A]	1859
Maxima [N/A]	1860
Giac [N/A]	1860
Mupad [N/A]	1860
Reduce [N/A]	1861

#### Optimal result

Integrand size = 7, antiderivative size = 7

$$\int \csc(\tanh(a + bx)) dx = -\frac{1}{2} \operatorname{Int}\left(\frac{\csc(\tanh(a + bx)) \operatorname{sech}^2(a + bx)}{1 - \tanh(a + bx)}, x\right) + \frac{1}{2} \operatorname{Int}\left(\frac{\csc(\tanh(a + bx)) \operatorname{sech}^2(a + bx)}{1 + \tanh(a + bx)}, x\right)$$

output `-1/2*Defer(Int)(-csc(tanh(b*x+a))*sech(b*x+a)^2/(1-tanh(b*x+a)),x)+1/2*Defer(Int)(csc(tanh(b*x+a))*sech(b*x+a)^2/(1+tanh(b*x+a)),x)`

#### Mathematica [N/A]

Not integrable

Time = 1.91 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(a + bx)) dx$$

input `Integrate[Csc[Tanh[a + b*x]],x]`

output `Integrate[Csc[Tanh[a + b*x]], x]`

**Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \csc(\tanh(a + bx)) dx \\
 \downarrow 4853 \\
 \frac{\int \frac{\csc(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b} \\
 \downarrow 7276 \\
 \frac{\int \left( \frac{\csc(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\csc(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)}{b} \\
 \downarrow 2009 \\
 \frac{\frac{1}{2} \int \frac{\csc(\tanh(a+bx))}{\tanh(a+bx)+1} d \tanh(a + bx) - \frac{1}{2} \int \frac{\csc(\tanh(a+bx))}{\tanh(a+bx)-1} d \tanh(a + bx)}{b}
 \end{array}$$

input `Int [Csc [Tanh [a + b*x] ] , x]`output `$Aborted`

**Maple [F(-1)]**

Timed out.

$$\int \csc(\tanh(bx + a)) dx$$

input `int(csc(tanh(b*x+a)),x)`output `int(csc(tanh(b*x+a)),x)`**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(bx + a)) dx$$

input `integrate(csc(tanh(b*x+a)),x, algorithm="fricas")`output `integral(csc(tanh(b*x + a)), x)`**Sympy [N/A]**

Not integrable

Time = 10.14 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(a + bx)) dx$$

input `integrate(csc(tanh(b*x+a)),x)`output `Integral(csc(tanh(a + b*x)), x)`

**Maxima [N/A]**

Not integrable

Time = 0.47 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(bx + a)) dx$$

input `integrate(csc(tanh(b*x+a)),x, algorithm="maxima")`output `integrate(csc(tanh(b*x + a)), x)`**Giac [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(bx + a)) dx$$

input `integrate(csc(tanh(b*x+a)),x, algorithm="giac")`output `integrate(csc(tanh(b*x + a)), x)`**Mupad [N/A]**

Not integrable

Time = 3.20 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.57

$$\int \csc(\tanh(a + bx)) dx = \int \frac{1}{\sin(\tanh(a + bx))} dx$$

input `int(1/sin(tanh(a + b*x)),x)`

output `int(1/sin(tanh(a + b*x)), x)`

**Reduce [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \csc(\tanh(a + bx)) dx = \int \csc(\tanh(bx + a)) dx$$

input `int(csc(tanh(b*x+a)), x)`

output `int(csc(tanh(a + b*x)), x)`

### 3.255 $\int \cos^3(\tanh(a + bx)) dx$

Optimal result	1862
Mathematica [A] (verified)	1863
Rubi [A] (verified)	1863
Maple [A] (verified)	1865
Fricas [C] (verification not implemented)	1865
Sympy [F]	1866
Maxima [F]	1867
Giac [F]	1867
Mupad [F(-1)]	1867
Reduce [F]	1868

#### Optimal result

Integrand size = 9, antiderivative size = 157

$$\int \cos^3(\tanh(a + bx)) dx = -\frac{\cos(3) \operatorname{CosIntegral}(3 - 3 \tanh(a + bx))}{8b} - \frac{3 \cos(1) \operatorname{CosIntegral}(1 - \tanh(a + bx))}{8b} + \frac{3 \cos(1) \operatorname{CosIntegral}(1 + \tanh(a + bx))}{8b} + \frac{\cos(3) \operatorname{CosIntegral}(3 + 3 \tanh(a + bx))}{8b} - \frac{\sin(3) \operatorname{Si}(3 - 3 \tanh(a + bx))}{8b} - \frac{3 \sin(1) \operatorname{Si}(1 - \tanh(a + bx))}{8b} + \frac{3 \sin(1) \operatorname{Si}(1 + \tanh(a + bx))}{8b} + \frac{\sin(3) \operatorname{Si}(3 + 3 \tanh(a + bx))}{8b}$$

output

```
-1/8*cos(3)*Ci(3-3*tanh(b*x+a))/b-3/8*cos(1)*Ci(1-tanh(b*x+a))/b+3/8*cos(1)*Ci(1+tanh(b*x+a))/b+1/8*cos(3)*Ci(3+3*tanh(b*x+a))/b+1/8*sin(3)*Si(-3+3*tanh(b*x+a))/b+3/8*sin(1)*Si(-1+tanh(b*x+a))/b+3/8*sin(1)*Si(1+tanh(b*x+a))/b+1/8*sin(3)*Si(3+3*tanh(b*x+a))/b
```

**Mathematica [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.79

$$\int \cos^3(\tanh(a + bx)) dx$$

$$= \frac{-2 \cos(3) \operatorname{CosIntegral}(3 - 3 \tanh(a + bx)) - 6 \cos(1) \operatorname{CosIntegral}(1 - \tanh(a + bx)) + 6 \cos(1) \operatorname{CosIntegral}(1 + \tanh(a + bx)) + 2 \cos(3) \operatorname{CosIntegral}(3 + 3 \tanh(a + bx)) - 2 \sin(3) \operatorname{SinIntegral}(3 - 3 \tanh(a + bx)) - 6 \sin(1) \operatorname{SinIntegral}(1 - \tanh(a + bx)) + 6 \sin(1) \operatorname{SinIntegral}(1 + \tanh(a + bx)) + 2 \sin(3) \operatorname{SinIntegral}(3 + 3 \tanh(a + bx))}{16b}$$

input `Integrate[Cos[Tanh[a + b*x]]^3,x]`output `(-2*Cos[3]*CosIntegral[3 - 3*Tanh[a + b*x]] - 6*Cos[1]*CosIntegral[1 - Tanh[a + b*x]] + 6*Cos[1]*CosIntegral[1 + Tanh[a + b*x]] + 2*Cos[3]*CosIntegral[3 + 3*Tanh[a + b*x]] - 2*Sin[3]*SinIntegral[3 - 3*Tanh[a + b*x]] - 6*Sin[1]*SinIntegral[1 - Tanh[a + b*x]] + 6*Sin[1]*SinIntegral[1 + Tanh[a + b*x]] + 2*Sin[3]*SinIntegral[3 + 3*Tanh[a + b*x]])/(16*b)`**Rubi [A] (verified)**Time = 0.61 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.87, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {4853, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(\tanh(a + bx)) dx$$

$$\downarrow 4853$$

$$\frac{\int \frac{\cos^3(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b}$$

$$\downarrow 7276$$

$$\frac{\int \left( \frac{\cos^3(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\cos^3(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)}{b}$$

$$\downarrow 2009$$



$$-\frac{1}{8} \cos(3) \operatorname{CosIntegral}(3 - 3 \tanh(a + bx)) - \frac{3}{8} \cos(1) \operatorname{CosIntegral}(1 - \tanh(a + bx)) + \frac{3}{8} \cos(1) \operatorname{CosIntegral}(\tanh(a + bx))$$

input `Int[Cos[Tanh[a + b*x]]^3,x]`

output `(-1/8*(Cos[3]*CosIntegral[3 - 3*Tanh[a + b*x]]) - (3*Cos[1]*CosIntegral[1 - Tanh[a + b*x]])/8 + (3*Cos[1]*CosIntegral[1 + Tanh[a + b*x]])/8 + (Cos[3]*CosIntegral[3 + 3*Tanh[a + b*x]])/8 - (Sin[3]*SinIntegral[3 - 3*Tanh[a + b*x]])/8 - (3*Ssin[1]*SinIntegral[1 - Tanh[a + b*x]])/8 + (3*Ssin[1]*SinIntegral[1 + Tanh[a + b*x]])/8 + (Sin[3]*SinIntegral[3 + 3*Tanh[a + b*x]])/8)/b`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4853 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x]]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

**Maple [A] (verified)**

Time = 1.72 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{\frac{\text{Si}(-3+3 \tanh(bx+a)) \sin(3) - \text{Ci}(-3+3 \tanh(bx+a)) \cos(3) + \frac{\text{Si}(3+3 \tanh(bx+a)) \sin(3) + \text{Ci}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{3 \text{Si}(-1+\tanh(bx+a)) \sin(1) - 3 \text{Ci}(-1+\tanh(bx+a)) \cos(1) + \frac{\text{Si}(1+\tanh(bx+a)) \sin(1) + \text{Ci}(1+\tanh(bx+a)) \cos(1)}{8}}{b}}{b}$
default	$\frac{\frac{\text{Si}(-3+3 \tanh(bx+a)) \sin(3) - \text{Ci}(-3+3 \tanh(bx+a)) \cos(3) + \frac{\text{Si}(3+3 \tanh(bx+a)) \sin(3) + \text{Ci}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{3 \text{Si}(-1+\tanh(bx+a)) \sin(1) - 3 \text{Ci}(-1+\tanh(bx+a)) \cos(1) + \frac{\text{Si}(1+\tanh(bx+a)) \sin(1) + \text{Ci}(1+\tanh(bx+a)) \cos(1)}{8}}{b}}{b}$
risch	$\frac{e^{3i} \exp\text{Integral}_1\left(\frac{6i}{1+e^{2bx+2a}}\right)}{16b} - \frac{e^{-3i} \exp\text{Integral}_1\left(\frac{6i}{1+e^{2bx+2a}} - 6i\right)}{16b} + \frac{3 e^i \exp\text{Integral}_1\left(\frac{2i}{1+e^{2bx+2a}}\right)}{16b} - \frac{3 e^{-i} \exp\text{Integral}_1\left(\frac{2i}{1+e^{2bx+2a}} - 6i\right)}{16b}$

input `int(cos(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`output `1/b*(1/8*Si(-3+3*tanh(b*x+a))*sin(3)-1/8*Ci(-3+3*tanh(b*x+a))*cos(3)+1/8*Si(3+3*tanh(b*x+a))*sin(3)+1/8*Ci(3+3*tanh(b*x+a))*cos(3)+3/8*Si(-1+tanh(b*x+a))*sin(1)-3/8*Ci(-1+tanh(b*x+a))*cos(1)+3/8*Si(1+tanh(b*x+a))*sin(1)+3/8*Ci(1+tanh(b*x+a))*cos(1))`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 698, normalized size of antiderivative = 4.45

$$\int \cos^3(\tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(cos(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```

1/16*((cos(3)^2*cos(1) - (cos(1) + I*sin(1))*sin(3)^2 + 2*I*(cos(3)*cos(1)
+ I*cos(3)*sin(1))*sin(3) + I*(cos(3)^2 + 1)*sin(1) + cos(1))*cos_integra
l(3*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) - 3*(-2*I*cos(3)*cos(1)
*sin(1) + cos(3)*sin(1)^2 - (cos(1)^2 + 1)*cos(3) - I*(cos(1)^2 + 2*I*cos(
1)*sin(1) - sin(1)^2 + 1)*sin(3))*cos_integral((cosh(b*x + a) + sinh(b*x +
a))/cosh(b*x + a)) - (cos(3)^2*cos(1) - (cos(1) + I*sin(1))*sin(3)^2 + 2*
I*(cos(3)*cos(1) + I*cos(3)*sin(1))*sin(3) + I*(cos(3)^2 + 1)*sin(1) + cos
(1))*cos_integral(6/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sin
h(b*x + a)^2 + 1)) - 3*(2*I*cos(3)*cos(1)*sin(1) - cos(3)*sin(1)^2 + (cos(
1)^2 + 1)*cos(3) + I*(cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*sin(3))
*cos_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*
x + a)^2 + 1)) + (-I*cos(3)^2*cos(1) - (-I*cos(1) + sin(1))*sin(3)^2 - 2*I
*(I*cos(3)*cos(1) - cos(3)*sin(1))*sin(3) + I*(-I*cos(3)^2 + I)*sin(1) + I
*cos(1))*sin_integral(3*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + 3
*(2*cos(3)*cos(1)*sin(1) + I*cos(3)*sin(1)^2 - (I*cos(1)^2 - I)*cos(3) - I
*(I*cos(1)^2 - 2*cos(1)*sin(1) - I*sin(1)^2 - I)*sin(3))*sin_integral((cos
h(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (I*cos(3)^2*cos(1) - (I*cos(1)
) - sin(1))*sin(3)^2 - 2*I*(-I*cos(3)*cos(1) + cos(3)*sin(1))*sin(3) + I*(
I*cos(3)^2 - I)*sin(1) - I*cos(1))*sin_integral(6/(cosh(b*x + a)^2 + 2*cos
h(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) - 3*(2*cos(3)*cos(1)*s...

```

## Sympy [F]

$$\int \cos^3(\tanh(a + bx)) dx = \int \cos^3(\tanh(a + bx)) dx$$

input

```
integrate(cos(tanh(b*x+a))**3,x)
```

output

```
Integral(cos(tanh(a + b*x))**3, x)
```

**Maxima [F]**

$$\int \cos^3(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^3 dx$$

input `integrate(cos(tanh(b*x+a))^3,x, algorithm="maxima")`

output `integrate(cos(tanh(b*x + a))^3, x)`

**Giac [F]**

$$\int \cos^3(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^3 dx$$

input `integrate(cos(tanh(b*x+a))^3,x, algorithm="giac")`

output `integrate(cos(tanh(b*x + a))^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \cos^3(\tanh(a + bx)) dx = \int \cos(\tanh(a + bx))^3 dx$$

input `int(cos(tanh(a + b*x))^3,x)`

output `int(cos(tanh(a + b*x))^3, x)`

**Reduce [F]**

$$\int \cos^3(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^3 dx$$

input `int(cos(tanh(b*x+a))^3,x)`

output `int(cos(tanh(a + b*x))**3,x)`

### 3.256 $\int \cos^2(\tanh(a + bx)) dx$

Optimal result	1869
Mathematica [A] (verified)	1870
Rubi [A] (verified)	1870
Maple [A] (verified)	1872
Fricas [C] (verification not implemented)	1872
Sympy [F]	1873
Maxima [F]	1873
Giac [F]	1874
Mupad [F(-1)]	1874
Reduce [F]	1874

#### Optimal result

Integrand size = 9, antiderivative size = 115

$$\int \cos^2(\tanh(a + bx)) dx = -\frac{\cos(2) \operatorname{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} + \frac{\cos(2) \operatorname{CosIntegral}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} - \frac{\sin(2) \operatorname{Si}(2 - 2 \tanh(a + bx))}{4b} + \frac{\sin(2) \operatorname{Si}(2 + 2 \tanh(a + bx))}{4b}$$

output

```
-1/4*cos(2)*Ci(2-2*tanh(b*x+a))/b+1/4*cos(2)*Ci(2+2*tanh(b*x+a))/b-1/4*ln(
1-tanh(b*x+a))/b+1/4*ln(1+tanh(b*x+a))/b+1/4*sin(2)*Si(-2+2*tanh(b*x+a))/b
+1/4*sin(2)*Si(2+2*tanh(b*x+a))/b
```

**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

$$\int \cos^2(\tanh(a + bx)) dx$$

$$= \frac{-\cos(2) \operatorname{CosIntegral}(2 - 2 \tanh(a + bx)) + \cos(2) \operatorname{CosIntegral}(2(1 + \tanh(a + bx))) - \log(1 - \tanh(a + bx)) + \log(1 + \tanh(a + bx)) - \operatorname{SinIntegral}(2 - 2 \tanh(a + bx)) + \operatorname{SinIntegral}(2(1 + \tanh(a + bx)))}{4b}$$

input `Integrate[Cos[Tanh[a + b*x]]^2,x]`

output `(-(Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]]) + Cos[2]*CosIntegral[2*(1 + Tanh[a + b*x])]) - Log[1 - Tanh[a + b*x]] + Log[1 + Tanh[a + b*x]] - Sin[2]*SinIntegral[2 - 2*Tanh[a + b*x]] + Sin[2]*SinIntegral[2*(1 + Tanh[a + b*x])])/(4*b)`

**Rubi [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.88, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {4853, 7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(\tanh(a + bx)) dx$$

$$\downarrow 4853$$

$$\int \frac{\cos^2(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)$$

$$\downarrow 7276$$

$$\int \left( \frac{\cos^2(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\cos^2(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)$$

$$\downarrow 2009$$

$$\frac{-\frac{1}{4} \cos(2) \operatorname{CosIntegral}(2 - 2 \tanh(a + bx)) + \frac{1}{4} \cos(2) \operatorname{CosIntegral}(2 \tanh(a + bx) + 2) - \frac{1}{4} \sin(2) \operatorname{Si}(2 - 2 \tanh(a + bx)) + \frac{1}{4} \sin(2) \operatorname{Si}(2 \tanh(a + bx) + 2)}{b}$$

input `Int[Cos[Tanh[a + b*x]]^2,x]`

output `(-1/4*(Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]]) + (Cos[2]*CosIntegral[2 + 2*Tanh[a + b*x]])/4 - Log[1 - Tanh[a + b*x]]/4 + Log[1 + Tanh[a + b*x]]/4 - (Sin[2]*SinIntegral[2 - 2*Tanh[a + b*x]])/4 + (Sin[2]*SinIntegral[2 + 2*Tanh[a + b*x]])/4)/b`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 4853 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x^2), Tan[v]/d, u, x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x]]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`



**Maple [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{\text{Si}(2+2 \tanh(bx+a)) \sin(2) + \text{Ci}(2+2 \tanh(bx+a)) \cos(2) + \text{Si}(-2+2 \tanh(bx+a)) \sin(2) - \text{Ci}(-2+2 \tanh(bx+a)) \cos(2) - \ln(-1+\tanh(bx+a))}{4b}$
default	$\frac{\text{Si}(2+2 \tanh(bx+a)) \sin(2) + \text{Ci}(2+2 \tanh(bx+a)) \cos(2) + \text{Si}(-2+2 \tanh(bx+a)) \sin(2) - \text{Ci}(-2+2 \tanh(bx+a)) \cos(2) - \ln(-1+\tanh(bx+a))}{4b}$
risch	$\frac{e^{-2i} \text{expIntegral}_1\left(-\frac{4i}{1+e^{2bx+2a}}\right)}{8b} - \frac{e^{-2i} \text{expIntegral}_1\left(\frac{4i}{1+e^{2bx+2a}}-4i\right)}{8b} - \frac{e^{2i} \text{expIntegral}_1\left(-\frac{4i}{1+e^{2bx+2a}}+4i\right)}{8b} +$

input `int(cos(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`output `1/b*(1/4*Si(2+2*tanh(b*x+a))*sin(2)+1/4*Ci(2+2*tanh(b*x+a))*cos(2)+1/4*Si(-2+2*tanh(b*x+a))*sin(2)-1/4*Ci(-2+2*tanh(b*x+a))*cos(2)-1/4*ln(-1+tanh(b*x+a))+1/4*ln(1+tanh(b*x+a)))`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.00

$$\int \cos^2(\tanh(a + bx)) dx$$

$$= \frac{4bx \cos(2) + 4i bx \sin(2) + (\cos(2)^2 + 2i \cos(2) \sin(2) - \sin(2)^2 + 1) \text{Ci}\left(\frac{2(\cosh(bx+a) + \sinh(bx+a))}{\cosh(bx+a)}\right) - (c$$

input `integrate(cos(tanh(b*x+a))^2,x, algorithm="fricas")`

output

```
1/8*(4*b*x*cos(2) + 4*I*b*x*sin(2) + (cos(2)^2 + 2*I*cos(2)*sin(2) - sin(2)^2 + 1)*cos_integral(2*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) - (cos(2)^2 + 2*I*cos(2)*sin(2) - sin(2)^2 + 1)*cos_integral(4/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) + (-I*cos(2)^2 + 2*cos(2)*sin(2) + I*sin(2)^2 + I)*sin_integral(2*(cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (I*cos(2)^2 - 2*cos(2)*sin(2) - I*sin(2)^2 - I)*sin_integral(4/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)))/(b*cos(2) + I*b*sin(2))
```

**Sympy [F]**

$$\int \cos^2(\tanh(a + bx)) dx = \int \cos^2(\tanh(a + bx)) dx$$

input

```
integrate(cos(tanh(b*x+a))**2,x)
```

output

```
Integral(cos(tanh(a + b*x))**2, x)
```

**Maxima [F]**

$$\int \cos^2(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^2 dx$$

input

```
integrate(cos(tanh(b*x+a))^2,x, algorithm="maxima")
```

output

```
1/2*x + 1/2*integrate(cos(2*(e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)), x)
```

**Giac [F]**

$$\int \cos^2(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^2 dx$$

input `integrate(cos(tanh(b*x+a))^2,x, algorithm="giac")`

output `integrate(cos(tanh(b*x + a))^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \cos^2(\tanh(a + bx)) dx = \int \cos(\tanh(a + bx))^2 dx$$

input `int(cos(tanh(a + b*x))^2,x)`

output `int(cos(tanh(a + b*x))^2, x)`

**Reduce [F]**

$$\int \cos^2(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a))^2 dx$$

input `int(cos(tanh(b*x+a))^2,x)`

output `int(cos(tanh(a + b*x))**2,x)`

### 3.257 $\int \cos(\tanh(a + bx)) dx$

Optimal result	1875
Mathematica [A] (verified)	1875
Rubi [A] (verified)	1876
Maple [A] (verified)	1877
Fricas [C] (verification not implemented)	1878
Sympy [F]	1878
Maxima [F]	1879
Giac [F]	1879
Mupad [F(-1)]	1879
Reduce [F]	1880

#### Optimal result

Integrand size = 7, antiderivative size = 77

$$\int \cos(\tanh(a + bx)) dx = -\frac{\cos(1) \operatorname{CosIntegral}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1) \operatorname{CosIntegral}(1 + \tanh(a + bx))}{2b} - \frac{\sin(1) \operatorname{Si}(1 - \tanh(a + bx))}{2b} + \frac{\sin(1) \operatorname{Si}(1 + \tanh(a + bx))}{2b}$$

output -1/2\*cos(1)\*Ci(1-tanh(b\*x+a))/b+1/2\*cos(1)\*Ci(1+tanh(b\*x+a))/b+1/2\*sin(1)\*Si(-1+tanh(b\*x+a))/b+1/2\*sin(1)\*Si(1+tanh(b\*x+a))/b

#### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.81

$$\int \cos(\tanh(a + bx)) dx = \frac{-\cos(1) \operatorname{CosIntegral}(1 - \tanh(a + bx)) + \cos(1) \operatorname{CosIntegral}(1 + \tanh(a + bx)) - \sin(1) \operatorname{Si}(1 - \tanh(a + bx)) + \sin(1) \operatorname{Si}(1 + \tanh(a + bx))}{2b}$$

input Integrate[Cos[Tanh[a + b\*x]],x]

output

```
(-(Cos[1]*CosIntegral[1 - Tanh[a + b*x]]) + Cos[1]*CosIntegral[1 + Tanh[a + b*x]] - Sin[1]*SinIntegral[1 - Tanh[a + b*x]] + Sin[1]*SinIntegral[1 + Tanh[a + b*x]])/(2*b)
```

**Rubi [A] (verified)**

Time = 0.59 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.90, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {4853, 3815, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(\tanh(a + bx)) dx$$

$$\downarrow 4853$$

$$\frac{\int \frac{\cos(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b}$$

$$\downarrow 3815$$

$$\frac{\int \left( \frac{\cos(\tanh(a+bx))}{2(1-\tanh(a+bx))} + \frac{\cos(\tanh(a+bx))}{2(\tanh(a+bx)+1)} \right) d \tanh(a + bx)}{b}$$

$$\downarrow 2009$$

$$\frac{-\frac{1}{2} \cos(1) \text{CosIntegral}(1 - \tanh(a + bx)) + \frac{1}{2} \cos(1) \text{CosIntegral}(\tanh(a + bx) + 1) - \frac{1}{2} \sin(1) \text{Si}(1 - \tanh(a + bx))}{b}$$

input

```
Int[Cos[Tanh[a + b*x]],x]
```

output

```
(-1/2*(Cos[1]*CosIntegral[1 - Tanh[a + b*x]]) + (Cos[1]*CosIntegral[1 + Tanh[a + b*x]])/2 - (Sin[1]*SinIntegral[1 - Tanh[a + b*x]])/2 + (Sin[1]*SinIntegral[1 + Tanh[a + b*x]])/2)/b
```

**Defintions of rubi rules used**

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3815 Int[Cos[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int
[ExpandIntegrand[Cos[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d},
x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])
```

```
rule 4853 Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Tan[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x, True] && TryPureTanSubst[ActivateTrig[u], x
]]
```

**Maple [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{\frac{\text{Si}(1+\tanh(bx+a)) \sin(1)}{2} + \frac{\text{Ci}(1+\tanh(bx+a)) \cos(1)}{2} + \frac{\text{Si}(-1+\tanh(bx+a)) \sin(1)}{2} - \frac{\text{Ci}(-1+\tanh(bx+a)) \cos(1)}{2}}{b}$
default	$\frac{\frac{\text{Si}(1+\tanh(bx+a)) \sin(1)}{2} + \frac{\text{Ci}(1+\tanh(bx+a)) \cos(1)}{2} + \frac{\text{Si}(-1+\tanh(bx+a)) \sin(1)}{2} - \frac{\text{Ci}(-1+\tanh(bx+a)) \cos(1)}{2}}{b}$
risch	$\frac{e^i \exp\text{Integral}_1\left(\frac{2i}{1+e^{2bx+2a}}\right)}{4b} - \frac{e^{-i} \exp\text{Integral}_1\left(\frac{2i}{1+e^{2bx+2a}} - 2i\right)}{4b} + \frac{e^{-i} \exp\text{Integral}_1\left(-\frac{2i}{1+e^{2bx+2a}}\right)}{4b} - \frac{e^i \exp\text{Integral}_1\left(\frac{2i}{1+e^{2bx+2a}}\right)}{4b}$

```
input int(cos(tanh(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output 1/b*(1/2*Si(1+tanh(b*x+a))*sin(1)+1/2*Ci(1+tanh(b*x+a))*cos(1)+1/2*Si(-1+t
anh(b*x+a))*sin(1)-1/2*Ci(-1+tanh(b*x+a))*cos(1))
```

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 216, normalized size of antiderivative = 2.81

$$\int \cos(\tanh(a + bx)) dx$$

$$= \frac{(\cos(1)^2 + 2i \cos(1) \sin(1) - \sin(1)^2 + 1) \operatorname{Ci}\left(\frac{\cosh(bx+a) + \sinh(bx+a)}{\cosh(bx+a)}\right) - (\cos(1)^2 + 2i \cos(1) \sin(1) - \sin(1)^2 + 1) \operatorname{Ci}\left(\frac{\cosh(bx+a) - \sinh(bx+a)}{\cosh(bx+a)}\right)}{2}$$

input `integrate(cos(tanh(b*x+a)),x, algorithm="fricas")`

output `1/4*((cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*cos_integral((cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) - (cos(1)^2 + 2*I*cos(1)*sin(1) - sin(1)^2 + 1)*cos_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)) + (-I*cos(1)^2 + 2*cos(1)*sin(1) + I*sin(1)^2 + I)*sin_integral((cosh(b*x + a) + sinh(b*x + a))/cosh(b*x + a)) + (I*cos(1)^2 - 2*cos(1)*sin(1) - I*sin(1)^2 - I)*sin_integral(2/(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)))/(b*cos(1) + I*b*sin(1))`

**Sympy [F]**

$$\int \cos(\tanh(a + bx)) dx = \int \cos(\tanh(a + bx)) dx$$

input `integrate(cos(tanh(b*x+a)),x)`

output `Integral(cos(tanh(a + b*x)), x)`

**Maxima [F]**

$$\int \cos(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a)) dx$$

input `integrate(cos(tanh(b*x+a)),x, algorithm="maxima")`

output `integrate(cos(tanh(b*x + a)), x)`

**Giac [F]**

$$\int \cos(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a)) dx$$

input `integrate(cos(tanh(b*x+a)),x, algorithm="giac")`

output `integrate(cos(tanh(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \cos(\tanh(a + bx)) dx = \int \cos(\tanh(a + bx)) dx$$

input `int(cos(tanh(a + b*x)),x)`

output `int(cos(tanh(a + b*x)), x)`



**Reduce [F]**

$$\int \cos(\tanh(a + bx)) dx = \int \cos(\tanh(bx + a)) dx$$

input `int(cos(tanh(b*x+a)),x)`

output `int(cos(tanh(a + b*x)),x)`

### 3.258 $\int \sec(\tanh(a + bx)) dx$

Optimal result	1881
Mathematica [N/A]	1881
Rubi [N/A]	1882
Maple [F(-1)]	1883
Fricas [N/A]	1883
Sympy [N/A]	1883
Maxima [N/A]	1884
Giac [N/A]	1884
Mupad [N/A]	1884
Reduce [N/A]	1885

#### Optimal result

Integrand size = 7, antiderivative size = 7

$$\int \sec(\tanh(a + bx)) dx = -\frac{1}{2} \operatorname{Int}\left(\frac{\sec(\tanh(a + bx)) \operatorname{sech}^2(a + bx)}{1 - \tanh(a + bx)}, x\right) + \frac{1}{2} \operatorname{Int}\left(\frac{\sec(\tanh(a + bx)) \operatorname{sech}^2(a + bx)}{1 + \tanh(a + bx)}, x\right)$$

output `-1/2*Defer(Int)(-sec(tanh(b*x+a))*sech(b*x+a)^2/(1-tanh(b*x+a)),x)+1/2*Defer(Int)(sec(tanh(b*x+a))*sech(b*x+a)^2/(1+tanh(b*x+a)),x)`

#### Mathematica [N/A]

Not integrable

Time = 3.65 (sec), antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(a + bx)) dx$$

input `Integrate[Sec[Tanh[a + b*x]],x]`

output `Integrate[Sec[Tanh[a + b*x]], x]`

**Rubi [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec(\tanh(a + bx)) dx \\
 \downarrow 4853 \\
 \frac{\int \frac{\sec(\tanh(a+bx))}{1-\tanh^2(a+bx)} d \tanh(a + bx)}{b} \\
 \downarrow 7276 \\
 \frac{\int \left( \frac{\sec(\tanh(a+bx))}{2(\tanh(a+bx)+1)} - \frac{\sec(\tanh(a+bx))}{2(\tanh(a+bx)-1)} \right) d \tanh(a + bx)}{b} \\
 \downarrow 2009 \\
 \frac{\frac{1}{2} \int \frac{\sec(\tanh(a+bx))}{\tanh(a+bx)+1} d \tanh(a + bx) - \frac{1}{2} \int \frac{\sec(\tanh(a+bx))}{\tanh(a+bx)-1} d \tanh(a + bx)}{b}
 \end{array}$$

input `Int [Sec [Tanh [a + b*x] ] , x]`

output `$Aborted`

**Maple [F(-1)]**

Timed out.

$$\int \sec(\tanh(bx + a)) dx$$

input `int(sec(tanh(b*x+a)),x)`output `int(sec(tanh(b*x+a)),x)`**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(bx + a)) dx$$

input `integrate(sec(tanh(b*x+a)),x, algorithm="fricas")`output `integral(sec(tanh(b*x + a)), x)`**Sympy [N/A]**

Not integrable

Time = 2.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(a + bx)) dx$$

input `integrate(sec(tanh(b*x+a)),x)`output `Integral(sec(tanh(a + b*x)), x)`

**Maxima [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(bx + a)) dx$$

input `integrate(sec(tanh(b*x+a)),x, algorithm="maxima")`output `integrate(sec(tanh(b*x + a)), x)`**Giac [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(bx + a)) dx$$

input `integrate(sec(tanh(b*x+a)),x, algorithm="giac")`output `integrate(sec(tanh(b*x + a)), x)`**Mupad [N/A]**

Not integrable

Time = 2.85 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.57

$$\int \sec(\tanh(a + bx)) dx = \int \frac{1}{\cos(\tanh(a + bx))} dx$$

input `int(1/cos(tanh(a + b*x)),x)`

output `int(1/cos(tanh(a + b*x)), x)`

**Reduce [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.29

$$\int \sec(\tanh(a + bx)) dx = \int \sec(\tanh(bx + a)) dx$$

input `int(sec(tanh(b*x+a)), x)`

output `int(sec(tanh(a + b*x)), x)`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	1886
4.2	Links to plain text integration problems used in this report for each CAS .	1904

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```



```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```



## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```



```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```



## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file