

Computer Algebra Independent Integration Tests

Summer 2024

6-Hyperbolic-functions/6-Miscellaneous/292-6.2

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 1:17am

Contents

1	Introduction	7
1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26
2	detailed summary tables of results	27
2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	76
3	Listing of integrals	82
3.1	$\int \sinh(x) \sinh(2x) dx$	87
3.2	$\int \sinh(x) \sinh(3x) dx$	92
3.3	$\int \sinh(x) \sinh(4x) dx$	97
3.4	$\int \sinh(x) \sinh(mx) dx$	102
3.5	$\int \sinh(a + bx) \sinh(c + bx) dx$	107
3.6	$\int \sinh(c - bx) \sinh(a + bx) dx$	113
3.7	$\int \sinh(a + bx) \sinh(c + dx) dx$	119

3.8	$\int \sinh(a + bx) \sinh^2(c + dx) dx$	125
3.9	$\int \sinh(a + bx) \sinh^3(c + dx) dx$	131
3.10	$\int \sinh^2(a + bx) \sinh^2(c + dx) dx$	138
3.11	$\int \sinh^2(a + bx) \sinh^3(c + dx) dx$	145
3.12	$\int \sinh^3(a + bx) \sinh^3(c + dx) dx$	154
3.13	$\int \cosh(2x) \sinh(x) dx$	163
3.14	$\int \cosh(3x) \sinh(x) dx$	168
3.15	$\int \cosh(4x) \sinh(x) dx$	173
3.16	$\int \cosh(mx) \sinh(x) dx$	178
3.17	$\int \cosh(c + dx) \sinh(a + bx) dx$	183
3.18	$\int \cosh^2(c + dx) \sinh(a + bx) dx$	189
3.19	$\int \cosh^3(c + dx) \sinh(a + bx) dx$	195
3.20	$\int \cosh(c + dx) \sinh^2(a + bx) dx$	202
3.21	$\int \cosh^2(c + dx) \sinh^2(a + bx) dx$	208
3.22	$\int \cosh^3(c + dx) \sinh^2(a + bx) dx$	215
3.23	$\int \cosh(c + dx) \sinh^3(a + bx) dx$	224
3.24	$\int \cosh^2(c + dx) \sinh^3(a + bx) dx$	231
3.25	$\int \cosh^3(c + dx) \sinh^3(a + bx) dx$	240
3.26	$\int \sinh(x) \tanh(2x) dx$	249
3.27	$\int \sinh(x) \tanh(3x) dx$	255
3.28	$\int \sinh(x) \tanh(4x) dx$	262
3.29	$\int \sinh(x) \tanh(5x) dx$	269
3.30	$\int \sinh(x) \tanh(6x) dx$	276
3.31	$\int \sinh(x) \tanh(nx) dx$	283
3.32	$\int \sinh(a + bx) \tanh(c + bx) dx$	288
3.33	$\int \sinh(a + bx) \tanh^2(c + bx) dx$	294
3.34	$\int \sinh(a + bx) \tanh^3(c + bx) dx$	302
3.35	$\int \sinh(a + bx) \tanh(c + dx) dx$	311
3.36	$\int \coth(2x) \sinh(x) dx$	316
3.37	$\int \coth(3x) \sinh(x) dx$	321
3.38	$\int \coth(4x) \sinh(x) dx$	327
3.39	$\int \coth(5x) \sinh(x) dx$	333
3.40	$\int \coth(6x) \sinh(x) dx$	340
3.41	$\int \coth(c + bx) \sinh(a + bx) dx$	346
3.42	$\int \coth^2(c + bx) \sinh(a + bx) dx$	353
3.43	$\int \coth^3(c + bx) \sinh(a + bx) dx$	361
3.44	$\int \coth(c + dx) \sinh(a + bx) dx$	370
3.45	$\int \operatorname{sech}(2x) \sinh(x) dx$	375
3.46	$\int \operatorname{sech}(3x) \sinh(x) dx$	381

3.47	$\int \operatorname{sech}(4x) \sinh(x) dx$	387
3.48	$\int \operatorname{sech}(5x) \sinh(x) dx$	396
3.49	$\int \operatorname{sech}(6x) \sinh(x) dx$	403
3.50	$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx$	413
3.51	$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx$	419
3.52	$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx$	426
3.53	$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx$	432
3.54	$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx$	440
3.55	$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx$	445
3.56	$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx$	451
3.57	$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx$	456
3.58	$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx$	461
3.59	$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx$	466
3.60	$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx$	472
3.61	$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx$	478
3.62	$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx$	484
3.63	$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx$	489
3.64	$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx$	495
3.65	$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx$	500
3.66	$\int \operatorname{csch}(2x) \sinh(x) dx$	506
3.67	$\int \operatorname{csch}(3x) \sinh(x) dx$	511
3.68	$\int \operatorname{csch}(4x) \sinh(x) dx$	516
3.69	$\int \operatorname{csch}(5x) \sinh(x) dx$	522
3.70	$\int \operatorname{csch}(6x) \sinh(x) dx$	529
3.71	$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx$	535
3.72	$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx$	541
3.73	$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx$	548
3.74	$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$	555
3.75	$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx$	564
3.76	$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx$	569
3.77	$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx$	575
3.78	$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$	580
3.79	$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$	586
3.80	$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx$	592
3.81	$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx$	598
3.82	$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx$	604
3.83	$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$	610
3.84	$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx$	616
3.85	$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx$	622

3.86	$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx$	628
3.87	$\int \cosh(x) \sinh(2x) dx$	634
3.88	$\int \cosh(x) \sinh(3x) dx$	639
3.89	$\int \cosh(x) \sinh(4x) dx$	644
3.90	$\int \cosh(x) \sinh(mx) dx$	649
3.91	$\int \cosh(x) \cosh(2x) dx$	654
3.92	$\int \cosh(x) \cosh(3x) dx$	659
3.93	$\int \cosh(x) \cosh(4x) dx$	664
3.94	$\int \cosh(x) \cosh(mx) dx$	669
3.95	$\int \cosh(a + bx) \cosh(c + bx) dx$	674
3.96	$\int \cosh(c - bx) \cosh(a + bx) dx$	680
3.97	$\int \cosh(a + bx) \cosh(c + dx) dx$	686
3.98	$\int \cosh(a + bx) \cosh^2(c + dx) dx$	692
3.99	$\int \cosh(a + bx) \cosh^3(c + dx) dx$	698
3.100	$\int \cosh^2(a + bx) \cosh^2(c + dx) dx$	705
3.101	$\int \cosh^2(a + bx) \cosh^3(c + dx) dx$	712
3.102	$\int \cosh^3(a + bx) \cosh^3(c + dx) dx$	721
3.103	$\int \cosh(x) \tanh(2x) dx$	730
3.104	$\int \cosh(x) \tanh(3x) dx$	736
3.105	$\int \cosh(x) \tanh(4x) dx$	742
3.106	$\int \cosh(x) \tanh(5x) dx$	750
3.107	$\int \cosh(x) \tanh(6x) dx$	757
3.108	$\int \cosh(a + bx) \tanh(c + bx) dx$	765
3.109	$\int \cosh(a + bx) \tanh^2(c + bx) dx$	771
3.110	$\int \cosh(a + bx) \tanh^3(c + bx) dx$	779
3.111	$\int \cosh(a + bx) \tanh(c + dx) dx$	788
3.112	$\int \cosh(x) \operatorname{coth}(2x) dx$	793
3.113	$\int \cosh(x) \operatorname{coth}(3x) dx$	799
3.114	$\int \cosh(x) \operatorname{coth}(4x) dx$	806
3.115	$\int \cosh(x) \operatorname{coth}(5x) dx$	813
3.116	$\int \cosh(x) \operatorname{coth}(6x) dx$	821
3.117	$\int \cosh(x) \operatorname{coth}(nx) dx$	828
3.118	$\int \cosh(a + bx) \operatorname{coth}(c + bx) dx$	833
3.119	$\int \cosh(a + bx) \operatorname{coth}^2(c + bx) dx$	840
3.120	$\int \cosh(a + bx) \operatorname{coth}^3(c + bx) dx$	848
3.121	$\int \cosh(a + bx) \operatorname{coth}(c + dx) dx$	858
3.122	$\int \cosh(x) \operatorname{sech}(2x) dx$	863
3.123	$\int \cosh(x) \operatorname{sech}(3x) dx$	869
3.124	$\int \cosh(x) \operatorname{sech}(4x) dx$	874

3.125	$\int \cosh(x)\operatorname{sech}(5x) dx$	881
3.126	$\int \cosh(x)\operatorname{sech}(6x) dx$	888
3.127	$\int \cosh(a+bx)\operatorname{sech}(c+bx) dx$	896
3.128	$\int \cosh(a+bx)\operatorname{sech}^2(c+bx) dx$	902
3.129	$\int \cosh(a+bx)\operatorname{sech}^3(c+bx) dx$	909
3.130	$\int \cosh(a+bx)\operatorname{sech}^4(c+bx) dx$	915
3.131	$\int \cosh(a+bx)\operatorname{sech}(c-bx) dx$	923
3.132	$\int \cosh(a+bx)\operatorname{sech}^2(c-bx) dx$	928
3.133	$\int \cosh(a+bx)\operatorname{sech}^3(c-bx) dx$	934
3.134	$\int \cosh(a+bx)\operatorname{sech}^4(c-bx) dx$	939
3.135	$\int \cosh^2(a+bx)\operatorname{sech}(c+bx) dx$	944
3.136	$\int \cosh^2(a+bx)\operatorname{sech}^2(c+bx) dx$	949
3.137	$\int \cosh^2(a+bx)\operatorname{sech}^3(c+bx) dx$	955
3.138	$\int \cosh^2(a+bx)\operatorname{sech}^4(c+bx) dx$	961
3.139	$\int \cosh^2(a+bx)\operatorname{sech}(c-bx) dx$	967
3.140	$\int \cosh^2(a+bx)\operatorname{sech}^2(c-bx) dx$	972
3.141	$\int \cosh^2(a+bx)\operatorname{sech}^3(c-bx) dx$	978
3.142	$\int \cosh^2(a+bx)\operatorname{sech}^4(c-bx) dx$	983
3.143	$\int \cosh(x)\operatorname{csch}(2x) dx$	989
3.144	$\int \cosh(x)\operatorname{csch}(3x) dx$	994
3.145	$\int \cosh(x)\operatorname{csch}(4x) dx$	1000
3.146	$\int \cosh(x)\operatorname{csch}(5x) dx$	1006
3.147	$\int \cosh(x)\operatorname{csch}(6x) dx$	1013
3.148	$\int \cosh(a+bx)\operatorname{csch}(c+bx) dx$	1020
3.149	$\int \cosh(a+bx)\operatorname{csch}^2(c+bx) dx$	1026
3.150	$\int \cosh(a+bx)\operatorname{csch}^3(c+bx) dx$	1033
3.151	$\int \cosh(a+bx)\operatorname{csch}^4(c+bx) dx$	1040
3.152	$\int \cosh(a+bx)\operatorname{csch}(c-bx) dx$	1049
3.153	$\int \cosh(a+bx)\operatorname{csch}^2(c-bx) dx$	1054
3.154	$\int \cosh(a+bx)\operatorname{csch}^3(c-bx) dx$	1060
3.155	$\int \cosh(a+bx)\operatorname{csch}^4(c-bx) dx$	1065
3.156	$\int \cosh^2(a+bx)\operatorname{csch}(c+bx) dx$	1071
3.157	$\int \cosh^2(a+bx)\operatorname{csch}^2(c+bx) dx$	1077
3.158	$\int \cosh^2(a+bx)\operatorname{csch}^3(c+bx) dx$	1083
3.159	$\int \cosh^2(a+bx)\operatorname{csch}^4(c+bx) dx$	1089
3.160	$\int \cosh^2(a+bx)\operatorname{csch}(c-bx) dx$	1095
3.161	$\int \cosh^2(a+bx)\operatorname{csch}^2(c-bx) dx$	1101
3.162	$\int \cosh^2(a+bx)\operatorname{csch}^3(c-bx) dx$	1107
3.163	$\int \cosh^2(a+bx)\operatorname{csch}^4(c-bx) dx$	1113

3.164	$\int \tanh(a + bx) \tanh(c + bx) dx$	1119
3.165	$\int \tanh(c - bx) \tanh(a + bx) dx$	1125
3.166	$\int \coth(a + bx) \coth(c + bx) dx$	1131
3.167	$\int \coth(c - bx) \coth(a + bx) dx$	1138
3.168	$\int \operatorname{sech}(a + bx) \operatorname{sech}(c + bx) dx$	1145
3.169	$\int \operatorname{sech}(c - bx) \operatorname{sech}(a + bx) dx$	1151
3.170	$\int \operatorname{csch}(a + bx) \operatorname{csch}(c + bx) dx$	1157
3.171	$\int \operatorname{csch}(c - bx) \operatorname{csch}(a + bx) dx$	1163
4	Appendix	1169
4.1	Listing of Grading functions	1169
4.2	Links to plain text integration problems used in this report for each CAS	187

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [171]. This is test number [292].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (171)	0.00 (0)
Maple	96.49 (165)	3.51 (6)
Fricas	96.49 (165)	3.51 (6)
Giac	96.49 (165)	3.51 (6)
Reduce	90.06 (154)	9.94 (17)
Mupad	75.44 (129)	24.56 (42)
Rubi	71.93 (123)	28.07 (48)
Maxima	70.18 (120)	29.82 (51)
Sympy	23.98 (41)	76.02 (130)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Mathematica	78.363	9.942	11.696	0.000
Rubi	61.988	0.585	9.357	28.070
Maple	37.427	32.749	26.316	3.509
Giac	34.503	61.988	0.000	3.509
Maxima	11.111	59.064	0.000	29.825
Fricas	7.602	88.889	0.000	3.509
Sympy	7.602	16.374	0.000	76.023
Mupad	0.000	75.439	0.000	24.561
Reduce	0.000	90.058	0.000	9.942

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Fricas	6	100.00	0.00	0.00
Maple	6	100.00	0.00	0.00
Giac	6	100.00	0.00	0.00
Reduce	17	100.00	0.00	0.00
Mupad	42	0.00	100.00	0.00
Rubi	48	100.00	0.00	0.00
Maxima	51	50.98	0.00	49.02
Sympy	130	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.07
Fricas	0.10
Giac	0.13
Reduce	0.24
Rubi	0.30
Mathematica	0.30
Mupad	0.81
Maple	1.54
Sympy	2.20

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	56.28	1.12	42.00	1.00
Mathematica	73.22	1.44	67.00	1.01
Giac	97.32	1.97	85.00	1.91
Maxima	110.99	2.71	99.00	2.44
Maple	118.94	2.68	84.00	2.26
Mupad	127.19	2.60	86.00	2.13
Reduce	174.12	3.19	102.00	3.00
Sympy	690.93	6.21	153.00	3.56
Fricas	766.90	13.18	246.00	4.73

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```


For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	76

2.1 List of integrals sorted by grade for each CAS

Rubi	28
Mma	29
Maple	29
Fricas	30
Maxima	30
Giac	31
Mupad	31
Sympy	32
Reduce	32

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 66, 67, 68, 69, 70, 72, 73, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 114, 115, 116, 117, 118, 119, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 143, 144, 145, 146, 147, 149, 150, 164, 165, 168, 169 }

B grade { 113 }

C grade { 13, 14, 15, 43, 71, 74, 87, 88, 89, 120, 148, 151, 166, 167, 170, 171 }

F normal fail { 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 34, 35, 36, 37, 38, 39, 40, 43, 46, 48, 50, 52, 53, 54, 56, 57, 58, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 73, 74, 75, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 110, 111, 115, 117, 120, 121, 122, 123, 124, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 144, 146, 148, 150, 151, 152, 154, 155, 156, 157, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

B grade { 32, 33, 44, 51, 55, 59, 63, 81, 108, 109, 112, 113, 128, 132, 136, 140, 158 }

C grade { 41, 42, 45, 47, 49, 72, 76, 103, 104, 105, 106, 107, 114, 116, 118, 119, 145, 147, 149, 153 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 36, 46, 48, 52, 56, 61, 65, 66, 73, 77, 82, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 129, 133, 138, 142, 143, 144, 146, 150, 154, 159, 163 }

B grade { 41, 42, 43, 45, 50, 54, 59, 63, 71, 72, 74, 75, 76, 78, 79, 80, 81, 83, 84, 85, 103, 104, 112, 113, 114, 115, 116, 118, 119, 120, 127, 131, 136, 140, 145, 147, 148, 149, 151, 152, 153, 155, 156, 157, 158, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171 }

C grade { 26, 27, 28, 29, 30, 32, 33, 34, 37, 38, 39, 40, 47, 49, 51, 53, 55, 57, 58, 60, 62, 64, 67, 68, 69, 70, 105, 106, 107, 108, 109, 110, 122, 123, 124, 125, 126, 128, 130, 132, 134, 135, 137, 139, 141 }

F normal fail { 31, 35, 44, 111, 117, 121 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 4, 7, 13, 16, 17, 20, 66, 90, 91, 92, 94, 97 }

B grade { 1, 3, 5, 6, 8, 9, 10, 11, 12, 14, 15, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 36, 37, 38, 39, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 93, 95, 96, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 112, 113, 114, 115, 116, 118, 119, 120, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

C grade { }

F normal fail { 31, 35, 44, 111, 117, 121 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 32, 36, 50, 51, 55, 59, 60, 63, 64, 66, 127, 128, 132, 136, 137, 140, 141, 168, 169 }

B grade { 1, 2, 3, 5, 6, 13, 14, 15, 26, 27, 33, 34, 37, 38, 41, 42, 43, 45, 46, 52, 53, 54, 56, 57, 58, 61, 62, 65, 67, 68, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 91, 92, 93, 95, 96, 103, 104, 108, 109, 110, 112, 113, 114, 118, 119, 120, 122, 123, 129, 130, 131, 133, 134, 135, 138, 139, 142, 143, 144, 145, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 170, 171 }

C grade { }

F normal fail { 28, 29, 30, 31, 35, 39, 40, 44, 47, 48, 49, 69, 70, 105, 106, 107, 111, 115, 116, 117, 121, 124, 125, 126, 146, 147 }

F(-1) timedout fail { }

F(-2) exception fail { 4, 7, 8, 9, 10, 11, 12, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 90, 94, 97, 98, 99, 100, 101, 102 }

Giac

A grade { 10, 11, 14, 20, 21, 22, 28, 29, 30, 32, 34, 36, 39, 50, 51, 52, 54, 55, 56, 57, 58, 60, 61, 64, 65, 66, 67, 69, 71, 73, 75, 77, 82, 86, 88, 100, 101, 108, 110, 123, 125, 127, 128, 129, 131, 133, 134, 135, 137, 138, 141, 142, 146, 148, 150, 152, 154, 159, 163 }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 15, 16, 17, 18, 19, 23, 24, 25, 26, 27, 33, 37, 38, 40, 41, 42, 43, 45, 46, 47, 48, 49, 53, 59, 62, 63, 68, 70, 72, 74, 76, 78, 79, 80, 81, 83, 84, 85, 87, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 102, 103, 104, 105, 106, 107, 109, 112, 113, 114, 115, 116, 118, 119, 120, 122, 124, 126, 130, 132, 136, 139, 140, 143, 144, 145, 147, 149, 151, 153, 155, 156, 157, 158, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171 }

C grade { }

F normal fail { 31, 35, 44, 111, 117, 121 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 36, 37, 38, 39, 40, 41, 42, 45, 46, 47, 48, 49, 50, 51, 54, 55, 58, 59, 62, 63, 66, 67, 68, 69, 70, 71, 72, 75, 76, 79, 80, 83, 84, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 112, 113, 114, 115, 116, 118, 119, 122, 123, 124, 125, 126, 127, 128, 131, 132, 135, 136, 139, 140, 143, 144, 145, 146, 147, 148, 149, 152, 153, 156, 157, 160, 161, 164, 165, 166, 167, 168, 169, 170, 171 }

C grade { }

F normal fail { }

F(-1) timedout fail { 31, 34, 35, 43, 44, 52, 53, 56, 57, 60, 61, 64, 65, 73, 74, 77, 78, 81, 82, 85, 86, 110, 111, 117, 120, 121, 129, 130, 133, 134, 137, 138, 141, 142, 150, 151, 154, 155, 158, 159, 162, 163 }

F(-2) exception fail { }

Sympy

A grade { 2, 3, 6, 13, 14, 15, 16, 88, 89, 90, 91, 92, 93 }

B grade { 1, 4, 5, 7, 8, 9, 10, 11, 12, 17, 18, 19, 20, 21, 22, 23, 24, 25, 87, 94, 95, 96, 97, 98, 99, 100, 101, 102 }

C grade { }

F normal fail { 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 30, 32, 33, 34, 36, 37, 38, 40, 41, 42, 43, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 107, 108, 109, 110, 112, 113, 114, 116, 118, 119, 120, 122, 123, 124, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

C grade { }

F normal fail { 12, 25, 29, 31, 35, 39, 44, 48, 69, 102, 106, 111, 115, 117, 121, 125, 146 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	15	15	12	27	17	20	25	17	6
N.S.	1	1.88	1.88	1.50	3.38	2.12	2.50	3.12	2.12	0.75
time (sec)	N/A	0.174	0.023	0.165	0.032	0.079	0.125	0.111	0.219	0.057

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	27	22	20	27	17	13
N.S.	1	1.00	1.00	0.82	1.59	1.29	1.18	1.59	1.00	0.76
time (sec)	N/A	0.180	0.020	0.192	0.025	0.075	0.126	0.107	0.227	0.934

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	27	36	20	27	17	14
N.S.	1	1.00	1.00	0.82	1.59	2.12	1.18	1.59	1.00	0.82
time (sec)	N/A	0.177	0.021	0.188	0.042	0.076	0.127	0.104	0.259	0.955

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	25	28	0	42	78	59	25	26
N.S.	1	1.00	0.71	0.80	0.00	1.20	2.23	1.69	0.71	0.74
time (sec)	N/A	0.215	0.034	0.236	0.000	0.090	0.262	0.118	0.221	0.102

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	26	24	58	87	58	68	50	23
N.S.	1	1.00	0.96	0.89	2.15	3.22	2.15	2.52	1.85	0.85
time (sec)	N/A	0.212	0.026	0.256	0.047	0.088	0.188	0.115	0.224	0.172

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	59	75	61	70	57	23
N.S.	1	1.00	1.00	0.89	2.19	2.78	2.26	2.59	2.11	0.85
time (sec)	N/A	0.215	0.032	0.197	0.053	0.104	0.191	0.107	0.215	0.155

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	40	0	72	153	85	42	42
N.S.	1	1.00	1.00	0.93	0.00	1.67	3.56	1.98	0.98	0.98
time (sec)	N/A	0.243	0.157	0.319	0.000	0.084	0.322	0.111	0.272	1.048

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	69	57	0	120	405	120	212	76
N.S.	1	1.00	1.11	0.92	0.00	1.94	6.53	1.94	3.42	1.23
time (sec)	N/A	0.278	0.507	1.207	0.000	0.098	0.689	0.117	0.228	1.148

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	86	84	0	218	918	179	574	182
N.S.	1	1.00	0.95	0.92	0.00	2.40	10.09	1.97	6.31	2.00
time (sec)	N/A	0.304	0.322	2.853	0.000	0.105	1.932	0.117	0.227	0.545

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	106	89	0	192	1027	156	357	152
N.S.	1	1.00	1.20	1.01	0.00	2.18	11.67	1.77	4.06	1.73
time (sec)	N/A	0.293	0.508	3.666	0.000	0.087	1.578	0.122	0.220	1.488

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	158	133	0	414	1999	260	865	337
N.S.	1	1.00	1.10	0.92	0.00	2.88	13.88	1.81	6.01	2.34
time (sec)	N/A	0.371	1.179	9.105	0.000	0.097	5.327	0.122	0.243	1.513

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	195	195	177	190	0	731	3577	373	19	906
N.S.	1	1.00	0.91	0.97	0.00	3.75	18.34	1.91	0.10	4.65
time (sec)	N/A	0.425	1.104	26.381	0.000	0.096	16.616	0.129	200.021	1.617

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	23	15	12	27	19	20	25	17	11
N.S.	1	1.53	1.00	0.80	1.80	1.27	1.33	1.67	1.13	0.73
time (sec)	N/A	0.187	0.036	0.129	0.036	0.088	0.117	0.108	0.226	0.073

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	25	17	14	27	33	20	26	17	11
N.S.	1	1.47	1.00	0.82	1.59	1.94	1.18	1.53	1.00	0.65
time (sec)	N/A	0.190	0.004	0.144	0.027	0.075	0.124	0.105	0.245	0.945

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	25	17	14	27	38	20	27	17	15
N.S.	1	1.47	1.00	0.82	1.59	2.24	1.18	1.59	1.00	0.88
time (sec)	N/A	0.191	0.031	0.156	0.043	0.080	0.115	0.105	0.226	0.074

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	25	28	0	42	37	59	25	26
N.S.	1	1.00	0.71	0.80	0.00	1.20	1.06	1.69	0.71	0.74
time (sec)	N/A	0.221	0.042	0.195	0.000	0.091	0.243	0.117	0.225	0.099

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	40	0	71	153	85	42	42
N.S.	1	1.00	1.00	0.93	0.00	1.65	3.56	1.98	0.98	0.98
time (sec)	N/A	0.238	0.147	0.304	0.000	0.089	0.315	0.116	0.239	0.157

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	69	57	0	119	408	120	212	68
N.S.	1	1.00	1.11	0.92	0.00	1.92	6.58	1.94	3.42	1.10
time (sec)	N/A	0.263	0.448	1.207	0.000	0.112	0.695	0.114	0.242	1.134

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	85	84	0	213	921	179	573	182
N.S.	1	1.00	0.93	0.92	0.00	2.34	10.12	1.97	6.30	2.00
time (sec)	N/A	0.290	0.331	2.522	0.000	0.086	1.951	0.118	0.212	1.404

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	74	63	0	114	408	124	216	76
N.S.	1	1.00	1.09	0.93	0.00	1.68	6.00	1.82	3.18	1.12
time (sec)	N/A	0.255	0.544	1.195	0.000	0.094	0.734	0.114	0.223	1.167

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	107	89	0	192	1027	156	357	135
N.S.	1	1.00	1.22	1.01	0.00	2.18	11.67	1.77	4.06	1.53
time (sec)	N/A	0.274	0.445	3.259	0.000	0.092	1.591	0.116	0.241	1.460

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	158	133	0	398	1999	260	865	337
N.S.	1	1.00	1.10	0.92	0.00	2.76	13.88	1.81	6.01	2.34
time (sec)	N/A	0.338	1.046	8.387	0.000	0.098	5.394	0.121	0.238	1.520

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	90	90	0	243	937	183	573	183
N.S.	1	1.00	0.93	0.93	0.00	2.51	9.66	1.89	5.91	1.89
time (sec)	N/A	0.297	0.358	2.530	0.000	0.099	1.966	0.123	0.241	0.553

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	138	153	127	0	443	2030	256	865	337
N.S.	1	1.00	1.11	0.92	0.00	3.21	14.71	1.86	6.27	2.44
time (sec)	N/A	0.341	1.089	8.028	0.000	0.117	5.339	0.128	0.238	1.526

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	195	195	176	190	0	729	3580	373	19	908
N.S.	1	1.00	0.90	0.97	0.00	3.74	18.36	1.91	0.10	4.66
time (sec)	N/A	0.416	1.044	24.372	0.000	0.113	16.481	0.127	200.027	1.644

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	27	19	54	53	115	0	36	58	47
N.S.	1	1.42	1.00	2.84	2.79	6.05	0.00	1.89	3.05	2.47
time (sec)	N/A	0.210	0.014	0.070	0.115	0.085	0.000	0.114	0.220	1.009

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	60	46	76	0	43	55	23
N.S.	1	1.00	1.00	3.16	2.42	4.00	0.00	2.26	2.89	1.21
time (sec)	N/A	0.256	0.021	0.083	0.117	0.094	0.000	0.111	0.239	0.076

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	80	69	42	0	287	0	71	146	71
N.S.	1	1.07	0.92	0.56	0.00	3.83	0.00	0.95	1.95	0.95
time (sec)	N/A	0.314	0.097	0.082	0.000	0.105	0.000	0.154	0.270	1.757

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	87	81	60	0	246	0	81	52	82
N.S.	1	1.04	0.96	0.71	0.00	2.93	0.00	0.96	0.62	0.98
time (sec)	N/A	0.416	0.135	0.095	0.000	0.102	0.000	0.134	0.264	2.219

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	93	87	84	0	375	0	100	254	98
N.S.	1	1.07	1.00	0.97	0.00	4.31	0.00	1.15	2.92	1.13
time (sec)	N/A	0.396	0.091	0.093	0.000	0.134	0.000	0.119	0.256	2.235

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	81	77	0	0	0	0	0	56	0
N.S.	1	1.21	1.15	0.00	0.00	0.00	0.00	0.00	0.84	0.00
time (sec)	N/A	0.280	0.184	0.000	0.000	0.000	0.000	0.000	0.230	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	86	167	57	327	0	49	70	133
N.S.	1	1.00	2.97	5.76	1.97	11.28	0.00	1.69	2.41	4.59
time (sec)	N/A	0.280	0.048	0.093	0.123	0.113	0.000	0.117	0.260	1.169

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	102	205	105	902	0	97	152	173
N.S.	1	1.00	2.27	4.56	2.33	20.04	0.00	2.16	3.38	3.84
time (sec)	N/A	0.442	0.079	0.102	0.120	0.107	0.000	0.112	0.239	1.157

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	86	70	240	149	1737	0	120	134	0
N.S.	1	1.19	0.97	3.33	2.07	24.12	0.00	1.67	1.86	0.00
time (sec)	N/A	0.757	0.241	0.141	0.140	0.125	0.000	0.125	0.263	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	100	121	103	0	0	0	0	0	91	0
N.S.	1	1.21	1.03	0.00	0.00	0.00	0.00	0.00	0.91	0.00
time (sec)	N/A	0.355	0.683	0.000	0.000	0.000	0.000	0.000	0.250	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	14	10	9	16	42	0	16	23	16
N.S.	1	1.40	1.00	0.90	1.60	4.20	0.00	1.60	2.30	1.60
time (sec)	N/A	0.205	0.008	0.076	0.126	0.091	0.000	0.111	0.266	0.060

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	54	49	118	0	36	56	47
N.S.	1	1.00	1.00	2.70	2.45	5.90	0.00	1.80	2.80	2.35
time (sec)	N/A	0.215	0.013	0.089	0.142	0.117	0.000	0.114	0.249	1.027

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	30	28	72	60	128	0	54	69	52
N.S.	1	1.07	1.00	2.57	2.14	4.57	0.00	1.93	2.46	1.86
time (sec)	N/A	0.279	0.026	0.098	0.162	0.121	0.000	0.110	0.220	0.996

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	82	76	42	0	303	0	75	47	141
N.S.	1	1.01	0.94	0.52	0.00	3.74	0.00	0.93	0.58	1.74
time (sec)	N/A	0.390	0.162	0.107	0.000	0.104	0.000	0.133	0.247	2.931

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	42	38	102	0	164	0	68	95	56
N.S.	1	1.11	1.00	2.68	0.00	4.32	0.00	1.79	2.50	1.47
time (sec)	N/A	0.311	0.040	0.125	0.000	0.097	0.000	0.116	0.264	1.202

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	93	155	94	439	0	93	114	139
N.S.	1	1.00	3.21	5.34	3.24	15.14	0.00	3.21	3.93	4.79
time (sec)	N/A	0.274	0.048	0.091	0.037	0.095	0.000	0.113	0.238	0.190

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	110	197	140	1237	0	136	247	181
N.S.	1	1.00	2.39	4.28	3.04	26.89	0.00	2.96	5.37	3.93
time (sec)	N/A	0.439	0.083	0.111	0.055	0.109	0.000	0.114	0.242	1.275

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	94	70	230	186	2372	0	169	180	0
N.S.	1	1.29	0.96	3.15	2.55	32.49	0.00	2.32	2.47	0.00
time (sec)	N/A	0.779	0.246	0.178	0.045	0.112	0.000	0.121	0.253	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	96	117	240	0	0	0	0	0	93	0
N.S.	1	1.22	2.50	0.00	0.00	0.00	0.00	0.00	0.97	0.00
time (sec)	N/A	0.347	2.060	0.000	0.000	0.000	0.000	0.000	0.231	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	42	39	42	35	0	38	36	35
N.S.	1	1.00	2.62	2.44	2.62	2.19	0.00	2.38	2.25	2.19
time (sec)	N/A	0.198	0.060	0.200	0.147	0.084	0.000	0.118	0.228	0.106

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	27	17	26	45	52	0	41	44	27
N.S.	1	1.29	0.81	1.24	2.14	2.48	0.00	1.95	2.10	1.29
time (sec)	N/A	0.210	0.008	0.208	0.150	0.083	0.000	0.115	0.261	0.091

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	110	40	0	215	0	115	221	251
N.S.	1	1.00	1.55	0.56	0.00	3.03	0.00	1.62	3.11	3.54
time (sec)	N/A	0.263	0.035	0.205	0.000	0.098	0.000	0.155	0.236	1.253

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	80	57	101	0	182	0	118	9	100
N.S.	1	1.13	0.80	1.42	0.00	2.56	0.00	1.66	0.13	1.41
time (sec)	N/A	0.329	0.060	0.292	0.000	0.093	0.000	0.135	0.235	1.197

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	269	78	0	250	0	154	260	288
N.S.	1	1.00	3.16	0.92	0.00	2.94	0.00	1.81	3.06	3.39
time (sec)	N/A	0.320	0.070	0.319	0.000	0.104	0.000	0.125	0.263	1.443

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	148	49	87	0	46	60	65
N.S.	1	1.00	1.00	5.69	1.88	3.35	0.00	1.77	2.31	2.50
time (sec)	N/A	0.239	0.086	0.303	0.041	0.091	0.000	0.103	0.252	0.270

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	83	181	70	405	0	70	120	150
N.S.	1	1.00	2.37	5.17	2.00	11.57	0.00	2.00	3.43	4.29
time (sec)	N/A	0.299	0.070	1.047	0.124	0.108	0.000	0.113	0.243	1.429

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	35	42	120	246	0	51	53	0
N.S.	1	1.00	0.92	1.11	3.16	6.47	0.00	1.34	1.39	0.00
time (sec)	N/A	0.327	0.121	0.971	0.047	0.081	0.000	0.116	0.247	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	63	105	245	155	1881	0	125	287	0
N.S.	1	0.94	1.57	3.66	2.31	28.07	0.00	1.87	4.28	0.00
time (sec)	N/A	0.407	0.232	7.201	0.128	0.102	0.000	0.109	0.272	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	0	23	146	49	146	0	46	53	65
N.S.	1	0.00	1.00	6.35	2.13	6.35	0.00	2.00	2.30	2.83
time (sec)	N/A	0.000	0.094	0.180	0.045	0.098	0.000	0.109	0.236	0.244

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	C	A	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	0	81	171	69	722	0	69	136	150
N.S.	1	0.00	2.38	5.03	2.03	21.24	0.00	2.03	4.00	4.41
time (sec)	N/A	0.000	0.070	0.540	0.131	0.094	0.000	0.118	0.245	0.261

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	0	33	44	119	447	0	50	54	0
N.S.	1	0.00	0.89	1.19	3.22	12.08	0.00	1.35	1.46	0.00
time (sec)	N/A	0.000	0.113	0.504	0.042	0.100	0.000	0.105	0.222	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	0	103	235	152	6299	0	124	319	0
N.S.	1	0.00	1.58	3.62	2.34	96.91	0.00	1.91	4.91	0.00
time (sec)	N/A	0.000	0.259	3.194	0.156	0.165	0.000	0.110	0.253	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	0	50	259	74	630	0	71	98	194
N.S.	1	0.00	1.39	7.19	2.06	17.50	0.00	1.97	2.72	5.39
time (sec)	N/A	0.000	0.096	0.931	0.121	0.110	0.000	0.114	0.222	0.422

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	A	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	0	177	247	97	692	0	102	187	85
N.S.	1	0.00	3.61	5.04	1.98	14.12	0.00	2.08	3.82	1.73
time (sec)	N/A	0.000	0.362	3.602	0.041	0.097	0.000	0.116	0.250	1.291

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	0	109	321	156	2187	0	143	303	0
N.S.	1	0.00	1.24	3.65	1.77	24.85	0.00	1.62	3.44	0.00
time (sec)	N/A	0.000	0.327	9.082	0.126	0.104	0.000	0.109	0.279	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	0	80	58	323	493	0	87	91	0
N.S.	1	0.00	0.95	0.69	3.85	5.87	0.00	1.04	1.08	0.00
time (sec)	N/A	0.000	0.192	6.338	0.046	0.092	0.000	0.118	0.237	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	0	45	243	73	427	0	74	109	194
N.S.	1	0.00	1.41	7.59	2.28	13.34	0.00	2.31	3.41	6.06
time (sec)	N/A	0.000	0.089	0.571	0.124	0.097	0.000	0.107	0.225	1.314

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	A	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	0	182	238	101	1304	0	106	179	83
N.S.	1	0.00	4.14	5.41	2.30	29.64	0.00	2.41	4.07	1.89
time (sec)	N/A	0.000	0.364	1.849	0.042	0.117	0.000	0.114	0.236	0.285

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	0	110	311	152	3915	0	142	339	0
N.S.	1	0.00	1.28	3.62	1.77	45.52	0.00	1.65	3.94	0.00
time (sec)	N/A	0.000	0.332	4.699	0.132	0.123	0.000	0.123	0.270	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	0	83	64	334	1015	0	86	97	0
N.S.	1	0.00	1.04	0.80	4.18	12.69	0.00	1.08	1.21	0.00
time (sec)	N/A	0.000	0.191	2.942	0.043	0.104	0.000	0.116	0.231	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	6	7	6	0	3	4	3
N.S.	1	1.00	1.00	0.86	1.00	0.86	0.00	0.43	0.57	0.43
time (sec)	N/A	0.235	0.001	0.073	0.140	0.091	0.000	0.113	0.231	0.056

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	40	39	31	0	19	33	19
N.S.	1	1.00	1.00	2.67	2.60	2.07	0.00	1.27	2.20	1.27
time (sec)	N/A	0.215	0.023	0.083	0.111	0.081	0.000	0.110	0.242	0.075

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	62	50	76	0	44	45	42
N.S.	1	1.00	1.00	2.38	1.92	2.92	0.00	1.69	1.73	1.62
time (sec)	N/A	0.217	0.019	0.084	0.110	0.097	0.000	0.140	0.244	0.077

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	87	84	41	0	111	0	68	9	282
N.S.	1	1.09	1.05	0.51	0.00	1.39	0.00	0.85	0.11	3.52
time (sec)	N/A	0.304	0.099	0.103	0.000	0.096	0.000	0.143	0.222	3.731

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	38	30	92	0	107	0	58	65	41
N.S.	1	1.06	0.83	2.56	0.00	2.97	0.00	1.61	1.81	1.14
time (sec)	N/A	0.258	0.020	0.102	0.000	0.095	0.000	0.143	0.247	0.218

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	30	26	150	84	86	0	48	88	65
N.S.	1	1.15	1.00	5.77	3.23	3.31	0.00	1.85	3.38	2.50
time (sec)	N/A	0.262	0.086	0.102	0.038	0.092	0.000	0.127	0.267	0.289

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	90	172	103	617	0	109	208	156
N.S.	1	1.00	2.50	4.78	2.86	17.14	0.00	3.03	5.78	4.33
time (sec)	N/A	0.311	0.068	0.216	0.037	0.103	0.000	0.139	0.272	1.163

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	35	57	131	246	0	53	55	0
N.S.	1	1.00	0.90	1.46	3.36	6.31	0.00	1.36	1.41	0.00
time (sec)	N/A	0.336	0.129	0.444	0.039	0.082	0.000	0.140	0.239	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	70	65	235	185	2696	0	161	478	0
N.S.	1	1.04	0.97	3.51	2.76	40.24	0.00	2.40	7.13	0.00
time (sec)	N/A	0.425	0.296	1.305	0.038	0.130	0.000	0.123	0.254	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	0	26	149	81	146	0	49	83	66
N.S.	1	0.00	1.04	5.96	3.24	5.84	0.00	1.96	3.32	2.64
time (sec)	N/A	0.000	0.091	0.084	0.045	0.092	0.000	0.126	0.286	0.242

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	0	87	164	103	1084	0	107	218	156
N.S.	1	0.00	2.64	4.97	3.12	32.85	0.00	3.24	6.61	4.73
time (sec)	N/A	0.000	0.074	0.161	0.047	0.128	0.000	0.116	0.230	1.141

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	0	34	57	132	449	0	53	56	0
N.S.	1	0.00	0.92	1.54	3.57	12.14	0.00	1.43	1.51	0.00
time (sec)	N/A	0.000	0.130	0.263	0.039	0.088	0.000	0.113	0.244	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	0	61	229	187	8922	0	159	496	0
N.S.	1	0.00	0.94	3.52	2.88	137.26	0.00	2.45	7.63	0.00
time (sec)	N/A	0.000	0.293	0.757	0.041	0.156	0.000	0.138	0.236	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	0	50	233	116	880	0	119	170	202
N.S.	1	0.00	1.39	6.47	3.22	24.44	0.00	3.31	4.72	5.61
time (sec)	N/A	0.000	0.105	0.292	0.039	0.099	0.000	0.188	0.236	1.254

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	0	53	258	132	710	0	103	254	88
N.S.	1	0.00	1.08	5.27	2.69	14.49	0.00	2.10	5.18	1.80
time (sec)	N/A	0.000	0.491	0.480	0.043	0.103	0.000	0.140	0.243	0.290

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	0	246	305	207	3234	0	195	520	0
N.S.	1	0.00	2.76	3.43	2.33	36.34	0.00	2.19	5.84	0.00
time (sec)	N/A	0.000	1.556	0.970	0.038	0.109	0.000	0.164	0.228	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	0	80	60	331	497	0	85	95	0
N.S.	1	0.00	0.95	0.71	3.94	5.92	0.00	1.01	1.13	0.00
time (sec)	N/A	0.000	0.216	1.846	0.044	0.094	0.000	0.143	0.262	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	0	50	221	118	665	0	117	174	202
N.S.	1	0.00	1.52	6.70	3.58	20.15	0.00	3.55	5.27	6.12
time (sec)	N/A	0.000	0.099	0.227	0.033	0.111	0.000	0.112	0.238	0.383

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	0	49	249	134	1311	0	111	256	86
N.S.	1	0.00	1.11	5.66	3.05	29.80	0.00	2.52	5.82	1.95
time (sec)	N/A	0.000	0.496	0.318	0.036	0.119	0.000	0.121	0.234	0.291

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	0	162	295	204	5785	0	189	538	0
N.S.	1	0.00	1.91	3.47	2.40	68.06	0.00	2.22	6.33	0.00
time (sec)	N/A	0.000	4.462	0.634	0.053	0.133	0.000	0.116	0.218	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	0	81	64	342	1007	0	86	103	0
N.S.	1	0.00	1.01	0.80	4.28	12.59	0.00	1.08	1.29	0.00
time (sec)	N/A	0.000	0.211	0.834	0.037	0.094	0.000	0.199	0.234	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	23	15	12	27	19	20	25	17	6
N.S.	1	2.88	1.88	1.50	3.38	2.38	2.50	3.12	2.12	0.75
time (sec)	N/A	0.191	0.004	0.193	0.033	0.082	0.120	0.122	0.242	1.188

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	25	17	14	27	33	20	26	17	11
N.S.	1	1.47	1.00	0.82	1.59	1.94	1.18	1.53	1.00	0.65
time (sec)	N/A	0.191	0.004	0.195	0.029	0.078	0.119	0.176	0.238	0.069

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	25	17	14	27	36	20	27	17	14
N.S.	1	1.47	1.00	0.82	1.59	2.12	1.18	1.59	1.00	0.82
time (sec)	N/A	0.192	0.004	0.220	0.040	0.082	0.118	0.119	0.245	1.010

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	25	28	0	42	42	59	25	26
N.S.	1	1.00	0.71	0.80	0.00	1.20	1.20	1.69	0.71	0.74
time (sec)	N/A	0.216	0.024	0.241	0.000	0.085	0.238	0.116	0.244	0.101

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	27	17	20	25	17	9
N.S.	1	1.00	1.00	0.80	1.80	1.13	1.33	1.67	1.13	0.60
time (sec)	N/A	0.182	0.004	0.131	0.034	0.064	0.119	0.177	0.243	0.062

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	27	20	20	27	17	20
N.S.	1	1.00	1.00	0.82	1.59	1.18	1.18	1.59	1.00	1.18
time (sec)	N/A	0.186	0.005	0.154	0.039	0.079	0.124	0.136	0.238	0.055

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	27	34	20	27	17	15
N.S.	1	1.00	1.00	0.82	1.59	2.00	1.18	1.59	1.00	0.88
time (sec)	N/A	0.181	0.004	0.159	0.037	0.069	0.117	0.158	0.278	0.062

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	25	28	0	42	56	59	25	26
N.S.	1	1.00	0.71	0.80	0.00	1.20	1.60	1.69	0.71	0.74
time (sec)	N/A	0.210	0.023	0.203	0.000	0.078	0.258	0.116	0.238	0.072

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	26	24	58	89	58	66	50	23
N.S.	1	1.00	0.96	0.89	2.15	3.30	2.15	2.44	1.85	0.85
time (sec)	N/A	0.207	0.018	0.234	0.041	0.077	0.163	0.140	0.239	0.168

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	26	24	59	77	58	66	56	23
N.S.	1	1.00	0.96	0.89	2.19	2.85	2.15	2.44	2.07	0.85
time (sec)	N/A	0.205	0.017	0.220	0.041	0.087	0.164	0.181	0.219	1.094

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	40	0	71	153	85	42	42
N.S.	1	1.00	1.00	0.93	0.00	1.65	3.56	1.98	0.98	0.98
time (sec)	N/A	0.233	0.127	0.326	0.000	0.081	0.313	0.180	0.256	0.165

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	69	57	0	115	408	120	215	68
N.S.	1	1.00	1.11	0.92	0.00	1.85	6.58	1.94	3.47	1.10
time (sec)	N/A	0.263	0.493	1.373	0.000	0.096	0.702	0.140	0.203	1.183

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	85	84	0	217	921	179	574	180
N.S.	1	1.00	0.93	0.92	0.00	2.38	10.12	1.97	6.31	1.98
time (sec)	N/A	0.300	0.299	2.882	0.000	0.085	1.909	0.119	0.247	0.542

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	105	89	0	192	1027	156	357	115
N.S.	1	1.00	1.19	1.01	0.00	2.18	11.67	1.77	4.06	1.31
time (sec)	N/A	0.287	0.476	3.654	0.000	0.091	1.534	0.135	0.221	1.499

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	144	144	158	133	0	397	2009	260	865	337
N.S.	1	1.00	1.10	0.92	0.00	2.76	13.95	1.81	6.01	2.34
time (sec)	N/A	0.356	1.073	9.081	0.000	0.097	5.321	0.133	0.260	1.537

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	195	195	176	190	0	726	3580	373	19	908
N.S.	1	1.00	0.90	0.97	0.00	3.72	18.36	1.91	0.10	4.66
time (sec)	N/A	0.415	1.131	28.763	0.000	0.119	16.481	0.166	200.022	1.625

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	27	51	49	52	73	0	45	58	48
N.S.	1	1.42	2.68	2.58	2.74	3.84	0.00	2.37	3.05	2.53
time (sec)	N/A	0.220	0.062	0.099	0.119	0.083	0.000	0.178	0.229	0.067

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	55	49	153	82	0	45	58	53
N.S.	1	1.00	2.75	2.45	7.65	4.10	0.00	2.25	2.90	2.65
time (sec)	N/A	0.213	0.051	0.106	0.135	0.087	0.000	0.156	0.257	0.951

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	80	113	42	0	213	0	119	132	133
N.S.	1	1.07	1.51	0.56	0.00	2.84	0.00	1.59	1.76	1.77
time (sec)	N/A	0.306	0.033	0.125	0.000	0.101	0.000	0.192	0.243	0.079

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	82	249	42	0	221	0	127	51	141
N.S.	1	1.01	3.07	0.52	0.00	2.73	0.00	1.57	0.63	1.74
time (sec)	N/A	0.383	0.033	0.122	0.000	0.098	0.000	0.149	0.238	0.094

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	93	281	79	0	258	0	157	236	170
N.S.	1	1.07	3.23	0.91	0.00	2.97	0.00	1.80	2.71	1.95
time (sec)	N/A	0.394	0.059	0.138	0.000	0.099	0.000	0.183	0.247	0.986

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	86	167	59	327	0	53	68	133
N.S.	1	1.00	2.97	5.76	2.03	11.28	0.00	1.83	2.34	4.59
time (sec)	N/A	0.278	0.041	0.131	0.130	0.106	0.000	0.118	0.229	0.216

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	102	207	103	902	0	97	154	173
N.S.	1	1.00	2.27	4.60	2.29	20.04	0.00	2.16	3.42	3.84
time (sec)	N/A	0.430	0.075	0.172	0.122	0.104	0.000	0.156	0.250	1.116

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	86	115	238	149	1737	0	122	234	0
N.S.	1	1.19	1.60	3.31	2.07	24.12	0.00	1.69	3.25	0.00
time (sec)	N/A	0.738	0.230	0.249	0.125	0.112	0.000	0.161	0.235	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	99	120	103	0	0	0	0	0	91	0
N.S.	1	1.21	1.04	0.00	0.00	0.00	0.00	0.00	0.92	0.00
time (sec)	N/A	0.346	0.584	0.000	0.000	0.000	0.000	0.000	0.237	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	14	25	26	29	52	0	26	35	29
N.S.	1	1.40	2.50	2.60	2.90	5.20	0.00	2.60	3.50	2.90
time (sec)	N/A	0.209	0.020	0.084	0.037	0.077	0.000	0.123	0.234	0.898

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	55	47	50	57	104	0	55	71	57
N.S.	1	2.75	2.35	2.50	2.85	5.20	0.00	2.75	3.55	2.85
time (sec)	N/A	0.310	0.025	0.098	0.112	0.104	0.000	0.167	0.228	0.064

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	30	73	63	70	101	0	67	80	71
N.S.	1	1.07	2.61	2.25	2.50	3.61	0.00	2.39	2.86	2.54
time (sec)	N/A	0.278	0.062	0.112	0.109	0.111	0.000	0.149	0.218	0.076

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	110	133	190	0	272	0	157	54	143
N.S.	1	1.31	1.58	2.26	0.00	3.24	0.00	1.87	0.64	1.70
time (sec)	N/A	0.426	0.109	0.117	0.000	0.088	0.000	0.137	0.253	0.101

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	42	95	87	0	157	0	89	113	101
N.S.	1	1.11	2.50	2.29	0.00	4.13	0.00	2.34	2.97	2.66
time (sec)	N/A	0.315	0.060	0.148	0.000	0.090	0.000	0.118	0.233	0.962

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	62	76	75	0	0	0	0	0	58	0
N.S.	1	1.23	1.21	0.00	0.00	0.00	0.00	0.00	0.94	0.00
time (sec)	N/A	0.284	0.301	0.000	0.000	0.000	0.000	0.000	0.224	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	93	155	90	439	0	91	112	139
N.S.	1	1.00	3.21	5.34	3.10	15.14	0.00	3.14	3.86	4.79
time (sec)	N/A	0.273	0.045	0.131	0.044	0.107	0.000	0.133	0.221	0.182

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	110	195	144	1237	0	142	245	183
N.S.	1	1.00	2.39	4.24	3.13	26.89	0.00	3.09	5.33	3.98
time (sec)	N/A	0.426	0.075	0.141	0.044	0.106	0.000	0.180	0.237	1.069

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	94	70	228	184	2372	0	167	386	0
N.S.	1	1.29	0.96	3.12	2.52	32.49	0.00	2.29	5.29	0.00
time (sec)	N/A	0.781	0.234	0.203	0.051	0.108	0.000	0.122	0.212	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	95	116	99	0	0	0	0	0	93	0
N.S.	1	1.22	1.04	0.00	0.00	0.00	0.00	0.00	0.98	0.00
time (sec)	N/A	0.348	0.808	0.000	0.000	0.000	0.000	0.000	0.219	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	44	43	68	0	39	35	32
N.S.	1	1.00	1.00	2.93	2.87	4.53	0.00	2.60	2.33	2.13
time (sec)	N/A	0.206	0.007	0.256	0.116	0.102	0.000	0.111	0.246	0.106

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	40	114	31	0	19	27	19
N.S.	1	1.00	1.00	2.67	7.60	2.07	0.00	1.27	1.80	1.27
time (sec)	N/A	0.210	0.021	0.164	0.128	0.092	0.000	0.106	0.266	0.948

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	67	40	0	217	0	135	249	126
N.S.	1	1.00	0.94	0.56	0.00	3.06	0.00	1.90	3.51	1.77
time (sec)	N/A	0.255	0.072	0.198	0.000	0.103	0.000	0.160	0.240	2.367

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	89	84	41	0	115	0	68	9	297
N.S.	1	1.22	1.15	0.56	0.00	1.58	0.00	0.93	0.12	4.07
time (sec)	N/A	0.321	0.091	0.187	0.000	0.095	0.000	0.131	0.218	3.555

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	81	83	0	239	0	177	287	206
N.S.	1	1.00	0.95	0.98	0.00	2.81	0.00	2.08	3.38	2.42
time (sec)	N/A	0.298	0.058	0.209	0.000	0.103	0.000	0.178	0.291	3.109

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	146	51	86	0	47	61	64
N.S.	1	1.00	1.00	5.62	1.96	3.31	0.00	1.81	2.35	2.46
time (sec)	N/A	0.247	0.082	0.218	0.050	0.087	0.000	0.163	0.215	0.268

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	83	183	70	405	0	70	116	148
N.S.	1	1.00	2.37	5.23	2.00	11.57	0.00	2.00	3.31	4.23
time (sec)	N/A	0.305	0.067	0.519	0.118	0.098	0.000	0.161	0.212	1.170

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	35	26	119	248	0	49	53	0
N.S.	1	1.00	0.92	0.68	3.13	6.53	0.00	1.29	1.39	0.00
time (sec)	N/A	0.323	0.122	0.506	0.041	0.080	0.000	0.116	0.219	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	63	82	245	151	1881	0	123	287	0
N.S.	1	0.94	1.22	3.66	2.25	28.07	0.00	1.84	4.28	0.00
time (sec)	N/A	0.405	0.443	4.343	0.123	0.111	0.000	0.137	0.251	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	0	23	144	49	146	0	45	55	64
N.S.	1	0.00	1.00	6.26	2.13	6.35	0.00	1.96	2.39	2.78
time (sec)	N/A	0.000	0.079	0.177	0.046	0.091	0.000	0.113	0.225	0.230

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	C	A	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	0	144	175	69	713	0	71	132	148
N.S.	1	0.00	4.24	5.15	2.03	20.97	0.00	2.09	3.88	4.35
time (sec)	N/A	0.000	0.074	0.522	0.128	0.097	0.000	0.123	0.221	1.167

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	0	33	28	118	454	0	48	56	0
N.S.	1	0.00	0.89	0.76	3.19	12.27	0.00	1.30	1.51	0.00
time (sec)	N/A	0.000	0.117	0.503	0.048	0.099	0.000	0.138	0.242	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	0	74	239	152	6243	0	124	319	0
N.S.	1	0.00	1.14	3.68	2.34	96.05	0.00	1.91	4.91	0.00
time (sec)	N/A	0.000	0.417	3.205	0.127	0.158	0.000	0.180	0.234	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	0	48	251	74	634	0	71	98	194
N.S.	1	0.00	1.33	6.97	2.06	17.61	0.00	1.97	2.72	5.39
time (sec)	N/A	0.000	0.085	0.703	0.123	0.108	0.000	0.187	0.236	0.380

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	A	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	0	177	248	97	694	0	102	187	85
N.S.	1	0.00	3.61	5.06	1.98	14.16	0.00	2.08	3.82	1.73
time (sec)	N/A	0.000	0.363	3.069	0.044	0.108	0.000	0.127	0.271	1.216

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	0	108	321	156	2173	0	143	303	0
N.S.	1	0.00	1.23	3.65	1.77	24.69	0.00	1.62	3.44	0.00
time (sec)	N/A	0.000	0.326	4.220	0.128	0.116	0.000	0.120	0.294	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	0	80	56	321	493	0	85	93	0
N.S.	1	0.00	0.95	0.67	3.82	5.87	0.00	1.01	1.11	0.00
time (sec)	N/A	0.000	0.191	3.000	0.036	0.075	0.000	0.161	0.253	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	0	47	231	73	431	0	74	109	194
N.S.	1	0.00	1.47	7.22	2.28	13.47	0.00	2.31	3.41	6.06
time (sec)	N/A	0.000	0.094	0.556	0.120	0.114	0.000	0.117	0.251	0.357

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	A	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	0	182	239	101	1306	0	106	179	83
N.S.	1	0.00	4.14	5.43	2.30	29.68	0.00	2.41	4.07	1.89
time (sec)	N/A	0.000	0.349	1.629	0.041	0.110	0.000	0.144	0.319	0.283

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	C	A	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	0	110	311	152	3893	0	142	339	0
N.S.	1	0.00	1.28	3.62	1.77	45.27	0.00	1.65	3.94	0.00
time (sec)	N/A	0.000	0.363	4.594	0.132	0.127	0.000	0.120	0.263	0.000

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	0	81	62	332	1007	0	84	99	0
N.S.	1	0.00	1.01	0.78	4.15	12.59	0.00	1.05	1.24	0.00
time (sec)	N/A	0.000	0.192	3.532	0.043	0.096	0.000	0.121	0.226	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	19	19	0	16	17	19
N.S.	1	1.00	1.00	1.14	2.71	2.71	0.00	2.29	2.43	2.71
time (sec)	N/A	0.227	0.001	0.097	0.034	0.083	0.000	0.145	0.304	0.061

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	27	21	24	47	52	0	40	45	29
N.S.	1	1.29	1.00	1.14	2.24	2.48	0.00	1.90	2.14	1.38
time (sec)	N/A	0.215	0.008	0.108	0.119	0.081	0.000	0.119	0.260	0.071

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	67	53	60	54	0	57	54	61
N.S.	1	1.00	2.58	2.04	2.31	2.08	0.00	2.19	2.08	2.35
time (sec)	N/A	0.227	0.061	0.110	0.131	0.092	0.000	0.181	0.232	0.064

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	80	57	101	0	180	0	108	9	104
N.S.	1	1.13	0.80	1.42	0.00	2.54	0.00	1.52	0.13	1.46
time (sec)	N/A	0.334	0.075	0.116	0.000	0.088	0.000	0.162	0.229	1.064

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	37	91	77	0	101	0	79	82	91
N.S.	1	1.03	2.53	2.14	0.00	2.81	0.00	2.19	2.28	2.53
time (sec)	N/A	0.281	0.060	0.129	0.000	0.104	0.000	0.128	0.299	0.089

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	30	26	152	80	87	0	47	86	66
N.S.	1	1.15	1.00	5.85	3.08	3.35	0.00	1.81	3.31	2.54
time (sec)	N/A	0.251	0.080	0.145	0.041	0.096	0.000	0.137	0.217	0.232

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	90	170	105	617	0	111	208	156
N.S.	1	1.00	2.50	4.72	2.92	17.14	0.00	3.08	5.78	4.33
time (sec)	N/A	0.308	0.067	0.321	0.036	0.096	0.000	0.129	0.233	1.108

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	35	36	132	243	0	51	54	0
N.S.	1	1.00	0.90	0.92	3.38	6.23	0.00	1.31	1.38	0.00
time (sec)	N/A	0.332	0.129	0.690	0.036	0.091	0.000	0.147	0.291	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	70	65	235	191	2696	0	165	478	0
N.S.	1	1.15	1.07	3.85	3.13	44.20	0.00	2.70	7.84	0.00
time (sec)	N/A	0.435	0.290	2.040	0.051	0.103	0.000	0.171	0.230	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	0	28	147	81	146	0	48	87	65
N.S.	1	0.00	1.12	5.88	3.24	5.84	0.00	1.92	3.48	2.60
time (sec)	N/A	0.000	0.090	0.112	0.048	0.103	0.000	0.195	0.261	0.220

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	0	149	164	103	1104	0	105	218	156
N.S.	1	0.00	4.52	4.97	3.12	33.45	0.00	3.18	6.61	4.73
time (sec)	N/A	0.000	0.073	0.214	0.043	0.094	0.000	0.125	0.246	0.194

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	0	34	38	131	453	0	51	54	0
N.S.	1	0.00	0.92	1.03	3.54	12.24	0.00	1.38	1.46	0.00
time (sec)	N/A	0.000	0.138	0.355	0.036	0.092	0.000	0.125	0.221	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	0	61	229	187	9016	0	159	496	0
N.S.	1	0.00	1.00	3.75	3.07	147.80	0.00	2.61	8.13	0.00
time (sec)	N/A	0.000	0.304	1.145	0.048	0.156	0.000	0.188	0.228	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	0	50	233	116	872	0	119	170	202
N.S.	1	0.00	1.39	6.47	3.22	24.22	0.00	3.31	4.72	5.61
time (sec)	N/A	0.000	0.103	0.430	0.044	0.109	0.000	0.117	0.240	0.370

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	0	53	257	132	708	0	103	254	88
N.S.	1	0.00	1.08	5.24	2.69	14.45	0.00	2.10	5.18	1.80
time (sec)	N/A	0.000	0.472	0.891	0.050	0.109	0.000	0.145	0.238	0.292

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	0	249	305	207	3254	0	195	520	0
N.S.	1	0.00	2.80	3.43	2.33	36.56	0.00	2.19	5.84	0.00
time (sec)	N/A	0.000	6.189	2.234	0.047	0.117	0.000	0.141	0.229	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	0	82	58	333	497	0	87	93	0
N.S.	1	0.00	0.98	0.69	3.96	5.92	0.00	1.04	1.11	0.00
time (sec)	N/A	0.000	0.212	3.848	0.036	0.112	0.000	0.126	0.236	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	0	52	221	118	657	0	117	174	202
N.S.	1	0.00	1.58	6.70	3.58	19.91	0.00	3.55	5.27	6.12
time (sec)	N/A	0.000	0.105	0.273	0.044	0.125	0.000	0.121	0.241	0.363

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	0	51	248	134	1313	0	111	256	86
N.S.	1	0.00	1.16	5.64	3.05	29.84	0.00	2.52	5.82	1.95
time (sec)	N/A	0.000	0.483	0.541	0.047	0.100	0.000	0.153	0.230	0.290

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	B	B	B	F	B	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	0	164	295	204	5829	0	189	538	0
N.S.	1	0.00	1.93	3.47	2.40	68.58	0.00	2.22	6.33	0.00
time (sec)	N/A	0.000	4.319	1.273	0.046	0.169	0.000	0.147	0.271	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	B	B	F	A	B	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	0	81	62	344	1015	0	88	101	0
N.S.	1	0.00	1.01	0.78	4.30	12.69	0.00	1.10	1.26	0.00
time (sec)	N/A	0.000	0.216	1.528	0.044	0.102	0.000	0.132	0.244	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	46	29	151	83	259	0	95	115	115
N.S.	1	1.24	0.78	4.08	2.24	7.00	0.00	2.57	3.11	3.11
time (sec)	N/A	0.341	0.326	0.116	0.121	0.097	0.000	0.128	0.239	1.554

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	42	30	149	87	216	0	90	109	121
N.S.	1	1.17	0.83	4.14	2.42	6.00	0.00	2.50	3.03	3.36
time (sec)	N/A	0.335	0.314	0.113	0.128	0.098	0.000	0.191	0.227	0.597

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	53	29	155	157	259	0	97	169	115
N.S.	1	1.43	0.78	4.19	4.24	7.00	0.00	2.62	4.57	3.11
time (sec)	N/A	0.328	0.320	0.112	0.039	0.096	0.000	0.132	0.231	1.423

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	50	32	153	160	216	0	94	164	121
N.S.	1	1.39	0.89	4.25	4.44	6.00	0.00	2.61	4.56	3.36
time (sec)	N/A	0.328	0.307	0.110	0.052	0.121	0.000	0.192	0.225	1.426

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	27	77	68	184	0	82	54	266
N.S.	1	1.00	0.75	2.14	1.89	5.11	0.00	2.28	1.50	7.39
time (sec)	N/A	0.258	0.148	0.372	0.132	0.083	0.000	0.168	0.227	2.002

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	27	75	67	156	0	72	52	268
N.S.	1	1.00	0.82	2.27	2.03	4.73	0.00	2.18	1.58	8.12
time (sec)	N/A	0.249	0.141	0.237	0.141	0.083	0.000	0.119	0.227	1.901

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	44	28	79	133	184	0	84	70	266
N.S.	1	1.22	0.78	2.19	3.69	5.11	0.00	2.33	1.94	7.39
time (sec)	N/A	0.265	0.153	0.100	0.040	0.092	0.000	0.122	0.280	0.322

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	41	29	77	129	156	0	76	70	269
N.S.	1	1.24	0.88	2.33	3.91	4.73	0.00	2.30	2.12	8.15
time (sec)	N/A	0.258	0.136	0.074	0.048	0.092	0.000	0.163	0.227	1.249

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [27] had the largest ratio of [1.1428599999999999]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.88	7	0.429
2	A	3	3	1.00	7	0.429
3	A	3	3	1.00	7	0.429
4	A	2	2	1.00	7	0.286
5	A	2	2	1.00	13	0.154
6	A	2	2	1.00	14	0.143
7	A	2	2	1.00	13	0.154
8	A	2	2	1.00	15	0.133
9	A	2	2	1.00	15	0.133
10	A	2	2	1.00	17	0.118
11	A	2	2	1.00	17	0.118
12	A	2	2	1.00	17	0.118
13	C	3	3	1.53	7	0.429
14	C	3	3	1.47	7	0.429
15	C	3	3	1.47	7	0.429
16	A	2	2	1.00	7	0.286
17	A	2	2	1.00	13	0.154
18	A	2	2	1.00	15	0.133
19	A	2	2	1.00	15	0.133
20	A	2	2	1.00	15	0.133
21	A	2	2	1.00	17	0.118

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	2	2	1.00	17	0.118
23	A	2	2	1.00	15	0.133
24	A	2	2	1.00	17	0.118
25	A	2	2	1.00	17	0.118
26	A	7	6	1.42	7	0.857
27	A	9	8	1.00	7	1.143
28	A	9	8	1.07	7	1.143
29	A	7	6	1.04	7	0.857
30	A	7	6	1.07	7	0.857
31	A	2	2	1.21	7	0.286
32	A	4	4	1.00	13	0.308
33	A	11	10	1.00	15	0.667
34	A	16	15	1.19	15	1.000
35	A	2	2	1.21	13	0.154
36	A	6	5	1.40	7	0.714
37	A	5	4	1.00	7	0.571
38	A	6	5	1.07	7	0.714
39	A	5	4	1.01	7	0.571
40	A	6	5	1.11	7	0.714
41	A	5	5	1.00	13	0.385
42	A	10	9	1.00	15	0.600
43	C	18	17	1.29	15	1.133
44	A	2	2	1.22	13	0.154
45	A	5	4	1.00	7	0.571
46	A	9	8	1.29	7	1.143
47	A	6	5	1.00	7	0.714
48	A	7	6	1.13	7	0.857
49	A	6	5	1.00	7	0.714
50	A	5	5	1.00	13	0.385
51	A	7	6	1.00	15	0.400
52	A	8	7	1.00	15	0.467
53	A	9	8	0.94	15	0.533
Continued on next page						

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	F	0	0	N/A	0.000	N/A
55	F	0	0	N/A	0.000	N/A
56	F	0	0	N/A	0.000	N/A
57	F	0	0	N/A	0.000	N/A
58	F	0	0	N/A	0.000	N/A
59	F	0	0	N/A	0.000	N/A
60	F	0	0	N/A	0.000	N/A
61	F	0	0	N/A	0.000	N/A
62	F	0	0	N/A	0.000	N/A
63	F	0	0	N/A	0.000	N/A
64	F	0	0	N/A	0.000	N/A
65	F	0	0	N/A	0.000	N/A
66	A	4	4	1.00	7	0.571
67	A	4	3	1.00	7	0.429
68	A	5	4	1.00	7	0.571
69	A	5	4	1.09	7	0.571
70	A	6	5	1.06	7	0.714
71	C	5	5	1.15	13	0.385
72	A	7	6	1.00	15	0.400
73	A	9	8	1.00	15	0.533
74	C	12	11	1.04	15	0.733
75	F	0	0	N/A	0.000	N/A
76	F	0	0	N/A	0.000	N/A
77	F	0	0	N/A	0.000	N/A
78	F	0	0	N/A	0.000	N/A
79	F	0	0	N/A	0.000	N/A
80	F	0	0	N/A	0.000	N/A
81	F	0	0	N/A	0.000	N/A
82	F	0	0	N/A	0.000	N/A
83	F	0	0	N/A	0.000	N/A

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
84	F	0	0	N/A	0.000	N/A
85	F	0	0	N/A	0.000	N/A
86	F	0	0	N/A	0.000	N/A
87	C	3	3	2.88	7	0.429
88	C	3	3	1.47	7	0.429
89	C	3	3	1.47	7	0.429
90	A	2	2	1.00	7	0.286
91	A	2	2	1.00	7	0.286
92	A	2	2	1.00	7	0.286
93	A	2	2	1.00	7	0.286
94	A	2	2	1.00	7	0.286
95	A	2	2	1.00	13	0.154
96	A	2	2	1.00	14	0.143
97	A	2	2	1.00	13	0.154
98	A	2	2	1.00	15	0.133
99	A	2	2	1.00	15	0.133
100	A	2	2	1.00	17	0.118
101	A	2	2	1.00	17	0.118
102	A	2	2	1.00	17	0.118
103	A	7	6	1.42	7	0.857
104	A	6	5	1.00	7	0.714
105	A	9	8	1.07	7	1.143
106	A	6	5	1.01	7	0.714
107	A	7	6	1.07	7	0.857
108	A	5	5	1.00	13	0.385
109	A	10	9	1.00	15	0.600
110	A	17	16	1.19	15	1.067
111	A	2	2	1.21	13	0.154
112	A	7	6	1.40	7	0.857
113	B	9	8	2.75	7	1.143
114	A	7	6	1.07	7	0.857
115	A	6	5	1.31	7	0.714

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
116	A	7	6	1.11	7	0.857
117	A	2	2	1.23	7	0.286
118	A	5	5	1.00	13	0.385
119	A	10	9	1.00	15	0.600
120	C	18	17	1.29	15	1.133
121	A	2	2	1.22	13	0.154
122	A	4	3	1.00	7	0.429
123	A	4	3	1.00	7	0.429
124	A	5	4	1.00	7	0.571
125	A	5	4	1.22	7	0.571
126	A	5	4	1.00	7	0.571
127	A	5	5	1.00	13	0.385
128	A	7	6	1.00	15	0.400
129	A	8	7	1.00	15	0.467
130	A	9	8	0.94	15	0.533
131	F	0	0	N/A	0.000	N/A
132	F	0	0	N/A	0.000	N/A
133	F	0	0	N/A	0.000	N/A
134	F	0	0	N/A	0.000	N/A
135	F	0	0	N/A	0.000	N/A
136	F	0	0	N/A	0.000	N/A
137	F	0	0	N/A	0.000	N/A
138	F	0	0	N/A	0.000	N/A
139	F	0	0	N/A	0.000	N/A
140	F	0	0	N/A	0.000	N/A
141	F	0	0	N/A	0.000	N/A
142	F	0	0	N/A	0.000	N/A
143	A	7	7	1.00	7	1.000
144	A	9	8	1.29	7	1.143
145	A	6	5	1.00	7	0.714
146	A	8	7	1.13	7	1.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
147	A	7	6	1.03	7	0.857
148	C	5	5	1.15	13	0.385
149	A	7	6	1.00	15	0.400
150	A	9	8	1.00	15	0.533
151	C	12	11	1.15	15	0.733
152	F	0	0	N/A	0.000	N/A
153	F	0	0	N/A	0.000	N/A
154	F	0	0	N/A	0.000	N/A
155	F	0	0	N/A	0.000	N/A
156	F	0	0	N/A	0.000	N/A
157	F	0	0	N/A	0.000	N/A
158	F	0	0	N/A	0.000	N/A
159	F	0	0	N/A	0.000	N/A
160	F	0	0	N/A	0.000	N/A
161	F	0	0	N/A	0.000	N/A
162	F	0	0	N/A	0.000	N/A
163	F	0	0	N/A	0.000	N/A
164	A	5	5	1.24	13	0.385
165	A	5	5	1.17	14	0.357
166	C	5	5	1.43	13	0.385
167	C	5	5	1.39	14	0.357
168	A	4	4	1.00	13	0.308
169	A	4	4	1.00	14	0.286
170	C	4	4	1.22	13	0.308
171	C	4	4	1.24	14	0.286

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \sinh(x) \sinh(2x) dx$	87
3.2	$\int \sinh(x) \sinh(3x) dx$	92
3.3	$\int \sinh(x) \sinh(4x) dx$	97
3.4	$\int \sinh(x) \sinh(mx) dx$	102
3.5	$\int \sinh(a + bx) \sinh(c + bx) dx$	107
3.6	$\int \sinh(c - bx) \sinh(a + bx) dx$	113
3.7	$\int \sinh(a + bx) \sinh(c + dx) dx$	119
3.8	$\int \sinh(a + bx) \sinh^2(c + dx) dx$	125
3.9	$\int \sinh(a + bx) \sinh^3(c + dx) dx$	131
3.10	$\int \sinh^2(a + bx) \sinh^2(c + dx) dx$	138
3.11	$\int \sinh^2(a + bx) \sinh^3(c + dx) dx$	145
3.12	$\int \sinh^3(a + bx) \sinh^3(c + dx) dx$	154
3.13	$\int \cosh(2x) \sinh(x) dx$	163
3.14	$\int \cosh(3x) \sinh(x) dx$	168
3.15	$\int \cosh(4x) \sinh(x) dx$	173
3.16	$\int \cosh(mx) \sinh(x) dx$	178
3.17	$\int \cosh(c + dx) \sinh(a + bx) dx$	183
3.18	$\int \cosh^2(c + dx) \sinh(a + bx) dx$	189
3.19	$\int \cosh^3(c + dx) \sinh(a + bx) dx$	195
3.20	$\int \cosh(c + dx) \sinh^2(a + bx) dx$	202
3.21	$\int \cosh^2(c + dx) \sinh^2(a + bx) dx$	208
3.22	$\int \cosh^3(c + dx) \sinh^2(a + bx) dx$	215
3.23	$\int \cosh(c + dx) \sinh^3(a + bx) dx$	224
3.24	$\int \cosh^2(c + dx) \sinh^3(a + bx) dx$	231
3.25	$\int \cosh^3(c + dx) \sinh^3(a + bx) dx$	240
3.26	$\int \sinh(x) \tanh(2x) dx$	249
3.27	$\int \sinh(x) \tanh(3x) dx$	255

3.28	$\int \sinh(x) \tanh(4x) dx$	262
3.29	$\int \sinh(x) \tanh(5x) dx$	269
3.30	$\int \sinh(x) \tanh(6x) dx$	276
3.31	$\int \sinh(x) \tanh(nx) dx$	283
3.32	$\int \sinh(a + bx) \tanh(c + bx) dx$	288
3.33	$\int \sinh(a + bx) \tanh^2(c + bx) dx$	294
3.34	$\int \sinh(a + bx) \tanh^3(c + bx) dx$	302
3.35	$\int \sinh(a + bx) \tanh(c + dx) dx$	311
3.36	$\int \coth(2x) \sinh(x) dx$	316
3.37	$\int \coth(3x) \sinh(x) dx$	321
3.38	$\int \coth(4x) \sinh(x) dx$	327
3.39	$\int \coth(5x) \sinh(x) dx$	333
3.40	$\int \coth(6x) \sinh(x) dx$	340
3.41	$\int \coth(c + bx) \sinh(a + bx) dx$	346
3.42	$\int \coth^2(c + bx) \sinh(a + bx) dx$	353
3.43	$\int \coth^3(c + bx) \sinh(a + bx) dx$	361
3.44	$\int \coth(c + dx) \sinh(a + bx) dx$	370
3.45	$\int \operatorname{sech}(2x) \sinh(x) dx$	375
3.46	$\int \operatorname{sech}(3x) \sinh(x) dx$	381
3.47	$\int \operatorname{sech}(4x) \sinh(x) dx$	387
3.48	$\int \operatorname{sech}(5x) \sinh(x) dx$	396
3.49	$\int \operatorname{sech}(6x) \sinh(x) dx$	403
3.50	$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx$	413
3.51	$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx$	419
3.52	$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx$	426
3.53	$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx$	432
3.54	$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx$	440
3.55	$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx$	445
3.56	$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx$	451
3.57	$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx$	456
3.58	$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx$	461
3.59	$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx$	466
3.60	$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx$	472
3.61	$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx$	478
3.62	$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx$	484
3.63	$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx$	489
3.64	$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx$	495
3.65	$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx$	500
3.66	$\int \operatorname{csch}(2x) \sinh(x) dx$	506

3.67	$\int \operatorname{csch}(3x) \sinh(x) dx$	511
3.68	$\int \operatorname{csch}(4x) \sinh(x) dx$	516
3.69	$\int \operatorname{csch}(5x) \sinh(x) dx$	522
3.70	$\int \operatorname{csch}(6x) \sinh(x) dx$	529
3.71	$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx$	535
3.72	$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx$	541
3.73	$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx$	548
3.74	$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$	555
3.75	$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx$	564
3.76	$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx$	569
3.77	$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx$	575
3.78	$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$	580
3.79	$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$	586
3.80	$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx$	592
3.81	$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx$	598
3.82	$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx$	604
3.83	$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$	610
3.84	$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx$	616
3.85	$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx$	622
3.86	$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx$	628
3.87	$\int \cosh(x) \sinh(2x) dx$	634
3.88	$\int \cosh(x) \sinh(3x) dx$	639
3.89	$\int \cosh(x) \sinh(4x) dx$	644
3.90	$\int \cosh(x) \sinh(mx) dx$	649
3.91	$\int \cosh(x) \cosh(2x) dx$	654
3.92	$\int \cosh(x) \cosh(3x) dx$	659
3.93	$\int \cosh(x) \cosh(4x) dx$	664
3.94	$\int \cosh(x) \cosh(mx) dx$	669
3.95	$\int \cosh(a + bx) \cosh(c + bx) dx$	674
3.96	$\int \cosh(c - bx) \cosh(a + bx) dx$	680
3.97	$\int \cosh(a + bx) \cosh(c + dx) dx$	686
3.98	$\int \cosh(a + bx) \cosh^2(c + dx) dx$	692
3.99	$\int \cosh(a + bx) \cosh^3(c + dx) dx$	698
3.100	$\int \cosh^2(a + bx) \cosh^2(c + dx) dx$	705
3.101	$\int \cosh^2(a + bx) \cosh^3(c + dx) dx$	712
3.102	$\int \cosh^3(a + bx) \cosh^3(c + dx) dx$	721
3.103	$\int \cosh(x) \tanh(2x) dx$	730
3.104	$\int \cosh(x) \tanh(3x) dx$	736
3.105	$\int \cosh(x) \tanh(4x) dx$	742

3.106	$\int \cosh(x) \tanh(5x) dx$	750
3.107	$\int \cosh(x) \tanh(6x) dx$	757
3.108	$\int \cosh(a + bx) \tanh(c + bx) dx$	765
3.109	$\int \cosh(a + bx) \tanh^2(c + bx) dx$	771
3.110	$\int \cosh(a + bx) \tanh^3(c + bx) dx$	779
3.111	$\int \cosh(a + bx) \tanh(c + dx) dx$	788
3.112	$\int \cosh(x) \coth(2x) dx$	793
3.113	$\int \cosh(x) \coth(3x) dx$	799
3.114	$\int \cosh(x) \coth(4x) dx$	806
3.115	$\int \cosh(x) \coth(5x) dx$	813
3.116	$\int \cosh(x) \coth(6x) dx$	821
3.117	$\int \cosh(x) \coth(nx) dx$	828
3.118	$\int \cosh(a + bx) \coth(c + bx) dx$	833
3.119	$\int \cosh(a + bx) \coth^2(c + bx) dx$	840
3.120	$\int \cosh(a + bx) \coth^3(c + bx) dx$	848
3.121	$\int \cosh(a + bx) \coth(c + dx) dx$	858
3.122	$\int \cosh(x) \operatorname{sech}(2x) dx$	863
3.123	$\int \cosh(x) \operatorname{sech}(3x) dx$	869
3.124	$\int \cosh(x) \operatorname{sech}(4x) dx$	874
3.125	$\int \cosh(x) \operatorname{sech}(5x) dx$	881
3.126	$\int \cosh(x) \operatorname{sech}(6x) dx$	888
3.127	$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx$	896
3.128	$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx$	902
3.129	$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx$	909
3.130	$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx$	915
3.131	$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx$	923
3.132	$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx$	928
3.133	$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx$	934
3.134	$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx$	939
3.135	$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx$	944
3.136	$\int \cosh^2(a + bx) \operatorname{sech}^2(c + bx) dx$	949
3.137	$\int \cosh^2(a + bx) \operatorname{sech}^3(c + bx) dx$	955
3.138	$\int \cosh^2(a + bx) \operatorname{sech}^4(c + bx) dx$	961
3.139	$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx$	967
3.140	$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx$	972
3.141	$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx$	978
3.142	$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx$	983
3.143	$\int \cosh(x) \operatorname{csch}(2x) dx$	989
3.144	$\int \cosh(x) \operatorname{csch}(3x) dx$	994

3.145	$\int \cosh(x)\operatorname{csch}(4x) dx$	1000
3.146	$\int \cosh(x)\operatorname{csch}(5x) dx$	1006
3.147	$\int \cosh(x)\operatorname{csch}(6x) dx$	1013
3.148	$\int \cosh(a + bx)\operatorname{csch}(c + bx) dx$	1020
3.149	$\int \cosh(a + bx)\operatorname{csch}^2(c + bx) dx$	1026
3.150	$\int \cosh(a + bx)\operatorname{csch}^3(c + bx) dx$	1033
3.151	$\int \cosh(a + bx)\operatorname{csch}^4(c + bx) dx$	1040
3.152	$\int \cosh(a + bx)\operatorname{csch}(c - bx) dx$	1049
3.153	$\int \cosh(a + bx)\operatorname{csch}^2(c - bx) dx$	1054
3.154	$\int \cosh(a + bx)\operatorname{csch}^3(c - bx) dx$	1060
3.155	$\int \cosh(a + bx)\operatorname{csch}^4(c - bx) dx$	1065
3.156	$\int \cosh^2(a + bx)\operatorname{csch}(c + bx) dx$	1071
3.157	$\int \cosh^2(a + bx)\operatorname{csch}^2(c + bx) dx$	1077
3.158	$\int \cosh^2(a + bx)\operatorname{csch}^3(c + bx) dx$	1083
3.159	$\int \cosh^2(a + bx)\operatorname{csch}^4(c + bx) dx$	1089
3.160	$\int \cosh^2(a + bx)\operatorname{csch}(c - bx) dx$	1095
3.161	$\int \cosh^2(a + bx)\operatorname{csch}^2(c - bx) dx$	1101
3.162	$\int \cosh^2(a + bx)\operatorname{csch}^3(c - bx) dx$	1107
3.163	$\int \cosh^2(a + bx)\operatorname{csch}^4(c - bx) dx$	1113
3.164	$\int \tanh(a + bx)\tanh(c + bx) dx$	1119
3.165	$\int \tanh(c - bx)\tanh(a + bx) dx$	1125
3.166	$\int \coth(a + bx)\coth(c + bx) dx$	1131
3.167	$\int \coth(c - bx)\coth(a + bx) dx$	1138
3.168	$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx$	1145
3.169	$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx$	1151
3.170	$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx$	1157
3.171	$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx$	1163

3.1 $\int \sinh(x) \sinh(2x) dx$

Optimal result	87
Mathematica [A] (verified)	87
Rubi [A] (verified)	88
Maple [A] (verified)	89
Fricas [B] (verification not implemented)	89
Sympy [B] (verification not implemented)	90
Maxima [B] (verification not implemented)	90
Giac [B] (verification not implemented)	90
Mupad [B] (verification not implemented)	91
Reduce [B] (verification not implemented)	91

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \sinh(x) \sinh(2x) dx = \frac{2 \sinh^3(x)}{3}$$

output

```
2/3*sinh(x)^3
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \sinh(x) \sinh(2x) dx = -\frac{\sinh(x)}{2} + \frac{1}{6} \sinh(3x)$$

input

```
Integrate[Sinh[x]*Sinh[2*x],x]
```

output

```
-1/2*Sinh[x] + Sinh[3*x]/6
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 25, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \sinh(2x) dx \\ & \quad \downarrow \text{3042} \\ & \int -\sin(ix) \sin(2ix) dx \\ & \quad \downarrow \text{25} \\ & -\int \sin(ix) \sin(2ix) dx \\ & \quad \downarrow \text{4770} \\ & \frac{1}{6} \sinh(3x) - \frac{\sinh(x)}{2} \end{aligned}$$

input `Int[Sinh[x]*Sinh[2*x],x]`

output `-1/2*Sinh[x] + Sinh[3*x]/6`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770

```
Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[
a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)
), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

method	result	size
default	$-\frac{\sinh(x)}{2} + \frac{\sinh(3x)}{6}$	12
parallelrisc	$-\frac{\sinh(x)}{2} + \frac{\sinh(3x)}{6}$	12
orering	$-\frac{\cosh(x)\sinh(2x)}{3} + \frac{2\cosh(2x)\sinh(x)}{3}$	18
risc	$\frac{e^{3x}}{12} - \frac{e^x}{4} + \frac{e^{-x}}{4} - \frac{e^{-3x}}{12}$	24

input

```
int(sinh(x)*sinh(2*x),x,method=_RETURNVERBOSE)
```

output

```
-1/2*sinh(x)+1/6*sinh(3*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. 2(6) = 12.

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \sinh(x) \sinh(2x) dx = \frac{1}{6} \sinh(x)^3 + \frac{1}{2} (\cosh(x)^2 - 1) \sinh(x)$$

input

```
integrate(sinh(x)*sinh(2*x),x, algorithm="fricas")
```

output

```
1/6*sinh(x)^3 + 1/2*(cosh(x)^2 - 1)*sinh(x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(7) = 14$.

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \sinh(x) \sinh(2x) dx = \frac{2 \sinh(x) \cosh(2x)}{3} - \frac{\sinh(2x) \cosh(x)}{3}$$

input `integrate(sinh(x)*sinh(2*x),x)`

output `2*sinh(x)*cosh(2*x)/3 - sinh(2*x)*cosh(x)/3`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(6) = 12$.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 3.38

$$\int \sinh(x) \sinh(2x) dx = -\frac{1}{12} (3e^{-2x} - 1)e^{3x} + \frac{1}{4} e^{-x} - \frac{1}{12} e^{-3x}$$

input `integrate(sinh(x)*sinh(2*x),x, algorithm="maxima")`

output `-1/12*(3*e^(-2*x) - 1)*e^(3*x) + 1/4*e^(-x) - 1/12*e^(-3*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(6) = 12$.

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 3.12

$$\int \sinh(x) \sinh(2x) dx = \frac{1}{12} (3e^{2x} - 1)e^{-3x} + \frac{1}{12} e^{3x} - \frac{1}{4} e^x$$

input `integrate(sinh(x)*sinh(2*x),x, algorithm="giac")`

output $1/12*(3*e^{(2*x)} - 1)*e^{(-3*x)} + 1/12*e^{(3*x)} - 1/4*e^x$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \sinh(x) \sinh(2x) dx = \frac{2 \sinh(x)^3}{3}$$

input `int(sinh(2*x)*sinh(x),x)`

output $(2*\sinh(x)^3)/3$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \sinh(x) \sinh(2x) dx = \frac{2 \cosh(2x) \sinh(x)}{3} - \frac{\cosh(x) \sinh(2x)}{3}$$

input `int(sinh(x)*sinh(2*x),x)`

output $(2*\cosh(2*x)*\sinh(x) - \cosh(x)*\sinh(2*x))/3$

3.2 $\int \sinh(x) \sinh(3x) dx$

Optimal result	92
Mathematica [A] (verified)	92
Rubi [A] (verified)	93
Maple [A] (verified)	94
Fricas [A] (verification not implemented)	94
Sympy [A] (verification not implemented)	95
Maxima [B] (verification not implemented)	95
Giac [B] (verification not implemented)	95
Mupad [B] (verification not implemented)	96
Reduce [B] (verification not implemented)	96

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \sinh(x) \sinh(3x) dx = -\frac{1}{4} \sinh(2x) + \frac{1}{8} \sinh(4x)$$

output `-1/4*sinh(2*x)+1/8*sinh(4*x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sinh(x) \sinh(3x) dx = -\frac{1}{4} \sinh(2x) + \frac{1}{8} \sinh(4x)$$

input `Integrate[Sinh[x]*Sinh[3*x],x]`

output `-1/4*Sinh[2*x] + Sinh[4*x]/8`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 25, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \sinh(3x) dx \\ & \quad \downarrow 3042 \\ & \int -\sin(ix) \sin(3ix) dx \\ & \quad \downarrow 25 \\ & -\int \sin(ix) \sin(3ix) dx \\ & \quad \downarrow 4770 \\ & \frac{1}{8} \sinh(4x) - \frac{1}{4} \sinh(2x) \end{aligned}$$

input `Int[Sinh[x]*Sinh[3*x],x]`

output `-1/4*Sinh[2*x] + Sinh[4*x]/8`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770

```
Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[
a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)
), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\sinh(2x)}{4} + \frac{\sinh(4x)}{8}$	14
parallelrisc	$-\frac{\sinh(2x)}{4} + \frac{\sinh(4x)}{8}$	14
orering	$-\frac{\cosh(x)\sinh(3x)}{8} + \frac{3\cosh(3x)\sinh(x)}{8}$	18
risch	$\frac{e^{4x}}{16} - \frac{e^{2x}}{8} + \frac{e^{-2x}}{8} - \frac{e^{-4x}}{16}$	26

input

```
int(sinh(x)*sinh(3*x),x,method=_RETURNVERBOSE)
```

output

```
-1/4*sinh(2*x)+1/8*sinh(4*x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \sinh(x) \sinh(3x) dx = \frac{1}{2} \cosh(x) \sinh(x)^3 + \frac{1}{2} (\cosh(x)^3 - \cosh(x)) \sinh(x)$$

input

```
integrate(sinh(x)*sinh(3*x),x, algorithm="fricas")
```

output

```
1/2*cosh(x)*sinh(x)^3 + 1/2*(cosh(x)^3 - cosh(x))*sinh(x)
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sinh(x) \sinh(3x) dx = \frac{3 \sinh(x) \cosh(3x)}{8} - \frac{\sinh(3x) \cosh(x)}{8}$$

input `integrate(sinh(x)*sinh(3*x),x)`

output `3*sinh(x)*cosh(3*x)/8 - sinh(3*x)*cosh(x)/8`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sinh(x) \sinh(3x) dx = -\frac{1}{16} (2e^{(-2x)} - 1)e^{(4x)} + \frac{1}{8} e^{(-2x)} - \frac{1}{16} e^{(-4x)}$$

input `integrate(sinh(x)*sinh(3*x),x, algorithm="maxima")`

output `-1/16*(2*e^(-2*x) - 1)*e^(4*x) + 1/8*e^(-2*x) - 1/16*e^(-4*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sinh(x) \sinh(3x) dx = \frac{1}{16} (2e^{(2x)} - 1)e^{(-4x)} + \frac{1}{16} e^{(4x)} - \frac{1}{8} e^{(2x)}$$

input `integrate(sinh(x)*sinh(3*x),x, algorithm="giac")`

output `1/16*(2*e^(2*x) - 1)*e^(-4*x) + 1/16*e^(4*x) - 1/8*e^(2*x)`

Mupad [B] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \sinh(x) \sinh(3x) dx = \frac{\sinh(4x)}{8} - \frac{\sinh(2x)}{4}$$

input `int(sinh(3*x)*sinh(x),x)`

output `sinh(4*x)/8 - sinh(2*x)/4`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sinh(x) \sinh(3x) dx = \frac{3 \cosh(3x) \sinh(x)}{8} - \frac{\cosh(x) \sinh(3x)}{8}$$

input `int(sinh(x)*sinh(3*x),x)`

output `(3*cosh(3*x)*sinh(x) - cosh(x)*sinh(3*x))/8`

3.3 $\int \sinh(x) \sinh(4x) dx$

Optimal result	97
Mathematica [A] (verified)	97
Rubi [A] (verified)	98
Maple [A] (verified)	99
Fricas [B] (verification not implemented)	99
Sympy [A] (verification not implemented)	100
Maxima [B] (verification not implemented)	100
Giac [B] (verification not implemented)	100
Mupad [B] (verification not implemented)	101
Reduce [B] (verification not implemented)	101

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \sinh(x) \sinh(4x) dx = -\frac{1}{6} \sinh(3x) + \frac{1}{10} \sinh(5x)$$

output `-1/6*sinh(3*x)+1/10*sinh(5*x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sinh(x) \sinh(4x) dx = -\frac{1}{6} \sinh(3x) + \frac{1}{10} \sinh(5x)$$

input `Integrate[Sinh[x]*Sinh[4*x],x]`

output `-1/6*Sinh[3*x] + Sinh[5*x]/10`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 25, 4770}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \sinh(4x) dx \\ & \quad \downarrow \text{3042} \\ & \int -\sin(ix) \sin(4ix) dx \\ & \quad \downarrow \text{25} \\ & -\int \sin(ix) \sin(4ix) dx \\ & \quad \downarrow \text{4770} \\ & \frac{1}{10} \sinh(5x) - \frac{1}{6} \sinh(3x) \end{aligned}$$

input `Int[Sinh[x]*Sinh[4*x],x]`

output `-1/6*Sinh[3*x] + Sinh[5*x]/10`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4770

```
Int[sin[(a_.) + (b_.)*(x_)]*sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[
a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Sin[a + c + (b + d)*x]/(2*(b + d)
), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\sinh(3x)}{6} + \frac{\sinh(5x)}{10}$	14
parallelsch	$-\frac{\sinh(3x)}{6} + \frac{\sinh(5x)}{10}$	14
orering	$-\frac{\cosh(x)\sinh(4x)}{15} + \frac{4\cosh(4x)\sinh(x)}{15}$	18
risch	$\frac{e^{5x}}{20} - \frac{e^{3x}}{12} + \frac{e^{-3x}}{12} - \frac{e^{-5x}}{20}$	26

input

```
int(sinh(x)*sinh(4*x),x,method=_RETURNVERBOSE)
```

output

```
-1/6*sinh(3*x)+1/10*sinh(5*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.12

$$\int \sinh(x) \sinh(4x) dx = \frac{1}{10} \sinh(x)^5 + \frac{1}{6} (6 \cosh(x)^2 - 1) \sinh(x)^3 + \frac{1}{2} (\cosh(x)^4 - \cosh(x)^2) \sinh(x)$$

input

```
integrate(sinh(x)*sinh(4*x),x, algorithm="fricas")
```

output

```
1/10*sinh(x)^5 + 1/6*(6*cosh(x)^2 - 1)*sinh(x)^3 + 1/2*(cosh(x)^4 - cosh(x)
)^2)*sinh(x)
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \sinh(x) \sinh(4x) dx = \frac{4 \sinh(x) \cosh(4x)}{15} - \frac{\sinh(4x) \cosh(x)}{15}$$

input `integrate(sinh(x)*sinh(4*x),x)`

output `4*sinh(x)*cosh(4*x)/15 - sinh(4*x)*cosh(x)/15`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sinh(x) \sinh(4x) dx = -\frac{1}{60} (5 e^{-2x} - 3) e^{5x} + \frac{1}{12} e^{-3x} - \frac{1}{20} e^{-5x}$$

input `integrate(sinh(x)*sinh(4*x),x, algorithm="maxima")`

output `-1/60*(5*e^(-2*x) - 3)*e^(5*x) + 1/12*e^(-3*x) - 1/20*e^(-5*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \sinh(x) \sinh(4x) dx = \frac{1}{60} (5 e^{2x} - 3) e^{-5x} + \frac{1}{20} e^{5x} - \frac{1}{12} e^{3x}$$

input `integrate(sinh(x)*sinh(4*x),x, algorithm="giac")`

output `1/60*(5*e^(2*x) - 3)*e^(-5*x) + 1/20*e^(5*x) - 1/12*e^(3*x)`

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \sinh(x) \sinh(4x) dx = \frac{4 \sinh(x)^3 (6 \sinh(x)^2 + 5)}{15}$$

input `int(sinh(4*x)*sinh(x),x)`

output `(4*sinh(x)^3*(6*sinh(x)^2 + 5))/15`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \sinh(x) \sinh(4x) dx = \frac{4 \cosh(4x) \sinh(x)}{15} - \frac{\cosh(x) \sinh(4x)}{15}$$

input `int(sinh(x)*sinh(4*x),x)`

output `(4*cosh(4*x)*sinh(x) - cosh(x)*sinh(4*x))/15`

3.4 $\int \sinh(x) \sinh(mx) dx$

Optimal result	102
Mathematica [A] (verified)	102
Rubi [A] (verified)	103
Maple [A] (verified)	104
Fricas [A] (verification not implemented)	104
Sympy [B] (verification not implemented)	105
Maxima [F(-2)]	105
Giac [B] (verification not implemented)	106
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 7, antiderivative size = 35

$$\int \sinh(x) \sinh(mx) dx = -\frac{\sinh((1-m)x)}{2(1-m)} + \frac{\sinh((1+m)x)}{2(1+m)}$$

output

```
-1/2*sinh((1-m)*x)/(1-m)+sinh((1+m)*x)/(2+2*m)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \sinh(x) \sinh(mx) dx = \frac{m \cosh(mx) \sinh(x) - \cosh(x) \sinh(mx)}{-1 + m^2}$$

input

```
Integrate[Sinh[x]*Sinh[m*x],x]
```

output

```
(m*Cosh[m*x]*Sinh[x] - Cosh[x]*Sinh[m*x])/(-1 + m^2)
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \sinh(mx) dx$$

$$\downarrow 6147$$

$$\int \left(\frac{1}{2} \cosh((m+1)x) - \frac{1}{2} \cosh((1-m)x) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh((m+1)x)}{2(m+1)} - \frac{\sinh((1-m)x)}{2(1-m)}$$

input `Int[Sinh[x]*Sinh[m*x],x]`

output `-1/2*Sinh[(1-m)*x]/(1-m) + Sinh[(1+m)*x]/(2*(1+m))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6147 `Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$-\frac{\sinh(x(-1+m))}{2(-1+m)} + \frac{\sinh((1+m)x)}{2+2m}$
parallelrisch	$\frac{(-1-m)\sinh(x(-1+m))+\sinh((1+m)x)(-1+m)}{2m^2-2}$
risch	$\frac{(m e^{2x}-e^{2x}-m-1)e^{x(-1+m)}}{4(1+m)(-1+m)} + \frac{(m e^{2x}+e^{2x}-m+1)e^{-(1+m)x}}{4(1+m)(-1+m)}$
orering	$\frac{2(m^2+1)(\cosh(x)\sinh(mx)+\sinh(x)m\cosh(mx))}{m^4-2m^2+1} - \frac{\cosh(x)\sinh(mx)+3\sinh(x)m\cosh(mx)+3\cosh(x)m^2\sinh(mx)+\sinh(x)m^3\cosh(mx)}{m^4-2m^2+1}$

input `int(sinh(x)*sinh(m*x),x,method=_RETURNVERBOSE)`output `-1/2/(-1+m)*sinh(x*(-1+m))+1/2/(1+m)*sinh((1+m)*x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \sinh(x) \sinh(mx) dx = \frac{m \cosh(mx) \sinh(x) - \cosh(x) \sinh(mx)}{(m^2 - 1) \cosh(x)^2 - (m^2 - 1) \sinh(x)^2}$$

input `integrate(sinh(x)*sinh(m*x),x, algorithm="fricas")`output `(m*cosh(m*x)*sinh(x) - cosh(x)*sinh(m*x))/((m^2 - 1)*cosh(x)^2 - (m^2 - 1)*sinh(x)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. $2(22) = 44$.

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.23

$$\int \sinh(x) \sinh(mx) dx = \begin{cases} -\frac{x \sinh^2(x)}{2} + \frac{x \cosh^2(x)}{2} - \frac{\sinh(x) \cosh(x)}{2} & \text{for } m = -1 \\ \frac{x \sinh^2(x)}{2} - \frac{x \cosh^2(x)}{2} + \frac{\sinh(x) \cosh(x)}{2} & \text{for } m = 1 \\ \frac{m \sinh(x) \cosh(mx)}{m^2-1} - \frac{\sinh(mx) \cosh(x)}{m^2-1} & \text{otherwise} \end{cases}$$

input `integrate(sinh(x)*sinh(m*x),x)`

output `Piecewise((-x*sinh(x)**2/2 + x*cosh(x)**2/2 - sinh(x)*cosh(x)/2, Eq(m, -1)), (x*sinh(x)**2/2 - x*cosh(x)**2/2 + sinh(x)*cosh(x)/2, Eq(m, 1)), (m*sinh(x)*cosh(m*x)/(m**2 - 1) - sinh(m*x)*cosh(x)/(m**2 - 1), True))`

Maxima [F(-2)]

Exception generated.

$$\int \sinh(x) \sinh(mx) dx = \text{Exception raised: ValueError}$$

input `integrate(sinh(x)*sinh(m*x),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see `assume?` for more details)Is`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \sinh(x) \sinh(mx) dx = \frac{e^{(mx+x)}}{4(m+1)} - \frac{e^{(mx-x)}}{4(m-1)} + \frac{e^{(-mx+x)}}{4(m-1)} - \frac{e^{(-mx-x)}}{4(m+1)}$$

input `integrate(sinh(x)*sinh(m*x),x, algorithm="giac")`

output `1/4*e^(m*x + x)/(m + 1) - 1/4*e^(m*x - x)/(m - 1) + 1/4*e^(-m*x + x)/(m - 1) - 1/4*e^(-m*x - x)/(m + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sinh(x) \sinh(mx) dx = -\frac{\sinh(mx) \cosh(x) - m \cosh(mx) \sinh(x)}{m^2 - 1}$$

input `int(sinh(m*x)*sinh(x),x)`

output `-(sinh(m*x)*cosh(x) - m*cosh(m*x)*sinh(x))/(m^2 - 1)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \sinh(x) \sinh(mx) dx = \frac{\cosh(mx) \sinh(x) m - \cosh(x) \sinh(mx)}{m^2 - 1}$$

input `int(sinh(x)*sinh(m*x),x)`

output `(cosh(m*x)*sinh(x)*m - cosh(x)*sinh(m*x))/(m**2 - 1)`

3.5 $\int \sinh(a + bx) \sinh(c + bx) dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [A] (verified)	109
Fricas [B] (verification not implemented)	109
Sympy [B] (verification not implemented)	110
Maxima [B] (verification not implemented)	110
Giac [B] (verification not implemented)	111
Mupad [B] (verification not implemented)	111
Reduce [B] (verification not implemented)	111

Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \sinh(a + bx) \sinh(c + bx) dx = -\frac{1}{2}x \cosh(a - c) + \frac{\sinh(a + c + 2bx)}{4b}$$

output `-1/2*x*cosh(a-c)+1/4*sinh(2*b*x+a+c)/b`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \sinh(a + bx) \sinh(c + bx) dx = \frac{-2bx \cosh(a - c) + \sinh(a + c + 2bx)}{4b}$$

input `Integrate[Sinh[a + b*x]*Sinh[c + b*x],x]`

output `(-2*b*x*Cosh[a - c] + Sinh[a + c + 2*b*x])/(4*b)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \sinh(bx + c) dx$$

$$\downarrow \text{6147}$$

$$\int \left(\frac{1}{2} \cosh(a + 2bx + c) - \frac{1}{2} \cosh(a - c) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sinh(a + 2bx + c)}{4b} - \frac{1}{2} x \cosh(a - c)$$

input

```
Int[Sinh[a + b*x]*Sinh[c + b*x],x]
```

output

```
-1/2*(x*Cosh[a - c]) + Sinh[a + c + 2*b*x]/(4*b)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6147

```
Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]^(p)*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result
default	$-\frac{x \cosh(a-c)}{2} + \frac{\sinh(2bx+a+c)}{4b}$
parallelrisch	$\frac{-2bx \cosh(a-c) + \sinh(2bx+a+c) - \sinh(a-c)}{4b}$
risch	$-\frac{x e^{a-c}}{4} - \frac{x e^{-a+c}}{4} + \frac{e^{2bx+a+c}}{8b} - \frac{e^{-2bx-a-c}}{8b}$
orering	$x \sinh(bx+a) \sinh(bx+c) + \frac{b \cosh(bx+a) \sinh(bx+c) + \sinh(bx+a) b \cosh(bx+c)}{4b^2} - \frac{x(2 \sinh(bx+c) \sinh(bx+a))}{4b^2}$

input `int(sinh(b*x+a)*sinh(b*x+c),x,method=_RETURNVERBOSE)`

output `-1/2*x*cosh(a-c)+1/4*sinh(2*b*x+a+c)/b`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(23) = 46.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.22

$$\int \sinh(a+bx) \sinh(c+bx) dx = \frac{2bx \cosh(-a+c) - 2 \cosh(bx+c) \cosh(-a+c) \sinh(bx+c) + \cosh(bx+c)^2 \sinh(-a+c) + \sinh(bx+c)^2 \sinh(-a+c)}{4(b \cosh(-a+c)^2 - b \sinh(-a+c)^2)}$$

input `integrate(sinh(b*x+a)*sinh(b*x+c),x, algorithm="fricas")`

output `-1/4*(2*b*x*cosh(-a+c) - 2*cosh(b*x+c)*cosh(-a+c)*sinh(b*x+c) + cosh(b*x+c)^2*sinh(-a+c) + sinh(b*x+c)^2*sinh(-a+c))/(b*cosh(-a+c)^2 - b*sinh(-a+c)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(20) = 40$.

Time = 0.19 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \sinh(a + bx) \sinh(c + bx) dx = \begin{cases} \frac{x \sinh(a+bx) \sinh(bx+c)}{2} - \frac{x \cosh(a+bx) \cosh(bx+c)}{2} + \frac{\sinh(bx+c) \cosh(a+bx)}{2b} & \text{for } b \neq 0 \\ x \sinh(a) \sinh(c) & \text{otherwise} \end{cases}$$

input `integrate(sinh(b*x+a)*sinh(b*x+c),x)`

output `Piecewise((x*sinh(a + b*x)*sinh(b*x + c)/2 - x*cosh(a + b*x)*cosh(b*x + c)/2 + sinh(b*x + c)*cosh(a + b*x)/(2*b), Ne(b, 0)), (x*sinh(a)*sinh(c), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(23) = 46$.

Time = 0.05 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \sinh(a + bx) \sinh(c + bx) dx = -\frac{(bx + a)(e^{2a} + e^{2c})e^{-a-c}}{4b} + \frac{e^{2bx+a+c}}{8b} - \frac{e^{-2bx-a-c}}{8b}$$

input `integrate(sinh(b*x+a)*sinh(b*x+c),x, algorithm="maxima")`

output `-1/4*(b*x + a)*(e^(2*a) + e^(2*c))*e^(-a - c)/b + 1/8*e^(2*b*x + a + c)/b - 1/8*e^(-2*b*x - a - c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(23) = 46$.

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.52

$$\int \sinh(a + bx) \sinh(c + bx) dx$$

$$= -\frac{(2bx(e^{2a} + e^{2c}) - (e^{2bx+2a} + e^{2bx+2c} - 1))e^{-2bx} - e^{(2bx+2a+2c)}e^{-a-c}}{8b}$$

input `integrate(sinh(b*x+a)*sinh(b*x+c),x, algorithm="giac")`

output `-1/8*(2*b*x*(e^(2*a) + e^(2*c)) - (e^(2*b*x + 2*a) + e^(2*b*x + 2*c) - 1)*
e^(-2*b*x) - e^(2*b*x + 2*a + 2*c))*e^(-a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \sinh(a + bx) \sinh(c + bx) dx = \frac{\sinh(a + c + 2bx)}{4b} - \frac{x \cosh(a - c)}{2}$$

input `int(sinh(a + b*x)*sinh(c + b*x),x)`

output `sinh(a + c + 2*b*x)/(4*b) - (x*cosh(a - c))/2`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.85

$$\int \sinh(a + bx) \sinh(c + bx) dx$$

$$= \frac{-\cosh(bx + c) \cosh(bx + a) bx + \cosh(bx + c) \sinh(bx + a) + \sinh(bx + c) \sinh(bx + a) bx}{2b}$$

input `int(sinh(b*x+a)*sinh(b*x+c),x)`

output `(- cosh(b*x + c)*cosh(a + b*x)*b*x + cosh(b*x + c)*sinh(a + b*x) + sinh(b*x + c)*sinh(a + b*x)*b*x)/(2*b)`

3.6 $\int \sinh(c - bx) \sinh(a + bx) dx$

Optimal result	113
Mathematica [A] (verified)	113
Rubi [A] (verified)	114
Maple [A] (verified)	115
Fricas [B] (verification not implemented)	115
Sympy [A] (verification not implemented)	116
Maxima [B] (verification not implemented)	116
Giac [B] (verification not implemented)	117
Mupad [B] (verification not implemented)	117
Reduce [B] (verification not implemented)	117

Optimal result

Integrand size = 14, antiderivative size = 27

$$\int \sinh(c - bx) \sinh(a + bx) dx = \frac{1}{2}x \cosh(a + c) - \frac{\sinh(a - c + 2bx)}{4b}$$

output `1/2*x*cosh(a+c)-1/4*sinh(2*b*x+a-c)/b`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \sinh(c - bx) \sinh(a + bx) dx = \frac{1}{2}x \cosh(a + c) - \frac{\sinh(a - c + 2bx)}{4b}$$

input `Integrate[Sinh[c - b*x]*Sinh[a + b*x],x]`

output `(x*Cosh[a + c])/2 - Sinh[a - c + 2*b*x]/(4*b)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \sinh(c - bx) dx$$

$$\downarrow \text{6147}$$

$$\int \left(\frac{1}{2} \cosh(a + c) - \frac{1}{2} \cosh(a + 2bx - c) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{1}{2}x \cosh(a + c) - \frac{\sinh(a + 2bx - c)}{4b}$$

input `Int[Sinh[c - b*x]*Sinh[a + b*x],x]`

output `(x*Cosh[a + c])/2 - Sinh[a - c + 2*b*x]/(4*b)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6147 `Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] :> Int[ExpandTrigReduce[Sinh[v]^p*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result
default	$\frac{x \cosh(a+c)}{2} - \frac{\sinh(2bx+a-c)}{4b}$
parallelrisc	$-\frac{-2bx \cosh(a+c) + \sinh(2bx+a-c) - \sinh(a+c)}{4b}$
risch	$\frac{x e^{a+c}}{4} + \frac{x e^{-a-c}}{4} - \frac{e^{2bx+a-c}}{8b} + \frac{e^{-2bx-a+c}}{8b}$
orering	$-x \sinh(bx - c) \sinh(bx + a) + \frac{-b \cosh(bx-c) \sinh(bx+a) - \sinh(bx-c) b \cosh(bx+a)}{4b^2} - \frac{x(-2 \sinh(bx+a) \sinh(a+c))}{4b^2}$

input `int(-sinh(b*x-c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*x*cosh(a+c)-1/4*sinh(2*b*x+a-c)/b`

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(23) = 46$.

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.78

$$\int \sinh(c - bx) \sinh(a + bx) dx$$

$$= \frac{2bx \cosh(a+c) - 2 \cosh(bx+a) \cosh(a+c) \sinh(bx+a) + \cosh(bx+a)^2 \sinh(a+c) + \sinh(bx+a)^2 \sinh(a+c)}{4(b \cosh(a+c)^2 - b \sinh(a+c)^2)}$$

input `integrate(-sinh(b*x-c)*sinh(b*x+a),x, algorithm="fricas")`

output `1/4*(2*b*x*cosh(a+c) - 2*cosh(b*x+a)*cosh(a+c)*sinh(b*x+a) + cosh(b*x+a)^2*sinh(a+c) + sinh(b*x+a)^2*sinh(a+c))/(b*cosh(a+c)^2 - b*sinh(a+c)^2)`

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.26

$$\int \sinh(c - bx) \sinh(a + bx) dx$$

$$= - \begin{cases} \frac{x \sinh(a + bx) \sinh(bx - c)}{2} - \frac{x \cosh(a + bx) \cosh(bx - c)}{2} + \frac{\sinh(bx - c) \cosh(a + bx)}{2b} & \text{for } b \neq 0 \\ -x \sinh(a) \sinh(c) & \text{otherwise} \end{cases}$$

input `integrate(-sinh(b*x-c)*sinh(b*x+a),x)`

output `-Piecewise((x*sinh(a + b*x)*sinh(b*x - c)/2 - x*cosh(a + b*x)*cosh(b*x - c)/2 + sinh(b*x - c)*cosh(a + b*x)/(2*b), Ne(b, 0)), (-x*sinh(a)*sinh(c), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(23) = 46.

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.19

$$\int \sinh(c - bx) \sinh(a + bx) dx = \frac{(bx + a)(e^{(2a+2c)} + 1)e^{(-a-c)}}{4b} - \frac{e^{(2bx+a-c)}}{8b} + \frac{e^{(-2bx-a+c)}}{8b}$$

input `integrate(-sinh(b*x-c)*sinh(b*x+a),x, algorithm="maxima")`

output `1/4*(b*x + a)*(e^(2*a + 2*c) + 1)*e^(-a - c)/b - 1/8*e^(2*b*x + a - c)/b + 1/8*e^(-2*b*x - a + c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(23) = 46$.

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.59

$$\int \sinh(c - bx) \sinh(a + bx) dx$$

$$= \frac{(2bx(e^{2a+2c} + 1) - (e^{2bx} + e^{2bx+2a+2c} - e^{2c}))e^{(-2bx)} - e^{(2bx+2a)}e^{(-a-c)}}{8b}$$

input `integrate(-sinh(b*x-c)*sinh(b*x+a),x, algorithm="giac")`

output `1/8*(2*b*x*(e^(2*a + 2*c) + 1) - (e^(2*b*x) + e^(2*b*x + 2*a + 2*c) - e^(2*c))*e^(-2*b*x) - e^(2*b*x + 2*a))*e^(-a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \sinh(c - bx) \sinh(a + bx) dx = \frac{x \cosh(a + c)}{2} - \frac{\sinh(a - c + 2bx)}{4b}$$

input `int(sinh(a + b*x)*sinh(c - b*x),x)`

output `(x*cosh(a + c))/2 - sinh(a - c + 2*b*x)/(4*b)`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int \sinh(c - bx) \sinh(a + bx) dx$$

$$= \frac{\cosh(bx - c) \cosh(bx + a) bx - \cosh(bx - c) \sinh(bx + a) - \sinh(bx - c) \sinh(bx + a) bx}{2b}$$

input `int(-sinh(b*x-c)*sinh(b*x+a),x)`

output `(cosh(b*x - c)*cosh(a + b*x)*b*x - cosh(b*x - c)*sinh(a + b*x) - sinh(b*x - c)*sinh(a + b*x)*b*x)/(2*b)`

3.7 $\int \sinh(a + bx) \sinh(c + dx) dx$

Optimal result	119
Mathematica [A] (verified)	119
Rubi [A] (verified)	120
Maple [A] (verified)	121
Fricas [A] (verification not implemented)	121
Sympy [B] (verification not implemented)	122
Maxima [F(-2)]	122
Giac [B] (verification not implemented)	123
Mupad [B] (verification not implemented)	123
Reduce [B] (verification not implemented)	124

Optimal result

Integrand size = 13, antiderivative size = 43

$$\int \sinh(a + bx) \sinh(c + dx) dx = -\frac{\sinh(a - c + (b - d)x)}{2(b - d)} + \frac{\sinh(a + c + (b + d)x)}{2(b + d)}$$

output `-1/2*sinh(a-c+(b-d)*x)/(b-d)+sinh(a+c+(b+d)*x)/(2*b+2*d)`

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \sinh(a + bx) \sinh(c + dx) dx = -\frac{\sinh(a - c + (b - d)x)}{2(b - d)} + \frac{\sinh(a + c + (b + d)x)}{2(b + d)}$$

input `Integrate[Sinh[a + b*x]*Sinh[c + d*x],x]`

output `-1/2*Sinh[a - c + (b - d)*x]/(b - d) + Sinh[a + c + (b + d)*x]/(2*(b + d))`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \sinh(c + dx) dx$$

$$\downarrow 6147$$

$$\int \left(\frac{1}{2} \cosh(a + x(b + d) + c) - \frac{1}{2} \cosh(a + x(b - d) - c) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh(a + x(b + d) + c)}{2(b + d)} - \frac{\sinh(a + x(b - d) - c)}{2(b - d)}$$

input `Int[Sinh[a + b*x]*Sinh[c + d*x],x]`

output `-1/2*Sinh[a - c + (b - d)*x]/(b - d) + Sinh[a + c + (b + d)*x]/(2*(b + d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6147 `Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

method	result
default	$-\frac{\sinh(a-c+(b-d)x)}{2(b-d)} + \frac{\sinh(a+c+(b+d)x)}{2b+2d}$
parallelrisch	$\frac{(-b-d)\sinh(a-c+(b-d)x)+\sinh(a+c+(b+d)x)(b-d)}{2b^2-2d^2}$
risch	$\frac{(be^{2bx+2a}-de^{2bx+2a}+b+d)e^{-bx+dx-a+c}}{4(b+d)(b-d)} - \frac{(be^{2bx+2a}+de^{2bx+2a}+b-d)e^{-bx-dx-a-c}}{4(b+d)(b-d)}$
orering	$\frac{2(b^2+d^2)(b\cosh(bx+a)\sinh(dx+c)+\sinh(bx+a)d\cosh(dx+c))}{b^4-2b^2d^2+d^4} - \frac{b^3\cosh(bx+a)\sinh(dx+c)+3b^2\sinh(bx+a)d\cosh(dx+c)}{b^4-2b^2d^2+d^4}$

input `int(sinh(b*x+a)*sinh(d*x+c),x,method=_RETURNVERBOSE)`output `-1/2*sinh(a-c+(b-d)*x)/(b-d)+1/2/(b+d)*sinh(a+c+(b+d)*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.67

$$\int \sinh(a+bx)\sinh(c+dx)dx$$

$$= -\frac{d\cosh(dx+c)\sinh(bx+a)-b\cosh(bx+a)\sinh(dx+c)}{(b^2-d^2)\cosh(bx+a)^2-(b^2-d^2)\sinh(bx+a)^2}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c),x,algorithm="fricas")`output `-(d*cosh(d*x + c)*sinh(b*x + a) - b*cosh(b*x + a)*sinh(d*x + c))/((b^2 - d^2)*cosh(b*x + a)^2 - (b^2 - d^2)*sinh(b*x + a)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(32) = 64$.

Time = 0.32 (sec) , antiderivative size = 153, normalized size of antiderivative = 3.56

$$\int \sinh(a + bx) \sinh(c + dx) dx$$

$$= \begin{cases} x \sinh(a) \sinh(c) & \text{for } b = 0 \wedge d = 0 \\ \frac{x \sinh(a-dx) \sinh(c+dx)}{2} + \frac{x \cosh(a-dx) \cosh(c+dx)}{2} - \frac{\sinh(c+dx) \cosh(a-dx)}{2d} & \text{for } b = -d \\ \frac{x \sinh(a+dx) \sinh(c+dx)}{2} - \frac{x \cosh(a+dx) \cosh(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(a+dx)}{2d} & \text{for } b = d \\ \frac{b \sinh(c+dx) \cosh(a+bx)}{b^2-d^2} - \frac{d \sinh(a+bx) \cosh(c+dx)}{b^2-d^2} & \text{otherwise} \end{cases}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c),x)`

output `Piecewise((x*sinh(a)*sinh(c), Eq(b, 0) & Eq(d, 0)), (x*sinh(a - d*x)*sinh(c + d*x)/2 + x*cosh(a - d*x)*cosh(c + d*x)/2 - sinh(c + d*x)*cosh(a - d*x)/(2*d), Eq(b, -d)), (x*sinh(a + d*x)*sinh(c + d*x)/2 - x*cosh(a + d*x)*cosh(c + d*x)/2 + sinh(c + d*x)*cosh(a + d*x)/(2*d), Eq(b, d)), (b*sinh(c + d*x)*cosh(a + b*x)/(b**2 - d**2) - d*sinh(a + b*x)*cosh(c + d*x)/(b**2 - d**2), True))`

Maxima [F(-2)]

Exception generated.

$$\int \sinh(a + bx) \sinh(c + dx) dx = \text{Exception raised: ValueError}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-d/b>0)', see `assume?` for more details)I`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(39) = 78.

Time = 0.11 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.98

$$\int \sinh(a + bx) \sinh(c + dx) dx = \frac{e^{(bx+dx+a+c)}}{4(b+d)} - \frac{e^{(bx-dx+a-c)}}{4(b-d)} + \frac{e^{(-bx+dx-a+c)}}{4(b-d)} - \frac{e^{(-bx-dx-a-c)}}{4(b+d)}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c),x, algorithm="giac")`

output `1/4*e^(b*x + d*x + a + c)/(b + d) - 1/4*e^(b*x - d*x + a - c)/(b - d) + 1/4*e^(-b*x + d*x - a + c)/(b - d) - 1/4*e^(-b*x - d*x - a - c)/(b + d)`

Mupad [B] (verification not implemented)

Time = 1.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \sinh(a + bx) \sinh(c + dx) dx = \frac{b \cosh(a + bx) \sinh(c + dx) - d \cosh(c + dx) \sinh(a + bx)}{b^2 - d^2}$$

input `int(sinh(a + b*x)*sinh(c + d*x),x)`

output `(b*cosh(a + b*x)*sinh(c + d*x) - d*cosh(c + d*x)*sinh(a + b*x))/(b^2 - d^2)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \sinh(a + bx) \sinh(c + dx) dx$$
$$= \frac{\cosh(bx + a) \sinh(dx + c) b - \cosh(dx + c) \sinh(bx + a) d}{b^2 - d^2}$$

input `int(sinh(b*x+a)*sinh(d*x+c),x)`

output `(cosh(a + b*x)*sinh(c + d*x)*b - cosh(c + d*x)*sinh(a + b*x)*d)/(b**2 - d**2)`

3.8 $\int \sinh(a + bx) \sinh^2(c + dx) dx$

Optimal result	125
Mathematica [A] (verified)	125
Rubi [A] (verified)	126
Maple [A] (verified)	127
Fricas [B] (verification not implemented)	127
Sympy [B] (verification not implemented)	128
Maxima [F(-2)]	129
Giac [B] (verification not implemented)	129
Mupad [B] (verification not implemented)	130
Reduce [B] (verification not implemented)	130

Optimal result

Integrand size = 15, antiderivative size = 62

$$\int \sinh(a + bx) \sinh^2(c + dx) dx = -\frac{\cosh(a + bx)}{2b} + \frac{\cosh(a - 2c + (b - 2d)x)}{4(b - 2d)} + \frac{\cosh(a + 2c + (b + 2d)x)}{4(b + 2d)}$$

output

```
-1/2*cosh(b*x+a)/b+cosh(a-2*c+(b-2*d)*x)/(4*b-8*d)+cosh(a+2*c+(b+2*d)*x)/(4*b+8*d)
```

Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

$$\int \sinh(a + bx) \sinh^2(c + dx) dx = \frac{1}{4} \left(-\frac{2 \cosh(a) \cosh(bx)}{b} + \frac{\cosh(a - 2c + bx - 2dx)}{b - 2d} + \frac{\cosh(a + 2c + bx + 2dx)}{b + 2d} - \frac{2 \sinh(a) \sinh(bx)}{b} \right)$$

input

```
Integrate[Sinh[a + b*x]*Sinh[c + d*x]^2,x]
```

output

$$\left(\frac{-2 \operatorname{Cosh}[a] \operatorname{Cosh}[b*x]}{b} + \frac{\operatorname{Cosh}[a - 2*c + b*x - 2*d*x]}{(b - 2*d)} + \frac{\operatorname{Cosh}[a + 2*c + b*x + 2*d*x]}{(b + 2*d)} - \frac{(2 \operatorname{Sinh}[a] \operatorname{Sinh}[b*x])}{b} \right) / 4$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \sinh^2(c + dx) dx$$

$$\downarrow 6147$$

$$\int \left(\frac{1}{4} \sinh(a + x(b - 2d) - 2c) + \frac{1}{4} \sinh(a + x(b + 2d) + 2c) - \frac{1}{2} \sinh(a + bx) \right) dx$$

$$\downarrow 2009$$

$$\frac{\cosh(a + x(b - 2d) - 2c)}{4(b - 2d)} + \frac{\cosh(a + x(b + 2d) + 2c)}{4(b + 2d)} - \frac{\cosh(a + bx)}{2b}$$

input

```
Int[Sinh[a + b*x]*Sinh[c + d*x]^2,x]
```

output

$$\frac{-1/2 \operatorname{Cosh}[a + b*x]}{b} + \frac{\operatorname{Cosh}[a - 2*c + (b - 2*d)*x]}{(4*(b - 2*d))} + \frac{\operatorname{Cosh}[a + 2*c + (b + 2*d)*x]}{(4*(b + 2*d))}$$

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6147

```
Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] :> Int[ExpandTrigReduce[Sinh[v]
]^(p)*Sinh[w]^(q), x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

method	result
default	$-\frac{\cosh(bx+a)}{2b} + \frac{\cosh(a-2c+(b-2d)x)}{4b-8d} + \frac{\cosh(a+2c+(b+2d)x)}{4b+8d}$
parallelrisc	$\frac{b(b+2d)\cosh(a-2c+(b-2d)x)+b(b-2d)\cosh(a+2c+(b+2d)x)+(-2b^2+8d^2)\cosh(bx+a)-8d^2}{4b^3-16bd^2}$
risc	$-\frac{e^{bx+a}}{4b} - \frac{e^{-bx-a}}{4b} + \frac{(be^{2bx+2a}-2de^{2bx+2a+b+2d})e^{-bx+2dx-a+2c}}{8(b+2d)(b-2d)} + \frac{(be^{2bx+2a}+2de^{2bx+2a+b-2d})e^{-bx-2dx-a-2c}}{8(b+2d)(b-2d)}$
orering	$\frac{(3b^4+16d^4)(b\cosh(bx+a)\sinh(dx+c)^2+2\sinh(bx+a)\sinh(dx+c)d\cosh(dx+c))}{b^2(b^4-8b^2d^2+16d^4)} - \frac{(3b^2+8d^2)(b^3\cosh(bx+a)\sinh(dx+c))}{b^2(b^4-8b^2d^2+16d^4)}$

input

```
int(sinh(b*x+a)*sinh(d*x+c)^2,x,method=_RETURNVERBOSE)
```

output

```
-1/2*cosh(b*x+a)/b+1/4/(b-2*d)*cosh(a-2*c+(b-2*d)*x)+1/4/(b+2*d)*cosh(a+2*
c+(b+2*d)*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.94

$$\int \sinh(a + bx) \sinh^2(c + dx) dx$$

$$= \frac{b^2 \cosh(bx + a) \cosh(dx + c)^2 - 4bd \cosh(dx + c) \sinh(bx + a) \sinh(dx + c) + b^2 \cosh(bx + a) \sinh(dx + c)}{2((b^3 - 4bd^2) \cosh(bx + a)^2 - (b^3 - 4bd^2) \sinh(bx + a)^2)}$$

input

```
integrate(sinh(b*x+a)*sinh(d*x+c)^2,x, algorithm="fricas")
```


output

```
1/2*(b^2*cosh(b*x + a)*cosh(d*x + c)^2 - 4*b*d*cosh(d*x + c)*sinh(b*x + a)
*sinh(d*x + c) + b^2*cosh(b*x + a)*sinh(d*x + c)^2 - (b^2 - 4*d^2)*cosh(b*
x + a))/((b^3 - 4*b*d^2)*cosh(b*x + a)^2 - (b^3 - 4*b*d^2)*sinh(b*x + a)^2
)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(49) = 98$.

Time = 0.69 (sec) , antiderivative size = 405, normalized size of antiderivative = 6.53

$$\int \sinh(a + bx) \sinh^2(c + dx) dx$$

$$= \begin{cases} x \sinh(a) \sinh^2(c) \\ \left(\frac{x \sinh^2(c+dx)}{2} - \frac{x \cosh^2(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right) \sinh(a) \\ \frac{x \sinh(a-2dx) \sinh^2(c+dx)}{4} + \frac{x \sinh(a-2dx) \cosh^2(c+dx)}{4} + \frac{x \sinh(c+dx) \cosh(a-2dx) \cosh(c+dx)}{2} - \frac{\sinh(a-2dx) \sinh(c+dx) \cosh(c+dx)}{4d} \\ \frac{x \sinh(a+2dx) \sinh^2(c+dx)}{4} + \frac{x \sinh(a+2dx) \cosh^2(c+dx)}{4} - \frac{x \sinh(c+dx) \cosh(a+2dx) \cosh(c+dx)}{2} - \frac{\sinh(a+2dx) \sinh(c+dx) \cosh(c+dx)}{4d} \\ \frac{b^2 \sinh^2(c+dx) \cosh(a+bx)}{b^3-4bd^2} - \frac{2bd \sinh(a+bx) \sinh(c+dx) \cosh(c+dx)}{b^3-4bd^2} - \frac{2d^2 \sinh^2(c+dx) \cosh(a+bx)}{b^3-4bd^2} + \frac{2d^2 \cosh(a+bx) \cosh^2(c+dx)}{b^3-4bd^2} \end{cases}$$

input

```
integrate(sinh(b*x+a)*sinh(d*x+c)**2,x)
```

output

```
Piecewise((x*sinh(a)*sinh(c)**2, Eq(b, 0) & Eq(d, 0)), ((x*sinh(c + d*x)**
2/2 - x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*sinh(a), E
q(b, 0)), (x*sinh(a - 2*d*x)*sinh(c + d*x)**2/4 + x*sinh(a - 2*d*x)*cosh(c
+ d*x)**2/4 + x*sinh(c + d*x)*cosh(a - 2*d*x)*cosh(c + d*x)/2 - sinh(a -
2*d*x)*sinh(c + d*x)*cosh(c + d*x)/(4*d) - sinh(c + d*x)**2*cosh(a - 2*d*x
)/(2*d), Eq(b, -2*d)), (x*sinh(a + 2*d*x)*sinh(c + d*x)**2/4 + x*sinh(a +
2*d*x)*cosh(c + d*x)**2/4 - x*sinh(c + d*x)*cosh(a + 2*d*x)*cosh(c + d*x)/
2 - sinh(a + 2*d*x)*sinh(c + d*x)*cosh(c + d*x)/(4*d) + sinh(c + d*x)**2*c
osh(a + 2*d*x)/(2*d), Eq(b, 2*d)), (b**2*sinh(c + d*x)**2*cosh(a + b*x)/(b
**3 - 4*b*d**2) - 2*b*d*sinh(a + b*x)*sinh(c + d*x)*cosh(c + d*x)/(b**3 -
4*b*d**2) - 2*d**2*sinh(c + d*x)**2*cosh(a + b*x)/(b**3 - 4*b*d**2) + 2*d*
**2*cosh(a + b*x)*cosh(c + d*x)**2/(b**3 - 4*b*d**2), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \sinh(a + bx) \sinh^2(c + dx) dx = \text{Exception raised: ValueError}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c)^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(2*d)/b>0)', see `assume?` for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 0.12 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.94

$$\int \sinh(a + bx) \sinh^2(c + dx) dx = \frac{e^{(bx+2dx+a+2c)}}{8(b+2d)} + \frac{e^{(bx-2dx+a-2c)}}{8(b-2d)} - \frac{e^{(bx+a)}}{4b} \\ + \frac{e^{(-bx+2dx-a+2c)}}{8(b-2d)} + \frac{e^{(-bx-2dx-a-2c)}}{8(b+2d)} - \frac{e^{(-bx-a)}}{4b}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c)^2,x, algorithm="giac")`

output `1/8*e^(b*x + 2*d*x + a + 2*c)/(b + 2*d) + 1/8*e^(b*x - 2*d*x + a - 2*c)/(b - 2*d) - 1/4*e^(b*x + a)/b + 1/8*e^(-b*x + 2*d*x - a + 2*c)/(b - 2*d) + 1/8*e^(-b*x - 2*d*x - a - 2*c)/(b + 2*d) - 1/4*e^(-b*x - a)/b`

Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.23

$$\int \sinh(a + bx) \sinh^2(c + dx) dx$$

$$= \frac{b^2 (\cosh(a + bx) - \cosh(a + bx) \cosh(c + dx)^2) - 2d^2 \cosh(a + bx) + 2bd \cosh(c + dx) \sinh(a + bx)}{4bd^2 - b^3}$$

input `int(sinh(a + b*x)*sinh(c + d*x)^2,x)`output `(b^2*(cosh(a + b*x) - cosh(a + b*x)*cosh(c + d*x)^2) - 2*d^2*cosh(a + b*x) + 2*b*d*cosh(c + d*x)*sinh(a + b*x))/(4*b*d^2 - b^3)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 212, normalized size of antiderivative = 3.42

$$\int \sinh(a + bx) \sinh^2(c + dx) dx$$

$$= \frac{e^{2bx+4dx+2a+4c}b^2 - 2e^{2bx+4dx+2a+4c}bd - 2e^{2bx+2dx+2a+2c}b^2 + 8e^{2bx+2dx+2a+2c}d^2 + e^{2bx+2a}b^2 + 2e^{2bx+2a}bd + 8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}{8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}$$

input `int(sinh(b*x+a)*sinh(d*x+c)^2,x)`output `(e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2 - 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d - 2*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2 + 8*e**(2*a + 2*b*x + 2*c + 2*d*x)*d**2 + e**(2*a + 2*b*x)*b**2 + 2*e**(2*a + 2*b*x)*b*d + e**(4*c + 4*d*x)*b**2 + 2*e**(4*c + 4*d*x)*b*d - 2*e**(2*c + 2*d*x)*b**2 + 8*e**(2*c + 2*d*x)*d**2 + b**2 - 2*b*d)/(8*e**(a + b*x + 2*c + 2*d*x)*b*(b**2 - 4*d**2))`

3.9 $\int \sinh(a + bx) \sinh^3(c + dx) dx$

Optimal result	131
Mathematica [A] (verified)	131
Rubi [A] (verified)	132
Maple [A] (verified)	133
Fricas [B] (verification not implemented)	133
Sympy [B] (verification not implemented)	134
Maxima [F(-2)]	135
Giac [B] (verification not implemented)	136
Mupad [B] (verification not implemented)	136
Reduce [B] (verification not implemented)	137

Optimal result

Integrand size = 15, antiderivative size = 91

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = -\frac{\sinh(a - 3c + (b - 3d)x)}{8(b - 3d)} + \frac{3 \sinh(a - c + (b - d)x)}{8(b - d)} - \frac{3 \sinh(a + c + (b + d)x)}{8(b + d)} + \frac{\sinh(a + 3c + (b + 3d)x)}{8(b + 3d)}$$

output

```
-1/8*sinh(a-3*c+(b-3*d)*x)/(b-3*d)+3*sinh(a-c+(b-d)*x)/(8*b-8*d)-3*sinh(a+c+(b+d)*x)/(8*b+8*d)+sinh(a+3*c+(b+3*d)*x)/(8*b+24*d)
```

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.95

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \frac{1}{8} \left(-\frac{\sinh(a - 3c + bx - 3dx)}{b - 3d} + \frac{3 \sinh(a - c + bx - dx)}{b - d} + \frac{\sinh(a + 3c + bx + 3dx)}{b + 3d} - \frac{3 \sinh(a + c + (b + d)x)}{b + d} \right)$$

input `Integrate[Sinh[a + b*x]*Sinh[c + d*x]^3,x]`

output $(-\frac{\text{Sinh}[a - 3c + b*x - 3*d*x]}{(b - 3*d)} + \frac{3*\text{Sinh}[a - c + b*x - d*x]}{(b - d)} + \frac{\text{Sinh}[a + 3*c + b*x + 3*d*x]}{(b + 3*d)} - \frac{3*\text{Sinh}[a + c + (b + d)*x]}{(b + d)})/8$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \sinh^3(c + dx) dx$$

$$\downarrow 6147$$

$$\int \left(-\frac{1}{8} \cosh(a + x(b - 3d) - 3c) + \frac{3}{8} \cosh(a + x(b - d) - c) - \frac{3}{8} \cosh(a + x(b + d) + c) + \frac{1}{8} \cosh(a + x(b + 3d) + 3c) \right) dx$$

$$\downarrow 2009$$

$$-\frac{\sinh(a + x(b - 3d) - 3c)}{8(b - 3d)} + \frac{3 \sinh(a + x(b - d) - c)}{8(b - d)} - \frac{3 \sinh(a + x(b + d) + c)}{8(b + d)} + \frac{\sinh(a + x(b + 3d) + 3c)}{8(b + 3d)}$$

input `Int[Sinh[a + b*x]*Sinh[c + d*x]^3,x]`

output $-1/8*\text{Sinh}[a - 3*c + (b - 3*d)*x]/(b - 3*d) + (3*\text{Sinh}[a - c + (b - d)*x])/(8*(b - d)) - (3*\text{Sinh}[a + c + (b + d)*x])/(8*(b + d)) + \text{Sinh}[a + 3*c + (b + 3*d)*x]/(8*(b + 3*d))$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6147 `Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]^(p)*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))`

Maple [A] (verified)

Time = 2.85 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.92

method	result
default	$-\frac{\sinh(a-3c+(b-3d)x)}{8(b-3d)} + \frac{3 \sinh(a-c+(b-d)x)}{8(b-d)} - \frac{3 \sinh(a+c+(b+d)x)}{8(b+d)} + \frac{\sinh(a+3c+(b+3d)x)}{8b+24d}$
risch	$\frac{(b e^{2bx+2a} - 3d e^{2bx+2a+b+3d}) e^{-bx+3dx-a+3c}}{16(b+3d)(b-3d)} - \frac{3(b e^{2bx+2a} - d e^{2bx+2a+b+d}) e^{-bx+dx-a+c}}{16(b+d)(b-d)} + \frac{3(b e^{2bx+2a} + d e^{2bx+2a})}{16(b+d)}$
parallelrisch	$\frac{-12d^3 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^6 \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) + 12d^2 b \left(1 + \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^2 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^5 + (-24b^2d + 36d^3) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)}{(b-d)(b+d)(b+3d)(b-3d)} \left(\tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^6$
orering	Expression too large to display

input `int(sinh(b*x+a)*sinh(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `-1/8*sinh(a-3*c+(b-3*d)*x)/(b-3*d)+3/8*sinh(a-c+(b-d)*x)/(b-d)-3/8/(b+d)*s
inh(a+c+(b+d)*x)+1/8/(b+3*d)*sinh(a+3*c+(b+3*d)*x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(83) = 166$.

Time = 0.11 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.40

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \frac{9(b^2d - d^3) \cosh(dx + c) \sinh(bx + a) \sinh(dx + c)^2 - (b^3 - bd^2) \cosh(bx + a) \sinh(dx + c)^3 + 3((b^2d - d^3) \cosh(dx + c) \sinh(bx + a) \sinh(dx + c) - (b^3 - bd^2) \cosh(bx + a) \sinh(dx + c)^2) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)}{4((b^4 - 10b^2d + 5d^2) \cosh^2\left(\frac{dx}{2} + \frac{c}{2}\right) - (b^2 - d^2) \sinh^2\left(\frac{dx}{2} + \frac{c}{2}\right))}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c)^3,x, algorithm="fricas")`

output `-1/4*(9*(b^2*d - d^3)*cosh(d*x + c)*sinh(b*x + a)*sinh(d*x + c)^2 - (b^3 - b*d^2)*cosh(b*x + a)*sinh(d*x + c)^3 + 3*((b^2*d - d^3)*cosh(d*x + c)^3 - (b^2*d - 9*d^3)*cosh(d*x + c))*sinh(b*x + a) - 3*((b^3 - b*d^2)*cosh(b*x + a)*cosh(d*x + c)^2 - (b^3 - 9*b*d^2)*cosh(b*x + a))*sinh(d*x + c))/((b^4 - 10*b^2*d^2 + 9*d^4)*cosh(b*x + a)^2 - (b^4 - 10*b^2*d^2 + 9*d^4)*sinh(b*x + a)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 918 vs. $2(76) = 152$.

Time = 1.93 (sec) , antiderivative size = 918, normalized size of antiderivative = 10.09

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \text{Too large to display}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c)**3,x)`

output

```
Piecewise((x*sinh(a)*sinh(c)**3, Eq(b, 0) & Eq(d, 0)), (x*sinh(a - 3*d*x)*
sinh(c + d*x)**3/8 + 3*x*sinh(a - 3*d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8
+ 3*x*sinh(c + d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)/8 + x*cosh(a - 3*d*x)
*cosh(c + d*x)**3/8 - sinh(a - 3*d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d)
+ sinh(a - 3*d*x)*cosh(c + d*x)**3/(24*d) - 3*sinh(c + d*x)**3*cosh(a - 3
*d*x)/(8*d), Eq(b, -3*d)), (3*x*sinh(a - d*x)*sinh(c + d*x)**3/8 - 3*x*sin
h(a - d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8 + 3*x*sinh(c + d*x)**2*cosh(a
- d*x)*cosh(c + d*x)/8 - 3*x*cosh(a - d*x)*cosh(c + d*x)**3/8 + 3*sinh(a -
d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d) - 3*sinh(a - d*x)*cosh(c + d*x)
**3/(8*d) + sinh(c + d*x)**3*cosh(a - d*x)/(8*d), Eq(b, -d)), (3*x*sinh(a
+ d*x)*sinh(c + d*x)**3/8 - 3*x*sinh(a + d*x)*sinh(c + d*x)*cosh(c + d*x)*
**2/8 - 3*x*sinh(c + d*x)**2*cosh(a + d*x)*cosh(c + d*x)/8 + 3*x*cosh(a + d
*x)*cosh(c + d*x)**3/8 + 3*sinh(a + d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4
*d) - 3*sinh(a + d*x)*cosh(c + d*x)**3/(8*d) - sinh(c + d*x)**3*cosh(a + d
*x)/(8*d), Eq(b, d)), (x*sinh(a + 3*d*x)*sinh(c + d*x)**3/8 + 3*x*sinh(a +
3*d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8 - 3*x*sinh(c + d*x)**2*cosh(a + 3
*d*x)*cosh(c + d*x)/8 - x*cosh(a + 3*d*x)*cosh(c + d*x)**3/8 - sinh(a + 3*
d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d) + sinh(a + 3*d*x)*cosh(c + d*x)*
**3/(24*d) + 3*sinh(c + d*x)**3*cosh(a + 3*d*x)/(8*d), Eq(b, 3*d)), (b**3*s
inh(c + d*x)**3*cosh(a + b*x)/(b**4 - 10*b**2*d**2 + 9*d**4) - 3*b**2*d...
```

Maxima [F(-2)]

Exception generated.

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(sinh(b*x+a)*sinh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for
more detail
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(83) = 166$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.97

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \frac{e^{(bx+3dx+a+3c)}}{16(b+3d)} - \frac{3e^{(bx+dx+a+c)}}{16(b+d)} + \frac{3e^{(bx-dx+a-c)}}{16(b-d)} - \frac{e^{(bx-3dx+a-3c)}}{16(b-3d)} + \frac{e^{(-bx+3dx-a+3c)}}{16(b-3d)} - \frac{3e^{(-bx+dx-a+c)}}{16(b-d)} + \frac{3e^{(-bx-dx-a-c)}}{16(b+d)} - \frac{e^{(-bx-3dx-a-3c)}}{16(b+3d)}$$

input `integrate(sinh(b*x+a)*sinh(d*x+c)^3,x, algorithm="giac")`

output $\frac{1}{16}e^{(bx+3dx+a+3c)}/(b+3d) - \frac{3}{16}e^{(bx+dx+a+c)}/(b+d) + \frac{3}{16}e^{(bx-dx+a-c)}/(b-d) - \frac{1}{16}e^{(bx-3dx+a-3c)}/(b-3d) + \frac{1}{16}e^{(-bx+3dx-a+3c)}/(b-3d) - \frac{3}{16}e^{(-bx+dx-a+c)}/(b-d) + \frac{3}{16}e^{(-bx-dx-a-c)}/(b+d) - \frac{1}{16}e^{(-bx-3dx-a-3c)}/(b+3d)$

Mupad [B] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.00

$$\int \sinh(a + bx) \sinh^3(c + dx) dx = \frac{6bd^2 \cosh(a + bx) \cosh(c + dx)^2 \sinh(c + dx)}{b^4 - 10b^2d^2 + 9d^4} - \frac{6d^3 \cosh(c + dx)^3 \sinh(a + bx)}{b^4 - 10b^2d^2 + 9d^4} - \frac{3d \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)^2 (b^2 - 3d^2)}{b^4 - 10b^2d^2 + 9d^4} - \frac{\cosh(a + bx) \sinh(c + dx)^3 (7bd^2 - b^3)}{b^4 - 10b^2d^2 + 9d^4}$$

input `int(sinh(a + b*x)*sinh(c + d*x)^3,x)`

output

$$\begin{aligned} & (6*b*d^2*cosh(a + b*x)*cosh(c + d*x)^2*sinh(c + d*x))/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (6*d^3*cosh(c + d*x)^3*sinh(a + b*x))/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (3*d*cosh(c + d*x)*sinh(a + b*x)*sinh(c + d*x)^2*(b^2 - 3*d^2))/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (cosh(a + b*x)*sinh(c + d*x)^3*(7*b*d^2 - b^3))/(b^4 + 9*d^4 - 10*b^2*d^2) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 574, normalized size of antiderivative = 6.31

$$\int \sinh(a + bx) \sinh^3(c + dx) dx$$

$$= \frac{-b^3 - 3d^3 + b d^2 - e^{6dx+6c} b d^2 + e^{2bx+2a} b d^2 - e^{2bx+6dx+2a+6c} b d^2 - 27e^{2bx+4dx+2a+4c} d^3 + 3e^{2bx+2dx+2a+2c} b d^2}{b^4 + 9d^4 - 10b^2d^2}$$

input

`int(sinh(b*x+a)*sinh(d*x+c)^3,x)`

output

$$\begin{aligned} & (e^{2a + 2bx + 6c + 6dx} b^3 - 3e^{2a + 2bx + 6c + 6dx} b^2 d - e^{2a + 2bx + 6c + 6dx} b d^2 + 3e^{2a + 2bx + 6c + 6dx} d^3 - 3e^{2a + 2bx + 4c + 4dx} b^3 + 3e^{2a + 2bx + 4c + 4dx} b^2 d + 27e^{2a + 2bx + 4c + 4dx} b d^2 - 27e^{2a + 2bx + 4c + 4dx} d^3 + 3e^{2a + 2bx + 2c + 2dx} b^3 + 3e^{2a + 2bx + 2c + 2dx} b^2 d - 27e^{2a + 2bx + 2c + 2dx} b d^2 - 27e^{2a + 2bx + 2c + 2dx} d^3 - e^{2a + 2bx} b^3 - 3e^{2a + 2bx} b^2 d + e^{2a + 2bx} b d^2 + 3e^{2a + 2bx} d^3 + e^{6c + 6dx} b^3 + 3e^{6c + 6dx} b^2 d - e^{6c + 6dx} b d^2 - 3e^{6c + 6dx} d^3 - 3e^{4c + 4dx} b^3 - 3e^{4c + 4dx} b^2 d + 27e^{4c + 4dx} b d^2 + 27e^{4c + 4dx} d^3 + 3e^{2c + 2dx} b^3 - 3e^{2c + 2dx} b^2 d - 27e^{2c + 2dx} b d^2 + 27e^{2c + 2dx} d^3 - b^3 + 3b^2 d + b d^2 - 3d^3)/(16e^{2a + 2bx + 6c + 6dx} (b^4 - 10b^2 d^2 + 9d^4)) \end{aligned}$$

3.10 $\int \sinh^2(a + bx) \sinh^2(c + dx) dx$

Optimal result	138
Mathematica [A] (verified)	138
Rubi [A] (verified)	139
Maple [A] (verified)	140
Fricas [B] (verification not implemented)	140
Sympy [B] (verification not implemented)	141
Maxima [F(-2)]	142
Giac [A] (verification not implemented)	143
Mupad [B] (verification not implemented)	143
Reduce [B] (verification not implemented)	144

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \frac{x}{4} - \frac{\sinh(2a + 2bx)}{8b} + \frac{\sinh(2(a - c) + 2(b - d)x)}{16(b - d)} - \frac{\sinh(2c + 2dx)}{8d} + \frac{\sinh(2(a + c) + 2(b + d)x)}{16(b + d)}$$

output

```
1/4*x-1/8*sinh(2*b*x+2*a)/b+sinh(2*a-2*c+2*(b-d)*x)/(16*b-16*d)-1/8*sinh(2*d*x+2*c)/d+sinh(2*a+2*c+2*(b+d)*x)/(16*b+16*d)
```

Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.20

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \frac{(-2b^2d + 2d^3) \sinh(2(a + bx)) + bd(b + d) \sinh(2(a - c + (b - d)x)) + b(b - d)(-2(b + d) \sinh(2(c + dx)))}{16b(b - d)d(b + d)}$$

input

```
Integrate[Sinh[a + b*x]^2*Sinh[c + d*x]^2,x]
```

output

$$\frac{((-2*b^2*d + 2*d^3)*\text{Sinh}[2*(a + b*x)] + b*d*(b + d)*\text{Sinh}[2*(a - c + (b - d)*x)] + b*(b - d)*(-2*(b + d)*\text{Sinh}[2*(c + d*x)] + d*(4*(b + d)*x + \text{Sinh}[2*(a + c + (b + d)*x)]))}{(16*b*(b - d)*d*(b + d))}$$
Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx$$

$$\downarrow \text{6147}$$

$$\int \left(\frac{1}{8} \cosh(2(a - c) + 2x(b - d)) + \frac{1}{8} \cosh(2(a + c) + 2x(b + d)) - \frac{1}{4} \cosh(2a + 2bx) - \frac{1}{4} \cosh(2c + 2dx) + \frac{1}{4} \right)$$

$$\downarrow \text{2009}$$

$$\frac{\sinh(2(a - c) + 2x(b - d))}{16(b - d)} + \frac{\sinh(2(a + c) + 2x(b + d))}{16(b + d)} - \frac{\sinh(2a + 2bx)}{8b} - \frac{\sinh(2c + 2dx)}{8d} + \frac{x}{4}$$

input

$$\text{Int}[\text{Sinh}[a + b*x]^2*\text{Sinh}[c + d*x]^2,x]$$

output

$$x/4 - \text{Sinh}[2*a + 2*b*x]/(8*b) + \text{Sinh}[2*(a - c) + 2*(b - d)*x]/(16*(b - d)) - \text{Sinh}[2*c + 2*d*x]/(8*d) + \text{Sinh}[2*(a + c) + 2*(b + d)*x]/(16*(b + d))$$

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6147 Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]^(p)*Sinh[w]^(q), x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 3.67 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

method	result
default	$\frac{x}{4} - \frac{\sinh(2bx+2a)}{8b} - \frac{\sinh(2dx+2c)}{8d} + \frac{\sinh((2b-2d)x+2a-2c)}{16b-16d} + \frac{\sinh((2b+2d)x+2a+2c)}{16b+16d}$
parallelrisch	$\frac{bd(b+d) \sinh((2b-2d)x+2a-2c)+4(b-d) \left(\frac{bd \sinh((2b+2d)x+2a+2c)}{4} + \left(-\frac{d \sinh(2bx+2a)}{2} + b \left(dx - \frac{\sinh(2dx+2c)}{2} \right) \right) \right) (b+d)}{16b^3d-16bd^3}$
risch	$\frac{x}{4} - \frac{e^{2bx+2a}}{16b} + \frac{e^{-2bx-2a}}{16b} - \frac{(-de^{4bx+4a}b+d^2e^{4bx+4a}+2b^2e^{2bx+2a}-2d^2e^{2bx+2a}+bd+d^2)e^{-2bx+2dx-2a+2c}}{32(b+d)(b-d)d} + \frac{(de^{4bx+4a}b+d^2e^{4bx+4a}+2b^2e^{2bx+2a}-2d^2e^{2bx+2a}+bd+d^2)e^{-2bx+2dx-2a+2c}}{32(b+d)(b-d)d}$
orering	Expression too large to display

```
input int(sinh(b*x+a)^2*sinh(d*x+c)^2,x,method=_RETURNVERBOSE)
```

```
output 1/4*x-1/8*sinh(2*b*x+2*a)/b-1/8*sinh(2*d*x+2*c)/d+1/8/(2*b-2*d)*sinh((2*b-2*d)*x+2*a-2*c)+1/8/(2*b+2*d)*sinh((2*b+2*d)*x+2*a+2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 192 vs. 2(78) = 156.

Time = 0.09 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.18

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx$$

$$= \frac{b^2d \cosh(bx + a) \sinh(bx + a) \sinh(dx + c)^2 + (b^3d - bd^3)x + (b^2d \cosh(bx + a) \cosh(dx + c)^2 - (b^2d - b^2d) \sinh(bx + a) \sinh(dx + c))}{4((b^3d - bd^3) \cosh(bx + a) \sinh(dx + c) + (b^2d \cosh(bx + a) \cosh(dx + c) - (b^2d - b^2d) \sinh(bx + a) \sinh(dx + c)))}$$

input `integrate(sinh(b*x+a)^2*sinh(d*x+c)^2,x, algorithm="fricas")`

output
$$\frac{1}{4}(b^2d\cosh(bx+a)\sinh(bx+a)\sinh(dx+c)^2 + (b^3d - b^2d^3)x + (b^2d\cosh(bx+a)\cosh(dx+c)^2 - (b^2d - d^3)\cosh(bx+a))\sinh(bx+a) - (bd^2\cosh(dx+c)\sinh(bx+a)^2 + (bd^2\cosh(bx+a)^2 + b^3 - bd^2)\cosh(dx+c))\sinh(dx+c)) / ((b^3d - b^2d^3)\cosh(bx+a)^2 - (b^3d - b^2d^3)\sinh(bx+a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1027 vs. $2(76) = 152$.

Time = 1.58 (sec) , antiderivative size = 1027, normalized size of antiderivative = 11.67

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \text{Too large to display}$$

input `integrate(sinh(b*x+a)**2*sinh(d*x+c)**2,x)`

output

```
Piecewise((x*sinh(a)**2*sinh(c)**2, Eq(b, 0) & Eq(d, 0)), ((x*sinh(c + d*x)
)**2/2 - x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*sinh(a)
**2, Eq(b, 0)), (3*x*sinh(a - d*x)**2*sinh(c + d*x)**2/8 - x*sinh(a - d*x)
**2*cosh(c + d*x)**2/8 + x*sinh(a - d*x)*sinh(c + d*x)*cosh(a - d*x)*cosh(
c + d*x)/2 - x*sinh(c + d*x)**2*cosh(a - d*x)**2/8 + 3*x*cosh(a - d*x)**2*
cosh(c + d*x)**2/8 - 5*sinh(a - d*x)*sinh(c + d*x)**2*cosh(a - d*x)/(8*d)
- sinh(a - d*x)*cosh(a - d*x)*cosh(c + d*x)**2/(8*d) - sinh(c + d*x)*cosh(
a - d*x)**2*cosh(c + d*x)/(2*d), Eq(b, -d)), (3*x*sinh(a + d*x)**2*sinh(c
+ d*x)**2/8 - x*sinh(a + d*x)**2*cosh(c + d*x)**2/8 - x*sinh(a + d*x)*sinh
(c + d*x)*cosh(a + d*x)*cosh(c + d*x)/2 - x*sinh(c + d*x)**2*cosh(a + d*x)
**2/8 + 3*x*cosh(a + d*x)**2*cosh(c + d*x)**2/8 + sinh(a + d*x)**2*sinh(c
+ d*x)*cosh(c + d*x)/(2*d) + sinh(a + d*x)*sinh(c + d*x)**2*cosh(a + d*x)/
(8*d) - 3*sinh(a + d*x)*cosh(a + d*x)*cosh(c + d*x)**2/(8*d), Eq(b, d)), (
(x*sinh(a + b*x)**2/2 - x*cosh(a + b*x)**2/2 + sinh(a + b*x)*cosh(a + b*x)
)/(2*b))*sinh(c)**2, Eq(d, 0)), (b**3*d*x*sinh(a + b*x)**2*sinh(c + d*x)**2
/(4*b**3*d - 4*b*d**3) - b**3*d*x*sinh(a + b*x)**2*cosh(c + d*x)**2/(4*b**
3*d - 4*b*d**3) - b**3*d*x*sinh(c + d*x)**2*cosh(a + b*x)**2/(4*b**3*d - 4
*b*d**3) + b**3*d*x*cosh(a + b*x)**2*cosh(c + d*x)**2/(4*b**3*d - 4*b*d**3
) + b**3*sinh(a + b*x)**2*sinh(c + d*x)*cosh(c + d*x)/(4*b**3*d - 4*b*d**3
) - b**3*sinh(c + d*x)*cosh(a + b*x)**2*cosh(c + d*x)/(4*b**3*d - 4*b*d...
```

Maxima [F(-2)]

Exception generated.

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(sinh(b*x+a)^2*sinh(d*x+c)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(1-(2*d)/b>0)', see `assume?` for
more deta
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.77

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \frac{1}{4}x + \frac{e^{(2bx+2dx+2a+2c)}}{32(b+d)} + \frac{e^{(2bx-2dx+2a-2c)}}{32(b-d)} - \frac{e^{(2bx+2a)}}{16b} - \frac{e^{(-2bx+2dx-2a+2c)}}{32(b-d)} - \frac{e^{(-2bx-2dx-2a-2c)}}{32(b+d)} + \frac{e^{(-2bx-2a)}}{16b} - \frac{e^{(2dx+2c)}}{16d} + \frac{e^{(-2dx-2c)}}{16d}$$

input `integrate(sinh(b*x+a)^2*sinh(d*x+c)^2,x, algorithm="giac")`output `1/4*x + 1/32*e^(2*b*x + 2*d*x + 2*a + 2*c)/(b + d) + 1/32*e^(2*b*x - 2*d*x + 2*a - 2*c)/(b - d) - 1/16*e^(2*b*x + 2*a)/b - 1/32*e^(-2*b*x + 2*d*x - 2*a + 2*c)/(b - d) - 1/32*e^(-2*b*x - 2*d*x - 2*a - 2*c)/(b + d) + 1/16*e^(-2*b*x - 2*a)/b - 1/16*e^(2*d*x + 2*c)/d + 1/16*e^(-2*d*x - 2*c)/d`**Mupad [B] (verification not implemented)**

Time = 1.49 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.73

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx = \frac{d^3 \cosh(a + bx) \sinh(a + bx) - b^3 \cosh(c + dx) \sinh(c + dx) - b d^3 x + b^3 dx - 2 b^2 d \cosh(a + bx) \sinh(c + dx) + 2 b^2 d \cosh(c + dx) \sinh(a + bx) - 2 b^2 d \cosh(a + bx) \cosh(c + dx) \sinh(a + bx) - 2 b^2 d \cosh(c + dx) \cosh(a + bx) \sinh(c + dx)}{(4 b^3 d^3 - 4 b^3 d)}$$

input `int(sinh(a + b*x)^2*sinh(c + d*x)^2,x)`output `-(d^3*cosh(a + b*x)*sinh(a + b*x) - b^3*cosh(c + d*x)*sinh(c + d*x) - b*d^3*x + b^3*d*x - 2*b^2*d*cosh(a + b*x)*sinh(a + b*x) + 2*b*d^2*cosh(c + d*x)*sinh(c + d*x) + 2*b^2*d*cosh(a + b*x)*cosh(c + d*x)^2*sinh(a + b*x) - 2*b*d^2*cosh(a + b*x)^2*cosh(c + d*x)*sinh(c + d*x))/(4*b*d^3 - 4*b^3*d)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.06

$$\int \sinh^2(a + bx) \sinh^2(c + dx) dx$$

$$= \frac{e^{4bx+4dx+4a+4c} b^2 d - e^{4bx+4dx+4a+4c} b d^2 - 2e^{4bx+2dx+4a+2c} b^2 d + 2e^{4bx+2dx+4a+2c} d^3 + e^{4bx+4a} b^2 d + e^{4bx+4a} b d^2}{32e^{2a+2bx+2c+2dx} (b^2 d - d^2)}$$

input

```
int(sinh(b*x+a)^2*sinh(d*x+c)^2,x)
```

output

```
(e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d - e**(4*a + 4*b*x + 4*c + 4*d*x)*b*
d**2 - 2*e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d + 2*e**(4*a + 4*b*x + 2*c +
2*d*x)*d**3 + e**(4*a + 4*b*x)*b**2*d + e**(4*a + 4*b*x)*b*d**2 - 2*e**(2
*a + 2*b*x + 4*c + 4*d*x)*b**3 + 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d**2 +
8*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**3*d*x - 8*e**(2*a + 2*b*x + 2*c + 2*d
*x)*b*d**3*x + 2*e**(2*a + 2*b*x)*b**3 - 2*e**(2*a + 2*b*x)*b*d**2 - e**(4
*c + 4*d*x)*b**2*d - e**(4*c + 4*d*x)*b*d**2 + 2*e**(2*c + 2*d*x)*b**2*d -
2*e**(2*c + 2*d*x)*d**3 - b**2*d + b*d**2)/(32*e**(2*a + 2*b*x + 2*c + 2*
d*x)*b*d*(b**2 - d**2))
```

3.11 $\int \sinh^2(a + bx) \sinh^3(c + dx) dx$

Optimal result	145
Mathematica [A] (verified)	146
Rubi [A] (verified)	146
Maple [A] (verified)	148
Fricas [B] (verification not implemented)	148
Sympy [B] (verification not implemented)	149
Maxima [F(-2)]	150
Giac [A] (verification not implemented)	151
Mupad [B] (verification not implemented)	152
Reduce [B] (verification not implemented)	153

Optimal result

Integrand size = 17, antiderivative size = 144

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx = -\frac{\cosh(2a - 3c + (2b - 3d)x)}{16(2b - 3d)} + \frac{3 \cosh(2a - c + (2b - d)x)}{16(2b - d)} + \frac{3 \cosh(c + dx)}{8d} - \frac{\cosh(3c + 3dx)}{24d} - \frac{3 \cosh(2a + c + (2b + d)x)}{16(2b + d)} + \frac{\cosh(2a + 3c + (2b + 3d)x)}{16(2b + 3d)}$$

output

```
-1/16*cosh(2*a-3*c+(2*b-3*d)*x)/(2*b-3*d)+3*cosh(2*a-c+(2*b-d)*x)/(32*b-16*d)+3/8*cosh(d*x+c)/d-1/24*cosh(3*d*x+3*c)/d-3*cosh(2*a+c+(2*b+d)*x)/(32*b+16*d)+cosh(2*a+3*c+(2*b+3*d)*x)/(32*b+48*d)
```

Mathematica [A] (verified)

Time = 1.18 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.10

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx = \frac{1}{48} \left(\frac{18 \cosh(c) \cosh(dx)}{d} - \frac{2 \cosh(3c) \cosh(3dx)}{d} - \frac{3 \cosh(2a - 3c + 2bx - 3dx)}{2b - 3d} + \frac{9 \cosh(2a - c + 2bx - dx)}{2b - d} - \frac{9 \cosh(2a + c + 2bx + dx)}{2b + d} + \frac{3 \cosh(2a + 3c + 2bx + 3dx)}{2b + 3d} + \frac{18 \sinh(c) \sinh(dx)}{d} - \frac{2 \sinh(3c) \sinh(3dx)}{d} \right)$$

input

```
Integrate[Sinh[a + b*x]^2*Sinh[c + d*x]^3,x]
```

output

```
((18*Cosh[c]*Cosh[d*x])/d - (2*Cosh[3*c]*Cosh[3*d*x])/d - (3*Cosh[2*a - 3*c + 2*b*x - 3*d*x])/(2*b - 3*d) + (9*Cosh[2*a - c + 2*b*x - d*x])/(2*b - d) - (9*Cosh[2*a + c + 2*b*x + d*x])/(2*b + d) + (3*Cosh[2*a + 3*c + 2*b*x + 3*d*x])/(2*b + 3*d) + (18*Sinh[c]*Sinh[d*x])/d - (2*Sinh[3*c]*Sinh[3*d*x])/d)/48
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx$$

↓ 6147

$$\int \left(-\frac{1}{16} \sinh(2a + x(2b - 3d) - 3c) + \frac{3}{16} \sinh(2a + x(2b - d) - c) - \frac{3}{16} \sinh(2a + x(2b + d) + c) + \frac{1}{16} \sinh(2a + x(2b + 3d) + 3c) + \frac{3}{8d} \cosh(c + dx) - \frac{\cosh(3c + 3dx)}{24d} \right) dx$$

↓ 2009

$$-\frac{\cosh(2a + x(2b - 3d) - 3c)}{16(2b - 3d)} + \frac{3 \cosh(2a + x(2b - d) - c)}{16(2b - d)} - \frac{3 \cosh(2a + x(2b + d) + c)}{16(2b + d)} + \frac{\cosh(2a + x(2b + 3d) + 3c)}{16(2b + 3d)} + \frac{3 \cosh(c + dx)}{8d} - \frac{\cosh(3c + 3dx)}{24d}$$

input `Int[Sinh[a + b*x]^2*Sinh[c + d*x]^3,x]`

output `-1/16*Cosh[2*a - 3*c + (2*b - 3*d)*x]/(2*b - 3*d) + (3*Cosh[2*a - c + (2*b - d)*x])/(16*(2*b - d)) + (3*Cosh[c + d*x])/(8*d) - Cosh[3*c + 3*d*x]/(24*d) - (3*Cosh[2*a + c + (2*b + d)*x])/(16*(2*b + d)) + Cosh[2*a + 3*c + (2*b + 3*d)*x]/(16*(2*b + 3*d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6147 `Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 9.10 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.92

method	result
default	$\frac{3 \cosh(dx+c)}{8d} - \frac{\cosh(3dx+3c)}{24d} - \frac{\cosh(2a-3c+(2b-3d)x)}{16(2b-3d)} + \frac{3 \cosh(2a-c+(2b-d)x)}{16(2b-d)} - \frac{3 \cosh(2a+c+(2b+d)x)}{16(2b+d)} + \dots$
parallelrisch	$\frac{(-24b^3d-36b^2d^2+6bd^3+9d^4) \cosh(2a-3c+(2b-3d)x) + (72b^3d+36b^2d^2-162bd^3-81d^4) \cosh(2a-c+(2b-d)x) + (24b^3d-36b^2d^2+6bd^3+9d^4) \cosh(2a+c+(2b+d)x)}{96(2b+3d)(2b-3d)d}$
risch	$-\frac{(-6de^{4bx+4a}b+9d^2e^{4bx+4a}+8b^2e^{2bx+2a}-18d^2e^{2bx+2a}+6bd+9d^2)e^{-2bx+3dx-2a+3c}}{96(2b+3d)(2b-3d)d} + \frac{3(-2de^{4bx+4a}b+d^2e^{4bx+4a}+8b^2e^{2bx+2a}-18d^2e^{2bx+2a}+6bd+9d^2)e^{-2bx+3dx-2a+3c}}{96(2b+3d)(2b-3d)d}$
orering	Expression too large to display

input `int(sinh(b*x+a)^2*sinh(d*x+c)^3,x,method=_RETURNVERBOSE)`

output `3/8*cosh(d*x+c)/d-1/24*cosh(3*d*x+3*c)/d-1/16*cosh(2*a-3*c+(2*b-3*d)*x)/(2*b-3*d)+3/16/(2*b-d)*cosh(2*a-c+(2*b-d)*x)-3/16/(2*b+d)*cosh(2*a+c+(2*b+d)*x)+1/16/(2*b+3*d)*cosh(2*a+3*c+(2*b+3*d)*x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 414 vs. 2(132) = 264.

Time = 0.10 (sec) , antiderivative size = 414, normalized size of antiderivative = 2.88

$$\int \sinh^2(a+bx) \sinh^3(c+dx) dx$$

$$= \frac{12(4b^3d - bd^3) \cosh(bx+a) \sinh(bx+a) \sinh(dx+c)^3 - (16b^4 - 40b^2d^2 + 9d^4 + 9(4b^2d^2 - d^4) \cosh(2a-c+(2b-d)x) + \dots}{96(2b+3d)(2b-3d)d}$$

input `integrate(sinh(b*x+a)^2*sinh(d*x+c)^3,x, algorithm="fricas")`

output

```
1/24*(12*(4*b^3*d - b*d^3)*cosh(b*x + a)*sinh(b*x + a)*sinh(d*x + c)^3 - (
16*b^4 - 40*b^2*d^2 + 9*d^4 + 9*(4*b^2*d^2 - d^4)*cosh(b*x + a)^2)*cosh(d*
x + c)^3 - 9*((4*b^2*d^2 - d^4)*cosh(d*x + c)^3 - (4*b^2*d^2 - 9*d^4)*cosh
(d*x + c))*sinh(b*x + a)^2 + 36*((4*b^3*d - b*d^3)*cosh(b*x + a)*cosh(d*x
+ c)^2 - (4*b^3*d - 9*b*d^3)*cosh(b*x + a))*sinh(b*x + a)*sinh(d*x + c) -
3*(9*(4*b^2*d^2 - d^4)*cosh(d*x + c)*sinh(b*x + a)^2 + (16*b^4 - 40*b^2*d^
2 + 9*d^4 + 9*(4*b^2*d^2 - d^4)*cosh(b*x + a)^2)*cosh(d*x + c))*sinh(d*x +
c)^2 + 9*(16*b^4 - 40*b^2*d^2 + 9*d^4 + (4*b^2*d^2 - 9*d^4)*cosh(b*x + a)
^2)*cosh(d*x + c))/((16*b^4*d - 40*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2 - (16*
b^4*d - 40*b^2*d^3 + 9*d^5)*sinh(b*x + a)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1999 vs. $2(116) = 232$.

Time = 5.33 (sec) , antiderivative size = 1999, normalized size of antiderivative = 13.88

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(sinh(b*x+a)**2*sinh(d*x+c)**3,x)
```

output

```
Piecewise((x*sinh(a)**2*sinh(c)**3, Eq(b, 0) & Eq(d, 0)), (x*sinh(a - 3*d*x/2)**2*sinh(c + d*x)**3/16 + 3*x*sinh(a - 3*d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/16 + 3*x*sinh(a - 3*d*x/2)*sinh(c + d*x)**2*cosh(a - 3*d*x/2)*cosh(c + d*x)/8 + x*sinh(a - 3*d*x/2)*cosh(a - 3*d*x/2)*cosh(c + d*x)**3/8 + x*sinh(c + d*x)**3*cosh(a - 3*d*x/2)**2/16 + 3*x*sinh(c + d*x)*cosh(a - 3*d*x/2)**2*cosh(c + d*x)**2/16 + sinh(a - 3*d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/d - 5*sinh(a - 3*d*x/2)**2*cosh(c + d*x)**3/(48*d) + sinh(a - 3*d*x/2)*sinh(c + d*x)**3*cosh(a - 3*d*x/2)/(24*d) + 5*sinh(a - 3*d*x/2)*sinh(c + d*x)*cosh(a - 3*d*x/2)*cosh(c + d*x)**2/(4*d) + 9*cosh(a - 3*d*x/2)**2*cosh(c + d*x)**3/(16*d), Eq(b, -3*d/2)), (3*x*sinh(a - d*x/2)**2*sinh(c + d*x)**3/16 - 3*x*sinh(a - d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/16 + 3*x*sinh(a - d*x/2)*sinh(c + d*x)**2*cosh(a - d*x/2)*cosh(c + d*x)/8 - 3*x*sinh(a - d*x/2)*cosh(a - d*x/2)*cosh(c + d*x)**3/8 + 3*x*sinh(c + d*x)**3*cosh(a - d*x/2)**2/16 - 3*x*sinh(c + d*x)*cosh(a - d*x/2)**2*cosh(c + d*x)**2/16 + sinh(a - d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/d - 31*sinh(a - d*x/2)**2*cosh(c + d*x)**3/(48*d) + 3*sinh(a - d*x/2)*sinh(c + d*x)**3*cosh(a - d*x/2)/(8*d) - sinh(a - d*x/2)*sinh(c + d*x)*cosh(a - d*x/2)*cosh(c + d*x)**2/(4*d) + cosh(a - d*x/2)**2*cosh(c + d*x)**3/(48*d), Eq(b, -d/2)), (3*x*sinh(a + d*x/2)**2*sinh(c + d*x)**3/16 - 3*x*sinh(a + d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/16 - 3*x*sinh(a + d*x/2)*sinh(c + d*x...
```

Maxima [F(-2)]

Exception generated.

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(sinh(b*x+a)^2*sinh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-(3*d)/b>0)', see `assume?` for more deta
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.81

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx = \frac{e^{(2bx+3dx+2a+3c)}}{32(2b+3d)} - \frac{3e^{(2bx+dx+2a+c)}}{32(2b+d)}$$

$$+ \frac{3e^{(2bx-dx+2a-c)}}{32(2b-d)} - \frac{e^{(2bx-3dx+2a-3c)}}{32(2b-3d)}$$

$$- \frac{e^{(-2bx+3dx-2a+3c)}}{32(2b-3d)} + \frac{3e^{(-2bx+dx-2a+c)}}{32(2b-d)}$$

$$- \frac{3e^{(-2bx-dx-2a-c)}}{32(2b+d)} + \frac{e^{(-2bx-3dx-2a-3c)}}{32(2b+3d)}$$

$$- \frac{e^{(3dx+3c)}}{48d} + \frac{3e^{(dx+c)}}{16d} + \frac{3e^{(-dx-c)}}{16d} - \frac{e^{(-3dx-3c)}}{48d}$$

input `integrate(sinh(b*x+a)^2*sinh(d*x+c)^3,x, algorithm="giac")`

output `1/32*e^(2*b*x + 3*d*x + 2*a + 3*c)/(2*b + 3*d) - 3/32*e^(2*b*x + d*x + 2*a + c)/(2*b + d) + 3/32*e^(2*b*x - d*x + 2*a - c)/(2*b - d) - 1/32*e^(2*b*x - 3*d*x + 2*a - 3*c)/(2*b - 3*d) - 1/32*e^(-2*b*x + 3*d*x - 2*a + 3*c)/(2*b - 3*d) + 3/32*e^(-2*b*x + d*x - 2*a + c)/(2*b - d) - 3/32*e^(-2*b*x - d*x - 2*a - c)/(2*b + d) + 1/32*e^(-2*b*x - 3*d*x - 2*a - 3*c)/(2*b + 3*d) - 1/48*e^(3*d*x + 3*c)/d + 3/16*e^(d*x + c)/d + 3/16*e^(-d*x - c)/d - 1/48*e^(-3*d*x - 3*c)/d`

Mupad [B] (verification not implemented)

Time = 1.51 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.34

$$\begin{aligned}
& \int \sinh^2(a + bx) \sinh^3(c + dx) dx \\
&= \frac{\cosh(c + dx) \sinh(a + bx)^2 \sinh(c + dx)^2 (8b^4 - 26b^2 d^2 + 9d^4)}{d (16b^4 - 40b^2 d^2 + 9d^4)} \\
&\quad - \cosh(c + dx)^3 \sinh(a + bx)^2 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} + \frac{1}{3d} \right) \\
&\quad - \frac{2 \cosh(a + bx) \sinh(a + bx) \sinh(c + dx)^3 (7bd^2 - 4b^3)}{16b^4 - 40b^2 d^2 + 9d^4} \\
&\quad - \frac{\cosh(a + bx)^2 \cosh(c + dx) \sinh(c + dx)^2 (8b^4 - 14b^2 d^2)}{d (16b^4 - 40b^2 d^2 + 9d^4)} \\
&\quad - \cosh(a + bx)^2 \cosh(c + dx)^3 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} - \frac{1}{3d} \right) \\
&\quad + \frac{12bd^2 \cosh(a + bx) \cosh(c + dx)^2 \sinh(a + bx) \sinh(c + dx)}{16b^4 - 40b^2 d^2 + 9d^4}
\end{aligned}$$

input `int(sinh(a + b*x)^2*sinh(c + d*x)^3,x)`output `(cosh(c + d*x)*sinh(a + b*x)^2*sinh(c + d*x)^2*(8*b^4 + 9*d^4 - 26*b^2*d^2))/ (d*(16*b^4 + 9*d^4 - 40*b^2*d^2)) - cosh(c + d*x)^3*sinh(a + b*x)^2*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) + 1/(3*d)) - (2*cosh(a + b*x)*sinh(a + b*x)*sinh(c + d*x)^3*(7*b*d^2 - 4*b^3))/(16*b^4 + 9*d^4 - 40*b^2*d^2) - (cosh(a + b*x)^2*cosh(c + d*x)*sinh(c + d*x)^2*(8*b^4 - 14*b^2*d^2))/(d*(16*b^4 + 9*d^4 - 40*b^2*d^2)) - cosh(a + b*x)^2*cosh(c + d*x)^3*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) - 1/(3*d)) + (12*b*d^2*cosh(a + b*x)*cosh(c + d*x)^2*sinh(a + b*x)*sinh(c + d*x))/(16*b^4 + 9*d^4 - 40*b^2*d^2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.01

$$\int \sinh^2(a + bx) \sinh^3(c + dx) dx$$

$$= \frac{36e^{4bx+4dx+4a+4c}b^2d^2 + 162e^{4bx+4dx+4a+4c}bd^3 + 72e^{4bx+2dx+4a+2c}b^3d + 36e^{4bx+2dx+4a+2c}b^2d^2 - 162e^{4bx+2dx+4a+2c}bd^3}{1}$$

input `int(sinh(b*x+a)^2*sinh(d*x+c)^3,x)`

output

```
(24***e**(4*a + 4*b*x + 6*c + 6*d*x)*b**3*d - 36***e**(4*a + 4*b*x + 6*c + 6*d*x)*b**2*d**2 - 6***e**(4*a + 4*b*x + 6*c + 6*d*x)*b*d**3 + 9***e**(4*a + 4*b*x + 6*c + 6*d*x)*d**4 - 72***e**(4*a + 4*b*x + 4*c + 4*d*x)*b**3*d + 36***e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d**2 + 162***e**(4*a + 4*b*x + 4*c + 4*d*x)*b*d**3 - 81***e**(4*a + 4*b*x + 4*c + 4*d*x)*d**4 + 72***e**(4*a + 4*b*x + 2*c + 2*d*x)*b**3*d + 36***e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d**2 - 162***e**(4*a + 4*b*x + 2*c + 2*d*x)*b*d**3 - 81***e**(4*a + 4*b*x + 2*c + 2*d*x)*d**4 - 24***e**(4*a + 4*b*x)*b**3*d - 36***e**(4*a + 4*b*x)*b**2*d**2 + 6***e**(4*a + 4*b*x)*b*d**3 + 9***e**(4*a + 4*b*x)*d**4 - 32***e**(2*a + 2*b*x + 6*c + 6*d*x)*b**4 + 80***e**(2*a + 2*b*x + 6*c + 6*d*x)*b**2*d**2 - 18***e**(2*a + 2*b*x + 6*c + 6*d*x)*d**4 + 288***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**4 - 720***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2*d**2 + 162***e**(2*a + 2*b*x + 4*c + 4*d*x)*d**4 + 288***e**(2*a + 2*b*x + 2*c + 2*d*x)*b**4 - 720***e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2*d**2 + 162***e**(2*a + 2*b*x + 2*c + 2*d*x)*d**4 - 32***e**(2*a + 2*b*x)*b**4 + 80***e**(2*a + 2*b*x)*b**2*d**2 - 18***e**(2*a + 2*b*x)*d**4 - 24***e**(6*c + 6*d*x)*b**3*d - 36***e**(6*c + 6*d*x)*b**2*d**2 + 6***e**(6*c + 6*d*x)*b*d**3 + 9***e**(6*c + 6*d*x)*d**4 + 72***e**(4*c + 4*d*x)*b**3*d + 36***e**(4*c + 4*d*x)*b**2*d**2 - 162***e**(4*c + 4*d*x)*b*d**3 - 81***e**(4*c + 4*d*x)*d**4 - 72***e**(2*c + 2*d*x)*b**3*d + 36***e**(2*c + 2*d*x)*b**2*d**2 + 162***e**(2*c + 2*d*x)*b*d**3 - 81***e**(2*c + 2*d*x)*d**4 + 24*b**3*d - 36*...
```

3.12 $\int \sinh^3(a + bx) \sinh^3(c + dx) dx$

Optimal result	154
Mathematica [A] (verified)	155
Rubi [A] (verified)	155
Maple [A] (verified)	157
Fricas [B] (verification not implemented)	157
Sympy [B] (verification not implemented)	158
Maxima [F(-2)]	159
Giac [B] (verification not implemented)	160
Mupad [B] (verification not implemented)	161
Reduce [F]	162

Optimal result

Integrand size = 17, antiderivative size = 195

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \frac{3 \sinh(a - 3c + (b - 3d)x)}{32(b - 3d)} - \frac{9 \sinh(a - c + (b - d)x)}{32(b - d)} - \frac{\sinh(3(a - c) + 3(b - d)x)}{96(b - d)} + \frac{3 \sinh(3a - c + (3b - d)x)}{32(3b - d)} + \frac{9 \sinh(a + c + (b + d)x)}{32(b + d)} + \frac{\sinh(3(a + c) + 3(b + d)x)}{96(b + d)} - \frac{3 \sinh(3a + c + (3b + d)x)}{32(3b + d)} - \frac{3 \sinh(a + 3c + (b + 3d)x)}{32(b + 3d)}$$

output

```
3*sinh(a-3*c+(b-3*d)*x)/(32*b-96*d)-9*sinh(a-c+(b-d)*x)/(32*b-32*d)-sinh(3
*a-3*c+3*(b-d)*x)/(96*b-96*d)+3*sinh(3*a-c+(3*b-d)*x)/(96*b-32*d)+9*sinh(a
+c+(b+d)*x)/(32*b+32*d)+sinh(3*a+3*c+3*(b+d)*x)/(96*b+96*d)-3*sinh(3*a+c+(
3*b+d)*x)/(96*b+32*d)-3*sinh(a+3*c+(b+3*d)*x)/(32*b+96*d)
```

Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.91

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \frac{1}{96} \left(\frac{9 \sinh(a - 3c + bx - 3dx)}{b - 3d} - \frac{27 \sinh(a - c + bx - dx)}{b - d} - \frac{\sinh(3(a - c + bx - dx))}{b - d} + \frac{9 \sinh(3a - c + 3bx - dx)}{3b - d} - \frac{9 \sinh(3a + c + 3bx + dx)}{3b + d} - \frac{9 \sinh(a + 3c + bx + 3dx)}{b + 3d} + \frac{27 \sinh(a + c + (b + d)x)}{b + d} + \frac{\sinh(3(a + c + (b + d)x))}{b + d} \right)$$

input `Integrate[Sinh[a + b*x]^3*Sinh[c + d*x]^3,x]`

output `((9*Sinh[a - 3*c + b*x - 3*d*x])/(b - 3*d) - (27*Sinh[a - c + b*x - d*x])/(b - d) - Sinh[3*(a - c + b*x - d*x)]/(b - d) + (9*Sinh[3*a - c + 3*b*x - d*x])/(3*b - d) - (9*Sinh[3*a + c + 3*b*x + d*x])/(3*b + d) - (9*Sinh[a + 3*c + b*x + 3*d*x])/(b + 3*d) + (27*Sinh[a + c + (b + d)*x])/(b + d) + Sinh[3*(a + c + (b + d)*x)]/(b + d))/96`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6147, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx$$

↓ 6147

$$\int \left(\frac{3}{32} \cosh(a + x(b - 3d) - 3c) - \frac{9}{32} \cosh(a + x(b - d) - c) - \frac{1}{32} \cosh(3(a - c) + 3x(b - d)) + \frac{3}{32} \cosh(3a + x(3b - d) - 3c) \right. \\ \left. + \frac{9}{32} \cosh(a + x(b + d) + c) - \frac{1}{32} \cosh(3(a + c) + 3x(b + d)) + \frac{3}{32} \cosh(3a + x(3b + d) + c) \right) dx$$

↓ 2009

$$\frac{3 \sinh(a + x(b - 3d) - 3c)}{32(b - 3d)} - \frac{9 \sinh(a + x(b - d) - c)}{32(b - d)} - \frac{\sinh(3(a - c) + 3x(b - d))}{96(b - d)} + \\ \frac{3 \sinh(3a + x(3b - d) - 3c)}{32(3b - d)} + \frac{9 \sinh(a + x(b + d) + c)}{32(b + d)} + \frac{\sinh(3(a + c) + 3x(b + d))}{96(b + d)} - \\ \frac{3 \sinh(3a + x(3b + d) + c)}{32(3b + d)} - \frac{3 \sinh(a + x(b + 3d) + 3c)}{32(b + 3d)}$$

input

```
Int[Sinh[a + b*x]^3*Sinh[c + d*x]^3,x]
```

output

```
(3*Sinh[a - 3*c + (b - 3*d)*x])/(32*(b - 3*d)) - (9*Sinh[a - c + (b - d)*x])/(32*(b - d)) - Sinh[3*(a - c) + 3*(b - d)*x]/(96*(b - d)) + (3*Sinh[3*a - c + (3*b - d)*x])/(32*(3*b - d)) + (9*Sinh[a + c + (b + d)*x])/(32*(b + d)) + Sinh[3*(a + c) + 3*(b + d)*x]/(96*(b + d)) - (3*Sinh[3*a + c + (3*b + d)*x])/(32*(3*b + d)) - (3*Sinh[a + 3*c + (b + 3*d)*x])/(32*(b + 3*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6147

```
Int[Sinh[v_]^(p_.)*Sinh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Sinh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))
```

Maple [A] (verified)

Time = 26.38 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.97

method	result
default	$\frac{3 \sinh(a-3c+(b-3d)x)}{32(b-3d)} - \frac{9 \sinh(a-c+(b-d)x)}{32(b-d)} + \frac{9 \sinh(a+c+(b+d)x)}{32(b+d)} - \frac{3 \sinh(a+3c+(b+3d)x)}{32(b+3d)} - \frac{\sinh((3b-3d)x+3a-3c)}{32(3b-3d)}$
parallelrisc	$\frac{9(b-d)(b+3d)(b-3d)(b+d)\left(b+\frac{d}{3}\right) \sinh(3a-c+(3b-d)x)}{32} - \frac{9\left(\frac{(b+3d)(b-3d)(b+d)\left(b+\frac{d}{3}\right) \sinh((3b-3d)x+3a-3c)}{3} - \frac{(b-d)(b+3d)(b-3d)(b+d)}{32}\right)}{192(b+d)(b+3d)(b-d)(b-3d)}$
risc	$\frac{(b^3 e^{6bx+6a} - b^2 d e^{6bx+6a} - 9b d^2 e^{6bx+6a} + 9d^3 e^{6bx+6a} - 9b^3 e^{4bx+4a} + 27b^2 d e^{4bx+4a} + 9b d^2 e^{4bx+4a} - 27d^3 e^{4bx+4a} - 9b^3 e^{2bx+2a} + 9b^2 d e^{2bx+2a} + 9bd^2 e^{2bx+2a} - 9d^3 e^{2bx+2a}) \sinh(3a-c+(3b-d)x)}{192(b+d)(b+3d)(b-d)(b-3d)}$
orering	Expression too large to display

input

```
int(sinh(b*x+a)^3*sinh(d*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
3/32*sinh(a-3*c+(b-3*d)*x)/(b-3*d)-9/32*sinh(a-c+(b-d)*x)/(b-d)+9/32/(b+d)*sinh(a+c+(b+d)*x)-3/32/(b+3*d)*sinh(a+3*c+(b+3*d)*x)-1/32/(3*b-3*d)*sinh((3*b-3*d)*x+3*a-3*c)+3/32/(3*b-d)*sinh(3*a-c+(3*b-d)*x)-3/32/(3*b+d)*sinh(3*a+c+(3*b+d)*x)+1/32/(3*b+3*d)*sinh((3*b+3*d)*x+3*a+3*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 731 vs. 2(179) = 358.

Time = 0.10 (sec) , antiderivative size = 731, normalized size of antiderivative = 3.75

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(sinh(b*x+a)^3*sinh(d*x+c)^3,x, algorithm="fricas")
```

output

```

-1/48*(((9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(d*x + c)^3 - 9*(b^4*d - 10*b^2
*d^3 + 9*d^5)*cosh(d*x + c))*sinh(b*x + a)^3 - ((9*b^5 - 82*b^3*d^2 + 9*b*
d^4)*cosh(b*x + a)^3 + 3*(9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)*sinh
(b*x + a)^2 - 9*(9*b^5 - 10*b^3*d^2 + b*d^4)*cosh(b*x + a))*sinh(d*x + c)^
3 + 3*((9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(d*x + c)*sinh(b*x + a)^3 - 3*(8
1*b^4*d - 90*b^2*d^3 + 9*d^5 - (9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a
)^2)*cosh(d*x + c)*sinh(b*x + a))*sinh(d*x + c)^2 - 3*((81*b^4*d - 90*b^2*
d^3 + 9*d^5 - (9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2)*cosh(d*x + c
)^3 - 9*(9*b^4*d - 82*b^2*d^3 + 9*d^5 - (b^4*d - 10*b^2*d^3 + 9*d^5)*cosh(
b*x + a)^2)*cosh(d*x + c))*sinh(b*x + a) + 3*(9*(b^5 - 10*b^3*d^2 + 9*b*d^
4)*cosh(b*x + a)^3 - ((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)^3 - 9*(
9*b^5 - 10*b^3*d^2 + b*d^4)*cosh(b*x + a))*cosh(d*x + c)^2 - 3*((9*b^5 - 8
2*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)*cosh(d*x + c)^2 - 9*(b^5 - 10*b^3*d^2 +
9*b*d^4)*cosh(b*x + a))*sinh(b*x + a)^2 - 9*(9*b^5 - 82*b^3*d^2 + 9*b*d^4
)*cosh(b*x + a))*sinh(d*x + c))/((9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)
*cosh(b*x + a)^4 - 2*(9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*cosh(b*x +
a)^2*sinh(b*x + a)^2 + (9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*sinh(b*x
+ a)^4)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3577 vs. $2(172) = 344$.

Time = 16.62 (sec) , antiderivative size = 3577, normalized size of antiderivative = 18.34

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(sinh(b*x+a)**3*sinh(d*x+c)**3,x)
```

output

```
Piecewise((x*sinh(a)**3*sinh(c)**3, Eq(b, 0) & Eq(d, 0)), (3*x*sinh(a - 3*d*x)**3*sinh(c + d*x)**3/32 + 9*x*sinh(a - 3*d*x)**3*sinh(c + d*x)*cosh(c + d*x)**2/32 + 9*x*sinh(a - 3*d*x)**2*sinh(c + d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)/32 + 3*x*sinh(a - 3*d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)**3/32 - 3*x*sinh(a - 3*d*x)*sinh(c + d*x)**3*cosh(a - 3*d*x)**2/32 - 9*x*sinh(a - 3*d*x)*sinh(c + d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**2/32 - 9*x*sinh(c + d*x)**2*cosh(a - 3*d*x)**3*cosh(c + d*x)/32 - 3*x*cosh(a - 3*d*x)**3*cosh(c + d*x)**3/32 - 61*sinh(a - 3*d*x)**3*sinh(c + d*x)**2*cosh(c + d*x)/(320*d) + sinh(a - 3*d*x)**3*cosh(c + d*x)**3/(30*d) - 117*sinh(a - 3*d*x)**2*sinh(c + d*x)**3*cosh(a - 3*d*x)/(320*d) + 3*sinh(a - 3*d*x)*sinh(c + d*x)**2*cosh(a - 3*d*x)**2*cosh(c + d*x)/(20*d) - 11*sinh(a - 3*d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**3/(320*d) + sinh(c + d*x)**3*cosh(a - 3*d*x)**3/(4*d) - 3*sinh(c + d*x)*cosh(a - 3*d*x)**3*cosh(c + d*x)**2/(320*d), Eq(b, -3*d)), (5*x*sinh(a - d*x)**3*sinh(c + d*x)**3/16 - 3*x*sinh(a - d*x)**3*sinh(c + d*x)*cosh(c + d*x)**2/16 + 9*x*sinh(a - d*x)**2*sinh(c + d*x)**2*cosh(a - d*x)*cosh(c + d*x)/16 - 3*x*sinh(a - d*x)**2*cosh(a - d*x)*cosh(c + d*x)**3/16 - 3*x*sinh(a - d*x)*sinh(c + d*x)**3*cosh(a - d*x)**2/16 + 9*x*sinh(a - d*x)*sinh(c + d*x)*cosh(a - d*x)**2*cosh(c + d*x)**2/16 - 3*x*sinh(c + d*x)**2*cosh(a - d*x)**3*cosh(c + d*x)/16 + 5*x*cosh(a - d*x)**3*cosh(c + d*x)**3/16 + 3*sinh(a - d*x)**3*sinh(c + d*x)**2*cosh(c + d...
```

Maxima [F(-2)]

Exception generated.

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(sinh(b*x+a)^3*sinh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for more detail
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(179) = 358$.

Time = 0.13 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.91

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \frac{e^{(3bx+3dx+3a+3c)}}{192(b+d)} - \frac{3e^{(3bx+dx+3a+c)}}{64(3b+d)} + \frac{3e^{(3bx-dx+3a-c)}}{64(3b-d)} - \frac{e^{(3bx-3dx+3a-3c)}}{192(b-d)} - \frac{3e^{(bx+3dx+a+3c)}}{64(b+3d)} + \frac{9e^{(bx+dx+a+c)}}{64(b+d)} - \frac{9e^{(bx-dx+a-c)}}{64(b-d)} + \frac{3e^{(bx-3dx+a-3c)}}{64(b-3d)} - \frac{3e^{(-bx+3dx-a+3c)}}{64(b-3d)} + \frac{9e^{(-bx+dx-a+c)}}{64(b-d)} - \frac{9e^{(-bx-dx-a-c)}}{64(b+d)} + \frac{3e^{(-bx-3dx-a-3c)}}{64(b+3d)} + \frac{e^{(-3bx+3dx-3a+3c)}}{192(b-d)} - \frac{3e^{(-3bx+dx-3a+c)}}{64(3b-d)} + \frac{3e^{(-3bx-dx-3a-c)}}{64(3b+d)} - \frac{e^{(-3bx-3dx-3a-3c)}}{192(b+d)}$$

input `integrate(sinh(b*x+a)^3*sinh(d*x+c)^3,x, algorithm="giac")`

output $\frac{1}{192}e^{(3bx+3dx+3a+3c)}/(b+d) - \frac{3}{64}e^{(3bx+dx+3a+c)}/(3b+d) + \frac{3}{64}e^{(3bx-dx+3a-c)}/(3b-d) - \frac{1}{192}e^{(3bx-3dx+3a-3c)}/(b-d) - \frac{3}{64}e^{(bx+3dx+a+3c)}/(b+3d) + \frac{9}{64}e^{(bx+dx+a+c)}/(b+d) - \frac{9}{64}e^{(bx-dx+a-c)}/(b-d) + \frac{3}{64}e^{(bx-3dx+a-3c)}/(b-3d) - \frac{3}{64}e^{(-bx+3dx-a+3c)}/(b-3d) + \frac{9}{64}e^{(-bx+dx-a+c)}/(b-d) - \frac{9}{64}e^{(-bx-dx-a-c)}/(b+d) + \frac{3}{64}e^{(-bx-3dx-a-3c)}/(b+3d) + \frac{1}{192}e^{(-3bx+3dx-3a+3c)}/(b-d) - \frac{3}{64}e^{(-3bx+dx-3a+c)}/(3b-d) + \frac{3}{64}e^{(-3bx-dx-3a-c)}/(3b+d) - \frac{1}{192}e^{(-3bx-3dx-3a-3c)}/(b+d)$

Mupad [B] (verification not implemented)

Time = 1.62 (sec) , antiderivative size = 906, normalized size of antiderivative = 4.65

$$\begin{aligned}
\int \sinh^3(a + bx) \sinh^3(c + dx) dx = & e^{3a+c+3bx+dx} \left(\frac{-9b^3 + 3b^2d + 9bd^2 - 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& + \frac{e^{-6a-6bx}(-9b^3 - 3b^2d + 9bd^2 + 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& - \frac{e^{-2a-2bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
- & e^{3a-c+3bx-dx} \left(\frac{-9b^3 - 3b^2d + 9bd^2 + 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& + \frac{e^{-6a-6bx}(-9b^3 + 3b^2d + 9bd^2 - 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& - \frac{e^{-2a-2bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
+ & e^{3a-3c+3bx-3dx} \left(\frac{-b^3 - b^2d + 9bd^2 + 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& + \frac{e^{-6a-6bx}(-b^3 + b^2d + 9bd^2 - 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& - \frac{e^{-2a-2bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right) \\
- & e^{3a+3c+3bx+3dx} \left(\frac{-b^3 + b^2d + 9bd^2 - 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& + \frac{e^{-6a-6bx}(-b^3 - b^2d + 9bd^2 + 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& - \frac{e^{-2a-2bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right)
\end{aligned}$$

input `int(sinh(a + b*x)^3*sinh(c + d*x)^3,x)`

output

```

exp(3*a + c + 3*b*x + d*x)*((9*b*d^2 + 3*b^2*d - 9*b^3 - 3*d^3)/(576*b^4 +
64*d^4 - 640*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b*d^2 - 3*b^2*d - 9*b^3 +
3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 2*a - 2*b*x)*(9*b*d^2 +
81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 4*a
- 4*b*x)*(9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*b^
2*d^2)) - exp(3*a - c + 3*b*x - d*x)*((9*b*d^2 - 3*b^2*d - 9*b^3 + 3*d^3)/
(576*b^4 + 64*d^4 - 640*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b*d^2 + 3*b^2*d
- 9*b^3 - 3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 2*a - 2*b*x)*(
9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (
exp(- 4*a - 4*b*x)*(9*b*d^2 + 81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*d^
4 - 640*b^2*d^2)) + exp(3*a - 3*c + 3*b*x - 3*d*x)*((9*b*d^2 - b^2*d - b^3
+ 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b*d
^2 + b^2*d - b^3 - 9*d^3))/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(- 2*
a - 2*b*x)*(9*b*d^2 - 27*b^2*d - 9*b^3 + 27*d^3))/(192*b^4 + 1728*d^4 - 19
20*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))/(
192*b^4 + 1728*d^4 - 1920*b^2*d^2)) - exp(3*a + 3*c + 3*b*x + 3*d*x)*((9*b
*d^2 + b^2*d - b^3 - 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) + (exp(- 6
*a - 6*b*x)*(9*b*d^2 - b^2*d - b^3 + 9*d^3))/(192*b^4 + 1728*d^4 - 1920*b^
2*d^2) - (exp(- 2*a - 2*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))/(192*b
^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 - 27*b^2*d...

```

Reduce [F]

$$\int \sinh^3(a + bx) \sinh^3(c + dx) dx = \int \sinh(bx + a)^3 \sinh(dx + c)^3 dx$$

input

```
int(sinh(b*x+a)^3*sinh(d*x+c)^3,x)
```

output

```
int(sinh(b*x+a)^3*sinh(d*x+c)^3,x)
```

3.13 $\int \cosh(2x) \sinh(x) dx$

Optimal result	163
Mathematica [A] (verified)	163
Rubi [C] (verified)	164
Maple [A] (verified)	165
Fricas [A] (verification not implemented)	165
Sympy [A] (verification not implemented)	166
Maxima [B] (verification not implemented)	166
Giac [B] (verification not implemented)	166
Mupad [B] (verification not implemented)	167
Reduce [B] (verification not implemented)	167

Optimal result

Integrand size = 7, antiderivative size = 15

$$\int \cosh(2x) \sinh(x) dx = -\frac{\cosh(x)}{2} + \frac{1}{6} \cosh(3x)$$

output `-1/2*cosh(x)+1/6*cosh(3*x)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cosh(2x) \sinh(x) dx = -\frac{\cosh(x)}{2} + \frac{1}{6} \cosh(3x)$$

input `Integrate[Cosh[2*x]*Sinh[x],x]`

output `-1/2*Cosh[x] + Cosh[3*x]/6`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \cosh(2x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(ix) \cos(2ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(2ix) \sin(ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{6} i \cosh(3x) - \frac{1}{2} i \cosh(x) \right) \end{aligned}$$

input `Int[Cosh[2*x]*Sinh[x],x]`

output `(-I)*((-1/2*I)*Cosh[x] + (I/6)*Cosh[3*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$-\frac{\cosh(x)}{2} + \frac{\cosh(3x)}{6}$	12
parallelrisch	$\frac{\cosh(3x)}{6} - \frac{\cosh(x)}{2} + \frac{1}{3}$	13
orering	$\frac{2 \sinh(x) \sinh(2x)}{3} - \frac{\cosh(x) \cosh(2x)}{3}$	18
risch	$\frac{e^{3x}}{12} - \frac{e^x}{4} - \frac{e^{-x}}{4} + \frac{e^{-3x}}{12}$	24

input

```
int(cosh(2*x)*sinh(x),x,method=_RETURNVERBOSE)
```

output

```
-1/2*cosh(x)+1/6*cosh(3*x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \cosh(2x) \sinh(x) dx = \frac{1}{6} \cosh(x)^3 + \frac{1}{2} \cosh(x) \sinh(x)^2 - \frac{1}{2} \cosh(x)$$

input

```
integrate(cosh(2*x)*sinh(x),x, algorithm="fricas")
```

output

```
1/6*cosh(x)^3 + 1/2*cosh(x)*sinh(x)^2 - 1/2*cosh(x)
```

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \cosh(2x) \sinh(x) dx = \frac{2 \sinh(x) \sinh(2x)}{3} - \frac{\cosh(x) \cosh(2x)}{3}$$

input `integrate(cosh(2*x)*sinh(x),x)`

output `2*sinh(x)*sinh(2*x)/3 - cosh(x)*cosh(2*x)/3`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(11) = 22.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \cosh(2x) \sinh(x) dx = -\frac{1}{12} (3 e^{(-2x)} - 1) e^{(3x)} - \frac{1}{4} e^{(-x)} + \frac{1}{12} e^{(-3x)}$$

input `integrate(cosh(2*x)*sinh(x),x, algorithm="maxima")`

output `-1/12*(3*e^(-2*x) - 1)*e^(3*x) - 1/4*e^(-x) + 1/12*e^(-3*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(11) = 22.

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.67

$$\int \cosh(2x) \sinh(x) dx = -\frac{1}{12} (3 e^{(2x)} - 1) e^{(-3x)} + \frac{1}{12} e^{(3x)} - \frac{1}{4} e^x$$

input `integrate(cosh(2*x)*sinh(x),x, algorithm="giac")`

output `-1/12*(3*e^(2*x) - 1)*e^(-3*x) + 1/12*e^(3*x) - 1/4*e^x`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \cosh(2x) \sinh(x) dx = \frac{2 \cosh(x)^3}{3} - \cosh(x)$$

input `int(cosh(2*x)*sinh(x),x)`output `(2*cosh(x)^3)/3 - cosh(x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cosh(2x) \sinh(x) dx = -\frac{\cosh(2x) \cosh(x)}{3} + \frac{2 \sinh(2x) \sinh(x)}{3}$$

input `int(cosh(2*x)*sinh(x),x)`output `(- cosh(2*x)*cosh(x) + 2*sinh(2*x)*sinh(x))/3`

3.14 $\int \cosh(3x) \sinh(x) dx$

Optimal result	168
Mathematica [A] (verified)	168
Rubi [C] (verified)	169
Maple [A] (verified)	170
Fricas [B] (verification not implemented)	170
Sympy [A] (verification not implemented)	171
Maxima [B] (verification not implemented)	171
Giac [A] (verification not implemented)	171
Mupad [B] (verification not implemented)	172
Reduce [B] (verification not implemented)	172

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(3x) \sinh(x) dx = -\frac{1}{4} \cosh(2x) + \frac{1}{8} \cosh(4x)$$

output `-1/4*cosh(2*x)+1/8*cosh(4*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(3x) \sinh(x) dx = -\frac{1}{2} \cosh^2(x) + \frac{1}{8} \cosh(4x)$$

input `Integrate[Cosh[3*x]*Sinh[x],x]`

output `-1/2*Cosh[x]^2 + Cosh[4*x]/8`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \cosh(3x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(ix) \cos(3ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(3ix) \sin(ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{8} i \cosh(4x) - \frac{1}{4} i \cosh(2x) \right) \end{aligned}$$

input `Int[Cosh[3*x]*Sinh[x],x]`

output `(-I)*((-1/4*I)*Cosh[2*x] + (I/8)*Cosh[4*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\cosh(2x)}{4} + \frac{\cosh(4x)}{8}$	14
parallelrisch	$\frac{\cosh(4x)}{8} + \frac{1}{8} - \frac{\cosh(2x)}{4}$	15
orering	$\frac{3 \sinh(x) \sinh(3x)}{8} - \frac{\cosh(x) \cosh(3x)}{8}$	18
risch	$\frac{e^{4x}}{16} - \frac{e^{2x}}{8} - \frac{e^{-2x}}{8} + \frac{e^{-4x}}{16}$	26

input

```
int(cosh(3*x)*sinh(x),x,method=_RETURNVERBOSE)
```

output

```
-1/4*cosh(2*x)+1/8*cosh(4*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.94

$$\int \cosh(3x) \sinh(x) dx = \frac{1}{8} \cosh(x)^4 + \frac{1}{8} \sinh(x)^4 + \frac{1}{4} (3 \cosh(x)^2 - 1) \sinh(x)^2 - \frac{1}{4} \cosh(x)^2$$

input

```
integrate(cosh(3*x)*sinh(x),x, algorithm="fricas")
```

output

```
1/8*cosh(x)^4 + 1/8*sinh(x)^4 + 1/4*(3*cosh(x)^2 - 1)*sinh(x)^2 - 1/4*cosh
(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(3x) \sinh(x) dx = \frac{3 \sinh(x) \sinh(3x)}{8} - \frac{\cosh(x) \cosh(3x)}{8}$$

input `integrate(cosh(3*x)*sinh(x),x)`

output `3*sinh(x)*sinh(3*x)/8 - cosh(x)*cosh(3*x)/8`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(3x) \sinh(x) dx = -\frac{1}{16} (2e^{(-2x)} - 1)e^{(4x)} - \frac{1}{8} e^{(-2x)} + \frac{1}{16} e^{(-4x)}$$

input `integrate(cosh(3*x)*sinh(x),x, algorithm="maxima")`

output `-1/16*(2*e^(-2*x) - 1)*e^(4*x) - 1/8*e^(-2*x) + 1/16*e^(-4*x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int \cosh(3x) \sinh(x) dx = \frac{1}{16} (e^{(2x)} + e^{(-2x)})^2 - \frac{1}{8} e^{(2x)} - \frac{1}{8} e^{(-2x)}$$

input `integrate(cosh(3*x)*sinh(x),x, algorithm="giac")`

output `1/16*(e^(2*x) + e^(-2*x))^2 - 1/8*e^(2*x) - 1/8*e^(-2*x)`

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \cosh(3x) \sinh(x) dx = \sinh(x)^4 + \frac{\sinh(x)^2}{2}$$

input `int(cosh(3*x)*sinh(x),x)`

output `sinh(x)^2/2 + sinh(x)^4`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(3x) \sinh(x) dx = -\frac{\cosh(3x) \cosh(x)}{8} + \frac{3 \sinh(3x) \sinh(x)}{8}$$

input `int(cosh(3*x)*sinh(x),x)`

output `(- cosh(3*x)*cosh(x) + 3*sinh(3*x)*sinh(x))/8`

3.15 $\int \cosh(4x) \sinh(x) dx$

Optimal result	173
Mathematica [A] (verified)	173
Rubi [C] (verified)	174
Maple [A] (verified)	175
Fricas [B] (verification not implemented)	175
Sympy [A] (verification not implemented)	176
Maxima [B] (verification not implemented)	176
Giac [B] (verification not implemented)	176
Mupad [B] (verification not implemented)	177
Reduce [B] (verification not implemented)	177

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(4x) \sinh(x) dx = -\frac{1}{6} \cosh(3x) + \frac{1}{10} \cosh(5x)$$

output `-1/6*cosh(3*x)+1/10*cosh(5*x)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(4x) \sinh(x) dx = -\frac{1}{6} \cosh(3x) + \frac{1}{10} \cosh(5x)$$

input `Integrate[Cosh[4*x]*Sinh[x],x]`

output `-1/6*Cosh[3*x] + Cosh[5*x]/10`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \cosh(4x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(ix) \cos(4ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(4ix) \sin(ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{10} i \cosh(5x) - \frac{1}{6} i \cosh(3x) \right) \end{aligned}$$

input `Int[Cosh[4*x]*Sinh[x],x]`

output `(-I)*((-1/6*I)*Cosh[3*x] + (I/10)*Cosh[5*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{\cosh(3x)}{6} + \frac{\cosh(5x)}{10}$	14
parallelrisch	$\frac{\cosh(5x)}{10} - \frac{\cosh(3x)}{6} + \frac{1}{15}$	15
orering	$\frac{4 \sinh(x) \sinh(4x)}{15} - \frac{\cosh(x) \cosh(4x)}{15}$	18
risch	$\frac{e^{5x}}{20} - \frac{e^{3x}}{12} - \frac{e^{-3x}}{12} + \frac{e^{-5x}}{20}$	26

input

```
int(cosh(4*x)*sinh(x),x,method=_RETURNVERBOSE)
```

output

```
-1/6*cosh(3*x)+1/10*cosh(5*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(13) = 26$.

Time = 0.08 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.24

$$\int \cosh(4x) \sinh(x) dx = \frac{1}{10} \cosh(x)^5 + \frac{1}{2} \cosh(x) \sinh(x)^4 - \frac{1}{6} \cosh(x)^3 + \frac{1}{2} (2 \cosh(x)^3 - \cosh(x)) \sinh(x)^2$$

input

```
integrate(cosh(4*x)*sinh(x),x, algorithm="fricas")
```

output

```
1/10*cosh(x)^5 + 1/2*cosh(x)*sinh(x)^4 - 1/6*cosh(x)^3 + 1/2*(2*cosh(x)^3
- cosh(x))*sinh(x)^2
```


Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(4x) \sinh(x) dx = \frac{4 \sinh(x) \sinh(4x)}{15} - \frac{\cosh(x) \cosh(4x)}{15}$$

input `integrate(cosh(4*x)*sinh(x),x)`

output `4*sinh(x)*sinh(4*x)/15 - cosh(x)*cosh(4*x)/15`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(4x) \sinh(x) dx = -\frac{1}{60} (5 e^{(-2x)} - 3) e^{(5x)} - \frac{1}{12} e^{(-3x)} + \frac{1}{20} e^{(-5x)}$$

input `integrate(cosh(4*x)*sinh(x),x, algorithm="maxima")`

output `-1/60*(5*e^(-2*x) - 3)*e^(5*x) - 1/12*e^(-3*x) + 1/20*e^(-5*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(4x) \sinh(x) dx = -\frac{1}{60} (5 e^{(2x)} - 3) e^{(-5x)} + \frac{1}{20} e^{(5x)} - \frac{1}{12} e^{(3x)}$$

input `integrate(cosh(4*x)*sinh(x),x, algorithm="giac")`

output `-1/60*(5*e^(2*x) - 3)*e^(-5*x) + 1/20*e^(5*x) - 1/12*e^(3*x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \cosh(4x) \sinh(x) dx = \frac{8 \cosh(x)^5}{5} - \frac{8 \cosh(x)^3}{3} + \cosh(x)$$

input `int(cosh(4*x)*sinh(x),x)`output `cosh(x) - (8*cosh(x)^3)/3 + (8*cosh(x)^5)/5`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(4x) \sinh(x) dx = -\frac{\cosh(4x) \cosh(x)}{15} + \frac{4 \sinh(4x) \sinh(x)}{15}$$

input `int(cosh(4*x)*sinh(x),x)`output `(- cosh(4*x)*cosh(x) + 4*sinh(4*x)*sinh(x))/15`

3.16 $\int \cosh(mx) \sinh(x) dx$

Optimal result	178
Mathematica [A] (verified)	178
Rubi [A] (verified)	179
Maple [A] (verified)	180
Fricas [A] (verification not implemented)	180
Sympy [A] (verification not implemented)	181
Maxima [F(-2)]	181
Giac [B] (verification not implemented)	181
Mupad [B] (verification not implemented)	182
Reduce [B] (verification not implemented)	182

Optimal result

Integrand size = 7, antiderivative size = 35

$$\int \cosh(mx) \sinh(x) dx = \frac{\cosh((1-m)x)}{2(1-m)} + \frac{\cosh((1+m)x)}{2(1+m)}$$

output

```
cosh((1-m)*x)/(2-2*m)+cosh((1+m)*x)/(2+2*m)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(mx) \sinh(x) dx = \frac{-\cosh(x) \cosh(mx) + m \sinh(x) \sinh(mx)}{-1 + m^2}$$

input

```
Integrate[Cosh[m*x]*Sinh[x],x]
```

output

```
(-(Cosh[x]*Cosh[m*x]) + m*Sinh[x]*Sinh[m*x])/(-1 + m^2)
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \cosh(mx) dx$$

$$\downarrow \text{6152}$$

$$\int \left(\frac{1}{2} \sinh((1-m)x) + \frac{1}{2} \sinh((m+1)x) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\cosh((1-m)x)}{2(1-m)} + \frac{\cosh((m+1)x)}{2(m+1)}$$

input `Int[Cosh[m*x]*Sinh[x],x]`

output `Cosh[(1 - m)*x]/(2*(1 - m)) + Cosh[(1 + m)*x]/(2*(1 + m))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$-\frac{\cosh(x(-1+m))}{2(-1+m)} + \frac{\cosh((1+m)x)}{2+2m}$
parallelrisch	$\frac{(-1-m)\cosh(x(-1+m))+2+(-1+m)\cosh((1+m)x)}{2m^2-2}$
risch	$\frac{(m e^{2x}-e^{2x}-m-1)e^{x(-1+m)}}{4(1+m)(-1+m)} - \frac{(m e^{2x}+e^{2x}-m+1)e^{-(1+m)x}}{4(1+m)(-1+m)}$
orering	$\frac{2(m^2+1)(m \sinh(mx) \sinh(x)+\cosh(x) \cosh(mx))}{m^4-2m^2+1} - \frac{m^3 \sinh(mx) \sinh(x)+3m^2 \cosh(mx) \cosh(x)+3m \sinh(mx) \sinh(x)}{m^4-2m^2+1}$

input `int(cosh(m*x)*sinh(x),x,method=_RETURNVERBOSE)`output `-1/2*cosh(x*(-1+m))/(-1+m)+1/2*cosh((1+m)*x)/(1+m)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \cosh(mx) \sinh(x) dx = \frac{m \sinh(mx) \sinh(x) - \cosh(mx) \cosh(x)}{(m^2 - 1) \cosh(x)^2 - (m^2 - 1) \sinh(x)^2}$$

input `integrate(cosh(m*x)*sinh(x),x, algorithm="fricas")`output `(m*sinh(m*x)*sinh(x) - cosh(m*x)*cosh(x))/((m^2 - 1)*cosh(x)^2 - (m^2 - 1)*sinh(x)^2)`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \cosh(mx) \sinh(x) dx = \begin{cases} \frac{\sinh^2(x)}{2} & \text{for } m = -1 \vee m = 1 \\ \frac{m \sinh(x) \sinh(mx)}{m^2-1} - \frac{\cosh(x) \cosh(mx)}{m^2-1} & \text{otherwise} \end{cases}$$

input `integrate(cosh(m*x)*sinh(x),x)`

output `Piecewise((sinh(x)**2/2, Eq(m, -1) | Eq(m, 1)), (m*sinh(x)*sinh(m*x)/(m**2 - 1) - cosh(x)*cosh(m*x)/(m**2 - 1), True))`

Maxima [F(-2)]

Exception generated.

$$\int \cosh(mx) \sinh(x) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(m*x)*sinh(x),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see `assume?` for more details)Is`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(28) = 56.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \cosh(mx) \sinh(x) dx = \frac{e^{(mx+x)}}{4(m+1)} - \frac{e^{(mx-x)}}{4(m-1)} - \frac{e^{(-mx+x)}}{4(m-1)} + \frac{e^{(-mx-x)}}{4(m+1)}$$

input `integrate(cosh(m*x)*sinh(x),x, algorithm="giac")`

output $\frac{1}{4}e^{(m+1)x}/(m+1) - \frac{1}{4}e^{(m-1)x}/(m-1) - \frac{1}{4}e^{(-m+1)x}/(m-1) + \frac{1}{4}e^{(-m-1)x}/(m+1)$

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \cosh(mx) \sinh(x) dx = -\frac{\cosh(mx) \cosh(x) - m \sinh(mx) \sinh(x)}{m^2 - 1}$$

input `int(cosh(m*x)*sinh(x),x)`

output $-(\cosh(m*x)*\cosh(x) - m*\sinh(m*x)*\sinh(x))/(m^2 - 1)$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(mx) \sinh(x) dx = \frac{-\cosh(mx) \cosh(x) + \sinh(mx) \sinh(x) m}{m^2 - 1}$$

input `int(cosh(m*x)*sinh(x),x)`

output $(-\cosh(m*x)*\cosh(x) + \sinh(m*x)*\sinh(x)*m)/(m**2 - 1)$

3.17 $\int \cosh(c + dx) \sinh(a + bx) dx$

Optimal result	183
Mathematica [A] (verified)	183
Rubi [A] (verified)	184
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	185
Sympy [B] (verification not implemented)	186
Maxima [F(-2)]	186
Giac [B] (verification not implemented)	187
Mupad [B] (verification not implemented)	187
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 13, antiderivative size = 43

$$\int \cosh(c + dx) \sinh(a + bx) dx = \frac{\cosh(a - c + (b - d)x)}{2(b - d)} + \frac{\cosh(a + c + (b + d)x)}{2(b + d)}$$

output

```
cosh(a-c+(b-d)*x)/(2*b-2*d)+cosh(a+c+(b+d)*x)/(2*b+2*d)
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \cosh(c + dx) \sinh(a + bx) dx = \frac{\cosh(a - c + (b - d)x)}{2(b - d)} + \frac{\cosh(a + c + (b + d)x)}{2(b + d)}$$

input

```
Integrate[Cosh[c + d*x]*Sinh[a + b*x],x]
```

output

```
Cosh[a - c + (b - d)*x]/(2*(b - d)) + Cosh[a + c + (b + d)*x]/(2*(b + d))
```


Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \cosh(c + dx) dx$$

$$\downarrow \text{6152}$$

$$\int \left(\frac{1}{2} \sinh(a + x(b - d) - c) + \frac{1}{2} \sinh(a + x(b + d) + c) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\cosh(a + x(b - d) - c)}{2(b - d)} + \frac{\cosh(a + x(b + d) + c)}{2(b + d)}$$

input `Int[Cosh[c + d*x]*Sinh[a + b*x],x]`

output `Cosh[a - c + (b - d)*x]/(2*(b - d)) + Cosh[a + c + (b + d)*x]/(2*(b + d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

method	result
default	$\frac{\cosh(a-c+(b-d)x)}{2b-2d} + \frac{\cosh(a+c+(b+d)x)}{2b+2d}$
risch	$\frac{(b e^{2bx+2a} - d e^{2bx+2a} + b+d)e^{-bx+dx-a+c}}{4(b+d)(b-d)} + \frac{(b e^{2bx+2a} + d e^{2bx+2a} + b-d)e^{-bx-dx-a-c}}{4(b+d)(b-d)}$
parallelrisch	$\frac{2b+2 \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^2 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^2 b - 4d \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)}{(b^2-d^2)\left(\tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^2 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1\right)}$
orering	$\frac{2(b^2+d^2)(d \sinh(dx+c) \sinh(bx+a) + \cosh(dx+c) b \cosh(bx+a))}{b^4-2b^2d^2+d^4} - \frac{d^3 \sinh(dx+c) \sinh(bx+a) + 3d^2 \cosh(dx+c) b \cosh(bx+a)}{b^4-2b^2d^2+d^4}$

input `int(cosh(d*x+c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*cosh(a-c+(b-d)*x)/(b-d)+1/2*cosh(a+c+(b+d)*x)/(b+d)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.65

$$\int \cosh(c+dx) \sinh(a+bx) dx$$

$$= \frac{b \cosh(bx+a) \cosh(dx+c) - d \sinh(bx+a) \sinh(dx+c)}{(b^2-d^2) \cosh(bx+a)^2 - (b^2-d^2) \sinh(bx+a)^2}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a),x, algorithm="fricas")`

output `(b*cosh(b*x + a)*cosh(d*x + c) - d*sinh(b*x + a)*sinh(d*x + c))/((b^2 - d^2)*cosh(b*x + a)^2 - (b^2 - d^2)*sinh(b*x + a)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(32) = 64$.

Time = 0.32 (sec) , antiderivative size = 153, normalized size of antiderivative = 3.56

$$\int \cosh(c + dx) \sinh(a + bx) dx$$

$$= \begin{cases} x \sinh(a) \cosh(c) & \text{for } b = 0 \wedge d = 0 \\ \frac{x \sinh(a-dx) \cosh(c+dx)}{2} + \frac{x \sinh(c+dx) \cosh(a-dx)}{2} + \frac{\sinh(a-dx) \sinh(c+dx)}{2d} & \text{for } b = -d \\ \frac{x \sinh(a+dx) \cosh(c+dx)}{2} - \frac{x \sinh(c+dx) \cosh(a+dx)}{2} + \frac{\sinh(a+dx) \sinh(c+dx)}{2d} & \text{for } b = d \\ \frac{b \cosh(a+bx) \cosh(c+dx)}{b^2-d^2} - \frac{d \sinh(a+bx) \sinh(c+dx)}{b^2-d^2} & \text{otherwise} \end{cases}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a),x)`

output `Piecewise((x*sinh(a)*cosh(c), Eq(b, 0) & Eq(d, 0)), (x*sinh(a - d*x)*cosh(c + d*x)/2 + x*sinh(c + d*x)*cosh(a - d*x)/2 + sinh(a - d*x)*sinh(c + d*x)/(2*d), Eq(b, -d)), (x*sinh(a + d*x)*cosh(c + d*x)/2 - x*sinh(c + d*x)*cosh(a + d*x)/2 + sinh(a + d*x)*sinh(c + d*x)/(2*d), Eq(b, d)), (b*cosh(a + b*x)*cosh(c + d*x)/(b**2 - d**2) - d*sinh(a + b*x)*sinh(c + d*x)/(b**2 - d**2), True))`

Maxima [F(-2)]

Exception generated.

$$\int \cosh(c + dx) \sinh(a + bx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-d/b>0)', see `assume?` for more details)I`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(39) = 78.

Time = 0.12 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.98

$$\int \cosh(c + dx) \sinh(a + bx) dx = \frac{e^{(bx+dx+a+c)}}{4(b+d)} + \frac{e^{(bx-dx+a-c)}}{4(b-d)} + \frac{e^{(-bx+dx-a+c)}}{4(b-d)} + \frac{e^{(-bx-dx-a-c)}}{4(b+d)}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a),x, algorithm="giac")`

output `1/4*e^(b*x + d*x + a + c)/(b + d) + 1/4*e^(b*x - d*x + a - c)/(b - d) + 1/4*e^(-b*x + d*x - a + c)/(b - d) + 1/4*e^(-b*x - d*x - a - c)/(b + d)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \cosh(c + dx) \sinh(a + bx) dx = \frac{b \cosh(a + bx) \cosh(c + dx) - d \sinh(a + bx) \sinh(c + dx)}{b^2 - d^2}$$

input `int(cosh(c + d*x)*sinh(a + b*x),x)`

output `(b*cosh(a + b*x)*cosh(c + d*x) - d*sinh(a + b*x)*sinh(c + d*x))/(b^2 - d^2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \cosh(c + dx) \sinh(a + bx) dx$$
$$= \frac{\cosh(bx + a) \cosh(dx + c) b - \sinh(bx + a) \sinh(dx + c) d}{b^2 - d^2}$$

input `int(cosh(d*x+c)*sinh(b*x+a),x)`

output `(cosh(a + b*x)*cosh(c + d*x)*b - sinh(a + b*x)*sinh(c + d*x)*d)/(b**2 - d**2)`

3.18 $\int \cosh^2(c + dx) \sinh(a + bx) dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	191
Fricas [B] (verification not implemented)	191
Sympy [B] (verification not implemented)	192
Maxima [F(-2)]	193
Giac [B] (verification not implemented)	193
Mupad [B] (verification not implemented)	194
Reduce [B] (verification not implemented)	194

Optimal result

Integrand size = 15, antiderivative size = 62

$$\int \cosh^2(c + dx) \sinh(a + bx) dx = \frac{\cosh(a + bx)}{2b} + \frac{\cosh(a - 2c + (b - 2d)x)}{4(b - 2d)} + \frac{\cosh(a + 2c + (b + 2d)x)}{4(b + 2d)}$$

output

```
1/2*cosh(b*x+a)/b+cosh(a-2*c+(b-2*d)*x)/(4*b-8*d)+cosh(a+2*c+(b+2*d)*x)/(4*b+8*d)
```

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

$$\int \cosh^2(c + dx) \sinh(a + bx) dx = \frac{1}{4} \left(\frac{2 \cosh(a) \cosh(bx)}{b} + \frac{\cosh(a - 2c + bx - 2dx)}{b - 2d} + \frac{\cosh(a + 2c + bx + 2dx)}{b + 2d} + \frac{2 \sinh(a) \sinh(bx)}{b} \right)$$

input

```
Integrate[Cosh[c + d*x]^2*Sinh[a + b*x],x]
```

output

$$\frac{((2*\text{Cosh}[a]*\text{Cosh}[b*x])/b + \text{Cosh}[a - 2*c + b*x - 2*d*x]/(b - 2*d) + \text{Cosh}[a + 2*c + b*x + 2*d*x]/(b + 2*d) + (2*\text{Sinh}[a]*\text{Sinh}[b*x])/b)/4}$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \cosh^2(c + dx) dx$$

$$\downarrow \text{6152}$$

$$\int \left(\frac{1}{4} \sinh(a + x(b - 2d) - 2c) + \frac{1}{4} \sinh(a + x(b + 2d) + 2c) + \frac{1}{2} \sinh(a + bx) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\cosh(a + x(b - 2d) - 2c)}{4(b - 2d)} + \frac{\cosh(a + x(b + 2d) + 2c)}{4(b + 2d)} + \frac{\cosh(a + bx)}{2b}$$

input

```
Int[Cosh[c + d*x]^2*Sinh[a + b*x],x]
```

output

```
Cosh[a + b*x]/(2*b) + Cosh[a - 2*c + (b - 2*d)*x]/(4*(b - 2*d)) + Cosh[a + 2*c + (b + 2*d)*x]/(4*(b + 2*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6152

```
Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] :> Int[ExpandTrigReduce[Sinh[v]
]^(p)*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

method	result
default	$\frac{\cosh(bx+a)}{2b} + \frac{\cosh(a-2c+(b-2d)x)}{4b-8d} + \frac{\cosh(a+2c+(b+2d)x)}{4b+8d}$
parallelrisch	$\frac{b(b+2d)\cosh(a-2c+(b-2d)x)+b(b-2d)\cosh(a+2c+(b+2d)x)+(2b^2-8d^2)\cosh(bx+a)+4b^2-8d^2}{4b^3-16bd^2}$
risch	$\frac{e^{bx+a}}{4b} + \frac{e^{-bx-a}}{4b} + \frac{(be^{2bx+2a}-2de^{2bx+2a}+b+2d)e^{-bx+2dx-a+2c}}{8(b+2d)(b-2d)} + \frac{(be^{2bx+2a}+2de^{2bx+2a}+b-2d)e^{-bx-2dx-a-2c}}{8(b+2d)(b-2d)}$
orering	$\frac{(3b^4+16d^4)\left(2\sinh(bx+a)\sinh(dx+c)d\cosh(dx+c)+\cosh(dx+c)^2b\cosh(bx+a)\right)}{(b^4-8b^2d^2+16d^4)b^2} - \frac{(3b^2+8d^2)\left(6b^2\sinh(bx+a)\cosh(dx+c)\right)}{(b^4-8b^2d^2+16d^4)b^2}$

input

```
int(cosh(d*x+c)^2*sinh(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
1/2*cosh(b*x+a)/b+1/4/(b-2*d)*cosh(a-2*c+(b-2*d)*x)+1/4/(b+2*d)*cosh(a+2*c
+(b+2*d)*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(56) = 112.

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.92

$$\int \cosh^2(c+dx)\sinh(a+bx)dx$$

$$= \frac{b^2 \cosh(bx+a)\cosh(dx+c)^2 - 4bd \cosh(dx+c)\sinh(bx+a)\sinh(dx+c) + b^2 \cosh(bx+a)\sinh(dx+c)}{2((b^3-4bd^2)\cosh(bx+a)^2 - (b^3-4bd^2)\sinh(bx+a)^2)}$$

input

```
integrate(cosh(d*x+c)^2*sinh(b*x+a), x, algorithm="fricas")
```


output

```
1/2*(b^2*cosh(b*x + a)*cosh(d*x + c)^2 - 4*b*d*cosh(d*x + c)*sinh(b*x + a)
*sinh(d*x + c) + b^2*cosh(b*x + a)*sinh(d*x + c)^2 + (b^2 - 4*d^2)*cosh(b*
x + a))/((b^3 - 4*b*d^2)*cosh(b*x + a)^2 - (b^3 - 4*b*d^2)*sinh(b*x + a)^2
)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(49) = 98$.

Time = 0.69 (sec) , antiderivative size = 408, normalized size of antiderivative = 6.58

$$\int \cosh^2(c + dx) \sinh(a + bx) dx$$

$$= \begin{cases} x \sinh(a) \cosh^2(c) \\ \left(-\frac{x \sinh^2(c+dx)}{2} + \frac{x \cosh^2(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right) \sinh(a) \\ \frac{x \sinh(a-2dx) \sinh^2(c+dx)}{4} + \frac{x \sinh(a-2dx) \cosh^2(c+dx)}{4} + \frac{x \sinh(c+dx) \cosh(a-2dx) \cosh(c+dx)}{2} + \frac{3 \sinh(a-2dx) \sinh(c+dx) \cosh(c+dx)}{4d} \\ \frac{x \sinh(a+2dx) \sinh^2(c+dx)}{4} + \frac{x \sinh(a+2dx) \cosh^2(c+dx)}{4} - \frac{x \sinh(c+dx) \cosh(a+2dx) \cosh(c+dx)}{2} + \frac{3 \sinh(a+2dx) \sinh(c+dx) \cosh(c+dx)}{4d} \\ \frac{b^2 \cosh(a+bx) \cosh^2(c+dx)}{b^3-4bd^2} - \frac{2bd \sinh(a+bx) \sinh(c+dx) \cosh(c+dx)}{b^3-4bd^2} + \frac{2d^2 \sinh^2(c+dx) \cosh(a+bx)}{b^3-4bd^2} - \frac{2d^2 \cosh(a+bx) \cosh^2(c+dx)}{b^3-4bd^2} \end{cases}$$

input

```
integrate(cosh(d*x+c)**2*sinh(b*x+a),x)
```

output

```
Piecewise((x*sinh(a)*cosh(c)**2, Eq(b, 0) & Eq(d, 0)), ((-x*sinh(c + d*x)*
**2/2 + x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*sinh(a),
Eq(b, 0)), (x*sinh(a - 2*d*x)*sinh(c + d*x)**2/4 + x*sinh(a - 2*d*x)*cosh(
c + d*x)**2/4 + x*sinh(c + d*x)*cosh(a - 2*d*x)*cosh(c + d*x)/2 + 3*sinh(a
- 2*d*x)*sinh(c + d*x)*cosh(c + d*x)/(4*d) + sinh(c + d*x)**2*cosh(a - 2*
d*x)/(2*d), Eq(b, -2*d)), (x*sinh(a + 2*d*x)*sinh(c + d*x)**2/4 + x*sinh(a
+ 2*d*x)*cosh(c + d*x)**2/4 - x*sinh(c + d*x)*cosh(a + 2*d*x)*cosh(c + d*
x)/2 + 3*sinh(a + 2*d*x)*sinh(c + d*x)*cosh(c + d*x)/(4*d) - sinh(c + d*x)
**2*cosh(a + 2*d*x)/(2*d), Eq(b, 2*d)), (b**2*cosh(a + b*x)*cosh(c + d*x)*
**2/(b**3 - 4*b*d**2) - 2*b*d*sinh(a + b*x)*sinh(c + d*x)*cosh(c + d*x)/(b*
*3 - 4*b*d**2) + 2*d**2*sinh(c + d*x)**2*cosh(a + b*x)/(b**3 - 4*b*d**2) -
2*d**2*cosh(a + b*x)*cosh(c + d*x)**2/(b**3 - 4*b*d**2), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^2(c + dx) \sinh(a + bx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(2*d)/b>0)', see `assume?` for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 0.11 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.94

$$\int \cosh^2(c + dx) \sinh(a + bx) dx = \frac{e^{(bx+2dx+a+2c)}}{8(b+2d)} + \frac{e^{(bx-2dx+a-2c)}}{8(b-2d)} + \frac{e^{(bx+a)}}{4b} \\ + \frac{e^{(-bx+2dx-a+2c)}}{8(b-2d)} + \frac{e^{(-bx-2dx-a-2c)}}{8(b+2d)} + \frac{e^{(-bx-a)}}{4b}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a),x, algorithm="giac")`

output $\frac{1}{8}e^{(bx + 2dx + a + 2c)}/(b + 2d) + \frac{1}{8}e^{(bx - 2dx + a - 2c)}/(b - 2d) + \frac{1}{4}e^{(bx + a)}/b + \frac{1}{8}e^{(-bx + 2dx - a + 2c)}/(b - 2d) + \frac{1}{8}e^{(-bx - 2dx - a - 2c)}/(b + 2d) + \frac{1}{4}e^{(-bx - a)}/b$

Mupad [B] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.10

$$\int \cosh^2(c + dx) \sinh(a + bx) dx$$

$$= \frac{2d^2 \cosh(a + bx) - b^2 \cosh(a + bx) \cosh(c + dx)^2 + 2bd \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)}{4bd^2 - b^3}$$

input `int(cosh(c + d*x)^2*sinh(a + b*x),x)`output `(2*d^2*cosh(a + b*x) - b^2*cosh(a + b*x)*cosh(c + d*x)^2 + 2*b*d*cosh(c + d*x)*sinh(a + b*x)*sinh(c + d*x))/(4*b*d^2 - b^3)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 212, normalized size of antiderivative = 3.42

$$\int \cosh^2(c + dx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+4dx+2a+4c}b^2 - 2e^{2bx+4dx+2a+4c}bd + 2e^{2bx+2dx+2a+2c}b^2 - 8e^{2bx+2dx+2a+2c}d^2 + e^{2bx+2a}b^2 + 2e^{2bx+2a}bd + 8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}{8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}$$

input `int(cosh(d*x+c)^2*sinh(b*x+a),x)`output `(e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2 - 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d + 2*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2 - 8*e**(2*a + 2*b*x + 2*c + 2*d*x)*d**2 + e**(2*a + 2*b*x)*b**2 + 2*e**(2*a + 2*b*x)*b*d + e**(4*c + 4*d*x)*b**2 + 2*e**(4*c + 4*d*x)*b*d + 2*e**(2*c + 2*d*x)*b**2 - 8*e**(2*c + 2*d*x)*d**2 + b**2 - 2*b*d)/(8*e**(a + b*x + 2*c + 2*d*x)*b*(b**2 - 4*d**2))`

3.19 $\int \cosh^3(c + dx) \sinh(a + bx) dx$

Optimal result	195
Mathematica [A] (verified)	195
Rubi [A] (verified)	196
Maple [A] (verified)	197
Fricas [B] (verification not implemented)	197
Sympy [B] (verification not implemented)	198
Maxima [F(-2)]	199
Giac [B] (verification not implemented)	200
Mupad [B] (verification not implemented)	200
Reduce [B] (verification not implemented)	201

Optimal result

Integrand size = 15, antiderivative size = 91

$$\int \cosh^3(c + dx) \sinh(a + bx) dx = \frac{\cosh(a - 3c + (b - 3d)x)}{8(b - 3d)} + \frac{3 \cosh(a - c + (b - d)x)}{8(b - d)} + \frac{3 \cosh(a + c + (b + d)x)}{8(b + d)} + \frac{\cosh(a + 3c + (b + 3d)x)}{8(b + 3d)}$$

output

```
cosh(a-3*c+(b-3*d)*x)/(8*b-24*d)+3*cosh(a-c+(b-d)*x)/(8*b-8*d)+3*cosh(a+c+(b+d)*x)/(8*b+8*d)+cosh(a+3*c+(b+3*d)*x)/(8*b+24*d)
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.93

$$\int \cosh^3(c + dx) \sinh(a + bx) dx = \frac{1}{8} \left(\frac{\cosh(a - 3c + bx - 3dx)}{b - 3d} + \frac{3 \cosh(a - c + bx - dx)}{b - d} + \frac{\cosh(a + 3c + bx + 3dx)}{b + 3d} + \frac{3 \cosh(a + c + (b + d)x)}{b + d} \right)$$

input `Integrate[Cosh[c + d*x]^3*Sinh[a + b*x],x]`

output $(\text{Cosh}[a - 3c + b*x - 3*d*x]/(b - 3*d) + (3*\text{Cosh}[a - c + b*x - d*x])/(b - d) + \text{Cosh}[a + 3*c + b*x + 3*d*x]/(b + 3*d) + (3*\text{Cosh}[a + c + (b + d)*x])/(b + d))/8$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \cosh^3(c + dx) dx$$

$$\downarrow 6152$$

$$\int \left(\frac{1}{8} \sinh(a + x(b - 3d) - 3c) + \frac{3}{8} \sinh(a + x(b - d) - c) + \frac{3}{8} \sinh(a + x(b + d) + c) + \frac{1}{8} \sinh(a + x(b + 3d) + 3c) \right) dx$$

$$\downarrow 2009$$

$$\frac{\cosh(a + x(b - 3d) - 3c)}{8(b - 3d)} + \frac{3 \cosh(a + x(b - d) - c)}{8(b - d)} + \frac{3 \cosh(a + x(b + d) + c)}{8(b + d)} + \frac{\cosh(a + x(b + 3d) + 3c)}{8(b + 3d)}$$

input `Int[Cosh[c + d*x]^3*Sinh[a + b*x],x]`

output $\text{Cosh}[a - 3*c + (b - 3*d)*x]/(8*(b - 3*d)) + (3*\text{Cosh}[a - c + (b - d)*x])/(8*(b - d)) + (3*\text{Cosh}[a + c + (b + d)*x])/(8*(b + d)) + \text{Cosh}[a + 3*c + (b + 3*d)*x]/(8*(b + 3*d))$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
]^(p)*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))`

Maple [A] (verified)

Time = 2.52 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.92

method	result
default	$\frac{\cosh(a-3c+(b-3d)x)}{8b-24d} + \frac{3 \cosh(a-c+(b-d)x)}{8(b-d)} + \frac{3 \cosh(a+c+(b+d)x)}{8(b+d)} + \frac{\cosh(a+3c+(b+3d)x)}{8b+24d}$
parallelrisc	$\frac{(b-d)(b+3d)(b+d) \cosh(a-3c+(b-3d)x)+3(b+3d)(b-3d)(b+d) \cosh(a-c+(b-d)x)+(b-d)(b-3d)(b+d) \cosh(a+3c+(b+3d)x)}{8b^4-80b^2d^2+72d^4}$
risc	$\frac{(b e^{2bx+2a}-3d e^{2bx+2a}+b+3d)e^{-bx+3dx-a+3c}}{16(b+3d)(b-3d)} + \frac{3(b e^{2bx+2a}-d e^{2bx+2a}+b+d)e^{-bx+dx-a+c}}{16(b+d)(b-d)} + \frac{3(b e^{2bx+2a}+d e^{2bx+2a})}{16(b+d)}$
orering	Expression too large to display

input `int(cosh(d*x+c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/8*cosh(a-3*c+(b-3*d)*x)/(b-3*d)+3/8*cosh(a-c+(b-d)*x)/(b-d)+3/8*cosh(a+c
+(b+d)*x)/(b+d)+1/8*cosh(a+3*c+(b+3*d)*x)/(b+3*d)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(83) = 166.

Time = 0.09 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.34

$$\int \cosh^3(c+dx) \sinh(a+bx) dx$$

$$= \frac{(b^3 - bd^2) \cosh(bx+a) \cosh(dx+c)^3 + 3(b^3 - bd^2) \cosh(bx+a) \cosh(dx+c) \sinh(dx+c)^2 - 3(b^2d}{4((b^4 - 10b^2d^2)}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a),x, algorithm="fricas")`

output `1/4*((b^3 - b*d^2)*cosh(b*x + a)*cosh(d*x + c)^3 + 3*(b^3 - b*d^2)*cosh(b*x + a)*cosh(d*x + c)*sinh(d*x + c)^2 - 3*(b^2*d - d^3)*sinh(b*x + a)*sinh(d*x + c)^3 + 3*(b^3 - 9*b*d^2)*cosh(b*x + a)*cosh(d*x + c) - 3*(b^2*d - 9*d^3 + 3*(b^2*d - d^3)*cosh(d*x + c)^2)*sinh(b*x + a)*sinh(d*x + c))/(b^4 - 10*b^2*d^2 + 9*d^4)*cosh(b*x + a)^2 - (b^4 - 10*b^2*d^2 + 9*d^4)*sinh(b*x + a)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 921 vs. $2(76) = 152$.

Time = 1.95 (sec) , antiderivative size = 921, normalized size of antiderivative = 10.12

$$\int \cosh^3(c + dx) \sinh(a + bx) dx = \text{Too large to display}$$

input `integrate(cosh(d*x+c)**3*sinh(b*x+a),x)`

output

```
Piecewise((x*sinh(a)*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (3*x*sinh(a - 3*d*x)
)*sinh(c + d*x)**2*cosh(c + d*x)/8 + x*sinh(a - 3*d*x)*cosh(c + d*x)**3/8
+ x*sinh(c + d*x)**3*cosh(a - 3*d*x)/8 + 3*x*sinh(c + d*x)*cosh(a - 3*d*x)
*cosh(c + d*x)**2/8 + sinh(a - 3*d*x)*sinh(c + d*x)**3/(8*d) + sinh(c + d*
x)**2*cosh(a - 3*d*x)*cosh(c + d*x)/(4*d) - 7*cosh(a - 3*d*x)*cosh(c + d*x
)**3/(24*d), Eq(b, -3*d)), (-3*x*sinh(a - d*x)*sinh(c + d*x)**2*cosh(c + d
*x)/8 + 3*x*sinh(a - d*x)*cosh(c + d*x)**3/8 - 3*x*sinh(c + d*x)**3*cosh(a
- d*x)/8 + 3*x*sinh(c + d*x)*cosh(a - d*x)*cosh(c + d*x)**2/8 + 3*sinh(a
- d*x)*sinh(c + d*x)**3/(8*d) + 3*sinh(c + d*x)**2*cosh(a - d*x)*cosh(c +
d*x)/(4*d) - 5*cosh(a - d*x)*cosh(c + d*x)**3/(8*d), Eq(b, -d)), (-3*x*si
nh(a + d*x)*sinh(c + d*x)**2*cosh(c + d*x)/8 + 3*x*sinh(a + d*x)*cosh(c + d
*x)**3/8 + 3*x*sinh(c + d*x)**3*cosh(a + d*x)/8 - 3*x*sinh(c + d*x)*cosh(a
+ d*x)*cosh(c + d*x)**2/8 + 3*sinh(a + d*x)*sinh(c + d*x)**3/(8*d) - 3*si
nh(c + d*x)**2*cosh(a + d*x)*cosh(c + d*x)/(4*d) + 5*cosh(a + d*x)*cosh(c
+ d*x)**3/(8*d), Eq(b, d)), (3*x*sinh(a + 3*d*x)*sinh(c + d*x)**2*cosh(c +
d*x)/8 + x*sinh(a + 3*d*x)*cosh(c + d*x)**3/8 - x*sinh(c + d*x)**3*cosh(a
+ 3*d*x)/8 - 3*x*sinh(c + d*x)*cosh(a + 3*d*x)*cosh(c + d*x)**2/8 + sinh(
a + 3*d*x)*sinh(c + d*x)**3/(8*d) - sinh(c + d*x)**2*cosh(a + 3*d*x)*cosh(
c + d*x)/(4*d) + 7*cosh(a + 3*d*x)*cosh(c + d*x)**3/(24*d), Eq(b, 3*d)), (
b**3*cosh(a + b*x)*cosh(c + d*x)**3/(b**4 - 10*b**2*d**2 + 9*d**4) - 3*...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^3(c + dx) \sinh(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)^3*sinh(b*x+a),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for
more detail
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(83) = 166$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.97

$$\int \cosh^3(c + dx) \sinh(a + bx) dx = \frac{e^{(bx+3dx+a+3c)}}{16(b+3d)} + \frac{3e^{(bx+dx+a+c)}}{16(b+d)} + \frac{3e^{(bx-dx+a-c)}}{16(b-d)} \\ + \frac{e^{(bx-3dx+a-3c)}}{16(b-3d)} + \frac{e^{(-bx+3dx-a+3c)}}{16(b-3d)} + \frac{3e^{(-bx+dx-a+c)}}{16(b-d)} \\ + \frac{3e^{(-bx-dx-a-c)}}{16(b+d)} + \frac{e^{(-bx-3dx-a-3c)}}{16(b+3d)}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a),x, algorithm="giac")`

output `1/16*e^(b*x + 3*d*x + a + 3*c)/(b + 3*d) + 3/16*e^(b*x + d*x + a + c)/(b + d) + 3/16*e^(b*x - d*x + a - c)/(b - d) + 1/16*e^(b*x - 3*d*x + a - 3*c)/(b - 3*d) + 1/16*e^(-b*x + 3*d*x - a + 3*c)/(b - 3*d) + 3/16*e^(-b*x + d*x - a + c)/(b - d) + 3/16*e^(-b*x - d*x - a - c)/(b + d) + 1/16*e^(-b*x - 3*d*x - a - 3*c)/(b + 3*d)`

Mupad [B] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.00

$$\int \cosh^3(c + dx) \sinh(a + bx) dx \\ = \frac{6bd^2 \cosh(a + bx) \cosh(c + dx) \sinh(c + dx)^2}{b^4 - 10b^2d^2 + 9d^4} - \frac{6d^3 \sinh(a + bx) \sinh(c + dx)^3}{b^4 - 10b^2d^2 + 9d^4} \\ - \frac{3d \cosh(c + dx)^2 \sinh(a + bx) \sinh(c + dx) (b^2 - 3d^2)}{b^4 - 10b^2d^2 + 9d^4} \\ - \frac{\cosh(a + bx) \cosh(c + dx)^3 (7bd^2 - b^3)}{b^4 - 10b^2d^2 + 9d^4}$$

input `int(cosh(c + d*x)^3*sinh(a + b*x),x)`

output

$$\begin{aligned} & (6*b*d^2*cosh(a + b*x)*cosh(c + d*x)*sinh(c + d*x)^2)/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (6*d^3*sinh(a + b*x)*sinh(c + d*x)^3)/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (3*d*cosh(c + d*x)^2*sinh(a + b*x)*sinh(c + d*x)*(b^2 - 3*d^2))/(b^4 + 9*d^4 - 10*b^2*d^2) \\ & - (cosh(a + b*x)*cosh(c + d*x)^3*(7*b*d^2 - b^3))/(b^4 + 9*d^4 - 10*b^2*d^2) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 573, normalized size of antiderivative = 6.30

$$\int \cosh^3(c + dx) \sinh(a + bx) dx$$

$$= \frac{b^3 + 3d^3 - b d^2 - e^{6dx+6c} b d^2 - e^{2bx+2a} b d^2 - e^{2bx+6dx+2a+6c} b d^2 + 27e^{2bx+4dx+2a+4c} d^3 + 3e^{2bx+2dx+2a+2c} b^3}{16e^{a+bx+3c+3d^2x}(b^4 - 10b^2d^2 + 9d^4)}$$

input

int(cosh(d*x+c)^3*sinh(b*x+a),x)

output

$$\begin{aligned} & (e^{2a+2bx+6c+6dx} b^3 - 3e^{2a+2bx+6c+6dx} b^2 d - e^{2a+2bx+6c+6dx} b d^2 + 3e^{2a+2bx+6c+6dx} d^3 + 3e^{2a+2bx+4c+4dx} b^3 - 3e^{2a+2bx+4c+4dx} b^2 d - 27e^{2a+2bx+4c+4dx} b d^2 + 27e^{2a+2bx+4c+4dx} d^3 + 3e^{2a+2bx+2c+2dx} b^3 + 3e^{2a+2bx+2c+2dx} b^2 d - 27e^{2a+2bx+2c+2dx} b d^2 - 27e^{2a+2bx+2c+2dx} d^3 + e^{2a+2bx} b^3 + 3e^{2a+2bx} b^2 d - e^{2a+2bx} b d^2 - 3e^{2a+2bx} d^3 + e^{6c+6dx} b^3 + 3e^{6c+6dx} b^2 d - e^{6c+6dx} b d^2 - 3e^{6c+6dx} d^3 + 3e^{4c+4dx} b^3 + 3e^{4c+4dx} b^2 d - 27e^{4c+4dx} b d^2 - 27e^{4c+4dx} d^3 + 3e^{2c+2dx} b^3 - 3e^{2c+2dx} b^2 d - 27e^{2c+2dx} b d^2 + 27e^{2c+2dx} d^3 + b^3 - 3b^2 d - b d^2 + 3d^3)/(16e^{a+bx+3c+3d^2x}(b^4 - 10b^2d^2 + 9d^4)) \end{aligned}$$

3.20 $\int \cosh(c + dx) \sinh^2(a + bx) dx$

Optimal result	202
Mathematica [A] (verified)	202
Rubi [A] (verified)	203
Maple [A] (verified)	204
Fricas [A] (verification not implemented)	204
Sympy [B] (verification not implemented)	205
Maxima [F(-2)]	206
Giac [A] (verification not implemented)	206
Mupad [B] (verification not implemented)	207
Reduce [B] (verification not implemented)	207

Optimal result

Integrand size = 15, antiderivative size = 68

$$\int \cosh(c + dx) \sinh^2(a + bx) dx = \frac{\sinh(2a - c + (2b - d)x)}{4(2b - d)} - \frac{\sinh(c + dx)}{2d} + \frac{\sinh(2a + c + (2b + d)x)}{4(2b + d)}$$

output

```
sinh(2*a-c+(2*b-d)*x)/(8*b-4*d)-1/2*sinh(d*x+c)/d+sinh(2*a+c+(2*b+d)*x)/(8*b+4*d)
```

Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.09

$$\int \cosh(c + dx) \sinh^2(a + bx) dx = \frac{1}{4} \left(-\frac{2 \cosh(dx) \sinh(c)}{d} - \frac{2 \cosh(c) \sinh(dx)}{d} + \frac{\sinh(2a - c + 2bx - dx)}{2b - d} + \frac{\sinh(2a + c + 2bx + dx)}{2b + d} \right)$$

input

```
Integrate[Cosh[c + d*x]*Sinh[a + b*x]^2,x]
```

output

$$\left(\frac{-2 \operatorname{Cosh}[d x] \operatorname{Sinh}[c]}{d} - \frac{2 \operatorname{Cosh}[c] \operatorname{Sinh}[d x]}{d} + \frac{\operatorname{Sinh}[2 a - c + 2 b x - d x]}{2 b - d} + \frac{\operatorname{Sinh}[2 a + c + 2 b x + d x]}{2 b + d} \right) / 4$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \cosh(c + dx) dx$$

$$\downarrow \text{6152}$$

$$\int \left(\frac{1}{4} \cosh(2a + x(2b - d) - c) + \frac{1}{4} \cosh(2a + x(2b + d) + c) - \frac{1}{2} \cosh(c + dx) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sinh(2a + x(2b - d) - c)}{4(2b - d)} + \frac{\sinh(2a + x(2b + d) + c)}{4(2b + d)} - \frac{\sinh(c + dx)}{2d}$$

input

$$\text{Int}[\operatorname{Cosh}[c + d x] \operatorname{Sinh}[a + b x]^2, x]$$

output

$$\frac{\operatorname{Sinh}[2 a - c + (2 b - d) x]}{4(2 b - d)} - \frac{\operatorname{Sinh}[c + d x]}{2 d} + \frac{\operatorname{Sinh}[2 a + c + (2 b + d) x]}{4(2 b + d)}$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp[IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 6152

```
Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] :> Int[ExpandTrigReduce[Sinh[v]
] ^p*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.93

method	result
default	$\frac{\sinh(2a-c+(2b-d)x)}{8b-4d} - \frac{\sinh(dx+c)}{2d} + \frac{\sinh(2a+c+(2b+d)x)}{8b+4d}$
parallelrisch	$\frac{(2bd+d^2) \sinh(2a-c+(2b-d)x) + (2bd-d^2) \sinh(2a+c+(2b+d)x) + (-8b^2+2d^2) \sinh(dx+c)}{16b^2d-4d^3}$
risch	$-\frac{(-2de^{4bx+4a}b+d^2e^{4bx+4a}+8b^2e^{2bx+2a}-2d^2e^{2bx+2a}+2bd+d^2)e^{-2bx+dx-2a+c}}{8(2b+d)(2b-d)d} + \frac{(2de^{4bx+4a}b+d^2e^{4bx+4a}+8b^2e^{2bx+2a}-2d^2e^{2bx+2a}+2bd+d^2)e^{-2bx+dx-2a+c}}{8(2b+d)(2b-d)d}$
orering	$\frac{(16b^4+3d^4) \left(d \sinh(dx+c) \sinh(bx+a)^2 + 2 \cosh(dx+c) \sinh(bx+a) b \cosh(bx+a) \right)}{d^2(16b^4-8b^2d^2+d^4)} - \frac{(8b^2+3d^2) \left(d^3 \sinh(dx+c) \sinh(bx+a) \right)}{d^2(16b^4-8b^2d^2+d^4)}$

input

```
int(cosh(d*x+c)*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
-1/2*sinh(d*x+c)/d+1/4*sinh(2*a-c+(2*b-d)*x)/(2*b-d)+1/4*sinh(2*a+c+(2*b+d)
)*x)/(2*b+d)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.68

$$\int \cosh(c+dx) \sinh^2(a+bx) dx$$

$$= \frac{4bd \cosh(bx+a) \cosh(dx+c) \sinh(bx+a) - (d^2 \cosh(bx+a)^2 + d^2 \sinh(bx+a)^2 + 4b^2 - d^2) \sinh(bx+a)}{2((4b^2d - d^3) \cosh(bx+a)^2 - (4b^2d - d^3) \sinh(bx+a)^2)}$$

input

```
integrate(cosh(d*x+c)*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

$$\frac{1}{2}(4*b*d*cosh(b*x + a)*cosh(d*x + c)*sinh(b*x + a) - (d^2*cosh(b*x + a)^2 + d^2*sinh(b*x + a)^2 + 4*b^2 - d^2)*sinh(d*x + c))/((4*b^2*d - d^3)*cosh(b*x + a)^2 - (4*b^2*d - d^3)*sinh(b*x + a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(49) = 98$.

Time = 0.73 (sec) , antiderivative size = 408, normalized size of antiderivative = 6.00

$$\int \cosh(c + dx) \sinh^2(a + bx) dx$$

$$= \begin{cases} x \sinh^2(a) \cosh(c) \\ \frac{x \sinh^2\left(a - \frac{dx}{2}\right) \cosh(c+dx)}{4} + \frac{x \sinh\left(a - \frac{dx}{2}\right) \sinh(c+dx) \cosh\left(a - \frac{dx}{2}\right)}{2} + \frac{x \cosh^2\left(a - \frac{dx}{2}\right) \cosh(c+dx)}{4} + \frac{\sinh^2\left(a - \frac{dx}{2}\right) \sinh(c+dx)}{d} \\ \frac{x \sinh^2\left(a + \frac{dx}{2}\right) \cosh(c+dx)}{4} - \frac{x \sinh\left(a + \frac{dx}{2}\right) \sinh(c+dx) \cosh\left(a + \frac{dx}{2}\right)}{2} + \frac{x \cosh^2\left(a + \frac{dx}{2}\right) \cosh(c+dx)}{4} + \frac{\sinh^2\left(a + \frac{dx}{2}\right) \sinh(c+dx)}{d} \\ \left(\frac{x \sinh^2(a+bx)}{2} - \frac{x \cosh^2(a+bx)}{2} + \frac{\sinh(a+bx) \cosh(a+bx)}{2b} \right) \cosh(c) \\ \frac{2b^2 \sinh^2(a+bx) \sinh(c+dx)}{4b^2d-d^3} - \frac{2b^2 \sinh(c+dx) \cosh^2(a+bx)}{4b^2d-d^3} + \frac{2bd \sinh(a+bx) \cosh(a+bx) \cosh(c+dx)}{4b^2d-d^3} - \frac{d^2 \sinh^2(a+bx) \sinh(c+dx)}{4b^2d-d^3} \end{cases}$$

input

```
integrate(cosh(d*x+c)*sinh(b*x+a)**2,x)
```

output

```
Piecewise((x*sinh(a)**2*cosh(c), Eq(b, 0) & Eq(d, 0)), (x*sinh(a - d*x/2)*
**2*cosh(c + d*x)/4 + x*sinh(a - d*x/2)*sinh(c + d*x)*cosh(a - d*x/2)/2 + x
*cosh(a - d*x/2)**2*cosh(c + d*x)/4 + sinh(a - d*x/2)**2*sinh(c + d*x)/d +
sinh(a - d*x/2)*cosh(a - d*x/2)*cosh(c + d*x)/(2*d), Eq(b, -d/2)), (x*sin
h(a + d*x/2)**2*cosh(c + d*x)/4 - x*sinh(a + d*x/2)*sinh(c + d*x)*cosh(a +
d*x/2)/2 + x*cosh(a + d*x/2)**2*cosh(c + d*x)/4 + sinh(a + d*x/2)**2*sinh
(c + d*x)/d - sinh(a + d*x/2)*cosh(a + d*x/2)*cosh(c + d*x)/(2*d), Eq(b, d
/2)), ((x*sinh(a + b*x)**2/2 - x*cosh(a + b*x)**2/2 + sinh(a + b*x)*cosh(a
+ b*x)/(2*b))*cosh(c), Eq(d, 0)), (2*b**2*sinh(a + b*x)**2*sinh(c + d*x)/
(4*b**2*d - d**3) - 2*b**2*sinh(c + d*x)*cosh(a + b*x)**2/(4*b**2*d - d**3
) + 2*b*d*sinh(a + b*x)*cosh(a + b*x)*cosh(c + d*x)/(4*b**2*d - d**3) - d
**2*sinh(a + b*x)**2*sinh(c + d*x)/(4*b**2*d - d**3), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh(c + dx) \sinh^2(a + bx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a)^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-d/b>0)', see `assume?` for more details)

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.82

$$\int \cosh(c + dx) \sinh^2(a + bx) dx = \frac{e^{(2bx+dx+2a+c)}}{8(2b+d)} + \frac{e^{(2bx-dx+2a-c)}}{8(2b-d)} - \frac{e^{(-2bx+dx-2a+c)}}{8(2b-d)} - \frac{e^{(-2bx-dx-2a-c)}}{8(2b+d)} - \frac{e^{(dx+c)}}{4d} + \frac{e^{(-dx-c)}}{4d}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a)^2,x, algorithm="giac")`

output `1/8*e^(2*b*x + d*x + 2*a + c)/(2*b + d) + 1/8*e^(2*b*x - d*x + 2*a - c)/(2*b - d) - 1/8*e^(-2*b*x + d*x - 2*a + c)/(2*b - d) - 1/8*e^(-2*b*x - d*x - 2*a - c)/(2*b + d) - 1/4*e^(d*x + c)/d + 1/4*e^(-d*x - c)/d`

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.12

$$\int \cosh(c + dx) \sinh^2(a + bx) dx$$

$$= \frac{d^2 (\sinh(c + dx) - \cosh(a + bx)^2 \sinh(c + dx)) - 2b^2 \sinh(c + dx) + 2bd \cosh(a + bx) \cosh(c + dx)}{4b^2 d - d^3}$$

input `int(cosh(c + d*x)*sinh(a + b*x)^2,x)`output `(d^2*(sinh(c + d*x) - cosh(a + b*x)^2*sinh(c + d*x)) - 2*b^2*sinh(c + d*x) + 2*b*d*cosh(a + b*x)*cosh(c + d*x))/(4*b^2*d - d^3)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 216, normalized size of antiderivative = 3.18

$$\int \cosh(c + dx) \sinh^2(a + bx) dx$$

$$= \frac{2e^{4bx+2dx+4a+2c}bd - e^{4bx+2dx+4a+2c}d^2 + 2e^{4bx+4a}bd + e^{4bx+4a}d^2 - 8e^{2bx+2dx+2a+2c}b^2 + 2e^{2bx+2dx+2a+2c}d^2 + 8e^{2bx+dx+2a+c}d(4b^2 - d^2)}{8e^{2bx+dx+2a+c}d(4b^2 - d^2)}$$

input `int(cosh(d*x+c)*sinh(b*x+a)^2,x)`output `(2*e**(4*a + 4*b*x + 2*c + 2*d*x)*b*d - e**(4*a + 4*b*x + 2*c + 2*d*x)*d**2 + 2*e**(4*a + 4*b*x)*b*d + e**(4*a + 4*b*x)*d**2 - 8*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2 + 2*e**(2*a + 2*b*x + 2*c + 2*d*x)*d**2 + 8*e**(2*a + 2*b*x)*b**2 - 2*e**(2*a + 2*b*x)*d**2 - 2*e**(2*c + 2*d*x)*b*d - e**(2*c + 2*d*x)*d**2 - 2*b*d + d**2)/(8*e**(2*a + 2*b*x + c + d*x)*d*(4*b**2 - d**2))`

3.21 $\int \cosh^2(c + dx) \sinh^2(a + bx) dx$

Optimal result	208
Mathematica [A] (verified)	208
Rubi [A] (verified)	209
Maple [A] (verified)	210
Fricas [B] (verification not implemented)	210
Sympy [B] (verification not implemented)	211
Maxima [F(-2)]	212
Giac [A] (verification not implemented)	213
Mupad [B] (verification not implemented)	213
Reduce [B] (verification not implemented)	214

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = -\frac{x}{4} + \frac{\sinh(2a + 2bx)}{8b} + \frac{\sinh(2(a - c) + 2(b - d)x)}{16(b - d)} - \frac{\sinh(2c + 2dx)}{8d} + \frac{\sinh(2(a + c) + 2(b + d)x)}{16(b + d)}$$

output

```
-1/4*x+1/8*sinh(2*b*x+2*a)/b+sinh(2*a-2*c+2*(b-d)*x)/(16*b-16*d)-1/8*sinh(2*d*x+2*c)/d+sinh(2*a+2*c+2*(b+d)*x)/(16*b+16*d)
```

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.22

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = \frac{2d(b^2 - d^2) \sinh(2(a + bx)) + bd(b + d) \sinh(2(a - c + (b - d)x)) - b(b - d)(4d(b + d)x + 2(b + d) \sinh(2(a + c) + 2(b + d)x))}{16b(b - d)d(b + d)}$$

input

```
Integrate[Cosh[c + d*x]^2*Sinh[a + b*x]^2,x]
```

output

$$\frac{(2*d*(b^2 - d^2)*\text{Sinh}[2*(a + b*x)] + b*d*(b + d)*\text{Sinh}[2*(a - c + (b - d)*x)] - b*(b - d)*(4*d*(b + d)*x + 2*(b + d)*\text{Sinh}[2*(c + d*x)] - d*\text{Sinh}[2*(a + c + (b + d)*x)])}{16*b*(b - d)*d*(b + d)}$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \cosh^2(c + dx) dx$$

↓ 6152

$$\int \left(\frac{1}{8} \cosh(2(a - c) + 2x(b - d)) + \frac{1}{8} \cosh(2(a + c) + 2x(b + d)) + \frac{1}{4} \cosh(2a + 2bx) - \frac{1}{4} \cosh(2c + 2dx) - \frac{1}{4} \right)$$

↓ 2009

$$\frac{\sinh(2(a - c) + 2x(b - d))}{16(b - d)} + \frac{\sinh(2(a + c) + 2x(b + d))}{16(b + d)} + \frac{\sinh(2a + 2bx)}{8b} - \frac{\sinh(2c + 2dx)}{8d} - \frac{x}{4}$$

input

$$\text{Int}[\text{Cosh}[c + d*x]^2*\text{Sinh}[a + b*x]^2,x]$$

output

$$-1/4*x + \text{Sinh}[2*a + 2*b*x]/(8*b) + \text{Sinh}[2*(a - c) + 2*(b - d)*x]/(16*(b - d)) - \text{Sinh}[2*c + 2*d*x]/(8*d) + \text{Sinh}[2*(a + c) + 2*(b + d)*x]/(16*(b + d))$$

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6152 Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]
] ^p*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 3.26 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

method	result
default	$-\frac{x}{4} + \frac{\sinh(2bx+2a)}{8b} - \frac{\sinh(2dx+2c)}{8d} + \frac{\sinh((2b-2d)x+2a-2c)}{16b-16d} + \frac{\sinh((2b+2d)x+2a+2c)}{16b+16d}$
parallelrisch	$\frac{bd(b+d) \sinh((2b-2d)x+2a-2c) - 4(b-d) \left(-\frac{bd \sinh((2b+2d)x+2a+2c)}{4} + \left(-\frac{d \sinh(2bx+2a)}{2} + b \left(dx + \frac{\sinh(2dx+2c)}{2} \right) \right) (b+d) \right)}{16b^3d - 16bd^3}$
risch	$-\frac{x}{4} + \frac{e^{2bx+2a}}{16b} - \frac{e^{-2bx-2a}}{16b} - \frac{(-de^{4bx+4a}b + d^2e^{4bx+4a} + 2b^2e^{2bx+2a} - 2d^2e^{2bx+2a} + bd + d^2)e^{-2bx+2dx-2a+2c}}{32(b+d)(b-d)d} + \frac{(de^{2bx+2a} - d^2e^{-2bx-2a})e^{2dx+2c}}{4(b-d)d}$
orering	Expression too large to display

```
input int(cosh(d*x+c)^2*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
output -1/4*x+1/8*sinh(2*b*x+2*a)/b-1/8*sinh(2*d*x+2*c)/d+1/8/(2*b-2*d)*sinh((2*b
-2*d)*x+2*a-2*c)+1/8/(2*b+2*d)*sinh((2*b+2*d)*x+2*a+2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 192 vs. 2(78) = 156.

Time = 0.09 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.18

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx$$

$$= \frac{b^2d \cosh(bx + a) \sinh(bx + a) \sinh(dx + c)^2 - (b^3d - bd^3)x + (b^2d \cosh(bx + a) \cosh(dx + c)^2 + (b^2d - d^3) \cosh(bx + a) \sinh(dx + c) \cosh(dx + c) - d^3 \sinh(bx + a) \sinh(dx + c) \cosh(dx + c) - d^3 \sinh(bx + a) \sinh(dx + c)^2)}{4((b^3d - bd^3) \cosh(dx + c) + (b^2d - d^3) \sinh(dx + c) \cosh(dx + c) - d^3 \sinh(dx + c)^2)}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{1}{4}(b^2d\cosh(bx+a)\sinh(bx+a)\sinh(dx+c)^2 - (b^3d - b^2d^3)x + (b^2d\cosh(bx+a)\cosh(dx+c)^2 + (b^2d - d^3)\cosh(bx+a))\sinh(bx+a) - (bd^2\cosh(dx+c)\sinh(bx+a)^2 + (bd^2\cosh(bx+a)^2 + b^3 - bd^2)\cosh(dx+c))\sinh(dx+c)) / ((b^3d - b^2d^3)\cosh(bx+a)^2 - (b^3d - b^2d^3)\sinh(bx+a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1027 vs. $2(76) = 152$.

Time = 1.59 (sec) , antiderivative size = 1027, normalized size of antiderivative = 11.67

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(cosh(d*x+c)**2*sinh(b*x+a)**2,x)`

output

```
Piecewise((x*sinh(a)**2*cosh(c)**2, Eq(b, 0) & Eq(d, 0)), ((-x*sinh(c + d*x)**2/2 + x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*sinh(a)**2, Eq(b, 0)), (-x*sinh(a - d*x)**2*sinh(c + d*x)**2/8 + 3*x*sinh(a - d*x)**2*cosh(c + d*x)**2/8 + x*sinh(a - d*x)*sinh(c + d*x)*cosh(a - d*x)*cosh(c + d*x)/2 + 3*x*sinh(c + d*x)**2*cosh(a - d*x)**2/8 - x*cosh(a - d*x)**2*cosh(c + d*x)**2/8 - sinh(a - d*x)*sinh(c + d*x)**2*cosh(a - d*x)/(8*d) - 5*sinh(a - d*x)*cosh(a - d*x)*cosh(c + d*x)**2/(8*d) - sinh(c + d*x)*cosh(a - d*x)**2*cosh(c + d*x)/(2*d), Eq(b, -d)), (-x*sinh(a + d*x)**2*sinh(c + d*x)**2/8 + 3*x*sinh(a + d*x)**2*cosh(c + d*x)**2/8 - x*sinh(a + d*x)*sinh(c + d*x)*cosh(a + d*x)*cosh(c + d*x)/2 + 3*x*sinh(c + d*x)**2*cosh(a + d*x)**2/8 - x*cosh(a + d*x)**2*cosh(c + d*x)**2/8 + sinh(a + d*x)**2*sinh(c + d*x)*cosh(c + d*x)/(2*d) - 3*sinh(a + d*x)*sinh(c + d*x)**2*cosh(a + d*x)/(8*d) + sinh(a + d*x)*cosh(a + d*x)*cosh(c + d*x)**2/(8*d), Eq(b, d)), ((x*sinh(a + b*x)**2/2 - x*cosh(a + b*x)**2/2 + sinh(a + b*x)*cosh(a + b*x)/(2*b))*cosh(c)**2, Eq(d, 0)), (-b**3*d*x*sinh(a + b*x)**2*sinh(c + d*x)**2/(4*b**3*d - 4*b*d**3) + b**3*d*x*sinh(a + b*x)**2*cosh(c + d*x)**2/(4*b**3*d - 4*b*d**3) + b**3*d*x*sinh(c + d*x)**2*cosh(a + b*x)**2/(4*b**3*d - 4*b*d**3) - b**3*d*x*cosh(a + b*x)**2*cosh(c + d*x)**2/(4*b**3*d - 4*b*d**3) + b**3*sinh(a + b*x)**2*sinh(c + d*x)*cosh(c + d*x)/(4*b**3*d - 4*b*d**3) - b**3*sinh(c + d*x)*cosh(a + b*x)**2*cosh(c + d*x)/(4*b**3*d - 4...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)^2*sinh(b*x+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-(2*d)/b>0)', see `assume?` for more deta
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.77

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = -\frac{1}{4}x + \frac{e^{(2bx+2dx+2a+2c)}}{32(b+d)} + \frac{e^{(2bx-2dx+2a-2c)}}{32(b-d)} + \frac{e^{(2bx+2a)}}{16b} - \frac{e^{(-2bx+2dx-2a+2c)}}{32(b-d)} - \frac{e^{(-2bx-2dx-2a-2c)}}{32(b+d)} - \frac{e^{(-2bx-2a)}}{16b} - \frac{e^{(2dx+2c)}}{16d} + \frac{e^{(-2dx-2c)}}{16d}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a)^2,x, algorithm="giac")`output `-1/4*x + 1/32*e^(2*b*x + 2*d*x + 2*a + 2*c)/(b + d) + 1/32*e^(2*b*x - 2*d*x + 2*a - 2*c)/(b - d) + 1/16*e^(2*b*x + 2*a)/b - 1/32*e^(-2*b*x + 2*d*x - 2*a + 2*c)/(b - d) - 1/32*e^(-2*b*x - 2*d*x - 2*a - 2*c)/(b + d) - 1/16*e^(-2*b*x - 2*a)/b - 1/16*e^(2*d*x + 2*c)/d + 1/16*e^(-2*d*x - 2*c)/d`**Mupad [B] (verification not implemented)**

Time = 1.46 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.53

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx = \frac{d^3 \cosh(a + bx) \sinh(a + bx) + b^3 \cosh(c + dx) \sinh(c + dx) - b d^3 x + b^3 dx - 2 b d^2 \cosh(c + dx) \sinh(c + dx)}{4}$$

input `int(cosh(c + d*x)^2*sinh(a + b*x)^2,x)`output `-(d^3*cosh(a + b*x)*sinh(a + b*x) + b^3*cosh(c + d*x)*sinh(c + d*x) - b*d^3*x + b^3*d*x - 2*b*d^2*cosh(c + d*x)*sinh(c + d*x) - 2*b^2*d*cosh(a + b*x)*cosh(c + d*x)^2*sinh(a + b*x) + 2*b*d^2*cosh(a + b*x)^2*cosh(c + d*x)*sinh(c + d*x))/(4*b*d*(b^2 - d^2))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.06

$$\int \cosh^2(c + dx) \sinh^2(a + bx) dx$$

$$= \frac{e^{4bx+4dx+4a+4c} b^2 d - e^{4bx+4dx+4a+4c} b d^2 + 2e^{4bx+2dx+4a+2c} b^2 d - 2e^{4bx+2dx+4a+2c} d^3 + e^{4bx+4a} b^2 d + e^{4bx+4a} b d^2}{32e^{2a+2bx+2c+2dx} (b^2 d^2 - d^2)}$$

input

```
int(cosh(d*x+c)^2*sinh(b*x+a)^2,x)
```

output

```
(e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d - e**(4*a + 4*b*x + 4*c + 4*d*x)*b*
d**2 + 2*e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d - 2*e**(4*a + 4*b*x + 2*c +
2*d*x)*d**3 + e**(4*a + 4*b*x)*b**2*d + e**(4*a + 4*b*x)*b*d**2 - 2*e**(2
*a + 2*b*x + 4*c + 4*d*x)*b**3 + 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d**2 -
8*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**3*d*x + 8*e**(2*a + 2*b*x + 2*c + 2*d
*x)*b*d**3*x + 2*e**(2*a + 2*b*x)*b**3 - 2*e**(2*a + 2*b*x)*b*d**2 - e**(4
*c + 4*d*x)*b**2*d - e**(4*c + 4*d*x)*b*d**2 - 2*e**(2*c + 2*d*x)*b**2*d +
2*e**(2*c + 2*d*x)*d**3 - b**2*d + b*d**2)/(32*e**(2*a + 2*b*x + 2*c + 2*
d*x)*b*d*(b**2 - d**2))
```

3.22 $\int \cosh^3(c + dx) \sinh^2(a + bx) dx$

Optimal result	215
Mathematica [A] (verified)	216
Rubi [A] (verified)	216
Maple [A] (verified)	218
Fricas [B] (verification not implemented)	218
Sympy [B] (verification not implemented)	219
Maxima [F(-2)]	220
Giac [A] (verification not implemented)	221
Mupad [B] (verification not implemented)	222
Reduce [B] (verification not implemented)	223

Optimal result

Integrand size = 17, antiderivative size = 144

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx = \frac{\sinh(2a - 3c + (2b - 3d)x)}{16(2b - 3d)} + \frac{3 \sinh(2a - c + (2b - d)x)}{16(2b - d)} - \frac{3 \sinh(c + dx)}{8d} - \frac{\sinh(3c + 3dx)}{24d} + \frac{3 \sinh(2a + c + (2b + d)x)}{16(2b + d)} + \frac{\sinh(2a + 3c + (2b + 3d)x)}{16(2b + 3d)}$$

output

```
sinh(2*a-3*c+(2*b-3*d)*x)/(32*b-48*d)+3*sinh(2*a-c+(2*b-d)*x)/(32*b-16*d)-
3/8*sinh(d*x+c)/d-1/24*sinh(3*d*x+3*c)/d+3*sinh(2*a+c+(2*b+d)*x)/(32*b+16*
d)+sinh(2*a+3*c+(2*b+3*d)*x)/(32*b+48*d)
```


Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.10

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx = \frac{1}{48} \left(-\frac{18 \cosh(dx) \sinh(c)}{d} - \frac{2 \cosh(3dx) \sinh(3c)}{d} - \frac{18 \cosh(c) \sinh(dx)}{d} - \frac{2 \cosh(3c) \sinh(3dx)}{d} + \frac{3 \sinh(2a - 3c + 2bx - 3dx)}{2b - 3d} + \frac{9 \sinh(2a - c + 2bx - dx)}{2b - d} + \frac{9 \sinh(2a + c + 2bx + dx)}{2b + d} + \frac{3 \sinh(2a + 3c + 2bx + 3dx)}{2b + 3d} \right)$$

input

```
Integrate[Cosh[c + d*x]^3*Sinh[a + b*x]^2,x]
```

output

```
((-18*Cosh[d*x]*Sinh[c])/d - (2*Cosh[3*d*x]*Sinh[3*c])/d - (18*Cosh[c]*Sinh[d*x])/d - (2*Cosh[3*c]*Sinh[3*d*x])/d + (3*Sinh[2*a - 3*c + 2*b*x - 3*d*x])/(2*b - 3*d) + (9*Sinh[2*a - c + 2*b*x - d*x])/(2*b - d) + (9*Sinh[2*a + c + 2*b*x + d*x])/(2*b + d) + (3*Sinh[2*a + 3*c + 2*b*x + 3*d*x])/(2*b + 3*d))/48
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \cosh^3(c + dx) dx$$

↓ 6152

$$\int \left(\frac{1}{16} \cosh(2a + x(2b - 3d) - 3c) + \frac{3}{16} \cosh(2a + x(2b - d) - c) + \frac{3}{16} \cosh(2a + x(2b + d) + c) + \frac{1}{16} \cosh(2a + x(2b + 3d) + 3c) - \frac{3 \sinh(c + dx)}{8d} - \frac{\sinh(3c + 3dx)}{24d} \right) dx$$

↓ 2009

$$\frac{\sinh(2a + x(2b - 3d) - 3c)}{16(2b - 3d)} + \frac{3 \sinh(2a + x(2b - d) - c)}{16(2b - d)} + \frac{3 \sinh(2a + x(2b + d) + c)}{16(2b + d)} + \frac{\sinh(2a + x(2b + 3d) + 3c)}{16(2b + 3d)} - \frac{3 \sinh(c + dx)}{8d} - \frac{\sinh(3c + 3dx)}{24d}$$

input `Int[Cosh[c + d*x]^3*Sinh[a + b*x]^2,x]`

output `Sinh[2*a - 3*c + (2*b - 3*d)*x]/(16*(2*b - 3*d)) + (3*Sinh[2*a - c + (2*b - d)*x])/(16*(2*b - d)) - (3*Sinh[c + d*x])/(8*d) - Sinh[3*c + 3*d*x]/(24*d) + (3*Sinh[2*a + c + (2*b + d)*x])/(16*(2*b + d)) + Sinh[2*a + 3*c + (2*b + 3*d)*x]/(16*(2*b + 3*d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 8.39 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.92

method	result
default	$-\frac{3 \sinh(dx+c)}{8d} - \frac{\sinh(3dx+3c)}{24d} + \frac{\sinh(2a-3c+(2b-3d)x)}{32b-48d} + \frac{3 \sinh(2a-c+(2b-d)x)}{16(2b-d)} + \frac{3 \sinh(2a+c+(2b+d)x)}{16(2b+d)} + \dots$
parallelrisc	$\frac{(24b^3d+36b^2d^2-6bd^3-9d^4) \sinh(2a-3c+(2b-3d)x)+72 \left(\left(b+\frac{3d}{2}\right) d \left(b+\frac{d}{2}\right) \sinh(2a-c+(2b-d)x) + \left(\frac{d \left(b+\frac{d}{2}\right) \sinh(2a+3c+(2b+d)x)}{3}\right)}{768b^4d-1920b^2d^3+432d^4}$
risc	$-\frac{(-6de^{4bx+4a}b+9d^2e^{4bx+4a}+8b^2e^{2bx+2a}-18d^2e^{2bx+2a}+6bd+9d^2)e^{-2bx+3dx-2a+3c}}{96(2b+3d)(2b-3d)d} - \frac{3(-2de^{4bx+4a}b+d^2e^{4bx+4a}+9d^2e^{2bx+2a}-18d^2e^{2bx+2a}+6bd+9d^2)e^{-2bx+3dx-2a+3c}}{96(2b+3d)(2b-3d)d}$
orering	Expression too large to display

input `int(cosh(d*x+c)^3*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$-\frac{3}{8} \frac{\sinh(dx+c)}{d} - \frac{1}{24} \frac{\sinh(3dx+3c)}{d} + \frac{1}{16} \frac{\sinh(2a-3c+(2b-3d)x)}{(2b-3d)} + \frac{3}{16} \frac{\sinh(2a-c+(2b-d)x)}{(2b-d)} + \frac{3}{16} \frac{\sinh(2a+c+(2b+d)x)}{(2b+d)} + \frac{1}{16} \frac{\sinh(2a+3c+(2b+3d)x)}{(2b+3d)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 398 vs. 2(132) = 264.

Time = 0.10 (sec) , antiderivative size = 398, normalized size of antiderivative = 2.76

$$\int \cosh^3(c+dx) \sinh^2(a+bx) dx$$

$$= \frac{36(4b^3d - bd^3) \cosh(bx+a) \cosh(dx+c) \sinh(bx+a) \sinh(dx+c)^2 - (16b^4 - 40b^2d^2 + 9d^4 + 9(4b^3d - bd^3)) \cosh^2(bx+a) \cosh^2(dx+c) \sinh(bx+a) \sinh(dx+c)}{72b^2d^2 - 36bd^3 + 9d^4}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/24*(36*(4*b^3*d - b*d^3)*cosh(b*x + a)*cosh(d*x + c)*sinh(b*x + a)*sinh(
d*x + c)^2 - (16*b^4 - 40*b^2*d^2 + 9*d^4 + 9*(4*b^2*d^2 - d^4)*cosh(b*x +
a)^2 + 9*(4*b^2*d^2 - d^4)*sinh(b*x + a)^2)*sinh(d*x + c)^3 + 12*((4*b^3*
d - b*d^3)*cosh(b*x + a)*cosh(d*x + c)^3 + 3*(4*b^3*d - 9*b*d^3)*cosh(b*x
+ a)*cosh(d*x + c)*sinh(b*x + a) - 3*(48*b^4 - 120*b^2*d^2 + 27*d^4 + 3*(
4*b^2*d^2 - 9*d^4)*cosh(b*x + a)^2 + (16*b^4 - 40*b^2*d^2 + 9*d^4 + 9*(4*b
^2*d^2 - d^4)*cosh(b*x + a)^2)*cosh(d*x + c)^2 + 3*(4*b^2*d^2 - 9*d^4 + 3*
(4*b^2*d^2 - d^4)*cosh(d*x + c)^2)*sinh(b*x + a)^2)*sinh(d*x + c))/((16*b^
4*d - 40*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2 - (16*b^4*d - 40*b^2*d^3 + 9*d^5
)*sinh(b*x + a)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1999 vs. 2(116) = 232.

Time = 5.39 (sec) , antiderivative size = 1999, normalized size of antiderivative = 13.88

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(d*x+c)**3*sinh(b*x+a)**2,x)
```

output

```
Piecewise((x*sinh(a)**2*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (3*x*sinh(a - 3*d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + x*sinh(a - 3*d*x/2)**2*cosh(c + d*x)**3/16 + x*sinh(a - 3*d*x/2)*sinh(c + d*x)**3*cosh(a - 3*d*x/2)/8 + 3*x*sinh(a - 3*d*x/2)*sinh(c + d*x)*cosh(a - 3*d*x/2)*cosh(c + d*x)**2/8 + 3*x*sinh(c + d*x)**2*cosh(a - 3*d*x/2)**2*cosh(c + d*x)/16 + x*cosh(a - 3*d*x/2)**2*cosh(c + d*x)**3/16 - 5*sinh(a - 3*d*x/2)**2*sinh(c + d*x)**3/(48*d) + sinh(a - 3*d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/d + 5*sinh(a - 3*d*x/2)*sinh(c + d*x)**2*cosh(a - 3*d*x/2)*cosh(c + d*x)/(4*d) + sinh(a - 3*d*x/2)*cosh(a - 3*d*x/2)*cosh(c + d*x)**3/(24*d) + 9*sinh(c + d*x)**3*cosh(a - 3*d*x/2)**2/(16*d), Eq(b, -3*d/2)), (-3*x*sinh(a - d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + 3*x*sinh(a - d*x/2)**2*cosh(c + d*x)**3/16 - 3*x*sinh(a - d*x/2)*sinh(c + d*x)**3*cosh(a - d*x/2)/8 + 3*x*sinh(a - d*x/2)*sinh(c + d*x)*cosh(a - d*x/2)*cosh(c + d*x)**2/8 - 3*x*sinh(c + d*x)**2*cosh(a - d*x/2)**2*cosh(c + d*x)/16 + 3*x*cosh(a - d*x/2)**2*cosh(c + d*x)**3/16 - 31*sinh(a - d*x/2)**2*sinh(c + d*x)**3/(48*d) + sinh(a - d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/d - sinh(a - d*x/2)*sinh(c + d*x)**2*cosh(a - d*x/2)*cosh(c + d*x)/(4*d) + 3*sinh(a - d*x/2)*cosh(a - d*x/2)*cosh(c + d*x)**3/(8*d) + sinh(c + d*x)**3*cosh(a - d*x/2)**2/(48*d), Eq(b, -d/2)), (-3*x*sinh(a + d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + 3*x*sinh(a + d*x/2)**2*cosh(c + d*x)**3/16 + 3*x*sinh(a + d*x/2)*sinh(c + d...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)^3*sinh(b*x+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-(3*d)/b>0)', see `assume?` for more deta
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.81

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx = \frac{e^{(2bx+3dx+2a+3c)}}{32(2b+3d)} + \frac{3e^{(2bx+dx+2a+c)}}{32(2b+d)}$$

$$+ \frac{3e^{(2bx-dx+2a-c)}}{32(2b-d)} + \frac{e^{(2bx-3dx+2a-3c)}}{32(2b-3d)}$$

$$- \frac{e^{(-2bx+3dx-2a+3c)}}{32(2b-3d)} - \frac{3e^{(-2bx+dx-2a+c)}}{32(2b-d)}$$

$$- \frac{3e^{(-2bx-dx-2a-c)}}{32(2b+d)} - \frac{e^{(-2bx-3dx-2a-3c)}}{32(2b+3d)}$$

$$- \frac{e^{(3dx+3c)}}{48d} - \frac{3e^{(dx+c)}}{16d} + \frac{3e^{(-dx-c)}}{16d} + \frac{e^{(-3dx-3c)}}{48d}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a)^2,x, algorithm="giac")`

output `1/32*e^(2*b*x + 3*d*x + 2*a + 3*c)/(2*b + 3*d) + 3/32*e^(2*b*x + d*x + 2*a + c)/(2*b + d) + 3/32*e^(2*b*x - d*x + 2*a - c)/(2*b - d) + 1/32*e^(2*b*x - 3*d*x + 2*a - 3*c)/(2*b - 3*d) - 1/32*e^(-2*b*x + 3*d*x - 2*a + 3*c)/(2*b - 3*d) - 3/32*e^(-2*b*x + d*x - 2*a + c)/(2*b - d) - 3/32*e^(-2*b*x - d*x - 2*a - c)/(2*b + d) - 1/32*e^(-2*b*x - 3*d*x - 2*a - 3*c)/(2*b + 3*d) - 1/48*e^(3*d*x + 3*c)/d - 3/16*e^(d*x + c)/d + 3/16*e^(-d*x - c)/d + 1/48*e^(-3*d*x - 3*c)/d`

Mupad [B] (verification not implemented)

Time = 1.52 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.34

$$\begin{aligned}
& \int \cosh^3(c + dx) \sinh^2(a + bx) dx \\
&= \frac{\cosh(c + dx)^2 \sinh(a + bx)^2 \sinh(c + dx) (8b^4 - 26b^2 d^2 + 9d^4)}{d (16b^4 - 40b^2 d^2 + 9d^4)} \\
&\quad - \sinh(a + bx)^2 \sinh(c + dx)^3 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} + \frac{1}{3d} \right) \\
&\quad - \frac{2 \cosh(a + bx) \cosh(c + dx)^3 \sinh(a + bx) (7bd^2 - 4b^3)}{16b^4 - 40b^2 d^2 + 9d^4} \\
&\quad - \frac{2 \cosh(a + bx)^2 \cosh(c + dx)^2 \sinh(c + dx) (4b^4 - 7b^2 d^2)}{d (16b^4 - 40b^2 d^2 + 9d^4)} \\
&\quad - \cosh(a + bx)^2 \sinh(c + dx)^3 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} - \frac{1}{3d} \right) \\
&\quad + \frac{12bd^2 \cosh(a + bx) \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)^2}{16b^4 - 40b^2 d^2 + 9d^4}
\end{aligned}$$

input `int(cosh(c + d*x)^3*sinh(a + b*x)^2,x)`output `(cosh(c + d*x)^2*sinh(a + b*x)^2*sinh(c + d*x)*(8*b^4 + 9*d^4 - 26*b^2*d^2))/ (d*(16*b^4 + 9*d^4 - 40*b^2*d^2)) - sinh(a + b*x)^2*sinh(c + d*x)^3*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) + 1/(3*d)) - (2*cosh(a + b*x)*cosh(c + d*x)^3*sinh(a + b*x)*(7*b*d^2 - 4*b^3))/(16*b^4 + 9*d^4 - 40*b^2*d^2) - (2*cosh(a + b*x)^2*cosh(c + d*x)^2*sinh(c + d*x)*(4*b^4 - 7*b^2*d^2))/(d*(16*b^4 + 9*d^4 - 40*b^2*d^2)) - cosh(a + b*x)^2*sinh(c + d*x)^3*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) - 1/(3*d)) + (12*b*d^2*cosh(a + b*x)*cosh(c + d*x)*sinh(a + b*x)*sinh(c + d*x)^2)/(16*b^4 + 9*d^4 - 40*b^2*d^2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.01

$$\int \cosh^3(c + dx) \sinh^2(a + bx) dx$$

$$= \frac{-36e^{4bx+4dx+4a+4c}b^2d^2 - 162e^{4bx+4dx+4a+4c}bd^3 + 72e^{4bx+2dx+4a+2c}b^3d + 36e^{4bx+2dx+4a+2c}b^2d^2 - 162e^{4bx+2c}}{b^2d^2}$$

input `int(cosh(d*x+c)^3*sinh(b*x+a)^2,x)`

output

```
(24*e**(4*a + 4*b*x + 6*c + 6*d*x)*b**3*d - 36*e**(4*a + 4*b*x + 6*c + 6*d*x)*b**2*d**2 - 6*e**(4*a + 4*b*x + 6*c + 6*d*x)*b*d**3 + 9*e**(4*a + 4*b*x + 6*c + 6*d*x)*d**4 + 72*e**(4*a + 4*b*x + 4*c + 4*d*x)*b**3*d - 36*e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d**2 - 162*e**(4*a + 4*b*x + 4*c + 4*d*x)*b*d**3 + 81*e**(4*a + 4*b*x + 4*c + 4*d*x)*d**4 + 72*e**(4*a + 4*b*x + 2*c + 2*d*x)*b**3*d + 36*e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d**2 - 162*e**(4*a + 4*b*x + 2*c + 2*d*x)*b*d**3 - 81*e**(4*a + 4*b*x + 2*c + 2*d*x)*d**4 + 24*e**(4*a + 4*b*x)*b**3*d + 36*e**(4*a + 4*b*x)*b**2*d**2 - 6*e**(4*a + 4*b*x)*b*d**3 - 9*e**(4*a + 4*b*x)*d**4 - 32*e**(2*a + 2*b*x + 6*c + 6*d*x)*b**4 + 80*e**(2*a + 2*b*x + 6*c + 6*d*x)*b**2*d**2 - 18*e**(2*a + 2*b*x + 6*c + 6*d*x)*d**4 - 288*e**(2*a + 2*b*x + 4*c + 4*d*x)*b**4 + 720*e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2*d**2 - 162*e**(2*a + 2*b*x + 4*c + 4*d*x)*d**4 + 288*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**4 - 720*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2*d**2 + 162*e**(2*a + 2*b*x + 2*c + 2*d*x)*d**4 + 32*e**(2*a + 2*b*x)*b**4 - 80*e**(2*a + 2*b*x)*b**2*d**2 + 18*e**(2*a + 2*b*x)*d**4 - 24*e**(6*c + 6*d*x)*b**3*d - 36*e**(6*c + 6*d*x)*b**2*d**2 + 6*e**(6*c + 6*d*x)*b*d**3 + 9*e**(6*c + 6*d*x)*d**4 - 72*e**(4*c + 4*d*x)*b**3*d - 36*e**(4*c + 4*d*x)*b**2*d**2 + 162*e**(4*c + 4*d*x)*b*d**3 + 81*e**(4*c + 4*d*x)*d**4 - 72*e**(2*c + 2*d*x)*b**3*d + 36*e**(2*c + 2*d*x)*b**2*d**2 + 162*e**(2*c + 2*d*x)*b*d**3 - 81*e**(2*c + 2*d*x)*d**4 - 24*b**3*d + 36*...
```


3.23 $\int \cosh(c + dx) \sinh^3(a + bx) dx$

Optimal result	224
Mathematica [A] (verified)	225
Rubi [A] (verified)	225
Maple [A] (verified)	226
Fricas [B] (verification not implemented)	227
Sympy [B] (verification not implemented)	227
Maxima [F(-2)]	228
Giac [B] (verification not implemented)	229
Mupad [B] (verification not implemented)	229
Reduce [B] (verification not implemented)	230

Optimal result

Integrand size = 15, antiderivative size = 97

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = -\frac{3 \cosh(a - c + (b - d)x)}{8(b - d)} + \frac{\cosh(3a - c + (3b - d)x)}{8(3b - d)} - \frac{3 \cosh(a + c + (b + d)x)}{8(b + d)} + \frac{\cosh(3a + c + (3b + d)x)}{8(3b + d)}$$

```
output -3*cosh(a-c+(b-d)*x)/(8*b-8*d)+cosh(3*a-c+(3*b-d)*x)/(24*b-8*d)-3*cosh(a+c+(b+d)*x)/(8*b+8*d)+cosh(3*a+c+(3*b+d)*x)/(24*b+8*d)
```

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.93

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = \frac{1}{8} \left(-\frac{3 \cosh(a - c + bx - dx)}{b - d} + \frac{\cosh(3a - c + 3bx - dx)}{3b - d} + \frac{\cosh(3a + c + 3bx + dx)}{3b + d} - \frac{3 \cosh(a + c + (b + d)x)}{b + d} \right)$$

input `Integrate[Cosh[c + d*x]*Sinh[a + b*x]^3,x]`

output $\frac{((-3*\text{Cosh}[a - c + b*x - d*x])/(b - d) + \text{Cosh}[3*a - c + 3*b*x - d*x]/(3*b - d) + \text{Cosh}[3*a + c + 3*b*x + d*x]/(3*b + d) - (3*\text{Cosh}[a + c + (b + d)*x])/(b + d))/8}$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^3(a + bx) \cosh(c + dx) dx$$

↓ 6152

$$\int \left(-\frac{3}{8} \sinh(a + x(b - d) - c) + \frac{1}{8} \sinh(3a + x(3b - d) - c) - \frac{3}{8} \sinh(a + x(b + d) + c) + \frac{1}{8} \sinh(3a + x(3b + d) + c) \right) dx$$

↓ 2009

$$-\frac{3 \cosh(a + x(b - d) - c)}{8(b - d)} + \frac{\cosh(3a + x(3b - d) - c)}{8(3b - d)} - \frac{3 \cosh(a + x(b + d) + c)}{8(b + d)} + \frac{\cosh(3a + x(3b + d) + c)}{8(3b + d)}$$

input `Int[Cosh[c + d*x]*Sinh[a + b*x]^3,x]`

output `(-3*Cosh[a - c + (b - d)*x])/(8*(b - d)) + Cosh[3*a - c + (3*b - d)*x]/(8*(3*b - d)) - (3*Cosh[a + c + (b + d)*x])/(8*(b + d)) + Cosh[3*a + c + (3*b + d)*x]/(8*(3*b + d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 2.53 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.93

method	result
default	$-\frac{3 \cosh(a-c+(b-d)x)}{8(b-d)} - \frac{3 \cosh(a+c+(b+d)x)}{8(b+d)} + \frac{\cosh(3a-c+(3b-d)x)}{24b-8d} + \frac{\cosh(3a+c+(3b+d)x)}{24b+8d}$
parallelrisch	$\frac{-12 \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^6 b^3 + 24b^2 d \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^5 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right) + \left(-12 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^2 b d^2 + 36b^3 - 12b d^2\right) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^4 + (-6 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^3 b d^2 + 36b^3 - 12b d^2) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^3 + \left(-12 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^2 b d^2 + 36b^3 - 12b d^2\right) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + (-6 \tanh\left(\frac{dx}{2} + \frac{c}{2}\right) b d^2 + 36b^3 - 12b d^2) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) + (-6 b d^2 + 36b^3 - 12b d^2) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) + (-6 b d^2 + 36b^3 - 12b d^2)}{9 \left(-1 + \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3 \left(1 + \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}$
risch	$\frac{(3b^3 e^{6bx+6a} - b^2 d e^{6bx+6a} - 3b d^2 e^{6bx+6a} + d^3 e^{6bx+6a} - 27b^3 e^{4bx+4a} + 27b^2 d e^{4bx+4a} + 3b d^2 e^{4bx+4a} - 3d^3 e^{4bx+4a} - 27b^3 e^{2bx+2a} + 27b^2 d e^{2bx+2a} + 27b d^2 e^{2bx+2a} - 27d^3 e^{2bx+2a}) \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)^6 + \dots}{16(3b+d)(b+d)(3b-d)(b-d)}$
orering	Expression too large to display

input `int(cosh(d*x+c)*sinh(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
-3/8*cosh(a-c+(b-d)*x)/(b-d)-3/8*cosh(a+c+(b+d)*x)/(b+d)+1/8*cosh(3*a-c+(3
*b-d)*x)/(3*b-d)+1/8*cosh(3*a+c+(3*b+d)*x)/(3*b+d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(89) = 178$.

Time = 0.10 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.51

$$\int \cosh(c + dx) \sinh^3(a + bx) dx$$

$$= \frac{9(b^3 - bd^2) \cosh(bx + a) \cosh(dx + c) \sinh(bx + a)^2 + 3((b^3 - bd^2) \cosh(bx + a)^3 - (9b^3 - bd^2) \cosh(bx + a) \sinh(bx + a)^2) \sinh(dx + c) + 3((b^3 - bd^2) \cosh(bx + a)^3 - (9b^3 - bd^2) \cosh(bx + a) \sinh(bx + a)^2) \sinh(dx + c)}{4((9b^4 - 10b^2d^2 + d^4) \cosh(bx + a)^4 - 2(9b^4 - 10b^2d^2 + d^4) \cosh(bx + a)^2 \sinh(bx + a)^2 + (9b^4 - 10b^2d^2 + d^4) \sinh(bx + a)^4)}$$

input

```
integrate(cosh(d*x+c)*sinh(b*x+a)^3,x, algorithm="fricas")
```

output

```
1/4*(9*(b^3 - b*d^2)*cosh(b*x + a)*cosh(d*x + c)*sinh(b*x + a)^2 + 3*((b^3
- b*d^2)*cosh(b*x + a)^3 - (9*b^3 - b*d^2)*cosh(b*x + a))*cosh(d*x + c) -
((b^2*d - d^3)*sinh(b*x + a)^3 - 3*(9*b^2*d - d^3 - (b^2*d - d^3)*cosh(b*
x + a)^2)*sinh(b*x + a))*sinh(d*x + c))/((9*b^4 - 10*b^2*d^2 + d^4)*cosh(b
*x + a)^4 - 2*(9*b^4 - 10*b^2*d^2 + d^4)*cosh(b*x + a)^2*sinh(b*x + a)^2 +
(9*b^4 - 10*b^2*d^2 + d^4)*sinh(b*x + a)^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs. $2(76) = 152$.

Time = 1.97 (sec) , antiderivative size = 937, normalized size of antiderivative = 9.66

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(d*x+c)*sinh(b*x+a)**3,x)
```

output

```
Piecewise((x*sinh(a)**3*cosh(c), Eq(b, 0) & Eq(d, 0)), (3*x*sinh(a - d*x)*
*3*cosh(c + d*x)/8 + 3*x*sinh(a - d*x)**2*sinh(c + d*x)*cosh(a - d*x)/8 -
3*x*sinh(a - d*x)*cosh(a - d*x)**2*cosh(c + d*x)/8 - 3*x*sinh(c + d*x)*cos
h(a - d*x)**3/8 + 5*sinh(a - d*x)**3*sinh(c + d*x)/(8*d) - 3*sinh(a - d*x)
*sinh(c + d*x)*cosh(a - d*x)**2/(4*d) - 3*cosh(a - d*x)**3*cosh(c + d*x)/(
8*d), Eq(b, -d)), (x*sinh(a - d*x/3)**3*cosh(c + d*x)/8 + 3*x*sinh(a - d*x
/3)**2*sinh(c + d*x)*cosh(a - d*x/3)/8 + 3*x*sinh(a - d*x/3)*cosh(a - d*x/
3)**2*cosh(c + d*x)/8 + x*sinh(c + d*x)*cosh(a - d*x/3)**3/8 + 7*sinh(a -
d*x/3)**3*sinh(c + d*x)/(8*d) - 3*sinh(a - d*x/3)*sinh(c + d*x)*cosh(a - d
*x/3)**2/(4*d) - 3*cosh(a - d*x/3)**3*cosh(c + d*x)/(8*d), Eq(b, -d/3)), (
x*sinh(a + d*x/3)**3*cosh(c + d*x)/8 - 3*x*sinh(a + d*x/3)**2*sinh(c + d*x
)*cosh(a + d*x/3)/8 + 3*x*sinh(a + d*x/3)*cosh(a + d*x/3)**2*cosh(c + d*x)
/8 - x*sinh(c + d*x)*cosh(a + d*x/3)**3/8 + 7*sinh(a + d*x/3)**3*sinh(c +
d*x)/(8*d) - 3*sinh(a + d*x/3)*sinh(c + d*x)*cosh(a + d*x/3)**2/(4*d) + 3*
cosh(a + d*x/3)**3*cosh(c + d*x)/(8*d), Eq(b, d/3)), (3*x*sinh(a + d*x)**3
*cosh(c + d*x)/8 - 3*x*sinh(a + d*x)**2*sinh(c + d*x)*cosh(a + d*x)/8 - 3*
x*sinh(a + d*x)*cosh(a + d*x)**2*cosh(c + d*x)/8 + 3*x*sinh(c + d*x)*cosh(
a + d*x)**3/8 - sinh(a + d*x)**3*sinh(c + d*x)/(8*d) + 3*sinh(a + d*x)**2*
cosh(a + d*x)*cosh(c + d*x)/(4*d) - 3*cosh(a + d*x)**3*cosh(c + d*x)/(8*d)
, Eq(b, d)), (9*b**3*sinh(a + b*x)**2*cosh(a + b*x)*cosh(c + d*x)/(9*b...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)*sinh(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(-d/b>0)', see `assume?` for more
details)I
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. $2(89) = 178$.

Time = 0.12 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.89

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = \frac{e^{(3bx+dx+3a+c)}}{16(3b+d)} + \frac{e^{(3bx-dx+3a-c)}}{16(3b-d)} - \frac{3e^{(bx+dx+a+c)}}{16(b+d)} - \frac{3e^{(bx-dx+a-c)}}{16(b-d)} - \frac{3e^{(-bx+dx-a+c)}}{16(b-d)} - \frac{3e^{(-bx-dx-a-c)}}{16(b+d)} + \frac{e^{(-3bx+dx-3a+c)}}{16(3b-d)} + \frac{e^{(-3bx-dx-3a-c)}}{16(3b+d)}$$

input `integrate(cosh(d*x+c)*sinh(b*x+a)^3,x, algorithm="giac")`

output `1/16*e^(3*b*x + d*x + 3*a + c)/(3*b + d) + 1/16*e^(3*b*x - d*x + 3*a - c)/(3*b - d) - 3/16*e^(b*x + d*x + a + c)/(b + d) - 3/16*e^(b*x - d*x + a - c)/(b - d) - 3/16*e^(-b*x + d*x - a + c)/(b - d) - 3/16*e^(-b*x - d*x - a - c)/(b + d) + 1/16*e^(-3*b*x + d*x - 3*a + c)/(3*b - d) + 1/16*e^(-3*b*x - d*x - 3*a - c)/(3*b + d)`

Mupad [B] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.89

$$\int \cosh(c + dx) \sinh^3(a + bx) dx = \frac{6b^2 d \cosh(a + bx)^2 \sinh(a + bx) \sinh(c + dx)}{9b^4 - 10b^2 d^2 + d^4} - \frac{d \sinh(a + bx)^3 \sinh(c + dx) (7b^2 - d^2)}{9b^4 - 10b^2 d^2 + d^4} - \frac{3 \cosh(a + bx) \cosh(c + dx) \sinh(a + bx)^2 (bd^2 - 3b^3)}{9b^4 - 10b^2 d^2 + d^4} - \frac{6b^3 \cosh(a + bx)^3 \cosh(c + dx)}{9b^4 - 10b^2 d^2 + d^4}$$

input `int(cosh(c + d*x)*sinh(a + b*x)^3,x)`

output

$$\begin{aligned} & (6*b^2*d*cosh(a + b*x)^2*sinh(a + b*x)*sinh(c + d*x))/(9*b^4 + d^4 - 10*b^2*d^2) - (d*sinh(a + b*x)^3*sinh(c + d*x)*(7*b^2 - d^2))/(9*b^4 + d^4 - 10*b^2*d^2) - (3*cosh(a + b*x)*cosh(c + d*x)*sinh(a + b*x)^2*(b*d^2 - 3*b^3))/(9*b^4 + d^4 - 10*b^2*d^2) - (6*b^3*cosh(a + b*x)^3*cosh(c + d*x))/(9*b^4 + d^4 - 10*b^2*d^2) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 573, normalized size of antiderivative = 5.91

$$\int \cosh(c + dx) \sinh^3(a + bx) dx$$

$$= \frac{27e^{4bx+2dx+4a+2c}b^2d + 3b^3 + d^3 - 3e^{6bx+2dx+6a+2c}bd^2 - 3e^{6bx+6a}bd^2 + 3e^{4bx+2dx+4a+2c}bd^2 - 3e^{4bx+2dx+4a+2c}bd^2}{}$$

input

int(cosh(d*x+c)*sinh(b*x+a)^3,x)

output

$$\begin{aligned} & (3*e^{(6*a + 6*b*x + 2*c + 2*d*x)*b**3} - e^{(6*a + 6*b*x + 2*c + 2*d*x)*b**2*d} - 3*e^{(6*a + 6*b*x + 2*c + 2*d*x)*b*d**2} + e^{(6*a + 6*b*x + 2*c + 2*d*x)*d**3} + 3*e^{(6*a + 6*b*x)*b**3} + e^{(6*a + 6*b*x)*b**2*d} - 3*e^{(6*a + 6*b*x)*b*d**2} - e^{(6*a + 6*b*x)*d**3} - 27*e^{(4*a + 4*b*x + 2*c + 2*d*x)*b**3} + 27*e^{(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d} + 3*e^{(4*a + 4*b*x + 2*c + 2*d*x)*b*d**2} - 3*e^{(4*a + 4*b*x + 2*c + 2*d*x)*d**3} - 27*e^{(4*a + 4*b*x)*b**3} - 27*e^{(4*a + 4*b*x)*b**2*d} + 3*e^{(4*a + 4*b*x)*b*d**2} + 3*e^{(4*a + 4*b*x)*d**3} - 27*e^{(2*a + 2*b*x + 2*c + 2*d*x)*b**3} - 27*e^{(2*a + 2*b*x + 2*c + 2*d*x)*b**2*d} + 3*e^{(2*a + 2*b*x + 2*c + 2*d*x)*b*d**2} + 3*e^{(2*a + 2*b*x + 2*c + 2*d*x)*d**3} - 27*e^{(2*a + 2*b*x)*b**3} + 27*e^{(2*a + 2*b*x)*b**2*d} + 3*e^{(2*a + 2*b*x)*b*d**2} - 3*e^{(2*a + 2*b*x)*d**3} + 3*e^{(2*c + 2*d*x)*b**3} + e^{(2*c + 2*d*x)*b**2*d} - 3*e^{(2*c + 2*d*x)*b*d**2} - e^{(2*c + 2*d*x)*d**3} + 3*b**3 - b**2*d - 3*b*d**2 + d**3)/(16*e^{(3*a + 3*b*x + c + d*x)*(9*b**4 - 10*b**2*d**2 + d**4)}) \end{aligned}$$

3.24 $\int \cosh^2(c + dx) \sinh^3(a + bx) dx$

Optimal result	231
Mathematica [A] (verified)	232
Rubi [A] (verified)	232
Maple [A] (verified)	234
Fricas [B] (verification not implemented)	234
Sympy [B] (verification not implemented)	235
Maxima [F(-2)]	236
Giac [B] (verification not implemented)	237
Mupad [B] (verification not implemented)	238
Reduce [B] (verification not implemented)	239

Optimal result

Integrand size = 17, antiderivative size = 138

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx = -\frac{3 \cosh(a + bx)}{8b} + \frac{\cosh(3a + 3bx)}{24b} - \frac{3 \cosh(a - 2c + (b - 2d)x)}{16(b - 2d)} + \frac{\cosh(3a - 2c + (3b - 2d)x)}{16(3b - 2d)} - \frac{3 \cosh(a + 2c + (b + 2d)x)}{16(b + 2d)} + \frac{\cosh(3a + 2c + (3b + 2d)x)}{16(3b + 2d)}$$

output

```
-3/8*cosh(b*x+a)/b+1/24*cosh(3*b*x+3*a)/b-3*cosh(a-2*c+(b-2*d)*x)/(16*b-32*d)+cosh(3*a-2*c+(3*b-2*d)*x)/(48*b-32*d)-3*cosh(a+2*c+(b+2*d)*x)/(16*b+32*d)+cosh(3*a+2*c+(3*b+2*d)*x)/(48*b+32*d)
```


Mathematica [A] (verified)

Time = 1.09 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.11

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx = \frac{1}{48} \left(-\frac{18 \cosh(a) \cosh(bx)}{b} + \frac{2 \cosh(3a) \cosh(3bx)}{b} - \frac{9 \cosh(a - 2c + bx - 2dx)}{b - 2d} + \frac{3 \cosh(3a - 2c + 3bx - 2dx)}{3b - 2d} - \frac{9 \cosh(a + 2c + bx + 2dx)}{b + 2d} + \frac{3 \cosh(3a + 2c + 3bx + 2dx)}{3b + 2d} - \frac{18 \sinh(a) \sinh(bx)}{b} + \frac{2 \sinh(3a) \sinh(3bx)}{b} \right)$$

input

```
Integrate[Cosh[c + d*x]^2*Sinh[a + b*x]^3,x]
```

output

```
((-18*Cosh[a]*Cosh[b*x])/b + (2*Cosh[3*a]*Cosh[3*b*x])/b - (9*Cosh[a - 2*c + b*x - 2*d*x])/(b - 2*d) + (3*Cosh[3*a - 2*c + 3*b*x - 2*d*x])/(3*b - 2*d) - (9*Cosh[a + 2*c + b*x + 2*d*x])/(b + 2*d) + (3*Cosh[3*a + 2*c + 3*b*x + 2*d*x])/(3*b + 2*d) - (18*Sinh[a]*Sinh[b*x])/b + (2*Sinh[3*a]*Sinh[3*b*x])/b)/48
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^3(a + bx) \cosh^2(c + dx) dx$$

↓ 6152

$$\int \left(-\frac{3}{16} \sinh(a + x(b - 2d) - 2c) + \frac{1}{16} \sinh(3a + x(3b - 2d) - 2c) - \frac{3}{16} \sinh(a + x(b + 2d) + 2c) + \frac{1}{16} \sinh(3a + x(3b + 2d) + 2c) \right) dx$$

↓ 2009

$$-\frac{3 \cosh(a + x(b - 2d) - 2c)}{16(b - 2d)} + \frac{\cosh(3a + x(3b - 2d) - 2c)}{16(3b - 2d)} - \frac{3 \cosh(a + x(b + 2d) + 2c)}{16(b + 2d)} + \frac{\cosh(3a + x(3b + 2d) + 2c)}{16(3b + 2d)} - \frac{3 \cosh(a + bx)}{8b} + \frac{\cosh(3a + 3bx)}{24b}$$

input `Int[Cosh[c + d*x]^2*Sinh[a + b*x]^3,x]`

output `(-3*Cosh[a + b*x])/(8*b) + Cosh[3*a + 3*b*x]/(24*b) - (3*Cosh[a - 2*c + (b - 2*d)*x])/(16*(b - 2*d)) + Cosh[3*a - 2*c + (3*b - 2*d)*x]/(16*(3*b - 2*d)) - (3*Cosh[a + 2*c + (b + 2*d)*x])/(16*(b + 2*d)) + Cosh[3*a + 2*c + (3*b + 2*d)*x]/(16*(3*b + 2*d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 8.03 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.92

method	result
default	$-\frac{3 \cosh(bx+a)}{8b} + \frac{\cosh(3bx+3a)}{24b} - \frac{3 \cosh(a-2c+(b-2d)x)}{16(b-2d)} - \frac{3 \cosh(a+2c+(b+2d)x)}{16(b+2d)} + \frac{\cosh(3a-2c+(3b-2d)x)}{48b-32d} +$
parallelrisch	$9\left(b+\frac{2d}{3}\right)b(b-2d)(b+2d) \cosh(3a-2c+(3b-2d)x)+9b(b-2d)(b+2d)\left(b-\frac{2d}{3}\right) \cosh(3a+2c+(3b+2d)x)-81\left(b+\frac{2d}{3}\right)b(b+2d)\left(b-\frac{2d}{3}\right)$
risch	$\frac{e^{3bx+3a}}{48b} - \frac{3e^{bx+a}}{16b} - \frac{3e^{-bx-a}}{16b} + \frac{e^{-3bx-3a}}{48b} + \frac{(3b^3e^{6bx+6a}-2b^2de^{6bx+6a}-12bd^2e^{6bx+6a}+8d^3e^{6bx+6a}-27b^3e^{4bx+4a})}{48b^3}$
orering	Expression too large to display

input `int(cosh(d*x+c)^2*sinh(b*x+a)^3,x,method=_RETURNVERBOSE)`output
$$-\frac{3}{8} \cosh(bx+a)/b + \frac{1}{24} \cosh(3bx+3a)/b - \frac{3}{16} \cosh(a-2c+(b-2d)x)/(b-2d) - \frac{3}{16} \cosh(a+2c+(b+2d)x)/(b+2d) + \frac{1}{16} \cosh(3a-2c+(3b-2d)x)/(3b-2d) + \frac{1}{16} \cosh(3a+2c+(3b+2d)x)/(3b+2d)$$
Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 443 vs. 2(126) = 252.

Time = 0.12 (sec) , antiderivative size = 443, normalized size of antiderivative = 3.21

$$\int \cosh^2(c+dx) \sinh^3(a+bx) dx$$

$$= \frac{(9b^4 - 40b^2d^2 + 16d^4) \cosh(bx+a)^3 + 9((b^4 - 4b^2d^2) \cosh(bx+a)^3 - (9b^4 - 4b^2d^2) \cosh(bx+a)) \cosh(bx+a)}{48b^3}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a)^3,x, algorithm="fricas")`

output

```

1/24*((9*b^4 - 40*b^2*d^2 + 16*d^4)*cosh(b*x + a)^3 + 9*((b^4 - 4*b^2*d^2)
*cosh(b*x + a)^3 - (9*b^4 - 4*b^2*d^2)*cosh(b*x + a))*cosh(d*x + c)^2 + 3*
(9*(b^4 - 4*b^2*d^2)*cosh(b*x + a)*cosh(d*x + c)^2 + (9*b^4 - 40*b^2*d^2 +
16*d^4)*cosh(b*x + a))*sinh(b*x + a)^2 + 9*((b^4 - 4*b^2*d^2)*cosh(b*x +
a)^3 + 3*(b^4 - 4*b^2*d^2)*cosh(b*x + a)*sinh(b*x + a)^2 - (9*b^4 - 4*b^2*
d^2)*cosh(b*x + a))*sinh(d*x + c)^2 - 9*(9*b^4 - 40*b^2*d^2 + 16*d^4)*cosh
(b*x + a) - 12*((b^3*d - 4*b*d^3)*cosh(d*x + c)*sinh(b*x + a)^3 - 3*(9*b^3
*d - 4*b*d^3 - (b^3*d - 4*b*d^3)*cosh(b*x + a)^2)*cosh(d*x + c)*sinh(b*x +
a))*sinh(d*x + c))/((9*b^5 - 40*b^3*d^2 + 16*b*d^4)*cosh(b*x + a)^4 - 2*(
9*b^5 - 40*b^3*d^2 + 16*b*d^4)*cosh(b*x + a)^2*sinh(b*x + a)^2 + (9*b^5 -
40*b^3*d^2 + 16*b*d^4)*sinh(b*x + a)^4)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2030 vs. $2(116) = 232$.

Time = 5.34 (sec) , antiderivative size = 2030, normalized size of antiderivative = 14.71

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(d*x+c)**2*sinh(b*x+a)**3,x)
```

output

```
Piecewise((x*sinh(a)**3*cosh(c)**2, Eq(b, 0) & Eq(d, 0)), ((-x*sinh(c + d*x)**2/2 + x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*sinh(a)**3, Eq(b, 0)), (3*x*sinh(a - 2*d*x)**3*sinh(c + d*x)**2/16 + 3*x*sinh(a - 2*d*x)**3*cosh(c + d*x)**2/16 + 3*x*sinh(a - 2*d*x)**2*sinh(c + d*x)*cosh(a - 2*d*x)*cosh(c + d*x)/8 - 3*x*sinh(a - 2*d*x)*sinh(c + d*x)**2*cosh(a - 2*d*x)**2/16 - 3*x*sinh(a - 2*d*x)*cosh(a - 2*d*x)**2*cosh(c + d*x)**2/16 - 3*x*sinh(c + d*x)*cosh(a - 2*d*x)**3*cosh(c + d*x)/8 + 13*sinh(a - 2*d*x)**3*sinh(c + d*x)*cosh(c + d*x)/(16*d) + sinh(a - 2*d*x)**2*sinh(c + d*x)**2*cosh(a - 2*d*x)/(2*d) - 7*sinh(a - 2*d*x)*sinh(c + d*x)*cosh(a - 2*d*x)**2*cosh(c + d*x)/(8*d) - 49*sinh(c + d*x)**2*cosh(a - 2*d*x)**3/(96*d) - 17*cosh(a - 2*d*x)**3*cosh(c + d*x)**2/(96*d), Eq(b, -2*d)), (x*sinh(a - 2*d*x/3)**3*sinh(c + d*x)**2/16 + x*sinh(a - 2*d*x/3)**3*cosh(c + d*x)**2/16 + 3*x*sinh(a - 2*d*x/3)**2*sinh(c + d*x)*cosh(a - 2*d*x/3)*cosh(c + d*x)/8 + 3*x*sinh(a - 2*d*x/3)*sinh(c + d*x)**2*cosh(a - 2*d*x/3)**2/16 + 3*x*sinh(a - 2*d*x/3)*cosh(a - 2*d*x/3)**2*cosh(c + d*x)**2/16 + x*sinh(c + d*x)*cosh(a - 2*d*x/3)**3*cosh(c + d*x)/8 + 15*sinh(a - 2*d*x/3)**3*sinh(c + d*x)*cosh(c + d*x)/(16*d) + 3*sinh(a - 2*d*x/3)**2*sinh(c + d*x)**2*cosh(a - 2*d*x/3)/(2*d) + 9*sinh(a - 2*d*x/3)*sinh(c + d*x)*cosh(a - 2*d*x/3)**2*cosh(c + d*x)/(8*d) - 11*sinh(c + d*x)**2*cosh(a - 2*d*x/3)**3/(32*d) + 21*cosh(a - 2*d*x/3)**3*cosh(c + d*x)**2/(32*d), Eq(b, -2*d/3)), (x...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)^2*sinh(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(2*d)/b>0)', see `assume?` for more detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 256 vs. $2(126) = 252$.

Time = 0.13 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.86

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx = \frac{e^{(3bx+2dx+3a+2c)}}{32(3b+2d)} + \frac{e^{(3bx-2dx+3a-2c)}}{32(3b-2d)} + \frac{e^{(3bx+3a)}}{48b}$$

$$- \frac{3e^{(bx+2dx+a+2c)}}{32(b+2d)} - \frac{3e^{(bx-2dx+a-2c)}}{32(b-2d)} - \frac{16b}{3e^{(bx+a)}}$$

$$- \frac{3e^{(-bx+2dx-a+2c)}}{32(b-2d)} - \frac{3e^{(-bx-2dx-a-2c)}}{32(b+2d)} - \frac{16b}{3e^{(-bx-a)}}$$

$$+ \frac{e^{(-3bx+2dx-3a+2c)}}{32(3b-2d)} + \frac{e^{(-3bx-2dx-3a-2c)}}{32(3b+2d)}$$

$$+ \frac{e^{(-3bx-3a)}}{48b}$$

input `integrate(cosh(d*x+c)^2*sinh(b*x+a)^3,x, algorithm="giac")`

output `1/32*e^(3*b*x + 2*d*x + 3*a + 2*c)/(3*b + 2*d) + 1/32*e^(3*b*x - 2*d*x + 3*a - 2*c)/(3*b - 2*d) + 1/48*e^(3*b*x + 3*a)/b - 3/32*e^(b*x + 2*d*x + a + 2*c)/(b + 2*d) - 3/32*e^(b*x - 2*d*x + a - 2*c)/(b - 2*d) - 3/16*e^(b*x + a)/b - 3/32*e^(-b*x + 2*d*x - a + 2*c)/(b - 2*d) - 3/32*e^(-b*x - 2*d*x - a - 2*c)/(b + 2*d) - 3/16*e^(-b*x - a)/b + 1/32*e^(-3*b*x + 2*d*x - 3*a + 2*c)/(3*b - 2*d) + 1/32*e^(-3*b*x - 2*d*x - 3*a - 2*c)/(3*b + 2*d) + 1/48*e^(-3*b*x - 3*a)/b`

Mupad [B] (verification not implemented)

Time = 1.53 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.44

$$\begin{aligned}
& \int \cosh^2(c + dx) \sinh^3(a + bx) dx \\
&= \frac{\cosh(a + bx) \cosh(c + dx)^2 \sinh(a + bx)^2 (9b^4 - 26b^2 d^2 + 8d^4)}{b (9b^4 - 40b^2 d^2 + 16d^4)} \\
&\quad - \cosh(a + bx)^3 \sinh(c + dx)^2 \left(\frac{3b^3}{9b^4 - 40b^2 d^2 + 16d^4} - \frac{1}{3b} \right) \\
&\quad - \cosh(a + bx)^3 \cosh(c + dx)^2 \left(\frac{3b^3}{9b^4 - 40b^2 d^2 + 16d^4} + \frac{1}{3b} \right) \\
&\quad - \frac{2d \cosh(c + dx) \sinh(a + bx)^3 \sinh(c + dx) (7b^2 - 4d^2)}{9b^4 - 40b^2 d^2 + 16d^4} \\
&\quad + \frac{12b^2 d \cosh(a + bx)^2 \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)}{9b^4 - 40b^2 d^2 + 16d^4} \\
&\quad + \frac{2d^2 \cosh(a + bx) \sinh(a + bx)^2 \sinh(c + dx)^2 (7b^2 - 4d^2)}{b (9b^4 - 40b^2 d^2 + 16d^4)}
\end{aligned}$$

input `int(cosh(c + d*x)^2*sinh(a + b*x)^3,x)`output `(cosh(a + b*x)*cosh(c + d*x)^2*sinh(a + b*x)^2*(9*b^4 + 8*d^4 - 26*b^2*d^2))/ (b*(9*b^4 + 16*d^4 - 40*b^2*d^2)) - cosh(a + b*x)^3*sinh(c + d*x)^2*((3*b^3)/(9*b^4 + 16*d^4 - 40*b^2*d^2) - 1/(3*b)) - cosh(a + b*x)^3*cosh(c + d*x)^2*((3*b^3)/(9*b^4 + 16*d^4 - 40*b^2*d^2) + 1/(3*b)) - (2*d*cosh(c + d*x)*sinh(a + b*x)^3*sinh(c + d*x)*(7*b^2 - 4*d^2))/(9*b^4 + 16*d^4 - 40*b^2*d^2) + (12*b^2*d*cosh(a + b*x)^2*cosh(c + d*x)*sinh(a + b*x)*sinh(c + d*x))/(9*b^4 + 16*d^4 - 40*b^2*d^2) + (2*d^2*cosh(a + b*x)*sinh(a + b*x)^2*sinh(c + d*x)^2*(7*b^2 - 4*d^2))/(b*(9*b^4 + 16*d^4 - 40*b^2*d^2))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.27

$$\int \cosh^2(c + dx) \sinh^3(a + bx) dx$$

$$= \frac{9b^4 + 36e^{4bx+4dx+4a+4c}b^2d^2 - 72e^{4bx+4dx+4a+4c}bd^3 + 720e^{4bx+2dx+4a+2c}b^2d^2 - 162e^{4bx+4a}b^3d + 36e^{4bx+4a}b^2d^2}{1}$$

input

```
int(cosh(d*x+c)^2*sinh(b*x+a)^3,x)
```

output

```
(9***e**(6*a + 6*b*x + 4*c + 4*d*x)*b**4 - 6***e**(6*a + 6*b*x + 4*c + 4*d*x)*
b**3*d - 36***e**(6*a + 6*b*x + 4*c + 4*d*x)*b**2*d**2 + 24***e**(6*a + 6*b*x
+ 4*c + 4*d*x)*b*d**3 + 18***e**(6*a + 6*b*x + 2*c + 2*d*x)*b**4 - 80***e**(6*
a + 6*b*x + 2*c + 2*d*x)*b**2*d**2 + 32***e**(6*a + 6*b*x + 2*c + 2*d*x)*d**
4 + 9***e**(6*a + 6*b*x)*b**4 + 6***e**(6*a + 6*b*x)*b**3*d - 36***e**(6*a + 6*b
*x)*b**2*d**2 - 24***e**(6*a + 6*b*x)*b*d**3 - 81***e**(4*a + 4*b*x + 4*c + 4*
d*x)*b**4 + 162***e**(4*a + 4*b*x + 4*c + 4*d*x)*b**3*d + 36***e**(4*a + 4*b*x
+ 4*c + 4*d*x)*b**2*d**2 - 72***e**(4*a + 4*b*x + 4*c + 4*d*x)*b*d**3 - 162
***e**(4*a + 4*b*x + 2*c + 2*d*x)*b**4 + 720***e**(4*a + 4*b*x + 2*c + 2*d*x)*
b**2*d**2 - 288***e**(4*a + 4*b*x + 2*c + 2*d*x)*d**4 - 81***e**(4*a + 4*b*x)*
b**4 - 162***e**(4*a + 4*b*x)*b**3*d + 36***e**(4*a + 4*b*x)*b**2*d**2 + 72***e
*(4*a + 4*b*x)*b*d**3 - 81***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**4 - 162***e**(2
*a + 2*b*x + 4*c + 4*d*x)*b**3*d + 36***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2*
d**2 + 72***e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d**3 - 162***e**(2*a + 2*b*x + 2*
c + 2*d*x)*b**4 + 720***e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2*d**2 - 288***e**(2
*a + 2*b*x + 2*c + 2*d*x)*d**4 - 81***e**(2*a + 2*b*x)*b**4 + 162***e**(2*a +
2*b*x)*b**3*d + 36***e**(2*a + 2*b*x)*b**2*d**2 - 72***e**(2*a + 2*b*x)*b*d**3
+ 9***e**(4*c + 4*d*x)*b**4 + 6***e**(4*c + 4*d*x)*b**3*d - 36***e**(4*c + 4*d*
x)*b**2*d**2 - 24***e**(4*c + 4*d*x)*b*d**3 + 18***e**(2*c + 2*d*x)*b**4 - 80*
e**(2*c + 2*d*x)*b**2*d**2 + 32***e**(2*c + 2*d*x)*d**4 + 9*b**4 - 6*b**3...
```


3.25 $\int \cosh^3(c + dx) \sinh^3(a + bx) dx$

Optimal result	240
Mathematica [A] (verified)	241
Rubi [A] (verified)	241
Maple [A] (verified)	243
Fricas [B] (verification not implemented)	243
Sympy [B] (verification not implemented)	244
Maxima [F(-2)]	245
Giac [B] (verification not implemented)	246
Mupad [B] (verification not implemented)	247
Reduce [F]	248

Optimal result

Integrand size = 17, antiderivative size = 195

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = -\frac{3 \cosh(a - 3c + (b - 3d)x)}{32(b - 3d)} - \frac{9 \cosh(a - c + (b - d)x)}{32(b - d)} + \frac{\cosh(3(a - c) + 3(b - d)x)}{96(b - d)} + \frac{3 \cosh(3a - c + (3b - d)x)}{32(3b - d)} - \frac{9 \cosh(a + c + (b + d)x)}{32(b + d)} + \frac{\cosh(3(a + c) + 3(b + d)x)}{96(b + d)} + \frac{3 \cosh(3a + c + (3b + d)x)}{32(3b + d)} - \frac{3 \cosh(a + 3c + (b + 3d)x)}{32(b + 3d)}$$

output

```
-3*cosh(a-3*c+(b-3*d)*x)/(32*b-96*d)-9*cosh(a-c+(b-d)*x)/(32*b-32*d)+cosh(
3*a-3*c+3*(b-d)*x)/(96*b-96*d)+3*cosh(3*a-c+(3*b-d)*x)/(96*b-32*d)-9*cosh(
a+c+(b+d)*x)/(32*b+32*d)+cosh(3*a+3*c+3*(b+d)*x)/(96*b+96*d)+3*cosh(3*a+c+
(3*b+d)*x)/(96*b+32*d)-3*cosh(a+3*c+(b+3*d)*x)/(32*b+96*d)
```

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.90

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \frac{1}{96} \left(-\frac{9 \cosh(a - 3c + bx - 3dx)}{b - 3d} - \frac{27 \cosh(a - c + bx - dx)}{b - d} + \frac{\cosh(3(a - c + bx - dx))}{b - d} + \frac{9 \cosh(3a - c + 3bx - dx)}{3b - d} + \frac{9 \cosh(3a + c + 3bx + dx)}{3b + d} - \frac{9 \cosh(a + 3c + bx + 3dx)}{b + 3d} - \frac{27 \cosh(a + c + (b + d)x)}{b + d} + \frac{\cosh(3(a + c + (b + d)x))}{b + d} \right)$$

input `Integrate[Cosh[c + d*x]^3*Sinh[a + b*x]^3,x]`

output `((-9*Cosh[a - 3*c + b*x - 3*d*x])/(b - 3*d) - (27*Cosh[a - c + b*x - d*x])/(b - d) + Cosh[3*(a - c + b*x - d*x)]/(b - d) + (9*Cosh[3*a - c + 3*b*x - d*x])/(3*b - d) + (9*Cosh[3*a + c + 3*b*x + d*x])/(3*b + d) - (9*Cosh[a + 3*c + b*x + 3*d*x])/(b + 3*d) - (27*Cosh[a + c + (b + d)*x])/(b + d) + Cosh[3*(a + c + (b + d)*x)]/(b + d))/96`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^3(a + bx) \cosh^3(c + dx) dx$$

↓ 6152

$$\int \left(-\frac{3}{32} \sinh(a + x(b - 3d) - 3c) - \frac{9}{32} \sinh(a + x(b - d) - c) + \frac{1}{32} \sinh(3(a - c) + 3x(b - d)) + \frac{3}{32} \sinh(3a + \dots \right)$$

↓ 2009

$$\begin{aligned} & -\frac{3 \cosh(a + x(b - 3d) - 3c)}{32(b - 3d)} - \frac{9 \cosh(a + x(b - d) - c)}{32(b - d)} + \frac{\cosh(3(a - c) + 3x(b - d))}{96(b - d)} + \\ & \frac{3 \cosh(3a + x(3b - d) - c)}{32(3b - d)} - \frac{9 \cosh(a + x(b + d) + c)}{32(b + d)} + \frac{\cosh(3(a + c) + 3x(b + d))}{96(b + d)} + \\ & \frac{3 \cosh(3a + x(3b + d) + c)}{32(3b + d)} - \frac{3 \cosh(a + x(b + 3d) + 3c)}{32(b + 3d)} \end{aligned}$$

input

```
Int[Cosh[c + d*x]^3*Sinh[a + b*x]^3,x]
```

output

```
(-3*Cosh[a - 3*c + (b - 3*d)*x])/(32*(b - 3*d)) - (9*Cosh[a - c + (b - d)*
x])/(32*(b - d)) + Cosh[3*(a - c) + 3*(b - d)*x]/(96*(b - d)) + (3*Cosh[3*
a - c + (3*b - d)*x])/(32*(3*b - d)) - (9*Cosh[a + c + (b + d)*x])/(32*(b
+ d)) + Cosh[3*(a + c) + 3*(b + d)*x]/(96*(b + d)) + (3*Cosh[3*a + c + (3*
b + d)*x])/(32*(3*b + d)) - (3*Cosh[a + 3*c + (b + 3*d)*x])/(32*(b + 3*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6152

```
Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v
]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 24.37 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.97

method	result
default	$-\frac{3 \cosh(a-3c+(b-3d)x)}{32(b-3d)} - \frac{9 \cosh(a-c+(b-d)x)}{32(b-d)} - \frac{9 \cosh(a+c+(b+d)x)}{32(b+d)} - \frac{3 \cosh(a+3c+(b+3d)x)}{32(b+3d)} + \frac{\cosh((3b-3d)x)}{96b-9}$
parallelrisch	$9(b-3d)\left(b-\frac{d}{3}\right)(b+d)\left(b+\frac{d}{3}\right)(b+3d) \cosh((3b-3d)x+3a-3c)+27(b-3d)(b+d)\left(b+\frac{d}{3}\right)(b-d)(b+3d) \cosh(3a-c+(3b-d)x)+9$
risch	$\frac{(b^3 e^{6bx+6a} - b^2 d e^{6bx+6a} - 9b d^2 e^{6bx+6a} + 9d^3 e^{6bx+6a} - 9b^3 e^{4bx+4a} + 27b^2 d e^{4bx+4a} + 9b d^2 e^{4bx+4a} - 27d^3 e^{4bx+4a} - 9b^3 e^{2bx+2a})}{192(b+d)(b+3d)(b-d)(b-3d)}$
orering	Expression too large to display

input `int(cosh(d*x+c)^3*sinh(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
-3/32*cosh(a-3*c+(b-3*d)*x)/(b-3*d)-9/32*cosh(a-c+(b-d)*x)/(b-d)-9/32*cosh
(a+c+(b+d)*x)/(b+d)-3/32*cosh(a+3*c+(b+3*d)*x)/(b+3*d)+1/32*cosh((3*b-3*d)
*x+3*a-3*c)/(3*b-3*d)+3/32*cosh(3*a-c+(3*b-d)*x)/(3*b-d)+3/32*cosh(3*a+c+(
3*b+d)*x)/(3*b+d)+1/32*cosh((3*b+3*d)*x+3*a+3*c)/(3*b+3*d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 729 vs. 2(179) = 358.

Time = 0.11 (sec) , antiderivative size = 729, normalized size of antiderivative = 3.74

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \text{Too large to display}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a)^3,x, algorithm="fricas")`

output

```

1/48*(((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)^3 - 9*(9*b^5 - 10*b^3*d^2 + b*d^4)*cosh(b*x + a))*cosh(d*x + c)^3 - ((9*b^4*d - 82*b^2*d^3 + 9*d^5)*sinh(b*x + a)^3 - 3*(81*b^4*d - 90*b^2*d^3 + 9*d^5 - (9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2)*sinh(b*x + a))*sinh(d*x + c)^3 + 3*((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)*cosh(d*x + c)^3 + 27*(b^5 - 10*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)*cosh(d*x + c))*sinh(b*x + a)^2 + 3*(3*(9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)*cosh(d*x + c)*sinh(b*x + a)^2 + ((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)^3 - 9*(9*b^5 - 10*b^3*d^2 + b*d^4)*cosh(b*x + a))*cosh(d*x + c))*sinh(d*x + c)^2 + 27*((b^5 - 10*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)^3 - (9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a))*cosh(d*x + c) - 3*((3*b^4*d - 30*b^2*d^3 + 27*d^5 + (9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(d*x + c)^2)*sinh(b*x + a)^3 - 3*(27*b^4*d - 246*b^2*d^3 + 27*d^5 - 3*(b^4*d - 10*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2 + (81*b^4*d - 90*b^2*d^3 + 9*d^5 - (9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2)*cosh(d*x + c)^2)*sinh(b*x + a))*sinh(d*x + c))/((9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*cosh(b*x + a)^4 - 2*(9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*cosh(b*x + a)^2*sinh(b*x + a)^2 + (9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*sinh(b*x + a)^4)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3580 vs. $2(172) = 344$.

Time = 16.48 (sec) , antiderivative size = 3580, normalized size of antiderivative = 18.36

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(d*x+c)**3*sinh(b*x+a)**3,x)
```

output

```
Piecewise((x*sinh(a)**3*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (9*x*sinh(a - 3*d*x)**3*sinh(c + d*x)**2*cosh(c + d*x)/32 + 3*x*sinh(a - 3*d*x)**3*cosh(c + d*x)**3/32 + 3*x*sinh(a - 3*d*x)**2*sinh(c + d*x)**3*cosh(a - 3*d*x)/32 + 9*x*sinh(a - 3*d*x)**2*sinh(c + d*x)*cosh(a - 3*d*x)*cosh(c + d*x)**2/32 - 9*x*sinh(a - 3*d*x)*sinh(c + d*x)**2*cosh(a - 3*d*x)**2*cosh(c + d*x)/32 - 3*x*sinh(a - 3*d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**3/32 - 3*x*sinh(c + d*x)**3*cosh(a - 3*d*x)**3/32 - 9*x*sinh(c + d*x)*cosh(a - 3*d*x)**3*cosh(c + d*x)**2/32 + sinh(a - 3*d*x)**3*sinh(c + d*x)**3/(30*d) - 61*sinh(a - 3*d*x)**3*sinh(c + d*x)*cosh(c + d*x)**2/(320*d) - 117*sinh(a - 3*d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)**3/(320*d) - 11*sinh(a - 3*d*x)*sinh(c + d*x)**3*cosh(a - 3*d*x)**2/(320*d) + 3*sinh(a - 3*d*x)*sinh(c + d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**2/(20*d) - 3*sinh(c + d*x)**2*cosh(a - 3*d*x)**3*cosh(c + d*x)/(320*d) + cosh(a - 3*d*x)**3*cosh(c + d*x)**3/(4*d), Eq(b, -3*d)), (-3*x*sinh(a - d*x)**3*sinh(c + d*x)**2*cosh(c + d*x)/16 + 5*x*sinh(a - d*x)**3*cosh(c + d*x)**3/16 - 3*x*sinh(a - d*x)**2*sinh(c + d*x)**3*cosh(a - d*x)/16 + 9*x*sinh(a - d*x)**2*sinh(c + d*x)*cosh(a - d*x)*cosh(c + d*x)**2/16 + 9*x*sinh(a - d*x)*sinh(c + d*x)**2*cosh(a - d*x)**2*cosh(c + d*x)/16 - 3*x*sinh(a - d*x)*cosh(a - d*x)**2*cosh(c + d*x)**3/16 + 5*x*sinh(c + d*x)**3*cosh(a - d*x)**3/16 - 3*x*sinh(c + d*x)*cosh(a - d*x)**3*cosh(c + d*x)**2/16 - 7*sinh(a - d*x)**3*sinh(c + d*x)**3/(48*d) + ...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(d*x+c)^3*sinh(b*x+a)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for more detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(179) = 358$.

Time = 0.13 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.91

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \frac{e^{(3bx+3dx+3a+3c)}}{192(b+d)} + \frac{3e^{(3bx+dx+3a+c)}}{64(3b+d)} + \frac{3e^{(3bx-dx+3a-c)}}{64(3b-d)} + \frac{e^{(3bx-3dx+3a-3c)}}{192(b-d)} - \frac{3e^{(bx+3dx+a+3c)}}{64(b+3d)} - \frac{9e^{(bx+dx+a+c)}}{64(b+d)} - \frac{9e^{(bx-dx+a-c)}}{64(b-d)} - \frac{3e^{(bx-3dx+a-3c)}}{64(b-3d)} - \frac{3e^{(-bx+3dx-a+3c)}}{64(b-3d)} - \frac{9e^{(-bx+dx-a+c)}}{64(b-d)} - \frac{9e^{(-bx-dx-a-c)}}{3e^{(-bx-3dx-a-3c)}} - \frac{64(b+d)}{e^{(-3bx+3dx-3a+3c)}} - \frac{64(b+3d)}{3e^{(-3bx+dx-3a+c)}} + \frac{192(b-d)}{3e^{(-3bx-dx-3a-c)}} + \frac{64(3b-d)}{e^{(-3bx-3dx-3a-3c)}} + \frac{64(3b+d)}{192(b+d)}$$

input `integrate(cosh(d*x+c)^3*sinh(b*x+a)^3,x, algorithm="giac")`

output

```
1/192*e^(3*b*x + 3*d*x + 3*a + 3*c)/(b + d) + 3/64*e^(3*b*x + d*x + 3*a +
c)/(3*b + d) + 3/64*e^(3*b*x - d*x + 3*a - c)/(3*b - d) + 1/192*e^(3*b*x -
3*d*x + 3*a - 3*c)/(b - d) - 3/64*e^(b*x + 3*d*x + a + 3*c)/(b + 3*d) - 9
/64*e^(b*x + d*x + a + c)/(b + d) - 9/64*e^(b*x - d*x + a - c)/(b - d) - 3
/64*e^(b*x - 3*d*x + a - 3*c)/(b - 3*d) - 3/64*e^(-b*x + 3*d*x - a + 3*c)/
(b - 3*d) - 9/64*e^(-b*x + d*x - a + c)/(b - d) - 9/64*e^(-b*x - d*x - a -
c)/(b + d) - 3/64*e^(-b*x - 3*d*x - a - 3*c)/(b + 3*d) + 1/192*e^(-3*b*x
+ 3*d*x - 3*a + 3*c)/(b - d) + 3/64*e^(-3*b*x + d*x - 3*a + c)/(3*b - d) +
3/64*e^(-3*b*x - d*x - 3*a - c)/(3*b + d) + 1/192*e^(-3*b*x - 3*d*x - 3*a
- 3*c)/(b + d)
```

Mupad [B] (verification not implemented)

Time = 1.64 (sec) , antiderivative size = 908, normalized size of antiderivative = 4.66

$$\begin{aligned}
\int \cosh^3(c + dx) \sinh^3(a + bx) dx = & -e^{3a+c+3bx+dx} \left(\frac{-9b^3 + 3b^2d + 9bd^2 - 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& + \frac{e^{-6a-6bx}(-9b^3 - 3b^2d + 9bd^2 + 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& - \frac{e^{-2a-2bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
& - e^{3a-c+3bx-dx} \left(\frac{-9b^3 - 3b^2d + 9bd^2 + 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& + \frac{e^{-6a-6bx}(-9b^3 + 3b^2d + 9bd^2 - 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& - \frac{e^{-2a-2bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
& - e^{3a-3c+3bx-3dx} \left(\frac{-b^3 - b^2d + 9bd^2 + 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& + \frac{e^{-6a-6bx}(-b^3 + b^2d + 9bd^2 - 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& - \frac{e^{-2a-2bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right) \\
& - e^{3a+3c+3bx+3dx} \left(\frac{-b^3 + b^2d + 9bd^2 - 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& + \frac{e^{-6a-6bx}(-b^3 - b^2d + 9bd^2 + 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& - \frac{e^{-2a-2bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right)
\end{aligned}$$

input `int(cosh(c + d*x)^3*sinh(a + b*x)^3,x)`

output

```

- exp(3*a + c + 3*b*x + d*x)*((9*b*d^2 + 3*b^2*d - 9*b^3 - 3*d^3)/(576*b^4
+ 64*d^4 - 640*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b*d^2 - 3*b^2*d - 9*b^3
+ 3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 2*a - 2*b*x)*(9*b*d^2
+ 81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 4*
a - 4*b*x)*(9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*
b^2*d^2)) - exp(3*a - c + 3*b*x - d*x)*((9*b*d^2 - 3*b^2*d - 9*b^3 + 3*d^3
)/(576*b^4 + 64*d^4 - 640*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b*d^2 + 3*b^2*
d - 9*b^3 - 3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 2*a - 2*b*x)
*(9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) -
(exp(- 4*a - 4*b*x)*(9*b*d^2 + 81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*
d^4 - 640*b^2*d^2)) - exp(3*a - 3*c + 3*b*x - 3*d*x)*((9*b*d^2 - b^2*d - b
^3 + 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) + (exp(- 6*a - 6*b*x)*(9*b
*d^2 + b^2*d - b^3 - 9*d^3))/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(-
2*a - 2*b*x)*(9*b*d^2 - 27*b^2*d - 9*b^3 + 27*d^3))/(192*b^4 + 1728*d^4 -
1920*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))
/(192*b^4 + 1728*d^4 - 1920*b^2*d^2)) - exp(3*a + 3*c + 3*b*x + 3*d*x)*((9
*b*d^2 + b^2*d - b^3 - 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) + (exp(-
6*a - 6*b*x)*(9*b*d^2 - b^2*d - b^3 + 9*d^3))/(192*b^4 + 1728*d^4 - 1920*
b^2*d^2) - (exp(- 2*a - 2*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))/(192
*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 - 27*b^2...

```

Reduce [F]

$$\int \cosh^3(c + dx) \sinh^3(a + bx) dx = \int \cosh(dx + c)^3 \sinh(bx + a)^3 dx$$

input

```
int(cosh(d*x+c)^3*sinh(b*x+a)^3,x)
```

output

```
int(cosh(d*x+c)^3*sinh(b*x+a)^3,x)
```

3.26 $\int \sinh(x) \tanh(2x) dx$

Optimal result	249
Mathematica [A] (verified)	249
Rubi [A] (verified)	250
Maple [C] (verified)	251
Fricas [B] (verification not implemented)	252
Sympy [F]	252
Maxima [B] (verification not implemented)	253
Giac [B] (verification not implemented)	253
Mupad [B] (verification not implemented)	254
Reduce [B] (verification not implemented)	254

Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \sinh(x) \tanh(2x) dx = -\frac{\arctan(\sqrt{2} \sinh(x))}{\sqrt{2}} + \sinh(x)$$

output `-1/2*arctan(sinh(x)*2^(1/2))*2^(1/2)+sinh(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sinh(x) \tanh(2x) dx = -\frac{\arctan(\sqrt{2} \sinh(x))}{\sqrt{2}} + \sinh(x)$$

input `Integrate[Sinh[x]*Tanh[2*x],x]`

output `-(ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2]) + Sinh[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 25, 4878, 27, 262, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \tanh(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin(ix) \tan(2ix) dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin(ix) \tan(2ix) dx \\
 & \quad \downarrow \text{4878} \\
 & -\int -\frac{2 \sinh^2(x)}{2 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int \frac{\sinh^2(x)}{2 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{262} \\
 & 2 \left(\frac{\sinh(x)}{2} - \frac{1}{2} \int \frac{1}{2 \sinh^2(x) + 1} d \sinh(x) \right) \\
 & \quad \downarrow \text{216} \\
 & 2 \left(\frac{\sinh(x)}{2} - \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}} \right)
 \end{aligned}$$

input

```
Int [Sinh [x] *Tanh [2*x] , x]
```

output

```
2*(-1/2*ArcTan[Sqrt [2] *Sinh [x]] /Sqrt [2] + Sinh [x] /2)
```

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.84

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x - 1)}{4} - \frac{i\sqrt{2} \ln(e^{2x} + i\sqrt{2}e^x - 1)}{4}$	54

input `int(sinh(x)*tanh(2*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/4*I*2^(1/2)*ln(exp(2*x)-I*2^(1/2)*exp(x)-1)-1/4*I*2^(1/2)*ln(exp(2*x)+I*2^(1/2)*exp(x)-1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(15) = 30$.

Time = 0.09 (sec) , antiderivative size = 115, normalized size of antiderivative = 6.05

$$\int \sinh(x) \tanh(2x) dx = \frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \arctan\left(\frac{1}{2} \sqrt{2} \cosh(x) + \frac{1}{2} \sqrt{2} \sinh(x)\right) - (\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \arctan\left(\frac{1}{2} \sqrt{2} \cosh(x) - \frac{1}{2} \sqrt{2} \sinh(x)\right)}{2(\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)*tanh(2*x),x, algorithm="fricas")`

output `-1/2*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)) - (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2)))/(cosh(x) - sinh(x))) - cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + 1)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \sinh(x) \tanh(2x) dx = \int \sinh(x) \tanh(2x) dx$$

input `integrate(sinh(x)*tanh(2*x),x)`

output `Integral(sinh(x)*tanh(2*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. $2(15) = 30$.

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.79

$$\int \sinh(x) \tanh(2x) dx = \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 e^{-x}) \right) + \frac{1}{2} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2 e^{-x}) \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(sinh(x)*tanh(2*x),x, algorithm="maxima")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(-x))) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^(-x))) - 1/2*e^(-x) + 1/2*e^x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(15) = 30$.

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.89

$$\int \sinh(x) \tanh(2x) dx = -\frac{1}{4} \sqrt{2} \left(\pi + 2 \arctan \left(\frac{1}{2} \sqrt{2} (e^{2x} - 1) e^{-x} \right) \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(sinh(x)*tanh(2*x),x, algorithm="giac")`

output `-1/4*sqrt(2)*(pi + 2*arctan(1/2*sqrt(2)*(e^(2*x) - 1)*e^(-x))) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.47

$$\int \sinh(x) \tanh(2x) dx = \frac{e^x}{2} - \frac{e^{-x}}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2} + \frac{\sqrt{2}e^{3x}}{2}\right)}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2}\right)}{2}$$

input `int(tanh(2*x)*sinh(x),x)`output `exp(x)/2 - exp(-x)/2 - (2^(1/2)*atan((2^(1/2)*exp(x))/2 + (2^(1/2)*exp(3*x))/2))/2 - (2^(1/2)*atan((2^(1/2)*exp(x))/2))/2`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 58, normalized size of antiderivative = 3.05

$$\int \sinh(x) \tanh(2x) dx = \frac{-e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) + e^{2x} - 1}{2e^x}$$

input `int(sinh(x)*tanh(2*x),x)`output `(- e**x*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - e**x*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) + e**(2*x) - 1)/(2*e**x)`

3.27 $\int \sinh(x) \tanh(3x) dx$

Optimal result	255
Mathematica [A] (verified)	255
Rubi [A] (verified)	256
Maple [C] (verified)	258
Fricas [B] (verification not implemented)	258
Sympy [F]	259
Maxima [B] (verification not implemented)	259
Giac [B] (verification not implemented)	260
Mupad [B] (verification not implemented)	260
Reduce [B] (verification not implemented)	260

Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \sinh(x) \tanh(3x) dx = -\frac{1}{3} \arctan(\sinh(x)) - \frac{1}{3} \arctan(2 \sinh(x)) + \sinh(x)$$

output `-1/3*arctan(sinh(x))-1/3*arctan(2*sinh(x))+sinh(x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \sinh(x) \tanh(3x) dx = -\frac{1}{3} \arctan(\sinh(x)) - \frac{1}{3} \arctan(2 \sinh(x)) + \sinh(x)$$

input `Integrate[Sinh[x]*Tanh[3*x],x]`

output `-1/3*ArcTan[Sinh[x]] - ArcTan[2*Sinh[x]]/3 + Sinh[x]`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 25, 4878, 25, 1602, 27, 1477, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \tanh(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin(ix) \tan(3ix) dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin(ix) \tan(3ix) dx \\
 & \quad \downarrow \text{4878} \\
 & -\int -\frac{\sinh^2(x) (4 \sinh^2(x) + 3)}{4 \sinh^4(x) + 5 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{\sinh^2(x) (4 \sinh^2(x) + 3)}{4 \sinh^4(x) + 5 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{1602} \\
 & \sinh(x) - \frac{1}{4} \int \frac{4(2 \sinh^2(x) + 1)}{4 \sinh^4(x) + 5 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & \sinh(x) - \int \frac{2 \sinh^2(x) + 1}{4 \sinh^4(x) + 5 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{1477} \\
 & -\frac{2}{3} \int \frac{1}{4 \sinh^2(x) + 1} d \sinh(x) - \frac{4}{3} \int \frac{1}{4 \sinh^2(x) + 4} d \sinh(x) + \sinh(x) \\
 & \quad \downarrow \text{216}
 \end{aligned}$$

$$-\frac{1}{3} \arctan(\sinh(x)) - \frac{1}{3} \arctan(2 \sinh(x)) + \sinh(x)$$

input `Int[Sinh[x]*Tanh[3*x],x]`

output `-1/3*ArcTan[Sinh[x]] - ArcTan[2*Sinh[x]]/3 + Sinh[x]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 1477 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(e/2 + (2*c*d - b*e)/(2*q)) Int[1/(b/2 - q/2 + c*x^2), x], x] + Simp[(e/2 - (2*c*d - b*e)/(2*q)) Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - a*e^2, 0] && GtQ[b^2 - 4*a*c, 0]`

rule 1602 `Int[((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[e*f*(f*x)^(m-1)*((a + b*x^2 + c*x^4)^(p+1)/(c*(m+4*p+3))), x] - Simp[f^2/(c*(m+4*p+3)) Int[(f*x)^(m-2)*(a + b*x^2 + c*x^4)^p*Simp[a*e*(m-1) + (b*e*(m+2*p+1) - c*d*(m+4*p+3))*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 1] && NeQ[m+4*p+3, 0] && IntegerQ[2*p] && (IntegerQ[p] | IntegerQ[m])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 3.16

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i \ln(e^x - i)}{3} - \frac{i \ln(e^x + i)}{3} + \frac{i \ln(e^{2x} - ie^x - 1)}{6} - \frac{i \ln(e^{2x} + ie^x - 1)}{6}$	60

input `int(sinh(x)*tanh(3*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/3*I*ln(exp(x)-I)-1/3*I*ln(exp(x)+I)+1/6*I*ln(exp(2*x)-I*exp(x)-1)-1/6*I*ln(exp(2*x)+I*exp(x)-1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(15) = 30$.

Time = 0.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 4.00

$$\int \sinh(x) \tanh(3x) dx$$

$$= \frac{2 (\cosh(x) + \sinh(x)) \arctan\left(-\frac{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2}{\cosh(x) - \sinh(x)}\right) - 6 (\cosh(x) + \sinh(x)) \arctan(\cosh(x))}{6 (\cosh(x) + \sinh(x))}$$

input `integrate(sinh(x)*tanh(3*x),x, algorithm="fricas")`

output

```
1/6*(2*(cosh(x) + sinh(x))*arctan(-(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)
)^2)/(cosh(x) - sinh(x))) - 6*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)
) + 3*cosh(x)^2 + 6*cosh(x)*sinh(x) + 3*sinh(x)^2 - 3)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \sinh(x) \tanh(3x) dx = \int \sinh(x) \tanh(3x) dx$$

input

```
integrate(sinh(x)*tanh(3*x),x)
```

output

```
Integral(sinh(x)*tanh(3*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(15) = 30$.

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.42

$$\int \sinh(x) \tanh(3x) dx = \frac{1}{3} \arctan(\sqrt{3} + 2e^{-x}) + \frac{1}{3} \arctan(-\sqrt{3} + 2e^{-x}) + \frac{2}{3} \arctan(e^{-x}) - \frac{1}{2}e^{-x} + \frac{1}{2}e^x$$

input

```
integrate(sinh(x)*tanh(3*x),x, algorithm="maxima")
```

output

```
1/3*arctan(sqrt(3) + 2*e^(-x)) + 1/3*arctan(-sqrt(3) + 2*e^(-x)) + 2/3*arc
tan(e^(-x)) - 1/2*e^(-x) + 1/2*e^x
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(15) = 30$.

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.26

$$\int \sinh(x) \tanh(3x) dx = -\frac{1}{3} \pi - \frac{1}{3} \arctan \left((e^{2x} - 1)e^{-x} \right) - \frac{1}{3} \arctan \left(\frac{1}{2} (e^{2x} - 1)e^{-x} \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(sinh(x)*tanh(3*x),x, algorithm="giac")`

output `-1/3*pi - 1/3*arctan((e^(2*x) - 1)*e^(-x)) - 1/3*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.21

$$\int \sinh(x) \tanh(3x) dx = \frac{e^x}{2} - \operatorname{atan}(e^x) - \frac{\operatorname{atan}(e^{3x})}{3} - \frac{e^{-x}}{2}$$

input `int(tanh(3*x)*sinh(x),x)`

output `exp(x)/2 - atan(exp(x)) - atan(exp(3*x))/3 - exp(-x)/2`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.89

$$\int \sinh(x) \tanh(3x) dx = \frac{-4e^x \operatorname{atan}(e^x) - 2e^x \operatorname{atan}(2e^x - \sqrt{3}) - 2e^x \operatorname{atan}(2e^x + \sqrt{3}) + 3e^{2x} - 3}{6e^x}$$

input `int(sinh(x)*tanh(3*x),x)`

output $(-4e^{3x}\operatorname{atan}(e^{3x}) - 2e^{3x}\operatorname{atan}(2e^{3x} - \sqrt{3})) - 2e^{3x}\operatorname{atan}(2e^{3x} + \sqrt{3}) + 3e^{2x} - 3)/(6e^{3x})$

3.28 $\int \sinh(x) \tanh(4x) dx$

Optimal result	262
Mathematica [A] (verified)	262
Rubi [A] (verified)	263
Maple [C] (verified)	265
Fricas [B] (verification not implemented)	266
Sympy [F]	266
Maxima [F]	267
Giac [A] (verification not implemented)	267
Mupad [B] (verification not implemented)	267
Reduce [B] (verification not implemented)	268

Optimal result

Integrand size = 7, antiderivative size = 75

$$\int \sinh(x) \tanh(4x) dx = -\frac{\arctan\left(\sqrt{2(2-\sqrt{2})} \sinh(x)\right)}{2\sqrt{2(2-\sqrt{2})}} - \frac{\arctan\left(\sqrt{2(2+\sqrt{2})} \sinh(x)\right)}{2\sqrt{2(2+\sqrt{2})}} + \sinh(x)$$

output `-1/2*arctan((4-2*2^(1/2))^(1/2)*sinh(x))/(4-2*2^(1/2))^(1/2)-1/2*arctan((4+2*2^(1/2))^(1/2)*sinh(x))/(4+2*2^(1/2))^(1/2)+sinh(x)`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.92

$$\int \sinh(x) \tanh(4x) dx = -\frac{1}{4}\sqrt{2-\sqrt{2}} \arctan\left(\frac{2\sinh(x)}{\sqrt{2-\sqrt{2}}}\right) - \frac{1}{4}\sqrt{2+\sqrt{2}} \arctan\left(\frac{2\sinh(x)}{\sqrt{2+\sqrt{2}}}\right) + \sinh(x)$$

input `Integrate[Sinh[x]*Tanh[4*x],x]`

output `-1/4*(Sqrt[2 - Sqrt[2]]*ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[2]]]) - (Sqrt[2 + Sqrt[2]]*ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[2]]])/4 + Sinh[x]`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 25, 4878, 27, 1602, 27, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \tanh(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin(ix) \tan(4ix) dx \\
 & \quad \downarrow \text{25} \\
 & -\int \sin(ix) \tan(4ix) dx \\
 & \quad \downarrow \text{4878} \\
 & -\int -\frac{4 \sinh^2(x) (2 \sinh^2(x) + 1)}{8 \sinh^4(x) + 8 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & 4 \int \frac{\sinh^2(x) (2 \sinh^2(x) + 1)}{8 \sinh^4(x) + 8 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{1602} \\
 & 4 \left(\frac{\sinh(x)}{4} - \frac{1}{8} \int \frac{2(4 \sinh^2(x) + 1)}{8 \sinh^4(x) + 8 \sinh^2(x) + 1} d \sinh(x) \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$4\left(\frac{\sinh(x)}{4} - \frac{1}{4} \int \frac{4 \sinh^2(x) + 1}{8 \sinh^4(x) + 8 \sinh^2(x) + 1} d \sinh(x)\right)$$

↓ 1480

$$4\left(\frac{1}{4}\left(-\left((2-\sqrt{2}) \int \frac{1}{8 \sinh^2(x) + 2(2-\sqrt{2})} d \sinh(x)\right) - (2+\sqrt{2}) \int \frac{1}{8 \sinh^2(x) + 2(2+\sqrt{2})} d \sinh(x)\right)\right) +$$

↓ 216

$$4\left(\frac{1}{4}\left(-\frac{1}{4}\sqrt{2-\sqrt{2}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2-\sqrt{2}}}\right) - \frac{1}{4}\sqrt{2+\sqrt{2}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2+\sqrt{2}}}\right)\right) + \frac{\sinh(x)}{4}\right)$$

input

```
Int[Sinh[x]*Tanh[4*x],x]
```

output

```
4*((-1/4*(Sqrt[2 - Sqrt[2]]*ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[2]]]) - (Sqrt[2 + Sqrt[2]]*ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[2]]]))/4)/4 + Sinh[x]/4
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 1480

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(e/2 + (2*c*d - b*e)/(2*q)) Int[1/(b/2 - q/2 + c*x^2), x], x] + Simp[(e/2 - (2*c*d - b*e)/(2*q)) Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]
```

rule 1602

```
Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[e*f*(f*x)^(m - 1)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 3))), x] - Simp[f^2/(c*(m + 4*p + 3)) Int[(f*x)^(m - 2)*(a + b*x^2 + c*x^4)^p*Simp[a*e*(m - 1) + (b*e*(m + 2*p + 1) - c*d*(m + 4*p + 3))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 1] && NeQ[m + 4*p + 3, 0] && IntegerQ[2*p] && (IntegerQ[p] | IntegerQ[m])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4878

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.56

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \left(\sum_{_R=\text{RootOf}(2048_Z^4+128_Z^2+1)} _R \ln(-8_R e^x + e^{2x} - 1) \right)$	42

input

```
int(sinh(x)*tanh(4*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)-1/2*exp(-x)+sum(_R*ln(-8*_R*exp(x)+exp(2*x)-1),_R=RootOf(2048*_Z^4+128*_Z^2+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(51) = 102$.

Time = 0.10 (sec) , antiderivative size = 287, normalized size of antiderivative = 3.83

$$\int \sinh(x) \tanh(4x) dx$$

$$= \frac{\sqrt{\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{2}\left((\sqrt{2} - 2)\cosh(x)^3 + 3(\sqrt{2} - 2)\cosh(x)\sinh(x)^2 + (\sqrt{2} - 2)\right)\right) + \sqrt{\sqrt{2} - 2}(\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{2}\left((\sqrt{2} + 2)\cosh(x)^3 + 3(\sqrt{2} + 2)\cosh(x)\sinh(x)^2 + (\sqrt{2} + 2)\right)\right) + \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{2}\left((\sqrt{2} - 2)\cosh(x) + (\sqrt{2} - 2)\sinh(x)\right)\right) - \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{2}\left((\sqrt{2} + 2)\cosh(x) + (\sqrt{2} + 2)\sinh(x)\right)\right) + 2\cosh(x)^2 + 4\cosh(x)\sinh(x) + 2\sinh(x)^2 - 2}{\cosh(x) + \sinh(x)}$$

input `integrate(sinh(x)*tanh(4*x),x, algorithm="fricas")`

output

```
1/4*(sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x))*arctan(1/2*((sqrt(2) - 2)*cosh(x)^3 + 3*(sqrt(2) - 2)*cosh(x)*sinh(x)^2 + (sqrt(2) - 2)*sinh(x)^3 + (3*(sqrt(2) - 2)*cosh(x)^2 - sqrt(2))*sinh(x) - sqrt(2)*cosh(x))*sqrt(sqrt(2) + 2)) + sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x))*arctan(1/2*((sqrt(2) - 2)*cosh(x) + (sqrt(2) - 2)*sinh(x))*sqrt(sqrt(2) + 2)) - sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x))*arctan(1/2*((sqrt(2) + 2)*cosh(x)^3 + 3*(sqrt(2) + 2)*cosh(x)*sinh(x)^2 + (sqrt(2) + 2)*sinh(x)^3 + (3*(sqrt(2) + 2)*cosh(x)^2 - sqrt(2))*sinh(x) - sqrt(2)*cosh(x))*sqrt(-sqrt(2) + 2)) - sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x))*arctan(1/2*((sqrt(2) + 2)*cosh(x) + (sqrt(2) + 2)*sinh(x))*sqrt(-sqrt(2) + 2)) + 2*cosh(x)^2 + 4*cosh(x)*sinh(x) + 2*sinh(x)^2 - 2)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \sinh(x) \tanh(4x) dx = \int \sinh(x) \tanh(4x) dx$$

input `integrate(sinh(x)*tanh(4*x),x)`

output `Integral(sinh(x)*tanh(4*x), x)`

Maxima [F]

$$\int \sinh(x) \tanh(4x) dx = \int \sinh(x) \tanh(4x) dx$$

input `integrate(sinh(x)*tanh(4*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) - 1)*e^(-x) - 1/2*integrate(2*(e^(7*x) + e^x)/(e^(8*x) + 1), x)`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.95

$$\int \sinh(x) \tanh(4x) dx = -\frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan\left(-\frac{e^{(-x)} - e^x}{\sqrt{\sqrt{2} + 2}}\right) - \frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan\left(-\frac{e^{(-x)} - e^x}{\sqrt{-\sqrt{2} + 2}}\right) - \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(sinh(x)*tanh(4*x),x, algorithm="giac")`

output `-1/4*sqrt(sqrt(2) + 2)*arctan(-(e^(-x) - e^x)/sqrt(sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(-(e^(-x) - e^x)/sqrt(-sqrt(2) + 2)) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 1.76 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.95

$$\int \sinh(x) \tanh(4x) dx = \frac{e^x}{2} - \frac{e^{-x}}{2} - \frac{\operatorname{atan}\left(\frac{e^{-x}(e^{2x}-1)}{\sqrt{\sqrt{2}+2}}\right) \sqrt{\sqrt{2}+2}}{4} - \frac{\operatorname{atan}\left(\frac{e^{-x}(e^{2x}-1)}{\sqrt{2-\sqrt{2}}}\right) \sqrt{2-\sqrt{2}}}{4}$$

input `int(tanh(4*x)*sinh(x),x)`

output $\frac{\exp(x)/2 - \exp(-x)/2 - (\operatorname{atan}((\exp(-x) * (\exp(2*x) - 1)) / (2^{(1/2)} + 2)^{(1/2)})) * (2^{(1/2)} + 2)^{(1/2)} / 4 - (\operatorname{atan}((\exp(-x) * (\exp(2*x) - 1)) / (2 - 2^{(1/2)})^{(1/2)})) * (2 - 2^{(1/2)})^{(1/2)} / 4}{4e^x}$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.95

$$\int \sinh(x) \tanh(4x) dx$$

$$= \frac{e^x \sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right) - e^x \sqrt{\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}+2e^x}{\sqrt{\sqrt{2}+2}}\right) + e^x \sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}-2e^x}{\sqrt{-\sqrt{2}+2}}\right) - e^x \sqrt{-\sqrt{2} + 2} \operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}+2e^x}{\sqrt{-\sqrt{2}+2}}\right)}{4e^x}$$

input `int(sinh(x)*tanh(4*x),x)`

output $(e^{**x} * \sqrt{\sqrt{2} + 2} * \operatorname{atan}((\sqrt{-\sqrt{2} + 2} - 2 * e^{**x}) / \sqrt{\sqrt{2} + 2})) - e^{**x} * \sqrt{\sqrt{2} + 2} * \operatorname{atan}((\sqrt{-\sqrt{2} + 2} + 2 * e^{**x}) / \sqrt{\sqrt{2} + 2})) + e^{**x} * \sqrt{-\sqrt{2} + 2} * \operatorname{atan}((\sqrt{\sqrt{2} + 2} - 2 * e^{**x}) / \sqrt{-\sqrt{2} + 2})) - e^{**x} * \sqrt{-\sqrt{2} + 2} * \operatorname{atan}((\sqrt{\sqrt{2} + 2} + 2 * e^{**x}) / \sqrt{-\sqrt{2} + 2})) + 2 * e^{**x} * (2 * x - 2) / (4 * e^{**x})$

3.29 $\int \sinh(x) \tanh(5x) dx$

Optimal result	269
Mathematica [A] (verified)	270
Rubi [A] (verified)	270
Maple [C] (verified)	272
Fricas [B] (verification not implemented)	273
Sympy [F]	273
Maxima [F]	274
Giac [A] (verification not implemented)	274
Mupad [B] (verification not implemented)	275
Reduce [F]	275

Optimal result

Integrand size = 7, antiderivative size = 84

$$\int \sinh(x) \tanh(5x) dx = -\frac{1}{5} \arctan(\sinh(x)) - \frac{1}{5} \sqrt{\frac{1}{2}(3 + \sqrt{5})} \arctan\left(\sqrt{2(3 - \sqrt{5})} \sinh(x)\right) - \frac{1}{5} \sqrt{\frac{2}{3 + \sqrt{5}}} \arctan\left(\sqrt{2(3 + \sqrt{5})} \sinh(x)\right) + \sinh(x)$$

```
output -1/5*arctan(sinh(x))-1/5*(1/2+1/2*5^(1/2))*arctan((5^(1/2)-1)*sinh(x))-1/5
*2^(1/2)/(3+5^(1/2))^(1/2)*arctan((5^(1/2)+1)*sinh(x))+sinh(x)
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.96

$$\int \sinh(x) \tanh(5x) dx = \frac{1}{10} \left(-2 \arctan(\sinh(x)) \right. \\ \left. - \sqrt{2(3 + \sqrt{5})} \arctan \left(2\sqrt{\frac{2}{3 + \sqrt{5}}} \sinh(x) \right) \right. \\ \left. - \sqrt{6 - 2\sqrt{5}} \arctan \left(\sqrt{2(3 + \sqrt{5})} \sinh(x) \right) + 10 \sinh(x) \right)$$

input

```
Integrate[Sinh[x]*Tanh[5*x],x]
```

output

```
(-2*ArcTan[Sinh[x]] - Sqrt[2*(3 + Sqrt[5])]*ArcTan[2*Sqrt[2/(3 + Sqrt[5])]]*Sinh[x] - Sqrt[6 - 2*Sqrt[5]]*ArcTan[Sqrt[2*(3 + Sqrt[5])]]*Sinh[x] + 10*Sinh[x])/10
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 25, 4878, 25, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \tanh(5x) dx \\ \downarrow \text{3042} \\ \int -\sin(ix) \tan(5ix) dx \\ \downarrow \text{25} \\ -\int \sin(ix) \tan(5ix) dx$$

$$\begin{aligned}
& \downarrow 4878 \\
& - \int - \frac{\sinh^2(x) (16 \sinh^4(x) + 20 \sinh^2(x) + 5)}{16 \sinh^6(x) + 28 \sinh^4(x) + 13 \sinh^2(x) + 1} d \sinh(x) \\
& \downarrow 25 \\
& \int \frac{\sinh^2(x) (16 \sinh^4(x) + 20 \sinh^2(x) + 5)}{16 \sinh^6(x) + 28 \sinh^4(x) + 13 \sinh^2(x) + 1} d \sinh(x) \\
& \downarrow 2460 \\
& \int \left(-\frac{1}{5 (\sinh^2(x) + 1)} - \frac{4(6 \sinh^2(x) + 1)}{5 (16 \sinh^4(x) + 12 \sinh^2(x) + 1)} + 1 \right) d \sinh(x) \\
& \downarrow 2009 \\
& -\frac{1}{5} \arctan(\sinh(x)) - \frac{1}{5} \sqrt{\frac{1}{2} (3 + \sqrt{5})} \arctan \left(2 \sqrt{\frac{2}{3 + \sqrt{5}}} \sinh(x) \right) - \\
& \quad \frac{1}{5} \sqrt{\frac{1}{2} (3 - \sqrt{5})} \arctan \left(\sqrt{2 (3 + \sqrt{5})} \sinh(x) \right) + \sinh(x)
\end{aligned}$$

input

```
Int [Sinh [x] *Tanh [5*x] , x]
```

output

```
-1/5*ArcTan [Sinh [x]] - (Sqrt [(3 + Sqrt [5])/2]*ArcTan [2*Sqrt [2/(3 + Sqrt [5])]]*Sinh [x])/5 - (Sqrt [(3 - Sqrt [5])/2]*ArcTan [Sqrt [2*(3 + Sqrt [5])]]*Sinh [x])/5 + Sinh [x]
```

Defintions of rubi rules used

rule 25

```
Int [-(Fx_), x_Symbol] :> Simp [Identity [-1] Int [Fx, x] , x]
```

rule 2009

```
Int [u_, x_Symbol] :> Simp [IntSum [u, x] , x] /; SumQ [u]
```


rule 2460

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]},
Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Q
x, x]]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] &&
!TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4878

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d,
u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeF
actors[Sin[v], x], u/Cos[v], x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

method	result
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i \ln(e^x - i)}{5} - \frac{i \ln(e^x + i)}{5} + \left(\sum_{R=\text{RootOf}(10000_Z^4+300_Z^2+1)} _R \ln(-10_R e^x + e^{2x} - 1) \right)$

input

```
int(sinh(x)*tanh(5*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)-1/2*exp(-x)+1/5*I*ln(exp(x)-I)-1/5*I*ln(exp(x)+I)+sum(_R*ln(-10
*_R*exp(x)+exp(2*x)-1),_R=RootOf(10000*_Z^4+300*_Z^2+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.93

$$\int \sinh(x) \tanh(5x) dx =$$

$$\frac{((\sqrt{5} - 1) \cosh(x) + (\sqrt{5} - 1) \sinh(x)) \arctan\left(\frac{1}{2}(\sqrt{5} + 1) \cosh(x) + \frac{1}{2}(\sqrt{5} + 1) \sinh(x)\right) + ((\sqrt{5} - 1) \cosh(x) + (\sqrt{5} - 1) \sinh(x)) \arctan\left(\frac{1}{2}(\sqrt{5} - 1) \cosh(x) + \frac{1}{2}(\sqrt{5} - 1) \sinh(x)\right) - ((\sqrt{5} + 1) \cosh(x) + (\sqrt{5} + 1) \sinh(x)) \arctan\left(\frac{1}{2}(\sqrt{5} - 1) \cosh(x) + \frac{1}{2}(\sqrt{5} - 1) \sinh(x)\right) - ((\sqrt{5} + 1) \cosh(x) + (\sqrt{5} + 1) \sinh(x)) \arctan\left(\frac{1}{2}(\sqrt{5} + 1) \cosh(x) + \frac{1}{2}(\sqrt{5} + 1) \sinh(x)\right) - 5 \cosh(x)^2 - 10 \cosh(x) \sinh(x) - 5 \sinh(x)^2 + 5}{(\cosh(x) - \sinh(x))(\cosh(x) + \sinh(x))}}$$

input `integrate(sinh(x)*tanh(5*x),x, algorithm="fricas")`

output `-1/10*(((sqrt(5) - 1)*cosh(x) + (sqrt(5) - 1)*sinh(x))*arctan(1/2*(sqrt(5) + 1)*cosh(x) + 1/2*(sqrt(5) + 1)*sinh(x)) + ((sqrt(5) + 1)*cosh(x) + (sqrt(5) + 1)*sinh(x))*arctan(1/2*(sqrt(5) - 1)*cosh(x) + 1/2*(sqrt(5) - 1)*sinh(x)) - ((sqrt(5) - 1)*cosh(x) + (sqrt(5) - 1)*sinh(x))*arctan(-1/2*((sqrt(5) + 1)*cosh(x)^2 + 2*(sqrt(5) + 1)*cosh(x)*sinh(x) + (sqrt(5) + 1)*sinh(x)^2 - 2)/(cosh(x) - sinh(x))) - ((sqrt(5) + 1)*cosh(x) + (sqrt(5) + 1)*sinh(x))*arctan(-1/2*((sqrt(5) - 1)*cosh(x)^2 + 2*(sqrt(5) - 1)*cosh(x)*sinh(x) + (sqrt(5) - 1)*sinh(x)^2 + 2)/(cosh(x) - sinh(x))) + 4*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)) - 5*cosh(x)^2 - 10*cosh(x)*sinh(x) - 5*sinh(x)^2 + 5)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \sinh(x) \tanh(5x) dx = \int \sinh(x) \tanh(5x) dx$$

input `integrate(sinh(x)*tanh(5*x),x)`

output `Integral(sinh(x)*tanh(5*x), x)`

Maxima [F]

$$\int \sinh(x) \tanh(5x) dx = \int \sinh(x) \tanh(5x) dx$$

input `integrate(sinh(x)*tanh(5*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) - 1)*e^(-x) - 2/5*arctan(e^x) - 1/2*integrate(2/5*(3*e^(7*x) - e^(5*x) - e^(3*x) + 3*e^x)/(e^(8*x) - e^(6*x) + e^(4*x) - e^(2*x) + 1), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.96

$$\begin{aligned} \int \sinh(x) \tanh(5x) dx &= -\frac{1}{10} \pi - \frac{1}{10} \left(\sqrt{5} + 1 \right) \arctan \left(-\frac{2(e^{-x} - e^x)}{\sqrt{5} + 1} \right) \\ &\quad - \frac{1}{10} \left(\sqrt{5} - 1 \right) \arctan \left(-\frac{2(e^{-x} - e^x)}{\sqrt{5} - 1} \right) \\ &\quad - \frac{1}{5} \arctan \left(\frac{1}{2} (e^{2x} - 1) e^{-x} \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(sinh(x)*tanh(5*x),x, algorithm="giac")`

output `-1/10*pi - 1/10*(sqrt(5) + 1)*arctan(-2*(e^(-x) - e^x)/(sqrt(5) + 1)) - 1/10*(sqrt(5) - 1)*arctan(-2*(e^(-x) - e^x)/(sqrt(5) - 1)) - 1/5*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 2.22 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.98

$$\int \sinh(x) \tanh(5x) dx = \frac{e^x}{2} - \frac{2 \operatorname{atan}(e^x)}{5} - \frac{e^{-x}}{2} - 2 \operatorname{atan}\left(\frac{e^{-x}(e^{2x}-1)}{10\sqrt{\frac{3}{200}-\frac{\sqrt{5}}{200}}}\right) \sqrt{\frac{3}{200}-\frac{\sqrt{5}}{200}} \\ - 2 \operatorname{atan}\left(\frac{e^{-x}(e^{2x}-1)}{10\sqrt{\frac{\sqrt{5}}{200}+\frac{3}{200}}}\right) \sqrt{\frac{\sqrt{5}}{200}+\frac{3}{200}}$$

input `int(tanh(5*x)*sinh(x),x)`output `exp(x)/2 - (2*atan(exp(x)))/5 - exp(-x)/2 - 2*atan((exp(-x)*(exp(2*x) - 1))/(10*(3/200 - 5^(1/2)/200)^(1/2)))*(3/200 - 5^(1/2)/200)^(1/2) - 2*atan((exp(-x)*(exp(2*x) - 1))/(10*(5^(1/2)/200 + 3/200)^(1/2)))*(5^(1/2)/200 + 3/200)^(1/2)`**Reduce [F]**

$$\int \sinh(x) \tanh(5x) dx = \frac{e^{2x} - 2e^x \left(\int \frac{e^x}{e^{10x}+1} dx \right) + 2e^x \left(\int \frac{1}{e^{11x}+e^x} dx \right) + 1}{2e^x}$$

input `int(sinh(x)*tanh(5*x),x)`output `(e**(2*x) - 2*e**x*int(e**x/(e**(10*x) + 1),x) + 2*e**x*int(1/(e**(11*x) + e**x),x) + 1)/(2*e**x)`

3.30 $\int \sinh(x) \tanh(6x) dx$

Optimal result	276
Mathematica [A] (verified)	276
Rubi [A] (verified)	277
Maple [C] (verified)	279
Fricas [B] (verification not implemented)	279
Sympy [F]	280
Maxima [F]	280
Giac [A] (verification not implemented)	281
Mupad [B] (verification not implemented)	281
Reduce [B] (verification not implemented)	282

Optimal result

Integrand size = 7, antiderivative size = 87

$$\int \sinh(x) \tanh(6x) dx = -\frac{\arctan(\sqrt{2} \sinh(x))}{3\sqrt{2}} - \frac{\arctan(2\sqrt{2 - \sqrt{3}} \sinh(x))}{6\sqrt{2 - \sqrt{3}}} - \frac{\arctan(2\sqrt{2 + \sqrt{3}} \sinh(x))}{6\sqrt{2 + \sqrt{3}}} + \sinh(x)$$

output

```
-1/6*arctan(sinh(x)*2^(1/2))*2^(1/2)-1/6*arctan(2*(1/2*6^(1/2)-1/2*2^(1/2))
)*sinh(x))/(1/2*6^(1/2)-1/2*2^(1/2))-1/6*arctan(2*(1/2*6^(1/2)+1/2*2^(1/2)
)*sinh(x))/(1/2*6^(1/2)+1/2*2^(1/2))+sinh(x)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00

$$\int \sinh(x) \tanh(6x) dx = -\frac{\arctan(\sqrt{2} \sinh(x))}{3\sqrt{2}} - \frac{1}{6} \sqrt{2 - \sqrt{3}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2 - \sqrt{3}}}\right) - \frac{1}{6} \sqrt{2 + \sqrt{3}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2 + \sqrt{3}}}\right) + \sinh(x)$$

input `Integrate[Sinh[x]*Tanh[6*x],x]`

output `-1/3*ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2] - (Sqrt[2 - Sqrt[3]]*ArcTan[(2*Sinh[x])
]/Sqrt[2 - Sqrt[3]])/6 - (Sqrt[2 + Sqrt[3]]*ArcTan[(2*Sinh[x])/Sqrt[2 +
Sqrt[3]]])/6 + Sinh[x]`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 25, 4878, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \tanh(6x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\sin(ix) \tan(6ix) dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sin(ix) \tan(6ix) dx \\
 & \quad \downarrow \text{4878} \\
 & - \int -\frac{2 \sinh^2(x) (16 \sinh^4(x) + 16 \sinh^2(x) + 3)}{32 \sinh^6(x) + 48 \sinh^4(x) + 18 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & 2 \int \frac{\sinh^2(x) (16 \sinh^4(x) + 16 \sinh^2(x) + 3)}{32 \sinh^6(x) + 48 \sinh^4(x) + 18 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{2460} \\
 & 2 \int \left(\frac{-8 \sinh^2(x) - 1}{3 (16 \sinh^4(x) + 16 \sinh^2(x) + 1)} - \frac{1}{6 (2 \sinh^2(x) + 1)} + \frac{1}{2} \right) d \sinh(x) \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$2 \left(-\frac{\arctan(\sqrt{2} \sinh(x))}{6\sqrt{2}} - \frac{1}{12} \sqrt{2-\sqrt{3}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2-\sqrt{3}}}\right) - \frac{1}{12} \sqrt{2+\sqrt{3}} \arctan\left(\frac{2 \sinh(x)}{\sqrt{2+\sqrt{3}}}\right) + \frac{\sinh(x)}{2} \right)$$

input `Int[Sinh[x]*Tanh[6*x],x]`

output `2*(-1/6*ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2] - (Sqrt[2 - Sqrt[3]]*ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[3]]])/12 - (Sqrt[2 + Sqrt[3]]*ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[3]]])/12 + Sinh[x]/2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.09 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.97

method	result
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x - 1)}{12} - \frac{i\sqrt{2} \ln(e^{2x} + i\sqrt{2}e^x - 1)}{12} + \left(\sum_{_R=\text{RootOf}(20736_Z^4+576_Z^2+1)} _R \ln(-12 \dots) \right)$

input `int(sinh(x)*tanh(6*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/12*I*2^(1/2)*ln(exp(2*x)-I*2^(1/2)*exp(x)-1)-1/12*I*2^(1/2)*ln(exp(2*x)+I*2^(1/2)*exp(x)-1)+sum(_R*ln(-12*_R*exp(x)+exp(2*x)-1),_R=RootOf(20736*_Z^4+576*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 375 vs. 2(63) = 126.

Time = 0.13 (sec) , antiderivative size = 375, normalized size of antiderivative = 4.31

$$\int \sinh(x) \tanh(6x) dx = \text{Too large to display}$$

input `integrate(sinh(x)*tanh(6*x),x, algorithm="fricas")`

output

```
1/6*(sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x))*arctan(((sqrt(3) - 2)*cosh(x)^3
+ 3*(sqrt(3) - 2)*cosh(x)*sinh(x)^2 + (sqrt(3) - 2)*sinh(x)^3 - (sqrt(3)
- 1)*cosh(x) + (3*(sqrt(3) - 2)*cosh(x)^2 - sqrt(3) + 1)*sinh(x))*sqrt(sqrt(3) + 2))
+ sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x))*arctan(((sqrt(3) - 2)*cosh(x) + (sqrt(3) - 2)*sinh(x))*sqrt(sqrt(3) + 2))
- sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x))*arctan(((sqrt(3) + 2)*cosh(x)^3 + 3*(sqrt(3) + 2)*cosh(x)
)*sinh(x)^2 + (sqrt(3) + 2)*sinh(x)^3 - (sqrt(3) + 1)*cosh(x) + (3*(sqrt(3)
+ 2)*cosh(x)^2 - sqrt(3) - 1)*sinh(x))*sqrt(-sqrt(3) + 2)) - sqrt(-sqrt(3)
+ 2)*(cosh(x) + sinh(x))*arctan(((sqrt(3) + 2)*cosh(x) + (sqrt(3) + 2)*sinh(x))*sqrt(-sqrt(3) + 2))
- (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)
+ (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x)
+ sqrt(2)*sinh(x)^2 + sqrt(2))/(cosh(x) - sinh(x))) + 3*cosh(x)^2 + 6*cosh(x)*sinh(x)
+ 3*sinh(x)^2 - 3)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \sinh(x) \tanh(6x) dx = \int \sinh(x) \tanh(6x) dx$$

input

```
integrate(sinh(x)*tanh(6*x),x)
```

output

```
Integral(sinh(x)*tanh(6*x), x)
```

Maxima [F]

$$\int \sinh(x) \tanh(6x) dx = \int \sinh(x) \tanh(6x) dx$$

input

```
integrate(sinh(x)*tanh(6*x),x, algorithm="maxima")
```

output

```
1/2*(e^(2*x) - 1)*e^(-x) - 1/6*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x))
- 1/6*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/2*integrate(2/
3*(2*e^(7*x) - e^(5*x) - e^(3*x) + 2*e^x)/(e^(8*x) - e^(4*x) + 1), x)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.15

$$\int \sinh(x) \tanh(6x) dx = -\frac{1}{12} (\sqrt{6} + \sqrt{2}) \arctan \left(-\frac{2(e^{-x} - e^x)}{\sqrt{6} + \sqrt{2}} \right) - \frac{1}{12} (\sqrt{6} - \sqrt{2}) \arctan \left(-\frac{2(e^{-x} - e^x)}{\sqrt{6} - \sqrt{2}} \right) - \frac{1}{12} \sqrt{2} \left(\pi + 2 \arctan \left(\frac{1}{2} \sqrt{2} (e^{2x} - 1) e^{-x} \right) \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(sinh(x)*tanh(6*x),x, algorithm="giac")`output `-1/12*(sqrt(6) + sqrt(2))*arctan(-2*(e^(-x) - e^x)/(sqrt(6) + sqrt(2))) - 1/12*(sqrt(6) - sqrt(2))*arctan(-2*(e^(-x) - e^x)/(sqrt(6) - sqrt(2))) - 1/12*sqrt(2)*(pi + 2*arctan(1/2*sqrt(2)*(e^(2*x) - 1)*e^(-x))) - 1/2*e^(-x) + 1/2*e^x`**Mupad [B] (verification not implemented)**

Time = 2.23 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.13

$$\int \sinh(x) \tanh(6x) dx = \frac{e^x}{2} - \frac{e^{-x}}{2} - \frac{\sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} e^{-x} (e^{2x} - 1)}{2} \right)}{6} - 2 \operatorname{atan} \left(\frac{e^{-x} (e^{2x} - 1)}{12 \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}}} \right) \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} - 2 \operatorname{atan} \left(\frac{e^{-x} (e^{2x} - 1)}{12 \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}}} \right) \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}}$$

input `int(tanh(6*x)*sinh(x),x)`

output

$$\begin{aligned} & \exp(x)/2 - \exp(-x)/2 - (2^{(1/2)} * \operatorname{atan}((2^{(1/2)} * \exp(-x) * (\exp(2*x) - 1))/2)) / \\ & 6 - 2 * \operatorname{atan}((\exp(-x) * (\exp(2*x) - 1)) / (12 * (1/72 - 3^{(1/2)} / 144)^{(1/2)})) * (1/72 \\ & - 3^{(1/2)} / 144)^{(1/2)} - 2 * \operatorname{atan}((\exp(-x) * (\exp(2*x) - 1)) / (12 * (3^{(1/2)} / 144 + \\ & 1/72)^{(1/2)})) * (3^{(1/2)} / 144 + 1/72)^{(1/2)} \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 254, normalized size of antiderivative = 2.92

$$\int \sinh(x) \tanh(6x) dx$$

$$= \frac{-2e^x \sqrt{-\sqrt{3} + 2} \operatorname{atan}\left(\frac{4e^x - \sqrt{6} - \sqrt{2}}{2\sqrt{-\sqrt{3} + 2}}\right) - 2e^x \sqrt{-\sqrt{3} + 2} \operatorname{atan}\left(\frac{4e^x + \sqrt{6} + \sqrt{2}}{2\sqrt{-\sqrt{3} + 2}}\right) - 2e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - 2e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{1}$$

input

`int(sinh(x)*tanh(6*x),x)`

output

$$\begin{aligned} & (-2 * e^{**x} * \sqrt{-\sqrt{3} + 2} * \operatorname{atan}((4 * e^{**x} - \sqrt{6} - \sqrt{2}) / (2 * \sqrt{-\sqrt{3} + 2}))) - 2 * e^{**x} * \sqrt{-\sqrt{3} + 2} * \operatorname{atan}((4 * e^{**x} + \sqrt{6} + \sqrt{2}) / (2 * \sqrt{-\sqrt{3} + 2}))) \\ & - 2 * e^{**x} * \sqrt{2} * \operatorname{atan}((2 * e^{**x} - \sqrt{2}) / \sqrt{2}) - 2 * e^{**x} * \sqrt{2} * \operatorname{atan}((2 * e^{**x} + \sqrt{2}) / \sqrt{2}) + e^{**x} * \sqrt{6} * \operatorname{atan}((2 * \sqrt{-\sqrt{3} + 2} - 4 * e^{**x}) / (\sqrt{6} + \sqrt{2}))) \\ & + e^{**x} * \sqrt{6} * \operatorname{atan}((2 * \sqrt{-\sqrt{3} + 2} + 4 * e^{**x}) / (\sqrt{6} + \sqrt{2}))) - e^{**x} * \sqrt{6} * \operatorname{atan}((2 * \sqrt{-\sqrt{3} + 2} - 4 * e^{**x}) / (\sqrt{6} + \sqrt{2}))) \\ & - e^{**x} * \sqrt{6} * \operatorname{atan}((2 * \sqrt{-\sqrt{3} + 2} + 4 * e^{**x}) / (\sqrt{6} + \sqrt{2}))) + 6 * e^{**x} * (2 * x - 6) / (12 * e^{**x}) \end{aligned}$$

3.31 $\int \sinh(x) \tanh(nx) dx$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [F]	285
Fricas [F]	285
Sympy [F]	286
Maxima [F]	286
Giac [F]	286
Mupad [F(-1)]	287
Reduce [F]	287

Optimal result

Integrand size = 7, antiderivative size = 67

$$\int \sinh(x) \tanh(nx) dx = \cosh(x) - e^{-x} \operatorname{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, -e^{2nx} \right) - e^x \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -e^{2nx} \right)$$

output

`cosh(x)-hypergeom([1, -1/2/n],[1-1/2/n],-exp(2*n*x))/exp(x)-exp(x)*hypergeom([1, 1/2/n],[1+1/2/n],-exp(2*n*x))`

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.15

$$\int \sinh(x) \tanh(nx) dx = \frac{1}{2} \left(e^{-x} + e^x - 2e^{-x} \operatorname{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, -e^{2nx} \right) - 2e^x \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2n}, 1 + \frac{1}{2n}, -e^{2nx} \right) \right)$$

input

`Integrate[Sinh[x]*Tanh[n*x],x]`

output

```
(E^(-x) + E^x - (2*Hypergeometric2F1[1, -1/2*1/n, 1 - 1/(2*n), -E^(2*n*x)]
)/E^x - 2*E^x*Hypergeometric2F1[1, 1/(2*n), 1 + 1/(2*n), -E^(2*n*x)])/2
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.21, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6135, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \tanh(nx) dx$$

$$\downarrow 6135$$

$$\int \left(\frac{e^{-x}}{e^{2nx} + 1} - \frac{e^x}{e^{2nx} + 1} - \frac{e^{-x}}{2} + \frac{e^x}{2} \right) dx$$

$$\downarrow 2009$$

$$-e^{-x} \text{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, -e^{2nx} \right) -$$

$$e^x \text{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), -e^{2nx} \right) + \frac{e^{-x}}{2} + \frac{e^x}{2}$$

input

```
Int [Sinh [x] *Tanh [n*x] , x]
```

output

```
1/(2*E^x) + E^x/2 - Hypergeometric2F1[1, -1/2*1/n, 1 - 1/(2*n), -E^(2*n*x)]
]/E^x - E^x*Hypergeometric2F1[1, 1/(2*n), (2 + n^(-1))/2, -E^(2*n*x)]
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6135 `Int[Sinh[(a_.) + (b_.)*(x_)]*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Int[-E^(-a + b*x)/2 + E^(a + b*x)/2 + 1/(E^(a + b*x)*(1 + E^(2*(c + d*x)))) - E^(a + b*x)/(1 + E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \sinh(x) \tanh(nx) dx$$

input `int(sinh(x)*tanh(n*x),x)`

output `int(sinh(x)*tanh(n*x),x)`

Fricas [F]

$$\int \sinh(x) \tanh(nx) dx = \int \sinh(x) \tanh(nx) dx$$

input `integrate(sinh(x)*tanh(n*x),x, algorithm="fricas")`

output `integral(sinh(x)*tanh(n*x), x)`

Sympy [F]

$$\int \sinh(x) \tanh(nx) dx = \int \sinh(x) \tanh(nx) dx$$

input `integrate(sinh(x)*tanh(n*x),x)`

output `Integral(sinh(x)*tanh(n*x), x)`

Maxima [F]

$$\int \sinh(x) \tanh(nx) dx = \int \sinh(x) \tanh(nx) dx$$

input `integrate(sinh(x)*tanh(n*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) + 1)*e^(-x) - 1/2*integrate(2*(e^(2*x) - 1)/(e^(2*n*x + x) + e^x), x)`

Giac [F]

$$\int \sinh(x) \tanh(nx) dx = \int \sinh(x) \tanh(nx) dx$$

input `integrate(sinh(x)*tanh(n*x),x, algorithm="giac")`

output `integrate(sinh(x)*tanh(n*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \sinh(x) \tanh(nx) dx = \int \tanh(nx) \sinh(x) dx$$

input `int(tanh(n*x)*sinh(x),x)`output `int(tanh(n*x)*sinh(x),x)`**Reduce [F]**

$$\int \sinh(x) \tanh(nx) dx = \frac{e^{2x} - 2e^x \left(\int \frac{e^x}{e^{2nx}+1} dx \right) + 2e^x \left(\int \frac{1}{e^{2nx+x}+e^x} dx \right) + 1}{2e^x}$$

input `int(sinh(x)*tanh(n*x),x)`output `(e**(2*x) - 2*e**x*int(e**x/(e**(2*n*x) + 1),x) + 2*e**x*int(1/(e**(2*n*x + x) + e**x),x) + 1)/(2*e**x)`

3.32 $\int \sinh(a + bx) \tanh(c + bx) dx$

Optimal result	288
Mathematica [B] (verified)	288
Rubi [A] (verified)	289
Maple [C] (verified)	290
Fricas [B] (verification not implemented)	291
Sympy [F]	291
Maxima [A] (verification not implemented)	292
Giac [A] (verification not implemented)	292
Mupad [B] (verification not implemented)	293
Reduce [B] (verification not implemented)	293

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \sinh(a + bx) \tanh(c + bx) dx = -\frac{\arctan(\sinh(c + bx)) \cosh(a - c)}{b} + \frac{\sinh(a + bx)}{b}$$

output `-arctan(sinh(b*x+c))*cosh(a-c)/b+sinh(b*x+a)/b`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 86 vs. $2(29) = 58$.

Time = 0.05 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.97

$$\begin{aligned} & \int \sinh(a + bx) \tanh(c + bx) dx \\ &= -\frac{2 \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right)\right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a - c)}{b} \\ & \quad + \frac{\cosh(bx) \sinh(a)}{b} + \frac{\cosh(a) \sinh(bx)}{b} \end{aligned}$$

input `Integrate[Sinh[a + b*x]*Tanh[c + b*x],x]`

output

```
(-2*ArcTan[((Cosh[c] - Sinh[c])*(Cosh[(b*x)/2]*Sinh[c] + Cosh[c]*Sinh[(b*x)/2]))/(Cosh[c]*Cosh[(b*x)/2] - Cosh[(b*x)/2]*Sinh[c])]*Cosh[a - c])/b + (Cosh[b*x]*Sinh[a])/b + (Cosh[a]*Sinh[b*x])/b
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {6154, 3042, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \tanh(bx + c) dx$$

$$\downarrow 6154$$

$$\int \cosh(a + bx) dx - \cosh(a - c) \int \operatorname{sech}(c + bx) dx$$

$$\downarrow 3042$$

$$\int \sin\left(ia + ibx + \frac{\pi}{2}\right) dx - \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx$$

$$\downarrow 3117$$

$$\frac{\sinh(a + bx)}{b} - \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx$$

$$\downarrow 4257$$

$$\frac{\sinh(a + bx)}{b} - \frac{\cosh(a - c) \arctan(\sinh(bx + c))}{b}$$

input

```
Int[Sinh[a + b*x]*Tanh[c + b*x],x]
```

output

```
-((ArcTan[Sinh[c + b*x])*Cosh[a - c])/b) + Sinh[a + b*x]/b
```

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6154 `Int[Sinh[v_]*Tanh[w_]^(n_.), x_Symbol] := Int[Cosh[v]*Tanh[w]^(n - 1), x] - Simp[Cosh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 167, normalized size of antiderivative = 5.76

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a} + ie^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{i \ln(e^{bx+a} + ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(sinh(b*x+a)*tanh(b*x+c),x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+a)-1/2/b*exp(-b*x-a)+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*a)+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*c)-1/2*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*a)-1/2*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. $2(29) = 58$.

Time = 0.11 (sec) , antiderivative size = 327, normalized size of antiderivative = 11.28

$$\int \sinh(a + bx) \tanh(c + bx) dx$$

$$= \frac{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c)^2 \cosh(-a + c) \sinh(-a + c) + \cosh(bx + c)^2 \sinh(-a + c)^2}{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c)^2 \cosh(-a + c) \sinh(-a + c) + \cosh(bx + c)^2 \sinh(-a + c)^2}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c),x, algorithm="fricas")`

output `1/2*(cosh(b*x + c)^2*cosh(-a + c)^2 - 2*cosh(b*x + c)^2*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)^2*sinh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^2 + 2*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c))*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 1)/(b*cosh(b*x + c)*cosh(-a + c) - b*cosh(b*x + c)*sinh(-a + c) + (b*cosh(-a + c) - b*sinh(-a + c))*sinh(b*x + c))`

Sympy [F]

$$\int \sinh(a + bx) \tanh(c + bx) dx = \int \sinh(a + bx) \tanh(bx + c) dx$$

input `integrate(sinh(b*x+a)*tanh(b*x+c),x)`

output `Integral(sinh(a + b*x)*tanh(b*x + c), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.97

$$\int \sinh(a + bx) \tanh(c + bx) dx = \frac{(e^{2a} + e^{2c}) \arctan(e^{-bx-c}) e^{(-a-c)}}{b} + \frac{e^{(bx+a)}}{2b} - \frac{e^{(-bx-a)}}{2b}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c),x, algorithm="maxima")`output `(e^(2*a) + e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b + 1/2*e^(b*x + a)/b - 1/2*e^(-b*x - a)/b`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.69

$$\int \sinh(a + bx) \tanh(c + bx) dx = -\frac{2(e^{2a} + e^{2c}) \arctan(e^{(bx+c)}) e^{(-a-c)} - e^{(bx+a)} + e^{(-bx-a)}}{2b}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c),x, algorithm="giac")`output `-1/2*(2*(e^(2*a) + e^(2*c))*arctan(e^(b*x + c))*e^(-a - c) - e^(b*x + a) + e^(-b*x - a))/b`

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 133, normalized size of antiderivative = 4.59

$$\int \sinh(a + bx) \tanh(c + bx) dx$$

$$= \frac{e^{a+bx}}{2b} - \frac{e^{-a-bx}}{2b}$$

$$- \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2 + e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{b^2}}$$

input `int(sinh(a + b*x)*tanh(c + b*x),x)`output `exp(a + b*x)/(2*b) - exp(- a - b*x)/(2*b) - (atan((exp(-a)*exp(2*c)*exp(b*x))*((b^2)^(1/2) + exp(2*a)*exp(-2*c)*(b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c)*(2*exp(2*a)*exp(-2*c) + exp(4*a)*exp(-4*c) + 1))^(1/2)))*(exp(2*c - 2*a)*(2*exp(2*a - 2*c) + exp(4*a - 4*c) + 1))^(1/2))/(b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.41

$$\int \sinh(a + bx) \tanh(c + bx) dx$$

$$= \frac{-2e^{bx+2a} \operatorname{atan}(e^{bx+c}) - 2e^{bx+2c} \operatorname{atan}(e^{bx+c}) + e^{2bx+2a+c} - e^c}{2e^{bx+a+c}b}$$

input `int(sinh(b*x+a)*tanh(b*x+c),x)`output `(- 2*e**(2*a + b*x)*atan(e**(b*x + c)) - 2*e**(b*x + 2*c)*atan(e**(b*x + c)) + e**(2*a + 2*b*x + c) - e**c)/(2*e**(a + b*x + c)*b)`

3.33 $\int \sinh(a + bx) \tanh^2(c + bx) dx$

Optimal result	294
Mathematica [B] (verified)	294
Rubi [A] (verified)	295
Maple [C] (verified)	297
Fricas [B] (verification not implemented)	298
Sympy [F]	299
Maxima [B] (verification not implemented)	299
Giac [B] (verification not implemented)	299
Mupad [B] (verification not implemented)	300
Reduce [B] (verification not implemented)	301

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \sinh(a + bx) \tanh^2(c + bx) dx = \frac{\cosh(a + bx)}{b} + \frac{\cosh(a - c)\operatorname{sech}(c + bx)}{b} - \frac{\arctan(\sinh(c + bx)) \sinh(a - c)}{b}$$

output cosh(b*x+a)/b+cosh(a-c)*sech(b*x+c)/b-arctan(sinh(b*x+c))*sinh(a-c)/b

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 102 vs. 2(45) = 90.

Time = 0.08 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.27

$$\begin{aligned} & \int \sinh(a + bx) \tanh^2(c + bx) dx \\ &= \frac{\cosh(a) \cosh(bx)}{b} + \frac{\cosh(a - c)\operatorname{sech}(c + bx)}{b} \\ & \quad - \frac{2 \arctan \left(\frac{(\cosh(c) - \sinh(c)) \left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right) \right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)} \right) \sinh(a - c)}{b} \\ & \quad + \frac{\sinh(a) \sinh(bx)}{b} \end{aligned}$$

input `Integrate[Sinh[a + b*x]*Tanh[c + b*x]^2,x]`

output
$$\frac{(\text{Cosh}[a]*\text{Cosh}[b*x])/b + (\text{Cosh}[a - c]*\text{Sech}[c + b*x])/b - (2*\text{ArcTan}[\frac{(\text{Cosh}[c] - \text{Sinh}[c])*(\text{Cosh}[(b*x)/2]*\text{Sinh}[c] + \text{Cosh}[c]*\text{Sinh}[(b*x)/2])}{(\text{Cosh}[c]*\text{Cosh}[(b*x)/2] - \text{Cosh}[(b*x)/2]*\text{Sinh}[c])}]*\text{Sinh}[a - c])/b + (\text{Sinh}[a]*\text{Sinh}[b*x])/b}{b}$$

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {6154, 3042, 26, 3086, 24, 6157, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(a + bx) \tanh^2(bx + c) dx \\ & \quad \downarrow 6154 \\ & \int \cosh(a + bx) \tanh(c + bx) dx - \cosh(a - c) \int \text{sech}(c + bx) \tanh(c + bx) dx \\ & \quad \downarrow 3042 \\ & \int \cosh(a + bx) \tanh(c + bx) dx - \cosh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx \\ & \quad \downarrow 26 \\ & \int \cosh(a + bx) \tanh(c + bx) dx + i \cosh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx \\ & \quad \downarrow 3086 \\ & \int \cosh(a + bx) \tanh(c + bx) dx + \frac{\cosh(a - c) \int 1 d\text{sech}(c + bx)}{b} \\ & \quad \downarrow 24 \\ & \int \cosh(a + bx) \tanh(c + bx) dx + \frac{\cosh(a - c) \text{sech}(bx + c)}{b} \\ & \quad \downarrow 6157 \end{aligned}$$

$$\begin{aligned}
& -\sinh(a-c) \int \operatorname{sech}(c+bx) dx + \int \sinh(a+bx) dx + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{3042} \\
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \int -i \sin(ia+ibx) dx + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{26} \\
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx - i \int \sin(ia+ibx) dx + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{3118} \\
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} + \frac{\cosh(a+bx)}{b} \\
& \quad \downarrow \text{4257} \\
& -\frac{\sinh(a-c) \arctan(\sinh(bx+c))}{b} + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} + \frac{\cosh(a+bx)}{b}
\end{aligned}$$

input `Int[Sinh[a + b*x]*Tanh[c + b*x]^2,x]`

output `Cosh[a + b*x]/b + (Cosh[a - c]*Sech[c + b*x])/b - (ArcTan[Sinh[c + b*x]]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^(n - 1)/2], x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6154 `Int[Sinh[v_]*Tanh[w_]^(n_.), x_Symbol] := Int[Cosh[v]*Tanh[w]^(n - 1), x] - Simp[Cosh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

rule 6157 `Int[Cosh[v_]*Tanh[w_]^(n_.), x_Symbol] := Int[Sinh[v]*Tanh[w]^(n - 1), x] - Simp[Sinh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 205, normalized size of antiderivative = 4.56

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(e^{2a}+e^{2c})}{b(e^{2bx+2a+2c}+e^{2a})} + \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(sinh(b*x+a)*tanh(b*x+c)^2,x,method=_RETURNVERBOSE)`

output

```
1/2/b*exp(b*x+a)+1/2/b*exp(-b*x-a)+1/b*exp(b*x+a)*(exp(2*a)+exp(2*c))/(exp
(2*b*x+2*a+2*c)+exp(2*a))+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(
a)^2-1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(c)^2-1/2*I*ln(exp(b*x
+a)+I*exp(a-c))/b*exp(-a-c)*exp(a)^2+1/2*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp
(-a-c)*exp(c)^2
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 902 vs. $2(45) = 90$.

Time = 0.11 (sec) , antiderivative size = 902, normalized size of antiderivative = 20.04

$$\int \sinh(a + bx) \tanh^2(c + bx) dx = \text{Too large to display}$$

input

```
integrate(sinh(b*x+a)*tanh(b*x+c)^2,x, algorithm="fricas")
```

output

```
1/2*(cosh(b*x + c)^4*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sin
h(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^4 + 4*(cosh(b*x + c)*cosh(-a + c
)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a +
c)^2)*sinh(b*x + c)^3 + 3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + 3*(2*cosh
(b*x + c)^2*cosh(-a + c)^2 + (2*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + cosh
(-a + c)^2 - 2*(2*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c
) + 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^4 + 3*cosh(b*x + c)^2)*sinh(-a + c
)^2 - 2*((cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-
a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x
+ c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a +
c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 + cosh(b*x +
c))*sinh(-a + c)^2 + (cosh(-a + c)^2 - 1)*cosh(b*x + c) + (3*(cosh(-a + c)
^2 - 1)*cosh(b*x + c)^2 + (3*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + cosh(-a
+ c)^2 - 2*(3*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c) -
1)*sinh(b*x + c) - 2*(cosh(b*x + c)^3*cosh(-a + c) + cosh(b*x + c)*cosh(-
a + c))*sinh(-a + c))*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(2*cosh(b*
x + c)^3*cosh(-a + c)^2 + (2*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a +
c)^2 + 3*(cosh(-a + c)^2 + 1)*cosh(b*x + c) - 2*(2*cosh(b*x + c)^3*cosh(-a
+ c) + 3*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c) - 2*(cos
h(b*x + c)^4*cosh(-a + c) + 3*cosh(b*x + c)^2*cosh(-a + c))*sinh(-a + c...
```

Sympy [F]

$$\int \sinh(a + bx) \tanh^2(c + bx) dx = \int \sinh(a + bx) \tanh^2(bx + c) dx$$

input `integrate(sinh(b*x+a)*tanh(b*x+c)**2,x)`

output `Integral(sinh(a + b*x)*tanh(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(45) = 90$.

Time = 0.12 (sec) , antiderivative size = 105, normalized size of antiderivative = 2.33

$$\int \sinh(a + bx) \tanh^2(c + bx) dx = \frac{(e^{(2a)} - e^{(2c)}) \arctan(e^{(-bx-c)}) e^{(-a-c)}}{b} + \frac{e^{(-bx-a)}}{2b} + \frac{(3e^{(2a)} + 2e^{(2c)})e^{(-2bx-2a)} + e^{(2c)}}{2b(e^{(-bx-a+2c)} + e^{(-3bx-a)})}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c)^2,x, algorithm="maxima")`

output `(e^(2*a) - e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b + 1/2*e^(-b*x - a)/b + 1/2*((3*e^(2*a) + 2*e^(2*c))*e^(-2*b*x - 2*a) + e^(2*c))/(b*(e^(-b*x - a + 2*c) + e^(-3*b*x - a)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(45) = 90$.

Time = 0.11 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.16

$$\int \sinh(a + bx) \tanh^2(c + bx) dx = -\frac{2(e^{(2a)} - e^{(2c)}) \arctan(e^{(bx+c)}) e^{(-a-c)} - \frac{2e^{(2bx+4a)} + 3e^{(2bx+2a+2c)} + e^{(2a)}}{e^{(3bx+3a+2c)} + e^{(bx+3a)}} - e^{(bx+a)}}{2b}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c)^2,x, algorithm="giac")`

output
$$-1/2*(2*(e^{(2*a)} - e^{(2*c)})*\arctan(e^{(b*x + c)})*e^{(-a - c)} - (2*e^{(2*b*x + 4*a)} + 3*e^{(2*b*x + 2*a + 2*c)} + e^{(2*a)})/(e^{(3*b*x + 3*a + 2*c)} + e^{(b*x + 3*a)}) - e^{(b*x + a)})/b$$

Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 173, normalized size of antiderivative = 3.84

$$\begin{aligned} & \int \sinh(a + bx) \tanh^2(c + bx) dx \\ &= \frac{e^{a+bx}}{2b} + \frac{e^{-a-bx}}{2b} \\ &+ \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2 - e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{b^2}} \\ &+ \frac{e^{a+bx} (e^{2a-2c} + 1)}{b (e^{2a-2c} + e^{2a+2bx})} \end{aligned}$$

input `int(sinh(a + b*x)*tanh(c + b*x)^2,x)`

output
$$\exp(a + b*x)/(2*b) + \exp(-a - b*x)/(2*b) + (\operatorname{atan}((\exp(-a)*\exp(2*c)*\exp(b*x))*((b^2)^{(1/2)} - \exp(2*a)*\exp(-2*c)*(b^2)^{(1/2)}))/(b*(\exp(-2*a)*\exp(2*c)*(\exp(4*a)*\exp(-4*c) - 2*\exp(2*a)*\exp(-2*c) + 1))^{(1/2)}))*(\exp(2*c - 2*a)*(\exp(4*a - 4*c) - 2*\exp(2*a - 2*c) + 1))^{(1/2)})/(b^2)^{(1/2)} + (\exp(a + b*x) * (\exp(2*a - 2*c) + 1))/(b*(\exp(2*a - 2*c) + \exp(2*a + 2*b*x)))$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 152, normalized size of antiderivative = 3.38

$$\int \sinh(a + bx) \tanh^2(c + bx) dx$$

$$= \frac{-2e^{3bx+2a+2c} \operatorname{atan}(e^{bx+c}) + 2e^{3bx+4c} \operatorname{atan}(e^{bx+c}) - 2e^{bx+2a} \operatorname{atan}(e^{bx+c}) + 2e^{bx+2c} \operatorname{atan}(e^{bx+c}) + e^{4bx+2a+3c}}{2e^{bx+a+c} b (e^{2bx+2c} + 1)}$$

input `int(sinh(b*x+a)*tanh(b*x+c)^2,x)`output `(- 2*e**(2*a + 3*b*x + 2*c)*atan(e**(b*x + c)) + 2*e**(3*b*x + 4*c)*atan(e**(b*x + c)) - 2*e**(2*a + b*x)*atan(e**(b*x + c)) + 2*e**(b*x + 2*c)*atan(e**(b*x + c)) + e**(2*a + 4*b*x + 3*c) + 3*e**(2*a + 2*b*x + c) + 3*e**(2*b*x + 3*c) + e**c)/(2*e**(a + b*x + c)*b*(e**(2*b*x + 2*c) + 1))`

3.34 $\int \sinh(a + bx) \tanh^3(c + bx) dx$

Optimal result	302
Mathematica [A] (verified)	302
Rubi [A] (verified)	303
Maple [C] (verified)	307
Fricas [B] (verification not implemented)	307
Sympy [F]	308
Maxima [B] (verification not implemented)	309
Giac [A] (verification not implemented)	309
Mupad [F(-1)]	310
Reduce [B] (verification not implemented)	310

Optimal result

Integrand size = 15, antiderivative size = 72

$$\int \sinh(a + bx) \tanh^3(c + bx) dx = -\frac{3 \arctan(\sinh(c + bx)) \cosh(a - c)}{2b} + \frac{\operatorname{sech}(c + bx) \sinh(a - c)}{b} + \frac{\sinh(a + bx)}{b} + \frac{\cosh(a - c) \operatorname{sech}(c + bx) \tanh(c + bx)}{2b}$$

output

```
-3/2*arctan(sinh(b*x+c))*cosh(a-c)/b+sech(b*x+c)*sinh(a-c)/b+sinh(b*x+a)/b
+1/2*cosh(a-c)*sech(b*x+c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.97

$$\int \sinh(a + bx) \tanh^3(c + bx) dx = \frac{-12 \arctan\left(\sinh(c) + \cosh(c) \tanh\left(\frac{bx}{2}\right)\right) \cosh(a - c) + \operatorname{sech}^2(c + bx)(2 \sinh(a - 2c - bx) + 5 \sinh(a + c))}{4b}$$

input

```
Integrate[Sinh[a + b*x]*Tanh[c + b*x]^3,x]
```

output

```
(-12*ArcTan[Sinh[c] + Cosh[c]*Tanh[(b*x)/2]]*Cosh[a - c] + Sech[c + b*x]^2
*(2*Sinh[a - 2*c - b*x] + 5*Sinh[a + b*x] + Sinh[a + 2*c + 3*b*x]))/(4*b)
```

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.19, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {6154, 3042, 25, 3091, 3042, 4257, 6157, 3042, 26, 3086, 24, 6154, 3042, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \tanh^3(bx + c) dx \\
 & \quad \downarrow \text{6154} \\
 & \int \cosh(a + bx) \tanh^2(c + bx) dx - \cosh(a - c) \int \operatorname{sech}(c + bx) \tanh^2(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cosh(a + bx) \tanh^2(c + bx) dx - \cosh(a - c) \int -\sec(ic + ibx) \tan(ic + ibx)^2 dx \\
 & \quad \downarrow \text{25} \\
 & \int \cosh(a + bx) \tanh^2(c + bx) dx + \cosh(a - c) \int \sec(ic + ibx) \tan(ic + ibx)^2 dx \\
 & \quad \downarrow \text{3091} \\
 & \int \cosh(a + bx) \tanh^2(c + bx) dx + \cosh(a - c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{1}{2} \int \operatorname{sech}(c + bx) dx \right) \\
 & \quad \downarrow \text{3042} \\
 & \int \cosh(a + bx) \tanh^2(c + bx) dx + \cosh(a - c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{1}{2} \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow \text{4257}
 \end{aligned}$$

$$\begin{aligned}
& \int \cosh(a + bx) \tanh^2(c + bx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{6157} \\
& \int \sinh(a + bx) \tanh(c + bx) dx - \sinh(a - c) \int \operatorname{sech}(c + bx) \tanh(c + bx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& \int \sinh(a + bx) \tanh(c + bx) dx - \sinh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{26} \\
& \int \sinh(a + bx) \tanh(c + bx) dx + i \sinh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{3086} \\
& \int \sinh(a + bx) \tanh(c + bx) dx + \frac{\sinh(a - c) \int 1 d \operatorname{sech}(c + bx)}{b} + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{24} \\
& \int \sinh(a + bx) \tanh(c + bx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) + \frac{\sinh(a - c) \operatorname{sech}(bx + c)}{b} \\
& \quad \downarrow \text{6154} \\
& - \cosh(a - c) \int \operatorname{sech}(c + bx) dx + \int \cosh(a + bx) dx + \cosh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) + \frac{\sinh(a - c) \operatorname{sech}(bx + c)}{b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
& -\cosh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \int \sin\left(ia+ibx+\frac{\pi}{2}\right) dx + \cosh(a-c) \\
& c) \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{3117} \\
& -\cosh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \cosh(a-c) \\
& c) \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} + \\
& \quad \frac{\sinh(a+bx)}{b} \\
& \quad \downarrow \text{4257} \\
& -\frac{\cosh(a-c)\arctan(\sinh(bx+c))}{b} + \cosh(a-c) \\
& c) \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} + \\
& \quad \frac{\sinh(a+bx)}{b}
\end{aligned}$$

input `Int[Sinh[a + b*x]*Tanh[c + b*x]^3,x]`

output `-((ArcTan[Sinh[c + b*x]]*Cosh[a - c])/b) + (Sech[c + b*x]*Sinh[a - c])/b + Sinh[a + b*x]/b + Cosh[a - c]*(-1/2*ArcTan[Sinh[c + b*x]]/b + (Sech[c + b*x]*Tanh[c + b*x]))/(2*b)`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 3091 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

rule 3117 `Int[sin[Pi/2 + (c_) + (d_)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6154 `Int[Sinh[v_]*Tanh[w_]^(n_), x_Symbol] := Int[Cosh[v]*Tanh[w]^(n - 1), x] - Simp[Cosh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

rule 6157 `Int[Cosh[v_]*Tanh[w_]^(n_), x_Symbol] := Int[Sinh[v]*Tanh[w]^(n - 1), x] - Simp[Sinh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 240, normalized size of antiderivative = 3.33

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(3e^{2bx+4a+2c} - e^{2bx+2a+4c} + e^{4a} - 3e^{2a+2c})}{2b(e^{2bx+2a+2c} + e^{2a})^2} + \frac{3i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2a}}{4b} + \frac{3i \ln(e^{bx+a} - ie^{a-c})}{4b}$

input

```
int(sinh(b*x+a)*tanh(b*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
1/2/b*exp(b*x+a)-1/2/b*exp(-b*x-a)+1/2*exp(b*x+a)*(3*exp(2*b*x+4*a+2*c)-exp(2*b*x+2*a+4*c)+exp(4*a)-3*exp(2*a+2*c))/b/(exp(2*b*x+2*a+2*c)+exp(2*a))^2+3/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*a)+3/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*c)-3/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*a)-3/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1737 vs. 2(68) = 136.

Time = 0.12 (sec) , antiderivative size = 1737, normalized size of antiderivative = 24.12

$$\int \sinh(a + bx) \tanh^3(c + bx) dx = \text{Too large to display}$$

input

```
integrate(sinh(b*x+a)*tanh(b*x+c)^3,x, algorithm="fricas")
```

output

```

1/2*(cosh(b*x + c)^6*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^6 + 6*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^5 + (5*cosh(-a + c)^2 - 2)*cosh(b*x + c)^4 + (15*cosh(b*x + c)^2*cosh(-a + c)^2 + 5*(3*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + 5*cosh(-a + c)^2 - 10*(3*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c) - 2)*sinh(b*x + c)^4 + 4*(5*cosh(b*x + c)^3*cosh(-a + c)^2 + 5*(cosh(b*x + c)^3 + cosh(b*x + c))*sinh(-a + c)^2 + (5*cosh(-a + c)^2 - 2)*cosh(b*x + c) - 10*(cosh(b*x + c)^3*cosh(-a + c) + cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^3 + (2*cosh(-a + c)^2 - 5)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^4*cosh(-a + c)^2 + 6*(5*cosh(-a + c)^2 - 2)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^4 + 30*cosh(b*x + c)^2 + 2)*sinh(-a + c)^2 + 2*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^4*cosh(-a + c) + 30*cosh(b*x + c)^2*cosh(-a + c) + 2*cosh(-a + c))*sinh(-a + c) - 5)*sinh(b*x + c)^2 + (cosh(b*x + c)^6 + 5*cosh(b*x + c)^4 + 2*cosh(b*x + c)^2)*sinh(-a + c)^2 - 3*((cosh(-a + c)^2 + 1)*cosh(b*x + c)^5 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^5 - 5*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^4 + 2*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + 2*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 + 1)*sinh(-a + ...

```

Sympy [F]

$$\int \sinh(a + bx) \tanh^3(c + bx) dx = \int \sinh(a + bx) \tanh^3(bx + c) dx$$

input

```
integrate(sinh(b*x+a)*tanh(b*x+c)**3,x)
```

output

```
Integral(sinh(a + b*x)*tanh(b*x + c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(68) = 136$.

Time = 0.14 (sec) , antiderivative size = 149, normalized size of antiderivative = 2.07

$$\int \sinh(a + bx) \tanh^3(c + bx) dx$$

$$= \frac{3(e^{2a} + e^{2c}) \arctan(e^{-bx-c}) e^{(-a-c)} - \frac{e^{(-bx-a)}}{2b}}{\frac{2b}{(5e^{2a+2c} - e^{4c})e^{-2bx-2a} + (2e^{4a} - 3e^{2a+2c})e^{-4bx-4a} + e^{4c}} + \frac{2b}{2b(e^{-bx-a+4c} + 2e^{-3bx-a+2c} + e^{-5bx-a})}}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c)^3,x, algorithm="maxima")`

output `3/2*(e^(2*a) + e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b - 1/2*e^(-b*x - a)/b + 1/2*((5*e^(2*a + 2*c) - e^(4*c))*e^(-2*b*x - 2*a) + (2*e^(4*a) - 3*e^(2*a + 2*c))*e^(-4*b*x - 4*a) + e^(4*c))/(b*(e^(-b*x - a + 4*c) + 2*e^(-3*b*x - a + 2*c) + e^(-5*b*x - a)))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.67

$$\int \sinh(a + bx) \tanh^3(c + bx) dx =$$

$$\frac{3(e^{2a} + e^{2c}) \arctan(e^{(bx+c)}) e^{(-a-c)} - \frac{3e^{(3bx+5a+2c)} - e^{(3bx+3a+4c)} + e^{(bx+5a)} - 3e^{(bx+3a+2c)}}{(e^{(2bx+2a+2c)} + e^{(2a)})^2} - e^{(bx+a)} + e^{(-bx-a)}}{2b}$$

input `integrate(sinh(b*x+a)*tanh(b*x+c)^3,x, algorithm="giac")`

output `-1/2*(3*(e^(2*a) + e^(2*c))*arctan(e^(b*x + c))*e^(-a - c) - (3*e^(3*b*x + 5*a + 2*c) - e^(3*b*x + 3*a + 4*c) + e^(b*x + 5*a) - 3*e^(b*x + 3*a + 2*c)))/(e^(2*b*x + 2*a + 2*c) + e^(2*a))^2 - e^(b*x + a) + e^(-b*x - a))/b`

Mupad [F(-1)]

Timed out.

$$\int \sinh(a + bx) \tanh^3(c + bx) dx = \int \sinh(a + bx) \tanh(c + bx)^3 dx$$

input `int(sinh(a + b*x)*tanh(c + b*x)^3,x)`output `int(sinh(a + b*x)*tanh(c + b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.86

$$\int \sinh(a + bx) \tanh^3(c + bx) dx$$

$$= \frac{-6e^{bx+2a} \operatorname{atan}(e^{bx+c}) - 6e^{bx+2c} \operatorname{atan}(e^{bx+c}) - 2e^{bx+a+c} \cosh(bx+a) \tanh(bx+c) + 3e^{2bx+2a+c} - 2e^{bx+a+c}}{4e^{bx+a+c}b}$$

input `int(sinh(b*x+a)*tanh(b*x+c)^3,x)`output `(- 6*e**(2*a + b*x)*atan(e**(b*x + c)) - 6*e**(b*x + 2*c)*atan(e**(b*x + c)) - 2*e**(a + b*x + c)*cosh(a + b*x)*tanh(b*x + c) + 3*e**(2*a + 2*b*x + c) - 2*e**(a + b*x + c)*sinh(a + b*x)*tanh(b*x + c)**2 + 2*e**(a + b*x + c)*sinh(a + b*x) - 3*e**c)/(4*e**(a + b*x + c)*b)`

3.35 $\int \sinh(a + bx) \tanh(c + dx) dx$

Optimal result	311
Mathematica [A] (verified)	311
Rubi [A] (verified)	312
Maple [F]	313
Fricas [F]	313
Sympy [F]	314
Maxima [F]	314
Giac [F]	314
Mupad [F(-1)]	315
Reduce [F]	315

Optimal result

Integrand size = 13, antiderivative size = 100

$$\int \sinh(a + bx) \tanh(c + dx) dx$$

$$= \frac{\cosh(a + bx)}{b} - \frac{e^{-a-bx} \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right)}{b}$$

$$- \frac{e^{a+bx} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, -e^{2(c+dx)}\right)}{b}$$

output

$\cosh(b*x+a)/b - \exp(-b*x-a)*\operatorname{hypergeom}\left([1, -1/2*b/d], [1-1/2*b/d], -\exp(2*d*x+2*c)\right)/b - \exp(b*x+a)*\operatorname{hypergeom}\left([1, 1/2*b/d], [1+1/2*b/d], -\exp(2*d*x+2*c)\right)/b$

Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.03

$$\int \sinh(a + bx) \tanh(c + dx) dx$$

$$= \frac{e^{-a-bx} (1 + e^{2(a+bx)}) - 2 \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right) - 2e^{2(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, -e^{2(c+dx)}\right)}{2b}$$

input

`Integrate[Sinh[a + b*x]*Tanh[c + d*x], x]`

output

$$\frac{(E^{-a - bx} * (1 + E^{2(a + bx)}) - 2 * \text{Hypergeometric2F1}[1, -1/2*b/d, 1 - b/(2*d), -E^{2(c + dx)}]) - 2 * E^{2(a + bx)} * \text{Hypergeometric2F1}[1, b/(2*d), 1 + b/(2*d), -E^{2(c + dx)}])}{(2*b)}$$
Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.21, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6135, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \tanh(c + dx) dx$$

$$\downarrow \text{6135}$$

$$\int \left(\frac{e^{-a-bx}}{e^{2(c+dx)} + 1} - \frac{e^{a+bx}}{e^{2(c+dx)} + 1} - \frac{1}{2}e^{-a-bx} + \frac{1}{2}e^{a+bx} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{e^{-a-bx} \text{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \text{Hypergeometric2F1}\left(1, \frac{b}{2d}, \frac{b}{2d} + 1, -e^{2(c+dx)}\right)}{b} + \frac{e^{-a-bx}}{2b} + \frac{e^{a+bx}}{2b}$$

input

$$\text{Int}[\text{Sinh}[a + b*x]*\text{Tanh}[c + d*x], x]$$

output

$$\frac{E^{-a - bx}}{(2*b)} + \frac{E^{a + bx}}{(2*b)} - \frac{(E^{-a - bx} * \text{Hypergeometric2F1}[1, -1/2*b/d, 1 - b/(2*d), -E^{2(c + dx)}])}{b} - \frac{(E^{a + bx} * \text{Hypergeometric2F1}[1, b/(2*d), 1 + b/(2*d), -E^{2(c + dx)}])}{b}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6135 `Int[Sinh[(a_.) + (b_.)*(x_)]*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Int[-E^(-a + b*x))/2 + E^(a + b*x)/2 + 1/(E^(a + b*x)*(1 + E^(2*(c + d*x)))) - E^(a + b*x)/(1 + E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \sinh (bx + a) \tanh (dx + c) dx$$

input `int(sinh(b*x+a)*tanh(d*x+c),x)`

output `int(sinh(b*x+a)*tanh(d*x+c),x)`

Fricas [F]

$$\int \sinh (a + bx) \tanh (c + dx) dx = \int \sinh (bx + a) \tanh (dx + c) dx$$

input `integrate(sinh(b*x+a)*tanh(d*x+c),x, algorithm="fricas")`

output `integral(sinh(b*x + a)*tanh(d*x + c), x)`

Sympy [F]

$$\int \sinh(a + bx) \tanh(c + dx) dx = \int \sinh(a + bx) \tanh(c + dx) dx$$

input `integrate(sinh(b*x+a)*tanh(d*x+c),x)`

output `Integral(sinh(a + b*x)*tanh(c + d*x), x)`

Maxima [F]

$$\int \sinh(a + bx) \tanh(c + dx) dx = \int \sinh(bx + a) \tanh(dx + c) dx$$

input `integrate(sinh(b*x+a)*tanh(d*x+c),x, algorithm="maxima")`

output `1/2*(e^(2*b*x + 2*a) + 1)*e^(-b*x - a)/b - 1/2*integrate(2*(e^(2*b*x + 2*a) - 1)/(e^(b*x + 2*d*x + a + 2*c) + e^(b*x + a)), x)`

Giac [F]

$$\int \sinh(a + bx) \tanh(c + dx) dx = \int \sinh(bx + a) \tanh(dx + c) dx$$

input `integrate(sinh(b*x+a)*tanh(d*x+c),x, algorithm="giac")`

output `integrate(sinh(b*x + a)*tanh(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \sinh(a + bx) \tanh(c + dx) dx = \int \sinh(a + bx) \tanh(c + dx) dx$$

input `int(sinh(a + b*x)*tanh(c + d*x),x)`output `int(sinh(a + b*x)*tanh(c + d*x), x)`**Reduce [F]**

$$\int \sinh(a + bx) \tanh(c + dx) dx$$

$$= \frac{e^{2bx+2a} - 2e^{bx+2a} \left(\int \frac{e^{bx}}{e^{2dx+2c}+1} dx \right) b + 2e^{bx} \left(\int \frac{1}{e^{bx+2dx+2c}+e^{bx}} dx \right) b + 1}{2e^{bx+ab}}$$

input `int(sinh(b*x+a)*tanh(d*x+c),x)`output `(e**(2*a + 2*b*x) - 2*e**(2*a + b*x)*int(e**(b*x)/(e**(2*c + 2*d*x) + 1),x) * b + 2*e**(b*x)*int(1/(e**(b*x + 2*c + 2*d*x) + e**(b*x)),x)*b + 1)/(2*e*(a + b*x)*b)`

3.36 $\int \coth(2x) \sinh(x) dx$

Optimal result	316
Mathematica [A] (verified)	316
Rubi [A] (verified)	317
Maple [A] (verified)	318
Fricas [B] (verification not implemented)	319
Sympy [F]	319
Maxima [A] (verification not implemented)	319
Giac [A] (verification not implemented)	320
Mupad [B] (verification not implemented)	320
Reduce [B] (verification not implemented)	320

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \coth(2x) \sinh(x) dx = -\frac{1}{2} \arctan(\sinh(x)) + \sinh(x)$$

output `-1/2*arctan(sinh(x))+sinh(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \coth(2x) \sinh(x) dx = -\frac{1}{2} \arctan(\sinh(x)) + \sinh(x)$$

input `Integrate[Coth[2*x]*Sinh[x],x]`

output `-1/2*ArcTan[Sinh[x]] + Sinh[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4878, 27, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \coth(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(2ix)}{\csc(ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{2 \sinh^2(x) + 1}{2 (\sinh^2(x) + 1)} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{2 \sinh^2(x) + 1}{\sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{299} \\
 & \frac{1}{2} \left(2 \sinh(x) - \int \frac{1}{\sinh^2(x) + 1} d \sinh(x) \right) \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} (2 \sinh(x) - \arctan(\sinh(x)))
 \end{aligned}$$

input `Int [Coth [2*x] *Sinh [x] , x]`

output `(-ArcTan [Sinh [x]] + 2*Sinh [x])/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
default	$-\frac{\arctan(\sinh(x))}{2} + \sinh(x)$	9
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i \ln(e^x - i)}{2} - \frac{i \ln(e^x + i)}{2}$	30

input `int(coth(2*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `-1/2*arctan(sinh(x))+sinh(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(8) = 16.

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 4.20

$$\int \coth(2x) \sinh(x) dx = \frac{2(\cosh(x) + \sinh(x)) \arctan(\cosh(x) + \sinh(x)) - \cosh(x)^2 - 2\cosh(x)\sinh(x) - \sinh(x)^2 + 1}{2(\cosh(x) + \sinh(x))}$$

input `integrate(coth(2*x)*sinh(x),x, algorithm="fricas")`

output `-1/2*(2*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)) - cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + 1)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \coth(2x) \sinh(x) dx = \int \sinh(x) \coth(2x) dx$$

input `integrate(coth(2*x)*sinh(x),x)`

output `Integral(sinh(x)*coth(2*x), x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \coth(2x) \sinh(x) dx = \arctan(e^{-x}) - \frac{1}{2}e^{-x} + \frac{1}{2}e^x$$

input `integrate(coth(2*x)*sinh(x),x, algorithm="maxima")`

output `arctan(e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \coth(2x) \sinh(x) dx = -\arctan(e^x) - \frac{1}{2}e^{-x} + \frac{1}{2}e^x$$

input `integrate(coth(2*x)*sinh(x),x, algorithm="giac")`

output `-arctan(e^x) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.60

$$\int \coth(2x) \sinh(x) dx = \frac{e^x}{2} - \operatorname{atan}(e^x) - \frac{e^{-x}}{2}$$

input `int(coth(2*x)*sinh(x),x)`

output `exp(x)/2 - atan(exp(x)) - exp(-x)/2`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.30

$$\int \coth(2x) \sinh(x) dx = \frac{-2e^x \operatorname{atan}(e^x) + e^{2x} - 1}{2e^x}$$

input `int(coth(2*x)*sinh(x),x)`

output `(- 2*e**x*atan(e**x) + e**(2*x) - 1)/(2*e**x)`

3.37 $\int \coth(3x) \sinh(x) dx$

Optimal result	321
Mathematica [A] (verified)	321
Rubi [A] (verified)	322
Maple [C] (verified)	323
Fricas [B] (verification not implemented)	324
Sympy [F]	324
Maxima [B] (verification not implemented)	325
Giac [B] (verification not implemented)	325
Mupad [B] (verification not implemented)	326
Reduce [B] (verification not implemented)	326

Optimal result

Integrand size = 7, antiderivative size = 20

$$\int \coth(3x) \sinh(x) dx = -\frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{3}}\right)}{\sqrt{3}} + \sinh(x)$$

output `-1/3*arctan(2/3*sinh(x)*3^(1/2))*3^(1/2)+sinh(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \coth(3x) \sinh(x) dx = -\frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{3}}\right)}{\sqrt{3}} + \sinh(x)$$

input `Integrate[Coth[3*x]*Sinh[x],x]`

output `-(ArcTan[(2*Sinh[x])/Sqrt[3]]/Sqrt[3]) + Sinh[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4878, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \coth(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(3ix)}{\csc(ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{4 \sinh^2(x) + 1}{4 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow \text{299} \\
 & \sinh(x) - 2 \int \frac{1}{4 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow \text{216} \\
 & \sinh(x) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{\sqrt{3}}
 \end{aligned}$$

input `Int [Coth [3*x] *Sinh [x] , x]`

output `-(ArcTan [(2*Sinh [x])/Sqrt [3]]/Sqrt [3]) + Sinh [x]`

Definitions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.70

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i\sqrt{3} \ln(e^{2x} - i\sqrt{3}e^x - 1)}{6} - \frac{i\sqrt{3} \ln(e^{2x} + i\sqrt{3}e^x - 1)}{6}$	54

input `int(coth(3*x)*sinh(x), x, method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/6*I*3^(1/2)*ln(exp(2*x)-I*3^(1/2)*exp(x)-1)-1/6*I*3^(1/2)*ln(exp(2*x)+I*3^(1/2)*exp(x)-1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(16) = 32$.

Time = 0.12 (sec) , antiderivative size = 118, normalized size of antiderivative = 5.90

$$\int \coth(3x) \sinh(x) dx = \frac{2(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)) \arctan\left(\frac{1}{3}\sqrt{3} \cosh(x) + \frac{1}{3}\sqrt{3} \sinh(x)\right) - 2(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x))}{6(\cosh(x) + \sinh(x))}$$

input `integrate(coth(3*x)*sinh(x),x, algorithm="fricas")`

output `-1/6*(2*(sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*arctan(1/3*sqrt(3)*cosh(x) + 1/3*sqrt(3)*sinh(x)) - 2*(sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*arctan(-1/3*(sqrt(3)*cosh(x)^2 + 2*sqrt(3)*cosh(x)*sinh(x) + sqrt(3)*sinh(x)^2 + 2*sqrt(3)))/(cosh(x) - sinh(x)) - 3*cosh(x)^2 - 6*cosh(x)*sinh(x) - 3*sinh(x)^2 + 3)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \coth(3x) \sinh(x) dx = \int \sinh(x) \coth(3x) dx$$

input `integrate(coth(3*x)*sinh(x),x)`

output `Integral(sinh(x)*coth(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(16) = 32$.

Time = 0.14 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.45

$$\int \coth(3x) \sinh(x) dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2e^{-x} + 1) \right) + \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2e^{-x} - 1) \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(coth(3*x)*sinh(x),x, algorithm="maxima")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(-x) + 1)) + 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(-x) - 1)) - 1/2*e^(-x) + 1/2*e^x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(16) = 32$.

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \coth(3x) \sinh(x) dx = -\frac{1}{6} \sqrt{3} \left(\pi + 2 \arctan \left(\frac{1}{3} \sqrt{3} (e^{2x} - 1) e^{-x} \right) \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(coth(3*x)*sinh(x),x, algorithm="giac")`

output `-1/6*sqrt(3)*(pi + 2*arctan(1/3*sqrt(3)*(e^(2*x) - 1)*e^(-x))) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \coth(3x) \sinh(x) dx = \frac{e^x}{2} - \frac{e^{-x}}{2} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}e^x}{3} + \frac{\sqrt{3}e^{3x}}{3}\right)}{3} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}e^x}{3}\right)}{3}$$

input `int(coth(3*x)*sinh(x),x)`output `exp(x)/2 - exp(-x)/2 - (3^(1/2)*atan((2*3^(1/2)*exp(x))/3 + (3^(1/2)*exp(3*x))/3))/3 - (3^(1/2)*atan((3^(1/2)*exp(x))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.80

$$\int \coth(3x) \sinh(x) dx = \frac{-2e^x \sqrt{3} \operatorname{atan}\left(\frac{2e^x-1}{\sqrt{3}}\right) - 2e^x \sqrt{3} \operatorname{atan}\left(\frac{2e^x+1}{\sqrt{3}}\right) + 3e^{2x} - 3}{6e^x}$$

input `int(coth(3*x)*sinh(x),x)`output `(- 2*e**x*sqrt(3)*atan((2*e**x - 1)/sqrt(3)) - 2*e**x*sqrt(3)*atan((2*e**x + 1)/sqrt(3)) + 3*e**(2*x) - 3)/(6*e**x)`

3.38 $\int \coth(4x) \sinh(x) dx$

Optimal result	327
Mathematica [A] (verified)	327
Rubi [A] (verified)	328
Maple [C] (verified)	329
Fricas [B] (verification not implemented)	330
Sympy [F]	330
Maxima [B] (verification not implemented)	331
Giac [B] (verification not implemented)	331
Mupad [B] (verification not implemented)	332
Reduce [B] (verification not implemented)	332

Optimal result

Integrand size = 7, antiderivative size = 28

$$\int \coth(4x) \sinh(x) dx = -\frac{1}{4} \arctan(\sinh(x)) - \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}} + \sinh(x)$$

output

```
-1/4*arctan(sinh(x))-1/4*arctan(sinh(x)*2^(1/2))*2^(1/2)+sinh(x)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \coth(4x) \sinh(x) dx = -\frac{1}{4} \arctan(\sinh(x)) - \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}} + \sinh(x)$$

input

```
Integrate[Coth[4*x]*Sinh[x],x]
```

output

```
-1/4*ArcTan[Sinh[x]] - ArcTan[Sqrt[2]*Sinh[x]]/(2*Sqrt[2]) + Sinh[x]
```


Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4878, 27, 2205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \coth(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(4ix)}{\csc(ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{8 \sinh^4(x) + 8 \sinh^2(x) + 1}{4(2 \sinh^4(x) + 3 \sinh^2(x) + 1)} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{4} \int \frac{8 \sinh^4(x) + 8 \sinh^2(x) + 1}{2 \sinh^4(x) + 3 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{2205} \\
 & \frac{1}{4} \int \left(4 - \frac{4 \sinh^2(x) + 3}{2 \sinh^4(x) + 3 \sinh^2(x) + 1} \right) d \sinh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{4} \left(-\arctan(\sinh(x)) - \sqrt{2} \arctan(\sqrt{2} \sinh(x)) + 4 \sinh(x) \right)
 \end{aligned}$$

input

```
Int [Coth [4*x]*Sinh [x] , x]
```

output

```
(-ArcTan [Sinh [x]] - Sqrt [2]*ArcTan [Sqrt [2]*Sinh [x]] + 4*Sinh [x])/4
```

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2205 `Int[(Px)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := Int[ExpandIntegrand[Px/(a + b*x^2 + c*x^4), x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.57

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i \ln(e^x - i)}{4} - \frac{i \ln(e^x + i)}{4} + \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x - 1)}{8} - \frac{i\sqrt{2} \ln(e^{2x} + i\sqrt{2}e^x - 1)}{8}$	72

input `int(coth(4*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/4*I*ln(exp(x)-I)-1/4*I*ln(exp(x)+I)+1/8*I*2^(1/2)*ln(exp(2*x)-I*2^(1/2)*exp(x)-1)-1/8*I*2^(1/2)*ln(exp(2*x)+I*2^(1/2)*exp(x)-1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. $2(20) = 40$.

Time = 0.12 (sec) , antiderivative size = 128, normalized size of antiderivative = 4.57

$$\int \coth(4x) \sinh(x) dx =$$

$$\frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \arctan\left(\frac{1}{2} \sqrt{2} \cosh(x) + \frac{1}{2} \sqrt{2} \sinh(x)\right) - (\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \arctan\left(\frac{1}{2} \sqrt{2} \cosh(x) - \frac{1}{2} \sqrt{2} \sinh(x)\right)}{2 \cosh(x) + 2 \sinh(x)}$$

input `integrate(coth(4*x)*sinh(x),x, algorithm="fricas")`

output `-1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)) - (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2)))/(cosh(x) - sinh(x))) + 2*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 + 2)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \coth(4x) \sinh(x) dx = \int \sinh(x) \coth(4x) dx$$

input `integrate(coth(4*x)*sinh(x),x)`

output `Integral(sinh(x)*coth(4*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(20) = 40$.

Time = 0.16 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.14

$$\int \coth(4x) \sinh(x) dx = \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2e^{-x}) \right) + \frac{1}{4} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2e^{-x}) \right) + \frac{1}{2} \arctan(e^{-x}) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(coth(4*x)*sinh(x),x, algorithm="maxima")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(-x))) + 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^(-x))) + 1/2*arctan(e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. $2(20) = 40$.

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.93

$$\int \coth(4x) \sinh(x) dx = -\frac{1}{8} \pi - \frac{1}{8} \sqrt{2} \left(\pi + 2 \arctan \left(\frac{1}{2} \sqrt{2} (e^{2x} - 1) e^{-x} \right) \right) - \frac{1}{4} \arctan \left(\frac{1}{2} (e^{2x} - 1) e^{-x} \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x$$

input `integrate(coth(4*x)*sinh(x),x, algorithm="giac")`

output `-1/8*pi - 1/8*sqrt(2)*(pi + 2*arctan(1/2*sqrt(2)*(e^(2*x) - 1)*e^(-x))) - 1/4*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.86

$$\int \coth(4x) \sinh(x) dx = \frac{e^x}{2} - \frac{\operatorname{atan}(e^x)}{2} - \frac{e^{-x}}{2} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2} + \frac{\sqrt{2}e^{3x}}{2}\right)}{4} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2}\right)}{4}$$

input `int(coth(4*x)*sinh(x),x)`output `exp(x)/2 - atan(exp(x))/2 - exp(-x)/2 - (2^(1/2)*atan((2^(1/2)*exp(x))/2 + (2^(1/2)*exp(3*x))/2))/4 - (2^(1/2)*atan((2^(1/2)*exp(x))/2))/4`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.46

$$\int \coth(4x) \sinh(x) dx = \frac{-2e^x \operatorname{atan}(e^x) - e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) - e^x \sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) + 2e^{2x} - 2}{4e^x}$$

input `int(coth(4*x)*sinh(x),x)`output `(- 2*e**x*atan(e**x) - e**x*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - e**x*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) + 2*e**(2*x) - 2)/(4*e**x)`

3.39 $\int \coth(5x) \sinh(x) dx$

Optimal result	333
Mathematica [A] (verified)	333
Rubi [A] (verified)	334
Maple [C] (verified)	335
Fricas [B] (verification not implemented)	336
Sympy [F]	337
Maxima [F]	337
Giac [A] (verification not implemented)	337
Mupad [B] (verification not implemented)	338
Reduce [F]	339

Optimal result

Integrand size = 7, antiderivative size = 81

$$\int \coth(5x) \sinh(x) dx = -\frac{1}{5} \sqrt{\frac{1}{2} (5 + \sqrt{5})} \arctan \left(\sqrt{\frac{2}{5} (5 - \sqrt{5})} \sinh(x) \right) - \sqrt{\frac{2}{5 (5 + \sqrt{5})}} \arctan \left(\sqrt{\frac{2}{5} (5 + \sqrt{5})} \sinh(x) \right) + \sinh(x)$$

output

```
-1/10*(10+2*5^(1/2))^(1/2)*arctan(1/5*(50-10*5^(1/2))^(1/2)*sinh(x))-2^(1/2)/(25+5*5^(1/2))^(1/2)*arctan(1/5*(50+10*5^(1/2))^(1/2)*sinh(x))+sinh(x)
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.94

$$\int \coth(5x) \sinh(x) dx = \frac{1}{10} \left(-\sqrt{10 - 2\sqrt{5}} \arctan \left(\sqrt{2 + \frac{2}{\sqrt{5}}} \sinh(x) \right) - \sqrt{2 (5 + \sqrt{5})} \arctan \left(2\sqrt{\frac{2}{5 + \sqrt{5}}} \sinh(x) \right) + 10 \sinh(x) \right)$$

input

```
Integrate[Coth[5*x]*Sinh[x],x]
```

output

$$\left(-\left(\sqrt{10 - 2\sqrt{5}} \operatorname{ArcTan}\left[\sqrt{2 + \frac{2}{\sqrt{5}}} \operatorname{Sinh}[x] \right] \right) - \sqrt{2(5 + \sqrt{5})} \right) \operatorname{ArcTan}\left[\frac{2\sqrt{2}}{5 + \sqrt{5}} \right] \operatorname{Sinh}[x] + 10 \operatorname{Sinh}[x] \Big/ 10$$
Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4878, 2205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \coth(5x) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cot(5ix)}{\csc(ix)} dx \\ & \quad \downarrow \text{4878} \\ & \int \frac{16 \sinh^4(x) + 12 \sinh^2(x) + 1}{16 \sinh^4(x) + 20 \sinh^2(x) + 5} d \sinh(x) \\ & \quad \downarrow \text{2205} \\ & \int \left(1 - \frac{4(2 \sinh^2(x) + 1)}{16 \sinh^4(x) + 20 \sinh^2(x) + 5} \right) d \sinh(x) \\ & \quad \downarrow \text{2009} \\ & -\frac{1}{5} \sqrt{\frac{1}{2}(5 + \sqrt{5})} \arctan \left(2 \sqrt{\frac{2}{5 + \sqrt{5}}} \sinh(x) \right) - \\ & \frac{1}{5} \sqrt{\frac{1}{2}(5 - \sqrt{5})} \arctan \left(\sqrt{\frac{2}{5}} (5 + \sqrt{5}) \sinh(x) \right) + \sinh(x) \end{aligned}$$

input

$$\text{Int}[\text{Coth}[5*x]*\text{Sinh}[x], x]$$

output

$$-1/5*(\text{Sqrt}[(5 + \text{Sqrt}[5])/2]*\text{ArcTan}[2*\text{Sqrt}[2/(5 + \text{Sqrt}[5])]*\text{Sinh}[x]]) - (\text{Sqrt}[(5 - \text{Sqrt}[5])/2]*\text{ArcTan}[\text{Sqrt}[(2*(5 + \text{Sqrt}[5]))/5]*\text{Sinh}[x]])/5 + \text{Sinh}[x]$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 2205

$$\text{Int}[(P_x)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[P_x/(a + b*x^2 + c*x^4), x], x] \text{ /; } \text{FreeQ}\{a, b, c\}, x \text{ \&\& } \text{PolyQ}[P_x, x^2] \text{ \&\& } \text{Expon}[P_x, x^2] > 1$$

rule 3042

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4878

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{With}\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}\{d = \text{FreeFactors}[\text{Sin}[v], x]\}, d/\text{Coefficient}[v, x, 1] \text{ Subst}[\text{Int}[\text{SubstFor}[1, \text{Sin}[v]/d, u/\text{Cos}[v], x], x], x, \text{Sin}[v]/d]], x] \text{ /; } \text{!FalseQ}[v] \text{ \&\& } \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Sin}[v], x], u/\text{Cos}[v], x]$$
Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.52

method	result	size
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \left(\sum_{R=\text{RootOf}(2000_Z^4+100_Z^2+1)} _R \ln(-10_R e^x + e^{2x} - 1) \right)$	42

input

$$\text{int}(\text{coth}(5*x)*\text{sinh}(x), x, \text{method}=_RETURNVERBOSE)$$

output `1/2*exp(x)-1/2*exp(-x)+sum(_R*ln(-10*_R*exp(x)+exp(2*x)-1),_R=RootOf(2000*_Z^4+100*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 303 vs. $2(56) = 112$.

Time = 0.10 (sec) , antiderivative size = 303, normalized size of antiderivative = 3.74

$$\int \operatorname{coth}(5x) \sinh(x) dx$$

$$= \frac{2 \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} (\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{10} ((\sqrt{5} - 5) \cosh(x))^3 + 3(\sqrt{5} - 5) \cosh(x) \sinh(x)^2 + (\sqrt{5} - 5) \sinh(x)^3 - (\sqrt{5} + 5) \cosh(x) + (3(\sqrt{5} - 5) \cosh(x)^2 - \sqrt{5} - 5) \sinh(x)) \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} + 2 \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} (\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{10} ((\sqrt{5} - 5) \cosh(x) + (\sqrt{5} - 5) \sinh(x)) \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}}\right) - 2 \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}} (\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{10} ((\sqrt{5} + 5) \cosh(x))^3 + 3(\sqrt{5} + 5) \cosh(x) \sinh(x)^2 + (\sqrt{5} + 5) \sinh(x)^3 - (\sqrt{5} - 5) \cosh(x) + (3(\sqrt{5} + 5) \cosh(x)^2 - \sqrt{5} + 5) \sinh(x)) \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}}\right) - 2 \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}} (\cosh(x) + \sinh(x)) \arctan\left(\frac{1}{10} ((\sqrt{5} + 5) \cosh(x) + (\sqrt{5} + 5) \sinh(x)) \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}}\right) + 5 \cosh(x)^2 + 10 \cosh(x) \sinh(x) + 5 \sinh(x)^2 - 5\right)}{\cosh(x) + \sinh(x)}$$

input `integrate(coth(5*x)*sinh(x),x, algorithm="fricas")`

output `1/10*(2*sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*arctan(1/10*((sqrt(5) - 5)*cosh(x)^3 + 3*(sqrt(5) - 5)*cosh(x)*sinh(x)^2 + (sqrt(5) - 5)*sinh(x)^3 - (sqrt(5) + 5)*cosh(x) + (3*(sqrt(5) - 5)*cosh(x)^2 - sqrt(5) - 5)*sinh(x))*sqrt(1/2*sqrt(5) + 5/2)) + 2*sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*arctan(1/10*((sqrt(5) - 5)*cosh(x) + (sqrt(5) - 5)*sinh(x))*sqrt(1/2*sqrt(5) + 5/2)) - 2*sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*arctan(1/10*((sqrt(5) + 5)*cosh(x)^3 + 3*(sqrt(5) + 5)*cosh(x)*sinh(x)^2 + (sqrt(5) + 5)*sinh(x)^3 - (sqrt(5) - 5)*cosh(x) + (3*(sqrt(5) + 5)*cosh(x)^2 - sqrt(5) + 5)*sinh(x))*sqrt(-1/2*sqrt(5) + 5/2)) - 2*sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*arctan(1/10*((sqrt(5) + 5)*cosh(x) + (sqrt(5) + 5)*sinh(x))*sqrt(-1/2*sqrt(5) + 5/2)) + 5*cosh(x)^2 + 10*cosh(x)*sinh(x) + 5*sinh(x)^2 - 5)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \coth(5x) \sinh(x) dx = \int \sinh(x) \coth(5x) dx$$

input `integrate(coth(5*x)*sinh(x),x)`

output `Integral(sinh(x)*coth(5*x), x)`

Maxima [F]

$$\int \coth(5x) \sinh(x) dx = \int \coth(5x) \sinh(x) dx$$

input `integrate(coth(5*x)*sinh(x),x, algorithm="maxima")`

output `1/2*(e^(2*x) - 1)*e^(-x) - 1/2*integrate((e^(3*x) + e^(2*x) + e^x)/(e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) - 1/2*integrate((e^(3*x) - e^(2*x) + e^x)/(e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.93

$$\int \coth(5x) \sinh(x) dx = -\frac{1}{10} \sqrt{2\sqrt{5} + 10} \arctan\left(-\frac{e^{(-x)} - e^x}{\sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}}}\right) - \frac{1}{10} \sqrt{-2\sqrt{5} + 10} \arctan\left(-\frac{e^{(-x)} - e^x}{\sqrt{-\frac{1}{2}\sqrt{5} + \frac{5}{2}}}\right) - \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(coth(5*x)*sinh(x),x, algorithm="giac")`

output

```
-1/10*sqrt(2*sqrt(5) + 10)*arctan(-(e^(-x) - e^x)/sqrt(1/2*sqrt(5) + 5/2))
- 1/10*sqrt(-2*sqrt(5) + 10)*arctan(-(e^(-x) - e^x)/sqrt(-1/2*sqrt(5) + 5
/2)) - 1/2*e^(-x) + 1/2*e^x
```

Mupad [B] (verification not implemented)

Time = 2.93 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.74

$$\int \coth(5x) \sinh(x) dx = \frac{e^x}{2} - \frac{e^{-x}}{2} + \ln \left(40 e^x \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40} - 4e^{2x} + 4} \right) \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}}$$

$$+ \ln \left(40 e^x \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40} - 4e^{2x} + 4} \right) \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}}$$

$$- \ln \left(4e^{2x} + 40e^x \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40} - 4} \right) \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}}$$

$$- \ln \left(4e^{2x} + 40e^x \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40} - 4} \right) \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}}$$

input

```
int(coth(5*x)*sinh(x),x)
```

output

```
exp(x)/2 - exp(-x)/2 + log(40*exp(x)*(- 5^(1/2)/200 - 1/40)^(1/2) - 4*exp(
2*x) + 4)*(- 5^(1/2)/200 - 1/40)^(1/2) + log(40*exp(x)*(5^(1/2)/200 - 1/40
)^(1/2) - 4*exp(2*x) + 4)*(5^(1/2)/200 - 1/40)^(1/2) - log(4*exp(2*x) + 40
*exp(x)*(- 5^(1/2)/200 - 1/40)^(1/2) - 4)*(- 5^(1/2)/200 - 1/40)^(1/2) - l
og(4*exp(2*x) + 40*exp(x)*(5^(1/2)/200 - 1/40)^(1/2) - 4)*(5^(1/2)/200 - 1
/40)^(1/2)
```

Reduce [F]

$$\int \coth(5x) \sinh(x) dx = \frac{e^{2x} + 2e^x \left(\int \frac{1}{e^{9x} + e^{7x} + e^{5x} + e^{3x} + e^x} dx \right) + 1}{2e^x}$$

input

```
int(coth(5*x)*sinh(x),x)
```

output

```
(e**(2*x) + 2*e**x*int(1/(e**(9*x) + e**(7*x) + e**(5*x) + e**(3*x) + e**x),x) + 1)/(2*e**x)
```

3.40 $\int \coth(6x) \sinh(x) dx$

Optimal result	340
Mathematica [A] (verified)	340
Rubi [A] (verified)	341
Maple [C] (verified)	342
Fricas [B] (verification not implemented)	343
Sympy [F]	343
Maxima [F]	344
Giac [B] (verification not implemented)	344
Mupad [B] (verification not implemented)	345
Reduce [B] (verification not implemented)	345

Optimal result

Integrand size = 7, antiderivative size = 38

$$\int \coth(6x) \sinh(x) dx = -\frac{1}{6} \arctan(\sinh(x)) - \frac{1}{6} \arctan(2 \sinh(x)) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{2\sqrt{3}} + \sinh(x)$$

output `-1/6*arctan(sinh(x))-1/6*arctan(2*sinh(x))-1/6*arctan(2/3*sinh(x)*3^(1/2))*3^(1/2)+sinh(x)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \coth(6x) \sinh(x) dx = -\frac{1}{6} \arctan(\sinh(x)) - \frac{1}{6} \arctan(2 \sinh(x)) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{2\sqrt{3}} + \sinh(x)$$

input `Integrate[Coth[6*x]*Sinh[x],x]`

output

```
-1/6*ArcTan[Sinh[x]] - ArcTan[2*Sinh[x]]/6 - ArcTan[(2*Sinh[x])/Sqrt[3]]/(2*Sqrt[3]) + Sinh[x]
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4878, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \coth(6x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(6ix)}{\csc(ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{32 \sinh^6(x) + 48 \sinh^4(x) + 18 \sinh^2(x) + 1}{2 (16 \sinh^6(x) + 32 \sinh^4(x) + 19 \sinh^2(x) + 3)} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{32 \sinh^6(x) + 48 \sinh^4(x) + 18 \sinh^2(x) + 1}{16 \sinh^6(x) + 32 \sinh^4(x) + 19 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow \text{2460} \\
 & \frac{1}{2} \int \left(-\frac{2}{3 (4 \sinh^2(x) + 1)} - \frac{2}{4 \sinh^2(x) + 3} + 2 - \frac{1}{3 (\sinh^2(x) + 1)} \right) d \sinh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(-\frac{1}{3} \arctan(\sinh(x)) - \frac{1}{3} \arctan(2 \sinh(x)) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{\sqrt{3}} + 2 \sinh(x) \right)
 \end{aligned}$$

input

```
Int [Coth [6*x] *Sinh [x] , x]
```

output
$$\frac{(-1/3 \operatorname{ArcTan}[\operatorname{Sinh}[x]] - \operatorname{ArcTan}[2 \operatorname{Sinh}[x]]/3 - \operatorname{ArcTan}[(2 \operatorname{Sinh}[x])/\sqrt{3}])/\sqrt{3} + 2 \operatorname{Sinh}[x])/2}$$

Defintions of rubi rules used

rule 27
$$\operatorname{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[Fx, (b_*)(Gx_)] /; \operatorname{FreeQ}[b, x]$$

rule 2009
$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 2460
$$\operatorname{Int}[(u_*)(Px_)^{(p_)}, x_Symbol] \rightarrow \operatorname{With}[\{Qx = \operatorname{Factor}[Px /. x \rightarrow \sqrt{x}]\}, \operatorname{Int}[\operatorname{ExpandIntegrand}[u*(Qx /. x \rightarrow x^2)^p, x], x] /; \ !\operatorname{SumQ}[\operatorname{NonfreeFactors}[Qx, x]] /; \operatorname{PolyQ}[Px, x^2] \ \&\& \ \operatorname{GtQ}[\operatorname{Expon}[Px, x], 2] \ \&\& \ !\operatorname{BinomialQ}[Px, x] \ \&\& \ !\operatorname{TrinomialQ}[Px, x] \ \&\& \ \operatorname{ILtQ}[p, 0] \ \&\& \ \operatorname{RationalFunctionQ}[u, x]$$

rule 3042
$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4878
$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{With}[\{v = \operatorname{FunctionOfTrig}[u, x]\}, \operatorname{Simp}[\operatorname{With}[\{d = \operatorname{FreeFactors}[\operatorname{Sin}[v], x]\}, d/\operatorname{Coefficient}[v, x, 1] \operatorname{Subst}[\operatorname{Int}[\operatorname{SubstFor}[1, \operatorname{Sin}[v]/d, u/\operatorname{Cos}[v], x], x], x, \operatorname{Sin}[v]/d]], x] /; \ !\operatorname{FalseQ}[v] \ \&\& \ \operatorname{FunctionOfQ}[\operatorname{NonfreeFactors}[\operatorname{Sin}[v], x], u/\operatorname{Cos}[v], x]]$$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.68

method	result
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2} + \frac{i \ln(e^x - i)}{6} - \frac{i \ln(e^x + i)}{6} + \frac{i\sqrt{3} \ln(e^{2x} - i\sqrt{3}e^x - 1)}{12} - \frac{i\sqrt{3} \ln(e^{2x} + i\sqrt{3}e^x - 1)}{12} + \frac{i \ln(e^{2x} - ie^x - 1)}{12} - \frac{i \ln(e^{2x} + ie^x - 1)}{12}$

input `int(coth(6*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)-1/2*exp(-x)+1/6*I*ln(exp(x)-I)-1/6*I*ln(exp(x)+I)+1/12*I*3^(1/2)*ln(exp(2*x)-I*3^(1/2)*exp(x)-1)-1/12*I*3^(1/2)*ln(exp(2*x)+I*3^(1/2)*exp(x)-1)+1/12*I*ln(exp(2*x)-I*exp(x)-1)-1/12*I*ln(exp(2*x)+I*exp(x)-1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 164 vs. $2(28) = 56$.

Time = 0.10 (sec) , antiderivative size = 164, normalized size of antiderivative = 4.32

$$\int \coth(6x) \sinh(x) dx =$$

$$\frac{(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)) \arctan\left(\frac{1}{3} \sqrt{3} \cosh(x) + \frac{1}{3} \sqrt{3} \sinh(x)\right) - (\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)) \arctan\left(\frac{1}{3} \sqrt{3} \cosh(x) - \frac{1}{3} \sqrt{3} \sinh(x)\right)}{3}$$

input `integrate(coth(6*x)*sinh(x),x, algorithm="fricas")`

output `-1/6*((sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*arctan(1/3*sqrt(3)*cosh(x) + 1/3*sqrt(3)*sinh(x)) - (sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*arctan(-1/3*(sqrt(3)*cosh(x)^2 + 2*sqrt(3)*cosh(x)*sinh(x) + sqrt(3)*sinh(x)^2 + 2*sqrt(3)))/(cosh(x) - sinh(x))) - (cosh(x) + sinh(x))*arctan(-(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)/(cosh(x) - sinh(x))) + 3*(cosh(x) + sinh(x))*arctan(cosh(x) + sinh(x)) - 3*cosh(x)^2 - 6*cosh(x)*sinh(x) - 3*sinh(x)^2 + 3)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \coth(6x) \sinh(x) dx = \int \sinh(x) \coth(6x) dx$$

input `integrate(coth(6*x)*sinh(x),x)`

output `Integral(sinh(x)*coth(6*x), x)`

Maxima [F]

$$\int \coth(6x) \sinh(x) dx = \int \coth(6x) \sinh(x) dx$$

input `integrate(coth(6*x)*sinh(x),x, algorithm="maxima")`

output `1/2*(e^(2*x) - 1)*e^(-x) - 1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) - 1/3*arctan(e^x) - 1/2*integrate(1/3*(e^(3*x) + e^x)/(e^(4*x) - e^(2*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(28) = 56$.

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.79

$$\begin{aligned} \int \coth(6x) \sinh(x) dx = & -\frac{1}{6} \pi - \frac{1}{12} \sqrt{3} \left(\pi + 2 \arctan \left(\frac{1}{3} \sqrt{3} (e^{2x} - 1) e^{-x} \right) \right) \\ & - \frac{1}{6} \arctan \left((e^{2x} - 1) e^{-x} \right) \\ & - \frac{1}{6} \arctan \left(\frac{1}{2} (e^{2x} - 1) e^{-x} \right) - \frac{1}{2} e^{-x} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(coth(6*x)*sinh(x),x, algorithm="giac")`

output `-1/6*pi - 1/12*sqrt(3)*(pi + 2*arctan(1/3*sqrt(3)*(e^(2*x) - 1)*e^(-x))) - 1/6*arctan((e^(2*x) - 1)*e^(-x)) - 1/6*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.47

$$\int \coth(6x) \sinh(x) dx = \frac{e^x}{2} - \frac{\operatorname{atan}(e^x)}{3} - \frac{e^{-x}}{2} - \frac{\operatorname{atan}\left(36 e^{-x} \left(\frac{e^{2x}}{36} - \frac{1}{36}\right)\right)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(4 \sqrt{3} e^{-x} \left(\frac{e^{2x}}{12} - \frac{1}{12}\right)\right)}{6}$$

input `int(coth(6*x)*sinh(x),x)`output `exp(x)/2 - atan(exp(x))/3 - exp(-x)/2 - atan(36*exp(-x)*(exp(2*x)/36 - 1/36))/6 - (3^(1/2)*atan(4*3^(1/2)*exp(-x)*(exp(2*x)/12 - 1/12)))/6`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.50

$$\int \coth(6x) \sinh(x) dx = \frac{-2e^x \operatorname{atan}(e^x) - e^x \operatorname{atan}(2e^x - \sqrt{3}) - e^x \operatorname{atan}(2e^x + \sqrt{3}) - e^x \sqrt{3} \operatorname{atan}\left(\frac{2e^x - 1}{\sqrt{3}}\right) - e^x \sqrt{3} \operatorname{atan}\left(\frac{2e^x + 1}{\sqrt{3}}\right) + 3}{6e^x}$$

input `int(coth(6*x)*sinh(x),x)`output `(- 2*e**x*atan(e**x) - e**x*atan(2*e**x - sqrt(3)) - e**x*atan(2*e**x + sqrt(3)) - e**x*sqrt(3)*atan((2*e**x - 1)/sqrt(3)) - e**x*sqrt(3)*atan((2*e**x + 1)/sqrt(3)) + 3*e**(2*x) - 3)/(6*e**x)`

3.41 $\int \coth(c + bx) \sinh(a + bx) dx$

Optimal result	346
Mathematica [C] (verified)	346
Rubi [A] (verified)	347
Maple [B] (verified)	348
Fricas [B] (verification not implemented)	349
Sympy [F]	350
Maxima [B] (verification not implemented)	350
Giac [B] (verification not implemented)	351
Mupad [B] (verification not implemented)	351
Reduce [B] (verification not implemented)	352

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \coth(c + bx) \sinh(a + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \sinh(a - c)}{b} + \frac{\sinh(a + bx)}{b}$$

output

$$-\operatorname{arctanh}(\cosh(b*x+c))*\sinh(a-c)/b+\sinh(b*x+a)/b$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.21

$$\begin{aligned} & \int \coth(c + bx) \sinh(a + bx) dx \\ &= \frac{\cosh(bx) \sinh(a)}{b} \\ & \quad - \frac{2i \operatorname{arctan} \left(\frac{(\cosh(c) - \sinh(c)) \left(\cosh(c) \cosh\left(\frac{bx}{2}\right) + \sinh(c) \sinh\left(\frac{bx}{2}\right) \right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)} \right) \sinh(a - c)}{b} \\ & \quad + \frac{\cosh(a) \sinh(bx)}{b} \end{aligned}$$

input `Integrate[Coth[c + b*x]*Sinh[a + b*x],x]`

output $(\text{Cosh}[b*x]*\text{Sinh}[a])/b - ((2*I)*\text{ArcTan}(((\text{Cosh}[c] - \text{Sinh}[c])*(\text{Cosh}[c]*\text{Cosh}[(b*x)/2] + \text{Sinh}[c]*\text{Sinh}[(b*x)/2]))/(I*\text{Cosh}[c]*\text{Cosh}[(b*x)/2] - I*\text{Cosh}[(b*x)/2]*\text{Sinh}[c]))*\text{Sinh}[a - c])/b + (\text{Cosh}[a]*\text{Sinh}[b*x])/b$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6156, 3042, 26, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \coth(bx + c) dx \\
 & \quad \downarrow \text{6156} \\
 & \sinh(a - c) \int \text{csch}(c + bx) dx + \int \cosh(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sinh(a - c) \int i \csc(ic + ibx) dx + \int \sin\left(ia + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & i \sinh(a - c) \int \csc(ic + ibx) dx + \int \sin\left(ia + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3117} \\
 & \frac{\sinh(a + bx)}{b} + i \sinh(a - c) \int \csc(ic + ibx) dx \\
 & \quad \downarrow \text{4257} \\
 & \frac{\sinh(a + bx)}{b} - \frac{\sinh(a - c) \text{arctanh}(\cosh(bx + c))}{b}
 \end{aligned}$$

input `Int[Coth[c + b*x]*Sinh[a + b*x],x]`

output `-((ArcTanh[Cosh[c + b*x]]*Sinh[a - c])/b) + Sinh[a + b*x]/b`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6156 `Int[Coth[w_]^(n_.)*Sinh[v_], x_Symbol] := Int[Cosh[v]*Coth[w]^(n - 1), x] + Simp[Sinh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 154 vs. $2(29) = 58$.

Time = 0.09 (sec) , antiderivative size = 155, normalized size of antiderivative = 5.34

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(coth(b*x+c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output

```
1/2/b*exp(b*x+a)-1/2/b*exp(-b*x-a)+1/2*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)
*exp(2*a)-1/2*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*c)-1/2*ln(exp(b*x+
a)+exp(a-c))/b*exp(-a-c)*exp(2*a)+1/2*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*
exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 439 vs. $2(29) = 58$.

Time = 0.10 (sec) , antiderivative size = 439, normalized size of antiderivative = 15.14

$$\int \coth(c + bx) \sinh(a + bx) dx$$

$$= \frac{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c)^2 \cosh(-a + c) \sinh(-a + c) + \cosh(bx + c)^2 \sinh(-a + c)^2}{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c) \cosh(-a + c) \sinh(-a + c) + \sinh(-a + c)^2}$$

input

```
integrate(coth(b*x+c)*sinh(b*x+a),x, algorithm="fricas")
```

output

```
1/2*(cosh(b*x + c)^2*cosh(-a + c)^2 - 2*cosh(b*x + c)^2*cosh(-a + c)*sinh(-
-a + c) + cosh(b*x + c)^2*sinh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)
)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^2 + (2*cosh(b*x + c)*cosh(-
a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*
cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a +
c)^2 - 1)*sinh(b*x + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) - (2*cosh(
b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(
-a + c)^2 - 1)*cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a +
c) + sinh(-a + c)^2 - 1)*sinh(b*x + c))*log(cosh(b*x + c) + sinh(b*x + c)
- 1) + 2*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh
(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 1)/(b*cosh(b*x +
c)*cosh(-a + c) - b*cosh(b*x + c)*sinh(-a + c) + (b*cosh(-a + c) - b*sinh(-
a + c))*sinh(b*x + c))
```

Sympy [F]

$$\int \coth(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \coth(bx + c) dx$$

input `integrate(coth(b*x+c)*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*coth(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(29) = 58$.

Time = 0.04 (sec) , antiderivative size = 94, normalized size of antiderivative = 3.24

$$\begin{aligned} \int \coth(c + bx) \sinh(a + bx) dx = & -\frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} \\ & + \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} \\ & + \frac{e^{(bx+a)}}{2b} - \frac{e^{(-bx-a)}}{2b} \end{aligned}$$

input `integrate(coth(b*x+c)*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/2*e^(b*x + a)/b - 1/2*e^(-b*x - a)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. $2(29) = 58$.

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.21

$$\int \coth(c + bx) \sinh(a + bx) dx = \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - (e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) - e^{(bx+a+c)}}{2b}$$

input `integrate(coth(b*x+c)*sinh(b*x+a),x, algorithm="giac")`

output
$$-1/2*((e^{(2*a + c)} - e^{(3*c)})*e^{(-a - 2*c)}*\log(e^{(b*x + a + c)} + e^a) - (e^{(2*a + c)} - e^{(3*c)})*e^{(-a - 2*c)}*\log(\text{abs}(e^{(b*x + a + c)} - e^a)) - e^{(b*x + a)} + e^{(-b*x - a)})/b$$

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 139, normalized size of antiderivative = 4.79

$$\int \coth(c + bx) \sinh(a + bx) dx = \frac{e^{a+bx}}{2b} - \frac{e^{-a-bx}}{2b} + \frac{\text{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2} - e^{2a} e^{-2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{-b^2}}$$

input `int(coth(c + b*x)*sinh(a + b*x),x)`

output
$$\exp(a + b*x)/(2*b) - \exp(-a - b*x)/(2*b) + (\text{atan}((\exp(-a)*\exp(2*c)*\exp(b*x))*((-b^2)^(1/2) - \exp(2*a)*\exp(-2*c)*(-b^2)^(1/2)))/(b*(\exp(-2*a)*\exp(2*c))*(\exp(4*a)*\exp(-4*c) - 2*\exp(2*a)*\exp(-2*c) + 1)^(1/2)))*(\exp(2*c - 2*a)*(\exp(4*a - 4*c) - 2*\exp(2*a - 2*c) + 1)^(1/2))/(-b^2)^(1/2)$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.93

$$\int \coth(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+2a+c} + e^{bx+2a} \log(e^{bx+c} - 1) - e^{bx+2a} \log(e^{bx+c} + 1) - e^{bx+2c} \log(e^{bx+c} - 1) + e^{bx+2c} \log(e^{bx+c} + 1) - e^{bx+2c}}{2e^{bx+a+cb}}$$

input

```
int(coth(b*x+c)*sinh(b*x+a),x)
```

output

```
(e**(2*a + 2*b*x + c) + e**(2*a + b*x)*log(e**(b*x + c) - 1) - e**(2*a + b*x)*log(e**(b*x + c) + 1) - e**(b*x + 2*c)*log(e**(b*x + c) - 1) + e**(b*x + 2*c)*log(e**(b*x + c) + 1) - e**c)/(2*e**(a + b*x + c)*b)
```

3.42 $\int \coth^2(c + bx) \sinh(a + bx) dx$

Optimal result	353
Mathematica [C] (verified)	353
Rubi [A] (verified)	354
Maple [B] (verified)	356
Fricas [B] (verification not implemented)	357
Sympy [F]	358
Maxima [B] (verification not implemented)	358
Giac [B] (verification not implemented)	359
Mupad [B] (verification not implemented)	359
Reduce [B] (verification not implemented)	360

Optimal result

Integrand size = 15, antiderivative size = 46

$$\int \coth^2(c + bx) \sinh(a + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(a - c)}{b} + \frac{\cosh(a + bx)}{b} - \frac{\operatorname{csch}(c + bx) \sinh(a - c)}{b}$$

output

```
-arctanh(cosh(b*x+c))*cosh(a-c)/b+cosh(b*x+a)/b-csch(b*x+c)*sinh(a-c)/b
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.39

$$\int \coth^2(c + bx) \sinh(a + bx) dx = -\frac{2i \arctan\left(\frac{(\cosh(c) - \sinh(c))(\cosh(c) \cosh(\frac{bx}{2}) + \sinh(c) \sinh(\frac{bx}{2}))}{i \cosh(c) \cosh(\frac{bx}{2}) - i \cosh(\frac{bx}{2}) \sinh(c)}\right) \cosh(a - c)}{b} + \frac{\cosh(a) \cosh(bx)}{b} - \frac{\operatorname{csch}(c + bx) \sinh(a - c)}{b} + \frac{\sinh(a) \sinh(bx)}{b}$$

input `Integrate[Coth[c + b*x]^2*Sinh[a + b*x],x]`

output
$$\frac{((-2*I)*\text{ArcTan}[\frac{(\text{Cosh}[c] - \text{Sinh}[c])*(\text{Cosh}[c]*\text{Cosh}[(b*x)/2] + \text{Sinh}[c]*\text{Sinh}[(b*x)/2])}{I*\text{Cosh}[c]*\text{Cosh}[(b*x)/2] - I*\text{Cosh}[(b*x)/2]*\text{Sinh}[c]}] * \text{Cosh}[a - c])}{b} + \frac{\text{Cosh}[a]*\text{Cosh}[b*x]}{b} - \frac{\text{Csch}[c + b*x]*\text{Sinh}[a - c]}{b} + \frac{\text{Sinh}[a]*\text{Sinh}[b*x]}{b}$$

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6156, 3042, 3086, 24, 6155, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(a + bx) \coth^2(bx + c) dx \\ & \quad \downarrow \text{6156} \\ & \int \cosh(a + bx) \coth(c + bx) dx + \sinh(a - c) \int \coth(c + bx) \text{csch}(c + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int \cosh(a + bx) \coth(c + bx) dx + \sinh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3086} \\ & \int \cosh(a + bx) \coth(c + bx) dx - \frac{i \sinh(a - c) \int 1d(-i\text{csch}(c + bx))}{b} \\ & \quad \downarrow \text{24} \\ & \int \cosh(a + bx) \coth(c + bx) dx - \frac{\sinh(a - c) \text{csch}(bx + c)}{b} \\ & \quad \downarrow \text{6155} \\ & \cosh(a - c) \int \text{csch}(c + bx) dx + \int \sinh(a + bx) dx - \frac{\sinh(a - c) \text{csch}(bx + c)}{b} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
& \cosh(a-c) \int i \csc(ic+ibx) dx + \int -i \sin(ia+ibx) dx - \frac{\sinh(a-c) \operatorname{csch}(bx+c)}{b} \\
& \quad \downarrow 26 \\
& i \cosh(a-c) \int \csc(ic+ibx) dx - i \int \sin(ia+ibx) dx - \frac{\sinh(a-c) \operatorname{csch}(bx+c)}{b} \\
& \quad \downarrow 3118 \\
& i \cosh(a-c) \int \csc(ic+ibx) dx - \frac{\sinh(a-c) \operatorname{csch}(bx+c)}{b} + \frac{\cosh(a+bx)}{b} \\
& \quad \downarrow 4257 \\
& -\frac{\cosh(a-c) \operatorname{arctanh}(\cosh(bx+c))}{b} - \frac{\sinh(a-c) \operatorname{csch}(bx+c)}{b} + \frac{\cosh(a+bx)}{b}
\end{aligned}$$

input `Int[Coth[c + b*x]^2*Sinh[a + b*x],x]`

output `-((ArcTanh[Cosh[c + b*x]]*Cosh[a - c])/b) + Cosh[a + b*x]/b - (Csch[c + b*x]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6155 `Int[Cosh[v_]*Coth[w_]^(n_), x_Symbol] := Int[Sinh[v]*Coth[w]^(n - 1), x] + Simp[Cosh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

rule 6156 `Int[Coth[w_]^(n_.)*Sinh[v_], x_Symbol] := Int[Cosh[v]*Coth[w]^(n - 1), x] + Simp[Sinh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. $2(46) = 92$.

Time = 0.11 (sec) , antiderivative size = 197, normalized size of antiderivative = 4.28

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(e^{2a}-e^{2c})}{b(-e^{2bx+2a+2c}+e^{2a})} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b}$

input `int(coth(b*x+c)^2*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output $\frac{1}{2} \frac{1}{b} \exp(bx+a) + \frac{1}{2} \frac{1}{b} \exp(-bx-a) + \frac{1}{b} \exp(bx+a) \frac{(\exp(2a) - \exp(2c))}{(-\exp(2bx+2a+2c) + \exp(2a))} - \frac{1}{2} \frac{\ln(\exp(bx+a) + \exp(a-c))}{b \exp(-a-c) \exp(2a)} - \frac{1}{2} \frac{\ln(\exp(bx+a) + \exp(a-c))}{b \exp(-a-c) \exp(2c)} + \frac{1}{2} \frac{\ln(\exp(bx+a) - \exp(a-c))}{b \exp(-a-c) \exp(2a)} + \frac{1}{2} \frac{\ln(\exp(bx+a) - \exp(a-c))}{b \exp(-a-c) \exp(2c)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1237 vs. $2(46) = 92$.

Time = 0.11 (sec) , antiderivative size = 1237, normalized size of antiderivative = 26.89

$$\int \coth^2(c + bx) \sinh(a + bx) dx = \text{Too large to display}$$

input `integrate(coth(b*x+c)^2*sinh(b*x+a),x, algorithm="fricas")`

output

```

1/2*(cosh(b*x + c)^4*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sin
h(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^4 + 4*(cosh(b*x + c)*cosh(-a + c
)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a +
c)^2)*sinh(b*x + c)^3 - 3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + 3*(2*cosh
(b*x + c)^2*cosh(-a + c)^2 + (2*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh
(-a + c)^2 - 2*(2*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c
) + 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^4 - 3*cosh(b*x + c)^2)*sinh(-a + c
)^2 - ((cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-a
+ c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x +
c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c
)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 - cosh(b*x + c)
)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c) + (3*(cosh(-a + c)^2
+ 1)*cosh(b*x + c)^2 + (3*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a +
c)^2 - 2*(3*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) - 1
)*sinh(b*x + c) - 2*(cosh(b*x + c)^3*cosh(-a + c) - cosh(b*x + c)*cosh(-a
+ c))*sinh(-a + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) + ((cosh(-a + c
)^2 + 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) +
sinh(-a + c)^2 + 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x + c)*cosh(-a + c)*sin
h(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x +
c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 - cosh(b*x + c))*sinh(-a + c)^2...
```

Sympy [F]

$$\int \coth^2(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \coth^2(bx + c) dx$$

input `integrate(coth(b*x+c)**2*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*coth(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. $2(46) = 92$.

Time = 0.06 (sec) , antiderivative size = 140, normalized size of antiderivative = 3.04

$$\begin{aligned} \int \coth^2(c + bx) \sinh(a + bx) dx = & -\frac{(e^{2a} + e^{2c})e^{(-a-c)} \log(e^{-bx} + e^c)}{2b} \\ & + \frac{(e^{2a} + e^{2c})e^{(-a-c)} \log(e^{-bx} - e^c)}{2b} \\ & + \frac{e^{(-bx-a)}}{2b} - \frac{(3e^{2a} - 2e^{2c})e^{(-2bx-2a)} - e^{2c}}{2b(e^{(-bx-a+2c)} - e^{(-3bx-a)})} \end{aligned}$$

input `integrate(coth(b*x+c)^2*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/2*e^(-b*x - a)/b - 1/2*((3*e^(2*a) - 2*e^(2*c))*e^(-2*b*x - 2*a) - e^(2*c))/(b*(e^(-b*x - a + 2*c) - e^(-3*b*x - a)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(46) = 92$.

Time = 0.11 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.96

$$\int \coth^2(c + bx) \sinh(a + bx) dx = \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - (e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) + \frac{2e^{(2bx+c)}}{e^{(3c)}}}{2b}$$

input `integrate(coth(b*x+c)^2*sinh(b*x+a),x, algorithm="giac")`

output
$$-1/2*((e^{(2*a + c)} + e^{(3*c)})*e^{(-a - 2*c)}*\log(e^{(b*x + a + c)} + e^a) - (e^{(2*a + c)} + e^{(3*c)})*e^{(-a - 2*c)}*\log(\text{abs}(e^{(b*x + a + c)} - e^a)) + (2*e^{(2*b*x + 4*a)} - 3*e^{(2*b*x + 2*a + 2*c)} + e^{(2*a)})/(e^{(3*b*x + 3*a + 2*c)} - e^{(b*x + 3*a)} - e^{(b*x + a)})/b$$

Mupad [B] (verification not implemented)

Time = 1.27 (sec) , antiderivative size = 181, normalized size of antiderivative = 3.93

$$\begin{aligned} & \int \coth^2(c + bx) \sinh(a + bx) dx \\ &= \frac{e^{a+bx}}{2b} + \frac{e^{-a-bx}}{2b} \\ & - \frac{\text{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2} + e^{2a} e^{-2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{-b^2}} \\ & + \frac{e^{a+bx} (e^{2a-2c} - 1)}{b (e^{2a-2c} - e^{2a+2bx})} \end{aligned}$$

input `int(coth(c + b*x)^2*sinh(a + b*x),x)`

output

```
exp(a + b*x)/(2*b) + exp(- a - b*x)/(2*b) - (atan((exp(-a)*exp(2*c)*exp(b*
x)*((-b^2)^(1/2) + exp(2*a)*exp(-2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c
)*(2*exp(2*a)*exp(-2*c) + exp(4*a)*exp(-4*c) + 1))^(1/2)))*(exp(2*c - 2*a)
*(2*exp(2*a - 2*c) + exp(4*a - 4*c) + 1))^(1/2))/(-b^2)^(1/2) + (exp(a + b
*x)*(exp(2*a - 2*c) - 1))/(b*(exp(2*a - 2*c) - exp(2*a + 2*b*x)))
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 247, normalized size of antiderivative = 5.37

$$\int \coth^2(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{4bx+2a+3c} + e^{3bx+2a+2c} \log(e^{bx+c} - 1) - e^{3bx+2a+2c} \log(e^{bx+c} + 1) + e^{3bx+4c} \log(e^{bx+c} - 1) - e^{3bx+4c} \log(e^{bx+c} + 1)}{b}$$

input

```
int(coth(b*x+c)^2*sinh(b*x+a),x)
```

output

```
(e**(2*a + 4*b*x + 3*c) + e**(2*a + 3*b*x + 2*c)*log(e**(b*x + c) - 1) - e
**(2*a + 3*b*x + 2*c)*log(e**(b*x + c) + 1) + e**(3*b*x + 4*c)*log(e**(b*x
+ c) - 1) - e**(3*b*x + 4*c)*log(e**(b*x + c) + 1) - 3*e**(2*a + 2*b*x +
c) + 3*e**(2*b*x + 3*c) - e**(2*a + b*x)*log(e**(b*x + c) - 1) + e**(2*a +
b*x)*log(e**(b*x + c) + 1) - e**(b*x + 2*c)*log(e**(b*x + c) - 1) + e**(b
*x + 2*c)*log(e**(b*x + c) + 1) - e**c)/(2*e**(a + b*x + c)*b*(e**(2*b*x +
2*c) - 1))
```

3.43 $\int \coth^3(c + bx) \sinh(a + bx) dx$

Optimal result	361
Mathematica [A] (verified)	361
Rubi [C] (verified)	362
Maple [B] (verified)	366
Fricas [B] (verification not implemented)	366
Sympy [F]	367
Maxima [B] (verification not implemented)	368
Giac [B] (verification not implemented)	368
Mupad [F(-1)]	369
Reduce [B] (verification not implemented)	369

Optimal result

Integrand size = 15, antiderivative size = 73

$$\int \coth^3(c + bx) \sinh(a + bx) dx = -\frac{\cosh(a - c)\operatorname{csch}(c + bx)}{b} - \frac{3\operatorname{arctanh}(\cosh(c + bx)) \sinh(a - c)}{2b} - \frac{\coth(c + bx)\operatorname{csch}(c + bx) \sinh(a - c)}{2b} + \frac{\sinh(a + bx)}{b}$$

output

$$-\cosh(a-c)*\operatorname{csch}(b*x+c)/b-3/2*\operatorname{arctanh}(\cosh(b*x+c))*\sinh(a-c)/b-1/2*\coth(b*x+c)*\operatorname{csch}(b*x+c)*\sinh(a-c)/b+\sinh(b*x+a)/b$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.96

$$\int \coth^3(c + bx) \sinh(a + bx) dx = \frac{-12\operatorname{arctanh}\left(\cosh(c) + \sinh(c) \tanh\left(\frac{bx}{2}\right)\right) \sinh(a - c) + \operatorname{csch}^2(c + bx)(2 \sinh(a - 2c - bx) - 5 \sinh(a + bx))}{4b}$$

input `Integrate[Coth[c + b*x]^3*Sinh[a + b*x],x]`

output `(-12*ArcTanh[Cosh[c] + Sinh[c]*Tanh[(b*x)/2]]*Sinh[a - c] + Csch[c + b*x]^2*(2*Sinh[a - 2*c - b*x] - 5*Sinh[a + b*x] + Sinh[a + 2*c + 3*b*x]))/(4*b)`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.29, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 1.133$, Rules used = {6156, 3042, 26, 3091, 26, 3042, 26, 4257, 6155, 3042, 3086, 24, 6156, 3042, 26, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \coth^3(bx + c) dx \\
 & \quad \downarrow \text{6156} \\
 & \int \cosh(a + bx) \coth^2(c + bx) dx + \sinh(a - c) \int \coth^2(c + bx) \operatorname{csch}(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cosh(a + bx) \coth^2(c + bx) dx + \sinh(a - c) \int -i \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{26} \\
 & \int \cosh(a + bx) \coth^2(c + bx) dx - i \sinh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right) \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right)^2 dx \\
 & \quad \downarrow \text{3091} \\
 & \int \cosh(a + bx) \coth^2(c + bx) dx - i \sinh(a - c) \left(-\frac{1}{2} \int -i \operatorname{csch}(c + bx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)
 \end{aligned}$$

↓ 26

$$c) \left(\int \cosh(a + bx) \coth^2(c + bx) dx - i \sinh(a - \frac{1}{2} i \int \operatorname{csch}(c + bx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3042

$$c) \left(\int \cosh(a + bx) \coth^2(c + bx) dx - i \sinh(a - \frac{1}{2} i \int i \operatorname{csc}(ic + ibx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 26

$$c) \left(-\frac{1}{2} \int \operatorname{csc}(ic + ibx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 4257

$$c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 6155

$$\int \coth(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int \coth(c + bx) \operatorname{csch}(c + bx) dx - i \sinh(a - c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3042

$$\int \coth(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx - i \sinh(a - c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3086

$$-\frac{i \cosh(a - c) \int 1d(-i \operatorname{csch}(c + bx))}{b} + \int \coth(c + bx) \sinh(a + bx) dx - i \sinh(a - c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 24

$$\begin{aligned}
& \int \coth(c + bx) \sinh(a + bx) dx - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow \text{6156} \\
& \sinh(a - c) \int \operatorname{csch}(c + bx) dx + \int \cosh(a + bx) dx - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow \text{3042} \\
& \sinh(a - c) \int i \csc(ic + ibx) dx + \int \sin\left(ia + ibx + \frac{\pi}{2}\right) dx - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow \text{26} \\
& i \sinh(a - c) \int \csc(ic + ibx) dx + \int \sin\left(ia + ibx + \frac{\pi}{2}\right) dx - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow \text{3117} \\
& i \sinh(a - c) \int \csc(ic + ibx) dx - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} + \\
& \quad \frac{\sinh(a + bx)}{b} \\
& \downarrow \text{4257} \\
& -\frac{\sinh(a - c) \operatorname{arctanh}(\cosh(bx + c))}{b} - i \sinh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} + \\
& \quad \frac{\sinh(a + bx)}{b}
\end{aligned}$$

input

```
Int[Coth[c + b*x]^3*Sinh[a + b*x],x]
```

output
$$-\left(\frac{\cosh[a - c] \operatorname{csch}[c + b x]}{b} - \frac{\operatorname{ArcTanh}[\cosh[c + b x]] \sinh[a - c]}{b}\right) - I \left(\frac{(-1/2 I) \operatorname{ArcTanh}[\cosh[c + b x]]}{b} - \frac{(I/2) \operatorname{Coth}[c + b x] \operatorname{csch}[c + b x]}{b}\right) \sinh[a - c] + \frac{\sinh[a + b x]}{b}$$

Defintions of rubi rules used

rule 24
$$\operatorname{Int}[a_, x_Symbol] \rightarrow \operatorname{Simp}[a x, x] \text{ ; FreeQ}[a, x]$$

rule 26
$$\operatorname{Int}[(\operatorname{Complex}[0, a_]) (F x_), x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$$

rule 3042
$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3086
$$\operatorname{Int}[(a_ \operatorname{sec}[e_] + (f_)(x_)]^{(m_)} ((b_)\operatorname{tan}[e_] + (f_)(x_)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[a/f \operatorname{Subst}[\operatorname{Int}[(a x)^{(m-1)} (-1+x^2)^{(n-1)/2}], x], x, \operatorname{Sec}[e + f x], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \operatorname{IntegerQ}[(n-1)/2] \ \&\& \ !(\operatorname{IntegerQ}[m/2] \ \&\& \ \operatorname{LtQ}[0, m, n+1])$$

rule 3091
$$\operatorname{Int}[(a_ \operatorname{sec}[e_] + (f_)(x_)]^{(m_)} ((b_)\operatorname{tan}[e_] + (f_)(x_)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[b (a \operatorname{Sec}[e + f x])^m ((b \operatorname{Tan}[e + f x])^{(n-1)} / (f (m + n - 1))), x] - \operatorname{Simp}[b^2 ((n-1) / (m + n - 1)) \operatorname{Int}[(a \operatorname{Sec}[e + f x])^m (b \operatorname{Tan}[e + f x])^{(n-2)}, x], x] \text{ ; FreeQ}[\{a, b, e, f, m\}, x] \ \&\& \ \operatorname{GtQ}[n, 1] \ \&\& \ \operatorname{NeQ}[m + n - 1, 0] \ \&\& \ \operatorname{IntegersQ}[2 m, 2 n]$$

rule 3117
$$\operatorname{Int}[\sin[\operatorname{Pi}/2 + (c_)] + (d_)(x_)], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Sin}[c + d x] / d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4257
$$\operatorname{Int}[\operatorname{csc}[(c_)] + (d_)(x_)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d x]] / d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 6155

```
Int[Cosh[v_]*Coth[w_]^(n_), x_Symbol] := Int[Sinh[v]*Coth[w]^(n - 1), x] +
Simp[Cosh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ
[w, v] && FreeQ[v - w, x]
```

rule 6156

```
Int[Coth[w_]^(n_)*Sinh[v_], x_Symbol] := Int[Cosh[v]*Coth[w]^(n - 1), x] +
Simp[Sinh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ
[w, v] && FreeQ[v - w, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. $2(69) = 138$.

Time = 0.18 (sec) , antiderivative size = 230, normalized size of antiderivative = 3.15

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(-3e^{2bx+4a+2c} - e^{2bx+2a+4c} + e^{4a} + 3e^{2a+2c})}{2b(-e^{2bx+2a+2c} + e^{2a})^2} + \frac{3 \ln(e^{bx+a} - e^{a-c})e^{-a-c}e^{2a}}{4b} - \frac{3 \ln(e^{bx+a} - e^{a-c})}{4b}$

input

```
int(coth(b*x+c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
1/2/b*exp(b*x+a)-1/2/b*exp(-b*x-a)+1/2*exp(b*x+a)*(-3*exp(2*b*x+4*a+2*c)-
exp(2*b*x+2*a+4*c)+exp(4*a)+3*exp(2*a+2*c))/b/(-exp(2*b*x+2*a+2*c)+exp(2*a)
)^2+3/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*a)-3/4*ln(exp(b*x+a)-exp
(a-c))/b*exp(-a-c)*exp(2*c)-3/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*
a)+3/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2372 vs. $2(69) = 138$.

Time = 0.11 (sec) , antiderivative size = 2372, normalized size of antiderivative = 32.49

$$\int \coth^3(c + bx) \sinh(a + bx) dx = \text{Too large to display}$$

input

```
integrate(coth(b*x+c)^3*sinh(b*x+a),x, algorithm="fricas")
```

output

```

1/4*(2*cosh(b*x + c)^6*cosh(-a + c)^2 + 2*(cosh(-a + c)^2 - 2*cosh(-a + c)
*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^6 + 12*(cosh(b*x + c)*cosh(-
a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(
-a + c)^2)*sinh(b*x + c)^5 - 2*(5*cosh(-a + c)^2 + 2)*cosh(b*x + c)^4 + 2*
(15*cosh(b*x + c)^2*cosh(-a + c)^2 + 5*(3*cosh(b*x + c)^2 - 1)*sinh(-a + c
)^2 - 5*cosh(-a + c)^2 - 10*(3*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c)
)*sinh(-a + c) - 2)*sinh(b*x + c)^4 + 8*(5*cosh(b*x + c)^3*cosh(-a + c)^2
+ 5*(cosh(b*x + c)^3 - cosh(b*x + c))*sinh(-a + c)^2 - (5*cosh(-a + c)^2 +
2)*cosh(b*x + c) - 10*(cosh(b*x + c)^3*cosh(-a + c) - cosh(b*x + c)*cosh(
-a + c))*sinh(-a + c))*sinh(b*x + c)^3 + 2*(2*cosh(-a + c)^2 + 5)*cosh(b*x
+ c)^2 + 2*(15*cosh(b*x + c)^4*cosh(-a + c)^2 - 6*(5*cosh(-a + c)^2 + 2)*
cosh(b*x + c)^2 + (15*cosh(b*x + c)^4 - 30*cosh(b*x + c)^2 + 2)*sinh(-a +
c)^2 + 2*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^4*cosh(-a + c) - 30*cosh(b*x
+ c)^2*cosh(-a + c) + 2*cosh(-a + c))*sinh(-a + c) + 5)*sinh(b*x + c)^2 +
2*(cosh(b*x + c)^6 - 5*cosh(b*x + c)^4 + 2*cosh(b*x + c)^2)*sinh(-a + c)^
2 - 3*((cosh(-a + c)^2 - 1)*cosh(b*x + c)^5 + (cosh(-a + c)^2 - 2*cosh(-a
+ c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^5 - 5*(2*cosh(b*x +
c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c
)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^4 - 2*(cosh(-a + c)^2 - 1)*cosh(b*x
+ c)^3 + 2*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2...

```

Sympy [F]

$$\int \coth^3(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \coth^3(bx + c) dx$$

input

```
integrate(coth(b*x+c)**3*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*coth(b*x + c)**3, x)
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(69) = 138$.

Time = 0.04 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.55

$$\int \coth^3(c + bx) \sinh(a + bx) dx$$

$$= -\frac{3(e^{2a} - e^{2c})e^{(-a-c)} \log(e^{-bx} + e^c)}{2b} + \frac{3(e^{2a} - e^{2c})e^{(-a-c)} \log(e^{-bx} - e^c)}{2b}$$

$$- \frac{e^{(-bx-a)}}{2b} - \frac{(5e^{(2a+2c)} + e^{(4c)})e^{(-2bx-2a)} - (2e^{(4a)} + 3e^{(2a+2c)})e^{(-4bx-4a)} - e^{(4c)}}{2b(e^{(-bx-a+4c)} - 2e^{(-3bx-a+2c)} + e^{(-5bx-a)})}$$

input `integrate(coth(b*x+c)^3*sinh(b*x+a),x, algorithm="maxima")`

output

```
-3/4*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 3/4*(e^(2*a) -
e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b - 1/2*e^(-b*x - a)/b - 1/2*((5*
e^(2*a + 2*c) + e^(4*c))*e^(-2*b*x - 2*a) - (2*e^(4*a) + 3*e^(2*a + 2*c))*
e^(-4*b*x - 4*a) - e^(4*c))/(b*(e^(-b*x - a + 4*c) - 2*e^(-3*b*x - a + 2*c)
) + e^(-5*b*x - a))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.32

$$\int \coth^3(c + bx) \sinh(a + bx) dx =$$

$$\frac{3(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - 3(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) + \frac{2}{3}}$$

$4b$

input `integrate(coth(b*x+c)^3*sinh(b*x+a),x, algorithm="giac")`

output

```
-1/4*(3*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(e^(b*x + a + c) + e^a) -
3*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + a + c) - e^a)) +
2*(3*e^(3*b*x + 5*a + 2*c) + e^(3*b*x + 3*a + 4*c) - e^(b*x + 5*a) - 3*e^(b
*x + 3*a + 2*c))/(e^(2*b*x + 2*a + 2*c) - e^(2*a))^2 - 2*e^(b*x + a) + 2*e
^(-b*x - a))/b
```

Mupad [F(-1)]

Timed out.

$$\int \coth^3(c + bx) \sinh(a + bx) dx = \int \coth(c + bx)^3 \sinh(a + bx) dx$$

input

```
int(coth(c + b*x)^3*sinh(a + b*x),x)
```

output

```
int(coth(c + b*x)^3*sinh(a + b*x), x)
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.47

$$\int \coth^3(c + bx) \sinh(a + bx) dx$$

$$= \frac{-2e^{bx+a+c} \cosh(bx + a) \coth(bx + c) - 2e^{bx+a+c} \coth(bx + c)^2 \sinh(bx + a) + 3e^{2bx+2a+c} + 3e^{bx+2a} \log(\dots)}{\dots}$$

input

```
int(coth(b*x+c)^3*sinh(b*x+a),x)
```

output

```
( - 2*e**(a + b*x + c)*cosh(a + b*x)*coth(b*x + c) - 2*e**(a + b*x + c)*co
th(b*x + c)**2*sinh(a + b*x) + 3*e**(2*a + 2*b*x + c) + 3*e**(2*a + b*x)*l
og(e**(b*x + c) - 1) - 3*e**(2*a + b*x)*log(e**(b*x + c) + 1) + 2*e**(a +
b*x + c)*sinh(a + b*x) - 3*e**(b*x + 2*c)*log(e**(b*x + c) - 1) + 3*e**(b
*x + 2*c)*log(e**(b*x + c) + 1) - 3*e**c)/(4*e**(a + b*x + c)*b)
```

3.44 $\int \coth(c + dx) \sinh(a + bx) dx$

Optimal result	370
Mathematica [B] (verified)	370
Rubi [A] (verified)	371
Maple [F]	372
Fricas [F]	373
Sympy [F]	373
Maxima [F]	373
Giac [F]	374
Mupad [F(-1)]	374
Reduce [F]	374

Optimal result

Integrand size = 13, antiderivative size = 96

$$\int \coth(c + dx) \sinh(a + bx) dx = \frac{\cosh(a + bx)}{b} - \frac{e^{-a-bx} \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, e^{2(c+dx)}\right)}{b}$$

output

```
cosh(b*x+a)/b-exp(-b*x-a)*hypergeom([1, -1/2*b/d], [1-1/2*b/d], exp(2*d*x+2*c))/b-exp(b*x+a)*hypergeom([1, 1/2*b/d], [1+1/2*b/d], exp(2*d*x+2*c))/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 240 vs. 2(96) = 192.

Time = 2.06 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.50

$$\int \coth(c + dx) \sinh(a + bx) dx = \frac{\cosh(a) \cosh(bx) \coth(c)}{b} + \frac{e^{-a+2c-bx} (b e^{2dx} \operatorname{Hypergeometric2F1}\left(1, 1 - \frac{b}{2d}, 2 - \frac{b}{2d}, e^{2(c+dx)}\right) - (b - 2d) \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 2 - \frac{b}{2d}, e^{2(c+dx)}\right))}{b(b - 2d)(-1 + e^{2c})} - \frac{e^{a+2c} \left(-\frac{e^{(b+2d)x} \operatorname{Hypergeometric2F1}\left(1, 1 + \frac{b}{2d}, 2 + \frac{b}{2d}, e^{2(c+dx)}\right)}{b+2d} + \frac{e^{bx} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, e^{2(c+dx)}\right)}{b} \right)}{-1 + e^{2c}} + \frac{\coth(c) \sinh(a) \sinh(bx)}{b}$$

input `Integrate[Coth[c + d*x]*Sinh[a + b*x],x]`

output `(Cosh[a]*Cosh[b*x]*Coth[c])/b + (E^(-a + 2*c - b*x)*(b*E^(2*d*x)*Hypergeometric2F1[1, 1 - b/(2*d), 2 - b/(2*d), E^(2*(c + d*x))] - (b - 2*d)*Hypergeometric2F1[1, -1/2*b/d, 1 - b/(2*d), E^(2*(c + d*x))])/(b*(b - 2*d)*(-1 + E^(2*c))) - (E^(a + 2*c)*(-(E^((b + 2*d)*x)*Hypergeometric2F1[1, 1 + b/(2*d), 2 + b/(2*d), E^(2*(c + d*x))])/(b + 2*d)) + (E^(b*x)*Hypergeometric2F1[1, b/(2*d), 1 + b/(2*d), E^(2*(c + d*x))])/b)/(-1 + E^(2*c)) + (Coth[c]*Sinh[a]*Sinh[b*x])/b`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.22, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6137, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \coth(c + dx) dx$$

↓ 6137

$$\int \left(\frac{e^{-a-bx}}{1 - e^{2(c+dx)}} - \frac{e^{a+bx}}{1 - e^{2(c+dx)}} - \frac{1}{2} e^{-a-bx} + \frac{1}{2} e^{a+bx} \right) dx$$

$$\begin{array}{c} \downarrow \text{2009} \\ \frac{e^{-a-bx} \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, e^{2(c+dx)}\right)}{b} + \frac{e^{-a-bx}}{2b} + \frac{e^{a+bx}}{2b} \end{array}$$

input `Int[Coth[c + d*x]*Sinh[a + b*x], x]`

output `E^(-a - b*x)/(2*b) + E^(a + b*x)/(2*b) - (E^(-a - b*x)*Hypergeometric2F1[1, -1/2*b/d, 1 - b/(2*d), E^(2*(c + d*x))])/b - (E^(a + b*x)*Hypergeometric2F1[1, b/(2*d), 1 + b/(2*d), E^(2*(c + d*x))])/b`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6137 `Int[Coth[(c_.) + (d_.)*(x_)]*Sinh[(a_.) + (b_.)*(x_)], x_Symbol] := Int[-E^(-(a + b*x))/2 + E^(a + b*x)/2 + 1/(E^(a + b*x)*(1 - E^(2*(c + d*x)))) - E^(a + b*x)/(1 - E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \coth(dx + c) \sinh(bx + a) dx$$

input `int(coth(d*x+c)*sinh(b*x+a), x)`

output `int(coth(d*x+c)*sinh(b*x+a), x)`

Fricas [F]

$$\int \coth(c + dx) \sinh(a + bx) dx = \int \coth(dx + c) \sinh(bx + a) dx$$

input `integrate(coth(d*x+c)*sinh(b*x+a),x, algorithm="fricas")`

output `integral(coth(d*x + c)*sinh(b*x + a), x)`

Sympy [F]

$$\int \coth(c + dx) \sinh(a + bx) dx = \int \sinh(a + bx) \coth(c + dx) dx$$

input `integrate(coth(d*x+c)*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*coth(c + d*x), x)`

Maxima [F]

$$\int \coth(c + dx) \sinh(a + bx) dx = \int \coth(dx + c) \sinh(bx + a) dx$$

input `integrate(coth(d*x+c)*sinh(b*x+a),x, algorithm="maxima")`

output `1/2*(e^(2*b*x + 2*a) + 1)*e^(-b*x - a)/b - 1/2*integrate((e^(2*b*x + 2*a) - 1)/(e^(b*x + d*x + a + c) + e^(b*x + a)), x) + 1/2*integrate((e^(2*b*x + 2*a) - 1)/(e^(b*x + d*x + a + c) - e^(b*x + a)), x)`

Giac [F]

$$\int \coth(c + dx) \sinh(a + bx) dx = \int \coth(dx + c) \sinh(bx + a) dx$$

input `integrate(coth(d*x+c)*sinh(b*x+a),x, algorithm="giac")`

output `integrate(coth(d*x + c)*sinh(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \coth(c + dx) \sinh(a + bx) dx = \int \coth(c + dx) \sinh(a + bx) dx$$

input `int(coth(c + d*x)*sinh(a + b*x),x)`

output `int(coth(c + d*x)*sinh(a + b*x), x)`

Reduce [F]

$$\int \coth(c + dx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+2a} + 2e^{bx+2a} \left(\int \frac{e^{bx}}{e^{2dx+2c}-1} dx \right) b - 2e^{bx} \left(\int \frac{1}{e^{bx+2dx+2c}-e^{bx}} dx \right) b + 1}{2e^{bx+ab}}$$

input `int(coth(d*x+c)*sinh(b*x+a),x)`

output `(e**(2*a + 2*b*x) + 2*e**(2*a + b*x)*int(e**(b*x)/(e**(2*c + 2*d*x) - 1),x)*b - 2*e**(b*x)*int(1/(e**(b*x + 2*c + 2*d*x) - e**(b*x)),x)*b + 1)/(2*e**(a + b*x)*b)`

3.45 $\int \operatorname{sech}(2x) \sinh(x) dx$

Optimal result	375
Mathematica [C] (verified)	375
Rubi [A] (verified)	376
Maple [B] (verified)	377
Fricas [B] (verification not implemented)	378
Sympy [F]	378
Maxima [B] (verification not implemented)	378
Giac [B] (verification not implemented)	379
Mupad [B] (verification not implemented)	379
Reduce [B] (verification not implemented)	380

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \operatorname{sech}(2x) \sinh(x) dx = -\frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{\sqrt{2}}$$

output `-1/2*arctanh(2^(1/2)*cosh(x))*2^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \operatorname{sech}(2x) \sinh(x) dx = -\frac{\operatorname{arctanh}(\sqrt{2} - i \tanh(\frac{x}{2})) + \operatorname{arctanh}(\sqrt{2} + i \tanh(\frac{x}{2}))}{\sqrt{2}}$$

input `Integrate[Sech[2*x]*Sinh[x],x]`

output `-((ArcTanh[Sqrt[2] - I*Tanh[x/2]] + ArcTanh[Sqrt[2] + I*Tanh[x/2]])/Sqrt[2])`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 26, 4857, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{sech}(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{\cos(2ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{\cos(2ix)} dx \\
 & \quad \downarrow \text{4857} \\
 & \int \frac{1}{2 \cosh^2(x) - 1} d \cosh(x) \\
 & \quad \downarrow \text{220} \\
 & -\frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{\sqrt{2}}
 \end{aligned}$$

input `Int [Sech [2*x] * Sinh [x] , x]`

output `-(ArcTanh [Sqrt [2] * Cosh [x]] / Sqrt [2])`

Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4857 `Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(12) = 24$.

Time = 0.20 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.44

method	result	size
risch	$\frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{4} - \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{4}$	39

input `int(sech(2*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `1/4*2^(1/2)*ln(exp(2*x)-2^(1/2)*exp(x)+1)-1/4*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(12) = 24$.

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.19

$$\int \operatorname{sech}(2x) \sinh(x) dx = \frac{1}{4} \sqrt{2} \log \left(\frac{\cosh(x)^2 + \sinh(x)^2 - 2\sqrt{2} \cosh(x) + 2}{\cosh(x)^2 + \sinh(x)^2} \right)$$

input `integrate(sech(2*x)*sinh(x),x, algorithm="fricas")`

output `1/4*sqrt(2)*log((cosh(x)^2 + sinh(x)^2 - 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2 + sinh(x)^2))`

Sympy [F]

$$\int \operatorname{sech}(2x) \sinh(x) dx = \int \sinh(x) \operatorname{sech}(2x) dx$$

input `integrate(sech(2*x)*sinh(x),x)`

output `Integral(sinh(x)*sech(2*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(12) = 24$.

Time = 0.15 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \operatorname{sech}(2x) \sinh(x) dx = -\frac{1}{4} \sqrt{2} \log \left(\sqrt{2} e^{-x} + e^{-2x} + 1 \right) + \frac{1}{4} \sqrt{2} \log \left(-\sqrt{2} e^{-x} + e^{-2x} + 1 \right)$$

input `integrate(sech(2*x)*sinh(x),x, algorithm="maxima")`

output $-1/4*\sqrt{2}*\log(\sqrt{2}*e^{-x} + e^{-2*x} + 1) + 1/4*\sqrt{2}*\log(-\sqrt{2}*e^{-x} + e^{-2*x} + 1)$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(12) = 24.

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.38

$$\int \operatorname{sech}(2x) \sinh(x) dx = -\frac{1}{4} \sqrt{2} \log(\sqrt{2}e^x + e^{2x} + 1) + \frac{1}{4} \sqrt{2} \log(-\sqrt{2}e^x + e^{2x} + 1)$$

input `integrate(sech(2*x)*sinh(x),x, algorithm="giac")`

output $-1/4*\sqrt{2}*\log(\sqrt{2}*e^x + e^{2*x} + 1) + 1/4*\sqrt{2}*\log(-\sqrt{2}*e^x + e^{2*x} + 1)$

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.19

$$\int \operatorname{sech}(2x) \sinh(x) dx = -\frac{\sqrt{2} (\ln(e^{2x} + \sqrt{2}e^x + 1) - \ln(e^{2x} - \sqrt{2}e^x + 1))}{4}$$

input `int(sinh(x)/cosh(2*x),x)`

output $-(2^{(1/2)}*(\log(\exp(2*x) + 2^{(1/2)}*\exp(x) + 1) - \log(\exp(2*x) - 2^{(1/2)}*\exp(x) + 1)))/4$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.25

$$\int \operatorname{sech}(2x) \sinh(x) dx = \frac{\sqrt{2} (\log(e^{2x} - e^x \sqrt{2} + 1) - \log(e^{2x} + e^x \sqrt{2} + 1))}{4}$$

input `int(sech(2*x)*sinh(x),x)`

output `(sqrt(2)*(log(e**(2*x) - e**x*sqrt(2) + 1) - log(e**(2*x) + e**x*sqrt(2) + 1)))/4`

3.46 $\int \operatorname{sech}(3x) \sinh(x) dx$

Optimal result	381
Mathematica [A] (verified)	381
Rubi [A] (verified)	382
Maple [A] (verified)	384
Fricas [B] (verification not implemented)	384
Sympy [F]	385
Maxima [B] (verification not implemented)	385
Giac [B] (verification not implemented)	385
Mupad [B] (verification not implemented)	386
Reduce [B] (verification not implemented)	386

Optimal result

Integrand size = 7, antiderivative size = 21

$$\int \operatorname{sech}(3x) \sinh(x) dx = -\frac{1}{3} \log(\cosh(x)) + \frac{1}{6} \log(3 - 4 \cosh^2(x))$$

output `-1/3*ln(cosh(x))+1/6*ln(3-4*cosh(x)^2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \operatorname{sech}(3x) \sinh(x) dx = -\frac{1}{3} \operatorname{arctanh}\left(\frac{1}{3}(5 + 8 \sinh^2(x))\right)$$

input `Integrate[Sech[3*x]*Sinh[x],x]`

output `-1/3*ArcTanh[(5 + 8*Sinh[x]^2)/3]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 26, 4857, 25, 243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{sech}(3x) dx \\
 & \quad \downarrow 3042 \\
 & \int -\frac{i \sin(ix)}{\cos(3ix)} dx \\
 & \quad \downarrow 26 \\
 & -i \int \frac{\sin(ix)}{\cos(3ix)} dx \\
 & \quad \downarrow 4857 \\
 & \int -\frac{\operatorname{sech}(x)}{3 - 4 \cosh^2(x)} d \cosh(x) \\
 & \quad \downarrow 25 \\
 & - \int \frac{\operatorname{sech}(x)}{3 - 4 \cosh^2(x)} d \cosh(x) \\
 & \quad \downarrow 243 \\
 & -\frac{1}{2} \int \frac{\operatorname{sech}(x)}{3 - 4 \cosh^2(x)} d \cosh^2(x) \\
 & \quad \downarrow 47 \\
 & \frac{1}{2} \left(-\frac{4}{3} \int \frac{1}{3 - 4 \cosh^2(x)} d \cosh^2(x) - \frac{1}{3} \int \operatorname{sech}(x) d \cosh^2(x) \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{2} \left(-\frac{4}{3} \int \frac{1}{3 - 4 \cosh^2(x)} d \cosh^2(x) - \frac{1}{3} \log(\cosh^2(x)) \right) \\
 & \quad \downarrow 16
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{3} \log(3 - 4 \cosh^2(x)) - \frac{1}{3} \log(\cosh^2(x)) \right)$$

input `Int[Sech[3*x]*Sinh[x],x]`

output `(-1/3*Log[Cosh[x]^2] + Log[3 - 4*Cosh[x]^2]/3)/2`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4857

```
Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

method	result	size
risch	$-\frac{\ln(e^{2x}+1)}{3} + \frac{\ln(e^{4x}-e^{2x}+1)}{6}$	26

input

```
int(sech(3*x)*sinh(x),x,method=_RETURNVERBOSE)
```

output

```
-1/3*ln(exp(2*x)+1)+1/6*ln(exp(4*x)-exp(2*x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(17) = 34.

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.48

$$\int \operatorname{sech}(3x) \sinh(x) dx = \frac{1}{6} \log \left(\frac{2 \cosh(x)^2 + 2 \sinh(x)^2 - 1}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2} \right) - \frac{1}{3} \log \left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)} \right)$$

input

```
integrate(sech(3*x)*sinh(x),x, algorithm="fricas")
```

output

```
1/6*log((2*cosh(x)^2 + 2*sinh(x)^2 - 1)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) - 1/3*log(2*cosh(x)/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \operatorname{sech}(3x) \sinh(x) dx = \int \sinh(x) \operatorname{sech}(3x) dx$$

input `integrate(sech(3*x)*sinh(x),x)`

output `Integral(sinh(x)*sech(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(17) = 34$.

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.14

$$\begin{aligned} \int \operatorname{sech}(3x) \sinh(x) dx &= \frac{1}{6} \log \left(\sqrt{3}e^{-x} + e^{-2x} + 1 \right) \\ &\quad + \frac{1}{6} \log \left(-\sqrt{3}e^{-x} + e^{-2x} + 1 \right) - \frac{1}{3} \log \left(e^{-2x} + 1 \right) \end{aligned}$$

input `integrate(sech(3*x)*sinh(x),x, algorithm="maxima")`

output `1/6*log(sqrt(3)*e^(-x) + e^(-2*x) + 1) + 1/6*log(-sqrt(3)*e^(-x) + e^(-2*x) + 1) - 1/3*log(e^(-2*x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(17) = 34$.

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.95

$$\begin{aligned} \int \operatorname{sech}(3x) \sinh(x) dx &= \frac{1}{6} \log \left(\sqrt{3}e^x + e^{2x} + 1 \right) \\ &\quad + \frac{1}{6} \log \left(-\sqrt{3}e^x + e^{2x} + 1 \right) - \frac{1}{3} \log \left(e^{2x} + 1 \right) \end{aligned}$$

input `integrate(sech(3*x)*sinh(x),x, algorithm="giac")`

output $\frac{1}{6} \log(\sqrt{3} e^x + e^{2x} + 1) + \frac{1}{6} \log(-\sqrt{3} e^x + e^{2x} + 1) - \frac{1}{3} \log(e^{2x} + 1)$

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

$$\int \operatorname{sech}(3x) \sinh(x) dx = \frac{\ln(e^{2x} - e^{4x} - 1)}{6} - \frac{\ln(3e^{2x} + 3)}{3}$$

input `int(sinh(x)/cosh(3*x),x)`

output $\log(\exp(2x) - \exp(4x) - 1)/6 - \log(3 \exp(2x) + 3)/3$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.10

$$\int \operatorname{sech}(3x) \sinh(x) dx = \frac{\log(e^{2x} - e^x \sqrt{3} + 1)}{6} + \frac{\log(e^{2x} + e^x \sqrt{3} + 1)}{6} - \frac{\log(e^{2x} + 1)}{3}$$

input `int(sech(3*x)*sinh(x),x)`

output $(\log(e^{2x} - e^x \sqrt{3} + 1) + \log(e^{2x} + e^x \sqrt{3} + 1) - 2 \log(e^{2x} + 1))/6$

3.47 $\int \operatorname{sech}(4x) \sinh(x) dx$

Optimal result	387
Mathematica [C] (verified)	387
Rubi [A] (verified)	388
Maple [C] (verified)	389
Fricas [B] (verification not implemented)	390
Sympy [F]	391
Maxima [F]	391
Giac [B] (verification not implemented)	392
Mupad [B] (verification not implemented)	393
Reduce [B] (verification not implemented)	394

Optimal result

Integrand size = 7, antiderivative size = 71

$$\int \operatorname{sech}(4x) \sinh(x) dx = \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2}(2-\sqrt{2})} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2}(2+\sqrt{2})}$$

output

```
1/2*arctanh(2*cosh(x)/(2-2^(1/2))^(1/2))/(4-2*2^(1/2))^(1/2)-1/2*arctanh(2
*cosh(x)/(2+2^(1/2))^(1/2))/(4+2*2^(1/2))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.03 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.55

$$\int \operatorname{sech}(4x) \sinh(x) dx = \frac{1}{16} \operatorname{RootSum}\left[1 + \#1^8 \&, \frac{-x - 2 \log\left(-\cosh\left(\frac{x}{2}\right) - \sinh\left(\frac{x}{2}\right) + \cosh\left(\frac{x}{2}\right) \#1 - \sinh\left(\frac{x}{2}\right) \#1\right) + x \#1^2 + 2 \log\left(-\cosh\left(\frac{x}{2}\right)\right)}{\#1^5}\right]$$

input

```
Integrate[Sech[4*x]*Sinh[x],x]
```

output

```
RootSum[1 + #1^8 & , (-x - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]**#1 - S
inh[x/2]**#1] + x**#1^2 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]**#1 - Sinh
[x/2]**#1]**#1^2)/#1^5 & ]/16
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4857, 1406, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{sech}(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{\cos(4ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{\cos(4ix)} dx \\
 & \quad \downarrow \text{4857} \\
 & \int \frac{1}{8 \cosh^4(x) - 8 \cosh^2(x) + 1} d \cosh(x) \\
 & \quad \downarrow \text{1406} \\
 & \sqrt{2} \int \frac{1}{8 \cosh^2(x) - 2(2 + \sqrt{2})} d \cosh(x) - \sqrt{2} \int \frac{1}{8 \cosh^2(x) - 2(2 - \sqrt{2})} d \cosh(x) \\
 & \quad \downarrow \text{220} \\
 & \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2}(2-\sqrt{2})} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2}(2+\sqrt{2})}
 \end{aligned}$$

input

```
Int [Sech [4*x] *Sinh [x] , x]
```

output $\text{ArcTanh}[(2*\text{Cosh}[x])/\text{Sqrt}[2 - \text{Sqrt}[2]]]/(2*\text{Sqrt}[2*(2 - \text{Sqrt}[2])]) - \text{ArcTanh}[(2*\text{Cosh}[x])/\text{Sqrt}[2 + \text{Sqrt}[2]]]/(2*\text{Sqrt}[2*(2 + \text{Sqrt}[2])])$

Defintions of rubi rules used

rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_{x_}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 220 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[b, 2])^{-1})*\text{ArcTanh}[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 1406 $\text{Int}[(a_ + (b_)*(x_)^2 + (c_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c/q \ \text{Int}[1/(b/2 - q/2 + c*x^2), x], x] - \text{Simp}[c/q \ \text{Int}[1/(b/2 + q/2 + c*x^2), x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[b^2 - 4*a*c]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4857 $\text{Int}[(u_)*(F_)[(c_)*((a_ + (b_)*(x_)))]], x_Symbol] \rightarrow \text{With}[\{d = \text{FreeFactors}[\text{Cos}[c*(a + b*x)], x]\}, \text{Simp}[-d/(b*c) \ \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Cos}[c*(a + b*x)]/d, u, x], x], x, \text{Cos}[c*(a + b*x)]/d, x] /; \text{FunctionOfQ}[\text{Cos}[c*(a + b*x)]/d, u, x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ (\text{EqQ}[F, \text{Sin}] \ || \ \text{EqQ}[F, \text{sin}])$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.56

method	result	size
risch	$2 \left(\sum_{_R=\text{RootOf}(32768_Z^4-512_Z^2+1)} _R \ln(e^{2x} + (4096_R^3 - 48_R) e^x + 1) \right)$	40

input `int(sech(4*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `2*sum(_R*ln(exp(2*x)+(4096*_R^3-48*_R)*exp(x)+1),_R=RootOf(32768*_Z^4-512*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. $2(49) = 98$.

Time = 0.10 (sec) , antiderivative size = 215, normalized size of antiderivative = 3.03

$$\begin{aligned} \int \operatorname{sech}(4x) \sinh(x) dx = & \frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ & \left. + \left((\sqrt{2} - 1) \cosh(x) + (\sqrt{2} - 1) \sinh(x) \right) \sqrt{\sqrt{2} + 2} + 1 \right) \\ & - \frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ & \left. - \left((\sqrt{2} - 1) \cosh(x) + (\sqrt{2} - 1) \sinh(x) \right) \sqrt{\sqrt{2} + 2} + 1 \right) \\ & - \frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ & \left. + \left((\sqrt{2} + 1) \cosh(x) + (\sqrt{2} + 1) \sinh(x) \right) \sqrt{-\sqrt{2} + 2} + 1 \right) \\ & + \frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ & \left. - \left((\sqrt{2} + 1) \cosh(x) + (\sqrt{2} + 1) \sinh(x) \right) \sqrt{-\sqrt{2} + 2} + 1 \right) \end{aligned}$$

input `integrate(sech(4*x)*sinh(x),x, algorithm="fricas")`

output

```
1/8*sqrt(sqrt(2) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + ((sqrt(2) - 1)*cosh(x) + (sqrt(2) - 1)*sinh(x))*sqrt(sqrt(2) + 2) + 1) - 1/8*sqrt(sqrt(2) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - ((sqrt(2) - 1)*cosh(x) + (sqrt(2) - 1)*sinh(x))*sqrt(sqrt(2) + 2) + 1) - 1/8*sqrt(-sqrt(2) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + ((sqrt(2) + 1)*cosh(x) + (sqrt(2) + 1)*sinh(x))*sqrt(-sqrt(2) + 2) + 1) + 1/8*sqrt(-sqrt(2) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - ((sqrt(2) + 1)*cosh(x) + (sqrt(2) + 1)*sinh(x))*sqrt(-sqrt(2) + 2) + 1)
```

Sympy [F]

$$\int \operatorname{sech}(4x) \sinh(x) dx = \int \sinh(x) \operatorname{sech}(4x) dx$$

input

```
integrate(sech(4*x)*sinh(x),x)
```

output

```
Integral(sinh(x)*sech(4*x), x)
```

Maxima [F]

$$\int \operatorname{sech}(4x) \sinh(x) dx = \int \operatorname{sech}(4x) \sinh(x) dx$$

input

```
integrate(sech(4*x)*sinh(x),x, algorithm="maxima")
```

output

```
integrate(sech(4*x)*sinh(x), x)
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(49) = 98$.

Time = 0.16 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.62

$$\begin{aligned} \int \operatorname{sech}(4x) \sinh(x) dx = & -\frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(\sqrt{\sqrt{2} + 2e^x + e^{(2x)}} + 1 \right) \\ & + \frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(-\sqrt{\sqrt{2} + 2e^x + e^{(2x)}} + 1 \right) \\ & + \frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(\sqrt{-\sqrt{2} + 2e^x + e^{(2x)}} + 1 \right) \\ & - \frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(-\sqrt{-\sqrt{2} + 2e^x + e^{(2x)}} + 1 \right) \end{aligned}$$

input `integrate(sech(4*x)*sinh(x),x, algorithm="giac")`

output `-1/8*sqrt(-sqrt(2) + 2)*log(sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/8*sqrt(-sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/8*sqrt(sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) - 1/8*sqrt(sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 251, normalized size of antiderivative = 3.54

$$\begin{aligned}
\int \operatorname{sech}(4x) \sinh(x) dx = & \ln \left(3e^{2x} - 2\sqrt{2} + 8e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} - 2\sqrt{2}e^{2x} \right. \\
& \left. - 8\sqrt{2}e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} + 3 \right) \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} \\
& - \ln \left(3e^{2x} - 2\sqrt{2} - 8e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} - 2\sqrt{2}e^{2x} \right. \\
& \left. + 8\sqrt{2}e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} + 3 \right) \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}} \\
& - \ln \left(3e^{2x} + 2\sqrt{2} - 8e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}} + 2\sqrt{2}e^{2x} \right. \\
& \left. - 8\sqrt{2}e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}} + 3 \right) \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}} \\
& + \ln \left(3e^{2x} + 2\sqrt{2} + 8e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}} + 2\sqrt{2}e^{2x} \right. \\
& \left. + 8\sqrt{2}e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}} + 3 \right) \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}}
\end{aligned}$$

input `int(sinh(x)/cosh(4*x),x)`

output

```

log(3*exp(2*x) - 2*2^(1/2) + 8*exp(x)*(1/32 - 2^(1/2)/64)^(1/2) - 2*2^(1/2)
)*exp(2*x) - 8*2^(1/2)*exp(x)*(1/32 - 2^(1/2)/64)^(1/2) + 3*(1/32 - 2^(1/2)
/64)^(1/2) - log(3*exp(2*x) - 2*2^(1/2) - 8*exp(x)*(1/32 - 2^(1/2)/64)^(1/2)
- 2*2^(1/2)*exp(2*x) + 8*2^(1/2)*exp(x)*(1/32 - 2^(1/2)/64)^(1/2) + 3
)*(1/32 - 2^(1/2)/64)^(1/2) - log(3*exp(2*x) + 2*2^(1/2) - 8*exp(x)*(2^(1/2)
/64 + 1/32)^(1/2) + 2*2^(1/2)*exp(2*x) - 8*2^(1/2)*exp(x)*(2^(1/2)/64 +
1/32)^(1/2) + 3*(2^(1/2)/64 + 1/32)^(1/2) + log(3*exp(2*x) + 2*2^(1/2) +
8*exp(x)*(2^(1/2)/64 + 1/32)^(1/2) + 2*2^(1/2)*exp(2*x) + 8*2^(1/2)*exp(x)
*(2^(1/2)/64 + 1/32)^(1/2) + 3*(2^(1/2)/64 + 1/32)^(1/2)

```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 221, normalized size of antiderivative = 3.11

$$\begin{aligned}
\int \operatorname{sech}(4x) \sinh(x) dx = & -\frac{\sqrt{-\sqrt{2}+2} \sqrt{2} \log\left(-e^x \sqrt{-\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& + \frac{\sqrt{-\sqrt{2}+2} \sqrt{2} \log\left(e^x \sqrt{-\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& - \frac{\sqrt{-\sqrt{2}+2} \log\left(-e^x \sqrt{-\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& + \frac{\sqrt{-\sqrt{2}+2} \log\left(e^x \sqrt{-\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& + \frac{\sqrt{\sqrt{2}+2} \sqrt{2} \log\left(-e^x \sqrt{\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& - \frac{\sqrt{\sqrt{2}+2} \sqrt{2} \log\left(e^x \sqrt{\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& - \frac{\sqrt{\sqrt{2}+2} \log\left(-e^x \sqrt{\sqrt{2}+2} + e^{2x} + 1\right)}{8} \\
& + \frac{\sqrt{\sqrt{2}+2} \log\left(e^x \sqrt{\sqrt{2}+2} + e^{2x} + 1\right)}{8}
\end{aligned}$$

input

```
int(sech(4*x)*sinh(x),x)
```

output

```
( - sqrt( - sqrt(2) + 2)*sqrt(2)*log( - e**x*sqrt( - sqrt(2) + 2) + e**(2*x) + 1) + sqrt( - sqrt(2) + 2)*sqrt(2)*log(e**x*sqrt( - sqrt(2) + 2) + e**(2*x) + 1) - sqrt( - sqrt(2) + 2)*log( - e**x*sqrt( - sqrt(2) + 2) + e**(2*x) + 1) + sqrt( - sqrt(2) + 2)*log(e**x*sqrt( - sqrt(2) + 2) + e**(2*x) + 1) + sqrt(sqrt(2) + 2)*sqrt(2)*log( - e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) - sqrt(sqrt(2) + 2)*sqrt(2)*log(e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) - sqrt(sqrt(2) + 2)*log( - e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1) + sqrt(sqrt(2) + 2)*log(e**x*sqrt(sqrt(2) + 2) + e**(2*x) + 1))/8
```

3.48 $\int \operatorname{sech}(5x) \sinh(x) dx$

Optimal result	396
Mathematica [A] (verified)	396
Rubi [A] (verified)	397
Maple [A] (verified)	399
Fricas [B] (verification not implemented)	399
Sympy [F]	400
Maxima [F]	400
Giac [B] (verification not implemented)	401
Mupad [B] (verification not implemented)	401
Reduce [F]	402

Optimal result

Integrand size = 7, antiderivative size = 71

$$\int \operatorname{sech}(5x) \sinh(x) dx = \frac{1}{5} \log(\cosh(x)) - \frac{\log(5 - \sqrt{5} - 8 \cosh^2(x))}{\sqrt{5}(5 - \sqrt{5})} + \frac{\log(5 + \sqrt{5} - 8 \cosh^2(x))}{\sqrt{5}(5 + \sqrt{5})}$$

output

```
1/5*ln(cosh(x))-1/5*ln(5-5^(1/2)-8*cosh(x)^2)*5^(1/2)/(5-5^(1/2))+1/5*ln(5+5^(1/2)-8*cosh(x)^2)*5^(1/2)/(5+5^(1/2))
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.80

$$\int \operatorname{sech}(5x) \sinh(x) dx = \frac{1}{20} \left(4 \log(\cosh(x)) + (-1 + \sqrt{5}) \log(3 - \sqrt{5} + 8 \sinh^2(x)) - (1 + \sqrt{5}) \log(3 + \sqrt{5} + 8 \sinh^2(x)) \right)$$

input

```
Integrate[Sech[5*x]*Sinh[x],x]
```

output

$$(4*\text{Log}[\text{Cosh}[x]] + (-1 + \text{Sqrt}[5])* \text{Log}[3 - \text{Sqrt}[5] + 8*\text{Sinh}[x]^2] - (1 + \text{Sqrt}[5])* \text{Log}[3 + \text{Sqrt}[5] + 8*\text{Sinh}[x]^2])/20$$
Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4857, 1434, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{sech}(5x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \sin(ix)}{\cos(5ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\sin(ix)}{\cos(5ix)} dx \\
 & \quad \downarrow \text{4857} \\
 & \int \frac{\operatorname{sech}(x)}{16 \cosh^4(x) - 20 \cosh^2(x) + 5} d \cosh(x) \\
 & \quad \downarrow \text{1434} \\
 & \frac{1}{2} \int \frac{\operatorname{sech}(x)}{16 \cosh^4(x) - 20 \cosh^2(x) + 5} d \cosh^2(x) \\
 & \quad \downarrow \text{1141} \\
 & 8 \int \left(\frac{\operatorname{sech}(x)}{80} + \frac{1}{\sqrt{5}(5-\sqrt{5})(-8 \cosh^2(x) - \sqrt{5} + 5)} - \frac{1}{\sqrt{5}(5+\sqrt{5})(-8 \cosh^2(x) + \sqrt{5} + 5)} \right) d \cosh^2(x) \\
 & \quad \downarrow \text{2009} \\
 & 8 \left(\frac{1}{80} \log(\cosh^2(x)) - \frac{\log(-8 \cosh^2(x) - \sqrt{5} + 5)}{8\sqrt{5}(5-\sqrt{5})} + \frac{\log(-8 \cosh^2(x) + \sqrt{5} + 5)}{8\sqrt{5}(5+\sqrt{5})} \right)
 \end{aligned}$$

input `Int[Sech[5*x]*Sinh[x],x]`

output `8*(Log[Cosh[x]^2]/80 - Log[5 - Sqrt[5] - 8*Cosh[x]^2]/(8*Sqrt[5]*(5 - Sqrt[5])) + Log[5 + Sqrt[5] - 8*Cosh[x]^2]/(8*Sqrt[5]*(5 + Sqrt[5])))`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 1141 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1434 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4857 `Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.42

method	result
risch	$\frac{\ln(e^{2x}+1)}{5} - \frac{\ln(e^{4x} + (-\frac{1}{2} - \frac{\sqrt{5}}{2})e^{2x}+1)}{20} + \frac{\ln(e^{4x} + (-\frac{1}{2} - \frac{\sqrt{5}}{2})e^{2x}+1)\sqrt{5}}{20} - \frac{\ln(e^{4x} + (\frac{\sqrt{5}}{2} - \frac{1}{2})e^{2x}+1)}{20} - \frac{\ln(e^{4x} + (\frac{\sqrt{5}}{2} - \frac{1}{2})e^{2x}+1)\sqrt{5}}{20}$

input `int(sech(5*x)*sinh(x),x,method=_RETURNVERBOSE)`output $1/5*\ln(\exp(2*x)+1)-1/20*\ln(\exp(4*x)+(-1/2-1/2*5^{(1/2)})*\exp(2*x)+1)+1/20*\ln(\exp(4*x)+(-1/2-1/2*5^{(1/2)})*\exp(2*x)+1)*5^{(1/2)}-1/20*\ln(\exp(4*x)+(1/2*5^{(1/2)}-1/2)*\exp(2*x)+1)-1/20*\ln(\exp(4*x)+(1/2*5^{(1/2)}-1/2)*\exp(2*x)+1)*5^{(1/2)}$ **Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(56) = 112.

Time = 0.09 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.56

$$\int \operatorname{sech}(5x) \sinh(x) dx$$

$$= \frac{1}{20} \sqrt{5} \log \left(\frac{4 \cosh(x)^4 + 4 \sinh(x)^4 - 4(\sqrt{5} + 1) \cosh(x)^2 + 4(6 \cosh(x)^2 - \sqrt{5} - 1) \sinh(x)^2 + \sqrt{5}}{2 \cosh(x)^4 + 2 \sinh(x)^4 + 2(6 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 1} \right)$$

$$- \frac{1}{20} \log \left(\frac{2 \cosh(x)^4 + 2 \sinh(x)^4 + 2(6 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 1}{\cosh(x)^4 - 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 - 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4} \right)$$

$$+ \frac{1}{5} \log \left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)} \right)$$

input `integrate(sech(5*x)*sinh(x),x, algorithm="fricas")`

output

```
1/20*sqrt(5)*log((4*cosh(x)^4 + 4*sinh(x)^4 - 4*(sqrt(5) + 1)*cosh(x)^2 +
4*(6*cosh(x)^2 - sqrt(5) - 1)*sinh(x)^2 + sqrt(5) + 7)/(2*cosh(x)^4 + 2*si
nh(x)^4 + 2*(6*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 1)) - 1/20*log((2*
cosh(x)^4 + 2*sinh(x)^4 + 2*(6*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 1)
/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh
(x)^3 + sinh(x)^4)) + 1/5*log(2*cosh(x)/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \operatorname{sech}(5x) \sinh(x) dx = \int \sinh(x) \operatorname{sech}(5x) dx$$

input

```
integrate(sech(5*x)*sinh(x),x)
```

output

```
Integral(sinh(x)*sech(5*x), x)
```

Maxima [F]

$$\int \operatorname{sech}(5x) \sinh(x) dx = \int \operatorname{sech}(5x) \sinh(x) dx$$

input

```
integrate(sech(5*x)*sinh(x),x, algorithm="maxima")
```

output

```
-2/5*integrate((e^(6*x) - e^(4*x) + e^(2*x) - 1)*e^(2*x)/(e^(8*x) - e^(6*x)
) + e^(4*x) - e^(2*x) + 1), x) + 2/5*integrate(e^(6*x)/(e^(8*x) - e^(6*x)
+ e^(4*x) - e^(2*x) + 1), x) + 1/5*integrate(e^(4*x)/(e^(8*x) - e^(6*x) +
e^(4*x) - e^(2*x) + 1), x) - 4/5*integrate(e^(2*x)/(e^(8*x) - e^(6*x) + e^
(4*x) - e^(2*x) + 1), x) + 1/5*log(e^(2*x) + 1)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(56) = 112$.

Time = 0.13 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.66

$$\begin{aligned} \int \operatorname{sech}(5x) \sinh(x) dx &= \frac{1}{20} (\sqrt{5} - 1) \log \left(\frac{1}{2} \sqrt{2\sqrt{5} + 10} e^x + e^{(2x)} + 1 \right) \\ &+ \frac{1}{20} (\sqrt{5} - 1) \log \left(-\frac{1}{2} \sqrt{2\sqrt{5} + 10} e^x + e^{(2x)} + 1 \right) \\ &- \frac{1}{20} (\sqrt{5} + 1) \log \left(\frac{1}{2} \sqrt{-2\sqrt{5} + 10} e^x + e^{(2x)} + 1 \right) \\ &- \frac{1}{20} (\sqrt{5} + 1) \log \left(-\frac{1}{2} \sqrt{-2\sqrt{5} + 10} e^x + e^{(2x)} + 1 \right) \\ &+ \frac{1}{5} \log (e^{(2x)} + 1) \end{aligned}$$

input `integrate(sech(5*x)*sinh(x),x, algorithm="giac")`

output `1/20*(sqrt(5) - 1)*log(1/2*sqrt(2*sqrt(5) + 10)*e^x + e^(2*x) + 1) + 1/20*(sqrt(5) - 1)*log(-1/2*sqrt(2*sqrt(5) + 10)*e^x + e^(2*x) + 1) - 1/20*(sqrt(5) + 1)*log(1/2*sqrt(-2*sqrt(5) + 10)*e^x + e^(2*x) + 1) - 1/20*(sqrt(5) + 1)*log(-1/2*sqrt(-2*sqrt(5) + 10)*e^x + e^(2*x) + 1) + 1/5*log(e^(2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.41

$$\begin{aligned} \int \operatorname{sech}(5x) \sinh(x) dx &= \frac{\ln(5e^{2x} + 5)}{5} - \ln \left(e^{2x} + 2e^{4x} \right. \\ &+ \left. \left(\frac{\sqrt{5}}{20} + \frac{1}{20} \right) (20e^{2x} + 30e^{4x} + 30) + 2 \right) \left(\frac{\sqrt{5}}{20} + \frac{1}{20} \right) \\ &+ \ln \left(e^{2x} + 2e^{4x} - \left(\frac{\sqrt{5}}{20} - \frac{1}{20} \right) (20e^{2x} + 30e^{4x} + 30) \right. \\ &\left. + 2 \right) \left(\frac{\sqrt{5}}{20} - \frac{1}{20} \right) \end{aligned}$$

input `int(sinh(x)/cosh(5*x),x)`

output `log(5*exp(2*x) + 5)/5 - log(exp(2*x) + 2*exp(4*x) + (5^(1/2)/20 + 1/20)*(20*exp(2*x) + 30*exp(4*x) + 30) + 2)*(5^(1/2)/20 + 1/20) + log(exp(2*x) + 2*exp(4*x) - (5^(1/2)/20 - 1/20)*(20*exp(2*x) + 30*exp(4*x) + 30) + 2)*(5^(1/2)/20 - 1/20)`

Reduce [F]

$$\int \operatorname{sech}(5x) \sinh(x) dx = \int \operatorname{sech}(5x) \sinh(x) dx$$

input `int(sech(5*x)*sinh(x),x)`

output `int(sech(5*x)*sinh(x),x)`

3.49 $\int \operatorname{sech}(6x) \sinh(x) dx$

Optimal result	403
Mathematica [C] (verified)	404
Rubi [A] (verified)	404
Maple [C] (verified)	406
Fricas [B] (verification not implemented)	407
Sympy [F]	408
Maxima [F]	408
Giac [B] (verification not implemented)	408
Mupad [B] (verification not implemented)	410
Reduce [B] (verification not implemented)	411

Optimal result

Integrand size = 7, antiderivative size = 85

$$\int \operatorname{sech}(6x) \sinh(x) dx = \frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{3\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2-\sqrt{3}}}\right)}{6\sqrt{2-\sqrt{3}}} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2+\sqrt{3}}}\right)}{6\sqrt{2+\sqrt{3}}}$$

output

```
1/6*arctanh(2^(1/2)*cosh(x))*2^(1/2)-1/6*arctanh(2*cosh(x)/(1/2*6^(1/2)-1/2*2^(1/2)))/(1/2*6^(1/2)-1/2*2^(1/2))-1/6*arctanh(2*cosh(x)/(1/2*6^(1/2)+1/2*2^(1/2)))/(1/2*6^(1/2)+1/2*2^(1/2))
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 269, normalized size of antiderivative = 3.16

$$\int \operatorname{sech}(6x) \sinh(x) dx$$

$$= \frac{1}{24} \left(4\sqrt{2} \left(\operatorname{arctanh} \left(\sqrt{2} - i \tanh \left(\frac{x}{2} \right) \right) + \operatorname{arctanh} \left(\sqrt{2} + i \tanh \left(\frac{x}{2} \right) \right) \right) \right. \\ \left. + \operatorname{RootSum} \left[1 - \#1^4 \right. \right. \\ \left. \left. + \#1^8 \&, \frac{-x - 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) + x \#1^2 + 2 \log \left(-\cosh \left(\frac{x}{2} \right) \right)}{\right. \right]$$

input

```
Integrate[Sech[6*x]*Sinh[x],x]
```

output

```
(4*Sqrt[2]*(ArcTanh[Sqrt[2] - I*Tanh[x/2]] + ArcTanh[Sqrt[2] + I*Tanh[x/2]]
) + RootSum[1 - #1^4 + #1^8 & , (-x - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh
[x/2]*#1 - Sinh[x/2]*#1] + x*#1^2 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[
x/2]*#1 - Sinh[x/2]*#1]*#1^2 - x*#1^4 - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cos
h[x/2]*#1 - Sinh[x/2]*#1]*#1^4 + x*#1^6 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cos
h[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/(-#1^3 + 2*#1^7) & ])/24
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4857, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \operatorname{sech}(6x) dx$$

↓ 3042

$$\begin{aligned}
& \int -\frac{i \sin(ix)}{\cos(6ix)} dx \\
& \quad \downarrow \text{26} \\
& -i \int \frac{\sin(ix)}{\cos(6ix)} dx \\
& \quad \downarrow \text{4857} \\
& \int \frac{1}{32 \cosh^6(x) - 48 \cosh^4(x) + 18 \cosh^2(x) - 1} d \cosh(x) \\
& \quad \downarrow \text{2460} \\
& \int \left(\frac{4(2 \cosh^2(x) - 1)}{3(16 \cosh^4(x) - 16 \cosh^2(x) + 1)} - \frac{1}{3(2 \cosh^2(x) - 1)} \right) d \cosh(x) \\
& \quad \downarrow \text{2009} \\
& \frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{3\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2-\sqrt{3}}}\right)}{6\sqrt{2-\sqrt{3}}} - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{2+\sqrt{3}}}\right)}{6\sqrt{2+\sqrt{3}}}
\end{aligned}$$

input `Int [Sech [6*x]*Sinh [x] ,x]`

output `ArcTanh[Sqrt[2]*Cosh[x]]/(3*Sqrt[2]) - ArcTanh[(2*Cosh[x])/Sqrt[2 - Sqrt[3]]]/(6*Sqrt[2 - Sqrt[3]]) - ArcTanh[(2*Cosh[x])/Sqrt[2 + Sqrt[3]]]/(6*Sqrt[2 + Sqrt[3]])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]},
Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Q
x, x]]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] &&
!TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4857 `Int[(u_.)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFacto
rs[Cos[c*(a + b*x)], x]}, Simp[-d/(b*c) Subst[Int[SubstFor[1, Cos[c*(a +
b*x)]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*
x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.92

method	result
risch	$2 \left(\sum_{R=\text{RootOf}(331776_Z^4-2304_Z^2+1)} -R \ln(e^{2x} + (13824_R^3 - 96_R) e^x + 1) \right) + \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2} e^x + 1)}{12}$

input `int(sech(6*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `2*sum(_R*ln(exp(2*x)+(13824*_R^3-96*_R)*exp(x)+1),_R=RootOf(331776*_Z^4-23
04*_Z^2+1))+1/12*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)-1/12*2^(1/2)*ln(exp
(2*x)-2^(1/2)*exp(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 250 vs. 2(67) = 134.

Time = 0.10 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.94

$$\int \operatorname{sech}(6x) \sinh(x) dx = \frac{1}{12} \sqrt{\sqrt{3} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. + \left((\sqrt{3} - 2) \cosh(x) + (\sqrt{3} - 2) \sinh(x) \right) \sqrt{\sqrt{3} + 2} + 1 \right) \\ - \frac{1}{12} \sqrt{\sqrt{3} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. - \left((\sqrt{3} - 2) \cosh(x) + (\sqrt{3} - 2) \sinh(x) \right) \sqrt{\sqrt{3} + 2} + 1 \right) \\ - \frac{1}{12} \sqrt{-\sqrt{3} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. + \left((\sqrt{3} + 2) \cosh(x) + (\sqrt{3} + 2) \sinh(x) \right) \sqrt{-\sqrt{3} + 2} + 1 \right) \\ + \frac{1}{12} \sqrt{-\sqrt{3} + 2} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. - \left((\sqrt{3} + 2) \cosh(x) + (\sqrt{3} + 2) \sinh(x) \right) \sqrt{-\sqrt{3} + 2} + 1 \right) \\ + \frac{1}{12} \sqrt{2} \log \left(\frac{\cosh(x)^2 + \sinh(x)^2 + 2\sqrt{2} \cosh(x) + 2}{\cosh(x)^2 + \sinh(x)^2} \right)$$

input `integrate(sech(6*x)*sinh(x),x, algorithm="fricas")`

output `1/12*sqrt(sqrt(3) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + ((s
qrt(3) - 2)*cosh(x) + (sqrt(3) - 2)*sinh(x))*sqrt(sqrt(3) + 2) + 1) - 1/12
*sqrt(sqrt(3) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - ((sqrt(
3) - 2)*cosh(x) + (sqrt(3) - 2)*sinh(x))*sqrt(sqrt(3) + 2) + 1) - 1/12*sqr
t(-sqrt(3) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + ((sqrt(3)
+ 2)*cosh(x) + (sqrt(3) + 2)*sinh(x))*sqrt(-sqrt(3) + 2) + 1) + 1/12*sqrt(
-sqrt(3) + 2)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - ((sqrt(3) +
2)*cosh(x) + (sqrt(3) + 2)*sinh(x))*sqrt(-sqrt(3) + 2) + 1) + 1/12*sqrt(2)
*log((cosh(x)^2 + sinh(x)^2 + 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2 + sinh(x)^
2))`

Sympy [F]

$$\int \operatorname{sech}(6x) \sinh(x) dx = \int \sinh(x) \operatorname{sech}(6x) dx$$

input `integrate(sech(6*x)*sinh(x),x)`

output `Integral(sinh(x)*sech(6*x), x)`

Maxima [F]

$$\int \operatorname{sech}(6x) \sinh(x) dx = \int \operatorname{sech}(6x) \sinh(x) dx$$

input `integrate(sech(6*x)*sinh(x),x, algorithm="maxima")`

output `1/12*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) - 1/12*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + integrate(1/3*(e^(7*x) - e^(5*x) + e^(3*x) - e^x)/(e^(8*x) - e^(4*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 154 vs. $2(67) = 134$.

Time = 0.12 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.81

$$\begin{aligned} \int \operatorname{sech}(6x) \sinh(x) dx = & -\frac{1}{24} (\sqrt{6} - \sqrt{2}) \log \left(\frac{1}{2} (\sqrt{6} + \sqrt{2}) e^x + e^{(2x)} + 1 \right) \\ & + \frac{1}{24} (\sqrt{6} - \sqrt{2}) \log \left(-\frac{1}{2} (\sqrt{6} + \sqrt{2}) e^x + e^{(2x)} + 1 \right) \\ & - \frac{1}{24} (\sqrt{6} + \sqrt{2}) \log \left(\frac{1}{2} (\sqrt{6} - \sqrt{2}) e^x + e^{(2x)} + 1 \right) \\ & + \frac{1}{24} (\sqrt{6} + \sqrt{2}) \log \left(-\frac{1}{2} (\sqrt{6} - \sqrt{2}) e^x + e^{(2x)} + 1 \right) \\ & + \frac{1}{12} \sqrt{2} \log (\sqrt{2} e^x + e^{(2x)} + 1) - \frac{1}{12} \sqrt{2} \log (-\sqrt{2} e^x + e^{(2x)} + 1) \end{aligned}$$

input `integrate(sech(6*x)*sinh(x),x, algorithm="giac")`

output
$$\begin{aligned} & -1/24*(\sqrt{6} - \sqrt{2})*\log(1/2*(\sqrt{6} + \sqrt{2})*e^x + e^{2x} + 1) + \\ & 1/24*(\sqrt{6} - \sqrt{2})*\log(-1/2*(\sqrt{6} + \sqrt{2})*e^x + e^{2x} + 1) \\ & - 1/24*(\sqrt{6} + \sqrt{2})*\log(1/2*(\sqrt{6} - \sqrt{2})*e^x + e^{2x} + 1) \\ & + 1/24*(\sqrt{6} + \sqrt{2})*\log(-1/2*(\sqrt{6} - \sqrt{2})*e^x + e^{2x} + 1) \\ & + 1/12*\sqrt{2}*\log(\sqrt{2}*e^x + e^{2x} + 1) - 1/12*\sqrt{2}*\log(-\sqrt{2} \\ & *e^x + e^{2x} + 1) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.44 (sec) , antiderivative size = 288, normalized size of antiderivative = 3.39

$$\begin{aligned}
\int \operatorname{sech}(6x) \sinh(x) dx = & \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{12} - \frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{12} \\
& + \ln \left(7e^{2x} - 4\sqrt{3} - 24e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} - 4\sqrt{3}e^{2x} \right. \\
& \quad \left. + 12\sqrt{3}e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144} + 7} \right) \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} \\
& - \ln \left(7e^{2x} - 4\sqrt{3} + 24e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} - 4\sqrt{3}e^{2x} \right. \\
& \quad \left. - 12\sqrt{3}e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144} + 7} \right) \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} \\
& + \ln \left(7e^{2x} + 4\sqrt{3} - 24e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}} + 4\sqrt{3}e^{2x} \right. \\
& \quad \left. - 12\sqrt{3}e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72} + 7} \right) \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}} \\
& - \ln \left(7e^{2x} + 4\sqrt{3} + 24e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}} + 4\sqrt{3}e^{2x} \right. \\
& \quad \left. + 12\sqrt{3}e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72} + 7} \right) \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}}
\end{aligned}$$

input

```
int(sinh(x)/cosh(6*x),x)
```

output

```
(2^(1/2)*log(exp(2*x) + 2^(1/2)*exp(x) + 1))/12 - (2^(1/2)*log(exp(2*x) -
2^(1/2)*exp(x) + 1))/12 + log(7*exp(2*x) - 4*3^(1/2) - 24*exp(x)*(1/72 - 3
^(1/2)/144)^(1/2) - 4*3^(1/2)*exp(2*x) + 12*3^(1/2)*exp(x)*(1/72 - 3^(1/2)
/144)^(1/2) + 7)*(1/72 - 3^(1/2)/144)^(1/2) - log(7*exp(2*x) - 4*3^(1/2) +
24*exp(x)*(1/72 - 3^(1/2)/144)^(1/2) - 4*3^(1/2)*exp(2*x) - 12*3^(1/2)*ex
p(x)*(1/72 - 3^(1/2)/144)^(1/2) + 7)*(1/72 - 3^(1/2)/144)^(1/2) + log(7*ex
p(2*x) + 4*3^(1/2) - 24*exp(x)*(3^(1/2)/144 + 1/72)^(1/2) + 4*3^(1/2)*exp(
2*x) - 12*3^(1/2)*exp(x)*(3^(1/2)/144 + 1/72)^(1/2) + 7)*(3^(1/2)/144 + 1/
72)^(1/2) - log(7*exp(2*x) + 4*3^(1/2) + 24*exp(x)*(3^(1/2)/144 + 1/72)^(1
/2) + 4*3^(1/2)*exp(2*x) + 12*3^(1/2)*exp(x)*(3^(1/2)/144 + 1/72)^(1/2) +
7)*(3^(1/2)/144 + 1/72)^(1/2)
```

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 260, normalized size of antiderivative = 3.06

$$\int \operatorname{sech}(6x) \sinh(x) dx = \frac{\sqrt{-\sqrt{3}+2} \sqrt{3} \log\left(-e^x \sqrt{-\sqrt{3}+2} + e^{2x} + 1\right)}{12}$$

$$- \frac{\sqrt{-\sqrt{3}+2} \sqrt{3} \log\left(e^x \sqrt{-\sqrt{3}+2} + e^{2x} + 1\right)}{12}$$

$$+ \frac{\sqrt{-\sqrt{3}+2} \log\left(-e^x \sqrt{-\sqrt{3}+2} + e^{2x} + 1\right)}{6}$$

$$- \frac{\sqrt{-\sqrt{3}+2} \log\left(e^x \sqrt{-\sqrt{3}+2} + e^{2x} + 1\right)}{6}$$

$$+ \frac{\sqrt{6} \log\left(e^{2x} - \frac{e^x \sqrt{6}}{2} - \frac{e^x \sqrt{2}}{2} + 1\right)}{24}$$

$$- \frac{\sqrt{6} \log\left(e^{2x} + \frac{e^x \sqrt{6}}{2} + \frac{e^x \sqrt{2}}{2} + 1\right)}{24} - \frac{\sqrt{2} \log\left(e^{2x} - e^x \sqrt{2} + 1\right)}{12}$$

$$+ \frac{\sqrt{2} \log\left(e^{2x} + e^x \sqrt{2} + 1\right)}{12} - \frac{\sqrt{2} \log\left(e^{2x} - \frac{e^x \sqrt{6}}{2} - \frac{e^x \sqrt{2}}{2} + 1\right)}{24}$$

$$+ \frac{\sqrt{2} \log\left(e^{2x} + \frac{e^x \sqrt{6}}{2} + \frac{e^x \sqrt{2}}{2} + 1\right)}{24}$$

input

```
int(sech(6*x)*sinh(x),x)
```

output

```
(2*sqrt(-sqrt(3)+2)*sqrt(3)*log(-e**x*sqrt(-sqrt(3)+2)+e**(2*x)
)+1)-2*sqrt(-sqrt(3)+2)*sqrt(3)*log(e**x*sqrt(-sqrt(3)+2)+e*
*(2*x)+1)+4*sqrt(-sqrt(3)+2)*log(-e**x*sqrt(-sqrt(3)+2)+e*
*(2*x)+1)-4*sqrt(-sqrt(3)+2)*log(e**x*sqrt(-sqrt(3)+2)+e**(2
*x)+1)+sqrt(6)*log((2*e**(2*x)-e**x*sqrt(6)-e**x*sqrt(2)+2)/2)-
sqrt(6)*log((2*e**(2*x)+e**x*sqrt(6)+e**x*sqrt(2)+2)/2)-2*sqrt(2)
*log(e**(2*x)-e**x*sqrt(2)+1)+2*sqrt(2)*log(e**(2*x)+e**x*sqrt(2)
+1)-sqrt(2)*log((2*e**(2*x)-e**x*sqrt(6)-e**x*sqrt(2)+2)/2)+sqr
t(2)*log((2*e**(2*x)+e**x*sqrt(6)+e**x*sqrt(2)+2)/2))/24
```

3.50 $\int \operatorname{sech}(c + bx) \sinh(a + bx) dx$

Optimal result	413
Mathematica [A] (verified)	413
Rubi [A] (verified)	414
Maple [B] (verified)	415
Fricas [B] (verification not implemented)	416
Sympy [F]	416
Maxima [A] (verification not implemented)	416
Giac [A] (verification not implemented)	417
Mupad [B] (verification not implemented)	417
Reduce [B] (verification not implemented)	418

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \frac{\cosh(a - c) \log(\cosh(c + bx))}{b} + x \sinh(a - c)$$

output `cosh(a-c)*ln(cosh(b*x+c))/b+x*sinh(a-c)`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \frac{\cosh(a - c) \log(\cosh(c + bx))}{b} + x \sinh(a - c)$$

input `Integrate[Sech[c + b*x]*Sinh[a + b*x],x]`

output `(Cosh[a - c]*Log[Cosh[c + b*x]])/b + x*Sinh[a - c]`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6158, 24, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{sech}(bx + c) dx \\
 & \quad \downarrow \text{6158} \\
 & \cosh(a - c) \int \tanh(c + bx) dx + \sinh(a - c) \int 1 dx \\
 & \quad \downarrow \text{24} \\
 & \cosh(a - c) \int \tanh(c + bx) dx + x \sinh(a - c) \\
 & \quad \downarrow \text{3042} \\
 & x \sinh(a - c) + \cosh(a - c) \int -i \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & x \sinh(a - c) - i \cosh(a - c) \int \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\cosh(a - c) \log(\cosh(bx + c))}{b} + x \sinh(a - c)
 \end{aligned}$$

input

```
Int[Sech[c + b*x]*Sinh[a + b*x],x]
```

output

```
(Cosh[a - c]*Log[Cosh[c + b*x]])/b + x*Sinh[a - c]
```

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6158 `Int[Sech[w_]^(n_.)*Sinh[v_], x_Symbol] := Simp[Cosh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Sinh[v - w] Int[Sech[w]^(n - 1), x], x] /; GTQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 147 vs. $2(26) = 52$.

Time = 0.30 (sec) , antiderivative size = 148, normalized size of antiderivative = 5.69

method	result
risch	$x e^{a-c} - e^{-a-c} e^{2a} x - e^{-a-c} e^{2c} x - \frac{e^{-a-c} e^{2a} a}{b} - \frac{e^{-a-c} e^{2c} a}{b} + \frac{\ln(e^{2bx+2a} + e^{2a-2c}) e^{-a-c} e^{2a}}{2b} + \frac{\ln(e^{2bx+2a} + e^{2a-2c})}{2b}$

input `int(sech(b*x+c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `x*exp(a-c)-exp(-a-c)*exp(2*a)*x-exp(-a-c)*exp(2*c)*x-1/b*exp(-a-c)*exp(2*a)*a-1/b*exp(-a-c)*exp(2*c)*a+1/2*ln(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-a-c)*exp(2*a)+1/2*ln(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(26) = 52$.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.35

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \frac{2bx - (\cosh(-a + c))^2 - 2\cosh(-a + c)\sinh(-a + c) + \sinh(-a + c)^2 + 1}{2(b\cosh(-a + c) - b\sinh(-a + c))} \log\left(\frac{2\cosh(bx+c)}{\cosh(bx+c) - \sinh(bx+c)}\right)$$

input `integrate(sech(b*x+c)*sinh(b*x+a),x, algorithm="fricas")`

output `-1/2*(2*b*x - (cosh(-a + c))^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c)))/(b*cosh(-a + c) - b*sinh(-a + c))`

Sympy [F]

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}(bx + c) dx$$

input `integrate(sech(b*x+c)*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*sech(b*x + c), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.88

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \frac{(e^{2a} + e^{2c})e^{(-a-c)} \log(e^{-2bx} + e^{2c})}{2b} + \frac{(bx + a)e^{(a-c)}}{b}$$

input `integrate(sech(b*x+c)*sinh(b*x+a),x, algorithm="maxima")`

output $\frac{1}{2}*(e^{(2*a)} + e^{(2*c)})*e^{(-a - c)}*\log(e^{(-2*b*x)} + e^{(2*c)})/b + (b*x + a)*e^{(a - c)}/b$

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.77

$$\int \operatorname{sech}(c+bx) \sinh(a+bx) dx = -xe^{(-a+c)} + \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(2bx+2c)} + 1)}{2b}$$

input `integrate(sech(b*x+c)*sinh(b*x+a),x, algorithm="giac")`

output $-x*e^{(-a + c)} + 1/2*(e^{(2*a + c)} + e^{(3*c)})*e^{(-a - 2*c)}*\log(e^{(2*b*x + 2*c)} + 1)/b$

Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.50

$$\int \operatorname{sech}(c+bx) \sinh(a+bx) dx = \frac{e^{2c-2a} \ln(e^{2a} e^{2bx} + e^{2a} e^{-2c}) (2b e^{3a-3c} + 2b e^{a-c})}{4b^2} - x e^{c-a}$$

input `int(sinh(a + b*x)/cosh(c + b*x),x)`

output $(\exp(2*c - 2*a)*\log(\exp(2*a)*\exp(2*b*x) + \exp(2*a)*\exp(-2*c))*(2*b*\exp(3*a - 3*c) + 2*b*\exp(a - c))/(4*b^2) - x*\exp(c - a)$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.31

$$\int \operatorname{sech}(c + bx) \sinh(a + bx) dx = \frac{e^{2a} \log(e^{2bx+2c} + 1) + e^{2c} \log(e^{2bx+2c} + 1) - 2e^{2c}bx}{2e^{a+cb}}$$

input `int(sech(b*x+c)*sinh(b*x+a),x)`

output `(e**(2*a)*log(e**(2*b*x + 2*c) + 1) + e**(2*c)*log(e**(2*b*x + 2*c) + 1) - 2*e**(2*c)*b*x)/(2*e**(a + c)*b)`

3.51 $\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx$

Optimal result	419
Mathematica [B] (verified)	419
Rubi [A] (verified)	420
Maple [C] (verified)	422
Fricas [B] (verification not implemented)	422
Sympy [F]	423
Maxima [A] (verification not implemented)	423
Giac [A] (verification not implemented)	424
Mupad [B] (verification not implemented)	424
Reduce [B] (verification not implemented)	425

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx = -\frac{\cosh(a - c)\operatorname{sech}(c + bx)}{b} + \frac{\arctan(\sinh(c + bx)) \sinh(a - c)}{b}$$

output

```
-cosh(a-c)*sech(b*x+c)/b+arctan(sinh(b*x+c))*sinh(a-c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 83 vs. 2(35) = 70.

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.37

$$\begin{aligned} & \int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx \\ &= -\frac{\cosh(a - c)\operatorname{sech}(c + bx)}{b} \\ &+ \frac{2 \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right)\right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \sinh(a - c)}{b} \end{aligned}$$

input `Integrate[Sech[c + b*x]^2*Sinh[a + b*x],x]`

output
$$-\left(\frac{\text{Cosh}[a - c] \text{Sech}[c + b*x]}{b}\right) + \frac{(2 \text{ArcTan}[\left(\frac{\text{Cosh}[c] - \text{Sinh}[c]}{\text{Cosh}[c] \text{Cosh}[(b*x)/2] - \text{Cosh}[(b*x)/2] \text{Sinh}[c]}\right)] \text{Sinh}[a - c])}{b}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6158, 3042, 26, 3086, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(a + bx) \operatorname{sech}^2(bx + c) dx \\ & \quad \downarrow 6158 \\ & \sinh(a - c) \int \operatorname{sech}(c + bx) dx + \cosh(a - c) \int \operatorname{sech}(c + bx) \tanh(c + bx) dx \\ & \quad \downarrow 3042 \\ & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx + \cosh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx \\ & \quad \downarrow 26 \\ & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx - i \cosh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx \\ & \quad \downarrow 3086 \\ & -\frac{\cosh(a - c) \int 1 d\operatorname{sech}(c + bx)}{b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\ & \quad \downarrow 24 \\ & -\frac{\cosh(a - c) \operatorname{sech}(bx + c)}{b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\ & \quad \downarrow 4257 \end{aligned}$$

$$\frac{\sinh(a - c) \arctan(\sinh(bx + c))}{b} - \frac{\cosh(a - c) \operatorname{sech}(bx + c)}{b}$$

input `Int[Sech[c + b*x]^2*Sinh[a + b*x],x]`

output `-((Cosh[a - c]*Sech[c + b*x])/b) + (ArcTan[Sinh[c + b*x]]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6158 `Int[Sech[w_]^(n_)*Sinh[v_], x_Symbol] := Simp[Cosh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Sinh[v - w] Int[Sech[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 181, normalized size of antiderivative = 5.17

method	result
risch	$-\frac{e^{bx+a}(e^{2a}+e^{2c})}{b(e^{2bx+2a+2c}+e^{2a})} + \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(sech(b*x+c)^2*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/b*\exp(b*x+a)*(exp(2*a)+exp(2*c))/(exp(2*b*x+2*a+2*c)+exp(2*a))+1/2*I*\ln \\ & (\exp(b*x+a)+I*\exp(a-c))/b*\exp(-a-c)*\exp(a)^2-1/2*I*\ln(\exp(b*x+a)+I*\exp(a-c)) \\ &)/b*\exp(-a-c)*\exp(c)^2-1/2*I*\ln(\exp(b*x+a)-I*\exp(a-c))/b*\exp(-a-c)*\exp(a) \\ & ^2+1/2*I*\ln(\exp(b*x+a)-I*\exp(a-c))/b*\exp(-a-c)*\exp(c)^2 \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(35) = 70$.

Time = 0.11 (sec) , antiderivative size = 405, normalized size of antiderivative = 11.57

$$\int \operatorname{sech}^2(c+bx) \sinh(a+bx) dx$$

$$= \frac{2 \cosh(bx+c) \cosh(-a+c) \sinh(-a+c) - \cosh(bx+c) \sinh(-a+c)^2 + ((\cosh(-a+c))^2 - 1) \cosh(bx+c)}{2b}$$

input `integrate(sech(b*x+c)^2*sinh(b*x+a),x, algorithm="fricas")`

output

```
(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2
+ ((cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)
*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^2 + 1
)*sinh(-a + c)^2 + cosh(-a + c)^2 - 2*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c)
)*sinh(b*x + c) - 2*(cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a
+ c) - 1)*arctan(cosh(b*x + c) + sinh(b*x + c)) - (cosh(-a + c)^2 + 1)*cos
h(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^
2 + 1)*sinh(b*x + c))/(b*cosh(b*x + c)^2*cosh(-a + c) + (b*cosh(-a + c) -
b*sinh(-a + c))*sinh(b*x + c)^2 + b*cosh(-a + c) + 2*(b*cosh(b*x + c)*cosh
(-a + c) - b*cosh(b*x + c)*sinh(-a + c))*sinh(b*x + c) - (b*cosh(b*x + c)^
2 + b)*sinh(-a + c))
```

Sympy [F]

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^2(bx + c) dx$$

input

```
integrate(sech(b*x+c)**2*sinh(b*x+a), x)
```

output

```
Integral(sinh(a + b*x)*sech(b*x + c)**2, x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.00

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx = -\frac{(e^{(2a)} - e^{(2c)}) \arctan(e^{(-bx-c)}) e^{(-a-c)}}{b} - \frac{(e^{(2a)} + e^{(2c)}) e^{(-bx-a)}}{b(e^{(-2bx)} + e^{(2c)})}$$

input

```
integrate(sech(b*x+c)^2*sinh(b*x+a), x, algorithm="maxima")
```

output

```
-(e^(2*a) - e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b - (e^(2*a) + e^(2*c)
))*e^(-b*x - a)/(b*(e^(-2*b*x) + e^(2*c)))
```


Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.00

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx = \frac{(e^{(2a)} - e^{(2c)}) \arctan(e^{(bx+c)}) e^{(-a-c)}}{b} - \frac{(e^{(bx+2a)} + e^{(bx+2c)}) e^{(-a)}}{b(e^{(2bx+2c)} + 1)}$$

input `integrate(sech(b*x+c)^2*sinh(b*x+a),x, algorithm="giac")`output $(e^{(2*a)} - e^{(2*c)}) * \arctan(e^{(b*x + c)}) * e^{(-a - c)} / b - (e^{(b*x + 2*a)} + e^{(b*x + 2*c)}) * e^{(-a)} / (b * (e^{(2*b*x + 2*c)} + 1))$ **Mupad [B] (verification not implemented)**

Time = 1.43 (sec) , antiderivative size = 150, normalized size of antiderivative = 4.29

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx$$

$$= - \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2 - e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{b^2}} - \frac{e^{a+bx} (e^{2a-2c} + 1)}{b (e^{2a-2c} + e^{2a+2bx})}$$

input `int(sinh(a + b*x)/cosh(c + b*x)^2,x)`output $- (\operatorname{atan}((\exp(-a) * \exp(2*c) * \exp(b*x) * ((b^2)^{(1/2)} - \exp(2*a) * \exp(-2*c) * (b^2)^{(1/2)})) / (b * (\exp(-2*a) * \exp(2*c) * (\exp(4*a) * \exp(-4*c) - 2 * \exp(2*a) * \exp(-2*c) + 1))^{(1/2)})) * (\exp(2*c - 2*a) * (\exp(4*a - 4*c) - 2 * \exp(2*a - 2*c) + 1))^{(1/2)}) / (b^2)^{(1/2)} - (\exp(a + b*x) * (\exp(2*a - 2*c) + 1)) / (b * (\exp(2*a - 2*c) + \exp(2*a + 2*b*x)))$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.43

$$\int \operatorname{sech}^2(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+2a+2c} \operatorname{atan}(e^{bx+c}) - e^{2bx+4c} \operatorname{atan}(e^{bx+c}) + e^{2a} \operatorname{atan}(e^{bx+c}) - e^{2c} \operatorname{atan}(e^{bx+c}) - e^{bx+2a+c} - e^{bx+3c}}{e^{a+cb}(e^{2bx+2c} + 1)}$$

input `int(sech(b*x+c)^2*sinh(b*x+a),x)`output `(e**(2*a + 2*b*x + 2*c)*atan(e**(b*x + c)) - e**(2*b*x + 4*c)*atan(e**(b*x + c)) + e**(2*a)*atan(e**(b*x + c)) - e**(2*c)*atan(e**(b*x + c)) - e**(2*a + b*x + c) - e**(b*x + 3*c))/(e**(a + c)*b*(e**(2*b*x + 2*c) + 1))`

3.52 $\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx$

Optimal result	426
Mathematica [A] (verified)	426
Rubi [A] (verified)	427
Maple [A] (verified)	429
Fricas [B] (verification not implemented)	429
Sympy [F]	430
Maxima [B] (verification not implemented)	430
Giac [A] (verification not implemented)	431
Mupad [F(-1)]	431
Reduce [B] (verification not implemented)	431

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = -\frac{\cosh(a - c)\operatorname{sech}^2(c + bx)}{2b} + \frac{\sinh(a - c)\tanh(c + bx)}{b}$$

output

```
-1/2*cosh(a-c)*sech(b*x+c)^2/b+sinh(a-c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\begin{aligned} & \int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx \\ &= -\frac{\operatorname{sech}(c)\operatorname{sech}^2(c + bx)(\cosh(a) - \sinh(a - c)\sinh(c + 2bx))}{2b} \end{aligned}$$

input

```
Integrate[Sech[c + b*x]^3*Sinh[a + b*x],x]
```

output

```
-1/2*(Sech[c]*Sech[c + b*x]^2*(Cosh[a] - Sinh[a - c]*Sinh[c + 2*b*x]))/b
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {6158, 3042, 26, 3086, 15, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{sech}^3(bx + c) dx \\
 & \quad \downarrow \text{6158} \\
 & \sinh(a - c) \int \operatorname{sech}^2(c + bx) dx + \cosh(a - c) \int \operatorname{sech}^2(c + bx) \tanh(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx + \cosh(a - c) \int -i \sec(ic + ibx)^2 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx - i \cosh(a - c) \int \sec(ic + ibx)^2 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3086} \\
 & -\frac{\cosh(a - c) \int \operatorname{sech}(c + bx) d\operatorname{sech}(c + bx)}{b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{\cosh(a - c) \operatorname{sech}^2(bx + c)}{2b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\cosh(a - c) \operatorname{sech}^2(bx + c)}{2b} + \frac{i \sinh(a - c) \int 1 d(-i \tanh(c + bx))}{b} \\
 & \quad \downarrow \text{24} \\
 & \frac{\sinh(a - c) \tanh(bx + c)}{b} - \frac{\cosh(a - c) \operatorname{sech}^2(bx + c)}{2b}
 \end{aligned}$$

input `Int[Sech[c + b*x]^3*Sinh[a + b*x],x]`

output $-1/2*(\text{Cosh}[a - c]*\text{Sech}[c + b*x]^2)/b + (\text{Sinh}[a - c]*\text{Tanh}[c + b*x])/b$

Defintions of rubi rules used

rule 15 $\text{Int}[(a_.)*(x_)^(m_.), x_Symbol] \rightarrow \text{Simp}[a*(x^(m + 1))/(m + 1), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F_x_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \ \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3086 $\text{Int}(((a_.)*\text{sec}[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol) \rightarrow \text{Simp}[a/f \ \text{Subst}[\text{Int}[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2)], x], x, \text{Sec}[e + f*x], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n - 1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n + 1])$

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[-d^(-1) \ \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^(n/2 - 1), x], x], x, \text{Cot}[c + d*x], x] \text{ ; FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 6158 $\text{Int}[\text{Sech}[w_]^(n_.)*\text{Sinh}[v_], x_Symbol] \rightarrow \text{Simp}[\text{Cosh}[v - w] \ \text{Int}[\text{Tanh}[w]*\text{Sech}[w]^(n - 1), x], x] + \text{Simp}[\text{Sinh}[v - w] \ \text{Int}[\text{Sech}[w]^(n - 1), x], x] \text{ ; GtQ}[n, 0] \ \&\& \ \text{NeQ}[w, v] \ \&\& \ \text{FreeQ}[v - w, x]$

Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

method	result	size
parallelrisch	$\frac{2 \cosh(2bx+a+c)-1-\cosh(2bx+2c)}{2b(\cosh(2bx+2c)+1)}$	42
risch	$-\frac{(2e^{2bx+2a+2c}+e^{2a}-e^{2c})e^{3a-c}}{(e^{2bx+2a+2c}+e^{2a})^2 b}$	58

input `int(sech(b*x+c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*(2*cosh(2*b*x+a+c)-1-cosh(2*b*x+2*c))/b/(cosh(2*b*x+2*c)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. 2(36) = 72.

Time = 0.08 (sec) , antiderivative size = 246, normalized size of antiderivative = 6.47

$$\int \operatorname{sech}^3(c+bx) \sinh(a+bx) dx =$$

$$-\frac{b \cosh(bx+c)^3 \cosh(-a+c)^2 + 3b \cosh(bx+c) \cosh(-a+c)^2 + (b \cosh(-a+c)^2 - b \sinh(-a+c)) \sinh(bx+c)^3 + 3(b \cosh(bx+c) \cosh(-a+c)^2 - b \cosh(bx+c) \sinh(-a+c)^2) \sinh(bx+c)^2 - (b \cosh(bx+c)^3 + 3b \cosh(bx+c)) \sinh(-a+c)^2 + (3b \cosh(bx+c)^2 \cosh(-a+c)^2 + b \cosh(-a+c)^2 - (3b \cosh(bx+c)^2 + b) \sinh(-a+c)^2) \sinh(bx+c)}{...}$$

input `integrate(sech(b*x+c)^3*sinh(b*x+a),x, algorithm="fricas")`

output `-2*(cosh(b*x + c)*cosh(-a + c) + cosh(-a + c)*sinh(b*x + c) - 2*cosh(b*x + c)*sinh(-a + c))/(b*cosh(b*x + c)^3*cosh(-a + c)^2 + 3*b*cosh(b*x + c)*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^3 + 3*(b*cosh(b*x + c)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^2 - (b*cosh(b*x + c)^3 + 3*b*cosh(b*x + c))*sinh(-a + c)^2 + (3*b*cosh(b*x + c)^2*cosh(-a + c)^2 + b*cosh(-a + c)^2 - (3*b*cosh(b*x + c)^2 + b)*sinh(-a + c)^2)*sinh(b*x + c)`

Sympy [F]

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^3(bx + c) dx$$

input `integrate(sech(b*x+c)**3*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*sech(b*x + c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. $2(36) = 72$.

Time = 0.05 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.16

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = -\frac{2e^{(-2bx+3c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})} + \frac{e^{(2a+3c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})} - \frac{e^{(5c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})}$$

input `integrate(sech(b*x+c)^3*sinh(b*x+a),x, algorithm="maxima")`

output `-2*e^(-2*b*x + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c))) + e^(2*a + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c))) - e^(5*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c)))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = -\frac{(2e^{(2bx+2a+2c)} + e^{(2a)} - e^{(2c)})e^{(-a-c)}}{b(e^{(2bx+2c)} + 1)^2}$$

input `integrate(sech(b*x+c)^3*sinh(b*x+a),x, algorithm="giac")`

output `-(2*e^(2*b*x + 2*a + 2*c) + e^(2*a) - e^(2*c))*e^(-a - c)/(b*(e^(2*b*x + 2*c) + 1)^2)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\cosh(c + bx)^3} dx$$

input `int(sinh(a + b*x)/cosh(c + b*x)^3,x)`

output `int(sinh(a + b*x)/cosh(c + b*x)^3, x)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int \operatorname{sech}^3(c + bx) \sinh(a + bx) dx = \frac{e^c(e^{4bx+2a+2c} + 1)}{e^{ab}(e^{4bx+4c} + 2e^{2bx+2c} + 1)}$$

input `int(sech(b*x+c)^3*sinh(b*x+a),x)`

output `(e**c*(e**(2*a + 4*b*x + 2*c) + 1))/(e**a*b*(e**(4*b*x + 4*c) + 2*e**(2*b*x + 2*c) + 1))`

3.53 $\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx$

Optimal result	432
Mathematica [A] (verified)	432
Rubi [A] (verified)	433
Maple [C] (verified)	435
Fricas [B] (verification not implemented)	436
Sympy [F]	437
Maxima [B] (verification not implemented)	437
Giac [B] (verification not implemented)	438
Mupad [F(-1)]	438
Reduce [B] (verification not implemented)	439

Optimal result

Integrand size = 15, antiderivative size = 67

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = -\frac{\cosh(a - c)\operatorname{sech}^3(c + bx)}{3b} + \frac{\arctan(\sinh(c + bx)) \sinh(a - c)}{2b} + \frac{\operatorname{sech}(c + bx) \sinh(a - c) \tanh(c + bx)}{2b}$$

output

```
-1/3*cosh(a-c)*sech(b*x+c)^3/b+1/2*arctan(sinh(b*x+c))*sinh(a-c)/b+1/2*sech(b*x+c)*sinh(a-c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.57

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = \frac{-3 \cosh(a - c - bx)\operatorname{sech}(c)\operatorname{sech}^2(c + bx) + 3 \cosh(a - c + bx)\operatorname{sech}(c)\operatorname{sech}^2(c + bx) - 4 \cosh(a - c)\operatorname{sech}^3(c + bx)}{12b}$$

input

```
Integrate[Sech[c + b*x]^4*Sinh[a + b*x],x]
```

output

```
(-3*Cosh[a - c - b*x]*Sech[c]*Sech[c + b*x]^2 + 3*Cosh[a - c + b*x]*Sech[c]*Sech[c + b*x]^2 - 4*Cosh[a - c]*Sech[c + b*x]^3 + 12*ArcTan[Sinh[c] + Cosh[c]*Tanh[(b*x)/2]]*Sinh[a - c] + 6*Sech[c + b*x]*Sinh[a - c]*Tanh[c])/(12*b)
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.94, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {6158, 3042, 26, 3086, 15, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{sech}^4(bx + c) dx \\
 & \quad \downarrow \text{6158} \\
 & \sinh(a - c) \int \operatorname{sech}^3(c + bx) dx + \cosh(a - c) \int \operatorname{sech}^3(c + bx) \tanh(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx + \cosh(a - c) \int -i \sec(ic + ibx)^3 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx - i \cosh(a - c) \int \sec(ic + ibx)^3 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3086} \\
 & -\frac{\cosh(a - c) \int \operatorname{sech}^2(c + bx) d\operatorname{sech}(c + bx)}{b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{\cosh(a - c) \operatorname{sech}^3(bx + c)}{3b} + \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{4255} \\
 & \sinh(a - c) \left(\frac{1}{2} \int \operatorname{sech}(c + bx) dx + \frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} \right) - \frac{\cosh(a - c) \operatorname{sech}^3(bx + c)}{3b}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & -\frac{\cosh(a-c)\operatorname{sech}^3(bx+c)}{3b} + \sinh(a-c) \\ & c) \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} + \frac{1}{2} \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4257 \\ & \sinh(a-c) \left(\frac{\arctan(\sinh(bx+c))}{2b} + \frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} \right) - \frac{\cosh(a-c)\operatorname{sech}^3(bx+c)}{3b} \end{aligned}$$

input `Int[Sech[c + b*x]^4*Sinh[a + b*x],x]`

output `-1/3*(Cosh[a - c]*Sech[c + b*x]^3)/b + Sinh[a - c]*(ArcTan[Sinh[c + b*x]]/(2*b) + (Sech[c + b*x]*Tanh[c + b*x])/(2*b))`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6158 `Int[Sech[w_]^(n_.)*Sinh[v_], x_Symbol] := Simp[Cosh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Sinh[v - w] Int[Sech[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.20 (sec) , antiderivative size = 245, normalized size of antiderivative = 3.66

method	result
risch	$-\frac{e^{bx+a}(-3e^{4bx+6a+4c}+3e^{4bx+4a+6c}+8e^{2bx+6a+2c}+8e^{2bx+4a+4c}+3e^{6a}-3e^{4a+2c})}{6b(e^{2bx+2a+2c}+e^{2a})^3} + \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2a}}{4b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{4b}$

input `int(sech(b*x+c)^4*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `-1/6*exp(b*x+a)*(-3*exp(4*b*x+6*a+4*c)+3*exp(4*b*x+4*a+6*c)+8*exp(2*b*x+6*a+2*c)+8*exp(2*b*x+4*a+4*c)+3*exp(6*a)-3*exp(4*a+2*c))/b/(exp(2*b*x+2*a+2*c)+exp(2*a))^3+1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(a)^2-1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(c)^2-1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(a)^2+1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(c)^2`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1881 vs. $2(61) = 122$.

Time = 0.10 (sec) , antiderivative size = 1881, normalized size of antiderivative = 28.07

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = \text{Too large to display}$$

input `integrate(sech(b*x+c)^4*sinh(b*x+a),x, algorithm="fricas")`

output

```
1/6*(3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^5 + 3*(cosh(-a + c)^2 - 2*cosh(-
a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^5 - 15*(2*cosh(b*x
+ c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a
+ c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^4 - 8*(cosh(-a + c)^2 + 1)*cosh(b
*x + c)^3 + 2*(15*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (15*cosh(b*x + c)
^2 - 4)*sinh(-a + c)^2 - 4*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^2*cosh(-a
+ c) - 4*cosh(-a + c))*sinh(-a + c) - 4)*sinh(b*x + c)^3 + 6*(5*(cosh(-a +
c)^2 - 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 - 4*cosh(b*x + c))*sinh(-a
+ c)^2 - 4*(cosh(-a + c)^2 + 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh
(-a + c) - 4*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^2 + (
3*cosh(b*x + c)^5 - 8*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 +
3*((cosh(-a + c)^2 - 1)*cosh(b*x + c)^6 + (cosh(-a + c)^2 - 2*cosh(-a + c)
*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^6 - 6*(2*cosh(b*x + c)*c
osh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2
- 1)*cosh(b*x + c))*sinh(b*x + c)^5 + 3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)
^4 + 3*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 + 1)*s
inh(-a + c)^2 + cosh(-a + c)^2 - 2*(5*cosh(b*x + c)^2*cosh(-a + c) + cosh(
-a + c))*sinh(-a + c) - 1)*sinh(b*x + c)^4 + 4*(5*(cosh(-a + c)^2 - 1)*cos
h(b*x + c)^3 + (5*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a + c)^2 + 3*(c
osh(-a + c)^2 - 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh(-a + c) + ...
```

Sympy [F]

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `integrate(sech(b*x+c)**4*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*sech(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(61) = 122$.

Time = 0.13 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.31

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = -\frac{(e^{2a} - e^{2c}) \arctan(e^{-bx-c}) e^{(-a-c)}}{2b} + \frac{3(e^{2a+4c} - e^{6c})e^{(-bx-a)} - 8(e^{4a+2c} + e^{2a+4c})e^{(-3bx-3a)} - 3(e^{6a} - e^{4a+2c})e^{(-5bx-5a)}}{6b(3e^{(-2bx+4c)} + 3e^{(-4bx+2c)} + e^{(-6bx)} + e^{6c})}$$

input `integrate(sech(b*x+c)^4*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a) - e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b + 1/6*(3*(e^(2*a + 4*c) - e^(6*c))*e^(-b*x - a) - 8*(e^(4*a + 2*c) + e^(2*a + 4*c))*e^(-3*b*x - 3*a) - 3*(e^(6*a) - e^(4*a + 2*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 4*c) + 3*e^(-4*b*x + 2*c) + e^(-6*b*x) + e^(6*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. $2(61) = 122$.

Time = 0.11 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.87

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = \frac{(e^{(2a)} - e^{(2c)}) \arctan(e^{(bx+c)}) e^{(-a-c)}}{2b} + \frac{(3e^{(5bx+2a+4c)} - 3e^{(5bx+6c)} - 8e^{(3bx+2a+2c)} - 8e^{(3bx+4c)} - 3e^{(bx+2a)} + 3e^{(bx+2c)})e^{(-a)}}{6b(e^{(2bx+2c)} + 1)^3}$$

input `integrate(sech(b*x+c)^4*sinh(b*x+a),x, algorithm="giac")`

output `1/2*(e^(2*a) - e^(2*c))*arctan(e^(b*x + c))*e^(-a - c)/b + 1/6*(3*e^(5*b*x + 2*a + 4*c) - 3*e^(5*b*x + 6*c) - 8*e^(3*b*x + 2*a + 2*c) - 8*e^(3*b*x + 4*c) - 3*e^(b*x + 2*a) + 3*e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) + 1)^3)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\cosh(c + bx)^4} dx$$

input `int(sinh(a + b*x)/cosh(c + b*x)^4,x)`

output `int(sinh(a + b*x)/cosh(c + b*x)^4, x)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 287, normalized size of antiderivative = 4.28

$$\int \operatorname{sech}^4(c + bx) \sinh(a + bx) dx$$

$$= \frac{3e^{6bx+2a+6c} \operatorname{atan}(e^{bx+c}) - 3e^{6bx+8c} \operatorname{atan}(e^{bx+c}) + 9e^{4bx+2a+4c} \operatorname{atan}(e^{bx+c}) - 9e^{4bx+6c} \operatorname{atan}(e^{bx+c}) + 9e^{2bx+2a+2c} \operatorname{atan}(e^{bx+c})}{(6e^{a+c}b(e^{6bx+6c}) + 3e^{4bx+4c}) + 3e^{2bx+2c} + 1}$$

input `int(sech(b*x+c)^4*sinh(b*x+a),x)`

output

```
(3*e**(2*a + 6*b*x + 6*c)*atan(e**(b*x + c)) - 3*e**(6*b*x + 8*c)*atan(e**
(b*x + c)) + 9*e**(2*a + 4*b*x + 4*c)*atan(e**(b*x + c)) - 9*e**(4*b*x + 6
*c)*atan(e**(b*x + c)) + 9*e**(2*a + 2*b*x + 2*c)*atan(e**(b*x + c)) - 9*e
**(2*b*x + 4*c)*atan(e**(b*x + c)) + 3*e**(2*a)*atan(e**(b*x + c)) - 3*e**
(2*c)*atan(e**(b*x + c)) + 3*e**(2*a + 5*b*x + 5*c) - 3*e**(5*b*x + 7*c) -
8*e**(2*a + 3*b*x + 3*c) - 8*e**(3*b*x + 5*c) - 3*e**(2*a + b*x + c) + 3*
e**(b*x + 3*c))/(6*e**(a + c)*b*(e**(6*b*x + 6*c) + 3*e**(4*b*x + 4*c) + 3
*e**(2*b*x + 2*c) + 1))
```


3.54 $\int \operatorname{sech}(c - bx) \sinh(a + bx) dx$

Optimal result	440
Mathematica [A] (verified)	440
Rubi [F]	441
Maple [B] (verified)	441
Fricas [B] (verification not implemented)	442
Sympy [F]	442
Maxima [B] (verification not implemented)	443
Giac [A] (verification not implemented)	443
Mupad [B] (verification not implemented)	444
Reduce [B] (verification not implemented)	444

Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \frac{\cosh(a + c) \log(\cosh(c - bx))}{b} + x \sinh(a + c)$$

output `cosh(a+c)*ln(cosh(b*x-c))/b+x*sinh(a+c)`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \frac{\cosh(a + c) \log(\cosh(c - bx))}{b} + x \sinh(a + c)$$

input `Integrate[Sech[c - b*x]*Sinh[a + b*x],x]`

output `(Cosh[a + c]*Log[Cosh[c - b*x]])/b + x*Sinh[a + c]`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{sech}(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \operatorname{sech}(c - bx) dx$$

input `Int[Sech[c - b*x]*Sinh[a + b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 145 vs. $2(24) = 48$.

Time = 0.18 (sec) , antiderivative size = 146, normalized size of antiderivative = 6.35

method	result
risch	$x e^{a+c} - e^{-a-c} e^{2a+2c} x - \frac{e^{-a-c} e^{2a+2c} a}{b} - x e^{-a-c} - \frac{e^{-a-c} a}{b} + \frac{\ln(e^{2a+2c} + e^{2bx+2a}) e^{-a-c} e^{2a+2c}}{2b} + \frac{\ln(e^{2a+2c} + e^{2bx+2a})}{b}$

input `int(sech(b*x-c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `x*exp(a+c)-exp(-a-c)*exp(2*a+2*c)*x-1/b*exp(-a-c)*exp(2*a+2*c)*a-x*exp(-a-c)-1/b*exp(-a-c)*a+1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)*exp(2*a+2*c)+1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(24) = 48$.

Time = 0.10 (sec) , antiderivative size = 146, normalized size of antiderivative = 6.35

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \frac{2bx \cosh(a + c)^2 - 4bx \cosh(a + c) \sinh(a + c) + 2bx \sinh(a + c)^2 - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log\left(\frac{2(\cosh(bx + a) \cosh(a + c) - \sinh(bx + a) \sinh(a + c))}{(\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c))}\right)}{2(b \cosh(a + c) - b \sinh(a + c))}$$

input `integrate(sech(b*x-c)*sinh(b*x+a),x, algorithm="fricas")`

output `-1/2*(2*b*x*cosh(a + c)^2 - 4*b*x*cosh(a + c)*sinh(a + c) + 2*b*x*sinh(a + c)^2 - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 + 1)*log(2*(cosh(b*x + a)*cosh(a + c) - sinh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c)))/b*cosh(a + c) - b*sinh(a + c))`

Sympy [F]

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}(bx - c) dx$$

input `integrate(sech(b*x-c)*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*sech(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(24) = 48$.

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.13

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx$$

$$= \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-2bx+2c)} + 1)}{2b} + \frac{(bx + a)e^{(a+c)}}{b}$$

input `integrate(sech(b*x-c)*sinh(b*x+a),x, algorithm="maxima")`

output `1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(-2*b*x + 2*c) + 1)/b + (b*x + a)*e^(a + c)/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.00

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = -xe^{(-a-c)} + \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(2bx)} + e^{(2c)})}{2b}$$

input `integrate(sech(b*x-c)*sinh(b*x+a),x, algorithm="giac")`

output `-x*e^(-a - c) + 1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(2*b*x) + e^(2*c))/b`

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.83

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \frac{e^{-2a-2c} \ln(e^{2a} e^{2bx} + e^{2a} e^{2c}) (2b e^{3a+3c} + 2b e^{a+c})}{4b^2 - x e^{-a-c}}$$

input `int(sinh(a + b*x)/cosh(c - b*x),x)`output `(exp(- 2*a - 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(2*c))*(2*b*exp(3*a + 3*c) + 2*b*exp(a + c)))/(4*b^2) - x*exp(- a - c)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.30

$$\int \operatorname{sech}(c - bx) \sinh(a + bx) dx = \frac{e^{2a+2c} \log(e^{2bx} + e^{2c}) + \log(e^{2bx} + e^{2c}) - 2bx}{2e^{a+cb}}$$

input `int(sech(b*x-c)*sinh(b*x+a),x)`output `(e**(2*a + 2*c)*log(e**(2*b*x) + e**(2*c)) + log(e**(2*b*x) + e**(2*c)) - 2*b*x)/(2*e**(a + c)*b)`

3.55 $\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx$

Optimal result	445
Mathematica [B] (verified)	445
Rubi [F]	446
Maple [C] (verified)	446
Fricas [B] (verification not implemented)	447
Sympy [F]	448
Maxima [A] (verification not implemented)	448
Giac [A] (verification not implemented)	448
Mupad [B] (verification not implemented)	449
Reduce [B] (verification not implemented)	449

Optimal result

Integrand size = 16, antiderivative size = 34

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx = -\frac{\cosh(a + c)\operatorname{sech}(c - bx)}{b} - \frac{\arctan(\sinh(c - bx)) \sinh(a + c)}{b}$$

output

```
-cosh(a+c)*sech(b*x-c)/b+arctan(sinh(b*x-c))*sinh(a+c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 81 vs. 2(34) = 68.

Time = 0.07 (sec) , antiderivative size = 81, normalized size of antiderivative = 2.38

$$\begin{aligned} & \int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx \\ &= -\frac{\cosh(a + c)\operatorname{sech}(c - bx)}{b} \\ & \quad + \frac{2 \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(-\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right)\right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \sinh(a + c)}{b} \end{aligned}$$

input `Integrate[Sech[c - b*x]^2*Sinh[a + b*x],x]`

output
$$-\left(\frac{\text{Cosh}[a + c] \text{Sech}[c - b*x]}{b}\right) + \frac{(2 \text{ArcTan}[\left(\frac{\text{Cosh}[c] - \text{Sinh}[c]}{\text{Cosh}[c] \text{Cosh}[(b*x)/2]} - \text{Cosh}[(b*x)/2] \text{Sinh}[c]\right)] + \text{Cosh}[c] \text{Sinh}[(b*x)/2])}{\text{Cosh}[c] \text{Cosh}[(b*x)/2] - \text{Cosh}[(b*x)/2] \text{Sinh}[c]}} \text{Sinh}[a + c]}{b}$$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \text{sech}^2(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \text{sech}^2(c - bx) dx$$

input `Int[Sech[c - b*x]^2*Sinh[a + b*x],x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 171, normalized size of antiderivative = 5.03

method	result
risch	$-\frac{e^{bx+a}(e^{2a+2c}+1)}{b(e^{2a+2c}+e^{2bx+2a})} + \frac{i \ln(e^{bx+a+ie^{a+c}})e^{-a-c}e^{2a+2c}}{2b} - \frac{i \ln(e^{bx+a+ie^{a+c}})e^{-a-c}}{2b} - \frac{i \ln(e^{bx+a-ie^{a+c}})e^{-a-c}e^{2a+2c}}{2b} +$

input `int(sech(b*x-c)^2*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output

```
-1/b*exp(b*x+a)*(exp(2*a+2*c)+1)/(exp(2*a+2*c)+exp(2*b*x+2*a))+1/2*I*ln(ex
p(b*x+a)+I*exp(a+c))/b*exp(-a-c)*exp(a+c)^2-1/2*I*ln(exp(b*x+a)+I*exp(a+c)
)/b*exp(-a-c)-1/2*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)*exp(a+c)^2+1/2*I
*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. $2(35) = 70$.

Time = 0.09 (sec) , antiderivative size = 722, normalized size of antiderivative = 21.24

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x-c)^2*sinh(b*x+a),x, algorithm="fricas")
```

output

```
-(3*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - cosh(b*x + a)*sinh(a + c)^3
- (3*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c) + (4*cosh(b*x + a)^2*cos
h(a + c)*sinh(a + c)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 -
cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)
)^3 + sinh(a + c)^4 + (6*cosh(a + c)^2 - 1)*sinh(a + c)^2 - cosh(a + c)^2
- 2*(2*cosh(a + c)^3 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*cos
h(a + c)^2 - 1)*cosh(b*x + a)^2 + 1)*sinh(a + c)^2 - cosh(a + c)^2 + 2*(4*
cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - (6
*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3 - cos
h(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 - cosh(a + c)^2)*cosh
(b*x + a))*sinh(b*x + a) + 2*((2*cosh(a + c)^3 - cosh(a + c))*cosh(b*x + a)
)^2 + cosh(a + c))*sinh(a + c) + 1)*arctan(-cosh(b*x + a)*cosh(a + c) - (c
osh(a + c) - sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c)) + (co
sh(a + c)^3 + cosh(a + c))*cosh(b*x + a) + (cosh(a + c)^3 + 3*cosh(a + c)*
sinh(a + c)^2 - sinh(a + c)^3 - (3*cosh(a + c)^2 + 1)*sinh(a + c) + cosh(a
+ c))*sinh(b*x + a))/(b*cosh(b*x + a)^2*cosh(a + c)^3 + 3*b*cosh(b*x + a)
^2*cosh(a + c)*sinh(a + c)^2 - b*cosh(b*x + a)^2*sinh(a + c)^3 + (b*cosh(a
+ c)^3 - 3*b*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(a + c)*sinh(a + c)^2 -
b*sinh(a + c)^3)*sinh(b*x + a)^2 + b*cosh(a + c) + 2*(b*cosh(b*x + a)*cosh
(a + c)^3 - 3*b*cosh(b*x + a)*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(b*x ...
```


Sympy [F]

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^2(bx - c) dx$$

input `integrate(sech(b*x-c)**2*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*sech(b*x - c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.03

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx = -\frac{(e^{(2a+2c)} - 1) \arctan(e^{(-bx+c)}) e^{(-a-c)}}{b} - \frac{(e^{(2a+2c)} + 1) e^{(-bx-a)}}{b(e^{(-2bx+2c)} + 1)}$$

input `integrate(sech(b*x-c)^2*sinh(b*x+a),x, algorithm="maxima")`

output `-(e^(2*a + 2*c) - 1)*arctan(e^(-b*x + c))*e^(-a - c)/b - (e^(2*a + 2*c) + 1)*e^(-b*x - a)/(b*(e^(-2*b*x + 2*c) + 1))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.03

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx = \frac{(e^{(2a+2c)} - 1) \arctan(e^{(bx-c)}) e^{(-a-c)}}{b} - \frac{(e^{(bx+2a+2c)} + e^{(bx)}) e^{(-a)}}{b(e^{(2bx)} + e^{(2c)})}$$

input `integrate(sech(b*x-c)^2*sinh(b*x+a),x, algorithm="giac")`

output

$$(e^{(2*a + 2*c)} - 1)*\arctan(e^{(b*x - c)})*e^{(-a - c)}/b - (e^{(b*x + 2*a + 2*c)} + e^{(b*x)})*e^{(-a)}/(b*(e^{(2*b*x)} + e^{(2*c)}))$$

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 150, normalized size of antiderivative = 4.41

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx$$

$$= -\frac{\operatorname{atan}\left(\frac{e^{-a} e^{-2c} e^{bx} (\sqrt{b^2 - e^{2a}} e^{2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{-2c} (e^{4a} e^{4c} - 2e^{2a} e^{2c} + 1)}}\right) \sqrt{e^{-2a-2c} (e^{4a+4c} - 2e^{2a+2c} + 1)}}{\sqrt{b^2}} - \frac{e^{a+bx} (e^{2a+2c} + 1)}{b (e^{2a+2c} + e^{2a+2bx})}$$

input

$$\operatorname{int}(\sinh(a + b*x)/\cosh(c - b*x)^2, x)$$

output

$$- (\operatorname{atan}((\exp(-a)*\exp(-2*c)*\exp(b*x)*((b^2)^{(1/2)} - \exp(2*a)*\exp(2*c)*(b^2)^{(1/2)}))/ (b*(\exp(-2*a)*\exp(-2*c)*(\exp(4*a)*\exp(4*c) - 2*\exp(2*a)*\exp(2*c) + 1))^{(1/2)})) * (\exp(-2*a - 2*c)*(\exp(4*a + 4*c) - 2*\exp(2*a + 2*c) + 1))^{(1/2)} / (b^2)^{(1/2)} - (\exp(a + b*x)*(\exp(2*a + 2*c) + 1)) / (b*(\exp(2*a + 2*c) + \exp(2*a + 2*b*x)))$$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 4.00

$$\int \operatorname{sech}^2(c - bx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{2bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{bx+2a+3c} - e^{bx+c}}{e^{a+cb} (e^{2bx} + e^{2c})}$$

input

$$\operatorname{int}(\operatorname{sech}(b*x-c)^2*\sinh(b*x+a), x)$$

output

```
(e**(2*a + 2*b*x + 2*c)*atan(e**(b*x)/e**c) - e**(2*b*x)*atan(e**(b*x)/e**c) + e**(2*a + 4*c)*atan(e**(b*x)/e**c) - e**(2*c)*atan(e**(b*x)/e**c) - e**(2*a + b*x + 3*c) - e**(b*x + c))/(e**(a + c)*b*(e**(2*b*x) + e**(2*c)))
```

3.56 $\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx$

Optimal result	451
Mathematica [A] (verified)	451
Rubi [F]	452
Maple [A] (verified)	452
Fricas [B] (verification not implemented)	453
Sympy [F]	453
Maxima [B] (verification not implemented)	454
Giac [A] (verification not implemented)	454
Mupad [F(-1)]	455
Reduce [B] (verification not implemented)	455

Optimal result

Integrand size = 16, antiderivative size = 37

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = -\frac{\cosh(a + c)\operatorname{sech}^2(c - bx)}{2b} - \frac{\sinh(a + c) \tanh(c - bx)}{b}$$

output

$$-1/2*\cosh(a+c)*\operatorname{sech}(b*x-c)^2/b+\sinh(a+c)*\tanh(b*x-c)/b$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\begin{aligned} & \int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx \\ &= -\frac{\operatorname{sech}(c)\operatorname{sech}^2(c - bx)(\cosh(a) + \sinh(a + c) \sinh(c - 2bx))}{2b} \end{aligned}$$

input

```
Integrate[Sech[c - b*x]^3*Sinh[a + b*x],x]
```

output

$$-1/2*(\operatorname{Sech}[c]*\operatorname{Sech}[c - b*x]^2*(\operatorname{Cosh}[a] + \operatorname{Sinh}[a + c]*\operatorname{Sinh}[c - 2*b*x]))/b$$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{sech}^3(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \operatorname{sech}^3(c - bx) dx$$

input `Int[Sech[c - b*x]^3*Sinh[a + b*x],x]`

output `$Aborted`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.19

method	result	size
parallelrisch	$\frac{-1 - \cosh(2bx - 2c) + 2 \cosh(2bx + a - c)}{2b(1 + \cosh(2bx - 2c))}$	44
risch	$-\frac{(e^{2a+2c} + 2e^{2bx+2a} - 1)e^{3a+3c}}{(e^{2a+2c} + e^{2bx+2a})^2 b}$	55

input `int(sech(b*x-c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/2/b*(-1-cosh(2*b*x-2*c)+2*cosh(2*b*x+a-c))/(1+cosh(2*b*x-2*c))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 447, normalized size of antiderivative = 12.08

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx =$$

$$\frac{-b \cosh(bx + a)^3 \cosh(a + c)^3 + 3b \cosh(bx + a) \cosh(a + c)^3 + (b \cosh(a + c)^3 - 3b \cosh(a + c)^2 \sinh(a + c)) \sinh(bx + a)}{\dots}$$

input `integrate(sech(b*x-c)^3*sinh(b*x+a),x, algorithm="fricas")`

output

```
-2*(cosh(b*x + a)*cosh(a + c)^2 + cosh(b*x + a)*cosh(a + c)*sinh(a + c) +
(cosh(a + c)^2 - cosh(a + c)*sinh(a + c) - 2*sinh(a + c)^2)*sinh(b*x + a))
/(b*cosh(b*x + a)^3*cosh(a + c)^3 + 3*b*cosh(b*x + a)*cosh(a + c)^3 + (b*c
osh(a + c)^3 - 3*b*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(a + c)*sinh(a + c)
^2 - b*sinh(a + c)^3)*sinh(b*x + a)^3 - (b*cosh(b*x + a)^3 - b*cosh(b*x +
a))*sinh(a + c)^3 + 3*(b*cosh(b*x + a)*cosh(a + c)^3 - 3*b*cosh(b*x + a)*c
osh(a + c)^2*sinh(a + c) + 3*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - b
*cosh(b*x + a)*sinh(a + c)^3)*sinh(b*x + a)^2 + 3*(b*cosh(b*x + a)^3*cosh(
a + c) - b*cosh(b*x + a)*cosh(a + c))*sinh(a + c)^2 + (3*b*cosh(b*x + a)^2
*cosh(a + c)^3 + b*cosh(a + c)^3 - 3*(b*cosh(b*x + a)^2 - b)*sinh(a + c)^3
+ (9*b*cosh(b*x + a)^2*cosh(a + c) - b*cosh(a + c))*sinh(a + c)^2 - 3*(3*
b*cosh(b*x + a)^2*cosh(a + c)^2 + b*cosh(a + c)^2)*sinh(a + c))*sinh(b*x +
a) - (3*b*cosh(b*x + a)^3*cosh(a + c)^2 + b*cosh(b*x + a)*cosh(a + c)^2)*
sinh(a + c))
```

Sympy [F]

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^3(bx - c) dx$$

input `integrate(sech(b*x-c)**3*sinh(b*x+a),x)`

output

`Integral(sinh(a + b*x)*sech(b*x - c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.22

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = -\frac{2e^{(-2bx+2c)}}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})} + \frac{e^{(2a+2c)}}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})} - \frac{1}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})}$$

input `integrate(sech(b*x-c)^3*sinh(b*x+a),x, algorithm="maxima")`

output

```
-2*e^(-2*b*x + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c))) + e^(2*a + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c))) - 1/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c)))
```

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.35

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = -\frac{(2e^{(2bx+2a+3c)} + e^{(2a+5c)} - e^{(3c)})e^{(-a)}}{b(e^{(2bx)} + e^{(2c)})^2}$$

input `integrate(sech(b*x-c)^3*sinh(b*x+a),x, algorithm="giac")`

output

```
-(2*e^(2*b*x + 2*a + 3*c) + e^(2*a + 5*c) - e^(3*c))*e^(-a)/(b*(e^(2*b*x) + e^(2*c))^2)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\cosh(c - bx)^3} dx$$

input `int(sinh(a + b*x)/cosh(c - b*x)^3,x)`output `int(sinh(a + b*x)/cosh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.46

$$\int \operatorname{sech}^3(c - bx) \sinh(a + bx) dx = \frac{e^c (e^{4bx+2a} + e^{2c})}{e^a b (e^{4bx} + 2e^{2bx+2c} + e^{4c})}$$

input `int(sech(b*x-c)^3*sinh(b*x+a),x)`output `(e**c*(e**(2*a + 4*b*x) + e**(2*c)))/(e**a*b*(e**(4*b*x) + 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.57 $\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx$

Optimal result	456
Mathematica [A] (verified)	456
Rubi [F]	457
Maple [C] (verified)	457
Fricas [B] (verification not implemented)	458
Sympy [F]	458
Maxima [B] (verification not implemented)	459
Giac [A] (verification not implemented)	459
Mupad [F(-1)]	460
Reduce [B] (verification not implemented)	460

Optimal result

Integrand size = 16, antiderivative size = 65

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = -\frac{\cosh(a + c)\operatorname{sech}^3(c - bx)}{3b} - \frac{\arctan(\sinh(c - bx)) \sinh(a + c)}{2b} - \frac{\operatorname{sech}(c - bx) \sinh(a + c) \tanh(c - bx)}{2b}$$

output

```
-1/3*cosh(a+c)*sech(b*x-c)^3/b+1/2*arctan(sinh(b*x-c))*sinh(a+c)/b+1/2*sech(b*x-c)*sinh(a+c)*tanh(b*x-c)/b
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.58

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = \frac{-3 \cosh(a + c - bx)\operatorname{sech}(c)\operatorname{sech}^2(c - bx) + 3 \cosh(a + c + bx)\operatorname{sech}(c)\operatorname{sech}^2(c - bx) - 2(2 \cosh(a + c)\operatorname{sech}(c - bx) \arctan(\sinh(c - bx)) \sinh(a + c) + \operatorname{sech}(c - bx) \sinh(a + c) \tanh(c - bx))}{12b}$$

input

```
Integrate[Sech[c - b*x]^4*Sinh[a + b*x],x]
```

output

```
(-3*Cosh[a + c - b*x]*Sech[c]*Sech[c - b*x]^2 + 3*Cosh[a + c + b*x]*Sech[c]*Sech[c - b*x]^2 - 2*(2*Cosh[a + c]*Sech[c - b*x]^3 + 6*ArcTan[Sinh[c] - Cosh[c]*Tanh[(b*x)/2]]*Sinh[a + c] + 3*Sech[c - b*x]*Sinh[a + c]*Tanh[c]))/(12*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx)\operatorname{sech}^4(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx)\operatorname{sech}^4(c - bx) dx$$

input

```
Int[Sech[c - b*x]^4*Sinh[a + b*x],x]
```

output

```
$Aborted
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.19 (sec) , antiderivative size = 235, normalized size of antiderivative = 3.62

method	result
risch	$-\frac{e^{bx+a}(3e^{6a+6c}+8e^{2bx+6a+4c}-3e^{4a+4c}-3e^{4bx+6a+2c}+8e^{2bx+4a+2c}+3e^{4bx+4a})}{6b(e^{2a+2c}+e^{2bx+2a})^3} + \frac{i \ln(e^{bx+a}+ie^{a+c})e^{-a-c}e^{2a+2c}}{4b} - \frac{i \ln(e^{bx+a}-ie^{a+c})e^{-a-c}e^{2a+2c}}{4b}$

input

```
int(sech(b*x-c)^4*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
-1/6*exp(b*x+a)*(3*exp(6*a+6*c)+8*exp(2*b*x+6*a+4*c)-3*exp(4*a+4*c)-3*exp(
4*b*x+6*a+2*c)+8*exp(2*b*x+4*a+2*c)+3*exp(4*b*x+4*a))/b/(exp(2*a+2*c)+exp(
2*b*x+2*a))^3+1/4*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-a-c)*exp(a+c)^2-1/4*I
*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-a-c)-1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*e
xp(-a-c)*exp(a+c)^2+1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6299 vs. $2(63) = 126$.

Time = 0.16 (sec) , antiderivative size = 6299, normalized size of antiderivative = 96.91

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x-c)^4*sinh(b*x+a),x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{sech}^4(bx - c) dx$$

input

```
integrate(sech(b*x-c)**4*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*sech(b*x - c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(63) = 126$.

Time = 0.16 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.34

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = -\frac{(e^{(2a+2c)} - 1) \arctan(e^{(-bx+c)}) e^{(-a-c)}}{2b} + \frac{3(e^{(2a+2c)} - 1)e^{(-bx-a)} - 8(e^{(4a+4c)} + e^{(2a+2c)})e^{(-3bx-3a)} - 3(e^{(6a+6c)} - e^{(4a+4c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+2c)} + 3e^{(-4bx+4c)} + e^{(-6bx+6c)} + 1)}$$

input `integrate(sech(b*x-c)^4*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) - 1)*arctan(e^(-b*x + c))*e^(-a - c)/b + 1/6*(3*(e^(2*a + 2*c) - 1)*e^(-b*x - a) - 8*(e^(4*a + 4*c) + e^(2*a + 2*c))*e^(-3*b*x - 3*a) - 3*(e^(6*a + 6*c) - e^(4*a + 4*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 2*c) + 3*e^(-4*b*x + 4*c) + e^(-6*b*x + 6*c) + 1))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.91

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = \frac{(e^{(2a+2c)} - 1) \arctan(e^{(bx-c)}) e^{(-a-c)}}{2b} - \frac{(3e^{(5bx)} - 3e^{(5bx+2a+2c)} + 8e^{(3bx+2a+4c)} + 8e^{(3bx+2c)} + 3e^{(bx+2a+6c)} - 3e^{(bx+4c)})e^{(-a)}}{6b(e^{(2bx)} + e^{(2c)})^3}$$

input `integrate(sech(b*x-c)^4*sinh(b*x+a),x, algorithm="giac")`

output `1/2*(e^(2*a + 2*c) - 1)*arctan(e^(b*x - c))*e^(-a - c)/b - 1/6*(3*e^(5*b*x) - 3*e^(5*b*x + 2*a + 2*c) + 8*e^(3*b*x + 2*a + 4*c) + 8*e^(3*b*x + 2*c) + 3*e^(b*x + 2*a + 6*c) - 3*e^(b*x + 4*c))*e^(-a)/(b*(e^(2*b*x) + e^(2*c))^3)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\cosh(c - bx)^4} dx$$

input `int(sinh(a + b*x)/cosh(c - b*x)^4,x)`output `int(sinh(a + b*x)/cosh(c - b*x)^4, x)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 319, normalized size of antiderivative = 4.91

$$\int \operatorname{sech}^4(c - bx) \sinh(a + bx) dx$$

$$= \frac{3e^{6bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - 3e^{6bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 9e^{4bx+2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - 9e^{4bx+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 9e^{2bx+2a+6c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right)}{6e^a}$$

input `int(sech(b*x-c)^4*sinh(b*x+a),x)`output `(3***e**(2*a + 6*b*x + 2*c)*atan(e**(b*x)/e**c) - 3***e**(6*b*x)*atan(e**(b*x)/e**c) + 9***e**(2*a + 4*b*x + 4*c)*atan(e**(b*x)/e**c) - 9***e**(4*b*x + 2*c)*atan(e**(b*x)/e**c) + 9***e**(2*a + 2*b*x + 6*c)*atan(e**(b*x)/e**c) - 9***e**(2*b*x + 4*c)*atan(e**(b*x)/e**c) + 3***e**(2*a + 8*c)*atan(e**(b*x)/e**c) - 3***e**(6*c)*atan(e**(b*x)/e**c) + 3***e**(2*a + 5*b*x + 3*c) - 3***e**(5*b*x + c) - 8***e**(2*a + 3*b*x + 5*c) - 8***e**(3*b*x + 3*c) - 3***e**(2*a + b*x + 7*c) + 3***e**(b*x + 5*c))/(6***e**(a + c)*b*(e**(6*b*x) + 3***e**(4*b*x + 2*c) + 3***e**(2*b*x + 4*c) + e**(6*c)))`

3.58 $\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx$

Optimal result	461
Mathematica [A] (verified)	461
Rubi [F]	462
Maple [C] (verified)	462
Fricas [B] (verification not implemented)	463
Sympy [F]	463
Maxima [B] (verification not implemented)	464
Giac [A] (verification not implemented)	464
Mupad [B] (verification not implemented)	465
Reduce [B] (verification not implemented)	465

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \operatorname{sech}(c+bx) \sinh^2(a+bx) dx = -\frac{\arctan(\sinh(c+bx)) \cosh^2(a-c)}{b} + \frac{\sinh(2a-c+bx)}{b}$$

output

```
-arctan(sinh(b*x+c))*cosh(a-c)^2/b+sinh(b*x+2*a-c)/b
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.39

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \frac{\arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) + \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) \cosh(2(a - c)) - \sinh(2a - c + bx)}{b}$$

input

```
Integrate[Sech[c + b*x]*Sinh[a + b*x]^2,x]
```

output

```
-((ArcTan[Tanh[(c + b*x)/2]] + ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(a - c)] - Sinh[2*a - c + b*x])/b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}(bx + c) dx$$

input `Int[Sech[c + b*x]*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 259, normalized size of antiderivative = 7.19

method	result
risch	$\frac{e^{bx+2a-c}}{2b} - \frac{e^{-bx-2a+c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a} e^{4a}}{4b} + \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a} e^{2a} e^{2c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a}}{4b}$

input `int(sech(b*x+c)*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+2*a-c)-1/2/b*exp(-b*x-2*a+c)+1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a)^2+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a)*exp(2*c)+1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*c)^2-1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a)^2-1/2*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a)*exp(2*c)-1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*c)^2`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 630 vs. $2(36) = 72$.

Time = 0.11 (sec) , antiderivative size = 630, normalized size of antiderivative = 17.50

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(sech(b*x+c)*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/2*(cosh(b*x + c)^2*cosh(-a + c)^4 - 4*cosh(b*x + c)^2*cosh(-a + c)^3*sin
h(-a + c) + 6*cosh(b*x + c)^2*cosh(-a + c)^2*sinh(-a + c)^2 - 4*cosh(b*x +
c)^2*cosh(-a + c)*sinh(-a + c)^3 + cosh(b*x + c)^2*sinh(-a + c)^4 + (cosh
(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a + c)^2*sinh(-a + c)
^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4)*sinh(b*x + c)^2 + (4*
cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 -
2*(3*cosh(-a + c)^2 + 1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a + c)^3
+ cosh(-a + c))*cosh(b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 + 2*cosh(-a
+ c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-a + c)^
3 + sinh(-a + c)^4 + 2*(3*cosh(-a + c)^2 + 1)*sinh(-a + c)^2 + 2*cosh(-a +
c)^2 - 4*(cosh(-a + c)^3 + cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c))
*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(cosh(b*x + c)*cosh(-a + c)^4 -
4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(b*x + c)*cosh(-a + c)
)^2*sinh(-a + c)^2 - 4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 + cosh(b*
x + c)*sinh(-a + c)^4)*sinh(b*x + c) - 1)/(b*cosh(b*x + c)*cosh(-a + c)^2
- 2*b*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a +
c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^
2)*sinh(b*x + c))
```

Sympy [F]

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}(bx + c) dx$$

input `integrate(sech(b*x+c)*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(36) = 72$.

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.06

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \frac{(e^{4a} + 2e^{(2a+2c)} + e^{4c}) \arctan(e^{-bx-c}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a-c)}}{2b} - \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(sech(b*x+c)*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))*arctan(e^(-b*x - c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a - c)/b - 1/2*e^(-b*x - 2*a + c)/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.97

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = -\frac{(e^{4a} + 2e^{(2a+2c)} + e^{4c}) \arctan(e^{(bx+c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a-c)}}{2b} - \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(sech(b*x+c)*sinh(b*x+a)^2,x, algorithm="giac")`

output `-1/2*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))*arctan(e^(b*x + c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a - c)/b - 1/2*e^(-b*x - 2*a + c)/b`

Mupad [B] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 194, normalized size of antiderivative = 5.39

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \frac{e^{2a-c+bx}}{2b} - \frac{e^{c-2a-bx}}{2b} - \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{3c} e^{bx} (\sqrt{b^2+2e^{2a} e^{-2c} \sqrt{b^2+e^{4a} e^{-4c} \sqrt{b^2}})}{b \sqrt{e^{-4a} e^{4c} (4e^{2a} e^{-2c} + 6e^{4a} e^{-4c} + 4e^{6a} e^{-6c} + e^{8a} e^{-8c} + 1)}}}\right) \sqrt{e^{4c-4a} (4e^{2a-2c} + 6e^{4a-4c} + 4e^{6a-6c} + e^{8a-8c} + 1)}}{2\sqrt{b^2}}$$

input `int(sinh(a + b*x)^2/cosh(c + b*x),x)`output `exp(2*a - c + b*x)/(2*b) - exp(c - 2*a - b*x)/(2*b) - (atan((exp(-2*a)*exp(3*c)*exp(b*x)*((b^2)^(1/2) + 2*exp(2*a)*exp(-2*c)*(b^2)^(1/2) + exp(4*a)*exp(-4*c)*(b^2)^(1/2)))/(b*(exp(-4*a)*exp(4*c)*(4*exp(2*a)*exp(-2*c) + 6*exp(4*a)*exp(-4*c) + 4*exp(6*a)*exp(-6*c) + exp(8*a)*exp(-8*c) + 1))^(1/2)))*(exp(4*c - 4*a)*(4*exp(2*a - 2*c) + 6*exp(4*a - 4*c) + 4*exp(6*a - 6*c) + exp(8*a - 8*c) + 1))^(1/2))/(2*(b^2)^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.72

$$\int \operatorname{sech}(c + bx) \sinh^2(a + bx) dx = \frac{-e^{bx+4a} \operatorname{atan}(e^{bx+c}) - 2e^{bx+2a+2c} \operatorname{atan}(e^{bx+c}) - e^{bx+4c} \operatorname{atan}(e^{bx+c}) + e^{2bx+4a+c} - e^{3c}}{2e^{bx+2a+2c}b}$$

input `int(sech(b*x+c)*sinh(b*x+a)^2,x)`output `(- e**(4*a + b*x)*atan(e**(b*x + c)) - 2*e**(2*a + b*x + 2*c)*atan(e**(b*x + c)) - e**(b*x + 4*c)*atan(e**(b*x + c)) + e**(4*a + 2*b*x + c) - e**(3*c))/(2*e**(2*a + b*x + 2*c)*b)`

3.59 $\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx$

Optimal result	466
Mathematica [B] (verified)	466
Rubi [F]	467
Maple [B] (verified)	467
Fricas [B] (verification not implemented)	468
Sympy [F]	469
Maxima [A] (verification not implemented)	469
Giac [B] (verification not implemented)	469
Mupad [B] (verification not implemented)	470
Reduce [B] (verification not implemented)	470

Optimal result

Integrand size = 17, antiderivative size = 49

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = x \cosh(2(a - c)) + \frac{\log(\cosh(c + bx)) \sinh(2(a - c))}{b} - \frac{\cosh^2(a - c) \tanh(c + bx)}{b}$$

output `x*cosh(2*a-2*c)+ln(cosh(b*x+c))*sinh(2*a-2*c)/b-cosh(a-c)^2*tanh(b*x+c)/b`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 177 vs. 2(49) = 98.

Time = 0.36 (sec) , antiderivative size = 177, normalized size of antiderivative = 3.61

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx$$

$$= \frac{\operatorname{sech}(c)\operatorname{sech}(c + bx)(bx \cosh(2a - 4c - bx) + bx \cosh(2a - 2c - bx) + bx \cosh(2a + bx) + bx \cosh(2a - 2c + bx))}{b}$$

input `Integrate[Sech[c + b*x]^2*Sinh[a + b*x]^2,x]`

output

```
(Sech[c]*Sech[c + b*x]*(b*x*Cosh[2*a - 4*c - b*x] + b*x*Cosh[2*a - 2*c - b*x] + b*x*Cosh[2*a + b*x] + b*x*Cosh[2*a - 2*c + b*x] - 2*Sinh[b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 4*c - b*x] + Sinh[2*a - 2*c - b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 2*c - b*x] + Log[Cosh[c + b*x]]*Sinh[2*a + b*x] - Sinh[2*a - 2*c + b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 2*c + b*x]))/(4*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

input

```
Int[Sech[c + b*x]^2*Sinh[a + b*x]^2,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(49) = 98$.

Time = 3.60 (sec) , antiderivative size = 247, normalized size of antiderivative = 5.04

method	result
risch	$x e^{2a-2c} - e^{-2c-2a} e^{4a} x + e^{-2c-2a} e^{4c} x - \frac{e^{-2c-2a} e^{4a} a}{b} + \frac{e^{-2c-2a} e^{4c} a}{b} + \frac{e^{-2c} e^{4a}}{2b(e^{2bx+2a+2c} + e^{2a})} + \frac{e^{-2c} e^{2a+2c}}{b(e^{2bx+2a+2c} + e^{2a})}$

input

```
int(sech(b*x+c)^2*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
x*exp(2*a-2*c)-exp(-2*c-2*a)*exp(4*a)*x+exp(-2*c-2*a)*exp(4*c)*x-1/b*exp(-
2*c-2*a)*exp(4*a)*a+1/b*exp(-2*c-2*a)*exp(4*c)*a+1/2/b*exp(-2*c)/(exp(2*b*
x+2*a+2*c)+exp(2*a))*exp(4*a)+1/b*exp(-2*c)/(exp(2*b*x+2*a+2*c)+exp(2*a))*
exp(2*a+2*c)+1/2/b*exp(-2*c)/(exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*c)+1/2*ln
(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*a)-1/2*ln(exp(2*b*x+2*
a)+exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 692 vs. $2(49) = 98$.

Time = 0.10 (sec) , antiderivative size = 692, normalized size of antiderivative = 14.12

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
1/2*(2*b*x*cosh(b*x + c)^2 + cosh(-a + c)^4 + 4*b*x*cosh(b*x + c)*sinh(b*x
+ c) + 2*b*x*sinh(b*x + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a +
c)^4 + 2*(3*cosh(-a + c)^2 + 1)*sinh(-a + c)^2 + 2*b*x + 2*cosh(-a + c)^2
+ ((cosh(b*x + c)^2 + 1)*sinh(-a + c)^4 + cosh(-a + c)^4 - 4*(cosh(b*x + c
)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c)^3 + (cosh(-a + c)^4 - 1)*cos
h(b*x + c)^2 + (cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a
+ c)^2*sinh(-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 -
1)*sinh(b*x + c)^2 + 6*(cosh(b*x + c)^2*cosh(-a + c)^2 + cosh(-a + c)^2)*s
inh(-a + c)^2 - 2*(4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) - 6*cosh(b*
x + c)*cosh(-a + c)^2*sinh(-a + c)^2 + 4*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - (cosh(-a + c)^4 - 1)*cosh(b*x +
c))*sinh(b*x + c) - 4*(cosh(b*x + c)^2*cosh(-a + c)^3 + cosh(-a + c)^3)*si
nh(-a + c) - 1)*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))) - 4*(
cosh(-a + c)^3 + cosh(-a + c))*sinh(-a + c) + 1)/(b*cosh(b*x + c)^2*cosh(-
a + c)^2 + b*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a
+ c) + b*sinh(-a + c)^2)*sinh(b*x + c)^2 + (b*cosh(b*x + c)^2 + b)*sinh(-
a + c)^2 + 2*(b*cosh(b*x + c)*cosh(-a + c)^2 - 2*b*cosh(b*x + c)*cosh(-a +
c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 2*(b*co
sh(b*x + c)^2*cosh(-a + c) + b*cosh(-a + c))*sinh(-a + c))
```

Sympy [F]

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

input `integrate(sech(b*x+c)**2*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x + c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.98

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{-2bx} + e^{2c})}{2b} + \frac{(bx + a)e^{(2a-2c)}}{b} - \frac{e^{4a} + 2e^{(2a+2c)} + e^{4c}}{2b(e^{(-2bx+2a)} + e^{(2a+2c)})}$$

input `integrate(sech(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-2*b*x) + e^(2*c))/b + (b*x + a)*e^(2*a - 2*c)/b - 1/2*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))/(b*(e^(-2*b*x + 2*a) + e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(49) = 98.

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.08

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = xe^{(-2a+2c)} + \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{(2bx+2c)} + 1)}{2b} - \frac{(e^{(2bx+4a)} - e^{(2bx+4c)} - 2e^{(2a)} - 2e^{(2c)})e^{(-2a)}}{2b(e^{(2bx+2c)} + 1)}$$

input `integrate(sech(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="giac")`

output `x*e^(-2*a + 2*c) + 1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(2*b*x + 2*c) + 1)/b - 1/2*(e^(2*b*x + 4*a) - e^(2*b*x + 4*c) - 2*e^(2*a) - 2*e^(2*c))*e^(-2*a)/(b*(e^(2*b*x + 2*c) + 1))`

Mupad [B] (verification not implemented)

Time = 1.29 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.73

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = x e^{2c-2a} + \frac{\sinh(2a - 2c) \ln(e^{2a} e^{2bx} + e^{2a} e^{-2c})}{b} + \frac{2 e^{2a-2c} \cosh(a - c)^2}{b (e^{2a-2c} + e^{2a+2bx})}$$

input `int(sinh(a + b*x)^2/cosh(c + b*x)^2,x)`

output `x*exp(2*c - 2*a) + (sinh(2*a - 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(-2*c)))/b + (2*exp(2*a - 2*c)*cosh(a - c)^2)/(b*(exp(2*a - 2*c) + exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 187, normalized size of antiderivative = 3.82

$$\int \operatorname{sech}^2(c + bx) \sinh^2(a + bx) dx = \frac{e^{2bx+4a+2c} \log(e^{2bx+2c} + 1) - e^{2bx+4a+2c} - 2e^{2bx+2a+4c} - e^{2bx+6c} \log(e^{2bx+2c} + 1) + 2e^{2bx+6c} bx - e^{2bx+6c}}{2e^{2a+2c} b (e^{2bx+2c} + 1)}$$

input `int(sech(b*x+c)^2*sinh(b*x+a)^2,x)`

output

```
(e**(4*a + 2*b*x + 2*c)*log(e**(2*b*x + 2*c) + 1) - e**(4*a + 2*b*x + 2*c)
- 2*e**(2*a + 2*b*x + 4*c) - e**(2*b*x + 6*c)*log(e**(2*b*x + 2*c) + 1) +
2*e**(2*b*x + 6*c)*b*x - e**(2*b*x + 6*c) + e**(4*a)*log(e**(2*b*x + 2*c)
+ 1) - e**(4*c)*log(e**(2*b*x + 2*c) + 1) + 2*e**(4*c)*b*x)/(2*e**(2*a +
2*c)*b*(e**(2*b*x + 2*c) + 1))
```


3.60 $\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx$

Optimal result	472
Mathematica [A] (verified)	472
Rubi [F]	473
Maple [C] (verified)	473
Fricas [B] (verification not implemented)	474
Sympy [F]	475
Maxima [A] (verification not implemented)	476
Giac [A] (verification not implemented)	476
Mupad [F(-1)]	477
Reduce [B] (verification not implemented)	477

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx = -\frac{\arctan(\sinh(c + bx)) \cosh^2(a - c)}{2b} + \frac{\arctan(\sinh(c + bx)) \cosh(2(a - c))}{b} - \frac{\operatorname{sech}(c + bx) \sinh(2(a - c))}{b} - \frac{\cosh^2(a - c) \operatorname{sech}(c + bx) \tanh(c + bx)}{2b}$$

```
output -1/2*arctan(sinh(b*x+c))*cosh(a-c)^2/b+arctan(sinh(b*x+c))*cosh(2*a-2*c)/b
-sech(b*x+c)*sinh(2*a-2*c)/b-1/2*cosh(a-c)^2*sech(b*x+c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.24

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx = \frac{12 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) \cosh(2(a - c)) - \operatorname{sech}^2(c + bx) \left(2 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) + 2 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right)\right)}{8b}$$

input `Integrate[Sech[c + b*x]^3*Sinh[a + b*x]^2,x]`

output `(12*ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(a - c)] - Sech[c + b*x]^2*(2*ArcTan[Tanh[(c + b*x)/2]] + 2*ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(c + b*x)] + 3*Sinh[2*a - 3*c - b*x] + 5*Sinh[2*a - c + b*x] + 2*Sinh[c + b*x]))/(8*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

input `Int[Sech[c + b*x]^3*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 9.08 (sec) , antiderivative size = 321, normalized size of antiderivative = 3.65

method	result
risch	$-\frac{(5e^{2bx+6a+2c}+2e^{2bx+4a+4c}-3e^{2bx+2a+6c}+3e^{6a}-2e^{4a+2c}-5e^{2a+4c})e^{bx-c}}{4(e^{2bx+2a+2c}+e^{2a})^2b} + \frac{3i \ln(e^{bx+a}+ie^{a-c})e^{-2c-2ae^{4a}}}{8b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-2c-2ae^{4a}}}{8b}$

input `int(sech(b*x+c)^3*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*b*x+2*a+2*c)+exp(2*a))^2/b*(5*exp(2*b*x+6*a+2*c)+2*exp(2*b*x+4
*a+4*c)-3*exp(2*b*x+2*a+6*c)+3*exp(6*a)-2*exp(4*a+2*c)-5*exp(2*a+4*c))*exp
(b*x-c)+3/8*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(4*a)-1/4*I*ln(
exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*I*ln(exp(b*x+a)+I*
exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)-3/8*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-
2*c-2*a)*exp(4*a)+1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+
2*c)-3/8*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2187 vs. $2(84) = 168$.

Time = 0.10 (sec) , antiderivative size = 2187, normalized size of antiderivative = 24.85

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
-1/4*((5*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a + c)^4 + (5*cosh(-a +
c)^4 + 2*cosh(-a + c)^2 - 3)*cosh(b*x + c)^3 + (5*cosh(-a + c)^4 - 20*cosh
(-a + c)*sinh(-a + c)^3 + 5*sinh(-a + c)^4 + 2*(15*cosh(-a + c)^2 + 1)*sin
h(-a + c)^2 + 2*cosh(-a + c)^2 - 4*(5*cosh(-a + c)^3 + cosh(-a + c))*sinh(
-a + c) - 3)*sinh(b*x + c)^3 - 4*(5*cosh(b*x + c)^3*cosh(-a + c) + 3*cosh(
b*x + c)*cosh(-a + c))*sinh(-a + c)^3 - 3*(20*cosh(b*x + c)*cosh(-a + c)*s
inh(-a + c)^3 - 5*cosh(b*x + c)*sinh(-a + c)^4 - 2*(15*cosh(-a + c)^2 + 1)
*cosh(b*x + c)*sinh(-a + c)^2 + 4*(5*cosh(-a + c)^3 + cosh(-a + c))*cosh(b
*x + c)*sinh(-a + c) - (5*cosh(-a + c)^4 + 2*cosh(-a + c)^2 - 3)*cosh(b*x
+ c))*sinh(b*x + c)^2 + 2*((15*cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + (9*c
osh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(-a + c)^2 - ((3*cosh(-a + c)^4 - 2*c
osh(-a + c)^2 + 3)*cosh(b*x + c)^4 + (3*cosh(-a + c)^4 - 12*cosh(-a + c)*s
inh(-a + c)^3 + 3*sinh(-a + c)^4 + 2*(9*cosh(-a + c)^2 - 1)*sinh(-a + c)^2
- 2*cosh(-a + c)^2 - 4*(3*cosh(-a + c)^3 - cosh(-a + c))*sinh(-a + c) + 3
)*sinh(b*x + c)^4 + 3*(cosh(b*x + c)^4 + 2*cosh(b*x + c)^2 + 1)*sinh(-a +
c)^4 + 3*cosh(-a + c)^4 - 4*(12*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3
- 3*cosh(b*x + c)*sinh(-a + c)^4 - 2*(9*cosh(-a + c)^2 - 1)*cosh(b*x + c)*
sinh(-a + c)^2 + 4*(3*cosh(-a + c)^3 - cosh(-a + c))*cosh(b*x + c)*sinh(-a
+ c) - (3*cosh(-a + c)^4 - 2*cosh(-a + c)^2 + 3)*cosh(b*x + c))*sinh(b*x
+ c)^3 - 12*(cosh(b*x + c)^4*cosh(-a + c) + 2*cosh(b*x + c)^2*cosh(-a + ...
```

Sympy [F]

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

input

```
integrate(sech(b*x+c)**3*sinh(b*x+a)**2,x)
```

output

```
Integral(sinh(a + b*x)**2*sech(b*x + c)**3, x)
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.77

$$\int \operatorname{sech}^3(c+bx) \sinh^2(a+bx) dx = -\frac{(3e^{4a} - 2e^{(2a+2c)} + 3e^{4c}) \arctan(e^{(-bx-c)}) e^{(-2a-2c)}}{4b} - \frac{(5e^{(4a+2c)} + 2e^{(2a+4c)} - 3e^{(6c)})e^{(-bx-a)} + (3e^{(6a)} - 2e^{(4a+2c)} - 5e^{(2a+4c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+c)} + e^{(a+5c)})}$$

input `integrate(sech(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="maxima")`output `-1/4*(3*e^(4*a) - 2*e^(2*a + 2*c) + 3*e^(4*c))*arctan(e^(-b*x - c))*e^(-2*a - 2*c)/b - 1/4*((5*e^(4*a + 2*c) + 2*e^(2*a + 4*c) - 3*e^(6*c))*e^(-b*x - a) + (3*e^(6*a) - 2*e^(4*a + 2*c) - 5*e^(2*a + 4*c))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + c) + e^(a + 5*c)))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.62

$$\int \operatorname{sech}^3(c+bx) \sinh^2(a+bx) dx = \frac{(3e^{4a} - 2e^{(2a+2c)} + 3e^{4c}) \arctan(e^{(bx+c)}) e^{(-2a-2c)}}{4b} - \frac{(5e^{(3bx+4a+2c)} + 2e^{(3bx+2a+4c)} - 3e^{(3bx+6c)} + 3e^{(bx+4a)} - 2e^{(bx+2a+2c)} - 5e^{(bx+4c)})e^{(-2a-c)}}{4b(e^{(2bx+2c)} + 1)^2}$$

input `integrate(sech(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="giac")`output `1/4*(3*e^(4*a) - 2*e^(2*a + 2*c) + 3*e^(4*c))*arctan(e^(b*x + c))*e^(-2*a - 2*c)/b - 1/4*(5*e^(3*b*x + 4*a + 2*c) + 2*e^(3*b*x + 2*a + 4*c) - 3*e^(3*b*x + 6*c) + 3*e^(b*x + 4*a) - 2*e^(b*x + 2*a + 2*c) - 5*e^(b*x + 4*c))*e^(-2*a - c)/(b*(e^(2*b*x + 2*c) + 1)^2)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\cosh(c + bx)^3} dx$$

input `int(sinh(a + b*x)^2/cosh(c + b*x)^3,x)`output `int(sinh(a + b*x)^2/cosh(c + b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 303, normalized size of antiderivative = 3.44

$$\int \operatorname{sech}^3(c + bx) \sinh^2(a + bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \operatorname{atan}(e^{bx+c}) - 2e^{4bx+2a+6c} \operatorname{atan}(e^{bx+c}) + 3e^{4bx+8c} \operatorname{atan}(e^{bx+c}) + 6e^{2bx+4a+2c} \operatorname{atan}(e^{bx+c}) - 4e^{2bx+4a+2c} \operatorname{atan}(e^{bx+c})}{(4e^{2bx+4a+2c} + 2e^{2bx+4a+2c} + 1)}$$

input `int(sech(b*x+c)^3*sinh(b*x+a)^2,x)`output `(3*e**(4*a + 4*b*x + 4*c)*atan(e**(b*x + c)) - 2*e**(2*a + 4*b*x + 6*c)*atan(e**(b*x + c)) + 3*e**(4*b*x + 8*c)*atan(e**(b*x + c)) + 6*e**(4*a + 2*b*x + 2*c)*atan(e**(b*x + c)) - 4*e**(2*a + 2*b*x + 4*c)*atan(e**(b*x + c)) + 6*e**(2*b*x + 6*c)*atan(e**(b*x + c)) + 3*e**(4*a)*atan(e**(b*x + c)) - 2*e**(2*a + 2*c)*atan(e**(b*x + c)) + 3*e**(4*c)*atan(e**(b*x + c)) - 5*e**(4*a + 3*b*x + 3*c) - 2*e**(2*a + 3*b*x + 5*c) + 3*e**(3*b*x + 7*c) - 3*e**(4*a + b*x + c) + 2*e**(2*a + b*x + 3*c) + 5*e**(b*x + 5*c))/(4*e**(2*a + 2*c)*b*(e**(4*b*x + 4*c) + 2*e**(2*b*x + 2*c) + 1))`

3.61 $\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx$

Optimal result	478
Mathematica [A] (verified)	478
Rubi [F]	479
Maple [A] (verified)	479
Fricas [B] (verification not implemented)	480
Sympy [F]	481
Maxima [B] (verification not implemented)	481
Giac [A] (verification not implemented)	482
Mupad [F(-1)]	482
Reduce [B] (verification not implemented)	483

Optimal result

Integrand size = 17, antiderivative size = 84

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = -\frac{\operatorname{sech}^2(c + bx) \sinh(2(a - c))}{2b} - \frac{\cosh^2(a - c) \tanh(c + bx)}{b} + \frac{\cosh(2(a - c)) \tanh(c + bx)}{b} + \frac{\cosh^2(a - c) \tanh^3(c + bx)}{3b}$$

```
output -1/2*sech(b*x+c)^2*sinh(2*a-2*c)/b-cosh(a-c)^2*tanh(b*x+c)/b+cosh(2*a-2*c)*tanh(b*x+c)/b+1/3*cosh(a-c)^2*tanh(b*x+c)^3/b
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.95

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}^3(c + bx)(3 \sinh(bx) + \sinh(2a - 4c - 3bx)) + 3 \sinh(2a - 2c - bx) + 3 \sinh(2a + bx) - \sinh(2a - 2c)}{12b}$$

input `Integrate[Sech[c + b*x]^4*Sinh[a + b*x]^2,x]`

output `-1/12*(Sech[c]*Sech[c + b*x]^3*(3*Sinh[b*x] + Sinh[2*a - 4*c - 3*b*x] + 3*Sinh[2*a - 2*c - b*x] + 3*Sinh[2*a + b*x] - Sinh[2*a + 3*b*x] + Sinh[2*c + 3*b*x]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `Int[Sech[c + b*x]^4*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [A] (verified)

Time = 6.34 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.69

method	result	size
parallelrisch	$\frac{2 \sinh(3bx+2a+c) - 3 \sinh(bx+c) - \sinh(3bx+3c)}{3b(\cosh(3bx+3c) + 3 \cosh(bx+c))}$	58
risch	$-\frac{2(3e^{4bx+4a+4c} + 3e^{2bx+4a+2c} - 3e^{2bx+2a+4c} + e^{4a} - e^{2a+2c} + e^{4c})e^{4a-2c}}{3(e^{2bx+2a+2c} + e^{2a})^3 b}$	94

input `int(sech(b*x+c)^4*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{3} \frac{(2 \sinh(3bx+2a+c) - 3 \sinh(bx+c) - \sinh(3bx+3c))}{b (\cosh(3bx+3c) + 3 \cosh(bx+c))}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(80) = 160$.

Time = 0.09 (sec) , antiderivative size = 493, normalized size of antiderivative = 5.87

$$\int \operatorname{sech}^4(c+bx) \sinh^2(a+bx) dx = \frac{2 \left((5 \cosh(-a+c))^2 - 1 \right) \cosh(bx+c) \sinh(bx+c)}{3 (b \cosh(bx+c))^4 \cosh(-a+c)^2 + 4b \cosh(bx+c)^2 \cosh(-a+c)^2 + (b \cosh(-a+c))^2 - b \sinh(-a+c)}$$

input `integrate(sech(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{-2/3 \left((5 \cosh(-a+c))^2 - 1 \right) \cosh(bx+c)^2 + (5 \cosh(-a+c))^2 - 6 \cosh(-a+c) \sinh(-a+c) + 5 \sinh(-a+c)^2 - 1 \right) \sinh(bx+c)^2 + (5 \cosh(bx+c)^2 + 3) \sinh(-a+c)^2 + 3 \cosh(-a+c)^2 - 2 \left(6 \cosh(bx+c) \cosh(-a+c) \sinh(-a+c) - \cosh(bx+c) \sinh(-a+c)^2 - (\cosh(-a+c))^2 + 1 \right) \cosh(bx+c) \sinh(bx+c) - 6 \left(\cosh(bx+c)^2 \cosh(-a+c) + \cosh(-a+c) \right) \sinh(-a+c) - 3}{(b \cosh(bx+c))^4 \cosh(-a+c)^2 + 4b \cosh(bx+c)^2 \cosh(-a+c)^2 + (b \cosh(-a+c))^2 - b \sinh(-a+c)^2} \sinh(bx+c)^4 + 4 \left(b \cosh(bx+c) \cosh(-a+c)^2 - b \cosh(bx+c) \sinh(-a+c)^2 \right) \sinh(bx+c)^3 + 3b \cosh(-a+c)^2 + 2 \left(3b \cosh(bx+c)^2 \cosh(-a+c)^2 + 2b \cosh(-a+c)^2 - (3b \cosh(bx+c)^2 + 2b) \sinh(-a+c)^2 \right) \sinh(bx+c)^2 - (b \cosh(bx+c))^4 + 4b \cosh(bx+c)^2 + 3b \sinh(-a+c)^2 + 4 \left(b \cosh(bx+c)^3 \cosh(-a+c)^2 + b \cosh(bx+c) \cosh(-a+c)^2 - (b \cosh(bx+c))^3 + b \cosh(bx+c) \right) \sinh(-a+c)^2 \sinh(bx+c)}$$

Sympy [F]

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `integrate(sech(b*x+c)**4*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. $2(80) = 160$.

Time = 0.05 (sec) , antiderivative size = 323, normalized size of antiderivative = 3.85

$$\begin{aligned} & \int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx \\ &= -\frac{2(e^{(4a+4c)} - e^{(2a+6c)})e^{(-2bx-2a)}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &- \frac{2e^{(2a+6c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(8c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \end{aligned}$$

input `integrate(sech(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned}
& -2*(e^{(4*a + 4*c)} - e^{(2*a + 6*c)})*e^{(-2*b*x - 2*a)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} + e^{(2*a + 6*c)}))} + 2 \\
& *e^{(-4*b*x + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} + e^{(2*a + 6*c)}))} + 2/3*e^{(4*a + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} + e^{(2*a + 6*c)}))} \\
& - 2/3*e^{(2*a + 6*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} + e^{(2*a + 6*c)}))} + 2/3*e^{(8*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} + e^{(2*a + 6*c)}))}
\end{aligned}$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.04

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = \frac{2(3e^{(4bx+4a+4c)} + 3e^{(2bx+4a+2c)} - 3e^{(2bx+2a+4c)} + e^{(4a)} - e^{(2a+2c)} + e^{(4c)})e^{(-2a-2c)}}{3b(e^{(2bx+2c)} + 1)^3}$$

input

```
integrate(sech(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="giac")
```

output

$$-2/3*(3*e^{(4*b*x + 4*a + 4*c)} + 3*e^{(2*b*x + 4*a + 2*c)} - 3*e^{(2*b*x + 2*a + 4*c)} + e^{(4*a)} - e^{(2*a + 2*c)} + e^{(4*c)})*e^{(-2*a - 2*c)/(b*(e^{(2*b*x + 2*c)} + 1)^3)}$$
Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\cosh(c + bx)^4} dx$$

input

```
int(sinh(a + b*x)^2/cosh(c + b*x)^4,x)
```

output

```
int(sinh(a + b*x)^2/cosh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.08

$$\int \operatorname{sech}^4(c + bx) \sinh^2(a + bx) dx = \frac{\frac{2e^{6bx+4a+4c}}{3} + 2e^{2bx+2a+2c} + \frac{2e^{2a}}{3} - \frac{2e^{2c}}{3}}{e^{2a}b(e^{6bx+6c} + 3e^{4bx+4c} + 3e^{2bx+2c} + 1)}$$

input `int(sech(b*x+c)^4*sinh(b*x+a)^2,x)`output `(2*(e**(4*a + 6*b*x + 4*c) + 3*e**(2*a + 2*b*x + 2*c) + e**(2*a) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x + 6*c) + 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) + 1))`

3.62 $\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx$

Optimal result	484
Mathematica [A] (verified)	484
Rubi [F]	485
Maple [C] (verified)	485
Fricas [B] (verification not implemented)	486
Sympy [F]	486
Maxima [B] (verification not implemented)	487
Giac [B] (verification not implemented)	487
Mupad [B] (verification not implemented)	488
Reduce [B] (verification not implemented)	488

Optimal result

Integrand size = 16, antiderivative size = 32

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \frac{\arctan(\sinh(c - bx)) \cosh^2(a + c)}{b} + \frac{\sinh(2a + c + bx)}{b}$$

output

```
-arctan(sinh(b*x-c))*cosh(a+c)^2/b+sinh(b*x+2*a+c)/b
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.41

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \frac{\arctan(\tanh(\frac{1}{2}(c - bx))) + \arctan(\tanh(\frac{1}{2}(c - bx))) \cosh(2(a + c)) + \sinh(2a + c + bx)}{b}$$

input

```
Integrate[Sech[c - b*x]*Sinh[a + b*x]^2,x]
```

output

```
(ArcTan[Tanh[(c - b*x)/2]] + ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(a + c)] + Sinh[2*a + c + b*x])/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}(c - bx) dx$$

$$\downarrow 7299$$

$$\int \sinh^2(a + bx) \operatorname{sech}(c - bx) dx$$

input `Int[Sech[c - b*x]*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 243, normalized size of antiderivative = 7.59

method	result
risch	$\frac{e^{bx+2a+c}}{2b} - \frac{e^{-bx-2a-c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a} e^{4a+4c}}{4b} + \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a} e^{2a+2c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a}}{4b}$

input `int(sech(b*x-c)*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+2*a+c)-1/2/b*exp(-b*x-2*a-c)+1/4*I*ln(exp(b*x+a)-I*exp(a+c))
/b*exp(-2*c-2*a)*exp(2*a+2*c)^2+1/2*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c
-2*a)*exp(2*a+2*c)+1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)-1/4*I*ln
n(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2*c)^2-1/2*I*ln(exp(b*x+a
) + I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2*c)-1/4*I*ln(exp(b*x+a)+I*exp(a+c))
/b*exp(-2*c-2*a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 427 vs. $2(34) = 68$.

Time = 0.10 (sec) , antiderivative size = 427, normalized size of antiderivative = 13.34

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(sech(b*x-c)*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/2*(cosh(b*x + a)^2*cosh(a + c) - cosh(a + c)^3 + (cosh(a + c) - sinh(a +
c))*sinh(b*x + a)^2 - 3*cosh(a + c)*sinh(a + c)^2 + sinh(a + c)^3 - (4*co
sh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - 2*(3
*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^2 + 4*(cosh(a + c)^3 + cosh(
a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 + 2*cosh(a + c)^2 + 1)*
cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + sinh(a + c)
^4 + 2*(3*cosh(a + c)^2 + 1)*sinh(a + c)^2 + 2*cosh(a + c)^2 - 4*(cosh(a +
c)^3 + cosh(a + c))*sinh(a + c) + 1)*sinh(b*x + a))*arctan(-cosh(b*x + a)
*cosh(a + c) - (cosh(a + c) - sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*s
inh(a + c)) + 2*(cosh(b*x + a)*cosh(a + c) - cosh(b*x + a)*sinh(a + c))*si
nh(b*x + a) - (cosh(b*x + a)^2 - 3*cosh(a + c)^2)*sinh(a + c))/(b*cosh(b*x
+ a)*cosh(a + c)^2 - 2*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c) + b*cosh(b
*x + a)*sinh(a + c)^2 + (b*cosh(a + c)^2 - 2*b*cosh(a + c)*sinh(a + c) + b
*sinh(a + c)^2)*sinh(b*x + a))
```

Sympy [F]

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}(bx - c) dx$$

input `integrate(sech(b*x-c)*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.28

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1) \arctan(e^{(-bx+c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(sech(b*x-c)*sinh(b*x+a)^2,x, algorithm="maxima")`

output $\frac{1}{2}*(e^{(4*a + 4*c)} + 2*e^{(2*a + 2*c)} + 1)*\arctan(e^{(-b*x + c)})*e^{(-2*a - 2*c)}/b + \frac{1}{2}*e^{(b*x + 2*a + c)}/b - \frac{1}{2}*e^{(-b*x - 2*a - c)}/b$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(34) = 68$.

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.31

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = -\frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1) \arctan(e^{(bx-c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(sech(b*x-c)*sinh(b*x+a)^2,x, algorithm="giac")`

output $-\frac{1}{2}*(e^{(4*a + 4*c)} + 2*e^{(2*a + 2*c)} + 1)*\arctan(e^{(b*x - c)})*e^{(-2*a - 2*c)}/b + \frac{1}{2}*e^{(b*x + 2*a + c)}/b - \frac{1}{2}*e^{(-b*x - 2*a - c)}/b$

Mupad [B] (verification not implemented)

Time = 1.31 (sec) , antiderivative size = 194, normalized size of antiderivative = 6.06

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \frac{e^{2a+c+bx}}{2b} - \frac{e^{-2a-c-bx}}{2b} - \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{-3c} e^{bx} (\sqrt{b^2+2e^{2a} e^{2c} \sqrt{b^2+e^{4a} e^{4c} \sqrt{b^2}})}{b \sqrt{e^{-4a} e^{-4c} (4e^{2a} e^{2c} + 6e^{4a} e^{4c} + 4e^{6a} e^{6c} + e^{8a} e^{8c} + 1)}}}\right) \sqrt{e^{-4a-4c} (4e^{2a+2c} + 6e^{4a+4c} + 4e^{6a+6c} + e^{8a+8c})}}{2\sqrt{b^2}}$$

input `int(sinh(a + b*x)^2/cosh(c - b*x),x)`output `exp(2*a + c + b*x)/(2*b) - exp(- 2*a - c - b*x)/(2*b) - (atan((exp(-2*a)*exp(-3*c)*exp(b*x)*((b^2)^(1/2) + 2*exp(2*a)*exp(2*c)*(b^2)^(1/2) + exp(4*a)*exp(4*c)*(b^2)^(1/2)))/(b*(exp(-4*a)*exp(-4*c)*(4*exp(2*a)*exp(2*c) + 6*exp(4*a)*exp(4*c) + 4*exp(6*a)*exp(6*c) + exp(8*a)*exp(8*c) + 1))^(1/2)))*(exp(- 4*a - 4*c)*(4*exp(2*a + 2*c) + 6*exp(4*a + 4*c) + 4*exp(6*a + 6*c) + exp(8*a + 8*c) + 1))^(1/2))/(2*(b^2)^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 109, normalized size of antiderivative = 3.41

$$\int \operatorname{sech}(c - bx) \sinh^2(a + bx) dx = \frac{-e^{bx+4a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - 2e^{bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2bx+4a+3c} - e^c}{2e^{bx+2a+2c} b}$$

input `int(sech(b*x-c)*sinh(b*x+a)^2,x)`output `(- e**(4*a + b*x + 4*c)*atan(e**(b*x)/e**c) - 2*e**(2*a + b*x + 2*c)*atan(e**(b*x)/e**c) - e**(b*x)*atan(e**(b*x)/e**c) + e**(4*a + 2*b*x + 3*c) - e**c)/(2*e**(2*a + b*x + 2*c)*b)`

3.63 $\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx$

Optimal result	489
Mathematica [B] (verified)	489
Rubi [F]	490
Maple [B] (verified)	490
Fricas [B] (verification not implemented)	491
Sympy [F]	492
Maxima [A] (verification not implemented)	492
Giac [B] (verification not implemented)	492
Mupad [B] (verification not implemented)	493
Reduce [B] (verification not implemented)	493

Optimal result

Integrand size = 18, antiderivative size = 44

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = x \cosh(2(a + c)) + \frac{\log(\cosh(c - bx)) \sinh(2(a + c))}{b} + \frac{\cosh^2(a + c) \tanh(c - bx)}{b}$$

output `x*cosh(2*a+2*c)+ln(cosh(b*x-c))*sinh(2*a+2*c)/b-cosh(a+c)^2*tanh(b*x-c)/b`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 182 vs. 2(44) = 88.

Time = 0.36 (sec) , antiderivative size = 182, normalized size of antiderivative = 4.14

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}(c - bx)(bx \cosh(2a + 2c - bx) + bx \cosh(2a + 4c - bx) + bx \cosh(2a + bx) + bx \cosh(2a + 2c - bx))}{b}$$

input `Integrate[Sech[c - b*x]^2*Sinh[a + b*x]^2,x]`

output

```
(Sech[c]*Sech[c - b*x]*(b*x*Cosh[2*a + 2*c - b*x] + b*x*Cosh[2*a + 4*c - b*x] + b*x*Cosh[2*a + b*x] + b*x*Cosh[2*a + 2*c + b*x] - 2*Sinh[b*x] + Sinh[2*a + 2*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 2*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 4*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + b*x] - Sinh[2*a + 2*c + b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 2*c + b*x]))/(4*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^2(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^2(c - bx) dx$$

input

```
Int[Sech[c - b*x]^2*Sinh[a + b*x]^2,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(51) = 102.

Time = 1.85 (sec) , antiderivative size = 238, normalized size of antiderivative = 5.41

method	result
risch	$x e^{2a+2c} - e^{-2c-2a} e^{4a+4c} x - \frac{e^{-2c-2a} a e^{4a+4c}}{b} + e^{-2c-2a} x + \frac{e^{-2c-2a} a}{b} + \frac{e^{4a+4c}}{2b(e^{2a+2c} + e^{2bx+2a})} + \frac{e^{2a+2c}}{b(e^{2a+2c} + e^{2bx+2a})}$

input

```
int(sech(b*x-c)^2*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
x*exp(2*a+2*c)-exp(-2*c-2*a)*exp(4*a+4*c)*x-1/b*exp(-2*c-2*a)*a*exp(4*a+4*c)+exp(-2*c-2*a)*x+1/b*exp(-2*c-2*a)*a+1/2/b/(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(4*a+4*c)+1/b/(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(2*a+2*c)+1/2/b/(exp(2*a+2*c)+exp(2*b*x+2*a))+1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1304 vs. $2(51) = 102$.

Time = 0.12 (sec) , antiderivative size = 1304, normalized size of antiderivative = 29.64

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
1/2*(2*b*x*cosh(b*x + a)^2*cosh(a + c)^6 - 12*b*x*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^5 + 2*b*x*cosh(b*x + a)^2*sinh(a + c)^6 + (2*b*x + 1)*cosh(a + c)^4 + (30*b*x*cosh(b*x + a)^2*cosh(a + c)^2 + 2*b*x + 1)*sinh(a + c)^4 - 4*(10*b*x*cosh(b*x + a)^2*cosh(a + c)^3 + (2*b*x + 1)*cosh(a + c))*sinh(a + c)^3 + 2*(b*x*cosh(a + c)^6 - 6*b*x*cosh(a + c)^5*sinh(a + c) + 15*b*x*cosh(a + c)^4*sinh(a + c)^2 - 20*b*x*cosh(a + c)^3*sinh(a + c)^3 + 15*b*x*cosh(a + c)^2*sinh(a + c)^4 - 6*b*x*cosh(a + c)*sinh(a + c)^5 + b*x*sinh(a + c)^6)*sinh(b*x + a)^2 + 2*(15*b*x*cosh(b*x + a)^2*cosh(a + c)^4 + 3*(2*b*x + 1)*cosh(a + c)^2 + 1)*sinh(a + c)^2 + 2*cosh(a + c)^2 + (6*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)^2*sinh(a + c)^6 - (15*cosh(b*x + a)^2*cosh(a + c)^2 + 1)*sinh(a + c)^4 - cosh(a + c)^4 + 4*(5*cosh(b*x + a)^2*cosh(a + c)^3 + cosh(a + c))*sinh(a + c)^3 - (cosh(a + c)^6 - cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^6 - 20*cosh(a + c)^3*sinh(a + c)^3 + 15*cosh(a + c)^2*sinh(a + c)^4 - 6*cosh(a + c)*sinh(a + c)^5 + sinh(a + c)^6 + (15*cosh(a + c)^4 - 1)*sinh(a + c)^2 - cosh(a + c)^2 - 2*(3*cosh(a + c)^5 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((15*cosh(a + c)^4 - 1)*cosh(b*x + a)^2 + 6*cosh(a + c)^2)*sinh(a + c)^2 + 2*(20*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c)^3 - 15*cosh(b*x + a)*cosh(a + c)^2*sinh(a + c)^4 + 6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)*sinh(a + c)^6 - (15*cosh(a + c)^4 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(3*...
```

Sympy [F]

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^2(bx - c) dx$$

input `integrate(sech(b*x-c)**2*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x - c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.30

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = \frac{(e^{4a+4c} - 1)e^{(-2a-2c)} \log(e^{(-2bx+2c)} + 1)}{2b} + \frac{(bx + a)e^{(2a+2c)}}{b} - \frac{e^{(4a+4c)} + 2e^{(2a+2c)} + 1}{2b(e^{(-2bx+2a+4c)} + e^{(2a+2c)})}$$

input `integrate(sech(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-2*b*x + 2*c) + 1)/b + (b*x + a)*e^(2*a + 2*c)/b - 1/2*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)/(b*(e^(-2*b*x + 2*a + 4*c) + e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(51) = 102.

Time = 0.11 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.41

$$\begin{aligned} & \int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx \\ &= x e^{(-2a-2c)} + \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(2bx)} + e^{(2c)})}{2b} \\ &+ \frac{(e^{(2bx)} - e^{(2bx+4a+4c)} + 2e^{(2a+4c)} + 2e^{(2c)})e^{(-2a-2c)}}{2b(e^{(2bx)} + e^{(2c)})} \end{aligned}$$

input `integrate(sech(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="giac")`

output `x*e^(-2*a - 2*c) + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(2*b*x) + e^(2*c))/b + 1/2*(e^(2*b*x) - e^(2*b*x + 4*a + 4*c) + 2*e^(2*a + 4*c) + 2*e^(2*c))*e^(-2*a - 2*c)/(b*(e^(2*b*x) + e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.89

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = x e^{-2a-2c} + \frac{\sinh(2a + 2c) \ln(e^{2a} e^{2bx} + e^{2a} e^{2c})}{b} + \frac{2 e^{2a+2c} \cosh(a + c)^2}{b (e^{2a+2c} + e^{2a+2bx})}$$

input `int(sinh(a + b*x)^2/cosh(c - b*x)^2,x)`

output `x*exp(- 2*a - 2*c) + (sinh(2*a + 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(2*c)))/b + (2*exp(2*a + 2*c)*cosh(a + c)^2)/(b*(exp(2*a + 2*c) + exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 179, normalized size of antiderivative = 4.07

$$\int \operatorname{sech}^2(c - bx) \sinh^2(a + bx) dx = \frac{e^{2bx+4a+4c} \log(e^{2bx} + e^{2c}) - e^{2bx+4a+4c} - 2e^{2bx+2a+2c} - e^{2bx} \log(e^{2bx} + e^{2c}) + 2e^{2bx} bx - e^{2bx} + e^{4a+6c} \log(e^{2bx} + e^{2c})}{2e^{2a+2c} b (e^{2bx} + e^{2c})}$$

input `int(sech(b*x-c)^2*sinh(b*x+a)^2,x)`

output

```
(e**(4*a + 2*b*x + 4*c)*log(e**(2*b*x) + e**(2*c)) - e**(4*a + 2*b*x + 4*c)
) - 2*e**(2*a + 2*b*x + 2*c) - e**(2*b*x)*log(e**(2*b*x) + e**(2*c)) + 2*e
**(2*b*x)*b*x - e**(2*b*x) + e**(4*a + 6*c)*log(e**(2*b*x) + e**(2*c)) - e
**(2*c)*log(e**(2*b*x) + e**(2*c)) + 2*e**(2*c)*b*x)/(2*e**(2*a + 2*c)*b*(
e**(2*b*x) + e**(2*c)))
```

3.64 $\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx$

Optimal result	495
Mathematica [A] (verified)	495
Rubi [F]	496
Maple [C] (verified)	496
Fricas [B] (verification not implemented)	497
Sympy [F]	497
Maxima [A] (verification not implemented)	498
Giac [A] (verification not implemented)	498
Mupad [F(-1)]	499
Reduce [B] (verification not implemented)	499

Optimal result

Integrand size = 18, antiderivative size = 86

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \frac{\arctan(\sinh(c - bx)) \cosh^2(a + c)}{2b} - \frac{\arctan(\sinh(c - bx)) \cosh(2(a + c))}{b} - \frac{\operatorname{sech}(c - bx) \sinh(2(a + c))}{b} + \frac{\cosh^2(a + c) \operatorname{sech}(c - bx) \tanh(c - bx)}{2b}$$

output

```
-1/2*arctan(sinh(b*x-c))*cosh(a+c)^2/b+arctan(sinh(b*x-c))*cosh(2*a+2*c)/b
-sech(b*x-c)*sinh(2*a+2*c)/b-1/2*cosh(a+c)^2*sech(b*x-c)*tanh(b*x-c)/b
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.28

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \frac{-12 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right) \cosh(2(a + c)) + \operatorname{sech}^2(c - bx) \left(2 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right)\right) + 2 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right)}{8b}$$

input `Integrate[Sech[c - b*x]^3*Sinh[a + b*x]^2,x]`

output `(-12*ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(a + c)] + Sech[c - b*x]^2*(2*ArcTan[Tanh[(c - b*x)/2]] + 2*ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(c - b*x)] + 2*Sinh[c - b*x] - 3*Sinh[2*a + 3*c - b*x] - 5*Sinh[2*a + c + b*x]))/(8*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

input `Int[Sech[c - b*x]^3*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.70 (sec) , antiderivative size = 311, normalized size of antiderivative = 3.62

method	result
risch	$-\frac{(3e^{6a+6c}+5e^{2bx+6a+4c}-2e^{4a+4c}+2e^{2bx+4a+2c}-5e^{2a+2c}-3e^{2bx+2a})e^{bx-c}}{4(e^{2a+2c}+e^{2bx+2a})^2b} + \frac{3i \ln(e^{bx+a}+ie^{a+c})e^{-2c-2ae^{4a+4c}}}{8b} - \frac{i \ln(e^{bx+a}-ie^{a+c})e^{-2c-2ae^{4a+4c}}}{8b}$

input `int(sech(b*x-c)^3*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*a+2*c)+exp(2*b*x+2*a))^2/b*(3*exp(6*a+6*c)+5*exp(2*b*x+6*a+4*c)
)-2*exp(4*a+4*c)+2*exp(2*b*x+4*a+2*c)-5*exp(2*a+2*c)-3*exp(2*b*x+2*a))*exp
(b*x-c)+3/8*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/4*I
*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*I*ln(exp(b*x+a
)+I*exp(a+c))/b*exp(-2*c-2*a)-3/8*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2
*a)*exp(4*a+4*c)+1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2
*c)-3/8*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3915 vs. $2(90) = 180$.

Time = 0.12 (sec) , antiderivative size = 3915, normalized size of antiderivative = 45.52

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^3(bx - c) dx$$

input

```
integrate(sech(b*x-c)**3*sinh(b*x+a)**2,x)
```

output

```
Integral(sinh(a + b*x)**2*sech(b*x - c)**3, x)
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.77

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx$$

$$= -\frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3) \arctan(e^{(-bx+c)}) e^{(-2a-2c)}}{4b}$$

$$-\frac{(5e^{(4a+4c)} + 2e^{(2a+2c)} - 3)e^{(-bx-a)} + (3e^{(6a+6c)} - 2e^{(4a+4c)} - 5e^{(2a+2c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})}$$

input `integrate(sech(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="maxima")`output `-1/4*(3*e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 3)*arctan(e^(-b*x + c))*e^(-2*a - 2*c)/b - 1/4*((5*e^(4*a + 4*c) + 2*e^(2*a + 2*c) - 3)*e^(-b*x - a) + (3*e^(6*a + 6*c) - 2*e^(4*a + 4*c) - 5*e^(2*a + 2*c))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c)))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.65

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3) \arctan(e^{(bx-c)}) e^{(-2a-2c)}}{4b}$$

$$+ \frac{(3e^{(3bx)} - 5e^{(3bx+4a+4c)} - 2e^{(3bx+2a+2c)} - 3e^{(bx+4a+6c)} + 2e^{(bx+2a+4c)} + 5e^{(bx+2c)})e^{(-2a-c)}}{4b(e^{(2bx)} + e^{(2c)})^2}$$

input `integrate(sech(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="giac")`output `1/4*(3*e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 3)*arctan(e^(b*x - c))*e^(-2*a - 2*c)/b + 1/4*(3*e^(3*b*x) - 5*e^(3*b*x + 4*a + 4*c) - 2*e^(3*b*x + 2*a + 2*c) - 3*e^(b*x + 4*a + 6*c) + 2*e^(b*x + 2*a + 4*c) + 5*e^(b*x + 2*c))*e^(-2*a - c)/(b*(e^(2*b*x) + e^(2*c))^2)`

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\cosh(c - bx)^3} dx$$

input `int(sinh(a + b*x)^2/cosh(c - b*x)^3,x)`output `int(sinh(a + b*x)^2/cosh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 339, normalized size of antiderivative = 3.94

$$\int \operatorname{sech}^3(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - 2e^{4bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 3e^{4bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 6e^{2bx+4a+6c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - 4e^{2bx+2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right)}{1}$$

input `int(sech(b*x-c)^3*sinh(b*x+a)^2,x)`output `(3*e**(4*a + 4*b*x + 4*c)*atan(e**(b*x)/e**c) - 2*e**(2*a + 4*b*x + 2*c)*atan(e**(b*x)/e**c) + 3*e**(4*b*x)*atan(e**(b*x)/e**c) + 6*e**(4*a + 2*b*x + 6*c)*atan(e**(b*x)/e**c) - 4*e**(2*a + 2*b*x + 4*c)*atan(e**(b*x)/e**c) + 6*e**(2*b*x + 2*c)*atan(e**(b*x)/e**c) + 3*e**(4*a + 8*c)*atan(e**(b*x)/e**c) - 2*e**(2*a + 6*c)*atan(e**(b*x)/e**c) + 3*e**(4*c)*atan(e**(b*x)/e**c) - 5*e**(4*a + 3*b*x + 5*c) - 2*e**(2*a + 3*b*x + 3*c) + 3*e**(3*b*x + c) - 3*e**(4*a + b*x + 7*c) + 2*e**(2*a + b*x + 5*c) + 5*e**(b*x + 3*c))/(4*e**(2*a + 2*c)*b*(e**(4*b*x) + 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.65 $\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx$

Optimal result	500
Mathematica [A] (verified)	500
Rubi [F]	501
Maple [A] (verified)	501
Fricas [B] (verification not implemented)	502
Sympy [F]	503
Maxima [B] (verification not implemented)	503
Giac [A] (verification not implemented)	504
Mupad [F(-1)]	504
Reduce [B] (verification not implemented)	505

Optimal result

Integrand size = 18, antiderivative size = 80

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = -\frac{\operatorname{sech}^2(c - bx) \sinh(2(a + c))}{2b} + \frac{\cosh^2(a + c) \tanh(c - bx)}{b} - \frac{\cosh(2(a + c)) \tanh(c - bx)}{b} - \frac{\cosh^2(a + c) \tanh^3(c - bx)}{3b}$$

```
output -1/2*sech(b*x-c)^2*sinh(2*a+2*c)/b-cosh(a+c)^2*tanh(b*x-c)/b+cosh(2*a+2*c)
*tanh(b*x-c)/b+1/3*cosh(a+c)^2*tanh(b*x-c)^3/b
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.04

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}^3(c - bx)(3 \sinh(bx) - \sinh(2c - 3bx) + \sinh(2a + 4c - 3bx) + 3 \sinh(2a + 2c - bx) + 3 \sinh(2a + 2c - bx))}{12b}$$

input `Integrate[Sech[c - b*x]^4*Sinh[a + b*x]^2,x]`

output `-1/12*(Sech[c]*Sech[c - b*x]^3*(3*Sinh[b*x] - Sinh[2*c - 3*b*x] + Sinh[2*a + 4*c - 3*b*x] + 3*Sinh[2*a + 2*c - b*x] + 3*Sinh[2*a + b*x] - Sinh[2*a + 3*b*x]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{sech}^4(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{sech}^4(c - bx) dx$$

input `Int[Sech[c - b*x]^4*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [A] (verified)

Time = 2.94 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.80

method	result	size
parallelrisch	$\frac{2 \sinh(3bx+2a-c) - 3 \sinh(bx-c) - \sinh(3bx-3c)}{3b(\cosh(3bx-3c) + 3 \cosh(bx-c))}$	64
risch	$-\frac{2(e^{4a+4c} + 3e^{2bx+4a+2c} - e^{2a+2c} + 3e^{4bx+4a} - 3e^{2bx+2a} + 1)e^{4a+4c}}{3(e^{2a+2c} + e^{2bx+2a})^3 b}$	90

input `int(sech(b*x-c)^4*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
1/3/b*(2*sinh(3*b*x+2*a-c)-3*sinh(b*x-c)-sinh(3*b*x-3*c))/(cosh(3*b*x-3*c)
+3*cosh(b*x-c))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1015 vs. $2(84) = 168$.

Time = 0.10 (sec) , antiderivative size = 1015, normalized size of antiderivative = 12.69

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(sech(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
-2/3*(4*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^3 + (5*cosh(b*x + a)^2 - 3)
)*sinh(a + c)^4 - 3*cosh(a + c)^4 + (5*cosh(a + c)^4 - cosh(a + c)^2)*cosh
(b*x + a)^2 + (5*cosh(a + c)^4 + 4*cosh(a + c)*sinh(a + c)^3 + 5*sinh(a +
c)^4 - (2*cosh(a + c)^2 + 1)*sinh(a + c)^2 - cosh(a + c)^2 + 2*(2*cosh(a +
c)^3 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((2*cosh(a + c)^2 + 1)
*cosh(b*x + a)^2 - 6*cosh(a + c)^2 - 3)*sinh(a + c)^2 + 3*cosh(a + c)^2 -
2*(4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4
+ (10*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3
- cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 + cosh(a + c)^2
)*cosh(b*x + a))*sinh(b*x + a) + 2*((2*cosh(a + c)^3 - cosh(a + c))*cosh(b
*x + a)^2 + 3*cosh(a + c))*sinh(a + c))/(b*cosh(b*x + a)^4*cosh(a + c)^4 +
4*b*cosh(b*x + a)^2*cosh(a + c)^4 + 3*b*cosh(a + c)^4 + (b*cosh(a + c)^4
- 4*b*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(a + c)^2*sinh(a + c)^2 - 4*b*co
sh(a + c)*sinh(a + c)^3 + b*sinh(a + c)^4)*sinh(b*x + a)^4 + (b*cosh(b*x +
a)^4 - 4*b*cosh(b*x + a)^2 + 3*b)*sinh(a + c)^4 + 4*(b*cosh(b*x + a)*cosh
(a + c)^4 - 4*b*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(b*x + a)
)*cosh(a + c)^2*sinh(a + c)^2 - 4*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^
3 + b*cosh(b*x + a)*sinh(a + c)^4)*sinh(b*x + a)^3 - 4*(b*cosh(b*x + a)^4*
cosh(a + c) - b*cosh(b*x + a)^2*cosh(a + c))*sinh(a + c)^3 + 2*(3*b*cosh(b
*x + a)^2*cosh(a + c)^4 + 18*b*cosh(b*x + a)^2*cosh(a + c)^2*sinh(a + c)...
```

Sympy [F]

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{sech}^4(bx - c) dx$$

input `integrate(sech(b*x-c)**4*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*sech(b*x - c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. $2(84) = 168$.

Time = 0.04 (sec) , antiderivative size = 334, normalized size of antiderivative = 4.18

$$\begin{aligned} & \int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx \\ &= -\frac{2(e^{4a+4c} - e^{2a+2c})e^{-2bx-2a}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} + e^{2a+2c})} \\ & \quad + \frac{2e^{-4bx+4c}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} + e^{2a+2c})} \\ & \quad + \frac{2e^{4a+4c}}{3b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} + e^{2a+2c})} \\ & \quad - \frac{2e^{2a+2c}}{3b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} + e^{2a+2c})} \\ & \quad + \frac{2}{3b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} + e^{2a+2c})} \end{aligned}$$

input `integrate(sech(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="maxima")`

output

```
-2*(e^(4*a + 4*c) - e^(2*a + 2*c))*e^(-2*b*x - 2*a)/(b*(3*e^(-2*b*x + 2*a + 4*c) + 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) + e^(2*a + 2*c))) + 2*e^(-4*b*x + 4*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) + 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) + e^(2*a + 2*c))) + 2/3*e^(4*a + 4*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) + 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) + e^(2*a + 2*c))) - 2/3*e^(2*a + 2*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) + 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) + e^(2*a + 2*c))) + 2/3/(b*(3*e^(-2*b*x + 2*a + 4*c) + 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) + e^(2*a + 2*c)))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.08

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \frac{2(3e^{(4bx+4a+4c)} + 3e^{(2bx+4a+6c)} - 3e^{(2bx+2a+4c)} + e^{(4a+8c)} - e^{(2a+6c)} + e^{(4c)})e^{(-2a)}}{3b(e^{(2bx)} + e^{(2c)})^3}$$

input

```
integrate(sech(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="giac")
```

output

```
-2/3*(3*e^(4*b*x + 4*a + 4*c) + 3*e^(2*b*x + 4*a + 6*c) - 3*e^(2*b*x + 2*a + 4*c) + e^(4*a + 8*c) - e^(2*a + 6*c) + e^(4*c))*e^(-2*a)/(b*(e^(2*b*x) + e^(2*c))^3)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\cosh(c - bx)^4} dx$$

input

```
int(sinh(a + b*x)^2/cosh(c - b*x)^4,x)
```

output

```
int(sinh(a + b*x)^2/cosh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.21

$$\int \operatorname{sech}^4(c - bx) \sinh^2(a + bx) dx = \frac{2e^{2c}(e^{6bx+4a} + 3e^{2bx+2a+2c} + e^{2a+4c} - e^{2c})}{3e^{2a}b(e^{6bx} + 3e^{4bx+2c} + 3e^{2bx+4c} + e^{6c})}$$

input `int(sech(b*x-c)^4*sinh(b*x+a)^2,x)`

output `(2*e**(2*c)*(e**(4*a + 6*b*x) + 3*e**(2*a + 2*b*x + 2*c) + e**(2*a + 4*c) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x) + 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) + e**(6*c)))`

3.66 $\int \operatorname{csch}(2x) \sinh(x) dx$

Optimal result	506
Mathematica [A] (verified)	506
Rubi [A] (verified)	507
Maple [A] (verified)	508
Fricas [A] (verification not implemented)	508
Sympy [F]	509
Maxima [A] (verification not implemented)	509
Giac [A] (verification not implemented)	509
Mupad [B] (verification not implemented)	510
Reduce [B] (verification not implemented)	510

Optimal result

Integrand size = 7, antiderivative size = 7

$$\int \operatorname{csch}(2x) \sinh(x) dx = \frac{1}{2} \arctan(\sinh(x))$$

output `1/2*arctan(sinh(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}(2x) \sinh(x) dx = -\frac{1}{2} \cot^{-1}(\sinh(x))$$

input `Integrate[Csch[2*x]*Sinh[x],x]`

output `-1/2*ArcCot[Sinh[x]]`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4776, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{csch}(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(ix)}{\sin(2ix)} dx \\
 & \quad \downarrow \text{4776} \\
 & \frac{\int \operatorname{sech}(x) dx}{2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \csc\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4257} \\
 & \frac{1}{2} \arctan(\sinh(x))
 \end{aligned}$$

input `Int [Csch [2*x] *Sinh [x] , x]`

output `ArcTan [Sinh [x]] /2`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4776 `Int[((f_.)*sin[(a_.) + (b_.)*(x_)])^(n_.)*sin[(c_.) + (d_.)*(x_)^(p_.), x_Symbol] := Simp[2^p/f^p Int[Cos[a + b*x]^p*(f*Sin[a + b*x])^(n + p), x], x] /; FreeQ[{a, b, c, d, f, n}, x] && EqQ[b*c - a*d, 0] && EqQ[d/b, 2] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{\arctan(\sinh(x))}{2}$	6
risch	$\frac{i \ln(e^x+i)}{2} - \frac{i \ln(e^x-i)}{2}$	20

input `int(csch(2*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `1/2*arctan(sinh(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \operatorname{csch}(2x) \sinh(x) dx = \arctan(\cosh(x) + \sinh(x))$$

input `integrate(csch(2*x)*sinh(x),x, algorithm="fricas")`

output `arctan(cosh(x) + sinh(x))`

Sympy [F]

$$\int \operatorname{csch}(2x) \sinh(x) dx = \int \sinh(x) \operatorname{csch}(2x) dx$$

input `integrate(csch(2*x)*sinh(x),x)`

output `Integral(sinh(x)*csch(2*x), x)`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}(2x) \sinh(x) dx = -\arctan(e^{-x})$$

input `integrate(csch(2*x)*sinh(x),x, algorithm="maxima")`

output `-arctan(e^(-x))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.43

$$\int \operatorname{csch}(2x) \sinh(x) dx = \arctan(e^x)$$

input `integrate(csch(2*x)*sinh(x),x, algorithm="giac")`

output `arctan(e^x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.43

$$\int \operatorname{csch}(2x) \sinh(x) dx = \operatorname{atan}(e^x)$$

input `int(sinh(x)/sinh(2*x),x)`

output `atan(exp(x))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.57

$$\int \operatorname{csch}(2x) \sinh(x) dx = \operatorname{atan}(e^x)$$

input `int(csch(2*x)*sinh(x),x)`

output `atan(e**x)`

3.67 $\int \operatorname{csch}(3x) \sinh(x) dx$

Optimal result	511
Mathematica [A] (verified)	511
Rubi [A] (verified)	512
Maple [C] (verified)	513
Fricas [B] (verification not implemented)	513
Sympy [F]	514
Maxima [B] (verification not implemented)	514
Giac [A] (verification not implemented)	514
Mupad [B] (verification not implemented)	515
Reduce [B] (verification not implemented)	515

Optimal result

Integrand size = 7, antiderivative size = 15

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{\arctan\left(\frac{\tanh(x)}{\sqrt{3}}\right)}{\sqrt{3}}$$

output `1/3*arctan(1/3*tanh(x)*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{\arctan\left(\frac{\tanh(x)}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Integrate[Csch[3*x]*Sinh[x],x]`

output `ArcTan[Tanh[x]/Sqrt[3]]/Sqrt[3]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4889, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(x) \operatorname{csch}(3x) dx$$

$$\downarrow 3042$$

$$\int \frac{\sin(ix)}{\sin(3ix)} dx$$

$$\downarrow 4889$$

$$\int \frac{1}{\tanh^2(x) + 3} d \tanh(x)$$

$$\downarrow 216$$

$$\frac{\arctan\left(\frac{\tanh(x)}{\sqrt{3}}\right)}{\sqrt{3}}$$

input `Int [Csch [3*x] *Sinh [x] , x]`

output `ArcTan [Tanh [x] /Sqrt [3]] /Sqrt [3]`

Defintions of rubi rules used

rule 216 `Int [((a_) + (b_) * (x_) ^ 2) ^ (-1), x_Symbol] := Simp [(1 / (Rt [a, 2] * Rt [b, 2])) * ArcTan [Rt [b, 2] * (x / Rt [a, 2])], x] /; FreeQ [{a, b}, x] && PosQ [a / b] && (GtQ [a, 0] || GtQ [b, 0])`

rule 3042 `Int [u_, x_Symbol] := Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinearQ [u, x]`

rule 4889

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.67

method	result	size
risch	$\frac{i\sqrt{3} \ln\left(e^{2x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}\right)}{6} - \frac{i\sqrt{3} \ln\left(e^{2x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}\right)}{6}$	40

input

```
int(csch(3*x)*sinh(x),x,method=_RETURNVERBOSE)
```

output

```
1/6*I*3^(1/2)*ln(exp(2*x)+1/2+1/2*I*3^(1/2))-1/6*I*3^(1/2)*ln(exp(2*x)+1/2
-1/2*I*3^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.07

$$\int \operatorname{csch}(3x) \sinh(x) dx = -\frac{1}{3} \sqrt{3} \arctan \left(-\frac{3\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)}{3(\cosh(x) - \sinh(x))} \right)$$

input

```
integrate(csch(3*x)*sinh(x),x, algorithm="fricas")
```

output

```
-1/3*sqrt(3)*arctan(-1/3*(3*sqrt(3)*cosh(x) + sqrt(3)*sinh(x))/(cosh(x) -
sinh(x)))
```

Sympy [F]

$$\int \operatorname{csch}(3x) \sinh(x) dx = \int \sinh(x) \operatorname{csch}(3x) dx$$

input `integrate(csch(3*x)*sinh(x),x)`

output `Integral(sinh(x)*csch(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(13) = 26$.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.60

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2e^{-x} + 1) \right) - \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2e^{-x} - 1) \right)$$

input `integrate(csch(3*x)*sinh(x),x, algorithm="maxima")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(-x) + 1)) - 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(-x) - 1))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} (2e^{2x} + 1) \right)$$

input `integrate(csch(3*x)*sinh(x),x, algorithm="giac")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(2*x) + 1))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2e^{2x}+1)}{3}\right)}{3}$$

input `int(sinh(x)/sinh(3*x),x)`output `(3^(1/2)*atan((3^(1/2)*(2*exp(2*x) + 1))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.20

$$\int \operatorname{csch}(3x) \sinh(x) dx = \frac{\sqrt{3} \left(\operatorname{atan}\left(\frac{2e^x-1}{\sqrt{3}}\right) - \operatorname{atan}\left(\frac{2e^x+1}{\sqrt{3}}\right) \right)}{3}$$

input `int(csch(3*x)*sinh(x),x)`output `(sqrt(3)*(atan((2*e**x - 1)/sqrt(3)) - atan((2*e**x + 1)/sqrt(3))))/3`

3.68 $\int \operatorname{csch}(4x) \sinh(x) dx$

Optimal result	516
Mathematica [A] (verified)	516
Rubi [A] (verified)	517
Maple [C] (verified)	518
Fricas [B] (verification not implemented)	519
Sympy [F]	519
Maxima [B] (verification not implemented)	520
Giac [B] (verification not implemented)	520
Mupad [B] (verification not implemented)	521
Reduce [B] (verification not implemented)	521

Optimal result

Integrand size = 7, antiderivative size = 26

$$\int \operatorname{csch}(4x) \sinh(x) dx = -\frac{1}{4} \arctan(\sinh(x)) + \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}}$$

output `-1/4*arctan(sinh(x))+1/4*arctan(sinh(x)*2^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}(4x) \sinh(x) dx = -\frac{1}{4} \arctan(\sinh(x)) + \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}}$$

input `Integrate[Csch[4*x]*Sinh[x],x]`

output `-1/4*ArcTan[Sinh[x]] + ArcTan[Sqrt[2]*Sinh[x]]/(2*Sqrt[2])`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4878, 1406, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{csch}(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(ix)}{\sin(4ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{1}{8 \sinh^4(x) + 12 \sinh^2(x) + 4} d \sinh(x) \\
 & \quad \downarrow \text{1406} \\
 & 2 \int \frac{1}{8 \sinh^2(x) + 4} d \sinh(x) - 2 \int \frac{1}{8 \sinh^2(x) + 8} d \sinh(x) \\
 & \quad \downarrow \text{216} \\
 & \frac{\arctan(\sqrt{2} \sinh(x))}{2\sqrt{2}} - \frac{1}{4} \arctan(\sinh(x))
 \end{aligned}$$

input `Int [Csch [4*x] *Sinh [x] , x]`

output `-1/4*ArcTan [Sinh [x]] + ArcTan [Sqrt [2] *Sinh [x]] / (2*Sqrt [2])`

Definitions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 1406 $\text{Int}[(a_ + (b_ \cdot x_)^2 + (c_ \cdot x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4 \cdot a \cdot c, 2]\}, \text{Simp}[c/q \ \text{Int}[1/(b/2 - q/2 + c \cdot x^2), x], x] - \text{Simp}[c/q \ \text{Int}[1/(b/2 + q/2 + c \cdot x^2), x], x] /;$ $\text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{PosQ}[b^2 - 4 \cdot a \cdot c]$

rule 3042 $\text{Int}[u_ , x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4878 $\text{Int}[u_ , x_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}\{d = \text{FreeFactors}[\text{Sin}[v], x]\}, d/\text{Coefficient}[v, x, 1] \ \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Sin}[v]/d, u/\text{Cos}[v], x], x], x, \text{Sin}[v]/d], x] /;$ $!\text{FalseQ}[v] \ \&\& \ \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Sin}[v], x], u/\text{Cos}[v], x]$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.38

method	result	size
risch	$\frac{i \ln(e^x - i)}{4} - \frac{i \ln(e^x + i)}{4} + \frac{i\sqrt{2} \ln(e^{2x} + i\sqrt{2}e^x - 1)}{8} - \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x - 1)}{8}$	62

input $\text{int}(\text{csch}(4 \cdot x) \cdot \sinh(x), x, \text{method}=_RETURNVERBOSE)$

output $1/4 \cdot I \cdot \ln(\exp(x) - I) - 1/4 \cdot I \cdot \ln(\exp(x) + I) + 1/8 \cdot I \cdot 2^{(1/2)} \cdot \ln(\exp(2 \cdot x) + I \cdot 2^{(1/2)} \cdot \exp(x) - 1) - 1/8 \cdot I \cdot 2^{(1/2)} \cdot \ln(\exp(2 \cdot x) - I \cdot 2^{(1/2)} \cdot \exp(x) - 1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(18) = 36$.

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.92

$$\begin{aligned} & \int \operatorname{csch}(4x) \sinh(x) dx \\ &= \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \cosh(x) + \frac{1}{2} \sqrt{2} \sinh(x) \right) \\ & \quad - \frac{1}{4} \sqrt{2} \arctan \left(-\frac{\sqrt{2} \cosh(x)^2 + 2 \sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}}{2 (\cosh(x) - \sinh(x))} \right) \\ & \quad - \frac{1}{2} \arctan (\cosh(x) + \sinh(x)) \end{aligned}$$

input `integrate(csch(4*x)*sinh(x),x, algorithm="fricas")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)) - 1/4*sqrt(2)*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2))/(cosh(x) - sinh(x))) - 1/2*arctan(cosh(x) + sinh(x))`

Sympy [F]

$$\int \operatorname{csch}(4x) \sinh(x) dx = \int \sinh(x) \operatorname{csch}(4x) dx$$

input `integrate(csch(4*x)*sinh(x),x)`

output `Integral(sinh(x)*csch(4*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. $2(18) = 36$.

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.92

$$\int \operatorname{csch}(4x) \sinh(x) dx = -\frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2e^{-x}) \right) - \frac{1}{4} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2e^{-x}) \right) + \frac{1}{2} \arctan (e^{-x})$$

input `integrate(csch(4*x)*sinh(x),x, algorithm="maxima")`

output `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(-x))) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^(-x))) + 1/2*arctan(e^(-x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(18) = 36$.

Time = 0.14 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.69

$$\int \operatorname{csch}(4x) \sinh(x) dx = -\frac{1}{8} \pi + \frac{1}{8} \sqrt{2} \left(\pi + 2 \arctan \left(\frac{1}{2} \sqrt{2} (e^{2x} - 1) e^{-x} \right) \right) - \frac{1}{4} \arctan \left(\frac{1}{2} (e^{2x} - 1) e^{-x} \right)$$

input `integrate(csch(4*x)*sinh(x),x, algorithm="giac")`

output `-1/8*pi + 1/8*sqrt(2)*(pi + 2*arctan(1/2*sqrt(2)*(e^(2*x) - 1)*e^(-x))) - 1/4*arctan(1/2*(e^(2*x) - 1)*e^(-x))`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.62

$$\int \operatorname{csch}(4x) \sinh(x) dx = \frac{\sqrt{2} \left(2 \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2} + \frac{\sqrt{2}e^{3x}}{2}\right) + 2 \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2}\right) \right)}{8} - \frac{\operatorname{atan}(e^x)}{2}$$

input `int(sinh(x)/sinh(4*x),x)`output `(2^(1/2)*(2*atan((2^(1/2)*exp(x))/2 + (2^(1/2)*exp(3*x))/2) + 2*atan((2^(1/2)*exp(x))/2)))/8 - atan(exp(x))/2`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.73

$$\int \operatorname{csch}(4x) \sinh(x) dx = -\frac{\operatorname{atan}(e^x)}{2} + \frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right)}{4} + \frac{\sqrt{2} \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right)}{4}$$

input `int(csch(4*x)*sinh(x),x)`output `(- 2*atan(e**x) + sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) + sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)))/4`

3.69 $\int \operatorname{csch}(5x) \sinh(x) dx$

Optimal result	522
Mathematica [A] (verified)	522
Rubi [A] (verified)	523
Maple [C] (verified)	524
Fricas [B] (verification not implemented)	525
Sympy [F]	525
Maxima [F]	526
Giac [A] (verification not implemented)	526
Mupad [B] (verification not implemented)	527
Reduce [F]	528

Optimal result

Integrand size = 7, antiderivative size = 80

$$\int \operatorname{csch}(5x) \sinh(x) dx = -\frac{1}{5} \sqrt{\frac{1}{2} (5 + \sqrt{5})} \arctan \left(\sqrt{\frac{1}{5} (5 - 2\sqrt{5})} \tanh(x) \right) + \sqrt{\frac{2}{5(5 + \sqrt{5})}} \arctan \left(\sqrt{\frac{1}{5} (5 + 2\sqrt{5})} \tanh(x) \right)$$

output

```
-1/10*(10+2*5^(1/2))^(1/2)*arctan(1/5*(25-10*5^(1/2))^(1/2)*tanh(x))+2^(1/2)/(25+5*5^(1/2))^(1/2)*arctan(1/5*(25+10*5^(1/2))^(1/2)*tanh(x))
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.05

$$\int \operatorname{csch}(5x) \sinh(x) dx = \frac{\sqrt{5 + \sqrt{5}} \arctan \left(\frac{(-3 + \sqrt{5}) \tanh(x)}{\sqrt{10 - 2\sqrt{5}}} \right) + \sqrt{5 - \sqrt{5}} \arctan \left(\frac{(3 + \sqrt{5}) \tanh(x)}{\sqrt{2(5 + \sqrt{5})}} \right)}{5\sqrt{2}}$$

input `Integrate[Csch[5*x]*Sinh[x],x]`

output $(\sqrt{5 + \sqrt{5}} \operatorname{ArcTan}[\frac{(-3 + \sqrt{5}) \operatorname{Tanh}[x]}{\sqrt{10 - 2\sqrt{5}}}] + \sqrt{5 - \sqrt{5}} \operatorname{ArcTan}[\frac{(3 + \sqrt{5}) \operatorname{Tanh}[x]}{\sqrt{2(5 + \sqrt{5})}}]) / (5\sqrt{2})$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4889, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(x) \operatorname{csch}(5x) dx \\ & \quad \downarrow 3042 \\ & \int \frac{\sin(ix)}{\sin(5ix)} dx \\ & \quad \downarrow 4889 \\ & \int \frac{1 - \tanh^2(x)}{\tanh^4(x) + 10 \tanh^2(x) + 5} d \tanh(x) \\ & \quad \downarrow 1480 \\ & -\frac{1}{10} (5 - 3\sqrt{5}) \int \frac{1}{\tanh^2(x) - 2\sqrt{5} + 5} d \tanh(x) - \\ & \quad \frac{1}{10} (5 + 3\sqrt{5}) \int \frac{1}{\tanh^2(x) + 2\sqrt{5} + 5} d \tanh(x) \\ & \quad \downarrow 216 \\ & -\frac{(5 - 3\sqrt{5}) \arctan\left(\frac{\tanh(x)}{\sqrt{5-2\sqrt{5}}}\right)}{10\sqrt{5-2\sqrt{5}}} - \frac{(5 + 3\sqrt{5}) \arctan\left(\frac{\tanh(x)}{\sqrt{5+2\sqrt{5}}}\right)}{10\sqrt{5+2\sqrt{5}}} \end{aligned}$$

input `Int[Csch[5*x]*Sinh[x],x]`

output

$$-1/10*((5 - 3\sqrt{5})\operatorname{ArcTan}[\operatorname{Tanh}[x]/\sqrt{5 - 2\sqrt{5}}])/\sqrt{5 - 2\sqrt{5}} - ((5 + 3\sqrt{5})\operatorname{ArcTan}[\operatorname{Tanh}[x]/\sqrt{5 + 2\sqrt{5}}])/(10\sqrt{5 + 2\sqrt{5}})$$
Defintions of rubi rules used

rule 216

$$\operatorname{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[b, 2])) \cdot \operatorname{ArcTan}[\operatorname{Rt}[b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \operatorname{GtQ}[b, 0])$$

rule 1480

$$\operatorname{Int}[(d_ + (e_ \cdot)(x_)^2)/((a_ + (b_ \cdot)(x_)^2 + (c_ \cdot)(x_)^4), x_Symbol] : > \operatorname{With}[\{q = \operatorname{Rt}[b^2 - 4ac, 2]\}, \operatorname{Simp}[(e/2 + (2cd - be)/(2q)) \operatorname{Int}[1/(b/2 - q/2 + cx^2), x], x] + \operatorname{Simp}[(e/2 - (2cd - be)/(2q)) \operatorname{Int}[1/(b/2 + q/2 + cx^2), x], x]] \text{ ; FreeQ}\{a, b, c, d, e\}, x \ \&\& \operatorname{NeQ}[b^2 - 4ac, 0] \ \&\& \operatorname{NeQ}[c^2 - ae^2, 0] \ \&\& \operatorname{PosQ}[b^2 - 4ac]$$

rule 3042

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4889

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{With}[\{v = \operatorname{FunctionOfTrig}[u, x]\}, \operatorname{With}[\{d = \operatorname{FreeFactors}[\operatorname{Tan}[v], x]\}, \operatorname{Simp}[d/\operatorname{Coefficient}[v, x, 1] \operatorname{Subst}[\operatorname{Int}[\operatorname{SubstFor}[1/(1 + d^2x^2), \operatorname{Tan}[v]/d, u, x], x], x, \operatorname{Tan}[v]/d], x]] \text{ ; !FalseQ}[v] \ \&\& \operatorname{FunctionOfQ}[\operatorname{NonfreeFactors}[\operatorname{Tan}[v], x], u, x]] \text{ ; InverseFunctionFreeQ}[u, x] \ \&\& \operatorname{!MatchQ}[u, (v_ \cdot)((c_ \cdot)\operatorname{tan}[w_]^{(n_ \cdot)}\operatorname{tan}[z_]^{(n_ \cdot)})^{(p_ \cdot)}] \text{ ; FreeQ}\{c, p\}, x \ \&\& \operatorname{IntegerQ}[n] \ \&\& \operatorname{LinearQ}[w, x] \ \&\& \operatorname{EqQ}[z, 2w]]$$
Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.51

method	result	size
risch	$2 \left(\sum_{R=\operatorname{RootOf}(32000Z^4+400Z^2+1)} -R \ln(4000R^3 - 200R^2 + e^{2x} + 30R - 1) \right)$	41

input `int(csch(5*x)*sinh(x),x,method=_RETURNVERBOSE)`

output `2*sum(_R*ln(4000*_R^3-200*_R^2+exp(2*x)+30*_R-1),_R=RootOf(32000*_Z^4+400*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(53) = 106$.

Time = 0.10 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.39

$$\int \operatorname{csch}(5x) \sinh(x) dx =$$

$$-\frac{1}{5} \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} \arctan \left(\frac{1}{10} \left(4 \sqrt{5} \cosh(x)^2 + 8 \sqrt{5} \cosh(x) \sinh(x) + 4 \sqrt{5} \sinh(x)^2 + \sqrt{5} + 5 \right) \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} \right)$$

$$+ \frac{1}{5} \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}} \arctan \left(\frac{1}{10} \left(4 \sqrt{5} \cosh(x)^2 + 8 \sqrt{5} \cosh(x) \sinh(x) + 4 \sqrt{5} \sinh(x)^2 + \sqrt{5} - 5 \right) \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}} \right)$$

input `integrate(csch(5*x)*sinh(x),x, algorithm="fricas")`

output `-1/5*sqrt(1/2*sqrt(5) + 5/2)*arctan(1/10*(4*sqrt(5)*cosh(x)^2 + 8*sqrt(5)*cosh(x)*sinh(x) + 4*sqrt(5)*sinh(x)^2 + sqrt(5) + 5)*sqrt(1/2*sqrt(5) + 5/2)) + 1/5*sqrt(-1/2*sqrt(5) + 5/2)*arctan(1/10*(4*sqrt(5)*cosh(x)^2 + 8*sqrt(5)*cosh(x)*sinh(x) + 4*sqrt(5)*sinh(x)^2 + sqrt(5) - 5)*sqrt(-1/2*sqrt(5) + 5/2))`

Sympy [F]

$$\int \operatorname{csch}(5x) \sinh(x) dx = \int \sinh(x) \operatorname{csch}(5x) dx$$

input `integrate(csch(5*x)*sinh(x),x)`

output `Integral(sinh(x)*csch(5*x), x)`

Maxima [F]

$$\int \operatorname{csch}(5x) \sinh(x) dx = \int \operatorname{csch}(5x) \sinh(x) dx$$

input `integrate(csch(5*x)*sinh(x),x, algorithm="maxima")`

output

```
1/10*(-1)^(3/5)*log((-1)^(1/5) + e^(-2*x)) + 1/10*sqrt(5)*(-1)^(3/5)*log((
sqrt(5)*(-1)^(1/5) + (-1)^(1/5)*sqrt(2*sqrt(5) - 10) + (-1)^(1/5) - 4*e^(-
2*x))/(sqrt(5)*(-1)^(1/5) - (-1)^(1/5)*sqrt(2*sqrt(5) - 10) + (-1)^(1/5) -
4*e^(-2*x)))/sqrt(2*sqrt(5) - 10) - 1/10*sqrt(5)*(-1)^(3/5)*log((sqrt(5)*
(-1)^(1/5) - (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*e^(-2*x))/(
sqrt(5)*(-1)^(1/5) + (-1)^(1/5)*sqrt(-2*sqrt(5) - 10) - (-1)^(1/5) + 4*e^(-
2*x)))/sqrt(-2*sqrt(5) - 10) - 1/10*log(-(sqrt(5)*(-1)^(1/5) + (-1)^(1/5)
)*e^(-2*x) + 2*(-1)^(2/5) + 2*e^(-4*x))/(sqrt(5)*(-1)^(2/5) + (-1)^(2/5))
+ 1/10*log((sqrt(5)*(-1)^(1/5) - (-1)^(1/5))*e^(-2*x) + 2*(-1)^(2/5) + 2*e
^(-4*x))/(sqrt(5)*(-1)^(2/5) - (-1)^(2/5)) - 1/10*integrate((e^(3*x) + 2*e
^(2*x) + 3*e^x + 4)*e^x/(e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) - 1/10
*integrate((e^(3*x) - 2*e^(2*x) + 3*e^x - 4)*e^x/(e^(4*x) - e^(3*x) + e^(2
*x) - e^x + 1), x) + 1/10*log(e^x + 1) + 1/10*log(e^x - 1)
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.85

$$\int \operatorname{csch}(5x) \sinh(x) dx = \frac{1}{10} \sqrt{-2\sqrt{5} + 10} \arctan \left(-\frac{\sqrt{5} - 4e^{(2x)} - 1}{\sqrt{2\sqrt{5} + 10}} \right) - \frac{1}{10} \sqrt{2\sqrt{5} + 10} \arctan \left(\frac{\sqrt{5} + 4e^{(2x)} + 1}{\sqrt{-2\sqrt{5} + 10}} \right)$$

input `integrate(csch(5*x)*sinh(x),x, algorithm="giac")`

output

```
1/10*sqrt(-2*sqrt(5) + 10)*arctan(-(sqrt(5) - 4*e^(2*x) - 1)/sqrt(2*sqrt(5)
) + 10)) - 1/10*sqrt(2*sqrt(5) + 10)*arctan((sqrt(5) + 4*e^(2*x) + 1)/sqrt
(-2*sqrt(5) + 10))
```

Mupad [B] (verification not implemented)

Time = 3.73 (sec) , antiderivative size = 282, normalized size of antiderivative = 3.52

$$\int \operatorname{csch}(5x) \sinh(x) dx$$

$$= 2 \operatorname{atan} \left(\frac{\frac{e^{2x}}{5} + \frac{9\sqrt{5}}{25} + \frac{6\sqrt{5}e^{2x}}{25} + \frac{4}{5}}{5e^{2x} \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}} + \frac{9\sqrt{5} \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}}}{5} + \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}} + \frac{9\sqrt{5}e^{2x} \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}}}{5}} \right) \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}}$$

$$+ \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} \left(\ln \left(\frac{9\sqrt{5}}{25} - \frac{e^{2x}}{5} + \frac{6\sqrt{5}e^{2x}}{25} - \frac{4}{5} - e^{2x} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 5i \right. \right.$$

$$\left. \left. + \frac{\sqrt{5} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 9i}{5} - \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 1i + \frac{\sqrt{5}e^{2x} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 9i}{5} \right) 1i \right.$$

$$- \ln \left(\frac{9\sqrt{5}}{25} - \frac{e^{2x}}{5} + \frac{6\sqrt{5}e^{2x}}{25} - \frac{4}{5} + e^{2x} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 5i - \frac{\sqrt{5} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 9i}{5} \right.$$

$$\left. \left. + \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 1i - \frac{\sqrt{5}e^{2x} \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} 9i}{5} \right) 1i \right)$$

input

```
int(sinh(x)/sinh(5*x),x)
```


output

```
2*atan((exp(2*x)/5 + (9*5^(1/2))/25 + (6*5^(1/2)*exp(2*x))/25 + 4/5)/(5*exp(2*x)*(5^(1/2)/200 + 1/40)^(1/2) + (9*5^(1/2)*(5^(1/2)/200 + 1/40)^(1/2))/5 + (5^(1/2)/200 + 1/40)^(1/2) + (9*5^(1/2)*exp(2*x)*(5^(1/2)/200 + 1/40)^(1/2))/5))*(5^(1/2)/200 + 1/40)^(1/2) + (1/40 - 5^(1/2)/200)^(1/2)*(log((5^(1/2)*(1/40 - 5^(1/2)/200)^(1/2)*9i)/5 - exp(2*x)*(1/40 - 5^(1/2)/200)^(1/2)*5i - exp(2*x)/5 + (9*5^(1/2))/25 - (1/40 - 5^(1/2)/200)^(1/2)*1i + (6*5^(1/2)*exp(2*x))/25 + (5^(1/2)*exp(2*x)*(1/40 - 5^(1/2)/200)^(1/2)*9i)/5 - 4/5)*1i - log(exp(2*x)*(1/40 - 5^(1/2)/200)^(1/2)*5i - exp(2*x)/5 - (5^(1/2)*(1/40 - 5^(1/2)/200)^(1/2)*9i)/5 + (9*5^(1/2))/25 + (1/40 - 5^(1/2)/200)^(1/2)*1i + (6*5^(1/2)*exp(2*x))/25 - (5^(1/2)*exp(2*x)*(1/40 - 5^(1/2)/200)^(1/2)*9i)/5 - 4/5)*1i)
```

Reduce [F]

$$\int \operatorname{csch}(5x) \sinh(x) dx = \int \operatorname{csch}(5x) \sinh(x) dx$$

input

```
int(csch(5*x)*sinh(x),x)
```

output

```
int(csch(5*x)*sinh(x),x)
```

3.70 $\int \operatorname{csch}(6x) \sinh(x) dx$

Optimal result	529
Mathematica [A] (verified)	529
Rubi [A] (verified)	530
Maple [C] (verified)	531
Fricas [B] (verification not implemented)	532
Sympy [F]	533
Maxima [F]	533
Giac [B] (verification not implemented)	533
Mupad [B] (verification not implemented)	534
Reduce [B] (verification not implemented)	534

Optimal result

Integrand size = 7, antiderivative size = 36

$$\int \operatorname{csch}(6x) \sinh(x) dx = \frac{1}{6} \arctan(\sinh(x)) + \frac{1}{6} \arctan(2 \sinh(x)) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{2\sqrt{3}}$$

output `1/6*arctan(sinh(x))+1/6*arctan(2*sinh(x))-1/6*arctan(2/3*sinh(x)*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \operatorname{csch}(6x) \sinh(x) dx = \frac{1}{6} \left(\arctan(\sinh(x)) + \arctan(2 \sinh(x)) - \sqrt{3} \arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right) \right)$$

input `Integrate[Csch[6*x]*Sinh[x],x]`

output `(ArcTan[Sinh[x]] + ArcTan[2*Sinh[x]] - Sqrt[3]*ArcTan[(2*Sinh[x])/Sqrt[3]])/6`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4878, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(x) \operatorname{csch}(6x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(ix)}{\sin(6ix)} dx \\
 & \quad \downarrow \text{4878} \\
 & \int \frac{1}{2(16 \sinh^6(x) + 32 \sinh^4(x) + 19 \sinh^2(x) + 3)} d \sinh(x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{1}{16 \sinh^6(x) + 32 \sinh^4(x) + 19 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow \text{2460} \\
 & \frac{1}{2} \int \left(\frac{2}{3(4 \sinh^2(x) + 1)} - \frac{2}{4 \sinh^2(x) + 3} + \frac{1}{3(\sinh^2(x) + 1)} \right) d \sinh(x) \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{1}{3} \arctan(\sinh(x)) + \frac{1}{3} \arctan(2 \sinh(x)) - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{3}}\right)}{\sqrt{3}} \right)
 \end{aligned}$$

input `Int [Csch [6*x]*Sinh [x] , x]`

output `(ArcTan [Sinh [x]]/3 + ArcTan [2*Sinh [x]]/3 - ArcTan [(2*Sinh [x])/Sqrt [3]]/Sqrt [3])/2`

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_)*(P_x_)^(p_), x_Symbol] := With[{Qx = Factor[P_x /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[P_x, x^2] && GtQ[Expon[P_x, x], 2] && !BinomialQ[P_x, x] && !TrinomialQ[P_x, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4878 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Sin[v], x]}, d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Sin[v]/d, u/Cos[v], x], x], x, Sin[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Sin[v], x], u/Cos[v], x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.56

method	result
risch	$\frac{i \ln(e^x+i)}{6} - \frac{i \ln(e^x-i)}{6} + \frac{i \ln(e^{2x}+ie^x-1)}{12} - \frac{i \ln(e^{2x}-ie^x-1)}{12} + \frac{i\sqrt{3} \ln(e^{2x}-i\sqrt{3}e^x-1)}{12} - \frac{i\sqrt{3} \ln(e^{2x}+i\sqrt{3}e^x-1)}{12}$

input `int(csch(6*x)*sinh(x), x, method=_RETURNVERBOSE)`

output

```
1/6*I*ln(exp(x)+I)-1/6*I*ln(exp(x)-I)+1/12*I*ln(exp(2*x)+I*exp(x)-1)-1/12*
I*ln(exp(2*x)-I*exp(x)-1)+1/12*I*3^(1/2)*ln(exp(2*x)-I*3^(1/2)*exp(x)-1)-1
/12*I*3^(1/2)*ln(exp(2*x)+I*3^(1/2)*exp(x)-1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(26) = 52$.

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.97

$$\int \operatorname{csch}(6x) \sinh(x) dx$$

$$= -\frac{1}{6} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \cosh(x) + \frac{1}{3} \sqrt{3} \sinh(x) \right)$$

$$+ \frac{1}{6} \sqrt{3} \arctan \left(-\frac{\sqrt{3} \cosh(x)^2 + 2\sqrt{3} \cosh(x) \sinh(x) + \sqrt{3} \sinh(x)^2 + 2\sqrt{3}}{3(\cosh(x) - \sinh(x))} \right)$$

$$- \frac{1}{6} \arctan \left(-\frac{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2}{\cosh(x) - \sinh(x)} \right)$$

$$+ \frac{1}{2} \arctan(\cosh(x) + \sinh(x))$$

input

```
integrate(csch(6*x)*sinh(x),x, algorithm="fricas")
```

output

```
-1/6*sqrt(3)*arctan(1/3*sqrt(3)*cosh(x) + 1/3*sqrt(3)*sinh(x)) + 1/6*sqrt(
3)*arctan(-1/3*(sqrt(3)*cosh(x)^2 + 2*sqrt(3)*cosh(x)*sinh(x) + sqrt(3)*si
nh(x)^2 + 2*sqrt(3))/(cosh(x) - sinh(x))) - 1/6*arctan(-(cosh(x)^2 + 2*cos
h(x)*sinh(x) + sinh(x)^2)/(cosh(x) - sinh(x))) + 1/2*arctan(cosh(x) + sinh
(x))
```

Sympy [F]

$$\int \operatorname{csch}(6x) \sinh(x) dx = \int \sinh(x) \operatorname{csch}(6x) dx$$

input `integrate(csch(6*x)*sinh(x),x)`

output `Integral(sinh(x)*csch(6*x), x)`

Maxima [F]

$$\int \operatorname{csch}(6x) \sinh(x) dx = \int \operatorname{csch}(6x) \sinh(x) dx$$

input `integrate(csch(6*x)*sinh(x),x, algorithm="maxima")`

output `-1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) + 1/3*arctan(e^x) + integrate(1/6*(e^(3*x) + e^x)/(e^(4*x) - e^(2*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. 2(26) = 52.

Time = 0.14 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.61

$$\int \operatorname{csch}(6x) \sinh(x) dx = \frac{1}{6} \pi - \frac{1}{12} \sqrt{3} \left(\pi + 2 \arctan \left(\frac{1}{3} \sqrt{3} (e^{2x} - 1) e^{-x} \right) \right) + \frac{1}{6} \arctan \left((e^{2x} - 1) e^{-x} \right) + \frac{1}{6} \arctan \left(\frac{1}{2} (e^{2x} - 1) e^{-x} \right)$$

input `integrate(csch(6*x)*sinh(x),x, algorithm="giac")`

output

$$\frac{1}{6}\pi - \frac{1}{12}\sqrt{3}(\pi + 2\arctan(\frac{1}{3}\sqrt{3}(e^{2x} - 1)e^{-x})) + \frac{1}{6}\arctan((e^{2x} - 1)e^{-x}) + \frac{1}{6}\arctan(\frac{1}{2}(e^{2x} - 1)e^{-x})$$
Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.14

$$\int \operatorname{csch}(6x) \sinh(x) dx = \frac{\operatorname{atan}(e^x)}{3} - \frac{\operatorname{atan}(e^{-x} - e^x)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}e^x}{3} - \frac{\sqrt{3}e^{-x}}{3}\right)}{6}$$

input

$$\operatorname{int}(\sinh(x)/\sinh(6*x), x)$$

output

$$\frac{\operatorname{atan}(\exp(x))}{3} - \frac{\operatorname{atan}(\exp(-x) - \exp(x))}{6} - \frac{(3^{1/2}) \operatorname{atan}((3^{1/2}) \exp(x))}{3} - \frac{(3^{1/2}) \operatorname{atan}(-x)}{6}$$
Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.81

$$\int \operatorname{csch}(6x) \sinh(x) dx = \frac{\operatorname{atan}(e^x)}{3} + \frac{\operatorname{atan}(2e^x - \sqrt{3})}{6} + \frac{\operatorname{atan}(2e^x + \sqrt{3})}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2e^x - 1}{\sqrt{3}}\right)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2e^x + 1}{\sqrt{3}}\right)}{6}$$

input

$$\operatorname{int}(\operatorname{csch}(6*x)*\sinh(x), x)$$

output

$$(2*\operatorname{atan}(e^{**x}) + \operatorname{atan}(2*e^{**x} - \operatorname{sqrt}(3)) + \operatorname{atan}(2*e^{**x} + \operatorname{sqrt}(3)) - \operatorname{sqrt}(3)*\operatorname{atan}((2*e^{**x} - 1)/\operatorname{sqrt}(3)) - \operatorname{sqrt}(3)*\operatorname{atan}((2*e^{**x} + 1)/\operatorname{sqrt}(3)))/6$$

3.71 $\int \operatorname{csch}(c + bx) \sinh(a + bx) dx$

Optimal result	535
Mathematica [A] (verified)	535
Rubi [C] (verified)	536
Maple [B] (verified)	537
Fricas [B] (verification not implemented)	538
Sympy [F]	538
Maxima [B] (verification not implemented)	538
Giac [A] (verification not implemented)	539
Mupad [B] (verification not implemented)	539
Reduce [B] (verification not implemented)	540

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = x \cosh(a - c) + \frac{\log(\sinh(c + bx)) \sinh(a - c)}{b}$$

output `x*cosh(a-c)+ln(sinh(b*x+c))*sinh(a-c)/b`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = x \cosh(a - c) + \frac{\log(\sinh(c + bx)) \sinh(a - c)}{b}$$

input `Integrate[Csch[c + b*x]*Sinh[a + b*x],x]`

output `x*Cosh[a - c] + (Log[Sinh[c + b*x]]*Sinh[a - c])/b`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6160, 24, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{csch}(bx + c) dx \\
 & \quad \downarrow \text{6160} \\
 & \sinh(a - c) \int \coth(c + bx) dx + \cosh(a - c) \int 1 dx \\
 & \quad \downarrow \text{24} \\
 & \sinh(a - c) \int \coth(c + bx) dx + x \cosh(a - c) \\
 & \quad \downarrow \text{3042} \\
 & x \cosh(a - c) + \sinh(a - c) \int -i \tan\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & x \cosh(a - c) - i \sinh(a - c) \int \tan\left(\frac{1}{2}(2ic + \pi) + ibx\right) dx \\
 & \quad \downarrow \text{3956} \\
 & x \cosh(a - c) + \frac{\sinh(a - c) \log(-i \sinh(bx + c))}{b}
 \end{aligned}$$

input

```
Int[Csch[c + b*x]*Sinh[a + b*x],x]
```

output

```
x*Cosh[a - c] + (Log[(-I)*Sinh[c + b*x]]*Sinh[a - c])/b
```

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6160 `Int[Csch[w_]^(n_.)*Sinh[v_], x_Symbol] := Simp[Sinh[v - w] Int[Coth[w]*Csch[w]^(n - 1), x], x] + Simp[Cosh[v - w] Int[Csch[w]^(n - 1), x], x] /; GTQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(26) = 52$.

Time = 0.10 (sec) , antiderivative size = 150, normalized size of antiderivative = 5.77

method	result
risch	$x e^{a-c} - e^{-a-c} e^{2a} x + e^{-a-c} e^{2c} x - \frac{e^{-a-c} e^{2a} a}{b} + \frac{e^{-a-c} e^{2c} a}{b} + \frac{\ln(e^{2bx+2a} - e^{2a-2c}) e^{-a-c} e^{2a}}{2b} - \frac{\ln(e^{2bx+2a} - e^{2a-2c})}{2b}$

input `int(csch(b*x+c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `x*exp(a-c)-exp(-a-c)*exp(2*a)*x+exp(-a-c)*exp(2*c)*x-1/b*exp(-a-c)*exp(2*a)*a+1/b*exp(-a-c)*exp(2*c)*a+1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-a-c)*exp(2*a)-1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(26) = 52$.

Time = 0.09 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.31

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx$$

$$= \frac{2bx + (\cosh(-a + c))^2 - 2\cosh(-a + c)\sinh(-a + c) + \sinh(-a + c)^2 - 1}{2(b\cosh(-a + c) - b\sinh(-a + c))} \log\left(\frac{2\sinh(bx+c)}{\cosh(bx+c) - \sinh(bx+c)}\right)$$

input `integrate(csch(b*x+c)*sinh(b*x+a),x, algorithm="fricas")`

output `1/2*(2*b*x + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c)))/(b*cosh(-a + c) - b*sinh(-a + c))`

Sympy [F]

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}(bx + c) dx$$

input `integrate(csch(b*x+c)*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*csch(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(26) = 52$.

Time = 0.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 3.23

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} + \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} + \frac{(bx + a)e^{(a-c)}}{b}$$

input `integrate(csch(b*x+c)*sinh(b*x+a),x, algorithm="maxima")`

output `1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + (b*x + a)*e^(a - c)/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.85

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = x e^{(-a+c)} + \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(2bx+2c)} - 1|)}{2b}$$

input `integrate(csch(b*x+c)*sinh(b*x+a),x, algorithm="giac")`

output `x*e^(-a + c) + 1/2*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(abs(e^(2*b*x + 2*c) - 1))/b`

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx = x e^{c-a} + \frac{e^{2c-2a} \ln(e^{2a} e^{2bx} - e^{2a} e^{-2c}) (2b e^{3a-3c} - 2b e^{a-c})}{4b^2}$$

input `int(sinh(a + b*x)/sinh(c + b*x),x)`

output `x*exp(c - a) + (exp(2*c - 2*a)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp(-2*c)))*(2*b*exp(3*a - 3*c) - 2*b*exp(a - c))/(4*b^2)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 3.38

$$\int \operatorname{csch}(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{2a} \log(e^{bx+c} - 1) + e^{2a} \log(e^{bx+c} + 1) - e^{2c} \log(e^{bx+c} - 1) - e^{2c} \log(e^{bx+c} + 1) + 2e^{2c}bx}{2e^{a+cb}}$$

input `int(csch(b*x+c)*sinh(b*x+a),x)`

output `(e**(2*a)*log(e**(b*x + c) - 1) + e**(2*a)*log(e**(b*x + c) + 1) - e**(2*c)*log(e**(b*x + c) - 1) - e**(2*c)*log(e**(b*x + c) + 1) + 2*e**(2*c)*b*x)/(2*e**(a + c)*b)`

3.72 $\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx$

Optimal result	541
Mathematica [C] (verified)	541
Rubi [A] (verified)	542
Maple [B] (verified)	544
Fricas [B] (verification not implemented)	544
Sympy [F]	545
Maxima [B] (verification not implemented)	546
Giac [B] (verification not implemented)	546
Mupad [B] (verification not implemented)	547
Reduce [B] (verification not implemented)	547

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(a - c)}{b} - \frac{\operatorname{csch}(c + bx) \sinh(a - c)}{b}$$

output

```
-arctanh(cosh(b*x+c))*cosh(a-c)/b-csch(b*x+c)*sinh(a-c)/b
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = -\frac{2i \operatorname{arctan}\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh(c) \cosh\left(\frac{bx}{2}\right) + \sinh(c) \sinh\left(\frac{bx}{2}\right)\right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a - c)}{b} - \frac{\operatorname{csch}(c + bx) \sinh(a - c)}{b}$$

input `Integrate[Csch[c + b*x]^2*Sinh[a + b*x],x]`

output
$$\frac{((-2*I)*ArcTan[((Cosh[c] - Sinh[c])*(Cosh[c]*Cosh[(b*x)/2] + Sinh[c]*Sinh[(b*x)/2]))/(I*Cosh[c]*Cosh[(b*x)/2] - I*Cosh[(b*x)/2]*Sinh[c])]*Cosh[a - c])/b - (Csch[c + b*x]*Sinh[a - c])/b}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6160, 3042, 26, 3086, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(a + bx) \operatorname{csch}^2(bx + c) dx \\ & \quad \downarrow 6160 \\ & \cosh(a - c) \int \operatorname{csch}(c + bx) dx + \sinh(a - c) \int \operatorname{coth}(c + bx) \operatorname{csch}(c + bx) dx \\ & \quad \downarrow 3042 \\ & \cosh(a - c) \int i \operatorname{csc}(ic + ibx) dx + \sinh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow 26 \\ & i \cosh(a - c) \int \operatorname{csc}(ic + ibx) dx + \sinh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow 3086 \\ & i \cosh(a - c) \int \operatorname{csc}(ic + ibx) dx - \frac{i \sinh(a - c) \int 1 d(-i \operatorname{csch}(c + bx))}{b} \\ & \quad \downarrow 24 \\ & -\frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} + i \cosh(a - c) \int \operatorname{csc}(ic + ibx) dx \\ & \quad \downarrow 4257 \end{aligned}$$

$$-\frac{\cosh(a-c)\operatorname{arctanh}(\cosh(bx+c))}{b} - \frac{\sinh(a-c)\operatorname{csch}(bx+c)}{b}$$

input `Int[Csch[c + b*x]^2*Sinh[a + b*x],x]`

output `-((ArcTanh[Cosh[c + b*x]]*Cosh[a - c])/b) - (Csch[c + b*x]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6160 `Int[Csch[w_]^(n_)*Sinh[v_], x_Symbol] := Simp[Sinh[v - w] Int[Coth[w]*Csch[w]^(n-1), x], x] + Simp[Cosh[v - w] Int[Csch[w]^(n-1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. $2(36) = 72$.

Time = 0.22 (sec) , antiderivative size = 172, normalized size of antiderivative = 4.78

method	result
risch	$\frac{e^{bx+a}(e^{2a}-e^{2c})}{b(-e^{2bx+2a+2c}+e^{2a})} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(csch(b*x+c)^2*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/b*exp(b*x+a)*(exp(2*a)-exp(2*c))/(-exp(2*b*x+2*a+2*c)+exp(2*a))+1/2*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*a)+1/2*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*c)-1/2*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*a)-1/2*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 617 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 617, normalized size of antiderivative = 17.14

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a),x, algorithm="fricas")`

output

```

1/2*(4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - 2*cosh(b*x + c)*sinh(-a +
c)^2 - 2*(cosh(-a + c)^2 - 1)*cosh(b*x + c) - ((cosh(-a + c)^2 + 1)*cosh(
b*x + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^
2 + 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a +
c)^2 - 2*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-
a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c) - 2*(cosh(b*x
+ c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) - 1)*log(cosh(b*x + c) +
sinh(b*x + c) + 1) + ((cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (cosh(-a + c
)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^2 +
(cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(2*cosh(b*x + c)
*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^
2 + 1)*cosh(b*x + c))*sinh(b*x + c) - 2*(cosh(b*x + c)^2*cosh(-a + c) - co
sh(-a + c))*sinh(-a + c) - 1)*log(cosh(b*x + c) + sinh(b*x + c) - 1) - 2*(
cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*
x + c))/(b*cosh(b*x + c)^2*cosh(-a + c) + (b*cosh(-a + c) - b*sinh(-a + c)
)*sinh(b*x + c)^2 - b*cosh(-a + c) + 2*(b*cosh(b*x + c)*cosh(-a + c) - b*c
osh(b*x + c)*sinh(-a + c))*sinh(b*x + c) - (b*cosh(b*x + c)^2 - b)*sinh(-a
+ c))

```

Sympy [F]

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}^2(bx + c) dx$$

input

```
integrate(csch(b*x+c)**2*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*csch(b*x + c)**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.86

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = -\frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} + \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} + \frac{(e^{(2a)} - e^{(2c)})e^{(-bx-a)}}{b(e^{(-2bx)} - e^{(2c)})}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + (e^(2*a) - e^(2*c))*e^(-b*x - a)/(b*(e^(-2*b*x) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 109 vs. $2(36) = 72$.

Time = 0.14 (sec) , antiderivative size = 109, normalized size of antiderivative = 3.03

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx = -\frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(bx+c)} + 1)}{2b} + \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+c)} - 1|)}{2b} - \frac{(e^{(bx+2a)} - e^{(bx+2c)})e^{(-a)}}{b(e^{(2bx+2c)} - 1)}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a),x, algorithm="giac")`

output `-1/2*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(e^(b*x + c) + 1)/b + 1/2*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + c) - 1))/b - (e^(b*x + 2*a) - e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) - 1))`

Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.33

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{a+bx} (e^{2a-2c} - 1)}{b (e^{2a-2c} - e^{2a+2bx})}$$

$$- \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2 + e^{2a} e^{-2c} \sqrt{-b^2}})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{-b^2}}$$

input `int(sinh(a + b*x)/sinh(c + b*x)^2,x)`output `(exp(a + b*x)*(exp(2*a - 2*c) - 1))/(b*(exp(2*a - 2*c) - exp(2*a + 2*b*x))) - (atan((exp(-a)*exp(2*c)*exp(b*x)*((-b^2)^(1/2) + exp(2*a)*exp(-2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c)*(2*exp(2*a)*exp(-2*c) + exp(4*a)*exp(-4*c) + 1))^(1/2)))*(exp(2*c - 2*a)*(2*exp(2*a - 2*c) + exp(4*a - 4*c) + 1))^(1/2))/(-b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 208, normalized size of antiderivative = 5.78

$$\int \operatorname{csch}^2(c + bx) \sinh(a + bx) dx$$

$$= \frac{e^{2bx+2a+2c} \log(e^{bx+c} - 1) - e^{2bx+2a+2c} \log(e^{bx+c} + 1) + e^{2bx+4c} \log(e^{bx+c} - 1) - e^{2bx+4c} \log(e^{bx+c} + 1) - 2e^{a+c} b (e^{2bx+2a+2c} - 1)}{2e^{a+c} b (e^{2bx+2a+2c} - 1)}$$

input `int(csch(b*x+c)^2*sinh(b*x+a),x)`output `(e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) - e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) + 1) + e**(2*b*x + 4*c)*log(e**(b*x + c) - 1) - e**(2*b*x + 4*c)*log(e**(b*x + c) + 1) - 2*e**(2*a + b*x + c) + 2*e**(b*x + 3*c) - e**(2*a)*log(e**(b*x + c) - 1) + e**(2*a)*log(e**(b*x + c) + 1) - e**(2*c)*log(e**(b*x + c) - 1) + e**(2*c)*log(e**(b*x + c) + 1))/(2*e**(a + c)*b*(e**(2*b*x + 2*c) - 1))`

3.73 $\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx$

Optimal result	548
Mathematica [A] (verified)	548
Rubi [A] (verified)	549
Maple [A] (verified)	551
Fricas [B] (verification not implemented)	551
Sympy [F]	552
Maxima [B] (verification not implemented)	552
Giac [A] (verification not implemented)	553
Mupad [F(-1)]	553
Reduce [B] (verification not implemented)	554

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = -\frac{\cosh(a - c) \coth(c + bx)}{b} - \frac{\operatorname{csch}^2(c + bx) \sinh(a - c)}{2b}$$

output

```
-cosh(a-c)*coth(b*x+c)/b-1/2*csch(b*x+c)^2*sinh(a-c)/b
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\begin{aligned} & \int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx \\ &= -\frac{(\cosh(a) - \cosh(a - c) \cosh(c + 2bx)) \operatorname{csch}(c) \operatorname{csch}^2(c + bx)}{2b} \end{aligned}$$

input

```
Integrate[Csch[c + b*x]^3*Sinh[a + b*x],x]
```

output

```
-1/2*((Cosh[a] - Cosh[a - c]*Cosh[c + 2*b*x])*Csch[c]*Csch[c + b*x]^2)/b
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {6160, 3042, 25, 26, 3086, 15, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{csch}^3(bx + c) dx \\
 & \quad \downarrow \text{6160} \\
 & \cosh(a - c) \int \operatorname{csch}^2(c + bx) dx + \sinh(a - c) \int \operatorname{coth}(c + bx) \operatorname{csch}^2(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \cosh(a - c) \int -\operatorname{csc}(ic + ibx)^2 dx + \sinh(a - c) \int i \sec\left(ic + ibx - \frac{\pi}{2}\right)^2 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & \sinh(a - c) \int i \sec\left(ic + ibx - \frac{\pi}{2}\right)^2 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx - \cosh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{26} \\
 & i \sinh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^2 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx - \cosh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{3086} \\
 & \frac{\sinh(a - c) \int -i \operatorname{csch}(c + bx) d(-i \operatorname{csch}(c + bx))}{b} - \cosh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{\sinh(a - c) \operatorname{csch}^2(bx + c)}{2b} - \cosh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sinh(a - c) \operatorname{csch}^2(bx + c)}{2b} - \frac{i \cosh(a - c) \int 1 d(-i \operatorname{coth}(c + bx))}{b} \\
 & \quad \downarrow \text{24}
 \end{aligned}$$

$$-\frac{\cosh(a-c)\coth(bx+c)}{b} - \frac{\sinh(a-c)\operatorname{csch}^2(bx+c)}{2b}$$

input `Int[Csch[c + b*x]^3*Sinh[a + b*x],x]`

output `-((Cosh[a - c]*Coth[c + b*x])/b) - (Csch[c + b*x]^2*Sinh[a - c])/(2*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 6160

```
Int[Csch[w_]^(n_.)*Sinh[v_], x_Symbol] := Simp[Sinh[v - w] Int[Coth[w]*Csch[w]^(n - 1), x], x] + Simp[Cosh[v - w] Int[Csch[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]
```

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.46

method	result	size
risch	$\frac{(-2e^{2bx+2a+2c}+e^{2a}+e^{2c})e^{3a-c}}{(-e^{2bx+2a+2c}+e^{2a})^2b}$	57
parallelrisch	$-\frac{\left(\sinh(bx+a)\left(-\frac{\operatorname{sech}\left(\frac{bx}{2}+\frac{c}{2}\right)^2}{2}+1\right)\operatorname{csch}\left(\frac{bx}{2}+\frac{c}{2}\right)+\operatorname{sech}\left(\frac{bx}{2}+\frac{c}{2}\right)\cosh(bx+a)\right)\operatorname{csch}\left(\frac{bx}{2}+\frac{c}{2}\right)}{4b}$	63

input

```
int(csch(b*x+c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
1/(-exp(2*b*x+2*a+2*c)+exp(2*a))^2/b*(-2*exp(2*b*x+2*a+2*c)+exp(2*a)+exp(2*c))*exp(3*a-c)
```

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(37) = 74$.

Time = 0.08 (sec) , antiderivative size = 246, normalized size of antiderivative = 6.31

$$\int \operatorname{csch}^3(c+bx) \sinh(a+bx) dx =$$

$$-\frac{b \cosh(bx+c)^3 \cosh(-a+c)^2 - b \cosh(bx+c) \cosh(-a+c)^2 + (b \cosh(-a+c)^2 - b \sinh(-a+c))}{4b^2}$$

input

```
integrate(csch(b*x+c)^3*sinh(b*x+a),x, algorithm="fricas")
```


output

```
-2*((2*cosh(-a + c) - sinh(-a + c))*sinh(b*x + c) - cosh(b*x + c)*sinh(-a
+ c))/(b*cosh(b*x + c)^3*cosh(-a + c)^2 - b*cosh(b*x + c)*cosh(-a + c)^2 +
(b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^3 + 3*(b*cosh(b*x + c
)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^2 - (b*co
sh(b*x + c)^3 - b*cosh(b*x + c))*sinh(-a + c)^2 + 3*(b*cosh(b*x + c)^2*cos
h(-a + c)^2 - b*cosh(-a + c)^2 - (b*cosh(b*x + c)^2 - b)*sinh(-a + c)^2)*s
inh(b*x + c))
```

Sympy [F]

$$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}^3(bx + c) dx$$

input

```
integrate(csch(b*x+c)**3*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*csch(b*x + c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(37) = 74$.

Time = 0.04 (sec) , antiderivative size = 131, normalized size of antiderivative = 3.36

$$\begin{aligned} \int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = & -\frac{2e^{(-2bx+3c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})} \\ & + \frac{e^{(2a+3c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})} \\ & + \frac{e^{(5c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})} \end{aligned}$$

input

```
integrate(csch(b*x+c)^3*sinh(b*x+a),x, algorithm="maxima")
```

output

```
-2*e^(-2*b*x + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c))) + e^(2*a + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c))) + e^(5*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c)))
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.36

$$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = -\frac{(2e^{(2bx+2a+2c)} - e^{(2a)} - e^{(2c)})e^{(-a-c)}}{b(e^{(2bx+2c)} - 1)^2}$$

input

```
integrate(csch(b*x+c)^3*sinh(b*x+a),x, algorithm="giac")
```

output

```
-(2*e^(2*b*x + 2*a + 2*c) - e^(2*a) - e^(2*c))*e^(-a - c)/(b*(e^(2*b*x + 2*c) - 1)^2)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\sinh(c + bx)^3} dx$$

input

```
int(sinh(a + b*x)/sinh(c + b*x)^3,x)
```

output

```
int(sinh(a + b*x)/sinh(c + b*x)^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.41

$$\int \operatorname{csch}^3(c + bx) \sinh(a + bx) dx = \frac{e^c(-e^{4bx+2a+2c} + 1)}{e^{ab}(e^{4bx+4c} - 2e^{2bx+2c} + 1)}$$

input `int(csch(b*x+c)^3*sinh(b*x+a),x)`

output `(e**c*(-e**(2*a + 4*b*x + 2*c) + 1))/(e**a*b*(e**(4*b*x + 4*c) - 2*e**(2*b*x + 2*c) + 1))`

3.74 $\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$

Optimal result	555
Mathematica [A] (verified)	555
Rubi [C] (verified)	556
Maple [B] (verified)	559
Fricas [B] (verification not implemented)	559
Sympy [F]	560
Maxima [B] (verification not implemented)	561
Giac [B] (verification not implemented)	561
Mupad [F(-1)]	562
Reduce [B] (verification not implemented)	562

Optimal result

Integrand size = 15, antiderivative size = 67

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx = \frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(a - c)}{2b} - \frac{\cosh(a - c) \coth(c + bx) \operatorname{csch}(c + bx)}{2b} - \frac{\operatorname{csch}^3(c + bx) \sinh(a - c)}{3b}$$

output `1/2*arctanh(cosh(b*x+c))*cosh(a-c)/b-1/2*cosh(a-c)*coth(b*x+c)*csch(b*x+c)/b-1/3*csch(b*x+c)^3*sinh(a-c)/b`

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.97

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx = \frac{12 \operatorname{arctanh}(\cosh(c) + \sinh(c) \tanh(\frac{bx}{2})) \cosh(a - c) - \operatorname{csch}^3(c + bx)(4 \sinh(a - c) + 3 \cosh(a - c) \sinh(a - c))}{12b}$$

input `Integrate[Csch[c + b*x]^4*Sinh[a + b*x],x]`

output

```
(12*ArcTanh[Cosh[c] + Sinh[c]*Tanh[(b*x)/2]]*Cosh[a - c] - Csch[c + b*x]^3
*(4*Sinh[a - c] + 3*Cosh[a - c]*Sinh[2*(c + b*x)]))/(12*b)
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {6160, 3042, 25, 26, 3086, 15, 4255, 26, 3042, 26, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sinh(a + bx) \operatorname{csch}^4(bx + c) dx \\
 & \quad \downarrow \text{6160} \\
 & \cosh(a - c) \int \operatorname{csch}^3(c + bx) dx + \sinh(a - c) \int \coth(c + bx) \operatorname{csch}^3(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \cosh(a - c) \int -i \operatorname{csc}(ic + ibx)^3 dx + \sinh(a - c) \int -\sec\left(ic + ibx - \frac{\pi}{2}\right)^3 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & \cosh(a - c) \int -i \operatorname{csc}(ic + ibx)^3 dx - \sinh(a - \\
 & \quad c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^3 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx \\
 & \quad \downarrow \text{26} \\
 & -i \cosh(a - c) \int \operatorname{csc}(ic + ibx)^3 dx - \sinh(a - \\
 & \quad c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^3 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx \\
 & \quad \downarrow \text{3086} \\
 & \frac{i \sinh(a - c) \int -\operatorname{csch}^2(c + bx) d(-i \operatorname{csch}(c + bx))}{b} - i \cosh(a - c) \int \operatorname{csc}(ic + ibx)^3 dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 15 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \int \csc(ic+ibx)^3 dx \\
& \downarrow 4255 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \\
& c) \left(\frac{1}{2} \int -i \operatorname{csch}(c+bx) dx - \frac{i \coth(bx+c) \operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 26 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \\
& c) \left(-\frac{1}{2} i \int \operatorname{csch}(c+bx) dx - \frac{i \coth(bx+c) \operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 3042 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \\
& c) \left(-\frac{1}{2} i \int i \csc(ic+ibx) dx - \frac{i \coth(bx+c) \operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 26 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \left(\frac{1}{2} \int \csc(ic+ibx) dx - \frac{i \coth(bx+c) \operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 4257 \\
& -\frac{\sinh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i \cosh(a-c) \\
& c) \left(\frac{i \operatorname{arctanh}(\cosh(bx+c))}{2b} - \frac{i \coth(bx+c) \operatorname{csch}(bx+c)}{2b} \right)
\end{aligned}$$

input

```
Int[Csch[c + b*x]^4*Sinh[a + b*x],x]
```

output

```
(-I)*Cosh[a - c]*(((I/2)*ArcTanh[Cosh[c + b*x]])/b - ((I/2)*Coth[c + b*x]*
Csch[c + b*x])/b) - (Csch[c + b*x]^3*Sinh[a - c])/(3*b)
```

Definitions of rubi rules used

- rule 15 $\text{Int}[(a_.)(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[a*(x^{(m+1)})/(m+1), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \ \text{Int}[Fx, x], x]$
- rule 26 $\text{Int}[(\text{Complex}[0, a_])*(Fx_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \ \text{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3086 $\text{Int}[(a_.)*\text{sec}[(e_.) + (f_.)(x_)]^{(m_.)}*((b_.)*\text{tan}[(e_.) + (f_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a/f \ \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e+f*x]], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$
- rule 4255 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c+d*x]*((b*\text{Csc}[c+d*x])^{(n-1)})/(d*(n-1)), x] + \text{Simp}[b^2*((n-2)/(n-1)) \ \text{Int}[(b*\text{Csc}[c+d*x])^{(n-2)}, x], x] \text{ ; FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c+d*x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$
- rule 6160 $\text{Int}[\text{Csch}[w_]^{(n_.)}*\text{Sinh}[v_], x_Symbol] \rightarrow \text{Simp}[\text{Sinh}[v-w] \ \text{Int}[\text{Coth}[w]*\text{Csch}[w]^{(n-1)}, x], x] + \text{Simp}[\text{Cosh}[v-w] \ \text{Int}[\text{Csch}[w]^{(n-1)}, x], x] \text{ ; GtQ}[n, 0] \ \&\& \ \text{NeQ}[w, v] \ \&\& \ \text{FreeQ}[v-w, x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(61) = 122$.

Time = 1.30 (sec) , antiderivative size = 235, normalized size of antiderivative = 3.51

method	result
risch	$-\frac{e^{bx+a}(-3e^{4bx+6a+4c}-3e^{4bx+4a+6c}-8e^{2bx+6a+2c}+8e^{2bx+4a+4c}+3e^{6a}+3e^{4a+2c})}{6b(-e^{2bx+2a+2c}+e^{2a})^3} + \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{4b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{4b}$

input `int(csch(b*x+c)^4*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output
$$-1/6*\exp(b*x+a)*(-3*\exp(4*b*x+6*a+4*c)-3*\exp(4*b*x+4*a+6*c)-8*\exp(2*b*x+6*a+2*c)+8*\exp(2*b*x+4*a+4*c)+3*\exp(6*a)+3*\exp(4*a+2*c))/b/(-\exp(2*b*x+2*a+2*c)+\exp(2*a))^3+1/4*\ln(\exp(b*x+a)+\exp(a-c))/b*\exp(-a-c)*\exp(2*a)+1/4*\ln(\exp(b*x+a)+\exp(a-c))/b*\exp(-a-c)*\exp(2*c)-1/4*\ln(\exp(b*x+a)-\exp(a-c))/b*\exp(-a-c)*\exp(2*a)-1/4*\ln(\exp(b*x+a)-\exp(a-c))/b*\exp(-a-c)*\exp(2*c)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2696 vs. $2(61) = 122$.

Time = 0.13 (sec) , antiderivative size = 2696, normalized size of antiderivative = 40.24

$$\int \operatorname{csch}^4(c+bx) \sinh(a+bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x+c)^4*sinh(b*x+a),x, algorithm="fricas")`

output

```
-1/12*(6*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^5 + 6*(cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^5 - 30*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^4 + 16*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + 4*(15*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^2 + 4)*sinh(-a + c)^2 + 4*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^2*cosh(-a + c) + 4*cosh(-a + c))*sinh(-a + c) - 4)*sinh(b*x + c)^3 + 12*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 + 4*cosh(b*x + c))*sinh(-a + c)^2 + 4*(cosh(-a + c)^2 - 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh(-a + c) + 4*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^2 + 2*(3*cosh(b*x + c)^5 + 8*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 - 6*(cosh(-a + c)^2 + 1)*cosh(b*x + c) - 3*((cosh(-a + c)^2 + 1)*cosh(b*x + c)^6 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^6 - 6*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^5 - 3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^4 + 3*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(5*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) - 1)*sinh(b*x + c)^4 + 4*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 - 3*(cosh(-a + c)^2 + 1)*cosh(b*...
```

SymPy [F]

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}^4(bx + c) dx$$

input

```
integrate(csch(b*x+c)**4*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*csch(b*x + c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(61) = 122$.

Time = 0.04 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.76

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$$

$$= \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{4b} - \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{4b}$$

$$+ \frac{3(e^{(2a+4c)} + e^{(6c)})e^{(-bx-a)} + 8(e^{(4a+2c)} - e^{(2a+4c)})e^{(-3bx-3a)} - 3(e^{(6a)} + e^{(4a+2c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+4c)} - 3e^{(-4bx+2c)} + e^{(-6bx)} - e^{(6c)})}$$

input `integrate(csch(b*x+c)^4*sinh(b*x+a),x, algorithm="maxima")`

output

```
1/4*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b - 1/4*(e^(2*a) +
e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/6*(3*(e^(2*a + 4*c) + e^(6*c)
))*e^(-b*x - a) + 8*(e^(4*a + 2*c) - e^(2*a + 4*c))*e^(-3*b*x - 3*a) - 3*(
e^(6*a) + e^(4*a + 2*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 4*c) - 3*e^(-
4*b*x + 2*c) + e^(-6*b*x) - e^(6*c)))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 161 vs. $2(61) = 122$.

Time = 0.12 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.40

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$$

$$= \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(bx+c)} + 1)}{4b} - \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+c)} - 1|)}{4b}$$

$$- \frac{(3e^{(5bx+2a+4c)} + 3e^{(5bx+6c)} + 8e^{(3bx+2a+2c)} - 8e^{(3bx+4c)} - 3e^{(bx+2a)} - 3e^{(bx+2c)})e^{(-a)}}{6b(e^{(2bx+2c)} - 1)^3}$$

input `integrate(csch(b*x+c)^4*sinh(b*x+a),x, algorithm="giac")`

output

```
1/4*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(e^(b*x + c) + 1)/b - 1/4*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + c) - 1))/b - 1/6*(3*e^(5*b*x + 2*a + 4*c) + 3*e^(5*b*x + 6*c) + 8*e^(3*b*x + 2*a + 2*c) - 8*e^(3*b*x + 4*c) - 3*e^(b*x + 2*a) - 3*e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) - 1)^3)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\sinh(c + bx)^4} dx$$

input

```
int(sinh(a + b*x)/sinh(c + b*x)^4,x)
```

output

```
int(sinh(a + b*x)/sinh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 478, normalized size of antiderivative = 7.13

$$\int \operatorname{csch}^4(c + bx) \sinh(a + bx) dx$$

$$= \frac{-3e^{6bx+2a+6c}\log(e^{bx+c} - 1) + 3e^{6bx+2a+6c}\log(e^{bx+c} + 1) - 3e^{6bx+8c}\log(e^{bx+c} - 1) + 3e^{6bx+8c}\log(e^{bx+c} + 1)}{b}$$

input

```
int(csch(b*x+c)^4*sinh(b*x+a),x)
```

output

```
( - 3*e**(2*a + 6*b*x + 6*c)*log(e**(b*x + c) - 1) + 3*e**(2*a + 6*b*x + 6*c)*log(e**(b*x + c) + 1) - 3*e**(6*b*x + 8*c)*log(e**(b*x + c) - 1) + 3*e**(6*b*x + 8*c)*log(e**(b*x + c) + 1) - 6*e**(2*a + 5*b*x + 5*c) - 6*e**(5*b*x + 7*c) + 9*e**(2*a + 4*b*x + 4*c)*log(e**(b*x + c) - 1) - 9*e**(2*a + 4*b*x + 4*c)*log(e**(b*x + c) + 1) + 9*e**(4*b*x + 6*c)*log(e**(b*x + c) - 1) - 9*e**(4*b*x + 6*c)*log(e**(b*x + c) + 1) - 16*e**(2*a + 3*b*x + 3*c) + 16*e**(3*b*x + 5*c) - 9*e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) + 9*e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) + 1) - 9*e**(2*b*x + 4*c)*log(e**(b*x + c) - 1) + 9*e**(2*b*x + 4*c)*log(e**(b*x + c) + 1) + 6*e**(2*a + b*x + c) + 6*e**(b*x + 3*c) + 3*e**(2*a)*log(e**(b*x + c) - 1) - 3*e**(2*a)*log(e**(b*x + c) + 1) + 3*e**(2*c)*log(e**(b*x + c) - 1) - 3*e**(2*c)*log(e**(b*x + c) + 1))/(12*e**(a + c)*b*(e**(6*b*x + 6*c) - 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) - 1))
```

3.75 $\int \operatorname{csch}(c - bx) \sinh(a + bx) dx$

Optimal result	564
Mathematica [A] (verified)	564
Rubi [F]	565
Maple [B] (verified)	565
Fricas [B] (verification not implemented)	566
Sympy [F]	566
Maxima [B] (verification not implemented)	567
Giac [A] (verification not implemented)	567
Mupad [B] (verification not implemented)	568
Reduce [B] (verification not implemented)	568

Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = -x \cosh(a + c) - \frac{\log(\sinh(c - bx)) \sinh(a + c)}{b}$$

output `-x*cosh(a+c)-ln(-sinh(b*x-c))*sinh(a+c)/b`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = -\frac{bx \cosh(a + c) + \log(\sinh(c - bx)) \sinh(a + c)}{b}$$

input `Integrate[Csch[c - b*x]*Sinh[a + b*x],x]`

output `-((b*x*Cosh[a + c] + Log[Sinh[c - b*x]]*Sinh[a + c])/b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{csch}(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \operatorname{csch}(c - bx) dx$$

input `Int[Csch[c - b*x]*Sinh[a + b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(28) = 56$.

Time = 0.08 (sec) , antiderivative size = 149, normalized size of antiderivative = 5.96

method	result
risch	$-x e^{a+c} + e^{-a-c} e^{2a+2c} x + \frac{e^{-a-c} e^{2a+2c} a}{b} - x e^{-a-c} - \frac{e^{-a-c} a}{b} - \frac{\ln(-e^{2a+2c} + e^{2bx+2a}) e^{-a-c} e^{2a+2c}}{2b} + \frac{\ln(-e^{2a+2c} + e^{2bx+2a})}{2b}$

input `int(-csch(b*x-c)*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `-x*exp(a+c)+exp(-a-c)*exp(2*a+2*c)*x+1/b*exp(-a-c)*exp(2*a+2*c)*a-x*exp(-a-c)-1/b*exp(-a-c)*a-1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)*exp(2*a+2*c)+1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(28) = 56$.

Time = 0.09 (sec) , antiderivative size = 146, normalized size of antiderivative = 5.84

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = \frac{2bx \cosh(a + c)^2 - 4bx \cosh(a + c) \sinh(a + c) + 2bx \sinh(a + c)^2 - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 - 1) \log(2(\cosh(a + c) \sinh(bx + a) - \cosh(bx + a) \sinh(a + c)) / (\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c)))}{2(b \cosh(a + c) - b \sinh(a + c))}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a),x, algorithm="fricas")`

output

```
-1/2*(2*b*x*cosh(a + c)^2 - 4*b*x*cosh(a + c)*sinh(a + c) + 2*b*x*sinh(a + c)^2 - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 - 1)*log(2*(cosh(a + c)*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c))))/(b*cosh(a + c) - b*sinh(a + c))
```

Sympy [F]

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = - \int \sinh(a + bx) \operatorname{csch}(bx - c) dx$$

input `integrate(-csch(b*x-c)*sinh(b*x+a),x)`

output

```
-Integral(sinh(a + b*x)*csch(b*x - c), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. $2(28) = 56$.

Time = 0.04 (sec) , antiderivative size = 81, normalized size of antiderivative = 3.24

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = -\frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{2b} - \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{2b} - \frac{(bx + a)e^{(a+c)}}{b}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(-b*x + c) + 1)/b - 1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(-b*x + c) - 1)/b - (b*x + a)*e^(a + c)/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.96

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = -xe^{(-a-c)} - \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(|e^{(2bx)} - e^{(2c)}|)}{2b}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a),x, algorithm="giac")`

output `-x*e^(-a - c) - 1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(abs(e^(2*b*x) - e^(2*c)))/b`

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.64

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = -x e^{-a-c} - \frac{e^{-2a-2c} \ln(e^{2a} e^{2bx} - e^{2a} e^{2c}) (2b e^{3a+3c} - 2b e^{a+c})}{4b^2}$$

input `int(sinh(a + b*x)/sinh(c - b*x),x)`output `- x*exp(- a - c) - (exp(- 2*a - 2*c)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp(2*c))*(2*b*exp(3*a + 3*c) - 2*b*exp(a + c)))/(4*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 83, normalized size of antiderivative = 3.32

$$\int \operatorname{csch}(c - bx) \sinh(a + bx) dx = \frac{-e^{2a+2c} \log(e^{bx} + e^c) - e^{2a+2c} \log(e^{bx} - e^c) + \log(e^{bx} + e^c) + \log(e^{bx} - e^c) - 2bx}{2e^{a+cb}}$$

input `int(-csch(b*x-c)*sinh(b*x+a),x)`output `(- e**(2*a + 2*c)*log(e**(b*x) + e**c) - e**(2*a + 2*c)*log(e**(b*x) - e**c) + log(e**(b*x) + e**c) + log(e**(b*x) - e**c) - 2*b*x)/(2*e**(a + c)*b)`

3.76 $\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx$

Optimal result	569
Mathematica [C] (verified)	569
Rubi [F]	570
Maple [B] (verified)	570
Fricas [B] (verification not implemented)	571
Sympy [F]	572
Maxima [B] (verification not implemented)	572
Giac [B] (verification not implemented)	573
Mupad [B] (verification not implemented)	573
Reduce [B] (verification not implemented)	574

Optimal result

Integrand size = 16, antiderivative size = 33

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh(a + c)}{b} + \frac{\operatorname{csch}(c - bx) \sinh(a + c)}{b}$$

output

`-arctanh(cosh(b*x-c))*cosh(a+c)/b-csch(b*x-c)*sinh(a+c)/b`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.64

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = -\frac{2i \operatorname{arctan}\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh(c) \cosh\left(\frac{bx}{2}\right) - \sinh(c) \sinh\left(\frac{bx}{2}\right)\right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a + c)}{b} + \frac{\operatorname{csch}(c - bx) \sinh(a + c)}{b}$$

input `Integrate[Csch[c - b*x]^2*Sinh[a + b*x],x]`

output `((-2*I)*ArcTan[((Cosh[c] - Sinh[c])*(Cosh[c]*Cosh[(b*x)/2] - Sinh[c]*Sinh[(b*x)/2]))/(I*Cosh[c]*Cosh[(b*x)/2] - I*Cosh[(b*x)/2]*Sinh[c])]*Cosh[a + c])/b + (Csch[c - b*x]*Sinh[a + c])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{csch}^2(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \operatorname{csch}^2(c - bx) dx$$

input `Int[Csch[c - b*x]^2*Sinh[a + b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(36) = 72$.

Time = 0.16 (sec) , antiderivative size = 164, normalized size of antiderivative = 4.97

method	result
risch	$\frac{e^{bx+a}(e^{2a+2c}-1)}{b(e^{2a+2c}-e^{2bx+2a})} - \frac{\ln(e^{a+c}+e^{bx+a})e^{-a-c}e^{2a+2c}}{2b} - \frac{\ln(e^{a+c}+e^{bx+a})e^{-a-c}}{2b} + \frac{\ln(-e^{a+c}+e^{bx+a})e^{-a-c}e^{2a+2c}}{2b} + \frac{\ln(-e^{a+c}+e^{bx+a})e^{-a-c}}{2b}$

input `int(csch(b*x-c)^2*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output

```
1/b*exp(b*x+a)*(exp(2*a+2*c)-1)/(exp(2*a+2*c)-exp(2*b*x+2*a))-1/2*ln(exp(a
+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)-1/2*ln(exp(a+c)+exp(b*x+a))/b*exp
(-a-c)+1/2*ln(-exp(a+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)+1/2*ln(-exp(a
+c)+exp(b*x+a))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1084 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 1084, normalized size of antiderivative = 32.85

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = \text{Too large to display}$$

input

```
integrate(csch(b*x-c)^2*sinh(b*x+a),x, algorithm="fricas")
```

output

```
1/2*(6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - 2*cosh(b*x + a)*sinh(a +
c)^3 - 2*(3*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c) + 2*(cosh(a + c)^
3 - cosh(a + c))*cosh(b*x + a) + (4*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c
)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 + cosh(a + c)^2)*cosh
(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + sinh(a + c)^4
+ (6*cosh(a + c)^2 + 1)*sinh(a + c)^2 + cosh(a + c)^2 - 2*(2*cosh(a + c)^
3 + cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*cosh(a + c)^2 + 1)*cos
h(b*x + a)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(4*cosh(b*x + a)*cosh(
a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - (6*cosh(a + c)^2 + 1)
*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3 + cosh(a + c))*cosh(b*x
+ a)*sinh(a + c) - (cosh(a + c)^4 + cosh(a + c)^2)*cosh(b*x + a))*sinh(b*x
+ a) + 2*((2*cosh(a + c)^3 + cosh(a + c))*cosh(b*x + a)^2 - cosh(a + c))*
sinh(a + c) + 1)*log(cosh(b*x + a)*cosh(a + c) + (cosh(a + c) - sinh(a + c
))*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c) + 1) - (4*cosh(b*x + a)^2*cos
h(a + c)*sinh(a + c)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 +
cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c
)^3 + sinh(a + c)^4 + (6*cosh(a + c)^2 + 1)*sinh(a + c)^2 + cosh(a + c)^2
- 2*(2*cosh(a + c)^3 + cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*cos
h(a + c)^2 + 1)*cosh(b*x + a)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(4*
cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - ...
```

Sympy [F]

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}^2(bx - c) dx$$

input `integrate(csch(b*x-c)**2*sinh(b*x+a),x)`

output `Integral(sinh(a + b*x)*csch(b*x - c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(36) = 72$.

Time = 0.05 (sec) , antiderivative size = 103, normalized size of antiderivative = 3.12

$$\begin{aligned} \int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = & -\frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{2b} \\ & + \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{2b} \\ & + \frac{(e^{(2a+2c)} - 1)e^{(-bx-a)}}{b(e^{(-2bx+2c)} - 1)} \end{aligned}$$

input `integrate(csch(b*x-c)^2*sinh(b*x+a),x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(-b*x + c) + 1)/b + 1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(-b*x + c) - 1)/b + (e^(2*a + 2*c) - 1)*e^(-b*x - a)/(b*(e^(-2*b*x + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(36) = 72$.

Time = 0.12 (sec) , antiderivative size = 107, normalized size of antiderivative = 3.24

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = -\frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(bx)} + e^c)}{2b} + \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(|e^{(bx)} - e^c|)}{2b} - \frac{(e^{(bx+2a+2c)} - e^{(bx)})e^{(-a)}}{b(e^{(2bx)} - e^{(2c)})}$$

input `integrate(csch(b*x-c)^2*sinh(b*x+a),x, algorithm="giac")`

output `-1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(b*x) + e^c)/b + 1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(abs(e^(b*x) - e^c))/b - (e^(b*x + 2*a + 2*c) - e^(b*x))*e^(-a)/(b*(e^(2*b*x) - e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.73

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx = \frac{e^{a+bx} (e^{2a+2c} - 1)}{b (e^{2a+2c} - e^{2a+2bx})} - \frac{\operatorname{atan}\left(\frac{e^{-a} e^{-2c} e^{bx} (\sqrt{-b^2 + e^{2a} e^{2c} \sqrt{-b^2}})}{b \sqrt{e^{-2a} e^{-2c} (2e^{2a} e^{2c} + e^{4a} e^{4c} + 1)}}\right) \sqrt{e^{-2a-2c} (2e^{2a+2c} + e^{4a+4c} + 1)}}{\sqrt{-b^2}}$$

input `int(sinh(a + b*x)/sinh(c - b*x)^2,x)`

output

```
(exp(a + b*x)*(exp(2*a + 2*c) - 1))/(b*(exp(2*a + 2*c) - exp(2*a + 2*b*x))
) - (atan((exp(-a)*exp(-2*c)*exp(b*x)*((-b^2)^(1/2) + exp(2*a)*exp(2*c)*(-
b^2)^(1/2)))/(b*(exp(-2*a)*exp(-2*c)*(2*exp(2*a)*exp(2*c) + exp(4*a)*exp(4
*c) + 1))^(1/2)))*(exp(- 2*a - 2*c)*(2*exp(2*a + 2*c) + exp(4*a + 4*c) + 1
))^(1/2))/(-b^2)^(1/2)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 218, normalized size of antiderivative = 6.61

$$\int \operatorname{csch}^2(c - bx) \sinh(a + bx) dx$$

$$= \frac{-e^{2bx+2a+2c} \log(e^{bx} + e^c) + e^{2bx+2a+2c} \log(e^{bx} - e^c) - e^{2bx} \log(e^{bx} + e^c) + e^{2bx} \log(e^{bx} - e^c) - 2e^{bx+2a+3c}}{2e^{a+cb}(e^{2bx} - e^{2c})}$$

input

```
int(csch(b*x-c)^2*sinh(b*x+a),x)
```

output

```
( - e**(2*a + 2*b*x + 2*c)*log(e**(b*x) + e**c) + e**(2*a + 2*b*x + 2*c)*l
og(e**(b*x) - e**c) - e**(2*b*x)*log(e**(b*x) + e**c) + e**(2*b*x)*log(e**
(b*x) - e**c) - 2*e**(2*a + b*x + 3*c) + 2*e**(b*x + c) + e**(2*a + 4*c)*l
og(e**(b*x) + e**c) - e**(2*a + 4*c)*log(e**(b*x) - e**c) + e**(2*c)*log(e
**(b*x) + e**c) - e**(2*c)*log(e**(b*x) - e**c))/(2*e**(a + c)*b*(e**(2*b*
x) - e**(2*c)))
```

3.77 $\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx$

Optimal result	575
Mathematica [A] (verified)	575
Rubi [F]	576
Maple [A] (verified)	576
Fricas [B] (verification not implemented)	577
Sympy [F]	577
Maxima [B] (verification not implemented)	578
Giac [A] (verification not implemented)	578
Mupad [F(-1)]	579
Reduce [B] (verification not implemented)	579

Optimal result

Integrand size = 16, antiderivative size = 37

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = -\frac{\cosh(a + c) \coth(c - bx)}{b} + \frac{\operatorname{csch}^2(c - bx) \sinh(a + c)}{2b}$$

output

$$\cosh(a+c)*\coth(b*x-c)/b+1/2*\operatorname{csch}(b*x-c)^2*\sinh(a+c)/b$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\begin{aligned} & \int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx \\ &= -\frac{(\cosh(a) - \cosh(a + c) \cosh(c - 2bx)) \operatorname{csch}(c) \operatorname{csch}^2(c - bx)}{2b} \end{aligned}$$

input

$$\text{Integrate}[\operatorname{Csch}[c - b*x]^3*\operatorname{Sinh}[a + b*x], x]$$

output

$$-1/2*((\operatorname{Cosh}[a] - \operatorname{Cosh}[a + c]*\operatorname{Cosh}[c - 2*b*x])*\operatorname{Csch}[c]*\operatorname{Csch}[c - b*x]^2)/b$$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{csch}^3(c - bx) dx$$

$$\downarrow 7299$$

$$\int \sinh(a + bx) \operatorname{csch}^3(c - bx) dx$$

input `Int[Csch[c - b*x]^3*Sinh[a + b*x],x]`

output `$Aborted`

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.54

method	result	size
risch	$-\frac{(e^{2a+2c}-2e^{2bx+2a}+1)e^{3a+3c}}{(e^{2a+2c}-e^{2bx+2a})^2 b}$	57
parallelrisch	$\frac{\left(\sinh(bx+a)\left(-\frac{\operatorname{sech}\left(\frac{bx}{2}-\frac{c}{2}\right)^2}{2}+1\right)\operatorname{csch}\left(\frac{bx}{2}-\frac{c}{2}\right)+\operatorname{sech}\left(\frac{bx}{2}-\frac{c}{2}\right)\cosh(bx+a)\right)\operatorname{csch}\left(\frac{bx}{2}-\frac{c}{2}\right)}{4b}$	63

input `int(-csch(b*x-c)^3*sinh(b*x+a),x,method=_RETURNVERBOSE)`

output `-1/(exp(2*a+2*c)-exp(2*b*x+2*a))^2/b*(exp(2*a+2*c)-2*exp(2*b*x+2*a)+1)*exp(3*a+3*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 449 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 449, normalized size of antiderivative = 12.14

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx =$$

$$\frac{-b \cosh(bx + a)^3 \cosh(a + c)^3 - b \cosh(bx + a) \cosh(a + c)^3 + (b \cosh(a + c)^3 - 3b \cosh(a + c)^2 \sinh(a + c)) \sinh(bx + a)}{\dots}$$

input `integrate(-csch(b*x-c)^3*sinh(b*x+a),x, algorithm="fricas")`

output

```
-2*(cosh(b*x + a)*cosh(a + c)*sinh(a + c) + cosh(b*x + a)*sinh(a + c)^2 -
(2*cosh(a + c)^2 + cosh(a + c)*sinh(a + c) - sinh(a + c)^2)*sinh(b*x + a))
/(b*cosh(b*x + a)^3*cosh(a + c)^3 - b*cosh(b*x + a)*cosh(a + c)^3 + (b*cos
h(a + c)^3 - 3*b*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(a + c)*sinh(a + c)^2
- b*sinh(a + c)^3)*sinh(b*x + a)^3 - (b*cosh(b*x + a)^3 + 3*b*cosh(b*x +
a))*sinh(a + c)^3 + 3*(b*cosh(b*x + a)*cosh(a + c)^3 - 3*b*cosh(b*x + a)*c
osh(a + c)^2*sinh(a + c) + 3*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - b
*cosh(b*x + a)*sinh(a + c)^3)*sinh(b*x + a)^2 + (3*b*cosh(b*x + a)^3*cosh(
a + c) + b*cosh(b*x + a)*cosh(a + c))*sinh(a + c)^2 + (3*b*cosh(b*x + a)^2
*cosh(a + c)^3 - 3*b*cosh(a + c)^3 - (3*b*cosh(b*x + a)^2 + b)*sinh(a + c)
^3 + 3*(3*b*cosh(b*x + a)^2*cosh(a + c) + b*cosh(a + c))*sinh(a + c)^2 - (
9*b*cosh(b*x + a)^2*cosh(a + c)^2 - b*cosh(a + c)^2)*sinh(a + c))*sinh(b*x
+ a) - 3*(b*cosh(b*x + a)^3*cosh(a + c)^2 - b*cosh(b*x + a)*cosh(a + c)^2
)*sinh(a + c))
```

Sympy [F]

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = - \int \sinh(a + bx) \operatorname{csch}^3(bx - c) dx$$

input `integrate(-csch(b*x-c)**3*sinh(b*x+a),x)`

output `-Integral(sinh(a + b*x)*csch(b*x - c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 3.57

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = \frac{2e^{(-2bx+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} - \frac{e^{(2a+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} - \frac{1}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})}$$

input `integrate(-csch(b*x-c)^3*sinh(b*x+a),x, algorithm="maxima")`

output
$$\frac{2e^{(-2bx+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} - \frac{e^{(2a+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} - \frac{1}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})}$$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = \frac{(2e^{(2bx+2a+3c)} - e^{(2a+5c)} - e^{(3c)})e^{(-a)}}{b(e^{(2bx)} - e^{(2c)})^2}$$

input `integrate(-csch(b*x-c)^3*sinh(b*x+a),x, algorithm="giac")`

output
$$(2e^{(2bx+2a+3c)} - e^{(2a+5c)} - e^{(3c)})e^{(-a)}/(b*(e^{(2bx)} - e^{(2c)})^2)$$

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\sinh(c - bx)^3} dx$$

input `int(sinh(a + b*x)/sinh(c - b*x)^3,x)`output `int(sinh(a + b*x)/sinh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.51

$$\int \operatorname{csch}^3(c - bx) \sinh(a + bx) dx = \frac{e^c (e^{4bx+2a} - e^{2c})}{e^a b (e^{4bx} - 2e^{2bx+2c} + e^{4c})}$$

input `int(-csch(b*x-c)^3*sinh(b*x+a),x)`output `(e**c*(e**(2*a + 4*b*x) - e**(2*c)))/(e**a*b*(e**(4*b*x) - 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.78 $\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$

Optimal result	580
Mathematica [A] (verified)	580
Rubi [F]	581
Maple [B] (verified)	581
Fricas [B] (verification not implemented)	582
Sympy [F]	582
Maxima [B] (verification not implemented)	583
Giac [B] (verification not implemented)	583
Mupad [F(-1)]	584
Reduce [B] (verification not implemented)	584

Optimal result

Integrand size = 16, antiderivative size = 65

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx = \frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh(a + c)}{2b} - \frac{\cosh(a + c) \coth(c - bx) \operatorname{csch}(c - bx)}{2b} + \frac{\operatorname{csch}^3(c - bx) \sinh(a + c)}{3b}$$

output

```
1/2*arctanh(cosh(b*x-c))*cosh(a+c)/b-1/2*cosh(a+c)*coth(b*x-c)*csch(b*x-c)/b-1/3*csch(b*x-c)^3*sinh(a+c)/b
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.94

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx = \frac{12 \operatorname{arctanh}(\cosh(c) - \sinh(c) \tanh(\frac{bx}{2})) \cosh(a + c) + \operatorname{csch}^3(c - bx)(4 \sinh(a + c) - 3 \cosh(a + c) \sinh(c))}{12b}$$

input

```
Integrate[Csch[c - b*x]^4*Sinh[a + b*x],x]
```

output

```
(12*ArcTanh[Cosh[c] - Sinh[c]*Tanh[(b*x)/2]]*Cosh[a + c] + Csch[c - b*x]^3
*(4*Sinh[a + c] - 3*Cosh[a + c]*Sinh[2*(c - b*x)]))/(12*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh(a + bx) \operatorname{csch}^4(c - bx) dx$$

↓ 7299

$$\int \sinh(a + bx) \operatorname{csch}^4(c - bx) dx$$

input

```
Int[Csch[c - b*x]^4*Sinh[a + b*x],x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. $2(63) = 126$.

Time = 0.76 (sec) , antiderivative size = 229, normalized size of antiderivative = 3.52

method	result
risch	$-\frac{e^{bx+a}(3e^{6a+6c}-8e^{2bx+6a+4c}+3e^{4a+4c}-3e^{4bx+6a+2c}+8e^{2bx+4a+2c}-3e^{4bx+4a})}{6b(e^{2a+2c}-e^{2bx+2a})^3} + \frac{\ln(e^a+c+e^{bx+a})e^{-a-c}e^{2a+2c}}{4b} + \frac{\ln(e^a-c+e^{bx+a})e^{-a-c}e^{2a+2c}}{4b}$

input

```
int(csch(b*x-c)^4*sinh(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
-1/6*exp(b*x+a)*(3*exp(6*a+6*c)-8*exp(2*b*x+6*a+4*c)+3*exp(4*a+4*c)-3*exp(
4*b*x+6*a+2*c)+8*exp(2*b*x+4*a+2*c)-3*exp(4*b*x+4*a))/b/(exp(2*a+2*c)-exp(
2*b*x+2*a))^3+1/4*ln(exp(a+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)+1/4*ln(
exp(a+c)+exp(b*x+a))/b*exp(-a-c)-1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-a-c)*
exp(2*a+2*c)-1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8922 vs. $2(63) = 126$.

Time = 0.16 (sec) , antiderivative size = 8922, normalized size of antiderivative = 137.26

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx = \text{Too large to display}$$

input

```
integrate(csch(b*x-c)^4*sinh(b*x+a),x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx = \int \sinh(a + bx) \operatorname{csch}^4(bx - c) dx$$

input

```
integrate(csch(b*x-c)**4*sinh(b*x+a),x)
```

output

```
Integral(sinh(a + b*x)*csch(b*x - c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(63) = 126$.

Time = 0.04 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.88

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$$

$$= \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{4b} - \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{4b}$$

$$+ \frac{3(e^{(2a+2c)} + 1)e^{(-bx-a)} + 8(e^{(4a+4c)} - e^{(2a+2c)})e^{(-3bx-3a)} - 3(e^{(6a+6c)} + e^{(4a+4c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+2c)} - 3e^{(-4bx+4c)} + e^{(-6bx+6c)} - 1)}$$

input `integrate(csch(b*x-c)^4*sinh(b*x+a),x, algorithm="maxima")`

output $\frac{1}{4}(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)/b - \frac{1}{4}(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)/b + \frac{1}{6}(3(e^{(2a+2c)} + 1)e^{(-bx-a)} + 8(e^{(4a+4c)} - e^{(2a+2c)})e^{(-3bx-3a)} - 3(e^{(6a+6c)} + e^{(4a+4c)})e^{(-5bx-5a)})/(b(3e^{(-2bx+2c)} - 3e^{(-4bx+4c)} + e^{(-6bx+6c)} - 1))$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. $2(63) = 126$.

Time = 0.14 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.45

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$$

$$= \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(bx)} + e^c)}{4b} - \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(|e^{(bx)} - e^c|)}{4b}$$

$$- \frac{(3e^{(5bx)} + 3e^{(5bx+2a+2c)} + 8e^{(3bx+2a+4c)} - 8e^{(3bx+2c)} - 3e^{(bx+2a+6c)} - 3e^{(bx+4c)})e^{(-a)}}{6b(e^{(2bx)} - e^{(2c)})^3}$$

input `integrate(csch(b*x-c)^4*sinh(b*x+a),x, algorithm="giac")`

output

```
1/4*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(b*x) + e^c)/b - 1/4*(e^(2*a + 2*
c) + 1)*e^(-a - c)*log(abs(e^(b*x) - e^c))/b - 1/6*(3*e^(5*b*x) + 3*e^(5*b
*x + 2*a + 2*c) + 8*e^(3*b*x + 2*a + 4*c) - 8*e^(3*b*x + 2*c) - 3*e^(b*x +
2*a + 6*c) - 3*e^(b*x + 4*c))*e^(-a)/(b*(e^(2*b*x) - e^(2*c))^3)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx = \int \frac{\sinh(a + bx)}{\sinh(c - bx)^4} dx$$

input

```
int(sinh(a + b*x)/sinh(c - b*x)^4,x)
```

output

```
int(sinh(a + b*x)/sinh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 496, normalized size of antiderivative = 7.63

$$\int \operatorname{csch}^4(c - bx) \sinh(a + bx) dx$$

$$= \frac{3e^{6bx+2a+2c} \log(e^{bx} + e^c) - 3e^{6bx+2a+2c} \log(e^{bx} - e^c) + 3e^{6bx} \log(e^{bx} + e^c) - 3e^{6bx} \log(e^{bx} - e^c) - 6e^{5bx+2a}}{\dots}$$

input

```
int(csch(b*x-c)^4*sinh(b*x+a),x)
```

output

```
(3*e**(2*a + 6*b*x + 2*c)*log(e**(b*x) + e**c) - 3*e**(2*a + 6*b*x + 2*c)*
log(e**(b*x) - e**c) + 3*e**(6*b*x)*log(e**(b*x) + e**c) - 3*e**(6*b*x)*lo
g(e**(b*x) - e**c) - 6*e**(2*a + 5*b*x + 3*c) - 6*e**(5*b*x + c) - 9*e**(2
*a + 4*b*x + 4*c)*log(e**(b*x) + e**c) + 9*e**(2*a + 4*b*x + 4*c)*log(e**(
b*x) - e**c) - 9*e**(4*b*x + 2*c)*log(e**(b*x) + e**c) + 9*e**(4*b*x + 2*c
)*log(e**(b*x) - e**c) - 16*e**(2*a + 3*b*x + 5*c) + 16*e**(3*b*x + 3*c) +
9*e**(2*a + 2*b*x + 6*c)*log(e**(b*x) + e**c) - 9*e**(2*a + 2*b*x + 6*c)*
log(e**(b*x) - e**c) + 9*e**(2*b*x + 4*c)*log(e**(b*x) + e**c) - 9*e**(2*b
*x + 4*c)*log(e**(b*x) - e**c) + 6*e**(2*a + b*x + 7*c) + 6*e**(b*x + 5*c)
- 3*e**(2*a + 8*c)*log(e**(b*x) + e**c) + 3*e**(2*a + 8*c)*log(e**(b*x) -
e**c) - 3*e**(6*c)*log(e**(b*x) + e**c) + 3*e**(6*c)*log(e**(b*x) - e**c)
)/(12*e**(a + c)*b*(e**(6*b*x) - 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) -
e**(6*c)))
```

3.79 $\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$

Optimal result	586
Mathematica [A] (verified)	586
Rubi [F]	587
Maple [B] (verified)	587
Fricas [B] (verification not implemented)	588
Sympy [F]	589
Maxima [B] (verification not implemented)	589
Giac [B] (verification not implemented)	590
Mupad [B] (verification not implemented)	590
Reduce [B] (verification not implemented)	591

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \operatorname{csch}(c+bx) \sinh^2(a+bx) dx = \frac{\cosh(2a-c+bx)}{b} - \frac{\operatorname{arctanh}(\cosh(c+bx)) \sinh^2(a-c)}{b}$$

output

```
cosh(b*x+2*a-c)/b-arctanh(cosh(b*x+c))*sinh(a-c)^2/b
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.39

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$$

$$= \frac{\cosh(2a - c + bx) + (-\log(\cosh(\frac{1}{2}(c + bx))) + \log(\sinh(\frac{1}{2}(c + bx)))) \sinh^2(a - c)}{b}$$

input

```
Integrate[Csch[c + b*x]*Sinh[a + b*x]^2,x]
```

output

```
(Cosh[2*a - c + b*x] + (-Log[Cosh[(c + b*x)/2]] + Log[Sinh[(c + b*x)/2]])*  
Sinh[a - c]^2)/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{csch}(bx + c) dx$$

$$\downarrow 7299$$

$$\int \sinh^2(a + bx) \operatorname{csch}(bx + c) dx$$

input `Int [Csch [c + b*x] * Sinh [a + b*x] ^2, x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. $2(36) = 72$.

Time = 0.29 (sec) , antiderivative size = 233, normalized size of antiderivative = 6.47

method	result
risch	$\frac{e^{bx+2a-c}}{2b} + \frac{e^{-bx-2a+c}}{2b} - \frac{\ln(e^{bx+a} + e^{a-c})e^{-2c-2a}e^{4a}}{4b} + \frac{\ln(e^{bx+a} + e^{a-c})e^{-2c-2a}e^{2a+2c}}{2b} - \frac{\ln(e^{bx+a} + e^{a-c})e^{-2c-2a}e^{4c}}{4b}$

input `int (csch (b*x+c) * sinh (b*x+a) ^2, x, method = _RETURNVERBOSE)`

output $\frac{1}{2} \frac{1}{b} \exp(bx+2a-c) + \frac{1}{2} \frac{1}{b} \exp(-bx-2a+c) - \frac{1}{4} \ln(\exp(bx+a) + \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(4a) + \frac{1}{2} \ln(\exp(bx+a) + \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(2a+2c) - \frac{1}{4} \ln(\exp(bx+a) + \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(4c) + \frac{1}{4} \ln(\exp(bx+a) - \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(4a) - \frac{1}{2} \ln(\exp(bx+a) - \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(2a+2c) + \frac{1}{4} \ln(\exp(bx+a) - \exp(a-c)) \frac{1}{b} \exp(-2c-2a) \exp(4c)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 880 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 880, normalized size of antiderivative = 24.44

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x+c)*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/4*(2*cosh(b*x + c)^2*cosh(-a + c)^4 - 8*cosh(b*x + c)^2*cosh(-a + c)^3*
sinh(-a + c) + 12*cosh(b*x + c)^2*cosh(-a + c)^2*sinh(-a + c)^2 - 8*cosh(b*
x + c)^2*cosh(-a + c)*sinh(-a + c)^3 + 2*cosh(b*x + c)^2*sinh(-a + c)^4 +
2*(cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a + c)^2*sinh(
-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4)*sinh(b*x + c)^
2 + (4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a +
c)^4 - 2*(3*cosh(-a + c)^2 - 1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a
+ c)^3 - cosh(-a + c))*cosh(b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 - 2*c
osh(-a + c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-
a + c)^3 + sinh(-a + c)^4 + 2*(3*cosh(-a + c)^2 - 1)*sinh(-a + c)^2 - 2*c
osh(-a + c)^2 - 4*(cosh(-a + c)^3 - cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*
x + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) - (4*cosh(b*x + c)*cosh(-a
+ c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - 2*(3*cosh(-a + c)^2 -
1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a + c)^3 - cosh(-a + c))*cosh(
b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 - 2*cosh(-a + c)^2 + 1)*cosh(b*x +
c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 + 2
*(3*cosh(-a + c)^2 - 1)*sinh(-a + c)^2 - 2*cosh(-a + c)^2 - 4*(cosh(-a + c
)^3 - cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c))*log(cosh(b*x + c) + s
inh(b*x + c) - 1) + 4*(cosh(b*x + c)*cosh(-a + c)^4 - 4*cosh(b*x + c)*cosh
(-a + c)^3*sinh(-a + c) + 6*cosh(b*x + c)*cosh(-a + c)^2*sinh(-a + c)^2...
```

Sympy [F]

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}(bx + c) dx$$

input `integrate(csch(b*x+c)*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.22

$$\begin{aligned} & \int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx \\ &= -\frac{(e^{4a} - 2e^{(2a+2c)} + e^{4c})e^{(-2a-2c)} \log(e^{-bx} + e^c)}{4b} \\ & \quad + \frac{(e^{4a} - 2e^{(2a+2c)} + e^{4c})e^{(-2a-2c)} \log(e^{-bx} - e^c)}{4b} + \frac{e^{(bx+2a-c)}}{2b} + \frac{e^{(-bx-2a+c)}}{2b} \end{aligned}$$

input `integrate(csch(b*x+c)*sinh(b*x+a)^2,x, algorithm="maxima")`

output `-1/4*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/4*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + 1/2*e^(b*x + 2*a - c)/b + 1/2*e^(-b*x - 2*a + c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(36) = 72$.

Time = 0.19 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.31

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$$

$$= -\frac{(e^{4a+c} - 2e^{2a+3c} + e^{5c})e^{(-2a-3c)} \log(e^{(bx+c)} + 1)}{4b}$$

$$+ \frac{(e^{4a+c} - 2e^{2a+3c} + e^{5c})e^{(-2a-3c)} \log(|e^{(bx+c)} - 1|)}{4b} + \frac{e^{(bx+2a-c)}}{2b} + \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(csch(b*x+c)*sinh(b*x+a)^2,x, algorithm="giac")`

output `-1/4*(e^(4*a + c) - 2*e^(2*a + 3*c) + e^(5*c))*e^(-2*a - 3*c)*log(e^(b*x + c) + 1)/b + 1/4*(e^(4*a + c) - 2*e^(2*a + 3*c) + e^(5*c))*e^(-2*a - 3*c)*log(abs(e^(b*x + c) - 1))/b + 1/2*e^(b*x + 2*a - c)/b + 1/2*e^(-b*x - 2*a + c)/b`

Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 202, normalized size of antiderivative = 5.61

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx = \frac{e^{c-2a-bx}}{2b} + \frac{e^{2a-c+bx}}{2b}$$

$$- \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{3c} e^{bx} (\sqrt{-b^2} - 2e^{2a} e^{-2c} \sqrt{-b^2} + e^{4a} e^{-4c} \sqrt{-b^2})}{b \sqrt{e^{-4a} e^{4c} (6e^{4a} e^{-4c} - 4e^{2a} e^{-2c} - 4e^{6a} e^{-6c} + e^{8a} e^{-8c} + 1)}}\right)}{2\sqrt{-b^2}} \sqrt{e^{4c-4a} (6e^{4a-4c} - 4e^{2a-2c} - 4e^{6a-6c} + e^{8a-8c} + 1)}$$

input `int(sinh(a + b*x)^2/sinh(c + b*x),x)`

output `exp(c - 2*a - b*x)/(2*b) + exp(2*a - c + b*x)/(2*b) - (atan((exp(-2*a)*exp(3*c)*exp(b*x))*((-b^2)^(1/2) - 2*exp(2*a)*exp(-2*c)*(-b^2)^(1/2) + exp(4*a)*exp(-4*c)*(-b^2)^(1/2)))/(b*(exp(-4*a)*exp(4*c)*(6*exp(4*a)*exp(-4*c) - 4*exp(2*a)*exp(-2*c) - 4*exp(6*a)*exp(-6*c) + exp(8*a)*exp(-8*c) + 1))^(1/2))*(exp(4*c - 4*a)*(6*exp(4*a - 4*c) - 4*exp(2*a - 2*c) - 4*exp(6*a - 6*c) + exp(8*a - 8*c) + 1))^(1/2))/(2*(-b^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 170, normalized size of antiderivative = 4.72

$$\int \operatorname{csch}(c + bx) \sinh^2(a + bx) dx$$

$$= \frac{2e^{2bx+4a+c} + e^{bx+4a} \log(e^{bx+c} - 1) - e^{bx+4a} \log(e^{bx+c} + 1) - 2e^{bx+2a+2c} \log(e^{bx+c} - 1) + 2e^{bx+2a+2c} \log(e^{bx+c} + 1)}{4e^{bx+2a+2c} b}$$

input

```
int(csch(b*x+c)*sinh(b*x+a)^2,x)
```

output

```
(2*e**(4*a + 2*b*x + c) + e**(4*a + b*x)*log(e**(b*x + c) - 1) - e**(4*a +
b*x)*log(e**(b*x + c) + 1) - 2*e**(2*a + b*x + 2*c)*log(e**(b*x + c) - 1)
+ 2*e**(2*a + b*x + 2*c)*log(e**(b*x + c) + 1) + e**(b*x + 4*c)*log(e**(b
*x + c) - 1) - e**(b*x + 4*c)*log(e**(b*x + c) + 1) + 2*e**(3*c))/(4*e**(2
*a + b*x + 2*c)*b)
```


3.80 $\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx$

Optimal result	592
Mathematica [A] (verified)	592
Rubi [F]	593
Maple [B] (verified)	593
Fricas [B] (verification not implemented)	594
Sympy [F]	594
Maxima [B] (verification not implemented)	595
Giac [B] (verification not implemented)	595
Mupad [B] (verification not implemented)	596
Reduce [B] (verification not implemented)	596

Optimal result

Integrand size = 17, antiderivative size = 49

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = x \cosh(2(a - c)) - \frac{\coth(c + bx) \sinh^2(a - c)}{b} + \frac{\log(\sinh(c + bx)) \sinh(2(a - c))}{b}$$

output `x*cosh(2*a-2*c)-coth(b*x+c)*sinh(a-c)^2/b+ln(sinh(b*x+c))*sinh(2*a-2*c)/b`

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = \frac{bx \cosh(2(a - c)) + \log(\sinh(c + bx)) \sinh(2(a - c)) + \operatorname{csch}(c) \operatorname{csch}(c + bx) \sinh^2(a - c) \sinh(bx)}{b}$$

input `Integrate[Csch[c + b*x]^2*Sinh[a + b*x]^2,x]`

output `(b*x*Cosh[2*(a - c)] + Log[Sinh[c + b*x]]*Sinh[2*(a - c)] + Csch[c]*Csch[c + b*x]*Sinh[a - c]^2*Sinh[b*x])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

input `Int[Csch[c + b*x]^2*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 257 vs. $2(49) = 98$.

Time = 0.48 (sec) , antiderivative size = 258, normalized size of antiderivative = 5.27

method	result
risch	$x e^{2a-2c} - e^{-2c-2a} e^{4a} x + e^{-2c-2a} e^{4c} x - \frac{e^{-2c-2a} e^{4a} a}{b} + \frac{e^{-2c-2a} e^{4c} a}{b} + \frac{e^{-2c} e^{4a}}{2b(-e^{2bx+2a+2c} + e^{2a})} - \frac{e^{-2c} e^{2a}}{b(-e^{2bx+2a} + e^{2c})}$

input `int(csch(b*x+c)^2*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `x*exp(2*a-2*c)-exp(-2*c-2*a)*exp(4*a)*x+exp(-2*c-2*a)*exp(4*c)*x-1/b*exp(-2*c-2*a)*exp(4*a)*a+1/b*exp(-2*c-2*a)*exp(4*c)*a+1/2/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*a)-1/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(2*a+2*c)+1/2/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*c)+1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*a)-1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 710 vs. $2(49) = 98$.

Time = 0.10 (sec) , antiderivative size = 710, normalized size of antiderivative = 14.49

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/2*(2*b*x*cosh(b*x + c)^2 - cosh(-a + c)^4 + 4*b*x*cosh(b*x + c)*sinh(b*x
+ c) + 2*b*x*sinh(b*x + c)^2 + 4*cosh(-a + c)*sinh(-a + c)^3 - sinh(-a +
c)^4 - 2*(3*cosh(-a + c)^2 - 1)*sinh(-a + c)^2 - 2*b*x + 2*cosh(-a + c)^2
+ ((cosh(b*x + c)^2 - 1)*sinh(-a + c)^4 - cosh(-a + c)^4 - 4*(cosh(b*x + c
)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c)^3 + (cosh(-a + c)^4 - 1)*cos
h(b*x + c)^2 + (cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a
+ c)^2*sinh(-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 -
1)*sinh(b*x + c)^2 + 6*(cosh(b*x + c)^2*cosh(-a + c)^2 - cosh(-a + c)^2)*s
inh(-a + c)^2 - 2*(4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) - 6*cosh(b*
x + c)*cosh(-a + c)^2*sinh(-a + c)^2 + 4*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - (cosh(-a + c)^4 - 1)*cosh(b*x +
c))*sinh(b*x + c) - 4*(cosh(b*x + c)^2*cosh(-a + c)^3 - cosh(-a + c)^3)*si
nh(-a + c) + 1)*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))) + 4*(
cosh(-a + c)^3 - cosh(-a + c))*sinh(-a + c) - 1)/(b*cosh(b*x + c)^2*cosh(-
a + c)^2 - b*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a
+ c) + b*sinh(-a + c)^2)*sinh(b*x + c)^2 + (b*cosh(b*x + c)^2 - b)*sinh(-
a + c)^2 + 2*(b*cosh(b*x + c)*cosh(-a + c)^2 - 2*b*cosh(b*x + c)*cosh(-a +
c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 2*(b*co
sh(b*x + c)^2*cosh(-a + c) - b*cosh(-a + c))*sinh(-a + c))
```

Sympy [F]

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

input `integrate(csch(b*x+c)**2*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(49) = 98$.

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.69

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{-bx} + e^c)}{2b} + \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{-bx} - e^c)}{2b} + \frac{(bx + a)e^{(2a-2c)}}{b} + \frac{e^{4a} - 2e^{(2a+2c)} + e^{4c}}{2b(e^{(-2bx+2a)} - e^{(2a+2c)})}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + (b*x + a)*e^(2*a - 2*c)/b + 1/2*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))/(b*(e^(-2*b*x + 2*a) - e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(49) = 98$.

Time = 0.14 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.10

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = xe^{(-2a+2c)} + \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(|e^{(2bx+2c)} - 1|)}{2b} - \frac{(e^{(2bx+4a)} - e^{(2bx+4c)} - 2e^{(2a)} + 2e^{(2c)})e^{(-2a)}}{2b(e^{(2bx+2c)} - 1)}$$

input `integrate(csch(b*x+c)^2*sinh(b*x+a)^2,x, algorithm="giac")`

output

```
x*e^(-2*a + 2*c) + 1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(abs(e^(2*b*x
+ 2*c) - 1))/b - 1/2*(e^(2*b*x + 4*a) - e^(2*b*x + 4*c) - 2*e^(2*a) + 2*
e^(2*c))*e^(-2*a)/(b*(e^(2*b*x + 2*c) - 1))
```

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.80

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = x e^{2c-2a} + \frac{\sinh(2a - 2c) \ln(e^{2a} e^{2bx} - e^{2a} e^{-2c})}{b} + \frac{2 e^{2a-2c} \sinh(a - c)^2}{b (e^{2a-2c} - e^{2a+2bx})}$$

input

```
int(sinh(a + b*x)^2/sinh(c + b*x)^2,x)
```

output

```
x*exp(2*c - 2*a) + (sinh(2*a - 2*c)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp
(-2*c)))/b + (2*exp(2*a - 2*c)*sinh(a - c)^2)/(b*(exp(2*a - 2*c) - exp(2*a
+ 2*b*x)))
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 254, normalized size of antiderivative = 5.18

$$\int \operatorname{csch}^2(c + bx) \sinh^2(a + bx) dx = \frac{e^{2bx+4a+2c} \log(e^{bx+c} - 1) + e^{2bx+4a+2c} \log(e^{bx+c} + 1) - e^{2bx+4a+2c} + 2e^{2bx+2a+4c} - e^{2bx+6c} \log(e^{bx+c} - 1) - e^{2bx+6c} \log(e^{bx+c} + 1) - 2e^{2bx+2a+4c} + 2e^{2bx+6c}}{b}$$

input

```
int(csch(b*x+c)^2*sinh(b*x+a)^2,x)
```

output

```
(e**(4*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) + e**(4*a + 2*b*x + 2*c)*log
(e**(b*x + c) + 1) - e**(4*a + 2*b*x + 2*c) + 2*e**(2*a + 2*b*x + 4*c) - e
**(2*b*x + 6*c)*log(e**(b*x + c) - 1) - e**(2*b*x + 6*c)*log(e**(b*x + c)
+ 1) + 2*e**(2*b*x + 6*c)*b*x - e**(2*b*x + 6*c) - e**(4*a)*log(e**(b*x +
c) - 1) - e**(4*a)*log(e**(b*x + c) + 1) + e**(4*c)*log(e**(b*x + c) - 1)
+ e**(4*c)*log(e**(b*x + c) + 1) - 2*e**(4*c)*b*x)/(2*e**(2*a + 2*c)*b*(e
*(2*b*x + 2*c) - 1))
```

3.81 $\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx$

Optimal result	598
Mathematica [B] (verified)	598
Rubi [F]	599
Maple [B] (verified)	599
Fricas [B] (verification not implemented)	600
Sympy [F]	600
Maxima [B] (verification not implemented)	601
Giac [B] (verification not implemented)	601
Mupad [F(-1)]	602
Reduce [B] (verification not implemented)	602

Optimal result

Integrand size = 17, antiderivative size = 89

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(2(a - c))}{b} + \frac{\operatorname{arctanh}(\cosh(c + bx)) \sinh^2(a - c)}{2b} - \frac{\operatorname{coth}(c + bx) \operatorname{csch}(c + bx) \sinh^2(a - c)}{2b} - \frac{\operatorname{csch}(c + bx) \sinh(2(a - c))}{b}$$

output `-arctanh(cosh(b*x+c))*cosh(2*a-2*c)/b+1/2*arctanh(cosh(b*x+c))*sinh(a-c)^2/b-1/2*coth(b*x+c)*csch(b*x+c)*sinh(a-c)^2/b-csch(b*x+c)*sinh(2*a-2*c)/b`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 246 vs. 2(89) = 178.

Time = 1.56 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.76

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx = \frac{\operatorname{csch}^2(\frac{1}{2}(c + bx)) - \cosh(2(a - c)) \operatorname{csch}^2(\frac{1}{2}(c + bx)) - 4 \log(\cosh(\frac{1}{2}(c + bx))) - 12 \cosh(2(a - c)) \log(\cosh(\frac{1}{2}(c + bx)))}{b}$$

input `Integrate[Csch[c + b*x]^3*Sinh[a + b*x]^2,x]`

output $(\text{Csch}[(c + b*x)/2]^2 - \text{Cosh}[2*(a - c)]*\text{Csch}[(c + b*x)/2]^2 - 4*\text{Log}[\text{Cosh}[(c + b*x)/2]] - 12*\text{Cosh}[2*(a - c)]*\text{Log}[\text{Cosh}[(c + b*x)/2]] + 4*\text{Log}[\text{Sinh}[(c + b*x)/2]] + 12*\text{Cosh}[2*(a - c)]*\text{Log}[\text{Sinh}[(c + b*x)/2]] + \text{Sech}[(c + b*x)/2]^2 - \text{Cosh}[2*(a - c)]*\text{Sech}[(c + b*x)/2]^2 - 4*\text{Cosh}[2*a - 2*c - (b*x)/2]*(\text{Csch}[c/2]*\text{Csch}[(c + b*x)/2] + \text{Sech}[c/2]*\text{Sech}[(c + b*x)/2]) + 4*\text{Cosh}[2*a - 2*c + (b*x)/2]*(\text{Csch}[c/2]*\text{Csch}[(c + b*x)/2] + \text{Sech}[c/2]*\text{Sech}[(c + b*x)/2]))/(16*b)$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \text{csch}^3(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \text{csch}^3(bx + c) dx$$

input `Int [Csch[c + b*x]^3*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(85) = 170$.

Time = 0.97 (sec) , antiderivative size = 305, normalized size of antiderivative = 3.43

method	result
risch	$\frac{(-5e^{2bx+6a+2c} + 2e^{2bx+4a+4c} + 3e^{2bx+2a+6c} + 3e^{6a} + 2e^{4a+2c} - 5e^{2a+4c})e^{bx-c}}{4(-e^{2bx+2a+2c} + e^{2a})^2 b} + \frac{3 \ln(e^{bx+a} - e^{a-c})e^{-2c-2a}e^{4a}}{8b} + \frac{\ln(e^{bx+a} - e^{a-c})}{8b}$

input `int(csch(b*x+c)^3*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/4/(-exp(2*b*x+2*a+2*c)+exp(2*a))^2/b*(-5*exp(2*b*x+6*a+2*c)+2*exp(2*b*x+4*a+4*c)+3*exp(2*b*x+2*a+6*c)+3*exp(6*a)+2*exp(4*a+2*c)-5*exp(2*a+4*c))*exp(b*x-c)+3/8*ln(exp(b*x+a)-exp(a-c))/b*exp(-2*c-2*a)*exp(4*a)+1/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*ln(exp(b*x+a)-exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)-3/8*ln(exp(b*x+a)+exp(a-c))/b*exp(-2*c-2*a)*exp(4*a)-1/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+2*c)-3/8*ln(exp(b*x+a)+exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3234 vs. $2(85) = 170$.

Time = 0.11 (sec) , antiderivative size = 3234, normalized size of antiderivative = 36.34

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}^3(bx + c) dx$$

input `integrate(csch(b*x+c)**3*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x + c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. $2(85) = 170$.

Time = 0.04 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.33

$$\int \operatorname{csch}^3(c+bx) \sinh^2(a+bx) dx$$

$$= -\frac{(3e^{4a} + 2e^{(2a+2c)} + 3e^{(4c)})e^{(-2a-2c)} \log(e^{(-bx)} + e^c)}{8b}$$

$$+ \frac{(3e^{4a} + 2e^{(2a+2c)} + 3e^{(4c)})e^{(-2a-2c)} \log(e^{(-bx)} - e^c)}{8b}$$

$$+ \frac{(5e^{(4a+2c)} - 2e^{(2a+4c)} - 3e^{(6c)})e^{(-bx-a)} - (3e^{(6a)} + 2e^{(4a+2c)} - 5e^{(2a+4c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+c)} - e^{(a+5c)})}$$

input `integrate(csch(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="maxima")`

output `-1/8*(3*e^(4*a) + 2*e^(2*a + 2*c) + 3*e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/8*(3*e^(4*a) + 2*e^(2*a + 2*c) + 3*e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + 1/4*((5*e^(4*a + 2*c) - 2*e^(2*a + 4*c) - 3*e^(6*c))*e^(-b*x - a) - (3*e^(6*a) + 2*e^(4*a + 2*c) - 5*e^(2*a + 4*c))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + c) - e^(a + 5*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. $2(85) = 170$.

Time = 0.16 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.19

$$\int \operatorname{csch}^3(c+bx) \sinh^2(a+bx) dx$$

$$= -\frac{(3e^{(4a+c)} + 2e^{(2a+3c)} + 3e^{(5c)})e^{(-2a-3c)} \log(e^{(bx+c)} + 1)}{8b}$$

$$+ \frac{(3e^{(4a+c)} + 2e^{(2a+3c)} + 3e^{(5c)})e^{(-2a-3c)} \log(|e^{(bx+c)} - 1|)}{8b}$$

$$- \frac{(5e^{(3bx+4a+2c)} - 2e^{(3bx+2a+4c)} - 3e^{(3bx+6c)} - 3e^{(bx+4a)} - 2e^{(bx+2a+2c)} + 5e^{(bx+4c)})e^{(-2a-c)}}{4b(e^{(2bx+2c)} - 1)^2}$$

input `integrate(csch(b*x+c)^3*sinh(b*x+a)^2,x, algorithm="giac")`

output

```
-1/8*(3*e^(4*a + c) + 2*e^(2*a + 3*c) + 3*e^(5*c))*e^(-2*a - 3*c)*log(e^(b
*x + c) + 1)/b + 1/8*(3*e^(4*a + c) + 2*e^(2*a + 3*c) + 3*e^(5*c))*e^(-2*a
- 3*c)*log(abs(e^(b*x + c) - 1))/b - 1/4*(5*e^(3*b*x + 4*a + 2*c) - 2*e^(
3*b*x + 2*a + 4*c) - 3*e^(3*b*x + 6*c) - 3*e^(b*x + 4*a) - 2*e^(b*x + 2*a
+ 2*c) + 5*e^(b*x + 4*c))*e^(-2*a - c)/(b*(e^(2*b*x + 2*c) - 1)^2)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\sinh(c + bx)^3} dx$$

input

```
int(sinh(a + b*x)^2/sinh(c + b*x)^3,x)
```

output

```
int(sinh(a + b*x)^2/sinh(c + b*x)^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 520, normalized size of antiderivative = 5.84

$$\int \operatorname{csch}^3(c + bx) \sinh^2(a + bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \log(e^{bx+c} - 1) - 3e^{4bx+4a+4c} \log(e^{bx+c} + 1) + 2e^{4bx+2a+6c} \log(e^{bx+c} - 1) - 2e^{4bx+2a+6c} \log(e^{bx+c} + 1)}{b}$$

input

```
int(csch(b*x+c)^3*sinh(b*x+a)^2,x)
```

output

$$\begin{aligned}
& (3e^{4a+4bx+4c} \log(e^{bx+c} - 1) - 3e^{4a+4bx+4c} \\
& \log(e^{bx+c} + 1) + 2e^{2a+4bx+6c} \log(e^{bx+c} - 1) - \\
& 2e^{2a+4bx+6c} \log(e^{bx+c} + 1) + 3e^{4bx+8c} \log(e^{bx+c} - 1) - \\
& 3e^{4bx+8c} \log(e^{bx+c} + 1) - 10e^{4a+3bx+3c} \\
& + 4e^{2a+3bx+5c} + 6e^{3bx+7c} - 6e^{4a+2bx+2c} \log(e^{bx+c} - 1) \\
& + 6e^{4a+2bx+2c} \log(e^{bx+c} + 1) - 4e^{2a+2bx+4c} \log(e^{bx+c} - 1) \\
& + 4e^{2a+2bx+4c} \log(e^{bx+c} + 1) - 6e^{2bx+6c} \log(e^{bx+c} - 1) \\
& + 6e^{2bx+6c} \log(e^{bx+c} + 1) + 6e^{4a+bx+c} + 4e^{2a+bx+3c} \\
& - 10e^{bx+5c} + 3e^{4a} \log(e^{bx+c} - 1) - 3e^{4a} \log(e^{bx+c} + 1) \\
& + 2e^{2a+2c} \log(e^{bx+c} - 1) - 2e^{2a+2c} \log(e^{bx+c} + 1) \\
& + 3e^{4c} \log(e^{bx+c} - 1) - 3e^{4c} \log(e^{bx+c} + 1)) / (8e^{2a+2c} b (e^{4bx+4c} - 2e^{2bx+2c} + 1))
\end{aligned}$$

3.82 $\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx$

Optimal result	604
Mathematica [A] (verified)	604
Rubi [F]	605
Maple [A] (verified)	605
Fricas [B] (verification not implemented)	606
Sympy [F]	607
Maxima [B] (verification not implemented)	607
Giac [A] (verification not implemented)	608
Mupad [F(-1)]	608
Reduce [B] (verification not implemented)	609

Optimal result

Integrand size = 17, antiderivative size = 84

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = -\frac{\cosh(2(a - c)) \coth(c + bx)}{b} + \frac{\coth(c + bx) \sinh^2(a - c)}{b} - \frac{\coth^3(c + bx) \sinh^2(a - c)}{3b} - \frac{\operatorname{csch}^2(c + bx) \sinh(2(a - c))}{2b}$$

output

```
-cosh(2*a-2*c)*coth(b*x+c)/b+coth(b*x+c)*sinh(a-c)^2/b-1/3*coth(b*x+c)^3*
sinh(a-c)^2/b-1/2*csch(b*x+c)^2*sinh(2*a-2*c)/b
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.95

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = \frac{\operatorname{csch}(c) \operatorname{csch}^3(c + bx) (-3 \sinh(bx) - \sinh(2a - 4c - 3bx)) + 3 \sinh(2a - 2c - bx) - 3 \sinh(2a + bx) + \sinh(2a - 2c)}{12b}$$

input `Integrate[Csch[c + b*x]^4*Sinh[a + b*x]^2,x]`

output $(\text{Csch}[c] * \text{Csch}[c + b*x]^3 * (-3 * \text{Sinh}[b*x] - \text{Sinh}[2*a - 4*c - 3*b*x] + 3 * \text{Sinh}[2*a - 2*c - b*x] - 3 * \text{Sinh}[2*a + b*x] + \text{Sinh}[2*a + 3*b*x] + \text{Sinh}[2*c + 3*b*x])) / (12*b)$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \text{csch}^4(bx + c) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \text{csch}^4(bx + c) dx$$

input `Int [Csch[c + b*x]^4*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [A] (verified)

Time = 1.85 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

method	result	size
parallelrisch	$\frac{\text{sech}\left(\frac{bx}{2} + \frac{c}{2}\right)^3 \text{csch}\left(\frac{bx}{2} + \frac{c}{2}\right)^3 (-\cosh(3bx+3c) - 2 \cosh(3bx+2a+c) + 3 \cosh(bx+c))}{96b}$	60
risch	$\frac{2(3e^{4bx+4a+4c} - 3e^{2bx+4a+2c} - 3e^{2bx+2a+4c} + e^{4a} + e^{2a+2c} + e^{4c})e^{4a-2c}}{3(-e^{2bx+2a+2c} + e^{2a})^3 b}$	94

input `int(csch(b*x+c)^4*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
1/96*sech(1/2*b*x+1/2*c)^3*csch(1/2*b*x+1/2*c)^3*(-cosh(3*b*x+3*c)-2*cosh(
3*b*x+2*a+c)+3*cosh(b*x+c))/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 497 vs. $2(80) = 160$.

Time = 0.09 (sec) , antiderivative size = 497, normalized size of antiderivative = 5.92

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx =$$

$$\frac{2 \left((5 \cosh(-a + c))^2 + 1 \right) \cosh(bx + c)^2 - 3 (b \cosh(bx + c)^4 \cosh(-a + c)^2 - 4b \cosh(bx + c)^2 \cosh(-a + c)^2 + (b \cosh(-a + c)^2 - b \sinh(-a + c)^2) \sinh(bx + c)^2}{3 (b \cosh(bx + c)^4 \cosh(-a + c)^2 - 4b \cosh(bx + c)^2 \cosh(-a + c)^2 + (b \cosh(-a + c)^2 - b \sinh(-a + c)^2) \sinh(bx + c)^2}$$

input

```
integrate(csch(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
-2/3*((5*cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (5*cosh(-a + c)^2 - 6*cosh(
-a + c)*sinh(-a + c) + 5*sinh(-a + c)^2 + 1)*sinh(b*x + c)^2 + (5*cosh(b*x
+ c)^2 - 3)*sinh(-a + c)^2 - 3*cosh(-a + c)^2 - 2*(6*cosh(b*x + c)*cosh(-
a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*
cosh(b*x + c))*sinh(b*x + c) - 6*(cosh(b*x + c)^2*cosh(-a + c) - cosh(-a +
c))*sinh(-a + c) - 3)/(b*cosh(b*x + c)^4*cosh(-a + c)^2 - 4*b*cosh(b*x +
c)^2*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^
4 + 4*(b*cosh(b*x + c)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*si
nh(b*x + c)^3 + 3*b*cosh(-a + c)^2 + 2*(3*b*cosh(b*x + c)^2*cosh(-a + c)^2
- 2*b*cosh(-a + c)^2 - (3*b*cosh(b*x + c)^2 - 2*b)*sinh(-a + c)^2)*sinh(b
*x + c)^2 - (b*cosh(b*x + c)^4 - 4*b*cosh(b*x + c)^2 + 3*b)*sinh(-a + c)^2
+ 4*(b*cosh(b*x + c)^3*cosh(-a + c)^2 - b*cosh(b*x + c)*cosh(-a + c)^2 -
(b*cosh(b*x + c)^3 - b*cosh(b*x + c))*sinh(-a + c)^2)*sinh(b*x + c))
```

Sympy [F]

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}^4(bx + c) dx$$

input `integrate(csch(b*x+c)**4*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 331 vs. $2(80) = 160$.

Time = 0.04 (sec) , antiderivative size = 331, normalized size of antiderivative = 3.94

$$\begin{aligned} & \int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx \\ &= -\frac{2(e^{(4a+4c)} + e^{(2a+6c)})e^{(-2bx-2a)}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(2a+6c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(8c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \end{aligned}$$

input `integrate(csch(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned}
& -2*(e^{(4*a + 4*c)} + e^{(2*a + 6*c)})*e^{(-2*b*x - 2*a)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a) - e^{(2*a + 6*c)}))} + 2 \\
& *e^{(-4*b*x + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a) - e^{(2*a + 6*c)}))} + 2/3*e^{(4*a + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a) - e^{(2*a + 6*c)}))} \\
&) + 2/3*e^{(2*a + 6*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a) - e^{(2*a + 6*c)}))} + 2/3*e^{(8*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a) - e^{(2*a + 6*c)}))}
\end{aligned}$$
Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.01

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = \frac{2(3e^{(4bx+4a+4c)} - 3e^{(2bx+4a+2c)} - 3e^{(2bx+2a+4c)} + e^{(4a)} + e^{(2a+2c)} + e^{(4c)})e^{(-2a-2c)}}{3b(e^{(2bx+2c)} - 1)^3}$$

input

```
integrate(csch(b*x+c)^4*sinh(b*x+a)^2,x, algorithm="giac")
```

output

$$-2/3*(3*e^{(4*b*x + 4*a + 4*c)} - 3*e^{(2*b*x + 4*a + 2*c)} - 3*e^{(2*b*x + 2*a + 4*c)} + e^{(4*a)} + e^{(2*a + 2*c)} + e^{(4*c)})*e^{(-2*a - 2*c)/(b*(e^{(2*b*x + 2*c)} - 1)^3)}$$
Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\sinh(c + bx)^4} dx$$

input

```
int(sinh(a + b*x)^2/sinh(c + b*x)^4,x)
```

output

```
int(sinh(a + b*x)^2/sinh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.13

$$\int \operatorname{csch}^4(c + bx) \sinh^2(a + bx) dx = \frac{-\frac{2e^{6bx+4a+4c}}{3} + 2e^{2bx+2a+2c} - \frac{2e^{2a}}{3} - \frac{2e^{2c}}{3}}{e^{2a}b(e^{6bx+6c} - 3e^{4bx+4c} + 3e^{2bx+2c} - 1)}$$

input `int(csch(b*x+c)^4*sinh(b*x+a)^2,x)`output `(2*(- e**(4*a + 6*b*x + 4*c) + 3*e**(2*a + 2*b*x + 2*c) - e**(2*a) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x + 6*c) - 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) - 1))`

3.83 $\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$

Optimal result	610
Mathematica [A] (verified)	610
Rubi [F]	611
Maple [B] (verified)	611
Fricas [B] (verification not implemented)	612
Sympy [F]	613
Maxima [B] (verification not implemented)	613
Giac [B] (verification not implemented)	614
Mupad [B] (verification not implemented)	614
Reduce [B] (verification not implemented)	615

Optimal result

Integrand size = 16, antiderivative size = 33

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx = -\frac{\cosh(2a + c + bx)}{b} + \frac{\operatorname{arctanh}(\cosh(c - bx)) \sinh^2(a + c)}{b}$$

output `-cosh(b*x+2*a+c)/b+arctanh(cosh(b*x-c))*sinh(a+c)^2/b`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.52

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx = \frac{-\cosh(2a + c + bx) + (\log(\cosh(\frac{1}{2}(c - bx))) - \log(\sinh(\frac{1}{2}(c - bx)))) \sinh^2(a + c)}{b}$$

input `Integrate[Csch[c - b*x]*Sinh[a + b*x]^2,x]`

output `(-Cosh[2*a + c + b*x] + (Log[Cosh[(c - b*x)/2]] - Log[Sinh[(c - b*x)/2]])*Sinh[a + c]^2)/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{csch}(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{csch}(c - bx) dx$$

input `Int[Csch[c - b*x]*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(34) = 68.

Time = 0.23 (sec) , antiderivative size = 221, normalized size of antiderivative = 6.70

method	result
risch	$-\frac{e^{bx+2a+c}}{2b} - \frac{e^{-bx-2a-c}}{2b} + \frac{\ln(e^{a+c}+e^{bx+a})e^{-2c-2a}e^{4a+4c}}{4b} - \frac{\ln(e^{a+c}+e^{bx+a})e^{-2c-2a}e^{2a+2c}}{2b} + \frac{\ln(e^{a+c}+e^{bx+a})e^{-2c-2a}}{4b}$

input `int(-csch(b*x-c)*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$-1/2/b*\exp(b*x+2*a+c)-1/2/b*\exp(-b*x-2*a-c)+1/4*\ln(\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)*\exp(4*a+4*c)-1/2*\ln(\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)*\exp(2*a+2*c)+1/4*\ln(\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)-1/4*\ln(-\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)*\exp(4*a+4*c)+1/2*\ln(-\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)*\exp(2*a+2*c)-1/4*\ln(-\exp(a+c)+\exp(b*x+a))/b*\exp(-2*c-2*a)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 665 vs. $2(34) = 68$.

Time = 0.11 (sec) , antiderivative size = 665, normalized size of antiderivative = 20.15

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
-1/4*(2*cosh(b*x + a)^2*cosh(a + c) + 2*cosh(a + c)^3 + 2*(cosh(a + c) - s
inh(a + c))*sinh(b*x + a)^2 + 6*cosh(a + c)*sinh(a + c)^2 - 2*sinh(a + c)^
3 + (4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)
^4 - 2*(3*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 4*(cosh(a + c)^
3 - cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 - 2*cosh(a + c)
)^2 + 1)*cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + si
nh(a + c)^4 + 2*(3*cosh(a + c)^2 - 1)*sinh(a + c)^2 - 2*cosh(a + c)^2 - 4*
(cosh(a + c)^3 - cosh(a + c))*sinh(a + c) + 1)*sinh(b*x + a))*log(cosh(b*x
+ a)*cosh(a + c) + (cosh(a + c) - sinh(a + c))*sinh(b*x + a) - cosh(b*x +
a)*sinh(a + c) + 1) - (4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b
*x + a)*sinh(a + c)^4 - 2*(3*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^
2 + 4*(cosh(a + c)^3 - cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a +
c)^4 - 2*cosh(a + c)^2 + 1)*cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)
)*sinh(a + c)^3 + sinh(a + c)^4 + 2*(3*cosh(a + c)^2 - 1)*sinh(a + c)^2 - 2
*cosh(a + c)^2 - 4*(cosh(a + c)^3 - cosh(a + c))*sinh(a + c) + 1)*sinh(b*x
+ a))*log(cosh(b*x + a)*cosh(a + c) + (cosh(a + c) - sinh(a + c))*sinh(b*
x + a) - cosh(b*x + a)*sinh(a + c) - 1) + 4*(cosh(b*x + a)*cosh(a + c) - c
osh(b*x + a)*sinh(a + c))*sinh(b*x + a) - 2*(cosh(b*x + a)^2 + 3*cosh(a +
c)^2)*sinh(a + c))/(b*cosh(b*x + a)*cosh(a + c)^2 - 2*b*cosh(b*x + a)*cosh
(a + c)*sinh(a + c) + b*cosh(b*x + a)*sinh(a + c)^2 + (b*cosh(a + c)^2 ...
```

Sympy [F]

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx = - \int \sinh^2(a + bx) \operatorname{csch}(bx - c) dx$$

input `integrate(-csch(b*x-c)*sinh(b*x+a)**2,x)`

output `-Integral(sinh(a + b*x)**2*csch(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(34) = 68$.

Time = 0.03 (sec) , antiderivative size = 118, normalized size of antiderivative = 3.58

$$\begin{aligned} & \int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx \\ &= \frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{4b} \\ & \quad - \frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{4b} - \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b} \end{aligned}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/4*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b - 1/4*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b - 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 117 vs. $2(34) = 68$.

Time = 0.11 (sec) , antiderivative size = 117, normalized size of antiderivative = 3.55

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(bx)} + e^c)}{4b} - \frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(|e^{(bx)} - e^c|)}{4b} - \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(-csch(b*x-c)*sinh(b*x+a)^2,x, algorithm="giac")`

output `1/4*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(b*x) + e^c)/b - 1/4*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(abs(e^(b*x) - e^c))/b - 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 202, normalized size of antiderivative = 6.12

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{-3c} e^{bx} (\sqrt{-b^2} - 2e^{2a} e^{2c} \sqrt{-b^2} + e^{4a} e^{4c} \sqrt{-b^2})}{b \sqrt{e^{-4a} e^{-4c} (6e^{4a} e^{4c} - 4e^{2a} e^{2c} - 4e^{6a} e^{6c} + e^{8a} e^{8c} + 1)}}\right) \sqrt{e^{-4a-4c} (6e^{4a+4c} - 4e^{2a+2c} - 4e^{6a+6c} + e^{8a+8c})}}{2\sqrt{-b^2}} - \frac{e^{-2a-c-bx}}{2b} - \frac{e^{2a+c+bx}}{2b}$$

input `int(sinh(a + b*x)^2/sinh(c - b*x),x)`

output

```
(atan((exp(-2*a)*exp(-3*c)*exp(b*x)*((-b^2)^(1/2) - 2*exp(2*a)*exp(2*c)*(-
b^2)^(1/2) + exp(4*a)*exp(4*c)*(-b^2)^(1/2)))/(b*(exp(-4*a)*exp(-4*c)*(6*exp
(4*a)*exp(4*c) - 4*exp(2*a)*exp(2*c) - 4*exp(6*a)*exp(6*c) + exp(8*a)*exp
(8*c) + 1))^(1/2)))*(exp(- 4*a - 4*c)*(6*exp(4*a + 4*c) - 4*exp(2*a + 2*c)
) - 4*exp(6*a + 6*c) + exp(8*a + 8*c) + 1))^(1/2))/(2*(-b^2)^(1/2)) - exp(
- 2*a - c - b*x)/(2*b) - exp(2*a + c + b*x)/(2*b)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 174, normalized size of antiderivative = 5.27

$$\int \operatorname{csch}(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{-2e^{2bx+4a+3c} + e^{bx+4a+4c} \log(e^{bx} + e^c) - e^{bx+4a+4c} \log(e^{bx} - e^c) - 2e^{bx+2a+2c} \log(e^{bx} + e^c) + 2e^{bx+2a+2c} \log(e^{bx} - e^c)}{4e^{bx+2a+2c} b}$$

input

```
int(-csch(b*x-c)*sinh(b*x+a)^2,x)
```

output

```
( - 2*e**(4*a + 2*b*x + 3*c) + e**(4*a + b*x + 4*c)*log(e**(b*x) + e**c) -
e**(4*a + b*x + 4*c)*log(e**(b*x) - e**c) - 2*e**(2*a + b*x + 2*c)*log(e*
*(b*x) + e**c) + 2*e**(2*a + b*x + 2*c)*log(e**(b*x) - e**c) + e**(b*x)*lo
g(e**(b*x) + e**c) - e**(b*x)*log(e**(b*x) - e**c) - 2*e**c)/(4*e**(2*a +
b*x + 2*c)*b)
```


3.84 $\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx$

Optimal result	616
Mathematica [A] (verified)	616
Rubi [F]	617
Maple [B] (verified)	617
Fricas [B] (verification not implemented)	618
Sympy [F]	619
Maxima [B] (verification not implemented)	619
Giac [B] (verification not implemented)	620
Mupad [B] (verification not implemented)	620
Reduce [B] (verification not implemented)	621

Optimal result

Integrand size = 18, antiderivative size = 44

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = x \cosh(2(a + c)) + \frac{\coth(c - bx) \sinh^2(a + c)}{b} + \frac{\log(\sinh(c - bx)) \sinh(2(a + c))}{b}$$

output `x*cosh(2*a+2*c)-coth(b*x-c)*sinh(a+c)^2/b+ln(-sinh(b*x-c))*sinh(2*a+2*c)/b`

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.11

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = \frac{bx \cosh(2(a + c)) + \log(\sinh(c - bx)) \sinh(2(a + c)) + \operatorname{csch}(c) \operatorname{csch}(c - bx) \sinh^2(a + c) \sinh(bx)}{b}$$

input `Integrate[Csch[c - b*x]^2*Sinh[a + b*x]^2,x]`

output `(b*x*Cosh[2*(a + c)] + Log[Sinh[c - b*x]]*Sinh[2*(a + c)] + Csch[c]*Csch[c - b*x]*Sinh[a + c]^2*Sinh[b*x])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

input `Int[Csch[c - b*x]^2*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. $2(53) = 106$.

Time = 0.32 (sec) , antiderivative size = 249, normalized size of antiderivative = 5.66

method	result
risch	$x e^{2a+2c} - e^{-2c-2a} e^{4a+4c} x - \frac{e^{-2c-2a} a e^{4a+4c}}{b} + e^{-2c-2a} x + \frac{e^{-2c-2a} a}{b} + \frac{e^{4a+4c}}{2b(e^{2a+2c} - e^{2bx+2a})} - \frac{e^{2a+2c}}{b(e^{2a+2c} - e^{2bx+2a})}$

input `int(csch(b*x-c)^2*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `x*exp(2*a+2*c)-exp(-2*c-2*a)*exp(4*a+4*c)*x-1/b*exp(-2*c-2*a)*a*exp(4*a+4*c)+exp(-2*c-2*a)*x+1/b*exp(-2*c-2*a)*a+1/2/b/(exp(2*a+2*c)-exp(2*b*x+2*a))*exp(4*a+4*c)-1/b/(exp(2*a+2*c)-exp(2*b*x+2*a))*exp(2*a+2*c)+1/2/b/(exp(2*a+2*c)-exp(2*b*x+2*a))+1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1311 vs. $2(53) = 106$.

Time = 0.12 (sec) , antiderivative size = 1311, normalized size of antiderivative = 29.80

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input `integrate(csch(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="fricas")`

output

```
1/2*(2*b*x*cosh(b*x + a)^2*cosh(a + c)^6 - 12*b*x*cosh(b*x + a)^2*cosh(a +
c)*sinh(a + c)^5 + 2*b*x*cosh(b*x + a)^2*sinh(a + c)^6 - (2*b*x + 1)*cosh
(a + c)^4 + (30*b*x*cosh(b*x + a)^2*cosh(a + c)^2 - 2*b*x - 1)*sinh(a + c)
^4 - 4*(10*b*x*cosh(b*x + a)^2*cosh(a + c)^3 - (2*b*x + 1)*cosh(a + c))*si
nh(a + c)^3 + 2*(b*x*cosh(a + c)^6 - 6*b*x*cosh(a + c)^5*sinh(a + c) + 15*
b*x*cosh(a + c)^4*sinh(a + c)^2 - 20*b*x*cosh(a + c)^3*sinh(a + c)^3 + 15*
b*x*cosh(a + c)^2*sinh(a + c)^4 - 6*b*x*cosh(a + c)*sinh(a + c)^5 + b*x*si
nh(a + c)^6)*sinh(b*x + a)^2 + 2*(15*b*x*cosh(b*x + a)^2*cosh(a + c)^4 - 3
*(2*b*x + 1)*cosh(a + c)^2 + 1)*sinh(a + c)^2 + 2*cosh(a + c)^2 + (6*cosh(
b*x + a)^2*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)^2*sinh(a + c)^6 - (15
*cosh(b*x + a)^2*cosh(a + c)^2 - 1)*sinh(a + c)^4 + cosh(a + c)^4 + 4*(5*c
osh(b*x + a)^2*cosh(a + c)^3 - cosh(a + c))*sinh(a + c)^3 - (cosh(a + c)^6
- cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^6 - 20*cosh(a + c)^3*sinh
(a + c)^3 + 15*cosh(a + c)^2*sinh(a + c)^4 - 6*cosh(a + c)*sinh(a + c)^5 +
sinh(a + c)^6 + (15*cosh(a + c)^4 - 1)*sinh(a + c)^2 - cosh(a + c)^2 - 2*
(3*cosh(a + c)^5 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((15*cosh(a
+ c)^4 - 1)*cosh(b*x + a)^2 - 6*cosh(a + c)^2)*sinh(a + c)^2 + 2*(20*cosh
(b*x + a)*cosh(a + c)^3*sinh(a + c)^3 - 15*cosh(b*x + a)*cosh(a + c)^2*si
nh(a + c)^4 + 6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)*sin
h(a + c)^6 - (15*cosh(a + c)^4 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(3*...
```

Sympy [F]

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}^2(bx - c) dx$$

input `integrate(csch(b*x-c)**2*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x - c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(53) = 106$.

Time = 0.04 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.05

$$\begin{aligned} \int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = & \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{2b} \\ & + \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{2b} \\ & + \frac{(bx + a)e^{(2a+2c)}}{b} + \frac{e^{(4a+4c)} - 2e^{(2a+2c)} + 1}{2b(e^{(-2bx+2a+4c)} - e^{(2a+2c)})} \end{aligned}$$

input `integrate(csch(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b + (b*x + a)*e^(2*a + 2*c)/b + 1/2*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)/(b*(e^(-2*b*x + 2*a + 4*c) - e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(53) = 106$.

Time = 0.12 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.52

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx$$

$$= x e^{(-2a-2c)} + \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(|e^{(2bx)} - e^{(2c)}|)}{2b}$$

$$+ \frac{(e^{(2bx)} - e^{(2bx+4a+4c)} + 2e^{(2a+4c)} - 2e^{(2c)})e^{(-2a-2c)}}{2b(e^{(2bx)} - e^{(2c)})}$$

input `integrate(csch(b*x-c)^2*sinh(b*x+a)^2,x, algorithm="giac")`

output `x*e^(-2*a - 2*c) + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(abs(e^(2*b*x) - e^(2*c)))/b + 1/2*(e^(2*b*x) - e^(2*b*x + 4*a + 4*c) + 2*e^(2*a + 4*c) - 2*e^(2*c))*e^(-2*a - 2*c)/(b*(e^(2*b*x) - e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.95

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx = x e^{-2a-2c} + \frac{\sinh(2a + 2c) \ln(e^{2a} e^{2bx} - e^{2a} e^{2c})}{b}$$

$$+ \frac{2e^{2a+2c} \sinh(a + c)^2}{b(e^{2a+2c} - e^{2a+2bx})}$$

input `int(sinh(a + b*x)^2/sinh(c - b*x)^2,x)`

output `x*exp(- 2*a - 2*c) + (sinh(2*a + 2*c)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp(2*c)))/b + (2*exp(2*a + 2*c)*sinh(a + c)^2)/(b*(exp(2*a + 2*c) - exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 256, normalized size of antiderivative = 5.82

$$\int \operatorname{csch}^2(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{e^{2bx+4a+4c} \log(e^{bx} + e^c) + e^{2bx+4a+4c} \log(e^{bx} - e^c) - e^{2bx+4a+4c} + 2e^{2bx+2a+2c} - e^{2bx} \log(e^{bx} + e^c) - e^{2bx} \log(e^{bx} - e^c)}{2e^c}$$

input `int(csch(b*x-c)^2*sinh(b*x+a)^2,x)`output `(e**(4*a + 2*b*x + 4*c)*log(e**(b*x) + e**c) + e**(4*a + 2*b*x + 4*c)*log(e**(b*x) - e**c) - e**(4*a + 2*b*x + 4*c) + 2*e**(2*a + 2*b*x + 2*c) - e**(2*b*x)*log(e**(b*x) + e**c) - e**(2*b*x)*log(e**(b*x) - e**c) + 2*e**(2*b*x)*b*x - e**(2*b*x) - e**(4*a + 6*c)*log(e**(b*x) + e**c) - e**(4*a + 6*c)*log(e**(b*x) - e**c) + e**(2*c)*log(e**(b*x) + e**c) + e**(2*c)*log(e**(b*x) - e**c) - 2*e**(2*c)*b*x)/(2*e**(2*a + 2*c)*b*(e**(2*b*x) - e**(2*c)))`

3.85 $\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx$

Optimal result	622
Mathematica [A] (verified)	622
Rubi [F]	623
Maple [B] (verified)	623
Fricas [B] (verification not implemented)	624
Sympy [F]	624
Maxima [B] (verification not implemented)	625
Giac [B] (verification not implemented)	625
Mupad [F(-1)]	626
Reduce [B] (verification not implemented)	626

Optimal result

Integrand size = 18, antiderivative size = 85

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = \frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh(2(a + c))}{b} - \frac{\operatorname{arctanh}(\cosh(c - bx)) \sinh^2(a + c)}{2b} + \frac{\operatorname{coth}(c - bx) \operatorname{csch}(c - bx) \sinh^2(a + c)}{2b} - \frac{\operatorname{csch}(c - bx) \sinh(2(a + c))}{b}$$

output

$$\operatorname{arctanh}(\cosh(b*x-c)) * \cosh(2*a+2*c) / b - 1/2 * \operatorname{arctanh}(\cosh(b*x-c)) * \sinh(a+c)^2 / b + 1/2 * \operatorname{coth}(b*x-c) * \operatorname{csch}(b*x-c) * \sinh(a+c)^2 / b + \operatorname{csch}(b*x-c) * \sinh(2*a+2*c) / b$$

Mathematica [A] (verified)

Time = 4.46 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.91

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = \frac{4(1 + 3 \cosh(2(a + c))) \log(\cosh(\frac{1}{2}(c - bx))) - 4(1 + 3 \cosh(2(a + c))) \log(\sinh(\frac{1}{2}(c - bx))) + 2 \operatorname{csch}^2(c - bx)}{b}$$

input `Integrate[Csch[c - b*x]^3*Sinh[a + b*x]^2,x]`

output $(4*(1 + 3*\text{Cosh}[2*(a + c)])*\text{Log}[\text{Cosh}[(c - b*x)/2]] - 4*(1 + 3*\text{Cosh}[2*(a + c)])*\text{Log}[\text{Sinh}[(c - b*x)/2]] + 2*\text{Csch}[(c - b*x)/2]^2*\text{Sinh}[a + c]^2 + 2*\text{Sech}[(c - b*x)/2]^2*\text{Sinh}[a + c]^2 - 8*\text{Csch}[c/2]*\text{Csch}[(c - b*x)/2]*\text{Sinh}[2*(a + c)]*\text{Sinh}[(b*x)/2] - 8*\text{Sech}[c/2]*\text{Sech}[(c - b*x)/2]*\text{Sinh}[2*(a + c)]*\text{Sinh}[(b*x)/2])/(16*b)$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \text{csch}^3(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \text{csch}^3(c - bx) dx$$

input `Int[Csch[c - b*x]^3*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(89) = 178.

Time = 0.63 (sec) , antiderivative size = 295, normalized size of antiderivative = 3.47

method	result
risch	$-\frac{(3e^{6a+6c}-5e^{2bx+6a+4c}+2e^{4a+4c}+2e^{2bx+4a+2c}-5e^{2a+2c}+3e^{2bx+2a})e^{bx-c}}{4(e^{2a+2c}-e^{2bx+2a})^2b} - \frac{3\ln(-e^{a+c}+e^{bx+a})e^{-2c-2a}e^{4a+4c}}{8b} - \frac{\ln(-$

input `int(-csch(b*x-c)^3*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*a+2*c)-exp(2*b*x+2*a))^2/b*(3*exp(6*a+6*c)-5*exp(2*b*x+6*a+4*c)
)+2*exp(4*a+4*c)+2*exp(2*b*x+4*a+2*c)-5*exp(2*a+2*c)+3*exp(2*b*x+2*a))*exp
(b*x-c)-3/8*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/4*ln(-
exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)-3/8*ln(-exp(a+c)+exp(b*x
+a))/b*exp(-2*c-2*a)+3/8*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4
*c)+1/4*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*ln(exp(a+
c)+exp(b*x+a))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5785 vs. $2(89) = 178$.

Time = 0.13 (sec) , antiderivative size = 5785, normalized size of antiderivative = 68.06

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(-csch(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = - \int \sinh^2(a + bx) \operatorname{csch}^3(bx - c) dx$$

input

```
integrate(-csch(b*x-c)**3*sinh(b*x+a)**2,x)
```

output

```
-Integral(sinh(a + b*x)**2*csch(b*x - c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. $2(89) = 178$.

Time = 0.05 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.40

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{8b}$$

$$- \frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{8b}$$

$$- \frac{(5e^{(4a+4c)} - 2e^{(2a+2c)} - 3)e^{(-bx-a)} - (3e^{(6a+6c)} + 2e^{(4a+4c)} - 5e^{(2a+2c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})}$$

input `integrate(-csch(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="maxima")`

output `1/8*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b - 1/8*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b - 1/4*((5*e^(4*a + 4*c) - 2*e^(2*a + 2*c) - 3)*e^(-b*x - a) - (3*e^(6*a + 6*c) + 2*e^(4*a + 4*c) - 5*e^(2*a + 2*c))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + 5*c) - e^(a + c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(89) = 178$.

Time = 0.12 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.22

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = \frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(bx)} + e^c)}{8b}$$

$$- \frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(|e^{(bx)} - e^c|)}{8b}$$

$$- \frac{(3e^{(3bx)} - 5e^{(3bx+4a+4c)} + 2e^{(3bx+2a+2c)} + 3e^{(bx+4a+6c)} + 2e^{(bx+2a+4c)} - 5e^{(bx+2c)})e^{(-2a-c)}}{4b(e^{(2bx)} - e^{(2c)})^2}$$

input `integrate(-csch(b*x-c)^3*sinh(b*x+a)^2,x, algorithm="giac")`

output

```
1/8*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(e^(b*x) + e
^c)/b - 1/8*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(abs
(e^(b*x) - e^c))/b - 1/4*(3*e^(3*b*x) - 5*e^(3*b*x + 4*a + 4*c) + 2*e^(3*b
*x + 2*a + 2*c) + 3*e^(b*x + 4*a + 6*c) + 2*e^(b*x + 2*a + 4*c) - 5*e^(b*x
+ 2*c))*e^(-2*a - c)/(b*(e^(2*b*x) - e^(2*c))^2)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\sinh(c - bx)^3} dx$$

input

```
int(sinh(a + b*x)^2/sinh(c - b*x)^3,x)
```

output

```
int(sinh(a + b*x)^2/sinh(c - b*x)^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 538, normalized size of antiderivative = 6.33

$$\int \operatorname{csch}^3(c - bx) \sinh^2(a + bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \log(e^{bx} + e^c) - 3e^{4bx+4a+4c} \log(e^{bx} - e^c) + 2e^{4bx+2a+2c} \log(e^{bx} + e^c) - 2e^{4bx+2a+2c} \log(e^{bx} - e^c)}{b}$$

input

```
int(-csch(b*x-c)^3*sinh(b*x+a)^2,x)
```

output

```
(3*e**(4*a + 4*b*x + 4*c)*log(e**(b*x) + e**c) - 3*e**(4*a + 4*b*x + 4*c)*
log(e**(b*x) - e**c) + 2*e**(2*a + 4*b*x + 2*c)*log(e**(b*x) + e**c) - 2*e
**(2*a + 4*b*x + 2*c)*log(e**(b*x) - e**c) + 3*e**(4*b*x)*log(e**(b*x) + e
**c) - 3*e**(4*b*x)*log(e**(b*x) - e**c) + 10*e**(4*a + 3*b*x + 5*c) - 4*e
**(2*a + 3*b*x + 3*c) - 6*e**(3*b*x + c) - 6*e**(4*a + 2*b*x + 6*c)*log(e*
*(b*x) + e**c) + 6*e**(4*a + 2*b*x + 6*c)*log(e**(b*x) - e**c) - 4*e**(2*a
+ 2*b*x + 4*c)*log(e**(b*x) + e**c) + 4*e**(2*a + 2*b*x + 4*c)*log(e**(b*
x) - e**c) - 6*e**(2*b*x + 2*c)*log(e**(b*x) + e**c) + 6*e**(2*b*x + 2*c)*
log(e**(b*x) - e**c) - 6*e**(4*a + b*x + 7*c) - 4*e**(2*a + b*x + 5*c) + 1
0*e**(b*x + 3*c) + 3*e**(4*a + 8*c)*log(e**(b*x) + e**c) - 3*e**(4*a + 8*c
)*log(e**(b*x) - e**c) + 2*e**(2*a + 6*c)*log(e**(b*x) + e**c) - 2*e**(2*a
+ 6*c)*log(e**(b*x) - e**c) + 3*e**(4*c)*log(e**(b*x) + e**c) - 3*e**(4*c
)*log(e**(b*x) - e**c))/(8*e**(2*a + 2*c)*b*(e**(4*b*x) - 2*e**(2*b*x + 2*
c) + e**(4*c)))
```

3.86 $\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx$

Optimal result	628
Mathematica [A] (verified)	628
Rubi [F]	629
Maple [A] (verified)	629
Fricas [B] (verification not implemented)	630
Sympy [F]	631
Maxima [B] (verification not implemented)	631
Giac [A] (verification not implemented)	632
Mupad [F(-1)]	632
Reduce [B] (verification not implemented)	633

Optimal result

Integrand size = 18, antiderivative size = 80

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \frac{\cosh(2(a + c)) \coth(c - bx)}{b} - \frac{\coth(c - bx) \sinh^2(a + c)}{b} + \frac{\coth^3(c - bx) \sinh^2(a + c)}{3b} - \frac{\operatorname{csch}^2(c - bx) \sinh(2(a + c))}{2b}$$

output

```
-cosh(2*a+2*c)*coth(b*x-c)/b+coth(b*x-c)*sinh(a+c)^2/b-1/3*coth(b*x-c)^3*
sinh(a+c)^2/b-1/2*csch(b*x-c)^2*sinh(2*a+2*c)/b
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \frac{\operatorname{csch}(c) \operatorname{csch}^3(c - bx) (3 \sinh(bx) + \sinh(2c - 3bx) + \sinh(2a + 4c - 3bx)) - 3 \sinh(2a + 2c - bx) + 3 \sinh(2a + 2c - bx)}{12b}$$

input `Integrate[Csch[c - b*x]^4*Sinh[a + b*x]^2,x]`

output `-1/12*(Csch[c]*Csch[c - b*x]^3*(3*Sinh[b*x] + Sinh[2*c - 3*b*x] + Sinh[2*a + 4*c - 3*b*x] - 3*Sinh[2*a + 2*c - b*x] + 3*Sinh[2*a + b*x] - Sinh[2*a + 3*b*x]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sinh^2(a + bx) \operatorname{csch}^4(c - bx) dx$$

↓ 7299

$$\int \sinh^2(a + bx) \operatorname{csch}^4(c - bx) dx$$

input `Int[Csch[c - b*x]^4*Sinh[a + b*x]^2,x]`

output `$Aborted`

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.80

method	result	size
parallelrisch	$\frac{\operatorname{sech}\left(\frac{bx}{2} - \frac{c}{2}\right)^3 \operatorname{csch}\left(\frac{bx}{2} - \frac{c}{2}\right)^3 (-2 \cosh(3bx + 2a - c) - \cosh(3bx - 3c) + 3 \cosh(bx - c))}{96b}$	64
risch	$\frac{2(e^{4a+4c} - 3e^{2bx+4a+2c} + e^{2a+2c} + 3e^{4bx+4a} - 3e^{2bx+2a} + 1)e^{4a+4c}}{3(e^{2a+2c} - e^{2bx+2a})^3 b}$	90

input `int(csch(b*x-c)^4*sinh(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
1/96*sech(1/2*b*x-1/2*c)^3*csch(1/2*b*x-1/2*c)^3*(-2*cosh(3*b*x+2*a-c)-cos
h(3*b*x-3*c)+3*cosh(b*x-c))/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1007 vs. $2(84) = 168$.

Time = 0.09 (sec) , antiderivative size = 1007, normalized size of antiderivative = 12.59

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \text{Too large to display}$$

input

```
integrate(csch(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="fricas")
```

output

```
-2/3*(4*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^3 + (5*cosh(b*x + a)^2 - 3
)*sinh(a + c)^4 - 3*cosh(a + c)^4 + (5*cosh(a + c)^4 + cosh(a + c)^2)*cosh
(b*x + a)^2 + (5*cosh(a + c)^4 + 4*cosh(a + c)*sinh(a + c)^3 + 5*sinh(a +
c)^4 - (2*cosh(a + c)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(2*cosh(a +
c)^3 + cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((2*cosh(a + c)^2 - 1)
*cosh(b*x + a)^2 - 6*cosh(a + c)^2 + 3)*sinh(a + c)^2 - 3*cosh(a + c)^2 -
2*(4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4
+ (10*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3
+ cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 - cosh(a + c)^2
)*cosh(b*x + a))*sinh(b*x + a) + 2*((2*cosh(a + c)^3 + cosh(a + c))*cosh(b
*x + a)^2 - 3*cosh(a + c))*sinh(a + c))/(b*cosh(b*x + a)^4*cosh(a + c)^4 -
4*b*cosh(b*x + a)^2*cosh(a + c)^4 + 3*b*cosh(a + c)^4 + (b*cosh(a + c)^4
- 4*b*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(a + c)^2*sinh(a + c)^2 - 4*b*co
sh(a + c)*sinh(a + c)^3 + b*sinh(a + c)^4)*sinh(b*x + a)^4 + (b*cosh(b*x +
a)^4 + 4*b*cosh(b*x + a)^2 + 3*b)*sinh(a + c)^4 + 4*(b*cosh(b*x + a)*cosh
(a + c)^4 - 4*b*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(b*x + a
)*cosh(a + c)^2*sinh(a + c)^2 - 4*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^
3 + b*cosh(b*x + a)*sinh(a + c)^4)*sinh(b*x + a)^3 - 4*(b*cosh(b*x + a)^4*
cosh(a + c) + b*cosh(b*x + a)^2*cosh(a + c))*sinh(a + c)^3 + 2*(3*b*cosh(b
*x + a)^2*cosh(a + c)^4 + 18*b*cosh(b*x + a)^2*cosh(a + c)^2*sinh(a + c)...
```

Sympy [F]

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \int \sinh^2(a + bx) \operatorname{csch}^4(bx - c) dx$$

input `integrate(csch(b*x-c)**4*sinh(b*x+a)**2,x)`

output `Integral(sinh(a + b*x)**2*csch(b*x - c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. $2(84) = 168$.

Time = 0.04 (sec) , antiderivative size = 342, normalized size of antiderivative = 4.28

$$\begin{aligned} & \int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx \\ &= -\frac{2(e^{4a+4c} + e^{2a+2c})e^{(-2bx-2a)}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} - e^{(2a+2c)})} \\ & \quad + \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} - e^{(2a+2c)})} \\ & \quad + \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} - e^{(2a+2c)})} \\ & \quad + \frac{2e^{(2a+2c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} - e^{(2a+2c)})} \\ & \quad + \frac{2}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} - e^{(2a+2c)})} \end{aligned}$$

input `integrate(csch(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="maxima")`

output

```
-2*(e^(4*a + 4*c) + e^(2*a + 2*c))*e^(-2*b*x - 2*a)/(b*(3*e^(-2*b*x + 2*a + 4*c) - 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) - e^(2*a + 2*c))) + 2*e^(-4*b*x + 4*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) - 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) - e^(2*a + 2*c))) + 2/3*e^(4*a + 4*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) - 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) - e^(2*a + 2*c))) + 2/3*e^(2*a + 2*c)/(b*(3*e^(-2*b*x + 2*a + 4*c) - 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) - e^(2*a + 2*c))) + 2/3/(b*(3*e^(-2*b*x + 2*a + 4*c) - 3*e^(-4*b*x + 2*a + 6*c) + e^(-6*b*x + 2*a + 8*c) - e^(2*a + 2*c)))
```

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.08

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \frac{2(3e^{(4bx+4a+4c)} - 3e^{(2bx+4a+6c)} - 3e^{(2bx+2a+4c)} + e^{(4a+8c)} + e^{(2a+6c)} + e^{(4c)})e^{(-2a)}}{3b(e^{(2bx)} - e^{(2c)})^3}$$

input

```
integrate(csch(b*x-c)^4*sinh(b*x+a)^2,x, algorithm="giac")
```

output

```
-2/3*(3*e^(4*b*x + 4*a + 4*c) - 3*e^(2*b*x + 4*a + 6*c) - 3*e^(2*b*x + 2*a + 4*c) + e^(4*a + 8*c) + e^(2*a + 6*c) + e^(4*c))*e^(-2*a)/(b*(e^(2*b*x) - e^(2*c))^3)
```

Mupad [F(-1)]

Timed out.

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \int \frac{\sinh(a + bx)^2}{\sinh(c - bx)^4} dx$$

input

```
int(sinh(a + b*x)^2/sinh(c - b*x)^4,x)
```

output

```
int(sinh(a + b*x)^2/sinh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.29

$$\int \operatorname{csch}^4(c - bx) \sinh^2(a + bx) dx = \frac{2e^{2c}(-e^{6bx+4a} + 3e^{2bx+2a+2c} - e^{2a+4c} - e^{2c})}{3e^{2ab}(e^{6bx} - 3e^{4bx+2c} + 3e^{2bx+4c} - e^{6c})}$$

input `int(csch(b*x-c)^4*sinh(b*x+a)^2,x)`output `(2*e**(2*c)*(- e**(4*a + 6*b*x) + 3*e**(2*a + 2*b*x + 2*c) - e**(2*a + 4*c) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x) - 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) - e**(6*c)))`

3.87 $\int \cosh(x) \sinh(2x) dx$

Optimal result	634
Mathematica [A] (verified)	634
Rubi [C] (verified)	635
Maple [A] (verified)	636
Fricas [B] (verification not implemented)	636
Sympy [B] (verification not implemented)	637
Maxima [B] (verification not implemented)	637
Giac [B] (verification not implemented)	637
Mupad [B] (verification not implemented)	638
Reduce [B] (verification not implemented)	638

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \cosh(x) \sinh(2x) dx = \frac{2 \cosh^3(x)}{3}$$

output `2/3*cosh(x)^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.88

$$\int \cosh(x) \sinh(2x) dx = \frac{\cosh(x)}{2} + \frac{1}{6} \cosh(3x)$$

input `Integrate[Cosh[x]*Sinh[2*x],x]`

output `Cosh[x]/2 + Cosh[3*x]/6`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.88, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(2x) \cosh(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(2ix) \cos(ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(ix) \sin(2ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{2} i \cosh(x) + \frac{1}{6} i \cosh(3x) \right) \end{aligned}$$

input `Int[Cosh[x]*Sinh[2*x],x]`

output `(-I)*((I/2)*Cosh[x] + (I/6)*Cosh[3*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.50

method	result	size
default	$\frac{\cosh(x)}{2} + \frac{\cosh(3x)}{6}$	12
paralletrisch	$\frac{\cosh(3x)}{6} + \frac{\cosh(x)}{2} - \frac{2}{3}$	13
orering	$-\frac{\sinh(x)\sinh(2x)}{3} + \frac{2\cosh(x)\cosh(2x)}{3}$	18
risch	$\frac{e^{3x}}{12} + \frac{e^x}{4} + \frac{e^{-x}}{4} + \frac{e^{-3x}}{12}$	24

input

```
int(cosh(x)*sinh(2*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*cosh(x)+1/6*cosh(3*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(6) = 12.

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \cosh(x) \sinh(2x) dx = \frac{1}{6} \cosh(x)^3 + \frac{1}{2} \cosh(x) \sinh(x)^2 + \frac{1}{2} \cosh(x)$$

input

```
integrate(cosh(x)*sinh(2*x),x, algorithm="fricas")
```

output

```
1/6*cosh(x)^3 + 1/2*cosh(x)*sinh(x)^2 + 1/2*cosh(x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(7) = 14$.

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 2.50

$$\int \cosh(x) \sinh(2x) dx = -\frac{\sinh(x) \sinh(2x)}{3} + \frac{2 \cosh(x) \cosh(2x)}{3}$$

input `integrate(cosh(x)*sinh(2*x),x)`

output `-sinh(x)*sinh(2*x)/3 + 2*cosh(x)*cosh(2*x)/3`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(6) = 12$.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 3.38

$$\int \cosh(x) \sinh(2x) dx = \frac{1}{12} (3 e^{(-2x)} + 1) e^{(3x)} + \frac{1}{4} e^{(-x)} + \frac{1}{12} e^{(-3x)}$$

input `integrate(cosh(x)*sinh(2*x),x, algorithm="maxima")`

output `1/12*(3*e^(-2*x) + 1)*e^(3*x) + 1/4*e^(-x) + 1/12*e^(-3*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(6) = 12$.

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 3.12

$$\int \cosh(x) \sinh(2x) dx = \frac{1}{12} (3 e^{(2x)} + 1) e^{(-3x)} + \frac{1}{12} e^{(3x)} + \frac{1}{4} e^x$$

input `integrate(cosh(x)*sinh(2*x),x, algorithm="giac")`

output $1/12*(3*e^{(2*x)} + 1)*e^{(-3*x)} + 1/12*e^{(3*x)} + 1/4*e^x$

Mupad [B] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \cosh(x) \sinh(2x) dx = \frac{2 \cosh(x)^3}{3}$$

input `int(sinh(2*x)*cosh(x),x)`

output $(2*\cosh(x)^3)/3$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.12

$$\int \cosh(x) \sinh(2x) dx = \frac{2 \cosh(2x) \cosh(x)}{3} - \frac{\sinh(2x) \sinh(x)}{3}$$

input `int(cosh(x)*sinh(2*x),x)`

output $(2*\cosh(2*x)*\cosh(x) - \sinh(2*x)*\sinh(x))/3$

3.88 $\int \cosh(x) \sinh(3x) dx$

Optimal result	639
Mathematica [A] (verified)	639
Rubi [C] (verified)	640
Maple [A] (verified)	641
Fricas [B] (verification not implemented)	641
Sympy [A] (verification not implemented)	642
Maxima [B] (verification not implemented)	642
Giac [A] (verification not implemented)	642
Mupad [B] (verification not implemented)	643
Reduce [B] (verification not implemented)	643

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(x) \sinh(3x) dx = \frac{1}{4} \cosh(2x) + \frac{1}{8} \cosh(4x)$$

output `1/4*cosh(2*x)+1/8*cosh(4*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \sinh(3x) dx = \frac{\cosh^2(x)}{2} + \frac{1}{8} \cosh(4x)$$

input `Integrate[Cosh[x]*Sinh[3*x],x]`

output `Cosh[x]^2/2 + Cosh[4*x]/8`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(3x) \cosh(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(3ix) \cos(ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(ix) \sin(3ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{4} i \cosh(2x) + \frac{1}{8} i \cosh(4x) \right) \end{aligned}$$

input `Int[Cosh[x]*Sinh[3*x],x]`

output `(-I)*((I/4)*Cosh[2*x] + (I/8)*Cosh[4*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\cosh(2x)}{4} + \frac{\cosh(4x)}{8}$	14
paralelrisch	$\frac{\cosh(4x)}{8} - \frac{3}{8} + \frac{\cosh(2x)}{4}$	15
orering	$-\frac{\sinh(x)\sinh(3x)}{8} + \frac{3\cosh(x)\cosh(3x)}{8}$	18
risch	$\frac{e^{4x}}{16} + \frac{e^{2x}}{8} + \frac{e^{-2x}}{8} + \frac{e^{-4x}}{16}$	26

input

```
int(cosh(x)*sinh(3*x),x,method=_RETURNVERBOSE)
```

output

```
1/4*cosh(2*x)+1/8*cosh(4*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.94

$$\int \cosh(x) \sinh(3x) dx = \frac{1}{8} \cosh(x)^4 + \frac{1}{8} \sinh(x)^4 + \frac{1}{4} (3 \cosh(x)^2 + 1) \sinh(x)^2 + \frac{1}{4} \cosh(x)^2$$

input

```
integrate(cosh(x)*sinh(3*x),x, algorithm="fricas")
```

output

```
1/8*cosh(x)^4 + 1/8*sinh(x)^4 + 1/4*(3*cosh(x)^2 + 1)*sinh(x)^2 + 1/4*cosh
(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \sinh(3x) dx = -\frac{\sinh(x) \sinh(3x)}{8} + \frac{3 \cosh(x) \cosh(3x)}{8}$$

input `integrate(cosh(x)*sinh(3*x),x)`

output `-sinh(x)*sinh(3*x)/8 + 3*cosh(x)*cosh(3*x)/8`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \sinh(3x) dx = \frac{1}{16} (2e^{-2x} + 1)e^{4x} + \frac{1}{8} e^{-2x} + \frac{1}{16} e^{-4x}$$

input `integrate(cosh(x)*sinh(3*x),x, algorithm="maxima")`

output `1/16*(2*e^(-2*x) + 1)*e^(4*x) + 1/8*e^(-2*x) + 1/16*e^(-4*x)`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

$$\int \cosh(x) \sinh(3x) dx = \frac{1}{16} (e^{2x} + e^{-2x})^2 + \frac{1}{8} e^{2x} + \frac{1}{8} e^{-2x}$$

input `integrate(cosh(x)*sinh(3*x),x, algorithm="giac")`

output `1/16*(e^(2*x) + e^(-2*x))^2 + 1/8*e^(2*x) + 1/8*e^(-2*x)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \cosh(x) \sinh(3x) dx = \cosh(x)^4 - \frac{\cosh(x)^2}{2}$$

input `int(sinh(3*x)*cosh(x),x)`

output `cosh(x)^4 - cosh(x)^2/2`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \sinh(3x) dx = \frac{3 \cosh(3x) \cosh(x)}{8} - \frac{\sinh(3x) \sinh(x)}{8}$$

input `int(cosh(x)*sinh(3*x),x)`

output `(3*cosh(3*x)*cosh(x) - sinh(3*x)*sinh(x))/8`

3.89 $\int \cosh(x) \sinh(4x) dx$

Optimal result	644
Mathematica [A] (verified)	644
Rubi [C] (verified)	645
Maple [A] (verified)	646
Fricas [B] (verification not implemented)	646
Sympy [A] (verification not implemented)	647
Maxima [B] (verification not implemented)	647
Giac [B] (verification not implemented)	647
Mupad [B] (verification not implemented)	648
Reduce [B] (verification not implemented)	648

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(x) \sinh(4x) dx = \frac{1}{6} \cosh(3x) + \frac{1}{10} \cosh(5x)$$

output `1/6*cosh(3*x)+1/10*cosh(5*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \sinh(4x) dx = \frac{1}{6} \cosh(3x) + \frac{1}{10} \cosh(5x)$$

input `Integrate[Cosh[x]*Sinh[4*x],x]`

output `Cosh[3*x]/6 + Cosh[5*x]/10`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 26, 4772}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sinh(4x) \cosh(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \sin(4ix) \cos(ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \cos(ix) \sin(4ix) dx \\ & \quad \downarrow \text{4772} \\ & -i \left(\frac{1}{6} i \cosh(3x) + \frac{1}{10} i \cosh(5x) \right) \end{aligned}$$

input `Int[Cosh[x]*Sinh[4*x],x]`

output `(-I)*((I/6)*Cosh[3*x] + (I/10)*Cosh[5*x])`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4772

```
Int[cos[(c_.) + (d_.)*(x_)]*sin[(a_.) + (b_.)*(x_)], x_Symbol] := Simp[-Cos
[a - c + (b - d)*x]/(2*(b - d)), x] - Simp[Cos[a + c + (b + d)*x]/(2*(b + d
)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\cosh(3x)}{6} + \frac{\cosh(5x)}{10}$	14
parallelsch	$-\frac{4}{15} + \frac{\cosh(5x)}{10} + \frac{\cosh(3x)}{6}$	15
orering	$-\frac{\sinh(x)\sinh(4x)}{15} + \frac{4\cosh(x)\cosh(4x)}{15}$	18
risch	$\frac{e^{5x}}{20} + \frac{e^{3x}}{12} + \frac{e^{-3x}}{12} + \frac{e^{-5x}}{20}$	26

input

```
int(cosh(x)*sinh(4*x),x,method=_RETURNVERBOSE)
```

output

```
1/6*cosh(3*x)+1/10*cosh(5*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.12

$$\int \cosh(x) \sinh(4x) dx = \frac{1}{10} \cosh(x)^5 + \frac{1}{2} \cosh(x) \sinh(x)^4 + \frac{1}{6} \cosh(x)^3 + \frac{1}{2} (2 \cosh(x)^3 + \cosh(x)) \sinh(x)^2$$

input

```
integrate(cosh(x)*sinh(4*x),x, algorithm="fricas")
```

output

```
1/10*cosh(x)^5 + 1/2*cosh(x)*sinh(x)^4 + 1/6*cosh(x)^3 + 1/2*(2*cosh(x)^3
+ cosh(x))*sinh(x)^2
```

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \sinh(4x) dx = -\frac{\sinh(x) \sinh(4x)}{15} + \frac{4 \cosh(x) \cosh(4x)}{15}$$

input `integrate(cosh(x)*sinh(4*x),x)`

output `-sinh(x)*sinh(4*x)/15 + 4*cosh(x)*cosh(4*x)/15`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \sinh(4x) dx = \frac{1}{60} (5 e^{(-2x)} + 3) e^{(5x)} + \frac{1}{12} e^{(-3x)} + \frac{1}{20} e^{(-5x)}$$

input `integrate(cosh(x)*sinh(4*x),x, algorithm="maxima")`

output `1/60*(5*e^(-2*x) + 3)*e^(5*x) + 1/12*e^(-3*x) + 1/20*e^(-5*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.12 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \sinh(4x) dx = \frac{1}{60} (5 e^{(2x)} + 3) e^{(-5x)} + \frac{1}{20} e^{(5x)} + \frac{1}{12} e^{(3x)}$$

input `integrate(cosh(x)*sinh(4*x),x, algorithm="giac")`

output `1/60*(5*e^(2*x) + 3)*e^(-5*x) + 1/20*e^(5*x) + 1/12*e^(3*x)`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \cosh(x) \sinh(4x) dx = \frac{4 \cosh(x)^3 (6 \cosh(x)^2 - 5)}{15}$$

input `int(sinh(4*x)*cosh(x),x)`

output `(4*cosh(x)^3*(6*cosh(x)^2 - 5))/15`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \sinh(4x) dx = \frac{4 \cosh(4x) \cosh(x)}{15} - \frac{\sinh(4x) \sinh(x)}{15}$$

input `int(cosh(x)*sinh(4*x),x)`

output `(4*cosh(4*x)*cosh(x) - sinh(4*x)*sinh(x))/15`

3.90 $\int \cosh(x) \sinh(mx) dx$

Optimal result	649
Mathematica [A] (verified)	649
Rubi [A] (verified)	650
Maple [A] (verified)	651
Fricas [A] (verification not implemented)	651
Sympy [A] (verification not implemented)	652
Maxima [F(-2)]	652
Giac [B] (verification not implemented)	653
Mupad [B] (verification not implemented)	653
Reduce [B] (verification not implemented)	653

Optimal result

Integrand size = 7, antiderivative size = 35

$$\int \cosh(x) \sinh(mx) dx = -\frac{\cosh((1-m)x)}{2(1-m)} + \frac{\cosh((1+m)x)}{2(1+m)}$$

output

```
-1/2*cosh((1-m)*x)/(1-m)+cosh((1+m)*x)/(2+2*m)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(x) \sinh(mx) dx = \frac{m \cosh(x) \cosh(mx) - \sinh(x) \sinh(mx)}{-1 + m^2}$$

input

```
Integrate[Cosh[x]*Sinh[m*x],x]
```

output

```
(m*Cosh[x]*Cosh[m*x] - Sinh[x]*Sinh[m*x])/(-1 + m^2)
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6152, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \sinh(mx) dx$$

$$\downarrow 6152$$

$$\int \left(\frac{1}{2} \sinh((m+1)x) - \frac{1}{2} \sinh((1-m)x) \right) dx$$

$$\downarrow 2009$$

$$\frac{\cosh((m+1)x)}{2(m+1)} - \frac{\cosh((1-m)x)}{2(1-m)}$$

input `Int[Cosh[x]*Sinh[m*x],x]`

output `-1/2*Cosh[(1 - m)*x]/(1 - m) + Cosh[(1 + m)*x]/(2*(1 + m))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6152 `Int[Cosh[w_]^(q_.)*Sinh[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sinh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$\frac{\cosh(x(-1+m))}{-2+2m} + \frac{\cosh((1+m)x)}{2+2m}$
parallelsch	$\frac{(1+m) \cosh(x(-1+m)) + (-1+m) \cosh((1+m)x) - 2m}{2m^2 - 2}$
risch	$\frac{(m e^{2x} - e^{2x} + m + 1) e^{x(-1+m)}}{4(1+m)(-1+m)} + \frac{(m e^{2x} + e^{2x} + m - 1) e^{-(1+m)x}}{4(1+m)(-1+m)}$
orering	$\frac{2(m^2+1)(\sinh(x) \sinh(mx) + \cosh(x)m \cosh(mx))}{m^4 - 2m^2 + 1} - \frac{\sinh(x) \sinh(mx) + 3 \cosh(x)m \cosh(mx) + 3 \sinh(x)m^2 \sinh(mx) + \cosh(x)m^2 \cosh(mx)}{m^4 - 2m^2 + 1}$

input `int(cosh(x)*sinh(m*x), x, method=_RETURNVERBOSE)`output `1/2*cosh(x*(-1+m))/(-1+m)+1/2*cosh((1+m)*x)/(1+m)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \cosh(x) \sinh(mx) dx = \frac{m \cosh(mx) \cosh(x) - \sinh(mx) \sinh(x)}{(m^2 - 1) \cosh(x)^2 - (m^2 - 1) \sinh(x)^2}$$

input `integrate(cosh(x)*sinh(m*x), x, algorithm="fricas")`output `(m*cosh(m*x)*cosh(x) - sinh(m*x)*sinh(x))/((m^2 - 1)*cosh(x)^2 - (m^2 - 1)*sinh(x)^2)`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \cosh(x) \sinh(mx) dx = \begin{cases} -\frac{\sinh^2(x)}{2} & \text{for } m = -1 \\ \frac{\sinh^2(x)}{2} & \text{for } m = 1 \\ \frac{m \cosh(x) \cosh(mx)}{m^2-1} - \frac{\sinh(x) \sinh(mx)}{m^2-1} & \text{otherwise} \end{cases}$$

input `integrate(cosh(x)*sinh(m*x),x)`output `Piecewise((-sinh(x)**2/2, Eq(m, -1)), (sinh(x)**2/2, Eq(m, 1)), (m*cosh(x)*cosh(m*x)/(m**2 - 1) - sinh(x)*sinh(m*x)/(m**2 - 1), True))`**Maxima [F(-2)]**

Exception generated.

$$\int \cosh(x) \sinh(mx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(x)*sinh(m*x),x, algorithm="maxima")`output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see `assume?` for more details)Is`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \cosh(x) \sinh(mx) dx = \frac{e^{(mx+x)}}{4(m+1)} + \frac{e^{(mx-x)}}{4(m-1)} + \frac{e^{(-mx+x)}}{4(m-1)} + \frac{e^{(-mx-x)}}{4(m+1)}$$

input `integrate(cosh(x)*sinh(m*x),x, algorithm="giac")`

output `1/4*e^(m*x + x)/(m + 1) + 1/4*e^(m*x - x)/(m - 1) + 1/4*e^(-m*x + x)/(m - 1) + 1/4*e^(-m*x - x)/(m + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \cosh(x) \sinh(mx) dx = -\frac{\sinh(mx) \sinh(x) - m \cosh(mx) \cosh(x)}{m^2 - 1}$$

input `int(sinh(m*x)*cosh(x),x)`

output `-(sinh(m*x)*sinh(x) - m*cosh(m*x)*cosh(x))/(m^2 - 1)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(x) \sinh(mx) dx = \frac{\cosh(mx) \cosh(x) m - \sinh(mx) \sinh(x)}{m^2 - 1}$$

input `int(cosh(x)*sinh(m*x),x)`

output `(cosh(m*x)*cosh(x)*m - sinh(m*x)*sinh(x))/(m**2 - 1)`

3.91 $\int \cosh(x) \cosh(2x) dx$

Optimal result	654
Mathematica [A] (verified)	654
Rubi [A] (verified)	655
Maple [A] (verified)	656
Fricas [A] (verification not implemented)	656
Sympy [A] (verification not implemented)	656
Maxima [B] (verification not implemented)	657
Giac [B] (verification not implemented)	657
Mupad [B] (verification not implemented)	658
Reduce [B] (verification not implemented)	658

Optimal result

Integrand size = 7, antiderivative size = 15

$$\int \cosh(x) \cosh(2x) dx = \frac{\sinh(x)}{2} + \frac{1}{6} \sinh(3x)$$

output `1/2*sinh(x)+1/6*sinh(3*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cosh(x) \cosh(2x) dx = \frac{\sinh(x)}{2} + \frac{1}{6} \sinh(3x)$$

input `Integrate[Cosh[x]*Cosh[2*x],x]`

output `Sinh[x]/2 + Sinh[3*x]/6`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4771}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \cosh(2x) dx$$

$$\downarrow 3042$$

$$\int \cos(ix) \cos(2ix) dx$$

$$\downarrow 4771$$

$$\frac{\sinh(x)}{2} + \frac{1}{6} \sinh(3x)$$

input `Int[Cosh[x]*Cosh[2*x],x]`

output `Sinh[x]/2 + Sinh[3*x]/6`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4771 `Int[cos[(a_.) + (b_.)*(x_)]*cos[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] + Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\sinh(x)}{2} + \frac{\sinh(3x)}{6}$	12
parallelsch	$\frac{\sinh(x)}{2} + \frac{\sinh(3x)}{6}$	12
orering	$-\frac{\cosh(2x)\sinh(x)}{3} + \frac{2\cosh(x)\sinh(2x)}{3}$	18
risch	$\frac{e^{3x}}{12} + \frac{e^x}{4} - \frac{e^{-x}}{4} - \frac{e^{-3x}}{12}$	24

input `int(cosh(x)*cosh(2*x),x,method=_RETURNVERBOSE)`output `1/2*sinh(x)+1/6*sinh(3*x)`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cosh(x) \cosh(2x) dx = \frac{1}{6} \sinh(x)^3 + \frac{1}{2} (\cosh(x)^2 + 1) \sinh(x)$$

input `integrate(cosh(x)*cosh(2*x),x, algorithm="fricas")`output `1/6*sinh(x)^3 + 1/2*(cosh(x)^2 + 1)*sinh(x)`**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \cosh(x) \cosh(2x) dx = -\frac{\sinh(x) \cosh(2x)}{3} + \frac{2 \sinh(2x) \cosh(x)}{3}$$

input `integrate(cosh(x)*cosh(2*x),x)`

output $-\sinh(x)\cosh(2x)/3 + 2\sinh(2x)\cosh(x)/3$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(11) = 22.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \cosh(x) \cosh(2x) dx = \frac{1}{12} (3e^{(-2x)} + 1)e^{(3x)} - \frac{1}{4}e^{(-x)} - \frac{1}{12}e^{(-3x)}$$

input `integrate(cosh(x)*cosh(2*x),x, algorithm="maxima")`

output $1/12*(3*e^{(-2*x)} + 1)*e^{(3*x)} - 1/4*e^{(-x)} - 1/12*e^{(-3*x)}$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(11) = 22.

Time = 0.18 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.67

$$\int \cosh(x) \cosh(2x) dx = -\frac{1}{12} (3e^{(2x)} + 1)e^{(-3x)} + \frac{1}{12}e^{(3x)} + \frac{1}{4}e^x$$

input `integrate(cosh(x)*cosh(2*x),x, algorithm="giac")`

output $-1/12*(3*e^{(2*x)} + 1)*e^{(-3*x)} + 1/12*e^{(3*x)} + 1/4*e^x$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.60

$$\int \cosh(x) \cosh(2x) dx = \frac{2 \sinh(x)^3}{3} + \sinh(x)$$

input `int(cosh(2*x)*cosh(x),x)`

output `sinh(x) + (2*sinh(x)^3)/3`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cosh(x) \cosh(2x) dx = -\frac{\cosh(2x) \sinh(x)}{3} + \frac{2 \cosh(x) \sinh(2x)}{3}$$

input `int(cosh(x)*cosh(2*x),x)`

output `(- cosh(2*x)*sinh(x) + 2*cosh(x)*sinh(2*x))/3`

3.92 $\int \cosh(x) \cosh(3x) dx$

Optimal result	659
Mathematica [A] (verified)	659
Rubi [A] (verified)	660
Maple [A] (verified)	661
Fricas [A] (verification not implemented)	661
Sympy [A] (verification not implemented)	661
Maxima [B] (verification not implemented)	662
Giac [B] (verification not implemented)	662
Mupad [B] (verification not implemented)	663
Reduce [B] (verification not implemented)	663

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(x) \cosh(3x) dx = \frac{1}{4} \sinh(2x) + \frac{1}{8} \sinh(4x)$$

output `1/4*sinh(2*x)+1/8*sinh(4*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \cosh(3x) dx = \frac{1}{4} \sinh(2x) + \frac{1}{8} \sinh(4x)$$

input `Integrate[Cosh[x]*Cosh[3*x],x]`

output `Sinh[2*x]/4 + Sinh[4*x]/8`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4771}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \cosh(3x) dx$$

$$\downarrow \text{3042}$$

$$\int \cos(ix) \cos(3ix) dx$$

$$\downarrow \text{4771}$$

$$\frac{1}{4} \sinh(2x) + \frac{1}{8} \sinh(4x)$$

input `Int[Cosh[x]*Cosh[3*x],x]`

output `Sinh[2*x]/4 + Sinh[4*x]/8`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4771 `Int[cos[(a_.) + (b_.)*(x_)]*cos[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] + Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\sinh(2x)}{4} + \frac{\sinh(4x)}{8}$	14
parallelrisc	$\frac{\sinh(2x)}{4} + \frac{\sinh(4x)}{8}$	14
orering	$-\frac{\cosh(3x)\sinh(x)}{8} + \frac{3\cosh(x)\sinh(3x)}{8}$	18
risc	$\frac{e^{4x}}{16} + \frac{e^{2x}}{8} - \frac{e^{-2x}}{8} - \frac{e^{-4x}}{16}$	26

input `int(cosh(x)*cosh(3*x),x,method=_RETURNVERBOSE)`output `1/4*sinh(2*x)+1/8*sinh(4*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \cosh(3x) dx = \frac{1}{2} \cosh(x) \sinh(x)^3 + \frac{1}{2} (\cosh(x)^3 + \cosh(x)) \sinh(x)$$

input `integrate(cosh(x)*cosh(3*x),x, algorithm="fricas")`output `1/2*cosh(x)*sinh(x)^3 + 1/2*(cosh(x)^3 + cosh(x))*sinh(x)`**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \cosh(3x) dx = -\frac{\sinh(x) \cosh(3x)}{8} + \frac{3 \sinh(3x) \cosh(x)}{8}$$

input `integrate(cosh(x)*cosh(3*x),x)`

output `-sinh(x)*cosh(3*x)/8 + 3*sinh(3*x)*cosh(x)/8`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \cosh(3x) dx = \frac{1}{16} (2e^{-2x} + 1)e^{4x} - \frac{1}{8} e^{-2x} - \frac{1}{16} e^{-4x}$$

input `integrate(cosh(x)*cosh(3*x),x, algorithm="maxima")`

output `1/16*(2*e^(-2*x) + 1)*e^(4*x) - 1/8*e^(-2*x) - 1/16*e^(-4*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \cosh(3x) dx = -\frac{1}{16} (2e^{2x} + 1)e^{-4x} + \frac{1}{16} e^{4x} + \frac{1}{8} e^{2x}$$

input `integrate(cosh(x)*cosh(3*x),x, algorithm="giac")`

output `-1/16*(2*e^(2*x) + 1)*e^(-4*x) + 1/16*e^(4*x) + 1/8*e^(2*x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \cosh(3x) dx = \frac{e^{-4x} (e^{2x} - 1) (e^{2x} + 1)^3}{16}$$

input `int(cosh(3*x)*cosh(x),x)`output `(exp(-4*x)*(exp(2*x) - 1)*(exp(2*x) + 1)^3)/16`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \cosh(3x) dx = -\frac{\cosh(3x) \sinh(x)}{8} + \frac{3 \cosh(x) \sinh(3x)}{8}$$

input `int(cosh(x)*cosh(3*x),x)`output `(- cosh(3*x)*sinh(x) + 3*cosh(x)*sinh(3*x))/8`

3.93 $\int \cosh(x) \cosh(4x) dx$

Optimal result	664
Mathematica [A] (verified)	664
Rubi [A] (verified)	665
Maple [A] (verified)	666
Fricas [B] (verification not implemented)	666
Sympy [A] (verification not implemented)	667
Maxima [B] (verification not implemented)	667
Giac [B] (verification not implemented)	667
Mupad [B] (verification not implemented)	668
Reduce [B] (verification not implemented)	668

Optimal result

Integrand size = 7, antiderivative size = 17

$$\int \cosh(x) \cosh(4x) dx = \frac{1}{6} \sinh(3x) + \frac{1}{10} \sinh(5x)$$

output `1/6*sinh(3*x)+1/10*sinh(5*x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \cosh(4x) dx = \frac{1}{6} \sinh(3x) + \frac{1}{10} \sinh(5x)$$

input `Integrate[Cosh[x]*Cosh[4*x],x]`

output `Sinh[3*x]/6 + Sinh[5*x]/10`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4771}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \cosh(4x) dx$$

$$\downarrow 3042$$

$$\int \cos(ix) \cos(4ix) dx$$

$$\downarrow 4771$$

$$\frac{1}{6} \sinh(3x) + \frac{1}{10} \sinh(5x)$$

input `Int[Cosh[x]*Cosh[4*x],x]`

output `Sinh[3*x]/6 + Sinh[5*x]/10`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4771 `Int[cos[(a_.) + (b_.)*(x_)]*cos[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[a - c + (b - d)*x]/(2*(b - d)), x] + Simp[Sin[a + c + (b + d)*x]/(2*(b + d)), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{\sinh(3x)}{6} + \frac{\sinh(5x)}{10}$	14
parallelrisch	$\frac{\sinh(3x)}{6} + \frac{\sinh(5x)}{10}$	14
orering	$-\frac{\cosh(4x)\sinh(x)}{15} + \frac{4\cosh(x)\sinh(4x)}{15}$	18
risch	$\frac{e^{5x}}{20} + \frac{e^{3x}}{12} - \frac{e^{-3x}}{12} - \frac{e^{-5x}}{20}$	26

input `int(cosh(x)*cosh(4*x),x,method=_RETURNVERBOSE)`

output `1/6*sinh(3*x)+1/10*sinh(5*x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(13) = 26.

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

$$\int \cosh(x) \cosh(4x) dx = \frac{1}{10} \sinh(x)^5 + \frac{1}{6} (6 \cosh(x)^2 + 1) \sinh(x)^3 + \frac{1}{2} (\cosh(x)^4 + \cosh(x)^2) \sinh(x)$$

input `integrate(cosh(x)*cosh(4*x),x, algorithm="fricas")`

output `1/10*sinh(x)^5 + 1/6*(6*cosh(x)^2 + 1)*sinh(x)^3 + 1/2*(cosh(x)^4 + cosh(x)^2)*sinh(x)`

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.18

$$\int \cosh(x) \cosh(4x) dx = -\frac{\sinh(x) \cosh(4x)}{15} + \frac{4 \sinh(4x) \cosh(x)}{15}$$

input `integrate(cosh(x)*cosh(4*x),x)`

output `-sinh(x)*cosh(4*x)/15 + 4*sinh(4*x)*cosh(x)/15`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \cosh(4x) dx = \frac{1}{60} (5 e^{(-2x)} + 3) e^{(5x)} - \frac{1}{12} e^{(-3x)} - \frac{1}{20} e^{(-5x)}$$

input `integrate(cosh(x)*cosh(4*x),x, algorithm="maxima")`

output `1/60*(5*e^(-2*x) + 3)*e^(5*x) - 1/12*e^(-3*x) - 1/20*e^(-5*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \cosh(x) \cosh(4x) dx = -\frac{1}{60} (5 e^{(2x)} + 3) e^{(-5x)} + \frac{1}{20} e^{(5x)} + \frac{1}{12} e^{(3x)}$$

input `integrate(cosh(x)*cosh(4*x),x, algorithm="giac")`

output `-1/60*(5*e^(2*x) + 3)*e^(-5*x) + 1/20*e^(5*x) + 1/12*e^(3*x)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \cosh(x) \cosh(4x) dx = \frac{8 \sinh(x)^5}{5} + \frac{8 \sinh(x)^3}{3} + \sinh(x)$$

input `int(cosh(4*x)*cosh(x),x)`

output `sinh(x) + (8*sinh(x)^3)/3 + (8*sinh(x)^5)/5`

Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \cosh(x) \cosh(4x) dx = -\frac{\cosh(4x) \sinh(x)}{15} + \frac{4 \cosh(x) \sinh(4x)}{15}$$

input `int(cosh(x)*cosh(4*x),x)`

output `(- cosh(4*x)*sinh(x) + 4*cosh(x)*sinh(4*x))/15`

3.94 $\int \cosh(x) \cosh(mx) dx$

Optimal result	669
Mathematica [A] (verified)	669
Rubi [A] (verified)	670
Maple [A] (verified)	671
Fricas [A] (verification not implemented)	671
Sympy [B] (verification not implemented)	672
Maxima [F(-2)]	672
Giac [B] (verification not implemented)	673
Mupad [B] (verification not implemented)	673
Reduce [B] (verification not implemented)	673

Optimal result

Integrand size = 7, antiderivative size = 35

$$\int \cosh(x) \cosh(mx) dx = \frac{\sinh((1-m)x)}{2(1-m)} + \frac{\sinh((1+m)x)}{2(1+m)}$$

output

```
sinh((1-m)*x)/(2-2*m)+sinh((1+m)*x)/(2+2*m)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(x) \cosh(mx) dx = \frac{-\cosh(mx) \sinh(x) + m \cosh(x) \sinh(mx)}{-1 + m^2}$$

input

```
Integrate[Cosh[x]*Cosh[m*x],x]
```

output

```
(-(Cosh[m*x]*Sinh[x]) + m*Cosh[x]*Sinh[m*x])/(-1 + m^2)
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \cosh(mx) dx$$

$$\downarrow \text{6148}$$

$$\int \left(\frac{1}{2} \cosh((1-m)x) + \frac{1}{2} \cosh((m+1)x) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sinh((1-m)x)}{2(1-m)} + \frac{\sinh((m+1)x)}{2(m+1)}$$

input `Int[Cosh[x]*Cosh[m*x],x]`

output `Sinh[(1 - m)*x]/(2*(1 - m)) + Sinh[(1 + m)*x]/(2*(1 + m))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6148 `Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$\frac{\sinh(x(-1+m))}{-2+2m} + \frac{\sinh((1+m)x)}{2+2m}$
parallelrisc	$\frac{(1+m) \sinh(x(-1+m)) + \sinh((1+m)x)(-1+m)}{2m^2-2}$
risch	$\frac{(m e^{2x} - e^{2x} + m + 1) e^{x(-1+m)}}{4(1+m)(-1+m)} - \frac{(m e^{2x} + e^{2x} + m - 1) e^{-(1+m)x}}{4(1+m)(-1+m)}$
orering	$\frac{2(m^2+1)(\cosh(mx) \sinh(x) + \cosh(x) \sinh(mx)m)}{m^4-2m^2+1} - \frac{3m^2 \cosh(mx) \sinh(x) + 3 \cosh(x) \sinh(mx)m + \cosh(mx) \sinh(x) + m}{m^4-2m^2+1}$

input `int(cosh(x)*cosh(m*x), x, method=_RETURNVERBOSE)`output `1/2/(-1+m)*sinh(x*(-1+m))+1/2/(1+m)*sinh((1+m)*x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \cosh(x) \cosh(mx) dx = \frac{m \cosh(x) \sinh(mx) - \cosh(mx) \sinh(x)}{(m^2 - 1) \cosh(x)^2 - (m^2 - 1) \sinh(x)^2}$$

input `integrate(cosh(x)*cosh(m*x), x, algorithm="fricas")`output `(m*cosh(x)*sinh(m*x) - cosh(m*x)*sinh(x))/((m^2 - 1)*cosh(x)^2 - (m^2 - 1)*sinh(x)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(22) = 44$.

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.60

$$\int \cosh(x) \cosh(mx) dx = \begin{cases} -\frac{x \sinh^2(x)}{2} + \frac{x \cosh^2(x)}{2} + \frac{\sinh(x) \cosh(x)}{2} & \text{for } m = -1 \vee m = 1 \\ \frac{m \sinh(mx) \cosh(x)}{m^2 - 1} - \frac{\sinh(x) \cosh(mx)}{m^2 - 1} & \text{otherwise} \end{cases}$$

input `integrate(cosh(x)*cosh(m*x),x)`

output `Piecewise((-x*sinh(x)**2/2 + x*cosh(x)**2/2 + sinh(x)*cosh(x)/2, Eq(m, -1) | Eq(m, 1)), (m*sinh(m*x)*cosh(x)/(m**2 - 1) - sinh(x)*cosh(m*x)/(m**2 - 1), True))`

Maxima [F(-2)]

Exception generated.

$$\int \cosh(x) \cosh(mx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(x)*cosh(m*x),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(m-2>0)', see `assume?` for more details)Is`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \cosh(x) \cosh(mx) dx = \frac{e^{(mx+x)}}{4(m+1)} + \frac{e^{(mx-x)}}{4(m-1)} - \frac{e^{(-mx+x)}}{4(m-1)} - \frac{e^{(-mx-x)}}{4(m+1)}$$

input `integrate(cosh(x)*cosh(m*x),x, algorithm="giac")`

output `1/4*e^(m*x + x)/(m + 1) + 1/4*e^(m*x - x)/(m - 1) - 1/4*e^(-m*x + x)/(m - 1) - 1/4*e^(-m*x - x)/(m + 1)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \cosh(x) \cosh(mx) dx = -\frac{\cosh(mx) \sinh(x) - m \sinh(mx) \cosh(x)}{m^2 - 1}$$

input `int(cosh(m*x)*cosh(x),x)`

output `-(cosh(m*x)*sinh(x) - m*sinh(m*x)*cosh(x))/(m^2 - 1)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \cosh(x) \cosh(mx) dx = \frac{-\cosh(mx) \sinh(x) + \cosh(x) \sinh(mx) m}{m^2 - 1}$$

input `int(cosh(x)*cosh(m*x),x)`

output `(- cosh(m*x)*sinh(x) + cosh(x)*sinh(m*x)*m)/(m**2 - 1)`

3.95 $\int \cosh(a + bx) \cosh(c + bx) dx$

Optimal result	674
Mathematica [A] (verified)	674
Rubi [A] (verified)	675
Maple [A] (verified)	676
Fricas [B] (verification not implemented)	676
Sympy [B] (verification not implemented)	677
Maxima [B] (verification not implemented)	677
Giac [B] (verification not implemented)	678
Mupad [B] (verification not implemented)	678
Reduce [B] (verification not implemented)	678

Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \cosh(a + bx) \cosh(c + bx) dx = \frac{1}{2}x \cosh(a - c) + \frac{\sinh(a + c + 2bx)}{4b}$$

output `1/2*x*cosh(a-c)+1/4*sinh(2*b*x+a+c)/b`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cosh(a + bx) \cosh(c + bx) dx = \frac{2bx \cosh(a - c) + \sinh(a + c + 2bx)}{4b}$$

input `Integrate[Cosh[a + b*x]*Cosh[c + b*x],x]`

output `(2*b*x*Cosh[a - c] + Sinh[a + c + 2*b*x])/(4*b)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \cosh(bx + c) dx$$

$$\downarrow 6148$$

$$\int \left(\frac{1}{2} \cosh(a + 2bx + c) + \frac{1}{2} \cosh(a - c) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh(a + 2bx + c)}{4b} + \frac{1}{2} x \cosh(a - c)$$

input

```
Int[Cosh[a + b*x]*Cosh[c + b*x],x]
```

output

```
(x*Cosh[a - c])/2 + Sinh[a + c + 2*b*x]/(4*b)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6148

```
Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]
]^(p)*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result
default	$\frac{x \cosh(a-c)}{2} + \frac{\sinh(2bx+a+c)}{4b}$
parallelrisch	$\frac{2bx \cosh(a-c) + \sinh(2bx+a+c) - \sinh(a-c)}{4b}$
risch	$\frac{x e^{a-c}}{4} + \frac{x e^{-a+c}}{4} + \frac{e^{2bx+a+c}}{8b} - \frac{e^{-2bx-a-c}}{8b}$
orering	$x \cosh(bx+a) \cosh(bx+c) + \frac{b \cosh(bx+a) \sinh(bx+c) + \sinh(bx+a) b \cosh(bx+c)}{4b^2} - \frac{x(2 \sinh(bx+c) \sinh(bx+a))}{4b^2}$

input `int(cosh(b*x+a)*cosh(b*x+c),x,method=_RETURNVERBOSE)`

output `1/2*x*cosh(a-c)+1/4*sinh(2*b*x+a+c)/b`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 89 vs. 2(23) = 46.

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 3.30

$$\int \cosh(a+bx) \cosh(c+bx) dx$$

$$= \frac{2bx \cosh(-a+c) + 2 \cosh(bx+c) \cosh(-a+c) \sinh(bx+c) - \cosh(bx+c)^2 \sinh(-a+c) - \sinh(bx+c)^2 \sinh(-a+c)}{4(b \cosh(-a+c)^2 - b \sinh(-a+c)^2)}$$

input `integrate(cosh(b*x+a)*cosh(b*x+c),x, algorithm="fricas")`

output `1/4*(2*b*x*cosh(-a+c) + 2*cosh(b*x+c)*cosh(-a+c)*sinh(b*x+c) - cosh(b*x+c)^2*sinh(-a+c) - sinh(b*x+c)^2*sinh(-a+c))/(b*cosh(-a+c)^2 - b*sinh(-a+c)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(20) = 40$.

Time = 0.16 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \cosh(a + bx) \cosh(c + bx) dx = \begin{cases} -\frac{x \sinh(a+bx) \sinh(bx+c)}{2} + \frac{x \cosh(a+bx) \cosh(bx+c)}{2} + \frac{\sinh(bx+c) \cosh(a+bx)}{2b} & \text{for } b \neq 0 \\ x \cosh(a) \cosh(c) & \text{otherwise} \end{cases}$$

input `integrate(cosh(b*x+a)*cosh(b*x+c), x)`

output `Piecewise((-x*sinh(a + b*x)*sinh(b*x + c)/2 + x*cosh(a + b*x)*cosh(b*x + c)/2 + sinh(b*x + c)*cosh(a + b*x)/(2*b), Ne(b, 0)), (x*cosh(a)*cosh(c), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(23) = 46$.

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \cosh(a + bx) \cosh(c + bx) dx = \frac{(bx + a)(e^{(2a)} + e^{(2c)})e^{(-a-c)}}{4b} + \frac{e^{(2bx+a+c)}}{8b} - \frac{e^{(-2bx-a-c)}}{8b}$$

input `integrate(cosh(b*x+a)*cosh(b*x+c), x, algorithm="maxima")`

output `1/4*(b*x + a)*(e^(2*a) + e^(2*c))*e^(-a - c)/b + 1/8*e^(2*b*x + a + c)/b - 1/8*e^(-2*b*x - a - c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(23) = 46$.

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.44

$$\int \cosh(a + bx) \cosh(c + bx) dx$$

$$= \frac{(2bx(e^{2a} + e^{2c}) - (e^{2bx+2a} + e^{2bx+2c} + 1)e^{-2bx} + e^{(2bx+2a+2c)})e^{-a-c}}{8b}$$

input `integrate(cosh(b*x+a)*cosh(b*x+c),x, algorithm="giac")`

output `1/8*(2*b*x*(e^(2*a) + e^(2*c)) - (e^(2*b*x + 2*a) + e^(2*b*x + 2*c) + 1)*e^(-2*b*x) + e^(2*b*x + 2*a + 2*c))*e^(-a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \cosh(a + bx) \cosh(c + bx) dx = \frac{x \cosh(a - c)}{2} + \frac{\sinh(a + c + 2bx)}{4b}$$

input `int(cosh(a + b*x)*cosh(c + b*x),x)`

output `(x*cosh(a - c))/2 + sinh(a + c + 2*b*x)/(4*b)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.85

$$\int \cosh(a + bx) \cosh(c + bx) dx$$

$$= \frac{\cosh(bx + c) \cosh(bx + a) bx + \cosh(bx + c) \sinh(bx + a) - \sinh(bx + c) \sinh(bx + a) bx}{2b}$$

input `int(cosh(b*x+a)*cosh(b*x+c),x)`

output `(cosh(b*x + c)*cosh(a + b*x)*b*x + cosh(b*x + c)*sinh(a + b*x) - sinh(b*x + c)*sinh(a + b*x)*b*x)/(2*b)`

3.96 $\int \cosh(c - bx) \cosh(a + bx) dx$

Optimal result	680
Mathematica [A] (verified)	680
Rubi [A] (verified)	681
Maple [A] (verified)	682
Fricas [B] (verification not implemented)	682
Sympy [B] (verification not implemented)	683
Maxima [B] (verification not implemented)	683
Giac [B] (verification not implemented)	684
Mupad [B] (verification not implemented)	684
Reduce [B] (verification not implemented)	684

Optimal result

Integrand size = 14, antiderivative size = 27

$$\int \cosh(c - bx) \cosh(a + bx) dx = \frac{1}{2}x \cosh(a + c) + \frac{\sinh(a - c + 2bx)}{4b}$$

output `1/2*x*cosh(a+c)+1/4*sinh(2*b*x+a-c)/b`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cosh(c - bx) \cosh(a + bx) dx = \frac{2bx \cosh(a + c) + \sinh(a - c + 2bx)}{4b}$$

input `Integrate[Cosh[c - b*x]*Cosh[a + b*x],x]`

output `(2*b*x*Cosh[a + c] + Sinh[a - c + 2*b*x])/(4*b)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \cosh(c - bx) dx$$

$$\downarrow \text{6148}$$

$$\int \left(\frac{1}{2} \cosh(a + 2bx - c) + \frac{1}{2} \cosh(a + c) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sinh(a + 2bx - c)}{4b} + \frac{1}{2} x \cosh(a + c)$$

input

```
Int[Cosh[c - b*x]*Cosh[a + b*x],x]
```

output

```
(x*Cosh[a + c])/2 + Sinh[a - c + 2*b*x]/(4*b)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6148

```
Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]
]^(p)*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result
default	$\frac{x \cosh(a+c)}{2} + \frac{\sinh(2bx+a-c)}{4b}$
parallelrisc	$\frac{2bx \cosh(a+c) + \sinh(2bx+a-c) - \sinh(a+c)}{4b}$
risch	$\frac{x e^{a+c}}{4} + \frac{x e^{-a-c}}{4} + \frac{e^{2bx+a-c}}{8b} - \frac{e^{-2bx-a+c}}{8b}$
orering	$x \cosh(bx-c) \cosh(bx+a) + \frac{\sinh(bx-c)b \cosh(bx+a) + b \cosh(bx-c) \sinh(bx+a)}{4b^2} - \frac{x(2 \cosh(bx-c) \cosh(bx+a) - \sinh(bx-c) \sinh(bx+a))}{4b^2}$

input `int(cosh(b*x-c)*cosh(b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*x*cosh(a+c)+1/4*sinh(2*b*x+a-c)/b`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(23) = 46.

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.85

$$\int \cosh(c - bx) \cosh(a + bx) dx$$

$$= \frac{2bx \cosh(a+c) + 2 \cosh(bx+a) \cosh(a+c) \sinh(bx+a) - \cosh(bx+a)^2 \sinh(a+c) - \sinh(bx+a) \cosh(a+c)}{4(b \cosh(a+c)^2 - b \sinh(a+c)^2)}$$

input `integrate(cosh(b*x-c)*cosh(b*x+a),x, algorithm="fricas")`

output `1/4*(2*b*x*cosh(a+c) + 2*cosh(b*x+a)*cosh(a+c)*sinh(b*x+a) - cosh(b*x+a)^2*sinh(a+c) - sinh(b*x+a)^2*sinh(a+c))/(b*cosh(a+c)^2 - b*sinh(a+c)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(20) = 40$.

Time = 0.16 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \cosh(c - bx) \cosh(a + bx) dx = \begin{cases} -\frac{x \sinh(a+bx) \sinh(bx-c)}{2} + \frac{x \cosh(a+bx) \cosh(bx-c)}{2} + \frac{\sinh(bx-c) \cosh(a+bx)}{2b} & \text{for } b \neq 0 \\ x \cosh(a) \cosh(c) & \text{otherwise} \end{cases}$$

input `integrate(cosh(b*x-c)*cosh(b*x+a), x)`

output `Piecewise((-x*sinh(a + b*x)*sinh(b*x - c)/2 + x*cosh(a + b*x)*cosh(b*x - c)/2 + sinh(b*x - c)*cosh(a + b*x)/(2*b), Ne(b, 0)), (x*cosh(a)*cosh(c), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(23) = 46$.

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.19

$$\int \cosh(c - bx) \cosh(a + bx) dx = \frac{(bx + a)(e^{(2a+2c)} + 1)e^{(-a-c)}}{4b} + \frac{e^{(2bx+a-c)}}{8b} - \frac{e^{(-2bx-a+c)}}{8b}$$

input `integrate(cosh(b*x-c)*cosh(b*x+a), x, algorithm="maxima")`

output `1/4*(b*x + a)*(e^(2*a + 2*c) + 1)*e^(-a - c)/b + 1/8*e^(2*b*x + a - c)/b - 1/8*e^(-2*b*x - a + c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(23) = 46$.

Time = 0.18 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.44

$$\int \cosh(c - bx) \cosh(a + bx) dx$$

$$= \frac{(2bx(e^{2a+2c} + 1) - (e^{2bx} + e^{2bx+2a+2c} + e^{2c})e^{-2bx} + e^{(2bx+2a)})e^{-a-c}}{8b}$$

input `integrate(cosh(b*x-c)*cosh(b*x+a),x, algorithm="giac")`

output $\frac{1}{8} * (2 * b * x * (e^{(2 * a + 2 * c)} + 1) - (e^{(2 * b * x)} + e^{(2 * b * x + 2 * a + 2 * c)} + e^{(2 * c)}) * e^{(-2 * b * x)} + e^{(2 * b * x + 2 * a)}) * e^{(-a - c)} / b$

Mupad [B] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \cosh(c - bx) \cosh(a + bx) dx = \frac{\sinh(a - c + 2bx)}{4b} + \frac{x \cosh(a + c)}{2}$$

input `int(cosh(a + b*x)*cosh(c - b*x),x)`

output $\frac{\sinh(a - c + 2 * b * x)}{(4 * b)} + (x * \cosh(a + c)) / 2$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.07

$$\int \cosh(c - bx) \cosh(a + bx) dx$$

$$= \frac{\cosh(bx - c) \cosh(bx + a) bx + \cosh(bx - c) \sinh(bx + a) - \sinh(bx - c) \sinh(bx + a) bx}{2b}$$

input `int(cosh(b*x-c)*cosh(b*x+a),x)`

output $(\cosh(b*x - c) * \cosh(a + b*x) * b*x + \cosh(b*x - c) * \sinh(a + b*x) - \sinh(b*x - c) * \sinh(a + b*x) * b*x) / (2*b)$

3.97 $\int \cosh(a + bx) \cosh(c + dx) dx$

Optimal result	686
Mathematica [A] (verified)	686
Rubi [A] (verified)	687
Maple [A] (verified)	688
Fricas [A] (verification not implemented)	688
Sympy [B] (verification not implemented)	689
Maxima [F(-2)]	689
Giac [B] (verification not implemented)	690
Mupad [B] (verification not implemented)	690
Reduce [B] (verification not implemented)	691

Optimal result

Integrand size = 13, antiderivative size = 43

$$\int \cosh(a + bx) \cosh(c + dx) dx = \frac{\sinh(a - c + (b - d)x)}{2(b - d)} + \frac{\sinh(a + c + (b + d)x)}{2(b + d)}$$

output

```
sinh(a-c+(b-d)*x)/(2*b-2*d)+sinh(a+c+(b+d)*x)/(2*b+2*d)
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \cosh(a + bx) \cosh(c + dx) dx = \frac{\sinh(a - c + (b - d)x)}{2(b - d)} + \frac{\sinh(a + c + (b + d)x)}{2(b + d)}$$

input

```
Integrate[Cosh[a + b*x]*Cosh[c + d*x],x]
```

output

```
Sinh[a - c + (b - d)*x]/(2*(b - d)) + Sinh[a + c + (b + d)*x]/(2*(b + d))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \cosh(c + dx) dx$$

$$\downarrow 6148$$

$$\int \left(\frac{1}{2} \cosh(a + x(b - d) - c) + \frac{1}{2} \cosh(a + x(b + d) + c) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh(a + x(b - d) - c)}{2(b - d)} + \frac{\sinh(a + x(b + d) + c)}{2(b + d)}$$

input `Int[Cosh[a + b*x]*Cosh[c + d*x],x]`

output `Sinh[a - c + (b - d)*x]/(2*(b - d)) + Sinh[a + c + (b + d)*x]/(2*(b + d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6148 `Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

method	result
default	$\frac{\sinh(a-c+(b-d)x)}{2b-2d} + \frac{\sinh(a+c+(b+d)x)}{2b+2d}$
parallelrisc	$\frac{(b+d)\sinh(a-c+(b-d)x)+\sinh(a+c+(b+d)x)(b-d)}{2b^2-2d^2}$
risch	$\frac{(b e^{2bx+2a}-d e^{2bx+2a}-b-d)e^{-bx+dx-a+c}}{4(b+d)(b-d)} + \frac{(b e^{2bx+2a}+d e^{2bx+2a}-b+d)e^{-bx-dx-a-c}}{4(b+d)(b-d)}$
orering	$\frac{2(b^2+d^2)(b\sinh(bx+a)\cosh(dx+c)+\cosh(bx+a)d\sinh(dx+c))}{b^4-2b^2d^2+d^4} - \frac{b^3\sinh(bx+a)\cosh(dx+c)+3b^2\cosh(bx+a)d\sinh(dx+c)}{b^4-2b^2d^2+d^4}$

input `int(cosh(b*x+a)*cosh(d*x+c),x,method=_RETURNVERBOSE)`

output `1/2*sinh(a-c+(b-d)*x)/(b-d)+1/2/(b+d)*sinh(a+c+(b+d)*x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.65

$$\int \cosh(a+bx)\cosh(c+dx)dx$$

$$= \frac{b\cosh(dx+c)\sinh(bx+a)-d\cosh(bx+a)\sinh(dx+c)}{(b^2-d^2)\cosh(bx+a)^2-(b^2-d^2)\sinh(bx+a)^2}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c),x, algorithm="fricas")`

output `(b*cosh(d*x + c)*sinh(b*x + a) - d*cosh(b*x + a)*sinh(d*x + c))/((b^2 - d^2)*cosh(b*x + a)^2 - (b^2 - d^2)*sinh(b*x + a)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(32) = 64$.

Time = 0.31 (sec) , antiderivative size = 153, normalized size of antiderivative = 3.56

$$\int \cosh(a + bx) \cosh(c + dx) dx$$

$$= \begin{cases} x \cosh(a) \cosh(c) & \text{for } b = 0 \wedge d = 0 \\ \frac{x \sinh(a-dx) \sinh(c+dx)}{2} + \frac{x \cosh(a-dx) \cosh(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(a-dx)}{2d} & \text{for } b = -d \\ -\frac{x \sinh(a+dx) \sinh(c+dx)}{2} + \frac{x \cosh(a+dx) \cosh(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(a+dx)}{2d} & \text{for } b = d \\ \frac{b \sinh(a+bx) \cosh(c+dx)}{b^2-d^2} - \frac{d \sinh(c+dx) \cosh(a+bx)}{b^2-d^2} & \text{otherwise} \end{cases}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c), x)`

output `Piecewise((x*cosh(a)*cosh(c), Eq(b, 0) & Eq(d, 0)), (x*sinh(a - d*x)*sinh(c + d*x)/2 + x*cosh(a - d*x)*cosh(c + d*x)/2 + sinh(c + d*x)*cosh(a - d*x)/(2*d), Eq(b, -d)), (-x*sinh(a + d*x)*sinh(c + d*x)/2 + x*cosh(a + d*x)*cosh(c + d*x)/2 + sinh(c + d*x)*cosh(a + d*x)/(2*d), Eq(b, d)), (b*sinh(a + b*x)*cosh(c + d*x)/(b**2 - d**2) - d*sinh(c + d*x)*cosh(a + b*x)/(b**2 - d**2), True))`

Maxima [F(-2)]

Exception generated.

$$\int \cosh(a + bx) \cosh(c + dx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c), x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-d/b>0)', see `assume?` for more details)I`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(39) = 78.

Time = 0.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.98

$$\int \cosh(a + bx) \cosh(c + dx) dx = \frac{e^{(bx+dx+a+c)}}{4(b+d)} + \frac{e^{(bx-dx+a-c)}}{4(b-d)} - \frac{e^{(-bx+dx-a+c)}}{4(b-d)} - \frac{e^{(-bx-dx-a-c)}}{4(b+d)}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c),x, algorithm="giac")`

output `1/4*e^(b*x + d*x + a + c)/(b + d) + 1/4*e^(b*x - d*x + a - c)/(b - d) - 1/4*e^(-b*x + d*x - a + c)/(b - d) - 1/4*e^(-b*x - d*x - a - c)/(b + d)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \cosh(a + bx) \cosh(c + dx) dx = \frac{b \cosh(c + dx) \sinh(a + bx) - d \cosh(a + bx) \sinh(c + dx)}{b^2 - d^2}$$

input `int(cosh(a + b*x)*cosh(c + d*x),x)`

output `(b*cosh(c + d*x)*sinh(a + b*x) - d*cosh(a + b*x)*sinh(c + d*x))/(b^2 - d^2)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \cosh(a + bx) \cosh(c + dx) dx$$
$$= \frac{-\cosh(bx + a) \sinh(dx + c) d + \cosh(dx + c) \sinh(bx + a) b}{b^2 - d^2}$$

input `int(cosh(b*x+a)*cosh(d*x+c),x)`

output `(- cosh(a + b*x)*sinh(c + d*x)*d + cosh(c + d*x)*sinh(a + b*x)*b)/(b**2 - d**2)`

3.98 $\int \cosh(a + bx) \cosh^2(c + dx) dx$

Optimal result	692
Mathematica [A] (verified)	692
Rubi [A] (verified)	693
Maple [A] (verified)	694
Fricas [B] (verification not implemented)	694
Sympy [B] (verification not implemented)	695
Maxima [F(-2)]	696
Giac [B] (verification not implemented)	696
Mupad [B] (verification not implemented)	697
Reduce [B] (verification not implemented)	697

Optimal result

Integrand size = 15, antiderivative size = 62

$$\int \cosh(a + bx) \cosh^2(c + dx) dx = \frac{\sinh(a + bx)}{2b} + \frac{\sinh(a - 2c + (b - 2d)x)}{4(b - 2d)} + \frac{\sinh(a + 2c + (b + 2d)x)}{4(b + 2d)}$$

output

```
1/2*sinh(b*x+a)/b+sinh(a-2*c+(b-2*d)*x)/(4*b-8*d)+sinh(a+2*c+(b+2*d)*x)/(4*b+8*d)
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

$$\int \cosh(a + bx) \cosh^2(c + dx) dx = \frac{1}{4} \left(\frac{2 \cosh(bx) \sinh(a)}{b} + \frac{2 \cosh(a) \sinh(bx)}{b} + \frac{\sinh(a - 2c + bx - 2dx)}{b - 2d} + \frac{\sinh(a + 2c + bx + 2dx)}{b + 2d} \right)$$

input

```
Integrate[Cosh[a + b*x]*Cosh[c + d*x]^2,x]
```

output

$$\frac{((2*\text{Cosh}[b*x]*\text{Sinh}[a])/b + (2*\text{Cosh}[a]*\text{Sinh}[b*x])/b + \text{Sinh}[a - 2*c + b*x - 2*d*x]/(b - 2*d) + \text{Sinh}[a + 2*c + b*x + 2*d*x]/(b + 2*d))/4}$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \cosh^2(c + dx) dx$$

$$\downarrow 6148$$

$$\int \left(\frac{1}{4} \cosh(a + x(b - 2d) - 2c) + \frac{1}{4} \cosh(a + x(b + 2d) + 2c) + \frac{1}{2} \cosh(a + bx) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh(a + x(b - 2d) - 2c)}{4(b - 2d)} + \frac{\sinh(a + x(b + 2d) + 2c)}{4(b + 2d)} + \frac{\sinh(a + bx)}{2b}$$

input

```
Int[Cosh[a + b*x]*Cosh[c + d*x]^2,x]
```

output

```
Sinh[a + b*x]/(2*b) + Sinh[a - 2*c + (b - 2*d)*x]/(4*(b - 2*d)) + Sinh[a + 2*c + (b + 2*d)*x]/(4*(b + 2*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6148

```
Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] :> Int[ExpandTrigReduce[Cosh[v]
]^(p)*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

method	result
default	$\frac{\sinh(bx+a)}{2b} + \frac{\sinh(a-2c+(b-2d)x)}{4b-8d} + \frac{\sinh(a+2c+(b+2d)x)}{4b+8d}$
parallelrisch	$\frac{b(b+2d) \sinh(a-2c+(b-2d)x) + 2 \left(\frac{b \sinh(a+2c+(b+2d)x)}{2} + \sinh(bx+a)(b+2d) \right) (b-2d)}{4b^3 - 16bd^2}$
risch	$\frac{e^{bx+a}}{4b} - \frac{e^{-bx-a}}{4b} + \frac{(be^{2bx+2a} - 2de^{2bx+2a} - b - 2d)e^{-bx+2dx-a+2c}}{8(b+2d)(b-2d)} + \frac{(be^{2bx+2a} + 2de^{2bx+2a} - b + 2d)e^{-bx-2dx-a-2c}}{8(b+2d)(b-2d)}$
orering	$\frac{(3b^4+16d^4) (b \sinh(bx+a) \cosh(dx+c)^2 + 2 \cosh(bx+a) \cosh(dx+c) d \sinh(dx+c))}{b^2(b^4 - 8b^2d^2 + 16d^4)} - \frac{(3b^2+8d^2) (b^3 \sinh(bx+a) \cosh(dx+c) + b^2 \cosh(bx+a) \sinh(dx+c))}{b^2(b^4 - 8b^2d^2 + 16d^4)}$

input

```
int(cosh(b*x+a)*cosh(d*x+c)^2,x,method=_RETURNVERBOSE)
```

output

```
1/2*sinh(b*x+a)/b+1/4*sinh(a-2*c+(b-2*d)*x)/(b-2*d)+1/4*sinh(a+2*c+(b+2*d)
*x)/(b+2*d)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(56) = 112.

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.85

$$\int \cosh(a + bx) \cosh^2(c + dx) dx = \frac{4bd \cosh(bx + a) \cosh(dx + c) \sinh(dx + c) - b^2 \sinh(bx + a) \sinh(dx + c)^2 - (b^2 \cosh(dx + c)^2 + b^2 \sinh(dx + c)^2)}{2((b^3 - 4bd^2) \cosh(bx + a)^2 - (b^3 - 4bd^2) \sinh(bx + a)^2)}$$

input

```
integrate(cosh(b*x+a)*cosh(d*x+c)^2,x, algorithm="fricas")
```

output

$$-1/2*(4*b*d*cosh(b*x + a)*cosh(d*x + c)*sinh(d*x + c) - b^2*sinh(b*x + a)*sinh(d*x + c)^2 - (b^2*cosh(d*x + c)^2 + b^2 - 4*d^2)*sinh(b*x + a))/((b^3 - 4*b*d^2)*cosh(b*x + a)^2 - (b^3 - 4*b*d^2)*sinh(b*x + a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(49) = 98$.

Time = 0.70 (sec) , antiderivative size = 408, normalized size of antiderivative = 6.58

$$\int \cosh(a + bx) \cosh^2(c + dx) dx$$

$$= \begin{cases} x \cosh(a) \cosh^2(c) \\ \left(-\frac{x \sinh^2(c+dx)}{2} + \frac{x \cosh^2(c+dx)}{2} + \frac{\sinh(c+dx) \cosh(c+dx)}{2d} \right) \cosh(a) \\ \frac{x \sinh(a-2dx) \sinh(c+dx) \cosh(c+dx)}{2} + \frac{x \sinh^2(c+dx) \cosh(a-2dx)}{4} + \frac{x \cosh(a-2dx) \cosh^2(c+dx)}{4} + \frac{\sinh(a-2dx) \sinh^2(c+dx)}{2d} \\ - \frac{x \sinh(a+2dx) \sinh(c+dx) \cosh(c+dx)}{2} + \frac{x \sinh^2(c+dx) \cosh(a+2dx)}{4} + \frac{x \cosh(a+2dx) \cosh^2(c+dx)}{4} - \frac{\sinh(a+2dx) \sinh^2(c+dx)}{2d} \\ \frac{b^2 \sinh(a+bx) \cosh^2(c+dx)}{b^3-4bd^2} - \frac{2bd \sinh(c+dx) \cosh(a+bx) \cosh(c+dx)}{b^3-4bd^2} + \frac{2d^2 \sinh(a+bx) \sinh^2(c+dx)}{b^3-4bd^2} - \frac{2d^2 \sinh(a+bx) \cosh^2(c+dx)}{b^3-4bd^2} \end{cases}$$

input

```
integrate(cosh(b*x+a)*cosh(d*x+c)**2,x)
```

output

```
Piecewise((x*cosh(a)*cosh(c)**2, Eq(b, 0) & Eq(d, 0)), ((-x*sinh(c + d*x)*
**2/2 + x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*cosh(a),
Eq(b, 0)), (x*sinh(a - 2*d*x)*sinh(c + d*x)*cosh(c + d*x)/2 + x*sinh(c + d
*x)**2*cosh(a - 2*d*x)/4 + x*cosh(a - 2*d*x)*cosh(c + d*x)**2/4 + sinh(a -
2*d*x)*sinh(c + d*x)**2/(2*d) + 3*sinh(c + d*x)*cosh(a - 2*d*x)*cosh(c +
d*x)/(4*d), Eq(b, -2*d)), (-x*sinh(a + 2*d*x)*sinh(c + d*x)*cosh(c + d*x)/
2 + x*sinh(c + d*x)**2*cosh(a + 2*d*x)/4 + x*cosh(a + 2*d*x)*cosh(c + d*x)
**2/4 - sinh(a + 2*d*x)*sinh(c + d*x)**2/(2*d) + 3*sinh(c + d*x)*cosh(a +
2*d*x)*cosh(c + d*x)/(4*d), Eq(b, 2*d)), (b**2*sinh(a + b*x)*cosh(c + d*x)
**2/(b**3 - 4*b*d**2) - 2*b*d*sinh(c + d*x)*cosh(a + b*x)*cosh(c + d*x)/(b
**3 - 4*b*d**2) + 2*d**2*sinh(a + b*x)*sinh(c + d*x)**2/(b**3 - 4*b*d**2)
- 2*d**2*sinh(a + b*x)*cosh(c + d*x)**2/(b**3 - 4*b*d**2), True))
```


Maxima [F(-2)]

Exception generated.

$$\int \cosh(a + bx) \cosh^2(c + dx) dx = \text{Exception raised: ValueError}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c)^2,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(2*d)/b>0)', see `assume?` for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 0.14 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.94

$$\int \cosh(a + bx) \cosh^2(c + dx) dx = \frac{e^{(bx+2dx+a+2c)}}{8(b+2d)} + \frac{e^{(bx-2dx+a-2c)}}{8(b-2d)} + \frac{e^{(bx+a)}}{4b} - \frac{e^{(-bx+2dx-a+2c)}}{8(b-2d)} - \frac{e^{(-bx-2dx-a-2c)}}{8(b+2d)} - \frac{e^{(-bx-a)}}{4b}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c)^2,x, algorithm="giac")`

output `1/8*e^(b*x + 2*d*x + a + 2*c)/(b + 2*d) + 1/8*e^(b*x - 2*d*x + a - 2*c)/(b - 2*d) + 1/4*e^(b*x + a)/b - 1/8*e^(-b*x + 2*d*x - a + 2*c)/(b - 2*d) - 1/8*e^(-b*x - 2*d*x - a - 2*c)/(b + 2*d) - 1/4*e^(-b*x - a)/b`

Mupad [B] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.10

$$\int \cosh(a + bx) \cosh^2(c + dx) dx$$

$$= \frac{2d^2 \sinh(a + bx) - b^2 \cosh(c + dx)^2 \sinh(a + bx) + 2bd \cosh(a + bx) \cosh(c + dx) \sinh(c + dx)}{4bd^2 - b^3}$$

input `int(cosh(a + b*x)*cosh(c + d*x)^2,x)`output `(2*d^2*sinh(a + b*x) - b^2*cosh(c + d*x)^2*sinh(a + b*x) + 2*b*d*cosh(a + b*x)*cosh(c + d*x)*sinh(c + d*x))/(4*b*d^2 - b^3)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 215, normalized size of antiderivative = 3.47

$$\int \cosh(a + bx) \cosh^2(c + dx) dx$$

$$= \frac{e^{2bx+4dx+2a+4c}b^2 - 2e^{2bx+4dx+2a+4c}bd + 2e^{2bx+2dx+2a+2c}b^2 - 8e^{2bx+2dx+2a+2c}d^2 + e^{2bx+2a}b^2 + 2e^{2bx+2a}bd - 8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}{8e^{bx+2dx+a+2c}b(b^2 - 4d^2)}$$

input `int(cosh(b*x+a)*cosh(d*x+c)^2,x)`output `(e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2 - 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d + 2*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2 - 8*e**(2*a + 2*b*x + 2*c + 2*d*x)*d**2 + e**(2*a + 2*b*x)*b**2 + 2*e**(2*a + 2*b*x)*b*d - e**(4*c + 4*d*x)*b**2 - 2*e**(4*c + 4*d*x)*b*d - 2*e**(2*c + 2*d*x)*b**2 + 8*e**(2*c + 2*d*x)*d**2 - b**2 + 2*b*d)/(8*e**(a + b*x + 2*c + 2*d*x)*b*(b**2 - 4*d**2))`

3.99 $\int \cosh(a + bx) \cosh^3(c + dx) dx$

Optimal result	698
Mathematica [A] (verified)	698
Rubi [A] (verified)	699
Maple [A] (verified)	700
Fricas [B] (verification not implemented)	700
Sympy [B] (verification not implemented)	701
Maxima [F(-2)]	702
Giac [B] (verification not implemented)	703
Mupad [B] (verification not implemented)	703
Reduce [B] (verification not implemented)	704

Optimal result

Integrand size = 15, antiderivative size = 91

$$\int \cosh(a + bx) \cosh^3(c + dx) dx = \frac{\sinh(a - 3c + (b - 3d)x)}{8(b - 3d)} + \frac{3 \sinh(a - c + (b - d)x)}{8(b - d)} + \frac{3 \sinh(a + c + (b + d)x)}{8(b + d)} + \frac{\sinh(a + 3c + (b + 3d)x)}{8(b + 3d)}$$

output

```
sinh(a-3*c+(b-3*d)*x)/(8*b-24*d)+3*sinh(a-c+(b-d)*x)/(8*b-8*d)+3*sinh(a+c+(b+d)*x)/(8*b+8*d)+sinh(a+3*c+(b+3*d)*x)/(8*b+24*d)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.93

$$\int \cosh(a + bx) \cosh^3(c + dx) dx = \frac{1}{8} \left(\frac{\sinh(a - 3c + bx - 3dx)}{b - 3d} + \frac{3 \sinh(a - c + bx - dx)}{b - d} + \frac{\sinh(a + 3c + bx + 3dx)}{b + 3d} + \frac{3 \sinh(a + c + (b + d)x)}{b + d} \right)$$

input `Integrate[Cosh[a + b*x]*Cosh[c + d*x]^3,x]`

output $(\text{Sinh}[a - 3c + b*x - 3d*x]/(b - 3d) + (3*\text{Sinh}[a - c + b*x - d*x])/(b - d) + \text{Sinh}[a + 3c + b*x + 3d*x]/(b + 3d) + (3*\text{Sinh}[a + c + (b + d)*x])/(b + d))/8$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \cosh^3(c + dx) dx$$

$$\downarrow 6148$$

$$\int \left(\frac{1}{8} \cosh(a + x(b - 3d) - 3c) + \frac{3}{8} \cosh(a + x(b - d) - c) + \frac{3}{8} \cosh(a + x(b + d) + c) + \frac{1}{8} \cosh(a + x(b + 3d) + 3c) \right) dx$$

$$\downarrow 2009$$

$$\frac{\sinh(a + x(b - 3d) - 3c)}{8(b - 3d)} + \frac{3 \sinh(a + x(b - d) - c)}{8(b - d)} + \frac{3 \sinh(a + x(b + d) + c)}{8(b + d)} + \frac{\sinh(a + x(b + 3d) + 3c)}{8(b + 3d)}$$

input `Int[Cosh[a + b*x]*Cosh[c + d*x]^3,x]`

output $\text{Sinh}[a - 3c + (b - 3d)*x]/(8*(b - 3d)) + (3*\text{Sinh}[a - c + (b - d)*x])/(8*(b - d)) + (3*\text{Sinh}[a + c + (b + d)*x])/(8*(b + d)) + \text{Sinh}[a + 3c + (b + 3d)*x]/(8*(b + 3d))$

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6148 Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]
]^(p)*Cosh[w]^q, x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 2.88 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.92

method	result
default	$\frac{\sinh(a-3c+(b-3d)x)}{8b-24d} + \frac{3 \sinh(a-c+(b-d)x)}{8(b-d)} + \frac{3 \sinh(a+c+(b+d)x)}{8(b+d)} + \frac{\sinh(a+3c+(b+3d)x)}{8b+24d}$
risch	$\frac{(b e^{2bx+2a} - 3d e^{2bx+2a-b-3d}) e^{-bx+3dx-a+3c}}{16(b+3d)(b-3d)} + \frac{3(b e^{2bx+2a-d} e^{2bx+2a-b-d}) e^{-bx+dx-a+c}}{16(b+d)(b-d)} + \frac{3(b e^{2bx+2a+d} e^{2bx+2a-b+d}) e^{-bx+dx-a+c}}{16(b+d)(b-d)}$
parallelrisch	$\frac{2b \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) (b^2 - 7d^2) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^6 - 6d \left(1 + \tanh\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^2 (b^2 - 3d^2) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^5 + 6b \tanh\left(\frac{bx}{2} + \frac{a}{2}\right) (b^2 + d^2) \tanh\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{(b-d)(b+d)(b+3d)(b-3d)}$
orering	Expression too large to display

```
input int(cosh(b*x+a)*cosh(d*x+c)^3,x,method=_RETURNVERBOSE)
```

```
output 1/8*sinh(a-3*c+(b-3*d)*x)/(b-3*d)+3/8*sinh(a-c+(b-d)*x)/(b-d)+3/8/(b+d)*si
nh(a+c+(b+d)*x)+1/8/(b+3*d)*sinh(a+3*c+(b+3*d)*x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 217 vs. 2(83) = 166.

Time = 0.08 (sec) , antiderivative size = 217, normalized size of antiderivative = 2.38

$$\int \cosh(a + bx) \cosh^3(c + dx) dx$$

$$= \frac{3(b^3 - bd^2) \cosh(dx + c) \sinh(bx + a) \sinh(dx + c)^2 - 3(b^2d - d^3) \cosh(bx + a) \sinh(dx + c)^3 + ((b^3 - 3bd^2) \cosh^2(dx + c) \sinh(bx + a) \sinh(dx + c) - (b^2d - d^3) \cosh^2(dx + c) \sinh^2(dx + c) + (b^3 - 3bd^2) \cosh(dx + c) \sinh^3(dx + c) - (b^2d - d^3) \sinh^3(dx + c))}{4((b^4 - 10b^2d^2 + 5d^4) \cosh^2(dx + c) \sinh^2(dx + c) + (b^3 - 3bd^2) \cosh(dx + c) \sinh^3(dx + c) - (b^2d - d^3) \sinh^3(dx + c))}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c)^3,x, algorithm="fricas")`

output
$$\frac{1}{4} \cdot (3 \cdot (b^3 - b \cdot d^2) \cdot \cosh(d \cdot x + c) \cdot \sinh(b \cdot x + a) \cdot \sinh(d \cdot x + c)^2 - 3 \cdot (b^2 \cdot d - d^3) \cdot \cosh(b \cdot x + a) \cdot \sinh(d \cdot x + c)^3 + ((b^3 - b \cdot d^2) \cdot \cosh(d \cdot x + c)^3 + 3 \cdot (b^3 - 9 \cdot b \cdot d^2) \cdot \cosh(d \cdot x + c)) \cdot \sinh(b \cdot x + a) - 3 \cdot (3 \cdot (b^2 \cdot d - d^3) \cdot \cosh(b \cdot x + a) \cdot \cosh(d \cdot x + c)^2 + (b^2 \cdot d - 9 \cdot d^3) \cdot \cosh(b \cdot x + a)) \cdot \sinh(d \cdot x + c)) / ((b^4 - 10 \cdot b^2 \cdot d^2 + 9 \cdot d^4) \cdot \cosh(b \cdot x + a)^2 - (b^4 - 10 \cdot b^2 \cdot d^2 + 9 \cdot d^4) \cdot \sinh(b \cdot x + a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 921 vs. $2(76) = 152$.

Time = 1.91 (sec) , antiderivative size = 921, normalized size of antiderivative = 10.12

$$\int \cosh(a + bx) \cosh^3(c + dx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c)**3,x)`

output

```
Piecewise((x*cosh(a)*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (x*sinh(a - 3*d*x)*
sinh(c + d*x)**3/8 + 3*x*sinh(a - 3*d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8
+ 3*x*sinh(c + d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)/8 + x*cosh(a - 3*d*x)
*cosh(c + d*x)**3/8 + sinh(a - 3*d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d)
- 7*sinh(a - 3*d*x)*cosh(c + d*x)**3/(24*d) + sinh(c + d*x)**3*cosh(a - 3
*d*x)/(8*d), Eq(b, -3*d)), (-3*x*sinh(a - d*x)*sinh(c + d*x)**3/8 + 3*x*si
nh(a - d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8 - 3*x*sinh(c + d*x)**2*cosh(a
- d*x)*cosh(c + d*x)/8 + 3*x*cosh(a - d*x)*cosh(c + d*x)**3/8 + 3*sinh(a
- d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d) - 5*sinh(a - d*x)*cosh(c + d*x
)**3/(8*d) + 3*sinh(c + d*x)**3*cosh(a - d*x)/(8*d), Eq(b, -d)), (3*x*sinh
(a + d*x)*sinh(c + d*x)**3/8 - 3*x*sinh(a + d*x)*sinh(c + d*x)*cosh(c + d*
x)**2/8 - 3*x*sinh(c + d*x)**2*cosh(a + d*x)*cosh(c + d*x)/8 + 3*x*cosh(a
+ d*x)*cosh(c + d*x)**3/8 - 3*sinh(a + d*x)*sinh(c + d*x)**2*cosh(c + d*x)
/(4*d) + 5*sinh(a + d*x)*cosh(c + d*x)**3/(8*d) + 3*sinh(c + d*x)**3*cosh(
a + d*x)/(8*d), Eq(b, d)), (-x*sinh(a + 3*d*x)*sinh(c + d*x)**3/8 - 3*x*si
nh(a + 3*d*x)*sinh(c + d*x)*cosh(c + d*x)**2/8 + 3*x*sinh(c + d*x)**2*cosh
(a + 3*d*x)*cosh(c + d*x)/8 + x*cosh(a + 3*d*x)*cosh(c + d*x)**3/8 - sinh(
a + 3*d*x)*sinh(c + d*x)**2*cosh(c + d*x)/(4*d) + 7*sinh(a + 3*d*x)*cosh(
c + d*x)**3/(24*d) + sinh(c + d*x)**3*cosh(a + 3*d*x)/(8*d), Eq(b, 3*d)), (
b**3*sinh(a + b*x)*cosh(c + d*x)**3/(b**4 - 10*b**2*d**2 + 9*d**4) - 3*...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh(a + bx) \cosh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(b*x+a)*cosh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for
more detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(83) = 166$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.97

$$\int \cosh(a + bx) \cosh^3(c + dx) dx = \frac{e^{(bx+3dx+a+3c)}}{16(b+3d)} + \frac{3e^{(bx+dx+a+c)}}{16(b+d)} + \frac{3e^{(bx-dx+a-c)}}{16(b-d)} \\ + \frac{e^{(bx-3dx+a-3c)}}{16(b-3d)} - \frac{e^{(-bx+3dx-a+3c)}}{16(b-3d)} - \frac{3e^{(-bx+dx-a+c)}}{16(b-d)} \\ - \frac{3e^{(-bx-dx-a-c)}}{16(b+d)} - \frac{e^{(-bx-3dx-a-3c)}}{16(b+3d)}$$

input `integrate(cosh(b*x+a)*cosh(d*x+c)^3,x, algorithm="giac")`

output `1/16*e^(b*x + 3*d*x + a + 3*c)/(b + 3*d) + 3/16*e^(b*x + d*x + a + c)/(b + d) + 3/16*e^(b*x - d*x + a - c)/(b - d) + 1/16*e^(b*x - 3*d*x + a - 3*c)/(b - 3*d) - 1/16*e^(-b*x + 3*d*x - a + 3*c)/(b - 3*d) - 3/16*e^(-b*x + d*x - a + c)/(b - d) - 3/16*e^(-b*x - d*x - a - c)/(b + d) - 1/16*e^(-b*x - 3*d*x - a - 3*c)/(b + 3*d)`

Mupad [B] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.98

$$\int \cosh(a + bx) \cosh^3(c + dx) dx \\ = \frac{b \cosh(c + dx)^3 \sinh(a + bx) (b^2 - 7d^2)}{b^4 - 10b^2d^2 + 9d^4} \\ - \frac{3 \cosh(a + bx) \cosh(c + dx)^2 \sinh(c + dx) (b^2d - 3d^3)}{b^4 - 10b^2d^2 + 9d^4} \\ - \frac{6d^3 \cosh(a + bx) \sinh(c + dx)^3}{b^4 - 10b^2d^2 + 9d^4} + \frac{6bd^2 \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)^2}{b^4 - 10b^2d^2 + 9d^4}$$

input `int(cosh(a + b*x)*cosh(c + d*x)^3,x)`

output

$$\begin{aligned} & (b \cosh(c + dx))^3 \sinh(a + bx) (b^2 - 7d^2) / (b^4 + 9d^4 - 10b^2d^2) \\ & - (3 \cosh(a + bx) \cosh(c + dx))^2 \sinh(c + dx) (b^2d - 3d^3) / (b^4 + \\ & 9d^4 - 10b^2d^2) - (6d^3 \cosh(a + bx) \sinh(c + dx)^3) / (b^4 + 9d^4 - \\ & 10b^2d^2) + (6bd^2 \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)^2) / (b^4 \\ & + 9d^4 - 10b^2d^2) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 574, normalized size of antiderivative = 6.31

$$\int \cosh(a + bx) \cosh^3(c + dx) dx$$

$$= \frac{-b^3 - 3d^3 + bd^2 + e^{6dx+6c}bd^2 - e^{2bx+2a}bd^2 - e^{2bx+6dx+2a+6c}bd^2 + 27e^{2bx+4dx+2a+4c}d^3 + 3e^{2bx+2dx+2a+2c}bd^2}{(b^4 + 9d^4 - 10b^2d^2)}$$

input

int(cosh(b*x+a)*cosh(d*x+c)^3,x)

output

$$\begin{aligned} & (e^{2a + 2bx + 6c + 6dx})b^3 - 3e^{2a + 2bx + 6c + 6dx})b^2d \\ & - e^{2a + 2bx + 6c + 6dx})bd^2 + 3e^{2a + 2bx + 6c + 6dx})d^3 \\ & + 3e^{2a + 2bx + 4c + 4dx})b^3 - 3e^{2a + 2bx + 4c + 4dx})b^2d \\ & - 27e^{2a + 2bx + 4c + 4dx})bd^2 + 27e^{2a + 2bx + 4c + 4dx})d^3 \\ & + 3e^{2a + 2bx + 2c + 2dx})b^3 + 3e^{2a + 2bx + 2c + 2dx})b^2d \\ & - 27e^{2a + 2bx + 2c + 2dx})bd^2 - 27e^{2a + 2bx + 2c + 2dx})d^3 \\ & + e^{2a + 2bx})b^3 + 3e^{2a + 2bx})b^2d - e^{2a + 2bx})bd^2 - 3e^{2a + 2bx})d^3 \\ & - e^{6c + 6dx})b^3 - 3e^{6c + 6dx})b^2d + e^{6c + 6dx})bd^2 \\ & + 3e^{6c + 6dx})d^3 - 3e^{4c + 4dx})b^3 - 3e^{4c + 4dx})b^2d \\ & + 27e^{4c + 4dx})bd^2 + 27e^{4c + 4dx})d^3 - 3e^{2c + 2dx})b^3 \\ & + 3e^{2c + 2dx})b^2d + 27e^{2c + 2dx})bd^2 - 27e^{2c + 2dx})d^3 \\ & - b^3 + 3b^2d + bd^2 - 3d^3) / (16e^{2a + 2bx + 6c + 6dx}) \\ & (b^4 - 10b^2d^2 + 9d^4) \end{aligned}$$

3.100 $\int \cosh^2(a + bx) \cosh^2(c + dx) dx$

Optimal result	705
Mathematica [A] (verified)	705
Rubi [A] (verified)	706
Maple [A] (verified)	707
Fricas [B] (verification not implemented)	707
Sympy [B] (verification not implemented)	708
Maxima [F(-2)]	709
Giac [A] (verification not implemented)	710
Mupad [B] (verification not implemented)	710
Reduce [B] (verification not implemented)	711

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \frac{x}{4} + \frac{\sinh(2a + 2bx)}{8b} + \frac{\sinh(2(a - c) + 2(b - d)x)}{16(b - d)} + \frac{\sinh(2c + 2dx)}{8d} + \frac{\sinh(2(a + c) + 2(b + d)x)}{16(b + d)}$$

output

```
1/4*x+1/8*sinh(2*b*x+2*a)/b+sinh(2*a-2*c+2*(b-d)*x)/(16*b-16*d)+1/8*sinh(2*d*x+2*c)/d+sinh(2*a+2*c+2*(b+d)*x)/(16*b+16*d)
```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.19

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \frac{2d(b^2 - d^2) \sinh(2(a + bx)) + bd(b + d) \sinh(2(a - c + (b - d)x)) + b(b - d)(2(b + d) \sinh(2(c + dx)))}{16b(b - d)d(b + d)}$$

input

```
Integrate[Cosh[a + b*x]^2*Cosh[c + d*x]^2,x]
```

output

```
(2*d*(b^2 - d^2)*Sinh[2*(a + b*x)] + b*d*(b + d)*Sinh[2*(a - c + (b - d)*x)
] + b*(b - d)*(2*(b + d)*Sinh[2*(c + d*x)] + d*(4*(b + d)*x + Sinh[2*(a +
c + (b + d)*x]])))/(16*b*(b - d)*d*(b + d))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx$$

↓ 6148

$$\int \left(\frac{1}{8} \cosh(2(a - c) + 2x(b - d)) + \frac{1}{8} \cosh(2(a + c) + 2x(b + d)) + \frac{1}{4} \cosh(2a + 2bx) + \frac{1}{4} \cosh(2c + 2dx) + \frac{1}{4} \right)$$

↓ 2009

$$\frac{\sinh(2(a - c) + 2x(b - d))}{16(b - d)} + \frac{\sinh(2(a + c) + 2x(b + d))}{16(b + d)} + \frac{\sinh(2a + 2bx)}{8b} + \frac{\sinh(2c + 2dx)}{8d} + \frac{x}{4}$$

input

```
Int[Cosh[a + b*x]^2*Cosh[c + d*x]^2,x]
```

output

```
x/4 + Sinh[2*a + 2*b*x]/(8*b) + Sinh[2*(a - c) + 2*(b - d)*x]/(16*(b - d))
+ Sinh[2*c + 2*d*x]/(8*d) + Sinh[2*(a + c) + 2*(b + d)*x]/(16*(b + d))
```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6148 Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]
]^(p)*Cosh[w]^(q), x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] &&
PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x
]))
```

Maple [A] (verified)

Time = 3.65 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

method	result
default	$\frac{x}{4} + \frac{\sinh(2bx+2a)}{8b} + \frac{\sinh(2dx+2c)}{8d} + \frac{\sinh((2b-2d)x+2a-2c)}{16b-16d} + \frac{\sinh((2b+2d)x+2a+2c)}{16b+16d}$
parallelrisch	$\frac{bd(b+d) \sinh((2b-2d)x+2a-2c)+4\left(\frac{bd \sinh((2b+2d)x+2a+2c)}{4} + \left(\frac{d \sinh(2bx+2a)}{2} + b\left(dx + \frac{\sinh(2dx+2c)}{2}\right)\right)(b+d)\right)(b-d)}{16b^3d-16bd^3}$
risch	$\frac{x}{4} + \frac{e^{2bx+2a}}{16b} - \frac{e^{-2bx-2a}}{16b} + \frac{(de^{4bx+4a}b-d^2e^{4bx+4a}+2b^2e^{2bx+2a}-2d^2e^{2bx+2a}-bd-d^2)e^{-2bx+2dx-2a+2c}}{32(b+d)(b-d)d} - \frac{(-de^{4bx+4a}b-d^2e^{4bx+4a}+2b^2e^{2bx+2a}-2d^2e^{2bx+2a}-bd-d^2)e^{-2bx-2a+2c}}{32(b+d)(b-d)d}$
orering	Expression too large to display

```
input int(cosh(b*x+a)^2*cosh(d*x+c)^2,x,method=_RETURNVERBOSE)
```

```
output 1/4*x+1/8*sinh(2*b*x+2*a)/b+1/8*sinh(2*d*x+2*c)/d+1/8/(2*b-2*d)*sinh((2*b-2*d)*x+2*a-2*c)+1/8/(2*b+2*d)*sinh((2*b+2*d)*x+2*a+2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 192 vs. 2(78) = 156.

Time = 0.09 (sec) , antiderivative size = 192, normalized size of antiderivative = 2.18

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx$$

$$= \frac{b^2d \cosh(bx + a) \sinh(bx + a) \sinh(dx + c)^2 + (b^3d - bd^3)x + (b^2d \cosh(bx + a) \cosh(dx + c)^2 + (b^2d - b^2d) \cosh(bx + a) \sinh(dx + c) \sinh(dx + c) + (b^2d - b^2d) \cosh(dx + c) \sinh(dx + c) \sinh(dx + c) + (b^2d - b^2d) \sinh(dx + c)^2)}{4((b^3d - bd^3) \cosh(bx + a) \sinh(bx + a) \sinh(dx + c) + (b^2d \cosh(bx + a) \cosh(dx + c)^2 + (b^2d - b^2d) \cosh(bx + a) \sinh(dx + c) \sinh(dx + c) + (b^2d - b^2d) \cosh(dx + c) \sinh(dx + c) \sinh(dx + c) + (b^2d - b^2d) \sinh(dx + c)^2)}$$

input `integrate(cosh(b*x+a)^2*cosh(d*x+c)^2,x, algorithm="fricas")`

output
$$\frac{1}{4}(b^2d\cosh(bx+a)\sinh(bx+a)\sinh(dx+c)^2 + (b^3d - b^2d^3)x + (b^2d\cosh(bx+a)\cosh(dx+c)^2 + (b^2d - d^3)\cosh(bx+a))\sinh(bx+a) - (b^2d^2\cosh(dx+c)\sinh(bx+a)^2 + (b^2d^2\cosh(bx+a)^2 - b^3 + b^2d^2)\cosh(dx+c))\sinh(dx+c)) / ((b^3d - b^2d^3)\cosh(bx+a)^2 - (b^3d - b^2d^3)\sinh(bx+a)^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1027 vs. $2(76) = 152$.

Time = 1.53 (sec) , antiderivative size = 1027, normalized size of antiderivative = 11.67

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)**2*cosh(d*x+c)**2,x)`

output

```
Piecewise((x*cosh(a)**2*cosh(c)**2, Eq(b, 0) & Eq(d, 0)), ((-x*sinh(c + d*x)**2/2 + x*cosh(c + d*x)**2/2 + sinh(c + d*x)*cosh(c + d*x)/(2*d))*cosh(a)**2, Eq(b, 0)), (3*x*sinh(a - d*x)**2*sinh(c + d*x)**2/8 - x*sinh(a - d*x)**2*cosh(c + d*x)**2/8 + x*sinh(a - d*x)*sinh(c + d*x)*cosh(a - d*x)*cosh(c + d*x)/2 - x*sinh(c + d*x)**2*cosh(a - d*x)**2/8 + 3*x*cosh(a - d*x)**2*cosh(c + d*x)**2/8 + 3*sinh(a - d*x)*sinh(c + d*x)**2*cosh(a - d*x)/(8*d) - sinh(a - d*x)*cosh(a - d*x)*cosh(c + d*x)**2/(8*d) + sinh(c + d*x)*cosh(a - d*x)**2*cosh(c + d*x)/(2*d), Eq(b, -d)), (3*x*sinh(a + d*x)**2*sinh(c + d*x)**2/8 - x*sinh(a + d*x)**2*cosh(c + d*x)**2/8 - x*sinh(a + d*x)*sinh(c + d*x)*cosh(a + d*x)*cosh(c + d*x)/2 - x*sinh(c + d*x)**2*cosh(a + d*x)**2/8 + 3*x*cosh(a + d*x)**2*cosh(c + d*x)**2/8 - sinh(a + d*x)**2*sinh(c + d*x)*cosh(c + d*x)/(2*d) + sinh(a + d*x)*sinh(c + d*x)**2*cosh(a + d*x)/(8*d) + 5*sinh(a + d*x)*cosh(a + d*x)*cosh(c + d*x)**2/(8*d), Eq(b, d)), ((-x*sinh(a + b*x)**2/2 + x*cosh(a + b*x)**2/2 + sinh(a + b*x)*cosh(a + b*x)/(2*b))*cosh(c)**2, Eq(d, 0)), (b**3*d*x*sinh(a + b*x)**2*sinh(c + d*x)**2/(4*b**3*d - 4*b*d**3) - b**3*d*x*sinh(a + b*x)**2*cosh(c + d*x)**2/(4*b**3*d - 4*b*d**3) - b**3*d*x*sinh(c + d*x)**2*cosh(a + b*x)**2/(4*b**3*d - 4*b*d**3) + b**3*d*x*cosh(a + b*x)**2*cosh(c + d*x)**2/(4*b**3*d - 4*b*d**3) - b**3*sinh(a + b*x)**2*sinh(c + d*x)*cosh(c + d*x)/(4*b**3*d - 4*b*d**3) + b**3*sinh(c + d*x)*cosh(a + b*x)**2*cosh(c + d*x)/(4*b**3*d - 4*b...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(b*x+a)^2*cosh(d*x+c)^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-(2*d)/b>0)', see `assume?` for more deta
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.77

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \frac{1}{4}x + \frac{e^{(2bx+2dx+2a+2c)}}{32(b+d)} + \frac{e^{(2bx-2dx+2a-2c)}}{32(b-d)} + \frac{e^{(2bx+2a)}}{16b} - \frac{e^{(-2bx+2dx-2a+2c)}}{32(b-d)} - \frac{e^{(-2bx-2dx-2a-2c)}}{32(b+d)} - \frac{e^{(-2bx-2a)}}{16b} + \frac{e^{(2dx+2c)}}{16d} - \frac{e^{(-2dx-2c)}}{16d}$$

input `integrate(cosh(b*x+a)^2*cosh(d*x+c)^2,x, algorithm="giac")`output `1/4*x + 1/32*e^(2*b*x + 2*d*x + 2*a + 2*c)/(b + d) + 1/32*e^(2*b*x - 2*d*x + 2*a - 2*c)/(b - d) + 1/16*e^(2*b*x + 2*a)/b - 1/32*e^(-2*b*x + 2*d*x - 2*a + 2*c)/(b - d) - 1/32*e^(-2*b*x - 2*d*x - 2*a - 2*c)/(b + d) - 1/16*e^(-2*b*x - 2*a)/b + 1/16*e^(2*d*x + 2*c)/d - 1/16*e^(-2*d*x - 2*c)/d`**Mupad [B] (verification not implemented)**

Time = 1.50 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.31

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx = \frac{d^3 \cosh(a + bx) \sinh(a + bx) - b^3 \cosh(c + dx) \sinh(c + dx) + b d^3 x - b^3 dx - 2 b^2 d \cosh(a + bx) \cosh(c + dx) + 2 b^2 d \sinh(a + bx) \sinh(c + dx)}{4 b d^3 - 4 b^3 d}$$

input `int(cosh(a + b*x)^2*cosh(c + d*x)^2,x)`output `(d^3*cosh(a + b*x)*sinh(a + b*x) - b^3*cosh(c + d*x)*sinh(c + d*x) + b*d^3*x - b^3*d*x - 2*b^2*d*cosh(a + b*x)*cosh(c + d*x)^2*sinh(a + b*x) + 2*b*d^2*cosh(a + b*x)^2*cosh(c + d*x)*sinh(c + d*x))/(4*b*d^3 - 4*b^3*d)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.06

$$\int \cosh^2(a + bx) \cosh^2(c + dx) dx$$

$$= \frac{e^{4bx+4dx+4a+4c} b^2 d - e^{4bx+4dx+4a+4c} b d^2 + 2e^{4bx+2dx+4a+2c} b^2 d - 2e^{4bx+2dx+4a+2c} d^3 + e^{4bx+4a} b^2 d + e^{4bx+4a} b d^2}{32e^{2a+2bx+2c+2dx} (b^2 d - d^3)}$$

input

```
int(cosh(b*x+a)^2*cosh(d*x+c)^2,x)
```

output

```
(e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d - e**(4*a + 4*b*x + 4*c + 4*d*x)*b*
d**2 + 2*e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d - 2*e**(4*a + 4*b*x + 2*c +
2*d*x)*d**3 + e**(4*a + 4*b*x)*b**2*d + e**(4*a + 4*b*x)*b*d**2 + 2*e**(2
*a + 2*b*x + 4*c + 4*d*x)*b**3 - 2*e**(2*a + 2*b*x + 4*c + 4*d*x)*b*d**2 +
8*e**(2*a + 2*b*x + 2*c + 2*d*x)*b**3*d*x - 8*e**(2*a + 2*b*x + 2*c + 2*d
*x)*b*d**3*x - 2*e**(2*a + 2*b*x)*b**3 + 2*e**(2*a + 2*b*x)*b*d**2 - e**(4
*c + 4*d*x)*b**2*d - e**(4*c + 4*d*x)*b*d**2 - 2*e**(2*c + 2*d*x)*b**2*d +
2*e**(2*c + 2*d*x)*d**3 - b**2*d + b*d**2)/(32*e**(2*a + 2*b*x + 2*c + 2*
d*x)*b*d*(b**2 - d**2))
```


3.101 $\int \cosh^2(a + bx) \cosh^3(c + dx) dx$

Optimal result	712
Mathematica [A] (verified)	713
Rubi [A] (verified)	713
Maple [A] (verified)	715
Fricas [B] (verification not implemented)	715
Sympy [B] (verification not implemented)	716
Maxima [F(-2)]	717
Giac [A] (verification not implemented)	718
Mupad [B] (verification not implemented)	719
Reduce [B] (verification not implemented)	720

Optimal result

Integrand size = 17, antiderivative size = 144

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx = \frac{\sinh(2a - 3c + (2b - 3d)x)}{16(2b - 3d)} + \frac{3 \sinh(2a - c + (2b - d)x)}{16(2b - d)} + \frac{3 \sinh(c + dx)}{8d} + \frac{\sinh(3c + 3dx)}{24d} + \frac{3 \sinh(2a + c + (2b + d)x)}{16(2b + d)} + \frac{\sinh(2a + 3c + (2b + 3d)x)}{16(2b + 3d)}$$

output

```
sinh(2*a-3*c+(2*b-3*d)*x)/(32*b-48*d)+3*sinh(2*a-c+(2*b-d)*x)/(32*b-16*d)+
3/8*sinh(d*x+c)/d+1/24*sinh(3*d*x+3*c)/d+3*sinh(2*a+c+(2*b+d)*x)/(32*b+16*
d)+sinh(2*a+3*c+(2*b+3*d)*x)/(32*b+48*d)
```

Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.10

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx = \frac{1}{48} \left(\frac{18 \cosh(dx) \sinh(c)}{d} + \frac{2 \cosh(3dx) \sinh(3c)}{d} + \frac{18 \cosh(c) \sinh(dx)}{d} + \frac{2 \cosh(3c) \sinh(3dx)}{d} + \frac{3 \sinh(2a - 3c + 2bx - 3dx)}{2b - 3d} + \frac{9 \sinh(2a - c + 2bx - dx)}{2b - d} + \frac{9 \sinh(2a + c + 2bx + dx)}{2b + d} + \frac{3 \sinh(2a + 3c + 2bx + 3dx)}{2b + 3d} \right)$$

input

```
Integrate[Cosh[a + b*x]^2*Cosh[c + d*x]^3,x]
```

output

```
((18*Cosh[d*x]*Sinh[c])/d + (2*Cosh[3*d*x]*Sinh[3*c])/d + (18*Cosh[c]*Sinh[d*x])/d + (2*Cosh[3*c]*Sinh[3*d*x])/d + (3*Sinh[2*a - 3*c + 2*b*x - 3*d*x])/((2*b - 3*d) + (9*Sinh[2*a - c + 2*b*x - d*x])/((2*b - d) + (9*Sinh[2*a + c + 2*b*x + d*x])/((2*b + d) + (3*Sinh[2*a + 3*c + 2*b*x + 3*d*x])/((2*b + 3*d)))/48
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx$$

↓ 6148

$$\int \left(\frac{1}{16} \cosh(2a + x(2b - 3d) - 3c) + \frac{3}{16} \cosh(2a + x(2b - d) - c) + \frac{3}{16} \cosh(2a + x(2b + d) + c) + \frac{1}{16} \cosh(2a + x(2b + 3d) + 3c) + \frac{3 \sinh(c + dx)}{8d} + \frac{\sinh(3c + 3dx)}{24d} \right) dx$$

↓ 2009

$$\frac{\sinh(2a + x(2b - 3d) - 3c)}{16(2b - 3d)} + \frac{3 \sinh(2a + x(2b - d) - c)}{16(2b - d)} + \frac{3 \sinh(2a + x(2b + d) + c)}{16(2b + d)} + \frac{\sinh(2a + x(2b + 3d) + 3c)}{16(2b + 3d)} + \frac{3 \sinh(c + dx)}{8d} + \frac{\sinh(3c + 3dx)}{24d}$$

input `Int[Cosh[a + b*x]^2*Cosh[c + d*x]^3,x]`

output `Sinh[2*a - 3*c + (2*b - 3*d)*x]/(16*(2*b - 3*d)) + (3*Sinh[2*a - c + (2*b - d)*x])/(16*(2*b - d)) + (3*Sinh[c + d*x])/(8*d) + Sinh[3*c + 3*d*x]/(24*d) + (3*Sinh[2*a + c + (2*b + d)*x])/(16*(2*b + d)) + Sinh[2*a + 3*c + (2*b + 3*d)*x]/(16*(2*b + 3*d))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6148 `Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 9.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.92

method	result
default	$\frac{3 \sinh(dx+c)}{8d} + \frac{\sinh(3dx+3c)}{24d} + \frac{\sinh(2a-3c+(2b-3d)x)}{32b-48d} + \frac{3 \sinh(2a-c+(2b-d)x)}{16(2b-d)} + \frac{3 \sinh(2a+c+(2b+d)x)}{16(2b+d)} + \frac{\sinh(2a-3c+(2b-3d)x)}{32b-48d}$
parallelrisc	$\frac{(24b^3d+36b^2d^2-6bd^3-9d^4) \sinh(2a-3c+(2b-3d)x)+72\left(b-\frac{3d}{2}\right) \left(\left(b+\frac{3d}{2}\right)d\left(b+\frac{d}{2}\right) \sinh(2a-c+(2b-d)x)+\left(b-\frac{d}{2}\right) \left(\frac{d(b+\frac{3d}{2})}{768b^4d-1920b^2d^3+432d^4}\right) \sinh(2a+c+(2b+d)x)\right)}{768b^4d-1920b^2d^3+432d^4}$
risc	$\frac{(6de^{4bx+4a}b-9d^2e^{4bx+4a}+8b^2e^{2bx+2a}-18d^2e^{2bx+2a}-6bd-9d^2)e^{-2bx+3dx-2a+3c}}{96(2b+3d)(2b-3d)d} + \frac{3(2de^{4bx+4a}b-d^2e^{4bx+4a}+8b^2e^{2bx+2a}-18d^2e^{2bx+2a}-6bd-9d^2)e^{-2bx+3dx-2a+3c}}{32(2b+3d)(2b-3d)d}$
orering	Expression too large to display

input `int(cosh(b*x+a)^2*cosh(d*x+c)^3,x,method=_RETURNVERBOSE)`

output
$$\frac{3}{8} \frac{\sinh(dx+c)}{d} + \frac{1}{24} \frac{\sinh(3dx+3c)}{d} + \frac{1}{16} \frac{\sinh(2a-3c+(2b-3d)x)}{(2b-3d)} + \frac{3}{16} \frac{\sinh(2a-c+(2b-d)x)}{(2b-d)} + \frac{3}{16} \frac{\sinh(2a+c+(2b+d)x)}{(2b+d)} + \frac{1}{16} \frac{\sinh(2a-3c+(2b-3d)x)}{(2b+3d)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(132) = 264.

Time = 0.10 (sec) , antiderivative size = 397, normalized size of antiderivative = 2.76

$$\int \cosh^2(a+bx) \cosh^3(c+dx) dx$$

$$= \frac{36(4b^3d-bd^3) \cosh(bx+a) \cosh(dx+c) \sinh(bx+a) \sinh(dx+c)^2 + (16b^4-40b^2d^2+9d^4-9(4b^3d-bd^3)) \cosh^2(bx+a) \cosh^2(dx+c) \sinh(bx+a) \sinh(dx+c) + (16b^4-40b^2d^2+9d^4-9(4b^3d-bd^3)) \cosh^3(bx+a) \cosh^3(dx+c)}{36(4b^3d-bd^3) \cosh(bx+a) \cosh(dx+c) \sinh(bx+a) \sinh(dx+c)^2 + (16b^4-40b^2d^2+9d^4-9(4b^3d-bd^3)) \cosh^2(bx+a) \cosh^2(dx+c) \sinh(bx+a) \sinh(dx+c) + (16b^4-40b^2d^2+9d^4-9(4b^3d-bd^3)) \cosh^3(bx+a) \cosh^3(dx+c)}$$

input `integrate(cosh(b*x+a)^2*cosh(d*x+c)^3,x, algorithm="fricas")`

output

```
1/24*(36*(4*b^3*d - b*d^3)*cosh(b*x + a)*cosh(d*x + c)*sinh(b*x + a)*sinh(
d*x + c)^2 + (16*b^4 - 40*b^2*d^2 + 9*d^4 - 9*(4*b^2*d^2 - d^4)*cosh(b*x +
a)^2 - 9*(4*b^2*d^2 - d^4)*sinh(b*x + a)^2)*sinh(d*x + c)^3 + 12*((4*b^3*
d - b*d^3)*cosh(b*x + a)*cosh(d*x + c)^3 + 3*(4*b^3*d - 9*b*d^3)*cosh(b*x
+ a)*cosh(d*x + c)*sinh(b*x + a) + 3*(48*b^4 - 120*b^2*d^2 + 27*d^4 - 3*(
4*b^2*d^2 - 9*d^4)*cosh(b*x + a)^2 + (16*b^4 - 40*b^2*d^2 + 9*d^4 - 9*(4*b
^2*d^2 - d^4)*cosh(b*x + a)^2)*cosh(d*x + c)^2 - 3*(4*b^2*d^2 - 9*d^4 + 3*
(4*b^2*d^2 - d^4)*cosh(d*x + c)^2)*sinh(b*x + a)^2)*sinh(d*x + c))/((16*b^
4*d - 40*b^2*d^3 + 9*d^5)*cosh(b*x + a)^2 - (16*b^4*d - 40*b^2*d^3 + 9*d^5
)*sinh(b*x + a)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2009 vs. 2(116) = 232.

Time = 5.32 (sec) , antiderivative size = 2009, normalized size of antiderivative = 13.95

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)**2*cosh(d*x+c)**3,x)
```

output

```
Piecewise((x*cosh(a)**2*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (3*x*sinh(a - 3*d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + x*sinh(a - 3*d*x/2)**2*cosh(c + d*x)**3/16 + x*sinh(a - 3*d*x/2)*sinh(c + d*x)**3*cosh(a - 3*d*x/2)/8 + 3*x*sinh(a - 3*d*x/2)*sinh(c + d*x)*cosh(a - 3*d*x/2)*cosh(c + d*x)**2/8 + 3*x*sinh(c + d*x)**2*cosh(a - 3*d*x/2)**2*cosh(c + d*x)/16 + x*cosh(a - 3*d*x/2)**2*cosh(c + d*x)**3/16 + 11*sinh(a - 3*d*x/2)**2*sinh(c + d*x)**3/(48*d) - sinh(a - 3*d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/d - 3*sinh(a - 3*d*x/2)*sinh(c + d*x)**2*cosh(a - 3*d*x/2)*cosh(c + d*x)/(4*d) - 5*sinh(a - 3*d*x/2)*cosh(a - 3*d*x/2)*cosh(c + d*x)**3/(8*d) - 7*sinh(c + d*x)**3*cosh(a - 3*d*x/2)**2/(16*d), Eq(b, -3*d/2)), (-3*x*sinh(a - d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + 3*x*sinh(a - d*x/2)**2*cosh(c + d*x)**3/16 - 3*x*sinh(a - d*x/2)*sinh(c + d*x)**3*cosh(a - d*x/2)/8 + 3*x*sinh(a - d*x/2)*sinh(c + d*x)*cosh(a - d*x/2)*cosh(c + d*x)**2/8 - 3*x*sinh(c + d*x)**2*cosh(a - d*x/2)**2*cosh(c + d*x)/16 + 3*x*cosh(a - d*x/2)**2*cosh(c + d*x)**3/16 + 49*sinh(a - d*x/2)**2*sinh(c + d*x)**3/(48*d) - sinh(a - d*x/2)**2*sinh(c + d*x)*cosh(c + d*x)**2/d + 7*sinh(a - d*x/2)*sinh(c + d*x)**2*cosh(a - d*x/2)*cosh(c + d*x)/(4*d) - 13*sinh(a - d*x/2)*cosh(a - d*x/2)*cosh(c + d*x)**3/(8*d) + 17*sinh(c + d*x)**3*cosh(a - d*x/2)**2/(48*d), Eq(b, -d/2)), (-3*x*sinh(a + d*x/2)**2*sinh(c + d*x)**2*cosh(c + d*x)/16 + 3*x*sinh(a + d*x/2)**2*cosh(c + d*x)**3/16 + 3*x*sinh(a + d*x/2)*si...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(b*x+a)^2*cosh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(1-(3*d)/b>0)', see `assume?` for more deta
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.81

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx = \frac{e^{(2bx+3dx+2a+3c)}}{32(2b+3d)} + \frac{3e^{(2bx+dx+2a+c)}}{32(2b+d)}$$

$$+ \frac{3e^{(2bx-dx+2a-c)}}{32(2b-d)} + \frac{e^{(2bx-3dx+2a-3c)}}{32(2b-3d)}$$

$$- \frac{e^{(-2bx+3dx-2a+3c)}}{32(2b-3d)} - \frac{3e^{(-2bx+dx-2a+c)}}{32(2b-d)}$$

$$- \frac{3e^{(-2bx-dx-2a-c)}}{32(2b+d)} - \frac{e^{(-2bx-3dx-2a-3c)}}{32(2b+3d)}$$

$$+ \frac{e^{(3dx+3c)}}{48d} + \frac{3e^{(dx+c)}}{16d} - \frac{3e^{(-dx-c)}}{16d} - \frac{e^{(-3dx-3c)}}{48d}$$

input `integrate(cosh(b*x+a)^2*cosh(d*x+c)^3,x, algorithm="giac")`

output `1/32*e^(2*b*x + 3*d*x + 2*a + 3*c)/(2*b + 3*d) + 3/32*e^(2*b*x + d*x + 2*a + c)/(2*b + d) + 3/32*e^(2*b*x - d*x + 2*a - c)/(2*b - d) + 1/32*e^(2*b*x - 3*d*x + 2*a - 3*c)/(2*b - 3*d) - 1/32*e^(-2*b*x + 3*d*x - 2*a + 3*c)/(2*b - 3*d) - 3/32*e^(-2*b*x + d*x - 2*a + c)/(2*b - d) - 3/32*e^(-2*b*x - d*x - 2*a - c)/(2*b + d) - 1/32*e^(-2*b*x - 3*d*x - 2*a - 3*c)/(2*b + 3*d) + 1/48*e^(3*d*x + 3*c)/d + 3/16*e^(d*x + c)/d - 3/16*e^(-d*x - c)/d - 1/48*e^(-3*d*x - 3*c)/d`

Mupad [B] (verification not implemented)

Time = 1.54 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.34

$$\begin{aligned}
& \int \cosh^2(a + bx) \cosh^3(c + dx) dx \\
&= \frac{\cosh(a + bx)^2 \cosh(c + dx)^2 \sinh(c + dx) (8b^4 - 26b^2 d^2 + 9d^4)}{d (16b^4 - 40b^2 d^2 + 9d^4)} \\
&\quad - \sinh(a + bx)^2 \sinh(c + dx)^3 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} - \frac{1}{3d} \right) \\
&\quad - \frac{2 \cosh(a + bx) \cosh(c + dx)^3 \sinh(a + bx) (7bd^2 - 4b^3)}{16b^4 - 40b^2 d^2 + 9d^4} \\
&\quad - \cosh(a + bx)^2 \sinh(c + dx)^3 \left(\frac{3d^3}{16b^4 - 40b^2 d^2 + 9d^4} + \frac{1}{3d} \right) \\
&\quad + \frac{12bd^2 \cosh(a + bx) \cosh(c + dx) \sinh(a + bx) \sinh(c + dx)^2}{16b^4 - 40b^2 d^2 + 9d^4} \\
&\quad - \frac{2b^2 \cosh(c + dx)^2 \sinh(a + bx)^2 \sinh(c + dx) (4b^2 - 7d^2)}{d (16b^4 - 40b^2 d^2 + 9d^4)}
\end{aligned}$$

input `int(cosh(a + b*x)^2*cosh(c + d*x)^3,x)`output `(cosh(a + b*x)^2*cosh(c + d*x)^2*sinh(c + d*x)*(8*b^4 + 9*d^4 - 26*b^2*d^2)) / (d*(16*b^4 + 9*d^4 - 40*b^2*d^2)) - sinh(a + b*x)^2*sinh(c + d*x)^3*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) - 1/(3*d)) - (2*cosh(a + b*x)*cosh(c + d*x)^3*sinh(a + b*x)*(7*b*d^2 - 4*b^3))/(16*b^4 + 9*d^4 - 40*b^2*d^2) - cosh(a + b*x)^2*sinh(c + d*x)^3*((3*d^3)/(16*b^4 + 9*d^4 - 40*b^2*d^2) + 1/(3*d)) + (12*b*d^2*cosh(a + b*x)*cosh(c + d*x)*sinh(a + b*x)*sinh(c + d*x)^2)/(16*b^4 + 9*d^4 - 40*b^2*d^2) - (2*b^2*cosh(c + d*x)^2*sinh(a + b*x)^2*sinh(c + d*x)*(4*b^2 - 7*d^2))/(d*(16*b^4 + 9*d^4 - 40*b^2*d^2))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 865, normalized size of antiderivative = 6.01

$$\int \cosh^2(a + bx) \cosh^3(c + dx) dx$$

$$= \frac{-36e^{4bx+4dx+4a+4c}b^2d^2 - 162e^{4bx+4dx+4a+4c}bd^3 + 72e^{4bx+2dx+4a+2c}b^3d + 36e^{4bx+2dx+4a+2c}b^2d^2 - 162e^{4bx+2c}}{d^4}$$

input `int(cosh(b*x+a)^2*cosh(d*x+c)^3,x)`

output

```
(24***e**(4*a + 4*b*x + 6*c + 6*d*x)*b**3*d - 36***e**(4*a + 4*b*x + 6*c + 6*d*x)*b**2*d**2 - 6***e**(4*a + 4*b*x + 6*c + 6*d*x)*b*d**3 + 9***e**(4*a + 4*b*x + 6*c + 6*d*x)*d**4 + 72***e**(4*a + 4*b*x + 4*c + 4*d*x)*b**3*d - 36***e**(4*a + 4*b*x + 4*c + 4*d*x)*b**2*d**2 - 162***e**(4*a + 4*b*x + 4*c + 4*d*x)*b*d**3 + 81***e**(4*a + 4*b*x + 4*c + 4*d*x)*d**4 + 72***e**(4*a + 4*b*x + 2*c + 2*d*x)*b**3*d + 36***e**(4*a + 4*b*x + 2*c + 2*d*x)*b**2*d**2 - 162***e**(4*a + 4*b*x + 2*c + 2*d*x)*b*d**3 - 81***e**(4*a + 4*b*x + 2*c + 2*d*x)*d**4 + 24***e**(4*a + 4*b*x)*b**3*d + 36***e**(4*a + 4*b*x)*b**2*d**2 - 6***e**(4*a + 4*b*x)*b*d**3 - 9***e**(4*a + 4*b*x)*d**4 + 32***e**(2*a + 2*b*x + 6*c + 6*d*x)*b**4 - 80***e**(2*a + 2*b*x + 6*c + 6*d*x)*b**2*d**2 + 18***e**(2*a + 2*b*x + 6*c + 6*d*x)*d**4 + 288***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**4 - 720***e**(2*a + 2*b*x + 4*c + 4*d*x)*b**2*d**2 + 162***e**(2*a + 2*b*x + 4*c + 4*d*x)*d**4 - 288***e**(2*a + 2*b*x + 2*c + 2*d*x)*b**4 + 720***e**(2*a + 2*b*x + 2*c + 2*d*x)*b**2*d**2 - 162***e**(2*a + 2*b*x + 2*c + 2*d*x)*d**4 - 32***e**(2*a + 2*b*x)*b**4 + 80***e**(2*a + 2*b*x)*b**2*d**2 - 18***e**(2*a + 2*b*x)*d**4 - 24***e**(6*c + 6*d*x)*b**3*d - 36***e**(6*c + 6*d*x)*b**2*d**2 + 6***e**(6*c + 6*d*x)*b*d**3 + 9***e**(6*c + 6*d*x)*d**4 - 72***e**(4*c + 4*d*x)*b**3*d - 36***e**(4*c + 4*d*x)*b**2*d**2 + 162***e**(4*c + 4*d*x)*b*d**3 + 81***e**(4*c + 4*d*x)*d**4 - 72***e**(2*c + 2*d*x)*b**3*d + 36***e**(2*c + 2*d*x)*b**2*d**2 + 162***e**(2*c + 2*d*x)*b*d**3 - 81***e**(2*c + 2*d*x)*d**4 - 24*b**3*d + 36*...
```

3.102 $\int \cosh^3(a + bx) \cosh^3(c + dx) dx$

Optimal result	721
Mathematica [A] (verified)	722
Rubi [A] (verified)	722
Maple [A] (verified)	724
Fricas [B] (verification not implemented)	724
Sympy [B] (verification not implemented)	725
Maxima [F(-2)]	726
Giac [B] (verification not implemented)	727
Mupad [B] (verification not implemented)	728
Reduce [F]	729

Optimal result

Integrand size = 17, antiderivative size = 195

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \frac{3 \sinh(a - 3c + (b - 3d)x)}{32(b - 3d)} + \frac{9 \sinh(a - c + (b - d)x)}{32(b - d)} + \frac{\sinh(3(a - c) + 3(b - d)x)}{96(b - d)} + \frac{3 \sinh(3a - c + (3b - d)x)}{32(3b - d)} + \frac{9 \sinh(a + c + (b + d)x)}{32(b + d)} + \frac{\sinh(3(a + c) + 3(b + d)x)}{96(b + d)} + \frac{3 \sinh(3a + c + (3b + d)x)}{32(3b + d)} + \frac{3 \sinh(a + 3c + (b + 3d)x)}{32(b + 3d)}$$

output

```
3*sinh(a-3*c+(b-3*d)*x)/(32*b-96*d)+9*sinh(a-c+(b-d)*x)/(32*b-32*d)+sinh(3
*a-3*c+3*(b-d)*x)/(96*b-96*d)+3*sinh(3*a-c+(3*b-d)*x)/(96*b-32*d)+9*sinh(a
+c+(b+d)*x)/(32*b+32*d)+sinh(3*a+3*c+3*(b+d)*x)/(96*b+96*d)+3*sinh(3*a+c+(
3*b+d)*x)/(96*b+32*d)+3*sinh(a+3*c+(b+3*d)*x)/(32*b+96*d)
```

Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.90

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \frac{1}{96} \left(\frac{9 \sinh(a - 3c + bx - 3dx)}{b - 3d} + \frac{27 \sinh(a - c + bx - dx)}{b - d} + \frac{\sinh(3(a - c + bx - dx))}{b - d} + \frac{9 \sinh(3a - c + 3bx - dx)}{3b - d} + \frac{9 \sinh(3a + c + 3bx + dx)}{3b + d} + \frac{9 \sinh(a + 3c + bx + 3dx)}{b + 3d} + \frac{27 \sinh(a + c + (b + d)x)}{b + d} + \frac{\sinh(3(a + c + (b + d)x))}{b + d} \right)$$

input `Integrate[Cosh[a + b*x]^3*Cosh[c + d*x]^3,x]`

output `((9*Sinh[a - 3*c + b*x - 3*d*x])/(b - 3*d) + (27*Sinh[a - c + b*x - d*x])/(b - d) + Sinh[3*(a - c + b*x - d*x)]/(b - d) + (9*Sinh[3*a - c + 3*b*x - d*x])/(3*b - d) + (9*Sinh[3*a + c + 3*b*x + d*x])/(3*b + d) + (9*Sinh[a + 3*c + b*x + 3*d*x])/(b + 3*d) + (27*Sinh[a + c + (b + d)*x])/(b + d) + Sinh[3*(a + c + (b + d)*x)]/(b + d))/96`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {6148, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx$$

↓ 6148

$$\int \left(\frac{3}{32} \cosh(a + x(b - 3d) - 3c) + \frac{9}{32} \cosh(a + x(b - d) - c) + \frac{1}{32} \cosh(3(a - c) + 3x(b - d)) + \frac{3}{32} \cosh(3a + x(3b - d) - c) \right. \\ \left. + \frac{9}{32} \cosh(a + x(b + d) + c) + \frac{1}{32} \cosh(3(a + c) + 3x(b + d)) + \frac{3}{32} \cosh(3a + x(3b + d) + c) \right) dx$$

↓ 2009

$$\frac{3 \sinh(a + x(b - 3d) - 3c)}{32(b - 3d)} + \frac{9 \sinh(a + x(b - d) - c)}{32(b - d)} + \frac{\sinh(3(a - c) + 3x(b - d))}{96(b - d)} + \\ \frac{3 \sinh(3a + x(3b - d) - c)}{32(3b - d)} + \frac{9 \sinh(a + x(b + d) + c)}{32(b + d)} + \frac{\sinh(3(a + c) + 3x(b + d))}{96(b + d)} + \\ \frac{3 \sinh(3a + x(3b + d) + c)}{32(3b + d)} + \frac{3 \sinh(a + x(b + 3d) + 3c)}{32(b + 3d)}$$

input

```
Int[Cosh[a + b*x]^3*Cosh[c + d*x]^3,x]
```

output

```
(3*Sinh[a - 3*c + (b - 3*d)*x])/(32*(b - 3*d)) + (9*Sinh[a - c + (b - d)*x])/(32*(b - d)) + Sinh[3*(a - c) + 3*(b - d)*x]/(96*(b - d)) + (3*Sinh[3*a - c + (3*b - d)*x])/(32*(3*b - d)) + (9*Sinh[a + c + (b + d)*x])/(32*(b + d)) + Sinh[3*(a + c) + 3*(b + d)*x]/(96*(b + d)) + (3*Sinh[3*a + c + (3*b + d)*x])/(32*(3*b + d)) + (3*Sinh[a + 3*c + (b + 3*d)*x])/(32*(b + 3*d))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 6148

```
Int[Cosh[v_]^(p_.)*Cosh[w_]^(q_.), x_Symbol] := Int[ExpandTrigReduce[Cosh[v]^p*Cosh[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))
```

Maple [A] (verified)

Time = 28.76 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.97

method	result
default	$\frac{3 \sinh(a-3c+(b-3d)x)}{32(b-3d)} + \frac{9 \sinh(a-c+(b-d)x)}{32(b-d)} + \frac{9 \sinh(a+c+(b+d)x)}{32(b+d)} + \frac{3 \sinh(a+3c+(b+3d)x)}{32(b+3d)} + \frac{\sinh((3b-3d)x+3a-3c)}{96b-96d}$
parallelrisc	$\frac{9(b-d)(b+3d)(b-3d)(b+d)\left(b+\frac{d}{3}\right) \sinh(3a-c+(3b-d)x)}{32} + \frac{9\left(\frac{(b+3d)(b-3d)(b+d)\left(b+\frac{d}{3}\right) \sinh((3b-3d)x+3a-3c)}{3} + \frac{(b-d)(b+3d)(b-3d)(b+d)}{192(b+d)(b+3d)(b-d)(b-3d)}\right)}{192(b+d)(b+3d)(b-d)(b-3d)}$
risc	$\frac{(b^3 e^{6bx+6a} - b^2 d e^{6bx+6a} - 9b d^2 e^{6bx+6a} + 9d^3 e^{6bx+6a} + 9b^3 e^{4bx+4a} - 27b^2 d e^{4bx+4a} - 9b d^2 e^{4bx+4a} + 27d^3 e^{4bx+4a} - 9b^3 e^{2bx+2a}) \sinh(3a-c+(3b-d)x)}{192(b+d)(b+3d)(b-d)(b-3d)}$
orering	Expression too large to display

input

```
int(cosh(b*x+a)^3*cosh(d*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
3/32*sinh(a-3*c+(b-3*d)*x)/(b-3*d)+9/32*sinh(a-c+(b-d)*x)/(b-d)+9/32/(b+d)
*sinh(a+c+(b+d)*x)+3/32/(b+3*d)*sinh(a+3*c+(b+3*d)*x)+1/32/(3*b-3*d)*sinh(
(3*b-3*d)*x+3*a-3*c)+3/32/(3*b-d)*sinh(3*a-c+(3*b-d)*x)+3/32/(3*b+d)*sinh(
3*a+c+(3*b+d)*x)+1/32/(3*b+3*d)*sinh((3*b+3*d)*x+3*a+3*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 726 vs. 2(179) = 358.

Time = 0.12 (sec) , antiderivative size = 726, normalized size of antiderivative = 3.72

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^3*cosh(d*x+c)^3,x, algorithm="fricas")
```

output

```

1/48*(((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(d*x + c)^3 + 27*(b^5 - 10*b^3*d
^2 + 9*b*d^4)*cosh(d*x + c))*sinh(b*x + a)^3 - ((9*b^4*d - 82*b^2*d^3 + 9*
d^5)*cosh(b*x + a)^3 + 3*(9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)*sinh
(b*x + a)^2 + 27*(9*b^4*d - 10*b^2*d^3 + d^5)*cosh(b*x + a))*sinh(d*x + c)
^3 + 3*((9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(d*x + c)*sinh(b*x + a)^3 + 3*(
27*b^5 - 30*b^3*d^2 + 3*b*d^4 + (9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x +
a)^2)*cosh(d*x + c)*sinh(b*x + a))*sinh(d*x + c)^2 + 3*((27*b^5 - 30*b^3*d
^2 + 3*b*d^4 + (9*b^5 - 82*b^3*d^2 + 9*b*d^4)*cosh(b*x + a)^2)*cosh(d*x +
c)^3 + 9*(9*b^5 - 82*b^3*d^2 + 9*b*d^4 + 3*(b^5 - 10*b^3*d^2 + 9*b*d^4)*co
sh(b*x + a)^2)*cosh(d*x + c))*sinh(b*x + a) - 3*(3*(b^4*d - 10*b^2*d^3 + 9
*d^5)*cosh(b*x + a)^3 + ((9*b^4*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)^3 +
27*(9*b^4*d - 10*b^2*d^3 + d^5)*cosh(b*x + a))*cosh(d*x + c)^2 + 3*((9*b^4
*d - 82*b^2*d^3 + 9*d^5)*cosh(b*x + a)*cosh(d*x + c)^2 + 3*(b^4*d - 10*b^2
*d^3 + 9*d^5)*cosh(b*x + a))*sinh(b*x + a)^2 + 9*(9*b^4*d - 82*b^2*d^3 + 9
*d^5)*cosh(b*x + a))*sinh(d*x + c))/((9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*
d^6)*cosh(b*x + a)^4 - 2*(9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*cosh(b*
x + a)^2*sinh(b*x + a)^2 + (9*b^6 - 91*b^4*d^2 + 91*b^2*d^4 - 9*d^6)*sinh(
b*x + a)^4)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3580 vs. $2(172) = 344$.

Time = 16.48 (sec) , antiderivative size = 3580, normalized size of antiderivative = 18.36

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)**3*cosh(d*x+c)**3,x)
```

output

```
Piecewise((x*cosh(a)**3*cosh(c)**3, Eq(b, 0) & Eq(d, 0)), (-3*x*sinh(a - 3*d*x)**3*sinh(c + d*x)**3/32 - 9*x*sinh(a - 3*d*x)**3*sinh(c + d*x)*cosh(c + d*x)**2/32 - 9*x*sinh(a - 3*d*x)**2*sinh(c + d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)/32 - 3*x*sinh(a - 3*d*x)**2*cosh(a - 3*d*x)*cosh(c + d*x)**3/32 + 3*x*sinh(a - 3*d*x)*sinh(c + d*x)**3*cosh(a - 3*d*x)**2/32 + 9*x*sinh(a - 3*d*x)*sinh(c + d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**2/32 + 9*x*sinh(c + d*x)**2*cosh(a - 3*d*x)**3*cosh(c + d*x)/32 + 3*x*cosh(a - 3*d*x)**3*cosh(c + d*x)**3/32 - 51*sinh(a - 3*d*x)**3*sinh(c + d*x)**2*cosh(c + d*x)/(320*d) + sinh(a - 3*d*x)**3*cosh(c + d*x)**3/(5*d) - 27*sinh(a - 3*d*x)**2*sinh(c + d*x)**3*cosh(a - 3*d*x)/(320*d) + 3*sinh(a - 3*d*x)*sinh(c + d*x)**2*cosh(a - 3*d*x)**2*cosh(c + d*x)/(20*d) - 101*sinh(a - 3*d*x)*cosh(a - 3*d*x)**2*cosh(c + d*x)**3/(320*d) + sinh(c + d*x)**3*cosh(a - 3*d*x)**3/(12*d) - 13*sinh(c + d*x)*cosh(a - 3*d*x)**3*cosh(c + d*x)**2/(320*d), Eq(b, -3*d)), (5*x*sinh(a - d*x)**3*sinh(c + d*x)**3/16 - 3*x*sinh(a - d*x)**3*sinh(c + d*x)*cosh(c + d*x)**2/16 + 9*x*sinh(a - d*x)**2*sinh(c + d*x)**2*cosh(a - d*x)*cosh(c + d*x)/16 - 3*x*sinh(a - d*x)**2*cosh(a - d*x)*cosh(c + d*x)**3/16 - 3*x*sinh(a - d*x)*sinh(c + d*x)**3*cosh(a - d*x)**2/16 + 9*x*sinh(a - d*x)*sinh(c + d*x)*cosh(a - d*x)**2*cosh(c + d*x)**2/16 - 3*x*sinh(c + d*x)**2*cosh(a - d*x)**3*cosh(c + d*x)/16 + 5*x*cosh(a - d*x)**3*cosh(c + d*x)**3/16 + 3*sinh(a - d*x)**3*sinh(c + d*x)**2*cosh(c + ...
```

Maxima [F(-2)]

Exception generated.

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \text{Exception raised: ValueError}$$

input

```
integrate(cosh(b*x+a)^3*cosh(d*x+c)^3,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-(3*d)/b>0)', see `assume?` for more detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(179) = 358$.

Time = 0.17 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.91

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \frac{e^{(3bx+3dx+3a+3c)}}{192(b+d)} + \frac{3e^{(3bx+dx+3a+c)}}{64(3b+d)}$$

$$+ \frac{3e^{(3bx-dx+3a-c)}}{64(3b-d)} + \frac{e^{(3bx-3dx+3a-3c)}}{192(b-d)}$$

$$+ \frac{3e^{(bx+3dx+a+3c)}}{64(b+3d)} + \frac{9e^{(bx+dx+a+c)}}{64(b+d)}$$

$$+ \frac{9e^{(bx-dx+a-c)}}{64(b-d)} + \frac{3e^{(bx-3dx+a-3c)}}{64(b-3d)}$$

$$- \frac{3e^{(-bx+3dx-a+3c)}}{64(b-3d)} - \frac{9e^{(-bx+dx-a+c)}}{64(b-d)}$$

$$- \frac{9e^{(-bx-dx-a-c)}}{64(b+d)} - \frac{3e^{(-bx-3dx-a-3c)}}{64(b+3d)}$$

$$- \frac{e^{(-3bx+3dx-3a+3c)}}{192(b-d)} - \frac{3e^{(-3bx+dx-3a+c)}}{64(3b-d)}$$

$$- \frac{3e^{(-3bx-dx-3a-c)}}{64(3b+d)} - \frac{e^{(-3bx-3dx-3a-3c)}}{192(b+d)}$$

input `integrate(cosh(b*x+a)^3*cosh(d*x+c)^3,x, algorithm="giac")`

output

```
1/192*e^(3*b*x + 3*d*x + 3*a + 3*c)/(b + d) + 3/64*e^(3*b*x + d*x + 3*a +
c)/(3*b + d) + 3/64*e^(3*b*x - d*x + 3*a - c)/(3*b - d) + 1/192*e^(3*b*x -
3*d*x + 3*a - 3*c)/(b - d) + 3/64*e^(b*x + 3*d*x + a + 3*c)/(b + 3*d) + 9
/64*e^(b*x + d*x + a + c)/(b + d) + 9/64*e^(b*x - d*x + a - c)/(b - d) + 3
/64*e^(b*x - 3*d*x + a - 3*c)/(b - 3*d) - 3/64*e^(-b*x + 3*d*x - a + 3*c)/
(b - 3*d) - 9/64*e^(-b*x + d*x - a + c)/(b - d) - 9/64*e^(-b*x - d*x - a -
c)/(b + d) - 3/64*e^(-b*x - 3*d*x - a - 3*c)/(b + 3*d) - 1/192*e^(-3*b*x
+ 3*d*x - 3*a + 3*c)/(b - d) - 3/64*e^(-3*b*x + d*x - 3*a + c)/(3*b - d) -
3/64*e^(-3*b*x - d*x - 3*a - c)/(3*b + d) - 1/192*e^(-3*b*x - 3*d*x - 3*a
- 3*c)/(b + d)
```


Mupad [B] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 908, normalized size of antiderivative = 4.66

$$\begin{aligned}
\int \cosh^3(a + bx) \cosh^3(c + dx) dx = & -e^{3a+c+3bx+dx} \left(\frac{-9b^3 + 3b^2d + 9bd^2 - 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& - \frac{e^{-6a-6bx}(-9b^3 - 3b^2d + 9bd^2 + 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& + \frac{e^{-2a-2bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
& - e^{3a-c+3bx-dx} \left(\frac{-9b^3 - 3b^2d + 9bd^2 + 3d^3}{576b^4 - 640b^2d^2 + 64d^4} \right. \\
& - \frac{e^{-6a-6bx}(-9b^3 + 3b^2d + 9bd^2 - 3d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& + \frac{e^{-2a-2bx}(-81b^3 - 81b^2d + 9bd^2 + 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \\
& \left. - \frac{e^{-4a-4bx}(-81b^3 + 81b^2d + 9bd^2 - 9d^3)}{576b^4 - 640b^2d^2 + 64d^4} \right) \\
& - e^{3a-3c+3bx-3dx} \left(\frac{-b^3 - b^2d + 9bd^2 + 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& - \frac{e^{-6a-6bx}(-b^3 + b^2d + 9bd^2 - 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& + \frac{e^{-2a-2bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right) \\
& - e^{3a+3c+3bx+3dx} \left(\frac{-b^3 + b^2d + 9bd^2 - 9d^3}{192b^4 - 1920b^2d^2 + 1728d^4} \right. \\
& - \frac{e^{-6a-6bx}(-b^3 - b^2d + 9bd^2 + 9d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& + \frac{e^{-2a-2bx}(-9b^3 + 27b^2d + 9bd^2 - 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \\
& \left. - \frac{e^{-4a-4bx}(-9b^3 - 27b^2d + 9bd^2 + 27d^3)}{192b^4 - 1920b^2d^2 + 1728d^4} \right)
\end{aligned}$$

input `int(cosh(a + b*x)^3*cosh(c + d*x)^3,x)`

output

```

- exp(3*a + c + 3*b*x + d*x)*((9*b*d^2 + 3*b^2*d - 9*b^3 - 3*d^3)/(576*b^4
+ 64*d^4 - 640*b^2*d^2) - (exp(- 6*a - 6*b*x)*(9*b*d^2 - 3*b^2*d - 9*b^3
+ 3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) + (exp(- 2*a - 2*b*x)*(9*b*d^2
+ 81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 4*
a - 4*b*x)*(9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*
b^2*d^2)) - exp(3*a - c + 3*b*x - d*x)*((9*b*d^2 - 3*b^2*d - 9*b^3 + 3*d^3
)/(576*b^4 + 64*d^4 - 640*b^2*d^2) - (exp(- 6*a - 6*b*x)*(9*b*d^2 + 3*b^2*
d - 9*b^3 - 3*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) + (exp(- 2*a - 2*b*x)
*(9*b*d^2 - 81*b^2*d - 81*b^3 + 9*d^3))/(576*b^4 + 64*d^4 - 640*b^2*d^2) -
(exp(- 4*a - 4*b*x)*(9*b*d^2 + 81*b^2*d - 81*b^3 - 9*d^3))/(576*b^4 + 64*
d^4 - 640*b^2*d^2)) - exp(3*a - 3*c + 3*b*x - 3*d*x)*((9*b*d^2 - b^2*d - b
^3 + 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(- 6*a - 6*b*x)*(9*b
*d^2 + b^2*d - b^3 - 9*d^3))/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) + (exp(-
2*a - 2*b*x)*(9*b*d^2 - 27*b^2*d - 9*b^3 + 27*d^3))/(192*b^4 + 1728*d^4 -
1920*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))
/(192*b^4 + 1728*d^4 - 1920*b^2*d^2)) - exp(3*a + 3*c + 3*b*x + 3*d*x)*((9
*b*d^2 + b^2*d - b^3 - 9*d^3)/(192*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(-
6*a - 6*b*x)*(9*b*d^2 - b^2*d - b^3 + 9*d^3))/(192*b^4 + 1728*d^4 - 1920*
b^2*d^2) + (exp(- 2*a - 2*b*x)*(9*b*d^2 + 27*b^2*d - 9*b^3 - 27*d^3))/(192
*b^4 + 1728*d^4 - 1920*b^2*d^2) - (exp(- 4*a - 4*b*x)*(9*b*d^2 - 27*b^2...

```

Reduce [F]

$$\int \cosh^3(a + bx) \cosh^3(c + dx) dx = \int \cosh(bx + a)^3 \cosh(dx + c)^3 dx$$

input

```
int(cosh(b*x+a)^3*cosh(d*x+c)^3,x)
```

output

```
int(cosh(b*x+a)^3*cosh(d*x+c)^3,x)
```

3.103 $\int \cosh(x) \tanh(2x) dx$

Optimal result	730
Mathematica [C] (verified)	730
Rubi [A] (verified)	731
Maple [B] (verified)	733
Fricas [B] (verification not implemented)	733
Sympy [F]	734
Maxima [B] (verification not implemented)	734
Giac [B] (verification not implemented)	734
Mupad [B] (verification not implemented)	735
Reduce [B] (verification not implemented)	735

Optimal result

Integrand size = 7, antiderivative size = 19

$$\int \cosh(x) \tanh(2x) dx = -\frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{\sqrt{2}} + \cosh(x)$$

output

```
-1/2*arctanh(2^(1/2)*cosh(x))*2^(1/2)+cosh(x)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.68

$$\int \cosh(x) \tanh(2x) dx = -\frac{\operatorname{arctanh}(\sqrt{2} - i \tanh(\frac{x}{2}))}{\sqrt{2}} - \frac{\operatorname{arctanh}(\sqrt{2} + i \tanh(\frac{x}{2}))}{\sqrt{2}} + \cosh(x)$$

input

```
Integrate[Cosh[x]*Tanh[2*x],x]
```

output

$$-(\text{ArcTanh}[\text{Sqrt}[2] - \text{I}*\text{Tanh}[x/2]]/\text{Sqrt}[2]) - \text{ArcTanh}[\text{Sqrt}[2] + \text{I}*\text{Tanh}[x/2]]/\text{Sqrt}[2] + \text{Cosh}[x]$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 262, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(x) \tanh(2x) dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{i \tan(2ix)}{\sec(ix)} dx \\ & \quad \downarrow \text{26} \\ & -i \int \frac{\tan(2ix)}{\sec(ix)} dx \\ & \quad \downarrow \text{4879} \\ & \int -\frac{2 \cosh^2(x)}{1 - 2 \cosh^2(x)} d \cosh(x) \\ & \quad \downarrow \text{27} \\ & -2 \int \frac{\cosh^2(x)}{1 - 2 \cosh^2(x)} d \cosh(x) \\ & \quad \downarrow \text{262} \\ & -2 \left(\frac{1}{2} \int \frac{1}{1 - 2 \cosh^2(x)} d \cosh(x) - \frac{\cosh(x)}{2} \right) \\ & \quad \downarrow \text{219} \\ & -2 \left(\frac{\text{arctanh}(\sqrt{2} \cosh(x))}{2\sqrt{2}} - \frac{\cosh(x)}{2} \right) \end{aligned}$$

input `Int[Cosh[x]*Tanh[2*x],x]`

output `-2*(ArcTanh[Sqrt[2]*Cosh[x]]/(2*Sqrt[2]) - Cosh[x]/2)`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 262 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(15) = 30$.

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.58

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{4} - \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{4}$	49

input `int(cosh(x)*tanh(2*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+1/4*2^(1/2)*ln(exp(2*x)-2^(1/2)*exp(x)+1)-1/4*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(15) = 30$.

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 3.84

$$\int \cosh(x) \tanh(2x) dx$$

$$= \frac{2 \cosh(x)^2 + (\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(\frac{\cosh(x)^2 + \sinh(x)^2 - 2\sqrt{2} \cosh(x) + 2}{\cosh(x)^2 + \sinh(x)^2}\right) + 4 \cosh(x) \sinh(x) + 2 \sinh(x)^2}{4(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)*tanh(2*x),x, algorithm="fricas")`

output `1/4*(2*cosh(x)^2 + (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log((cosh(x)^2 + sinh(x)^2 - 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2 + sinh(x)^2)) + 4*cosh(x)*sinh(x) + 2*sinh(x)^2 + 2)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \cosh(x) \tanh(2x) dx = \int \cosh(x) \tanh(2x) dx$$

input `integrate(cosh(x)*tanh(2*x),x)`

output `Integral(cosh(x)*tanh(2*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(15) = 30.

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.74

$$\int \cosh(x) \tanh(2x) dx = -\frac{1}{4} \sqrt{2} \log \left(\sqrt{2} e^{(-x)} + e^{(-2x)} + 1 \right) \\ + \frac{1}{4} \sqrt{2} \log \left(-\sqrt{2} e^{(-x)} + e^{(-2x)} + 1 \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(cosh(x)*tanh(2*x),x, algorithm="maxima")`

output `-1/4*sqrt(2)*log(sqrt(2)*e^(-x) + e^(-2*x) + 1) + 1/4*sqrt(2)*log(-sqrt(2)*e^(-x) + e^(-2*x) + 1) + 1/2*e^(-x) + 1/2*e^x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(15) = 30.

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.37

$$\int \cosh(x) \tanh(2x) dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{\sqrt{2} - e^{(-x)} - e^x}{\sqrt{2} + e^{(-x)} + e^x} \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(cosh(x)*tanh(2*x),x, algorithm="giac")`

output $\frac{1}{4}\sqrt{2}\log(-\sqrt{2} - e^{-x} - e^x)/(\sqrt{2} + e^{-x} + e^x) + \frac{1}{2}e^{-x} + \frac{1}{2}e^x$

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.53

$$\int \cosh(x) \tanh(2x) dx = \frac{e^{-x}}{2} + \frac{e^x}{2} - \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{4} + \frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{4}$$

input `int(tanh(2*x)*cosh(x),x)`

output $\frac{\exp(-x)/2 + \exp(x)/2 - (2^{(1/2)}\log(\exp(2*x) + 2^{(1/2)}\exp(x) + 1))/4 + (2^{(1/2)}\log(\exp(2*x) - 2^{(1/2)}\exp(x) + 1))/4}$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 58, normalized size of antiderivative = 3.05

$$\int \cosh(x) \tanh(2x) dx = \frac{2e^{2x} + e^x\sqrt{2}\log(e^{2x} - e^x\sqrt{2} + 1) - e^x\sqrt{2}\log(e^{2x} + e^x\sqrt{2} + 1) + 2}{4e^x}$$

input `int(cosh(x)*tanh(2*x),x)`

output $(2*e^{2*x} + e^{*x}\sqrt{2}*\log(e^{2*x} - e^{*x}\sqrt{2} + 1) - e^{*x}\sqrt{2}*\log(e^{2*x} + e^{*x}\sqrt{2} + 1) + 2)/(4*e^{*x})$

3.104 $\int \cosh(x) \tanh(3x) dx$

Optimal result	736
Mathematica [C] (verified)	736
Rubi [A] (verified)	737
Maple [B] (verified)	738
Fricas [B] (verification not implemented)	739
Sympy [F]	739
Maxima [B] (verification not implemented)	740
Giac [B] (verification not implemented)	740
Mupad [B] (verification not implemented)	741
Reduce [B] (verification not implemented)	741

Optimal result

Integrand size = 7, antiderivative size = 20

$$\int \cosh(x) \tanh(3x) dx = -\frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{\sqrt{3}} + \cosh(x)$$

output `-1/3*arctanh(2/3*cosh(x)*3^(1/2))*3^(1/2)+cosh(x)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \cosh(x) \tanh(3x) dx = -\frac{\operatorname{arctanh}\left(\frac{2-i \tanh\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{2+i \tanh\left(\frac{x}{2}\right)}{\sqrt{3}}\right)}{\sqrt{3}} + \cosh(x)$$

input `Integrate[Cosh[x]*Tanh[3*x],x]`

output `-(ArcTanh[(2 - I*Tanh[x/2])/Sqrt[3]]/Sqrt[3]) - ArcTanh[(2 + I*Tanh[x/2])/Sqrt[3]]/Sqrt[3] + Cosh[x]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4879, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \tanh(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(3ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(3ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{4879} \\
 & \int \frac{1 - 4 \cosh^2(x)}{3 - 4 \cosh^2(x)} d \cosh(x) \\
 & \quad \downarrow \text{299} \\
 & \cosh(x) - 2 \int \frac{1}{3 - 4 \cosh^2(x)} d \cosh(x) \\
 & \quad \downarrow \text{219} \\
 & \cosh(x) - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{\sqrt{3}}
 \end{aligned}$$

input `Int [Cosh [x] *Tanh [3*x] , x]`

output `-(ArcTanh [(2*Cosh [x])/Sqrt [3]]/Sqrt [3]) + Cosh [x]`

Definitions of rubi rules used

rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_{x_}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 219 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 299 $\text{Int}[(a_ + (b_)*(x_)^2)^{p_}*((c_ + (d_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^2)^{p+1}/(b*(2*p+3))), x] - \text{Simp}[(a*d - b*c*(2*p+3))/(b*(2*p+3)) \text{Int}[(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[2*p+3, 0]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4879 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}\{d = \text{FreeFactors}[\text{Cos}[v], x]\}, -d/\text{Coefficient}[v, x, 1] \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Cos}[v]/d, u/\text{Sin}[v], x], x], x, \text{Cos}[v]/d]], x] /; \text{!FalseQ}[v] \ \&\& \ \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Cos}[v], x], u/\text{Sin}[v], x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(16) = 32$.

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.45

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\sqrt{3} \ln(e^{2x} - \sqrt{3}e^x + 1)}{6} - \frac{\sqrt{3} \ln(e^{2x} + \sqrt{3}e^x + 1)}{6}$	49

input `int(cosh(x)*tanh(3*x),x,method=_RETURNVERBOSE)`

output

```
1/2*exp(x)+1/2*exp(-x)+1/6*3^(1/2)*ln(exp(2*x)-3^(1/2)*exp(x)+1)-1/6*3^(1/2)*ln(exp(2*x)+3^(1/2)*exp(x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(16) = 32$.

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 4.10

$$\int \cosh(x) \tanh(3x) dx$$

$$= \frac{3 \cosh(x)^2 + (\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)) \log\left(\frac{2 \cosh(x)^2 + 2 \sinh(x)^2 - 4\sqrt{3} \cosh(x) + 5}{2 \cosh(x)^2 + 2 \sinh(x)^2 - 1}\right) + 6 \cosh(x) \sinh(x) + 3}{6 (\cosh(x) + \sinh(x))}$$

input

```
integrate(cosh(x)*tanh(3*x),x, algorithm="fricas")
```

output

```
1/6*(3*cosh(x)^2 + (sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*log((2*cosh(x)^2 + 2*sinh(x)^2 - 4*sqrt(3)*cosh(x) + 5)/(2*cosh(x)^2 + 2*sinh(x)^2 - 1)) + 6*cosh(x)*sinh(x) + 3*sinh(x)^2 + 3)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \cosh(x) \tanh(3x) dx = \int \cosh(x) \tanh(3x) dx$$

input

```
integrate(cosh(x)*tanh(3*x),x)
```

output

```
Integral(cosh(x)*tanh(3*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(16) = 32$.

Time = 0.14 (sec) , antiderivative size = 153, normalized size of antiderivative = 7.65

$$\begin{aligned} \int \cosh(x) \tanh(3x) dx = & -\frac{1}{12} \sqrt{3} \log \left(\sqrt{3} e^{(-x)} + e^{(-2x)} + 1 \right) \\ & + \frac{1}{12} \sqrt{3} \log \left(-\sqrt{3} e^{(-x)} + e^{(-2x)} + 1 \right) \\ & - \frac{1}{12} \sqrt{3} \log \left(\sqrt{3} e^x + e^{(2x)} + 1 \right) \\ & + \frac{1}{12} \sqrt{3} \log \left(-\sqrt{3} e^x + e^{(2x)} + 1 \right) \\ & + \frac{1}{6} \arctan \left(\sqrt{3} + 2 e^{(-x)} \right) + \frac{1}{6} \arctan \left(\sqrt{3} + 2 e^x \right) \\ & + \frac{1}{6} \arctan \left(-\sqrt{3} + 2 e^{(-x)} \right) + \frac{1}{6} \arctan \left(-\sqrt{3} + 2 e^x \right) \\ & + \frac{1}{3} \arctan \left(e^{(-x)} \right) + \frac{1}{3} \arctan \left(e^x \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(cosh(x)*tanh(3*x),x, algorithm="maxima")`

output `-1/12*sqrt(3)*log(sqrt(3)*e^(-x) + e^(-2*x) + 1) + 1/12*sqrt(3)*log(-sqrt(3)*e^(-x) + e^(-2*x) + 1) - 1/12*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/12*sqrt(3)*log(-sqrt(3)*e^x + e^(2*x) + 1) + 1/6*arctan(sqrt(3) + 2*e^(-x)) + 1/6*arctan(sqrt(3) + 2*e^x) + 1/6*arctan(-sqrt(3) + 2*e^(-x)) + 1/6*arctan(-sqrt(3) + 2*e^x) + 1/3*arctan(e^(-x)) + 1/3*arctan(e^x) + 1/2*e^(-x) + 1/2*e^x`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(16) = 32$.

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.25

$$\int \cosh(x) \tanh(3x) dx = \frac{1}{6} \sqrt{3} \log \left(-\frac{\sqrt{3} - e^{(-x)} - e^x}{\sqrt{3} + e^{(-x)} + e^x} \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(cosh(x)*tanh(3*x),x, algorithm="giac")`

output $\frac{1}{6}\sqrt{3}\log(-\sqrt{3} - e^{-x} - e^x)/(\sqrt{3} + e^{-x} + e^x) + \frac{1}{2}e^{-x} + \frac{1}{2}e^x$

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.65

$$\int \cosh(x) \tanh(3x) dx = \frac{e^{-x}}{2} + \frac{e^x}{2} + \frac{\sqrt{3} \ln\left(\frac{e^{2x}}{3} - \frac{\sqrt{3}e^x}{3} + \frac{1}{3}\right)}{6} - \frac{\sqrt{3} \ln\left(\frac{e^{2x}}{3} + \frac{\sqrt{3}e^x}{3} + \frac{1}{3}\right)}{6}$$

input `int(tanh(3*x)*cosh(x),x)`

output $\frac{\exp(-x)/2 + \exp(x)/2 + (3^{(1/2)}\log(\exp(2*x)/3 - (3^{(1/2)}\exp(x))/3 + 1/3))}{6} - (3^{(1/2)}\log(\exp(2*x)/3 + (3^{(1/2)}\exp(x))/3 + 1/3))/6$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.90

$$\int \cosh(x) \tanh(3x) dx = \frac{3e^{2x} + e^x\sqrt{3}\log(e^{2x} - e^x\sqrt{3} + 1) - e^x\sqrt{3}\log(e^{2x} + e^x\sqrt{3} + 1) + 3}{6e^x}$$

input `int(cosh(x)*tanh(3*x),x)`

output $(3e^{2x} + e^x\sqrt{3}\log(e^{2x} - e^x\sqrt{3} + 1) - e^x\sqrt{3}\log(e^{2x} + e^x\sqrt{3} + 1) + 3)/(6e^x)$

3.105 $\int \cosh(x) \tanh(4x) dx$

Optimal result	742
Mathematica [C] (verified)	742
Rubi [A] (verified)	743
Maple [C] (verified)	745
Fricas [B] (verification not implemented)	746
Sympy [F]	746
Maxima [F]	747
Giac [B] (verification not implemented)	747
Mupad [B] (verification not implemented)	748
Reduce [B] (verification not implemented)	748

Optimal result

Integrand size = 7, antiderivative size = 75

$$\int \cosh(x) \tanh(4x) dx = -\frac{\operatorname{arctanh}\left(\sqrt{2(2-\sqrt{2})} \cosh(x)\right)}{2\sqrt{2(2-\sqrt{2})}} - \frac{\operatorname{arctanh}\left(\sqrt{2(2+\sqrt{2})} \cosh(x)\right)}{2\sqrt{2(2+\sqrt{2})}} + \cosh(x)$$

output

$$-1/2*\operatorname{arctanh}((4-2*2^{(1/2)})^{(1/2)}*\cosh(x))/(4-2*2^{(1/2)})^{(1/2)}-1/2*\operatorname{arctanh}((4+2*2^{(1/2)})^{(1/2)}*\cosh(x))/(4+2*2^{(1/2)})^{(1/2)}+\cosh(x)$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.03 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.51

$$\int \cosh(x) \tanh(4x) dx = \cosh(x) + \frac{1}{16} \operatorname{RootSum}\left[1 + \#1^8 \&, \frac{-x - 2 \log\left(-\cosh\left(\frac{x}{2}\right) - \sinh\left(\frac{x}{2}\right) + \cosh\left(\frac{x}{2}\right) \#1 - \sinh\left(\frac{x}{2}\right) \#1\right) + x \#1^6 + 2 \log\left(-\cosh\left(\frac{x}{2}\right)\right)}{\#1^7}\right]$$

input `Integrate[Cosh[x]*Tanh[4*x],x]`

output `Cosh[x] + RootSum[1 + #1^8 & , (-x - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1] + x*#1^6 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/#1^7 &]/16`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 26, 4879, 27, 1602, 27, 1480, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \tanh(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(4ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(4ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{4879} \\
 & \int -\frac{4 \cosh^2(x) (1 - 2 \cosh^2(x))}{8 \cosh^4(x) - 8 \cosh^2(x) + 1} d \cosh(x) \\
 & \quad \downarrow \text{27} \\
 & -4 \int \frac{\cosh^2(x) (1 - 2 \cosh^2(x))}{8 \cosh^4(x) - 8 \cosh^2(x) + 1} d \cosh(x) \\
 & \quad \downarrow \text{1602} \\
 & -4 \left(-\frac{1}{8} \int -\frac{2(1 - 4 \cosh^2(x))}{8 \cosh^4(x) - 8 \cosh^2(x) + 1} d \cosh(x) - \frac{\cosh(x)}{4} \right)
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& -4 \left(\frac{1}{4} \int \frac{1 - 4 \cosh^2(x)}{8 \cosh^4(x) - 8 \cosh^2(x) + 1} d \cosh(x) - \frac{\cosh(x)}{4} \right) \\
& \downarrow 1480 \\
& -4 \left(\frac{1}{4} \left(- \left((2 - \sqrt{2}) \int \frac{1}{8 \cosh^2(x) - 2(2 - \sqrt{2})} d \cosh(x) \right) - (2 + \sqrt{2}) \int \frac{1}{8 \cosh^2(x) - 2(2 + \sqrt{2})} d \cosh(x) \right) \right) \\
& \downarrow 220 \\
& -4 \left(\frac{1}{4} \left(\frac{1}{4} \sqrt{2 - \sqrt{2}} \operatorname{arctanh} \left(\frac{2 \cosh(x)}{\sqrt{2 - \sqrt{2}}} \right) + \frac{1}{4} \sqrt{2 + \sqrt{2}} \operatorname{arctanh} \left(\frac{2 \cosh(x)}{\sqrt{2 + \sqrt{2}}} \right) \right) - \frac{\cosh(x)}{4} \right)
\end{aligned}$$

input `Int [Cosh[x]*Tanh[4*x], x]`

output `-4*(((Sqrt[2 - Sqrt[2]]*ArcTanh[(2*Cosh[x])/Sqrt[2 - Sqrt[2]]])/4 + (Sqrt[2 + Sqrt[2]]*ArcTanh[(2*Cosh[x])/Sqrt[2 + Sqrt[2]]])/4)/4 - Cosh[x]/4)`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 1480

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :
> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(e/2 + (2*c*d - b*e)/(2*q)) Int[1/(
b/2 - q/2 + c*x^2), x], x] + Simp[(e/2 - (2*c*d - b*e)/(2*q)) Int[1/(b/2
+ q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0]
&& NeQ[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]
```

rule 1602

```
Int(((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(
x_)^4)^(p_), x_Symbol] := Simp[e*f*(f*x)^(m - 1)*((a + b*x^2 + c*x^4)^(p +
1)/(c*(m + 4*p + 3))), x] - Simp[f^2/(c*(m + 4*p + 3)) Int[(f*x)^(m - 2)*
(a + b*x^2 + c*x^4)^p*Simp[a*e*(m - 1) + (b*e*(m + 2*p + 1) - c*d*(m + 4*p
+ 3))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && NeQ[b^2 - 4*a*c
, 0] && GtQ[m, 1] && NeQ[m + 4*p + 3, 0] && IntegerQ[2*p] && (IntegerQ[p] |
IntegerQ[m])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4879

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d
, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[Nonfree
Factors[Cos[v], x], u/Sin[v], x]]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.56

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \left(\sum_{R=\text{RootOf}(2048_Z^4-128_Z^2+1)} -R \ln(-8_R e^x + e^{2x} + 1) \right)$	42

input

```
int(cosh(x)*tanh(4*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)+1/2*exp(-x)+sum(_R*ln(-8*_R*exp(x)+exp(2*x)+1),_R=RootOf(2048*_Z^4-128*_Z^2+1))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. $2(51) = 102$.

Time = 0.10 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.84

$$\int \cosh(x) \tanh(4x) dx = \frac{\sqrt{\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + \sqrt{\sqrt{2} + 2}(\cosh(x) + \sinh(x))) - \sqrt{\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - \sqrt{\sqrt{2} + 2}(\cosh(x) + \sinh(x))) + \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x))) - \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x)) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - \sqrt{-\sqrt{2} + 2}(\cosh(x) + \sinh(x))) - 4 \cosh(x)^2 - 8 \cosh(x) \sinh(x) - 4 \sinh(x)^2 - 4}{(\cosh(x) + \sinh(x))^2}$$

input

```
integrate(cosh(x)*tanh(4*x),x, algorithm="fricas")
```

output

```
-1/8*(sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x)) + 1) - sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(sqrt(2) + 2)*(cosh(x) + sinh(x)) + 1) + sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x)) + 1) - sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(-sqrt(2) + 2)*(cosh(x) + sinh(x)) + 1) - 4*cosh(x)^2 - 8*cosh(x)*sinh(x) - 4*sinh(x)^2 - 4)/(cosh(x) + sinh(x))^2
```

Sympy [F]

$$\int \cosh(x) \tanh(4x) dx = \int \cosh(x) \tanh(4x) dx$$

input

```
integrate(cosh(x)*tanh(4*x),x)
```

output

```
Integral(cosh(x)*tanh(4*x), x)
```

Maxima [F]

$$\int \cosh(x) \tanh(4x) dx = \int \cosh(x) \tanh(4x) dx$$

input `integrate(cosh(x)*tanh(4*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) + 1)*e^(-x) + 1/2*integrate(2*(e^(7*x) - e^x)/(e^(8*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(51) = 102$.

Time = 0.19 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.59

$$\begin{aligned} \int \cosh(x) \tanh(4x) dx = & -\frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(\sqrt{\sqrt{2} + 2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{8} \sqrt{\sqrt{2} + 2} \log \left(-\sqrt{\sqrt{2} + 2} + e^{(-x)} + e^x \right) \\ & - \frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(\sqrt{-\sqrt{2} + 2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{8} \sqrt{-\sqrt{2} + 2} \log \left(-\sqrt{-\sqrt{2} + 2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(cosh(x)*tanh(4*x),x, algorithm="giac")`

output `-1/8*sqrt(sqrt(2) + 2)*log(sqrt(sqrt(2) + 2) + e^(-x) + e^x) + 1/8*sqrt(sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2) + e^(-x) + e^x) - 1/8*sqrt(-sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2) + e^(-x) + e^x) + 1/8*sqrt(-sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2) + e^(-x) + e^x) + 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.77

$$\int \cosh(x) \tanh(4x) dx = \frac{e^{-x}}{2} + \frac{e^x}{2} + \ln \left(e^{2x} - 8e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64} + 1} \right) \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}}$$

$$- \ln \left(e^{2x} + 8e^x \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64} + 1} \right) \sqrt{\frac{1}{32} - \frac{\sqrt{2}}{64}}$$

$$+ \ln \left(e^{2x} - 8e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32} + 1} \right) \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}}$$

$$- \ln \left(e^{2x} + 8e^x \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32} + 1} \right) \sqrt{\frac{\sqrt{2}}{64} + \frac{1}{32}}$$

input `int(tanh(4*x)*cosh(x), x)`output `exp(-x)/2 + exp(x)/2 + log(exp(2*x) - 8*exp(x)*(1/32 - 2^(1/2)/64)^(1/2) + 1)*(1/32 - 2^(1/2)/64)^(1/2) - log(exp(2*x) + 8*exp(x)*(1/32 - 2^(1/2)/64)^(1/2) + 1)*(1/32 - 2^(1/2)/64)^(1/2) + log(exp(2*x) - 8*exp(x)*(2^(1/2)/64 + 1/32)^(1/2) + 1)*(2^(1/2)/64 + 1/32)^(1/2) - log(exp(2*x) + 8*exp(x)*(2^(1/2)/64 + 1/32)^(1/2) + 1)*(2^(1/2)/64 + 1/32)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.76

$$\int \cosh(x) \tanh(4x) dx$$

$$= \frac{e^x \sqrt{-\sqrt{2} + 2} \log\left(-e^x \sqrt{-\sqrt{2} + 2} + e^{2x} + 1\right) - e^x \sqrt{-\sqrt{2} + 2} \log\left(e^x \sqrt{-\sqrt{2} + 2} + e^{2x} + 1\right) + e^x \sqrt{\sqrt{2}}}{8e^x}$$

input `int(cosh(x)*tanh(4*x), x)`

output

```
(e**x*sqrt(-sqrt(2)+2)*log(-e**x*sqrt(-sqrt(2)+2)+e**(2*x)+1) - e**x*sqrt(-sqrt(2)+2)*log(e**x*sqrt(-sqrt(2)+2)+e**(2*x)+1) + e**x*sqrt(sqrt(2)+2)*log(-e**x*sqrt(sqrt(2)+2)+e**(2*x)+1) - e**x*sqrt(sqrt(2)+2)*log(e**x*sqrt(sqrt(2)+2)+e**(2*x)+1) + 4*e**(2*x)+4)/(8*e**x)
```

3.106 $\int \cosh(x) \tanh(5x) dx$

Optimal result	750
Mathematica [C] (verified)	750
Rubi [A] (verified)	751
Maple [C] (verified)	753
Fricas [B] (verification not implemented)	753
Sympy [F]	754
Maxima [F]	754
Giac [B] (verification not implemented)	755
Mupad [B] (verification not implemented)	756
Reduce [F]	756

Optimal result

Integrand size = 7, antiderivative size = 81

$$\int \cosh(x) \tanh(5x) dx = -\frac{1}{5} \sqrt{\frac{1}{2} (5 + \sqrt{5})} \operatorname{arctanh} \left(\sqrt{\frac{2}{5} (5 - \sqrt{5})} \cosh(x) \right) - \sqrt{\frac{2}{5 (5 + \sqrt{5})}} \operatorname{arctanh} \left(\sqrt{\frac{2}{5} (5 + \sqrt{5})} \cosh(x) \right) + \cosh(x)$$

output

```
-1/10*(10+2*5^(1/2))^(1/2)*arctanh(1/5*(50-10*5^(1/2))^(1/2)*cosh(x))-2^(1/2)/(25+5*5^(1/2))^(1/2)*arctanh(1/5*(50+10*5^(1/2))^(1/2)*cosh(x))+cosh(x)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.03 (sec) , antiderivative size = 249, normalized size of antiderivative = 3.07

$$\int \cosh(x) \tanh(5x) dx = \cosh(x) + \frac{1}{4} \operatorname{RootSum} \left[1 - \#1^2 + \#1^4 - \#1^6 + \#1^8 \&, \frac{-x - 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) + x \#1^2 + 2 \log \left(-\cosh \left(\frac{x}{2} \right) \right)}{\dots} \right]$$

input `Integrate[Cosh[x]*Tanh[5*x],x]`

output `Cosh[x] + RootSum[1 - #1^2 + #1^4 - #1^6 + #1^8 & , (-x - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1] + x*#1^2 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^2 - x*#1^4 - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^4 + x*#1^6 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/(-#1 + 2*#1^3 - 3*#1^5 + 4*#1^7) &]/4`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4879, 2205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \tanh(5x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{i \tan(5ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{26} \\
 & -i \int \frac{\tan(5ix)}{\sec(ix)} dx \\
 & \quad \downarrow \text{4879} \\
 & \int \frac{16 \cosh^4(x) - 12 \cosh^2(x) + 1}{16 \cosh^4(x) - 20 \cosh^2(x) + 5} d \cosh(x) \\
 & \quad \downarrow \text{2205} \\
 & \int \left(1 - \frac{4(1 - 2 \cosh^2(x))}{16 \cosh^4(x) - 20 \cosh^2(x) + 5} \right) d \cosh(x) \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$-\frac{1}{5}\sqrt{\frac{1}{2}(5+\sqrt{5})}\operatorname{arctanh}\left(2\sqrt{\frac{2}{5+\sqrt{5}}}\cosh(x)\right) - \frac{1}{5}\sqrt{\frac{1}{2}(5-\sqrt{5})}\operatorname{arctanh}\left(\sqrt{\frac{2}{5}}(5+\sqrt{5})\cosh(x)\right) + \cosh(x)$$

input `Int[Cosh[x]*Tanh[5*x],x]`

output `-1/5*(Sqrt[(5 + Sqrt[5])/2]*ArcTanh[2*Sqrt[2/(5 + Sqrt[5])]*Cosh[x]]) - (Sqrt[(5 - Sqrt[5])/2]*ArcTanh[Sqrt[(2*(5 + Sqrt[5]))/5]*Cosh[x]])/5 + Cosh[x]`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2205 `Int[(Px_)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := Int[ExpandIntegrand[Px/(a + b*x^2 + c*x^4), x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.52

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \left(\sum_{_R=\text{RootOf}(2000_Z^4-100_Z^2+1)} _R \ln(-10_R e^x + e^{2x} + 1) \right)$	42

input `int(cosh(x)*tanh(5*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+sum(_R*ln(-10*_R*exp(x)+exp(2*x)+1),_R=RootOf(2000*_Z^4-100*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(56) = 112$.

Time = 0.10 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.73

$$\int \cosh(x) \tanh(5x) dx =$$

$$\frac{\sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}}(\cosh(x) + \sinh(x)) \log\left(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + \sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}}(\cosh(x) + \sinh(x))\right)}{\sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}}}$$

input `integrate(cosh(x)*tanh(5*x),x, algorithm="fricas")`

output

```
-1/10*(sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x)) + 1) - sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x)) + 1) + sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x)) + 1) - sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(-1/2*sqrt(5) + 5/2)*(cosh(x) + sinh(x)) + 1) - 5*cosh(x)^2 - 10*cosh(x)*sinh(x) - 5*sinh(x)^2 - 5)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \cosh(x) \tanh(5x) dx = \int \cosh(x) \tanh(5x) dx$$

input

```
integrate(cosh(x)*tanh(5*x),x)
```

output

```
Integral(cosh(x)*tanh(5*x), x)
```

Maxima [F]

$$\int \cosh(x) \tanh(5x) dx = \int \cosh(x) \tanh(5x) dx$$

input

```
integrate(cosh(x)*tanh(5*x),x, algorithm="maxima")
```

output

```
1/2*(e^(2*x) + 1)*e^(-x) + 1/2*integrate(2*(e^(7*x) - e^(5*x) + e^(3*x) - e^x)/(e^(8*x) - e^(6*x) + e^(4*x) - e^(2*x) + 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. $2(56) = 112$.

Time = 0.15 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.57

$$\begin{aligned} \int \cosh(x) \tanh(5x) dx = & -\frac{1}{20} \sqrt{2\sqrt{5} + 10} \log \left(\sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}} + e^{(-x)} + e^x \right) \\ & + \frac{1}{20} \sqrt{2\sqrt{5} + 10} \log \left(-\sqrt{\frac{1}{2}\sqrt{5} + \frac{5}{2}} + e^{(-x)} + e^x \right) \\ & - \frac{1}{20} \sqrt{-2\sqrt{5} + 10} \log \left(\sqrt{-\frac{1}{2}\sqrt{5} + \frac{5}{2}} + e^{(-x)} + e^x \right) \\ & + \frac{1}{20} \sqrt{-2\sqrt{5} + 10} \log \left(-\sqrt{-\frac{1}{2}\sqrt{5} + \frac{5}{2}} + e^{(-x)} + e^x \right) \\ & + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(cosh(x)*tanh(5*x),x, algorithm="giac")`

output `-1/20*sqrt(2*sqrt(5) + 10)*log(sqrt(1/2*sqrt(5) + 5/2) + e^(-x) + e^x) + 1/20*sqrt(2*sqrt(5) + 10)*log(-sqrt(1/2*sqrt(5) + 5/2) + e^(-x) + e^x) - 1/20*sqrt(-2*sqrt(5) + 10)*log(sqrt(-1/2*sqrt(5) + 5/2) + e^(-x) + e^x) + 1/20*sqrt(-2*sqrt(5) + 10)*log(-sqrt(-1/2*sqrt(5) + 5/2) + e^(-x) + e^x) + 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.74

$$\int \cosh(x) \tanh(5x) dx = \frac{e^{-x}}{2} + \frac{e^x}{2} + \ln \left(4e^{2x} - 40e^x \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} + 4 \right) \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}}$$

$$- \ln \left(4e^{2x} + 40e^x \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}} + 4 \right) \sqrt{\frac{1}{40} - \frac{\sqrt{5}}{200}}$$

$$+ \ln \left(4e^{2x} - 40e^x \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}} + 4 \right) \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}}$$

$$- \ln \left(4e^{2x} + 40e^x \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}} + 4 \right) \sqrt{\frac{\sqrt{5}}{200} + \frac{1}{40}}$$

input `int(tanh(5*x)*cosh(x),x)`output `exp(-x)/2 + exp(x)/2 + log(4*exp(2*x) - 40*exp(x)*(1/40 - 5^(1/2)/200)^(1/2) + 4)*(1/40 - 5^(1/2)/200)^(1/2) - log(4*exp(2*x) + 40*exp(x)*(1/40 - 5^(1/2)/200)^(1/2) + 4)*(1/40 - 5^(1/2)/200)^(1/2) + log(4*exp(2*x) - 40*exp(x)*(5^(1/2)/200 + 1/40)^(1/2) + 4)*(5^(1/2)/200 + 1/40)^(1/2) - log(4*exp(2*x) + 40*exp(x)*(5^(1/2)/200 + 1/40)^(1/2) + 4)*(5^(1/2)/200 + 1/40)^(1/2)`**Reduce [F]**

$$\int \cosh(x) \tanh(5x) dx = \frac{e^{2x} - 2e^x \left(\int \frac{1}{e^{9x} - e^{7x} + e^{5x} - e^{3x} + e^x} dx \right) - 1}{2e^x}$$

input `int(cosh(x)*tanh(5*x),x)`output `(e**(2*x) - 2*e**x*int(1/(e**(9*x) - e**(7*x) + e**(5*x) - e**(3*x) + e**x),x) - 1)/(2*e**x)`

3.107 $\int \cosh(x) \tanh(6x) dx$

Optimal result	757
Mathematica [C] (verified)	757
Rubi [A] (verified)	758
Maple [C] (verified)	760
Fricas [B] (verification not implemented)	761
Sympy [F]	761
Maxima [F]	762
Giac [B] (verification not implemented)	762
Mupad [B] (verification not implemented)	763
Reduce [B] (verification not implemented)	763

Optimal result

Integrand size = 7, antiderivative size = 87

$$\int \cosh(x) \tanh(6x) dx = -\frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{3\sqrt{2}} - \frac{\operatorname{arctanh}(2\sqrt{2 - \sqrt{3}} \cosh(x))}{6\sqrt{2 - \sqrt{3}}} - \frac{\operatorname{arctanh}(2\sqrt{2 + \sqrt{3}} \cosh(x))}{6\sqrt{2 + \sqrt{3}}} + \cosh(x)$$

output

```
-1/6*arctanh(2^(1/2)*cosh(x))*2^(1/2)-1/6*arctanh(2*(1/2*6^(1/2)-1/2*2^(1/2))*cosh(x))/(1/2*6^(1/2)-1/2*2^(1/2))-1/6*arctanh(2*(1/2*6^(1/2)+1/2*2^(1/2))*cosh(x))/(1/2*6^(1/2)+1/2*2^(1/2))+cosh(x)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 281, normalized size of antiderivative = 3.23

$$\int \cosh(x) \tanh(6x) dx$$

$$= \frac{1}{24} \left(-4 \left(\sqrt{2} \operatorname{arctanh} \left(\sqrt{2} - i \tanh \left(\frac{x}{2} \right) \right) + \sqrt{2} \operatorname{arctanh} \left(\sqrt{2} + i \tanh \left(\frac{x}{2} \right) \right) - 6 \cosh(x) \right) \right. \\ \left. + \operatorname{RootSum} \left[1 - \#1^4 \right. \right. \\ \left. \left. + \#1^8 \&, \frac{-2x - 4 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) - x \#1^2 - 2 \log \left(-\cosh \left(\frac{x}{2} \right) \right)}{\#1^3 + 2 \#1^7} \right] \right)$$

input `Integrate[Cosh[x]*Tanh[6*x],x]`

output `(-4*(Sqrt[2]*ArcTanh[Sqrt[2] - I*Tanh[x/2]] + Sqrt[2]*ArcTanh[Sqrt[2] + I*Tanh[x/2]] - 6*Cosh[x]) + RootSum[1 - #1^4 + #1^8 & , (-2*x - 4*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1] - x*#1^2 - 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^2 + x*#1^4 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^4 + 2*x*#1^6 + 4*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/(-#1^3 + 2*#1^7) &])/24`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \tanh(6x) dx$$

$$\downarrow \text{3042}$$

$$\int -\frac{i \tan(6ix)}{\sec(ix)} dx$$

$$\begin{aligned}
& \downarrow 26 \\
& -i \int \frac{\tan(6ix)}{\sec(ix)} dx \\
& \downarrow 4879 \\
& \int -\frac{2 \cosh^2(x) (16 \cosh^4(x) - 16 \cosh^2(x) + 3)}{-32 \cosh^6(x) + 48 \cosh^4(x) - 18 \cosh^2(x) + 1} d \cosh(x) \\
& \downarrow 27 \\
& -2 \int \frac{\cosh^2(x) (16 \cosh^4(x) - 16 \cosh^2(x) + 3)}{-32 \cosh^6(x) + 48 \cosh^4(x) - 18 \cosh^2(x) + 1} d \cosh(x) \\
& \downarrow 2460 \\
& -2 \int \left(\frac{1 - 8 \cosh^2(x)}{3 (16 \cosh^4(x) - 16 \cosh^2(x) + 1)} - \frac{1}{6 (2 \cosh^2(x) - 1)} - \frac{1}{2} \right) d \cosh(x) \\
& \downarrow 2009 \\
& -2 \left(\frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{6\sqrt{2}} + \frac{1}{12} \sqrt{2 - \sqrt{3}} \operatorname{arctanh} \left(\frac{2 \cosh(x)}{\sqrt{2 - \sqrt{3}}} \right) + \frac{1}{12} \sqrt{2 + \sqrt{3}} \operatorname{arctanh} \left(\frac{2 \cosh(x)}{\sqrt{2 + \sqrt{3}}} \right) - \frac{\cosh(x)}{2} \right)
\end{aligned}$$

input `Int [Cosh [x] *Tanh [6*x] , x]`

output `-2*(ArcTanh[Sqrt[2]*Cosh[x]]/(6*Sqrt[2]) + (Sqrt[2 - Sqrt[3]]*ArcTanh[(2*Cosh[x])/Sqrt[2 - Sqrt[3]]])/12 + (Sqrt[2 + Sqrt[3]]*ArcTanh[(2*Cosh[x])/Sqrt[2 + Sqrt[3]]])/12 - Cosh[x]/2)`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.14 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.91

method	result
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{12} - \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{12} + \left(\sum_{R=\text{RootOf}(20736_Z^4 - 576_Z^2 + 1)} -R \ln(-12 - R) \right)$

input `int(cosh(x)*tanh(6*x), x, method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+1/12*2^(1/2)*ln(exp(2*x)-2^(1/2)*exp(x)+1)-1/12*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)+sum(_R*ln(-12*_R*exp(x)+exp(2*x)+1), _R=RootOf(20736*_Z^4-576*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 258 vs. $2(63) = 126$.

Time = 0.10 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.97

$$\int \cosh(x) \tanh(6x) dx = \frac{\sqrt{\sqrt{3} + 2}(\cosh(x) + \sinh(x)) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + \sqrt{\sqrt{3} + 2}(\cosh(x) + \sinh(x)))}{\dots}$$

input `integrate(cosh(x)*tanh(6*x),x, algorithm="fricas")`

output

```
-1/12*(sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x)) + 1) - sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x)) + 1) + sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x)) + 1) - sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x))*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x)) + 1) - 6*cosh(x)^2 - (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log((cosh(x)^2 + sinh(x)^2 - 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2 + sinh(x)^2)) - 12*cosh(x)*sinh(x) - 6*sinh(x)^2 - 6)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \cosh(x) \tanh(6x) dx = \int \cosh(x) \tanh(6x) dx$$

input `integrate(cosh(x)*tanh(6*x),x)`

output

`Integral(cosh(x)*tanh(6*x), x)`

Maxima [F]

$$\int \cosh(x) \tanh(6x) dx = \int \cosh(x) \tanh(6x) dx$$

input `integrate(cosh(x)*tanh(6*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) + 1)*e^(-x) - 1/12*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/12*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + 1/2*integrate(2/3*(2*e^(7*x) + e^(5*x) - e^(3*x) - 2*e^x)/(e^(8*x) - e^(4*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. 2(63) = 126.

Time = 0.18 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.80

$$\begin{aligned} \int \cosh(x) \tanh(6x) dx = & -\frac{1}{24} (\sqrt{6} + \sqrt{2}) \log \left(\frac{1}{2} \sqrt{6} + \frac{1}{2} \sqrt{2} + e^{(-x)} + e^x \right) \\ & - \frac{1}{24} (\sqrt{6} - \sqrt{2}) \log \left(\frac{1}{2} \sqrt{6} - \frac{1}{2} \sqrt{2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{24} (\sqrt{6} - \sqrt{2}) \log \left(-\frac{1}{2} \sqrt{6} + \frac{1}{2} \sqrt{2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{24} (\sqrt{6} + \sqrt{2}) \log \left(-\frac{1}{2} \sqrt{6} - \frac{1}{2} \sqrt{2} + e^{(-x)} + e^x \right) \\ & + \frac{1}{12} \sqrt{2} \log \left(-\frac{\sqrt{2} - e^{(-x)} - e^x}{\sqrt{2} + e^{(-x)} + e^x} \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \end{aligned}$$

input `integrate(cosh(x)*tanh(6*x),x, algorithm="giac")`

output `-1/24*(sqrt(6) + sqrt(2))*log(1/2*sqrt(6) + 1/2*sqrt(2) + e^(-x) + e^x) - 1/24*(sqrt(6) - sqrt(2))*log(1/2*sqrt(6) - 1/2*sqrt(2) + e^(-x) + e^x) + 1/24*(sqrt(6) - sqrt(2))*log(-1/2*sqrt(6) + 1/2*sqrt(2) + e^(-x) + e^x) + 1/24*(sqrt(6) + sqrt(2))*log(-1/2*sqrt(6) - 1/2*sqrt(2) + e^(-x) + e^x) + 1/12*sqrt(2)*log(-(sqrt(2) - e^(-x) - e^x)/(sqrt(2) + e^(-x) + e^x)) + 1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.95

$$\int \cosh(x) \tanh(6x) dx = \frac{e^{-x}}{2} + \frac{e^x}{2} - \frac{\sqrt{2} \ln(e^{2x} + \sqrt{2}e^x + 1)}{12} + \frac{\sqrt{2} \ln(e^{2x} - \sqrt{2}e^x + 1)}{12}$$

$$+ \ln\left(e^{2x} - 12e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} + 1\right) \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}}$$

$$- \ln\left(e^{2x} + 12e^x \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}} + 1\right) \sqrt{\frac{1}{72} - \frac{\sqrt{3}}{144}}$$

$$+ \ln\left(e^{2x} - 12e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}} + 1\right) \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}}$$

$$- \ln\left(e^{2x} + 12e^x \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}} + 1\right) \sqrt{\frac{\sqrt{3}}{144} + \frac{1}{72}}$$

input `int(tanh(6*x)*cosh(x),x)`output

```
exp(-x)/2 + exp(x)/2 - (2^(1/2)*log(exp(2*x) + 2^(1/2)*exp(x) + 1))/12 + (
2^(1/2)*log(exp(2*x) - 2^(1/2)*exp(x) + 1))/12 + log(exp(2*x) - 12*exp(x)*
(1/72 - 3^(1/2)/144)^(1/2) + 1)*(1/72 - 3^(1/2)/144)^(1/2) - log(exp(2*x)
+ 12*exp(x)*(1/72 - 3^(1/2)/144)^(1/2) + 1)*(1/72 - 3^(1/2)/144)^(1/2) + 1
og(exp(2*x) - 12*exp(x)*(3^(1/2)/144 + 1/72)^(1/2) + 1)*(3^(1/2)/144 + 1/7
2)^(1/2) - log(exp(2*x) + 12*exp(x)*(3^(1/2)/144 + 1/72)^(1/2) + 1)*(3^(1/
2)/144 + 1/72)^(1/2)
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 236, normalized size of antiderivative = 2.71

$$\int \cosh(x) \tanh(6x) dx$$

$$= \frac{2e^x \sqrt{-\sqrt{3}} + 2 \log\left(-e^x \sqrt{-\sqrt{3}} + 2 + e^{2x} + 1\right) - 2e^x \sqrt{-\sqrt{3}} + 2 \log\left(e^x \sqrt{-\sqrt{3}} + 2 + e^{2x} + 1\right) + 12e^{2x}}$$

input `int(cosh(x)*tanh(6*x),x)`

output $(2e^{2x}\sqrt{-\sqrt{3}+2})\log(-e^{2x}\sqrt{-\sqrt{3}+2}+e^{2x}+1) - 2e^{2x}\sqrt{-\sqrt{3}+2}\log(e^{2x}\sqrt{-\sqrt{3}+2}+e^{2x}+1) + 12e^{2x} + e^{2x}\sqrt{6}\log((2e^{2x}-e^{2x}\sqrt{6}-e^{2x}\sqrt{2}+2)/2) - e^{2x}\sqrt{6}\log((2e^{2x}+e^{2x}\sqrt{6}+e^{2x}\sqrt{2}+2)/2) + 2e^{2x}\sqrt{2}\log(e^{2x}-e^{2x}\sqrt{2}+1) - 2e^{2x}\sqrt{2}\log(e^{2x}+e^{2x}\sqrt{2}+1) + e^{2x}\sqrt{2}\log((2e^{2x}-e^{2x}\sqrt{6}-e^{2x}\sqrt{2}+2)/2) - e^{2x}\sqrt{2}\log((2e^{2x}+e^{2x}\sqrt{6}+e^{2x}\sqrt{2}+2)/2) + 12)/(24e^{2x})$

3.108 $\int \cosh(a + bx) \tanh(c + bx) dx$

Optimal result	765
Mathematica [B] (verified)	765
Rubi [A] (verified)	766
Maple [C] (verified)	767
Fricas [B] (verification not implemented)	768
Sympy [F]	769
Maxima [B] (verification not implemented)	769
Giac [A] (verification not implemented)	769
Mupad [B] (verification not implemented)	770
Reduce [B] (verification not implemented)	770

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \cosh(a + bx) \tanh(c + bx) dx = \frac{\cosh(a + bx)}{b} - \frac{\arctan(\sinh(c + bx)) \sinh(a - c)}{b}$$

output `cosh(b*x+a)/b-arctan(sinh(b*x+c))*sinh(a-c)/b`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 86 vs. 2(29) = 58.

Time = 0.04 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.97

$$\begin{aligned} & \int \cosh(a + bx) \tanh(c + bx) dx \\ &= \frac{\cosh(a) \cosh(bx)}{b} \\ & \quad - \frac{2 \arctan \left(\frac{(\cosh(c) - \sinh(c)) \left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right) \right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)} \right) \sinh(a - c)}{b} \\ & \quad + \frac{\sinh(a) \sinh(bx)}{b} \end{aligned}$$

input `Integrate[Cosh[a + b*x]*Tanh[c + b*x],x]`

output $(\text{Cosh}[a]*\text{Cosh}[b*x])/b - (2*\text{ArcTan}[(\text{Cosh}[c] - \text{Sinh}[c])*(\text{Cosh}[(b*x)/2]*\text{Sinh}[c] + \text{Cosh}[c]*\text{Sinh}[(b*x)/2])]/(\text{Cosh}[c]*\text{Cosh}[(b*x)/2] - \text{Cosh}[(b*x)/2]*\text{Sinh}[c]))*\text{Sinh}[a - c])/b + (\text{Sinh}[a]*\text{Sinh}[b*x])/b$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6157, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \tanh(bx + c) dx \\
 & \quad \downarrow \text{6157} \\
 & \int \sinh(a + bx) dx - \sinh(a - c) \int \operatorname{sech}(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -i \sin(ia + ibx) dx - \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & -\sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx - i \int \sin(ia + ibx) dx \\
 & \quad \downarrow \text{3118} \\
 & \frac{\cosh(a + bx)}{b} - \sinh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4257} \\
 & \frac{\cosh(a + bx)}{b} - \frac{\sinh(a - c) \arctan(\sinh(bx + c))}{b}
 \end{aligned}$$

input `Int[Cosh[a + b*x]*Tanh[c + b*x],x]`

output `Cosh[a + b*x]/b - (ArcTan[Sinh[c + b*x]]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6157 `Int[Cosh[v_]*Tanh[w_]^(n_), x_Symbol] := Int[Sinh[v]*Tanh[w]^(n - 1), x] - Simp[Sinh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 167, normalized size of antiderivative = 5.76

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a} + ie^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{i \ln(e^{bx+a} + ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*tanh(b*x+c),x,method=_RETURNVERBOSE)`

output

```
1/2/b*exp(b*x+a)+1/2/b*exp(-b*x-a)+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-
a-c)*exp(a)^2-1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(c)^2-1/2*I*ln
n(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(a)^2+1/2*I*ln(exp(b*x+a)+I*exp(a-
c))/b*exp(-a-c)*exp(c)^2
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. $2(29) = 58$.

Time = 0.11 (sec) , antiderivative size = 327, normalized size of antiderivative = 11.28

$$\int \cosh(a + bx) \tanh(c + bx) dx$$

$$= \frac{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c)^2 \cosh(-a + c) \sinh(-a + c) + \cosh(bx + c)^2 \sinh(-a + c)^2}{2}$$

input

```
integrate(cosh(b*x+a)*tanh(b*x+c),x, algorithm="fricas")
```

output

```
1/2*(cosh(b*x + c)^2*cosh(-a + c)^2 - 2*cosh(b*x + c)^2*cosh(-a + c)*sinh(-
a + c) + cosh(b*x + c)^2*sinh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)
)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^2 + 2*(2*cosh(b*x + c)*cosh
(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1
)*cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a
+ c)^2 - 1)*sinh(b*x + c))*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(cosh
(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cos
h(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) + 1)/(b*cosh(b*x + c)*cosh(-a + c
) - b*cosh(b*x + c)*sinh(-a + c) + (b*cosh(-a + c) - b*sinh(-a + c))*sinh(
b*x + c))
```

Sympy [F]

$$\int \cosh(a + bx) \tanh(c + bx) dx = \int \cosh(a + bx) \tanh(bx + c) dx$$

input `integrate(cosh(b*x+a)*tanh(b*x+c),x)`

output `Integral(cosh(a + b*x)*tanh(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(29) = 58.

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.03

$$\int \cosh(a + bx) \tanh(c + bx) dx = \frac{(e^{(2a)} - e^{(2c)}) \arctan(e^{(-bx-c)}) e^{(-a-c)}}{b} + \frac{e^{(bx+a)}}{2b} + \frac{e^{(-bx-a)}}{2b}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c),x, algorithm="maxima")`

output `(e^(2*a) - e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b + 1/2*e^(b*x + a)/b + 1/2*e^(-b*x - a)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.83

$$\int \cosh(a + bx) \tanh(c + bx) dx = -\frac{2(e^{(2a)} - e^{(2c)}) \arctan(e^{(bx+c)}) e^{(-a-c)} - e^{(bx+a)} - e^{(-bx-a)}}{2b}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c),x, algorithm="giac")`

output

$$-1/2*(2*(e^{(2*a)} - e^{(2*c)})*\arctan(e^{(b*x + c)})*e^{(-a - c)} - e^{(b*x + a)} - e^{(-b*x - a)})/b$$

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 133, normalized size of antiderivative = 4.59

$$\begin{aligned} & \int \cosh(a + bx) \tanh(c + bx) dx \\ &= \frac{e^{a+bx}}{2b} + \frac{e^{-a-bx}}{2b} \\ &+ \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2 - e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{b^2}} \end{aligned}$$

input

$$\operatorname{int}(\cosh(a + b*x)*\tanh(c + b*x), x)$$

output

$$\begin{aligned} & \exp(a + b*x)/(2*b) + \exp(-a - b*x)/(2*b) + (\operatorname{atan}((\exp(-a)*\exp(2*c)*\exp(b*x) \\ & *((b^2)^{(1/2)} - \exp(2*a)*\exp(-2*c)*(b^2)^{(1/2)}))/ (b*(\exp(-2*a)*\exp(2*c)* \\ & (\exp(4*a)*\exp(-4*c) - 2*\exp(2*a)*\exp(-2*c) + 1))^{(1/2)})) * (\exp(2*c - 2*a)* \\ & (\exp(4*a - 4*c) - 2*\exp(2*a - 2*c) + 1))^{(1/2)}) / (b^2)^{(1/2)} \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.34

$$\begin{aligned} & \int \cosh(a + bx) \tanh(c + bx) dx \\ &= \frac{-2e^{bx+2a} \operatorname{atan}(e^{bx+c}) + 2e^{bx+2c} \operatorname{atan}(e^{bx+c}) + e^{2bx+2a+c} + e^c}{2e^{bx+a+c} b} \end{aligned}$$

input

$$\operatorname{int}(\cosh(b*x+a)*\tanh(b*x+c), x)$$

output

$$(-2*e^{(2*a + b*x)}*\operatorname{atan}(e^{(b*x + c)}) + 2*e^{(b*x + 2*c)}*\operatorname{atan}(e^{(b*x + c)}) + e^{(2*a + 2*b*x + c)} + e^c)/(2*e^{(a + b*x + c)}*b)$$

3.109 $\int \cosh(a + bx) \tanh^2(c + bx) dx$

Optimal result	771
Mathematica [B] (verified)	771
Rubi [A] (verified)	772
Maple [C] (verified)	774
Fricas [B] (verification not implemented)	775
Sympy [F]	776
Maxima [B] (verification not implemented)	776
Giac [B] (verification not implemented)	776
Mupad [B] (verification not implemented)	777
Reduce [B] (verification not implemented)	778

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = -\frac{\arctan(\sinh(c + bx)) \cosh(a - c)}{b} + \frac{\operatorname{sech}(c + bx) \sinh(a - c)}{b} + \frac{\sinh(a + bx)}{b}$$

output

```
-arctan(sinh(b*x+c))*cosh(a-c)/b+sech(b*x+c)*sinh(a-c)/b+sinh(b*x+a)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 102 vs. 2(45) = 90.

Time = 0.08 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.27

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = -\frac{2 \arctan\left(\frac{(\cosh(c) - \sinh(c)) \left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right)\right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a - c)}{b} + \frac{\cosh(bx) \sinh(a)}{b} + \frac{\operatorname{sech}(c + bx) \sinh(a - c)}{b} + \frac{\cosh(a) \sinh(bx)}{b}$$

input `Integrate[Cosh[a + b*x]*Tanh[c + b*x]^2,x]`

output `(-2*ArcTan[((Cosh[c] - Sinh[c])*(Cosh[(b*x)/2]*Sinh[c] + Cosh[c]*Sinh[(b*x)/2]))/(Cosh[c]*Cosh[(b*x)/2] - Cosh[(b*x)/2]*Sinh[c])]*Cosh[a - c])/b + (Cosh[b*x]*Sinh[a])/b + (Sech[c + b*x]*Sinh[a - c])/b + (Cosh[a]*Sinh[b*x])/b`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6157, 3042, 26, 3086, 24, 6154, 3042, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \tanh^2(bx + c) dx \\
 & \quad \downarrow \text{6157} \\
 & \int \sinh(a + bx) \tanh(c + bx) dx - \sinh(a - c) \int \operatorname{sech}(c + bx) \tanh(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sinh(a + bx) \tanh(c + bx) dx - \sinh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & \int \sinh(a + bx) \tanh(c + bx) dx + i \sinh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3086} \\
 & \int \sinh(a + bx) \tanh(c + bx) dx + \frac{\sinh(a - c) \int 1 d\operatorname{sech}(c + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & \int \sinh(a + bx) \tanh(c + bx) dx + \frac{\sinh(a - c) \operatorname{sech}(bx + c)}{b} \\
 & \quad \downarrow \text{6154}
 \end{aligned}$$

$$\begin{aligned}
& -\cosh(a-c) \int \operatorname{sech}(c+bx) dx + \int \cosh(a+bx) dx + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{3042} \\
& -\cosh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \int \sin\left(ia+ibx+\frac{\pi}{2}\right) dx + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow \text{3117} \\
& -\cosh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} + \frac{\sinh(a+bx)}{b} \\
& \quad \downarrow \text{4257} \\
& -\frac{\cosh(a-c) \arctan(\sinh(bx+c))}{b} + \frac{\sinh(a-c)\operatorname{sech}(bx+c)}{b} + \frac{\sinh(a+bx)}{b}
\end{aligned}$$

input `Int[Cosh[a + b*x]*Tanh[c + b*x]^2,x]`

output `-((ArcTan[Sinh[c + b*x]]*Cosh[a - c])/b) + (Sech[c + b*x]*Sinh[a - c])/b + Sinh[a + b*x]/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]`
`/; FreeQ[{c, d}, x]`

rule 6154 `Int[Sinh[v_]*Tanh[w_]^(n_.), x_Symbol] := Int[Cosh[v]*Tanh[w]^(n - 1), x] -`
`Simp[Cosh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ`
`[w, v] && FreeQ[v - w, x]`

rule 6157 `Int[Cosh[v_]*Tanh[w_]^(n_.), x_Symbol] := Int[Sinh[v]*Tanh[w]^(n - 1), x] -`
`Simp[Sinh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ`
`[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 207, normalized size of antiderivative = 4.60

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(e^{2a}-e^{2c})}{b(e^{2bx+2a+2c}+e^{2a})} + \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*tanh(b*x+c)^2,x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+a)-1/2/b*exp(-b*x-a)+1/b*exp(b*x+a)*(exp(2*a)-exp(2*c))/(exp`
`(2*b*x+2*a+2*c)+exp(2*a))+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(`
`2*a)+1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*c)-1/2*I*ln(exp(b*x`
`+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*a)-1/2*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp`
`(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 902 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 902, normalized size of antiderivative = 20.04

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)^2,x, algorithm="fricas")`

output

```
1/2*(cosh(b*x + c)^4*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^4 + 4*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^3 + 3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + 3*(2*cosh(b*x + c)^2*cosh(-a + c)^2 + (2*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + cosh(-a + c)^2 - 2*(2*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c) - 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^4 + 3*cosh(b*x + c)^2)*sinh(-a + c)^2 - 2*((cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 + cosh(b*x + c))*sinh(-a + c)^2 + (cosh(-a + c)^2 + 1)*cosh(b*x + c) + (3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (3*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + cosh(-a + c)^2 - 2*(3*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c) - 2*(cosh(b*x + c)^3*cosh(-a + c) + cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(2*cosh(b*x + c)^3*cosh(-a + c)^2 + (2*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a + c)^2 + 3*(cosh(-a + c)^2 - 1)*cosh(b*x + c) - 2*(2*cosh(b*x + c)^3*cosh(-a + c) + 3*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c) - 2*(cosh(b*x + c)^4*cosh(-a + c) + 3*cosh(b*x + c)^2*cosh(-a + c))*sinh(-a + c)...
```


Sympy [F]

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = \int \cosh(a + bx) \tanh^2(bx + c) dx$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)**2,x)`

output `Integral(cosh(a + b*x)*tanh(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(45) = 90$.

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.29

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = \frac{(e^{2a} + e^{2c}) \arctan(e^{-bx-c}) e^{-a-c}}{b} - \frac{e^{-bx-a}}{2b} + \frac{(3e^{2a} - 2e^{2c})e^{-2bx-2a} + e^{2c}}{2b(e^{-bx-a+2c} + e^{-3bx-a})}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)^2,x, algorithm="maxima")`

output `(e^(2*a) + e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b - 1/2*e^(-b*x - a)/b + 1/2*((3*e^(2*a) - 2*e^(2*c))*e^(-2*b*x - 2*a) + e^(2*c))/(b*(e^(-b*x - a + 2*c) + e^(-3*b*x - a)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(45) = 90$.

Time = 0.16 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.16

$$\int \cosh(a + bx) \tanh^2(c + bx) dx = -\frac{2(e^{2a} + e^{2c}) \arctan(e^{(bx+c)}) e^{-a-c} - \frac{2e^{(2bx+4a)} - 3e^{(2bx+2a+2c)} - e^{(2a)}}{e^{(3bx+3a+2c)} + e^{(bx+3a)}} - e^{(bx+a)}}{2b}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)^2,x, algorithm="giac")`

output
$$-1/2*(2*(e^{(2*a)} + e^{(2*c)})*\arctan(e^{(b*x + c)})*e^{(-a - c)} - (2*e^{(2*b*x + 4*a)} - 3*e^{(2*b*x + 2*a + 2*c)} - e^{(2*a)})/(e^{(3*b*x + 3*a + 2*c)} + e^{(b*x + 3*a)}) - e^{(b*x + a)})/b$$

Mupad [B] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 173, normalized size of antiderivative = 3.84

$$\begin{aligned} & \int \cosh(a + bx) \tanh^2(c + bx) dx \\ &= \frac{e^{a+bx}}{2b} - \frac{e^{-a-bx}}{2b} \\ & \quad - \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2+e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{b^2}} \\ & \quad + \frac{e^{a+bx} (e^{2a-2c} - 1)}{b (e^{2a-2c} + e^{2a+2bx})} \end{aligned}$$

input `int(cosh(a + b*x)*tanh(c + b*x)^2,x)`

output
$$\begin{aligned} & \exp(a + b*x)/(2*b) - \exp(-a - b*x)/(2*b) - (\operatorname{atan}((\exp(-a)*\exp(2*c)*\exp(b*x) \\ & *((b^2)^{(1/2)} + \exp(2*a)*\exp(-2*c)*(b^2)^{(1/2)}))/(b*(\exp(-2*a)*\exp(2*c)* \\ & (2*\exp(2*a)*\exp(-2*c) + \exp(4*a)*\exp(-4*c) + 1))^{(1/2)}))*(\exp(2*c - 2*a)* \\ & (2*\exp(2*a - 2*c) + \exp(4*a - 4*c) + 1))^{(1/2)})/(b^2)^{(1/2)} + (\exp(a + b*x) \\ & *(\exp(2*a - 2*c) - 1))/(b*(\exp(2*a - 2*c) + \exp(2*a + 2*b*x))) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 154, normalized size of antiderivative = 3.42

$$\int \cosh(a + bx) \tanh^2(c + bx) dx$$

$$= \frac{-2e^{3bx+2a+2c} \operatorname{atan}(e^{bx+c}) - 2e^{3bx+4c} \operatorname{atan}(e^{bx+c}) - 2e^{bx+2a} \operatorname{atan}(e^{bx+c}) - 2e^{bx+2c} \operatorname{atan}(e^{bx+c}) + e^{4bx+2a+3c}}{2e^{bx+a+c} b (e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)*tanh(b*x+c)^2,x)`output `(- 2*e**(2*a + 3*b*x + 2*c)*atan(e**(b*x + c)) - 2*e**(3*b*x + 4*c)*atan(e**(b*x + c)) - 2*e**(2*a + b*x)*atan(e**(b*x + c)) - 2*e**(b*x + 2*c)*atan(e**(b*x + c)) + e**(2*a + 4*b*x + 3*c) + 3*e**(2*a + 2*b*x + c) - 3*e**(2*b*x + 3*c) - e**c)/(2*e**(a + b*x + c)*b*(e**(2*b*x + 2*c) + 1))`

3.110 $\int \cosh(a + bx) \tanh^3(c + bx) dx$

Optimal result	779
Mathematica [A] (verified)	779
Rubi [A] (verified)	780
Maple [C] (verified)	784
Fricas [B] (verification not implemented)	784
Sympy [F]	785
Maxima [B] (verification not implemented)	786
Giac [A] (verification not implemented)	786
Mupad [F(-1)]	787
Reduce [B] (verification not implemented)	787

Optimal result

Integrand size = 15, antiderivative size = 72

$$\int \cosh(a + bx) \tanh^3(c + bx) dx = \frac{\cosh(a + bx)}{b} + \frac{\cosh(a - c)\operatorname{sech}(c + bx)}{b} - \frac{3 \arctan(\sinh(c + bx)) \sinh(a - c)}{2b} + \frac{\operatorname{sech}(c + bx) \sinh(a - c) \tanh(c + bx)}{2b}$$

output

```
cosh(b*x+a)/b+cosh(a-c)*sech(b*x+c)/b-3/2*arctan(sinh(b*x+c))*sinh(a-c)/b+1/2*sech(b*x+c)*sinh(a-c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.60

$$\int \cosh(a + bx) \tanh^3(c + bx) dx = \frac{\cosh(a - 2c)\operatorname{sech}(c)\operatorname{sech}(c + bx) - \cosh(a - c - bx)\operatorname{sech}(c)\operatorname{sech}^2(c + bx) + \cosh(a - c + bx)\operatorname{sech}(c)\operatorname{sech}(c + bx)}{b}$$

input

```
Integrate[Cosh[a + b*x]*Tanh[c + b*x]^3,x]
```

output

```
(Cosh[a - 2*c]*Sech[c]*Sech[c + b*x] - Cosh[a - c - b*x]*Sech[c]*Sech[c +
b*x]^2 + Cosh[a - c + b*x]*Sech[c]*Sech[c + b*x]^2 + Cosh[a]*(4*Cosh[b*x]
+ 3*Sech[c]*Sech[c + b*x]) - 12*ArcTan[Sinh[c] + Cosh[c]*Tanh[(b*x)/2]]*Si
nh[a - c] + 4*Sinh[a]*Sinh[b*x])/(4*b)
```

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.19, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.067$, Rules used = {6157, 3042, 25, 3091, 3042, 4257, 6154, 3042, 26, 3086, 24, 6157, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \tanh^3(bx + c) dx \\
 & \quad \downarrow 6157 \\
 & \int \sinh(a + bx) \tanh^2(c + bx) dx - \sinh(a - c) \int \operatorname{sech}(c + bx) \tanh^2(c + bx) dx \\
 & \quad \downarrow 3042 \\
 & \int \sinh(a + bx) \tanh^2(c + bx) dx - \sinh(a - c) \int -\sec(ic + ibx) \tan(ic + ibx)^2 dx \\
 & \quad \downarrow 25 \\
 & \int \sinh(a + bx) \tanh^2(c + bx) dx + \sinh(a - c) \int \sec(ic + ibx) \tan(ic + ibx)^2 dx \\
 & \quad \downarrow 3091 \\
 & \int \sinh(a + bx) \tanh^2(c + bx) dx + \sinh(a - c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{1}{2} \int \operatorname{sech}(c + bx) dx \right) \\
 & \quad \downarrow 3042 \\
 & \int \sinh(a + bx) \tanh^2(c + bx) dx + \sinh(a - c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{1}{2} \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow 4257
 \end{aligned}$$

$$\begin{aligned}
& \int \sinh(a + bx) \tanh^2(c + bx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{6154} \\
& \int \cosh(a + bx) \tanh(c + bx) dx - \cosh(a - c) \int \operatorname{sech}(c + bx) \tanh(c + bx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& \int \cosh(a + bx) \tanh(c + bx) dx - \cosh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{26} \\
& \int \cosh(a + bx) \tanh(c + bx) dx + i \cosh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{3086} \\
& \int \cosh(a + bx) \tanh(c + bx) dx + \frac{\cosh(a - c) \int 1 d\operatorname{sech}(c + bx)}{b} + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) \\
& \quad \downarrow \text{24} \\
& \int \cosh(a + bx) \tanh(c + bx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) + \frac{\cosh(a - c) \operatorname{sech}(bx + c)}{b} \\
& \quad \downarrow \text{6157} \\
& - \sinh(a - c) \int \operatorname{sech}(c + bx) dx + \int \sinh(a + bx) dx + \sinh(a - \\
& c) \left(\frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} - \frac{\arctan(\sinh(bx + c))}{2b} \right) + \frac{\cosh(a - c) \operatorname{sech}(bx + c)}{b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \int -i \sin(ia+ibx) dx + \sinh(a-c) \\
c) & \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow 26 \\
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx - i \int \sin(ia+ibx) dx + \sinh(a-c) \\
c) & \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} \\
& \quad \downarrow 3118 \\
& -\sinh(a-c) \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx + \sinh(a-c) \\
c) & \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} + \\
& \quad \frac{\cosh(a+bx)}{b} \\
& \quad \downarrow 4257 \\
& -\frac{\sinh(a-c)\arctan(\sinh(bx+c))}{b} + \sinh(a-c) \\
c) & \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} - \frac{\arctan(\sinh(bx+c))}{2b} \right) + \frac{\cosh(a-c)\operatorname{sech}(bx+c)}{b} + \\
& \quad \frac{\cosh(a+bx)}{b}
\end{aligned}$$

input

```
Int[Cosh[a + b*x]*Tanh[c + b*x]^3,x]
```

output

```
Cosh[a + b*x]/b + (Cosh[a - c]*Sech[c + b*x])/b - (ArcTan[Sinh[c + b*x]]*Sinh[a - c])/b + Sinh[a - c]*(-1/2*ArcTan[Sinh[c + b*x]]/b + (Sech[c + b*x]*Tanh[c + b*x])/(2*b))
```

Definitions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`
- rule 3091 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Simp[b^2*((n - 1)/(m + n - 1)) Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]`
- rule 3118 `Int[sin[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 6154 `Int[Sinh[v_]*Tanh[w_]^(n_), x_Symbol] := Int[Cosh[v]*Tanh[w]^(n - 1), x] - Simp[Cosh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

rule 6157

```
Int[Cosh[v_]*Tanh[w_]^(n_), x_Symbol] := Int[Sinh[v]*Tanh[w]^(n - 1), x] -
Simp[Sinh[v - w] Int[Sech[w]*Tanh[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ
[w, v] && FreeQ[v - w, x]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 238, normalized size of antiderivative = 3.31

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(3e^{2bx+4a+2c} + e^{2bx+2a+4c} + e^{4a} + 3e^{2a+2c})}{2b(e^{2bx+2a+2c} + e^{2a})^2} + \frac{3i \ln(e^{bx+a} - ie^{a-c})e^{-a-c}e^{2a}}{4b} - \frac{3i \ln(e^{bx+a} - ie^{a-c})}{4b}$

input

```
int(cosh(b*x+a)*tanh(b*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
1/2/b*exp(b*x+a)+1/2/b*exp(-b*x-a)+1/2*exp(b*x+a)*(3*exp(2*b*x+4*a+2*c)+ex
p(2*b*x+2*a+4*c)+exp(4*a)+3*exp(2*a+2*c))/b/(exp(2*b*x+2*a+2*c)+exp(2*a))^
2+3/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(a)^2-3/4*I*ln(exp(b*x+a)
-I*exp(a-c))/b*exp(-a-c)*exp(c)^2-3/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a
-c)*exp(a)^2+3/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(c)^2
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1737 vs. 2(68) = 136.

Time = 0.11 (sec) , antiderivative size = 1737, normalized size of antiderivative = 24.12

$$\int \cosh(a + bx) \tanh^3(c + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*tanh(b*x+c)^3,x, algorithm="fricas")
```

output

```

1/2*(cosh(b*x + c)^6*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^6 + 6*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^5 + (5*cosh(-a + c)^2 + 2)*cosh(b*x + c)^4 + (15*cosh(b*x + c)^2*cosh(-a + c)^2 + 5*(3*cosh(b*x + c)^2 + 1)*sinh(-a + c)^2 + 5*cosh(-a + c)^2 - 10*(3*cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a + c) + 2)*sinh(b*x + c)^4 + 4*(5*cosh(b*x + c)^3*cosh(-a + c)^2 + 5*(cosh(b*x + c)^3 + cosh(b*x + c))*sinh(-a + c)^2 + (5*cosh(-a + c)^2 + 2)*cosh(b*x + c) - 10*(cosh(b*x + c)^3*cosh(-a + c) + cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^3 + (2*cosh(-a + c)^2 + 5)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^4*cosh(-a + c)^2 + 6*(5*cosh(-a + c)^2 + 2)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^4 + 30*cosh(b*x + c)^2 + 2)*sinh(-a + c)^2 + 2*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^4*cosh(-a + c) + 30*cosh(b*x + c)^2*cosh(-a + c) + 2*cosh(-a + c))*sinh(-a + c) + 5)*sinh(b*x + c)^2 + (cosh(b*x + c)^6 + 5*cosh(b*x + c)^4 + 2*cosh(b*x + c)^2)*sinh(-a + c)^2 - 3*((cosh(-a + c)^2 - 1)*cosh(b*x + c)^5 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^5 - 5*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^4 + 2*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + 2*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 + 1)*sinh(-a + ...

```

Sympy [F]

$$\int \cosh(a + bx) \tanh^3(c + bx) dx = \int \cosh(a + bx) \tanh^3(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*tanh(b*x+c)**3,x)
```

output

```
Integral(cosh(a + b*x)*tanh(b*x + c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(68) = 136$.

Time = 0.12 (sec) , antiderivative size = 149, normalized size of antiderivative = 2.07

$$\int \cosh(a + bx) \tanh^3(c + bx) dx$$

$$= \frac{3(e^{2a} - e^{2c}) \arctan(e^{-bx-c}) e^{(-a-c)} + \frac{e^{-bx-a}}{2b}}{\frac{2b}{(5e^{2a+2c} + e^{4c})e^{-2bx-2a} + (2e^{4a} + 3e^{2a+2c})e^{-4bx-4a} + e^{4c}} + \frac{2b}{2b(e^{-bx-a+4c} + 2e^{-3bx-a+2c} + e^{-5bx-a})}}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)^3,x, algorithm="maxima")`

output
$$\frac{3/2*(e^{2*a} - e^{2*c})*\arctan(e^{-b*x - c})*e^{-a - c}/b + 1/2*e^{-b*x - a}/b + 1/2*((5*e^{2*a + 2*c} + e^{4*c})*e^{-2*b*x - 2*a} + (2*e^{4*a} + 3*e^{2*a + 2*c})*e^{-4*b*x - 4*a} + e^{4*c})/(b*(e^{-b*x - a + 4*c} + 2*e^{-3*b*x - a + 2*c} + e^{-5*b*x - a}))}{2b}$$

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.69

$$\int \cosh(a + bx) \tanh^3(c + bx) dx =$$

$$\frac{3(e^{2a} - e^{2c}) \arctan(e^{(bx+c)}) e^{(-a-c)} - \frac{3e^{(3bx+5a+2c)+e^{(3bx+3a+4c)+e^{(bx+5a)+3e^{(bx+3a+2c)}}}{(e^{2bx+2a+2c}+e^{2a})^2} - e^{(bx+a)} - e^{(-bx-a)}}{2b}}$$

input `integrate(cosh(b*x+a)*tanh(b*x+c)^3,x, algorithm="giac")`

output
$$\frac{-1/2*(3*(e^{2*a} - e^{2*c})*\arctan(e^{(b*x + c)})*e^{-a - c} - (3*e^{(3*b*x + 5*a + 2*c)} + e^{(3*b*x + 3*a + 4*c)} + e^{(b*x + 5*a)} + 3*e^{(b*x + 3*a + 2*c)}))/(e^{(2*b*x + 2*a + 2*c)} + e^{(2*a)})^2 - e^{(b*x + a)} - e^{(-b*x - a)})/b}{2b}$$

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \tanh^3(c + bx) dx = \int \cosh(a + bx) \tanh(c + bx)^3 dx$$

input `int(cosh(a + b*x)*tanh(c + b*x)^3,x)`output `int(cosh(a + b*x)*tanh(c + b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 234, normalized size of antiderivative = 3.25

$$\int \cosh(a + bx) \tanh^3(c + bx) dx$$

$$= \frac{-3e^{5bx+2a+4c} \operatorname{atan}(e^{bx+c}) + 3e^{5bx+6c} \operatorname{atan}(e^{bx+c}) - 6e^{3bx+2a+2c} \operatorname{atan}(e^{bx+c}) + 6e^{3bx+4c} \operatorname{atan}(e^{bx+c}) - 3e^{bx+c}}{2e^{bx+a+c} b (e^{4bx+4c})}$$

input `int(cosh(b*x+a)*tanh(b*x+c)^3,x)`output `(- 3*e**(2*a + 5*b*x + 4*c)*atan(e**(b*x + c)) + 3*e**(5*b*x + 6*c)*atan(e**(b*x + c)) - 6*e**(2*a + 3*b*x + 2*c)*atan(e**(b*x + c)) + 6*e**(3*b*x + 4*c)*atan(e**(b*x + c)) - 3*e**(2*a + b*x)*atan(e**(b*x + c)) + 3*e**(b*x + 2*c)*atan(e**(b*x + c)) + e**(2*a + 6*b*x + 5*c) + 5*e**(2*a + 4*b*x + 3*c) + 2*e**(4*b*x + 5*c) + 2*e**(2*a + 2*b*x + c) + 5*e**(2*b*x + 3*c) + e**c)/(2*e**(a + b*x + c)*b*(e**(4*b*x + 4*c) + 2*e**(2*b*x + 2*c) + 1))`

3.111 $\int \cosh(a + bx) \tanh(c + dx) dx$

Optimal result	788
Mathematica [A] (verified)	788
Rubi [A] (verified)	789
Maple [F]	790
Fricas [F]	790
Sympy [F]	791
Maxima [F]	791
Giac [F]	791
Mupad [F(-1)]	792
Reduce [F]	792

Optimal result

Integrand size = 13, antiderivative size = 99

$$\int \cosh(a + bx) \tanh(c + dx) dx = \frac{e^{-a-bx} \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, -e^{2(c+dx)}\right)}{b} + \frac{\sinh(a + bx)}{b}$$

```
output exp(-b*x-a)*hypergeom([1, -1/2*b/d], [1-1/2*b/d], -exp(2*d*x+2*c))/b-exp(b*x+a)*hypergeom([1, 1/2*b/d], [1+1/2*b/d], -exp(2*d*x+2*c))/b+sinh(b*x+a)/b
```

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.04

$$\int \cosh(a + bx) \tanh(c + dx) dx = \frac{e^{-a-bx}(-1 + e^{2(a+bx)} + 2 \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right) - 2e^{2(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, -e^{2(c+dx)}\right))}{2b}$$

input `Integrate[Cosh[a + b*x]*Tanh[c + d*x],x]`

output $(E^{-a - b*x}*(-1 + E^{2*(a + b*x)}) + 2*Hypergeometric2F1[1, -1/2*b/d, 1 - b/(2*d), -E^{2*(c + d*x)}]) - 2*E^{2*(a + b*x)}*Hypergeometric2F1[1, b/(2*d), 1 + b/(2*d), -E^{2*(c + d*x)}])/(2*b)$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.21, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6138, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \tanh(c + dx) dx$$

$$\downarrow 6138$$

$$\int \left(-\frac{e^{-a-bx}}{e^{2(c+dx)} + 1} - \frac{e^{a+bx}}{e^{2(c+dx)} + 1} + \frac{1}{2}e^{-a-bx} + \frac{1}{2}e^{a+bx} \right) dx$$

$$\downarrow 2009$$

$$\frac{e^{-a-bx} \text{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, -e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \text{Hypergeometric2F1}\left(1, \frac{b}{2d}, \frac{b}{2d} + 1, -e^{2(c+dx)}\right)}{b} - \frac{e^{-a-bx}}{2b} + \frac{e^{a+bx}}{2b}$$

input `Int[Cosh[a + b*x]*Tanh[c + d*x],x]`

output $-1/2*E^{-a - b*x}/b + E^{a + b*x}/(2*b) + (E^{-a - b*x}*Hypergeometric2F1[1, -1/2*b/d, 1 - b/(2*d), -E^{2*(c + d*x)}])/b - (E^{a + b*x}*Hypergeometric2F1[1, b/(2*d), 1 + b/(2*d), -E^{2*(c + d*x)}])/b$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6138 `Int[Cosh[(a_.) + (b_.)*(x_)]*Tanh[(c_.) + (d_.)*(x_)], x_Symbol] := Int[1/(E^(a + b*x)*2) + E^(a + b*x)/2 - 1/(E^(a + b*x)*(1 + E^(2*(c + d*x)))) - E^(a + b*x)/(1 + E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \cosh (bx + a) \tanh (dx + c) dx$$

input `int(cosh(b*x+a)*tanh(d*x+c),x)`

output `int(cosh(b*x+a)*tanh(d*x+c),x)`

Fricas [F]

$$\int \cosh(a + bx) \tanh(c + dx) dx = \int \cosh (bx + a) \tanh (dx + c) dx$$

input `integrate(cosh(b*x+a)*tanh(d*x+c),x, algorithm="fricas")`

output `integral(cosh(b*x + a)*tanh(d*x + c), x)`

Sympy [F]

$$\int \cosh(a + bx) \tanh(c + dx) dx = \int \cosh(a + bx) \tanh(c + dx) dx$$

input `integrate(cosh(b*x+a)*tanh(d*x+c),x)`

output `Integral(cosh(a + b*x)*tanh(c + d*x), x)`

Maxima [F]

$$\int \cosh(a + bx) \tanh(c + dx) dx = \int \cosh(bx + a) \tanh(dx + c) dx$$

input `integrate(cosh(b*x+a)*tanh(d*x+c),x, algorithm="maxima")`

output `1/2*(e^(2*b*x + 2*a) - 1)*e^(-b*x - a)/b - 1/2*integrate(2*(e^(2*b*x + 2*a) + 1)/(e^(b*x + 2*d*x + a + 2*c) + e^(b*x + a)), x)`

Giac [F]

$$\int \cosh(a + bx) \tanh(c + dx) dx = \int \cosh(bx + a) \tanh(dx + c) dx$$

input `integrate(cosh(b*x+a)*tanh(d*x+c),x, algorithm="giac")`

output `integrate(cosh(b*x + a)*tanh(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \tanh(c + dx) dx = \int \cosh(a + bx) \tanh(c + dx) dx$$

input `int(cosh(a + b*x)*tanh(c + d*x),x)`output `int(cosh(a + b*x)*tanh(c + d*x), x)`**Reduce [F]**

$$\int \cosh(a + bx) \tanh(c + dx) dx$$

$$= \frac{e^{2bx+2a} - 2e^{bx+2a} \left(\int \frac{e^{bx}}{e^{2dx+2c}+1} dx \right) b - 2e^{bx} \left(\int \frac{1}{e^{bx+2dx+2c}+e^{bx}} dx \right) b - 1}{2e^{bx+ab}}$$

input `int(cosh(b*x+a)*tanh(d*x+c),x)`output `(e**(2*a + 2*b*x) - 2*e**(2*a + b*x)*int(e**(b*x)/(e**(2*c + 2*d*x) + 1),x)*b - 2*e**(b*x)*int(1/(e**(b*x + 2*c + 2*d*x) + e**(b*x)),x)*b - 1)/(2*e*(a + b*x)*b)`

3.112 $\int \cosh(x) \coth(2x) dx$

Optimal result	793
Mathematica [B] (verified)	793
Rubi [A] (verified)	794
Maple [B] (verified)	795
Fricas [B] (verification not implemented)	796
Sympy [F]	796
Maxima [B] (verification not implemented)	797
Giac [B] (verification not implemented)	797
Mupad [B] (verification not implemented)	797
Reduce [B] (verification not implemented)	798

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \cosh(x) \coth(2x) dx = -\frac{1}{2} \operatorname{arctanh}(\cosh(x)) + \cosh(x)$$

output `-1/2*arctanh(cosh(x))+cosh(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. $2(10) = 20$.

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.50

$$\int \cosh(x) \coth(2x) dx = \cosh(x) - \frac{1}{2} \log\left(\cosh\left(\frac{x}{2}\right)\right) + \frac{1}{2} \log\left(\sinh\left(\frac{x}{2}\right)\right)$$

input `Integrate[Cosh[x]*Coth[2*x],x]`

output `Cosh[x] - Log[Cosh[x/2]]/2 + Log[Sinh[x/2]]/2`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.40, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \coth(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \cos(ix) \cot(2ix) dx \\
 & \quad \downarrow \text{26} \\
 & i \int \cos(ix) \cot(2ix) dx \\
 & \quad \downarrow \text{4879} \\
 & - \int -\frac{1 - 2 \cosh^2(x)}{2(1 - \cosh^2(x))} d \cosh(x) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \frac{1 - 2 \cosh^2(x)}{1 - \cosh^2(x)} d \cosh(x) \\
 & \quad \downarrow \text{299} \\
 & \frac{1}{2} \left(2 \cosh(x) - \int \frac{1}{1 - \cosh^2(x)} d \cosh(x) \right) \\
 & \quad \downarrow \text{219} \\
 & \frac{1}{2} (2 \cosh(x) - \operatorname{arctanh}(\cosh(x)))
 \end{aligned}$$

input

```
Int [Cosh[x]*Coth[2*x], x]
```

output

```
(-ArcTanh[Cosh[x]] + 2*Cosh[x])/2
```

Defintions of rubi rules used

- rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F_x), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 219 $\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 299 $\text{Int}[((a_) + (b_)*(x_)^2)^{(p_)}*((c_) + (d_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^2)^{(p + 1)}/(b*(2*p + 3))), x] - \text{Simp}[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) \text{Int}[(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[2*p + 3, 0]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4879 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}\{d = \text{FreeFactors}[\text{Cos}[v], x]\}, -d/\text{Coefficient}[v, x, 1] \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Cos}[v]/d, u/\text{Sin}[v], x], x], x, \text{Cos}[v]/d]], x] /; \text{!FalseQ}[v] \ \&\& \ \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Cos}[v], x], u/\text{Sin}[v], x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(8) = 16$.

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.60

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\ln(e^x-1)}{2} - \frac{\ln(e^x+1)}{2}$	26

input `int(cosh(x)*coth(2*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+1/2*ln(exp(x)-1)-1/2*ln(exp(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(8) = 16$.

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 5.20

$$\int \cosh(x) \coth(2x) dx$$

$$= \frac{\cosh(x)^2 - (\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) + 1) + (\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) - 1)}{2(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)*coth(2*x),x, algorithm="fricas")`

output `1/2*(cosh(x)^2 - (cosh(x) + sinh(x))*log(cosh(x) + sinh(x) + 1) + (cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \cosh(x) \coth(2x) dx = \int \cosh(x) \coth(2x) dx$$

input `integrate(cosh(x)*coth(2*x),x)`

output `Integral(cosh(x)*coth(2*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(8) = 16$.

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.90

$$\int \cosh(x) \coth(2x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x - \frac{1}{2} \log(e^{(-x)} + 1) + \frac{1}{2} \log(e^{(-x)} - 1)$$

input `integrate(cosh(x)*coth(2*x),x, algorithm="maxima")`

output `1/2*e^(-x) + 1/2*e^x - 1/2*log(e^(-x) + 1) + 1/2*log(e^(-x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(8) = 16$.

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.60

$$\int \cosh(x) \coth(2x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(cosh(x)*coth(2*x),x, algorithm="giac")`

output `1/2*e^(-x) + 1/2*e^x - 1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

Mupad [B] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.90

$$\int \cosh(x) \coth(2x) dx = \frac{\ln(1 - e^x)}{2} - \frac{\ln(-e^x - 1)}{2} + \frac{e^{-x}}{2} + \frac{e^x}{2}$$

input `int(coth(2*x)*cosh(x),x)`

output `log(1 - exp(x))/2 - log(- exp(x) - 1)/2 + exp(-x)/2 + exp(x)/2`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 3.50

$$\int \cosh(x) \coth(2x) dx = \frac{e^{2x} + e^x \log(e^x - 1) - e^x \log(e^x + 1) + 1}{2e^x}$$

input `int(cosh(x)*coth(2*x),x)`

output `(e**(2*x) + e**x*log(e**x - 1) - e**x*log(e**x + 1) + 1)/(2*e**x)`

3.113 $\int \cosh(x) \coth(3x) dx$

Optimal result	799
Mathematica [B] (verified)	799
Rubi [B] (verified)	800
Maple [B] (verified)	802
Fricas [B] (verification not implemented)	803
Sympy [F]	803
Maxima [B] (verification not implemented)	804
Giac [B] (verification not implemented)	804
Mupad [B] (verification not implemented)	805
Reduce [B] (verification not implemented)	805

Optimal result

Integrand size = 7, antiderivative size = 20

$$\int \cosh(x) \coth(3x) dx = -\frac{1}{3} \operatorname{arctanh}\left(\frac{3 \cosh(x)}{2 + \cosh(2x)}\right) + \cosh(x)$$

output

```
-1/3*arctanh(3*cosh(x)/(2+cosh(2*x)))+cosh(x)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 47 vs. 2(20) = 40.

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \cosh(x) \coth(3x) dx = \cosh(x) - \frac{1}{3} \log\left(\cosh\left(\frac{x}{2}\right)\right) + \frac{1}{6} \log(1 - 2 \cosh(x)) - \frac{1}{6} \log(1 + 2 \cosh(x)) + \frac{1}{3} \log\left(\sinh\left(\frac{x}{2}\right)\right)$$

input

```
Integrate[Cosh[x]*Coth[3*x],x]
```

output

```
Cosh[x] - Log[Cosh[x/2]]/3 + Log[1 - 2*Cosh[x]]/6 - Log[1 + 2*Cosh[x]]/6 + Log[Sinh[x/2]]/3
```


Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 55 vs. $2(20) = 40$.

Time = 0.31 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 26, 4879, 1602, 27, 1475, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \coth(3x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int i \cos(ix) \cot(3ix) dx \\
 & \quad \downarrow \text{26} \\
 & i \int \cos(ix) \cot(3ix) dx \\
 & \quad \downarrow \text{4879} \\
 & - \int \frac{\cosh^2(x) (3 - 4 \cosh^2(x))}{4 \cosh^4(x) - 5 \cosh^2(x) + 1} d \cosh(x) \\
 & \quad \downarrow \text{1602} \\
 & \frac{1}{4} \int - \frac{4(1 - 2 \cosh^2(x))}{4 \cosh^4(x) - 5 \cosh^2(x) + 1} d \cosh(x) + \cosh(x) \\
 & \quad \downarrow \text{27} \\
 & \cosh(x) - \int \frac{1 - 2 \cosh^2(x)}{4 \cosh^4(x) - 5 \cosh^2(x) + 1} d \cosh(x) \\
 & \quad \downarrow \text{1475} \\
 & \frac{1}{4} \int \frac{1}{\cosh^2(x) - \frac{\cosh(x)}{2} - \frac{1}{2}} d \cosh(x) + \frac{1}{4} \int \frac{1}{\cosh^2(x) + \frac{\cosh(x)}{2} - \frac{1}{2}} d \cosh(x) + \cosh(x) \\
 & \quad \downarrow \text{1081}
 \end{aligned}$$

$$\frac{1}{4} \int \left(-\frac{2}{3(\cosh(x)+1)} - \frac{4}{3(1-2\cosh(x))} \right) d\cosh(x) +$$

$$\frac{1}{4} \int \left(-\frac{4}{3(2\cosh(x)+1)} - \frac{2}{3(1-\cosh(x))} \right) d\cosh(x) + \cosh(x)$$

↓ 2009

$$\cosh(x) + \frac{1}{4} \left(\frac{2}{3} \log(1-2\cosh(x)) - \frac{2}{3} \log(\cosh(x)+1) \right) +$$

$$\frac{1}{4} \left(\frac{2}{3} \log(1-\cosh(x)) - \frac{2}{3} \log(2\cosh(x)+1) \right)$$

input `Int[Cosh[x]*Coth[3*x],x]`

output `Cosh[x] + ((2*Log[1 - 2*Cosh[x]])/3 - (2*Log[1 + Cosh[x]])/3)/4 + ((2*Log[1 - Cosh[x]])/3 - (2*Log[1 + 2*Cosh[x]])/3)/4`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1475 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e) - b/c, 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - a*e^2, 0] && (GtQ[2*(d/e) - b/c, 0] || (!LtQ[2*(d/e) - b/c, 0] && EqQ[d - e*Rt[a/c, 2], 0]))`

rule 1602

```
Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[e*f*(f*x)^(m - 1)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 3))), x] - Simp[f^2/(c*(m + 4*p + 3)) Int[(f*x)^(m - 2)*(a + b*x^2 + c*x^4)^p*Simp[a*e*(m - 1) + (b*e*(m + 2*p + 1) - c*d*(m + 4*p + 3))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 1] && NeQ[m + 4*p + 3, 0] && IntegerQ[2*p] && (IntegerQ[p] | IntegerQ[m])
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4879

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(18) = 36$.

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.50

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\ln(e^x-1)}{3} - \frac{\ln(e^x+1)}{3} - \frac{\ln(e^{2x}+e^x+1)}{6} + \frac{\ln(e^{2x}-e^x+1)}{6}$	50

input

```
int(cosh(x)*coth(3*x),x,method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)+1/2*exp(-x)+1/3*ln(exp(x)-1)-1/3*ln(exp(x)+1)-1/6*ln(exp(2*x)+exp(x)+1)+1/6*ln(exp(2*x)-exp(x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(18) = 36$.

Time = 0.10 (sec) , antiderivative size = 104, normalized size of antiderivative = 5.20

$$\int \cosh(x) \coth(3x) dx$$

$$= \frac{3 \cosh(x)^2 - (\cosh(x) + \sinh(x)) \log\left(\frac{2 \cosh(x)+1}{\cosh(x)-\sinh(x)}\right) + (\cosh(x) + \sinh(x)) \log\left(\frac{2 \cosh(x)-1}{\cosh(x)-\sinh(x)}\right) - 2(\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x))}{6}$$

input `integrate(cosh(x)*coth(3*x),x, algorithm="fricas")`

output `1/6*(3*cosh(x)^2 - (cosh(x) + sinh(x))*log((2*cosh(x) + 1)/(cosh(x) - sinh(x))) + (cosh(x) + sinh(x))*log((2*cosh(x) - 1)/(cosh(x) - sinh(x)))) - 2*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) + 1) + 2*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) + 6*cosh(x)*sinh(x) + 3*sinh(x)^2 + 3)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \cosh(x) \coth(3x) dx = \int \cosh(x) \coth(3x) dx$$

input `integrate(cosh(x)*coth(3*x),x)`

output `Integral(cosh(x)*coth(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(18) = 36$.

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.85

$$\int \cosh(x) \coth(3x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x - \frac{1}{6} \log(e^{(-x)} + e^{(-2x)} + 1) - \frac{1}{3} \log(e^{(-x)} + 1) \\ + \frac{1}{3} \log(e^{(-x)} - 1) + \frac{1}{6} \log(-e^{(-x)} + e^{(-2x)} + 1)$$

input `integrate(cosh(x)*coth(3*x),x, algorithm="maxima")`

output `1/2*e^(-x) + 1/2*e^x - 1/6*log(e^(-x) + e^(-2*x) + 1) - 1/3*log(e^(-x) + 1) + 1/3*log(e^(-x) - 1) + 1/6*log(-e^(-x) + e^(-2*x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55 vs. $2(18) = 36$.

Time = 0.17 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \cosh(x) \coth(3x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x - \frac{1}{6} \log(e^{(-x)} + e^x + 2) - \frac{1}{6} \log(e^{(-x)} + e^x + 1) \\ + \frac{1}{6} \log(e^{(-x)} + e^x - 1) + \frac{1}{6} \log(e^{(-x)} + e^x - 2)$$

input `integrate(cosh(x)*coth(3*x),x, algorithm="giac")`

output `1/2*e^(-x) + 1/2*e^x - 1/6*log(e^(-x) + e^x + 2) - 1/6*log(e^(-x) + e^x + 1) + 1/6*log(e^(-x) + e^x - 1) + 1/6*log(e^(-x) + e^x - 2)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.85

$$\int \cosh(x) \coth(3x) dx = \frac{\ln(6 - 6e^x)}{3} - \frac{\ln(-6e^x - 6)}{3} + \frac{e^{-x}}{2} + \frac{\ln(e^x - e^{2x} - 1)}{6} - \frac{\ln(-e^{2x} - e^x - 1)}{6} + \frac{e^x}{2}$$

input `int(coth(3*x)*cosh(x),x)`output `log(6 - 6*exp(x))/3 - log(- 6*exp(x) - 6)/3 + exp(-x)/2 + log(exp(x) - exp(2*x) - 1)/6 - log(- exp(2*x) - exp(x) - 1)/6 + exp(x)/2`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 3.55

$$\int \cosh(x) \coth(3x) dx = \frac{3e^{2x} - e^x \log(e^{2x} + e^x + 1) + e^x \log(e^{2x} - e^x + 1) + 2e^x \log(e^x - 1) - 2e^x \log(e^x + 1) + 3}{6e^x}$$

input `int(cosh(x)*coth(3*x),x)`output `(3***e**(2*x) - e**x*log(e**(2*x) + e**x + 1) + e**x*log(e**(2*x) - e**x + 1) + 2*e**x*log(e**x - 1) - 2*e**x*log(e**x + 1) + 3)/(6*e**x)`

3.114 $\int \cosh(x) \coth(4x) dx$

Optimal result	806
Mathematica [C] (verified)	806
Rubi [A] (verified)	807
Maple [B] (verified)	809
Fricas [B] (verification not implemented)	809
Sympy [F]	810
Maxima [B] (verification not implemented)	810
Giac [B] (verification not implemented)	811
Mupad [B] (verification not implemented)	811
Reduce [B] (verification not implemented)	812

Optimal result

Integrand size = 7, antiderivative size = 28

$$\int \cosh(x) \coth(4x) dx = -\frac{1}{4} \operatorname{arctanh}(\cosh(x)) - \frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{2\sqrt{2}} + \cosh(x)$$

output

```
-1/4*arctanh(cosh(x))-1/4*arctanh(2^(1/2)*cosh(x))*2^(1/2)+cosh(x)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.61

$$\begin{aligned} \int \cosh(x) \coth(4x) dx = \frac{1}{4} & \left(-\sqrt{2} \operatorname{arctanh} \left(\sqrt{2} - i \tanh \left(\frac{x}{2} \right) \right) \right. \\ & - \sqrt{2} \operatorname{arctanh} \left(\sqrt{2} + i \tanh \left(\frac{x}{2} \right) \right) + 4 \cosh(x) \\ & \left. - \log \left(\cosh \left(\frac{x}{2} \right) \right) + \log \left(\sinh \left(\frac{x}{2} \right) \right) \right) \end{aligned}$$

input

```
Integrate[Cosh[x]*Coth[4*x],x]
```

output

$$\frac{(-(\text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[2]] - I*\text{Tanh}[x/2])) - \text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[2]] + I*\text{Tanh}[x/2] + 4*\text{Cosh}[x] - \text{Log}[\text{Cosh}[x/2]] + \text{Log}[\text{Sinh}[x/2]])}{4}$$
Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 2205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(x) \coth(4x) dx \\ & \quad \downarrow \text{3042} \\ & \int i \cos(ix) \cot(4ix) dx \\ & \quad \downarrow \text{26} \\ & i \int \cos(ix) \cot(4ix) dx \\ & \quad \downarrow \text{4879} \\ & - \int -\frac{8 \cosh^4(x) - 8 \cosh^2(x) + 1}{4(2 \cosh^4(x) - 3 \cosh^2(x) + 1)} d \cosh(x) \\ & \quad \downarrow \text{27} \\ & \frac{1}{4} \int \frac{8 \cosh^4(x) - 8 \cosh^2(x) + 1}{2 \cosh^4(x) - 3 \cosh^2(x) + 1} d \cosh(x) \\ & \quad \downarrow \text{2205} \\ & \frac{1}{4} \int \left(4 - \frac{3 - 4 \cosh^2(x)}{2 \cosh^4(x) - 3 \cosh^2(x) + 1} \right) d \cosh(x) \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \left(-\operatorname{arctanh}(\cosh(x)) - \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cosh(x)) + 4 \cosh(x) \right) \end{aligned}$$

input `Int[Cosh[x]*Coth[4*x],x]`

output `(-ArcTanh[Cosh[x]] - Sqrt[2]*ArcTanh[Sqrt[2]*Cosh[x]] + 4*Cosh[x])/4`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2205 `Int[(Px_)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := Int[ExpandIntegrand[Px/(a + b*x^2 + c*x^4), x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Px, x^2] && Expon[Px, x^2] > 1`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. $2(20) = 40$.

Time = 0.11 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.25

method	result	size
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\ln(e^x-1)}{4} - \frac{\ln(e^x+1)}{4} + \frac{\sqrt{2} \ln(e^{2x}-\sqrt{2}e^x+1)}{8} - \frac{\sqrt{2} \ln(e^{2x}+\sqrt{2}e^x+1)}{8}$	63

input `int(cosh(x)*coth(4*x),x,method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+1/4*ln(exp(x)-1)-1/4*ln(exp(x)+1)+1/8*2^(1/2)*ln(exp(2*x)-2^(1/2)*exp(x)+1)-1/8*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(20) = 40$.

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 3.61

$$\int \cosh(x) \coth(4x) dx$$

$$= \frac{4 \cosh(x)^2 + (\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(\frac{\cosh(x)^2 + \sinh(x)^2 - 2\sqrt{2} \cosh(x) + 2}{\cosh(x)^2 + \sinh(x)^2}\right) - 2(\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) + 1)}{8(\cosh(x) + \sinh(x))}$$

input `integrate(cosh(x)*coth(4*x),x, algorithm="fricas")`

output `1/8*(4*cosh(x)^2 + (sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log((cosh(x)^2 + sinh(x)^2 - 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2 + sinh(x)^2)) - 2*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) + 1) + 2*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) + 8*cosh(x)*sinh(x) + 4*sinh(x)^2 + 4)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \cosh(x) \coth(4x) dx = \int \cosh(x) \coth(4x) dx$$

input `integrate(cosh(x)*coth(4*x),x)`

output `Integral(cosh(x)*coth(4*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(20) = 40$.

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.50

$$\begin{aligned} \int \cosh(x) \coth(4x) dx &= -\frac{1}{8} \sqrt{2} \log \left(\sqrt{2} e^{(-x)} + e^{(-2x)} + 1 \right) \\ &\quad + \frac{1}{8} \sqrt{2} \log \left(-\sqrt{2} e^{(-x)} + e^{(-2x)} + 1 \right) + \frac{1}{2} e^{(-x)} \\ &\quad + \frac{1}{2} e^x - \frac{1}{4} \log \left(e^{(-x)} + 1 \right) + \frac{1}{4} \log \left(e^{(-x)} - 1 \right) \end{aligned}$$

input `integrate(cosh(x)*coth(4*x),x, algorithm="maxima")`

output `-1/8*sqrt(2)*log(sqrt(2)*e^(-x) + e^(-2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^(-x) + e^(-2*x) + 1) + 1/2*e^(-x) + 1/2*e^x - 1/4*log(e^(-x) + 1) + 1/4*log(e^(-x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. $2(20) = 40$.

Time = 0.15 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.39

$$\int \cosh(x) \coth(4x) dx = \frac{1}{8} \sqrt{2} \log \left(-\frac{\sqrt{2} - e^{(-x)} - e^x}{\sqrt{2} + e^{(-x)} + e^x} \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \\ - \frac{1}{8} \log(e^{(-x)} + e^x + 2) + \frac{1}{8} \log(e^{(-x)} + e^x - 2)$$

input `integrate(cosh(x)*coth(4*x),x, algorithm="giac")`

output `1/8*sqrt(2)*log(-(sqrt(2) - e^(-x) - e^x)/(sqrt(2) + e^(-x) + e^x)) + 1/2*
e^(-x) + 1/2*e^x - 1/8*log(e^(-x) + e^x + 2) + 1/8*log(e^(-x) + e^x - 2)`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.54

$$\int \cosh(x) \coth(4x) dx = \frac{\ln\left(\frac{1}{2} - \frac{e^x}{2}\right)}{4} - \frac{\ln\left(-\frac{e^x}{2} - \frac{1}{2}\right)}{4} + \frac{e^{-x}}{2} + \frac{e^x}{2} \\ - \frac{\sqrt{2} \ln\left(-\frac{e^{2x}}{8} - \frac{\sqrt{2}e^x}{8} - \frac{1}{8}\right)}{8} + \frac{\sqrt{2} \ln\left(\frac{\sqrt{2}e^x}{8} - \frac{e^{2x}}{8} - \frac{1}{8}\right)}{8}$$

input `int(coth(4*x)*cosh(x),x)`

output `log(1/2 - exp(x)/2)/4 - log(- exp(x)/2 - 1/2)/4 + exp(-x)/2 + exp(x)/2 - (
2^(1/2)*log(- exp(2*x)/8 - (2^(1/2)*exp(x))/8 - 1/8))/8 + (2^(1/2)*log((2^
(1/2)*exp(x))/8 - exp(2*x)/8 - 1/8))/8`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.86

$$\int \cosh(x) \coth(4x) dx$$
$$= \frac{4e^{2x} + e^x\sqrt{2}\log(e^{2x} - e^x\sqrt{2} + 1) - e^x\sqrt{2}\log(e^{2x} + e^x\sqrt{2} + 1) + 2e^x\log(e^x - 1) - 2e^x\log(e^x + 1) + 4}{8e^x}$$

input `int(cosh(x)*coth(4*x),x)`

output `(4*e**(2*x) + e**x*sqrt(2)*log(e**(2*x) - e**x*sqrt(2) + 1) - e**x*sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1) + 2*e**x*log(e**x - 1) - 2*e**x*log(e**x + 1) + 4)/(8*e**x)`

3.115 $\int \cosh(x) \coth(5x) dx$

Optimal result	813
Mathematica [A] (verified)	814
Rubi [A] (verified)	814
Maple [B] (verified)	816
Fricas [B] (verification not implemented)	817
Sympy [F]	817
Maxima [F]	818
Giac [B] (verification not implemented)	818
Mupad [B] (verification not implemented)	819
Reduce [F]	820

Optimal result

Integrand size = 7, antiderivative size = 84

$$\int \cosh(x) \coth(5x) dx = -\frac{1}{5} \operatorname{arctanh}(\cosh(x)) - \frac{1}{5} \sqrt{\frac{1}{2}(3 + \sqrt{5})} \operatorname{arctanh}\left(\sqrt{2(3 - \sqrt{5})} \cosh(x)\right) - \frac{1}{5} \sqrt{\frac{2}{3 + \sqrt{5}}} \operatorname{arctanh}\left(\sqrt{2(3 + \sqrt{5})} \cosh(x)\right) + \cosh(x)$$

output

$-1/5*\operatorname{arctanh}(\cosh(x))-1/5*(1/2+1/2*5^{(1/2)})*\operatorname{arctanh}((5^{(1/2)}-1)*\cosh(x))-1/5*2^{(1/2)/(3+5^{(1/2)})^{(1/2)}}*\operatorname{arctanh}((5^{(1/2)}+1)*\cosh(x))+\cosh(x)$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.58

$$\int \cosh(x) \coth(5x) dx = \frac{1}{100} \left(100 \cosh(x) - 20 \log \left(\cosh \left(\frac{x}{2} \right) \right) \right. \\ \left. + \sqrt{5} (-5 + \sqrt{5}) \log \left(1 - \sqrt{5} - 4 \cosh(x) \right) \right. \\ \left. + \sqrt{5} (5 + \sqrt{5}) \log \left(1 + \sqrt{5} - 4 \cosh(x) \right) \right. \\ \left. - \sqrt{5} (-5 + \sqrt{5}) \log \left(1 - \sqrt{5} + 4 \cosh(x) \right) \right. \\ \left. - \sqrt{5} (5 + \sqrt{5}) \log \left(1 + \sqrt{5} + 4 \cosh(x) \right) + 20 \log \left(\sinh \left(\frac{x}{2} \right) \right) \right)$$

input `Integrate[Cosh[x]*Coth[5*x],x]`

output `(100*Cosh[x] - 20*Log[Cosh[x/2]] + Sqrt[5]*(-5 + Sqrt[5])*Log[1 - Sqrt[5] - 4*Cosh[x]] + Sqrt[5]*(5 + Sqrt[5])*Log[1 + Sqrt[5] - 4*Cosh[x]] - Sqrt[5]*(-5 + Sqrt[5])*Log[1 - Sqrt[5] + 4*Cosh[x]] - Sqrt[5]*(5 + Sqrt[5])*Log[1 + Sqrt[5] + 4*Cosh[x]] + 20*Log[Sinh[x/2]])/100`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.31, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4879, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \coth(5x) dx \\ \downarrow 3042 \\ \int i \cos(ix) \cot(5ix) dx \\ \downarrow 26$$

$$\begin{aligned}
& i \int \cos(ix) \cot(5ix) dx \\
& \quad \downarrow \text{4879} \\
& - \int \frac{\cosh^2(x) (16 \cosh^4(x) - 20 \cosh^2(x) + 5)}{-16 \cosh^6(x) + 28 \cosh^4(x) - 13 \cosh^2(x) + 1} d \cosh(x) \\
& \quad \downarrow \text{2460} \\
& - \int \left(\frac{2(\cosh(x) - 1)}{5(4 \cosh^2(x) + 2 \cosh(x) - 1)} - \frac{1}{5(\cosh^2(x) - 1)} - \frac{2(\cosh(x) + 1)}{5(4 \cosh^2(x) - 2 \cosh(x) - 1)} - 1 \right) d \cosh(x) \\
& \quad \downarrow \text{2009} \\
& -\frac{1}{5} \operatorname{arctanh}(\cosh(x)) + \cosh(x) + \frac{1}{20} (1 - \sqrt{5}) \log(-4 \cosh(x) - \sqrt{5} + 1) + \\
& \frac{1}{20} (1 + \sqrt{5}) \log(-4 \cosh(x) + \sqrt{5} + 1) - \frac{1}{20} (1 - \sqrt{5}) \log(4 \cosh(x) - \sqrt{5} + 1) - \\
& \frac{1}{20} (1 + \sqrt{5}) \log(4 \cosh(x) + \sqrt{5} + 1)
\end{aligned}$$

input `Int[Cosh[x]*Coth[5*x],x]`

output `-1/5*ArcTanh[Cosh[x]] + Cosh[x] + ((1 - Sqrt[5])*Log[1 - Sqrt[5] - 4*Cosh[x]])/20 + ((1 + Sqrt[5])*Log[1 + Sqrt[5] - 4*Cosh[x]])/20 - ((1 - Sqrt[5])*Log[1 - Sqrt[5] + 4*Cosh[x]])/20 - ((1 + Sqrt[5])*Log[1 + Sqrt[5] + 4*Cosh[x]])/20`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]},
Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Q
x, x]]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] &&
!TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4879 `Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d
, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[Nonfree
Factors[Cos[v], x], u/Sin[v], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(47) = 94$.

Time = 0.12 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.26

method	result
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\ln(e^x-1)}{5} - \frac{\ln(e^x+1)}{5} - \frac{\ln\left(e^{2x} + \left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)e^x + 1\right)}{20} + \frac{\ln\left(e^{2x} + \left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)e^x + 1\right)\sqrt{5}}{20} - \frac{\ln\left(e^{2x} + \left(\frac{1}{2} + \frac{\sqrt{5}}{2}\right)e^x + 1\right)}{20}$

input `int(cosh(x)*coth(5*x), x, method=_RETURNVERBOSE)`

output `1/2*exp(x)+1/2*exp(-x)+1/5*ln(exp(x)-1)-1/5*ln(exp(x)+1)-1/20*ln(exp(2*x)+
(1/2-1/2*5^(1/2))*exp(x)+1)+1/20*ln(exp(2*x)+(1/2-1/2*5^(1/2))*exp(x)+1)*5
^(1/2)-1/20*ln(exp(2*x)+(1/2+1/2*5^(1/2))*exp(x)+1)-1/20*ln(exp(2*x)+(1/2+
1/2*5^(1/2))*exp(x)+1)*5^(1/2)+1/20*ln(exp(2*x)+(-1/2-1/2*5^(1/2))*exp(x)+
1)+1/20*ln(exp(2*x)+(-1/2-1/2*5^(1/2))*exp(x)+1)*5^(1/2)+1/20*ln(exp(2*x)+
(1/2*5^(1/2)-1/2)*exp(x)+1)-1/20*ln(exp(2*x)+(1/2*5^(1/2)-1/2)*exp(x)+1)*5
^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. $2(45) = 90$.

Time = 0.09 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.24

$$\int \cosh(x) \coth(5x) dx$$

$$= \frac{10 \cosh(x)^2 + (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x)) \log\left(-\frac{4(\sqrt{5}-1)\cosh(x)-4\cosh(x)^2-4\sinh(x)^2+\sqrt{5}-7}{2\cosh(x)^2+2\sinh(x)^2+2\cosh(x)+1}\right) + (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x)) \log\left(-\frac{4(\sqrt{5}+1)\cosh(x)-4\cosh(x)^2-4\sinh(x)^2-\sqrt{5}-7}{2\cosh(x)^2+2\sinh(x)^2-2\cosh(x)+1}\right) - (\cosh(x) + \sinh(x)) \log\left(\frac{2\cosh(x)^2+2\sinh(x)^2+2\cosh(x)+1}{\cosh(x)^2-2\cosh(x)\sinh(x)+\sinh(x)^2}\right) + (\cosh(x) + \sinh(x)) \log\left(\frac{2\cosh(x)^2+2\sinh(x)^2-2\cosh(x)+1}{\cosh(x)^2-2\cosh(x)\sinh(x)+\sinh(x)^2}\right) - 4(\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) - 1) + 20\cosh(x)\sinh(x) + 10\sinh(x)^2 + 10}{\cosh(x) + \sinh(x)}$$

input `integrate(cosh(x)*coth(5*x),x, algorithm="fricas")`

output `1/20*(10*cosh(x)^2 + (sqrt(5)*cosh(x) + sqrt(5)*sinh(x))*log(-(4*(sqrt(5) - 1)*cosh(x) - 4*cosh(x)^2 - 4*sinh(x)^2 + sqrt(5) - 7)/(2*cosh(x)^2 + 2*sinh(x)^2 + 2*cosh(x) + 1)) + (sqrt(5)*cosh(x) + sqrt(5)*sinh(x))*log(-(4*(sqrt(5) + 1)*cosh(x) - 4*cosh(x)^2 - 4*sinh(x)^2 - sqrt(5) - 7)/(2*cosh(x)^2 + 2*sinh(x)^2 - 2*cosh(x) + 1)) - (cosh(x) + sinh(x))*log((2*cosh(x)^2 + 2*sinh(x)^2 + 2*cosh(x) + 1)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + (cosh(x) + sinh(x))*log((2*cosh(x)^2 + 2*sinh(x)^2 - 2*cosh(x) + 1)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) - 4*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) + 20*cosh(x)*sinh(x) + 10*sinh(x)^2 + 10)/(cosh(x) + sinh(x))`

Sympy [F]

$$\int \cosh(x) \coth(5x) dx = \int \cosh(x) \coth(5x) dx$$

input `integrate(cosh(x)*coth(5*x),x)`

output `Integral(cosh(x)*coth(5*x), x)`

Maxima [F]

$$\int \cosh(x) \coth(5x) dx = \int \cosh(x) \coth(5x) dx$$

input `integrate(cosh(x)*coth(5*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) + 1)*e^(-x) - 1/5*integrate((e^(3*x) + e^(2*x) + e^x + 1)*e^x / (e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) + 1/5*integrate((e^(3*x) - e^(2*x) + e^x - 1)*e^x / (e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x) + 3/10*integrate(e^(3*x) / (e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) + 3/10*integrate(e^(3*x) / (e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x) + 1/10*integrate(e^(2*x) / (e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) - 1/10*integrate(e^(2*x) / (e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x) - 1/10*integrate(e^x / (e^(4*x) + e^(3*x) + e^(2*x) + e^x + 1), x) - 1/10*integrate(e^x / (e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x) - 1/5*log(e^x + 1) + 1/5*log(e^x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(45) = 90$.

Time = 0.14 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.87

$$\begin{aligned} \int \cosh(x) \coth(5x) dx &= \frac{1}{20} \sqrt{5} \log \left(-\frac{\sqrt{5} - 2e^{(-x)} - 2e^x + 1}{\sqrt{5} + 2e^{(-x)} + 2e^x - 1} \right) \\ &+ \frac{1}{20} \sqrt{5} \log \left(-\frac{\sqrt{5} - 2e^{(-x)} - 2e^x - 1}{\sqrt{5} + 2e^{(-x)} + 2e^x + 1} \right) + \frac{1}{2} e^{(-x)} \\ &+ \frac{1}{2} e^x - \frac{1}{20} \log \left((e^{(-x)} + e^x)^2 + e^{(-x)} + e^x - 1 \right) \\ &+ \frac{1}{20} \log \left((e^{(-x)} + e^x)^2 - e^{(-x)} - e^x - 1 \right) \\ &- \frac{1}{10} \log (e^{(-x)} + e^x + 2) + \frac{1}{10} \log (e^{(-x)} + e^x - 2) \end{aligned}$$

input `integrate(cosh(x)*coth(5*x),x, algorithm="giac")`

output

```
1/20*sqrt(5)*log(-(sqrt(5) - 2*e^(-x) - 2*e^x + 1)/(sqrt(5) + 2*e^(-x) + 2
*e^x - 1)) + 1/20*sqrt(5)*log(-(sqrt(5) - 2*e^(-x) - 2*e^x - 1)/(sqrt(5) +
2*e^(-x) + 2*e^x + 1)) + 1/2*e^(-x) + 1/2*e^x - 1/20*log((e^(-x) + e^x)^2
+ e^(-x) + e^x - 1) + 1/20*log((e^(-x) + e^x)^2 - e^(-x) - e^x - 1) - 1/1
0*log(e^(-x) + e^x + 2) + 1/10*log(e^(-x) + e^x - 2)
```

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.70

$$\int \cosh(x) \coth(5x) dx = \frac{\ln(10 - 10e^x)}{5} - \frac{\ln(-10e^x - 10)}{5} + \frac{e^{-x}}{2} + \frac{e^x}{2} - \ln\left(-e^{2x} - 10e^x\left(\frac{\sqrt{5}}{20} - \frac{1}{20}\right) - 1\right)\left(\frac{\sqrt{5}}{20} - \frac{1}{20}\right) + \ln\left(10e^x\left(\frac{\sqrt{5}}{20} - \frac{1}{20}\right) - e^{2x} - 1\right)\left(\frac{\sqrt{5}}{20} - \frac{1}{20}\right) - \ln\left(-e^{2x} - 10e^x\left(\frac{\sqrt{5}}{20} + \frac{1}{20}\right) - 1\right)\left(\frac{\sqrt{5}}{20} + \frac{1}{20}\right) + \ln\left(10e^x\left(\frac{\sqrt{5}}{20} + \frac{1}{20}\right) - e^{2x} - 1\right)\left(\frac{\sqrt{5}}{20} + \frac{1}{20}\right)$$

input

```
int(coth(5*x)*cosh(x),x)
```

output

```
log(10 - 10*exp(x))/5 - log(- 10*exp(x) - 10)/5 + exp(-x)/2 + exp(x)/2 - 1
og(- exp(2*x) - 10*exp(x)*(5^(1/2)/20 - 1/20) - 1)*(5^(1/2)/20 - 1/20) + 1
og(10*exp(x)*(5^(1/2)/20 - 1/20) - exp(2*x) - 1)*(5^(1/2)/20 - 1/20) - log
(- exp(2*x) - 10*exp(x)*(5^(1/2)/20 + 1/20) - 1)*(5^(1/2)/20 + 1/20) + log
(10*exp(x)*(5^(1/2)/20 + 1/20) - exp(2*x) - 1)*(5^(1/2)/20 + 1/20)
```

Reduce [F]

$$\int \cosh(x) \coth(5x) dx = \frac{e^{2x} + 2e^x \left(\int \frac{e^x}{e^{10x}-1} dx \right) + 2e^x \left(\int \frac{1}{e^{11x}-e^x} dx \right) - 1}{2e^x}$$

input

```
int(cosh(x)*coth(5*x),x)
```

output

```
(e**(2*x) + 2*e**x*int(e**x/(e**(10*x) - 1),x) + 2*e**x*int(1/(e**(11*x) - e**x),x) - 1)/(2*e**x)
```

3.116 $\int \cosh(x) \coth(6x) dx$

Optimal result	821
Mathematica [C] (verified)	822
Rubi [A] (verified)	822
Maple [B] (verified)	824
Fricas [B] (verification not implemented)	825
Sympy [F]	825
Maxima [F]	826
Giac [B] (verification not implemented)	826
Mupad [B] (verification not implemented)	827
Reduce [B] (verification not implemented)	827

Optimal result

Integrand size = 7, antiderivative size = 38

$$\int \cosh(x) \coth(6x) dx = -\frac{1}{6} \operatorname{arctanh}(\cosh(x)) - \frac{1}{6} \operatorname{arctanh}(2 \cosh(x)) - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{2\sqrt{3}} + \cosh(x)$$

output

```
-1/6*arctanh(cosh(x))-1/6*arctanh(2*cosh(x))-1/6*arctanh(2/3*cosh(x)*3^(1/2))*3^(1/2)+cosh(x)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.50

$$\int \cosh(x) \coth(6x) dx = \frac{1}{12} \left(-2\sqrt{3} \operatorname{arctanh} \left(\frac{2 - i \tanh \left(\frac{x}{2} \right)}{\sqrt{3}} \right) - 2\sqrt{3} \operatorname{arctanh} \left(\frac{2 + i \tanh \left(\frac{x}{2} \right)}{\sqrt{3}} \right) + 12 \cosh(x) - 2 \log \left(\cosh \left(\frac{x}{2} \right) \right) + \log(1 - 2 \cosh(x)) - \log(1 + 2 \cosh(x)) + 2 \log \left(\sinh \left(\frac{x}{2} \right) \right) \right)$$

input

```
Integrate[Cosh[x]*Coth[6*x],x]
```

output

```
(-2*Sqrt[3]*ArcTanh[(2 - I*Tanh[x/2])/Sqrt[3]] - 2*Sqrt[3]*ArcTanh[(2 + I*Tanh[x/2])/Sqrt[3]] + 12*Cosh[x] - 2*Log[Cosh[x/2]] + Log[1 - 2*Cosh[x]] - Log[1 + 2*Cosh[x]] + 2*Log[Sinh[x/2]])/12
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \coth(6x) dx$$

↓ 3042

$$\int i \cos(ix) \cot(6ix) dx$$

↓ 26

$$\begin{aligned}
 & i \int \cos(ix) \cot(6ix) dx \\
 & \quad \downarrow 4879 \\
 & - \int - \frac{-32 \cosh^6(x) + 48 \cosh^4(x) - 18 \cosh^2(x) + 1}{2(-16 \cosh^6(x) + 32 \cosh^4(x) - 19 \cosh^2(x) + 3)} d \cosh(x) \\
 & \quad \downarrow 27 \\
 & \frac{1}{2} \int \frac{-32 \cosh^6(x) + 48 \cosh^4(x) - 18 \cosh^2(x) + 1}{-16 \cosh^6(x) + 32 \cosh^4(x) - 19 \cosh^2(x) + 3} d \cosh(x) \\
 & \quad \downarrow 2460 \\
 & \frac{1}{2} \int \left(\frac{2}{4 \cosh^2(x) - 3} + \frac{2}{3(4 \cosh^2(x) - 1)} + 2 + \frac{1}{3(\cosh^2(x) - 1)} \right) d \cosh(x) \\
 & \quad \downarrow 2009 \\
 & \frac{1}{2} \left(-\frac{1}{3} \operatorname{arctanh}(\cosh(x)) - \frac{1}{3} \operatorname{arctanh}(2 \cosh(x)) - \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{\sqrt{3}} + 2 \cosh(x) \right)
 \end{aligned}$$

input `Int[Cosh[x]*Coth[6*x],x]`

output `(-1/3*ArcTanh[Cosh[x]] - ArcTanh[2*Cosh[x]]/3 - ArcTanh[(2*Cosh[x])/Sqrt[3]]/Sqrt[3] + 2*Cosh[x])/2`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]},
Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Q
x, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] &&
!TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4879

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFa
ctors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d
, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[Nonfree
Factors[Cos[v], x], u/Sin[v], x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(28) = 56$.

Time = 0.15 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.29

method	result
risch	$\frac{e^x}{2} + \frac{e^{-x}}{2} + \frac{\ln(e^x-1)}{6} - \frac{\ln(e^x+1)}{6} + \frac{\ln(e^{2x}-e^x+1)}{12} - \frac{\ln(e^{2x}+e^x+1)}{12} + \frac{\sqrt{3} \ln(e^{2x}-\sqrt{3}e^x+1)}{12} - \frac{\sqrt{3} \ln(e^{2x}+\sqrt{3}e^x+1)}{12}$

input

```
int(cosh(x)*coth(6*x), x, method=_RETURNVERBOSE)
```

output

```
1/2*exp(x)+1/2*exp(-x)+1/6*ln(exp(x)-1)-1/6*ln(exp(x)+1)+1/12*ln(exp(2*x)-
exp(x)+1)-1/12*ln(exp(2*x)+exp(x)+1)+1/12*3^(1/2)*ln(exp(2*x)-3^(1/2)*exp(
x)+1)-1/12*3^(1/2)*ln(exp(2*x)+3^(1/2)*exp(x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(28) = 56$.

Time = 0.09 (sec) , antiderivative size = 157, normalized size of antiderivative = 4.13

$$\int \cosh(x) \coth(6x) dx$$

$$= \frac{6 \cosh(x)^2 + (\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x)) \log\left(\frac{2 \cosh(x)^2 + 2 \sinh(x)^2 - 4\sqrt{3} \cosh(x) + 5}{2 \cosh(x)^2 + 2 \sinh(x)^2 - 1}\right) - (\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x))}{1}$$

input `integrate(cosh(x)*coth(6*x),x, algorithm="fricas")`

output

```
1/12*(6*cosh(x)^2 + (sqrt(3)*cosh(x) + sqrt(3)*sinh(x))*log((2*cosh(x)^2 +
2*sinh(x)^2 - 4*sqrt(3)*cosh(x) + 5)/(2*cosh(x)^2 + 2*sinh(x)^2 - 1)) - (
cosh(x) + sinh(x))*log((2*cosh(x) + 1)/(cosh(x) - sinh(x))) + (cosh(x) + s
inh(x))*log((2*cosh(x) - 1)/(cosh(x) - sinh(x))) - 2*(cosh(x) + sinh(x))*l
og(cosh(x) + sinh(x) + 1) + 2*(cosh(x) + sinh(x))*log(cosh(x) + sinh(x) -
1) + 12*cosh(x)*sinh(x) + 6*sinh(x)^2 + 6)/(cosh(x) + sinh(x))
```

Sympy [F]

$$\int \cosh(x) \coth(6x) dx = \int \cosh(x) \coth(6x) dx$$

input `integrate(cosh(x)*coth(6*x),x)`

output

```
Integral(cosh(x)*coth(6*x), x)
```

Maxima [F]

$$\int \cosh(x) \coth(6x) dx = \int \cosh(x) \coth(6x) dx$$

input `integrate(cosh(x)*coth(6*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) + 1)*e^(-x) + 1/2*integrate((e^(3*x) - e^x)/(e^(4*x) - e^(2*x) + 1), x) - 1/12*log(e^(2*x) + e^x + 1) + 1/12*log(e^(2*x) - e^x + 1) - 1/6*log(e^x + 1) + 1/6*log(e^x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 89 vs. 2(28) = 56.

Time = 0.12 (sec) , antiderivative size = 89, normalized size of antiderivative = 2.34

$$\begin{aligned} \int \cosh(x) \coth(6x) dx = & \frac{1}{12} \sqrt{3} \log \left(-\frac{\sqrt{3} - e^{(-x)} - e^x}{\sqrt{3} + e^{(-x)} + e^x} \right) + \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x \\ & - \frac{1}{12} \log(e^{(-x)} + e^x + 2) - \frac{1}{12} \log(e^{(-x)} + e^x + 1) \\ & + \frac{1}{12} \log(e^{(-x)} + e^x - 1) + \frac{1}{12} \log(e^{(-x)} + e^x - 2) \end{aligned}$$

input `integrate(cosh(x)*coth(6*x),x, algorithm="giac")`

output `1/12*sqrt(3)*log(-(sqrt(3) - e^(-x) - e^x)/(sqrt(3) + e^(-x) + e^x)) + 1/2*e^(-x) + 1/2*e^x - 1/12*log(e^(-x) + e^x + 2) - 1/12*log(e^(-x) + e^x + 1) + 1/12*log(e^(-x) + e^x - 1) + 1/12*log(e^(-x) + e^x - 2)`

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.66

$$\int \cosh(x) \coth(6x) dx = \frac{\ln\left(\frac{1}{3} - \frac{e^x}{3}\right)}{6} - \frac{\ln\left(-\frac{e^x}{3} - \frac{1}{3}\right)}{6} + \frac{e^{-x}}{2}$$

$$- \frac{\ln\left(-\frac{e^{2x}}{36} - \frac{e^x}{36} - \frac{1}{36}\right)}{12} + \frac{\ln\left(\frac{e^x}{36} - \frac{e^{2x}}{36} - \frac{1}{36}\right)}{12} + \frac{e^x}{2}$$

$$- \frac{\sqrt{3} \ln\left(-\frac{e^{2x}}{12} - \frac{\sqrt{3}e^x}{12} - \frac{1}{12}\right)}{12} + \frac{\sqrt{3} \ln\left(\frac{\sqrt{3}e^x}{12} - \frac{e^{2x}}{12} - \frac{1}{12}\right)}{12}$$

input `int(coth(6*x)*cosh(x), x)`output `log(1/3 - exp(x)/3)/6 - log(- exp(x)/3 - 1/3)/6 + exp(-x)/2 - log(- exp(2*x)/36 - exp(x)/36 - 1/36)/12 + log(exp(x)/36 - exp(2*x)/36 - 1/36)/12 + exp(x)/2 - (3^(1/2)*log(- exp(2*x)/12 - (3^(1/2)*exp(x))/12 - 1/12))/12 + (3^(1/2)*log((3^(1/2)*exp(x))/12 - exp(2*x)/12 - 1/12))/12`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.97

$$\int \cosh(x) \coth(6x) dx$$

$$= \frac{6e^{2x} + e^x\sqrt{3}\log(e^{2x} - e^x\sqrt{3} + 1) - e^x\sqrt{3}\log(e^{2x} + e^x\sqrt{3} + 1) - e^x\log(e^{2x} + e^x + 1) + e^x\log(e^{2x} - e^x + 1)}{12e^x}$$

input `int(cosh(x)*coth(6*x), x)`output `(6*e**(2*x) + e**x*sqrt(3)*log(e**(2*x) - e**x*sqrt(3) + 1) - e**x*sqrt(3)*log(e**(2*x) + e**x*sqrt(3) + 1) - e**x*log(e**(2*x) + e**x + 1) + e**x*log(e**(2*x) - e**x + 1) + 2*e**x*log(e**x - 1) - 2*e**x*log(e**x + 1) + 6)/(12*e**x)`

3.117 $\int \cosh(x) \coth(nx) dx$

Optimal result	828
Mathematica [A] (verified)	828
Rubi [A] (verified)	829
Maple [F]	830
Fricas [F]	830
Sympy [F]	831
Maxima [F]	831
Giac [F]	831
Mupad [F(-1)]	832
Reduce [F]	832

Optimal result

Integrand size = 7, antiderivative size = 62

$$\int \cosh(x) \coth(nx) dx = e^{-x} \operatorname{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, e^{2nx} \right) - e^x \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), e^{2nx} \right) + \sinh(x)$$

output `hypergeom([1, -1/2/n], [1-1/2/n], exp(2*n*x))/exp(x)-exp(x)*hypergeom([1, 1/2/n], [1+1/2/n], exp(2*n*x))+sinh(x)`

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.21

$$\int \cosh(x) \coth(nx) dx = \frac{1}{2} \left(-e^{-x} + e^x + 2e^{-x} \operatorname{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, e^{2nx} \right) - 2e^x \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2n}, 1 + \frac{1}{2n}, e^{2nx} \right) \right)$$

input `Integrate[Cosh[x]*Coth[n*x], x]`

output

$$\frac{(-E^{-x}) + E^x + (2 \operatorname{Hypergeometric2F1}[1, -1/2*1/n, 1 - 1/(2*n), E^{(2*n*x)}]) / E^x - 2 * E^x * \operatorname{Hypergeometric2F1}[1, 1/(2*n), 1 + 1/(2*n), E^{(2*n*x)}])}{2}$$
Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.23, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6136, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \coth(nx) dx$$

$$\downarrow 6136$$

$$\int \left(-\frac{e^{-x}}{1 - e^{2nx}} - \frac{e^x}{1 - e^{2nx}} + \frac{e^{-x}}{2} + \frac{e^x}{2} \right) dx$$

$$\downarrow 2009$$

$$e^{-x} \operatorname{Hypergeometric2F1} \left(1, -\frac{1}{2n}, 1 - \frac{1}{2n}, e^{2nx} \right) - e^x \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2n}, \frac{1}{2} \left(2 + \frac{1}{n} \right), e^{2nx} \right) - \frac{e^{-x}}{2} + \frac{e^x}{2}$$

input

$$\text{Int}[\text{Cosh}[x] * \text{Coth}[n*x], x]$$

output

$$-1/2*1/E^x + E^x/2 + \operatorname{Hypergeometric2F1}[1, -1/2*1/n, 1 - 1/(2*n), E^{(2*n*x)}] / E^x - E^x * \operatorname{Hypergeometric2F1}[1, 1/(2*n), (2 + n^{(-1)})/2, E^{(2*n*x)}]$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6136 `Int[Cosh[(a_.) + (b_.)*(x_)]*Coth[(c_.) + (d_.)*(x_)], x_Symbol] := Int[1/(E^(a + b*x)*2) + E^(a + b*x)/2 - 1/(E^(a + b*x)*(1 - E^(2*(c + d*x)))) - E^(a + b*x)/(1 - E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \cosh(x) \coth(nx) dx$$

input `int(cosh(x)*coth(n*x),x)`

output `int(cosh(x)*coth(n*x),x)`

Fricas [F]

$$\int \cosh(x) \coth(nx) dx = \int \cosh(x) \coth(nx) dx$$

input `integrate(cosh(x)*coth(n*x),x, algorithm="fricas")`

output `integral(cosh(x)*coth(n*x), x)`

Sympy [F]

$$\int \cosh(x) \coth(nx) dx = \int \cosh(x) \coth(nx) dx$$

input `integrate(cosh(x)*coth(n*x),x)`

output `Integral(cosh(x)*coth(n*x), x)`

Maxima [F]

$$\int \cosh(x) \coth(nx) dx = \int \cosh(x) \coth(nx) dx$$

input `integrate(cosh(x)*coth(n*x),x, algorithm="maxima")`

output `1/2*(e^(2*x) - 1)*e^(-x) - 1/2*integrate((e^(2*x) + 1)/(e^(n*x + x) + e^x), x) + 1/2*integrate((e^(2*x) + 1)/(e^(n*x + x) - e^x), x)`

Giac [F]

$$\int \cosh(x) \coth(nx) dx = \int \cosh(x) \coth(nx) dx$$

input `integrate(cosh(x)*coth(n*x),x, algorithm="giac")`

output `integrate(cosh(x)*coth(n*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(x) \coth(nx) dx = \int \coth(nx) \cosh(x) dx$$

input `int(coth(n*x)*cosh(x),x)`output `int(coth(n*x)*cosh(x),x)`**Reduce [F]**

$$\int \cosh(x) \coth(nx) dx = \frac{e^{2x} + 2e^x \left(\int \frac{e^x}{e^{2nx}-1} dx \right) + 2e^x \left(\int \frac{1}{e^{2nx+x}-e^x} dx \right) - 1}{2e^x}$$

input `int(cosh(x)*coth(n*x),x)`output `(e**(2*x) + 2*e**x*int(e**x/(e**(2*n*x) - 1),x) + 2*e**x*int(1/(e**(2*n*x + x) - e**x),x) - 1)/(2*e**x)`

3.118 $\int \cosh(a + bx) \coth(c + bx) dx$

Optimal result	833
Mathematica [C] (verified)	833
Rubi [A] (verified)	834
Maple [B] (verified)	835
Fricas [B] (verification not implemented)	836
Sympy [F]	837
Maxima [B] (verification not implemented)	837
Giac [B] (verification not implemented)	838
Mupad [B] (verification not implemented)	838
Reduce [B] (verification not implemented)	839

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \cosh(a + bx) \coth(c + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(a - c)}{b} + \frac{\cosh(a + bx)}{b}$$

output `-arctanh(cosh(b*x+c))*cosh(a-c)/b+cosh(b*x+a)/b`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.21

$$\int \cosh(a + bx) \coth(c + bx) dx$$

$$= -\frac{2i \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh(c) \cosh\left(\frac{bx}{2}\right) + \sinh(c) \sinh\left(\frac{bx}{2}\right)\right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a - c)}{b} + \frac{\cosh(a) \cosh(bx)}{b} + \frac{\sinh(a) \sinh(bx)}{b}$$

input `Integrate[Cosh[a + b*x]*Coth[c + b*x],x]`

output

$$\frac{((-2*I)*\text{ArcTan}[\frac{(\text{Cosh}[c] - \text{Sinh}[c])*(\text{Cosh}[c]*\text{Cosh}[(b*x)/2] + \text{Sinh}[c]*\text{Sinh}[(b*x)/2])}{I*\text{Cosh}[c]*\text{Cosh}[(b*x)/2] - I*\text{Cosh}[(b*x)/2]*\text{Sinh}[c]})*\text{Cosh}[a - c]}{b} + \frac{\text{Cosh}[a]*\text{Cosh}[b*x]}{b} + \frac{\text{Sinh}[a]*\text{Sinh}[b*x]}{b}$$
Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6155, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(a + bx) \coth(bx + c) dx \\ & \quad \downarrow \text{6155} \\ & \cosh(a - c) \int \text{csch}(c + bx) dx + \int \sinh(a + bx) dx \\ & \quad \downarrow \text{3042} \\ & \cosh(a - c) \int i \csc(ic + ibx) dx + \int -i \sin(ia + ibx) dx \\ & \quad \downarrow \text{26} \\ & i \cosh(a - c) \int \csc(ic + ibx) dx - i \int \sin(ia + ibx) dx \\ & \quad \downarrow \text{3118} \\ & \frac{\cosh(a + bx)}{b} + i \cosh(a - c) \int \csc(ic + ibx) dx \\ & \quad \downarrow \text{4257} \\ & \frac{\cosh(a + bx)}{b} - \frac{\cosh(a - c) \text{arctanh}(\cosh(bx + c))}{b} \end{aligned}$$

input

$$\text{Int}[\text{Cosh}[a + b*x]*\text{Coth}[c + b*x], x]$$

output $-\left(\frac{\text{ArcTanh}[\text{Cosh}[c + b*x]]*\text{Cosh}[a - c]}{b}\right) + \text{Cosh}[a + b*x]/b$

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6155 `Int[Cosh[v_] * Coth[w_]^(n_), x_Symbol] := Int[Sinh[v] * Coth[w]^(n - 1), x] + Simp[Cosh[v - w] Int[Csch[w] * Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 154 vs. $2(29) = 58$.

Time = 0.13 (sec) , antiderivative size = 155, normalized size of antiderivative = 5.34

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*coth(b*x+c),x,method=_RETURNVERBOSE)`

output

```
1/2/b*exp(b*x+a)+1/2/b*exp(-b*x-a)-1/2*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)
*exp(2*a)-1/2*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*c)+1/2*ln(exp(b*x+
a)-exp(a-c))/b*exp(-a-c)*exp(2*a)+1/2*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*
exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 439 vs. $2(29) = 58$.

Time = 0.11 (sec) , antiderivative size = 439, normalized size of antiderivative = 15.14

$$\int \cosh(a + bx) \coth(c + bx) dx$$

$$= \frac{\cosh(bx + c)^2 \cosh(-a + c)^2 - 2 \cosh(bx + c)^2 \cosh(-a + c) \sinh(-a + c) + \cosh(bx + c)^2 \sinh(-a + c)^2}{\cosh(bx + c)^2 + \sinh(bx + c)^2}$$

input

```
integrate(cosh(b*x+a)*coth(b*x+c),x, algorithm="fricas")
```

output

```
1/2*(cosh(b*x + c)^2*cosh(-a + c)^2 - 2*cosh(b*x + c)^2*cosh(-a + c)*sinh(
-a + c) + cosh(b*x + c)^2*sinh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)
)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^2 + (2*cosh(b*x + c)*cosh(-
a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*
cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a +
c)^2 + 1)*sinh(b*x + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) - (2*cosh(
b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(
-a + c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a +
c) + sinh(-a + c)^2 + 1)*sinh(b*x + c))*log(cosh(b*x + c) + sinh(b*x + c)
- 1) + 2*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh
(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) + 1)/(b*cosh(b*x +
c)*cosh(-a + c) - b*cosh(b*x + c)*sinh(-a + c) + (b*cosh(-a + c) - b*sinh(
-a + c))*sinh(b*x + c))
```

Sympy [F]

$$\int \cosh(a + bx) \coth(c + bx) dx = \int \cosh(a + bx) \coth(bx + c) dx$$

input `integrate(cosh(b*x+a)*coth(b*x+c),x)`

output `Integral(cosh(a + b*x)*coth(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. $2(29) = 58$.

Time = 0.04 (sec) , antiderivative size = 90, normalized size of antiderivative = 3.10

$$\begin{aligned} \int \cosh(a + bx) \coth(c + bx) dx = & -\frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} \\ & + \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} \\ & + \frac{e^{(bx+a)}}{2b} + \frac{e^{(-bx-a)}}{2b} \end{aligned}$$

input `integrate(cosh(b*x+a)*coth(b*x+c),x, algorithm="maxima")`

output `-1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/2*e^(b*x + a)/b + 1/2*e^(-b*x - a)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 91 vs. $2(29) = 58$.

Time = 0.13 (sec) , antiderivative size = 91, normalized size of antiderivative = 3.14

$$\int \cosh(a + bx) \coth(c + bx) dx = \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - (e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) - e^{(bx+a)}}{2b}$$

input `integrate(cosh(b*x+a)*coth(b*x+c),x, algorithm="giac")`

output `-1/2*((e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(e^(b*x + a + c) + e^a) - (e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + a + c) - e^a)) - e^(b*x + a) - e^(-b*x - a))/b`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 139, normalized size of antiderivative = 4.79

$$\int \cosh(a + bx) \coth(c + bx) dx = \frac{e^{a+bx}}{2b} + \frac{e^{-a-bx}}{2b} - \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2} + e^{2a} e^{-2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{-b^2}}$$

input `int(cosh(a + b*x)*coth(c + b*x),x)`

output `exp(a + b*x)/(2*b) + exp(- a - b*x)/(2*b) - (atan((exp(-a)*exp(2*c))*exp(b*x)*((-b^2)^(1/2) + exp(2*a)*exp(-2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c))*(2*exp(2*a)*exp(-2*c) + exp(4*a)*exp(-4*c) + 1)^(1/2)))*(exp(2*c - 2*a)*(2*exp(2*a - 2*c) + exp(4*a - 4*c) + 1)^(1/2))/(-b^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 112, normalized size of antiderivative = 3.86

$$\int \cosh(a + bx) \coth(c + bx) dx$$

$$= \frac{e^{2bx+2a+c} + e^{bx+2a} \log(e^{bx+c} - 1) - e^{bx+2a} \log(e^{bx+c} + 1) + e^{bx+2c} \log(e^{bx+c} - 1) - e^{bx+2c} \log(e^{bx+c} + 1) + e^{2c}}{2e^{bx+a+cb}}$$

input `int(cosh(b*x+a)*coth(b*x+c),x)`output `(e**(2*a + 2*b*x + c) + e**(2*a + b*x)*log(e**(b*x + c) - 1) - e**(2*a + b*x)*log(e**(b*x + c) + 1) + e**(b*x + 2*c)*log(e**(b*x + c) - 1) - e**(b*x + 2*c)*log(e**(b*x + c) + 1) + e**c)/(2*e**(a + b*x + c)*b)`

3.119 $\int \cosh(a + bx) \coth^2(c + bx) dx$

Optimal result	840
Mathematica [C] (verified)	840
Rubi [A] (verified)	841
Maple [B] (verified)	843
Fricas [B] (verification not implemented)	844
Sympy [F]	845
Maxima [B] (verification not implemented)	845
Giac [B] (verification not implemented)	846
Mupad [B] (verification not implemented)	846
Reduce [B] (verification not implemented)	847

Optimal result

Integrand size = 15, antiderivative size = 46

$$\int \cosh(a + bx) \coth^2(c + bx) dx = -\frac{\cosh(a - c)\operatorname{csch}(c + bx)}{b} - \frac{\operatorname{arctanh}(\cosh(c + bx)) \sinh(a - c)}{b} + \frac{\sinh(a + bx)}{b}$$

output `-cosh(a-c)*csch(b*x+c)/b-arctanh(cosh(b*x+c))*sinh(a-c)/b+sinh(b*x+a)/b`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.39

$$\begin{aligned} & \int \cosh(a + bx) \coth^2(c + bx) dx \\ &= -\frac{\cosh(a - c)\operatorname{csch}(c + bx)}{b} + \frac{\cosh(bx) \sinh(a)}{b} \\ & \quad - \frac{2i \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh(c) \cosh\left(\frac{bx}{2}\right) + \sinh(c) \sinh\left(\frac{bx}{2}\right)\right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \sinh(a - c)}{b} \\ & \quad + \frac{\cosh(a) \sinh(bx)}{b} \end{aligned}$$

input `Integrate[Cosh[a + b*x]*Coth[c + b*x]^2,x]`

output
$$-\left(\frac{\text{Cosh}[a - c] \cdot \text{Csch}[c + b*x]}{b}\right) + \frac{\text{Cosh}[b*x] \cdot \text{Sinh}[a]}{b} - \left(\frac{2 \cdot \text{I} \cdot \text{ArcTan}\left[\frac{(\text{Cosh}[c] - \text{Sinh}[c]) \cdot (\text{Cosh}[c] \cdot \text{Cosh}[(b*x)/2] + \text{Sinh}[c] \cdot \text{Sinh}[(b*x)/2])}{\text{I} \cdot \text{Cosh}[c] \cdot \text{Cosh}[(b*x)/2] - \text{I} \cdot \text{Cosh}[(b*x)/2] \cdot \text{Sinh}[c]}\right]}{b} + \frac{\text{Cosh}[a] \cdot \text{Sinh}[b*x]}{b}\right)$$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {6155, 3042, 3086, 24, 6156, 3042, 26, 3117, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(a + bx) \coth^2(bx + c) dx \\ & \quad \downarrow \text{6155} \\ & \int \coth(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int \coth(c + bx) \operatorname{csch}(c + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int \coth(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3086} \\ & \int \coth(c + bx) \sinh(a + bx) dx - \frac{i \cosh(a - c) \int 1d(-i \operatorname{csch}(c + bx))}{b} \\ & \quad \downarrow \text{24} \\ & \int \coth(c + bx) \sinh(a + bx) dx - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\ & \quad \downarrow \text{6156} \\ & \sinh(a - c) \int \operatorname{csch}(c + bx) dx + \int \cosh(a + bx) dx - \frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
& \sinh(a-c) \int i \csc(ic+ibx) dx + \int \sin\left(ia+ibx+\frac{\pi}{2}\right) dx - \frac{\cosh(a-c)\operatorname{csch}(bx+c)}{b} \\
& \quad \downarrow 26 \\
& i \sinh(a-c) \int \csc(ic+ibx) dx + \int \sin\left(ia+ibx+\frac{\pi}{2}\right) dx - \frac{\cosh(a-c)\operatorname{csch}(bx+c)}{b} \\
& \quad \downarrow 3117 \\
& i \sinh(a-c) \int \csc(ic+ibx) dx - \frac{\cosh(a-c)\operatorname{csch}(bx+c)}{b} + \frac{\sinh(a+bx)}{b} \\
& \quad \downarrow 4257 \\
& -\frac{\sinh(a-c)\operatorname{arctanh}(\cosh(bx+c))}{b} - \frac{\cosh(a-c)\operatorname{csch}(bx+c)}{b} + \frac{\sinh(a+bx)}{b}
\end{aligned}$$

input `Int[Cosh[a + b*x]*Coth[c + b*x]^2,x]`

output `-((Cosh[a - c]*Csch[c + b*x])/b) - (ArcTanh[Cosh[c + b*x]]*Sinh[a - c])/b + Sinh[a + b*x]/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]`
`/; FreeQ[{c, d}, x]`

rule 6155 `Int[Cosh[v_]*Coth[w_]^(n_.), x_Symbol] := Int[Sinh[v]*Coth[w]^(n - 1), x] +`
`Simp[Cosh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ`
`[w, v] && FreeQ[v - w, x]`

rule 6156 `Int[Coth[w_]^(n_.)*Sinh[v_], x_Symbol] := Int[Cosh[v]*Coth[w]^(n - 1), x] +`
`Simp[Sinh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ`
`[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(46) = 92$.

Time = 0.14 (sec) , antiderivative size = 195, normalized size of antiderivative = 4.24

method	result
risch	$\frac{e^{bx+a}}{2b} - \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(e^{2a}+e^{2c})}{b(-e^{2bx+2a+2c}+e^{2a})} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*coth(b*x+c)^2,x,method=_RETURNVERBOSE)`

output $\frac{1}{2}/b*\exp(b*x+a)-1/2/b*\exp(-b*x-a)+1/b*\exp(b*x+a)*(\exp(2*a)+\exp(2*c))/(-\exp(2*b*x+2*a+2*c)+\exp(2*a))+1/2*\ln(\exp(b*x+a)-\exp(a-c))/b*\exp(-a-c)*\exp(2*a)-1/2*\ln(\exp(b*x+a)-\exp(a-c))/b*\exp(-a-c)*\exp(2*c)-1/2*\ln(\exp(b*x+a)+\exp(a-c))/b*\exp(-a-c)*\exp(2*a)+1/2*\ln(\exp(b*x+a)+\exp(a-c))/b*\exp(-a-c)*\exp(2*c)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1237 vs. $2(46) = 92$.

Time = 0.11 (sec) , antiderivative size = 1237, normalized size of antiderivative = 26.89

$$\int \cosh(a + bx) \coth^2(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)*coth(b*x+c)^2,x, algorithm="fricas")`

output

```

1/2*(cosh(b*x + c)^4*cosh(-a + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^4 + 4*(cosh(b*x + c)*cosh(-a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^3 - 3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + 3*(2*cosh(b*x + c)^2*cosh(-a + c)^2 + (2*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(2*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) - 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^4 - 3*cosh(b*x + c)^2)*sinh(-a + c)^2 - ((cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 - cosh(b*x + c))*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c) + (3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (3*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(3*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c) - 2*(cosh(b*x + c)^3*cosh(-a + c) - cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) + ((cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^3 - 3*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^2 + (cosh(b*x + c)^3 - cosh(b*x + c))*sinh(-a + c)^2...
```

Sympy [F]

$$\int \cosh(a + bx) \coth^2(c + bx) dx = \int \cosh(a + bx) \coth^2(bx + c) dx$$

input `integrate(cosh(b*x+a)*coth(b*x+c)**2,x)`

output `Integral(cosh(a + b*x)*coth(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. $2(46) = 92$.

Time = 0.04 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.13

$$\int \cosh(a + bx) \coth^2(c + bx) dx = -\frac{(e^{2a} - e^{2c})e^{(-a-c)} \log(e^{-bx} + e^c)}{2b} + \frac{(e^{2a} - e^{2c})e^{(-a-c)} \log(e^{-bx} - e^c)}{2b} - \frac{e^{(-bx-a)}}{2b} - \frac{(3e^{2a} + 2e^{2c})e^{(-2bx-2a)} - e^{2c}}{2b(e^{(-bx-a+2c)} - e^{(-3bx-a)})}$$

input `integrate(cosh(b*x+a)*coth(b*x+c)^2,x, algorithm="maxima")`

output `-1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b - 1/2*e^(-b*x - a)/b - 1/2*((3*e^(2*a) + 2*e^(2*c))*e^(-2*b*x - 2*a) - e^(2*c))/(b*(e^(-b*x - a + 2*c) - e^(-3*b*x - a)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. $2(46) = 92$.

Time = 0.18 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.09

$$\int \cosh(a + bx) \coth^2(c + bx) dx = \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - (e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) + \frac{2e^{(2bx+a+c)}}{e^{(3c)}}}{2b}$$

input `integrate(cosh(b*x+a)*coth(b*x+c)^2,x, algorithm="giac")`

output
$$-1/2*((e^{(2*a + c)} - e^{(3*c)})*e^{(-a - 2*c)}*\log(e^{(b*x + a + c)} + e^a) - (e^{(2*a + c)} - e^{(3*c)})*e^{(-a - 2*c)}*\log(\text{abs}(e^{(b*x + a + c)} - e^a)) + (2*e^{(2*b*x + 4*a)} + 3*e^{(2*b*x + 2*a + 2*c)} - e^{(2*a)})/(e^{(3*b*x + 3*a + 2*c)} - e^{(b*x + 3*a)} - e^{(b*x + a)})/b$$

Mupad [B] (verification not implemented)

Time = 1.07 (sec) , antiderivative size = 183, normalized size of antiderivative = 3.98

$$\int \cosh(a + bx) \coth^2(c + bx) dx = \frac{e^{a+bx}}{2b} - \frac{e^{-a-bx}}{2b} - \frac{\text{atan}\left(-\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2} - e^{2a} e^{-2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{-b^2}} + \frac{e^{a+bx} (e^{2a-2c} + 1)}{b (e^{2a-2c} - e^{2a+2bx})}$$

input `int(cosh(a + b*x)*coth(c + b*x)^2,x)`

output

```
exp(a + b*x)/(2*b) - exp(- a - b*x)/(2*b) - (atan(-(exp(-a)*exp(2*c)*exp(b
*x)*((-b^2)^(1/2) - exp(2*a)*exp(-2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*
c)*(exp(4*a)*exp(-4*c) - 2*exp(2*a)*exp(-2*c) + 1))^(1/2)))*(exp(2*c - 2*a
)*(exp(4*a - 4*c) - 2*exp(2*a - 2*c) + 1))^(1/2))/(-b^2)^(1/2) + (exp(a +
b*x)*(exp(2*a - 2*c) + 1))/(b*(exp(2*a - 2*c) - exp(2*a + 2*b*x)))
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 245, normalized size of antiderivative = 5.33

$$\int \cosh(a + bx) \coth^2(c + bx) dx$$

$$= \frac{e^{4bx+2a+3c} + e^{3bx+2a+2c} \log(e^{bx+c} - 1) - e^{3bx+2a+2c} \log(e^{bx+c} + 1) - e^{3bx+4c} \log(e^{bx+c} - 1) + e^{3bx+4c} \log(e^{bx+c} + 1)}{b}$$

input

```
int(cosh(b*x+a)*coth(b*x+c)^2,x)
```

output

```
(e**(2*a + 4*b*x + 3*c) + e**(2*a + 3*b*x + 2*c)*log(e**(b*x + c) - 1) - e
**(2*a + 3*b*x + 2*c)*log(e**(b*x + c) + 1) - e**(3*b*x + 4*c)*log(e**(b*x
+ c) - 1) + e**(3*b*x + 4*c)*log(e**(b*x + c) + 1) - 3*e**(2*a + 2*b*x +
c) - 3*e**(2*b*x + 3*c) - e**(2*a + b*x)*log(e**(b*x + c) - 1) + e**(2*a +
b*x)*log(e**(b*x + c) + 1) + e**(b*x + 2*c)*log(e**(b*x + c) - 1) - e**(b
*x + 2*c)*log(e**(b*x + c) + 1) + e**c)/(2*e**(a + b*x + c)*b*(e**(2*b*x +
2*c) - 1))
```


3.120 $\int \cosh(a + bx) \coth^3(c + bx) dx$

Optimal result	848
Mathematica [A] (verified)	848
Rubi [C] (verified)	849
Maple [B] (verified)	853
Fricas [B] (verification not implemented)	853
Sympy [F]	854
Maxima [B] (verification not implemented)	855
Giac [B] (verification not implemented)	855
Mupad [F(-1)]	856
Reduce [B] (verification not implemented)	856

Optimal result

Integrand size = 15, antiderivative size = 73

$$\int \cosh(a + bx) \coth^3(c + bx) dx = -\frac{3\operatorname{arctanh}(\cosh(c + bx)) \cosh(a - c)}{2b} + \frac{\cosh(a + bx)}{b} - \frac{\cosh(a - c) \coth(c + bx) \operatorname{csch}(c + bx)}{2b} - \frac{\operatorname{csch}(c + bx) \sinh(a - c)}{b}$$

output

```
-3/2*arctanh(cosh(b*x+c))*cosh(a-c)/b+cosh(b*x+a)/b-1/2*cosh(a-c)*coth(b*x+c)*csch(b*x+c)/b-csch(b*x+c)*sinh(a-c)/b
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.96

$$\int \cosh(a + bx) \coth^3(c + bx) dx = \frac{-12\operatorname{arctanh}\left(\cosh(c) + \sinh(c) \tanh\left(\frac{bx}{2}\right)\right) \cosh(a - c) + (2 \cosh(a - 2c - bx) - 5 \cosh(a + bx) + \cosh(a + 2c + bx))}{4b}$$

input

```
Integrate[Cosh[a + b*x]*Coth[c + b*x]^3,x]
```

output

$$(-12*\text{ArcTanh}[\text{Cosh}[c] + \text{Sinh}[c]*\text{Tanh}[(b*x)/2]]*\text{Cosh}[a - c] + (2*\text{Cosh}[a - 2*c - b*x] - 5*\text{Cosh}[a + b*x] + \text{Cosh}[a + 2*c + 3*b*x])*\text{Csch}[c + b*x]^2)/(4*b)$$
Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.29, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 1.133$, Rules used = {6155, 3042, 26, 3091, 26, 3042, 26, 4257, 6156, 3042, 3086, 24, 6155, 3042, 26, 3118, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \coth^3(bx + c) dx$$

$$\downarrow 6155$$

$$\int \coth^2(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int \coth^2(c + bx) \text{csch}(c + bx) dx$$

$$\downarrow 3042$$

$$\int \coth^2(c + bx) \sinh(a + bx) dx + \cosh(a - c) \int -i \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right)^2 dx$$

$$\downarrow 26$$

$$\int \coth^2(c + bx) \sinh(a + bx) dx - i \cosh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right) \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right)^2 dx$$

$$\downarrow 3091$$

$$\int \coth^2(c + bx) \sinh(a + bx) dx - i \cosh(a - c) \left(-\frac{1}{2} \int -i \text{csch}(c + bx) dx - \frac{i \coth(bx + c) \text{csch}(bx + c)}{2b}\right)$$

$$\downarrow 26$$

$$c) \left(\int \coth^2(c + bx) \sinh(a + bx) dx - i \cosh(a - \right. \\ \left. \frac{1}{2} \int \operatorname{csch}(c + bx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3042

$$c) \left(\int \coth^2(c + bx) \sinh(a + bx) dx - i \cosh(a - \right. \\ \left. \frac{1}{2} \int i \operatorname{csc}(ic + ibx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 26

$$c) \left(-\frac{1}{2} \int \operatorname{csc}(ic + ibx) dx - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 4257

$$c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 6156

$$\int \cosh(a + bx) \coth(c + bx) dx + \sinh(a - c) \int \coth(c + bx) \operatorname{csch}(c + bx) dx - i \cosh(a - \\ c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3042

$$\int \cosh(a + bx) \coth(c + bx) dx + \sinh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx - \\ i \cosh(a - c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 3086

$$\int \cosh(a + bx) \coth(c + bx) dx - \frac{i \sinh(a - c) \int 1d(-i \operatorname{csch}(c + bx))}{b} - i \cosh(a - \\ c) \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right)$$

↓ 24

$$\begin{aligned}
& \int \cosh(a + bx) \coth(c + bx) dx - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow 6155 \\
& \cosh(a - c) \int \operatorname{csch}(c + bx) dx + \int \sinh(a + bx) dx - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow 3042 \\
& \cosh(a - c) \int i \csc(ic + ibx) dx + \int -i \sin(ia + ibx) dx - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow 26 \\
& i \cosh(a - c) \int \csc(ic + ibx) dx - i \int \sin(ia + ibx) dx - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} \\
& \downarrow 3118 \\
& i \cosh(a - c) \int \csc(ic + ibx) dx - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} + \\
& \frac{\cosh(a + bx)}{b} \\
& \downarrow 4257 \\
& -\frac{\cosh(a - c) \operatorname{arctanh}(\cosh(bx + c))}{b} - i \cosh(a - \\
c) & \left(-\frac{i \operatorname{arctanh}(\cosh(bx + c))}{2b} - \frac{i \coth(bx + c) \operatorname{csch}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{csch}(bx + c)}{b} + \\
& \frac{\cosh(a + bx)}{b}
\end{aligned}$$

input

```
Int[Cosh[a + b*x]*Coth[c + b*x]^3,x]
```

output
$$-\left(\frac{\text{ArcTanh}[\text{Cosh}[c + b*x]]*\text{Cosh}[a - c]}{b}\right) + \frac{\text{Cosh}[a + b*x]}{b} - I*\text{Cosh}[a - c] * \left(\frac{(-1/2*I)*\text{ArcTanh}[\text{Cosh}[c + b*x]]}{b} - \left(\frac{I}{2}\right)*\text{Coth}[c + b*x]*\text{Csch}[c + b*x]\right)/b - \left(\frac{\text{Csch}[c + b*x]*\text{Sinh}[a - c]}{b}\right)$$

Defintions of rubi rules used

rule 24
$$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$$

rule 26
$$\text{Int}[(\text{Complex}[0, a_])*(F_x), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$$

rule 3042
$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3086
$$\text{Int}[(a_)*\text{sec}[(e_)] + (f_)*(x_)]^{(m_)}*((b_)*\text{tan}[(e_)] + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[a/f \text{ Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{(n-1)/2}], x], x, \text{Sec}[e+f*x], x] \text{ ; FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$$

rule 3091
$$\text{Int}[(a_)*\text{sec}[(e_)] + (f_)*(x_)]^{(m_)}*((b_)*\text{tan}[(e_)] + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Sec}[e+f*x])^m*((b*\text{Tan}[e+f*x])^{(n-1)/(f*(m+n-1))}), x] - \text{Simp}[b^2*((n-1)/(m+n-1)) \text{ Int}[(a*\text{Sec}[e+f*x])^m*(b*\text{Tan}[e+f*x])^{(n-2)}, x], x] \text{ ; FreeQ}\{a, b, e, f, m\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{NeQ}[m+n-1, 0] \ \&\& \ \text{IntegersQ}[2*m, 2*n]$$

rule 3118
$$\text{Int}[\sin[(c_)] + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c+d*x]/d, x] \text{ ; FreeQ}\{c, d\}, x]$$

rule 4257
$$\text{Int}[\text{csc}[(c_)] + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c+d*x]]/d, x] \text{ ; FreeQ}\{c, d\}, x]$$

rule 6155

```
Int[Cosh[v_]*Coth[w_]^(n_), x_Symbol] := Int[Sinh[v]*Coth[w]^(n - 1), x] +
Simp[Cosh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ
[w, v] && FreeQ[v - w, x]
```

rule 6156

```
Int[Coth[w_]^(n_)*Sinh[v_], x_Symbol] := Int[Cosh[v]*Coth[w]^(n - 1), x] +
Simp[Sinh[v - w] Int[Csch[w]*Coth[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ
[w, v] && FreeQ[v - w, x]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(69) = 138.

Time = 0.20 (sec) , antiderivative size = 228, normalized size of antiderivative = 3.12

method	result
risch	$\frac{e^{bx+a}}{2b} + \frac{e^{-bx-a}}{2b} + \frac{e^{bx+a}(-3e^{2bx+4a+2c} + e^{2bx+2a+4c} + e^{4a} - 3e^{2a+2c})}{2b(-e^{2bx+2a+2c} + e^{2a})^2} - \frac{3 \ln(e^{bx+a} + e^{a-c})e^{-a-c}e^{2a}}{4b} - \frac{3 \ln(e^{bx+a} + e^{a-c})}{4b}$

input

```
int(cosh(b*x+a)*coth(b*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
1/2/b*exp(b*x+a)+1/2/b*exp(-b*x-a)+1/2*exp(b*x+a)*(-3*exp(2*b*x+4*a+2*c)+
xp(2*b*x+2*a+4*c)+exp(4*a)-3*exp(2*a+2*c))/b/(-exp(2*b*x+2*a+2*c)+exp(2*a)
)^2-3/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*a)-3/4*ln(exp(b*x+a)+exp
(a-c))/b*exp(-a-c)*exp(2*c)+3/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*
a)+3/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2372 vs. 2(69) = 138.

Time = 0.11 (sec) , antiderivative size = 2372, normalized size of antiderivative = 32.49

$$\int \cosh(a + bx) \coth^3(c + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*coth(b*x+c)^3,x, algorithm="fricas")
```

output

```

1/4*(2*cosh(b*x + c)^6*cosh(-a + c)^2 + 2*(cosh(-a + c)^2 - 2*cosh(-a + c)
*sinh(-a + c) + sinh(-a + c)^2)*sinh(b*x + c)^6 + 12*(cosh(b*x + c)*cosh(-
a + c)^2 - 2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + cosh(b*x + c)*sinh(
-a + c)^2)*sinh(b*x + c)^5 - 2*(5*cosh(-a + c)^2 - 2)*cosh(b*x + c)^4 + 2*
(15*cosh(b*x + c)^2*cosh(-a + c)^2 + 5*(3*cosh(b*x + c)^2 - 1)*sinh(-a + c
)^2 - 5*cosh(-a + c)^2 - 10*(3*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c)
)*sinh(-a + c) + 2)*sinh(b*x + c)^4 + 8*(5*cosh(b*x + c)^3*cosh(-a + c)^2
+ 5*(cosh(b*x + c)^3 - cosh(b*x + c))*sinh(-a + c)^2 - (5*cosh(-a + c)^2 -
2)*cosh(b*x + c) - 10*(cosh(b*x + c)^3*cosh(-a + c) - cosh(b*x + c)*cosh(
-a + c))*sinh(-a + c))*sinh(b*x + c)^3 + 2*(2*cosh(-a + c)^2 - 5)*cosh(b*x
+ c)^2 + 2*(15*cosh(b*x + c)^4*cosh(-a + c)^2 - 6*(5*cosh(-a + c)^2 - 2)*
cosh(b*x + c)^2 + (15*cosh(b*x + c)^4 - 30*cosh(b*x + c)^2 + 2)*sinh(-a +
c)^2 + 2*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^4*cosh(-a + c) - 30*cosh(b*x
+ c)^2*cosh(-a + c) + 2*cosh(-a + c))*sinh(-a + c) - 5)*sinh(b*x + c)^2 +
2*(cosh(b*x + c)^6 - 5*cosh(b*x + c)^4 + 2*cosh(b*x + c)^2)*sinh(-a + c)^
2 - 3*((cosh(-a + c)^2 + 1)*cosh(b*x + c)^5 + (cosh(-a + c)^2 - 2*cosh(-a
+ c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^5 - 5*(2*cosh(b*x +
c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c
)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^4 - 2*(cosh(-a + c)^2 + 1)*cosh(b*x
+ c)^3 + 2*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2...

```

Sympy [F]

$$\int \cosh(a + bx) \coth^3(c + bx) dx = \int \cosh(a + bx) \coth^3(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*coth(b*x+c)**3,x)
```

output

```
Integral(cosh(a + b*x)*coth(b*x + c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(69) = 138$.

Time = 0.05 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.52

$$\int \cosh(a + bx) \coth^3(c + bx) dx$$

$$= -\frac{3(e^{2a} + e^{2c})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{4b} + \frac{3(e^{2a} + e^{2c})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{4b}$$

$$+ \frac{e^{(-bx-a)}}{2b} - \frac{(5e^{(2a+2c)} - e^{(4c)})e^{(-2bx-2a)} - (2e^{(4a)} - 3e^{(2a+2c)})e^{(-4bx-4a)} - e^{(4c)}}{2b(e^{(-bx-a+4c)} - 2e^{(-3bx-a+2c)} + e^{(-5bx-a)})}$$

input `integrate(cosh(b*x+a)*coth(b*x+c)^3,x, algorithm="maxima")`

output `-3/4*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 3/4*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/2*e^(-b*x - a)/b - 1/2*((5*e^(2*a + 2*c) - e^(4*c))*e^(-2*b*x - 2*a) - (2*e^(4*a) - 3*e^(2*a + 2*c))*e^(-4*b*x - 4*a) - e^(4*c))/(b*(e^(-b*x - a + 4*c) - 2*e^(-3*b*x - a + 2*c) + e^(-5*b*x - a)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 167, normalized size of antiderivative = 2.29

$$\int \cosh(a + bx) \coth^3(c + bx) dx =$$

$$\frac{3(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(e^{(bx+a+c)} + e^a) - 3(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+a+c)} - e^a|) + \frac{2}{3}}{4b}$$

input `integrate(cosh(b*x+a)*coth(b*x+c)^3,x, algorithm="giac")`

output

```
-1/4*(3*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(e^(b*x + a + c) + e^a) -
3*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + a + c) - e^a)) + 2
*(3*e^(3*b*x + 5*a + 2*c) - e^(3*b*x + 3*a + 4*c) - e^(b*x + 5*a) + 3*e^(b
*x + 3*a + 2*c))/(e^(2*b*x + 2*a + 2*c) - e^(2*a))^2 - 2*e^(b*x + a) - 2*e
^(-b*x - a))/b
```

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \coth^3(c + bx) dx = \int \cosh(a + bx) \coth(c + bx)^3 dx$$

input

```
int(cosh(a + b*x)*coth(c + b*x)^3,x)
```

output

```
int(cosh(a + b*x)*coth(c + b*x)^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 386, normalized size of antiderivative = 5.29

$$\int \cosh(a + bx) \coth^3(c + bx) dx$$

$$= \frac{2e^{6bx+2a+5c} + 3e^{5bx+2a+4c}\log(e^{bx+c} - 1) - 3e^{5bx+2a+4c}\log(e^{bx+c} + 1) + 3e^{5bx+6c}\log(e^{bx+c} - 1) - 3e^{5bx+6c}\log(e^{bx+c} + 1)}{b}$$

input

```
int(cosh(b*x+a)*coth(b*x+c)^3,x)
```

output

```
(2*e**(2*a + 6*b*x + 5*c) + 3*e**(2*a + 5*b*x + 4*c)*log(e**(b*x + c) - 1)
- 3*e**(2*a + 5*b*x + 4*c)*log(e**(b*x + c) + 1) + 3*e**(5*b*x + 6*c)*log
(e**(b*x + c) - 1) - 3*e**(5*b*x + 6*c)*log(e**(b*x + c) + 1) - 10*e**(2*a
+ 4*b*x + 3*c) + 4*e**(4*b*x + 5*c) - 6*e**(2*a + 3*b*x + 2*c)*log(e**(b*
x + c) - 1) + 6*e**(2*a + 3*b*x + 2*c)*log(e**(b*x + c) + 1) - 6*e**(3*b*x
+ 4*c)*log(e**(b*x + c) - 1) + 6*e**(3*b*x + 4*c)*log(e**(b*x + c) + 1) +
4*e**(2*a + 2*b*x + c) - 10*e**(2*b*x + 3*c) + 3*e**(2*a + b*x)*log(e**(b
*x + c) - 1) - 3*e**(2*a + b*x)*log(e**(b*x + c) + 1) + 3*e**(b*x + 2*c)*l
og(e**(b*x + c) - 1) - 3*e**(b*x + 2*c)*log(e**(b*x + c) + 1) + 2*e**c)/(4
*e**(a + b*x + c)*b*(e**(4*b*x + 4*c) - 2*e**(2*b*x + 2*c) + 1))
```

3.121 $\int \cosh(a + bx) \coth(c + dx) dx$

Optimal result	858
Mathematica [A] (verified)	858
Rubi [A] (verified)	859
Maple [F]	860
Fricas [F]	860
Sympy [F]	861
Maxima [F]	861
Giac [F]	861
Mupad [F(-1)]	862
Reduce [F]	862

Optimal result

Integrand size = 13, antiderivative size = 95

$$\int \cosh(a + bx) \coth(c + dx) dx = \frac{e^{-a-bx} \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, e^{2(c+dx)}\right)}{b} + \frac{\sinh(a + bx)}{b}$$

output

$\exp(-b*x-a)*\operatorname{hypergeom}([1, -1/2*b/d], [1-1/2*b/d], \exp(2*d*x+2*c))/b - \exp(b*x+a)*\operatorname{hypergeom}([1, 1/2*b/d], [1+1/2*b/d], \exp(2*d*x+2*c))/b + \sinh(b*x+a)/b$

Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.04

$$\int \cosh(a + bx) \coth(c + dx) dx = \frac{e^{-a-bx}(-1 + e^{2(a+bx)}) + 2 \operatorname{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, e^{2(c+dx)}\right) - 2e^{2(a+bx)} \operatorname{Hypergeometric2F1}\left(1, \frac{b}{2d}, 1 + \frac{b}{2d}, e^{2(c+dx)}\right)}{2b}$$

input

`Integrate[Cosh[a + b*x]*Coth[c + d*x], x]`

output

$$\frac{(E^{-a - bx}) * (-1 + E^{2(a + bx)}) + 2 * \text{Hypergeometric2F1}[1, -1/2 * b/d, 1 - b/(2 * d), E^{2(c + dx)}] - 2 * E^{2(a + bx)} * \text{Hypergeometric2F1}[1, b/(2 * d), 1 + b/(2 * d), E^{2(c + dx)}])}{(2 * b)}$$
Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.22, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6136, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \coth(c + dx) dx$$

$$\downarrow \text{6136}$$

$$\int \left(-\frac{e^{-a-bx}}{1 - e^{2(c+dx)}} - \frac{e^{a+bx}}{1 - e^{2(c+dx)}} + \frac{1}{2}e^{-a-bx} + \frac{1}{2}e^{a+bx} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{e^{-a-bx} \text{Hypergeometric2F1}\left(1, -\frac{b}{2d}, 1 - \frac{b}{2d}, e^{2(c+dx)}\right)}{b} - \frac{e^{a+bx} \text{Hypergeometric2F1}\left(1, \frac{b}{2d}, \frac{b}{2d} + 1, e^{2(c+dx)}\right)}{b} - \frac{e^{-a-bx}}{2b} + \frac{e^{a+bx}}{2b}$$

input

$$\text{Int}[\text{Cosh}[a + b * x] * \text{Coth}[c + d * x], x]$$

output

$$\frac{-1/2 * E^{-a - bx}/b + E^{a + bx}/(2 * b) + (E^{-a - bx} * \text{Hypergeometric2F1}[1, -1/2 * b/d, 1 - b/(2 * d), E^{2(c + dx)}]) / b - (E^{a + bx} * \text{Hypergeometric2F1}[1, b/(2 * d), 1 + b/(2 * d), E^{2(c + dx)}]) / b}{b}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6136 `Int[Cosh[(a_.) + (b_.)*(x_)]*Coth[(c_.) + (d_.)*(x_)], x_Symbol] := Int[1/(E^(a + b*x)*2) + E^(a + b*x)/2 - 1/(E^(a + b*x)*(1 - E^(2*(c + d*x)))) - E^(a + b*x)/(1 - E^(2*(c + d*x))), x] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - d^2, 0]`

Maple [F]

$$\int \cosh (bx + a) \coth (dx + c) dx$$

input `int(cosh(b*x+a)*coth(d*x+c),x)`

output `int(cosh(b*x+a)*coth(d*x+c),x)`

Fricas [F]

$$\int \cosh (a + bx) \coth (c + dx) dx = \int \cosh (bx + a) \coth (dx + c) dx$$

input `integrate(cosh(b*x+a)*coth(d*x+c),x, algorithm="fricas")`

output `integral(cosh(b*x + a)*coth(d*x + c), x)`

Sympy [F]

$$\int \cosh(a + bx) \coth(c + dx) dx = \int \cosh(a + bx) \coth(c + dx) dx$$

input `integrate(cosh(b*x+a)*coth(d*x+c),x)`

output `Integral(cosh(a + b*x)*coth(c + d*x), x)`

Maxima [F]

$$\int \cosh(a + bx) \coth(c + dx) dx = \int \cosh(bx + a) \coth(dx + c) dx$$

input `integrate(cosh(b*x+a)*coth(d*x+c),x, algorithm="maxima")`

output `1/2*(e^(2*b*x + 2*a) - 1)*e^(-b*x - a)/b - 1/2*integrate((e^(2*b*x + 2*a) + 1)/(e^(b*x + d*x + a + c) + e^(b*x + a)), x) + 1/2*integrate((e^(2*b*x + 2*a) + 1)/(e^(b*x + d*x + a + c) - e^(b*x + a)), x)`

Giac [F]

$$\int \cosh(a + bx) \coth(c + dx) dx = \int \cosh(bx + a) \coth(dx + c) dx$$

input `integrate(cosh(b*x+a)*coth(d*x+c),x, algorithm="giac")`

output `integrate(cosh(b*x + a)*coth(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \coth(c + dx) dx = \int \cosh(a + bx) \coth(c + dx) dx$$

input `int(cosh(a + b*x)*coth(c + d*x),x)`output `int(cosh(a + b*x)*coth(c + d*x), x)`**Reduce [F]**

$$\int \cosh(a + bx) \coth(c + dx) dx$$

$$= \frac{e^{2bx+2a} + 2e^{bx+2a} \left(\int \frac{e^{bx}}{e^{2dx+2c}-1} dx \right) b + 2e^{bx} \left(\int \frac{1}{e^{bx+2dx+2c}-e^{bx}} dx \right) b - 1}{2e^{bx+ab}}$$

input `int(cosh(b*x+a)*coth(d*x+c),x)`output `(e**(2*a + 2*b*x) + 2*e**(2*a + b*x)*int(e**(b*x)/(e**(2*c + 2*d*x) - 1),x)*b + 2*e**(b*x)*int(1/(e**(b*x + 2*c + 2*d*x) - e**(b*x)),x)*b - 1)/(2*e*(a + b*x)*b)`

3.122 $\int \cosh(x)\operatorname{sech}(2x) dx$

Optimal result	863
Mathematica [A] (verified)	863
Rubi [A] (verified)	864
Maple [C] (verified)	865
Fricas [B] (verification not implemented)	865
Sympy [F]	866
Maxima [B] (verification not implemented)	866
Giac [B] (verification not implemented)	867
Mupad [B] (verification not implemented)	867
Reduce [B] (verification not implemented)	868

Optimal result

Integrand size = 7, antiderivative size = 15

$$\int \cosh(x)\operatorname{sech}(2x) dx = \frac{\arctan(\sqrt{2}\sinh(x))}{\sqrt{2}}$$

output

```
1/2*arctan(sinh(x)*2^(1/2))*2^(1/2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cosh(x)\operatorname{sech}(2x) dx = \frac{\arctan(\sqrt{2}\sinh(x))}{\sqrt{2}}$$

input

```
Integrate[Cosh[x]*Sech[2*x],x]
```

output

```
ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2]
```


Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4856, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x)\operatorname{sech}(2x) dx$$

↓ 3042

$$\int \frac{\cos(ix)}{\cos(2ix)} dx$$

↓ 4856

$$\int \frac{1}{2 \sinh^2(x) + 1} d \sinh(x)$$

↓ 216

$$\frac{\arctan(\sqrt{2} \sinh(x))}{\sqrt{2}}$$

input `Int[Cosh[x]*Sech[2*x],x]`

output `ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2]`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4856

```
Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.93

method	result	size
risch	$\frac{i\sqrt{2} \ln(e^{2x} + i\sqrt{2}e^x - 1)}{4} - \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x - 1)}{4}$	44

input

```
int(cosh(x)*sech(2*x), x, method=_RETURNVERBOSE)
```

output

```
1/4*I*2^(1/2)*ln(exp(2*x)+I*2^(1/2)*exp(x)-1)-1/4*I*2^(1/2)*ln(exp(2*x)-I*2^(1/2)*exp(x)-1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(12) = 24$.

Time = 0.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 4.53

$$\begin{aligned} & \int \cosh(x) \operatorname{sech}(2x) dx \\ &= \frac{1}{2} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \cosh(x) + \frac{1}{2} \sqrt{2} \sinh(x) \right) \\ & \quad - \frac{1}{2} \sqrt{2} \arctan \left(-\frac{\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}}{2(\cosh(x) - \sinh(x))} \right) \end{aligned}$$

input

```
integrate(cosh(x)*sech(2*x), x, algorithm="fricas")
```

output

```
1/2*sqrt(2)*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)) - 1/2*sqrt(2)
)*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sin
h(x)^2 + sqrt(2))/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \cosh(x)\operatorname{sech}(2x) dx = \int \cosh(x) \operatorname{sech}(2x) dx$$

input

```
integrate(cosh(x)*sech(2*x),x)
```

output

```
Integral(cosh(x)*sech(2*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(12) = 24$.

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.87

$$\int \cosh(x)\operatorname{sech}(2x) dx = -\frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2e^{-x}\right)\right) - \frac{1}{2}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2e^{-x}\right)\right)$$

input

```
integrate(cosh(x)*sech(2*x),x, algorithm="maxima")
```

output

```
-1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(-x))) - 1/2*sqrt(2)*arctan
(-1/2*sqrt(2)*(sqrt(2) - 2*e^(-x)))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(12) = 24$.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.60

$$\int \cosh(x)\operatorname{sech}(2x) dx = \frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2e^x)\right) + \frac{1}{2}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2e^x)\right)$$

input `integrate(cosh(x)*sech(2*x),x, algorithm="giac")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x))`

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 2.13

$$\int \cosh(x)\operatorname{sech}(2x) dx = \frac{\sqrt{2}\left(\operatorname{atan}\left(\frac{\sqrt{2}e^x}{2} + \frac{\sqrt{2}e^{3x}}{2}\right) + \operatorname{atan}\left(\frac{\sqrt{2}e^x}{2}\right)\right)}{2}$$

input `int(cosh(x)/cosh(2*x),x)`

output `(2^(1/2)*(atan((2^(1/2)*exp(x))/2 + (2^(1/2)*exp(3*x))/2) + atan((2^(1/2)*exp(x))/2)))/2`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.33

$$\int \cosh(x)\operatorname{sech}(2x) dx = \frac{\sqrt{2} \left(\operatorname{atan}\left(\frac{2e^x - \sqrt{2}}{\sqrt{2}}\right) + \operatorname{atan}\left(\frac{2e^x + \sqrt{2}}{\sqrt{2}}\right) \right)}{2}$$

input `int(cosh(x)*sech(2*x),x)`output `(sqrt(2)*(atan((2*e**x - sqrt(2))/sqrt(2)) + atan((2*e**x + sqrt(2))/sqrt(2))))/2`

3.123 $\int \cosh(x)\operatorname{sech}(3x) dx$

Optimal result	869
Mathematica [A] (verified)	869
Rubi [A] (verified)	870
Maple [C] (verified)	871
Fricas [B] (verification not implemented)	871
Sympy [F]	872
Maxima [B] (verification not implemented)	872
Giac [A] (verification not implemented)	873
Mupad [B] (verification not implemented)	873
Reduce [B] (verification not implemented)	873

Optimal result

Integrand size = 7, antiderivative size = 15

$$\int \cosh(x)\operatorname{sech}(3x) dx = \frac{\arctan(\sqrt{3}\tanh(x))}{\sqrt{3}}$$

output `1/3*arctan(tanh(x)*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cosh(x)\operatorname{sech}(3x) dx = \frac{\arctan(\sqrt{3}\tanh(x))}{\sqrt{3}}$$

input `Integrate[Cosh[x]*Sech[3*x],x]`

output `ArcTan[Sqrt[3]*Tanh[x]]/Sqrt[3]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4889, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(x) \operatorname{sech}(3x) dx$$

$$\downarrow 3042$$

$$\int \frac{\cos(ix)}{\cos(3ix)} dx$$

$$\downarrow 4889$$

$$\int \frac{1}{3 \tanh^2(x) + 1} d \tanh(x)$$

$$\downarrow 216$$

$$\frac{\arctan(\sqrt{3} \tanh(x))}{\sqrt{3}}$$

input `Int[Cosh[x]*Sech[3*x],x]`

output `ArcTan[Sqrt[3]*Tanh[x]]/Sqrt[3]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4889

```
Int[u_, x_Symbol] := With[{v = FunctionOfTrig[u, x]}, With[{d = FreeFactors
[Tan[v], x]}, Simp[d/Coefficient[v, x, 1] Subst[Int[SubstFor[1/(1 + d^2*x
^2), Tan[v]/d, u, x], x], x, Tan[v]/d], x]] /; !FalseQ[v] && FunctionOfQ[N
onfreeFactors[Tan[v], x], u, x]] /; InverseFunctionFreeQ[u, x] && !MatchQ[
u, (v_.)*((c_.)*tan[w_]^(n_.)*tan[z_]^(n_.))^ (p_.) /; FreeQ[{c, p}, x] && I
ntegerQ[n] && LinearQ[w, x] && EqQ[z, 2*w]]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.67

method	result	size
risch	$\frac{i\sqrt{3} \ln\left(e^{2x} - \frac{1}{2} + \frac{i\sqrt{3}}{2}\right)}{6} - \frac{i\sqrt{3} \ln\left(e^{2x} - \frac{1}{2} - \frac{i\sqrt{3}}{2}\right)}{6}$	40

input

```
int(cosh(x)*sech(3*x), x, method=_RETURNVERBOSE)
```

output

```
1/6*I*3^(1/2)*ln(exp(2*x)-1/2+1/2*I*3^(1/2))-1/6*I*3^(1/2)*ln(exp(2*x)-1/2
-1/2*I*3^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(12) = 24.

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.07

$$\int \cosh(x) \operatorname{sech}(3x) dx = -\frac{1}{3} \sqrt{3} \arctan \left(-\frac{\sqrt{3} \cosh(x) + 3\sqrt{3} \sinh(x)}{3(\cosh(x) - \sinh(x))} \right)$$

input

```
integrate(cosh(x)*sech(3*x), x, algorithm="fricas")
```

output

```
-1/3*sqrt(3)*arctan(-1/3*(sqrt(3)*cosh(x) + 3*sqrt(3)*sinh(x))/(cosh(x) -
sinh(x)))
```


Sympy [F]

$$\int \cosh(x)\operatorname{sech}(3x) dx = \int \cosh(x) \operatorname{sech}(3x) dx$$

input `integrate(cosh(x)*sech(3*x),x)`

output `Integral(cosh(x)*sech(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(12) = 24$.

Time = 0.13 (sec) , antiderivative size = 114, normalized size of antiderivative = 7.60

$$\begin{aligned} \int \cosh(x)\operatorname{sech}(3x) dx = & -\frac{1}{6}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^{-2x}-1)\right) \\ & -\frac{1}{6}\sqrt{3}\arctan(\sqrt{3}+2e^x) + \frac{1}{6}\sqrt{3}\arctan(-\sqrt{3}+2e^x) \\ & + \frac{1}{12}\log(\sqrt{3}e^x+e^{2x}+1) + \frac{1}{12}\log(-\sqrt{3}e^x+e^{2x}+1) \\ & - \frac{1}{6}\log(e^{2x}+1) + \frac{1}{6}\log(e^{-2x}+1) \\ & - \frac{1}{12}\log(-e^{-2x}+e^{-4x}+1) \end{aligned}$$

input `integrate(cosh(x)*sech(3*x),x, algorithm="maxima")`

output `-1/6*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(-2*x) - 1)) - 1/6*sqrt(3)*arctan(sqrt(3) + 2*e^x) + 1/6*sqrt(3)*arctan(-sqrt(3) + 2*e^x) + 1/12*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/12*log(-sqrt(3)*e^x + e^(2*x) + 1) - 1/6*log(e^(2*x) + 1) + 1/6*log(e^(-2*x) + 1) - 1/12*log(-e^(-2*x) + e^(-4*x) + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \cosh(x)\operatorname{sech}(3x) dx = \frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^{(2x)} - 1)\right)$$

input `integrate(cosh(x)*sech(3*x),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^(2*x) - 1))`**Mupad [B] (verification not implemented)**

Time = 0.95 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int \cosh(x)\operatorname{sech}(3x) dx = \frac{\sqrt{3}\operatorname{atan}\left(\frac{\sqrt{3}(2e^{2x}-1)}{3}\right)}{3}$$

input `int(cosh(x)/cosh(3*x),x)`output `(3^(1/2)*atan((3^(1/2)*(2*exp(2*x) - 1))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \cosh(x)\operatorname{sech}(3x) dx = \frac{\sqrt{3}(\operatorname{atan}(2e^x - \sqrt{3}) - \operatorname{atan}(2e^x + \sqrt{3}))}{3}$$

input `int(cosh(x)*sech(3*x),x)`output `(sqrt(3)*(atan(2*e**x - sqrt(3)) - atan(2*e**x + sqrt(3))))/3`

3.124 $\int \cosh(x)\operatorname{sech}(4x) dx$

Optimal result	874
Mathematica [A] (verified)	874
Rubi [A] (verified)	875
Maple [C] (verified)	876
Fricas [B] (verification not implemented)	877
Sympy [F]	877
Maxima [F]	878
Giac [B] (verification not implemented)	878
Mupad [B] (verification not implemented)	879
Reduce [B] (verification not implemented)	880

Optimal result

Integrand size = 7, antiderivative size = 71

$$\int \cosh(x)\operatorname{sech}(4x) dx = \frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{2}-\sqrt{2}}\right)}{2\sqrt{2}(2-\sqrt{2})} - \frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{2}+\sqrt{2}}\right)}{2\sqrt{2}(2+\sqrt{2})}$$

output

```
1/2*arctan(2*sinh(x)/(2-2^(1/2))^(1/2))/(4-2*2^(1/2))^(1/2)-1/2*arctan(2*si
inh(x)/(2+2^(1/2))^(1/2))/(4+2*2^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.94

$$\int \cosh(x)\operatorname{sech}(4x) dx = \frac{1}{4}\sqrt{2+\sqrt{2}}\arctan\left(\frac{2\sinh(x)}{\sqrt{2}-\sqrt{2}}\right) - \frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{2}+\sqrt{2}}\right)}{2\sqrt{2}(2+\sqrt{2})}$$

input

```
Integrate[Cosh[x]*Sech[4*x],x]
```

output

```
(Sqrt[2 + Sqrt[2]]*ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[2]]])/4 - ArcTan[(2*Si
nh[x])/Sqrt[2 + Sqrt[2]]]/(2*Sqrt[2*(2 + Sqrt[2])])
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4856, 1406, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \operatorname{sech}(4x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(ix)}{\cos(4ix)} dx \\
 & \quad \downarrow \text{4856} \\
 & \int \frac{1}{8 \sinh^4(x) + 8 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{1406} \\
 & \sqrt{2} \int \frac{1}{8 \sinh^2(x) + 2(2 - \sqrt{2})} d \sinh(x) - \sqrt{2} \int \frac{1}{8 \sinh^2(x) + 2(2 + \sqrt{2})} d \sinh(x) \\
 & \quad \downarrow \text{216} \\
 & \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{2} - \sqrt{2}}\right)}{2\sqrt{2}(2 - \sqrt{2})} - \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{2} + \sqrt{2}}\right)}{2\sqrt{2}(2 + \sqrt{2})}
 \end{aligned}$$

input `Int[Cosh[x]*Sech[4*x],x]`

output `ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[2]]]/(2*Sqrt[2*(2 - Sqrt[2])]) - ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[2]]]/(2*Sqrt[2*(2 + Sqrt[2])])`

Defintions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_ \cdot x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 1406 $\text{Int}[(a_ + (b_ \cdot x_)^2 + (c_ \cdot x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4 \cdot a \cdot c, 2]\}, \text{Simp}[c/q \ \text{Int}[1/(b/2 - q/2 + c \cdot x^2), x], x] - \text{Simp}[c/q \ \text{Int}[1/(b/2 + q/2 + c \cdot x^2), x], x] /;$ $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{PosQ}[b^2 - 4 \cdot a \cdot c]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4856 $\text{Int}[(u_ \cdot (F_))[(c_ \cdot (a_ + (b_ \cdot x_))], x_Symbol] \rightarrow \text{With}\{d = \text{FreeFactors}[\text{Sin}[c \cdot (a + b \cdot x)], x]\}, \text{Simp}[d/(b \cdot c) \ \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Sin}[c \cdot (a + b \cdot x)]]/d, u, x], x], x, \text{Sin}[c \cdot (a + b \cdot x)]/d, x] /;$ $\text{FunctionOfQ}[\text{Sin}[c \cdot (a + b \cdot x)]/d, u, x] /;$ $\text{FreeQ}\{a, b, c, x\} \ \&\& \ (\text{EqQ}[F, \text{Cos}] \ || \ \text{EqQ}[F, \text{cos}])$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.20 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.56

method	result	size
risch	$2 \left(\sum_{R=\text{RootOf}(32768_Z^4+512_Z^2+1)} _R \ln(e^{2x} + (-4096_R^3 - 48_R) e^x - 1) \right)$	40

input `int(cosh(x)*sech(4*x),x,method=_RETURNVERBOSE)`

output `2*sum(_R*ln(exp(2*x)+(-4096*_R^3-48*_R)*exp(x)-1),_R=RootOf(32768*_Z^4+512*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 217 vs. $2(49) = 98$.

Time = 0.10 (sec) , antiderivative size = 217, normalized size of antiderivative = 3.06

$$\int \cosh(x) \operatorname{sech}(4x) dx$$

$$= \frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan \left(\frac{1}{2} \left(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (\sqrt{2} - 2) \cosh(x) + (\sqrt{2} + 2) \sinh(x) \right) \sqrt{\sqrt{2} + 2} \right)$$

$$+ \frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan \left(\frac{1}{2} \left(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) \right) \sqrt{\sqrt{2} + 2} \right)$$

$$- \frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan \left(\frac{1}{2} \left(\sqrt{2} \cosh(x)^3 + 3\sqrt{2} \cosh(x) \sinh(x)^2 + \sqrt{2} \sinh(x)^3 + (\sqrt{2} + 2) \cosh(x) + (\sqrt{2} - 2) \sinh(x) \right) \sqrt{-\sqrt{2} + 2} \right)$$

$$- \frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan \left(\frac{1}{2} \left(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x) \right) \sqrt{-\sqrt{2} + 2} \right)$$

input `integrate(cosh(x)*sech(4*x),x, algorithm="fricas")`

output `1/4*sqrt(sqrt(2) + 2)*arctan(1/2*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (sqrt(2) - 2)*cosh(x) + (3*sqrt(2)*cosh(x)^2 + sqrt(2) - 2)*sinh(x))*sqrt(sqrt(2) + 2)) + 1/4*sqrt(sqrt(2) + 2)*arctan(1/2*(sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*sqrt(sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(1/2*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3 + (sqrt(2) + 2)*cosh(x) + (3*sqrt(2)*cosh(x)^2 + sqrt(2) + 2)*sinh(x))*sqrt(-sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(1/2*(sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*sqrt(-sqrt(2) + 2))`

Sympy [F]

$$\int \cosh(x) \operatorname{sech}(4x) dx = \int \cosh(x) \operatorname{sech}(4x) dx$$

input `integrate(cosh(x)*sech(4*x),x)`

output `Integral(cosh(x)*sech(4*x), x)`

Maxima [F]

$$\int \cosh(x)\operatorname{sech}(4x) dx = \int \cosh(x) \operatorname{sech}(4x) dx$$

input `integrate(cosh(x)*sech(4*x),x, algorithm="maxima")`

output `integrate(cosh(x)*sech(4*x), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 135 vs. $2(49) = 98$.

Time = 0.16 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.90

$$\begin{aligned} \int \cosh(x)\operatorname{sech}(4x) dx &= \frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan\left(\frac{\sqrt{\sqrt{2} + 2} + 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\ &+ \frac{1}{4} \sqrt{\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{\sqrt{2} + 2} - 2e^x}{\sqrt{-\sqrt{2} + 2}}\right) \\ &- \frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan\left(\frac{\sqrt{-\sqrt{2} + 2} + 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \\ &- \frac{1}{4} \sqrt{-\sqrt{2} + 2} \arctan\left(-\frac{\sqrt{-\sqrt{2} + 2} - 2e^x}{\sqrt{\sqrt{2} + 2}}\right) \end{aligned}$$

input `integrate(cosh(x)*sech(4*x),x, algorithm="giac")`

output `1/4*sqrt(sqrt(2) + 2)*arctan((sqrt(sqrt(2) + 2) + 2*e^x)/sqrt(-sqrt(2) + 2)) + 1/4*sqrt(sqrt(2) + 2)*arctan(-(sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan((sqrt(-sqrt(2) + 2) + 2*e^x)/sqrt(sqrt(2) + 2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(-(sqrt(-sqrt(2) + 2) - 2*e^x)/sqrt(sqrt(2) + 2))`

Mupad [B] (verification not implemented)

Time = 2.37 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.77

$$\int \cosh(x)\operatorname{sech}(4x) dx = \frac{\operatorname{atan}\left(\frac{3e^{2x}-2\sqrt{2}+2\sqrt{2}e^{2x}-3}{e^x\sqrt{2+2}+\sqrt{2}e^x\sqrt{2+2}}\right)\sqrt{\sqrt{2}+2}}{4} + \frac{\operatorname{atan}\left(\frac{3e^{2x}+2\sqrt{2}-2\sqrt{2}e^{2x}-3}{e^x\sqrt{2-\sqrt{2}}-\sqrt{2}e^x\sqrt{2-\sqrt{2}}}\right)\sqrt{2-\sqrt{2}}}{4}$$

input `int(cosh(x)/cosh(4*x), x)`output `(atan((3*exp(2*x) - 2*2^(1/2) + 2*2^(1/2)*exp(2*x) - 3)/(exp(x)*(2^(1/2) + 2)^(1/2) + 2^(1/2)*exp(x)*(2^(1/2) + 2)^(1/2)))*(2^(1/2) + 2)^(1/2))/4 + (atan((3*exp(2*x) + 2*2^(1/2) - 2*2^(1/2)*exp(2*x) - 3)/(exp(x)*(2 - 2^(1/2))^(1/2) - 2^(1/2)*exp(x)*(2 - 2^(1/2))^(1/2)))*(2 - 2^(1/2))^(1/2))/4`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 249, normalized size of antiderivative = 3.51

$$\int \cosh(x)\operatorname{sech}(4x) dx = \frac{\sqrt{\sqrt{2}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{\sqrt{2}+2}\operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{\sqrt{2}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}+2e^x}{\sqrt{\sqrt{2}+2}}\right)}{4} + \frac{\sqrt{\sqrt{2}+2}\operatorname{atan}\left(\frac{\sqrt{-\sqrt{2}+2}+2e^x}{\sqrt{\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{-\sqrt{2}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}-2e^x}{\sqrt{-\sqrt{2}+2}}\right)}{4} - \frac{\sqrt{-\sqrt{2}+2}\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}-2e^x}{\sqrt{-\sqrt{2}+2}}\right)}{4} + \frac{\sqrt{-\sqrt{2}+2}\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}+2e^x}{\sqrt{-\sqrt{2}+2}}\right)}{4} + \frac{\sqrt{-\sqrt{2}+2}\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}+2e^x}{\sqrt{-\sqrt{2}+2}}\right)}{4}$$

input `int(cosh(x)*sech(4*x),x)`output `(sqrt(sqrt(2) + 2)*sqrt(2)*atan((sqrt(-sqrt(2) + 2) - 2*e**x)/sqrt(sqrt(2) + 2)) - sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) - 2*e**x)/sqrt(sqrt(2) + 2)) - sqrt(sqrt(2) + 2)*sqrt(2)*atan((sqrt(-sqrt(2) + 2) + 2*e**x)/sqrt(sqrt(2) + 2)) + sqrt(sqrt(2) + 2)*atan((sqrt(-sqrt(2) + 2) + 2*e**x)/sqrt(sqrt(2) + 2)) - sqrt(-sqrt(2) + 2)*sqrt(2)*atan((sqrt(sqrt(2) + 2) - 2*e**x)/sqrt(-sqrt(2) + 2)) - sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) - 2*e**x)/sqrt(-sqrt(2) + 2)) + sqrt(-sqrt(2) + 2)*sqrt(2)*atan((sqrt(sqrt(2) + 2) + 2*e**x)/sqrt(-sqrt(2) + 2)) + sqrt(-sqrt(2) + 2)*atan((sqrt(sqrt(2) + 2) + 2*e**x)/sqrt(-sqrt(2) + 2)))/4`

3.125 $\int \cosh(x)\operatorname{sech}(5x) dx$

Optimal result	881
Mathematica [A] (verified)	881
Rubi [A] (verified)	882
Maple [C] (verified)	883
Fricas [B] (verification not implemented)	884
Sympy [F]	884
Maxima [F]	885
Giac [A] (verification not implemented)	885
Mupad [B] (verification not implemented)	886
Reduce [F]	887

Optimal result

Integrand size = 7, antiderivative size = 73

$$\int \cosh(x)\operatorname{sech}(5x) dx = -\sqrt{\frac{2}{5(5+\sqrt{5})}} \arctan\left(\sqrt{5-2\sqrt{5}} \tanh(x)\right) + \frac{1}{5}\sqrt{\frac{1}{2}(5+\sqrt{5})} \arctan\left(\sqrt{5+2\sqrt{5}} \tanh(x)\right)$$

output `-2^(1/2)/(25+5*5^(1/2))^(1/2)*arctan((5-2*5^(1/2))^(1/2)*tanh(x))+1/10*(10+2*5^(1/2))^(1/2)*arctan((5+2*5^(1/2))^(1/2)*tanh(x))`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.15

$$\int \cosh(x)\operatorname{sech}(5x) dx = \frac{\sqrt{5+\sqrt{5}} \arctan\left(\frac{(5+\sqrt{5}) \tanh(x)}{\sqrt{10-2\sqrt{5}}}\right) + \sqrt{5-\sqrt{5}} \arctan\left(\frac{(-5+\sqrt{5}) \tanh(x)}{\sqrt{2(5+\sqrt{5})}}\right)}{5\sqrt{2}}$$

input `Integrate[Cosh[x]*Sech[5*x],x]`

output

```
(Sqrt[5 + Sqrt[5]]*ArcTan[((5 + Sqrt[5])*Tanh[x])/Sqrt[10 - 2*Sqrt[5]]] +
Sqrt[5 - Sqrt[5]]*ArcTan[((-5 + Sqrt[5])*Tanh[x])/Sqrt[2*(5 + Sqrt[5])]])/
(5*Sqrt[2])
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.22, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4889, 1480, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \operatorname{sech}(5x) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\cos(ix)}{\cos(5ix)} dx \\
 & \quad \downarrow 4889 \\
 & \int \frac{1 - \tanh^2(x)}{5 \tanh^4(x) + 10 \tanh^2(x) + 1} d \tanh(x) \\
 & \quad \downarrow 1480 \\
 & -\frac{1}{2}(1 - \sqrt{5}) \int \frac{1}{5 \tanh^2(x) - 2\sqrt{5} + 5} d \tanh(x) - \\
 & \quad \frac{1}{2}(1 + \sqrt{5}) \int \frac{1}{5 \tanh^2(x) + 2\sqrt{5} + 5} d \tanh(x) \\
 & \quad \downarrow 216 \\
 & \frac{(1 + \sqrt{5}) \arctan(\sqrt{5 - 2\sqrt{5}} \tanh(x))}{2\sqrt{5}(5 + 2\sqrt{5})} - \frac{(1 - \sqrt{5}) \arctan(\sqrt{5 + 2\sqrt{5}} \tanh(x))}{2\sqrt{5}(5 - 2\sqrt{5})}
 \end{aligned}$$

input

```
Int[Cosh[x]*Sech[5*x], x]
```

output

$$-1/2*((1 + \sqrt{5})\operatorname{ArcTan}[\sqrt{5 - 2\sqrt{5}}\operatorname{Tanh}[x]])/\sqrt{5*(5 + 2\sqrt{5})} - ((1 - \sqrt{5})\operatorname{ArcTan}[\sqrt{5 + 2\sqrt{5}}\operatorname{Tanh}[x]])/(2*\sqrt{5*(5 - 2*\sqrt{5})}))$$
Defintions of rubi rules used

rule 216

$$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b\}, x] \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$$

rule 1480

$$\operatorname{Int}[(d_ + (e_)*(x_)^2)/((a_ + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] : > \operatorname{With}[\{q = \operatorname{Rt}[b^2 - 4*a*c, 2]\}, \operatorname{Simp}[(e/2 + (2*c*d - b*e)/(2*q)) \operatorname{Int}[1/(b/2 - q/2 + c*x^2), x], x] + \operatorname{Simp}[(e/2 - (2*c*d - b*e)/(2*q)) \operatorname{Int}[1/(b/2 + q/2 + c*x^2), x], x]] \text{ ; FreeQ}\{a, b, c, d, e\}, x] \ \&\& \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \operatorname{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \operatorname{PosQ}[b^2 - 4*a*c]$$

rule 3042

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4889

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{With}[\{v = \operatorname{FunctionOfTrig}[u, x]\}, \operatorname{With}[\{d = \operatorname{FreeFactors}[\operatorname{Tan}[v], x]\}, \operatorname{Simp}[d/\operatorname{Coefficient}[v, x, 1] \operatorname{Subst}[\operatorname{Int}[\operatorname{SubstFor}[1/(1 + d^2*x^2), \operatorname{Tan}[v]/d, u, x], x], x, \operatorname{Tan}[v]/d], x]] \text{ ; !FalseQ}[v] \ \&\& \operatorname{FunctionOfQ}[\operatorname{NonfreeFactors}[\operatorname{Tan}[v], x], u, x]] \text{ ; InverseFunctionFreeQ}[u, x] \ \&\& \operatorname{!MatchQ}[u, (v_)*((c_)*\operatorname{tan}[w_]^(n_)*\operatorname{tan}[z_]^(n_))^(p_)] \text{ ; FreeQ}\{c, p\}, x] \ \&\& \operatorname{IntegerQ}[n] \ \&\& \operatorname{LinearQ}[w, x] \ \&\& \operatorname{EqQ}[z, 2*w]]$$
Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.56

method	result	size
risch	$2 \left(\sum_{R=\operatorname{RootOf}(32000_Z^4+400_Z^2+1)} -R \ln(-4000_R^3 + 200_R^2 + e^{2x} - 30_R + 1) \right)$	41

input `int(cosh(x)*sech(5*x),x,method=_RETURNVERBOSE)`

output `2*sum(_R*ln(-4000*_R^3+200*_R^2+exp(2*x)-30*_R+1),_R=RootOf(32000*_Z^4+400*_Z^2+1))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(52) = 104$.

Time = 0.09 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.58

$$\int \cosh(x) \operatorname{sech}(5x) dx$$

$$= \frac{1}{5} \sqrt{\frac{1}{2} \sqrt{5} + \frac{5}{2}} \arctan \left(\frac{1}{10} \left(4 \sqrt{5} \cosh(x)^2 + 8 \sqrt{5} \cosh(x) \sinh(x) + 4 \sqrt{5} \sinh(x)^2 - \sqrt{5} - 5 \right) \sqrt{\frac{1}{2} \sqrt{5}} \right) - \frac{1}{5} \sqrt{-\frac{1}{2} \sqrt{5} + \frac{5}{2}} \arctan \left(\frac{1}{10} \left(4 \sqrt{5} \cosh(x)^2 + 8 \sqrt{5} \cosh(x) \sinh(x) + 4 \sqrt{5} \sinh(x)^2 - \sqrt{5} + 5 \right) \sqrt{-\frac{1}{2} \sqrt{5}} \right)$$

input `integrate(cosh(x)*sech(5*x),x, algorithm="fricas")`

output `1/5*sqrt(1/2*sqrt(5) + 5/2)*arctan(1/10*(4*sqrt(5)*cosh(x)^2 + 8*sqrt(5)*cosh(x)*sinh(x) + 4*sqrt(5)*sinh(x)^2 - sqrt(5) - 5)*sqrt(1/2*sqrt(5) + 5/2)) - 1/5*sqrt(-1/2*sqrt(5) + 5/2)*arctan(1/10*(4*sqrt(5)*cosh(x)^2 + 8*sqrt(5)*cosh(x)*sinh(x) + 4*sqrt(5)*sinh(x)^2 - sqrt(5) + 5)*sqrt(-1/2*sqrt(5) + 5/2))`

Sympy [F]

$$\int \cosh(x) \operatorname{sech}(5x) dx = \int \cosh(x) \operatorname{sech}(5x) dx$$

input `integrate(cosh(x)*sech(5*x),x)`

output `Integral(cosh(x)*sech(5*x), x)`

Maxima [F]

$$\int \cosh(x)\operatorname{sech}(5x) dx = \int \cosh(x) \operatorname{sech}(5x) dx$$

input `integrate(cosh(x)*sech(5*x),x, algorithm="maxima")`

output `1/5*sqrt(5)*arctan((sqrt(5) + 4*e^(-2*x) - 1)/sqrt(2*sqrt(5) + 10))/sqrt(2*sqrt(5) + 10) - 1/5*sqrt(5)*arctan(-(sqrt(5) - 4*e^(-2*x) + 1)/sqrt(-2*sqrt(5) + 10))/sqrt(-2*sqrt(5) + 10) - 1/10*log(-(sqrt(5) + 1)*e^(-2*x) + 2*e^(-4*x) + 2)/(sqrt(5) + 1) + 1/10*log((sqrt(5) - 1)*e^(-2*x) + 2*e^(-4*x) + 2)/(sqrt(5) - 1) - 1/5*integrate((e^(7*x) - 2*e^(5*x) - 2*e^(3*x) + e^x)*e^x/(e^(8*x) - e^(6*x) + e^(4*x) - e^(2*x) + 1), x) + 1/10*log(e^(2*x) + 1) - 1/10*log(e^(-2*x) + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.93

$$\int \cosh(x)\operatorname{sech}(5x) dx = -\frac{1}{10} \sqrt{-2\sqrt{5} + 10} \arctan\left(\frac{\sqrt{5} + 4e^{2x} - 1}{\sqrt{2\sqrt{5} + 10}}\right) + \frac{1}{10} \sqrt{2\sqrt{5} + 10} \arctan\left(-\frac{\sqrt{5} - 4e^{2x} + 1}{\sqrt{-2\sqrt{5} + 10}}\right)$$

input `integrate(cosh(x)*sech(5*x),x, algorithm="giac")`

output `-1/10*sqrt(-2*sqrt(5) + 10)*arctan((sqrt(5) + 4*e^(2*x) - 1)/sqrt(2*sqrt(5) + 10)) + 1/10*sqrt(2*sqrt(5) + 10)*arctan(-(sqrt(5) - 4*e^(2*x) + 1)/sqrt(-2*sqrt(5) + 10))`

Mupad [B] (verification not implemented)

Time = 3.55 (sec) , antiderivative size = 297, normalized size of antiderivative = 4.07

$$\begin{aligned}
\int \cosh(x)\operatorname{sech}(5x) dx = & \ln \left(1 - \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(4e^{2x} \right. \right. \\
& + \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(48e^{2x} + \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} (360e^{2x} - 360) - 72 \right) \\
& \left. \left. - 8 \right) - e^{2x} \right) \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} - \ln \left(\sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(4e^{2x} \right. \right. \\
& + \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(\sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} (360e^{2x} - 360) - 48e^{2x} + 72 \right) \\
& \left. \left. - 8 \right) - e^{2x} + 1 \right) \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} - \ln \left(\sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(4e^{2x} \right. \right. \\
& + \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(\sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} (360e^{2x} - 360) - 48e^{2x} + 72 \right) \\
& \left. \left. - 8 \right) - e^{2x} + 1 \right) \sqrt{-\frac{\sqrt{5}}{200} - \frac{1}{40}} + \ln \left(1 - \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(4e^{2x} \right. \right. \\
& + \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} \left(48e^{2x} + \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}} (360e^{2x} - 360) - 72 \right) \\
& \left. \left. - 8 \right) - e^{2x} \right) \sqrt{\frac{\sqrt{5}}{200} - \frac{1}{40}}
\end{aligned}$$

input `int(cosh(x)/cosh(5*x), x)`

output

```

log(1 - (- 5^(1/2)/200 - 1/40)^(1/2)*(4*exp(2*x) + (- 5^(1/2)/200 - 1/40)^(1/2)*(48*exp(2*x) + (- 5^(1/2)/200 - 1/40)^(1/2)*(360*exp(2*x) - 360) - 72) - 8) - exp(2*x))*(- 5^(1/2)/200 - 1/40)^(1/2) - log((5^(1/2)/200 - 1/40)^(1/2)*(4*exp(2*x) + (5^(1/2)/200 - 1/40)^(1/2)*((5^(1/2)/200 - 1/40)^(1/2)*(360*exp(2*x) - 360) - 48*exp(2*x) + 72) - 8) - exp(2*x) + 1)*(5^(1/2)/200 - 1/40)^(1/2) - log((- 5^(1/2)/200 - 1/40)^(1/2)*(4*exp(2*x) + (- 5^(1/2)/200 - 1/40)^(1/2)*((- 5^(1/2)/200 - 1/40)^(1/2)*(360*exp(2*x) - 360) - 48*exp(2*x) + 72) - 8) - exp(2*x) + 1)*(- 5^(1/2)/200 - 1/40)^(1/2) + log(1 - (5^(1/2)/200 - 1/40)^(1/2)*(4*exp(2*x) + (5^(1/2)/200 - 1/40)^(1/2)*(48*exp(2*x) + (5^(1/2)/200 - 1/40)^(1/2)*(360*exp(2*x) - 360) - 72) - 8) - exp(2*x))*(5^(1/2)/200 - 1/40)^(1/2)

```

Reduce [F]

$$\int \cosh(x)\operatorname{sech}(5x) dx = \int \cosh(x) \operatorname{sech}(5x) dx$$

input

```
int(cosh(x)*sech(5*x),x)
```

output

```
int(cosh(x)*sech(5*x),x)
```


3.126 $\int \cosh(x)\operatorname{sech}(6x) dx$

Optimal result	888
Mathematica [A] (verified)	888
Rubi [A] (verified)	889
Maple [C] (verified)	890
Fricas [B] (verification not implemented)	891
Sympy [F]	892
Maxima [F]	892
Giac [B] (verification not implemented)	893
Mupad [B] (verification not implemented)	894
Reduce [B] (verification not implemented)	895

Optimal result

Integrand size = 7, antiderivative size = 85

$$\int \cosh(x)\operatorname{sech}(6x) dx = -\frac{\arctan(\sqrt{2}\sinh(x))}{3\sqrt{2}} + \frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{2}-\sqrt{3}}\right)}{6\sqrt{2}-\sqrt{3}} + \frac{\arctan\left(\frac{2\sinh(x)}{\sqrt{2}+\sqrt{3}}\right)}{6\sqrt{2}+\sqrt{3}}$$

output

```
-1/6*arctan(sinh(x)*2^(1/2))*2^(1/2)+1/6*arctan(2*sinh(x)/(1/2*6^(1/2)-1/2
*2^(1/2)))/(1/2*6^(1/2)-1/2*2^(1/2))+1/6*arctan(2*sinh(x)/(1/2*6^(1/2)+1/2
*2^(1/2)))/(1/2*6^(1/2)+1/2*2^(1/2))
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.95

$$\int \cosh(x)\operatorname{sech}(6x) dx = \frac{1}{6} \left(-\sqrt{2} \arctan(\sqrt{2}\sinh(x)) + \sqrt{2+\sqrt{3}} \arctan\left(\frac{2\sinh(x)}{\sqrt{2}-\sqrt{3}}\right) + \sqrt{2-\sqrt{3}} \arctan\left(\frac{2\sinh(x)}{\sqrt{2}+\sqrt{3}}\right) \right)$$

input

```
Integrate[Cosh[x]*Sech[6*x],x]
```

output

```
(-(Sqrt[2]*ArcTan[Sqrt[2]*Sinh[x]]) + Sqrt[2 + Sqrt[3]]*ArcTan[(2*Sinh[x])
/Sqrt[2 - Sqrt[3]]] + Sqrt[2 - Sqrt[3]]*ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[3]
]])/6
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 4856, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \operatorname{sech}(6x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(ix)}{\cos(6ix)} dx \\
 & \quad \downarrow \text{4856} \\
 & \int \frac{1}{32 \sinh^6(x) + 48 \sinh^4(x) + 18 \sinh^2(x) + 1} d \sinh(x) \\
 & \quad \downarrow \text{2460} \\
 & \int \left(\frac{4(2 \sinh^2(x) + 1)}{3(16 \sinh^4(x) + 16 \sinh^2(x) + 1)} - \frac{1}{3(2 \sinh^2(x) + 1)} \right) d \sinh(x) \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\arctan(\sqrt{2} \sinh(x))}{3\sqrt{2}} + \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{2-\sqrt{3}}}\right)}{6\sqrt{2-\sqrt{3}}} + \frac{\arctan\left(\frac{2 \sinh(x)}{\sqrt{2+\sqrt{3}}}\right)}{6\sqrt{2+\sqrt{3}}}
 \end{aligned}$$

input

```
Int[Cosh[x]*Sech[6*x],x]
```

output

```
-1/3*ArcTan[Sqrt[2]*Sinh[x]]/Sqrt[2] + ArcTan[(2*Sinh[x])/Sqrt[2 - Sqrt[3]]]/(6*Sqrt[2 - Sqrt[3]]) + ArcTan[(2*Sinh[x])/Sqrt[2 + Sqrt[3]]]/(6*Sqrt[2 + Sqrt[3]])
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2460

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4856

```
Int[(u_.)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.21 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.98

method	result
risch	$2 \left(\sum_{R=\text{RootOf}(331776_Z^4+2304_Z^2+1)} -R \ln(e^{2x} + (13824_R^3 + 96_R) e^x - 1) \right) + \frac{i\sqrt{2} \ln(e^{2x} - i\sqrt{2}e^x)}{12}$

input

```
int(cosh(x)*sech(6*x), x, method=_RETURNVERBOSE)
```

output

```
2*sum(_R*ln(exp(2*x)+(13824*_R^3+96*_R)*exp(x)-1),_R=RootOf(331776*_Z^4+2304*_Z^2+1))+1/12*I*2^(1/2)*ln(exp(2*x)-I*2^(1/2)*exp(x)-1)-1/12*I*2^(1/2)*ln(exp(2*x)+I*2^(1/2)*exp(x)-1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(67) = 134$.

Time = 0.10 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.81

$$\int \cosh(x) \operatorname{sech}(6x) dx$$

$$= \frac{1}{6} \sqrt{\sqrt{3} + 2} \arctan \left(\left(\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + \sinh(x)^3 - (\sqrt{3} - 1) \cosh(x) + (3 \cosh(x)^2 - \sinh(x)) \right) \right)$$

$$+ \frac{1}{6} \sqrt{-\sqrt{3} + 2} \arctan \left(\left(\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + \sinh(x)^3 + (\sqrt{3} + 1) \cosh(x) + (3 \cosh(x)^2 + \sinh(x)) \right) \right)$$

$$+ \frac{1}{6} \sqrt{\sqrt{3} + 2} \arctan \left(\sqrt{\sqrt{3} + 2} (\cosh(x) + \sinh(x)) \right)$$

$$+ \frac{1}{6} \sqrt{-\sqrt{3} + 2} \arctan \left(\sqrt{-\sqrt{3} + 2} (\cosh(x) + \sinh(x)) \right)$$

$$- \frac{1}{6} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \cosh(x) + \frac{1}{2} \sqrt{2} \sinh(x) \right)$$

$$+ \frac{1}{6} \sqrt{2} \arctan \left(-\frac{\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 + \sqrt{2}}{2(\cosh(x) - \sinh(x))} \right)$$

input

```
integrate(cosh(x)*sech(6*x),x, algorithm="fricas")
```

output

```
1/6*sqrt(sqrt(3) + 2)*arctan((cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + sinh(x)^3 - (sqrt(3) - 1)*cosh(x) + (3*cosh(x)^2 - sqrt(3) + 1)*sinh(x))*sqrt(sqrt(3) + 2)) + 1/6*sqrt(-sqrt(3) + 2)*arctan((cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + sinh(x)^3 + (sqrt(3) + 1)*cosh(x) + (3*cosh(x)^2 + sqrt(3) + 1)*sinh(x))*sqrt(-sqrt(3) + 2)) + 1/6*sqrt(sqrt(3) + 2)*arctan(sqrt(sqrt(3) + 2)*(cosh(x) + sinh(x))) + 1/6*sqrt(-sqrt(3) + 2)*arctan(sqrt(-sqrt(3) + 2)*(cosh(x) + sinh(x))) - 1/6*sqrt(2)*arctan(1/2*sqrt(2)*cosh(x) + 1/2*sqrt(2)*sinh(x)) + 1/6*sqrt(2)*arctan(-1/2*(sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2))/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \cosh(x)\operatorname{sech}(6x) dx = \int \cosh(x) \operatorname{sech}(6x) dx$$

input `integrate(cosh(x)*sech(6*x),x)`

output `Integral(cosh(x)*sech(6*x), x)`

Maxima [F]

$$\int \cosh(x)\operatorname{sech}(6x) dx = \int \cosh(x) \operatorname{sech}(6x) dx$$

input `integrate(cosh(x)*sech(6*x),x, algorithm="maxima")`

output `-1/6*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/6*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) + integrate(1/3*(e^(7*x) + e^(5*x) + e^(3*x) + e^x)/(e^(8*x) - e^(4*x) + 1), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 177 vs. $2(67) = 134$.

Time = 0.18 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.08

$$\begin{aligned} \int \cosh(x)\operatorname{sech}(6x) dx &= \frac{1}{12} (\sqrt{6} - \sqrt{2}) \arctan \left(\frac{\sqrt{6} - \sqrt{2} + 4e^x}{\sqrt{6} + \sqrt{2}} \right) \\ &+ \frac{1}{12} (\sqrt{6} - \sqrt{2}) \arctan \left(-\frac{\sqrt{6} - \sqrt{2} - 4e^x}{\sqrt{6} + \sqrt{2}} \right) \\ &+ \frac{1}{12} (\sqrt{6} + \sqrt{2}) \arctan \left(\frac{\sqrt{6} + \sqrt{2} + 4e^x}{\sqrt{6} - \sqrt{2}} \right) \\ &+ \frac{1}{12} (\sqrt{6} + \sqrt{2}) \arctan \left(-\frac{\sqrt{6} + \sqrt{2} - 4e^x}{\sqrt{6} - \sqrt{2}} \right) \\ &- \frac{1}{6} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2e^x) \right) \\ &- \frac{1}{6} \sqrt{2} \arctan \left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2e^x) \right) \end{aligned}$$

input `integrate(cosh(x)*sech(6*x),x, algorithm="giac")`

output `1/12*(sqrt(6) - sqrt(2))*arctan((sqrt(6) - sqrt(2) + 4*e^x)/(sqrt(6) + sqrt(2))) + 1/12*(sqrt(6) - sqrt(2))*arctan(-(sqrt(6) - sqrt(2) - 4*e^x)/(sqrt(6) + sqrt(2))) + 1/12*(sqrt(6) + sqrt(2))*arctan((sqrt(6) + sqrt(2) + 4*e^x)/(sqrt(6) - sqrt(2))) + 1/12*(sqrt(6) + sqrt(2))*arctan(-(sqrt(6) + sqrt(2) - 4*e^x)/(sqrt(6) - sqrt(2))) - 1/6*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/6*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x))`

Mupad [B] (verification not implemented)

Time = 3.11 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.42

$$\int \cosh(x)\operatorname{sech}(6x) dx = \frac{\sqrt{2} \operatorname{atan}\left(\frac{7e^{2x} + 4\sqrt{3} - 4\sqrt{3}e^{2x} - 7}{\frac{5\sqrt{2}e^x}{2} - \frac{3\sqrt{6}e^x}{2}}\right)}{12} + \frac{\sqrt{2} \operatorname{atan}\left(\frac{7e^{2x} - 4\sqrt{3} + 4\sqrt{3}e^{2x} - 7}{\frac{5\sqrt{2}e^x}{2} + \frac{3\sqrt{6}e^x}{2}}\right)}{12} - \frac{\sqrt{6} \operatorname{atan}\left(\frac{7e^{2x} + 4\sqrt{3} - 4\sqrt{3}e^{2x} - 7}{\frac{5\sqrt{2}e^x}{2} - \frac{3\sqrt{6}e^x}{2}}\right)}{12} + \frac{\sqrt{6} \operatorname{atan}\left(\frac{7e^{2x} - 4\sqrt{3} + 4\sqrt{3}e^{2x} - 7}{\frac{5\sqrt{2}e^x}{2} + \frac{3\sqrt{6}e^x}{2}}\right)}{12} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}e^{-x}(e^{2x} - 1)}{2}\right)}{6}$$

input `int(cosh(x)/cosh(6*x),x)`

output

```
(2^(1/2)*atan((7*exp(2*x) + 4*3^(1/2) - 4*3^(1/2)*exp(2*x) - 7)/((5*2^(1/2)*exp(x))/2 - (3*6^(1/2)*exp(x))/2))/12 + (2^(1/2)*atan((7*exp(2*x) - 4*3^(1/2) + 4*3^(1/2)*exp(2*x) - 7)/((5*2^(1/2)*exp(x))/2 + (3*6^(1/2)*exp(x))/2))/12 - (6^(1/2)*atan((7*exp(2*x) + 4*3^(1/2) - 4*3^(1/2)*exp(2*x) - 7)/((5*2^(1/2)*exp(x))/2 - (3*6^(1/2)*exp(x))/2))/12 + (6^(1/2)*atan((7*exp(2*x) - 4*3^(1/2) + 4*3^(1/2)*exp(2*x) - 7)/((5*2^(1/2)*exp(x))/2 + (3*6^(1/2)*exp(x))/2))/12 - (2^(1/2)*atan((2^(1/2)*exp(-x)*(exp(2*x) - 1))/2))/6
```

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 287, normalized size of antiderivative = 3.38

$$\begin{aligned}
\int \cosh(x)\operatorname{sech}(6x) dx = & \frac{\sqrt{-\sqrt{3}+2}\sqrt{3}\operatorname{atan}\left(\frac{4e^x-\sqrt{6}-\sqrt{2}}{2\sqrt{-\sqrt{3}+2}}\right)}{6} \\
& + \frac{\sqrt{-\sqrt{3}+2}\operatorname{atan}\left(\frac{4e^x-\sqrt{6}-\sqrt{2}}{2\sqrt{-\sqrt{3}+2}}\right)}{3} \\
& + \frac{\sqrt{-\sqrt{3}+2}\sqrt{3}\operatorname{atan}\left(\frac{4e^x+\sqrt{6}+\sqrt{2}}{2\sqrt{-\sqrt{3}+2}}\right)}{6} \\
& + \frac{\sqrt{-\sqrt{3}+2}\operatorname{atan}\left(\frac{4e^x+\sqrt{6}+\sqrt{2}}{2\sqrt{-\sqrt{3}+2}}\right)}{3} \\
& - \frac{\sqrt{2}\operatorname{atan}\left(\frac{2e^x-\sqrt{2}}{\sqrt{2}}\right)}{6} - \frac{\sqrt{2}\operatorname{atan}\left(\frac{2e^x+\sqrt{2}}{\sqrt{2}}\right)}{6} \\
& - \frac{\sqrt{6}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}-4e^x}{\sqrt{6}+\sqrt{2}}\right)}{12} + \frac{\sqrt{2}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}-4e^x}{\sqrt{6}+\sqrt{2}}\right)}{12} \\
& + \frac{\sqrt{6}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}+4e^x}{\sqrt{6}+\sqrt{2}}\right)}{12} - \frac{\sqrt{2}\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{3}+2}+4e^x}{\sqrt{6}+\sqrt{2}}\right)}{12}
\end{aligned}$$

input `int(cosh(x)*sech(6*x),x)`

output

```

(2*sqrt(-sqrt(3)+2)*sqrt(3)*atan((4*e**x - sqrt(6) - sqrt(2))/(2*sqrt(-sqrt(3)+2))) + 4*sqrt(-sqrt(3)+2)*atan((4*e**x - sqrt(6) - sqrt(2))/(2*sqrt(-sqrt(3)+2))) + 2*sqrt(-sqrt(3)+2)*sqrt(3)*atan((4*e**x + sqrt(6) + sqrt(2))/(2*sqrt(-sqrt(3)+2))) + 4*sqrt(-sqrt(3)+2)*atan((4*e**x + sqrt(6) + sqrt(2))/(2*sqrt(-sqrt(3)+2))) - 2*sqrt(2)*atan((2*e**x - sqrt(2))/sqrt(2)) - 2*sqrt(2)*atan((2*e**x + sqrt(2))/sqrt(2)) - sqrt(6)*atan((2*sqrt(-sqrt(3)+2) - 4*e**x)/(sqrt(6) + sqrt(2))) + sqrt(2)*atan((2*sqrt(-sqrt(3)+2) - 4*e**x)/(sqrt(6) + sqrt(2))) + sqrt(6)*atan((2*sqrt(-sqrt(3)+2) + 4*e**x)/(sqrt(6) + sqrt(2))) - sqrt(2)*atan((2*sqrt(-sqrt(3)+2) + 4*e**x)/(sqrt(6) + sqrt(2))))/12

```


3.127 $\int \cosh(a + bx)\operatorname{sech}(c + bx) dx$

Optimal result	896
Mathematica [A] (verified)	896
Rubi [A] (verified)	897
Maple [B] (verified)	898
Fricas [B] (verification not implemented)	899
Sympy [F]	899
Maxima [A] (verification not implemented)	899
Giac [A] (verification not implemented)	900
Mupad [B] (verification not implemented)	900
Reduce [B] (verification not implemented)	901

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \cosh(a + bx)\operatorname{sech}(c + bx) dx = x \cosh(a - c) + \frac{\log(\cosh(c + bx)) \sinh(a - c)}{b}$$

output `x*cosh(a-c)+ln(cosh(b*x+c))*sinh(a-c)/b`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \cosh(a + bx)\operatorname{sech}(c + bx) dx = x \cosh(a - c) + \frac{\log(\cosh(c + bx)) \sinh(a - c)}{b}$$

input `Integrate[Cosh[a + b*x]*Sech[c + b*x],x]`

output `x*Cosh[a - c] + (Log[Cosh[c + b*x]]*Sinh[a - c])/b`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6161, 24, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{sech}(bx + c) dx \\
 & \quad \downarrow \text{6161} \\
 & \sinh(a - c) \int \tanh(c + bx) dx + \cosh(a - c) \int 1 dx \\
 & \quad \downarrow \text{24} \\
 & \sinh(a - c) \int \tanh(c + bx) dx + x \cosh(a - c) \\
 & \quad \downarrow \text{3042} \\
 & x \cosh(a - c) + \sinh(a - c) \int -i \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & x \cosh(a - c) - i \sinh(a - c) \int \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\sinh(a - c) \log(\cosh(bx + c))}{b} + x \cosh(a - c)
 \end{aligned}$$

input

```
Int[Cosh[a + b*x]*Sech[c + b*x],x]
```

output

```
x*Cosh[a - c] + (Log[Cosh[c + b*x]]*Sinh[a - c])/b
```

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6161 `Int[Cosh[v_]*Sech[w_]^(n_.), x_Symbol] := Simp[Sinh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Cosh[v - w] Int[Sech[w]^(n - 1), x], x] /; GetQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 145 vs. $2(26) = 52$.

Time = 0.22 (sec) , antiderivative size = 146, normalized size of antiderivative = 5.62

method	result
risch	$x e^{a-c} - e^{-a-c} e^{2a} x + e^{-a-c} e^{2c} x - \frac{e^{-a-c} e^{2a} a}{b} + \frac{e^{-a-c} e^{2c} a}{b} + \frac{\ln(e^{2bx+2a} + e^{2a-2c}) e^{-a-c} e^{2a}}{2b} - \frac{\ln(e^{2bx+2a} + e^{2a-2c})}{2b}$

input `int(cosh(b*x+a)*sech(b*x+c),x,method=_RETURNVERBOSE)`

output `x*exp(a-c)-exp(-a-c)*exp(2*a)*x+exp(-a-c)*exp(2*c)*x-1/b*exp(-a-c)*exp(2*a)*a+1/b*exp(-a-c)*exp(2*c)*a+1/2*ln(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-a-c)*exp(2*a)-1/2*ln(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(26) = 52$.

Time = 0.09 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.31

$$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx$$

$$= \frac{2bx + (\cosh(-a + c))^2 - 2 \cosh(-a + c) \sinh(-a + c) + \sinh(-a + c)^2 - 1}{2(b \cosh(-a + c) - b \sinh(-a + c))} \log\left(\frac{2 \cosh(bx + c)}{\cosh(bx + c) - \sinh(bx + c)}\right)$$

input `integrate(cosh(b*x+a)*sech(b*x+c),x, algorithm="fricas")`

output `1/2*(2*b*x + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c)))/(b*cosh(-a + c) - b*sinh(-a + c))`

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx = \int \cosh(a + bx) \operatorname{sech}(bx + c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x+c),x)`

output `Integral(cosh(a + b*x)*sech(b*x + c), x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.96

$$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx$$

$$= \frac{(e^{2a} - e^{2c})e^{(-a-c)} \log(e^{(-2bx)} + e^{(2c)})}{2b} + \frac{(bx + a)e^{(a-c)}}{b}$$

input `integrate(cosh(b*x+a)*sech(b*x+c),x, algorithm="maxima")`

output $\frac{1}{2}(e^{2a} - e^{2c})e^{-a-c} \log(e^{-2bx} + e^{2c})/b + (bx + a)e^{a-c}/b$

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.81

$$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx = x e^{(-a+c)} + \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(2bx+2c)} + 1)}{2b}$$

input `integrate(cosh(b*x+a)*sech(b*x+c),x, algorithm="giac")`

output $x e^{-a+c} + \frac{1}{2}(e^{2a+c} - e^{3c})e^{-a-2c} \log(e^{2bx+2c} + 1)/b$

Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.46

$$\int \cosh(a + bx) \operatorname{sech}(c + bx) dx = x e^{c-a} + \frac{e^{2c-2a} \ln(e^{2a} e^{2bx} + e^{2a} e^{-2c}) (2b e^{3a-3c} - 2b e^{a-c})}{4b^2}$$

input `int(cosh(a + b*x)/cosh(c + b*x),x)`

output $x \exp(c - a) + (\exp(2c - 2a) \log(\exp(2a) \exp(2bx) + \exp(2a) \exp(-2c))) * (2b \exp(3a - 3c) - 2b \exp(a - c)) / (4b^2)$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.35

$$\int \cosh(a + bx)\operatorname{sech}(c + bx) dx = \frac{e^{2a}\log(e^{2bx+2c} + 1) - e^{2c}\log(e^{2bx+2c} + 1) + 2e^{2c}bx}{2e^{a+cb}}$$

input `int(cosh(b*x+a)*sech(b*x+c),x)`

output `(e**(2*a)*log(e**(2*b*x + 2*c) + 1) - e**(2*c)*log(e**(2*b*x + 2*c) + 1) + 2*e**(2*c)*b*x)/(2*e**(a + c)*b)`

3.128 $\int \cosh(a + bx)\operatorname{sech}^2(c + bx) dx$

Optimal result	902
Mathematica [B] (verified)	902
Rubi [A] (verified)	903
Maple [C] (verified)	905
Fricas [B] (verification not implemented)	905
Sympy [F]	906
Maxima [A] (verification not implemented)	906
Giac [A] (verification not implemented)	907
Mupad [B] (verification not implemented)	907
Reduce [B] (verification not implemented)	908

Optimal result

Integrand size = 15, antiderivative size = 35

$$\int \cosh(a + bx)\operatorname{sech}^2(c + bx) dx = \frac{\arctan(\sinh(c + bx)) \cosh(a - c)}{b} - \frac{\operatorname{sech}(c + bx) \sinh(a - c)}{b}$$

output

```
arctan(sinh(b*x+c))*cosh(a-c)/b-sech(b*x+c)*sinh(a-c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 83 vs. 2(35) = 70.

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.37

$$\int \cosh(a + bx)\operatorname{sech}^2(c + bx) dx = \frac{2 \arctan\left(\frac{(\cosh(c) - \sinh(c))\left(\cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh(c) \sinh\left(\frac{bx}{2}\right)\right)}{\cosh(c) \cosh\left(\frac{bx}{2}\right) - \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right) \cosh(a - c)}{b} - \frac{\operatorname{sech}(c + bx) \sinh(a - c)}{b}$$

input `Integrate[Cosh[a + b*x]*Sech[c + b*x]^2,x]`

output `(2*ArcTan[((Cosh[c] - Sinh[c])*(Cosh[(b*x)/2]*Sinh[c] + Cosh[c]*Sinh[(b*x)/2]))/(Cosh[c]*Cosh[(b*x)/2] - Cosh[(b*x)/2]*Sinh[c])]*Cosh[a - c])/b - (Sech[c + b*x]*Sinh[a - c])/b`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6161, 3042, 26, 3086, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{sech}^2(bx + c) dx \\
 & \quad \downarrow 6161 \\
 & \cosh(a - c) \int \operatorname{sech}(c + bx) dx + \sinh(a - c) \int \operatorname{sech}(c + bx) \tanh(c + bx) dx \\
 & \quad \downarrow 3042 \\
 & \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx + \sinh(a - c) \int -i \sec(ic + ibx) \tan(ic + ibx) dx \\
 & \quad \downarrow 26 \\
 & \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx - i \sinh(a - c) \int \sec(ic + ibx) \tan(ic + ibx) dx \\
 & \quad \downarrow 3086 \\
 & -\frac{\sinh(a - c) \int 1 d\operatorname{sech}(c + bx)}{b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow 24 \\
 & -\frac{\sinh(a - c) \operatorname{sech}(bx + c)}{b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow 4257
 \end{aligned}$$

$$\frac{\cosh(a - c) \arctan(\sinh(bx + c))}{b} - \frac{\sinh(a - c) \operatorname{sech}(bx + c)}{b}$$

input `Int[Cosh[a + b*x]*Sech[c + b*x]^2,x]`

output `(ArcTan[Sinh[c + b*x]]*Cosh[a - c])/b - (Sech[c + b*x]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(F_x_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6161 `Int[Cosh[v_]*Sech[w_]^(n_), x_Symbol] := Simp[Sinh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Cosh[v - w] Int[Sech[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 183, normalized size of antiderivative = 5.23

method	result
risch	$-\frac{e^{bx+a}(e^{2a}-e^{2c})}{b(e^{2bx+2a+2c}+e^{2a})} + \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2c}}{2b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*sech(b*x+c)^2,x,method=_RETURNVERBOSE)`

output
$$-1/b*\exp(b*x+a)*(exp(2*a)-exp(2*c))/(exp(2*b*x+2*a+2*c)+exp(2*a))+1/2*I*\ln(\exp(b*x+a)+I*\exp(a-c))/b*\exp(-a-c)*\exp(2*a)+1/2*I*\ln(\exp(b*x+a)+I*\exp(a-c))/b*\exp(-a-c)*\exp(2*c)-1/2*I*\ln(\exp(b*x+a)-I*\exp(a-c))/b*\exp(-a-c)*\exp(2*a)-1/2*I*\ln(\exp(b*x+a)-I*\exp(a-c))/b*\exp(-a-c)*\exp(2*c)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(35) = 70$.

Time = 0.10 (sec) , antiderivative size = 405, normalized size of antiderivative = 11.57

$$\int \cosh(a+bx)\operatorname{sech}^2(c+bx) dx$$

$$= \frac{2 \cosh(bx+c) \cosh(-a+c) \sinh(-a+c) - \cosh(bx+c) \sinh(-a+c)^2 + ((\cosh(-a+c))^2 + 1) \cosh(-a+c)}{2b}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^2,x, algorithm="fricas")`

output

```
(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2
+ ((cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)
*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^2 + 1
)*sinh(-a + c)^2 + cosh(-a + c)^2 - 2*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*cosh(b*x + c)
)*sinh(b*x + c) - 2*(cosh(b*x + c)^2*cosh(-a + c) + cosh(-a + c))*sinh(-a
+ c) + 1)*arctan(cosh(b*x + c) + sinh(b*x + c)) - (cosh(-a + c)^2 - 1)*cos
h(b*x + c) - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^
2 - 1)*sinh(b*x + c))/(b*cosh(b*x + c)^2*cosh(-a + c) + (b*cosh(-a + c) -
b*sinh(-a + c))*sinh(b*x + c)^2 + b*cosh(-a + c) + 2*(b*cosh(b*x + c)*cosh
(-a + c) - b*cosh(b*x + c)*sinh(-a + c))*sinh(b*x + c) - (b*cosh(b*x + c)^
2 + b)*sinh(-a + c))
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx = \int \cosh(a + bx) \operatorname{sech}^2(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*sech(b*x+c)**2,x)
```

output

```
Integral(cosh(a + b*x)*sech(b*x + c)**2, x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.00

$$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx = -\frac{(e^{(2a)} + e^{(2c)}) \arctan(e^{(-bx-c)}) e^{(-a-c)}}{b} - \frac{(e^{(2a)} - e^{(2c)}) e^{(-bx-a)}}{b(e^{(-2bx)} + e^{(2c)})}$$

input

```
integrate(cosh(b*x+a)*sech(b*x+c)^2,x, algorithm="maxima")
```

output

```
-(e^(2*a) + e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b - (e^(2*a) - e^(2*c)
))*e^(-b*x - a)/(b*(e^(-2*b*x) + e^(2*c)))
```

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.00

$$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx = \frac{(e^{(2a)} + e^{(2c)}) \arctan(e^{(bx+c)}) e^{(-a-c)}}{b} - \frac{(e^{(bx+2a)} - e^{(bx+2c)}) e^{(-a)}}{b(e^{(2bx+2c)} + 1)}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^2,x, algorithm="giac")`output `(e^(2*a) + e^(2*c))*arctan(e^(b*x + c))*e^(-a - c)/b - (e^(b*x + 2*a) - e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) + 1))`**Mupad [B] (verification not implemented)**

Time = 1.17 (sec) , antiderivative size = 148, normalized size of antiderivative = 4.23

$$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx = \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{b^2 + e^{2a}} e^{-2c} \sqrt{b^2})}{b \sqrt{e^{-2a} e^{2c} (2e^{2a} e^{-2c} + e^{4a} e^{-4c} + 1)}}\right) \sqrt{e^{2c-2a} (2e^{2a-2c} + e^{4a-4c} + 1)}}{\sqrt{b^2}} - \frac{e^{a+bx} (e^{2a-2c} - 1)}{b (e^{2a-2c} + e^{2a+2bx})}$$

input `int(cosh(a + b*x)/cosh(c + b*x)^2,x)`output `(atan((exp(-a)*exp(2*c)*exp(b*x)*((b^2)^(1/2) + exp(2*a)*exp(-2*c)*(b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c)*(2*exp(2*a)*exp(-2*c) + exp(4*a)*exp(-4*c) + 1))^(1/2)))*(exp(2*c - 2*a)*(2*exp(2*a - 2*c) + exp(4*a - 4*c) + 1))^(1/2))/((b^2)^(1/2) - (exp(a + b*x)*(exp(2*a - 2*c) - 1))/(b*(exp(2*a - 2*c) + exp(2*a + 2*b*x))))`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.31

$$\int \cosh(a + bx) \operatorname{sech}^2(c + bx) dx$$

$$= \frac{e^{2bx+2a+2c} \operatorname{atan}(e^{bx+c}) + e^{2bx+4c} \operatorname{atan}(e^{bx+c}) + e^{2a} \operatorname{atan}(e^{bx+c}) + e^{2c} \operatorname{atan}(e^{bx+c}) - e^{bx+2a+c} + e^{bx+3c}}{e^{a+c} b (e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)*sech(b*x+c)^2,x)`output `(e**(2*a + 2*b*x + 2*c)*atan(e**(b*x + c)) + e**(2*b*x + 4*c)*atan(e**(b*x + c)) + e**(2*a)*atan(e**(b*x + c)) + e**(2*c)*atan(e**(b*x + c)) - e**(2*a + b*x + c) + e**(b*x + 3*c))/(e**(a + c)*b*(e**(2*b*x + 2*c) + 1))`

3.129 $\int \cosh(a + bx)\operatorname{sech}^3(c + bx) dx$

Optimal result	909
Mathematica [A] (verified)	909
Rubi [A] (verified)	910
Maple [A] (verified)	912
Fricas [B] (verification not implemented)	912
Sympy [F]	913
Maxima [B] (verification not implemented)	913
Giac [A] (verification not implemented)	914
Mupad [F(-1)]	914
Reduce [B] (verification not implemented)	914

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \cosh(a + bx)\operatorname{sech}^3(c + bx) dx = -\frac{\operatorname{sech}^2(c + bx)\sinh(a - c)}{2b} + \frac{\cosh(a - c)\tanh(c + bx)}{b}$$

output

```
-1/2*sech(b*x+c)^2*sinh(a-c)/b+cosh(a-c)*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

$$\int \cosh(a + bx)\operatorname{sech}^3(c + bx) dx = -\frac{\operatorname{sech}(c)\operatorname{sech}^2(c + bx)(\sinh(a) - \cosh(a - c)\sinh(c + 2bx))}{2b}$$

input

```
Integrate[Cosh[a + b*x]*Sech[c + b*x]^3,x]
```

output

```
-1/2*(Sech[c]*Sech[c + b*x]^2*(Sinh[a] - Cosh[a - c]*Sinh[c + 2*b*x]))/b
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {6161, 3042, 26, 3086, 15, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{sech}^3(bx + c) dx \\
 & \quad \downarrow \text{6161} \\
 & \cosh(a - c) \int \operatorname{sech}^2(c + bx) dx + \sinh(a - c) \int \operatorname{sech}^2(c + bx) \tanh(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx + \sinh(a - c) \int -i \sec(ic + ibx)^2 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{26} \\
 & \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx - i \sinh(a - c) \int \sec(ic + ibx)^2 \tan(ic + ibx) dx \\
 & \quad \downarrow \text{3086} \\
 & -\frac{\sinh(a - c) \int \operatorname{sech}(c + bx) d\operatorname{sech}(c + bx)}{b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{\sinh(a - c) \operatorname{sech}^2(bx + c)}{2b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\sinh(a - c) \operatorname{sech}^2(bx + c)}{2b} + \frac{i \cosh(a - c) \int 1 d(-i \tanh(c + bx))}{b} \\
 & \quad \downarrow \text{24} \\
 & \frac{\cosh(a - c) \tanh(bx + c)}{b} - \frac{\sinh(a - c) \operatorname{sech}^2(bx + c)}{2b}
 \end{aligned}$$

input `Int[Cosh[a + b*x]*Sech[c + b*x]^3,x]`

output $-1/2*(\text{Sech}[c + b*x]^2*\text{Sinh}[a - c])/b + (\text{Cosh}[a - c]*\text{Tanh}[c + b*x])/b$

Defintions of rubi rules used

rule 15 $\text{Int}[(a_.)*(x_)^(m_.), x_Symbol] \rightarrow \text{Simp}[a*(x^(m + 1))/(m + 1), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

rule 26 $\text{Int}[(\text{Complex}[0, a_])*(Fx_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \ \text{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3086 $\text{Int}(((a_.)*\text{sec}[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol) \rightarrow \text{Simp}[a/f \ \text{Subst}[\text{Int}[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2)], x], x, \text{Sec}[e + f*x], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n - 1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n + 1])$

rule 4254 $\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[-d^(-1) \ \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^(n/2 - 1), x], x], x, \text{Cot}[c + d*x], x] \text{ ; FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

rule 6161 $\text{Int}[\text{Cosh}[v_] * \text{Sech}[w_]^(n_.), x_Symbol] \rightarrow \text{Simp}[\text{Sinh}[v - w] \ \text{Int}[\text{Tanh}[w] * \text{Sech}[w]^(n - 1), x], x] + \text{Simp}[\text{Cosh}[v - w] \ \text{Int}[\text{Sech}[w]^(n - 1), x], x] \text{ ; GtQ}[n, 0] \ \&\& \ \text{NeQ}[w, v] \ \&\& \ \text{FreeQ}[v - w, x]$

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.68

method	result	size
parallelrisc	$\frac{\sinh(2bx+a+c)}{b(\cosh(2bx+2c)+1)}$	26
risch	$-\frac{(2e^{2bx+2a+2c}+e^{2a}+e^{2c})e^{3a-c}}{(e^{2bx+2a+2c}+e^{2a})^2b}$	56

input `int(cosh(b*x+a)*sech(b*x+c)^3,x,method=_RETURNVERBOSE)`

output `1/b*sinh(2*b*x+a+c)/(cosh(2*b*x+2*c)+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(36) = 72.

Time = 0.08 (sec) , antiderivative size = 248, normalized size of antiderivative = 6.53

$$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx =$$

$$-\frac{b \cosh(bx + c)^3 \cosh(-a + c)^2 + 3b \cosh(bx + c) \cosh(-a + c)^2 + (b \cosh(-a + c)^2 - b \sinh(-a + c) \cosh(bx + c)) \sinh(bx + c)}{(b \cosh(bx + c) \cosh(-a + c) + \sinh(bx + c))^3}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^3,x, algorithm="fricas")`

output `-2*(2*cosh(b*x + c)*cosh(-a + c) - cosh(b*x + c)*sinh(-a + c) - sinh(b*x + c)*sinh(-a + c))/(b*cosh(b*x + c)^3*cosh(-a + c)^2 + 3*b*cosh(b*x + c)*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^3 + 3*(b*cosh(b*x + c)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^2 - (b*cosh(b*x + c)^3 + 3*b*cosh(b*x + c))*sinh(-a + c)^2 + (3*b*cosh(b*x + c)^2*cosh(-a + c)^2 + b*cosh(-a + c)^2 - (3*b*cosh(b*x + c)^2 + b)*sinh(-a + c)^2)*sinh(b*x + c)`

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx = \int \cosh(a + bx) \operatorname{sech}^3(bx + c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x+c)**3,x)`

output `Integral(cosh(a + b*x)*sech(b*x + c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.13

$$\begin{aligned} \int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx &= \frac{2e^{(-2bx+3c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})} \\ &+ \frac{e^{(2a+3c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})} \\ &+ \frac{e^{(5c)}}{b(2e^{(-2bx+a+2c)} + e^{(-4bx+a)} + e^{(a+4c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^3,x, algorithm="maxima")`

output `2*e^(-2*b*x + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c))) + e^(2*a + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c))) + e^(5*c)/(b*(2*e^(-2*b*x + a + 2*c) + e^(-4*b*x + a) + e^(a + 4*c)))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx = -\frac{(2e^{(2bx+2a+2c)} + e^{(2a)} + e^{(2c)})e^{(-a-c)}}{b(e^{(2bx+2c)} + 1)^2}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^3,x, algorithm="giac")`output `-(2*e^(2*b*x + 2*a + 2*c) + e^(2*a) + e^(2*c))*e^(-a - c)/(b*(e^(2*b*x + 2*c) + 1)^2)`**Mupad [F(-1)]**

Timed out.

$$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx = \int \frac{\cosh(a + bx)}{\cosh(c + bx)^3} dx$$

input `int(cosh(a + b*x)/cosh(c + b*x)^3,x)`output `int(cosh(a + b*x)/cosh(c + b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int \cosh(a + bx) \operatorname{sech}^3(c + bx) dx = \frac{e^c(e^{4bx+2a+2c} - 1)}{e^{ab}(e^{4bx+4c} + 2e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)*sech(b*x+c)^3,x)`output `(e**c*(e**(2*a + 4*b*x + 2*c) - 1))/(e**a*b*(e**(4*b*x + 4*c) + 2*e**(2*b*x + 2*c) + 1))`

3.130 $\int \cosh(a + bx)\operatorname{sech}^4(c + bx) dx$

Optimal result	915
Mathematica [A] (verified)	915
Rubi [A] (verified)	916
Maple [C] (verified)	918
Fricas [B] (verification not implemented)	919
Sympy [F]	920
Maxima [B] (verification not implemented)	920
Giac [B] (verification not implemented)	921
Mupad [F(-1)]	921
Reduce [B] (verification not implemented)	922

Optimal result

Integrand size = 15, antiderivative size = 67

$$\int \cosh(a + bx)\operatorname{sech}^4(c + bx) dx = \frac{\arctan(\sinh(c + bx)) \cosh(a - c)}{2b} - \frac{\operatorname{sech}^3(c + bx) \sinh(a - c)}{3b} + \frac{\cosh(a - c)\operatorname{sech}(c + bx) \tanh(c + bx)}{2b}$$

output `1/2*arctan(sinh(b*x+c))*cosh(a-c)/b-1/3*sech(b*x+c)^3*sinh(a-c)/b+1/2*cosh(a-c)*sech(b*x+c)*tanh(b*x+c)/b`

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.22

$$\int \cosh(a + bx)\operatorname{sech}^4(c + bx) dx = \frac{-\frac{1}{2}\operatorname{sech}(c)\operatorname{sech}^3(c + bx)(7 \sinh(a) + \sinh(a - 2c) - 6 \cosh(a - c) \sinh(c + 2bx)) + 6 \cosh(a - c) (2 \arctan(\sinh(c + bx)))}{12b}$$

input `Integrate[Cosh[a + b*x]*Sech[c + b*x]^4,x]`

output

```
(-1/2*(Sech[c]*Sech[c + b*x]^3*(7*Sinh[a] + Sinh[a - 2*c] - 6*Cosh[a - c]*
Sinh[c + 2*b*x])) + 6*Cosh[a - c]*(2*ArcTan[Sinh[c] + Cosh[c]*Tanh[(b*x)/2
]] + Sech[c + b*x]*Tanh[c]))/(12*b)
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.94, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {6161, 3042, 26, 3086, 15, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \cosh(a + bx) \operatorname{sech}^4(bx + c) dx \\
& \quad \downarrow \text{6161} \\
& \cosh(a - c) \int \operatorname{sech}^3(c + bx) dx + \sinh(a - c) \int \operatorname{sech}^3(c + bx) \tanh(c + bx) dx \\
& \quad \downarrow \text{3042} \\
& \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx + \sinh(a - c) \int -i \sec(ic + ibx)^3 \tan(ic + ibx) dx \\
& \quad \downarrow \text{26} \\
& \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx - i \sinh(a - c) \int \sec(ic + ibx)^3 \tan(ic + ibx) dx \\
& \quad \downarrow \text{3086} \\
& -\frac{\sinh(a - c) \int \operatorname{sech}^2(c + bx) d\operatorname{sech}(c + bx)}{b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx \\
& \quad \downarrow \text{15} \\
& -\frac{\sinh(a - c) \operatorname{sech}^3(bx + c)}{3b} + \cosh(a - c) \int \csc\left(ic + ibx + \frac{\pi}{2}\right)^3 dx \\
& \quad \downarrow \text{4255} \\
& \cosh(a - c) \left(\frac{1}{2} \int \operatorname{sech}(c + bx) dx + \frac{\tanh(bx + c) \operatorname{sech}(bx + c)}{2b} \right) - \frac{\sinh(a - c) \operatorname{sech}^3(bx + c)}{3b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$-\frac{\sinh(a-c)\operatorname{sech}^3(bx+c)}{3b} + \cosh(a-c) \left(\frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} + \frac{1}{2} \int \csc\left(ic+ibx+\frac{\pi}{2}\right) dx \right)$$

↓ 4257

$$\cosh(a-c) \left(\frac{\arctan(\sinh(bx+c))}{2b} + \frac{\tanh(bx+c)\operatorname{sech}(bx+c)}{2b} \right) - \frac{\sinh(a-c)\operatorname{sech}^3(bx+c)}{3b}$$

input `Int[Cosh[a + b*x]*Sech[c + b*x]^4,x]`

output `-1/3*(Sech[c + b*x]^3*Sinh[a - c])/b + Cosh[a - c]*(ArcTan[Sinh[c + b*x]]/(2*b) + (Sech[c + b*x]*Tanh[c + b*x])/(2*b))`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4255 `Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6161 `Int[Cosh[v_]*Sech[w_]^(n_.), x_Symbol] := Simp[Sinh[v - w] Int[Tanh[w]*Sech[w]^(n - 1), x], x] + Simp[Cosh[v - w] Int[Sech[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.34 (sec) , antiderivative size = 245, normalized size of antiderivative = 3.66

method	result
risch	$-\frac{e^{bx+a}(-3e^{4bx+6a+4c}-3e^{4bx+4a+6c}+8e^{2bx+6a+2c}-8e^{2bx+4a+4c}+3e^{6a}+3e^{4a+2c})}{6b(e^{2bx+2a+2c}+e^{2a})^3} + \frac{i \ln(e^{bx+a}+ie^{a-c})e^{-a-c}e^{2a}}{4b} + \frac{i \ln(e^{bx+a}-ie^{a-c})e^{-a-c}e^{2a}}{4b}$

input `int(cosh(b*x+a)*sech(b*x+c)^4,x,method=_RETURNVERBOSE)`

output `-1/6*exp(b*x+a)*(-3*exp(4*b*x+6*a+4*c)-3*exp(4*b*x+4*a+6*c)+8*exp(2*b*x+6*a+2*c)-8*exp(2*b*x+4*a+4*c)+3*exp(6*a)+3*exp(4*a+2*c))/b/(exp(2*b*x+2*a+2*c)+exp(2*a))^3+1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*a)+1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-a-c)*exp(2*c)-1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*a)-1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1881 vs. $2(61) = 122$.

Time = 0.11 (sec) , antiderivative size = 1881, normalized size of antiderivative = 28.07

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^4,x, algorithm="fricas")`

output

```
1/6*(3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^5 + 3*(cosh(-a + c)^2 - 2*cosh(-
a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^5 - 15*(2*cosh(b*x
+ c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a
+ c)^2 + 1)*cosh(b*x + c))*sinh(b*x + c)^4 - 8*(cosh(-a + c)^2 - 1)*cosh(b
*x + c)^3 + 2*(15*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (15*cosh(b*x + c)
^2 - 4)*sinh(-a + c)^2 - 4*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^2*cosh(-a
+ c) - 4*cosh(-a + c))*sinh(-a + c) + 4)*sinh(b*x + c)^3 + 6*(5*(cosh(-a +
c)^2 + 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 - 4*cosh(b*x + c))*sinh(-a
+ c)^2 - 4*(cosh(-a + c)^2 - 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh
(-a + c) - 4*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^2 + (
3*cosh(b*x + c)^5 - 8*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 +
3*((cosh(-a + c)^2 + 1)*cosh(b*x + c)^6 + (cosh(-a + c)^2 - 2*cosh(-a + c)
)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*x + c)^6 - 6*(2*cosh(b*x + c)*c
osh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2
+ 1)*cosh(b*x + c))*sinh(b*x + c)^5 + 3*(cosh(-a + c)^2 + 1)*cosh(b*x + c)
^4 + 3*(5*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 + 1)*s
inh(-a + c)^2 + cosh(-a + c)^2 - 2*(5*cosh(b*x + c)^2*cosh(-a + c) + cosh(
-a + c))*sinh(-a + c) + 1)*sinh(b*x + c)^4 + 4*(5*(cosh(-a + c)^2 + 1)*cos
h(b*x + c)^3 + (5*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a + c)^2 + 3*(c
osh(-a + c)^2 + 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh(-a + c) + ...
```


Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx = \int \cosh(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x+c)**4,x)`

output `Integral(cosh(a + b*x)*sech(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(61) = 122$.

Time = 0.12 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.25

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx = -\frac{(e^{2a} + e^{2c}) \arctan(e^{-bx-c}) e^{(-a-c)}}{2b} + \frac{3(e^{2a+4c} + e^{6c})e^{-bx-a} - 8(e^{4a+2c} - e^{2a+4c})e^{-3bx-3a} - 3(e^{6a} + e^{4a+2c})e^{-5bx-5a}}{6b(3e^{-2bx+4c} + 3e^{-4bx+2c} + e^{-6bx} + e^{6c})}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^4,x, algorithm="maxima")`

output `-1/2*(e^(2*a) + e^(2*c))*arctan(e^(-b*x - c))*e^(-a - c)/b + 1/6*(3*(e^(2*a + 4*c) + e^(6*c))*e^(-b*x - a) - 8*(e^(4*a + 2*c) - e^(2*a + 4*c))*e^(-3*b*x - 3*a) - 3*(e^(6*a) + e^(4*a + 2*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 4*c) + 3*e^(-4*b*x + 2*c) + e^(-6*b*x) + e^(6*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(61) = 122$.

Time = 0.14 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.84

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx = \frac{(e^{2a} + e^{2c}) \arctan(e^{(bx+c)}) e^{(-a-c)}}{2b} + \frac{(3e^{(5bx+2a+4c)} + 3e^{(5bx+6c)} - 8e^{(3bx+2a+2c)} + 8e^{(3bx+4c)} - 3e^{(bx+2a)} - 3e^{(bx+2c)})e^{(-a)}}{6b(e^{(2bx+2c)} + 1)^3}$$

input `integrate(cosh(b*x+a)*sech(b*x+c)^4,x, algorithm="giac")`

output `1/2*(e^(2*a) + e^(2*c))*arctan(e^(b*x + c))*e^(-a -c)/b + 1/6*(3*e^(5*b*x + 2*a + 4*c) + 3*e^(5*b*x + 6*c) - 8*e^(3*b*x + 2*a + 2*c) + 8*e^(3*b*x + 4*c) - 3*e^(b*x + 2*a) - 3*e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) + 1)^3)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx = \int \frac{\cosh(a + bx)}{\cosh(c + bx)^4} dx$$

input `int(cosh(a + b*x)/cosh(c + b*x)^4,x)`

output `int(cosh(a + b*x)/cosh(c + b*x)^4, x)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 287, normalized size of antiderivative = 4.28

$$\int \cosh(a + bx) \operatorname{sech}^4(c + bx) dx$$

$$= \frac{3e^{6bx+2a+6c} \operatorname{atan}(e^{bx+c}) + 3e^{6bx+8c} \operatorname{atan}(e^{bx+c}) + 9e^{4bx+2a+4c} \operatorname{atan}(e^{bx+c}) + 9e^{4bx+6c} \operatorname{atan}(e^{bx+c}) + 9e^{2bx+2a+2c} \operatorname{atan}(e^{bx+c})}{(6e^{a+c}b(e^{6bx+6c} + 3e^{4bx+4c} + 3e^{2bx+2c} + 1))}$$

input `int(cosh(b*x+a)*sech(b*x+c)^4,x)`

output

```
(3*e**(2*a + 6*b*x + 6*c)*atan(e**(b*x + c)) + 3*e**(6*b*x + 8*c)*atan(e**
(b*x + c)) + 9*e**(2*a + 4*b*x + 4*c)*atan(e**(b*x + c)) + 9*e**(4*b*x + 6
*c)*atan(e**(b*x + c)) + 9*e**(2*a + 2*b*x + 2*c)*atan(e**(b*x + c)) + 9*e
**(2*b*x + 4*c)*atan(e**(b*x + c)) + 3*e**(2*a)*atan(e**(b*x + c)) + 3*e**
(2*c)*atan(e**(b*x + c)) + 3*e**(2*a + 5*b*x + 5*c) + 3*e**(5*b*x + 7*c) -
8*e**(2*a + 3*b*x + 3*c) + 8*e**(3*b*x + 5*c) - 3*e**(2*a + b*x + c) - 3*
e**(b*x + 3*c))/(6*e**(a + c)*b*(e**(6*b*x + 6*c) + 3*e**(4*b*x + 4*c) + 3
*e**(2*b*x + 2*c) + 1))
```

3.131 $\int \cosh(a + bx)\operatorname{sech}(c - bx) dx$

Optimal result	923
Mathematica [A] (verified)	923
Rubi [F]	924
Maple [B] (verified)	924
Fricas [B] (verification not implemented)	925
Sympy [F]	925
Maxima [B] (verification not implemented)	926
Giac [A] (verification not implemented)	926
Mupad [B] (verification not implemented)	927
Reduce [B] (verification not implemented)	927

Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \cosh(a + bx)\operatorname{sech}(c - bx) dx = x \cosh(a + c) + \frac{\log(\cosh(c - bx)) \sinh(a + c)}{b}$$

output `x*cosh(a+c)+ln(cosh(b*x-c))*sinh(a+c)/b`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \cosh(a + bx)\operatorname{sech}(c - bx) dx = x \cosh(a + c) + \frac{\log(\cosh(c - bx)) \sinh(a + c)}{b}$$

input `Integrate[Cosh[a + b*x]*Sech[c - b*x],x]`

output `x*Cosh[a + c] + (Log[Cosh[c - b*x]]*Sinh[a + c])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx$$

input `Int[Cosh[a + b*x]*Sech[c - b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. $2(24) = 48$.

Time = 0.18 (sec) , antiderivative size = 144, normalized size of antiderivative = 6.26

method	result
risch	$x e^{a+c} - e^{-a-c} e^{2a+2c} x - \frac{e^{-a-c} e^{2a+2c} a}{b} + x e^{-a-c} + \frac{e^{-a-c} a}{b} + \frac{\ln(e^{2a+2c} + e^{2bx+2a}) e^{-a-c} e^{2a+2c}}{2b} - \frac{\ln(e^{2a+2c} + e^{2bx+2a})}{2b}$

input `int(cosh(b*x+a)*sech(b*x-c),x,method=_RETURNVERBOSE)`

output `x*exp(a+c)-exp(-a-c)*exp(2*a+2*c)*x-1/b*exp(-a-c)*exp(2*a+2*c)*a+x*exp(-a-c)+1/b*exp(-a-c)*a+1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)*exp(2*a+2*c)-1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(24) = 48$.

Time = 0.09 (sec) , antiderivative size = 146, normalized size of antiderivative = 6.35

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx$$

$$= \frac{2bx \cosh(a + c)^2 - 4bx \cosh(a + c) \sinh(a + c) + 2bx \sinh(a + c)^2 - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 - 1) \log(2(\cosh(bx + a) \cosh(a + c) - \sinh(bx + a) \sinh(a + c)) / (\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c)))}{2(b \cosh(a + c) - b \sinh(a + c))}$$

input `integrate(cosh(b*x+a)*sech(b*x-c),x, algorithm="fricas")`

output `1/2*(2*b*x*cosh(a + c)^2 - 4*b*x*cosh(a + c)*sinh(a + c) + 2*b*x*sinh(a + c)^2 - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 - 1)*log(2*(cosh(b*x + a)*cosh(a + c) - sinh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c))))/(b*cosh(a + c) - b*sinh(a + c))`

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx = \int \cosh(a + bx) \operatorname{sech}(bx - c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x-c),x)`

output `Integral(cosh(a + b*x)*sech(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(24) = 48$.

Time = 0.05 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.13

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx$$

$$= \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-2bx+2c)} + 1)}{2b} + \frac{(bx + a)e^{(a+c)}}{b}$$

input `integrate(cosh(b*x+a)*sech(b*x-c),x, algorithm="maxima")`

output `1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(-2*b*x + 2*c) + 1)/b + (b*x + a)*e^(a + c)/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.96

$$\int \cosh(a + bx) \operatorname{sech}(c - bx) dx = xe^{(-a-c)} + \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(2bx)} + e^{(2c)})}{2b}$$

input `integrate(cosh(b*x+a)*sech(b*x-c),x, algorithm="giac")`

output `x*e^(-a - c) + 1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(2*b*x) + e^(2*c))/b`

Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.78

$$\int \cosh(a + bx)\operatorname{sech}(c - bx) dx = x e^{-a-c} + \frac{e^{-2a-2c} \ln(e^{2a} e^{2bx} + e^{2a} e^{2c}) (2b e^{3a+3c} - 2b e^{a+c})}{4b^2}$$

input `int(cosh(a + b*x)/cosh(c - b*x),x)`output `x*exp(- a - c) + (exp(- 2*a - 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(2*c))*(2*b*exp(3*a + 3*c) - 2*b*exp(a + c)))/(4*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.39

$$\int \cosh(a + bx)\operatorname{sech}(c - bx) dx = \frac{e^{2a+2c} \log(e^{2bx} + e^{2c}) - \log(e^{2bx} + e^{2c}) + 2bx}{2e^{a+cb}}$$

input `int(cosh(b*x+a)*sech(b*x-c),x)`output `(e**(2*a + 2*c)*log(e**(2*b*x) + e**(2*c)) - log(e**(2*b*x) + e**(2*c)) + 2*b*x)/(2*e**(a + c)*b)`

3.132 $\int \cosh(a + bx)\operatorname{sech}^2(c - bx) dx$

Optimal result	928
Mathematica [B] (verified)	928
Rubi [F]	929
Maple [C] (verified)	929
Fricas [B] (verification not implemented)	930
Sympy [F]	931
Maxima [A] (verification not implemented)	931
Giac [B] (verification not implemented)	931
Mupad [B] (verification not implemented)	932
Reduce [B] (verification not implemented)	932

Optimal result

Integrand size = 16, antiderivative size = 34

$$\int \cosh(a + bx)\operatorname{sech}^2(c - bx) dx = -\frac{\arctan(\sinh(c - bx)) \cosh(a + c)}{b} - \frac{\operatorname{sech}(c - bx) \sinh(a + c)}{b}$$

output

```
arctan(sinh(b*x-c))*cosh(a+c)/b-sech(b*x-c)*sinh(a+c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 144 vs. 2(34) = 68.

Time = 0.07 (sec) , antiderivative size = 144, normalized size of antiderivative = 4.24

$$\int \cosh(a + bx)\operatorname{sech}^2(c - bx) dx$$

$$= \frac{2 \arctan\left(\frac{\cosh\left(\frac{bx}{2}\right) - \cosh^2(c) \cosh\left(\frac{bx}{2}\right) - 2 \cosh(c) \cosh\left(\frac{bx}{2}\right) \sinh(c) - \cosh\left(\frac{bx}{2}\right) \sinh^2(c) + \sinh\left(\frac{bx}{2}\right) + \cosh^2(c) \sinh\left(\frac{bx}{2}\right) + 2 \cosh(c) \sinh\left(\frac{bx}{2}\right)}{2 \cosh(c) \cosh\left(\frac{bx}{2}\right) + 2 \cosh\left(\frac{bx}{2}\right) \sinh(c)}\right)}{b} - \frac{\operatorname{sech}(c - bx) \sinh(a + c)}{b}$$

input `Integrate[Cosh[a + b*x]*Sech[c - b*x]^2,x]`

output $(2*\text{ArcTan}[(\text{Cosh}[(b*x)/2] - \text{Cosh}[c]^2*\text{Cosh}[(b*x)/2] - 2*\text{Cosh}[c]*\text{Cosh}[(b*x)/2]*\text{Sinh}[c] - \text{Cosh}[(b*x)/2]*\text{Sinh}[c]^2 + \text{Sinh}[(b*x)/2] + \text{Cosh}[c]^2*\text{Sinh}[(b*x)/2]) + 2*\text{Cosh}[c]*\text{Sinh}[c]*\text{Sinh}[(b*x)/2] + \text{Sinh}[c]^2*\text{Sinh}[(b*x)/2]) / (2*\text{Cosh}[c]*\text{Cosh}[(b*x)/2] + 2*\text{Cosh}[(b*x)/2]*\text{Sinh}[c]) * \text{Cosh}[a + c] / b - (\text{Sech}[c - b*x]*\text{Sinh}[a + c]) / b$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx$$

input `Int[Cosh[a + b*x]*Sech[c - b*x]^2,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 175, normalized size of antiderivative = 5.15

method	result
risch	$-\frac{e^{bx+a}(e^{2a+2c}-1)}{b(e^{2a+2c}+e^{2bx+2a})} + \frac{i \ln(e^{bx+a}+ie^{a+c})e^{-a-c}e^{2a+2c}}{2b} + \frac{i \ln(e^{bx+a}+ie^{a+c})e^{-a-c}}{2b} - \frac{i \ln(e^{bx+a}-ie^{a+c})e^{-a-c}e^{2a+2c}}{2b} -$

input `int(cosh(b*x+a)*sech(b*x-c)^2,x,method=_RETURNVERBOSE)`

output

```
-1/b*exp(b*x+a)*(exp(2*a+2*c)-1)/(exp(2*a+2*c)+exp(2*b*x+2*a))+1/2*I*ln(ex
p(b*x+a)+I*exp(a+c))/b*exp(-a-c)*exp(2*a+2*c)+1/2*I*ln(exp(b*x+a)+I*exp(a+
c))/b*exp(-a-c)-1/2*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)*exp(2*a+2*c)-1
/2*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 713 vs. $2(35) = 70$.

Time = 0.10 (sec) , antiderivative size = 713, normalized size of antiderivative = 20.97

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*sech(b*x-c)^2,x, algorithm="fricas")
```

output

```
(3*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - cosh(b*x + a)*sinh(a + c)^3 -
(3*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c) + (4*cosh(b*x + a)^2*cosh
(a + c)*sinh(a + c)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 + c
osh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)
^3 + sinh(a + c)^4 + (6*cosh(a + c)^2 + 1)*sinh(a + c)^2 + cosh(a + c)^2 -
2*(2*cosh(a + c)^3 + cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*cosh
(a + c)^2 + 1)*cosh(b*x + a)^2 + 1)*sinh(a + c)^2 - cosh(a + c)^2 + 2*(4*c
osh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - (6*
cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3 + cosh
(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 + cosh(a + c)^2)*cosh(
b*x + a)*sinh(b*x + a) + 2*((2*cosh(a + c)^3 + cosh(a + c))*cosh(b*x + a)
^2 + cosh(a + c))*sinh(a + c) - 1)*arctan(-cosh(b*x + a)*cosh(a + c) - (co
sh(a + c) - sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c)) + (cos
h(a + c)^3 - cosh(a + c))*cosh(b*x + a) + (cosh(a + c)^3 + 3*cosh(a + c)*s
inh(a + c)^2 - sinh(a + c)^3 - (3*cosh(a + c)^2 - 1)*sinh(a + c) - cosh(a
+ c))*sinh(b*x + a)/(b*cosh(b*x + a)^2*cosh(a + c)^3 + 3*b*cosh(b*x + a)^
2*cosh(a + c)*sinh(a + c)^2 - b*cosh(b*x + a)^2*sinh(a + c)^3 + (b*cosh(a
+ c)^3 - 3*b*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(a + c)*sinh(a + c)^2 - b
*sinh(a + c)^3)*sinh(b*x + a)^2 + b*cosh(a + c) + 2*(b*cosh(b*x + a)*cosh(
a + c)^3 - 3*b*cosh(b*x + a)*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(b*x + ...
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx = \int \cosh(a + bx) \operatorname{sech}^2(bx - c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x-c)**2,x)`

output `Integral(cosh(a + b*x)*sech(b*x - c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.03

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx = -\frac{(e^{(2a+2c)} + 1) \arctan(e^{(-bx+c)}) e^{(-a-c)}}{b} - \frac{(e^{(2a+2c)} - 1) e^{(-bx-a)}}{b(e^{(-2bx+2c)} + 1)}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^2,x, algorithm="maxima")`

output `-(e^(2*a + 2*c) + 1)*arctan(e^(-b*x + c))*e^(-a - c)/b - (e^(2*a + 2*c) - 1)*e^(-b*x - a)/(b*(e^(-2*b*x + 2*c) + 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(35) = 70.

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.09

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx = \frac{(e^{(2a+2c)} + 1) \arctan(e^{(bx-c)}) e^{(-a-c)}}{b} - \frac{(e^{(bx+2a+2c)} - e^{(bx)}) e^{(-a)}}{b(e^{(2bx)} + e^{(2c)})}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^2,x, algorithm="giac")`

output $(e^{(2a + 2c)} + 1) \arctan(e^{(bx - c)}) e^{(-a - c)/b} - (e^{(bx + 2a + 2c)} - e^{(bx)}) e^{(-a)/(b(e^{(2bx)} + e^{(2c)}))}$

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 148, normalized size of antiderivative = 4.35

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx$$

$$= \frac{\operatorname{atan}\left(\frac{e^{-a} e^{-2c} e^{bx} (\sqrt{b^2 + e^{2a} e^{2c} \sqrt{b^2}})}{b \sqrt{e^{-2a} e^{-2c} (2e^{2a} e^{2c} + e^{4a} e^{4c} + 1)}}\right) \sqrt{e^{-2a-2c} (2e^{2a+2c} + e^{4a+4c} + 1)}}{\sqrt{b^2}} - \frac{e^{a+bx} (e^{2a+2c} - 1)}{b (e^{2a+2c} + e^{2a+2bx})}$$

input `int(cosh(a + b*x)/cosh(c - b*x)^2,x)`

output $(\operatorname{atan}((\exp(-a) \exp(-2c) \exp(bx) * ((b^2)^{(1/2)} + \exp(2a) \exp(2c) * (b^2)^{(1/2)})) / (b * (\exp(-2a) \exp(-2c) * (2 \exp(2a) \exp(2c) + \exp(4a) \exp(4c) + 1))^{(1/2)})) * (\exp(-2a - 2c) * (2 \exp(2a) \exp(2c) + \exp(4a) \exp(4c) + 1))^{(1/2)}) / (b^2)^{(1/2)} - (\exp(a + bx) * (\exp(2a) \exp(2c) - 1)) / (b * (\exp(2a) \exp(2c) + \exp(2a + 2bx))))$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 132, normalized size of antiderivative = 3.88

$$\int \cosh(a + bx) \operatorname{sech}^2(c - bx) dx$$

$$= \frac{e^{2bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{bx+2a+3c} + e^{bx+c}}{e^{a+cb} (e^{2bx} + e^{2c})}$$

input `int(cosh(b*x+a)*sech(b*x-c)^2,x)`

output

```
(e**(2*a + 2*b*x + 2*c)*atan(e**(b*x)/e**c) + e**(2*b*x)*atan(e**(b*x)/e**c) + e**(2*a + 4*c)*atan(e**(b*x)/e**c) + e**(2*c)*atan(e**(b*x)/e**c) - e**(2*a + b*x + 3*c) + e**(b*x + c))/(e**(a + c)*b*(e**(2*b*x) + e**(2*c)))
```

3.133 $\int \cosh(a + bx)\operatorname{sech}^3(c - bx) dx$

Optimal result	934
Mathematica [A] (verified)	934
Rubi [F]	935
Maple [A] (verified)	935
Fricas [B] (verification not implemented)	936
Sympy [F]	936
Maxima [B] (verification not implemented)	937
Giac [A] (verification not implemented)	937
Mupad [F(-1)]	938
Reduce [B] (verification not implemented)	938

Optimal result

Integrand size = 16, antiderivative size = 37

$$\int \cosh(a + bx)\operatorname{sech}^3(c - bx) dx = -\frac{\operatorname{sech}^2(c - bx)\sinh(a + c)}{2b} - \frac{\cosh(a + c)\tanh(c - bx)}{b}$$

output

```
-1/2*sech(b*x-c)^2*sinh(a+c)/b+cosh(a+c)*tanh(b*x-c)/b
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\begin{aligned} & \int \cosh(a + bx)\operatorname{sech}^3(c - bx) dx \\ &= -\frac{\operatorname{sech}(c)\operatorname{sech}^2(c - bx)(\sinh(a) + \cosh(a + c)\sinh(c - 2bx))}{2b} \end{aligned}$$

input

```
Integrate[Cosh[a + b*x]*Sech[c - b*x]^3,x]
```

output

```
-1/2*(Sech[c]*Sech[c - b*x]^2*(Sinh[a] + Cosh[a + c]*Sinh[c - 2*b*x]))/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx$$

input `Int[Cosh[a + b*x]*Sech[c - b*x]^3,x]`

output `$Aborted`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

method	result	size
parallelrisch	$\frac{\sinh(2bx+a-c)}{b(1+\cosh(2bx-2c))}$	28
risch	$-\frac{(e^{2a+2c}+2e^{2bx+2a}+1)e^{3a+3c}}{(e^{2a+2c}+e^{2bx+2a})^2b}$	55

input `int(cosh(b*x+a)*sech(b*x-c)^3,x,method=_RETURNVERBOSE)`

output `1/b*sinh(2*b*x+a-c)/(1+cosh(2*b*x-2*c))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 454 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 454, normalized size of antiderivative = 12.27

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx =$$

$$\frac{-b \cosh(bx + a)^3 \cosh(a + c)^3 + 3b \cosh(bx + a) \cosh(a + c)^3 + (b \cosh(a + c)^3 - 3b \cosh(a + c)^2 \sinh(a + c)) \sinh(bx + a)}{\cosh(a + c)^3 + 3b \cosh(bx + a) \cosh(a + c)^2 \sinh(a + c) + 3b^2 \cosh(bx + a)^2 \cosh(a + c) \sinh(a + c) - b^3 \cosh(bx + a)^3 \sinh(a + c)}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^3,x, algorithm="fricas")`

output

$$\frac{-2*(2*\cosh(b*x + a)*\cosh(a + c)^2 + \cosh(b*x + a)*\cosh(a + c)*\sinh(a + c) - \cosh(b*x + a)*\sinh(a + c)^2 - (\cosh(a + c)*\sinh(a + c) + \sinh(a + c)^2)*\sinh(b*x + a))/(b*\cosh(b*x + a)^3*\cosh(a + c)^3 + 3*b*\cosh(b*x + a)*\cosh(a + c)^3 + (b*\cosh(a + c)^3 - 3*b*\cosh(a + c)^2*\sinh(a + c) + 3*b*\cosh(a + c)*\sinh(a + c)^2 - b*\sinh(a + c)^3)*\sinh(b*x + a)^3 - (b*\cosh(b*x + a)^3 - b*\cosh(b*x + a)*\sinh(a + c)^3 + 3*(b*\cosh(b*x + a)*\cosh(a + c)^3 - 3*b*\cosh(b*x + a)*\cosh(a + c)^2*\sinh(a + c) + 3*b*\cosh(b*x + a)*\cosh(a + c)*\sinh(a + c)^2 - b*\cosh(b*x + a)*\sinh(a + c)^3)*\sinh(b*x + a)^2 + 3*(b*\cosh(b*x + a)^3*\cosh(a + c) - b*\cosh(b*x + a)*\cosh(a + c))*\sinh(a + c)^2 + (3*b*\cosh(b*x + a)^2*\cosh(a + c)^3 + b*\cosh(a + c)^3 - 3*(b*\cosh(b*x + a)^2 - b)*\sinh(a + c)^3 + (9*b*\cosh(b*x + a)^2*\cosh(a + c) - b*\cosh(a + c))*\sinh(a + c)^2 - 3*(3*b*\cosh(b*x + a)^2*\cosh(a + c)^2 + b*\cosh(a + c)^2)*\sinh(a + c))*\sinh(b*x + a) - (3*b*\cosh(b*x + a)^3*\cosh(a + c)^2 + b*\cosh(b*x + a)*\cosh(a + c)^2)*\sinh(a + c)}$$
Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx = \int \cosh(a + bx) \operatorname{sech}^3(bx - c) dx$$

input `integrate(cosh(b*x+a)*sech(b*x-c)**3,x)`

output `Integral(cosh(a + b*x)*sech(b*x - c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(36) = 72$.

Time = 0.05 (sec) , antiderivative size = 118, normalized size of antiderivative = 3.19

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx = \frac{2e^{(-2bx+2c)}}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})} + \frac{e^{(2a+2c)}}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})} + \frac{1}{b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^3,x, algorithm="maxima")`

output `2*e^(-2*b*x + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c))) + e^(2*a + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c))) + 1/(b*(2*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c)))`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.30

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx = -\frac{(2e^{(2bx+2a+3c)} + e^{(2a+5c)} + e^{(3c)})e^{(-a)}}{b(e^{(2bx)} + e^{(2c)})^2}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^3,x, algorithm="giac")`

output `-(2*e^(2*b*x + 2*a + 3*c) + e^(2*a + 5*c) + e^(3*c))*e^(-a)/(b*(e^(2*b*x) + e^(2*c))^2)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx = \int \frac{\cosh(a + bx)}{\cosh(c - bx)^3} dx$$

input `int(cosh(a + b*x)/cosh(c - b*x)^3,x)`output `int(cosh(a + b*x)/cosh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.51

$$\int \cosh(a + bx) \operatorname{sech}^3(c - bx) dx = \frac{e^c (e^{4bx+2a} - e^{2c})}{e^a b (e^{4bx} + 2e^{2bx+2c} + e^{4c})}$$

input `int(cosh(b*x+a)*sech(b*x-c)^3,x)`output `(e**c*(e**(2*a + 4*b*x) - e**(2*c)))/(e**a*b*(e**(4*b*x) + 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.134 $\int \cosh(a + bx)\operatorname{sech}^4(c - bx) dx$

Optimal result	939
Mathematica [A] (verified)	939
Rubi [F]	940
Maple [C] (verified)	940
Fricas [B] (verification not implemented)	941
Sympy [F]	941
Maxima [B] (verification not implemented)	942
Giac [A] (verification not implemented)	942
Mupad [F(-1)]	943
Reduce [B] (verification not implemented)	943

Optimal result

Integrand size = 16, antiderivative size = 65

$$\int \cosh(a + bx)\operatorname{sech}^4(c - bx) dx = -\frac{\arctan(\sinh(c - bx)) \cosh(a + c)}{2b} - \frac{\operatorname{sech}^3(c - bx) \sinh(a + c)}{3b} - \frac{\cosh(a + c)\operatorname{sech}(c - bx) \tanh(c - bx)}{2b}$$

output `1/2*arctan(sinh(b*x-c))*cosh(a+c)/b-1/3*sech(b*x-c)^3*sinh(a+c)/b+1/2*cosh(a+c)*sech(b*x-c)*tanh(b*x-c)/b`

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.14

$$\int \cosh(a + bx)\operatorname{sech}^4(c - bx) dx = \frac{2\operatorname{sech}^3(c - bx) \sinh(a + c) + 3 \cosh(a + c) (2 \arctan(\sinh(c - bx)) - \cosh(c - bx) \tanh(\frac{bx}{2})) - \operatorname{sech}(c - bx)\operatorname{sech}^2(c - bx)}{6b}$$

input `Integrate[Cosh[a + b*x]*Sech[c - b*x]^4,x]`

output

```
-1/6*(2*Sech[c - b*x]^3*Sinh[a + c] + 3*Cosh[a + c]*(2*ArcTan[Sinh[c] - Co
sh[c]*Tanh[(b*x)/2]] - Sech[c]*Sech[c - b*x]^2*Sinh[b*x] + Sech[c - b*x]*T
anh[c]))/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx$$

input

```
Int[Cosh[a + b*x]*Sech[c - b*x]^4,x]
```

output

```
$Aborted
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.20 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.68

method	result
risch	$-\frac{e^{bx+a}(3e^{6a+6c}+8e^{2bx+6a+4c}+3e^{4a+4c}-3e^{4bx+6a+2c}-8e^{2bx+4a+2c}-3e^{4bx+4a})}{6b(e^{2a+2c}+e^{2bx+2a})^3} + \frac{i \ln(e^{bx+a}+ie^{a+c})e^{-a-c}e^{2a+2c}}{4b} + \frac{i \ln(e^{bx+a}-ie^{a+c})e^{-a-c}e^{2a+2c}}{4b}$

input

```
int(cosh(b*x+a)*sech(b*x-c)^4,x,method=_RETURNVERBOSE)
```

output

```
-1/6*exp(b*x+a)*(3*exp(6*a+6*c)+8*exp(2*b*x+6*a+4*c)+3*exp(4*a+4*c)-3*exp(
4*b*x+6*a+2*c)-8*exp(2*b*x+4*a+2*c)-3*exp(4*b*x+4*a))/b/(exp(2*a+2*c)+exp(
2*b*x+2*a))^3+1/4*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-a-c)*exp(2*a+2*c)+1/4
*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-a-c)-1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b
*exp(-a-c)*exp(2*a+2*c)-1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6243 vs. $2(63) = 126$.

Time = 0.16 (sec) , antiderivative size = 6243, normalized size of antiderivative = 96.05

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*sech(b*x-c)^4,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx = \int \cosh(a + bx) \operatorname{sech}^4(bx - c) dx$$

input

```
integrate(cosh(b*x+a)*sech(b*x-c)**4,x)
```

output

```
Integral(cosh(a + b*x)*sech(b*x - c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(63) = 126$.

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.34

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx = -\frac{(e^{(2a+2c)} + 1) \arctan(e^{(-bx+c)}) e^{(-a-c)}}{2b} + \frac{3(e^{(2a+2c)} + 1)e^{(-bx-a)} - 8(e^{(4a+4c)} - e^{(2a+2c)})e^{(-3bx-3a)} - 3(e^{(6a+6c)} + e^{(4a+4c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+2c)} + 3e^{(-4bx+4c)} + e^{(-6bx+6c)} + 1)}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^4,x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) + 1)*arctan(e^(-b*x + c))*e^(-a - c)/b + 1/6*(3*(e^(2*a + 2*c) + 1)*e^(-b*x - a) - 8*(e^(4*a + 4*c) - e^(2*a + 2*c))*e^(-3*b*x - 3*a) - 3*(e^(6*a + 6*c) + e^(4*a + 4*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 2*c) + 3*e^(-4*b*x + 4*c) + e^(-6*b*x + 6*c) + 1))`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.91

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx = \frac{(e^{(2a+2c)} + 1) \arctan(e^{(bx-c)}) e^{(-a-c)}}{2b} + \frac{(3e^{(5bx)} + 3e^{(5bx+2a+2c)} - 8e^{(3bx+2a+4c)} + 8e^{(3bx+2c)} - 3e^{(bx+2a+6c)} - 3e^{(bx+4c)})e^{(-a)}}{6b(e^{(2bx)} + e^{(2c)})^3}$$

input `integrate(cosh(b*x+a)*sech(b*x-c)^4,x, algorithm="giac")`

output `1/2*(e^(2*a + 2*c) + 1)*arctan(e^(b*x - c))*e^(-a - c)/b + 1/6*(3*e^(5*b*x) + 3*e^(5*b*x + 2*a + 2*c) - 8*e^(3*b*x + 2*a + 4*c) + 8*e^(3*b*x + 2*c) - 3*e^(b*x + 2*a + 6*c) - 3*e^(b*x + 4*c))*e^(-a)/(b*(e^(2*b*x) + e^(2*c))^3)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx = \int \frac{\cosh(a + bx)}{\cosh(c - bx)^4} dx$$

input `int(cosh(a + b*x)/cosh(c - b*x)^4, x)`output `int(cosh(a + b*x)/cosh(c - b*x)^4, x)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 319, normalized size of antiderivative = 4.91

$$\int \cosh(a + bx) \operatorname{sech}^4(c - bx) dx$$

$$= \frac{3e^{6bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 3e^{6bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 9e^{4bx+2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 9e^{4bx+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 9e^{2bx+2a+6c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right)}{6e^a}$$

input `int(cosh(b*x+a)*sech(b*x-c)^4, x)`output `(3***e**(2*a + 6*b*x + 2*c)*atan(e**(b*x)/e**c) + 3***e**(6*b*x)*atan(e**(b*x)/e**c) + 9***e**(2*a + 4*b*x + 4*c)*atan(e**(b*x)/e**c) + 9***e**(4*b*x + 2*c)*atan(e**(b*x)/e**c) + 9***e**(2*a + 2*b*x + 6*c)*atan(e**(b*x)/e**c) + 9***e**(2*b*x + 4*c)*atan(e**(b*x)/e**c) + 3***e**(2*a + 8*c)*atan(e**(b*x)/e**c) + 3***e**(6*c)*atan(e**(b*x)/e**c) + 3***e**(2*a + 5*b*x + 3*c) + 3***e**(5*b*x + c) - 8***e**(2*a + 3*b*x + 5*c) + 8***e**(3*b*x + 3*c) - 3***e**(2*a + b*x + 7*c) - 3***e**(b*x + 5*c))/(6***e**(a + c)*b*(e**(6*b*x) + 3***e**(4*b*x + 2*c) + 3***e**(2*b*x + 4*c) + e**(6*c)))`

3.135 $\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx$

Optimal result	944
Mathematica [A] (verified)	944
Rubi [F]	945
Maple [C] (verified)	945
Fricas [B] (verification not implemented)	946
Sympy [F]	946
Maxima [B] (verification not implemented)	947
Giac [A] (verification not implemented)	947
Mupad [B] (verification not implemented)	948
Reduce [B] (verification not implemented)	948

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = -\frac{\arctan(\sinh(c + bx)) \sinh^2(a - c)}{b} + \frac{\sinh(2a - c + bx)}{b}$$

output

```
-arctan(sinh(b*x+c))*sinh(a-c)^2/b+sinh(b*x+2*a-c)/b
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.33

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx$$

$$= \frac{\arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) - \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) \cosh(2(a - c)) + \sinh(2a - c + bx)}{b}$$

input

```
Integrate[Cosh[a + b*x]^2*Sech[c + b*x],x]
```

output

```
(ArcTan[Tanh[(c + b*x)/2]] - ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(a - c)] + Sinh[2*a - c + b*x])/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c + b*x],x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 251, normalized size of antiderivative = 6.97

method	result
risch	$\frac{e^{bx+2a-c}}{2b} - \frac{e^{-bx-2a+c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a} e^{4a}}{4b} - \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a} e^{2a} e^{2c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a-c}) e^{-2c-2a}}{4b}$

input `int(cosh(b*x+a)^2*sech(b*x+c),x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+2*a-c)-1/2/b*exp(-b*x-2*a+c)+1/4*I*ln(exp(b*x+a)-I*exp(a-c))
/b*exp(-2*c-2*a)*exp(a)^4-1/2*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*
exp(a)^2*exp(c)^2+1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(c)^4
-1/4*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(a)^4+1/2*I*ln(exp(b*x
+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(a)^2*exp(c)^2-1/4*I*ln(exp(b*x+a)+I*ex
p(a-c))/b*exp(-2*c-2*a)*exp(c)^4`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 634 vs. $2(36) = 72$.

Time = 0.11 (sec) , antiderivative size = 634, normalized size of antiderivative = 17.61

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c),x, algorithm="fricas")`

output

```
1/2*(cosh(b*x + c)^2*cosh(-a + c)^4 - 4*cosh(b*x + c)^2*cosh(-a + c)^3*sin
h(-a + c) + 6*cosh(b*x + c)^2*cosh(-a + c)^2*sinh(-a + c)^2 - 4*cosh(b*x +
c)^2*cosh(-a + c)*sinh(-a + c)^3 + cosh(b*x + c)^2*sinh(-a + c)^4 + (cosh
(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a + c)^2*sinh(-a + c)
^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4)*sinh(b*x + c)^2 + (4*
cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 -
2*(3*cosh(-a + c)^2 - 1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a + c)^3
- cosh(-a + c))*cosh(b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 - 2*cosh(-a
+ c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-a + c)^
3 + sinh(-a + c)^4 + 2*(3*cosh(-a + c)^2 - 1)*sinh(-a + c)^2 - 2*cosh(-a +
c)^2 - 4*(cosh(-a + c)^3 - cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c))
*arctan(cosh(b*x + c) + sinh(b*x + c)) + 2*(cosh(b*x + c)*cosh(-a + c)^4 -
4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(b*x + c)*cosh(-a + c
)^2*sinh(-a + c)^2 - 4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 + cosh(b*
x + c)*sinh(-a + c)^4)*sinh(b*x + c) - 1)/(b*cosh(b*x + c)*cosh(-a + c)^2
- 2*b*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a +
c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^
2)*sinh(b*x + c))
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = \int \cosh^2(a + bx) \operatorname{sech}(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x+c),x)`

output `Integral(cosh(a + b*x)**2*sech(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(36) = 72$.

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.06

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = \frac{(e^{4a} - 2e^{(2a+2c)} + e^{4c}) \arctan(e^{-bx-c}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a-c)}}{2b} - \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c),x, algorithm="maxima")`

output `1/2*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))*arctan(e^(-b*x - c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a - c)/b - 1/2*e^(-b*x - 2*a + c)/b`

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.97

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = -\frac{(e^{4a} - 2e^{(2a+2c)} + e^{4c}) \arctan(e^{(bx+c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a-c)}}{2b} - \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c),x, algorithm="giac")`

output `-1/2*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))*arctan(e^(b*x + c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a - c)/b - 1/2*e^(-b*x - 2*a + c)/b`

Mupad [B] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 194, normalized size of antiderivative = 5.39

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = \frac{e^{2a-c+bx}}{2b} - \frac{e^{c-2a-bx}}{2b} - \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{3c} e^{bx} (\sqrt{b^2-2} e^{2a} e^{-2c} \sqrt{b^2} + e^{4a} e^{-4c} \sqrt{b^2})}{b \sqrt{e^{-4a} e^{4c} (6e^{4a} e^{-4c} - 4e^{2a} e^{-2c} - 4e^{6a} e^{-6c} + e^{8a} e^{-8c} + 1)}}}\right) \sqrt{e^{4c-4a} (6e^{4a-4c} - 4e^{2a-2c} - 4e^{6a-6c} + e^{8a-8c} + 1)}}{2\sqrt{b^2}}$$

input `int(cosh(a + b*x)^2/cosh(c + b*x),x)`output
$$\frac{\exp(2*a - c + b*x)/(2*b) - \exp(c - 2*a - b*x)/(2*b) - (\operatorname{atan}((\exp(-2*a)*\exp(3*c)*\exp(b*x)*(\sqrt{b^2} - 2*\exp(2*a)*\exp(-2*c)*\sqrt{b^2} + \exp(4*a)*\exp(-4*c)*\sqrt{b^2}))/(\sqrt{b^2}*(\exp(-4*a)*\exp(4*c)*(6*\exp(4*a)*\exp(-4*c) - 4*\exp(2*a)*\exp(-2*c) - 4*\exp(6*a)*\exp(-6*c) + \exp(8*a)*\exp(-8*c) + 1))^{1/2}))*(\exp(4*c - 4*a)*(6*\exp(4*a - 4*c) - 4*\exp(2*a - 2*c) - 4*\exp(6*a - 6*c) + \exp(8*a - 8*c) + 1))^{1/2})/(2*(b^2)^{1/2})$$
Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.72

$$\int \cosh^2(a + bx) \operatorname{sech}(c + bx) dx = \frac{-e^{bx+4a} \operatorname{atan}(e^{bx+c}) + 2e^{bx+2a+2c} \operatorname{atan}(e^{bx+c}) - e^{bx+4c} \operatorname{atan}(e^{bx+c}) + e^{2bx+4a+c} - e^{3c}}{2e^{bx+2a+2c}b}$$

input `int(cosh(b*x+a)^2*sech(b*x+c),x)`output
$$(-e^{4a+b*x}*\operatorname{atan}(e^{b*x+c}) + 2e^{2a+b*x+2c}*\operatorname{atan}(e^{b*x+c}) - e^{b*x+4c}*\operatorname{atan}(e^{b*x+c}) + e^{4a+2*b*x+c} - e^{3c})/(2e^{2a+b*x+2c}*b)$$

3.136 $\int \cosh^2(a + bx)\operatorname{sech}^2(c + bx) dx$

Optimal result	949
Mathematica [B] (verified)	949
Rubi [F]	950
Maple [B] (verified)	950
Fricas [B] (verification not implemented)	951
Sympy [F]	952
Maxima [A] (verification not implemented)	952
Giac [B] (verification not implemented)	952
Mupad [B] (verification not implemented)	953
Reduce [B] (verification not implemented)	953

Optimal result

Integrand size = 17, antiderivative size = 49

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c + bx) dx = x \cosh(2(a - c)) + \frac{\log(\cosh(c + bx)) \sinh(2(a - c))}{b} - \frac{\sinh^2(a - c) \tanh(c + bx)}{b}$$

output

```
x*cosh(2*a-2*c)+ln(cosh(b*x+c))*sinh(2*a-2*c)/b-sinh(a-c)^2*tanh(b*x+c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 177 vs. 2(49) = 98.

Time = 0.36 (sec) , antiderivative size = 177, normalized size of antiderivative = 3.61

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c + bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}(c + bx)(bx \cosh(2a - 4c - bx) + bx \cosh(2a - 2c - bx) + bx \cosh(2a + bx) + bx \cosh(2a - 2c + bx))}{b}$$

input

```
Integrate[Cosh[a + b*x]^2*Sech[c + b*x]^2,x]
```

output

```
(Sech[c]*Sech[c + b*x]*(b*x*Cosh[2*a - 4*c - b*x] + b*x*Cosh[2*a - 2*c - b*x] + b*x*Cosh[2*a + b*x] + b*x*Cosh[2*a - 2*c + b*x] + 2*Sinh[b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 4*c - b*x] + Sinh[2*a - 2*c - b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 2*c - b*x] + Log[Cosh[c + b*x]]*Sinh[2*a + b*x] - Sinh[2*a - 2*c + b*x] + Log[Cosh[c + b*x]]*Sinh[2*a - 2*c + b*x]))/(4*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

input

```
Int[Cosh[a + b*x]^2*Sech[c + b*x]^2,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(49) = 98$.

Time = 3.07 (sec) , antiderivative size = 248, normalized size of antiderivative = 5.06

method	result
risch	$x e^{2a-2c} - e^{-2c-2a} e^{4a} x + e^{-2c-2a} e^{4c} x - \frac{e^{-2c-2a} e^{4a} a}{b} + \frac{e^{-2c-2a} e^{4c} a}{b} + \frac{e^{-2c} e^{4a}}{2b(e^{2bx+2a+2c} + e^{2a})} - \frac{e^{-2c} e^{2a+2c}}{b(e^{2bx+2a+2c} + e^{2a})}$

input

```
int(cosh(b*x+a)^2*sech(b*x+c)^2,x,method=_RETURNVERBOSE)
```

output

```
x*exp(2*a-2*c)-exp(-2*c-2*a)*exp(4*a)*x+exp(-2*c-2*a)*exp(4*c)*x-1/b*exp(-
2*c-2*a)*exp(4*a)*a+1/b*exp(-2*c-2*a)*exp(4*c)*a+1/2/b*exp(-2*c)/(exp(2*b*
x+2*a+2*c)+exp(2*a))*exp(4*a)-1/b*exp(-2*c)/(exp(2*b*x+2*a+2*c)+exp(2*a))*
exp(2*a+2*c)+1/2/b*exp(-2*c)/(exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*c)+1/2*ln
(exp(2*b*x+2*a)+exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*a)-1/2*ln(exp(2*b*x+2*
a)+exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 694 vs. $2(49) = 98$.

Time = 0.11 (sec) , antiderivative size = 694, normalized size of antiderivative = 14.16

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x+c)^2,x, algorithm="fricas")
```

output

```
1/2*(2*b*x*cosh(b*x + c)^2 + cosh(-a + c)^4 + 4*b*x*cosh(b*x + c)*sinh(b*x
+ c) + 2*b*x*sinh(b*x + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a +
c)^4 + 2*(3*cosh(-a + c)^2 - 1)*sinh(-a + c)^2 + 2*b*x - 2*cosh(-a + c)^2
+ ((cosh(b*x + c)^2 + 1)*sinh(-a + c)^4 + cosh(-a + c)^4 - 4*(cosh(b*x + c
)^2*cosh(-a + c) + cosh(-a + c)*sinh(-a + c)^3 + (cosh(-a + c)^4 - 1)*cos
h(b*x + c)^2 + (cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a
+ c)^2*sinh(-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 -
1)*sinh(b*x + c)^2 + 6*(cosh(b*x + c)^2*cosh(-a + c)^2 + cosh(-a + c)^2)*s
inh(-a + c)^2 - 2*(4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) - 6*cosh(b*
x + c)*cosh(-a + c)^2*sinh(-a + c)^2 + 4*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - (cosh(-a + c)^4 - 1)*cosh(b*x +
c))*sinh(b*x + c) - 4*(cosh(b*x + c)^2*cosh(-a + c)^3 + cosh(-a + c)^3)*si
nh(-a + c) - 1)*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))) - 4*(
cosh(-a + c)^3 - cosh(-a + c))*sinh(-a + c) + 1)/(b*cosh(b*x + c)^2*cosh(-
a + c)^2 + b*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a
+ c) + b*sinh(-a + c)^2)*sinh(b*x + c)^2 + (b*cosh(b*x + c)^2 + b)*sinh(-
a + c)^2 + 2*(b*cosh(b*x + c)*cosh(-a + c)^2 - 2*b*cosh(b*x + c)*cosh(-a +
c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 2*(b*co
sh(b*x + c)^2*cosh(-a + c) + b*cosh(-a + c))*sinh(-a + c))
```


Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c + bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^2(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x+c)**2,x)`

output `Integral(cosh(a + b*x)**2*sech(b*x + c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.98

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c + bx) dx = \frac{(e^{4a} - e^{4c})e^{-2a-2c} \log(e^{-2bx} + e^{2c})}{2b} + \frac{(bx + a)e^{2a-2c}}{b} - \frac{e^{4a} - 2e^{2a+2c} + e^{4c}}{2b(e^{-2bx+2a} + e^{2a+2c})}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-2*b*x) + e^(2*c))/b + (b*x + a)*e^(2*a - 2*c)/b - 1/2*(e^(4*a) - 2*e^(2*a + 2*c) + e^(4*c))/(b*(e^(-2*b*x + 2*a) + e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(49) = 98.

Time = 0.13 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.08

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c + bx) dx = xe^{(-2a+2c)} + \frac{(e^{4a} - e^{4c})e^{-2a-2c} \log(e^{2bx+2c} + 1)}{2b} - \frac{(e^{2bx+4a} - e^{2bx+4c}) + 2e^{2a} - 2e^{2c}}{2b(e^{2bx+2c} + 1)}e^{(-2a)}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^2,x, algorithm="giac")`

output `x*e^(-2*a + 2*c) + 1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(2*b*x + 2*c) + 1)/b - 1/2*(e^(2*b*x + 4*a) - e^(2*b*x + 4*c) + 2*e^(2*a) - 2*e^(2*c))*e^(-2*a)/(b*(e^(2*b*x + 2*c) + 1))`

Mupad [B] (verification not implemented)

Time = 1.22 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.73

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c + bx) dx = x e^{2c-2a} + \frac{\sinh(2a - 2c) \ln(e^{2a} e^{2bx} + e^{2a} e^{-2c})}{b} + \frac{2 e^{2a-2c} \sinh(a - c)^2}{b (e^{2a-2c} + e^{2a+2bx})}$$

input `int(cosh(a + b*x)^2/cosh(c + b*x)^2,x)`

output `x*exp(2*c - 2*a) + (sinh(2*a - 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(-2*c)))/b + (2*exp(2*a - 2*c)*sinh(a - c)^2)/(b*(exp(2*a - 2*c) + exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 187, normalized size of antiderivative = 3.82

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c + bx) dx = \frac{e^{2bx+4a+2c}\log(e^{2bx+2c} + 1) - e^{2bx+4a+2c} + 2e^{2bx+2a+4c} - e^{2bx+6c}\log(e^{2bx+2c} + 1) + 2e^{2bx+6c}bx - e^{2bx+6c}}{2e^{2a+2c}b(e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)^2*sech(b*x+c)^2,x)`

output

```
(e**(4*a + 2*b*x + 2*c)*log(e**(2*b*x + 2*c) + 1) - e**(4*a + 2*b*x + 2*c)
+ 2*e**(2*a + 2*b*x + 4*c) - e**(2*b*x + 6*c)*log(e**(2*b*x + 2*c) + 1) +
2*e**(2*b*x + 6*c)*b*x - e**(2*b*x + 6*c) + e**(4*a)*log(e**(2*b*x + 2*c)
+ 1) - e**(4*c)*log(e**(2*b*x + 2*c) + 1) + 2*e**(4*c)*b*x)/(2*e**(2*a +
2*c)*b*(e**(2*b*x + 2*c) + 1))
```

3.137 $\int \cosh^2(a + bx)\operatorname{sech}^3(c + bx) dx$

Optimal result	955
Mathematica [A] (verified)	955
Rubi [F]	956
Maple [C] (verified)	956
Fricas [B] (verification not implemented)	957
Sympy [F]	958
Maxima [A] (verification not implemented)	959
Giac [A] (verification not implemented)	959
Mupad [F(-1)]	960
Reduce [B] (verification not implemented)	960

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \cosh^2(a + bx)\operatorname{sech}^3(c + bx) dx = \frac{\arctan(\sinh(c + bx)) \cosh(2(a - c))}{b} - \frac{\arctan(\sinh(c + bx)) \sinh^2(a - c)}{2b} - \frac{\operatorname{sech}(c + bx) \sinh(2(a - c))}{b} - \frac{\operatorname{sech}(c + bx) \sinh^2(a - c) \tanh(c + bx)}{2b}$$

output

```
arctan(sinh(b*x+c))*cosh(2*a-2*c)/b-1/2*arctan(sinh(b*x+c))*sinh(a-c)^2/b-
sech(b*x+c)*sinh(2*a-2*c)/b-1/2*sech(b*x+c)*sinh(a-c)^2*tanh(b*x+c)/b
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.23

$$\int \cosh^2(a + bx)\operatorname{sech}^3(c + bx) dx = \frac{12 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right) \cosh(2(a - c)) + \operatorname{sech}^2(c + bx) \left(2 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right)\right) + 2 \arctan\left(\tanh\left(\frac{1}{2}(c + bx)\right)\right)}{8b}$$

input `Integrate[Cosh[a + b*x]^2*Sech[c + b*x]^3,x]`

output `(12*ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(a - c)] + Sech[c + b*x]^2*(2*ArcTan[Tanh[(c + b*x)/2]] + 2*ArcTan[Tanh[(c + b*x)/2]]*Cosh[2*(c + b*x)] - 3*Sinh[2*a - 3*c - b*x] - 5*Sinh[2*a - c + b*x] + 2*Sinh[c + b*x]))/(8*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c + b*x]^3,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.22 (sec) , antiderivative size = 321, normalized size of antiderivative = 3.65

method	result
risch	$-\frac{(5e^{2bx+6a+2c}-2e^{2bx+4a+4c}-3e^{2bx+2a+6c}+3e^{6a}+2e^{4a+2c}-5e^{2a+4c})e^{bx-c}}{4(e^{2bx+2a+2c}+e^{2a})^2b} + \frac{3i \ln(e^{bx+a+ie^{a-c}})e^{-2c-2ae^{4a}}}{8b} + \frac{i \ln(e^{bx+a+ie^{a-c}})}{8b}$

input `int(cosh(b*x+a)^2*sech(b*x+c)^3,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*b*x+2*a+2*c)+exp(2*a))^2/b*(5*exp(2*b*x+6*a+2*c)-2*exp(2*b*x+4
*a+4*c)-3*exp(2*b*x+2*a+6*c)+3*exp(6*a)+2*exp(4*a+2*c)-5*exp(2*a+4*c))*exp
(b*x-c)+3/8*I*ln(exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(4*a)+1/4*I*ln(
exp(b*x+a)+I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*I*ln(exp(b*x+a)+I*
exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)-3/8*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-
2*c-2*a)*exp(4*a)-1/4*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(2*a+
2*c)-3/8*I*ln(exp(b*x+a)-I*exp(a-c))/b*exp(-2*c-2*a)*exp(4*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2173 vs. $2(84) = 168$.

Time = 0.12 (sec) , antiderivative size = 2173, normalized size of antiderivative = 24.69

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x+c)^3,x, algorithm="fricas")
```

output

```
-1/4*((5*cosh(b*x + c)^3 + 3*cosh(b*x + c))*sinh(-a + c)^4 + (5*cosh(-a +
c)^4 - 2*cosh(-a + c)^2 - 3)*cosh(b*x + c)^3 + (5*cosh(-a + c)^4 - 20*cosh
(-a + c)*sinh(-a + c)^3 + 5*sinh(-a + c)^4 + 2*(15*cosh(-a + c)^2 - 1)*sin
h(-a + c)^2 - 2*cosh(-a + c)^2 - 4*(5*cosh(-a + c)^3 - cosh(-a + c))*sinh(
-a + c) - 3)*sinh(b*x + c)^3 - 4*(5*cosh(b*x + c)^3*cosh(-a + c) + 3*cosh(
b*x + c)*cosh(-a + c))*sinh(-a + c)^3 - 3*(20*cosh(b*x + c)*cosh(-a + c)*s
inh(-a + c)^3 - 5*cosh(b*x + c)*sinh(-a + c)^4 - 2*(15*cosh(-a + c)^2 - 1)
*cosh(b*x + c)*sinh(-a + c)^2 + 4*(5*cosh(-a + c)^3 - cosh(-a + c))*cosh(b
*x + c)*sinh(-a + c) - (5*cosh(-a + c)^4 - 2*cosh(-a + c)^2 - 3)*cosh(b*x
+ c))*sinh(b*x + c)^2 + 2*((15*cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (9*co
sh(-a + c)^2 + 1)*cosh(b*x + c))*sinh(-a + c)^2 - ((3*cosh(-a + c)^4 + 2*c
osh(-a + c)^2 + 3)*cosh(b*x + c)^4 + (3*cosh(-a + c)^4 - 12*cosh(-a + c)*s
inh(-a + c)^3 + 3*sinh(-a + c)^4 + 2*(9*cosh(-a + c)^2 + 1)*sinh(-a + c)^2
+ 2*cosh(-a + c)^2 - 4*(3*cosh(-a + c)^3 + cosh(-a + c))*sinh(-a + c) + 3
)*sinh(b*x + c)^4 + 3*(cosh(b*x + c)^4 + 2*cosh(b*x + c)^2 + 1)*sinh(-a +
c)^4 + 3*cosh(-a + c)^4 - 4*(12*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3
- 3*cosh(b*x + c)*sinh(-a + c)^4 - 2*(9*cosh(-a + c)^2 + 1)*cosh(b*x + c)*
sinh(-a + c)^2 + 4*(3*cosh(-a + c)^3 + cosh(-a + c))*cosh(b*x + c)*sinh(-a
+ c) - (3*cosh(-a + c)^4 + 2*cosh(-a + c)^2 + 3)*cosh(b*x + c))*sinh(b*x
+ c)^3 - 12*(cosh(b*x + c)^4*cosh(-a + c) + 2*cosh(b*x + c)^2*cosh(-a + ...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c + bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^3(bx + c) dx$$

input

```
integrate(cosh(b*x+a)**2*sech(b*x+c)**3,x)
```

output

```
Integral(cosh(a + b*x)**2*sech(b*x + c)**3, x)
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.77

$$\int \cosh^2(a+bx)\operatorname{sech}^3(c+bx) dx = -\frac{(3e^{4a} + 2e^{(2a+2c)} + 3e^{4c}) \arctan(e^{(-bx-c)}) e^{(-2a-2c)}}{4b} - \frac{(5e^{(4a+2c)} - 2e^{(2a+4c)} - 3e^{(6c)})e^{(-bx-a)} + (3e^{(6a)} + 2e^{(4a+2c)} - 5e^{(2a+4c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+c)} + e^{(a+5c)})}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^3,x, algorithm="maxima")`

output

$$-1/4*(3*e^{(4*a)} + 2*e^{(2*a + 2*c)} + 3*e^{(4*c)})*\arctan(e^{(-b*x - c)})*e^{(-2*a - 2*c)/b} - 1/4*((5*e^{(4*a + 2*c)} - 2*e^{(2*a + 4*c)} - 3*e^{(6*c)})*e^{(-b*x - a)} + (3*e^{(6*a)} + 2*e^{(4*a + 2*c)} - 5*e^{(2*a + 4*c)})*e^{(-3*b*x - 3*a)})/(b*(2*e^{(-2*b*x + a + 3*c)} + e^{(-4*b*x + a + c)} + e^{(a + 5*c)}))$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.62

$$\int \cosh^2(a+bx)\operatorname{sech}^3(c+bx) dx = \frac{(3e^{4a} + 2e^{(2a+2c)} + 3e^{4c}) \arctan(e^{(bx+c)}) e^{(-2a-2c)}}{4b} - \frac{(5e^{(3bx+4a+2c)} - 2e^{(3bx+2a+4c)} - 3e^{(3bx+6c)} + 3e^{(bx+4a)} + 2e^{(bx+2a+2c)} - 5e^{(bx+4c)})e^{(-2a-c)}}{4b(e^{(2bx+2c)} + 1)^2}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^3,x, algorithm="giac")`

output

$$1/4*(3*e^{(4*a)} + 2*e^{(2*a + 2*c)} + 3*e^{(4*c)})*\arctan(e^{(b*x + c)})*e^{(-2*a - 2*c)/b} - 1/4*(5*e^{(3*b*x + 4*a + 2*c)} - 2*e^{(3*b*x + 2*a + 4*c)} - 3*e^{(3*b*x + 6*c)} + 3*e^{(b*x + 4*a)} + 2*e^{(b*x + 2*a + 2*c)} - 5*e^{(b*x + 4*c)})*e^{(-2*a - c)/(b*(e^{(2*b*x + 2*c)} + 1)^2)}$$

Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c + bx) dx = \int \frac{\cosh(a + bx)^2}{\cosh(c + bx)^3} dx$$

input `int(cosh(a + b*x)^2/cosh(c + b*x)^3,x)`output `int(cosh(a + b*x)^2/cosh(c + b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 303, normalized size of antiderivative = 3.44

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c + bx) dx = \frac{3e^{4bx+4a+4c} \operatorname{atan}(e^{bx+c}) + 2e^{4bx+2a+6c} \operatorname{atan}(e^{bx+c}) + 3e^{4bx+8c} \operatorname{atan}(e^{bx+c}) + 6e^{2bx+4a+2c} \operatorname{atan}(e^{bx+c}) + 4e^{2bx+4a+2c} \operatorname{atan}(e^{bx+c})}{(4e^{2bx+4a+2c} + 1)}$$

input `int(cosh(b*x+a)^2*sech(b*x+c)^3,x)`output `(3*e**(4*a + 4*b*x + 4*c)*atan(e**(b*x + c)) + 2*e**(2*a + 4*b*x + 6*c)*atan(e**(b*x + c)) + 3*e**(4*b*x + 8*c)*atan(e**(b*x + c)) + 6*e**(4*a + 2*b*x + 2*c)*atan(e**(b*x + c)) + 4*e**(2*a + 2*b*x + 4*c)*atan(e**(b*x + c)) + 6*e**(2*b*x + 6*c)*atan(e**(b*x + c)) + 3*e**(4*a)*atan(e**(b*x + c)) + 2*e**(2*a + 2*c)*atan(e**(b*x + c)) + 3*e**(4*c)*atan(e**(b*x + c)) - 5*e**(4*a + 3*b*x + 3*c) + 2*e**(2*a + 3*b*x + 5*c) + 3*e**(3*b*x + 7*c) - 3*e**(4*a + b*x + c) - 2*e**(2*a + b*x + 3*c) + 5*e**(b*x + 5*c))/(4*e**(2*a + 2*c)*b*(e**(4*b*x + 4*c) + 2*e**(2*b*x + 2*c) + 1))`

3.138 $\int \cosh^2(a + bx)\operatorname{sech}^4(c + bx) dx$

Optimal result	961
Mathematica [A] (verified)	961
Rubi [F]	962
Maple [A] (verified)	962
Fricas [B] (verification not implemented)	963
Sympy [F]	964
Maxima [B] (verification not implemented)	964
Giac [A] (verification not implemented)	965
Mupad [F(-1)]	965
Reduce [B] (verification not implemented)	966

Optimal result

Integrand size = 17, antiderivative size = 84

$$\int \cosh^2(a + bx)\operatorname{sech}^4(c + bx) dx = -\frac{\operatorname{sech}^2(c + bx) \sinh(2(a - c))}{2b} + \frac{\cosh(2(a - c)) \tanh(c + bx)}{b} - \frac{\sinh^2(a - c) \tanh(c + bx)}{b} + \frac{\sinh^2(a - c) \tanh^3(c + bx)}{3b}$$

```
output -1/2*sech(b*x+c)^2*sinh(2*a-2*c)/b+cosh(2*a-2*c)*tanh(b*x+c)/b-sinh(a-c)^2
*tanh(b*x+c)/b+1/3*sinh(a-c)^2*tanh(b*x+c)^3/b
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.95

$$\int \cosh^2(a + bx)\operatorname{sech}^4(c + bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}^3(c + bx)(3 \sinh(bx) - \sinh(2a - 4c - 3bx) - 3 \sinh(2a - 2c - bx) - 3 \sinh(2a + bx) + \sinh(2a - 2c - bx))}{12b}$$

input `Integrate[Cosh[a + b*x]^2*Sech[c + b*x]^4,x]`

output `(Sech[c]*Sech[c + b*x]^3*(3*Sinh[b*x] - Sinh[2*a - 4*c - 3*b*x] - 3*Sinh[2*a - 2*c - b*x] - 3*Sinh[2*a + b*x] + Sinh[2*a + 3*b*x] + Sinh[2*c + 3*b*x]))/(12*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c + b*x]^4,x]`

output `$Aborted`

Maple [A] (verified)

Time = 3.00 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.67

method	result	size
parallelrisch	$\frac{\sinh(3bx+3c)+2\sinh(3bx+2a+c)+3\sinh(bx+c)}{3b(\cosh(3bx+3c)+3\cosh(bx+c))}$	56
risch	$-\frac{2(3e^{4bx+4a+4c}+3e^{2bx+4a+2c}+3e^{2bx+2a+4c}+e^{4a}+e^{2a+2c}+e^{4c})e^{4a-2c}}{3(e^{2bx+2a+2c}+e^{2a})^3b}$	92

input `int(cosh(b*x+a)^2*sech(b*x+c)^4,x,method=_RETURNVERBOSE)`

output $\frac{1}{3} \frac{1}{b} (\sinh(3bx+3c) + 2\sinh(3bx+2a+c) + 3\sinh(bx+c)) / (\cosh(3bx+3c) + 3\cosh(bx+c))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(80) = 160$.

Time = 0.07 (sec) , antiderivative size = 493, normalized size of antiderivative = 5.87

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c + bx) dx = \frac{2 \left((5 \cosh(-a + c))^2 + 1 \right) \cosh(bx + c) \sinh(bx + c)}{3 (b \cosh(bx + c))^4 \cosh(-a + c)^2 + 4b \cosh(bx + c)^2 \cosh(-a + c)^2 + (b \cosh(-a + c))^2 - b \sinh(-a + c)}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^4,x, algorithm="fricas")`

output
$$\frac{-2/3 \left((5 \cosh(-a + c))^2 + 1 \right) \cosh(bx + c)^2 + (5 \cosh(-a + c))^2 - 6 \cosh(-a + c) \sinh(-a + c) + 5 \sinh(-a + c)^2 + 1 \right) \sinh(bx + c)^2 + (5 \cosh(bx + c)^2 + 3) \sinh(-a + c)^2 + 3 \cosh(-a + c)^2 - 2 \left(6 \cosh(bx + c) \cosh(-a + c) \sinh(-a + c) - \cosh(bx + c) \sinh(-a + c)^2 - (\cosh(-a + c))^2 - 1 \right) \cosh(bx + c) \sinh(bx + c) - 6 \left(\cosh(bx + c)^2 \cosh(-a + c) + \cosh(-a + c) \right) \sinh(-a + c) + 3}{(b \cosh(bx + c))^4 \cosh(-a + c)^2 + 4b \cosh(bx + c)^2 \cosh(-a + c)^2 + (b \cosh(-a + c))^2 - b \sinh(-a + c)^2} \sinh(bx + c)^4 + 4 \left(b \cosh(bx + c) \cosh(-a + c)^2 - b \cosh(bx + c) \sinh(-a + c)^2 \right) \sinh(bx + c)^3 + 3b \cosh(-a + c)^2 + 2 \left(3b \cosh(bx + c)^2 \cosh(-a + c)^2 + 2b \cosh(-a + c)^2 - (3b \cosh(bx + c)^2 + 2b) \sinh(-a + c)^2 \right) \sinh(bx + c)^2 - (b \cosh(bx + c))^4 + 4b \cosh(bx + c)^2 + 3b \sinh(-a + c)^2 + 4 \left(b \cosh(bx + c) \cosh(-a + c)^2 + b \cosh(bx + c) \cosh(-a + c)^2 - (b \cosh(bx + c))^3 + b \cosh(bx + c) \right) \sinh(-a + c)^2 \sinh(bx + c)}$$

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c + bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^4(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x+c)**4,x)`

output `Integral(cosh(a + b*x)**2*sech(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. $2(80) = 160$.

Time = 0.04 (sec) , antiderivative size = 321, normalized size of antiderivative = 3.82

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{sech}^4(c + bx) dx \\ &= \frac{2(e^{4a+4c} + e^{2a+6c})e^{-2bx-2a}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(2a+6c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \\ &+ \frac{2e^{(8c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} + e^{(2a+6c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*sech(b*x+c)^4,x, algorithm="maxima")`

output

$$\frac{2(e^{4a+4c} + e^{2a+6c})e^{-2bx-2a}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+2c} + e^{-6bx+2a} + e^{2a+6c})} + \frac{2e^{-4bx+4c}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+2c} + e^{-6bx+2a} + e^{2a+6c})} + \frac{2/3e^{4a+4c}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+2c} + e^{-6bx+2a} + e^{2a+6c})} + \frac{2/3e^{2a+6c}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+2c} + e^{-6bx+2a} + e^{2a+6c})} + \frac{2/3e^{8c}}{b(3e^{-2bx+2a+4c} + 3e^{-4bx+2a+2c} + e^{-6bx+2a} + e^{2a+6c})}$$
Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.01

$$\int \cosh^2(a+bx)\operatorname{sech}^4(c+bx) dx = \frac{2(3e^{4bx+4a+4c} + 3e^{2bx+4a+2c} + 3e^{2bx+2a+4c} + e^{4a} + e^{2a+2c} + e^{4c})e^{-2a-2c}}{3b(e^{2bx+2c} + 1)^3}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x+c)^4,x, algorithm="giac")
```

output

$$-\frac{2}{3} \frac{(3e^{4bx+4a+4c} + 3e^{2bx+4a+2c} + 3e^{2bx+2a+4c} + e^{4a} + e^{2a+2c} + e^{4c})e^{-2a-2c}}{(e^{2bx+2c} + 1)^3}$$
Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a+bx)\operatorname{sech}^4(c+bx) dx = \int \frac{\cosh(a+bx)^2}{\cosh(c+bx)^4} dx$$

input

```
int(cosh(a + b*x)^2/cosh(c + b*x)^4,x)
```

output

```
int(cosh(a + b*x)^2/cosh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.11

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c + bx) dx = \frac{\frac{2e^{6bx+4a+4c}}{3} - 2e^{2bx+2a+2c} - \frac{2e^{2a}}{3} - \frac{2e^{2c}}{3}}{e^{2a}b(e^{6bx+6c} + 3e^{4bx+4c} + 3e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)^2*sech(b*x+c)^4,x)`output `(2*(e**(4*a + 6*b*x + 4*c) - 3*e**(2*a + 2*b*x + 2*c) - e**(2*a) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x + 6*c) + 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) + 1))`

3.139 $\int \cosh^2(a + bx)\operatorname{sech}(c - bx) dx$

Optimal result	967
Mathematica [A] (verified)	967
Rubi [F]	968
Maple [C] (verified)	968
Fricas [B] (verification not implemented)	969
Sympy [F]	969
Maxima [B] (verification not implemented)	970
Giac [B] (verification not implemented)	970
Mupad [B] (verification not implemented)	971
Reduce [B] (verification not implemented)	971

Optimal result

Integrand size = 16, antiderivative size = 32

$$\int \cosh^2(a + bx)\operatorname{sech}(c - bx) dx = \frac{\arctan(\sinh(c - bx)) \sinh^2(a + c)}{b} + \frac{\sinh(2a + c + bx)}{b}$$

output

```
-arctan(sinh(b*x-c))*sinh(a+c)^2/b+sinh(b*x+2*a+c)/b
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.47

$$\int \cosh^2(a + bx)\operatorname{sech}(c - bx) dx = \frac{-\arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right) + \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right) \cosh(2(a + c)) + \sinh(2a + c + bx)}{b}$$

input

```
Integrate[Cosh[a + b*x]^2*Sech[c - b*x],x]
```

output

```
(-ArcTan[Tanh[(c - b*x)/2]] + ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(a + c)] + Sinh[2*a + c + b*x])/b
```


Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c - b*x],x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 231, normalized size of antiderivative = 7.22

method	result
risch	$\frac{e^{bx+2a+c}}{2b} - \frac{e^{-bx-2a-c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a} e^{4a+4c}}{4b} - \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a} e^{2a+2c}}{2b} + \frac{i \ln(e^{bx+a} - ie^{a+c}) e^{-2c-2a}}{4b}$

input `int(cosh(b*x+a)^2*sech(b*x-c),x,method=_RETURNVERBOSE)`

output `1/2/b*exp(b*x+2*a+c)-1/2/b*exp(-b*x-2*a-c)+1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)*exp(a+c)^4-1/2*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)*exp(a+c)^2+1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)-1/4*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(a+c)^4+1/2*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(a+c)^2-1/4*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 431 vs. $2(34) = 68$.

Time = 0.11 (sec) , antiderivative size = 431, normalized size of antiderivative = 13.47

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c),x, algorithm="fricas")`

output

```
1/2*(cosh(b*x + a)^2*cosh(a + c) - cosh(a + c)^3 + (cosh(a + c) - sinh(a +
c))*sinh(b*x + a)^2 - 3*cosh(a + c)*sinh(a + c)^2 + sinh(a + c)^3 - (4*co
sh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - 2*(3
*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 4*(cosh(a + c)^3 - cosh(
a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 - 2*cosh(a + c)^2 + 1)*
cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + sinh(a + c)
^4 + 2*(3*cosh(a + c)^2 - 1)*sinh(a + c)^2 - 2*cosh(a + c)^2 - 4*(cosh(a +
c)^3 - cosh(a + c))*sinh(a + c) + 1)*sinh(b*x + a))*arctan(-cosh(b*x + a)
*cosh(a + c) - (cosh(a + c) - sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*s
inh(a + c)) + 2*(cosh(b*x + a)*cosh(a + c) - cosh(b*x + a)*sinh(a + c))*si
nh(b*x + a) - (cosh(b*x + a)^2 - 3*cosh(a + c)^2)*sinh(a + c))/(b*cosh(b*x
+ a)*cosh(a + c)^2 - 2*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c) + b*cosh(b
*x + a)*sinh(a + c)^2 + (b*cosh(a + c)^2 - 2*b*cosh(a + c)*sinh(a + c) + b
*sinh(a + c)^2)*sinh(b*x + a))
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = \int \cosh^2(a + bx) \operatorname{sech}(bx - c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x-c),x)`

output `Integral(cosh(a + b*x)**2*sech(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.28

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = \frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1) \arctan(e^{(-bx+c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c),x, algorithm="maxima")`

output `1/2*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*arctan(e^(-b*x + c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.31

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = -\frac{(e^{(4a+4c)} - 2e^{(2a+2c)} + 1) \arctan(e^{(bx-c)}) e^{(-2a-2c)}}{2b} + \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c),x, algorithm="giac")`

output `-1/2*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)*arctan(e^(b*x - c))*e^(-2*a - 2*c)/b + 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 194, normalized size of antiderivative = 6.06

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = \frac{e^{2a+c+bx}}{2b} - \frac{e^{-2a-c-bx}}{2b} - \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{-3c} e^{bx} (\sqrt{b^2-2e^{2a} e^{2c} \sqrt{b^2} + e^{4a} e^{4c} \sqrt{b^2})}{b \sqrt{e^{-4a} e^{-4c} (6e^{4a} e^{4c} - 4e^{2a} e^{2c} - 4e^{6a} e^{6c} + e^{8a} e^{8c} + 1)}}}\right) \sqrt{e^{-4a-4c} (6e^{4a+4c} - 4e^{2a+2c} - 4e^{6a+6c} + e^{8a+8c} + 1)}}{2\sqrt{b^2}}$$

input `int(cosh(a + b*x)^2/cosh(c - b*x),x)`output
$$\frac{\exp(2*a + c + b*x)/(2*b) - \exp(-2*a - c - b*x)/(2*b) - (\operatorname{atan}((\exp(-2*a)*\exp(-3*c)*\exp(b*x)*((b^2)^{(1/2)} - 2*\exp(2*a)*\exp(2*c)*(b^2)^{(1/2)} + \exp(4*a)*\exp(4*c)*(b^2)^{(1/2)}))/(b*(\exp(-4*a)*\exp(-4*c)*(6*\exp(4*a)*\exp(4*c) - 4*\exp(2*a)*\exp(2*c) - 4*\exp(6*a)*\exp(6*c) + \exp(8*a)*\exp(8*c) + 1))^{(1/2)})) * (\exp(-4*a - 4*c)*(6*\exp(4*a + 4*c) - 4*\exp(2*a + 2*c) - 4*\exp(6*a + 6*c) + \exp(8*a + 8*c) + 1))^{(1/2)})/(2*(b^2)^{(1/2)})}$$
Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 109, normalized size of antiderivative = 3.41

$$\int \cosh^2(a + bx) \operatorname{sech}(c - bx) dx = \frac{-e^{bx+4a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 2e^{bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) - e^{bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + e^{2bx+4a+3c} - e^c}{2e^{bx+2a+2c} b}$$

input `int(cosh(b*x+a)^2*sech(b*x-c),x)`output
$$(-e^{4a + b*x + 4c} \operatorname{atan}(e^{b*x}/e^c) + 2e^{2a + b*x + 2c} \operatorname{atan}(e^{b*x}/e^c) - e^{b*x} \operatorname{atan}(e^{b*x}/e^c) + e^{4a + 2*b*x + 3c} - e^c)/(2e^{2a + b*x + 2c} * b)$$

3.140 $\int \cosh^2(a + bx)\operatorname{sech}^2(c - bx) dx$

Optimal result	972
Mathematica [B] (verified)	972
Rubi [F]	973
Maple [B] (verified)	973
Fricas [B] (verification not implemented)	974
Sympy [F]	975
Maxima [A] (verification not implemented)	975
Giac [B] (verification not implemented)	975
Mupad [B] (verification not implemented)	976
Reduce [B] (verification not implemented)	976

Optimal result

Integrand size = 18, antiderivative size = 44

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c - bx) dx = x \cosh(2(a + c)) + \frac{\log(\cosh(c - bx)) \sinh(2(a + c))}{b} + \frac{\sinh^2(a + c) \tanh(c - bx)}{b}$$

output

```
x*cosh(2*a+2*c)+ln(cosh(b*x-c))*sinh(2*a+2*c)/b-sinh(a+c)^2*tanh(b*x-c)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 182 vs. 2(44) = 88.

Time = 0.35 (sec) , antiderivative size = 182, normalized size of antiderivative = 4.14

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c - bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}(c - bx)(bx \cosh(2a + 2c - bx) + bx \cosh(2a + 4c - bx) + bx \cosh(2a + bx) + bx \cosh(2a + 2c - bx))}{b}$$

input

```
Integrate[Cosh[a + b*x]^2*Sech[c - b*x]^2,x]
```

output

```
(Sech[c]*Sech[c - b*x]*(b*x*Cosh[2*a + 2*c - b*x] + b*x*Cosh[2*a + 4*c - b*x] + b*x*Cosh[2*a + b*x] + b*x*Cosh[2*a + 2*c + b*x] + 2*Sinh[b*x] + Sinh[2*a + 2*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 2*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 4*c - b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + b*x] - Sinh[2*a + 2*c + b*x] + Log[Cosh[c - b*x]]*Sinh[2*a + 2*c + b*x]))/(4*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx)\operatorname{sech}^2(c - bx) dx$$

input

```
Int[Cosh[a + b*x]^2*Sech[c - b*x]^2,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. 2(51) = 102.

Time = 1.63 (sec) , antiderivative size = 239, normalized size of antiderivative = 5.43

method	result
risch	$x e^{2a+2c} - e^{-2c-2a} e^{4a+4c} x - \frac{e^{-2c-2a} a e^{4a+4c}}{b} + e^{-2c-2a} x + \frac{e^{-2c-2a} a}{b} + \frac{e^{4a+4c}}{2b(e^{2a+2c} + e^{2bx+2a})} - \frac{e^{2a+2c}}{b(e^{2a+2c} + e^{2bx+2a})}$

input

```
int(cosh(b*x+a)^2*sech(b*x-c)^2,x,method=_RETURNVERBOSE)
```

output

```
x*exp(2*a+2*c)-exp(-2*c-2*a)*exp(4*a+4*c)*x-1/b*exp(-2*c-2*a)*a*exp(4*a+4*c)+exp(-2*c-2*a)*x+1/b*exp(-2*c-2*a)*a+1/2/b/(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(4*a+4*c)-1/b/(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(2*a+2*c)+1/2/b/(exp(2*a+2*c)+exp(2*b*x+2*a))+1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/2*ln(exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1306 vs. $2(51) = 102$.

Time = 0.11 (sec) , antiderivative size = 1306, normalized size of antiderivative = 29.68

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x-c)^2,x, algorithm="fricas")
```

output

```
1/2*(2*b*x*cosh(b*x + a)^2*cosh(a + c)^6 - 12*b*x*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^5 + 2*b*x*cosh(b*x + a)^2*sinh(a + c)^6 + (2*b*x + 1)*cosh(a + c)^4 + (30*b*x*cosh(b*x + a)^2*cosh(a + c)^2 + 2*b*x + 1)*sinh(a + c)^4 - 4*(10*b*x*cosh(b*x + a)^2*cosh(a + c)^3 + (2*b*x + 1)*cosh(a + c))*sinh(a + c)^3 + 2*(b*x*cosh(a + c)^6 - 6*b*x*cosh(a + c)^5*sinh(a + c) + 15*b*x*cosh(a + c)^4*sinh(a + c)^2 - 20*b*x*cosh(a + c)^3*sinh(a + c)^3 + 15*b*x*cosh(a + c)^2*sinh(a + c)^4 - 6*b*x*cosh(a + c)*sinh(a + c)^5 + b*x*sinh(a + c)^6)*sinh(b*x + a)^2 + 2*(15*b*x*cosh(b*x + a)^2*cosh(a + c)^4 + 3*(2*b*x + 1)*cosh(a + c)^2 - 1)*sinh(a + c)^2 - 2*cosh(a + c)^2 + (6*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)^2*sinh(a + c)^6 - (15*cosh(b*x + a)^2*cosh(a + c)^2 + 1)*sinh(a + c)^4 - cosh(a + c)^4 + 4*(5*cosh(b*x + a)^2*cosh(a + c)^3 + cosh(a + c))*sinh(a + c)^3 - (cosh(a + c)^6 - cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^6 - 20*cosh(a + c)^3*sinh(a + c)^3 + 15*cosh(a + c)^2*sinh(a + c)^4 - 6*cosh(a + c)*sinh(a + c)^5 + sinh(a + c)^6 + (15*cosh(a + c)^4 - 1)*sinh(a + c)^2 - cosh(a + c)^2 - 2*(3*cosh(a + c)^5 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((15*cosh(a + c)^4 - 1)*cosh(b*x + a)^2 + 6*cosh(a + c)^2)*sinh(a + c)^2 + 2*(20*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c)^3 - 15*cosh(b*x + a)*cosh(a + c)^2*sinh(a + c)^4 + 6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)*sinh(a + c)^6 - (15*cosh(a + c)^4 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(3*...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^2(bx - c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x-c)**2,x)`

output `Integral(cosh(a + b*x)**2*sech(b*x - c)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.30

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(-2bx+2c)} + 1)}{2b} + \frac{(bx + a)e^{(2a+2c)}}{b} - \frac{e^{(4a+4c)} - 2e^{(2a+2c)} + 1}{2b(e^{(-2bx+2a+4c)} + e^{(2a+2c)})}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-2*b*x + 2*c) + 1)/b + (b*x + a)*e^(2*a + 2*c)/b - 1/2*(e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 1)/(b*(e^(-2*b*x + 2*a + 4*c) + e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(51) = 102.

Time = 0.14 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.41

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = xe^{(-2a-2c)} + \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(2bx)} + e^{(2c)})}{2b} + \frac{(e^{(2bx)} - e^{(2bx+4a+4c)} - 2e^{(2a+4c)} + 2e^{(2c)})e^{(-2a-2c)}}{2b(e^{(2bx)} + e^{(2c)})}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c)^2,x, algorithm="giac")`

output `x*e^(-2*a - 2*c) + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(2*b*x) + e^(2*c))/b + 1/2*(e^(2*b*x) - e^(2*b*x + 4*a + 4*c) - 2*e^(2*a + 4*c) + 2*e^(2*c))*e^(-2*a - 2*c)/(b*(e^(2*b*x) + e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.89

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = x e^{-2a-2c} + \frac{\sinh(2a + 2c) \ln(e^{2a} e^{2bx} + e^{2a} e^{2c})}{b} + \frac{2 e^{2a+2c} \sinh(a + c)^2}{b (e^{2a+2c} + e^{2a+2bx})}$$

input `int(cosh(a + b*x)^2/cosh(c - b*x)^2,x)`

output `x*exp(- 2*a - 2*c) + (sinh(2*a + 2*c)*log(exp(2*a)*exp(2*b*x) + exp(2*a)*exp(2*c)))/b + (2*exp(2*a + 2*c)*sinh(a + c)^2)/(b*(exp(2*a + 2*c) + exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 179, normalized size of antiderivative = 4.07

$$\int \cosh^2(a + bx) \operatorname{sech}^2(c - bx) dx = \frac{e^{2bx+4a+4c} \log(e^{2bx} + e^{2c}) - e^{2bx+4a+4c} + 2e^{2bx+2a+2c} - e^{2bx} \log(e^{2bx} + e^{2c}) + 2e^{2bx} bx - e^{2bx} + e^{4a+6c} \log(e^{2bx} + e^{2c})}{2e^{2a+2c} b (e^{2bx} + e^{2c})}$$

input `int(cosh(b*x+a)^2*sech(b*x-c)^2,x)`

output

```
(e**(4*a + 2*b*x + 4*c)*log(e**(2*b*x) + e**(2*c)) - e**(4*a + 2*b*x + 4*c)
) + 2*e**(2*a + 2*b*x + 2*c) - e**(2*b*x)*log(e**(2*b*x) + e**(2*c)) + 2*e
**(2*b*x)*b*x - e**(2*b*x) + e**(4*a + 6*c)*log(e**(2*b*x) + e**(2*c)) - e
**(2*c)*log(e**(2*b*x) + e**(2*c)) + 2*e**(2*c)*b*x)/(2*e**(2*a + 2*c)*b*(
e**(2*b*x) + e**(2*c)))
```

3.141 $\int \cosh^2(a + bx)\operatorname{sech}^3(c - bx) dx$

Optimal result	978
Mathematica [A] (verified)	978
Rubi [F]	979
Maple [C] (verified)	979
Fricas [B] (verification not implemented)	980
Sympy [F]	980
Maxima [A] (verification not implemented)	981
Giac [A] (verification not implemented)	981
Mupad [F(-1)]	982
Reduce [B] (verification not implemented)	982

Optimal result

Integrand size = 18, antiderivative size = 86

$$\int \cosh^2(a + bx)\operatorname{sech}^3(c - bx) dx = -\frac{\arctan(\sinh(c - bx)) \cosh(2(a + c))}{b} + \frac{\arctan(\sinh(c - bx)) \sinh^2(a + c)}{2b} - \frac{\operatorname{sech}(c - bx) \sinh(2(a + c))}{b} + \frac{\operatorname{sech}(c - bx) \sinh^2(a + c) \tanh(c - bx)}{2b}$$

output

```
arctan(sinh(b*x-c))*cosh(2*a+2*c)/b-1/2*arctan(sinh(b*x-c))*sinh(a+c)^2/b-
sech(b*x-c)*sinh(2*a+2*c)/b-1/2*sech(b*x-c)*sinh(a+c)^2*tanh(b*x-c)/b
```

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.28

$$\int \cosh^2(a + bx)\operatorname{sech}^3(c - bx) dx = \frac{12 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right) \cosh(2(a + c)) + \operatorname{sech}^2(c - bx) \left(2 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right)\right) + 2 \arctan\left(\tanh\left(\frac{1}{2}(c - bx)\right)\right)}{8b}$$

input `Integrate[Cosh[a + b*x]^2*Sech[c - b*x]^3,x]`

output `-1/8*(12*ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(a + c)] + Sech[c - b*x]^2*(2*ArcTan[Tanh[(c - b*x)/2]] + 2*ArcTan[Tanh[(c - b*x)/2]]*Cosh[2*(c - b*x)] + 2*Sinh[c - b*x] + 3*Sinh[2*a + 3*c - b*x] + 5*Sinh[2*a + c + b*x]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c - b*x]^3,x]`

output `$Aborted`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 4.59 (sec) , antiderivative size = 311, normalized size of antiderivative = 3.62

method	result
risch	$-\frac{(3e^{6a+6c}+5e^{2bx+6a+4c}+2e^{4a+4c}-2e^{2bx+4a+2c}-5e^{2a+2c}-3e^{2bx+2a})e^{bx-c}}{4(e^{2a+2c}+e^{2bx+2a})^2b} + \frac{3i \ln(e^{bx+a}+ie^{a+c})e^{-2c-2ae^{4a+4c}}}{8b} + \frac{i \ln(e^{bx+a}-ie^{a+c})e^{-2c-2ae^{4a+4c}}}{8b}$

input `int(cosh(b*x+a)^2*sech(b*x-c)^3,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*a+2*c)+exp(2*b*x+2*a))^2/b*(3*exp(6*a+6*c)+5*exp(2*b*x+6*a+4*c)
)+2*exp(4*a+4*c)-2*exp(2*b*x+4*a+2*c)-5*exp(2*a+2*c)-3*exp(2*b*x+2*a))*exp
(b*x-c)+3/8*I*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(4*a+4*c)+1/4*I
*ln(exp(b*x+a)+I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*I*ln(exp(b*x+a
)+I*exp(a+c))/b*exp(-2*c-2*a)-3/8*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2
*a)*exp(4*a+4*c)-1/4*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)*exp(2*a+2
*c)-3/8*I*ln(exp(b*x+a)-I*exp(a+c))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3893 vs. $2(90) = 180$.

Time = 0.13 (sec) , antiderivative size = 3893, normalized size of antiderivative = 45.27

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x-c)^3,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^3(bx - c) dx$$

input

```
integrate(cosh(b*x+a)**2*sech(b*x-c)**3,x)
```

output

```
Integral(cosh(a + b*x)**2*sech(b*x - c)**3, x)
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.77

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

$$= -\frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3) \arctan(e^{(-bx+c)}) e^{(-2a-2c)}}{4b}$$

$$-\frac{(5e^{(4a+4c)} - 2e^{(2a+2c)} - 3)e^{(-bx-a)} + (3e^{(6a+6c)} + 2e^{(4a+4c)} - 5e^{(2a+2c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} + e^{(-4bx+a+5c)} + e^{(a+c)})}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c)^3,x, algorithm="maxima")`

output

```
-1/4*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*arctan(e^(-b*x + c))*e^(-2*a
- 2*c)/b - 1/4*((5*e^(4*a + 4*c) - 2*e^(2*a + 2*c) - 3)*e^(-b*x - a) + (3*
e^(6*a + 6*c) + 2*e^(4*a + 4*c) - 5*e^(2*a + 2*c))*e^(-3*b*x - 3*a))/(b*(2
*e^(-2*b*x + a + 3*c) + e^(-4*b*x + a + 5*c) + e^(a + c)))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.65

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx = \frac{(3e^{(4a+4c)} + 2e^{(2a+2c)} + 3) \arctan(e^{(bx-c)}) e^{(-2a-2c)}}{4b}$$

$$+ \frac{(3e^{(3bx)} - 5e^{(3bx+4a+4c)} + 2e^{(3bx+2a+2c)} - 3e^{(bx+4a+6c)} - 2e^{(bx+2a+4c)} + 5e^{(bx+2c)})e^{(-2a-c)}}{4b(e^{(2bx)} + e^{(2c)})^2}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c)^3,x, algorithm="giac")`

output

```
1/4*(3*e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 3)*arctan(e^(b*x - c))*e^(-2*a -
2*c)/b + 1/4*(3*e^(3*b*x) - 5*e^(3*b*x + 4*a + 4*c) + 2*e^(3*b*x + 2*a +
2*c) - 3*e^(b*x + 4*a + 6*c) - 2*e^(b*x + 2*a + 4*c) + 5*e^(b*x + 2*c))*e^(
-2*a - c)/(b*(e^(2*b*x) + e^(2*c))^2)
```

Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx = \int \frac{\cosh(a + bx)^2}{\cosh(c - bx)^3} dx$$

input `int(cosh(a + b*x)^2/cosh(c - b*x)^3,x)`output `int(cosh(a + b*x)^2/cosh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 339, normalized size of antiderivative = 3.94

$$\int \cosh^2(a + bx) \operatorname{sech}^3(c - bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 2e^{4bx+2a+2c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 3e^{4bx} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 6e^{2bx+4a+6c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right) + 4e^{2bx+2a+4c} \operatorname{atan}\left(\frac{e^{bx}}{e^c}\right)}{4e^{2a+2c} b (e^{4bx} + 2e^{2bx+2c} + e^{4c})}$$

input `int(cosh(b*x+a)^2*sech(b*x-c)^3,x)`output `(3*e**(4*a + 4*b*x + 4*c)*atan(e**(b*x)/e**c) + 2*e**(2*a + 4*b*x + 2*c)*atan(e**(b*x)/e**c) + 3*e**(4*b*x)*atan(e**(b*x)/e**c) + 6*e**(4*a + 2*b*x + 6*c)*atan(e**(b*x)/e**c) + 4*e**(2*a + 2*b*x + 4*c)*atan(e**(b*x)/e**c) + 6*e**(2*b*x + 2*c)*atan(e**(b*x)/e**c) + 3*e**(4*a + 8*c)*atan(e**(b*x)/e**c) + 2*e**(2*a + 6*c)*atan(e**(b*x)/e**c) + 3*e**(4*c)*atan(e**(b*x)/e**c) - 5*e**(4*a + 3*b*x + 5*c) + 2*e**(2*a + 3*b*x + 3*c) + 3*e**(3*b*x + c) - 3*e**(4*a + b*x + 7*c) - 2*e**(2*a + b*x + 5*c) + 5*e**(b*x + 3*c))/(4*e**(2*a + 2*c)*b*(e**(4*b*x) + 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.142 $\int \cosh^2(a + bx)\operatorname{sech}^4(c - bx) dx$

Optimal result	983
Mathematica [A] (verified)	983
Rubi [F]	984
Maple [A] (verified)	984
Fricas [B] (verification not implemented)	985
Sympy [F]	986
Maxima [B] (verification not implemented)	986
Giac [A] (verification not implemented)	987
Mupad [F(-1)]	987
Reduce [B] (verification not implemented)	988

Optimal result

Integrand size = 18, antiderivative size = 80

$$\int \cosh^2(a + bx)\operatorname{sech}^4(c - bx) dx = -\frac{\operatorname{sech}^2(c - bx) \sinh(2(a + c))}{2b} - \frac{\cosh(2(a + c)) \tanh(c - bx)}{b} + \frac{\sinh^2(a + c) \tanh(c - bx)}{b} - \frac{\sinh^2(a + c) \tanh^3(c - bx)}{3b}$$

```
output -1/2*sech(b*x-c)^2*sinh(2*a+2*c)/b+cosh(2*a+2*c)*tanh(b*x-c)/b-sinh(a+c)^2
*tanh(b*x-c)/b+1/3*sinh(a+c)^2*tanh(b*x-c)^3/b
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \cosh^2(a + bx)\operatorname{sech}^4(c - bx) dx = \frac{\operatorname{sech}(c)\operatorname{sech}^3(c - bx)(-3 \sinh(bx) + \sinh(2c - 3bx)) + \sinh(2a + 4c - 3bx) + 3 \sinh(2a + 2c - bx) + 3}{12b}$$

input `Integrate[Cosh[a + b*x]^2*Sech[c - b*x]^4,x]`

output `-1/12*(Sech[c]*Sech[c - b*x]^3*(-3*Sinh[b*x] + Sinh[2*c - 3*b*x] + Sinh[2*a + 4*c - 3*b*x] + 3*Sinh[2*a + 2*c - b*x] + 3*Sinh[2*a + b*x] - Sinh[2*a + 3*b*x]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Sech[c - b*x]^4,x]`

output `$Aborted`

Maple [A] (verified)

Time = 3.53 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.78

method	result	size
parallelrisch	$\frac{\sinh(3bx-3c)+3\sinh(bx-c)+2\sinh(3bx+2a-c)}{3b(\cosh(3bx-3c)+3\cosh(bx-c))}$	62
risch	$-\frac{2(e^{4a+4c}+3e^{2bx+4a+2c}+e^{2a+2c}+3e^{4bx+4a}+3e^{2bx+2a}+1)e^{4a+4c}}{3(e^{2a+2c}+e^{2bx+2a})^3b}$	88

input `int(cosh(b*x+a)^2*sech(b*x-c)^4,x,method=_RETURNVERBOSE)`

output

```
1/3/b*(sinh(3*b*x-3*c)+3*sinh(b*x-c)+2*sinh(3*b*x+2*a-c))/(cosh(3*b*x-3*c)
+3*cosh(b*x-c))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1007 vs. $2(84) = 168$.

Time = 0.10 (sec) , antiderivative size = 1007, normalized size of antiderivative = 12.59

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x-c)^4,x, algorithm="fricas")
```

output

```
-2/3*(4*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^3 + (5*cosh(b*x + a)^2 + 3
)*sinh(a + c)^4 + 3*cosh(a + c)^4 + (5*cosh(a + c)^4 + cosh(a + c)^2)*cosh
(b*x + a)^2 + (5*cosh(a + c)^4 + 4*cosh(a + c)*sinh(a + c)^3 + 5*sinh(a +
c)^4 - (2*cosh(a + c)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(2*cosh(a +
c)^3 + cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((2*cosh(a + c)^2 - 1)
*cosh(b*x + a)^2 + 6*cosh(a + c)^2 - 3)*sinh(a + c)^2 + 3*cosh(a + c)^2 -
2*(4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4
+ (10*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3
+ cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 - cosh(a + c)^2
)*cosh(b*x + a))*sinh(b*x + a) + 2*((2*cosh(a + c)^3 + cosh(a + c))*cosh(b
*x + a)^2 + 3*cosh(a + c))*sinh(a + c))/(b*cosh(b*x + a)^4*cosh(a + c)^4 +
4*b*cosh(b*x + a)^2*cosh(a + c)^4 + 3*b*cosh(a + c)^4 + (b*cosh(a + c)^4
- 4*b*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(a + c)^2*sinh(a + c)^2 - 4*b*co
sh(a + c)*sinh(a + c)^3 + b*sinh(a + c)^4)*sinh(b*x + a)^4 + (b*cosh(b*x +
a)^4 - 4*b*cosh(b*x + a)^2 + 3*b)*sinh(a + c)^4 + 4*(b*cosh(b*x + a)*cosh
(a + c)^4 - 4*b*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(b*x + a
)*cosh(a + c)^2*sinh(a + c)^2 - 4*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^
3 + b*cosh(b*x + a)*sinh(a + c)^4)*sinh(b*x + a)^3 - 4*(b*cosh(b*x + a)^4*
cosh(a + c) - b*cosh(b*x + a)^2*cosh(a + c))*sinh(a + c)^3 + 2*(3*b*cosh(b
*x + a)^2*cosh(a + c)^4 + 18*b*cosh(b*x + a)^2*cosh(a + c)^2*sinh(a + c)...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx = \int \cosh^2(a + bx) \operatorname{sech}^4(bx - c) dx$$

input `integrate(cosh(b*x+a)**2*sech(b*x-c)**4,x)`

output `Integral(cosh(a + b*x)**2*sech(b*x - c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(84) = 168$.

Time = 0.04 (sec) , antiderivative size = 332, normalized size of antiderivative = 4.15

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx \\ &= \frac{2(e^{(4a+4c)} + e^{(2a+2c)})e^{(-2bx-2a)}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} + e^{(2a+2c)})} \\ &+ \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} + e^{(2a+2c)})} \\ &+ \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} + e^{(2a+2c)})} \\ &+ \frac{2e^{(2a+2c)}}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} + e^{(2a+2c)})} \\ &+ \frac{2}{3b(3e^{(-2bx+2a+4c)} + 3e^{(-4bx+2a+6c)} + e^{(-6bx+2a+8c)} + e^{(2a+2c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*sech(b*x-c)^4,x, algorithm="maxima")`

output

$$2*(e^{(4*a + 4*c)} + e^{(2*a + 2*c)})*e^{(-2*b*x - 2*a)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} + e^{(2*a + 2*c)})} + 2*e^{(-4*b*x + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} + e^{(2*a + 2*c)})} + 2/3*e^{(4*a + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} + e^{(2*a + 2*c)})} + 2/3*e^{(2*a + 2*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} + e^{(2*a + 2*c)})} + 2/3/(b*(3*e^{(-2*b*x + 2*a + 4*c)} + 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} + e^{(2*a + 2*c)}))$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.05

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx = \frac{2(3e^{(4bx+4a+4c)} + 3e^{(2bx+4a+6c)} + 3e^{(2bx+2a+4c)} + e^{(4a+8c)} + e^{(2a+6c)} + e^{(4c)})e^{(-2a)}}{3b(e^{(2bx)} + e^{(2c)})^3}$$

input

```
integrate(cosh(b*x+a)^2*sech(b*x-c)^4,x, algorithm="giac")
```

output

$$-2/3*(3*e^{(4*b*x + 4*a + 4*c)} + 3*e^{(2*b*x + 4*a + 6*c)} + 3*e^{(2*b*x + 2*a + 4*c)} + e^{(4*a + 8*c)} + e^{(2*a + 6*c)} + e^{(4*c)})*e^{(-2*a)/(b*(e^{(2*b*x)} + e^{(2*c)})^3)}$$
Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx = \int \frac{\cosh(a + bx)^2}{\cosh(c - bx)^4} dx$$

input

```
int(cosh(a + b*x)^2/cosh(c - b*x)^4,x)
```

output

```
int(cosh(a + b*x)^2/cosh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.24

$$\int \cosh^2(a + bx) \operatorname{sech}^4(c - bx) dx = \frac{2e^{2c}(e^{6bx+4a} - 3e^{2bx+2a+2c} - e^{2a+4c} - e^{2c})}{3e^{2a}b(e^{6bx} + 3e^{4bx+2c} + 3e^{2bx+4c} + e^{6c})}$$

input `int(cosh(b*x+a)^2*sech(b*x-c)^4,x)`output `(2*e**(2*c)*(e**(4*a + 6*b*x) - 3*e**(2*a + 2*b*x + 2*c) - e**(2*a + 4*c) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x) + 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) + e**(6*c)))`

3.143 $\int \cosh(x) \operatorname{csch}(2x) dx$

Optimal result	989
Mathematica [A] (verified)	989
Rubi [A] (verified)	990
Maple [A] (verified)	991
Fricas [B] (verification not implemented)	992
Sympy [F]	992
Maxima [B] (verification not implemented)	992
Giac [B] (verification not implemented)	993
Mupad [B] (verification not implemented)	993
Reduce [B] (verification not implemented)	993

Optimal result

Integrand size = 7, antiderivative size = 7

$$\int \cosh(x) \operatorname{csch}(2x) dx = -\frac{1}{2} \operatorname{arctanh}(\cosh(x))$$

output `-1/2*arctanh(cosh(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \cosh(x) \operatorname{csch}(2x) dx = -\frac{1}{2} \operatorname{arctanh}(\cosh(x))$$

input `Integrate[Cosh[x]*Csch[2*x],x]`

output `-1/2*ArcTanh[Cosh[x]]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 26, 4775, 26, 3042, 26, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \operatorname{csch}(2x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{i \cos(ix)}{\sin(2ix)} dx \\
 & \quad \downarrow \text{26} \\
 & i \int \frac{\cos(ix)}{\sin(2ix)} dx \\
 & \quad \downarrow \text{4775} \\
 & \frac{1}{2} i \int -i \operatorname{csch}(x) dx \\
 & \quad \downarrow \text{26} \\
 & \frac{\int \operatorname{csch}(x) dx}{2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int i \operatorname{csc}(ix) dx \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{2} i \int \operatorname{csc}(ix) dx \\
 & \quad \downarrow \text{4257} \\
 & -\frac{1}{2} \operatorname{arctanh}(\cosh(x))
 \end{aligned}$$

input

Int [Cosh[x]*Csch[2*x], x]

output `-1/2*ArcTanh[Cosh[x]]`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4775 `Int[(cos[(a_.) + (b_.)*(x_)])*(e_.))^(m_.)*sin[(c_.) + (d_.)*(x_)^(p_.), x_Symbol] := Simp[2^p/e^p Int[(e*Cos[a + b*x])^(m + p)*Sin[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b*c - a*d, 0] && EqQ[d/b, 2] && IntegerQ[p]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
default	$\frac{\ln(\tanh(\frac{x}{2}))}{2}$	8
risch	$\frac{\ln(e^x-1)}{2} - \frac{\ln(e^x+1)}{2}$	16

input `int(cosh(x)*csch(2*x),x,method=_RETURNVERBOSE)`

output `1/2*ln(tanh(1/2*x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.71

$$\int \cosh(x)\operatorname{csch}(2x) dx = -\frac{1}{2} \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{2} \log(\cosh(x) + \sinh(x) - 1)$$

input `integrate(cosh(x)*csch(2*x),x, algorithm="fricas")`

output `-1/2*log(cosh(x) + sinh(x) + 1) + 1/2*log(cosh(x) + sinh(x) - 1)`

Sympy [F]

$$\int \cosh(x)\operatorname{csch}(2x) dx = \int \cosh(x) \operatorname{csch}(2x) dx$$

input `integrate(cosh(x)*csch(2*x),x)`

output `Integral(cosh(x)*csch(2*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.71

$$\int \cosh(x)\operatorname{csch}(2x) dx = -\frac{1}{2} \log(e^{-x} + 1) + \frac{1}{2} \log(e^{-x} - 1)$$

input `integrate(cosh(x)*csch(2*x),x, algorithm="maxima")`

output `-1/2*log(e^(-x) + 1) + 1/2*log(e^(-x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 16 vs. $2(5) = 10$.

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 2.29

$$\int \cosh(x) \operatorname{csch}(2x) dx = -\frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

input `integrate(cosh(x)*csch(2*x),x, algorithm="giac")`

output `-1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.71

$$\int \cosh(x) \operatorname{csch}(2x) dx = \frac{\ln(1 - e^x)}{2} - \frac{\ln(-e^x - 1)}{2}$$

input `int(cosh(x)/sinh(2*x),x)`

output `log(1 - exp(x))/2 - log(- exp(x) - 1)/2`

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.43

$$\int \cosh(x) \operatorname{csch}(2x) dx = \frac{\log(e^x - 1)}{2} - \frac{\log(e^x + 1)}{2}$$

input `int(cosh(x)*csch(2*x),x)`

output `(log(e**x - 1) - log(e**x + 1))/2`

3.144 $\int \cosh(x) \operatorname{csch}(3x) dx$

Optimal result	994
Mathematica [A] (verified)	994
Rubi [A] (verified)	995
Maple [A] (verified)	997
Fricas [B] (verification not implemented)	997
Sympy [F]	998
Maxima [B] (verification not implemented)	998
Giac [B] (verification not implemented)	998
Mupad [B] (verification not implemented)	999
Reduce [B] (verification not implemented)	999

Optimal result

Integrand size = 7, antiderivative size = 21

$$\int \cosh(x) \operatorname{csch}(3x) dx = \frac{1}{3} \log(\sinh(x)) - \frac{1}{6} \log(3 + 4 \sinh^2(x))$$

output `1/3*ln(sinh(x))-1/6*ln(3+4*sinh(x)^2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \cosh(x) \operatorname{csch}(3x) dx = \frac{1}{3} \log(\sinh(x)) - \frac{1}{6} \log(3 + 4 \sinh^2(x))$$

input `Integrate[Cosh[x]*Csch[3*x],x]`

output `Log[Sinh[x]]/3 - Log[3 + 4*Sinh[x]^2]/6`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {3042, 26, 4856, 26, 243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x) \operatorname{csch}(3x) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{i \cos(ix)}{\sin(3ix)} dx \\
 & \quad \downarrow 26 \\
 & i \int \frac{\cos(ix)}{\sin(3ix)} dx \\
 & \quad \downarrow 4856 \\
 & i \int -\frac{i \operatorname{csch}(x)}{4 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow 26 \\
 & \int \frac{\operatorname{csch}(x)}{4 \sinh^2(x) + 3} d \sinh(x) \\
 & \quad \downarrow 243 \\
 & \frac{1}{2} \int \frac{\operatorname{csch}(x)}{4 \sinh^2(x) + 3} d \sinh^2(x) \\
 & \quad \downarrow 47 \\
 & \frac{1}{2} \left(\frac{1}{3} \int \operatorname{csch}(x) d \sinh^2(x) - \frac{4}{3} \int \frac{1}{4 \sinh^2(x) + 3} d \sinh^2(x) \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{2} \left(\frac{1}{3} \log(\sinh^2(x)) - \frac{4}{3} \int \frac{1}{4 \sinh^2(x) + 3} d \sinh^2(x) \right) \\
 & \quad \downarrow 16
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{3} \log(\sinh^2(x)) - \frac{1}{3} \log(4\sinh^2(x) + 3) \right)$$

input `Int[Cosh[x]*Csch[3*x],x]`

output `(Log[Sinh[x]^2]/3 - Log[3 + 4*Sinh[x]^2]/3)/2`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :=> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :=> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :=> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] :=> Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :=> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4856

```
Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

method	result	size
risch	$\frac{\ln(e^{2x}-1)}{3} - \frac{\ln(e^{4x}+e^{2x}+1)}{6}$	24

input

```
int(cosh(x)*csch(3*x),x,method=_RETURNVERBOSE)
```

output

```
1/3*ln(exp(2*x)-1)-1/6*ln(exp(4*x)+exp(2*x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(17) = 34.

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.48

$$\int \cosh(x) \operatorname{csch}(3x) dx = -\frac{1}{6} \log \left(\frac{2 \cosh(x)^2 + 2 \sinh(x)^2 + 1}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2} \right) + \frac{1}{3} \log \left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)} \right)$$

input

```
integrate(cosh(x)*csch(3*x),x, algorithm="fricas")
```

output

```
-1/6*log((2*cosh(x)^2 + 2*sinh(x)^2 + 1)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1/3*log(2*sinh(x)/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \cosh(x) \operatorname{csch}(3x) dx = \int \cosh(x) \operatorname{csch}(3x) dx$$

input `integrate(cosh(x)*csch(3*x),x)`

output `Integral(cosh(x)*csch(3*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(17) = 34$.

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.24

$$\begin{aligned} \int \cosh(x) \operatorname{csch}(3x) dx &= -\frac{1}{6} \log(e^{-x} + e^{-2x} + 1) + \frac{1}{3} \log(e^{-x} + 1) \\ &\quad + \frac{1}{3} \log(e^{-x} - 1) - \frac{1}{6} \log(-e^{-x} + e^{-2x} + 1) \end{aligned}$$

input `integrate(cosh(x)*csch(3*x),x, algorithm="maxima")`

output `-1/6*log(e^(-x) + e^(-2*x) + 1) + 1/3*log(e^(-x) + 1) + 1/3*log(e^(-x) - 1) - 1/6*log(-e^(-x) + e^(-2*x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(17) = 34$.

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.90

$$\begin{aligned} \int \cosh(x) \operatorname{csch}(3x) dx &= -\frac{1}{6} \log(e^{2x} + e^x + 1) - \frac{1}{6} \log(e^{2x} - e^x + 1) \\ &\quad + \frac{1}{3} \log(e^x + 1) + \frac{1}{3} \log(|e^x - 1|) \end{aligned}$$

input `integrate(cosh(x)*csch(3*x),x, algorithm="giac")`

output `-1/6*log(e^(2*x) + e^x + 1) - 1/6*log(e^(2*x) - e^x + 1) + 1/3*log(e^x + 1) + 1/3*log(abs(e^x - 1))`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int \cosh(x)\operatorname{csch}(3x) dx = \frac{\ln(3e^{2x} - 3)}{3} - \frac{\ln(-e^{2x} - e^{4x} - 1)}{6}$$

input `int(cosh(x)/sinh(3*x),x)`

output `log(3*exp(2*x) - 3)/3 - log(- exp(2*x) - exp(4*x) - 1)/6`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.14

$$\int \cosh(x)\operatorname{csch}(3x) dx = -\frac{\log(e^{2x} + e^x + 1)}{6} - \frac{\log(e^{2x} - e^x + 1)}{6} + \frac{\log(e^x - 1)}{3} + \frac{\log(e^x + 1)}{3}$$

input `int(cosh(x)*csch(3*x),x)`

output `(- log(e**(2*x) + e**x + 1) - log(e**(2*x) - e**x + 1) + 2*log(e**x - 1) + 2*log(e**x + 1))/6`

3.145 $\int \cosh(x) \operatorname{csch}(4x) dx$

Optimal result	1000
Mathematica [C] (verified)	1000
Rubi [A] (verified)	1001
Maple [B] (verified)	1002
Fricas [B] (verification not implemented)	1003
Sympy [F]	1003
Maxima [B] (verification not implemented)	1004
Giac [B] (verification not implemented)	1004
Mupad [B] (verification not implemented)	1005
Reduce [B] (verification not implemented)	1005

Optimal result

Integrand size = 7, antiderivative size = 26

$$\int \cosh(x) \operatorname{csch}(4x) dx = -\frac{1}{4} \operatorname{arctanh}(\cosh(x)) + \frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{2\sqrt{2}}$$

output

```
-1/4*arctanh(cosh(x))+1/4*arctanh(2^(1/2)*cosh(x))*2^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.58

$$\begin{aligned} \int \cosh(x) \operatorname{csch}(4x) dx = & \frac{1}{4} \left(\sqrt{2} \operatorname{arctanh} \left(\sqrt{2} - i \tanh \left(\frac{x}{2} \right) \right) \right. \\ & + \sqrt{2} \operatorname{arctanh} \left(\sqrt{2} + i \tanh \left(\frac{x}{2} \right) \right) - \log \left(\cosh \left(\frac{x}{2} \right) \right) \\ & \left. + \log \left(\sinh \left(\frac{x}{2} \right) \right) \right) \end{aligned}$$

input

```
Integrate[Cosh[x]*Csch[4*x],x]
```

output

$$\frac{(\text{Sqrt}[2] \cdot \text{ArcTanh}[\text{Sqrt}[2] - \text{I} \cdot \text{Tanh}[x/2]] + \text{Sqrt}[2] \cdot \text{ArcTanh}[\text{Sqrt}[2] + \text{I} \cdot \text{Tanh}[x/2]] - \text{Log}[\text{Cosh}[x/2]] + \text{Log}[\text{Sinh}[x/2]])}{4}$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 26, 4879, 1406, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(x) \operatorname{csch}(4x) dx \\ & \quad \downarrow 3042 \\ & \int \frac{i \cos(ix)}{\sin(4ix)} dx \\ & \quad \downarrow 26 \\ & i \int \frac{\cos(ix)}{\sin(4ix)} dx \\ & \quad \downarrow 4879 \\ & - \int \frac{1}{-8 \cosh^4(x) + 12 \cosh^2(x) - 4} d \cosh(x) \\ & \quad \downarrow 1406 \\ & 2 \int \frac{1}{4 - 8 \cosh^2(x)} d \cosh(x) - 2 \int \frac{1}{8 - 8 \cosh^2(x)} d \cosh(x) \\ & \quad \downarrow 219 \\ & \frac{\operatorname{arctanh}(\sqrt{2} \cosh(x))}{2\sqrt{2}} - \frac{1}{4} \operatorname{arctanh}(\cosh(x)) \end{aligned}$$

input

$$\text{Int}[\text{Cosh}[x] \cdot \text{Csch}[4 \cdot x], x]$$

output

$$-1/4 \cdot \text{ArcTanh}[\text{Cosh}[x]] + \text{ArcTanh}[\text{Sqrt}[2] \cdot \text{Cosh}[x]] / (2 \cdot \text{Sqrt}[2])$$

Defintions of rubi rules used

- rule 26 $\text{Int}[(\text{Complex}[0, a_])*(F x_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 219 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1406 $\text{Int}[(a_ + (b_)*(x_)^2 + (c_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c/q \ \text{Int}[1/(b/2 - q/2 + c*x^2), x], x] - \text{Simp}[c/q \ \text{Int}[1/(b/2 + q/2 + c*x^2), x], x]] /; \text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[b^2 - 4*a*c]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4879 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}\{d = \text{FreeFactors}[\text{Cos}[v], x]\}, -d/\text{Coefficient}[v, x, 1] \ \text{Subst}[\text{Int}[\text{SubstFor}[1, \text{Cos}[v]/d, u/\text{Sin}[v], x], x], x, \text{Cos}[v]/d]], x] /; \text{!FalseQ}[v] \ \&\& \ \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Cos}[v], x], u/\text{Sin}[v], x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(18) = 36$.

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.04

method	result	size
risch	$\frac{\ln(e^x-1)}{4} - \frac{\ln(e^x+1)}{4} + \frac{\sqrt{2} \ln(e^{2x}+\sqrt{2}e^x+1)}{8} - \frac{\sqrt{2} \ln(e^{2x}-\sqrt{2}e^x+1)}{8}$	53

input `int(cosh(x)*csch(4*x),x,method=_RETURNVERBOSE)`

output

```
1/4*ln(exp(x)-1)-1/4*ln(exp(x)+1)+1/8*2^(1/2)*ln(exp(2*x)+2^(1/2)*exp(x)+1)
)-1/8*2^(1/2)*ln(exp(2*x)-2^(1/2)*exp(x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(18) = 36.

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.08

$$\int \cosh(x) \operatorname{csch}(4x) dx = \frac{1}{8} \sqrt{2} \log \left(\frac{\cosh(x)^2 + \sinh(x)^2 + 2\sqrt{2} \cosh(x) + 2}{\cosh(x)^2 + \sinh(x)^2} \right) - \frac{1}{4} \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{4} \log(\cosh(x) + \sinh(x) - 1)$$

input

```
integrate(cosh(x)*csch(4*x),x, algorithm="fricas")
```

output

```
1/8*sqrt(2)*log((cosh(x)^2 + sinh(x)^2 + 2*sqrt(2)*cosh(x) + 2)/(cosh(x)^2
+ sinh(x)^2)) - 1/4*log(cosh(x) + sinh(x) + 1) + 1/4*log(cosh(x) + sinh(x)
) - 1)
```

Sympy [F]

$$\int \cosh(x) \operatorname{csch}(4x) dx = \int \cosh(x) \operatorname{csch}(4x) dx$$

input

```
integrate(cosh(x)*csch(4*x),x)
```

output

```
Integral(cosh(x)*csch(4*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(18) = 36$.

Time = 0.13 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.31

$$\int \cosh(x)\operatorname{csch}(4x) dx = \frac{1}{8} \sqrt{2} \log \left(\sqrt{2}e^{(-x)} + e^{(-2x)} + 1 \right) \\ - \frac{1}{8} \sqrt{2} \log \left(-\sqrt{2}e^{(-x)} + e^{(-2x)} + 1 \right) \\ - \frac{1}{4} \log \left(e^{(-x)} + 1 \right) + \frac{1}{4} \log \left(e^{(-x)} - 1 \right)$$

input `integrate(cosh(x)*csch(4*x),x, algorithm="maxima")`

output `1/8*sqrt(2)*log(sqrt(2)*e^(-x) + e^(-2*x) + 1) - 1/8*sqrt(2)*log(-sqrt(2)*e^(-x) + e^(-2*x) + 1) - 1/4*log(e^(-x) + 1) + 1/4*log(e^(-x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(18) = 36$.

Time = 0.18 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.19

$$\int \cosh(x)\operatorname{csch}(4x) dx = -\frac{1}{8} \sqrt{2} \log \left(-\frac{\sqrt{2} - e^{(-x)} - e^x}{\sqrt{2} + e^{(-x)} + e^x} \right) \\ - \frac{1}{8} \log \left(e^{(-x)} + e^x + 2 \right) + \frac{1}{8} \log \left(e^{(-x)} + e^x - 2 \right)$$

input `integrate(cosh(x)*csch(4*x),x, algorithm="giac")`

output `-1/8*sqrt(2)*log(-(sqrt(2) - e^(-x) - e^x)/(sqrt(2) + e^(-x) + e^x)) - 1/8*log(e^(-x) + e^x + 2) + 1/8*log(e^(-x) + e^x - 2)`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.35

$$\int \cosh(x) \operatorname{csch}(4x) dx = \frac{\ln\left(\frac{1}{2} - \frac{e^x}{2}\right)}{4} - \frac{\ln\left(-\frac{e^x}{2} - \frac{1}{2}\right)}{4} + \frac{\sqrt{2} \ln\left(-\frac{e^{2x}}{8} - \frac{\sqrt{2}e^x}{8} - \frac{1}{8}\right)}{8} - \frac{\sqrt{2} \ln\left(\frac{\sqrt{2}e^x}{8} - \frac{e^{2x}}{8} - \frac{1}{8}\right)}{8}$$

input `int(cosh(x)/sinh(4*x),x)`output `log(1/2 - exp(x)/2)/4 - log(- exp(x)/2 - 1/2)/4 + (2^(1/2)*log(- exp(2*x)/8 - (2^(1/2)*exp(x))/8 - 1/8))/8 - (2^(1/2)*log((2^(1/2)*exp(x))/8 - exp(2*x)/8 - 1/8))/8`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.08

$$\int \cosh(x) \operatorname{csch}(4x) dx = -\frac{\sqrt{2} \log(e^{2x} - e^x \sqrt{2} + 1)}{8} + \frac{\sqrt{2} \log(e^{2x} + e^x \sqrt{2} + 1)}{8} + \frac{\log(e^x - 1)}{4} - \frac{\log(e^x + 1)}{4}$$

input `int(cosh(x)*csch(4*x),x)`output `(- sqrt(2)*log(e**(2*x) - e**x*sqrt(2) + 1) + sqrt(2)*log(e**(2*x) + e**x*sqrt(2) + 1) + 2*log(e**x - 1) - 2*log(e**x + 1))/8`

3.146 $\int \cosh(x) \operatorname{csch}(5x) dx$

Optimal result	1006
Mathematica [A] (verified)	1006
Rubi [A] (verified)	1007
Maple [A] (verified)	1009
Fricas [B] (verification not implemented)	1009
Sympy [F]	1010
Maxima [F]	1010
Giac [A] (verification not implemented)	1011
Mupad [B] (verification not implemented)	1011
Reduce [F]	1012

Optimal result

Integrand size = 7, antiderivative size = 71

$$\int \cosh(x) \operatorname{csch}(5x) dx = \frac{1}{5} \log(\sinh(x)) - \frac{\log(5 - \sqrt{5} + 8 \sinh^2(x))}{\sqrt{5}(5 - \sqrt{5})} + \frac{\log(5 + \sqrt{5} + 8 \sinh^2(x))}{\sqrt{5}(5 + \sqrt{5})}$$

output

```
1/5*ln(sinh(x))-1/5*ln(5-5^(1/2)+8*sinh(x)^2)*5^(1/2)/(5-5^(1/2))+1/5*ln(5+5^(1/2)+8*sinh(x)^2)*5^(1/2)/(5+5^(1/2))
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.80

$$\int \cosh(x) \operatorname{csch}(5x) dx = \frac{1}{20} \left(- \left((1 + \sqrt{5}) \log(1 - \sqrt{5} + 4 \cosh(2x)) \right) + \left(-1 + \sqrt{5} \right) \log(1 + \sqrt{5} + 4 \cosh(2x)) + 4 \log(\sinh(x)) \right)$$

input

```
Integrate[Cosh[x]*Csch[5*x],x]
```

output

$$\frac{(-((1 + \sqrt{5}) \cdot \text{Log}[1 - \sqrt{5} + 4 \cdot \text{Cosh}[2 \cdot x]]) + (-1 + \sqrt{5}) \cdot \text{Log}[1 + \sqrt{5} + 4 \cdot \text{Cosh}[2 \cdot x]] + 4 \cdot \text{Log}[\text{Sinh}[x]])}{20}$$
Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.13, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 26, 4856, 26, 1434, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(x) \operatorname{csch}(5x) dx \\ & \quad \downarrow 3042 \\ & \int \frac{i \cos(ix)}{\sin(5ix)} dx \\ & \quad \downarrow 26 \\ & i \int \frac{\cos(ix)}{\sin(5ix)} dx \\ & \quad \downarrow 4856 \\ & i \int -\frac{i \operatorname{csch}(x)}{16 \sinh^4(x) + 20 \sinh^2(x) + 5} d \sinh(x) \\ & \quad \downarrow 26 \\ & \int \frac{\operatorname{csch}(x)}{16 \sinh^4(x) + 20 \sinh^2(x) + 5} d \sinh(x) \\ & \quad \downarrow 1434 \\ & \frac{1}{2} \int \frac{\operatorname{csch}(x)}{16 \sinh^4(x) + 20 \sinh^2(x) + 5} d \sinh^2(x) \\ & \quad \downarrow 1141 \\ & 8 \int \left(\frac{\operatorname{csch}(x)}{80} - \frac{1}{\sqrt{5} (5 - \sqrt{5}) (8 \sinh^2(x) - \sqrt{5} + 5)} + \frac{1}{\sqrt{5} (5 + \sqrt{5}) (8 \sinh^2(x) + \sqrt{5} + 5)} \right) d \sinh^2(x) \end{aligned}$$

$$8 \left(\frac{1}{80} \log(\sinh^2(x)) - \frac{\log(8 \sinh^2(x) - \sqrt{5} + 5)}{8\sqrt{5}(5 - \sqrt{5})} + \frac{\log(8 \sinh^2(x) + \sqrt{5} + 5)}{8\sqrt{5}(5 + \sqrt{5})} \right)$$

input `Int[Cosh[x]*Csch[5*x],x]`

output `8*(Log[Sinh[x]^2]/80 - Log[5 - Sqrt[5] + 8*Sinh[x]^2]/(8*Sqrt[5]*(5 - Sqrt[5])) + Log[5 + Sqrt[5] + 8*Sinh[x]^2]/(8*Sqrt[5]*(5 + Sqrt[5])))`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 1141 `Int[((d._) + (e._)*(x_)^(m._))*((a._) + (b._)*(x_) + (c._)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1434 `Int[(x_)^(m_)*((a_) + (b._)*(x_)^2 + (c._)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4856

```
Int[(u_)*(F_)[(c_)*((a_) + (b_)*(x_))], x_Symbol] := With[{d = FreeFactors[Sin[c*(a + b*x)], x]}, Simp[d/(b*c) Subst[Int[SubstFor[1, Sin[c*(a + b*x)]]/d, u, x], x], x, Sin[c*(a + b*x)]/d, x] /; FunctionOfQ[Sin[c*(a + b*x)]/d, u, x] /; FreeQ[{a, b, c}, x] && (EqQ[F, Cos] || EqQ[F, cos])
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.42

method	result
risch	$\frac{\ln(e^{2x}-1)}{5} - \frac{\ln(e^{4x} + (\frac{1}{2} + \frac{\sqrt{5}}{2})e^{2x} + 1)}{20} + \frac{\ln(e^{4x} + (\frac{1}{2} + \frac{\sqrt{5}}{2})e^{2x} + 1)\sqrt{5}}{20} - \frac{\ln(e^{4x} + (\frac{1}{2} - \frac{\sqrt{5}}{2})e^{2x} + 1)}{20} - \frac{\ln(e^{4x} + (\frac{1}{2} - \frac{\sqrt{5}}{2})e^{2x} + 1)}{20}$

input

```
int(cosh(x)*csch(5*x),x,method=_RETURNVERBOSE)
```

output

```
1/5*ln(exp(2*x)-1)-1/20*ln(exp(4*x)+(1/2+1/2*5^(1/2))*exp(2*x)+1)+1/20*ln(exp(4*x)+(1/2+1/2*5^(1/2))*exp(2*x)+1)*5^(1/2)-1/20*ln(exp(4*x)+(1/2-1/2*5^(1/2))*exp(2*x)+1)-1/20*ln(exp(4*x)+(1/2-1/2*5^(1/2))*exp(2*x)+1)*5^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 180 vs. 2(56) = 112.

Time = 0.09 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.54

$$\int \cosh(x) \operatorname{csch}(5x) dx$$

$$= \frac{1}{20} \sqrt{5} \log \left(\frac{4 \cosh(x)^4 + 4 \sinh(x)^4 + 4(\sqrt{5} + 1) \cosh(x)^2 + 4(6 \cosh(x)^2 + \sqrt{5} + 1) \sinh(x)^2 + \sqrt{5}}{2 \cosh(x)^4 + 2 \sinh(x)^4 + 2(6 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 1} \right)$$

$$- \frac{1}{20} \log \left(\frac{2 \cosh(x)^4 + 2 \sinh(x)^4 + 2(6 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 1}{\cosh(x)^4 - 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 - 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4} \right)$$

$$+ \frac{1}{5} \log \left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)} \right)$$

input

```
integrate(cosh(x)*csch(5*x),x, algorithm="fricas")
```

output

```
1/20*sqrt(5)*log((4*cosh(x)^4 + 4*sinh(x)^4 + 4*(sqrt(5) + 1)*cosh(x)^2 +
4*(6*cosh(x)^2 + sqrt(5) + 1)*sinh(x)^2 + sqrt(5) + 7)/(2*cosh(x)^4 + 2*si
nh(x)^4 + 2*(6*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 1)) - 1/20*log((2*
cosh(x)^4 + 2*sinh(x)^4 + 2*(6*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 1)
/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh
(x)^3 + sinh(x)^4)) + 1/5*log(2*sinh(x)/(cosh(x) - sinh(x)))
```

Sympy [F]

$$\int \cosh(x) \operatorname{csch}(5x) dx = \int \cosh(x) \operatorname{csch}(5x) dx$$

input

```
integrate(cosh(x)*csch(5*x),x)
```

output

```
Integral(cosh(x)*csch(5*x), x)
```

Maxima [F]

$$\int \cosh(x) \operatorname{csch}(5x) dx = \int \cosh(x) \operatorname{csch}(5x) dx$$

input

```
integrate(cosh(x)*csch(5*x),x, algorithm="maxima")
```

output

```
-1/5*integrate((e^(3*x) + e^(2*x) + e^x + 1)*e^x/(e^(4*x) + e^(3*x) + e^(2
*x) + e^x + 1), x) - 1/5*integrate((e^(3*x) - e^(2*x) + e^x - 1)*e^x/(e^(4
*x) - e^(3*x) + e^(2*x) - e^x + 1), x) + 3/10*integrate(e^(3*x)/(e^(4*x) +
e^(3*x) + e^(2*x) + e^x + 1), x) - 3/10*integrate(e^(3*x)/(e^(4*x) - e^(3
*x) + e^(2*x) - e^x + 1), x) + 1/10*integrate(e^(2*x)/(e^(4*x) + e^(3*x) +
e^(2*x) + e^x + 1), x) + 1/10*integrate(e^(2*x)/(e^(4*x) - e^(3*x) + e^(2
*x) - e^x + 1), x) - 1/10*integrate(e^x/(e^(4*x) + e^(3*x) + e^(2*x) + e^x
+ 1), x) + 1/10*integrate(e^x/(e^(4*x) - e^(3*x) + e^(2*x) - e^x + 1), x)
+ 1/5*log(e^x + 1) + 1/5*log(e^x - 1)
```

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.52

$$\int \cosh(x) \operatorname{csch}(5x) dx = -\frac{1}{20} (\sqrt{5} + 1) \log \left(\frac{1}{2} (\sqrt{5} + 1) e^x + e^{(2x)} + 1 \right) \\ - \frac{1}{20} (\sqrt{5} + 1) \log \left(-\frac{1}{2} (\sqrt{5} + 1) e^x + e^{(2x)} + 1 \right) \\ + \frac{1}{20} (\sqrt{5} - 1) \log \left(\frac{1}{2} (\sqrt{5} - 1) e^x + e^{(2x)} + 1 \right) \\ + \frac{1}{20} (\sqrt{5} - 1) \log \left(-\frac{1}{2} (\sqrt{5} - 1) e^x + e^{(2x)} + 1 \right) \\ + \frac{1}{5} \log(e^x + 1) + \frac{1}{5} \log(|e^x - 1|)$$

input `integrate(cosh(x)*csch(5*x),x, algorithm="giac")`output `-1/20*(sqrt(5) + 1)*log(1/2*(sqrt(5) + 1)*e^x + e^(2*x) + 1) - 1/20*(sqrt(5) + 1)*log(-1/2*(sqrt(5) + 1)*e^x + e^(2*x) + 1) + 1/20*(sqrt(5) - 1)*log(1/2*(sqrt(5) - 1)*e^x + e^(2*x) + 1) + 1/20*(sqrt(5) - 1)*log(-1/2*(sqrt(5) - 1)*e^x + e^(2*x) + 1) + 1/5*log(e^x + 1) + 1/5*log(abs(e^x - 1))`**Mupad [B] (verification not implemented)**

Time = 1.06 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.46

$$\int \cosh(x) \operatorname{csch}(5x) dx = \frac{\ln(5 - 5e^{2x})}{5} - \ln \left(2e^{4x} - e^{2x} \right. \\ \left. + \left(\frac{\sqrt{5}}{20} + \frac{1}{20} \right) (30e^{4x} - 20e^{2x} + 30) + 2 \right) \left(\frac{\sqrt{5}}{20} + \frac{1}{20} \right) \\ + \ln \left(2e^{4x} - e^{2x} - \left(\frac{\sqrt{5}}{20} - \frac{1}{20} \right) (30e^{4x} - 20e^{2x} + 30) \right. \\ \left. + 2 \right) \left(\frac{\sqrt{5}}{20} - \frac{1}{20} \right)$$

input `int(cosh(x)/sinh(5*x),x)`

output

```
log(5 - 5*exp(2*x))/5 - log(2*exp(4*x) - exp(2*x) + (5^(1/2)/20 + 1/20)*(3
0*exp(4*x) - 20*exp(2*x) + 30) + 2)*(5^(1/2)/20 + 1/20) + log(2*exp(4*x) -
exp(2*x) - (5^(1/2)/20 - 1/20)*(30*exp(4*x) - 20*exp(2*x) + 30) + 2)*(5^(
1/2)/20 - 1/20)
```

Reduce [F]

$$\int \cosh(x)\operatorname{csch}(5x) dx = \int \cosh(x) \operatorname{csch}(5x) dx$$

input

```
int(cosh(x)*csch(5*x),x)
```

output

```
int(cosh(x)*csch(5*x),x)
```

3.147 $\int \cosh(x) \operatorname{csch}(6x) dx$

Optimal result	1013
Mathematica [C] (verified)	1013
Rubi [A] (verified)	1014
Maple [B] (verified)	1016
Fricas [B] (verification not implemented)	1016
Sympy [F]	1017
Maxima [F]	1017
Giac [B] (verification not implemented)	1017
Mupad [B] (verification not implemented)	1018
Reduce [B] (verification not implemented)	1018

Optimal result

Integrand size = 7, antiderivative size = 36

$$\int \cosh(x) \operatorname{csch}(6x) dx = -\frac{1}{6} \operatorname{arctanh}(\cosh(x)) - \frac{1}{6} \operatorname{arctanh}(2 \cosh(x)) + \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{2\sqrt{3}}$$

output

`-1/6*arctanh(cosh(x))-1/6*arctanh(2*cosh(x))+1/6*arctanh(2/3*cosh(x)*3^(1/2))*3^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.53

$$\int \cosh(x) \operatorname{csch}(6x) dx = \frac{1}{12} \left(2\sqrt{3} \operatorname{arctanh}\left(\frac{2 - i \tanh\left(\frac{x}{2}\right)}{\sqrt{3}}\right) + 2\sqrt{3} \operatorname{arctanh}\left(\frac{2 + i \tanh\left(\frac{x}{2}\right)}{\sqrt{3}}\right) - 2 \log\left(\cosh\left(\frac{x}{2}\right)\right) + \log(1 - 2 \cosh(x)) - \log(1 + 2 \cosh(x)) + 2 \log\left(\sinh\left(\frac{x}{2}\right)\right) \right)$$

input `Integrate[Cosh[x]*Csch[6*x],x]`

output $(2\sqrt{3}\operatorname{ArcTanh}[(2 - I\tanh[x/2])/\sqrt{3}] + 2\sqrt{3}\operatorname{ArcTanh}[(2 + I\tanh[x/2])/\sqrt{3}] - 2\operatorname{Log}[\operatorname{Cosh}[x/2]] + \operatorname{Log}[1 - 2\operatorname{Cosh}[x]] - \operatorname{Log}[1 + 2\operatorname{Cosh}[x]] + 2\operatorname{Log}[\operatorname{Sinh}[x/2]])/12$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 26, 4879, 27, 2460, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(x)\operatorname{csch}(6x) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{i \cos(ix)}{\sin(6ix)} dx \\
 & \quad \downarrow 26 \\
 & i \int \frac{\cos(ix)}{\sin(6ix)} dx \\
 & \quad \downarrow 4879 \\
 & - \int \frac{1}{2(-16 \cosh^6(x) + 32 \cosh^4(x) - 19 \cosh^2(x) + 3)} d \cosh(x) \\
 & \quad \downarrow 27 \\
 & - \frac{1}{2} \int \frac{1}{-16 \cosh^6(x) + 32 \cosh^4(x) - 19 \cosh^2(x) + 3} d \cosh(x) \\
 & \quad \downarrow 2460 \\
 & - \frac{1}{2} \int \left(\frac{2}{4 \cosh^2(x) - 3} - \frac{2}{3(4 \cosh^2(x) - 1)} - \frac{1}{3(\cosh^2(x) - 1)} \right) d \cosh(x) \\
 & \quad \downarrow 2009
 \end{aligned}$$

$$\frac{1}{2} \left(-\frac{1}{3} \operatorname{arctanh}(\cosh(x)) - \frac{1}{3} \operatorname{arctanh}(2 \cosh(x)) + \frac{\operatorname{arctanh}\left(\frac{2 \cosh(x)}{\sqrt{3}}\right)}{\sqrt{3}} \right)$$

input `Int[Cosh[x]*Csch[6*x],x]`

output `(-1/3*ArcTanh[Cosh[x]] - ArcTanh[2*Cosh[x]]/3 + ArcTanh[(2*Cosh[x])/Sqrt[3]]/Sqrt[3])/2`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2460 `Int[(u_.)*(Px_)^(p_), x_Symbol] :> With[{Qx = Factor[Px /. x -> Sqrt[x]]}, Int[ExpandIntegrand[u*(Qx /. x -> x^2)^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x^2] && GtQ[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0] && RationalFunctionQ[u, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4879 `Int[u_, x_Symbol] :> With[{v = FunctionOfTrig[u, x]}, Simp[With[{d = FreeFactors[Cos[v], x]}, -d/Coefficient[v, x, 1] Subst[Int[SubstFor[1, Cos[v]/d, u/Sin[v], x], x], x, Cos[v]/d]], x] /; !FalseQ[v] && FunctionOfQ[NonfreeFactors[Cos[v], x], u/Sin[v], x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(26) = 52$.

Time = 0.13 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.14

method	result	size
risch	$-\frac{\ln(e^x+1)}{6} + \frac{\ln(e^x-1)}{6} + \frac{\sqrt{3} \ln(e^{2x}+\sqrt{3}e^x+1)}{12} - \frac{\sqrt{3} \ln(e^{2x}-\sqrt{3}e^x+1)}{12} + \frac{\ln(e^{2x}-e^x+1)}{12} - \frac{\ln(e^{2x}+e^x+1)}{12}$	77

input `int(cosh(x)*csch(6*x),x,method=_RETURNVERBOSE)`

output
$$-1/6*\ln(\exp(x)+1)+1/6*\ln(\exp(x)-1)+1/12*3^{(1/2)}*\ln(\exp(2*x)+3^{(1/2)}*\exp(x)+1)-1/12*3^{(1/2)}*\ln(\exp(2*x)-3^{(1/2)}*\exp(x)+1)+1/12*\ln(\exp(2*x)-\exp(x)+1)-1/12*\ln(\exp(2*x)+\exp(x)+1)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(26) = 52$.

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.81

$$\int \cosh(x)\operatorname{csch}(6x) dx = \frac{1}{12} \sqrt{3} \log \left(\frac{2 \cosh(x)^2 + 2 \sinh(x)^2 + 4 \sqrt{3} \cosh(x) + 5}{2 \cosh(x)^2 + 2 \sinh(x)^2 - 1} \right) - \frac{1}{12} \log \left(\frac{2 \cosh(x) + 1}{\cosh(x) - \sinh(x)} \right) + \frac{1}{12} \log \left(\frac{2 \cosh(x) - 1}{\cosh(x) - \sinh(x)} \right) - \frac{1}{6} \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{6} \log(\cosh(x) + \sinh(x) - 1)$$

input `integrate(cosh(x)*csch(6*x),x, algorithm="fricas")`

output
$$1/12*\sqrt{3}*\log((2*\cosh(x)^2 + 2*\sinh(x)^2 + 4*\sqrt{3}*\cosh(x) + 5)/(2*\cosh(x)^2 + 2*\sinh(x)^2 - 1)) - 1/12*\log((2*\cosh(x) + 1)/(\cosh(x) - \sinh(x))) + 1/12*\log((2*\cosh(x) - 1)/(\cosh(x) - \sinh(x))) - 1/6*\log(\cosh(x) + \sinh(x) + 1) + 1/6*\log(\cosh(x) + \sinh(x) - 1)$$

Sympy [F]

$$\int \cosh(x) \operatorname{csch}(6x) dx = \int \cosh(x) \operatorname{csch}(6x) dx$$

input `integrate(cosh(x)*csch(6*x),x)`

output `Integral(cosh(x)*csch(6*x), x)`

Maxima [F]

$$\int \cosh(x) \operatorname{csch}(6x) dx = \int \cosh(x) \operatorname{csch}(6x) dx$$

input `integrate(cosh(x)*csch(6*x),x, algorithm="maxima")`

output `-integrate(1/2*(e^(3*x) - e^x)/(e^(4*x) - e^(2*x) + 1), x) - 1/12*log(e^(2*x) + e^x + 1) + 1/12*log(e^(2*x) - e^x + 1) - 1/6*log(e^x + 1) + 1/6*log(e^x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 79 vs. $2(26) = 52$.

Time = 0.13 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.19

$$\begin{aligned} \int \cosh(x) \operatorname{csch}(6x) dx = & -\frac{1}{12} \sqrt{3} \log \left(-\frac{\sqrt{3} - e^{(-x)} - e^x}{\sqrt{3} + e^{(-x)} + e^x} \right) \\ & - \frac{1}{12} \log(e^{(-x)} + e^x + 2) - \frac{1}{12} \log(e^{(-x)} + e^x + 1) \\ & + \frac{1}{12} \log(e^{(-x)} + e^x - 1) + \frac{1}{12} \log(e^{(-x)} + e^x - 2) \end{aligned}$$

input `integrate(cosh(x)*csch(6*x),x, algorithm="giac")`

output

```
-1/12*sqrt(3)*log(-(sqrt(3) - e^(-x) - e^x)/(sqrt(3) + e^(-x) + e^x)) - 1/
12*log(e^(-x) + e^x + 2) - 1/12*log(e^(-x) + e^x + 1) + 1/12*log(e^(-x) +
e^x - 1) + 1/12*log(e^(-x) + e^x - 2)
```

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.53

$$\int \cosh(x) \operatorname{csch}(6x) dx = \frac{\ln\left(\frac{1}{3} - \frac{e^x}{3}\right)}{6} - \frac{\ln\left(-\frac{e^x}{3} - \frac{1}{3}\right)}{6} - \frac{\ln\left(-\frac{e^{2x}}{36} - \frac{e^x}{36} - \frac{1}{36}\right)}{12}$$

$$+ \frac{\ln\left(\frac{e^x}{36} - \frac{e^{2x}}{36} - \frac{1}{36}\right)}{12} + \frac{\sqrt{3} \ln\left(-\frac{e^{2x}}{12} - \frac{\sqrt{3}e^x}{12} - \frac{1}{12}\right)}{12}$$

$$- \frac{\sqrt{3} \ln\left(\frac{\sqrt{3}e^x}{12} - \frac{e^{2x}}{12} - \frac{1}{12}\right)}{12}$$

input

```
int(cosh(x)/sinh(6*x),x)
```

output

```
log(1/3 - exp(x)/3)/6 - log(- exp(x)/3 - 1/3)/6 - log(- exp(2*x)/36 - exp(
x)/36 - 1/36)/12 + log(exp(x)/36 - exp(2*x)/36 - 1/36)/12 + (3^(1/2)*log(-
exp(2*x)/12 - (3^(1/2)*exp(x))/12 - 1/12))/12 - (3^(1/2)*log((3^(1/2)*exp
(x))/12 - exp(2*x)/12 - 1/12))/12
```

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.28

$$\int \cosh(x) \operatorname{csch}(6x) dx = -\frac{\sqrt{3} \log(e^{2x} - e^x \sqrt{3} + 1)}{12} + \frac{\sqrt{3} \log(e^{2x} + e^x \sqrt{3} + 1)}{12}$$

$$- \frac{\log(e^{2x} + e^x + 1)}{12} + \frac{\log(e^{2x} - e^x + 1)}{12}$$

$$+ \frac{\log(e^x - 1)}{6} - \frac{\log(e^x + 1)}{6}$$

input

```
int(cosh(x)*csch(6*x),x)
```

output

```
( - sqrt(3)*log(e**(2*x) - e**x*sqrt(3) + 1) + sqrt(3)*log(e**(2*x) + e**x
*sqrt(3) + 1) - log(e**(2*x) + e**x + 1) + log(e**(2*x) - e**x + 1) + 2*lo
g(e**x - 1) - 2*log(e**x + 1))/12
```

3.148 $\int \cosh(a + bx) \operatorname{csch}(c + bx) dx$

Optimal result	1020
Mathematica [A] (verified)	1020
Rubi [C] (verified)	1021
Maple [B] (verified)	1022
Fricas [B] (verification not implemented)	1023
Sympy [F]	1023
Maxima [B] (verification not implemented)	1023
Giac [A] (verification not implemented)	1024
Mupad [B] (verification not implemented)	1024
Reduce [B] (verification not implemented)	1025

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \frac{\cosh(a - c) \log(\sinh(c + bx))}{b} + x \sinh(a - c)$$

output `cosh(a-c)*ln(sinh(b*x+c))/b+x*sinh(a-c)`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \frac{\cosh(a - c) \log(\sinh(c + bx))}{b} + x \sinh(a - c)$$

input `Integrate[Cosh[a + b*x]*Csch[c + b*x],x]`

output `(Cosh[a - c]*Log[Sinh[c + b*x]])/b + x*Sinh[a - c]`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6159, 24, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{csch}(bx + c) dx \\
 & \quad \downarrow \text{6159} \\
 & \cosh(a - c) \int \coth(c + bx) dx + \sinh(a - c) \int 1 dx \\
 & \quad \downarrow \text{24} \\
 & \cosh(a - c) \int \coth(c + bx) dx + x \sinh(a - c) \\
 & \quad \downarrow \text{3042} \\
 & x \sinh(a - c) + \cosh(a - c) \int -i \tan\left(ic + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & x \sinh(a - c) - i \cosh(a - c) \int \tan\left(\frac{1}{2}(2ic + \pi) + ibx\right) dx \\
 & \quad \downarrow \text{3956} \\
 & x \sinh(a - c) + \frac{\cosh(a - c) \log(-i \sinh(bx + c))}{b}
 \end{aligned}$$

input `Int[Cosh[a + b*x]*Csch[c + b*x],x]`

output `(Cosh[a - c]*Log[(-I)*Sinh[c + b*x]])/b + x*Sinh[a - c]`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6159 `Int[Cosh[v_]*Csch[w_]^(n_.), x_Symbol] := Simp[Cosh[v - w] Int[Coth[w]*Csch[w]^(n - 1), x], x] + Simp[Sinh[v - w] Int[Csch[w]^(n - 1), x], x] /; GTQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(26) = 52$.

Time = 0.14 (sec) , antiderivative size = 152, normalized size of antiderivative = 5.85

method	result
risch	$x e^{a-c} - e^{-a-c} e^{2a} x - e^{-a-c} e^{2c} x - \frac{e^{-a-c} e^{2a} a}{b} - \frac{e^{-a-c} e^{2c} a}{b} + \frac{\ln(e^{2bx+2a} - e^{2a-2c}) e^{-a-c} e^{2a}}{2b} + \frac{\ln(e^{2bx+2a} - e^{2a-2c})}{2b}$

input `int(cosh(b*x+a)*csch(b*x+c),x,method=_RETURNVERBOSE)`

output `x*exp(a-c)-exp(-a-c)*exp(2*a)*x-exp(-a-c)*exp(2*c)*x-1/b*exp(-a-c)*exp(2*a)*a-1/b*exp(-a-c)*exp(2*c)*a+1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-a-c)*exp(2*a)+1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-a-c)*exp(2*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(26) = 52$.

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.35

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \frac{2bx - (\cosh(-a + c))^2 - 2\cosh(-a + c)\sinh(-a + c) + \sinh(-a + c)^2 + 1}{2(b\cosh(-a + c) - b\sinh(-a + c))} \log\left(\frac{2\sinh(bx + c)}{\cosh(bx + c) - \sinh(bx + c)}\right)$$

input `integrate(cosh(b*x+a)*csch(b*x+c),x, algorithm="fricas")`

output `-1/2*(2*b*x - (cosh(-a + c))^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c)))/(b*cosh(-a + c) - b*sinh(-a + c))`

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \int \cosh(a + bx) \operatorname{csch}(bx + c) dx$$

input `integrate(cosh(b*x+a)*csch(b*x+c),x)`

output `Integral(cosh(a + b*x)*csch(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(26) = 52$.

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.08

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} + \frac{(e^{(2a)} + e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} + \frac{(bx + a)e^{(a-c)}}{b}$$

input `integrate(cosh(b*x+a)*csch(b*x+c),x, algorithm="maxima")`

output `1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) + e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + (b*x + a)*e^(a - c)/b`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.81

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = -xe^{(-a+c)} + \frac{(e^{(2a+c)} + e^{(3c)})e^{(-a-2c)} \log(|e^{(2bx+2c)} - 1|)}{2b}$$

input `integrate(cosh(b*x+a)*csch(b*x+c),x, algorithm="giac")`

output `-x*e^(-a + c) + 1/2*(e^(2*a + c) + e^(3*c))*e^(-a - 2*c)*log(abs(e^(2*b*x + 2*c) - 1))/b`

Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.54

$$\int \cosh(a + bx) \operatorname{csch}(c + bx) dx = \frac{e^{2c-2a} \ln(e^{2a} e^{2bx} - e^{2a} e^{-2c}) (2b e^{3a-3c} + 2b e^{a-c})}{4b^2} - x e^{c-a}$$

input `int(cosh(a + b*x)/sinh(c + b*x),x)`

output $(\exp(2*c - 2*a)*\log(\exp(2*a)*\exp(2*b*x) - \exp(2*a)*\exp(-2*c))*(2*b*\exp(3*a - 3*c) + 2*b*\exp(a - c))/(4*b^2) - x*\exp(c - a)$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.31

$$\int \cosh(a + bx)\operatorname{csch}(c + bx) dx$$

$$= \frac{e^{2a}\log(e^{bx+c} - 1) + e^{2a}\log(e^{bx+c} + 1) + e^{2c}\log(e^{bx+c} - 1) + e^{2c}\log(e^{bx+c} + 1) - 2e^{2c}bx}{2e^{a+cb}}$$

input `int(cosh(b*x+a)*csch(b*x+c),x)`

output $(e^{2*a}*\log(e^{b*x + c} - 1) + e^{2*a}*\log(e^{b*x + c} + 1) + e^{2*c}*\log(e^{b*x + c} - 1) + e^{2*c}*\log(e^{b*x + c} + 1) - 2*e^{2*c}*b*x)/(2*e^{(a + c)*b})$

3.149 $\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx$

Optimal result	1026
Mathematica [C] (verified)	1026
Rubi [A] (verified)	1027
Maple [B] (verified)	1029
Fricas [B] (verification not implemented)	1029
Sympy [F]	1030
Maxima [B] (verification not implemented)	1031
Giac [B] (verification not implemented)	1031
Mupad [B] (verification not implemented)	1032
Reduce [B] (verification not implemented)	1032

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = -\frac{\cosh(a - c) \operatorname{csch}(c + bx)}{b} - \frac{\operatorname{arctanh}(\cosh(c + bx)) \sinh(a - c)}{b}$$

output

```
-cosh(a-c)*csch(b*x+c)/b-arctanh(cosh(b*x+c))*sinh(a-c)/b
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.50

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = -\frac{\cosh(a - c) \operatorname{csch}(c + bx)}{b} - \frac{2i \operatorname{arctan} \left(\frac{(\cosh(c) - \sinh(c)) \left(\cosh(c) \cosh\left(\frac{bx}{2}\right) + \sinh(c) \sinh\left(\frac{bx}{2}\right) \right)}{i \cosh(c) \cosh\left(\frac{bx}{2}\right) - i \cosh\left(\frac{bx}{2}\right) \sinh(c)} \right) \sinh(a - c)}{b}$$

input `Integrate[Cosh[a + b*x]*Csch[c + b*x]^2,x]`

output
$$-\left(\frac{\text{Cosh}[a - c] \cdot \text{Csch}[c + b*x]}{b}\right) - \left(\frac{(2*I) \cdot \text{ArcTan}\left[\frac{(\text{Cosh}[c] - \text{Sinh}[c]) \cdot (\text{Cosh}[c] \cdot \text{Cosh}[(b*x)/2] + \text{Sinh}[c] \cdot \text{Sinh}[(b*x)/2])}{I \cdot \text{Cosh}[c] \cdot \text{Cosh}[(b*x)/2] - I \cdot \text{Cosh}[(b*x)/2] \cdot \text{Sinh}[c]}\right]}{\text{Sinh}[a - c]}\right)}{b}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {6159, 3042, 26, 3086, 24, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh(a + bx) \operatorname{csch}^2(bx + c) dx \\ & \quad \downarrow \text{6159} \\ & \sinh(a - c) \int \operatorname{csch}(c + bx) dx + \cosh(a - c) \int \operatorname{coth}(c + bx) \operatorname{csch}(c + bx) dx \\ & \quad \downarrow \text{3042} \\ & \sinh(a - c) \int i \operatorname{csc}(ic + ibx) dx + \cosh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{26} \\ & i \sinh(a - c) \int \operatorname{csc}(ic + ibx) dx + \cosh(a - c) \int \sec\left(ic + ibx - \frac{\pi}{2}\right) \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3086} \\ & i \sinh(a - c) \int \operatorname{csc}(ic + ibx) dx - \frac{i \cosh(a - c) \int 1 d(-i \operatorname{csch}(c + bx))}{b} \\ & \quad \downarrow \text{24} \\ & -\frac{\cosh(a - c) \operatorname{csch}(bx + c)}{b} + i \sinh(a - c) \int \operatorname{csc}(ic + ibx) dx \\ & \quad \downarrow \text{4257} \end{aligned}$$

$$-\frac{\sinh(a-c)\operatorname{arctanh}(\cosh(bx+c))}{b} - \frac{\cosh(a-c)\operatorname{csch}(bx+c)}{b}$$

input `Int[Cosh[a + b*x]*Csch[c + b*x]^2,x]`

output `-((Cosh[a - c]*Csch[c + b*x])/b) - (ArcTanh[Cosh[c + b*x]]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 26 `Int[(Complex[0, a_]*(F_x_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])`

rule 4257 `Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6159 `Int[Cosh[v_]*Csch[w_]^(n_), x_Symbol] := Simp[Cosh[v - w] Int[Coth[w]*Csch[w]^(n-1), x], x] + Simp[Sinh[v - w] Int[Csch[w]^(n-1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(36) = 72$.

Time = 0.32 (sec) , antiderivative size = 170, normalized size of antiderivative = 4.72

method	result
risch	$\frac{e^{bx+a}(e^{2a}+e^{2c})}{b(-e^{2bx+2a+2c}+e^{2a})} - \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{2b} + \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2c}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{2b} - \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2c}}{2b}$

input `int(cosh(b*x+a)*csch(b*x+c)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{b} \frac{\exp(bx+a)(\exp(2a)+\exp(2c))}{(-\exp(2bx+2a+2c)+\exp(2a))-1/2 \ln(\exp(bx+a)+\exp(a-c))} + \frac{1}{b} \frac{\exp(-a-c)\exp(2a)+1/2 \ln(\exp(bx+a)+\exp(a-c))}{\exp(-a-c)\exp(2c)+1/2 \ln(\exp(bx+a)-\exp(a-c))} - \frac{1}{b} \frac{\exp(-a-c)\exp(2c)+1/2 \ln(\exp(bx+a)-\exp(a-c))}{\exp(-a-c)\exp(2a)-1/2 \ln(\exp(bx+a)-\exp(a-c))} + \frac{1}{b} \frac{\exp(-a-c)\exp(2a)-1/2 \ln(\exp(bx+a)-\exp(a-c))}{\exp(-a-c)\exp(2c)-1/2 \ln(\exp(bx+a)+\exp(a-c))}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 617 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 617, normalized size of antiderivative = 17.14

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)*csch(b*x+c)^2,x, algorithm="fricas")`

output

```

1/2*(4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - 2*cosh(b*x + c)*sinh(-a +
c)^2 - 2*(cosh(-a + c)^2 + 1)*cosh(b*x + c) - ((cosh(-a + c)^2 - 1)*cosh(
b*x + c)^2 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^
2 - 1)*sinh(b*x + c)^2 + (cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a +
c)^2 - 2*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-
a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c) - 2*(cosh(b*x
+ c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) + 1)*log(cosh(b*x + c) +
sinh(b*x + c) + 1) + ((cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (cosh(-a + c
)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^2 +
(cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(2*cosh(b*x + c)
*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^
2 - 1)*cosh(b*x + c))*sinh(b*x + c) - 2*(cosh(b*x + c)^2*cosh(-a + c) - co
sh(-a + c))*sinh(-a + c) + 1)*log(cosh(b*x + c) + sinh(b*x + c) - 1) - 2*(
cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 + 1)*sinh(b*
x + c))/(b*cosh(b*x + c)^2*cosh(-a + c) + (b*cosh(-a + c) - b*sinh(-a + c)
)*sinh(b*x + c)^2 - b*cosh(-a + c) + 2*(b*cosh(b*x + c)*cosh(-a + c) - b*c
osh(b*x + c)*sinh(-a + c))*sinh(b*x + c) - (b*cosh(b*x + c)^2 - b)*sinh(-a
+ c))

```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = \int \cosh(a + bx) \operatorname{csch}^2(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)**2,x)
```

output

```
Integral(cosh(a + b*x)*csch(b*x + c)**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 105, normalized size of antiderivative = 2.92

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = -\frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{2b} + \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{2b} + \frac{(e^{(2a)} + e^{(2c)})e^{(-bx-a)}}{b(e^{(-2bx)} - e^{(2c)})}$$

input `integrate(cosh(b*x+a)*csch(b*x+c)^2,x, algorithm="maxima")`

output `-1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + (e^(2*a) + e^(2*c))*e^(-b*x - a)/(b*(e^(-2*b*x) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 111, normalized size of antiderivative = 3.08

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx = -\frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(bx+c)} + 1)}{2b} + \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+c)} - 1|)}{2b} - \frac{(e^{(bx+2a)} + e^{(bx+2c)})e^{(-a)}}{b(e^{(2bx+2c)} - 1)}$$

input `integrate(cosh(b*x+a)*csch(b*x+c)^2,x, algorithm="giac")`

output `-1/2*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(e^(b*x + c) + 1)/b + 1/2*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + c) - 1))/b - (e^(b*x + 2*a) + e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) - 1))`

Mupad [B] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.33

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx$$

$$= \frac{\operatorname{atan}\left(\frac{e^{-a} e^{2c} e^{bx} (\sqrt{-b^2 - e^{2a}} e^{-2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{2c} (e^{4a} e^{-4c} - 2e^{2a} e^{-2c} + 1)}}\right) \sqrt{e^{2c-2a} (e^{4a-4c} - 2e^{2a-2c} + 1)}}{\sqrt{-b^2}} + \frac{e^{a+bx} (e^{2a-2c} + 1)}{b (e^{2a-2c} - e^{2a+2bx})}$$

input `int(cosh(a + b*x)/sinh(c + b*x)^2,x)`output `(atan((exp(-a)*exp(2*c)*exp(b*x)*((-b^2)^(1/2) - exp(2*a)*exp(-2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(2*c)*(exp(4*a)*exp(-4*c) - 2*exp(2*a)*exp(-2*c) + 1))^(1/2)))*(exp(2*c - 2*a)*(exp(4*a - 4*c) - 2*exp(2*a - 2*c) + 1))^(1/2))/(-b^2)^(1/2) + (exp(a + b*x)*(exp(2*a - 2*c) + 1))/(b*(exp(2*a - 2*c) - exp(2*a + 2*b*x)))`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 208, normalized size of antiderivative = 5.78

$$\int \cosh(a + bx) \operatorname{csch}^2(c + bx) dx$$

$$= \frac{e^{2bx+2a+2c} \log(e^{bx+c} - 1) - e^{2bx+2a+2c} \log(e^{bx+c} + 1) - e^{2bx+4c} \log(e^{bx+c} - 1) + e^{2bx+4c} \log(e^{bx+c} + 1) - 2e^{a+c} b (e^{2bx+2a+2c} - 1)}{2e^{a+c} b (e^{2bx+2a+2c} - 1)}$$

input `int(cosh(b*x+a)*csch(b*x+c)^2,x)`output `(e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) - e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) + 1) - e**(2*b*x + 4*c)*log(e**(b*x + c) - 1) + e**(2*b*x + 4*c)*log(e**(b*x + c) + 1) - 2*e**(2*a + b*x + c) - 2*e**(b*x + 3*c) - e**(2*a)*log(e**(b*x + c) - 1) + e**(2*a)*log(e**(b*x + c) + 1) + e**(2*c)*log(e**(b*x + c) - 1) - e**(2*c)*log(e**(b*x + c) + 1))/(2*e**(a + c)*b*(e**(2*b*x + 2*c) - 1))`

3.150 $\int \cosh(a + bx)\operatorname{csch}^3(c + bx) dx$

Optimal result	1033
Mathematica [A] (verified)	1033
Rubi [A] (verified)	1034
Maple [A] (verified)	1036
Fricas [B] (verification not implemented)	1036
Sympy [F]	1037
Maxima [B] (verification not implemented)	1037
Giac [A] (verification not implemented)	1038
Mupad [F(-1)]	1038
Reduce [B] (verification not implemented)	1039

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \cosh(a + bx)\operatorname{csch}^3(c + bx) dx = -\frac{\cosh(a - c)\operatorname{csch}^2(c + bx)}{2b} - \frac{\operatorname{coth}(c + bx)\sinh(a - c)}{b}$$

output

$$-1/2*\cosh(a-c)*\operatorname{csch}(b*x+c)^2/b-\operatorname{coth}(b*x+c)*\sinh(a-c)/b$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\begin{aligned} &\int \cosh(a + bx)\operatorname{csch}^3(c + bx) dx \\ &= -\frac{\operatorname{csch}(c)\operatorname{csch}^2(c + bx)(\sinh(a) - \cosh(c + 2bx)\sinh(a - c))}{2b} \end{aligned}$$

input

$$\text{Integrate}[\text{Cosh}[a + b*x]*\text{Csch}[c + b*x]^3,x]$$

output

$$-1/2*(\text{Csch}[c]*\text{Csch}[c + b*x]^2*(\text{Sinh}[a] - \text{Cosh}[c + 2*b*x]*\text{Sinh}[a - c]))/b$$

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {6159, 3042, 25, 26, 3086, 15, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{csch}^3(bx + c) dx \\
 & \quad \downarrow \text{6159} \\
 & \sinh(a - c) \int \operatorname{csch}^2(c + bx) dx + \cosh(a - c) \int \coth(c + bx) \operatorname{csch}^2(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sinh(a - c) \int -\operatorname{csc}(ic + ibx)^2 dx + \cosh(a - c) \int i \sec\left(ic + ibx - \frac{\pi}{2}\right)^2 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & \cosh(a - c) \int i \sec\left(ic + ibx - \frac{\pi}{2}\right)^2 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx - \sinh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{26} \\
 & i \cosh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^2 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx - \sinh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{3086} \\
 & \frac{\cosh(a - c) \int -i \operatorname{csch}(c + bx) d(-i \operatorname{csch}(c + bx))}{b} - \sinh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{\cosh(a - c) \operatorname{csch}^2(bx + c)}{2b} - \sinh(a - c) \int \operatorname{csc}(ic + ibx)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -\frac{\cosh(a - c) \operatorname{csch}^2(bx + c)}{2b} - \frac{i \sinh(a - c) \int 1 d(-i \coth(c + bx))}{b} \\
 & \quad \downarrow \text{24}
 \end{aligned}$$

$$-\frac{\cosh(a-c)\operatorname{csch}^2(bx+c)}{2b} - \frac{\sinh(a-c)\operatorname{coth}(bx+c)}{b}$$

input `Int[Cosh[a + b*x]*Csch[c + b*x]^3,x]`

output `-1/2*(Cosh[a - c]*Csch[c + b*x]^2)/b - (Coth[c + b*x]*Sinh[a - c])/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_)), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

rule 6159

```
Int[Cosh[v_]*Csch[w_]^(n_), x_Symbol] := Simp[Cosh[v - w] Int[Coth[w]*Csch[w]^(n - 1), x], x] + Simp[Sinh[v - w] Int[Csch[w]^(n - 1), x], x] /; GtQ[n, 0] && NeQ[w, v] && FreeQ[v - w, x]
```

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

method	result	size
parallelrisch	$-\frac{\operatorname{sech}\left(\frac{bx}{2} + \frac{c}{2}\right)^2 \operatorname{csch}\left(\frac{bx}{2} + \frac{c}{2}\right)^2 \cosh(2bx+a+c)}{8b}$	36
risch	$\frac{(-2e^{2bx+2a+2c} + e^{2a} - e^{2c})e^{3a-c}}{(-e^{2bx+2a+2c} + e^{2a})^2 b}$	59

input

```
int(cosh(b*x+a)*csch(b*x+c)^3,x,method=_RETURNVERBOSE)
```

output

```
-1/8/b*sech(1/2*b*x+1/2*c)^2*csch(1/2*b*x+1/2*c)^2*cosh(2*b*x+a+c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(37) = 74$.

Time = 0.09 (sec) , antiderivative size = 243, normalized size of antiderivative = 6.23

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx =$$

$$-\frac{b \cosh(bx + c)^3 \cosh(-a + c)^2 - b \cosh(bx + c) \cosh(-a + c)^2 + (b \cosh(-a + c)^2 - b \sinh(-a + c))}{\dots}$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)^3,x, algorithm="fricas")
```

output

```
-2*(cosh(b*x + c)*cosh(-a + c) + (cosh(-a + c) - 2*sinh(-a + c))*sinh(b*x
+ c))/(b*cosh(b*x + c)^3*cosh(-a + c)^2 - b*cosh(b*x + c)*cosh(-a + c)^2 +
(b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^3 + 3*(b*cosh(b*x + c
)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c)^2 - (b*co
sh(b*x + c)^3 - b*cosh(b*x + c))*sinh(-a + c)^2 + 3*(b*cosh(b*x + c)^2*cos
h(-a + c)^2 - b*cosh(-a + c)^2 - (b*cosh(b*x + c)^2 - b)*sinh(-a + c)^2)*s
inh(b*x + c))
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx = \int \cosh(a + bx) \operatorname{csch}^3(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)**3,x)
```

output

```
Integral(cosh(a + b*x)*csch(b*x + c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(37) = 74$.

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 3.38

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx = \frac{2e^{(-2bx+3c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})} + \frac{e^{(2a+3c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})} - \frac{e^{(5c)}}{b(2e^{(-2bx+a+2c)} - e^{(-4bx+a)} - e^{(a+4c)})}$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)^3,x, algorithm="maxima")
```

output

```
2*e^(-2*b*x + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c))) + e^(2*a + 3*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c))) - e^(5*c)/(b*(2*e^(-2*b*x + a + 2*c) - e^(-4*b*x + a) - e^(a + 4*c)))
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.31

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx = -\frac{(2e^{(2bx+2a+2c)} - e^{(2a)} + e^{(2c)})e^{(-a-c)}}{b(e^{(2bx+2c)} - 1)^2}$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)^3,x, algorithm="giac")
```

output

```
-(2*e^(2*b*x + 2*a + 2*c) - e^(2*a) + e^(2*c))*e^(-a - c)/(b*(e^(2*b*x + 2*c) - 1)^2)
```

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx = \int \frac{\cosh(a + bx)}{\sinh(c + bx)^3} dx$$

input

```
int(cosh(a + b*x)/sinh(c + b*x)^3,x)
```

output

```
int(cosh(a + b*x)/sinh(c + b*x)^3, x)
```

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.38

$$\int \cosh(a + bx) \operatorname{csch}^3(c + bx) dx = -\frac{e^c (e^{4bx+2a+2c} + 1)}{e^{ab} (e^{4bx+4c} - 2e^{2bx+2c} + 1)}$$

input `int(cosh(b*x+a)*csch(b*x+c)^3,x)`

output `(- e**c*(e**(2*a + 4*b*x + 2*c) + 1))/(e**a*b*(e**(4*b*x + 4*c) - 2*e**(2*b*x + 2*c) + 1))`

3.151 $\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx$

Optimal result	1040
Mathematica [A] (verified)	1040
Rubi [C] (verified)	1041
Maple [B] (verified)	1044
Fricas [B] (verification not implemented)	1044
Sympy [F]	1045
Maxima [B] (verification not implemented)	1046
Giac [B] (verification not implemented)	1046
Mupad [F(-1)]	1047
Reduce [B] (verification not implemented)	1047

Optimal result

Integrand size = 15, antiderivative size = 61

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx = -\frac{\cosh(a - c) \operatorname{csch}^3(c + bx)}{3b} + \frac{1}{2} \left(\frac{\operatorname{arctanh}(\cosh(c + bx))}{b} - \frac{\operatorname{coth}(c + bx) \operatorname{csch}(c + bx)}{b} \right) \sinh(a - c)$$

output

```
-1/3*cosh(a-c)*csch(b*x+c)^3/b+1/2*(arctanh(cosh(b*x+c))/b-coth(b*x+c)*csch(b*x+c)/b)*sinh(a-c)
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx = \frac{12 \operatorname{arctanh}(\cosh(c) + \sinh(c) \tanh(\frac{bx}{2})) \sinh(a - c) - \operatorname{csch}^3(c + bx) (4 \cosh(a - c) + 3 \sinh(a - c) \sinh(2c))}{12b}$$

input

```
Integrate[Cosh[a + b*x]*Csch[c + b*x]^4,x]
```

output

```
(12*ArcTanh[Cosh[c] + Sinh[c]*Tanh[(b*x)/2]]*Sinh[a - c] - Csch[c + b*x]^3
*(4*Cosh[a - c] + 3*Sinh[a - c]*Sinh[2*(c + b*x)]))/(12*b)
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.15, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {6159, 3042, 25, 26, 3086, 15, 4255, 26, 3042, 26, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cosh(a + bx) \operatorname{csch}^4(bx + c) dx \\
 & \quad \downarrow \text{6159} \\
 & \sinh(a - c) \int \operatorname{csch}^3(c + bx) dx + \cosh(a - c) \int \coth(c + bx) \operatorname{csch}^3(c + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \sinh(a - c) \int -i \csc(ic + ibx)^3 dx + \cosh(a - c) \int -\sec\left(ic + ibx - \frac{\pi}{2}\right)^3 \tan\left(ic + ibx - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & \sinh(a - c) \int -i \csc(ic + ibx)^3 dx - \cosh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^3 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx \\
 & \quad \downarrow \text{26} \\
 & -i \sinh(a - c) \int \csc(ic + ibx)^3 dx - \cosh(a - c) \int \sec\left(\frac{1}{2}(2ic - \pi) + ibx\right)^3 \tan\left(\frac{1}{2}(2ic - \pi) + ibx\right) dx \\
 & \quad \downarrow \text{3086} \\
 & \frac{i \cosh(a - c) \int -\operatorname{csch}^2(c + bx) d(-i \operatorname{csch}(c + bx))}{b} - i \sinh(a - c) \int \csc(ic + ibx)^3 dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 15 \\
& -\frac{\cosh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i\sinh(a-c) \int \csc(ic+ibx)^3 dx \\
& \downarrow 4255 \\
& c) \left(\frac{1}{2} \int -i\operatorname{csch}(c+bx) dx - \frac{i\coth(bx+c)\operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 26 \\
& c) \left(-\frac{1}{2}i \int \operatorname{csch}(c+bx) dx - \frac{i\coth(bx+c)\operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 3042 \\
& c) \left(-\frac{1}{2}i \int i\csc(ic+ibx) dx - \frac{i\coth(bx+c)\operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 26 \\
& -\frac{\cosh(a-c)\operatorname{csch}^3(bx+c)}{3b} - i\sinh(a-c) \left(\frac{1}{2} \int \csc(ic+ibx) dx - \frac{i\coth(bx+c)\operatorname{csch}(bx+c)}{2b} \right) \\
& \downarrow 4257 \\
& c) \left(\frac{i\operatorname{arctanh}(\cosh(bx+c))}{2b} - \frac{i\coth(bx+c)\operatorname{csch}(bx+c)}{2b} \right)
\end{aligned}$$

input

```
Int[Cosh[a + b*x]*Csch[c + b*x]^4,x]
```

output

```
-1/3*(Cosh[a - c]*Csch[c + b*x]^3)/b - I*(((I/2)*ArcTanh[Cosh[c + b*x]])/b
- ((I/2)*Coth[c + b*x]*Csch[c + b*x])/b)*Sinh[a - c]
```

Defintions of rubi rules used

- rule 15 $\text{Int}[(a_.)(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[a*(x^{(m+1)})/(m+1), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \ \text{Int}[Fx, x], x]$
- rule 26 $\text{Int}[(\text{Complex}[0, a_])*(Fx_), x_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \ \text{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3086 $\text{Int}[(a_.)*\text{sec}[(e_.) + (f_.)(x_)]^{(m_.)}*((b_.)*\text{tan}[(e_.) + (f_.)(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a/f \ \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e+f*x]], x] \text{ ; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$
- rule 4255 $\text{Int}[(\text{csc}[(c_.) + (d_.)(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c+d*x]*((b*\text{Csc}[c+d*x])^{(n-1)})/(d*(n-1)), x] + \text{Simp}[b^2*((n-2)/(n-1)) \ \text{Int}[(b*\text{Csc}[c+d*x])^{(n-2)}, x], x] \text{ ; FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 4257 $\text{Int}[\text{csc}[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c+d*x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$
- rule 6159 $\text{Int}[\text{Cosh}[v_]*\text{Csch}[w_]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Cosh}[v-w] \ \text{Int}[\text{Coth}[w]*\text{Csch}[w]^{(n-1)}, x], x] + \text{Simp}[\text{Sinh}[v-w] \ \text{Int}[\text{Csch}[w]^{(n-1)}, x], x] \text{ ; GtQ}[n, 0] \ \&\& \ \text{NeQ}[w, v] \ \&\& \ \text{FreeQ}[v-w, x]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. $2(57) = 114$.

Time = 2.04 (sec) , antiderivative size = 235, normalized size of antiderivative = 3.85

method	result
risch	$-\frac{e^{bx+a}(-3e^{4bx+6a+4c}+3e^{4bx+4a+6c}-8e^{2bx+6a+2c}-8e^{2bx+4a+4c}+3e^{6a}-3e^{4a+2c})}{6b(-e^{2bx+2a+2c}+e^{2a})^3} + \frac{\ln(e^{bx+a}+e^{a-c})e^{-a-c}e^{2a}}{4b} - \frac{\ln(e^{bx+a}-e^{a-c})e^{-a-c}e^{2a}}{4b}$

input

```
int(cosh(b*x+a)*csch(b*x+c)^4,x,method=_RETURNVERBOSE)
```

output

```
-1/6*exp(b*x+a)*(-3*exp(4*b*x+6*a+4*c)+3*exp(4*b*x+4*a+6*c)-8*exp(2*b*x+6*a+2*c)-8*exp(2*b*x+4*a+4*c)+3*exp(6*a)-3*exp(4*a+2*c))/b/(-exp(2*b*x+2*a+2*c)+exp(2*a))^3+1/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*a)-1/4*ln(exp(b*x+a)+exp(a-c))/b*exp(-a-c)*exp(2*c)-1/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*a)+1/4*ln(exp(b*x+a)-exp(a-c))/b*exp(-a-c)*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2696 vs. $2(57) = 114$.

Time = 0.10 (sec) , antiderivative size = 2696, normalized size of antiderivative = 44.20

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)^4,x, algorithm="fricas")
```

output

```
-1/12*(6*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^5 + 6*(cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^5 - 30*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^4 + 16*(cosh(-a + c)^2 + 1)*cosh(b*x + c)^3 + 4*(15*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (15*cosh(b*x + c)^2 + 4)*sinh(-a + c)^2 + 4*cosh(-a + c)^2 - 2*(15*cosh(b*x + c)^2*cosh(-a + c) + 4*cosh(-a + c))*sinh(-a + c) + 4)*sinh(b*x + c)^3 + 12*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 + 4*cosh(b*x + c))*sinh(-a + c)^2 + 4*(cosh(-a + c)^2 + 1)*cosh(b*x + c) - 2*(5*cosh(b*x + c)^3*cosh(-a + c) + 4*cosh(b*x + c)*cosh(-a + c))*sinh(-a + c))*sinh(b*x + c)^2 + 2*(3*cosh(b*x + c)^5 + 8*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 - 6*(cosh(-a + c)^2 - 1)*cosh(b*x + c) - 3*((cosh(-a + c)^2 - 1)*cosh(b*x + c)^6 + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 - 1)*sinh(b*x + c)^6 - 6*(2*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 - 1)*cosh(b*x + c))*sinh(b*x + c)^5 - 3*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^4 + 3*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (5*cosh(b*x + c)^2 - 1)*sinh(-a + c)^2 - cosh(-a + c)^2 - 2*(5*cosh(b*x + c)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c)^4 + 4*(5*(cosh(-a + c)^2 - 1)*cosh(b*x + c)^3 + (5*cosh(b*x + c)^3 - 3*cosh(b*x + c))*sinh(-a + c)^2 - 3*(cosh(-a + c)^2 - 1)*cosh(b*...
```

SymPy [F]

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx = \int \cosh(a + bx) \operatorname{csch}^4(bx + c) dx$$

input

```
integrate(cosh(b*x+a)*csch(b*x+c)**4,x)
```

output

```
Integral(cosh(a + b*x)*csch(b*x + c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(57) = 114$.

Time = 0.05 (sec) , antiderivative size = 191, normalized size of antiderivative = 3.13

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx$$

$$= \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} + e^c)}{4b} - \frac{(e^{(2a)} - e^{(2c)})e^{(-a-c)} \log(e^{(-bx)} - e^c)}{4b}$$

$$+ \frac{3(e^{(2a+4c)} - e^{(6c)})e^{(-bx-a)} + 8(e^{(4a+2c)} + e^{(2a+4c)})e^{(-3bx-3a)} - 3(e^{(6a)} - e^{(4a+2c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+4c)} - 3e^{(-4bx+2c)} + e^{(-6bx)} - e^{(6c)})}$$

input `integrate(cosh(b*x+a)*csch(b*x+c)^4,x, algorithm="maxima")`

output

```
1/4*(e^(2*a) - e^(2*c))*e^(-a - c)*log(e^(-b*x) + e^c)/b - 1/4*(e^(2*a) -
e^(2*c))*e^(-a - c)*log(e^(-b*x) - e^c)/b + 1/6*(3*(e^(2*a + 4*c) - e^(6*c)
))*e^(-b*x - a) + 8*(e^(4*a + 2*c) + e^(2*a + 4*c))*e^(-3*b*x - 3*a) - 3*(
e^(6*a) - e^(4*a + 2*c))*e^(-5*b*x - 5*a))/(b*(3*e^(-2*b*x + 4*c) - 3*e^(-
4*b*x + 2*c) + e^(-6*b*x) - e^(6*c)))
```

Giacc [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. $2(57) = 114$.

Time = 0.17 (sec) , antiderivative size = 165, normalized size of antiderivative = 2.70

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx$$

$$= \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(e^{(bx+c)} + 1)}{4b} - \frac{(e^{(2a+c)} - e^{(3c)})e^{(-a-2c)} \log(|e^{(bx+c)} - 1|)}{4b}$$

$$- \frac{(3e^{(5bx+2a+4c)} - 3e^{(5bx+6c)} + 8e^{(3bx+2a+2c)} + 8e^{(3bx+4c)} - 3e^{(bx+2a)} + 3e^{(bx+2c)})e^{(-a)}}{6b(e^{(2bx+2c)} - 1)^3}$$

input `integrate(cosh(b*x+a)*csch(b*x+c)^4,x, algorithm="giac")`

output

```
1/4*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(e^(b*x + c) + 1)/b - 1/4*(e^(2*a + c) - e^(3*c))*e^(-a - 2*c)*log(abs(e^(b*x + c) - 1))/b - 1/6*(3*e^(5*b*x + 2*a + 4*c) - 3*e^(5*b*x + 6*c) + 8*e^(3*b*x + 2*a + 2*c) + 8*e^(3*b*x + 4*c) - 3*e^(b*x + 2*a) + 3*e^(b*x + 2*c))*e^(-a)/(b*(e^(2*b*x + 2*c) - 1)^3)
```

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx = \int \frac{\cosh(a + bx)}{\sinh(c + bx)^4} dx$$

input

```
int(cosh(a + b*x)/sinh(c + b*x)^4,x)
```

output

```
int(cosh(a + b*x)/sinh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 478, normalized size of antiderivative = 7.84

$$\int \cosh(a + bx) \operatorname{csch}^4(c + bx) dx$$

$$= \frac{-3e^{6bx+2a+6c} \log(e^{bx+c} - 1) + 3e^{6bx+2a+6c} \log(e^{bx+c} + 1) + 3e^{6bx+8c} \log(e^{bx+c} - 1) - 3e^{6bx+8c} \log(e^{bx+c} + 1)}{b}$$

input

```
int(cosh(b*x+a)*csch(b*x+c)^4,x)
```


output

```
( - 3*e**(2*a + 6*b*x + 6*c)*log(e**(b*x + c) - 1) + 3*e**(2*a + 6*b*x + 6*c)*log(e**(b*x + c) + 1) + 3*e**(6*b*x + 8*c)*log(e**(b*x + c) - 1) - 3*e**(6*b*x + 8*c)*log(e**(b*x + c) + 1) - 6*e**(2*a + 5*b*x + 5*c) + 6*e**(5*b*x + 7*c) + 9*e**(2*a + 4*b*x + 4*c)*log(e**(b*x + c) - 1) - 9*e**(2*a + 4*b*x + 4*c)*log(e**(b*x + c) + 1) - 9*e**(4*b*x + 6*c)*log(e**(b*x + c) - 1) + 9*e**(4*b*x + 6*c)*log(e**(b*x + c) + 1) - 16*e**(2*a + 3*b*x + 3*c) - 16*e**(3*b*x + 5*c) - 9*e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) + 9*e**(2*a + 2*b*x + 2*c)*log(e**(b*x + c) + 1) + 9*e**(2*b*x + 4*c)*log(e**(b*x + c) - 1) - 9*e**(2*b*x + 4*c)*log(e**(b*x + c) + 1) + 6*e**(2*a + b*x + c) - 6*e**(b*x + 3*c) + 3*e**(2*a)*log(e**(b*x + c) - 1) - 3*e**(2*a)*log(e**(b*x + c) + 1) - 3*e**(2*c)*log(e**(b*x + c) - 1) + 3*e**(2*c)*log(e**(b*x + c) + 1))/(12*e**(a + c)*b*(e**(6*b*x + 6*c) - 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) - 1))
```

3.152 $\int \cosh(a + bx) \operatorname{csch}(c - bx) dx$

Optimal result	1049
Mathematica [A] (verified)	1049
Rubi [F]	1050
Maple [B] (verified)	1050
Fricas [B] (verification not implemented)	1051
Sympy [F]	1051
Maxima [B] (verification not implemented)	1052
Giac [A] (verification not implemented)	1052
Mupad [B] (verification not implemented)	1053
Reduce [B] (verification not implemented)	1053

Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = -\frac{\cosh(a + c) \log(\sinh(c - bx))}{b} - x \sinh(a + c)$$

output

```
-cosh(a+c)*ln(-sinh(b*x-c))/b-x*sinh(a+c)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.12

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = -\frac{\cosh(a + c) \log(-\sinh(c - bx)) + bx \sinh(a + c)}{b}$$

input

```
Integrate[Cosh[a + b*x]*Csch[c - b*x],x]
```

output

```
-((Cosh[a + c]*Log[-Sinh[c - b*x]] + b*x*Sinh[a + c])/b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx$$

input `Int[Cosh[a + b*x]*Csch[c - b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(28) = 56$.

Time = 0.11 (sec) , antiderivative size = 147, normalized size of antiderivative = 5.88

method	result
risch	$-x e^{a+c} + e^{-a-c} e^{2a+2c} x + \frac{e^{-a-c} e^{2a+2c} a}{b} + x e^{-a-c} + \frac{e^{-a-c} a}{b} - \frac{\ln(-e^{2a+2c} + e^{2bx+2a}) e^{-a-c} e^{2a+2c}}{2b} - \frac{\ln(-e^{2a+2c} + e^{2bx+2a})}{2b}$

input `int(-cosh(b*x+a)*csch(b*x-c),x,method=_RETURNVERBOSE)`

output `-x*exp(a+c)+exp(-a-c)*exp(2*a+2*c)*x+1/b*exp(-a-c)*exp(2*a+2*c)*a+x*exp(-a-c)+1/b*exp(-a-c)*a-1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)*exp(2*a+2*c)-1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-a-c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(28) = 56$.

Time = 0.10 (sec) , antiderivative size = 146, normalized size of antiderivative = 5.84

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx$$

$$= \frac{2bx \cosh(a + c)^2 - 4bx \cosh(a + c) \sinh(a + c) + 2bx \sinh(a + c)^2 - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log(2(\cosh(a + c) \sinh(bx + a) - \cosh(bx + a) \sinh(a + c)) / (\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c)))}{2(b \cosh(a + c) - b \sinh(a + c))}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c),x, algorithm="fricas")`

output `1/2*(2*b*x*cosh(a + c)^2 - 4*b*x*cosh(a + c)*sinh(a + c) + 2*b*x*sinh(a + c)^2 - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 + 1)*log(2*(cosh(a + c)*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c)))/b*cosh(a + c) - b*sinh(a + c))`

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = - \int \cosh(a + bx) \operatorname{csch}(bx - c) dx$$

input `integrate(-cosh(b*x+a)*csch(b*x-c),x)`

output `-Integral(cosh(a + b*x)*csch(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. $2(28) = 56$.

Time = 0.05 (sec) , antiderivative size = 81, normalized size of antiderivative = 3.24

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = -\frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{2b} - \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{2b} - \frac{(bx + a)e^{(a+c)}}{b}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c),x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(-b*x + c) + 1)/b - 1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(e^(-b*x + c) - 1)/b - (b*x + a)*e^(a + c)/b`

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.92

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = xe^{(-a-c)} - \frac{(e^{(2a+2c)} + 1)e^{(-a-c)} \log(|e^{(2bx)} - e^{(2c)}|)}{2b}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c),x, algorithm="giac")`

output `x*e^(-a - c) - 1/2*(e^(2*a + 2*c) + 1)*e^(-a - c)*log(abs(e^(2*b*x) - e^(2*c)))/b`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.60

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = x e^{-a-c} - \frac{e^{-2a-2c} \ln(e^{2a} e^{2bx} - e^{2a} e^{2c}) (2b e^{3a+3c} + 2b e^{a+c})}{4b^2}$$

input `int(cosh(a + b*x)/sinh(c - b*x),x)`output `x*exp(- a - c) - (exp(- 2*a - 2*c)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp(2*c))*(2*b*exp(3*a + 3*c) + 2*b*exp(a + c)))/(4*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.48

$$\int \cosh(a + bx) \operatorname{csch}(c - bx) dx = \frac{-e^{2a+2c} \log(e^{bx} + e^c) - e^{2a+2c} \log(e^{bx} - e^c) - \log(e^{bx} + e^c) - \log(e^{bx} - e^c) + 2bx}{2e^{a+cb}}$$

input `int(-cosh(b*x+a)*csch(b*x-c),x)`output `(- e**(2*a + 2*c)*log(e**(b*x) + e**c) - e**(2*a + 2*c)*log(e**(b*x) - e**c) - log(e**(b*x) + e**c) - log(e**(b*x) - e**c) + 2*b*x)/(2*e**(a + c)*b)`

3.153 $\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx$

Optimal result	1054
Mathematica [C] (verified)	1054
Rubi [F]	1055
Maple [B] (verified)	1055
Fricas [B] (verification not implemented)	1056
Sympy [F]	1057
Maxima [B] (verification not implemented)	1057
Giac [B] (verification not implemented)	1058
Mupad [B] (verification not implemented)	1058
Reduce [B] (verification not implemented)	1059

Optimal result

Integrand size = 16, antiderivative size = 33

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = \frac{\cosh(a + c) \operatorname{csch}(c - bx)}{b} - \frac{\operatorname{arctanh}(\cosh(c - bx)) \sinh(a + c)}{b}$$

output

```
-cosh(a+c)*csch(b*x-c)/b-arctanh(cosh(b*x-c))*sinh(a+c)/b
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 149, normalized size of antiderivative = 4.52

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = \frac{\cosh(a + c) \operatorname{csch}(c - bx)}{b} - \frac{2i \operatorname{arctan} \left(\frac{\cosh\left(\frac{bx}{2}\right) + \cosh^2(c) \cosh\left(\frac{bx}{2}\right) + 2 \cosh(c) \cosh\left(\frac{bx}{2}\right) \sinh(c) + \cosh\left(\frac{bx}{2}\right) \sinh^2(c) + \sinh\left(\frac{bx}{2}\right) - \cosh^2(c) \sinh\left(\frac{bx}{2}\right) - 2 \cosh(c) \sinh\left(\frac{bx}{2}\right)}{2i \cosh(c) \cosh\left(\frac{bx}{2}\right) + 2i \cosh\left(\frac{bx}{2}\right) \sinh(c)} \right)}{b}$$

input

```
Integrate[Cosh[a + b*x]*Csch[c - b*x]^2,x]
```

output

```
(Cosh[a + c]*Csch[c - b*x])/b - ((2*I)*ArcTan[(Cosh[(b*x)/2] + Cosh[c]^2*Cosh[(b*x)/2] + 2*Cosh[c]*Cosh[(b*x)/2]*Sinh[c] + Cosh[(b*x)/2]*Sinh[c]^2 + Sinh[(b*x)/2] - Cosh[c]^2*Sinh[(b*x)/2] - 2*Cosh[c]*Sinh[c]*Sinh[(b*x)/2] - Sinh[c]^2*Sinh[(b*x)/2])/((2*I)*Cosh[c]*Cosh[(b*x)/2] + (2*I)*Cosh[(b*x)/2]*Sinh[c])]*Sinh[a + c])/b
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx$$

input

```
Int[Cosh[a + b*x]*Csch[c - b*x]^2,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(36) = 72$.

Time = 0.21 (sec) , antiderivative size = 164, normalized size of antiderivative = 4.97

method	result
risch	$\frac{e^{bx+a}(e^{2a+2c}+1)}{b(e^{2a+2c}-e^{2bx+2a})} + \frac{\ln(-e^{a+c}+e^{bx+a})e^{-a-c}e^{2a+2c}}{2b} - \frac{\ln(-e^{a+c}+e^{bx+a})e^{-a-c}}{2b} - \frac{\ln(e^{a+c}+e^{bx+a})e^{-a-c}e^{2a+2c}}{2b} + \frac{\ln(e^{a+c}+e^{bx+a})e^{-a-c}}{2b}$

input

```
int(cosh(b*x+a)*csch(b*x-c)^2,x,method=_RETURNVERBOSE)
```


output

```
1/b*exp(b*x+a)*(exp(2*a+2*c)+1)/(exp(2*a+2*c)-exp(2*b*x+2*a))+1/2*ln(-exp(
a+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)-1/2*ln(-exp(a+c)+exp(b*x+a))/b*e
xp(-a-c)-1/2*ln(exp(a+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)+1/2*ln(exp(a
+c)+exp(b*x+a))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1104 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 1104, normalized size of antiderivative = 33.45

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*csch(b*x-c)^2,x, algorithm="fricas")
```

output

```
-1/2*(6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^2 - 2*cosh(b*x + a)*sinh(a +
c)^3 - 2*(3*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c) + 2*(cosh(a + c)
^3 + cosh(a + c))*cosh(b*x + a) + (4*cosh(b*x + a)^2*cosh(a + c)*sinh(a +
c)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 - cosh(a + c)^2)*cos
h(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + sinh(a + c)^
4 + (6*cosh(a + c)^2 - 1)*sinh(a + c)^2 - cosh(a + c)^2 - 2*(2*cosh(a + c)
^3 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*cosh(a + c)^2 - 1)*co
sh(b*x + a)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(4*cosh(b*x + a)*cosh
(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 - (6*cosh(a + c)^2 - 1)
)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3 - cosh(a + c))*cosh(b*x
+ a)*sinh(a + c) - (cosh(a + c)^4 - cosh(a + c)^2)*cosh(b*x + a)*sinh(b*
x + a) + 2*((2*cosh(a + c)^3 - cosh(a + c))*cosh(b*x + a)^2 - cosh(a + c))
*sinh(a + c) - 1)*log(cosh(b*x + a)*cosh(a + c) + (cosh(a + c) - sinh(a +
c))*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c) + 1) - (4*cosh(b*x + a)^2*co
sh(a + c)*sinh(a + c)^3 - cosh(b*x + a)^2*sinh(a + c)^4 - (cosh(a + c)^4 -
cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a +
c)^3 + sinh(a + c)^4 + (6*cosh(a + c)^2 - 1)*sinh(a + c)^2 - cosh(a + c)^2
- 2*(2*cosh(a + c)^3 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((6*co
sh(a + c)^2 - 1)*cosh(b*x + a)^2 - 1)*sinh(a + c)^2 + cosh(a + c)^2 + 2*(4
*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4 ...
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = \int \cosh(a + bx) \operatorname{csch}^2(bx - c) dx$$

input `integrate(cosh(b*x+a)*csch(b*x-c)**2,x)`

output `Integral(cosh(a + b*x)*csch(b*x - c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 103, normalized size of antiderivative = 3.12

$$\begin{aligned} \int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = & -\frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{2b} \\ & + \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{2b} \\ & + \frac{(e^{(2a+2c)} + 1)e^{(-bx-a)}}{b(e^{(-2bx+2c)} - 1)} \end{aligned}$$

input `integrate(cosh(b*x+a)*csch(b*x-c)^2,x, algorithm="maxima")`

output `-1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(-b*x + c) + 1)/b + 1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(-b*x + c) - 1)/b + (e^(2*a + 2*c) + 1)*e^(-b*x - a)/(b*(e^(-2*b*x + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.18

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = -\frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(bx)} + e^c)}{2b} + \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(|e^{(bx)} - e^c|)}{2b} - \frac{(e^{(bx+2a+2c)} + e^{(bx)})e^{(-a)}}{b(e^{(2bx)} - e^{(2c)})}$$

input `integrate(cosh(b*x+a)*csch(b*x-c)^2,x, algorithm="giac")`

output `-1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(b*x) + e^c)/b + 1/2*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(abs(e^(b*x) - e^c))/b - (e^(b*x + 2*a + 2*c) + e^(b*x))*e^(-a)/(b*(e^(2*b*x) - e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.73

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx = \frac{\operatorname{atan}\left(\frac{e^{-a} e^{-2c} e^{bx} (\sqrt{-b^2} - e^{2a} e^{2c} \sqrt{-b^2})}{b \sqrt{e^{-2a} e^{-2c} (e^{4a} e^{4c} - 2e^{2a} e^{2c} + 1)}}\right) \sqrt{e^{-2a-2c} (e^{4a+4c} - 2e^{2a+2c} + 1)}}{\sqrt{-b^2}} + \frac{e^{a+bx} (e^{2a+2c} + 1)}{b (e^{2a+2c} - e^{2a+2bx})}$$

input `int(cosh(a + b*x)/sinh(c - b*x)^2,x)`

output

```
(atan((exp(-a)*exp(-2*c)*exp(b*x)*((-b^2)^(1/2) - exp(2*a)*exp(2*c)*(-b^2)^(1/2)))/(b*(exp(-2*a)*exp(-2*c)*(exp(4*a)*exp(4*c) - 2*exp(2*a)*exp(2*c) + 1))^(1/2)))*(exp(- 2*a - 2*c)*(exp(4*a + 4*c) - 2*exp(2*a + 2*c) + 1))^(1/2))/(-b^2)^(1/2) + (exp(a + b*x)*(exp(2*a + 2*c) + 1))/(b*(exp(2*a + 2*c) - exp(2*a + 2*b*x)))
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 218, normalized size of antiderivative = 6.61

$$\int \cosh(a + bx) \operatorname{csch}^2(c - bx) dx$$

$$= \frac{-e^{2bx+2a+2c} \log(e^{bx} + e^c) + e^{2bx+2a+2c} \log(e^{bx} - e^c) + e^{2bx} \log(e^{bx} + e^c) - e^{2bx} \log(e^{bx} - e^c) - 2e^{bx+2a+3c}}{2e^{a+cb}(e^{2bx} - e^{2c})}$$

input

```
int(cosh(b*x+a)*csch(b*x-c)^2,x)
```

output

```
( - e**(2*a + 2*b*x + 2*c)*log(e**(b*x) + e**c) + e**(2*a + 2*b*x + 2*c)*log(e**(b*x) - e**c) + e**(2*b*x)*log(e**(b*x) + e**c) - e**(2*b*x)*log(e**(b*x) - e**c) - 2*e**(2*a + b*x + 3*c) - 2*e**(b*x + c) + e**(2*a + 4*c)*log(e**(b*x) + e**c) - e**(2*a + 4*c)*log(e**(b*x) - e**c) - e**(2*c)*log(e**(b*x) + e**c) + e**(2*c)*log(e**(b*x) - e**c))/(2*e**(a + c)*b*(e**(2*b*x) - e**(2*c)))
```

3.154 $\int \cosh(a + bx)\operatorname{csch}^3(c - bx) dx$

Optimal result	1060
Mathematica [A] (verified)	1060
Rubi [F]	1061
Maple [A] (verified)	1061
Fricas [B] (verification not implemented)	1062
Sympy [F]	1062
Maxima [B] (verification not implemented)	1063
Giac [A] (verification not implemented)	1063
Mupad [F(-1)]	1064
Reduce [B] (verification not implemented)	1064

Optimal result

Integrand size = 16, antiderivative size = 37

$$\int \cosh(a + bx)\operatorname{csch}^3(c - bx) dx = \frac{\cosh(a + c)\operatorname{csch}^2(c - bx)}{2b} - \frac{\operatorname{coth}(c - bx)\sinh(a + c)}{b}$$

output

$$1/2*\cosh(a+c)*\operatorname{csch}(b*x-c)^2/b+\operatorname{coth}(b*x-c)*\sinh(a+c)/b$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\begin{aligned} &\int \cosh(a + bx)\operatorname{csch}^3(c - bx) dx \\ &= -\frac{\operatorname{csch}(c)\operatorname{csch}^2(c - bx)(\sinh(a) - \cosh(c - 2bx)\sinh(a + c))}{2b} \end{aligned}$$

input

$$\operatorname{Integrate}[\operatorname{Cosh}[a + b*x]*\operatorname{Csch}[c - b*x]^3,x]$$

output

$$-1/2*(\operatorname{Csch}[c]*\operatorname{Csch}[c - b*x]^2*(\operatorname{Sinh}[a] - \operatorname{Cosh}[c - 2*b*x]*\operatorname{Sinh}[a + c]))/b$$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx$$

input `Int[Cosh[a + b*x]*Csch[c - b*x]^3,x]`

output `$Aborted`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

method	result	size
parallelrisch	$\frac{\operatorname{sech}\left(\frac{bx}{2} - \frac{c}{2}\right)^2 \operatorname{csch}\left(\frac{bx}{2} - \frac{c}{2}\right)^2 \cosh(2bx + a - c)}{8b}$	38
risch	$-\frac{(e^{2a+2c} - 2e^{2bx+2a} - 1)e^{3a+3c}}{(e^{2a+2c} - e^{2bx+2a})^2 b}$	57

input `int(-cosh(b*x+a)*csch(b*x-c)^3,x,method=_RETURNVERBOSE)`

output `1/8/b*sech(1/2*b*x-1/2*c)^2*csch(1/2*b*x-1/2*c)^2*cosh(2*b*x+a-c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 453 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 453, normalized size of antiderivative = 12.24

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx$$

$$= \frac{b \cosh(bx + a)^3 \cosh(a + c)^3 - b \cosh(bx + a) \cosh(a + c)^3 + (b \cosh(a + c)^3 - 3b \cosh(a + c)^2 \sinh(a + c)) \sinh(bx + a)}{b^3 \cosh(bx + a)^3 \cosh(a + c)^3 - b^3 \cosh(bx + a) \cosh(a + c)^3 + (b^3 \cosh(a + c)^3 - 3b^3 \cosh(a + c)^2 \sinh(a + c)) \sinh(bx + a)}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c)^3,x, algorithm="fricas")`

output

```
2*(cosh(b*x + a)*cosh(a + c)^2 - cosh(b*x + a)*cosh(a + c)*sinh(a + c) - 2
*cosh(b*x + a)*sinh(a + c)^2 + (cosh(a + c)^2 + cosh(a + c)*sinh(a + c))*s
inh(b*x + a))/(b*cosh(b*x + a)^3*cosh(a + c)^3 - b*cosh(b*x + a)*cosh(a +
c)^3 + (b*cosh(a + c)^3 - 3*b*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(a + c)*
sinh(a + c)^2 - b*sinh(a + c)^3)*sinh(b*x + a)^3 - (b*cosh(b*x + a)^3 + 3*
b*cosh(b*x + a))*sinh(a + c)^3 + 3*(b*cosh(b*x + a)*cosh(a + c)^3 - 3*b*co
sh(b*x + a)*cosh(a + c)^2*sinh(a + c) + 3*b*cosh(b*x + a)*cosh(a + c)*sinh
(a + c)^2 - b*cosh(b*x + a)*sinh(a + c)^3)*sinh(b*x + a)^2 + (3*b*cosh(b*x
+ a)^3*cosh(a + c) + b*cosh(b*x + a)*cosh(a + c))*sinh(a + c)^2 + (3*b*co
sh(b*x + a)^2*cosh(a + c)^3 - 3*b*cosh(a + c)^3 - (3*b*cosh(b*x + a)^2 + b
)*sinh(a + c)^3 + 3*(3*b*cosh(b*x + a)^2*cosh(a + c) + b*cosh(a + c))*sinh
(a + c)^2 - (9*b*cosh(b*x + a)^2*cosh(a + c)^2 - b*cosh(a + c)^2)*sinh(a +
c))*sinh(b*x + a) - 3*(b*cosh(b*x + a)^3*cosh(a + c)^2 - b*cosh(b*x + a)*
cosh(a + c)^2)*sinh(a + c))
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx = - \int \cosh(a + bx) \operatorname{csch}^3(bx - c) dx$$

input `integrate(-cosh(b*x+a)*csch(b*x-c)**3,x)`

output `-Integral(cosh(a + b*x)*csch(b*x - c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 131, normalized size of antiderivative = 3.54

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx = -\frac{2e^{(-2bx+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} - \frac{e^{(2a+2c)}}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})} + \frac{1}{b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c)^3,x, algorithm="maxima")`

output `-2*e^(-2*b*x + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + 5*c) - e^(a + c))) - e^(2*a + 2*c)/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + 5*c) - e^(a + c))) + 1/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + 5*c) - e^(a + c)))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.38

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx = \frac{(2e^{(2bx+2a+3c)} - e^{(2a+5c)} + e^{(3c)})e^{(-a)}}{b(e^{(2bx)} - e^{(2c)})^2}$$

input `integrate(-cosh(b*x+a)*csch(b*x-c)^3,x, algorithm="giac")`

output `(2*e^(2*b*x + 2*a + 3*c) - e^(2*a + 5*c) + e^(3*c))*e^(-a)/(b*(e^(2*b*x) - e^(2*c))^2)`

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx = \int \frac{\cosh(a + bx)}{\sinh(c - bx)^3} dx$$

input `int(cosh(a + b*x)/sinh(c - b*x)^3,x)`output `int(cosh(a + b*x)/sinh(c - b*x)^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.46

$$\int \cosh(a + bx) \operatorname{csch}^3(c - bx) dx = \frac{e^c (e^{4bx+2a} + e^{2c})}{eab (e^{4bx} - 2e^{2bx+2c} + e^{4c})}$$

input `int(-cosh(b*x+a)*csch(b*x-c)^3,x)`output `(e**c*(e**(2*a + 4*b*x) + e**(2*c)))/(e**a*b*(e**(4*b*x) - 2*e**(2*b*x + 2*c) + e**(4*c)))`

3.155 $\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$

Optimal result	1065
Mathematica [A] (verified)	1065
Rubi [F]	1066
Maple [B] (verified)	1066
Fricas [B] (verification not implemented)	1067
Sympy [F]	1067
Maxima [B] (verification not implemented)	1068
Giac [B] (verification not implemented)	1068
Mupad [F(-1)]	1069
Reduce [B] (verification not implemented)	1069

Optimal result

Integrand size = 16, antiderivative size = 61

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx = \frac{\cosh(a + c) \operatorname{csch}^3(c - bx)}{3b} + \frac{1}{2} \left(\frac{\operatorname{arctanh}(\cosh(c - bx))}{b} - \frac{\operatorname{coth}(c - bx) \operatorname{csch}(c - bx)}{b} \right) \sinh(a + c)$$

output

```
-1/3*cosh(a+c)*csch(b*x-c)^3/b+1/2*(arctanh(cosh(b*x-c))/b-coth(b*x-c)*csc
h(b*x-c)/b)*sinh(a+c)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx = \frac{12 \operatorname{arctanh}(\cosh(c) - \sinh(c) \tanh(\frac{bx}{2})) \sinh(a + c) + \operatorname{csch}^3(c - bx) (4 \cosh(a + c) - 3 \sinh(a + c) \sinh(2(c - bx)))}{12b}$$

input

```
Integrate[Cosh[a + b*x]*Csch[c - b*x]^4,x]
```

output

```
(12*ArcTanh[Cosh[c] - Sinh[c]*Tanh[(b*x)/2]]*Sinh[a + c] + Csch[c - b*x]^3
*(4*Cosh[a + c] - 3*Sinh[a + c]*Sinh[2*(c - b*x)]))/(12*b)
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$$

↓ 7299

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$$

input

```
Int[Cosh[a + b*x]*Csch[c - b*x]^4,x]
```

output

```
$Aborted
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. $2(61) = 122$.

Time = 1.14 (sec) , antiderivative size = 229, normalized size of antiderivative = 3.75

method	result
risch	$-\frac{e^{bx+a}(3e^{6a+6c}-8e^{2bx+6a+4c}-3e^{4a+4c}-3e^{4bx+6a+2c}-8e^{2bx+4a+2c}+3e^{4bx+4a})}{6b(e^{2a+2c}-e^{2bx+2a})^3} + \frac{\ln(e^{a+c}+e^{bx+a})e^{-a-c}e^{2a+2c}}{4b} - \frac{\ln(e^{a+c}-e^{bx+a})e^{-a-c}e^{2a+2c}}{4b}$

input

```
int(cosh(b*x+a)*csch(b*x-c)^4,x,method=_RETURNVERBOSE)
```

output

```
-1/6*exp(b*x+a)*(3*exp(6*a+6*c)-8*exp(2*b*x+6*a+4*c)-3*exp(4*a+4*c)-3*exp(
4*b*x+6*a+2*c)-8*exp(2*b*x+4*a+2*c)+3*exp(4*b*x+4*a))/b/(exp(2*a+2*c)-exp(
2*b*x+2*a))^3+1/4*ln(exp(a+c)+exp(b*x+a))/b*exp(-a-c)*exp(2*a+2*c)-1/4*ln(
exp(a+c)+exp(b*x+a))/b*exp(-a-c)-1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-a-c)*
exp(2*a+2*c)+1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-a-c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9016 vs. $2(61) = 122$.

Time = 0.16 (sec) , antiderivative size = 9016, normalized size of antiderivative = 147.80

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)*csch(b*x-c)^4,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx = \int \cosh(a + bx) \operatorname{csch}^4(bx - c) dx$$

input

```
integrate(cosh(b*x+a)*csch(b*x-c)**4,x)
```

output

```
Integral(cosh(a + b*x)*csch(b*x - c)**4, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(61) = 122$.

Time = 0.05 (sec) , antiderivative size = 187, normalized size of antiderivative = 3.07

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$$

$$= \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} + 1)}{4b} - \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(-bx+c)} - 1)}{4b}$$

$$+ \frac{3(e^{(2a+2c)} - 1)e^{(-bx-a)} + 8(e^{(4a+4c)} + e^{(2a+2c)})e^{(-3bx-3a)} - 3(e^{(6a+6c)} - e^{(4a+4c)})e^{(-5bx-5a)}}{6b(3e^{(-2bx+2c)} - 3e^{(-4bx+4c)} + e^{(-6bx+6c)} - 1)}$$

input `integrate(cosh(b*x+a)*csch(b*x-c)^4,x, algorithm="maxima")`

output $\frac{1}{4}*(e^{(2*a + 2*c)} - 1)*e^{(-a - c)}*\log(e^{(-b*x + c)} + 1)/b - \frac{1}{4}*(e^{(2*a + 2*c)} - 1)*e^{(-a - c)}*\log(e^{(-b*x + c)} - 1)/b + \frac{1}{6}*(3*(e^{(2*a + 2*c)} - 1)*e^{(-b*x - a)} + 8*(e^{(4*a + 4*c)} + e^{(2*a + 2*c)})*e^{(-3*b*x - 3*a)} - 3*(e^{(6*a + 6*c)} - e^{(4*a + 4*c)})*e^{(-5*b*x - 5*a)})/(b*(3*e^{(-2*b*x + 2*c)} - 3*e^{(-4*b*x + 4*c)} + e^{(-6*b*x + 6*c)} - 1))$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. $2(61) = 122$.

Time = 0.19 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.61

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$$

$$= \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(e^{(bx)} + e^c)}{4b} - \frac{(e^{(2a+2c)} - 1)e^{(-a-c)} \log(|e^{(bx)} - e^c|)}{4b}$$

$$+ \frac{(3e^{(5bx)} - 3e^{(5bx+2a+2c)} - 8e^{(3bx+2a+4c)} - 8e^{(3bx+2c)} + 3e^{(bx+2a+6c)} - 3e^{(bx+4c)})e^{(-a)}}{6b(e^{(2bx)} - e^{(2c)})^3}$$

input `integrate(cosh(b*x+a)*csch(b*x-c)^4,x, algorithm="giac")`

output

```
1/4*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(e^(b*x) + e^c)/b - 1/4*(e^(2*a + 2*c) - 1)*e^(-a - c)*log(abs(e^(b*x) - e^c))/b + 1/6*(3*e^(5*b*x) - 3*e^(5*b*x + 2*a + 2*c) - 8*e^(3*b*x + 2*a + 4*c) - 8*e^(3*b*x + 2*c) + 3*e^(b*x + 2*a + 6*c) - 3*e^(b*x + 4*c))*e^(-a)/(b*(e^(2*b*x) - e^(2*c))^3)
```

Mupad [F(-1)]

Timed out.

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx = \int \frac{\cosh(a + bx)}{\sinh(c - bx)^4} dx$$

input

```
int(cosh(a + b*x)/sinh(c - b*x)^4,x)
```

output

```
int(cosh(a + b*x)/sinh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 496, normalized size of antiderivative = 8.13

$$\int \cosh(a + bx) \operatorname{csch}^4(c - bx) dx$$

$$= \frac{3e^{6bx+2a+2c} \log(e^{bx} + e^c) - 3e^{6bx+2a+2c} \log(e^{bx} - e^c) - 3e^{6bx} \log(e^{bx} + e^c) + 3e^{6bx} \log(e^{bx} - e^c) - 6e^{5bx+2a}}{\dots}$$

input

```
int(cosh(b*x+a)*csch(b*x-c)^4,x)
```

output

```
(3*e**(2*a + 6*b*x + 2*c)*log(e**(b*x) + e**c) - 3*e**(2*a + 6*b*x + 2*c)*
log(e**(b*x) - e**c) - 3*e**(6*b*x)*log(e**(b*x) + e**c) + 3*e**(6*b*x)*lo
g(e**(b*x) - e**c) - 6*e**(2*a + 5*b*x + 3*c) + 6*e**(5*b*x + c) - 9*e**(2
*a + 4*b*x + 4*c)*log(e**(b*x) + e**c) + 9*e**(2*a + 4*b*x + 4*c)*log(e**(
b*x) - e**c) + 9*e**(4*b*x + 2*c)*log(e**(b*x) + e**c) - 9*e**(4*b*x + 2*c
)*log(e**(b*x) - e**c) - 16*e**(2*a + 3*b*x + 5*c) - 16*e**(3*b*x + 3*c) +
9*e**(2*a + 2*b*x + 6*c)*log(e**(b*x) + e**c) - 9*e**(2*a + 2*b*x + 6*c)*
log(e**(b*x) - e**c) - 9*e**(2*b*x + 4*c)*log(e**(b*x) + e**c) + 9*e**(2*b
*x + 4*c)*log(e**(b*x) - e**c) + 6*e**(2*a + b*x + 7*c) - 6*e**(b*x + 5*c)
- 3*e**(2*a + 8*c)*log(e**(b*x) + e**c) + 3*e**(2*a + 8*c)*log(e**(b*x) -
e**c) + 3*e**(6*c)*log(e**(b*x) + e**c) - 3*e**(6*c)*log(e**(b*x) - e**c)
)/(12*e**(a + c)*b*(e**(6*b*x) - 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) -
e**(6*c)))
```

3.156 $\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx$

Optimal result	1071
Mathematica [A] (verified)	1071
Rubi [F]	1072
Maple [B] (verified)	1072
Fricas [B] (verification not implemented)	1073
Sympy [F]	1074
Maxima [B] (verification not implemented)	1074
Giac [B] (verification not implemented)	1075
Mupad [B] (verification not implemented)	1075
Reduce [B] (verification not implemented)	1076

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx = -\frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh^2(a - c)}{b} + \frac{\cosh(2a - c + bx)}{b}$$

output `-arctanh(cosh(b*x+c))*cosh(a-c)^2/b+cosh(b*x+2*a-c)/b`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.39

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx = \frac{\cosh(2a - c + bx) + \cosh^2(a - c) \left(-\log \left(\cosh \left(\frac{1}{2}(c + bx) \right) \right) + \log \left(\sinh \left(\frac{1}{2}(c + bx) \right) \right) \right)}{b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c + b*x],x]`

output `(Cosh[2*a - c + b*x] + Cosh[a - c]^2*(-Log[Cosh[(c + b*x)/2]] + Log[Sinh[(c + b*x)/2]]))/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c + b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. $2(36) = 72$.

Time = 0.43 (sec) , antiderivative size = 233, normalized size of antiderivative = 6.47

method	result
risch	$\frac{e^{bx+2a-c}}{2b} + \frac{e^{-bx-2a+c}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-2c-2a}e^{4a}}{4b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-2c-2a}e^{2a+2c}}{2b} + \frac{\ln(e^{bx+a}-e^{a-c})e^{-2c-2a}e^{4c}}{4b}$

input `int(cosh(b*x+a)^2*csch(b*x+c),x,method=_RETURNVERBOSE)`

output $\frac{1}{2} \frac{1}{b} \exp(bx+2a-c) + \frac{1}{2} \frac{1}{b} \exp(-bx-2a+c) + \frac{1}{4} \ln(\exp(bx+a) - \exp(a-c)) / b \exp(-2c-2a) \exp(4a) + \frac{1}{2} \ln(\exp(bx+a) - \exp(a-c)) / b \exp(-2c-2a) \exp(2a+2c) + \frac{1}{4} \ln(\exp(bx+a) - \exp(a-c)) / b \exp(-2c-2a) \exp(4c) - \frac{1}{4} \ln(\exp(bx+a) + \exp(a-c)) / b \exp(-2c-2a) \exp(4a) - \frac{1}{2} \ln(\exp(bx+a) + \exp(a-c)) / b \exp(-2c-2a) \exp(2a+2c) - \frac{1}{4} \ln(\exp(bx+a) + \exp(a-c)) / b \exp(-2c-2a) \exp(4c)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 872 vs. $2(36) = 72$.

Time = 0.11 (sec) , antiderivative size = 872, normalized size of antiderivative = 24.22

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*cscch(b*x+c),x, algorithm="fricas")`

output

```
1/4*(2*cosh(b*x + c)^2*cosh(-a + c)^4 - 8*cosh(b*x + c)^2*cosh(-a + c)^3*
sinh(-a + c) + 12*cosh(b*x + c)^2*cosh(-a + c)^2*sinh(-a + c)^2 - 8*cosh(b*
x + c)^2*cosh(-a + c)*sinh(-a + c)^3 + 2*cosh(b*x + c)^2*sinh(-a + c)^4 +
2*(cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a + c)^2*sinh(
-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4)*sinh(b*x + c)^
2 + (4*cosh(b*x + c)*cosh(-a + c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a +
c)^4 - 2*(3*cosh(-a + c)^2 + 1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a
+ c)^3 + cosh(-a + c))*cosh(b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 + 2*c
osh(-a + c)^2 + 1)*cosh(b*x + c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-
a + c)^3 + sinh(-a + c)^4 + 2*(3*cosh(-a + c)^2 + 1)*sinh(-a + c)^2 + 2*co
sh(-a + c)^2 - 4*(cosh(-a + c)^3 + cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*
x + c))*log(cosh(b*x + c) + sinh(b*x + c) + 1) - (4*cosh(b*x + c)*cosh(-a
+ c)*sinh(-a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - 2*(3*cosh(-a + c)^2 +
1)*cosh(b*x + c)*sinh(-a + c)^2 + 4*(cosh(-a + c)^3 + cosh(-a + c))*cosh(
b*x + c)*sinh(-a + c) - (cosh(-a + c)^4 + 2*cosh(-a + c)^2 + 1)*cosh(b*x +
c) - (cosh(-a + c)^4 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 + 2
*(3*cosh(-a + c)^2 + 1)*sinh(-a + c)^2 + 2*cosh(-a + c)^2 - 4*(cosh(-a + c
)^3 + cosh(-a + c))*sinh(-a + c) + 1)*sinh(b*x + c))*log(cosh(b*x + c) + s
inh(b*x + c) - 1) + 4*(cosh(b*x + c)*cosh(-a + c)^4 - 4*cosh(b*x + c)*cosh
(-a + c)^3*sinh(-a + c) + 6*cosh(b*x + c)*cosh(-a + c)^2*sinh(-a + c)^2...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx = \int \cosh^2(a + bx) \operatorname{csch}(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x+c), x)`

output `Integral(cosh(a + b*x)**2*csch(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.22

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx \\ &= -\frac{(e^{4a} + 2e^{(2a+2c)} + e^{4c})e^{(-2a-2c)} \log(e^{-bx} + e^c)}{4b} \\ & \quad + \frac{(e^{4a} + 2e^{(2a+2c)} + e^{4c})e^{(-2a-2c)} \log(e^{-bx} - e^c)}{4b} + \frac{e^{(bx+2a-c)}}{2b} + \frac{e^{(-bx-2a+c)}}{2b} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c), x, algorithm="maxima")`

output `-1/4*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/4*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + 1/2*e^(b*x + 2*a - c)/b + 1/2*e^(-b*x - 2*a + c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(36) = 72$.

Time = 0.12 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.31

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx$$

$$= -\frac{(e^{4a+c} + 2e^{2a+3c} + e^{5c})e^{(-2a-3c)} \log(e^{(bx+c)} + 1)}{4b}$$

$$+ \frac{(e^{4a+c} + 2e^{2a+3c} + e^{5c})e^{(-2a-3c)} \log(|e^{(bx+c)} - 1|)}{4b} + \frac{e^{(bx+2a-c)}}{2b} + \frac{e^{(-bx-2a+c)}}{2b}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c),x, algorithm="giac")`

output `-1/4*(e^(4*a + c) + 2*e^(2*a + 3*c) + e^(5*c))*e^(-2*a - 3*c)*log(e^(b*x + c) + 1)/b + 1/4*(e^(4*a + c) + 2*e^(2*a + 3*c) + e^(5*c))*e^(-2*a - 3*c)*log(abs(e^(b*x + c) - 1))/b + 1/2*e^(b*x + 2*a - c)/b + 1/2*e^(-b*x - 2*a + c)/b`

Mupad [B] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 202, normalized size of antiderivative = 5.61

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx = \frac{e^{c-2a-bx}}{2b} + \frac{e^{2a-c+bx}}{2b}$$

$$- \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{3c} e^{bx} (\sqrt{-b^2} + 2e^{2a} e^{-2c} \sqrt{-b^2} + e^{4a} e^{-4c} \sqrt{-b^2})}{b \sqrt{e^{-4a} e^{4c} (4e^{2a} e^{-2c} + 6e^{4a} e^{-4c} + 4e^{6a} e^{-6c} + e^{8a} e^{-8c} + 1)}}\right)}{2\sqrt{-b^2}} \sqrt{e^{4c-4a} (4e^{2a-2c} + 6e^{4a-4c} + 4e^{6a-6c} + e^{8a-8c} + 1)}$$

input `int(cosh(a + b*x)^2/sinh(c + b*x),x)`

output `exp(c - 2*a - b*x)/(2*b) + exp(2*a - c + b*x)/(2*b) - (atan((exp(-2*a)*exp(3*c)*exp(b*x))*((-b^2)^(1/2) + 2*exp(2*a)*exp(-2*c)*((-b^2)^(1/2) + exp(4*a)*exp(-4*c)*((-b^2)^(1/2)))/(b*(exp(-4*a)*exp(4*c)*(4*exp(2*a)*exp(-2*c) + 6*exp(4*a)*exp(-4*c) + 4*exp(6*a)*exp(-6*c) + exp(8*a)*exp(-8*c) + 1))^(1/2)))*(exp(4*c - 4*a)*(4*exp(2*a - 2*c) + 6*exp(4*a - 4*c) + 4*exp(6*a - 6*c) + exp(8*a - 8*c) + 1))^(1/2))/(2*(-b^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 170, normalized size of antiderivative = 4.72

$$\int \cosh^2(a + bx) \operatorname{csch}(c + bx) dx$$

$$= \frac{2e^{2bx+4a+c} + e^{bx+4a} \log(e^{bx+c} - 1) - e^{bx+4a} \log(e^{bx+c} + 1) + 2e^{bx+2a+2c} \log(e^{bx+c} - 1) - 2e^{bx+2a+2c} \log(e^{bx+c} + 1)}{4e^{bx+2a+2c} b}$$

input

```
int(cosh(b*x+a)^2*csch(b*x+c),x)
```

output

```
(2*e**(4*a + 2*b*x + c) + e**(4*a + b*x)*log(e**(b*x + c) - 1) - e**(4*a +
b*x)*log(e**(b*x + c) + 1) + 2*e**(2*a + b*x + 2*c)*log(e**(b*x + c) - 1)
- 2*e**(2*a + b*x + 2*c)*log(e**(b*x + c) + 1) + e**(b*x + 4*c)*log(e**(b
*x + c) - 1) - e**(b*x + 4*c)*log(e**(b*x + c) + 1) + 2*e**(3*c))/(4*e**(2
*a + b*x + 2*c)*b)
```

3.157 $\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx$

Optimal result	1077
Mathematica [A] (verified)	1077
Rubi [F]	1078
Maple [B] (verified)	1078
Fricas [B] (verification not implemented)	1079
Sympy [F]	1079
Maxima [B] (verification not implemented)	1080
Giac [B] (verification not implemented)	1080
Mupad [B] (verification not implemented)	1081
Reduce [B] (verification not implemented)	1081

Optimal result

Integrand size = 17, antiderivative size = 49

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = x \cosh(2(a - c)) - \frac{\cosh^2(a - c) \coth(c + bx)}{b} + \frac{\log(\sinh(c + bx)) \sinh(2(a - c))}{b}$$

output `x*cosh(2*a-2*c)-cosh(a-c)^2*coth(b*x+c)/b+ln(sinh(b*x+c))*sinh(2*a-2*c)/b`

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = \frac{bx \cosh(2(a - c)) + \log(\sinh(c + bx)) \sinh(2(a - c)) + \cosh^2(a - c) \operatorname{csch}(c) \operatorname{csch}(c + bx) \sinh(bx)}{b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c + b*x]^2,x]`

output `(b*x*Cosh[2*(a - c)] + Log[Sinh[c + b*x]]*Sinh[2*(a - c)] + Cosh[a - c]^2*Csch[c]*Csch[c + b*x]*Sinh[b*x])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c + b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 256 vs. $2(49) = 98$.

Time = 0.89 (sec) , antiderivative size = 257, normalized size of antiderivative = 5.24

method	result
risch	$x e^{2a-2c} - e^{-2c-2a} e^{4a} x + e^{-2c-2a} e^{4c} x - \frac{e^{-2c-2a} e^{4a} a}{b} + \frac{e^{-2c-2a} e^{4c} a}{b} + \frac{e^{-2c} e^{4a}}{2b(-e^{2bx+2a+2c} + e^{2a})} + \frac{e^{-2c} e^{2a}}{b(-e^{2bx+2a} + e^{2c})}$

input `int(cosh(b*x+a)^2*csch(b*x+c)^2,x,method=_RETURNVERBOSE)`

output `x*exp(2*a-2*c)-exp(-2*c-2*a)*exp(4*a)*x+exp(-2*c-2*a)*exp(4*c)*x-1/b*exp(-2*c-2*a)*exp(4*a)*a+1/b*exp(-2*c-2*a)*exp(4*c)*a+1/2/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*a)+1/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(2*a+2*c)+1/2/b*exp(-2*c)/(-exp(2*b*x+2*a+2*c)+exp(2*a))*exp(4*c)+1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*a)-1/2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/b*exp(-2*c-2*a)*exp(4*c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 708 vs. $2(49) = 98$.

Time = 0.11 (sec) , antiderivative size = 708, normalized size of antiderivative = 14.45

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^2,x, algorithm="fricas")`

output

```
1/2*(2*b*x*cosh(b*x + c)^2 - cosh(-a + c)^4 + 4*b*x*cosh(b*x + c)*sinh(b*x
+ c) + 2*b*x*sinh(b*x + c)^2 + 4*cosh(-a + c)*sinh(-a + c)^3 - sinh(-a +
c)^4 - 2*(3*cosh(-a + c)^2 + 1)*sinh(-a + c)^2 - 2*b*x - 2*cosh(-a + c)^2
+ ((cosh(b*x + c)^2 - 1)*sinh(-a + c)^4 - cosh(-a + c)^4 - 4*(cosh(b*x + c
)^2*cosh(-a + c) - cosh(-a + c))*sinh(-a + c)^3 + (cosh(-a + c)^4 - 1)*cos
h(b*x + c)^2 + (cosh(-a + c)^4 - 4*cosh(-a + c)^3*sinh(-a + c) + 6*cosh(-a
+ c)^2*sinh(-a + c)^2 - 4*cosh(-a + c)*sinh(-a + c)^3 + sinh(-a + c)^4 -
1)*sinh(b*x + c)^2 + 6*(cosh(b*x + c)^2*cosh(-a + c)^2 - cosh(-a + c)^2)*s
inh(-a + c)^2 - 2*(4*cosh(b*x + c)*cosh(-a + c)^3*sinh(-a + c) - 6*cosh(b*
x + c)*cosh(-a + c)^2*sinh(-a + c)^2 + 4*cosh(b*x + c)*cosh(-a + c)*sinh(-
a + c)^3 - cosh(b*x + c)*sinh(-a + c)^4 - (cosh(-a + c)^4 - 1)*cosh(b*x +
c))*sinh(b*x + c) - 4*(cosh(b*x + c)^2*cosh(-a + c)^3 - cosh(-a + c)^3)*si
nh(-a + c) + 1)*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))) + 4*(
cosh(-a + c)^3 + cosh(-a + c))*sinh(-a + c) - 1)/(b*cosh(b*x + c)^2*cosh(-
a + c)^2 - b*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a
+ c) + b*sinh(-a + c)^2)*sinh(b*x + c)^2 + (b*cosh(b*x + c)^2 - b)*sinh(-
a + c)^2 + 2*(b*cosh(b*x + c)*cosh(-a + c)^2 - 2*b*cosh(b*x + c)*cosh(-a +
c)*sinh(-a + c) + b*cosh(b*x + c)*sinh(-a + c)^2)*sinh(b*x + c) - 2*(b*co
sh(b*x + c)^2*cosh(-a + c) - b*cosh(-a + c))*sinh(-a + c))
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = \int \cosh^2(a + bx) \operatorname{csch}^2(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x+c)**2,x)`

output `Integral(cosh(a + b*x)**2*csch(b*x + c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(49) = 98$.

Time = 0.05 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.69

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{-bx} + e^c)}{2b} + \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(e^{-bx} - e^c)}{2b} + \frac{(bx + a)e^{(2a-2c)}}{b} + \frac{e^{4a} + 2e^{(2a+2c)} + e^{4c}}{2b(e^{(-2bx+2a)} - e^{(2a+2c)})}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/2*(e^(4*a) - e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + (b*x + a)*e^(2*a - 2*c)/b + 1/2*(e^(4*a) + 2*e^(2*a + 2*c) + e^(4*c))/(b*(e^(-2*b*x + 2*a) - e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(49) = 98$.

Time = 0.14 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.10

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c + bx) dx = xe^{(-2a+2c)} + \frac{(e^{4a} - e^{4c})e^{(-2a-2c)} \log(|e^{(2bx+2c)} - 1|)}{2b} - \frac{(e^{(2bx+4a)} - e^{(2bx+4c)} + 2e^{(2a)} + 2e^{(2c)})e^{(-2a)}}{2b(e^{(2bx+2c)} - 1)}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^2,x, algorithm="giac")`

output

$$x e^{-2a+2c} + \frac{1}{2} (e^{4a} - e^{4c}) e^{-2a-2c} \log(\operatorname{abs}(e^{2bx+2c} - 1)) / b - \frac{1}{2} (e^{2bx+4a} - e^{2bx+4c} + 2e^{2a} + 2e^{2c}) e^{-2a} / (b(e^{2bx+2c} - 1))$$

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.80

$$\int \cosh^2(a+bx) \operatorname{csch}^2(c+bx) dx = x e^{2c-2a} + \frac{\sinh(2a-2c) \ln(e^{2a} e^{2bx} - e^{2a} e^{-2c})}{b} + \frac{2 e^{2a-2c} \cosh(a-c)^2}{b(e^{2a-2c} - e^{2a+2bx})}$$

input

$$\operatorname{int}(\cosh(a+b*x)^2/\sinh(c+b*x)^2,x)$$

output

$$x \exp(2c-2a) + (\sinh(2a-2c) \log(\exp(2a) \exp(2bx) - \exp(2a) \exp(-2c))) / b + (2 \exp(2a-2c) \cosh(a-c)^2) / (b(\exp(2a-2c) - \exp(2a+2bx)))$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 254, normalized size of antiderivative = 5.18

$$\int \cosh^2(a+bx) \operatorname{csch}^2(c+bx) dx = \frac{e^{2bx+4a+2c} \log(e^{bx+c} - 1) + e^{2bx+4a+2c} \log(e^{bx+c} + 1) - e^{2bx+4a+2c} - 2e^{2bx+2a+4c} - e^{2bx+6c} \log(e^{bx+c} - 1) - e^{2bx+6c} \log(e^{bx+c} + 1)}{2b}$$

input

$$\operatorname{int}(\cosh(b*x+a)^2*\operatorname{csch}(b*x+c)^2,x)$$

output

```
(e**(4*a + 2*b*x + 2*c)*log(e**(b*x + c) - 1) + e**(4*a + 2*b*x + 2*c)*log
(e**(b*x + c) + 1) - e**(4*a + 2*b*x + 2*c) - 2*e**(2*a + 2*b*x + 4*c) - e
**(2*b*x + 6*c)*log(e**(b*x + c) - 1) - e**(2*b*x + 6*c)*log(e**(b*x + c)
+ 1) + 2*e**(2*b*x + 6*c)*b*x - e**(2*b*x + 6*c) - e**(4*a)*log(e**(b*x +
c) - 1) - e**(4*a)*log(e**(b*x + c) + 1) + e**(4*c)*log(e**(b*x + c) - 1)
+ e**(4*c)*log(e**(b*x + c) + 1) - 2*e**(4*c)*b*x)/(2*e**(2*a + 2*c)*b*(e
*(2*b*x + 2*c) - 1))
```

3.158 $\int \cosh^2(a + bx)\operatorname{csch}^3(c + bx) dx$

Optimal result	1083
Mathematica [B] (verified)	1083
Rubi [F]	1084
Maple [B] (verified)	1085
Fricas [B] (verification not implemented)	1085
Sympy [F]	1086
Maxima [B] (verification not implemented)	1086
Giac [B] (verification not implemented)	1087
Mupad [F(-1)]	1087
Reduce [B] (verification not implemented)	1088

Optimal result

Integrand size = 17, antiderivative size = 89

$$\int \cosh^2(a + bx)\operatorname{csch}^3(c + bx) dx = \frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh^2(a - c)}{2b} - \frac{\operatorname{arctanh}(\cosh(c + bx)) \cosh(2(a - c))}{b} - \frac{\cosh^2(a - c) \operatorname{coth}(c + bx)\operatorname{csch}(c + bx)}{2b} - \frac{\operatorname{csch}(c + bx) \sinh(2(a - c))}{b}$$

output

$1/2*\operatorname{arctanh}(\cosh(b*x+c))*\cosh(a-c)^2/b-\operatorname{arctanh}(\cosh(b*x+c))*\cosh(2*a-2*c)/b-1/2*\cosh(a-c)^2*\operatorname{coth}(b*x+c)*\operatorname{csch}(b*x+c)/b-\operatorname{csch}(b*x+c)*\sinh(2*a-2*c)/b$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 249 vs. 2(89) = 178.

Time = 6.19 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.80

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx \\ &= \frac{(-\cosh(2a - 2c - \frac{bx}{2}) + \cosh(2a - 2c + \frac{bx}{2})) \operatorname{csch}(\frac{c}{2}) \operatorname{csch}(\frac{c}{2} + \frac{bx}{2})}{4b} \\ &+ \frac{(-1 - \cosh(2a - 2c)) \operatorname{csch}^2(\frac{c}{2} + \frac{bx}{2})}{16b} + \frac{(1 - 3 \cosh(2a - 2c)) \log(\cosh(\frac{c}{2} + \frac{bx}{2}))}{4b} \\ &+ \frac{(-1 + 3 \cosh(2a - 2c)) \log(\sinh(\frac{c}{2} + \frac{bx}{2}))}{4b} \\ &+ \frac{(-\cosh(2a - 2c - \frac{bx}{2}) + \cosh(2a - 2c + \frac{bx}{2})) \operatorname{sech}(\frac{c}{2}) \operatorname{sech}(\frac{c}{2} + \frac{bx}{2})}{4b} \\ &+ \frac{(-1 - \cosh(2a - 2c)) \operatorname{sech}^2(\frac{c}{2} + \frac{bx}{2})}{16b} \end{aligned}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c + b*x]^3,x]`

output `((-Cosh[2*a - 2*c - (b*x)/2] + Cosh[2*a - 2*c + (b*x)/2])*Csch[c/2]*Csch[c/2 + (b*x)/2])/(4*b) + ((-1 - Cosh[2*a - 2*c])*Csch[c/2 + (b*x)/2]^2)/(16*b) + ((1 - 3*Cosh[2*a - 2*c])*Log[Cosh[c/2 + (b*x)/2]])/(4*b) + ((-1 + 3*Cosh[2*a - 2*c])*Log[Sinh[c/2 + (b*x)/2]])/(4*b) + ((-Cosh[2*a - 2*c - (b*x)/2] + Cosh[2*a - 2*c + (b*x)/2])*Sech[c/2]*Sech[c/2 + (b*x)/2])/(4*b) + ((-1 - Cosh[2*a - 2*c])*Sech[c/2 + (b*x)/2]^2)/(16*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}^3(bx + c) dx \\ & \quad \downarrow 7299 \\ & \int \cosh^2(a + bx) \operatorname{csch}^3(bx + c) dx \end{aligned}$$

input `Int[Cosh[a + b*x]^2*Csch[c + b*x]^3,x]`

output \$Aborted

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(85) = 170$.

Time = 2.23 (sec) , antiderivative size = 305, normalized size of antiderivative = 3.43

method	result
risch	$\frac{(-5 e^{2bx+6a+2c} - 2 e^{2bx+4a+4c} + 3 e^{2bx+2a+6c} + 3 e^{6a} - 2 e^{4a+2c} - 5 e^{2a+4c}) e^{bx-c}}{4(-e^{2bx+2a+2c} + e^{2a})^2 b} + \frac{3 \ln(e^{bx+a} - e^{a-c}) e^{-2c-2a} e^{4a}}{8b} - \frac{\ln(e^{bx+a} - e^{a-c})}{b}$

input `int (cosh(b*x+a)^2*csch(b*x+c)^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{4} / (-\exp(2bx+2a+2c) + \exp(2a))^2 / b * (-5 * \exp(2bx+6a+2c) - 2 * \exp(2bx+4a+4c) + 3 * \exp(2bx+2a+6c) + 3 * \exp(6a) - 2 * \exp(4a+2c) - 5 * \exp(2a+4c)) * \exp(bx-c) + 3/8 * \ln(\exp(bx+a) - \exp(a-c)) / b * \exp(-2c-2a) * \exp(4a) - 1/4 * \ln(\exp(bx+a) - \exp(a-c)) / b * \exp(-2c-2a) * \exp(2a+2c) + 3/8 * \ln(\exp(bx+a) - \exp(a-c)) / b * \exp(-2c-2a) * \exp(4c) - 3/8 * \ln(\exp(bx+a) + \exp(a-c)) / b * \exp(-2c-2a) * \exp(4a) + 1/4 * \ln(\exp(bx+a) + \exp(a-c)) / b * \exp(-2c-2a) * \exp(2a+2c) - 3/8 * \ln(\exp(bx+a) + \exp(a-c)) / b * \exp(-2c-2a) * \exp(4c)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3254 vs. $2(85) = 170$.

Time = 0.12 (sec) , antiderivative size = 3254, normalized size of antiderivative = 36.56

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^3,x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx = \int \cosh^2(a + bx) \operatorname{csch}^3(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x+c)**3,x)`

output `Integral(cosh(a + b*x)**2*csch(b*x + c)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. 2(85) = 170.

Time = 0.05 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.33

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx \\ &= -\frac{(3e^{4a} - 2e^{2a+2c} + 3e^{4c})e^{(-2a-2c)} \log(e^{-bx} + e^c)}{8b} \\ & \quad + \frac{(3e^{4a} - 2e^{2a+2c} + 3e^{4c})e^{(-2a-2c)} \log(e^{-bx} - e^c)}{8b} \\ & \quad + \frac{(5e^{4a+2c} + 2e^{2a+4c} - 3e^{6c})e^{(-bx-a)} - (3e^{6a} - 2e^{4a+2c} - 5e^{2a+4c})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+c)} - e^{(a+5c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^3,x, algorithm="maxima")`

output `-1/8*(3*e^(4*a) - 2*e^(2*a + 2*c) + 3*e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) + e^c)/b + 1/8*(3*e^(4*a) - 2*e^(2*a + 2*c) + 3*e^(4*c))*e^(-2*a - 2*c)*log(e^(-b*x) - e^c)/b + 1/4*((5*e^(4*a + 2*c) + 2*e^(2*a + 4*c) - 3*e^(6*c))*e^(-b*x - a) - (3*e^(6*a) - 2*e^(4*a + 2*c) - 5*e^(2*a + 4*c))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + c) - e^(a + 5*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. $2(85) = 170$.

Time = 0.14 (sec) , antiderivative size = 195, normalized size of antiderivative = 2.19

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx$$

$$= -\frac{(3e^{4a+c} - 2e^{(2a+3c)} + 3e^{(5c)})e^{(-2a-3c)} \log(e^{(bx+c)} + 1)}{8b}$$

$$+ \frac{(3e^{4a+c} - 2e^{(2a+3c)} + 3e^{(5c)})e^{(-2a-3c)} \log(|e^{(bx+c)} - 1|)}{8b}$$

$$- \frac{(5e^{(3bx+4a+2c)} + 2e^{(3bx+2a+4c)} - 3e^{(3bx+6c)} - 3e^{(bx+4a)} + 2e^{(bx+2a+2c)} + 5e^{(bx+4c)})e^{(-2a-c)}}{4b(e^{(2bx+2c)} - 1)^2}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^3,x, algorithm="giac")`

output `-1/8*(3*e^(4*a + c) - 2*e^(2*a + 3*c) + 3*e^(5*c))*e^(-2*a - 3*c)*log(e^(b*x + c) + 1)/b + 1/8*(3*e^(4*a + c) - 2*e^(2*a + 3*c) + 3*e^(5*c))*e^(-2*a - 3*c)*log(abs(e^(b*x + c) - 1))/b - 1/4*(5*e^(3*b*x + 4*a + 2*c) + 2*e^(3*b*x + 2*a + 4*c) - 3*e^(3*b*x + 6*c) - 3*e^(b*x + 4*a) + 2*e^(b*x + 2*a + 2*c) + 5*e^(b*x + 4*c))*e^(-2*a - c)/(b*(e^(2*b*x + 2*c) - 1)^2)`

Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx = \int \frac{\cosh(a + bx)^2}{\sinh(c + bx)^3} dx$$

input `int(cosh(a + b*x)^2/sinh(c + b*x)^3,x)`

output `int(cosh(a + b*x)^2/sinh(c + b*x)^3, x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 520, normalized size of antiderivative = 5.84

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c + bx) dx$$

$$= \frac{3e^{4bx+4a+4c} \log(e^{bx+c} - 1) - 3e^{4bx+4a+4c} \log(e^{bx+c} + 1) - 2e^{4bx+2a+6c} \log(e^{bx+c} - 1) + 2e^{4bx+2a+6c} \log(e^{bx+c} + 1)}{b}$$

input `int(cosh(b*x+a)^2*csch(b*x+c)^3,x)`

output

```
(3***e**(4*a + 4*b*x + 4*c)*log(e**(b*x + c) - 1) - 3***e**(4*a + 4*b*x + 4*c)
*log(e**(b*x + c) + 1) - 2***e**(2*a + 4*b*x + 6*c)*log(e**(b*x + c) - 1) +
2***e**(2*a + 4*b*x + 6*c)*log(e**(b*x + c) + 1) + 3***e**(4*b*x + 8*c)*log(e*
*(b*x + c) - 1) - 3***e**(4*b*x + 8*c)*log(e**(b*x + c) + 1) - 10***e**(4*a +
3*b*x + 3*c) - 4***e**(2*a + 3*b*x + 5*c) + 6***e**(3*b*x + 7*c) - 6***e**(4*a +
2*b*x + 2*c)*log(e**(b*x + c) - 1) + 6***e**(4*a + 2*b*x + 2*c)*log(e**(b*x
+ c) + 1) + 4***e**(2*a + 2*b*x + 4*c)*log(e**(b*x + c) - 1) - 4***e**(2*a +
2*b*x + 4*c)*log(e**(b*x + c) + 1) - 6***e**(2*b*x + 6*c)*log(e**(b*x + c) -
1) + 6***e**(2*b*x + 6*c)*log(e**(b*x + c) + 1) + 6***e**(4*a + b*x + c) - 4*
e**(2*a + b*x + 3*c) - 10***e**(b*x + 5*c) + 3***e**(4*a)*log(e**(b*x + c) - 1
) - 3***e**(4*a)*log(e**(b*x + c) + 1) - 2***e**(2*a + 2*c)*log(e**(b*x + c) -
1) + 2***e**(2*a + 2*c)*log(e**(b*x + c) + 1) + 3***e**(4*c)*log(e**(b*x + c)
- 1) - 3***e**(4*c)*log(e**(b*x + c) + 1))/(8***e**(2*a + 2*c)*b*(e**(4*b*x +
4*c) - 2***e**(2*b*x + 2*c) + 1))
```

3.159 $\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx$

Optimal result	1089
Mathematica [A] (verified)	1089
Rubi [F]	1090
Maple [A] (verified)	1090
Fricas [B] (verification not implemented)	1091
Sympy [F]	1092
Maxima [B] (verification not implemented)	1092
Giac [A] (verification not implemented)	1093
Mupad [F(-1)]	1093
Reduce [B] (verification not implemented)	1094

Optimal result

Integrand size = 17, antiderivative size = 84

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \frac{\cosh^2(a - c) \operatorname{coth}(c + bx)}{b} - \frac{\cosh(2(a - c)) \operatorname{coth}(c + bx)}{b} - \frac{\cosh^2(a - c) \operatorname{coth}^3(c + bx)}{3b} - \frac{\operatorname{csch}^2(c + bx) \sinh(2(a - c))}{2b}$$

```
output cosh(a-c)^2*coth(b*x+c)/b-cosh(2*a-2*c)*coth(b*x+c)/b-1/3*cosh(a-c)^2*coth
(b*x+c)^3/b-1/2*csch(b*x+c)^2*sinh(2*a-2*c)/b
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.98

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \frac{\operatorname{csch}(c) \operatorname{csch}^3(c + bx) (3 \sinh(bx) - \sinh(2a - 4c - 3bx)) + 3 \sinh(2a - 2c - bx) - 3 \sinh(2a + bx) + \sinh(2a - 2c)}{12b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c + b*x]^4,x]`

output `(Csch[c]*Csch[c + b*x]^3*(3*Sinh[b*x] - Sinh[2*a - 4*c - 3*b*x] + 3*Sinh[2*a - 2*c - b*x] - 3*Sinh[2*a + b*x] + Sinh[2*a + 3*b*x] - Sinh[2*c + 3*b*x]))/(12*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}^4(bx + c) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}^4(bx + c) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c + b*x]^4,x]`

output `$Aborted`

Maple [A] (verified)

Time = 3.85 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.69

method	result	size
parallelrisch	$\frac{\operatorname{sech}\left(\frac{bx}{2} + \frac{c}{2}\right)^3 \operatorname{csch}\left(\frac{bx}{2} + \frac{c}{2}\right)^3 (\cosh(3bx+3c) - 2 \cosh(3bx+2a+c) - 3 \cosh(bx+c))}{96b}$	58
risch	$\frac{2(3e^{4bx+4a+4c} - 3e^{2bx+4a+2c} + 3e^{2bx+2a+4c} + e^{4a} - e^{2a+2c} + e^{4c})e^{4a-2c}}{3(-e^{2bx+2a+2c} + e^{2a})^3 b}$	96

input `int(cosh(b*x+a)^2*csch(b*x+c)^4,x,method=_RETURNVERBOSE)`

output

```
1/96*sech(1/2*b*x+1/2*c)^3*csch(1/2*b*x+1/2*c)^3*(cosh(3*b*x+3*c)-2*cosh(3
*b*x+2*a+c)-3*cosh(b*x+c))/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 497 vs. $2(80) = 160$.

Time = 0.11 (sec) , antiderivative size = 497, normalized size of antiderivative = 5.92

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx =$$

$$\frac{2 \left((5 \cosh(-a + c))^2 - 1 \right) \cosh(bx + c)^2}{3 (b \cosh(bx + c))^4 \cosh(-a + c)^2 - 4b \cosh(bx + c)^2 \cosh(-a + c)^2 + (b \cosh(-a + c))^2 - b \sinh(-a + c)^2}$$

input

```
integrate(cosh(b*x+a)^2*csch(b*x+c)^4,x, algorithm="fricas")
```

output

```
-2/3*((5*cosh(-a + c)^2 - 1)*cosh(b*x + c)^2 + (5*cosh(-a + c)^2 - 6*cosh(
-a + c)*sinh(-a + c) + 5*sinh(-a + c)^2 - 1)*sinh(b*x + c)^2 + (5*cosh(b*x
+ c)^2 - 3)*sinh(-a + c)^2 - 3*cosh(-a + c)^2 - 2*(6*cosh(b*x + c)*cosh(-
a + c)*sinh(-a + c) - cosh(b*x + c)*sinh(-a + c)^2 - (cosh(-a + c)^2 + 1)*
cosh(b*x + c))*sinh(b*x + c) - 6*(cosh(b*x + c)^2*cosh(-a + c) - cosh(-a +
c))*sinh(-a + c) + 3)/(b*cosh(b*x + c)^4*cosh(-a + c)^2 - 4*b*cosh(b*x +
c)^2*cosh(-a + c)^2 + (b*cosh(-a + c)^2 - b*sinh(-a + c)^2)*sinh(b*x + c)^
4 + 4*(b*cosh(b*x + c)*cosh(-a + c)^2 - b*cosh(b*x + c)*sinh(-a + c)^2)*si
nh(b*x + c)^3 + 3*b*cosh(-a + c)^2 + 2*(3*b*cosh(b*x + c)^2*cosh(-a + c)^2
- 2*b*cosh(-a + c)^2 - (3*b*cosh(b*x + c)^2 - 2*b)*sinh(-a + c)^2)*sinh(b
*x + c)^2 - (b*cosh(b*x + c)^4 - 4*b*cosh(b*x + c)^2 + 3*b)*sinh(-a + c)^2
+ 4*(b*cosh(b*x + c)^3*cosh(-a + c)^2 - b*cosh(b*x + c)*cosh(-a + c)^2 -
(b*cosh(b*x + c)^3 - b*cosh(b*x + c))*sinh(-a + c)^2)*sinh(b*x + c))
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \int \cosh^2(a + bx) \operatorname{csch}^4(bx + c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x+c)**4,x)`

output `Integral(cosh(a + b*x)**2*csch(b*x + c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 333 vs. $2(80) = 160$.

Time = 0.04 (sec) , antiderivative size = 333, normalized size of antiderivative = 3.96

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx \\ &= \frac{2(e^{4a+4c} - e^{2a+6c})e^{-2bx-2a}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(-4bx+4c)}}{b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(4a+4c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &- \frac{2e^{(2a+6c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \\ &+ \frac{2e^{(8c)}}{3b(3e^{(-2bx+2a+4c)} - 3e^{(-4bx+2a+2c)} + e^{(-6bx+2a)} - e^{(2a+6c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*csch(b*x+c)^4,x, algorithm="maxima")`

output

$$\begin{aligned} & 2*(e^{(4*a + 4*c)} - e^{(2*a + 6*c)})*e^{(-2*b*x - 2*a)}/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} - e^{(2*a + 6*c)})) + 2* \\ & e^{(-4*b*x + 4*c)}/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} - e^{(2*a + 6*c)})) + 2/3*e^{(4*a + 4*c)}/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} - e^{(2*a + 6*c)})) \\ & - 2/3*e^{(2*a + 6*c)}/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} - e^{(2*a + 6*c)})) + 2/3*e^{(8*c)}/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 2*c)} + e^{(-6*b*x + 2*a)} - e^{(2*a + 6*c)})) \end{aligned}$$
Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.04

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \frac{2(3e^{(4bx+4a+4c)} - 3e^{(2bx+4a+2c)} + 3e^{(2bx+2a+4c)} + e^{(4a)} - e^{(2a+2c)} + e^{(4c)})e^{(-2a-2c)}}{3b(e^{(2bx+2c)} - 1)^3}$$

input

```
integrate(cosh(b*x+a)^2*csch(b*x+c)^4,x, algorithm="giac")
```

output

$$-2/3*(3*e^{(4*b*x + 4*a + 4*c)} - 3*e^{(2*b*x + 4*a + 2*c)} + 3*e^{(2*b*x + 2*a + 4*c)} + e^{(4*a)} - e^{(2*a + 2*c)} + e^{(4*c)})*e^{(-2*a - 2*c)}/(b*(e^{(2*b*x + 2*c)} - 1)^3)$$
Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \int \frac{\cosh(a + bx)^2}{\sinh(c + bx)^4} dx$$

input

```
int(cosh(a + b*x)^2/sinh(c + b*x)^4,x)
```

output

```
int(cosh(a + b*x)^2/sinh(c + b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.11

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c + bx) dx = \frac{-\frac{2e^{6bx+4a+4c}}{3} - 2e^{2bx+2a+2c} + \frac{2e^{2a}}{3} - \frac{2e^{2c}}{3}}{e^{2a}b(e^{6bx+6c} - 3e^{4bx+4c} + 3e^{2bx+2c} - 1)}$$

input `int(cosh(b*x+a)^2*csch(b*x+c)^4,x)`

output `(2*(- e**(4*a + 6*b*x + 4*c) - 3*e**(2*a + 2*b*x + 2*c) + e**(2*a) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x + 6*c) - 3*e**(4*b*x + 4*c) + 3*e**(2*b*x + 2*c) - 1))`

3.160 $\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$

Optimal result	1095
Mathematica [A] (verified)	1095
Rubi [F]	1096
Maple [B] (verified)	1096
Fricas [B] (verification not implemented)	1097
Sympy [F]	1098
Maxima [B] (verification not implemented)	1098
Giac [B] (verification not implemented)	1099
Mupad [B] (verification not implemented)	1099
Reduce [B] (verification not implemented)	1100

Optimal result

Integrand size = 16, antiderivative size = 33

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx = \frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh^2(a + c)}{b} - \frac{\cosh(2a + c + bx)}{b}$$

output

```
arctanh(cosh(b*x-c))*cosh(a+c)^2/b-cosh(b*x+2*a+c)/b
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.58

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx = \frac{-\cosh(2a + c + bx) + \cosh^2(a + c) (\log(\cosh(\frac{1}{2}(c - bx))) - \log(-\sinh(\frac{1}{2}(c - bx))))}{b}$$

input

```
Integrate[Cosh[a + b*x]^2*Csch[c - b*x],x]
```

output

```
(-Cosh[2*a + c + b*x] + Cosh[a + c]^2*(Log[Cosh[(c - b*x)/2]] - Log[-Sinh[(c - b*x)/2]]))/b
```


Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c - b*x],x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(34) = 68.

Time = 0.27 (sec) , antiderivative size = 221, normalized size of antiderivative = 6.70

method	result
risch	$-\frac{e^{bx+2a+c}}{2b} - \frac{e^{-bx-2a-c}}{2b} - \frac{\ln(-e^{a+c}+e^{bx+a})e^{-2c-2a}e^{4a+4c}}{4b} - \frac{\ln(-e^{a+c}+e^{bx+a})e^{-2c-2a}e^{2a+2c}}{2b} - \frac{\ln(-e^{a+c}+e^{bx+a})e^{-2c-2a}e^{4a+4c}}{4b}$

input `int(-cosh(b*x+a)^2*csch(b*x-c),x,method=_RETURNVERBOSE)`

output `-1/2/b*exp(b*x+2*a+c)-1/2/b*exp(-b*x-2*a-c)-1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/2*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)-1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)+1/4*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4*c)+1/2*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)+1/4*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 657 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 657, normalized size of antiderivative = 19.91

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx = \text{Too large to display}$$

input `integrate(-cosh(b*x+a)^2*csh(b*x-c),x, algorithm="fricas")`

output

```
-1/4*(2*cosh(b*x + a)^2*cosh(a + c) + 2*cosh(a + c)^3 + 2*(cosh(a + c) - s
inh(a + c))*sinh(b*x + a)^2 + 6*cosh(a + c)*sinh(a + c)^2 - 2*sinh(a + c)^
3 + (4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)
^4 - 2*(3*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^2 + 4*(cosh(a + c)^
3 + cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 + 2*cosh(a + c
)^2 + 1)*cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)*sinh(a + c)^3 + si
nh(a + c)^4 + 2*(3*cosh(a + c)^2 + 1)*sinh(a + c)^2 + 2*cosh(a + c)^2 - 4*
(cosh(a + c)^3 + cosh(a + c))*sinh(a + c) + 1)*sinh(b*x + a))*log(cosh(b*x
+ a)*cosh(a + c) + (cosh(a + c) - sinh(a + c))*sinh(b*x + a) - cosh(b*x +
a)*sinh(a + c) + 1) - (4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b
*x + a)*sinh(a + c)^4 - 2*(3*cosh(a + c)^2 + 1)*cosh(b*x + a)*sinh(a + c)^
2 + 4*(cosh(a + c)^3 + cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a +
c)^4 + 2*cosh(a + c)^2 + 1)*cosh(b*x + a) - (cosh(a + c)^4 - 4*cosh(a + c)
*sinh(a + c)^3 + sinh(a + c)^4 + 2*(3*cosh(a + c)^2 + 1)*sinh(a + c)^2 + 2
*cosh(a + c)^2 - 4*(cosh(a + c)^3 + cosh(a + c))*sinh(a + c) + 1)*sinh(b*x
+ a))*log(cosh(b*x + a)*cosh(a + c) + (cosh(a + c) - sinh(a + c))*sinh(b*
x + a) - cosh(b*x + a)*sinh(a + c) - 1) + 4*(cosh(b*x + a)*cosh(a + c) - c
osh(b*x + a)*sinh(a + c))*sinh(b*x + a) - 2*(cosh(b*x + a)^2 + 3*cosh(a +
c)^2)*sinh(a + c))/(b*cosh(b*x + a)*cosh(a + c)^2 - 2*b*cosh(b*x + a)*cosh
(a + c)*sinh(a + c) + b*cosh(b*x + a)*sinh(a + c)^2 + (b*cosh(a + c)^2 ...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx = - \int \cosh^2(a + bx) \operatorname{csch}(bx - c) dx$$

input `integrate(-cosh(b*x+a)**2*csch(b*x-c), x)`

output `-Integral(cosh(a + b*x)**2*csch(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(34) = 68$.

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 3.58

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx \\ &= \frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{4b} \\ & \quad - \frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{4b} - \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b} \end{aligned}$$

input `integrate(-cosh(b*x+a)^2*csch(b*x-c), x, algorithm="maxima")`

output `1/4*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b - 1/4*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b - 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 117 vs. $2(34) = 68$.

Time = 0.12 (sec) , antiderivative size = 117, normalized size of antiderivative = 3.55

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$$

$$= \frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(e^{(bx)} + e^c)}{4b} - \frac{(e^{(4a+4c)} + 2e^{(2a+2c)} + 1)e^{(-2a-2c)} \log(|e^{(bx)} - e^c|)}{4b} - \frac{e^{(bx+2a+c)}}{2b} - \frac{e^{(-bx-2a-c)}}{2b}$$

input `integrate(-cosh(b*x+a)^2*csh(b*x-c),x, algorithm="giac")`

output `1/4*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(e^(b*x) + e^c)/b - 1/4*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)*e^(-2*a - 2*c)*log(abs(e^(b*x) - e^c))/b - 1/2*e^(b*x + 2*a + c)/b - 1/2*e^(-b*x - 2*a - c)/b`

Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 202, normalized size of antiderivative = 6.12

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$$

$$= \frac{\operatorname{atan}\left(\frac{e^{-2a} e^{-3c} e^{bx} (\sqrt{-b^2} + 2e^{2a} e^{2c} \sqrt{-b^2} + e^{4a} e^{4c} \sqrt{-b^2})}{b \sqrt{e^{-4a} e^{-4c} (4e^{2a} e^{2c} + 6e^{4a} e^{4c} + 4e^{6a} e^{6c} + e^{8a} e^{8c} + 1)}}\right) \sqrt{e^{-4a-4c} (4e^{2a+2c} + 6e^{4a+4c} + 4e^{6a+6c} + e^{8a+8c})}}{2\sqrt{-b^2}} - \frac{e^{-2a-c-bx}}{2b} - \frac{e^{2a+c+bx}}{2b}$$

input `int(cosh(a + b*x)^2/sinh(c - b*x),x)`

output

```
(atan((exp(-2*a)*exp(-3*c)*exp(b*x)*((-b^2)^(1/2) + 2*exp(2*a)*exp(2*c)*(-
b^2)^(1/2) + exp(4*a)*exp(4*c)*(-b^2)^(1/2)))/(b*(exp(-4*a)*exp(-4*c)*(4*exp(2*a)*exp(2*c) + 6*exp(4*a)*exp(4*c) + 4*exp(6*a)*exp(6*c) + exp(8*a)*exp(8*c) + 1))^(1/2)))*(exp(-4*a - 4*c)*(4*exp(2*a + 2*c) + 6*exp(4*a + 4*c) + 4*exp(6*a + 6*c) + exp(8*a + 8*c) + 1))^(1/2))/(2*(-b^2)^(1/2)) - exp(-2*a - c - b*x)/(2*b) - exp(2*a + c + b*x)/(2*b)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 174, normalized size of antiderivative = 5.27

$$\int \cosh^2(a + bx) \operatorname{csch}(c - bx) dx$$

$$= \frac{-2e^{2bx+4a+3c} + e^{bx+4a+4c} \log(e^{bx} + e^c) - e^{bx+4a+4c} \log(e^{bx} - e^c) + 2e^{bx+2a+2c} \log(e^{bx} + e^c) - 2e^{bx+2a+2c} \log(e^{bx} - e^c)}{4e^{bx+2a+2c} b}$$

input

```
int(-cosh(b*x+a)^2*csc(b*x-c),x)
```

output

```
( - 2*e**(4*a + 2*b*x + 3*c) + e**(4*a + b*x + 4*c)*log(e**(b*x) + e**c) - e**(4*a + b*x + 4*c)*log(e**(b*x) - e**c) + 2*e**(2*a + b*x + 2*c)*log(e**(b*x) + e**c) - 2*e**(2*a + b*x + 2*c)*log(e**(b*x) - e**c) + e**(b*x)*log(e**(b*x) + e**c) - e**(b*x)*log(e**(b*x) - e**c) - 2*e**c)/(4*e**(2*a + b*x + 2*c)*b)
```

3.161 $\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx$

Optimal result	1101
Mathematica [A] (verified)	1101
Rubi [F]	1102
Maple [B] (verified)	1102
Fricas [B] (verification not implemented)	1103
Sympy [F]	1104
Maxima [B] (verification not implemented)	1104
Giac [B] (verification not implemented)	1105
Mupad [B] (verification not implemented)	1105
Reduce [B] (verification not implemented)	1106

Optimal result

Integrand size = 18, antiderivative size = 44

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = x \cosh(2(a + c)) + \frac{\cosh^2(a + c) \coth(c - bx)}{b} + \frac{\log(\sinh(c - bx)) \sinh(2(a + c))}{b}$$

output `x*cosh(2*a+2*c)-cosh(a+c)^2*coth(b*x-c)/b+ln(-sinh(b*x-c))*sinh(2*a+2*c)/b`

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.16

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = \frac{bx \cosh(2(a + c)) + \log(-\sinh(c - bx)) \sinh(2(a + c)) + \cosh^2(a + c) \operatorname{csch}(c) \operatorname{csch}(c - bx) \sinh(bx)}{b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c - b*x]^2,x]`

output `(b*x*Cosh[2*(a + c)] + Log[-Sinh[c - b*x]]*Sinh[2*(a + c)] + Cosh[a + c]^2*Csch[c]*Csch[c - b*x]*Sinh[b*x])/b`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c - b*x]^2,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. 2(53) = 106.

Time = 0.54 (sec) , antiderivative size = 248, normalized size of antiderivative = 5.64

method	result
risch	$x e^{2a+2c} - e^{-2c-2a} e^{4a+4c} x - \frac{e^{-2c-2a} a e^{4a+4c}}{b} + e^{-2c-2a} x + \frac{e^{-2c-2a} a}{b} + \frac{e^{4a+4c}}{2b(e^{2a+2c} - e^{2bx+2a})} + \frac{e^{2a+2c}}{b(e^{2a+2c} - e^{2bx+2a})}$

input `int(cosh(b*x+a)^2*csch(b*x-c)^2,x,method=_RETURNVERBOSE)`

output `x*exp(2*a+2*c)-exp(-2*c-2*a)*exp(4*a+4*c)*x-1/b*exp(-2*c-2*a)*a*exp(4*a+4*c)+exp(-2*c-2*a)*x+1/b*exp(-2*c-2*a)*a+1/2/b/(exp(2*a+2*c)-exp(2*b*x+2*a))*exp(4*a+4*c)+1/b/(exp(2*a+2*c)-exp(2*b*x+2*a))*exp(2*a+2*c)+1/2/b/(exp(2*a+2*c)-exp(2*b*x+2*a))+1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/2*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))/b*exp(-2*c-2*a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1313 vs. $2(53) = 106$.

Time = 0.10 (sec) , antiderivative size = 1313, normalized size of antiderivative = 29.84

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = \text{Too large to display}$$

input `integrate(cosh(b*x+a)^2*cosh(b*x-c)^2,x, algorithm="fricas")`

output

```

1/2*(2*b*x*cosh(b*x + a)^2*cosh(a + c)^6 - 12*b*x*cosh(b*x + a)^2*cosh(a +
c)*sinh(a + c)^5 + 2*b*x*cosh(b*x + a)^2*sinh(a + c)^6 - (2*b*x + 1)*cosh
(a + c)^4 + (30*b*x*cosh(b*x + a)^2*cosh(a + c)^2 - 2*b*x - 1)*sinh(a + c)
^4 - 4*(10*b*x*cosh(b*x + a)^2*cosh(a + c)^3 - (2*b*x + 1)*cosh(a + c))*si
nh(a + c)^3 + 2*(b*x*cosh(a + c)^6 - 6*b*x*cosh(a + c)^5*sinh(a + c) + 15*
b*x*cosh(a + c)^4*sinh(a + c)^2 - 20*b*x*cosh(a + c)^3*sinh(a + c)^3 + 15*
b*x*cosh(a + c)^2*sinh(a + c)^4 - 6*b*x*cosh(a + c)*sinh(a + c)^5 + b*x*si
nh(a + c)^6)*sinh(b*x + a)^2 + 2*(15*b*x*cosh(b*x + a)^2*cosh(a + c)^4 - 3
*(2*b*x + 1)*cosh(a + c)^2 - 1)*sinh(a + c)^2 - 2*cosh(a + c)^2 + (6*cosh(
b*x + a)^2*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)^2*sinh(a + c)^6 - (15
*cosh(b*x + a)^2*cosh(a + c)^2 - 1)*sinh(a + c)^4 + cosh(a + c)^4 + 4*(5*c
osh(b*x + a)^2*cosh(a + c)^3 - cosh(a + c))*sinh(a + c)^3 - (cosh(a + c)^6
- cosh(a + c)^2)*cosh(b*x + a)^2 - (cosh(a + c)^6 - 20*cosh(a + c)^3*sinh
(a + c)^3 + 15*cosh(a + c)^2*sinh(a + c)^4 - 6*cosh(a + c)*sinh(a + c)^5 +
sinh(a + c)^6 + (15*cosh(a + c)^4 - 1)*sinh(a + c)^2 - cosh(a + c)^2 - 2*
(3*cosh(a + c)^5 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((15*cosh(a
+ c)^4 - 1)*cosh(b*x + a)^2 - 6*cosh(a + c)^2)*sinh(a + c)^2 + 2*(20*cosh
(b*x + a)*cosh(a + c)^3*sinh(a + c)^3 - 15*cosh(b*x + a)*cosh(a + c)^2*si
nh(a + c)^4 + 6*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^5 - cosh(b*x + a)*sin
h(a + c)^6 - (15*cosh(a + c)^4 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(3*...

```


Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = \int \cosh^2(a + bx) \operatorname{csch}^2(bx - c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x-c)**2,x)`

output `Integral(cosh(a + b*x)**2*csch(b*x - c)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(53) = 106$.

Time = 0.05 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.05

$$\begin{aligned} \int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = & \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{2b} \\ & + \frac{(e^{(4a+4c)} - 1)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{2b} \\ & + \frac{(bx + a)e^{(2a+2c)}}{b} + \frac{e^{(4a+4c)} + 2e^{(2a+2c)} + 1}{2b(e^{(-2bx+2a+4c)} - e^{(2a+2c)})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*csch(b*x-c)^2,x, algorithm="maxima")`

output `1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b + (b*x + a)*e^(2*a + 2*c)/b + 1/2*(e^(4*a + 4*c) + 2*e^(2*a + 2*c) + 1)/(b*(e^(-2*b*x + 2*a + 4*c) - e^(2*a + 2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(53) = 106$.

Time = 0.15 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.52

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

$$= x e^{(-2a-2c)} + \frac{(e^{(4a+4c)} - 1) e^{(-2a-2c)} \log(|e^{(2bx)} - e^{(2c)}|)}{2b}$$

$$+ \frac{(e^{(2bx)} - e^{(2bx+4a+4c)} - 2e^{(2a+4c)} - 2e^{(2c)}) e^{(-2a-2c)}}{2b(e^{(2bx)} - e^{(2c)})}$$

input `integrate(cosh(b*x+a)^2*csch(b*x-c)^2,x, algorithm="giac")`

output `x*e^(-2*a - 2*c) + 1/2*(e^(4*a + 4*c) - 1)*e^(-2*a - 2*c)*log(abs(e^(2*b*x) - e^(2*c)))/b + 1/2*(e^(2*b*x) - e^(2*b*x + 4*a + 4*c) - 2*e^(2*a + 4*c) - 2*e^(2*c))*e^(-2*a - 2*c)/(b*(e^(2*b*x) - e^(2*c)))`

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.95

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx = x e^{-2a-2c} + \frac{\sinh(2a + 2c) \ln(e^{2a} e^{2bx} - e^{2a} e^{2c})}{b}$$

$$+ \frac{2 e^{2a+2c} \cosh(a + c)^2}{b (e^{2a+2c} - e^{2a+2bx})}$$

input `int(cosh(a + b*x)^2/sinh(c - b*x)^2,x)`

output `x*exp(- 2*a - 2*c) + (sinh(2*a + 2*c)*log(exp(2*a)*exp(2*b*x) - exp(2*a)*exp(2*c)))/b + (2*exp(2*a + 2*c)*cosh(a + c)^2)/(b*(exp(2*a + 2*c) - exp(2*a + 2*b*x)))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 256, normalized size of antiderivative = 5.82

$$\int \cosh^2(a + bx) \operatorname{csch}^2(c - bx) dx$$

$$= \frac{e^{2bx+4a+4c} \log(e^{bx} + e^c) + e^{2bx+4a+4c} \log(e^{bx} - e^c) - e^{2bx+4a+4c} - 2e^{2bx+2a+2c} - e^{2bx} \log(e^{bx} + e^c) - e^{2bx} \log(e^{bx} - e^c)}{2e^c}$$

input `int(cosh(b*x+a)^2*csch(b*x-c)^2,x)`

output

```
(e**(4*a + 2*b*x + 4*c)*log(e**(b*x) + e**c) + e**(4*a + 2*b*x + 4*c)*log(
e**(b*x) - e**c) - e**(4*a + 2*b*x + 4*c) - 2*e**(2*a + 2*b*x + 2*c) - e**
(2*b*x)*log(e**(b*x) + e**c) - e**(2*b*x)*log(e**(b*x) - e**c) + 2*e**(2*b
*x)*b*x - e**(2*b*x) - e**(4*a + 6*c)*log(e**(b*x) + e**c) - e**(4*a + 6*c
)*log(e**(b*x) - e**c) + e**(2*c)*log(e**(b*x) + e**c) + e**(2*c)*log(e**
(b*x) - e**c) - 2*e**(2*c)*b*x)/(2*e**(2*a + 2*c)*b*(e**(2*b*x) - e**(2*c))
)
```

3.162 $\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx$

Optimal result	1107
Mathematica [A] (verified)	1107
Rubi [F]	1108
Maple [B] (verified)	1108
Fricas [B] (verification not implemented)	1109
Sympy [F]	1109
Maxima [B] (verification not implemented)	1110
Giac [B] (verification not implemented)	1110
Mupad [F(-1)]	1111
Reduce [B] (verification not implemented)	1111

Optimal result

Integrand size = 18, antiderivative size = 85

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = -\frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh^2(a + c)}{2b} + \frac{\operatorname{arctanh}(\cosh(c - bx)) \cosh(2(a + c))}{b} + \frac{\cosh^2(a + c) \operatorname{coth}(c - bx) \operatorname{csch}(c - bx)}{2b} - \frac{\operatorname{csch}(c - bx) \sinh(2(a + c))}{b}$$

output

$$-1/2*\operatorname{arctanh}(\cosh(b*x-c))*\cosh(a+c)^2/b+\operatorname{arctanh}(\cosh(b*x-c))*\cosh(2*a+2*c)/b+1/2*\cosh(a+c)^2*\operatorname{coth}(b*x-c)*\operatorname{csch}(b*x-c)/b+\operatorname{csch}(b*x-c)*\sinh(2*a+2*c)/b$$

Mathematica [A] (verified)

Time = 4.32 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.93

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = \frac{2 \cosh^2(a + c) \operatorname{csch}^2\left(\frac{1}{2}(c - bx)\right) + 4(-1 + 3 \cosh(2(a + c))) \log\left(\cosh\left(\frac{1}{2}(c - bx)\right)\right) + 4(1 - 3 \cosh(2(a + c))) \operatorname{csch}^2\left(\frac{1}{2}(c - bx)\right)}{2b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c - b*x]^3,x]`

output $(2*\text{Cosh}[a + c]^2*\text{Csch}[(c - b*x)/2]^2 + 4*(-1 + 3*\text{Cosh}[2*(a + c)])*\text{Log}[\text{Cosh}[(c - b*x)/2]] + 4*(1 - 3*\text{Cosh}[2*(a + c)])*\text{Log}[-\text{Sinh}[(c - b*x)/2]] + 2*\text{Cosh}[a + c]^2*\text{Sech}[(c - b*x)/2]^2 - 8*\text{Csch}[c/2]*\text{Csch}[(c - b*x)/2]*\text{Sinh}[2*(a + c)]*\text{Sinh}[(b*x)/2] - 8*\text{Sech}[c/2]*\text{Sech}[(c - b*x)/2]*\text{Sinh}[2*(a + c)]*\text{Sinh}[(b*x)/2])/(16*b)$

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c - b*x]^3,x]`

output `$Aborted`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(89) = 178.

Time = 1.27 (sec) , antiderivative size = 295, normalized size of antiderivative = 3.47

method	result
risch	$-\frac{(3e^{6a+6c}-5e^{2bx+6a+4c}-2e^{4a+4c}-2e^{2bx+4a+2c}-5e^{2a+2c}+3e^{2bx+2a})e^{bx-c}}{4(e^{2a+2c}-e^{2bx+2a})^2b} + \frac{3\ln(e^a+c+e^{bx+a})e^{-2c-2a}e^{4a+4c}}{8b} - \frac{\ln(e^a+c)}{8b}$

input `int(-cosh(b*x+a)^2*csch(b*x-c)^3,x,method=_RETURNVERBOSE)`

output

```
-1/4/(exp(2*a+2*c)-exp(2*b*x+2*a))^2/b*(3*exp(6*a+6*c)-5*exp(2*b*x+6*a+4*c)
)-2*exp(4*a+4*c)-2*exp(2*b*x+4*a+2*c)-5*exp(2*a+2*c)+3*exp(2*b*x+2*a))*exp
(b*x-c)+3/8*ln(exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4*c)-1/4*ln(ex
p(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)+3/8*ln(exp(a+c)+exp(b*x+a)
)/b*exp(-2*c-2*a)-3/8*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(4*a+4*c
)+1/4*ln(-exp(a+c)+exp(b*x+a))/b*exp(-2*c-2*a)*exp(2*a+2*c)-3/8*ln(-exp(a+
c)+exp(b*x+a))/b*exp(-2*c-2*a)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5829 vs. 2(89) = 178.

Time = 0.17 (sec) , antiderivative size = 5829, normalized size of antiderivative = 68.58

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = \text{Too large to display}$$

input

```
integrate(-cosh(b*x+a)^2*csch(b*x-c)^3,x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = - \int \cosh^2(a + bx) \operatorname{csch}^3(bx - c) dx$$

input

```
integrate(-cosh(b*x+a)**2*csch(b*x-c)**3,x)
```

output

```
-Integral(cosh(a + b*x)**2*csch(b*x - c)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. $2(89) = 178$.

Time = 0.05 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.40

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx$$

$$= \frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(-bx+c)} + 1)}{8b}$$

$$- \frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(-bx+c)} - 1)}{8b}$$

$$- \frac{(5e^{(4a+4c)} + 2e^{(2a+2c)} - 3)e^{(-bx-a)} - (3e^{(6a+6c)} - 2e^{(4a+4c)} - 5e^{(2a+2c)})e^{(-3bx-3a)}}{4b(2e^{(-2bx+a+3c)} - e^{(-4bx+a+5c)} - e^{(a+c)})}$$

input `integrate(-cosh(b*x+a)^2*csch(b*x-c)^3,x, algorithm="maxima")`

output `1/8*(3*e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(e^(-b*x + c) + 1)/b - 1/8*(3*e^(4*a + 4*c) - 2*e^(2*a + 2*c) + 3)*e^(-2*a - 2*c)*log(e^(-b*x + c) - 1)/b - 1/4*((5*e^(4*a + 4*c) + 2*e^(2*a + 2*c) - 3)*e^(-b*x - a) - (3*e^(6*a + 6*c) - 2*e^(4*a + 4*c) - 5*e^(2*a + 2*c)))*e^(-3*b*x - 3*a))/(b*(2*e^(-2*b*x + a + 3*c) - e^(-4*b*x + a + 5*c) - e^(a + c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(89) = 178$.

Time = 0.15 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.22

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = \frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(bx)} + e^c)}{8b}$$

$$- \frac{(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(|e^{(bx)} - e^c|)}{8b}$$

$$- \frac{(3e^{(3bx)} - 5e^{(3bx+4a+4c)} - 2e^{(3bx+2a+2c)} + 3e^{(bx+4a+6c)} - 2e^{(bx+2a+4c)} - 5e^{(bx+2c)})e^{(-2a-c)}}{4b(e^{(2bx)} - e^{(2c)})^2}$$

input `integrate(-cosh(b*x+a)^2*csch(b*x-c)^3,x, algorithm="giac")`

output

$$\frac{1}{8}(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(e^{(bx)} + e^c)/b - \frac{1}{8}(3e^{(4a+4c)} - 2e^{(2a+2c)} + 3)e^{(-2a-2c)} \log(\text{abs}(e^{(bx)} - e^c))/b - \frac{1}{4}(3e^{(3bx)} - 5e^{(3bx+4a+4c)} - 2e^{(3bx+2a+2c)} + 3e^{(bx+4a+6c)} - 2e^{(bx+2a+4c)} - 5e^{(bx+2c)})e^{(-2a-c)}/(b(e^{(2bx)} - e^{(2c)}))^2$$
Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = \int \frac{\cosh(a + bx)^2}{\sinh(c - bx)^3} dx$$

input

`int(cosh(a + b*x)^2/sinh(c - b*x)^3, x)`

output

`int(cosh(a + b*x)^2/sinh(c - b*x)^3, x)`
Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 538, normalized size of antiderivative = 6.33

$$\int \cosh^2(a + bx) \operatorname{csch}^3(c - bx) dx = \frac{3e^{4bx+4a+4c} \log(e^{bx} + e^c) - 3e^{4bx+4a+4c} \log(e^{bx} - e^c) - 2e^{4bx+2a+2c} \log(e^{bx} + e^c) + 2e^{4bx+2a+2c} \log(e^{bx} - e^c)}{b}$$

input

`int(-cosh(b*x+a)^2*csch(b*x-c)^3, x)`

output

```
(3*e**(4*a + 4*b*x + 4*c)*log(e**(b*x) + e**c) - 3*e**(4*a + 4*b*x + 4*c)*
log(e**(b*x) - e**c) - 2*e**(2*a + 4*b*x + 2*c)*log(e**(b*x) + e**c) + 2*e
**(2*a + 4*b*x + 2*c)*log(e**(b*x) - e**c) + 3*e**(4*b*x)*log(e**(b*x) + e
**c) - 3*e**(4*b*x)*log(e**(b*x) - e**c) + 10*e**(4*a + 3*b*x + 5*c) + 4*e
**(2*a + 3*b*x + 3*c) - 6*e**(3*b*x + c) - 6*e**(4*a + 2*b*x + 6*c)*log(e*
*(b*x) + e**c) + 6*e**(4*a + 2*b*x + 6*c)*log(e**(b*x) - e**c) + 4*e**(2*a
+ 2*b*x + 4*c)*log(e**(b*x) + e**c) - 4*e**(2*a + 2*b*x + 4*c)*log(e**(b*
x) - e**c) - 6*e**(2*b*x + 2*c)*log(e**(b*x) + e**c) + 6*e**(2*b*x + 2*c)*
log(e**(b*x) - e**c) - 6*e**(4*a + b*x + 7*c) + 4*e**(2*a + b*x + 5*c) + 1
0*e**(b*x + 3*c) + 3*e**(4*a + 8*c)*log(e**(b*x) + e**c) - 3*e**(4*a + 8*c
)*log(e**(b*x) - e**c) - 2*e**(2*a + 6*c)*log(e**(b*x) + e**c) + 2*e**(2*a
+ 6*c)*log(e**(b*x) - e**c) + 3*e**(4*c)*log(e**(b*x) + e**c) - 3*e**(4*c
)*log(e**(b*x) - e**c))/(8*e**(2*a + 2*c)*b*(e**(4*b*x) - 2*e**(2*b*x + 2*
c) + e**(4*c)))
```

3.163 $\int \cosh^2(a + bx)\operatorname{csch}^4(c - bx) dx$

Optimal result	1113
Mathematica [A] (verified)	1113
Rubi [F]	1114
Maple [A] (verified)	1114
Fricas [B] (verification not implemented)	1115
Sympy [F]	1116
Maxima [B] (verification not implemented)	1116
Giac [A] (verification not implemented)	1117
Mupad [F(-1)]	1117
Reduce [B] (verification not implemented)	1118

Optimal result

Integrand size = 18, antiderivative size = 80

$$\int \cosh^2(a + bx)\operatorname{csch}^4(c - bx) dx = -\frac{\cosh^2(a + c)\operatorname{coth}(c - bx)}{b} + \frac{\cosh(2(a + c))\operatorname{coth}(c - bx)}{b} + \frac{\cosh^2(a + c)\operatorname{coth}^3(c - bx)}{3b} - \frac{\operatorname{csch}^2(c - bx)\sinh(2(a + c))}{2b}$$

```
output cosh(a+c)^2*coth(b*x-c)/b-cosh(2*a+2*c)*coth(b*x-c)/b-1/3*cosh(a+c)^2*coth
(b*x-c)^3/b-1/2*csch(b*x-c)^2*sinh(2*a+2*c)/b
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \cosh^2(a + bx)\operatorname{csch}^4(c - bx) dx = \frac{\operatorname{csch}(c)\operatorname{csch}^3(c - bx)(3\sinh(bx) + \sinh(2c - 3bx)) - \sinh(2a + 4c - 3bx) + 3\sinh(2a + 2c - bx) - 3\sinh(2a + c - bx)}{12b}$$

input `Integrate[Cosh[a + b*x]^2*Csch[c - b*x]^4,x]`

output `(Csch[c]*Csch[c - b*x]^3*(3*Sinh[b*x] + Sinh[2*c - 3*b*x] - Sinh[2*a + 4*c - 3*b*x] + 3*Sinh[2*a + 2*c - b*x] - 3*Sinh[2*a + b*x] + Sinh[2*a + 3*b*x]))/(12*b)`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx$$

↓ 7299

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx$$

input `Int[Cosh[a + b*x]^2*Csch[c - b*x]^4,x]`

output `$Aborted`

Maple [A] (verified)

Time = 1.53 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.78

method	result	size
parallelrisch	$\frac{\operatorname{sech}\left(\frac{bx}{2} - \frac{c}{2}\right)^3 \operatorname{csch}\left(\frac{bx}{2} - \frac{c}{2}\right)^3 (-2 \cosh(3bx + 2a - c) + \cosh(3bx - 3c) - 3 \cosh(bx - c))}{96b}$	62
risch	$\frac{2(e^{4a+4c} - 3e^{2bx+4a+2c} - e^{2a+2c} + 3e^{4bx+4a} + 3e^{2bx+2a} + 1)e^{4a+4c}}{3(e^{2a+2c} - e^{2bx+2a})^3 b}$	92

input `int(cosh(b*x+a)^2*csch(b*x-c)^4,x,method=_RETURNVERBOSE)`

output

```
1/96*sech(1/2*b*x-1/2*c)^3*csc(1/2*b*x-1/2*c)^3*(-2*cosh(3*b*x+2*a-c)+cos
h(3*b*x-3*c)-3*cosh(b*x-c))/b
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1015 vs. $2(84) = 168$.

Time = 0.10 (sec) , antiderivative size = 1015, normalized size of antiderivative = 12.69

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx = \text{Too large to display}$$

input

```
integrate(cosh(b*x+a)^2*csc(b*x-c)^4,x, algorithm="fricas")
```

output

```
-2/3*(4*cosh(b*x + a)^2*cosh(a + c)*sinh(a + c)^3 + (5*cosh(b*x + a)^2 + 3
)*sinh(a + c)^4 + 3*cosh(a + c)^4 + (5*cosh(a + c)^4 - cosh(a + c)^2)*cosh
(b*x + a)^2 + (5*cosh(a + c)^4 + 4*cosh(a + c)*sinh(a + c)^3 + 5*sinh(a +
c)^4 - (2*cosh(a + c)^2 + 1)*sinh(a + c)^2 - cosh(a + c)^2 + 2*(2*cosh(a +
c)^3 - cosh(a + c))*sinh(a + c))*sinh(b*x + a)^2 - ((2*cosh(a + c)^2 + 1)
*cosh(b*x + a)^2 + 6*cosh(a + c)^2 + 3)*sinh(a + c)^2 - 3*cosh(a + c)^2 -
2*(4*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^3 - cosh(b*x + a)*sinh(a + c)^4
+ (10*cosh(a + c)^2 - 1)*cosh(b*x + a)*sinh(a + c)^2 + 2*(2*cosh(a + c)^3
- cosh(a + c))*cosh(b*x + a)*sinh(a + c) - (cosh(a + c)^4 + cosh(a + c)^2
)*cosh(b*x + a))*sinh(b*x + a) + 2*((2*cosh(a + c)^3 - cosh(a + c))*cosh(b
*x + a)^2 - 3*cosh(a + c))*sinh(a + c))/(b*cosh(b*x + a)^4*cosh(a + c)^4 -
4*b*cosh(b*x + a)^2*cosh(a + c)^4 + 3*b*cosh(a + c)^4 + (b*cosh(a + c)^4
- 4*b*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(a + c)^2*sinh(a + c)^2 - 4*b*co
sh(a + c)*sinh(a + c)^3 + b*sinh(a + c)^4)*sinh(b*x + a)^4 + (b*cosh(b*x +
a)^4 + 4*b*cosh(b*x + a)^2 + 3*b)*sinh(a + c)^4 + 4*(b*cosh(b*x + a)*cosh
(a + c)^4 - 4*b*cosh(b*x + a)*cosh(a + c)^3*sinh(a + c) + 6*b*cosh(b*x + a
)*cosh(a + c)^2*sinh(a + c)^2 - 4*b*cosh(b*x + a)*cosh(a + c)*sinh(a + c)^
3 + b*cosh(b*x + a)*sinh(a + c)^4)*sinh(b*x + a)^3 - 4*(b*cosh(b*x + a)^4*
cosh(a + c) + b*cosh(b*x + a)^2*cosh(a + c))*sinh(a + c)^3 + 2*(3*b*cosh(b
*x + a)^2*cosh(a + c)^4 + 18*b*cosh(b*x + a)^2*cosh(a + c)^2*sinh(a + c)...
```

Sympy [F]

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx = \int \cosh^2(a + bx) \operatorname{csch}^4(bx - c) dx$$

input `integrate(cosh(b*x+a)**2*csch(b*x-c)**4,x)`

output `Integral(cosh(a + b*x)**2*csch(b*x - c)**4, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 344 vs. $2(84) = 168$.

Time = 0.04 (sec) , antiderivative size = 344, normalized size of antiderivative = 4.30

$$\begin{aligned} & \int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx \\ &= \frac{2(e^{4a+4c} - e^{2a+2c})e^{-2bx-2a}}{b(3e^{-2bx+2a+4c} - 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} - e^{2a+2c})} \\ &+ \frac{2e^{-4bx+4c}}{b(3e^{-2bx+2a+4c} - 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} - e^{2a+2c})} \\ &+ \frac{2e^{4a+4c}}{3b(3e^{-2bx+2a+4c} - 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} - e^{2a+2c})} \\ &- \frac{2e^{2a+2c}}{3b(3e^{-2bx+2a+4c} - 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} - e^{2a+2c})} \\ &+ \frac{2}{3b(3e^{-2bx+2a+4c} - 3e^{-4bx+2a+6c} + e^{-6bx+2a+8c} - e^{2a+2c})} \end{aligned}$$

input `integrate(cosh(b*x+a)^2*csch(b*x-c)^4,x, algorithm="maxima")`

output

$$2*(e^{(4*a + 4*c)} - e^{(2*a + 2*c)})*e^{(-2*b*x - 2*a)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} - e^{(2*a + 2*c)})} + 2*e^{(-4*b*x + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} - e^{(2*a + 2*c)})} + 2/3*e^{(4*a + 4*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} - e^{(2*a + 2*c)})} - 2/3*e^{(2*a + 2*c)/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} - e^{(2*a + 2*c)})} + 2/3/(b*(3*e^{(-2*b*x + 2*a + 4*c)} - 3*e^{(-4*b*x + 2*a + 6*c)} + e^{(-6*b*x + 2*a + 8*c)} - e^{(2*a + 2*c)}))$$
Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.10

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx = \frac{2(3e^{(4bx+4a+4c)} - 3e^{(2bx+4a+6c)} + 3e^{(2bx+2a+4c)} + e^{(4a+8c)} - e^{(2a+6c)} + e^{(4c)})e^{(-2a)}}{3b(e^{(2bx)} - e^{(2c)})^3}$$

input

```
integrate(cosh(b*x+a)^2*csch(b*x-c)^4,x, algorithm="giac")
```

output

$$-2/3*(3*e^{(4*b*x + 4*a + 4*c)} - 3*e^{(2*b*x + 4*a + 6*c)} + 3*e^{(2*b*x + 2*a + 4*c)} + e^{(4*a + 8*c)} - e^{(2*a + 6*c)} + e^{(4*c)})*e^{(-2*a)/(b*(e^{(2*b*x)} - e^{(2*c)})^3)}$$
Mupad [F(-1)]

Timed out.

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx = \int \frac{\cosh(a + bx)^2}{\sinh(c - bx)^4} dx$$

input

```
int(cosh(a + b*x)^2/sinh(c - b*x)^4,x)
```

output

```
int(cosh(a + b*x)^2/sinh(c - b*x)^4, x)
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.26

$$\int \cosh^2(a + bx) \operatorname{csch}^4(c - bx) dx = \frac{2e^{2c}(-e^{6bx+4a} - 3e^{2bx+2a+2c} + e^{2a+4c} - e^{2c})}{3e^{2a}b(e^{6bx} - 3e^{4bx+2c} + 3e^{2bx+4c} - e^{6c})}$$

input `int(cosh(b*x+a)^2*csch(b*x-c)^4,x)`output `(2*e**(2*c)*(- e**(4*a + 6*b*x) - 3*e**(2*a + 2*b*x + 2*c) + e**(2*a + 4*c) - e**(2*c)))/(3*e**(2*a)*b*(e**(6*b*x) - 3*e**(4*b*x + 2*c) + 3*e**(2*b*x + 4*c) - e**(6*c)))`

3.164 $\int \tanh(a + bx) \tanh(c + bx) dx$

Optimal result	1119
Mathematica [A] (verified)	1119
Rubi [A] (verified)	1120
Maple [B] (verified)	1121
Fricas [B] (verification not implemented)	1122
Sympy [F]	1122
Maxima [B] (verification not implemented)	1123
Giac [B] (verification not implemented)	1123
Mupad [B] (verification not implemented)	1124
Reduce [B] (verification not implemented)	1124

Optimal result

Integrand size = 13, antiderivative size = 37

$$\int \tanh(a + bx) \tanh(c + bx) dx = x - \frac{\coth(a - c) \log(\cosh(a + bx))}{b} + \frac{\coth(a - c) \log(\cosh(c + bx))}{b}$$

output `x-coth(a-c)*ln(cosh(b*x+a))/b+coth(a-c)*ln(cosh(b*x+c))/b`

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \tanh(a + bx) \tanh(c + bx) dx = x + \frac{\coth(a - c)(-\log(\cosh(a + bx)) + \log(\cosh(c + bx)))}{b}$$

input `Integrate[Tanh[a + b*x]*Tanh[c + b*x],x]`

output `x + (Coth[a - c]*(-Log[Cosh[a + b*x]] + Log[Cosh[c + b*x]]))/b`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6180, 6178, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(a + bx) \tanh(bx + c) dx \\
 & \quad \downarrow \text{6180} \\
 & x - \cosh(a - c) \int \operatorname{sech}(a + bx) \operatorname{sech}(c + bx) dx \\
 & \quad \downarrow \text{6178} \\
 & x - \cosh(a - c) (\operatorname{csch}(a - c) \int \tanh(a + bx) dx - \operatorname{csch}(a - c) \int \tanh(c + bx) dx) \\
 & \quad \downarrow \text{3042} \\
 & x - \cosh(a - c) (\operatorname{csch}(a - c) \int -i \tan(ia + ibx) dx - \operatorname{csch}(a - c) \int -i \tan(ic + ibx) dx) \\
 & \quad \downarrow \text{26} \\
 & x - \cosh(a - c) (i \operatorname{csch}(a - c) \int \tan(ic + ibx) dx - i \operatorname{csch}(a - c) \int \tan(ia + ibx) dx) \\
 & \quad \downarrow \text{3956} \\
 & x - \cosh(a - c) \left(\frac{\operatorname{csch}(a - c) \log(\cosh(a + bx))}{b} - \frac{\operatorname{csch}(a - c) \log(\cosh(bx + c))}{b} \right)
 \end{aligned}$$

input

```
Int[Tanh[a + b*x]*Tanh[c + b*x],x]
```

output

```
x - Cosh[a - c]*((Csch[a - c]*Log[Cosh[a + b*x]])/b - (Csch[a - c]*Log[Cosh[c + b*x]])/b)
```

Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 6178 `Int[Sech[(a_.) + (b_.)*(x_)]*Sech[(c_) + (d_.)*(x_)], x_Symbol] := Simp[-Csch[(b*c - a*d)/d] Int[Tanh[a + b*x], x], x] + Simp[Csch[(b*c - a*d)/b] Int[Tanh[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`
- rule 6180 `Int[Tanh[(a_.) + (b_.)*(x_)]*Tanh[(c_) + (d_.)*(x_)], x_Symbol] := Simp[b*(x/d), x] - Simp[(b/d)*Cosh[(b*c - a*d)/d] Int[Sech[a + b*x]*Sech[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. $2(37) = 74$.

Time = 0.12 (sec) , antiderivative size = 151, normalized size of antiderivative = 4.08

method	result	size
risch	$x - \frac{\ln(e^{2bx+2a}+1)e^{2a}}{b(e^{2a}-e^{2c})} - \frac{\ln(e^{2bx+2a}+1)e^{2c}}{b(e^{2a}-e^{2c})} + \frac{\ln(e^{2bx+2a}+e^{2a-2c})e^{2a}}{b(e^{2a}-e^{2c})} + \frac{\ln(e^{2bx+2a}+e^{2a-2c})e^{2c}}{b(e^{2a}-e^{2c})}$	151

input `int(tanh(b*x+a)*tanh(b*x+c),x,method=_RETURNVERBOSE)`

output

```
x-1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2*a)+1)*exp(2*a)-1/b/(exp(2*a)-exp(
2*c))*ln(exp(2*b*x+2*a)+1)*exp(2*c)+1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2
*a)+exp(2*a-2*c))*exp(2*a)+1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2*a)+exp(2
*a-2*c))*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. $2(37) = 74$.

Time = 0.10 (sec) , antiderivative size = 259, normalized size of antiderivative = 7.00

$$\int \tanh(a + bx) \tanh(c + bx) dx$$

$$= \frac{bx \cosh(-a + c)^2 - 2bx \cosh(-a + c) \sinh(-a + c) + bx \sinh(-a + c)^2 - bx - (\cosh(-a + c)^2 - 2 \cosh(-a + c) \sinh(-a + c) + \sinh(-a + c)^2 + 1) \log(2(\cosh(bx + c) \cosh(-a + c) - \sinh(bx + c) \sinh(-a + c)) / (\cosh(bx + c) \cosh(-a + c) - (\cosh(-a + c) + \sinh(-a + c)) \sinh(bx + c) + \cosh(bx + c) \sinh(-a + c))) + (\cosh(-a + c)^2 - 2 \cosh(-a + c) \sinh(-a + c) + \sinh(-a + c)^2 + 1) \log(2 \cosh(bx + c) / (\cosh(bx + c) - \sinh(bx + c)))}{(b \cosh(-a + c)^2 - 2b \cosh(-a + c) \sinh(-a + c) + b \sinh(-a + c)^2 - b)}$$

input

```
integrate(tanh(b*x+a)*tanh(b*x+c),x, algorithm="fricas")
```

output

```
(b*x*cosh(-a + c)^2 - 2*b*x*cosh(-a + c)*sinh(-a + c) + b*x*sinh(-a + c)^2
- b*x - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 +
1)*log(2*(cosh(b*x + c)*cosh(-a + c) - sinh(b*x + c)*sinh(-a + c))/(cosh(b
*x + c)*cosh(-a + c) - (cosh(-a + c) + sinh(-a + c))*sinh(b*x + c) + cosh(
b*x + c)*sinh(-a + c))) + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) +
sinh(-a + c)^2 + 1)*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c)))/
(b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^2 - b)
```

Sympy [F]

$$\int \tanh(a + bx) \tanh(c + bx) dx = \int \tanh(a + bx) \tanh(bx + c) dx$$

input

```
integrate(tanh(b*x+a)*tanh(b*x+c),x)
```

output

```
Integral(tanh(a + b*x)*tanh(b*x + c), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(37) = 74.

Time = 0.12 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.24

$$\int \tanh(a + bx) \tanh(c + bx) dx = x + \frac{a}{b} - \frac{(e^{(2a)} + e^{(2c)}) \log(e^{(-2bx-2a)} + 1)}{b(e^{(2a)} - e^{(2c)})} + \frac{(e^{(2a)} + e^{(2c)}) \log(e^{(-2bx)} + e^{(2c)})}{b(e^{(2a)} - e^{(2c)})}$$

input `integrate(tanh(b*x+a)*tanh(b*x+c),x, algorithm="maxima")`

output `x + a/b - (e^(2*a) + e^(2*c))*log(e^(-2*b*x - 2*a) + 1)/(b*(e^(2*a) - e^(2*c))) + (e^(2*a) + e^(2*c))*log(e^(-2*b*x) + e^(2*c))/(b*(e^(2*a) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(37) = 74.

Time = 0.13 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.57

$$\int \tanh(a + bx) \tanh(c + bx) dx = x - \frac{(e^{(4a)} + e^{(2a+2c)}) \log(e^{(2bx+2a)} + 1)}{be^{(4a)} - be^{(2a+2c)}} + \frac{(e^{(2a+2c)} + e^{(4c)}) \log(e^{(2bx+2c)} + 1)}{be^{(2a+2c)} - be^{(4c)}}$$

input `integrate(tanh(b*x+a)*tanh(b*x+c),x, algorithm="giac")`

output `x - (e^(4*a) + e^(2*a + 2*c))*log(e^(2*b*x + 2*a) + 1)/(b*e^(4*a) - b*e^(2*a + 2*c)) + (e^(2*a + 2*c) + e^(4*c))*log(e^(2*b*x + 2*c) + 1)/(b*e^(2*a + 2*c) - b*e^(4*c))`

Mupad [B] (verification not implemented)

Time = 1.55 (sec) , antiderivative size = 115, normalized size of antiderivative = 3.11

$$\int \tanh(a + bx) \tanh(c + bx) dx$$

$$= x - \frac{\ln(4e^{4a} + 4e^{6a}e^{2bx} + 4e^{2a}e^{2c} + 4e^{4a}e^{2c}e^{2bx}) \coth(a - c)}{b}$$

$$+ \frac{\ln(4e^{4a} + 4e^{2a}e^{2c} + 4e^{2a}e^{4c}e^{2bx} + 4e^{4a}e^{2c}e^{2bx}) \coth(a - c)}{b}$$

input `int(tanh(a + b*x)*tanh(c + b*x),x)`output `x - (log(4*exp(4*a) + 4*exp(6*a)*exp(2*b*x) + 4*exp(2*a)*exp(2*c) + 4*exp(4*a)*exp(2*c)*exp(2*b*x))*coth(a - c))/b + (log(4*exp(4*a) + 4*exp(2*a)*exp(2*c) + 4*exp(2*a)*exp(4*c)*exp(2*b*x) + 4*exp(4*a)*exp(2*c)*exp(2*b*x))*coth(a - c))/b`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 115, normalized size of antiderivative = 3.11

$$\int \tanh(a + bx) \tanh(c + bx) dx$$

$$= \frac{-e^{2a} \log(e^{2bx+2a} + 1) + e^{2a} \log(e^{2bx+2c} + 1) + e^{2a}bx - e^{2c} \log(e^{2bx+2a} + 1) + e^{2c} \log(e^{2bx+2c} + 1) - e^{2c}bx}{b(e^{2a} - e^{2c})}$$

input `int(tanh(b*x+a)*tanh(b*x+c),x)`output `(- e**(2*a)*log(e**(2*a + 2*b*x) + 1) + e**(2*a)*log(e**(2*b*x + 2*c) + 1) + e**(2*a)*b*x - e**(2*c)*log(e**(2*a + 2*b*x) + 1) + e**(2*c)*log(e**(2*b*x + 2*c) + 1) - e**(2*c)*b*x)/(b*(e**(2*a) - e**(2*c)))`

3.165 $\int \tanh(c - bx) \tanh(a + bx) dx$

Optimal result	1125
Mathematica [A] (verified)	1125
Rubi [A] (verified)	1126
Maple [B] (verified)	1127
Fricas [B] (verification not implemented)	1128
Sympy [F]	1128
Maxima [B] (verification not implemented)	1129
Giac [B] (verification not implemented)	1129
Mupad [B] (verification not implemented)	1130
Reduce [B] (verification not implemented)	1130

Optimal result

Integrand size = 14, antiderivative size = 36

$$\int \tanh(c - bx) \tanh(a + bx) dx = -x - \frac{\coth(a + c) \log(\cosh(c - bx))}{b} + \frac{\coth(a + c) \log(\cosh(a + bx))}{b}$$

output

```
-x-coth(a+c)*ln(cosh(b*x-c))/b+coth(a+c)*ln(cosh(b*x+a))/b
```

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \tanh(c - bx) \tanh(a + bx) dx = -x + \frac{\coth(a + c)(-\log(\cosh(c - bx)) + \log(\cosh(a + bx)))}{b}$$

input

```
Integrate[Tanh[c - b*x]*Tanh[a + b*x],x]
```

output

```
-x + (Coth[a + c]*(-Log[Cosh[c - b*x]] + Log[Cosh[a + b*x]]))/b
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.17, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6180, 6178, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tanh(a + bx) \tanh(c - bx) dx \\
 & \quad \downarrow \text{6180} \\
 & \cosh(a + c) \int \operatorname{sech}(c - bx) \operatorname{sech}(a + bx) dx - x \\
 & \quad \downarrow \text{6178} \\
 & \cosh(a + c) (\operatorname{csch}(a + c) \int \tanh(c - bx) dx + \operatorname{csch}(a + c) \int \tanh(a + bx) dx) - x \\
 & \quad \downarrow \text{3042} \\
 & -x + \cosh(a + c) (\operatorname{csch}(a + c) \int -i \tan(ic - ibx) dx + \operatorname{csch}(a + c) \int -i \tan(ia + ibx) dx) \\
 & \quad \downarrow \text{26} \\
 & -x + \cosh(a + c) (-i \operatorname{csch}(a + c) \int \tan(ic - ibx) dx - i \operatorname{csch}(a + c) \int \tan(ia + ibx) dx) \\
 & \quad \downarrow \text{3956} \\
 & \cosh(a + c) \left(\frac{\operatorname{csch}(a + c) \log(\cosh(a + bx))}{b} - \frac{\operatorname{csch}(a + c) \log(\cosh(c - bx))}{b} \right) - x
 \end{aligned}$$

input

```
Int[Tanh[c - b*x]*Tanh[a + b*x],x]
```

output

```
-x + Cosh[a + c]*(-((Csch[a + c]*Log[Cosh[c - b*x]])/b) + (Csch[a + c]*Log[Cosh[a + b*x]])/b)
```

Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 6178 `Int[Sech[(a_.) + (b_.)*(x_)]*Sech[(c_) + (d_.)*(x_)], x_Symbol] := Simp[-Csch[(b*c - a*d)/d] Int[Tanh[a + b*x], x], x] + Simp[Csch[(b*c - a*d)/b] Int[Tanh[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`
- rule 6180 `Int[Tanh[(a_.) + (b_.)*(x_)]*Tanh[(c_) + (d_.)*(x_)], x_Symbol] := Simp[b*(x/d), x] - Simp[(b/d)*Cosh[(b*c - a*d)/d] Int[Sech[a + b*x]*Sech[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(37) = 74$.

Time = 0.11 (sec) , antiderivative size = 149, normalized size of antiderivative = 4.14

method	result	size
risch	$-x - \frac{\ln(e^{2a+2c} + e^{2bx+2a})e^{2a+2c}}{b(e^{2a+2c}-1)} - \frac{\ln(e^{2a+2c} + e^{2bx+2a})}{b(e^{2a+2c}-1)} + \frac{\ln(e^{2bx+2a} + 1)e^{2a+2c}}{b(e^{2a+2c}-1)} + \frac{\ln(e^{2bx+2a} + 1)}{b(e^{2a+2c}-1)}$	149

input `int(-tanh(b*x-c)*tanh(b*x+a), x, method=_RETURNVERBOSE)`

output

```
-x-1/b/(exp(2*a+2*c)-1)*ln(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(2*a+2*c)-1/b/(
exp(2*a+2*c)-1)*ln(exp(2*a+2*c)+exp(2*b*x+2*a))+1/b/(exp(2*a+2*c)-1)*ln(ex
p(2*b*x+2*a)+1)*exp(2*a+2*c)+1/b/(exp(2*a+2*c)-1)*ln(exp(2*b*x+2*a)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. $2(37) = 74$.

Time = 0.10 (sec) , antiderivative size = 216, normalized size of antiderivative = 6.00

$$\int \tanh(c - bx) \tanh(a + bx) dx = \frac{bx \cosh(a + c)^2 - 2bx \cosh(a + c) \sinh(a + c) + bx \sinh(a + c)^2 - bx - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log\left(\frac{2(\cosh(bx + a) \cosh(a + c) - \sinh(bx + a) \sinh(a + c))}{\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c)}\right) + (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log\left(\frac{2 \cosh(bx + a)}{\cosh(bx + a) - \sinh(bx + a)}\right)}{(b \cosh(a + c)^2 - 2b \cosh(a + c) \sinh(a + c) + b \sinh(a + c)^2 - b)}$$

input

```
integrate(-tanh(b*x-c)*tanh(b*x+a),x, algorithm="fricas")
```

output

```
-(b*x*cosh(a + c)^2 - 2*b*x*cosh(a + c)*sinh(a + c) + b*x*sinh(a + c)^2 -
b*x - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 + 1)*log(
2*(cosh(b*x + a)*cosh(a + c) - sinh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*c
osh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sin
h(a + c))) + (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 +
1)*log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/(b*cosh(a + c)^2
- 2*b*cosh(a + c)*sinh(a + c) + b*sinh(a + c)^2 - b)
```

Sympy [F]

$$\int \tanh(c - bx) \tanh(a + bx) dx = - \int \tanh(a + bx) \tanh(bx - c) dx$$

input

```
integrate(-tanh(b*x-c)*tanh(b*x+a),x)
```

output

```
-Integral(tanh(a + b*x)*tanh(b*x - c), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(37) = 74.

Time = 0.13 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.42

$$\int \tanh(c - bx) \tanh(a + bx) dx = -x - \frac{a}{b} + \frac{(e^{(2a+2c)} + 1) \log(e^{(-2bx-2a)} + 1)}{b(e^{(2a+2c)} - 1)} - \frac{(e^{(2a+2c)} + 1) \log(e^{(-2bx+2c)} + 1)}{b(e^{(2a+2c)} - 1)}$$

input `integrate(-tanh(b*x-c)*tanh(b*x+a),x, algorithm="maxima")`

output `-x - a/b + (e^(2*a + 2*c) + 1)*log(e^(-2*b*x - 2*a) + 1)/(b*(e^(2*a + 2*c) - 1)) - (e^(2*a + 2*c) + 1)*log(e^(-2*b*x + 2*c) + 1)/(b*(e^(2*a + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. 2(37) = 74.

Time = 0.19 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.50

$$\int \tanh(c - bx) \tanh(a + bx) dx = -x - \frac{(e^{(2a+2c)} + 1) \log(e^{(2bx)} + e^{(2c)})}{be^{(2a+2c)} - b} - \frac{(e^{(2a)} + e^{(4a+2c)}) \log(e^{(2bx+2a)} + 1)}{be^{(2a)} - be^{(4a+2c)}}$$

input `integrate(-tanh(b*x-c)*tanh(b*x+a),x, algorithm="giac")`

output `-x - (e^(2*a + 2*c) + 1)*log(e^(2*b*x) + e^(2*c))/(b*e^(2*a + 2*c) - b) - (e^(2*a) + e^(4*a + 2*c))*log(e^(2*b*x + 2*a) + 1)/(b*e^(2*a) - b*e^(4*a + 2*c))`

Mupad [B] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 121, normalized size of antiderivative = 3.36

$$\int \tanh(c - bx) \tanh(a + bx) dx$$

$$= \frac{\coth(a + c) \ln(4e^{2a} e^{2c} + 4e^{4a} e^{4c} + 4e^{4a} e^{2c} e^{2bx} + 4e^{6a} e^{4c} e^{2bx})}{b} - \frac{\coth(a + c) \ln(4e^{2a} e^{2bx} + 4e^{2a} e^{2c} + 4e^{4a} e^{4c} + 4e^{4a} e^{2c} e^{2bx})}{b} - x$$

input `int(tanh(a + b*x)*tanh(c - b*x),x)`output `(coth(a + c)*log(4*exp(2*a)*exp(2*c) + 4*exp(4*a)*exp(4*c) + 4*exp(4*a)*exp(2*c)*exp(2*b*x) + 4*exp(6*a)*exp(4*c)*exp(2*b*x)))/b - (coth(a + c)*log(4*exp(2*a)*exp(2*b*x) + 4*exp(2*a)*exp(2*c) + 4*exp(4*a)*exp(4*c) + 4*exp(4*a)*exp(2*c)*exp(2*b*x)))/b - x`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 109, normalized size of antiderivative = 3.03

$$\int \tanh(c - bx) \tanh(a + bx) dx$$

$$= \frac{e^{2a+2c} \log(e^{2bx+2a} + 1) - e^{2a+2c} \log(e^{2bx} + e^{2c}) - e^{2a+2c} bx + \log(e^{2bx+2a} + 1) - \log(e^{2bx} + e^{2c}) + bx}{b(e^{2a+2c} - 1)}$$

input `int(-tanh(b*x-c)*tanh(b*x+a),x)`output `(e**(2*a + 2*c)*log(e**(2*a + 2*b*x) + 1) - e**(2*a + 2*c)*log(e**(2*b*x) + e**(2*c)) - e**(2*a + 2*c)*b*x + log(e**(2*a + 2*b*x) + 1) - log(e**(2*b*x) + e**(2*c)) + b*x)/(b*(e**(2*a + 2*c) - 1))`

3.166 $\int \coth(a + bx) \coth(c + bx) dx$

Optimal result	1131
Mathematica [A] (verified)	1131
Rubi [C] (verified)	1132
Maple [B] (verified)	1133
Fricas [B] (verification not implemented)	1134
Sympy [F]	1134
Maxima [B] (verification not implemented)	1135
Giac [B] (verification not implemented)	1135
Mupad [B] (verification not implemented)	1136
Reduce [B] (verification not implemented)	1136

Optimal result

Integrand size = 13, antiderivative size = 37

$$\int \coth(a + bx) \coth(c + bx) dx = x - \frac{\coth(a - c) \log(\sinh(a + bx))}{b} + \frac{\coth(a - c) \log(\sinh(c + bx))}{b}$$

output `x-coth(a-c)*ln(sinh(b*x+a))/b+coth(a-c)*ln(sinh(b*x+c))/b`

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \coth(a + bx) \coth(c + bx) dx = x + \frac{\coth(a - c)(-\log(\sinh(a + bx)) + \log(\sinh(c + bx)))}{b}$$

input `Integrate[Coth[a + b*x]*Coth[c + b*x],x]`

output `x + (Coth[a - c]*(-Log[Sinh[a + b*x]] + Log[Sinh[c + b*x]]))/b`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {6181, 6179, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth(a + bx) \coth(bx + c) dx \\
 & \quad \downarrow \text{6181} \\
 & \cosh(a - c) \int \operatorname{csch}(a + bx) \operatorname{csch}(c + bx) dx + x \\
 & \quad \downarrow \text{6179} \\
 & \cosh(a - c) (\operatorname{csch}(a - c) \int \coth(c + bx) dx - \operatorname{csch}(a - c) \int \coth(a + bx) dx) + x \\
 & \quad \downarrow \text{3042} \\
 & c) \left(\operatorname{csch}(a - c) \int -i \tan\left(ic + ibx + \frac{\pi}{2}\right) dx - \operatorname{csch}(a - c) \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow \text{26} \\
 & c) \left(\operatorname{icsch}(a - c) \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right) dx - \operatorname{icsch}(a - c) \int \tan\left(\frac{1}{2}(2ic + \pi) + ibx\right) dx \right) \\
 & \quad \downarrow \text{3956} \\
 & x + \cosh(a - c) \left(\frac{\operatorname{csch}(a - c) \log(-i \sinh(bx + c))}{b} - \frac{\operatorname{csch}(a - c) \log(-i \sinh(a + bx))}{b} \right)
 \end{aligned}$$

input `Int[Coth[a + b*x]*Coth[c + b*x],x]`

output `x + Cosh[a - c]*(-((Csch[a - c]*Log[(-I)*Sinh[a + b*x]])/b) + (Csch[a - c]*Log[(-I)*Sinh[c + b*x]])/b)`

Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6179 `Int[Csch[(a_.) + (b_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Csch[(b*c - a*d)/b] Int[Coth[a + b*x], x], x] - Simp[Csch[(b*c - a*d)/d] Int[Coth[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

rule 6181 `Int[Coth[(a_.) + (b_.)*(x_)]*Coth[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[b*(x/d), x] + Simp[Cosh[(b*c - a*d)/d] Int[Csch[a + b*x]*Csch[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 154 vs. $2(37) = 74$.

Time = 0.11 (sec) , antiderivative size = 155, normalized size of antiderivative = 4.19

method	result	size
risch	$x - \frac{\ln(e^{2bx+2a}-1)e^{2a}}{b(e^{2a}-e^{2c})} - \frac{\ln(e^{2bx+2a}-1)e^{2c}}{b(e^{2a}-e^{2c})} + \frac{\ln(e^{2bx+2a}-e^{2a-2c})e^{2a}}{b(e^{2a}-e^{2c})} + \frac{\ln(e^{2bx+2a}-e^{2a-2c})e^{2c}}{b(e^{2a}-e^{2c})}$	155

input `int(coth(b*x+a)*coth(b*x+c),x,method=_RETURNVERBOSE)`

output

```
x-1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2*a)-1)*exp(2*a)-1/b/(exp(2*a)-exp(
2*c))*ln(exp(2*b*x+2*a)-1)*exp(2*c)+1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2
*a)-exp(2*a-2*c))*exp(2*a)+1/b/(exp(2*a)-exp(2*c))*ln(exp(2*b*x+2*a)-exp(2
*a-2*c))*exp(2*c)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. $2(37) = 74$.

Time = 0.10 (sec) , antiderivative size = 259, normalized size of antiderivative = 7.00

$$\int \coth(a + bx) \coth(c + bx) dx$$

$$= \frac{bx \cosh(-a + c)^2 - 2bx \cosh(-a + c) \sinh(-a + c) + bx \sinh(-a + c)^2 - bx - (\cosh(-a + c)^2 - 2 \coth(-a + c) \sinh(-a + c) + \sinh(-a + c)^2)}{b^2 \cosh(-a + c)^2 - 2b \cosh(-a + c) \sinh(-a + c) + b \sinh(-a + c)^2 - b}$$

input

```
integrate(coth(b*x+a)*coth(b*x+c),x, algorithm="fricas")
```

output

```
(b*x*cosh(-a + c)^2 - 2*b*x*cosh(-a + c)*sinh(-a + c) + b*x*sinh(-a + c)^2
- b*x - (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) + sinh(-a + c)^2 +
1)*log(2*(cosh(-a + c)*sinh(b*x + c) - cosh(b*x + c)*sinh(-a + c))/(cosh(b
*x + c)*cosh(-a + c) - (cosh(-a + c) + sinh(-a + c))*sinh(b*x + c) + cosh(
b*x + c)*sinh(-a + c))) + (cosh(-a + c)^2 - 2*cosh(-a + c)*sinh(-a + c) +
sinh(-a + c)^2 + 1)*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))))/
(b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^2 - b)
```

Sympy [F]

$$\int \coth(a + bx) \coth(c + bx) dx = \int \coth(a + bx) \coth(bx + c) dx$$

input

```
integrate(coth(b*x+a)*coth(b*x+c),x)
```

output

```
Integral(coth(a + b*x)*coth(b*x + c), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(37) = 74$.

Time = 0.04 (sec) , antiderivative size = 157, normalized size of antiderivative = 4.24

$$\int \coth(a + bx) \coth(c + bx) dx = x + \frac{a}{b} - \frac{(e^{(2a)} + e^{(2c)}) \log(e^{-bx-a} + 1)}{b(e^{(2a)} - e^{(2c)})} - \frac{(e^{(2a)} + e^{(2c)}) \log(e^{-bx-a} - 1)}{b(e^{(2a)} - e^{(2c)})} + \frac{(e^{(2a)} + e^{(2c)}) \log(e^{-bx} + e^c)}{b(e^{(2a)} - e^{(2c)})} + \frac{(e^{(2a)} + e^{(2c)}) \log(e^{-bx} - e^c)}{b(e^{(2a)} - e^{(2c)})}$$

input `integrate(coth(b*x+a)*coth(b*x+c),x, algorithm="maxima")`

output `x + a/b - (e^(2*a) + e^(2*c))*log(e^(-b*x - a) + 1)/(b*(e^(2*a) - e^(2*c))) - (e^(2*a) + e^(2*c))*log(e^(-b*x - a) - 1)/(b*(e^(2*a) - e^(2*c))) + (e^(2*a) + e^(2*c))*log(e^(-b*x) + e^c)/(b*(e^(2*a) - e^(2*c))) + (e^(2*a) + e^(2*c))*log(e^(-b*x) - e^c)/(b*(e^(2*a) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. $2(37) = 74$.

Time = 0.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.62

$$\int \coth(a + bx) \coth(c + bx) dx = x - \frac{(e^{(4a)} + e^{(2a+2c)}) \log(|e^{(2bx+2a)} - 1|)}{be^{(4a)} - be^{(2a+2c)}} + \frac{(e^{(2a+2c)} + e^{(4c)}) \log(|e^{(2bx+2c)} - 1|)}{be^{(2a+2c)} - be^{(4c)}}$$

input `integrate(coth(b*x+a)*coth(b*x+c),x, algorithm="giac")`

output

$$x - (e^{(4*a)} + e^{(2*a + 2*c)}) * \log(\text{abs}(e^{(2*b*x + 2*a)} - 1)) / (b * e^{(4*a)} - b * e^{(2*a + 2*c)}) + (e^{(2*a + 2*c)} + e^{(4*c)}) * \log(\text{abs}(e^{(2*b*x + 2*c)} - 1)) / (b * e^{(2*a + 2*c)} - b * e^{(4*c)})$$

Mupad [B] (verification not implemented)

Time = 1.42 (sec) , antiderivative size = 115, normalized size of antiderivative = 3.11

$$\int \coth(a + bx) \coth(c + bx) dx$$

$$= x - \frac{\ln(4e^{4a} - 4e^{6a}e^{2bx} + 4e^{2a}e^{2c} - 4e^{4a}e^{2c}e^{2bx}) \coth(a - c)}{b}$$

$$+ \frac{\ln(4e^{4a} + 4e^{2a}e^{2c} - 4e^{2a}e^{4c}e^{2bx} - 4e^{4a}e^{2c}e^{2bx}) \coth(a - c)}{b}$$

input

```
int(coth(a + b*x)*coth(c + b*x),x)
```

output

$$x - (\log(4 * \exp(4 * a) - 4 * \exp(6 * a) * \exp(2 * b * x) + 4 * \exp(2 * a) * \exp(2 * c) - 4 * \exp(4 * a) * \exp(2 * c) * \exp(2 * b * x)) * \coth(a - c)) / b + (\log(4 * \exp(4 * a) + 4 * \exp(2 * a) * \exp(2 * c) - 4 * \exp(2 * a) * \exp(4 * c) * \exp(2 * b * x) - 4 * \exp(4 * a) * \exp(2 * c) * \exp(2 * b * x)) * \coth(a - c)) / b$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 169, normalized size of antiderivative = 4.57

$$\int \coth(a + bx) \coth(c + bx) dx$$

$$= \frac{e^{2a} \log(e^{bx+c} - 1) + e^{2a} \log(e^{bx+c} + 1) - e^{2a} \log(e^{bx+a} - 1) - e^{2a} \log(e^{bx+a} + 1) + e^{2a} bx + e^{2c} \log(e^{bx+c} - 1)}{b(e^{2a} - e^{2c})}$$

input

```
int(coth(b*x+a)*coth(b*x+c),x)
```

output

```
(e**(2*a)*log(e**(b*x + c) - 1) + e**(2*a)*log(e**(b*x + c) + 1) - e**(2*a)
)*log(e**(a + b*x) - 1) - e**(2*a)*log(e**(a + b*x) + 1) + e**(2*a)*b*x +
e**(2*c)*log(e**(b*x + c) - 1) + e**(2*c)*log(e**(b*x + c) + 1) - e**(2*c)
*log(e**(a + b*x) - 1) - e**(2*c)*log(e**(a + b*x) + 1) - e**(2*c)*b*x)/(b
*(e**(2*a) - e**(2*c)))
```

3.167 $\int \coth(c - bx) \coth(a + bx) dx$

Optimal result	1138
Mathematica [A] (verified)	1138
Rubi [C] (verified)	1139
Maple [B] (verified)	1140
Fricas [B] (verification not implemented)	1141
Sympy [F]	1141
Maxima [B] (verification not implemented)	1142
Giac [B] (verification not implemented)	1142
Mupad [B] (verification not implemented)	1143
Reduce [B] (verification not implemented)	1143

Optimal result

Integrand size = 14, antiderivative size = 36

$$\int \coth(c - bx) \coth(a + bx) dx = -x - \frac{\coth(a + c) \log(\sinh(c - bx))}{b} + \frac{\coth(a + c) \log(\sinh(a + bx))}{b}$$

output `-x-coth(a+c)*ln(-sinh(b*x-c))/b+coth(a+c)*ln(sinh(b*x+a))/b`

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

$$\int \coth(c - bx) \coth(a + bx) dx = -x + \frac{\coth(a + c)(-\log(\sinh(c - bx)) + \log(-\sinh(a + bx)))}{b}$$

input `Integrate[Coth[c - b*x]*Coth[a + b*x],x]`

output `-x + (Coth[a + c]*(-Log[Sinh[c - b*x]] + Log[-Sinh[a + b*x]]))/b`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.39, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {6181, 6179, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \coth(a + bx) \coth(c - bx) dx \\
 & \quad \downarrow \text{6181} \\
 & \cosh(a + c) \int \operatorname{csch}(c - bx) \operatorname{csch}(a + bx) dx - x \\
 & \quad \downarrow \text{6179} \\
 & \cosh(a + c) (\operatorname{csch}(a + c) \int \coth(c - bx) dx + \operatorname{csch}(a + c) \int \coth(a + bx) dx) - x \\
 & \quad \downarrow \text{3042} \\
 & c) \left(\operatorname{csch}(a + c) \int -i \tan \left(ic - ibx + \frac{\pi}{2} \right) dx + \operatorname{csch}(a + c) \int -i \tan \left(ia + ibx + \frac{\pi}{2} \right) dx \right) \\
 & \quad \downarrow \text{26} \\
 & c) \left(-i \operatorname{csch}(a + c) \int \tan \left(\frac{1}{2}(2ic + \pi) - ibx \right) dx - i \operatorname{csch}(a + c) \int \tan \left(\frac{1}{2}(2ia + \pi) + ibx \right) dx \right) \\
 & \quad \downarrow \text{3956} \\
 & -x + \cosh(a + c) \left(\frac{\operatorname{csch}(a + c) \log(-i \sinh(a + bx))}{b} - \frac{\operatorname{csch}(a + c) \log(-i \sinh(c - bx))}{b} \right)
 \end{aligned}$$

input `Int[Coth[c - b*x]*Coth[a + b*x],x]`

output `-x + Cosh[a + c]*(-((Csch[a + c]*Log[(-I)*Sinh[c - b*x]])/b) + (Csch[a + c]*Log[(-I)*Sinh[a + b*x]])/b)`

Definitions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 6179 `Int[Csch[(a_.) + (b_.)*(x_)]*Csch[(c_) + (d_.)*(x_)], x_Symbol] := Simp[Csch[(b*c - a*d)/b] Int[Coth[a + b*x], x], x] - Simp[Csch[(b*c - a*d)/d] Int[Coth[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`
- rule 6181 `Int[Coth[(a_.) + (b_.)*(x_)]*Coth[(c_) + (d_.)*(x_)], x_Symbol] := Simp[b*(x/d), x] + Simp[Cosh[(b*c - a*d)/d] Int[Csch[a + b*x]*Csch[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(39) = 78$.

Time = 0.11 (sec) , antiderivative size = 153, normalized size of antiderivative = 4.25

method	result	size
risch	$-x - \frac{\ln(-e^{2a+2c} + e^{2bx+2a})e^{2a+2c}}{b(e^{2a+2c}-1)} - \frac{\ln(-e^{2a+2c} + e^{2bx+2a})}{b(e^{2a+2c}-1)} + \frac{\ln(e^{2bx+2a}-1)e^{2a+2c}}{b(e^{2a+2c}-1)} + \frac{\ln(e^{2bx+2a}-1)}{b(e^{2a+2c}-1)}$	153

input `int(-coth(b*x-c)*coth(b*x+a),x,method=_RETURNVERBOSE)`

output

```
-x-1/b/(exp(2*a+2*c)-1)*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))*exp(2*a+2*c)-1/b/
(exp(2*a+2*c)-1)*ln(-exp(2*a+2*c)+exp(2*b*x+2*a))+1/b/(exp(2*a+2*c)-1)*ln(
exp(2*b*x+2*a)-1)*exp(2*a+2*c)+1/b/(exp(2*a+2*c)-1)*ln(exp(2*b*x+2*a)-1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(39) = 78.

Time = 0.12 (sec) , antiderivative size = 216, normalized size of antiderivative = 6.00

$$\int \coth(c - bx) \coth(a + bx) dx =$$

$$\frac{bx \cosh(a + c)^2 - 2bx \cosh(a + c) \sinh(a + c) + bx \sinh(a + c)^2 - bx - (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log(2 * (\cosh(a + c) \sinh(bx + a) - \cosh(bx + a) \sinh(a + c)) / (\cosh(bx + a) \cosh(a + c) - (\cosh(a + c) + \sinh(a + c)) \sinh(bx + a) + \cosh(bx + a) \sinh(a + c))) + (\cosh(a + c)^2 - 2 \cosh(a + c) \sinh(a + c) + \sinh(a + c)^2 + 1) \log(2 * \sinh(bx + a) / (\cosh(bx + a) - \sinh(bx + a)))}{(b \cosh(a + c)^2 - 2 * b \cosh(a + c) \sinh(a + c) + b \sinh(a + c)^2 - b)}$$

input

```
integrate(-coth(b*x-c)*coth(b*x+a),x, algorithm="fricas")
```

output

```
-(b*x*cosh(a + c)^2 - 2*b*x*cosh(a + c)*sinh(a + c) + b*x*sinh(a + c)^2 -
b*x - (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 + 1)*log(
2*(cosh(a + c)*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*c
osh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sin
h(a + c))) + (cosh(a + c)^2 - 2*cosh(a + c)*sinh(a + c) + sinh(a + c)^2 +
1)*log(2*sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/(b*cosh(a + c)^2
- 2*b*cosh(a + c)*sinh(a + c) + b*sinh(a + c)^2 - b)
```

Sympy [F]

$$\int \coth(c - bx) \coth(a + bx) dx = - \int \coth(a + bx) \coth(bx - c) dx$$

input

```
integrate(-coth(b*x-c)*coth(b*x+a),x)
```

output

```
-Integral(coth(a + b*x)*coth(b*x - c), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(39) = 78$.

Time = 0.05 (sec) , antiderivative size = 160, normalized size of antiderivative = 4.44

$$\int \coth(c - bx) \coth(a + bx) dx = -x - \frac{a}{b} + \frac{(e^{(2a+2c)} + 1) \log(e^{(-bx-a)} + 1)}{b(e^{(2a+2c)} - 1)} + \frac{(e^{(2a+2c)} + 1) \log(e^{(-bx-a)} - 1)}{b(e^{(2a+2c)} - 1)} - \frac{(e^{(2a+2c)} + 1) \log(e^{(-bx+c)} + 1)}{b(e^{(2a+2c)} - 1)} - \frac{(e^{(2a+2c)} + 1) \log(e^{(-bx+c)} - 1)}{b(e^{(2a+2c)} - 1)}$$

input `integrate(-coth(b*x-c)*coth(b*x+a),x, algorithm="maxima")`

output `-x - a/b + (e^(2*a + 2*c) + 1)*log(e^(-b*x - a) + 1)/(b*(e^(2*a + 2*c) - 1)) + (e^(2*a + 2*c) + 1)*log(e^(-b*x - a) - 1)/(b*(e^(2*a + 2*c) - 1)) - (e^(2*a + 2*c) + 1)*log(e^(-b*x + c) + 1)/(b*(e^(2*a + 2*c) - 1)) - (e^(2*a + 2*c) + 1)*log(e^(-b*x + c) - 1)/(b*(e^(2*a + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(39) = 78$.

Time = 0.19 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.61

$$\int \coth(c - bx) \coth(a + bx) dx = -x - \frac{(e^{(2a+2c)} + 1) \log(|e^{(2bx)} - e^{(2c)}|)}{be^{(2a+2c)} - b} - \frac{(e^{(2a)} + e^{(4a+2c)}) \log(|e^{(2bx+2a)} - 1|)}{be^{(2a)} - be^{(4a+2c)}}$$

input `integrate(-coth(b*x-c)*coth(b*x+a),x, algorithm="giac")`

output

$$-x - (e^{(2*a + 2*c)} + 1)*\log(\text{abs}(e^{(2*b*x)} - e^{(2*c)}))/(b*e^{(2*a + 2*c)} - b) - (e^{(2*a)} + e^{(4*a + 2*c)})*\log(\text{abs}(e^{(2*b*x + 2*a)} - 1))/(b*e^{(2*a)} - b*e^{(4*a + 2*c)})$$

Mupad [B] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 121, normalized size of antiderivative = 3.36

$$\int \coth(c - bx) \coth(a + bx) dx$$

$$= \frac{\coth(a + c) \ln(4e^{2a}e^{2c} + 4e^{4a}e^{4c} - 4e^{4a}e^{2c}e^{2bx} - 4e^{6a}e^{4c}e^{2bx})}{b} - \frac{\coth(a + c) \ln(4e^{2a}e^{2bx} - 4e^{2a}e^{2c} - 4e^{4a}e^{4c} + 4e^{4a}e^{2c}e^{2bx})}{b} - x$$

input

```
int(coth(a + b*x)*coth(c - b*x),x)
```

output

$$\frac{(\coth(a + c)*\log(4*\exp(2*a)*\exp(2*c) + 4*\exp(4*a)*\exp(4*c) - 4*\exp(4*a)*\exp(2*c)*\exp(2*b*x) - 4*\exp(6*a)*\exp(4*c)*\exp(2*b*x))}{b} - (\coth(a + c)*\log(4*\exp(2*a)*\exp(2*b*x) - 4*\exp(2*a)*\exp(2*c) - 4*\exp(4*a)*\exp(4*c) + 4*\exp(4*a)*\exp(2*c)*\exp(2*b*x))}{b} - x$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 164, normalized size of antiderivative = 4.56

$$\int \coth(c - bx) \coth(a + bx) dx$$

$$= \frac{e^{2a+2c}\log(e^{bx+a} - 1) + e^{2a+2c}\log(e^{bx+a} + 1) - e^{2a+2c}\log(e^{bx} + e^c) - e^{2a+2c}\log(e^{bx} - e^c) - e^{2a+2c}bx + \log(b)}{b(e^{2a+2c} - 1)}$$

input

```
int(-coth(b*x-c)*coth(b*x+a),x)
```


output

```
(e**(2*a + 2*c)*log(e**(a + b*x) - 1) + e**(2*a + 2*c)*log(e**(a + b*x) +
1) - e**(2*a + 2*c)*log(e**(b*x) + e**c) - e**(2*a + 2*c)*log(e**(b*x) - e
**c) - e**(2*a + 2*c)*b*x + log(e**(a + b*x) - 1) + log(e**(a + b*x) + 1)
- log(e**(b*x) + e**c) - log(e**(b*x) - e**c) + b*x)/(b*(e**(2*a + 2*c) -
1))
```

3.168 $\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx$

Optimal result	1145
Mathematica [A] (verified)	1145
Rubi [A] (verified)	1146
Maple [B] (verified)	1147
Fricas [B] (verification not implemented)	1148
Sympy [F]	1148
Maxima [A] (verification not implemented)	1149
Giac [B] (verification not implemented)	1149
Mupad [B] (verification not implemented)	1150
Reduce [B] (verification not implemented)	1150

Optimal result

Integrand size = 13, antiderivative size = 36

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx = \frac{\operatorname{csch}(a - c) \log(\cosh(a + bx))}{b} - \frac{\operatorname{csch}(a - c) \log(\cosh(c + bx))}{b}$$

output

```
csch(a-c)*ln(cosh(b*x+a))/b-csch(a-c)*ln(cosh(b*x+c))/b
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx = \frac{\operatorname{csch}(a - c)(\log(\cosh(a + bx)) - \log(\cosh(c + bx)))}{b}$$

input

```
Integrate[Sech[a + b*x]*Sech[c + b*x],x]
```

output

```
(Csch[a - c]*(Log[Cosh[a + b*x]] - Log[Cosh[c + b*x]]))/b
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {6178, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(bx + c) dx$$

$$\downarrow 6178$$

$$\operatorname{csch}(a - c) \int \tanh(a + bx) dx - \operatorname{csch}(a - c) \int \tanh(c + bx) dx$$

$$\downarrow 3042$$

$$\operatorname{csch}(a - c) \int -i \tan(ia + ibx) dx - \operatorname{csch}(a - c) \int -i \tan(ic + ibx) dx$$

$$\downarrow 26$$

$$i \operatorname{csch}(a - c) \int \tan(ic + ibx) dx - i \operatorname{csch}(a - c) \int \tan(ia + ibx) dx$$

$$\downarrow 3956$$

$$\frac{\operatorname{csch}(a - c) \log(\cosh(a + bx))}{b} - \frac{\operatorname{csch}(a - c) \log(\cosh(bx + c))}{b}$$

input `Int[Sech[a + b*x]*Sech[c + b*x],x]`

output `(Csch[a - c]*Log[Cosh[a + b*x]])/b - (Csch[a - c]*Log[Cosh[c + b*x]])/b`

Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6178 `Int[Sech[(a_.) + (b_.)*(x_)]*Sech[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Csch[(b*c - a*d)/d] Int[Tanh[a + b*x], x], x] + Simp[Csch[(b*c - a*d)/b] Int[Tanh[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(36) = 72$.

Time = 0.37 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.14

method	result	size
risch	$\frac{2 \ln(e^{2bx+2a}+1)e^{a+c}}{(e^{2a}-e^{2c})b} - \frac{2 \ln(e^{2bx+2a}+e^{2a-2c})e^{a+c}}{(e^{2a}-e^{2c})b}$	77

input `int(sech(b*x+a)*sech(b*x+c),x,method=_RETURNVERBOSE)`

output `2*ln(exp(2*b*x+2*a)+1)/(exp(2*a)-exp(2*c))/b*exp(a+c)-2*ln(exp(2*b*x+2*a)+exp(2*a-2*c))/(exp(2*a)-exp(2*c))/b*exp(a+c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(36) = 72$.

Time = 0.08 (sec) , antiderivative size = 184, normalized size of antiderivative = 5.11

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx$$

$$= \frac{2 \left((\cosh(-a + c) - \sinh(-a + c)) \log \left(\frac{2 (\cosh(bx+c) \cosh(-a+c) - \sinh(bx+c) \sinh(-a+c))}{\cosh(bx+c) \cosh(-a+c) - (\cosh(-a+c) + \sinh(-a+c)) \sinh(bx+c) + \cosh(bx+c) \sinh(-a+c)} \right) \right)}{b \cosh(-a + c)^2 - 2b \cosh(-a + c) \sinh(-a + c) + b \sinh(-a + c)^2}$$

input `integrate(sech(b*x+a)*sech(b*x+c),x, algorithm="fricas")`

output `2*((cosh(-a + c) - sinh(-a + c))*log(2*(cosh(b*x + c)*cosh(-a + c) - sinh(b*x + c)*sinh(-a + c))/(cosh(b*x + c)*cosh(-a + c) - (cosh(-a + c) + sinh(-a + c))*sinh(b*x + c) + cosh(b*x + c)*sinh(-a + c))) - (cosh(-a + c) - sinh(-a + c))*log(2*cosh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))))/(b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^2 - b)`

Sympy [F]

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx = \int \operatorname{sech}(a + bx) \operatorname{sech}(bx + c) dx$$

input `integrate(sech(b*x+a)*sech(b*x+c),x)`

output `Integral(sech(a + b*x)*sech(b*x + c), x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.89

$$\int \operatorname{sech}(a+bx)\operatorname{sech}(c+bx) dx = \frac{2e^{(a+c)} \log(e^{-2bx-2a} + 1)}{b(e^{2a} - e^{2c})} - \frac{2e^{(a+c)} \log(e^{-2bx} + e^{2c})}{b(e^{2a} - e^{2c})}$$

input `integrate(sech(b*x+a)*sech(b*x+c),x, algorithm="maxima")`

output `2*e^(a + c)*log(e^(-2*b*x - 2*a) + 1)/(b*(e^(2*a) - e^(2*c))) - 2*e^(a + c)*log(e^(-2*b*x) + e^(2*c))/(b*(e^(2*a) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(36) = 72.

Time = 0.17 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.28

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx = 2 \left(\frac{e^{(2a)} \log(e^{(2bx+2a)} + 1)}{be^{(4a)} - be^{(2a+2c)}} - \frac{e^{(2c)} \log(e^{(2bx+2c)} + 1)}{be^{(2a+2c)} - be^{(4c)}} \right) e^{(a+c)}$$

input `integrate(sech(b*x+a)*sech(b*x+c),x, algorithm="giac")`

output `2*(e^(2*a)*log(e^(2*b*x + 2*a) + 1)/(b*e^(4*a) - b*e^(2*a + 2*c)) - e^(2*c)*log(e^(2*b*x + 2*c) + 1)/(b*e^(2*a + 2*c) - b*e^(4*c)))*e^(a + c)`

Mupad [B] (verification not implemented)

Time = 2.00 (sec) , antiderivative size = 266, normalized size of antiderivative = 7.39

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx$$

$$= \frac{4\sqrt{e^{2a-2c}} \operatorname{atan}\left(\frac{b(e^{-a}e^c + e^{-3a}e^{3c})(e^{2a}e^{-2c})^{3/2}}{\sqrt{-b^2(e^{2a}e^{-2c}-1)^2}} + \frac{e^{2a}e^{2bx}\left(\frac{2e^{-c}e^a}{b(e^{2a}e^{-2c})^{3/2}} + \frac{2(e^{-a}e^c + e^{-3a}e^{3c})(b\sqrt{e^{2a}e^{-2c}+b}(e^{2a}e^{-2c})^{3/2})}{\sqrt{-b^2(e^{2a}e^{-2c}-1)^2}\sqrt{2b^2e^{2a}e^{-2c}-b^2-b^2e^{4a}e^{4c}}}\right)}{4}\right)}{\sqrt{2b^2e^{2a-2c}-b^2e^{4a-4c}-b^2}}$$

input `int(1/(cosh(a + b*x)*cosh(c + b*x)),x)`output `(4*exp(2*a - 2*c)^(1/2)*atan((b*(exp(-a)*exp(c) + exp(-3*a)*exp(3*c))*(exp(2*a)*exp(-2*c))^(3/2))/(-b^2*(exp(2*a)*exp(-2*c) - 1)^2)^(1/2) + (exp(2*a)*exp(2*b*x)*((2*exp(-c)*exp(a))/(b*(exp(2*a)*exp(-2*c))^(3/2)) + (2*(exp(-a)*exp(c) + exp(-3*a)*exp(3*c))*(b*(exp(2*a)*exp(-2*c))^(1/2) + b*(exp(2*a)*exp(-2*c))^(3/2)))/((-b^2*(exp(2*a)*exp(-2*c) - 1)^2)^(1/2)*(2*b^2*exp(2*a)*exp(-2*c) - b^2 - b^2*exp(4*a)*exp(-4*c))^(1/2)))/(2*b^2*exp(2*a)*exp(-2*c) - b^2 - b^2*exp(4*a)*exp(-4*c))^(1/2))/4)/(2*b^2*exp(2*a - 2*c) - b^2*exp(4*a - 4*c) - b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.50

$$\int \operatorname{sech}(a + bx)\operatorname{sech}(c + bx) dx = \frac{2e^{a+c}(\log(e^{2bx+2a} + 1) - \log(e^{2bx+2c} + 1))}{b(e^{2a} - e^{2c})}$$

input `int(sech(b*x+a)*sech(b*x+c),x)`output `(2*e**(a + c)*(log(e**(2*a + 2*b*x) + 1) - log(e**(2*b*x + 2*c) + 1)))/(b*(e**(2*a) - e**(2*c)))`

3.169 $\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx$

Optimal result	1151
Mathematica [A] (verified)	1151
Rubi [A] (verified)	1152
Maple [B] (verified)	1153
Fricas [B] (verification not implemented)	1154
Sympy [F]	1154
Maxima [A] (verification not implemented)	1155
Giac [B] (verification not implemented)	1155
Mupad [B] (verification not implemented)	1156
Reduce [B] (verification not implemented)	1156

Optimal result

Integrand size = 14, antiderivative size = 33

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx = -\frac{\operatorname{csch}(a + c) \log(\cosh(c - bx))}{b} + \frac{\operatorname{csch}(a + c) \log(\cosh(a + bx))}{b}$$

output

```
-csch(a+c)*ln(cosh(b*x-c))/b+csch(a+c)*ln(cosh(b*x+a))/b
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx = -\frac{\operatorname{csch}(a + c)(\log(\cosh(c - bx)) - \log(\cosh(a + bx)))}{b}$$

input

```
Integrate[Sech[c - b*x]*Sech[a + b*x],x]
```

output

```
-((Csch[a + c]*(Log[Cosh[c - b*x]] - Log[Cosh[a + b*x]]))/b)
```


Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6178, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{sech}(a + bx)\operatorname{sech}(c - bx) dx \\
 & \quad \downarrow \text{6178} \\
 & \operatorname{csch}(a + c) \int \tanh(c - bx)dx + \operatorname{csch}(a + c) \int \tanh(a + bx)dx \\
 & \quad \downarrow \text{3042} \\
 & \operatorname{csch}(a + c) \int -i \tan(ic - ibx)dx + \operatorname{csch}(a + c) \int -i \tan(ia + ibx)dx \\
 & \quad \downarrow \text{26} \\
 & -i\operatorname{csch}(a + c) \int \tan(ic - ibx)dx - i\operatorname{csch}(a + c) \int \tan(ia + ibx)dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\operatorname{csch}(a + c) \log(\cosh(a + bx))}{b} - \frac{\operatorname{csch}(a + c) \log(\cosh(c - bx))}{b}
 \end{aligned}$$

input `Int[Sech[c - b*x]*Sech[a + b*x],x]`

output `-((Csch[a + c]*Log[Cosh[c - b*x]])/b) + (Csch[a + c]*Log[Cosh[a + b*x]])/b`

Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6178 `Int[Sech[(a_.) + (b_.)*(x_)]*Sech[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Csch[(b*c - a*d)/d] Int[Tanh[a + b*x], x], x] + Simp[Csch[(b*c - a*d)/b] Int[Tanh[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(34) = 68$.

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.27

method	result	size
risch	$-\frac{2 \ln(e^{2a+2c} + e^{2bx+2a})e^{a+c}}{b(e^{2a+2c}-1)} + \frac{2 \ln(e^{2bx+2a} + 1)e^{a+c}}{b(e^{2a+2c}-1)}$	75

input `int(sech(b*x-c)*sech(b*x+a), x, method=_RETURNVERBOSE)`

output `-2/b/(exp(2*a+2*c)-1)*ln(exp(2*a+2*c)+exp(2*b*x+2*a))*exp(a+c)+2/b/(exp(2*a+2*c)-1)*ln(exp(2*b*x+2*a)+1)*exp(a+c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(34) = 68$.

Time = 0.08 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.73

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx$$

$$= \frac{2 \left((\cosh(a + c) - \sinh(a + c)) \log \left(\frac{2 (\cosh(bx+a) \cosh(a+c) - \sinh(bx+a) \sinh(a+c))}{\cosh(bx+a) \cosh(a+c) - (\cosh(a+c) + \sinh(a+c)) \sinh(bx+a) + \cosh(bx+a) \sinh(a+c)} \right) - (\cosh(a + c) - \sinh(a + c)) \log \left(\frac{2 \cosh(bx+a)}{\cosh(bx+a) - \sinh(bx+a)} \right) \right)}{b \cosh(a + c)^2 - 2 b \cosh(a + c) \sinh(a + c) + b \sinh(a + c)^2}$$

input `integrate(sech(b*x-c)*sech(b*x+a),x, algorithm="fricas")`

output `2*((cosh(a + c) - sinh(a + c))*log(2*(cosh(b*x + a)*cosh(a + c) - sinh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c))) - (cosh(a + c) - sinh(a + c))*log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/(b*cosh(a + c)^2 - 2*b*cosh(a + c)*sinh(a + c) + b*sinh(a + c)^2 - b)`

Sympy [F]

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx = \int \operatorname{sech}(a + bx) \operatorname{sech}(bx - c) dx$$

input `integrate(sech(b*x-c)*sech(b*x+a),x)`

output `Integral(sech(a + b*x)*sech(b*x - c), x)`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.03

$$\int \operatorname{sech}(c-bx)\operatorname{sech}(a+bx) dx = \frac{2e^{(a+c)} \log(e^{(-2bx-2a)} + 1)}{b(e^{(2a+2c)} - 1)} - \frac{2e^{(a+c)} \log(e^{(-2bx+2c)} + 1)}{b(e^{(2a+2c)} - 1)}$$

input `integrate(sech(b*x-c)*sech(b*x+a),x, algorithm="maxima")`

output `2*e^(a + c)*log(e^(-2*b*x - 2*a) + 1)/(b*(e^(2*a + 2*c) - 1)) - 2*e^(a + c)*log(e^(-2*b*x + 2*c) + 1)/(b*(e^(2*a + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(34) = 68.

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.18

$$\int \operatorname{sech}(c-bx)\operatorname{sech}(a+bx) dx = -2 \left(\frac{e^{(2a)} \log(e^{(2bx+2a)} + 1)}{be^{(2a)} - be^{(4a+2c)}} + \frac{\log(e^{(2bx)} + e^{(2c)})}{be^{(2a+2c)} - b} \right) e^{(a+c)}$$

input `integrate(sech(b*x-c)*sech(b*x+a),x, algorithm="giac")`

output `-2*(e^(2*a)*log(e^(2*b*x + 2*a) + 1)/(b*e^(2*a) - b*e^(4*a + 2*c)) + log(e^(2*b*x) + e^(2*c))/(b*e^(2*a + 2*c) - b))*e^(a + c)`

Mupad [B] (verification not implemented)

Time = 1.90 (sec) , antiderivative size = 268, normalized size of antiderivative = 8.12

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx$$

$$= \frac{4 \operatorname{atan} \left(\frac{e^{2a} e^{2bx} \left(\frac{2e^a e^c}{b(e^{2a} e^{2c})^{3/2}} + \frac{2e^{-3a} e^{-3c} (e^{2a} e^{2c} + 1) (b\sqrt{e^{2a} e^{2c} + b(e^{2a} e^{2c})^{3/2}})}{\sqrt{-b^2 (e^{2a} e^{2c} - 1)^2} \sqrt{2b^2 e^{2a} e^{2c} - b^2 - b^2 e^{4a} e^{4c}}} \right)}{\sqrt{2b^2 e^{2a} e^{2c} - b^2 - b^2 e^{4a} e^{4c}}} \right) + \frac{be^{-3a} e^{-3c}}{\sqrt{-b^2 (e^{2a} e^{2c} - 1)^2}}}{\sqrt{2b^2 e^{2a+2c} - b^2 e^{4a+4c} - b^2}}$$

input `int(1/(cosh(a + b*x)*cosh(c - b*x)),x)`output `(4*atan((exp(2*a)*exp(2*b*x)*((2*exp(a)*exp(c))/(b*(exp(2*a)*exp(2*c))^(3/2)) + (2*exp(-3*a)*exp(-3*c)*(exp(2*a)*exp(2*c) + 1)*(b*(exp(2*a)*exp(2*c))^(1/2) + b*(exp(2*a)*exp(2*c))^(3/2)))/((-b^2*(exp(2*a)*exp(2*c) - 1)^2)^(1/2)*(2*b^2*exp(2*a)*exp(2*c) - b^2 - b^2*exp(4*a)*exp(4*c))^(1/2)))*(2*b^2*exp(2*a)*exp(2*c) - b^2 - b^2*exp(4*a)*exp(4*c))^(1/2))/4 + (b*exp(-3*a)*exp(-3*c)*(exp(2*a)*exp(2*c) + 1)*(exp(2*a)*exp(2*c))^(3/2))/(-b^2*(exp(2*a)*exp(2*c) - 1)^2)^(1/2))*exp(2*a + 2*c)^(1/2))/(2*b^2*exp(2*a + 2*c) - b^2*exp(4*a + 4*c) - b^2)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.58

$$\int \operatorname{sech}(c - bx)\operatorname{sech}(a + bx) dx = \frac{2e^{a+c}(\log(e^{2bx+2a} + 1) - \log(e^{2bx} + e^{2c}))}{b(e^{2a+2c} - 1)}$$

input `int(sech(b*x-c)*sech(b*x+a),x)`output `(2*e**(a + c)*(log(e**(2*a + 2*b*x) + 1) - log(e**(2*b*x) + e**(2*c))))/(b*(e**(2*a + 2*c) - 1))`

3.170 $\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx$

Optimal result	1157
Mathematica [A] (verified)	1157
Rubi [C] (verified)	1158
Maple [B] (verified)	1159
Fricas [B] (verification not implemented)	1160
Sympy [F]	1160
Maxima [B] (verification not implemented)	1161
Giac [B] (verification not implemented)	1161
Mupad [B] (verification not implemented)	1162
Reduce [B] (verification not implemented)	1162

Optimal result

Integrand size = 13, antiderivative size = 36

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx = -\frac{\operatorname{csch}(a - c)\log(\sinh(a + bx))}{b} + \frac{\operatorname{csch}(a - c)\log(\sinh(c + bx))}{b}$$

output

```
-csch(a-c)*ln(sinh(b*x+a))/b+csch(a-c)*ln(sinh(b*x+c))/b
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx = -\frac{\operatorname{csch}(a - c)(\log(\sinh(a + bx)) - \log(\sinh(c + bx)))}{b}$$

input

```
Integrate[Csch[a + b*x]*Csch[c + b*x],x]
```

output

```
-((Csch[a - c]*(Log[Sinh[a + b*x]] - Log[Sinh[c + b*x]]))/b)
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.22, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {6179, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{csch}(a + bx)\operatorname{csch}(bx + c) dx \\
 & \quad \downarrow \text{6179} \\
 & \operatorname{csch}(a - c) \int \operatorname{coth}(c + bx) dx - \operatorname{csch}(a - c) \int \operatorname{coth}(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \operatorname{csch}(a - c) \int -i \tan\left(ic + ibx + \frac{\pi}{2}\right) dx - \operatorname{csch}(a - c) \int -i \tan\left(ia + ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & i\operatorname{csch}(a - c) \int \tan\left(\frac{1}{2}(2ia + \pi) + ibx\right) dx - i\operatorname{csch}(a - c) \int \tan\left(\frac{1}{2}(2ic + \pi) + ibx\right) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\operatorname{csch}(a - c) \log(-i \sinh(bx + c))}{b} - \frac{\operatorname{csch}(a - c) \log(-i \sinh(a + bx))}{b}
 \end{aligned}$$

input

```
Int[Csch[a + b*x]*Csch[c + b*x],x]
```

output

```
-((Csch[a - c]*Log[(-I)*Sinh[a + b*x]])/b) + (Csch[a - c]*Log[(-I)*Sinh[c + b*x]])/b
```

Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6179 `Int[Csch[(a_.) + (b_.)*(x_)]*Csch[(c_) + (d_.)*(x_)], x_Symbol] := Simp[Csch[(b*c - a*d)/b] Int[Coth[a + b*x], x], x] - Simp[Csch[(b*c - a*d)/d] Int[Coth[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.19

method	result	size
risch	$\frac{2 \ln(e^{2bx+2a} - e^{2a-2c})e^{a+c}}{(e^{2a} - e^{2c})b} - \frac{2 \ln(e^{2bx+2a} - 1)e^{a+c}}{(e^{2a} - e^{2c})b}$	79

input `int(csch(b*x+a)*csch(b*x+c),x,method=_RETURNVERBOSE)`

output `2*ln(exp(2*b*x+2*a)-exp(2*a-2*c))/(exp(2*a)-exp(2*c))/b*exp(a+c)-2*ln(exp(2*b*x+2*a)-1)/(exp(2*a)-exp(2*c))/b*exp(a+c)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 184, normalized size of antiderivative = 5.11

$$\int \operatorname{csch}(a + bx) \operatorname{csch}(c + bx) dx = \frac{2 \left((\cosh(-a + c) - \sinh(-a + c)) \log \left(\frac{2 (\cosh(-a + c) \sinh(bx + c) - \cosh(bx + c) \sinh(-a + c))}{\cosh(bx + c) \cosh(-a + c) - (\cosh(-a + c) + \sinh(-a + c)) \sinh(bx + c) + \cosh(bx + c) \sinh(-a + c)} \right) \right)}{b \cosh(-a + c)^2 - 2b \cosh(-a + c) \sinh(-a + c) + b^2 \sinh(-a + c)^2 - b}$$

input `integrate(csch(b*x+a)*csch(b*x+c),x, algorithm="fricas")`

output `-2*((cosh(-a + c) - sinh(-a + c))*log(2*(cosh(-a + c)*sinh(b*x + c) - cosh(b*x + c)*sinh(-a + c))/(cosh(b*x + c)*cosh(-a + c) - (cosh(-a + c) + sinh(-a + c))*sinh(b*x + c) + cosh(b*x + c)*sinh(-a + c))) - (cosh(-a + c) - sinh(-a + c))*log(2*sinh(b*x + c)/(cosh(b*x + c) - sinh(b*x + c))))/(b*cosh(-a + c)^2 - 2*b*cosh(-a + c)*sinh(-a + c) + b*sinh(-a + c)^2 - b)`

Sympy [F]

$$\int \operatorname{csch}(a + bx) \operatorname{csch}(c + bx) dx = \int \operatorname{csch}(a + bx) \operatorname{csch}(bx + c) dx$$

input `integrate(csch(b*x+a)*csch(b*x+c),x)`

output `Integral(csch(a + b*x)*csch(b*x + c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. $2(36) = 72$.

Time = 0.04 (sec) , antiderivative size = 133, normalized size of antiderivative = 3.69

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx = -\frac{2e^{(a+c)} \log(e^{-bx-a} + 1)}{b(e^{2a} - e^{2c})} - \frac{2e^{(a+c)} \log(e^{-bx-a} - 1)}{b(e^{2a} - e^{2c})} \\ + \frac{2e^{(a+c)} \log(e^{-bx} + e^c)}{b(e^{2a} - e^{2c})} + \frac{2e^{(a+c)} \log(e^{-bx} - e^c)}{b(e^{2a} - e^{2c})}$$

input `integrate(csch(b*x+a)*csch(b*x+c),x, algorithm="maxima")`

output `-2*e^(a + c)*log(e^(-b*x - a) + 1)/(b*(e^(2*a) - e^(2*c))) - 2*e^(a + c)*log(e^(-b*x - a) - 1)/(b*(e^(2*a) - e^(2*c))) + 2*e^(a + c)*log(e^(-b*x) + e^c)/(b*(e^(2*a) - e^(2*c))) + 2*e^(a + c)*log(e^(-b*x) - e^c)/(b*(e^(2*a) - e^(2*c)))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. $2(36) = 72$.

Time = 0.12 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.33

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx \\ = -2 \left(\frac{e^{(2a)} \log(|e^{(2bx+2a)} - 1|)}{be^{(4a)} - be^{(2a+2c)}} - \frac{e^{(2c)} \log(|e^{(2bx+2c)} - 1|)}{be^{(2a+2c)} - be^{(4c)}} \right) e^{(a+c)}$$

input `integrate(csch(b*x+a)*csch(b*x+c),x, algorithm="giac")`

output `-2*(e^(2*a)*log(abs(e^(2*b*x + 2*a) - 1)))/(b*e^(4*a) - b*e^(2*a + 2*c)) - e^(2*c)*log(abs(e^(2*b*x + 2*c) - 1))/(b*e^(2*a + 2*c) - b*e^(4*c))*e^(a + c)`

Mupad [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 266, normalized size of antiderivative = 7.39

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx =$$

$$4\sqrt{e^{2a-2c}} \operatorname{atan}\left(\frac{b(e^{-a}e^c + e^{-3a}e^{3c})(e^{2a}e^{-2c})^{3/2}}{\sqrt{-b^2(e^{2a}e^{-2c}-1)^2}} - \frac{e^{2a}e^{2bx}\left(\frac{2e^{-c}e^a}{b(e^{2a}e^{-2c})^{3/2}} + \frac{2(e^{-a}e^c + e^{-3a}e^{3c})(b\sqrt{e^{2a}e^{-2c}+b}(e^{2a}e^{-2c})^{3/2})}{\sqrt{-b^2(e^{2a}e^{-2c}-1)^2}\sqrt{2b^2e^{2a}e^{-2c}-b^2-b^2e^{4a-4c}}}\right)}{4}\right)}{\sqrt{2b^2e^{2a-2c}-b^2e^{4a-4c}-b^2}}$$

input `int(1/(sinh(a + b*x)*sinh(c + b*x)),x)`output
$$\frac{-(4\exp(2a-2c)^{(1/2)}\operatorname{atan}((b(\exp(-a)\exp(c) + \exp(-3a)\exp(3c))\exp(2a)\exp(-2c))^{(3/2)})/(-b^2(\exp(2a)\exp(-2c)-1)^{(1/2)} - (\exp(2a)\exp(2bx)\exp(-2c)\exp(a)/b(\exp(2a)\exp(-2c))^{(3/2)} + (2(\exp(-a)\exp(c) + \exp(-3a)\exp(3c))\exp(2a)\exp(-2c))^{(1/2)} + b(\exp(2a)\exp(-2c))^{(3/2)}))/((-b^2(\exp(2a)\exp(-2c)-1)^{(1/2)}(2b^2\exp(2a)\exp(-2c) - b^2 - b^2\exp(4a)\exp(-4c))^{(1/2)}))\exp(2a)\exp(-2c) - b^2 - b^2\exp(4a)\exp(-4c))^{(1/2)}/4}}{(2b^2\exp(2a-2c) - b^2\exp(4a-4c) - b^2)^{(1/2)}}$$
Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.94

$$\int \operatorname{csch}(a + bx)\operatorname{csch}(c + bx) dx = \frac{2e^{a+c}(\log(e^{bx+c}-1) + \log(e^{bx+c}+1) - \log(e^{bx+a}-1) - \log(e^{bx+a}+1))}{b(e^{2a}-e^{2c})}$$

input `int(csch(b*x+a)*csch(b*x+c),x)`output
$$(2e^{a+c}(\log(e^{bx+c}-1) + \log(e^{bx+c}+1) - \log(e^{bx+a}-1) - \log(e^{bx+a}+1)))/(b(e^{2a}-e^{2c})))$$

3.171 $\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx$

Optimal result	1163
Mathematica [A] (verified)	1163
Rubi [C] (verified)	1164
Maple [B] (verified)	1165
Fricas [B] (verification not implemented)	1166
Sympy [F]	1166
Maxima [B] (verification not implemented)	1167
Giac [B] (verification not implemented)	1167
Mupad [B] (verification not implemented)	1168
Reduce [B] (verification not implemented)	1168

Optimal result

Integrand size = 14, antiderivative size = 33

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx = -\frac{\operatorname{csch}(a + c) \log(\sinh(c - bx))}{b} + \frac{\operatorname{csch}(a + c) \log(\sinh(a + bx))}{b}$$

output `-csch(a+c)*ln(-sinh(b*x-c))/b+csch(a+c)*ln(sinh(b*x+a))/b`

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx = -\frac{\operatorname{csch}(a + c)(\log(\sinh(c - bx)) - \log(-\sinh(a + bx)))}{b}$$

input `Integrate[Csch[c - b*x]*Csch[a + b*x],x]`

output `-((Csch[a + c]*(Log[Sinh[c - b*x]] - Log[-Sinh[a + b*x]]))/b)`

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {6179, 3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{csch}(a+bx)\operatorname{csch}(c-bx) dx \\
 & \quad \downarrow \text{6179} \\
 & \operatorname{csch}(a+c) \int \operatorname{coth}(c-bx) dx + \operatorname{csch}(a+c) \int \operatorname{coth}(a+bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \operatorname{csch}(a+c) \int -i \tan\left(ic-ibx+\frac{\pi}{2}\right) dx + \operatorname{csch}(a+c) \int -i \tan\left(ia+ibx+\frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{26} \\
 & -i\operatorname{csch}(a+c) \int \tan\left(\frac{1}{2}(2ic+\pi)-ibx\right) dx - i\operatorname{csch}(a+c) \int \tan\left(\frac{1}{2}(2ia+\pi)+ibx\right) dx \\
 & \quad \downarrow \text{3956} \\
 & \frac{\operatorname{csch}(a+c) \log(-i \sinh(a+bx))}{b} - \frac{\operatorname{csch}(a+c) \log(-i \sinh(c-bx))}{b}
 \end{aligned}$$

input

```
Int[Csch[c - b*x]*Csch[a + b*x],x]
```

output

```
-((Csch[a + c]*Log[(-I)*Sinh[c - b*x]])/b) + (Csch[a + c]*Log[(-I)*Sinh[a + b*x]])/b
```

Definitions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 6179 `Int[Csch[(a_.) + (b_.)*(x_)]*Csch[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Csch[(b*c - a*d)/b] Int[Coth[a + b*x], x], x] - Simp[Csch[(b*c - a*d)/d] Int[Coth[c + d*x], x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b^2 - d^2, 0] && NeQ[b*c - a*d, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(36) = 72$.

Time = 0.07 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.33

method	result	size
risch	$\frac{2 \ln(e^{2bx+2a}-1)e^{a+c}}{b(e^{2a+2c}-1)} - \frac{2 \ln(-e^{2a+2c}+e^{2bx+2a})e^{a+c}}{b(e^{2a+2c}-1)}$	77

input `int(-csch(b*x-c)*csch(b*x+a),x,method=_RETURNVERBOSE)`

output $\frac{2}{b} / (\exp(2*a+2*c)-1) * \ln(\exp(2*b*x+2*a)-1) * \exp(a+c) - \frac{2}{b} / (\exp(2*a+2*c)-1) * \ln(-\exp(2*a+2*c)+\exp(2*b*x+2*a)) * \exp(a+c)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 156, normalized size of antiderivative = 4.73

$$\int \operatorname{csch}(c - bx) \operatorname{csch}(a + bx) dx$$

$$= \frac{2 \left((\cosh(a + c) - \sinh(a + c)) \log \left(\frac{2 (\cosh(a+c) \sinh(bx+a) - \cosh(bx+a) \sinh(a+c))}{\cosh(bx+a) \cosh(a+c) - (\cosh(a+c) + \sinh(a+c)) \sinh(bx+a) + \cosh(bx+a) \sinh(a+c)} \right) - (\cosh(a + c) - \sinh(a + c)) \log(2 \sinh(bx + a) / (\cosh(bx + a) - \sinh(bx + a))) \right)}{b \cosh(a + c)^2 - 2 b \cosh(a + c) \sinh(a + c) + b \sinh(a + c)^2}$$

input `integrate(-csch(b*x-c)*csch(b*x+a),x, algorithm="fricas")`

output `2*((cosh(a + c) - sinh(a + c))*log(2*(cosh(a + c)*sinh(b*x + a) - cosh(b*x + a)*sinh(a + c))/(cosh(b*x + a)*cosh(a + c) - (cosh(a + c) + sinh(a + c))*sinh(b*x + a) + cosh(b*x + a)*sinh(a + c))) - (cosh(a + c) - sinh(a + c))*log(2*sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/(b*cosh(a + c)^2 - 2*b*cosh(a + c)*sinh(a + c) + b*sinh(a + c)^2 - b)`

Sympy [F]

$$\int \operatorname{csch}(c - bx) \operatorname{csch}(a + bx) dx = - \int \operatorname{csch}(a + bx) \operatorname{csch}(bx - c) dx$$

input `integrate(-csch(b*x-c)*csch(b*x+a),x)`

output `-Integral(csch(a + b*x)*csch(b*x - c), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 129 vs. $2(36) = 72$.

Time = 0.05 (sec) , antiderivative size = 129, normalized size of antiderivative = 3.91

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx = \frac{2e^{(a+c)} \log(e^{-bx-a} + 1)}{b(e^{2a+2c} - 1)} + \frac{2e^{(a+c)} \log(e^{-bx-a} - 1)}{b(e^{2a+2c} - 1)} - \frac{2e^{(a+c)} \log(e^{-bx+c} + 1)}{b(e^{2a+2c} - 1)} - \frac{2e^{(a+c)} \log(e^{-bx+c} - 1)}{b(e^{2a+2c} - 1)}$$

input `integrate(-csch(b*x-c)*csch(b*x+a),x, algorithm="maxima")`

output `2*e^(a + c)*log(e^(-b*x - a) + 1)/(b*(e^(2*a + 2*c) - 1)) + 2*e^(a + c)*log(e^(-b*x - a) - 1)/(b*(e^(2*a + 2*c) - 1)) - 2*e^(a + c)*log(e^(-b*x + c) + 1)/(b*(e^(2*a + 2*c) - 1)) - 2*e^(a + c)*log(e^(-b*x + c) - 1)/(b*(e^(2*a + 2*c) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(36) = 72$.

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.30

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx = -2 \left(\frac{e^{(2a)} \log(|e^{(2bx+2a)} - 1|)}{be^{(2a)} - be^{(4a+2c)}} + \frac{\log(|e^{(2bx)} - e^{(2c)}|)}{be^{(2a+2c)} - b} \right) e^{(a+c)}$$

input `integrate(-csch(b*x-c)*csch(b*x+a),x, algorithm="giac")`

output `-2*(e^(2*a)*log(abs(e^(2*b*x + 2*a) - 1)))/(b*e^(2*a) - b*e^(4*a + 2*c)) + log(abs(e^(2*b*x) - e^(2*c)))/(b*e^(2*a + 2*c) - b)*e^(a + c)`

Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 269, normalized size of antiderivative = 8.15

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx =$$

$$4 \operatorname{atan} \left(\frac{e^{2a} e^{2bx} \left(\frac{2e^a e^c}{b(e^{2a} e^{2c})^{3/2}} + \frac{2e^{-3a} e^{-3c} (e^{2a} e^{2c} + 1) (b\sqrt{e^{2a} e^{2c} + b(e^{2a} e^{2c})^{3/2}})}{\sqrt{-b^2 (e^{2a} e^{2c} - 1)^2} \sqrt{2b^2 e^{2a} e^{2c} - b^2 - b^2 e^{4a} e^{4c}}} \right)}{\sqrt{2b^2 e^{2a+2c} - b^2 e^{4a+4c} - b^2}} \right) - \frac{b e^{-3a} e^{-3c}}{\sqrt{2b^2 e^{2a+2c} - b^2 e^{4a+4c} - b^2}}$$

input `int(1/(sinh(a + b*x)*sinh(c - b*x)),x)`output
$$-(4*\operatorname{atan}((\exp(2*a)*\exp(2*b*x))*((2*\exp(a)*\exp(c))/(b*(\exp(2*a)*\exp(2*c))^{3/2})) + (2*\exp(-3*a)*\exp(-3*c)*(\exp(2*a)*\exp(2*c) + 1)*(b*(\exp(2*a)*\exp(2*c))^{1/2} + b*(\exp(2*a)*\exp(2*c))^{3/2}))/((-b^2*(\exp(2*a)*\exp(2*c) - 1)^{2/2}*(2*b^2*\exp(2*a)*\exp(2*c) - b^2 - b^2*\exp(4*a)*\exp(4*c))^{1/2}))* (2*b^2*\exp(2*a)*\exp(2*c) - b^2 - b^2*\exp(4*a)*\exp(4*c))^{1/2})/4 - (b*\exp(-3*a)*\exp(-3*c)*(\exp(2*a)*\exp(2*c) + 1)*(\exp(2*a)*\exp(2*c))^{3/2}))/(-b^2*(\exp(2*a)*\exp(2*c) - 1)^{2/2})^{1/2})*\exp(2*a + 2*c)^{1/2})/(2*b^2*\exp(2*a + 2*c) - b^2*\exp(4*a + 4*c) - b^2)^{1/2})$$
Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.12

$$\int \operatorname{csch}(c - bx)\operatorname{csch}(a + bx) dx = \frac{2e^{a+c}(\log(e^{bx+a} - 1) + \log(e^{bx+a} + 1) - \log(e^{bx} + e^c) - \log(e^{bx} - e^c))}{b(e^{2a+2c} - 1)}$$

input `int(-csch(b*x-c)*csch(b*x+a),x)`output
$$(2e^{a+c}(\log(e^{a+b*x} - 1) + \log(e^{a+b*x} + 1) - \log(e^{b*x} + e^c) - \log(e^{b*x} - e^c)))/(b*(e^{2a+2c} - 1)))$$

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	1169
4.2	Links to plain text integration problems used in this report for each CAS .	1187

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
      ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
    ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]], 2]],
      Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result   := ExpnType(result);
      ExpnType_optimal  := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                convert(leaf_count_result,string)," $ vs. $2(",
                convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
            convert(ExpnType_result,string)," vs. order ",
            convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```



```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```



```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file