

# Computer Algebra Independent Integration Tests

Summer 2024

7-Inverse-hyperbolic-functions/7.3-Inverse-hyperbolic-  
tangent/336-7.3

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 7:35am

# Contents

<b>1</b>	<b>Introduction</b>	<b>14</b>
1.1	Listing of CAS systems tested . . . . .	15
1.2	Results . . . . .	16
1.3	Time and leaf size Performance . . . . .	20
1.4	Performance based on number of rules Rubi used . . . . .	22
1.5	Performance based on number of steps Rubi used . . . . .	23
1.6	Solved integrals histogram based on leaf size of result . . . . .	24
1.7	Solved integrals histogram based on CPU time used . . . . .	25
1.8	Leaf size vs. CPU time used . . . . .	26
1.9	list of integrals with no known antiderivative . . . . .	27
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	27
1.11	list of integrals solved by CAS but failed verification . . . . .	27
1.12	Timing . . . . .	28
1.13	Verification . . . . .	28
1.14	Important notes about some of the results . . . . .	29
1.15	Current tree layout of integration tests . . . . .	32
1.16	Design of the test system . . . . .	33
<b>2</b>	<b>detailed summary tables of results</b>	<b>34</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	35
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	42
2.3	Detailed conclusion table specific for Rubi results . . . . .	133
<b>3</b>	<b>Listing of integrals</b>	<b>145</b>
3.1	$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$ . . . . .	157
3.2	$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$ . . . . .	165
3.3	$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$ . . . . .	171
3.4	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx$ . . . . .	177

3.5	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$	186
3.6	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$	191
3.7	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$	196
3.8	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$	202
3.9	$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	208
3.10	$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	215
3.11	$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	221
3.12	$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	227
3.13	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$	232
3.14	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$	238
3.15	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$	244
3.16	$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	251
3.17	$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	258
3.18	$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	264
3.19	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$	270
3.20	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$	276
3.21	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$	282
3.22	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$	289
3.23	$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	296
3.24	$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	304
3.25	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$	311
3.26	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$	318
3.27	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$	326
3.28	$\int x^3 \operatorname{arctanh}(a + bx^4) dx$	335

3.29	$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx$	342
3.30	$\int \frac{(a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^n}{1-c^2x^2} dx$	348
3.31	$\int \frac{(a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^3}{1-c^2x^2} dx$	353
3.32	$\int \frac{(a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^2}{1-c^2x^2} dx$	362
3.33	$\int \frac{a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}})}{1-c^2x^2} dx$	371
3.34	$\int \frac{1}{(1-c^2x^2)(a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))} dx$	377
3.35	$\int \frac{1}{(1-c^2x^2)(a + b \operatorname{arctanh}(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^2} dx$	382
3.36	$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$	388
3.37	$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx$	394
3.38	$\int x \operatorname{arctanh}(\tanh(a + bx)) dx$	399
3.39	$\int \operatorname{arctanh}(\tanh(a + bx)) dx$	404
3.40	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx$	409
3.41	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx$	414
3.42	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx$	419
3.43	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx$	424
3.44	$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$	429
3.45	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx$	435
3.46	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx$	440
3.47	$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx$	445
3.48	$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx$	450
3.49	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx$	455
3.50	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx$	461
3.51	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx$	467
3.52	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx$	472
3.53	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx$	477
3.54	$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$	483
3.55	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx$	491
3.56	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx$	497
3.57	$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx$	503
3.58	$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx$	509
3.59	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx$	515
3.60	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx$	522
3.61	$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx$	529



3.62	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^4} dx$	536
3.63	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^5} dx$	542
3.64	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^6} dx$	547
3.65	$\int x^m \operatorname{arctanh}(\tanh(a+bx))^4 dx$	553
3.66	$\int x^6 \operatorname{arctanh}(\tanh(a+bx))^4 dx$	561
3.67	$\int x^5 \operatorname{arctanh}(\tanh(a+bx))^4 dx$	568
3.68	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^4 dx$	575
3.69	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^4 dx$	581
3.70	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^4 dx$	587
3.71	$\int x \operatorname{arctanh}(\tanh(a+bx))^4 dx$	593
3.72	$\int \operatorname{arctanh}(\tanh(a+bx))^4 dx$	599
3.73	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx$	605
3.74	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^2} dx$	612
3.75	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^3} dx$	618
3.76	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^4} dx$	625
3.77	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^5} dx$	632
3.78	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^6} dx$	638
3.79	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^7} dx$	644
3.80	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^8} dx$	650
3.81	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx$	656
3.82	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{10}} dx$	663
3.83	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{11}} dx$	670
3.84	$\int x \operatorname{arctanh}(\tanh(a+bx))^6 dx$	677
3.85	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))} dx$	684
3.86	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx$	689
3.87	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx$	695
3.88	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx$	701
3.89	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx$	706
3.90	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx$	711
3.91	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))} dx$	716
3.92	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))} dx$	722
3.93	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	729
3.94	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	734
3.95	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	742

3.96	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	749
3.97	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	755
3.98	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	760
3.99	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^2} dx$	765
3.100	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx$	771
3.101	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^2} dx$	778
3.102	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	786
3.103	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	791
3.104	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	799
3.105	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	806
3.106	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	812
3.107	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	817
3.108	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^3} dx$	822
3.109	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^3} dx$	829
3.110	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^3} dx$	837
3.111	$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	846
3.112	$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	854
3.113	$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	861
3.114	$\int x \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	867
3.115	$\int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	872
3.116	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$	877
3.117	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx$	883
3.118	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx$	889
3.119	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$	896
3.120	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	904
3.121	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	911
3.122	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	918
3.123	$\int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	924
3.124	$\int \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	930
3.125	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x} dx$	935
3.126	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^2} dx$	941
3.127	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^3} dx$	948
3.128	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^4} dx$	955

3.129	$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$	962
3.130	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$	969
3.131	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$	976
3.132	$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$	983
3.133	$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$	989
3.134	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx$	995
3.135	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^2} dx$	1002
3.136	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^3} dx$	1009
3.137	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx$	1016
3.138	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^5} dx$	1023
3.139	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx$	1031
3.140	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$	1040
3.141	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx$	1047
3.142	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx$	1054
3.143	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$	1060
3.144	$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1066
3.145	$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1071
3.146	$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1076
3.147	$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1083
3.148	$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1090
3.149	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1099
3.150	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1107
3.151	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1114
3.152	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1120
3.153	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1126
3.154	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1131
3.155	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1137
3.156	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1144
3.157	$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1152
3.158	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1161
3.159	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1168

3.160	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1175
3.161	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1181
3.162	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1186
3.163	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1191
3.164	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1198
3.165	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1205
3.166	$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1213
3.167	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1222
3.168	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1227
3.169	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1232
3.170	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx)) dx$	1237
3.171	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx$	1242
3.172	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{3/2}} dx$	1247
3.173	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{5/2}} dx$	1252
3.174	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{7/2}} dx$	1257
3.175	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1262
3.176	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1267
3.177	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1273
3.178	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1278
3.179	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx$	1283
3.180	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{3/2}} dx$	1288
3.181	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{5/2}} dx$	1294
3.182	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{7/2}} dx$	1300
3.183	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1306
3.184	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1312
3.185	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1318
3.186	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1324
3.187	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} dx$	1330
3.188	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx$	1336
3.189	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{5/2}} dx$	1342
3.190	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{7/2}} dx$	1348
3.191	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1354
3.192	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1362
3.193	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1369

3.194	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx$	1375
3.195	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx$	1381
3.196	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1386
3.197	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1392
3.198	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1398
3.199	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1405
3.200	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1413
3.201	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1421
3.202	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1427
3.203	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1433
3.204	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1440
3.205	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1447
3.206	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1455
3.207	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1465
3.208	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1473
3.209	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1480
3.210	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1486
3.211	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1493
3.212	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1501
3.213	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1510
3.214	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1521
3.215	$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	1534
3.216	$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	1541
3.217	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx$	1547
3.218	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx$	1553
3.219	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx$	1559
3.220	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx$	1564
3.221	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx$	1570
3.222	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx$	1576
3.223	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	1583
3.224	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	1590

3.225	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}} dx$	1597
3.226	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}} dx$	1603
3.227	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{5/2}} dx$	1609
3.228	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx$	1615
3.229	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx$	1620
3.230	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx$	1625
3.231	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{13/2}} dx$	1631
3.232	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	1638
3.233	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx$	1645
3.234	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{3/2}} dx$	1652
3.235	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}} dx$	1659
3.236	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{7/2}} dx$	1666
3.237	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx$	1673
3.238	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx$	1678
3.239	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx$	1683
3.240	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{15/2}} dx$	1689
3.241	$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1696
3.242	$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1702
3.243	$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1708
3.244	$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1714
3.245	$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1719
3.246	$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1724
3.247	$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1730
3.248	$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1736
3.249	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1743
3.250	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1751
3.251	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1758
3.252	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1764
3.253	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1770
3.254	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1775

3.255	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1780
3.256	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1786
3.257	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1793
3.258	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1801
3.259	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1808
3.260	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1814
3.261	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1819
3.262	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1824
3.263	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1830
3.264	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1837
3.265	$\int x^m \operatorname{arctanh}(\tanh(a+bx))^n dx$	1845
3.266	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1850
3.267	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1859
3.268	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1867
3.269	$\int x \operatorname{arctanh}(\tanh(a+bx))^n dx$	1874
3.270	$\int \operatorname{arctanh}(\tanh(a+bx))^n dx$	1880
3.271	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x} dx$	1886
3.272	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^2} dx$	1891
3.273	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^3} dx$	1896
3.274	$\int x^m \coth^{-1}(\tanh(a+bx)) dx$	1901
3.275	$\int x^2 \operatorname{arctanh}(\coth(a+bx)) dx$	1907
3.276	$\int x \operatorname{arctanh}(\coth(a+bx)) dx$	1912
3.277	$\int \operatorname{arctanh}(\coth(a+bx)) dx$	1917
3.278	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x} dx$	1922
3.279	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^2} dx$	1927
3.280	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^3} dx$	1932
3.281	$\int \operatorname{arctanh}(\cosh(x)) dx$	1937
3.282	$\int x \operatorname{arctanh}(\cosh(x)) dx$	1943
3.283	$\int x^2 \operatorname{arctanh}(\cosh(x)) dx$	1950
3.284	$\int x^2 \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1958
3.285	$\int x \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1968
3.286	$\int \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1977
3.287	$\int \frac{\operatorname{arctanh}(c+d \tanh(a+bx))}{x} dx$	1985
3.288	$\int x^3 \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	1990
3.289	$\int x^2 \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	1999

3.290	$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$	2008
3.291	$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$	2016
3.292	$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(a + bx))}{x} dx$	2022
3.293	$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$	2027
3.294	$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$	2036
3.295	$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$	2045
3.296	$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$	2053
3.297	$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx$	2059
3.298	$\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx$	2064
3.299	$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx$	2074
3.300	$\int \operatorname{arctanh}(c + d \coth(a + bx)) dx$	2083
3.301	$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx$	2091
3.302	$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$	2096
3.303	$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$	2105
3.304	$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$	2114
3.305	$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$	2122
3.306	$\int \frac{\operatorname{arctanh}(1 + d + d \coth(a + bx))}{x} dx$	2128
3.307	$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$	2133
3.308	$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$	2142
3.309	$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$	2151
3.310	$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$	2159
3.311	$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx$	2165
3.312	$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx$	2170
3.313	$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx$	2181
3.314	$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx$	2191
3.315	$\int \operatorname{arctanh}(\tan(a + bx)) dx$	2199
3.316	$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx$	2206
3.317	$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2211
3.318	$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2223
3.319	$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2232
3.320	$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx$	2240
3.321	$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2245
3.322	$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2254
3.323	$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2262
3.324	$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx$	2270
3.325	$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2275
3.326	$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2284



3.327	$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2292
3.328	$\int \frac{\operatorname{arctanh}(1+id-d \tan(a+bx))}{x} dx$	2299
3.329	$\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx$	2304
3.330	$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx$	2315
3.331	$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx$	2325
3.332	$\int \operatorname{arctanh}(\cot(a + bx)) dx$	2333
3.333	$\int \frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx} dx$	2340
3.334	$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2345
3.335	$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2357
3.336	$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2366
3.337	$\int \frac{\operatorname{arctanh}(c+d \cot(a+bx))}{x} dx$	2374
3.338	$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2379
3.339	$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2387
3.340	$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2395
3.341	$\int \frac{\operatorname{arctanh}(1+id+d \cot(a+bx))}{x} dx$	2402
3.342	$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2407
3.343	$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2415
3.344	$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2423
3.345	$\int \frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x} dx$	2430
3.346	$\int \operatorname{arctanh}(e^x) dx$	2435
3.347	$\int x \operatorname{arctanh}(e^x) dx$	2440
3.348	$\int x^2 \operatorname{arctanh}(e^x) dx$	2446
3.349	$\int \operatorname{arctanh}(e^{a+bx}) dx$	2452
3.350	$\int x \operatorname{arctanh}(e^{a+bx}) dx$	2457
3.351	$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx$	2463
3.352	$\int \operatorname{arctanh}(a + bf^{c+dx}) dx$	2470
3.353	$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$	2478
3.354	$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx$	2486
3.355	$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$	2495
3.356	$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx$	2503
3.357	$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx$	2510
3.358	$\int e^{c(a+bx)} \operatorname{arctanh}(\coth(ac + bcx)) dx$	2515
3.359	$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx$	2520
3.360	$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx$	2527
3.361	$\int \frac{(a+b \operatorname{arctanh}(cx^n))^{(d+e \log(fx^m))}}{x} dx$	2535

<b>4</b>	<b>Appendix</b>	<b>2541</b>
4.1	Listing of Grading functions	2541

4.2 Links to plain text integration problems used in this report for each CA9559

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	15
1.2	Results . . . . .	16
1.3	Time and leaf size Performance . . . . .	20
1.4	Performance based on number of rules Rubi used . . . . .	22
1.5	Performance based on number of steps Rubi used . . . . .	23
1.6	Solved integrals histogram based on leaf size of result . . . . .	24
1.7	Solved integrals histogram based on CPU time used . . . . .	25
1.8	Leaf size vs. CPU time used . . . . .	26
1.9	list of integrals with no known antiderivative . . . . .	27
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	27
1.11	list of integrals solved by CAS but failed verification . . . . .	27
1.12	Timing . . . . .	28
1.13	Verification . . . . .	28
1.14	Important notes about some of the results . . . . .	29
1.15	Current tree layout of integration tests . . . . .	32
1.16	Design of the test system . . . . .	33

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 361 ]. This is test number [ 336 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 361 )	0.00 ( 0 )
Mathematica	99.45 ( 359 )	0.55 ( 2 )
Fricas	96.95 ( 350 )	3.05 ( 11 )
Maple	94.74 ( 342 )	5.26 ( 19 )
Maxima	74.24 ( 268 )	25.76 ( 93 )
Giac	72.30 ( 261 )	27.70 ( 100 )
Mupad	66.20 ( 239 )	33.80 ( 122 )
Sympy	27.15 ( 98 )	72.85 ( 263 )
Reduce	20.50 ( 74 )	79.50 ( 287 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	93.906	0.277	1.108	4.709
Mathematica	87.535	3.601	3.601	5.263
Fricas	70.914	19.114	2.216	7.756
Maple	63.435	15.235	11.357	9.972
Maxima	62.050	7.479	0.000	30.471
Giac	60.111	7.202	0.277	32.410
Sympy	18.837	3.601	0.000	77.562
Mupad	0.000	61.496	0.000	38.504
Reduce	0.000	15.789	0.000	84.211

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.



System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	2	100.00	0.00	0.00
Fricas	11	100.00	0.00	0.00
Maple	19	100.00	0.00	0.00
Maxima	93	100.00	0.00	0.00
Giac	100	85.00	14.00	1.00
Mupad	122	0.00	100.00	0.00
Sympy	263	86.69	12.93	0.38
Reduce	287	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.09
Reduce	0.18
Maxima	0.28
Mathematica	0.38
Rubi	0.42
Giac	2.09
Maple	2.85
Mupad	3.72
Sympy	4.85

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	39.64	1.17	26.00	1.00
Sympy	51.48	1.39	49.00	1.13
Giac	60.79	0.96	46.00	0.64
Maxima	75.48	1.08	54.00	0.86
Mathematica	93.79	1.04	71.00	0.89
Rubi	109.37	1.09	88.00	1.05
Fricas	159.11	1.38	72.00	1.02
Mupad	367.64	4.36	211.00	3.17
Maple	410.93	2.80	95.00	1.03

Table 1.6: Leaf size performance for each CAS

# 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{30, 34, 35, 287, 292, 297, 301, 306, 311, 316, 320, 324, 328, 333, 337, 341, 345}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {28, 29, 355, 360}

Mathematica {319, 323, 327, 336, 340, 344}



**Maple** {282, 283, 284, 285, 288, 289, 290, 293, 294, 295, 298, 299, 302, 303, 304, 307, 308, 309, 312, 313, 314, 317, 318, 321, 322, 325, 326, 329, 330, 331, 334, 335, 338, 339, 342, 343, 355, 356, 359, 360, 361}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals.

These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'beselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	35
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	42
2.3	Detailed conclusion table specific for Rubi results . . . . .	133

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	35
Mma . . . . .	36
Maple . . . . .	36
Fricas . . . . .	37
Maxima . . . . .	38
Giac . . . . .	39
Mupad . . . . .	39
Sympy . . . . .	40
Reduce . . . . .	41

### Rubi

**A grade** { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31, 32, 33, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 338, 339, 340, 342, 343, 344, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361 }

**B grade** { 81 }

**C grade** { 4, 281, 282, 283 }

**F normal fail** { }

**F(-1) timedout fail** { }



**F(-2) exception fail { }**

## **Mma**

**A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 28, 29, 33, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 72, 73, 74, 75, 76, 77, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 313, 314, 315, 317, 318, 319, 321, 322, 325, 326, 330, 331, 332, 334, 335, 338, 339, 342, 343, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360 }**

**B grade { 47, 57, 71, 78, 84, 312, 323, 327, 329, 336, 340, 344, 346 }**

**C grade { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 361 }**

**F normal fail { 31, 32 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## **Maple**

**A grade { 4, 6, 7, 8, 13, 14, 28, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 72, 74, 75, 76, 77, 79, 80, 81, 82, 83, 87, 88, 89, 90, 91, 92, 96, 97, 98, 99, 100, 101, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 152, 153, 154, 155, 156, 157, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 193, 194, 195, 196, 197, 198, 200, 201, 202, 203, 204, 205, 206, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 219, 220, 221, 222, 223, 224, 225, 226, 228, 229, 230, 231, 232, 233, 234, 237, 238, 239,**

240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 261, 262, 263, 264, 268, 269, 270, 274, 275, 276, 277, 278, 279, 280, 281, 315, 332, 346, 347, 348, 349, 352, 357, 358 }

**B grade** { 1, 2, 3, 5, 9, 10, 11, 12, 15, 29, 31, 32, 33, 71, 73, 78, 84, 86, 94, 95, 103, 104, 134, 149, 150, 151, 158, 159, 191, 192, 199, 207, 218, 227, 235, 236, 260, 266, 267, 286, 291, 296, 300, 305, 310, 319, 323, 327, 336, 340, 344, 350, 351, 353, 354 }

**C grade** { 282, 283, 284, 285, 288, 289, 290, 293, 294, 295, 298, 299, 302, 303, 304, 307, 308, 309, 312, 313, 314, 317, 318, 321, 322, 325, 326, 329, 330, 331, 334, 335, 338, 339, 342, 343, 355, 356, 359, 360, 361 }

**F normal fail** { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 85, 93, 102, 265, 271, 272, 273 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 36, 37, 38, 39, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52, 53, 55, 56, 57, 59, 60, 61, 62, 63, 64, 66, 67, 68, 69, 70, 71, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 269, 270, 283, 338, 339, 340, 342, 343, 344, 352, 354, 356, 357, 359 }

**B grade** { 13, 14, 15, 29, 44, 54, 58, 65, 72, 84, 133, 162, 266, 267, 268, 281, 282, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 346, 347, 348, 349, 350, 351, 353, 355, 360, 361 }

**C grade** { 274, 275, 276, 277, 278, 279, 280, 358 }

**F normal fail** { 4, 31, 32, 33, 85, 93, 102, 265, 271, 272, 273 }

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## **Maxima**

**A grade { 5, 6, 7, 8, 9, 10, 11, 28, 29, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 73, 74, 75, 76, 77, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 120, 121, 122, 123, 124, 129, 130, 131, 132, 133, 140, 141, 142, 143, 144, 149, 150, 151, 152, 153, 158, 159, 160, 161, 162, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 218, 219, 220, 221, 222, 228, 229, 230, 231, 237, 238, 239, 240, 245, 246, 247, 248, 254, 255, 256, 262, 263, 264, 266, 267, 268, 269, 270, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 348, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359 }**

**B grade { 12, 48, 58, 71, 72, 78, 84, 315, 319, 321, 322, 323, 325, 326, 327, 332, 336, 338, 339, 340, 342, 343, 344, 346, 347, 349, 360 }**

**C grade { }**

**F normal fail { 1, 2, 3, 4, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 85, 93, 102, 116, 117, 118, 119, 125, 126, 127, 128, 134, 135, 136, 137, 138, 139, 145, 146, 147, 148, 154, 155, 156, 157, 163, 164, 165, 166, 215, 216, 217, 223, 224, 225, 226, 227, 232, 233, 234, 235, 236, 241, 242, 243, 244, 249, 250, 251, 252, 253, 257, 258, 259, 260, 261, 265, 271, 272, 273, 312, 313, 314, 317, 318, 329, 330, 331, 334, 335, 361 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## Giac

**A grade** { 12, 36, 37, 38, 39, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52, 53, 55, 56, 57, 59, 60, 61, 62, 63, 64, 66, 67, 68, 69, 70, 71, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 115, 116, 117, 118, 119, 125, 126, 127, 128, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 267, 268, 269, 270, 355, 356, 357, 358, 360 }

**B grade** { 28, 29, 58, 72, 84, 113, 114, 120, 121, 122, 123, 124, 129, 130, 131, 132, 133, 219, 266, 274, 275, 276, 277, 279, 280, 359 }

**C grade** { 278 }

**F normal fail** { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 44, 54, 65, 85, 93, 102, 265, 271, 272, 273, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 338, 339, 340, 342, 343, 344, 346, 347, 348, 349, 350, 351, 353, 354, 361 }

**F(-1) timedout fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15 }

**F(-2) exception fail** { 352 }

## Mupad

**A grade** { }

**B grade** { 12, 28, 29, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 219, 220, 221, 222, 228, 229,

230, 231, 237, 238, 239, 240, 245, 246, 247, 248, 253, 254, 255, 256, 260, 261, 262, 263, 264, 266, 267, 268, 269, 270, 274, 275, 276, 277, 278, 279, 280, 355, 356, 357, 358, 359, 360 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 85, 93, 102, 215, 216, 217, 218, 223, 224, 225, 226, 227, 232, 233, 234, 235, 236, 241, 242, 243, 244, 249, 250, 251, 252, 257, 258, 259, 265, 271, 272, 273, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 338, 339, 340, 342, 343, 344, 346, 347, 348, 349, 350, 351, 352, 353, 354, 361 }

**F(-2) exception fail** { }

## Sympy

**A grade** { 1, 2, 3, 9, 10, 11, 12, 28, 37, 38, 39, 41, 42, 43, 46, 47, 48, 51, 52, 53, 55, 56, 57, 58, 62, 64, 66, 67, 68, 69, 70, 71, 72, 77, 79, 80, 81, 82, 83, 84, 89, 107, 115, 123, 124, 130, 131, 132, 133, 144, 152, 153, 159, 160, 161, 162, 168, 169, 172, 173, 174, 176, 181, 182, 184, 189, 190, 357 }

**B grade** { 45, 63, 78, 97, 98, 105, 106, 270, 275, 276, 277, 279, 280 }

**C grade** { }

**F normal fail** { 4, 5, 6, 7, 8, 13, 14, 15, 17, 18, 19, 20, 24, 25, 26, 31, 32, 33, 36, 40, 44, 49, 50, 54, 59, 60, 61, 65, 73, 74, 75, 76, 85, 86, 87, 88, 90, 91, 92, 93, 94, 95, 96, 99, 100, 101, 102, 103, 104, 108, 109, 110, 111, 112, 113, 114, 116, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 145, 146, 147, 148, 149, 150, 151, 154, 155, 156, 157, 158, 163, 164, 165, 166, 170, 171, 177, 178, 179, 180, 185, 186, 187, 188, 191, 192, 193, 194, 195, 196, 197, 198, 200, 201, 202, 203, 204, 205, 206, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 223, 224, 225, 226, 227, 233, 234, 235, 241, 242, 243, 244, 245, 246, 250, 251, 252, 253, 254, 255, 258, 259, 260, 261, 262, 265, 266, 267, 268, 269, 271, 272, 273, 274, 278, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 338, 339, 340, 342, 343, 344, 346, 347, 348, 349, 350, 351, 352, 353, 354, 356, 358, 360 }

**F(-1) timedout fail** { 16, 21, 22, 23, 27, 129, 167, 175, 183, 199, 207, 220, 221, 222, 228, 229, 230, 231, 232, 236, 237, 238, 239, 240, 247, 248, 249, 256, 257, 263, 264, 355, 359, 361 }

}

**F(-2) exception fail { 29 }**

## **Reduce**

**A grade { }**

**B grade { 3, 12, 28, 29, 36, 39, 41, 42, 43, 48, 51, 52, 53, 58, 62, 63, 64, 72, 77, 78, 79, 80, 81, 82, 83, 89, 97, 98, 105, 106, 107, 114, 115, 123, 124, 132, 133, 143, 144, 152, 153, 161, 162, 172, 173, 174, 181, 182, 190, 269, 270, 274, 277, 279, 280, 357, 358 }**

**C grade { }**

**F normal fail { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 37, 38, 40, 44, 45, 46, 47, 49, 50, 54, 55, 56, 57, 59, 60, 61, 65, 66, 67, 68, 69, 70, 71, 73, 74, 75, 76, 84, 85, 86, 87, 88, 90, 91, 92, 93, 94, 95, 96, 99, 100, 101, 102, 103, 104, 108, 109, 110, 111, 112, 113, 116, 117, 118, 119, 120, 121, 122, 125, 126, 127, 128, 129, 130, 131, 134, 135, 136, 137, 138, 139, 140, 141, 142, 145, 146, 147, 148, 149, 150, 151, 154, 155, 156, 157, 158, 159, 160, 163, 164, 165, 166, 167, 168, 169, 170, 171, 175, 176, 177, 178, 179, 180, 183, 184, 185, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 271, 272, 273, 275, 276, 278, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 293, 294, 295, 296, 298, 299, 300, 302, 303, 304, 305, 307, 308, 309, 310, 312, 313, 314, 315, 317, 318, 319, 321, 322, 323, 325, 326, 327, 329, 330, 331, 332, 334, 335, 336, 338, 339, 340, 342, 343, 344, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 359, 360, 361 }**

**F(-1) timedout fail { }**

**F(-2) exception fail { }**

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	A	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	145	99	253	0	86	121	0	21	0
N.S.	1	1.14	0.78	1.99	0.00	0.68	0.95	0.00	0.17	0.00
time (sec)	N/A	0.269	0.055	0.052	0.000	0.095	0.908	0.000	0.164	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	A	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	115	88	205	0	75	95	0	21	0
N.S.	1	1.14	0.87	2.03	0.00	0.74	0.94	0.00	0.21	0.00
time (sec)	N/A	0.245	0.036	0.034	0.000	0.094	0.430	0.000	0.166	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	A	<b>F(-1)</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	85	76	157	0	59	66	0	132	0
N.S.	1	1.13	1.01	2.09	0.00	0.79	0.88	0.00	1.76	0.00
time (sec)	N/A	0.224	0.024	0.033	0.000	0.089	0.213	0.000	0.193	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	238	185	167	209	0	0	0	0	21	0
N.S.	1	0.78	0.70	0.88	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	0.676	1.537	0.217	0.000	0.000	0.000	0.000	0.171	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	50	111	51	54	0	0	21	0
N.S.	1	1.00	0.94	2.09	0.96	1.02	0.00	0.00	0.40	0.00
time (sec)	N/A	0.203	0.024	0.036	0.068	0.098	0.000	0.000	0.166	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	80	63	62	61	67	0	0	21	0
N.S.	1	1.01	0.80	0.78	0.77	0.85	0.00	0.00	0.27	0.00
time (sec)	N/A	0.254	0.030	0.039	0.036	0.098	0.000	0.000	0.174	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	110	74	110	102	78	0	0	21	0
N.S.	1	1.05	0.70	1.05	0.97	0.74	0.00	0.00	0.20	0.00
time (sec)	N/A	0.328	0.034	0.036	0.035	0.120	0.000	0.000	0.161	0.000



Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	140	85	158	125	89	0	0	21	0
N.S.	1	1.07	0.65	1.21	0.95	0.68	0.00	0.00	0.16	0.00
time (sec)	N/A	0.353	0.039	0.036	0.037	0.114	0.000	0.000	0.170	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	112	79	224	155	88	116	0	21	0
N.S.	1	0.98	0.69	1.96	1.36	0.77	1.02	0.00	0.18	0.00
time (sec)	N/A	0.388	0.039	0.037	0.034	0.102	1.433	0.000	0.187	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	93	68	176	127	77	90	0	21	0
N.S.	1	1.02	0.75	1.93	1.40	0.85	0.99	0.00	0.23	0.00
time (sec)	N/A	0.405	0.040	0.037	0.033	0.111	0.613	0.000	0.173	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	72	56	128	99	65	65	0	21	0
N.S.	1	1.06	0.82	1.88	1.46	0.96	0.96	0.00	0.31	0.00
time (sec)	N/A	0.346	0.036	0.036	0.044	0.104	0.301	0.000	0.178	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	76	65	51	36	59	61	32
N.S.	1	1.00	1.00	1.90	1.62	1.28	0.90	1.48	1.52	0.80
time (sec)	N/A	0.320	0.008	0.024	0.032	0.117	0.201	0.148	0.177	2.977

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	61	84	0	273	0	0	21	0
N.S.	1	1.00	1.11	1.53	0.00	4.96	0.00	0.00	0.38	0.00
time (sec)	N/A	0.339	0.030	0.036	0.000	0.138	0.000	0.000	0.167	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	81	92	123	0	340	0	0	21	0
N.S.	1	0.95	1.08	1.45	0.00	4.00	0.00	0.00	0.25	0.00
time (sec)	N/A	0.262	0.059	0.034	0.000	0.137	0.000	0.000	0.176	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	107	171	0	383	0	0	21	0
N.S.	1	1.00	0.96	1.54	0.00	3.45	0.00	0.00	0.19	0.00
time (sec)	N/A	0.284	0.070	0.035	0.000	0.141	0.000	0.000	0.171	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	196	214	161	0	0	94	0	0	23	0
N.S.	1	1.09	0.82	0.00	0.00	0.48	0.00	0.00	0.12	0.00
time (sec)	N/A	0.321	0.354	0.000	0.000	0.118	0.000	0.000	0.331	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	168	182	147	0	0	83	0	0	23	0
N.S.	1	1.08	0.88	0.00	0.00	0.49	0.00	0.00	0.14	0.00
time (sec)	N/A	0.298	0.263	0.000	0.000	0.125	0.000	0.000	0.285	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	142	150	135	0	0	69	0	0	20	0
N.S.	1	1.06	0.95	0.00	0.00	0.49	0.00	0.00	0.14	0.00
time (sec)	N/A	0.269	0.200	0.000	0.000	0.112	0.000	0.000	0.224	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	111	0	0	50	0	0	23	0
N.S.	1	1.00	0.98	0.00	0.00	0.44	0.00	0.00	0.20	0.00
time (sec)	N/A	0.252	0.093	0.000	0.000	0.127	0.000	0.000	0.197	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	145	150	142	0	0	75	0	0	23	0
N.S.	1	1.03	0.98	0.00	0.00	0.52	0.00	0.00	0.16	0.00
time (sec)	N/A	0.272	0.153	0.000	0.000	0.099	0.000	0.000	0.204	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	173	182	154	0	0	90	0	0	23	0
N.S.	1	1.05	0.89	0.00	0.00	0.52	0.00	0.00	0.13	0.00
time (sec)	N/A	0.295	0.225	0.000	0.000	0.103	0.000	0.000	0.210	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	201	214	163	0	0	101	0	0	23	0
N.S.	1	1.06	0.81	0.00	0.00	0.50	0.00	0.00	0.11	0.00
time (sec)	N/A	0.319	0.311	0.000	0.000	0.105	0.000	0.000	0.212	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	297	317	124	0	0	92	0	0	23	0
N.S.	1	1.07	0.42	0.00	0.00	0.31	0.00	0.00	0.08	0.00
time (sec)	N/A	0.425	0.077	0.000	0.000	0.132	0.000	0.000	0.311	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	269	285	109	0	0	77	0	0	21	0
N.S.	1	1.06	0.41	0.00	0.00	0.29	0.00	0.00	0.08	0.00
time (sec)	N/A	0.526	0.067	0.000	0.000	0.111	0.000	0.000	0.240	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	232	249	85	0	0	52	0	0	23	0
N.S.	1	1.07	0.37	0.00	0.00	0.22	0.00	0.00	0.10	0.00
time (sec)	N/A	0.555	0.074	0.000	0.000	0.112	0.000	0.000	0.182	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	272	281	118	0	0	82	0	0	23	0
N.S.	1	1.03	0.43	0.00	0.00	0.30	0.00	0.00	0.08	0.00
time (sec)	N/A	0.604	0.074	0.000	0.000	0.103	0.000	0.000	0.216	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	302	313	131	0	0	98	0	0	23	0
N.S.	1	1.04	0.43	0.00	0.00	0.32	0.00	0.00	0.08	0.00
time (sec)	N/A	0.656	0.069	0.000	0.000	0.108	0.000	0.000	0.222	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	36	39	39	37	59	60	223	377	90
N.S.	1	0.82	0.89	0.89	0.84	1.34	1.36	5.07	8.57	2.05
time (sec)	N/A	0.316	0.015	0.357	0.027	0.096	0.766	0.124	0.221	3.347

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F(-2)	B	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	38	42	121	40	109	0	124	56	56
N.S.	1	0.81	0.89	2.57	0.85	2.32	0.00	2.64	1.19	1.19
time (sec)	N/A	0.299	0.027	4.930	0.027	0.094	0.000	0.139	0.180	3.739

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	39	38	34	38	39	39
N.S.	1	1.00	1.05	0.90	0.98	0.95	0.85	0.95	0.98	0.98
time (sec)	N/A	0.252	0.137	0.404	0.550	0.123	9.707	0.526	0.294	3.637

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	409	402	0	1623	0	0	0	0	150	0
N.S.	1	0.98	0.00	3.97	0.00	0.00	0.00	0.00	0.37	0.00
time (sec)	N/A	1.200	0.000	1.837	0.000	0.000	0.000	0.000	0.273	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	268	271	0	893	0	0	0	0	108	0
N.S.	1	1.01	0.00	3.33	0.00	0.00	0.00	0.00	0.40	0.00
time (sec)	N/A	0.838	0.000	0.608	0.000	0.000	0.000	0.000	0.242	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	84	43	357	0	0	0	0	64	0
N.S.	1	0.94	0.48	4.01	0.00	0.00	0.00	0.00	0.72	0.00
time (sec)	N/A	0.483	0.266	0.329	0.000	0.000	0.000	0.000	0.189	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	39	48	61	38	63	39
N.S.	1	1.00	1.05	0.90	0.98	1.20	1.52	0.95	1.58	0.98
time (sec)	N/A	0.400	0.220	0.376	0.186	0.078	3.874	0.261	0.270	3.363

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	246	91	126	38	123	39
N.S.	1	1.00	1.05	0.90	6.15	2.28	3.15	0.95	3.08	0.98
time (sec)	N/A	0.390	0.830	0.385	0.237	0.089	11.367	0.604	0.370	4.089

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	34	41	38	62	0	43	38	96
N.S.	1	1.00	0.92	1.11	1.03	1.68	0.00	1.16	1.03	2.59
time (sec)	N/A	0.301	0.041	0.341	0.035	0.079	0.000	0.117	0.159	3.663

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	20	19	13	19	13	13	19
N.S.	1	1.00	0.87	0.87	0.83	0.57	0.83	0.57	0.57	0.83
time (sec)	N/A	0.264	0.014	0.168	0.076	0.063	0.112	0.106	0.167	0.137

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	20	19	13	19	13	11	19
N.S.	1	1.00	0.87	0.87	0.83	0.57	0.83	0.57	0.48	0.83
time (sec)	N/A	0.294	0.013	0.165	0.080	0.066	0.080	0.115	0.158	3.151

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	15	16	10	19	10	14	16
N.S.	1	1.00	1.12	0.94	1.00	0.62	1.19	0.62	0.88	1.00
time (sec)	N/A	0.211	0.007	0.145	0.071	0.071	0.070	0.111	0.161	0.046



Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	19	23	34	8	0	9	13	58
N.S.	1	1.00	0.90	1.10	1.62	0.38	0.00	0.43	0.62	2.76
time (sec)	N/A	0.218	0.011	0.155	0.033	0.077	0.000	0.111	0.158	3.177

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	18	18	17	13	14	12	19	17
N.S.	1	1.00	1.06	1.06	1.00	0.76	0.82	0.71	1.12	1.00
time (sec)	N/A	0.161	0.013	0.166	0.079	0.066	0.089	0.118	0.154	0.091

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	17	19	11	19	11	19	16
N.S.	1	1.00	0.78	0.74	0.83	0.48	0.83	0.48	0.83	0.70
time (sec)	N/A	0.164	0.012	0.158	0.079	0.078	0.193	0.116	0.180	3.147

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	19	19	13	20	13	19	19
N.S.	1	1.00	0.87	0.83	0.83	0.57	0.87	0.57	0.83	0.83
time (sec)	N/A	0.156	0.012	0.164	0.079	0.083	0.235	0.118	0.167	0.072

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	66	62	98	73	161	0	0	15	203
N.S.	1	0.93	0.87	1.38	1.03	2.27	0.00	0.00	0.21	2.86
time (sec)	N/A	0.225	0.108	0.613	0.089	0.084	0.000	0.000	0.164	3.289

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	45	37	37	36	24	78	24	15	36
N.S.	1	1.07	0.88	0.88	0.86	0.57	1.86	0.57	0.36	0.86
time (sec)	N/A	0.205	0.021	30.847	0.128	0.067	0.367	0.117	0.187	3.233

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	45	37	37	36	24	60	24	15	36
N.S.	1	1.07	0.88	0.88	0.86	0.57	1.43	0.57	0.36	0.86
time (sec)	N/A	0.197	0.033	27.172	0.122	0.079	0.272	0.118	0.171	3.205

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	74	37	36	24	41	24	13	36
N.S.	1	1.00	2.18	1.09	1.06	0.71	1.21	0.71	0.38	1.06
time (sec)	N/A	0.210	0.166	26.163	0.119	0.081	0.207	0.116	0.172	3.130

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	33	20	20	20	14	33
N.S.	1	1.00	1.00	0.94	2.06	1.25	1.25	1.25	0.88	2.06
time (sec)	N/A	0.155	0.006	0.247	0.120	0.078	0.091	0.107	0.167	3.090

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	50	53	67	20	20	0	21	15	183
N.S.	1	1.02	1.08	1.37	0.41	0.41	0.00	0.43	0.31	3.73
time (sec)	N/A	0.207	0.081	0.184	0.324	0.073	0.000	0.112	0.190	3.274

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	37	41	54	24	0	21	15	198
N.S.	1	1.00	0.95	1.05	1.38	0.62	0.00	0.54	0.38	5.08
time (sec)	N/A	0.200	0.031	0.193	0.085	0.081	0.000	0.116	0.174	0.198

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	42	35	34	26	32	22	38	34
N.S.	1	1.00	1.17	0.97	0.94	0.72	0.89	0.61	1.06	0.94
time (sec)	N/A	0.197	0.025	0.266	0.131	0.080	0.196	0.112	0.172	3.120

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	34	33	36	22	37	22	36	32
N.S.	1	1.00	1.10	1.06	1.16	0.71	1.19	0.71	1.16	1.03
time (sec)	N/A	0.164	0.029	0.257	0.131	0.068	0.237	0.118	0.181	3.079

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	64	37	36	36	24	39	24	36	36
N.S.	1	1.52	0.88	0.86	0.86	0.57	0.93	0.57	0.86	0.86
time (sec)	N/A	0.228	0.021	0.296	0.124	0.067	0.311	0.111	0.181	3.088

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	95	97	177	109	300	0	0	15	332
N.S.	1	0.86	0.88	1.61	0.99	2.73	0.00	0.00	0.14	3.02
time (sec)	N/A	0.278	0.108	4.467	0.138	0.084	0.000	0.000	0.201	3.420

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	67	54	56	54	35	80	35	15	53
N.S.	1	1.10	0.89	0.92	0.89	0.57	1.31	0.57	0.25	0.87
time (sec)	N/A	0.245	0.021	0.122	0.178	0.068	0.502	0.106	0.191	3.267

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	61	54	56	54	35	60	35	15	53
N.S.	1	1.15	1.02	1.06	1.02	0.66	1.13	0.66	0.28	1.00
time (sec)	N/A	0.251	0.018	0.196	0.181	0.069	0.359	0.114	0.188	3.272

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	99	54	54	34	41	34	13	53
N.S.	1	1.00	2.91	1.59	1.59	1.00	1.21	1.00	0.38	1.56
time (sec)	N/A	0.271	0.214	43.933	0.177	0.067	0.270	0.112	0.166	0.132

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	51	31	20	31	14	47
N.S.	1	1.00	1.00	0.94	3.19	1.94	1.25	1.94	0.88	2.94
time (sec)	N/A	0.233	0.006	44.282	0.165	0.074	0.122	0.110	0.177	3.202

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	79	104	147	31	31	0	32	15	306
N.S.	1	1.03	1.35	1.91	0.40	0.40	0.00	0.42	0.19	3.97
time (sec)	N/A	0.428	0.088	0.408	0.328	0.075	0.000	0.113	0.189	0.155

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	62	85	65	36	0	33	15	415
N.S.	1	1.00	0.91	1.25	0.96	0.53	0.00	0.49	0.22	6.10
time (sec)	N/A	0.397	0.029	0.411	0.366	0.068	0.000	0.112	0.181	3.396

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	61	66	59	72	37	0	31	15	365
N.S.	1	1.02	1.10	0.98	1.20	0.62	0.00	0.52	0.25	6.08
time (sec)	N/A	0.405	0.031	0.430	0.138	0.073	0.000	0.110	0.179	0.223

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	60	52	52	37	51	35	55	51
N.S.	1	1.00	1.09	0.95	0.95	0.67	0.93	0.64	1.00	0.93
time (sec)	N/A	0.398	0.017	1.226	0.195	0.070	0.255	0.112	0.167	3.301

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	50	49	53	33	56	33	53	48
N.S.	1	1.00	1.61	1.58	1.71	1.06	1.81	1.06	1.71	1.55
time (sec)	N/A	0.256	0.019	1.273	0.181	0.077	0.305	0.106	0.167	3.231

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	54	53	54	35	60	35	53	53
N.S.	1	1.00	0.84	0.83	0.84	0.55	0.94	0.55	0.83	0.83
time (sec)	N/A	0.225	0.025	1.296	0.188	0.066	0.423	0.111	0.169	3.308

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	124	137	278	145	483	0	0	15	479
N.S.	1	0.81	0.89	1.81	0.94	3.14	0.00	0.00	0.10	3.11
time (sec)	N/A	0.349	0.122	46.254	0.193	0.087	0.000	0.000	0.175	3.660

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	89	71	74	72	46	134	46	15	242
N.S.	1	1.11	0.89	0.92	0.90	0.58	1.68	0.58	0.19	3.02
time (sec)	N/A	0.295	0.042	0.215	0.220	0.069	2.573	0.118	0.167	3.332

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	89	71	74	72	46	117	46	15	242
N.S.	1	1.11	0.89	0.92	0.90	0.58	1.46	0.58	0.19	3.02
time (sec)	N/A	0.294	0.023	0.216	0.222	0.068	1.591	0.114	0.167	3.491

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	89	71	74	72	46	100	46	15	77
N.S.	1	1.11	0.89	0.92	0.90	0.58	1.25	0.58	0.19	0.96
time (sec)	N/A	0.299	0.026	0.207	0.226	0.066	1.132	0.107	0.167	3.468

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	88	71	74	72	45	78	45	15	70
N.S.	1	1.22	0.99	1.03	1.00	0.62	1.08	0.62	0.21	0.97
time (sec)	N/A	0.296	0.018	0.132	0.229	0.074	0.729	0.111	0.164	3.887

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	61	71	74	72	45	60	45	15	70
N.S.	1	1.15	1.34	1.40	1.36	0.85	1.13	0.85	0.28	1.32
time (sec)	N/A	0.241	0.036	0.131	0.222	0.067	0.493	0.114	0.170	0.182

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	125	71	72	46	41	46	13	70
N.S.	1	1.00	3.68	2.09	2.12	1.35	1.21	1.35	0.38	2.06
time (sec)	N/A	0.187	0.190	27.618	0.212	0.069	0.379	0.117	0.178	3.481



Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	69	42	20	42	14	67
N.S.	1	1.00	1.00	0.94	4.31	2.62	1.25	2.62	0.88	4.19
time (sec)	N/A	0.154	0.006	29.598	0.220	0.069	0.181	0.105	0.177	0.129

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	108	175	265	42	42	0	43	15	423
N.S.	1	1.03	1.67	2.52	0.40	0.40	0.00	0.41	0.14	4.03
time (sec)	N/A	0.334	0.111	0.309	0.326	0.070	0.000	0.117	0.177	0.165

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	97	85	165	77	47	0	44	15	553
N.S.	1	1.02	0.89	1.74	0.81	0.49	0.00	0.46	0.16	5.82
time (sec)	N/A	0.307	0.052	0.332	0.371	0.076	0.000	0.111	0.175	0.179

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	88	81	103	83	47	0	43	15	672
N.S.	1	1.01	0.93	1.18	0.95	0.54	0.00	0.49	0.17	7.72
time (sec)	N/A	0.319	0.027	0.369	0.412	0.085	0.000	0.111	0.184	3.848

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	83	82	77	91	48	0	42	15	571
N.S.	1	1.08	1.06	1.00	1.18	0.62	0.00	0.55	0.19	7.42
time (sec)	N/A	0.502	0.029	0.618	0.183	0.085	0.000	0.111	0.170	4.105

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	78	69	72	48	70	46	72	68
N.S.	1	1.00	1.05	0.93	0.97	0.65	0.95	0.62	0.97	0.92
time (sec)	N/A	0.493	0.023	2.241	0.228	0.067	0.320	0.114	0.179	3.814

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	66	65	70	44	75	44	70	64
N.S.	1	1.00	2.13	2.10	2.26	1.42	2.42	1.42	2.26	2.06
time (sec)	N/A	0.284	0.037	1.991	0.232	0.073	0.442	0.115	0.182	3.529

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	71	70	72	46	78	46	70	70
N.S.	1	1.00	1.11	1.09	1.12	0.72	1.22	0.72	1.09	1.09
time (sec)	N/A	0.365	0.025	2.075	0.230	0.068	0.612	0.106	0.181	3.790

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	116	71	70	72	46	80	46	70	70
N.S.	1	1.18	0.72	0.71	0.73	0.47	0.82	0.47	0.71	0.71
time (sec)	N/A	0.515	0.032	2.178	0.235	0.077	0.891	0.112	0.179	3.317

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	168	71	70	72	46	76	46	70	70
N.S.	1	2.10	0.89	0.88	0.90	0.58	0.95	0.58	0.88	0.88
time (sec)	N/A	0.429	0.024	2.442	0.240	0.064	1.208	0.112	0.185	3.496

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	89	71	70	72	46	76	46	70	70
N.S.	1	1.11	0.89	0.88	0.90	0.58	0.95	0.58	0.88	0.88
time (sec)	N/A	0.297	0.041	2.663	0.225	0.077	1.882	0.109	0.176	3.529

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	89	71	70	72	46	78	46	70	70
N.S.	1	1.11	0.89	0.88	0.90	0.58	0.98	0.58	0.88	0.88
time (sec)	N/A	0.297	0.025	3.049	0.244	0.071	2.885	0.114	0.175	3.247

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	177	105	110	68	41	68	13	104
N.S.	1	1.00	5.21	3.09	3.24	2.00	1.21	2.00	0.38	3.06
time (sec)	N/A	0.192	0.231	41.608	0.326	0.072	0.762	0.111	0.179	3.387

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	51	0	0	0	0	0	15	0
N.S.	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.197	0.080	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	87	79	163	42	41	0	43	15	354
N.S.	1	1.07	0.98	2.01	0.52	0.51	0.00	0.53	0.19	4.37
time (sec)	N/A	0.312	0.033	1.040	0.183	0.077	0.000	0.122	0.160	3.399

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	59	55	86	29	29	0	30	15	234
N.S.	1	1.05	0.98	1.54	0.52	0.52	0.00	0.54	0.27	4.18
time (sec)	N/A	0.246	0.026	0.287	0.184	0.073	0.000	0.112	0.192	0.287

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	32	18	17	0	19	13	108
N.S.	1	1.00	1.00	1.03	0.58	0.55	0.00	0.61	0.42	3.48
time (sec)	N/A	0.191	0.019	0.174	0.184	0.076	0.000	0.112	0.197	0.160

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	13	10	17	11	12	12
N.S.	1	1.00	1.00	1.08	1.08	0.83	1.42	0.92	1.00	1.00
time (sec)	N/A	0.146	0.096	0.129	0.117	0.070	0.258	0.107	0.162	0.165

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	29	43	18	16	0	20	15	107
N.S.	1	1.00	0.66	0.98	0.41	0.36	0.00	0.45	0.34	2.43
time (sec)	N/A	0.211	0.017	9.249	0.192	0.077	0.000	0.106	0.179	5.906

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	81	45	64	28	26	0	30	15	210
N.S.	1	1.25	0.69	0.98	0.43	0.40	0.00	0.46	0.23	3.23
time (sec)	N/A	0.280	0.019	0.089	0.181	0.078	0.000	0.116	0.178	5.934

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	121	66	87	40	41	0	45	15	286
N.S.	1	1.32	0.72	0.95	0.43	0.45	0.00	0.49	0.16	3.11
time (sec)	N/A	0.349	0.022	0.095	0.190	0.076	0.000	0.109	0.189	5.661

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	51	0	0	0	0	0	46	0
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.00	0.00	0.71	0.00
time (sec)	N/A	0.232	0.836	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	110	106	262	70	73	0	62	15	669
N.S.	1	1.12	1.08	2.67	0.71	0.74	0.00	0.63	0.15	6.83
time (sec)	N/A	0.611	0.062	1.221	0.352	0.081	0.000	0.113	0.176	3.261

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	82	83	169	59	62	0	48	15	490
N.S.	1	1.09	1.11	2.25	0.79	0.83	0.00	0.64	0.20	6.53
time (sec)	N/A	0.490	0.037	0.340	0.338	0.077	0.000	0.114	0.169	3.330

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	54	56	86	44	47	0	34	15	302
N.S.	1	1.08	1.12	1.72	0.88	0.94	0.00	0.68	0.30	6.04
time (sec)	N/A	0.398	0.048	0.230	0.342	0.074	0.000	0.113	0.171	3.324

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	27	35	26	28	94	24	34	28
N.S.	1	1.00	0.96	1.25	0.93	1.00	3.36	0.86	1.21	1.00
time (sec)	N/A	0.319	0.098	0.145	0.332	0.075	13.323	0.108	0.168	0.090

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	12	13	71	12	14	14
N.S.	1	1.00	1.00	1.07	0.86	0.93	5.07	0.86	1.00	1.00
time (sec)	N/A	0.251	0.006	0.116	0.112	0.077	13.106	0.111	0.163	0.149

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	88	53	67	28	39	0	31	15	359
N.S.	1	1.26	0.76	0.96	0.40	0.56	0.00	0.44	0.21	5.13
time (sec)	N/A	0.350	0.093	0.115	0.358	0.073	0.000	0.119	0.172	5.443

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	135	70	91	45	63	0	45	15	432
N.S.	1	1.32	0.69	0.89	0.44	0.62	0.00	0.44	0.15	4.24
time (sec)	N/A	0.410	0.039	0.077	0.356	0.083	0.000	0.109	0.185	5.257

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	187	92	116	64	86	0	64	15	660
N.S.	1	1.31	0.64	0.81	0.45	0.60	0.00	0.45	0.10	4.62
time (sec)	N/A	0.476	0.032	0.085	0.350	0.084	0.000	0.108	0.183	5.114

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	94	97	51	0	0	0	0	0	49	0
N.S.	1	1.03	0.54	0.00	0.00	0.00	0.00	0.00	0.52	0.00
time (sec)	N/A	0.283	0.864	0.000	0.000	0.000	0.000	0.000	0.191	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	107	114	266	81	95	0	61	15	867
N.S.	1	1.16	1.24	2.89	0.88	1.03	0.00	0.66	0.16	9.42
time (sec)	N/A	0.372	0.030	0.369	0.594	0.084	0.000	0.118	0.174	3.241



Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	81	86	169	69	83	0	44	15	620
N.S.	1	1.14	1.21	2.38	0.97	1.17	0.00	0.62	0.21	8.73
time (sec)	N/A	0.301	0.048	0.231	0.590	0.085	0.000	0.117	0.164	3.310

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	52	49	54	48	61	112	37	53	46
N.S.	1	1.11	1.04	1.15	1.02	1.30	2.38	0.79	1.13	0.98
time (sec)	N/A	0.243	0.030	0.164	0.591	0.075	26.934	0.114	0.166	3.033

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	27	26	32	32	100	18	28	25
N.S.	1	1.00	0.79	0.76	0.94	0.94	2.94	0.53	0.82	0.74
time (sec)	N/A	0.190	0.099	0.148	0.594	0.074	25.883	0.113	0.166	0.084

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	12	24	24	12	14	14
N.S.	1	1.00	1.00	0.94	0.75	1.50	1.50	0.75	0.88	0.88
time (sec)	N/A	0.154	0.007	0.106	0.111	0.071	25.585	0.105	0.169	3.013

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	134	74	92	51	80	0	43	15	645
N.S.	1	1.38	0.76	0.95	0.53	0.82	0.00	0.44	0.15	6.65
time (sec)	N/A	0.350	0.113	0.080	0.585	0.081	0.000	0.111	0.186	5.427

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	181	93	117	69	109	0	60	15	804
N.S.	1	1.38	0.71	0.89	0.53	0.83	0.00	0.46	0.11	6.14
time (sec)	N/A	0.535	0.034	0.113	0.596	0.083	0.000	0.112	0.169	5.111

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	231	107	145	86	130	0	73	15	909
N.S.	1	1.36	0.63	0.85	0.51	0.76	0.00	0.43	0.09	5.35
time (sec)	N/A	0.779	0.029	0.092	0.605	0.084	0.000	0.116	0.176	4.821

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	125	83	154	64	64	0	150	14	811
N.S.	1	1.24	0.82	1.52	0.63	0.63	0.00	1.49	0.14	8.03
time (sec)	N/A	0.581	0.027	0.277	0.189	0.077	0.000	0.118	0.180	3.031

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	94	66	124	53	53	0	125	14	648
N.S.	1	1.18	0.82	1.55	0.66	0.66	0.00	1.56	0.18	8.10
time (sec)	N/A	0.387	0.024	0.246	0.186	0.081	0.000	0.119	0.171	3.085

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	67	49	69	42	42	0	102	14	485
N.S.	1	1.14	0.83	1.17	0.71	0.71	0.00	1.73	0.24	8.22
time (sec)	N/A	0.244	0.024	0.243	0.200	0.079	0.000	0.121	0.168	3.103

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	32	42	30	30	0	75	34	151
N.S.	1	1.00	0.84	1.11	0.79	0.79	0.00	1.97	0.89	3.97
time (sec)	N/A	0.192	0.154	0.243	0.191	0.078	0.000	0.112	0.163	3.186

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	12	12	26	18	20	95
N.S.	1	1.00	1.00	0.83	0.67	0.67	1.44	1.00	1.11	5.28
time (sec)	N/A	0.148	0.007	0.252	0.168	0.079	0.138	0.122	0.179	3.191

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	61	54	0	70	0	40	14	308
N.S.	1	1.00	0.97	0.86	0.00	1.11	0.00	0.63	0.22	4.89
time (sec)	N/A	0.239	0.103	0.224	0.000	0.086	0.000	0.116	0.169	4.422

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	65	63	0	90	0	48	42	341
N.S.	1	1.00	0.98	0.95	0.00	1.36	0.00	0.73	0.64	5.17
time (sec)	N/A	0.217	0.027	0.214	0.000	0.083	0.000	0.113	0.210	9.223

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	124	89	92	0	116	0	75	44	741
N.S.	1	0.99	0.71	0.74	0.00	0.93	0.00	0.60	0.35	5.93
time (sec)	N/A	0.326	0.071	0.223	0.000	0.103	0.000	0.105	0.216	7.994

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	179	196	115	185	0	142	0	81	44	964
N.S.	1	1.09	0.64	1.03	0.00	0.79	0.00	0.45	0.25	5.39
time (sec)	N/A	0.450	0.066	0.208	0.000	0.093	0.000	0.119	0.215	7.568

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	125	83	154	64	75	0	241	21	1813
N.S.	1	1.24	0.82	1.52	0.63	0.74	0.00	2.39	0.21	17.95
time (sec)	N/A	0.344	0.026	0.248	0.193	0.087	0.000	0.115	0.184	3.230

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	96	66	124	53	64	0	205	21	1483
N.S.	1	1.20	0.82	1.55	0.66	0.80	0.00	2.56	0.26	18.54
time (sec)	N/A	0.284	0.026	0.244	0.180	0.081	0.000	0.117	0.196	3.241

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	67	49	69	42	53	0	168	21	1153
N.S.	1	1.14	0.83	1.17	0.71	0.90	0.00	2.85	0.36	19.54
time (sec)	N/A	0.331	0.021	0.236	0.191	0.076	0.000	0.115	0.177	3.182

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	32	42	31	41	49	131	36	823
N.S.	1	1.00	0.84	1.11	0.82	1.08	1.29	3.45	0.95	21.66
time (sec)	N/A	0.314	0.165	0.237	0.189	0.083	1.652	0.115	0.171	3.200

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	12	28	26	84	22	97
N.S.	1	1.00	1.00	0.83	0.67	1.56	1.44	4.67	1.22	5.39
time (sec)	N/A	0.224	0.007	0.240	0.169	0.089	0.811	0.114	0.173	3.241

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	94	80	131	0	85	0	57	21	501
N.S.	1	1.03	0.88	1.44	0.00	0.93	0.00	0.63	0.23	5.51
time (sec)	N/A	0.384	0.095	0.192	0.000	0.080	0.000	0.114	0.177	7.581

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	85	79	85	0	99	0	63	21	459
N.S.	1	1.05	0.98	1.05	0.00	1.22	0.00	0.78	0.26	5.67
time (sec)	N/A	0.416	0.027	0.202	0.000	0.076	0.000	0.111	0.166	4.272

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	90	88	91	0	121	0	73	66	609
N.S.	1	0.98	0.96	0.99	0.00	1.32	0.00	0.79	0.72	6.62
time (sec)	N/A	0.370	0.043	0.198	0.000	0.087	0.000	0.122	0.199	8.040

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	148	117	116	0	142	0	81	66	1019
N.S.	1	1.01	0.80	0.79	0.00	0.97	0.00	0.55	0.45	6.98
time (sec)	N/A	0.418	0.061	0.217	0.000	0.099	0.000	0.114	0.188	7.470

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	125	83	154	64	86	0	344	23	2681
N.S.	1	1.24	0.82	1.52	0.63	0.85	0.00	3.41	0.23	26.54
time (sec)	N/A	0.339	0.028	0.547	0.195	0.082	0.000	0.114	0.199	3.416

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	96	66	124	53	75	94	296	23	2235
N.S.	1	1.20	0.82	1.55	0.66	0.94	1.18	3.70	0.29	27.94
time (sec)	N/A	0.280	0.026	0.345	0.187	0.077	76.158	0.117	0.196	3.348

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	67	49	69	42	64	71	248	23	1789
N.S.	1	1.14	0.83	1.17	0.71	1.08	1.20	4.20	0.39	30.32
time (sec)	N/A	0.228	0.021	0.262	0.197	0.089	40.758	0.116	0.193	3.245

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	32	42	31	52	49	197	36	773
N.S.	1	1.00	0.84	1.11	0.82	1.37	1.29	5.18	0.95	20.34
time (sec)	N/A	0.188	0.174	0.241	0.185	0.086	23.007	0.111	0.177	3.224

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	12	39	26	136	22	337
N.S.	1	1.00	1.00	0.83	0.67	2.17	1.44	7.56	1.22	18.72
time (sec)	N/A	0.147	0.007	0.242	0.183	0.074	12.155	0.113	0.169	3.318

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	125	99	222	0	111	0	73	23	789
N.S.	1	1.03	0.82	1.83	0.00	0.92	0.00	0.60	0.19	6.52
time (sec)	N/A	0.322	0.094	0.197	0.000	0.091	0.000	0.113	0.174	7.137

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	116	106	193	0	123	0	80	23	616
N.S.	1	1.05	0.96	1.75	0.00	1.12	0.00	0.73	0.21	5.60
time (sec)	N/A	0.322	0.041	0.209	0.000	0.090	0.000	0.124	0.200	7.472



Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	109	108	142	0	130	0	92	23	614
N.S.	1	0.99	0.98	1.29	0.00	1.18	0.00	0.84	0.21	5.58
time (sec)	N/A	0.421	0.032	0.260	0.000	0.090	0.000	0.121	0.180	4.321

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	114	107	144	0	143	0	76	91	669
N.S.	1	1.01	0.95	1.27	0.00	1.27	0.00	0.67	0.81	5.92
time (sec)	N/A	0.436	0.049	0.342	0.000	0.087	0.000	0.120	0.207	7.743

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	172	134	169	0	164	0	108	91	1069
N.S.	1	1.03	0.80	1.01	0.00	0.98	0.00	0.65	0.54	6.40
time (sec)	N/A	0.642	0.076	0.626	0.000	0.080	0.000	0.119	0.205	7.845

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	221	244	150	262	0	186	0	105	91	1292
N.S.	1	1.10	0.68	1.19	0.00	0.84	0.00	0.48	0.41	5.85
time (sec)	N/A	0.724	0.082	1.373	0.000	0.091	0.000	0.128	0.194	8.655

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	119	83	153	64	53	0	61	23	496
N.S.	1	1.20	0.84	1.55	0.65	0.54	0.00	0.62	0.23	5.01
time (sec)	N/A	0.343	0.027	0.204	0.193	0.071	0.000	0.111	0.176	3.122

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	92	66	123	53	42	0	49	23	385
N.S.	1	1.21	0.87	1.62	0.70	0.55	0.00	0.64	0.30	5.07
time (sec)	N/A	0.294	0.027	0.208	0.189	0.084	0.000	0.114	0.170	3.010

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	63	49	68	42	31	0	37	23	211
N.S.	1	1.11	0.86	1.19	0.74	0.54	0.00	0.65	0.40	3.70
time (sec)	N/A	0.239	0.023	0.233	0.190	0.078	0.000	0.121	0.169	3.045

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	32	56	30	19	0	23	27	105
N.S.	1	1.00	0.89	1.56	0.83	0.53	0.00	0.64	0.75	2.92
time (sec)	N/A	0.187	0.108	0.203	0.190	0.071	0.000	0.120	0.186	3.112

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	12	12	24	12	13	52
N.S.	1	1.00	1.00	0.94	0.75	0.75	1.50	0.75	0.81	3.25
time (sec)	N/A	0.148	0.007	0.294	0.168	0.070	0.286	0.118	0.177	3.124

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	47	42	0	53	0	21	23	285
N.S.	1	1.00	0.96	0.86	0.00	1.08	0.00	0.43	0.47	5.82
time (sec)	N/A	0.181	0.087	0.282	0.000	0.089	0.000	0.110	0.180	9.158

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	100	78	95	0	90	0	44	23	570
N.S.	1	1.06	0.83	1.01	0.00	0.96	0.00	0.47	0.24	6.06
time (sec)	N/A	0.270	0.042	0.204	0.000	0.096	0.000	0.118	0.180	8.706

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	172	98	148	0	120	0	69	23	802
N.S.	1	1.09	0.62	0.94	0.00	0.76	0.00	0.44	0.15	5.08
time (sec)	N/A	0.398	0.057	0.217	0.000	0.092	0.000	0.116	0.186	7.993

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	212	244	117	200	0	142	0	72	23	1086
N.S.	1	1.15	0.55	0.94	0.00	0.67	0.00	0.34	0.11	5.12
time (sec)	N/A	0.594	0.075	0.203	0.000	0.099	0.000	0.118	0.176	7.744

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	117	83	319	64	63	0	82	23	1057
N.S.	1	1.23	0.87	3.36	0.67	0.66	0.00	0.86	0.24	11.13
time (sec)	N/A	0.586	0.037	0.198	0.186	0.080	0.000	0.115	0.187	3.700

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	88	66	201	52	51	0	61	23	660
N.S.	1	1.19	0.89	2.72	0.70	0.69	0.00	0.82	0.31	8.92
time (sec)	N/A	0.436	0.029	0.201	0.191	0.072	0.000	0.122	0.188	3.374

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	61	49	106	41	40	0	51	23	259
N.S.	1	1.11	0.89	1.93	0.75	0.73	0.00	0.93	0.42	4.71
time (sec)	N/A	0.390	0.026	0.198	0.187	0.077	0.000	0.112	0.178	3.432

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	29	40	30	29	46	29	36	152
N.S.	1	1.00	0.85	1.18	0.88	0.85	1.35	0.85	1.06	4.47
time (sec)	N/A	0.279	0.108	0.190	0.190	0.076	0.549	0.113	0.176	3.465

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	12	20	26	12	22	97
N.S.	1	1.00	1.00	0.94	0.75	1.25	1.62	0.75	1.38	6.06
time (sec)	N/A	0.145	0.007	0.243	0.170	0.073	0.536	0.118	0.174	3.437

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	75	68	0	107	0	37	23	614
N.S.	1	1.00	0.96	0.87	0.00	1.37	0.00	0.47	0.29	7.87
time (sec)	N/A	0.272	0.123	0.273	0.000	0.084	0.000	0.113	0.172	8.325

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	148	91	105	0	148	0	64	23	807
N.S.	1	1.19	0.73	0.85	0.00	1.19	0.00	0.52	0.19	6.51
time (sec)	N/A	0.345	0.056	0.256	0.000	0.089	0.000	0.114	0.181	8.904

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	191	220	115	131	0	186	0	80	23	1028
N.S.	1	1.15	0.60	0.69	0.00	0.97	0.00	0.42	0.12	5.38
time (sec)	N/A	0.502	0.080	0.209	0.000	0.099	0.000	0.123	0.186	8.624

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	245	292	133	186	0	208	0	95	23	1258
N.S.	1	1.19	0.54	0.76	0.00	0.85	0.00	0.39	0.09	5.13
time (sec)	N/A	0.648	0.079	0.213	0.000	0.097	0.000	0.116	0.187	7.348

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	117	83	295	64	74	0	79	23	817
N.S.	1	1.18	0.84	2.98	0.65	0.75	0.00	0.80	0.23	8.25
time (sec)	N/A	0.358	0.033	0.207	0.204	0.080	0.000	0.118	0.181	3.514

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	88	65	186	52	62	90	59	23	533
N.S.	1	1.16	0.86	2.45	0.68	0.82	1.18	0.78	0.30	7.01
time (sec)	N/A	0.284	0.030	0.201	0.188	0.081	2.414	0.110	0.173	3.254

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	63	48	91	42	52	71	43	23	259
N.S.	1	1.07	0.81	1.54	0.71	0.88	1.20	0.73	0.39	4.39
time (sec)	N/A	0.333	0.026	0.202	0.190	0.074	2.385	0.117	0.177	3.300

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	31	42	31	41	51	20	36	152
N.S.	1	1.00	0.82	1.11	0.82	1.08	1.34	0.53	0.95	4.00
time (sec)	N/A	0.313	0.115	0.191	0.191	0.081	2.305	0.117	0.173	3.344

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	15	12	31	27	12	22	103
N.S.	1	1.00	1.00	0.83	0.67	1.72	1.50	0.67	1.22	5.72
time (sec)	N/A	0.213	0.007	0.241	0.167	0.072	2.187	0.115	0.186	3.363

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	126	91	93	0	174	0	45	23	886
N.S.	1	1.17	0.84	0.86	0.00	1.61	0.00	0.42	0.21	8.20
time (sec)	N/A	0.505	0.166	0.202	0.000	0.081	0.000	0.121	0.184	8.055

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	196	113	130	0	218	0	65	23	1230
N.S.	1	1.26	0.73	0.84	0.00	1.41	0.00	0.42	0.15	7.94
time (sec)	N/A	0.691	0.086	0.213	0.000	0.087	0.000	0.124	0.202	8.823

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	224	268	133	157	0	252	0	93	23	1514
N.S.	1	1.20	0.59	0.70	0.00	1.12	0.00	0.42	0.10	6.76
time (sec)	N/A	0.639	0.075	0.216	0.000	0.088	0.000	0.116	0.190	9.220

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	340	150	211	0	274	0	115	23	2359
N.S.	1	1.22	0.54	0.76	0.00	0.99	0.00	0.41	0.08	8.49
time (sec)	N/A	0.759	0.130	0.208	0.000	0.097	0.000	0.116	0.174	7.920

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	23	20	19	18	0	13	15	57
N.S.	1	1.00	0.85	0.74	0.70	0.67	0.00	0.48	0.56	2.11
time (sec)	N/A	0.150	0.026	0.260	0.035	0.077	0.000	0.113	0.181	3.447



Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	23	20	19	18	26	13	15	57
N.S.	1	1.00	0.85	0.74	0.70	0.67	0.96	0.48	0.56	2.11
time (sec)	N/A	0.150	0.020	0.257	0.036	0.076	11.963	0.107	0.210	3.291

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	23	20	19	18	26	13	13	57
N.S.	1	1.00	0.85	0.74	0.70	0.67	0.96	0.48	0.48	2.11
time (sec)	N/A	0.151	0.020	0.260	0.037	0.072	1.098	0.106	0.216	3.366

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	23	20	19	16	0	13	12	57
N.S.	1	1.00	0.85	0.74	0.70	0.59	0.00	0.48	0.44	2.11
time (sec)	N/A	0.152	0.020	0.269	0.035	0.068	0.000	0.114	0.170	3.326

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	23	20	19	12	0	13	14	56
N.S.	1	1.00	0.92	0.80	0.76	0.48	0.00	0.52	0.56	2.24
time (sec)	N/A	0.155	0.016	0.283	0.035	0.074	0.000	0.110	0.184	3.365

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	20	19	12	22	13	20	56
N.S.	1	1.00	0.87	0.87	0.83	0.52	0.96	0.57	0.87	2.43
time (sec)	N/A	0.148	0.016	0.262	0.035	0.078	0.276	0.113	0.178	3.348

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	21	20	19	11	27	11	23	52
N.S.	1	1.00	0.78	0.74	0.70	0.41	1.00	0.41	0.85	1.93
time (sec)	N/A	0.150	0.016	0.342	0.038	0.076	1.290	0.113	0.179	3.426

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	23	20	19	13	27	13	23	57
N.S.	1	1.00	0.85	0.74	0.70	0.48	1.00	0.48	0.85	2.11
time (sec)	N/A	0.148	0.016	0.306	0.037	0.088	13.671	0.114	0.178	3.311

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	51	40	38	36	29	0	24	17	122
N.S.	1	1.06	0.83	0.79	0.75	0.60	0.00	0.50	0.35	2.54
time (sec)	N/A	0.197	0.047	0.467	0.043	0.087	0.000	0.112	0.181	3.294

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	51	40	38	36	29	48	24	17	122
N.S.	1	1.06	0.83	0.79	0.75	0.60	1.00	0.50	0.35	2.54
time (sec)	N/A	0.329	0.031	0.412	0.044	0.085	22.354	0.116	0.189	3.217

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	51	40	38	36	29	0	24	15	122
N.S.	1	1.06	0.83	0.79	0.75	0.60	0.00	0.50	0.31	2.54
time (sec)	N/A	0.320	0.030	0.399	0.043	0.087	0.000	0.117	0.188	3.162

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	51	40	38	36	27	0	24	14	122
N.S.	1	1.06	0.83	0.79	0.75	0.56	0.00	0.50	0.29	2.54
time (sec)	N/A	0.337	0.040	0.392	0.049	0.085	0.000	0.116	0.176	3.166

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	47	40	47	36	24	0	24	16	122
N.S.	1	1.02	0.87	1.02	0.78	0.52	0.00	0.52	0.35	2.65
time (sec)	N/A	0.308	0.022	0.362	0.043	0.068	0.000	0.114	0.185	3.094

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	45	40	37	36	23	0	24	19	122
N.S.	1	1.02	0.91	0.84	0.82	0.52	0.00	0.55	0.43	2.77
time (sec)	N/A	0.320	0.033	0.394	0.043	0.077	0.000	0.112	0.177	3.063

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	47	40	38	36	24	48	23	40	122
N.S.	1	0.98	0.83	0.79	0.75	0.50	1.00	0.48	0.83	2.54
time (sec)	N/A	0.332	0.034	0.401	0.044	0.085	1.288	0.109	0.192	3.118

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	51	40	38	36	24	49	24	40	122
N.S.	1	1.06	0.83	0.79	0.75	0.50	1.02	0.50	0.83	2.54
time (sec)	N/A	0.326	0.032	0.402	0.049	0.071	13.091	0.113	0.255	3.119

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	75	57	56	55	40	0	35	17	182
N.S.	1	1.09	0.83	0.81	0.80	0.58	0.00	0.51	0.25	2.64
time (sec)	N/A	0.235	0.030	1.052	0.059	0.074	0.000	0.112	0.230	3.134

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	75	57	56	55	40	70	35	17	182
N.S.	1	1.09	0.83	0.81	0.80	0.58	1.01	0.51	0.25	2.64
time (sec)	N/A	0.242	0.034	1.047	0.057	0.070	41.579	0.109	0.229	3.130

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	75	57	56	55	40	0	35	15	182
N.S.	1	1.09	0.83	0.81	0.80	0.58	0.00	0.51	0.22	2.64
time (sec)	N/A	0.237	0.023	0.970	0.057	0.081	0.000	0.110	0.182	3.144

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	73	57	56	55	38	0	35	14	182
N.S.	1	1.06	0.83	0.81	0.80	0.55	0.00	0.51	0.20	2.64
time (sec)	N/A	0.239	0.023	0.979	0.058	0.066	0.000	0.113	0.225	3.136

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	71	57	69	55	35	0	35	16	182
N.S.	1	1.09	0.88	1.06	0.85	0.54	0.00	0.54	0.25	2.80
time (sec)	N/A	0.238	0.022	0.994	0.055	0.091	0.000	0.115	0.162	3.120

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	67	57	64	55	34	0	35	19	182
N.S.	1	1.06	0.90	1.02	0.87	0.54	0.00	0.56	0.30	2.89
time (sec)	N/A	0.238	0.025	1.062	0.056	0.074	0.000	0.115	0.169	3.145

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	67	55	55	55	34	66	34	19	182
N.S.	1	1.03	0.85	0.85	0.85	0.52	1.02	0.52	0.29	2.80
time (sec)	N/A	0.241	0.022	0.987	0.056	0.077	1.285	0.112	0.170	3.157

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	71	57	56	55	35	70	34	57	182
N.S.	1	1.03	0.83	0.81	0.80	0.51	1.01	0.49	0.83	2.64
time (sec)	N/A	0.248	0.025	0.968	0.059	0.078	12.814	0.114	0.166	3.164

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	154	129	262	65	153	0	70	17	475
N.S.	1	1.08	0.90	1.83	0.45	1.07	0.00	0.49	0.12	3.32
time (sec)	N/A	0.413	0.052	0.351	0.121	0.076	0.000	0.112	0.158	3.720

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	124	108	201	54	132	0	59	17	415
N.S.	1	1.07	0.93	1.73	0.47	1.14	0.00	0.51	0.15	3.58
time (sec)	N/A	0.332	0.075	0.266	0.117	0.087	0.000	0.112	0.170	3.345

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	94	86	120	42	103	0	45	15	354
N.S.	1	1.06	0.97	1.35	0.47	1.16	0.00	0.51	0.17	3.98
time (sec)	N/A	0.278	0.059	0.253	0.118	0.078	0.000	0.112	0.167	3.847

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	62	66	31	85	0	31	14	296
N.S.	1	1.00	0.97	1.03	0.48	1.33	0.00	0.48	0.22	4.62
time (sec)	N/A	0.358	0.031	0.254	0.120	0.073	0.000	0.118	0.160	4.089

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	51	41	18	68	0	18	16	347
N.S.	1	1.00	0.96	0.77	0.34	1.28	0.00	0.34	0.30	6.55
time (sec)	N/A	0.287	0.021	0.257	0.116	0.083	0.000	0.111	0.153	5.913

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	73	76	31	87	0	31	19	464
N.S.	1	1.00	0.96	1.00	0.41	1.14	0.00	0.41	0.25	6.11
time (sec)	N/A	0.372	0.042	0.261	0.119	0.094	0.000	0.111	0.166	3.952

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	118	89	98	41	113	0	41	19	642
N.S.	1	1.17	0.88	0.97	0.41	1.12	0.00	0.41	0.19	6.36
time (sec)	N/A	0.488	0.113	0.251	0.116	0.090	0.000	0.110	0.170	3.863

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	160	107	120	52	139	0	52	19	822
N.S.	1	1.25	0.84	0.94	0.41	1.09	0.00	0.41	0.15	6.42
time (sec)	N/A	0.616	0.103	0.264	0.122	0.093	0.000	0.111	0.176	3.569

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	151	144	280	75	188	0	76	17	523
N.S.	1	1.12	1.07	2.07	0.56	1.39	0.00	0.56	0.13	3.87
time (sec)	N/A	0.434	0.147	1.642	0.124	0.101	0.000	0.112	0.170	3.513



Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	121	119	181	64	161	0	65	17	463
N.S.	1	1.12	1.10	1.68	0.59	1.49	0.00	0.60	0.16	4.29
time (sec)	N/A	0.317	0.099	1.276	0.129	0.090	0.000	0.118	0.167	3.464

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	91	81	95	50	134	0	46	15	403
N.S.	1	1.10	0.98	1.14	0.60	1.61	0.00	0.55	0.18	4.86
time (sec)	N/A	0.258	0.048	1.154	0.119	0.089	0.000	0.124	0.166	3.896

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	70	61	37	115	0	36	43	344
N.S.	1	1.00	0.96	0.84	0.51	1.58	0.00	0.49	0.59	4.71
time (sec)	N/A	0.213	0.041	1.303	0.123	0.086	0.000	0.117	0.182	3.933

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	103	80	82	35	116	0	35	16	516
N.S.	1	1.06	0.82	0.85	0.36	1.20	0.00	0.36	0.16	5.32
time (sec)	N/A	0.270	0.043	1.499	0.120	0.085	0.000	0.109	0.187	4.302

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	145	104	93	51	143	0	49	19	705
N.S.	1	1.21	0.87	0.78	0.42	1.19	0.00	0.41	0.16	5.88
time (sec)	N/A	0.342	0.080	1.311	0.122	0.092	0.000	0.113	0.211	4.018

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	187	120	115	64	179	0	58	19	871
N.S.	1	1.29	0.83	0.79	0.44	1.23	0.00	0.40	0.13	6.01
time (sec)	N/A	0.423	0.137	1.410	0.120	0.090	0.000	0.115	0.188	4.287

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	172	229	139	137	75	205	0	70	19	1051
N.S.	1	1.33	0.81	0.80	0.44	1.19	0.00	0.41	0.11	6.11
time (sec)	N/A	0.500	0.165	1.548	0.122	0.084	0.000	0.118	0.189	4.286

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	150	147	255	86	227	0	77	17	571
N.S.	1	1.11	1.09	1.89	0.64	1.68	0.00	0.57	0.13	4.23
time (sec)	N/A	0.617	0.078	1.351	0.130	0.083	0.000	0.114	0.173	4.191

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	120	104	153	73	200	0	59	17	511
N.S.	1	1.09	0.95	1.39	0.66	1.82	0.00	0.54	0.15	4.65
time (sec)	N/A	0.447	0.067	1.242	0.123	0.087	0.000	0.114	0.170	4.087

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	102	96	85	61	185	0	47	59	667
N.S.	1	1.04	0.98	0.87	0.62	1.89	0.00	0.48	0.60	6.81
time (sec)	N/A	0.437	0.053	1.180	0.125	0.090	0.000	0.118	0.180	4.108

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	132	107	98	64	186	0	52	45	580
N.S.	1	1.06	0.86	0.78	0.51	1.49	0.00	0.42	0.36	4.64
time (sec)	N/A	0.471	0.100	1.230	0.125	0.086	0.000	0.113	0.212	4.038

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	174	118	112	60	186	0	47	16	741
N.S.	1	1.14	0.78	0.74	0.39	1.22	0.00	0.31	0.11	4.88
time (sec)	N/A	0.446	0.055	1.316	0.125	0.080	0.000	0.113	0.185	4.093

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	176	216	141	114	73	209	0	59	19	1077
N.S.	1	1.23	0.80	0.65	0.41	1.19	0.00	0.34	0.11	6.12
time (sec)	N/A	0.507	0.120	1.383	0.130	0.087	0.000	0.112	0.183	4.136

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	258	156	136	86	245	0	71	19	1362
N.S.	1	1.28	0.78	0.68	0.43	1.22	0.00	0.35	0.09	6.78
time (sec)	N/A	0.584	0.180	1.586	0.128	0.085	0.000	0.111	0.185	4.699

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	228	300	174	158	97	271	0	80	19	2151
N.S.	1	1.32	0.76	0.69	0.43	1.19	0.00	0.35	0.08	9.43
time (sec)	N/A	0.677	0.218	1.894	0.132	0.098	0.000	0.111	0.192	4.071

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	145	104	121	0	138	0	60	14	0
N.S.	1	1.02	0.73	0.85	0.00	0.97	0.00	0.42	0.10	0.00
time (sec)	N/A	0.333	0.059	0.405	0.000	0.090	0.000	0.116	0.192	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	101	84	86	0	111	0	48	13	0
N.S.	1	0.97	0.81	0.83	0.00	1.07	0.00	0.46	0.12	0.00
time (sec)	N/A	0.307	0.053	0.391	0.000	0.080	0.000	0.120	0.185	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	62	49	0	90	0	36	16	0
N.S.	1	1.00	1.02	0.80	0.00	1.48	0.00	0.59	0.26	0.00
time (sec)	N/A	0.355	0.030	0.407	0.000	0.092	0.000	0.119	0.240	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	52	97	40	86	0	57	45	0
N.S.	1	1.00	1.06	1.98	0.82	1.76	0.00	1.16	0.92	0.00
time (sec)	N/A	0.329	0.032	0.402	0.151	0.074	0.000	0.123	0.241	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	34	29	15	15	0	59	50	210
N.S.	1	1.00	0.97	0.83	0.43	0.43	0.00	1.69	1.43	6.00
time (sec)	N/A	0.279	0.027	0.389	0.125	0.075	0.000	0.117	0.242	3.969

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	48	59	34	34	0	112	52	174
N.S.	1	1.00	0.67	0.82	0.47	0.47	0.00	1.56	0.72	2.42
time (sec)	N/A	0.313	0.029	0.418	0.119	0.078	0.000	0.120	0.242	3.593

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	128	66	105	45	45	0	138	52	234
N.S.	1	1.16	0.60	0.95	0.41	0.41	0.00	1.25	0.47	2.13
time (sec)	N/A	0.504	0.049	0.422	0.121	0.075	0.000	0.123	0.242	3.836

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	184	82	151	56	56	0	166	52	294
N.S.	1	1.24	0.55	1.02	0.38	0.38	0.00	1.12	0.35	1.99
time (sec)	N/A	0.398	0.049	0.417	0.118	0.089	0.000	0.125	0.247	3.745

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	177	183	122	150	0	160	0	147	21	0
N.S.	1	1.03	0.69	0.85	0.00	0.90	0.00	0.83	0.12	0.00
time (sec)	N/A	0.417	0.066	0.409	0.000	0.089	0.000	0.141	0.266	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	139	105	115	0	137	0	122	20	0
N.S.	1	1.00	0.76	0.83	0.00	0.99	0.00	0.88	0.14	0.00
time (sec)	N/A	0.333	0.055	0.381	0.000	0.089	0.000	0.132	0.256	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	99	83	80	0	116	0	97	23	0
N.S.	1	0.98	0.82	0.79	0.00	1.15	0.00	0.96	0.23	0.00
time (sec)	N/A	0.270	0.045	0.399	0.000	0.096	0.000	72.365	0.205	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	83	77	126	0	106	0	90	44	0
N.S.	1	1.02	0.95	1.56	0.00	1.31	0.00	1.11	0.54	0.00
time (sec)	N/A	0.254	0.039	0.389	0.000	0.086	0.000	72.697	0.259	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	72	74	172	0	106	0	91	72	0
N.S.	1	1.03	1.06	2.46	0.00	1.51	0.00	1.30	1.03	0.00
time (sec)	N/A	0.245	0.041	0.405	0.000	0.103	0.000	73.162	0.255	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	34	29	15	31	0	33	73	332
N.S.	1	1.00	0.97	0.83	0.43	0.89	0.00	0.94	2.09	9.49
time (sec)	N/A	0.171	0.032	0.396	0.124	0.079	0.000	0.133	0.245	3.797

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	48	59	34	45	0	61	74	228
N.S.	1	1.00	0.67	0.82	0.47	0.62	0.00	0.85	1.03	3.17
time (sec)	N/A	0.230	0.045	0.434	0.120	0.072	0.000	0.138	0.252	3.769

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	128	66	105	45	56	0	78	74	288
N.S.	1	1.16	0.60	0.95	0.41	0.51	0.00	0.71	0.67	2.62
time (sec)	N/A	0.301	0.038	0.399	0.125	0.092	0.000	0.137	0.249	3.944

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	184	82	151	56	67	0	99	73	348
N.S.	1	1.24	0.55	1.02	0.38	0.45	0.00	0.67	0.49	2.35
time (sec)	N/A	0.605	0.056	0.425	0.119	0.079	0.000	0.132	0.247	4.157



Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	177	121	144	0	159	0	204	22	0
N.S.	1	1.02	0.70	0.83	0.00	0.91	0.00	1.17	0.13	0.00
time (sec)	N/A	0.618	0.061	0.397	0.000	0.104	0.000	0.151	0.309	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	136	137	101	109	0	138	0	116	25	0
N.S.	1	1.01	0.74	0.80	0.00	1.01	0.00	0.85	0.18	0.00
time (sec)	N/A	0.392	0.042	0.384	0.000	0.085	0.000	73.066	0.221	0.000

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	101	155	0	134	0	108	25	0
N.S.	1	1.00	0.83	1.28	0.00	1.11	0.00	0.89	0.21	0.00
time (sec)	N/A	0.315	0.052	0.446	0.000	0.085	0.000	73.411	0.195	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	109	97	201	0	135	0	109	72	0
N.S.	1	1.03	0.92	1.90	0.00	1.27	0.00	1.03	0.68	0.00
time (sec)	N/A	0.303	0.054	0.385	0.000	0.099	0.000	73.119	0.274	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F(-1)</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	95	95	247	0	134	0	111	99	0
N.S.	1	1.02	1.02	2.66	0.00	1.44	0.00	1.19	1.06	0.00
time (sec)	N/A	0.298	0.039	0.384	0.000	0.083	0.000	73.144	0.257	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	34	29	15	42	0	33	98	396
N.S.	1	1.00	0.97	0.83	0.43	1.20	0.00	0.94	2.80	11.31
time (sec)	N/A	0.164	0.045	0.385	0.121	0.075	0.000	0.142	0.256	3.814

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	48	59	34	56	0	59	99	293
N.S.	1	1.00	0.67	0.82	0.47	0.78	0.00	0.82	1.38	4.07
time (sec)	N/A	0.224	0.038	0.422	0.122	0.088	0.000	0.144	0.260	3.794

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	128	66	105	45	67	0	80	99	353
N.S.	1	1.16	0.60	0.95	0.41	0.61	0.00	0.73	0.90	3.21
time (sec)	N/A	0.290	0.040	0.436	0.120	0.080	0.000	0.138	0.252	3.801

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	184	82	151	56	78	0	97	99	413
N.S.	1	1.24	0.55	1.02	0.38	0.53	0.00	0.66	0.67	2.79
time (sec)	N/A	0.397	0.059	0.503	0.118	0.080	0.000	0.148	0.247	3.930

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	151	105	124	0	137	0	64	25	0
N.S.	1	1.04	0.72	0.86	0.00	0.94	0.00	0.44	0.17	0.00
time (sec)	N/A	0.452	0.064	0.414	0.000	0.085	0.000	0.118	0.184	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	88	89	0	116	0	52	23	0
N.S.	1	1.00	0.82	0.83	0.00	1.08	0.00	0.49	0.21	0.00
time (sec)	N/A	0.469	0.059	0.401	0.000	0.083	0.000	0.118	0.184	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	66	53	0	88	0	38	22	0
N.S.	1	1.00	1.05	0.84	0.00	1.40	0.00	0.60	0.35	0.00
time (sec)	N/A	0.357	0.040	0.440	0.000	0.082	0.000	0.123	0.225	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	33	24	0	54	0	23	25	0
N.S.	1	1.00	1.10	0.80	0.00	1.80	0.00	0.77	0.83	0.00
time (sec)	N/A	0.267	0.025	0.412	0.000	0.085	0.000	0.126	0.191	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	29	15	15	0	30	25	101
N.S.	1	1.00	0.97	0.88	0.45	0.45	0.00	0.91	0.76	3.06
time (sec)	N/A	0.276	0.032	0.415	0.127	0.078	0.000	0.112	0.188	3.736

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	46	59	33	23	0	55	25	218
N.S.	1	1.00	0.64	0.82	0.46	0.32	0.00	0.76	0.35	3.03
time (sec)	N/A	0.370	0.040	0.402	0.129	0.087	0.000	0.121	0.186	3.817

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	128	66	105	45	34	0	77	25	227
N.S.	1	1.16	0.60	0.95	0.41	0.31	0.00	0.70	0.23	2.06
time (sec)	N/A	0.296	0.036	0.415	0.121	0.083	0.000	0.125	0.185	3.590

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	184	82	151	55	45	0	103	25	287
N.S.	1	1.24	0.55	1.02	0.37	0.30	0.00	0.70	0.17	1.94
time (sec)	N/A	0.409	0.051	0.441	0.118	0.080	0.000	0.121	0.194	3.537

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F(-1)	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	178	122	146	0	193	0	75	25	0
N.S.	1	1.07	0.73	0.88	0.00	1.16	0.00	0.45	0.15	0.00
time (sec)	N/A	0.411	0.079	0.402	0.000	0.088	0.000	0.126	0.199	0.000

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	134	104	111	0	172	0	63	25	0
N.S.	1	1.05	0.81	0.87	0.00	1.34	0.00	0.49	0.20	0.00
time (sec)	N/A	0.324	0.074	0.392	0.000	0.088	0.000	0.125	0.192	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	90	81	75	0	142	0	48	58	0
N.S.	1	1.05	0.94	0.87	0.00	1.65	0.00	0.56	0.67	0.00
time (sec)	N/A	0.257	0.052	0.392	0.000	0.093	0.000	0.120	0.260	0.000

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	55	42	0	116	0	39	59	0
N.S.	1	1.00	1.06	0.81	0.00	2.23	0.00	0.75	1.13	0.00
time (sec)	N/A	0.204	0.038	0.391	0.000	0.090	0.000	0.117	0.271	0.000

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	29	0	22	0	15	25	163
N.S.	1	1.00	0.97	0.88	0.00	0.67	0.00	0.45	0.76	4.94
time (sec)	N/A	0.166	0.025	0.387	0.000	0.087	0.000	0.112	0.216	3.680

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	43	59	32	34	0	50	25	281
N.S.	1	1.00	0.63	0.87	0.47	0.50	0.00	0.74	0.37	4.13
time (sec)	N/A	0.222	0.037	0.399	0.127	0.087	0.000	0.120	0.215	3.443

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	124	64	105	45	49	0	107	25	286
N.S.	1	1.13	0.58	0.95	0.41	0.45	0.00	0.97	0.23	2.60
time (sec)	N/A	0.304	0.044	0.400	0.135	0.085	0.000	0.131	0.219	3.707

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	148	180	80	151	54	58	0	161	25	346
N.S.	1	1.22	0.54	1.02	0.36	0.39	0.00	1.09	0.17	2.34
time (sec)	N/A	0.566	0.051	0.384	0.120	0.078	0.000	0.142	0.210	3.746

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	153	165	121	133	0	238	0	75	25	0
N.S.	1	1.08	0.79	0.87	0.00	1.56	0.00	0.49	0.16	0.00
time (sec)	N/A	0.539	0.082	0.396	0.000	0.094	0.000	0.126	0.215	0.000

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	121	101	97	0	211	0	61	25	0
N.S.	1	1.09	0.91	0.87	0.00	1.90	0.00	0.55	0.23	0.00
time (sec)	N/A	0.536	0.071	0.386	0.000	0.111	0.000	0.124	0.222	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	80	78	59	0	183	0	49	84	0
N.S.	1	1.07	1.04	0.79	0.00	2.44	0.00	0.65	1.12	0.00
time (sec)	N/A	0.388	0.045	0.420	0.000	0.093	0.000	0.126	0.268	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	34	92	0	33	0	15	62	229
N.S.	1	1.00	0.97	2.63	0.00	0.94	0.00	0.43	1.77	6.54
time (sec)	N/A	0.161	0.038	0.401	0.000	0.095	0.000	0.122	0.279	3.827

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	47	58	0	43	0	25	25	346
N.S.	1	1.00	0.66	0.82	0.00	0.61	0.00	0.35	0.35	4.87
time (sec)	N/A	0.244	0.031	0.391	0.000	0.082	0.000	0.118	0.202	3.852

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	123	66	104	45	58	0	62	25	348
N.S.	1	1.16	0.62	0.98	0.42	0.55	0.00	0.58	0.24	3.28
time (sec)	N/A	0.292	0.042	0.401	0.124	0.093	0.000	0.129	0.207	3.788

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	177	79	150	56	71	0	119	25	406
N.S.	1	1.21	0.54	1.03	0.38	0.49	0.00	0.82	0.17	2.78
time (sec)	N/A	0.384	0.050	0.392	0.119	0.087	0.000	0.142	0.203	3.931



Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	186	233	100	196	67	82	0	173	25	466
N.S.	1	1.25	0.54	1.05	0.36	0.44	0.00	0.93	0.13	2.51
time (sec)	N/A	0.481	0.054	0.505	0.122	0.091	0.000	0.151	0.201	4.058

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	71	0	0	0	0	0	15	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.210	0.114	0.000	0.000	0.000	0.000	0.000	0.183	0.000

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	150	146	406	139	374	0	332	15	546
N.S.	1	0.91	0.88	2.46	0.84	2.27	0.00	2.01	0.09	3.31
time (sec)	N/A	0.424	0.074	3.497	0.195	0.081	0.000	0.118	0.173	3.864

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	116	106	277	101	255	0	226	15	418
N.S.	1	0.96	0.88	2.29	0.83	2.11	0.00	1.87	0.12	3.45
time (sec)	N/A	0.336	0.058	1.491	0.190	0.082	0.000	0.112	0.172	3.556

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	71	163	68	168	0	140	15	304
N.S.	1	1.00	0.87	1.99	0.83	2.05	0.00	1.71	0.18	3.71
time (sec)	N/A	0.272	0.044	0.803	0.186	0.082	0.000	0.113	0.181	3.520

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	41	77	42	91	0	76	48	205
N.S.	1	1.00	0.85	1.60	0.88	1.90	0.00	1.58	1.00	4.27
time (sec)	N/A	0.301	0.032	0.503	0.188	0.083	0.000	0.114	0.171	3.434

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	21	21	39	51	28	25	121
N.S.	1	1.00	1.00	1.05	1.05	1.95	2.55	1.40	1.25	6.05
time (sec)	N/A	0.261	0.013	0.458	0.175	0.085	0.281	0.127	0.175	3.363

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	60	0	0	0	0	0	15	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.256	0.090	0.000	0.000	0.000	0.000	0.000	0.180	0.000

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	67	0	0	0	0	0	44	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.62	0.00
time (sec)	N/A	0.395	0.037	0.000	0.000	0.000	0.000	0.000	0.219	0.000

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	101	98	67	0	0	0	0	0	47	0
N.S.	1	0.97	0.66	0.00	0.00	0.00	0.00	0.00	0.47	0.00
time (sec)	N/A	0.468	0.051	0.000	0.000	0.000	0.000	0.000	0.199	0.000

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	<b>F</b>	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	34	49	38	81	0	90	37	96
N.S.	1	1.00	0.92	1.32	1.03	2.19	0.00	2.43	1.00	2.59
time (sec)	N/A	0.287	0.039	0.242	0.036	0.100	0.000	0.124	0.215	3.943

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	19	19	19	76	71	13	19
N.S.	1	1.00	0.87	0.83	0.83	0.83	3.30	3.09	0.57	0.83
time (sec)	N/A	0.272	0.018	0.281	0.075	0.087	4.798	0.124	0.184	0.077

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	19	19	19	76	71	11	19
N.S.	1	1.00	0.87	0.83	0.83	0.83	3.30	3.09	0.48	0.83
time (sec)	N/A	0.294	0.012	0.185	0.077	0.082	2.893	0.120	0.171	3.503

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	15	16	14	68	69	14	16
N.S.	1	1.00	1.12	0.94	1.00	0.88	4.25	4.31	0.88	1.00
time (sec)	N/A	0.149	0.005	0.191	0.073	0.076	1.588	0.117	0.179	0.021

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	<b>F</b>	C	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	19	23	34	15	0	15	13	59
N.S.	1	1.00	0.90	1.10	1.62	0.71	0.00	0.71	0.62	2.81
time (sec)	N/A	0.206	0.013	0.164	0.033	0.074	0.000	0.130	0.193	0.179

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	18	18	17	18	68	70	19	17
N.S.	1	1.00	1.06	1.06	1.00	1.06	4.00	4.12	1.12	1.00
time (sec)	N/A	0.170	0.013	0.181	0.084	0.084	3.102	0.117	0.169	0.073

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	18	20	19	16	80	71	19	16
N.S.	1	1.00	0.78	0.87	0.83	0.70	3.48	3.09	0.83	0.70
time (sec)	N/A	0.161	0.012	0.177	0.077	0.069	5.343	0.132	0.173	0.061

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	40	39	36	33	58	0	0	5	0
N.S.	1	1.48	1.44	1.33	1.22	2.15	0.00	0.00	0.19	0.00
time (sec)	N/A	0.299	0.015	0.490	0.073	0.079	0.000	0.000	0.169	0.000

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	51	69	70	400	56	88	0	0	7	0
N.S.	1	1.35	1.37	7.84	1.10	1.73	0.00	0.00	0.14	0.00
time (sec)	N/A	0.421	0.012	0.398	0.075	0.091	0.000	0.000	0.173	0.000

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	77	99	91	422	78	118	0	0	9	0
N.S.	1	1.29	1.18	5.48	1.01	1.53	0.00	0.00	0.12	0.00
time (sec)	N/A	0.567	0.017	0.421	0.080	0.089	0.000	0.000	0.197	0.000

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	307	395	265	5316	281	900	0	0	17	0
N.S.	1	1.29	0.86	17.32	0.92	2.93	0.00	0.00	0.06	0.00
time (sec)	N/A	1.418	0.572	13.133	0.306	0.120	0.000	0.000	0.187	0.000

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	231	305	199	5012	215	746	0	0	15	0
N.S.	1	1.32	0.86	21.70	0.93	3.23	0.00	0.00	0.06	0.00
time (sec)	N/A	1.181	0.510	3.922	0.303	0.132	0.000	0.000	0.178	0.000

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	207	131	348	142	552	0	0	13	0
N.S.	1	1.38	0.87	2.32	0.95	3.68	0.00	0.00	0.09	0.00
time (sec)	N/A	1.059	3.343	3.828	0.299	0.125	0.000	0.000	0.182	0.000

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.468	9.748	0.379	0.749	0.082	0.954	0.517	0.182	3.640

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	155	199	148	1721	149	451	0	0	18	0
N.S.	1	1.28	0.95	11.10	0.96	2.91	0.00	0.00	0.12	0.00
time (sec)	N/A	1.336	0.138	3.169	0.591	0.097	0.000	0.000	0.170	0.000

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	128	165	122	1662	125	382	0	0	18	0
N.S.	1	1.29	0.95	12.98	0.98	2.98	0.00	0.00	0.14	0.00
time (sec)	N/A	0.817	0.078	2.698	0.568	0.099	0.000	0.000	0.173	0.000

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	101	131	91	1579	101	323	0	0	16	0
N.S.	1	1.30	0.90	15.63	1.00	3.20	0.00	0.00	0.16	0.00
time (sec)	N/A	0.656	0.078	2.483	0.593	0.095	0.000	0.000	0.178	0.000

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	90	63	255	72	239	0	0	14	0
N.S.	1	1.30	0.91	3.70	1.04	3.46	0.00	0.00	0.20	0.00
time (sec)	N/A	0.449	0.792	2.299	0.586	0.091	0.000	0.000	0.175	0.000

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12	1.12
time (sec)	N/A	0.243	3.553	0.289	0.634	0.089	0.616	0.174	0.180	3.330

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	168	218	148	1725	146	424	0	0	20	0
N.S.	1	1.30	0.88	10.27	0.87	2.52	0.00	0.00	0.12	0.00
time (sec)	N/A	1.092	0.142	3.296	0.580	0.087	0.000	0.000	0.183	0.000

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	139	182	123	1668	123	360	0	0	20	0
N.S.	1	1.31	0.88	12.00	0.88	2.59	0.00	0.00	0.14	0.00
time (sec)	N/A	0.867	0.085	3.184	0.622	0.100	0.000	0.000	0.182	0.000

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	110	146	93	1587	100	306	0	0	18	0
N.S.	1	1.33	0.85	14.43	0.91	2.78	0.00	0.00	0.16	0.00
time (sec)	N/A	0.678	0.087	2.731	0.606	0.093	0.000	0.000	0.191	0.000



Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	103	66	272	73	228	0	0	16	0
N.S.	1	1.36	0.87	3.58	0.96	3.00	0.00	0.00	0.21	0.00
time (sec)	N/A	0.464	0.786	2.559	0.595	0.094	0.000	0.000	0.195	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	17	20	19	17	19	20	19
N.S.	1	1.00	1.11	0.89	1.05	1.00	0.89	1.00	1.05	1.00
time (sec)	N/A	0.237	4.269	0.312	0.722	0.086	0.634	0.203	0.180	3.348

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	303	391	265	5248	277	880	0	0	17	0
N.S.	1	1.29	0.87	17.32	0.91	2.90	0.00	0.00	0.06	0.00
time (sec)	N/A	1.421	0.548	15.175	0.308	0.122	0.000	0.000	0.184	0.000

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	229	301	199	4944	213	730	0	0	15	0
N.S.	1	1.31	0.87	21.59	0.93	3.19	0.00	0.00	0.07	0.00
time (sec)	N/A	1.046	0.513	5.155	0.299	0.123	0.000	0.000	0.184	0.000

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	207	131	348	142	540	0	0	13	0
N.S.	1	1.38	0.87	2.32	0.95	3.60	0.00	0.00	0.09	0.00
time (sec)	N/A	0.646	3.114	3.962	0.304	0.120	0.000	0.000	0.169	0.000

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.420	9.914	0.349	0.761	0.095	2.250	0.514	0.184	3.383

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	152	195	145	1693	146	424	0	0	18	0
N.S.	1	1.28	0.95	11.14	0.96	2.79	0.00	0.00	0.12	0.00
time (sec)	N/A	1.503	0.139	3.672	0.578	0.110	0.000	0.000	0.174	0.000

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	126	162	120	1636	123	360	0	0	18	0
N.S.	1	1.29	0.95	12.98	0.98	2.86	0.00	0.00	0.14	0.00
time (sec)	N/A	1.277	0.088	3.221	0.584	0.111	0.000	0.000	0.194	0.000

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	100	128	90	1555	100	306	0	0	16	0
N.S.	1	1.28	0.90	15.55	1.00	3.06	0.00	0.00	0.16	0.00
time (sec)	N/A	0.714	0.081	2.760	0.572	0.105	0.000	0.000	0.179	0.000

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	89	63	255	72	227	0	0	14	0
N.S.	1	1.29	0.91	3.70	1.04	3.29	0.00	0.00	0.20	0.00
time (sec)	N/A	0.470	0.699	2.766	0.583	0.106	0.000	0.000	0.223	0.000

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	18	18	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	1.12	1.12	1.12
time (sec)	N/A	0.251	3.522	0.394	0.661	0.078	1.948	0.167	0.191	3.258

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	165	214	151	1753	149	451	0	0	20	0
N.S.	1	1.30	0.92	10.62	0.90	2.73	0.00	0.00	0.12	0.00
time (sec)	N/A	1.095	0.143	3.308	0.571	0.111	0.000	0.000	0.204	0.000

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	137	179	125	1694	125	382	0	0	20	0
N.S.	1	1.31	0.91	12.36	0.91	2.79	0.00	0.00	0.15	0.00
time (sec)	N/A	0.859	0.082	3.262	0.600	0.102	0.000	0.000	0.195	0.000

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	109	143	94	1611	101	323	0	0	18	0
N.S.	1	1.31	0.86	14.78	0.93	2.96	0.00	0.00	0.17	0.00
time (sec)	N/A	0.689	0.084	2.524	0.599	0.099	0.000	0.000	0.186	0.000

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	102	66	272	73	240	0	0	16	0
N.S.	1	1.34	0.87	3.58	0.96	3.16	0.00	0.00	0.21	0.00
time (sec)	N/A	0.463	0.669	2.573	0.565	0.129	0.000	0.000	0.173	0.000

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	17	20	19	17	19	20	19
N.S.	1	1.00	1.11	0.89	1.05	1.00	0.89	1.00	1.05	1.00
time (sec)	N/A	0.247	3.604	0.304	0.703	0.091	1.813	0.194	0.178	3.237

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	302	355	654	3640	0	1809	0	0	67	0
N.S.	1	1.18	2.17	12.05	0.00	5.99	0.00	0.00	0.22	0.00
time (sec)	N/A	1.137	0.977	16.670	0.000	0.153	0.000	0.000	0.202	0.000

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	234	271	409	2719	0	1283	0	0	46	0
N.S.	1	1.16	1.75	11.62	0.00	5.48	0.00	0.00	0.20	0.00
time (sec)	N/A	0.853	0.510	9.737	0.000	0.142	0.000	0.000	0.200	0.000

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	162	182	295	1818	0	835	0	0	25	0
N.S.	1	1.12	1.82	11.22	0.00	5.15	0.00	0.00	0.15	0.00
time (sec)	N/A	0.599	0.371	3.458	0.000	0.132	0.000	0.000	0.180	0.000

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	86	127	120	182	499	0	0	9	0
N.S.	1	1.09	1.61	1.52	2.30	6.32	0.00	0.00	0.11	0.00
time (sec)	N/A	0.368	0.020	1.771	0.144	0.125	0.000	0.000	0.174	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.219	6.964	0.368	1.283	0.092	0.602	0.409	0.179	3.349

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	395	522	349	6917	0	2165	0	0	17	0
N.S.	1	1.32	0.88	17.51	0.00	5.48	0.00	0.00	0.04	0.00
time (sec)	N/A	1.789	3.196	46.012	0.000	0.170	0.000	0.000	0.179	0.000

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	295	402	259	6525	0	1689	0	0	15	0
N.S.	1	1.36	0.88	22.12	0.00	5.73	0.00	0.00	0.05	0.00
time (sec)	N/A	1.830	2.587	5.254	0.000	0.172	0.000	0.000	0.178	0.000

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	287	365	557	372	1185	0	0	13	0
N.S.	1	1.48	1.88	2.87	1.92	6.11	0.00	0.00	0.07	0.00
time (sec)	N/A	0.725	3.726	3.424	0.200	0.153	0.000	0.000	0.176	0.000

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.308	2.046	0.315	3.113	0.104	0.825	0.718	0.165	3.972

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	170	218	155	2277	343	345	0	0	21	0
N.S.	1	1.28	0.91	13.39	2.02	2.03	0.00	0.00	0.12	0.00
time (sec)	N/A	0.890	0.268	4.358	0.063	0.104	0.000	0.000	0.172	0.000

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	133	171	119	2187	248	293	0	0	19	0
N.S.	1	1.29	0.89	16.44	1.86	2.20	0.00	0.00	0.14	0.00
time (sec)	N/A	0.699	0.166	3.319	0.053	0.096	0.000	0.000	0.181	0.000

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	123	766	307	260	218	0	0	17	0
N.S.	1	1.32	8.24	3.30	2.80	2.34	0.00	0.00	0.18	0.00
time (sec)	N/A	0.473	7.496	3.374	0.117	0.102	0.000	0.000	0.173	0.000

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	144	37	17	20	21	21
N.S.	1	1.00	1.10	0.90	7.20	1.85	0.85	1.00	1.05	1.05
time (sec)	N/A	0.259	0.814	0.494	3.796	0.087	0.800	0.460	0.175	3.605

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	171	223	156	2387	342	345	0	0	23	0
N.S.	1	1.30	0.91	13.96	2.00	2.02	0.00	0.00	0.13	0.00
time (sec)	N/A	0.929	0.233	4.478	0.059	0.105	0.000	0.000	0.188	0.000

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	134	176	120	2289	247	293	0	0	21	0
N.S.	1	1.31	0.90	17.08	1.84	2.19	0.00	0.00	0.16	0.00
time (sec)	N/A	0.692	0.159	3.814	0.050	0.106	0.000	0.000	0.181	0.000

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	128	723	332	262	219	0	0	19	0
N.S.	1	1.36	7.69	3.53	2.79	2.33	0.00	0.00	0.20	0.00
time (sec)	N/A	0.483	4.727	3.731	0.123	0.098	0.000	0.000	0.206	0.000



Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	143	37	17	21	23	22
N.S.	1	1.00	1.10	0.90	6.81	1.76	0.81	1.00	1.10	1.05
time (sec)	N/A	0.261	0.787	0.493	3.956	0.094	0.867	0.476	0.187	3.579

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	302	355	654	3640	0	1567	0	0	67	0
N.S.	1	1.18	2.17	12.05	0.00	5.19	0.00	0.00	0.22	0.00
time (sec)	N/A	1.708	0.192	14.017	0.000	0.193	0.000	0.000	0.207	0.000

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	234	271	409	2719	0	1085	0	0	46	0
N.S.	1	1.16	1.75	11.62	0.00	4.64	0.00	0.00	0.20	0.00
time (sec)	N/A	1.303	0.118	10.923	0.000	0.173	0.000	0.000	0.198	0.000

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	162	182	295	1819	0	681	0	0	25	0
N.S.	1	1.12	1.82	11.23	0.00	4.20	0.00	0.00	0.15	0.00
time (sec)	N/A	0.644	0.199	3.641	0.000	0.142	0.000	0.000	0.194	0.000

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	86	127	120	184	389	0	0	9	0
N.S.	1	1.09	1.61	1.52	2.33	4.92	0.00	0.00	0.11	0.00
time (sec)	N/A	0.347	0.019	2.013	0.157	0.134	0.000	0.000	0.173	0.000

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.217	0.088	0.385	1.227	0.096	0.706	0.265	0.183	3.429

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	391	514	341	6725	0	1799	0	0	17	0
N.S.	1	1.31	0.87	17.20	0.00	4.60	0.00	0.00	0.04	0.00
time (sec)	N/A	1.501	3.335	49.451	0.000	0.268	0.000	0.000	0.183	0.000

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	293	394	255	6375	0	1463	0	0	15	0
N.S.	1	1.34	0.87	21.76	0.00	4.99	0.00	0.00	0.05	0.00
time (sec)	N/A	1.104	2.751	6.407	0.000	0.248	0.000	0.000	0.176	0.000

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	283	390	563	392	1099	0	0	13	0
N.S.	1	1.46	2.01	2.90	2.02	5.66	0.00	0.00	0.07	0.00
time (sec)	N/A	1.093	3.751	4.053	0.193	0.224	0.000	0.000	0.172	0.000

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.470	1.881	0.351	3.073	0.087	1.115	0.504	0.174	4.177

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	168	219	155	2387	344	180	0	0	20	0
N.S.	1	1.30	0.92	14.21	2.05	1.07	0.00	0.00	0.12	0.00
time (sec)	N/A	1.319	0.252	4.770	0.061	0.095	0.000	0.000	0.184	0.000

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	132	172	119	2289	249	157	0	0	18	0
N.S.	1	1.30	0.90	17.34	1.89	1.19	0.00	0.00	0.14	0.00
time (sec)	N/A	0.767	0.173	3.644	0.049	0.096	0.000	0.000	0.188	0.000

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	126	709	307	288	122	0	0	16	0
N.S.	1	1.35	7.62	3.30	3.10	1.31	0.00	0.00	0.17	0.00
time (sec)	N/A	0.497	9.927	3.872	0.115	0.106	0.000	0.000	0.185	0.000

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	143	37	17	20	20	21
N.S.	1	1.00	1.10	0.90	7.15	1.85	0.85	1.00	1.00	1.05
time (sec)	N/A	0.257	0.804	0.578	4.324	0.089	1.098	0.292	0.174	4.473

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	169	216	155	2277	345	180	0	0	22	0
N.S.	1	1.28	0.92	13.47	2.04	1.07	0.00	0.00	0.13	0.00
time (sec)	N/A	0.892	0.245	4.576	0.068	0.115	0.000	0.000	0.197	0.000

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	133	169	119	2187	250	157	0	0	20	0
N.S.	1	1.27	0.89	16.44	1.88	1.18	0.00	0.00	0.15	0.00
time (sec)	N/A	0.707	0.175	3.550	0.060	0.090	0.000	0.000	0.195	0.000

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	123	605	332	286	122	0	0	18	0
N.S.	1	1.31	6.44	3.53	3.04	1.30	0.00	0.00	0.19	0.00
time (sec)	N/A	0.480	8.977	4.677	0.117	0.087	0.000	0.000	0.188	0.000

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	144	37	19	21	22	22
N.S.	1	1.00	1.10	0.90	6.86	1.76	0.90	1.00	1.05	1.05
time (sec)	N/A	0.259	0.783	0.707	4.416	0.089	1.124	0.289	0.195	3.916

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	51	18	58	65	0	0	6	0
N.S.	1	1.00	2.43	0.86	2.76	3.10	0.00	0.00	0.29	0.00
time (sec)	N/A	0.194	0.074	0.156	0.031	0.097	0.000	0.000	0.176	0.000

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	42	71	32	59	95	0	0	8	0
N.S.	1	0.98	1.65	0.74	1.37	2.21	0.00	0.00	0.19	0.00
time (sec)	N/A	0.328	0.017	0.145	0.032	0.092	0.000	0.000	0.179	0.000

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	70	93	49	76	120	0	0	10	0
N.S.	1	1.21	1.60	0.84	1.31	2.07	0.00	0.00	0.17	0.00
time (sec)	N/A	0.452	0.018	0.214	0.031	0.098	0.000	0.000	0.165	0.000

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	33	68	32	107	138	0	0	10	0
N.S.	1	0.94	1.94	0.91	3.06	3.94	0.00	0.00	0.29	0.00
time (sec)	N/A	0.206	0.016	0.394	0.032	0.107	0.000	0.000	0.171	0.000

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	113	155	108	199	0	0	12	0
N.S.	1	1.00	1.59	2.18	1.52	2.80	0.00	0.00	0.17	0.00
time (sec)	N/A	0.384	0.032	0.378	0.043	0.090	0.000	0.000	0.170	0.000

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	118	149	197	142	248	0	0	14	0
N.S.	1	1.17	1.48	1.95	1.41	2.46	0.00	0.00	0.14	0.00
time (sec)	N/A	0.529	0.032	0.447	0.041	0.096	0.000	0.000	0.177	0.000

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	149	108	160	202	284	0	0	14	0
N.S.	1	0.89	0.64	0.95	1.20	1.69	0.00	0.00	0.08	0.00
time (sec)	N/A	0.526	0.475	1.500	0.035	0.096	0.000	0.000	0.172	0.000

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	211	211	177	596	194	396	0	0	16	0
N.S.	1	1.00	0.84	2.82	0.92	1.88	0.00	0.00	0.08	0.00
time (sec)	N/A	0.669	0.076	0.856	0.058	0.102	0.000	0.000	0.176	0.000

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	289	235	672	254	480	0	0	18	0
N.S.	1	1.09	0.89	2.55	0.96	1.82	0.00	0.00	0.07	0.00
time (sec)	N/A	0.979	0.052	1.260	0.061	0.107	0.000	0.000	0.163	0.000

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	<b>F(-1)</b>	A	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	112	91	153	868	184	234	0	157	25	179
N.S.	1	0.81	1.37	7.75	1.64	2.09	0.00	1.40	0.22	1.60
time (sec)	N/A	0.486	0.096	5.077	0.113	0.107	0.000	0.279	0.181	4.103

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	49	46	60	887	64	93	0	94	25	111
N.S.	1	0.94	1.22	18.10	1.31	1.90	0.00	1.92	0.51	2.27
time (sec)	N/A	0.323	0.044	0.773	0.035	0.090	0.000	0.129	0.212	3.930

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	41	46	37	43	46	56	35	29	28
N.S.	1	0.91	1.02	0.82	0.96	1.02	1.24	0.78	0.64	0.62
time (sec)	N/A	0.306	0.043	0.371	0.039	0.078	0.649	0.112	0.198	0.158

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	41	46	27	43	57	0	40	29	28
N.S.	1	0.91	1.02	0.60	0.96	1.27	0.00	0.89	0.64	0.62
time (sec)	N/A	0.306	0.046	0.431	0.040	0.083	0.000	0.117	0.191	3.617

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	<b>F(-1)</b>	B	<b>F</b>	B
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	49	46	59	872	64	92	0	98	25	119
N.S.	1	0.94	1.20	17.80	1.31	1.88	0.00	2.00	0.51	2.43
time (sec)	N/A	0.330	0.045	0.718	0.037	0.084	0.000	0.202	0.216	4.113



Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	107	91	150	842	184	233	0	167	25	187
N.S.	1	0.85	1.40	7.87	1.72	2.18	0.00	1.56	0.23	1.75
time (sec)	N/A	0.505	0.086	4.468	0.118	0.108	0.000	0.283	0.202	4.010

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD	TBD
size	110	136	114	327	0	327	0	0	64	0
N.S.	1	1.24	1.04	2.97	0.00	2.97	0.00	0.00	0.58	0.00
time (sec)	N/A	0.749	0.183	179.308	0.000	0.098	0.000	0.000	0.192	0.000

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [281] had the largest ratio of [2.33333000000000013]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	6	1.14	23	0.261
2	A	6	5	1.14	23	0.217
3	A	5	4	1.13	21	0.190
4	C	12	11	0.78	23	0.478
5	A	2	2	1.00	23	0.087
6	A	3	3	1.01	23	0.130
7	A	4	4	1.05	23	0.174
8	A	5	5	1.07	23	0.217
9	A	5	4	0.98	23	0.174
10	A	5	4	1.02	23	0.174
11	A	5	4	1.06	23	0.174
12	A	2	2	1.00	19	0.105
13	A	5	4	1.00	23	0.174
14	A	6	5	0.95	23	0.217
15	A	7	6	1.00	23	0.261
16	A	7	6	1.09	25	0.240
17	A	6	5	1.08	25	0.200
18	A	5	4	1.06	25	0.160
19	A	4	3	1.00	25	0.120
20	A	5	4	1.03	25	0.160
21	A	6	5	1.05	25	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	7	6	1.06	25	0.240
23	A	9	8	1.07	25	0.320
24	A	8	7	1.06	25	0.280
25	A	7	6	1.07	25	0.240
26	A	8	7	1.03	25	0.280
27	A	9	8	1.04	25	0.320
28	A	5	4	0.82	12	0.333
29	A	5	4	0.81	14	0.286
30	N/A	1	0	1.00	40	0.000
31	A	7	6	0.98	40	0.150
32	A	6	5	1.01	40	0.125
33	A	3	2	0.94	38	0.053
34	N/A	1	0	1.00	40	0.000
35	N/A	1	0	1.00	40	0.000
36	A	2	2	1.00	11	0.182
37	A	2	2	1.00	11	0.182
38	A	2	2	1.00	9	0.222
39	A	3	2	1.00	7	0.286
40	A	2	2	1.00	11	0.182
41	A	2	2	1.00	11	0.182
42	A	2	2	1.00	11	0.182
43	A	2	2	1.00	11	0.182
44	A	3	3	0.93	13	0.231
45	A	3	3	1.07	13	0.231
46	A	3	3	1.07	13	0.231
47	A	4	3	1.00	11	0.273
48	A	3	2	1.00	9	0.222
49	A	3	3	1.02	13	0.231
50	A	3	3	1.00	13	0.231
51	A	3	3	1.00	13	0.231
52	A	1	1	1.00	13	0.077
53	A	2	2	1.52	13	0.154

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	4	4	0.86	13	0.308
55	A	4	4	1.10	13	0.308
56	A	5	4	1.15	13	0.308
57	A	4	3	1.00	11	0.273
58	A	3	2	1.00	9	0.222
59	A	4	4	1.03	13	0.308
60	A	4	4	1.00	13	0.308
61	A	4	4	1.02	13	0.308
62	A	4	4	1.00	13	0.308
63	A	1	1	1.00	13	0.077
64	A	2	2	1.00	13	0.154
65	A	5	5	0.81	13	0.385
66	A	5	5	1.11	13	0.385
67	A	5	5	1.11	13	0.385
68	A	5	5	1.11	13	0.385
69	A	6	5	1.22	13	0.385
70	A	5	4	1.15	13	0.308
71	A	4	3	1.00	11	0.273
72	A	3	2	1.00	9	0.222
73	A	5	5	1.03	13	0.385
74	A	5	5	1.02	13	0.385
75	A	5	5	1.01	13	0.385
76	A	5	5	1.08	13	0.385
77	A	5	5	1.00	13	0.385
78	A	1	1	1.00	13	0.077
79	A	2	2	1.00	13	0.154
80	A	3	3	1.18	13	0.231
81	B	4	4	2.10	13	0.308
82	A	5	5	1.11	13	0.385
83	A	5	5	1.11	13	0.385
84	A	4	3	1.00	11	0.273
85	A	1	1	1.00	13	0.077

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	6	5	1.07	13	0.385
87	A	5	4	1.05	13	0.308
88	A	4	3	1.00	11	0.273
89	A	3	2	1.00	9	0.222
90	A	5	4	1.00	13	0.308
91	A	6	5	1.25	13	0.385
92	A	7	6	1.32	13	0.462
93	A	2	2	1.00	13	0.154
94	A	7	6	1.12	13	0.462
95	A	6	5	1.09	13	0.385
96	A	5	4	1.08	13	0.308
97	A	4	3	1.00	11	0.273
98	A	3	2	1.00	9	0.222
99	A	6	5	1.26	13	0.385
100	A	7	6	1.32	13	0.462
101	A	8	7	1.31	13	0.538
102	A	3	3	1.03	13	0.231
103	A	7	6	1.16	13	0.462
104	A	6	5	1.14	13	0.385
105	A	5	4	1.11	13	0.308
106	A	4	3	1.00	11	0.273
107	A	3	2	1.00	9	0.222
108	A	7	6	1.38	13	0.462
109	A	8	7	1.38	13	0.538
110	A	9	8	1.36	13	0.615
111	A	7	6	1.24	15	0.400
112	A	6	5	1.18	15	0.333
113	A	5	4	1.14	15	0.267
114	A	4	3	1.00	13	0.231
115	A	3	2	1.00	11	0.182
116	A	2	2	1.00	15	0.133
117	A	2	2	1.00	15	0.133

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	4	4	0.99	15	0.267
119	A	6	6	1.09	15	0.400
120	A	7	6	1.24	15	0.400
121	A	6	5	1.20	15	0.333
122	A	5	4	1.14	15	0.267
123	A	4	3	1.00	13	0.231
124	A	3	2	1.00	11	0.182
125	A	3	3	1.03	15	0.200
126	A	3	3	1.05	15	0.200
127	A	3	3	0.98	15	0.200
128	A	5	5	1.01	15	0.333
129	A	7	6	1.24	15	0.400
130	A	6	5	1.20	15	0.333
131	A	5	4	1.14	15	0.267
132	A	4	3	1.00	13	0.231
133	A	3	2	1.00	11	0.182
134	A	4	4	1.03	15	0.267
135	A	4	4	1.05	15	0.267
136	A	4	4	0.99	15	0.267
137	A	4	4	1.01	15	0.267
138	A	6	6	1.03	15	0.400
139	A	8	8	1.10	15	0.533
140	A	7	6	1.20	15	0.400
141	A	6	5	1.21	15	0.333
142	A	5	4	1.11	15	0.267
143	A	4	3	1.00	13	0.231
144	A	3	2	1.00	11	0.182
145	A	1	1	1.00	15	0.067
146	A	3	3	1.06	15	0.200
147	A	5	5	1.09	15	0.333
148	A	7	7	1.15	15	0.467
149	A	7	6	1.23	15	0.400

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	6	5	1.19	15	0.333
151	A	5	4	1.11	15	0.267
152	A	4	3	1.00	13	0.231
153	A	3	2	1.00	11	0.182
154	A	2	2	1.00	15	0.133
155	A	4	4	1.19	15	0.267
156	A	6	6	1.15	15	0.400
157	A	8	8	1.19	15	0.533
158	A	7	6	1.18	15	0.400
159	A	6	5	1.16	15	0.333
160	A	5	4	1.07	15	0.267
161	A	4	3	1.00	13	0.231
162	A	3	2	1.00	11	0.182
163	A	3	3	1.17	15	0.200
164	A	5	5	1.26	15	0.333
165	A	7	7	1.20	15	0.467
166	A	9	9	1.22	15	0.600
167	A	2	2	1.00	13	0.154
168	A	2	2	1.00	13	0.154
169	A	2	2	1.00	13	0.154
170	A	2	2	1.00	13	0.154
171	A	2	2	1.00	13	0.154
172	A	2	2	1.00	13	0.154
173	A	2	2	1.00	13	0.154
174	A	2	2	1.00	13	0.154
175	A	3	3	1.06	15	0.200
176	A	3	3	1.06	15	0.200
177	A	3	3	1.06	15	0.200
178	A	3	3	1.06	15	0.200
179	A	3	3	1.02	15	0.200
180	A	3	3	1.02	15	0.200
181	A	3	3	0.98	15	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
182	A	3	3	1.06	15	0.200
183	A	4	4	1.09	15	0.267
184	A	4	4	1.09	15	0.267
185	A	4	4	1.09	15	0.267
186	A	4	4	1.06	15	0.267
187	A	4	4	1.09	15	0.267
188	A	4	4	1.06	15	0.267
189	A	4	4	1.03	15	0.267
190	A	4	4	1.03	15	0.267
191	A	5	5	1.08	15	0.333
192	A	4	4	1.07	15	0.267
193	A	3	3	1.06	15	0.200
194	A	2	2	1.00	15	0.133
195	A	1	1	1.00	15	0.067
196	A	2	2	1.00	15	0.133
197	A	3	3	1.17	15	0.200
198	A	4	4	1.25	15	0.267
199	A	5	5	1.12	15	0.333
200	A	4	4	1.12	15	0.267
201	A	3	3	1.10	15	0.200
202	A	2	2	1.00	15	0.133
203	A	3	3	1.06	15	0.200
204	A	4	4	1.21	15	0.267
205	A	5	5	1.29	15	0.333
206	A	6	6	1.33	15	0.400
207	A	5	5	1.11	15	0.333
208	A	4	4	1.09	15	0.267
209	A	3	3	1.04	15	0.200
210	A	4	4	1.06	15	0.267
211	A	5	5	1.14	15	0.333
212	A	6	6	1.23	15	0.400
213	A	7	7	1.28	15	0.467
Continued on next page						



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
214	A	8	8	1.32	15	0.533
215	A	4	4	1.02	17	0.235
216	A	3	3	0.97	17	0.176
217	A	2	2	1.00	17	0.118
218	A	2	2	1.00	17	0.118
219	A	1	1	1.00	17	0.059
220	A	2	2	1.00	17	0.118
221	A	3	3	1.16	17	0.176
222	A	4	4	1.24	17	0.235
223	A	5	5	1.03	17	0.294
224	A	4	4	1.00	17	0.235
225	A	3	3	0.98	17	0.176
226	A	3	3	1.02	17	0.176
227	A	3	3	1.03	17	0.176
228	A	1	1	1.00	17	0.059
229	A	2	2	1.00	17	0.118
230	A	3	3	1.16	17	0.176
231	A	4	4	1.24	17	0.235
232	A	5	5	1.02	17	0.294
233	A	4	4	1.01	17	0.235
234	A	4	4	1.00	17	0.235
235	A	4	4	1.03	17	0.235
236	A	4	4	1.02	17	0.235
237	A	1	1	1.00	17	0.059
238	A	2	2	1.00	17	0.118
239	A	3	3	1.16	17	0.176
240	A	4	4	1.24	17	0.235
241	A	4	4	1.04	17	0.235
242	A	3	3	1.00	17	0.176
243	A	2	2	1.00	17	0.118
244	A	1	1	1.00	17	0.059
245	A	1	1	1.00	17	0.059

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
246	A	2	2	1.00	17	0.118
247	A	3	3	1.16	17	0.176
248	A	4	4	1.24	17	0.235
249	A	5	5	1.07	17	0.294
250	A	4	4	1.05	17	0.235
251	A	3	3	1.05	17	0.176
252	A	2	2	1.00	17	0.118
253	A	1	1	1.00	17	0.059
254	A	2	2	1.00	17	0.118
255	A	3	3	1.13	17	0.176
256	A	4	4	1.22	17	0.235
257	A	5	5	1.08	17	0.294
258	A	4	4	1.09	17	0.235
259	A	3	3	1.07	17	0.176
260	A	1	1	1.00	17	0.059
261	A	2	2	1.00	17	0.118
262	A	3	3	1.16	17	0.176
263	A	4	4	1.21	17	0.235
264	A	5	5	1.25	17	0.294
265	A	1	1	1.00	13	0.077
266	A	7	6	0.91	13	0.462
267	A	6	5	0.96	13	0.385
268	A	5	4	1.00	13	0.308
269	A	4	3	1.00	11	0.273
270	A	3	2	1.00	9	0.222
271	A	1	1	1.00	13	0.077
272	A	2	2	1.00	13	0.154
273	A	3	3	0.97	13	0.231
274	A	2	2	1.00	11	0.182
275	A	2	2	1.00	11	0.182
276	A	2	2	1.00	9	0.222
277	A	3	2	1.00	7	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
278	A	2	2	1.00	11	0.182
279	A	2	2	1.00	11	0.182
280	A	2	2	1.00	11	0.182
281	C	8	7	1.48	3	2.333
282	C	9	8	1.35	5	1.600
283	C	10	9	1.29	7	1.286
284	A	7	6	1.29	15	0.400
285	A	6	5	1.32	13	0.385
286	A	5	4	1.38	11	0.364
287	N/A	1	0	1.00	15	0.000
288	A	9	8	1.28	16	0.500
289	A	8	7	1.29	16	0.438
290	A	7	6	1.30	14	0.429
291	A	6	5	1.30	12	0.417
292	N/A	1	0	1.00	16	0.000
293	A	9	8	1.30	19	0.421
294	A	8	7	1.31	19	0.368
295	A	7	6	1.33	17	0.353
296	A	6	5	1.36	15	0.333
297	N/A	1	0	1.00	19	0.000
298	A	7	6	1.29	15	0.400
299	A	6	5	1.31	13	0.385
300	A	5	4	1.38	11	0.364
301	N/A	1	0	1.00	15	0.000
302	A	9	8	1.28	16	0.500
303	A	8	7	1.29	16	0.438
304	A	7	6	1.28	14	0.429
305	A	6	5	1.29	12	0.417
306	N/A	1	0	1.00	16	0.000
307	A	9	8	1.30	19	0.421
308	A	8	7	1.31	19	0.368
309	A	7	6	1.31	17	0.353

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
310	A	6	5	1.34	15	0.333
311	N/A	1	0	1.00	19	0.000
312	A	9	8	1.18	15	0.533
313	A	8	7	1.16	15	0.467
314	A	7	6	1.12	13	0.462
315	A	6	5	1.09	7	0.714
316	N/A	1	0	1.00	15	0.000
317	A	7	6	1.32	15	0.400
318	A	6	5	1.36	13	0.385
319	A	5	4	1.48	11	0.364
320	N/A	1	0	1.00	15	0.000
321	A	8	7	1.28	20	0.350
322	A	7	6	1.29	18	0.333
323	A	6	5	1.32	16	0.312
324	N/A	1	0	1.00	20	0.000
325	A	8	7	1.30	21	0.333
326	A	7	6	1.31	19	0.316
327	A	6	5	1.36	17	0.294
328	N/A	1	0	1.00	21	0.000
329	A	9	8	1.18	15	0.533
330	A	8	7	1.16	15	0.467
331	A	7	6	1.12	13	0.462
332	A	6	5	1.09	7	0.714
333	N/A	1	0	1.00	15	0.000
334	A	7	6	1.31	15	0.400
335	A	6	5	1.34	13	0.385
336	A	5	4	1.46	11	0.364
337	N/A	1	0	1.00	15	0.000
338	A	8	7	1.30	20	0.350
339	A	7	6	1.30	18	0.333
340	A	6	5	1.35	16	0.312

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
341	N/A	1	0	1.00	20	0.000
342	A	8	7	1.28	21	0.333
343	A	7	6	1.27	19	0.316
344	A	6	5	1.31	17	0.294
345	N/A	1	0	1.00	21	0.000
346	A	3	2	1.00	4	0.500
347	A	5	4	0.98	6	0.667
348	A	6	5	1.21	8	0.625
349	A	3	2	0.94	8	0.250
350	A	5	4	1.00	10	0.400
351	A	6	5	1.17	12	0.417
352	A	9	8	0.89	12	0.667
353	A	6	5	1.00	14	0.357
354	A	7	6	1.09	16	0.375
355	A	8	7	0.81	20	0.350
356	A	7	6	0.94	20	0.300
357	A	4	3	0.91	20	0.150
358	A	4	3	0.91	20	0.150
359	A	7	6	0.94	20	0.300
360	A	9	8	0.85	20	0.400
361	A	2	2	1.24	24	0.083

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	157
3.2	$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	165
3.3	$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	171
3.4	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx \dots\dots\dots$	177
3.5	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx \dots\dots\dots$	186
3.6	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx \dots\dots\dots$	191
3.7	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx \dots\dots\dots$	196
3.8	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx \dots\dots\dots$	202
3.9	$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	208
3.10	$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	215
3.11	$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	221
3.12	$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	227
3.13	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx \dots\dots\dots$	232
3.14	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx \dots\dots\dots$	238
3.15	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx \dots\dots\dots$	244
3.16	$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	251
3.17	$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \dots\dots\dots$	258

3.18	$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	264
3.19	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$	270
3.20	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$	276
3.21	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$	282
3.22	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$	289
3.23	$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	296
3.24	$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$	304
3.25	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$	311
3.26	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$	318
3.27	$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$	326
3.28	$\int x^3 \operatorname{arctanh}(a + bx^4) dx$	335
3.29	$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx$	342
3.30	$\int \frac{(a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^n}{1-c^2x^2} dx$	348
3.31	$\int \frac{(a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^3}{1-c^2x^2} dx$	353
3.32	$\int \frac{(a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^2}{1-c^2x^2} dx$	362
3.33	$\int \frac{a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1-c^2x^2} dx$	371
3.34	$\int \frac{1}{(1-c^2x^2)(a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))} dx$	377
3.35	$\int \frac{1}{(1-c^2x^2)(a+b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^2} dx$	382
3.36	$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$	388
3.37	$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx$	394
3.38	$\int x \operatorname{arctanh}(\tanh(a + bx)) dx$	399
3.39	$\int \operatorname{arctanh}(\tanh(a + bx)) dx$	404
3.40	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} dx$	409
3.41	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^2} dx$	414
3.42	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^3} dx$	419
3.43	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx$	424
3.44	$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$	429
3.45	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx$	435

3.46	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx$	440
3.47	$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx$	445
3.48	$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx$	450
3.49	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x} dx$	455
3.50	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^2} dx$	461
3.51	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^3} dx$	467
3.52	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^4} dx$	472
3.53	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^5} dx$	477
3.54	$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$	483
3.55	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx$	491
3.56	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx$	497
3.57	$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx$	503
3.58	$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx$	509
3.59	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x} dx$	515
3.60	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^2} dx$	522
3.61	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^3} dx$	529
3.62	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^4} dx$	536
3.63	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^5} dx$	542
3.64	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^6} dx$	547
3.65	$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$	553
3.66	$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$	561
3.67	$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$	568
3.68	$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx$	575
3.69	$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx$	581
3.70	$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx$	587
3.71	$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx$	593
3.72	$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx$	599
3.73	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx$	605
3.74	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^2} dx$	612
3.75	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^3} dx$	618
3.76	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^4} dx$	625
3.77	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^5} dx$	632
3.78	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^6} dx$	638
3.79	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^7} dx$	644
3.80	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^8} dx$	650



3.81	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx$	656
3.82	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{10}} dx$	663
3.83	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{11}} dx$	670
3.84	$\int x \operatorname{arctanh}(\tanh(a+bx))^6 dx$	677
3.85	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))} dx$	684
3.86	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx$	689
3.87	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx$	695
3.88	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx$	701
3.89	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx$	706
3.90	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx$	711
3.91	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))} dx$	716
3.92	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))} dx$	722
3.93	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	729
3.94	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	734
3.95	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	742
3.96	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	749
3.97	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	755
3.98	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	760
3.99	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^2} dx$	765
3.100	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx$	771
3.101	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^2} dx$	778
3.102	$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	786
3.103	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	791
3.104	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	799
3.105	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	806
3.106	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	812
3.107	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	817
3.108	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^3} dx$	822
3.109	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^3} dx$	829
3.110	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^3} dx$	837
3.111	$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	846
3.112	$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	854
3.113	$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	861

3.114	$\int x \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	867
3.115	$\int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	872
3.116	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$	877
3.117	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx$	883
3.118	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx$	889
3.119	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$	896
3.120	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	904
3.121	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	911
3.122	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	918
3.123	$\int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	924
3.124	$\int \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	930
3.125	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x} dx$	935
3.126	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^2} dx$	941
3.127	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^3} dx$	948
3.128	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^4} dx$	955
3.129	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	962
3.130	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	969
3.131	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	976
3.132	$\int x \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	983
3.133	$\int \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	989
3.134	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx$	995
3.135	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^2} dx$	1002
3.136	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^3} dx$	1009
3.137	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx$	1016
3.138	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^5} dx$	1023
3.139	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx$	1031
3.140	$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1040
3.141	$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1047
3.142	$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1054
3.143	$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1060
3.144	$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1066
3.145	$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1071

3.146	$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1076
3.147	$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1083
3.148	$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1090
3.149	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1099
3.150	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1107
3.151	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1114
3.152	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1120
3.153	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1126
3.154	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1131
3.155	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1137
3.156	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1144
3.157	$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1152
3.158	$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1161
3.159	$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1168
3.160	$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1175
3.161	$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1181
3.162	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1186
3.163	$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1191
3.164	$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1198
3.165	$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1205
3.166	$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1213
3.167	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1222
3.168	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1227
3.169	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx)) dx$	1232
3.170	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx)) dx$	1237
3.171	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx$	1242
3.172	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{3/2}} dx$	1247
3.173	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{5/2}} dx$	1252
3.174	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{7/2}} dx$	1257
3.175	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1262
3.176	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1267
3.177	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1273
3.178	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2 dx$	1278

3.179	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx$	1283
3.180	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{3/2}} dx$	1288
3.181	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{5/2}} dx$	1294
3.182	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{7/2}} dx$	1300
3.183	$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1306
3.184	$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1312
3.185	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1318
3.186	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3 dx$	1324
3.187	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} dx$	1330
3.188	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx$	1336
3.189	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{5/2}} dx$	1342
3.190	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{7/2}} dx$	1348
3.191	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx)) x^{7/2}} dx$	1354
3.192	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx)) x^{5/2}} dx$	1362
3.193	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx)) x^{3/2}} dx$	1369
3.194	$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx)) \sqrt{x}} dx$	1375
3.195	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx$	1381
3.196	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1386
3.197	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1392
3.198	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx$	1398
3.199	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1405
3.200	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1413
3.201	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1421
3.202	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$	1427
3.203	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1433
3.204	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1440
3.205	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1447
3.206	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$	1455
3.207	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1465
3.208	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1473
3.209	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1480
3.210	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$	1486

3.211	$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1493
3.212	$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1501
3.213	$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1510
3.214	$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$	1521
3.215	$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	1534
3.216	$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx$	1541
3.217	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx$	1547
3.218	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx$	1553
3.219	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx$	1559
3.220	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx$	1564
3.221	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx$	1570
3.222	$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx$	1576
3.223	$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	1583
3.224	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx$	1590
3.225	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}} dx$	1597
3.226	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}} dx$	1603
3.227	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{5/2}} dx$	1609
3.228	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx$	1615
3.229	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx$	1620
3.230	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx$	1625
3.231	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{13/2}} dx$	1631
3.232	$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx$	1638
3.233	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx$	1645
3.234	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{3/2}} dx$	1652
3.235	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}} dx$	1659
3.236	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{7/2}} dx$	1666
3.237	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx$	1673
3.238	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx$	1678
3.239	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx$	1683
3.240	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{15/2}} dx$	1689
3.241	$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))} x^{5/2}} dx$	1696

3.242	$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1702
3.243	$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1708
3.244	$\int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1714
3.245	$\int \frac{1}{x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1719
3.246	$\int \frac{1}{x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1724
3.247	$\int \frac{1}{x^{7/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1730
3.248	$\int \frac{1}{x^{9/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$	1736
3.249	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1743
3.250	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1751
3.251	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1758
3.252	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1764
3.253	$\int \frac{1}{\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1770
3.254	$\int \frac{1}{x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1775
3.255	$\int \frac{1}{x^{5/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1780
3.256	$\int \frac{1}{x^{7/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$	1786
3.257	$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1793
3.258	$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1801
3.259	$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1808
3.260	$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1814
3.261	$\int \frac{1}{\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1819
3.262	$\int \frac{1}{x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1824
3.263	$\int \frac{1}{x^{5/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1830
3.264	$\int \frac{1}{x^{7/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$	1837
3.265	$\int x^m \operatorname{arctanh}(\tanh(a+bx))^n dx$	1845
3.266	$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1850
3.267	$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1859
3.268	$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^n dx$	1867
3.269	$\int x \operatorname{arctanh}(\tanh(a+bx))^n dx$	1874
3.270	$\int \operatorname{arctanh}(\tanh(a+bx))^n dx$	1880
3.271	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x} dx$	1886
3.272	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^2} dx$	1891

3.273	$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^3} dx$	1896
3.274	$\int x^m \coth^{-1}(\tanh(a+bx)) dx$	1901
3.275	$\int x^2 \operatorname{arctanh}(\coth(a+bx)) dx$	1907
3.276	$\int x \operatorname{arctanh}(\coth(a+bx)) dx$	1912
3.277	$\int \operatorname{arctanh}(\coth(a+bx)) dx$	1917
3.278	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x} dx$	1922
3.279	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^2} dx$	1927
3.280	$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^3} dx$	1932
3.281	$\int \operatorname{arctanh}(\cosh(x)) dx$	1937
3.282	$\int x \operatorname{arctanh}(\cosh(x)) dx$	1943
3.283	$\int x^2 \operatorname{arctanh}(\cosh(x)) dx$	1950
3.284	$\int x^2 \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1958
3.285	$\int x \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1968
3.286	$\int \operatorname{arctanh}(c+d \tanh(a+bx)) dx$	1977
3.287	$\int \frac{\operatorname{arctanh}(c+d \tanh(a+bx))}{x} dx$	1985
3.288	$\int x^3 \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	1990
3.289	$\int x^2 \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	1999
3.290	$\int x \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	2008
3.291	$\int \operatorname{arctanh}(1+d+d \tanh(a+bx)) dx$	2016
3.292	$\int \frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x} dx$	2022
3.293	$\int x^3 \operatorname{arctanh}(1-d-d \tanh(a+bx)) dx$	2027
3.294	$\int x^2 \operatorname{arctanh}(1-d-d \tanh(a+bx)) dx$	2036
3.295	$\int x \operatorname{arctanh}(1-d-d \tanh(a+bx)) dx$	2045
3.296	$\int \operatorname{arctanh}(1-d-d \tanh(a+bx)) dx$	2053
3.297	$\int \frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x} dx$	2059
3.298	$\int x^2 \operatorname{arctanh}(c+d \coth(a+bx)) dx$	2064
3.299	$\int x \operatorname{arctanh}(c+d \coth(a+bx)) dx$	2074
3.300	$\int \operatorname{arctanh}(c+d \coth(a+bx)) dx$	2083
3.301	$\int \frac{\operatorname{arctanh}(c+d \coth(a+bx))}{x} dx$	2091
3.302	$\int x^3 \operatorname{arctanh}(1+d+d \coth(a+bx)) dx$	2096
3.303	$\int x^2 \operatorname{arctanh}(1+d+d \coth(a+bx)) dx$	2105
3.304	$\int x \operatorname{arctanh}(1+d+d \coth(a+bx)) dx$	2114
3.305	$\int \operatorname{arctanh}(1+d+d \coth(a+bx)) dx$	2122
3.306	$\int \frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x} dx$	2128
3.307	$\int x^3 \operatorname{arctanh}(1-d-d \coth(a+bx)) dx$	2133
3.308	$\int x^2 \operatorname{arctanh}(1-d-d \coth(a+bx)) dx$	2142
3.309	$\int x \operatorname{arctanh}(1-d-d \coth(a+bx)) dx$	2151

3.310	$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$	2159
3.311	$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx$	2165
3.312	$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx$	2170
3.313	$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx$	2181
3.314	$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx$	2191
3.315	$\int \operatorname{arctanh}(\tan(a + bx)) dx$	2199
3.316	$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx$	2206
3.317	$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2211
3.318	$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2223
3.319	$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx$	2232
3.320	$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx$	2240
3.321	$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2245
3.322	$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2254
3.323	$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$	2262
3.324	$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx$	2270
3.325	$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2275
3.326	$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2284
3.327	$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$	2292
3.328	$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx$	2299
3.329	$\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx$	2304
3.330	$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx$	2315
3.331	$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx$	2325
3.332	$\int \operatorname{arctanh}(\cot(a + bx)) dx$	2333
3.333	$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx$	2340
3.334	$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2345
3.335	$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2357
3.336	$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx$	2366
3.337	$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx$	2374
3.338	$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2379
3.339	$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2387
3.340	$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$	2395
3.341	$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx$	2402
3.342	$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2407
3.343	$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2415
3.344	$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$	2423
3.345	$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx$	2430
3.346	$\int \operatorname{arctanh}(e^x) dx$	2435



3.347	$\int x \operatorname{arctanh}(e^x) dx$	2440
3.348	$\int x^2 \operatorname{arctanh}(e^x) dx$	2446
3.349	$\int \operatorname{arctanh}(e^{a+bx}) dx$	2452
3.350	$\int x \operatorname{arctanh}(e^{a+bx}) dx$	2457
3.351	$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx$	2463
3.352	$\int \operatorname{arctanh}(a + bf^{c+dx}) dx$	2470
3.353	$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$	2478
3.354	$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx$	2486
3.355	$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$	2495
3.356	$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx$	2503
3.357	$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx$	2510
3.358	$\int e^{c(a+bx)} \operatorname{arctanh}(\coth(ac + bcx)) dx$	2515
3.359	$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx$	2520
3.360	$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx$	2527
3.361	$\int \frac{(a+b \operatorname{arctanh}(cx^n))^{(d+e \log(fx^m))}}{x} dx$	2535

### 3.1 $\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	157
Mathematica [A] (verified)	157
Rubi [A] (verified)	158
Maple [B] (verified)	161
Fricas [A] (verification not implemented)	162
Sympy [A] (verification not implemented)	162
Maxima [F]	163
Giac [F(-1)]	163
Mupad [F(-1)]	164
Reduce [F]	164

#### Optimal result

Integrand size = 23, antiderivative size = 127

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{5d^2x\sqrt{d+ex^2}}{96e^{5/2}} + \frac{5dx^3\sqrt{d+ex^2}}{144e^{3/2}} - \frac{x^5\sqrt{d+ex^2}}{36\sqrt{e}}$$

$$+ \frac{5d^3\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{96e^3} + \frac{1}{6}x^6\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
-5/96*d^2*x*(e*x^2+d)^(1/2)/e^(5/2)+5/144*d*x^3*(e*x^2+d)^(1/2)/e^(3/2)-1/36*x^5*(e*x^2+d)^(1/2)/e^(1/2)+5/96*d^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/e^3+1/6*x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.78

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{\sqrt{ex}\sqrt{d+ex^2}(-15d^2 + 10dex^2 - 8e^2x^4) + 48e^3x^6\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + 15d^3 \log(\sqrt{ex} + \sqrt{d+ex^2})}{288e^3}$$

input `Integrate[x^5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(Sqrt[e]*x*Sqrt[d + e*x^2]*(-15*d^2 + 10*d*e*x^2 - 8*e^2*x^4) + 48*e^3*x^6*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] + 15*d^3*Log[Sqrt[e]*x + Sqrt[d + e*x^2]])/(288*e^3)`

### Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6775, 262, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow 6775 \\
 & \frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \int \frac{x^6}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow 262 \\
 & \frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \left( \frac{x^5\sqrt{d+ex^2}}{6e} - \frac{5d \int \frac{x^4}{\sqrt{ex^2+d}} dx}{6e} \right) \\
 & \quad \downarrow 262 \\
 & \frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \left( \frac{x^5\sqrt{d+ex^2}}{6e} - \frac{5d \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \int \frac{x^2}{\sqrt{ex^2+d}} dx}{4e} \right)}{6e} \right) \\
 & \quad \downarrow 262
 \end{aligned}$$

$$\frac{1}{6}\sqrt{e} \left( \frac{x^5\sqrt{d+ex^2}}{6e} - \frac{\frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 5d \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right)}{6e} \right)$$

↓ 224

$$\frac{1}{6}\sqrt{e} \left( \frac{x^5\sqrt{d+ex^2}}{6e} - \frac{\frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 5d \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1-\frac{ex^2}{ex^2+d}} \frac{dx}{\sqrt{ex^2+d}}} \right)}{4e} \right)}{6e} \right)$$

↓ 219

$$\frac{1}{6}\sqrt{e} \left( \frac{x^5\sqrt{d+ex^2}}{6e} - \frac{\frac{1}{6}x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 5d \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}} \right)}{4e} \right)}{6e} \right)$$

input `Int[x^5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output

$$\frac{(x^6 \operatorname{ArcTanh}[\frac{\sqrt{e}x}{\sqrt{d+ex^2}}])/6 - (\sqrt{e}((x^5 \sqrt{d+ex^2})/(6e) - (5d((x^3 \sqrt{d+ex^2})/(4e) - (3d((x \sqrt{d+ex^2})/(2e) - (d \operatorname{ArcTanh}[\frac{\sqrt{e}x}{\sqrt{d+ex^2}}])/(2e^{3/2}))))/(4e)))/(6e)))/6$$
**Defintions of rubi rules used**

rule 219

$$\operatorname{Int}[(a_ + (b_)(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2](x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 224

$$\operatorname{Int}[1/\sqrt{(a_ + (b_)(x_)^2)}, x\_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\sqrt{a + b*x^2}] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ !\operatorname{GtQ}[a, 0]$$

rule 262

$$\operatorname{Int}[(c_)(x_)^{m_}((a_ + (b_)(x_)^2)^{p_}), x\_Symbol] \rightarrow \operatorname{Simp}[c*(c*x)^{m-1}((a + b*x^2)^{p+1}/(b*(m + 2*p + 1))), x] - \operatorname{Simp}[a*c^2*(m-1)/(b*(m + 2*p + 1)) \operatorname{Int}[(c*x)^{m-2}(a + b*x^2)^p, x], x] /; \operatorname{FreeQ}[\{a, b, c, p\}, x] \ \&\& \operatorname{GtQ}[m, 2 - 1] \ \&\& \operatorname{NeQ}[m + 2*p + 1, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 6775

$$\operatorname{Int}[\operatorname{ArcTanh}[(c_)(x_)/\sqrt{(a_ + (b_)(x_)^2)}]((d_)(x_)^{m_}), x\_Symbol] \rightarrow \operatorname{Simp}[(d*x)^{m+1}(\operatorname{ArcTanh}[(c*x)/\sqrt{a + b*x^2}]/(d*(m + 1))), x] - \operatorname{Simp}[c/(d*(m + 1)) \operatorname{Int}[(d*x)^{m+1}/\sqrt{a + b*x^2}, x], x] /; \operatorname{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \operatorname{EqQ}[b, c^2] \ \&\& \operatorname{NeQ}[m, -1]$$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(97) = 194.

Time = 0.05 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.99

method	result
default	$\frac{x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{6} + \frac{e^{\frac{3}{2}}}{6d} \left( \frac{x^7 \sqrt{ex^2+d}}{8e} - \frac{7d \left( \frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left( \frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left( \frac{x \sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{6e} \right)}{8e} \right)$
parts	$\frac{x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{6} + \frac{e^{\frac{3}{2}}}{6d} \left( \frac{x^7 \sqrt{ex^2+d}}{8e} - \frac{7d \left( \frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left( \frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left( \frac{x \sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{6e} \right)}{8e} \right)$

input

```
int(x^5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
1/6*x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+1/6*e^(3/2)/d*(1/8*x^7/e*(e*x^2+d)^(1/2)-7/8*d/e*(1/6*x^5/e*(e*x^2+d)^(1/2)-5/6*d/e*(1/4*x^3/e*(e*x^2+d)^(1/2)-3/4*d/e*(1/2*x/e*(e*x^2+d)^(1/2)-1/2*d/e^(3/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2)))))-1/6*e^(1/2)/d*(1/8*x^5*(e*x^2+d)^(3/2)/e-5/8*d/e*(1/6*x^3*(e*x^2+d)^(3/2)/e-1/2*d/e*(1/4*x*(e*x^2+d)^(3/2)/e-1/4*d/e*(1/2*x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.68

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2(8e^2x^5 - 10dex^3 + 15d^2x)\sqrt{ex^2+d}\sqrt{e} - 3(16e^3x^6 + 5d^3)\log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right)}{576e^3}$$

input

```
integrate(x^5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
-1/576*(2*(8*e^2*x^5 - 10*d*e*x^3 + 15*d^2*x)*sqrt(e*x^2 + d)*sqrt(e) - 3*(16*e^3*x^6 + 5*d^3)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/e^3
```

**Sympy [A] (verification not implemented)**

Time = 0.91 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.95

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} \frac{5d^3 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{96e^3} - \frac{5d^2x\sqrt{d+ex^2}}{96e^{\frac{5}{2}}} + \frac{5dx^3\sqrt{d+ex^2}}{144e^{\frac{3}{2}}} + \frac{x^6 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6} - \frac{x^5\sqrt{d+ex^2}}{36\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input

```
integrate(x**5*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

```
Piecewise((5*d**3*atanh(sqrt(e)*x/sqrt(d + e*x**2))/(96*e**3) - 5*d**2*x*sqrt(d + e*x**2)/(96*e**(5/2)) + 5*d*x**3*sqrt(d + e*x**2)/(144*e**(3/2)) + x**6*atanh(sqrt(e)*x/sqrt(d + e*x**2))/6 - x**5*sqrt(d + e*x**2)/(36*sqrt(e)), Ne(e, 0)), (0, True))
```

**Maxima [F]**

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^5 \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x^5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/12*x^6*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/12*x^6*log(-sqrt(e)*x + sqrt(e*x^2 + d)) - 1/2*d*sqrt(e)*integrate(-1/3*sqrt(e*x^2 + d)*x^6/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)
```

**Giac [F(-1)]**

Timed out.

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input

```
integrate(x^5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
Timed out
```



**Mupad [F(-1)]**

Timed out.

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^5 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^5*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`output `int(x^5*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`**Reduce [F]**

$$\int x^5 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^5 dx$$

input `int(x^5*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**5,x)`

### 3.2 $\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (verified)	166
Maple [B] (verified)	168
Fricas [A] (verification not implemented)	168
Sympy [A] (verification not implemented)	169
Maxima [F]	169
Giac [F(-1)]	170
Mupad [F(-1)]	170
Reduce [F]	170

#### Optimal result

Integrand size = 23, antiderivative size = 101

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{3dx\sqrt{d+ex^2}}{32e^{3/2}} - \frac{x^3\sqrt{d+ex^2}}{16\sqrt{e}} - \frac{3d^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{32e^2} + \frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
3/32*d*x*(e*x^2+d)^(1/2)/e^(3/2)-1/16*x^3*(e*x^2+d)^(1/2)/e^(1/2)-3/32*d^2
*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/e^2+1/4*x^4*arctanh(e^(1/2)*x/(e*x^2+d)
)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.87

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{ex}(3d-2ex^2)\sqrt{d+ex^2} + 8e^2x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 3d^2 \log(\sqrt{ex} + \sqrt{d+ex^2})}{32e^2}$$

input `Integrate[x^3*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(Sqrt[e]*x*(3*d - 2*e*x^2)*Sqrt[d + e*x^2] + 8*e^2*x^4*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - 3*d^2*Log[Sqrt[e]*x + Sqrt[d + e*x^2]])/(32*e^2)`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {6775, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{6775} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{e} \int \frac{x^4}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{e} \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \int \frac{x^2}{\sqrt{ex^2+d}} dx}{4e} \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{e} \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right) \\
 & \quad \downarrow \text{224} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{e} \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1-\frac{ex^2}{ex^2+d}} d \frac{x}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right)
 \end{aligned}$$

$$\frac{1}{4}x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{e} \left( \frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}} \right)}{4e} \right)$$

input `Int[x^3*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(x^4*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/4 - (Sqrt[e]*((x^3*Sqrt[d + e*x^2])/(4*e) - (3*d*((x*Sqrt[d + e*x^2])/(2*e) - (d*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(2*e^(3/2))))/(4*e)))/4`

### Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 6775 `Int[ArcTanh[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(77) = 154.

Time = 0.03 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.03

method	result
default	$\frac{x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{4} + \frac{e^{\frac{3}{2}} \left( \frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left( \frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left( \frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{6e} \right)}{4d} - \frac{\sqrt{e} \left( \frac{x^3 (ex^2+d)^{\frac{3}{2}}}{6e} \right)}{4d}$
parts	$\frac{x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{4} + \frac{e^{\frac{3}{2}} \left( \frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left( \frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left( \frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{6e} \right)}{4d} - \frac{\sqrt{e} \left( \frac{x^3 (ex^2+d)^{\frac{3}{2}}}{6e} \right)}{4d}$

input `int(x^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output `1/4*x^4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+1/4*e^(3/2)/d*(1/6*x^5/e*(e*x^2+d)^(1/2)-5/6*d/e*(1/4*x^3/e*(e*x^2+d)^(1/2)-3/4*d/e*(1/2*x/e*(e*x^2+d)^(1/2)-1/2*d/e^(3/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))-1/4*e^(1/2)/d*(1/6*x^3*(e*x^2+d)^(3/2)/e-1/2*d/e*(1/4*x*(e*x^2+d)^(3/2)/e-1/4*d/e*(1/2*x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx$$

$$= -\frac{2(2ex^3 - 3dx)\sqrt{ex^2+d}\sqrt{e} - (8e^2x^4 - 3d^2) \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{e}x+d}{d}\right)}{64e^2}$$

input `integrate(x^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `-1/64*(2*(2*e*x^3 - 3*d*x)*sqrt(e*x^2 + d)*sqrt(e) - (8*e^2*x^4 - 3*d^2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/e^2`

### Sympy [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.94

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} -\frac{3d^2 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{32e^2} + \frac{3dx\sqrt{d+ex^2}}{32e^{\frac{3}{2}}} + \frac{x^4 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4} - \frac{x^3\sqrt{d+ex^2}}{16\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**3*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((-3*d**2*atanh(sqrt(e)*x/sqrt(d + e*x**2))/(32*e**2) + 3*d*x*sqrt(d + e*x**2)/(32*e**(3/2)) + x**4*atanh(sqrt(e)*x/sqrt(d + e*x**2))/4 - x**3*sqrt(d + e*x**2)/(16*sqrt(e)), Ne(e, 0)), (0, True))`

### Maxima [F]

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^3 \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `1/8*x^4*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/8*x^4*log(-sqrt(e)*x + sqrt(e*x^2 + d)) - 1/2*d*sqrt(e)*integrate(-1/2*sqrt(e*x^2 + d)*x^4/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

**Giac [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input `integrate(x^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^3 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^3*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^3*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^3 dx$$

input `int(x^3*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**3,x)`

### 3.3 $\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [B] (verified)	173
Fricas [A] (verification not implemented)	174
Sympy [A] (verification not implemented)	174
Maxima [F]	175
Giac [F(-1)]	175
Mupad [F(-1)]	176
Reduce [B] (verification not implemented)	176

#### Optimal result

Integrand size = 21, antiderivative size = 75

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{x\sqrt{d+ex^2}}{4\sqrt{e}} + \frac{\operatorname{darctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4e} + \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
-1/4*x*(e*x^2+d)^(1/2)/e^(1/2)+1/4*d*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/e+
1/2*x^2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.01

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{x\sqrt{d+ex^2}}{4\sqrt{e}} + \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + \frac{d \log(\sqrt{ex} + \sqrt{d+ex^2})}{4e}$$

input

```
Integrate[x*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```



output

$$-1/4*(x*\text{Sqrt}[d + e*x^2])/ \text{Sqrt}[e] + (x^2*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]])/2 + (d*\text{Log}[\text{Sqrt}[e]*x + \text{Sqrt}[d + e*x^2]])/(4*e)$$

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6775, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{6775} \\ & \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{e} \int \frac{x^2}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{e} \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right) \\ & \quad \downarrow \text{224} \\ & \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{e} \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1-\frac{ex^2}{ex^2+d}} d \frac{x}{\sqrt{ex^2+d}}}{2e} \right) \\ & \quad \downarrow \text{219} \\ & \frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{e} \left( \frac{x\sqrt{d+ex^2}}{2e} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}} \right) \end{aligned}$$

input

$$\text{Int}[x*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]], x]$$

output

$$(x^2*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]])/2 - (\text{Sqrt}[e]*((x*\text{Sqrt}[d + e*x^2])/2 - (d*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]])/(2*e^(3/2))))/2$$

**Defintions of rubi rules used**

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 224 Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

```
rule 262 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

```
rule 6775 Int[ArcTanh[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_))^(m_), x_
Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))),
x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; Fre
eQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. 2(57) = 114.

Time = 0.03 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.09

method	result
default	$\frac{x^2 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{2} + \frac{e^{\frac{3}{2}} \left( \frac{x^3 \sqrt{e x^2+d}}{4e} - \frac{3d \left( \frac{x \sqrt{e x^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{e x^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{2d} - \frac{\sqrt{e} \left( \frac{x(e x^2+d)^{\frac{3}{2}}}{4e} - \frac{d \left( \frac{x \sqrt{e x^2+d}}{2} + \frac{d \ln(\sqrt{e}x + \sqrt{e x^2+d})}{4e} \right)}{2d} \right)}{2d}$
parts	$\frac{x^2 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{2} + \frac{e^{\frac{3}{2}} \left( \frac{x^3 \sqrt{e x^2+d}}{4e} - \frac{3d \left( \frac{x \sqrt{e x^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{e x^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{2d} - \frac{\sqrt{e} \left( \frac{x(e x^2+d)^{\frac{3}{2}}}{4e} - \frac{d \left( \frac{x \sqrt{e x^2+d}}{2} + \frac{d \ln(\sqrt{e}x + \sqrt{e x^2+d})}{4e} \right)}{4e} \right)}{2d}$

input `int(x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}x^2 \operatorname{arctanh}\left(\frac{e^{1/2}x}{(e x^2 + d)^{1/2}}\right) + \frac{1}{2}e^{3/2}/d \left( \frac{1}{4}x^3/e^{(e x^2 + d)^{1/2}} - \frac{3}{4}d/e^{1/2}x/e^{(e x^2 + d)^{1/2}} - \frac{1}{2}d/e^{3/2} \ln\left(\frac{e^{1/2}x + (e x^2 + d)^{1/2}}{d}\right) \right) - \frac{1}{2}e^{1/2}/d \left( \frac{1}{4}x(e x^2 + d)^{3/2}/e - \frac{1}{4}d/e^{1/2}x(e x^2 + d)^{1/2} + \frac{1}{2}d/e^{1/2} \ln\left(\frac{e^{1/2}x + (e x^2 + d)^{1/2}}{d}\right) \right)$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.79

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = -\frac{2\sqrt{ex^2+d}\sqrt{e}x - (2ex^2+d)\log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{e}x+d}{d}\right)}{8e}$$

input `integrate(x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output  $-1/8*(2*\sqrt{e*x^2+d}*\sqrt{e}*x - (2*e*x^2+d)*\log((2*e*x^2+d)*\sqrt{e}x+d)/d)/e$

### Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.88

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \begin{cases} \frac{d \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{4e} + \frac{x^2 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{2} - \frac{x\sqrt{d+ex^2}}{4\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output

```
Piecewise((d*atanh(sqrt(e)*x/sqrt(d + e*x**2))/(4*e) + x**2*atanh(sqrt(e)*
x/sqrt(d + e*x**2))/2 - x*sqrt(d + e*x**2)/(4*sqrt(e)), Ne(e, 0)), (0, Tru
e))
```

**Maxima [F]**

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/4*x^2*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/4*x^2*log(-sqrt(e)*x + sqrt(e
*x^2 + d)) - 1/2*d*sqrt(e)*integrate(-sqrt(e*x^2 + d)*x^2/(e^2*x^4 + d*e*x
^2 - (e*x^2 + d)^2), x)
```

**Giac [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input

```
integrate(x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
Timed out
```

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

output `int(x*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.76

$$\int x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{2 \operatorname{atanh}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+d+ex^2}}\right) d + 2 \operatorname{atanh}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+d+ex^2}}\right) ex^2 - \sqrt{e}\sqrt{ex^2+d}x - \log\left(\frac{\sqrt{ex^2+d}+\sqrt{ex}}{\sqrt{d}}\right) d}{4e}$$

input `int(x*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)), x)`

output `(2*atanh((sqrt(e)*sqrt(d + e*x**2)*x + e*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d + 2*atanh((sqrt(e)*sqrt(d + e*x**2)*x + e*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e*x**2 - sqrt(e)*sqrt(d + e*x**2)*x - log((sqrt(d + e*x**2) + sqrt(e)*x)/sqrt(d))*d)/(4*e)`

$$3.4 \quad \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx$$

Optimal result	177
Mathematica [A] (verified)	178
Rubi [C] (verified)	178
Maple [A] (verified)	183
Fricas [F]	183
Sympy [F]	184
Maxima [F]	184
Giac [F(-1)]	184
Mupad [F(-1)]	185
Reduce [F]	185

### Optimal result

Integrand size = 23, antiderivative size = 238

$$\begin{aligned} \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = & -\frac{\sqrt{d}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2}{2\sqrt{d+ex^2}} \\ & + \frac{\sqrt{d}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)}{\sqrt{d+ex^2}} \\ & - \frac{\sqrt{d}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log(x)}{\sqrt{d+ex^2}} \\ & + \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)\log(x) \\ & + \frac{\sqrt{d}\sqrt{1+\frac{ex^2}{d}}\operatorname{PolyLog}\left(2, e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)}{2\sqrt{d+ex^2}} \end{aligned}$$

output

$$-1/2*d^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)*x/d^{(1/2)}})^2/(e*x^2+d)^{(1/2)}+d^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)*x/d^{(1/2)}})*\ln(1-(e^{(1/2)*x/d^{(1/2)}})^2)/(e*x^2+d)^{(1/2)}-d^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)*x/d^{(1/2)}})*\ln(x)/(e*x^2+d)^{(1/2)}+\operatorname{arctanh}(e^{(1/2)*x/(e*x^2+d)^{(1/2)}})*\ln(x)+1/2*d^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{polylog}(2,(e^{(1/2)*x/d^{(1/2)}}+(1+e*x^2/d)^{(1/2)})^2)/(e*x^2+d)^{(1/2)}$$
**Mathematica [A] (verified)**

Time = 1.54 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \log(x) + \frac{\sqrt{e}\sqrt{1+\frac{ex^2}{d}}\left(\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right)^2 + 2\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right) \log\left(1 - e^{-2\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right)}\right) - 2\log(x) \log\left(\sqrt{\frac{e}{d}}x + \sqrt{d+ex^2}\right)\right)}{2\sqrt{\frac{e}{d}}\sqrt{d+ex^2}}$$

input

`Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x,x]`

output

`ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]*Log[x] + (Sqrt[e]*Sqrt[1 + (e*x^2)/d] * (ArcSinh[Sqrt[e/d]*x]^2 + 2*ArcSinh[Sqrt[e/d]*x]*Log[1 - E^(-2*ArcSinh[Sqrt[e/d]*x]])] - 2*Log[x]*Log[Sqrt[e/d]*x + Sqrt[1 + (e*x^2)/d]] - PolyLog[2, E^(-2*ArcSinh[Sqrt[e/d]*x]]))/(2*Sqrt[e/d]*Sqrt[d + e*x^2])`
**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.78, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {6773, 2764, 2762, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx \\
& \quad \downarrow \text{6773} \\
& \log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \sqrt{e} \int \frac{\log(x)}{\sqrt{ex^2+d}} dx \\
& \quad \downarrow \text{2764} \\
& \log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{e}\sqrt{\frac{ex^2}{d}+1} \int \frac{\log(x)}{\sqrt{\frac{ex^2}{d}+1}} dx}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{2762} \\
& \log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{e}\sqrt{\frac{ex^2}{d}+1} \left( \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) dx}{\sqrt{e}}}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{6190} \\
& \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \sqrt{e}\sqrt{\frac{ex^2}{d}+1} \left( \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}\sqrt{\frac{ex^2}{d}+1}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{ex}\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \sqrt{e}\sqrt{\frac{ex^2}{d}+1} \left( \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \frac{\sqrt{d} \int -i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) \tan\left(i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{26} \\
& \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \sqrt{e}\sqrt{\frac{ex^2}{d}+1} \left( \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d} \int \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) \tan\left(i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{4199}
\end{aligned}$$



$$\sqrt{e}\sqrt{\frac{ex^2}{d} + 1} \left( \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(2i\int \frac{e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2\right)}{1-e}}{\sqrt{e}} \right)$$

---


$$\sqrt{d+ex^2}$$

↓ 25

$$\sqrt{e}\sqrt{\frac{ex^2}{d} + 1} \left( \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\int \frac{e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2\right)}{1-e}}{\sqrt{e}} \right)$$

---


$$\sqrt{d+ex^2}$$

↓ 2620

$$\sqrt{e}\sqrt{\frac{ex^2}{d} + 1} \left( \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(\frac{1}{2}\int \log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right)}{1-e}}{\sqrt{e}} \right)$$

---


$$\sqrt{d+ex^2}$$

↓ 2715

$$\sqrt{e}\sqrt{\frac{ex^2}{d} + 1} \left( \frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(\frac{1}{4}\int e^{-2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right) de^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\right)}{1-e}}{\sqrt{e}} \right)$$

---


$$\sqrt{d+ex^2}$$

↓ 2838

$$\frac{\log(x)\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \sqrt{e}\sqrt{\frac{ex^2}{d}+1}\left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2,e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right) - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right)}{\sqrt{e}}\right)}{\sqrt{d+ex^2}}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x,x]`

output `ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]*Log[x] - (Sqrt[e]*Sqrt[1 + (e*x^2)/d] * ((Sqrt[d]*ArcSinh[(Sqrt[e]*x)/Sqrt[d]]*Log[x])/Sqrt[e] + (I*Sqrt[d]*((-1/2*I)*ArcSinh[(Sqrt[e]*x)/Sqrt[d]]^2 - (2*I)*(-1/2*(ArcSinh[(Sqrt[e]*x)/Sqrt[d]]*Log[1 - E^(2*ArcSinh[(Sqrt[e]*x)/Sqrt[d]])]) - PolyLog[2, E^(2*ArcSinh[(Sqrt[e]*x)/Sqrt[d]])/4]))/Sqrt[e]))/Sqrt[d + e*x^2]`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2762 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[e, 2]*(x/Sqrt[d])]*((a + b*Log[c*x^n])/Rt[e, 2]), x] - Simp[b*(n/Rt[e, 2]) Int[ArcSinh[Rt[e, 2]*(x/Sqrt[d])]/x, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && GtQ[d, 0] && PosQ[e]`

rule 2764 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[Sqrt[1 + (e/d)*x^2]/Sqrt[d + e*x^2] Int[(a + b*Log[c*x^n])/Sqrt[1 + (e/d)*x^2], x], x] /; FreeQ[{a, b, c, d, e, n}, x] && !GtQ[d, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6190 `Int[((a_.) + ArcSinh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6773 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]/(x_), x_Symbol] := Simp[ArcTanh[c*(x/Sqrt[a + b*x^2])*Log[x], x] - Simp[c Int[Log[x]/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c}, x] && EqQ[b, c^2]`

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.88

method	result
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)^2}{2} + \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) \ln\left(1 + \frac{\frac{\sqrt{e}x}{\sqrt{ex^2+d}}+1}{\sqrt{1-\frac{ex^2}{ex^2+d}}}\right) + \operatorname{polylog}\left(2, -\frac{\frac{\sqrt{e}x}{\sqrt{ex^2+d}}+1}{\sqrt{1-\frac{ex^2}{ex^2+d}}}\right) + \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) \ln\left(1 - \frac{\frac{\sqrt{e}x}{\sqrt{ex^2+d}}+1}{\sqrt{1-\frac{ex^2}{ex^2+d}}}\right) + \operatorname{polylog}\left(2, \frac{\frac{\sqrt{e}x}{\sqrt{ex^2+d}}+1}{\sqrt{1-\frac{ex^2}{ex^2+d}}}\right)$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x,x,method=_RETURNVERBOSE)`

output `-1/2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))^2+arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))*ln(1+(e^(1/2)*x/(e*x^2+d)^(1/2)+1)/(1-e*x^2/(e*x^2+d)^(1/2))+polylog(2,-(e^(1/2)*x/(e*x^2+d)^(1/2)+1)/(1-e*x^2/(e*x^2+d)^(1/2))+arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))*ln(1-(e^(1/2)*x/(e*x^2+d)^(1/2)+1)/(1-e*x^2/(e*x^2+d)^(1/2))+polylog(2,(e^(1/2)*x/(e*x^2+d)^(1/2)+1)/(1-e*x^2/(e*x^2+d)^(1/2)))`

**Fricas [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="fricas")`

output `integral(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="maxima")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x, x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x,x)`output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x, x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x,x)`output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x,x)`

$$3.5 \quad \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$$

Optimal result	186
Mathematica [A] (verified)	186
Rubi [A] (verified)	187
Maple [B] (verified)	188
Fricas [A] (verification not implemented)	188
Sympy [F]	189
Maxima [A] (verification not implemented)	189
Giac [F(-1)]	189
Mupad [F(-1)]	190
Reduce [F]	190

### Optimal result

Integrand size = 23, antiderivative size = 53

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{2dx} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2x^2}$$

output

```
-1/2*e^(1/2)*(e*x^2+d)^(1/2)/d/x-1/2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2
```

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\sqrt{ex}\sqrt{d+ex^2} + d\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2dx^2}$$

input

```
Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^3,x]
```

output

```
-1/2*(Sqrt[e]*x*Sqrt[d + e*x^2] + d*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(d*x^2)
```

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {6775, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$$

↓ 6775

$$\frac{1}{2}\sqrt{e} \int \frac{1}{x^2\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2x^2}$$

↓ 242

$$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2x^2} - \frac{\sqrt{e}\sqrt{d+ex^2}}{2dx}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^3,x]`

output `-1/2*(Sqrt[e]*Sqrt[d + e*x^2])/(d*x) - ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(2*x^2)`

**Defintions of rubi rules used**

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`



**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 110 vs.  $2(41) = 82$ .

Time = 0.04 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.09

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{2x^2} - \frac{e \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2d} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{dx} + \frac{2e \left( \frac{x\sqrt{ex^2+d}}{2} + \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2\sqrt{e}} \right)}{d} \right)}{2d}$	111
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{2x^2} - \frac{e \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2d} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{dx} + \frac{2e \left( \frac{x\sqrt{ex^2+d}}{2} + \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2\sqrt{e}} \right)}{d} \right)}{2d}$	111

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2-1/2*e/d*ln(e^(1/2)*x+(e*x^2+d)^(1/2))+1/2*e^(1/2)/d*(-1/d/x*(e*x^2+d)^(3/2)+2*e/d*(1/2*x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{2\sqrt{ex^2+d}\sqrt{e}x + d \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{e}x+d}{d}\right)}{4dx^2}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="fricas")`

output `-1/4*(2*sqrt(e*x^2 + d)*sqrt(e)*x + d*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d*x^2)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**3,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{2x^2} - \frac{e^{\frac{3}{2}}x^2 + d\sqrt{e}}{2\sqrt{ex^2+d}dx}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="maxima")`

output `-1/2*arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^2 - 1/2*(e^(3/2)*x^2 + d*sqrt(e))/(sqrt(e*x^2 + d)*d*x)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^3} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^3,x)`output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^3, x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^3} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x)`output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**3,x)`

**3.6**  $\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$

Optimal result	191
Mathematica [A] (verified)	191
Rubi [A] (verified)	192
Maple [A] (verified)	193
Fricas [A] (verification not implemented)	194
Sympy [F]	194
Maxima [A] (verification not implemented)	194
Giac [F(-1)]	195
Mupad [F(-1)]	195
Reduce [F]	195

**Optimal result**

Integrand size = 23, antiderivative size = 79

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{12dx^3} + \frac{e^{3/2}\sqrt{d+ex^2}}{6d^2x} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

output

`-1/12*e^(1/2)*(e*x^2+d)^(1/2)/d/x^3+1/6*e^(3/2)*(e*x^2+d)^(1/2)/d^2/x-1/4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4`

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \frac{\sqrt{ex}\sqrt{d+ex^2}(-d+2ex^2) - 3d^2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{12d^2x^4}$$

input

`Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^5,x]`

output

`(Sqrt[e]*x*Sqrt[d + e*x^2]*(-d + 2*e*x^2) - 3*d^2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(12*d^2*x^4)`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {6775, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$$

↓ 6775

$$\frac{1}{4}\sqrt{e} \int \frac{1}{x^4\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

↓ 245

$$\frac{1}{4}\sqrt{e} \left( -\frac{2e \int \frac{1}{x^2\sqrt{ex^2+d}} dx}{3d} - \frac{\sqrt{d+ex^2}}{3dx^3} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

↓ 242

$$\frac{1}{4}\sqrt{e} \left( \frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^5,x]`

output `(Sqrt[e]*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2])/(3*d^2*x)))/4 - ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(4*x^4)`

## Definitions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a + b*x^2)^(p+1)/(a*c*(m+1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 245 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m+1)*((a + b*x^2)^(p+1)/(a*(m+1))), x] - Simp[b*((m + 2*(p + 1) + 1)/(a*(m + 1)) Int[x^(m+2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m+1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m+1))), x] - Simp[c/(d*(m+1)) Int[(d*x)^(m+1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.78

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{4x^4} + \frac{e^{\frac{3}{2}}\sqrt{ex^2+d}}{4d^2x} - \frac{\sqrt{e}(ex^2+d)^{\frac{3}{2}}}{12d^2x^3}$	62
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{4x^4} + \frac{e^{\frac{3}{2}}\sqrt{ex^2+d}}{4d^2x} - \frac{\sqrt{e}(ex^2+d)^{\frac{3}{2}}}{12d^2x^3}$	62

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x,method=_RETURNVERBOSE)`

output 
$$-1/4*\operatorname{arctanh}(e^{1/2}*x/(e*x^2+d)^{1/2})/x^4+1/4*e^{3/2}*(e*x^2+d)^{1/2}/d^2/x-1/12*e^{1/2}/d^2/x^3*(e*x^2+d)^{3/2}$$

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{3d^2 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2(2ex^3 - dx)\sqrt{ex^2+d}\sqrt{e}}{24d^2x^4}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="fricas")`

output `-1/24*(3*d^2*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - 2*(2*e*x^3 - d*x)*sqrt(e*x^2 + d)*sqrt(e))/(d^2*x^4)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**5,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**5, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.77

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \frac{\sqrt{ex^2+d}e^{\frac{3}{2}}}{4d^2x} - \frac{(ex^2+d)^{\frac{3}{2}}\sqrt{e}}{12d^2x^3} - \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{4x^4}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="maxima")`

output `1/4*sqrt(e*x^2 + d)*e^(3/2)/(d^2*x) - 1/12*(e*x^2 + d)^(3/2)*sqrt(e)/(d^2*x^3) - 1/4*arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^4`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^5} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^5,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^5, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^5} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**5,x)`



**3.7** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

Optimal result	196
Mathematica [A] (verified)	196
Rubi [A] (verified)	197
Maple [A] (verified)	198
Fricas [A] (verification not implemented)	199
Sympy [F]	199
Maxima [A] (verification not implemented)	200
Giac [F(-1)]	200
Mupad [F(-1)]	200
Reduce [F]	201

**Optimal result**

Integrand size = 23, antiderivative size = 105

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{30dx^5} + \frac{2e^{3/2}\sqrt{d+ex^2}}{45d^2x^3} - \frac{4e^{5/2}\sqrt{d+ex^2}}{45d^3x} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

output 
$$-1/30*e^{(1/2)}*(e*x^2+d)^{(1/2)}/d/x^5+2/45*e^{(3/2)}*(e*x^2+d)^{(1/2)}/d^2/x^3-4/45*e^{(5/2)}*(e*x^2+d)^{(1/2)}/d^3/x-1/6*\operatorname{arctanh}(e^{(1/2)}*x/(e*x^2+d)^{(1/2)})/x^6$$

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \frac{\sqrt{ex}\sqrt{d+ex^2}(-3d^2+4dex^2-8e^2x^4)-15d^3\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{90d^3x^6}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^7,x]`

output `(Sqrt[e]*x*Sqrt[d + e*x^2]*(-3*d^2 + 4*d*e*x^2 - 8*e^2*x^4) - 15*d^3*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(90*d^3*x^6)`

### Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {6775, 245, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

$$\downarrow 6775$$

$$\frac{1}{6}\sqrt{e} \int \frac{1}{x^6\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 245$$

$$\frac{1}{6}\sqrt{e} \left( -\frac{4e \int \frac{1}{x^4\sqrt{ex^2+d}} dx}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 245$$

$$\frac{1}{6}\sqrt{e} \left( -\frac{4e \left( -\frac{2e \int \frac{1}{x^2\sqrt{ex^2+d}} dx}{3d} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 242$$

$$\frac{1}{6}\sqrt{e} \left( -\frac{4e \left( \frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^7,x]`

output `(Sqrt[e]*(-1/5*Sqrt[d + e*x^2]/(d*x^5) - (4*e*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2])/(3*d^2*x)))/(5*d))/6 - ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(6*x^6)`

### Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 245 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*((m + 2*(p + 1) + 1)/(a*(m + 1))) Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

### Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.05

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{6x^6} - \frac{e^{\frac{3}{2}}\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{6d} + \frac{\sqrt{e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{6d}$	110
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{6x^6} - \frac{e^{\frac{3}{2}}\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{6d} + \frac{\sqrt{e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{6d}$	110

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x,method=_RETURNVERBOSE)`

output `-1/6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^6-1/6*e^(3/2)/d*(-1/3/d/x^3*(e*x^2+d)^(1/2)+2/3*e/d^2/x*(e*x^2+d)^(1/2))+1/6*e^(1/2)/d*(-1/5/d/x^5*(e*x^2+d)^(3/2)+2/15*e/d^2/x^3*(e*x^2+d)^(3/2))`

### Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.74

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

$$= -\frac{15d^3 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) + 2(8e^2x^5 - 4dex^3 + 3d^2x)\sqrt{ex^2+d}\sqrt{e}}{180d^3x^6}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="fricas")`

output `-1/180*(15*d^3*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 2*(8*e^2*x^5 - 4*d*e*x^3 + 3*d^2*x)*sqrt(e*x^2 + d)*sqrt(e))/(d^3*x^6)`

### Sympy [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**7,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**7, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = -\frac{(2e^2x^4 + dex^2 - d^2)e^{\frac{3}{2}}}{18\sqrt{ex^2+d}d^3x^3} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{6x^6} + \frac{(2e^2x^4 - dex^2 - 3d^2)\sqrt{ex^2+d}\sqrt{e}}{90d^3x^5}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="maxima")`

output `-1/18*(2*e^2*x^4 + d*e*x^2 - d^2)*e^(3/2)/(sqrt(e*x^2 + d)*d^3*x^3) - 1/6*arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^6 + 1/90*(2*e^2*x^4 - d*e*x^2 - 3*d^2)*sqrt(e*x^2 + d)*sqrt(e)/(d^3*x^5)`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^7} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^7,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^7, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^7} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^7} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^7, x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**7, x)`

**3.8**  $\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$

Optimal result	202
Mathematica [A] (verified)	202
Rubi [A] (verified)	203
Maple [A] (verified)	205
Fricas [A] (verification not implemented)	205
Sympy [F]	206
Maxima [A] (verification not implemented)	206
Giac [F(-1)]	207
Mupad [F(-1)]	207
Reduce [F]	207

**Optimal result**

Integrand size = 23, antiderivative size = 131

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{56dx^7} + \frac{3e^{3/2}\sqrt{d+ex^2}}{140d^2x^5} - \frac{e^{5/2}\sqrt{d+ex^2}}{35d^3x^3} + \frac{2e^{7/2}\sqrt{d+ex^2}}{35d^4x} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

output `-1/56*e^(1/2)*(e*x^2+d)^(1/2)/d/x^7+3/140*e^(3/2)*(e*x^2+d)^(1/2)/d^2/x^5-1/35*e^(5/2)*(e*x^2+d)^(1/2)/d^3/x^3+2/35*e^(7/2)*(e*x^2+d)^(1/2)/d^4/x-1/8*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^8`

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.65

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \frac{\sqrt{ex}\sqrt{d+ex^2}(-5d^3 + 6d^2ex^2 - 8de^2x^4 + 16e^3x^6) - 35d^4\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{280d^4x^8}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^9,x]`

output `(Sqrt[e]*x*Sqrt[d + e*x^2]*(-5*d^3 + 6*d^2*e*x^2 - 8*d*e^2*x^4 + 16*e^3*x^6) - 35*d^4*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(280*d^4*x^8)`

### Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {6775, 245, 245, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

↓ 6775

$$\frac{1}{8}\sqrt{e} \int \frac{1}{x^8\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

↓ 245

$$\frac{1}{8}\sqrt{e} \left( -\frac{6e \int \frac{1}{x^6\sqrt{ex^2+d}} dx}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

↓ 245

$$\frac{1}{8}\sqrt{e} \left( -\frac{6e \left( -\frac{4e \int \frac{1}{x^4\sqrt{ex^2+d}} dx}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

↓ 245



$$\frac{1}{8}\sqrt{e} \left( -\frac{6e \left( -\frac{4e \left( -\frac{2e \int \frac{1}{x^2 \sqrt{ex^2+d}} dx - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8} \right)$$

↓ 242

$$\frac{1}{8}\sqrt{e} \left( -\frac{6e \left( -\frac{4e \left( \frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^9,x]`

output `(Sqrt[e]*(-1/7*Sqrt[d + e*x^2]/(d*x^7) - (6*e*(-1/5*Sqrt[d + e*x^2]/(d*x^5) - (4*e*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2])/(3*d^2*x)))/(5*d)))/(7*d)))/8 - ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(8*x^8)`

### Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 245 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*(m + 2*(p + 1) + 1)/(a*(m + 1)) Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`

rule 6775

```
Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_
Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))),
x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; Fre
eQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]
```

### Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.21

method	result
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{e^{\frac{3}{2}}\left(-\frac{\sqrt{ex^2+d}}{5dx^5} - \frac{4e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{5d}\right)}{8d} + \frac{\sqrt{e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{7dx^7} - \frac{4e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{7d}\right)}{8d}$
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{e^{\frac{3}{2}}\left(-\frac{\sqrt{ex^2+d}}{5dx^5} - \frac{4e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{5d}\right)}{8d} + \frac{\sqrt{e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{7dx^7} - \frac{4e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{7d}\right)}{8d}$

input

```
int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x,method=_RETURNVERBOSE)
```

output

```
-1/8*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^8-1/8*e^(3/2)/d*(-1/5/d/x^5*(e*x
^2+d)^(1/2)-4/5*e/d*(-1/3/d/x^3*(e*x^2+d)^(1/2)+2/3*e/d^2/x*(e*x^2+d)^(1/2
)))+1/8*e^(1/2)/d*(-1/7/d/x^7*(e*x^2+d)^(3/2)-4/7*e/d*(-1/5/d/x^5*(e*x^2+d
)^(3/2)+2/15*e/d^2/x^3*(e*x^2+d)^(3/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.68

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^9} dx = \frac{35d^4 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2(16e^3x^7 - 8de^2x^5 + 6d^2ex^3 - 5d^3x)\sqrt{ex^2+d}\sqrt{e}}{560d^4x^8}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="fricas")`

output 
$$-1/560*(35*d^4*\log((2*e*x^2 + 2*\sqrt{e*x^2 + d}*\sqrt{e}*x + d)/d) - 2*(16*e^3*x^7 - 8*d*e^2*x^5 + 6*d^2*e*x^3 - 5*d^3*x)*\sqrt{e*x^2 + d}*\sqrt{e})/(d^4*x^8)$$

## Sympy [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**9,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**9, x)`

## Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \frac{(8e^3x^6 + 4de^2x^4 - d^2ex^2 + 3d^3)e^{\frac{3}{2}}}{120\sqrt{ex^2 + d}d^4x^5} - \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{(8e^3x^6 - 4de^2x^4 + 3d^2ex^2 + 15d^3)\sqrt{ex^2 + d}\sqrt{e}}{840d^4x^7}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="maxima")`

output 
$$1/120*(8*e^3*x^6 + 4*d*e^2*x^4 - d^2*e*x^2 + 3*d^3)*e^(3/2)/(sqrt(e*x^2 + d)*d^4*x^5) - 1/8*arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^8 - 1/840*(8*e^3*x^6 - 4*d*e^2*x^4 + 3*d^2*e*x^2 + 15*d^3)*sqrt(e*x^2 + d)*sqrt(e)/(d^4*x^7)$$

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^9} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^9,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^9, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^9} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**9,x)`

### 3.9 $\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	208
Mathematica [A] (verified)	208
Rubi [A] (verified)	209
Maple [B] (verified)	210
Fricas [A] (verification not implemented)	211
Sympy [A] (verification not implemented)	212
Maxima [A] (verification not implemented)	212
Giac [F(-1)]	213
Mupad [F(-1)]	213
Reduce [F]	214

#### Optimal result

Integrand size = 23, antiderivative size = 114

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{d^3 \sqrt{d+ex^2}}{7e^{7/2}} - \frac{d^2 (d+ex^2)^{3/2}}{7e^{7/2}} + \frac{3d(d+ex^2)^{5/2}}{35e^{7/2}} - \frac{(d+ex^2)^{7/2}}{49e^{7/2}} + \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
1/7*d^3*(e*x^2+d)^(1/2)/e^(7/2)-1/7*d^2*(e*x^2+d)^(3/2)/e^(7/2)+3/35*d*(e*x^2+d)^(5/2)/e^(7/2)-1/49*(e*x^2+d)^(7/2)/e^(7/2)+1/7*x^7*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.69

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{d+ex^2}(16d^3 - 8d^2ex^2 + 6de^2x^4 - 5e^3x^6)}{245e^{7/2}} + \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

input

```
Integrate[x^6*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

$$\frac{(\sqrt{d + ex^2} * (16d^3 - 8d^2 * ex^2 + 6d * e^2 * x^4 - 5e^3 * x^6)) / (245 * e^{7/2}) + (x^7 * \text{ArcTanh}[(\sqrt{e} * x) / \sqrt{d + ex^2}]) / 7}{1}$$
**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {6775, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^6 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{6775} \\ & \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{7} \sqrt{e} \int \frac{x^7}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{14} \sqrt{e} \int \frac{x^6}{\sqrt{ex^2+d}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\ & \frac{1}{14} \sqrt{e} \int \left( -\frac{d^3}{e^3 \sqrt{ex^2+d}} + \frac{3\sqrt{ex^2+d}d^2}{e^3} - \frac{3(ex^2+d)^{3/2}d}{e^3} + \frac{(ex^2+d)^{5/2}}{e^3} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{7} x^7 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\ & \frac{1}{14} \sqrt{e} \left( -\frac{2d^3 \sqrt{d+ex^2}}{e^4} + \frac{2d^2 (d+ex^2)^{3/2}}{e^4} + \frac{2(d+ex^2)^{7/2}}{7e^4} - \frac{6d(d+ex^2)^{5/2}}{5e^4} \right) \end{aligned}$$

input

$$\text{Int}[x^6 * \text{ArcTanh}[(\sqrt{e} * x) / \sqrt{d + ex^2}], x]$$

output

$$-1/14*(\text{Sqrt}[e]*((-2*d^3*\text{Sqrt}[d + e*x^2])/e^4 + (2*d^2*(d + e*x^2)^{(3/2)})/e^4 - (6*d*(d + e*x^2)^{(5/2)})/(5*e^4) + (2*(d + e*x^2)^{(7/2)})/(7*e^4))) + (x^7*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]])/7$$
**Defintions of rubi rules used**

rule 53

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^{(n_.)}), x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$$

rule 243

$$\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 2009

$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 6775

$$\text{Int}[\text{ArcTanh}[(c_.)*(x_)]/\text{Sqrt}[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^{(m_.)}), x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTanh}[c*x]/\text{Sqrt}[a + b*x^2])/(d*(m+1)), x] - \text{Simp}[c/(d*(m+1)) \ \text{Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x \ \&\& \ \text{EqQ}[b, c^2] \ \&\& \ \text{NeQ}[m, -1]$$
**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 223 vs.  $2(84) = 168$ .

Time = 0.04 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.96

method	result
default	$\frac{x^7 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{7} + \frac{e^{\frac{3}{2}} \left( \frac{x^8 \sqrt{ex^2+d}}{9e} - \frac{8d \left( \frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left( \frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left( \frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{9e} \right)}{7d} - \frac{\sqrt{e} \left( \frac{x^6 (e}{\dots} \right)}{\dots}$
parts	$\frac{x^7 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{7} + \frac{e^{\frac{3}{2}} \left( \frac{x^8 \sqrt{ex^2+d}}{9e} - \frac{8d \left( \frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left( \frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left( \frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{9e} \right)}{7d} - \frac{\sqrt{e} \left( \frac{x^6 (e}{\dots} \right)}{\dots}$

```
input int(x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 1/7*x^7*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+1/7*e^(3/2)/d*(1/9*x^8/e*(e*x^2+d)^(1/2)-8/9*d/e*(1/7*x^6/e*(e*x^2+d)^(1/2)-6/7*d/e*(1/5*x^4/e*(e*x^2+d)^(1/2)-4/5*d/e*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1/2))))-1/7*e^(1/2)/d*(1/9*x^6*(e*x^2+d)^(3/2)/e-2/3*d/e*(1/7*x^4*(e*x^2+d)^(3/2)/e-4/7*d/e*(1/5*x^2*(e*x^2+d)^(3/2)/e-2/15*d/e^2*(e*x^2+d)^(3/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.77

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \frac{35 e^4 x^7 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2(5e^3x^6 - 6de^2x^4 + 8d^2ex^2 - 16d^3)\sqrt{ex^2+d}\sqrt{e}}{490e^4}$$



input `integrate(x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output 
$$\frac{1}{490} \cdot (35e^4 x^7 \log\left(\frac{2e x^2 + 2\sqrt{e x^2 + d} \sqrt{e} x + d}{d}\right) - 2(5e^3 x^6 - 6d e^2 x^4 + 8d^2 e x^2 - 16d^3) \sqrt{e x^2 + d} \sqrt{e}) / e^4$$

### Sympy [A] (verification not implemented)

Time = 1.43 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.02

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx$$

$$= \begin{cases} \frac{16d^3\sqrt{d+ex^2}}{245e^{\frac{7}{2}}} - \frac{8d^2x^2\sqrt{d+ex^2}}{245e^{\frac{5}{2}}} + \frac{6dx^4\sqrt{d+ex^2}}{245e^{\frac{3}{2}}} + \frac{x^7 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{7} - \frac{x^6\sqrt{d+ex^2}}{49\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**6*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((16*d**3*sqrt(d + e*x**2)/(245*e**(7/2)) - 8*d**2*x**2*sqrt(d + e*x**2)/(245*e**(5/2)) + 6*d*x**4*sqrt(d + e*x**2)/(245*e**(3/2)) + x**7*atanh(sqrt(e)*x/sqrt(d + e*x**2))/7 - x**6*sqrt(d + e*x**2)/(49*sqrt(e)), Ne(e, 0)), (0, True))`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.36

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \frac{1}{7} x^7 \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)$$

$$- \frac{35(ex^2+d)^{\frac{9}{2}} - 135(ex^2+d)^{\frac{7}{2}}d + 189(ex^2+d)^{\frac{5}{2}}d^2 - 105(ex^2+d)^{\frac{3}{2}}d^3}{2205de^{\frac{7}{2}}}$$

$$+ \frac{35(ex^2+d)^{\frac{9}{2}} - 180(ex^2+d)^{\frac{7}{2}}d + 378(ex^2+d)^{\frac{5}{2}}d^2 - 420(ex^2+d)^{\frac{3}{2}}d^3 + 315\sqrt{ex^2+d}d^4}{2205de^{\frac{7}{2}}}$$

input `integrate(x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `1/7*x^7*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)) - 1/2205*(35*(e*x^2 + d)^(9/2) - 135*(e*x^2 + d)^(7/2)*d + 189*(e*x^2 + d)^(5/2)*d^2 - 105*(e*x^2 + d)^(3/2)*d^3)/(d*e^(7/2)) + 1/2205*(35*(e*x^2 + d)^(9/2) - 180*(e*x^2 + d)^(7/2)*d + 378*(e*x^2 + d)^(5/2)*d^2 - 420*(e*x^2 + d)^(3/2)*d^3 + 315*sqrt(e*x^2 + d)*d^4)/(d*e^(7/2))`

### Giac [F(-1)]

Timed out.

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input `integrate(x^6*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `Timed out`

### Mupad [F(-1)]

Timed out.

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^6 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^6*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^6*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int x^6 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^6 dx$$

input `int(x^6*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**6,x)`

### 3.10 $\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	215
Mathematica [A] (verified)	215
Rubi [A] (verified)	216
Maple [B] (verified)	217
Fricas [A] (verification not implemented)	218
Sympy [A] (verification not implemented)	218
Maxima [A] (verification not implemented)	219
Giac [F(-1)]	219
Mupad [F(-1)]	220
Reduce [F]	220

#### Optimal result

Integrand size = 23, antiderivative size = 91

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{d^2 \sqrt{d+ex^2}}{5e^{5/2}} + \frac{2d(d+ex^2)^{3/2}}{15e^{5/2}} - \frac{(d+ex^2)^{5/2}}{25e^{5/2}} + \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
-1/5*d^2*(e*x^2+d)^(1/2)/e^(5/2)+2/15*d*(e*x^2+d)^(3/2)/e^(5/2)-1/25*(e*x^2+d)^(5/2)/e^(5/2)+1/5*x^5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.75

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{\sqrt{d+ex^2}(8d^2-4dex^2+3e^2x^4)}{75e^{5/2}} + \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

input

```
Integrate[x^4*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
-1/75*(Sqrt[d + e*x^2]*(8*d^2 - 4*d*e*x^2 + 3*e^2*x^4))/e^(5/2) + (x^5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/5
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {6775, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow 6775 \\ & \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{5}\sqrt{e} \int \frac{x^5}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow 243 \\ & \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{e} \int \frac{x^4}{\sqrt{ex^2+d}} dx^2 \\ & \quad \downarrow 53 \\ & \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{e} \int \left( \frac{d^2}{e^2\sqrt{ex^2+d}} - \frac{2\sqrt{ex^2+d}d}{e^2} + \frac{(ex^2+d)^{3/2}}{e^2} \right) dx^2 \\ & \quad \downarrow 2009 \\ & \frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{e} \left( \frac{2d^2\sqrt{d+ex^2}}{e^3} + \frac{2(d+ex^2)^{5/2}}{5e^3} - \frac{4d(d+ex^2)^{3/2}}{3e^3} \right) \end{aligned}$$

input

```
Int[x^4*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
-1/10*(Sqrt[e]*((2*d^2*Sqrt[d + e*x^2])/e^3 - (4*d*(d + e*x^2)^(3/2))/(3*e^3) + (2*(d + e*x^2)^(5/2))/(5*e^3))) + (x^5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/5
```



input `int(x^4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{5}x^5 \operatorname{arctanh}\left(\frac{e^{1/2}x}{(e x^2+d)^{1/2}}\right) + \frac{1}{5}e^{3/2}/d \left( \frac{1}{7}x^6/e \cdot (e x^2+d)^{1/2} - \frac{6}{7}d/e \cdot (1/5 x^4/e \cdot (e x^2+d)^{1/2}) - \frac{4}{5}d/e \cdot (1/3 x^2/e \cdot (e x^2+d)^{1/2}) - \frac{2}{3}d/e^2 \cdot (e x^2+d)^{1/2} \right) - \frac{1}{5}e^{1/2}/d \left( \frac{1}{7}x^4 \cdot (e x^2+d)^{3/2} - \frac{4}{7}d/e \cdot (1/5 x^2 \cdot (e x^2+d)^{3/2}) - \frac{2}{15}d/e^2 \cdot (e x^2+d)^{3/2} \right)$$

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.85

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \frac{15e^3x^5 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2(3e^2x^4 - 4dex^2 + 8d^2)\sqrt{ex^2+d}\sqrt{e}}{150e^3}$$

input `integrate(x^4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output 
$$\frac{1}{150} \left( 15e^3x^5 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2(3e^2x^4 - 4dex^2 + 8d^2)\sqrt{ex^2+d}\sqrt{e} \right) / e^3$$

### Sympy [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.99

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \begin{cases} -\frac{8d^2\sqrt{d+ex^2}}{75e^{5/2}} + \frac{4dx^2\sqrt{d+ex^2}}{75e^{3/2}} + \frac{x^5 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{5} - \frac{x^4\sqrt{d+ex^2}}{25\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**4*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output

```
Piecewise((-8*d**2*sqrt(d + e*x**2)/(75*e**(5/2)) + 4*d*x**2*sqrt(d + e*x**2)/(75*e**(3/2)) + x**5*atanh(sqrt(e)*x/sqrt(d + e*x**2))/5 - x**4*sqrt(d + e*x**2)/(25*sqrt(e)), Ne(e, 0)), (0, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.40

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{1}{5} x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) - \frac{15(ex^2+d)^{\frac{7}{2}} - 42(ex^2+d)^{\frac{5}{2}}d + 35(ex^2+d)^{\frac{3}{2}}d^2}{525de^{\frac{5}{2}}}$$

$$+ \frac{5(ex^2+d)^{\frac{7}{2}} - 21(ex^2+d)^{\frac{5}{2}}d + 35(ex^2+d)^{\frac{3}{2}}d^2 - 35\sqrt{ex^2+d}d^3}{175de^{\frac{5}{2}}}$$

input

```
integrate(x^4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/5*x^5*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)) - 1/525*(15*(e*x^2 + d)^(7/2) - 42*(e*x^2 + d)^(5/2)*d + 35*(e*x^2 + d)^(3/2)*d^2)/(d*e^(5/2)) + 1/175*(5*(e*x^2 + d)^(7/2) - 21*(e*x^2 + d)^(5/2)*d + 35*(e*x^2 + d)^(3/2)*d^2 - 35*sqrt(e*x^2 + d)*d^3)/(d*e^(5/2))
```

**Giac [F(-1)]**

Timed out.

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input

```
integrate(x^4*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
Timed out
```



**Mupad [F(-1)]**

Timed out.

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^4 \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^4*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`output `int(x^4*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`**Reduce [F]**

$$\int x^4 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^4 dx$$

input `int(x^4*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**4,x)`

### 3.11 $\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	221
Mathematica [A] (verified)	221
Rubi [A] (verified)	222
Maple [B] (verified)	223
Fricas [A] (verification not implemented)	224
Sympy [A] (verification not implemented)	224
Maxima [A] (verification not implemented)	225
Giac [F(-1)]	225
Mupad [F(-1)]	225
Reduce [F]	226

#### Optimal result

Integrand size = 23, antiderivative size = 68

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{d\sqrt{d+ex^2}}{3e^{3/2}} - \frac{(d+ex^2)^{3/2}}{9e^{3/2}} + \frac{1}{3}x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output

```
1/3*d*(e*x^2+d)^(1/2)/e^(3/2)-1/9*(e*x^2+d)^(3/2)/e^(3/2)+1/3*x^3*arctanh(
e^(1/2)*x/(e*x^2+d)^(1/2))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.82

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{9} \left( \frac{(2d-ex^2)\sqrt{d+ex^2}}{e^{3/2}} + 3x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \right)$$

input

```
Integrate[x^2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
((2*d - e*x^2)*Sqrt[d + e*x^2])/e^(3/2) + 3*x^3*ArcTanh[(Sqrt[e]*x)/Sqrt[
d + e*x^2]]/9
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {6775, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{6775} \\
 & \frac{1}{3}x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{3}\sqrt{e} \int \frac{x^3}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3}x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \int \frac{x^2}{\sqrt{ex^2+d}} dx^2 \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{3}x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \int \left(\frac{\sqrt{ex^2+d}}{e} - \frac{d}{e\sqrt{ex^2+d}}\right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{e} \left(\frac{2(d+ex^2)^{3/2}}{3e^2} - \frac{2d\sqrt{d+ex^2}}{e^2}\right)
 \end{aligned}$$

input `Int [x^2*ArcTanh[(Sqrt [e]*x)/Sqrt [d + e*x^2]],x]`

output `-1/6*(Sqrt [e]*((-2*d*Sqrt [d + e*x^2])/e^2 + (2*(d + e*x^2)^(3/2))/(3*e^2)) + (x^3*ArcTanh[(Sqrt [e]*x)/Sqrt [d + e*x^2]])/3`

## Definitions of rubi rules used

rule 53  $\text{Int}[(a_.) + (b_.)(x_)^{(m_.)}*((c_.) + (d_.)(x_)^{(n_.)}, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

rule 243  $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \ \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 2009  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6775  $\text{Int}[\text{ArcTanh}[(c_.)(x_)]/\text{Sqrt}[(a_.) + (b_.)(x_)^2]*((d_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTanh}[c*x]/\text{Sqrt}[a + b*x^2])/(d*(m+1)), x] - \text{Simp}[c/(d*(m+1)) \ \text{Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b, c^2] \ \&\& \ \text{NeQ}[m, -1]$

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs.  $2(50) = 100$ .

Time = 0.04 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.88

method	result	size
default	$\frac{x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{3} + \frac{e^{\frac{3}{2}} \left( \frac{x^4 \sqrt{e x^2+d}}{5e} - \frac{4d \left( \frac{x^2 \sqrt{e x^2+d}}{3e} - \frac{2d \sqrt{e x^2+d}}{3e^2} \right)}{5e} \right)}{3d} - \frac{\sqrt{e} \left( \frac{x^2 (e x^2+d)^{\frac{3}{2}}}{5e} - \frac{2d (e x^2+d)^{\frac{3}{2}}}{15e^2} \right)}{3d}$	128
parts	$\frac{x^3 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{3} + \frac{e^{\frac{3}{2}} \left( \frac{x^4 \sqrt{e x^2+d}}{5e} - \frac{4d \left( \frac{x^2 \sqrt{e x^2+d}}{3e} - \frac{2d \sqrt{e x^2+d}}{3e^2} \right)}{5e} \right)}{3d} - \frac{\sqrt{e} \left( \frac{x^2 (e x^2+d)^{\frac{3}{2}}}{5e} - \frac{2d (e x^2+d)^{\frac{3}{2}}}{15e^2} \right)}{3d}$	128

input  $\text{int}(x^2*\operatorname{arctanh}(e^{(1/2)}*x/(e*x^2+d)^{(1/2)}),x,\text{method}=\_RETURNVERBOSE)$

output

```
1/3*x^3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+1/3*e^(3/2)/d*(1/5*x^4/e*(e*x^2+d)^(1/2)-4/5*d/e*(1/3*x^2/e*(e*x^2+d)^(1/2))-2/3*d/e^2*(e*x^2+d)^(1/2))-1/3*e^(1/2)/d*(1/5*x^2*(e*x^2+d)^(3/2)/e-2/15*d/e^2*(e*x^2+d)^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.96

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{3e^2x^3 \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2\sqrt{ex^2+d}(ex^2-2d)\sqrt{e}}{18e^2}$$

input

```
integrate(x^2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
1/18*(3*e^2*x^3*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - 2*sqrt(e*x^2 + d)*(e*x^2 - 2*d)*sqrt(e))/e^2
```

**Sympy [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.96

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} \frac{2d\sqrt{d+ex^2}}{9e^{\frac{3}{2}}} + \frac{x^3 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3} - \frac{x^2\sqrt{d+ex^2}}{9\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input

```
integrate(x**2*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

```
Piecewise((2*d*sqrt(d + e*x**2)/(9*e**(3/2)) + x**3*atanh(sqrt(e)*x/sqrt(d + e*x**2))/3 - x**2*sqrt(d + e*x**2)/(9*sqrt(e)), Ne(e, 0)), (0, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.46

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{3} x^3 \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) - \frac{3(ex^2+d)^{\frac{5}{2}} - 5(ex^2+d)^{\frac{3}{2}}d}{45de^{\frac{3}{2}}} + \frac{3(ex^2+d)^{\frac{5}{2}} - 10(ex^2+d)^{\frac{3}{2}}d + 15\sqrt{ex^2+d}d^2}{45de^{\frac{3}{2}}}$$

input `integrate(x^2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)) - 1/45*(3*(e*x^2 + d)^(5/2) - 5*(e*x^2 + d)^(3/2)*d)/(d*e^(3/2)) + 1/45*(3*(e*x^2 + d)^(5/2) - 10*(e*x^2 + d)^(3/2)*d + 15*sqrt(e*x^2 + d)*d^2)/(d*e^(3/2))`

**Giac [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input `integrate(x^2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^2 \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^2*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^2*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^2 dx$$

input `int(x^2*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**2,x)`

## 3.12 $\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	227
Mathematica [A] (verified)	227
Rubi [A] (verified)	228
Maple [B] (verified)	229
Fricas [A] (verification not implemented)	229
Sympy [A] (verification not implemented)	230
Maxima [B] (verification not implemented)	230
Giac [A] (verification not implemented)	231
Mupad [B] (verification not implemented)	231
Reduce [B] (verification not implemented)	231

### Optimal result

Integrand size = 19, antiderivative size = 40

$$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{\sqrt{d+ex^2}}{\sqrt{e}} + x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

output `-(e*x^2+d)^(1/2)/e^(1/2)+x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{\sqrt{d+ex^2}}{\sqrt{e}} + x \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `-(Sqrt[d + e*x^2]/Sqrt[e]) + x*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]`



**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {6771, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx$$

$$\downarrow 6771$$

$$x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \sqrt{e} \int \frac{x}{\sqrt{ex^2+d}} dx$$

$$\downarrow 241$$

$$x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d+ex^2}}{\sqrt{e}}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]], x]`

output `-(Sqrt[d + e*x^2]/Sqrt[e]) + x*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]`

**Defintions of rubi rules used**

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 6771 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]], x_Symbol] :> Simp[x*ArcTanh[(c*x)/Sqrt[a + b*x^2]], x] - Simp[c Int[x/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c}, x] && EqQ[b, c^2]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs.  $2(32) = 64$ .

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.90

method	result	size
default	$x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) + \frac{e^{\frac{3}{2}}\left(\frac{x^2\sqrt{ex^2+d}}{3e} - \frac{2d\sqrt{ex^2+d}}{3e^2}\right)}{d} - \frac{(ex^2+d)^{\frac{3}{2}}}{3\sqrt{e}d}$	76
parts	$x \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) + \frac{e^{\frac{3}{2}}\left(\frac{x^2\sqrt{ex^2+d}}{3e} - \frac{2d\sqrt{ex^2+d}}{3e^2}\right)}{d} - \frac{(ex^2+d)^{\frac{3}{2}}}{3\sqrt{e}d}$	76

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output `x*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+e^(3/2)/d*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1/2))-1/3/e^(1/2)/d*(e*x^2+d)^(3/2)`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.28

$$\int \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \frac{ex \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 2\sqrt{ex^2+d}\sqrt{e}}{2e}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `1/2*(e*x*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - 2*sqrt(e*x^2 + d)*sqrt(e))/e`

**Sympy [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \begin{cases} x \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d+ex^2}}{\sqrt{e}} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((x*atanh(sqrt(e)*x/sqrt(d + e*x**2)) - sqrt(d + e*x**2)/sqrt(e), Ne(e, 0)), (0, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.62

$$\int \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = x \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) - \frac{(ex^2+d)^{\frac{3}{2}}}{3d\sqrt{e}} + \frac{(ex^2+d)^{\frac{3}{2}} - 3\sqrt{ex^2+dd}}{3d\sqrt{e}}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `x*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)) - 1/3*(e*x^2 + d)^(3/2)/(d*sqrt(e)) + 1/3*((e*x^2 + d)^(3/2) - 3*sqrt(e*x^2 + d)*d)/(d*sqrt(e))`

**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.48

$$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{2} x \log\left(-\frac{\frac{\sqrt{ex}}{\sqrt{ex^2+d}}+1}{\frac{\sqrt{ex}}{\sqrt{ex^2+d}}-1}\right) - \frac{\sqrt{e^2x^2+de}}{e}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`output `1/2*x*log(-(sqrt(e)*x/sqrt(e*x^2 + d) + 1)/(sqrt(e)*x/sqrt(e*x^2 + d) - 1)) - sqrt(e^2*x^2 + d*e)/e`**Mupad [B] (verification not implemented)**

Time = 2.98 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.80

$$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = x \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) - \frac{\sqrt{ex^2+d}}{\sqrt{e}}$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`output `x*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)) - (d + e*x^2)^(1/2)/e^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.52

$$\int \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+ex^2}}\right) ex - \sqrt{e}\sqrt{ex^2+d}}{e}$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`output `(atanh((sqrt(e)*sqrt(d + e*x**2)*x + e*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e*x - sqrt(e)*sqrt(d + e*x**2))/e`

**3.13** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

Optimal result	232
Mathematica [A] (verified)	232
Rubi [A] (verified)	233
Maple [A] (verified)	234
Fricas [B] (verification not implemented)	235
Sympy [F]	235
Maxima [F]	236
Giac [F(-1)]	236
Mupad [F(-1)]	236
Reduce [F]	237

**Optimal result**

Integrand size = 23, antiderivative size = 55

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} - \frac{\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{\sqrt{d}}$$

output

$$-\operatorname{arctanh}\left(\frac{e^{1/2}x}{(e x^2+d)^{1/2}}\right)/x - e^{1/2}\operatorname{arctanh}\left(\frac{(e x^2+d)^{1/2}}{d^{1/2}}\right)/d^{1/2}$$

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} + \frac{\sqrt{e}\left(\log(x) - \log\left(d + \sqrt{d}\sqrt{d+ex^2}\right)\right)}{\sqrt{d}}$$

input

`Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^2,x]`

output

$$-\left(\frac{\operatorname{ArcTanh}\left[\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right]}{x}\right) + \frac{\sqrt{e}\left(\log[x] - \log\left[d + \sqrt{d}\sqrt{d+ex^2}\right]\right)}{\sqrt{d}}$$

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$ , Rules used = {6775, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx \\
 & \quad \downarrow \text{6775} \\
 & \sqrt{e} \int \frac{1}{x\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2}\sqrt{e} \int \frac{1}{x^2\sqrt{ex^2+d}} dx^2 - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} \\
 & \quad \downarrow \text{73} \\
 & \frac{\int \frac{1}{\frac{x^4}{e} - \frac{d}{e}} d\sqrt{ex^2+d}}{\sqrt{e}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} \\
 & \quad \downarrow \text{221} \\
 & -\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x} - \frac{\sqrt{e}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{\sqrt{d}}
 \end{aligned}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^2,x]`

output `-(ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x) - (Sqrt[e]*ArcTanh[Sqrt[d + e*x^2]/Sqrt[d]])/Sqrt[d]`

## Definitions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[  
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +  
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt  
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL  
 inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x  
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int  
 t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I  
 ntegerQ[(m - 1)/2]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_  
 Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))),  
 x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; Fre  
 eQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.53

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x} - \frac{\sqrt{e}\sqrt{ex^2+d}}{d} + \frac{\sqrt{e}\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{d}$	84
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x} - \frac{\sqrt{e}\sqrt{ex^2+d}}{d} + \frac{\sqrt{e}\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{d}$	84

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x,method=_RETURNVERBOSE)`

output `-arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x-e^(1/2)/d*(e*x^2+d)^(1/2)+e^(1/2)/d*  
 ((e*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 135 vs.  $2(43) = 86$ .

Time = 0.14 (sec) , antiderivative size = 273, normalized size of antiderivative = 4.96

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

$$= \frac{\left[ x\sqrt{\frac{e}{d}} \log\left(-\frac{e^2x^2 - 2\sqrt{ex^2+d}d\sqrt{e}\sqrt{\frac{e}{d}} + 2de}{x^2}\right) + (x-1) \log\left(\frac{2ex^2 + 2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - x \log\left(\frac{ex + \sqrt{ex^2+d}\sqrt{e}}{x}\right) + x \log\left(\frac{ex - \sqrt{ex^2+d}\sqrt{e}}{x}\right) \right]}{2x}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="fricas")`

output `[1/2*(x*sqrt(e/d)*log(-(e^2*x^2 - 2*sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(e/d) + 2*d*e)/x^2) + (x - 1)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - x*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) + x*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x))/x, 1/2*(2*x*sqrt(-e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(-e/d)/(e^2*x^2 + d*e)) + (x - 1)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - x*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) + x*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x))/x]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**2,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**2, x)`



**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="maxima")`

output `d*sqrt(e)*integrate(-sqrt(e*x^2 + d)/(e^2*x^5 + d*e*x^3 - (e*x^3 + d*x)*(e*x^2 + d)), x) - 1/2*(log(sqrt(e)*x + sqrt(e*x^2 + d)) - log(-sqrt(e)*x + sqrt(e*x^2 + d)))/x`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^2,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^2, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**2,x)`

**3.14** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

Optimal result	238
Mathematica [A] (verified)	238
Rubi [A] (verified)	239
Maple [A] (verified)	241
Fricas [B] (verification not implemented)	241
Sympy [F]	242
Maxima [F]	242
Giac [F(-1)]	243
Mupad [F(-1)]	243
Reduce [F]	243

**Optimal result**

Integrand size = 23, antiderivative size = 85

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{6dx^2} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3} + \frac{e^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{6d^{3/2}}$$

output `-1/6*e^(1/2)*(e*x^2+d)^(1/2)/d/x^2-1/3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3+1/6*e^(3/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2))/d^(3/2)`

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + \frac{\sqrt{ex}\left(\sqrt{d}\sqrt{d+ex^2}+ex^2\log(x)-ex^2\log\left(d+\sqrt{d}\sqrt{d+ex^2}\right)\right)}{d^{3/2}}}{6x^3}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^4,x]`

output

```
-1/6*(2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] + (Sqrt[e]*x*(Sqrt[d]*Sqrt[d
+ e*x^2] + e*x^2*Log[x] - e*x^2*Log[d + Sqrt[d]*Sqrt[d + e*x^2]]))/d^(3/2)
)/x^3
```

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {6775, 243, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

$$\downarrow 6775$$

$$\frac{1}{3}\sqrt{e} \int \frac{1}{x^3\sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

$$\downarrow 243$$

$$\frac{1}{6}\sqrt{e} \int \frac{1}{x^4\sqrt{ex^2+d}} dx^2 - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

$$\downarrow 52$$

$$\frac{1}{6}\sqrt{e} \left( -\frac{e \int \frac{1}{x^2\sqrt{ex^2+d}} dx^2}{2d} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

$$\downarrow 73$$

$$\frac{1}{6}\sqrt{e} \left( -\frac{\int \frac{1}{\frac{x^4}{e} - \frac{d}{e}} d\sqrt{ex^2+d}}{d} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

$$\downarrow 221$$

$$\frac{1}{6}\sqrt{e} \left( \frac{e \operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^4,x]`

output `-1/3*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^3 + (Sqrt[e]*(-(Sqrt[d + e*x^2]/(d*x^2)) + (e*ArcTanh[Sqrt[d + e*x^2]/Sqrt[d]])/d^(3/2)))/6`

### Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

### Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.45

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{3x^3} + \frac{e^{\frac{3}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{3d^{\frac{3}{2}}} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{2d} \right)}{3d}$	123
parts	$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{3x^3} + \frac{e^{\frac{3}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{3d^{\frac{3}{2}}} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{2d} \right)}{3d}$	123

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^3+1/3*e^(3/2)/d^(3/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x)+1/3*e^(1/2)/d*(-1/2/d/x^2*(e*x^2+d)^(3/2)+1/2*e/d*((e*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x)))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 167 vs. 2(63) = 126.

Time = 0.14 (sec) , antiderivative size = 340, normalized size of antiderivative = 4.00

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

$$= \frac{\left[ ex^3 \sqrt{\frac{e}{d}} \log\left(-\frac{e^2x^2+2\sqrt{ex^2+dd}\sqrt{e}\sqrt{\frac{e}{d}+2de}}{x^2}\right) - 2dx^3 \log\left(\frac{ex+\sqrt{ex^2+d}\sqrt{e}}{x}\right) + 2dx^3 \log\left(\frac{ex-\sqrt{ex^2+d}\sqrt{e}}{x}\right) - 2\sqrt{ex^2+d}\sqrt{e} \right]}{12dx^3}$$

$$- \frac{ex^3 \sqrt{-\frac{e}{d}} \arctan\left(\frac{\sqrt{ex^2+dd}\sqrt{e}\sqrt{-\frac{e}{d}}}{e^2x^2+de}\right) + dx^3 \log\left(\frac{ex+\sqrt{ex^2+d}\sqrt{e}}{x}\right) - dx^3 \log\left(\frac{ex-\sqrt{ex^2+d}\sqrt{e}}{x}\right) + \sqrt{ex^2+d}\sqrt{e}}{6dx^3}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="fricas")`

output

```
[1/12*(e*x^3*sqrt(e/d)*log(-(e^2*x^2 + 2*sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(e/d) + 2*d*e)/x^2) - 2*d*x^3*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) + 2*d*x^3*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x) - 2*sqrt(e*x^2 + d)*sqrt(e)*x + 2*(d*x^3 - d)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d*x^3), -1/6*(e*x^2*sqrt(-e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(-e/d)/(e^2*x^2 + d*e)) + d*x^3*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) - d*x^3*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x) + sqrt(e*x^2 + d)*sqrt(e)*x - (d*x^3 - d)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d*x^3)]
```

### Sympy [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

input

```
integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**4,x)
```

output

```
Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**4, x)
```

### Maxima [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input

```
integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="maxima")
```

output

```
d*sqrt(e)*integrate(-1/3*sqrt(e*x^2 + d)/(e^2*x^7 + d*e*x^5 - (e*x^5 + d*x^3)*(e*x^2 + d)), x) - 1/6*(log(sqrt(e)*x + sqrt(e*x^2 + d)) - log(-sqrt(e)*x + sqrt(e*x^2 + d)))/x^3
```

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^4,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^4, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**4,x)`



**3.15** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

Optimal result	244
Mathematica [A] (verified)	244
Rubi [A] (verified)	245
Maple [B] (verified)	247
Fricas [B] (verification not implemented)	248
Sympy [F]	248
Maxima [F]	249
Giac [F(-1)]	249
Mupad [F(-1)]	249
Reduce [F]	250

**Optimal result**

Integrand size = 23, antiderivative size = 111

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = -\frac{\sqrt{e}\sqrt{d+ex^2}}{20dx^4} + \frac{3e^{3/2}\sqrt{d+ex^2}}{40d^2x^2} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5} - \frac{3e^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{40d^{5/2}}$$

output `-1/20*e^(1/2)*(e*x^2+d)^(1/2)/d/x^4+3/40*e^(3/2)*(e*x^2+d)^(1/2)/d^2/x^2-1/5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^5-3/40*e^(5/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2))/d^(5/2)`

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \frac{-8\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + \frac{\sqrt{ex}\left(\sqrt{d}\sqrt{d+ex^2}(-2d+3ex^2)+3e^2x^4\log(x)-3e^2x^4\log\left(d+\sqrt{d}\sqrt{d+ex^2}\right)\right)}{d^{5/2}}}{40x^5}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^6,x]`

output `(-8*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] + (Sqrt[e]*x*(Sqrt[d]*Sqrt[d + e*x^2]*(-2*d + 3*e*x^2) + 3*e^2*x^4*Log[x] - 3*e^2*x^4*Log[d + Sqrt[d]*Sqrt[d + e*x^2]]))/d^(5/2))/(40*x^5)`

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6775, 243, 52, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx \\
 & \quad \downarrow \text{6775} \\
 & \frac{1}{5} \sqrt{e} \int \frac{1}{x^5 \sqrt{ex^2+d}} dx - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{10} \sqrt{e} \int \frac{1}{x^6 \sqrt{ex^2+d}} dx^2 - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{10} \sqrt{e} \left( -\frac{3e \int \frac{1}{x^4 \sqrt{ex^2+d}} dx^2}{4d} - \frac{\sqrt{d+ex^2}}{2dx^4} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{10} \sqrt{e} \left( -\frac{3e \left( -\frac{e \int \frac{1}{x^2 \sqrt{ex^2+d}} dx^2}{2d} - \frac{\sqrt{d+ex^2}}{dx^2} \right)}{4d} - \frac{\sqrt{d+ex^2}}{2dx^4} \right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{10}\sqrt{e}\left(-\frac{3e\left(-\frac{\int\frac{1}{e}-\frac{d}{e}d\sqrt{ex^2+d}}{d}-\frac{\sqrt{d+ex^2}}{dx^2}\right)}{4d}-\frac{\sqrt{d+ex^2}}{2dx^4}\right)-\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5}$$

↓ 221

$$\frac{1}{10}\sqrt{e}\left(-\frac{3e\left(\frac{e\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{d^{3/2}}-\frac{\sqrt{d+ex^2}}{dx^2}\right)}{4d}-\frac{\sqrt{d+ex^2}}{2dx^4}\right)-\frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^5}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^6,x]`

output `-1/5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^5 + (Sqrt[e]*(-1/2*Sqrt[d + e*x^2]/(d*x^4) - (3*e*(-(Sqrt[d + e*x^2]/(d*x^2)) + (e*ArcTanh[Sqrt[d + e*x^2]/Sqrt[d]))/d^(3/2)))/(4*d))/10`

### Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243  $\text{Int}[(x_)^{(m\_)}*((a_) + (b\_)*(x_)^2)^{(p\_)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m-1)/2)*(a+b*x)^p}, x], x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

rule 6775  $\text{Int}[\text{ArcTanh}[(c\_)*(x_)]/\text{Sqrt}[(a_) + (b\_)*(x_)^2]]*((d\_)*(x_)^{(m\_)}), x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTanh}[c*x]/\text{Sqrt}[a+b*x^2])/(d*(m+1)), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a+b*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b, c^2] \ \&\& \ \text{NeQ}[m, -1]$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs.  $2(83) = 166$ .

Time = 0.04 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.54

method	result
default	$-\frac{\text{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{5x^5} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{e \left( \frac{\sqrt{ex^2+d}-\sqrt{d} \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{2d}\right)\right)}{4d} \right)}{5d} - \frac{e^{\frac{3}{2}} \left( -\frac{\sqrt{ex^2+d}}{2dx^2} + \dots \right)}{5d}$
parts	$-\frac{\text{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{5x^5} + \frac{\sqrt{e} \left( -\frac{(ex^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{e \left( \frac{\sqrt{ex^2+d}-\sqrt{d} \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{2d}\right)\right)}{4d} \right)}{5d} - \frac{e^{\frac{3}{2}} \left( -\frac{\sqrt{ex^2+d}}{2dx^2} + \dots \right)}{5d}$

input  $\text{int}(\text{arctanh}(e^{(1/2)*x}/(e*x^2+d)^{(1/2)})/x^6, x, \text{method}=\_RETURNVERBOSE)$

output  $-1/5*\text{arctanh}(e^{(1/2)*x}/(e*x^2+d)^{(1/2)})/x^5 + 1/5*e^{(1/2)}/d*(-1/4/d/x^4*(e*x^2+d)^{(3/2)} - 1/4*e/d*(-1/2/d/x^2*(e*x^2+d)^{(3/2)} + 1/2*e/d*((e*x^2+d)^{(1/2)} - d^{(1/2)}*\ln((2*d+2*d^{(1/2)}*(e*x^2+d)^{(1/2)})/x))) - 1/5*e^{(3/2)}/d*(-1/2/d/x^2*(e*x^2+d)^{(1/2)} + 1/2*e/d^{(3/2)}*\ln((2*d+2*d^{(1/2)}*(e*x^2+d)^{(1/2)})/x))$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 189 vs.  $2(83) = 166$ .

Time = 0.14 (sec) , antiderivative size = 383, normalized size of antiderivative = 3.45

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

$$= \frac{\left[ 3e^2x^5\sqrt{\frac{e}{d}}\log\left(-\frac{e^2x^2-2\sqrt{ex^2+d}d\sqrt{e}\sqrt{\frac{e}{d}}+2de}{x^2}\right) - 8d^2x^5\log\left(\frac{ex+\sqrt{ex^2+d}\sqrt{e}}{x}\right) + 8d^2x^5\log\left(\frac{ex-\sqrt{ex^2+d}\sqrt{e}}{x}\right) + 2 \right]}{80d^2x^5}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="fricas")`

output `[1/80*(3*e^2*x^5*sqrt(e/d)*log(-(e^2*x^2 - 2*sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(e/d) + 2*d*e)/x^2) - 8*d^2*x^5*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) + 8*d^2*x^5*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x) + 2*(3*e*x^3 - 2*d*x)*sqrt(e*x^2 + d)*sqrt(e) + 8*(d^2*x^5 - d^2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d^2*x^5), 1/40*(3*e^2*x^5*sqrt(-e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(e)*sqrt(-e/d)/(e^2*x^2 + d*e)) - 4*d^2*x^5*log((e*x + sqrt(e*x^2 + d)*sqrt(e))/x) + 4*d^2*x^5*log((e*x - sqrt(e*x^2 + d)*sqrt(e))/x) + (3*e*x^3 - 2*d*x)*sqrt(e*x^2 + d)*sqrt(e) + 4*(d^2*x^5 - d^2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d^2*x^5)]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**6,x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**6, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="maxima")`

output `d*sqrt(e)*integrate(-1/5*sqrt(e*x^2 + d)/(e^2*x^9 + d*e*x^7 - (e*x^7 + d*x^5)*(e*x^2 + d)), x) - 1/10*(log(sqrt(e)*x + sqrt(e*x^2 + d)) - log(-sqrt(e)*x + sqrt(e*x^2 + d)))/x^5`

**Giac [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \text{Timed out}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="giac")`

output `Timed out`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^6,x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^6, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x)`

output `int(atanh((sqrt(e)*x)/sqrt(d + e*x**2))/x**6,x)`

### 3.16 $\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	251
Mathematica [C] (verified)	252
Rubi [A] (verified)	252
Maple [F]	255
Fricas [A] (verification not implemented)	255
Sympy [F(-1)]	256
Maxima [F]	256
Giac [F]	256
Mupad [F(-1)]	257
Reduce [F]	257

#### Optimal result

Integrand size = 25, antiderivative size = 196

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{60d^2\sqrt{x}\sqrt{d+ex^2}}{847e^{5/2}} + \frac{36dx^{5/2}\sqrt{d+ex^2}}{847e^{3/2}} - \frac{4x^{9/2}\sqrt{d+ex^2}}{121\sqrt{e}} + \frac{2}{11}x^{11/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + \frac{30d^{11/4}(\sqrt{d} + \sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{847e^{11/4}\sqrt{d+ex^2}}$$

output

```
-60/847*d^2*x^(1/2)*(e*x^2+d)^(1/2)/e^(5/2)+36/847*d*x^(5/2)*(e*x^2+d)^(1/2)/e^(3/2)-4/121*x^(9/2)*(e*x^2+d)^(1/2)/e^(1/2)+2/11*x^(11/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+30/847*d^(11/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(11/4)/(e*x^2+d)^(1/2)
```



**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.82

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2}{847} \sqrt{x} \left( -\frac{2\sqrt{d+ex^2}(15d^2 - 9dex^2 + 7e^2x^4)}{e^{5/2}} \right. \\ \left. + 77x^5 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \right) \\ + \frac{60d^{5/2} \sqrt{\frac{i\sqrt{d}}{\sqrt{e}}} \sqrt{1 + \frac{d}{ex^2}} x \operatorname{EllipticF}\left(i \operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{847e^2 \sqrt{d+ex^2}}$$

input

```
Integrate[x^(9/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
(2*Sqrt[x]*((-2*Sqrt[d + e*x^2]*(15*d^2 - 9*d*e*x^2 + 7*e^2*x^4))/e^(5/2)
+ 77*x^5*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]))/847 + (60*d^(5/2)*Sqrt[(I*
Sqrt[d])/Sqrt[e]]*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d]
)/Sqrt[e]]/Sqrt[x]], -1])/(847*e^2*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6775, 262, 262, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ \downarrow 6775 \\ \frac{2}{11} x^{11/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11} \sqrt{e} \int \frac{x^{11/2}}{\sqrt{ex^2+d}} dx$$

$$\begin{aligned}
 & \downarrow 262 \\
 & \frac{2}{11}x^{11/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{e}\left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d\int\frac{x^{7/2}}{\sqrt{ex^2+d}}dx}{11e}\right) \\
 & \downarrow 262 \\
 & \frac{2}{11}x^{11/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{e}\left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\int\frac{x^{3/2}}{\sqrt{ex^2+d}}dx}{7e}\right)}{11e}\right) \\
 & \downarrow 262 \\
 & \frac{2}{11}x^{11/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\
 & \left(\frac{2}{11}\sqrt{e}\left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d\int\frac{1}{\sqrt{x}\sqrt{ex^2+d}}dx}{3e}\right)}{7e}\right)}{11e}\right)\right) \\
 & \downarrow 266 \\
 & \frac{2}{11}x^{11/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\
 & \left(\frac{2}{11}\sqrt{e}\left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{3e}\right)}{7e}\right)}{11e}\right)\right) \\
 & \downarrow 761
 \end{aligned}$$

$$\frac{2}{11}\sqrt{e} \left( \frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d \left( \frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left( \frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)\right)}{3e^{5/4}\sqrt{d+ex^2}} \right)}{7e} \right)}{11e}$$

```
input Int[x^(9/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]], x]
```

```
output (2*x^(11/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/11 - (2*Sqrt[e]*((2*x^(9/2)*Sqrt[d + e*x^2])/(11*e) - (9*d*((2*x^(5/2)*Sqrt[d + e*x^2])/(7*e) - (5*d*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*e^(5/4)*Sqrt[d + e*x^2])))/(7*e)))/(11*e)))/11
```

**Defintions of rubi rules used**

```
rule 262 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 266 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 761

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

rule 6775

```
Int[ArcTanh[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]
```

**Maple [F]**

$$\int x^{\frac{9}{2}} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2 + d}}\right) dx$$

input

```
int(x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)
```

output

```
int(x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)
```

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.48

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d + e x^2}}\right) dx = \frac{77 e^3 x^{\frac{11}{2}} \log\left(\frac{2 e x^2 + 2 \sqrt{e x^2 + d} \sqrt{e} x + d}{d}\right) + 60 d^3 \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)}{847 e^3}$$

input

```
integrate(x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
1/847*(77*e^3*x^(11/2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 60*d^3*weierstrassPInverse(-4*d/e, 0, x) - 4*(7*e^2*x^4 - 9*d*e*x^2 + 15*d^2)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/e^3
```

**Sympy [F(-1)]**

Timed out.

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input `integrate(x**(9/2)*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output Timed out

**Maxima [F]**

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `1/11*x^(11/2)*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/11*x^(11/2)*log(-sqrt(e)*x + sqrt(e*x^2 + d)) - 2*d*sqrt(e)*integrate(-1/11*x*e^(1/2*log(e*x^2 + d) + 9/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

**Giac [F]**

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(9/2)*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(9/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(9/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^4 dx$$

input `int(x^(9/2)*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**4,x)`

### 3.17 $\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	258
Mathematica [C] (verified)	258
Rubi [A] (verified)	259
Maple [F]	261
Fricas [A] (verification not implemented)	262
Sympy [F]	262
Maxima [F]	262
Giac [F]	263
Mupad [F(-1)]	263
Reduce [F]	263

#### Optimal result

Integrand size = 25, antiderivative size = 168

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{20d\sqrt{x}\sqrt{d+ex^2}}{147e^{3/2}} - \frac{4x^{5/2}\sqrt{d+ex^2}}{49\sqrt{e}} + \frac{2}{7}x^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{10d^{7/4}(\sqrt{d} + \sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d} + \sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{147e^{7/4}\sqrt{d+ex^2}}$$

output

```
20/147*d*x^(1/2)*(e*x^2+d)^(1/2)/e^(3/2)-4/49*x^(5/2)*(e*x^2+d)^(1/2)/e^(1/2)+2/7*x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))-10/147*d^(7/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(7/4)/(e*x^2+d)^(1/2)
```

#### Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.88

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2}{147} \sqrt{x} \left( \frac{2(5d-3ex^2)\sqrt{d+ex^2}}{e^{3/2}} \right. \\ \left. + 21x^3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \right) \\ + \frac{20\sqrt{d} \left(\frac{i\sqrt{d}}{\sqrt{e}}\right)^{5/2} \sqrt{1+\frac{d}{ex^2}} x \operatorname{EllipticF}\left(i \operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{147\sqrt{d+ex^2}}$$

input

```
Integrate[x^(5/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]], x]
```

output

```
(2*Sqrt[x]*((2*(5*d - 3*e*x^2)*Sqrt[d + e*x^2])/e^(3/2) + 21*x^3*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]))/147 + (20*Sqrt[d]*((I*Sqrt[d])/Sqrt[e])^(5/2)*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(147*Sqrt[d + e*x^2]))
```

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6775, 262, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ \downarrow 6775 \\ \frac{2}{7} x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7} \sqrt{e} \int \frac{x^{7/2}}{\sqrt{ex^2+d}} dx \\ \downarrow 262$$



$$\begin{aligned}
& \frac{2}{7}x^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{e}\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\int\frac{x^{3/2}}{\sqrt{ex^2+d}}dx}{7e}\right) \\
& \quad \downarrow 262 \\
& \frac{2}{7}x^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{e}\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d\int\frac{1}{\sqrt{x}\sqrt{ex^2+d}}dx}{3e}\right)}{7e}\right) \\
& \quad \downarrow 266 \\
& \frac{2}{7}x^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{e}\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{3e}\right)}{7e}\right) \\
& \quad \downarrow 761 \\
& \frac{2}{7}x^{7/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{e}\left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d^{3/4}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{3e^{5/4}\sqrt{d+ex^2}}\right)}{7e}\right)
\end{aligned}$$

input `Int [x^(5/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output

```
(2*x^(7/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/7 - (2*Sqrt[e]*((2*x^(5/2)
)*Sqrt[d + e*x^2])/(7*e) - (5*d*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3
/4)*(Sqrt[d + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d + Sqrt[e]*x]^2)*Ellipt
icF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*e^(5/4)*Sqrt[d + e*x^2]
)))/(7*e)))/7
```

## Definitions of rubi rules used

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int x^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

**Fricas [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.49

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{21 e^2 x^{7/2} \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 20 d^2 \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)}{147 e^2}$$

input `integrate(x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `1/147*(21*e^2*x^(7/2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - 20*d^2*weierstrassPInverse(-4*d/e, 0, x) - 4*(3*e*x^2 - 5*d)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/e^2`

**Sympy [F]**

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

input `integrate(x**(5/2)*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Integral(x**(5/2)*atanh(sqrt(e)*x/sqrt(d + e*x**2)), x)`

**Maxima [F]**

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output  $1/7*x^{(7/2)}*\log(\sqrt{e}*x + \sqrt{e*x^2 + d}) - 1/7*x^{(7/2)}*\log(-\sqrt{e}*x + \sqrt{e*x^2 + d}) - 2*d*\sqrt{e}*integrate(-1/7*x*e^{(1/2)*\log(e*x^2 + d)} + 5/2*\log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x$

### Giac [F]

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(5/2)*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)), x)`

### Mupad [F(-1)]

Timed out.

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(5/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(5/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

### Reduce [F]

$$\int x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^2 dx$$

input `int(x^(5/2)*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**2,x)`

### 3.18 $\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	264
Mathematica [C] (verified)	265
Rubi [A] (verified)	265
Maple [F]	267
Fricas [A] (verification not implemented)	267
Sympy [F]	268
Maxima [F]	268
Giac [F]	268
Mupad [F(-1)]	269
Reduce [F]	269

#### Optimal result

Integrand size = 25, antiderivative size = 142

$$\begin{aligned} & \int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ &= -\frac{4\sqrt{x}\sqrt{d+ex^2}}{9\sqrt{e}} + \frac{2}{3}x^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \\ & \quad + \frac{2d^{3/4}(\sqrt{d} + \sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{9e^{3/4}\sqrt{d+ex^2}} \end{aligned}$$

output

```
-4/9*x^(1/2)*(e*x^2+d)^(1/2)/e^(1/2)+2/3*x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+2/9*d^(3/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(3/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.95

$$\begin{aligned} & \int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ &= \frac{2}{9} \sqrt{x} \left( -\frac{2\sqrt{d+ex^2}}{\sqrt{e}} + 3 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \right) \\ & \quad + \frac{4\sqrt{d} \sqrt{\frac{i\sqrt{d}}{\sqrt{e}}} \sqrt{1 + \frac{d}{ex^2}} x \operatorname{EllipticF}\left(i \operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{9\sqrt{d+ex^2}} \end{aligned}$$

input `Integrate[Sqrt[x]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]], x]`

output `(2*Sqrt[x]*((-2*Sqrt[d + e*x^2])/Sqrt[e] + 3*x*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]))/9 + (4*Sqrt[d]*Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(9*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6775, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{6775} \\ & \frac{2}{3} x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3} \sqrt{e} \int \frac{x^{3/2}}{\sqrt{ex^2+d}} dx \end{aligned}$$

$$\begin{aligned}
& \downarrow 262 \\
& \frac{2}{3}x^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3}\sqrt{e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d\int\frac{1}{\sqrt{x}\sqrt{ex^2+d}}dx}{3e}\right) \\
& \downarrow 266 \\
& \frac{2}{3}x^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3}\sqrt{e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{3e}\right) \\
& \downarrow 761 \\
& \frac{2}{3}x^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\
& \frac{2}{3}\sqrt{e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{3e^{5/4}\sqrt{d+ex^2}}\right)
\end{aligned}$$

input `Int[Sqrt[x]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*x^(3/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/3 - (2*Sqrt[e]*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*e^(5/4)*Sqrt[d + e*x^2])))`

### Defintions of rubi rules used

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 6775 `Int[ArcTanh[(c_.)*(x_)]/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2 + d}}\right) dx$$

input `int(x^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

## Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.49

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d + e x^2}}\right) dx$$

$$= \frac{3 e x^{\frac{3}{2}} \log\left(\frac{2 e x^2 + 2 \sqrt{e x^2 + d} \sqrt{e x + d}}{d}\right) + 4 d \operatorname{weierstrassPInverse}\left(-\frac{4 d}{e}, 0, x\right) - 4 \sqrt{e x^2 + d} \sqrt{e} \sqrt{x}}{9 e}$$

input `integrate(x^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `1/9*(3*e*x^(3/2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 4*d*weierstrassPInverse(-4*d/e, 0, x) - 4*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/e`



**Sympy [F]**

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

input `integrate(x**(1/2)*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Integral(sqrt(x)*atanh(sqrt(e)*x/sqrt(d + e*x**2)), x)`

**Maxima [F]**

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `-2*d*sqrt(e)*integrate(-1/3*x*e^(1/2*log(e*x^2 + d) + 1/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x) + 1/3*x^(3/2)*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/3*x^(3/2)*log(-sqrt(e)*x + sqrt(e*x^2 + d))`

**Giac [F]**

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(sqrt(x)*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(1/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(1/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int \sqrt{x} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(1/2)*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)),x)`

**3.19** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$$

Optimal result	270
Mathematica [C] (verified)	271
Rubi [A] (verified)	271
Maple [F]	273
Fricas [A] (verification not implemented)	273
Sympy [F]	273
Maxima [F]	274
Giac [F]	274
Mupad [F(-1)]	274
Reduce [F]	275

**Optimal result**

Integrand size = 25, antiderivative size = 113

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} + \frac{2\sqrt[4]{e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{d}\sqrt{d+ex^2}}$$

output

```
-2*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2)+2*e^(1/4)*(d^(1/2)+e^(1/2)*x)
*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)
*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(1/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} + \frac{4i\sqrt{e}\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{\sqrt{\frac{d}{e}}}{\sqrt{x}}\right), -1\right)}{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}\sqrt{d+ex^2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(3/2),x]`

output `(-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/Sqrt[x] + ((4*I)*Sqrt[e]*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$ , Rules used = {6775, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$$

↓ 6775

$$2\sqrt{e} \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

↓ 266

$$4\sqrt{e} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

↓ 761

$$\frac{2\sqrt[4]{e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\operatorname{arctan}\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{d}\sqrt{d+ex^2}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

input

```
Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(3/2),x]
```

output

```
(-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/Sqrt[x] + (2*e^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(d^(1/4)*Sqrt[d + e*x^2])
```

### Defintions of rubi rules used

rule 266

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 761

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

rule 6775

```
Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]
```

**Maple [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{x^{\frac{3}{2}}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.44

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+e x^2}}\right)}{x^{3/2}} dx = \frac{4x \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - \sqrt{x} \log\left(\frac{2e x^2 + 2\sqrt{e x^2+d}\sqrt{e}x+d}{d}\right)}{x}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="fricas")`

output `(4*x*weierstrassPInverse(-4*d/e, 0, x) - sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/x`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+e x^2}}\right)}{x^{3/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{d+e x^2}}\right)}{x^{\frac{3}{2}}} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(3/2),x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**(3/2), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{3/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(3/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 3/2*log(x))), x) - log(sqrt(e)*x + sqrt(e*x^2 + d))/sqrt(x) + log(-sqrt(e)*x + sqrt(e*x^2 + d))/sqrt(x)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{3/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{3/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(3/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(3/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**2,x)`



**3.20** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$$

Optimal result	276
Mathematica [C] (verified)	277
Rubi [A] (verified)	277
Maple [F]	279
Fricas [A] (verification not implemented)	279
Sympy [F]	280
Maxima [F]	280
Giac [F]	280
Mupad [F(-1)]	281
Reduce [F]	281

**Optimal result**

Integrand size = 25, antiderivative size = 145

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{15dx^{3/2}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} - \frac{2e^{5/4}\left(\sqrt{d} + \sqrt{ex}\right) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\operatorname{arctan}\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{15d^{5/4}\sqrt{d+ex^2}}$$

output

```
-4/15*e^(1/2)*(e*x^2+d)^(1/2)/d/x^(3/2)-2/5*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2)-2/15*e^(5/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(5/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = -\frac{2\left(2\sqrt{ex}\sqrt{d+ex^2} + 3d\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)\right)}{15dx^{5/2}} - \frac{4\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}e^2\sqrt{1+\frac{d}{ex^2}}x\operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{15d^{3/2}\sqrt{d+ex^2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(7/2),x]`

output `(-2*(2*Sqrt[e]*x*Sqrt[d + e*x^2] + 3*d*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]))/(15*d*x^(5/2)) - (4*Sqrt[(I*Sqrt[d])/Sqrt[e]]*e^2*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(15*d^(3/2)*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6775, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$$

↓ 6775

$$\frac{2}{5}\sqrt{e} \int \frac{1}{x^{5/2}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}}$$

↓ 264

$$\begin{aligned}
& \frac{2}{5}\sqrt{e}\left(-\frac{e\int\frac{1}{\sqrt{x}\sqrt{ex^2+d}}dx}{3d}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} \\
& \quad \downarrow 266 \\
& \frac{2}{5}\sqrt{e}\left(-\frac{2e\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{3d}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} \\
& \quad \downarrow 761 \\
& \frac{2}{5}\sqrt{e}\left(-\frac{e^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)- \\
& \quad \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}}
\end{aligned}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(7/2),x]`

output `(-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(5*x^(5/2)) + (2*Sqrt[e]*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2)) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]))/(3*d^(5/4)*Sqrt[d + e*x^2]))/5`

### Defintions of rubi rules used

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 6775 `Int[ArcTanh[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)`

## Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \frac{4ex^3 \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) + 4\sqrt{ex^2+d}\sqrt{ex}^{\frac{3}{2}} + 3d\sqrt{x} \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right)}{15dx^3}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x, algorithm="fricas")`

output `-1/15*(4*e*x^3*weierstrassPInverse(-4*d/e, 0, x) + 4*sqrt(e*x^2 + d)*sqrt(e)*x^(3/2) + 3*d*sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d*x^3)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(7/2), x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**(7/2), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2), x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-1/5*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(7/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 7/2*log(x))), x) - 1/5*log(sqrt(e)*x + sqrt(e*x^2 + d))/x^(5/2) + 1/5*log(-sqrt(e)*x + sqrt(e*x^2 + d))/x^(5/2)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2), x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(7/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(7/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(7/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**4,x)`

**3.21** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$$

Optimal result	282
Mathematica [C] (verified)	283
Rubi [A] (verified)	283
Maple [F]	285
Fricas [A] (verification not implemented)	286
Sympy [F(-1)]	286
Maxima [F]	286
Giac [F]	287
Mupad [F(-1)]	287
Reduce [F]	287

**Optimal result**

Integrand size = 25, antiderivative size = 173

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{63dx^{7/2}} + \frac{20e^{3/2}\sqrt{d+ex^2}}{189d^2x^{3/2}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} + \frac{10e^{9/4}\left(\sqrt{d} + \sqrt{ex}\right) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{189d^{9/4}\sqrt{d+ex^2}}$$

output

```
-4/63*e^(1/2)*(e*x^2+d)^(1/2)/d/x^(7/2)+20/189*e^(3/2)*(e*x^2+d)^(1/2)/d^2/x^(3/2)-2/9*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2)+10/189*e^(9/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(9/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \frac{4\sqrt{ex}\sqrt{d+ex^2}(-3d+5ex^2) - 42d^2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{189d^2x^{9/2}} + \frac{20\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}e^3\sqrt{1+\frac{d}{ex^2}}x\operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{189d^{5/2}\sqrt{d+ex^2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(11/2), x]`

output `(4*Sqrt[e]*x*Sqrt[d + e*x^2]*(-3*d + 5*e*x^2) - 42*d^2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(189*d^2*x^(9/2)) + (20*Sqrt[(I*Sqrt[d])/Sqrt[e]]*e^3*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(189*d^(5/2)*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6775, 264, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$$

↓ 6775

$$\frac{2}{9}\sqrt{e} \int \frac{1}{x^{9/2}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}}$$

↓ 264



$$\begin{aligned}
& \frac{2}{9}\sqrt{e}\left(-\frac{5e\int\frac{1}{x^{5/2}\sqrt{ex^2+d}}dx}{7d}-\frac{2\sqrt{d+ex^2}}{7dx^{7/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \quad \downarrow 264 \\
& \frac{2}{9}\sqrt{e}\left(-\frac{5e\left(-\frac{e\int\frac{1}{\sqrt{x}\sqrt{ex^2+d}}dx}{3d}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)}{7d}-\frac{2\sqrt{d+ex^2}}{7dx^{7/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \quad \downarrow 266 \\
& \frac{2}{9}\sqrt{e}\left(-\frac{5e\left(-\frac{2e\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{3d}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)}{7d}-\frac{2\sqrt{d+ex^2}}{7dx^{7/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \quad \downarrow 761 \\
& \frac{2}{9}\sqrt{e}\left(-\frac{5e\left(-\frac{e^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right),\frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}}-\frac{2\sqrt{d+ex^2}}{3dx^{3/2}}\right)}{7d}-\frac{2\sqrt{d+ex^2}}{7dx^{7/2}}\right)-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}}
\end{aligned}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(11/2),x]`

output `(-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(9*x^(9/2)) + (2*Sqrt[e]*((-2*Sqrt[d + e*x^2])/(7*d*x^(7/2)) - (5*e*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2)) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(3*d^(5/4)*Sqrt[d + e*x^2])))/(7*d)))/9`

## Definitions of rubi rules used

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^{\frac{11}{2}}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \frac{20 e^2 x^5 \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - 21 d^2 \sqrt{x} \log\left(\frac{2ex^2 + 2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) + 4}{189 d^2 x^5}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="fricas")`

output `1/189*(20*e^2*x^5*weierstrassPInverse(-4*d/e, 0, x) - 21*d^2*sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 4*(5*e*x^3 - 3*d*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/(d^2*x^5)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \text{Timed out}$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(11/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{11}{2}}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-1/9*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(11/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 11/2*log(x))), x) - 1/9*log(sqrt(e)*x + sqrt(e*x^2 + d))/x^(9/2) + 1/9*log(-sqrt(e)*x + sqrt(e*x^2 + d))/x^(9/2)`

### Giac [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{11/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(11/2), x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{11/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(11/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(11/2), x)`

### Reduce [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**6,x)`

**3.22** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$$

Optimal result	289
Mathematica [C] (verified)	290
Rubi [A] (verified)	290
Maple [F]	293
Fricas [A] (verification not implemented)	293
Sympy [F(-1)]	294
Maxima [F]	294
Giac [F]	295
Mupad [F(-1)]	295
Reduce [F]	295

**Optimal result**

Integrand size = 25, antiderivative size = 201

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{143dx^{11/2}} + \frac{36e^{3/2}\sqrt{d+ex^2}}{1001d^2x^{7/2}} - \frac{60e^{5/2}\sqrt{d+ex^2}}{1001d^3x^{3/2}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} - \frac{30e^{13/4}\left(\sqrt{d} + \sqrt{ex}\right) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{1001d^{13/4}\sqrt{d+ex^2}}$$

output

```
-4/143*e^(1/2)*(e*x^2+d)^(1/2)/d/x^(11/2)+36/1001*e^(3/2)*(e*x^2+d)^(1/2)/d^2/x^(7/2)-60/1001*e^(5/2)*(e*x^2+d)^(1/2)/d^3/x^(3/2)-2/13*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(13/2)-30/1001*e^(13/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(13/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.81

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \frac{2 \left( -\frac{2\sqrt{ex}\sqrt{d+ex^2}(7d^2-9dex^2+15e^2x^4)}{d^3} - 77\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{30\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}e^4\sqrt{1+\frac{d}{ex^2}}x^{15/2}\operatorname{EllipticF}\left[\operatorname{ArcSinh}\left[\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right], -1\right]}{d^{7/2}} \right)}{1001x^{13/2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(15/2), x]`

output `(2*((-2*Sqrt[e]*x*Sqrt[d + e*x^2]*(7*d^2 - 9*d*e*x^2 + 15*e^2*x^4))/d^3 - 77*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - (30*Sqrt[(I*Sqrt[d])/Sqrt[e]]*e^4*Sqrt[1 + d/(e*x^2)]*x^(15/2)*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(d^(7/2)*Sqrt[d + e*x^2])))/(1001*x^(13/2))`

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6775, 264, 264, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$$

↓ 6775

$$\frac{2}{13}\sqrt{e} \int \frac{1}{x^{13/2}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}}$$

↓ 264

$$\begin{aligned}
 & \frac{2}{13} \sqrt{e} \left( -\frac{9e \int \frac{1}{x^{9/2} \sqrt{ex^2+d}} dx}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} \\
 & \quad \downarrow 264 \\
 & \frac{2}{13} \sqrt{e} \left( -\frac{9e \left( -\frac{5e \int \frac{1}{x^{5/2} \sqrt{ex^2+d}} dx}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} \\
 & \quad \downarrow 264 \\
 & \frac{2}{13} \sqrt{e} \left( -\frac{9e \left( -\frac{5e \left( -\frac{e \int \frac{1}{\sqrt{x} \sqrt{ex^2+d}} dx}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \\
 & \quad \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} \\
 & \quad \downarrow 266 \\
 & \frac{2}{13} \sqrt{e} \left( -\frac{9e \left( -\frac{5e \left( -\frac{2e \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \\
 & \quad \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} \\
 & \quad \downarrow 761
 \end{aligned}$$



$$\frac{2}{13}\sqrt{e} \left( \frac{9e \left( \frac{5e \left( \frac{e^{3/4}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d} + \sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}}$$

```
input Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(15/2),x]
```

```
output (-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(13*x^(13/2)) + (2*Sqrt[e]*((-2*Sqrt[d + e*x^2])/(11*d*x^(11/2)) - (9*e*((-2*Sqrt[d + e*x^2])/(7*d*x^(7/2)) - (5*e*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2)) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*d^(5/4)*Sqrt[d + e*x^2])))/(7*d)))/(11*d))/13
```

**Defintions of rubi rules used**

```
rule 264 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

## Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx =$$

$$\frac{60 e^3 x^7 \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) + 77 d^3 \sqrt{x} \log\left(\frac{2ex^2 + 2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) + 4(15e^2x^5 - 9dex^3 + 7d^2x)}{1001 d^3 x^7}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="fricas")`

output `-1/1001*(60*e^3*x^7*weierstrassPInverse(-4*d/e, 0, x) + 77*d^3*sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 4*(15*e^2*x^5 - 9*d*e*x^3 + 7*d^2*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/(d^3*x^7)`

### Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \text{Timed out}$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(15/2),x)`

output Timed out

### Maxima [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-1/13*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(15/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 15/2*log(x))), x) - 1/13*log(sqrt(e)*x + sqrt(e*x^2 + d))/x^(13/2) + 1/13*log(-sqrt(e)*x + sqrt(e*x^2 + d))/x^(13/2)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(15/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(15/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(15/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^8} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**8,x)`

### 3.23 $\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	296
Mathematica [C] (verified)	297
Rubi [A] (verified)	297
Maple [F]	301
Fricas [A] (verification not implemented)	301
Sympy [F(-1)]	302
Maxima [F]	302
Giac [F]	303
Mupad [F(-1)]	303
Reduce [F]	303

#### Optimal result

Integrand size = 25, antiderivative size = 297

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{28dx^{3/2}\sqrt{d+ex^2}}{405e^{3/2}} - \frac{4x^{7/2}\sqrt{d+ex^2}}{81\sqrt{e}} - \frac{28d^2\sqrt{x}\sqrt{d+ex^2}}{135e^2(\sqrt{d}+\sqrt{ex})} + \frac{2}{9}x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + \frac{28d^{9/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{135e^{9/4}\sqrt{d+ex^2}} - \frac{14d^{9/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{135e^{9/4}\sqrt{d+ex^2}}$$

output

```
28/405*d*x^(3/2)*(e*x^2+d)^(1/2)/e^(3/2)-4/81*x^(7/2)*(e*x^2+d)^(1/2)/e^(1/2)-28/135*d^2*x^(1/2)*(e*x^2+d)^(1/2)/e^2/(d^(1/2)+e^(1/2)*x)+2/9*x^(9/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))+28/135*d^(9/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/e^(9/4)/(e*x^2+d)^(1/2)-14/135*d^(9/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(9/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.08 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.42

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{3/2}\left(14d^2 + 4dex^2 - 10e^2x^4 + 45e^{3/2}x^3\sqrt{d+ex^2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \dots\right)}{405e^{3/2}\sqrt{d+ex^2}}$$

input

```
Integrate[x^(7/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
(2*x^(3/2)*(14*d^2 + 4*d*e*x^2 - 10*e^2*x^4 + 45*e^(3/2)*x^3*Sqrt[d + e*x^2]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - 14*d^2*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)])/(405*e^(3/2)*Sqrt[d + e*x^2])
```

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {6775, 262, 262, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{6775} \\ & \frac{2}{9}x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{e} \int \frac{x^{9/2}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{2}{9}x^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{e} \left( \frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \int \frac{x^{5/2}}{\sqrt{ex^2+d}} dx}{9e} \right) \\ & \quad \downarrow \text{262} \end{aligned}$$

$$\frac{2}{9}x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{e}\left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{3d\int\frac{\sqrt{x}}{\sqrt{ex^2+d}}dx}{5e}\right)}{9e}\right)$$

↓ 266

$$\frac{2}{9}x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{e}\left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\int\frac{x}{\sqrt{ex^2+d}}d\sqrt{x}}{5e}\right)}{9e}\right)$$

↓ 834

$$\frac{2}{9}\sqrt{e}\left(\frac{2x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 7d\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\left(\frac{\sqrt{d}\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d}\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right)}{5e}\right)}{9e}\right)$$

↓ 27

$$\frac{2}{9}\sqrt{e}\left(\frac{2x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 7d\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\left(\frac{\sqrt{d}\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right)}{5e}\right)}{9e}\right)$$

↓ 761

$$\frac{2}{9}\sqrt{e} \left( \frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left( \frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left( \frac{\frac{2}{9}x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{4\sqrt{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int \frac{\sqrt{d-\sqrt{ex}}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right)}{9e} \right)$$

1510

$$\frac{2}{9}\sqrt{e} \left( \frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left( \frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left( \frac{\frac{2}{9}x^{9/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{4\sqrt{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{4\sqrt{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}}}{5e} \right)}{9e} \right)}{9e} \right)$$



input `Int[x^(7/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*x^(9/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/9 - (2*Sqrt[e]*((2*x^(7/2)*Sqrt[d + e*x^2])/(9*e) - (7*d*((2*x^(3/2)*Sqrt[d + e*x^2])/(5*e) - (6*d*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2])))/(5*e)))/(9*e))/9`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 834 `Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[1/q Int[1/Sqrt[a + b*x^4], x], x] - Simp[1/q Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 6775 `Int[ArcTanh[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int x^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

## Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.31

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \frac{45 e^2 x^{\frac{9}{2}} \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) + 84 d^2 \operatorname{weierstrassZeta}\left(-\frac{4d}{e}, 0, \operatorname{weierstrassZeta}\right)}{405 e^2}$$

input `integrate(x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output

```
1/405*(45*e^2*x^(9/2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) +
84*d^2*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) - 4*
(5*e*x^3 - 7*d*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/e^2
```

**Sympy [F(-1)]**

Timed out.

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input

```
integrate(x**(7/2)*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

Timed out

**Maxima [F]**

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima"
)
```

output

```
1/9*x^(9/2)*log(sqrt(e)*x + sqrt(e*x^2 + d)) - 1/9*x^(9/2)*log(-sqrt(e)*x
+ sqrt(e*x^2 + d)) - 2*d*sqrt(e)*integrate(-1/9*x*e^(1/2*log(e*x^2 + d) +
7/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)
```

**Giac [F]**

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(7/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(7/2)*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(7/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(7/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

**Reduce [F]**

$$\int x^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x^3 dx$$

input `int(x^(7/2)*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x**3,x)`

### 3.24 $\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	304
Mathematica [C] (verified)	305
Rubi [A] (verified)	305
Maple [F]	308
Fricas [A] (verification not implemented)	309
Sympy [F]	309
Maxima [F]	309
Giac [F]	310
Mupad [F(-1)]	310
Reduce [F]	310

#### Optimal result

Integrand size = 25, antiderivative size = 269

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{4x^{3/2}\sqrt{d+ex^2}}{25\sqrt{e}} + \frac{12d\sqrt{x}\sqrt{d+ex^2}}{25e(\sqrt{d}+\sqrt{ex})} + \frac{2}{5}x^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{12d^{5/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{25e^{5/4}\sqrt{d+ex^2}} + \frac{6d^{5/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{25e^{5/4}\sqrt{d+ex^2}}$$

output

```
-4/25*x^(3/2)*(e*x^2+d)^(1/2)/e^(1/2)+12/25*d*x^(1/2)*(e*x^2+d)^(1/2)/e/(d
^(1/2)+e^(1/2)*x)+2/5*x^(5/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))-12/25*d^(
5/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE
(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/e^(5/4)/(e*x^2+d)^(1/
2)+6/25*d^(5/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2
)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(5/4)/(
e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.41

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{3/2}\left(-2(d+ex^2) + 5\sqrt{ex}\sqrt{d+ex^2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) + 2d\sqrt{1+\frac{ex^2}{d}}\right)}{25\sqrt{e}\sqrt{d+ex^2}}$$

input `Integrate[x^(3/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*x^(3/2)*(-2*(d + e*x^2) + 5*Sqrt[e]*x*Sqrt[d + e*x^2]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] + 2*d*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)])/(25*Sqrt[e]*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6775, 262, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{6775} \\ & \frac{2}{5}x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e} \int \frac{x^{5/2}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{2}{5}x^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e} \left( \frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{3d \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{5e} \right) \\ & \quad \downarrow \text{266} \end{aligned}$$

$$\frac{2}{5}x^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e}\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\int\frac{x}{\sqrt{ex^2+d}}d\sqrt{x}}{5e}\right)$$

↓ 834

$$\frac{2}{5}x^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e}\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\left(\frac{\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right)}{5e}\right)$$

↓ 27

$$\frac{2}{5}x^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e}\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\left(\frac{\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right)}{5e}\right)$$

↓ 761

$$\frac{2}{5}x^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{e}\left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d\left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right),\frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right)}{5e}\right)$$

↓ 1510

$$\frac{2}{5}\sqrt{e} \left( \frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left( \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} \right) - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}}}{5e} \right)$$

input `Int[x^(3/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]], x]`

output `(2*x^(5/2)*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/5 - (2*Sqrt[e]*((2*x^(3/2)*Sqrt[d + e*x^2])/(5*e) - (6*d*(-((-((Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)^2)*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)^2)*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2])))/(5*e))/5`

**Defintions of rubi rules used**

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`



rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 834 `Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[1/q Int[1/Sqrt[a + b*x^4], x], x] - Simp[1/q Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple **[F]**

$$\int x^{\frac{3}{2}} \operatorname{arctanh} \left( \frac{\sqrt{e} x}{\sqrt{e x^2 + d}} \right) dx$$

input `int(x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.29

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \frac{5ex^{5/2} \log\left(\frac{2ex^2+2\sqrt{ex^2+d}\sqrt{ex+d}}{d}\right) - 4\sqrt{ex^2+d}\sqrt{ex}^{3/2} - 12d\operatorname{weierstrassZeta}\left(\frac{x}{\sqrt{d+ex^2}}\right)}{25e}$$

input `integrate(x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `1/25*(5*e*x^(5/2)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) - 4*sqrt(e*x^2 + d)*sqrt(e)*x^(3/2) - 12*d*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)))/e`

**Sympy [F]**

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx$$

input `integrate(x**(3/2)*atanh(e**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Integral(x**(3/2)*atanh(sqrt(e)*x/sqrt(d + e*x**2)), x)`

**Maxima [F]**

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output  $1/5*x^{(5/2)}*\log(\sqrt{e}*x + \sqrt{e*x^2 + d}) - 1/5*x^{(5/2)}*\log(-\sqrt{e}*x + \sqrt{e*x^2 + d}) - 2*d*\sqrt{e}*integrate(-1/5*x*e^{(1/2)*\log(e*x^2 + d)} + 3/2*\log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x$

### Giac [F]

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \operatorname{artanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(3/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(3/2)*arctanh(sqrt(e)*x/sqrt(e*x^2 + d)), x)`

### Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(3/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(3/2)*atanh((e^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

### Reduce [F]

$$\int x^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right) x dx$$

input `int(x^(3/2)*atanh(e^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2))*x,x)`

**3.25** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$$

Optimal result	311
Mathematica [C] (verified)	312
Rubi [A] (verified)	312
Maple [F]	315
Fricas [A] (verification not implemented)	315
Sympy [F]	316
Maxima [F]	316
Giac [F]	316
Mupad [F(-1)]	317
Reduce [F]	317

**Optimal result**

Integrand size = 25, antiderivative size = 232

$$\begin{aligned} & \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx \\ &= -\frac{4\sqrt{x}\sqrt{d+ex^2}}{\sqrt{d}+\sqrt{ex}} + 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) \\ & \quad + \frac{4\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} \\ & \quad - \frac{2\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} \end{aligned}$$

output

```
-4*x^(1/2)*(e*x^2+d)^(1/2)/(d^(1/2)+e^(1/2)*x)+2*x^(1/2)*arctanh(e^(1/2)*x
/(e*x^2+d)^(1/2))+4*d^(1/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)
)*x)^2^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2)
)/e^(1/4)/(e*x^2+d)^(1/2)-2*d^(1/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)
)+e^(1/2)*x)^2^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/
2*2^(1/2))/e^(1/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.37

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \frac{4\sqrt{e}x^{3/2}\sqrt{1+\frac{ex^2}{d}}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{ex^2}{d}\right)}{3\sqrt{d+ex^2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/Sqrt[x], x]`

output `2*Sqrt[x]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - (4*Sqrt[e]*x^(3/2)*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)]/(3*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6775, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx \\ & \quad \downarrow \text{6775} \\ & 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 2\sqrt{e} \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{266} \\ & 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{e} \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 834 \\
& 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{e}\left(\frac{\sqrt{d}\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d}\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right) \\
& \downarrow 27 \\
& 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{e}\left(\frac{\sqrt{d}\int\frac{1}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right) \\
& \downarrow 761 \\
& 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\
& 4\sqrt{e}\left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int\frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}}d\sqrt{x}}{\sqrt{e}}\right) \\
& \downarrow 1510 \\
& 2\sqrt{x}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - \\
& 4\sqrt{e}\left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}}\right)
\end{aligned}$$

input `Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/Sqrt[x],x]`

output `2*Sqrt[x]*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - 4*Sqrt[e]*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2]))`

## Definitions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 266  $\text{Int}[((c_*)(x_))^{(m_)}*((a_)+(b_*)(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a+b*(x^{(2*k)/c^2)})^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 761  $\text{Int}[1/\text{Sqrt}[(a_)+(b_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1+q^2*x^2)*(\text{Sqrt}[a+b*x^4]/(a*(1+q^2*x^2)^2)]/(2*q*\text{Sqrt}[a+b*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 834  $\text{Int}[(x_)^2/\text{Sqrt}[(a_)+(b_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 2]\}, \text{Simp}[1/q \text{ Int}[1/\text{Sqrt}[a+b*x^4], x], x] - \text{Simp}[1/q \text{ Int}[(1-q*x^2)/\text{Sqrt}[a+b*x^4], x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 1510  $\text{Int}[((d_)+(e_*)(x_)^2)/\text{Sqrt}[(a_)+(c_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a+c*x^4]/(a*(1+q^2*x^2))), x] + \text{Simp}[d*(1+q^2*x^2)*(\text{Sqrt}[a+c*x^4]/(a*(1+q^2*x^2)^2)]/(q*\text{Sqrt}[a+c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e+d*q^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$
- rule 6775  $\text{Int}[\text{ArcTanh}[(c_*)(x_)/\text{Sqrt}[(a_)+(b_*)(x_)^2]]*((d_*)(x_))^{(m_)}, x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTanh}[(c*x)/\text{Sqrt}[a+b*x^2]]/(d*(m+1))), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a+b*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b, c^2] \ \&\& \ \text{NeQ}[m, -1]$

**Maple [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.22

$$\begin{aligned} & \int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx \\ &= \sqrt{x} \log\left(\frac{2ex^2 + 2\sqrt{ex^2+d}\sqrt{e}x + d}{d}\right) \\ & \quad + 4 \operatorname{weierstrassZeta}\left(-\frac{4d}{e}, 0, \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right) \end{aligned}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="fricas")`

output `sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 4*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x))`



**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(1/2),x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/sqrt(x), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="maxima")`

output `-2*d*sqrt(e)*integrate(sqrt(e*x^2 + d)*x/((e*x^2 + d)*e^(log(e*x^2 + d) + 1/2*log(x)) - (e^2*x^4 + d*e*x^2)*sqrt(x)), x) + sqrt(x)*log(sqrt(e)*x + sqrt(e*x^2 + d)) - sqrt(x)*log(-sqrt(e)*x + sqrt(e*x^2 + d))`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/sqrt(x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(1/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(1/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x,x)`

**3.26** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$$

Optimal result	318
Mathematica [C] (verified)	319
Rubi [A] (verified)	319
Maple [F]	322
Fricas [A] (verification not implemented)	323
Sympy [F]	323
Maxima [F]	324
Giac [F]	324
Mupad [F(-1)]	324
Reduce [F]	325

**Optimal result**

Integrand size = 25, antiderivative size = 272

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{3d\sqrt{x}} + \frac{4e\sqrt{x}\sqrt{d+ex^2}}{3d(\sqrt{d}+\sqrt{ex})}$$

$$-\frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} - \frac{4e^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{3d^{3/4}\sqrt{d+ex^2}}$$

$$+\frac{2e^{3/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{3d^{3/4}\sqrt{d+ex^2}}$$

output

```
-4/3*e^(1/2)*(e*x^2+d)^(1/2)/d/x^(1/2)+4/3*e*x^(1/2)*(e*x^2+d)^(1/2)/d/(d^(1/2)+e^(1/2)*x)-2/3*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2)-4/3*e^(3/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/d^(3/4)/(e*x^2+d)^(1/2)+2/3*e^(3/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(3/4)/(e*x^2+d)^(1/2)
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{3d\sqrt{x}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} + \frac{4e^{3/2}x^{3/2}\sqrt{1+\frac{ex^2}{d}}\operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{ex^2}{d}\right)}{9d\sqrt{d+ex^2}}$$

input `Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(5/2),x]`

output `(-4*Sqrt[e]*Sqrt[d + e*x^2])/(3*d*Sqrt[x]) - (2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(3*x^(3/2)) + (4*e^(3/2)*x^(3/2)*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -(e*x^2)/d])/(9*d*Sqrt[d + e*x^2])`

**Rubi [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6775, 264, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$$

↓ 6775

$$\frac{2}{3}\sqrt{e} \int \frac{1}{x^{3/2}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}}$$

↓ 264

$$\begin{aligned}
& \frac{2}{3}\sqrt{e} \left( \frac{e \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow 266 \\
& \frac{2}{3}\sqrt{e} \left( \frac{2e \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow 834 \\
& \frac{2}{3}\sqrt{e} \left( \frac{2e \left( \frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow 27 \\
& \frac{2}{3}\sqrt{e} \left( \frac{2e \left( \frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow 761 \\
& \frac{2}{3}\sqrt{e} \left( \frac{2e \left( \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right) - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}}}{2e^{3/4}\sqrt{d+ex^2}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \\
& \quad \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow 1510
\end{aligned}$$

$$\frac{2}{3}\sqrt{e} \left( \frac{2e \left( \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right) \middle| \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} \right)}{d} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \right)$$

```
input Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(5/2),x]
```

```
output (-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(3*x^(3/2)) + (2*Sqrt[e]*((-2*Sqrt[d + e*x^2])/(d*Sqrt[x]) + (2*e*(-((-((Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2])))/d)/3
```

**Defintions of rubi rules used**

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 264 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 834 `Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[1/q Int[1/Sqrt[a + b*x^4], x], x] - Simp[1/q Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 6775 `Int[ArcTanh[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTanh[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b, c^2] && NeQ[m, -1]`

## Maple [F]

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{x^{\frac{5}{2}}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.30

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \frac{4ex^2 \operatorname{weierstrassZeta}\left(-\frac{4d}{e}, 0, \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right) + 4\sqrt{ex^2+d}\sqrt{ex}^{\frac{3}{2}} + d\sqrt{x} \log\left(\frac{2ex^2+2\sqrt{ex}}{d}\right)}{3dx^2}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="fricas")`

output `-1/3*(4*e*x^2*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) + 4*sqrt(e*x^2 + d)*sqrt(e)*x^(3/2) + d*sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d))/(d*x^2)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{\frac{5}{2}}} dx$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(5/2),x)`

output `Integral(atanh(sqrt(e)*x/sqrt(d + e*x**2))/x**(5/2), x)`



**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-1/3*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(5/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 5/2*log(x))), x) - 1/3*log(sqrt(e)*x + sqrt(e*x^2 + d))/x^(3/2) + 1/3*log(-sqrt(e)*x + sqrt(e*x^2 + d))/x^(3/2)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(5/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(5/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(5/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^3} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**3,x)`

**3.27** 
$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$$

Optimal result	326
Mathematica [C] (verified)	327
Rubi [A] (verified)	327
Maple [F]	332
Fricas [A] (verification not implemented)	332
Sympy [F(-1)]	332
Maxima [F]	333
Giac [F]	333
Mupad [F(-1)]	333
Reduce [F]	334

**Optimal result**

Integrand size = 25, antiderivative size = 302

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = -\frac{4\sqrt{e}\sqrt{d+ex^2}}{35dx^{5/2}} + \frac{12e^{3/2}\sqrt{d+ex^2}}{35d^2\sqrt{x}}$$

$$-\frac{12e^2\sqrt{x}\sqrt{d+ex^2}}{35d^2(\sqrt{d}+\sqrt{ex})} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

$$+ \frac{12e^{7/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{35d^{7/4}\sqrt{d+ex^2}}$$

$$- \frac{6e^{7/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{35d^{7/4}\sqrt{d+ex^2}}$$

output

```
-4/35*e^(1/2)*(e*x^2+d)^(1/2)/d/x^(5/2)+12/35*e^(3/2)*(e*x^2+d)^(1/2)/d^2/
x^(1/2)-12/35*e^2*x^(1/2)*(e*x^2+d)^(1/2)/d^2/(d^(1/2)+e^(1/2)*x)-2/7*arct
anh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2)+12/35*e^(7/4)*(d^(1/2)+e^(1/2)*x)*
(e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1
/2)/d^(1/4))),1/2*2^(1/2))/d^(7/4)/(e*x^2+d)^(1/2)-6/35*e^(7/4)*(d^(1/2)+e
^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan
(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(7/4)/(e*x^2+d)^(1/2)
```

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \frac{4\sqrt{ex}(-d^2 + 2dex^2 + 3e^2x^4) - 10d^2\sqrt{d+ex^2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right) - 4e^{5/2}x^5\sqrt{1 + \frac{ex}{d}}}{35d^2x^{7/2}\sqrt{d+ex^2}}$$

input

```
Integrate[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(9/2),x]
```

output

```
(4*Sqrt[e]*x*(-d^2 + 2*d*e*x^2 + 3*e^2*x^4) - 10*d^2*Sqrt[d + e*x^2]*ArcTa
nh[(Sqrt[e]*x)/Sqrt[d + e*x^2]] - 4*e^(5/2)*x^5*Sqrt[1 + (e*x^2)/d]*Hyperg
eometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)]/(35*d^2*x^(7/2)*Sqrt[d + e*x^2])
```

### Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$ , Rules used = {6775, 264, 264, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$$

↓ 6775

$$\begin{aligned}
& \frac{2}{7}\sqrt{e} \int \frac{1}{x^{7/2}\sqrt{ex^2+d}} dx - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
& \quad \downarrow 264 \\
& \frac{2}{7}\sqrt{e} \left( -\frac{3e \int \frac{1}{x^{3/2}\sqrt{ex^2+d}} dx}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
& \quad \downarrow 264 \\
& \frac{2}{7}\sqrt{e} \left( -\frac{3e \left( \frac{e \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
& \quad \downarrow 266 \\
& \frac{2}{7}\sqrt{e} \left( -\frac{3e \left( \frac{2e \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
& \quad \downarrow 834 \\
& \frac{2}{7}\sqrt{e} \left( -\frac{3e \left( \frac{2e \left( \frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \\
& \quad \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
& \quad \downarrow 27
\end{aligned}$$

$$\frac{2}{7}\sqrt{e} \left( \frac{3e \left( \frac{2e \left( \frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

↓ 761

$$\frac{2}{7}\sqrt{e} \left( \frac{3e \left( \frac{2e \left( \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right) \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{2e^{3/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

↓ 1510

$$\left( \frac{2e}{3e} \left( \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}}) \sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}}) \sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right) \middle| \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} - \sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \right) - \frac{2\sqrt[7]{e}}{5d} \right) - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

input

```
Int[ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/x^(9/2),x]
```

output

```
(-2*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]]/(7*x^(7/2)) + (2*Sqrt[e]*((-2*Sqrt[d + e*x^2])/(5*d*x^(5/2)) - (3*e*(-2*Sqrt[d + e*x^2])/(d*Sqrt[x]) + (2*e*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(e^(1/4)*Sqrt[d + e*x^2])/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(2*e^(3/4)*Sqrt[d + e*x^2])))/d)/(5*d)))/7
```

## Definitions of rubi rules used

- rule 27  $\text{Int}[(a_*)(Fx_), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 264  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^2*(m+1))) \text{ Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 266  $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 761  $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^4]/(a*(1 + q^2*x^2)^2)]/(2*q*\text{Sqrt}[a + b*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 834  $\text{Int}[(x_)^2/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 2]\}, \text{Simp}[1/q \text{ Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Simp}[1/q \text{ Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 1510  $\text{Int}[((d_*) + (e_*)(x_)^2)/\text{Sqrt}[(a_*) + (c_*)(x_)^4], x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*\text{Sqrt}[a + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$
- rule 6775  $\text{Int}[\text{ArcTanh}[(c_*)(x_)/\text{Sqrt}[(a_*) + (b_*)(x_)^2]]*((d_*)(x_))^{(m_*)}, x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTanh}[(c*x)/\text{Sqrt}[a + b*x^2]]/(d*(m+1))), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b, c^2] \ \&\& \ \text{NeQ}[m, -1]$



**Maple [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{e x^2+d}}\right)}{x^{\frac{9}{2}}} dx$$

input `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

output `int(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.32

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \frac{12 e^2 x^4 \operatorname{weierstrassZeta}\left(-\frac{4d}{e}, 0, \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right) - 5 d^2 \sqrt{x} \log\left(\frac{\sqrt{d+ex^2} + \sqrt{e}x}{\sqrt{d+ex^2} - \sqrt{e}x}\right)}{35 d^2 x^4}$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="fricas")`

output `1/35*(12*e^2*x^4*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) - 5*d^2*sqrt(x)*log((2*e*x^2 + 2*sqrt(e*x^2 + d)*sqrt(e)*x + d)/d) + 4*(3*e*x^3 - d*x)*sqrt(e*x^2 + d)*sqrt(e)*sqrt(x))/(d^2*x^4)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \text{Timed out}$$

input `integrate(atanh(e**(1/2)*x/(e*x**2+d)**(1/2))/x**(9/2),x)`

output `Timed out`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="maxima")`

output `2*d*sqrt(e)*integrate(-1/7*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(9/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 9/2*log(x))), x) - 1/7*log(sqrt(e)*x + sqrt(e*x^2 + d))/x^(7/2) + 1/7*log(-sqrt(e)*x + sqrt(e*x^2 + d))/x^(7/2)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\operatorname{artanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `integrate(arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="giac")`

output `integrate(arctanh(sqrt(e)*x/sqrt(e*x^2 + d))/x^(9/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\operatorname{atanh}\left(\frac{\sqrt{ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(9/2),x)`

output `int(atanh((e^(1/2)*x)/(d + e*x^2)^(1/2))/x^(9/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{e}x}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\sqrt{x} \operatorname{atanh}\left(\frac{\sqrt{e}x}{\sqrt{ex^2+d}}\right)}{x^5} dx$$

input `int(atanh(e^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

output `int((sqrt(x)*atanh((sqrt(e)*x)/sqrt(d + e*x**2)))/x**5,x)`

## 3.28 $\int x^3 \operatorname{arctanh}(a + bx^4) dx$

Optimal result	335
Mathematica [A] (verified)	335
Rubi [A] (warning: unable to verify)	336
Maple [A] (verified)	337
Fricas [A] (verification not implemented)	338
Sympy [A] (verification not implemented)	338
Maxima [A] (verification not implemented)	339
Giac [B] (verification not implemented)	339
Mupad [B] (verification not implemented)	340
Reduce [B] (verification not implemented)	340

### Optimal result

Integrand size = 12, antiderivative size = 44

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx = \frac{(a + bx^4) \operatorname{arctanh}(a + bx^4)}{4b} + \frac{\log(1 - (a + bx^4)^2)}{8b}$$

output  $1/4*(b*x^4+a)*\operatorname{arctanh}(b*x^4+a)/b+1/8*\ln(1-(b*x^4+a)^2)/b$

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx = \frac{2(a + bx^4) \operatorname{arctanh}(a + bx^4) + \log(1 - (a + bx^4)^2)}{8b}$$

input  $\operatorname{Integrate}[x^3*\operatorname{ArcTanh}[a + b*x^4], x]$

output  $(2*(a + b*x^4)*\operatorname{ArcTanh}[a + b*x^4] + \operatorname{Log}[1 - (a + b*x^4)^2])/(8*b)$

**Rubi [A] (warning: unable to verify)**

Time = 0.32 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {7266, 6653, 6436, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}(a + bx^4) dx \\
 & \quad \downarrow \text{7266} \\
 & \frac{1}{4} \int \operatorname{arctanh}(bx^4 + a) dx^4 \\
 & \quad \downarrow \text{6653} \\
 & \frac{\int \operatorname{arctanh}(bx^4 + a) d(bx^4 + a)}{4b} \\
 & \quad \downarrow \text{6436} \\
 & \frac{(a + bx^4) \operatorname{arctanh}(a + bx^4) - \int \frac{bx^4 + a}{1 - x^8} d(bx^4 + a)}{4b} \\
 & \quad \downarrow \text{240} \\
 & \frac{(a + bx^4) \operatorname{arctanh}(a + bx^4) + \frac{1}{2} \log(1 - x^8)}{4b}
 \end{aligned}$$

input `Int[x^3*ArcTanh[a + b*x^4],x]`

output `((a + b*x^4)*ArcTanh[a + b*x^4] + Log[1 - x^8]/2)/(4*b)`

**Defintions of rubi rules used**

rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 6436 `Int[((a_) + ArcTanh[(c_)*(x_)^(n_)]*(b_))^(p_), x_Symbol] := Simp[x*(a + b*ArcTanh[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTanh[c*x^n])^(p - 1)/(1 - c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 6653 `Int[((a_) + ArcTanh[(c_) + (d_)*(x_)]*(b_))^(p_), x_Symbol] := Simp[1/d Subst[Int[(a + b*ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]`

rule 7266 `Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

**Maple [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{(bx^4+a) \operatorname{arctanh}(bx^4+a) + \frac{\ln(1-(bx^4+a)^2)}{2}}{4b}$
default	$\frac{(bx^4+a) \operatorname{arctanh}(bx^4+a) + \frac{\ln(1-(bx^4+a)^2)}{2}}{4b}$
parts	$\frac{x^4 \operatorname{arctanh}(bx^4+a)}{4} - b \left( \frac{(-a-1) \ln(bx^4+a+1)}{8b^2} + \frac{(a-1) \ln(bx^4+a-1)}{8b^2} \right)$
parallelrisch	$-\frac{\operatorname{arctanh}(bx^4+a)x^4b^2 - \operatorname{arctanh}(bx^4+a)ab - \ln(bx^4+a-1)b - \operatorname{arctanh}(bx^4+a)b}{4b^2}$
risch	$\frac{x^4 \ln(bx^4+a+1)}{8} - \frac{x^4 \ln(-bx^4-a+1)}{8} + \frac{\ln(bx^4+a+1)a}{8b} - \frac{\ln(-bx^4-a+1)a}{8b} + \frac{\ln(bx^4+a+1)}{8b} + \frac{\ln(-bx^4-a+1)}{8b}$

input `int(x^3*arctanh(b*x^4+a), x, method=_RETURNVERBOSE)`

output `1/4/b*((b*x^4+a)*arctanh(b*x^4+a)+1/2*ln(1-(b*x^4+a)^2))`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.34

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx$$

$$= \frac{bx^4 \log\left(-\frac{bx^4+a+1}{bx^4+a-1}\right) + (a+1) \log(bx^4+a+1) - (a-1) \log(bx^4+a-1)}{8b}$$

input `integrate(x^3*arctanh(b*x^4+a),x, algorithm="fricas")`

output `1/8*(b*x^4*log(-(b*x^4 + a + 1)/(b*x^4 + a - 1)) + (a + 1)*log(b*x^4 + a + 1) - (a - 1)*log(b*x^4 + a - 1))/b`

### Sympy [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.36

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx$$

$$= \begin{cases} \frac{a \operatorname{atanh}(a+bx^4)}{4b} + \frac{x^4 \operatorname{atanh}(a+bx^4)}{4} + \frac{\log(a+bx^4+1)}{4b} - \frac{\operatorname{atanh}(a+bx^4)}{4b} & \text{for } b \neq 0 \\ \frac{x^4 \operatorname{atanh}(a)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*atanh(b*x**4+a),x)`

output `Piecewise((a*atanh(a + b*x**4)/(4*b) + x**4*atanh(a + b*x**4)/4 + log(a + b*x**4 + 1)/(4*b) - atanh(a + b*x**4)/(4*b), Ne(b, 0)), (x**4*atanh(a)/4, True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx = \frac{2(bx^4 + a) \operatorname{artanh}(bx^4 + a) + \log\left(-(bx^4 + a)^2 + 1\right)}{8b}$$

input `integrate(x^3*arctanh(b*x^4+a),x, algorithm="maxima")`

output `1/8*(2*(b*x^4 + a)*arctanh(b*x^4 + a) + log(-(b*x^4 + a)^2 + 1))/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(40) = 80.

Time = 0.12 (sec) , antiderivative size = 223, normalized size of antiderivative = 5.07

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx$$

$$= \frac{1}{8} ((a+1)b - (a-1)b) \left( \frac{\log\left(\frac{|-bx^4-a-1|}{|bx^4+a-1|}\right)}{b^2} - \frac{\log\left(\left|-\frac{bx^4+a+1}{bx^4+a-1} + 1\right|\right)}{b^2} + \frac{\log\left(\frac{a - \frac{\left(\frac{bx^4+a+1}{bx^4+a-1}\right)(a-1) - a-1}{(bx^4+a+1)b - b}}{\frac{\left(\frac{bx^4+a+1}{bx^4+a-1}\right)(a-1) - a-1}{(bx^4+a+1)b - b}} - 1\right)}{b^2 \left(\frac{bx^4+a+1}{bx^4+a-1} - 1\right)} \right)$$

input `integrate(x^3*arctanh(b*x^4+a),x, algorithm="giac")`



output

```
1/8*((a + 1)*b - (a - 1)*b)*(log(abs(-b*x^4 - a - 1)/abs(b*x^4 + a - 1))/b
^2 - log(abs(-(b*x^4 + a + 1)/(b*x^4 + a - 1) + 1))/b^2 + log(-(a - ((b*x^
4 + a + 1)*(a - 1)/(b*x^4 + a - 1) - a - 1)*b/((b*x^4 + a + 1)*b/(b*x^4 +
a - 1) - b) + 1)/(a - ((b*x^4 + a + 1)*(a - 1)/(b*x^4 + a - 1) - a - 1)*b/
((b*x^4 + a + 1)*b/(b*x^4 + a - 1) - b) - 1))/(b^2*((b*x^4 + a + 1)/(b*x^4
+ a - 1) - 1)))
```

**Mupad [B] (verification not implemented)**

Time = 3.35 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.05

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx = \frac{\ln(bx^4 + a - 1)}{8b} - \frac{x^4 \ln(-bx^4 - a + 1)}{8} + \frac{\ln(bx^4 + a + 1)}{8b} + \frac{x^4 \ln(bx^4 + a + 1)}{8} - \frac{a \ln(bx^4 + a - 1)}{8b} + \frac{a \ln(bx^4 + a + 1)}{8b}$$

input

```
int(x^3*atanh(a + b*x^4),x)
```

output

```
log(a + b*x^4 - 1)/(8*b) - (x^4*log(1 - b*x^4 - a))/8 + log(a + b*x^4 + 1)
/(8*b) + (x^4*log(a + b*x^4 + 1))/8 - (a*log(a + b*x^4 - 1))/(8*b) + (a*log
(a + b*x^4 + 1))/(8*b)
```

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 377, normalized size of antiderivative = 8.57

$$\int x^3 \operatorname{arctanh}(a + bx^4) dx$$

$$= \frac{2 \operatorname{atanh}(bx^4 + a) a + 2 \operatorname{atanh}(bx^4 + a) bx^4 + \log\left(\frac{\sqrt{a-1} (a^2-1)^{\frac{1}{4}} - b^{\frac{1}{4}} \sqrt{2(a^2-1)^{\frac{1}{4}} a - 2(a^2-1)^{\frac{1}{4}} - \sqrt{-a+1} \sqrt{a^2-1} + \sqrt{-a+1}}}{\sqrt{a-1}}\right)}{\dots}$$

input

```
int(x^3*atanh(b*x^4+a),x)
```

output

```
(2*atanh(a + b*x**4)*a + 2*atanh(a + b*x**4)*b*x**4 + log((sqrt(a - 1)*(a**2 - 1)**(1/4) - b**(1/4)*sqrt(2*(a**2 - 1)**(1/4)*a - 2*(a**2 - 1)**(1/4) - sqrt(-a + 1)*sqrt(a**2 - 1) + sqrt(-a + 1)*a - sqrt(-a + 1))*x + sqrt(b)*sqrt(a - 1)*x**2)/sqrt(a - 1)) + log((sqrt(a - 1)*(a**2 - 1)**(1/4) - b**(1/4)*sqrt(2*(a**2 - 1)**(1/4)*a - 2*(a**2 - 1)**(1/4) + sqrt(-a + 1)*sqrt(a**2 - 1) - sqrt(-a + 1)*a + sqrt(-a + 1))*x + sqrt(b)*sqrt(a - 1)*x**2)/sqrt(a - 1)) + log((sqrt(a - 1)*(a**2 - 1)**(1/4) + b**(1/4)*sqrt(2*(a**2 - 1)**(1/4)*a - 2*(a**2 - 1)**(1/4) - sqrt(-a + 1)*sqrt(a**2 - 1) + sqrt(-a + 1)*a - sqrt(-a + 1))*x + sqrt(b)*sqrt(a - 1)*x**2)/sqrt(a - 1)) + log((sqrt(a - 1)*(a**2 - 1)**(1/4) + b**(1/4)*sqrt(2*(a**2 - 1)**(1/4)*a - 2*(a**2 - 1)**(1/4) + sqrt(-a + 1)*sqrt(a**2 - 1) - sqrt(-a + 1)*a + sqrt(-a + 1))*x + sqrt(b)*sqrt(a - 1)*x**2)/sqrt(a - 1)))/(8*b)
```

### 3.29 $\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx$

Optimal result	342
Mathematica [A] (verified)	342
Rubi [A] (warning: unable to verify)	343
Maple [B] (verified)	344
Fricas [B] (verification not implemented)	345
Sympy [F(-2)]	345
Maxima [A] (verification not implemented)	346
Giac [B] (verification not implemented)	346
Mupad [B] (verification not implemented)	347
Reduce [B] (verification not implemented)	347

#### Optimal result

Integrand size = 14, antiderivative size = 47

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \frac{(a + bx^n) \operatorname{arctanh}(a + bx^n)}{bn} + \frac{\log(1 - (a + bx^n)^2)}{2bn}$$

output  $(a+b*x^n)*\operatorname{arctanh}(a+b*x^n)/b/n+1/2*\ln(1-(a+b*x^n)^2)/b/n$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \frac{2(a + bx^n) \operatorname{arctanh}(a + bx^n) + \log(1 - (a + bx^n)^2)}{2bn}$$

input  $\operatorname{Integrate}[x^{(-1 + n)}*\operatorname{ArcTanh}[a + b*x^n], x]$

output  $(2*(a + b*x^n)*\operatorname{ArcTanh}[a + b*x^n] + \operatorname{Log}[1 - (a + b*x^n)^2])/(2*b*n)$

**Rubi [A] (warning: unable to verify)**

Time = 0.30 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {7266, 6653, 6436, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x^{n-1} \operatorname{arctanh}(a + bx^n) dx \\
 \downarrow 7266 \\
 \frac{\int \operatorname{arctanh}(bx^n + a) dx^n}{n} \\
 \downarrow 6653 \\
 \frac{\int \operatorname{arctanh}(bx^n + a) d(bx^n + a)}{bn} \\
 \downarrow 6436 \\
 \frac{(a + bx^n) \operatorname{arctanh}(a + bx^n) - \int \frac{bx^n + a}{1 - x^{2n}} d(bx^n + a)}{bn} \\
 \downarrow 240 \\
 \frac{(a + bx^n) \operatorname{arctanh}(a + bx^n) + \frac{1}{2} \log(1 - x^{2n})}{bn}
 \end{array}$$

input `Int[x^(-1 + n)*ArcTanh[a + b*x^n], x]`

output `((a + b*x^n)*ArcTanh[a + b*x^n] + Log[1 - x^(2*n)]/2)/(b*n)`

## Definitions of rubi rules used

rule 240  $\text{Int}[(x_)/((a_) + (b_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}[\{a, b\}, x]$

rule 6436  $\text{Int}[(a_) + \text{ArcTanh}[(c_)*(x_)^(n_)]*(b_)]^(p_), x\_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcTanh}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \text{Int}[x^n*((a + b*\text{ArcTanh}[c*x^n])^(p-1)/(1 - c^2*x^(2*n))), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[n, 1] \mid \mid \text{EqQ}[p, 1])$

rule 6653  $\text{Int}[(a_) + \text{ArcTanh}[(c_) + (d_)*(x_)]*(b_)]^(p_), x\_Symbol] \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(a + b*\text{ArcTanh}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{IGtQ}[p, 0]$

rule 7266  $\text{Int}[(u_)*(x_)^(m_), x\_Symbol] \rightarrow \text{Simp}[1/(m + 1) \text{Subst}[\text{Int}[\text{SubstFor}[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; \text{FreeQ}[m, x] \&\& \text{NeQ}[m, -1] \&\& \text{FunctionOfQ}[x^(m + 1), u, x]$

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs.  $2(45) = 90$ .

Time = 4.93 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.57

method	result	size
risch	$\frac{x^n \ln(1+a+bx^n)}{2n} - \frac{x^n \ln(1-a-bx^n)}{2n} + \frac{\ln(x^n + \frac{1+a}{b})a}{2nb} - \frac{\ln(x^n + \frac{a-1}{b})a}{2nb} + \frac{\ln(x^n + \frac{1+a}{b})}{2nb} + \frac{\ln(x^n + \frac{a-1}{b})}{2nb}$	121

input `int(x^(-1+n)*arctanh(a+b*x^n),x,method=_RETURNVERBOSE)`

output  $\frac{1}{2/n*x^n*\ln(1+a+b*x^n)} - \frac{1}{2/n*x^n*\ln(1-a-b*x^n)} + \frac{1}{2/n/b*\ln(x^n+(1+a)/b)}*a - \frac{1}{2/n/b*\ln(x^n+(a-1)/b)}*a + \frac{1}{2/n/b*\ln(x^n+(1+a)/b)} + \frac{1}{2/n/b*\ln(x^n+(a-1)/b)}$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 109 vs.  $2(45) = 90$ .

Time = 0.09 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.32

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx$$

$$= \frac{(a + 1) \log(b \cosh(n \log(x)) + b \sinh(n \log(x)) + a + 1) - (a - 1) \log(b \cosh(n \log(x)) + b \sinh(n \log(x)) + a - 1)}{2bn}$$

input `integrate(x^(-1+n)*arctanh(a+b*x^n),x, algorithm="fricas")`

output `1/2*((a + 1)*log(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a + 1) - (a - 1)*log(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a - 1) + (b*cosh(n*log(x)) + b*sinh(n*log(x)))*log(-(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a + 1)/(b*cosh(n*log(x)) + b*sinh(n*log(x)) + a - 1)))/(b*n)`

**Sympy [F(-2)]**

Exception generated.

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate(x**(-1+n)*atanh(a+b*x**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.85

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \frac{2(bx^n + a) \operatorname{artanh}(bx^n + a) + \log(-(bx^n + a)^2 + 1)}{2bn}$$

input `integrate(x^(-1+n)*arctanh(a+b*x^n),x, algorithm="maxima")`

output `1/2*(2*(b*x^n + a)*arctanh(b*x^n + a) + log(-(b*x^n + a)^2 + 1))/(b*n)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(45) = 90.

Time = 0.14 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.64

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx$$

$$= \frac{((a + 1)b - (a - 1)b) \left( \frac{\log\left(\frac{|-bx^n - a - 1|}{|bx^n + a - 1|}\right)}{b^2} - \frac{\log\left(\left|-\frac{bx^n + a + 1}{bx^n + a - 1} + 1\right|\right)}{b^2} + \frac{\log\left(-\frac{bx^n + a + 1}{bx^n + a - 1}\right)}{b^2 \left(\frac{bx^n + a + 1}{bx^n + a - 1} - 1\right)} \right)}{2n}$$

input `integrate(x^(-1+n)*arctanh(a+b*x^n),x, algorithm="giac")`

output `1/2*((a + 1)*b - (a - 1)*b)*(log(abs(-b*x^n - a - 1)/abs(b*x^n + a - 1))/b^2 - log(abs(-(b*x^n + a + 1)/(b*x^n + a - 1) + 1))/b^2 + log(-(b*x^n + a + 1)/(b*x^n + a - 1))/(b^2*((b*x^n + a + 1)/(b*x^n + a - 1) - 1)))/n`

**Mupad [B] (verification not implemented)**

Time = 3.74 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.19

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \frac{x^n \operatorname{atanh}(a + bx^n)}{n} - \frac{\ln(a + bx^n - 1)(a - 1)}{2bn} + \frac{\ln(a + bx^n + 1)(a + 1)}{2bn}$$

input `int(x^(n - 1)*atanh(a + b*x^n),x)`output `(x^n*atanh(a + b*x^n))/n - (log(a + b*x^n - 1)*(a - 1))/(2*b*n) + (log(a + b*x^n + 1)*(a + 1))/(2*b*n)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.19

$$\int x^{-1+n} \operatorname{arctanh}(a + bx^n) dx = \frac{2x^n \operatorname{atanh}(x^n b + a) b + 2 \operatorname{atanh}(x^n b + a) a + \log(x^{2n} b^2 + 2x^n a b + a^2 - 1)}{2bn}$$

input `int(x^(-1+n)*atanh(a+b*x^n),x)`output `(2*x**n*atanh(x**n*b + a)*b + 2*atanh(x**n*b + a)*a + log(x**(2*n)*b**2 + 2*x**n*a*b + a**2 - 1))/(2*b*n)`



$$3.30 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx$$

Optimal result	348
Mathematica [N/A]	348
Rubi [N/A]	349
Maple [N/A]	349
Fricas [N/A]	350
Sympy [N/A]	350
Maxima [N/A]	351
Giac [N/A]	351
Mupad [N/A]	352
Reduce [N/A]	352

### Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx = \operatorname{Int}\left(\frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2}, x\right)$$

output `Defer(Int)((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x)`

### Mathematica [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx = \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx$$

input `Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2),x]`

output `Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2), x]`

**Rubi [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{1 - c^2 x^2} dx$$

↓ 7234

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{1 - c^2 x^2} dx$$

input

```
Int[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2), x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 0.40 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)\right)^n}{-c^2 x^2 + 1} dx$$

input

```
int((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1), x)
```

output

```
int((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1), x)
```

**Fricas [N/A]**

Not integrable

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, algorithm="fricas")`

output `integral(-(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

**Sympy [N/A]**

Not integrable

Time = 9.71 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = - \int \frac{\left(a + b \operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**n/(-c**2*x**2+1),x)`

output `-Integral((a + b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1)))**n/(c**2*x**2 - 1), x)`

**Maxima [N/A]**

Not integrable

Time = 0.55 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, algorithm="maxima")`

output `-integrate((b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

**Giac [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

**Mupad [N/A]**

Not integrable

Time = 3.64 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2 x^2} dx = - \int \frac{\left(a + b \operatorname{atanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{c^2 x^2 - 1} dx$$

input `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^n/(c^2*x^2 - 1),x)`

output `-int((a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^n/(c^2*x^2 - 1), x)`

**Reduce [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2 x^2} dx = - \left( \int \frac{\left(\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) b + a\right)^n}{c^2 x^2 - 1} dx \right)$$

input `int((a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x)`

output `- int((atanh(sqrt(- c*x + 1)/sqrt(c*x + 1))*b + a)**n/(c**2*x**2 - 1),x)`

$$3.31 \quad \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1-c^2x^2} dx$$

Optimal result	353
Mathematica [F]	354
Rubi [A] (verified)	354
Maple [B] (verified)	357
Fricas [F]	358
Sympy [F]	359
Maxima [F]	359
Giac [F]	360
Mupad [F(-1)]	360
Reduce [F]	361

### Optimal result

Integrand size = 40, antiderivative size = 409

$$\begin{aligned} & \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1-c^2x^2} dx \\ &= -\frac{2\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad + \frac{3b\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad - \frac{3b\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad - \frac{3b^2\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad + \frac{3b^2\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad + \frac{3b^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{4c} - \frac{3b^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{4c} \end{aligned}$$

output

```
2*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3*arctanh(-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/2*b*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(2,1-2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/2*b*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(2,-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/2*b^2*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))*polylog(3,1-2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/2*b^2*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))*polylog(3,-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/4*b^3*polylog(4,1-2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/4*b^3*polylog(4,-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c
```

**Mathematica [F]**

$$\int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx$$

input

```
Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2), x]
```

output

```
Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2), x]
```

**Rubi [A] (verified)**

Time = 1.20 (sec) , antiderivative size = 402, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {7232, 6448, 6614, 6620, 6624, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^3}{1 - c^2x^2} dx$$

↓ 7232

$$\int \frac{\sqrt{cx+1} \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}$$

c  
↓ 6448

$$\frac{2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \int \frac{\left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right)}{1 - \frac{1-cx}{cx+1}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}{c}$$

↓ 6614

$$\frac{2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left( \frac{1}{2} \int \frac{\left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2 \log \left( 2 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right)}{1 - \frac{1-cx}{cx+1}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right)}{c}$$

↓ 6620

$$\frac{2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left( \frac{1}{2} \left( \frac{1}{2} \operatorname{PolyLog} \left( 2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

↓ 6624

$$\frac{2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left( \frac{1}{2} \left( \frac{1}{2} \operatorname{PolyLog} \left( 2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

↓ 7164

$$\frac{2 \operatorname{arctanh} \left( 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left( \frac{1}{2} \left( \frac{1}{2} \operatorname{PolyLog} \left( 2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \right) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

input

`Int[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2), x]`



output

```

-((2*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3*ArcTanh[1 - 2/(1 - Sqr
t[1 - c*x]/Sqrt[1 + c*x]]) - 6*b*(((a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 +
c*x]])^2*PolyLog[2, 1 - 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x])))/2 - b*((a +
b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[3, 1 - 2/(1 - Sqrt[1 - c*
x]/Sqrt[1 + c*x])))/2 - (b*PolyLog[4, 1 - 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*
x])))/4))/2 + (-1/2*((a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2*PolyLo
g[2, -1 + 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x])) + b*((a + b*ArcTanh[Sqrt[
1 - c*x]/Sqrt[1 + c*x]])*PolyLog[3, -1 + 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x
])))/2 - (b*PolyLog[4, -1 + 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]))/4))/2))/
c)

```

### Defintions of rubi rules used

rule 6448

```

Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_)/(x_), x_Symbol] := Simp[2*(a +
b*ArcTanh[c*x])^p*ArcTanh[1 - 2/(1 - c*x)], x] - Simp[2*b*c*p Int[(a + b
*ArcTanh[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 - c*x)]/(1 - c^2*x^2)), x], x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]

```

rule 6614

```

Int[(ArcTanh[u_]*((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(
x_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTanh[c*x])^p/(d +
e*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTanh[c*x])^p/(d + e
*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d + e,
0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]

```

rule 6620

```

Int[(Log[u_]*((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(x_)^
2), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d))
, x] + Simp[b*(p/2 Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d
+ e, 0] && EqQ[(1 - u)^2 - (1 - 2/(1 - c*x))^2, 0]

```

rule 6624

```

Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^p_*PolyLog[k_, u_]/((d_) + (e_.
)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^p*(PolyLog[k + 1, u]/(2*
c*d)), x] - Simp[b*(p/2 Int[(a + b*ArcTanh[c*x])^(p - 1)*(PolyLog[k + 1,
u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] && E
qQ[c^2*d + e, 0] && EqQ[u^2 - (1 - 2/(1 - c*x))^2, 0]

```

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] := With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

rule 7232 `Int[((a_.) + (b_.)*(F_)[((c_.)*Sqrt[(d_.) + (e_.)*(x_)])/Sqrt[(f_.) + (g_.)*(x_)])^(n_.)/((A_.) + (C_.)*(x_)^2), x_Symbol] := Simp[2*e*(g/(C*(e*f - d*g))) Subst[Int[(a + b*F[c*x])^n/x, x], x, Sqrt[d + e*x]/Sqrt[f + g*x]], x] /; FreeQ[{a, b, c, d, e, f, g, A, C, F}, x] && EqQ[C*d*f - A*e*g, 0] && EqQ[e*f + d*g, 0] && IGtQ[n, 0]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1622 vs.  $2(349) = 698$ .

Time = 1.84 (sec) , antiderivative size = 1623, normalized size of antiderivative = 3.97

method	result	size
default	Expression too large to display	1623
parts	Expression too large to display	1623

input `int((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x,method=_RETURNERVERBOSE)`

output

```

1/2*a^3/c*ln(c*x+1)-1/2*a^3/c*ln(c*x-1)-b^3*(1/c*arctanh((-c*x+1)^(1/2)/(c
*x+1)^(1/2))^3*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(
1/2))+3/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))^2*polylog(2,-((-c*x+1)^(1
/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-6/c*arctanh((-c*x+1)^(1/2
)/(c*x+1)^(1/2))*polylog(3,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(
c*x+1))^(1/2))+6/c*polylog(4,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)
/(c*x+1))^(1/2))-1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))^3*ln(1+((-c*x+1
)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))^2-3/2/c*arctanh((-c*x+1)^(1
/2)/(c*x+1)^(1/2))^2*polylog(2,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*
x+1)/(c*x+1)))^2+3/2/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(3,-((-c
*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))^2-3/4/c*polylog(4,-((-c
*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))^2+1/c*arctanh((-c*x+1)^(
1/2)/(c*x+1)^(1/2))^3*ln(1-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(
c*x+1))^(1/2))+3/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))^2*polylog(2,((-c*
x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-6/c*arctanh((-c*x+
1)^(1/2)/(c*x+1)^(1/2))*polylog(3,((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*
x+1)/(c*x+1))^(1/2))+6/c*polylog(4,((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c
*x+1)/(c*x+1))^(1/2)))^2-3*a*b^2*(1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))^
2*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))+2/c*ar
ctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-((-c*x+1)^(1/2)/(c*x+1))^...

```

**Fricas [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2 x^2} dx = \int -\frac{\left(b \operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2 x^2 - 1} dx$$

input

```

integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, al
gorithm="fricas")

```

output

```

integral(-(b^3*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1))^3 + 3*a*b^2*arctanh(s
qrt(-c*x + 1)/sqrt(c*x + 1))^2 + 3*a^2*b*arctanh(sqrt(-c*x + 1)/sqrt(c*x +
1)) + a^3)/(c^2*x^2 - 1), x)

```

**Sympy [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = - \int \frac{a^3}{c^2x^2 - 1} dx - \int \frac{b^3 \operatorname{atanh}^3\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

$$- \int \frac{3ab^2 \operatorname{atanh}^2\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

$$- \int \frac{3a^2b \operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input `integrate((a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**3/(-c**2*x**2+1),x)`

output `-Integral(a**3/(c**2*x**2 - 1), x) - Integral(b**3*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))**3/(c**2*x**2 - 1), x) - Integral(3*a*b**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1), x) - Integral(3*a**2*b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)`

**Maxima [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int - \frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, algorithm="maxima")`

output

```
1/2*a^3*(log(c*x + 1)/c - log(c*x - 1)/c) - 1/16*(b^3*log(c*x + 1) - b^3*log(-c*x + 1))*log(sqrt(c*x + 1) - sqrt(-c*x + 1))^3/c - integrate(1/32*(4*(sqrt(c*x + 1)*b^3 - sqrt(-c*x + 1)*b^3)*log(sqrt(c*x + 1) + sqrt(-c*x + 1))^3 + 24*(sqrt(c*x + 1)*a*b^2 - sqrt(-c*x + 1)*a*b^2)*log(sqrt(c*x + 1) + sqrt(-c*x + 1))^2 + 3*(4*(sqrt(c*x + 1)*b^3 - sqrt(-c*x + 1)*b^3)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) + (8*a*b^2 - (b^3*c*x - b^3)*log(c*x + 1) + (b^3*c*x - b^3)*log(-c*x + 1))*sqrt(c*x + 1) - (8*a*b^2 - (b^3*c*x + b^3)*log(c*x + 1) + (b^3*c*x + b^3)*log(-c*x + 1))*sqrt(-c*x + 1))*log(sqrt(c*x + 1) - sqrt(-c*x + 1))^2 + 48*(sqrt(c*x + 1)*a^2*b - sqrt(-c*x + 1)*a^2*b)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) - 12*(4*sqrt(c*x + 1)*a^2*b - 4*sqrt(-c*x + 1)*a^2*b + (sqrt(c*x + 1)*b^3 - sqrt(-c*x + 1)*b^3)*log(sqrt(c*x + 1) + sqrt(-c*x + 1))^2 + 4*(sqrt(c*x + 1)*a*b^2 - sqrt(-c*x + 1)*a*b^2)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)))*log(sqrt(c*x + 1) - sqrt(-c*x + 1)))/((c^2*x^2 - 1)*sqrt(c*x + 1) - (c^2*x^2 - 1)*sqrt(-c*x + 1)), x)
```

**Giac [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int -\frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2x^2 - 1} dx$$

input

```
integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, algorithm="giac")
```

output

```
integrate(-(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^3/(c^2*x^2 - 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int -\frac{\left(a + b \operatorname{atanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^3}{c^2x^2 - 1} dx$$

input

```
int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^3/(c^2*x^2 - 1),x)
```

output `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^3/(c^2*x^2 - 1), x)`

**Reduce [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx$$

$$= \frac{-6 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2-1} dx \right) a^2bc - 2 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^3}{c^2x^2-1} dx \right) b^3c - 6 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2}{c^2x^2-1} dx \right) ab^2c - \log(c^2x - c)}{2c}$$

input `int((a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1), x)`

output `( - 6*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1),x)*a**2*b*c - 2*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))**3/(c**2*x**2 - 1),x)*b**3*c - 6*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1),x)*a*b**2*c - log(c**2*x - c)*a**3 + log(c**2*x + c)*a**3)/(2*c)`

**3.32** 
$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1-c^2x^2} dx$$

Optimal result	362
Mathematica [F]	363
Rubi [A] (verified)	363
Maple [B] (verified)	366
Fricas [F]	367
Sympy [F]	368
Maxima [F]	368
Giac [F]	369
Mupad [F(-1)]	369
Reduce [F]	369

**Optimal result**

Integrand size = 40, antiderivative size = 268

$$\begin{aligned} & \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1-c^2x^2} dx \\ &= -\frac{2\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad + \frac{b\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad - \frac{b\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad - \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} + \frac{b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \end{aligned}$$

output

```
2*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*arctanh(-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+b*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,1-2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-b*(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-1/2*b^2*polylog(3,1-2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+1/2*b^2*polylog(3,-1+2/(1-(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c
```

**Mathematica [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2 x^2} dx = \int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2 x^2} dx$$

input

```
Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]
```

output

```
Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]
```

**Rubi [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {7232, 6448, 6614, 6620, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{1 - c^2 x^2} dx$$

↓ 7232

$$\int \frac{\sqrt{cx+1} \left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}$$

c

↓ 6448



$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \int \frac{\left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right) \operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) d\frac{\sqrt{1-cx}}{\sqrt{cx+1}}}{1 - \frac{1-cx}{cx+1}}}{c}$$

↓ 6614

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \int \frac{\left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right) \log\left(2 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) d\frac{\sqrt{1-cx}}{\sqrt{cx+1}}}{1 - \frac{1-cx}{cx+1}} - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)\right)}{c}$$

↓ 6620

$$2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)\right)$$

↓ 7164

$$2\operatorname{arctanh}\left(1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)\right)$$

input `Int[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]`

output `-((2*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2*ArcTanh[1 - 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]]) - 4*b*(((a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[2, 1 - 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]])/2 - (b*PolyLog[3, 1 - 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]])/4)/2 + (-1/2*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[2, -1 + 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]]) + (b*PolyLog[3, -1 + 2/(1 - Sqrt[1 - c*x]/Sqrt[1 + c*x]])/4)/2))/c)`

## Definitions of rubi rules used

rule 6448  $\text{Int}[(a + \text{ArcTanh}[c \cdot x] \cdot b)^p / x, x\_Symbol] \rightarrow \text{Simp}[2 \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot \text{ArcTanh}[1 - 2/(1 - c \cdot x)], x] - \text{Simp}[2 \cdot b \cdot c \cdot p \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{ArcTanh}[1 - 2/(1 - c \cdot x)] / (1 - c^2 \cdot x^2)), x], x] /;$   
 $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$

rule 6614  $\text{Int}[(\text{ArcTanh}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b))^p / ((d + e \cdot x^2)), x\_Symbol] \rightarrow \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 + u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)), x], x] - \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 - u] \cdot (a + b \cdot \text{ArcTanh}[c \cdot x])^p / (d + e \cdot x^2)), x], x] /;$   
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[u^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 6620  $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTanh}[c \cdot x] \cdot b))^p / ((d + e \cdot x^2)), x\_Symbol] \rightarrow \text{Simp}[(-a + b \cdot \text{ArcTanh}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot (p/2) \cdot \text{Int}[(a + b \cdot \text{ArcTanh}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2)), x], x] /;$   
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2/(1 - c \cdot x))^2, 0]$

rule 7164  $\text{Int}[u \cdot \text{PolyLog}[n, v], x\_Symbol] \rightarrow \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$   
 $! \text{FalseQ}[w] /;$   
 $\text{FreeQ}[n, x]$

rule 7232  $\text{Int}[(a + b \cdot (F)[(c \cdot \text{Sqrt}[d + e \cdot x]) / \text{Sqrt}[f + g \cdot x]) \cdot x]^n / ((A + C \cdot x^2)), x\_Symbol] \rightarrow \text{Simp}[2 \cdot e \cdot (g / (C \cdot (e \cdot f - d \cdot g))) \cdot \text{Subst}[\text{Int}[(a + b \cdot F[c \cdot x])^n / x, x], x, \text{Sqrt}[d + e \cdot x] / \text{Sqrt}[f + g \cdot x]], x] /;$   
 $\text{FreeQ}\{a, b, c, d, e, f, g, A, C, F, x\} \ \&\& \ \text{EqQ}[C \cdot d \cdot f - A \cdot e \cdot g, 0] \ \&\& \ \text{EqQ}[e \cdot f + d \cdot g, 0] \ \&\& \ \text{IGtQ}[n, 0]$

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 892 vs. 2(232) = 464.

Time = 0.61 (sec) , antiderivative size = 893, normalized size of antiderivative = 3.33

method	result
default	$\frac{a^2 \ln(cx+1)}{2c} - \frac{a^2 \ln(cx-1)}{2c} - b^2 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2 \ln\left(1 + \frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \operatorname{polylog}\left(2, -\frac{\sqrt{-cx+1}}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} \right)$
parts	$\frac{a^2 \ln(cx+1)}{2c} - \frac{a^2 \ln(cx-1)}{2c} - b^2 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2 \ln\left(1 + \frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \operatorname{polylog}\left(2, -\frac{\sqrt{-cx+1}}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} \right)$

input

```
int((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x,method=_R
ETURNVERBOSE)
```

output

```

1/2*a^2/c*ln(c*x+1)-1/2*a^2/c*ln(c*x-1)-b^2*(1/c*arctanh((-c*x+1)^(1/2)/(c
*x+1)^(1/2))^2*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(
1/2))+2/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-((-c*x+1)^(1/2)
)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-2/c*polylog(3,-((-c*x+1)^(1
/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-1/c*arctanh((-c*x+1)^(1/2)
)/(c*x+1)^(1/2))^2*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*
x+1)))-1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-((-c*x+1)^(1/2)
)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))+1/2/c*polylog(3,-((-c*x+1)^(1/2)
)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))+1/c*arctanh((-c*x+1)^(1/2)/(c*x
+1)^(1/2))^2*ln(1-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1
/2))+2/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,((-c*x+1)^(1/2)/(
c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-2/c*polylog(3,((-c*x+1)^(1/2)/
(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2)))-2*a*b*(1/c*arctanh((-c*x+1)^(
1/2)/(c*x+1)^(1/2))*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*
x+1)^(1/2))+1/c*polylog(2,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(
c*x+1))^(1/2))-1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1+((-c*x+1)^(1
/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))-1/2/c*polylog(2,-((-c*x+1)^(1
/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))+1/c*arctanh((-c*x+1)^(1/2)/(c
*x+1)^(1/2))*ln(1-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1
/2))+1/c*polylog(2,((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1)...

```

**Fricas [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2 x^2} dx = \int -\frac{\left(b \operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2 x^2 - 1} dx$$

input

```

integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, al
gorithm="fricas")

```

output

```

integral(-(b^2*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1))^2 + 2*a*b*arctanh(sqrt
(-c*x + 1)/sqrt(c*x + 1)) + a^2)/(c^2*x^2 - 1), x)

```

**Sympy [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = - \int \frac{a^2}{c^2x^2 - 1} dx - \int \frac{b^2 \operatorname{atanh}^2\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx - \int \frac{2ab \operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input `integrate((a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**2/(-c**2*x**2+1),x)`

output `-Integral(a**2/(c**2*x**2 - 1), x) - Integral(b**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1), x) - Integral(2*a*b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)`

**Maxima [F]**

$$\int \frac{\left(a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int - \frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, algorithm="maxima")`

output `1/2*a^2*(log(c*x + 1)/c - log(c*x - 1)/c) + 1/8*(b^2*log(c*x + 1) - b^2*log(-c*x + 1))*log(sqrt(c*x + 1) - sqrt(-c*x + 1))^2/c + integrate(-1/8*(2*(sqrt(c*x + 1)*b^2 - sqrt(-c*x + 1)*b^2)*log(sqrt(c*x + 1) + sqrt(-c*x + 1))^2 + 8*(sqrt(c*x + 1)*a*b - sqrt(-c*x + 1)*a*b)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) - (4*(sqrt(c*x + 1)*b^2 - sqrt(-c*x + 1)*b^2)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) + (8*a*b - (b^2*c*x - b^2)*log(c*x + 1) + (b^2*c*x - b^2)*log(-c*x + 1))*sqrt(c*x + 1) - (8*a*b - (b^2*c*x + b^2)*log(c*x + 1) + (b^2*c*x + b^2)*log(-c*x + 1))*sqrt(-c*x + 1))*log(sqrt(c*x + 1) - sqrt(-c*x + 1)))/((c^2*x^2 - 1)*sqrt(c*x + 1) - (c^2*x^2 - 1)*sqrt(-c*x + 1)), x)`

**Giac [F]**

$$\int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int -\frac{\left(b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, algorithm="giac")`

output `integrate(-(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^2/(c^2*x^2 - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int -\frac{\left(a + b \operatorname{atanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{c^2x^2 - 1} dx$$

input `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2/(c^2*x^2 - 1), x)`

output `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2/(c^2*x^2 - 1), x)`

**Reduce [F]**

$$\int \frac{\left(a + \operatorname{barctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx$$

$$= \frac{-4 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2-1} dx \right) abc - 2 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2}{c^2x^2-1} dx \right) b^2c - \log(c^2x - c) a^2 + \log(c^2x + c) a^2}{2c}$$

input `int((a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1), x)`

output

```
( - 4*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1),x)*a*b*c -  
2*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1),x)*b**2*c  
- log(c**2*x - c)*a**2 + log(c**2*x + c)*a**2)/(2*c)
```

**3.33**  $\int \frac{a+b\operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1-c^2x^2} dx$

Optimal result	371
Mathematica [A] (verified)	371
Rubi [A] (verified)	372
Maple [B] (verified)	373
Fricas [F]	374
Sympy [F]	374
Maxima [F]	375
Giac [F]	375
Mupad [F(-1)]	376
Reduce [F]	376

**Optimal result**

Integrand size = 38, antiderivative size = 89

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = -\frac{a \log\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{c} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{2c} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{2c}$$

output

```
-a*ln((-c*x+1)^(1/2)/(c*x+1)^(1/2))/c+1/2*b*polylog(2,-(c*x+1)^(1/2)/(c*x+1)^(1/2))/c-1/2*b*polylog(2,(c*x+1)^(1/2)/(c*x+1)^(1/2))/c
```

**Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.48

$$\int \frac{a + b\operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \frac{a\operatorname{arctanh}(cx)}{c} + \frac{b(\operatorname{PolyLog}(2, -e^{-\operatorname{arctanh}(cx)}) - \operatorname{PolyLog}(2, e^{-\operatorname{arctanh}(cx)}))}{2c}$$

input

```
Integrate[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])/(1 - c^2*x^2),x]
```



output

```
(a*ArcTanh[c*x])/c + (b*(PolyLog[2, -E^(-ArcTanh[c*x])] - PolyLog[2, E^(-ArcTanh[c*x])]))/(2*c)
```

**Rubi [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.94, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$ , Rules used = {7232, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{1 - c^2 x^2} dx$$

↓ 7232

$$\int \frac{\sqrt{cx+1} \left( a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right) \right)}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}$$

↓ 6446

$$\frac{a \log\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right) - \frac{1}{2} b \operatorname{PolyLog}\left(2, -\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right) + \frac{1}{2} b \operatorname{PolyLog}\left(2, \frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{c}$$

input

```
Int[(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])/(1 - c^2*x^2), x]
```

output

```
-((a*Log[Sqrt[1 - c*x]/Sqrt[1 + c*x]] - (b*PolyLog[2, -(Sqrt[1 - c*x]/Sqrt[1 + c*x])]))/2 + (b*PolyLog[2, Sqrt[1 - c*x]/Sqrt[1 + c*x]]/2)/c)
```

## Definitions of rubi rules used

rule 6446

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x]
+ (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /
; FreeQ[{a, b, c}, x]
```

rule 7232

```
Int[((a_.) + (b_.)*(F_)[((c_.)*Sqrt[(d_.) + (e_.)*(x_)])/Sqrt[(f_.) + (g_.)
*(x_)])^(n_.)/((A_.) + (C_.)*(x_)^2), x_Symbol] := Simp[2*e*(g/(C*(e*f - d
*g))) Subst[Int[(a + b*F[c*x])^n/x, x], x, Sqrt[d + e*x]/Sqrt[f + g*x]],
x] /; FreeQ[{a, b, c, d, e, f, g, A, C, F}, x] && EqQ[C*d*f - A*e*g, 0] &&
EqQ[e*f + d*g, 0] && IGtQ[n, 0]
```

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs.  $2(73) = 146$ .

Time = 0.33 (sec) , antiderivative size = 357, normalized size of antiderivative = 4.01

method	result
default	$\frac{a \ln(cx+1)}{2c} - \frac{a \ln(cx-1)}{2c} - b \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \ln\left(1 + \frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} + \frac{\operatorname{polylog}\left(2, -\frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$
parts	$\frac{a \ln(cx+1)}{2c} - \frac{a \ln(cx-1)}{2c} - b \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \ln\left(1 + \frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} + \frac{\operatorname{polylog}\left(2, -\frac{\sqrt{-cx+1}+1}{\sqrt{1-\frac{-cx+1}{cx+1}}}\right)}{c} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$

input

```
int((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1), x, method=_RET
URNVERBOSE)
```

output

```
1/2*a/c*ln(c*x+1)-1/2*a/c*ln(c*x-1)-b*(1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))+1/c*polylog(2,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))-1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1+((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))-1/2/c*polylog(2,-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)^2/(1-(-c*x+1)/(c*x+1)))+1/c*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1-((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))+1/c*polylog(2,((-c*x+1)^(1/2)/(c*x+1)^(1/2)+1)/(1-(-c*x+1)/(c*x+1))^(1/2))
```

**Fricas [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2 x^2} dx = \int -\frac{b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2 x^2 - 1} dx$$

input

```
integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x, algorithm="fricas")
```

output

```
integral(-(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)/(c^2*x^2 - 1), x)
```

**Sympy [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2 x^2} dx = -\int \frac{a}{c^2 x^2 - 1} dx - \int \frac{b \operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2 x^2 - 1} dx$$

input

```
integrate((a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2)))/(-c**2*x**2+1),x)
```

output

```
-Integral(a/(c**2*x**2 - 1), x) - Integral(b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)
```

**Maxima [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x, algorith="maxima")`

output `1/4*b*((log(c*x + 1) - log(-c*x + 1))*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) - (log(c*x + 1) - log(-c*x + 1))*log(sqrt(c*x + 1) - sqrt(-c*x + 1)))/c - 2*integrate(-1/2*sqrt(c*x + 1)*(log(c*x + 1) - log(-c*x + 1))/((c^2*x^2 - 1)*sqrt(c*x + 1) + (c^2*x^2 - 1)*sqrt(-c*x + 1)), x) - 2*integrate(1/2*sqrt(c*x + 1)*(log(c*x + 1) - log(-c*x + 1))/((c^2*x^2 - 1)*sqrt(c*x + 1) - (c^2*x^2 - 1)*sqrt(-c*x + 1)), x) + 1/2*a*(log(c*x + 1)/c - log(c*x - 1)/c)`

**Giac [F]**

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{b \operatorname{artanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x, algorith="giac")`

output `integrate(-(b*arctanh(sqrt(-c*x + 1))/sqrt(c*x + 1)) + a)/(c^2*x^2 - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2 x^2} dx = \int -\frac{a + b \operatorname{atanh}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{c^2 x^2 - 1} dx$$

input `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))/(c^2*x^2 - 1), x)`

output `int(-(a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))/(c^2*x^2 - 1), x)`

**Reduce [F]**

$$\begin{aligned} & \int \frac{a + b \operatorname{arctanh}\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2 x^2} dx \\ & - 2 \left( \int \frac{\operatorname{atanh}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2 x^2 - 1} dx \right) bc - \log(c^2 x - c) a + \log(c^2 x + c) a \\ & = \frac{\hspace{10em}}{2c} \end{aligned}$$

input `int((a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1), x)`

output `( - 2*int(atanh(sqrt( - c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)*b*c - 1  
og(c**2*x - c)*a + log(c**2*x + c)*a)/(2*c)`

$$3.34 \quad \int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx$$

Optimal result	377
Mathematica [N/A]	377
Rubi [N/A]	378
Maple [N/A]	378
Fricas [N/A]	379
Sympy [N/A]	379
Maxima [N/A]	380
Giac [N/A]	380
Mupad [N/A]	381
Reduce [N/A]	381

### Optimal result

Integrand size = 40, antiderivative size = 40

$$\begin{aligned} & \int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx \\ &= \operatorname{Int} \left( \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)}, x \right) \end{aligned}$$

output `Defer(Int)(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)`

### Mathematica [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx$$

input `Integrate[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])),x]`

output

```
Integrate[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])),
x]
```

**Rubi [N/A]**

Not integrable

Time = 0.40 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)} dx$$

↓ 7234

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)} dx$$

input

```
Int[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])),x]
```

output

```
$Aborted
```

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-c^2 x^2 + 1) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{-cx + 1}}{\sqrt{cx + 1}} \right) \right)} dx$$

input

```
int(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)
```

output `int(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)`

### Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.20

$$\int \frac{1}{(1 - c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int -\frac{1}{(c^2x^2 - 1) \left( b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="fricas")`

output `integral(-1/(a*c^2*x^2 + (b*c^2*x^2 - b)*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) - a), x)`

### Sympy [N/A]

Not integrable

Time = 3.87 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.52

$$\int \frac{1}{(1 - c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx$$

$$= - \int \frac{1}{ac^2x^2 - a + bc^2x^2 \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - b \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)} dx$$

input `integrate(1/(-c**2*x**2+1)/(a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2))),x)`

output `-Integral(1/(a*c**2*x**2 - a + b*c**2*x**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) - b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))), x)`



**Maxima [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx = \int -\frac{1}{(c^2 x^2 - 1) \left( b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="maxima")`

output `-integrate(1/((c^2*x^2 - 1)*(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)), x)`

**Giac [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx = \int -\frac{1}{(c^2 x^2 - 1) \left( b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="giac")`

output `integrate(-1/((c^2*x^2 - 1)*(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)), x)`

**Mupad [N/A]**

Not integrable

Time = 3.36 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx = - \int \frac{1}{\left( a + b \operatorname{atanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right) (c^2 x^2 - 1)} dx$$

input `int(-1/((a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))*(c^2*x^2 - 1)),x)`

output `-int(1/((a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))*(c^2*x^2 - 1)), x)`

**Reduce [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.58

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx$$

$$= - \left( \int \frac{1}{\operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) b c^2 x^2 - \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) b + a c^2 x^2 - a} dx \right)$$

input `int(1/(-c^2*x^2+1)/(a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)`

output `-int(1/(atanh(sqrt(-c*x+1)/sqrt(c*x+1))*b*c**2*x**2 - atanh(sqrt(-c*x+1)/sqrt(c*x+1))*b + a*c**2*x**2 - a),x)`

$$3.35 \quad \int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx$$

Optimal result	382
Mathematica [N/A]	382
Rubi [N/A]	383
Maple [N/A]	384
Fricas [N/A]	384
Sympy [N/A]	385
Maxima [N/A]	385
Giac [N/A]	386
Mupad [N/A]	386
Reduce [N/A]	387

### Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx$$

$$= \operatorname{Int} \left( \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2}, x \right)$$

output `Defer(Int)(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

### Mathematica [N/A]

Not integrable

Time = 0.83 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx = \int \frac{1}{(1-c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx$$

input `Integrate[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2), x]`

output `Integrate[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2), x]`

### Rubi [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2x^2) \left( a + \text{barctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2} dx$$

↓ 7234

$$\int \frac{1}{(1 - c^2x^2) \left( a + \text{barctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2} dx$$

input `Int[1/((1 - c^2*x^2)*(a + b*ArcTanh[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2), x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-c^2x^2 + 1) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) \right)^2} dx$$

input `int(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

output `int(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.28

$$\begin{aligned} & \int \frac{1}{(1 - c^2x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx \\ &= \int -\frac{1}{(c^2x^2 - 1) \left( b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx \end{aligned}$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="fricas")`

output `integral(-1/(a^2*c^2*x^2 + (b^2*c^2*x^2 - b^2)*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1))^2 - a^2 + 2*(a*b*c^2*x^2 - a*b)*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1))), x)`

**Sympy [N/A]**

Not integrable

Time = 11.37 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.15

$$\int \frac{1}{(1 - c^2 x^2) \left( a + \operatorname{barctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx =$$

$$- \int \frac{1}{a^2 c^2 x^2 - a^2 + 2abc^2 x^2 \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - 2ab \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + b^2 c^2 x^2 \operatorname{atanh}^2 \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - b^2 \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)} dx$$

input `integrate(1/(-c**2*x**2+1)/(a+b*atanh((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**2, x)`

output `-Integral(1/(a**2*c**2*x**2 - a**2 + 2*a*b*c**2*x**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) - 2*a*b*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + b**2*c**2*x**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))**2 - b**2*atanh(sqrt(-c*x + 1)/sqrt(c*x + 1))**2), x)`

**Maxima [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 246, normalized size of antiderivative = 6.15

$$\int \frac{1}{(1 - c^2 x^2) \left( a + \operatorname{barctanh} \left( \frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx$$

$$= \int -\frac{1}{(c^2 x^2 - 1) \left( b \operatorname{artanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="maxima")`

output

```
4*c*x/(sqrt(c*x + 1)*sqrt(-c*x + 1)*b^2*c*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) - sqrt(c*x + 1)*sqrt(-c*x + 1)*b^2*c*log(sqrt(c*x + 1) - sqrt(-c*x + 1)) + 2*sqrt(c*x + 1)*sqrt(-c*x + 1)*a*b*c) - integrate(-4/((b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*log(sqrt(c*x + 1) + sqrt(-c*x + 1)) - (b^2*c^2*x^2 - b^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*log(sqrt(c*x + 1) - sqrt(-c*x + 1)) + 2*(a*b*c^2*x^2 - a*b)*sqrt(c*x + 1)*sqrt(-c*x + 1)), x)
```

**Giac [N/A]**

Not integrable

Time = 0.60 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx$$

$$= \int -\frac{1}{(c^2 x^2 - 1) \left( b \operatorname{arctanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx$$

input

```
integrate(1/(-c^2*x^2+1)/(a+b*arctanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="giac")
```

output

```
integrate(-1/((c^2*x^2 - 1)*(b*arctanh(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^2), x)
```

**Mupad [N/A]**

Not integrable

Time = 4.09 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx$$

$$= - \int \frac{1}{\left( a + b \operatorname{atanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)^2 (c^2 x^2 - 1)} dx$$

input `int(-1/((a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2*(c^2*x^2 - 1)),x)`

output `-int(1/((a + b*atanh((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2*(c^2*x^2 - 1)), x)`

### Reduce [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 123, normalized size of antiderivative = 3.08

$$\int \frac{1}{(1 - c^2 x^2) \left( a + b \operatorname{arctanh} \left( \frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx =$$

$$- \left( \int \frac{1}{\operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)^2 b^2 c^2 x^2 - \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)^2 b^2 + 2 \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) ab c^2 x^2 - 2 \operatorname{atanh} \left( \frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) a} \right)$$

input `int(1/(-c^2*x^2+1)/(a+b*atanh((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

output `-int(1/(atanh(sqrt(-c*x+1)/sqrt(c*x+1)))**2*b**2*c**2*x**2 - atanh(sqrt(-c*x+1)/sqrt(c*x+1))**2*b**2 + 2*atanh(sqrt(-c*x+1)/sqrt(c*x+1))*a*b*c**2*x**2 - 2*atanh(sqrt(-c*x+1)/sqrt(c*x+1))*a*b + a**2*c**2*x**2 - a**2),x)`



### 3.36 $\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result . . . . .	388
Mathematica [A] (verified) . . . . .	388
Rubi [A] (verified) . . . . .	389
Maple [A] (verified) . . . . .	390
Fricas [A] (verification not implemented) . . . . .	390
Sympy [F] . . . . .	391
Maxima [A] (verification not implemented) . . . . .	391
Giac [A] (verification not implemented) . . . . .	392
Mupad [B] (verification not implemented) . . . . .	392
Reduce [B] (verification not implemented) . . . . .	392

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^{2+m}}{2 + 3m + m^2} + \frac{x^{1+m} \operatorname{arctanh}(\tanh(a + bx))}{1 + m}$$

output

```
-b*x^(2+m)/(m^2+3*m+2)+x^(1+m)*arctanh(tanh(b*x+a))/(1+m)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx = x^m \left( \frac{bx^2}{2 + m} + \frac{x(-bx + \operatorname{arctanh}(\tanh(a + bx)))}{1 + m} \right)$$

input

```
Integrate[x^m*ArcTanh[Tanh[a + b*x]],x]
```

output

```
x^m*((b*x^2)/(2 + m) + (x*(-(b*x) + ArcTanh[Tanh[a + b*x]])))/(1 + m))
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))}{m+1} - \frac{b \int x^{m+1} dx}{m+1}$$

$$\downarrow \text{15}$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))}{m+1} - \frac{bx^{m+2}}{(m+1)(m+2)}$$

input `Int[x^m*ArcTanh[Tanh[a + b*x]],x]`

output `-((b*x^(2 + m))/((1 + m)*(2 + m))) + (x^(1 + m)*ArcTanh[Tanh[a + b*x]])/(1 + m)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

method	result
default	$\frac{b x^2 e^{m \ln(x)}}{2+m} + \frac{(\operatorname{arctanh}(\tanh(bx+a)) - bx) x e^{m \ln(x)}}{1+m}$
parallelrisch	$-\frac{-x x^m \operatorname{arctanh}(\tanh(bx+a)) m + b x^m x^2 - 2 \operatorname{arctanh}(\tanh(bx+a)) x x^m}{m^2 + 3m + 2}$
risch	$\frac{x x^m \ln(e^{bx+a})}{1+m} - \frac{x \left( 2i\pi \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^3 - i\pi \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 m + 4bx + i\pi \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^3 m + i\pi \right)}{m^2 + 3m + 2}$

input `int(x^m*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `b/(2+m)*x^2*exp(m*ln(x))+(arctanh(tanh(b*x+a))-b*x)/(1+m)*x*exp(m*ln(x))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.68

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$= \frac{((bm + b)x^2 + (am + 2a)x) \cosh(m \log(x)) + ((bm + b)x^2 + (am + 2a)x) \sinh(m \log(x))}{m^2 + 3m + 2}$$

input `integrate(x^m*arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `((b*m + b)*x^2 + (a*m + 2*a)*x)*cosh(m*log(x)) + ((b*m + b)*x^2 + (a*m + 2*a)*x)*sinh(m*log(x))/(m^2 + 3*m + 2)`

**Sympy [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$= \begin{cases} b \log(x) - \frac{\operatorname{atanh}(\tanh(a + bx))}{x} & \text{for } m = -2 \\ \int \frac{\operatorname{atanh}(\tanh(a + bx))}{x} dx & \text{for } m = -1 \\ -\frac{bx^2 x^m}{m^2 + 3m + 2} + \frac{m x x^m \operatorname{atanh}(\tanh(a + bx))}{m^2 + 3m + 2} + \frac{2 x x^m \operatorname{atanh}(\tanh(a + bx))}{m^2 + 3m + 2} & \text{otherwise} \end{cases}$$

input `integrate(x**m*atanh(tanh(b*x+a)),x)`

output `Piecewise((b*log(x) - atanh(tanh(a + b*x))/x, Eq(m, -2)), (Integral(atanh(tanh(a + b*x))/x, x), Eq(m, -1)), (-b*x**2*x**m/(m**2 + 3*m + 2) + m*x*x**m*atanh(tanh(a + b*x))/(m**2 + 3*m + 2) + 2*x*x**m*atanh(tanh(a + b*x))/(m**2 + 3*m + 2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^2 x^m}{(m + 2)(m + 1)} + \frac{x^{m+1} \operatorname{arctanh}(\tanh(bx + a))}{m + 1}$$

input `integrate(x^m*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `-b*x^2*x^m/((m + 2)*(m + 1)) + x^(m + 1)*arctanh(tanh(b*x + a))/(m + 1)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{bx^2 x^m + amx x^m + bx^2 x^m + 2axx^m}{m^2 + 3m + 2}$$

input `integrate(x^m*arctanh(tanh(b*x+a)),x, algorithm="giac")`output `(b*m*x^2*x^m + a*m*x*x^m + b*x^2*x^m + 2*a*x*x^m)/(m^2 + 3*m + 2)`**Mupad [B] (verification not implemented)**

Time = 3.66 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.59

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$= \frac{2bx^m x^2 (m+1)}{2m^2 + 6m + 4} - \frac{xx^m (m+2) \left( \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2bx \right)}{2m^2 + 6m + 4}$$

input `int(x^m*atanh(tanh(a + b*x)),x)`output `(2*b*x^m*x^2*(m + 1))/(6*m + 2*m^2 + 4) - (x*x^m*(m + 2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(6*m + 2*m^2 + 4)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int x^m \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$= \frac{x^m x (\operatorname{atanh}(\tanh(bx + a)) m + 2 \operatorname{atanh}(\tanh(bx + a)) - bx)}{m^2 + 3m + 2}$$

input `int(x^m*atanh(tanh(b*x+a)),x)`

output `(x**m*x*(atanh(tanh(a + b*x))*m + 2*atanh(tanh(a + b*x)) - b*x)/(m**2 + 3*m + 2)`

### 3.37 $\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	394
Mathematica [A] (verified)	394
Rubi [A] (verified)	395
Maple [A] (verified)	396
Fricas [A] (verification not implemented)	396
Sympy [A] (verification not implemented)	396
Maxima [A] (verification not implemented)	397
Giac [A] (verification not implemented)	397
Mupad [B] (verification not implemented)	397
Reduce [F]	398

#### Optimal result

Integrand size = 11, antiderivative size = 23

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(\tanh(a + bx))$$

output `-1/12*b*x^4+1/3*x^3*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{1}{12}x^3(bx - 4\operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[x^2*ArcTanh[Tanh[a + b*x]],x]`

output `-1/12*(x^3*(b*x - 4*ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{1}{3} x^3 \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^3 dx}{3}$$

$$\downarrow \text{15}$$

$$\frac{1}{3} x^3 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^4}{12}$$

input `Int[x^2*ArcTanh[Tanh[a + b*x]],x]`

output `-1/12*(b*x^4) + (x^3*ArcTanh[Tanh[a + b*x]])/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n])`



**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
default	$-\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))}{3}$
parallelrisch	$-\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))}{3}$
parts	$-\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))}{3}$
risch	$\frac{x^3 \ln(e^{bx+a})}{3} - \frac{bx^4}{12} + \frac{i\pi x^3 \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2}{6} - \frac{i\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)}{12}$

input `int(x^2*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `-1/12*b*x^4+1/3*x^3*arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{4} bx^4 + \frac{1}{3} ax^3$$

input `integrate(x^2*arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `1/4*b*x^4 + 1/3*a*x^3`**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^4}{12} + \frac{x^3 \operatorname{atanh}(\tanh(a + bx))}{3}$$

input `integrate(x**2*atanh(tanh(b*x+a)),x)`

output `-b*x**4/12 + x**3*atanh(tanh(a + b*x))/3`

### Maxima [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{1}{12} bx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(x^2*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `-1/12*b*x^4 + 1/3*x^3*arctanh(tanh(b*x + a))`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{4} bx^4 + \frac{1}{3} ax^3$$

input `integrate(x^2*arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `1/4*b*x^4 + 1/3*a*x^3`

### Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^3 \operatorname{atanh}(\tanh(a + bx))}{3} - \frac{bx^4}{12}$$

input `int(x^2*atanh(tanh(a + b*x)),x)`

output `(x^3*atanh(tanh(a + b*x)))/3 - (b*x^4)/12`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a)) x^2 dx$$

input `int(x^2*atanh(tanh(b*x+a)),x)`

output `int(atanh(tanh(a + b*x))*x**2,x)`

### 3.38 $\int x \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	399
Mathematica [A] (verified)	399
Rubi [A] (verified)	400
Maple [A] (verified)	401
Fricas [A] (verification not implemented)	401
Sympy [A] (verification not implemented)	401
Maxima [A] (verification not implemented)	402
Giac [A] (verification not implemented)	402
Mupad [B] (verification not implemented)	402
Reduce [F]	403

#### Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(\tanh(a + bx))$$

output `-1/6*b*x^3+1/2*x^2*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{1}{6}x^2(bx - 3 \operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[x*ArcTanh[Tanh[a + b*x]],x]`

output `-1/6*(x^2*(b*x - 3*ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6793, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{2}b \int -x^2 dx + \frac{1}{2}x^2 \operatorname{arctanh}(\tanh(a + bx))$$

$$\downarrow \text{15}$$

$$\frac{1}{2}x^2 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^3}{6}$$

input `Int[x*ArcTanh[Tanh[a + b*x]],x]`

output `-1/6*(b*x^3) + (x^2*ArcTanh[Tanh[a + b*x]])/2`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6793 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
default	$-\frac{bx^3}{6} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))}{2}$
parallelrisc	$-\frac{bx^3}{6} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))}{2}$
parts	$-\frac{bx^3}{6} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))}{2}$
risch	$\frac{x^2 \ln(e^{bx+a})}{2} - \frac{bx^3}{6} + \frac{i\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2}{8} - \frac{i\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)}{8}$

input `int(x*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `-1/6*b*x^3+1/2*x^2*arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{3} bx^3 + \frac{1}{2} ax^2$$

input `integrate(x*arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `1/3*b*x^3 + 1/2*a*x^2`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^3}{6} + \frac{x^2 \operatorname{atanh}(\tanh(a + bx))}{2}$$

input `integrate(x*atanh(tanh(b*x+a)),x)`

output `-b*x**3/6 + x**2*atanh(tanh(a + b*x))/2`

### Maxima [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{1}{6}bx^3 + \frac{1}{2}x^2 \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(x*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `-1/6*b*x^3 + 1/2*x^2*arctanh(tanh(b*x + a))`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{3}bx^3 + \frac{1}{2}ax^2$$

input `integrate(x*arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `1/3*b*x^3 + 1/2*a*x^2`

### Mupad [B] (verification not implemented)

Time = 3.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^2 \operatorname{atanh}(\tanh(a + bx))}{2} - \frac{bx^3}{6}$$

input `int(x*atanh(tanh(a + b*x)),x)`

output `(x^2*atanh(tanh(a + b*x)))/2 - (b*x^3)/6`

**Reduce [F]**

$$\int x \operatorname{arctanh}(\tanh(a + bx)) dx = \int x \operatorname{atanh}(\tanh(bx + a)) dx$$

input `int(x*atanh(tanh(b*x+a)),x)`

output `int(atanh(tanh(a + b*x))*x,x)`



### 3.39 $\int \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	404
Mathematica [A] (verified)	404
Rubi [A] (verified)	405
Maple [A] (verified)	406
Fricas [A] (verification not implemented)	406
Sympy [A] (verification not implemented)	407
Maxima [A] (verification not implemented)	407
Giac [A] (verification not implemented)	407
Mupad [B] (verification not implemented)	408
Reduce [B] (verification not implemented)	408

#### Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2b}$$

output `1/2*arctanh(tanh(b*x+a))^2/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{bx^2}{2} + x \operatorname{arctanh}(\tanh(a + bx))$$

input `Integrate[ArcTanh[Tanh[a + b*x]], x]`

output `-1/2*(b*x^2) + x*ArcTanh[Tanh[a + b*x]]`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2588$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx)) d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow 15$$

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2b}$$

input `Int[ArcTanh[Tanh[a + b*x]],x]`

output `ArcTanh[Tanh[a + b*x]]^2/(2*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2b}$
default	$\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2b}$
parallelrisch	$-\frac{x^2b}{2} + x \operatorname{arctanh}(\tanh(bx+a))$
parts	$-\frac{x^2b}{2} + x \operatorname{arctanh}(\tanh(bx+a))$
risch	$x \ln(e^{bx+a}) - \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) x}{4} + \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)}{4}$

input `int(arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*arctanh(tanh(b*x+a))^2/b`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{2} bx^2 + ax$$

input `integrate(arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `1/2*b*x^2 + a*x`

**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = \begin{cases} \frac{\operatorname{atanh}^2(\tanh(a + bx))}{2b} & \text{for } b \neq 0 \\ x \operatorname{atanh}(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a)),x)`output `Piecewise((atanh(tanh(a + b*x))**2/(2*b), Ne(b, 0)), (x*atanh(tanh(a)), True))`**Maxima [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{1}{2}bx^2 + x \operatorname{artanh}(\tanh(bx + a))$$

input `integrate(arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-1/2*b*x^2 + x*arctanh(tanh(b*x + a))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{1}{2}bx^2 + ax$$

input `integrate(arctanh(tanh(b*x+a)),x, algorithm="giac")`output `1/2*b*x^2 + a*x`

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = x \operatorname{atanh}(\tanh(a + bx)) - \frac{bx^2}{2}$$

input `int(atanh(tanh(a + b*x)),x)`output `x*atanh(tanh(a + b*x)) - (b*x^2)/2`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{\operatorname{atanh}(\tanh(bx + a))^2}{2b}$$

input `int(atanh(tanh(b*x+a)),x)`output `atanh(tanh(a + b*x))**2/(2*b)`

### 3.40 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} dx$

Optimal result	409
Mathematica [A] (verified)	409
Rubi [A] (verified)	410
Maple [A] (verified)	411
Fricas [A] (verification not implemented)	411
Sympy [F]	411
Maxima [A] (verification not implemented)	412
Giac [A] (verification not implemented)	412
Mupad [B] (verification not implemented)	412
Reduce [F]	413

#### Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} dx = bx - (bx - \operatorname{arctanh}(\tanh(a+bx))) \log(x)$$

output `b*x-(b*x-arctanh(tanh(b*x+a)))*ln(x)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} dx = bx + (-bx + \operatorname{arctanh}(\tanh(a+bx))) \log(x)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]/x,x]`

output `b*x + (-(b*x) + ArcTanh[Tanh[a + b*x]])*Log[x]`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx$$

↓ 2589

$$bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx$$

↓ 14

$$bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx)))$$

input `Int[ArcTanh[Tanh[a + b*x]]/x,x]`

output `b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x]`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

method	result
default	$\ln(x) \operatorname{arctanh}(\tanh(bx + a)) - b(x \ln(x) - x)$
parts	$\ln(x) \operatorname{arctanh}(\tanh(bx + a)) - b(x \ln(x) - x)$
risch	$\ln(x) \ln(e^{bx+a}) - \ln(x)xb + bx - \frac{i\pi \left( \operatorname{csgn}\left(\frac{i}{e^{2bx+2a+1}}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a+1}}\right) - \operatorname{csgn}\left(\frac{i}{e^{2bx+2a+1}}\right) \operatorname{csgn}\right)}{e^{2bx+2a+1}}$

input `int(arctanh(tanh(b*x+a))/x,x,method=_RETURNVERBOSE)`output `ln(x)*arctanh(tanh(b*x+a))-b*(x*ln(x)-x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.38

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = bx + a \log(x)$$

input `integrate(arctanh(tanh(b*x+a))/x,x, algorithm="fricas")`output `b*x + a*log(x)`**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))/x,x)`output `Integral(atanh(tanh(a + b*x))/x, x)`



**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = -b \left( x + \frac{a}{b} \right) \log(x) + b \left( x + \frac{a \log(x)}{b} \right) + \operatorname{arctanh}(\tanh(bx + a)) \log(x)$$

input `integrate(arctanh(tanh(b*x+a))/x,x, algorithm="maxima")`output `-b*(x + a/b)*log(x) + b*(x + a*log(x)/b) + arctanh(tanh(b*x + a))*log(x)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = bx + a \log(|x|)$$

input `integrate(arctanh(tanh(b*x+a))/x,x, algorithm="giac")`output `b*x + a*log(abs(x))`**Mupad [B] (verification not implemented)**

Time = 3.18 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.76

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = bx - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln(x)}{2} + \frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \ln(x)}{2} - bx \ln(x)$$

input `int(atanh(tanh(a + b*x))/x,x)`

output `b*x - (log(1/(exp(2*a)*exp(2*b*x) + 1))*log(x))/2 + (log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x))/2 - b*x*log(x)`

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))}{x} dx$$

input `int(atanh(tanh(b*x+a))/x,x)`

output `int(atanh(tanh(a + b*x))/x,x)`

### 3.41 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^2} dx$

Optimal result	414
Mathematica [A] (verified)	414
Rubi [A] (verified)	415
Maple [A] (verified)	416
Fricas [A] (verification not implemented)	416
Sympy [A] (verification not implemented)	416
Maxima [A] (verification not implemented)	417
Giac [A] (verification not implemented)	417
Mupad [B] (verification not implemented)	417
Reduce [B] (verification not implemented)	418

#### Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^2} dx = -\frac{\operatorname{arctanh}(\tanh(a+bx))}{x} + b \log(x)$$

output `-arctanh(tanh(b*x+a))/x+b*ln(x)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^2} dx = b - \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} + b \log(x)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]/x^2,x]`

output `b - ArcTanh[Tanh[a + b*x]]/x + b*Log[x]`

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx$$

↓ 2599

$$b \int \frac{1}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x}$$

↓ 14

$$b \log(x) - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]/x^2,x]
```

output

```
-(ArcTanh[Tanh[a + b*x]]/x) + b*Log[x]
```

**Defintions of rubi rules used**

rule 14

```
Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} + b \ln(x)$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} + b \ln(x)$
paralelrisch	$\frac{\ln(x)xb - \operatorname{arctanh}(\tanh(bx+a))}{x}$
risch	$-\frac{\ln(e^{bx+a})}{x} + \frac{i\pi \operatorname{csgn}(ie^{2bx+2a})^3 - 2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 + i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+a}}{e^{2bx+a}+1}\right)}{x}$

input `int(arctanh(tanh(b*x+a))/x^2,x,method=_RETURNVERBOSE)`output `-arctanh(tanh(b*x+a))/x+b*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = \frac{bx \log(x) - a}{x}$$

input `integrate(arctanh(tanh(b*x+a))/x^2,x, algorithm="fricas")`output `(b*x*log(x) - a)/x`**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = b \log(x) - \frac{\operatorname{atanh}(\tanh(a + bx))}{x}$$

input `integrate(atanh(tanh(b*x+a))/x**2,x)`

output `b*log(x) - atanh(tanh(a + b*x))/x`

### Maxima [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = b \log(x) - \frac{\operatorname{arctanh}(\tanh(bx + a))}{x}$$

input `integrate(arctanh(tanh(b*x+a))/x^2,x, algorithm="maxima")`

output `b*log(x) - arctanh(tanh(b*x + a))/x`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = b \log(|x|) - \frac{a}{x}$$

input `integrate(arctanh(tanh(b*x+a))/x^2,x, algorithm="giac")`

output `b*log(abs(x)) - a/x`

### Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = b \ln(x) - \frac{\operatorname{atanh}(\tanh(a + bx))}{x}$$

input `int(atanh(tanh(a + b*x))/x^2,x)`

output `b*log(x) - atanh(tanh(a + b*x))/x`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx = \frac{-\operatorname{atanh}(\tanh(bx + a)) + \log(x) bx}{x}$$

input `int(atanh(tanh(b*x+a))/x^2,x)`

output `( - atanh(tanh(a + b*x)) + log(x)*b*x)/x`

### 3.42 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^3} dx$

Optimal result . . . . .	419
Mathematica [A] (verified) . . . . .	419
Rubi [A] (verified) . . . . .	420
Maple [A] (verified) . . . . .	421
Fricas [A] (verification not implemented) . . . . .	421
Sympy [A] (verification not implemented) . . . . .	421
Maxima [A] (verification not implemented) . . . . .	422
Giac [A] (verification not implemented) . . . . .	422
Mupad [B] (verification not implemented) . . . . .	422
Reduce [B] (verification not implemented) . . . . .	423

#### Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(a + bx))}{2x^2}$$

output

```
-1/2*b/x-1/2*arctanh(tanh(b*x+a))/x^2
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{bx + \operatorname{arctanh}(\tanh(a + bx))}{2x^2}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]/x^3,x]
```

output

```
-1/2*(b*x + ArcTanh[Tanh[a + b*x]])/x^2
```



**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx$$

↓ 2599

$$\frac{1}{2}b \int \frac{1}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{2x^2}$$

↓ 15

$$-\frac{\operatorname{arctanh}(\tanh(a + bx))}{2x^2} - \frac{b}{2x}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]/x^3,x]
```

output

```
-1/2*b/x - ArcTanh[Tanh[a + b*x]]/(2*x^2)
```

**Defintions of rubi rules used**

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

method	result
parallelrisc	$-\frac{\operatorname{arctanh}(\tanh(bx+a))+bx}{2x^2}$
default	$-\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2}$
parts	$-\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2}$
risch	$-\frac{\ln(e^{bx+a})}{2x^2} - \frac{4bx - i\pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}(ie^{2bx+2a}) + 2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 - i\pi \operatorname{csgn}(ie^{2bx+2a})^3 - i\pi \operatorname{csgn}(ie^{2bx+2a})}{2x^2}$

input `int(arctanh(tanh(b*x+a))/x^3,x,method=_RETURNVERBOSE)`output `-1/2*(arctanh(tanh(b*x+a))+b*x)/x^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{2bx + a}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))/x^3,x, algorithm="fricas")`output `-1/2*(2*b*x + a)/x^2`**Sympy [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{b}{2x} - \frac{\operatorname{atanh}(\tanh(a + bx))}{2x^2}$$

input `integrate(atanh(tanh(b*x+a))/x**3,x)`

output `-b/(2*x) - atanh(tanh(a + b*x))/(2*x**2)`

### Maxima [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx + a))}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))/x^3,x, algorithm="maxima")`

output `-1/2*b/x - 1/2*arctanh(tanh(b*x + a))/x^2`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{2bx + a}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))/x^3,x, algorithm="giac")`

output `-1/2*(2*b*x + a)/x^2`

### Mupad [B] (verification not implemented)

Time = 3.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = -\frac{\operatorname{atanh}(\tanh(a + bx)) + bx}{2x^2}$$

input `int(atanh(tanh(a + b*x))/x^3,x)`

output `-(atanh(tanh(a + b*x)) + b*x)/(2*x^2)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^3} dx = \frac{-\operatorname{atanh}(\tanh(bx + a)) - bx}{2x^2}$$

input `int(atanh(tanh(b*x+a))/x^3,x)`

output `( - (atanh(tanh(a + b*x)) + b*x))/(2*x**2)`

### 3.43 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx$

Optimal result	424
Mathematica [A] (verified)	424
Rubi [A] (verified)	425
Maple [A] (verified)	426
Fricas [A] (verification not implemented)	426
Sympy [A] (verification not implemented)	426
Maxima [A] (verification not implemented)	427
Giac [A] (verification not implemented)	427
Mupad [B] (verification not implemented)	427
Reduce [B] (verification not implemented)	428

#### Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx = -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(a+bx))}{3x^3}$$

output

```
-1/6*b/x^2-1/3*arctanh(tanh(b*x+a))/x^3
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx = -\frac{bx + 2\operatorname{arctanh}(\tanh(a+bx))}{6x^3}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]/x^4,x]
```

output

```
-1/6*(b*x + 2*ArcTanh[Tanh[a + b*x]])/x^3
```

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx$$

↓ 2599

$$\frac{1}{3}b \int \frac{1}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{3x^3}$$

↓ 15

$$-\frac{\operatorname{arctanh}(\tanh(a + bx))}{3x^3} - \frac{b}{6x^2}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]/x^4,x]
```

output

```
-1/6*b/x^2 - ArcTanh[Tanh[a + b*x]]/(3*x^3)
```

**Defintions of rubi rules used**

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result
parallelrisc	$-\frac{bx+2 \operatorname{arctanh}(\tanh(bx+a))}{6x^3}$
default	$-\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^3}$
parts	$-\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^3}$
risc	$-\frac{\ln(e^{bx+a})}{3x^3} - \frac{2bx+2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 + i\pi \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)}{3x^3}$

input `int(arctanh(tanh(b*x+a))/x^4,x,method=_RETURNVERBOSE)`output `-1/6*(b*x+2*arctanh(tanh(b*x+a)))/x^3`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx = -\frac{3bx+2a}{6x^3}$$

input `integrate(arctanh(tanh(b*x+a))/x^4,x, algorithm="fricas")`output `-1/6*(3*b*x + 2*a)/x^3`**Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^4} dx = -\frac{b}{6x^2} - \frac{\operatorname{atanh}(\tanh(a+bx))}{3x^3}$$

input `integrate(atanh(tanh(b*x+a))/x**4,x)`

output  $-b/(6*x**2) - \operatorname{atanh}(\tanh(a + b*x))/(3*x**3)$

### Maxima [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx = -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx + a))}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))/x^4,x, algorithm="maxima")`

output  $-1/6*b/x^2 - 1/3*\operatorname{arctanh}(\tanh(b*x + a))/x^3$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx = -\frac{3bx + 2a}{6x^3}$$

input `integrate(arctanh(tanh(b*x+a))/x^4,x, algorithm="giac")`

output  $-1/6*(3*b*x + 2*a)/x^3$

### Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))}{3x^3} - \frac{b}{6x^2}$$

input `int(atanh(tanh(a + b*x))/x^4,x)`

output  $-\operatorname{atanh}(\tanh(a + b*x))/(3*x^3) - b/(6*x^2)$



**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^4} dx = \frac{-2\operatorname{atanh}(\tanh(bx + a)) - bx}{6x^3}$$

input `int(atanh(tanh(b*x+a))/x^4,x)`

output `( - 2*atanh(tanh(a + b*x)) - b*x)/(6*x**3)`

### 3.44 $\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	429
Mathematica [A] (verified)	429
Rubi [A] (verified)	430
Maple [A] (verified)	431
Fricas [B] (verification not implemented)	432
Sympy [F]	432
Maxima [A] (verification not implemented)	433
Giac [F]	433
Mupad [B] (verification not implemented)	434
Reduce [F]	434

#### Optimal result

Integrand size = 13, antiderivative size = 71

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2b^2 x^{3+m}}{6 + 11m + 6m^2 + m^3} - \frac{2bx^{2+m} \operatorname{arctanh}(\tanh(a + bx))}{2 + 3m + m^2} + \frac{x^{1+m} \operatorname{arctanh}(\tanh(a + bx))^2}{1 + m}$$

output

$$\frac{2*b^2*x^(3+m)/(m^3+6*m^2+11*m+6)-2*b*x^(2+m)*\operatorname{arctanh}(\tanh(b*x+a))/(m^2+3*m+2)+x^(1+m)*\operatorname{arctanh}(\tanh(b*x+a))^2/(1+m)}$$

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.87

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x^{1+m}(2b^2x^2 - 2b(3 + m)x \operatorname{arctanh}(\tanh(a + bx)) + (6 + 5m + m^2) \operatorname{arctanh}(\tanh(a + bx))^2)}{(1 + m)(2 + m)(3 + m)}$$

input

```
Integrate[x^m*ArcTanh[Tanh[a + b*x]]^2,x]
```

output

$$(x^{(1+m)}*(2*b^2*x^2 - 2*b*(3+m)*x*ArcTanh[Tanh[a+b*x]] + (6+5*m+m^2)*ArcTanh[Tanh[a+b*x]]^2))/((1+m)*(2+m)*(3+m))$$

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.93, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \operatorname{arctanh}(\tanh(a+bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^2}{m+1} - \frac{2b \int x^{m+1} \operatorname{arctanh}(\tanh(a+bx)) dx}{m+1}$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^2}{m+1} - \frac{2b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))}{m+2} - \frac{b \int x^{m+2} dx}{m+2} \right)}{m+1}$$

$$\downarrow 15$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^2}{m+1} - \frac{2b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))}{m+2} - \frac{bx^{m+3}}{(m+2)(m+3)} \right)}{m+1}$$

input

$$\text{Int}[x^m \operatorname{ArcTanh}[\operatorname{Tanh}[a+b*x]]^2, x]$$

output

$$(x^{(1+m)}*ArcTanh[Tanh[a+b*x]]^2)/(1+m) - (2*b*(-((b*x^(3+m)))/((2+m)*(3+m))) + (x^(2+m)*ArcTanh[Tanh[a+b*x]])/(2+m))/(1+m)$$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.38

method	result
default	$\frac{b^2 x^3 e^{m \ln(x)}}{3+m} + \frac{(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2) x e^{m \ln(x)}}{1+m} + \frac{2b(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{(1+m)(m^2 + 5m + 6)}$
parallelrisc	$-\frac{-2b^2 x^m x^3 - 6 \operatorname{arctanh}(\tanh(bx+a))^2 x x^m - x x^m \operatorname{arctanh}(\tanh(bx+a))^2 m^2 - 5x x^m \operatorname{arctanh}(\tanh(bx+a))^2 m + 6b \operatorname{arctanh}(\tanh(bx+a))^2}{(1+m)(m^2 + 5m + 6)}$
risc	Expression too large to display

input `int(x^m*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output `b^2/(3+m)*x^3*exp(m*ln(x))+(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/(1+m)*x*exp(m*ln(x))+2*b*(arctanh(tanh(b*x+a))-b*x)/(2+m)*x^2*exp(m*ln(x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 161 vs.  $2(71) = 142$ .

Time = 0.08 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.27

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$= \frac{((b^2 m^2 + 3 b^2 m + 2 b^2)x^3 + 2(abm^2 + 4 abm + 3 ab)x^2 + (a^2 m^2 + 5 a^2 m + 6 a^2)x) \cosh(m \log(x)) + ((b^2 m^2 + 3 b^2 m + 2 b^2)x^3 + 2(abm^2 + 4 abm + 3 ab)x^2 + (a^2 m^2 + 5 a^2 m + 6 a^2)x) \sinh(m \log(x))}{m^3 + 6 m^2 + 11 m + 6}$$

input `integrate(x^m*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `((b^2*m^2 + 3*b^2*m + 2*b^2)*x^3 + 2*(a*b*m^2 + 4*a*b*m + 3*a*b)*x^2 + (a^2*m^2 + 5*a^2*m + 6*a^2)*x)*cosh(m*log(x)) + ((b^2*m^2 + 3*b^2*m + 2*b^2)*x^3 + 2*(a*b*m^2 + 4*a*b*m + 3*a*b)*x^2 + (a^2*m^2 + 5*a^2*m + 6*a^2)*x)*sinh(m*log(x))/(m^3 + 6*m^2 + 11*m + 6)`

**Sympy [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$= \begin{cases} b^2 \log(x) - \frac{b \operatorname{atanh}(\tanh(a+bx))}{x} - \frac{\operatorname{atanh}^2(\tanh(a+bx))}{2x^2} \\ \int \frac{\operatorname{atanh}^2(\tanh(a+bx))}{x^2} dx \\ \int \frac{\operatorname{atanh}^2(\tanh(a+bx))}{x} dx \\ \frac{2b^2 x^3 x^m}{m^3 + 6m^2 + 11m + 6} - \frac{2bm x^2 x^m \operatorname{atanh}(\tanh(a+bx))}{m^3 + 6m^2 + 11m + 6} - \frac{6bx^2 x^m \operatorname{atanh}(\tanh(a+bx))}{m^3 + 6m^2 + 11m + 6} + \frac{m^2 x x^m \operatorname{atanh}^2(\tanh(a+bx))}{m^3 + 6m^2 + 11m + 6} + \frac{5m x x^m \operatorname{atanh}(\tanh(a+bx))}{m^3 + 6m^2 + 11m + 6} \end{cases}$$

input `integrate(x**m*atanh(tanh(b*x+a))**2,x)`

output

```
Piecewise((b**2*log(x) - b*atanh(tanh(a + b*x))/x - atanh(tanh(a + b*x))**
2/(2*x**2), Eq(m, -3)), (Integral(atanh(tanh(a + b*x))**2/x**2, x), Eq(m,
-2)), (Integral(atanh(tanh(a + b*x))**2/x, x), Eq(m, -1)), (2*b**2*x**3*x*
*m/(m**3 + 6*m**2 + 11*m + 6) - 2*b*m*x**2*x**m*atanh(tanh(a + b*x))/(m**3
+ 6*m**2 + 11*m + 6) - 6*b*x**2*x**m*atanh(tanh(a + b*x))/(m**3 + 6*m**2
+ 11*m + 6) + m**2*x*x**m*atanh(tanh(a + b*x))**2/(m**3 + 6*m**2 + 11*m +
6) + 5*m*x*x**m*atanh(tanh(a + b*x))**2/(m**3 + 6*m**2 + 11*m + 6) + 6*x*x
**m*atanh(tanh(a + b*x))**2/(m**3 + 6*m**2 + 11*m + 6), True))
```

### Maxima [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.03

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2b^2x^3x^m}{(m+3)(m+2)(m+1)} - \frac{2bx^2x^m \operatorname{arctanh}(\tanh(bx+a))}{(m+2)(m+1)} + \frac{x^{m+1} \operatorname{arctanh}(\tanh(bx+a))^2}{m+1}$$

input

```
integrate(x^m*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")
```

output

```
2*b^2*x^3*x^m/((m + 3)*(m + 2)*(m + 1)) - 2*b*x^2*x^m*arctanh(tanh(b*x + a
))/((m + 2)*(m + 1)) + x^(m + 1)*arctanh(tanh(b*x + a))^2/(m + 1)
```

### Giac [F]

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int x^m \operatorname{arctanh}(\tanh(bx + a))^2 dx$$

input

```
integrate(x^m*arctanh(tanh(b*x+a))^2,x, algorithm="giac")
```

output

```
integrate(x^m*arctanh(tanh(b*x + a))^2, x)
```

**Mupad [B] (verification not implemented)**

Time = 3.29 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.86

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$= \frac{4b^2 x^m x^3 (m^2 + 3m + 2)}{4m^3 + 24m^2 + 44m + 24}$$

$$+ \frac{x x^m \left( \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx \right)^2 (m^2 + 5m + 6)}{4m^3 + 24m^2 + 44m + 24}$$

$$- \frac{4bx^m x^2 \left( \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx \right) (m^2 + 4m + 3)}{4m^3 + 24m^2 + 44m + 24}$$

input `int(x^m*atanh(tanh(a + b*x))^2,x)`output `(4*b^2*x^m*x^3*(3*m + m^2 + 2))/(44*m + 24*m^2 + 4*m^3 + 24) + (x*x^m*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2*(5*m + m^2 + 6))/(44*m + 24*m^2 + 4*m^3 + 24) - (4*b*x^m*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*(4*m + m^2 + 3))/(44*m + 24*m^2 + 4*m^3 + 24)`**Reduce [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int x^m \operatorname{atanh}(\tanh(bx + a))^2 dx$$

input `int(x^m*atanh(tanh(b*x+a))^2,x)`output `int(x**m*atanh(tanh(a + b*x))**2,x)`

### 3.45 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	435
Mathematica [A] (verified)	435
Rubi [A] (verified)	436
Maple [A] (verified)	437
Fricas [A] (verification not implemented)	437
Sympy [B] (verification not implemented)	438
Maxima [A] (verification not implemented)	438
Giac [A] (verification not implemented)	439
Mupad [B] (verification not implemented)	439
Reduce [F]	439

#### Optimal result

Integrand size = 13, antiderivative size = 42

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{b^2 x^6}{60} - \frac{1}{10} b x^5 \operatorname{arctanh}(\tanh(a + bx)) + \frac{1}{4} x^4 \operatorname{arctanh}(\tanh(a + bx))^2$$

output

```
1/60*b^2*x^6-1/10*b*x^5*arctanh(tanh(b*x+a))+1/4*x^4*arctanh(tanh(b*x+a))^2
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{60} x^4 (b^2 x^2 - 6bx \operatorname{arctanh}(\tanh(a + bx)) + 15 \operatorname{arctanh}(\tanh(a + bx))^2)$$

input

```
Integrate[x^3*ArcTanh[Tanh[a + b*x]]^2,x]
```

output

```
(x^4*(b^2*x^2 - 6*b*x*ArcTanh[Tanh[a + b*x]] + 15*ArcTanh[Tanh[a + b*x]]^2))/60
```



**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{1}{2}b \int x^4 \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{1}{2}b \left( \frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^5 dx}{5} \right)$$

$$\downarrow 15$$

$$\frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{1}{2}b \left( \frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^6}{30} \right)$$

input `Int[x^3*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(x^4*ArcTanh[Tanh[a + b*x]]^2)/4 - (b*(-1/30*(b*x^6) + (x^5*ArcTanh[Tanh[a + b*x]])/5))/2`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 30.85 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

method	result	size
parallelrisch	$\frac{b^2 x^6}{60} - \frac{b x^5 \operatorname{arctanh}(\tanh(bx+a))}{10} + \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^2}{4}$	37
default	$\frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^2}{4} - \frac{b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))}{5} - \frac{b x^6}{30} \right)}{2}$	38
parts	$\frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^2}{4} - \frac{b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))}{5} - \frac{b x^6}{30} \right)}{2}$	38
risch	Expression too large to display	2083

input

```
int(x^3*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
1/60*b^2*x^6-1/10*b*x^5*arctanh(tanh(b*x+a))+1/4*x^4*arctanh(tanh(b*x+a))^2
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{6} b^2 x^6 + \frac{2}{5} abx^5 + \frac{1}{4} a^2 x^4$$

input

```
integrate(x^3*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output  $1/6*b^2*x^6 + 2/5*a*b*x^5 + 1/4*a^2*x^4$

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(37) = 74.

Time = 0.37 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.86

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$= \begin{cases} \frac{x^3 \operatorname{atanh}^3(\tanh(a+bx))}{3b} - \frac{x^2 \operatorname{atanh}^4(\tanh(a+bx))}{4b^2} + \frac{x \operatorname{atanh}^5(\tanh(a+bx))}{10b^3} - \frac{\operatorname{atanh}^6(\tanh(a+bx))}{60b^4} & \text{for } b \neq 0 \\ \frac{x^4 \operatorname{atanh}^2(\tanh(a))}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*atanh(tanh(b*x+a))**2,x)`

output `Piecewise((x**3*atanh(tanh(a + b*x))**3/(3*b) - x**2*atanh(tanh(a + b*x))*  
*4/(4*b**2) + x*atanh(tanh(a + b*x))**5/(10*b**3) - atanh(tanh(a + b*x))**  
6/(60*b**4), Ne(b, 0)), (x**4*atanh(tanh(a))**2/4, True))`

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{60} b^2 x^6 - \frac{1}{10} b x^5 \operatorname{artanh}(\tanh(bx + a))$$

$$+ \frac{1}{4} x^4 \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x^3*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output  $1/60*b^2*x^6 - 1/10*b*x^5*\operatorname{arctanh}(\tanh(b*x + a)) + 1/4*x^4*\operatorname{arctanh}(\tanh(b*x + a))^2$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{6} b^2 x^6 + \frac{2}{5} abx^5 + \frac{1}{4} a^2 x^4$$

input `integrate(x^3*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `1/6*b^2*x^6 + 2/5*a*b*x^5 + 1/4*a^2*x^4`**Mupad [B] (verification not implemented)**

Time = 3.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{b^2 x^6}{60} - \frac{b x^5 \operatorname{atanh}(\tanh(a + bx))}{10} + \frac{x^4 \operatorname{atanh}(\tanh(a + bx))^2}{4}$$

input `int(x^3*atanh(tanh(a + b*x))^2,x)`output `(x^4*atanh(tanh(a + b*x))^2)/4 + (b^2*x^6)/60 - (b*x^5*atanh(tanh(a + b*x)))/10`**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \operatorname{atanh}(\tanh(bx + a))^2 x^3 dx$$

input `int(x^3*atanh(tanh(b*x+a))^2,x)`output `int(atanh(tanh(a + b*x))**2*x**3,x)`

### 3.46 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	440
Mathematica [A] (verified)	440
Rubi [A] (verified)	441
Maple [A] (verified)	442
Fricas [A] (verification not implemented)	442
Sympy [A] (verification not implemented)	443
Maxima [A] (verification not implemented)	443
Giac [A] (verification not implemented)	444
Mupad [B] (verification not implemented)	444
Reduce [F]	444

#### Optimal result

Integrand size = 13, antiderivative size = 42

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{b^2 x^5}{30} - \frac{1}{6} b x^4 \operatorname{arctanh}(\tanh(a + bx)) + \frac{1}{3} x^3 \operatorname{arctanh}(\tanh(a + bx))^2$$

output `1/30*b^2*x^5-1/6*b*x^4*arctanh(tanh(b*x+a))+1/3*x^3*arctanh(tanh(b*x+a))^2`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{30} x^3 (b^2 x^2 - 5bx \operatorname{arctanh}(\tanh(a + bx)) + 10 \operatorname{arctanh}(\tanh(a + bx))^2)$$

input `Integrate[x^2*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(x^3*(b^2*x^2 - 5*b*x*ArcTanh[Tanh[a + b*x]] + 10*ArcTanh[Tanh[a + b*x]]^2))/30`

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{3}b \int x^3 \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{3}b \left( \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^4 dx}{4} \right)$$

$$\downarrow 15$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{3}b \left( \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^5}{20} \right)$$

input `Int[x^2*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(x^3*ArcTanh[Tanh[a + b*x]]^2)/3 - (2*b*(-1/20*(b*x^5) + (x^4*ArcTanh[Tanh[a + b*x]])/4))/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 27.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

method	result	size
parallelrisch	$\frac{b^2 x^5}{30} - \frac{b x^4 \operatorname{arctanh}(\tanh(bx+a))}{6} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^2}{3}$	37
default	$\frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{2b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))}{4} - \frac{b x^5}{20} \right)}{3}$	38
parts	$\frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{2b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))}{4} - \frac{b x^5}{20} \right)}{3}$	38
risch	Expression too large to display	2083

input

```
int(x^2*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
1/30*b^2*x^5-1/6*b*x^4*arctanh(tanh(b*x+a))+1/3*x^3*arctanh(tanh(b*x+a))^2
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{5} b^2 x^5 + \frac{1}{2} abx^4 + \frac{1}{3} a^2 x^3$$

input

```
integrate(x^2*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
1/5*b^2*x^5 + 1/2*a*b*x^4 + 1/3*a^2*x^3
```

**Sympy [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.43

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$= \begin{cases} \frac{x^2 \operatorname{atanh}^3(\tanh(a+bx))}{3b} - \frac{x \operatorname{atanh}^4(\tanh(a+bx))}{6b^2} + \frac{\operatorname{atanh}^5(\tanh(a+bx))}{30b^3} & \text{for } b \neq 0 \\ \frac{x^3 \operatorname{atanh}^2(\tanh(a))}{3} & \text{otherwise} \end{cases}$$

input `integrate(x**2*atanh(tanh(b*x+a))**2,x)`output `Piecewise((x**2*atanh(tanh(a + b*x))**3/(3*b) - x*atanh(tanh(a + b*x))**4/(6*b**2) + atanh(tanh(a + b*x))**5/(30*b**3), Ne(b, 0)), (x**3*atanh(tanh(a))**2/3, True))`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{30} b^2 x^5 - \frac{1}{6} b x^4 \operatorname{artanh}(\tanh(bx + a)) + \frac{1}{3} x^3 \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x^2*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `1/30*b^2*x^5 - 1/6*b*x^4*arctanh(tanh(b*x + a)) + 1/3*x^3*arctanh(tanh(b*x + a))^2`



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{5} b^2 x^5 + \frac{1}{2} abx^4 + \frac{1}{3} a^2 x^3$$

input `integrate(x^2*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `1/5*b^2*x^5 + 1/2*a*b*x^4 + 1/3*a^2*x^3`

**Mupad [B] (verification not implemented)**

Time = 3.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{b^2 x^5}{30} - \frac{b x^4 \operatorname{atanh}(\tanh(a + bx))}{6} + \frac{x^3 \operatorname{atanh}(\tanh(a + bx))^2}{3}$$

input `int(x^2*atanh(tanh(a + b*x))^2,x)`

output `(x^3*atanh(tanh(a + b*x))^2)/3 + (b^2*x^5)/30 - (b*x^4*atanh(tanh(a + b*x)))/6`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \operatorname{atanh}(\tanh(bx + a))^2 x^2 dx$$

input `int(x^2*atanh(tanh(b*x+a))^2,x)`

output `int(atanh(tanh(a + b*x))**2*x**2,x)`

### 3.47 $\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	445
Mathematica [B] (verified)	445
Rubi [A] (verified)	446
Maple [A] (verified)	447
Fricas [A] (verification not implemented)	447
Sympy [A] (verification not implemented)	448
Maxima [A] (verification not implemented)	448
Giac [A] (verification not implemented)	448
Mupad [B] (verification not implemented)	449
Reduce [F]	449

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x \operatorname{arctanh}(\tanh(a + bx))^3}{3b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{12b^2}$$

output `1/3*x*arctanh(tanh(b*x+a))^3/b-1/12*arctanh(tanh(b*x+a))^4/b^2`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 74 vs. 2(34) = 68.

Time = 0.17 (sec) , antiderivative size = 74, normalized size of antiderivative = 2.18

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{(a + bx) (-((3a - bx)(a + bx)^2) + 4(2a^2 + abx - b^2x^2) \operatorname{arctanh}(\tanh(a + bx)) - 6(a - bx) \operatorname{arctanh}(\tanh(a + bx)))}{12b^2}$$

input `Integrate[x*ArcTanh[Tanh[a + b*x]]^2,x]`

output `((a + b*x)*(-((3*a - b*x)*(a + b*x)^2) + 4*(2*a^2 + a*b*x - b^2*x^2)*ArcTanh[Tanh[a + b*x]] - 6*(a - b*x)*ArcTanh[Tanh[a + b*x]]^2))/(12*b^2)`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^3}{3b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^3 dx}{3b}$$

$$\downarrow 2588$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^3}{3b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^3 d \operatorname{arctanh}(\tanh(a + bx))}{3b^2}$$

$$\downarrow 15$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^3}{3b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{12b^2}$$

input `Int[x*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(x*ArcTanh[Tanh[a + b*x]]^3)/(3*b) - ArcTanh[Tanh[a + b*x]]^4/(12*b^2)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n])
```

**Maple [A] (verified)**

Time = 26.16 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.09

method	result	size
parallelrisch	$\frac{b^2 x^4}{12} - \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))b}{3} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^2}{2}$	37
default	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^2}{2} - b \left( -\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))}{3} \right)$	38
parts	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^2}{2} - b \left( -\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))}{3} \right)$	38
risch	Expression too large to display	2083

input

```
int(x*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
1/12*b^2*x^4-1/3*x^3*arctanh(tanh(b*x+a))*b+1/2*x^2*arctanh(tanh(b*x+a))^2
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{4} b^2 x^4 + \frac{2}{3} abx^3 + \frac{1}{2} a^2 x^2$$

input

```
integrate(x*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
1/4*b^2*x^4 + 2/3*a*b*x^3 + 1/2*a^2*x^2
```

**Sympy [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \begin{cases} \frac{x \operatorname{atanh}^3(\tanh(a + bx))}{3b} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{12b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^2(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**2,x)`output `Piecewise((x*atanh(tanh(a + b*x))**3/(3*b) - atanh(tanh(a + b*x))**4/(12*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**2/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{12} b^2 x^4 - \frac{1}{3} b x^3 \operatorname{artanh}(\tanh(bx + a)) + \frac{1}{2} x^2 \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `1/12*b^2*x^4 - 1/3*b*x^3*arctanh(tanh(b*x + a)) + 1/2*x^2*arctanh(tanh(b*x + a))^2`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{4} b^2 x^4 + \frac{2}{3} a b x^3 + \frac{1}{2} a^2 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output  $1/4*b^2*x^4 + 2/3*a*b*x^3 + 1/2*a^2*x^2$

### Mupad [B] (verification not implemented)

Time = 3.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{b^2 x^4}{12} - \frac{b x^3 \operatorname{atanh}(\tanh(a + bx))}{3} + \frac{x^2 \operatorname{atanh}(\tanh(a + bx))^2}{2}$$

input `int(x*atanh(tanh(a + b*x))^2,x)`

output  $(x^2*\operatorname{atanh}(\tanh(a + b*x))^2)/2 + (b^2*x^4)/12 - (b*x^3*\operatorname{atanh}(\tanh(a + b*x)))/3$

### Reduce [F]

$$\int x \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \operatorname{atanh}(\tanh(bx + a))^2 x dx$$

input `int(x*atanh(tanh(b*x+a))^2,x)`

output `int(atanh(tanh(a + b*x))**2*x,x)`

### 3.48 $\int \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	450
Mathematica [A] (verified)	450
Rubi [A] (verified)	451
Maple [A] (verified)	452
Fricas [A] (verification not implemented)	452
Sympy [A] (verification not implemented)	452
Maxima [B] (verification not implemented)	453
Giac [A] (verification not implemented)	453
Mupad [B] (verification not implemented)	454
Reduce [B] (verification not implemented)	454

#### Optimal result

Integrand size = 9, antiderivative size = 16

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3b}$$

output `1/3*arctanh(tanh(b*x+a))^3/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3b}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^2,x]`

output `ArcTanh[Tanh[a + b*x]]^3/(3*b)`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2588$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^2 d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow 15$$

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^2,x]`

output `ArcTanh[Tanh[a + b*x]]^3/(3*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



**Maple [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{3b}$	15
default	$\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{3b}$	15
parallelrisch	$\frac{b^2x^3}{3} - x^2 \operatorname{arctanh}(\tanh(bx+a)) b + x \operatorname{arctanh}(\tanh(bx+a))^2$	34
risch	Expression too large to display	6270

input `int(arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output `1/3*arctanh(tanh(b*x+a))^3/b`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{3} b^2 x^3 + abx^2 + a^2 x$$

input `integrate(arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `1/3*b^2*x^3 + a*b*x^2 + a^2*x`

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \begin{cases} \frac{\operatorname{atanh}^3(\tanh(a+bx))}{3b} & \text{for } b \neq 0 \\ x \operatorname{atanh}^2(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**2,x)`

output `Piecewise((atanh(tanh(a + b*x))**3/(3*b), Ne(b, 0)), (x*atanh(tanh(a))**2, True))`

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(14) = 28$ .

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{3} b^2 x^3 - bx^2 \operatorname{arctanh}(\tanh(bx + a)) + x \operatorname{arctanh}(\tanh(bx + a))^2$$

input `integrate(arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `1/3*b^2*x^3 - b*x^2*arctanh(tanh(b*x + a)) + x*arctanh(tanh(b*x + a))^2`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{1}{3} b^2 x^3 + abx^2 + a^2 x$$

input `integrate(arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `1/3*b^2*x^3 + a*b*x^2 + a^2*x`

**Mupad [B] (verification not implemented)**

Time = 3.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \operatorname{arctanh}(\tanh(a+bx))^2 dx = \frac{b^2 x^3}{3} - b x^2 \operatorname{atanh}(\tanh(a+bx)) + x \operatorname{atanh}(\tanh(a+bx))^2$$

input `int(atanh(tanh(a + b*x))^2,x)`

output `x*atanh(tanh(a + b*x))^2 + (b^2*x^3)/3 - b*x^2*atanh(tanh(a + b*x))`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\tanh(a+bx))^2 dx = \frac{\operatorname{atanh}(\tanh(bx+a))^3}{3b}$$

input `int(atanh(tanh(b*x+a))^2,x)`

output `atanh(tanh(a + b*x))**3/(3*b)`

### 3.49 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x} dx$

Optimal result	455
Mathematica [A] (verified)	455
Rubi [A] (verified)	456
Maple [A] (verified)	457
Fricas [A] (verification not implemented)	458
Sympy [F]	458
Maxima [A] (verification not implemented)	458
Giac [A] (verification not implemented)	459
Mupad [B] (verification not implemented)	459
Reduce [F]	460

#### Optimal result

Integrand size = 13, antiderivative size = 49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = -bx(bx - \operatorname{arctanh}(\tanh(a + bx))) + \frac{1}{2}\operatorname{arctanh}(\tanh(a + bx))^2 + (bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \log(x)$$

output

```
-b*x*(b*x-arctanh(tanh(b*x+a)))+1/2*arctanh(tanh(b*x+a))^2+(b*x-arctanh(tanh(b*x+a)))^2*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \frac{1}{2}(a + bx)^2 - (a + bx)(a + 2bx - 2\operatorname{arctanh}(\tanh(a + bx))) + (-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log(bx)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^2/x,x]
```

output

$$(a + b*x)^2/2 - (a + b*x)*(a + 2*b*x - 2*ArcTanh[Tanh[a + b*x]]) + -(b*x) + ArcTanh[Tanh[a + b*x]]^2*Log[b*x]$$

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx$$

↓ 2590

$$\frac{1}{2}\operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx$$

↓ 2589

$$\frac{1}{2}\operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right)$$

↓ 14

$$\frac{1}{2}\operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx)))(bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx))))$$

input

$$\text{Int}[\text{ArcTanh}[\text{Tanh}[a + b*x]]^2/x, x]$$

output

$$\text{ArcTanh}[\text{Tanh}[a + b*x]]^2/2 - (b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])*(b*x - (b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])*\text{Log}[x])$$

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

## Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.37

method	result
default	$\ln(x) \operatorname{arctanh}(\tanh(bx + a))^2 - 2b \left( b \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + a(x \ln(x) - x) + \operatorname{arctanh}(\tanh(bx + a)) \right)$
parts	$\ln(x) \operatorname{arctanh}(\tanh(bx + a))^2 - 2b \left( b \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + a(x \ln(x) - x) + \operatorname{arctanh}(\tanh(bx + a)) \right)$
risch	$\ln(x) \ln(e^{bx+a})^2 + b^2 x^2 \ln(x) - \frac{3b^2 x^2}{2} - 2b \ln(e^{bx+a}) \ln(x) x + 2b \ln(e^{bx+a}) x - \frac{\pi^2 \left( \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}}\right) \right)}{2}$

input `int(arctanh(tanh(b*x+a))^2/x,x,method=_RETURNVERBOSE)`

output `ln(x)*arctanh(tanh(b*x+a))^2-2*b*(b*(1/2*x^2*ln(x)-1/4*x^2)+a*(x*ln(x)-x)+arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \frac{1}{2} b^2 x^2 + 2 abx + a^2 \log(x)$$

input `integrate(arctanh(tanh(b*x+a))^2/x,x, algorithm="fricas")`

output `1/2*b^2*x^2 + 2*a*b*x + a^2*log(x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \int \frac{\operatorname{atanh}^2(\tanh(a + bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))**2/x,x)`

output `Integral(atanh(tanh(a + b*x))**2/x, x)`

**Maxima [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \frac{1}{2} b^2 x^2 + 2 abx + a^2 \log(x)$$

input `integrate(arctanh(tanh(b*x+a))^2/x,x, algorithm="maxima")`

output `1/2*b^2*x^2 + 2*a*b*x + a^2*log(x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \frac{1}{2} b^2 x^2 + 2 abx + a^2 \log(|x|)$$

input `integrate(arctanh(tanh(b*x+a))^2/x,x, algorithm="giac")`

output `1/2*b^2*x^2 + 2*a*b*x + a^2*log(abs(x))`

**Mupad [B] (verification not implemented)**

Time = 3.27 (sec) , antiderivative size = 183, normalized size of antiderivative = 3.73

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx \\ &= \ln(x) \left( \frac{\left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2}{4} \right. \\ & \quad \left. - a \left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right) + a^2 \right) \\ & \quad + \frac{b^2 x^2}{2} - bx \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right) \end{aligned}$$

input `int(atanh(tanh(a + b*x))^2/x,x)`

output `log(x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/4 - a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + a^2) + (b^2*x^2)/2 - b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)`



**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^2}{x} dx$$

input `int(atanh(tanh(b*x+a))^2/x,x)`

output `int(atanh(tanh(a + b*x))**2/x,x)`

### 3.50 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^2} dx$

Optimal result . . . . .	461
Mathematica [A] (verified) . . . . .	461
Rubi [A] (verified) . . . . .	462
Maple [A] (verified) . . . . .	463
Fricas [A] (verification not implemented) . . . . .	463
Sympy [F] . . . . .	464
Maxima [A] (verification not implemented) . . . . .	464
Giac [A] (verification not implemented) . . . . .	464
Mupad [B] (verification not implemented) . . . . .	465
Reduce [F] . . . . .	466

#### Optimal result

Integrand size = 13, antiderivative size = 39

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = 2b^2x - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} - 2b(bx - \operatorname{arctanh}(\tanh(a + bx))) \log(x)$$

output

```
2*b^2*x-arctanh(tanh(b*x+a))^2/x-2*b*(b*x-arctanh(tanh(b*x+a)))*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = -\frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} - 2b^2x \log(x) + 2b \operatorname{arctanh}(\tanh(a + bx))(1 + \log(x))$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^2/x^2,x]
```

output

```
-(ArcTanh[Tanh[a + b*x]]^2/x) - 2*b^2*x*Log[x] + 2*b*ArcTanh[Tanh[a + b*x]]*(1 + Log[x])
```

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx$$

$$\downarrow \text{2599}$$

$$2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x}$$

$$\downarrow \text{2589}$$

$$2b \left( bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x}$$

$$\downarrow \text{14}$$

$$2b(bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx)))) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x}$$

input `Int[ArcTanh[Tanh[a + b*x]]^2/x^2,x]`

output `-(ArcTanh[Tanh[a + b*x]]^2/x) + 2*b*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x])`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x) \operatorname{arctanh}(\tanh(bx+a)) - b(x \ln(x) - x))$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x) \operatorname{arctanh}(\tanh(bx+a)) - b(x \ln(x) - x))$
risch	$-\frac{\ln(e^{bx+a})^2}{x} - 2 \ln(x) x b^2 + 2 \ln(x) \ln(e^{bx+a}) b + 2b^2 x + \frac{\pi^2 \left( \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) \right)}{e^{2bx+2a}+1}$

input

```
int(arctanh(tanh(b*x+a))^2/x^2,x,method=_RETURNVERBOSE)
```

output

```
-arctanh(tanh(b*x+a))^2/x+2*b*(ln(x)*arctanh(tanh(b*x+a))-b*(x*ln(x)-x))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = \frac{b^2 x^2 + 2 abx \log(x) - a^2}{x}$$

input

```
integrate(arctanh(tanh(b*x+a))^2/x^2,x, algorithm="fricas")
```

output

```
(b^2*x^2 + 2*a*b*x*log(x) - a^2)/x
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = \int \frac{\operatorname{atanh}^2(\tanh(a + bx))}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**2/x**2,x)`

output `Integral(atanh(tanh(a + b*x))**2/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.38

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = 2b \operatorname{arctanh}(\tanh(bx + a)) \log(x) - 2 \left( b \left( x + \frac{a}{b} \right) \log(x) - b \left( x + \frac{a \log(x)}{b} \right) \right) b - \frac{\operatorname{arctanh}(\tanh(bx + a))^2}{x}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^2,x, algorithm="maxima")`

output `2*b*arctanh(tanh(b*x + a))*log(x) - 2*(b*(x + a/b)*log(x) - b*(x + a*log(x)/b))*b - arctanh(tanh(b*x + a))^2/x`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = b^2 x + 2ab \log(|x|) - \frac{a^2}{x}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^2,x, algorithm="giac")`

output  $b^2x + 2ab \log(\text{abs}(x)) - a^2/x$

### Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 198, normalized size of antiderivative = 5.08

$$\int \frac{\text{arctanh}(\tanh(a + bx))^2}{x^2} dx = b \ln \left( \frac{e^{2bx}}{e^{2a} e^{2bx} + 1} \right) - \frac{\ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)^2}{4x}$$

$$- b \ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right) - \frac{\ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right)^2}{4x}$$

$$+ b \ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) \ln(x) - 2b^2 x \ln(x)$$

$$- b \ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right) \ln(x)$$

$$+ \frac{\ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right) \ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)}{2x}$$

input `int(atanh(tanh(a + b*x))^2/x^2,x)`

output `b*log(exp(2*b*x)/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2/(4*x) - b*log(1/(exp(2*a)*exp(2*b*x) + 1)) - log(1/(exp(2*a)*exp(2*b*x) + 1))^2/(4*x) + b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x) - 2*b^2*x*log(x) - b*log(1/(exp(2*a)*exp(2*b*x) + 1))*log(x) + (log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(2*x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^2}{x^2} dx$$

input `int(atanh(tanh(b*x+a))^2/x^2,x)`

output `int(atanh(tanh(a + b*x))**2/x**2,x)`

### 3.51 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^3} dx$

Optimal result . . . . .	467
Mathematica [A] (verified) . . . . .	467
Rubi [A] (verified) . . . . .	468
Maple [A] (verified) . . . . .	469
Fricas [A] (verification not implemented) . . . . .	470
Sympy [A] (verification not implemented) . . . . .	470
Maxima [A] (verification not implemented) . . . . .	470
Giac [A] (verification not implemented) . . . . .	471
Mupad [B] (verification not implemented) . . . . .	471
Reduce [B] (verification not implemented) . . . . .	471

#### Optimal result

Integrand size = 13, antiderivative size = 36

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^3} dx = -\frac{b \operatorname{arctanh}(\tanh(a+bx))}{x} - \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{2x^2} + b^2 \log(x)$$

output

```
-b*arctanh(tanh(b*x+a))/x-1/2*arctanh(tanh(b*x+a))^2/x^2+b^2*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^3} dx = -\frac{2bx \operatorname{arctanh}(\tanh(a+bx)) + \operatorname{arctanh}(\tanh(a+bx))^2 - b^2 x^2 (3 + 2 \log(x))}{2x^2}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^2/x^3,x]
```



output

$$-1/2*(2*b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2 - b^2*x^2*(3 + 2*Log[x]))/x^2$$

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx \\ & \quad \downarrow \text{2599} \\ & b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \\ & \quad \downarrow \text{2599} \\ & b \left( b \int \frac{1}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \\ & \quad \downarrow \text{14} \\ & b \left( b \log(x) - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \end{aligned}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^2/x^3,x]
```

output

```
-1/2*ArcTanh[Tanh[a + b*x]]^2/x^2 + b*(-(ArcTanh[Tanh[a + b*x]]/x) + b*Log[x])
```

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.97

method	result	size
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2x^2} + b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} + b \ln(x)\right)$	35
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2x^2} + b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} + b \ln(x)\right)$	35
parallelrisch	$\frac{2b^2x^2 \ln(x) - 2x \operatorname{arctanh}(\tanh(bx+a))b - \operatorname{arctanh}(\tanh(bx+a))^2}{2x^2}$	39
risch	Expression too large to display	1974

input `int(arctanh(tanh(b*x+a))^2/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arctanh(tanh(b*x+a))^2/x^2+b*(-arctanh(tanh(b*x+a))/x+b*ln(x))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = \frac{2b^2x^2 \log(x) - 4abx - a^2}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^3,x, algorithm="fricas")`output `1/2*(2*b^2*x^2*log(x) - 4*a*b*x - a^2)/x^2`**Sympy [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = b^2 \log(x) - \frac{b \operatorname{atanh}(\tanh(a + bx))}{x} - \frac{\operatorname{atanh}^2(\tanh(a + bx))}{2x^2}$$

input `integrate(atanh(tanh(b*x+a))**2/x**3,x)`output `b**2*log(x) - b*atanh(tanh(a + b*x))/x - atanh(tanh(a + b*x))**2/(2*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = b^2 \log(x) - \frac{b \operatorname{artanh}(\tanh(bx + a))}{x} - \frac{\operatorname{artanh}(\tanh(bx + a))^2}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^3,x, algorithm="maxima")`output `b^2*log(x) - b*arctanh(tanh(b*x + a))/x - 1/2*arctanh(tanh(b*x + a))^2/x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.61

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = b^2 \log(|x|) - \frac{4abx + a^2}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^3,x, algorithm="giac")`output `b^2*log(abs(x)) - 1/2*(4*a*b*x + a^2)/x^2`**Mupad [B] (verification not implemented)**

Time = 3.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = b^2 \ln(x) - \frac{\operatorname{atanh}(\tanh(a + bx))^2}{2} + \frac{bx \operatorname{atanh}(\tanh(a + bx))}{x^2}$$

input `int(atanh(tanh(a + b*x))^2/x^3,x)`output `b^2*log(x) - (atanh(tanh(a + b*x))^2/2 + b*x*atanh(tanh(a + b*x)))/x^2`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx = \frac{-\operatorname{atanh}(\tanh(bx + a))^2 - 2\operatorname{atanh}(\tanh(bx + a))bx + 2\log(x)b^2x^2}{2x^2}$$

input `int(atanh(tanh(b*x+a))^2/x^3,x)`output `( - atanh(tanh(a + b*x))**2 - 2*atanh(tanh(a + b*x))*b*x + 2*log(x)*b**2*x**2)/(2*x**2)`

### 3.52 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^4} dx$

Optimal result	472
Mathematica [A] (verified)	472
Rubi [A] (verified)	473
Maple [A] (verified)	473
Fricas [A] (verification not implemented)	474
Sympy [A] (verification not implemented)	474
Maxima [A] (verification not implemented)	475
Giac [A] (verification not implemented)	475
Mupad [B] (verification not implemented)	475
Reduce [B] (verification not implemented)	476

#### Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^4} dx = \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{3x^3(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `1/3*arctanh(tanh(b*x+a))^3/x^3/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^4} dx = -\frac{b^2x^2 + bx\operatorname{arctanh}(\tanh(a+bx)) + \operatorname{arctanh}(\tanh(a+bx))^2}{3x^3}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^2/x^4,x]`

output `-1/3*(b^2*x^2 + b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2)/x^3`

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^2/x^4,x]`

output `ArcTanh[Tanh[a + b*x]]^3/(3*x^3*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

method	result	size
parallelrisch	$-\frac{b^2x^2+x \operatorname{arctanh}(\tanh(bx+a))b+\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3}$	33
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b\left(-\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2}\right)}{3}$	38
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b\left(-\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2}\right)}{3}$	38
risch	Expression too large to display	1978

input `int(arctanh(tanh(b*x+a))^2/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*(b^2*x^2+x*arctanh(tanh(b*x+a))*b+arctanh(tanh(b*x+a))^2)/x^3`

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx = -\frac{3b^2x^2 + 3abx + a^2}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^4,x, algorithm="fricas")`

output `-1/3*(3*b^2*x^2 + 3*a*b*x + a^2)/x^3`

### Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx = -\frac{b^2}{3x} - \frac{b \operatorname{atanh}(\tanh(a + bx))}{3x^2} - \frac{\operatorname{atanh}^2(\tanh(a + bx))}{3x^3}$$

input `integrate(atanh(tanh(b*x+a))**2/x**4,x)`

output `-b**2/(3*x) - b*atanh(tanh(a + b*x))/(3*x**2) - atanh(tanh(a + b*x))**2/(3*x**3)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx$$

$$= -\frac{b^2}{3x} - \frac{b \operatorname{arctanh}(\tanh(bx + a))}{3x^2} - \frac{\operatorname{arctanh}(\tanh(bx + a))^2}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^4,x, algorithm="maxima")`output `-1/3*b^2/x - 1/3*b*arctanh(tanh(b*x + a))/x^2 - 1/3*arctanh(tanh(b*x + a))^2/x^3`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx = -\frac{3b^2x^2 + 3abx + a^2}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^4,x, algorithm="giac")`output `-1/3*(3*b^2*x^2 + 3*a*b*x + a^2)/x^3`**Mupad [B] (verification not implemented)**

Time = 3.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx$$

$$= -\frac{b^2x^2 + bx \operatorname{atanh}(\tanh(a + bx)) + \operatorname{atanh}(\tanh(a + bx))^2}{3x^3}$$

input `int(atanh(tanh(a + b*x))^2/x^4,x)`



output  $-(\operatorname{atanh}(\tanh(a + b*x))^2 + b^2*x^2 + b*x*\operatorname{atanh}(\tanh(a + b*x)))/(3*x^3)$

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx$$

$$= \frac{-\operatorname{atanh}(\tanh(bx + a))^2 - \operatorname{atanh}(\tanh(bx + a))bx - b^2x^2}{3x^3}$$

input `int(atanh(tanh(b*x+a))^2/x^4,x)`

output  $( - (\operatorname{atanh}(\tanh(a + b*x))^{**2} + \operatorname{atanh}(\tanh(a + b*x))*b*x + b^{**2}*x^{**2}))/ (3*x^{**3})$

### 3.53 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^5} dx$

Optimal result	477
Mathematica [A] (verified)	477
Rubi [A] (verified)	478
Maple [A] (verified)	479
Fricas [A] (verification not implemented)	480
Sympy [A] (verification not implemented)	480
Maxima [A] (verification not implemented)	480
Giac [A] (verification not implemented)	481
Mupad [B] (verification not implemented)	481
Reduce [B] (verification not implemented)	482

#### Optimal result

Integrand size = 13, antiderivative size = 42

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^5} dx = -\frac{b^2}{12x^2} - \frac{b \operatorname{arctanh}(\tanh(a+bx))}{6x^3} - \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{4x^4}$$

output `-1/12*b^2/x^2-1/6*b*arctanh(tanh(b*x+a))/x^3-1/4*arctanh(tanh(b*x+a))^2/x^4`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^5} dx = -\frac{b^2x^2 + 2bx \operatorname{arctanh}(\tanh(a+bx)) + 3 \operatorname{arctanh}(\tanh(a+bx))^2}{12x^4}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^2/x^5,x]`

output

```
-1/12*(b^2*x^2 + 2*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2
)/x^4
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.52, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx$$

↓ 2602

$$\frac{b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^4} dx}{4(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{4x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^3}{4x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{b \operatorname{arctanh}(\tanh(a + bx))^3}{12x^3(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^2/x^5, x]
```

output

```
(b*ArcTanh[Tanh[a + b*x]]^3)/(12*x^3*(b*x - ArcTanh[Tanh[a + b*x]])^2) + A
rcTanh[Tanh[a + b*x]]^3/(4*x^4*(b*x - ArcTanh[Tanh[a + b*x]]))
```

## Definitions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

## Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

method	result	size
parallelrisch	$-\frac{b^2 x^2 + 2x \operatorname{arctanh}(\tanh(bx+a))b + 3 \operatorname{arctanh}(\tanh(bx+a))^2}{12x^4}$	36
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b\left(-\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^3}\right)}{2}$	38
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b\left(-\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^3}\right)}{2}$	38
risch	Expression too large to display	1978

input `int(arctanh(tanh(b*x+a))^2/x^5,x,method=_RETURNVERBOSE)`

output `-1/12*(b^2*x^2+2*x*arctanh(tanh(b*x+a))*b+3*arctanh(tanh(b*x+a))^2)/x^4`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx = -\frac{6b^2x^2 + 8abx + 3a^2}{12x^4}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^5,x, algorithm="fricas")`output `-1/12*(6*b^2*x^2 + 8*a*b*x + 3*a^2)/x^4`**Sympy [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93

$$\begin{aligned} \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx \\ = -\frac{b^2}{12x^2} - \frac{b \operatorname{atanh}(\tanh(a + bx))}{6x^3} - \frac{\operatorname{atanh}^2(\tanh(a + bx))}{4x^4} \end{aligned}$$

input `integrate(atanh(tanh(b*x+a))**2/x**5,x)`output `-b**2/(12*x**2) - b*atanh(tanh(a + b*x))/(6*x**3) - atanh(tanh(a + b*x))**2/(4*x**4)`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx = -\frac{b^2}{12x^2} - \frac{b \operatorname{artanh}(\tanh(bx + a))}{6x^3} - \frac{\operatorname{artanh}(\tanh(bx + a))^2}{4x^4}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^5,x, algorithm="maxima")`

output

$$-1/12*b^2/x^2 - 1/6*b*arctanh(tanh(b*x + a))/x^3 - 1/4*arctanh(tanh(b*x + a))^2/x^4$$

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx = -\frac{6b^2x^2 + 8abx + 3a^2}{12x^4}$$

input

```
integrate(arctanh(tanh(b*x+a))^2/x^5,x, algorithm="giac")
```

output

$$-1/12*(6*b^2*x^2 + 8*a*b*x + 3*a^2)/x^4$$

**Mupad [B] (verification not implemented)**

Time = 3.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx \\ &= -\frac{\operatorname{atanh}(\tanh(a + bx))^2}{4x^4} - \frac{b^2}{12x^2} - \frac{b \operatorname{atanh}(\tanh(a + bx))}{6x^3} \end{aligned}$$

input

```
int(atanh(tanh(a + b*x))^2/x^5,x)
```

output

$$- \operatorname{atanh}(\tanh(a + b*x))^2/(4*x^4) - b^2/(12*x^2) - (b*\operatorname{atanh}(\tanh(a + b*x)))/(6*x^3)$$

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^5} dx$$
$$= \frac{-3 \operatorname{atanh}(\tanh(bx + a))^2 - 2 \operatorname{atanh}(\tanh(bx + a)) bx - b^2 x^2}{12x^4}$$

input `int(atanh(tanh(b*x+a))^2/x^5,x)`output `( - 3*atanh(tanh(a + b*x))**2 - 2*atanh(tanh(a + b*x))*b*x - b**2*x**2)/(12*x**4)`

### 3.54 $\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	483
Mathematica [A] (verified)	484
Rubi [A] (verified)	484
Maple [A] (verified)	486
Fricas [B] (verification not implemented)	486
Sympy [F]	487
Maxima [A] (verification not implemented)	488
Giac [F]	489
Mupad [B] (verification not implemented)	489
Reduce [F]	490

#### Optimal result

Integrand size = 13, antiderivative size = 110

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{6b^3 x^{4+m}}{(1+m)(24+26m+9m^2+m^3)} + \frac{6b^2 x^{3+m} \operatorname{arctanh}(\tanh(a + bx))}{6+11m+6m^2+m^3} - \frac{3bx^{2+m} \operatorname{arctanh}(\tanh(a + bx))^2}{2+3m+m^2} + \frac{x^{1+m} \operatorname{arctanh}(\tanh(a + bx))^3}{1+m}$$

output

```
-6*b^3*x^(4+m)/(1+m)/(m^3+9*m^2+26*m+24)+6*b^2*x^(3+m)*arctanh(tanh(b*x+a))
)/(m^3+6*m^2+11*m+6)-3*b*x^(2+m)*arctanh(tanh(b*x+a))^2/(m^2+3*m+2)+x^(1+m)
)*arctanh(tanh(b*x+a))^3/(1+m)
```



**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.88

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$= \frac{x^{1+m}(-6b^3x^3 + 6b^2(4+m)x^2 \operatorname{arctanh}(\tanh(a + bx)) - 3b(12 + 7m + m^2)x \operatorname{arctanh}(\tanh(a + bx))^2 + (1+m)(2+m)(3+m)(4+m))}{(1+m)(2+m)(3+m)(4+m)}$$

input

```
Integrate[x^m*ArcTanh[Tanh[a + b*x]]^3,x]
```

output

```
(x^(1 + m)*(-6*b^3*x^3 + 6*b^2*(4 + m)*x^2*ArcTanh[Tanh[a + b*x]] - 3*b*(1
2 + 7*m + m^2)*x*ArcTanh[Tanh[a + b*x]]^2 + (24 + 26*m + 9*m^2 + m^3)*ArcT
anh[Tanh[a + b*x]]^3))/((1 + m)*(2 + m)*(3 + m)*(4 + m))
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^3}{m+1} - \frac{3b \int x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^2 dx}{m+1}$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^3}{m+1} - \frac{3b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a + bx))^2}{m+2} - \frac{2b \int x^{m+2} \operatorname{arctanh}(\tanh(a + bx)) dx}{m+2} \right)}{m+1}$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^3}{m+1} - \frac{3b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))^2}{m+2} - \frac{2b \left( \frac{x^{m+3} \operatorname{arctanh}(\tanh(a+bx))}{m+3} - \frac{b \int x^{m+3} dx}{m+3} \right)}{m+2} \right)}{m+1}$$

↓ 15

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^3}{m+1} - \frac{3b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))^2}{m+2} - \frac{2b \left( \frac{x^{m+3} \operatorname{arctanh}(\tanh(a+bx))}{m+3} - \frac{bx^{m+4}}{(m+3)(m+4)} \right)}{m+2} \right)}{m+1}$$

input `Int[x^m*ArcTanh[Tanh[a + b*x]]^3,x]`

output `(x^(1 + m)*ArcTanh[Tanh[a + b*x]]^3)/(1 + m) - (3*b*((x^(2 + m)*ArcTanh[Tanh[a + b*x]]^2)/(2 + m) - (2*b*(-((b*x^(4 + m))/((3 + m)*(4 + m)))) + (x^(3 + m)*ArcTanh[Tanh[a + b*x]])/(3 + m)))/(2 + m))/(1 + m)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 4.47 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.61

method	result
default	$\frac{b^3 x^4 e^{m \ln(x)}}{4+m} + \frac{(a^3 + 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3)}{1+m}$
parallelrisch	$-\frac{24b^2 \operatorname{arctanh}(\tanh(bx+a))x^m x^3 + 36b \operatorname{arctanh}(\tanh(bx+a))^2 x^2 x^m - x x^m \operatorname{arctanh}(\tanh(bx+a))^3 m^3 - 9x x^m \operatorname{arctanh}(\tanh(bx+a))}{(4+m)^3}$
risch	Expression too large to display

input `int(x^m*arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`output 
$$\frac{b^3}{(4+m)}x^4 \exp(m \ln(x)) + \frac{(a^3 + 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3)}{(1+m)}x \exp(m \ln(x)) + \frac{3b(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2)}{(2+m)}x^2 \exp(m \ln(x)) + \frac{3b^2(\operatorname{arctanh}(\tanh(bx+a)) - bx)}{(3+m)}x^3 \exp(m \ln(x))$$
**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 300 vs. 2(110) = 220.

Time = 0.08 (sec) , antiderivative size = 300, normalized size of antiderivative = 2.73

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{((b^3 m^3 + 6 b^3 m^2 + 11 b^3 m + 6 b^3)x^4 + 3(ab^2 m^3 + 7 ab^2 m^2 + 14 ab^2 m + 8 ab^2)x^3 + 3(a^2 b m^3 + 8 a^2 b m^2 + 3 a^2 b m + 3 a^2 b)x^2 + 3(a^2 b m^2 + 4 a^2 b m + 3 a^2 b)x + 3 a^2 b)}{(4+m)^3}$$

input `integrate(x^m*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```
((b^3*m^3 + 6*b^3*m^2 + 11*b^3*m + 6*b^3)*x^4 + 3*(a*b^2*m^3 + 7*a*b^2*m^2 + 14*a*b^2*m + 8*a*b^2)*x^3 + 3*(a^2*b*m^3 + 8*a^2*b*m^2 + 19*a^2*b*m + 12*a^2*b)*x^2 + (a^3*m^3 + 9*a^3*m^2 + 26*a^3*m + 24*a^3)*x)*cosh(m*log(x)) + ((b^3*m^3 + 6*b^3*m^2 + 11*b^3*m + 6*b^3)*x^4 + 3*(a*b^2*m^3 + 7*a*b^2*m^2 + 14*a*b^2*m + 8*a*b^2)*x^3 + 3*(a^2*b*m^3 + 8*a^2*b*m^2 + 19*a^2*b*m + 12*a^2*b)*x^2 + (a^3*m^3 + 9*a^3*m^2 + 26*a^3*m + 24*a^3)*x)*sinh(m*log(x)))/(m^4 + 10*m^3 + 35*m^2 + 50*m + 24)
```

SymPy [F]

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$= \begin{cases} b^3 \log(x) - \frac{b^2 \operatorname{atanh}(\tanh(a+bx))}{x} - \frac{b \operatorname{atanh}^2(\tanh(a+bx))}{2x^2} - \frac{\operatorname{atanh}^3(\tanh(a+bx))}{3x^3} \\ \int \frac{\operatorname{atanh}^3(\tanh(a+bx))}{x^3} dx \\ \int \frac{\operatorname{atanh}^3(\tanh(a+bx))}{x^2} dx \\ \int \frac{\operatorname{atanh}^3(\tanh(a+bx))}{x} dx \\ -\frac{6b^3x^4x^m}{m^4+10m^3+35m^2+50m+24} + \frac{6b^2mx^3x^m \operatorname{atanh}(\tanh(a+bx))}{m^4+10m^3+35m^2+50m+24} + \frac{24b^2x^3x^m \operatorname{atanh}(\tanh(a+bx))}{m^4+10m^3+35m^2+50m+24} - \frac{3bm^2x^2x^m \operatorname{atanh}^2(\tanh(a+bx))}{m^4+10m^3+35m^2+50m+24} \end{cases}$$

input

```
integrate(x**m*atanh(tanh(b*x+a))**3, x)
```

output

```
Piecewise((b**3*log(x) - b**2*atanh(tanh(a + b*x))/x - b*atanh(tanh(a + b*x))**2/(2*x**2) - atanh(tanh(a + b*x))**3/(3*x**3), Eq(m, -4)), (Integral(atanh(tanh(a + b*x))**3/x**3, x), Eq(m, -3)), (Integral(atanh(tanh(a + b*x))**3/x**2, x), Eq(m, -2)), (Integral(atanh(tanh(a + b*x))**3/x, x), Eq(m, -1)), (-6*b**3*x**4*x**m/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + 6*b**2*m*x**3*x**m*atanh(tanh(a + b*x))/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + 24*b**2*x**3*x**m*atanh(tanh(a + b*x))/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) - 3*b**2*x**2*x**m*atanh(tanh(a + b*x))**2/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) - 21*b*m*x**2*x**m*atanh(tanh(a + b*x))**2/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) - 36*b*x**2*x**m*atanh(tanh(a + b*x))**2/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + m**3*x*x**m*atanh(tanh(a + b*x))**3/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + 9*m**2*x*x**m*atanh(tanh(a + b*x))**3/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + 26*m*x*x**m*atanh(tanh(a + b*x))**3/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24) + 24*x*x**m*atanh(tanh(a + b*x))**3/(m**4 + 10*m**3 + 35*m**2 + 50*m + 24), True))
```

### Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.99

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{3bx^2x^m \operatorname{arctanh}(\tanh(bx + a))^2}{(m+2)(m+1)} + \frac{x^{m+1} \operatorname{arctanh}(\tanh(bx + a))^3}{m+1} - \frac{6 \left( \frac{b^2x^4x^m}{(m+4)(m+3)(m+2)} - \frac{bx^3x^m \operatorname{arctanh}(\tanh(bx+a))}{(m+3)(m+2)} \right) b}{m+1}$$

input

```
integrate(x^m*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")
```

output

```
-3*b*x^2*x^m*arctanh(tanh(b*x + a))^2/((m + 2)*(m + 1)) + x^(m + 1)*arctanh(tanh(b*x + a))^3/(m + 1) - 6*(b^2*x^4*x^m/((m + 4)*(m + 3)*(m + 2)) - b*x^3*x^m*arctanh(tanh(b*x + a))/((m + 3)*(m + 2)))*b/(m + 1)
```

**Giac [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int x^m \operatorname{artanh}(\tanh(bx + a))^3 dx$$

input `integrate(x^m*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `integrate(x^m*arctanh(tanh(b*x + a))^3, x)`

**Mupad [B] (verification not implemented)**

Time = 3.42 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.02

$$\begin{aligned} & \int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx \\ &= \frac{8b^3 x^m x^4 (m^3 + 6m^2 + 11m + 6)}{8m^4 + 80m^3 + 280m^2 + 400m + 192} \\ & \quad - \frac{x x^m \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3 (m^3 + 9m^2 + 26m + 24)}{8m^4 + 80m^3 + 280m^2 + 400m + 192} \\ & \quad - \frac{12b^2 x^m x^3 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right) (m^3 + 7m^2 + 14m + 8)}{8m^4 + 80m^3 + 280m^2 + 400m + 192} \\ & \quad + \frac{6b x^m x^2 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2 (m^3 + 8m^2 + 19m + 12)}{8m^4 + 80m^3 + 280m^2 + 400m + 192} \end{aligned}$$

input `int(x^m*atanh(tanh(a + b*x))^3,x)`

output `(8*b^3*x^m*x^4*(11*m + 6*m^2 + m^3 + 6))/(400*m + 280*m^2 + 80*m^3 + 8*m^4 + 192) - (x*x^m*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3*(26*m + 9*m^2 + m^3 + 24))/(400*m + 280*m^2 + 80*m^3 + 8*m^4 + 192) - (12*b^2*x^m*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*(14*m + 7*m^2 + m^3 + 8))/(400*m + 280*m^2 + 80*m^3 + 8*m^4 + 192) + (6*b*x^m*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2*(19*m + 8*m^2 + m^3 + 12))/(400*m + 280*m^2 + 80*m^3 + 8*m^4 + 192)`

**Reduce [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int x^m \operatorname{atanh}(\tanh(bx + a))^3 dx$$

input `int(x^m*atanh(tanh(b*x+a))^3,x)`

output `int(x**m*atanh(tanh(a + b*x))**3,x)`

### 3.55 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	491
Mathematica [A] (verified)	491
Rubi [A] (verified)	492
Maple [A] (verified)	493
Fricas [A] (verification not implemented)	494
Sympy [A] (verification not implemented)	494
Maxima [A] (verification not implemented)	494
Giac [A] (verification not implemented)	495
Mupad [B] (verification not implemented)	495
Reduce [F]	496

#### Optimal result

Integrand size = 13, antiderivative size = 61

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{1}{140}b^3x^7 + \frac{1}{20}b^2x^6 \operatorname{arctanh}(\tanh(a + bx)) - \frac{3}{20}bx^5 \operatorname{arctanh}(\tanh(a + bx))^2 + \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^3$$

output

```
-1/140*b^3*x^7+1/20*b^2*x^6*arctanh(tanh(b*x+a))-3/20*b*x^5*arctanh(tanh(b*x+a))^2+1/4*x^4*arctanh(tanh(b*x+a))^3
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.89

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{1}{140}x^4(b^3x^3 - 7b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 21bx \operatorname{arctanh}(\tanh(a + bx))^2 - 35 \operatorname{arctanh}(\tanh(a + bx))^3)$$

input

```
Integrate[x^3*ArcTanh[Tanh[a + b*x]]^3,x]
```



output

```
-1/140*(x^4*(b^3*x^3 - 7*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 21*b*x*ArcTanh[Tanh[a + b*x]]^2 - 35*ArcTanh[Tanh[a + b*x]]^3))
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{4}b \int x^4 \operatorname{arctanh}(\tanh(a + bx))^2 dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{3}{4}b \left( \frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{5}b \int x^5 \operatorname{arctanh}(\tanh(a + bx)) dx \right) \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{3}{4}b \left( \frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{5}b \left( \frac{1}{6}x^6 \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^6 dx}{6} \right) \right) \\
 & \quad \downarrow \text{15} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{3}{4}b \left( \frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{5}b \left( \frac{1}{6}x^6 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^7}{42} \right) \right)
 \end{aligned}$$

input

```
Int[x^3*ArcTanh[Tanh[a + b*x]]^3,x]
```

output  $(x^4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3)/4 - (3*b*((x^5 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2)/5 - (2*b*(-1/42*(b*x^7) + (x^6 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]))/6))/5)/4$

### Defintions of rubi rules used

rule 15  $\operatorname{Int}[(a_.)*(x_)^(m_.), x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^(m + 1)/(m + 1)), x] \;/; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$

rule 2599  $\operatorname{Int}[(u_)^(m_.)*(v_)^(n_.), x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^(m + 1)*(v^n/(a*(m + 1))), x] - \operatorname{Simp}[b*(n/(a*(m + 1))) \operatorname{Int}[u^(m + 1)*v^(n - 1), x], x] \;/; \operatorname{NeQ}[b*u - a*v, 0] \;/; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m + n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n + m + 1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$

### Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.92

$$\frac{x^4 \operatorname{arctanh}(\operatorname{tanh}(bx + a))^3}{4} - \frac{3b \left( \frac{x^5 \operatorname{arctanh}(\operatorname{tanh}(bx + a))^2}{5} - \frac{2b \left( \frac{x^6 \operatorname{arctanh}(\operatorname{tanh}(bx + a))}{6} - \frac{bx^7}{42} \right)}{5} \right)}{4}$$

input  $\operatorname{int}(x^3 \operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3, x)$

output  $1/4*x^4*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3-3/4*b*(1/5*x^5*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^2-2/5*b*(1/6*x^6*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))-1/42*b*x^7))$

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.57

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{7} b^3 x^7 + \frac{1}{2} a b^2 x^6 + \frac{3}{5} a^2 b x^5 + \frac{1}{4} a^3 x^4$$

input `integrate(x^3*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `1/7*b^3*x^7 + 1/2*a*b^2*x^6 + 3/5*a^2*b*x^5 + 1/4*a^3*x^4`**Sympy [A] (verification not implemented)**

Time = 0.50 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.31

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$= \begin{cases} \frac{x^3 \operatorname{atanh}^4(\tanh(a + bx))}{4b} - \frac{3x^2 \operatorname{atanh}^5(\tanh(a + bx))}{20b^2} + \frac{x \operatorname{atanh}^6(\tanh(a + bx))}{20b^3} - \frac{\operatorname{atanh}^7(\tanh(a + bx))}{140b^4} & \text{for } b \neq 0 \\ \frac{x^4 \operatorname{atanh}^3(\tanh(a))}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*atanh(tanh(b*x+a))**3,x)`output `Piecewise((x**3*atanh(tanh(a + b*x))**4/(4*b) - 3*x**2*atanh(tanh(a + b*x))**5/(20*b**2) + x*atanh(tanh(a + b*x))**6/(20*b**3) - atanh(tanh(a + b*x))**7/(140*b**4), Ne(b, 0)), (x**4*atanh(tanh(a))**3/4, True))`**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.89

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{3}{20} b x^5 \operatorname{artanh}(\tanh(bx + a))^2$$

$$+ \frac{1}{4} x^4 \operatorname{artanh}(\tanh(bx + a))^3$$

$$- \frac{1}{140} (b^2 x^7 - 7 b x^6 \operatorname{artanh}(\tanh(bx + a))) b$$

input `integrate(x^3*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output 
$$-3/20*b*x^5*arctanh(tanh(b*x + a))^2 + 1/4*x^4*arctanh(tanh(b*x + a))^3 - 1/140*(b^2*x^7 - 7*b*x^6*arctanh(tanh(b*x + a)))*b$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.57

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{7} b^3 x^7 + \frac{1}{2} a b^2 x^6 + \frac{3}{5} a^2 b x^5 + \frac{1}{4} a^3 x^4$$

input `integrate(x^3*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output 
$$1/7*b^3*x^7 + 1/2*a*b^2*x^6 + 3/5*a^2*b*x^5 + 1/4*a^3*x^4$$

### Mupad [B] (verification not implemented)

Time = 3.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{b^3 x^7}{140} + \frac{b^2 x^6 \operatorname{atanh}(\tanh(a + bx))}{20} - \frac{3 b x^5 \operatorname{atanh}(\tanh(a + bx))^2}{20} + \frac{x^4 \operatorname{atanh}(\tanh(a + bx))^3}{4}$$

input `int(x^3*atanh(tanh(a + b*x))^3,x)`

output 
$$(x^4*\operatorname{atanh}(\tanh(a + b*x))^3)/4 - (b^3*x^7)/140 - (3*b*x^5*\operatorname{atanh}(\tanh(a + b*x))^2)/20 + (b^2*x^6*\operatorname{atanh}(\tanh(a + b*x)))/20$$

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \operatorname{atanh}(\tanh(bx + a))^3 x^3 dx$$

input `int(x^3*atanh(tanh(b*x+a))^3,x)`

output `int(atanh(tanh(a + b*x))**3*x**3,x)`

### 3.56 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	497
Mathematica [A] (verified)	497
Rubi [A] (verified)	498
Maple [A] (verified)	499
Fricas [A] (verification not implemented)	500
Sympy [A] (verification not implemented)	500
Maxima [A] (verification not implemented)	500
Giac [A] (verification not implemented)	501
Mupad [B] (verification not implemented)	501
Reduce [F]	502

#### Optimal result

Integrand size = 13, antiderivative size = 53

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{10b^2} + \frac{\operatorname{arctanh}(\tanh(a + bx))^6}{60b^3}$$

output

$1/4*x^2*\operatorname{arctanh}(\tanh(b*x+a))^4/b-1/10*x*\operatorname{arctanh}(\tanh(b*x+a))^5/b^2+1/60*\operatorname{arctanh}(\tanh(b*x+a))^6/b^3$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{1}{60}x^3(b^3x^3 - 6b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 15bx \operatorname{arctanh}(\tanh(a + bx))^2 - 20 \operatorname{arctanh}(\tanh(a + bx))^3)$$

input

`Integrate[x^2*ArcTanh[Tanh[a + b*x]]^3,x]`

output

```
-1/60*(x^3*(b^3*x^3 - 6*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 15*b*x*ArcTanh[Ta
nh[a + b*x]]^2 - 20*ArcTanh[Tanh[a + b*x]]^3))
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2599$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx}{2b}$$

$$\downarrow 2599$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^5 dx}{2b}$$

$$\downarrow 2588$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^5 d \operatorname{arctanh}(\tanh(a + bx))}{2b}$$

$$\downarrow 15$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^6}{30b^2}$$

input

```
Int[x^2*ArcTanh[Tanh[a + b*x]]^3,x]
```

output

```
(x^2*ArcTanh[Tanh[a + b*x]]^4)/(4*b) - ((x*ArcTanh[Tanh[a + b*x]]^5)/(5*b)
- ArcTanh[Tanh[a + b*x]]^6/(30*b^2))/(2*b)
```

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :=> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :=> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :=> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.06

$$\frac{x^3 \operatorname{arctanh}(\tanh(bx + a))^3}{3} - b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx + a))^2}{4} - \frac{b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx + a))}{5} - \frac{bx^6}{30} \right)}{2} \right)$$

input `int(x^2*arctanh(tanh(b*x+a))^3,x)`

output `1/3*x^3*arctanh(tanh(b*x+a))^3-b*(1/4*x^4*arctanh(tanh(b*x+a))^2-1/2*b*(1/5*x^5*arctanh(tanh(b*x+a))-1/30*b*x^6))`



**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.66

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{6} b^3 x^6 + \frac{3}{5} a b^2 x^5 + \frac{3}{4} a^2 b x^4 + \frac{1}{3} a^3 x^3$$

input `integrate(x^2*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `1/6*b^3*x^6 + 3/5*a*b^2*x^5 + 3/4*a^2*b*x^4 + 1/3*a^3*x^3`

**Sympy [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.13

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \begin{cases} \frac{x^2 \operatorname{atanh}^4(\tanh(a+bx))}{4b} - \frac{x \operatorname{atanh}^5(\tanh(a+bx))}{10b^2} + \frac{\operatorname{atanh}^6(\tanh(a+bx))}{60b^3} & \text{for } b \neq 0 \\ \frac{x^3 \operatorname{atanh}^3(\tanh(a))}{3} & \text{otherwise} \end{cases}$$

input `integrate(x**2*atanh(tanh(b*x+a))**3,x)`

output `Piecewise((x**2*atanh(tanh(a + b*x))**4/(4*b) - x*atanh(tanh(a + b*x))**5/(10*b**2) + atanh(tanh(a + b*x))**6/(60*b**3), Ne(b, 0)), (x**3*atanh(tanh(a))**3/3, True))`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{1}{4} b x^4 \operatorname{artanh}(\tanh(bx + a))^2 + \frac{1}{3} x^3 \operatorname{artanh}(\tanh(bx + a))^3 - \frac{1}{60} (b^2 x^6 - 6 b x^5 \operatorname{artanh}(\tanh(bx + a))) b$$

input `integrate(x^2*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output 
$$-1/4*b*x^4*arctanh(tanh(b*x + a))^2 + 1/3*x^3*arctanh(tanh(b*x + a))^3 - 1/60*(b^2*x^6 - 6*b*x^5*arctanh(tanh(b*x + a)))*b$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.66

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{6} b^3 x^6 + \frac{3}{5} a b^2 x^5 + \frac{3}{4} a^2 b x^4 + \frac{1}{3} a^3 x^3$$

input `integrate(x^2*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output 
$$1/6*b^3*x^6 + 3/5*a*b^2*x^5 + 3/4*a^2*b*x^4 + 1/3*a^3*x^3$$

### Mupad [B] (verification not implemented)

Time = 3.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{b^3 x^6}{60} + \frac{b^2 x^5 \operatorname{atanh}(\tanh(a + bx))}{10} - \frac{b x^4 \operatorname{atanh}(\tanh(a + bx))^2}{4} + \frac{x^3 \operatorname{atanh}(\tanh(a + bx))^3}{3}$$

input `int(x^2*atanh(tanh(a + b*x))^3,x)`

output 
$$(x^3*\operatorname{atanh}(\tanh(a + b*x))^3)/3 - (b^3*x^6)/60 - (b*x^4*\operatorname{atanh}(\tanh(a + b*x))^2)/4 + (b^2*x^5*\operatorname{atanh}(\tanh(a + b*x)))/10$$

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \operatorname{atanh}(\tanh(bx + a))^3 x^2 dx$$

input `int(x^2*atanh(tanh(b*x+a))^3,x)`

output `int(atanh(tanh(a + b*x))**3*x**2,x)`

### 3.57 $\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	503
Mathematica [B] (verified)	503
Rubi [A] (verified)	504
Maple [A] (verified)	505
Fricas [A] (verification not implemented)	506
Sympy [A] (verification not implemented)	506
Maxima [A] (verification not implemented)	506
Giac [A] (verification not implemented)	507
Mupad [B] (verification not implemented)	507
Reduce [F]	508

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{x \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{20b^2}$$

output

$$1/4*x*\operatorname{arctanh}(\tanh(b*x+a))^4/b-1/20*\operatorname{arctanh}(\tanh(b*x+a))^5/b^2$$

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 99 vs. 2(34) = 68.

Time = 0.21 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.91

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{(a + bx) ((4a - bx)(a + bx)^3 - 5(3a - bx)(a + bx)^2 \operatorname{arctanh}(\tanh(a + bx)) + 10(2a^2 + abx - b^2x^2) \operatorname{arctanh}(\tanh(a + bx)))}{20b^2}$$

input

$$\operatorname{Integrate}[x*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3,x]$$

output

```
((a + b*x)*((4*a - b*x)*(a + b*x)^3 - 5*(3*a - b*x)*(a + b*x)^2*ArcTanh[Tanh[a + b*x]] + 10*(2*a^2 + a*b*x - b^2*x^2)*ArcTanh[Tanh[a + b*x]]^2 - 10*(a - b*x)*ArcTanh[Tanh[a + b*x]]^3))/(20*b^2)
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2599$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^4 dx}{4b}$$

$$\downarrow 2588$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^4 d \operatorname{arctanh}(\tanh(a + bx))}{4b^2}$$

$$\downarrow 15$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^4}{4b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{20b^2}$$

input

```
Int[x*ArcTanh[Tanh[a + b*x]]^3,x]
```

output

```
(x*ArcTanh[Tanh[a + b*x]]^4)/(4*b) - ArcTanh[Tanh[a + b*x]]^5/(20*b^2)
```

## Definitions of rubi rules used

rule 15  $\text{Int}[(a_.)(x_)^(m_.), x\_Symbol] \rightarrow \text{Simp}[a*(x^(m + 1)/(m + 1)), x] \text{ /; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 2588  $\text{Int}[(u_)^(m_.), x\_Symbol] \rightarrow \text{With}[\{c = \text{Simplify}[D[u, x]]\}, \text{Simp}[1/c \ \text{Subst}[\text{Int}[x^m, x], x, u], x]] \text{ /; FreeQ}[m, x] \ \&\& \ \text{PiecewiseLinearQ}[u, x]$

rule 2599  $\text{Int}[(u_)^(m_.)(v_)^(n_.), x\_Symbol] \rightarrow \text{With}[\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[u^(m + 1)*(v^n/(a*(m + 1))), x] - \text{Simp}[b*(n/(a*(m + 1))) \ \text{Int}[u^(m + 1)*v^(n - 1), x], x] \text{ /; NeQ}[b*u - a*v, 0] \text{ /; FreeQ}[\{m, n\}, x] \ \&\& \ \text{PiecewiseLinearQ}[u, v, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ ((\text{LtQ}[m, -1] \ \&\& \ \text{GtQ}[n, 0]) \ \&\& \ !(\text{ILtQ}[m + n, -2] \ \&\& \ (\text{FractionQ}[m] \ || \ \text{GeQ}[2*n + m + 1, 0]))) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LeQ}[n, m]) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ !\text{IntegerQ}[m]) \ || \ (\text{ILtQ}[m, 0] \ \&\& \ !\text{IntegerQ}[n]))$

## Maple [A] (verified)

Time = 43.93 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.59

method	result	size
parallelrisch	$\frac{b^2 \operatorname{arctanh}(\tanh(bx+a))x^4}{4} - \frac{b \operatorname{arctanh}(\tanh(bx+a))^2 x^3}{2} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^3}{2} - \frac{b^3 x^5}{20}$	54
default	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^3}{2} - \frac{3b \left( \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{2b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))}{4} - \frac{bx^5}{20} \right)}{3} \right)}{2}$	56
parts	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^3}{2} - \frac{3b \left( \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{2b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))}{4} - \frac{bx^5}{20} \right)}{3} \right)}{2}$	56
risch	Expression too large to display	8165

input  $\text{int}(x*\operatorname{arctanh}(\tanh(b*x+a))^3, x, \text{method}=\_RETURNVERBOSE)$

output  $1/4*b^2*\operatorname{arctanh}(\tanh(b*x+a))*x^4-1/2*b*\operatorname{arctanh}(\tanh(b*x+a))^2*x^3+1/2*x^2*\operatorname{arctanh}(\tanh(b*x+a))^3-1/20*b^3*x^5$

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{5} b^3 x^5 + \frac{3}{4} a b^2 x^4 + a^2 b x^3 + \frac{1}{2} a^3 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `1/5*b^3*x^5 + 3/4*a*b^2*x^4 + a^2*b*x^3 + 1/2*a^3*x^2`**Sympy [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = \begin{cases} \frac{x \operatorname{atanh}^4(\tanh(a+bx))}{4b} - \frac{\operatorname{atanh}^5(\tanh(a+bx))}{20b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^3(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**3,x)`output `Piecewise((x*atanh(tanh(a + b*x))**4/(4*b) - atanh(tanh(a + b*x))**5/(20*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**3/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.59

$$\begin{aligned} \int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = & -\frac{1}{2} b x^3 \operatorname{artanh}(\tanh(bx + a))^2 \\ & + \frac{1}{2} x^2 \operatorname{artanh}(\tanh(bx + a))^3 \\ & - \frac{1}{20} (b^2 x^5 - 5 b x^4 \operatorname{artanh}(\tanh(bx + a))) b \end{aligned}$$

input `integrate(x*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output 
$$-1/2*b*x^3*\operatorname{arctanh}(\operatorname{tanh}(b*x + a))^2 + 1/2*x^2*\operatorname{arctanh}(\operatorname{tanh}(b*x + a))^3 - 1/20*(b^2*x^5 - 5*b*x^4*\operatorname{arctanh}(\operatorname{tanh}(b*x + a)))*b$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int x \operatorname{arctanh}(\operatorname{tanh}(a + bx))^3 dx = \frac{1}{5} b^3 x^5 + \frac{3}{4} a b^2 x^4 + a^2 b x^3 + \frac{1}{2} a^3 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output 
$$1/5*b^3*x^5 + 3/4*a*b^2*x^4 + a^2*b*x^3 + 1/2*a^3*x^2$$

### Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.56

$$\int x \operatorname{arctanh}(\operatorname{tanh}(a + bx))^3 dx = -\frac{b^3 x^5}{20} + \frac{b^2 x^4 \operatorname{atanh}(\operatorname{tanh}(a + bx))}{4} - \frac{b x^3 \operatorname{atanh}(\operatorname{tanh}(a + bx))^2}{2} + \frac{x^2 \operatorname{atanh}(\operatorname{tanh}(a + bx))^3}{2}$$

input `int(x*atanh(tanh(a + b*x))^3,x)`

output 
$$(x^2*\operatorname{atanh}(\operatorname{tanh}(a + b*x))^3)/2 - (b^3*x^5)/20 - (b*x^3*\operatorname{atanh}(\operatorname{tanh}(a + b*x))^2)/2 + (b^2*x^4*\operatorname{atanh}(\operatorname{tanh}(a + b*x)))/4$$



**Reduce [F]**

$$\int x \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int x \operatorname{atanh}(\tanh(bx + a))^3 dx$$

input `int(x*atanh(tanh(b*x+a))^3,x)`

output `int(atanh(tanh(a + b*x))**3*x,x)`

### 3.58 $\int \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	509
Mathematica [A] (verified)	509
Rubi [A] (verified)	510
Maple [A] (verified)	511
Fricas [B] (verification not implemented)	511
Sympy [A] (verification not implemented)	512
Maxima [B] (verification not implemented)	512
Giac [B] (verification not implemented)	513
Mupad [B] (verification not implemented)	513
Reduce [B] (verification not implemented)	513

#### Optimal result

Integrand size = 9, antiderivative size = 16

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4b}$$

output `1/4*arctanh(tanh(b*x+a))^4/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4b}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^3,x]`

output `ArcTanh[Tanh[a + b*x]]^4/(4*b)`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2588$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^3 d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow 15$$

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^3,x]`

output `ArcTanh[Tanh[a + b*x]]^4/(4*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 44.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{4b}$
default	$\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{4b}$
parallelrisc	$-\frac{b^3x^4}{4} + b^2 \operatorname{arctanh}(\tanh(bx+a))x^3 - \frac{3b \operatorname{arctanh}(\tanh(bx+a))^2x^2}{2} + x \operatorname{arctanh}(\tanh(bx+a))$
risc	Expression too large to display

input `int(arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `1/4*arctanh(tanh(b*x+a))^4/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{4}b^3x^4 + ab^2x^3 + \frac{3}{2}a^2bx^2 + a^3x$$

input `integrate(arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `1/4*b^3*x^4 + a*b^2*x^3 + 3/2*a^2*b*x^2 + a^3*x`

**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \begin{cases} \frac{\operatorname{atanh}^4(\tanh(a+bx))}{4b} & \text{for } b \neq 0 \\ x \operatorname{atanh}^3(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**3,x)`

output `Piecewise((atanh(tanh(a + b*x))**4/(4*b), Ne(b, 0)), (x*atanh(tanh(a))**3, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(14) = 28.

Time = 0.16 (sec) , antiderivative size = 51, normalized size of antiderivative = 3.19

$$\begin{aligned} \int \operatorname{arctanh}(\tanh(a + bx))^3 dx &= -\frac{3}{2} bx^2 \operatorname{arctanh}(\tanh(bx + a))^2 \\ &\quad + x \operatorname{arctanh}(\tanh(bx + a))^3 \\ &\quad - \frac{1}{4} (b^2 x^4 - 4 bx^3 \operatorname{arctanh}(\tanh(bx + a)))b \end{aligned}$$

input `integrate(arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `-3/2*b*x^2*arctanh(tanh(b*x + a))^2 + x*arctanh(tanh(b*x + a))^3 - 1/4*(b^2*x^4 - 4*b*x^3*arctanh(tanh(b*x + a)))*b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs.  $2(14) = 28$ .

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{1}{2} (bx^2 + 2ax)a^2 + \frac{1}{4} (bx^2 + 2ax)^2 b$$

input `integrate(arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `1/2*(b*x^2 + 2*a*x)*a^2 + 1/4*(b*x^2 + 2*a*x)^2*b`

**Mupad [B] (verification not implemented)**

Time = 3.20 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.94

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{x(2 \operatorname{atanh}(\tanh(a + bx)) - bx)(b^2 x^2 - 2bx \operatorname{atanh}(\tanh(a + bx)) + 2 \operatorname{atanh}(\tanh(a + bx))^2)}{4}$$

input `int(atanh(tanh(a + b*x))^3,x)`

output `(x*(2*atanh(tanh(a + b*x)) - b*x)*(2*atanh(tanh(a + b*x))^2 + b^2*x^2 - 2*b*x*atanh(tanh(a + b*x))))/4`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{\operatorname{atanh}(\tanh(bx + a))^4}{4b}$$

input `int(atanh(tanh(b*x+a))^3,x)`

output `atanh(tanh(a + b*x))**4/(4*b)`

### 3.59 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x} dx$

Optimal result . . . . .	515
Mathematica [A] (verified) . . . . .	515
Rubi [A] (verified) . . . . .	516
Maple [A] (verified) . . . . .	517
Fricas [A] (verification not implemented) . . . . .	518
Sympy [F] . . . . .	518
Maxima [A] (verification not implemented) . . . . .	519
Giac [A] (verification not implemented) . . . . .	519
Mupad [B] (verification not implemented) . . . . .	520
Reduce [F] . . . . .	521

#### Optimal result

Integrand size = 13, antiderivative size = 77

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = bx(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))^2 + \frac{1}{3}\operatorname{arctanh}(\tanh(a + bx))^3 - (bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \log(x)$$

output

```
b*x*(b*x-arctanh(tanh(b*x+a)))^2-1/2*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^2+1/3*arctanh(tanh(b*x+a))^3-(b*x-arctanh(tanh(b*x+a)))^3*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.35

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \frac{1}{3}(a + bx)^3 + (a + bx)(a^2 - 3a(a + bx - \operatorname{arctanh}(\tanh(a + bx))) + 3(a + bx - \operatorname{arctanh}(\tanh(a + bx)))^2) - \frac{1}{2}(a + bx)^2(2a + 3bx - 3\operatorname{arctanh}(\tanh(a + bx))) + (-bx + \operatorname{arctanh}(\tanh(a + bx)))^3 \log(bx)$$



input `Integrate[ArcTanh[Tanh[a + b*x]]^3/x, x]`

output  $(a + b*x)^3/3 + (a + b*x)*(a^2 - 3*a*(a + b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]) + 3*(a + b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])^2) - ((a + b*x)^2*(2*a + 3*b*x - 3*\text{ArcTanh}[\text{Tanh}[a + b*x]]))/2 + (-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]])^3*\text{Log}[b*x]$

## Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2590, 2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\text{arctanh}(\tanh(a + bx))^3}{x} dx$$

$$\downarrow 2590$$

$$\frac{1}{3}\text{arctanh}(\tanh(a + bx))^3 - (bx - \text{arctanh}(\tanh(a + bx))) \int \frac{\text{arctanh}(\tanh(a + bx))^2}{x} dx$$

$$\downarrow 2590$$

$$\frac{1}{3}\text{arctanh}(\tanh(a + bx))^3 - (bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{1}{2}\text{arctanh}(\tanh(a + bx))^2 - (bx - \text{arctanh}(\tanh(a + bx))) \int \frac{\text{arctanh}(\tanh(a + bx))}{x} dx \right)$$

$$\downarrow 2589$$

$$\frac{1}{3}\text{arctanh}(\tanh(a + bx))^3 - (bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{1}{2}\text{arctanh}(\tanh(a + bx))^2 - (bx - \text{arctanh}(\tanh(a + bx))) \left( bx - (bx - \text{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) \right)$$

$$\downarrow 14$$

$$\frac{1}{3}\text{arctanh}(\tanh(a + bx))^3 - (bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{1}{2}\text{arctanh}(\tanh(a + bx))^2 - (bx - \text{arctanh}(\tanh(a + bx)))(bx - \log(x)(bx - \text{arctanh}(\tanh(a + bx)))) \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^3/x,x]`

output `ArcTanh[Tanh[a + b*x]]^3/3 - (b*x - ArcTanh[Tanh[a + b*x]])*(ArcTanh[Tanh[a + b*x]]^2/2 - (b*x - ArcTanh[Tanh[a + b*x]])*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x]))`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

### Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.91

method	result
default	$\ln(x) \operatorname{arctanh}(\tanh(bx + a))^3 - 3b \left( b^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 2ab \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + 2b \operatorname{arctanh}(\tanh(bx + a)) \right)$
parts	$\ln(x) \operatorname{arctanh}(\tanh(bx + a))^3 - 3b \left( b^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 2ab \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + 2b \operatorname{arctanh}(\tanh(bx + a)) \right)$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^3/x,x,method=_RETURNVERBOSE)`

output

```
ln(x)*arctanh(tanh(b*x+a))^3-3*b*(b^2*(1/3*x^3*ln(x)-1/9*x^3)+2*a*b*(1/2*x^2*ln(x)-1/4*x^2)+2*b*(arctanh(tanh(b*x+a))-b*x-a)*(1/2*x^2*ln(x)-1/4*x^2)+a^2*(x*ln(x)-x)+2*a*(arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x)+(arctanh(tanh(b*x+a))-b*x-a)^2*(x*ln(x)-x))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.40

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \frac{1}{3} b^3 x^3 + \frac{3}{2} ab^2 x^2 + 3a^2 bx + a^3 \log(x)$$

input

```
integrate(arctanh(tanh(b*x+a))^3/x,x, algorithm="fricas")
```

output

```
1/3*b^3*x^3 + 3/2*a*b^2*x^2 + 3*a^2*b*x + a^3*log(x)
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \int \frac{\operatorname{atanh}^3(\tanh(a + bx))}{x} dx$$

input

```
integrate(atanh(tanh(b*x+a))**3/x,x)
```

output

```
Integral(atanh(tanh(a + b*x))**3/x, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.40

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \frac{1}{3} b^3 x^3 + \frac{3}{2} ab^2 x^2 + 3 a^2 bx + a^3 \log(x)$$

input `integrate(arctanh(tanh(b*x+a))^3/x,x, algorithm="maxima")`

output `1/3*b^3*x^3 + 3/2*a*b^2*x^2 + 3*a^2*b*x + a^3*log(x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.42

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \frac{1}{3} b^3 x^3 + \frac{3}{2} ab^2 x^2 + 3 a^2 bx + a^3 \log(|x|)$$

input `integrate(arctanh(tanh(b*x+a))^3/x,x, algorithm="giac")`

output `1/3*b^3*x^3 + 3/2*a*b^2*x^2 + 3*a^2*b*x + a^3*log(abs(x))`

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 306, normalized size of antiderivative = 3.97

$$\begin{aligned}
& \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx \\
&= \frac{b^3 x^3}{3} - \ln(x) \left( \frac{\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^3}{8} - a^3 \right. \\
&\quad \left. - \frac{3a\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{4} \right. \\
&\quad \left. + \frac{3a^2\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{2} \right) \\
&\quad - \frac{3b^2x^2\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{4} \\
&\quad + \frac{3bx\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{4}
\end{aligned}$$

input `int(atanh(tanh(a + b*x))^3/x,x)`

output

```

(b^3*x^3)/3 - log(x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3/8 - a^3 - (3*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/4 + (3*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/2) - (3*b^2*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/4 + (3*b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/4

```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{x} dx$$

input `int(atanh(tanh(b*x+a))^3/x,x)`

output `int(atanh(tanh(a + b*x))**3/x,x)`

### 3.60 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^2} dx$

Optimal result . . . . .	522
Mathematica [A] (verified) . . . . .	522
Rubi [A] (verified) . . . . .	523
Maple [A] (verified) . . . . .	525
Fricas [A] (verification not implemented) . . . . .	525
Sympy [F] . . . . .	525
Maxima [A] (verification not implemented) . . . . .	526
Giac [A] (verification not implemented) . . . . .	526
Mupad [B] (verification not implemented) . . . . .	527
Reduce [F] . . . . .	528

#### Optimal result

Integrand size = 13, antiderivative size = 68

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx = -3b^2x(bx - \operatorname{arctanh}(\tanh(a + bx))) + \frac{3}{2}b\operatorname{arctanh}(\tanh(a + bx))^2 - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} + 3b(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \log(x)$$

output

```
-3*b^2*x*(b*x-arctanh(tanh(b*x+a)))+3/2*b*arctanh(tanh(b*x+a))^2-arctanh(tanh(b*x+a))^3/x+3*b*(b*x-arctanh(tanh(b*x+a)))^2*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.91

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx = -\frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} - 6b^2x\operatorname{arctanh}(\tanh(a + bx)) \log(x) + 3b\operatorname{arctanh}(\tanh(a + bx))^2(1 + \log(x)) + \frac{3}{2}b^3x^2(-1 + 2\log(x))$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^3/x^2, x]`

output `-(ArcTanh[Tanh[a + b*x]]^3/x) - 6*b^2*x*ArcTanh[Tanh[a + b*x]]*Log[x] + 3*b*ArcTanh[Tanh[a + b*x]]^2*(1 + Log[x]) + (3*b^3*x^2*(-1 + 2*Log[x]))/2`

### Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx$$

$$\downarrow \text{2599}$$

$$3b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x}$$

$$\downarrow \text{2590}$$

$$3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x}$$

$$\downarrow \text{2589}$$

$$3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x}$$

$$\downarrow \text{14}$$



$$3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx)))(bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx)))) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x}$$

input `Int[ArcTanh[Tanh[a + b*x]]^3/x^2,x]`

output `-(ArcTanh[Tanh[a + b*x]]^3/x) + 3*b*(ArcTanh[Tanh[a + b*x]]^2/2 - (b*x - ArcTanh[Tanh[a + b*x]])*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x]))`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.25

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{x} + 3b \left( \ln(x) \operatorname{arctanh}(\tanh(bx+a))^2 - 2b \left( b \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + a(x \ln(x) - x) \right) \right)$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{x} + 3b \left( \ln(x) \operatorname{arctanh}(\tanh(bx+a))^2 - 2b \left( b \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{4} \right) + a(x \ln(x) - x) \right) \right)$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^3/x^2,x,method=_RETURNVERBOSE)`

output `-arctanh(tanh(b*x+a))^3/x+3*b*(ln(x)*arctanh(tanh(b*x+a))^2-2*b*(b*(1/2*x^2*ln(x)-1/4*x^2)+a*(x*ln(x)-x)+(arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x)))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.53

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^2} dx = \frac{b^3 x^3 + 6ab^2 x^2 + 6a^2 bx \log(x) - 2a^3}{2x}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^2,x, algorithm="fricas")`

output `1/2*(b^3*x^3 + 6*a*b^2*x^2 + 6*a^2*b*x*log(x) - 2*a^3)/x`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^2} dx = \int \frac{\operatorname{atanh}^3(\tanh(a+bx))}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**3/x**2,x)`

output `Integral(atanh(tanh(a + b*x))**3/x**2, x)`

### Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx$$

$$= 3b \operatorname{arctanh}(\tanh(bx + a))^2 \log(x)$$

$$+ \frac{3}{2} (b^2 x^2 + 4abx + 2a^2 \log(x) - 2 \operatorname{arctanh}(\tanh(bx + a))^2 \log(x)) b$$

$$- \frac{\operatorname{arctanh}(\tanh(bx + a))^3}{x}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^2,x, algorithm="maxima")`

output `3*b*arctanh(tanh(b*x + a))^2*log(x) + 3/2*(b^2*x^2 + 4*a*b*x + 2*a^2*log(x)) - 2*arctanh(tanh(b*x + a))^2*log(x))*b - arctanh(tanh(b*x + a))^3/x`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx = \frac{1}{2} b^3 x^2 + 3ab^2 x + 3a^2 b \log(|x|) - \frac{a^3}{x}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^2,x, algorithm="giac")`

output `1/2*b^3*x^2 + 3*a*b^2*x + 3*a^2*b*log(abs(x)) - a^3/x`

**Mupad [B] (verification not implemented)**

Time = 3.40 (sec) , antiderivative size = 415, normalized size of antiderivative = 6.10

$$\begin{aligned}
\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx = & \frac{3b \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^2}{4} - \frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^3}{8x} \\
& + \frac{3b \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^2}{4} + \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^3}{8x} \\
& - \frac{3b^3 x^2}{2} + \frac{3b \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^2 \ln(x)}{4} \\
& + \frac{3 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^2}{8x} \\
& - \frac{3 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^2 \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{8x} \\
& + \frac{3b \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^2 \ln(x)}{4} \\
& - \frac{3b \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} + 3b^3 x^2 \ln(x) \\
& - \frac{3b \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \ln(x)}{2} \\
& + 3b^2 x \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln(x) \\
& - 3b^2 x \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \ln(x)
\end{aligned}$$

input `int(atanh(tanh(a + b*x))^3/x^2,x)`

output

```
(3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))^2)/4 - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3/(8*x) + (3*b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/4 + log(1/(exp(2*a)*exp(2*b*x) + 1))^3/(8*x) - (3*b^3*x^2)/2 + (3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log(x))/4 + (3*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/(8*x) - (3*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(8*x) + (3*b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2*log(x))/4 - (3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + 3*b^3*x^2*log(x) - (3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x))/2 + 3*b^2*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*log(x) - 3*b^2*x*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{x^2} dx$$

input

```
int(atanh(tanh(b*x+a))^3/x^2,x)
```

output

```
int(atanh(tanh(a + b*x))**3/x**2,x)
```

### 3.61 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^3} dx$

Optimal result	529
Mathematica [A] (verified)	529
Rubi [A] (verified)	530
Maple [A] (verified)	531
Fricas [A] (verification not implemented)	532
Sympy [F]	532
Maxima [A] (verification not implemented)	532
Giac [A] (verification not implemented)	533
Mupad [B] (verification not implemented)	534
Reduce [F]	535

#### Optimal result

Integrand size = 13, antiderivative size = 60

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = 3b^3x - \frac{3b\operatorname{arctanh}(\tanh(a + bx))^2}{2x} - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{2x^2} - 3b^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \log(x)$$

output

$3*b^3*x-3/2*b*\operatorname{arctanh}(\tanh(b*x+a))^2/x-1/2*\operatorname{arctanh}(\tanh(b*x+a))^3/x^2-3*b^2*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))*\ln(x)$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = b^3x - \frac{3b(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}{x} - \frac{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3}{2x^2} + 3b^2(-bx + \operatorname{arctanh}(\tanh(a + bx))) \log(x)$$

input

`Integrate[ArcTanh[Tanh[a + b*x]]^3/x^3,x]`

output

$$b^3 x - (3b(-bx) + \text{ArcTanh}[\text{Tanh}[a + bx]])^2/x - (-bx) + \text{ArcTanh}[\text{Tanh}[a + bx]]^3/(2x^2) + 3b^2(-bx) + \text{ArcTanh}[\text{Tanh}[a + bx]] * \text{Log}[x]$$
**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\text{arctanh}(\tanh(a + bx))^3}{x^3} dx$$

$$\downarrow 2599$$

$$\frac{3}{2}b \int \frac{\text{arctanh}(\tanh(a + bx))^2}{x^2} dx - \frac{\text{arctanh}(\tanh(a + bx))^3}{2x^2}$$

$$\downarrow 2599$$

$$\frac{3}{2}b \left( 2b \int \frac{\text{arctanh}(\tanh(a + bx))}{x} dx - \frac{\text{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\text{arctanh}(\tanh(a + bx))^3}{2x^2}$$

$$\downarrow 2589$$

$$\frac{3}{2}b \left( 2b \left( bx - (bx - \text{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) - \frac{\text{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\text{arctanh}(\tanh(a + bx))^3}{2x^2}$$

$$\downarrow 14$$

$$\frac{3}{2}b \left( 2b(bx - \log(x)(bx - \text{arctanh}(\tanh(a + bx)))) - \frac{\text{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\text{arctanh}(\tanh(a + bx))^3}{2x^2}$$

input

$$\text{Int}[\text{ArcTanh}[\text{Tanh}[a + bx]]^3/x^3, x]$$

output

$$-1/2*\text{ArcTanh}[\text{Tanh}[a + b*x]]^3/x^2 + (3*b*(-(\text{ArcTanh}[\text{Tanh}[a + b*x]]^2/x) + 2*b*(b*x - (b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])*\text{Log}[x]))) / 2$$

### Defintions of rubi rules used

rule 14

$$\text{Int}[(a\_)/(x\_), x\_Symbol] \text{ :> } \text{Simp}[a*\text{Log}[x], x] \text{ /; } \text{FreeQ}[a, x]$$

rule 2589

$$\text{Int}[(v\_)/(u\_), x\_Symbol] \text{ :> } \text{With}[\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[b*(x/a), x] - \text{Simp}[(b*u - a*v)/a \text{ Int}[1/u, x], x] \text{ /; } \text{NeQ}[b*u - a*v, 0] \text{ /; } \text{PiecewiseLinearQ}[u, v, x]$$

rule 2599

$$\text{Int}[(u_)^{(m)}*(v_)^{(n)}, x\_Symbol] \text{ :> } \text{With}[\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[u^{(m+1)}*(v^{(n/(a*(m+1))}), x] - \text{Simp}[b*(n/(a*(m+1))) \text{ Int}[u^{(m+1)}*v^{(n-1)}, x], x] \text{ /; } \text{NeQ}[b*u - a*v, 0] \text{ /; } \text{FreeQ}[\{m, n\}, x] \ \&\& \ \text{PiecewiseLinearQ}[u, v, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ ((\text{LtQ}[m, -1] \ \&\& \ \text{GtQ}[n, 0]) \ \&\& \ !(\text{ILtQ}[m+n, -2] \ \&\& \ (\text{FractionQ}[m] \ || \ \text{GeQ}[2*n+m+1, 0]))) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LeQ}[n, m]) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ !\text{IntegerQ}[m]) \ || \ (\text{ILtQ}[m, 0] \ \&\& \ !\text{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.98

method	result	size
default	$-\frac{\text{arctanh}(\tanh(bx+a))^3}{2x^2} + \frac{3b\left(-\frac{\text{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x)\text{arctanh}(\tanh(bx+a)) - b(x\ln(x) - x))\right)}{2}$	59
parts	$-\frac{\text{arctanh}(\tanh(bx+a))^3}{2x^2} + \frac{3b\left(-\frac{\text{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x)\text{arctanh}(\tanh(bx+a)) - b(x\ln(x) - x))\right)}{2}$	59
risch	Expression too large to display	2008

input

$$\text{int}(\text{arctanh}(\tanh(b*x+a))^3/x^3, x, \text{method}=\_RETURNVERBOSE)$$



output

```
-1/2*arctanh(tanh(b*x+a))^3/x^2+3/2*b*(-arctanh(tanh(b*x+a))^2/x+2*b*(ln(x)
)*arctanh(tanh(b*x+a))-b*(x*ln(x)-x))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = \frac{2b^3x^3 + 6ab^2x^2 \log(x) - 6a^2bx - a^3}{2x^2}$$

input

```
integrate(arctanh(tanh(b*x+a))^3/x^3,x, algorithm="fricas")
```

output

```
1/2*(2*b^3*x^3 + 6*a*b^2*x^2*log(x) - 6*a^2*b*x - a^3)/x^2
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = \int \frac{\operatorname{atanh}^3(\tanh(a + bx))}{x^3} dx$$

input

```
integrate(atanh(tanh(b*x+a))**3/x**3,x)
```

output

```
Integral(atanh(tanh(a + b*x))**3/x**3, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.20

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx \\ &= 3 \left( b \operatorname{artanh}(\tanh(bx + a)) \log(x) - \left( b \left( x + \frac{a}{b} \right) \log(x) - b \left( x + \frac{a \log(x)}{b} \right) \right) b \right) b \\ & \quad - \frac{3b \operatorname{artanh}(\tanh(bx + a))^2}{2x} - \frac{\operatorname{artanh}(\tanh(bx + a))^3}{2x^2} \end{aligned}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^3,x, algorithm="maxima")`

output  $3*(b*\operatorname{arctanh}(\tanh(b*x + a))*\log(x) - (b*(x + a/b)*\log(x) - b*(x + a*\log(x)/b))*b - 3/2*b*\operatorname{arctanh}(\tanh(b*x + a))^2/x - 1/2*\operatorname{arctanh}(\tanh(b*x + a))^3/x^2$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = b^3 x + 3 ab^2 \log(|x|) - \frac{6 a^2 bx + a^3}{2 x^2}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^3,x, algorithm="giac")`

output  $b^3*x + 3*a*b^2*\log(\operatorname{abs}(x)) - 1/2*(6*a^2*b*x + a^3)/x^2$

**Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 365, normalized size of antiderivative = 6.08

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = \frac{9b^2 \ln\left(\frac{e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{4} - \frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)^3}{16x^2}$$

$$- \frac{9b^2 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{4} - \frac{3b^3 x}{2} + \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)^3}{16x^2}$$

$$- \frac{3b \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)^2}{8x} - \frac{3b^2 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) \ln(x)}{2}$$

$$+ \frac{3 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)^2}{16x^2}$$

$$- \frac{3 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)^2 \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{16x^2}$$

$$- \frac{3b \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)^2}{8x} + \frac{3b^2 \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) \ln(x)}{2}$$

$$- 3b^3 x \ln(x) + \frac{3b \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{4x}$$

input `int(atanh(tanh(a + b*x))^3/x^3,x)`output

```
(9*b^2*log(exp(2*b*x)/(exp(2*a)*exp(2*b*x) + 1)))/4 - log((exp(2*a)*exp(2*
b*x))/(exp(2*a)*exp(2*b*x) + 1))^3/(16*x^2) - (9*b^2*log(1/(exp(2*a)*exp(2
*b*x) + 1)))/4 - (3*b^3*x)/2 + log(1/(exp(2*a)*exp(2*b*x) + 1))^3/(16*x^2)
- (3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))^2)/(8*x) - (3*b^2*log(1/(exp(2*a)
*exp(2*b*x) + 1))*log(x))/2 + (3*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp
(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/(16*x^2) - (3*log(1/(exp(2
*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)
))/(16*x^2) - (3*b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)
/(8*x) + (3*b^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x)
)/2 - 3*b^3*x*log(x) + (3*b*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)
)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(4*x)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{x^3} dx$$

input `int(atanh(tanh(b*x+a))^3/x^3,x)`

output `int(atanh(tanh(a + b*x))**3/x**3,x)`

### 3.62 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^4} dx$

Optimal result . . . . .	536
Mathematica [A] (verified) . . . . .	536
Rubi [A] (verified) . . . . .	537
Maple [A] (verified) . . . . .	538
Fricas [A] (verification not implemented) . . . . .	539
Sympy [A] (verification not implemented) . . . . .	539
Maxima [A] (verification not implemented) . . . . .	539
Giac [A] (verification not implemented) . . . . .	540
Mupad [B] (verification not implemented) . . . . .	540
Reduce [B] (verification not implemented) . . . . .	541

#### Optimal result

Integrand size = 13, antiderivative size = 55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = -\frac{b^2 \operatorname{arctanh}(\tanh(a + bx))}{x} - \frac{b \operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3} + b^3 \log(x)$$

output `-b^2*arctanh(tanh(b*x+a))/x-1/2*b*arctanh(tanh(b*x+a))^2/x^2-1/3*arctanh(tanh(b*x+a))^3/x^3+b^3*ln(x)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.09

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = \frac{-6b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) - 3bx \operatorname{arctanh}(\tanh(a + bx))^2 - 2 \operatorname{arctanh}(\tanh(a + bx))^3 + b^3x^3(11 + 6 \ln(x))}{6x^3}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^3/x^4,x]`

output

```
(-6*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 3*b*x*ArcTanh[Tanh[a + b*x]]^2 - 2*ArcTanh[Tanh[a + b*x]]^3 + b^3*x^3*(11 + 6*Log[x]))/(6*x^3)
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2599, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx$$

$$\downarrow 2599$$

$$b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3}$$

$$\downarrow 2599$$

$$b \left( b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3}$$

$$\downarrow 2599$$

$$b \left( b \left( b \int \frac{1}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3}$$

$$\downarrow 14$$

$$b \left( b \left( b \log(x) - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^3/x^4,x]
```

output

$$-1/3 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3/x^3 + b*(-1/2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2/x^2 + b*(-\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]/x) + b*\operatorname{Log}[x]))$$

### Defintions of rubi rules used

rule 14

$$\operatorname{Int}[(a_.)/(x_), x\_Symbol] \rightarrow \operatorname{Simp}[a*\operatorname{Log}[x], x] \;/; \operatorname{FreeQ}[a, x]$$

rule 2599

$$\operatorname{Int}[(u_)^{(m_*)}*(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}*(v^n/(a*(m+1))), x] - \operatorname{Simp}[b*(n/(a*(m+1))) \operatorname{Int}[u^{(m+1)}*v^{(n-1)}, x], x] \;/; \operatorname{NeQ}[b*u - a*v, 0] \;/; \operatorname{FreeQ}[\{m, n\}, x] \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \&\& \operatorname{NeQ}[m, -1] \&\& ((\operatorname{LtQ}[m, -1] \&\& \operatorname{GtQ}[n, 0]) \&\& !(\operatorname{ILtQ}[m+n, -2] \&\& (\operatorname{FractionQ}[m] \|\operatorname{GeQ}[2*n+m+1, 0]))) \|\operatorname{IGtQ}[n, 0] \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{LeQ}[n, m]) \|\operatorname{IGtQ}[n, 0] \&\& !\operatorname{IntegerQ}[m]) \|\operatorname{ILtQ}[m, 0] \&\& !\operatorname{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{3x^3} + b\left(-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{2x^2} + b\left(-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))}{x} + b \ln(x)\right)\right)$	52
parts	$-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{3x^3} + b\left(-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{2x^2} + b\left(-\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))}{x} + b \ln(x)\right)\right)$	52
paralelrisch	$\frac{6b^3x^3 \ln(x) - 6b^2x^2 \operatorname{arctanh}(\operatorname{tanh}(bx+a)) - 3x \operatorname{arctanh}(\operatorname{tanh}(bx+a))^2 b - 2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{6x^3}$	56
risch	Expression too large to display	7816

input

$$\operatorname{int}(\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3/x^4, x, \operatorname{method}=\_RETURNVERBOSE)$$

output

$$-1/3 \operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3/x^3 + b*(-1/2 \operatorname{arctanh}(\operatorname{tanh}(b*x+a))^2/x^2 + b*(-\operatorname{arctanh}(\operatorname{tanh}(b*x+a))/x + b*\ln(x)))$$

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = \frac{6b^3x^3 \log(x) - 18ab^2x^2 - 9a^2bx - 2a^3}{6x^3}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^4,x, algorithm="fricas")`

output `1/6*(6*b^3*x^3*log(x) - 18*a*b^2*x^2 - 9*a^2*b*x - 2*a^3)/x^3`

**Sympy [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = b^3 \log(x) - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))}{x} - \frac{b \operatorname{atanh}^2(\tanh(a + bx))}{2x^2} - \frac{\operatorname{atanh}^3(\tanh(a + bx))}{3x^3}$$

input `integrate(atanh(tanh(b*x+a))**3/x**4,x)`

output `b**3*log(x) - b**2*atanh(tanh(a + b*x))/x - b*atanh(tanh(a + b*x))**2/(2*x**2) - atanh(tanh(a + b*x))**3/(3*x**3)`

**Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = \left( b^2 \log(x) - \frac{b \operatorname{artanh}(\tanh(bx + a))}{x} \right) b - \frac{b \operatorname{artanh}(\tanh(bx + a))^2}{2x^2} - \frac{\operatorname{artanh}(\tanh(bx + a))^3}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^4,x, algorithm="maxima")`



output  $(b^2 \log(x) - b \operatorname{arctanh}(\tanh(bx + a))/x) * b - 1/2 * b \operatorname{arctanh}(\tanh(bx + a))^2 / x^2 - 1/3 * \operatorname{arctanh}(\tanh(bx + a))^3 / x^3$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.64

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx = b^3 \log(|x|) - \frac{18ab^2x^2 + 9a^2bx + 2a^3}{6x^3}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^4,x, algorithm="giac")`

output  $b^3 \log(\operatorname{abs}(x)) - 1/6 * (18 * a * b^2 * x^2 + 9 * a^2 * b * x + 2 * a^3) / x^3$

### Mupad [B] (verification not implemented)

Time = 3.30 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx$$

$$= b^3 \ln(x) - \frac{b^2 x^2 \operatorname{atanh}(\tanh(a + bx)) + \frac{bx \operatorname{atanh}(\tanh(a + bx))^2}{2} + \frac{\operatorname{atanh}(\tanh(a + bx))^3}{3}}{x^3}$$

input `int(atanh(tanh(a + b*x))^3/x^4,x)`

output  $b^3 \log(x) - (\operatorname{atanh}(\tanh(a + b*x))^3 / 3 + (b*x * \operatorname{atanh}(\tanh(a + b*x))^2) / 2 + b^2 * x^2 * \operatorname{atanh}(\tanh(a + b*x))) / x^3$

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx$$

$$= \frac{-2 \operatorname{atanh}(\tanh(bx + a))^3 - 3 \operatorname{atanh}(\tanh(bx + a))^2 bx - 6 \operatorname{atanh}(\tanh(bx + a)) b^2 x^2 + 6 \log(x) b^3 x^3}{6x^3}$$

input `int(atanh(tanh(b*x+a))^3/x^4,x)`output `( - 2*atanh(tanh(a + b*x))**3 - 3*atanh(tanh(a + b*x))**2*b*x - 6*atanh(tanh(a + b*x))*b**2*x**2 + 6*log(x)*b**3*x**3)/(6*x**3)`

### 3.63 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^5} dx$

Optimal result	542
Mathematica [A] (verified)	542
Rubi [A] (verified)	543
Maple [A] (verified)	543
Fricas [A] (verification not implemented)	544
Sympy [B] (verification not implemented)	544
Maxima [A] (verification not implemented)	545
Giac [A] (verification not implemented)	545
Mupad [B] (verification not implemented)	546
Reduce [B] (verification not implemented)	546

#### Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^5} dx = \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{4x^4(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `1/4*arctanh(tanh(b*x+a))^4/x^4/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.61

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^5} dx = \frac{b^3x^3 + b^2x^2\operatorname{arctanh}(\tanh(a+bx)) + bx\operatorname{arctanh}(\tanh(a+bx))^2 + \operatorname{arctanh}(\tanh(a+bx))^3}{4x^4}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^3/x^5,x]`

output `-1/4*(b^3*x^3 + b^2*x^2*ArcTanh[Tanh[a + b*x]] + b*x*ArcTanh[Tanh[a + b*x]]^2 + ArcTanh[Tanh[a + b*x]]^3)/x^4`

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^3/x^5,x]`

output `ArcTanh[Tanh[a + b*x]]^4/(4*x^4*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 1.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

method	result	size
parallelsch	$-\frac{b^3x^3 + b^2x^2 \operatorname{arctanh}(\tanh(bx+a)) + x \operatorname{arctanh}(\tanh(bx+a))^2 b + \operatorname{arctanh}(\tanh(bx+a))^3}{4x^4}$	49
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{4x^4} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b \left( -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2} \right)}{3} \right)}{4}$	56
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{4x^4} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b \left( -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2x^2} \right)}{3} \right)}{4}$	56
risch	Expression too large to display	7814

input `int(arctanh(tanh(b*x+a))^3/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*(b^3*x^3+b^2*x^2*arctanh(tanh(b*x+a))+x*arctanh(tanh(b*x+a))^2*b+arctanh(tanh(b*x+a))^3)/x^4`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = -\frac{4b^3x^3 + 6ab^2x^2 + 4a^2bx + a^3}{4x^4}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^5,x, algorithm="fricas")`

output `-1/4*(4*b^3*x^3 + 6*a*b^2*x^2 + 4*a^2*b*x + a^3)/x^4`

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(26) = 52.

Time = 0.31 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.81

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = -\frac{b^3}{4x} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))}{4x^2} - \frac{b \operatorname{atanh}^2(\tanh(a + bx))}{4x^3} - \frac{\operatorname{atanh}^3(\tanh(a + bx))}{4x^4}$$

input `integrate(atanh(tanh(b*x+a))**3/x**5,x)`

output 
$$-b^3/(4x) - b^2 \operatorname{atanh}(\tanh(a + bx))/(4x^2) - b \operatorname{atanh}(\tanh(a + bx)) * 2/(4x^3) - \operatorname{atanh}(\tanh(a + bx))^3/(4x^4)$$

### Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = -\frac{1}{4} b \left( \frac{b^2}{x} + \frac{b \operatorname{arctanh}(\tanh(bx + a))}{x^2} \right) - \frac{b \operatorname{arctanh}(\tanh(bx + a))^2}{4x^3} - \frac{\operatorname{arctanh}(\tanh(bx + a))^3}{4x^4}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^5,x, algorithm="maxima")`

output 
$$-1/4*b*(b^2/x + b*\operatorname{arctanh}(\tanh(b*x + a))/x^2) - 1/4*b*\operatorname{arctanh}(\tanh(b*x + a))^2/x^3 - 1/4*\operatorname{arctanh}(\tanh(b*x + a))^3/x^4$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = -\frac{4b^3x^3 + 6ab^2x^2 + 4a^2bx + a^3}{4x^4}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^5,x, algorithm="giac")`

output 
$$-1/4*(4*b^3*x^3 + 6*a*b^2*x^2 + 4*a^2*b*x + a^3)/x^4$$

**Mupad [B] (verification not implemented)**

Time = 3.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = \frac{b^3 x^3 + b^2 x^2 \operatorname{atanh}(\tanh(a + bx)) + bx \operatorname{atanh}(\tanh(a + bx))^2 + \operatorname{atanh}(\tanh(a + bx))^3}{4x^4}$$

input `int(atanh(tanh(a + b*x))^3/x^5,x)`output `-(atanh(tanh(a + b*x))^3 + b^3*x^3 + b*x*atanh(tanh(a + b*x))^2 + b^2*x^2*atanh(tanh(a + b*x)))/(4*x^4)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx = \frac{-\operatorname{atanh}(\tanh(bx + a))^3 - \operatorname{atanh}(\tanh(bx + a))^2 bx - \operatorname{atanh}(\tanh(bx + a)) b^2 x^2 - b^3 x^3}{4x^4}$$

input `int(atanh(tanh(b*x+a))^3/x^5,x)`output `( - (atanh(tanh(a + b*x)))**3 + atanh(tanh(a + b*x))**2*b*x + atanh(tanh(a + b*x))*b**2*x**2 + b**3*x**3)/(4*x**4)`

### 3.64 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^6} dx$

Optimal result . . . . .	547
Mathematica [A] (verified) . . . . .	547
Rubi [A] (verified) . . . . .	548
Maple [A] (verified) . . . . .	549
Fricas [A] (verification not implemented) . . . . .	550
Sympy [A] (verification not implemented) . . . . .	550
Maxima [A] (verification not implemented) . . . . .	550
Giac [A] (verification not implemented) . . . . .	551
Mupad [B] (verification not implemented) . . . . .	551
Reduce [B] (verification not implemented) . . . . .	552

#### Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{20x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))^2} + \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{5x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output  $\frac{1}{20}b*\operatorname{arctanh}(\tanh(b*x+a))^4/x^4/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2+1/5*\operatorname{arctanh}(\tanh(b*x+a))^4/x^5/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.84

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = \frac{b^3x^3 + 2b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 3bx\operatorname{arctanh}(\tanh(a + bx))^2 + 4\operatorname{arctanh}(\tanh(a + bx))^3}{20x^5}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^3/x^6,x]`



output

$$-1/20*(b^3*x^3 + 2*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 3*b*x*ArcTanh[Tanh[a + b*x]]^2 + 4*ArcTanh[Tanh[a + b*x]]^3)/x^5$$

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx$$

↓ 2602

$$\frac{b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^5} dx}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{5x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^4}{5x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{b \operatorname{arctanh}(\tanh(a + bx))^4}{20x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^3/x^6, x]
```

output

```
(b*ArcTanh[Tanh[a + b*x]]^4)/(20*x^4*(b*x - ArcTanh[Tanh[a + b*x]])^2) + ArcTanh[Tanh[a + b*x]]^4/(5*x^5*(b*x - ArcTanh[Tanh[a + b*x]]))
```

**Defintions of rubi rules used**

```
rule 2598 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 2602 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 1.30 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

method	result	size
parallelrisc	$-\frac{b^3x^3+2b^2x^2 \operatorname{arctanh}(\tanh(bx+a))+3x \operatorname{arctanh}(\tanh(bx+a))^2b+4 \operatorname{arctanh}(\tanh(bx+a))^3}{20x^5}$	53
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{5x^5} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b \left( -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2 \cdot 3x^3} \right)}{5} \right)}{5}$	56
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{5x^5} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b \left( -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2 \cdot 3x^3} \right)}{5} \right)}{5}$	56
risc	Expression too large to display	7813

```
input int(arctanh(tanh(b*x+a))^3/x^6,x,method=_RETURNVERBOSE)
```

```
output -1/20*(b^3*x^3+2*b^2*x^2*arctanh(tanh(b*x+a))+3*x*arctanh(tanh(b*x+a))^2*b+4*arctanh(tanh(b*x+a))^3)/x^5
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = -\frac{10b^3x^3 + 20ab^2x^2 + 15a^2bx + 4a^3}{20x^5}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^6,x, algorithm="fricas")`

output `-1/20*(10*b^3*x^3 + 20*a*b^2*x^2 + 15*a^2*b*x + 4*a^3)/x^5`

**Sympy [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = -\frac{b^3}{20x^2} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))}{10x^3} - \frac{3b \operatorname{atanh}^2(\tanh(a + bx))}{20x^4} - \frac{\operatorname{atanh}^3(\tanh(a + bx))}{5x^5}$$

input `integrate(atanh(tanh(b*x+a))**3/x**6,x)`

output `-b**3/(20*x**2) - b**2*atanh(tanh(a + b*x))/(10*x**3) - 3*b*atanh(tanh(a + b*x))**2/(20*x**4) - atanh(tanh(a + b*x))**3/(5*x**5)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.84

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = -\frac{1}{20} b \left( \frac{b^2}{x^2} + \frac{2b \operatorname{atanh}(\tanh(bx + a))}{x^3} \right) - \frac{3b \operatorname{atanh}(\tanh(bx + a))^2}{20x^4} - \frac{\operatorname{atanh}(\tanh(bx + a))^3}{5x^5}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^6,x, algorithm="maxima")`

output

```
-1/20*b*(b^2/x^2 + 2*b*arctanh(tanh(b*x + a))/x^3) - 3/20*b*arctanh(tanh(b*x + a))^2/x^4 - 1/5*arctanh(tanh(b*x + a))^3/x^5
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = -\frac{10b^3x^3 + 20ab^2x^2 + 15a^2bx + 4a^3}{20x^5}$$

input

```
integrate(arctanh(tanh(b*x+a))^3/x^6,x, algorithm="giac")
```

output

```
-1/20*(10*b^3*x^3 + 20*a*b^2*x^2 + 15*a^2*b*x + 4*a^3)/x^5
```

**Mupad [B] (verification not implemented)**

Time = 3.31 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^3}{5x^5} - \frac{b^3}{20x^2} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))}{10x^3} - \frac{3b \operatorname{atanh}(\tanh(a + bx))^2}{20x^4}$$

input

```
int(atanh(tanh(a + b*x))^3/x^6,x)
```

output

```
- atanh(tanh(a + b*x))^3/(5*x^5) - b^3/(20*x^2) - (b^2*atanh(tanh(a + b*x)))/(10*x^3) - (3*b*atanh(tanh(a + b*x))^2)/(20*x^4)
```

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^6} dx$$

$$= \frac{-4\operatorname{atanh}(\tanh(bx + a))^3 - 3\operatorname{atanh}(\tanh(bx + a))^2 bx - 2\operatorname{atanh}(\tanh(bx + a)) b^2 x^2 - b^3 x^3}{20x^5}$$

input `int(atanh(tanh(b*x+a))^3/x^6,x)`output `( - 4*atanh(tanh(a + b*x))**3 - 3*atanh(tanh(a + b*x))**2*b*x - 2*atanh(tanh(a + b*x))*b**2*x**2 - b**3*x**3)/(20*x**5)`

### 3.65 $\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	553
Mathematica [A] (verified)	554
Rubi [A] (verified)	554
Maple [A] (verified)	556
Fricas [B] (verification not implemented)	557
Sympy [F]	557
Maxima [A] (verification not implemented)	558
Giac [F]	559
Mupad [B] (verification not implemented)	559
Reduce [F]	560

#### Optimal result

Integrand size = 13, antiderivative size = 154

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{24b^4 x^{5+m}}{(1+m)(2+m)(3+m)(20+9m+m^2)} - \frac{24b^3 x^{4+m} \operatorname{arctanh}(\tanh(a + bx))}{(1+m)(24+26m+9m^2+m^3)} + \frac{12b^2 x^{3+m} \operatorname{arctanh}(\tanh(a + bx))^2}{6+11m+6m^2+m^3} - \frac{4bx^{2+m} \operatorname{arctanh}(\tanh(a + bx))^3}{2+3m+m^2} + \frac{x^{1+m} \operatorname{arctanh}(\tanh(a + bx))^4}{1+m}$$

output

```
24*b^4*x^(5+m)/(1+m)/(2+m)/(3+m)/(m^2+9*m+20)-24*b^3*x^(4+m)*arctanh(tanh(b*x+a))/(1+m)/(m^3+9*m^2+26*m+24)+12*b^2*x^(3+m)*arctanh(tanh(b*x+a))^2/(m^3+6*m^2+11*m+6)-4*b*x^(2+m)*arctanh(tanh(b*x+a))^3/(m^2+3*m+2)+x^(1+m)*arctanh(tanh(b*x+a))^4/(1+m)
```

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.89

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{x^{1+m}(24b^4x^4 - 24b^3(5+m)x^3 \operatorname{arctanh}(\tanh(a + bx)) + 12b^2(20 + 9m + m^2)x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 4b(60 + 47m + 12m^2 + m^3)x \operatorname{arctanh}(\tanh(a + bx))^3 + (120 + 154m + 71m^2 + 14m^3 + m^4) \operatorname{arctanh}(\tanh(a + bx))^4)}{(1+m)(2+m)(3+m)(4+m)(5+m)}$$

input

Integrate[x^m\*ArcTanh[Tanh[a + b\*x]]^4,x]

output

$$\frac{(x^{1+m}(24b^4x^4 - 24b^3(5+m)x^3 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]] + 12b^2(20 + 9m + m^2)x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2 - 4b(60 + 47m + 12m^2 + m^3)x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3 + (120 + 154m + 71m^2 + 14m^3 + m^4) \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^4))}{(1+m)(2+m)(3+m)(4+m)(5+m)}$$
**Rubi [A] (verified)**Time = 0.35 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.81, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^4}{m+1} - \frac{4b \int x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^3 dx}{m+1}$$

$$\downarrow 2599$$

$$\frac{x^{m+1} \operatorname{arctanh}(\tanh(a + bx))^4}{m+1} - \frac{4b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a + bx))^3}{m+2} - \frac{3b \int x^{m+2} \operatorname{arctanh}(\tanh(a + bx))^2 dx}{m+2} \right)}{m+1}$$

$$\downarrow 2599$$

$$\begin{array}{c}
 \frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^4}{m+1} - \\
 4b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))^3}{m+2} - \frac{3b \left( \frac{x^{m+3} \operatorname{arctanh}(\tanh(a+bx))^2}{m+3} - \frac{2b \int x^{m+3} \operatorname{arctanh}(\tanh(a+bx)) dx}{m+3} \right)}{m+2} \right) \\
 \hline
 m+1 \\
 \downarrow 2599 \\
 \frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^4}{m+1} - \\
 4b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))^3}{m+2} - \frac{3b \left( \frac{x^{m+3} \operatorname{arctanh}(\tanh(a+bx))^2}{m+3} - \frac{2b \left( \frac{x^{m+4} \operatorname{arctanh}(\tanh(a+bx))}{m+4} - \frac{b \int x^{m+4} dx}{m+4} \right)}{m+3} \right)}{m+2} \right) \\
 \hline
 m+1 \\
 \downarrow 15 \\
 \frac{x^{m+1} \operatorname{arctanh}(\tanh(a+bx))^4}{m+1} - \\
 4b \left( \frac{x^{m+2} \operatorname{arctanh}(\tanh(a+bx))^3}{m+2} - \frac{3b \left( \frac{x^{m+3} \operatorname{arctanh}(\tanh(a+bx))^2}{m+3} - \frac{2b \left( \frac{x^{m+4} \operatorname{arctanh}(\tanh(a+bx))}{m+4} - \frac{bx^{m+5}}{(m+4)(m+5)} \right)}{m+3} \right)}{m+2} \right) \\
 \hline
 m+1
 \end{array}$$

input `Int [x^m*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^(1 + m)*ArcTanh[Tanh[a + b*x]]^4)/(1 + m) - (4*b*((x^(2 + m)*ArcTanh[Tanh[a + b*x]]^3)/(2 + m) - (3*b*((x^(3 + m)*ArcTanh[Tanh[a + b*x]]^2)/(3 + m) - (2*b*(-((b*x^(5 + m))/((4 + m)*(5 + m))) + (x^(4 + m)*ArcTanh[Tanh[a + b*x]])/(4 + m)))/(3 + m)))/(2 + m))/(1 + m)`



## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 46.25 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.81

method	result
default	$\frac{b^4 x^5 e^{m \ln(x)}}{5+m} + \frac{(a^4 + 4a^3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 6a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + 4a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3 + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^4)}{1+m} x^m$
parallelrisch	$-\frac{240b \operatorname{arctanh}(\tanh(bx+a))^3 x^m x^2 - 240b^2 \operatorname{arctanh}(\tanh(bx+a))^2 x^m x^3 + 120b^3 \operatorname{arctanh}(\tanh(bx+a)) x^m x^4 + 188x^2 x^m \operatorname{arctanh}(\tanh(bx+a))}{(1+m)x^m}$
risch	Expression too large to display

input `int(x^m*arctanh(tanh(b*x+a))^4,x,method=_RETURNVERBOSE)`

output  $b^4/(5+m)x^5 \exp(m \ln(x)) + (a^4 + 4a^3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 6a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + 4a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3 + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^4)/(1+m)x^m \exp(m \ln(x)) + 4*b*(a^3 + 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3)/(2+m)x^2 \exp(m \ln(x)) + 6*b^2*(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2)/(3+m)x^3 \exp(m \ln(x)) + 4*b^3*(\operatorname{arctanh}(\tanh(bx+a)) - bx)/(4+m)x^4 \exp(m \ln(x))$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 483 vs.  $2(154) = 308$ .

Time = 0.09 (sec) , antiderivative size = 483, normalized size of antiderivative = 3.14

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{((b^4 m^4 + 10 b^4 m^3 + 35 b^4 m^2 + 50 b^4 m + 24 b^4) x^5 + 4 (ab^3 m^4 + 11 ab^3 m^3 + 41 ab^3 m^2 + 61 ab^3 m + 30 ab^3$$

input `integrate(x^m*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`

output

```
((b^4*m^4 + 10*b^4*m^3 + 35*b^4*m^2 + 50*b^4*m + 24*b^4)*x^5 + 4*(a*b^3*m^4 + 11*a*b^3*m^3 + 41*a*b^3*m^2 + 61*a*b^3*m + 30*a*b^3)*x^4 + 6*(a^2*b^2*m^4 + 12*a^2*b^2*m^3 + 49*a^2*b^2*m^2 + 78*a^2*b^2*m + 40*a^2*b^2)*x^3 + 4*(a^3*b*m^4 + 13*a^3*b*m^3 + 59*a^3*b*m^2 + 107*a^3*b*m + 60*a^3*b)*x^2 + (a^4*m^4 + 14*a^4*m^3 + 71*a^4*m^2 + 154*a^4*m + 120*a^4)*x)*cosh(m*log(x)) + ((b^4*m^4 + 10*b^4*m^3 + 35*b^4*m^2 + 50*b^4*m + 24*b^4)*x^5 + 4*(a*b^3*m^4 + 11*a*b^3*m^3 + 41*a*b^3*m^2 + 61*a*b^3*m + 30*a*b^3)*x^4 + 6*(a^2*b^2*m^4 + 12*a^2*b^2*m^3 + 49*a^2*b^2*m^2 + 78*a^2*b^2*m + 40*a^2*b^2)*x^3 + 4*(a^3*b*m^4 + 13*a^3*b*m^3 + 59*a^3*b*m^2 + 107*a^3*b*m + 60*a^3*b)*x^2 + (a^4*m^4 + 14*a^4*m^3 + 71*a^4*m^2 + 154*a^4*m + 120*a^4)*x)*sinh(m*log(x)))/(m^5 + 15*m^4 + 85*m^3 + 225*m^2 + 274*m + 120)
```

**Sympy [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx = \text{Too large to display}$$

input `integrate(x**m*atanh(tanh(b*x+a))**4,x)`

output

```
Piecewise((b**4*log(x) - b**3*atanh(tanh(a + b*x))/x - b**2*atanh(tanh(a +
b*x))**2/(2*x**2) - b*atanh(tanh(a + b*x))**3/(3*x**3) - atanh(tanh(a + b
*x))**4/(4*x**4), Eq(m, -5)), (Integral(atanh(tanh(a + b*x))**4/x**4, x),
Eq(m, -4)), (Integral(atanh(tanh(a + b*x))**4/x**3, x), Eq(m, -3)), (Integ
ral(atanh(tanh(a + b*x))**4/x**2, x), Eq(m, -2)), (Integral(atanh(tanh(a +
b*x))**4/x, x), Eq(m, -1)), (24*b**4*x**5*x**m/(m**5 + 15*m**4 + 85*m**3
+ 225*m**2 + 274*m + 120) - 24*b**3*m*x**4*x**m*atanh(tanh(a + b*x))/(m**5
+ 15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120) - 120*b**3*x**4*x**m*atanh(
tanh(a + b*x))/(m**5 + 15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120) + 12*b*
**2*m**2*x**3*x**m*atanh(tanh(a + b*x))**2/(m**5 + 15*m**4 + 85*m**3 + 225*
m**2 + 274*m + 120) + 108*b**2*m*x**3*x**m*atanh(tanh(a + b*x))**2/(m**5 +
15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120) + 240*b**2*x**3*x**m*atanh(ta
nh(a + b*x))**2/(m**5 + 15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120) - 4*b*
m**3*x**2*x**m*atanh(tanh(a + b*x))**3/(m**5 + 15*m**4 + 85*m**3 + 225*m**
2 + 274*m + 120) - 48*b*m**2*x**2*x**m*atanh(tanh(a + b*x))**3/(m**5 + 15*
m**4 + 85*m**3 + 225*m**2 + 274*m + 120) - 188*b*m*x**2*x**m*atanh(tanh(a
+ b*x))**3/(m**5 + 15*m**4 + 85*m**3 + 225*m**2 + 274*m + 120) - 240*b*x**
2*x**m*atanh(tanh(a + b*x))**3/(m**5 + 15*m**4 + 85*m**3 + 225*m**2 + 274*
m + 120) + m**4*x*x**m*atanh(tanh(a + b*x))**4/(m**5 + 15*m**4 + 85*m**3 +
225*m**2 + 274*m + 120) + 14*m**3*x*x**m*atanh(tanh(a + b*x))**4/(m**5...
```

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.94

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= -\frac{4bx^2 x^m \operatorname{arctanh}(\tanh(bx + a))^3}{(m+2)(m+1)} + \frac{x^{m+1} \operatorname{arctanh}(\tanh(bx + a))^4}{m+1}$$

$$+ \frac{12 \left( \frac{bx^3 x^m \operatorname{arctanh}(\tanh(bx+a))^2}{(m+3)(m+2)} + \frac{2 \left( \frac{b^2 x^5 x^m}{(m+5)(m+4)(m+3)} - \frac{bx^4 x^m \operatorname{arctanh}(\tanh(bx+a))}{(m+4)(m+3)} \right) b}{m+2} \right)}{m+1}$$

input

```
integrate(x^m*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")
```

output

```
-4*b*x^2*x^m*arctanh(tanh(b*x + a))^3/((m + 2)*(m + 1)) + x^(m + 1)*arctan
h(tanh(b*x + a))^4/(m + 1) + 12*(b*x^3*x^m*arctanh(tanh(b*x + a))^2/((m +
3)*(m + 2)) + 2*(b^2*x^5*x^m/((m + 5)*(m + 4)*(m + 3)) - b*x^4*x^m*arctanh
(tanh(b*x + a))/((m + 4)*(m + 3)))*b/(m + 2))*b/(m + 1)
```

**Giac [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int x^m \operatorname{artanh}(\tanh(bx + a))^4 dx$$

input

```
integrate(x^m*arctanh(tanh(b*x+a))^4,x, algorithm="giac")
```

output

```
integrate(x^m*arctanh(tanh(b*x + a))^4, x)
```

**Mupad [B] (verification not implemented)**

Time = 3.66 (sec) , antiderivative size = 479, normalized size of antiderivative = 3.11

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{x x^m \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^4 (m^4 + 14m^3 + 71m^2 + 154m + 120)}{16m^5 + 240m^4 + 1360m^3 + 3600m^2 + 4384m + 1920}$$

$$+ \frac{16b^4 x^m x^5 (m^4 + 10m^3 + 35m^2 + 50m + 24)}{16m^5 + 240m^4 + 1360m^3 + 3600m^2 + 4384m + 1920}$$

$$+ \frac{24b^2 x^m x^3 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2 (m^4 + 12m^3 + 49m^2 + 78m + 40)}{16m^5 + 240m^4 + 1360m^3 + 3600m^2 + 4384m + 1920}$$

$$- \frac{32b^3 x^m x^4 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right) (m^4 + 11m^3 + 41m^2 + 61m + 30)}{16m^5 + 240m^4 + 1360m^3 + 3600m^2 + 4384m + 1920}$$

$$- \frac{8bx^m x^2 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3 (m^4 + 13m^3 + 59m^2 + 107m + 60)}{16m^5 + 240m^4 + 1360m^3 + 3600m^2 + 4384m + 1920}$$

input

```
int(x^m*atanh(tanh(a + b*x))^4,x)
```

output

```
(x*x^m*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4*(154*m + 71*m^2 + 14*m^3 + m^4 + 120))/(4384*m + 3600*m^2 + 1360*m^3 + 240*m^4 + 16*m^5 + 1920) + (16*b^4*x^m*x^5*(50*m + 35*m^2 + 10*m^3 + m^4 + 24))/(4384*m + 3600*m^2 + 1360*m^3 + 240*m^4 + 16*m^5 + 1920) + (24*b^2*x^m*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2*(78*m + 49*m^2 + 12*m^3 + m^4 + 40))/(4384*m + 3600*m^2 + 1360*m^3 + 240*m^4 + 16*m^5 + 1920) - (32*b^3*x^m*x^4*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*(61*m + 41*m^2 + 11*m^3 + m^4 + 30))/(4384*m + 3600*m^2 + 1360*m^3 + 240*m^4 + 16*m^5 + 1920) - (8*b*x^m*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3*(107*m + 59*m^2 + 13*m^3 + m^4 + 60))/(4384*m + 3600*m^2 + 1360*m^3 + 240*m^4 + 16*m^5 + 1920)
```

**Reduce [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int x^m \operatorname{atanh}(\tanh(bx + a))^4 dx$$

input

```
int(x^m*atanh(tanh(b*x+a))^4,x)
```

output

```
int(x**m*atanh(tanh(a + b*x))**4,x)
```

### 3.66 $\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	561
Mathematica [A] (verified)	562
Rubi [A] (verified)	562
Maple [A] (verified)	564
Fricas [A] (verification not implemented)	564
Sympy [A] (verification not implemented)	565
Maxima [A] (verification not implemented)	565
Giac [A] (verification not implemented)	566
Mupad [B] (verification not implemented)	566
Reduce [F]	567

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^{11}}{2310} - \frac{1}{210} b^3 x^{10} \operatorname{arctanh}(\tanh(a + bx))$$

$$+ \frac{1}{42} b^2 x^9 \operatorname{arctanh}(\tanh(a + bx))^2$$

$$- \frac{1}{14} b x^8 \operatorname{arctanh}(\tanh(a + bx))^3$$

$$+ \frac{1}{7} x^7 \operatorname{arctanh}(\tanh(a + bx))^4$$

output

```
1/2310*b^4*x^11-1/210*b^3*x^10*arctanh(tanh(b*x+a))+1/42*b^2*x^9*arctanh(tanh(b*x+a))^2-1/14*b*x^8*arctanh(tanh(b*x+a))^3+1/7*x^7*arctanh(tanh(b*x+a))^4
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{x^7(b^4x^4 - 11b^3x^3 \operatorname{arctanh}(\tanh(a + bx)) + 55b^2x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 165bx \operatorname{arctanh}(\tanh(a + bx)))}{2310}$$

input `Integrate[x^6*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^7*(b^4*x^4 - 11*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 55*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 165*b*x*ArcTanh[Tanh[a + b*x]]^3 + 330*ArcTanh[Tanh[a + b*x]]^4))/2310`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$\downarrow 2599$$

$$\frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{7}b \int x^7 \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2599$$

$$\frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{8}b \int x^8 \operatorname{arctanh}(\tanh(a + bx))^2 dx \right)$$

$$\downarrow 2599$$

$$\frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{8}b \left( \frac{1}{9}x^9 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{9}b \int x^9 \operatorname{arctanh}(\tanh(a + bx)) dx \right) \right)$$

↓ 2599

$$\frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{8}b \left( \frac{1}{9}x^9 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{9}b \left( \frac{1}{10}x^{10} \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^{10} \operatorname{arctanh}(\tanh(a + bx)) dx}{10} \right) \right) \right)$$

↓ 15

$$\frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{8}b \left( \frac{1}{9}x^9 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{9}b \left( \frac{1}{10}x^{10} \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^{11}}{110} \right) \right) \right)$$

input `Int[x^6*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^7*ArcTanh[Tanh[a + b*x]]^4)/7 - (4*b*((x^8*ArcTanh[Tanh[a + b*x]]^3)/8 - (3*b*((x^9*ArcTanh[Tanh[a + b*x]]^2)/9 - (2*b*(-1/110*(b*x^11) + (x^10*ArcTanh[Tanh[a + b*x]])/10))/9))/8)/7`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`



**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

$$\frac{x^7 \operatorname{arctanh}(\tanh(bx+a))^4}{7} - \frac{4b \left( \frac{x^8 \operatorname{arctanh}(\tanh(bx+a))^3}{8} - \frac{3b \left( \frac{x^9 \operatorname{arctanh}(\tanh(bx+a))^2}{9} - \frac{2b \left( \frac{x^{10} \operatorname{arctanh}(\tanh(bx+a))}{10} - \frac{x^{11}b}{110} \right)}{9} \right)}{8} \right)}{7}$$

input `int(x^6*arctanh(tanh(b*x+a))^4,x)`output `1/7*x^7*arctanh(tanh(b*x+a))^4-4/7*b*(1/8*x^8*arctanh(tanh(b*x+a))^3-3/8*b*(1/9*x^9*arctanh(tanh(b*x+a))^2-2/9*b*(1/10*x^10*arctanh(tanh(b*x+a))-1/10*x^11*b)))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^6 \operatorname{arctanh}(\tanh(a+bx))^4 dx = \frac{1}{11} b^4 x^{11} + \frac{2}{5} ab^3 x^{10} + \frac{2}{3} a^2 b^2 x^9 + \frac{1}{2} a^3 b x^8 + \frac{1}{7} a^4 x^7$$

input `integrate(x^6*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`output `1/11*b^4*x^11 + 2/5*a*b^3*x^10 + 2/3*a^2*b^2*x^9 + 1/2*a^3*b*x^8 + 1/7*a^4*x^7`

**Sympy [A] (verification not implemented)**

Time = 2.57 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.68

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \begin{cases} \frac{x^6 \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{x^5 \operatorname{atanh}^6(\tanh(a+bx))}{5b^2} + \frac{x^4 \operatorname{atanh}^7(\tanh(a+bx))}{7b^3} - \frac{x^3 \operatorname{atanh}^8(\tanh(a+bx))}{14b^4} + \frac{x^2 \operatorname{atanh}^9(\tanh(a+bx))}{42b^5} \\ \frac{x^7 \operatorname{atanh}^4(\tanh(a))}{7} \end{cases}$$

input `integrate(x**6*atanh(tanh(b*x+a))**4,x)`output `Piecewise((x**6*atanh(tanh(a + b*x))**5/(5*b) - x**5*atanh(tanh(a + b*x))**6/(5*b**2) + x**4*atanh(tanh(a + b*x))**7/(7*b**3) - x**3*atanh(tanh(a + b*x))**8/(14*b**4) + x**2*atanh(tanh(a + b*x))**9/(42*b**5) - x*atanh(tanh(a + b*x))**10/(210*b**6) + atanh(tanh(a + b*x))**11/(2310*b**7), Ne(b, 0)), (x**7*atanh(tanh(a))**4/7, True))`**Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= -\frac{1}{14} bx^8 \operatorname{artanh}(\tanh(bx + a))^3 + \frac{1}{7} x^7 \operatorname{artanh}(\tanh(bx + a))^4$$

$$+ \frac{1}{2310} (55 bx^9 \operatorname{artanh}(\tanh(bx + a))^2 + (b^2 x^{11} - 11 bx^{10} \operatorname{artanh}(\tanh(bx + a)))b)b$$

input `integrate(x^6*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`output `-1/14*b*x^8*arctanh(tanh(b*x + a))^3 + 1/7*x^7*arctanh(tanh(b*x + a))^4 + 1/2310*(55*b*x^9*arctanh(tanh(b*x + a))^2 + (b^2*x^11 - 11*b*x^10*arctanh(tanh(b*x + a)))b)*b`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{11} b^4 x^{11} + \frac{2}{5} ab^3 x^{10} + \frac{2}{3} a^2 b^2 x^9 + \frac{1}{2} a^3 b x^8 + \frac{1}{7} a^4 x^7$$

input `integrate(x^6*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/11*b^4*x^11 + 2/5*a*b^3*x^10 + 2/3*a^2*b^2*x^9 + 1/2*a^3*b*x^8 + 1/7*a^4*x^7`

**Mupad [B] (verification not implemented)**

Time = 3.33 (sec) , antiderivative size = 242, normalized size of antiderivative = 3.02

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{x^7 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^4}{112} + \frac{b^4 x^{11}}{11} - \frac{bx^8 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{16} - \frac{b^3 x^{10} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{5} + \frac{b^2 x^9 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{6}$$

input `int(x^6*atanh(tanh(a + b*x))^4,x)`

output `(x^7*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)/112 + (b^4*x^11)/11 - (b*x^8*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/16 - (b^3*x^10*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/5 + (b^2*x^9*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/6`

**Reduce [F]**

$$\int x^6 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x^6 dx$$

input `int(x^6*atanh(tanh(b*x+a))^4,x)`

output `int(atanh(tanh(a + b*x))**4*x**6,x)`

### 3.67 $\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	568
Mathematica [A] (verified)	569
Rubi [A] (verified)	569
Maple [A] (verified)	571
Fricas [A] (verification not implemented)	571
Sympy [A] (verification not implemented)	572
Maxima [A] (verification not implemented)	572
Giac [A] (verification not implemented)	573
Mupad [B] (verification not implemented)	573
Reduce [F]	574

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^{10}}{1260} - \frac{1}{126} b^3 x^9 \operatorname{arctanh}(\tanh(a + bx)) + \frac{1}{28} b^2 x^8 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{21} b x^7 \operatorname{arctanh}(\tanh(a + bx))^3 + \frac{1}{6} x^6 \operatorname{arctanh}(\tanh(a + bx))^4$$

output

```
1/1260*b^4*x^10-1/126*b^3*x^9*arctanh(tanh(b*x+a))+1/28*b^2*x^8*arctanh(tanh(b*x+a))^2-2/21*b*x^7*arctanh(tanh(b*x+a))^3+1/6*x^6*arctanh(tanh(b*x+a))^4
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{x^6(b^4x^4 - 10b^3x^3 \operatorname{arctanh}(\tanh(a + bx))) + 45b^2x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 120bx \operatorname{arctanh}(\tanh(a + bx))}{1260}$$

input `Integrate[x^5*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^6*(b^4*x^4 - 10*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 45*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 120*b*x*ArcTanh[Tanh[a + b*x]]^3 + 210*ArcTanh[Tanh[a + b*x]]^4))/1260`

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$\downarrow 2599$$

$$\frac{1}{6}x^6 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{2}{3}b \int x^6 \operatorname{arctanh}(\tanh(a + bx))^3 dx$$

$$\downarrow 2599$$

$$\frac{1}{6}x^6 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{2}{3}b \left( \frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{3}{7}b \int x^7 \operatorname{arctanh}(\tanh(a + bx))^2 dx \right)$$

$$\downarrow 2599$$

$$\frac{2}{3}b \left( \frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a+bx))^3 - \frac{3}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a+bx))^2 - \frac{1}{4}b \int x^8 \operatorname{arctanh}(\tanh(a+bx)) dx \right) \right)$$

↓ 2599

$$\frac{2}{3}b \left( \frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a+bx))^3 - \frac{3}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a+bx))^2 - \frac{1}{4}b \left( \frac{1}{9}x^9 \operatorname{arctanh}(\tanh(a+bx)) - \frac{b \int x^9 dx}{9} \right) \right) \right)$$

↓ 15

$$\frac{2}{3}b \left( \frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a+bx))^3 - \frac{3}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a+bx))^2 - \frac{1}{4}b \left( \frac{1}{9}x^9 \operatorname{arctanh}(\tanh(a+bx)) - \frac{bx^{10}}{90} \right) \right) \right)$$

input `Int[x^5*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^6*ArcTanh[Tanh[a + b*x]]^4)/6 - (2*b*((x^7*ArcTanh[Tanh[a + b*x]]^3)/7 - (3*b*((x^8*ArcTanh[Tanh[a + b*x]]^2)/8 - (b*(-1/90*(b*x^10) + (x^9*ArcTanh[Tanh[a + b*x]])/9))/4))/7)/3`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

$$\frac{x^6 \operatorname{arctanh}(\tanh(bx+a))^4}{6} - \frac{2b \left( \frac{x^7 \operatorname{arctanh}(\tanh(bx+a))^3}{7} - \frac{3b \left( \frac{x^8 \operatorname{arctanh}(\tanh(bx+a))^2}{8} - \frac{b \left( \frac{x^9 \operatorname{arctanh}(\tanh(bx+a))}{9} - \frac{x^{10}b}{90} \right)}{4} \right)}{7} \right)}{3}$$

input `int(x^5*arctanh(tanh(b*x+a))^4,x)`output `1/6*x^6*arctanh(tanh(b*x+a))^4-2/3*b*(1/7*x^7*arctanh(tanh(b*x+a))^3-3/7*b*(1/8*x^8*arctanh(tanh(b*x+a))^2-1/4*b*(1/9*x^9*arctanh(tanh(b*x+a))-1/90*x^10*b))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^5 \operatorname{arctanh}(\tanh(a+bx))^4 dx = \frac{1}{10} b^4 x^{10} + \frac{4}{9} a b^3 x^9 + \frac{3}{4} a^2 b^2 x^8 + \frac{4}{7} a^3 b x^7 + \frac{1}{6} a^4 x^6$$

input `integrate(x^5*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`output `1/10*b^4*x^10 + 4/9*a*b^3*x^9 + 3/4*a^2*b^2*x^8 + 4/7*a^3*b*x^7 + 1/6*a^4*x^6`



**Sympy [A] (verification not implemented)**

Time = 1.59 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.46

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \begin{cases} \frac{x^5 \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{x^4 \operatorname{atanh}^6(\tanh(a+bx))}{6b^2} + \frac{2x^3 \operatorname{atanh}^7(\tanh(a+bx))}{21b^3} - \frac{x^2 \operatorname{atanh}^8(\tanh(a+bx))}{28b^4} + \frac{x \operatorname{atanh}^9(\tanh(a+bx))}{126b^5} \\ \frac{x^6 \operatorname{atanh}^4(\tanh(a))}{6} \end{cases}$$

input `integrate(x**5*atanh(tanh(b*x+a))**4,x)`output `Piecewise((x**5*atanh(tanh(a + b*x))**5/(5*b) - x**4*atanh(tanh(a + b*x))**6/(6*b**2) + 2*x**3*atanh(tanh(a + b*x))**7/(21*b**3) - x**2*atanh(tanh(a + b*x))**8/(28*b**4) + x*atanh(tanh(a + b*x))**9/(126*b**5) - atanh(tanh(a + b*x))**10/(1260*b**6), Ne(b, 0)), (x**6*atanh(tanh(a))**4/6, True))`**Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= -\frac{2}{21} bx^7 \operatorname{artanh}(\tanh(bx + a))^3 + \frac{1}{6} x^6 \operatorname{artanh}(\tanh(bx + a))^4$$

$$+ \frac{1}{1260} (45 bx^8 \operatorname{artanh}(\tanh(bx + a))^2 + (b^2 x^{10} - 10 bx^9 \operatorname{artanh}(\tanh(bx + a)))b)b$$

input `integrate(x^5*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`output `-2/21*b*x^7*arctanh(tanh(b*x + a))^3 + 1/6*x^6*arctanh(tanh(b*x + a))^4 + 1/1260*(45*b*x^8*arctanh(tanh(b*x + a))^2 + (b^2*x^10 - 10*b*x^9*arctanh(tanh(b*x + a))))*b)*b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{10} b^4 x^{10} + \frac{4}{9} ab^3 x^9 + \frac{3}{4} a^2 b^2 x^8 + \frac{4}{7} a^3 b x^7 + \frac{1}{6} a^4 x^6$$

input `integrate(x^5*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`output `1/10*b^4*x^10 + 4/9*a*b^3*x^9 + 3/4*a^2*b^2*x^8 + 4/7*a^3*b*x^7 + 1/6*a^4*x^6`**Mupad [B] (verification not implemented)**

Time = 3.49 (sec) , antiderivative size = 242, normalized size of antiderivative = 3.02

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{x^6 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^4}{96} + \frac{b^4 x^{10} - bx^7 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{10} - \frac{14 b^3 x^9 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{9} + \frac{3b^2 x^8 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{16}$$

input `int(x^5*atanh(tanh(a + b*x))^4,x)`output `(x^6*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)/96 + (b^4*x^10)/10 - (b*x^7*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/14 - (2*b^3*x^9*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/9 + (3*b^2*x^8*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/16`

**Reduce [F]**

$$\int x^5 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x^5 dx$$

input `int(x^5*atanh(tanh(b*x+a))^4,x)`

output `int(atanh(tanh(a + b*x))**4*x**5,x)`

### 3.68 $\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	575
Mathematica [A] (verified)	575
Rubi [A] (verified)	576
Maple [A] (verified)	577
Fricas [A] (verification not implemented)	578
Sympy [A] (verification not implemented)	578
Maxima [A] (verification not implemented)	579
Giac [A] (verification not implemented)	579
Mupad [B] (verification not implemented)	580
Reduce [F]	580

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^9}{630} - \frac{1}{70} b^3 x^8 \operatorname{arctanh}(\tanh(a + bx))$$

$$+ \frac{2}{35} b^2 x^7 \operatorname{arctanh}(\tanh(a + bx))^2$$

$$- \frac{2}{15} b x^6 \operatorname{arctanh}(\tanh(a + bx))^3$$

$$+ \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(a + bx))^4$$

output

```
1/630*b^4*x^9-1/70*b^3*x^8*arctanh(tanh(b*x+a))+2/35*b^2*x^7*arctanh(tanh(
b*x+a))^2-2/15*b*x^6*arctanh(tanh(b*x+a))^3+1/5*x^5*arctanh(tanh(b*x+a))^4
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{630} x^5 (b^4 x^4 - 9b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)))$$

$$+ 36b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2$$

$$- 84bx \operatorname{arctanh}(\tanh(a + bx))^3$$

$$+ 126 \operatorname{arctanh}(\tanh(a + bx))^4$$

input `Integrate[x^4*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^5*(b^4*x^4 - 9*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 36*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 84*b*x*ArcTanh[Tanh[a + b*x]]^3 + 126*ArcTanh[Tanh[a + b*x]]^4))/630`

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{5} b \int x^5 \operatorname{arctanh}(\tanh(a + bx))^3 dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(a + bx))^4 - \\
 & \frac{4}{5} b \left( \frac{1}{6} x^6 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{1}{2} b \int x^6 \operatorname{arctanh}(\tanh(a + bx))^2 dx \right) \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(a + bx))^4 - \\
 & \frac{4}{5} b \left( \frac{1}{6} x^6 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{1}{2} b \left( \frac{1}{7} x^7 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{7} b \int x^7 \operatorname{arctanh}(\tanh(a + bx)) dx \right) \right) \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(a + bx))^4 - \\
 & \frac{4}{5} b \left( \frac{1}{6} x^6 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{1}{2} b \left( \frac{1}{7} x^7 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{7} b \left( \frac{1}{8} x^8 \operatorname{arctanh}(\tanh(a + bx)) - \frac{b \int x^8 dx}{8} \right) \right) \right) \\
 & \quad \downarrow \text{15}
 \end{aligned}$$

$$\frac{1}{5}x^5 \operatorname{arctanh}(\tanh(a + bx))^4 - \frac{4}{5}b \left( \frac{1}{6}x^6 \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{1}{2}b \left( \frac{1}{7}x^7 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2}{7}b \left( \frac{1}{8}x^8 \operatorname{arctanh}(\tanh(a + bx)) - \frac{bx^9}{72} \right) \right) \right)$$

input `Int[x^4*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^5*ArcTanh[Tanh[a + b*x]]^4)/5 - (4*b*((x^6*ArcTanh[Tanh[a + b*x]]^3)/6 - (b*((x^7*ArcTanh[Tanh[a + b*x]]^2)/7 - (2*b*(-1/72*(b*x^9) + (x^8*ArcTanh[Tanh[a + b*x]])/8))/7))/2))/5`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

$$\frac{x^5 \operatorname{arctanh}(\tanh(bx + a))^4}{5} - \frac{4b \left( \frac{x^6 \operatorname{arctanh}(\tanh(bx + a))^3}{6} - \frac{b \left( \frac{x^7 \operatorname{arctanh}(\tanh(bx + a))^2}{7} - \frac{2b \left( \frac{x^8 \operatorname{arctanh}(\tanh(bx + a))}{8} - \frac{x^9 b}{72} \right)}{7} \right)}{2} \right)}{5}$$

input `int(x^4*arctanh(tanh(b*x+a))^4,x)`

output

```
1/5*x^5*arctanh(tanh(b*x+a))^4-4/5*b*(1/6*x^6*arctanh(tanh(b*x+a))^3-1/2*b
*(1/7*x^7*arctanh(tanh(b*x+a))^2-2/7*b*(1/8*x^8*arctanh(tanh(b*x+a))-1/72*
x^9*b)))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{9} b^4 x^9 + \frac{1}{2} a b^3 x^8 + \frac{6}{7} a^2 b^2 x^7 + \frac{2}{3} a^3 b x^6 + \frac{1}{5} a^4 x^5$$

input

```
integrate(x^4*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")
```

output

```
1/9*b^4*x^9 + 1/2*a*b^3*x^8 + 6/7*a^2*b^2*x^7 + 2/3*a^3*b*x^6 + 1/5*a^4*x^5
```

**Sympy [A] (verification not implemented)**

Time = 1.13 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.25

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \begin{cases} \frac{x^4 \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{2x^3 \operatorname{atanh}^6(\tanh(a+bx))}{15b^2} + \frac{2x^2 \operatorname{atanh}^7(\tanh(a+bx))}{35b^3} - \frac{x \operatorname{atanh}^8(\tanh(a+bx))}{70b^4} + \frac{\operatorname{atanh}^9(\tanh(a+bx))}{630b^5} \\ \frac{x^5 \operatorname{atanh}^4(\tanh(a))}{5} \end{cases}$$

input

```
integrate(x**4*atanh(tanh(b*x+a))**4,x)
```

output

```
Piecewise((x**4*atanh(tanh(a + b*x))**5/(5*b) - 2*x**3*atanh(tanh(a + b*x))
)**6/(15*b**2) + 2*x**2*atanh(tanh(a + b*x))**7/(35*b**3) - x*atanh(tanh(a
+ b*x))**8/(70*b**4) + atanh(tanh(a + b*x))**9/(630*b**5), Ne(b, 0)), (x*
**5*atanh(tanh(a))**4/5, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= -\frac{2}{15} bx^6 \operatorname{arctanh}(\tanh(bx + a))^3 + \frac{1}{5} x^5 \operatorname{arctanh}(\tanh(bx + a))^4$$

$$+ \frac{1}{630} (36 bx^7 \operatorname{arctanh}(\tanh(bx + a))^2 + (b^2 x^9 - 9 bx^8 \operatorname{arctanh}(\tanh(bx + a)))b)b$$

input `integrate(x^4*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`

output `-2/15*b*x^6*arctanh(tanh(b*x + a))^3 + 1/5*x^5*arctanh(tanh(b*x + a))^4 + 1/630*(36*b*x^7*arctanh(tanh(b*x + a))^2 + (b^2*x^9 - 9*b*x^8*arctanh(tanh(b*x + a)))*b)*b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{9} b^4 x^9 + \frac{1}{2} ab^3 x^8 + \frac{6}{7} a^2 b^2 x^7 + \frac{2}{3} a^3 b x^6 + \frac{1}{5} a^4 x^5$$

input `integrate(x^4*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/9*b^4*x^9 + 1/2*a*b^3*x^8 + 6/7*a^2*b^2*x^7 + 2/3*a^3*b*x^6 + 1/5*a^4*x^5`



**Mupad [B] (verification not implemented)**

Time = 3.47 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.96

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= \frac{\operatorname{atanh}(\tanh(a + bx))^5 (126 b^4 x^4 - 84 b^3 x^3 \operatorname{atanh}(\tanh(a + bx)) + 36 b^2 x^2 \operatorname{atanh}(\tanh(a + bx))^2 - 9 b x \operatorname{atanh}(\tanh(a + bx))^3 - 84 b^3 x^3 \operatorname{atanh}(\tanh(a + bx)))}{630 b^5}$$

input `int(x^4*atanh(tanh(a + b*x))^4,x)`output `(atanh(tanh(a + b*x))^5*(atanh(tanh(a + b*x))^4 + 126*b^4*x^4 + 36*b^2*x^2*atanh(tanh(a + b*x))^2 - 9*b*x*atanh(tanh(a + b*x))^3 - 84*b^3*x^3*atanh(tanh(a + b*x))))/(630*b^5)`**Reduce [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x^4 dx$$

input `int(x^4*atanh(tanh(b*x+a))^4,x)`output `int(atanh(tanh(a + b*x))**4*x**4,x)`

### 3.69 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	581
Mathematica [A] (verified)	581
Rubi [A] (verified)	582
Maple [A] (verified)	584
Fricas [A] (verification not implemented)	584
Sympy [A] (verification not implemented)	584
Maxima [A] (verification not implemented)	585
Giac [A] (verification not implemented)	585
Mupad [B] (verification not implemented)	586
Reduce [F]	586

#### Optimal result

Integrand size = 13, antiderivative size = 72

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{10b^2} + \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{35b^3} - \frac{\operatorname{arctanh}(\tanh(a + bx))^8}{280b^4}$$

output

```
1/5*x^3*arctanh(tanh(b*x+a))^5/b-1/10*x^2*arctanh(tanh(b*x+a))^6/b^2+1/35*
x*arctanh(tanh(b*x+a))^7/b^3-1/280*arctanh(tanh(b*x+a))^8/b^4
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.99

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{280} x^4 (b^4 x^4 - 8b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) + 28b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 56bx \operatorname{arctanh}(\tanh(a + bx))^3 + 70 \operatorname{arctanh}(\tanh(a + bx))^4)$$

input `Integrate[x^3*ArcTanh[Tanh[a + b*x]]^4,x]`

output  $(x^4*(b^4*x^4 - 8*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 28*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 56*b*x*ArcTanh[Tanh[a + b*x]]^3 + 70*ArcTanh[Tanh[a + b*x]]^4)/280$

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx \\
 & \quad \downarrow 2599 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^5 dx}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx}{3b} \right)}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 dx}{3b} \right)}{5b} \\
 & \quad \downarrow 2588 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 dx}{3b} \right)}{5b} \\
 & \quad \downarrow 2588 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 dx}{3b} \right)}{5b} \\
 & \quad \downarrow 2588 \\
 & \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 dx}{3b} \right)}{5b}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 15 \\
 \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} \\
 \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^8}{56b^2} \right)}{5b}
 \end{array}$$

input `Int [x^3*ArcTanh [Tanh [a + b*x]]^4, x]`

output `(x^3*ArcTanh [Tanh [a + b*x]]^5)/(5*b) - (3*((x^2*ArcTanh [Tanh [a + b*x]]^6)/(6*b) - ((x*ArcTanh [Tanh [a + b*x]]^7)/(7*b) - ArcTanh [Tanh [a + b*x]]^8/(56*b^2)))/(3*b)))/(5*b)`

### Defintions of rubi rules used

rule 15 `Int [(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int [(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst [Int [x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int [(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03

$$\frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^4}{4} - b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))^3}{5} - \frac{3b \left( \frac{x^6 \operatorname{arctanh}(\tanh(bx+a))^2}{6} - \frac{b \left( \frac{x^7 \operatorname{arctanh}(\tanh(bx+a))}{7} - \frac{b \operatorname{arctanh}(\tanh(bx+a))}{3} \right)}{5} \right)}{5} \right)$$

input `int(x^3*arctanh(tanh(b*x+a))^4,x)`output `1/4*x^4*arctanh(tanh(b*x+a))^4-b*(1/5*x^5*arctanh(tanh(b*x+a))^3-3/5*b*(1/6*x^6*arctanh(tanh(b*x+a))^2-1/3*b*(1/7*x^7*arctanh(tanh(b*x+a))-1/56*b*x^8)))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^4 dx = \frac{1}{8} b^4 x^8 + \frac{4}{7} ab^3 x^7 + a^2 b^2 x^6 + \frac{4}{5} a^3 b x^5 + \frac{1}{4} a^4 x^4$$

input `integrate(x^3*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`output `1/8*b^4*x^8 + 4/7*a*b^3*x^7 + a^2*b^2*x^6 + 4/5*a^3*b*x^5 + 1/4*a^4*x^4`**Sympy [A] (verification not implemented)**

Time = 0.73 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.08

$$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^4 dx = \begin{cases} \frac{x^3 \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{x^2 \operatorname{atanh}^6(\tanh(a+bx))}{10b^2} + \frac{x \operatorname{atanh}^7(\tanh(a+bx))}{35b^3} - \frac{\operatorname{atanh}^8(\tanh(a+bx))}{280b^4} & \text{for } b \neq 0 \\ \frac{x^4 \operatorname{atanh}^4(\tanh(a))}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*atanh(tanh(b*x+a))**4,x)`

output `Piecewise((x**3*atanh(tanh(a + b*x))**5/(5*b) - x**2*atanh(tanh(a + b*x))*  
*6/(10*b**2) + x*atanh(tanh(a + b*x))**7/(35*b**3) - atanh(tanh(a + b*x))*  
*8/(280*b**4), Ne(b, 0)), (x**4*atanh(tanh(a))**4/4, True))`

### Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00

$$\begin{aligned} \int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx \\ = -\frac{1}{5} bx^5 \operatorname{arctanh}(\tanh(bx + a))^3 + \frac{1}{4} x^4 \operatorname{arctanh}(\tanh(bx + a))^4 \\ + \frac{1}{280} (28 bx^6 \operatorname{arctanh}(\tanh(bx + a))^2 + (b^2 x^8 - 8 bx^7 \operatorname{arctanh}(\tanh(bx + a)))b) b \end{aligned}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`

output `-1/5*b*x^5*arctanh(tanh(b*x + a))^3 + 1/4*x^4*arctanh(tanh(b*x + a))^4 + 1  
/280*(28*b*x^6*arctanh(tanh(b*x + a))^2 + (b^2*x^8 - 8*b*x^7*arctanh(tanh(  
b*x + a))))*b)*b`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{8} b^4 x^8 + \frac{4}{7} ab^3 x^7 + a^2 b^2 x^6 + \frac{4}{5} a^3 b x^5 + \frac{1}{4} a^4 x^4$$

input `integrate(x^3*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/8*b^4*x^8 + 4/7*a*b^3*x^7 + a^2*b^2*x^6 + 4/5*a^3*b*x^5 + 1/4*a^4*x^4`

**Mupad [B] (verification not implemented)**

Time = 3.89 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.97

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^8}{280} - \frac{b^3 x^7 \operatorname{atanh}(\tanh(a + bx))}{35} + \frac{b^2 x^6 \operatorname{atanh}(\tanh(a + bx))^2}{10} - \frac{b x^5 \operatorname{atanh}(\tanh(a + bx))^3}{5} + \frac{x^4 \operatorname{atanh}(\tanh(a + bx))^4}{4}$$

input `int(x^3*atanh(tanh(a + b*x))^4,x)`output `(x^4*atanh(tanh(a + b*x))^4)/4 + (b^4*x^8)/280 + (b^2*x^6*atanh(tanh(a + b*x))^2)/10 - (b*x^5*atanh(tanh(a + b*x))^3)/5 - (b^3*x^7*atanh(tanh(a + b*x)))/35`**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x^3 dx$$

input `int(x^3*atanh(tanh(b*x+a))^4,x)`output `int(atanh(tanh(a + b*x))**4*x**3,x)`

### 3.70 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	587
Mathematica [A] (verified)	587
Rubi [A] (verified)	588
Maple [A] (verified)	589
Fricas [A] (verification not implemented)	590
Sympy [A] (verification not implemented)	590
Maxima [A] (verification not implemented)	591
Giac [A] (verification not implemented)	591
Mupad [B] (verification not implemented)	592
Reduce [F]	592

#### Optimal result

Integrand size = 13, antiderivative size = 53

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{x \operatorname{arctanh}(\tanh(a + bx))^6}{15b^2} + \frac{\operatorname{arctanh}(\tanh(a + bx))^7}{105b^3}$$

output `1/5*x^2*arctanh(tanh(b*x+a))^5/b-1/15*x*arctanh(tanh(b*x+a))^6/b^2+1/105*arctanh(tanh(b*x+a))^7/b^3`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.34

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{105} x^3 (b^4 x^4 - 7b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) + 21b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 35bx \operatorname{arctanh}(\tanh(a + bx))^3 + 35 \operatorname{arctanh}(\tanh(a + bx))^4)$$



input `Integrate[x^2*ArcTanh[Tanh[a + b*x]]^4,x]`

output `(x^3*(b^4*x^4 - 7*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 21*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 35*b*x*ArcTanh[Tanh[a + b*x]]^3 + 35*ArcTanh[Tanh[a + b*x]]^4)/105`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.15, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx \\
 & \quad \downarrow 2599 \\
 & \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{2 \int x \operatorname{arctanh}(\tanh(a + bx))^5 dx}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^6 dx}{6b} \right)}{5b} \\
 & \quad \downarrow 2588 \\
 & \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^6 d \operatorname{arctanh}(\tanh(a + bx))}{6b^2} \right)}{5b} \\
 & \quad \downarrow 15 \\
 & \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^6}{6b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^7}{42b^2} \right)}{5b}
 \end{aligned}$$

input `Int[x^2*ArcTanh[Tanh[a + b*x]]^4,x]`

output  $(x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^5)/(5*b) - (2*((x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^6)/(6*b) - \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^7/(42*b^2)))/(5*b)$

### Defintions of rubi rules used

rule 15  $\operatorname{Int}[(a_.)*(x_)^(m_.), x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^(m + 1)/(m + 1)), x] \;/; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$

rule 2588  $\operatorname{Int}[(u_)^(m_.), x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] \;/; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$

rule 2599  $\operatorname{Int}[(u_)^(m_.)*(v_)^(n_.), x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^(m + 1)*(v^n/(a*(m + 1))), x] - \operatorname{Simp}[b*(n/(a*(m + 1))) \operatorname{Int}[u^(m + 1)*v^(n - 1), x], x] \;/; \operatorname{NeQ}[b*u - a*v, 0] \;/; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m + n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n + m + 1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$

### Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.40

$$\frac{x^3 \operatorname{arctanh}(\tanh(bx + a))^4}{3} - \frac{4b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx + a))^3}{4} - \frac{3b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx + a))^2}{5} - \frac{2b \left( \frac{x^6 \operatorname{arctanh}(\tanh(bx + a))}{6} - \frac{bx^7}{42} \right)}{5} \right)}{4} \right)}{3}$$

input  $\operatorname{int}(x^2 \operatorname{arctanh}(\tanh(b*x+a))^4, x)$

output

```
1/3*x^3*arctanh(tanh(b*x+a))^4-4/3*b*(1/4*x^4*arctanh(tanh(b*x+a))^3-3/4*b
*(1/5*x^5*arctanh(tanh(b*x+a))^2-2/5*b*(1/6*x^6*arctanh(tanh(b*x+a))-1/42*
b*x^7)))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.85

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{7} b^4 x^7 + \frac{2}{3} a b^3 x^6 + \frac{6}{5} a^2 b^2 x^5 + a^3 b x^4 + \frac{1}{3} a^4 x^3$$

input

```
integrate(x^2*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")
```

output

```
1/7*b^4*x^7 + 2/3*a*b^3*x^6 + 6/5*a^2*b^2*x^5 + a^3*b*x^4 + 1/3*a^4*x^3
```

**Sympy [A] (verification not implemented)**

Time = 0.49 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.13

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \begin{cases} \frac{x^2 \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{x \operatorname{atanh}^6(\tanh(a+bx))}{15b^2} + \frac{\operatorname{atanh}^7(\tanh(a+bx))}{105b^3} & \text{for } b \neq 0 \\ \frac{x^3 \operatorname{atanh}^4(\tanh(a))}{3} & \text{otherwise} \end{cases}$$

input

```
integrate(x**2*atanh(tanh(b*x+a))**4,x)
```

output

```
Piecewise((x**2*atanh(tanh(a + b*x))**5/(5*b) - x*atanh(tanh(a + b*x))**6/
(15*b**2) + atanh(tanh(a + b*x))**7/(105*b**3), Ne(b, 0)), (x**3*atanh(tan
h(a))**4/3, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.36

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$= -\frac{1}{3} bx^4 \operatorname{arctanh}(\tanh(bx + a))^3 + \frac{1}{3} x^3 \operatorname{arctanh}(\tanh(bx + a))^4$$

$$+ \frac{1}{105} (21 bx^5 \operatorname{arctanh}(\tanh(bx + a))^2 + (b^2 x^7 - 7 bx^6 \operatorname{arctanh}(\tanh(bx + a)))b) b$$

input `integrate(x^2*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`

output `-1/3*b*x^4*arctanh(tanh(b*x + a))^3 + 1/3*x^3*arctanh(tanh(b*x + a))^4 + 1/105*(21*b*x^5*arctanh(tanh(b*x + a))^2 + (b^2*x^7 - 7*b*x^6*arctanh(tanh(b*x + a))))*b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.85

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{7} b^4 x^7 + \frac{2}{3} ab^3 x^6 + \frac{6}{5} a^2 b^2 x^5 + a^3 b x^4 + \frac{1}{3} a^4 x^3$$

input `integrate(x^2*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/7*b^4*x^7 + 2/3*a*b^3*x^6 + 6/5*a^2*b^2*x^5 + a^3*b*x^4 + 1/3*a^4*x^3`

**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.32

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^7}{105} - \frac{b^3 x^6 \operatorname{atanh}(\tanh(a + bx))}{15} + \frac{b^2 x^5 \operatorname{atanh}(\tanh(a + bx))^2}{5} - \frac{b x^4 \operatorname{atanh}(\tanh(a + bx))^3}{3} + \frac{x^3 \operatorname{atanh}(\tanh(a + bx))^4}{3}$$

input `int(x^2*atanh(tanh(a + b*x))^4,x)`output `(x^3*atanh(tanh(a + b*x))^4)/3 + (b^4*x^7)/105 + (b^2*x^5*atanh(tanh(a + b*x))^2)/5 - (b*x^4*atanh(tanh(a + b*x))^3)/3 - (b^3*x^6*atanh(tanh(a + b*x)))`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x^2 dx$$

input `int(x^2*atanh(tanh(b*x+a))^4,x)`output `int(atanh(tanh(a + b*x))**4*x**2,x)`

### 3.71 $\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	593
Mathematica [B] (verified)	593
Rubi [A] (verified)	594
Maple [B] (verified)	595
Fricas [A] (verification not implemented)	596
Sympy [A] (verification not implemented)	596
Maxima [B] (verification not implemented)	597
Giac [A] (verification not implemented)	597
Mupad [B] (verification not implemented)	598
Reduce [F]	598

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^6}{30b^2}$$

output

```
1/5*x*arctanh(tanh(b*x+a))^5/b-1/30*arctanh(tanh(b*x+a))^6/b^2
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 125 vs. 2(34) = 68.

Time = 0.19 (sec) , antiderivative size = 125, normalized size of antiderivative = 3.68

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{(a + bx) ((5a - bx)(a + bx)^4 - 6(4a - bx)(a + bx)^3 \operatorname{arctanh}(\tanh(a + bx)) + 15(3a - bx)(a + bx)^2 \operatorname{arctanh}(\tanh(a + bx))^2 - \dots)}{\dots}$$

input

```
Integrate[x*ArcTanh[Tanh[a + b*x]]^4,x]
```

output

```
-1/30*((a + b*x)*((5*a - b*x)*(a + b*x)^4 - 6*(4*a - b*x)*(a + b*x)^3*ArcTanh[Tanh[a + b*x]] + 15*(3*a - b*x)*(a + b*x)^2*ArcTanh[Tanh[a + b*x]]^2 - 20*(2*a^2 + a*b*x - b^2*x^2)*ArcTanh[Tanh[a + b*x]]^3 + 15*(a - b*x)*ArcTanh[Tanh[a + b*x]]^4))/b^2
```

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$\downarrow 2599$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^5 dx}{5b}$$

$$\downarrow 2588$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^5 d \operatorname{arctanh}(\tanh(a + bx))}{5b^2}$$

$$\downarrow 15$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^5}{5b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^6}{30b^2}$$

input

```
Int[x*ArcTanh[Tanh[a + b*x]]^4,x]
```

output

```
(x*ArcTanh[Tanh[a + b*x]]^5)/(5*b) - ArcTanh[Tanh[a + b*x]]^6/(30*b^2)
```

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs.  $2(30) = 60$ .

Time = 27.62 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.09

method	result
parallelrisch	$\frac{b^4 x^6}{30} - \frac{b^3 \operatorname{arctanh}(\tanh(bx+a))x^5}{5} + \frac{b^2 \operatorname{arctanh}(\tanh(bx+a))^2 x^4}{2} - \frac{2b \operatorname{arctanh}(\tanh(bx+a))^3 x^3}{3} + \frac{x^2 \operatorname{arctanh}(\tanh(bx+a))}{2}$
default	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^4}{2} - 2b \left( \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^3}{3} - b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^2}{4} - \frac{b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))}{5} \right)}{2} \right) \right)$
parts	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^4}{2} - 2b \left( \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^3}{3} - b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^2}{4} - \frac{b \left( \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))}{5} \right)}{2} \right) \right)$
risch	Expression too large to display

input `int(x*arctanh(tanh(b*x+a))^4,x,method=_RETURNVERBOSE)`



output  $1/30*b^4*x^6-1/5*b^3*\operatorname{arctanh}(\tanh(b*x+a))*x^5+1/2*b^2*\operatorname{arctanh}(\tanh(b*x+a))^2*x^4-2/3*b*\operatorname{arctanh}(\tanh(b*x+a))^3*x^3+1/2*x^2*\operatorname{arctanh}(\tanh(b*x+a))^4$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{6} b^4 x^6 + \frac{4}{5} a b^3 x^5 + \frac{3}{2} a^2 b^2 x^4 + \frac{4}{3} a^3 b x^3 + \frac{1}{2} a^4 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`

output  $1/6*b^4*x^6 + 4/5*a*b^3*x^5 + 3/2*a^2*b^2*x^4 + 4/3*a^3*b*x^3 + 1/2*a^4*x^2$

### Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \begin{cases} \frac{x \operatorname{atanh}^5(\tanh(a+bx))}{5b} - \frac{\operatorname{atanh}^6(\tanh(a+bx))}{30b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^4(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**4,x)`

output `Piecewise((x*atanh(tanh(a + b*x))**5/(5*b) - atanh(tanh(a + b*x))**6/(30*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**4/2, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 72 vs.  $2(30) = 60$ .

Time = 0.21 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.12

$$\begin{aligned} \int x \operatorname{arctanh}(\tanh(a + bx))^4 dx \\ = -\frac{2}{3} bx^3 \operatorname{arctanh}(\tanh(bx + a))^3 + \frac{1}{2} x^2 \operatorname{arctanh}(\tanh(bx + a))^4 \\ + \frac{1}{30} (15 bx^4 \operatorname{arctanh}(\tanh(bx + a))^2 + (b^2 x^6 - 6 bx^5 \operatorname{arctanh}(\tanh(bx + a)))b) b \end{aligned}$$

input `integrate(x*arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`

output `-2/3*b*x^3*arctanh(tanh(b*x + a))^3 + 1/2*x^2*arctanh(tanh(b*x + a))^4 + 1/30*(15*b*x^4*arctanh(tanh(b*x + a))^2 + (b^2*x^6 - 6*b*x^5*arctanh(tanh(b*x + a)))*b)*b`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{6} b^4 x^6 + \frac{4}{5} a b^3 x^5 + \frac{3}{2} a^2 b^2 x^4 + \frac{4}{3} a^3 b x^3 + \frac{1}{2} a^4 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/6*b^4*x^6 + 4/5*a*b^3*x^5 + 3/2*a^2*b^2*x^4 + 4/3*a^3*b*x^3 + 1/2*a^4*x^2`

**Mupad [B] (verification not implemented)**

Time = 3.48 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.06

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{b^4 x^6}{30} - \frac{b^3 x^5 \operatorname{atanh}(\tanh(a + bx))}{5} + \frac{b^2 x^4 \operatorname{atanh}(\tanh(a + bx))^2}{2} - \frac{2 b x^3 \operatorname{atanh}(\tanh(a + bx))^3}{3} + \frac{x^2 \operatorname{atanh}(\tanh(a + bx))^4}{2}$$

input `int(x*atanh(tanh(a + b*x))^4,x)`output `(x^2*atanh(tanh(a + b*x))^4)/2 + (b^4*x^6)/30 + (b^2*x^4*atanh(tanh(a + b*x))^2)/2 - (2*b*x^3*atanh(tanh(a + b*x))^3)/3 - (b^3*x^5*atanh(tanh(a + b*x)))/5`**Reduce [F]**

$$\int x \operatorname{arctanh}(\tanh(a + bx))^4 dx = \int \operatorname{atanh}(\tanh(bx + a))^4 x dx$$

input `int(x*atanh(tanh(b*x+a))^4,x)`output `int(atanh(tanh(a + b*x))**4*x,x)`

## 3.72 $\int \operatorname{arctanh}(\tanh(a + bx))^4 dx$

Optimal result	599
Mathematica [A] (verified)	599
Rubi [A] (verified)	600
Maple [A] (verified)	601
Fricas [B] (verification not implemented)	601
Sympy [A] (verification not implemented)	602
Maxima [B] (verification not implemented)	602
Giac [B] (verification not implemented)	603
Mupad [B] (verification not implemented)	603
Reduce [B] (verification not implemented)	604

### Optimal result

Integrand size = 9, antiderivative size = 16

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{5b}$$

output `1/5*arctanh(tanh(b*x+a))^5/b`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{5b}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4,x]`

output `ArcTanh[Tanh[a + b*x]]^5/(5*b)`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^4 d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^5}{5b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4,x]`

output `ArcTanh[Tanh[a + b*x]]^5/(5*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 29.60 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\operatorname{arctanh}(\tanh(bx+a))^5}{5b}$
default	$\frac{\operatorname{arctanh}(\tanh(bx+a))^5}{5b}$
parallelrisch	$-b^3 \operatorname{arctanh}(\tanh(bx+a)) x^4 - 2x^2 b \operatorname{arctanh}(\tanh(bx+a))^3 + x \operatorname{arctanh}(\tanh(bx+a))$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4,x,method=_RETURNVERBOSE)`

output `1/5*arctanh(tanh(b*x+a))^5/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(14) = 28.

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{5} b^4 x^5 + ab^3 x^4 + 2a^2 b^2 x^3 + 2a^3 b x^2 + a^4 x$$

input `integrate(arctanh(tanh(b*x+a))^4,x, algorithm="fricas")`

output `1/5*b^4*x^5 + a*b^3*x^4 + 2*a^2*b^2*x^3 + 2*a^3*b*x^2 + a^4*x`

**Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \begin{cases} \frac{\operatorname{atanh}^5(\tanh(a+bx))}{5b} & \text{for } b \neq 0 \\ x \operatorname{atanh}^4(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**4,x)`

output `Piecewise((atanh(tanh(a + b*x))**5/(5*b), Ne(b, 0)), (x*atanh(tanh(a))**4, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. 2(14) = 28.

Time = 0.22 (sec) , antiderivative size = 69, normalized size of antiderivative = 4.31

$$\begin{aligned} & \int \operatorname{arctanh}(\tanh(a + bx))^4 dx \\ &= -2bx^2 \operatorname{artanh}(\tanh(bx + a))^3 + x \operatorname{artanh}(\tanh(bx + a))^4 \\ & \quad + \frac{1}{5} (10bx^3 \operatorname{artanh}(\tanh(bx + a))^2 + (b^2x^5 - 5bx^4 \operatorname{artanh}(\tanh(bx + a)))b) \end{aligned}$$

input `integrate(arctanh(tanh(b*x+a))^4,x, algorithm="maxima")`

output `-2*b*x^2*arctanh(tanh(b*x + a))^3 + x*arctanh(tanh(b*x + a))^4 + 1/5*(10*b*x^3*arctanh(tanh(b*x + a))^2 + (b^2*x^5 - 5*b*x^4*arctanh(tanh(b*x + a)))*b)*b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 42 vs.  $2(14) = 28$ .

Time = 0.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{1}{5} b^4 x^5 + ab^3 x^4 + 2a^2 b^2 x^3 + 2a^3 b x^2 + a^4 x$$

input `integrate(arctanh(tanh(b*x+a))^4,x, algorithm="giac")`

output `1/5*b^4*x^5 + a*b^3*x^4 + 2*a^2*b^2*x^3 + 2*a^3*b*x^2 + a^4*x`

**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 67, normalized size of antiderivative = 4.19

$$\begin{aligned} \int \operatorname{arctanh}(\tanh(a + bx))^4 dx &= \frac{b^4 x^5}{5} - b^3 x^4 \operatorname{atanh}(\tanh(a + bx)) \\ &\quad + 2b^2 x^3 \operatorname{atanh}(\tanh(a + bx))^2 \\ &\quad - 2b x^2 \operatorname{atanh}(\tanh(a + bx))^3 \\ &\quad + x \operatorname{atanh}(\tanh(a + bx))^4 \end{aligned}$$

input `int(atanh(tanh(a + b*x))^4,x)`

output `x*atanh(tanh(a + b*x))^4 + (b^4*x^5)/5 + 2*b^2*x^3*atanh(tanh(a + b*x))^2 - 2*b*x^2*atanh(tanh(a + b*x))^3 - b^3*x^4*atanh(tanh(a + b*x))`



**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\tanh(a + bx))^4 dx = \frac{\operatorname{atanh}(\tanh(bx + a))^5}{5b}$$

input

```
int(atanh(tanh(b*x+a))^4,x)
```

output

```
atanh(tanh(a + b*x))**5/(5*b)
```

### 3.73 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx$

Optimal result . . . . .	605
Mathematica [A] (verified) . . . . .	606
Rubi [A] (verified) . . . . .	606
Maple [B] (verified) . . . . .	608
Fricas [A] (verification not implemented) . . . . .	609
Sympy [F] . . . . .	609
Maxima [A] (verification not implemented) . . . . .	609
Giac [A] (verification not implemented) . . . . .	610
Mupad [B] (verification not implemented) . . . . .	610
Reduce [F] . . . . .	611

#### Optimal result

Integrand size = 13, antiderivative size = 105

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} dx = -bx(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 + \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{1}{3}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^3 + \frac{1}{4} \operatorname{arctanh}(\tanh(a + bx))^4 + (bx - \operatorname{arctanh}(\tanh(a + bx)))^4 \log(x)$$

output

```
-b*x*(b*x-arctanh(tanh(b*x+a)))^3+1/2*(b*x-arctanh(tanh(b*x+a)))^2*arctanh
(tanh(b*x+a))^2-1/3*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^3+1/4*
arctanh(tanh(b*x+a))^4+(b*x-arctanh(tanh(b*x+a)))^4*ln(x)
```

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.67

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx = \frac{1}{4}(a+bx)^4 + \frac{1}{2}(a+bx)^2(a^2 - 4a(a+bx - \operatorname{arctanh}(\tanh(a+bx)))) + 6(a+bx - \operatorname{arctanh}(\tanh(a+bx)))^2 + (a+bx)(a^3 - 4a^2(a+bx - \operatorname{arctanh}(\tanh(a+bx)))) + 6a(a+bx - \operatorname{arctanh}(\tanh(a+bx)))^2 - 4(a+bx - \operatorname{arctanh}(\tanh(a+bx)))^3 - \frac{1}{3}(a+bx)^3(3a + 4bx - 4\operatorname{arctanh}(\tanh(a+bx))) + (-bx + \operatorname{arctanh}(\tanh(a+bx)))^4 \log(bx)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x,x]`

output `(a + b*x)^4/4 + ((a + b*x)^2*(a^2 - 4*a*(a + b*x - ArcTanh[Tanh[a + b*x]])) + 6*(a + b*x - ArcTanh[Tanh[a + b*x]])^2)/2 + (a + b*x)*(a^3 - 4*a^2*(a + b*x - ArcTanh[Tanh[a + b*x]])) + 6*a*(a + b*x - ArcTanh[Tanh[a + b*x]])^2 - 4*(a + b*x - ArcTanh[Tanh[a + b*x]])^3 - ((a + b*x)^3*(3*a + 4*b*x - 4*ArcTanh[Tanh[a + b*x]]))/3 + (-b*x + ArcTanh[Tanh[a + b*x]])^4*Log[b*x]`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2590, 2590, 2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx$$

↓ 2590

$$\begin{aligned}
& \frac{1}{4} \operatorname{arctanh}(\tanh(a+bx))^4 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x} dx \\
& \quad \downarrow \text{2590} \\
& \frac{1}{4} \operatorname{arctanh}(\tanh(a+bx))^4 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a+bx))^3 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x} dx \right) \\
& \quad \downarrow \text{2590} \\
& \frac{1}{4} \operatorname{arctanh}(\tanh(a+bx))^4 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a+bx))^3 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a+bx))^2 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x} dx \right) \right) \\
& \quad \downarrow \text{2589} \\
& \frac{1}{4} \operatorname{arctanh}(\tanh(a+bx))^4 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a+bx))^3 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a+bx))^2 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \log|x| \right) \right) \\
& \quad \downarrow \text{14} \\
& \frac{1}{4} \operatorname{arctanh}(\tanh(a+bx))^4 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a+bx))^3 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a+bx))^2 - (bx - \operatorname{arctanh}(\tanh(a+bx))) \log|x| \right) \right)
\end{aligned}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^4/x, x]
```

output

```
ArcTanh[Tanh[a + b*x]]^4/4 - (b*x - ArcTanh[Tanh[a + b*x]])*(ArcTanh[Tanh[a + b*x]]^3/3 - (b*x - ArcTanh[Tanh[a + b*x]])*(ArcTanh[Tanh[a + b*x]]^2/2 - (b*x - ArcTanh[Tanh[a + b*x]])*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x])))
```

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

## Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 264 vs.  $2(99) = 198$ .

Time = 0.31 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.52

method	result
default	$\ln(x) \operatorname{arctanh}(\tanh(bx+a))^4 - 4b \left( b^3 \left( \frac{x^4 \ln(x)}{4} - \frac{x^4}{16} \right) + 3ab^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 3b^2 \operatorname{arctanh}(\tanh(bx+a)) \right)$
parts	$\ln(x) \operatorname{arctanh}(\tanh(bx+a))^4 - 4b \left( b^3 \left( \frac{x^4 \ln(x)}{4} - \frac{x^4}{16} \right) + 3ab^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 3b^2 \operatorname{arctanh}(\tanh(bx+a)) \right)$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x,x,method=_RETURNVERBOSE)`

output `ln(x)*arctanh(tanh(b*x+a))^4-4*b*(b^3*(1/4*x^4*ln(x)-1/16*x^4)+3*a*b^2*(1/3*x^3*ln(x)-1/9*x^3)+3*b^2*(arctanh(tanh(b*x+a))-b*x-a)*(1/3*x^3*ln(x)-1/9*x^3)+3*a^2*b*(1/2*x^2*ln(x)-1/4*x^2)+6*a*b*(arctanh(tanh(b*x+a))-b*x-a)*(1/2*x^2*ln(x)-1/4*x^2)+3*b*(arctanh(tanh(b*x+a))-b*x-a)^2*(1/2*x^2*ln(x)-1/4*x^2)+a^3*(x*ln(x)-x)+3*a^2*(arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x)+3*a*(arctanh(tanh(b*x+a))-b*x-a)^2*(x*ln(x)-x)+(arctanh(tanh(b*x+a))-b*x-a)^3*(x*ln(x)-x)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.40

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx = \frac{1}{4} b^4 x^4 + \frac{4}{3} ab^3 x^3 + 3a^2 b^2 x^2 + 4a^3 bx + a^4 \log(x)$$

input `integrate(arctanh(tanh(b*x+a))^4/x,x, algorithm="fricas")`output `1/4*b^4*x^4 + 4/3*a*b^3*x^3 + 3*a^2*b^2*x^2 + 4*a^3*b*x + a^4*log(x)`**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx = \int \frac{\operatorname{atanh}^4(\tanh(a+bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))**4/x,x)`output `Integral(atanh(tanh(a + b*x))**4/x, x)`**Maxima [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.40

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} dx = \frac{1}{4} b^4 x^4 + \frac{4}{3} ab^3 x^3 + 3a^2 b^2 x^2 + 4a^3 bx + a^4 \log(x)$$

input `integrate(arctanh(tanh(b*x+a))^4/x,x, algorithm="maxima")`output `1/4*b^4*x^4 + 4/3*a*b^3*x^3 + 3*a^2*b^2*x^2 + 4*a^3*b*x + a^4*log(x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} dx = \frac{1}{4} b^4 x^4 + \frac{4}{3} a b^3 x^3 + 3 a^2 b^2 x^2 + 4 a^3 b x + a^4 \log(|x|)$$

input `integrate(arctanh(tanh(b*x+a))^4/x,x, algorithm="giac")`

output `1/4*b^4*x^4 + 4/3*a*b^3*x^3 + 3*a^2*b^2*x^2 + 4*a^3*b*x + a^4*log(abs(x))`

**Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 423, normalized size of antiderivative = 4.03

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} dx \\ &= \ln(x) \left( \frac{\left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^4}{16} \right. \\ & \quad + \frac{3a^2 \left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2}{2} + a^4 \\ & \quad \left. - \frac{a \left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^3}{2} \right. \\ & \quad \left. - 2a^3 \left( 2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx \right) \right) \\ & + \frac{b^4 x^4}{4} - \frac{2b^3 x^3 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)}{3} \\ & + \frac{3b^2 x^2 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2}{4} \\ & - \frac{bx \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^3}{2} \end{aligned}$$

input `int(atanh(tanh(a + b*x))^4/x,x)`

output `log(x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4/16 + (3*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 + a^4 - (a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/2 - 2*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (b^4*x^4)/4 - (2*b^3*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/3 + (3*b^2*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/4 - (b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/2`

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^4}{x} dx$$

input `int(atanh(tanh(b*x+a))^4/x,x)`

output `int(atanh(tanh(a + b*x))**4/x,x)`



### 3.74 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^2} dx$

Optimal result	612
Mathematica [A] (verified)	612
Rubi [A] (verified)	613
Maple [A] (verified)	615
Fricas [A] (verification not implemented)	615
Sympy [F]	616
Maxima [A] (verification not implemented)	616
Giac [A] (verification not implemented)	616
Mupad [B] (verification not implemented)	617
Reduce [F]	617

#### Optimal result

Integrand size = 13, antiderivative size = 95

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^2} dx = 4b^2x(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 - 2b(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^2 + \frac{4}{3}b\operatorname{arctanh}(\tanh(a+bx))^3 - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} - 4b(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log(x)$$

output

```
4*b^2*x*(b*x-arctanh(tanh(b*x+a)))^2-2*b*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^2+4/3*b*arctanh(tanh(b*x+a))^3-arctanh(tanh(b*x+a))^4/x-4*b*(b*x-arctanh(tanh(b*x+a)))^3*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^2} dx = -\frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x} + \frac{2}{3}b^4x^3(5 - 6\log(x)) - 12b^2x\operatorname{arctanh}(\tanh(a+bx))^2 \log(x) + 4b\operatorname{arctanh}(\tanh(a+bx))^3(1 + \log(x)) + 6b^3x^2\operatorname{arctanh}(\tanh(a+bx))(-1 + 2\log(x))$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^2, x]`

output `-(ArcTanh[Tanh[a + b*x]]^4/x) + (2*b^4*x^3*(5 - 6*Log[x]))/3 - 12*b^2*x*ArcTanh[Tanh[a + b*x]]^2*Log[x] + 4*b*ArcTanh[Tanh[a + b*x]]^3*(1 + Log[x]) + 6*b^3*x^2*ArcTanh[Tanh[a + b*x]]*(-1 + 2*Log[x])`

### Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2590, 2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^2} dx$$

$$\downarrow 2599$$

$$4b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x}$$

$$\downarrow 2590$$

$$4b \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a + bx))^3 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x}$$

$$\downarrow 2590$$

$$4b \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a + bx))^3 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx \right) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x}$$

$$\downarrow 2589$$

$$4b \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a + bx))^3 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} \right)$$

↓ 14

$$4b \left( \frac{1}{3} \operatorname{arctanh}(\tanh(a + bx))^3 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^2,x]`

output `-(ArcTanh[Tanh[a + b*x]]^4/x) + 4*b*(ArcTanh[Tanh[a + b*x]]^3/3 - (b*x - ArcTanh[Tanh[a + b*x]])*(ArcTanh[Tanh[a + b*x]]^2/2 - (b*x - ArcTanh[Tanh[a + b*x]])*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x])))`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.74

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{x} + 4b \left( \ln(x) \operatorname{arctanh}(\tanh(bx+a))^3 - 3b \left( b^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 2ab \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{2} \right) \right) \right)$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{x} + 4b \left( \ln(x) \operatorname{arctanh}(\tanh(bx+a))^3 - 3b \left( b^2 \left( \frac{x^3 \ln(x)}{3} - \frac{x^3}{9} \right) + 2ab \left( \frac{x^2 \ln(x)}{2} - \frac{x^2}{2} \right) \right) \right)$
risch	Expression too large to display

input

```
int(arctanh(tanh(b*x+a))^4/x^2,x,method=_RETURNVERBOSE)
```

output

```
-arctanh(tanh(b*x+a))^4/x+4*b*(ln(x)*arctanh(tanh(b*x+a))^3-3*b*(b^2*(1/3*x^3*ln(x)-1/9*x^3)+2*a*b*(1/2*x^2*ln(x)-1/4*x^2)+2*b*(arctanh(tanh(b*x+a))-b*x-a)*(1/2*x^2*ln(x)-1/4*x^2)+a^2*(x*ln(x)-x)+2*a*(arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x)+(arctanh(tanh(b*x+a))-b*x-a)^2*(x*ln(x)-x)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^2} dx = \frac{b^4 x^4 + 6 a b^3 x^3 + 18 a^2 b^2 x^2 + 12 a^3 b x \log(x) - 3 a^4}{3 x}$$

input

```
integrate(arctanh(tanh(b*x+a))^4/x^2,x, algorithm="fricas")
```

output

```
1/3*(b^4*x^4 + 6*a*b^3*x^3 + 18*a^2*b^2*x^2 + 12*a^3*b*x*log(x) - 3*a^4)/x
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^2} dx = \int \frac{\operatorname{atanh}^4(\tanh(a + bx))}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**4/x**2,x)`

output `Integral(atanh(tanh(a + b*x))**4/x**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^2} dx \\ &= 4b \operatorname{artanh}(\tanh(bx + a))^3 \log(x) - \frac{\operatorname{artanh}(\tanh(bx + a))^4}{x} \\ & \quad + \frac{2}{3} (2b^3x^3 + 9ab^2x^2 + 18a^2bx + 6a^3 \log(x) - 6 \operatorname{artanh}(\tanh(bx + a))^3 \log(x))b \end{aligned}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^2,x, algorithm="maxima")`

output `4*b*arctanh(tanh(b*x + a))^3*log(x) - arctanh(tanh(b*x + a))^4/x + 2/3*(2*b^3*x^3 + 9*a*b^2*x^2 + 18*a^2*b*x + 6*a^3*log(x) - 6*arctanh(tanh(b*x + a))^3*log(x))*b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.46

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^2} dx = \frac{1}{3} b^4 x^3 + 2ab^3x^2 + 6a^2b^2x + 4a^3b \log(|x|) - \frac{a^4}{x}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^2,x, algorithm="giac")`

output  $1/3*b^4*x^3 + 2*a*b^3*x^2 + 6*a^2*b^2*x + 4*a^3*b*\log(\text{abs}(x)) - a^4/x$

### Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 553, normalized size of antiderivative = 5.82

$$\int \frac{\arctanh(\tanh(a + bx))^4}{x^2} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^4/x^2,x)`

output  $\log(x)*(4*a^3*b - (b*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3)/2 + 3*a*b*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 - 6*a^2*b*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) - ((2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^4 + 24*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + 16*a^4 - 8*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - 32*a^3*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)/(16*x) + (b^4*x^3)/3 + (3*b^2*x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2)/2 - b^3*x^2*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)$

### Reduce [F]

$$\int \frac{\arctanh(\tanh(a + bx))^4}{x^2} dx = \int \frac{\text{atanh}(\tanh(bx + a))^4}{x^2} dx$$

input `int(atanh(tanh(b*x+a))^4/x^2,x)`

output `int(atanh(tanh(a + b*x))**4/x**2,x)`

### 3.75 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^3} dx$

Optimal result . . . . .	618
Mathematica [A] (verified) . . . . .	618
Rubi [A] (verified) . . . . .	619
Maple [A] (verified) . . . . .	621
Fricas [A] (verification not implemented) . . . . .	621
Sympy [F] . . . . .	622
Maxima [A] (verification not implemented) . . . . .	622
Giac [A] (verification not implemented) . . . . .	622
Mupad [B] (verification not implemented) . . . . .	623
Reduce [F] . . . . .	624

#### Optimal result

Integrand size = 13, antiderivative size = 87

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = -6b^3x(bx - \operatorname{arctanh}(\tanh(a + bx))) + 3b^2\operatorname{arctanh}(\tanh(a + bx))^2 - \frac{2b\operatorname{arctanh}(\tanh(a + bx))^3}{x} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2} + 6b^2(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \log(x)$$

output

```
-6*b^3*x*(b*x-arctanh(tanh(b*x+a)))+3*b^2*arctanh(tanh(b*x+a))^2-2*b*arctanh(tanh(b*x+a))^3/x-1/2*arctanh(tanh(b*x+a))^4/x^2+6*b^2*(b*x-arctanh(tanh(b*x+a)))^2*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = -\frac{2b\operatorname{arctanh}(\tanh(a + bx))^3}{x} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2} + 6b^4x^2 \log(x) - 6b^3x\operatorname{arctanh}(\tanh(a + bx))(1 + 2 \log(x)) + 3b^2\operatorname{arctanh}(\tanh(a + bx))^2(3 + 2 \log(x))$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^3, x]`

output  $(-2*b*ArcTanh[Tanh[a + b*x]]^3)/x - ArcTanh[Tanh[a + b*x]]^4/(2*x^2) + 6*b^4*x^2*Log[x] - 6*b^3*x*ArcTanh[Tanh[a + b*x]]*(1 + 2*Log[x]) + 3*b^2*ArcTanh[Tanh[a + b*x]]^2*(3 + 2*Log[x])$

### Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2590, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx$$

$$\downarrow 2599$$

$$2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2}$$

$$\downarrow 2599$$

$$2b \left( 3b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2}$$

$$\downarrow 2590$$

$$2b \left( 3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x}$$

$$\downarrow 2589$$

$$2b \left( 3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x}$$



↓ 14

$$2b \left( 3b \left( \frac{1}{2} \operatorname{arctanh}(\tanh(a + bx))^2 - (bx - \operatorname{arctanh}(\tanh(a + bx)))(bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx)))) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{2x^2} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^3,x]`

output `-1/2*ArcTanh[Tanh[a + b*x]]^4/x^2 + 2*b*(-(ArcTanh[Tanh[a + b*x]]^3/x) + 3*b*(ArcTanh[Tanh[a + b*x]]^2/2 - (b*x - ArcTanh[Tanh[a + b*x]])*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x])))`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.18

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{2x^2} + 2b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{x} + 3b\left(\ln(x) \operatorname{arctanh}(\tanh(bx+a))^2 - 2b\left(b\left(\frac{x^2 \ln(x)}{2}\right.\right.\right.\right.$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{2x^2} + 2b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{x} + 3b\left(\ln(x) \operatorname{arctanh}(\tanh(bx+a))^2 - 2b\left(b\left(\frac{x^2 \ln(x)}{2}\right.\right.\right.\right.$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arctanh(tanh(b*x+a))^4/x^2+2*b*(-arctanh(tanh(b*x+a))^3/x+3*b*(ln(x)*arctanh(tanh(b*x+a))^2-2*b*(b*(1/2*x^2*ln(x)-1/4*x^2)+a*(x*ln(x)-x)+(arctanh(tanh(b*x+a))-b*x-a)*(x*ln(x)-x))))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.54

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^3} dx = \frac{b^4 x^4 + 8ab^3 x^3 + 12a^2 b^2 x^2 \log(x) - 8a^3 bx - a^4}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^3,x, algorithm="fricas")`

output `1/2*(b^4*x^4 + 8*a*b^3*x^3 + 12*a^2*b^2*x^2*log(x) - 8*a^3*b*x - a^4)/x^2`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = \int \frac{\operatorname{atanh}^4(\tanh(a + bx))}{x^3} dx$$

input `integrate(atanh(tanh(b*x+a))**4/x**3,x)`

output `Integral(atanh(tanh(a + b*x))**4/x**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = -\frac{2b \operatorname{arctanh}(\tanh(bx + a))^3}{x} + 3(2b \operatorname{arctanh}(\tanh(bx + a))^2 \log(x) + (b^2x^2 + 4abx + 2a^2 \log(x) - 2 \operatorname{arctanh}(\tanh(bx + a))^2 \log(x)) \operatorname{arctanh}(\tanh(bx + a)) - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^3,x, algorithm="maxima")`

output `-2*b*arctanh(tanh(b*x + a))^3/x + 3*(2*b*arctanh(tanh(b*x + a))^2*log(x) + (b^2*x^2 + 4*a*b*x + 2*a^2*log(x) - 2*arctanh(tanh(b*x + a))^2*log(x))*b *b - 1/2*arctanh(tanh(b*x + a))^4/x^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = \frac{1}{2} b^4 x^2 + 4ab^3x + 6a^2b^2 \log(|x|) - \frac{8a^3bx + a^4}{2x^2}$$

input `integrate(arctanh(tanh(b*x+a))**4/x^3,x, algorithm="giac")`

output  $\frac{1}{2}b^4x^2 + 4ab^3x + 6a^2b^2\log(\text{abs}(x)) - \frac{1}{2}(8a^3bx + a^4)/x^2$

### Mupad [B] (verification not implemented)

Time = 3.85 (sec) , antiderivative size = 672, normalized size of antiderivative = 7.72

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^4/x^3,x)`

output  $(9b^2\log(1/(\exp(2a)\exp(2bx) + 1))^2)/4 - \log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1))^4/(32x^2) - \log(1/(\exp(2a)\exp(2bx) + 1))^4/(32x^2) + (9b^2\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1))^2)/4 - 3b^3x\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + (b\log(1/(\exp(2a)\exp(2bx) + 1))^3)/(4x) + (\log(1/(\exp(2a)\exp(2bx) + 1)))\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1))^3/(8x^2) + (\log(1/(\exp(2a)\exp(2bx) + 1)))^3\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/(8x^2) - (b\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))^3/(4x) + (3b^2\log(1/(\exp(2a)\exp(2bx) + 1))^2\log(x))/2 - (3\log(1/(\exp(2a)\exp(2bx) + 1)))^2\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1))^2)/(16x^2) + (3b^2\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))^2\log(x))/2 + 6b^4x^2\log(x) - (9b^2\log(1/(\exp(2a)\exp(2bx) + 1)))\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/2 + 3b^3x\log(1/(\exp(2a)\exp(2bx) + 1)) + (3b\log(1/(\exp(2a)\exp(2bx) + 1)))\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1))^2)/(4x) - (3b\log(1/(\exp(2a)\exp(2bx) + 1)))^2\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/(4x) - 3b^2\log(1/(\exp(2a)\exp(2bx) + 1))\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))\log(x) + 6b^3x\log(1/(\exp(2a)\exp(2bx) + 1)))\log(x) - 6b^3x\log((\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))\log(x)$

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^3} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^4}{x^3} dx$$

input `int(atanh(tanh(b*x+a))^4/x^3,x)`

output `int(atanh(tanh(a + b*x))**4/x**3,x)`

### 3.76 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^4} dx$

Optimal result	625
Mathematica [A] (verified)	625
Rubi [A] (verified)	626
Maple [A] (verified)	628
Fricas [A] (verification not implemented)	628
Sympy [F]	629
Maxima [A] (verification not implemented)	629
Giac [A] (verification not implemented)	630
Mupad [B] (verification not implemented)	630
Reduce [F]	631

#### Optimal result

Integrand size = 13, antiderivative size = 77

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^4} dx = 4b^4x - \frac{2b^2\operatorname{arctanh}(\tanh(a+bx))^2}{x} - \frac{2b\operatorname{arctanh}(\tanh(a+bx))^3}{3x^2} - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{3x^3} - 4b^3(bx - \operatorname{arctanh}(\tanh(a+bx)))\log(x)$$

output

$$4*b^4*x-2*b^2*\operatorname{arctanh}(\tanh(b*x+a))^2/x-2/3*b*\operatorname{arctanh}(\tanh(b*x+a))^3/x^2-1/3*\operatorname{arctanh}(\tanh(b*x+a))^4/x^3-4*b^3*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))*\ln(x)$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^4} dx = \frac{6b^2x^2\operatorname{arctanh}(\tanh(a+bx))^2 + 2bx\operatorname{arctanh}(\tanh(a+bx))^3 + \operatorname{arctanh}(\tanh(a+bx))^4 + 2b^4x^4(5+6\operatorname{arctanh}(\tanh(a+bx)))}{3x^3}$$

input

$$\operatorname{Integrate}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^4/x^4, x]$$

output

$$\frac{-1/3*(6*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 2*b*x*ArcTanh[Tanh[a + b*x]]^3 + ArcTanh[Tanh[a + b*x]]^4 + 2*b^4*x^4*(5 + 6*Log[x]) - 2*b^3*x^3*ArcTanh[Tanh[a + b*x]]*(11 + 6*Log[x]))}{x^3}$$
**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx$$

$$\downarrow 2599$$

$$\frac{4}{3}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{3x^3}$$

$$\downarrow 2599$$

$$\frac{4}{3}b \left( \frac{3}{2}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{3x^3}$$

$$\downarrow 2599$$

$$\frac{4}{3}b \left( \frac{3}{2}b \left( 2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{3x^3}$$

$$\downarrow 2589$$

$$\frac{4}{3}b \left( \frac{3}{2}b \left( 2b \left( bx - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{3x^3}$$

$$\downarrow 14$$

$$\frac{4}{3}b \left( \frac{3}{2}b \left( 2b(bx - \log(x)(bx - \operatorname{arctanh}(\tanh(a + bx)))) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{2x^2} \right) \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{3x^3}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^4,x]`

output `-1/3*ArcTanh[Tanh[a + b*x]]^4/x^3 + (4*b*(-1/2*ArcTanh[Tanh[a + b*x]]^3/x^2 + (3*b*(-(ArcTanh[Tanh[a + b*x]]^2/x) + 2*b*(b*x - (b*x - ArcTanh[Tanh[a + b*x]])*Log[x])))/2))/3`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`



**Maple [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{3x^3} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{2x^2} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x) \operatorname{arctanh}(\tanh(bx+a)) - b(x \ln(x) - x)) \right)}{2} \right)}{3}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{3x^3} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{2x^2} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{x} + 2b(\ln(x) \operatorname{arctanh}(\tanh(bx+a)) - b(x \ln(x) - x)) \right)}{2} \right)}{3}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*arctanh(tanh(b*x+a))^4/x^3+4/3*b*(-1/2*arctanh(tanh(b*x+a))^3/x^2+3/2*b*(-arctanh(tanh(b*x+a))^2/x+2*b*(ln(x)*arctanh(tanh(b*x+a))-b*(x*ln(x)-x))))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx = \frac{3b^4x^4 + 12ab^3x^3 \log(x) - 18a^2b^2x^2 - 6a^3bx - a^4}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^4,x, algorithm="fricas")`

output `1/3*(3*b^4*x^4 + 12*a*b^3*x^3*log(x) - 18*a^2*b^2*x^2 - 6*a^3*b*x - a^4)/x^3`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx = \int \frac{\operatorname{atanh}^4(\tanh(a + bx))}{x^4} dx$$

input `integrate(atanh(tanh(b*x+a))**4/x**4, x)`

output `Integral(atanh(tanh(a + b*x))**4/x**4, x)`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx$$

$$= 2 \left( 2 \left( b \operatorname{artanh}(\tanh(bx + a)) \log(x) - \left( b \left( x + \frac{a}{b} \right) \log(x) - b \left( x + \frac{a \log(x)}{b} \right) \right) b \right) b - \frac{b \operatorname{artanh}(\tanh(bx + a))}{x} \right. \\ \left. - \frac{2 b \operatorname{artanh}(\tanh(bx + a))^3}{3 x^2} - \frac{\operatorname{artanh}(\tanh(bx + a))^4}{3 x^3} \right)$$

input `integrate(arctanh(tanh(b*x+a))^4/x^4,x, algorithm="maxima")`

output `2*(2*(b*arctanh(tanh(b*x + a))*log(x) - (b*(x + a/b)*log(x) - b*(x + a*log(x)/b))*b)*b - b*arctanh(tanh(b*x + a))^2/x*b - 2/3*b*arctanh(tanh(b*x + a))^3/x^2 - 1/3*arctanh(tanh(b*x + a))^4/x^3`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx = b^4 x + 4ab^3 \log(|x|) - \frac{18a^2 b^2 x^2 + 6a^3 bx + a^4}{3x^3}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^4,x, algorithm="giac")`

output `b^4*x + 4*a*b^3*log(abs(x)) - 1/3*(18*a^2*b^2*x^2 + 6*a^3*b*x + a^4)/x^3`

**Mupad [B] (verification not implemented)**

Time = 4.11 (sec) , antiderivative size = 571, normalized size of antiderivative = 7.42

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^4/x^4,x)`

output `(11*b^3*log(exp(2*b*x)/(exp(2*a)*exp(2*b*x) + 1)))/3 - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^4/(48*x^3) - (11*b^3*log(1/(exp(2*a)*exp(2*b*x) + 1)))/3 - (10*b^4*x)/3 - log(1/(exp(2*a)*exp(2*b*x) + 1))^4/(48*x^3) + (b*log(1/(exp(2*a)*exp(2*b*x) + 1))^3)/(12*x^2) - 2*b^3*log(1/(exp(2*a)*exp(2*b*x) + 1))*log(x) + (log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3)/(12*x^3) + (log(1/(exp(2*a)*exp(2*b*x) + 1))^3*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(12*x^3) - (b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3)/(12*x^2) + 2*b^3*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(x) - 4*b^4*x*log(x) - (b^2*log(1/(exp(2*a)*exp(2*b*x) + 1))^2)/(2*x) - (log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/(8*x^3) - (b^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/(2*x) + (b^2*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/x + (b*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2)/(4*x^2) - (b*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(4*x^2)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^4} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^4}{x^4} dx$$

input `int(atanh(tanh(b*x+a))^4/x^4,x)`

output `int(atanh(tanh(a + b*x))**4/x**4,x)`

### 3.77 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^5} dx$

Optimal result . . . . .	632
Mathematica [A] (verified) . . . . .	632
Rubi [A] (verified) . . . . .	633
Maple [A] (verified) . . . . .	634
Fricas [A] (verification not implemented) . . . . .	635
Sympy [A] (verification not implemented) . . . . .	635
Maxima [A] (verification not implemented) . . . . .	636
Giac [A] (verification not implemented) . . . . .	636
Mupad [B] (verification not implemented) . . . . .	637
Reduce [B] (verification not implemented) . . . . .	637

#### Optimal result

Integrand size = 13, antiderivative size = 74

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^5} dx = -\frac{b^3 \operatorname{arctanh}(\tanh(a+bx))}{x} - \frac{b^2 \operatorname{arctanh}(\tanh(a+bx))^2}{2x^2} - \frac{b \operatorname{arctanh}(\tanh(a+bx))^3}{3x^3} - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{4x^4} + b^4 \log(x)$$

output

```
-b^3*arctanh(tanh(b*x+a))/x-1/2*b^2*arctanh(tanh(b*x+a))^2/x^2-1/3*b*arctanh(tanh(b*x+a))^3/x^3-1/4*arctanh(tanh(b*x+a))^4/x^4+b^4*ln(x)
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^5} dx = \frac{12b^3x^3\operatorname{arctanh}(\tanh(a+bx)) + 6b^2x^2\operatorname{arctanh}(\tanh(a+bx))^2 + 4bx\operatorname{arctanh}(\tanh(a+bx))^3 + 3\operatorname{arctanh}(\tanh(a+bx))^4}{12x^4}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^4/x^5,x]
```

output

$$-1/12*(12*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 6*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 4*b*x*ArcTanh[Tanh[a + b*x]]^3 + 3*ArcTanh[Tanh[a + b*x]]^4 - b^4*x^4*(25 + 12*Log[x]))/x^4$$
**Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx$$

$$\downarrow 2599$$

$$b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^4} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4}$$

$$\downarrow 2599$$

$$b \left( b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4}$$

$$\downarrow 2599$$

$$b \left( b \left( b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4}$$

$$\downarrow 2599$$

$$b \left( b \left( b \left( b \int \frac{1}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4}$$

$$\downarrow 14$$

$$b \left( b \left( b \left( b \log(x) - \frac{\operatorname{arctanh}(\tanh(a + bx))}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{4x^4}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^5,x]`

output `-1/4*ArcTanh[Tanh[a + b*x]]^4/x^4 + b*(-1/3*ArcTanh[Tanh[a + b*x]]^3/x^3 + b*(-1/2*ArcTanh[Tanh[a + b*x]]^2/x^2 + b*(-(ArcTanh[Tanh[a + b*x]]/x) + b*Log[x])))`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 2.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.93

method	result
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{4x^4} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{3x^3} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2x^2} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} \right) \right) \right)$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{4x^4} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{3x^3} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{2x^2} + b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{x} \right) \right) \right)$
paralelrisch	$\frac{12b^4x^4 \ln(x) - 12b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 - 6b^2x^2 \operatorname{arctanh}(\tanh(bx+a))^2 - 4x \operatorname{arctanh}(\tanh(bx+a))^3 b - 3 \operatorname{arctanh}(\tanh(bx+a))}{12x^4}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*arctanh(tanh(b*x+a))^4/x^4+b*(-1/3*arctanh(tanh(b*x+a))^3/x^3+b*(-1/2*arctanh(tanh(b*x+a))^2/x^2+b*(-arctanh(tanh(b*x+a))/x+b*ln(x))))`

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.65

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx = \frac{12b^4x^4 \log(x) - 48ab^3x^3 - 36a^2b^2x^2 - 16a^3bx - 3a^4}{12x^4}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^5,x, algorithm="fricas")`

output `1/12*(12*b^4*x^4*log(x) - 48*a*b^3*x^3 - 36*a^2*b^2*x^2 - 16*a^3*b*x - 3*a^4)/x^4`

### Sympy [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx = b^4 \log(x) - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{x} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{2x^2} - \frac{b \operatorname{atanh}^3(\tanh(a + bx))}{3x^3} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{4x^4}$$

input `integrate(atanh(tanh(b*x+a))**4/x**5,x)`

output `b**4*log(x) - b**3*atanh(tanh(a + b*x))/x - b**2*atanh(tanh(a + b*x))**2/(2*x**2) - b*atanh(tanh(a + b*x))**3/(3*x**3) - atanh(tanh(a + b*x))**4/(4*x**4)`



**Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx$$

$$= \frac{1}{2} \left( 2 \left( b^2 \log(x) - \frac{b \operatorname{arctanh}(\tanh(bx + a))}{x} \right) b - \frac{b \operatorname{arctanh}(\tanh(bx + a))^2}{x^2} \right) b$$

$$- \frac{b \operatorname{arctanh}(\tanh(bx + a))^3}{3x^3} - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{4x^4}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^5,x, algorithm="maxima")`

output `1/2*(2*(b^2*log(x) - b*arctanh(tanh(b*x + a))/x)*b - b*arctanh(tanh(b*x + a))^2/x^2)*b - 1/3*b*arctanh(tanh(b*x + a))^3/x^3 - 1/4*arctanh(tanh(b*x + a))^4/x^4`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx = b^4 \log(|x|) - \frac{48ab^3x^3 + 36a^2b^2x^2 + 16a^3bx + 3a^4}{12x^4}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^5,x, algorithm="giac")`

output `b^4*log(abs(x)) - 1/12*(48*a*b^3*x^3 + 36*a^2*b^2*x^2 + 16*a^3*b*x + 3*a^4)/x^4`

**Mupad [B] (verification not implemented)**

Time = 3.81 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx = b^4 \ln(x) - \frac{\operatorname{atanh}(\tanh(a + bx))^4}{4x^4} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))^2}{2x^2} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{x} - \frac{b \operatorname{atanh}(\tanh(a + bx))^3}{3x^3}$$

input `int(atanh(tanh(a + b*x))^4/x^5,x)`output `b^4*log(x) - atanh(tanh(a + b*x))^4/(4*x^4) - (b^2*atanh(tanh(a + b*x))^2)/(2*x^2) - (b^3*atanh(tanh(a + b*x)))/x - (b*atanh(tanh(a + b*x))^3)/(3*x^3)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^5} dx = \frac{-3 \operatorname{atanh}(\tanh(bx + a))^4 - 4 \operatorname{atanh}(\tanh(bx + a))^3 bx - 6 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 12 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 + 12 \log(x) b^4 x^4}{12x^4}$$

input `int(atanh(tanh(b*x+a))^4/x^5,x)`output `( - 3*atanh(tanh(a + b*x))**4 - 4*atanh(tanh(a + b*x))**3*b*x - 6*atanh(tanh(a + b*x))**2*b**2*x**2 - 12*atanh(tanh(a + b*x))*b**3*x**3 + 12*log(x)*b**4*x**4)/(12*x**4)`

### 3.78 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^6} dx$

Optimal result . . . . .	638
Mathematica [B] (verified) . . . . .	638
Rubi [A] (verified) . . . . .	639
Maple [B] (verified) . . . . .	639
Fricas [A] (verification not implemented) . . . . .	640
Sympy [B] (verification not implemented) . . . . .	641
Maxima [B] (verification not implemented) . . . . .	641
Giac [A] (verification not implemented) . . . . .	642
Mupad [B] (verification not implemented) . . . . .	642
Reduce [B] (verification not implemented) . . . . .	642

#### Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{5x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output `1/5*arctanh(tanh(b*x+a))^5/x^5/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 66 vs. 2(31) = 62.

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.13

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = \frac{b^4x^4 + b^3x^3\operatorname{arctanh}(\tanh(a + bx)) + b^2x^2\operatorname{arctanh}(\tanh(a + bx))^2 + bx\operatorname{arctanh}(\tanh(a + bx))^3 + \operatorname{arctanh}(\tanh(a + bx))^4}{5x^5}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^6,x]`

output `-1/5*(b^4*x^4 + b^3*x^3*ArcTanh[Tanh[a + b*x]] + b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + b*x*ArcTanh[Tanh[a + b*x]]^3 + ArcTanh[Tanh[a + b*x]]^4)/x^5`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^5}{5x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^6,x]`

output `ArcTanh[Tanh[a + b*x]]^5/(5*x^5*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 64 vs.  $2(29) = 58$ .

Time = 1.99 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.10

method	result
parallelrisch	$-\frac{b^4 x^4 + b^3 \operatorname{arctanh}(\tanh(bx+a)) x^3 + b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + x \operatorname{arctanh}(\tanh(bx+a))^3 + \operatorname{arctanh}(\tanh(bx+a))^4}{5x^5}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{5x^5} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{4x^4} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b \left( -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3 \cdot 2x^2} \right)}{3} \right)}{4} \right)}{5}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{5x^5} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{4x^4} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^3} + \frac{2b \left( -\frac{b}{2x} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{3 \cdot 2x^2} \right)}{3} \right)}{4} \right)}{5}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^6,x,method=_RETURNVERBOSE)`

output 
$$-1/5*(b^4*x^4+b^3*\operatorname{arctanh}(\tanh(b*x+a))*x^3+b^2*x^2*\operatorname{arctanh}(\tanh(b*x+a))^2+x*\operatorname{arctanh}(\tanh(b*x+a))^3+b+\operatorname{arctanh}(\tanh(b*x+a))^4)/x^5$$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = -\frac{5b^4x^4 + 10ab^3x^3 + 10a^2b^2x^2 + 5a^3bx + a^4}{5x^5}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^6,x, algorithm="fricas")`

output 
$$-1/5*(5*b^4*x^4 + 10*a*b^3*x^3 + 10*a^2*b^2*x^2 + 5*a^3*b*x + a^4)/x^5$$

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs.  $2(26) = 52$ .

Time = 0.44 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.42

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = -\frac{b^4}{5x} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{5x^2} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{5x^3} - \frac{b \operatorname{atanh}^3(\tanh(a + bx))}{5x^4} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{5x^5}$$

input `integrate(atanh(tanh(b*x+a))**4/x**6,x)`

output `-b**4/(5*x) - b**3*atanh(tanh(a + b*x))/(5*x**2) - b**2*atanh(tanh(a + b*x))**2/(5*x**3) - b*atanh(tanh(a + b*x))**3/(5*x**4) - atanh(tanh(a + b*x))**4/(5*x**5)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 70 vs.  $2(29) = 58$ .

Time = 0.23 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.26

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = -\frac{1}{5} \left( b \left( \frac{b^2}{x} + \frac{b \operatorname{artanh}(\tanh(bx + a))}{x^2} \right) + \frac{b \operatorname{artanh}(\tanh(bx + a))^2}{x^3} \right) b - \frac{b \operatorname{artanh}(\tanh(bx + a))^3}{5x^4} - \frac{\operatorname{artanh}(\tanh(bx + a))^4}{5x^5}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^6,x, algorithm="maxima")`

output `-1/5*(b*(b^2/x + b*arctanh(tanh(b*x + a)))/x^2) + b*arctanh(tanh(b*x + a))^2/x^3)*b - 1/5*b*arctanh(tanh(b*x + a))^3/x^4 - 1/5*arctanh(tanh(b*x + a))^4/x^5`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = -\frac{5b^4x^4 + 10ab^3x^3 + 10a^2b^2x^2 + 5a^3bx + a^4}{5x^5}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^6,x, algorithm="giac")`output `-1/5*(5*b^4*x^4 + 10*a*b^3*x^3 + 10*a^2*b^2*x^2 + 5*a^3*b*x + a^4)/x^5`**Mupad [B] (verification not implemented)**

Time = 3.53 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.06

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = \frac{b^4x^4 + b^3x^3 \operatorname{atanh}(\tanh(a + bx)) + b^2x^2 \operatorname{atanh}(\tanh(a + bx))^2 + bx \operatorname{atanh}(\tanh(a + bx))^3 + \operatorname{atanh}(\tanh(a + bx))^4}{5x^5}$$

input `int(atanh(tanh(a + b*x))^4/x^6,x)`output `-(atanh(tanh(a + b*x))^4 + b^4*x^4 + b^2*x^2*atanh(tanh(a + b*x))^2 + b*x*atanh(tanh(a + b*x))^3 + b^3*x^3*atanh(tanh(a + b*x)))/(5*x^5)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.26

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx = \frac{-\operatorname{atanh}(\tanh(bx + a))^4 - \operatorname{atanh}(\tanh(bx + a))^3 bx - \operatorname{atanh}(\tanh(bx + a))^2 b^2x^2 - \operatorname{atanh}(\tanh(bx + a)) b^3x^3 - \operatorname{atanh}(\tanh(bx + a)) b^4x^4}{5x^5}$$

input `int(atanh(tanh(b*x+a))^4/x^6,x)`

output

```
( - (atanh(tanh(a + b*x))**4 + atanh(tanh(a + b*x))**3*b*x + atanh(tanh(a
+ b*x))**2*b**2*x**2 + atanh(tanh(a + b*x))*b**3*x**3 + b**4*x**4))/(5*x**
5)
```



### 3.79 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^7} dx$

Optimal result . . . . .	644
Mathematica [A] (verified) . . . . .	644
Rubi [A] (verified) . . . . .	645
Maple [A] (verified) . . . . .	646
Fricas [A] (verification not implemented) . . . . .	647
Sympy [A] (verification not implemented) . . . . .	647
Maxima [A] (verification not implemented) . . . . .	648
Giac [A] (verification not implemented) . . . . .	648
Mupad [B] (verification not implemented) . . . . .	649
Reduce [B] (verification not implemented) . . . . .	649

#### Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{30x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))^2} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output `1/30*b*arctanh(tanh(b*x+a))^5/x^5/(b*x-arctanh(tanh(b*x+a)))^2+1/6*arctanh(tanh(b*x+a))^5/x^6/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = \frac{b^4x^4 + 2b^3x^3\operatorname{arctanh}(\tanh(a + bx)) + 3b^2x^2\operatorname{arctanh}(\tanh(a + bx))^2 + 4bx\operatorname{arctanh}(\tanh(a + bx))^3 + 5\operatorname{arctanh}(\tanh(a + bx))^4}{30x^6}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^7, x]`

output

```
-1/30*(b^4*x^4 + 2*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 3*b^2*x^2*ArcTanh[Tanh
[a + b*x]]^2 + 4*b*x*ArcTanh[Tanh[a + b*x]]^3 + 5*ArcTanh[Tanh[a + b*x]]^4
)/x^6
```

**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^7} dx$$

↓ 2602

$$\frac{b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^6} dx}{6(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2598

$$\frac{\operatorname{arctanh}(\tanh(a+bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \operatorname{arctanh}(\tanh(a+bx))^5}{30x^5(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^4/x^7,x]
```

output

```
(b*ArcTanh[Tanh[a + b*x]]^5)/(30*x^5*(b*x - ArcTanh[Tanh[a + b*x]])^2) + A
rcTanh[Tanh[a + b*x]]^5/(6*x^6*(b*x - ArcTanh[Tanh[a + b*x]]))
```

## Definitions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

## Maple [A] (verified)

Time = 2.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.09

method	result
parallelrisc	$-\frac{b^4 x^4 + 2b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 + 3b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + 4x \operatorname{arctanh}(\tanh(bx+a))^3 + 5 \operatorname{arctanh}(\tanh(bx+a))^4}{30x^6}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{6x^6} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{5x^5} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b \left( -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2 \cdot 3x^3} \right)}{5} \right)}{3} \right)}{3}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{6x^6} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{5x^5} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{4x^4} + \frac{b \left( -\frac{b}{6x^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))}{2 \cdot 3x^3} \right)}{5} \right)}{3} \right)}{3}$
risc	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^7,x,method=_RETURNVERBOSE)`

output `-1/30*(b^4*x^4+2*b^3*arctanh(tanh(b*x+a))*x^3+3*b^2*x^2*arctanh(tanh(b*x+a))^2+4*x*arctanh(tanh(b*x+a))^3*b+5*arctanh(tanh(b*x+a))^4)/x^6`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = -\frac{15b^4x^4 + 40ab^3x^3 + 45a^2b^2x^2 + 24a^3bx + 5a^4}{30x^6}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^7,x, algorithm="fricas")`

output `-1/30*(15*b^4*x^4 + 40*a*b^3*x^3 + 45*a^2*b^2*x^2 + 24*a^3*b*x + 5*a^4)/x^6`

**Sympy [A] (verification not implemented)**

Time = 0.61 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.22

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = -\frac{b^4}{30x^2} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{15x^3} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{10x^4} - \frac{2b \operatorname{atanh}^3(\tanh(a + bx))}{15x^5} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{6x^6}$$

input `integrate(atanh(tanh(b*x+a))**4/x**7,x)`

output `-b**4/(30*x**2) - b**3*atanh(tanh(a + b*x))/(15*x**3) - b**2*atanh(tanh(a + b*x))**2/(10*x**4) - 2*b*atanh(tanh(a + b*x))**3/(15*x**5) - atanh(tanh(a + b*x))**4/(6*x**6)`

**Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx$$

$$= -\frac{1}{30} \left( b \left( \frac{b^2}{x^2} + \frac{2b \operatorname{arctanh}(\tanh(bx + a))}{x^3} \right) + \frac{3b \operatorname{arctanh}(\tanh(bx + a))^2}{x^4} \right) b$$

$$- \frac{2b \operatorname{arctanh}(\tanh(bx + a))^3}{15x^5} - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{6x^6}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^7,x, algorithm="maxima")`

output `-1/30*(b*(b^2/x^2 + 2*b*arctanh(tanh(b*x + a)))/x^3) + 3*b*arctanh(tanh(b*x + a))^2/x^4)*b - 2/15*b*arctanh(tanh(b*x + a))^3/x^5 - 1/6*arctanh(tanh(b*x + a))^4/x^6`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = -\frac{15b^4x^4 + 40ab^3x^3 + 45a^2b^2x^2 + 24a^3bx + 5a^4}{30x^6}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^7,x, algorithm="giac")`

output `-1/30*(15*b^4*x^4 + 40*a*b^3*x^3 + 45*a^2*b^2*x^2 + 24*a^3*b*x + 5*a^4)/x^6`

**Mupad [B] (verification not implemented)**

Time = 3.79 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.09

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^4}{6x^6} - \frac{b^4}{30x^2}$$

$$-\frac{b^2 \operatorname{atanh}(\tanh(a + bx))^2}{10x^4} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{15x^3}$$

$$-\frac{2b \operatorname{atanh}(\tanh(a + bx))^3}{15x^5}$$

input `int(atanh(tanh(a + b*x))^4/x^7,x)`output `- atanh(tanh(a + b*x))^4/(6*x^6) - b^4/(30*x^2) - (b^2*atanh(tanh(a + b*x))^2)/(10*x^4) - (b^3*atanh(tanh(a + b*x)))/(15*x^3) - (2*b*atanh(tanh(a + b*x))^3)/(15*x^5)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.09

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx$$

$$= \frac{-5 \operatorname{atanh}(\tanh(bx + a))^4 - 4 \operatorname{atanh}(\tanh(bx + a))^3 bx - 3 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 2 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 - b^4 x^4}{30x^6}$$

input `int(atanh(tanh(b*x+a))^4/x^7,x)`output `( - 5*atanh(tanh(a + b*x))**4 - 4*atanh(tanh(a + b*x))**3*b*x - 3*atanh(tanh(a + b*x))**2*b**2*x**2 - 2*atanh(tanh(a + b*x))*b**3*x**3 - b**4*x**4)/(30*x**6)`

### 3.80 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^8} dx$

Optimal result . . . . .	650
Mathematica [A] (verified) . . . . .	650
Rubi [A] (verified) . . . . .	651
Maple [A] (verified) . . . . .	652
Fricas [A] (verification not implemented) . . . . .	653
Sympy [A] (verification not implemented) . . . . .	653
Maxima [A] (verification not implemented) . . . . .	654
Giac [A] (verification not implemented) . . . . .	654
Mupad [B] (verification not implemented) . . . . .	655
Reduce [B] (verification not implemented) . . . . .	655

#### Optimal result

Integrand size = 13, antiderivative size = 98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = \frac{b^2 \operatorname{arctanh}(\tanh(a + bx))^5}{105x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))^3} + \frac{b \operatorname{arctanh}(\tanh(a + bx))^5}{21x^6(bx - \operatorname{arctanh}(\tanh(a + bx)))^2} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output

```
1/105*b^2*arctanh(tanh(b*x+a))^5/x^5/(b*x-arctanh(tanh(b*x+a)))^3+1/21*b*a
rctanh(tanh(b*x+a))^5/x^6/(b*x-arctanh(tanh(b*x+a)))^2+1/7*arctanh(tanh(b*
x+a))^5/x^7/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.72

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = \frac{b^4 x^4 + 3b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) + 6b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2 + 10bx \operatorname{arctanh}(\tanh(a + bx))^3 + \operatorname{arctanh}(\tanh(a + bx))^4}{105x^7}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^8, x]`

output `-1/105*(b^4*x^4 + 3*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 6*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 10*b*x*ArcTanh[Tanh[a + b*x]]^3 + 15*ArcTanh[Tanh[a + b*x]]^4)/x^7`

### Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.18, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx \\
 & \quad \downarrow \text{2602} \\
 & \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^7} dx}{7(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow \text{2602} \\
 & \frac{2b \left( \frac{b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^6} dx}{6(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{\frac{7(bx - \operatorname{arctanh}(\tanh(a + bx)))}{\operatorname{arctanh}(\tanh(a + bx))^5}} + \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow \text{2598} \\
 & \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \\
 & \frac{2b \left( \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{b \operatorname{arctanh}(\tanh(a + bx))^5}{30x^5(bx - \operatorname{arctanh}(\tanh(a + bx)))^2} \right)}{7(bx - \operatorname{arctanh}(\tanh(a + bx)))}
 \end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^8, x]`



output

```
ArcTanh[Tanh[a + b*x]]^5/(7*x^7*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*b*((b
*ArcTanh[Tanh[a + b*x]]^5)/(30*x^5*(b*x - ArcTanh[Tanh[a + b*x]]^2) + Arc
Tanh[Tanh[a + b*x]]^5/(6*x^6*(b*x - ArcTanh[Tanh[a + b*x]]))))/(7*(b*x - A
rcTanh[Tanh[a + b*x]]))
```

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1)/((m + 1)*(b*u - a*v))), x] /;
NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[
m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1)/((m + 1)*(b*u - a*v))), x] + S
imp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; Ne
Q[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m
, -1]
```

**Maple [A] (verified)**

Time = 2.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

method	result
parallelrisch	$-\frac{b^4 x^4 + 3b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 + 6b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + 10x \operatorname{arctanh}(\tanh(bx+a))^3 b + 15 \operatorname{arctanh}(\tanh(bx+a))}{105x^7}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{7x^7} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{6x^6} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{5x^5} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{4x^4} - \frac{b}{12x^3} \right)}{5} \right)}{2} \right)}{7}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{7x^7} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{6x^6} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{5x^5} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{4x^4} - \frac{b}{12x^3} \right)}{5} \right)}{2} \right)}{7}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^8,x,method=_RETURNVERBOSE)`

output `-1/105*(b^4*x^4+3*b^3*arctanh(tanh(b*x+a))*x^3+6*b^2*x^2*arctanh(tanh(b*x+a))^2+10*x*arctanh(tanh(b*x+a))^3*b+15*arctanh(tanh(b*x+a))^4)/x^7`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.47

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = -\frac{35b^4x^4 + 105ab^3x^3 + 126a^2b^2x^2 + 70a^3bx + 15a^4}{105x^7}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^8,x, algorithm="fricas")`

output `-1/105*(35*b^4*x^4 + 105*a*b^3*x^3 + 126*a^2*b^2*x^2 + 70*a^3*b*x + 15*a^4)/x^7`

### Sympy [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = -\frac{b^4}{105x^3} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{35x^4} - \frac{2b^2 \operatorname{atanh}^2(\tanh(a + bx))}{35x^5} - \frac{2b \operatorname{atanh}^3(\tanh(a + bx))}{21x^6} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{7x^7}$$

input `integrate(atanh(tanh(b*x+a))**4/x**8,x)`

output `-b**4/(105*x**3) - b**3*atanh(tanh(a + b*x))/(35*x**4) - 2*b**2*atanh(tanh(a + b*x))**2/(35*x**5) - 2*b*atanh(tanh(a + b*x))**3/(21*x**6) - atanh(tanh(a + b*x))**4/(7*x**7)`

**Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx$$

$$= -\frac{1}{105} \left( b \left( \frac{b^2}{x^3} + \frac{3b \operatorname{arctanh}(\tanh(bx + a))}{x^4} \right) + \frac{6b \operatorname{arctanh}(\tanh(bx + a))^2}{x^5} \right) b$$

$$- \frac{2b \operatorname{arctanh}(\tanh(bx + a))^3}{21x^6} - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{7x^7}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^8,x, algorithm="maxima")`

output `-1/105*(b*(b^2/x^3 + 3*b*arctanh(tanh(b*x + a))/x^4) + 6*b*arctanh(tanh(b*x + a))^2/x^5)*b - 2/21*b*arctanh(tanh(b*x + a))^3/x^6 - 1/7*arctanh(tanh(b*x + a))^4/x^7`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.47

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = -\frac{35b^4x^4 + 105ab^3x^3 + 126a^2b^2x^2 + 70a^3bx + 15a^4}{105x^7}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^8,x, algorithm="giac")`

output `-1/105*(35*b^4*x^4 + 105*a*b^3*x^3 + 126*a^2*b^2*x^2 + 70*a^3*b*x + 15*a^4)/x^7`

**Mupad [B] (verification not implemented)**

Time = 3.32 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^4}{7x^7} - \frac{b^4}{105x^3} - \frac{2b^2 \operatorname{atanh}(\tanh(a + bx))^2}{35x^5} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{35x^4} - \frac{2b \operatorname{atanh}(\tanh(a + bx))^3}{21x^6}$$

input `int(atanh(tanh(a + b*x))^4/x^8,x)`output `- atanh(tanh(a + b*x))^4/(7*x^7) - b^4/(105*x^3) - (2*b^2*atanh(tanh(a + b*x))^2)/(35*x^5) - (b^3*atanh(tanh(a + b*x)))/(35*x^4) - (2*b*atanh(tanh(a + b*x))^3)/(21*x^6)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^8} dx = \frac{-15 \operatorname{atanh}(\tanh(bx + a))^4 - 10 \operatorname{atanh}(\tanh(bx + a))^3 bx - 6 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 3 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 - b^4 x^4}{105x^7}$$

input `int(atanh(tanh(b*x+a))^4/x^8,x)`output `( - 15*atanh(tanh(a + b*x))**4 - 10*atanh(tanh(a + b*x))**3*b*x - 6*atanh(tanh(a + b*x))**2*b**2*x**2 - 3*atanh(tanh(a + b*x))*b**3*x**3 - b**4*x**4 )/(105*x**7)`

### 3.81 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx$

Optimal result . . . . .	656
Mathematica [A] (verified) . . . . .	656
Rubi [B] (verified) . . . . .	657
Maple [A] (verified) . . . . .	659
Fricas [A] (verification not implemented) . . . . .	659
Sympy [A] (verification not implemented) . . . . .	660
Maxima [A] (verification not implemented) . . . . .	660
Giac [A] (verification not implemented) . . . . .	661
Mupad [B] (verification not implemented) . . . . .	661
Reduce [B] (verification not implemented) . . . . .	662

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx = -\frac{b^4}{280x^4} - \frac{b^3 \operatorname{arctanh}(\tanh(a+bx))}{70x^5} - \frac{b^2 \operatorname{arctanh}(\tanh(a+bx))^2}{28x^6} - \frac{b \operatorname{arctanh}(\tanh(a+bx))^3}{14x^7} - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{8x^8}$$

output

```
-1/280*b^4/x^4-1/70*b^3*arctanh(tanh(b*x+a))/x^5-1/28*b^2*arctanh(tanh(b*x+a))^2/x^6-1/14*b*arctanh(tanh(b*x+a))^3/x^7-1/8*arctanh(tanh(b*x+a))^4/x^8
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx = \frac{b^4 x^4 + 4b^3 x^3 \operatorname{arctanh}(\tanh(a+bx)) + 10b^2 x^2 \operatorname{arctanh}(\tanh(a+bx))^2 + 20bx \operatorname{arctanh}(\tanh(a+bx))^3 - \operatorname{arctanh}(\tanh(a+bx))^4}{280x^8}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^9, x]`

output 
$$-1/280*(b^4*x^4 + 4*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 10*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 20*b*x*ArcTanh[Tanh[a + b*x]]^3 + 35*ArcTanh[Tanh[a + b*x]]^4)/x^8$$

### Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 168 vs.  $2(80) = 160$ .

Time = 0.43 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.10, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx$$

$$\downarrow 2602$$

$$\frac{3b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^8} dx}{8(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{8x^8(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2602$$

$$\frac{3b \left( \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^7} dx}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{\frac{8(bx - \operatorname{arctanh}(\tanh(a + bx)))}{\operatorname{arctanh}(\tanh(a + bx))^5}} + \frac{\operatorname{arctanh}(\tanh(a + bx))^5}{8x^8(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2602$$

$$\begin{aligned}
& 3b \left( \frac{2b \left( \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^6} dx + \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{8(bx - \operatorname{arctanh}(\tanh(a+bx)))}{\operatorname{arctanh}(\tanh(a+bx))^5} \\
& \frac{8x^8(bx - \operatorname{arctanh}(\tanh(a+bx)))}{\operatorname{arctanh}(\tanh(a+bx))^5} \\
& \quad \downarrow \text{2598} \\
& \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{8x^8(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \\
& 3b \left( \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{7x^7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2b \left( \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{6x^6(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{\operatorname{arctanh}(\tanh(a+bx))^5}{30x^5(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right)}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{8(bx - \operatorname{arctanh}(\tanh(a+bx)))}{\operatorname{arctanh}(\tanh(a+bx))^5}
\end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^9,x]`

output `ArcTanh[Tanh[a + b*x]]^5/(8*x^8*(b*x - ArcTanh[Tanh[a + b*x]])) + (3*b*(ArcTanh[Tanh[a + b*x]]^5/(7*x^7*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*b*((b*ArcTanh[Tanh[a + b*x]]^5)/(30*x^5*(b*x - ArcTanh[Tanh[a + b*x]]^2) + ArcTanh[Tanh[a + b*x]]^5/(6*x^6*(b*x - ArcTanh[Tanh[a + b*x]])))))/(7*(b*x - ArcTanh[Tanh[a + b*x]])))/(8*(b*x - ArcTanh[Tanh[a + b*x]]))`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v))] Int[u^(m + 1)*v^n, x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

**Maple [A] (verified)**

Time = 2.44 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

method	result
parallelrisch	$-\frac{b^4 x^4 + 4b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 + 10b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + 20x \operatorname{arctanh}(\tanh(bx+a))^3 b + 35 \operatorname{arctanh}(\tanh(bx+a))^4}{280x^8}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{8x^8} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{7x^7} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{6x^6} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{5x^5} - \frac{b}{20x^4} \right)}{3} \right)}{7} \right)}{2}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{8x^8} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{7x^7} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{6x^6} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{5x^5} - \frac{b}{20x^4} \right)}{3} \right)}{7} \right)}{2}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^9,x,method=_RETURNVERBOSE)`

output 
$$-1/280*(b^4*x^4+4*b^3*\operatorname{arctanh}(\tanh(b*x+a))*x^3+10*b^2*x^2*\operatorname{arctanh}(\tanh(b*x+a))^2+20*x*\operatorname{arctanh}(\tanh(b*x+a))^3*b+35*\operatorname{arctanh}(\tanh(b*x+a))^4)/x^8$$

**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^9} dx = -\frac{70b^4x^4 + 224ab^3x^3 + 280a^2b^2x^2 + 160a^3bx + 35a^4}{280x^8}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^9,x, algorithm="fricas")`

output 
$$-1/280*(70*b^4*x^4 + 224*a*b^3*x^3 + 280*a^2*b^2*x^2 + 160*a^3*b*x + 35*a^4)/x^8$$



**Sympy [A] (verification not implemented)**

Time = 1.21 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx = -\frac{b^4}{280x^4} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{70x^5} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{28x^6} - \frac{b \operatorname{atanh}^3(\tanh(a + bx))}{14x^7} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{8x^8}$$

input `integrate(atanh(tanh(b*x+a))**4/x**9,x)`output `-b**4/(280*x**4) - b**3*atanh(tanh(a + b*x))/(70*x**5) - b**2*atanh(tanh(a + b*x))**2/(28*x**6) - b*atanh(tanh(a + b*x))**3/(14*x**7) - atanh(tanh(a + b*x))**4/(8*x**8)`**Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx = -\frac{1}{280} \left( b \left( \frac{b^2}{x^4} + \frac{4b \operatorname{artanh}(\tanh(bx + a))}{x^5} \right) + \frac{10b \operatorname{artanh}(\tanh(bx + a))^2}{x^6} \right) b - \frac{b \operatorname{artanh}(\tanh(bx + a))^3}{14x^7} - \frac{\operatorname{artanh}(\tanh(bx + a))^4}{8x^8}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^9,x, algorithm="maxima")`output `-1/280*(b*(b^2/x^4 + 4*b*arctanh(tanh(b*x + a)))/x^5) + 10*b*arctanh(tanh(b*x + a))^2/x^6)*b - 1/14*b*arctanh(tanh(b*x + a))^3/x^7 - 1/8*arctanh(tanh(b*x + a))^4/x^8`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx = -\frac{70b^4x^4 + 224ab^3x^3 + 280a^2b^2x^2 + 160a^3bx + 35a^4}{280x^8}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^9,x, algorithm="giac")`

output `-1/280*(70*b^4*x^4 + 224*a*b^3*x^3 + 280*a^2*b^2*x^2 + 160*a^3*b*x + 35*a^4)/x^8`

**Mupad [B] (verification not implemented)**

Time = 3.50 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^4}{8x^8} - \frac{b^4}{280x^4} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))^2}{28x^6} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{70x^5} - \frac{b \operatorname{atanh}(\tanh(a + bx))^3}{14x^7}$$

input `int(atanh(tanh(a + b*x))^4/x^9,x)`

output `- atanh(tanh(a + b*x))^4/(8*x^8) - b^4/(280*x^4) - (b^2*atanh(tanh(a + b*x)))^2/(28*x^6) - (b^3*atanh(tanh(a + b*x)))/(70*x^5) - (b*atanh(tanh(a + b*x)))^3/(14*x^7)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^9} dx$$

$$= \frac{-35 \operatorname{atanh}(\tanh(bx + a))^4 - 20 \operatorname{atanh}(\tanh(bx + a))^3 bx - 10 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 4 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 - b^4 x^4}{280 x^8}$$

input `int(atanh(tanh(b*x+a))^4/x^9,x)`output `( - 35*atanh(tanh(a + b*x))**4 - 20*atanh(tanh(a + b*x))**3*b*x - 10*atanh(tanh(a + b*x))**2*b**2*x**2 - 4*atanh(tanh(a + b*x))*b**3*x**3 - b**4*x**4)/(280*x**8)`

### 3.82 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{10}} dx$

Optimal result . . . . .	663
Mathematica [A] (verified) . . . . .	663
Rubi [A] (verified) . . . . .	664
Maple [A] (verified) . . . . .	666
Fricas [A] (verification not implemented) . . . . .	666
Sympy [A] (verification not implemented) . . . . .	667
Maxima [A] (verification not implemented) . . . . .	667
Giac [A] (verification not implemented) . . . . .	668
Mupad [B] (verification not implemented) . . . . .	668
Reduce [B] (verification not implemented) . . . . .	669

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{b^4}{630x^5} - \frac{b^3 \operatorname{arctanh}(\tanh(a + bx))}{126x^6} - \frac{b^2 \operatorname{arctanh}(\tanh(a + bx))^2}{42x^7} - \frac{b \operatorname{arctanh}(\tanh(a + bx))^3}{18x^8} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{9x^9}$$

output

```
-1/630*b^4/x^5-1/126*b^3*arctanh(tanh(b*x+a))/x^6-1/42*b^2*arctanh(tanh(b*x+a))^2/x^7-1/18*b*arctanh(tanh(b*x+a))^3/x^8-1/9*arctanh(tanh(b*x+a))^4/x^9
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = \frac{b^4 x^4 + 5b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) + 15b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2 + 35bx \operatorname{arctanh}(\tanh(a + bx))^3 - \operatorname{arctanh}(\tanh(a + bx))^4}{630x^9}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^10,x]`

output 
$$-1/630*(b^4*x^4 + 5*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 15*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 35*b*x*ArcTanh[Tanh[a + b*x]]^3 + 70*ArcTanh[Tanh[a + b*x]]^4)/x^9$$

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx \\ & \quad \downarrow 2599 \\ & \frac{4}{9}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^9} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{9x^9} \\ & \quad \downarrow 2599 \\ & \frac{4}{9}b \left( \frac{3}{8}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^8} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{9x^9} \\ & \quad \downarrow 2599 \\ & \frac{4}{9}b \left( \frac{3}{8}b \left( \frac{2}{7}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^7} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{7x^7} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{9x^9} \\ & \quad \downarrow 2599 \\ & \frac{4}{9}b \left( \frac{3}{8}b \left( \frac{2}{7}b \left( \frac{1}{6}b \int \frac{1}{x^6} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{6x^6} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{7x^7} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{9x^9} \end{aligned}$$

↓ 15

$$\frac{4}{9}b \left( \frac{3}{8}b \left( \frac{2}{7}b \left( -\frac{\operatorname{arctanh}(\tanh(a+bx))}{6x^6} - \frac{b}{30x^5} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{7x^7} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{9x^9}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^10,x]`

output `-1/9*ArcTanh[Tanh[a + b*x]]^4/x^9 + (4*b*(-1/8*ArcTanh[Tanh[a + b*x]]^3/x^8 + (3*b*(-1/7*ArcTanh[Tanh[a + b*x]]^2/x^7 + (2*b*(-1/30*b/x^5 - ArcTanh[Tanh[a + b*x]]/(6*x^6)))/7))/8)/9`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 2.66 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

method	result
parallelrisch	$-\frac{b^4 x^4 + 5b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 + 15b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + 35x \operatorname{arctanh}(\tanh(bx+a))^3 b + 70 \operatorname{arctanh}(\tanh(bx+a))^4}{630x^9}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{9x^9} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{8x^8} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{7x^7} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{6x^6} - \frac{b}{30x^5} \right)}{7} \right)}{8} \right)}{9}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{9x^9} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{8x^8} + \frac{3b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{7x^7} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{6x^6} - \frac{b}{30x^5} \right)}{7} \right)}{8} \right)}{9}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^10,x,method=_RETURNVERBOSE)`

output `-1/630*(b^4*x^4+5*b^3*arctanh(tanh(b*x+a))*x^3+15*b^2*x^2*arctanh(tanh(b*x+a))^2+35*x*arctanh(tanh(b*x+a))^3*b+70*arctanh(tanh(b*x+a))^4)/x^9`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{126 b^4 x^4 + 420 a b^3 x^3 + 540 a^2 b^2 x^2 + 315 a^3 b x + 70 a^4}{630 x^9}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^10,x, algorithm="fricas")`

output `-1/630*(126*b^4*x^4 + 420*a*b^3*x^3 + 540*a^2*b^2*x^2 + 315*a^3*b*x + 70*a^4)/x^9`

**Sympy [A] (verification not implemented)**

Time = 1.88 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{b^4}{630x^5} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{126x^6} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{42x^7} - \frac{b \operatorname{atanh}^3(\tanh(a + bx))}{18x^8} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{9x^9}$$

input `integrate(atanh(tanh(b*x+a))**4/x**10,x)`output `-b**4/(630*x**5) - b**3*atanh(tanh(a + b*x))/(126*x**6) - b**2*atanh(tanh(a + b*x))**2/(42*x**7) - b*atanh(tanh(a + b*x))**3/(18*x**8) - atanh(tanh(a + b*x))**4/(9*x**9)`**Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{1}{630} \left( b \left( \frac{b^2}{x^5} + \frac{5b \operatorname{arctanh}(\tanh(bx + a))}{x^6} \right) + \frac{15b \operatorname{arctanh}(\tanh(bx + a))^2}{x^7} \right) b - \frac{b \operatorname{arctanh}(\tanh(bx + a))^3}{18x^8} - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{9x^9}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^10,x, algorithm="maxima")`output `-1/630*(b*(b^2/x^5 + 5*b*arctanh(tanh(b*x + a)))/x^6) + 15*b*arctanh(tanh(b*x + a))^2/x^7)*b - 1/18*b*arctanh(tanh(b*x + a))^3/x^8 - 1/9*arctanh(tanh(b*x + a))^4/x^9`



**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{126 b^4 x^4 + 420 ab^3 x^3 + 540 a^2 b^2 x^2 + 315 a^3 b x + 70 a^4}{630 x^9}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^10,x, algorithm="giac")`

output `-1/630*(126*b^4*x^4 + 420*a*b^3*x^3 + 540*a^2*b^2*x^2 + 315*a^3*b*x + 70*a^4)/x^9`

**Mupad [B] (verification not implemented)**

Time = 3.53 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^4}{9 x^9} - \frac{b^4}{630 x^5} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))^2}{42 x^7} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{126 x^6} - \frac{b \operatorname{atanh}(\tanh(a + bx))^3}{18 x^8}$$

input `int(atanh(tanh(a + b*x))^4/x^10,x)`

output `- atanh(tanh(a + b*x))^4/(9*x^9) - b^4/(630*x^5) - (b^2*atanh(tanh(a + b*x)))^2/(42*x^7) - (b^3*atanh(tanh(a + b*x)))/(126*x^6) - (b*atanh(tanh(a + b*x)))^3/(18*x^8)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{10}} dx$$

$$= \frac{-70 \operatorname{atanh}(\tanh(bx + a))^4 - 35 \operatorname{atanh}(\tanh(bx + a))^3 bx - 15 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 5 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 - b^4 x^4}{630 x^9}$$

input `int(atanh(tanh(b*x+a))^4/x^10,x)`output `( - 70*atanh(tanh(a + b*x))**4 - 35*atanh(tanh(a + b*x))**3*b*x - 15*atanh(tanh(a + b*x))**2*b**2*x**2 - 5*atanh(tanh(a + b*x))*b**3*x**3 - b**4*x**4)/(630*x**9)`

### 3.83 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{x^{11}} dx$

Optimal result . . . . .	670
Mathematica [A] (verified) . . . . .	670
Rubi [A] (verified) . . . . .	671
Maple [A] (verified) . . . . .	673
Fricas [A] (verification not implemented) . . . . .	673
Sympy [A] (verification not implemented) . . . . .	674
Maxima [A] (verification not implemented) . . . . .	674
Giac [A] (verification not implemented) . . . . .	675
Mupad [B] (verification not implemented) . . . . .	675
Reduce [B] (verification not implemented) . . . . .	676

#### Optimal result

Integrand size = 13, antiderivative size = 80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{b^4}{1260x^6} - \frac{b^3 \operatorname{arctanh}(\tanh(a + bx))}{210x^7} - \frac{b^2 \operatorname{arctanh}(\tanh(a + bx))^2}{60x^8} - \frac{2b \operatorname{arctanh}(\tanh(a + bx))^3}{45x^9} - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{10x^{10}}$$

output

```
-1/1260*b^4/x^6-1/210*b^3*arctanh(tanh(b*x+a))/x^7-1/60*b^2*arctanh(tanh(b*x+a))^2/x^8-2/45*b*arctanh(tanh(b*x+a))^3/x^9-1/10*arctanh(tanh(b*x+a))^4/x^10
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = \frac{b^4 x^4 + 6b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) + 21b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^2 + 56bx \operatorname{arctanh}(\tanh(a + bx))^3 - \operatorname{arctanh}(\tanh(a + bx))^4}{1260x^{10}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^4/x^11,x]`

output 
$$-1/1260*(b^4*x^4 + 6*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 21*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 56*b*x*ArcTanh[Tanh[a + b*x]]^3 + 126*ArcTanh[Tanh[a + b*x]]^4)/x^{10}$$

### Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx \\ & \quad \downarrow 2599 \\ & \frac{2}{5}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{10}} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{10x^{10}} \\ & \quad \downarrow 2599 \\ & \frac{2}{5}b \left( \frac{1}{3}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^9} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{9x^9} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{10x^{10}} \\ & \quad \downarrow 2599 \\ & \frac{2}{5}b \left( \frac{1}{3}b \left( \frac{1}{4}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^8} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{9x^9} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{10x^{10}} \\ & \quad \downarrow 2599 \\ & \frac{2}{5}b \left( \frac{1}{3}b \left( \frac{1}{4}b \left( \frac{1}{7}b \int \frac{1}{x^7} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))}{7x^7} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{9x^9} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{10x^{10}} \end{aligned}$$

↓ 15

$$\frac{2}{5}b \left( \frac{1}{3}b \left( \frac{1}{4}b \left( -\frac{\operatorname{arctanh}(\tanh(a+bx))}{7x^7} - \frac{b}{42x^6} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{8x^8} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{9x^9} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^4}{10x^{10}}$$

input `Int[ArcTanh[Tanh[a + b*x]]^4/x^11,x]`

output `-1/10*ArcTanh[Tanh[a + b*x]]^4/x^10 + (2*b*(-1/9*ArcTanh[Tanh[a + b*x]]^3/x^9 + (b*(-1/8*ArcTanh[Tanh[a + b*x]]^2/x^8 + (b*(-1/42*b/x^6 - ArcTanh[Tanh[a + b*x]]/(7*x^7)))/4))/3)/5`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 3.05 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

method	result
parallelrisch	$-\frac{b^4 x^4 + 6b^3 \operatorname{arctanh}(\tanh(bx+a))x^3 + 21b^2 x^2 \operatorname{arctanh}(\tanh(bx+a))^2 + 56x \operatorname{arctanh}(\tanh(bx+a))^3 b + 126 \operatorname{arctanh}(\tanh(bx+a))^4}{1260x^{10}}$
default	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{10x^{10}} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{9x^9} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{8x^8} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{7x^7} - \frac{b}{42x^6} \right)}{4} \right)}{3} \right)}{5}$
parts	$-\frac{\operatorname{arctanh}(\tanh(bx+a))^4}{10x^{10}} + \frac{2b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^3}{9x^9} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{8x^8} + \frac{b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{7x^7} - \frac{b}{42x^6} \right)}{4} \right)}{3} \right)}{5}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^4/x^11,x,method=_RETURNVERBOSE)`

output `-1/1260*(b^4*x^4+6*b^3*arctanh(tanh(b*x+a))*x^3+21*b^2*x^2*arctanh(tanh(b*x+a))^2+56*x*arctanh(tanh(b*x+a))^3*b+126*arctanh(tanh(b*x+a))^4)/x^10`

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{210 b^4 x^4 + 720 a b^3 x^3 + 945 a^2 b^2 x^2 + 560 a^3 b x + 126 a^4}{1260 x^{10}}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^11,x, algorithm="fricas")`

output `-1/1260*(210*b^4*x^4 + 720*a*b^3*x^3 + 945*a^2*b^2*x^2 + 560*a^3*b*x + 126*a^4)/x^10`

**Sympy [A] (verification not implemented)**

Time = 2.89 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{b^4}{1260x^6} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{210x^7} - \frac{b^2 \operatorname{atanh}^2(\tanh(a + bx))}{60x^8} - \frac{2b \operatorname{atanh}^3(\tanh(a + bx))}{45x^9} - \frac{\operatorname{atanh}^4(\tanh(a + bx))}{10x^{10}}$$

input `integrate(atanh(tanh(b*x+a))**4/x**11,x)`output `-b**4/(1260*x**6) - b**3*atanh(tanh(a + b*x))/(210*x**7) - b**2*atanh(tanh(a + b*x))**2/(60*x**8) - 2*b*atanh(tanh(a + b*x))**3/(45*x**9) - atanh(tanh(a + b*x))**4/(10*x**10)`**Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{1}{1260} \left( b \left( \frac{b^2}{x^6} + \frac{6b \operatorname{arctanh}(\tanh(bx + a))}{x^7} \right) + \frac{21b \operatorname{arctanh}(\tanh(bx + a))^2}{x^8} \right) b - \frac{2b \operatorname{arctanh}(\tanh(bx + a))^3}{45x^9} - \frac{\operatorname{arctanh}(\tanh(bx + a))^4}{10x^{10}}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^11,x, algorithm="maxima")`output `-1/1260*(b*(b^2/x^6 + 6*b*arctanh(tanh(b*x + a))/x^7) + 21*b*arctanh(tanh(b*x + a))^2/x^8)*b - 2/45*b*arctanh(tanh(b*x + a))^3/x^9 - 1/10*arctanh(tanh(b*x + a))^4/x^10`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{210 b^4 x^4 + 720 a b^3 x^3 + 945 a^2 b^2 x^2 + 560 a^3 b x + 126 a^4}{1260 x^{10}}$$

input `integrate(arctanh(tanh(b*x+a))^4/x^11,x, algorithm="giac")`

output `-1/1260*(210*b^4*x^4 + 720*a*b^3*x^3 + 945*a^2*b^2*x^2 + 560*a^3*b*x + 126*a^4)/x^10`

**Mupad [B] (verification not implemented)**

Time = 3.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx = -\frac{\operatorname{atanh}(\tanh(a + bx))^4}{10 x^{10}} - \frac{b^4}{1260 x^6} - \frac{b^2 \operatorname{atanh}(\tanh(a + bx))^2}{60 x^8} - \frac{b^3 \operatorname{atanh}(\tanh(a + bx))}{210 x^7} - \frac{2 b \operatorname{atanh}(\tanh(a + bx))^3}{45 x^9}$$

input `int(atanh(tanh(a + b*x))^4/x^11,x)`

output `- atanh(tanh(a + b*x))^4/(10*x^10) - b^4/(1260*x^6) - (b^2*atanh(tanh(a + b*x))^2)/(60*x^8) - (b^3*atanh(tanh(a + b*x)))/(210*x^7) - (2*b*atanh(tanh(a + b*x))^3)/(45*x^9)`



**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^4}{x^{11}} dx$$

$$= \frac{-126 \operatorname{atanh}(\tanh(bx + a))^4 - 56 \operatorname{atanh}(\tanh(bx + a))^3 bx - 21 \operatorname{atanh}(\tanh(bx + a))^2 b^2 x^2 - 6 \operatorname{atanh}(\tanh(bx + a)) b^3 x^3 - b^4 x^4}{1260 x^{10}}$$

input `int(atanh(tanh(b*x+a))^4/x^11,x)`output `( - 126*atanh(tanh(a + b*x))**4 - 56*atanh(tanh(a + b*x))**3*b*x - 21*atanh(tanh(a + b*x))**2*b**2*x**2 - 6*atanh(tanh(a + b*x))*b**3*x**3 - b**4*x**4)/(1260*x**10)`

### 3.84 $\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx$

Optimal result	677
Mathematica [B] (verified)	677
Rubi [A] (verified)	678
Maple [B] (verified)	679
Fricas [B] (verification not implemented)	681
Sympy [A] (verification not implemented)	681
Maxima [B] (verification not implemented)	682
Giac [B] (verification not implemented)	682
Mupad [B] (verification not implemented)	683
Reduce [F]	683

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^8}{56b^2}$$

output

```
1/7*x*arctanh(tanh(b*x+a))^7/b-1/56*arctanh(tanh(b*x+a))^8/b^2
```

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 177 vs. 2(34) = 68.

Time = 0.23 (sec) , antiderivative size = 177, normalized size of antiderivative = 5.21

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \frac{(a + bx) ((7a - bx)(a + bx)^6 - 8(6a - bx)(a + bx)^5 \operatorname{arctanh}(\tanh(a + bx)) + 28(5a - bx)(a + bx)^4 \operatorname{arctanh}(\tanh(a + bx))^2 - \dots)}{\dots}$$

input

```
Integrate[x*ArcTanh[Tanh[a + b*x]]^6,x]
```

output

```
-1/56*((a + b*x)*((7*a - b*x)*(a + b*x)^6 - 8*(6*a - b*x)*(a + b*x)^5*ArcTanh[Tanh[a + b*x]] + 28*(5*a - b*x)*(a + b*x)^4*ArcTanh[Tanh[a + b*x]]^2 - 56*(4*a - b*x)*(a + b*x)^3*ArcTanh[Tanh[a + b*x]]^3 + 70*(3*a - b*x)*(a + b*x)^2*ArcTanh[Tanh[a + b*x]]^4 - 56*(2*a^2 + a*b*x - b^2*x^2)*ArcTanh[Tanh[a + b*x]]^5 + 28*(a - b*x)*ArcTanh[Tanh[a + b*x]]^6))/b^2
```

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx$$

$$\downarrow 2599$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 dx}{7b}$$

$$\downarrow 2588$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^7 d \operatorname{arctanh}(\tanh(a + bx))}{7b^2}$$

$$\downarrow 15$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^7}{7b} - \frac{\operatorname{arctanh}(\tanh(a + bx))^8}{56b^2}$$

input

```
Int [x*ArcTanh[Tanh[a + b*x]]^6, x]
```

output

```
(x*ArcTanh[Tanh[a + b*x]]^7)/(7*b) - ArcTanh[Tanh[a + b*x]]^8/(56*b^2)
```

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_.)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 104 vs.  $2(30) = 60$ .

Time = 41.61 (sec) , antiderivative size = 105, normalized size of antiderivative = 3.09

method	result
parallelrisch	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^6}{2} + \frac{b^6 x^8}{56} - b \operatorname{arctanh}(\tanh(bx+a))^5 x^3 - \frac{b^5 \operatorname{arctanh}(\tanh(bx+a))x^7}{7} + \frac{b^4 \operatorname{arctanh}(\tanh(bx+a))^4 x^4}{4} - b^3 \operatorname{arctanh}(\tanh(bx+a))^3 x^2 + \frac{b^2 \operatorname{arctanh}(\tanh(bx+a))^2 x^5}{2} - \frac{b \operatorname{arctanh}(\tanh(bx+a))x^8}{8} + \frac{x^6}{6}$
default	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^6}{2} - 3b \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^5}{3} - \frac{5b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^4}{4} - b \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))^3}{5} \right)}{1} - \frac{b^2 \operatorname{arctanh}(\tanh(bx+a))^2 x^5}{2} - \frac{b^3 \operatorname{arctanh}(\tanh(bx+a))x^8}{8} + \frac{x^6}{6}$
parts	$\frac{x^2 \operatorname{arctanh}(\tanh(bx+a))^6}{2} - 3b \frac{x^3 \operatorname{arctanh}(\tanh(bx+a))^5}{3} - \frac{5b \left( \frac{x^4 \operatorname{arctanh}(\tanh(bx+a))^4}{4} - b \frac{x^5 \operatorname{arctanh}(\tanh(bx+a))^3}{5} \right)}{1} - \frac{b^2 \operatorname{arctanh}(\tanh(bx+a))^2 x^5}{2} - \frac{b^3 \operatorname{arctanh}(\tanh(bx+a))x^8}{8} + \frac{x^6}{6}$
risch	Expression too large to display

```
input int(x*arctanh(tanh(b*x+a))^6,x,method=_RETURNVERBOSE)
```

```
output 1/2*x^2*arctanh(tanh(b*x+a))^6+1/56*b^6*x^8-b*arctanh(tanh(b*x+a))^5*x^3-1/7*b^5*arctanh(tanh(b*x+a))*x^7+1/2*b^4*arctanh(tanh(b*x+a))^2*x^6-b^3*arctanh(tanh(b*x+a))^3*x^5+5/4*b^2*arctanh(tanh(b*x+a))^4*x^4
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 68 vs.  $2(30) = 60$ .

Time = 0.07 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.00

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \frac{1}{8} b^6 x^8 + \frac{6}{7} ab^5 x^7 + \frac{5}{2} a^2 b^4 x^6 + 4 a^3 b^3 x^5 + \frac{15}{4} a^4 b^2 x^4 + 2 a^5 b x^3 + \frac{1}{2} a^6 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^6,x, algorithm="fricas")`

output `1/8*b^6*x^8 + 6/7*a*b^5*x^7 + 5/2*a^2*b^4*x^6 + 4*a^3*b^3*x^5 + 15/4*a^4*b^2*x^4 + 2*a^5*b*x^3 + 1/2*a^6*x^2`

**Sympy [A] (verification not implemented)**

Time = 0.76 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \begin{cases} \frac{x \operatorname{atanh}^7(\tanh(a+bx))}{7b} - \frac{\operatorname{atanh}^8(\tanh(a+bx))}{56b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^6(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**6,x)`

output `Piecewise((x*atanh(tanh(a + b*x))**7/(7*b) - atanh(tanh(a + b*x))**8/(56*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**6/2, True))`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 110 vs.  $2(30) = 60$ .

Time = 0.33 (sec) , antiderivative size = 110, normalized size of antiderivative = 3.24

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx$$

$$= -bx^3 \operatorname{arctanh}(\tanh(bx + a))^5 + \frac{1}{2} x^2 \operatorname{arctanh}(\tanh(bx + a))^6$$

$$+ \frac{1}{56} (70bx^4 \operatorname{arctanh}(\tanh(bx + a))^4 - (56bx^5 \operatorname{arctanh}(\tanh(bx + a)))^3 - (28bx^6 \operatorname{arctanh}(\tanh(bx + a)))^2 + (b^2x^8 - 8bx^7 \operatorname{arctanh}(\tanh(bx + a))) * b) * b) * b$$

input `integrate(x*arctanh(tanh(b*x+a))^6,x, algorithm="maxima")`

output `-b*x^3*arctanh(tanh(b*x + a))^5 + 1/2*x^2*arctanh(tanh(b*x + a))^6 + 1/56*(70*b*x^4*arctanh(tanh(b*x + a))^4 - (56*b*x^5*arctanh(tanh(b*x + a)))^3 - (28*b*x^6*arctanh(tanh(b*x + a)))^2 + (b^2*x^8 - 8*b*x^7*arctanh(tanh(b*x + a))) * b) * b) * b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 68 vs.  $2(30) = 60$ .

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.00

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \frac{1}{8} b^6 x^8 + \frac{6}{7} ab^5 x^7 + \frac{5}{2} a^2 b^4 x^6 + 4 a^3 b^3 x^5$$

$$+ \frac{15}{4} a^4 b^2 x^4 + 2 a^5 b x^3 + \frac{1}{2} a^6 x^2$$

input `integrate(x*arctanh(tanh(b*x+a))^6,x, algorithm="giac")`

output `1/8*b^6*x^8 + 6/7*a*b^5*x^7 + 5/2*a^2*b^4*x^6 + 4*a^3*b^3*x^5 + 15/4*a^4*b^2*x^4 + 2*a^5*b*x^3 + 1/2*a^6*x^2`

**Mupad [B] (verification not implemented)**

Time = 3.39 (sec) , antiderivative size = 104, normalized size of antiderivative = 3.06

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \frac{b^6 x^8}{56} - \frac{b^5 x^7 \operatorname{atanh}(\tanh(a + bx))}{7} + \frac{b^4 x^6 \operatorname{atanh}(\tanh(a + bx))^2}{2} - b^3 x^5 \operatorname{atanh}(\tanh(a + bx))^3 + \frac{5 b^2 x^4 \operatorname{atanh}(\tanh(a + bx))^4}{4} - b x^3 \operatorname{atanh}(\tanh(a + bx))^5 + \frac{x^2 \operatorname{atanh}(\tanh(a + bx))^6}{2}$$

input `int(x*atanh(tanh(a + b*x))^6,x)`output `(x^2*atanh(tanh(a + b*x))^6)/2 + (b^6*x^8)/56 + (5*b^2*x^4*atanh(tanh(a + b*x))^4)/4 - b^3*x^5*atanh(tanh(a + b*x))^3 + (b^4*x^6*atanh(tanh(a + b*x))^2)/2 - b*x^3*atanh(tanh(a + b*x))^5 - (b^5*x^7*atanh(tanh(a + b*x)))/7`**Reduce [F]**

$$\int x \operatorname{arctanh}(\tanh(a + bx))^6 dx = \int \operatorname{atanh}(\tanh(bx + a))^6 x dx$$

input `int(x*atanh(tanh(b*x+a))^6,x)`output `int(atanh(tanh(a + b*x))^6*x,x)`



### 3.85 $\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	684
Mathematica [A] (verified)	684
Rubi [A] (verified)	685
Maple [F]	686
Fricas [F]	686
Sympy [F]	686
Maxima [F]	687
Giac [F]	687
Mupad [F(-1)]	687
Reduce [F]	688

#### Optimal result

Integrand size = 13, antiderivative size = 53

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{bx}{bx-\operatorname{arctanh}(\tanh(a+bx))}\right)}{(1+m)(bx-\operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
-x^(1+m)*hypergeom([1, 1+m],[2+m],b*x/(b*x-arctanh(tanh(b*x+a))))/(1+m)/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.96

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, -\frac{bx}{-bx+\operatorname{arctanh}(\tanh(a+bx))}\right)}{(1+m)(-bx+\operatorname{arctanh}(\tanh(a+bx)))}$$

input

```
Integrate[x^m/ArcTanh[Tanh[a + b*x]],x]
```

output

```
(x^(1 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, -((b*x)/(-b*x) + ArcTanh[Tanh[a + b*x]])]) / ((1 + m)*(-b*x) + ArcTanh[Tanh[a + b*x]])
```

### Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

↓ 2595

$$-\frac{x^{m+1} \operatorname{Hypergeometric2F1}\left(1, m+1, m+2, \frac{bx}{bx - \operatorname{arctanh}(\tanh(a + bx))}\right)}{(m+1)(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input

```
Int[x^m/ArcTanh[Tanh[a + b*x]],x]
```

output

```
-((x^(1 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (b*x)/(b*x - ArcTanh[Tanh[a + b*x]])]) / ((1 + m)*(b*x - ArcTanh[Tanh[a + b*x]])))
```

### Defintions of rubi rules used

rule 2595

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(v^(n + 1)/((n + 1)*(b*u - a*v)))*Hypergeometric2F1[1, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && !IntegerQ[n]
```

**Maple [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(bx + a))} dx$$

input `int(x^m/arctanh(tanh(b*x+a)),x)`

output `int(x^m/arctanh(tanh(b*x+a)),x)`

**Fricas [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `integral(x^m/arctanh(tanh(b*x + a)), x)`

**Sympy [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**m/atanh(tanh(b*x+a)),x)`

output `Integral(x**m/atanh(tanh(a + b*x)), x)`

**Maxima [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `integrate(x^m/arctanh(tanh(b*x + a)), x)`

**Giac [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x^m/arctanh(tanh(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `int(x^m/atanh(tanh(a + b*x)),x)`

output `int(x^m/atanh(tanh(a + b*x)), x)`

Reduce [F]

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^m}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(xm/atanh(tanh(b*x+a)),x)`

output `int(x**m/atanh(tanh(a + b*x)),x)`

### 3.86 $\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	689
Mathematica [A] (verified)	689
Rubi [A] (verified)	690
Maple [B] (verified)	692
Fricas [A] (verification not implemented)	692
Sympy [F]	693
Maxima [A] (verification not implemented)	693
Giac [A] (verification not implemented)	693
Mupad [B] (verification not implemented)	694
Reduce [F]	694

#### Optimal result

Integrand size = 13, antiderivative size = 81

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx$$

$$= \frac{x^3}{3b} + \frac{x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2b^2} + \frac{x(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{b^3}$$

$$+ \frac{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

output

```
1/3*x^3/b+1/2*x^2*(b*x-arctanh(tanh(b*x+a)))/b^2+x*(b*x-arctanh(tanh(b*x+a)))^2/b^3+(b*x-arctanh(tanh(b*x+a)))^3*ln(arctanh(tanh(b*x+a)))/b^4
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.98

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx$$

$$= \frac{x^3}{3b} - \frac{x^2(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{2b^2} + \frac{x(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}{b^3}$$

$$- \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

input `Integrate[x^3/ArcTanh[Tanh[a + b*x]], x]`

output `x^3/(3*b) - (x^2*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/(2*b^2) + (x*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)/b^3 - ((-(b*x) + ArcTanh[Tanh[a + b*x]])^3*Log[ArcTanh[Tanh[a + b*x]]])/b^4`

### Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2590, 2590, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx \\
 & \quad \downarrow \text{2590} \\
 & \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx + \frac{x^3}{3b} \\
 & \quad \downarrow \text{2590} \\
 & \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx + \frac{x^2}{2b} \right) + \frac{x^3}{3b} \\
 & \quad \downarrow \text{2589} \\
 & \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx + \frac{x}{b} \right) + \frac{x^2}{2b} \right) + \frac{x^3}{3b} \\
 & \quad \downarrow \text{2588} \\
 & \frac{x^3}{3b}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{b} \right)}{b} \\
 & \frac{x^3}{3b} \\
 & \quad \downarrow 14 \\
 & \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \log(\operatorname{arctanh}(\tanh(a + bx))) + \frac{x}{b}}{b^2} \right) + \frac{x^2}{2b}}{b} \right) + \frac{x^2}{2b}}{b} \\
 & \frac{x^3}{3b}
 \end{aligned}$$

input `Int[x^3/ArcTanh[Tanh[a + b*x]],x]`

output `x^3/(3*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x^2/(2*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2))/b))/b`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`



rule 2590

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^n/(a^n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x
] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n,
1]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 162 vs.  $2(77) = 154$ .

Time = 1.04 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.01

method	result
default	$\frac{b^2 x^3 - \frac{x^2 ab}{2} - \frac{b x^2 (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{2} + x a^2 + 2 x a (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + x (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^3} + \dots$
risch	Expression too large to display

input

```
int(x^3/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/b^3*(1/3*b^2*x^3-1/2*x^2*a*b-1/2*b*x^2*(arctanh(tanh(b*x+a))-b*x-a)+x*a^
2+2*x*a*(arctanh(tanh(b*x+a))-b*x-a)+x*(arctanh(tanh(b*x+a))-b*x-a)^2+(-a
^3-3*a^2*(arctanh(tanh(b*x+a))-b*x-a)-3*a*(arctanh(tanh(b*x+a))-b*x-a)^2-(
arctanh(tanh(b*x+a))-b*x-a)^3)/b^4*ln(arctanh(tanh(b*x+a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.51

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2b^3x^3 - 3ab^2x^2 + 6a^2bx - 6a^3 \log(bx + a)}{6b^4}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
1/6*(2*b^3*x^3 - 3*a*b^2*x^2 + 6*a^2*b*x - 6*a^3*log(b*x + a))/b^4
```

**Sympy [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^3}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**3/atanh(tanh(b*x+a)),x)`

output `Integral(x**3/atanh(tanh(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.52

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{a^3 \log(bx + a)}{b^4} + \frac{2b^2x^3 - 3abx^2 + 6a^2x}{6b^3}$$

input `integrate(x^3/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `-a^3*log(b*x + a)/b^4 + 1/6*(2*b^2*x^3 - 3*a*b*x^2 + 6*a^2*x)/b^3`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.53

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{a^3 \log(|bx + a|)}{b^4} + \frac{2b^2x^3 - 3abx^2 + 6a^2x}{6b^3}$$

input `integrate(x^3/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `-a^3*log(abs(b*x + a))/b^4 + 1/6*(2*b^2*x^3 - 3*a*b*x^2 + 6*a^2*x)/b^3`

**Mupad [B] (verification not implemented)**

Time = 3.40 (sec) , antiderivative size = 354, normalized size of antiderivative = 4.37

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{x^3}{3b} + \frac{x^2 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{4b^2}$$

$$+ \frac{x \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{4b^3}$$

$$+ \frac{\ln \left( \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) \right) \left( \left( 2a - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3 - 8a^3 - 6a \right)}{8b^4}$$

input `int(x^3/atanh(tanh(a + b*x)),x)`

output

```
x^3/(3*b) + (x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(4*b^2) + (x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^3) + (log(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(8*b^4)
```

**Reduce [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^3}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^3/atanh(tanh(b*x+a)),x)`output `int(x**3/atanh(tanh(a + b*x)),x)`

### 3.87 $\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	695
Mathematica [A] (verified)	695
Rubi [A] (verified)	696
Maple [A] (verified)	697
Fricas [A] (verification not implemented)	698
Sympy [F]	698
Maxima [A] (verification not implemented)	698
Giac [A] (verification not implemented)	699
Mupad [B] (verification not implemented)	699
Reduce [F]	700

#### Optimal result

Integrand size = 13, antiderivative size = 56

$$\begin{aligned} & \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx \\ &= \frac{x^2}{2b} + \frac{x(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^2} \\ & \quad + \frac{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^3} \end{aligned}$$

output

$1/2*x^2/b+x*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/b^2+(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2*\ln(\operatorname{arctanh}(\tanh(b*x+a)))/b^3$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\begin{aligned} & \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx \\ &= \frac{x^2}{2b} - \frac{x(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{b^2} \\ & \quad + \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^3} \end{aligned}$$

input `Integrate[x^2/ArcTanh[Tanh[a + b*x]], x]`

output  $x^2/(2*b) - (x*(-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]))/b^2 + ((-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]])^2*\text{Log}[\text{ArcTanh}[\text{Tanh}[a + b*x]]])/b^3$

### Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2590, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\text{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2590$$

$$\frac{(bx - \text{arctanh}(\tanh(a + bx))) \int \frac{x}{\text{arctanh}(\tanh(a + bx))} dx}{b} + \frac{x^2}{2b}$$

$$\downarrow 2589$$

$$\frac{(bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \text{arctanh}(\tanh(a + bx))) \int \frac{1}{\text{arctanh}(\tanh(a + bx))} dx}{b} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b}$$

$$\downarrow 2588$$

$$\frac{(bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \text{arctanh}(\tanh(a + bx))) \int \frac{1}{\text{arctanh}(\tanh(a + bx))} d\text{arctanh}(\tanh(a + bx))}{b^2} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b}$$

$$\downarrow 14$$

$$\frac{(bx - \text{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \text{arctanh}(\tanh(a + bx))) \log(\text{arctanh}(\tanh(a + bx)))}{b^2} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b}$$

input `Int[x^2/ArcTanh[Tanh[a + b*x]],x]`

output `x^2/(2*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2))/b`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

### Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.54

method	result
default	$\frac{\frac{x^2 b}{2} - ax - x(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{b^2} + \frac{(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2) \ln(\operatorname{arctanh}(\tanh(bx+a)))}{b^3}$
risch	Expression too large to display

input `int(x^2/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/b^2*(1/2*x^2*b-a*x-x*(arctanh(tanh(b*x+a))-b*x-a))+(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/b^3*ln(arctanh(tanh(b*x+a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.52

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{b^2 x^2 - 2 abx + 2 a^2 \log(bx + a)}{2 b^3}$$

input

```
integrate(x^2/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
1/2*(b^2*x^2 - 2*a*b*x + 2*a^2*log(b*x + a))/b^3
```

**Sympy [F]**

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^2}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input

```
integrate(x**2/atanh(tanh(b*x+a)),x)
```

output

```
Integral(x**2/atanh(tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.52

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{a^2 \log(bx + a)}{b^3} + \frac{bx^2 - 2 ax}{2 b^2}$$

input

```
integrate(x^2/arctanh(tanh(b*x+a)),x, algorithm="maxima")
```

output  $a^2 \log(bx + a)/b^3 + 1/2*(bx^2 - 2ax)/b^2$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.54

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{a^2 \log(|bx + a|)}{b^3} + \frac{bx^2 - 2ax}{2b^2}$$

input `integrate(x^2/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output  $a^2 \log(\operatorname{abs}(bx + a))/b^3 + 1/2*(bx^2 - 2ax)/b^2$

### Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 234, normalized size of antiderivative = 4.18

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{x^2}{2b} + \frac{x \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{2b^2}$$

$$+ \frac{\ln \left( \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) \right) \left( \left( 2a - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2 - 4a \left( 2a - \right) \right)}{4b^3}$$

input `int(x^2/atanh(tanh(a + b*x)),x)`

output  $x^2/(2*b) + (x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))/(2*b^2) + (\log(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))))*((2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))^2 - 4*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) + 4*a^2))/(4*b^3)$



**Reduce [F]**

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^2}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^2/atanh(tanh(b*x+a)),x)`

output `int(x**2/atanh(tanh(a + b*x)),x)`

### 3.88 $\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	701
Mathematica [A] (verified)	701
Rubi [A] (verified)	702
Maple [A] (verified)	703
Fricas [A] (verification not implemented)	703
Sympy [F]	704
Maxima [A] (verification not implemented)	704
Giac [A] (verification not implemented)	704
Mupad [B] (verification not implemented)	705
Reduce [F]	705

#### Optimal result

Integrand size = 11, antiderivative size = 31

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx$$

$$= \frac{x}{b} + \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2}$$

output `x/b+(b*x-arctanh(tanh(b*x+a)))*ln(arctanh(tanh(b*x+a)))/b^2`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx$$

$$= \frac{x}{b} - \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2}$$

input `Integrate[x/ArcTanh[Tanh[a + b*x]],x]`

output `x/b - ((-(b*x) + ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2`

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow \text{2589}$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{x}{b}$$

$$\downarrow \text{2588}$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{b^2} + \frac{x}{b}$$

$$\downarrow \text{14}$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \log(\operatorname{arctanh}(\tanh(a + bx)))}{b^2} + \frac{x}{b}$$

input `Int[x/ArcTanh[Tanh[a + b*x]],x]`

output `x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst [Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589

```
Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v,
x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u -
a*v, 0]] /; PiecewiseLinearQ[u, v, x]
```

**Maple [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

method	result	size
default	$\frac{x}{b} + \frac{(bx - \operatorname{arctanh}(\tanh(bx+a))) \ln(\operatorname{arctanh}(\tanh(bx+a)))}{b^2}$	32
risch	Expression too large to display	2837

input

```
int(x/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
x/b+(b*x-arctanh(tanh(b*x+a)))*ln(arctanh(tanh(b*x+a)))/b^2
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.55

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{bx - a \log(bx + a)}{b^2}$$

input

```
integrate(x/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
(b*x - a*log(b*x + a))/b^2
```

**Sympy [F]**

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x/atanh(tanh(b*x+a)),x)`

output `Integral(x/atanh(tanh(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.58

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{x}{b} - \frac{a \log(bx + a)}{b^2}$$

input `integrate(x/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `x/b - a*log(b*x + a)/b^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{x}{b} - \frac{a \log(|bx + a|)}{b^2}$$

input `integrate(x/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `x/b - a*log(abs(b*x + a))/b^2`

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 108, normalized size of antiderivative = 3.48

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{x}{b} + \frac{\ln\left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)\right) \left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{2b^2}$$

input `int(x/atanh(tanh(a + b*x)),x)`output `x/b + (log(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b^2)`**Reduce [F]**

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x/atanh(tanh(b*x+a)),x)`output `int(x/atanh(tanh(a + b*x)),x)`

$$3.89 \quad \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx$$

Optimal result	706
Mathematica [A] (verified)	706
Rubi [A] (verified)	707
Maple [A] (verified)	708
Fricas [A] (verification not implemented)	708
Sympy [A] (verification not implemented)	709
Maxima [A] (verification not implemented)	709
Giac [A] (verification not implemented)	709
Mupad [B] (verification not implemented)	710
Reduce [B] (verification not implemented)	710

### Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{b}$$

output `ln(arctanh(tanh(b*x+a)))/b`

### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{b}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(-1),x]`

output `Log[ArcTanh[Tanh[a + b*x]]]/b`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2588$$

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))$$

$$\frac{\phantom{\int} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow 14$$

$$\frac{\log(\operatorname{arctanh}(\tanh(a + bx)))}{b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(-1), x]`

output `Log[ArcTanh[Tanh[a + b*x]]]/b`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst [Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



**Maple [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result
derivativedivides	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{b}$
default	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{b}$
parallelrisch	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{b}$
risch	$\ln\left(\ln(e^{bx+a}) + \frac{i\pi\left(-\operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)\operatorname{csgn}(ie^{2bx+2a})\operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) + \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)\operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)\right)^2 - \operatorname{csgn}\right)}{b}$

input `int(1/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `ln(arctanh(tanh(b*x+a)))/b`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{\log(bx+a)}{b}$$

input `integrate(1/arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `log(b*x + a)/b`

**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx = \begin{cases} \frac{\log(\operatorname{atanh}(\tanh(a + bx)))}{b} & \text{for } b \neq 0 \\ \frac{x}{\operatorname{atanh}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a)),x)`output `Piecewise((log(atanh(tanh(a + b*x)))/b, Ne(b, 0)), (x/atanh(tanh(a)), True))`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\log(-bx - a)}{b}$$

input `integrate(1/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `log(-b*x - a)/b`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\log(|bx + a|)}{b}$$

input `integrate(1/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `log(abs(b*x + a))/b`

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\ln(\operatorname{atanh}(\tanh(a + bx)))}{b}$$

input `int(1/atanh(tanh(a + b*x)),x)`

output `log(atanh(tanh(a + b*x)))/b`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\log(\operatorname{atanh}(\tanh(bx + a)))}{b}$$

input `int(1/atanh(tanh(b*x+a)),x)`

output `log(atanh(tanh(a + b*x)))/b`

### 3.90 $\int \frac{1}{x \mathbf{arctanh}(\tanh(a+bx))} dx$

Optimal result	711
Mathematica [A] (verified)	711
Rubi [A] (verified)	712
Maple [A] (verified)	713
Fricas [A] (verification not implemented)	713
Sympy [F]	714
Maxima [A] (verification not implemented)	714
Giac [A] (verification not implemented)	714
Mupad [B] (verification not implemented)	715
Reduce [F]	715

#### Optimal result

Integrand size = 13, antiderivative size = 44

$$\int \frac{1}{x \mathbf{arctanh}(\tanh(a + bx))} dx = -\frac{\log(x)}{bx - \mathbf{arctanh}(\tanh(a + bx))} + \frac{\log(\mathbf{arctanh}(\tanh(a + bx)))}{bx - \mathbf{arctanh}(\tanh(a + bx))}$$

output `-ln(x)/(b*x-arctanh(tanh(b*x+a)))+ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.66

$$\int \frac{1}{x \mathbf{arctanh}(\tanh(a + bx))} dx = \frac{-\log(x) + \log(\mathbf{arctanh}(\tanh(a + bx)))}{bx - \mathbf{arctanh}(\tanh(a + bx))}$$

input `Integrate[1/(x*ArcTanh[Tanh[a + b*x]]),x]`

output `(-Log[x] + Log[ArcTanh[Tanh[a + b*x]]])/(b*x - ArcTanh[Tanh[a + b*x]])`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx \\
 & \quad \downarrow \text{2591} \\
 & \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} \\
 & \quad \downarrow \text{14} \\
 & \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \\
 & \quad \downarrow \text{2588} \\
 & \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \\
 & \quad \downarrow \text{14} \\
 & \frac{\log(\operatorname{arctanh}(\tanh(a + bx)))}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))}
 \end{aligned}$$

input `Int[1/(x*ArcTanh[Tanh[a + b*x]]),x]`

output `-(Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst  
[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D  
[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1  
/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 9.25 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.98

method	result
default	$\frac{\ln(x)}{\operatorname{arctanh}(\tanh(bx+a))-bx} - \frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{\operatorname{arctanh}(\tanh(bx+a))-bx}$
risch	$\frac{\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) - \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + \pi \operatorname{csgn}(ie^{2bx+2a})^3 - 2\pi \operatorname{csgn}(ie^{2bx+2a})}{\operatorname{arctanh}(\tanh(bx+a))-bx}$

input `int(1/x/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `1/(arctanh(tanh(b*x+a))-b*x)*ln(x)-1/(arctanh(tanh(b*x+a))-b*x)*ln(arctanh  
(tanh(b*x+a)))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.36

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{\log(bx+a) - \log(x)}{a}$$

input `integrate(1/x/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output  $-(\log(b*x + a) - \log(x))/a$

### Sympy [F]

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{x \operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(1/x/atanh(tanh(b*x+a)),x)`

output `Integral(1/(x*atanh(tanh(a + b*x))), x)`

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.41

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{\log(bx + a)}{a} + \frac{\log(x)}{a}$$

input `integrate(1/x/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output  $-\log(b*x + a)/a + \log(x)/a$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.45

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{\log(|bx + a|)}{a} + \frac{\log(|x|)}{a}$$

input `integrate(1/x/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output  $-\log(\operatorname{abs}(b*x + a))/a + \log(\operatorname{abs}(x))/a$

**Mupad [B] (verification not implemented)**

Time = 5.91 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.43

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{4 \operatorname{atanh}\left(\frac{4bx}{\ln\left(\frac{1}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx} - 1\right)}{\ln\left(\frac{1}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx}$$

input `int(1/(x*atanh(tanh(a + b*x))),x)`output `-(4*atanh((4*b*x)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 1))/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)`**Reduce [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a)) x} dx$$

input `int(1/x/atanh(tanh(b*x+a)),x)`output `int(1/(atanh(tanh(a + b*x))*x),x)`



### 3.91 $\int \frac{1}{x^2 \mathbf{arctanh}(\tanh(a+bx))} dx$

Optimal result	716
Mathematica [A] (verified)	716
Rubi [A] (verified)	717
Maple [A] (verified)	719
Fricas [A] (verification not implemented)	719
Sympy [F]	719
Maxima [A] (verification not implemented)	720
Giac [A] (verification not implemented)	720
Mupad [B] (verification not implemented)	720
Reduce [F]	721

#### Optimal result

Integrand size = 13, antiderivative size = 65

$$\int \frac{1}{x^2 \mathbf{arctanh}(\tanh(a + bx))} dx = \frac{1}{x(bx - \mathbf{arctanh}(\tanh(a + bx)))} - \frac{b \log(x)}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^2} + \frac{b \log(\mathbf{arctanh}(\tanh(a + bx)))}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^2}$$

output

```
1/x/(b*x-arctanh(tanh(b*x+a)))-b*ln(x)/(b*x-arctanh(tanh(b*x+a)))^2+b*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^2
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.69

$$\int \frac{1}{x^2 \mathbf{arctanh}(\tanh(a + bx))} dx = \frac{-\mathbf{arctanh}(\tanh(a + bx)) + bx(1 - \log(x) + \log(\mathbf{arctanh}(\tanh(a + bx))))}{x(-bx + \mathbf{arctanh}(\tanh(a + bx)))^2}$$

input

```
Integrate[1/(x^2*ArcTanh[Tanh[a + b*x]]),x]
```

output

```
(-ArcTanh[Tanh[a + b*x]] + b*x*(1 - Log[x] + Log[ArcTanh[Tanh[a + b*x]]]))
/(x*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.25, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx \\
 & \quad \downarrow 2594 \\
 & \frac{b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow 2591 \\
 & \frac{b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow 14 \\
 & \frac{b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow 2588 \\
 & \frac{b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \\
 & \quad \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\
 & \quad \downarrow 14
 \end{aligned}$$

$$\frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{b \left( \frac{\log(\operatorname{arctanh}(\tanh(a + bx)))}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))}$$

input `Int[1/(x^2*ArcTanh[Tanh[a + b*x]]),x]`

output `1/(x*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(-(Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

**Maple [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98

$$-\frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)x} - \frac{b \ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2} + \frac{b \ln(\operatorname{arctanh}(\tanh(bx+a)))}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2}$$

input `int(1/x^2/arctanh(tanh(b*x+a)),x)`output `-1/(arctanh(tanh(b*x+a))-b*x)/x-1/(arctanh(tanh(b*x+a))-b*x)^2*b*ln(x)+1/(arctanh(tanh(b*x+a))-b*x)^2*b*ln(arctanh(tanh(b*x+a)))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.40

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))} dx = \frac{bx \log(bx+a) - bx \log(x) - a}{a^2 x}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `(b*x*log(b*x + a) - b*x*log(x) - a)/(a^2*x)`**Sympy [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))} dx = \int \frac{1}{x^2 \operatorname{atanh}(\tanh(a+bx))} dx$$

input `integrate(1/x**2/atanh(tanh(b*x+a)),x)`output `Integral(1/(x**2*atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.43

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{b \log(bx + a)}{a^2} - \frac{b \log(x)}{a^2} - \frac{1}{ax}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `b*log(b*x + a)/a^2 - b*log(x)/a^2 - 1/(a*x)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.46

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{b \log(|bx + a|)}{a^2} - \frac{b \log(|x|)}{a^2} - \frac{1}{ax}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `b*log(abs(b*x + a))/a^2 - b*log(abs(x))/a^2 - 1/(a*x)`**Mupad [B] (verification not implemented)**

Time = 5.93 (sec) , antiderivative size = 210, normalized size of antiderivative = 3.23

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2 \ln\left(\frac{1}{e^{2a} e^{2bx+1}}\right) - 2 \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 4bx + bx \operatorname{atan}\left(\frac{-\ln\left(\frac{1}{e^{2a} e^{2bx+1}}\right) + \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + bx}{\ln\left(\frac{1}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx}\right)}{x \left(\ln\left(\frac{1}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx\right)^2} \quad 8i$$

input `int(1/(x^2*atanh(tanh(a + b*x))),x)`

output

```
(2*log(1/(exp(2*a)*exp(2*b*x) + 1)) - 2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x + b*x*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*8i)/(x*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)
```

**Reduce [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a)) x^2} dx$$

input

```
int(1/x^2/atanh(tanh(b*x+a)),x)
```

output

```
int(1/(atanh(tanh(a + b*x))*x**2),x)
```

### 3.92 $\int \frac{1}{x^3 \mathbf{arctanh}(\tanh(a+bx))} dx$

Optimal result	722
Mathematica [A] (verified)	723
Rubi [A] (verified)	723
Maple [A] (verified)	725
Fricas [A] (verification not implemented)	726
Sympy [F]	726
Maxima [A] (verification not implemented)	726
Giac [A] (verification not implemented)	727
Mupad [B] (verification not implemented)	727
Reduce [F]	728

#### Optimal result

Integrand size = 13, antiderivative size = 92

$$\int \frac{1}{x^3 \mathbf{arctanh}(\tanh(a + bx))} dx = \frac{b}{x(bx - \mathbf{arctanh}(\tanh(a + bx)))^2} + \frac{1}{2x^2(bx - \mathbf{arctanh}(\tanh(a + bx)))} - \frac{b^2 \log(x)}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^3} + \frac{b^2 \log(\mathbf{arctanh}(\tanh(a + bx)))}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^3}$$

output

```
b/x/(b*x-arctanh(tanh(b*x+a)))^2+1/2/x^2/(b*x-arctanh(tanh(b*x+a)))-b^2*ln(x)/(b*x-arctanh(tanh(b*x+a)))^3+b^2*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^3
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.72

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{-4bx \operatorname{arctanh}(\tanh(a + bx)) + \operatorname{arctanh}(\tanh(a + bx))^2 + b^2 x^2 (3 - 2 \log(x) + 2 \log(\operatorname{arctanh}(\tanh(a + bx))))}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx)))^3}$$

input

```
Integrate[1/(x^3*ArcTanh[Tanh[a + b*x]]),x]
```

output

```
(-4*b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2 + b^2*x^2*(3 - 2*Log[x] + 2*Log[ArcTanh[Tanh[a + b*x]]]))/(2*x^2*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2594, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2594$$

$$\frac{b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2594$$

$$\frac{b \left( \frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x (bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2591$$



$$\begin{aligned}
& b \left( \frac{b \int \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} dx - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow 14 \\
& b \left( \frac{b \int \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} dx - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow 2588 \\
& b \left( \frac{\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} d\operatorname{arctanh}(\tanh(a+bx)) - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow 14 \\
& \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \\
& b \left( \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}
\end{aligned}$$

input

```
Int [1/(x^3*ArcTanh[Tanh[a + b*x]]), x]
```

output

```
1/(2*x^2*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(1/(x*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(-(Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])
```

### Defintions of rubi rules used

rule 14

```
Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]
```

rule 2588

```
Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]
```

rule 2591

```
Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

### Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.95

$$-\frac{1}{2(\operatorname{arctanh}(\tanh(bx+a)) - bx)x^2} + \frac{b^2 \ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3} + \frac{b}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2}$$

input

```
int(1/x^3/arctanh(tanh(b*x+a)),x)
```

output

```
-1/2/(arctanh(tanh(b*x+a))-b*x)/x^2+1/(arctanh(tanh(b*x+a))-b*x)^3*b^2*ln(x)+1/(arctanh(tanh(b*x+a))-b*x)^2*b/x-1/(arctanh(tanh(b*x+a))-b*x)^3*b^2*ln(arctanh(tanh(b*x+a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b^2x^2 \log(bx + a) - 2b^2x^2 \log(x) - 2abx + a^2}{2a^3x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `-1/2*(2*b^2*x^2*log(b*x + a) - 2*b^2*x^2*log(x) - 2*a*b*x + a^2)/(a^3*x^2)`**Sympy [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{x^3 \operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(1/x**3/atanh(tanh(b*x+a)),x)`output `Integral(1/(x**3*atanh(tanh(a + b*x))), x)`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.43

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{b^2 \log(bx + a)}{a^3} + \frac{b^2 \log(x)}{a^3} + \frac{2bx - a}{2a^2x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-b^2*log(b*x + a)/a^3 + b^2*log(x)/a^3 + 1/2*(2*b*x - a)/(a^2*x^2)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.49

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{b^2 \log(|bx + a|)}{a^3} + \frac{b^2 \log(|x|)}{a^3} + \frac{2abx - a^2}{2a^3x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `-b^2*log(abs(b*x + a))/a^3 + b^2*log(abs(x))/a^3 + 1/2*(2*a*b*x - a^2)/(a^3*x^2)`

**Mupad [B] (verification not implemented)**

Time = 5.66 (sec) , antiderivative size = 286, normalized size of antiderivative = 3.11

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)^2 - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) \left(2 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) + 8bx\right) + \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)^2 + 12b^2x^2 + 8bx \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{x^2 \left(\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}$$

input `int(1/(x^3*atanh(tanh(a + b*x))),x)`

output `(log(1/(exp(2*a)*exp(2*b*x) + 1)))^2 - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(2*log(1/(exp(2*a)*exp(2*b*x) + 1)) + 8*b*x) + log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 12*b^2*x^2 + 8*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1)) + b^2*x^2*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*16i)/(x^2*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))^3`

Reduce [F]

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a)) x^3} dx$$

input `int(1/x^3/atanh(tanh(b*x+a)),x)`

output `int(1/(atanh(tanh(a + b*x))*x**3),x)`

### 3.93 $\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	729
Mathematica [A] (verified)	729
Rubi [A] (verified)	730
Maple [F]	731
Fricas [F]	731
Sympy [F]	732
Maxima [F]	732
Giac [F]	732
Mupad [F(-1)]	733
Reduce [F]	733

#### Optimal result

Integrand size = 13, antiderivative size = 65

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$= -\frac{x^m}{b \operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{x^m \operatorname{Hypergeometric2F1}\left(1, m, 1+m, \frac{bx}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{b(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

```
output -x^m/b/arctanh(tanh(b*x+a))-x^m*hypergeom([1, m],[1+m],b*x/(b*x-arctanh(tanh(b*x+a))))/b/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.78

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$= \frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(2, 1+m, 2+m, -\frac{bx}{-bx + \operatorname{arctanh}(\tanh(a+bx))}\right)}{(1+m)(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input `Integrate[x^m/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(x^(1 + m)*Hypergeometric2F1[2, 1 + m, 2 + m, -((b*x)/(-b*x) + ArcTanh[Tanh[a + b*x]])])/((1 + m)*(-b*x) + ArcTanh[Tanh[a + b*x]]^2)`

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow \text{2599}$$

$$\frac{m \int \frac{x^{m-1}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} - \frac{x^m}{b \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow \text{2595}$$

$$-\frac{x^m \operatorname{Hypergeometric2F1}\left(1, m, m + 1, \frac{bx}{bx - \operatorname{arctanh}(\tanh(a + bx))}\right)}{b(bx - \operatorname{arctanh}(\tanh(a + bx)))} - \frac{x^m}{b \operatorname{arctanh}(\tanh(a + bx))}$$

input `Int[x^m/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(x^m/(b*ArcTanh[Tanh[a + b*x]])) - (x^m*Hypergeometric2F1[1, m, 1 + m, (b*x)/(b*x - ArcTanh[Tanh[a + b*x]])]/(b*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2595 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(v^(n + 1)/((n + 1)*(b*u - a*v))*Hypergeometric2F1[1, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && !IntegerQ[n]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(bx + a))^2} dx$$

input `int(x^m/arctanh(tanh(b*x+a))^2,x)`

output `int(x^m/arctanh(tanh(b*x+a))^2,x)`

**Fricas [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^2} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `integral(x^m/arctanh(tanh(b*x + a))^2, x)`



**Sympy [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^m}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**m/atanh(tanh(b*x+a))**2,x)`

output `Integral(x**m/atanh(tanh(a + b*x))**2, x)`

**Maxima [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^2} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `integrate(x^m/arctanh(tanh(b*x + a))^2, x)`

**Giac [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^2} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `integrate(x^m/arctanh(tanh(b*x + a))^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^m}{\operatorname{atanh}(\tanh(a + bx))^2} dx$$

input `int(x^m/atanh(tanh(a + b*x))^2,x)`output `int(x^m/atanh(tanh(a + b*x))^2, x)`**Reduce [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\operatorname{atanh}(\tanh(bx + a)) \left( \int \frac{x^m}{\operatorname{atanh}(\tanh(bx+a))x} dx \right) m - x^m}{\operatorname{atanh}(\tanh(bx + a)) b}$$

input `int(x^m/atanh(tanh(b*x+a))^2,x)`output `(atanh(tanh(a + b*x))*int(x**m/(atanh(tanh(a + b*x))*x),x)*m - x**m)/(atanh(tanh(a + b*x))*b)`

### 3.94 $\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	734
Mathematica [A] (verified)	735
Rubi [A] (verified)	735
Maple [B] (verified)	738
Fricas [A] (verification not implemented)	739
Sympy [F]	739
Maxima [A] (verification not implemented)	739
Giac [A] (verification not implemented)	740
Mupad [B] (verification not implemented)	740
Reduce [F]	741

#### Optimal result

Integrand size = 13, antiderivative size = 98

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$= \frac{4x^3}{3b^2} + \frac{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^3}$$

$$+ \frac{4x(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{b^4} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^5}$$

output

```
4/3*x^3/b^2+2*x^2*(b*x-arctanh(tanh(b*x+a)))/b^3+4*x*(b*x-arctanh(tanh(b*x+a)))^2/b^4-x^4/b/arctanh(tanh(b*x+a))+4*(b*x-arctanh(tanh(b*x+a)))^3*ln(arctanh(tanh(b*x+a)))/b^5
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.08

$$\begin{aligned} & \int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx \\ &= \frac{x^3}{3b^2} - \frac{x^2(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{b^3} \\ &+ \frac{3x(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}{b^4} - \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^4}{b^5 \operatorname{arctanh}(\tanh(a+bx))} \\ &- \frac{4(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^5} \end{aligned}$$

input `Integrate[x^4/ArcTanh[Tanh[a + b*x]]^2,x]`

output  $x^3/(3*b^2) - (x^2*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]))/b^3 + (3*x*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2)/b^4 - (-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^4/(b^5 * \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])) - (4*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3 * \operatorname{Log}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]])/b^5$

**Rubi [A] (verified)**

Time = 0.61 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2599, 2590, 2590, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^2} dx \\ & \quad \downarrow \text{2599} \\ & \frac{4 \int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))} \\ & \quad \downarrow \text{2590} \end{aligned}$$

$$\begin{aligned}
 & \frac{4 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{x^3}{3b}}{b} \right)}{b} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2590} \\
 & \frac{4 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{x^2}{2b}}{b} \right) + \frac{x^3}{3b}}{b} \right)}{b} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2589} \\
 & \frac{4 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{x}{b}}{b} \right) + \frac{x^2}{2b}}{b} \right) + \frac{x^3}{3b}}{b} \right)}{b} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2588} \\
 & \frac{4 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d \operatorname{arctanh}(\tanh(a+bx))}{b^2} + \frac{\operatorname{arctanh}(\tanh(a+bx))}{b} \right) + \frac{x^2}{2b}}{b} \right) + \frac{x^3}{3b}}{b} \right)}{b} - \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{14} \\
 & \frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))}
 \end{aligned}$$

$$4 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx))) + \frac{x}{b}}{b^2} + \frac{x^2}{2b} \right)}{b} \right) + \frac{x^3}{3b} \right)$$


---


$$\frac{x^4}{b \operatorname{arctanh}(\tanh(a+bx))}$$

input `Int [x^4/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(x^4/(b*ArcTanh[Tanh[a + b*x]])) + (4*(x^3/(3*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x^2/(2*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]]/b^2))/b))/b)/b`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n-1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 261 vs.  $2(96) = 192$ .

Time = 1.22 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.67

method	result
default	$\frac{b^2 x^3 - x^2 a b - b x^2 (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3x a^2 + 6xa (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3x (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^4}$
risch	Expression too large to display

input

```
int(x^4/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
1/b^4*(1/3*b^2*x^3-x^2*a*b-b*x^2*(arctanh(tanh(b*x+a))-b*x-a)+3*x*a^2+6*x*a*(arctanh(tanh(b*x+a))-b*x-a)+3*x*(arctanh(tanh(b*x+a))-b*x-a)^2)-(a^4+4*a^3*(arctanh(tanh(b*x+a))-b*x-a)+6*a^2*(arctanh(tanh(b*x+a))-b*x-a)^2+4*a*(arctanh(tanh(b*x+a))-b*x-a)^3+(arctanh(tanh(b*x+a))-b*x-a)^4)/b^5/arctanh(tanh(b*x+a))+(-4*a^3-12*a^2*(arctanh(tanh(b*x+a))-b*x-a)-12*a*(arctanh(tanh(b*x+a))-b*x-a)^2-4*(arctanh(tanh(b*x+a))-b*x-a)^3)/b^5*ln(arctanh(tanh(b*x+a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^4 x^4 - 2ab^3 x^3 + 6a^2 b^2 x^2 + 9a^3 bx - 3a^4 - 12(a^3 bx + a^4) \log(bx + a)}{3(b^6 x + ab^5)}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`output `1/3*(b^4*x^4 - 2*a*b^3*x^3 + 6*a^2*b^2*x^2 + 9*a^3*b*x - 3*a^4 - 12*(a^3*b*x + a^4)*log(b*x + a))/(b^6*x + a*b^5)`**Sympy [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^4}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**4/atanh(tanh(b*x+a))**2,x)`output `Integral(x**4/atanh(tanh(a + b*x))**2, x)`**Maxima [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.71

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^4 x^4 - 2ab^3 x^3 + 6a^2 b^2 x^2 + 9a^3 bx - 3a^4}{3(b^6 x + ab^5)} - \frac{4a^3 \log(bx + a)}{b^5}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`



output  $1/3*(b^4*x^4 - 2*a*b^3*x^3 + 6*a^2*b^2*x^2 + 9*a^3*b*x - 3*a^4)/(b^6*x + a*b^5) - 4*a^3*log(b*x + a)/b^5$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{4a^3 \log(|bx + a|)}{b^5} - \frac{a^4}{(bx + a)b^5} + \frac{b^4x^3 - 3ab^3x^2 + 9a^2b^2x}{3b^6}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output  $-4*a^3*log(abs(b*x + a))/b^5 - a^4/((b*x + a)*b^5) + 1/3*(b^4*x^3 - 3*a*b^3*x^2 + 9*a^2*b^2*x)/b^6$

### Mupad [B] (verification not implemented)

Time = 3.26 (sec) , antiderivative size = 669, normalized size of antiderivative = 6.83

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(x^4/atanh(tanh(a + b*x))^2,x)`

output

```

x^3/(3*b^2) - ((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 + 24*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 16*a^4 - 8*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 32*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/(2*b*(8*a*b^4 + 8*b^5*x - 4*b^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))) + (x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b^3) + (3*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^4) + (log(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b^5)

```

**Reduce [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^4}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input

```
int(x^4/atanh(tanh(b*x+a))^2,x)
```

output

```
int(x**4/atanh(tanh(a + b*x))**2,x)
```

### 3.95 $\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	742
Mathematica [A] (verified)	742
Rubi [A] (verified)	743
Maple [B] (verified)	745
Fricas [A] (verification not implemented)	746
Sympy [F]	746
Maxima [A] (verification not implemented)	746
Giac [A] (verification not implemented)	747
Mupad [B] (verification not implemented)	747
Reduce [F]	748

#### Optimal result

Integrand size = 13, antiderivative size = 75

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{3x^2}{2b^2} + \frac{3x(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^3} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} + \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

output

```
3/2*x^2/b^2+3*x*(b*x-arctanh(tanh(b*x+a)))/b^3-x^3/b/arctanh(tanh(b*x+a))+
3*(b*x-arctanh(tanh(b*x+a)))^2*ln(arctanh(tanh(b*x+a)))/b^4
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.11

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{x^2}{2b^2} - \frac{2x(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{b^3} + \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3}{b^4 \operatorname{arctanh}(\tanh(a+bx))} + \frac{3(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

input `Integrate[x^3/ArcTanh[Tanh[a + b*x]]^2,x]`

output `x^2/(2*b^2) - (2*x*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/b^3 + (-(b*x) + ArcTanh[Tanh[a + b*x]])^3/(b^4*ArcTanh[Tanh[a + b*x]]) + (3*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[ArcTanh[Tanh[a + b*x]]])/b^4`

### Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2590, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^2} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{3 \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2590} \\
 & \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2589} \\
 & \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2588} \\
 & \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))}
 \end{aligned}$$

$$\begin{array}{c}
 3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} \operatorname{arctanh}(\tanh(a+bx))}{b^2} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right) \\
 \hline
 \frac{x^3 b}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 \downarrow 14 \\
 3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx))) + \frac{x}{b}}{b^2} + \frac{x^2}{2b} \right)}{b} + \frac{x^2}{2b} \right) \\
 \hline
 \frac{b}{x^3} \\
 \frac{b}{b \operatorname{arctanh}(\tanh(a+bx))}
 \end{array}$$

input `Int [x^3/ArcTanh [Tanh [a + b*x]]^2,x]`

output `-(x^3/(b*ArcTanh [Tanh [a + b*x]])) + (3*(x^2/(2*b) + ((b*x - ArcTanh [Tanh [a + b*x]])*(x/b + ((b*x - ArcTanh [Tanh [a + b*x]])*Log [ArcTanh [Tanh [a + b*x]]])/b^2))/b)/b`

### Defintions of rubi rules used

rule 14 `Int [(a_)/(x_), x_Symbol] :> Simp [a*Log [x], x] /; FreeQ [a, x]`

rule 2588 `Int [(u_)^(m_), x_Symbol] :> With [{c = Simplify [D [u, x]]}, Simp [1/c Subst [Int [x^m, x], x, u], x]] /; FreeQ [m, x] && PiecewiseLinearQ [u, x]`

rule 2589 `Int [(v_)/(u_), x_Symbol] :> With [{a = Simplify [D [u, x]], b = Simplify [D [v, x]]}, Simp [b*(x/a), x] - Simp [(b*u - a*v)/a Int [1/u, x], x] /; NeQ [b*u - a*v, 0] /; PiecewiseLinearQ [u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs.  $2(73) = 146$ .

Time = 0.34 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.25

method	result
default	$\frac{\frac{x^2 b}{2} - 2ax - 2x(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{b^3} - \frac{-a^3 - 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) - 3a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 - (a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a))^3}{b^4 \operatorname{arctanh}(\tanh(bx+a))}$
risch	Expression too large to display

input `int(x^3/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{b^3} \left( \frac{1}{2} x^2 b - 2ax - 2x(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) - (-a^3 - 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) - 3a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 - (a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a))^3) / b^4 \operatorname{arctanh}(\tanh(bx+a)) + (3a^2 + 6a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2) / b^4 \ln(\operatorname{arctanh}(\tanh(bx+a))) \right)$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.83

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^3 x^3 - 3ab^2 x^2 - 4a^2 bx + 2a^3 + 6(a^2 bx + a^3) \log(bx + a)}{2(b^5 x + ab^4)}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`output `1/2*(b^3*x^3 - 3*a*b^2*x^2 - 4*a^2*b*x + 2*a^3 + 6*(a^2*b*x + a^3)*log(b*x + a))/(b^5*x + a*b^4)`**Sympy [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^3}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**3/atanh(tanh(b*x+a))**2,x)`output `Integral(x**3/atanh(tanh(a + b*x))**2, x)`**Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.79

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^3 x^3 - 3ab^2 x^2 - 4a^2 bx + 2a^3}{2(b^5 x + ab^4)} + \frac{3a^2 \log(bx + a)}{b^4}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `1/2*(b^3*x^3 - 3*a*b^2*x^2 - 4*a^2*b*x + 2*a^3)/(b^5*x + a*b^4) + 3*a^2*log(b*x + a)/b^4`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.64

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{3a^2 \log(|bx + a|)}{b^4} + \frac{a^3}{(bx + a)b^4} + \frac{b^2 x^2 - 4abx}{2b^4}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `3*a^2*log(abs(b*x + a))/b^4 + a^3/((b*x + a)*b^4) + 1/2*(b^2*x^2 - 4*a*b*x)/b^4`**Mupad [B] (verification not implemented)**

Time = 3.33 (sec) , antiderivative size = 490, normalized size of antiderivative = 6.53

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{x^2}{2b^2} + \frac{\ln\left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)\right) \left(3\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2 - 12a\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)\right)}{4b^4} - \frac{\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^3 - 8a^3 - 6a\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{4b\left(2ab^3 + 2b^4x - b^3\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)\right)} + \frac{x\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{b^3}$$

input `int(x^3/atanh(tanh(a + b*x))^2,x)`



output

```

x^2/(2*b^2) + (log(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
- log(2/(exp(2*a)*exp(2*b*x) + 1)))*(3*(2*a - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 -
12*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(
2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 12*a^2))/(4*b^4) - ((2*a - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*
x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(
2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a
^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(
exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(4*b*(2*a*b^3 + 2*b^4*x - b^3*(2*a - 1
og((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*ex
p(2*b*x) + 1)) + 2*b*x))) + (x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b^3

```

**Reduce [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^3}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input

```
int(x^3/atanh(tanh(b*x+a))^2,x)
```

output

```
int(x**3/atanh(tanh(a + b*x))**2,x)
```

### 3.96 $\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	749
Mathematica [A] (verified)	749
Rubi [A] (verified)	750
Maple [A] (verified)	751
Fricas [A] (verification not implemented)	752
Sympy [F]	752
Maxima [A] (verification not implemented)	753
Giac [A] (verification not implemented)	753
Mupad [B] (verification not implemented)	753
Reduce [F]	754

#### Optimal result

Integrand size = 13, antiderivative size = 50

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{2x}{b^2} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} + \frac{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^3}$$

output `2*x/b^2-x^2/b/arctanh(tanh(b*x+a))+2*(b*x-arctanh(tanh(b*x+a)))*ln(arctanh(tanh(b*x+a)))/b^3`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{bx - \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}{\operatorname{arctanh}(\tanh(a+bx))} + 2(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^3}$$

input `Integrate[x^2/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(b*x - (-(b*x) + ArcTanh[Tanh[a + b*x]])^2/ArcTanh[Tanh[a + b*x]] + 2*(b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^3`

### Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{2 \int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2589} \\
 & \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2588} \\
 & \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d \operatorname{arctanh}(\tanh(a+bx))}{b^2} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{14} \\
 & \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))}
 \end{aligned}$$

input `Int[x^2/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(x^2/(b*ArcTanh[Tanh[a + b*x]])) + (2*(x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2))/b`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :=> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_), x_Symbol] :=> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :=> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] :=> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.72

method	result
default	$\frac{x}{b^2} - \frac{a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^3 \operatorname{arctanh}(\tanh(bx+a))} + \frac{(2bx - 2 \operatorname{arctanh}(\tanh(bx+a))) \ln(\operatorname{arctanh}(\tanh(bx+a)))}{b^3}$
risch	Expression too large to display

input `int(x^2/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output `x/b^2-(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/b^3/arctanh(tanh(b*x+a))+(2*b*x-2*arctanh(tanh(b*x+a)))/b^3*ln(arctanh(tanh(b*x+a)))`

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^2 x^2 + abx - a^2 - 2(abx + a^2) \log(bx + a)}{b^4 x + ab^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `(b^2*x^2 + a*b*x - a^2 - 2*(a*b*x + a^2)*log(b*x + a))/(b^4*x + a*b^3)`

### Sympy [F]

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^2}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**2/atanh(tanh(b*x+a))**2,x)`

output `Integral(x**2/atanh(tanh(a + b*x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{b^2 x^2 + abx - a^2}{b^4 x + ab^3} - \frac{2a \log(bx + a)}{b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `(b^2*x^2 + a*b*x - a^2)/(b^4*x + a*b^3) - 2*a*log(b*x + a)/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{x}{b^2} - \frac{2a \log(|bx + a|)}{b^3} - \frac{a^2}{(bx + a)b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `x/b^2 - 2*a*log(abs(b*x + a))/b^3 - a^2/((b*x + a)*b^3)`**Mupad [B] (verification not implemented)**

Time = 3.32 (sec) , antiderivative size = 302, normalized size of antiderivative = 6.04

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{x}{b^2} - \frac{\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2 - 4a\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{2b\left(2ab^2 + 2b^3x - b^2\left(2a - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)\right)} + \frac{\ln\left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)\right)\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{b^3}$$

input `int(x^2/atanh(tanh(a + b*x))^2,x)`

output `x/b^2 - ((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 4*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 4*a^2)/(2*b*(2*a*b^2 + 2*b^3*x - b^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))) + (log(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b^3`

### Reduce [F]

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^2}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^2/atanh(tanh(b*x+a))^2,x)`

output `int(x**2/atanh(tanh(a + b*x))**2,x)`

### 3.97 $\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result . . . . .	755
Mathematica [A] (verified) . . . . .	755
Rubi [A] (verified) . . . . .	756
Maple [A] (verified) . . . . .	757
Fricas [A] (verification not implemented) . . . . .	757
Sympy [B] (verification not implemented) . . . . .	758
Maxima [A] (verification not implemented) . . . . .	758
Giac [A] (verification not implemented) . . . . .	759
Mupad [B] (verification not implemented) . . . . .	759
Reduce [B] (verification not implemented) . . . . .	759

#### Optimal result

Integrand size = 11, antiderivative size = 28

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{x}{b \operatorname{arctanh}(\tanh(a+bx))} + \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2}$$

output `-x/b/arctanh(tanh(b*x+a))+ln(arctanh(tanh(b*x+a)))/b^2`

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{1 - \frac{bx}{\operatorname{arctanh}(\tanh(a+bx))} + \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2}$$

input `Integrate[x/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(1 - (b*x)/ArcTanh[Tanh[a + b*x]] + Log[ArcTanh[Tanh[a + b*x]]])/b^2`



**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow \text{2599}$$

$$\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} - \frac{x}{b \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow \text{2588}$$

$$\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} d \operatorname{arctanh}(\tanh(a + bx))}{b^2} - \frac{x}{b \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow \text{14}$$

$$\frac{\log(\operatorname{arctanh}(\tanh(a + bx)))}{b^2} - \frac{x}{b \operatorname{arctanh}(\tanh(a + bx))}$$

input `Int [x/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(x/(b*ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/b^2`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst [Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

method	result
parallelrisch	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a)) - bx}{b^2 \operatorname{arctanh}(\tanh(bx+a))}$
default	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{b^2} - \frac{bx - \operatorname{arctanh}(\tanh(bx+a))}{b^2 \operatorname{arctanh}(\tanh(bx+a))}$
risch	$b \left( -\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) + \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 - \pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) \right)$

input `int(x/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`output `(ln(arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))-b*x)/b^2/arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{(bx + a) \log(bx + a) + a}{b^3x + ab^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output  $((b*x + a)*\log(b*x + a) + a)/(b^3*x + a*b^2)$

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs.  $2(24) = 48$ .

Time = 13.32 (sec) , antiderivative size = 94, normalized size of antiderivative = 3.36

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \begin{cases} \frac{x^2}{2 \operatorname{atanh}^2(\tanh(a))} & \text{for } b = 0 \\ \frac{x^2}{2 \operatorname{atanh}^2(\tanh(bx + \log(-e^{-bx})))} & \text{for } a = \log(-e^{-bx}) \\ \frac{x^2}{2 \operatorname{atanh}^2(\tanh(bx + \log(e^{-bx})))} & \text{for } a = \log(e^{-bx}) \\ -\frac{x}{b \operatorname{atanh}(\tanh(a + bx))} + \frac{\log(\operatorname{atanh}(\tanh(a + bx)))}{b^2} & \text{otherwise} \end{cases}$$

input `integrate(x/atanh(tanh(b*x+a))**2,x)`

output `Piecewise((x**2/(2*atanh(tanh(a))**2), Eq(b, 0)), (x**2/(2*atanh(tanh(b*x + log(-exp(-b*x))))**2), Eq(a, log(-exp(-b*x)))), (x**2/(2*atanh(tanh(b*x + log(exp(-b*x))))**2), Eq(a, log(exp(-b*x)))), (-x/(b*atanh(tanh(a + b*x))) + log(atanh(tanh(a + b*x)))/b**2, True))`

### Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{a}{b^3x + ab^2} + \frac{\log(bx + a)}{b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output  $a/(b^3*x + a*b^2) + \log(b*x + a)/b^2$

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\log(|bx + a|)}{b^2} + \frac{a}{(bx + a)b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `log(abs(b*x + a))/b^2 + a/((b*x + a)*b^2)`

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\ln(\operatorname{atanh}(\tanh(a + bx)))}{b^2} - \frac{x}{b \operatorname{atanh}(\tanh(a + bx))}$$

input `int(x/atanh(tanh(a + b*x))^2,x)`

output `log(atanh(tanh(a + b*x)))/b^2 - x/(b*atanh(tanh(a + b*x)))`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\operatorname{atanh}(\tanh(bx + a)) \log(\operatorname{atanh}(\tanh(bx + a))) - bx}{\operatorname{atanh}(\tanh(bx + a)) b^2}$$

input `int(x/atanh(tanh(b*x+a))^2,x)`

output `(atanh(tanh(a + b*x))*log(atanh(tanh(a + b*x))) - b*x)/(atanh(tanh(a + b*x))*b**2)`

### 3.98 $\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	760
Mathematica [A] (verified)	760
Rubi [A] (verified)	761
Maple [A] (verified)	762
Fricas [A] (verification not implemented)	762
Sympy [B] (verification not implemented)	762
Maxima [A] (verification not implemented)	763
Giac [A] (verification not implemented)	763
Mupad [B] (verification not implemented)	764
Reduce [B] (verification not implemented)	764

#### Optimal result

Integrand size = 9, antiderivative size = 14

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{1}{b \operatorname{arctanh}(\tanh(a+bx))}$$

output `-1/b/arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{1}{b \operatorname{arctanh}(\tanh(a+bx))}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(-2), x]`

output `-(1/(b*ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

↓ 2588

$$\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

↓ 15

$$-\frac{1}{b\operatorname{arctanh}(\tanh(a + bx))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(-2), x]`

output `-(1/(b*ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result
derivativedivides	$-\frac{1}{b \operatorname{arctanh}(\tanh(bx+a))}$
default	$-\frac{1}{b \operatorname{arctanh}(\tanh(bx+a))}$
parallelrisc	$-\frac{1}{b \operatorname{arctanh}(\tanh(bx+a))}$
risc	$b \left( -\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) + \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 - \pi \operatorname{csgn}(ie^{bx+a}) \right)$

input `int(1/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`output `-1/b/arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{1}{b^2x+ab}$$

input `integrate(1/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`output `-1/(b^2*x + a*b)`**Sympy [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(12) = 24$ .

Time = 13.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 5.07

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \begin{cases} \frac{x}{\operatorname{atanh}^2(\tanh(a))} & \text{for } b = 0 \\ \frac{x}{\operatorname{atanh}^2(\tanh(bx + \log(-e^{-bx})))} & \text{for } a = \log(-e^{-bx}) \\ \frac{x}{\operatorname{atanh}^2(\tanh(bx + \log(e^{-bx})))} & \text{for } a = \log(e^{-bx}) \\ -\frac{1}{b \operatorname{atanh}(\tanh(a + bx))} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a))**2,x)`

output `Piecewise((x/atanh(tanh(a))**2, Eq(b, 0)), (x/atanh(tanh(b*x + log(-exp(-b*x))))**2, Eq(a, log(-exp(-b*x)))), (x/atanh(tanh(b*x + log(exp(-b*x))))**2, Eq(a, log(exp(-b*x)))), (-1/(b*atanh(tanh(a + b*x))), True))`

### Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{1}{(bx + a)b}$$

input `integrate(1/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `-1/((b*x + a)*b)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{1}{(bx + a)b}$$

input `integrate(1/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`



output `-1/((b*x + a)*b)`

### Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{1}{b \operatorname{atanh}(\tanh(a + bx))}$$

input `int(1/atanh(tanh(a + b*x))^2,x)`

output `-1/(b*atanh(tanh(a + b*x)))`

### Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{1}{\operatorname{atanh}(\tanh(bx + a)) b}$$

input `int(1/atanh(tanh(b*x+a))^2,x)`

output `( - 1)/(atanh(tanh(a + b*x))*b)`

### 3.99 $\int \frac{1}{x \mathbf{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	765
Mathematica [A] (verified)	766
Rubi [A] (verified)	766
Maple [A] (verified)	768
Fricas [A] (verification not implemented)	768
Sympy [F]	769
Maxima [A] (verification not implemented)	769
Giac [A] (verification not implemented)	769
Mupad [B] (verification not implemented)	770
Reduce [F]	770

#### Optimal result

Integrand size = 13, antiderivative size = 70

$$\int \frac{1}{x \mathbf{arctanh}(\tanh(a + bx))^2} dx =$$

$$\frac{1}{(bx - \mathbf{arctanh}(\tanh(a + bx))) \mathbf{arctanh}(\tanh(a + bx))}$$

$$+ \frac{\log(x)}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^2}$$

$$- \frac{\log(\mathbf{arctanh}(\tanh(a + bx)))}{(bx - \mathbf{arctanh}(\tanh(a + bx)))^2}$$

output

```
-1/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))+ln(x)/(b*x-arctanh(tanh
(b*x+a)))^2-ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^2
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.76

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \frac{-bx + \operatorname{arctanh}(\tanh(a + bx))(1 + \log(bx) - \log(\operatorname{arctanh}(\tanh(a + bx))))}{\operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Integrate[1/(x*ArcTanh[Tanh[a + b*x]]^2),x]`

output `(-(b*x) + ArcTanh[Tanh[a + b*x]]*(1 + Log[b*x] - Log[ArcTanh[Tanh[a + b*x]]]))/(ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.26, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow 2594$$

$$\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow 2591$$

$$\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} -$$

$$\frac{1}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow 14$$

$$\begin{aligned}
& \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))} \\
& \quad \downarrow \text{2588} \\
& \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))} \\
& \quad \downarrow \text{14} \\
& \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
& \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}
\end{aligned}$$

input `Int[1/(x*ArcTanh[Tanh[a + b*x]]^2), x]`

output `-(1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (- (Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n +
1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew
iseLinearQ[u, v, x] && LtQ[n, -1]
```

**Maple [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.96

$$\frac{\ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2} - \frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2} + \frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx) \operatorname{arctanh}(\tanh(bx+a))}$$

input

```
int(1/x/arctanh(tanh(b*x+a))^2,x)
```

output

```
1/(arctanh(tanh(b*x+a))-b*x)^2*ln(x)-1/(arctanh(tanh(b*x+a))-b*x)^2*ln(arc
tanh(tanh(b*x+a)))+1/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.56

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{(bx+a) \log(bx+a) - (bx+a) \log(x) - a}{a^2 bx + a^3}$$

input

```
integrate(1/x/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
-((b*x + a)*log(b*x + a) - (b*x + a)*log(x) - a)/(a^2*b*x + a^3)
```

**Sympy [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{x \operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(1/x/atanh(tanh(b*x+a))**2,x)`

output `Integral(1/(x*atanh(tanh(a + b*x))**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.40

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{1}{abx + a^2} - \frac{\log(bx + a)}{a^2} + \frac{\log(x)}{a^2}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `1/(a*b*x + a^2) - log(b*x + a)/a^2 + log(x)/a^2`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.44

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{\log(|bx + a|)}{a^2} + \frac{\log(|x|)}{a^2} + \frac{1}{(bx + a)a}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `-log(abs(b*x + a))/a^2 + log(abs(x))/a^2 + 1/((b*x + a)*a)`

**Mupad [B] (verification not implemented)**

Time = 5.44 (sec) , antiderivative size = 359, normalized size of antiderivative = 5.13

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \frac{8bx - \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \left(-4 + \operatorname{atan}\left(\frac{-\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) 1i + \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) 1i + bx 2i}{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx}\right) 8i\right) + \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \left(-4 + \right)}{\left(\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)\right) \left(\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)\right)}$$

input `int(1/(x*atanh(tanh(a + b*x))^2),x)`output `(8*b*x - log(1/(exp(2*a)*exp(2*b*x) + 1))*(atan((log((exp(2*a)*exp(2*b*x)) / (exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*8i - 4) + log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*8i - 4)/((log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))* (log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)`**Reduce [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^2 x} dx$$

input `int(1/x/atanh(tanh(b*x+a))^2,x)`output `int(1/(atanh(tanh(a + b*x))**2*x),x)`

### 3.100 $\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	771
Mathematica [A] (verified)	772
Rubi [A] (verified)	772
Maple [A] (verified)	775
Fricas [A] (verification not implemented)	775
Sympy [F]	775
Maxima [A] (verification not implemented)	776
Giac [A] (verification not implemented)	776
Mupad [B] (verification not implemented)	776
Reduce [F]	777

#### Optimal result

Integrand size = 13, antiderivative size = 102

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$= -\frac{2b}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} + \frac{2b \log(x)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} - \frac{2b \log(\operatorname{arctanh}(\tanh(a+bx)))}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

output

```
-2*b/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))+1/x/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))+2*b*ln(x)/(b*x-arctanh(tanh(b*x+a)))^3-2*b*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^3
```



**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.69

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \frac{-b^2 x^2 + \operatorname{arctanh}(\tanh(a + bx))^2 + 2bx \operatorname{arctanh}(\tanh(a + bx))(\log(x) - \log(\operatorname{arctanh}(\tanh(a + bx))))}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \operatorname{arctanh}(\tanh(a + bx))}$$

input

```
Integrate[1/(x^2*ArcTanh[Tanh[a + b*x]]^2),x]
```

output

```
(-(b^2*x^2) + ArcTanh[Tanh[a + b*x]]^2 + 2*b*x*ArcTanh[Tanh[a + b*x]]*(Log[x] - Log[ArcTanh[Tanh[a + b*x]]]))/(x*(bx - ArcTanh[Tanh[a + b*x]])^3*ArcTanh[Tanh[a + b*x]])
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2602, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow \text{2602}$$

$$2b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow \text{2594}$$

$$2b \left( \frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))} \right) + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 2591

$$2b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right) +$$

$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

$$\frac{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}{1}$$

↓ 14

$$2b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right) +$$

$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

$$\frac{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}{1}$$

↓ 2588

$$2b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right) +$$

$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

$$\frac{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}{1}$$

↓ 14

$$\frac{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}{1} +$$

$$2b \left( -\frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

input

`Int [1/(x^2*ArcTanh[Tanh[a + b*x]]^2), x]`

output

```
1/(x*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]) + (2*b*(-(1/((
b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (-(Log[x]/(b*x -
ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh
[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]
])
```

### Defintions of rubi rules used

rule 14

```
Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]
```

rule 2588

```
Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst
[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]
```

rule 2591

```
Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D
[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1
/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n +
1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew
iseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1)/((m + 1)*(b*u - a*v))), x] + S
imp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; Ne
Q[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m
, -1]
```

**Maple [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.89

$$\frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2 x} - \frac{2b \ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3} - \frac{b}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2 \operatorname{arctanh}(\tanh(bx+a))}$$

input `int(1/x^2/arctanh(tanh(b*x+a))^2,x)`output `-1/(arctanh(tanh(b*x+a))-b*x)^2/x-2/(arctanh(tanh(b*x+a))-b*x)^3*b*ln(x)-1/(arctanh(tanh(b*x+a))-b*x)^2*b/arctanh(tanh(b*x+a))+2/(arctanh(tanh(b*x+a))-b*x)^3*b*ln(arctanh(tanh(b*x+a)))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.62

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{2abx + a^2 - 2(b^2x^2 + abx) \log(bx+a) + 2(b^2x^2 + abx) \log(x)}{a^3bx^2 + a^4x}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`output `-(2*a*b*x + a^2 - 2*(b^2*x^2 + a*b*x)*log(b*x + a) + 2*(b^2*x^2 + a*b*x)*log(x))/(a^3*b*x^2 + a^4*x)`**Sympy [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^2} dx = \int \frac{1}{x^2 \operatorname{atanh}^2(\tanh(a+bx))} dx$$

input `integrate(1/x**2/atanh(tanh(b*x+a))**2,x)`

output `Integral(1/(x**2*atanh(tanh(a + b*x))**2), x)`

### Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{2bx + a}{a^2 bx^2 + a^3 x} + \frac{2b \log(bx + a)}{a^3} - \frac{2b \log(x)}{a^3}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `-(2*b*x + a)/(a^2*b*x^2 + a^3*x) + 2*b*log(b*x + a)/a^3 - 2*b*log(x)/a^3`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2b \log(|bx + a|)}{a^3} - \frac{2b \log(|x|)}{a^3} - \frac{2bx + a}{(bx^2 + ax)a^2}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `2*b*log(abs(b*x + a))/a^3 - 2*b*log(abs(x))/a^3 - (2*b*x + a)/((b*x^2 + a*x)*a^2)`

### Mupad [B] (verification not implemented)

Time = 5.26 (sec) , antiderivative size = 432, normalized size of antiderivative = 4.24

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{4 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^2 - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \left(8 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) + bx \operatorname{atan}\left(\frac{-\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) + \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx}\right)}{x \left(\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)\right)}$$

input `int(1/(x^2*atanh(tanh(a + b*x))^2),x)`

output `-(4*log(1/(exp(2*a)*exp(2*b*x) + 1))^2 - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(8*log(1/(exp(2*a)*exp(2*b*x) + 1)) + b*x*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*32i) + 4*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 - 16*b^2*x^2 + b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*32i)/(x*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)`

## Reduce [F]

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^2 x^2} dx$$

input `int(1/x^2/atanh(tanh(b*x+a))^2,x)`

output `int(1/(atanh(tanh(a + b*x))^2*x**2),x)`

### 3.101 $\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	778
Mathematica [A] (verified)	779
Rubi [A] (verified)	779
Maple [A] (verified)	782
Fricas [A] (verification not implemented)	783
Sympy [F]	783
Maxima [A] (verification not implemented)	784
Giac [A] (verification not implemented)	784
Mupad [B] (verification not implemented)	784
Reduce [F]	785

#### Optimal result

Integrand size = 13, antiderivative size = 143

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$= -\frac{3b^2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{3b}{2x(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{3b^2 \log(x)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} - \frac{3b^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^4}$$

output

```
-3*b^2/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))+3/2*b/x/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))+1/2/x^2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))+3*b^2*ln(x)/(b*x-arctanh(tanh(b*x+a)))^4-3*b^2*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^4
```

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.64

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2b^3x^3 - 6bx \operatorname{arctanh}(\tanh(a + bx))^2 + \operatorname{arctanh}(\tanh(a + bx))^3 - 3b^2x^2 \operatorname{arctanh}(\tanh(a + bx))(-1 + 2 \operatorname{arctanh}(\tanh(a + bx)))}{2x^2 \operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input `Integrate[1/(x^3*ArcTanh[Tanh[a + b*x]]^2),x]`

output `-1/2*(2*b^3*x^3 - 6*b*x*ArcTanh[Tanh[a + b*x]]^2 + ArcTanh[Tanh[a + b*x]]^3 - 3*b^2*x^2*ArcTanh[Tanh[a + b*x]]*(-1 + 2*Log[x] - 2*Log[ArcTanh[Tanh[a + b*x]]]))/(x^2*ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)`

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.31, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {2602, 2602, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

↓ 2602

$$\frac{3b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^2} dx}{2(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 2602

$$\frac{3b \left( \frac{2b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))} \right)}{2(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$



↓ 2594

$$3b \left( \frac{2b \left( \frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$

---


$$\frac{2(bx - \operatorname{arctanh}(\tanh(a + bx)))}{1} \\ \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 2591

$$3b \left( \frac{2b \left( \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$

---


$$\frac{2(bx - \operatorname{arctanh}(\tanh(a + bx)))}{1} \\ \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 14

$$3b \left( \frac{2b \left( \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$

---


$$\frac{2(bx - \operatorname{arctanh}(\tanh(a + bx)))}{1} \\ \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 2588

$$\begin{aligned}
 & \left( \frac{2b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} \operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) \\
 & \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
 & \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow 14 \\
 & \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} + \\
 & \left( \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} + \frac{2b \left( -\frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) \\
 & \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx)))}
 \end{aligned}$$

input `Int [1/(x^3*ArcTanh[Tanh[a + b*x]]^2),x]`

output `1/(2*x^2*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]) + (3*b*(1/(x*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]) + (2*b*(-(1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (-Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])`

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1)/((m + 1)*(b*u - a*v))), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

## Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.81

$$-\frac{1}{2(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2 x^2} + \frac{3b^2 \ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4} + \frac{2b}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3}$$

input `int(1/x^3/arctanh(tanh(b*x+a))^2,x)`

output

```
-1/2/(arctanh(tanh(b*x+a))-b*x)^2/x^2+3/(arctanh(tanh(b*x+a))-b*x)^4*b^2*ln(x)+2/(arctanh(tanh(b*x+a))-b*x)^3*b/x-3/(arctanh(tanh(b*x+a))-b*x)^4*b^2*ln(arctanh(tanh(b*x+a)))+1/(arctanh(tanh(b*x+a))-b*x)^3*b^2/arctanh(tanh(b*x+a))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{6ab^2x^2 + 3a^2bx - a^3 - 6(b^3x^3 + ab^2x^2) \log(bx + a) + 6(b^3x^3 + ab^2x^2) \log(x)}{2(a^4bx^3 + a^5x^2)}$$

input

```
integrate(1/x^3/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
1/2*(6*a*b^2*x^2 + 3*a^2*b*x - a^3 - 6*(b^3*x^3 + a*b^2*x^2)*log(b*x + a) + 6*(b^3*x^3 + a*b^2*x^2)*log(x))/(a^4*b*x^3 + a^5*x^2)
```

**Sympy [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{x^3 \operatorname{atanh}^2(\tanh(a + bx))} dx$$

input

```
integrate(1/x**3/atanh(tanh(b*x+a))**2,x)
```

output

```
Integral(1/(x**3*atanh(tanh(a + b*x))**2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{6b^2x^2 + 3abx - a^2}{2(a^3bx^3 + a^4x^2)} - \frac{3b^2 \log(bx + a)}{a^4} + \frac{3b^2 \log(x)}{a^4}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `1/2*(6*b^2*x^2 + 3*a*b*x - a^2)/(a^3*b*x^3 + a^4*x^2) - 3*b^2*log(b*x + a)/a^4 + 3*b^2*log(x)/a^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{3b^2 \log(|bx + a|)}{a^4} + \frac{3b^2 \log(|x|)}{a^4} + \frac{6ab^2x^2 + 3a^2bx - a^3}{2(bx + a)a^4x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `-3*b^2*log(abs(b*x + a))/a^4 + 3*b^2*log(abs(x))/a^4 + 1/2*(6*a*b^2*x^2 + 3*a^2*b*x - a^3)/((b*x + a)*a^4*x^2)`**Mupad [B] (verification not implemented)**

Time = 5.11 (sec) , antiderivative size = 660, normalized size of antiderivative = 4.62

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(1/(x^3*atanh(tanh(a + b*x))^2),x)`

output

```

-(6*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 - 6*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*log(1/(exp(2*a)*exp(2*b*x) + 1))^3 - 2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3 - 32*b^3*x^3 + 24*b*x*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 24*b^2*x^2*log(1/(exp(2*a)*exp(2*b*x) + 1)) - 24*b^2*x^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 24*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))^2 + b^2*x^2*log(1/(exp(2*a)*exp(2*b*x) + 1))*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*96i - b^2*x^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*96i - 48*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(x^2*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)

```

**Reduce [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^2 x^3} dx$$

input

```
int(1/x^3/atanh(tanh(b*x+a))^2,x)
```

output

```
int(1/(atanh(tanh(a + b*x))^2*x**3),x)
```

### 3.102 $\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	786
Mathematica [A] (verified)	786
Rubi [A] (verified)	787
Maple [F]	788
Fricas [F]	789
Sympy [F]	789
Maxima [F]	789
Giac [F]	790
Mupad [F(-1)]	790
Reduce [F]	790

#### Optimal result

Integrand size = 13, antiderivative size = 94

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= -\frac{x^m}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{mx^{-1+m}}{2b^2\operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{mx^{-1+m} \operatorname{Hypergeometric2F1}\left(1, -1+m, m, \frac{bx}{bx-\operatorname{arctanh}(\tanh(a+bx))}\right)}{2b^2(bx-\operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
-1/2*x^m/b/arctanh(tanh(b*x+a))^2-1/2*m*x^(-1+m)/b^2/arctanh(tanh(b*x+a))-
1/2*m*x^(-1+m)*hypergeom([1,-1+m],[m],b*x/(b*x-arctanh(tanh(b*x+a))))/b^2
/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.86 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.54

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= \frac{x^{1+m} \operatorname{Hypergeometric2F1}\left(3, 1+m, 2+m, -\frac{bx}{-bx+\operatorname{arctanh}(\tanh(a+bx))}\right)}{(1+m)(-bx+\operatorname{arctanh}(\tanh(a+bx)))^3}$$

input `Integrate[x^m/ArcTanh[Tanh[a + b*x]]^3,x]`

output `(x^(1 + m)*Hypergeometric2F1[3, 1 + m, 2 + m, -((b*x)/(-b*x) + ArcTanh[Tanh[a + b*x]])])/((1 + m)*(-b*x) + ArcTanh[Tanh[a + b*x]])^3)`

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{m \int \frac{x^{m-1}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx}{2b} - \frac{x^m}{2b \operatorname{arctanh}(\tanh(a + bx))^2} \\
 & \quad \downarrow \text{2599} \\
 & \frac{m \left( -\frac{(1-m) \int \frac{x^{m-2}}{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} - \frac{x^{m-1}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{2b} - \frac{x^m}{2b \operatorname{arctanh}(\tanh(a + bx))^2} \\
 & \quad \downarrow \text{2595} \\
 & \frac{m \left( -\frac{x^{m-1} \operatorname{Hypergeometric2F1}\left(1, m-1, m, \frac{bx}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{b(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{x^{m-1}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{2b} - \frac{x^m}{2b \operatorname{arctanh}(\tanh(a + bx))^2}
 \end{aligned}$$

input `Int[x^m/ArcTanh[Tanh[a + b*x]]^3,x]`



output

```
-1/2*x^m/(b*ArcTanh[Tanh[a + b*x]]^2) + (m*(-(x^(-1 + m))/(b*ArcTanh[Tanh[a + b*x]])) - (x^(-1 + m)*Hypergeometric2F1[1, -1 + m, m, (b*x)/(b*x - ArcTanh[Tanh[a + b*x]])]/(b*(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b)
```

### Defintions of rubi rules used

rule 2595

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(v^(n + 1)/((n + 1)*(b*u - a*v))*Hypergeometric2F1[1, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && !IntegerQ[n]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [F]

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(bx + a))^3} dx$$

input

```
int(x^m/arctanh(tanh(b*x+a))^3,x)
```

output

```
int(x^m/arctanh(tanh(b*x+a))^3,x)
```

**Fricas [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^3} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `integral(x^m/arctanh(tanh(b*x + a))^3, x)`

**Sympy [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^m}{\operatorname{atanh}^3(\tanh(a + bx))} dx$$

input `integrate(x**m/atanh(tanh(b*x+a))**3,x)`

output `Integral(x**m/atanh(tanh(a + b*x))**3, x)`

**Maxima [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^3} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `integrate(x^m/arctanh(tanh(b*x + a))^3, x)`

**Giac [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^m}{\operatorname{artanh}(\tanh(bx + a))^3} dx$$

input `integrate(x^m/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `integrate(x^m/arctanh(tanh(b*x + a))^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^m}{\operatorname{atanh}(\tanh(a + bx))^3} dx$$

input `int(x^m/atanh(tanh(a + b*x))^3,x)`

output `int(x^m/atanh(tanh(a + b*x))^3, x)`

**Reduce [F]**

$$\int \frac{x^m}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{\operatorname{atanh}(\tanh(bx + a))^2 \left( \int \frac{x^m}{\operatorname{atanh}(\tanh(bx+a))^2 x} dx \right) m - x^m}{2 \operatorname{atanh}(\tanh(bx + a))^2 b}$$

input `int(x^m/atanh(tanh(b*x+a))^3,x)`

output `(atanh(tanh(a + b*x))**2*int(x**m/(atanh(tanh(a + b*x))**2*x),x)*m - x**m)/(2*atanh(tanh(a + b*x))**2*b)`

### 3.103 $\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	791
Mathematica [A] (verified)	792
Rubi [A] (verified)	792
Maple [B] (verified)	795
Fricas [A] (verification not implemented)	796
Sympy [F]	796
Maxima [A] (verification not implemented)	796
Giac [A] (verification not implemented)	797
Mupad [B] (verification not implemented)	797
Reduce [F]	798

#### Optimal result

Integrand size = 13, antiderivative size = 92

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= \frac{3x^2}{b^3} + \frac{6x(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

$$- \frac{x^4}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{2x^3}{b^2\operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{6(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^5}$$

output

```
3*x^2/b^3+6*x*(b*x-arctanh(tanh(b*x+a)))/b^4-1/2*x^4/b/arctanh(tanh(b*x+a))
^2-2*x^3/b^2/arctanh(tanh(b*x+a))+6*(b*x-arctanh(tanh(b*x+a)))^2*ln(arcta
nh(tanh(b*x+a)))/b^5
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.24

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{x^2}{2b^3} - \frac{3x(-bx + \operatorname{arctanh}(\tanh(a + bx)))}{b^4}$$

$$+ \frac{4(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3}{b^5 \operatorname{arctanh}(\tanh(a + bx))} - \frac{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}{2b^5 \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$+ \frac{6(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log(\operatorname{arctanh}(\tanh(a + bx)))}{b^5}$$

input `Integrate[x^4/ArcTanh[Tanh[a + b*x]]^3,x]`

output `x^2/(2*b^3) - (3*x*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/b^4 + (4*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3)/(b^5*ArcTanh[Tanh[a + b*x]]) - (-(b*x) + ArcTanh[Tanh[a + b*x]])^4/(2*b^5*ArcTanh[Tanh[a + b*x]]^2) + (6*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[ArcTanh[Tanh[a + b*x]]])/b^5`

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2599, 2599, 2590, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2599$$

$$2 \int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^2} dx - \frac{x^4}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2599$$

$$\begin{aligned}
 & \frac{2 \left( \frac{3 \int \frac{x^2 \operatorname{arctanh}(\tanh(a+bx)) dx}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{b} - \frac{x^4}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2590} \\
 & \frac{2 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x \operatorname{arctanh}(\tanh(a+bx)) dx}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{b} - \frac{x^4}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2589} \\
 & \frac{2 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1 \operatorname{arctanh}(\tanh(a+bx)) dx}{b} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{b} - \frac{x^4}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2588} \\
 & \frac{2 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{b^2} \operatorname{arctanh}(\tanh(a+bx)) dx}{b} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{b} - \frac{x^4}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{14}
 \end{aligned}$$

$$2 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2} + \frac{x}{b} \right)}{b} + \frac{x^2}{2b} \right)}{b} - \frac{x^3}{b \operatorname{arctanh}(\tanh(a+bx))} \right) \frac{x^4}{2b \operatorname{arctanh}(\tanh(a+bx))^2}$$

input `Int[x^4/ArcTanh[Tanh[a + b*x]]^3,x]`

output `-1/2*x^4/(b*ArcTanh[Tanh[a + b*x]]^2) + (2*(-(x^3/(b*ArcTanh[Tanh[a + b*x]])) + (3*(x^2/(2*b) + ((b*x - ArcTanh[Tanh[a + b*x]])*(x/b + ((b*x - ArcTanh[Tanh[a + b*x]])*Log[ArcTanh[Tanh[a + b*x]]])/b^2))/b))/b)/b`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n-1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 265 vs.  $2(90) = 180$ .

Time = 0.37 (sec) , antiderivative size = 266, normalized size of antiderivative = 2.89

method	result
default	$\frac{\frac{x^2 b}{2} - 3ax - 3x(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{b^4} - \frac{a^4 + 4a^3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 6a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + 4a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3}{2b^5 \operatorname{arctanh}(\tanh(bx+a))}$
risch	Expression too large to display

input

```
int(x^4/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
1/b^4*(1/2*x^2*b-3*a*x-3*x*(arctanh(tanh(b*x+a))-b*x-a))-1/2*(a^4+4*a^3*(arctanh(tanh(b*x+a))-b*x-a)+6*a^2*(arctanh(tanh(b*x+a))-b*x-a)^2+4*a*(arctanh(tanh(b*x+a))-b*x-a)^3+(arctanh(tanh(b*x+a))-b*x-a)^4)/b^5/arctanh(tanh(b*x+a))^2+(6*a^2+12*a*(arctanh(tanh(b*x+a))-b*x-a)+6*(arctanh(tanh(b*x+a))-b*x-a)^2)/b^5*ln(arctanh(tanh(b*x+a)))-(-4*a^3-12*a^2*(arctanh(tanh(b*x+a))-b*x-a)-12*a*(arctanh(tanh(b*x+a))-b*x-a)^2-4*(arctanh(tanh(b*x+a))-b*x-a)^3)/b^5/arctanh(tanh(b*x+a))
```



**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.03

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{b^4 x^4 - 4ab^3 x^3 - 11a^2 b^2 x^2 + 2a^3 bx + 7a^4 + 12(a^2 b^2 x^2 + 2a^3 bx + a^4) \log(bx + a)}{2(b^7 x^2 + 2ab^6 x + a^2 b^5)}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `1/2*(b^4*x^4 - 4*a*b^3*x^3 - 11*a^2*b^2*x^2 + 2*a^3*b*x + 7*a^4 + 12*(a^2*b^2*x^2 + 2*a^3*b*x + a^4)*log(b*x + a))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5)`**Sympy [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^4}{\operatorname{atanh}^3(\tanh(a + bx))} dx$$

input `integrate(x**4/atanh(tanh(b*x+a))**3,x)`output `Integral(x**4/atanh(tanh(a + b*x))**3, x)`**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.88

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{b^4 x^4 - 4ab^3 x^3 - 11a^2 b^2 x^2 + 2a^3 bx + 7a^4}{2(b^7 x^2 + 2ab^6 x + a^2 b^5)} + \frac{6a^2 \log(bx + a)}{b^5}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output

$$\frac{1}{2}(b^4x^4 - 4ab^3x^3 - 11a^2b^2x^2 + 2a^3bx + 7a^4)/(b^7x^2 + 2ab^6x + a^2b^5) + 6a^2\log(bx + a)/b^5$$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.66

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{6a^2 \log(|bx + a|)}{b^5} + \frac{b^3x^2 - 6ab^2x}{2b^6} + \frac{8a^3bx + 7a^4}{2(bx + a)^2b^5}$$

input

```
integrate(x^4/arctanh(tanh(b*x+a))^3,x, algorithm="giac")
```

output

$$6a^2\log(\operatorname{abs}(bx + a))/b^5 + 1/2*(b^3x^2 - 6ab^2x)/b^6 + 1/2*(8a^3bx + 7a^4)/((bx + a)^2b^5)$$

**Mupad [B] (verification not implemented)**

Time = 3.24 (sec) , antiderivative size = 867, normalized size of antiderivative = 9.42

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input

```
int(x^4/atanh(tanh(a + b*x))^3,x)
```

output

```

((7*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2
/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 + 24*a^2*(2*a - log((2*exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2
*b*x)^2 + 16*a^4 - 8*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*
b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 32*a^3*(2*a - l
og((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*ex
p(2*b*x) + 1)) + 2*b*x)))/(4*b) - x*(4*(2*a - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 -
32*a^3 - 24*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1
)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 48*a^2*(2*a - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x)))/(2*b^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(
2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + x*(16*a*b^5 -
8*b^5*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log
(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + 8*a^2*b^4 + 8*b^6*x^2 - 8*a*b^4*
(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + x^2/(2*b^3) + (log(log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(3*
(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 12*a*(2*a - log((2*exp(2*a)*exp(2*b...

```

**Reduce [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^4}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input

```
int(x^4/atanh(tanh(b*x+a))^3,x)
```

output

```
int(x**4/atanh(tanh(a + b*x))**3,x)
```

### 3.104 $\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	799
Mathematica [A] (verified)	799
Rubi [A] (verified)	800
Maple [B] (verified)	802
Fricas [A] (verification not implemented)	802
Sympy [F]	803
Maxima [A] (verification not implemented)	803
Giac [A] (verification not implemented)	803
Mupad [B] (verification not implemented)	804
Reduce [F]	805

#### Optimal result

Integrand size = 13, antiderivative size = 71

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{3x}{b^3} - \frac{x^3}{2b \operatorname{arctanh}(\tanh(a+bx))^2} - \frac{3x^2}{2b^2 \operatorname{arctanh}(\tanh(a+bx))} + \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^4}$$

output

```
3*x/b^3-1/2*x^3/b/arctanh(tanh(b*x+a))^2-3/2*x^2/b^2/arctanh(tanh(b*x+a))+
3*(b*x-arctanh(tanh(b*x+a)))*ln(arctanh(tanh(b*x+a)))/b^4
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.21

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{b^3 x^3 + 3b^2 x^2 \operatorname{arctanh}(\tanh(a+bx)) + \operatorname{arctanh}(\tanh(a+bx))^3 (5 + 6 \log(\operatorname{arctanh}(\tanh(a+bx)))) - b^4 \operatorname{arctanh}(\tanh(a+bx))^2}{2b^4 \operatorname{arctanh}(\tanh(a+bx))^2}$$

input

```
Integrate[x^3/ArcTanh[Tanh[a + b*x]]^3,x]
```

output

$$\frac{-1/2*(b^3*x^3 + 3*b^2*x^2*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^3*(5 + 6*Log[ArcTanh[Tanh[a + b*x]]]) - b*x*ArcTanh[Tanh[a + b*x]]^2*(11 + 6*Log[ArcTanh[Tanh[a + b*x]]]))}{(b^4*ArcTanh[Tanh[a + b*x]]^2)}$$

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2589, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2599$$

$$\frac{3 \int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^2} dx}{2b} - \frac{x^3}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2599$$

$$\frac{3 \left( \frac{2 \int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a + bx))} \right)}{2b} - \frac{x^3}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2589$$

$$\frac{3 \left( \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a + bx))} \right)}{2b} - \frac{x^3}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2588$$

$$\begin{array}{c}
 3 \left( \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d\operatorname{arctanh}(\tanh(a+bx))}{b^2} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} \right) \\
 \hline
 \frac{2b}{x^3} \\
 \frac{2b \operatorname{arctanh}(\tanh(a+bx))^2}{x^3} \\
 \downarrow 14 \\
 3 \left( \frac{2 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \log(\operatorname{arctanh}(\tanh(a+bx)))}{b^2} + \frac{x}{b} \right)}{b} - \frac{x^2}{b \operatorname{arctanh}(\tanh(a+bx))} \right) \\
 \hline
 \frac{2b}{x^3} \\
 \frac{2b \operatorname{arctanh}(\tanh(a+bx))^2}{x^3}
 \end{array}$$

input `Int [x^3/ArcTanh [Tanh [a + b*x]]^3,x]`

output `-1/2*x^3/(b*ArcTanh [Tanh [a + b*x]]^2) + (3*(-(x^2/(b*ArcTanh [Tanh [a + b*x]])) + (2*(x/b + ((b*x - ArcTanh [Tanh [a + b*x]])*Log [ArcTanh [Tanh [a + b*x]]])/b^2))/b)/(2*b)`

### Defintions of rubi rules used

rule 14 `Int [(a_.)/(x_), x_Symbol] := Simp [a*Log [x], x] /; FreeQ [a, x]`

rule 2588 `Int [(u_)^(m_.), x_Symbol] := With [{c = Simplify [D [u, x]]}, Simp [1/c Subst [Int [x^m, x], x, u], x]] /; FreeQ [m, x] && PiecewiseLinearQ [u, x]`

rule 2589 `Int [(v_)/(u_), x_Symbol] := With [{a = Simplify [D [u, x]], b = Simplify [D [v, x]]}, Simp [b*(x/a), x] - Simp [(b*u - a*v)/a Int [1/u, x], x] /; NeQ [b*u - a*v, 0] /; PiecewiseLinearQ [u, v, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs.  $2(67) = 134$ .

Time = 0.23 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.38

method	result
default	$\frac{x}{b^3} - \frac{3a^2 + 6a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^4 \operatorname{arctanh}(\tanh(bx+a))} - \frac{-a^3 - 3a^2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) - 3a}{2b^4}$
risch	Expression too large to display

input

```
int(x^3/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
x/b^3-(3*a^2+6*a*(arctanh(tanh(b*x+a))-b*x-a)+3*(arctanh(tanh(b*x+a))-b*x-a)^2)/b^4/arctanh(tanh(b*x+a))-1/2*(-a^3-3*a^2*(arctanh(tanh(b*x+a))-b*x-a)-3*a*(arctanh(tanh(b*x+a))-b*x-a)^2-(arctanh(tanh(b*x+a))-b*x-a)^3)/b^4/arctanh(tanh(b*x+a))^2+(-3*arctanh(tanh(b*x+a))+3*b*x)/b^4*ln(arctanh(tanh(b*x+a)))
```

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.17

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{2b^3x^3 + 4ab^2x^2 - 4a^2bx - 5a^3 - 6(ab^2x^2 + 2a^2bx + a^3) \log(bx + a)}{2(b^6x^2 + 2ab^5x + a^2b^4)}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

$$\frac{1}{2} \frac{(2b^3x^3 + 4ab^2x^2 - 4a^2bx - 5a^3 - 6(ab^2x^2 + 2a^2bx + a^3)) \log(bx + a)}{(b^6x^2 + 2ab^5x + a^2b^4)}$$

**Sympy [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^3}{\operatorname{atanh}^3(\tanh(a + bx))} dx$$

input

```
integrate(x**3/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(x**3/atanh(tanh(a + b*x))**3, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.97

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{2b^3x^3 + 4ab^2x^2 - 4a^2bx - 5a^3}{2(b^6x^2 + 2ab^5x + a^2b^4)} - \frac{3a \log(bx + a)}{b^4}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")
```

output

$$\frac{1}{2} \frac{(2b^3x^3 + 4ab^2x^2 - 4a^2bx - 5a^3)}{(b^6x^2 + 2ab^5x + a^2b^4)} - \frac{3a \log(bx + a)}{b^4}$$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.62

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{x}{b^3} - \frac{3a \log(|bx + a|)}{b^4} - \frac{6a^2bx + 5a^3}{2(bx + a)^2b^4}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^3,x, algorithm="giac")
```



output  $x/b^3 - 3*a*\log(\text{abs}(b*x + a))/b^4 - 1/2*(6*a^2*b*x + 5*a^3)/((b*x + a)^2*b^4)$

### Mupad [B] (verification not implemented)

Time = 3.31 (sec) , antiderivative size = 620, normalized size of antiderivative = 8.73

$$\int \frac{x^3}{\text{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(x^3/atanh(tanh(a + b*x))^3,x)`

output  $x/b^3 - (x*(3*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 - 12*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) + 12*a^2) - (5*((2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)))/(4*b))/(b^3*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + x*(8*a*b^4 - 4*b^4*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)) + 4*a^2*b^3 + 4*b^5*x^2 - 4*a*b^3*(2*a - \log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)) + (\log(\log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))*(3*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - 3*\log((2*\exp(2*a)*\exp(2*b*x)))/(\exp(2*a)*\exp(2*b*x) + 1)) + 6*b*x))/(2*b^4)$

**Reduce [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{x^3}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^3/atanh(tanh(b*x+a))^3,x)`

output `int(x**3/atanh(tanh(a + b*x))**3,x)`

### 3.105 $\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	806
Mathematica [A] (verified)	806
Rubi [A] (verified)	807
Maple [A] (verified)	808
Fricas [A] (verification not implemented)	809
Sympy [B] (verification not implemented)	809
Maxima [A] (verification not implemented)	810
Giac [A] (verification not implemented)	810
Mupad [B] (verification not implemented)	810
Reduce [B] (verification not implemented)	811

#### Optimal result

Integrand size = 13, antiderivative size = 47

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{x^2}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{x}{b^2\operatorname{arctanh}(\tanh(a+bx))} + \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{b^3}$$

output

$$-1/2*x^2/b/\operatorname{arctanh}(\tanh(b*x+a))^2 - x/b^2/\operatorname{arctanh}(\tanh(b*x+a)) + \ln(\operatorname{arctanh}(\tanh(b*x+a)))/b^3$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{3 - \frac{b^2 x^2}{\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{2bx}{\operatorname{arctanh}(\tanh(a+bx))} + 2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{2b^3}$$

input

`Integrate[x^2/ArcTanh[Tanh[a + b*x]]^3, x]`

output

$$(3 - (b^2 x^2)/\text{ArcTanh}[\text{Tanh}[a + b x]]^2 - (2 b x)/\text{ArcTanh}[\text{Tanh}[a + b x]] + 2 \text{Log}[\text{ArcTanh}[\text{Tanh}[a + b x]]])/(2 b^3)$$

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\text{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2599$$

$$\frac{\int \frac{x}{\text{arctanh}(\tanh(a + bx))^2} dx}{b} - \frac{x^2}{2b \text{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2599$$

$$\frac{\int \frac{1}{\text{arctanh}(\tanh(a + bx))} dx}{b} - \frac{x}{b \text{arctanh}(\tanh(a + bx))} - \frac{x^2}{2b \text{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2588$$

$$\frac{\int \frac{1}{\text{arctanh}(\tanh(a + bx))} d\text{arctanh}(\tanh(a + bx))}{b^2} - \frac{x}{b \text{arctanh}(\tanh(a + bx))} - \frac{x^2}{2b \text{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 14$$

$$\frac{\log(\text{arctanh}(\tanh(a + bx)))}{b^2} - \frac{x}{b \text{arctanh}(\tanh(a + bx))} - \frac{x^2}{2b \text{arctanh}(\tanh(a + bx))^2}$$

input

$$\text{Int}[x^2/\text{ArcTanh}[\text{Tanh}[a + b x]]^3, x]$$

output

$$-1/2 x^2/(b \text{ArcTanh}[\text{Tanh}[a + b x]]^2) + (-x/(b \text{ArcTanh}[\text{Tanh}[a + b x]])) + \text{Log}[\text{ArcTanh}[\text{Tanh}[a + b x]]]/b^2/b$$

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.15

method	result
parallelrisch	$\frac{-b^2 x^2 + 2 \ln(\operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^2 - 2x \operatorname{arctanh}(\tanh(bx+a))b}{2b^3 \operatorname{arctanh}(\tanh(bx+a))^2}$
default	$\frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{b^3} - \frac{2bx - 2 \operatorname{arctanh}(\tanh(bx+a))}{b^3 \operatorname{arctanh}(\tanh(bx+a))} - \frac{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2}{2b^3 \operatorname{arctanh}(\tanh(bx+a))^2}$
risch	$-\frac{4i \left( \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) x - \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 x + \pi \operatorname{csgn}(ie^{bx+a}) \right)}{b^2 \left( \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) - \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + \pi \operatorname{csgn}(ie^{bx+a}) \right)}$

input `int(x^2/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `1/2*(-b^2*x^2+2*ln(arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^2-2*x*arctanh(tanh(b*x+a))*b)/b^3/arctanh(tanh(b*x+a))^2`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.30

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{4abx + 3a^2 + 2(b^2x^2 + 2abx + a^2) \log(bx + a)}{2(b^5x^2 + 2ab^4x + a^2b^3)}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `1/2*(4*a*b*x + 3*a^2 + 2*(b^2*x^2 + 2*a*b*x + a^2)*log(b*x + a))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(42) = 84.

Time = 26.93 (sec) , antiderivative size = 112, normalized size of antiderivative = 2.38

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \begin{cases} \frac{x^3}{3 \operatorname{atanh}^3(\tanh(a))} & \text{for } b = 0 \\ \frac{x^3}{3 \operatorname{atanh}^3(\tanh(bx + \log(-e^{-bx})))} & \text{for } a = \log(-e^{-bx}) \\ \frac{x^3}{3 \operatorname{atanh}^3(\tanh(bx + \log(e^{-bx})))} & \text{for } a = \log(e^{-bx}) \\ -\frac{x^2}{2b \operatorname{atanh}^2(\tanh(a + bx))} - \frac{x}{b^2 \operatorname{atanh}(\tanh(a + bx))} + \frac{\log(\operatorname{atanh}(\tanh(a + bx)))}{b^3} & \text{otherwise} \end{cases}$$

input `integrate(x**2/atanh(tanh(b*x+a))**3,x)`

output `Piecewise((x**3/(3*atanh(tanh(a))**3), Eq(b, 0)), (x**3/(3*atanh(tanh(b*x + log(-exp(-b*x))))**3), Eq(a, log(-exp(-b*x)))), (x**3/(3*atanh(tanh(b*x + log(exp(-b*x))))**3), Eq(a, log(exp(-b*x)))), (-x**2/(2*b*atanh(tanh(a + b*x))**2) - x/(b**2*atanh(tanh(a + b*x))) + log(atanh(tanh(a + b*x)))/b**3, True))`

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.02

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{4abx + 3a^2}{2(b^5x^2 + 2ab^4x + a^2b^3)} + \frac{\log(bx + a)}{b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output `1/2*(4*a*b*x + 3*a^2)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3) + log(b*x + a)/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{\log(|bx + a|)}{b^3} + \frac{4ax + \frac{3a^2}{b}}{2(bx + a)^2b^2}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`output `log(abs(b*x + a))/b^3 + 1/2*(4*a*x + 3*a^2/b)/((b*x + a)^2*b^2)`**Mupad [B] (verification not implemented)**

Time = 3.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{\ln(\operatorname{atanh}(\tanh(a + bx)))}{b^3} - \frac{\frac{b^2x^2}{2} + bx \operatorname{atanh}(\tanh(a + bx))}{b^3 \operatorname{atanh}(\tanh(a + bx))^2}$$

input `int(x^2/atanh(tanh(a + b*x))^3,x)`output `log(atanh(tanh(a + b*x)))/b^3 - ((b^2*x^2)/2 + b*x*atanh(tanh(a + b*x)))/(b^3*atanh(tanh(a + b*x))^2)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.13

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{2 \operatorname{atanh}(\tanh(bx + a))^2 \log(\operatorname{atanh}(\tanh(bx + a))) - 2 \operatorname{atanh}(\tanh(bx + a)) bx - b^2 x^2}{2 \operatorname{atanh}(\tanh(bx + a))^2 b^3}$$

input

```
int(x^2/atanh(tanh(b*x+a))^3,x)
```

output

```
(2*atanh(tanh(a + b*x))**2*log(atanh(tanh(a + b*x))) - 2*atanh(tanh(a + b*x))*b*x - b**2*x**2)/(2*atanh(tanh(a + b*x))**2*b**3)
```



### 3.106 $\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	812
Mathematica [A] (verified)	812
Rubi [A] (verified)	813
Maple [A] (verified)	814
Fricas [A] (verification not implemented)	814
Sympy [B] (verification not implemented)	815
Maxima [A] (verification not implemented)	815
Giac [A] (verification not implemented)	816
Mupad [B] (verification not implemented)	816
Reduce [B] (verification not implemented)	816

#### Optimal result

Integrand size = 11, antiderivative size = 34

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{x}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{1}{2b^2\operatorname{arctanh}(\tanh(a+bx))}$$

output `-1/2*x/b/arctanh(tanh(b*x+a))^2-1/2/b^2/arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{bx + \operatorname{arctanh}(\tanh(a+bx))}{2b^2\operatorname{arctanh}(\tanh(a+bx))^2}$$

input `Integrate[x/ArcTanh[Tanh[a + b*x]]^3,x]`

output `-1/2*(b*x + ArcTanh[Tanh[a + b*x]])/(b^2*ArcTanh[Tanh[a + b*x]]^2)`

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2599$$

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} dx - \frac{x}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2588$$

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^2} d\operatorname{arctanh}(\tanh(a + bx)) - \frac{x}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 15$$

$$-\frac{1}{2b^2 \operatorname{arctanh}(\tanh(a + bx))} - \frac{x}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

input `Int[x/ArcTanh[Tanh[a + b*x]]^3,x]`

output `-1/2*x/(b*ArcTanh[Tanh[a + b*x]]^2) - 1/(2*b^2*ArcTanh[Tanh[a + b*x]])`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```

Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))

```

**Maple [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

method	result
parallelisch	$-\frac{\operatorname{arctanh}(\tanh(bx+a))+bx}{2b^2 \operatorname{arctanh}(\tanh(bx+a))^2}$
default	$-\frac{1}{b^2 \operatorname{arctanh}(\tanh(bx+a))} - \frac{bx - \operatorname{arctanh}(\tanh(bx+a))}{2b^2 \operatorname{arctanh}(\tanh(bx+a))^2}$
risch	$-\frac{2i \left( \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) - \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + \pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \right)}{b^2 \left( \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) - \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + \pi \operatorname{csgn}(ie^{bx+a})^2 \right)}$

input

```
int(x/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*(arctanh(tanh(b*x+a))+b*x)/b^2/arctanh(tanh(b*x+a))^2
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{2bx + a}{2(b^4x^2 + 2ab^3x + a^2b^2)}$$

input

```
integrate(x/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

```
-1/2*(2*b*x + a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 100 vs.  $2(31) = 62$ .

Time = 25.88 (sec) , antiderivative size = 100, normalized size of antiderivative = 2.94

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \begin{cases} \frac{x^2}{2 \operatorname{atanh}^3(\tanh(a))} & \text{for } b = 0 \\ \frac{x^2}{2 \operatorname{atanh}^3(\tanh(bx + \log(-e^{-bx})))} & \text{for } a = \log(-e^{-bx}) \\ \frac{x^2}{2 \operatorname{atanh}^3(\tanh(bx + \log(e^{-bx})))} & \text{for } a = \log(e^{-bx}) \\ -\frac{x}{2b \operatorname{atanh}^2(\tanh(a + bx))} - \frac{1}{2b^2 \operatorname{atanh}(\tanh(a + bx))} & \text{otherwise} \end{cases}$$

input `integrate(x/atanh(tanh(b*x+a))**3,x)`

output `Piecewise((x**2/(2*atanh(tanh(a))**3), Eq(b, 0)), (x**2/(2*atanh(tanh(b*x + log(-exp(-b*x))))**3), Eq(a, log(-exp(-b*x)))), (x**2/(2*atanh(tanh(b*x + log(exp(-b*x))))**3), Eq(a, log(exp(-b*x)))), (-x/(2*b*atanh(tanh(a + b*x)**2) - 1/(2*b**2*atanh(tanh(a + b*x))), True))`

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{2bx + a}{2(b^4x^2 + 2ab^3x + a^2b^2)}$$

input `integrate(x/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `-1/2*(2*b*x + a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.53

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{2bx + a}{2(bx + a)^2 b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `-1/2*(2*b*x + a)/((b*x + a)^2*b^2)`

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{\operatorname{atanh}(\tanh(a + bx)) + bx}{2b^2 \operatorname{atanh}(\tanh(a + bx))^2}$$

input `int(x/atanh(tanh(a + b*x))^3,x)`

output `-(atanh(tanh(a + b*x)) + b*x)/(2*b^2*atanh(tanh(a + b*x))^2)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{-\operatorname{atanh}(\tanh(bx + a)) - bx}{2\operatorname{atanh}(\tanh(bx + a))^2 b^2}$$

input `int(x/atanh(tanh(b*x+a))^3,x)`

output `( - (atanh(tanh(a + b*x)) + b*x))/(2*atanh(tanh(a + b*x))*2*b**2)`

### 3.107 $\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	817
Mathematica [A] (verified)	817
Rubi [A] (verified)	818
Maple [A] (verified)	819
Fricas [A] (verification not implemented)	819
Sympy [A] (verification not implemented)	820
Maxima [A] (verification not implemented)	820
Giac [A] (verification not implemented)	820
Mupad [B] (verification not implemented)	821
Reduce [B] (verification not implemented)	821

#### Optimal result

Integrand size = 9, antiderivative size = 16

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{1}{2b\operatorname{arctanh}(\tanh(a+bx))^2}$$

output `-1/2/b/arctanh(tanh(b*x+a))^2`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{1}{2b\operatorname{arctanh}(\tanh(a+bx))^2}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(-3), x]`

output `-1/2*1/(b*ArcTanh[Tanh[a + b*x]]^2)`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

↓ 2588

$$\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

↓ 15

$$-\frac{1}{2b\operatorname{arctanh}(\tanh(a + bx))^2}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(-3), x]`

output `-1/2*1/(b*ArcTanh[Tanh[a + b*x]]^2)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{1}{2b \operatorname{arctanh}(\tanh(bx+a))^2}$
default	$-\frac{1}{2b \operatorname{arctanh}(\tanh(bx+a))^2}$
parallelrisc	$-\frac{1}{2b \operatorname{arctanh}(\tanh(bx+a))^2}$
risc	$b \left( -\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) + \pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 - \pi \operatorname{csgn}(ie^{bx+2a}) \right)$

input `int(1/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`output `-1/2/b/arctanh(tanh(b*x+a))^2`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{1}{2(b^3x^2 + 2ab^2x + a^2b)}$$

input `integrate(1/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `-1/2/(b^3*x^2 + 2*a*b^2*x + a^2*b)`



**Sympy [A] (verification not implemented)**

Time = 25.58 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \begin{cases} -\frac{1}{2b \operatorname{atanh}^2(\tanh(a + bx))} & \text{for } b \neq 0 \\ \frac{x}{\operatorname{atanh}^3(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a))**3,x)`output `Piecewise((-1/(2*b*atanh(tanh(a + b*x))**2), Ne(b, 0)), (x/atanh(tanh(a))*  
*3, True))`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{1}{2(bx + a)^2 b}$$

input `integrate(1/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output `-1/2/((b*x + a)^2*b)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{1}{2(bx + a)^2 b}$$

input `integrate(1/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`output `-1/2/((b*x + a)^2*b)`

**Mupad [B] (verification not implemented)**

Time = 3.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{1}{2b \operatorname{atanh}(\tanh(a + bx))^2}$$

input `int(1/atanh(tanh(a + b*x))^3,x)`output `-1/(2*b*atanh(tanh(a + b*x))^2)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{1}{2 \operatorname{atanh}(\tanh(bx + a))^2 b}$$

input `int(1/atanh(tanh(b*x+a))^3,x)`output `( - 1)/(2*atanh(tanh(a + b*x))**2*b)`

### 3.108 $\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	822
Mathematica [A] (verified)	823
Rubi [A] (verified)	823
Maple [A] (verified)	825
Fricas [A] (verification not implemented)	826
Sympy [F]	826
Maxima [A] (verification not implemented)	826
Giac [A] (verification not implemented)	827
Mupad [B] (verification not implemented)	827
Reduce [F]	828

#### Optimal result

Integrand size = 13, antiderivative size = 97

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= -\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{\log(x)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

output

```
-1/2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^2+1/(b*x-arctanh(tanh
(b*x+a))^2/arctanh(tanh(b*x+a))-ln(x)/(b*x-arctanh(tanh(b*x+a)))^3+ln(arc
tanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^3
```

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{b^2 x^2 - 4bx \operatorname{arctanh}(\tanh(a + bx)) + \operatorname{arctanh}(\tanh(a + bx))^2 (3 + 2 \log(bx)) - 2 \log(\operatorname{arctanh}(\tanh(a + bx)))}{2 \operatorname{arctanh}(\tanh(a + bx))^2 (-bx + \operatorname{arctanh}(\tanh(a + bx)))^3}$$

input

```
Integrate[1/(x*ArcTanh[Tanh[a + b*x]]^3),x]
```

output

```
(b^2*x^2 - 4*b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2*(3 + 2*
Log[b*x] - 2*Log[ArcTanh[Tanh[a + b*x]]]))/(2*ArcTanh[Tanh[a + b*x]]^2*(-(
b*x) + ArcTanh[Tanh[a + b*x]]^3)
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.38, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2594, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2594$$

$$-\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2594$$

$$-\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))}$$

$$\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

↓ 2591

$$\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx - \int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2588

$$\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} d \operatorname{arctanh}(\tanh(a+bx)) - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$\frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{\frac{\log(\operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))}}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

input `Int [1/(x*ArcTanh[Tanh[a + b*x]]^3), x]`

output `-1/2*1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) - (-1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (-Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])/(b*x - ArcTanh[Tanh[a + b*x]])/(b*x - ArcTanh[Tanh[a + b*x]])`

## Definitions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst  
[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D  
[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1  
/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D  
[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n +  
1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew  
iseLinearQ[u, v, x] && LtQ[n, -1]`

## Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.95

$$\frac{\ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3} - \frac{\ln(\operatorname{arctanh}(\tanh(bx+a)))}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3} + \frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2 \operatorname{arctanh}(\tanh(bx+a))}$$

input `int(1/x/arctanh(tanh(b*x+a))^3,x)`

output `1/(arctanh(tanh(b*x+a))-b*x)^3*ln(x)-1/(arctanh(tanh(b*x+a))-b*x)^3*ln(arc  
tanh(tanh(b*x+a)))+1/(arctanh(tanh(b*x+a))-b*x)^2/arctanh(tanh(b*x+a))+1/2  
/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^2`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{2 abx + 3 a^2 - 2 (b^2 x^2 + 2 abx + a^2) \log (bx + a) + 2 (b^2 x^2 + 2 abx + a^2) \log (x)}{2 (a^3 b^2 x^2 + 2 a^4 bx + a^5)}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `1/2*(2*a*b*x + 3*a^2 - 2*(b^2*x^2 + 2*a*b*x + a^2)*log(b*x + a) + 2*(b^2*x^2 + 2*a*b*x + a^2)*log(x))/(a^3*b^2*x^2 + 2*a^4*b*x + a^5)`**Sympy [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input `integrate(1/x/atanh(tanh(b*x+a))**3,x)`output `Integral(1/(x*atanh(tanh(a + b*x))**3), x)`**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.53

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{2 bx + 3 a}{2 (a^2 b^2 x^2 + 2 a^3 bx + a^4)} - \frac{\log (bx + a)}{a^3} + \frac{\log (x)}{a^3}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output `1/2*(2*b*x + 3*a)/(a^2*b^2*x^2 + 2*a^3*b*x + a^4) - log(b*x + a)/a^3 + log(x)/a^3`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.44

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{\log(|bx + a|)}{a^3} + \frac{\log(|x|)}{a^3} + \frac{2abx + 3a^2}{2(bx + a)^2 a^3}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `-log(abs(b*x + a))/a^3 + log(abs(x))/a^3 + 1/2*(2*a*b*x + 3*a^2)/((b*x + a)^2*a^3)`

**Mupad [B] (verification not implemented)**

Time = 5.43 (sec) , antiderivative size = 645, normalized size of antiderivative = 6.65

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(1/(x*atanh(tanh(a + b*x))^3),x)`



output

```

-(12*log(1/(exp(2*a)*exp(2*b*x) + 1))^2 - 24*log(1/(exp(2*a)*exp(2*b*x) +
1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*e
xp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2*atan((log((exp(2*a)*exp(2*b*x))/(e
xp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i
)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x))*16i - log(1/(exp(2*a)*exp(2*b*x) + 1))^2*atan((
log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*
exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp
(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*16i + 12*log((exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 16*b^2*x^2 + log(1/(exp(2*a
)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*at
an((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2
*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log(
(exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*32i + b*x*(32*log
(1/(exp(2*a)*exp(2*b*x) + 1)) - 32*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*e
xp(2*b*x) + 1))))/((log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2
*b*x))/(exp(2*a)*exp(2*b*x) + 1)))^2*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log
((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)

```

**Reduce [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^3 x} dx$$

input

```
int(1/x/atanh(tanh(b*x+a))^3,x)
```

output

```
int(1/(atanh(tanh(a + b*x))**3*x),x)
```

### 3.109 $\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result . . . . .	829
Mathematica [A] (verified) . . . . .	830
Rubi [A] (verified) . . . . .	830
Maple [A] (verified) . . . . .	833
Fricas [A] (verification not implemented) . . . . .	833
Sympy [F] . . . . .	834
Maxima [A] (verification not implemented) . . . . .	834
Giac [A] (verification not implemented) . . . . .	834
Mupad [B] (verification not implemented) . . . . .	835
Reduce [F] . . . . .	836

#### Optimal result

Integrand size = 13, antiderivative size = 131

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= -\frac{3b}{2(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{3b}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{3b \log(x)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{3b \log(\operatorname{arctanh}(\tanh(a+bx)))}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^4}$$

output

```
-3/2*b/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^2+1/x/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^2+3*b/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))-3*b*ln(x)/(b*x-arctanh(tanh(b*x+a)))^4+3*b*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^4
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.71

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{b^3 x^3 - 6b^2 x^2 \operatorname{arctanh}(\tanh(a + bx)) + 2 \operatorname{arctanh}(\tanh(a + bx))^3 + 3bx \operatorname{arctanh}(\tanh(a + bx))^2 (1 + 2 \operatorname{Log}[x]) - 2 \operatorname{Log}[\operatorname{ArcTanh}[\tanh(a + bx)]]}{2x \operatorname{arctanh}(\tanh(a + bx))^2 (-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input `Integrate[1/(x^2*ArcTanh[Tanh[a + b*x]]^3),x]`

output `-1/2*(b^3*x^3 - 6*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 2*ArcTanh[Tanh[a + b*x]]^3 + 3*b*x*ArcTanh[Tanh[a + b*x]]^2*(1 + 2*Log[x] - 2*Log[ArcTanh[Tanh[a + b*x]]]))/(x*ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)`

**Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.38, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {2602, 2594, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

↓ 2602

$$\frac{3b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

↓ 2594

$$\frac{3b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^2} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

↓ 2594

$$3b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$


---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{1}$$


---


$$\frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2591

$$3b \left( -\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$


---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{1}$$


---


$$\frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$3b \left( -\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$


---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{1}$$


---


$$\frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2588

$$3b \left( -\frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} \operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)$$


---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{1}$$


---


$$\frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$3b \left( \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))^2} + \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))^2} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))} - \frac{\log(\operatorname{arctanh}(\tanh(a + bx)))}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right) \frac{1}{bx - \operatorname{arctanh}(\tanh(a + bx))}$$

input `Int[1/(x^2*ArcTanh[Tanh[a + b*x]]^3),x]`

output `1/(x*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) + (3*b*(-1/2 *1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) - (-1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (-Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst [Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*(m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.89

$$\frac{1}{(\operatorname{arctanh}(\tanh(bx + a)) - bx)^3 x} - \frac{3b \ln(x)}{(\operatorname{arctanh}(\tanh(bx + a)) - bx)^4} - \frac{b}{2(\operatorname{arctanh}(\tanh(bx + a)) - bx)^2}$$

input

```
int(1/x^2/arctanh(tanh(b*x+a))^3,x)
```

output

```
-1/(arctanh(tanh(b*x+a))-b*x)^3/x-3/(arctanh(tanh(b*x+a))-b*x)^4*b*ln(x)-1/2/(arctanh(tanh(b*x+a))-b*x)^2/arctanh(tanh(b*x+a))^2*b+3/(arctanh(tanh(b*x+a))-b*x)^4*b*ln(arctanh(tanh(b*x+a)))-2/(arctanh(tanh(b*x+a))-b*x)^3*b/arctanh(tanh(b*x+a))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.83

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{6ab^2x^2 + 9a^2bx + 2a^3 - 6(b^3x^3 + 2ab^2x^2 + a^2bx) \log(bx + a) + 6(b^3x^3 + 2ab^2x^2 + a^2bx) \log(x)}{2(a^4b^2x^3 + 2a^5bx^2 + a^6x)}$$

input

```
integrate(1/x^2/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

```
-1/2*(6*a*b^2*x^2 + 9*a^2*b*x + 2*a^3 - 6*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*log(b*x + a) + 6*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*log(x))/(a^4*b^2*x^3 + 2*a^5*b*x^2 + a^6*x)
```

**Sympy [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x^2 \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input `integrate(1/x**2/atanh(tanh(b*x+a))**3,x)`

output `Integral(1/(x**2*atanh(tanh(a + b*x))**3), x)`

**Maxima [A] (verification not implemented)**

Time = 0.60 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.53

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{6b^2x^2 + 9abx + 2a^2}{2(a^3b^2x^3 + 2a^4bx^2 + a^5x)} + \frac{3b \log(bx + a)}{a^4} - \frac{3b \log(x)}{a^4}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `-1/2*(6*b^2*x^2 + 9*a*b*x + 2*a^2)/(a^3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x) + 3*b*log(b*x + a)/a^4 - 3*b*log(x)/a^4`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.46

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{3b \log(|bx + a|)}{a^4} - \frac{3b \log(|x|)}{a^4} - \frac{6ab^2x^2 + 9a^2bx + 2a^3}{2(bx + a)^2 a^4 x}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output

```
3*b*log(abs(b*x + a))/a^4 - 3*b*log(abs(x))/a^4 - 1/2*(6*a*b^2*x^2 + 9*a^2
*b*x + 2*a^3)/((b*x + a)^2*a^4*x)
```

**Mupad [B] (verification not implemented)**

Time = 5.11 (sec) , antiderivative size = 804, normalized size of antiderivative = 6.14

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input

```
int(1/(x^2*atanh(tanh(a + b*x))^3),x)
```

output

```
-(24*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)
)*exp(2*b*x) + 1)) - 24*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 - 8*log(1/(exp(2*a)*exp(2*b*x) + 1))
^3 + 8*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3 + 32*b^3*x^3
+ 24*b*x*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 96*b^2*
x^2*log(1/(exp(2*a)*exp(2*b*x) + 1)) - 96*b^2*x^2*log((exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1)) + 24*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))^2 -
b*x*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2*atan((log((exp
(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*
x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*ex
p(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*96i - b*x*log(1/(exp(2*a)*e
xp(2*b*x) + 1))^2*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1
))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp
(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b
*x))*96i - 48*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x)
))/(exp(2*a)*exp(2*b*x) + 1)) + b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*atan((log((exp(2*a)*exp(2*
b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i
+ b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(e
xp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*192i)/(x*(log(1/(exp(2*a)*exp(2*b*x)...

```



**Reduce [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^3 x^2} dx$$

input `int(1/x^2/atanh(tanh(b*x+a))^3,x)`

output `int(1/(atanh(tanh(a + b*x))**3*x**2),x)`

### 3.110 $\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result . . . . .	837
Mathematica [A] (verified) . . . . .	838
Rubi [A] (verified) . . . . .	838
Maple [A] (verified) . . . . .	842
Fricas [A] (verification not implemented) . . . . .	842
Sympy [F] . . . . .	843
Maxima [A] (verification not implemented) . . . . .	843
Giac [A] (verification not implemented) . . . . .	843
Mupad [B] (verification not implemented) . . . . .	844
Reduce [F] . . . . .	845

#### Optimal result

Integrand size = 13, antiderivative size = 170

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= -\frac{3b^2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{2b}{x(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$+ \frac{6b^2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^4 \operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{6b^2 \log(x)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^5} + \frac{6b^2 \log(\operatorname{arctanh}(\tanh(a+bx)))}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^5}$$

output

```
-3*b^2/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^2+2*b/x/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^2+1/2/x^2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^2+6*b^2/(b*x-arctanh(tanh(b*x+a)))^4/arctanh(tanh(b*x+a))-6*b^2*ln(x)/(b*x-arctanh(tanh(b*x+a)))^5+6*b^2*ln(arctanh(tanh(b*x+a)))/(b*x-arctanh(tanh(b*x+a)))^5
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.63

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{-b^4 x^4 + 8b^3 x^3 \operatorname{arctanh}(\tanh(a + bx)) - 8bx \operatorname{arctanh}(\tanh(a + bx))^3 + \operatorname{arctanh}(\tanh(a + bx))^4 - 12b^2 x^2}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx)))^5 \operatorname{arctanh}(\tanh(a + bx))}$$

input

```
Integrate[1/(x^3*ArcTanh[Tanh[a + b*x]]^3),x]
```

output

```
(-(b^4*x^4) + 8*b^3*x^3*ArcTanh[Tanh[a + b*x]] - 8*b*x*ArcTanh[Tanh[a + b*x]]^3 + ArcTanh[Tanh[a + b*x]]^4 - 12*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2*(Log[x] - Log[ArcTanh[Tanh[a + b*x]]]))/(2*x^2*(b*x - ArcTanh[Tanh[a + b*x]])^5*ArcTanh[Tanh[a + b*x]]^2)
```

**Rubi [A] (verified)**Time = 0.78 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.36, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$ , Rules used = {2602, 2602, 2594, 2594, 2591, 14, 2588, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2602$$

$$\frac{2b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^3} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2602$$

$$\frac{2b \left( \frac{3b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^3} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{x (bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{2x^2 (bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^2}$$

↓ 2594

$$2b \left( \frac{3b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^2} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))}$$

---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2594

$$2b \left( \frac{3b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2591

$$2b \left( \frac{3b \left( -\frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{\int \frac{1}{x} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

---


$$\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$2b \left( \frac{3b \left( \frac{b \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} dx - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$\frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2588

$$2b \left( \frac{3b \left( \frac{\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))} a \operatorname{arctanh}(\tanh(a+bx)) - \frac{\log(x)}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$\frac{1}{2x^2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 14

$$2b \left( \frac{1}{x(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2} + \frac{3b \left( \frac{1}{2(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^2} - \frac{1}{(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$bx - \operatorname{arctanh}(\tanh(a+bx))$$

input `Int[1/(x^3*ArcTanh[Tanh[a + b*x]]^3),x]`

output `1/(2*x^2*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) + (2*b*(1/(x*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) + (3*b*(-1/2*1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^2) - (-1/((b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]])) - (-Log[x]/(b*x - ArcTanh[Tanh[a + b*x]])) + Log[ArcTanh[Tanh[a + b*x]]]/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])`

### Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2591 `Int[1/((u_)*(v_)), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b/(b*u - a*v) Int[1/v, x], x] - Simp[a/(b*u - a*v) Int[1/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1)/((m + 1)*(b*u - a*v))), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

**Maple [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.85

$$-\frac{1}{2(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3 x^2} + \frac{6b^2 \ln(x)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^5} + \frac{3b}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4}$$

input `int(1/x^3/arctanh(tanh(b*x+a))^3,x)`output `-1/2/(arctanh(tanh(b*x+a))-b*x)^3/x^2+6/(arctanh(tanh(b*x+a))-b*x)^5*b^2*ln(x)+3/(arctanh(tanh(b*x+a))-b*x)^4*b/x-6/(arctanh(tanh(b*x+a))-b*x)^5*b^2*ln(arctanh(tanh(b*x+a)))+3/(arctanh(tanh(b*x+a))-b*x)^4*b^2/arctanh(tanh(b*x+a))+1/2/(arctanh(tanh(b*x+a))-b*x)^3*b^2/arctanh(tanh(b*x+a))^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= \frac{12ab^3x^3 + 18a^2b^2x^2 + 4a^3bx - a^4 - 12(b^4x^4 + 2ab^3x^3 + a^2b^2x^2) \log(bx+a) + 12(b^4x^4 + 2ab^3x^3 + a^2b^2x^2) \log(x)}{2(a^5b^2x^4 + 2a^6bx^3 + a^7x^2)}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `1/2*(12*a*b^3*x^3 + 18*a^2*b^2*x^2 + 4*a^3*b*x - a^4 - 12*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*log(b*x + a) + 12*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*log(x))/(a^5*b^2*x^4 + 2*a^6*b*x^3 + a^7*x^2)`

**Sympy [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x^3 \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input `integrate(1/x**3/atanh(tanh(b*x+a))**3,x)`

output `Integral(1/(x**3*atanh(tanh(a + b*x))**3), x)`

**Maxima [A] (verification not implemented)**

Time = 0.61 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.51

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{12b^3x^3 + 18ab^2x^2 + 4a^2bx - a^3}{2(a^4b^2x^4 + 2a^5bx^3 + a^6x^2)} - \frac{6b^2 \log(bx + a)}{a^5} + \frac{6b^2 \log(x)}{a^5}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `1/2*(12*b^3*x^3 + 18*a*b^2*x^2 + 4*a^2*b*x - a^3)/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2) - 6*b^2*log(b*x + a)/a^5 + 6*b^2*log(x)/a^5`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.43

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{6b^2 \log(|bx + a|)}{a^5} + \frac{6b^2 \log(|x|)}{a^5} + \frac{12b^3x^3 + 18ab^2x^2 + 4a^2bx - a^3}{2(bx^2 + ax)^2 a^4}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`



output

$$-6*b^2*\log(\text{abs}(b*x + a))/a^5 + 6*b^2*\log(\text{abs}(x))/a^5 + 1/2*(12*b^3*x^3 + 18*a*b^2*x^2 + 4*a^2*b*x - a^3)/((b*x^2 + a*x)^2*a^4)$$
**Mupad [B] (verification not implemented)**

Time = 4.82 (sec) , antiderivative size = 909, normalized size of antiderivative = 5.35

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input

```
int(1/(x^3*atanh(tanh(a + b*x))^3),x)
```

output

```
(4*log(1/(exp(2*a)*exp(2*b*x) + 1))^4 - 16*log(1/(exp(2*a)*exp(2*b*x) + 1))^3*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - 16*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3 + 4*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^4 + 24*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 - 64*b^4*x^4 - 64*b*x*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3 - 256*b^3*x^3*log(1/(exp(2*a)*exp(2*b*x) + 1)) + 256*b^3*x^3*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 64*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))^3 + 192*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 - 192*b*x*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + b^2*x^2*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*384i + b^2*x^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2*atan((log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*1i - log(1/(exp(2*a)*exp(2*b*x) + 1))*1i + b*x*2i)/(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*384i - b^2*x^2*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*atan((log((exp(2*a)*exp(2*b*x))/(e...
```

Reduce [F]

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\operatorname{atanh}(\tanh(bx + a))^3 x^3} dx$$

input `int(1/x^3/atanh(tanh(b*x+a))^3,x)`

output `int(1/(atanh(tanh(a + b*x))**3*x**3),x)`

### 3.111 $\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	846
Mathematica [A] (verified)	847
Rubi [A] (verified)	847
Maple [A] (verified)	849
Fricas [A] (verification not implemented)	850
Sympy [F]	850
Maxima [A] (verification not implemented)	851
Giac [A] (verification not implemented)	851
Mupad [B] (verification not implemented)	852
Reduce [F]	852

#### Optimal result

Integrand size = 15, antiderivative size = 101

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{16x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{15b^2} + \frac{32x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^3} - \frac{128x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{315b^4} + \frac{256 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{3465b^5}$$

output

```
2/3*x^4*arctanh(tanh(b*x+a))^(3/2)/b-16/15*x^3*arctanh(tanh(b*x+a))^(5/2)/
b^2+32/35*x^2*arctanh(tanh(b*x+a))^(7/2)/b^3-128/315*x*arctanh(tanh(b*x+a)
)^(9/2)/b^4+256/3465*arctanh(tanh(b*x+a))^(11/2)/b^5
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2} (1155b^4x^4 - 1848b^3x^3\operatorname{arctanh}(\tanh(a + bx)) + 1584b^2x^2\operatorname{arctanh}(\tanh(a + bx)) - 704bx\operatorname{arctanh}(\tanh(a + bx))^2 + 128\operatorname{arctanh}(\tanh(a + bx))^4)}{3465b^5}$$

input `Integrate[x^4*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2)*(1155*b^4*x^4 - 1848*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 1584*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 704*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4))/(3465*b^5)`

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2599$$

$$\frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{8 \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{3b}$$

$$\downarrow 2599$$

$$\frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{6 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b} \right)}{3b}$$

$$\downarrow 2599$$

$$\begin{array}{c}
 \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \\
 8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a+bx))^{7/2} dx}{5b} \right)}{5b} \right) \\
 \hline
 \frac{3b}{\downarrow} \text{2599} \\
 \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \\
 8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{9/2} dx}{7b} \right)}{5b} \right)}{5b} \right) \\
 \hline
 \frac{3b}{\downarrow} \text{2588} \\
 \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \\
 8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{9/2} dx}{7b} - \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{9/2} dx}{9b^2} \right)}{5b} \right)}{5b} \right) \\
 \hline
 \frac{3b}{\downarrow} \text{15} \\
 \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \\
 8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{99b^2} \right)}{5b} \right)}{5b} \right) \\
 \hline
 \frac{3b}{\downarrow}
 \end{array}$$

input `Int [x^4*sqrt [ArcTanh [Tanh [a + b*x]]], x]`

output

$$\frac{(2x^4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{3/2})}{(3b)} - \frac{(8((2x^3 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{5/2}))}{(5b)} - \frac{(6((2x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{7/2}))}{(7b)} - \frac{(4((2x \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{9/2}))}{(9b)} - \frac{(4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{11/2})}{(99b^2)) / (7b)) / (5b)) / (3b)}$$

### Defintions of rubi rules used

rule 15

$$\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^{(m+1)})/(m+1), x] /; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$$

rule 2588

$$\operatorname{Int}[(u_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$$

rule 2599

$$\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}(v^{(n/(a*(m+1))})], x] - \operatorname{Simp}[b*(n/(a*(m+1))) \operatorname{Int}[u^{(m+1)}v^{(n-1)}, x], x] /; \operatorname{NeQ}[b*u - a*v, 0] /; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m+n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.52

method	result
default	$\frac{2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{11/2}}{11} + \frac{2(-4 \operatorname{arctanh}(\operatorname{tanh}(bx+a))+4bx) \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{9/2}}{9} + \frac{2(2(bx-\operatorname{arctanh}(\operatorname{tanh}(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))) \operatorname{arctanh}(\operatorname{tanh}(bx+a)))^{7/2}}{7}$

input

$$\operatorname{int}(x^4 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{1/2}, x, \operatorname{method}=\_RETURNVERBOSE)$$

output

```
2/b^5*(1/11*arctanh(tanh(b*x+a))^(11/2)+1/9*(-4*arctanh(tanh(b*x+a))+4*b*x
)*arctanh(tanh(b*x+a))^(9/2)+1/7*(2*(b*x-arctanh(tanh(b*x+a)))^2+(2*b*x-2*
arctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(7/2)+2/5*(b*x-arctanh(tanh(
b*x+a)))^2*(2*b*x-2*arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(5/2)+1/3*(
b*x-arctanh(tanh(b*x+a)))^4*arctanh(tanh(b*x+a))^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2(315b^5x^5 + 35ab^4x^4 - 40a^2b^3x^3 + 48a^3b^2x^2 - 64a^4bx + 128a^5)\sqrt{bx+a}}{3465b^5}$$

input

```
integrate(x^4*arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
2/3465*(315*b^5*x^5 + 35*a*b^4*x^4 - 40*a^2*b^3*x^3 + 48*a^3*b^2*x^2 - 64*
a^4*b*x + 128*a^5)*sqrt(b*x + a)/b^5
```

**Sympy [F]**

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x^4 \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input

```
integrate(x**4*atanh(tanh(b*x+a))**(1/2),x)
```

output

```
Integral(x**4*sqrt(atanh(tanh(a + b*x))), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2(315b^5x^5 + 35ab^4x^4 - 40a^2b^3x^3 + 48a^3b^2x^2 - 64a^4bx + 128a^5)\sqrt{bx+a}}{3465b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`output `2/3465*(315*b^5*x^5 + 35*a*b^4*x^4 - 40*a^2*b^3*x^3 + 48*a^3*b^2*x^2 - 64*a^4*b*x + 128*a^5)*sqrt(b*x + a)/b^5`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.49

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\sqrt{2} \left( \frac{11\sqrt{2}(35(bx+a)^{\frac{9}{2}} - 180(bx+a)^{\frac{7}{2}}a + 378(bx+a)^{\frac{5}{2}}a^2 - 420(bx+a)^{\frac{3}{2}}a^3 + 315\sqrt{bx+aa^4})a}{b^4} + \frac{5\sqrt{2}(63(bx+a)^{\frac{11}{2}} - 385(bx+a)^{\frac{9}{2}}a + 990(bx+a)^{\frac{7}{2}}a^2 - 1386(bx+a)^{\frac{5}{2}}a^3 + 1155(bx+a)^{\frac{3}{2}}a^4 - 693\sqrt{bx+a})a^5}{b^4} \right)}{3465b}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`output `1/3465*sqrt(2)*(11*sqrt(2)*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)*a/b^4 + 5*sqrt(2)*(63*(b*x + a)^(11/2) - 385*(b*x + a)^(9/2)*a + 990*(b*x + a)^(7/2)*a^2 - 1386*(b*x + a)^(5/2)*a^3 + 1155*(b*x + a)^(3/2)*a^4 - 693*sqrt(b*x + a)*a^5)/b^4)/b`



**Mupad [B] (verification not implemented)**

Time = 3.03 (sec) , antiderivative size = 811, normalized size of antiderivative = 8.03

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \text{Too large to display}$$

input `int(x^4*atanh(tanh(a + b*x))^(1/2),x)`

output

```
(2*x^5*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)/11 - (x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/11 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/11 + (2*b*x)/11)/(9*b) - (128*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^4*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/11 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/11 + (2*b*x)/11)/(315*b^5) - (8*x^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/11 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/11 + (2*b*x)/11)/(63*b^2) - (64*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/11 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/11 + (2*b*x)/11)/(315*b^4) - (16*x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp...
```

**Reduce [F]**

$$\int x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^4 dx$$

input `int(x^4*atanh(tanh(b*x+a))^(1/2),x)`

output `int(sqrt(atanh(tanh(a + b*x)))*x**4,x)`

### 3.112 $\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result . . . . .	854
Mathematica [A] (verified) . . . . .	855
Rubi [A] (verified) . . . . .	855
Maple [A] (verified) . . . . .	857
Fricas [A] (verification not implemented) . . . . .	857
Sympy [F] . . . . .	858
Maxima [A] (verification not implemented) . . . . .	858
Giac [A] (verification not implemented) . . . . .	859
Mupad [B] (verification not implemented) . . . . .	859
Reduce [F] . . . . .	860

#### Optimal result

Integrand size = 15, antiderivative size = 80

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{4x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b^2} + \frac{16x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^3} - \frac{32 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{315b^4}$$

output

```
2/3*x^3*arctanh(tanh(b*x+a))^(3/2)/b-4/5*x^2*arctanh(tanh(b*x+a))^(5/2)/b^2+16/35*x*arctanh(tanh(b*x+a))^(7/2)/b^3-32/315*arctanh(tanh(b*x+a))^(9/2)/b^4
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.82

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2} (105b^3x^3 - 126b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 72bx\operatorname{arctanh}(\tanh(a + bx))^2 - 16\operatorname{arctanh}(\tanh(a + bx))^3) - 16\operatorname{arctanh}(\tanh(a + bx))^3}{315b^4}$$

input `Integrate[x^3*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2)*(105*b^3*x^3 - 126*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 72*b*x*ArcTanh[Tanh[a + b*x]]^2 - 16*ArcTanh[Tanh[a + b*x]]^3))/ (315*b^4)`

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2599$$

$$\frac{2x^3\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \int x^2\operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{b}$$

$$\downarrow 2599$$

$$\frac{2x^3\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \left( \frac{2x^2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \int x\operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b} \right)}{b}$$

$$\downarrow 2599$$

$$\begin{array}{c}
 \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b} - \frac{3b}{5b} \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{7/2} dx}{7b} \right)}{5b} \right) \\
 \downarrow \text{2588} \\
 \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b} - \frac{3b}{5b} \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{7/2} d \operatorname{arctanh}(\tanh(a+bx))}{7b^2} \right)}{5b} \right) \\
 \downarrow \text{15} \\
 \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b} - \frac{3b}{5b} \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{63b^2} \right)}{5b} \right)
 \end{array}$$

input `Int[x^3*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*x^3*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (2*((2*x^2*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (4*ArcTanh[Tanh[a + b*x]]^(9/2))/(63*b^2)))/(5*b)))/b`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.55

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}} + 2(-3 \operatorname{arctanh}(\tanh(bx+a)) + 3bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}} + 2((bx - \operatorname{arctanh}(\tanh(bx+a)))(2bx - 2 \operatorname{arctanh}(\tanh(bx+a)))}{b^4}$

input

```
int(x^3*arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^4*(1/9*arctanh(tanh(b*x+a))^(9/2)+1/7*(-3*arctanh(tanh(b*x+a))+3*b*x)*arctanh(tanh(b*x+a))^(7/2)+1/5*((b*x-arctanh(tanh(b*x+a)))*(2*b*x-2*arctanh(tanh(b*x+a)))+(b*x-arctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(5/2)+1/3*(b*x-arctanh(tanh(b*x+a)))^3*arctanh(tanh(b*x+a))^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.66

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{2(35b^4x^4 + 5ab^3x^3 - 6a^2b^2x^2 + 8a^3bx - 16a^4)\sqrt{bx + a}}{315b^4}$$

input

```
integrate(x^3*arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output  $2/315*(35*b^4*x^4 + 5*a*b^3*x^3 - 6*a^2*b^2*x^2 + 8*a^3*b*x - 16*a^4)*\text{sqrt}(b*x + a)/b^4$

### Sympy [F]

$$\int x^3 \sqrt{\text{arctanh}(\tanh(a + bx))} dx = \int x^3 \sqrt{\text{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**3*atanh(tanh(b*x+a))**(1/2), x)`

output `Integral(x**3*sqrt(atanh(tanh(a + b*x))), x)`

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.66

$$\begin{aligned} & \int x^3 \sqrt{\text{arctanh}(\tanh(a + bx))} dx \\ &= \frac{2(35b^4x^4 + 5ab^3x^3 - 6a^2b^2x^2 + 8a^3bx - 16a^4)\sqrt{bx + a}}{315b^4} \end{aligned}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")`

output  $2/315*(35*b^4*x^4 + 5*a*b^3*x^3 - 6*a^2*b^2*x^2 + 8*a^3*b*x - 16*a^4)*\text{sqrt}(b*x + a)/b^4$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.56

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\sqrt{2} \left( \frac{9\sqrt{2} \left( 5(bx+a)^{\frac{7}{2}} - 21(bx+a)^{\frac{5}{2}}a + 35(bx+a)^{\frac{3}{2}}a^2 - 35\sqrt{bx+aa^3} \right) a}{b^3} + \frac{\sqrt{2} \left( 35(bx+a)^{\frac{9}{2}} - 180(bx+a)^{\frac{7}{2}}a + 378(bx+a)^{\frac{5}{2}}a^2 - 420(bx+a)^{\frac{3}{2}}a^3 \right)}{b^3} \right)}{315b}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `1/315*sqrt(2)*(9*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a/b^3 + sqrt(2)*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)/b^3)/b`

**Mupad [B] (verification not implemented)**

Time = 3.08 (sec) , antiderivative size = 648, normalized size of antiderivative = 8.10

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \text{Too large to display}$$

input `int(x^3*atanh(tanh(a + b*x))^(1/2),x)`



output

```
(2*x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/9 - (x^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/9 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/9 + (2*b*x)/9))/(7*b) - (16*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^3*(log(2/(exp(2*a)*exp(2*b*x) + 1))/9 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/9 + (2*b*x)/9))/(35*b^4) - (6*x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/9 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/9 + (2*b*x)/9))/(35*b^2) - (8*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^2*(log(2/(exp(2*a)*exp(2*b*x) + 1))/9 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/9 + (2*b*x)/9))/(35*b^3)
```

**Reduce [F]**

$$\int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3 dx$$

input

```
int(x^3*atanh(tanh(b*x+a))^(1/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*x**3,x)
```

### 3.113 $\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	861
Mathematica [A] (verified)	861
Rubi [A] (verified)	862
Maple [A] (verified)	863
Fricas [A] (verification not implemented)	864
Sympy [F]	864
Maxima [A] (verification not implemented)	864
Giac [B] (verification not implemented)	865
Mupad [B] (verification not implemented)	865
Reduce [F]	866

#### Optimal result

Integrand size = 15, antiderivative size = 59

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{8x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{15b^2} + \frac{16 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{105b^3}$$

output

$2/3*x^2*\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}/b-8/15*x*\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}/b^2+16/105*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b^3$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} (35b^2x^2 - 28bx \operatorname{arctanh}(\tanh(a + bx)) + 8 \operatorname{arctanh}(\tanh(a + bx))^2)}{105b^3}$$

input

`Integrate[x^2*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output

$$\frac{(2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{3/2} (35 b^2 x^2 - 28 b x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]] + 8 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^2))}{(105 b^3)}$$
**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{\operatorname{arctanh}(\operatorname{tanh}(a + b x))} dx$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2}}{3b} - \frac{4 \int x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2} dx}{3b}$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2} dx}{5b} \right)}{3b}$$

$$\downarrow 2588$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2} d \operatorname{arctanh}(\operatorname{tanh}(a + b x))}{5b^2} \right)}{3b}$$

$$\downarrow 15$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2}}{35b^2} \right)}{3b}$$

input

$$\operatorname{Int}[x^2 \operatorname{Sqrt}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]], x]$$

output

$$\frac{(2 x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{3/2})}{(3 b)} - \frac{(4 ((2 x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{5/2}) / (5 b) - (4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{7/2}) / (35 b^2)))}{(3 b)}$$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_.)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}} + 2(2bx-2 \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}} + 2(bx-\operatorname{arctanh}(\tanh(bx+a)))^2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{b^3}$	69

input `int(x^2*arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2/b^3*(1/7*arctanh(tanh(b*x+a))^(7/2)+1/5*(2*b*x-2*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^(5/2)+1/3*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(3/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.71

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(15b^3x^3 + 3ab^2x^2 - 4a^2bx + 8a^3)\sqrt{bx + a}}{105b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`output `2/105*(15*b^3*x^3 + 3*a*b^2*x^2 - 4*a^2*b*x + 8*a^3)*sqrt(b*x + a)/b^3`**Sympy [F]**

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x^2 \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**2*atanh(tanh(b*x+a))**(1/2),x)`output `Integral(x**2*sqrt(atanh(tanh(a + b*x))), x)`**Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.71

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(15b^3x^3 + 3ab^2x^2 - 4a^2bx + 8a^3)\sqrt{bx + a}}{105b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`output `2/105*(15*b^3*x^3 + 3*a*b^2*x^2 - 4*a^2*b*x + 8*a^3)*sqrt(b*x + a)/b^3`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 102 vs.  $2(47) = 94$ .

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.73

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\sqrt{2} \left( \frac{7\sqrt{2} \left( 3(bx+a)^{\frac{5}{2}} - 10(bx+a)^{\frac{3}{2}}a + 15\sqrt{bx+aa^2} \right) a}{b^2} + \frac{3\sqrt{2} \left( 5(bx+a)^{\frac{7}{2}} - 21(bx+a)^{\frac{5}{2}}a + 35(bx+a)^{\frac{3}{2}}a^2 - 35\sqrt{bx+aa^3} \right)}{b^2} \right)}{105b}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `1/105*sqrt(2)*(7*sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2)*a/b^2 + 3*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)/b^2)/b`

**Mupad [B] (verification not implemented)**

Time = 3.10 (sec) , antiderivative size = 485, normalized size of antiderivative = 8.22

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \text{Too large to display}$$

input `int(x^2*atanh(tanh(a + b*x))^(1/2),x)`

output

```
(2*x^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/7 - (x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/7 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/7 + (2*b*x)/7))/(5*b) - (8*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^2*(log(2/(exp(2*a)*exp(2*b*x) + 1))/7 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/7 + (2*b*x)/7))/(15*b^3) - (4*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/7 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/7 + (2*b*x)/7))/(15*b^2)
```

**Reduce [F]**

$$\int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2 dx$$

input

```
int(x^2*atanh(tanh(b*x+a))^(1/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*x**2,x)
```

### 3.114 $\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	867
Mathematica [A] (verified)	867
Rubi [A] (verified)	868
Maple [A] (verified)	869
Fricas [A] (verification not implemented)	869
Sympy [F]	870
Maxima [A] (verification not implemented)	870
Giac [B] (verification not implemented)	870
Mupad [B] (verification not implemented)	871
Reduce [B] (verification not implemented)	871

#### Optimal result

Integrand size = 13, antiderivative size = 38

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{15b^2}$$

output `2/3*x*arctanh(tanh(b*x+a))^(3/2)/b-4/15*arctanh(tanh(b*x+a))^(5/2)/b^2`

#### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(5bx - 2 \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{15b^2}$$

input `Integrate[x*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*(5*b*x - 2*ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))/(15*b^2)`



**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2599$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{3b}$$

$$\downarrow 2588$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{3/2} d \operatorname{arctanh}(\tanh(a + bx))}{3b^2}$$

$$\downarrow 15$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{15b^2}$$

input `Int[x*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*x*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (4*ArcTanh[Tanh[a + b*x]]^(5/2))/(15*b^2)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5} + \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{b^2}$	42

input

```
int(x*arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^2*(1/5*arctanh(tanh(b*x+a))^(5/2)+1/3*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(3b^2x^2 + abx - 2a^2)\sqrt{bx + a}}{15b^2}$$

input

```
integrate(x*arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
2/15*(3*b^2*x^2 + a*b*x - 2*a^2)*sqrt(b*x + a)/b^2
```

**Sympy [F]**

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x*atanh(tanh(b*x+a))**(1/2), x)`

output `Integral(x*sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(3b^2x^2 + abx - 2a^2)\sqrt{bx + a}}{15b^2}$$

input `integrate(x*arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")`

output `2/15*(3*b^2*x^2 + a*b*x - 2*a^2)*sqrt(b*x + a)/b^2`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs. 2(30) = 60.

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.97

$$\begin{aligned} & \int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx \\ &= \frac{\sqrt{2} \left( \frac{5\sqrt{2}((bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa})a}{b} + \frac{\sqrt{2}(3(bx+a)^{\frac{5}{2}} - 10(bx+a)^{\frac{3}{2}}a + 15\sqrt{bx+aa^2})}{b} \right)}{15b} \end{aligned}$$

input `integrate(x*arctanh(tanh(b*x+a))^(1/2), x, algorithm="giac")`

output

$$\frac{1}{15}\sqrt{2}\left(5\sqrt{2}\left((bx+a)^{3/2}-3\sqrt{bx+a}\right)a/b+\sqrt{2}\right)\left(3(bx+a)^{5/2}-10(bx+a)^{3/2}a+15\sqrt{bx+a}a^2\right)/b$$

**Mupad [B] (verification not implemented)**

Time = 3.19 (sec) , antiderivative size = 151, normalized size of antiderivative = 3.97

$$\int x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}dx$$

$$= \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2}-\frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}\left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)-\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)\right)\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)-\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)\right)+5}{15b^2}$$

input

```
int(x*atanh(tanh(a + b*x))^(1/2),x)
```

output

$$\left(\frac{\log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx)+1}\right)}{2}-\log\left(\frac{2}{\exp(2a)\exp(2bx)+1}\right)\right)^{1/2}\left(\log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx)+1}\right)-\log\left(\frac{2}{\exp(2a)\exp(2bx)+1}\right)\right)\left(\log\left(\frac{2}{\exp(2a)\exp(2bx)+1}\right)-\log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx)+1}\right)+5bx\right)/\left(15b^2\right)$$

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}dx$$

$$= \frac{2\sqrt{\operatorname{atanh}(\tanh(bx+a))}\operatorname{atanh}(\tanh(bx+a))(-2\operatorname{atanh}(\tanh(bx+a))+5bx)}{15b^2}$$

input

```
int(x*atanh(tanh(b*x+a))^(1/2),x)
```

output

$$\left(2\sqrt{\operatorname{atanh}(\tanh(a+b*x))}\operatorname{atanh}(\tanh(a+b*x))\left(-2\operatorname{atanh}(\tanh(a+b*x))+5b*x\right)\right)/\left(15b^2\right)$$

### 3.115 $\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	872
Mathematica [A] (verified)	872
Rubi [A] (verified)	873
Maple [A] (verified)	874
Fricas [A] (verification not implemented)	874
Sympy [A] (verification not implemented)	874
Maxima [A] (verification not implemented)	875
Giac [A] (verification not implemented)	875
Mupad [B] (verification not implemented)	875
Reduce [B] (verification not implemented)	876

#### Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b}$$

output

```
2/3*arctanh(tanh(b*x+a))^(3/2)/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b}$$

input

```
Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]],x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b)
```

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b}$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3b}$	15
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3b}$	15

input `int(arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`output `2/3*arctanh(tanh(b*x+a))^(3/2)/b`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(bx + a)^{\frac{3}{2}}}{3b}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`output `2/3*(b*x + a)^(3/2)/b`**Sympy [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.44

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \begin{cases} \frac{2 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a+bx))}{3b} & \text{for } b \neq 0 \\ x \sqrt{\operatorname{atanh}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**(1/2),x)`

output `Piecewise((2*atanh(tanh(a + b*x))**(3/2)/(3*b), Ne(b, 0)), (x*sqrt(atanh(tanh(a))), True))`

### Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2(bx + a)^{\frac{3}{2}}}{3b}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `2/3*(b*x + a)^(3/2)/b`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\sqrt{2}(2bx + 2a)^{\frac{3}{2}}}{6b}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `1/6*sqrt(2)*(2*b*x + 2*a)^(3/2)/b`

### Mupad [B] (verification not implemented)

Time = 3.19 (sec) , antiderivative size = 95, normalized size of antiderivative = 5.28

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= - \frac{\left( \ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) \right) \sqrt{\frac{\ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)}{2} - \frac{\ln \left( \frac{1}{e^{2a} e^{2bx} + 1} \right)}{2}}}{3b}$$



input `int(atanh(tanh(a + b*x))^(1/2),x)`

output `-((log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*b)`

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{3b}$$

input `int(atanh(tanh(b*x+a))^(1/2),x)`

output `(2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)))/(3*b)`

**3.116**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$

Optimal result	877
Mathematica [A] (verified)	877
Rubi [A] (verified)	878
Maple [A] (verified)	879
Fricas [A] (verification not implemented)	879
Sympy [F]	880
Maxima [F]	880
Giac [A] (verification not implemented)	880
Mupad [B] (verification not implemented)	881
Reduce [F]	881

**Optimal result**

Integrand size = 15, antiderivative size = 63

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$$

$$= -2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ 2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}$$

output

`-2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(1/2)+2*arctanh(tanh(b*x+a))^(1/2)`

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx$$

$$= 2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}$$

$$- 2\operatorname{arctanh} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}$$

input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x,x]`

output `2*Sqrt[ArcTanh[Tanh[a + b*x]]] - 2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx$$

$$\downarrow \text{2590}$$

$$2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow \text{2592}$$

$$2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))} \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x,x]`

output `-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]] *Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]]`

**Defintions of rubi rules used**

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a]] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86

method	result
default	$2\sqrt{\operatorname{arctanh}(\tanh(bx+a))} - 2\sqrt{\operatorname{arctanh}(\tanh(bx+a)) - bx} \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a)) - bx}}\right)$

input `int(arctanh(tanh(b*x+a))^(1/2)/x,x,method=_RETURNVERBOSE)`

output `2*arctanh(tanh(b*x+a))^(1/2)-2*(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} dx = \left[ \sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) + 2\sqrt{bx+a}, 2\sqrt{-a} \operatorname{arctan}\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + 2\sqrt{bx+a} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x,x, algorithm="fricas")`

output

```
[sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*sqrt(b*x + a), 2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + 2*sqrt(b*x + a)]
```

**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{x} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(1/2)/x,x)
```

output

```
Integral(sqrt(atanh(tanh(a + b*x)))/x, x)
```

**Maxima [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx = \int \frac{\sqrt{\operatorname{artanh}(\tanh(bx + a))}}{x} dx$$

input

```
integrate(arctanh(tanh(b*x+a))^(1/2)/x,x, algorithm="maxima")
```

output

```
integrate(sqrt(arctanh(tanh(b*x + a)))/x, x)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx = \sqrt{2} \left( \frac{\sqrt{2}a \operatorname{arctan}\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + \sqrt{2}\sqrt{bx+a} \right)$$

input

```
integrate(arctanh(tanh(b*x+a))^(1/2)/x,x, algorithm="giac")
```

output

```
sqrt(2)*(sqrt(2)*a*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + sqrt(2)*sqrt(b*x + a))
```

**Mupad [B] (verification not implemented)**

Time = 4.42 (sec) , antiderivative size = 308, normalized size of antiderivative = 4.89

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx = 2 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}$$

$$+ \ln\left(-\frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2 \sqrt{\frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{2}}{\frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{2}}}{x \sqrt{\frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{2}} - bx}\right)$$

input

```
int(atanh(tanh(a + b*x))^(1/2)/x,x)
```

output

```
2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2) + log(-log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2 - b*x)^(1/2) + b*x/(x*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2 - b*x)^(1/2))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2 - b*x)^(1/2)
```

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x} dx$$

input

```
int(atanh(tanh(b*x+a))^(1/2)/x,x)
```

output `int(sqrt(atanh(tanh(a + b*x)))/x,x)`

**3.117**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx$

Optimal result . . . . .	883
Mathematica [A] (verified) . . . . .	883
Rubi [A] (verified) . . . . .	884
Maple [A] (verified) . . . . .	885
Fricas [A] (verification not implemented) . . . . .	886
Sympy [F] . . . . .	886
Maxima [F] . . . . .	886
Giac [A] (verification not implemented) . . . . .	887
Mupad [B] (verification not implemented) . . . . .	887
Reduce [F] . . . . .	888

**Optimal result**

Integrand size = 15, antiderivative size = 66

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = \frac{b \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x}$$

output `b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a))^(1/2)-arctanh(tanh(b*x+a))^(1/2)/x`

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.98

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = -\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} - \frac{b \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}$$



input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^2,x]`

output `-(Sqrt[ArcTanh[Tanh[a + b*x]]]/x) - (b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]`

### Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2599, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx$$

↓ 2599

$$\frac{1}{2}b \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x}$$

↓ 2592

$$\frac{b \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x}$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^2,x]`

output `(b*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] - Sqrt[ArcTanh[Tanh[a + b*x]]]/x`

## Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.95

method	result	size
default	$2b \left( -\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2bx} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}} \right)$	63

input `int(arctanh(tanh(b*x+a))^(1/2)/x^2,x,method=_RETURNVERBOSE)`

output `2*b*(-1/2*arctanh(tanh(b*x+a))^(1/2)/b/x-1/2/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.36

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx$$

$$= \left[ \frac{\sqrt{abx} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2\sqrt{bx+aa}}{2ax}, \frac{\sqrt{-abx} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) - \sqrt{bx+aa}}{ax} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^2,x, algorithm="fricas")`output `[1/2*(sqrt(a)*b*x*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*sqrt(b*x + a)*a)/(a*x), (sqrt(-a)*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) - sqrt(b*x + a)*a)/(a*x)]`**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**2,x)`output `Integral(sqrt(atanh(tanh(a + b*x)))/x**2, x)`**Maxima [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = \int \frac{\sqrt{\operatorname{artanh}(\tanh(bx + a))}}{x^2} dx$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^2,x, algorithm="maxima")`output `integrate(sqrt(arctanh(tanh(b*x + a)))/x^2, x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = \frac{1}{2} \sqrt{2} b \left( \frac{\sqrt{2} \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} - \frac{\sqrt{2} \sqrt{bx+a}}{bx} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^2,x, algorithm="giac")`output `1/2*sqrt(2)*b*(sqrt(2)*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) - sqrt(2)*sqrt(b*x + a)/(b*x))`**Mupad [B] (verification not implemented)**

Time = 9.22 (sec) , antiderivative size = 341, normalized size of antiderivative = 5.17

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx = -\frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{x}$$

$$+ \frac{\sqrt{2} b \ln \left( \frac{\sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)} - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \left( \sqrt{2}bx - \sqrt{2} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right) + \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2}} \right)}{x} \right)}{2 \sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)} - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx}$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^2,x)`

output

```
(2^(1/2)*b*log((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*
b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*((log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/
2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*
a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x
) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)
+ 2^(1/2)*b*x)*1i)/x)*1i)/(2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)) - (log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*
x) + 1))/2)^(1/2)/x
```

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx$$

$$= \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \left( \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))x} dx \right) bx}{2x}$$

input

```
int(atanh(tanh(b*x+a))^(1/2)/x^2,x)
```

output

```
( - 2*sqrt(atanh(tanh(a + b*x))) + int(sqrt(atanh(tanh(a + b*x)))/(atanh(t
anh(a + b*x))*x),x)*b*x)/(2*x)
```

**3.118**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx$

Optimal result . . . . .	889
Mathematica [A] (verified) . . . . .	890
Rubi [A] (verified) . . . . .	890
Maple [A] (verified) . . . . .	892
Fricas [A] (verification not implemented) . . . . .	893
Sympy [F] . . . . .	893
Maxima [F] . . . . .	894
Giac [A] (verification not implemented) . . . . .	894
Mupad [B] (verification not implemented) . . . . .	894
Reduce [F] . . . . .	895

**Optimal result**

Integrand size = 15, antiderivative size = 125

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx$$

$$= \frac{b^2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx-\operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{b}{4x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{b^2}{4(bx-\operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$- \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2}$$

output

```
1/4*b^2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^(3/2)-1/4*b/x/arctanh(tanh(b*x+a))^(1/2)+1/4*
b^2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)-1/2*arctanh(tanh
(b*x+a))^(1/2)/x^2
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.71

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx$$

$$= \frac{1}{4} \left( \frac{\left(-2 + \frac{bx}{bx - \operatorname{arctanh}(\tanh(a + bx))}\right) \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} + \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} \right)$$

input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^3,x]`

output `(((-2 + (b*x)/(b*x - ArcTanh[Tanh[a + b*x]]))*Sqrt[ArcTanh[Tanh[a + b*x]]])/x^2 + (b^2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2))/4`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx$$

$$\downarrow \text{2599}$$

$$\frac{1}{4} b \int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2x^2}$$

$$\begin{aligned}
 & \downarrow 2599 \\
 & \frac{1}{4}b \left( -\frac{1}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
 & \qquad \qquad \qquad \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2} \\
 & \downarrow 2594 \\
 & \frac{1}{4}b \left( -\frac{1}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
 & \qquad \qquad \qquad \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2} \\
 & \downarrow 2592 \\
 & \frac{1}{4}b \left( -\frac{1}{2}b \left( -\frac{2 \arctan \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
 & \qquad \qquad \qquad \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2}
 \end{aligned}$$

input `Int [Sqrt [ArcTanh [Tanh [a + b*x]]]/x^3,x]`

output `(b*(-1/2*(b*((-2*ArcTan [Sqrt [ArcTanh [Tanh [a + b*x]]]/Sqrt [b*x - ArcTanh [Tanh [a + b*x]]]))/(b*x - ArcTanh [Tanh [a + b*x]]^(3/2) - 2/((b*x - ArcTanh [Tanh [a + b*x]])*Sqrt [ArcTanh [Tanh [a + b*x]]])) - 1/(x*Sqrt [ArcTanh [Tanh [a + b*x]]])))/4 - Sqrt [ArcTanh [Tanh [a + b*x]]]/(2*x^2)`



## Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.74

method	result	size
default	$2b^2 \left( \frac{-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{8(\operatorname{arctanh}(\tanh(bx+a))-bx)} - \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{8}}{b^2x^2} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{8(\operatorname{arctanh}(\tanh(bx+a))-bx)^{\frac{3}{2}}}\right)$	92

input `int(arctanh(tanh(b*x+a))^(1/2)/x^3,x,method=_RETURNVERBOSE)`

output `2*b^2*((-1/8/(arctanh(tanh(b*x+a))-b*x)*arctanh(tanh(b*x+a))^(3/2)-1/8*arctanh(tanh(b*x+a))^(1/2))/b^2/x^2+1/8/(arctanh(tanh(b*x+a))-b*x)^(3/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx$$

$$= \left[ \frac{\sqrt{ab^2x^2} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(abx + 2a^2)\sqrt{bx+a}}{8a^2x^2}, \right. \\ \left. - \frac{\sqrt{-ab^2x^2} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (abx + 2a^2)\sqrt{bx+a}}{4a^2x^2} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^3,x, algorithm="fricas")`

output `[1/8*(sqrt(a)*b^2*x^2*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(a*b*x + 2*a^2)*sqrt(b*x + a))/(a^2*x^2), -1/4*(sqrt(-a)*b^2*x^2*arctan(sqrt(-a)/sqrt(b*x + a)) + (a*b*x + 2*a^2)*sqrt(b*x + a))/(a^2*x^2)]`

**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{x^3} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**3,x)`

output `Integral(sqrt(atanh(tanh(a + b*x)))/x**3, x)`

**Maxima [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx = \int \frac{\sqrt{\operatorname{artanh}(\tanh(bx + a))}}{x^3} dx$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^3,x, algorithm="maxima")`

output `integrate(sqrt(arctanh(tanh(b*x + a)))/x^3, x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx = -\frac{\sqrt{2} \left( \frac{\sqrt{2} b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{\sqrt{2} \left( (bx+a)^{\frac{3}{2}} b^3 + \sqrt{bx+aab^3} \right)}{ab^2 x^2} \right)}{8b}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^3,x, algorithm="giac")`

output `-1/8*sqrt(2)*(sqrt(2)*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a) + sqrt(2)*((b*x + a)^(3/2)*b^3 + sqrt(b*x + a)*a*b^3)/(a*b^2*x^2))/b`

**Mupad [B] (verification not implemented)**

Time = 7.99 (sec) , antiderivative size = 741, normalized size of antiderivative = 5.93

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^3,x)`

output

```
(b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(2*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) - ((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(x^2*(2*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x)) + (2^(1/2)*b^2*log(((2*2^(1/2)*a + log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*4i)/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2))*i)/(4*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/...
```

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \left( \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))x^2} dx \right) b x^2}{4x^2}$$

input

```
int(atanh(tanh(b*x+a))^(1/2)/x^3,x)
```

output

```
( - 2*sqrt(atanh(tanh(a + b*x))) + int(sqrt(atanh(tanh(a + b*x)))/(atanh(atanh(a + b*x))*x**2),x)*b*x**2)/(4*x**2)
```

**3.119**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$

Optimal result	896
Mathematica [A] (verified)	897
Rubi [A] (verified)	897
Maple [A] (verified)	899
Fricas [A] (verification not implemented)	900
Sympy [F]	900
Maxima [F]	901
Giac [A] (verification not implemented)	901
Mupad [B] (verification not implemented)	902
Reduce [F]	902

**Optimal result**

Integrand size = 15, antiderivative size = 179

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$$

$$= \frac{b^3 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} + \frac{b^2}{24x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{24(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b}$$

$$- \frac{12x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^3}$$

$$+ \frac{8(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$- \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^3}$$

output

```
1/8*b^3*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))
)/(b*x-arctanh(tanh(b*x+a)))^(5/2)+1/24*b^2/x/arctanh(tanh(b*x+a))^(3/2)-1
/24*b^3/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)-1/12*b/x^2/a
rctanh(tanh(b*x+a))^(1/2)+1/8*b^3/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tan
h(b*x+a))^(1/2)-1/3*arctanh(tanh(b*x+a))^(1/2)/x^3
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx = \frac{1}{24} \left( -\frac{3b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} \right. \\ \left. + \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}(-3b^2x^2 + 14bx \operatorname{arctanh}(\tanh(a+bx)) - 8 \operatorname{arctanh}(\tanh(a+bx))^2)}{x^3(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2} \right)$$

input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^4,x]`

output `((-3*b^3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2) + (Sqrt[ArcTanh[Tanh[a + b*x]]]*(-3*b^2*x^2 + 14*b*x*ArcTanh[Tanh[a + b*x]] - 8*ArcTanh[Tanh[a + b*x]]^2))/(x^3*(-(b*x) + ArcTanh[Tanh[a + b*x]]^2))/24`

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx \\ \downarrow 2599 \\ \frac{1}{6}b \int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^3} \\ \downarrow 2599$$

$$\frac{1}{6}b \left( -\frac{1}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2599

$$\frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2594

$$\frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2594

$$\frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2592

$$\frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input Int [Sqrt [ArcTanh [Tanh [a + b\*x]]] /x^4, x]

output

$$\begin{aligned} & (b*(-1/4*(b*((-3*b*(-(((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - \\ & ArcTanh[Tanh[a + b*x]]]])))/(b*x - ArcTanh[Tanh[a + b*x]])^{(3/2)} - 2/((b*x - \\ & ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])))/(b*x - ArcTanh[Tan \\ & h[a + b*x])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]] \\ & ^{(3/2)})))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^{(3/2)}) - 1/(2*x^2*Sqrt[ArcTanh \\ & [Tanh[a + b*x]]])))/6 - Sqrt[ArcTanh[Tanh[a + b*x]]]/(3*x^3) \end{aligned}$$

### Defintions of rubi rules used

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.03

method	result
default	$2b^3 \left( \frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{16a^2+32a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)+16(\operatorname{arctanh}(\tanh(bx+a))-bx-a)^2} - \frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{6(\operatorname{arctanh}(\tanh(bx+a))-bx)} - \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{16} \right)$



input `int(arctanh(tanh(b*x+a))^(1/2)/x^4,x,method=_RETURNVERBOSE)`

output `2*b^3*((1/16/(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(5/2)-1/6/(arctanh(tanh(b*x+a))-b*x)*arctanh(tanh(b*x+a))^(3/2)-1/16*arctanh(tanh(b*x+a))^(1/2))/b^3/x^3-1/16/(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx$$

$$= \left[ \frac{3\sqrt{ab^3x^3} \log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(3ab^2x^2 - 2a^2bx - 8a^3)\sqrt{bx+a}}{48a^3x^3}, \frac{3\sqrt{-ab^3x^3} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (3a^2bx^2 - 2a^2bx - 8a^3)\sqrt{bx+a}}{48a^3x^3} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^4,x, algorithm="fricas")`

output `[1/48*(3*sqrt(a)*b^3*x^3*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(3*a*b^2*x^2 - 2*a^2*b*x - 8*a^3)*sqrt(b*x + a))/(a^3*x^3), 1/24*(3*sqrt(-a)*b^3*x^3*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b^2*x^2 - 2*a^2*b*x - 8*a^3)*sqrt(b*x + a))/(a^3*x^3)]`

### Sympy [F]

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a+bx))}}{x^4} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**4,x)`

output `Integral(sqrt(atanh(tanh(a + b*x)))/x**4, x)`

### Maxima [F]

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^4} dx = \int \frac{\sqrt{\operatorname{arctanh}(\tanh(bx + a))}}{x^4} dx$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^4,x, algorithm="maxima")`

output `integrate(sqrt(arctanh(tanh(b*x + a)))/x^4, x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.45

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^4} dx$$

$$= \frac{1}{48} \sqrt{2} b^3 \left( \frac{3 \sqrt{2} \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} + \frac{\sqrt{2} \left( 3 (bx + a)^{\frac{5}{2}} - 8 (bx + a)^{\frac{3}{2}} a - 3 \sqrt{bx + aa^2} \right)}{a^2 b^3 x^3} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^4,x, algorithm="giac")`

output `1/48*sqrt(2)*b^3*(3*sqrt(2)*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) + sqrt(2)*(3*(b*x + a)^(5/2) - 8*(b*x + a)^(3/2)*a - 3*sqrt(b*x + a)*a^2)/(a^2*b^3*x^3)`

**Mupad [B] (verification not implemented)**

Time = 7.57 (sec) , antiderivative size = 964, normalized size of antiderivative = 5.39

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^4} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^4,x)`

output

$$\begin{aligned} & (b \cdot (\log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) / 2 - \log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) / 2^{(1/2)} / (3x^2 \cdot (2 \cdot \log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) \\ & - 2 \cdot \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1)) + 4bx) + (b^2 \cdot (\log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) / 2 - \log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) / 2^{(1/2)} / (2x \cdot (\log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) - 1 \\ & \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx)^2 + (2^{(1/2)} \cdot b^3 \cdot \log(((\log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) / 2 - 1 \\ & \log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) / 2^{(1/2)} \cdot (\log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) \\ & - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx)^{(1/2)} \cdot 2 \\ & i - 2^{(1/2)} \cdot (\log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx) + 2^{(1/2)} \cdot b \cdot ((2a - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) + \log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) + 2bx)^5 + 40a^2 \cdot (2a - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) + \log(2 / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx)^3 - 80a^3 \cdot (2a - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) + \log(2 / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx)^2 - 32a^5 - 10a \cdot (2a - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) + \log(2 / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx)^4 + 80a^4 \cdot (2a - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1))) + \log(2 / (\exp(2a) \cdot \exp(2bx) + 1)) + 2bx) \cdot 4i / (x \cdot (\log(2 / (\exp(2a) \cdot \exp(2bx) + 1))) - \log((2 \cdot \exp(2a) \cdot \exp(2bx)) / (\exp(2a) \cdot \exp(2bx) + 1)) + 2 \dots \end{aligned}$$
**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^4} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{6x^3} + \left( \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))x^3} dx \right) b x^3$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^4,x)`

output  $(-2\sqrt{\operatorname{atanh}(\tanh(a + bx))} + \int(\sqrt{\operatorname{atanh}(\tanh(a + bx))}/(\operatorname{atanh}(\operatorname{atanh}(a + bx))x^3), x) * b * x^3)/(6 * x^3)$

### 3.120 $\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	904
Mathematica [A] (verified)	904
Rubi [A] (verified)	905
Maple [A] (verified)	907
Fricas [A] (verification not implemented)	908
Sympy [F]	908
Maxima [A] (verification not implemented)	908
Giac [B] (verification not implemented)	909
Mupad [B] (verification not implemented)	909
Reduce [F]	910

#### Optimal result

Integrand size = 15, antiderivative size = 101

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{16x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^2} + \frac{32x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{105b^3} - \frac{128x \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{1155b^4} + \frac{256 \operatorname{arctanh}(\tanh(a + bx))^{13/2}}{15015b^5}$$

output

$2/5*x^4*\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}/b-16/35*x^3*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b^2+32/105*x^2*\operatorname{arctanh}(\tanh(b*x+a))^{(9/2)}/b^3-128/1155*x*\operatorname{arctanh}(\tanh(b*x+a))^{(11/2)}/b^4+256/15015*\operatorname{arctanh}(\tanh(b*x+a))^{(13/2)}/b^5$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} (3003b^4x^4 - 3432b^3x^3 \operatorname{arctanh}(\tanh(a + bx)) + 2288b^2x^2 \operatorname{arctanh}(\tanh(a + bx)))}{15015b^5}$$

input `Integrate[x^4*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2)*(3003*b^4*x^4 - 3432*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 2288*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 832*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4)/(15015*b^5)`

### Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{8 \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{6 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b} \right)}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{9/2} dx}{9b} \right)}{7b} \right)}{5b} \\
 & \quad \downarrow 2599
 \end{aligned}$$

$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{11/2} dx}{9b} \right)}{7b} \right)}{7b} \right)$$

5b

↓ 2588

$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{11/2} dx}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{11/2} dx}{11b^2} \right)}{7b} \right)}{7b} \right)$$

5b

↓ 15

$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{13/2}}{143b^2} \right)}{7b} \right)}{7b} \right)$$

5b

input

`Int [x^4*ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output

`(2*x^4*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (8*((2*x^3*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (6*((2*x^2*ArcTanh[Tanh[a + b*x]]^(9/2))/(9*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(11/2))/(11*b) - (4*ArcTanh[Tanh[a + b*x]]^(13/2))/(143*b^2)))/(9*b)))/(7*b)))/(5*b)`

## Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.52

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{13}{2}}}{13} + \frac{2(-4 \operatorname{arctanh}(\tanh(bx+a))+4bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{11}{2}}}{11} + \frac{2(2(bx-\operatorname{arctanh}(\tanh(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2)}{9}$

input `int(x^4*arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2/b^5*(1/13*arctanh(tanh(b*x+a))^(13/2)+1/11*(-4*arctanh(tanh(b*x+a))+4*b*x)*arctanh(tanh(b*x+a))^(11/2)+1/9*(2*(b*x-arctanh(tanh(b*x+a)))^2+(2*b*x-2*arctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(9/2)+2/7*(b*x-arctanh(tanh(b*x+a)))^2*(2*b*x-2*arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(7/2)+1/5*(b*x-arctanh(tanh(b*x+a)))^4*arctanh(tanh(b*x+a))^(5/2))`



**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.74

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(1155b^6x^6 + 1470ab^5x^5 + 35a^2b^4x^4 - 40a^3b^3x^3 + 48a^4b^2x^2 - 64a^5bx + 128a^6)\sqrt{bx+a}}{15015b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `2/15015*(1155*b^6*x^6 + 1470*a*b^5*x^5 + 35*a^2*b^4*x^4 - 40*a^3*b^3*x^3 + 48*a^4*b^2*x^2 - 64*a^5*b*x + 128*a^6)*sqrt(b*x + a)/b^5`

**Sympy [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int x^4 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx)) dx$$

input `integrate(x**4*atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(x**4*atanh(tanh(a + b*x))**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(1155b^5x^5 + 315ab^4x^4 - 280a^2b^3x^3 + 240a^3b^2x^2 - 192a^4bx + 128a^5)(bx+a)^{\frac{3}{2}}}{15015b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output

$$2/15015*(1155*b^5*x^5 + 315*a*b^4*x^4 - 280*a^2*b^3*x^3 + 240*a^3*b^2*x^2 - 192*a^4*b*x + 128*a^5)*(b*x + a)^{(3/2)}/b^5$$

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 241 vs.  $2(81) = 162$ .

Time = 0.11 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.39

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{2} \left( \frac{143 \sqrt{2} (35 (bx+a)^{9/2} - 180 (bx+a)^{7/2} a + 378 (bx+a)^{5/2} a^2 - 420 (bx+a)^{3/2} a^3 + 315 \sqrt{bx+aa^4}) a^2}{b^4} + \frac{130 \sqrt{2} (63 (bx+a)^{11/2} - 385 (bx+a)^{9/2} a + 990 (bx+a)^{7/2} a^2 - 1386 (bx+a)^{5/2} a^3 + 1155 (bx+a)^{3/2} a^4 - 693 \sqrt{bx+a} a^5) a}{b^4} + 15 \sqrt{2} (231 (bx+a)^{13/2} - 1638 (bx+a)^{11/2} a + 5005 (bx+a)^{9/2} a^2 - 8580 (bx+a)^{7/2} a^3 + 9009 (bx+a)^{5/2} a^4 - 6006 (bx+a)^{3/2} a^5 + 3003 \sqrt{bx+a} a^6) / b^4 \right)}{b}$$

input

```
integrate(x^4*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")
```

output

$$\frac{1/45045*\sqrt{2}*(143*\sqrt{2}*(35*(b*x + a)^{(9/2)} - 180*(b*x + a)^{(7/2)}*a + 378*(b*x + a)^{(5/2)}*a^2 - 420*(b*x + a)^{(3/2)}*a^3 + 315*\sqrt{b*x + a}*a^4)*a^2/b^4 + 130*\sqrt{2}*(63*(b*x + a)^{(11/2)} - 385*(b*x + a)^{(9/2)}*a + 990*(b*x + a)^{(7/2)}*a^2 - 1386*(b*x + a)^{(5/2)}*a^3 + 1155*(b*x + a)^{(3/2)}*a^4 - 693*\sqrt{b*x + a}*a^5)*a/b^4 + 15*\sqrt{2}*(231*(b*x + a)^{(13/2)} - 1638*(b*x + a)^{(11/2)}*a + 5005*(b*x + a)^{(9/2)}*a^2 - 8580*(b*x + a)^{(7/2)}*a^3 + 9009*(b*x + a)^{(5/2)}*a^4 - 6006*(b*x + a)^{(3/2)}*a^5 + 3003*\sqrt{b*x + a}*a^6)/b^4)/b}{b}$$

**Mupad [B] (verification not implemented)**

Time = 3.23 (sec) , antiderivative size = 1813, normalized size of antiderivative = 17.95

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \text{Too large to display}$$

input

```
int(x^4*atanh(tanh(a + b*x))^(3/2),x)
```

output

```
(2*b*x^6*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2
/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/13 + (x^5*(log((2*exp(2*a)*exp(2*b*x
))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)
)*((24*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))
/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/13 - 2*b*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))
/(11*b) + (x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 -
log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) +
1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/2
+ (10*((24*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/13 - 2*b*(log(2/(exp(2*a)*exp(2*
b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*
x))*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp
(2*a)*exp(2*b*x) + 1))/2 + b*x))/(11*b)))/(9*b) + (128*(log((2*exp(2*a)*ex
p(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/
2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))
/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/2 + (10*((24*b*(log(2/(exp(2*a)*exp(
2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2
+ b*x))/13 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2
*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x)...
```

**Reduce [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a)) x^4 dx$$

input

```
int(x^4*atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*x**4,x)
```

### 3.121 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	911
Mathematica [A] (verified)	911
Rubi [A] (verified)	912
Maple [A] (verified)	914
Fricas [A] (verification not implemented)	914
Sympy [F]	915
Maxima [A] (verification not implemented)	915
Giac [B] (verification not implemented)	915
Mupad [B] (verification not implemented)	916
Reduce [F]	917

#### Optimal result

Integrand size = 15, antiderivative size = 80

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{12x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^2} + \frac{16x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{105b^3} - \frac{32 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{1155b^4}$$

output

```
2/5*x^3*arctanh(tanh(b*x+a))^(5/2)/b-12/35*x^2*arctanh(tanh(b*x+a))^(7/2)/
b^2+16/105*x*arctanh(tanh(b*x+a))^(9/2)/b^3-32/1155*arctanh(tanh(b*x+a))^(
11/2)/b^4
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.82

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} (231b^3x^3 - 198b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 88bx \operatorname{arctanh}(\tanh(a + bx))) + 1155b^4}{1155b^4}$$

input `Integrate[x^3*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2)*(231*b^3*x^3 - 198*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 88*b*x*ArcTanh[Tanh[a + b*x]]^2 - 16*ArcTanh[Tanh[a + b*x]]^3))/(1155*b^4)`

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.20, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{6 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b} \right)}{5b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{9/2} dx}{9b} \right)}{7b} \right)}{5b} \\
 & \quad \downarrow 2588
 \end{aligned}$$

$$\frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{9/2} d \operatorname{arctanh}(\tanh(a+bx))}{9b^2} \right)}{7b} \right)}{5b}$$

$\downarrow$  15

$$\frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{99b^2} \right)}{7b} \right)}{5b}$$

input `Int[x^3*ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output `(2*x^3*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (6*((2*x^2*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(9/2))/(9*b) - (4*ArcTanh[Tanh[a + b*x]]^(11/2))/(99*b^2)))/(7*b)))/(5*b)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.55

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{11}{2}}}{11} + \frac{2(-3 \operatorname{arctanh}(\tanh(bx+a))+3bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}}}{9} + \frac{2((bx-\operatorname{arctanh}(\tanh(bx+a)))(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^{\frac{7}{2}}}{b^4}$

input `int(x^3*arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{2}{b^4} \left( \frac{1}{11} \operatorname{arctanh}(\tanh(bx+a))^{\frac{11}{2}} + \frac{1}{9} (-3 \operatorname{arctanh}(\tanh(bx+a)) + 3bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}} + \frac{1}{7} ((bx - \operatorname{arctanh}(\tanh(bx+a))) (2bx - 2 \operatorname{arctanh}(\tanh(bx+a))))^{\frac{7}{2}} + \frac{1}{5} (bx - \operatorname{arctanh}(\tanh(bx+a)))^3 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}} \right)$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.80

$$\int x^3 \operatorname{arctanh}(\tanh(ax + bx))^{3/2} dx = \frac{2(105b^5x^5 + 140ab^4x^4 + 5a^2b^3x^3 - 6a^3b^2x^2 + 8a^4bx - 16a^5)\sqrt{bx+a}}{1155b^4}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output 
$$\frac{2}{1155} (105b^5x^5 + 140a^2b^3x^3 - 6a^3b^2x^2 + 8a^4bx - 16a^5) \operatorname{sqrt}(bx+a) / b^4$$

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int x^3 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx)) dx$$

input `integrate(x**3*atanh(tanh(b*x+a))**(3/2), x)`

output `Integral(x**3*atanh(tanh(a + b*x))**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.66

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(105b^4x^4 + 35ab^3x^3 - 30a^2b^2x^2 + 24a^3bx - 16a^4)(bx + a)^{\frac{3}{2}}}{1155b^4}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(3/2), x, algorithm="maxima")`

output `2/1155*(105*b^4*x^4 + 35*a*b^3*x^3 - 30*a^2*b^2*x^2 + 24*a^3*b*x - 16*a^4) * (b*x + a)^(3/2)/b^4`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 205 vs.  $2(64) = 128$ .

Time = 0.12 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.56

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{2} \left( \frac{99\sqrt{2} \left( 5(bx+a)^{\frac{7}{2}} - 21(bx+a)^{\frac{5}{2}}a + 35(bx+a)^{\frac{3}{2}}a^2 - 35\sqrt{bx+aa^3} \right) a^2}{b^3} + \frac{22\sqrt{2} \left( 35(bx+a)^{\frac{9}{2}} - 180(bx+a)^{\frac{7}{2}}a + 378(bx+a)^{\frac{5}{2}}a^2 - 126(bx+a)^{\frac{3}{2}}a^3 + 35a^4 \right)}{b^3} \right)}{1155b^4}$$



input `integrate(x^3*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/3465*sqrt(2)*(99*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a^2/b^3 + 22*sqrt(2)*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)*a/b^3 + 5*sqrt(2)*(63*(b*x + a)^(11/2) - 385*(b*x + a)^(9/2)*a + 990*(b*x + a)^(7/2)*a^2 - 1386*(b*x + a)^(5/2)*a^3 + 1155*(b*x + a)^(3/2)*a^4 - 693*sqrt(b*x + a)*a^5)/b^3)/b`

### Mupad [B] (verification not implemented)

Time = 3.24 (sec) , antiderivative size = 1483, normalized size of antiderivative = 18.54

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \text{Too large to display}$$

input `int(x^3*atanh(tanh(a + b*x))^(3/2),x)`

output

```
(2*b*x^5*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/11 + (x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))*((20*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/11 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)))/(9*b) + (x^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^2/2 + (8*((20*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/11 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/(9*b)))/(7*b) + (16*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^2/2 + (8*((20*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/11 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1))...
```

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a)) x^3 dx$$

input

```
int(x^3*atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*x**3,x)
```

### 3.122 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result . . . . .	918
Mathematica [A] (verified) . . . . .	918
Rubi [A] (verified) . . . . .	919
Maple [A] (verified) . . . . .	920
Fricas [A] (verification not implemented) . . . . .	921
Sympy [F] . . . . .	921
Maxima [A] (verification not implemented) . . . . .	921
Giac [B] (verification not implemented) . . . . .	922
Mupad [B] (verification not implemented) . . . . .	922
Reduce [F] . . . . .	923

#### Optimal result

Integrand size = 15, antiderivative size = 59

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{8x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^2} + \frac{16 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{315b^3}$$

output

```
2/5*x^2*arctanh(tanh(b*x+a))^(5/2)/b-8/35*x*arctanh(tanh(b*x+a))^(7/2)/b^2
+16/315*arctanh(tanh(b*x+a))^(9/2)/b^3
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} (63b^2x^2 - 36bx \operatorname{arctanh}(\tanh(a + bx)) + 8 \operatorname{arctanh}(\tanh(a + bx)))}{315b^3}$$

input

```
Integrate[x^2*ArcTanh[Tanh[a + b*x]]^(3/2),x]
```

output

$$\frac{(2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{5/2} (63 b^2 x^2 - 36 b x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]] + 8 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^2))}{315 b^3}$$

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{3/2} dx$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{4 \int x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2} dx}{5b}$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2} dx}{7b} \right)}{5b}$$

$$\downarrow 2588$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2} d \operatorname{arctanh}(\operatorname{tanh}(a + b x))}{7b^2} \right)}{5b}$$

$$\downarrow 15$$

$$\frac{2x^2 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{7/2}}{7b} - \frac{4 \operatorname{arctanh}(\operatorname{tanh}(a + b x))^{9/2}}{63b^2} \right)}{5b}$$

input

$$\operatorname{Int}[x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{3/2}, x]$$

output

$$\frac{(2 x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{5/2})}{(5 b)} - \frac{(4 ((2 x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{7/2}) / (7 b) - (4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{9/2}) / (63 b^2)))}{(5 b)}$$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_.)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}} + 2(2bx-2 \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}} + 2(bx-\operatorname{arctanh}(\tanh(bx+a)))^2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{b^3}$	69

input `int(x^2*arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2/b^3*(1/9*arctanh(tanh(b*x+a))^(9/2)+1/7*(2*b*x-2*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^(7/2)+1/5*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(5/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(35b^4x^4 + 50ab^3x^3 + 3a^2b^2x^2 - 4a^3bx + 8a^4)\sqrt{bx+a}}{315b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`output `2/315*(35*b^4*x^4 + 50*a*b^3*x^3 + 3*a^2*b^2*x^2 - 4*a^3*b*x + 8*a^4)*sqrt(b*x + a)/b^3`**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int x^2 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx)) dx$$

input `integrate(x**2*atanh(tanh(b*x+a))**(3/2),x)`output `Integral(x**2*atanh(tanh(a + b*x))**(3/2), x)`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.71

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(35b^3x^3 + 15ab^2x^2 - 12a^2bx + 8a^3)(bx+a)^{\frac{3}{2}}}{315b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`output `2/315*(35*b^3*x^3 + 15*a*b^2*x^2 - 12*a^2*b*x + 8*a^3)*(b*x + a)^(3/2)/b^3`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 168 vs.  $2(47) = 94$ .

Time = 0.12 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.85

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{2} \left( \frac{21\sqrt{2}(3(bx+a)^{5/2} - 10(bx+a)^{3/2}a + 15\sqrt{bx+aa^2})a^2}{b^2} + \frac{18\sqrt{2}(5(bx+a)^{7/2} - 21(bx+a)^{5/2}a + 35(bx+a)^{3/2}a^2 - 35\sqrt{bx+aa^3})}{b^2} \right)}{315b}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/315*sqrt(2)*(21*sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2)*a^2/b^2 + 18*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a/b^2 + sqrt(2)*(3*5*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)/b^2)/b`

**Mupad [B] (verification not implemented)**

Time = 3.18 (sec) , antiderivative size = 1153, normalized size of antiderivative = 19.54

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \text{Too large to display}$$

input `int(x^2*atanh(tanh(a + b*x))^(3/2),x)`

output

```
(2*b*x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2
/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/9 + (x^3*(log((2*exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)
*((16*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/9 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) +
1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/((
7*b) + (x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - lo
g(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) + 1))
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/2 + (
6*((16*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/9 - 2*b*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*
(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
*exp(2*b*x) + 1))/2 + b*x))/(7*b)))/(5*b) + (8*(log((2*exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)
*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
)*exp(2*b*x) + 1)) + 2*b*x)^2/2 + (6*((16*b*(log(2/(exp(2*a)*exp(2*b*x) +
1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/9
- 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(ex
p(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - ...
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a)) x^2 dx$$

input

```
int(x^2*atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*x**2,x)
```



### 3.123 $\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	924
Mathematica [A] (verified)	924
Rubi [A] (verified)	925
Maple [A] (verified)	926
Fricas [A] (verification not implemented)	926
Sympy [A] (verification not implemented)	927
Maxima [A] (verification not implemented)	927
Giac [B] (verification not implemented)	927
Mupad [B] (verification not implemented)	928
Reduce [B] (verification not implemented)	929

#### Optimal result

Integrand size = 13, antiderivative size = 38

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^2}$$

output  $2/5*x*\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}/b-4/35*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b^2$

#### Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(7bx - 2 \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{35b^2}$$

input  $\operatorname{Integrate}[x*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(3/2)}, x]$

output  $(2*(7*b*x - 2*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(5/2)})/(35*b^2)$

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$$

$$\downarrow 2599$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b}$$

$$\downarrow 2588$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{5/2} d \operatorname{arctanh}(\tanh(a + bx))}{5b^2}$$

$$\downarrow 15$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{35b^2}$$

input `Int[x*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(2*x*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (4*ArcTanh[Tanh[a + b*x]]^(7/2))/(35*b^2)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```

Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))

```

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{7} + \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{b^2}$	42

input

```
int(x*arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^2*(1/7*arctanh(tanh(b*x+a))^(7/2)+1/5*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(5/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(5b^3x^3 + 8ab^2x^2 + a^2bx - 2a^3)\sqrt{bx + a}}{35b^2}$$

input

```
integrate(x*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
2/35*(5*b^3*x^3 + 8*a*b^2*x^2 + a^2*b*x - 2*a^3)*sqrt(b*x + a)/b^2
```

**Sympy [A] (verification not implemented)**

Time = 1.65 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \begin{cases} \frac{2x \operatorname{atanh}^{\frac{5}{2}}(\tanh(a+bx))}{5b} - \frac{4 \operatorname{atanh}^{\frac{7}{2}}(\tanh(a+bx))}{35b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**(3/2), x)`

output `Piecewise(((2*x*atanh(tanh(a + b*x))**(5/2)/(5*b) - 4*atanh(tanh(a + b*x))**  
*(7/2)/(35*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**(3/2)/2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \frac{2(5b^2x^2 + 3abx - 2a^2)(bx+a)^{\frac{3}{2}}}{35b^2}$$

input `integrate(x*arctanh(tanh(b*x+a))^(3/2), x, algorithm="maxima")`

output `2/35*(5*b^2*x^2 + 3*a*b*x - 2*a^2)*(b*x + a)^(3/2)/b^2`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 131 vs.  $2(30) = 60$ .

Time = 0.11 (sec) , antiderivative size = 131, normalized size of antiderivative = 3.45

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \frac{\sqrt{2} \left( \frac{35\sqrt{2}((bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa})a^2}{b} + \frac{14\sqrt{2}(3(bx+a)^{\frac{5}{2}} - 10(bx+a)^{\frac{3}{2}}a + 15\sqrt{bx+aa^2})a}{b} + \frac{3\sqrt{2}(5(bx+a)^{\frac{7}{2}} - 21(bx+a) \right)}{105b}$$

input `integrate(x*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output 
$$\frac{1}{105}\sqrt{2}*(35\sqrt{2}*((b*x + a)^{3/2} - 3\sqrt{b*x + a})*a^2/b + 14*\sqrt{2}*(3*(b*x + a)^{5/2} - 10*(b*x + a)^{3/2})*a + 15*\sqrt{b*x + a})*a^2*a/b + 3*\sqrt{2}*(5*(b*x + a)^{7/2} - 21*(b*x + a)^{5/2})*a + 35*(b*x + a)^{3/2})*a^2 - 35*\sqrt{b*x + a})*a^3)/b)/b$$

### Mupad [B] (verification not implemented)

Time = 3.20 (sec) , antiderivative size = 823, normalized size of antiderivative = 21.66

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \text{Too large to display}$$

input `int(x*atanh(tanh(a + b*x))^(3/2),x)`

output 
$$\begin{aligned} & (2*b*x^3*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{1/2})/7 + (x^2*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2) \\ & *((12*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x))/7 - 2*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))/((5*b) \\ & + (x*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{1/2}*((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{2/2} + (4*((12*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x))/7 - 2*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x))/(5*b)))/(3*b) + (2*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{1/2}*((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{2/2} + (4*((12*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x))/7 - 2*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - 1... \end{aligned}$$

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 (-2\operatorname{atanh}(\tanh(bx + a)) + 7bx)}{35b^2}$$

input `int(x*atanh(tanh(b*x+a))^(3/2),x)`

output `(2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2*(- 2*atanh(tanh(a + b*x)) + 7*b*x))/(35*b**2)`

### 3.124 $\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	930
Mathematica [A] (verified)	930
Rubi [A] (verified)	931
Maple [A] (verified)	932
Fricas [A] (verification not implemented)	932
Sympy [A] (verification not implemented)	932
Maxima [A] (verification not implemented)	933
Giac [B] (verification not implemented)	933
Mupad [B] (verification not implemented)	934
Reduce [B] (verification not implemented)	934

#### Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b}$$

output

```
2/5*arctanh(tanh(b*x+a))^(5/2)/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b)
```

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5b}$	15
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5b}$	15

input `int(arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2/5*arctanh(tanh(b*x+a))^(5/2)/b`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.56

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(b^2x^2 + 2abx + a^2)\sqrt{bx + a}}{5b}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `2/5*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(b*x + a)/b`

**Sympy [A] (verification not implemented)**

Time = 0.81 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.44

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \begin{cases} \frac{2 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a+bx))}{5b} & \text{for } b \neq 0 \\ x \operatorname{atanh}^{\frac{3}{2}}(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**(3/2),x)`

output `Piecewise((2*atanh(tanh(a + b*x))**(5/2)/(5*b), Ne(b, 0)), (x*atanh(tanh(a))**(3/2), True))`

### Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2(bx + a)^{5/2}}{5b}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `2/5*(b*x + a)^(5/2)/b`

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs.  $2(14) = 28$ .

Time = 0.11 (sec) , antiderivative size = 84, normalized size of antiderivative = 4.67

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{2} \left( 15 \sqrt{2} \sqrt{bx + a} a^2 + 10 \sqrt{2} \left( (bx + a)^{3/2} - 3 \sqrt{bx + a} a \right) a + \sqrt{2} \left( 3 (bx + a)^{5/2} - 10 (bx + a)^{3/2} a + 15 \sqrt{bx + a} a^2 \right) \right)}{15b}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/15*sqrt(2)*(15*sqrt(2)*sqrt(b*x + a)*a^2 + 10*sqrt(2)*((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)*a + sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2))/b`

**Mupad [B] (verification not implemented)**

Time = 3.24 (sec) , antiderivative size = 97, normalized size of antiderivative = 5.39

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\left(\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)\right)^2 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{10b}$$

input `int(atanh(tanh(a + b*x))^(3/2),x)`output `((log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))^2*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(10*b)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{5b}$$

input `int(atanh(tanh(b*x+a))^(3/2),x)`output `(2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2)/(5*b)`

### 3.125 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x} dx$

Optimal result	935
Mathematica [A] (verified)	936
Rubi [A] (verified)	936
Maple [A] (verified)	938
Fricas [A] (verification not implemented)	938
Sympy [F]	939
Maxima [F]	939
Giac [A] (verification not implemented)	939
Mupad [B] (verification not implemented)	940
Reduce [F]	940

#### Optimal result

Integrand size = 15, antiderivative size = 91

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = 2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2} - 2(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{3}\operatorname{arctanh}(\tanh(a + bx))^{3/2}$$

output

```
2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x
-arctanh(tanh(b*x+a)))^(3/2)-2*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x
+a))^(1/2)+2/3*arctanh(tanh(b*x+a))^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx =$$

$$-\frac{2}{3} \left( 3bx \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - 4 \operatorname{arctanh}(\tanh(a + bx))^{3/2} \right.$$

$$\left. + 3 \operatorname{arctanh} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}} \right) (-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2} \right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x,x]
```

output

```
(-2*(3*b*x*Sqrt[ArcTanh[Tanh[a + b*x]]] - 4*ArcTanh[Tanh[a + b*x]]^(3/2) +
3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]
]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^(3/2)))/3
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2590, 2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx$$

$$\downarrow 2590$$

$$\frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx$$

$$\downarrow 2590$$

$$\frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \right)$$

↓ 2592

$$\frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))} \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} \right) \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x,x]`

output `-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/3`

### Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.44

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3} + 2\sqrt{\operatorname{arctanh}(\tanh(bx+a))} a + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}$

input `int(arctanh(tanh(b*x+a))^(3/2)/x,x,method=_RETURNVERBOSE)`

output `2/3*arctanh(tanh(b*x+a))^(3/2)+2*arctanh(tanh(b*x+a))^(1/2)*a+2*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2)-2*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x} dx = \left[ a^{\frac{3}{2}} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) + \frac{2}{3}(bx+4a)\sqrt{bx+a}, 2\sqrt{-a}a \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + \frac{2}{3}(bx+4a)\sqrt{bx+a} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x,x, algorithm="fricas")`

output `[a^(3/2)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2/3*(b*x + 4*a)*sqrt(b*x + a), 2*sqrt(-a)*a*arctan(sqrt(-a)/sqrt(b*x + a)) + 2/3*(b*x + 4*a)*sqrt(b*x + a)]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x,x)`

output `Integral(atanh(tanh(a + b*x))**(3/2)/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}}{x} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/x, x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.63

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = \frac{1}{3} \sqrt{2} \left( \frac{3 \sqrt{2} a^2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + \sqrt{2}(bx + a)^{\frac{3}{2}} + 3 \sqrt{2} \sqrt{bx + a} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x,x, algorithm="giac")`

output `1/3*sqrt(2)*(3*sqrt(2)*a^2*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + sqrt(2)*(b*x + a)^(3/2) + 3*sqrt(2)*sqrt(b*x + a)*a)`



**Mupad [B] (verification not implemented)**

Time = 7.58 (sec) , antiderivative size = 501, normalized size of antiderivative = 5.51

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x,x)`

output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((4*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + b*x))/3 - 2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b + (2^(1/2)*log((((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*4i)/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2)*1i)/4 + (2*b*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2))/3
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x} dx$$

input `int(atanh(tanh(b*x+a))^(3/2)/x,x)`

output

```
int((sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)))/x,x)
```

### 3.126 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^2} dx$

Optimal result . . . . .	941
Mathematica [A] (verified) . . . . .	942
Rubi [A] (verified) . . . . .	942
Maple [A] (verified) . . . . .	944
Fricas [A] (verification not implemented) . . . . .	944
Sympy [F] . . . . .	945
Maxima [F] . . . . .	945
Giac [A] (verification not implemented) . . . . .	945
Mupad [B] (verification not implemented) . . . . .	946
Reduce [F] . . . . .	947

#### Optimal result

Integrand size = 15, antiderivative size = 81

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx =$$

$$-3b \arctan \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}$$

$$+ 3b \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x}$$

output

```
-3*b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(
b*x-arctanh(tanh(b*x+a)))^(1/2)+3*b*arctanh(tanh(b*x+a))^(1/2)-arctanh(tan
h(b*x+a))^(3/2)/x
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = 3b\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} - 3b\operatorname{arctanh}\left(\frac{x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^2,x]`

output `3*b*Sqrt[ArcTanh[Tanh[a + b*x]]] - ArcTanh[Tanh[a + b*x]]^(3/2)/x - 3*b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]`

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.05, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx$$

$$\downarrow \text{2599}$$

$$\frac{3}{2}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x}$$

$$\downarrow \text{2590}$$

$$\frac{3}{2}b \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - (bx - \operatorname{arctanh}(\tanh(a+bx)))}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \right) -$$

$x$   
 $\downarrow$  2592

$$\frac{3}{2}b \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \arctan \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \right) -$$

$x$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^2,x]`

output `(3*b*(-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]])/2 - ArcTanh[Tanh[a + b*x]]^(3/2)/x`

### Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.05

method	result
default	$2b \left( \sqrt{\operatorname{arctanh}(\tanh(bx+a))} + \frac{\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))}{2} + \frac{bx}{2}\right) \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{bx} - \frac{3\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2bx} \right)$

input

```
int(arctanh(tanh(b*x+a))^(3/2)/x^2,x,method=_RETURNVERBOSE)
```

output

```
2*b*(arctanh(tanh(b*x+a))^(1/2))+(-1/2*arctanh(tanh(b*x+a))+1/2*b*x)*arctanh(tanh(b*x+a))^(1/2)/b/x-3/2*(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.22

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = \left[ \frac{3\sqrt{abx} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(2bx - a)\sqrt{bx+a}}{2x}, \frac{3\sqrt{-abx} \operatorname{arctanh}\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right)}{2x} \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^2,x, algorithm="fricas")
```

output

```
[1/2*(3*sqrt(a)*b*x*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(2*b*x - a)*sqrt(b*x + a))/x, (3*sqrt(-a)*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) + (2*b*x - a)*sqrt(b*x + a))/x]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{x^2} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(3/2)/x**2,x)
```

output

```
Integral(atanh(tanh(a + b*x))**(3/2)/x**2, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}}{x^2} dx$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^2,x, algorithm="maxima")
```

output

```
integrate(arctanh(tanh(b*x + a))^(3/2)/x^2, x)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = \frac{1}{2} \sqrt{2} \left( \frac{3 \sqrt{2} a \operatorname{arctan}\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + 2 \sqrt{2} \sqrt{bx+a} - \frac{\sqrt{2} \sqrt{bx+aa}}{bx} \right) b$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^2,x, algorithm="giac")
```

output

```
1/2*sqrt(2)*(3*sqrt(2)*a*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 2*sqrt(2)*sqrt(b*x + a) - sqrt(2)*sqrt(b*x + a)*a/(b*x))*b
```

**Mupad [B] (verification not implemented)**

Time = 4.27 (sec) , antiderivative size = 459, normalized size of antiderivative = 5.67

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = 3b \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}$$

$$+ \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{2x}$$

$$- \frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{2x}$$

$$+ b \ln \left( \frac{4\sqrt{2} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - 4\sqrt{2} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 8 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}} \sqrt{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}}{x \sqrt{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)} - 2bx} \right)$$

input

```
int(atanh(tanh(a + b*x))^(3/2)/x^2,x)
```

output

```
3*b*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2) + (log(1/(exp(2*a)*exp(2*b*x) + 1))*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(2*x) - (log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(2*x) + b*log(-(4*2^(1/2)*log(1/(exp(2*a)*exp(2*b*x) + 1)) - 4*2^(1/2)*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 8*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(1/(exp(2*a)*exp(2*b*x) + 1)) - 2*b*x)^(1/2) + 4*2^(1/2)*b*x)/(x*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(1/(exp(2*a)*exp(2*b*x) + 1)) - 2*b*x)^(1/2))*((9*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/8 - (9*log(1/(exp(2*a)*exp(2*b*x) + 1)))/8 - (9*b*x)/4)^(1/2)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^2} dx$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^2,x)`

output `int((sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)))/x**2,x)`



### 3.127 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^3} dx$

Optimal result	948
Mathematica [A] (verified)	949
Rubi [A] (verified)	949
Maple [A] (verified)	951
Fricas [A] (verification not implemented)	951
Sympy [F]	952
Maxima [F]	952
Giac [A] (verification not implemented)	952
Mupad [B] (verification not implemented)	953
Reduce [F]	953

#### Optimal result

Integrand size = 15, antiderivative size = 92

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^3} dx = \frac{3b^2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} - \frac{3b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{4x} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2}$$

output

```
3/4*b^2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^(1/2)-3/4*b*arctanh(tanh(b*x+a))^(1/2)/x-1/2*
arctanh(tanh(b*x+a))^(3/2)/x^2
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \frac{1}{4} \left( -\frac{3b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} - \frac{3b^2\operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}} \right)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^3,x]`

output  $((-3*b*\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]])/x - (2*\text{ArcTanh}[\text{Tanh}[a + b*x]]^{(3/2)})/x^2 - (3*b^2*\text{ArcTanh}[\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]]/\text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]])/ \text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]])/4$

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx$$

↓ 2599

$$\frac{3}{4}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{2x^2}$$

↓ 2599

$$\frac{3}{4}b \left( \frac{1}{2}b \int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2}$$

↓ 2592

$$\frac{3}{4}b \left( \frac{b \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^3,x]`

output `(3*b*((b*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] - Sqrt[ArcTanh[Tanh[a + b*x]]]/x))/4 - ArcTanh[Tanh[a + b*x]]^(3/2)/(2*x^2)`

### Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.99

method	result
default	$2b^2 \left( \frac{-\frac{5}{8} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{b^2 x^2} + \left( \frac{3}{8} \operatorname{arctanh}(\tanh(bx+a)) - \frac{3bx}{8} \right) \sqrt{\operatorname{arctanh}(\tanh(bx+a))} \right) - \frac{3 \operatorname{arctanh} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a)) - bx}} \right)}{8 \sqrt{\operatorname{arctanh}(\tanh(bx+a)) - bx}}$

input `int(arctanh(tanh(b*x+a))^(3/2)/x^3,x,method=_RETURNVERBOSE)`

output `2*b^2*((-5/8*arctanh(tanh(b*x+a))^(3/2)+(3/8*arctanh(tanh(b*x+a))-3/8*b*x)*arctanh(tanh(b*x+a))^(1/2))/b^2/x^2-3/8/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.32

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \left[ \frac{3 \sqrt{ab^2 x^2} \log \left( \frac{bx - 2 \sqrt{bx+a} \sqrt{a} + 2a}{x} \right) - 2(5abx + 2a^2) \sqrt{bx+a}}{8ax^2}, \frac{3 \sqrt{-ab^2 x^2}}{8ax^2} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^3,x, algorithm="fricas")`

output `[1/8*(3*sqrt(a)*b^2*x^2*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(5*a*b*x + 2*a^2)*sqrt(b*x + a))/(a*x^2), 1/4*(3*sqrt(-a)*b^2*x^2*arctan(sqrt(-a)/sqrt(b*x + a)) - (5*a*b*x + 2*a^2)*sqrt(b*x + a))/(a*x^2)]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{x^3} dx$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**3,x)`

output `Integral(atanh(tanh(a + b*x))**(3/2)/x**3, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}}{x^3} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^3,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/x^3, x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.79

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \frac{\sqrt{2} \left( \frac{3\sqrt{2}b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} - \frac{\sqrt{2}(5(bx+a)^{\frac{3}{2}}b^3 - 3\sqrt{bx+ab^3})}{b^2x^2} \right)}{8b}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^3,x, algorithm="giac")`

output `1/8*sqrt(2)*(3*sqrt(2)*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) - sqrt(2)*(5*(b*x + a)^(3/2)*b^3 - 3*sqrt(b*x + a)*a*b^3)/(b^2*x^2))/b`

**Mupad [B] (verification not implemented)**

Time = 8.04 (sec) , antiderivative size = 609, normalized size of antiderivative = 6.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^3,x)`

output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*x^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x)) + (2^(1/2)*b^2*log(((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x*16i)/x)*3i)/(8*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)) - (b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((5*log(2/(exp(2*a)*exp(2*b*x) + 1)))/4 - (5*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/4 + (5*b*x)/2))/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^3} dx = \frac{-4\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 6\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{8x^2}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^3,x)`

output

```
( - 4*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 6*sqrt(atanh(tanh(a + b*x)))*b*x + 3*int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x),x)*b**2*x**2)/(8*x**2)
```

**3.128**  $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^4} dx$

Optimal result . . . . .	955
Mathematica [A] (verified) . . . . .	956
Rubi [A] (verified) . . . . .	956
Maple [A] (verified) . . . . .	958
Fricas [A] (verification not implemented) . . . . .	959
Sympy [F] . . . . .	959
Maxima [F] . . . . .	960
Giac [A] (verification not implemented) . . . . .	960
Mupad [B] (verification not implemented) . . . . .	960
Reduce [F] . . . . .	961

**Optimal result**

Integrand size = 15, antiderivative size = 146

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \frac{b^3 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8(bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} - \frac{8x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b^3} + \frac{8(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{4x^2} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^3}$$

output

```
1/8*b^3*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^(3/2)-1/8*b^2/x/arctanh(tanh(b*x+a))^(1/2)+1/
8*b^3/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)-1/4*b*arctanh(
tanh(b*x+a))^(1/2)/x^2-1/3*arctanh(tanh(b*x+a))^(3/2)/x^3
```



**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \frac{b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{8(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} + \sqrt{\operatorname{arctanh}(\tanh(a + bx))} \left( -\frac{7b}{12x^2} - \frac{b^2}{8x(-bx + \operatorname{arctanh}(\tanh(a + bx)))} - \frac{-bx + \operatorname{arctanh}(\tanh(a + bx))}{3x^3} \right)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^4,x]`

output  $(b^3 \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]]/\operatorname{Sqrt}[-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]]])/(8*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])^{(3/2)}) + \operatorname{Sqrt}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]]*((-7*b)/(12*x^2) - b^2/(8*x*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])) - (-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])/(3*x^3))$

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx \\ & \quad \downarrow \text{2599} \\ & \frac{1}{2}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^3} \\ & \quad \downarrow \text{2599} \\ & \frac{1}{2}b \left( \frac{1}{4}b \int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^3} \end{aligned}$$

↓ 2599

$$\frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3}$$

↓ 2594

$$\frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \left( \frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3} \right) - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3}$$

↓ 2592

$$\frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \left( \frac{2 \arctan \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3} \right) - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^4, x]
```

output

```
(b*((b*(-1/2*(b*((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])) - 1/(x*Sqrt[ArcTanh[Tanh[a + b*x]]])))/4 - Sqrt[ArcTanh[Tanh[a + b*x]]/(2*x^2))/2 - ArcTanh[Tanh[a + b*x]]^(3/2)/(3*x^3)
```

## Definitions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/(n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.79

method	result
default	$2b^3 \left( \frac{-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{16(\operatorname{arctanh}(\tanh(bx+a))-bx)} - \frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{6} + \left( \frac{\operatorname{arctanh}(\tanh(bx+a))}{16} - \frac{bx}{16} \right) \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b^3 x^3} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}\right)}{16(\operatorname{arctanh}(\tanh(bx+a))-bx)} \right)$

input `int(arctanh(tanh(b*x+a))^(3/2)/x^4,x,method=_RETURNVERBOSE)`

output `2*b^3*((-1/16/(arctanh(tanh(b*x+a))-b*x)*arctanh(tanh(b*x+a))^(5/2)-1/6*arctanh(tanh(b*x+a))^(3/2)+(1/16*arctanh(tanh(b*x+a))-1/16*b*x)*arctanh(tanh(b*x+a))^(1/2))/b^3/x^3+1/16/(arctanh(tanh(b*x+a))-b*x)^(3/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \left[ \frac{3\sqrt{ab^3}x^3 \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(3ab^2x^2 + 14a^2bx + 8a^3)\sqrt{bx+a}}{48a^2x^3} - \frac{3\sqrt{-ab^3}x^3 \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (3ab^2x^2 + 14a^2bx + 8a^3)\sqrt{bx+a}}{24a^2x^3} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^4,x, algorithm="fricas")`

output `[1/48*(3*sqrt(a)*b^3*x^3*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(3*a*b^2*x^2 + 14*a^2*b*x + 8*a^3)*sqrt(b*x + a))/(a^2*x^3), -1/24*(3*sqrt(-a)*b^3*x^3*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b^2*x^2 + 14*a^2*b*x + 8*a^3)*sqrt(b*x + a))/(a^2*x^3)]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{x^4} dx$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**4, x)`

output `Integral(atanh(tanh(a + b*x))**(3/2)/x**4, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{3/2}}{x^4} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^4,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/x^4, x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx =$$

$$-\frac{1}{48} \sqrt{2} b^3 \left( \frac{3 \sqrt{2} \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{\sqrt{2} \left( 3 (bx + a)^{5/2} + 8 (bx + a)^{3/2} a - 3 \sqrt{bx + aa^2} \right)}{ab^3 x^3} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^4,x, algorithm="giac")`

output `-1/48*sqrt(2)*b^3*(3*sqrt(2)*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a) + sqrt(2)*(3*(b*x + a)^(5/2) + 8*(b*x + a)^(3/2)*a - 3*sqrt(b*x + a)*a^2)/(a*b^3*x^3)`

**Mupad [B] (verification not implemented)**

Time = 7.47 (sec) , antiderivative size = 1019, normalized size of antiderivative = 6.98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^4,x)`

output

```
(11*b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/
(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(12*x*(log(2/(exp(2*a)*exp(2*b*x) + 1
)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + ((
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)
*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(
2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*x^3*(3*log(2/(e
xp(2*a)*exp(2*b*x) + 1)) - 3*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b
*x) + 1)) + 6*b*x)) - (2*b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*
b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(x*(3*log(2/(exp
(2*a)*exp(2*b*x) + 1)) - 3*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x
) + 1)) + 6*b*x)) + (2^(1/2)*b^3*log(((2*2^(1/2)*a + log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)
^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(ex
p(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(2*a - log((2*exp(2*a)
*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1))
+ 2*b*x) + 2^(1/2)*b*x*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp
(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*
(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b...
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx = \frac{-8\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 6\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{24x^3}$$

input

```
int(atanh(tanh(b*x+a))^(3/2)/x^4,x)
```

output

```
( - 8*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 6*sqrt(atanh(tanh(
a + b*x)))*b*x + 3*int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x*
*2),x)*b**2*x**3)/(24*x**3)
```

### 3.129 $\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	962
Mathematica [A] (verified)	962
Rubi [A] (verified)	963
Maple [A] (verified)	965
Fricas [A] (verification not implemented)	966
Sympy [F(-1)]	966
Maxima [A] (verification not implemented)	966
Giac [B] (verification not implemented)	967
Mupad [B] (verification not implemented)	967
Reduce [F]	968

#### Optimal result

Integrand size = 15, antiderivative size = 101

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{16x^3 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{63b^2} + \frac{32x^2 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{231b^3} - \frac{128x \operatorname{arctanh}(\tanh(a + bx))^{13/2}}{3003b^4} + \frac{256 \operatorname{arctanh}(\tanh(a + bx))^{15/2}}{45045b^5}$$

output  $2/7*x^4*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b-16/63*x^3*\operatorname{arctanh}(\tanh(b*x+a))^{(9/2)}/b^2+32/231*x^2*\operatorname{arctanh}(\tanh(b*x+a))^{(11/2)}/b^3-128/3003*x*\operatorname{arctanh}(\tanh(b*x+a))^{(13/2)}/b^4+256/45045*\operatorname{arctanh}(\tanh(b*x+a))^{(15/2)}/b^5$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{7/2} (6435b^4x^4 - 5720b^3x^3 \operatorname{arctanh}(\tanh(a + bx)) + 3120b^2x^2 \operatorname{arctanh}(\tanh(a + bx))) + 45045b^5}{45045b^5}$$

input `Integrate[x^4*ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(7/2)*(6435*b^4*x^4 - 5720*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 3120*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 960*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4)/(45045*b^5)`

### Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{8 \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{2 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2} dx}{3b} \right)}{7b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{11b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{11/2} dx}{11b} \right)}{3b} \right)}{7b} \\
 & \quad \downarrow 2599
 \end{aligned}$$



$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{13/2}}{13b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{13/2} dx}{11b} \right)}{11b} \right)}{3b} \right)$$

2588

$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{13/2}}{13b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{13/2} dx}{11b} \right)}{11b} \right)}{3b} \right)$$

15

$$8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{2x^4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{13/2}}{13b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{15/2}}{195b^2} \right)}{11b} \right)}{3b} \right)$$

input `Int [x^4*ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(2*x^4*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (8*((2*x^3*ArcTanh[Tanh[a + b*x]]^(9/2))/(9*b) - (2*((2*x^2*ArcTanh[Tanh[a + b*x]]^(11/2))/(11*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(13/2))/(13*b) - (4*ArcTanh[Tanh[a + b*x]]^(15/2))/(195*b^2)))/(11*b)))/(3*b)))/(7*b)`

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.52

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{15}}{15} + \frac{2(-4 \operatorname{arctanh}(\tanh(bx+a))+4bx) \operatorname{arctanh}(\tanh(bx+a))^{13}}{13} + \frac{2(2(bx-\operatorname{arctanh}(\tanh(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2+(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2)}{11}$

input `int(x^4*arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `2/b^5*(1/15*arctanh(tanh(b*x+a))^(15/2)+1/13*(-4*arctanh(tanh(b*x+a))+4*b*x)*arctanh(tanh(b*x+a))^(13/2)+1/11*(2*(b*x-arctanh(tanh(b*x+a)))^2+(2*b*x-2*arctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(11/2)+2/9*(b*x-arctanh(tanh(b*x+a)))^2*(2*b*x-2*arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(9/2)+1/7*(b*x-arctanh(tanh(b*x+a)))^4*arctanh(tanh(b*x+a))^(7/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.85

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(3003 b^7 x^7 + 7161 a b^6 x^6 + 4473 a^2 b^5 x^5 + 35 a^3 b^4 x^4 - 40 a^4 b^3 x^3 + 48 a^5 b^2 x^2 - 64 a^6 b x + 128 a^7)}{45045 b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `2/45045*(3003*b^7*x^7 + 7161*a*b^6*x^6 + 4473*a^2*b^5*x^5 + 35*a^3*b^4*x^4 - 40*a^4*b^3*x^3 + 48*a^5*b^2*x^2 - 64*a^6*b*x + 128*a^7)*sqrt(b*x + a)/b^5`

**Sympy [F(-1)]**

Timed out.

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Timed out}$$

input `integrate(x**4*atanh(tanh(b*x+a))**(5/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.63

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(3003 b^5 x^5 + 1155 a b^4 x^4 - 840 a^2 b^3 x^3 + 560 a^3 b^2 x^2 - 320 a^4 b x + 128 a^5)(bx + a)^{5/2}}{45045 b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output  $2/45045*(3003*b^5*x^5 + 1155*a*b^4*x^4 - 840*a^2*b^3*x^3 + 560*a^3*b^2*x^2 - 320*a^4*b*x + 128*a^5)*(b*x + a)^{(5/2)}/b^5$

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 344 vs.  $2(81) = 162$ .

Time = 0.11 (sec) , antiderivative size = 344, normalized size of antiderivative = 3.41

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\sqrt{2} \left( \frac{143 \sqrt{2} (35 (bx+a)^{9/2} - 180 (bx+a)^{7/2} a + 378 (bx+a)^{5/2} a^2 - 420 (bx+a)^{3/2} a^3 + 315 \sqrt{bx+aa^4}) a^3}{b^4} + \frac{195 \sqrt{2} (63 (bx+a)^{11/2} - 385 (bx+a)^{9/2} a + 990 (bx+a)^{7/2} a^2 - 1386 (bx+a)^{5/2} a^3 + 1155 (bx+a)^{3/2} a^4 - 693 \sqrt{bx+a} a^5) a^2}{b^4} + 45 \sqrt{2} (231 (bx+a)^{13/2} - 1638 (bx+a)^{11/2} a + 5005 (bx+a)^{9/2} a^2 - 8580 (bx+a)^{7/2} a^3 + 9009 (bx+a)^{5/2} a^4 - 6006 (bx+a)^{3/2} a^5 + 3003 \sqrt{bx+a} a^6) a}{b^4} + 7 \sqrt{2} (429 (bx+a)^{15/2} - 3465 (bx+a)^{13/2} a + 12285 (bx+a)^{11/2} a^2 - 25025 (bx+a)^{9/2} a^3 + 32175 (bx+a)^{7/2} a^4 - 27027 (bx+a)^{5/2} a^5 + 15015 (bx+a)^{3/2} a^6 - 6435 \sqrt{bx+a} a^7) / b^4}{b}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output  $1/45045*\sqrt{2}*(143*\sqrt{2}*(35*(b*x + a)^{(9/2)} - 180*(b*x + a)^{(7/2)}*a + 378*(b*x + a)^{(5/2)}*a^2 - 420*(b*x + a)^{(3/2)}*a^3 + 315*\sqrt{b*x + a}*a^4)*a^3/b^4 + 195*\sqrt{2}*(63*(b*x + a)^{(11/2)} - 385*(b*x + a)^{(9/2)}*a + 990*(b*x + a)^{(7/2)}*a^2 - 1386*(b*x + a)^{(5/2)}*a^3 + 1155*(b*x + a)^{(3/2)}*a^4 - 693*\sqrt{b*x + a}*a^5)*a^2/b^4 + 45*\sqrt{2}*(231*(b*x + a)^{(13/2)} - 1638*(b*x + a)^{(11/2)}*a + 5005*(b*x + a)^{(9/2)}*a^2 - 8580*(b*x + a)^{(7/2)}*a^3 + 9009*(b*x + a)^{(5/2)}*a^4 - 6006*(b*x + a)^{(3/2)}*a^5 + 3003*\sqrt{b*x + a}*a^6)*a/b^4 + 7*\sqrt{2}*(429*(b*x + a)^{(15/2)} - 3465*(b*x + a)^{(13/2)}*a + 12285*(b*x + a)^{(11/2)}*a^2 - 25025*(b*x + a)^{(9/2)}*a^3 + 32175*(b*x + a)^{(7/2)}*a^4 - 27027*(b*x + a)^{(5/2)}*a^5 + 15015*(b*x + a)^{(3/2)}*a^6 - 6435*\sqrt{b*x + a}*a^7)/b^4)/b$

### Mupad [B] (verification not implemented)

Time = 3.42 (sec) , antiderivative size = 2681, normalized size of antiderivative = 26.54

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Too large to display}$$

input `int(x^4*atanh(tanh(a + b*x))^(5/2),x)`

output

```
(2*b^2*x^7*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log
(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/15 + (x^5*(log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1
/2)*((3*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 - (12*(3*b^2*(log(2/(exp(2*a)*exp
(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2
*b*x) - (28*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/15*(log(2/(exp(2*a)*exp(2*b*
x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*
x))/(13*b)))/(11*b) - (x^6*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log(
(2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (28*b^2*(log
(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*ex
p(2*b*x) + 1))/2 + b*x))/15*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*
b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(13*b) - (x^4*(1
og((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*
exp(2*b*x) + 1))/2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(
2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3/4 - (10*((3*b*(log(
2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2
*b*x) + 1)) + 2*b*x)^2)/2 - (12*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) -
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (28*...
```

**Reduce [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a))^2 x^4 dx$$

input

```
int(x^4*atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2*x**4,x)
```

### 3.130 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	969
Mathematica [A] (verified)	969
Rubi [A] (verified)	970
Maple [A] (verified)	972
Fricas [A] (verification not implemented)	972
Sympy [A] (verification not implemented)	973
Maxima [A] (verification not implemented)	973
Giac [B] (verification not implemented)	974
Mupad [B] (verification not implemented)	974
Reduce [F]	975

#### Optimal result

Integrand size = 15, antiderivative size = 80

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{21b^2} + \frac{16x \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{231b^3} - \frac{32 \operatorname{arctanh}(\tanh(a + bx))^{13/2}}{3003b^4}$$

output

```
2/7*x^3*arctanh(tanh(b*x+a))^(7/2)/b-4/21*x^2*arctanh(tanh(b*x+a))^(9/2)/b
^2+16/231*x*arctanh(tanh(b*x+a))^(11/2)/b^3-32/3003*arctanh(tanh(b*x+a))^(
13/2)/b^4
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.82

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{7/2} (429b^3 x^3 - 286b^2 x^2 \operatorname{arctanh}(\tanh(a + bx)) + 104bx \operatorname{arctanh}(\tanh(a + bx))) + 104b^2 x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2} - 104b^2 x \operatorname{arctanh}(\tanh(a + bx))^{11/2} + 104b^2 \operatorname{arctanh}(\tanh(a + bx))^{13/2}}{3003b^4}$$

input `Integrate[x^3*ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(7/2)*(429*b^3*x^3 - 286*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 104*b*x*ArcTanh[Tanh[a + b*x]]^2 - 16*ArcTanh[Tanh[a + b*x]]^3))/(3003*b^4)`

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.20, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{6 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{9/2} dx}{9b} \right)}{7b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{11b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{11/2} dx}{11b} \right)}{9b} \right)}{7b} \\
 & \quad \downarrow 2588
 \end{aligned}$$

$$\frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{11/2} d \operatorname{arctanh}(\tanh(a+bx))}{11b^2} \right)}{9b} \right)}{7b}$$

$\downarrow$  15

$$\frac{2x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{9b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{11/2}}{11b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{13/2}}{143b^2} \right)}{9b} \right)}{7b}$$

input `Int[x^3*ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(2*x^3*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (6*((2*x^2*ArcTanh[Tanh[a + b*x]]^(9/2))/(9*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(11/2))/(11*b) - (4*ArcTanh[Tanh[a + b*x]]^(13/2))/(143*b^2)))/(9*b)))/(7*b)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`



**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.55

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{13}}{13} + \frac{2(-3 \operatorname{arctanh}(\tanh(bx+a))+3bx) \operatorname{arctanh}(\tanh(bx+a))^{11}}{11} + \frac{2((bx-\operatorname{arctanh}(\tanh(bx+a)))(2bx-2 \operatorname{arctanh}(\tanh(bx+a)))^2}{b^4}$

input `int(x^3*arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{2}{b^4} \left( \frac{1}{13} \operatorname{arctanh}(\tanh(bx+a))^{13/2} + \frac{1}{11} (-3 \operatorname{arctanh}(\tanh(bx+a)) + 3bx) \operatorname{arctanh}(\tanh(bx+a))^{11/2} + \frac{1}{9} ((bx - \operatorname{arctanh}(\tanh(bx+a)))^2 (2bx - 2 \operatorname{arctanh}(\tanh(bx+a))) + (bx - \operatorname{arctanh}(\tanh(bx+a)))^2) \operatorname{arctanh}(\tanh(bx+a))^{9/2} + \frac{1}{7} (bx - \operatorname{arctanh}(\tanh(bx+a)))^3 \operatorname{arctanh}(\tanh(bx+a))^{7/2} \right)$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.94

$$\int x^3 \operatorname{arctanh}(\tanh(ax+bx))^{5/2} dx = \frac{2(231b^6x^6 + 567ab^5x^5 + 371a^2b^4x^4 + 5a^3b^3x^3 - 6a^4b^2x^2 + 8a^5bx - 16a^6)\sqrt{bx+a}}{3003b^4}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output 
$$\frac{2}{3003} (231b^6x^6 + 567a^5b^5x^5 + 371a^2b^4x^4 + 5a^3b^3x^3 - 6a^4b^2x^2 + 8a^5bx - 16a^6) \operatorname{sqrt}(bx+a) / b^4$$

**Sympy [A] (verification not implemented)**

Time = 76.16 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.18

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \begin{cases} \frac{2x^3 \operatorname{atanh}^{\frac{7}{2}}(\tanh(a+bx))}{7b} - \frac{4x^2 \operatorname{atanh}^{\frac{9}{2}}(\tanh(a+bx))}{21b^2} + \frac{16x \operatorname{atanh}^{\frac{11}{2}}(\tanh(a+bx))}{231b^3} - \frac{32 \operatorname{atanh}^{\frac{13}{2}}(\tanh(a+bx))}{3003b^4} \\ \frac{x^4 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a))}{4} \end{cases}$$

input `integrate(x**3*atanh(tanh(b*x+a))**(5/2),x)`output `Piecewise((2*x**3*atanh(tanh(a + b*x))**(7/2)/(7*b) - 4*x**2*atanh(tanh(a + b*x))**(9/2)/(21*b**2) + 16*x*atanh(tanh(a + b*x))**(11/2)/(231*b**3) - 32*atanh(tanh(a + b*x))**(13/2)/(3003*b**4), Ne(b, 0)), (x**4*atanh(tanh(a))**(5/2)/4, True))`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.66

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(231b^4x^4 + 105ab^3x^3 - 70a^2b^2x^2 + 40a^3bx - 16a^4)(bx + a)^{5/2}}{3003b^4}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`output `2/3003*(231*b^4*x^4 + 105*a*b^3*x^3 - 70*a^2*b^2*x^2 + 40*a^3*b*x - 16*a^4)*(b*x + a)^(5/2)/b^4`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 296 vs.  $2(64) = 128$ .

Time = 0.12 (sec) , antiderivative size = 296, normalized size of antiderivative = 3.70

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\sqrt{2} \left( \frac{429 \sqrt{2} (5 (bx+a)^{7/2} - 21 (bx+a)^{5/2} a + 35 (bx+a)^{3/2} a^2 - 35 \sqrt{bx+aa^3}) a^3}{b^3} + \frac{143 \sqrt{2} (35 (bx+a)^{9/2} - 180 (bx+a)^{7/2} a + 378 (bx+a)^{5/2} a^2 - 420 (bx+a)^{3/2} a^3 + 315 \sqrt{bx+a} a^4) a^2}{b^3} + 65 \sqrt{2} (63 (bx+a)^{11/2} - 385 (bx+a)^{9/2} a + 990 (bx+a)^{7/2} a^2 - 1386 (bx+a)^{5/2} a^3 + 1155 (bx+a)^{3/2} a^4 - 693 \sqrt{bx+a} a^5) a/b^3 + 5 \sqrt{2} (231 (bx+a)^{13/2} - 1638 (bx+a)^{11/2} a + 5005 (bx+a)^{9/2} a^2 - 8580 (bx+a)^{7/2} a^3 + 9009 (bx+a)^{5/2} a^4 - 6006 (bx+a)^{3/2} a^5 + 3003 \sqrt{bx+a} a^6) / b^3 \right)}{b^3}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/15015*sqrt(2)*(429*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a^3/b^3 + 143*sqrt(2)*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)*a^2/b^3 + 65*sqrt(2)*(63*(b*x + a)^(11/2) - 385*(b*x + a)^(9/2)*a + 990*(b*x + a)^(7/2)*a^2 - 1386*(b*x + a)^(5/2)*a^3 + 1155*(b*x + a)^(3/2)*a^4 - 693*sqrt(b*x + a)*a^5)*a/b^3 + 5*sqrt(2)*(231*(b*x + a)^(13/2) - 1638*(b*x + a)^(11/2)*a + 5005*(b*x + a)^(9/2)*a^2 - 8580*(b*x + a)^(7/2)*a^3 + 9009*(b*x + a)^(5/2)*a^4 - 6006*(b*x + a)^(3/2)*a^5 + 3003*sqrt(b*x + a)*a^6)/b^3/b`

**Mupad [B] (verification not implemented)**

Time = 3.35 (sec) , antiderivative size = 2235, normalized size of antiderivative = 27.94

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Too large to display}$$

input `int(x^3*atanh(tanh(a + b*x))^(5/2),x)`

output

```
(2*b^2*x^6*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log
(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/13 + (x^4*(log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1
/2)*((3*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 - (10*(3*b^2*(log(2/(exp(2*a)*exp
(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2
*b*x) - (24*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/13*(log(2/(exp(2*a)*exp(2*b*
x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*
x))/(11*b)))/(9*b) - (x^5*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((
2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (24*b^2*(log(
2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp
(2*b*x) + 1))/2 + b*x))/13*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b
*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(11*b) - (x^3*(lo
g((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*e
xp(2*b*x) + 1))/2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3/4 - (8*((3*b*(log(2/
(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b
*x) + 1)) + 2*b*x)^2)/2 - (10*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - 1
og((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (24*b^...
```

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a))^2 x^3 dx$$

input

```
int(x^3*atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2*x**3,x)
```

### 3.131 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	976
Mathematica [A] (verified)	976
Rubi [A] (verified)	977
Maple [A] (verified)	978
Fricas [A] (verification not implemented)	979
Sympy [A] (verification not implemented)	979
Maxima [A] (verification not implemented)	980
Giac [B] (verification not implemented)	980
Mupad [B] (verification not implemented)	981
Reduce [F]	981

#### Optimal result

Integrand size = 15, antiderivative size = 59

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{8x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{63b^2} + \frac{16 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{693b^3}$$

output

```
2/7*x^2*arctanh(tanh(b*x+a))^(7/2)/b-8/63*x*arctanh(tanh(b*x+a))^(9/2)/b^2
+16/693*arctanh(tanh(b*x+a))^(11/2)/b^3
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{7/2} (99b^2 x^2 - 44bx \operatorname{arctanh}(\tanh(a + bx)) + 8 \operatorname{arctanh}(\tanh(a + bx)))}{693b^3}$$

input

```
Integrate[x^2*ArcTanh[Tanh[a + b*x]]^(5/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2)*(99*b^2*x^2 - 44*b*x*ArcTanh[Tanh[a + b*x]] + 8*ArcTanh[Tanh[a + b*x]]^2))/(693*b^3)
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b}$$

$$\downarrow 2599$$

$$\frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{9/2} dx}{9b} \right)}{7b}$$

$$\downarrow 2588$$

$$\frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{9/2} d \operatorname{arctanh}(\tanh(a + bx))}{9b^2} \right)}{7b}$$

$$\downarrow 15$$

$$\frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{9b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{11/2}}{99b^2} \right)}{7b}$$

input

```
Int[x^2*ArcTanh[Tanh[a + b*x]]^(5/2), x]
```

output

```
(2*x^2*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(9/2))/(9*b) - (4*ArcTanh[Tanh[a + b*x]]^(11/2))/(99*b^2)))/(7*b)
```

## Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_.)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.17

method	result	S
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{11}{2}} + 2(2bx-2 \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}} + 2(bx-\operatorname{arctanh}(\tanh(bx+a)))^2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{b^3}$	6

input `int(x^2*arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `2/b^3*(1/11*arctanh(tanh(b*x+a))^(11/2)+1/9*(2*b*x-2*arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(9/2)+1/7*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(7/2))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.08

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(63b^5x^5 + 161ab^4x^4 + 113a^2b^3x^3 + 3a^3b^2x^2 - 4a^4bx + 8a^5)\sqrt{bx+a}}{693b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `2/693*(63*b^5*x^5 + 161*a*b^4*x^4 + 113*a^2*b^3*x^3 + 3*a^3*b^2*x^2 - 4*a^4*b*x + 8*a^5)*sqrt(b*x + a)/b^3`

**Sympy [A] (verification not implemented)**

Time = 40.76 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.20

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \begin{cases} \frac{2x^2 \operatorname{atanh}^{\frac{7}{2}}(\tanh(a+bx))}{7b} - \frac{8x \operatorname{atanh}^{\frac{9}{2}}(\tanh(a+bx))}{63b^2} + \frac{16 \operatorname{atanh}^{\frac{11}{2}}(\tanh(a+bx))}{693b^3} & \text{for } b \neq 0 \\ \frac{x^3 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a))}{3} & \text{otherwise} \end{cases}$$

input `integrate(x**2*atanh(tanh(b*x+a))**(5/2),x)`

output `Piecewise((2*x**2*atanh(tanh(a + b*x))**(7/2)/(7*b) - 8*x*atanh(tanh(a + b*x))**(9/2)/(63*b**2) + 16*atanh(tanh(a + b*x))**(11/2)/(693*b**3), Ne(b, 0)), (x**3*atanh(tanh(a))**(5/2)/3, True))`



**Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.71

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(63b^3x^3 + 35ab^2x^2 - 20a^2bx + 8a^3)(bx + a)^{5/2}}{693b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `2/693*(63*b^3*x^3 + 35*a*b^2*x^2 - 20*a^2*b*x + 8*a^3)*(b*x + a)^(5/2)/b^3`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(47) = 94.

Time = 0.12 (sec) , antiderivative size = 248, normalized size of antiderivative = 4.20

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\sqrt{2} \left( \frac{231\sqrt{2}(3(bx+a)^{5/2} - 10(bx+a)^{3/2}a + 15\sqrt{bx+aa^2})a^3}{b^2} + \frac{297\sqrt{2}(5(bx+a)^{7/2} - 21(bx+a)^{5/2}a + 35(bx+a)^{3/2}a^2 - 35\sqrt{bx+aa^2})a^3}{b^2} \right)}{693b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/3465*sqrt(2)*(231*sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2)*a^3/b^2 + 297*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a^2/b^2 + 33*sqrt(2)*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)*a/b^2 + 5*sqrt(2)*(63*(b*x + a)^(11/2) - 385*(b*x + a)^(9/2)*a + 990*(b*x + a)^(7/2)*a^2 - 1386*(b*x + a)^(5/2)*a^3 + 1155*(b*x + a)^(3/2)*a^4 - 693*sqrt(b*x + a)*a^5)/b^3`

**Mupad [B] (verification not implemented)**

Time = 3.25 (sec) , antiderivative size = 1789, normalized size of antiderivative = 30.32

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Too large to display}$$

input `int(x^2*atanh(tanh(a + b*x))^(5/2),x)`

output

$$\begin{aligned} & (2*b^2*x^5*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log \\ & (2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/11 + (x^3*(\log((2*\exp(2*a)*\exp(2*b \\ & *x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1 \\ & /2)*((3*b*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/ \\ & (\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2)/2 - (8*(3*b^2*(\log(2/(\exp(2*a)*\exp( \\ & 2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2* \\ & b*x) - (20*b^2*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2 \\ & *b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x))/11*(\log(2/(\exp(2*a)*\exp(2*b*x \\ & ) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 + b*x \\ & ))/(9*b)))/(7*b) - (x^4*(3*b^2*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2* \\ & \exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) - (20*b^2*(\log(2/ \\ & (\exp(2*a)*\exp(2*b*x) + 1))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2 \\ & *b*x) + 1))/2 + b*x))/11*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x \\ & ) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/(9*b) - (x^2*(\log(( \\ & 2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp( \\ & 2*b*x) + 1))/2)^{(1/2)*((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a) \\ & *\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{3/4} - (6*((3*b*(\log(2/(ex \\ & p(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) \\ & + 1)) + 2*b*x)^2)/2 - (8*(3*b^2*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(( \\ & 2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) - (20*b^2*(1... \end{aligned}$$
**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \int \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 x^2 dx$$

input `int(x^2*atanh(tanh(b*x+a))^(5/2),x)`

output `int(sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2*x**2,x)`

### 3.132 $\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	983
Mathematica [A] (verified)	983
Rubi [A] (verified)	984
Maple [A] (verified)	985
Fricas [A] (verification not implemented)	985
Sympy [A] (verification not implemented)	986
Maxima [A] (verification not implemented)	986
Giac [B] (verification not implemented)	986
Mupad [B] (verification not implemented)	987
Reduce [B] (verification not implemented)	988

#### Optimal result

Integrand size = 13, antiderivative size = 38

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{63b^2}$$

output  $2/7*x*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b-4/63*\operatorname{arctanh}(\tanh(b*x+a))^{(9/2)}/b^2$

#### Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(9bx - 2 \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{63b^2}$$

input  $\operatorname{Integrate}[x*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(5/2)}, x]$

output  $(2*(9*b*x - 2*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(7/2)})/(63*b^2)$

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$$

$$\downarrow 2599$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{7b}$$

$$\downarrow 2588$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{7/2} d \operatorname{arctanh}(\tanh(a + bx))}{7b^2}$$

$$\downarrow 15$$

$$\frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{63b^2}$$

input `Int[x*ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(2*x*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b) - (4*ArcTanh[Tanh[a + b*x]]^(9/2))/(63*b^2)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{9}{2}}}{9} + \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{b^2}$	42

input

```
int(x*arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^2*(1/9*arctanh(tanh(b*x+a))^(9/2)+1/7*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(7/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.37

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx = \frac{2(7b^4x^4 + 19ab^3x^3 + 15a^2b^2x^2 + a^3bx - 2a^4)\sqrt{bx+a}}{63b^2}$$

input

```
integrate(x*arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
2/63*(7*b^4*x^4 + 19*a*b^3*x^3 + 15*a^2*b^2*x^2 + a^3*b*x - 2*a^4)*sqrt(b*x + a)/b^2
```

**Sympy [A] (verification not implemented)**

Time = 23.01 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.29

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx = \begin{cases} \frac{2x \operatorname{atanh}^{\frac{7}{2}}(\tanh(a+bx))}{7b} - \frac{4 \operatorname{atanh}^{\frac{9}{2}}(\tanh(a+bx))}{63b^2} & \text{for } b \neq 0 \\ \frac{x^2 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a))}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**(5/2),x)`

output `Piecewise(((2*x*atanh(tanh(a + b*x))**(7/2)/(7*b) - 4*atanh(tanh(a + b*x))**  
*(9/2)/(63*b**2), Ne(b, 0)), (x**2*atanh(tanh(a))**(5/2)/2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int x \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx = \frac{2(7b^2x^2 + 5abx - 2a^2)(bx+a)^{5/2}}{63b^2}$$

input `integrate(x*arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `2/63*(7*b^2*x^2 + 5*a*b*x - 2*a^2)*(b*x + a)^(5/2)/b^2`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 197 vs.  $2(30) = 60$ .

Time = 0.11 (sec) , antiderivative size = 197, normalized size of antiderivative = 5.18

$$\int x \operatorname{arctanh}(\tanh(a$$

$$+bx))^{5/2} dx = \frac{\sqrt{2} \left( \frac{105\sqrt{2}((bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa})a^3}{b} + \frac{63\sqrt{2}(3(bx+a)^{\frac{5}{2}} - 10(bx+a)^{\frac{3}{2}}a + 15\sqrt{bx+aa^2})a^2}{b} + \frac{27\sqrt{2}(5(bx+a)^{\frac{7}{2}} - 21(bx$$

input `integrate(x*arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/315*sqrt(2)*(105*sqrt(2)*((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)*a^3/b + 6  
3*sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2  
) *a^2/b + 27*sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(5/2)*a + 35*(b*x +  
a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3)*a/b + sqrt(2)*(35*(b*x + a)^(9/2) -  
180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3  
+ 315*sqrt(b*x + a)*a^4)/b)/b`

### Mupad [B] (verification not implemented)

Time = 3.22 (sec) , antiderivative size = 773, normalized size of antiderivative = 20.34

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Too large to display}$$

input `int(x*atanh(tanh(a + b*x))^(5/2),x)`



output

```
(log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(
2*b*x) + 1))^3*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - 1
og(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(63*b^2) - (log((exp(2*a)*exp(2*
b*x))/(exp(2*a)*exp(2*b*x) + 1))^4*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*ex
p(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(252*b^2) -
(log(1/(exp(2*a)*exp(2*b*x) + 1))^4*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*e
xp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(252*b^2) +
(log(1/(exp(2*a)*exp(2*b*x) + 1))^3*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*e
xp(2*b*x) + 1))*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 -
log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(63*b^2) - (x*log(1/(exp(2*a)*e
xp(2*b*x) + 1))^3*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2
- log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b) + (x*log((exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^3*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)
*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b) -
(log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*ex
p(2*b*x) + 1))^2*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 -
log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(42*b^2) - (3*x*log(1/(exp(2*a)
)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2*
(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*
exp(2*b*x) + 1))/2)^(1/2))/(28*b) + (3*x*log(1/(exp(2*a)*exp(2*b*x) + 1...
```

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^3 (-2\operatorname{atanh}(\tanh(bx + a)) + 9bx)}{63b^2}$$

input

```
int(x*atanh(tanh(b*x+a))^(5/2),x)
```

output

```
(2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)))^3*(- 2*atanh(tanh(a +
b*x)) + 9*b*x))/(63*b**2)
```

### 3.133 $\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	989
Mathematica [A] (verified)	989
Rubi [A] (verified)	990
Maple [A] (verified)	991
Fricas [B] (verification not implemented)	991
Sympy [A] (verification not implemented)	991
Maxima [A] (verification not implemented)	992
Giac [B] (verification not implemented)	992
Mupad [B] (verification not implemented)	993
Reduce [B] (verification not implemented)	993

#### Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b}$$

output

```
2/7*arctanh(tanh(b*x+a))^(7/2)/b
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b)
```

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{7b}$	15
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{7b}$	15

input `int(arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `2/7*arctanh(tanh(b*x+a))^(7/2)/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(14) = 28.

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.17

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)\sqrt{bx + a}}{7b}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `2/7*(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(b*x + a)/b`

**Sympy [A] (verification not implemented)**

Time = 12.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.44

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \begin{cases} \frac{2 \operatorname{atanh}^{\frac{7}{2}}(\tanh(a+bx))}{7b} & \text{for } b \neq 0 \\ x \operatorname{atanh}^{\frac{5}{2}}(\tanh(a)) & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**(5/2),x)`

output

```
Piecewise((2*atanh(tanh(a + b*x))**(7/2)/(7*b), Ne(b, 0)), (x*atanh(tanh(a
))**(5/2), True))
```

### Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2(bx + a)^{7/2}}{7b}$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")
```

output

```
2/7*(b*x + a)^(7/2)/b
```

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs.  $2(14) = 28$ .

Time = 0.11 (sec) , antiderivative size = 136, normalized size of antiderivative = 7.56

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\sqrt{2} \left( 35 \sqrt{2} \sqrt{bx + a} a^3 + 35 \sqrt{2} \left( (bx + a)^{3/2} - 3 \sqrt{bx + a} a \right) a^2 + 7 \sqrt{2} \left( 3 (bx + a)^{5/2} - 10 (bx + a)^{3/2} \right) a + 15 \sqrt{2} \sqrt{bx + a} a^2 + \sqrt{2} \left( 5 (bx + a)^{7/2} - 21 (bx + a)^{5/2} a + 35 (bx + a)^{3/2} a^2 - 35 \sqrt{2} \sqrt{bx + a} a^3 \right) \right)}{b}$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")
```

output

```
1/35*sqrt(2)*(35*sqrt(2)*sqrt(b*x + a)*a^3 + 35*sqrt(2)*((b*x + a)^(3/2) -
3*sqrt(b*x + a)*a)*a^2 + 7*sqrt(2)*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)
)*a + 15*sqrt(b*x + a)*a^2)*a + sqrt(2)*(5*(b*x + a)^(7/2) - 21*(b*x + a)^(
5/2)*a + 35*(b*x + a)^(3/2)*a^2 - 35*sqrt(b*x + a)*a^3))/b
```

**Mupad [B] (verification not implemented)**

Time = 3.32 (sec) , antiderivative size = 337, normalized size of antiderivative = 18.72

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^3 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{28b}$$

$$- \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^3 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{28b}$$

$$- \frac{3 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)^2 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{28b}$$

$$+ \frac{3 \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)^2 \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{28b}$$

input `int(atanh(tanh(a + b*x))^(5/2), x)`output `(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))^3*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b) - (log(1/(exp(2*a)*exp(2*b*x) + 1)))^3*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b) - (3*log(1/(exp(2*a)*exp(2*b*x) + 1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))^2*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b) + (3*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(28*b)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^3}{7b}$$

input `int(atanh(tanh(b*x+a))^(5/2), x)`

output  $(2\sqrt{\operatorname{atanh}(\tanh(a + b*x))}*\operatorname{atanh}(\tanh(a + b*x))^{**3})/(7*b)$

### 3.134 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx$

Optimal result	995
Mathematica [A] (verified)	996
Rubi [A] (verified)	996
Maple [B] (verified)	998
Fricas [A] (verification not implemented)	998
Sympy [F]	999
Maxima [F]	999
Giac [A] (verification not implemented)	999
Mupad [B] (verification not implemented)	1000
Reduce [F]	1000

#### Optimal result

Integrand size = 15, antiderivative size = 121

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx =$$

$$-2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^{5/2}$$

$$+ 2(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}$$

$$- \frac{2}{3}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}$$

$$+ \frac{2}{5} \operatorname{arctanh}(\tanh(a + bx))^{5/2}$$

output

```
-2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(5/2)+2*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)-2/3*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(3/2)+2/5*arctanh(tanh(b*x+a))^(5/2)
```



**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx = \frac{2}{15} \left( 15b^2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} - 35bx \operatorname{arctanh}(\tanh(a+bx))^{3/2} + 23 \operatorname{arctanh}(\tanh(a+bx))^{5/2} - 15 \operatorname{arctanh} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}} \right) (-bx + \operatorname{arctanh}(\tanh(a+bx))) \right)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x,x]`

output

```
(2*(15*b^2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]] - 35*b*x*ArcTanh[Tanh[a + b*x]]
)^(3/2) + 23*ArcTanh[Tanh[a + b*x]]^(5/2) - 15*ArcTanh[Sqrt[ArcTanh[Tanh[a
+ b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a
+ b*x]])^(5/2))/15
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2590, 2590, 2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx$$

↓ 2590

$$\frac{2}{5} \operatorname{arctanh}(\tanh(a+bx))^{5/2} - (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x} dx$$

↓ 2590

$$\frac{2}{5} \operatorname{arctanh}(\tanh(a + bx))^{5/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx \right)$$

↓ 2590

$$\frac{2}{5} \operatorname{arctanh}(\tanh(a + bx))^{5/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) \right)$$

↓ 2592

$$\frac{2}{5} \operatorname{arctanh}(\tanh(a + bx))^{5/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))} \right) \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x,x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2))/5 - (b*x - ArcTanh[Tanh[a + b*x]])*(-((-2 *ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/3`

### Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 221 vs.  $2(105) = 210$ .

Time = 0.20 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.83

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5} + \frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}} a}{3} + \frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{3} + 2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}$

input `int(arctanh(tanh(b*x+a))^(5/2)/x,x,method=_RETURNVERBOSE)`

output

```
2/5*arctanh(tanh(b*x+a))^(5/2)+2/3*arctanh(tanh(b*x+a))^(3/2)*a+2/3*arctanh(tanh(b*x+a))^(3/2)*(arctanh(tanh(b*x+a))-b*x-a)+2*arctanh(tanh(b*x+a))^(1/2)*a^2+4*a*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2)+2*(arctanh(tanh(b*x+a))-b*x-a)^2*arctanh(tanh(b*x+a))^(1/2)-2*(a^3+3*a^2*(arctanh(tanh(b*x+a))-b*x-a)+3*a*(arctanh(tanh(b*x+a))-b*x-a)^2+(arctanh(tanh(b*x+a))-b*x-a)^3)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x} dx = \left[ a^{\frac{5}{2}} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) + \frac{2}{15} (3b^2x^2 + 11abx + 23a^2)\sqrt{bx+a}, 2\sqrt{-aa^2} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + \frac{2}{15} (3b^2x^2 + 11abx + 23a^2)\sqrt{bx+a} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x,x, algorithm="fricas")`

output

```
[a^(5/2)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2/15*(3*b^2*x^2 + 11*a*b*x + 23*a^2)*sqrt(b*x + a), 2*sqrt(-a)*a^2*arctan(sqrt(-a)/sqrt(b*x + a)) + 2/15*(3*b^2*x^2 + 11*a*b*x + 23*a^2)*sqrt(b*x + a)]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx = \int \frac{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x,x)`

output `Integral(atanh(tanh(a + b*x))**(5/2)/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{5}{2}}}{x} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x, x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx = \frac{1}{15} \sqrt{2} \left( \frac{15 \sqrt{2} a^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + 3 \sqrt{2} (bx + a)^{\frac{5}{2}} + 5 \sqrt{2} (bx + a)^{\frac{3}{2}} a + \dots \right)$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x,x, algorithm="giac")`

output `1/15*sqrt(2)*(15*sqrt(2)*a^3*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 3*sqrt(2)*(b*x + a)^(5/2) + 5*sqrt(2)*(b*x + a)^(3/2)*a + 15*sqrt(2)*sqrt(b*x + a)*a^2)`

**Mupad [B] (verification not implemented)**

Time = 7.14 (sec) , antiderivative size = 789, normalized size of antiderivative = 6.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x,x)`

output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((3*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 - (2*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (8*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + b*x))/5*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + b*x)/(3*b))/b + (2*b^2*x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2))/5 + (2^(1/2)*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i + 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 2^(1/2)*b*x)*16i)/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2)*1i)/8 - (x*(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (8*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + b*x))/...
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{x} dx$$

input `int(atanh(tanh(b*x+a))^(5/2)/x,x)`

output `int((sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2)/x,x)`

### 3.135 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^2} dx$

Optimal result	1002
Mathematica [A] (verified)	1003
Rubi [A] (verified)	1003
Maple [A] (verified)	1005
Fricas [A] (verification not implemented)	1006
Sympy [F]	1006
Maxima [F]	1006
Giac [A] (verification not implemented)	1007
Mupad [B] (verification not implemented)	1007
Reduce [F]	1008

#### Optimal result

Integrand size = 15, antiderivative size = 110

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = 5b \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2} - 5b(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{5}{3}b\operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x}$$

output

```
5*b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(3/2)-5*b*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)+5/3*b*arctanh(tanh(b*x+a))^(3/2)-arctanh(tanh(b*x+a))^(5/2)/x
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx =$$

$$-5b \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right) (-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}$$

$$+ \sqrt{\operatorname{arctanh}(\tanh(a + bx))} \left(\frac{2b^2x}{3} + \frac{14}{3}b(-bx + \operatorname{arctanh}(\tanh(a + bx))) - \frac{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}{x}\right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^2,x]
```

output

```
-5*b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2) + Sqrt[ArcTanh[Tanh[a + b*x]]]*((2*b^2*x)/3 + (14*b*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/3 - (-(b*x) + ArcTanh[Tanh[a + b*x]])^2/x)
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2590, 2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx$$

$$\downarrow \text{2599}$$

$$\frac{5}{2}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x}$$

$$\downarrow \text{2590}$$



$$\frac{5}{2}b \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x}$$

↓ 2590

$$\frac{5}{2}b \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x}$$

↓ 2592

$$\frac{5}{2}b \left( \frac{2}{3} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( 2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))} \right) \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^2, x]`

output `-(ArcTanh[Tanh[a + b*x]]^(5/2)/x) + (5*b*(-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/3)/2`

### Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

## Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.75

method	result
default	$2b \left( \frac{\operatorname{arctanh}(\tanh(bx+a))^{3/2}}{3} + 2\sqrt{\operatorname{arctanh}(\tanh(bx+a))} a + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) \sqrt{\operatorname{arctanh}(\tanh(bx+a))} \right)$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^2,x,method=_RETURNVERBOSE)
```

output

```
2*b*(1/3*arctanh(tanh(b*x+a))^(3/2)+2*arctanh(tanh(b*x+a))^(1/2)*a+2*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2)+(-1/2*a^2-a*(arctanh(tanh(b*x+a))-b*x-a)-1/2*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2)/b/x-5/2*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \left[ \frac{15 a^{\frac{3}{2}} bx \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(2b^2x^2 + 14abx - 3a^2)\sqrt{bx+a}}{6x}, \dots \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^2,x, algorithm="fricas")`

output `[1/6*(15*a^(3/2)*b*x*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(2*b^2*x^2 + 14*a*b*x - 3*a^2)*sqrt(b*x + a))/x, 1/3*(15*sqrt(-a)*a*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) + (2*b^2*x^2 + 14*a*b*x - 3*a^2)*sqrt(b*x + a))/x]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \int \frac{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**2,x)`

output `Integral(atanh(tanh(a + b*x))**(5/2)/x**2, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{5}{2}}}{x^2} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^2,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^2, x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \frac{1}{6} \sqrt{2} \left( \frac{15 \sqrt{2} a^2 \operatorname{arctan} \left( \frac{\sqrt{bx+a}}{\sqrt{-a}} \right)}{\sqrt{-a}} + 2 \sqrt{2} (bx + a)^{3/2} + 12 \sqrt{2} \sqrt{bx + a} a - 3 \sqrt{2} \sqrt{bx + a} a^2 / (bx) \right) b$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^2,x, algorithm="giac")`

output `1/6*sqrt(2)*(15*sqrt(2)*a^2*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 2*sqrt(2)*(b*x + a)^(3/2) + 12*sqrt(2)*sqrt(b*x + a)*a - 3*sqrt(2)*sqrt(b*x + a)*a^2/(b*x))*b`

**Mupad [B] (verification not implemented)**

Time = 7.47 (sec) , antiderivative size = 616, normalized size of antiderivative = 5.60

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^2,x)`

output

```
(2*b^2*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/3 - ((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*x) - ((3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (4*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/b + (2^(1/2)*b*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x)*16i)/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2)*5i)/8
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^2} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{x^2} dx$$

input

```
int(atanh(tanh(b*x+a))^(5/2)/x^2,x)
```

output

```
int((sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2)/x**2,x)
```

### 3.136 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^3} dx$

Optimal result	1009
Mathematica [A] (verified)	1010
Rubi [A] (verified)	1010
Maple [A] (verified)	1012
Fricas [A] (verification not implemented)	1012
Sympy [F]	1013
Maxima [F]	1013
Giac [A] (verification not implemented)	1013
Mupad [B] (verification not implemented)	1014
Reduce [F]	1015

#### Optimal result

Integrand size = 15, antiderivative size = 110

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^3} dx =$$

$$-\frac{15}{4}b^2 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

$$+ \frac{15}{4}b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{5b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{4x} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{2x^2}$$

output

```
-15/4*b^2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(1/2)+15/4*b^2*arctanh(tanh(b*x+a))^(1/2)-5/4*b*arctanh(tanh(b*x+a))^(3/2)/x-1/2*arctanh(tanh(b*x+a))^(5/2)/x^2
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx =$$

$$-15b^2x^2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + 5bx\operatorname{arctanh}(\tanh(a + bx))^{3/2} + 2\operatorname{arctanh}(\tanh(a + bx))^{5/2} + 15b^2x^2$$


---


$$4x^2$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^3,x]
```

output

```
-1/4*(-15*b^2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]] + 5*b*x*ArcTanh[Tanh[a + b*x]]^(3/2) + 2*ArcTanh[Tanh[a + b*x]]^(5/2) + 15*b^2*x^2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])/x^2
```

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2590, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx$$

$$\downarrow 2599$$

$$\frac{5}{4}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{2x^2}$$

$$\downarrow 2599$$

$$\frac{5}{4}b \left( \frac{3}{2}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{2x^2}$$

↓ 2590

$$\frac{5}{4}b \left( \frac{3}{2}b \left( 2\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{2x^2}$$

↓ 2592

$$\frac{5}{4}b \left( \frac{3}{2}b \left( 2\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - 2\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))} \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{2x^2} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^3,x]`

output `-1/2*ArcTanh[Tanh[a + b*x]]^(5/2)/x^2 + (5*b*((3*b*(-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] + 2*Sqrt[ArcTanh[Tanh[a + b*x]]]))/2 - ArcTanh[Tanh[a + b*x]]^(3/2)/x))/4`

### Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`



rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.29

method	result
default	$2b^2 \left( \sqrt{\operatorname{arctanh}(\tanh(bx+a))} + \frac{\left(-\frac{9}{8} \operatorname{arctanh}(\tanh(bx+a)) + \frac{9bx}{8}\right) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}} + \left(\frac{7a^2}{8} + \frac{7a \operatorname{arctanh}(\tanh(bx+a))}{4}\right)}{b^2 x^2} \right)$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^3,x,method=_RETURNVERBOSE)
```

output

```
2*b^2*(arctanh(tanh(b*x+a))^(1/2))+((-9/8*arctanh(tanh(b*x+a))+9/8*b*x)*arctanh(tanh(b*x+a))^(3/2)+(7/8*a^2+7/4*a*(arctanh(tanh(b*x+a))-b*x-a)+7/8*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2))/b^2/x^2-15/8*(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.18

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \left[ \frac{15 \sqrt{ab^2 x^2} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(8b^2 x^2 - 9abx - 2a^2)\sqrt{bx+a}}{8x^2}, \dots \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^3,x, algorithm="fricas")
```

output

```
[1/8*(15*sqrt(a)*b^2*x^2*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*
(8*b^2*x^2 - 9*a*b*x - 2*a^2)*sqrt(b*x + a))/x^2, 1/4*(15*sqrt(-a)*b^2*x^2
*arctan(sqrt(-a)/sqrt(b*x + a)) + (8*b^2*x^2 - 9*a*b*x - 2*a^2)*sqrt(b*x +
a))/x^2]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \int \frac{\operatorname{atanh}^{5/2}(\tanh(a + bx))}{x^3} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(5/2)/x**3,x)
```

output

```
Integral(atanh(tanh(a + b*x))**(5/2)/x**3, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^3} dx$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^3,x, algorithm="maxima")
```

output

```
integrate(arctanh(tanh(b*x + a))^(5/2)/x^3, x)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.84

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \frac{\sqrt{2} \left( \frac{15\sqrt{2}ab^3 \operatorname{arctan}\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + 8\sqrt{2}\sqrt{bx+ab^3} - \frac{\sqrt{2}(9(bx+a)^{\frac{3}{2}}ab^3 - 7\sqrt{bx+aa^2}b^3)}{b^2x^2} \right)}{8b}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^3,x, algorithm="giac")`

output  $\frac{1}{8}\sqrt{2}\cdot(15\sqrt{2}\cdot a\cdot b^3\cdot\arctan(\sqrt{b\cdot x+a}/\sqrt{-a})/\sqrt{-a} + 8\sqrt{2}\cdot\sqrt{b\cdot x+a}\cdot b^3 - \sqrt{2}\cdot(9\cdot(b\cdot x+a)^{(3/2)}\cdot a\cdot b^3 - 7\sqrt{b\cdot x+a}\cdot a^2\cdot b^3)/(b^2\cdot x^2))/b$

### Mupad [B] (verification not implemented)

Time = 4.32 (sec) , antiderivative size = 614, normalized size of antiderivative = 5.58

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^3,x)`

output  $2\cdot b^2\cdot(\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))/2 - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))^{1/2} + b^2\cdot\log((64\cdot(2\cdot 2^{1/2})\cdot a - 2\cdot(\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))/2 - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))^{1/2})\cdot(\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - 2\cdot b\cdot x)^{1/2} - 2^{1/2}\cdot(2\cdot a - \log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) + \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) + 2\cdot b\cdot x + 2^{1/2}\cdot b\cdot x))/(x\cdot(\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - 2\cdot b\cdot x)^{1/2})))\cdot((225\cdot\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)))/128 - (225\cdot\log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)))/128 - (225\cdot b\cdot x)/64)^{1/2} - ((\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)))/2 - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))^{1/2})\cdot(\log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - \log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) + 2\cdot b\cdot x)^3)/(4\cdot x^2\cdot(2\cdot\log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - 2\cdot\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) + 4\cdot b\cdot x)) + (9\cdot b\cdot(\log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)))/2 - \log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1))^{1/2})\cdot(\log(2/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) - \log((2\cdot\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x))/(\exp(2\cdot a)\cdot\exp(2\cdot b\cdot x) + 1)) + 2\cdot b\cdot x))/(8\cdot x)$

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^3} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{x^3} dx$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^3,x)`

output `int((sqrt(atanh(tanh(a + b*x))))*atanh(tanh(a + b*x))**2)/x**3,x)`

### 3.137 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx$

Optimal result	1016
Mathematica [A] (verified)	1017
Rubi [A] (verified)	1017
Maple [A] (verified)	1019
Fricas [A] (verification not implemented)	1019
Sympy [F]	1020
Maxima [F]	1020
Giac [A] (verification not implemented)	1020
Mupad [B] (verification not implemented)	1021
Reduce [F]	1022

#### Optimal result

Integrand size = 15, antiderivative size = 113

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx = \frac{5b^3 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} - \frac{5b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{8x} - \frac{5b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{12x^2} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^3}$$

output

```
5/8*b^3*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^(1/2)-5/8*b^2*arctanh(tanh(b*x+a))^(1/2)/x-5/
12*b*arctanh(tanh(b*x+a))^(3/2)/x^2-1/3*arctanh(tanh(b*x+a))^(5/2)/x^3
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx = \frac{1}{24} \left( -\frac{15b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} - \frac{10b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^2} - \frac{8 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^3} - \frac{15b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}} \right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^4, x]
```

output

```
((-15*b^2*Sqrt[ArcTanh[Tanh[a + b*x]])]/x - (10*b*ArcTanh[Tanh[a + b*x]]^(3/2))/x^2 - (8*ArcTanh[Tanh[a + b*x]]^(5/2))/x^3 - (15*b^3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])/24
```

**Rubi [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^4} dx$$

↓ 2599

$$\frac{5}{6}b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^3}$$

$$\frac{5}{6}b \left( \frac{3}{4}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^3}$$

↓ 2599

$$\frac{5}{6}b \left( \frac{3}{4}b \left( \frac{1}{2}b \int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^3}$$

↓ 2599

$$\frac{5}{6}b \left( \frac{3}{4}b \left( \frac{b \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^3}$$

↓ 2592

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^4, x]`

output `-1/3*ArcTanh[Tanh[a + b*x]]^(5/2)/x^3 + (5*b*((3*b*((b*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]] - Sqrt[ArcTanh[Tanh[a + b*x]]/x])/4 - ArcTanh[Tanh[a + b*x]]^(3/2)/(2*x^2)))/6`

### Defintions of rubi rules used

rule 2592 `Int[1/((u)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLine arQ[u, v, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.27

method	result
default	$2b^3 \left( \frac{-\frac{11}{16} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}} + \left( \frac{5}{6} \operatorname{arctanh}(\tanh(bx+a)) - \frac{5bx}{6} \right) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}} + \left( -\frac{5a^2}{16} - \frac{5a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{8} \right) \operatorname{arctanh}(\tanh(bx+a))^{\frac{1}{2}}}{b^3 x^3} \right)$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^4,x,method=_RETURNVERBOSE)
```

output

```
2*b^3*((-11/16*arctanh(tanh(b*x+a))^(5/2)+(5/6*arctanh(tanh(b*x+a))-5/6*b*x)*arctanh(tanh(b*x+a))^(3/2)+(-5/16*a^2-5/8*a*(arctanh(tanh(b*x+a))-b*x-a)-5/16*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2))/b^3/x^3-5/16/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.27

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \left[ \frac{15 \sqrt{ab^3} x^3 \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(33ab^2x^2 + 26a^2bx + 8a^3)\sqrt{bx - a}}{48ax^3} \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^4,x, algorithm="fricas")
```



output

```
[1/48*(15*sqrt(a)*b^3*x^3*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2
*(33*a*b^2*x^2 + 26*a^2*b*x + 8*a^3)*sqrt(b*x + a))/(a*x^3), 1/24*(15*sqrt
(-a)*b^3*x^3*arctan(sqrt(-a)/sqrt(b*x + a)) - (33*a*b^2*x^2 + 26*a^2*b*x +
8*a^3)*sqrt(b*x + a))/(a*x^3)]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \int \frac{\operatorname{atanh}^{5/2}(\tanh(a + bx))}{x^4} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(5/2)/x**4, x)
```

output

```
Integral(atanh(tanh(a + b*x))**(5/2)/x**4, x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^4} dx$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^4, x, algorithm="maxima")
```

output

```
integrate(arctanh(tanh(b*x + a))^(5/2)/x^4, x)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \frac{1}{48} \sqrt{2} b^3 \left( \frac{15 \sqrt{2} \operatorname{arctan} \left( \frac{\sqrt{bx+a}}{\sqrt{-a}} \right)}{\sqrt{-a}} - \frac{\sqrt{2} \left( 33 (bx + a)^{5/2} - 40 (bx + a)^{3/2} a + \dots \right)}{b^3 x^3} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^4,x, algorithm="giac")`

output  $\frac{1}{48}\sqrt{2}b^3(15\sqrt{2})\arctan(\sqrt{bx+a}/\sqrt{-a})/\sqrt{-a} - \sqrt{2}(33(bx+a)^{5/2} - 40(bx+a)^{3/2}a + 15\sqrt{2}(bx+a)a^2)/(b^3x^3)$

### Mupad [B] (verification not implemented)

Time = 7.74 (sec) , antiderivative size = 669, normalized size of antiderivative = 5.92

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^4,x)`

output  $(2^{1/2}b^3\log((\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx)^{1/2} * ((\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/2 - \log(2/(\exp(2a)\exp(2bx) + 1)))/2)^{1/2} * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx)^{1/2} * 2i - 2^{1/2} * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx) + 2^{1/2} * bx * 64i/x * 5i)/(16 * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx)^{1/2}) - ((\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/2 - \log(2/(\exp(2a)\exp(2bx) + 1)))/2)^{1/2} * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx)^3/(4x^3 * (3\log(2/(\exp(2a)\exp(2bx) + 1)) - 3\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 6bx) - (11b^2 * (\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/2 - \log(2/(\exp(2a)\exp(2bx) + 1)))/2)^{1/2})/(8x) + (13b * (\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)))/2 - \log(2/(\exp(2a)\exp(2bx) + 1)))/2)^{1/2} * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 2bx)^2/(12x^2 * (2\log(2/(\exp(2a)\exp(2bx) + 1)) - 2\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) + 4bx))$

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^4} dx = \frac{-16\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 20\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^3} - \frac{16\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^2} - \frac{20\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x} + \frac{16\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{48x^3}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^4,x)`

output `( - 16*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 20*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 30*sqrt(atanh(tanh(a + b*x)))*b**2*x**2 + 15*int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x),x)*b**3*x**3)/(48*x**3)`

### 3.138 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^5} dx$

Optimal result	1023
Mathematica [A] (verified)	1024
Rubi [A] (verified)	1024
Maple [A] (verified)	1026
Fricas [A] (verification not implemented)	1027
Sympy [F]	1028
Maxima [F]	1028
Giac [A] (verification not implemented)	1028
Mupad [B] (verification not implemented)	1029
Reduce [F]	1029

#### Optimal result

Integrand size = 15, antiderivative size = 167

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^5} dx = \frac{5b^4 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{64(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{5b^3}{64x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{5b^4}{64(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{5b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{32x^2} - \frac{5b\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{24x^3} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

output

```
5/64*b^4*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)))/(b*x-arctanh(tanh(b*x+a)))^(3/2)-5/64*b^3/x/arctanh(tanh(b*x+a))^(1/2)+5/64*b^4/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)-5/32*b^2*arctanh(tanh(b*x+a))^(1/2)/x^2-5/24*b*arctanh(tanh(b*x+a))^(3/2)/x^3-1/4*arctanh(tanh(b*x+a))^(5/2)/x^4
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \frac{5b^4 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{64(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))(15b^3x^3 + 10b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 8bx \operatorname{arctanh}(\tanh(a + bx))^2 - 48 \operatorname{arctanh}(\tanh(a + bx)))}}{192x^4(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^5,x]
```

output

```
(5*b^4*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(64*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2)) - (Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^3*x^3 + 10*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 8*b*x*ArcTanh[Tanh[a + b*x]]^2 - 48*ArcTanh[Tanh[a + b*x]]^3))/(192*x^4*(b*x - ArcTanh[Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.64 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx$$

↓ 2599

$$\frac{5}{8}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^4} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{4x^4}$$

↓ 2599

$$\frac{5}{8}b \left( \frac{1}{2}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^3} dx - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

↓ 2599

$$\frac{5}{8}b \left( \frac{1}{2}b \left( \frac{1}{4}b \int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

↓ 2599

$$\frac{5}{8}b \left( \frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2x^2} \right) \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

↓ 2594

$$\frac{5}{8}b \left( \frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \right) \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

↓ 2592

$$\frac{5}{8}b \left( \frac{1}{2}b \left( \frac{1}{4}b \left( -\frac{1}{2}b \left( -\frac{2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \right) \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{4x^4}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^5, x]
```

output

```
-1/4*ArcTanh[Tanh[a + b*x]]^(5/2)/x^4 + (5*b*((b*((b*(-1/2*(b*(-2*ArcTan[
Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]))/(b*x - A
rcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcT
anh[Tanh[a + b*x]]])) - 1/(x*Sqrt[ArcTanh[Tanh[a + b*x]]])))/4 - Sqrt[Arc
Tanh[Tanh[a + b*x]]/(2*x^2))/2 - ArcTanh[Tanh[a + b*x]]^(3/2)/(3*x^3))/8
```

### Defintions of rubi rules used

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simpli
fy[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v
)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLine
arQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n +
1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew
iseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m +
1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n
}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

### Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.01

method	result
default	$2b^4 \left( \frac{5 \operatorname{arctanh}(\tanh(bx+a)) \frac{7}{2}}{128(\operatorname{arctanh}(\tanh(bx+a))-bx)} - \frac{73 \operatorname{arctanh}(\tanh(bx+a)) \frac{5}{2}}{384} + \left( \frac{55 \operatorname{arctanh}(\tanh(bx+a))}{384} - \frac{55bx}{384} \right) \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} + \left( -\frac{5a^2}{128} - \frac{5a}{128} \right) \frac{1}{b^4 x^4} \right)$

input `int(arctanh(tanh(b*x+a))^(5/2)/x^5,x,method=_RETURNVERBOSE)`

output `2*b^4*((-5/128/(arctanh(tanh(b*x+a))-b*x)*arctanh(tanh(b*x+a))^(7/2)-73/384*arctanh(tanh(b*x+a))^(5/2)+(55/384*arctanh(tanh(b*x+a))-55/384*b*x)*arctanh(tanh(b*x+a))^(3/2)+(-5/128*a^2-5/64*a*(arctanh(tanh(b*x+a))-b*x-a)-5/128*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2))/b^4/x^4+5/128/(arctanh(tanh(b*x+a))-b*x)^(3/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \left[ \frac{15 \sqrt{ab^4} x^4 \log\left(\frac{bx + 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) - 2(15ab^3x^3 + 118a^2b^2x^2 + 136a^3bx + 48a^4)\sqrt{bx+a}}{384a^2x^4} - \frac{15\sqrt{-ab^4}x^4 \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (15ab^3x^3 + 118a^2b^2x^2 + 136a^3bx + 48a^4)\sqrt{bx+a}}{192a^2x^4} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^5,x, algorithm="fricas")`

output `[1/384*(15*sqrt(a)*b^4*x^4*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(15*a*b^3*x^3 + 118*a^2*b^2*x^2 + 136*a^3*b*x + 48*a^4)*sqrt(b*x + a))/(a^2*x^4), -1/192*(15*sqrt(-a)*b^4*x^4*arctan(sqrt(-a)/sqrt(b*x + a)) + (15*a*b^3*x^3 + 118*a^2*b^2*x^2 + 136*a^3*b*x + 48*a^4)*sqrt(b*x + a))/(a^2*x^4)]`



**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \int \frac{\operatorname{atanh}^{5/2}(\tanh(a + bx))}{x^5} dx$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**5,x)`

output `Integral(atanh(tanh(a + b*x))**(5/2)/x**5, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^5} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^5,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^5, x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.65

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \frac{\sqrt{2} \left( \frac{15\sqrt{2}b^5 \operatorname{arctan}\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + \frac{\sqrt{2}(15(bx+a)^{7/2}b^5 + 73(bx+a)^{5/2}ab^5 - 55(bx+a)^{3/2}a^2b^5 + 15\sqrt{bx+aa^3}b^5)}{ab^4x^4} \right)}{384b}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^5,x, algorithm="giac")`

output `-1/384*sqrt(2)*(15*sqrt(2)*b^5*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a) + sqrt(2)*(15*(b*x + a)^(7/2)*b^5 + 73*(b*x + a)^(5/2)*a*b^5 - 55*(b*x + a)^(3/2)*a^2*b^5 + 15*sqrt(b*x + a)*a^3*b^5)/(a*b^4*x^4)/b`

**Mupad [B] (verification not implemented)**

Time = 7.84 (sec) , antiderivative size = 1069, normalized size of antiderivative = 6.40

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^5,x)`

output

$$\begin{aligned} & (5*b^3*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/(32*x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)) - ((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3}/(4*x^4*(4*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - 4*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 8*b*x)) + (2^{(1/2)}*b^4*\log(((2*2^{(1/2)}*a + (\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)*2i} - 2^{(1/2)}*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) + 2^{(1/2)}*b*x*((2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))*1024i)/(x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)}))*5i)/(64*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))... \end{aligned}$$
**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^5} dx = \frac{-48\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 40\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x^5}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^5,x)`

output

```
( - 48*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 40*sqrt(atanh(
tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 30*sqrt(atanh(tanh(a + b*x)))*b
**2*x**2 + 15*int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x**2),x
)*b**3*x**4)/(192*x**4)
```

**3.139**      $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx$

Optimal result . . . . .	1031
Mathematica [A] (verified) . . . . .	1032
Rubi [A] (verified) . . . . .	1033
Maple [A] (verified) . . . . .	1036
Fricas [A] (verification not implemented) . . . . .	1036
Sympy [F] . . . . .	1037
Maxima [F] . . . . .	1037
Giac [A] (verification not implemented) . . . . .	1037
Mupad [B] (verification not implemented) . . . . .	1038
Reduce [F] . . . . .	1038

**Optimal result**

Integrand size = 15, antiderivative size = 221

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx = \frac{3b^5 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{128(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}}$$

$$+ \frac{b^4}{128x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{b^5}{128(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{64x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b^5}$$

$$+ \frac{128(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{16x^3}$$

$$- \frac{b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{8x^4} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^5}$$

output

```
3/128*b^5*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)))/(b*x-arctanh(tanh(b*x+a)))^(5/2)+1/128*b^4/x/arctanh(tanh(b*x+a))^(3/2)-1/128*b^5/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)-1/64*b^3/x^2/arctanh(tanh(b*x+a))^(1/2)+3/128*b^5/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(1/2)-1/16*b^2*arctanh(tanh(b*x+a))^(1/2)/x^3-1/8*b*arctanh(tanh(b*x+a))^(3/2)/x^4-1/5*arctanh(tanh(b*x+a))^(5/2)/x^5
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.68

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \frac{1}{640} \left( -\frac{15b^5 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} \right.$$

$$\left. - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}(15b^4x^4 + 10b^3x^3 \operatorname{arctanh}(\tanh(a + bx)) + 8b^2x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 176b^2x \operatorname{arctanh}(\tanh(a + bx)) + 128b \operatorname{arctanh}(\tanh(a + bx))^3 - 128 \operatorname{arctanh}(\tanh(a + bx))^4)}{x^5(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2} \right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^6,x]
```

output

```
((-15*b^5*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2) - (Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^4*x^4 + 10*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 8*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 176*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4))/(x^5*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2))/640
```

**Rubi [A] (verified)**

Time = 0.72 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$ , Rules used = {2599, 2599, 2599, 2599, 2599, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^5} dx - \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^5} \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}b \left( \frac{3}{8}b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^4} dx - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{4x^4} \right) - \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^5} \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^3} \right) - \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{4x^4} \right) - \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^5} \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \left( -\frac{1}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \right) - \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^3} \right) - \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^5} \\
 & \quad \downarrow \text{2599}
 \end{aligned}$$

$$\frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) \frac{1}{5x^5}$$

↓ 2594

$$\frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))} \right) - \frac{2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) \right) \frac{1}{5x^5}$$

↓ 2594

$$\frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{2}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) - \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) \frac{1}{5x^5}$$

↓ 2592

$$\frac{1}{2}b \left( \frac{3}{8}b \left( \frac{1}{6}b \left( -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) - \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) \frac{1}{5x^5}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^6, x]
```

output

```
-1/5*ArcTanh[Tanh[a + b*x]]^(5/2)/x^5 + (b*((3*b*((b*(-1/4*(b*((-3*b*(-(((
-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])
)/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])
*Sqrt[ArcTanh[Tanh[a + b*x]]])))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(3*(b*
x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))))/2 - 1/(x*ArcTa
nh[Tanh[a + b*x]]^(3/2)))) - 1/(2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]])))/6 -
Sqrt[ArcTanh[Tanh[a + b*x]]/(3*x^3))/8 - ArcTanh[Tanh[a + b*x]]^(3/2)/(4
*x^4))/2
```

### Defintions of rubi rules used

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simpli
fy[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v
)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLine
arQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n +
1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew
iseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```



**Maple [A] (verified)**

Time = 1.37 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.19

method	result
default	$2b^5 \left( \frac{3 \operatorname{arctanh}(\tanh(bx+a)) \frac{9}{2}}{256(a^2+2a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)+(\operatorname{arctanh}(\tanh(bx+a))-bx-a)^2)} - \frac{7 \operatorname{arctanh}(\tanh(bx+a)) \frac{7}{2}}{128(\operatorname{arctanh}(\tanh(bx+a))-bx)} - \frac{\operatorname{arctanh}(\tanh(bx+a)) \frac{5}{2}}{10} + \dots \right)$

input `int(arctanh(tanh(b*x+a))^(5/2)/x^6,x,method=_RETURNVERBOSE)`

output  $2*b^5*((3/256/(a^2+2*a*(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)+(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)^2)*\operatorname{arctanh}(\tanh(b*x+a))^(9/2)-7/128/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)*\operatorname{arctanh}(\tanh(b*x+a))^(7/2)-1/10*\operatorname{arctanh}(\tanh(b*x+a))^(5/2)+(7/128*\operatorname{arctanh}(\tanh(b*x+a))-7/128*b*x)*\operatorname{arctanh}(\tanh(b*x+a))^(3/2)+(-3/256*a^2-3/128*a*(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)-3/256*(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)^2)*\operatorname{arctanh}(\tanh(b*x+a))^(1/2))/b^5/x^5-3/256/(a^2+2*a*(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)+(\operatorname{arctanh}(\tanh(b*x+a))-b*x-a)^2)/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^(1/2)*\operatorname{arctanh}(\operatorname{arctanh}(\tanh(b*x+a))^(1/2)/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^(1/2)))$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.84

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^6} dx = \left[ \frac{15 \sqrt{ab^5} x^5 \log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(15ab^4x^4 - 10a^2b^3x^3 - 248a^3b^2x^2 - 336a^4bx - 128a^5)\sqrt{bx+a}}{1280a^3x^5} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^6,x, algorithm="fricas")`

output  $[1/1280*(15*\sqrt{a}*b^5*x^5*\log((b*x-2*\sqrt{b*x+a})*\sqrt{a+2*a})/x)+2*(15*a*b^4*x^4-10*a^2*b^3*x^3-248*a^3*b^2*x^2-336*a^4*b*x-128*a^5)*\sqrt{b*x+a})/(a^3*x^5), 1/640*(15*\sqrt{-a}*b^5*x^5*\operatorname{arctan}(\sqrt{-a}/\sqrt{b*x+a})+(15*a*b^4*x^4-10*a^2*b^3*x^3-248*a^3*b^2*x^2-336*a^4*b*x-128*a^5)*\sqrt{b*x+a})/(a^3*x^5)]$

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \int \frac{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))}{x^6} dx$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**6,x)`

output `Integral(atanh(tanh(a + b*x))**(5/2)/x**6, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{5}{2}}}{x^6} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^6,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^6, x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \frac{1}{1280} \sqrt{2} b^5 \left( \frac{15 \sqrt{2} \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} + \frac{\sqrt{2} \left( 15 (bx + a)^{\frac{9}{2}} - 70 (bx + a)^{\frac{7}{2}} a \right)}{\dots} \right)$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^6,x, algorithm="giac")`

output `1/1280*sqrt(2)*b^5*(15*sqrt(2)*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) + sqrt(2)*(15*(b*x + a)^(9/2) - 70*(b*x + a)^(7/2)*a - 128*(b*x + a)^(5/2)*a^2 + 70*(b*x + a)^(3/2)*a^3 - 15*sqrt(b*x + a)*a^4)/(a^2*b^5*x^5)`

**Mupad [B] (verification not implemented)**

Time = 8.65 (sec) , antiderivative size = 1292, normalized size of antiderivative = 5.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \text{Too large to display}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^6,x)`

output

$$\begin{aligned} & (3*b^4*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/(32*x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2) - \\ & (\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3/(4*x^5*(5*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - 5*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 10*b*x)) + \\ & (b^3*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/(16*x^2*(2*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - 2*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 4*b*x)) + \\ & (2^{(1/2)}*b^5*\log(((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)}*2i - 2^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) + \\ & 2^{(1/2)}*b*x)*((2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^5 + 40*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - \\ & 80*a^3*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 - 32*a^5 - 10*... \end{aligned}$$
**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^6} dx = \frac{-32\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 20\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x^6}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^6,x)`

output

```
( - 32*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 20*sqrt(atanh(
tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 10*sqrt(atanh(tanh(a + b*x)))*b
**2*x**2 + 5*int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x**3),x)
*b**3*x**5)/(160*x**5)
```

$$3.140 \quad \int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1040
Mathematica [A] (verified)	1041
Rubi [A] (verified)	1041
Maple [A] (verified)	1043
Fricas [A] (verification not implemented)	1044
Sympy [F]	1044
Maxima [A] (verification not implemented)	1045
Giac [A] (verification not implemented)	1045
Mupad [B] (verification not implemented)	1046
Reduce [F]	1046

### Optimal result

Integrand size = 15, antiderivative size = 99

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{16x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^2} + \frac{32x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b^3} - \frac{128x \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^4} + \frac{256 \operatorname{arctanh}(\tanh(a+bx))^{9/2}}{315b^5}$$

output

```
2*x^4*arctanh(tanh(b*x+a))^(1/2)/b-16/3*x^3*arctanh(tanh(b*x+a))^(3/2)/b^2
+32/5*x^2*arctanh(tanh(b*x+a))^(5/2)/b^3-128/35*x*arctanh(tanh(b*x+a))^(7/
2)/b^4+256/315*arctanh(tanh(b*x+a))^(9/2)/b^5
```

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.84

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))(315b^4x^4 - 840b^3x^3\operatorname{arctanh}(\tanh(a + bx)) + 1008b^2x^2\operatorname{arctanh}(\tanh(a + bx)))}{315b^5}$$

input `Integrate[x^4/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(315*b^4*x^4 - 840*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 1008*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 576*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4))/(315*b^5)`

### Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.20, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow 2599$$

$$\frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \int x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx}{b}$$

$$\downarrow 2599$$

$$\frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{b} \right)}{b}$$

$$\downarrow 2599$$

$$\begin{aligned}
 & \frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx}{5b} \right)}{b} \right)}{b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{5b} \right)}{b} \right)}{b} \right)}{b} \\
 & \quad \downarrow 2588 \\
 & \frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{7/2} dx}{5b} \right) \operatorname{arctanh}(\tanh(a + bx))}{7b^2} \right)}{b} \right)}{b} \\
 & \quad \downarrow 15 \\
 & \frac{2x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{8 \left( \frac{2x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7b} - \frac{4 \operatorname{arctanh}(\tanh(a + bx))^{9/2}}{63b^2} \right)}{5b} \right)}{b} \right)}{b}
 \end{aligned}$$

input `Int [x^4/Sqrt [ArcTanh [Tanh [a + b*x]]] , x]`

output

$$\frac{(2x^4 \sqrt{\operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]})/b - (8((2x^3 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])^{(3/2)})/(3b) - (2((2x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])^{(5/2)})/(5b) - (4((2x \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])^{(7/2)})/(7b) - (4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{(9/2)})/(63b^2)))/(5b)))/b$$

### Defintions of rubi rules used

rule 15

$$\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^{(m+1)})/(m+1), x] \;/; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$$

rule 2588

$$\operatorname{Int}[(u_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] \;/; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$$

rule 2599

$$\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}(v^{(n/(a*(m+1))})], x] - \operatorname{Simp}[b*(n/(a*(m+1))) \operatorname{Int}[u^{(m+1)}v^{(n-1)}, x], x] \;/; \operatorname{NeQ}[b*u - a*v, 0] \;/; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m+n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.55

method	result
default	$\frac{2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{\frac{9}{2}}}{9} + \frac{2(-4 \operatorname{arctanh}(\operatorname{tanh}(bx+a))+4bx) \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{\frac{7}{2}}}{7} + \frac{2(2(bx - \operatorname{arctanh}(\operatorname{tanh}(bx+a)))^2 + (2bx - 2 \operatorname{arctanh}(\operatorname{tanh}(bx+a)))^2)}{5}$

input

$$\operatorname{int}(x^4/\operatorname{arctanh}(\operatorname{tanh}(bx+a))^{(1/2)}, x, \operatorname{method}=\_RETURNVERBOSE)$$



output

```
2/b^5*(1/9*arctanh(tanh(b*x+a))^(9/2)+1/7*(-4*arctanh(tanh(b*x+a))+4*b*x)*
arctanh(tanh(b*x+a))^(7/2)+1/5*(2*(b*x-arctanh(tanh(b*x+a)))^2+(2*b*x-2*ar
ctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(5/2)+2/3*(b*x-arctanh(tanh(b*
x+a)))^2*(2*b*x-2*arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(3/2)+(b*x-ar
ctanh(tanh(b*x+a)))^4*arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.54

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$= \frac{2(35b^4x^4 - 40ab^3x^3 + 48a^2b^2x^2 - 64a^3bx + 128a^4)\sqrt{bx+a}}{315b^5}$$

input

```
integrate(x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
2/315*(35*b^4*x^4 - 40*a*b^3*x^3 + 48*a^2*b^2*x^2 - 64*a^3*b*x + 128*a^4)*
sqrt(b*x + a)/b^5
```

**Sympy [F]**

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{x^4}{\sqrt{\operatorname{atanh}(\tanh(a+bx))}} dx$$

input

```
integrate(x**4/atanh(tanh(b*x+a))**(1/2),x)
```

output

```
Integral(x**4/sqrt(atanh(tanh(a + b*x))), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2(35b^5x^5 - 5ab^4x^4 + 8a^2b^3x^3 - 16a^3b^2x^2 + 64a^4bx + 128a^5)}{315\sqrt{bx + ab^5}}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`output `2/315*(35*b^5*x^5 - 5*a*b^4*x^4 + 8*a^2*b^3*x^3 - 16*a^3*b^2*x^2 + 64*a^4*b*x + 128*a^5)/(sqrt(b*x + a)*b^5)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.62

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2\left(35(bx + a)^{\frac{9}{2}} - 180(bx + a)^{\frac{7}{2}}a + 378(bx + a)^{\frac{5}{2}}a^2 - 420(bx + a)^{\frac{3}{2}}a^3 + 315\sqrt{bx + a}a^4\right)}{315b^5}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`output `2/315*(35*(b*x + a)^(9/2) - 180*(b*x + a)^(7/2)*a + 378*(b*x + a)^(5/2)*a^2 - 420*(b*x + a)^(3/2)*a^3 + 315*sqrt(b*x + a)*a^4)/b^5`

**Mupad [B] (verification not implemented)**

Time = 3.12 (sec) , antiderivative size = 496, normalized size of antiderivative = 5.01

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \text{Too large to display}$$

input `int(x^4/atanh(tanh(a + b*x))^(1/2),x)`

output

```
(2*x^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(9*b) + (256*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^4)/(315*b^5) + (16*x^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/(63*b^2) + (128*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^3)/(315*b^4) + (32*x^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)^2)/(105*b^3)
```

**Reduce [F]**

$$\int \frac{x^4}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^4}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^4/atanh(tanh(b*x+a))^(1/2),x)`output `int((sqrt(atanh(tanh(a + b*x)))**4)/atanh(tanh(a + b*x)),x)`

**3.141**  $\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1047
Mathematica [A] (verified)	1048
Rubi [A] (verified)	1048
Maple [A] (verified)	1050
Fricas [A] (verification not implemented)	1050
Sympy [F]	1051
Maxima [A] (verification not implemented)	1051
Giac [A] (verification not implemented)	1051
Mupad [B] (verification not implemented)	1052
Reduce [F]	1053

**Optimal result**

Integrand size = 15, antiderivative size = 76

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b^2} + \frac{16x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b^3} - \frac{32 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^4}$$

output

$2*x^3*\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}/b-4*x^2*\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}/b^2+16/5*x*\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}/b^3-32/35*\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}/b^4$

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))(35b^3x^3 - 70b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 56bx\operatorname{arctanh}(\tanh(a + bx))^2 - 16\operatorname{arctanh}(\tanh(a + bx))^3)}{35b^4}$$

input `Integrate[x^3/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(35*b^3*x^3 - 70*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 56*b*x*ArcTanh[Tanh[a + b*x]]^2 - 16*ArcTanh[Tanh[a + b*x]]^3))/(35*b^4)`

### Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow 2599$$

$$\frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{6 \int x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx}{b}$$

$$\downarrow 2599$$

$$\frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{3b} \right)}{b}$$

$$\downarrow 2599$$

$$\begin{array}{c}
 \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \\
 6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx}{5b} \right)}{3b} \right) \\
 \hline
 \downarrow \text{2588} \\
 \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \\
 6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{5/2} d \operatorname{arctanh}(\tanh(a+bx))}{5b^2} \right)}{3b} \right) \\
 \hline
 \downarrow \text{15} \\
 \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \\
 6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^2} \right)}{3b} \right) \\
 \hline
 \downarrow \\
 \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \\
 6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^2} \right)}{3b} \right)
 \end{array}$$

input `Int[x^3/Sqrt[ArcTanh[Tanh[a + b*x]]], x]`

output `(2*x^3*Sqrt[ArcTanh[Tanh[a + b*x]]])/b - (6*((2*x^2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (4*ArcTanh[Tanh[a + b*x]]^(7/2))/(35*b^2)))/(3*b)))/b`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.62

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}} + 2(-3 \operatorname{arctanh}(\tanh(bx+a)) + 3bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}} + 2((bx - \operatorname{arctanh}(\tanh(bx+a)))(2bx - 2 \operatorname{arctanh}(\tanh(bx+a)))}{b^4}$

input

```
int(x^3/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^4*(1/7*arctanh(tanh(b*x+a))^(7/2)+1/5*(-3*arctanh(tanh(b*x+a))+3*b*x)*arctanh(tanh(b*x+a))^(5/2)+1/3*((b*x-arctanh(tanh(b*x+a)))*(2*b*x-2*arctanh(tanh(b*x+a)))+(b*x-arctanh(tanh(b*x+a)))^2)*arctanh(tanh(b*x+a))^(3/2)+(b*x-arctanh(tanh(b*x+a)))^3*arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(5b^3x^3 - 6ab^2x^2 + 8a^2bx - 16a^3)\sqrt{bx + a}}{35b^4}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
2/35*(5*b^3*x^3 - 6*a*b^2*x^2 + 8*a^2*b*x - 16*a^3)*sqrt(b*x + a)/b^4
```

**Sympy [F]**

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^3}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(x**3/atanh(tanh(b*x+a))**(1/2), x)`

output `Integral(x**3/sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.70

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(5b^4x^4 - ab^3x^3 + 2a^2b^2x^2 - 8a^3bx - 16a^4)}{35\sqrt{bx + ab^4}}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")`

output `2/35*(5*b^4*x^4 - a*b^3*x^3 + 2*a^2*b^2*x^2 - 8*a^3*b*x - 16*a^4)/(sqrt(b*x + a)*b^4)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.64

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\left(5(bx + a)^{\frac{7}{2}} - 21(bx + a)^{\frac{5}{2}}a + 35(bx + a)^{\frac{3}{2}}a^2 - 35\sqrt{bx + a}a^3\right)}{35b^4}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(1/2), x, algorithm="giac")`



output

$$2/35*(5*(b*x + a)^{(7/2)} - 21*(b*x + a)^{(5/2)*a + 35*(b*x + a)^{(3/2)*a^2 - 35*\sqrt{b*x + a}*a^3)/b^4$$

**Mupad [B] (verification not implemented)**

Time = 3.01 (sec) , antiderivative size = 385, normalized size of antiderivative = 5.07

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2x^3 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{7b}$$

$$+ \frac{32 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left(\frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} + bx\right)^3}{35b^4}$$

$$+ \frac{12x^2 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left(\frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} + bx\right)}{35b^2}$$

$$+ \frac{16x \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left(\frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} + bx\right)^2}{35b^3}$$

input

$$\operatorname{int}(x^3/\operatorname{atanh}(\tanh(a + b*x))^{(1/2)}, x)$$

output

$$(2*x^3*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)}/(7*b) + (32*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x)^3/(35*b^4) + (12*x^2*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x)/(35*b^2) + (16*x*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x)^2/(35*b^3)$$

**Reduce [F]**

$$\int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^3/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(atanh(tanh(a + b*x))))*x**3)/atanh(tanh(a + b*x)),x)`

**3.142**  $\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1054
Mathematica [A] (verified)	1054
Rubi [A] (verified)	1055
Maple [A] (verified)	1056
Fricas [A] (verification not implemented)	1057
Sympy [F]	1057
Maxima [A] (verification not implemented)	1057
Giac [A] (verification not implemented)	1058
Mupad [B] (verification not implemented)	1058
Reduce [F]	1059

**Optimal result**

Integrand size = 15, antiderivative size = 57

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{8x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^2} + \frac{16 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{15b^3}$$

output

```
2*x^2*arctanh(tanh(b*x+a))^(1/2)/b-8/3*x*arctanh(tanh(b*x+a))^(3/2)/b^2+16/15*arctanh(tanh(b*x+a))^(5/2)/b^3
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.86

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))(15b^2x^2 - 20bx \operatorname{arctanh}(\tanh(a+bx)) + 8 \operatorname{arctanh}(\tanh(a+bx))^2)}}{15b^3}$$

input `Integrate[x^2/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^2*x^2 - 20*b*x*ArcTanh[Tanh[a + b*x]] + 8*ArcTanh[Tanh[a + b*x]]^2))/(15*b^3)`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \int x \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx}{3b} \right)}{b} \\
 & \quad \downarrow 2588 \\
 & \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{3/2} d \operatorname{arctanh}(\tanh(a+bx))}{3b^2} \right)}{b} \\
 & \quad \downarrow 15 \\
 & \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{15b^2} \right)}{b}
 \end{aligned}$$

input `Int[x^2/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output  $(2x^2\sqrt{\operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]})/b - (4((2x\operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])^{(3/2)})/(3b) - (4\operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^{(5/2)})/(15b^2)))/b$

### Defintions of rubi rules used

rule 15  $\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^{(m+1)})/(m+1), x] /; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$

rule 2588  $\operatorname{Int}[(u_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$

rule 2599  $\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}(v^n/(a*(m+1))), x] - \operatorname{Simp}[b*(n/(a*(m+1))) \operatorname{Int}[u^{(m+1)}v^{(n-1)}, x], x] /; \operatorname{NeQ}[b*u - a*v, 0] /; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m+n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$

### Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.19

method	result
default	$\frac{2\operatorname{arctanh}(\operatorname{tanh}(bx+a))^{\frac{5}{2}}}{5} + \frac{2(2bx-2\operatorname{arctanh}(\operatorname{tanh}(bx+a)))\operatorname{arctanh}(\operatorname{tanh}(bx+a))^{\frac{3}{2}}}{3} + 2(bx-\operatorname{arctanh}(\operatorname{tanh}(bx+a)))^2\sqrt{\operatorname{arctanh}(\operatorname{tanh}(bx+a))}$

input `int(x^2/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output  $2/b^3*(1/5*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{(5/2)}+1/3*(2*b*x-2*\operatorname{arctanh}(\operatorname{tanh}(b*x+a)))*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{(3/2)}+(b*x-\operatorname{arctanh}(\operatorname{tanh}(b*x+a)))^2*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{(1/2)})$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.54

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(3b^2x^2 - 4abx + 8a^2)\sqrt{bx + a}}{15b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `2/15*(3*b^2*x^2 - 4*a*b*x + 8*a^2)*sqrt(b*x + a)/b^3`

**Sympy [F]**

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^2}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(x**2/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(x**2/sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.74

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(3b^3x^3 - ab^2x^2 + 4a^2bx + 8a^3)}{15\sqrt{bx + ab^3}}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `2/15*(3*b^3*x^3 - a*b^2*x^2 + 4*a^2*b*x + 8*a^3)/(sqrt(b*x + a)*b^3)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.65

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2 \left( 3 (bx + a)^{\frac{5}{2}} - 10 (bx + a)^{\frac{3}{2}} a + 15 \sqrt{bx + aa^2} \right)}{15 b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `2/15*(3*(b*x + a)^(5/2) - 10*(b*x + a)^(3/2)*a + 15*sqrt(b*x + a)*a^2)/b^3`

**Mupad [B] (verification not implemented)**

Time = 3.05 (sec) , antiderivative size = 211, normalized size of antiderivative = 3.70

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( 15b^2x^2 - 10bx \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 10bx \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2 \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) \right)}{15b^3}$$

input `int(x^2/atanh(tanh(a + b*x))^(1/2),x)`

output `(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(2*log(2/(exp(2*a)*exp(2*b*x) + 1))^2 - 4*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 15*b^2*x^2 - 10*b*x*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 10*b*x*log(2/(exp(2*a)*exp(2*b*x) + 1)))/(15*b^3)`

**Reduce [F]**

$$\int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^2/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(atanh(tanh(a + b*x))))*x**2)/atanh(tanh(a + b*x)),x)`



$$3.143 \quad \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1060
Mathematica [A] (verified)	1060
Rubi [A] (verified)	1061
Maple [A] (verified)	1062
Fricas [A] (verification not implemented)	1063
Sympy [F]	1063
Maxima [A] (verification not implemented)	1063
Giac [A] (verification not implemented)	1064
Mupad [B] (verification not implemented)	1064
Reduce [B] (verification not implemented)	1065

### Optimal result

Integrand size = 13, antiderivative size = 36

$$\begin{aligned} & \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\ &= \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^2} \end{aligned}$$

output `2*x*arctanh(tanh(b*x+a))^(1/2)/b-4/3*arctanh(tanh(b*x+a))^(3/2)/b^2`

### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.89

$$\begin{aligned} & \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\ &= \frac{2(3bx - 2\operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b^2} \end{aligned}$$

input `Integrate[x/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output

```
(2*(3*b*x - 2*ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]/(3*b^2
)
```

**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} \\
 & \quad \downarrow \text{2588} \\
 & \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a + bx))} d\operatorname{arctanh}(\tanh(a + bx))}{b^2} \\
 & \quad \downarrow \text{15} \\
 & \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{4\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b^2}
 \end{aligned}$$

input

```
Int [x/Sqrt [ArcTanh [Tanh [a + b*x]]] , x]
```

output

```
(2*x*Sqrt [ArcTanh [Tanh [a + b*x]]])/b - (4*ArcTanh [Tanh [a + b*x]]^(3/2))/(3
*b^2)
```

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.56

method	result	size
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{3/2} - 2\sqrt{\operatorname{arctanh}(\tanh(bx+a))} a - 2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b^2}$	56

input `int(x/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2/b^2*(1/3*arctanh(tanh(b*x+a))^(3/2)-arctanh(tanh(b*x+a))^(1/2)*a-(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{bx + a}(bx - 2a)}{3b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `2/3*sqrt(b*x + a)*(b*x - 2*a)/b^2`

**Sympy [F]**

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(x/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(x/sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(b^2x^2 - abx - 2a^2)}{3\sqrt{bx + ab^2}}$$

input `integrate(x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `2/3*(b^2*x^2 - a*b*x - 2*a^2)/(sqrt(b*x + a)*b^2)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2 \left( (bx + a)^{\frac{3}{2}} - 3 \sqrt{bx + a} a \right)}{3 b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`output `2/3*((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)/b^2`**Mupad [B] (verification not implemented)**

Time = 3.11 (sec) , antiderivative size = 105, normalized size of antiderivative = 2.92

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}{\frac{2}{e^{2a}e^{2bx}+1}}}}{\frac{2}{e^{2a}e^{2bx}+1}} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 3bx \right)$$

input `int(x/atanh(tanh(a + b*x))^(1/2),x)`output `(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 3*b*x))/(3*b^2)`

**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))}(-2\operatorname{atanh}(\tanh(bx + a)) + 3bx)}{3b^2}$$

input `int(x/atanh(tanh(b*x+a))^(1/2),x)`output `(2*sqrt(atanh(tanh(a + b*x)))*(- 2*atanh(tanh(a + b*x)) + 3*b*x))/(3*b**2)`

$$3.144 \quad \int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1066
Mathematica [A] (verified)	1066
Rubi [A] (verified)	1067
Maple [A] (verified)	1068
Fricas [A] (verification not implemented)	1068
Sympy [A] (verification not implemented)	1068
Maxima [A] (verification not implemented)	1069
Giac [A] (verification not implemented)	1069
Mupad [B] (verification not implemented)	1069
Reduce [B] (verification not implemented)	1070

### Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

output `2*arctanh(tanh(b*x+a))^(1/2)/b`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

input `Integrate[1/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]]])/b`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2588

$$\frac{\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

↓ 15

$$\frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b}$$

input `Int[1/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]]])/b`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



**Maple [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b}$	15
default	$\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b}$	15

input `int(1/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2*arctanh(tanh(b*x+a))^(1/2)/b`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{bx+a}}{b}$$

input `integrate(1/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `2*sqrt(b*x + a)/b`

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \begin{cases} \frac{2\sqrt{\operatorname{atanh}(\tanh(a+bx))}}{b} & \text{for } b \neq 0 \\ \frac{x}{\sqrt{\operatorname{atanh}(\tanh(a))}} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a))**(1/2),x)`

output `Piecewise((2*sqrt(atanh(tanh(a + b*x)))/b, Ne(b, 0)), (x/sqrt(atanh(tanh(a))), True))`

### Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{bx + a}}{b}$$

input `integrate(1/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `2*sqrt(b*x + a)/b`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{bx + a}}{b}$$

input `integrate(1/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `2*sqrt(b*x + a)/b`

### Mupad [B] (verification not implemented)

Time = 3.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 3.25

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{\frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{2}}}{b}$$

input `int(1/atanh(tanh(a + b*x))^(1/2),x)`

output `(2*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/b`

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{b}$$

input `int(1/atanh(tanh(b*x+a))^(1/2),x)`

output `(2*sqrt(atanh(tanh(a + b*x))))/b`

**3.145** 
$$\int \frac{1}{x \sqrt{\mathbf{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1071
Mathematica [A] (verified)	1071
Rubi [A] (verified)	1072
Maple [A] (verified)	1073
Fricas [A] (verification not implemented)	1073
Sympy [F]	1073
Maxima [F]	1074
Giac [A] (verification not implemented)	1074
Mupad [B] (verification not implemented)	1075
Reduce [F]	1075

**Optimal result**

Integrand size = 15, antiderivative size = 49

$$\int \frac{1}{x \sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = \frac{2 \arctan\left(\frac{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \mathbf{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{bx - \mathbf{arctanh}(\tanh(a + bx))}}$$

output

$2*\arctan(\mathbf{arctanh}(\tanh(b*x+a))^{(1/2)}/(b*x-\mathbf{arctanh}(\tanh(b*x+a)))^{(1/2)})/(b*x-\mathbf{arctanh}(\tanh(b*x+a)))^{(1/2)}$

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.96

$$\int \frac{1}{x \sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = -\frac{2\mathbf{arctanh}\left(\frac{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}{\sqrt{-bx+\mathbf{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{-bx + \mathbf{arctanh}(\tanh(a + bx))}}$$

input

`Integrate[1/(x*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output  $(-2*\text{ArcTanh}[\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]]/\text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]])/\text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]]$

### Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \sqrt{\text{arctanh}(\tanh(a + bx))}} dx$$

↓ 2592

$$\frac{2 \arctan\left(\frac{\sqrt{\text{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \text{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{bx - \text{arctanh}(\tanh(a + bx))}}$$

input  $\text{Int}[1/(x*\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]]), x]$

output  $(2*\text{ArcTan}[\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]]/\text{Sqrt}[b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]])/\text{Sqrt}[b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]]$

### Defintions of rubi rules used

rule 2592  $\text{Int}[1/((u_)*\text{Sqrt}[v_]), x\_Symbol] \text{ :> With}[\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[2*(\text{ArcTan}[\text{Sqrt}[v]/\text{Rt}[(b*u - a*v)/a, 2]]/(a*\text{Rt}[(b*u - a*v)/a, 2])), x] \text{ /; NeQ}[b*u - a*v, 0] \ \&\& \ \text{PosQ}[(b*u - a*v)/a] \text{ /; PiecewiseLine arQ}[u, v, x]$

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}$	42

input `int(1/x/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \left[ \frac{\log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right)}{\sqrt{a}}, \frac{2\sqrt{-a}\arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right)}{a} \right]$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x)/sqrt(a), 2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a))/a]`

**Sympy [F]**

$$\int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{1}{x\sqrt{\operatorname{atanh}(\tanh(a+bx))}} dx$$

input `integrate(1/x/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(1/(x*sqrt(atanh(tanh(a + b*x)))), x)`

### Maxima [F]

$$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(bx + a))}} dx$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/(x*sqrt(arctanh(tanh(b*x + a)))), x)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.43

$$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `2*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a)`

**Mupad [B] (verification not implemented)**

Time = 9.16 (sec) , antiderivative size = 285, normalized size of antiderivative = 5.82

$$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{\sqrt{2} \ln \left( \frac{\sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx} \left( \frac{\sqrt{2} bx}{2} - \frac{\sqrt{2} \left( \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx \right)}{2} + \sqrt{\frac{\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right)}{2} - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)} \right)}{x} \right)}{\sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx}}$$

input `int(1/(x*atanh(tanh(a + b*x))^(1/2)),x)`output

```
(2^(1/2)*log(((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*((log((2*exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)
*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*1i - (2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/
2 + (2^(1/2)*b*x)/2)*1i)/x)*1i)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2
*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)
```

**Reduce [F]**

$$\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x} dx$$

input `int(1/x/atanh(tanh(b*x+a))^(1/2),x)`output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x),x)`



**3.146**  $\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1076
Mathematica [A] (verified)	1077
Rubi [A] (verified)	1077
Maple [A] (verified)	1079
Fricas [A] (verification not implemented)	1079
Sympy [F]	1080
Maxima [F]	1080
Giac [A] (verification not implemented)	1080
Mupad [B] (verification not implemented)	1081
Reduce [F]	1081

**Optimal result**

Integrand size = 15, antiderivative size = 94

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{b \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

$$+ \frac{b}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

output

```
b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x
-arctanh(tanh(b*x+a)))^(3/2)-1/x/arctanh(tanh(b*x+a))^(1/2)+b/(b*x-arctanh
(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.83

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{\operatorname{barctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x(-bx + \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Integrate[1/(x^2*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `(b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2) - Sqrt[ArcTanh[Tanh[a + b*x]]]/(x*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow \text{2599}$$

$$-\frac{1}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

$$\downarrow \text{2594}$$

$$\begin{aligned}
& -\frac{1}{2}b \left( -\frac{\int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
& \qquad \qquad \qquad \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
& \qquad \qquad \qquad \downarrow \text{2592} \\
& -\frac{1}{2}b \left( -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
& \qquad \qquad \qquad \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}
\end{aligned}$$

input `Int[1/(x^2*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `-1/2*(b*((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])) - 1/(x*Sqrt[ArcTanh[Tanh[a + b*x]]]))`

### Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.01

method	result	size
default	$2b \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)bx} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}} \right)$	95

input

```
int(1/x^2/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2*b*(2*arctanh(tanh(b*x+a))^(1/2)/(-4*arctanh(tanh(b*x+a))+4*b*x)/b/x-2/(-4*arctanh(tanh(b*x+a))+4*b*x)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.96

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \left[ \frac{\sqrt{abx} \log\left(\frac{bx + 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2\sqrt{bx+aa}}{2a^2x}, -\frac{\sqrt{-abx} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + \sqrt{bx+aa}}{a^2x} \right]$$

input

```
integrate(1/x^2/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
[1/2*(sqrt(a)*b*x*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*sqrt(b*x + a)*a)/(a^2*x), -(sqrt(-a)*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) + sqrt(b*x + a)*a)/(a^2*x)]
```

**Sympy [F]**

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^2 \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input

```
integrate(1/x**2/atanh(tanh(b*x+a))**(1/2), x)
```

output

```
Integral(1/(x**2*sqrt(atanh(tanh(a + b*x)))), x)
```

**Maxima [F]**

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^2 \sqrt{\operatorname{artanh}(\tanh(bx + a))}} dx$$

input

```
integrate(1/x^2/arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")
```

output

```
integrate(1/(x^2*sqrt(arctanh(tanh(b*x + a)))), x)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.47

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -b \left( \frac{\arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{\sqrt{bx+a}}{abx} \right)$$

input

```
integrate(1/x^2/arctanh(tanh(b*x+a))^(1/2), x, algorithm="giac")
```

output  $-b*(\arctan(\sqrt{b*x + a}/\sqrt{-a})/(\sqrt{-a}*a) + \sqrt{b*x + a}/(a*b*x))$

### Mupad [B] (verification not implemented)

Time = 8.71 (sec) , antiderivative size = 570, normalized size of antiderivative = 6.06

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \text{Too large to display}$$

input  $\text{int}(1/(x^2*\operatorname{atanh}(\tanh(a + b*x))^(1/2)),x)$

output  $(2*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}/(x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)) + (2^{(1/2)*b} * \log(((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)} * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)} * 2i - 2^{(1/2)} * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x) + 2^{(1/2)*b*x} * ((2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)) * i) / (2*x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)}) * i) / (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(3/2)}$

### Reduce [F]

$$\int \frac{1}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^2} dx$$

input  $\text{int}(1/x^2/\operatorname{atanh}(\tanh(b*x+a))^(1/2),x)$

output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x**2),x)`

$$3.147 \quad \int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result . . . . .	1083
Mathematica [A] (verified) . . . . .	1084
Rubi [A] (verified) . . . . .	1084
Maple [A] (verified) . . . . .	1086
Fricas [A] (verification not implemented) . . . . .	1087
Sympy [F] . . . . .	1087
Maxima [F] . . . . .	1088
Giac [A] (verification not implemented) . . . . .	1088
Mupad [B] (verification not implemented) . . . . .	1088
Reduce [F] . . . . .	1089

**Optimal result**

Integrand size = 15, antiderivative size = 158

$$\begin{aligned} & \int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\ &= \frac{3b^2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx-\operatorname{arctanh}(\tanh(a+bx)))^{5/2}} + \frac{b}{4x\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\ & \quad - \frac{1}{4(bx-\operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\ & \quad - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\ & \quad + \frac{3b^2}{4(bx-\operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \end{aligned}$$

output

```

3/4*b^2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^(5/2)+1/4*b/x/arctanh(tanh(b*x+a))^(3/2)-1/4*
b^2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)-1/2/x^2/arctanh(
tanh(b*x+a))^(1/2)+3/4*b^2/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a
))^(1/2)
    
```



**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.62

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \frac{1}{4} \left( -\frac{3b^2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} + \frac{(5bx - 2\operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^2(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2} \right)$$

input `Integrate[1/(x^3*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `((-3*b^2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2) + ((5*b*x - 2*ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]/(x^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2))/4`

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow \text{2599}$$

$$-\frac{1}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx - \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

$$\begin{aligned}
 & \downarrow 2599 \\
 & -\frac{1}{4}b \left( -\frac{3}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2594 \\
 & -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2594 \\
 & -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2592 \\
 & -\frac{1}{4}b \left( -\frac{3}{2}b \left( -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}
 \end{aligned}$$

input

```
Int [1/(x^3*sqrt [ArcTanh [Tanh [a + b*x]]] ), x]
```

output

```
-1/4*(b*((-3*b*(-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(3/2))) - 1/(2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]])
```

### Defintions of rubi rules used

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.94

method	result
default	$2b^2 \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(-4 \operatorname{arctanh}(\tanh(bx+a))+4bx)b^2x^2} + \frac{6 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(-4 \operatorname{arctanh}(\tanh(bx+a))+4bx)bx} - \frac{6 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{(-4 \operatorname{arctanh}(\tanh(bx+a))+4bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}} \right)$

input `int(1/x^3/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2*b^2*(arctanh(tanh(b*x+a))^(1/2)/(-4*arctanh(tanh(b*x+a))+4*b*x)/b^2/x^2+3/(-4*arctanh(tanh(b*x+a))+4*b*x)*(2*arctanh(tanh(b*x+a))^(1/2)/(-4*arctanh(tanh(b*x+a))+4*b*x)/b/x-2/(-4*arctanh(tanh(b*x+a))+4*b*x)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \left[ \frac{3\sqrt{ab^2x^2} \log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(3abx - 2a^2)\sqrt{bx+a}}{8a^3x^2}, \frac{3\sqrt{-ab^2x^2} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (3abx - 2a^2)}{4a^3x^2} \right]$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[1/8*(3*sqrt(a)*b^2*x^2*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(3*a*b*x - 2*a^2)*sqrt(b*x + a))/(a^3*x^2), 1/4*(3*sqrt(-a)*b^2*x^2*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b*x - 2*a^2)*sqrt(b*x + a))/(a^3*x^2)]`

### Sympy [F]

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^3 \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(1/x**3/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(1/(x**3*sqrt(atanh(tanh(a + b*x)))), x)`

**Maxima [F]**

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(bx + a))}} dx$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/(x^3*sqrt(arctanh(tanh(b*x + a)))) , x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{3b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} + \frac{3(bx+a)^{\frac{3}{2}}b^3 - 5\sqrt{bx+aa}b^3}{4b}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `1/4*(3*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) + (3*(b*x + a)^(3/2)*b^3 - 5*sqrt(b*x + a)*a*b^3)/(a^2*b^2*x^2))/b`

**Mupad [B] (verification not implemented)**

Time = 7.99 (sec) , antiderivative size = 802, normalized size of antiderivative = 5.08

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \text{Too large to display}$$

input `int(1/(x^3*atanh(tanh(a + b*x))^(1/2)),x)`

output

```
(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(x^2*(2*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x) + (3*b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (2^(1/2)*b^2*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5 + 40*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 80*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 32*a^5 - 10*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 + 80*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*1i)/(x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x...
```

**Reduce [F]**

$$\int \frac{1}{x^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^3} dx$$

input

```
int(1/x^3/atanh(tanh(b*x+a))^(1/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x**3),x)
```

**3.148** 
$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1090
Mathematica [A] (verified)	1091
Rubi [A] (verified)	1091
Maple [A] (verified)	1094
Fricas [A] (verification not implemented)	1095
Sympy [F]	1095
Maxima [F]	1096
Giac [A] (verification not implemented)	1096
Mupad [B] (verification not implemented)	1097
Reduce [F]	1097

**Optimal result**

Integrand size = 15, antiderivative size = 212

$$\begin{aligned} & \int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\ &= \frac{5b^3 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} - \frac{b^2}{8x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\ & \quad + \frac{b}{8(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\ & \quad + \frac{12x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5b^3} \\ & \quad - \frac{24(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{1} \\ & \quad - \frac{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5b^3} \\ & \quad + \frac{8(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5b^3} \end{aligned}$$

output

$$\frac{5/8*b^3*\arctan(\operatorname{arctanh}(\tanh(b*x+a))^{1/2}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{1/2})}{(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{7/2}-1/8*b^2/x/\operatorname{arctanh}(\tanh(b*x+a))^{5/2}+1/8*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/\operatorname{arctanh}(\tanh(b*x+a))^{5/2}+1/12*b/x^2/\operatorname{arctanh}(\tanh(b*x+a))^{3/2}-5/24*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2/\operatorname{arctanh}(\tanh(b*x+a))^{3/2}-1/3/x^3/\operatorname{arctanh}(\tanh(b*x+a))^{1/2}+5/8*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^3/\operatorname{arctanh}(\tanh(b*x+a))^{1/2}}$$
**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.55

$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{5b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} + \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}(33b^2x^2 - 26bx \operatorname{arctanh}(\tanh(a+bx)) + 8 \operatorname{arctanh}(\tanh(a+bx))^2)}{24x^3(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input

`Integrate[1/(x^4*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output

$$\frac{(5*b^3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(8*(-(b*x) + ArcTanh[Tanh[a + b*x]])^{7/2}) + (Sqrt[ArcTanh[Tanh[a + b*x]]]*(33*b^2*x^2 - 26*b*x*ArcTanh[Tanh[a + b*x]] + 8*ArcTanh[Tanh[a + b*x]]^2))/(24*x^3*(b*x - ArcTanh[Tanh[a + b*x]])^3)}$$
**Rubi [A] (verified)**Time = 0.59 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.15, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {2599, 2599, 2599, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$



$$\begin{aligned}
 & \downarrow 2599 \\
 & -\frac{1}{6}b \int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx - \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2599 \\
 & -\frac{1}{6}b \left( -\frac{3}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \\
 & \quad \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2599 \\
 & -\frac{1}{6}b \left( -\frac{3}{4}b \left( -\frac{5}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \\
 & \quad \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2594 \\
 & -\frac{1}{6}b \left( -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) - \\
 & \quad \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2594 \\
 & -\frac{1}{6}b \left( -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) \right) \right) - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \frac{1}{3x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \downarrow 2594
 \end{aligned}$$

$$-\frac{1}{6}b \left( -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \right) \right) \frac{1}{3x^3\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2592

$$-\frac{1}{6}b \left( -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{2 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \right) \right) \frac{1}{3x^3\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Int[1/(x^4*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `-1/6*(b*((-3*b*((-5*b*(-((-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) - 2/(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2))))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(5/2))))/4 - 1/(2*x^2*ArcTanh[Tanh[a + b*x]]^(3/2)))) - 1/(3*x^3*Sqrt[ArcTanh[Tanh[a + b*x]]])`

## Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*m + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.94

method	result
default	$2b^3 \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)b^3x^3} + \frac{10\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)b^2x^2} + \frac{10 \left( \frac{6\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)bx} - \frac{1}{(-4\operatorname{arctanh}(\tanh(bx+a))+4bx)} \right)}{-4\operatorname{arctanh}(\tanh(bx+a))+4bx} \right)$

input `int(1/x^4/arctanh(tanh(b*x+a))^(1/2), x, method=_RETURNVERBOSE)`

output

```
2*b^3*(2/3*arctanh(tanh(b*x+a))^(1/2)/(-4*arctanh(tanh(b*x+a))+4*b*x)/b^3/
x^3+10/3/(-4*arctanh(tanh(b*x+a))+4*b*x)*(arctanh(tanh(b*x+a))^(1/2)/(-4*
arctanh(tanh(b*x+a))+4*b*x)/b^2/x^2+3/(-4*arctanh(tanh(b*x+a))+4*b*x)*(2*ar
ctanh(tanh(b*x+a))^(1/2)/(-4*arctanh(tanh(b*x+a))+4*b*x)/b/x-2/(-4*arctanh
(tanh(b*x+a))+4*b*x)/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh
(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.67

$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$= \left[ \frac{15 \sqrt{ab^3} x^3 \log\left(\frac{bx + 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(15ab^2x^2 - 10a^2bx + 8a^3)\sqrt{bx+a}}{48a^4x^3}, \right.$$

$$\left. - \frac{15\sqrt{-ab^3}x^3 \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (15ab^2x^2 - 10a^2bx + 8a^3)\sqrt{bx+a}}{24a^4x^3} \right]$$

input

```
integrate(1/x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
[1/48*(15*sqrt(a)*b^3*x^3*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2
*(15*a*b^2*x^2 - 10*a^2*b*x + 8*a^3)*sqrt(b*x + a))/(a^4*x^3), -1/24*(15*s
qrt(-a)*b^3*x^3*arctan(sqrt(-a)/sqrt(b*x + a)) + (15*a*b^2*x^2 - 10*a^2*b*
x + 8*a^3)*sqrt(b*x + a))/(a^4*x^3)]
```

**Sympy [F]**

$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^4 \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input

```
integrate(1/x**4/atanh(tanh(b*x+a))**(1/2),x)
```

output `Integral(1/(x**4*sqrt(atanh(tanh(a + b*x)))), x)`

### Maxima [F]

$$\int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(bx + a))}} dx$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/(x^4*sqrt(arctanh(tanh(b*x + a)))), x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.34

$$\begin{aligned} & \int \frac{1}{x^4 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \\ &= -\frac{1}{24} b^3 \left( \frac{15 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a} a^3} + \frac{15 (bx+a)^{\frac{5}{2}} - 40 (bx+a)^{\frac{3}{2}} a + 33 \sqrt{bx+aa^2}}{a^3 b^3 x^3} \right) \end{aligned}$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `-1/24*b^3*(15*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^3) + (15*(b*x + a)^(5/2) - 40*(b*x + a)^(3/2)*a + 33*sqrt(b*x + a)*a^2)/(a^3*b^3*x^3)`



output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))*x**4),x)`

**3.149**       $\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1099
Mathematica [A] (verified)	1099
Rubi [A] (verified)	1100
Maple [B] (verified)	1102
Fricas [A] (verification not implemented)	1103
Sympy [F]	1103
Maxima [A] (verification not implemented)	1104
Giac [A] (verification not implemented)	1104
Mupad [B] (verification not implemented)	1105
Reduce [F]	1105

**Optimal result**

Integrand size = 15, antiderivative size = 95

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2x^4}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{16x^3\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2} - \frac{32x^2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b^3} + \frac{128x\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b^4} - \frac{256\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^5}$$

output

```
-2*x^4/b/arctanh(tanh(b*x+a))^(1/2)+16*x^3*arctanh(tanh(b*x+a))^(1/2)/b^2-
32*x^2*arctanh(tanh(b*x+a))^(3/2)/b^3+128/5*x*arctanh(tanh(b*x+a))^(5/2)/b
^4-256/35*arctanh(tanh(b*x+a))^(7/2)/b^5
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.87

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2(35b^4x^4 - 280b^3x^3\operatorname{arctanh}(\tanh(a+bx)) + 560b^2x^2\operatorname{arctanh}(\tanh(a+bx))^2 - 448bx\operatorname{arctanh}(\tanh(a+bx)) - 256\operatorname{arctanh}(\tanh(a+bx))^3)}{35b^5\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$



input `Integrate[x^4/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(-2*(35*b^4*x^4 - 280*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 560*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 448*b*x*ArcTanh[Tanh[a + b*x]]^3 + 128*ArcTanh[Tanh[a + b*x]]^4)/(35*b^5*Sqrt[ArcTanh[Tanh[a + b*x]]])`

### Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.23, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx \\
 & \quad \downarrow 2599 \\
 & \frac{8 \int \frac{x^3}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^4}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow 2599 \\
 & \frac{8 \left( \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{6 \int x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} \right)}{b} - \frac{2x^4}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow 2599 \\
 & \frac{8 \left( \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \int x \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx}{3b} \right)}{b} \right)}{b} - \frac{2x^4}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow 2599 \\
 & \frac{2x^4}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}
 \end{aligned}$$

$$8 \left( \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx}{5b} \right)}{3b} \right)}{b} \right)$$

$$\frac{2x^4}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2588

$$8 \left( \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{5/2} dx}{5b^2} \right)}{3b} \right)}{b} \right)$$

$$\frac{2x^4}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 15

$$8 \left( \frac{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{6 \left( \frac{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{35b^2} \right)}{3b} \right)}{b} \right)$$

$$\frac{2x^4}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Int [x^4/ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output `(-2*x^4)/(b*sqrt[ArcTanh[Tanh[a + b*x]]]) + (8*((2*x^3*sqrt[ArcTanh[Tanh[a + b*x]]])/b - (6*((2*x^2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*b) - (4*ArcTanh[Tanh[a + b*x]]^(7/2))/(35*b^2)))/(3*b)))/b)`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :=> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :=> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :=> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 318 vs.  $2(81) = 162$ .

Time = 0.20 (sec) , antiderivative size = 319, normalized size of antiderivative = 3.36

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a)) \frac{7}{2} - 8 \operatorname{arctanh}(\tanh(bx+a)) \frac{5}{2} a - 8 \operatorname{arctanh}(\tanh(bx+a)) \frac{5}{2} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 4 \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} a^2 + \dots}{\dots}$

input `int(x^4/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output

```
2/b^5*(1/7*arctanh(tanh(b*x+a))^(7/2)-4/5*arctanh(tanh(b*x+a))^(5/2)*a-4/5
*arctanh(tanh(b*x+a))^(5/2)*(arctanh(tanh(b*x+a))-b*x-a)+2*arctanh(tanh(b*
x+a))^(3/2)*a^2+4*arctanh(tanh(b*x+a))^(3/2)*a*(arctanh(tanh(b*x+a))-b*x-a
)+2*arctanh(tanh(b*x+a))^(3/2)*(arctanh(tanh(b*x+a))-b*x-a)^2-4*a^3*arctan
h(tanh(b*x+a))^(1/2)-12*a^2*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+
a))^(1/2)-12*a*(arctanh(tanh(b*x+a))-b*x-a)^2*arctanh(tanh(b*x+a))^(1/2)-4
*(arctanh(tanh(b*x+a))-b*x-a)^3*arctanh(tanh(b*x+a))^(1/2)-(a^4+4*a^3*(arc
tanh(tanh(b*x+a))-b*x-a)+6*a^2*(arctanh(tanh(b*x+a))-b*x-a)^2+4*a*(arctanh
(tanh(b*x+a))-b*x-a)^3+(arctanh(tanh(b*x+a))-b*x-a)^4)/arctanh(tanh(b*x+a)
)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.66

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(5b^4x^4 - 8ab^3x^3 + 16a^2b^2x^2 - 64a^3bx - 128a^4)\sqrt{bx + a}}{35(b^6x + ab^5)}$$

input

```
integrate(x^4/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
2/35*(5*b^4*x^4 - 8*a*b^3*x^3 + 16*a^2*b^2*x^2 - 64*a^3*b*x - 128*a^4)*sqr
t(b*x + a)/(b^6*x + a*b^5)
```

**Sympy [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^4}{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input

```
integrate(x**4/atanh(tanh(b*x+a))**(3/2),x)
```

output

```
Integral(x**4/atanh(tanh(a + b*x))**(3/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.67

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(5b^5x^5 - 3ab^4x^4 + 8a^2b^3x^3 - 48a^3b^2x^2 - 192a^4bx - 128a^5)}{35(bx + a)^{\frac{3}{2}}b^5}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `2/35*(5*b^5*x^5 - 3*a*b^4*x^4 + 8*a^2*b^3*x^3 - 48*a^3*b^2*x^2 - 192*a^4*b*x - 128*a^5)/((b*x + a)^(3/2)*b^5)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.86

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2 \left( \frac{35a^4}{\sqrt{bx+ab}} - \frac{5(bx+a)^{\frac{7}{2}}b^6 - 28(bx+a)^{\frac{5}{2}}ab^6 + 70(bx+a)^{\frac{3}{2}}a^2b^6 - 140\sqrt{bx+aa^3b^6}}{b^7} \right)}{35b^4}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `-2/35*(35*a^4/(sqrt(b*x + a)*b) - (5*(b*x + a)^(7/2)*b^6 - 28*(b*x + a)^(5/2)*a*b^6 + 70*(b*x + a)^(3/2)*a^2*b^6 - 140*sqrt(b*x + a)*a^3*b^6)/b^4`



output `int((sqrt(atanh(tanh(a + b*x)))*x**4)/atanh(tanh(a + b*x))**2,x)`

### 3.150 $\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1107
Mathematica [A] (verified)	1107
Rubi [A] (verified)	1108
Maple [B] (verified)	1110
Fricas [A] (verification not implemented)	1110
Sympy [F]	1111
Maxima [A] (verification not implemented)	1111
Giac [A] (verification not implemented)	1111
Mupad [B] (verification not implemented)	1112
Reduce [F]	1113

#### Optimal result

Integrand size = 15, antiderivative size = 74

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2x^3}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{12x^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2} - \frac{16x\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{b^3} + \frac{32\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5b^4}$$

output

$$-2*x^3/b/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}+12*x^2*\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}/b^2-16*x*\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}/b^3+32/5*\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}/b^4$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.89

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2(-5b^3x^3 + 30b^2x^2\operatorname{arctanh}(\tanh(a+bx)) - 40bx\operatorname{arctanh}(\tanh(a+bx)))}{5b^4\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input

```
Integrate[x^3/ArcTanh[Tanh[a + b*x]]^(3/2), x]
```



output

$(2*(-5*b^3*x^3 + 30*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 40*b*x*ArcTanh[Tanh[a + b*x]])^2 + 16*ArcTanh[Tanh[a + b*x]]^3)/(5*b^4*sqrt[ArcTanh[Tanh[a + b*x]]])$

### Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

↓ 2599

$$\frac{6 \int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{b} - \frac{2x^3}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

↓ 2599

$$\frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{4 \int x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} \right)}{b} - \frac{2x^3}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

↓ 2599

$$\frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx}{3b} \right)}{b} \right)}{b} - \frac{2x^3}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

↓ 2588

$$\frac{b}{2x^3} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}$$

$$\frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{3/2} d \operatorname{arctanh}(\tanh(a+bx))}{3b^2} \right)}{b} \right)}{2x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 15

$$\frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{15b^2} \right)}{b} \right)}{\frac{b}{2x^3} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Int [x^3/ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output `(-2*x^3)/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (6*((2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]])/b - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (4*ArcTanh[Tanh[a + b*x]]^(5/2))/(15*b^2))/b))/b`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs.  $2(64) = 128$ .

Time = 0.20 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.72

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{5/2}}{5} - 2 \operatorname{arctanh}(\tanh(bx+a))^{3/2} a - 2 \operatorname{arctanh}(\tanh(bx+a))^{3/2} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 6 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}$

input

```
int(x^3/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

output

$$\frac{2}{b^4} \left( \frac{1}{5} \operatorname{arctanh}(\tanh(bx+a))^{5/2} - \operatorname{arctanh}(\tanh(bx+a))^{3/2} a - \operatorname{arctanh}(\tanh(bx+a))^{3/2} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3 \operatorname{arctanh}(\tanh(bx+a))^{1/2} a^2 + 6 a \operatorname{arctanh}(\tanh(bx+a))^{1/2} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 3 (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 \operatorname{arctanh}(\tanh(bx+a))^{1/2} - (-a^3 - 3 a^2 (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) - 3 a (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 - (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^3) / \operatorname{arctanh}(\tanh(bx+a))^{1/2} \right)$$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.69

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(b^3 x^3 - 2ab^2 x^2 + 8a^2 bx + 16a^3) \sqrt{bx + a}}{5(b^5 x + ab^4)}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output  $2/5*(b^3*x^3 - 2*a*b^2*x^2 + 8*a^2*b*x + 16*a^3)*\text{sqrt}(b*x + a)/(b^5*x + a*b^4)$

### Sympy [F]

$$\int \frac{x^3}{\text{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^3}{\text{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(x**3/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(x**3/atanh(tanh(a + b*x))**(3/2), x)`

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.70

$$\int \frac{x^3}{\text{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(b^4x^4 - ab^3x^3 + 6a^2b^2x^2 + 24a^3bx + 16a^4)}{5(bx + a)^{\frac{3}{2}}b^4}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output  $2/5*(b^4*x^4 - a*b^3*x^3 + 6*a^2*b^2*x^2 + 24*a^3*b*x + 16*a^4)/((b*x + a)^{(3/2)}*b^4)$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.82

$$\int \frac{x^3}{\text{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2a^3}{\sqrt{bx + ab^4}} + \frac{2\left((bx + a)^{\frac{5}{2}}b^{16} - 5(bx + a)^{\frac{3}{2}}ab^{16} + 15\sqrt{bx + a}a^2b^{16}\right)}{5b^{20}}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output  $2a^3/(\sqrt{bx+a}b^4) + 2/5*((bx+a)^{5/2}b^{16} - 5*(bx+a)^{3/2}ab^{16} + 15*\sqrt{bx+a}a^2b^{16})/b^{20}$

### Mupad [B] (verification not implemented)

Time = 3.37 (sec) , antiderivative size = 660, normalized size of antiderivative = 8.92

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \text{Too large to display}$$

input `int(x^3/atanh(tanh(a + b*x))^(3/2),x)`

output 
$$\begin{aligned} & ((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)} * ((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + 2*b*x)^2 / (2*b^3) + (2 * ((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + 2*b*x)/b^2 + (8 * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x) / (5*b^2)) * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x) / (3*b)) / b + (2*x^2 * (\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)} / (5*b^2) + (x * ((\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + 2*b*x) / b^2 + (8 * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + b*x) / (5*b^2)) * (\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)} / (3*b) - ((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/2)^{(1/2)} * (\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 + 2*b*x)^3 / (2*b^4 * (\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))) \end{aligned}$$

**Reduce [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^3/atanh(tanh(b*x+a))^(3/2),x)`

output `int((sqrt(atanh(tanh(a + b*x)))*x**3)/atanh(tanh(a + b*x))**2,x)`

### 3.151 $\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1114
Mathematica [A] (verified)	1114
Rubi [A] (verified)	1115
Maple [B] (verified)	1116
Fricas [A] (verification not implemented)	1117
Sympy [F]	1117
Maxima [A] (verification not implemented)	1118
Giac [A] (verification not implemented)	1118
Mupad [B] (verification not implemented)	1118
Reduce [F]	1119

#### Optimal result

Integrand size = 15, antiderivative size = 55

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2x^2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{8x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2} - \frac{16\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^3}$$

output

$-2*x^2/b/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}+8*x*\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}/b^2-16/3*\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}/b^3$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2(3b^2x^2 - 12bx\operatorname{arctanh}(\tanh(a+bx)) + 8\operatorname{arctanh}(\tanh(a+bx))^2)}{3b^3\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input

`Integrate[x^2/ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output

```
(-2*(3*b^2*x^2 - 12*b*x*ArcTanh[Tanh[a + b*x]] + 8*ArcTanh[Tanh[a + b*x]]^2)/(3*b^3*Sqrt[ArcTanh[Tanh[a + b*x]]])
```

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{4 \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow \text{2599} \\
 & \frac{4 \left( \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} \right)}{b} - \frac{2x^2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow \text{2588} \\
 & \frac{4 \left( \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} d\operatorname{arctanh}(\tanh(a+bx))}{b^2} \right)}{b} - \frac{2x^2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow \text{15} \\
 & \frac{4 \left( \frac{2x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^2} \right)}{b} - \frac{2x^2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}
 \end{aligned}$$



input `Int[x^2/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(-2*x^2)/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (4*((2*x*Sqrt[ArcTanh[Tanh[a + b*x]]])/b - (4*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b^2)))/b`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs.  $2(47) = 94$ .

Time = 0.20 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.93

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} - 4 \sqrt{\operatorname{arctanh}(\tanh(bx+a))} a - 4(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) \sqrt{\operatorname{arctanh}(\tanh(bx+a))} - \frac{2(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)))}{b^3}}$

input `int(x^2/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output

```
2/b^3*(1/3*arctanh(tanh(b*x+a))^(3/2)-2*arctanh(tanh(b*x+a))^(1/2)*a-2*(ar
ctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2)-(a^2+2*a*(arctanh(tan
h(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/arctanh(tanh(b*x+a))^(1/2
))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(b^2x^2 - 4abx - 8a^2)\sqrt{bx + a}}{3(b^4x + ab^3)}$$

input

```
integrate(x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
2/3*(b^2*x^2 - 4*a*b*x - 8*a^2)*sqrt(b*x + a)/(b^4*x + a*b^3)
```

**Sympy [F]**

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^2}{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input

```
integrate(x**2/atanh(tanh(b*x+a))**(3/2),x)
```

output

```
Integral(x**2/atanh(tanh(a + b*x))**(3/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.75

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(b^3 x^3 - 3ab^2 x^2 - 12a^2 bx - 8a^3)}{3(bx + a)^{3/2} b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`output `2/3*(b^3*x^3 - 3*a*b^2*x^2 - 12*a^2*b*x - 8*a^3)/((b*x + a)^(3/2)*b^3)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2\left(\frac{3a^2}{\sqrt{bx+ab}} - \frac{(bx+a)^{3/2} b^2 - 6\sqrt{bx+ab} b^2}{b^3}\right)}{3b^2}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`output `-2/3*(3*a^2/(sqrt(b*x + a)*b) - ((b*x + a)^(3/2)*b^2 - 6*sqrt(b*x + a)*a*b^2)/b^3)/b^2`**Mupad [B] (verification not implemented)**

Time = 3.43 (sec) , antiderivative size = 259, normalized size of antiderivative = 4.71

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{4\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)}{2}} \left(3b^2 x^2 - 6bx \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 6bx \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) + 2\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right)\right)}{3b^3 \left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)\right)}$$

input `int(x^2/atanh(tanh(a + b*x))^(3/2),x)`

output `-(4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2)^(1/2)*(2*log(2/(exp(2*a)*exp(2*b*x) + 1))^2 - 4*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2 + 3*b^2*x^2 - 6*b*x*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 6*b*x*log(2/(exp(2*a)*exp(2*b*x) + 1))))/(3*b^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))`

**Reduce [F]**

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^2/atanh(tanh(b*x+a))^(3/2),x)`

output `int((sqrt(atanh(tanh(a + b*x)))*x**2)/atanh(tanh(a + b*x))**2,x)`

### 3.152 $\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1120
Mathematica [A] (verified)	1120
Rubi [A] (verified)	1121
Maple [A] (verified)	1122
Fricas [A] (verification not implemented)	1123
Sympy [A] (verification not implemented)	1123
Maxima [A] (verification not implemented)	1123
Giac [A] (verification not implemented)	1124
Mupad [B] (verification not implemented)	1124
Reduce [B] (verification not implemented)	1125

#### Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2x}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{4\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2}$$

output `-2*x/b/arctanh(tanh(b*x+a))^(1/2)+4*arctanh(tanh(b*x+a))^(1/2)/b^2`

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{-2bx + 4\operatorname{arctanh}(\tanh(a+bx))}{b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Integrate[x/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(-2*b*x + 4*ArcTanh[Tanh[a + b*x]])/(b^2*Sqrt[ArcTanh[Tanh[a + b*x]]])`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$$

$$\downarrow 2599$$

$$\frac{2 \int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$\downarrow 2588$$

$$\frac{2 \int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} d\operatorname{arctanh}(\tanh(a+bx))}{b^2} - \frac{2x}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$\downarrow 15$$

$$\frac{4\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2} - \frac{2x}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Int[x/ArcTanh[Tanh[a + b*x]]^(3/2), x]`

output `(-2*x)/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (4*Sqrt[ArcTanh[Tanh[a + b*x]]])/b^2`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.18

method	result	size
default	$\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))} - \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a)))}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}}{b^2}$	40

input `int(x/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2/b^2*(arctanh(tanh(b*x+a))^(1/2)-(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(bx + 2a)\sqrt{bx + a}}{b^3x + ab^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`output `2*(b*x + 2*a)*sqrt(b*x + a)/(b^3*x + a*b^2)`**Sympy [A] (verification not implemented)**

Time = 0.55 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.35

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \begin{cases} -\frac{2x}{b\sqrt{\operatorname{atanh}(\tanh(a + bx))}} + \frac{4\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{b^2} & \text{for } b \neq 0 \\ \frac{x^2}{2\operatorname{atanh}^{\frac{3}{2}}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(x/atanh(tanh(b*x+a))**(3/2),x)`output `Piecewise((-2*x/(b*sqrt(atanh(tanh(a + b*x)))) + 4*sqrt(atanh(tanh(a + b*x)))/b**2, Ne(b, 0)), (x**2/(2*atanh(tanh(a))**(3/2)), True))`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(b^2x^2 + 3abx + 2a^2)}{(bx + a)^{\frac{3}{2}}b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`output `2*(b^2*x^2 + 3*a*b*x + 2*a^2)/((b*x + a)^(3/2)*b^2)`



**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2 \left( \frac{\sqrt{bx+a}}{b} + \frac{a}{\sqrt{bx+ab}} \right)}{b}$$

input `integrate(x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`output `2*(sqrt(b*x + a)/b + a/(sqrt(b*x + a)*b))/b`**Mupad [B] (verification not implemented)**

Time = 3.47 (sec) , antiderivative size = 152, normalized size of antiderivative = 4.47

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{4 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + bx \right)}{b^2 \left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) \right)}$$

input `int(x/atanh(tanh(a + b*x))^(3/2),x)`output `-(4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + b*x))/(b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))} (2\operatorname{atanh}(\tanh(bx + a)) - bx)}{\operatorname{atanh}(\tanh(bx + a)) b^2}$$

input `int(x/atanh(tanh(b*x+a))^(3/2),x)`

output `(2*sqrt(atanh(tanh(a + b*x)))*(2*atanh(tanh(a + b*x)) - b*x))/(atanh(tanh(a + b*x))*b**2)`

### 3.153 $\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1126
Mathematica [A] (verified)	1126
Rubi [A] (verified)	1127
Maple [A] (verified)	1128
Fricas [A] (verification not implemented)	1128
Sympy [A] (verification not implemented)	1128
Maxima [A] (verification not implemented)	1129
Giac [A] (verification not implemented)	1129
Mupad [B] (verification not implemented)	1129
Reduce [B] (verification not implemented)	1130

#### Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output `-2/b/arctanh(tanh(b*x+a))^(1/2)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(-3/2), x]`

output `-2/(b*Sqrt[ArcTanh[Tanh[a + b*x]]])`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

$$\downarrow \text{2588}$$

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} d\operatorname{arctanh}(\tanh(a + bx))$$

$$\downarrow \text{15}$$

$$-\frac{2}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(-3/2), x]`

output `-2/(b*Sqrt[ArcTanh[Tanh[a + b*x]]])`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$-\frac{2}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	15
default	$-\frac{2}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	15

input `int(1/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/b/arctanh(tanh(b*x+a))^(1/2)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2\sqrt{bx+a}}{b^2x+ab}$$

input `integrate(1/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `-2*sqrt(b*x + a)/(b^2*x + a*b)`

**Sympy [A] (verification not implemented)**

Time = 0.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \begin{cases} -\frac{2}{b\sqrt{\operatorname{atanh}(\tanh(a+bx))}} & \text{for } b \neq 0 \\ \frac{x}{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a))**(3/2),x)`

output `Piecewise((-2/(b*sqrt(atanh(tanh(a + b*x))))), Ne(b, 0)), (x/atanh(tanh(a))  
**(3/2), True))`

### Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2}{\sqrt{bx + ab}}$$

input `integrate(1/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `-2/(sqrt(b*x + a)*b)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2}{\sqrt{bx + ab}}$$

input `integrate(1/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `-2/(sqrt(b*x + a)*b)`

### Mupad [B] (verification not implemented)

Time = 3.44 (sec) , antiderivative size = 97, normalized size of antiderivative = 6.06

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{4 \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{b \left( \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) \right)}$$

input `int(1/atanh(tanh(a + b*x))^(3/2),x)`

output `(4*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(b*(log(1/(exp(2*a)*exp(2*b*x) + 1)) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))))`

### Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))b}$$

input `int(1/atanh(tanh(b*x+a))^(3/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*b)`

### 3.154 $\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1131
Mathematica [A] (verified)	1131
Rubi [A] (verified)	1132
Maple [A] (verified)	1133
Fricas [A] (verification not implemented)	1133
Sympy [F]	1134
Maxima [F]	1134
Giac [A] (verification not implemented)	1134
Mupad [B] (verification not implemented)	1135
Reduce [F]	1136

#### Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(3/2)-2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} + \frac{2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}$$



input `Integrate[1/(x*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]]^(3/2) + 2/(Sqrt[ArcTanh[Tanh[a + b*x]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

### Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

↓ 2594

$$-\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

↓ 2592

$$\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

input `Int[1/(x*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])`

### Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

### Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^{\frac{3}{2}}}$	68

input `int(1/x/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-2/(arctanh(tanh(b*x+a))-b*x)^(3/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.37

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \left[ \frac{(bx + a)\sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) + 2\sqrt{bx + aa}}{a^2bx + a^3}, \frac{2((bx + a)\sqrt{-a} \operatorname{arctanh}(\sqrt{-a} \operatorname{arctanh}(\tanh(a + bx))))}{a^2bx + a^3} \right]$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[((b*x + a)*sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*sqrt(b*x + a)*a)/(a^2*b*x + a^3), 2*((b*x + a)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + sqrt(b*x + a)*a)/(a^2*b*x + a^3)]`

### Sympy [F]

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(1/(x*atanh(tanh(a + b*x))**(3/2)), x)`

### Maxima [F]

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x \operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(1/(x*arctanh(tanh(b*x + a))^(3/2)), x)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.47

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{2}{\sqrt{bx+aa}}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output  $2*\arctan(\sqrt{b*x + a}/\sqrt{-a})/(\sqrt{-a}*a) + 2/(\sqrt{b*x + a}*a)$

### Mupad [B] (verification not implemented)

Time = 8.32 (sec) , antiderivative size = 614, normalized size of antiderivative = 7.87

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Too large to display}$$

input  $\text{int}(1/(x*\operatorname{atanh}(\tanh(a + b*x))^{(3/2)}),x)$

output  $(2^{(1/2)}*\log(((\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)} * 2i + 2^{(1/2)}*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x - 2^{(1/2)}*b*x)*((2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^3 - 8*a^3 - 6*a*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^2 + 12*a^2*(2*a - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + \log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)*i)/(2*x*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(1/2)})) * 2i)/(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x)^{(3/2)} - (8*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)})/(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))*(\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) - \log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x))$

**Reduce [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x} dx$$

input `int(1/x/atanh(tanh(b*x+a))^(3/2),x)`

output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**2*x),x)`

### 3.155 $\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1137
Mathematica [A] (verified)	1138
Rubi [A] (verified)	1138
Maple [A] (verified)	1140
Fricas [A] (verification not implemented)	1141
Sympy [F]	1141
Maxima [F]	1142
Giac [A] (verification not implemented)	1142
Mupad [B] (verification not implemented)	1142
Reduce [F]	1143

#### Optimal result

Integrand size = 15, antiderivative size = 124

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$\frac{3b \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{b}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{3b}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-3*b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(
b*x-arctanh(tanh(b*x+a)))^(5/2)-1/x/arctanh(tanh(b*x+a))^(3/2)+b/(b*x-arct
anh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)-3*b/(b*x-arctanh(tanh(b*x+a))
)^2/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.73

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{3b \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} - \frac{2bx + \operatorname{arctanh}(\tanh(a + bx))}{x \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Integrate[1/(x^2*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(3*b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2) - (2*b*x + ArcTanh[Tanh[a + b*x]])/(x*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.19, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

↓ 2599

$$-\frac{3}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2594

$$-\frac{3}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

↓ 2594

$$-\frac{3}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

↓ 2592

$$-\frac{3}{2}b \left( -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Int[1/(x^2*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-3*b*(-(((2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(3/2))`



## Definitions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.85

method	result
default	$2b \left( -\frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2bx} - \frac{3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))} - bx}\right)}{2\sqrt{\operatorname{arctanh}(\tanh(bx+a)) - bx}} \right)$

input `int(1/x^2/arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)`

output `2*b*(-1/(arctanh(tanh(b*x+a))-b*x)^2/arctanh(tanh(b*x+a))^(1/2)-1/(arctanh(tanh(b*x+a))-b*x)^2*(1/2*arctanh(tanh(b*x+a))^(1/2)/b/x-3/2/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \left[ \frac{3(b^2x^2 + abx)\sqrt{a} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(3abx + a^2)\sqrt{bx+a}}{2(a^3bx^2 + a^4x)}, \right. \\ \left. - \frac{3(b^2x^2 + abx)\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (3abx + a^2)\sqrt{bx+a}}{a^3bx^2 + a^4x} \right]$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[1/2*(3*(b^2*x^2 + a*b*x)*sqrt(a)*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(3*a*b*x + a^2)*sqrt(b*x + a))/(a^3*b*x^2 + a^4*x), -(3*(b^2*x^2 + a*b*x)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b*x + a^2)*sqrt(b*x + a))/(a^3*b*x^2 + a^4*x)]`

**Sympy [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^2 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**2/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(1/(x**2*atanh(tanh(a + b*x))**(3/2)), x)`

**Maxima [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^2 \operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(1/(x^2*arctanh(tanh(b*x + a))^(3/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.52

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{3b \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} - \frac{3(bx+a)b - 2ab}{\left((bx+a)^{\frac{3}{2}} - \sqrt{bx+aa}\right)a^2}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `-3*b*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) - (3*(b*x + a)*b - 2*a*b)/(((b*x + a)^(3/2) - sqrt(b*x + a))*a^2)`

**Mupad [B] (verification not implemented)**

Time = 8.90 (sec) , antiderivative size = 807, normalized size of antiderivative = 6.51

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `int(1/(x^2*atanh(tanh(a + b*x))^(3/2)),x)`

output

```

((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1)))/2)^(1/2)*(4/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2
*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (24*b*x)/(log(
2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2
*b*x) + 1)) + 2*b*x)^2)/(2*b*x^2 - x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) -
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (2^(1/2
)*b*log((((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(
2/(exp(2*a)*exp(2*b*x) + 1)))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) -
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i +
2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(
exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*
exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1))
+ 2*b*x)^5 + 40*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)
) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 80*a^3*(2*a - log(
(2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2
*b*x) + 1)) + 2*b*x)^2 - 32*a^5 - 10*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 +
80*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + lo
g(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*1i)/(2*x*(log(2/(exp(2*a)*exp(2*b
*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*...

```

**Reduce [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x^2} dx$$

input

```
int(1/x^2/atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**2*x**2),x)
```

**3.156**  $\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1144
Mathematica [A] (verified)	1145
Rubi [A] (verified)	1145
Maple [A] (verified)	1148
Fricas [A] (verification not implemented)	1148
Sympy [F]	1149
Maxima [F]	1149
Giac [A] (verification not implemented)	1149
Mupad [B] (verification not implemented)	1150
Reduce [F]	1150

**Optimal result**

Integrand size = 15, antiderivative size = 191

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$\frac{15b^2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} + \frac{3b}{4x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

$$- \frac{3b^2}{4(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

$$- \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{5b^2}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{15b^2}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-15/4*b^2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(7/2)+3/4*b/x/arctanh(tanh(b*x+a))^(5/2)-3/4*b^2/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(5/2)-1/2/x^2/arctanh(tanh(b*x+a))^(3/2)+5/4*b^2/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(3/2)-15/4*b^2/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{1}{4} \left( -\frac{15b^2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{7/2}} \right. \\ \left. + \frac{8b^2x^2 + 9bx \operatorname{arctanh}(\tanh(a + bx)) - 2 \operatorname{arctanh}(\tanh(a + bx))^2}{x^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (-bx + \operatorname{arctanh}(\tanh(a + bx)))^3} \right)$$

input `Integrate[1/(x^3*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `((-15*b^2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]]^(7/2) + (8*b^2*x^2 + 9*b*x*ArcTanh[Tanh[a + b*x]] - 2*ArcTanh[Tanh[a + b*x]]^2)/(x^2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^3))/4`

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.15, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx \\ \downarrow 2599 \\ -\frac{3}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} \\ \downarrow 2599$$

$$\begin{aligned}
 & -\frac{3}{4}b \left( -\frac{5}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow 2594 \\
 & -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow 2594 \\
 & -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow 2594 \\
 & -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \qquad \qquad \qquad \downarrow 2592 \\
 & -\frac{3}{4}b \left( -\frac{5}{2}b \left( -\frac{2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) - \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) - \\
 & \qquad \qquad \qquad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}
 \end{aligned}$$

input `Int[1/(x^3*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-3*b*((-5*b*(-((-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2)))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(5/2)))/4 - 1/(2*x^2*ArcTanh[Tanh[a + b*x]]^(3/2))`

### Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`



**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.69

method	result
default	$2b^2 \left( \frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{\frac{7 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{8} + \left(-\frac{9 \operatorname{arctanh}(\tanh(bx+a))}{8} + \frac{9bx}{8}\right) \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b^2 x^2} \right) (\operatorname{arctanh}(\tanh(bx+a)))$

input `int(1/x^3/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2*b^2*(1/(arctanh(tanh(b*x+a))-b*x)^3/arctanh(tanh(b*x+a))^(1/2)+1/(arctanh(tanh(b*x+a))-b*x)^3*((7/8*arctanh(tanh(b*x+a))^(3/2)+(-9/8*arctanh(tanh(b*x+a))+9/8*b*x)*arctanh(tanh(b*x+a))^(1/2))/b^2/x^2-15/8/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.97

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{15(b^3 x^3 + ab^2 x^2) \sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(15ab^2 x^2 + 5a^2 bx - 2a^3) \sqrt{bx+a}}{8(a^4 bx^3 + a^5 x^2)}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[1/8*(15*(b^3*x^3 + a*b^2*x^2)*sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(15*a*b^2*x^2 + 5*a^2*b*x - 2*a^3)*sqrt(b*x + a))/(a^4*b*x^3 + a^5*x^2), 1/4*(15*(b^3*x^3 + a*b^2*x^2)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (15*a*b^2*x^2 + 5*a^2*b*x - 2*a^3)*sqrt(b*x + a))/(a^4*b*x^3 + a^5*x^2)]`

**Sympy [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^3 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**3/atanh(tanh(b*x+a))**(3/2), x)`

output `Integral(1/(x**3*atanh(tanh(a + b*x))**(3/2)), x)`

**Maxima [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^3 \operatorname{artanh}^{\frac{3}{2}}(\tanh(bx + a))} dx$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(3/2), x, algorithm="maxima")`

output `integrate(1/(x^3*arctanh(tanh(b*x + a))^(3/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{15 b^2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{4 \sqrt{-a} a^3} + \frac{2 b^2}{\sqrt{bx+a} a^3} + \frac{7 (bx+a)^{\frac{3}{2}} b^2 - 9 \sqrt{bx+a} a b^2}{4 a^3 b^2 x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(3/2), x, algorithm="giac")`

output `15/4*b^2*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^3) + 2*b^2/(sqrt(b*x + a)*a^3) + 1/4*(7*(b*x + a)^(3/2)*b^2 - 9*sqrt(b*x + a)*a*b^2)/(a^3*b^2*x^2)`

**Mupad [B] (verification not implemented)**

Time = 8.62 (sec) , antiderivative size = 1028, normalized size of antiderivative = 5.38

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `int(1/(x^3*atanh(tanh(a + b*x))^(3/2)),x)`

output

$$\begin{aligned} & (2^{(1/2)}b^2 \log\left(\frac{\log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1}\right) / 2 - \log(2/(\exp(2a)\exp(2bx) + 1)) / 2)^{(1/2)} \left( \log(2/(\exp(2a)\exp(2bx) + 1)) - \log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) + 2bx \right)^{(1/2)} \\ & + 2i + 2^{(1/2)} \left( \log(2/(\exp(2a)\exp(2bx) + 1)) - \log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) + 2bx \right) - 2^{(1/2)}bx \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^7 \\ & + 84a^2(2a - \log(2\exp(2a)\exp(2bx)) / (\exp(2a)\exp(2bx) + 1)) + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx)^5 - 280a^3 \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^4 \\ & + 560a^4 \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^3 - 672a^5 \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^2 \\ & - 128a^7 - 14a \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^6 + 448a^6 \left( \frac{2a - \log(2\exp(2a)\exp(2bx))}{\exp(2a)\exp(2bx) + 1} + \log(2/(\exp(2a)\exp(2bx) + 1)) + 2bx \right)^5 \\ & + 11i / (2x \left( \log(2/(\exp(2a)\exp(2bx) + 1)) - \log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) + 2bx \right)^{(1/2)}) + 15i / \left( \log(2/(\exp(2a)\exp(2bx) + 1)) - \log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) + 2bx \right)^{(7/2)} - 2 \left( \log\left(\frac{2\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) / (\exp(2a)\exp(2bx) + 1) / 2 - \log(2/(\dots \right) \end{aligned}$$
**Reduce [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x^3} dx$$

input `int(1/x^3/atanh(tanh(b*x+a))^(3/2),x)`

output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**2*x**3),x)`

**3.157**  $\int \frac{1}{x^4 \mathbf{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1152
Mathematica [A] (verified)	1153
Rubi [A] (verified)	1153
Maple [A] (verified)	1157
Fricas [A] (verification not implemented)	1157
Sympy [F]	1158
Maxima [F]	1158
Giac [A] (verification not implemented)	1159
Mupad [B] (verification not implemented)	1159
Reduce [F]	1160

**Optimal result**

Integrand size = 15, antiderivative size = 245

$$\int \frac{1}{x^4 \mathbf{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$\frac{35b^3 \arctan\left(\frac{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \mathbf{arctanh}(\tanh(a+bx))}}\right)}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^{9/2}} - \frac{5b^2}{8x \mathbf{arctanh}(\tanh(a+bx))^{7/2}}$$

$$+ \frac{5b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx))) \mathbf{arctanh}(\tanh(a+bx))^{7/2}}$$

$$+ \frac{b}{4x^2 \mathbf{arctanh}(\tanh(a+bx))^{5/2}}$$

$$- \frac{7b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^2 \mathbf{arctanh}(\tanh(a+bx))^{5/2}}$$

$$- \frac{1}{3x^3 \mathbf{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{35b^3}{24(bx - \mathbf{arctanh}(\tanh(a+bx)))^3 \mathbf{arctanh}(\tanh(a+bx))^{3/2}}$$

$$- \frac{35b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^4 \sqrt{\mathbf{arctanh}(\tanh(a+bx))}}$$

output

$$\begin{aligned}
& -35/8*b^3*\arctan(\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)})/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)} \\
& -5/8*b^2/x/\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)}+5/8*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/\operatorname{arctanh}(\tanh(b*x+a))^{(7/2)} \\
& +1/4*b/x^2/\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)}-7/8*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2/\operatorname{arctanh}(\tanh(b*x+a))^{(5/2)} \\
& -1/3/x^3/\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}+35/24*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^3/\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)} \\
& -35/8*b^3/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^4/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}
\end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.54

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{35b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{8(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{9/2}} - \frac{48b^3x^3 + 87b^2x^2 \operatorname{arctanh}(\tanh(a+bx)) - 38bx \operatorname{arctanh}(\tanh(a+bx))^2 + 8 \operatorname{arctanh}(\tanh(a+bx))^3}{24x^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^4}$$

input

`Integrate[1/(x^4*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output

$$\begin{aligned}
& (35*b^3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(8*(-(b*x) + ArcTanh[Tanh[a + b*x]])^{(9/2)}) - (48*b^3*x^3 + 87*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 38*b*x*ArcTanh[Tanh[a + b*x]]^2 + 8*ArcTanh[Tanh[a + b*x]]^3)/(24*x^3*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)
\end{aligned}$$
**Rubi [A] (verified)**

Time = 0.65 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.19, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$ , Rules used = {2599, 2599, 2599, 2594, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx \\
 & \quad \downarrow \text{2599} \\
 & -\frac{1}{2}b \int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx - \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2599} \\
 & -\frac{1}{2}b \left( -\frac{5}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2599} \\
 & -\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{9/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} \right) - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2594} \\
 & -\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{7(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{7/2}} \right) \right) \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2594} \\
 & -\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \right) \right) - \frac{1}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2594}
 \end{aligned}$$

$$-\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

2594

$$-\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

2592

$$-\frac{1}{2}b \left( -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{2 \operatorname{arctan} \left( \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input

```
Int [1/(x^4*ArcTanh[Tanh[a + b*x]]^(3/2)),x]
```



output

```
-1/2*(b*((-5*b*((-7*b*(-((-((-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])))/(b*x - ArcTanh[Tanh[a + b*x]]))^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(7*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(7/2)))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(7/2)))/4 - 1/(2*x^2*ArcTanh[Tanh[a + b*x]]^(5/2))) - 1/(3*x^3*ArcTanh[Tanh[a + b*x]]^(3/2))
```

### Defintions of rubi rules used

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.76

method	result
default	$2b^3 \left( -\frac{1}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{\frac{19 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{16} + \left( -\frac{17 \operatorname{arctanh}(\tanh(bx+a))}{6} + \frac{17bx}{6} \right) \operatorname{arctanh}(\tanh(bx+a))}{\dots} \right)$

input `int(1/x^4/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2*b^3*(-1/(arctanh(tanh(b*x+a))-b*x)^4/arctanh(tanh(b*x+a))^(1/2)-1/(arctanh(tanh(b*x+a))-b*x)^4*((19/16*arctanh(tanh(b*x+a))^(5/2)+(-17/6*arctanh(tanh(b*x+a))+17/6*b*x)*arctanh(tanh(b*x+a))^(3/2)+(29/16*a^2+29/8*a*(arctanh(tanh(b*x+a))-b*x-a)+29/16*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2))/b^3/x^3-35/16/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.85

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \left[ \frac{105(b^4x^4 + ab^3x^3)\sqrt{a} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(105ab^3x^3 + 35a^2b^2x^2)}{48(a^5bx^4 + a^6x^3)} - \frac{105(b^4x^4 + ab^3x^3)\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (105ab^3x^3 + 35a^2b^2x^2 - 14a^3bx + 8a^4)\sqrt{bx+a}}{24(a^5bx^4 + a^6x^3)} \right]$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output

```
[1/48*(105*(b^4*x^4 + a*b^3*x^3)*sqrt(a)*log((b*x + 2*sqrt(b*x + a)*sqrt(a)
+ 2*a)/x) - 2*(105*a*b^3*x^3 + 35*a^2*b^2*x^2 - 14*a^3*b*x + 8*a^4)*sqrt
(b*x + a))/(a^5*b*x^4 + a^6*x^3), -1/24*(105*(b^4*x^4 + a*b^3*x^3)*sqrt(-a
)*arctan(sqrt(-a)/sqrt(b*x + a)) + (105*a*b^3*x^3 + 35*a^2*b^2*x^2 - 14*a^
3*b*x + 8*a^4)*sqrt(b*x + a))/(a^5*b*x^4 + a^6*x^3)]
```

**Sympy [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^4 \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input

```
integrate(1/x**4/atanh(tanh(b*x+a))**(3/2), x)
```

output

```
Integral(1/(x**4*atanh(tanh(a + b*x))**(3/2)), x)
```

**Maxima [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^4 \operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}} dx$$

input

```
integrate(1/x^4/arctanh(tanh(b*x+a))^(3/2), x, algorithm="maxima")
```

output

```
integrate(1/(x^4*arctanh(tanh(b*x + a))^(3/2)), x)
```

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.39

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{35 b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{8 \sqrt{-a} a^4} - \frac{2 b^3}{\sqrt{bx+aa^4}} - \frac{57 (bx+a)^{5/2} b^3 - 136 (bx+a)^{3/2} a b^3 + 87 \sqrt{bx+aa^2} b^3}{24 a^4 b^3 x^3}$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `-35/8*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^4) - 2*b^3/(sqrt(b*x + a)*a^4) - 1/24*(57*(b*x + a)^(5/2)*b^3 - 136*(b*x + a)^(3/2)*a*b^3 + 87*sqrt(b*x + a)*a^2*b^3)/(a^4*b^3*x^3)`

**Mupad [B] (verification not implemented)**

Time = 7.35 (sec) , antiderivative size = 1258, normalized size of antiderivative = 5.13

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Too large to display}$$

input `int(1/(x^4*atanh(tanh(a + b*x))^(3/2)),x)`

output

```

(((38*b^2)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))
/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - (140*b^3*x)/(log(2/(exp(2*a)*exp(
2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*
b*x)^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/
(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(x*(log((2*exp(2*a)*exp(2*b*x))/(exp(
2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) - (4*(log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*
x) + 1))/2)^(1/2))/(3*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (2^(1/2)*b^3*log(
(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2
*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*2i + 2^(1/2)
*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
*exp(2*b*x) + 1)) + 2*b*x) - 2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)
^9 + 144*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
+ log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^7 - 672*a^3*(2*a - log((2*exp
(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x)^6 + 2016*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*e
xp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5 - 4032*a^...

```

**Reduce [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x^4} dx$$

input

```
int(1/x^4/atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**2*x**4),x)
```

**3.158**  $\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1161
Mathematica [A] (verified)	1161
Rubi [A] (verified)	1162
Maple [B] (verified)	1164
Fricas [A] (verification not implemented)	1165
Sympy [F]	1165
Maxima [A] (verification not implemented)	1166
Giac [A] (verification not implemented)	1166
Mupad [B] (verification not implemented)	1166
Reduce [F]	1167

**Optimal result**

Integrand size = 15, antiderivative size = 99

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2x^4}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{16x^3}{3b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{32x^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^3} - \frac{128x\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^4} + \frac{256\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{15b^5}$$

output

```
-2/3*x^4/b/arctanh(tanh(b*x+a))^(3/2)-16/3*x^3/b^2/arctanh(tanh(b*x+a))^(1/2)+32*x^2*arctanh(tanh(b*x+a))^(1/2)/b^3-128/3*x*arctanh(tanh(b*x+a))^(3/2)/b^4+256/15*arctanh(tanh(b*x+a))^(5/2)/b^5
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.84

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2(5b^4x^4 + 40b^3x^3\operatorname{arctanh}(\tanh(a+bx)) - 240b^2x^2\operatorname{arctanh}(\tanh(a+bx))^2 + 320bx\operatorname{arctanh}(\tanh(a+bx)) - 15b^5\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{15b^5\operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Integrate[x^4/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output  $(-2*(5*b^4*x^4 + 40*b^3*x^3*ArcTanh[Tanh[a + b*x]] - 240*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 + 320*b*x*ArcTanh[Tanh[a + b*x]]^3 - 128*ArcTanh[Tanh[a + b*x]]^4)/(15*b^5*ArcTanh[Tanh[a + b*x]]^(3/2))$

### Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.18, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx \\
 & \quad \downarrow 2599 \\
 & \frac{8 \int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3b} - \frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599 \\
 & \frac{8 \left( \frac{6 \int \frac{x^2}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^3}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599 \\
 & \frac{8 \left( \frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \int x \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} \right)}{b} - \frac{2x^3}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599 \\
 & \frac{3b}{2x^4} - \frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599
 \end{aligned}$$

$$8 \left( \frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx}{b} \right)}{b} \right)}{b} - \frac{2x^3}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)$$

$$\frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

↓ 2588

$$8 \left( \frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{2 \int \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx}{b} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))}{3b^2} \right)}{b} \right)}{b} - \frac{2x^3}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)$$

$$\frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

↓ 15

$$8 \left( \frac{6 \left( \frac{2x^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \left( \frac{2x \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{15b^2} \right)}{b} \right)}{b} - \frac{2x^3}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)$$

$$\frac{2x^4}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Int [x^4/ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(-2*x^4)/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2)) + (8*((-2*x^3)/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (6*((2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]])/b - (4*((2*x*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b) - (4*ArcTanh[Tanh[a + b*x]]^(5/2))/(15*b^2)))/b))/b)/(3*b)`



### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :=> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] :=> With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :=> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs.  $2(81) = 162$ .

Time = 0.21 (sec) , antiderivative size = 295, normalized size of antiderivative = 2.98

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a)) \frac{5}{2} - 8 \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} a - 8 \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} (\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + 12 \sqrt{\operatorname{arctanh}(\tanh(bx+a))} a^2}{\dots}$

input `int(x^4/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output

```
2/b^5*(1/5*arctanh(tanh(b*x+a))^(5/2)-4/3*arctanh(tanh(b*x+a))^(3/2)*a-4/3
*arctanh(tanh(b*x+a))^(3/2)*(arctanh(tanh(b*x+a))-b*x-a)+6*arctanh(tanh(b*
x+a))^(1/2)*a^2+12*a*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/
2)+6*(arctanh(tanh(b*x+a))-b*x-a)^2*arctanh(tanh(b*x+a))^(1/2)-(-4*a^3-12*
a^2*(arctanh(tanh(b*x+a))-b*x-a)-12*a*(arctanh(tanh(b*x+a))-b*x-a)^2-4*(ar
ctanh(tanh(b*x+a))-b*x-a)^3)/arctanh(tanh(b*x+a))^(1/2)-1/3*(a^4+4*a^3*(ar
ctanh(tanh(b*x+a))-b*x-a)+6*a^2*(arctanh(tanh(b*x+a))-b*x-a)^2+4*a*(arctan
h(tanh(b*x+a))-b*x-a)^3+(arctanh(tanh(b*x+a))-b*x-a)^4)/arctanh(tanh(b*x+a
))^(3/2))
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.75

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(3b^4x^4 - 8ab^3x^3 + 48a^2b^2x^2 + 192a^3bx + 128a^4)\sqrt{bx + a}}{15(b^7x^2 + 2ab^6x + a^2b^5)}$$

input

```
integrate(x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
2/15*(3*b^4*x^4 - 8*a*b^3*x^3 + 48*a^2*b^2*x^2 + 192*a^3*b*x + 128*a^4)*sq
rt(b*x + a)/(b^7*x^2 + 2*a*b^6*x + a^2*b^5)
```

### Sympy [F]

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^4}{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))} dx$$

input

```
integrate(x**4/atanh(tanh(b*x+a))**(5/2),x)
```

output

```
Integral(x**4/atanh(tanh(a + b*x))**(5/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(3b^5x^5 - 5ab^4x^4 + 40a^2b^3x^3 + 240a^3b^2x^2 + 320a^4bx + 128a^5)}{15(bx + a)^{5/2}b^5}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`output `2/15*(3*b^5*x^5 - 5*a*b^4*x^4 + 40*a^2*b^3*x^3 + 240*a^3*b^2*x^2 + 320*a^4*b*x + 128*a^5)/((b*x + a)^(5/2)*b^5)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.80

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2 \left( \frac{5(12(bx+a)a^3 - a^4)}{(bx+a)^{3/2}b} + \frac{3(bx+a)^{5/2}b^4 - 20(bx+a)^{3/2}ab^4 + 90\sqrt{bx+aa^2b^4}}{b^5} \right)}{15b^4}$$

input `integrate(x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`output `2/15*(5*(12*(b*x + a)*a^3 - a^4)/((b*x + a)^(3/2)*b) + (3*(b*x + a)^(5/2)*b^4 - 20*(b*x + a)^(3/2)*a*b^4 + 90*sqrt(b*x + a)*a^2*b^4)/b^5)/b^4`**Mupad [B] (verification not implemented)**

Time = 3.51 (sec) , antiderivative size = 817, normalized size of antiderivative = 8.25

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(x^4/atanh(tanh(a + b*x))^(5/2),x)`

output

```

((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^(1/2)*((3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((
2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*b^4) + (2
*((2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(
2*a)*exp(2*b*x) + 1)) + 2*b*x))/b^3 + (8*(log(2/(exp(2*a)*exp(2*b*x) + 1))
/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x))/(5*b
^3))*(log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(ex
p(2*a)*exp(2*b*x) + 1))/2 + b*x))/(3*b))/b + (2*x^2*(log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)
^(1/2))/(5*b^3) + (x*((2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*
a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b^3 + (8*(log(2/(exp(2
*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)
+ 1))/2 + b*x))/(5*b^3))*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)
+ 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*b) - (2*(log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b
*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(b^5*(log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) -
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2...

```

**Reduce [F]**

$$\int \frac{x^4}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^4}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input

```
int(x^4/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int((sqrt(atanh(tanh(a + b*x))))*x**4)/atanh(tanh(a + b*x))**3,x)
```

**3.159**      $\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1168
Mathematica [A] (verified)	1168
Rubi [A] (verified)	1169
Maple [B] (verified)	1171
Fricas [A] (verification not implemented)	1171
Sympy [A] (verification not implemented)	1172
Maxima [A] (verification not implemented)	1172
Giac [A] (verification not implemented)	1173
Mupad [B] (verification not implemented)	1173
Reduce [F]	1174

**Optimal result**

Integrand size = 15, antiderivative size = 76

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2x^3}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{4x^2}{b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{16x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^3} - \frac{32\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^4}$$

output

```
-2/3*x^3/b/arctanh(tanh(b*x+a))^(3/2)-4*x^2/b^2/arctanh(tanh(b*x+a))^(1/2)
+16*x*arctanh(tanh(b*x+a))^(1/2)/b^3-32/3*arctanh(tanh(b*x+a))^(3/2)/b^4
```

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.86

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2(b^3x^3 + 6b^2x^2\operatorname{arctanh}(\tanh(a+bx)) - 24bx\operatorname{arctanh}(\tanh(a+bx))^2 + 16\operatorname{arctanh}(\tanh(a+bx))^3)}{3b^4\operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Integrate[x^3/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(-2*(b^3*x^3 + 6*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 24*b*x*ArcTanh[Tanh[a + b*x]]^2 + 16*ArcTanh[Tanh[a + b*x]]^3)/(3*b^4*ArcTanh[Tanh[a + b*x]]^(3/2))`

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx \\
 & \quad \downarrow 2599 \\
 & \frac{2 \int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{b} - \frac{2x^3}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599 \\
 & \frac{2 \left( \frac{4 \int \frac{x}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^2}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{b} - \frac{2x^3}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2599 \\
 & \frac{2 \left( \frac{4 \left( \frac{2x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} dx}{b} \right)}{b} - \frac{2x^2}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{b} - \frac{2x^3}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2588 \\
 & \frac{b}{2x^3} - \frac{2x^3}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}
 \end{aligned}$$

$$\begin{array}{c}
 2 \left( \frac{4 \left( \frac{2x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{2 \int \sqrt{\operatorname{arctanh}(\tanh(a+bx))} \operatorname{arctanh}(\tanh(a+bx))}{b^2} \right)}{b} - \frac{2x^2}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \\
 \hline
 \frac{b}{2x^3} \\
 \frac{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^3} \\
 \downarrow 15 \\
 2 \left( \frac{4 \left( \frac{2x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} - \frac{4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3b^2} \right)}{b} - \frac{2x^2}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \\
 \hline
 \frac{b}{2x^3} \\
 \frac{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2x^3}
 \end{array}$$

input `Int [x^3/ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(-2*x^3)/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2)) + (2*((-2*x^2)/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (4*((2*x*Sqrt[ArcTanh[Tanh[a + b*x]]])/b - (4*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*b^2)))/b)/b`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs.  $2(64) = 128$ .

Time = 0.20 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.45

method	result
default	$\frac{2 \operatorname{arctanh}(\tanh(bx+a)) \frac{3}{2} - 6 \sqrt{\operatorname{arctanh}(\tanh(bx+a))} a - 6(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) \sqrt{\operatorname{arctanh}(\tanh(bx+a))} - \frac{2(3a^2 + 6a \operatorname{arctanh}(\tanh(bx+a)))}{3}}$

input

```
int(x^3/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
2/b^4*(1/3*arctanh(tanh(b*x+a))^(3/2)-3*arctanh(tanh(b*x+a))^(1/2)*a-3*(arctanh(tanh(b*x+a))-b*x-a)*arctanh(tanh(b*x+a))^(1/2)-(3*a^2+6*a*(arctanh(tanh(b*x+a))-b*x-a)+3*(arctanh(tanh(b*x+a))-b*x-a)^2)/arctanh(tanh(b*x+a))^(1/2)-1/3*(-a^3-3*a^2*(arctanh(tanh(b*x+a))-b*x-a)-3*a*(arctanh(tanh(b*x+a))-b*x-a)^2-(arctanh(tanh(b*x+a))-b*x-a)^3)/arctanh(tanh(b*x+a))^(3/2))
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.82

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(b^3 x^3 - 6ab^2 x^2 - 24a^2 bx - 16a^3) \sqrt{bx + a}}{3(b^6 x^2 + 2ab^5 x + a^2 b^4)}$$

input

```
integrate(x^3/arctanh(tanh(b*x+a))^(5/2), x, algorithm="fricas")
```



output  $\frac{2}{3}(b^3x^3 - 6ab^2x^2 - 24a^2bx - 16a^3)\sqrt{bx + a}/(b^6x^2 + 2ab^5x + a^2b^4)$

### Sympy [A] (verification not implemented)

Time = 2.41 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.18

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \begin{cases} -\frac{2x^3}{3b \operatorname{atanh}^{\frac{3}{2}}(\tanh(a+bx))} - \frac{4x^2}{b^2 \sqrt{\operatorname{atanh}(\tanh(a+bx))}} + \frac{16x \sqrt{\operatorname{atanh}(\tanh(a+bx))}}{b^3} - \frac{32}{4 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a))} \end{cases}$$

input `integrate(x**3/atanh(tanh(b*x+a))**(5/2), x)`

output `Piecewise((-2*x**3/(3*b*atanh(tanh(a + b*x))**(3/2)) - 4*x**2/(b**2*sqrt(a*tanh(tanh(a + b*x)))) + 16*x*sqrt(atanh(tanh(a + b*x)))/b**3 - 32*atanh(tanh(a + b*x))**(3/2)/(3*b**4), Ne(b, 0)), (x**4/(4*atanh(tanh(a))**(5/2)), True))`

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.68

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(b^4x^4 - 5ab^3x^3 - 30a^2b^2x^2 - 40a^3bx - 16a^4)}{3(bx + a)^{\frac{5}{2}}b^4}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(5/2), x, algorithm="maxima")`

output  $\frac{2}{3}(b^4x^4 - 5a^3b^3x^3 - 30a^2b^2x^2 - 40a^3bx - 16a^4)/((bx + a)^{5/2}b^4)$

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2(9(bx + a)a^2 - a^3)}{3(bx + a)^{3/2}b^4} + \frac{2\left((bx + a)^{3/2}b^8 - 9\sqrt{bx + a}ab^8\right)}{3b^{12}}$$

input `integrate(x^3/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-2/3*(9*(b*x + a)*a^2 - a^3)/((b*x + a)^(3/2)*b^4) + 2/3*((b*x + a)^(3/2)*b^8 - 9*sqrt(b*x + a)*a*b^8)/b^12`

**Mupad [B] (verification not implemented)**

Time = 3.25 (sec) , antiderivative size = 533, normalized size of antiderivative = 7.01

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(x^3/atanh(tanh(a + b*x))^(5/2),x)`

output

```
(2*x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*b^3) + (((2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/b^3 + (4*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 + b*x))/(3*b^3))*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/b - (3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(b^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) - ((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(3*b^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^2)
```

**Reduce [F]**

$$\int \frac{x^3}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input

```
int(x^3/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int((sqrt(atanh(tanh(a + b*x)))*x**3)/atanh(tanh(a + b*x))**3,x)
```

### 3.160 $\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1175
Mathematica [A] (verified)	1175
Rubi [A] (verified)	1176
Maple [A] (verified)	1177
Fricas [A] (verification not implemented)	1178
Sympy [A] (verification not implemented)	1178
Maxima [A] (verification not implemented)	1178
Giac [A] (verification not implemented)	1179
Mupad [B] (verification not implemented)	1179
Reduce [F]	1180

#### Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2x^2}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{8x}{3b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{16\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b^3}$$

output `-2/3*x^2/b/arctanh(tanh(b*x+a))^(3/2)-8/3*x/b^2/arctanh(tanh(b*x+a))^(1/2)+16/3*arctanh(tanh(b*x+a))^(1/2)/b^3`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.81

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2(b^2x^2 + 4bx\operatorname{arctanh}(\tanh(a+bx)) - 8\operatorname{arctanh}(\tanh(a+bx))^2)}{3b^3\operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Integrate[x^2/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output  $(-2*(b^2*x^2 + 4*b*x*ArcTanh[Tanh[a + b*x]] - 8*ArcTanh[Tanh[a + b*x]]^2)) / (3*b^3*ArcTanh[Tanh[a + b*x]]^(3/2))$

### Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$$

$$\downarrow 2599$$

$$\frac{4 \int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3b} - \frac{2x^2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$\downarrow 2599$$

$$\frac{4 \left( \frac{2 \int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$\downarrow 2588$$

$$\frac{4 \left( \frac{2 \int \frac{1}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} d \operatorname{arctanh}(\tanh(a+bx))}{b^2} - \frac{2x}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$\downarrow 15$$

$$\frac{4 \left( \frac{4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2} - \frac{2x}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input  $\text{Int}[x^2/\text{ArcTanh}[\text{Tanh}[a + b*x]]^{(5/2)}, x]$

output  $(4*((-2*x)/(b*\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]]) + (4*\text{Sqrt}[\text{ArcTanh}[\text{Tanh}[a + b*x]]])/b^2))/(3*b) - (2*x^2)/(3*b*\text{ArcTanh}[\text{Tanh}[a + b*x]]^{(3/2)})$

**Defintions of rubi rules used**

rule 15  $\text{Int}[(a_.)*(x_)^(m_.), x\_Symbol] \rightarrow \text{Simp}[a*(x^(m + 1)/(m + 1)), x] /; \text{FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 2588  $\text{Int}[(u_)^(m_.), x\_Symbol] \rightarrow \text{With}[\{c = \text{Simplify}[D[u, x]]\}, \text{Simp}[1/c \ \text{Subst}[\text{Int}[x^m, x], x, u], x]] /; \text{FreeQ}[m, x] \ \&\& \ \text{PiecewiseLinearQ}[u, x]$

rule 2599  $\text{Int}[(u_)^(m_.)*(v_)^(n_.), x\_Symbol] \rightarrow \text{With}[\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[u^(m + 1)*(v^n/(a*(m + 1))), x] - \text{Simp}[b*(n/(a*(m + 1))) \ \text{Int}[u^(m + 1)*v^(n - 1), x], x] /; \text{NeQ}[b*u - a*v, 0] /; \text{FreeQ}[\{m, n\}, x] \ \&\& \ \text{PiecewiseLinearQ}[u, v, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ ((\text{LtQ}[m, -1] \ \&\& \ \text{GtQ}[n, 0]) \ \&\& \ !(\text{ILtQ}[m + n, -2] \ \&\& \ (\text{FractionQ}[m] \ || \ \text{GeQ}[2*n + m + 1, 0]))) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LeQ}[n, m]) \ || \ (\text{IGtQ}[n, 0] \ \&\& \ !\text{IntegerQ}[m]) \ || \ (\text{ILtQ}[m, 0] \ \&\& \ !\text{IntegerQ}[n]))$

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.54

method	result	size
default	$\frac{2\sqrt{\text{arctanh}(\tanh(bx+a))} - \frac{2(2bx-2 \text{arctanh}(\tanh(bx+a)))}{\sqrt{\text{arctanh}(\tanh(bx+a))}} - \frac{2(a^2+2a(\text{arctanh}(\tanh(bx+a))-bx-a)+(\text{arctanh}(\tanh(bx+a))-bx-a)^2)}{3 \text{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}}{b^3}$	91

input `int(x^2/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output  $2/b^3*(\text{arctanh}(\tanh(b*x+a))^{(1/2)} - (2*b*x - 2*\text{arctanh}(\tanh(b*x+a)))/\text{arctanh}(\tanh(b*x+a))^{(1/2)} - 1/3*(a^2 + 2*a*(\text{arctanh}(\tanh(b*x+a)) - b*x - a) + (\text{arctanh}(\tanh(b*x+a)) - b*x - a)^2)/\text{arctanh}(\tanh(b*x+a))^{(3/2)})$

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(3b^2x^2 + 12abx + 8a^2)\sqrt{bx + a}}{3(b^5x^2 + 2ab^4x + a^2b^3)}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`output `2/3*(3*b^2*x^2 + 12*a*b*x + 8*a^2)*sqrt(b*x + a)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)`**Sympy [A] (verification not implemented)**

Time = 2.38 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.20

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \begin{cases} -\frac{2x^2}{3b \operatorname{atanh}^{\frac{3}{2}}(\tanh(a+bx))} - \frac{8x}{3b^2 \sqrt{\operatorname{atanh}(\tanh(a+bx))}} + \frac{16 \sqrt{\operatorname{atanh}(\tanh(a+bx))}}{3b^3} \\ \frac{x^3}{3 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a))} \end{cases}$$

input `integrate(x**2/atanh(tanh(b*x+a))**(5/2),x)`output `Piecewise((-2*x**2/(3*b*atanh(tanh(a + b*x))**(3/2)) - 8*x/(3*b**2*sqrt(atanh(tanh(a + b*x)))) + 16*sqrt(atanh(tanh(a + b*x)))/(3*b**3), Ne(b, 0)), (x**3/(3*atanh(tanh(a))**(5/2)), True))`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.71

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(3b^3x^3 + 15ab^2x^2 + 20a^2bx + 8a^3)}{3(bx + a)^{\frac{5}{2}}b^3}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output  $2/3*(3*b^3*x^3 + 15*a*b^2*x^2 + 20*a^2*b*x + 8*a^3)/((b*x + a)^{(5/2)}*b^3)$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.73

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2 \left( \frac{3\sqrt{bx+a}}{b} + \frac{6(bx+a)a-a^2}{(bx+a)^{3/2}b} \right)}{3b^2}$$

input `integrate(x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output  $2/3*(3*\sqrt{b*x + a}/b + (6*(b*x + a)*a - a^2)/((b*x + a)^{(3/2)}*b))/b^2$

### Mupad [B] (verification not implemented)

Time = 3.30 (sec) , antiderivative size = 259, normalized size of antiderivative = 4.39

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{8 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{3b^3 \left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)} \left( -b^2 x^2 - 2bx \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)$$

input `int(x^2/atanh(tanh(a + b*x))^(5/2),x)`

output  $(8*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)))/2 - \log(2/(\exp(2*a)*\exp(2*b*x) + 1))/2)^{(1/2)}*(2*\log(2/(\exp(2*a)*\exp(2*b*x) + 1))^2 - 4*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))^2 - b^2*x^2 - 2*b*x*\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) + 2*b*x*\log(2/(\exp(2*a)*\exp(2*b*x) + 1)))/((3*b^3*(\log((2*\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1)) - \log(2/(\exp(2*a)*\exp(2*b*x) + 1)))^2)$



**Reduce [F]**

$$\int \frac{x^2}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^2/atanh(tanh(b*x+a))^(5/2),x)`

output `int((sqrt(atanh(tanh(a + b*x)))*x**2)/atanh(tanh(a + b*x))**3,x)`

### 3.161 $\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1181
Mathematica [A] (verified)	1181
Rubi [A] (verified)	1182
Maple [A] (verified)	1183
Fricas [A] (verification not implemented)	1183
Sympy [A] (verification not implemented)	1184
Maxima [A] (verification not implemented)	1184
Giac [A] (verification not implemented)	1184
Mupad [B] (verification not implemented)	1185
Reduce [B] (verification not implemented)	1185

#### Optimal result

Integrand size = 13, antiderivative size = 38

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2x}{3\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{4}{3b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output `-2/3*x/b/arctanh(tanh(b*x+a))^(3/2)-4/3/b^2/arctanh(tanh(b*x+a))^(1/2)`

#### Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2(bx + 2\operatorname{arctanh}(\tanh(a+bx)))}{3b^2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Integrate[x/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(-2*(b*x + 2*ArcTanh[Tanh[a + b*x]]))/(3*b^2*ArcTanh[Tanh[a + b*x]]^(3/2))`

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

$$\downarrow 2599$$

$$\frac{2 \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{3b} - \frac{2x}{3b \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

$$\downarrow 2588$$

$$\frac{2 \int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} d\operatorname{arctanh}(\tanh(a + bx))}{3b^2} - \frac{2x}{3b \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

$$\downarrow 15$$

$$-\frac{4}{3b^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{2x}{3b \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input `Int [x/ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(-2*x)/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2)) - 4/(3*b^2*Sqrt[ArcTanh[Tanh[a + b*x]]])`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

method	result	size
default	$-\frac{2}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a)))}{3 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$	42

input

```
int(x/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
2/b^2*(-1/arctanh(tanh(b*x+a))^(1/2)-1/3*(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2(3bx + 2a)\sqrt{bx + a}}{3(b^4x^2 + 2ab^3x + a^2b^2)}$$

input

```
integrate(x/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
-2/3*(3*b*x + 2*a)*sqrt(b*x + a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)
```

**Sympy [A] (verification not implemented)**

Time = 2.30 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \begin{cases} -\frac{2x}{3b \operatorname{atanh}^{3/2}(\tanh(a+bx))} - \frac{4}{3b^2 \sqrt{\operatorname{atanh}(\tanh(a+bx))}} & \text{for } b \neq 0 \\ \frac{x^2}{2 \operatorname{atanh}^{5/2}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(x/atanh(tanh(b*x+a))**(5/2), x)`output `Piecewise((-2*x/(3*b*atanh(tanh(a + b*x))**(3/2)) - 4/(3*b**2*sqrt(atanh(tanh(a + b*x))))), Ne(b, 0)), (x**2/(2*atanh(tanh(a))**(5/2)), True))`**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2(3b^2x^2 + 5abx + 2a^2)}{3(bx + a)^{5/2}b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(5/2), x, algorithm="maxima")`output `-2/3*(3*b^2*x^2 + 5*a*b*x + 2*a^2)/((b*x + a)^(5/2)*b^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.53

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2(3bx + 2a)}{3(bx + a)^{3/2}b^2}$$

input `integrate(x/arctanh(tanh(b*x+a))^(5/2), x, algorithm="giac")`output `-2/3*(3*b*x + 2*a)/((b*x + a)^(3/2)*b^2)`

**Mupad [B] (verification not implemented)**

Time = 3.34 (sec) , antiderivative size = 152, normalized size of antiderivative = 4.00

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{8 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + bx \right)}{3b^2 \left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) \right)^2}$$

input `int(x/atanh(tanh(a + b*x))^(5/2), x)`output `-(8*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)) + b*x))/(3*b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^2)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int \frac{x}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))}(-2\operatorname{atanh}(\tanh(bx + a)) - bx)}{3\operatorname{atanh}(\tanh(bx + a))^2 b^2}$$

input `int(x/atanh(tanh(b*x+a))^(5/2), x)`output `(2*sqrt(atanh(tanh(a + b*x)))*(- 2*atanh(tanh(a + b*x)) - b*x))/(3*atanh(tanh(a + b*x))^2*b**2)`

$$3.162 \quad \int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$$

Optimal result	1186
Mathematica [A] (verified)	1186
Rubi [A] (verified)	1187
Maple [A] (verified)	1188
Fricas [B] (verification not implemented)	1188
Sympy [A] (verification not implemented)	1188
Maxima [A] (verification not implemented)	1189
Giac [A] (verification not implemented)	1189
Mupad [B] (verification not implemented)	1189
Reduce [B] (verification not implemented)	1190

### Optimal result

Integrand size = 11, antiderivative size = 18

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

output `-2/3/b/arctanh(tanh(b*x+a))^(3/2)`

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(-5/2), x]`

output `-2/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2))`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

$$\downarrow \text{2588}$$

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} d\operatorname{arctanh}(\tanh(a + bx))$$

$$\downarrow \text{15}$$

$$-\frac{2}{3b\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(-5/2), x]`

output `-2/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2))`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2}{3b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$	15
default	$-\frac{2}{3b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$	15

input `int(1/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/3/b/arctanh(tanh(b*x+a))^(3/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2\sqrt{bx+a}}{3(b^3x^2 + 2ab^2x + a^2b)}$$

input `integrate(1/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `-2/3*sqrt(b*x + a)/(b^3*x^2 + 2*a*b^2*x + a^2*b)`

**Sympy [A] (verification not implemented)**

Time = 2.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.50

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \begin{cases} -\frac{2}{3b \operatorname{atanh}^{\frac{3}{2}}(\tanh(a+bx))} & \text{for } b \neq 0 \\ \frac{x}{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a))} & \text{otherwise} \end{cases}$$

input `integrate(1/atanh(tanh(b*x+a))**(5/2),x)`

output `Piecewise((-2/(3*b*atanh(tanh(a + b*x))**(3/2)), Ne(b, 0)), (x/atanh(tanh(a))**(5/2), True))`

### Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2}{3(bx + a)^{3/2}b}$$

input `integrate(1/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `-2/3/((b*x + a)^(3/2)*b)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.67

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2}{3(bx + a)^{3/2}b}$$

input `integrate(1/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-2/3/((b*x + a)^(3/2)*b)`

### Mupad [B] (verification not implemented)

Time = 3.36 (sec) , antiderivative size = 103, normalized size of antiderivative = 5.72

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{8 \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{3b \left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) \right)^2}$$

input `int(1/atanh(tanh(a + b*x))^(5/2),x)`

output `-(8*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))^2)`

### Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{3\operatorname{atanh}(\tanh(bx + a))^2 b}$$

input `int(1/atanh(tanh(b*x+a))^(5/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x)))/3*atanh(tanh(a + b*x))**2*b)`

### 3.163 $\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1191
Mathematica [A] (verified)	1192
Rubi [A] (verified)	1192
Maple [A] (verified)	1194
Fricas [A] (verification not implemented)	1194
Sympy [F]	1195
Maxima [F]	1195
Giac [A] (verification not implemented)	1195
Mupad [B] (verification not implemented)	1196
Reduce [F]	1196

#### Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} - \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2} + \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
2*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x
-arctanh(tanh(b*x+a)))^(5/2)-2/3/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b
*x+a))^(3/2)+2/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.84

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} + \frac{2(-bx + 4 \operatorname{arctanh}(\tanh(a + bx)))}{3 \operatorname{arctanh}(\tanh(a + bx))^{3/2} (-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Integrate[1/(x*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(-2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]]^(5/2) + (2*(-(b*x) + 4*ArcTanh[Tanh[a + b*x]]))/(3*ArcTanh[Tanh[a + b*x]]^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.17, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2594

$$-\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2594

$$\begin{aligned}
 & \int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \\
 & - \frac{2}{bx - \operatorname{arctanh}(\tanh(a+bx)) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \frac{2}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow \text{2592} \\
 & - \frac{2 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \\
 & - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & - \frac{2}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}
 \end{aligned}$$

input `Int[1/(x*ArcTanh[Tanh[a + b*x]]^(5/2)), x]`

output `-((( -2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))`

### Defintions of rubi rules used

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.86

method	result
default	$\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{2}{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^{3/2}} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a}}{\sqrt{a}}\right)}{(\operatorname{arctanh}(t$

input `int(1/x/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `2/(arctanh(tanh(b*x+a))-b*x)^2/arctanh(tanh(b*x+a))^(1/2)+2/3/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-2/(arctanh(tanh(b*x+a))-b*x)^(5/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.61

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \left[ \frac{3(b^2x^2 + 2abx + a^2)\sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(3abx + 4a^2)\sqrt{bx+a}}{3(a^3b^2x^2 + 2a^4bx + a^5)} \right]$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `[1/3*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(3*a*b*x + 4*a^2)*sqrt(b*x + a))/(a^3*b^2*x^2 + 2*a^4*b*x + a^5), 2/3*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b*x + 4*a^2)*sqrt(b*x + a))/(a^3*b^2*x^2 + 2*a^4*b*x + a^5)]`

**Sympy [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x \operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(1/(x*atanh(tanh(a + b*x))**(5/2)), x)`

**Maxima [F]**

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x \operatorname{artanh}(\tanh(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(x*arctanh(tanh(b*x + a))^(5/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.42

$$\int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}a^2} + \frac{2(3bx + 4a)}{3(bx + a)^{\frac{3}{2}}a^2}$$

input `integrate(1/x/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `2*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) + 2/3*(3*b*x + 4*a)/((b*x + a)^(3/2)*a^2)`



**Mupad [B] (verification not implemented)**

Time = 8.05 (sec) , antiderivative size = 886, normalized size of antiderivative = 8.20

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(1/(x*atanh(tanh(a + b*x))^(5/2)),x)`

output

```
(2^(1/2)*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 -
log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)
)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)
*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x)*((2*a - log((2*exp(
2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) +
1)) + 2*b*x)^5 + 40*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(
2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 80*a^3*(2*a -
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)^2 - 32*a^5 - 10*a*(2*a - log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x
)^4 + 80*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
+ log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*1i)/(2*x*(log(2/(exp(2*a)*ex
p(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) +
2*b*x)^(1/2)))*4i)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2) + (16*(log((2*exp(2*a)*
exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)
)/2)^(1/2))/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log
(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) - (16*(log(...
```

**Reduce [F]**

$$\int \frac{1}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x} dx$$

input `int(1/x/atanh(tanh(b*x+a))^(5/2),x)`

output `int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**3*x),x)`

**3.164**  $\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1198
Mathematica [A] (verified)	1199
Rubi [A] (verified)	1199
Maple [A] (verified)	1201
Fricas [A] (verification not implemented)	1202
Sympy [F]	1202
Maxima [F]	1203
Giac [A] (verification not implemented)	1203
Mupad [B] (verification not implemented)	1203
Reduce [F]	1204

**Optimal result**

Integrand size = 15, antiderivative size = 155

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{5b \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} + \frac{b}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} - \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5b} + \frac{5b}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
5*b*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(7/2)-1/x/arctanh(tanh(b*x+a))^(5/2)+b/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(5/2)-5/3*b/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(3/2)+5*b/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{5b \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{7/2}} + \frac{-2b^2x^2 + 14bx \operatorname{arctanh}(\tanh(a + bx)) + 3 \operatorname{arctanh}(\tanh(a + bx))^2}{3x(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input

```
Integrate[1/(x^2*ArcTanh[Tanh[a + b*x]]^(5/2)),x]
```

output

```
(5*b*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(7/2) + (-2*b^2*x^2 + 14*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2)/(3*x*(b*x - ArcTanh[Tanh[a + b*x]])^3*ArcTanh[Tanh[a + b*x]]^(3/2))
```

**Rubi [A] (verified)**

Time = 0.69 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.26, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2599

$$-\frac{5}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{7/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a + bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2592

$$-\frac{5}{2}b \left( -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

input

`Int [1/(x^2*ArcTanh[Tanh[a + b*x]]^(5/2)), x]`

output

```
(-5*b*(-((-(2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2))))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(5/2))
```

**Defintions of rubi rules used**

rule 2592

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

method	result
default	$2b \left( -\frac{1}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{\dots} \right)$

input `int(1/x^2/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `2*b*(-1/3/(arctanh(tanh(b*x+a))-b*x)^2/arctanh(tanh(b*x+a))^(3/2)-2/(arctanh(tanh(b*x+a))-b*x)^3/arctanh(tanh(b*x+a))^(1/2)-1/(arctanh(tanh(b*x+a))-b*x)^3*(1/2*arctanh(tanh(b*x+a))^(1/2)/b/x-5/2/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.41

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \left[ \frac{15(b^3x^3 + 2ab^2x^2 + a^2bx)\sqrt{a} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(15ab^2x^2 + 15a^2bx + 3a^3)\sqrt{bx+a}}{6(a^4b^2x^3 + 2a^5bx^2 + a^6x)} - \frac{15(b^3x^3 + 2ab^2x^2 + a^2bx)\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (15ab^2x^2 + 20a^2bx + 3a^3)\sqrt{bx+a}}{3(a^4b^2x^3 + 2a^5bx^2 + a^6x)} \right]$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `[1/6*(15*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*sqrt(a)*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(15*a*b^2*x^2 + 20*a^2*b*x + 3*a^3)*sqrt(b*x + a))/(a^4*b^2*x^3 + 2*a^5*b*x^2 + a^6*x), -1/3*(15*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (15*a*b^2*x^2 + 20*a^2*b*x + 3*a^3)*sqrt(b*x + a))/(a^4*b^2*x^3 + 2*a^5*b*x^2 + a^6*x)]`

### Sympy [F]

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^2 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**2/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(1/(x**2*atanh(tanh(a + b*x))**(5/2)), x)`

### Maxima [F]

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(x^2*arctanh(tanh(b*x + a))^(5/2)), x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{5b \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}a^3} - \frac{2(6(bx+a)b+ab)}{3(bx+a)^{3/2}a^3} - \frac{\sqrt{bx+a}}{a^3x}$$

input `integrate(1/x^2/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-5*b*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^3) - 2/3*(6*(b*x + a)*b + a*b)/((b*x + a)^(3/2)*a^3) - sqrt(b*x + a)/(a^3*x)`

### Mupad [B] (verification not implemented)

Time = 8.82 (sec) , antiderivative size = 1230, normalized size of antiderivative = 7.94

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(1/(x^2*atanh(tanh(a + b*x))^(5/2)),x)`



output

```
(2^(1/2)*b*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2
- log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) +
1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/
2)*2i - 2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*
b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 2^(1/2)*b*x*((2*a - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x)^7 + 84*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*ex
p(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5 - 280*a^3*(2*
a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*
a)*exp(2*b*x) + 1)) + 2*b*x)^4 + 560*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x)
))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3
- 672*a^5*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) +
log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 128*a^7 - 14*a*(2*a - log((
2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*
b*x) + 1)) + 2*b*x)^6 + 448*a^6*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*
a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*1i)/(2*x*
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)^(1/2))) * 20i) / (log(2/(exp(2*a)*exp(2*b*x) + 1)) -
log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(7/2) - (
32*b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/...
```

**Reduce [F]**

$$\int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^2} dx$$

input

```
int(1/x^2/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**3*x**2),x)
```

**3.165**  $\int \frac{1}{x^3 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1205
Mathematica [A] (verified)	1206
Rubi [A] (verified)	1206
Maple [A] (verified)	1209
Fricas [A] (verification not implemented)	1210
Sympy [F]	1210
Maxima [F]	1211
Giac [A] (verification not implemented)	1211
Mupad [B] (verification not implemented)	1211
Reduce [F]	1212

**Optimal result**

Integrand size = 15, antiderivative size = 224

$$\int \frac{1}{x^3 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{35b^2 \arctan\left(\frac{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \mathbf{arctanh}(\tanh(a+bx))}}\right)}{4(bx - \mathbf{arctanh}(\tanh(a+bx)))^{9/2}}$$

$$+ \frac{5b}{4x \mathbf{arctanh}(\tanh(a+bx))^{7/2}}$$

$$- \frac{5b^2}{4(bx - \mathbf{arctanh}(\tanh(a+bx))) \mathbf{arctanh}(\tanh(a+bx))^{7/2}}$$

$$- \frac{1}{2x^2 \mathbf{arctanh}(\tanh(a+bx))^{5/2}}$$

$$+ \frac{7b^2}{4(bx - \mathbf{arctanh}(\tanh(a+bx)))^2 \mathbf{arctanh}(\tanh(a+bx))^{5/2}}$$

$$- \frac{35b^2}{12(bx - \mathbf{arctanh}(\tanh(a+bx)))^3 \mathbf{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{35b^2}{4(bx - \mathbf{arctanh}(\tanh(a+bx)))^4 \sqrt{\mathbf{arctanh}(\tanh(a+bx))}}$$

output

$$\begin{aligned} & 35/4*b^2*\arctan(\operatorname{arctanh}(\tanh(b*x+a))^{1/2}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{1/2}) \\ & )/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{9/2}+5/4*b/x/\operatorname{arctanh}(\tanh(b*x+a))^{7/2}-5/4 \\ & *b^2/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/\operatorname{arctanh}(\tanh(b*x+a))^{7/2}-1/2/x^2/\operatorname{arctanh} \\ & (\tanh(b*x+a))^{5/2}+7/4*b^2/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2/\operatorname{arctanh}(\tanh(b*x+ \\ & a))^{5/2}-35/12*b^2/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^3/\operatorname{arctanh}(\tanh(b*x+a))^{3/2} \\ & )+35/4*b^2/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^4/\operatorname{arctanh}(\tanh(b*x+a))^{1/2} \end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.59

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{35b^2 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(-bx+\operatorname{arctanh}(\tanh(a+bx)))^{9/2}} + \frac{-8b^3x^3+80b^2x^2\operatorname{arctanh}(\tanh(a+bx))+39bx\operatorname{arctanh}(\tanh(a+bx))^2-6\operatorname{arctanh}(\tanh(a+bx))^3}{12x^2\operatorname{arctanh}(\tanh(a+bx))^{3/2}(-bx+\operatorname{arctanh}(\tanh(a+bx)))^4}$$

input

`Integrate[1/(x^3*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output

$$\begin{aligned} & (-35*b^2*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[a \\ & + b*x]]]])/(4*(-(b*x) + ArcTanh[Tanh[a + b*x]])^{9/2}) + (-8*b^3*x^3 + 80 \\ & *b^2*x^2*ArcTanh[Tanh[a + b*x]] + 39*b*x*ArcTanh[Tanh[a + b*x]]^2 - 6*ArcT \\ & anh[Tanh[a + b*x]]^3)/(12*x^2*ArcTanh[Tanh[a + b*x]]^{3/2}*(-(b*x) + ArcTa \\ & nh[Tanh[a + b*x]])^4 \end{aligned}$$
**Rubi [A] (verified)**Time = 0.64 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.20, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {2599, 2599, 2594, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx \\
 & \quad \downarrow 2599 \\
 & -\frac{5}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow 2599 \\
 & -\frac{5}{4}b \left( -\frac{7}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{9/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} \right) - \\
 & \quad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow 2594 \\
 & -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{7(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{7/2}} \right) - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) \\
 & \quad \downarrow 2594 \\
 & -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \right) - \frac{1}{7(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right) \\
 & \quad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow 2594 \\
 & -\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{1/2}} \right) \right) \\
 & \quad \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow 2594
 \end{aligned}$$

$$-\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{\int \frac{1}{x\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))} \right) \right)$$

$$\frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2592

$$-\frac{5}{4}b \left( -\frac{7}{2}b \left( -\frac{2 \operatorname{arctan}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))} \right) \right)$$

$$\frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

input `Int[1/(x^3*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(-5*b*((-7*b*(-((-((-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]])]))/(b*x - ArcTanh[Tanh[a + b*x]]))^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(7*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(7/2)))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(7/2)))/4 - 1/(2*x^2*ArcTanh[Tanh[a + b*x]]^(5/2))`

**Defintions of rubi rules used**

```
rule 2592 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2594 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.70

method	result
default	$2b^2 \left( \frac{3}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{1}{3(\operatorname{arctanh}(\tanh(bx+a)) - bx)^3 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{11 \operatorname{arctanh}(\tanh(bx+a))}{\dots} \right)$

```
input int(1/x^3/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
2*b^2*(3/(arctanh(tanh(b*x+a))-b*x)^4/arctanh(tanh(b*x+a))^(1/2)+1/3/(arctanh(tanh(b*x+a))-b*x)^3/arctanh(tanh(b*x+a))^(3/2)+1/(arctanh(tanh(b*x+a))-b*x)^4*((11/8*arctanh(tanh(b*x+a))^(3/2)+(-13/8*arctanh(tanh(b*x+a))+13/8*b*x)*arctanh(tanh(b*x+a))^(1/2))/b^2/x^2-35/8/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \left[ \frac{105 (b^4 x^4 + 2 a b^3 x^3 + a^2 b^2 x^2) \sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2 (105 a b^3 x^4 + 140 a^2 b^2 x^3 + 21 a^3 b x^2 - 6 a^4) \sqrt{bx+a}}{24 (a^5 b^2 x^4 + 2 a^6 b x^3 + a^7 x^2)} \right]$$

input

```
integrate(1/x^3/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
[1/24*(105*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(105*a*b^3*x^3 + 140*a^2*b^2*x^2 + 21*a^3*b*x - 6*a^4)*sqrt(b*x + a))/(a^5*b^2*x^4 + 2*a^6*b*x^3 + a^7*x^2), 1/12*(105*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (105*a*b^3*x^3 + 140*a^2*b^2*x^2 + 21*a^3*b*x - 6*a^4)*sqrt(b*x + a))/(a^5*b^2*x^4 + 2*a^6*b*x^3 + a^7*x^2)]
```

**Sympy [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^3 \operatorname{atanh}^{5/2}(\tanh(a + bx))} dx$$

input

```
integrate(1/x**3/atanh(tanh(b*x+a))**(5/2),x)
```

output

```
Integral(1/(x**3*atanh(tanh(a + b*x))**(5/2)), x)
```

**Maxima [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^3 \operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(x^3*arctanh(tanh(b*x + a))^(5/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{35 b^2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{4 \sqrt{-a} a^4} + \frac{2(9(bx+a)b^2 + ab^2)}{3(bx+a)^{3/2} a^4} + \frac{11(bx+a)^{3/2} b^2 - 13 \sqrt{bx+a} a b^2}{4 a^4 b^2 x^2}$$

input `integrate(1/x^3/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `35/4*b^2*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^4) + 2/3*(9*(b*x + a)*b^2 + a*b^2)/((b*x + a)^(3/2)*a^4) + 1/4*(11*(b*x + a)^(3/2)*b^2 - 13*sqrt(b*x + a)*a*b^2)/(a^4*b^2*x^2)`

**Mupad [B] (verification not implemented)**

Time = 9.22 (sec) , antiderivative size = 1514, normalized size of antiderivative = 6.76

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(1/(x^3*atanh(tanh(a + b*x))^(5/2)),x)`



output

```

((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^(1/2)*((4*(2*b*(2*log(2/(exp(2*a)*exp(2*b*x) + 1))
- 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x - 7*b*
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)))/(3*(2*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x)
)))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))
*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
*exp(2*b*x) + 1)) + 2*b*x)) + (56*b^2*x)/(3*(2*a*b - b*(2*a - log((2*exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) +
1)) + 2*b*x))*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(2*b*x^2 - x*(log(2/(exp(2*a)*e
xp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) +
2*b*x))^2 - ((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 -
log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*((140*b)/(3*(log(2/(exp(2*a)*exp
(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2
*b*x)^3 - (280*b^2*x)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)
*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4))/(2*b*x^2 - x*(log(2/(
exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*
x) + 1)) + 2*b*x)) + (2^(1/2)*b^2*log(((log((2*exp(2*a)*exp(2*b*x))/(exp(
2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(1o...

```

**Reduce [F]**

$$\int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^3} dx$$

input

```
int(1/x^3/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**3*x**3),x)
```

**3.166**  $\int \frac{1}{x^4 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1213
Mathematica [A] (verified)	1214
Rubi [A] (verified)	1215
Maple [A] (verified)	1218
Fricas [A] (verification not implemented)	1219
Sympy [F]	1219
Maxima [F]	1220
Giac [A] (verification not implemented)	1220
Mupad [B] (verification not implemented)	1220
Reduce [F]	1221

**Optimal result**

Integrand size = 15, antiderivative size = 278

$$\int \frac{1}{x^4 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{105b^3 \arctan\left(\frac{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}{\sqrt{bx - \mathbf{arctanh}(\tanh(a+bx))}}\right)}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^{11/2}} - \frac{24x \mathbf{arctanh}(\tanh(a+bx))^{9/2}}{35b^2} + \frac{24(bx - \mathbf{arctanh}(\tanh(a+bx))) \mathbf{arctanh}(\tanh(a+bx))^{9/2}}{35b^3} + \frac{12x^2 \mathbf{arctanh}(\tanh(a+bx))^{7/2}}{5b} - \frac{15b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^2 \mathbf{arctanh}(\tanh(a+bx))^{7/2}} - \frac{1}{3x^3 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} + \frac{21b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^3 \mathbf{arctanh}(\tanh(a+bx))^{5/2}} - \frac{105b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^4 \mathbf{arctanh}(\tanh(a+bx))^{3/2}} + \frac{105b^3}{8(bx - \mathbf{arctanh}(\tanh(a+bx)))^5 \sqrt{\mathbf{arctanh}(\tanh(a+bx))}}$$

output

```
105/8*b^3*arctan(arctanh(tanh(b*x+a))^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/
(b*x-arctanh(tanh(b*x+a)))^(11/2)-35/24*b^2/x/arctanh(tanh(b*x+a))^(9/2)+
35/24*b^3/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(9/2)+5/12*b/x^2/
arctanh(tanh(b*x+a))^(7/2)-15/8*b^3/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(
tanh(b*x+a))^(7/2)-1/3/x^3/arctanh(tanh(b*x+a))^(5/2)+21/8*b^3/(b*x-arc
tanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(5/2)-35/8*b^3/(b*x-arctanh(tanh
(b*x+a)))^4/arctanh(tanh(b*x+a))^(3/2)+105/8*b^3/(b*x-arctanh(tanh(b*x+a))
)^5/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.54

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{1}{24} \left( \frac{315b^3 \operatorname{arctanh}\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{11/2}} \right)$$

$$+ \frac{-16b^4x^4 + 208b^3x^3 \operatorname{arctanh}(\tanh(a + bx)) + 165b^2x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 50bx \operatorname{arctanh}(\tanh(a + bx))}{x^3(bx - \operatorname{arctanh}(\tanh(a + bx)))^5 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input

```
Integrate[1/(x^4*ArcTanh[Tanh[a + b*x]]^(5/2)),x]
```

output

```
((315*b^3*ArcTanh[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[-(b*x) + ArcTanh[Tanh[
a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(11/2) + (-16*b^4*x^4 + 208
*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 165*b^2*x^2*ArcTanh[Tanh[a + b*x]]^2 - 5
0*b*x*ArcTanh[Tanh[a + b*x]]^3 + 8*ArcTanh[Tanh[a + b*x]]^4)/(x^3*(b*x - A
rcTanh[Tanh[a + b*x]])^5*ArcTanh[Tanh[a + b*x]]^(3/2))/24
```

**Rubi [A] (verified)**

Time = 0.76 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.22, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {2599, 2599, 2599, 2594, 2594, 2594, 2594, 2594, 2592}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx \\
 & \quad \downarrow \text{2599} \\
 & -\frac{5}{6}b \int \frac{1}{x^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx - \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow \text{2599} \\
 & -\frac{5}{6}b \left( -\frac{7}{4}b \int \frac{1}{x^2 \operatorname{arctanh}(\tanh(a+bx))^{9/2}} dx - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}} \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow \text{2599} \\
 & -\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{11/2}} dx - \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{9/2}} \right) - \frac{1}{2x^2 \operatorname{arctanh}(\tanh(a+bx))} \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow \text{2594} \\
 & -\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{9/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{9(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{9/2}} \right) \right) \right) - \\
 & \quad \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}} \\
 & \quad \downarrow \text{2594}
 \end{aligned}$$

$$-\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{7/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{7(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{7/2}} - \frac{1}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{5/2}} - \frac{1}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{\int \frac{1}{x \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{1}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2594

$$-\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{\int \frac{1}{x \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{1}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) \right) \right) \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

↓ 2592

$$-\frac{5}{6}b \left( -\frac{7}{4}b \left( -\frac{9}{2}b \left( -\frac{2 \arctan\left(\frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx-\operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{(bx-\operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{3(bx-\operatorname{arctanh}(\tanh(a+bx)))} - \frac{2}{bx-\operatorname{arctanh}(\tanh(a+bx))} - \frac{2}{bx-\operatorname{arctanh}(\tanh(a+bx))} \right) \right) \right) - \frac{1}{3x^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}$$

input `Int[1/(x^4*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(-5*b*((-7*b*((-9*b*(-((-((-((-2*ArcTan[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) - 2/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]])) - 2/(5*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(7*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(7/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) - 2/(9*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(9/2)))/2 - 1/(x*ArcTanh[Tanh[a + b*x]]^(9/2)))/4 - 1/(2*x^2*ArcTanh[Tanh[a + b*x]]^(7/2)))/6 - 1/(3*x^3*ArcTanh[Tanh[a + b*x]]^(5/2))`

**Defintions of rubi rules used**

rule 2592 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[2*(ArcTan[Sqrt[v]/Rt[(b*u - a*v)/a, 2]]/(a*Rt[(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && PosQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v)))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.76

method	result
default	$2b^3 \left( \frac{\frac{41 \operatorname{arctanh}(\tanh(bx+a))^{5/2}}{16} + \left( -\frac{35 \operatorname{arctanh}(\tanh(bx+a))}{6} + \frac{35bx}{6} \right) \operatorname{arctanh}(\tanh(bx+a))^{3/2} + \left( \frac{55a^2}{16} + \frac{55a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{8} \right)}{b^3 x^3} \right) (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)$

input `int(1/x^4/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)`

output `2*b^3*(-1/(arctanh(tanh(b*x+a))-b*x)^5*((41/16*arctanh(tanh(b*x+a))^(5/2)+(-35/6*arctanh(tanh(b*x+a))+35/6*b*x)*arctanh(tanh(b*x+a))^(3/2)+(55/16*a^2+55/8*a*(arctanh(tanh(b*x+a))-b*x-a)+55/16*(arctanh(tanh(b*x+a))-b*x-a)^2)*arctanh(tanh(b*x+a))^(1/2))/b^3/x^3-105/16/(arctanh(tanh(b*x+a))-b*x)^(1/2)*arctanh(arctanh(tanh(b*x+a))^(1/2)/(arctanh(tanh(b*x+a))-b*x)^(1/2)))-1/3/(arctanh(tanh(b*x+a))-b*x)^4/arctanh(tanh(b*x+a))^(3/2)-4/(arctanh(tanh(b*x+a))-b*x)^5/arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 274, normalized size of antiderivative = 0.99

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{315 (b^5 x^5 + 2 ab^4 x^4 + a^2 b^3 x^3) \sqrt{a} \log\left(\frac{bx + 2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2 (315 ab^4 x^4 + 420 a^2 b^3 x^3 + 63 a^3 b^2 x^2 - 18 a^4 bx + 8 a^5) \sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (315 ab^4 x^4 + 420 a^2 b^3 x^3 + 63 a^3 b^2 x^2 - 18 a^4 bx + 8 a^5) \sqrt{a}}{48 (a^6 b^2 x^5 + 2 a^7 b x^4 + a^8 x^3)}$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `[1/48*(315*(b^5*x^5 + 2*a*b^4*x^4 + a^2*b^3*x^3)*sqrt(a)*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*(315*a*b^4*x^4 + 420*a^2*b^3*x^3 + 63*a^3*b^2*x^2 - 18*a^4*b*x + 8*a^5)*sqrt(b*x + a))/(a^6*b^2*x^5 + 2*a^7*b*x^4 + a^8*x^3), -1/24*(315*(b^5*x^5 + 2*a*b^4*x^4 + a^2*b^3*x^3)*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + (315*a*b^4*x^4 + 420*a^2*b^3*x^3 + 63*a^3*b^2*x^2 - 18*a^4*b*x + 8*a^5)*sqrt(b*x + a))/(a^6*b^2*x^5 + 2*a^7*b*x^4 + a^8*x^3)]`

**Sympy [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^4 \operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**4/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(1/(x**4*atanh(tanh(a + b*x))**(5/2)), x)`



**Maxima [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{x^4 \operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(x^4*arctanh(tanh(b*x + a))^(5/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{105 b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{8 \sqrt{-a} a^5} - \frac{315 (bx + a)^4 b^3 - 840 (bx + a)^3 a b^3 + 693 (bx + a)^2 a^2 b^3 - 144 (bx + a) a^3 b^3 - 16 a^4 b^3}{24 \left( (bx + a)^{3/2} - \sqrt{bx + aa} \right)^3 a^5}$$

input `integrate(1/x^4/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-105/8*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^5) - 1/24*(315*(b*x + a)^4*b^3 - 840*(b*x + a)^3*a*b^3 + 693*(b*x + a)^2*a^2*b^3 - 144*(b*x + a)*a^3*b^3 - 16*a^4*b^3)/(((b*x + a)^(3/2) - sqrt(b*x + a)*a)^3*a^5)`

**Mupad [B] (verification not implemented)**

Time = 7.92 (sec) , antiderivative size = 2359, normalized size of antiderivative = 8.49

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Too large to display}$$

input `int(1/(x^4*atanh(tanh(a + b*x))^(5/2)),x)`

output

```
(8*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (((140*b^2)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 - (840*b^3*x)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(x*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) - ((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(x*(((1232*b^4)/(9*(2*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (80*b^3*(2*b*(2*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 4*b*x) - 7*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(9*(2*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4))*(log(2/(exp(2*a)*exp(2*b*x) + 1...))
```

**Reduce [F]**

$$\int \frac{1}{x^4 \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^4} dx$$

input

```
int(1/x^4/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int(sqrt(atanh(tanh(a + b*x)))/(atanh(tanh(a + b*x))**3*x**4),x)
```

### 3.167 $\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	1222
Mathematica [A] (verified)	1222
Rubi [A] (verified)	1223
Maple [A] (verified)	1224
Fricas [A] (verification not implemented)	1224
Sympy [F(-1)]	1224
Maxima [A] (verification not implemented)	1225
Giac [A] (verification not implemented)	1225
Mupad [B] (verification not implemented)	1225
Reduce [F]	1226

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{99}bx^{11/2} + \frac{2}{9}x^{9/2} \operatorname{arctanh}(\tanh(a + bx))$$

output `-4/99*b*x^(11/2)+2/9*x^(9/2)*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{99}x^{9/2}(-2bx + 11\operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[x^(7/2)*ArcTanh[Tanh[a + b*x]],x]`

output `(2*x^(9/2)*(-2*b*x + 11*ArcTanh[Tanh[a + b*x]]))/99`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{9} b \int x^{9/2} dx$$

$$\downarrow \text{15}$$

$$\frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{99} bx^{11/2}$$

input `Int[x^(7/2)*ArcTanh[Tanh[a + b*x]], x]`

output `(-4*b*x^(11/2))/99 + (2*x^(9/2)*ArcTanh[Tanh[a + b*x]])/9`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{4bx^{\frac{11}{2}}}{99} + \frac{2x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9}$
default	$-\frac{4bx^{\frac{11}{2}}}{99} + \frac{2x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9}$
risch	$\frac{2x^{\frac{9}{2}} \ln(e^{bx+a})}{9} - \frac{\left(11i\pi x^4 \operatorname{csgn}(ie^{2bx+2a})^3 + 11i\pi x^4 \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}(ie^{2bx+2a}) - 11i\pi x^4 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)\right) \operatorname{csgn}(ie^{bx+a})}{9}$

input `int(x^(7/2)*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `-4/99*b*x^(11/2)+2/9*x^(9/2)*arctanh(tanh(b*x+a))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{99} (9bx^5 + 11ax^4) \sqrt{x}$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `2/99*(9*b*x^5 + 11*a*x^4)*sqrt(x)`

**Sympy [F(-1)]**

Timed out.

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \text{Timed out}$$

input `integrate(x**(7/2)*atanh(tanh(b*x+a)),x)`

output Timed out

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{99} bx^{11/2} + \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `-4/99*b*x^(11/2) + 2/9*x^(9/2)*arctanh(tanh(b*x + a))`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{11} bx^{11/2} + \frac{2}{9} ax^{9/2}$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `2/11*b*x^(11/2) + 2/9*a*x^(9/2)`

### Mupad [B] (verification not implemented)

Time = 3.45 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^{9/2} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{9} - \frac{4bx^{11/2}}{99} - \frac{x^{9/2} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{9}$$

input `int(x^(7/2)*atanh(tanh(a + b*x)),x)`

output  $(x^{9/2} \log((\exp(2a) \exp(2bx)) / (\exp(2a) \exp(2bx) + 1))) / 9 - (4bx^{11/2}) / 99 - (x^{9/2} \log(1 / (\exp(2a) \exp(2bx) + 1))) / 9$

### Reduce [F]

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x^3 dx$$

input `int(x^(7/2)*atanh(tanh(b*x+a)),x)`

output `int(sqrt(x)*atanh(tanh(a + b*x))*x**3,x)`

### 3.168 $\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	1227
Mathematica [A] (verified)	1227
Rubi [A] (verified)	1228
Maple [A] (verified)	1229
Fricas [A] (verification not implemented)	1229
Sympy [A] (verification not implemented)	1229
Maxima [A] (verification not implemented)	1230
Giac [A] (verification not implemented)	1230
Mupad [B] (verification not implemented)	1230
Reduce [F]	1231

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{63}bx^{9/2} + \frac{2}{7}x^{7/2}\operatorname{arctanh}(\tanh(a + bx))$$

output `-4/63*b*x^(9/2)+2/7*x^(7/2)*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{63}x^{7/2}(-2bx + 9\operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[x^(5/2)*ArcTanh[Tanh[a + b*x]],x]`

output `(2*x^(7/2)*(-2*b*x + 9*ArcTanh[Tanh[a + b*x]]))/63`



**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{7} b \int x^{7/2} dx$$

$$\downarrow \text{15}$$

$$\frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{63} bx^{9/2}$$

input `Int[x^(5/2)*ArcTanh[Tanh[a + b*x]], x]`

output `(-4*b*x^(9/2))/63 + (2*x^(7/2)*ArcTanh[Tanh[a + b*x]])/7`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{4bx^{\frac{9}{2}}}{63} + \frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7}$
default	$-\frac{4bx^{\frac{9}{2}}}{63} + \frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7}$
risch	$\frac{2x^{\frac{7}{2}} \ln(e^{bx+a})}{7} - \frac{\left(-9i\pi x^3 \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + 9i\pi x^3 \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^3 + 9i\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)\right)}{7}$

input `int(x^(5/2)*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `-4/63*b*x^(9/2)+2/7*x^(7/2)*arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{63} (7bx^4 + 9ax^3) \sqrt{x}$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `2/63*(7*b*x^4 + 9*a*x^3)*sqrt(x)`**Sympy [A] (verification not implemented)**

Time = 11.96 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4bx^{\frac{9}{2}}}{63} + \frac{2x^{\frac{7}{2}} \operatorname{atanh}(\tanh(a + bx))}{7}$$

input `integrate(x**(5/2)*atanh(tanh(b*x+a)),x)`

output  $-4*b*x^{(9/2)}/63 + 2*x^{(7/2)}*atanh(tanh(a + b*x))/7$

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{63} bx^{9/2} + \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output  $-4/63*b*x^{(9/2)} + 2/7*x^{(7/2)}*arctanh(tanh(b*x + a))$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{9} bx^{9/2} + \frac{2}{7} ax^{7/2}$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a)),x, algorithm="giac")`

output  $2/9*b*x^{(9/2)} + 2/7*a*x^{(7/2)}$

### Mupad [B] (verification not implemented)

Time = 3.29 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^{7/2} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{7} - \frac{4bx^{9/2}}{63} - \frac{x^{7/2} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{7}$$

input `int(x^(5/2)*atanh(tanh(a + b*x)),x)`

output  $(x^{7/2} \log((\exp(2a) \exp(2bx)) / (\exp(2a) \exp(2bx) + 1))) / 7 - (4bx^{9/2}) / 63 - (x^{7/2} \log(1 / (\exp(2a) \exp(2bx) + 1))) / 7$

### Reduce [F]

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x^2 dx$$

input `int(x^(5/2)*atanh(tanh(b*x+a)),x)`

output `int(sqrt(x)*atanh(tanh(a + b*x))*x**2,x)`

### 3.169 $\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	1232
Mathematica [A] (verified)	1232
Rubi [A] (verified)	1233
Maple [A] (verified)	1234
Fricas [A] (verification not implemented)	1234
Sympy [A] (verification not implemented)	1234
Maxima [A] (verification not implemented)	1235
Giac [A] (verification not implemented)	1235
Mupad [B] (verification not implemented)	1235
Reduce [F]	1236

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{35}bx^{7/2} + \frac{2}{5}x^{5/2}\operatorname{arctanh}(\tanh(a + bx))$$

output `-4/35*b*x^(7/2)+2/5*x^(5/2)*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{35}x^{5/2}(-2bx + 7\operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[x^(3/2)*ArcTanh[Tanh[a + b*x]],x]`

output `(2*x^(5/2)*(-2*b*x + 7*ArcTanh[Tanh[a + b*x]]))/35`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{5} b \int x^{5/2} dx$$

$$\downarrow \text{15}$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{35} bx^{7/2}$$

input `Int[x^(3/2)*ArcTanh[Tanh[a + b*x]], x]`

output `(-4*b*x^(7/2))/35 + (2*x^(5/2)*ArcTanh[Tanh[a + b*x]])/5`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{4bx^{\frac{7}{2}}}{35} + \frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))}{5}$
default	$-\frac{4bx^{\frac{7}{2}}}{35} + \frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))}{5}$
risch	$\frac{2x^{\frac{5}{2}} \ln(e^{bx+a})}{5} - \frac{\left(7i\pi x^2 \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^3 + 7i\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) - 7i\pi x^2\right)}{5}$

input `int(x^(3/2)*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`output `-4/35*b*x^(7/2)+2/5*x^(5/2)*arctanh(tanh(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{35} (5bx^3 + 7ax^2) \sqrt{x}$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `2/35*(5*b*x^3 + 7*a*x^2)*sqrt(x)`**Sympy [A] (verification not implemented)**

Time = 1.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4bx^{\frac{7}{2}}}{35} + \frac{2x^{\frac{5}{2}} \operatorname{atanh}(\tanh(a + bx))}{5}$$

input `integrate(x**(3/2)*atanh(tanh(b*x+a)),x)`

output  $-4*b*x^{(7/2)}/35 + 2*x^{(5/2)}*atanh(tanh(a + b*x))/5$

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{35} bx^{7/2} + \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output  $-4/35*b*x^{(7/2)} + 2/5*x^{(5/2)}*arctanh(tanh(b*x + a))$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{7} bx^{7/2} + \frac{2}{5} ax^{5/2}$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a)),x, algorithm="giac")`

output  $2/7*b*x^{(7/2)} + 2/5*a*x^{(5/2)}$

### Mupad [B] (verification not implemented)

Time = 3.37 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^{5/2} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{5} - \frac{4bx^{7/2}}{35} - \frac{x^{5/2} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{5}$$

input `int(x^(3/2)*atanh(tanh(a + b*x)),x)`



output `(x^(5/2)*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/5 - (4*b*x^(7/2))/35 - (x^(5/2)*log(1/(exp(2*a)*exp(2*b*x) + 1)))/5`

### Reduce [F]

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x dx$$

input `int(x^(3/2)*atanh(tanh(b*x+a)),x)`

output `int(sqrt(x)*atanh(tanh(a + b*x))*x,x)`

### 3.170 $\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx$

Optimal result	1237
Mathematica [A] (verified)	1237
Rubi [A] (verified)	1238
Maple [A] (verified)	1239
Fricas [A] (verification not implemented)	1239
Sympy [F]	1239
Maxima [A] (verification not implemented)	1240
Giac [A] (verification not implemented)	1240
Mupad [B] (verification not implemented)	1240
Reduce [F]	1241

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{15}bx^{5/2} + \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx))$$

output `-4/15*b*x^(5/2)+2/3*x^(3/2)*arctanh(tanh(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{15}x^{3/2}(-2bx + 5\operatorname{arctanh}(\tanh(a + bx)))$$

input `Integrate[Sqrt[x]*ArcTanh[Tanh[a + b*x]],x]`

output `(2*x^(3/2)*(-2*b*x + 5*ArcTanh[Tanh[a + b*x]]))/15`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{3} b \int x^{3/2} dx$$

$$\downarrow \text{15}$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{15} bx^{5/2}$$

input `Int[Sqrt[x]*ArcTanh[Tanh[a + b*x]], x]`

output `(-4*b*x^(5/2))/15 + (2*x^(3/2)*ArcTanh[Tanh[a + b*x]])/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{4bx^{\frac{5}{2}}}{15} + \frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))}{3}$
default	$-\frac{4bx^{\frac{5}{2}}}{15} + \frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))}{3}$
risch	$\frac{2x^{\frac{3}{2}} \ln(e^{bx+a})}{3} - \frac{\left(5i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}(ie^{2bx+2a}) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) x + 5i\pi x \operatorname{csgn}(ie^{2bx+2a})^3 - 5i\pi x \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)\right)}{3}$

input `int(x^(1/2)*arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `-4/15*b*x^(5/2)+2/3*x^(3/2)*arctanh(tanh(b*x+a))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.59

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{15} (3bx^2 + 5ax) \sqrt{x}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `2/15*(3*b*x^2 + 5*a*x)*sqrt(x)`

**Sympy [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \int \sqrt{x} \operatorname{atanh}(\tanh(a + bx)) dx$$

input `integrate(x**(1/2)*atanh(tanh(b*x+a)),x)`

output `Integral(sqrt(x)*atanh(tanh(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = -\frac{4}{15} bx^{\frac{5}{2}} + \frac{2}{3} x^{\frac{3}{2}} \operatorname{artanh}(\tanh(bx + a))$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-4/15*b*x^(5/2) + 2/3*x^(3/2)*arctanh(tanh(b*x + a))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{2}{5} bx^{\frac{5}{2}} + \frac{2}{3} ax^{\frac{3}{2}}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a)),x, algorithm="giac")`output `2/5*b*x^(5/2) + 2/3*a*x^(3/2)`**Mupad [B] (verification not implemented)**

Time = 3.33 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \frac{x^{3/2} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{3} - \frac{4bx^{5/2}}{15} - \frac{x^{3/2} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{3}$$

input `int(x^(1/2)*atanh(tanh(a + b*x)),x)`output `(x^(3/2)*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/3 - (4*b*x^(5/2))/15 - (x^(3/2)*log(1/(exp(2*a)*exp(2*b*x) + 1)))/3`

**Reduce [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a)) dx$$

input `int(x^(1/2)*atanh(tanh(b*x+a)),x)`

output `int(sqrt(x)*atanh(tanh(a + b*x)),x)`

### 3.171 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx$

Optimal result	1242
Mathematica [A] (verified)	1242
Rubi [A] (verified)	1243
Maple [A] (verified)	1244
Fricas [A] (verification not implemented)	1244
Sympy [F]	1244
Maxima [A] (verification not implemented)	1245
Giac [A] (verification not implemented)	1245
Mupad [B] (verification not implemented)	1245
Reduce [F]	1246

#### Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx = -\frac{4}{3}bx^{3/2} + 2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))$$

output

```
-4/3*b*x^(3/2)+2*x^(1/2)*arctanh(tanh(b*x+a))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx = \frac{2}{3}\sqrt{x}(-2bx + 3\operatorname{arctanh}(\tanh(a+bx)))$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]/Sqrt[x], x]
```

output

```
(2*Sqrt[x]*(-2*b*x + 3*ArcTanh[Tanh[a + b*x]]))/3
```

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx$$

↓ 2599

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - 2b \int \sqrt{x} dx$$

↓ 15

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{3}bx^{3/2}$$

input `Int[ArcTanh[Tanh[a + b*x]]/Sqrt[x],x]`

output `(-4*b*x^(3/2))/3 + 2*Sqrt[x]*ArcTanh[Tanh[a + b*x]]`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`



**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

method	result
derivativedivides	$-\frac{4bx^{\frac{3}{2}}}{3} + 2\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))$
default	$-\frac{4bx^{\frac{3}{2}}}{3} + 2\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))$
risch	$2\sqrt{x} \ln(e^{bx+a}) - \frac{(-6i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 - 3i\pi \operatorname{csgn}(\frac{i}{e^{2bx+2a}+1}) \operatorname{csgn}(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1})^2 + 3i\pi \operatorname{csgn}(\frac{i}{e^{2bx+2a}+1}) \operatorname{csgn}(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1})^2)}{e^{2bx+2a}+1}}$

input `int(arctanh(tanh(b*x+a))/x^(1/2),x,method=_RETURNVERBOSE)`

output `-4/3*b*x^(3/2)+2*x^(1/2)*arctanh(tanh(b*x+a))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx = \frac{2}{3} (bx+3a)\sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))/x^(1/2),x, algorithm="fricas")`

output `2/3*(b*x + 3*a)*sqrt(x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(a+bx))}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))/x**(1/2),x)`

output `Integral(atanh(tanh(a + b*x))/sqrt(x), x)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx = -\frac{4}{3} bx^{\frac{3}{2}} + 2\sqrt{x} \operatorname{arctanh}(\tanh(bx + a))$$

input `integrate(arctanh(tanh(b*x+a))/x^(1/2),x, algorithm="maxima")`

output `-4/3*b*x^(3/2) + 2*sqrt(x)*arctanh(tanh(b*x + a))`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx = \frac{2}{3} bx^{\frac{3}{2}} + 2a\sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))/x^(1/2),x, algorithm="giac")`

output `2/3*b*x^(3/2) + 2*a*sqrt(x)`

### Mupad [B] (verification not implemented)

Time = 3.37 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.24

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx = \sqrt{x} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \frac{4bx^{3/2}}{3} - \sqrt{x} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)$$

input `int(atanh(tanh(a + b*x))/x^(1/2),x)`

output  $x^{1/2} \log\left(\frac{\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) - (4bx^{3/2})/3 - x^{1/2} \log(1/(\exp(2a)\exp(2bx) + 1))$

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))}{\sqrt{x}} dx$$

input `int(atanh(tanh(b*x+a))/x^(1/2),x)`

output `int(atanh(tanh(a + b*x))/sqrt(x),x)`

### 3.172 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{3/2}} dx$

Optimal result	1247
Mathematica [A] (verified)	1247
Rubi [A] (verified)	1248
Maple [A] (verified)	1249
Fricas [A] (verification not implemented)	1249
Sympy [A] (verification not implemented)	1249
Maxima [A] (verification not implemented)	1250
Giac [A] (verification not implemented)	1250
Mupad [B] (verification not implemented)	1250
Reduce [B] (verification not implemented)	1251

#### Optimal result

Integrand size = 13, antiderivative size = 23

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{3/2}} dx = 4b\sqrt{x} - \frac{2\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}}$$

output `4*b*x^(1/2)-2*arctanh(tanh(b*x+a))/x^(1/2)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{3/2}} dx = \frac{4bx - 2\operatorname{arctanh}(\tanh(a+bx))}{\sqrt{x}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]/x^(3/2),x]`

output `(4*b*x - 2*ArcTanh[Tanh[a + b*x]])/Sqrt[x]`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx$$

$$\downarrow \text{2599}$$

$$2b \int \frac{1}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}}$$

$$\downarrow \text{15}$$

$$4b\sqrt{x} - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}}$$

input `Int[ArcTanh[Tanh[a + b*x]]/x^(3/2),x]`

output `4*b*Sqrt[x] - (2*ArcTanh[Tanh[a + b*x]])/Sqrt[x]`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
derivativedivides	$4b\sqrt{x} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}}$
default	$4b\sqrt{x} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}}$
risch	$-\frac{2 \ln(e^{bx+a})}{\sqrt{x}} + \frac{-2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 - i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 + i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}}\right)}{\sqrt{x}}$

input `int(arctanh(tanh(b*x+a))/x^(3/2),x,method=_RETURNVERBOSE)`output `4*b*x^(1/2)-2*arctanh(tanh(b*x+a))/x^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = \frac{2(bx - a)}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(3/2),x, algorithm="fricas")`output `2*(b*x - a)/sqrt(x)`**Sympy [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = 4b\sqrt{x} - \frac{2 \operatorname{atanh}(\tanh(a + bx))}{\sqrt{x}}$$

input `integrate(atanh(tanh(b*x+a))/x**(3/2),x)`

output  $4*b*\text{sqrt}(x) - 2*\text{atanh}(\tanh(a + b*x))/\text{sqrt}(x)$

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = 4b\sqrt{x} - \frac{2 \text{artanh}(\tanh(bx + a))}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(3/2),x, algorithm="maxima")`

output  $4*b*\text{sqrt}(x) - 2*\text{arctanh}(\tanh(b*x + a))/\text{sqrt}(x)$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = 2b\sqrt{x} - \frac{2a}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(3/2),x, algorithm="giac")`

output  $2*b*\text{sqrt}(x) - 2*a/\text{sqrt}(x)$

### Mupad [B] (verification not implemented)

Time = 3.35 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.43

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{\sqrt{x}} + 4b\sqrt{x} - \frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{\sqrt{x}}$$

input `int(atanh(tanh(a + b*x))/x^(3/2),x)`

output  $\log(1/(\exp(2*a)*\exp(2*b*x) + 1))/x^{(1/2)} + 4*b*x^{(1/2)} - \log((\exp(2*a)*\exp(2*b*x))/(\exp(2*a)*\exp(2*b*x) + 1))/x^{(1/2)}$

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx = \frac{-2a \operatorname{atanh}(\tanh(bx + a)) + 4bx}{\sqrt{x}}$$

input `int(atanh(tanh(b*x+a))/x^(3/2),x)`

output  $(2*(-\operatorname{atanh}(\tanh(a + b*x)) + 2*b*x))/\operatorname{sqrt}(x)$



### 3.173 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{5/2}} dx$

Optimal result	1252
Mathematica [A] (verified)	1252
Rubi [A] (verified)	1253
Maple [A] (verified)	1254
Fricas [A] (verification not implemented)	1254
Sympy [A] (verification not implemented)	1254
Maxima [A] (verification not implemented)	1255
Giac [A] (verification not implemented)	1255
Mupad [B] (verification not implemented)	1255
Reduce [B] (verification not implemented)	1256

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{5/2}} dx = -\frac{4b}{3\sqrt{x}} - \frac{2\operatorname{arctanh}(\tanh(a+bx))}{3x^{3/2}}$$

output  $-4/3*b/x^{(1/2)}-2/3*\operatorname{arctanh}(\tanh(b*x+a))/x^{(3/2)}$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{5/2}} dx = -\frac{2(2bx + \operatorname{arctanh}(\tanh(a+bx)))}{3x^{3/2}}$$

input  $\operatorname{Integrate}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]/x^{(5/2)}, x]$

output  $(-2*(2*b*x + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]))/(3*x^{(3/2)})$

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(ax + b))}{x^{5/2}} dx$$

↓ 2599

$$\frac{2}{3}b \int \frac{1}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(ax + b))}{3x^{3/2}}$$

↓ 15

$$-\frac{2\operatorname{arctanh}(\tanh(ax + b))}{3x^{3/2}} - \frac{4b}{3\sqrt{x}}$$

input `Int[ArcTanh[Tanh[a + b*x]]/x^(5/2),x]`

output `(-4*b)/(3*Sqrt[x]) - (2*ArcTanh[Tanh[a + b*x]])/(3*x^(3/2))`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{4b}{3\sqrt{x}} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{3x^{\frac{3}{2}}}$
default	$-\frac{4b}{3\sqrt{x}} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{3x^{\frac{3}{2}}}$
risch	$-\frac{2 \ln(e^{bx+a})}{3x^{\frac{3}{2}}} - \frac{2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 + i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 - i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)}{3x^{\frac{3}{2}}}$

input `int(arctanh(tanh(b*x+a))/x^(5/2),x,method=_RETURNVERBOSE)`

output `-4/3*b/x^(1/2)-2/3*arctanh(tanh(b*x+a))/x^(3/2)`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = -\frac{2(3bx + a)}{3x^{\frac{3}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(5/2),x, algorithm="fricas")`

output `-2/3*(3*b*x + a)/x^(3/2)`

**Sympy [A] (verification not implemented)**

Time = 1.29 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = -\frac{4b}{3\sqrt{x}} - \frac{2 \operatorname{atanh}(\tanh(a + bx))}{3x^{\frac{3}{2}}}$$

input `integrate(atanh(tanh(b*x+a))/x**(5/2),x)`

output  $-4*b/(3*\text{sqrt}(x)) - 2*\text{atanh}(\tanh(a + b*x))/(3*x**(3/2))$

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = -\frac{4b}{3\sqrt{x}} - \frac{2 \text{arctanh}(\tanh(bx + a))}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(5/2),x, algorithm="maxima")`

output  $-4/3*b/\text{sqrt}(x) - 2/3*\text{arctanh}(\tanh(b*x + a))/x^(3/2)$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.41

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = -\frac{2(3bx + a)}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(5/2),x, algorithm="giac")`

output  $-2/3*(3*b*x + a)/x^(3/2)$

### Mupad [B] (verification not implemented)

Time = 3.43 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.93

$$\int \frac{\text{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = -\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) + 4bx}{3x^{3/2}}$$

input `int(atanh(tanh(a + b*x))/x^(5/2),x)`

output  $-\left(\log\left(\frac{\exp(2a)\exp(2bx)}{\exp(2a)\exp(2bx) + 1}\right) - \log\left(\frac{1}{\exp(2a)\exp(2bx) + 1}\right) + 4bx\right)/(3x^{3/2})$

### Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{5/2}} dx = \frac{-\frac{2\operatorname{atanh}(\tanh(bx+a))}{3} - \frac{4bx}{3}}{\sqrt{x} x}$$

input `int(atanh(tanh(b*x+a))/x^(5/2),x)`

output  $(2*(-\operatorname{atanh}(\tanh(a + bx)) - 2bx))/(3\sqrt{x}x)$

### 3.174 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))}{x^{7/2}} dx$

Optimal result	1257
Mathematica [A] (verified)	1257
Rubi [A] (verified)	1258
Maple [A] (verified)	1259
Fricas [A] (verification not implemented)	1259
Sympy [A] (verification not implemented)	1259
Maxima [A] (verification not implemented)	1260
Giac [A] (verification not implemented)	1260
Mupad [B] (verification not implemented)	1260
Reduce [B] (verification not implemented)	1261

#### Optimal result

Integrand size = 13, antiderivative size = 27

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{4b}{15x^{3/2}} - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{5x^{5/2}}$$

output `-4/15*b/x^(3/2)-2/5*arctanh(tanh(b*x+a))/x^(5/2)`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{2(2bx + 3\operatorname{arctanh}(\tanh(a + bx)))}{15x^{5/2}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]/x^(7/2),x]`

output `(-2*(2*b*x + 3*ArcTanh[Tanh[a + b*x]]))/(15*x^(5/2))`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx$$

$$\downarrow \text{2599}$$

$$\frac{2}{5}b \int \frac{1}{x^{5/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{5x^{5/2}}$$

$$\downarrow \text{15}$$

$$-\frac{2\operatorname{arctanh}(\tanh(a + bx))}{5x^{5/2}} - \frac{4b}{15x^{3/2}}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]/x^(7/2), x]
```

output

```
(-4*b)/(15*x^(3/2)) - (2*ArcTanh[Tanh[a + b*x]])/(5*x^(5/2))
```

**Defintions of rubi rules used**

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result
derivativdivides	$-\frac{4b}{15x^{\frac{3}{2}}} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{5x^{\frac{5}{2}}}$
default	$-\frac{4b}{15x^{\frac{3}{2}}} - \frac{2 \operatorname{arctanh}(\tanh(bx+a))}{5x^{\frac{5}{2}}}$
risch	$-\frac{2 \ln(e^{bx+a})}{5x^{\frac{5}{2}}} - \frac{6i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 + 3i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)^2 - 3i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}+1}\right)}{5x^{\frac{5}{2}}}$

input `int(arctanh(tanh(b*x+a))/x^(7/2),x,method=_RETURNVERBOSE)`output `-4/15*b/x^(3/2)-2/5*arctanh(tanh(b*x+a))/x^(5/2)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{2(5bx + 3a)}{15x^{\frac{5}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(7/2),x, algorithm="fricas")`output `-2/15*(5*b*x + 3*a)/x^(5/2)`**Sympy [A] (verification not implemented)**

Time = 13.67 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{4b}{15x^{\frac{3}{2}}} - \frac{2 \operatorname{atanh}(\tanh(a + bx))}{5x^{\frac{5}{2}}}$$

input `integrate(atanh(tanh(b*x+a))/x**(7/2),x)`



output `-4*b/(15*x**(3/2)) - 2*atanh(tanh(a + b*x))/(5*x**(5/2))`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{4b}{15x^{3/2}} - \frac{2 \operatorname{arctanh}(\tanh(bx + a))}{5x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(7/2),x, algorithm="maxima")`

output `-4/15*b/x^(3/2) - 2/5*arctanh(tanh(b*x + a))/x^(5/2)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = -\frac{2(5bx + 3a)}{15x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))/x^(7/2),x, algorithm="giac")`

output `-2/15*(5*b*x + 3*a)/x^(5/2)`

### Mupad [B] (verification not implemented)

Time = 3.31 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = \frac{\ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{5x^{5/2}} - \frac{4b}{15x^{3/2}} - \frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{5x^{5/2}}$$

input `int(atanh(tanh(a + b*x))/x^(7/2),x)`

output

```
log(1/(exp(2*a)*exp(2*b*x) + 1))/(5*x^(5/2)) - (4*b)/(15*x^(3/2)) - log((e
xp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/(5*x^(5/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{7/2}} dx = \frac{-\frac{2 \operatorname{atanh}(\tanh(bx+a))}{5} - \frac{4bx}{15}}{\sqrt{x} x^2}$$

input

```
int(atanh(tanh(b*x+a))/x^(7/2),x)
```

output

```
(2*( - 3*atanh(tanh(a + b*x)) - 2*b*x))/(15*sqrt(x)*x**2)
```

### 3.175 $\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	1262
Mathematica [A] (verified)	1262
Rubi [A] (verified)	1263
Maple [A] (verified)	1264
Fricas [A] (verification not implemented)	1264
Sympy [F(-1)]	1265
Maxima [A] (verification not implemented)	1265
Giac [A] (verification not implemented)	1265
Mupad [B] (verification not implemented)	1266
Reduce [F]	1266

#### Optimal result

Integrand size = 15, antiderivative size = 48

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16b^2 x^{13/2}}{1287} - \frac{8}{99} b x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) + \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2$$

output `16/1287*b^2*x^(13/2)-8/99*b*x^(11/2)*arctanh(tanh(b*x+a))+2/9*x^(9/2)*arctanh(tanh(b*x+a))^2`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2x^{9/2}(8b^2x^2 - 52bx \operatorname{arctanh}(\tanh(a + bx)) + 143 \operatorname{arctanh}(\tanh(a + bx)))}{1287}$$

input `Integrate[x^(7/2)*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(9/2)*(8*b^2*x^2 - 52*b*x*ArcTanh[Tanh[a + b*x]] + 143*ArcTanh[Tanh[a + b*x]]^2))/1287`

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \int x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \left( \frac{2}{11} x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{11} b \int x^{11/2} dx \right)$$

$$\downarrow 15$$

$$\frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \left( \frac{2}{11} x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{143} b x^{13/2} \right)$$

input `Int[x^(7/2)*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(9/2)*ArcTanh[Tanh[a + b*x]]^2)/9 - (4*b*((-4*b*x^(13/2))/143 + (2*x^(11/2)*ArcTanh[Tanh[a + b*x]])/11))/9`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{2x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{9} - \frac{8b \left( \frac{x^{\frac{11}{2}} \operatorname{arctanh}(\tanh(bx+a))}{11} - \frac{2x^{\frac{13}{2}} b}{143} \right)}{9}$	38
default	$\frac{2x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{9} - \frac{8b \left( \frac{x^{\frac{11}{2}} \operatorname{arctanh}(\tanh(bx+a))}{11} - \frac{2x^{\frac{13}{2}} b}{143} \right)}{9}$	38
risch	Expression too large to display	2093

input

```
int(x^(7/2)*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
2/9*x^(9/2)*arctanh(tanh(b*x+a))^2-8/9*b*(1/11*x^(11/2)*arctanh(tanh(b*x+a)))-2/143*x^(13/2)*b)
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.60

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{1287} (99b^2x^6 + 234abx^5 + 143a^2x^4) \sqrt{x}$$

input

```
integrate(x^(7/2)*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
2/1287*(99*b^2*x^6 + 234*a*b*x^5 + 143*a^2*x^4)*sqrt(x)
```

**Sympy [F(-1)]**

Timed out.

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \text{Timed out}$$

input `integrate(x**(7/2)*atanh(tanh(b*x+a))**2,x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{1287} b^2 x^{\frac{13}{2}} - \frac{8}{99} b x^{\frac{11}{2}} \operatorname{arctanh}(\tanh(bx + a)) + \frac{2}{9} x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx + a))^2$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `16/1287*b^2*x^(13/2) - 8/99*b*x^(11/2)*arctanh(tanh(b*x + a)) + 2/9*x^(9/2)*arctanh(tanh(b*x + a))^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{13} b^2 x^{\frac{13}{2}} + \frac{4}{11} a b x^{\frac{11}{2}} + \frac{2}{9} a^2 x^{\frac{9}{2}}$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `2/13*b^2*x^(13/2) + 4/11*a*b*x^(11/2) + 2/9*a^2*x^(9/2)`

**Mupad [B] (verification not implemented)**

Time = 3.29 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x^{9/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{18} + \frac{2b^2 x^{13/2}}{13} - \frac{2bx^{11/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{11}$$

input `int(x^(7/2)*atanh(tanh(a + b*x))^2,x)`output `(x^(9/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/18 + (2*b^2*x^(13/2))/13 - (2*b*x^(11/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/11`**Reduce [F]**

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x^3 dx$$

input `int(x^(7/2)*atanh(tanh(b*x+a))^2,x)`output `int(sqrt(x)*atanh(tanh(a + b*x))*2*x**3,x)`

### 3.176 $\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	1267
Mathematica [A] (verified)	1267
Rubi [A] (verified)	1268
Maple [A] (verified)	1269
Fricas [A] (verification not implemented)	1269
Sympy [A] (verification not implemented)	1270
Maxima [A] (verification not implemented)	1270
Giac [A] (verification not implemented)	1271
Mupad [B] (verification not implemented)	1271
Reduce [F]	1272

#### Optimal result

Integrand size = 15, antiderivative size = 48

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{693} b^2 x^{11/2} - \frac{8}{63} b x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) + \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2$$

output `16/693*b^2*x^(11/2)-8/63*b*x^(9/2)*arctanh(tanh(b*x+a))+2/7*x^(7/2)*arctanh(tanh(b*x+a))^2`

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{693} x^{7/2} (8b^2 x^2 - 44bx \operatorname{arctanh}(\tanh(a + bx)) + 99 \operatorname{arctanh}(\tanh(a + bx))^2)$$

input `Integrate[x^(5/2)*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(7/2)*(8*b^2*x^2 - 44*b*x*ArcTanh[Tanh[a + b*x]] + 99*ArcTanh[Tanh[a + b*x]]^2))/693`



**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{9} b \int x^{9/2} dx \right)$$

$$\downarrow 15$$

$$\frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{99} b x^{11/2} \right)$$

input `Int[x^(5/2)*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(7/2)*ArcTanh[Tanh[a + b*x]]^2)/7 - (4*b*((-4*b*x^(11/2))/99 + (2*x^(9/2)*ArcTanh[Tanh[a + b*x]])/9))/7`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{7} - \frac{8b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9} - \frac{2bx^{\frac{11}{2}}}{99} \right)}{7}$	38
default	$\frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{7} - \frac{8b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9} - \frac{2bx^{\frac{11}{2}}}{99} \right)}{7}$	38
risch	Expression too large to display	2093

input

```
int(x^(5/2)*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
2/7*x^(7/2)*arctanh(tanh(b*x+a))^2-8/7*b*(1/9*x^(9/2)*arctanh(tanh(b*x+a))-2/99*b*x^(11/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.60

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{693} (63b^2x^5 + 154abx^4 + 99a^2x^3) \sqrt{x}$$

input

```
integrate(x^(5/2)*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
2/693*(63*b^2*x^5 + 154*a*b*x^4 + 99*a^2*x^3)*sqrt(x)
```

**Sympy [A] (verification not implemented)**

Time = 22.35 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16b^2 x^{11/2}}{693} - \frac{8bx^{9/2} \operatorname{atanh}(\tanh(a + bx))}{63} + \frac{2x^{7/2} \operatorname{atanh}^2(\tanh(a + bx))}{7}$$

input `integrate(x**(5/2)*atanh(tanh(b*x+a))**2,x)`output `16*b**2*x**(11/2)/693 - 8*b*x**(9/2)*atanh(tanh(a + b*x))/63 + 2*x**(7/2)*atanh(tanh(a + b*x))**2/7`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{693} b^2 x^{11/2} - \frac{8}{63} bx^{9/2} \operatorname{artanh}(\tanh(bx + a)) + \frac{2}{7} x^{7/2} \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `16/693*b^2*x^(11/2) - 8/63*b*x^(9/2)*arctanh(tanh(b*x + a)) + 2/7*x^(7/2)*arctanh(tanh(b*x + a))^2`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{11} b^2 x^{11/2} + \frac{4}{9} abx^{9/2} + \frac{2}{7} a^2 x^{7/2}$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `2/11*b^2*x^(11/2) + 4/9*a*b*x^(9/2) + 2/7*a^2*x^(7/2)`

**Mupad [B] (verification not implemented)**

Time = 3.22 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x^{7/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{14} + \frac{2b^2 x^{11/2}}{11} - \frac{2bx^{9/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{9}$$

input `int(x^(5/2)*atanh(tanh(a + b*x))^2,x)`

output `(x^(7/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/14 + (2*b^2*x^(11/2))/11 - (2*b*x^(9/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/9`

**Reduce [F]**

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x^2 dx$$

input `int(x^(5/2)*atanh(tanh(b*x+a))^2,x)`

output `int(sqrt(x)*atanh(tanh(a + b*x))**2*x**2,x)`

### 3.177 $\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	1273
Mathematica [A] (verified)	1273
Rubi [A] (verified)	1274
Maple [A] (verified)	1275
Fricas [A] (verification not implemented)	1275
Sympy [F]	1276
Maxima [A] (verification not implemented)	1276
Giac [A] (verification not implemented)	1276
Mupad [B] (verification not implemented)	1277
Reduce [F]	1277

#### Optimal result

Integrand size = 15, antiderivative size = 48

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{315} b^2 x^{9/2} - \frac{8}{35} b x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) + \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2$$

output

```
16/315*b^2*x^(9/2)-8/35*b*x^(7/2)*arctanh(tanh(b*x+a))+2/5*x^(5/2)*arctanh
(tanh(b*x+a))^2
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{315} x^{5/2} (8b^2 x^2 - 36bx \operatorname{arctanh}(\tanh(a + bx)) + 63 \operatorname{arctanh}(\tanh(a + bx))^2)$$

input

```
Integrate[x^(3/2)*ArcTanh[Tanh[a + b*x]]^2,x]
```

output

```
(2*x^(5/2)*(8*b^2*x^2 - 36*b*x*ArcTanh[Tanh[a + b*x]] + 63*ArcTanh[Tanh[a
+ b*x]]^2))/315
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{7} b \int x^{7/2} dx \right)$$

$$\downarrow 15$$

$$\frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{63} b x^{9/2} \right)$$

input `Int[x^(3/2)*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(5/2)*ArcTanh[Tanh[a + b*x]]^2)/5 - (4*b*((-4*b*x^(9/2))/63 + (2*x^(7/2)*ArcTanh[Tanh[a + b*x]])/7))/5`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```

Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))

```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{5} - \frac{8b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7} - \frac{2bx^{\frac{9}{2}}}{63} \right)}{5}$	38
default	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{5} - \frac{8b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7} - \frac{2bx^{\frac{9}{2}}}{63} \right)}{5}$	38
risch	Expression too large to display	2093

input

```
int(x^(3/2)*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
2/5*x^(5/2)*arctanh(tanh(b*x+a))^2-8/5*b*(1/7*x^(7/2)*arctanh(tanh(b*x+a))-2/63*b*x^(9/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.60

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{315} (35b^2x^4 + 90abx^3 + 63a^2x^2) \sqrt{x}$$

input

```
integrate(x^(3/2)*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
2/315*(35*b^2*x^4 + 90*a*b*x^3 + 63*a^2*x^2)*sqrt(x)
```



**Sympy [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int x^{3/2} \operatorname{atanh}^2(\tanh(a + bx)) dx$$

input `integrate(x**(3/2)*atanh(tanh(b*x+a))**2,x)`

output `Integral(x**(3/2)*atanh(tanh(a + b*x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{315} b^2 x^{9/2} - \frac{8}{35} b x^{7/2} \operatorname{artanh}(\tanh(bx + a)) + \frac{2}{5} x^{5/2} \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `16/315*b^2*x^(9/2) - 8/35*b*x^(7/2)*arctanh(tanh(b*x + a)) + 2/5*x^(5/2)*arctanh(tanh(b*x + a))^2`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{9} b^2 x^{9/2} + \frac{4}{7} a b x^{7/2} + \frac{2}{5} a^2 x^{5/2}$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `2/9*b^2*x^(9/2) + 4/7*a*b*x^(7/2) + 2/5*a^2*x^(5/2)`

**Mupad [B] (verification not implemented)**

Time = 3.16 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{10} + \frac{2b^2 x^{9/2}}{9} - \frac{2bx^{7/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{7}$$

input `int(x^(3/2)*atanh(tanh(a + b*x))^2,x)`output `(x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/10 + (2*b^2*x^(9/2))/9 - (2*b*x^(7/2))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/7`**Reduce [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x dx$$

input `int(x^(3/2)*atanh(tanh(b*x+a))^2,x)`output `int(sqrt(x)*atanh(tanh(a + b*x))*2*x,x)`

### 3.178 $\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx$

Optimal result	1278
Mathematica [A] (verified)	1278
Rubi [A] (verified)	1279
Maple [A] (verified)	1280
Fricas [A] (verification not implemented)	1280
Sympy [F]	1281
Maxima [A] (verification not implemented)	1281
Giac [A] (verification not implemented)	1281
Mupad [B] (verification not implemented)	1282
Reduce [F]	1282

#### Optimal result

Integrand size = 15, antiderivative size = 48

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{105} b^2 x^{7/2} - \frac{8}{15} b x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) + \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2$$

output

```
16/105*b^2*x^(7/2)-8/15*b*x^(5/2)*arctanh(tanh(b*x+a))+2/3*x^(3/2)*arctanh
(tanh(b*x+a))^2
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{105} x^{3/2} (8b^2 x^2 - 28bx \operatorname{arctanh}(\tanh(a + bx)) + 35 \operatorname{arctanh}(\tanh(a + bx))^2)$$

input

```
Integrate[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2,x]
```

output

```
(2*x^(3/2)*(8*b^2*x^2 - 28*b*x*ArcTanh[Tanh[a + b*x]] + 35*ArcTanh[Tanh[a
+ b*x]]^2))/105
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3} b \int x^{3/2} \operatorname{arctanh}(\tanh(a + bx)) dx$$

$$\downarrow 2599$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3} b \left( \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{5} b \int x^{5/2} dx \right)$$

$$\downarrow 15$$

$$\frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3} b \left( \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{35} b x^{7/2} \right)$$

input `Int[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(3/2)*ArcTanh[Tanh[a + b*x]]^2)/3 - (4*b*((-4*b*x^(7/2))/35 + (2*x^(5/2)*ArcTanh[Tanh[a + b*x]]/5)))/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{8b \left( \frac{x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))}{5} - \frac{2bx^{\frac{7}{2}}}{35} \right)}{3}$	38
default	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{3} - \frac{8b \left( \frac{x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))}{5} - \frac{2bx^{\frac{7}{2}}}{35} \right)}{3}$	38
risch	Expression too large to display	2027

input

```
int(x^(1/2)*arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
2/3*x^(3/2)*arctanh(tanh(b*x+a))^2-8/3*b*(1/5*x^(5/2)*arctanh(tanh(b*x+a))-2/35*b*x^(7/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.56

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{105} (15b^2x^3 + 42abx^2 + 35a^2x) \sqrt{x}$$

input

```
integrate(x^(1/2)*arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
2/105*(15*b^2*x^3 + 42*a*b*x^2 + 35*a^2*x)*sqrt(x)
```

**Sympy [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \sqrt{x} \operatorname{atanh}^2(\tanh(a + bx)) dx$$

input `integrate(x**(1/2)*atanh(tanh(b*x+a))**2,x)`

output `Integral(sqrt(x)*atanh(tanh(a + b*x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{16}{105} b^2 x^{\frac{7}{2}} - \frac{8}{15} b x^{\frac{5}{2}} \operatorname{artanh}(\tanh(bx + a)) + \frac{2}{3} x^{\frac{3}{2}} \operatorname{artanh}(\tanh(bx + a))^2$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `16/105*b^2*x^(7/2) - 8/15*b*x^(5/2)*arctanh(tanh(b*x + a)) + 2/3*x^(3/2)*arctanh(tanh(b*x + a))^2`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{2}{7} b^2 x^{\frac{7}{2}} + \frac{4}{5} a b x^{\frac{5}{2}} + \frac{2}{3} a^2 x^{\frac{3}{2}}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `2/7*b^2*x^(7/2) + 4/5*a*b*x^(5/2) + 2/3*a^2*x^(3/2)`

**Mupad [B] (verification not implemented)**

Time = 3.17 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \frac{x^{3/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{6} + \frac{2b^2 x^{7/2}}{7} - \frac{2bx^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{5}$$

input `int(x^(1/2)*atanh(tanh(a + b*x))^2,x)`output `(x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/6 + (2*b^2*x^(7/2))/7 - (2*b*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/5`**Reduce [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 dx$$

input `int(x^(1/2)*atanh(tanh(b*x+a))^2,x)`output `int(sqrt(x)*atanh(tanh(a + b*x))^2,x)`

$$3.179 \quad \int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx$$

Optimal result	1283
Mathematica [A] (verified)	1283
Rubi [A] (verified)	1284
Maple [A] (verified)	1285
Fricas [A] (verification not implemented)	1285
Sympy [F]	1286
Maxima [A] (verification not implemented)	1286
Giac [A] (verification not implemented)	1286
Mupad [B] (verification not implemented)	1287
Reduce [F]	1287

### Optimal result

Integrand size = 15, antiderivative size = 46

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx = \frac{16}{15}b^2x^{5/2} - \frac{8}{3}bx^{3/2}\operatorname{arctanh}(\tanh(a+bx)) + 2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^2$$

output `16/15*b^2*x^(5/2)-8/3*b*x^(3/2)*arctanh(tanh(b*x+a))+2*x^(1/2)*arctanh(tanh(b*x+a))^2`

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx = \frac{2}{15}\sqrt{x}(8b^2x^2 - 20bx\operatorname{arctanh}(\tanh(a+bx)) + 15\operatorname{arctanh}(\tanh(a+bx))^2)$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^2/Sqrt[x],x]`

output `(2*Sqrt[x]*(8*b^2*x^2 - 20*b*x*ArcTanh[Tanh[a + b*x]] + 15*ArcTanh[Tanh[a + b*x]]^2))/15`



**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx$$

↓ 2599

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2 - 4b \int \sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) dx$$

↓ 2599

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2 - 4b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{3}b \int x^{3/2} dx \right)$$

↓ 15

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2 - 4b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{15}bx^{5/2} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^2/Sqrt[x], x]`

output `2*Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2 - 4*b*((-4*b*x^(5/2))/15 + (2*x^(3/2)*ArcTanh[Tanh[a + b*x]])/3)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$\frac{2b^2x^{\frac{5}{2}}}{5} + \frac{4(\operatorname{arctanh}(\tanh(bx+a))-bx)bx^{\frac{3}{2}}}{3} + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2\sqrt{x}$	47
default	$\frac{2b^2x^{\frac{5}{2}}}{5} + \frac{4(\operatorname{arctanh}(\tanh(bx+a))-bx)bx^{\frac{3}{2}}}{3} + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx)^2\sqrt{x}$	47
risch	Expression too large to display	1978

input

```
int(arctanh(tanh(b*x+a))^2/x^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2/5*b^2*x^(5/2)+4/3*(arctanh(tanh(b*x+a))-b*x)*b*x^(3/2)+2*(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \frac{2}{15} (3b^2x^2 + 10abx + 15a^2)\sqrt{x}$$

input

```
integrate(arctanh(tanh(b*x+a))^2/x^(1/2),x, algorithm="fricas")
```

output

```
2/15*(3*b^2*x^2 + 10*a*b*x + 15*a^2)*sqrt(x)
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}^2(\tanh(a + bx))}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))**2/x**(1/2), x)`

output `Integral(atanh(tanh(a + b*x))**2/sqrt(x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \frac{16}{15} b^2 x^{\frac{5}{2}} - \frac{8}{3} b x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx + a)) + 2\sqrt{x} \operatorname{arctanh}(\tanh(bx + a))^2$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(1/2), x, algorithm="maxima")`

output `16/15*b^2*x^(5/2) - 8/3*b*x^(3/2)*arctanh(tanh(b*x + a)) + 2*sqrt(x)*arctanh(tanh(b*x + a))^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \frac{2}{5} b^2 x^{\frac{5}{2}} + \frac{4}{3} a b x^{\frac{3}{2}} + 2 a^2 \sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(1/2), x, algorithm="giac")`

output `2/5*b^2*x^(5/2) + 4/3*a*b*x^(3/2) + 2*a^2*sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 3.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.65

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \frac{\sqrt{x} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{2} + \frac{2b^2 x^{5/2}}{5} - \frac{2bx^{3/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{3}$$

input `int(atanh(tanh(a + b*x))^2/x^(1/2),x)`output `(x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 + (2*b^2*x^(5/2))/5 - (2*b*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/3`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^2}{\sqrt{x}} dx$$

input `int(atanh(tanh(b*x+a))^2/x^(1/2),x)`output `int(atanh(tanh(a + b*x))^2/sqrt(x),x)`

### 3.180 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{3/2}} dx$

Optimal result	1288
Mathematica [A] (verified)	1288
Rubi [A] (verified)	1289
Maple [A] (verified)	1290
Fricas [A] (verification not implemented)	1291
Sympy [F]	1291
Maxima [A] (verification not implemented)	1291
Giac [A] (verification not implemented)	1292
Mupad [B] (verification not implemented)	1292
Reduce [F]	1293

#### Optimal result

Integrand size = 15, antiderivative size = 44

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = -\frac{16}{3}b^2x^{3/2} + 8b\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}}$$

output

$$-16/3*b^2*x^(3/2)+8*b*x^(1/2)*\operatorname{arctanh}(\tanh(b*x+a))-2*\operatorname{arctanh}(\tanh(b*x+a))^2/x^(1/2)$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.91

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \frac{2(8b^2x^2 - 12bx\operatorname{arctanh}(\tanh(a + bx)) + 3\operatorname{arctanh}(\tanh(a + bx))^2)}{3\sqrt{x}}$$

input

$$\operatorname{Integrate}[\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2/x^(3/2), x]$$

output

$$\frac{(-2*(8*b^2*x^2 - 12*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2))/(3*Sqrt[x])}$$
**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx \\ & \quad \downarrow \text{2599} \\ & 4b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \\ & \quad \downarrow \text{2599} \\ & 4b \left( 2\sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) - 2b \int \sqrt{x} dx \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \\ & \quad \downarrow \text{15} \\ & 4b \left( 2\sqrt{x} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{3}bx^{3/2} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \end{aligned}$$

input

$$\text{Int}[ArcTanh[Tanh[a + b*x]]^2/x^(3/2), x]$$

output

$$\frac{(-2*ArcTanh[Tanh[a + b*x]]^2)/Sqrt[x] + 4*b*((-4*b*x^(3/2))/3 + 2*Sqrt[x]*ArcTanh[Tanh[a + b*x]])}$$

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{\sqrt{x}} + 8b \left( \sqrt{x} \operatorname{arctanh}(\tanh(bx+a)) - \frac{2bx^{\frac{3}{2}}}{3} \right)$	37
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{\sqrt{x}} + 8b \left( \sqrt{x} \operatorname{arctanh}(\tanh(bx+a)) - \frac{2bx^{\frac{3}{2}}}{3} \right)$	37
risch	Expression too large to display	1977

input `int(arctanh(tanh(b*x+a))^2/x^(3/2),x,method=_RETURNVERBOSE)`

output `-2*arctanh(tanh(b*x+a))^2/x^(1/2)+8*b*(x^(1/2)*arctanh(tanh(b*x+a))-2/3*b*x^(3/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \frac{2(b^2x^2 + 6abx - 3a^2)}{3\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(3/2),x, algorithm="fricas")`output `2/3*(b^2*x^2 + 6*a*b*x - 3*a^2)/sqrt(x)`**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \int \frac{\operatorname{atanh}^2(\tanh(a + bx))}{x^{3/2}} dx$$

input `integrate(atanh(tanh(b*x+a))**2/x**(3/2),x)`output `Integral(atanh(tanh(a + b*x))**2/x**(3/2), x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = -\frac{16}{3}b^2x^{3/2} + 8b\sqrt{x} \operatorname{artanh}(\tanh(bx + a)) - \frac{2 \operatorname{artanh}(\tanh(bx + a))^2}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(3/2),x, algorithm="maxima")`output `-16/3*b^2*x^(3/2) + 8*b*sqrt(x)*arctanh(tanh(b*x + a)) - 2*arctanh(tanh(b*x + a))^2/sqrt(x)`



**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \frac{2}{3} b^2 x^{3/2} + 4ab\sqrt{x} - \frac{2a^2}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(3/2),x, algorithm="giac")`

output `2/3*b^2*x^(3/2) + 4*a*b*sqrt(x) - 2*a^2/sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 3.06 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.77

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \frac{2b^2 x^{3/2}}{3} - \frac{\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)^2}{2\sqrt{x}} - 2b\sqrt{x} \left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)$$

input `int(atanh(tanh(a + b*x))^2/x^(3/2),x)`

output `(2*b^2*x^(3/2))/3 - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(2*x^(1/2)) - 2*b*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^2}{\sqrt{x} x} dx$$

input `int(atanh(tanh(b*x+a))^2/x^(3/2),x)`

output `int(atanh(tanh(a + b*x))**2/(sqrt(x)*x),x)`

**3.181**  $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{5/2}} dx$

Optimal result	1294
Mathematica [A] (verified)	1294
Rubi [A] (verified)	1295
Maple [A] (verified)	1296
Fricas [A] (verification not implemented)	1296
Sympy [A] (verification not implemented)	1297
Maxima [A] (verification not implemented)	1297
Giac [A] (verification not implemented)	1298
Mupad [B] (verification not implemented)	1298
Reduce [B] (verification not implemented)	1299

**Optimal result**

Integrand size = 15, antiderivative size = 48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{16b^2\sqrt{x}}{3} - \frac{8b\operatorname{arctanh}(\tanh(a + bx))}{3\sqrt{x}} - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}}$$

output `16/3*b^2*x^(1/2)-8/3*b*arctanh(tanh(b*x+a))/x^(1/2)-2/3*arctanh(tanh(b*x+a))^2/x^(3/2)`

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{2(8b^2x^2 - 4bx\operatorname{arctanh}(\tanh(a + bx)) - \operatorname{arctanh}(\tanh(a + bx))^2)}{3x^{3/2}}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^2/x^(5/2),x]`

output `(2*(8*b^2*x^2 - 4*b*x*ArcTanh[Tanh[a + b*x]] - ArcTanh[Tanh[a + b*x]]^2))/(3*x^(3/2))`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx$$

$$\downarrow 2599$$

$$\frac{4}{3}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}}$$

$$\downarrow 2599$$

$$\frac{4}{3}b \left( 2b \int \frac{1}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}}$$

$$\downarrow 15$$

$$\frac{4}{3}b \left( 4b\sqrt{x} - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}}$$

input `Int[ArcTanh[Tanh[a + b*x]]^2/x^(5/2), x]`

output `(-2*ArcTanh[Tanh[a + b*x]]^2)/(3*x^(3/2)) + (4*b*(4*b*Sqrt[x] - (2*ArcTanh[Tanh[a + b*x]])/Sqrt[x]))/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n])
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{3x^{\frac{3}{2}}} + \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}} + 2b\sqrt{x} \right)}{3}$	38
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{3x^{\frac{3}{2}}} + \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}} + 2b\sqrt{x} \right)}{3}$	38
risch	Expression too large to display	1972

input

```
int(arctanh(tanh(b*x+a))^2/x^(5/2), x, method=_RETURNVERBOSE)
```

output

```
-2/3*arctanh(tanh(b*x+a))^2/x^(3/2)+8/3*b*(-arctanh(tanh(b*x+a))/x^(1/2)+*b*x^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{2(3b^2x^2 - 6abx - a^2)}{3x^{\frac{3}{2}}}$$

input

```
integrate(arctanh(tanh(b*x+a))^2/x^(5/2), x, algorithm="fricas")
```

output

```
2/3*(3*b^2*x^2 - 6*a*b*x - a^2)/x^(3/2)
```

**Sympy [A] (verification not implemented)**

Time = 1.29 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{16b^2\sqrt{x}}{3} - \frac{8b \operatorname{artanh}(\tanh(a + bx))}{3\sqrt{x}} - \frac{2 \operatorname{artanh}^2(\tanh(a + bx))}{3x^{3/2}}$$

input `integrate(atanh(tanh(b*x+a))**2/x**(5/2), x)`

output `16*b**2*sqrt(x)/3 - 8*b*atanh(tanh(a + b*x))/(3*sqrt(x)) - 2*atanh(tanh(a + b*x))**2/(3*x**(3/2))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{16}{3} b^2 \sqrt{x} - \frac{8b \operatorname{artanh}(\tanh(bx + a))}{3\sqrt{x}} - \frac{2 \operatorname{artanh}(\tanh(bx + a))^2}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(5/2), x, algorithm="maxima")`

output `16/3*b^2*sqrt(x) - 8/3*b*arctanh(tanh(b*x + a))/sqrt(x) - 2/3*arctanh(tanh(b*x + a))^2/x^(3/2)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = 2b^2\sqrt{x} - \frac{2(6abx + a^2)}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(5/2),x, algorithm="giac")`

output `2*b^2*sqrt(x) - 2/3*(6*a*b*x + a^2)/x^(3/2)`

**Mupad [B] (verification not implemented)**

Time = 3.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = 2b^2\sqrt{x} - \frac{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{6x^{3/2}} + \frac{2b\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{\sqrt{x}}$$

input `int(atanh(tanh(a + b*x))^2/x^(5/2),x)`

output `2*b^2*x^(1/2) - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(6*x^(3/2)) + (2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/x^(1/2)`

**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx = \frac{-\frac{2\operatorname{atanh}(\tanh(bx+a))^2}{3} - \frac{8\operatorname{atanh}(\tanh(bx+a))bx}{3} + \frac{16b^2x^2}{3}}{\sqrt{x} x}$$

input `int(atanh(tanh(b*x+a))^2/x^(5/2),x)`output `(2*( - atanh(tanh(a + b*x))**2 - 4*atanh(tanh(a + b*x))*b*x + 8*b**2*x**2) )/(3*sqrt(x)*x)`



**3.182**  $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{x^{7/2}} dx$

Optimal result	1300
Mathematica [A] (verified)	1300
Rubi [A] (verified)	1301
Maple [A] (verified)	1302
Fricas [A] (verification not implemented)	1303
Sympy [A] (verification not implemented)	1303
Maxima [A] (verification not implemented)	1303
Giac [A] (verification not implemented)	1304
Mupad [B] (verification not implemented)	1304
Reduce [B] (verification not implemented)	1305

**Optimal result**

Integrand size = 15, antiderivative size = 48

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{16b^2}{15\sqrt{x}} - \frac{8b\operatorname{arctanh}(\tanh(a + bx))}{15x^{3/2}} - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{5x^{5/2}}$$

output

`-16/15*b^2/x^(1/2)-8/15*b*arctanh(tanh(b*x+a))/x^(3/2)-2/5*arctanh(tanh(b*x+a))^2/x^(5/2)`

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{2(8b^2x^2 + 4bx\operatorname{arctanh}(\tanh(a + bx)) + 3\operatorname{arctanh}(\tanh(a + bx))^2)}{15x^{5/2}}$$

input

`Integrate[ArcTanh[Tanh[a + b*x]]^2/x^(7/2), x]`

output

```
(-2*(8*b^2*x^2 + 4*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2
))/(15*x^(5/2))
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx$$

$$\downarrow 2599$$

$$\frac{4}{5}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{5/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{5x^{5/2}}$$

$$\downarrow 2599$$

$$\frac{4}{5}b \left( \frac{2}{3}b \int \frac{1}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{5x^{5/2}}$$

$$\downarrow 15$$

$$\frac{4}{5}b \left( -\frac{2\operatorname{arctanh}(\tanh(a + bx))}{3x^{3/2}} - \frac{4b}{3\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{5x^{5/2}}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^2/x^(7/2), x]
```

output

```
(-2*ArcTanh[Tanh[a + b*x]]^2)/(5*x^(5/2)) + (4*b*((-4*b)/(3*Sqrt[x]) - (2*
ArcTanh[Tanh[a + b*x]])/(3*x^(3/2))))/5
```

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{5x^{\frac{5}{2}}} + \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^{\frac{3}{2}}} - \frac{2b}{3\sqrt{x}} \right)}{5}$	38
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^2}{5x^{\frac{5}{2}}} + \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{3x^{\frac{3}{2}}} - \frac{2b}{3\sqrt{x}} \right)}{5}$	38
risch	Expression too large to display	1978

input `int(arctanh(tanh(b*x+a))^2/x^(7/2),x,method=_RETURNVERBOSE)`

output `-2/5*arctanh(tanh(b*x+a))^2/x^(5/2)+8/5*b*(-1/3*arctanh(tanh(b*x+a))/x^(3/2)-2/3*b/x^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{2(15b^2x^2 + 10abx + 3a^2)}{15x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(7/2),x, algorithm="fricas")`

output `-2/15*(15*b^2*x^2 + 10*a*b*x + 3*a^2)/x^(5/2)`

**Sympy [A] (verification not implemented)**

Time = 13.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{16b^2}{15\sqrt{x}} - \frac{8b \operatorname{atanh}(\tanh(a + bx))}{15x^{3/2}} - \frac{2 \operatorname{atanh}^2(\tanh(a + bx))}{5x^{5/2}}$$

input `integrate(atanh(tanh(b*x+a))**2/x**(7/2),x)`

output `-16*b**2/(15*sqrt(x)) - 8*b*atanh(tanh(a + b*x))/(15*x**(3/2)) - 2*atanh(atanh(a + b*x))**2/(5*x**(5/2))`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.75

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{16b^2}{15\sqrt{x}} - \frac{8b \operatorname{artanh}(\tanh(bx + a))}{15x^{3/2}} - \frac{2 \operatorname{artanh}(\tanh(bx + a))^2}{5x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^2/x^(7/2),x, algorithm="maxima")`

output

```
-16/15*b^2/sqrt(x) - 8/15*b*arctanh(tanh(b*x + a))/x^(3/2) - 2/5*arctanh(tanh(b*x + a))^2/x^(5/2)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.50

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = -\frac{2(15b^2x^2 + 10abx + 3a^2)}{15x^{5/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^2/x^(7/2),x, algorithm="giac")
```

output

```
-2/15*(15*b^2*x^2 + 10*a*b*x + 3*a^2)/x^(5/2)
```

**Mupad [B] (verification not implemented)**

Time = 3.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = \frac{2b \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{3x^{3/2}} - \frac{2b^2}{\sqrt{x}} - \frac{\left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{10x^{5/2}}$$

input

```
int(atanh(tanh(a + b*x))^2/x^(7/2),x)
```

output

```
(2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(3*x^(3/2)) - (2*b^2)/x^(1/2) - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(10*x^(5/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{7/2}} dx = \frac{-\frac{2\operatorname{atanh}(\tanh(bx+a))^2}{5} - \frac{8\operatorname{atanh}(\tanh(bx+a))bx}{15} - \frac{16b^2x^2}{15}}{\sqrt{x}x^2}$$

input `int(atanh(tanh(b*x+a))^2/x^(7/2),x)`output `(2*( - 3*atanh(tanh(a + b*x))**2 - 4*atanh(tanh(a + b*x))*b*x - 8*b**2*x**2))/(15*sqrt(x)*x**2)`

### 3.183 $\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	1306
Mathematica [A] (verified)	1306
Rubi [A] (verified)	1307
Maple [A] (verified)	1308
Fricas [A] (verification not implemented)	1309
Sympy [F(-1)]	1309
Maxima [A] (verification not implemented)	1309
Giac [A] (verification not implemented)	1310
Mupad [B] (verification not implemented)	1310
Reduce [F]	1311

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{32b^3 x^{15/2}}{6435} + \frac{16}{429} b^2 x^{13/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{33} b x^{11/2} \operatorname{arctanh}(\tanh(a + bx))^2 + \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^3$$

output

$$-32/6435*b^3*x^(15/2)+16/429*b^2*x^(13/2)*\operatorname{arctanh}(\tanh(b*x+a))-4/33*b*x^(11/2)*\operatorname{arctanh}(\tanh(b*x+a))^2+2/9*x^(9/2)*\operatorname{arctanh}(\tanh(b*x+a))^3$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2x^{9/2}(16b^3x^3 - 120b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 390bx \operatorname{arctanh}(\tanh(a + bx))^2 - 715 \operatorname{arctanh}(\tanh(a + bx))^3)}{6435}$$

input

$$\operatorname{Integrate}[x^{(7/2)}*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3,x]$$

output

$$\frac{(-2x^{9/2})(16b^3x^3 - 120b^2x^2\text{ArcTanh}[\text{Tanh}[a + bx]] + 390bxx\text{ArcTanh}[\text{Tanh}[a + bx]]^2 - 715\text{ArcTanh}[\text{Tanh}[a + bx]]^3)}{6435}$$
**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{7/2} \text{arctanh}(\tanh(a + bx))^3 dx \\ & \quad \downarrow \text{2599} \\ & \frac{2}{9} x^{9/2} \text{arctanh}(\tanh(a + bx))^3 - \frac{2}{3} b \int x^{9/2} \text{arctanh}(\tanh(a + bx))^2 dx \\ & \quad \downarrow \text{2599} \\ & \frac{2}{9} x^{9/2} \text{arctanh}(\tanh(a + bx))^3 - \\ & \frac{2}{3} b \left( \frac{2}{11} x^{11/2} \text{arctanh}(\tanh(a + bx))^2 - \frac{4}{11} b \int x^{11/2} \text{arctanh}(\tanh(a + bx)) dx \right) \\ & \quad \downarrow \text{2599} \\ & \frac{2}{9} x^{9/2} \text{arctanh}(\tanh(a + bx))^3 - \\ & \frac{2}{3} b \left( \frac{2}{11} x^{11/2} \text{arctanh}(\tanh(a + bx))^2 - \frac{4}{11} b \left( \frac{2}{13} x^{13/2} \text{arctanh}(\tanh(a + bx)) - \frac{2}{13} b \int x^{13/2} dx \right) \right) \\ & \quad \downarrow \text{15} \\ & \frac{2}{9} x^{9/2} \text{arctanh}(\tanh(a + bx))^3 - \\ & \frac{2}{3} b \left( \frac{2}{11} x^{11/2} \text{arctanh}(\tanh(a + bx))^2 - \frac{4}{11} b \left( \frac{2}{13} x^{13/2} \text{arctanh}(\tanh(a + bx)) - \frac{4}{195} bx^{15/2} \right) \right) \end{aligned}$$

input

$$\text{Int}[x^{(7/2)} \text{ArcTanh}[\text{Tanh}[a + bx]]^3, x]$$



output  $(2x^{9/2} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^3)/9 - (2b((2x^{11/2} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^2)/11 - (4b((-4bx^{15/2}))/195 + (2x^{13/2} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])/13))/11)/3$

**Defintions of rubi rules used**

rule 15  $\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a(x^{(m + 1)})/(m + 1), x] /; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$

rule 2599  $\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m + 1)}(v^n/(a(m + 1))), x] - \operatorname{Simp}[b(n/(a(m + 1))) \operatorname{Int}[u^{(m + 1)}v^{(n - 1)}, x], x] /; \operatorname{NeQ}[b u - a v, 0] /; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m + n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2n + m + 1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))]$

**Maple [A] (verified)**

Time = 1.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2x^{9/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{9} - \frac{4b \left( \frac{x^{11/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{11} - \frac{4b \left( \frac{x^{13/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))}{13} - \frac{2x^{15/2} b}{195} \right)}{11} \right)}{3}$	56
default	$\frac{2x^{9/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{9} - \frac{4b \left( \frac{x^{11/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{11} - \frac{4b \left( \frac{x^{13/2} \operatorname{arctanh}(\operatorname{tanh}(bx+a))}{13} - \frac{2x^{15/2} b}{195} \right)}{11} \right)}{3}$	56
risch	Expression too large to display	8179

input `int(x^(7/2)*arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output  $2/9*x^{(9/2)}*\operatorname{arctanh}(\tanh(b*x+a))^3-4/3*b*(1/11*x^{(11/2)}*\operatorname{arctanh}(\tanh(b*x+a))^2-4/11*b*(1/13*x^{(13/2)}*\operatorname{arctanh}(\tanh(b*x+a))-2/195*x^{(15/2)*b})$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.58

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = \frac{2}{6435} (429 b^3 x^7 + 1485 a b^2 x^6 + 1755 a^2 b x^5 + 715 a^3 x^4) \sqrt{x}$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output  $2/6435*(429*b^3*x^7 + 1485*a*b^2*x^6 + 1755*a^2*b*x^5 + 715*a^3*x^4)*\operatorname{sqrt}(x)$

### Sympy [F(-1)]

Timed out.

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = \text{Timed out}$$

input `integrate(x**(7/2)*atanh(tanh(b*x+a))**3,x)`

output Timed out

### Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.80

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = -\frac{4}{33} b x^{\frac{11}{2}} \operatorname{artanh}(\tanh(bx+a))^2 + \frac{2}{9} x^{\frac{9}{2}} \operatorname{artanh}(\tanh(bx+a))^3 - \frac{16}{6435} \left( 2 b^2 x^{\frac{15}{2}} - 15 b x^{\frac{13}{2}} \operatorname{artanh}(\tanh(bx+a)) \right) b$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `-4/33*b*x^(11/2)*arctanh(tanh(b*x + a))^2 + 2/9*x^(9/2)*arctanh(tanh(b*x + a))^3 - 16/6435*(2*b^2*x^(15/2) - 15*b*x^(13/2)*arctanh(tanh(b*x + a)))*b`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2}{15} b^3 x^{15/2} + \frac{6}{13} ab^2 x^{13/2} + \frac{6}{11} a^2 b x^{11/2} + \frac{2}{9} a^3 x^{9/2}$$

input `integrate(x^(7/2)*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `2/15*b^3*x^(15/2) + 6/13*a*b^2*x^(13/2) + 6/11*a^2*b*x^(11/2) + 2/9*a^3*x^(9/2)`

### Mupad [B] (verification not implemented)

Time = 3.13 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.64

$$\begin{aligned} \int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx &= \frac{2b^3 x^{15/2}}{15} \\ &- \frac{x^{9/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{36} \\ &+ \frac{3bx^{11/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{22} \\ &- \frac{3b^2 x^{13/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{13} \end{aligned}$$

input `int(x^(7/2)*atanh(tanh(a + b*x))^3,x)`

output

```
(2*b^3*x^(15/2))/15 - (x^(9/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/36 + (3*b*x^(1
1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(
2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/22 - (3*b^2*x^(13/2)*(log(2/(exp(2*a)*ex
p(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) +
2*b*x))/13
```

**Reduce [F]**

$$\int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x^3 dx$$

input

```
int(x^(7/2)*atanh(tanh(b*x+a))^3,x)
```

output

```
int(sqrt(x)*atanh(tanh(a + b*x))*3*x**3,x)
```

### 3.184 $\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	1312
Mathematica [A] (verified)	1312
Rubi [A] (verified)	1313
Maple [A] (verified)	1314
Fricas [A] (verification not implemented)	1315
Sympy [A] (verification not implemented)	1315
Maxima [A] (verification not implemented)	1315
Giac [A] (verification not implemented)	1316
Mupad [B] (verification not implemented)	1316
Reduce [F]	1317

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{32b^3 x^{13/2}}{3003} + \frac{16}{231} b^2 x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{21} b x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 + \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3$$

output 
$$-32/3003*b^3*x^{13/2}+16/231*b^2*x^{11/2}*\operatorname{arctanh}(\tanh(b*x+a))-4/21*b*x^{9/2}*\operatorname{arctanh}(\tanh(b*x+a))^2+2/7*x^{7/2}*\operatorname{arctanh}(\tanh(b*x+a))^3$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2x^{7/2}(-16b^3x^3 + 104b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) - 286bx \operatorname{arctanh}(\tanh(a + bx))) - 286b^3x^3 \operatorname{arctanh}(\tanh(a + bx))}{3003}$$

input 
$$\operatorname{Integrate}[x^{5/2}*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3,x]$$

output 
$$(2*x^{7/2}*(-16*b^3*x^3 + 104*b^2*x^2*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]] - 286*b*x*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2 + 429*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3))/3003$$

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx \\
 & \quad \downarrow 2599 \\
 & \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{6}{7} b \int x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx \\
 & \quad \downarrow 2599 \\
 & \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{6}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \int x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) dx \right) \\
 & \quad \downarrow 2599 \\
 & \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{6}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \left( \frac{2}{11} x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{11} b \int x^{11/2} dx \right) \right) \\
 & \quad \downarrow 15 \\
 & \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\
 & \frac{6}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{9} b \left( \frac{2}{11} x^{11/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{143} b x^{13/2} \right) \right)
 \end{aligned}$$

input `Int [x^(5/2)*ArcTanh [Tanh [a + b*x]]^3, x]`

output

`(2*x^(7/2)*ArcTanh [Tanh [a + b*x]]^3)/7 - (6*b*((2*x^(9/2)*ArcTanh [Tanh [a + b*x]]^2)/9 - (4*b*((-4*b*x^(13/2))/143 + (2*x^(11/2)*ArcTanh [Tanh [a + b*x]])/11))/9))/7`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 1.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{7} - \frac{12b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{9} - \frac{4b \left( \frac{x^{\frac{11}{2}} \operatorname{arctanh}(\tanh(bx+a))}{11} - \frac{2x^{\frac{13}{2}} b}{143} \right)}{9} \right)}{7}$	56
default	$\frac{2x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{7} - \frac{12b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{9} - \frac{4b \left( \frac{x^{\frac{11}{2}} \operatorname{arctanh}(\tanh(bx+a))}{11} - \frac{2x^{\frac{13}{2}} b}{143} \right)}{9} \right)}{7}$	56
risch	Expression too large to display	8179

input `int(x^(5/2)*arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `2/7*x^(7/2)*arctanh(tanh(b*x+a))^3-12/7*b*(1/9*x^(9/2)*arctanh(tanh(b*x+a))^2-4/9*b*(1/11*x^(11/2)*arctanh(tanh(b*x+a))-2/143*x^(13/2)*b)`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.58

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = \frac{2}{3003} (231 b^3 x^6 + 819 a b^2 x^5 + 1001 a^2 b x^4 + 429 a^3 x^3) \sqrt{x}$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`output `2/3003*(231*b^3*x^6 + 819*a*b^2*x^5 + 1001*a^2*b*x^4 + 429*a^3*x^3)*sqrt(x)`**Sympy [A] (verification not implemented)**

Time = 41.58 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = -\frac{32b^3 x^{13/2}}{3003} + \frac{16b^2 x^{11/2} \operatorname{atanh}(\tanh(a+bx))}{231} - \frac{4bx^{9/2} \operatorname{atanh}^2(\tanh(a+bx))}{21} + \frac{2x^{7/2} \operatorname{atanh}^3(\tanh(a+bx))}{7}$$

input `integrate(x**(5/2)*atanh(tanh(b*x+a))**3,x)`output `-32*b**3*x**(13/2)/3003 + 16*b**2*x**(11/2)*atanh(tanh(a + b*x))/231 - 4*b*x**(9/2)*atanh(tanh(a + b*x))**2/21 + 2*x**(7/2)*atanh(tanh(a + b*x))**3/7`**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.80

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3 dx = -\frac{4}{21} b x^{9/2} \operatorname{artanh}(\tanh(bx+a))^2 + \frac{2}{7} x^{7/2} \operatorname{artanh}(\tanh(bx+a))^3 - \frac{16}{3003} \left( 2b^2 x^{13/2} - 13bx^{11/2} \operatorname{artanh}(\tanh(bx+a)) \right) b$$



input `integrate(x^(5/2)*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output 
$$-4/21*b*x^{9/2}*arctanh(tanh(b*x + a))^2 + 2/7*x^{7/2}*arctanh(tanh(b*x + a))^3 - 16/3003*(2*b^2*x^{13/2} - 13*b*x^{11/2}*arctanh(tanh(b*x + a))) * b$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2}{13} b^3 x^{13/2} + \frac{6}{11} ab^2 x^{11/2} + \frac{2}{3} a^2 b x^{9/2} + \frac{2}{7} a^3 x^{7/2}$$

input `integrate(x^(5/2)*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output 
$$2/13*b^3*x^{13/2} + 6/11*a*b^2*x^{11/2} + 2/3*a^2*b*x^{9/2} + 2/7*a^3*x^{7/2}$$

### Mupad [B] (verification not implemented)

Time = 3.13 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.64

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2b^3 x^{13/2}}{13} - \frac{x^{7/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{28} + \frac{bx^{9/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{6} - \frac{3b^2 x^{11/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{11}$$

input `int(x^(5/2)*atanh(tanh(a + b*x))^3,x)`

output

```
(2*b^3*x^(13/2))/13 - (x^(7/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/28 + (b*x^(9/2)
)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)
)*exp(2*b*x) + 1)) + 2*b*x)^2)/6 - (3*b^2*x^(11/2)*(log(2/(exp(2*a)*exp(2*
b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*
x))/11
```

**Reduce [F]**

$$\int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x^2 dx$$

input

```
int(x^(5/2)*atanh(tanh(b*x+a))^3,x)
```

output

```
int(sqrt(x)*atanh(tanh(a + b*x))*3*x**2,x)
```

### 3.185 $\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	1318
Mathematica [A] (verified)	1318
Rubi [A] (verified)	1319
Maple [A] (verified)	1320
Fricas [A] (verification not implemented)	1321
Sympy [F]	1321
Maxima [A] (verification not implemented)	1321
Giac [A] (verification not implemented)	1322
Mupad [B] (verification not implemented)	1322
Reduce [F]	1323

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = -\frac{32b^3 x^{11/2}}{1155} + \frac{16}{105} b^2 x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{12}{35} b x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 + \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3$$

output

$$-32/1155*b^3*x^(11/2)+16/105*b^2*x^(9/2)*\operatorname{arctanh}(\tanh(b*x+a))-12/35*b*x^(7/2)*\operatorname{arctanh}(\tanh(b*x+a))^2+2/5*x^(5/2)*\operatorname{arctanh}(\tanh(b*x+a))^3$$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2x^{5/2}(16b^3x^3 - 88b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) + 198bx \operatorname{arctanh}(\tanh(a + bx))^2 - 231 \operatorname{arctanh}(\tanh(a + bx))^3)}{1155}$$

input

$$\operatorname{Integrate}[x^(3/2)*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3, x]$$

output

$$\frac{(-2x^{5/2}(16b^3x^3 - 88b^2x^2\text{ArcTanh}[\text{Tanh}[a + bx]] + 198bxx\text{ArcTanh}[\text{Tanh}[a + bx]]^2 - 231\text{ArcTanh}[\text{Tanh}[a + bx]]^3))/1155}$$
**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx \\ & \quad \downarrow 2599 \\ & \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \frac{6}{5} b \int x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx \\ & \quad \downarrow 2599 \\ & \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & \frac{6}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \int x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) dx \right) \\ & \quad \downarrow 2599 \\ & \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & \frac{6}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{9} b \int x^{9/2} dx \right) \right) \\ & \quad \downarrow 15 \\ & \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & \frac{6}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{7} b \left( \frac{2}{9} x^{9/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{99} bx^{11/2} \right) \right) \end{aligned}$$

input

$$\text{Int}[x^{(3/2)} \text{ArcTanh}[\text{Tanh}[a + bx]]^3, x]$$

```
output (2*x^(5/2)*ArcTanh[Tanh[a + b*x]]^3)/5 - (6*b*((2*x^(7/2)*ArcTanh[Tanh[a +
b*x]]^2)/7 - (4*b*((-4*b*x^(11/2))/99 + (2*x^(9/2)*ArcTanh[Tanh[a + b*x]]
)/9))/7))/5
```

**Defintions of rubi rules used**

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{5} - \frac{12b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{7} - \frac{4b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9} - \frac{2bx^{\frac{11}{2}}}{99} \right)}{7} \right)}{5}$	56
default	$\frac{2x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{5} - \frac{12b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{7} - \frac{4b \left( \frac{x^{\frac{9}{2}} \operatorname{arctanh}(\tanh(bx+a))}{9} - \frac{2bx^{\frac{11}{2}}}{99} \right)}{7} \right)}{5}$	56
risch	Expression too large to display	8179

```
input int(x^(3/2)*arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output  $2/5*x^{(5/2)}*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3-12/5*b*(1/7*x^{(7/2)}*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^2-4/7*b*(1/9*x^{(9/2)}*\operatorname{arctanh}(\operatorname{tanh}(b*x+a))-2/99*b*x^{(11/2)}))$

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.58

$$\int x^{3/2} \operatorname{arctanh}(\operatorname{tanh}(a+bx))^3 dx = \frac{2}{1155} (105 b^3 x^5 + 385 a b^2 x^4 + 495 a^2 b x^3 + 231 a^3 x^2) \sqrt{x}$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output  $2/1155*(105*b^3*x^5 + 385*a*b^2*x^4 + 495*a^2*b*x^3 + 231*a^3*x^2)*\operatorname{sqrt}(x)$

### Sympy [F]

$$\int x^{3/2} \operatorname{arctanh}(\operatorname{tanh}(a+bx))^3 dx = \int x^{3/2} \operatorname{atanh}^3(\operatorname{tanh}(a+bx)) dx$$

input `integrate(x**(3/2)*atanh(tanh(b*x+a))**3,x)`

output `Integral(x**(3/2)*atanh(tanh(a + b*x))**3, x)`

### Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.80

$$\int x^{3/2} \operatorname{arctanh}(\operatorname{tanh}(a+bx))^3 dx = -\frac{12}{35} b x^{7/2} \operatorname{artanh}(\operatorname{tanh}(bx+a))^2 + \frac{2}{5} x^{5/2} \operatorname{artanh}(\operatorname{tanh}(bx+a))^3 - \frac{16}{1155} \left( 2 b^2 x^{11/2} - 11 b x^{9/2} \operatorname{artanh}(\operatorname{tanh}(bx+a)) \right) b$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output

$$-12/35*b*x^{(7/2)}*arctanh(\tanh(b*x + a))^2 + 2/5*x^{(5/2)}*arctanh(\tanh(b*x + a))^3 - 16/1155*(2*b^2*x^{(11/2)} - 11*b*x^{(9/2)}*arctanh(\tanh(b*x + a)))*b$$
**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2}{11} b^3 x^{11/2} + \frac{2}{3} ab^2 x^{9/2} + \frac{6}{7} a^2 b x^{7/2} + \frac{2}{5} a^3 x^{5/2}$$

input

```
integrate(x^(3/2)*arctanh(tanh(b*x+a))^3,x, algorithm="giac")
```

output

$$2/11*b^3*x^{(11/2)} + 2/3*a*b^2*x^{(9/2)} + 6/7*a^2*b*x^{(7/2)} + 2/5*a^3*x^{(5/2)}$$
**Mupad [B] (verification not implemented)**

Time = 3.14 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.64

$$\begin{aligned} \int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx &= \frac{2b^3 x^{11/2}}{11} \\ &- \frac{x^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{20} \\ &+ \frac{3bx^{7/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{14} \\ &- \frac{b^2 x^{9/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{3} \end{aligned}$$

input

```
int(x^(3/2)*atanh(tanh(a + b*x))^3,x)
```

output

```
(2*b^3*x^(11/2))/11 - (x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/20 + (3*b*x^(7
/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2
*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/14 - (b^2*x^(9/2)*(log(2/(exp(2*a)*exp(2*
b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*
x))/3
```

**Reduce [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x dx$$

input

```
int(x^(3/2)*atanh(tanh(b*x+a))^3,x)
```

output

```
int(sqrt(x)*atanh(tanh(a + b*x))*3*x,x)
```



### 3.186 $\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx$

Optimal result	1324
Mathematica [A] (verified)	1324
Rubi [A] (verified)	1325
Maple [A] (verified)	1326
Fricas [A] (verification not implemented)	1327
Sympy [F]	1327
Maxima [A] (verification not implemented)	1327
Giac [A] (verification not implemented)	1328
Mupad [B] (verification not implemented)	1328
Reduce [F]	1329

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\begin{aligned} & \int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx \\ &= -\frac{32}{315} b^3 x^{9/2} + \frac{16}{35} b^2 x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) \\ & \quad - \frac{4}{5} b x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 + \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 \end{aligned}$$

output

```
-32/315*b^3*x^(9/2)+16/35*b^2*x^(7/2)*arctanh(tanh(b*x+a))-4/5*b*x^(5/2)*a
rctanh(tanh(b*x+a))^2+2/3*x^(3/2)*arctanh(tanh(b*x+a))^3
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\begin{aligned} \int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx &= -\frac{2}{315} x^{3/2} (16b^3 x^3 - 72b^2 x^2 \operatorname{arctanh}(\tanh(a + bx)) \\ & \quad + 126bx \operatorname{arctanh}(\tanh(a + bx))^2 \\ & \quad - 105 \operatorname{arctanh}(\tanh(a + bx))^3) \end{aligned}$$

input

```
Integrate[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^3,x]
```

output

$$\frac{(-2x^{3/2}(16b^3x^3 - 72b^2x^2\text{ArcTanh}[\text{Tanh}[a + bx]] + 126bxx\text{ArcTanh}[\text{Tanh}[a + bx]]^2 - 105\text{ArcTanh}[\text{Tanh}[a + bx]]^3))/315}$$
**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx \\ & \quad \downarrow 2599 \\ & \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 - 2b \int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2 dx \\ & \quad \downarrow 2599 \\ & \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & 2b \left( \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \int x^{5/2} \operatorname{arctanh}(\tanh(a + bx)) dx \right) \\ & \quad \downarrow 2599 \\ & \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & 2b \left( \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{7} b \int x^{7/2} dx \right) \right) \\ & \quad \downarrow 15 \\ & \frac{2}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3 - \\ & 2b \left( \frac{2}{5} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{5} b \left( \frac{2}{7} x^{7/2} \operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{63} bx^{9/2} \right) \right) \end{aligned}$$

input

$$\text{Int}[\text{Sqrt}[x] * \text{ArcTanh}[\text{Tanh}[a + b*x]]^3, x]$$

```
output (2*x^(3/2)*ArcTanh[Tanh[a + b*x]]^3)/3 - 2*b*((2*x^(5/2)*ArcTanh[Tanh[a +
b*x]]^2)/5 - (4*b*((-4*b*x^(9/2))/63 + (2*x^(7/2)*ArcTanh[Tanh[a + b*x]])/
7))/5)
```

**Defintions of rubi rules used**

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.98 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{3} - 4b \left( \frac{x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{5} - \frac{4b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7} - \frac{2bx^{\frac{9}{2}}}{63} \right)}{5} \right)$	56
default	$\frac{2x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^3}{3} - 4b \left( \frac{x^{\frac{5}{2}} \operatorname{arctanh}(\tanh(bx+a))^2}{5} - \frac{4b \left( \frac{x^{\frac{7}{2}} \operatorname{arctanh}(\tanh(bx+a))}{7} - \frac{2bx^{\frac{9}{2}}}{63} \right)}{5} \right)$	56
risch	Expression too large to display	798

```
input int(x^(1/2)*arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output  $2/3*x^{(3/2)}*\operatorname{arctanh}(\tanh(b*x+a))^3-4*b*(1/5*x^{(5/2)}*\operatorname{arctanh}(\tanh(b*x+a))^2-4/5*b*(1/7*x^{(7/2)}*\operatorname{arctanh}(\tanh(b*x+a))-2/63*b*x^{(9/2)}))$

### Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.55

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2}{315} (35 b^3 x^4 + 135 a b^2 x^3 + 189 a^2 b x^2 + 105 a^3 x) \sqrt{x}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output  $2/315*(35*b^3*x^4 + 135*a*b^2*x^3 + 189*a^2*b*x^2 + 105*a^3*x)*\operatorname{sqrt}(x)$

### Sympy [F]

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \sqrt{x} \operatorname{atanh}^3(\tanh(a + bx)) dx$$

input `integrate(x**(1/2)*atanh(tanh(b*x+a))**3,x)`

output `Integral(sqrt(x)*atanh(tanh(a + b*x))**3, x)`

### Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.80

$$\begin{aligned} \int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = & -\frac{4}{5} b x^{\frac{5}{2}} \operatorname{artanh}(\tanh(bx + a))^2 \\ & + \frac{2}{3} x^{\frac{3}{2}} \operatorname{artanh}(\tanh(bx + a))^3 \\ & - \frac{16}{315} \left( 2 b^2 x^{\frac{9}{2}} - 9 b x^{\frac{7}{2}} \operatorname{artanh}(\tanh(bx + a)) \right) b \end{aligned}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output 
$$-4/5*b*x^{(5/2)}*arctanh(tanh(b*x + a))^2 + 2/3*x^{(3/2)}*arctanh(tanh(b*x + a))^3 - 16/315*(2*b^2*x^{(9/2)} - 9*b*x^{(7/2)}*arctanh(tanh(b*x + a)))*b$$

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2}{9} b^3 x^{\frac{9}{2}} + \frac{6}{7} ab^2 x^{\frac{7}{2}} + \frac{6}{5} a^2 b x^{\frac{5}{2}} + \frac{2}{3} a^3 x^{\frac{3}{2}}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output 
$$2/9*b^3*x^{(9/2)} + 6/7*a*b^2*x^{(7/2)} + 6/5*a^2*b*x^{(5/2)} + 2/3*a^3*x^{(3/2)}$$

### Mupad [B] (verification not implemented)

Time = 3.14 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.64

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \frac{2b^3 x^{9/2}}{9} - \frac{x^{3/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{12} + \frac{3bx^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{10} - \frac{3b^2 x^{7/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{7}$$

input `int(x^(1/2)*atanh(tanh(a + b*x))^3,x)`

output

```
(2*b^3*x^(9/2))/9 - (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^3)/12 + (3*b*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^2)/10 - (3*b^2*x^(7/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x))/7
```

**Reduce [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3 dx = \int \sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 dx$$

input

```
int(x^(1/2)*atanh(tanh(b*x+a))^3,x)
```

output

```
int(sqrt(x)*atanh(tanh(a + b*x))**3,x)
```

**3.187**      $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} dx$

Optimal result	1330
Mathematica [A] (verified)	1330
Rubi [A] (verified)	1331
Maple [A] (verified)	1332
Fricas [A] (verification not implemented)	1333
Sympy [F]	1333
Maxima [A] (verification not implemented)	1333
Giac [A] (verification not implemented)	1334
Mupad [B] (verification not implemented)	1334
Reduce [F]	1335

**Optimal result**

Integrand size = 15, antiderivative size = 65

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx$$

$$= -\frac{32}{35}b^3x^{7/2} + \frac{16}{5}b^2x^{5/2}\operatorname{arctanh}(\tanh(a + bx))$$

$$- 4bx^{3/2}\operatorname{arctanh}(\tanh(a + bx))^2 + 2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^3$$

output

`-32/35*b^3*x^(7/2)+16/5*b^2*x^(5/2)*arctanh(tanh(b*x+a))-4*b*x^(3/2)*arctanh(tanh(b*x+a))^2+2*x^(1/2)*arctanh(tanh(b*x+a))^3`

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.88

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \frac{2}{35}\sqrt{x}(-16b^3x^3 + 56b^2x^2\operatorname{arctanh}(\tanh(a + bx))$$

$$- 70bx\operatorname{arctanh}(\tanh(a + bx))^2 + 35\operatorname{arctanh}(\tanh(a + bx))^3)$$

input

`Integrate[ArcTanh[Tanh[a + b*x]]^3/Sqrt[x], x]`

output

```
(2*Sqrt[x]*(-16*b^3*x^3 + 56*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 70*b*x*ArcTanh[Tanh[a + b*x]]^2 + 35*ArcTanh[Tanh[a + b*x]]^3))/35
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx$$

$$\downarrow 2599$$

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^3 - 6b \int \sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2 dx$$

$$\downarrow 2599$$

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^3 - 6b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3}b \int x^{3/2}\operatorname{arctanh}(\tanh(a + bx)) dx \right)$$

$$\downarrow 2599$$

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^3 - 6b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3}b \left( \frac{2}{5}x^{5/2}\operatorname{arctanh}(\tanh(a + bx)) - \frac{2}{5}b \int x^{5/2} dx \right) \right)$$

$$\downarrow 15$$

$$2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^3 - 6b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a + bx))^2 - \frac{4}{3}b \left( \frac{2}{5}x^{5/2}\operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{35}bx^{7/2} \right) \right)$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^3/Sqrt[x], x]
```



output

$$2\sqrt{x} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^3 - 6b \left( (2x^{3/2} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^2)/3 - (4b \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]]^3)/35 + (2x^{5/2} \operatorname{ArcTanh}[\operatorname{Tanh}[a + bx]])/5 \right) / 3$$

### Defintions of rubi rules used

rule 15

$$\operatorname{Int}[(a \cdot x)^m, x\_Symbol] \rightarrow \operatorname{Simp}[a \cdot (x^{m+1}) / (m+1), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \operatorname{NeQ}[m, -1]$$

rule 2599

$$\operatorname{Int}[(u \cdot v)^n, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{m+1} \cdot (v^n / (a \cdot (m+1))), x] - \operatorname{Simp}[b \cdot (n / (a \cdot (m+1))) \operatorname{Int}[u^{m+1} \cdot v^{n-1}, x], x] \text{ ; NeQ}[b \cdot u - a \cdot v, 0] \text{ ; FreeQ}[\{m, n\}, x] \ \&\& \ \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \ \operatorname{NeQ}[m, -1] \ \&\& \ ((\operatorname{LtQ}[m, -1] \ \&\& \ \operatorname{GtQ}[n, 0]) \ \&\& \ !(\operatorname{ILtQ}[m+n, -2] \ \&\& \ (\operatorname{FractionQ}[m] \ || \ \operatorname{GeQ}[2 \cdot n + m + 1, 0]))) \ || \ (\operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{IGtQ}[m, 0] \ \&\& \ \operatorname{LeQ}[n, m]) \ || \ (\operatorname{IGtQ}[n, 0] \ \&\& \ !\operatorname{IntegerQ}[m]) \ || \ (\operatorname{ILtQ}[m, 0] \ \&\& \ !\operatorname{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

method	result
derivativedivides	$\frac{2b^3x^{7/2}}{7} + \frac{6(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)b^2x^{5/2}}{5} + 2(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)^2bx^{3/2} + 2(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)^3x^{1/2}$
default	$\frac{2b^3x^{7/2}}{7} + \frac{6(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)b^2x^{5/2}}{5} + 2(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)^2bx^{3/2} + 2(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)^3x^{1/2}$
risch	Expression too large to display

input

$$\operatorname{int}(\operatorname{arctanh}(\operatorname{tanh}(b \cdot x + a))^3 / x^{1/2}, x, \operatorname{method} = \_RETURNVERBOSE)$$

output

$$2/7 \cdot b^3 \cdot x^{7/2} + 6/5 \cdot (\operatorname{arctanh}(\operatorname{tanh}(b \cdot x + a)) - b \cdot x) \cdot b^2 \cdot x^{5/2} + 2 \cdot (\operatorname{arctanh}(\operatorname{tanh}(b \cdot x + a)) - b \cdot x)^2 \cdot b \cdot x^{3/2} + 2 \cdot (\operatorname{arctanh}(\operatorname{tanh}(b \cdot x + a)) - b \cdot x)^3 \cdot x^{1/2}$$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \frac{2}{35} (5b^3x^3 + 21ab^2x^2 + 35a^2bx + 35a^3)\sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(1/2),x, algorithm="fricas")`

output `2/35*(5*b^3*x^3 + 21*a*b^2*x^2 + 35*a^2*b*x + 35*a^3)*sqrt(x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}^3(\tanh(a + bx))}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))**3/x**(1/2),x)`

output `Integral(atanh(tanh(a + b*x))**3/sqrt(x), x)`

**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

$$\begin{aligned} \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = & -4bx^{\frac{3}{2}} \operatorname{artanh}(\tanh(bx + a))^2 \\ & + 2\sqrt{x} \operatorname{artanh}(\tanh(bx + a))^3 \\ & - \frac{16}{35} \left( 2b^2x^{\frac{7}{2}} - 7bx^{\frac{5}{2}} \operatorname{artanh}(\tanh(bx + a)) \right) b \end{aligned}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(1/2),x, algorithm="maxima")`

output `-4*b*x^(3/2)*arctanh(tanh(b*x + a))^2 + 2*sqrt(x)*arctanh(tanh(b*x + a))^3 - 16/35*(2*b^2*x^(7/2) - 7*b*x^(5/2)*arctanh(tanh(b*x + a)))*b`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \frac{2}{7} b^3 x^{7/2} + \frac{6}{5} ab^2 x^{5/2} + 2a^2 b x^{3/2} + 2a^3 \sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(1/2),x, algorithm="giac")`

output `2/7*b^3*x^(7/2) + 6/5*a*b^2*x^(5/2) + 2*a^2*b*x^(3/2) + 2*a^3*sqrt(x)`

**Mupad [B] (verification not implemented)**

Time = 3.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \frac{2b^3 x^{7/2}}{7} - \frac{\sqrt{x} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{4} + \frac{bx^{3/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{2} - \frac{3b^2 x^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{5}$$

input `int(atanh(tanh(a + b*x))^3/x^(1/2),x)`

output `(2*b^3*x^(7/2))/7 - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/4 + (b*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/2 - (3*b^2*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/5`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{\sqrt{x}} dx$$

input `int(atanh(tanh(b*x+a))^3/x^(1/2),x)`

output `int(atanh(tanh(a + b*x))**3/sqrt(x),x)`

### 3.188 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx$

Optimal result	1336
Mathematica [A] (verified)	1336
Rubi [A] (verified)	1337
Maple [A] (verified)	1338
Fricas [A] (verification not implemented)	1339
Sympy [F]	1339
Maxima [A] (verification not implemented)	1339
Giac [A] (verification not implemented)	1340
Mupad [B] (verification not implemented)	1340
Reduce [F]	1341

#### Optimal result

Integrand size = 15, antiderivative size = 63

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx = \frac{32}{5}b^3x^{5/2} - 16b^2x^{3/2}\operatorname{arctanh}(\tanh(a+bx)) + 12b\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^2 - \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}}$$

output

```
32/5*b^3*x^(5/2)-16*b^2*x^(3/2)*arctanh(tanh(b*x+a))+12*b*x^(1/2)*arctanh(
tanh(b*x+a))^2-2*arctanh(tanh(b*x+a))^3/x^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx = \frac{2(16b^3x^3 - 40b^2x^2\operatorname{arctanh}(\tanh(a+bx)) + 30bx\operatorname{arctanh}(\tanh(a+bx))^2)}{5\sqrt{x}}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^3/x^(3/2),x]
```

output

```
(2*(16*b^3*x^3 - 40*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 30*b*x*ArcTanh[Tanh[a
+ b*x]]^2 - 5*ArcTanh[Tanh[a + b*x]]^3))/(5*Sqrt[x])
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{3/2}} dx \\
 & \quad \downarrow \text{2599} \\
 & 6b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} \\
 & \quad \downarrow \text{2599} \\
 & 6b \left( 2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^2 - 4b \int \sqrt{x}\operatorname{arctanh}(\tanh(a+bx)) dx \right) - \\
 & \quad \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} \\
 & \quad \downarrow \text{2599} \\
 & 6b \left( 2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^2 - 4b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx)) - \frac{2}{3}b \int x^{3/2} dx \right) \right) - \\
 & \quad \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}} \\
 & \quad \downarrow \text{15} \\
 & 6b \left( 2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^2 - 4b \left( \frac{2}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx)) - \frac{4}{15}bx^{5/2} \right) \right) - \\
 & \quad \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{\sqrt{x}}
 \end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^3/x^(3/2), x]`

output `(-2*ArcTanh[Tanh[a + b*x]]^3)/Sqrt[x] + 6*b*(2*Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2 - 4*b*((-4*b*x^(5/2))/15 + (2*x^(3/2)*ArcTanh[Tanh[a + b*x]])/3)`

## Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^3}{\sqrt{x}} + 12b \left( \frac{b^2 x^{\frac{5}{2}}}{5} + \frac{2(\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}}}{3} + (\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}} \right) + (\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^3}{\sqrt{x}} + 12b \left( \frac{b^2 x^{\frac{5}{2}}}{5} + \frac{2(\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}}}{3} + (\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}} \right) + (\operatorname{arctanh}(\tanh(bx+a)) - bx) b x^{\frac{3}{2}}$
risch	Expression too large to display

input `int(arctanh(tanh(b*x+a))^3/x^(3/2),x,method=_RETURNVERBOSE)`

output `-2*arctanh(tanh(b*x+a))^3/x^(1/2)+12*b*(1/5*b^2*x^(5/2)+2/3*(arctanh(tanh(b*x+a))-b*x)*b*x^(3/2)+(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.54

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = \frac{2(b^3x^3 + 5ab^2x^2 + 15a^2bx - 5a^3)}{5\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(3/2),x, algorithm="fricas")`

output `2/5*(b^3*x^3 + 5*a*b^2*x^2 + 15*a^2*b*x - 5*a^3)/sqrt(x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = \int \frac{\operatorname{atanh}^3(\tanh(a + bx))}{x^{3/2}} dx$$

input `integrate(atanh(tanh(b*x+a))**3/x**(3/2),x)`

output `Integral(atanh(tanh(a + b*x))**3/x**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.87

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = 12b\sqrt{x} \operatorname{artanh}(\tanh(bx + a))^2 - \frac{2 \operatorname{artanh}(\tanh(bx + a))^3}{\sqrt{x}} + \frac{16}{5} \left( 2b^2x^{5/2} - 5bx^{3/2} \operatorname{artanh}(\tanh(bx + a)) \right) b$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(3/2),x, algorithm="maxima")`

output `12*b*sqrt(x)*arctanh(tanh(b*x + a))^2 - 2*arctanh(tanh(b*x + a))^3/sqrt(x) + 16/5*(2*b^2*x^(5/2) - 5*b*x^(3/2)*arctanh(tanh(b*x + a)))*b`



**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.56

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = \frac{2}{5} b^3 x^{5/2} + 2 ab^2 x^{3/2} + 6 a^2 b \sqrt{x} - \frac{2 a^3}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(3/2),x, algorithm="giac")`output `2/5*b^3*x^(5/2) + 2*a*b^2*x^(3/2) + 6*a^2*b*sqrt(x) - 2*a^3/sqrt(x)`**Mupad [B] (verification not implemented)**

Time = 3.15 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.89

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = \frac{\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)^3}{4\sqrt{x}} + \frac{2b^3 x^{5/2}}{5} + \frac{3b\sqrt{x} \left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)^2}{2} - b^2 x^{3/2} \left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)$$

input `int(atanh(tanh(a + b*x))^3/x^(3/2),x)`output `(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3/(4*x^(1/2)) + (2*b^3*x^(5/2))/5 + (3*b*x^(1/2))*log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/2 - b^2*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{3/2}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{\sqrt{x} x} dx$$

input `int(atanh(tanh(b*x+a))^3/x^(3/2),x)`

output `int(atanh(tanh(a + b*x))**3/(sqrt(x)*x),x)`

**3.189**  $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{5/2}} dx$

Optimal result	1342
Mathematica [A] (verified)	1342
Rubi [A] (verified)	1343
Maple [A] (verified)	1344
Fricas [A] (verification not implemented)	1345
Sympy [A] (verification not implemented)	1345
Maxima [A] (verification not implemented)	1345
Giac [A] (verification not implemented)	1346
Mupad [B] (verification not implemented)	1346
Reduce [F]	1347

**Optimal result**

Integrand size = 15, antiderivative size = 65

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = -\frac{32}{3}b^3x^{3/2} + 16b^2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - \frac{4b\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{3x^{3/2}}$$

output

```
-32/3*b^3*x^(3/2)+16*b^2*x^(1/2)*arctanh(tanh(b*x+a))-4*b*arctanh(tanh(b*x+a))^2/x^(1/2)-2/3*arctanh(tanh(b*x+a))^3/x^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = \frac{2(16b^3x^3 - 24b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 6bx\operatorname{arctanh}(\tanh(a + bx))^2 + \operatorname{arctanh}(\tanh(a + bx))^3)}{3x^{3/2}}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^3/x^(5/2), x]
```

output

$$\frac{(-2*(16*b^3*x^3 - 24*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 6*b*x*ArcTanh[Tanh[a + b*x]])^2 + ArcTanh[Tanh[a + b*x]]^3)/(3*x^(3/2))}{}$$
**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx \\ & \quad \downarrow \text{2599} \\ & 2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{3x^{3/2}} \\ & \quad \downarrow \text{2599} \\ & 2b \left( 4b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{3x^{3/2}} \\ & \quad \downarrow \text{2599} \\ & 2b \left( 4b \left( 2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - 2b \int \sqrt{x} dx \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{3x^{3/2}} \\ & \quad \downarrow \text{15} \\ & 2b \left( 4b \left( 2\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - \frac{4}{3}bx^{3/2} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{3x^{3/2}} \end{aligned}$$

input

$$\text{Int}[ArcTanh[Tanh[a + b*x]]^3/x^(5/2), x]$$

output 
$$\frac{(-2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^3)/(3 x^{3/2}) + 2 b ((-2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^2)/\sqrt{x} + 4 b ((-4 b x^{3/2})/3 + 2 \sqrt{x} \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]))}{1}$$

### Defintions of rubi rules used

rule 15 
$$\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a(x^{(m+1)})/(m+1), x] \text{ /; FreeQ}[\{a, m\}, x] \ \&\& \ \operatorname{NeQ}[m, -1]$$

rule 2599 
$$\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}(v^n/(a(m+1))), x] - \operatorname{Simp}[b(n/(a(m+1))) \operatorname{Int}[u^{(m+1)}v^{(n-1)}, x], x] \text{ /; NeQ}[b u - a v, 0] \text{ /; FreeQ}[\{m, n\}, x] \ \&\& \ \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \ \operatorname{NeQ}[m, -1] \ \&\& \ ((\operatorname{LtQ}[m, -1] \ \&\& \ \operatorname{GtQ}[n, 0]) \ \&\& \ !(\operatorname{ILtQ}[m+n, -2] \ \&\& \ (\operatorname{FractionQ}[m] \ || \ \operatorname{GeQ}[2n+m+1, 0]))) \ || \ (\operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{IGtQ}[m, 0] \ \&\& \ \operatorname{LeQ}[n, m]) \ || \ (\operatorname{IGtQ}[n, 0] \ \&\& \ !\operatorname{IntegerQ}[m]) \ || \ (\operatorname{ILtQ}[m, 0] \ \&\& \ !\operatorname{IntegerQ}[n]))]$$

### Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{3x^{3/2}} + 4b \left( -\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{\sqrt{x}} + 4b \left( \sqrt{x} \operatorname{arctanh}(\operatorname{tanh}(bx+a)) - \frac{2b}{3} \right) \right)$
default	$-\frac{2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^3}{3x^{3/2}} + 4b \left( -\frac{\operatorname{arctanh}(\operatorname{tanh}(bx+a))^2}{\sqrt{x}} + 4b \left( \sqrt{x} \operatorname{arctanh}(\operatorname{tanh}(bx+a)) - \frac{2b}{3} \right) \right)$
risch	Expression too large to display

input 
$$\operatorname{int}(\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3/x^{(5/2)}, x, \operatorname{method}=\_RETURNVERBOSE)$$

output 
$$-2/3 \operatorname{arctanh}(\operatorname{tanh}(b*x+a))^3/x^{(3/2)} + 4*b*(-\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^2/x^{(1/2)} + 4*b*(x^{(1/2)}*\operatorname{arctanh}(\operatorname{tanh}(b*x+a)) - 2/3*b*x^{(3/2)}))$$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = \frac{2(b^3x^3 + 9ab^2x^2 - 9a^2bx - a^3)}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(5/2),x, algorithm="fricas")`

output `2/3*(b^3*x^3 + 9*a*b^2*x^2 - 9*a^2*b*x - a^3)/x^(3/2)`

**Sympy [A] (verification not implemented)**

Time = 1.28 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = -\frac{32b^3x^{3/2}}{3} + 16b^2\sqrt{x} \operatorname{atanh}(\tanh(a + bx)) - \frac{4b \operatorname{atanh}^2(\tanh(a + bx))}{\sqrt{x}} - \frac{2 \operatorname{atanh}^3(\tanh(a + bx))}{3x^{3/2}}$$

input `integrate(atanh(tanh(b*x+a))**3/x**(5/2),x)`

output `-32*b**3*x**(3/2)/3 + 16*b**2*sqrt(x)*atanh(tanh(a + b*x)) - 4*b*atanh(tanh(a + b*x))**2/sqrt(x) - 2*atanh(tanh(a + b*x))**3/(3*x**(3/2))`

**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = -\frac{4b \operatorname{artanh}(\tanh(bx + a))^2}{\sqrt{x}} - \frac{16}{3} \left( 2b^2x^{3/2} - 3b\sqrt{x} \operatorname{artanh}(\tanh(bx + a)) \right) b - \frac{2 \operatorname{artanh}(\tanh(bx + a))^3}{3x^{3/2}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(5/2),x, algorithm="maxima")`

output

```
-4*b*arctanh(tanh(b*x + a))^2/sqrt(x) - 16/3*(2*b^2*x^(3/2) - 3*b*sqrt(x)*
arctanh(tanh(b*x + a)))*b - 2/3*arctanh(tanh(b*x + a))^3/x^(3/2)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = \frac{2}{3} b^3 x^{\frac{3}{2}} + 6 ab^2 \sqrt{x} - \frac{2(9 a^2 bx + a^3)}{3 x^{\frac{3}{2}}}$$

input

```
integrate(arctanh(tanh(b*x+a))^3/x^(5/2),x, algorithm="giac")
```

output

```
2/3*b^3*x^(3/2) + 6*a*b^2*sqrt(x) - 2/3*(9*a^2*b*x + a^3)/x^(3/2)
```

**Mupad [B] (verification not implemented)**

Time = 3.16 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = \frac{\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)^3}{12x^{3/2}} + \frac{2b^3 x^{3/2}}{3} - \frac{3b\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)^2}{2\sqrt{x}} - 3b^2 \sqrt{x} \left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)$$

input

```
int(atanh(tanh(a + b*x))^3/x^(5/2),x)
```

output

```
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)^3/(12*x^(3/2)) + (2*b^3*x^(3/2))/3 - (3*b*(log(2
/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*
b*x) + 1)) + 2*b*x)^2)/(2*x^(1/2)) - 3*b^2*x^(1/2)*(log(2/(exp(2*a)*exp(2*
b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*
x)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{5/2}} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^3}{\sqrt{x} x^2} dx$$

input `int(atanh(tanh(b*x+a))^3/x^(5/2),x)`

output `int(atanh(tanh(a + b*x))**3/(sqrt(x)*x**2),x)`



### 3.190 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{7/2}} dx$

Optimal result	1348
Mathematica [A] (verified)	1348
Rubi [A] (verified)	1349
Maple [A] (verified)	1350
Fricas [A] (verification not implemented)	1351
Sympy [A] (verification not implemented)	1351
Maxima [A] (verification not implemented)	1351
Giac [A] (verification not implemented)	1352
Mupad [B] (verification not implemented)	1352
Reduce [B] (verification not implemented)	1353

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{7/2}} dx = \frac{32b^3\sqrt{x}}{5} - \frac{16b^2\operatorname{arctanh}(\tanh(a+bx))}{5\sqrt{x}} - \frac{4b\operatorname{arctanh}(\tanh(a+bx))^2}{5x^{3/2}} - \frac{2\operatorname{arctanh}(\tanh(a+bx))^3}{5x^{5/2}}$$

output

```
32/5*b^3*x^(1/2)-16/5*b^2*arctanh(tanh(b*x+a))/x^(1/2)-4/5*b*arctanh(tanh(b*x+a))^2/x^(3/2)-2/5*arctanh(tanh(b*x+a))^3/x^(5/2)
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^3}{x^{7/2}} dx = \frac{2(16b^3x^3 - 8b^2x^2\operatorname{arctanh}(\tanh(a+bx)) - 2bx\operatorname{arctanh}(\tanh(a+bx))^2 - \operatorname{arctanh}(\tanh(a+bx))^3)}{5x^{5/2}}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^3/x^(7/2),x]
```

output

```
(2*(16*b^3*x^3 - 8*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 2*b*x*ArcTanh[Tanh[a + b*x]]^2 - ArcTanh[Tanh[a + b*x]]^3)/(5*x^(5/2))
```

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx$$

↓ 2599

$$\frac{6}{5}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^2}{x^{5/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{5x^{5/2}}$$

↓ 2599

$$\frac{6}{5}b \left( \frac{4}{3}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{5x^{5/2}}$$

↓ 2599

$$\frac{6}{5}b \left( \frac{4}{3}b \left( 2b \int \frac{1}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{5x^{5/2}}$$

↓ 15

$$\frac{6}{5}b \left( \frac{4}{3}b \left( 4b\sqrt{x} - \frac{2\operatorname{arctanh}(\tanh(a + bx))}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^2}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^3}{5x^{5/2}}$$

input `Int [ArcTanh [Tanh [a + b*x]] ^3/x^(7/2) , x]`

output `(-2*ArcTanh [Tanh [a + b*x]] ^3)/(5*x^(5/2)) + (6*b*((-2*ArcTanh [Tanh [a + b*x]] ^2)/(3*x^(3/2)) + (4*b*(4*b*Sqrt [x] - (2*ArcTanh [Tanh [a + b*x]])/Sqrt [x]))/3))/5`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^3}{5x^{\frac{5}{2}}} + \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^{\frac{3}{2}}} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}} + 2b\sqrt{x} \right)}{3} \right)}{5}$	56
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^3}{5x^{\frac{5}{2}}} + \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^2}{3x^{\frac{3}{2}}} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{\sqrt{x}} + 2b\sqrt{x} \right)}{3} \right)}{5}$	56
risch	Expression too large to display	7814

input `int(arctanh(tanh(b*x+a))^3/x^(7/2),x,method=_RETURNVERBOSE)`

output `-2/5*arctanh(tanh(b*x+a))^3/x^(5/2)+12/5*b*(-1/3*arctanh(tanh(b*x+a))^2/x^(3/2)+4/3*b*(-arctanh(tanh(b*x+a))/x^(1/2)+2*b*x^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = \frac{2(5b^3x^3 - 15ab^2x^2 - 5a^2bx - a^3)}{5x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(7/2),x, algorithm="fricas")`

output `2/5*(5*b^3*x^3 - 15*a*b^2*x^2 - 5*a^2*b*x - a^3)/x^(5/2)`

**Sympy [A] (verification not implemented)**

Time = 12.81 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = \frac{32b^3\sqrt{x}}{5} - \frac{16b^2 \operatorname{atanh}(\tanh(a + bx))}{5\sqrt{x}} - \frac{4b \operatorname{atanh}^2(\tanh(a + bx))}{5x^{3/2}} - \frac{2 \operatorname{atanh}^3(\tanh(a + bx))}{5x^{5/2}}$$

input `integrate(atanh(tanh(b*x+a))**3/x**(7/2),x)`

output `32*b**3*sqrt(x)/5 - 16*b**2*atanh(tanh(a + b*x))/(5*sqrt(x)) - 4*b*atanh(atanh(a + b*x))**2/(5*x**(3/2)) - 2*atanh(tanh(a + b*x))**3/(5*x**(5/2))`

**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.80

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = \frac{16}{5} \left( 2b^2\sqrt{x} - \frac{b \operatorname{artanh}(\tanh(bx + a))}{\sqrt{x}} \right) b - \frac{4b \operatorname{artanh}(\tanh(bx + a))^2}{5x^{3/2}} - \frac{2 \operatorname{artanh}(\tanh(bx + a))^3}{5x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^3/x^(7/2),x, algorithm="maxima")`

output

```
16/5*(2*b^2*sqrt(x) - b*arctanh(tanh(b*x + a))/sqrt(x))*b - 4/5*b*arctanh(
tanh(b*x + a))^2/x^(3/2) - 2/5*arctanh(tanh(b*x + a))^3/x^(5/2)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = 2b^3\sqrt{x} - \frac{2(15ab^2x^2 + 5a^2bx + a^3)}{5x^{5/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^3/x^(7/2),x, algorithm="giac")
```

output

```
2*b^3*sqrt(x) - 2/5*(15*a*b^2*x^2 + 5*a^2*b*x + a^3)/x^(5/2)
```

**Mupad [B] (verification not implemented)**

Time = 3.16 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.64

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = \frac{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^3}{20x^{5/2}} + 2b^3\sqrt{x} + \frac{3b^2\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{\sqrt{x}} - \frac{b\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{2x^{3/2}}$$

input

```
int(atanh(tanh(a + b*x))^3/x^(7/2),x)
```

output

```
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)^3/(20*x^(5/2)) + 2*b^3*x^(1/2) + (3*b^2*(log(2/(
exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*
x) + 1)) + 2*b*x))/x^(1/2) - (b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2
*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*x^(3/2))
```

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^3}{x^{7/2}} dx = \frac{-\frac{2\operatorname{atanh}(\tanh(bx+a))^3}{5} - \frac{4\operatorname{atanh}(\tanh(bx+a))^2 bx}{5} - \frac{16\operatorname{atanh}(\tanh(bx+a))b^2 x^2}{5} + \frac{32b^3 x^3}{5}}{\sqrt{x} x^2}$$

input

```
int(atanh(tanh(b*x+a))^3/x^(7/2),x)
```

output

```
(2*( - atanh(tanh(a + b*x))**3 - 2*atanh(tanh(a + b*x))**2*b*x - 8*atanh(tanh(a + b*x))*b**2*x**2 + 16*b**3*x**3))/(5*sqrt(x)*x**2)
```

### 3.191 $\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1354
Mathematica [A] (verified)	1355
Rubi [A] (verified)	1355
Maple [B] (verified)	1357
Fricas [A] (verification not implemented)	1358
Sympy [F]	1358
Maxima [A] (verification not implemented)	1359
Giac [A] (verification not implemented)	1359
Mupad [B] (verification not implemented)	1360
Reduce [F]	1361

#### Optimal result

Integrand size = 15, antiderivative size = 143

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2x^{7/2}}{7b} + \frac{2x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{5b^2} + \frac{2x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{3b^3} + \frac{2\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{b^4} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}}{b^{9/2}}$$

output

```
2/7*x^(7/2)/b+2/5*x^(5/2)*(b*x-arctanh(tanh(b*x+a)))/b^2+2/3*x^(3/2)*(b*x-
arctanh(tanh(b*x+a)))^2/b^3+2*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^3/b^4-2*a
rctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh
(b*x+a)))^(7/2)/b^(9/2)
```

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.90

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \left( 176b^{7/2}x^{7/2} - 406b^{5/2}x^{5/2}\operatorname{arctanh}(\tanh(a + bx)) + 350b^{3/2}x^{3/2}\operatorname{arctanh}(\tanh(a + bx)) - 105\sqrt{b}\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2 + 105\operatorname{ArcTan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-(bx) + \operatorname{arctanh}(\tanh(a + bx))}}\right) \right)}{105b^{9/2}}$$

input `Integrate[x^(7/2)/ArcTanh[Tanh[a + b*x]],x]`

output  $(2*(176*b^{(7/2)}*x^{(7/2)} - 406*b^{(5/2)}*x^{(5/2)}*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]] + 350*b^{(3/2)}*x^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^2 - 105*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[x]*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3 + 105*\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[x])/\operatorname{Sqrt}[-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]])*(-(b*x) + \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]])^{(7/2)})/(105*b^{(9/2)})$

### Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2590, 2590, 2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2x^{7/2}}{7b}$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2x^{5/2}}{5b} \right)}{b} + \frac{2x^{7/2}}{7b}$$



↓ 2590

$$(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx + \frac{2x^{3/2}}{3b} \right)}{b} \right)}{b} + 2$$

$\frac{2x^{7/2}}{7b}$   
↓ 2590

$$(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} \right)}{b} \right)}{b}$$

$\frac{2x^{7/2}}{7b}$   
↓ 2593

$$\left( \left( \left( \frac{2\sqrt{x}}{b} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\frac{b^{3/2}}{b}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))} \right) (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) + \frac{2x^{3/2}}{3b} (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) / b$$

$\frac{2x^{7/2}}{7b}$

input `Int[x^(7/2)/ArcTanh[Tanh[a + b*x]],x]`

output 
$$\frac{(2x^{7/2})/(7b) + (((2x^{5/2})/(5b) + (((2x^{3/2})/(3b) + ((2\sqrt{x})/b - 2\text{ArcTanh}[\sqrt{b}\sqrt{x}]/\sqrt{bx - \text{ArcTanh}[\text{Tanh}[a + b*x]])]*\sqrt{bx - \text{ArcTanh}[\text{Tanh}[a + b*x]])})/b^{3/2})*(bx - \text{ArcTanh}[\text{Tanh}[a + b*x]])/b)*(bx - \text{ArcTanh}[\text{Tanh}[a + b*x]])/b)*(bx - \text{ArcTanh}[\text{Tanh}[a + b*x]])/b$$

**Defintions of rubi rules used**

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 261 vs. 2(119) = 238.

Time = 0.35 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.83

method	result
derivativedivides	$2 \left( -\frac{b^3 x^{\frac{7}{2}}}{7} + \frac{(\text{arctanh}(\tanh(bx+a)) - bx)b^2 x^{\frac{5}{2}}}{5} - \frac{(a^2 + 2a(\text{arctanh}(\tanh(bx+a)) - bx - a) + (\text{arctanh}(\tanh(bx+a)) - bx - a)^2)x^{\frac{3}{2}}b}{3} \right) \frac{1}{b^4}$
default	$2 \left( -\frac{b^3 x^{\frac{7}{2}}}{7} + \frac{(\text{arctanh}(\tanh(bx+a)) - bx)b^2 x^{\frac{5}{2}}}{5} - \frac{(a^2 + 2a(\text{arctanh}(\tanh(bx+a)) - bx - a) + (\text{arctanh}(\tanh(bx+a)) - bx - a)^2)x^{\frac{3}{2}}b}{3} \right) \frac{1}{b^4}$

input `int(x^(7/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-2/b^4*(-1/7*b^3*x^(7/2)+1/5*(arctanh(tanh(b*x+a))-b*x)*b^2*x^(5/2)-1/3*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)*x^(3/2)*b+(arctanh(tanh(b*x+a))-b*x)*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)*x^(1/2))+2*(a^4+4*a^3*(arctanh(tanh(b*x+a))-b*x-a)+6*a^2*(arctanh(tanh(b*x+a))-b*x-a)^2+4*a*(arctanh(tanh(b*x+a))-b*x-a)^3+(arctanh(tanh(b*x+a))-b*x-a)^4)/b^4/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.07

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{105 a^3 \sqrt{-\frac{a}{b}} \log\left(\frac{bx + 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(15b^3x^3 - 21ab^2x^2 + 35a^2bx - 105a^3)\sqrt{x}}{105b^4}$$

input

```
integrate(x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
[1/105*(105*a^3*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(15*b^3*x^3 - 21*a*b^2*x^2 + 35*a^2*b*x - 105*a^3)*sqrt(x))/b^4, 2/105*(105*a^3*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) + (15*b^3*x^3 - 21*a*b^2*x^2 + 35*a^2*b*x - 105*a^3)*sqrt(x))/b^4]
```

**Sympy [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^{7/2}}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input

```
integrate(x**(7/2)/atanh(tanh(b*x+a)),x)
```

output

```
Integral(x**(7/2)/atanh(tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.45

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 a^4 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}b^4} + \frac{2 \left(15 b^3 x^{7/2} - 21 ab^2 x^{5/2} + 35 a^2 b x^{3/2} - 105 a^3 \sqrt{x}\right)}{105 b^4}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `2*a^4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^4) + 2/105*(15*b^3*x^(7/2) - 21*a*b^2*x^(5/2) + 35*a^2*b*x^(3/2) - 105*a^3*sqrt(x))/b^4`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.49

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 a^4 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}b^4} + \frac{2 \left(15 b^6 x^{7/2} - 21 ab^5 x^{5/2} + 35 a^2 b^4 x^{3/2} - 105 a^3 b^3 \sqrt{x}\right)}{105 b^7}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `2*a^4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^4) + 2/105*(15*b^6*x^(7/2) - 21*a*b^5*x^(5/2) + 35*a^2*b^4*x^(3/2) - 105*a^3*b^3*sqrt(x))/b^7`

**Mupad [B] (verification not implemented)**

Time = 3.72 (sec) , antiderivative size = 475, normalized size of antiderivative = 3.32

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x^{7/2}}{7b} + \frac{x^{5/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{5b^2}$$

$$+ \frac{x^{3/2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{6b^3}$$

$$+ \frac{\sqrt{x} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{4b^4}$$

$$+ \frac{\sqrt{2} \ln \left( \frac{64b^{19/2} \left( \sqrt{2} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx + 2\sqrt{2}bx} \right)}{\left( \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) \right) \sqrt{\ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx}} \right)}{16b^{9/2}} \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)$$

input `int(x^(7/2)/atanh(tanh(a + b*x)),x)`

output

```
(2*x^(7/2))/(7*b) + (x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(5*b^2) + (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(6*b^3) + (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(4*b^4) + (2^(1/2)*log((64*b^(19/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x^(1/2) + 2*2^(1/2)*b*x))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x^(1/2)))/(16*b^(9/2))
```

Reduce [F]

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{\sqrt{x} x^3}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(7/2)/atanh(tanh(b*x+a)),x)`

output `int((sqrt(x)*x**3)/atanh(tanh(a + b*x)),x)`

### 3.192 $\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1362
Mathematica [A] (verified)	1363
Rubi [A] (verified)	1363
Maple [B] (verified)	1365
Fricas [A] (verification not implemented)	1366
Sympy [F]	1366
Maxima [A] (verification not implemented)	1367
Giac [A] (verification not implemented)	1367
Mupad [B] (verification not implemented)	1367
Reduce [F]	1368

#### Optimal result

Integrand size = 15, antiderivative size = 116

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2x^{5/2}}{5b} + \frac{2x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{3b^2} + \frac{2\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{b^3} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}}{b^{7/2}}$$

output

```
2/5*x^(5/2)/b+2/3*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))/b^2+2*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2/b^3-2*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a))))^(1/2)*(b*x-arctanh(tanh(b*x+a)))^(5/2)/b^(7/2)
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.93

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \left( 23b^{5/2}x^{5/2} - 35b^{3/2}x^{3/2}\operatorname{arctanh}(\tanh(a + bx)) + 15\sqrt{b}\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) \right)}{15b^{7/2}}$$

input `Integrate[x^(5/2)/ArcTanh[Tanh[a + b*x]],x]`

output `(2*(23*b^(5/2)*x^(5/2) - 35*b^(3/2)*x^(3/2)*ArcTanh[Tanh[a + b*x]] + 15*sqrt[b]*sqrt[x]*ArcTanh[Tanh[a + b*x]]^2 - 15*ArcTan[(sqrt[b]*sqrt[x])/sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2))/(15*b^(7/2))`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2590, 2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2x^{5/2}}{5b}$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2x^{3/2}}{3b} \right)}{b} + \frac{2x^{5/2}}{5b}$$



$$\begin{aligned}
 & \downarrow 2590 \\
 & (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx + \frac{2\sqrt{x}}{b}}{b} \right) + \\
 & \frac{2x^{5/2}}{5b} \\
 & \downarrow 2593 \\
 & \left( \frac{\left( \frac{2\sqrt{x}}{b} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{b^{3/2}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}{b} (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) + \frac{2x^{3/2}}{3b} (bx - \operatorname{arctanh}(\tanh(a + bx)))}{\frac{2x^{5/2}}{5b}}
 \end{aligned}$$

input `Int[x^(5/2)/ArcTanh[Tanh[a + b*x]], x]`

output `(2*x^(5/2))/(5*b) + (((2*x^(3/2))/(3*b) + (((2*sqrt[x])/b - (2*ArcTanh[(sqrt[b]*sqrt[x])/sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2))*(b*x - ArcTanh[Tanh[a + b*x]]))/b*(b*x - ArcTanh[Tanh[a + b*x]]))/b`

**Defintions of rubi rules used**

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 200 vs.  $2(96) = 192$ .

Time = 0.27 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.73

method	result
derivativedivides	$\frac{2b^2x^{\frac{5}{2}}}{5} - \frac{2bx^{\frac{3}{2}}a}{3} - \frac{2bx^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{3} + \frac{2a^2\sqrt{x} + 4a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x} + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2\sqrt{x}}{b^3}$
default	$\frac{2b^2x^{\frac{5}{2}}}{5} - \frac{2bx^{\frac{3}{2}}a}{3} - \frac{2bx^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{3} + \frac{2a^2\sqrt{x} + 4a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x} + 2(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2\sqrt{x}}{b^3}$

input

```
int(x^(5/2)/arctanh(tanh(b*x+a)), x, method=_RETURNVERBOSE)
```

output

```
2/b^3*(1/5*b^2*x^(5/2)-1/3*b*x^(3/2)*a-1/3*b*x^(3/2)*(arctanh(tanh(b*x+a))-b*x-a)+a^2*x^(1/2)+2*a*(arctanh(tanh(b*x+a))-b*x-a)*x^(1/2)+(arctanh(tanh(b*x+a))-b*x-a)^2*x^(1/2))+2*(-a^3-3*a^2*(arctanh(tanh(b*x+a))-b*x-a)-3*a*(arctanh(tanh(b*x+a))-b*x-a)^2-(arctanh(tanh(b*x+a))-b*x-a)^3)/b^3/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.14

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{15 a^2 \sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(3b^2x^2 - 5abx + 15a^2)\sqrt{x}}{15b^3}, \right. \\ \left. - \frac{2\left(15a^2\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (3b^2x^2 - 5abx + 15a^2)\sqrt{x}\right)}{15b^3} \right]$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `[1/15*(15*a^2*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(3*b^2*x^2 - 5*a*b*x + 15*a^2)*sqrt(x))/b^3, -2/15*(15*a^2*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - (3*b^2*x^2 - 5*a*b*x + 15*a^2)*sqrt(x))/b^3]`

**Sympy [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^{5/2}}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**(5/2)/atanh(tanh(b*x+a)),x)`

output `Integral(x**(5/2)/atanh(tanh(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.47

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2a^3 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} + \frac{2\left(3b^2x^{5/2} - 5abx^{3/2} + 15a^2\sqrt{x}\right)}{15b^3}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-2*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2/15*(3*b^2*x^(5/2) - 5*a*b*x^(3/2) + 15*a^2*sqrt(x))/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.51

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2a^3 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} + \frac{2\left(3b^4x^{5/2} - 5ab^3x^{3/2} + 15a^2b^2\sqrt{x}\right)}{15b^5}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `-2*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2/15*(3*b^4*x^(5/2) - 5*a*b^3*x^(3/2) + 15*a^2*b^2*sqrt(x))/b^5`**Mupad [B] (verification not implemented)**

Time = 3.35 (sec) , antiderivative size = 415, normalized size of antiderivative = 3.58

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2x^{5/2}}{5b} + \frac{x^{3/2} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)}{3b^2}$$

$$+ \frac{\sqrt{x} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2}{2b^3}$$

$$+ \frac{\sqrt{2} \ln \left( \frac{16b^{15/2} \left( \sqrt{2} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx + 2\sqrt{2}bx} \right)}{\left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) \right) \sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx}} \right)}{8b^{7/2}} \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)$$

input `int(x^(5/2)/atanh(tanh(a + b*x)),x)`

output `(2*x^(5/2))/(5*b) + (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(3*b^2) + (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*b^3) + (2^(1/2)*log((16*b^(15/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2))/(8*b^(7/2))`

## Reduce [F]

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{\sqrt{x} x^2}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a)),x)`

output `int((sqrt(x)*x**2)/atanh(tanh(a + b*x)),x)`

### 3.193 $\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1369
Mathematica [A] (verified)	1369
Rubi [A] (verified)	1370
Maple [A] (verified)	1371
Fricas [A] (verification not implemented)	1372
Sympy [F]	1372
Maxima [A] (verification not implemented)	1373
Giac [A] (verification not implemented)	1373
Mupad [B] (verification not implemented)	1373
Reduce [F]	1374

#### Optimal result

Integrand size = 15, antiderivative size = 89

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2x^{3/2}}{3b} + \frac{2\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^2} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}}{b^{5/2}}$$

output

$$\frac{2}{3}x^{3/2}/b + 2x^{1/2}*(b*x - \operatorname{arctanh}(\tanh(b*x+a)))/b^2 - 2*\operatorname{arctanh}(b^{1/2}*x^{1/2}/(b*x - \operatorname{arctanh}(\tanh(b*x+a))))^{1/2}*(b*x - \operatorname{arctanh}(\tanh(b*x+a)))^{3/2}/b^{5/2}$$

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.97

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2x^{3/2}}{3b} - \frac{2\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{b^2} + \frac{2\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{3/2}}{b^{5/2}}$$

input `Integrate[x^(3/2)/ArcTanh[Tanh[a + b*x]], x]`

output  $(2*x^{(3/2)})/(3*b) - (2*sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/b^2 + (2*ArcTan[(sqrt[b]*sqrt[x])/sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2))/b^{(5/2)}$

### Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2x^{3/2}}{3b}$$

$$\downarrow 2590$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx}{b} + \frac{2\sqrt{x}}{b} \right)}{b} + \frac{2x^{3/2}}{3b}$$

$$\downarrow 2593$$

$$\frac{\left( \frac{2\sqrt{x}}{b} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right) \sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}{b^{3/2}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{b} + \frac{2x^{3/2}}{3b}$$

input `Int[x^(3/2)/ArcTanh[Tanh[a + b*x]],x]`

output `(2*x^(3/2))/(3*b) + (((2*Sqrt[x])/b - (2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2))*(b*x - ArcTanh[Tanh[a + b*x]])/b`

**Defintions of rubi rules used**

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.35

method	result
derivativedivides	$-\frac{2\left(-\frac{bx^{\frac{3}{2}}}{3} + a\sqrt{x} + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x}\right)}{b^2} + \frac{2\left(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + x}\right)}{b^2\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + x}}$
default	$-\frac{2\left(-\frac{bx^{\frac{3}{2}}}{3} + a\sqrt{x} + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x}\right)}{b^2} + \frac{2\left(a^2 + 2a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a) + (\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + x}\right)}{b^2\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2 + x}}$

input `int(x^(3/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`



output

```
-2/b^2*(-1/3*b*x^(3/2)+a*x^(1/2)+(arctanh(tanh(b*x+a))-b*x-a)*x^(1/2))+2*(
a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/b^2/(
(arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a)
)-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.16

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{3a\sqrt{-\frac{a}{b}} \log\left(\frac{bx+2b\sqrt{x}\sqrt{-\frac{a}{b}}-a}{bx+a}\right) + 2(bx-3a)\sqrt{x}}{3b^2}, \frac{2\left(3a\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}}{a}\right) + (bx-3a)\sqrt{x}\right)}{3b^2} \right]$$

input

```
integrate(x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
[1/3*(3*a*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2
*(b*x - 3*a)*sqrt(x))/b^2, 2/3*(3*a*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a
) + (b*x - 3*a)*sqrt(x))/b^2]
```

**Sympy [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{x^{\frac{3}{2}}}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input

```
integrate(x**(3/2)/atanh(tanh(b*x+a)),x)
```

output

```
Integral(x**(3/2)/atanh(tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.47

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{2\left(bx^{\frac{3}{2}} - 3a\sqrt{x}\right)}{3b^2}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `2*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + 2/3*(b*x^(3/2) - 3*a*sqrt(x))/b^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.51

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{2\left(b^2x^{\frac{3}{2}} - 3ab\sqrt{x}\right)}{3b^3}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `2*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + 2/3*(b^2*x^(3/2) - 3*a*b*sqrt(x))/b^3`**Mupad [B] (verification not implemented)**

Time = 3.85 (sec) , antiderivative size = 354, normalized size of antiderivative = 3.98

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2x^{3/2}}{3b} + \frac{\sqrt{x} \left( \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx \right)}{b^2}$$

$$+ \frac{\sqrt{2} \ln \left( \frac{4b^{11/2} \left( \sqrt{2} \left( \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx \right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx + 2\sqrt{2}bx} \right)}{\left( \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) \right) \sqrt{\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx}} \right)}{4b^{5/2}} \left( \ln \right)$$

input `int(x^(3/2)/atanh(tanh(a + b*x)),x)`

output `(2*x^(3/2))/(3*b) + (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/b^2 + (2^(1/2)*log((4*b^(11/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2))/(4*b^(5/2))`

### Reduce [F]

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{\sqrt{x} x}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(3/2)/atanh(tanh(b*x+a)),x)`

output `int((sqrt(x)*x)/atanh(tanh(a + b*x)),x)`

### 3.194 $\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1375
Mathematica [A] (verified)	1375
Rubi [A] (verified)	1376
Maple [A] (verified)	1377
Fricas [A] (verification not implemented)	1378
Sympy [F]	1378
Maxima [A] (verification not implemented)	1379
Giac [A] (verification not implemented)	1379
Mupad [B] (verification not implemented)	1379
Reduce [F]	1380

#### Optimal result

Integrand size = 15, antiderivative size = 64

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2\sqrt{x}}{b} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right) \sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}}$$

output

```
2*x^(1/2)/b-2*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(1/2)/b^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2\sqrt{x}}{b} - \frac{2\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}\right) \sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}}$$

input

```
Integrate[Sqrt[x]/ArcTanh[Tanh[a + b*x]], x]
```

output

$$(2\sqrt{x})/b - (2\text{ArcTan}[(\sqrt{b}\sqrt{x})/\sqrt{-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]}]]*\sqrt{-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]})/b^{(3/2)}$$
**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\text{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow \text{2590}$$

$$\frac{(bx - \text{arctanh}(\tanh(a + bx)))}{b} \int \frac{1}{\sqrt{x}\text{arctanh}(\tanh(a + bx))} dx + \frac{2\sqrt{x}}{b}$$

$$\downarrow \text{2593}$$

$$\frac{2\sqrt{x}}{b} - \frac{2\text{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \text{arctanh}(\tanh(a + bx))}}\right) \sqrt{bx - \text{arctanh}(\tanh(a + bx))}}{b^{3/2}}$$

input

$$\text{Int}[\sqrt{x}/\text{ArcTanh}[\text{Tanh}[a + b*x]], x]$$

output

$$(2\sqrt{x})/b - (2\text{ArcTanh}[(\sqrt{b}\sqrt{x})/\sqrt{b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]}]]*\sqrt{b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]})/b^{(3/2)}$$

## Defintions of rubi rules used

rule 2590 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]`

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

## Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b} + \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{b\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}$	66
default	$\frac{2\sqrt{x}}{b} + \frac{2(bx - \operatorname{arctanh}(\tanh(bx+a))) \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{b\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}$	66

input `int(x^(1/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output  $2*x^{(1/2)}/b+2*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/b/((\operatorname{arctanh}(\tanh(b*x+a))-b*x)*b)^{(1/2)}*\operatorname{arctan}(b*x^{(1/2)}/((\operatorname{arctanh}(\tanh(b*x+a))-b*x)*b)^{(1/2)})$

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.33

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{\sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2\sqrt{x}}{b}, \right. \\ \left. - \frac{2\left(\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - \sqrt{x}\right)}{b} \right]$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `[(sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*sqrt(x))/b, -2*(sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - sqrt(x))/b]`

**Sympy [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a)),x)`

output `Integral(sqrt(x)/atanh(tanh(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.48

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2a \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-2*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) + 2*sqrt(x)/b`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.48

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2a \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `-2*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) + 2*sqrt(x)/b`**Mupad [B] (verification not implemented)**

Time = 4.09 (sec) , antiderivative size = 296, normalized size of antiderivative = 4.62

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2\sqrt{x}}{b} + \frac{\sqrt{2} \ln\left(\frac{b^{7/2} \left(\sqrt{2} \left(\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx\right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx + 2\sqrt{2}bx}\right)}{\left(\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)\right) \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx}}\right)}{2b^{3/2}} \sqrt{\dots}$$

input `int(x^(1/2)/atanh(tanh(a + b*x)),x)`



output

```
(2*x^(1/2))/b + (2^(1/2)*log((b^(7/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) -
4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x))/((log
((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(
2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x
) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^
(1/2))/(2*b^(3/2))
```

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input

```
int(x^(1/2)/atanh(tanh(b*x+a)),x)
```

output

```
int(sqrt(x)/atanh(tanh(a + b*x)),x)
```

### 3.195 $\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1381
Mathematica [A] (verified)	1381
Rubi [A] (verified)	1382
Maple [A] (verified)	1383
Fricas [A] (verification not implemented)	1383
Sympy [F]	1384
Maxima [A] (verification not implemented)	1384
Giac [A] (verification not implemented)	1384
Mupad [B] (verification not implemented)	1385
Reduce [F]	1385

#### Optimal result

Integrand size = 15, antiderivative size = 53

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{b}\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}$$

output

$-2 \operatorname{arctanh}(b^{(1/2)} * x^{(1/2)} / (b * x - \operatorname{arctanh}(\tanh(b * x + a)))^{(1/2)}) / b^{(1/2)} / (b * x - \operatorname{arctanh}(\tanh(b * x + a)))^{(1/2)}$

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{b}\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}$$

input

`Integrate[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]),x]`

output

```
(2*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(Sqrt[b]*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx$$

↓ 2593

$$\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{b}\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}$$

input

```
Int[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]),x]
```

output

```
(-2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(Sqrt[b]*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])
```

**Defintions of rubi rules used**

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

**Maple [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$\frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$	41
default	$\frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$	41

input `int(1/x^(1/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `2/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.28

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \left[ -\frac{\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right)}{ab}, -\frac{2\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{ab} \right]$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `[-sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a))/(a*b), -2*sqrt(a*b)*arctan(sqrt(a*b)/(b*sqrt(x)))/(a*b)]`

**Sympy [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(1/x**(1/2)/atanh(tanh(b*x+a)),x)`

output `Integral(1/(sqrt(x)*atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.34

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`

output `2*arctan(b*sqrt(x)/sqrt(a*b))/sqrt(a*b)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.34

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`

output `2*arctan(b*sqrt(x)/sqrt(a*b))/sqrt(a*b)`

**Mupad [B] (verification not implemented)**

Time = 5.91 (sec) , antiderivative size = 347, normalized size of antiderivative = 6.55

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\sqrt{2} \ln \left( \frac{b^2 \left( \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2bx \right) \left( 2\sqrt{2}a + 4\sqrt{x} \sqrt{b \left( \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2bx \right) - \sqrt{2} \left( 2a - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) \right)}{2\sqrt{b \left( \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2bx \right) \left( \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) \right)}}{\sqrt{b \ln \left( \frac{1}{e^{2a} e^{2bx+1}} \right) - b \ln \left( \frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2b^2 x}} \right)}{2\sqrt{b \left( \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) + 2bx \right) \left( \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}} \right) - \ln \left( \frac{2}{e^{2a} e^{2bx+1}} \right) \right)}$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))),x)`output `(2^(1/2)*log((b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*(2*2^(1/2)*a + 4*x^(1/2)*(b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) - 2^(1/2)*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 2*2^(1/2)*b*x))/(2*(b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(b*log(1/(exp(2*a)*exp(2*b*x) + 1)) - b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b^2*x)^(1/2)`**Reduce [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(1/x^(1/2)/atanh(tanh(b*x+a)),x)`output `int(1/(sqrt(x)*atanh(tanh(a + b*x))),x)`

### 3.196 $\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1386
Mathematica [A] (verified)	1386
Rubi [A] (verified)	1387
Maple [A] (verified)	1388
Fricas [A] (verification not implemented)	1389
Sympy [F]	1389
Maxima [A] (verification not implemented)	1390
Giac [A] (verification not implemented)	1390
Mupad [B] (verification not implemented)	1390
Reduce [F]	1391

#### Optimal result

Integrand size = 15, antiderivative size = 76

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

$-2*b^{(1/2)}*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)})/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(3/2)}+2/x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2\sqrt{b} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{2}{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}$$

input

`Integrate[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]),x]`

output

```
(-2*Sqrt[b]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]
)/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2) - 2/(Sqrt[x]*(-(b*x) + ArcTanh[
Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2594$$

$$\frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2593$$

$$\frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2}}$$

input

```
Int[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]),x]
```

output

```
(-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])
/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a +
b*x]]))
```



## Defintions of rubi rules used

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

## Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} - \frac{2b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$	76
default	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} - \frac{2b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$	76

input

```
int(1/x^(3/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-2/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)-2*b/(arctanh(tanh(b*x+a))-b*x)/((arc  
tanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*  
x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.14

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{x \sqrt{-\frac{b}{a}} \log\left(\frac{bx - 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a}\right) - 2\sqrt{x}}{ax}, \right. \\ \left. - \frac{2\left(x\sqrt{\frac{b}{a}} \arctan\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + \sqrt{x}\right)}{ax} \right]$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")`output `[(x*sqrt(-b/a)*log((b*x - 2*a*sqrt(x)*sqrt(-b/a) - a)/(b*x + a)) - 2*sqrt(x))/(a*x), -2*(x*sqrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + sqrt(x))/(a*x)]`**Sympy [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{x^{3/2} \operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(1/x**(3/2)/atanh(tanh(b*x+a)),x)`output `Integral(1/(x**(3/2)*atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}} - \frac{2}{a\sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-2*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a) - 2/(a*sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}} - \frac{2}{a\sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `-2*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a) - 2/(a*sqrt(x))`**Mupad [B] (verification not implemented)**

Time = 3.95 (sec) , antiderivative size = 464, normalized size of antiderivative = 6.11

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{4}{\sqrt{x} \left( \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx \right)}$$

$$+ \frac{2\sqrt{2}\sqrt{b} \ln\left(\frac{\sqrt{b}\sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right)} - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx}{\sqrt{2}\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right) - 4\sqrt{b}\sqrt{x}\sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right)}}\right)}{2\left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right)} - \frac{2}{a\sqrt{x}}$$

input `int(1/(x^(3/2)*atanh(tanh(a + b*x))),x)`

output `4/(x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (2*2^(1/2)*b^(1/2)*log((b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 4*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 4*a^2))/(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2)`

### Reduce [F]

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x} dx$$

input `int(1/x^(3/2)/atanh(tanh(b*x+a)),x)`

output `int(1/(sqrt(x)*atanh(tanh(a + b*x))*x),x)`

### 3.197 $\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1392
Mathematica [A] (verified)	1392
Rubi [A] (verified)	1393
Maple [A] (verified)	1394
Fricas [A] (verification not implemented)	1395
Sympy [F]	1395
Maxima [A] (verification not implemented)	1396
Giac [A] (verification not implemented)	1396
Mupad [B] (verification not implemented)	1396
Reduce [F]	1397

#### Optimal result

Integrand size = 15, antiderivative size = 101

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} + \frac{2b}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
-2*b^(3/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-
arctanh(tanh(b*x+a)))^(5/2)+2*b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/3/x
^(3/2)/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx = \frac{2b^{3/2} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} + \frac{2(4bx - \operatorname{arctanh}(\tanh(a+bx)))}{3x^{3/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input `Integrate[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]),x]`

output 
$$\frac{(2*b^{(3/2)}*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^{(5/2)} + (2*(4*b*x - ArcTanh[Tanh[a + b*x]])))/(3*x^{(3/2)}*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2}$$

### Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.17, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx \\ & \quad \downarrow 2594 \\ & \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2594 \\ & \frac{b \left( \frac{\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \\ & \quad \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2593 \\ & \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \\ & \frac{b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} \end{aligned}$$

input `Int[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]),x]`

output 
$$\frac{2}{(3x^{3/2}(bx - \text{ArcTanh}[\text{Tanh}[a + bx]])) + (b((-2\sqrt{b})\text{ArcTanh}[\sqrt{b}\sqrt{x}]/\sqrt{bx - \text{ArcTanh}[\text{Tanh}[a + bx]]]))/(bx - \text{ArcTanh}[\text{Tanh}[a + bx]])^{3/2} + 2/(\sqrt{x}(bx - \text{ArcTanh}[\text{Tanh}[a + bx]])))/(bx - \text{ArcTanh}[\text{Tanh}[a + bx]])$$

### Defintions of rubi rules used

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

### Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2}{3(\text{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{2b}{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}} + \frac{2b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}}}\right)}{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}}$
default	$-\frac{2}{3(\text{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{2b}{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}} + \frac{2b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}}}\right)}{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{(\text{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}}$

input `int(1/x^(5/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-2/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)+2/(arctanh(tanh(b*x+a))-b*x)^2*b/x
^(1/2)+2*b^2/(arctanh(tanh(b*x+a))-b*x)^2/((arctanh(tanh(b*x+a))-b*x)*b)^(
1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{3bx^2 \sqrt{-\frac{b}{a}} \log\left(\frac{bx+2a\sqrt{x}\sqrt{-\frac{b}{a}}-a}{bx+a}\right) + 2(3bx-a)\sqrt{x}}{3a^2x^2}, \frac{2\left(3bx^2\sqrt{\frac{b}{a}} \operatorname{arctan}\left(\frac{\sqrt{x}}{\sqrt{b/a}}\right) + (3bx-a)\sqrt{x}\right)}{3a^2x^2} \right]$$

input

```
integrate(1/x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")
```

output

```
[1/3*(3*b*x^2*sqrt(-b/a)*log((b*x + 2*a*sqrt(x)*sqrt(-b/a) - a)/(b*x + a))
+ 2*(3*b*x - a)*sqrt(x))/(a^2*x^2), 2/3*(3*b*x^2*sqrt(b/a)*arctan(sqrt(x)
*sqrt(b/a)) + (3*b*x - a)*sqrt(x))/(a^2*x^2)]
```

**Sympy [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{x^{5/2} \operatorname{atanh}(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(5/2)/atanh(tanh(b*x+a)),x)
```

output

```
Integral(1/(x**(5/2)*atanh(tanh(a + b*x))), x)
```



**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}a^2} + \frac{2(3bx - a)}{3a^2x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `2*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2) + 2/3*(3*b*x - a)/(a^2*x^(3/2))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \frac{2b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}a^2} + \frac{2(3bx - a)}{3a^2x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `2*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2) + 2/3*(3*b*x - a)/(a^2*x^(3/2))`**Mupad [B] (verification not implemented)**

Time = 3.86 (sec) , antiderivative size = 642, normalized size of antiderivative = 6.36

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \text{Too large to display}$$

input `int(1/(x^(5/2)*atanh(tanh(a + b*x))),x)`

output

```

4/(3*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
)))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (8*b)/(x^(1/2)*(log(2/(exp(2*a)*
exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
+ 2*b*x)^2) + (4*2^(1/2)*b^(3/2)*log((b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1
/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
)))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)
)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)
) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp
(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 + 24*
a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/
(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 16*a^4 - 8*a*(2*a - log((2*exp(2*a)
)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)
) + 2*b*x)^3 - 32*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b
*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*(log((2*exp(2*a)
)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)
))))/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2)

```

**Reduce [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x^2} dx$$

input

```
int(1/x^(5/2)/atanh(tanh(b*x+a)),x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))*x**2),x)
```

**3.198**       $\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx$

Optimal result	1398
Mathematica [A] (verified)	1399
Rubi [A] (verified)	1399
Maple [A] (verified)	1401
Fricas [A] (verification not implemented)	1402
Sympy [F]	1402
Maxima [A] (verification not implemented)	1403
Giac [A] (verification not implemented)	1403
Mupad [B] (verification not implemented)	1403
Reduce [F]	1404

**Optimal result**

Integrand size = 15, antiderivative size = 128

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx = -\frac{2b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} + \frac{2b^2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{2b}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
-2*b^(5/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-
arctanh(tanh(b*x+a)))^(7/2)+2*b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3+2/3
*b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/5/x^(5/2)/(b*x-arctanh(tanh(b*x+
a)))
```

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.84

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b^{5/2} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{7/2}} + \frac{2(23b^2x^2 - 11bx \operatorname{arctanh}(\tanh(a + bx)) + 3 \operatorname{arctanh}(\tanh(a + bx))^2)}{15x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))^3}$$

input

```
Integrate[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]), x]
```

output

```
(-2*b^(5/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(7/2) + (2*(23*b^2*x^2 - 11*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2))/(15*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.25, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2594$$

$$\frac{b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

$$\downarrow 2594$$

$$\begin{aligned}
 & \frac{b \left( \frac{\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2}} + \\
 & \frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2594} \\
 & \frac{b \left( \frac{b \left( \frac{\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2}} + \\
 & \frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2593} \\
 & \frac{b \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{\frac{bx - \operatorname{arctanh}(\tanh(a+bx))}{2}} + \\
 & \frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2}
 \end{aligned}$$

input `Int[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]),x]`

output `2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]])])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))))/(b*x - ArcTanh[Tanh[a + b*x]]))/(b*x - ArcTanh[Tanh[a + b*x]])`

## Defintions of rubi rules used

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

## Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} - \frac{2b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}} + \frac{2b}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2x^{\frac{3}{2}}} - \frac{2b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}}$
default	$-\frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} - \frac{2b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}} + \frac{2b}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2x^{\frac{3}{2}}} - \frac{2b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}}$

input

```
int(1/x^(7/2)/arctanh(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-2/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)-2/(arctanh(tanh(b*x+a))-b*x)^3*b^2/x^(1/2)+2/3/(arctanh(tanh(b*x+a))-b*x)^2*b/x^(3/2)-2*b^3/(arctanh(tanh(b*x+a))-b*x)^3/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.09

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{15 b^2 x^3 \sqrt{-\frac{b}{a}} \log\left(\frac{bx - 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a}\right) - 2(15 b^2 x^2 - 5 abx + 3 a^2)\sqrt{x}}{15 a^3 x^3}, \right. \\ \left. - \frac{2\left(15 b^2 x^3 \sqrt{\frac{b}{a}} \arctan\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + (15 b^2 x^2 - 5 abx + 3 a^2)\sqrt{x}\right)}{15 a^3 x^3} \right]$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="fricas")`

output `[1/15*(15*b^2*x^3*sqrt(-b/a)*log((b*x - 2*a*sqrt(x)*sqrt(-b/a) - a)/(b*x + a)) - 2*(15*b^2*x^2 - 5*a*b*x + 3*a^2)*sqrt(x))/(a^3*x^3), -2/15*(15*b^2*x^3*sqrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + (15*b^2*x^2 - 5*a*b*x + 3*a^2)*sqrt(x))/(a^3*x^3)]`

**Sympy [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{x^{7/2} \operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(1/x**(7/2)/atanh(tanh(b*x+a)),x)`

output `Integral(1/(x**(7/2)*atanh(tanh(a + b*x))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba^3}} - \frac{2(15b^2x^2 - 5abx + 3a^2)}{15a^3x^{5/2}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="maxima")`output `-2*b^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3) - 2/15*(15*b^2*x^2 - 5*a*b*x + 3*a^2)/(a^3*x^(5/2))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = -\frac{2b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba^3}} - \frac{2(15b^2x^2 - 5abx + 3a^2)}{15a^3x^{5/2}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a)),x, algorithm="giac")`output `-2*b^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3) - 2/15*(15*b^2*x^2 - 5*a*b*x + 3*a^2)/(a^3*x^(5/2))`**Mupad [B] (verification not implemented)**

Time = 3.57 (sec) , antiderivative size = 822, normalized size of antiderivative = 6.42

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \text{Too large to display}$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))),x)`



output

```

4/(5*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
))/((exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (8*b)/(3*x^(3/2)*(log(2/(exp(2*a)
)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)
) + 2*b*x)^2) + (16*b^2)/(x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log(
(2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (8*2^(1/2)
)*b^(5/2)*log((b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)
)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(ex
p(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - lo
g((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(
1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1))
+ log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^6 + 60*a^2*(2*a - log((2*exp(2
*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) +
1)) + 2*b*x)^4 - 160*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(
2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 + 240*a^4*(2*a
- log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)
)*exp(2*b*x) + 1)) + 2*b*x)^2 + 64*a^6 - 12*a*(2*a - log((2*exp(2*a)*exp(2*
b*x))/((exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*
x)^5 - 192*a^5*(2*a - log((2*exp(2*a)*exp(2*b*x))/((exp(2*a)*exp(2*b*x) + 1
)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(2*(log((2*exp(2*a)*ex...

```

**Reduce [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a)) x^3} dx$$

input

```
int(1/x^(7/2)/atanh(tanh(b*x+a)),x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))*x**3),x)
```

**3.199**       $\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1405
Mathematica [A] (verified)	1406
Rubi [A] (verified)	1406
Maple [B] (verified)	1409
Fricas [A] (verification not implemented)	1410
Sympy [F(-1)]	1410
Maxima [A] (verification not implemented)	1411
Giac [A] (verification not implemented)	1411
Mupad [B] (verification not implemented)	1412
Reduce [F]	1412

**Optimal result**

Integrand size = 15, antiderivative size = 135

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{7x^{5/2}}{5b^2} + \frac{7x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{3b^3} + \frac{7\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{b^4} - \frac{7\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}}{b^{9/2}} - \frac{x^{7/2}}{b\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
7/5*x^(5/2)/b^2+7/3*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))/b^3+7*x^(1/2)*(b*x-
arctanh(tanh(b*x+a)))^2/b^4-7*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*
x+a))))^(1/2)*(b*x-arctanh(tanh(b*x+a)))^(5/2)/b^(9/2)-x^(7/2)/b/arctanh(t
anh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.07

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2x^{5/2}}{5b^2} - \frac{4x^{3/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))}{3b^3} + \frac{6\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}{b^4} - \frac{7 \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right) (-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}}{b^{9/2}} + \frac{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3}{b^4 \operatorname{arctanh}(\tanh(a + bx))}$$

input

```
Integrate[x^(7/2)/ArcTanh[Tanh[a + b*x]]^2,x]
```

output

```
(2*x^(5/2))/(5*b^2) - (4*x^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/(3*b^3) + (6*Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^2)/b^4 - (7*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^(5/2))/b^(9/2) + (Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^3)/(b^4*ArcTanh[Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.12, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2590, 2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

↓ 2599

$$\frac{7 \int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))} dx}{2b} - \frac{x^{7/2}}{b \operatorname{arctanh}(\tanh(a + bx))}$$

$$\begin{aligned}
 & \downarrow 2590 \\
 & \frac{7 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2x^{5/2}}{5b}}{b} \right)}{2b} - \frac{x^{7/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \downarrow 2590 \\
 & \frac{7 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2x^{3/2}}{3b}}{b} \right) + \frac{2x^{5/2}}{5b}}{b} \right)}{2b} - \frac{x^{7/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \downarrow 2590 \\
 & \frac{7 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2\sqrt{x}}{b}}{b} \right) + \frac{2x^{3/2}}{3b}}{b} \right) + \frac{2x^{5/2}}{5b}}{b} \right)}{2b} - \frac{x^{7/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \downarrow 2593
 \end{aligned}$$

$$\frac{\left( \frac{\left( \frac{2\sqrt{x}}{b} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}} \right) \sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)^2 + \frac{2x^{3/2}}{3b} \right) \operatorname{arctanh}(\tanh(a+bx))}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

```
input Int [x^(7/2)/ArcTanh[Tanh[a + b*x]]^2,x]
```

```
output (7*((2*x^(5/2))/(5*b) + ((2*x^(3/2))/(3*b) + ((2*Sqrt[x])/b - (2*ArcTanh
[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]*Sqrt[b*x - ArcTanh[
Tanh[a + b*x]]])/b^(3/2))*(b*x - ArcTanh[Tanh[a + b*x]]))/b)*(b*x - ArcTan
h[Tanh[a + b*x]]))/b)/(2*b) - x^(7/2)/(b*ArcTanh[Tanh[a + b*x]])
```

**Defintions of rubi rules used**

```
rule 2590 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x
] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n,
1]
```

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simpli
fy[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u -
a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; Piecewise
LinearQ[u, v, x]
```

rule 2599

```

Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))

```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs.  $2(113) = 226$ .

Time = 1.64 (sec) , antiderivative size = 280, normalized size of antiderivative = 2.07

method	result
derivativedivides	$\frac{\frac{2b^2x^{\frac{5}{2}}}{5} - \frac{4bx^{\frac{3}{2}}a}{3} - \frac{4bx^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{3} + 6a^2\sqrt{x} + 12a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x} + 6(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^4}$
default	$\frac{2b^2x^{\frac{5}{2}}}{5} - \frac{4bx^{\frac{3}{2}}a}{3} - \frac{4bx^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{3} + 6a^2\sqrt{x} + 12a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x} + 6(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{b^4}$
risch	Expression too large to display

input

```
int(x^(7/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```

2/b^4*(1/5*b^2*x^(5/2)-2/3*b*x^(3/2)*a-2/3*b*x^(3/2)*(arctanh(tanh(b*x+a))-b*x-a)+3*a^2*x^(1/2)+6*a*(arctanh(tanh(b*x+a))-b*x-a)*x^(1/2)+3*(arctanh(tanh(b*x+a))-b*x-a)^2*x^(1/2))-2/b^4*((-1/2*a^3-3/2*a^2*(arctanh(tanh(b*x+a))-b*x-a)-3/2*a*(arctanh(tanh(b*x+a))-b*x-a)^2-1/2*(arctanh(tanh(b*x+a))-b*x-a)^3)*x^(1/2)/arctanh(tanh(b*x+a))+7/2*(a^3+3*a^2*(arctanh(tanh(b*x+a))-b*x-a)+3*a*(arctanh(tanh(b*x+a))-b*x-a)^2+(arctanh(tanh(b*x+a))-b*x-a)^3)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))

```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.39

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \left[ \frac{105 (a^2bx + a^3) \sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(6b^3x^3 - 14ab^2x^2 + 70a^2bx + 105a^3)\sqrt{x}}{30(b^5x + ab^4)} \right. \\ \left. - \frac{105(a^2bx + a^3)\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (6b^3x^3 - 14ab^2x^2 + 70a^2bx + 105a^3)\sqrt{x}}{15(b^5x + ab^4)} \right]$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `[1/30*(105*(a^2*b*x + a^3)*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(6*b^3*x^3 - 14*a*b^2*x^2 + 70*a^2*b*x + 105*a^3)*sqrt(x))/ (b^5*x + a*b^4), -1/15*(105*(a^2*b*x + a^3)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - (6*b^3*x^3 - 14*a*b^2*x^2 + 70*a^2*b*x + 105*a^3)*sqrt(x))/ (b^5*x + a*b^4)]`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Timed out}$$

input `integrate(x**(7/2)/atanh(tanh(b*x+a))**2,x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.56

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{6b^3x^{7/2} - 14ab^2x^{5/2} + 70a^2bx^{3/2} + 105a^3\sqrt{x}}{15(b^5x + ab^4)} - \frac{7a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^4}}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `1/15*(6*b^3*x^(7/2) - 14*a*b^2*x^(5/2) + 70*a^2*b*x^(3/2) + 105*a^3*sqrt(x))/ (b^5*x + a*b^4) - 7*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^4)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.56

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{7a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^4}} + \frac{a^3\sqrt{x}}{(bx+a)b^4} + \frac{2\left(3b^8x^{5/2} - 10ab^7x^{3/2} + 45a^2b^6\sqrt{x}\right)}{15b^{10}}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `-7*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^4) + a^3*sqrt(x)/((b*x + a)*b^4) + 2/15*(3*b^8*x^(5/2) - 10*a*b^7*x^(3/2) + 45*a^2*b^6*sqrt(x))/b^10`



**Mupad [B] (verification not implemented)**

Time = 3.51 (sec) , antiderivative size = 523, normalized size of antiderivative = 3.87

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(x^(7/2)/atanh(tanh(a + b*x))^2,x)`

output

```
(2*x^(5/2))/(5*b^2) + (2*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(3*b^3) + (3*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*b^4) + (7*2^(1/2)*log((64*b^(19/2)*2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2))/(16*b^(9/2)) - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(4*b^4*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))
```

**Reduce [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{\sqrt{x} x^3}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^(7/2)/atanh(tanh(b*x+a))^2,x)`

output

```
int((sqrt(x)*x**3)/atanh(tanh(a + b*x))**2,x)
```

### 3.200 $\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1413
Mathematica [A] (verified)	1414
Rubi [A] (verified)	1414
Maple [A] (verified)	1416
Fricas [A] (verification not implemented)	1417
Sympy [F]	1417
Maxima [A] (verification not implemented)	1418
Giac [A] (verification not implemented)	1418
Mupad [B] (verification not implemented)	1419
Reduce [F]	1420

#### Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{5x^{3/2}}{3b^2} + \frac{5\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^3} - \frac{5\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}}{b^{7/2}} - \frac{x^{5/2}}{b\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
5/3*x^(3/2)/b^2+5*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))/b^3-5*arctanh(b^(1/2)
*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(3/2
)/b^(7/2)-x^(5/2)/b/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.10

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{2x^{3/2}}{3b^2} - \frac{4\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}{b^3}$$

$$+ \frac{5 \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right) (-bx + \operatorname{arctanh}(\tanh(a+bx)))^{3/2}}{b^{7/2}}$$

$$- \frac{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}{b^3 \operatorname{arctanh}(\tanh(a+bx))}$$

input `Integrate[x^(5/2)/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(2*x^(3/2))/(3*b^2) - (4*Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))/b^3 + (5*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]^(3/2))/b^(7/2) - (Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))^2/(b^3*ArcTanh[Tanh[a + b*x]])`

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.12, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$\downarrow 2599$$

$$\frac{5 \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))}$$

$$\downarrow 2590$$

$$\begin{aligned}
 & \frac{5 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2x^{3/2}}{3b}}{b} \right)}{2b} - \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2590} \\
 & \frac{5 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2\sqrt{x}}{b}}{b} \right)}{b} + \frac{2x^{3/2}}{3b} \right)}{2b} \\
 & \quad \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2593} \\
 & \frac{5 \left( \frac{\left( \frac{2 \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{\frac{2\sqrt{x}}{b} - \frac{2 \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}}} \right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b} + \frac{2x^{3/2}}{3b} \right)}{x^{5/2}} \\
 & \quad \frac{2b}{b \operatorname{arctanh}(\tanh(a+bx))}
 \end{aligned}$$

input `Int [x^(5/2)/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(5*((2*x^(3/2))/(3*b) + (((2*Sqrt[x])/b - (2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2))*(b*x - ArcTanh[Tanh[a + b*x]]))/b)/(2*b) - x^(5/2)/(b*ArcTanh[Tanh[a + b*x]])`

Defintions of rubi rules used

```
rule 2590 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x
] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n,
1]
```

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simpli
fy[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u -
a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; Piecewise
LinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

Maple [A] (verified)

Time = 1.28 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.68

method	result
derivativedivides	$\frac{2\left(-\frac{a^2}{2}-a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)-\frac{(\operatorname{arctanh}(\tanh(\frac{bx+a}{2}))-bx-a)^2}{2}\right)\sqrt{x}}{\operatorname{arctanh}(\tanh(bx+a))} + \frac{5\left(a^2+2a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)+(\operatorname{arctanh}(\tanh(\frac{bx+a}{2}))-bx-a)^2\right)\sqrt{x}}{b^3}$
default	$\frac{2\left(-\frac{a^2}{2}-a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)-\frac{(\operatorname{arctanh}(\tanh(\frac{bx+a}{2}))-bx-a)^2}{2}\right)\sqrt{x}}{\operatorname{arctanh}(\tanh(bx+a))} + \frac{5\left(a^2+2a(\operatorname{arctanh}(\tanh(bx+a))-bx-a)+(\operatorname{arctanh}(\tanh(\frac{bx+a}{2}))-bx-a)^2\right)\sqrt{x}}{b^3}$
risch	Expression too large to display

```
input int(x^(5/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
2/b^3*((-1/2*a^2-a*(arctanh(tanh(b*x+a))-b*x-a)-1/2*(arctanh(tanh(b*x+a))-
b*x-a)^2)*x^(1/2)/arctanh(tanh(b*x+a))+5/2*(a^2+2*a*(arctanh(tanh(b*x+a))-
b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2
)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))-2/b^3*(-1/3*b*x^
(3/2)+2*a*x^(1/2)+2*(arctanh(tanh(b*x+a))-b*x-a)*x^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.49

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{15(abx + a^2)\sqrt{-\frac{a}{b}} \log\left(\frac{bx + 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(2b^2x^2 - 10abx - 15a^2)\sqrt{x}}{6(b^4x + ab^3)}$$

input

```
integrate(x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
[1/6*(15*(a*b*x + a^2)*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(
b*x + a)) + 2*(2*b^2*x^2 - 10*a*b*x - 15*a^2)*sqrt(x))/(b^4*x + a*b^3), 1/
3*(15*(a*b*x + a^2)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) + (2*b^2*x^2 -
10*a*b*x - 15*a^2)*sqrt(x))/(b^4*x + a*b^3)]
```

**Sympy [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^{5/2}}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input

```
integrate(x**(5/2)/atanh(tanh(b*x+a))**2,x)
```

output

```
Integral(x**(5/2)/atanh(tanh(a + b*x))**2, x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.59

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2b^2x^{5/2} - 10abx^{3/2} - 15a^2\sqrt{x}}{3(b^4x + ab^3)} + \frac{5a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `1/3*(2*b^2*x^(5/2) - 10*a*b*x^(3/2) - 15*a^2*sqrt(x))/(b^4*x + a*b^3) + 5*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.60

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{5a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} - \frac{a^2\sqrt{x}}{(bx + a)b^3} + \frac{2(b^4x^{3/2} - 6ab^3\sqrt{x})}{3b^6}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `5*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) - a^2*sqrt(x)/((b*x + a)*b^3) + 2/3*(b^4*x^(3/2) - 6*a*b^3*sqrt(x))/b^6`

**Mupad [B] (verification not implemented)**

Time = 3.46 (sec) , antiderivative size = 463, normalized size of antiderivative = 4.29

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2x^{3/2}}{3b^2} + \frac{2\sqrt{x} \left( \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx \right)}{b^3} + \frac{5\sqrt{2} \ln\left(\frac{16b^{15/2} \left(\sqrt{2} \left(\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx\right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx + 2\sqrt{2}bx}\right)}{\left(\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right)\right) \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx}}\right)}{8b^{7/2}} - \frac{\sqrt{x} \left( \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) + 2bx \right)^2}{2b^3 \left( \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) \right)}$$

input `int(x^(5/2)/atanh(tanh(a + b*x))^2,x)`

output

```
(2*x^(3/2))/(3*b^2) + (2*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b^3 + (5*2^(1/2)*log((16*b^(15/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x))/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2))/(8*b^(7/2)) - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(2*b^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))
```



**Reduce [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{\sqrt{x} x^2}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a))^2,x)`

output `int((sqrt(x)*x**2)/atanh(tanh(a + b*x))**2,x)`

### 3.201 $\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1421
Mathematica [A] (verified)	1421
Rubi [A] (verified)	1422
Maple [A] (verified)	1423
Fricas [A] (verification not implemented)	1424
Sympy [F]	1424
Maxima [A] (verification not implemented)	1425
Giac [A] (verification not implemented)	1425
Mupad [B] (verification not implemented)	1425
Reduce [F]	1426

#### Optimal result

Integrand size = 15, antiderivative size = 83

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{3\sqrt{x}}{b^2} - \frac{3\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}{b^{5/2}} - \frac{x^{3/2}}{b\operatorname{arctanh}(\tanh(a+bx))}$$

output

$3*x^{(1/2)}/b^2-3*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)})*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)}/b^{(5/2)}-x^{(3/2)}/b/\operatorname{arctanh}(\tanh(b*x+a))$

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.98

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{3\sqrt{x}}{b^2} - \frac{x^{3/2}}{b\operatorname{arctanh}(\tanh(a+bx))} - \frac{3\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}\right)\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}{b^{5/2}}$$

input `Integrate[x^(3/2)/ArcTanh[Tanh[a + b*x]]^2,x]`

output `(3*sqrt[x])/b^2 - x^(3/2)/(b*ArcTanh[Tanh[a + b*x]]) - (3*ArcTan[(sqrt[b]*sqrt[x])/sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])/b^(5/2)`

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.10, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{3 \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a + bx))} \\
 & \quad \downarrow \text{2590} \\
 & \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{2\sqrt{x}}{b} \right)}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a + bx))} \\
 & \quad \downarrow \text{2593} \\
 & \frac{3 \left( \frac{2\sqrt{x}}{b} - \frac{2 \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}} \right)}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a + bx))}
 \end{aligned}$$

input `Int[x^(3/2)/ArcTanh[Tanh[a + b*x]]^2,x]`

```
output (3*((2*Sqrt[x])/b - (2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2))/(2*b) - x^(3/2)/(b*ArcTanh[Tanh[a + b*x]])
```

**Defintions of rubi rules used**

```
rule 2590 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]
```

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 1.15 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b^2} - \frac{2 \left( \frac{\left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{2} + \frac{bx}{2} \right) \sqrt{x}}{\operatorname{arctanh}(\tanh(bx+a))} + \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx) \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}} \right)}{b^2}$	95
default	$\frac{2\sqrt{x}}{b^2} - \frac{2 \left( \frac{\left( -\frac{\operatorname{arctanh}(\tanh(bx+a))}{2} + \frac{bx}{2} \right) \sqrt{x}}{\operatorname{arctanh}(\tanh(bx+a))} + \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx) \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}} \right)}{b^2}$	95
risch	Expression too large to display	10

input `int(x^(3/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output `2*x^(1/2)/b^2-2/b^2*((-1/2*arctanh(tanh(b*x+a))+1/2*b*x)*x^(1/2)/arctanh(tanh(b*x+a))+3/2*(arctanh(tanh(b*x+a))-b*x)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.61

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \left[ \frac{3(bx + a)\sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(2bx + 3a)\sqrt{x}}{2(b^3x + ab^2)}, \right. \\ \left. - \frac{3(bx + a)\sqrt{\frac{a}{b}} \operatorname{arctan}\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (2bx + 3a)\sqrt{x}}{b^3x + ab^2} \right]$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `[1/2*(3*(b*x + a)*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(2*b*x + 3*a)*sqrt(x))/(b^3*x + a*b^2), -(3*(b*x + a)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - (2*b*x + 3*a)*sqrt(x))/(b^3*x + a*b^2)]`

### Sympy [F]

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{x^{3/2}}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**(3/2)/atanh(tanh(b*x+a))**2,x)`

output `Integral(x**(3/2)/atanh(tanh(a + b*x))**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.60

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2bx^{3/2} + 3a\sqrt{x}}{b^3x + ab^2} - \frac{3a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output  $(2*b*x^{(3/2)} + 3*a*\sqrt{x})/(b^3*x + a*b^2) - 3*a*\arctan(b*\sqrt{x}/\sqrt{a*b})/(\sqrt{a*b}*b^2)$ **Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.55

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{3a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{a\sqrt{x}}{(bx + a)b^2} + \frac{2\sqrt{x}}{b^2}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output  $-3*a*\arctan(b*\sqrt{x}/\sqrt{a*b})/(\sqrt{a*b}*b^2) + a*\sqrt{x}/((b*x + a)*b^2) + 2*\sqrt{x}/b^2$ **Mupad [B] (verification not implemented)**

Time = 3.90 (sec) , antiderivative size = 403, normalized size of antiderivative = 4.86

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{2\sqrt{x}}{b^2} - \frac{\sqrt{x} \left( \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx \right)}{b^2 \left( \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) \right)}$$

$$+ \frac{3\sqrt{2} \ln\left( \frac{4b^{11/2} \left( \sqrt{2} \left( \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx \right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx + 2\sqrt{2}bx} \right)}{\left( \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) \right) \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx}} \right)}{4b^{5/2}}$$

input `int(x^(3/2)/atanh(tanh(a + b*x))^2,x)`

output `(2*x^(1/2))/b^2 - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x))/(b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) + (3*2^(1/2)*log((4*b^(11/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x))/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2)))/(4*b^(5/2))`

### Reduce [F]

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{\sqrt{x} x}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^(3/2)/atanh(tanh(b*x+a))^2,x)`

output `int((sqrt(x)*x)/atanh(tanh(a + b*x))*2,x)`

### 3.202 $\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1427
Mathematica [A] (verified)	1427
Rubi [A] (verified)	1428
Maple [A] (verified)	1429
Fricas [A] (verification not implemented)	1430
Sympy [F]	1430
Maxima [A] (verification not implemented)	1430
Giac [A] (verification not implemented)	1431
Mupad [B] (verification not implemented)	1431
Reduce [F]	1432

#### Optimal result

Integrand size = 15, antiderivative size = 73

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}} - \frac{\sqrt{x}}{b\operatorname{arctanh}(\tanh(a+bx))}$$

output

$$-\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)})/b^{(3/2)/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(1/2)}-x^{(1/2)/b/\operatorname{arctanh}(\tanh(b*x+a))}$$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{\sqrt{x}}{b\operatorname{arctanh}(\tanh(a+bx))} + \frac{\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}$$



input `Integrate[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(Sqrt[x]/(b*ArcTanh[Tanh[a + b*x]])) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]/(b^(3/2)*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])`

### Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {2599, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

↓ 2599

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))} dx - \frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(a + bx))}$$

↓ 2593

$$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}}\right)}{b^{3/2}\sqrt{bx - \operatorname{arctanh}(\tanh(a + bx))}} - \frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(a + bx))}$$

input `Int[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^2,x]`

output `-(ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]/(b^(3/2)*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]) - Sqrt[x]/(b*ArcTanh[Tanh[a + b*x]])`

## Definitions of rubi rules used

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$-\frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(bx+a))} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}b}\right)}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}b}$	61
default	$-\frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(bx+a))} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}b}\right)}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}b}$	61
risch	Expression too large to display	807

input `int(x^(1/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

output `-x^(1/2)/b/arctanh(tanh(b*x+a))+1/b/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.58

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \left[ -\frac{2ab\sqrt{x} + \sqrt{-ab}(bx + a) \log\left(\frac{bx - a - 2\sqrt{-ab}\sqrt{x}}{bx + a}\right)}{2(ab^3x + a^2b^2)}, \right. \\ \left. -\frac{ab\sqrt{x} + \sqrt{ab}(bx + a) \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{ab^3x + a^2b^2} \right]$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`output `[-1/2*(2*a*b*sqrt(x) + sqrt(-a*b)*(b*x + a)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)))/(a*b^3*x + a^2*b^2), -(a*b*sqrt(x) + sqrt(a*b)*(b*x + a)*arctan(sqrt(a*b)/(b*sqrt(x))))/(a*b^3*x + a^2*b^2)]`**Sympy [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a))**2,x)`output `Integral(sqrt(x)/atanh(tanh(a + b*x))**2, x)`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.51

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{\sqrt{x}}{b^2x + ab} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `-sqrt(x)/(b^2*x + a*b) + arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.49

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} - \frac{\sqrt{x}}{(bx + a)b}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) - sqrt(x)/((b*x + a)*b)`

### Mupad [B] (verification not implemented)

Time = 3.93 (sec) , antiderivative size = 344, normalized size of antiderivative = 4.71

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \frac{\sqrt{2} \ln \left( \frac{b^{7/2} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)} - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx \left( \sqrt{2} \left( \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx \right) - 4\sqrt{b}\sqrt{x} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)} - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) \right)}{\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)} \right)}{2b^{3/2} \sqrt{\ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) + 2bx}} - \frac{2\sqrt{x}}{b \left( \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) \right)}$$

input `int(x^(1/2)/atanh(tanh(a + b*x))^2,x)`

output

```
(2^(1/2)*log((b^(7/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*
exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp
(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log
((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(
1/2)*b*x))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2
/(exp(2*a)*exp(2*b*x) + 1)))))/(2*b^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)
) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2))
- (2*x^(1/2))/(b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
- log(2/(exp(2*a)*exp(2*b*x) + 1))))
```

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\operatorname{atanh}(\tanh(bx + a)) \left( \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx+a))} dx \right) - 2\sqrt{x}}{2 \operatorname{atanh}(\tanh(bx + a)) b}$$

input

```
int(x^(1/2)/atanh(tanh(b*x+a))^2,x)
```

output

```
(atanh(tanh(a + b*x))*int(1/(sqrt(x)*atanh(tanh(a + b*x))),x) - 2*sqrt(x))
/(2*atanh(tanh(a + b*x))*b)
```

### 3.203 $\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1433
Mathematica [A] (verified)	1434
Rubi [A] (verified)	1434
Maple [A] (verified)	1436
Fricas [A] (verification not implemented)	1436
Sympy [F]	1437
Maxima [A] (verification not implemented)	1437
Giac [A] (verification not implemented)	1438
Mupad [B] (verification not implemented)	1438
Reduce [F]	1439

#### Optimal result

Integrand size = 15, antiderivative size = 97

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{b}(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{1}{b\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))}$$

output

```
arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/b^(1/2)/(b*x-arc
tanh(tanh(b*x+a)))^(3/2)-1/b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))-1/b/x^(1/2
)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$= \frac{\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{\sqrt{b}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} + \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Integrate[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2),x]`

output `ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]/(Sqrt[b]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2)) + Sqrt[x]/(ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow \text{2599}$$

$$-\frac{\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))} dx}{2b} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow \text{2594}$$

$$\begin{aligned}
& -\frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} \\
& \quad \downarrow \text{2593} \\
& -\frac{\frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}}{\frac{2b}{1}} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \\
& \quad \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))}
\end{aligned}$$

input `Int[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2), x]`

output `-1/2*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])))/b - 1/(b*Sqrt[x]*ArcTanh[Tanh[a + b*x]])`

### Defintions of rubi rules used

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v)))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`



rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [A] (verified)

Time = 1.50 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$
default	$\frac{\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}$
risch	Expression too large to display

input

```
int(1/x^(1/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))+1/(arctanh(tanh(b*x+a))-b*x)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.20

$$\int \frac{1}{\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^2} dx = \left[ \frac{2ab\sqrt{x} - \sqrt{-ab}(bx + a) \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right)}{2(a^2b^2x + a^3b)}, \frac{ab\sqrt{x} - \sqrt{ab}(bx + a) \operatorname{arctan}\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{a^2b^2x + a^3b} \right]$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `[1/2*(2*a*b*sqrt(x) - sqrt(-a*b)*(b*x + a)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)))/(a^2*b^2*x + a^3*b), (a*b*sqrt(x) - sqrt(a*b)*(b*x + a)*arctan(sqrt(a*b)/(b*sqrt(x)))/(a^2*b^2*x + a^3*b)]`

### Sympy [F]

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(1/x**(1/2)/atanh(tanh(b*x+a))**2,x)`

output `Integral(1/(sqrt(x)*atanh(tanh(a + b*x))**2), x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.36

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\sqrt{x}}{abx + a^2} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `sqrt(x)/(a*b*x + a^2) + arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.36

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}} + \frac{\sqrt{x}}{(bx + a)a}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a) + sqrt(x)/((b*x + a)*a)`

**Mupad [B] (verification not implemented)**

Time = 4.30 (sec) , antiderivative size = 516, normalized size of antiderivative = 5.32

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))^2),x)`

output `(2^(1/2)*log(-(b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*(4*a^2*b + b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - 4*a*b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2)) - (4*x^(1/2))/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))`

Reduce [F]

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(1/x^(1/2)/atanh(tanh(b*x+a))^2,x)`

output `int(1/(sqrt(x)*atanh(tanh(a + b*x))**2),x)`

### 3.204 $\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1440
Mathematica [A] (verified)	1441
Rubi [A] (verified)	1441
Maple [A] (verified)	1443
Fricas [A] (verification not implemented)	1444
Sympy [F]	1444
Maxima [A] (verification not implemented)	1445
Giac [A] (verification not implemented)	1445
Mupad [B] (verification not implemented)	1445
Reduce [F]	1446

#### Optimal result

Integrand size = 15, antiderivative size = 120

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{3\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} - \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{1} - \frac{1}{bx^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))}$$

output  $3*b^{(1/2)}*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a))))^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{(5/2)}-3/x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2-1/b/x^{(3/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))-1/b/x^{(3/2)}/\operatorname{arctanh}(\tanh(b*x+a))$

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.87

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{3\sqrt{b} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} - \frac{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}{b\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Integrate[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^2),x]`

output `(-3*Sqrt[b]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2) - 2/(Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))^2 - (b*Sqrt[x])/(ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))^2)`

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx$$

$$\downarrow 2599$$

$$-\frac{3 \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))} dx}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a + bx))}$$

$$\downarrow 2594$$

$$\frac{3 \left( \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$\frac{3 \left( \frac{b \left( \frac{\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2593

$$\frac{3 \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{1} - \frac{2b}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))}$$

input `Int [1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^2), x]`

output `(-3*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]))^3/2 + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))))/(b*x - ArcTanh[Tanh[a + b*x]]))/(2*b) - 1/(b*x^(3/2)*ArcTanh[Tanh[a + b*x]])`

Defintions of rubi rules used

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2594 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}} - \frac{2b\left(\frac{\sqrt{x}}{2\operatorname{arctanh}(\tanh(bx+a))} + \frac{3\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2}$	93
default	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}} - \frac{2b\left(\frac{\sqrt{x}}{2\operatorname{arctanh}(\tanh(bx+a))} + \frac{3\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2}$	93
risch	Expression too large to display	1584

```
input int(1/x^(3/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```



output

```
-2/(arctanh(tanh(b*x+a))-b*x)^2/x^(1/2)-2/(arctanh(tanh(b*x+a))-b*x)^2*b*(
1/2*x^(1/2)/arctanh(tanh(b*x+a))+3/2/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*
arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \left[ \frac{3(bx^2 + ax) \sqrt{-\frac{b}{a}} \log\left(\frac{bx - 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a}\right) - 2(3bx + 2a)\sqrt{x}}{2(a^2bx^2 + a^3x)}, \right. \\ \left. - \frac{3(bx^2 + ax) \sqrt{\frac{b}{a}} \arctan\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + (3bx + 2a)\sqrt{x}}{a^2bx^2 + a^3x} \right]$$

input

```
integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")
```

output

```
[1/2*(3*(b*x^2 + a*x)*sqrt(-b/a)*log((b*x - 2*a*sqrt(x)*sqrt(-b/a) - a)/(b
*x + a)) - 2*(3*b*x + 2*a)*sqrt(x))/(a^2*b*x^2 + a^3*x), -(3*(b*x^2 + a*x)
*sqrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + (3*b*x + 2*a)*sqrt(x))/(a^2*b*x^2 +
a^3*x)]
```

**Sympy [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{x^{3/2} \operatorname{atanh}^2(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(3/2)/atanh(tanh(b*x+a))**2,x)
```

output

```
Integral(1/(x**(3/2)*atanh(tanh(a + b*x))**2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{3bx + 2a}{a^2 b x^{3/2} + a^3 \sqrt{x}} - \frac{3b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab} a^2}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `-(3*b*x + 2*a)/(a^2*b*x^(3/2) + a^3*sqrt(x)) - 3*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{3b \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab} a^2} - \frac{3bx + 2a}{(bx^{3/2} + a\sqrt{x})a^2}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `-3*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2) - (3*b*x + 2*a)/((b*x^(3/2) + a*sqrt(x))*a^2)`**Mupad [B] (verification not implemented)**

Time = 4.02 (sec) , antiderivative size = 705, normalized size of antiderivative = 5.88

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(1/(x^(3/2)*atanh(tanh(a + b*x))^2),x)`

output

```
(x^(1/2)*(8/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
)/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - (24*b*x)/(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)
^2))/(2*b*x^2 - x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(
2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (6*2^(1/2)*b^(1/2)*log(-(b^(
1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x)
+ 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) +
4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp
(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*((2*a
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)
*exp(2*b*x) + 1)) + 2*b*x)^4 + 24*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(
exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 +
16*a^4 - 8*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
+ log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - 32*a^3*(2*a - log((2*exp(
2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) +
1)) + 2*b*x)))/(2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))
- log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(log(2/(exp(2*a)*exp(2*b*x) + 1))
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(5/2)
```

**Reduce [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x} dx$$

input

```
int(1/x^(3/2)/atanh(tanh(b*x+a))^2,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))^2*x),x)
```

**3.205**  $\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1447
Mathematica [A] (verified)	1448
Rubi [A] (verified)	1448
Maple [A] (verified)	1451
Fricas [A] (verification not implemented)	1451
Sympy [F]	1452
Maxima [A] (verification not implemented)	1452
Giac [A] (verification not implemented)	1453
Mupad [B] (verification not implemented)	1453
Reduce [F]	1454

**Optimal result**

Integrand size = 15, antiderivative size = 145

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{5b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} - \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{1} - \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{1} - \frac{bx^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{1} - \frac{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))}{1}$$

output

```
5*b^(3/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(7/2)-5*b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3-5/3/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2-1/b/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))-1/b/x^(5/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.83

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{2(-7bx + \operatorname{arctanh}(\tanh(a+bx)))}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{5b^{3/2} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} + \frac{b^2\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input `Integrate[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^2), x]`

output `(2*(-7*b*x + ArcTanh[Tanh[a + b*x]]))/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3) + (5*b^(3/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(7/2) + (b^2*Sqrt[x])/(ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3)`

**Rubi [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.29, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$\downarrow 2599$$

$$-\frac{5 \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))}$$

$$\downarrow 2594$$

$$\frac{5 \left( \frac{b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$5 \left( \frac{b \left( \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)$$

$$\frac{2b}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$5 \left( \frac{b \left( \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$$\frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))} \quad 2b$$

↓ 2593

$$\left( \frac{b \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{1}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)$$


---


$$\frac{1}{bx^{5/2}\operatorname{arctanh}(\tanh(a+bx))}$$

input `Int[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^2), x]`

output `(-5*(2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(5/2)*ArcTanh[Tanh[a + b*x]]))`

### Defintions of rubi rules used

rule 2593 `Int[1/((u)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v)) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [A] (verified)

Time = 1.41 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.79

method	result
derivativedivides	$-\frac{2}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}} + \frac{4b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 \sqrt{x}} + \frac{2b^2 \left( \frac{\sqrt{x}}{2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{5 \operatorname{arctan}\left(\frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}\right)}{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} \right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 \sqrt{x}}$
default	$-\frac{2}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}} + \frac{4b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 \sqrt{x}} + \frac{2b^2 \left( \frac{\sqrt{x}}{2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{5 \operatorname{arctan}\left(\frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}\right)}{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} \right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 \sqrt{x}}$
risch	Expression too large to display

input

```
int(1/x^(5/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output

```
-2/3/(arctanh(tanh(b*x+a))-b*x)^2/x^(3/2)+4/(arctanh(tanh(b*x+a))-b*x)^3*b/x^(1/2)+2/(arctanh(tanh(b*x+a))-b*x)^3*b^2*(1/2*x^(1/2)/arctanh(tanh(b*x+a))+5/2/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.23

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{15(b^2 x^3 + abx^2) \sqrt{-\frac{b}{a}} \log\left(\frac{bx + 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a}\right) + 2(15b^2 x^2 + 10abx - 5a^2)}{6(a^3 bx^3 + a^4 x^2)}$$



input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output `[1/6*(15*(b^2*x^3 + a*b*x^2)*sqrt(-b/a)*log((b*x + 2*a*sqrt(x)*sqrt(-b/a) - a)/(b*x + a)) + 2*(15*b^2*x^2 + 10*a*b*x - 2*a^2)*sqrt(x))/(a^3*b*x^3 + a^4*x^2), 1/3*(15*(b^2*x^3 + a*b*x^2)*sqrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + (15*b^2*x^2 + 10*a*b*x - 2*a^2)*sqrt(x))/(a^3*b*x^3 + a^4*x^2)]`

### Sympy [F]

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{x^{5/2} \operatorname{atanh}^2(\tanh(a + bx))} dx$$

input `integrate(1/x**(5/2)/atanh(tanh(b*x+a))**2,x)`

output `Integral(1/(x**(5/2)*atanh(tanh(a + b*x))**2), x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{15b^2x^2 + 10abx - 2a^2}{3(a^3bx^{5/2} + a^4x^{3/2})} + \frac{5b^2 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba^3}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`

output `1/3*(15*b^2*x^2 + 10*a*b*x - 2*a^2)/(a^3*b*x^(5/2) + a^4*x^(3/2)) + 5*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.40

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \frac{5b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}a^3} + \frac{b^2\sqrt{x}}{(bx+a)a^3} + \frac{2(6bx-a)}{3a^3x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`

output `5*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3) + b^2*sqrt(x)/((b*x + a)*a^3) + 2/3*(6*b*x - a)/(a^3*x^(3/2))`

**Mupad [B] (verification not implemented)**

Time = 4.29 (sec) , antiderivative size = 871, normalized size of antiderivative = 6.01

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(1/(x^(5/2)*atanh(tanh(a + b*x))^2),x)`

output

```

((32*b)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 - (80*b^2*x)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(x^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) - 8/(3*x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (20*2^(1/2)*b^(3/2)*log(-(b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^6 + 60*a^2*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 - 160*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 + 240*a^4*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 64*a^6 - 12*a*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5 - 192*a^5*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2...

```

**Reduce [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x^2} dx$$

input

```
int(1/x^(5/2)/atanh(tanh(b*x+a))^2,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))**2*x**2),x)
```

**3.206**  $\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$

Optimal result	1455
Mathematica [A] (verified)	1456
Rubi [A] (verified)	1456
Maple [A] (verified)	1460
Fricas [A] (verification not implemented)	1461
Sympy [F]	1461
Maxima [A] (verification not implemented)	1462
Giac [A] (verification not implemented)	1462
Mupad [B] (verification not implemented)	1463
Reduce [F]	1463

**Optimal result**

Integrand size = 15, antiderivative size = 172

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \frac{7b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{9/2}} - \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4}{7} - \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{7b} - \frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{1} - \frac{1}{bx^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

output

```
7*b^(5/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(9/2)-7*b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^4-7/3*b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^3-7/5/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^2-1/b/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))-1/b/x^(7/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.81

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = -\frac{7b^{5/2} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{9/2}} - \frac{b^3\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))(-bx + \operatorname{arctanh}(\tanh(a+bx)))^4} - \frac{2(58b^2x^2 - 16bx \operatorname{arctanh}(\tanh(a+bx)) + 3 \operatorname{arctanh}(\tanh(a+bx))^2)}{15x^{5/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^4}$$

input `Integrate[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^2), x]`

output `(-7*b^(5/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(9/2) - (b^3*Sqrt[x])/(ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4) - (2*(58*b^2*x^2 - 16*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2))/(15*x^(5/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)`

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.33, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2594, 2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx$$

$$\downarrow 2599$$

$$-\frac{7 \int \frac{1}{x^{9/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

$$\downarrow 2594$$

$$\frac{7 \left( \frac{b \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$\frac{7 \left( \frac{b \left( \frac{b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$\frac{7 \left( \frac{b \left( \frac{b \left( \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

↓ 2594

$$\left( \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{7}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$

---


$$\frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))}$$

$2b$

↓ 2593

$$\begin{aligned}
 & \left( \frac{b}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) \\
 & \left. \frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{5x^{5/2}(bx)} \right) \\
 & \frac{7}{bx - \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{1}{bx^{7/2}\operatorname{arctanh}(\tanh(a+bx))} \qquad 2b
 \end{aligned}$$

```
input Int [1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^2), x]
```

```
output (-7*(2/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]))/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(7/2)*ArcTanh[Tanh[a + b*x]])
```



Defintions of rubi rules used

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2594 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

Maple [A] (verified)

Time = 1.55 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.80

method	result
derivativedivides	$-\frac{2b^3 \left( \frac{\sqrt{x}}{2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{7 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}} \right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^4} - \frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{5}{2}}} - \dots$
default	$-\frac{2b^3 \left( \frac{\sqrt{x}}{2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{7 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}}\right)}{2\sqrt{(\operatorname{arctanh}(\tanh(bx+a))-bx)b}} \right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^4} - \frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{5}{2}}} - \dots$
risch	Expression too large to display

```
input int(1/x^(7/2)/arctanh(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)
```

output 
$$\begin{aligned} & -2/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^4*b^3*(1/2*x^{(1/2)}/\operatorname{arctanh}(\tanh(b*x+a))+7/2/ \\ & ((\operatorname{arctanh}(\tanh(b*x+a))-b*x)*b)^{(1/2)}*\operatorname{arctan}(b*x^{(1/2)}/((\operatorname{arctanh}(\tanh(b*x+a) \\ & ))-b*x)*b)^{(1/2)}))-2/5/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^2/x^{(5/2)}-6/(\operatorname{arctanh}(\tan \\ & h(b*x+a))-b*x)^4*b^2/x^{(1/2)}+4/3/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^3*b/x^{(3/2)} \end{aligned}$$

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \left[ \frac{105(b^3x^4 + ab^2x^3) \sqrt{-\frac{b}{a}} \log\left(\frac{bx - 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx+a}\right) - 2(105b^3x^3 + 70ab}{30(a^4bx^4 + a^5x^3)} \right. \\ \left. - \frac{105(b^3x^4 + ab^2x^3) \sqrt{\frac{b}{a}} \operatorname{arctan}\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + (105b^3x^3 + 70ab^2x^2 - 14a^2bx + 6a^3)\sqrt{x}}{15(a^4bx^4 + a^5x^3)} \right]$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="fricas")`

output 
$$\begin{aligned} & [1/30*(105*(b^3*x^4 + a*b^2*x^3)*\operatorname{sqrt}(-b/a)*\log((b*x - 2*a*\operatorname{sqrt}(x))*\operatorname{sqrt}(-b \\ & /a) - a)/(b*x + a) - 2*(105*b^3*x^3 + 70*a*b^2*x^2 - 14*a^2*b*x + 6*a^3)* \\ & \operatorname{sqrt}(x))/(a^4*b*x^4 + a^5*x^3), -1/15*(105*(b^3*x^4 + a*b^2*x^3)*\operatorname{sqrt}(b/a) \\ & *\operatorname{arctan}(\operatorname{sqrt}(x)*\operatorname{sqrt}(b/a)) + (105*b^3*x^3 + 70*a*b^2*x^2 - 14*a^2*b*x + 6* \\ & a^3)*\operatorname{sqrt}(x))/(a^4*b*x^4 + a^5*x^3)] \end{aligned}$$

### Sympy [F]

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx = \int \frac{1}{x^{7/2} \operatorname{atanh}^2(\tanh(a+bx))} dx$$

input `integrate(1/x**(7/2)/atanh(tanh(b*x+a))**2,x)`

output `Integral(1/(x**(7/2)*atanh(tanh(a + b*x))**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{105 b^3 x^3 + 70 a b^2 x^2 - 14 a^2 b x + 6 a^3}{15 (a^4 b x^{7/2} + a^5 x^{5/2})} - \frac{7 b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab} a^4}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="maxima")`output `-1/15*(105*b^3*x^3 + 70*a*b^2*x^2 - 14*a^2*b*x + 6*a^3)/(a^4*b*x^(7/2) + a^5*x^(5/2)) - 7*b^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = -\frac{7 b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab} a^4} - \frac{b^3 \sqrt{x}}{(bx + a)a^4} - \frac{2(45 b^2 x^2 - 10 abx + 3 a^2)}{15 a^4 x^{5/2}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^2,x, algorithm="giac")`output `-7*b^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^4) - b^3*sqrt(x)/((b*x + a)*a^4) - 2/15*(45*b^2*x^2 - 10*a*b*x + 3*a^2)/(a^4*x^(5/2))`

**Mupad [B] (verification not implemented)**

Time = 4.29 (sec) , antiderivative size = 1051, normalized size of antiderivative = 6.11

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \text{Too large to display}$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))^2),x)`

output

```
((96*b^2)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 - (224*b^3*x)/(log(2/(exp(2*a)*exp(2
*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + 2*b
*x)^4)/(x^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) -
log(2/(exp(2*a)*exp(2*b*x) + 1)))) - (32*b)/(3*x^(3/2)*(log(2/(exp(2*a)*ex
p(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) +
2*b*x)^3) - 8/(5*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*
a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (56*2^(1/2)*b^(5/2
)*log(-(b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*
b*x))/exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*
exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1))
+ 2*b*x) + 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*ex
p(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b
*x)*((2*a - log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + log(2
/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^8 + 112*a^2*(2*a - log((2*exp(2*a)*ex
p(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) +
2*b*x)^6 - 448*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x)
+ 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5 + 1120*a^4*(2*a - log
((2*exp(2*a)*exp(2*b*x))/exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(
2*b*x) + 1)) + 2*b*x)^4 - 1792*a^5*(2*a - log((2*exp(2*a)*exp(2*b*x))/(...
```

**Reduce [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^2 x^3} dx$$

input `int(1/x^(7/2)/atanh(tanh(b*x+a))^2,x)`

output `int(1/(sqrt(x)*atanh(tanh(a + b*x))**2*x**3),x)`

### 3.207 $\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1465
Mathematica [A] (verified)	1466
Rubi [A] (verified)	1466
Maple [B] (verified)	1469
Fricas [A] (verification not implemented)	1470
Sympy [F(-1)]	1470
Maxima [A] (verification not implemented)	1471
Giac [A] (verification not implemented)	1471
Mupad [B] (verification not implemented)	1472
Reduce [F]	1472

#### Optimal result

Integrand size = 15, antiderivative size = 135

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{35x^{3/2}}{12b^3} + \frac{35\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{4b^4}$$

$$- \frac{35\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}}{4b^{9/2}}$$

$$- \frac{x^{7/2}}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{7x^{5/2}}{4b^2\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
35/12*x^(3/2)/b^3+35/4*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))/b^4-35/4*arctanh
(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a
)))^(3/2)/b^(9/2)-1/2*x^(7/2)/b/arctanh(tanh(b*x+a))^2-7/4*x^(5/2)/b^2/arc
tanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.09

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx =$$

$$6b^{7/2}x^{7/2} + 21b^{5/2}x^{5/2}\operatorname{arctanh}(\tanh(a + bx)) - 140b^{3/2}x^{3/2}\operatorname{arctanh}(\tanh(a + bx))^2 + 105\sqrt{b}\sqrt{x}\operatorname{arctanh}(\tanh(a + bx)) - 12b^{9/2}\operatorname{arctanh}(\tanh(a + bx))^3$$

input `Integrate[x^(7/2)/ArcTanh[Tanh[a + b*x]]^3,x]`

output `-1/12*(6*b^(7/2)*x^(7/2) + 21*b^(5/2)*x^(5/2)*ArcTanh[Tanh[a + b*x]] - 140*b^(3/2)*x^(3/2)*ArcTanh[Tanh[a + b*x]]^2 + 105*Sqrt[b]*Sqrt[x]*ArcTanh[Tanh[a + b*x]]^3 - 105*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2))/(b^(9/2)*ArcTanh[Tanh[a + b*x]]^2)`

**Rubi [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2590, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow 2599$$

$$\frac{7 \int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^2} dx}{4b} - \frac{x^{7/2}}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\downarrow 2599$$

$$\begin{aligned}
 & \frac{7 \left( \frac{5 \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{7/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2590} \\
 & \frac{7 \left( \frac{5 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2x^{3/2}}{3b} \right)}{2b} - \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{7/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2590} \\
 & \frac{7 \left( \frac{5 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{b \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2\sqrt{x}}{b} \right) + \frac{2x^{3/2}}{3b} \right)}{2b} - \frac{x^{5/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{7/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2593}
 \end{aligned}$$



$$\frac{\left( \frac{\frac{2\sqrt{x}}{b} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}}}{b} \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{b} + \frac{2x^{3/2}}{3b} \right)}{2b} - \frac{x^{5/2}}{b\operatorname{arctanh}(\tanh(a+bx))} \right)}{x^{7/2}} \frac{4b}{2b\operatorname{arctanh}(\tanh(a+bx))^2}$$

```
input Int [x^(7/2)/ArcTanh[Tanh[a + b*x]]^3,x]
```

```
output (7*((5*((2*x^(3/2))/(3*b) + ((2*Sqrt[x])/b - (2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2))*(b*x - ArcTanh[Tanh[a + b*x]])/b)/(2*b) - x^(5/2)/(b*ArcTanh[Tanh[a + b*x]]))/(4*b) - x^(7/2)/(2*b*ArcTanh[Tanh[a + b*x]]^2)
```

**Defintions of rubi rules used**

```
rule 2590 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]
```

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(107) = 214.

Time = 1.35 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.89

method	result
derivativedivides	$-\frac{2\left(-\frac{bx^{\frac{3}{2}}}{3} + 3a\sqrt{x} + 3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x}\right)}{b^4} + \frac{2\left(\left(-\frac{13a^2b}{8} - \frac{13ab(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{4} - \frac{13b(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{8}\right)\sqrt{x}\right)}{b^4}$
default	$-\frac{2\left(-\frac{bx^{\frac{3}{2}}}{3} + 3a\sqrt{x} + 3(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)\sqrt{x}\right)}{b^4} + \frac{2\left(\left(-\frac{13a^2b}{8} - \frac{13ab(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{4} - \frac{13b(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)^2}{8}\right)\sqrt{x}\right)}{b^4}$
risch	Expression too large to display

input

```
int(x^(7/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
-2/b^4*(-1/3*b*x^(3/2)+3*a*x^(1/2)+3*(arctanh(tanh(b*x+a))-b*x-a)*x^(1/2))
+2/b^4*((( -13/8*a^2*b-13/4*a*b*(arctanh(tanh(b*x+a))-b*x-a)-13/8*b*(arctanh(tanh(b*x+a))-b*x-a)^2)*x^(3/2)+(-11/8*a^3-33/8*a^2*(arctanh(tanh(b*x+a))-b*x-a)-33/8*a*(arctanh(tanh(b*x+a))-b*x-a)^2-11/8*(arctanh(tanh(b*x+a))-b*x-a)^3)*x^(1/2))/arctanh(tanh(b*x+a))^2+35/8*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.68

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \left[ \frac{105(ab^2x^2 + 2a^2bx + a^3)\sqrt{-\frac{a}{b}} \log\left(\frac{bx + 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2(8b^3x^3 - 56ab^2x^2 - 175a^2bx - 105a^3)\sqrt{x}}{24(b^6x^2 + 2ab^5x + a^2b^4)} \right]$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `[1/24*(105*(a*b^2*x^2 + 2*a^2*b*x + a^3)*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(8*b^3*x^3 - 56*a*b^2*x^2 - 175*a^2*b*x - 105*a^3)*sqrt(x))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4), 1/12*(105*(a*b^2*x^2 + 2*a^2*b*x + a^3)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) + (8*b^3*x^3 - 56*a*b^2*x^2 - 175*a^2*b*x - 105*a^3)*sqrt(x))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4)]`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Timed out}$$

input `integrate(x**(7/2)/atanh(tanh(b*x+a))**3,x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.64

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{8b^3x^{7/2} - 56ab^2x^{5/2} - 175a^2bx^{3/2} - 105a^3\sqrt{x}}{12(b^6x^2 + 2ab^5x + a^2b^4)} + \frac{35a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^4}}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output  $\frac{1}{12} \cdot (8 \cdot b^3 \cdot x^{7/2} - 56 \cdot a \cdot b^2 \cdot x^{5/2} - 175 \cdot a^2 \cdot b \cdot x^{3/2} - 105 \cdot a^3 \cdot \sqrt{x}) / (b^6 \cdot x^2 + 2 \cdot a \cdot b^5 \cdot x + a^2 \cdot b^4) + 35/4 \cdot a^2 \cdot \arctan(b \cdot \sqrt{x} / \sqrt{a \cdot b}) / (\sqrt{a \cdot b} \cdot b^4)$ **Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.57

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{35a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^4}} - \frac{13a^2bx^{3/2} + 11a^3\sqrt{x}}{4(bx+a)^2b^4} + \frac{2(b^6x^{3/2} - 9ab^5\sqrt{x})}{3b^9}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`output  $\frac{35}{4} \cdot a^2 \cdot \arctan(b \cdot \sqrt{x} / \sqrt{a \cdot b}) / (\sqrt{a \cdot b} \cdot b^4) - 1/4 \cdot (13 \cdot a^2 \cdot b \cdot x^{3/2} + 11 \cdot a^3 \cdot \sqrt{x}) / ((b \cdot x + a)^2 \cdot b^4) + 2/3 \cdot (b^6 \cdot x^{3/2} - 9 \cdot a \cdot b^5 \cdot \sqrt{x}) / b^9$

**Mupad [B] (verification not implemented)**

Time = 4.19 (sec) , antiderivative size = 571, normalized size of antiderivative = 4.23

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(x^(7/2)/atanh(tanh(a + b*x))^3,x)`

output

$$\begin{aligned} & (2x^{3/2})/(3b^3) + (3x^{1/2}) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx))/b^4 + (35 \cdot 2^{1/2} * \log((256b^{19/2}) * (2^{1/2}) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx) - 4b^{1/2}) * x^{1/2}) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx)^{1/2} + 2 \cdot 2^{1/2} * bx)) / ((\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) - \log(2/(\exp(2a)\exp(2bx) + 1))) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx)^{1/2})) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx)^{3/2}) / (32b^{9/2}) - (13x^{1/2}) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx)^2) / (8b^4 * (\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) - \log(2/(\exp(2a)\exp(2bx) + 1)))) - (x^{1/2}) * (\log(2/(\exp(2a)\exp(2bx) + 1)) - \log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1) + 2bx)^3) / (4b^4 * (\log((2\exp(2a)\exp(2bx))/(\exp(2a)\exp(2bx) + 1)) - \log(2/(\exp(2a)\exp(2bx) + 1)))^2 \end{aligned}$$
**Reduce [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{\sqrt{x} x^3}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^(7/2)/atanh(tanh(b*x+a))^3,x)`output `int((sqrt(x)*x**3)/atanh(tanh(a + b*x))**3,x)`

### 3.208 $\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1473
Mathematica [A] (verified)	1474
Rubi [A] (verified)	1474
Maple [A] (verified)	1476
Fricas [A] (verification not implemented)	1477
Sympy [F]	1477
Maxima [A] (verification not implemented)	1478
Giac [A] (verification not implemented)	1478
Mupad [B] (verification not implemented)	1479
Reduce [F]	1479

#### Optimal result

Integrand size = 15, antiderivative size = 110

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{15\sqrt{x}}{4b^3} - \frac{15\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}{4b^{7/2}} - \frac{x^{5/2}}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{5x^{3/2}}{4b^2\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
15/4*x^(1/2)/b^3-15/4*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^(1/2)/b^(7/2)-1/2*x^(5/2)/b/arctanh(tanh(b*x+a))^2-5/4*x^(3/2)/b^2/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.95

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{1}{4} \left( \frac{15\sqrt{x}}{b^3} - \frac{2x^{5/2}}{b \operatorname{arctanh}(\tanh(a + bx))^2} - \frac{5x^{3/2}}{b^2 \operatorname{arctanh}(\tanh(a + bx))} - \frac{15 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right) \sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}{b^{7/2}} \right)$$

input `Integrate[x^(5/2)/ArcTanh[Tanh[a + b*x]]^3,x]`

output `((15*Sqrt[x])/b^3 - (2*x^(5/2))/(b*ArcTanh[Tanh[a + b*x]]^2) - (5*x^(3/2))/(b^2*ArcTanh[Tanh[a + b*x]]) - (15*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]*Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]])/b^(7/2))/4`

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2590, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

↓ 2599

$$\begin{aligned}
 & \frac{5 \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^2} dx}{4b} - \frac{x^{5/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2599} \\
 & \frac{5 \left( \frac{3 \int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{5/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2590} \\
 & \frac{5 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{b} + \frac{2\sqrt{x}}{b} \right)}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{5/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2593} \\
 & \frac{5 \left( \frac{3 \left( \frac{2\sqrt{x}}{b} - \frac{2 \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}} \right) \sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}{b^{3/2}} \right)}{2b} - \frac{x^{3/2}}{b \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{5/2}}{2b \operatorname{arctanh}(\tanh(a+bx))^2}
 \end{aligned}$$

input

`Int[x^(5/2)/ArcTanh[Tanh[a + b*x]]^3,x]`

output

`(5*((3*((2*Sqrt[x])/b - (2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])*Sqrt[b*x - ArcTanh[Tanh[a + b*x]]])/b^(3/2)))/(2*b) - x^(3/2)/(b*ArcTanh[Tanh[a + b*x]]))/(4*b) - x^(5/2)/(2*b*ArcTanh[Tanh[a + b*x]]^2)`



Defintions of rubi rules used

```
rule 2590 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^n/(a*n), x] - Simp[(b*u - a*v)/a Int[v^(n - 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && GtQ[n, 0] && NeQ[n, 1]
```

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.39

method	result
derivativedivides	$\frac{2\sqrt{x}}{b^3} - \frac{2 \left( \left( -\frac{9ab}{8} - \frac{9b(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{8} \right) x^{\frac{3}{2}} + \left( -\frac{7a^2}{8} - \frac{7a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{4} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))}{8} \right) \operatorname{arctanh}(\tanh(bx+a))^2 \right)}{b^3}$
default	$\frac{2\sqrt{x}}{b^3} - \frac{2 \left( \left( -\frac{9ab}{8} - \frac{9b(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{8} \right) x^{\frac{3}{2}} + \left( -\frac{7a^2}{8} - \frac{7a(\operatorname{arctanh}(\tanh(bx+a)) - bx - a)}{4} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))}{8} \right) \operatorname{arctanh}(\tanh(bx+a))^2 \right)}{b^3}$
risch	Expression too large to display

```
input int(x^(5/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
2*x^(1/2)/b^3-2/b^3*(((9/8*a*b-9/8*b*(arctanh(tanh(b*x+a))-b*x-a))*x^(3/2)
)+(-7/8*a^2-7/4*a*(arctanh(tanh(b*x+a))-b*x-a)-7/8*(arctanh(tanh(b*x+a))-b
*x-a)^2)*x^(1/2))/arctanh(tanh(b*x+a))^2+15/8*(arctanh(tanh(b*x+a))-b*x)/(
(arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a)
)-b*x)*b)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.82

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \left[ \frac{15(b^2x^2 + 2abx + a^2)\sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx+a}\right) + 2(8b^2x^2 + 25abx + 15a^2)\sqrt{x}}{8(b^5x^2 + 2ab^4x + a^2b^3)} - \frac{15(b^2x^2 + 2abx + a^2)\sqrt{\frac{a}{b}} \operatorname{arctan}\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (8b^2x^2 + 25abx + 15a^2)\sqrt{x}}{4(b^5x^2 + 2ab^4x + a^2b^3)} \right]$$

input

```
integrate(x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

```
[1/8*(15*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(
-a/b) - a)/(b*x + a)) + 2*(8*b^2*x^2 + 25*a*b*x + 15*a^2)*sqrt(x))/(b^5*x^
2 + 2*a*b^4*x + a^2*b^3), -1/4*(15*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a/b)*arc
tan(b*sqrt(x)*sqrt(a/b)/a) - (8*b^2*x^2 + 25*a*b*x + 15*a^2)*sqrt(x))/(b^5
*x^2 + 2*a*b^4*x + a^2*b^3)]
```

**Sympy [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \int \frac{x^{5/2}}{\operatorname{atanh}^3(\tanh(a+bx))} dx$$

input

```
integrate(x**(5/2)/atanh(tanh(b*x+a))**3,x)
```

output `Integral(x**(5/2)/atanh(tanh(a + b*x))**3, x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.66

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{8b^2x^{\frac{5}{2}} + 25abx^{\frac{3}{2}} + 15a^2\sqrt{x}}{4(b^5x^2 + 2ab^4x + a^2b^3)} - \frac{15a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^3}}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `1/4*(8*b^2*x^(5/2) + 25*a*b*x^(3/2) + 15*a^2*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3) - 15/4*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3)`

### Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.54

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{15a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^3}} + \frac{2\sqrt{x}}{b^3} + \frac{9abx^{\frac{3}{2}} + 7a^2\sqrt{x}}{4(bx + a)^2b^3}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `-15/4*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2*sqrt(x)/b^3 + 1/4*(9*a*b*x^(3/2) + 7*a^2*sqrt(x))/((b*x + a)^2*b^3)`

**Mupad [B] (verification not implemented)**

Time = 4.09 (sec) , antiderivative size = 511, normalized size of antiderivative = 4.65

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(x^(5/2)/atanh(tanh(a + b*x))^3,x)`

output

```
(2*x^(1/2))/b^3 - (9*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x))/(4*b^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) + (15*2^(1/2)*log((64*b^(15/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^(1/2)))/(16*b^(7/2)) - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) + 2*b*x)^2)/(2*b^3*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^2)
```

**Reduce [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{\sqrt{x} x^2}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a))^3,x)`output `int((sqrt(x)*x**2)/atanh(tanh(a + b*x))**3,x)`

### 3.209 $\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1480
Mathematica [A] (verified)	1480
Rubi [A] (verified)	1481
Maple [A] (verified)	1482
Fricas [A] (verification not implemented)	1483
Sympy [F]	1483
Maxima [A] (verification not implemented)	1484
Giac [A] (verification not implemented)	1484
Mupad [B] (verification not implemented)	1484
Reduce [F]	1485

#### Optimal result

Integrand size = 15, antiderivative size = 98

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{3\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4b^{5/2}\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}} - \frac{x^{3/2}}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{3\sqrt{x}}{4b^2\operatorname{arctanh}(\tanh(a+bx))}$$

output

$$-3/4*\operatorname{arctanh}(b^{1/2}*x^{1/2}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{1/2})/b^{5/2}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^{1/2}-1/2*x^{3/2}/b/\operatorname{arctanh}(\tanh(b*x+a))^2-3/4*x^{1/2}/b^2/\operatorname{arctanh}(\tanh(b*x+a))$$

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.98

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{x^{3/2}}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{3\sqrt{x}}{4b^2\operatorname{arctanh}(\tanh(a+bx))} + \frac{3\operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4b^{5/2}\sqrt{-bx+\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Integrate[x^(3/2)/ArcTanh[Tanh[a + b*x]]^3,x]`

output 
$$-1/2*x^{(3/2)}/(b*\text{ArcTanh}[\text{Tanh}[a + b*x]]^2) - (3*\text{Sqrt}[x])/(4*b^2*\text{ArcTanh}[\text{Tanh}[a + b*x]]) + (3*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[x])/\text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]]])/(4*b^{(5/2)}*\text{Sqrt}[-(b*x) + \text{ArcTanh}[\text{Tanh}[a + b*x]]])$$

### Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {2599, 2599, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{3/2}}{\text{arctanh}(\tanh(a + bx))^3} dx$$

↓ 2599

$$\frac{3 \int \frac{\sqrt{x}}{\text{arctanh}(\tanh(a+bx))^2} dx}{4b} - \frac{x^{3/2}}{2b \text{arctanh}(\tanh(a + bx))^2}$$

↓ 2599

$$\frac{3 \left( \frac{\int \frac{1}{\sqrt{x} \text{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{\sqrt{x}}{b \text{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{3/2}}{2b \text{arctanh}(\tanh(a + bx))^2}$$

↓ 2593

$$\frac{3 \left( -\frac{\text{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \text{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}\sqrt{bx - \text{arctanh}(\tanh(a+bx))}} - \frac{\sqrt{x}}{b \text{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{x^{3/2}}{2b \text{arctanh}(\tanh(a + bx))^2}$$

input `Int[x^(3/2)/ArcTanh[Tanh[a + b*x]]^3,x]`

output

$$\frac{(3*(-\text{ArcTanh}[\text{Sqrt}[b]*\text{Sqrt}[x]]/\text{Sqrt}[b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]])/(b^{(3/2)*\text{Sqrt}[b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]]) - \text{Sqrt}[x]/(b*\text{ArcTanh}[\text{Tanh}[a + b*x]])))/(4*b) - x^{(3/2)/(2*b*\text{ArcTanh}[\text{Tanh}[a + b*x]]^2)}$$
**Defintions of rubi rules used**

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 1.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$\frac{-\frac{5x^{\frac{3}{2}}}{4b} - \frac{3(\text{arctanh}(\tanh(bx+a)) - bx)\sqrt{x}}{4b^2}}{\text{arctanh}(\tanh(bx+a))^2} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\text{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{4b^2 \sqrt{(\text{arctanh}(\tanh(bx+a)) - bx)b}}$	85
default	$\frac{-\frac{5x^{\frac{3}{2}}}{4b} - \frac{3(\text{arctanh}(\tanh(bx+a)) - bx)\sqrt{x}}{4b^2}}{\text{arctanh}(\tanh(bx+a))^2} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{(\text{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{4b^2 \sqrt{(\text{arctanh}(\tanh(bx+a)) - bx)b}}$	85
risch	Expression too large to display	1075

input

```
int(x^(3/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
2*(-5/8*x^(3/2)/b-3/8*(arctanh(tanh(b*x+a))-b*x)/b^2*x^(1/2))/arctanh(tanh
(b*x+a))^2+3/4/b^2/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/
(arctanh(tanh(b*x+a))-b*x)*b)^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.89

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \left[ -\frac{3(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right) + 2(5ab^2x + 3a^2b)\sqrt{x}}{8(ab^5x^2 + 2a^2b^4x + a^3b^3)} \right. \\ \left. - \frac{3(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) + (5ab^2x + 3a^2b)\sqrt{x}}{4(ab^5x^2 + 2a^2b^4x + a^3b^3)} \right]$$

input

```
integrate(x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

```
[-1/8*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*
sqrt(x))/(b*x + a)) + 2*(5*a*b^2*x + 3*a^2*b)*sqrt(x))/(a*b^5*x^2 + 2*a^2*
b^4*x + a^3*b^3), -1/4*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(
a*b)/(b*sqrt(x))) + (5*a*b^2*x + 3*a^2*b)*sqrt(x))/(a*b^5*x^2 + 2*a^2*b^4*
x + a^3*b^3)]
```

**Sympy [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \int \frac{x^{\frac{3}{2}}}{\operatorname{atanh}^3(\tanh(a+bx))} dx$$

input

```
integrate(x**(3/2)/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(x**(3/2)/atanh(tanh(a + b*x))**3, x)
```



**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.62

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{5bx^{3/2} + 3a\sqrt{x}}{4(b^4x^2 + 2ab^3x + a^2b^2)} + \frac{3 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^2}}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output `-1/4*(5*b*x^(3/2) + 3*a*sqrt(x))/(b^4*x^2 + 2*a*b^3*x + a^2*b^2) + 3/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.48

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{3 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^2}} - \frac{5bx^{3/2} + 3a\sqrt{x}}{4(bx + a)^2b^2}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`output `3/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) - 1/4*(5*b*x^(3/2) + 3*a*sqrt(x))/((b*x + a)^2*b^2)`**Mupad [B] (verification not implemented)**

Time = 4.11 (sec) , antiderivative size = 667, normalized size of antiderivative = 6.81

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(x^(3/2)/atanh(tanh(a + b*x))^3,x)`

output

```
(3*2^(1/2)*log((16*b^(11/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(8*b^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)) - (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(b^2*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^2) - (x^(1/2)*(1/(b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (8*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 8*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 16*b*x)/(2*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1))))))
```

**Reduce [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{3 \operatorname{atanh}(\tanh(bx + a))^2 \left( \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx+a))} dx \right) - 6\sqrt{x} \operatorname{atanh}(\tanh(bx + a))}{8 \operatorname{atanh}(\tanh(bx + a))^2 b^2}$$

input

```
int(x^(3/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
(3*atanh(tanh(a + b*x))**2*int(1/(sqrt(x)*atanh(tanh(a + b*x))),x) - 6*sqrt(x)*atanh(tanh(a + b*x)) - 4*sqrt(x)*b*x)/(8*atanh(tanh(a + b*x))**2*b**2)
```

**3.210**  $\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1486
Mathematica [A] (verified)	1487
Rubi [A] (verified)	1487
Maple [A] (verified)	1489
Fricas [A] (verification not implemented)	1490
Sympy [F]	1490
Maxima [A] (verification not implemented)	1491
Giac [A] (verification not implemented)	1491
Mupad [B] (verification not implemented)	1491
Reduce [F]	1492

**Optimal result**

Integrand size = 15, antiderivative size = 125

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx-\operatorname{arctanh}(\tanh(a+bx))}}\right)}{4b^{3/2}(bx-\operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{1}{4b^2\sqrt{x}(bx-\operatorname{arctanh}(\tanh(a+bx)))} - \frac{\sqrt{x}}{2b\operatorname{arctanh}(\tanh(a+bx))^2} - \frac{1}{4b^2\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))}$$

```
output 1/4*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/b^(3/2)/(b*x
-arctanh(tanh(b*x+a)))^(3/2)-1/4/b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))-1/
2*x^(1/2)/b/arctanh(tanh(b*x+a))^2-1/4/b^2/x^(1/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{1}{4} \left( -\frac{2\sqrt{x}}{b \operatorname{arctanh}(\tanh(a + bx))^2} + \frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{b^{3/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{3/2}} \right. \\ \left. + \frac{\sqrt{x}}{-b^2 x \operatorname{arctanh}(\tanh(a + bx)) + b \operatorname{arctanh}(\tanh(a + bx))^2} \right)$$

input `Integrate[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^3,x]`

output `((-2*Sqrt[x])/(b*ArcTanh[Tanh[a + b*x]]^2) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]]/(b^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(3/2)) + Sqrt[x]/(-(b^2*x*ArcTanh[Tanh[a + b*x]]) + b*ArcTanh[Tanh[a + b*x]]^2))/4`

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$ , Rules used = {2599, 2599, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow \text{2599}$$

$$\frac{\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2} dx}{4b} - \frac{\sqrt{x}}{2b \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\begin{aligned}
 & \int \frac{\frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} - \frac{\sqrt{x}}{2b \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \quad \downarrow \text{2599} \\
 & \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{b x - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \text{2594} \\
 & \frac{4b}{\sqrt{x}} \\
 & \frac{2b \operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}} \\
 & \quad \downarrow \text{2593} \\
 & \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} - \frac{1}{b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \quad \downarrow \\
 & \frac{4b}{\sqrt{x}} \\
 & \frac{2b \operatorname{arctanh}(\tanh(a+bx))^2}{\sqrt{x}}
 \end{aligned}$$

input `Int[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^3,x]`

output 
$$\frac{(-1/2*((-2*\text{Sqrt}[b]*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[x])/ \text{Sqrt}[b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]]])/(b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])^{3/2} + 2/(\text{Sqrt}[x]*(b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]])))/b - 1/(b*\text{Sqrt}[x]*\text{ArcTanh}[\text{Tanh}[a + b*x]]))/(4*b) - \text{Sqrt}[x]/(2*b*\text{ArcTanh}[\text{Tanh}[a + b*x]]^2)}$$

**Defintions of rubi rules used**

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{\frac{2x^{\frac{3}{2}}}{8 \operatorname{arctanh}(\tanh(bx+a)) - 8bx} - \frac{\sqrt{x}}{4b}}{\operatorname{arctanh}(\tanh(bx+a))^2} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{4(\operatorname{arctanh}(\tanh(bx+a)) - bx)b\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}$	98
default	$\frac{\frac{2x^{\frac{3}{2}}}{8 \operatorname{arctanh}(\tanh(bx+a)) - 8bx} - \frac{\sqrt{x}}{4b}}{\operatorname{arctanh}(\tanh(bx+a))^2} + \frac{\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{4(\operatorname{arctanh}(\tanh(bx+a)) - bx)b\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}$	98
risch	Expression too large to display	1595

input `int(x^(1/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `2*(1/8/(arctanh(tanh(b*x+a))-b*x)*x^(3/2)-1/8*x^(1/2)/b)/arctanh(tanh(b*x+a))^2+1/4/(arctanh(tanh(b*x+a))-b*x)/b/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.49

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= \left[ -\frac{(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right) - 2(ab^2x - a^2b)\sqrt{x}}{8(a^2b^4x^2 + 2a^3b^3x + a^4b^2)}, \right. \\ \left. -\frac{(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) - (ab^2x - a^2b)\sqrt{x}}{4(a^2b^4x^2 + 2a^3b^3x + a^4b^2)} \right]$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output `[-1/8*((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)) - 2*(a*b^2*x - a^2*b)*sqrt(x))/(a^2*b^4*x^2 + 2*a^3*b^3*x + a^4*b^2), -1/4*((b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(a*b)/(b*sqrt(x))) - (a*b^2*x - a^2*b)*sqrt(x))/(a^2*b^4*x^2 + 2*a^3*b^3*x + a^4*b^2)]`

**Sympy [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}^3(\tanh(a+bx))} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a))**3,x)`

output `Integral(sqrt(x)/atanh(tanh(a + b*x))**3, x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.51

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{bx^{\frac{3}{2}} - a\sqrt{x}}{4(ab^3x^2 + 2a^2b^2x + a^3b)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abab}}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`

output `1/4*(b*x^(3/2) - a*sqrt(x))/(a*b^3*x^2 + 2*a^2*b^2*x + a^3*b) + 1/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a*b)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.42

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abab}} + \frac{bx^{\frac{3}{2}} - a\sqrt{x}}{4(bx+a)^2ab}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `1/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a*b) + 1/4*(b*x^(3/2) - a*sqrt(x))/((b*x + a)^2*a*b)`

**Mupad [B] (verification not implemented)**

Time = 4.04 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.64

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^3} dx = \text{Too large to display}$$

input `int(x^(1/2)/atanh(tanh(a + b*x))^3,x)`



output

```
(2^(1/2)*log(-4*b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*(b^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + 4*a^2*b^3 - 4*a*b^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))))/(4*b^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(3/2)) - (2*x^(1/2))/(b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^2) - x^(1/2)/(b*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))
```

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{\operatorname{atanh}(\tanh(bx + a))^2 \left( \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx+a))^2} dx \right) - 2\sqrt{x}}{4 \operatorname{atanh}(\tanh(bx + a))^2 b}$$

input

```
int(x^(1/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
(atanh(tanh(a + b*x))**2*int(1/(sqrt(x)*atanh(tanh(a + b*x))**2),x) - 2*sqrt(x))/(4*atanh(tanh(a + b*x))**2*b)
```

**3.211**  $\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1493
Mathematica [A] (verified)	1494
Rubi [A] (verified)	1494
Maple [A] (verified)	1497
Fricas [A] (verification not implemented)	1497
Sympy [F]	1498
Maxima [A] (verification not implemented)	1498
Giac [A] (verification not implemented)	1499
Mupad [B] (verification not implemented)	1499
Reduce [F]	1500

**Optimal result**

Integrand size = 15, antiderivative size = 152

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{3 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4\sqrt{b}(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2}} + \frac{3}{4b\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{1}{4b^2x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{1}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} + \frac{1}{4b^2x^{3/2} \operatorname{arctanh}(\tanh(a+bx))}$$

output

```
-3/4*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/b^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(5/2)+3/4/b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2+1/4/b^2/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))-1/2/b/x^(1/2)/arctanh(tanh(b*x+a))^2+1/4/b^2/x^(3/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.78

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$= \frac{3 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{4\sqrt{b}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{5/2}} + \frac{3\sqrt{x}}{4\operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2} + \frac{\sqrt{x}}{2\operatorname{arctanh}(\tanh(a + bx))^2(-bx + \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Integrate[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^3),x]`

output `(3*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(4*Sqrt[b]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(5/2)) + (3*Sqrt[x])/(4*ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2) + Sqrt[x]/(2*ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {2599, 2599, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\downarrow \text{2599}$$

$$-\frac{\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^2} dx}{4b} - \frac{1}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^2}$$

$$\begin{aligned}
 & \downarrow 2599 \\
 & \frac{3 \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{4b \operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2594 \\
 & \frac{3 \left( \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{4b}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2594 \\
 & \frac{3 \left( \frac{b \left( \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{4b}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2593 \\
 & \frac{3 \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{2b} - \frac{1}{bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{4b}{2b\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^2}
 \end{aligned}$$

input

```
Int [1/(Sqrt [x]*ArcTanh [Tanh [a + b*x]]^3), x]
```

output

```
-1/4*((-3*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*
ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]]))/(b*x - ArcT
anh[Tanh[a + b*x]]^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])))))/(
b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(3/2)*ArcTanh[Tanh[a + b*x]
]))/b - 1/(2*b*Sqrt[x]*ArcTanh[Tanh[a + b*x]]^2)
```

### Defintions of rubi rules used

rule 2593

```
Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simpli
fy[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u -
a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; Piecewise
LinearQ[u, v, x]
```

rule 2594

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n +
1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; Piecew
iseLinearQ[u, v, x] && LtQ[n, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 1.32 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{\sqrt{x}}{2(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^2} + \frac{3\sqrt{x}}{4(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{1}{4(a$
default	$\frac{\sqrt{x}}{2(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^2} + \frac{3\sqrt{x}}{4(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \operatorname{arctanh}(\tanh(bx+a))} + \frac{1}{4(a$
risch	Expression too large to display

input `int(1/x^(1/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}x^{1/2}/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)/\operatorname{arctanh}(\tanh(b*x+a))^2+3/4/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^2*x^{1/2}/\operatorname{arctanh}(\tanh(b*x+a))+3/4/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^2/((\operatorname{arctanh}(\tanh(b*x+a))-b*x)*b)^{(1/2)}*\arctan(b*x^{1/2}/((\operatorname{arctanh}(\tanh(b*x+a))-b*x)*b)^{(1/2)})$

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.22

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

$$= \left[ \frac{3(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right) - 2(3ab^2x + 5a^2b)\sqrt{x}}{8(a^3b^3x^2 + 2a^4b^2x + a^5b)}, \right.$$

$$\left. \frac{3(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) - (3ab^2x + 5a^2b)\sqrt{x}}{4(a^3b^3x^2 + 2a^4b^2x + a^5b)} \right]$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```
[-1/8*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)) - 2*(3*a*b^2*x + 5*a^2*b)*sqrt(x))/(a^3*b^3*x^2 + 2*a^4*b^2*x + a^5*b), -1/4*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(a*b)/(b*sqrt(x))) - (3*a*b^2*x + 5*a^2*b)*sqrt(x))/(a^3*b^3*x^2 + 2*a^4*b^2*x + a^5*b)]
```

**Sympy [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(1/2)/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(1/(sqrt(x)*atanh(tanh(a + b*x))**3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.39

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{3bx^{\frac{3}{2}} + 5a\sqrt{x}}{4(a^2b^2x^2 + 2a^3bx + a^4)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{aba^2}}$$

input

```
integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")
```

output

```
1/4*(3*b*x^(3/2) + 5*a*sqrt(x))/(a^2*b^2*x^2 + 2*a^3*b*x + a^4) + 3/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.31

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4 \sqrt{ab} a^2} + \frac{3bx^{\frac{3}{2}} + 5a\sqrt{x}}{4(bx + a)^2 a^2}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `3/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2) + 1/4*(3*b*x^(3/2) + 5*a*sqrt(x))/((b*x + a)^2*a^2)`

**Mupad [B] (verification not implemented)**

Time = 4.09 (sec) , antiderivative size = 741, normalized size of antiderivative = 4.88

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))^3),x)`



output

```
(6*x^(1/2))/((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log
(2/(exp(2*a)*exp(2*b*x) + 1)))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) - (4*x^(1/2))/
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)
*exp(2*b*x) + 1)))^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*e
xp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (3*2^(1/2)*log((b^(1/2)*
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1))
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(
1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x)
)))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x)*(16*a^4*b +
b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(e
xp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 - 8*a*b*(2*a - log((2*exp(2*a)*exp(2*b
*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)
)^3 - 32*a^3*b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1
)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) + 24*a^2*b*(2*a - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x)^2))/(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)
) - log(2/(exp(2*a)*exp(2*b*x) + 1))))/(2*b^(1/2)*(log(2/(exp(2*a)*exp(2*
b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2...
```

**Reduce [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3} dx$$

input

```
int(1/x^(1/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))^3),x)
```

**3.212**  $\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1501
Mathematica [A] (verified)	1502
Rubi [A] (verified)	1502
Maple [A] (verified)	1506
Fricas [A] (verification not implemented)	1507
Sympy [F]	1507
Maxima [A] (verification not implemented)	1508
Giac [A] (verification not implemented)	1508
Mupad [B] (verification not implemented)	1508
Reduce [F]	1509

**Optimal result**

Integrand size = 15, antiderivative size = 176

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{15\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} + \frac{15}{4\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{15}{4bx^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{4b^2x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2bx^{3/2}\operatorname{arctanh}(\tanh(a+bx))^2} + \frac{3}{4b^2x^{5/2}\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
-15/4*b^(1/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(7/2)+15/4/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3+5/4/b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2+3/4/b^2/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))-1/2/b/x^(3/2)/arctanh(tanh(b*x+a))^2+3/4/b^2/x^(5/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.80

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{15\sqrt{b} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(-bx + \operatorname{arctanh}(\tanh(a+bx)))^{7/2}} - \frac{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3}{7b\sqrt{x}} - \frac{4\operatorname{arctanh}(\tanh(a+bx))(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3}{b\sqrt{x}} - \frac{2\operatorname{arctanh}(\tanh(a+bx))^2(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}{b\sqrt{x}}$$

input `Integrate[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^3), x]`

output `(-15*Sqrt[b]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(4*(-(b*x) + ArcTanh[Tanh[a + b*x]])^(7/2)) - 2/(Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3) - (7*b*Sqrt[x])/(4*ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3) - (b*Sqrt[x])/(2*ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.23, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {2599, 2599, 2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

↓ 2599

$$-\frac{3 \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx}{4b} - \frac{1}{2bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

$$\frac{3 \left( -\frac{5 \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{1}{2bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

2599

$$\frac{3 \left( -\frac{5 \left( \frac{b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b}$$

2594

$$\frac{1}{2bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

2594

$$\frac{3 \left( -\frac{5 \left( \frac{b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{4b} - \frac{1}{bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))}$$

$$\frac{1}{2bx^{3/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

2594

$$\left( \frac{b \int \frac{\arctanh(\tanh(a+bx)) dx}{\sqrt{x} (bx - \arctanh(\tanh(a+bx)))} + \frac{2}{\sqrt{x} (bx - \arctanh(\tanh(a+bx)))}}{bx - \arctanh(\tanh(a+bx))} + \frac{2}{3x^{3/2} (bx - \arctanh(\tanh(a+bx)))} \right) + \frac{2}{5x^{5/2} (bx - \arctanh(\tanh(a+bx)))}$$


---

$\frac{1}{2bx^{3/2} \arctanh(\tanh(a+bx))^2}$   
 $\downarrow$  2593

$$\frac{\frac{b \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{1}{5x^{5/2}}}{3} \cdot \frac{1}{2b} = \frac{1}{2bx^{3/2}\operatorname{arctanh}(\tanh(a+bx))^2} + \frac{4b}{4b}$$

input `Int [1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^3), x]`

output `(-3*((-5*(2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]])^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])))))/(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(5/2)*ArcTanh[Tanh[a + b*x]]))/(4*b) - 1/(2*b*x^(3/2)*ArcTanh[Tanh[a + b*x]]^2)`

Defintions of rubi rules used

```
rule 2593 Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2594 Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

Maple [A] (verified)

Time = 1.38 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.65

method	result
derivativedivides	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}} - \frac{2b\left(\frac{7bx^{\frac{3}{2}}}{8} + \left(\frac{9\operatorname{arctanh}(\tanh(bx+a)) - 9bx}{8}\right)\sqrt{x} + \frac{15\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{8\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3}$
default	$-\frac{2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3\sqrt{x}} - \frac{2b\left(\frac{7bx^{\frac{3}{2}}}{8} + \left(\frac{9\operatorname{arctanh}(\tanh(bx+a)) - 9bx}{8}\right)\sqrt{x} + \frac{15\operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{8\sqrt{\operatorname{arctanh}(\tanh(bx+a))-bx}}\right)}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^3}$
risch	Expression too large to display

```
input int(1/x^(3/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)
```

output

```
-2/(arctanh(tanh(b*x+a))-b*x)^3/x^(1/2)-2/(arctanh(tanh(b*x+a))-b*x)^3*b*(
(7/8*b*x^(3/2)+(9/8*arctanh(tanh(b*x+a))-9/8*b*x)*x^(1/2))/arctanh(tanh(b*
x+a))^2+15/8/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arcta
nh(tanh(b*x+a))-b*x)*b)^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \left[ \frac{15(b^2x^3 + 2abx^2 + a^2x) \sqrt{-\frac{b}{a}} \log\left(\frac{bx - 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a}\right) - 2(15b^2x^2 + 15abx + 8a^2)\sqrt{x}}{8(a^3b^2x^3 + 2a^4bx^2 + a^5x)} \right. \\ \left. - \frac{15(b^2x^3 + 2abx^2 + a^2x) \sqrt{\frac{b}{a}} \operatorname{arctan}\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + (15b^2x^2 + 25abx + 8a^2)\sqrt{x}}{4(a^3b^2x^3 + 2a^4bx^2 + a^5x)} \right]$$

input

```
integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")
```

output

```
[1/8*(15*(b^2*x^3 + 2*a*b*x^2 + a^2*x)*sqrt(-b/a)*log((b*x - 2*a*sqrt(x)*s
qrt(-b/a) - a)/(b*x + a)) - 2*(15*b^2*x^2 + 25*a*b*x + 8*a^2)*sqrt(x))/(a^
3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x), -1/4*(15*(b^2*x^3 + 2*a*b*x^2 + a^2*x)*s
qrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + (15*b^2*x^2 + 25*a*b*x + 8*a^2)*sqrt(x
))/(a^3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x)]
```

**Sympy [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x^{3/2} \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(3/2)/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(1/(x**(3/2)*atanh(tanh(a + b*x))**3), x)
```



**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{15b^2x^2 + 25abx + 8a^2}{4(a^3b^2x^{5/2} + 2a^4bx^{3/2} + a^5\sqrt{x})} - \frac{15b \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{ab}a^3}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")`output `-1/4*(15*b^2*x^2 + 25*a*b*x + 8*a^2)/(a^3*b^2*x^(5/2) + 2*a^4*b*x^(3/2) + a^5*sqrt(x)) - 15/4*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.34

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{15b \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{ab}a^3} - \frac{2}{a^3\sqrt{x}} - \frac{7b^2x^{3/2} + 9ab\sqrt{x}}{4(bx + a)^2a^3}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`output `-15/4*b*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^3) - 2/(a^3*sqrt(x)) - 1/4*(7*b^2*x^(3/2) + 9*a*b*sqrt(x))/((b*x + a)^2*a^3)`**Mupad [B] (verification not implemented)**

Time = 4.14 (sec) , antiderivative size = 1077, normalized size of antiderivative = 6.12

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(1/(x^(3/2)*atanh(tanh(a + b*x))^3),x)`

output

```
(x*((12*b)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))
/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 + (3*b*(16*log(2/(exp(2*a)*exp(2*b*
x) + 1)) - 16*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 32*
b*x))/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp
(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4 - (16*log(2/(exp(2*a)*exp(2*b*x) + 1))
- 16*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 32*b*x)/(log
(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(
2*b*x) + 1)) + 2*b*x)^3)/(x^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*e
xp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))) + (15*2^(1/2)*b^(1/2)
*log((b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*ex
p(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) +
2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(
2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2) + 2*2^(1/2)*b*x
)*((2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(
exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^6 + 60*a^2*(2*a - log((2*exp(2*a)*exp(2
*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b
*x)^4 - 160*a^3*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) +
1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 + 240*a^4*(2*a - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2...
```

**Reduce [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x} dx$$

input

```
int(1/x^(3/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))**3*x),x)
```

### 3.213 $\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1510
Mathematica [A] (verified)	1511
Rubi [A] (verified)	1511
Maple [A] (verified)	1517
Fricas [A] (verification not implemented)	1517
Sympy [F]	1518
Maxima [A] (verification not implemented)	1518
Giac [A] (verification not implemented)	1519
Mupad [B] (verification not implemented)	1519
Reduce [F]	1520

#### Optimal result

Integrand size = 15, antiderivative size = 201

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx = -\frac{35b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^{9/2}} + \frac{35b}{4\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{12x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{7} + \frac{4bx^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{5} + \frac{4b^2x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{5} - \frac{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2}{1} + \frac{4b^2x^{7/2} \operatorname{arctanh}(\tanh(a+bx))}{5}$$

output

```
-35/4*b^(3/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b*x-arctanh(tanh(b*x+a)))^(9/2)+35/4*b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^4+35/12/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^3+7/4/b/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^2+5/4/b^2/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))-1/2/b/x^(5/2)/arctanh(tanh(b*x+a))^2+5/4/b^2/x^(7/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.78

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{1}{12} \left( \frac{105b^{3/2} \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{9/2}} \right. \\ \left. + \frac{80bx - 8\operatorname{arctanh}(\tanh(a + bx))}{x^{3/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4} \right. \\ \left. + \frac{33b^2\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4} \right. \\ \left. + \frac{6b^2\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3} \right)$$

input

```
Integrate[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^3),x]
```

output

```
((105*b^(3/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(9/2) + (80*b*x - 8*ArcTanh[Tanh[a + b*x]])/(x^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4) + (33*b^2*Sqrt[x])/((ArcTanh[Tanh[a + b*x]]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4) + (6*b^2*Sqrt[x])/(ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3))/12
```

**Rubi [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.28, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {2599, 2599, 2594, 2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx$$

$$\begin{aligned}
 & \downarrow 2599 \\
 & \frac{5 \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx}{4b} - \frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2599 \\
 & \frac{5 \left( -\frac{7 \int \frac{1}{x^{9/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2594 \\
 & \frac{5 \left( -\frac{7 \left( \frac{b \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} \\
 & \frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2594 \\
 & \frac{5 \left( \frac{7 \left( \frac{b \left( \frac{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} \right)}{4b} - \frac{1}{bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))} \\
 & \frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2} \\
 & \downarrow 2594
 \end{aligned}$$

$$\left( \frac{b \int \frac{x^{3/2} \operatorname{arctanh}(\tanh(a+bx)) \, dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2594

$$\left( \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---


$$\frac{2b}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{1}{2bx^{5/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2593

$$\left( \frac{b}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{b}{5x^5}$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))}$$


---


$$\frac{b}{2b}$$



input `Int[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^3),x]`

output `(-5*((-7*(2/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]])])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])))))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(7/2)*ArcTanh[Tanh[a + b*x]]))/(4*b) - 1/(2*b*x^(5/2)*ArcTanh[Tanh[a + b*x]]^2)`

### Defintions of rubi rules used

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*(n + 1)/((n + 1)*(b*u - a*v))] Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 1.59 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.68

method	result
derivativdivides	$\frac{2b^2 \left( \frac{11bx^{\frac{3}{2}}}{8} + \frac{(13 \operatorname{arctanh}(\tanh(bx+a)) - 13bx)\sqrt{x}}{8} + \frac{35 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{8\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}} \right)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4} - \frac{2}{3(\operatorname{arctanh}(\tanh(bx+a)) - bx)}$
default	$\frac{2b^2 \left( \frac{11bx^{\frac{3}{2}}}{8} + \frac{(13 \operatorname{arctanh}(\tanh(bx+a)) - 13bx)\sqrt{x}}{8} + \frac{35 \operatorname{arctan}\left(\frac{b\sqrt{x}}{\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}}\right)}{8\sqrt{(\operatorname{arctanh}(\tanh(bx+a)) - bx)b}} \right)}{(\operatorname{arctanh}(\tanh(bx+a)) - bx)^4} - \frac{2}{3(\operatorname{arctanh}(\tanh(bx+a)) - bx)}$
risch	Expression too large to display

input `int(1/x^(5/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{2}{(\operatorname{arctanh}(\tanh(b*x+a)) - b*x)^4} b^2 * \left( \frac{11}{8} b*x^{(3/2)} + \frac{13}{8} * \operatorname{arctanh}(\tanh(b*x+a)) - 13/8 * b*x \right) * x^{(1/2)} / \operatorname{arctanh}(\tanh(b*x+a))^2 + 35/8 / \left( (\operatorname{arctanh}(\tanh(b*x+a)) - b*x) * b \right)^{(1/2)} * \operatorname{arctan}(b*x^{(1/2)} / \left( (\operatorname{arctanh}(\tanh(b*x+a)) - b*x) * b \right)^{(1/2)}) - 2/3 / (\operatorname{arctanh}(\tanh(b*x+a)) - b*x)^3 / x^{(3/2)} + 6 / (\operatorname{arctanh}(\tanh(b*x+a)) - b*x)^4 * b / x^{(1/2)}$$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.22

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \left[ \frac{105 (b^3 x^4 + 2 ab^2 x^3 + a^2 b x^2) \sqrt{-\frac{b}{a}} \log \left( \frac{bx + 2a\sqrt{x}\sqrt{-\frac{b}{a}} - a}{bx + a} \right) + 2 (105 b^3 x^4 + 210 a b^2 x^3 + 105 a^2 b x^2)}{24 (a^4 b^2 x^4 + 2 a^5 b x^3 + a^6 x^2)} \right]$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```
[1/24*(105*(b^3*x^4 + 2*a*b^2*x^3 + a^2*b*x^2)*sqrt(-b/a)*log((b*x + 2*a*sqrt(x)*sqrt(-b/a) - a)/(b*x + a)) + 2*(105*b^3*x^3 + 175*a*b^2*x^2 + 56*a^2*b*x - 8*a^3)*sqrt(x))/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2), 1/12*(105*(b^3*x^4 + 2*a*b^2*x^3 + a^2*b*x^2)*sqrt(b/a)*arctan(sqrt(x)*sqrt(b/a)) + (105*b^3*x^3 + 175*a*b^2*x^2 + 56*a^2*b*x - 8*a^3)*sqrt(x))/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2)]
```

**Sympy [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x^{5/2} \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(5/2)/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(1/(x**(5/2)*atanh(tanh(a + b*x))**3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.43

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{105 b^3 x^3 + 175 a b^2 x^2 + 56 a^2 b x - 8 a^3}{12 \left( a^4 b^2 x^{7/2} + 2 a^5 b x^{5/2} + a^6 x^{3/2} \right)} + \frac{35 b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4 \sqrt{ab} a^4}$$

input

```
integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")
```

output

```
1/12*(105*b^3*x^3 + 175*a*b^2*x^2 + 56*a^2*b*x - 8*a^3)/(a^4*b^2*x^(7/2) + 2*a^5*b*x^(5/2) + a^6*x^(3/2)) + 35/4*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^4)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.35

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{35 b^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4 \sqrt{ab} a^4} + \frac{2(9bx - a)}{3 a^4 x^{3/2}} + \frac{11 b^3 x^{3/2} + 13 ab^2 \sqrt{x}}{4 (bx + a)^2 a^4}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `35/4*b^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^4) + 2/3*(9*b*x - a)/(a^4*x^(3/2)) + 1/4*(11*b^3*x^(3/2) + 13*a*b^2*sqrt(x))/((b*x + a)^2*a^4)`

**Mupad [B] (verification not implemented)**

Time = 4.70 (sec) , antiderivative size = 1362, normalized size of antiderivative = 6.78

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(1/(x^(5/2)*atanh(tanh(a + b*x))^3),x)`

output

```
(x^(1/2)*((2*(2*b*(3*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 3*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 6*b*x) - 14*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(3*(2*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (56*b^2*x)/(3*(2*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))))/(2*b*x^2 - x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))^2 - (x^(1/2)*((280*b)/(3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (280*b^2*x)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4))/(2*b*x^2 - x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (70*2^(1/2)*b^(3/2)*log((b^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^(1/2)*(2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x) - 4*b^(1/2)*x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*...
```

**Reduce [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x^2} dx$$

input

```
int(1/x^(5/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))^3*x**2),x)
```

### 3.214 $\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$

Optimal result	1521
Mathematica [A] (verified)	1522
Rubi [A] (verified)	1522
Maple [A] (verified)	1530
Fricas [A] (verification not implemented)	1530
Sympy [F]	1531
Maxima [A] (verification not implemented)	1531
Giac [A] (verification not implemented)	1532
Mupad [B] (verification not implemented)	1532
Reduce [F]	1533

#### Optimal result

Integrand size = 15, antiderivative size = 228

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx =$$

$$\frac{63b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{4(bx - \operatorname{arctanh}(\tanh(a+bx)))^{11/2}}$$

$$+ \frac{63b^2}{4\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^5}$$

$$+ \frac{21b}{4x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{63}{20x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

$$+ \frac{4bx^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{9} + \frac{4b^2x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{7}$$

$$- \frac{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}{1} + \frac{4b^2x^{9/2} \operatorname{arctanh}(\tanh(a+bx))}{7}$$

output

```
-63/4*b^(5/2)*arctanh(b^(1/2)*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^(1/2))/(b
*x-arctanh(tanh(b*x+a)))^(11/2)+63/4*b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a))
)^(5)+21/4*b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^4+63/20/x^(5/2)/(b*x-arctanh
(tanh(b*x+a)))^3+9/4/b/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^2+7/4/b^2/x^(9/2
)/(b*x-arctanh(tanh(b*x+a)))-1/2/b/x^(7/2)/arctanh(tanh(b*x+a))^2+7/4/b^2/
x^(9/2)/arctanh(tanh(b*x+a))
```

**Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{1}{20} \left( \frac{75b^3 \sqrt{x}}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^5 \operatorname{arctanh}(\tanh(a + bx))} \right. \\ - \frac{315b^{5/2} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{-bx + \operatorname{arctanh}(\tanh(a + bx))}}\right)}{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^{11/2}} \\ - \frac{10b^3 \sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^2 (-bx + \operatorname{arctanh}(\tanh(a + bx)))^4} \\ \left. + \frac{8(36b^2x^2 - 7bx \operatorname{arctanh}(\tanh(a + bx)) + \operatorname{arctanh}(\tanh(a + bx))^2)}{x^{5/2} (bx - \operatorname{arctanh}(\tanh(a + bx)))^5} \right)$$

input `Integrate[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^3),x]`

output `((75*b^3*Sqrt[x])/((b*x - ArcTanh[Tanh[a + b*x]])^5*ArcTanh[Tanh[a + b*x]]) - (315*b^(5/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[-(b*x) + ArcTanh[Tanh[a + b*x]]]])/(-(b*x) + ArcTanh[Tanh[a + b*x]])^(11/2) - (10*b^3*Sqrt[x])/(ArcTanh[Tanh[a + b*x]]^2*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4) + (8*(36*b^2*x^2 - 7*b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2))/(x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^5))/20`

**Rubi [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.32, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$ , Rules used = {2599, 2599, 2594, 2594, 2594, 2594, 2594, 2593}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx$$

↓ 2599

$$\frac{7 \int \frac{1}{x^{9/2} \operatorname{arctanh}(\tanh(a+bx))^2} dx}{4b} - \frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2599

$$\frac{7 \left( -\frac{9 \int \frac{1}{x^{11/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{2b} - \frac{1}{bx^{9/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2594

$$\frac{7 \left( -\frac{9 \left( \frac{b \int \frac{1}{x^{9/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{9x^{9/2} (bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{9/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2594

$$\frac{7 \left( -\frac{9 \left( \frac{b \left( \frac{b \int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2} (bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{9x^{9/2} (bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{2b} - \frac{1}{bx^{9/2} \operatorname{arctanh}(\tanh(a+bx))} \right)}{4b} - \frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$

↓ 2594

$$\frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$$



$$\left( \frac{b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))} dx + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$


---

7

---

$\frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$   
 $\downarrow$  2594

$$\left( \int \frac{b \left( \frac{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{2}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$\frac{1}{2bx^{7/2} \operatorname{arctanh}(\tanh(a+bx))^2}$   
 $\downarrow$  2594

7	$\int \frac{b \left( \frac{b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$	2b
9	$\int \frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))}$	bx - arctanh(tanh(a+bx))

↓ 2593

$$\left( \frac{b}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{b \left( \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{bx - \operatorname{arctanh}(\tanh(a+bx))}}\right)}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \dots$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \dots$$


---


$$\frac{b}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \dots$$


---


$$\frac{9}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \dots$$

input `Int[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^3),x]`

output `(-7*((-9*(2/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*(2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (b*((-2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[b*x - ArcTanh[Tanh[a + b*x]]]])/(b*x - ArcTanh[Tanh[a + b*x]]^(3/2) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(b*x - ArcTanh[Tanh[a + b*x]])))/(2*b) - 1/(b*x^(9/2)*ArcTanh[Tanh[a + b*x]])))/(4*b) - 1/(2*b*x^(7/2)*ArcTanh[Tanh[a + b*x]]^2)`

### Defintions of rubi rules used

rule 2593 `Int[1/((u_)*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[-2*(ArcTanh[Sqrt[v]/Rt[-(b*u - a*v)/a, 2]]/(a*Rt[-(b*u - a*v)/a, 2])), x] /; NeQ[b*u - a*v, 0] && NegQ[(b*u - a*v)/a] /; PiecewiseLinearQ[u, v, x]`

rule 2594 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[v^(n + 1)/((n + 1)*(b*u - a*v)), x] - Simp[a*((n + 1)/((n + 1)*(b*u - a*v))) Int[v^(n + 1)/u, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && LtQ[n, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [A] (verified)

Time = 1.89 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.69

method	result
derivativdivides	$-\frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 x^{\frac{5}{2}}} - \frac{12b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^5 \sqrt{x}} + \frac{2b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^4 x^{\frac{3}{2}}}$
default	$-\frac{2}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)^3 x^{\frac{5}{2}}} - \frac{12b^2}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^5 \sqrt{x}} + \frac{2b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^4 x^{\frac{3}{2}}}$
risch	Expression too large to display

input `int(1/x^(7/2)/arctanh(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

output `-2/5/(arctanh(tanh(b*x+a))-b*x)^3/x^(5/2)-12/(arctanh(tanh(b*x+a))-b*x)^5*b^2/x^(1/2)+2/(arctanh(tanh(b*x+a))-b*x)^4*b/x^(3/2)-2/(arctanh(tanh(b*x+a))-b*x)^5*b^3*((15/8*b*x^(3/2)+(17/8*arctanh(tanh(b*x+a))-17/8*b*x)*x^(1/2))/arctanh(tanh(b*x+a))^2+63/8/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)*arctan(b*x^(1/2)/((arctanh(tanh(b*x+a))-b*x)*b)^(1/2)))`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.19

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^3} dx = \left[ \frac{315(b^4x^5 + 2ab^3x^4 + a^2b^2x^3)\sqrt{-\frac{b}{a}} \log\left(\frac{bx-2a\sqrt{x}\sqrt{-\frac{b}{a}}-a}{bx+a}\right) - 2(315b^4x^5 + 2ab^3x^4 + a^2b^2x^3)\sqrt{-\frac{b}{a}} \operatorname{arctan}\left(\frac{bx-2a\sqrt{x}\sqrt{-\frac{b}{a}}-a}{bx+a}\right)}{40(a^5b^2x^5 + 2a^6bx^4 + a^7x^3)} \right. \\ \left. - \frac{315(b^4x^5 + 2ab^3x^4 + a^2b^2x^3)\sqrt{\frac{b}{a}} \operatorname{arctan}\left(\sqrt{x}\sqrt{\frac{b}{a}}\right) + (315b^4x^4 + 525ab^3x^3 + 168a^2b^2x^2 - 24a^3bx + 8a^4)}{20(a^5b^2x^5 + 2a^6bx^4 + a^7x^3)} \right]$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="fricas")`

output

```
[1/40*(315*(b^4*x^5 + 2*a*b^3*x^4 + a^2*b^2*x^3)*sqrt(-b/a)*log((b*x - 2*a
*sqrt(x)*sqrt(-b/a) - a)/(b*x + a)) - 2*(315*b^4*x^4 + 525*a*b^3*x^3 + 168
*a^2*b^2*x^2 - 24*a^3*b*x + 8*a^4)*sqrt(x))/(a^5*b^2*x^5 + 2*a^6*b*x^4 + a
^7*x^3), -1/20*(315*(b^4*x^5 + 2*a*b^3*x^4 + a^2*b^2*x^3)*sqrt(b/a)*arctan
(sqrt(x)*sqrt(b/a)) + (315*b^4*x^4 + 525*a*b^3*x^3 + 168*a^2*b^2*x^2 - 24*
a^3*b*x + 8*a^4)*sqrt(x))/(a^5*b^2*x^5 + 2*a^6*b*x^4 + a^7*x^3)]
```

**Sympy [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{x^{7/2} \operatorname{atanh}^3(\tanh(a + bx))} dx$$

input

```
integrate(1/x**(7/2)/atanh(tanh(b*x+a))**3,x)
```

output

```
Integral(1/(x**(7/2)*atanh(tanh(a + b*x))**3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.43

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \frac{315 b^4 x^4 + 525 a b^3 x^3 + 168 a^2 b^2 x^2 - 24 a^3 b x + 8 a^4}{20 \left( a^5 b^2 x^{\frac{9}{2}} + 2 a^6 b x^{\frac{7}{2}} + a^7 x^{\frac{5}{2}} \right)} - \frac{63 b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4 \sqrt{ab} a^5}$$

input

```
integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="maxima")
```

output

```
-1/20*(315*b^4*x^4 + 525*a*b^3*x^3 + 168*a^2*b^2*x^2 - 24*a^3*b*x + 8*a^4)
/(a^5*b^2*x^(9/2) + 2*a^6*b*x^(7/2) + a^7*x^(5/2)) - 63/4*b^3*arctan(b*sqrt
t(x)/sqrt(a*b))/(sqrt(a*b)*a^5)
```



**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.35

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = -\frac{63 b^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4 \sqrt{ab} a^5} - \frac{15 b^4 x^{\frac{3}{2}} + 17 ab^3 \sqrt{x}}{4 (bx + a)^2 a^5} - \frac{2 (30 b^2 x^2 - 5 abx + a^2)}{5 a^5 x^{\frac{5}{2}}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^3,x, algorithm="giac")`

output `-63/4*b^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^5) - 1/4*(15*b^4*x^(3/2) + 17*a*b^3*sqrt(x))/((b*x + a)^2*a^5) - 2/5*(30*b^2*x^2 - 5*a*b*x + a^2)/(a^5*x^(5/2))`

**Mupad [B] (verification not implemented)**

Time = 4.07 (sec) , antiderivative size = 2151, normalized size of antiderivative = 9.43

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \text{Too large to display}$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))^3),x)`

output

```

16/(5*x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (x*(((448*b^4)/(3*(2*a*b - b
*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(ex
p(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log(
(2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (16*b^3*(
2*b*(3*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 3*log((2*exp(2*a)*exp(2*b*x))/(e
xp(2*a)*exp(2*b*x) + 1)) + 6*b*x) - 14*b*(log(2/(exp(2*a)*exp(2*b*x) + 1))
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/(3*(2
*a*b - b*(2*a - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 1
og(2/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)
) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4))*(1
og(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*ex
p(2*b*x) + 1)) + 2*b*x))/(2*b) - (112*b^3)/((2*a*b - b*(2*a - log((2*exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2*b*x) +
1)) + 2*b*x))*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*
x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (8*b^2*(2*b*(3*log(2/(exp(2*a
)*exp(2*b*x) + 1)) - 3*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) +
1)) + 6*b*x) - 14*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*ex
p(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)))/((2*a*b - b*(2*a - log((2*
exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + log(2/(exp(2*a)*exp(2...

```

**Reduce [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^3} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}(\tanh(bx + a))^3 x^3} dx$$

input

```
int(1/x^(7/2)/atanh(tanh(b*x+a))^3,x)
```

output

```
int(1/(sqrt(x)*atanh(tanh(a + b*x))^3*x**3),x)
```

### 3.215 $\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	1534
Mathematica [A] (verified)	1535
Rubi [A] (verified)	1535
Maple [A] (verified)	1537
Fricas [A] (verification not implemented)	1538
Sympy [F]	1538
Maxima [F]	1539
Giac [A] (verification not implemented)	1539
Mupad [F(-1)]	1539
Reduce [F]	1540

#### Optimal result

Integrand size = 17, antiderivative size = 142

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx =$$

$$\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^3}{8b^{5/2}}$$

$$+ \frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}$$

$$- \frac{x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{12b}$$

$$- \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{8b^2}$$

output

```
-1/8*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^3/b^(5/2)+1/3*x^(5/2)*arctanh(tanh(b*x+a))^(1/2)-1/12*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b-1/8*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)/b^2
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.73

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (3b^2 x^2 + 8bx \operatorname{arctanh}(\tanh(a + bx)) - 3(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3 \log(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))})}{24b^2} + \frac{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^3 \log(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))})}{8b^{5/2}}$$

input

```
Integrate[x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]],x]
```

output

```
(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(3*b^2*x^2 + 8*b*x*ArcTanh[Tanh[a + b*x]] - 3*ArcTanh[Tanh[a + b*x]]^2))/(24*b^2) + ((-(b*x) + ArcTanh[Tanh[a + b*x]])^3*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(8*b^(5/2))
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

↓ 2600

$$\frac{1}{3} x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{6} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2600

$$bx))) \left( \frac{\frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{4b} + \frac{x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)$$

↓ 2600

$$bx))) \left( \frac{\frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{2b} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{4b} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)$$

↓ 2596

$$\frac{1}{6} \left( \frac{\frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - 3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{4b} \right) (bx - \operatorname{arctanh}(\tanh(a+bx)))$$

input `Int [x^(3/2)*Sqrt [ArcTanh [Tanh [a + b*x]]], x]`

output `-1/6*(((3*((ArcTanh [(Sqrt [b]*Sqrt [x])/Sqrt [ArcTanh [Tanh [a + b*x]]])*(b*x - ArcTanh [Tanh [a + b*x]]))/b^(3/2) + (Sqrt [x]*Sqrt [ArcTanh [Tanh [a + b*x]]])/b)*(b*x - ArcTanh [Tanh [a + b*x]]))/(4*b) + (x^(3/2)*Sqrt [ArcTanh [Tanh [a + b*x]]])/(2*b))*(b*x - ArcTanh [Tanh [a + b*x]]) + (x^(5/2)*Sqrt [ArcTanh [Tanh [a + b*x]]])/3`

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Si
mplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v
]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u -
a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /;
PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n +
1, 0] && !(IGtQ[m, 0] && ( !IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n,
-2]
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x}}{b} \right)}{b} \right)}{b}$
default	$\frac{x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x}}{b} \right)}{b} \right)}{b}$

```
input int(x^(3/2)*arctanh(tanh(b*x+a))^(1/2), x, method=_RETURNVERBOSE)
```

```
output 1/3*x^(3/2)*arctanh(tanh(b*x+a))^(3/2)/b-(arctanh(tanh(b*x+a))-b*x)/b*(1/4
*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)/b-1/4*(arctanh(tanh(b*x+a))-b*x)/b*(1/
2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x
)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.97

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \left[ \frac{3 a^3 \sqrt{b} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2(8b^3x^2 + 2ab^2x - 3a^2b)\sqrt{bx+a}\sqrt{x}}{48b^3} - \frac{3a^3\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) - (8b^3x^2 + 2ab^2x - 3a^2b)\sqrt{bx+a}\sqrt{x}}{24b^3} \right]$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[1/48*(3*a^3*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(8*b^3*x^2 + 2*a*b^2*x - 3*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^3, -1/24*(3*a^3*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (8*b^3*x^2 + 2*a*b^2*x - 3*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^3]`

**Sympy [F]**

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x^{3/2} \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**(3/2)*atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(x**(3/2)*sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [F]**

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x^{3/2} \sqrt{\operatorname{artanh}(\tanh(bx + a))} dx$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(x^(3/2)*sqrt(arctanh(tanh(b*x + a))), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.42

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{1}{24} \sqrt{bx + a} \left( 2 \left( 4x + \frac{a}{b} \right) x - \frac{3a^2}{b^2} \right) \sqrt{x} - \frac{a^3 \log \left( \left| -\sqrt{b}\sqrt{x} + \sqrt{bx + a} \right| \right)}{8b^{5/2}}$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `1/24*sqrt(b*x + a)*(2*(4*x + a/b)*x - 3*a^2/b^2)*sqrt(x) - 1/8*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2)`

**Mupad [F(-1)]**

Timed out.

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int x^{3/2} \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `int(x^(3/2)*atanh(tanh(a + b*x))^(1/2),x)`

output `int(x^(3/2)*atanh(tanh(a + b*x))^(1/2), x)`



**Reduce [F]**

$$\int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x dx$$

input `int(x^(3/2)*atanh(tanh(b*x+a))^(1/2),x)`

output `int(sqrt(x)*sqrt(atanh(tanh(a + b*x)))*x,x)`

### 3.216 $\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$

Optimal result	1541
Mathematica [A] (verified)	1542
Rubi [A] (verified)	1542
Maple [A] (verified)	1544
Fricas [A] (verification not implemented)	1544
Sympy [F]	1545
Maxima [F]	1545
Giac [A] (verification not implemented)	1545
Mupad [F(-1)]	1546
Reduce [F]	1546

#### Optimal result

Integrand size = 17, antiderivative size = 104

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^2}{4b^{3/2}}$$

$$+ \frac{1}{2}x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}$$

$$- \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{4b}$$

output

```
-1/4*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh
(b*x+a)))^2/b^(3/2)+1/2*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)-1/4*x^(1/2)*(b*
x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.81

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$= \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (bx + \operatorname{arctanh}(\tanh(a + bx)))}{4b} - \frac{(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log\left(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)}{4b^{3/2}}$$

input `Integrate[Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(b*x + ArcTanh[Tanh[a + b*x]]))/(4*b) - (((-b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(4*b^(3/2))`

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2600$$

$$\frac{1}{2} x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow 2600$$

$$\begin{aligned}
 & \left( \frac{1}{2} x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right) \\
 & \quad \downarrow \text{2596} \\
 & \frac{1}{4} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{b^{3/2}} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))
 \end{aligned}$$

input `Int[Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `-1/4*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]])*(b*x - ArcTanh[Tanh[a + b*x]])/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b*(b*x - ArcTanh[Tanh[a + b*x]]) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/2`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u]*Sqrt[v]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1)) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`



**Sympy [F]**

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `integrate(x**(1/2)*atanh(tanh(b*x+a))**(1/2), x)`

output `Integral(sqrt(x)*sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [F]**

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{x} \sqrt{\operatorname{artanh}(\tanh(bx + a))} dx$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")`

output `integrate(sqrt(x)*sqrt(arctanh(tanh(b*x + a))), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.46

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \frac{1}{4} \sqrt{bx + a} \left( 2x + \frac{a}{b} \right) \sqrt{x} + \frac{a^2 \log \left( \left| -\sqrt{b} \sqrt{x} + \sqrt{bx + a} \right| \right)}{4 b^{\frac{3}{2}}}$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(1/2), x, algorithm="giac")`

output `1/4*sqrt(b*x + a)*(2*x + a/b)*sqrt(x) + 1/4*a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(3/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(a + bx))} dx$$

input `int(x^(1/2)*atanh(tanh(a + b*x))^(1/2), x)`output `int(x^(1/2)*atanh(tanh(a + b*x))^(1/2), x)`**Reduce [F]**

$$\int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(1/2)*atanh(tanh(b*x+a))^(1/2), x)`output `int(sqrt(x)*sqrt(atanh(tanh(a + b*x))), x)`

**3.217**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx$

Optimal result	1547
Mathematica [A] (verified)	1547
Rubi [A] (verified)	1548
Maple [A] (verified)	1549
Fricas [A] (verification not implemented)	1550
Sympy [F]	1550
Maxima [F]	1551
Giac [A] (verification not implemented)	1551
Mupad [F(-1)]	1551
Reduce [F]	1552

**Optimal result**

Integrand size = 17, antiderivative size = 61

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx$$

$$= -\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{\sqrt{b} + \sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

$-\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)})*(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/b^{(1/2)}+x^{(1/2)}*\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}$

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.02

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx$$

$$= \sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} + \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx))) \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{\sqrt{b}}$$



input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[x], x]`

output `Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]] + ((-(b*x) + ArcTanh[Tanh[a + b*x]])*  
Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]])/Sqrt[b]`

### Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx$$

↓ 2600

$$\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2596

$$\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a + bx)))}{\sqrt{b}}$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/Sqrt[x], x]`

output `-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]]*(b*x - ArcTanh[  
Tanh[a + b*x]]))/Sqrt[b] + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]`

## Definitions of rubi rules used

rule 2596

```
Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

## Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{\sqrt{b}}$	49
default	$\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{\sqrt{b}}$	49

input

```
int(arctanh(tanh(b*x+a))^(1/2)/x^(1/2), x, method=_RETURNVERBOSE)
```

output

```
x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.48

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx$$

$$= \left[ \frac{a\sqrt{b} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2\sqrt{bx+a}b\sqrt{x}}{2b}, \right.$$

$$\left. - \frac{a\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) - \sqrt{bx+a}b\sqrt{x}}{b} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(1/2),x, algorithm="fricas")`

output `[1/2*(a*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*sqrt(b*x + a)*b*sqrt(x))/b, -(a*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - sqrt(b*x + a)*b*sqrt(x))/b]`

**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**(1/2),x)`

output `Integral(sqrt(atanh(tanh(a + b*x)))/sqrt(x), x)`

**Maxima [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx = \int \frac{\sqrt{\operatorname{artanh}(\tanh(bx + a))}}{\sqrt{x}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(arctanh(tanh(b*x + a)))/sqrt(x), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx = -\frac{a \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{\sqrt{b}} + \sqrt{bx + a}\sqrt{x}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(1/2),x, algorithm="giac")`

output `-a*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/sqrt(b) + sqrt(b*x + a)*sqrt(x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{\sqrt{x}} dx$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(1/2),x)`

output `int(atanh(tanh(a + b*x))^(1/2)/x^(1/2), x)`

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x} dx$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/x,x)`

$$3.218 \quad \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx$$

Optimal result	1553
Mathematica [A] (verified)	1553
Rubi [A] (verified)	1554
Maple [B] (verified)	1555
Fricas [A] (verification not implemented)	1556
Sympy [F]	1556
Maxima [A] (verification not implemented)	1556
Giac [A] (verification not implemented)	1557
Mupad [F(-1)]	1557
Reduce [F]	1558

### Optimal result

Integrand size = 17, antiderivative size = 49

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx = 2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}}$$

output

```
2*b^(1/2)*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))-2*arctanh(tanh(b*x+a))^(1/2)/x^(1/2)
```

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx = -\frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} + 2\sqrt{b} \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)$$

input

```
Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(3/2),x]
```

output

```
(-2*Sqrt[ArcTanh[Tanh[a + b*x]])/Sqrt[x] + 2*Sqrt[b]*Log[b*Sqrt[x] + Sqrt
[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2599, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx$$

↓ 2599

$$b \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}}$$

↓ 2596

$$2\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}}$$

input

```
Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(3/2), x]
```

output

```
2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]] - (2*Sqr
t[ArcTanh[Tanh[a + b*x]])/Sqrt[x]
```

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Si
mplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v
]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 96 vs. 2(37) = 74.

Time = 0.40 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.98

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{4b \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{2\sqrt{b}} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{4b \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{2\sqrt{b}} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx}$

```
input int(arctanh(tanh(b*x+a))^(1/2)/x^(3/2), x, method=_RETURNVERBOSE)
```

```
output -2/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+4*b/(arct
anh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*
(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```



**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.76

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx = \left[ \frac{\sqrt{bx} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) - 2\sqrt{bx+a}\sqrt{x}}{x}, \right. \\ \left. - \frac{2\left(\sqrt{-bx} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) + \sqrt{bx+a}\sqrt{x}\right)}{x} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(3/2),x, algorithm="fricas")`

output `[(sqrt(b)*x*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) - 2*sqrt(b*x + a)*sqrt(x))/x, -2*(sqrt(-b)*x*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + sqrt(b*x + a)*sqrt(x))/x]`

**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a+bx))}}{x^{3/2}} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**(3/2),x)`

output `Integral(sqrt(atanh(tanh(a + b*x)))/x**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{3/2}} dx = 2\sqrt{b} \log\left(\frac{b\sqrt{x}}{\sqrt{ab}} + \sqrt{\frac{bx}{a} + 1}\right) - \frac{2\sqrt{bx+a}}{\sqrt{x}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(3/2),x, algorithm="maxima")`

output  $2*\sqrt{b}*\log(b*\sqrt{x}/\sqrt{a*b} + \sqrt{b*x/a + 1}) - 2*\sqrt{b*x + a}/\sqrt{t(x)}$

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx = -\sqrt{b} \log\left(\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2\right) + \frac{4a\sqrt{b}}{\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2 - a}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(3/2),x, algorithm="giac")`

output  $-\sqrt{b}*\log((\sqrt{b}*\sqrt{x} - \sqrt{b*x + a})^2) + 4*a*\sqrt{b}/((\sqrt{b}*\sqrt{x} - \sqrt{b*x + a})^2 - a)$

### Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a + bx))}}{x^{3/2}} dx$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(3/2),x)`

output `int(atanh(tanh(a + b*x))^(1/2)/x^(3/2), x)`

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \sqrt{x} \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x} dx \right) b}{\sqrt{x}}$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(3/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x))) + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x),x)*b)/sqrt(x)`

$$3.219 \quad \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx$$

Optimal result	1559
Mathematica [A] (verified)	1559
Rubi [A] (verified)	1560
Maple [A] (verified)	1561
Fricas [A] (verification not implemented)	1561
Sympy [F]	1561
Maxima [A] (verification not implemented)	1562
Giac [B] (verification not implemented)	1562
Mupad [B] (verification not implemented)	1562
Reduce [F]	1563

### Optimal result

Integrand size = 17, antiderivative size = 35

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `2/3*arctanh(tanh(b*x+a))^(3/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))`

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}(3bx - 3\operatorname{arctanh}(\tanh(a+bx)))}$$

input `Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(5/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(x^(3/2)*(3*b*x - 3*ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(5/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}$	29
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}$	29

input `int(arctanh(tanh(b*x+a))^(1/2)/x^(5/2),x,method=_RETURNVERBOSE)`output `-2/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh(tanh(b*x+a))^(3/2)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.43

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx = -\frac{2(bx+a)^{\frac{3}{2}}}{3ax^{\frac{3}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(5/2),x, algorithm="fricas")`output `-2/3*(b*x + a)^(3/2)/(a*x^(3/2))`**Sympy [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx = \int \frac{\sqrt{\operatorname{atanh}(\tanh(a+bx))}}{x^{\frac{5}{2}}} dx$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**(5/2),x)`output `Integral(sqrt(atanh(tanh(a + b*x)))/x**(5/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.43

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx = -\frac{2(bx + a)^{\frac{3}{2}}}{3ax^{\frac{3}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(5/2),x, algorithm="maxima")`

output `-2/3*(b*x + a)^(3/2)/(a*x^(3/2))`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(29) = 58.

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx = \frac{4 \left( 3b^{\frac{3}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^4 + a^2b^{\frac{3}{2}} \right)}{3 \left( \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a \right)^3}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(5/2),x, algorithm="giac")`

output `4/3*(3*b^(3/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 + a^2*b^(3/2))/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^3`

**Mupad [B] (verification not implemented)**

Time = 3.97 (sec) , antiderivative size = 210, normalized size of antiderivative = 6.00

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx = \frac{2 \ln \left( \frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1} \right) \sqrt{\frac{\ln \left( \frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1} \right)}{2} - \frac{\ln \left( \frac{2}{e^{2a}e^{2bx}+1} \right)}{2}} - 2 \ln \left( \frac{2}{e^{2a}e^{2bx}+1} \right) \sqrt{\frac{\ln \left( \frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1} \right)}{2} - \frac{\ln \left( \frac{2}{e^{2a}e^{2bx}+1} \right)}{2}}}{\sqrt{x} \left( 3x \ln \left( \frac{2}{e^{2a}e^{2bx}+1} \right) - 3x \ln \left( \frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1} \right) + 6 \right)}$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(5/2),x)`

output `(2*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2) - 2*log(2/(exp(2*a)*exp(2*b*x) + 1))*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2))/(x^(1/2)*(3*x*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 3*x*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 6*b*x^2))`

### Reduce [F]

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \sqrt{x} \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x^2} dx \right) bx}{3\sqrt{x} x}$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(5/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x))) + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**2),x)*b*x)/(3*sqrt(x)*x)`



$$3.220 \quad \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx$$

Optimal result	1564
Mathematica [A] (verified)	1564
Rubi [A] (verified)	1565
Maple [A] (verified)	1566
Fricas [A] (verification not implemented)	1567
Sympy [F(-1)]	1567
Maxima [A] (verification not implemented)	1567
Giac [A] (verification not implemented)	1568
Mupad [B] (verification not implemented)	1568
Reduce [F]	1569

### Optimal result

Integrand size = 17, antiderivative size = 72

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx = \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
4/15*b*arctanh(tanh(b*x+a))^(3/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/5
*arctanh(tanh(b*x+a))^(3/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))
```

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.67

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx = \frac{2(5bx - 3\operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{15x^{5/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input

```
Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(7/2),x]
```

output

```
(2*(5*b*x - 3*ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))/(15*x^(5/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)
```

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx$$

↓ 2602

$$\frac{2b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{5/2}} dx}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input

```
Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(7/2), x]
```

output

```
(4*b*ArcTanh[Tanh[a + b*x]]^(3/2))/(15*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]]))
```

## Definitions of rubi rules used

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

## Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}}$	59
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}}$	59

input

```
int(arctanh(tanh(b*x+a))^(1/2)/x^(7/2), x, method=_RETURNVERBOSE)
```

output

```
-2/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(3/2)+4/15*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(3/2)*arctanh(tanh(b*x+a))^(3/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.47

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \frac{2(2b^2x^2 - abx - 3a^2)\sqrt{bx + a}}{15a^2x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(7/2),x, algorithm="fricas")`

output `2/15*(2*b^2*x^2 - a*b*x - 3*a^2)*sqrt(b*x + a)/(a^2*x^(5/2))`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**(7/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.47

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \frac{2(2b^2x^2 - abx - 3a^2)\sqrt{bx + a}}{15a^2x^{5/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(7/2),x, algorithm="maxima")`

output `2/15*(2*b^2*x^2 - a*b*x - 3*a^2)*sqrt(b*x + a)/(a^2*x^(5/2))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.56

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \frac{8 \left( 15 b^{5/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^6 + 5 a b^{5/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^4 + 5 a^2 b^{5/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a^3 b^{5/2} \right)}{15 \left( \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a \right)^5}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(7/2),x, algorithm="giac")`

output `8/15*(15*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 + 5*a*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 + 5*a^2*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a^3*b^(5/2))/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^5`

**Mupad [B] (verification not implemented)**

Time = 3.59 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.42

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2}} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}{15 \left( \frac{16b^2x^2}{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2} + \dots \right)} x^{5/2}$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(7/2),x)`

output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((16*b^2*x^2)/(15*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (4*b*x)/(15*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) - 2/5))/x^(5/2)`

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \sqrt{x} \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x^3} dx \right) b x^2}{5\sqrt{x} x^2}$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(7/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x))) + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**3),x)*b*x**2)/(5*sqrt(x)*x**2)`

**3.221**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx$

Optimal result	1570
Mathematica [A] (verified)	1570
Rubi [A] (verified)	1571
Maple [A] (verified)	1572
Fricas [A] (verification not implemented)	1573
Sympy [F(-1)]	1573
Maxima [A] (verification not implemented)	1574
Giac [A] (verification not implemented)	1574
Mupad [B] (verification not implemented)	1575
Reduce [F]	1575

**Optimal result**

Integrand size = 17, antiderivative size = 110

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{105x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{8b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
16/105*b^2*arctanh(tanh(b*x+a))^(3/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^3
+8/35*b*arctanh(tanh(b*x+a))^(3/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/
7*arctanh(tanh(b*x+a))^(3/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{3/2} (35b^2x^2 - 42bx \operatorname{arctanh}(\tanh(a+bx)) + 15a)}{105x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input

```
Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(9/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(3/2)*(35*b^2*x^2 - 42*b*x*ArcTanh[Tanh[a + b*x]] + 15*ArcTanh[Tanh[a + b*x]]^2))/(105*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx$$

↓ 2602

$$\frac{4b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{7/2}} dx}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2602

$$\frac{4b \left( \frac{2b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + 4b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) \frac{1}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

input

```
Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(9/2), x]
```



output

```
(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]]))
+ (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(3/2))/(15*x^(3/2)*(b*x - ArcTanh[Ta
nh[a + b*x]]^2) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/(5*x^(5/2)*(b*x - ArcT
anh[Tanh[a + b*x]])))))/(7*(b*x - ArcTanh[Tanh[a + b*x]]))
```

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /;
NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[
m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + S
imp[b*((m + n + 2)/((m + 1)*(b*u - a*v))) Int[u^(m + 1)*v^n, x], x] /; Ne
Q[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m
, -1]
```

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} - \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \frac{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}} \right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	105
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} - \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \frac{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{3}{2}}} \right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	105

input

```
int(arctanh(tanh(b*x+a))^(1/2)/x^(9/2), x, method=_RETURNVERBOSE)
```

output

```
-2/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(3/2)-8/7*b/(
arctanh(tanh(b*x+a))-b*x)*(-1/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh
(tanh(b*x+a))^(3/2)+2/15*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(3/2)*arctanh(ta
nh(b*x+a))^(3/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = -\frac{2(8b^3x^3 - 4ab^2x^2 + 3a^2bx + 15a^3)\sqrt{bx+a}}{105a^3x^{7/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(1/2)/x^(9/2),x, algorithm="fricas")
```

output

```
-2/105*(8*b^3*x^3 - 4*a*b^2*x^2 + 3*a^2*b*x + 15*a^3)*sqrt(b*x + a)/(a^3*x
^(7/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = \text{Timed out}$$

input

```
integrate(atanh(tanh(b*x+a))**(1/2)/x**(9/2),x)
```

output

```
Timed out
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = -\frac{2(8b^3x^3 - 4ab^2x^2 + 3a^2bx + 15a^3)\sqrt{bx+a}}{105a^3x^{7/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(9/2),x, algorithm="maxima")`

output `-2/105*(8*b^3*x^3 - 4*a*b^2*x^2 + 3*a^2*b*x + 15*a^3)*sqrt(b*x + a)/(a^3*x^(7/2))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.25

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{9/2}} dx = \frac{32 \left( 70 b^{7/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx+a} \right)^8 + 35 ab^{7/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx+a} \right)^6 + 21 a^2 b^{7/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx+a} \right)^4 - 7 a^3 b^{7/2} \left( \sqrt{b}\sqrt{x} - \sqrt{bx+a} \right)^2 + a^4 b^{7/2} \right)}{105 \left( \left( \sqrt{b}\sqrt{x} - \sqrt{bx+a} \right)^2 - a \right)^7}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(9/2),x, algorithm="giac")`

output `32/105*(70*b^(7/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^8 + 35*a*b^(7/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 + 21*a^2*b^(7/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 - 7*a^3*b^(7/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 + a^4*b^(7/2))/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^7`

**Mupad [B] (verification not implemented)**

Time = 3.84 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.13

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{9/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{105 \left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2} +$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(9/2), x)`output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((32*b^2*x^2)/(105*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (128*b^3*x^3)/(105*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (4*b*x)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) - 2/7)/x^(7/2)`**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{9/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \sqrt{x} \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x^4} dx \right) b x^3}{7\sqrt{x} x^3}$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(9/2), x)`output `( - 2*sqrt(atanh(tanh(a + b*x))) + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**4), x)*b*x**3)/(7*sqrt(x)*x**3)`

**3.222**  $\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx$

Optimal result	1576
Mathematica [A] (verified)	1577
Rubi [A] (verified)	1577
Maple [A] (verified)	1579
Fricas [A] (verification not implemented)	1580
Sympy [F(-1)]	1580
Maxima [A] (verification not implemented)	1580
Giac [A] (verification not implemented)	1581
Mupad [B] (verification not implemented)	1581
Reduce [F]	1582

**Optimal result**

Integrand size = 17, antiderivative size = 148

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx = \frac{32b^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{315x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{105x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{4b \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{21x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

```
output 32/315*b^3*arctanh(tanh(b*x+a))^(3/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^4
+16/105*b^2*arctanh(tanh(b*x+a))^(3/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^3
+3+4/21*b*arctanh(tanh(b*x+a))^(3/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^2+2
/9*arctanh(tanh(b*x+a))^(3/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))
```

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.55

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{11/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2} (105b^3x^3 - 189b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 135bx\operatorname{arctanh}(\tanh(a + bx)) - 35\operatorname{arctanh}(\tanh(a + bx))^3)}{315x^{9/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input

```
Integrate[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(11/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(3/2)*(105*b^3*x^3 - 189*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 135*b*x*ArcTanh[Tanh[a + b*x]]^2 - 35*ArcTanh[Tanh[a + b*x]]^3))/(315*x^(9/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)
```

### Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{11/2}} dx$$

↓ 2602

$$\frac{2b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{9/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2602

$$\frac{2b \left( \frac{4b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{7/2}} dx}{7(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2602

$$\begin{aligned}
& 2b \left( \frac{4b \left( \frac{2b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{5/2}} dx}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \\
& \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow \text{2598} \\
& 2b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right)}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}
\end{aligned}$$

input `Int[Sqrt[ArcTanh[Tanh[a + b*x]]]/x^(11/2), x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(3/2))/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*b*((2*ArcTanh[Tanh[a + b*x]]^(3/2))/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) + (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(3/2))/(15*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]]^2) + (2*ArcTanh[Tanh[a + b*x]]^(3/2))/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])))))/(7*(b*x - ArcTanh[Tanh[a + b*x]])))/(3*(b*x - ArcTanh[Tanh[a + b*x]]))`

Defintions of rubi rules used

```
rule 2598 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 2602 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}}-\frac{4b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}-\frac{4b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{2}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}\right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}\right)}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}}-\frac{4b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}-\frac{4b\left(-\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{2}{15(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}\right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}\right)}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}$

```
input int(arctanh(tanh(b*x+a))^(1/2)/x^(11/2), x, method=_RETURNVERBOSE)
```

```
output -2/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arctanh(tanh(b*x+a))^(3/2)-4/3*b/(arctanh(tanh(b*x+a))-b*x)*(-1/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(3/2)-4/7*b/(arctanh(tanh(b*x+a))-b*x)*(-1/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(3/2)+2/15*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(3/2)*arctanh(tanh(b*x+a))^(3/2))
```



**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.38

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx = \frac{2(16b^4x^4 - 8ab^3x^3 + 6a^2b^2x^2 - 5a^3bx - 35a^4)\sqrt{bx+a}}{315a^4x^{\frac{9}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(11/2),x, algorithm="fricas")`

output `2/315*(16*b^4*x^4 - 8*a*b^3*x^3 + 6*a^2*b^2*x^2 - 5*a^3*b*x - 35*a^4)*sqrt(b*x + a)/(a^4*x^(9/2))`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(1/2)/x**(11/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.38

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{x^{11/2}} dx = \frac{2(16b^4x^4 - 8ab^3x^3 + 6a^2b^2x^2 - 5a^3bx - 35a^4)\sqrt{bx+a}}{315a^4x^{\frac{9}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(11/2),x, algorithm="maxima")`

output `2/315*(16*b^4*x^4 - 8*a*b^3*x^3 + 6*a^2*b^2*x^2 - 5*a^3*b*x - 35*a^4)*sqrt(b*x + a)/(a^4*x^(9/2))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{11/2}} dx = \frac{64 \left( 315 b^{\frac{9}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^{10} + 189 ab^{\frac{9}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^8 + 84 a^2 b^{\frac{9}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^6 - 36 a^3 b^{\frac{9}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^4 + 9 a^4 b^{\frac{9}{2}} \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a^5 b^{\frac{9}{2}} \right)}{\left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a^9}$$

315

input `integrate(arctanh(tanh(b*x+a))^(1/2)/x^(11/2),x, algorithm="giac")`

output `64/315*(315*b^(9/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^10 + 189*a*b^(9/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^8 + 84*a^2*b^(9/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 - 36*a^3*b^(9/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 + 9*a^4*b^(9/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a^5*b^(9/2))/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^9`

**Mupad [B] (verification not implemented)**

Time = 3.74 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.99

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{11/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{105 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2} + \dots$$

input `int(atanh(tanh(a + b*x))^(1/2)/x^(11/2),x)`

output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((16*b^2*x^2)/(105*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (128*b^3*x^3)/(315*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (512*b^4*x^4)/(315*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4) + (4*b*x)/(63*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) - 2/9)/x^(9/2)`

**Reduce [F]**

$$\int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{11/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} + \sqrt{x} \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x^5} dx \right) b x^4}{9\sqrt{x} x^4}$$

input `int(atanh(tanh(b*x+a))^(1/2)/x^(11/2),x)`

output `( - 2*sqrt(atanh(tanh(a + b*x))) + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**5),x)*b*x**4)/(9*sqrt(x)*x**4)`

### 3.223 $\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	1583
Mathematica [A] (verified)	1583
Rubi [A] (verified)	1584
Maple [A] (verified)	1586
Fricas [A] (verification not implemented)	1587
Sympy [F]	1587
Maxima [F]	1588
Giac [A] (verification not implemented)	1588
Mupad [F(-1)]	1588
Reduce [F]	1589

#### Optimal result

Integrand size = 17, antiderivative size = 177

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{3 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^4}{64b^{5/2}} - \frac{1}{8} x^{5/2} (bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{x^{3/2} (bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{32b}$$

output

```
3/64*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^4/b^(5/2)-1/8*x^(5/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)+1/32*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)/b+3/64*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^3*arctanh(tanh(b*x+a))^(1/2)/b^2+1/4*x^(5/2)*arctanh(tanh(b*x+a))^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.69

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{b}\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}(-3b^3x^3 + 11b^2x^2 \operatorname{arctanh}(\tanh(a + bx)))}{32b^2}$$

input

```
Integrate[x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2),x]
```

output

```
(Sqrt[b]*Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-3*b^3*x^3 + 11*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 11*b*x*ArcTanh[Tanh[a + b*x]]^2 - 3*ArcTanh[Tanh[a + b*x]]^3) + 3*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(64*b^(5/2))
```

### Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2600, 2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx \\
 & \quad \downarrow 2600 \\
 & \frac{1}{4} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{8} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx \\
 & \quad \downarrow 2600 \\
 & \frac{1}{4} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{8} (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{1}{3} x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{6} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \right) \\
 & \quad \downarrow 2600 \\
 & \frac{1}{4} x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{8} (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{1}{3} x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{6} (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{a}} dx}{4b} \right) \right) \\
 & \quad \downarrow 2600
 \end{aligned}$$

$$\begin{aligned}
 & \frac{1}{4}x^{5/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{3}{8}(bx - \operatorname{arctanh}(\tanh(a+bx))) \\
 & \left( \frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \right) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}{\operatorname{arctanh}(\tanh(a+bx))} \right) \\
 & \quad \downarrow \text{2596} \\
 & \frac{1}{4}x^{5/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \\
 & \left( \frac{3}{8} \left( \frac{1}{3}x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{6} \left( \frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{4b} \right)}{\operatorname{arctanh}(\tanh(a+bx))} \right) \right) \right)
 \end{aligned}$$

input `Int[x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(-3*(-1/6*(((3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b)*(b*x - ArcTanh[Tanh[a + b*x]]))/(4*b) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/(2*b))*(b*x - ArcTanh[Tanh[a + b*x]]) + (x^(5/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/3*(b*x - ArcTanh[Tanh[a + b*x]]))/8 + (x^(5/2)*ArcTanh[Tanh[a + b*x]]^(3/2))/4`

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Si
mplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v
]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u -
a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /;
PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n +
1, 0] && !(IGtQ[m, 0] && ( !IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n,
-2]
```

**Maple [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{4b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6b}}{\frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6b}}$
default	$\frac{x^{\frac{3}{2}} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{4b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6b}}{\frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6b}}$

```
input int(x^(3/2)*arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)
```

output

```
1/4*x^(3/2)*arctanh(tanh(b*x+a))^(5/2)/b-3/4*(arctanh(tanh(b*x+a))-b*x)/b*
(1/6*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)/b-1/6*(arctanh(tanh(b*x+a))-b*x)/b
*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1
/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*
x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.90

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \left[ \frac{3 a^4 \sqrt{b} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2(16b^4x^3 + 24ab^3x^2 + 12a^2b^2x - 3a^3b)\sqrt{bx+a}\sqrt{x}}{128b^3} \right]$$

input

```
integrate(x^(3/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
[1/128*(3*a^4*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2
*(16*b^4*x^3 + 24*a*b^3*x^2 + 2*a^2*b^2*x - 3*a^3*b)*sqrt(b*x + a)*sqrt(x)
)/b^3, -1/64*(3*a^4*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (16*
b^4*x^3 + 24*a*b^3*x^2 + 2*a^2*b^2*x - 3*a^3*b)*sqrt(b*x + a)*sqrt(x))/b^3
]
```

**Sympy [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int x^{3/2} \operatorname{atanh}^{3/2}(\tanh(a+bx)) dx$$

input

```
integrate(x**(3/2)*atanh(tanh(b*x+a))**(3/2),x)
```

output

```
Integral(x**(3/2)*atanh(tanh(a + b*x))**(3/2), x)
```



**Maxima [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int x^{3/2} \operatorname{artanh}(\tanh(bx + a))^{3/2} dx$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(x^(3/2)*arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.83

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{1}{384} \sqrt{2} \left( 8 \sqrt{2} \left( \sqrt{bx + a} \left( 2 \left( 4x + \frac{a}{b} \right) x - \frac{3a^2}{b^2} \right) \sqrt{x} - \frac{3a^3 \log \left( \left| -\sqrt{bx + a} \right| \right)}{\sqrt{x}} \right) \right)$$

input `integrate(x^(3/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/384*sqrt(2)*(8*sqrt(2)*(sqrt(b*x + a)*(2*(4*x + a/b)*x - 3*a^2/b^2)*sqrt(x) - 3*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2))*a + sqrt(2)*((2*(4*(6*x + a/b)*x - 5*a^2/b^2)*x + 15*a^3/b^3)*sqrt(b*x + a)*sqrt(x) + 15*a^4*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(7/2))*b)`

**Mupad [F(-1)]**

Timed out.

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int x^{3/2} \operatorname{atanh}(\tanh(a + bx))^{3/2} dx$$

input `int(x^(3/2)*atanh(tanh(a + b*x))^(3/2),x)`

output `int(x^(3/2)*atanh(tanh(a + b*x))^(3/2), x)`

**Reduce [F]**

$$\int x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a)) x dx$$

input `int(x^(3/2)*atanh(tanh(b*x+a))^(3/2),x)`

output `int(sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*x,x)`

### 3.224 $\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$

Optimal result	1590
Mathematica [A] (verified)	1591
Rubi [A] (verified)	1591
Maple [A] (verified)	1593
Fricas [A] (verification not implemented)	1593
Sympy [F]	1594
Maxima [F]	1594
Giac [A] (verification not implemented)	1595
Mupad [F(-1)]	1595
Reduce [F]	1596

#### Optimal result

Integrand size = 17, antiderivative size = 139

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^3}{8b^{3/2}} - \frac{1}{4}x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{8b}$$

output

```
1/8*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^3/b^(3/2)-1/4*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)+1/8*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)/b+1/3*x^(3/2)*arctanh(tanh(b*x+a))^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (-3b^2 x^2 + 8bx \operatorname{arctanh}(\tanh(a + bx)) + 3 \operatorname{arctanh}(\tanh(a + bx)))}{24b} + \frac{(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \log(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))})}{8b^{3/2}}$$

input `Integrate[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-3*b^2*x^2 + 8*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2))/(24*b) + ((b*x - ArcTanh[Tanh[a + b*x]])^3*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(8*b^(3/2))`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$$

$$\downarrow 2600$$

$$\frac{1}{3} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{1}{2} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} dx$$

$$\downarrow 2600$$

$$\begin{aligned}
 & \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx \right) \\
 & \quad \downarrow \text{2600} \\
 & \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{2b} \right) \right) \\
 & \quad \downarrow \text{2596} \\
 & \frac{1}{2} \left( \frac{1}{2}x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{4} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}} (bx - \operatorname{arctanh}(\tanh(a+bx))) + \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))} \right) \right)
 \end{aligned}$$

input `Int[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `-1/2*((-1/4*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b*(b*x - ArcTanh[Tanh[a + b*x]])) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/2*(b*x - ArcTanh[Tanh[a + b*x]])) + (x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2))/3`

**Defintions of rubi rules used**

rule 2596 `Int[1/(Sqrt[u]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1)) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$
default	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$

input

```
int(x^(1/2)*arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)
```

output

```
1/3*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)/b-1/3*(arctanh(tanh(b*x+a))-b*x)/b*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(ax + bx))^{3/2} dx = \left[ \frac{3 a^3 \sqrt{b} \log \left( 2 b x - 2 \sqrt{b x + a} \sqrt{b} \sqrt{x} + a \right) + 2 (8 b^3 x^2 + 14 a b^2 x + 3 a^2 b) \sqrt{b x + a} \sqrt{x}}{48 b^2}, \frac{3 a^3 \sqrt{b}}{\dots} \right]$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[1/48*(3*a^3*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(8*b^3*x^2 + 14*a*b^2*x + 3*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^2, 1/24*(3*a^3*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (8*b^3*x^2 + 14*a*b^2*x + 3*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^2]`

### Sympy [F]

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int \sqrt{x} \operatorname{artanh}^{\frac{3}{2}}(\tanh(a + bx)) dx$$

input `integrate(x**(1/2)*atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(sqrt(x)*atanh(tanh(a + b*x))**(3/2), x)`

### Maxima [F]

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int \sqrt{x} \operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}} dx$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(sqrt(x)*arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.88

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \frac{1}{48} \sqrt{2} \left( 6 \sqrt{2} \left( \sqrt{bx+a} \left( 2x + \frac{a}{b} \right) \sqrt{x} + \frac{a^2 \log \left( \left| -\sqrt{b} \sqrt{x} + \sqrt{bx+a} \right| \right)}{b^{3/2}} \right) a + \sqrt{2} \left( \sqrt{bx+a} \right) \right)$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/48*sqrt(2)*(6*sqrt(2)*(sqrt(b*x + a)*(2*x + a/b)*sqrt(x) + a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(3/2))*a + sqrt(2)*(sqrt(b*x + a)*(2*(4*x + a/b)*x - 3*a^2/b^2)*sqrt(x) - 3*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2))*b)`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx = \int \sqrt{x} \operatorname{atanh}(\tanh(a + bx))^{3/2} dx$$

input `int(x^(1/2)*atanh(tanh(a + b*x))^(3/2),x)`

output `int(x^(1/2)*atanh(tanh(a + b*x))^(3/2), x)`



**Reduce [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))} \operatorname{atanh}(\tanh(bx+a)) dx$$

input `int(x^(1/2)*atanh(tanh(b*x+a))^(3/2),x)`

output `int(sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)),x)`

**3.225**  $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}} dx$

Optimal result	1597
Mathematica [A] (verified)	1598
Rubi [A] (verified)	1598
Maple [A] (verified)	1600
Fricas [A] (verification not implemented)	1600
Sympy [F]	1601
Maxima [F]	1601
Giac [A] (verification not implemented)	1601
Mupad [F(-1)]	1602
Reduce [F]	1602

**Optimal result**

Integrand size = 17, antiderivative size = 101

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \frac{3\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^2}{4\sqrt{b}} - \frac{3}{4}\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{1}{2}\sqrt{x}\operatorname{arctanh}(\tanh(a + bx))^{3/2}$$

output

```
3/4*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^2/b^(1/2)-3/4*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)+1/2*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \frac{1}{4} \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (-3bx + 5\operatorname{arctanh}(\tanh(a + bx))) \right. \\ \left. + \frac{3(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log \left( b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} \right)}{\sqrt{b}} \right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/Sqrt[x], x]
```

output

```
(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-3*b*x + 5*ArcTanh[Tanh[a + b*x]])
+ (3*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcT
anh[Tanh[a + b*x]]]])/Sqrt[b])/4
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx$$

↓ 2600

$$\frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx$$

↓ 2600

$$\frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \right)$$

$$\begin{array}{c}
 \downarrow \text{2596} \\
 \frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \\
 \frac{3}{4} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{\sqrt{b}} \right. \\
 \left. \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\operatorname{arctanh}(\tanh(a + bx))} \right) (bx -
 \end{array}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/Sqrt[x], x]`

output `(-3*(-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])/Sqrt[b]) + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/4 + (Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2))/2`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*(b*u - a*v)/(a*(m + n + 1)) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{2} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \dots\right)}{2}\right)}{2}\right)}{2}\right)}{2}$
default	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{2} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} + \dots\right)}{2}\right)}{2}\right)}{2}$

input `int(arctanh(tanh(b*x+a))^(3/2)/x^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/2*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.15

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}} dx = \left[ \frac{3a^2\sqrt{b} \log\left(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a\right) + 2(2b^2x + 5ab)\sqrt{bx+a}\sqrt{x}}{8b} - \frac{3a^2\sqrt{-b} \operatorname{arctan}\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) - (2b^2x + 5ab)\sqrt{bx+a}\sqrt{x}}{4b} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(1/2),x, algorithm="fricas")`

output `[1/8*(3*a^2*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(2*b^2*x + 5*a*b)*sqrt(b*x + a)*sqrt(x))/b, -1/4*(3*a^2*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (2*b^2*x + 5*a*b)*sqrt(b*x + a)*sqrt(x))/b]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**(1/2),x)`

output `Integral(atanh(tanh(a + b*x))**(3/2)/sqrt(x), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}}{\sqrt{x}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(1/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/sqrt(x), x)`

**Giac [A] (verification not implemented)**

Time = 72.36 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.96

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \frac{\sqrt{2} \left( \frac{3\sqrt{2}a^2 \log\left(\left| \frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}} \right| \right) - \sqrt{(bx+a)b-ab}\sqrt{bx+a} \left( \frac{2\sqrt{2}(bx+a)}{b} + \frac{3\sqrt{2}a}{b} \right)}{8|b|} \right) b}{8|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(1/2),x, algorithm="giac")`

output

```
-1/8*sqrt(2)*(3*sqrt(2)*a^2*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)
)*b - a*b)))/sqrt(b) - sqrt((b*x + a)*b - a*b)*sqrt(b*x + a)*(2*sqrt(2)*(b
*x + a)/b + 3*sqrt(2)*a/b))*b/abs(b)
```

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx$$

input

```
int(atanh(tanh(a + b*x))^(3/2)/x^(1/2), x)
```

output

```
int(atanh(tanh(a + b*x))^(3/2)/x^(1/2), x)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x} dx$$

input

```
int(atanh(tanh(b*x+a))^(3/2)/x^(1/2), x)
```

output

```
int((sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)))/x,x)
```

### 3.226 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}} dx$

Optimal result	1603
Mathematica [A] (verified)	1603
Rubi [A] (verified)	1604
Maple [A] (verified)	1606
Fricas [A] (verification not implemented)	1606
Sympy [F]	1607
Maxima [F]	1607
Giac [A] (verification not implemented)	1607
Mupad [F(-1)]	1608
Reduce [F]	1608

#### Optimal result

Integrand size = 17, antiderivative size = 81

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}} dx =$$

$$-3\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))$$

$$+ 3b\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{2\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}}$$

output

```
-3*b^(1/2)*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))+3*b*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)-2*arctanh(tanh(b*x+a))^(3/2)/x^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.95

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{3/2}} dx = \frac{(3bx - 2\operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}}$$

$$+ 3\sqrt{b}(-bx + \operatorname{arctanh}(\tanh(a+bx)))\log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)$$



input `Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(3/2),x]`

output `((3*b*x - 2*ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]])/Sqrt[x] + 3*Sqrt[b]*(-(b*x) + ArcTanh[Tanh[a + b*x]])*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]`

### Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2599, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx$$

$$\downarrow \text{2599}$$

$$3b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}}$$

$$\downarrow \text{2600}$$

$$3b \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}}$$

$$\downarrow \text{2596}$$

$$3b \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{\sqrt{b}} - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(3/2),x]`

output `3*b*(-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/Sqrt[b]) + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]] - (2*ArcTanh[Tanh[a + b*x]]^(3/2))/Sqrt[x]`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

### Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.56

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{8b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{8b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx} \right)}{\operatorname{arctanh}(\tanh(bx+a))-bx}$

input `int(arctanh(tanh(b*x+a))^(3/2)/x^(3/2), x, method=_RETURNVERBOSE)`

output `-2/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(tanh(b*x+a))^(5/2)+8*b/(arctanh(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.31

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \left[ \frac{3 a \sqrt{bx} \log \left( 2 bx + 2 \sqrt{bx + a} \sqrt{b} \sqrt{x} + a \right) + 2 \sqrt{bx + a} (bx - 2 a) \sqrt{x}}{2 x}, \dots \right]$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(3/2), x, algorithm="fricas")`

output `[1/2*(3*a*sqrt(b)*x*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*sqrt(b*x + a)*(b*x - 2*a)*sqrt(x))/x, -(3*a*sqrt(-b)*x*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - sqrt(b*x + a)*(b*x - 2*a)*sqrt(x))/x]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \int \frac{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))}{x^{\frac{3}{2}}} dx$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**(3/2), x)`

output `Integral(atanh(tanh(a + b*x))**(3/2)/x**(3/2), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}}{x^{\frac{3}{2}}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(3/2), x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/x^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 72.70 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \frac{\sqrt{2} \left( \frac{3\sqrt{2}a \log\left(\left| \frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}} \right| \right) - \frac{\sqrt{bx+a}(\sqrt{2}(bx+a) - 3\sqrt{2}a)}{\sqrt{(bx+a)b-ab}}}{2|b|} \right) b^2}{2|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(3/2), x, algorithm="giac")`

output `-1/2*sqrt(2)*(3*sqrt(2)*a*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*  
b - a*b)))/sqrt(b) - sqrt(b*x + a)*(sqrt(2)*(b*x + a) - 3*sqrt(2)*a)/sqrt(  
(b*x + a)*b - a*b))*b^2/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(3/2), x)`output `int(atanh(tanh(a + b*x))^(3/2)/x^(3/2), x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx = \frac{-2\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) + 3 \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x} dx \right)}{x}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^(3/2), x)`output `( - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) + 3*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/x,x)*b*x)/x`

### 3.227 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{5/2}} dx$

Optimal result	1609
Mathematica [A] (verified)	1609
Rubi [A] (verified)	1610
Maple [B] (verified)	1611
Fricas [A] (verification not implemented)	1612
Sympy [F]	1612
Maxima [F]	1613
Giac [A] (verification not implemented)	1613
Mupad [F(-1)]	1613
Reduce [F]	1614

#### Optimal result

Integrand size = 17, antiderivative size = 70

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = 2b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) - \frac{2b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}}$$

output

$2*b^{(3/2)}*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)})-2*b*\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}/x^{(1/2)}-2/3*\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}/x^{(3/2)}$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \frac{2\left(3bx\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \operatorname{arctanh}(\tanh(a + bx))^{3/2} - 3b^{3/2}x^{3/2} \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)\right)}{3x^{3/2}}$$

input

`Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(5/2), x]`

output

```
(-2*(3*b*x*Sqrt[ArcTanh[Tanh[a + b*x]]] + ArcTanh[Tanh[a + b*x]]^(3/2) - 3
*b^(3/2)*x^(3/2)*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(
3*x^(3/2))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2599, 2599, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx$$

$$\downarrow 2599$$

$$b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}}$$

$$\downarrow 2599$$

$$b \left( b \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}}$$

$$\downarrow 2596$$

$$b \left( 2\sqrt{b} \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \right) - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(5/2), x]
```

output

```
b*(2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]] - (2*
Sqrt[ArcTanh[Tanh[a + b*x]]])/Sqrt[x] - (2*ArcTanh[Tanh[a + b*x]]^(3/2))/
(3*x^(3/2))
```

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. 2(52) = 104.

Time = 0.40 (sec) , antiderivative size = 172, normalized size of antiderivative = 2.46

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a)))^{\frac{3}{2}}}{4} \right)}{3}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a)))^{\frac{3}{2}}}{4} \right)}{3}$

```
input int(arctanh(tanh(b*x+a))^(3/2)/x^(5/2), x, method=_RETURNVERBOSE)
```



output

```
-2/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh(tanh(b*x+a))^(5/2)+4/3*b/(
arctanh(tanh(b*x+a))-b*x)*(-1/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(t
anh(b*x+a))^(5/2)+4*b/(arctanh(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh
(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*
x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arct
anh(tanh(b*x+a))^(1/2))))))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.51

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \left[ \frac{3 b^{3/2} x^2 \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) - 2(4bx + a)\sqrt{bx+a}\sqrt{x}}{3x^2}, - \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^(5/2),x, algorithm="fricas")
```

output

```
[1/3*(3*b^(3/2)*x^2*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) - 2*(
4*b*x + a)*sqrt(b*x + a)*sqrt(x))/x^2, -2/3*(3*sqrt(-b)*b*x^2*arctan(sqrt(
-b)*sqrt(x)/sqrt(b*x + a)) + (4*b*x + a)*sqrt(b*x + a)*sqrt(x))/x^2]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \int \frac{\operatorname{atanh}^{3/2}(\tanh(a + bx))}{x^{5/2}} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(3/2)/x**(5/2),x)
```

output

```
Integral(atanh(tanh(a + b*x))**(3/2)/x**(5/2), x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{3/2}}{x^{5/2}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(5/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(3/2)/x^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 73.16 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \frac{\sqrt{2}b^3 \left( \frac{3\sqrt{2} \log\left(\left| \frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}} \right| \right)}{\sqrt{b}} + \frac{(4\sqrt{2}(bx+a)b - 3\sqrt{2}ab)\sqrt{bx+a}}{((bx+a)b-ab)^{3/2}} \right)}{3|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(5/2),x, algorithm="giac")`

output `-1/3*sqrt(2)*b^3*(3*sqrt(2)*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*b - a*b)))/sqrt(b) + (4*sqrt(2)*(b*x + a)*b - 3*sqrt(2)*a*b)*sqrt(b*x + a)/((b*x + a)*b - a*b)^(3/2))/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(5/2),x)`

output `int(atanh(tanh(a + b*x))^(3/2)/x^(5/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 6\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{3\sqrt{x} x}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^(5/2), x)`

output `( - 2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 6*sqrt(atanh(tanh(a + b*x)))*b*x + 3*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x), x)*b**2*x)/(3*sqrt(x)*x)`

$$3.228 \quad \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx$$

Optimal result	1615
Mathematica [A] (verified)	1615
Rubi [A] (verified)	1616
Maple [A] (verified)	1616
Fricas [A] (verification not implemented)	1617
Sympy [F(-1)]	1617
Maxima [A] (verification not implemented)	1618
Giac [A] (verification not implemented)	1618
Mupad [B] (verification not implemented)	1618
Reduce [F]	1619

### Optimal result

Integrand size = 17, antiderivative size = 35

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
2/5*arctanh(tanh(b*x+a))^(5/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))
```

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}(5bx - 5\operatorname{arctanh}(\tanh(a+bx)))}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(7/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(x^(5/2)*(5*b*x - 5*ArcTanh[Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{7/2}} dx$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(7/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}$	29
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}$	29

input `int(arctanh(tanh(b*x+a))^(3/2)/x^(7/2),x,method=_RETURNVERBOSE)`

output `-2/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(5/2)`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx = -\frac{2(b^2x^2 + 2abx + a^2)\sqrt{bx+a}}{5ax^{\frac{5}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(7/2),x, algorithm="fricas")`

output `-2/5*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(b*x + a)/(a*x^(5/2))`

### Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**(7/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{7/2}} dx = -\frac{2(bx + a)^{\frac{5}{2}}}{5ax^{\frac{5}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(7/2),x, algorithm="maxima")`output `-2/5*(b*x + a)^(5/2)/(a*x^(5/2))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{7/2}} dx = -\frac{2(bx + a)^{\frac{5}{2}}b^6}{5((bx + a)b - ab)^{\frac{5}{2}}a|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(7/2),x, algorithm="giac")`output `-2/5*(b*x + a)^(5/2)*b^6/(((b*x + a)*b - a*b)^(5/2)*a*abs(b))`**Mupad [B] (verification not implemented)**

Time = 3.80 (sec) , antiderivative size = 332, normalized size of antiderivative = 9.49

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{7/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{5 \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - 5 \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 10bx} + \frac{2bx}{5} \right)}{x^{7/2}}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(7/2),x)`

output

```
(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(5*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 5*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 10*b*x) + (4*b^2*x^2)/(5*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 5*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 10*b*x) - (4*b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(5*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 5*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 10*b*x))/x^(5/2)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{7/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{5\sqrt{x} x^2}$$

input

```
int(atanh(tanh(b*x+a))^(3/2)/x^(7/2),x)
```

output

```
( - 2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 2*sqrt(atanh(tanh(a + b*x)))*b*x + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(atanh(a + b*x))*x**2),x)*b**2*x**2)/(5*sqrt(x)*x**2)
```



### 3.229 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx$

Optimal result	1620
Mathematica [A] (verified)	1620
Rubi [A] (verified)	1621
Maple [A] (verified)	1622
Fricas [A] (verification not implemented)	1622
Sympy [F(-1)]	1623
Maxima [A] (verification not implemented)	1623
Giac [A] (verification not implemented)	1623
Mupad [B] (verification not implemented)	1624
Reduce [F]	1624

#### Optimal result

Integrand size = 17, antiderivative size = 72

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx = \frac{4\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `4/35*b*arctanh(tanh(b*x+a))^(5/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/7*arctanh(tanh(b*x+a))^(5/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx = \frac{2(7bx - 5\operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{35x^{7/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(9/2),x]`

output `(2*(7*b*x - 5*ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(5/2))/(35*x^(7/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx$$

↓ 2602

$$\frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx}{7(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(9/2), x]`

output `(4*b*ArcTanh[Tanh[a + b*x]]^(5/2))/(35*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(5/2))/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{35(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{5}{2}}}$	59
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{35(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{5}{2}}}$	59

input

```
int(arctanh(tanh(b*x+a))^(3/2)/x^(9/2), x, method=_RETURNVERBOSE)
```

output

```
-2/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(5/2)+4/35*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(5/2)*arctanh(tanh(b*x+a))^(5/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \frac{2(2b^3x^3 - ab^2x^2 - 8a^2bx - 5a^3)\sqrt{bx + a}}{35a^2x^{\frac{7}{2}}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^(9/2), x, algorithm="fricas")
```

output

```
2/35*(2*b^3*x^3 - a*b^2*x^2 - 8*a^2*b*x - 5*a^3)*sqrt(b*x + a)/(a^2*x^(7/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**(9/2), x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.47

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \frac{2(2b^2x^2 - 3abx - 5a^2)(bx + a)^{\frac{3}{2}}}{35a^2x^{\frac{7}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(9/2), x, algorithm="maxima")`

output `2/35*(2*b^2*x^2 - 3*a*b*x - 5*a^2)*(b*x + a)^(3/2)/(a^2*x^(7/2))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \frac{\sqrt{2} \left( \frac{2\sqrt{2}(bx+a)b^3}{a^2} - \frac{7\sqrt{2}b^3}{a} \right) (bx + a)^{\frac{5}{2}} b^5}{35((bx + a)b - ab)^{\frac{7}{2}} |b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(9/2), x, algorithm="giac")`

output `1/35*sqrt(2)*(2*sqrt(2)*(b*x + a)*b^3/a^2 - 7*sqrt(2)*b^3/a)*(b*x + a)^(5/2)*b^5/(((b*x + a)*b - a*b)^(7/2)*abs(b))`

**Mupad [B] (verification not implemented)**

Time = 3.77 (sec) , antiderivative size = 228, normalized size of antiderivative = 3.17

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{7} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{7} - \frac{6bx}{35} + \frac{4b^2x^2}{35(\ln(2/(e^{2a}e^{2bx}+1)) - \log((2e^{2a}e^{2bx})/(e^{2a}e^{2bx}+1)) + 2bx))} + \frac{16b^3x^3}{35(\ln(2/(e^{2a}e^{2bx}+1)) - \log((2e^{2a}e^{2bx})/(e^{2a}e^{2bx}+1)) + 2bx)^2)} \right)}{x^{7/2}}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(9/2), x)`output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/7 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/7 - (6*b*x)/35 + (4*b^2*x^2)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (16*b^3*x^3)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(7/2)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx = \frac{-10\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 6\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{35\sqrt{x}x^3}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^(9/2), x)`output `( - 10*sqrt(atanh(tanh(a + b*x))) * atanh(tanh(a + b*x)) - 6*sqrt(atanh(tanh(a + b*x))) * b*x + 3*sqrt(x) * int((sqrt(x) * sqrt(atanh(tanh(a + b*x)))) / (atanh(tanh(a + b*x)) * x**3), x) * b**2 * x**3) / (35*sqrt(x) * x**3)`

### 3.230 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx$

Optimal result	1625
Mathematica [A] (verified)	1625
Rubi [A] (verified)	1626
Maple [A] (verified)	1627
Fricas [A] (verification not implemented)	1628
Sympy [F(-1)]	1628
Maxima [A] (verification not implemented)	1629
Giac [A] (verification not implemented)	1629
Mupad [B] (verification not implemented)	1630
Reduce [F]	1630

#### Optimal result

Integrand size = 17, antiderivative size = 110

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx = \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{315x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{8b \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{63x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
16/315*b^2*arctanh(tanh(b*x+a))^(5/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^3
+8/63*b*arctanh(tanh(b*x+a))^(5/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/
9*arctanh(tanh(b*x+a))^(5/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx = \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2} (63b^2x^2 - 90bx \operatorname{arctanh}(\tanh(a+bx)) + 35a)}{315x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(11/2), x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(5/2)*(63*b^2*x^2 - 90*b*x*ArcTanh[Tanh[a + b*x]] + 35*ArcTanh[Tanh[a + b*x]]^2))/(315*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{11/2}} dx$$

↓ 2602

$$\frac{4b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{9/2}} dx}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2602

$$\frac{4b \left( \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + 4b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) / 9(bx - \operatorname{arctanh}(\tanh(a+bx)))$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(11/2), x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]]))
+ (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(5/2))/(35*x^(5/2)*(b*x - ArcTanh[Ta
nh[a + b*x]]^2) + (2*ArcTanh[Tanh[a + b*x]]^(5/2))/(7*x^(7/2)*(b*x - ArcT
anh[Tanh[a + b*x]])))))/(9*(b*x - ArcTanh[Tanh[a + b*x]]))
```

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /;
NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[
m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + S
imp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; Ne
Q[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m
, -1]
```

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}}$	105
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}}$	105

input

```
int(arctanh(tanh(b*x+a))^(3/2)/x^(11/2), x, method=_RETURNVERBOSE)
```



output

```
-2/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arctanh(tanh(b*x+a))^(5/2)-8/9*b/(
arctanh(tanh(b*x+a))-b*x)*(-1/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh
(tanh(b*x+a))^(5/2)+2/35*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(5/2)*arctanh(ta
nh(b*x+a))^(5/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.51

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = \frac{2(8b^4x^4 - 4ab^3x^3 + 3a^2b^2x^2 + 50a^3bx + 35a^4)\sqrt{bx + a}}{315a^3x^{9/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^(11/2),x, algorithm="fricas")
```

output

```
-2/315*(8*b^4*x^4 - 4*a*b^3*x^3 + 3*a^2*b^2*x^2 + 50*a^3*b*x + 35*a^4)*sqrt
t(b*x + a)/(a^3*x^(9/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = \text{Timed out}$$

input

```
integrate(atanh(tanh(b*x+a))**(3/2)/x**(11/2),x)
```

output

```
Timed out
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = -\frac{2(8b^3x^3 - 12ab^2x^2 + 15a^2bx + 35a^3)(bx + a)^{3/2}}{315a^3x^{9/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(11/2),x, algorithm="maxima")`output `-2/315*(8*b^3*x^3 - 12*a*b^2*x^2 + 15*a^2*b*x + 35*a^3)*(b*x + a)^(3/2)/(a^3*x^(9/2))`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = -\frac{\sqrt{2}\left(\frac{63\sqrt{2}b^9}{a} + 4\left(\frac{2\sqrt{2}(bx+a)b^9}{a^3} - \frac{9\sqrt{2}b^9}{a^2}\right)(bx + a)\right)(bx + a)^{5/2}b}{315((bx + a)b - ab)^{9/2}|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(11/2),x, algorithm="giac")`output `-1/315*sqrt(2)*(63*sqrt(2)*b^9/a + 4*(2*sqrt(2)*(b*x + a)*b^9/a^3 - 9*sqrt(2)*b^9/a^2)*(b*x + a))*(b*x + a)^(5/2)*b/(((b*x + a)*b - a*b)^(9/2)*abs(b))`

**Mupad [B] (verification not implemented)**

Time = 3.94 (sec) , antiderivative size = 288, normalized size of antiderivative = 2.62

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{9} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{9} - \frac{2bx}{21} + \frac{1}{1} \right)}{1}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(11/2), x)`output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)))/9 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/9 - (2*b*x)/21 + (4*b^2*x^2)/(105*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (32*b^3*x^3)/(315*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (128*b^4*x^4)/(315*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3))/x^(9/2)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx = \frac{-14\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 6\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{63\sqrt{x} x^4}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^(11/2), x)`output

```
( - 14*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 6*sqrt(atanh(tanh(a + b*x)))*b*x + 3*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**4), x)*b**2*x**4)/(63*sqrt(x)*x**4)
```

### 3.231 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{13/2}} dx$

Optimal result	1631
Mathematica [A] (verified)	1632
Rubi [A] (verified)	1632
Maple [A] (verified)	1634
Fricas [A] (verification not implemented)	1634
Sympy [F(-1)]	1635
Maxima [A] (verification not implemented)	1635
Giac [A] (verification not implemented)	1635
Mupad [B] (verification not implemented)	1636
Reduce [F]	1636

#### Optimal result

Integrand size = 17, antiderivative size = 148

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{13/2}} dx = \frac{32b^3 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{1155x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{231x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{4b \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{33x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
32/1155*b^3*arctanh(tanh(b*x+a))^(5/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^4+16/231*b^2*arctanh(tanh(b*x+a))^(5/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^3+4/33*b*arctanh(tanh(b*x+a))^(5/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/11*arctanh(tanh(b*x+a))^(5/2)/x^(11/2)/(b*x-arctanh(tanh(b*x+a)))
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2} (231b^3x^3 - 495b^2x^2\operatorname{arctanh}(\tanh(a + bx)) + 1155x^{11/2}(-bx + \operatorname{arctanh}(\tanh(a + bx))))}{1155x^{11/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(13/2), x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(5/2)*(231*b^3*x^3 - 495*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 385*b*x*ArcTanh[Tanh[a + b*x]]^2 - 105*ArcTanh[Tanh[a + b*x]]^3)/(1155*x^(11/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]]))^4)
```

**Rubi [A] (verified)**

Time = 0.61 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx \\ & \quad \downarrow 2602 \\ & \frac{6b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{11/2}} dx}{11(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2602 \\ & \frac{6b \left( \frac{4b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{9/2}} dx}{9(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{11(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \\ & \quad \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2602 \end{aligned}$$

$$\begin{aligned}
& 6b \left( \frac{4b \left( \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{x^{7/2}} dx}{7(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{11(bx - \operatorname{arctanh}(\tanh(a+bx)))}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow \text{2598} \\
& 6b \left( \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b \left( \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b \operatorname{arctanh}(\tanh(a+bx))^{5/2}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right)}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{11(bx - \operatorname{arctanh}(\tanh(a+bx)))}{11(bx - \operatorname{arctanh}(\tanh(a+bx)))}
\end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(3/2)/x^(13/2), x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(5/2))/(11*x^(11/2)*(b*x - ArcTanh[Tanh[a + b*x]]) + (6*b*((2*ArcTanh[Tanh[a + b*x]]^(5/2))/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) + (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(5/2))/(35*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(5/2))/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])))))/(9*(b*x - ArcTanh[Tanh[a + b*x]])))/(11*(b*x - ArcTanh[Tanh[a + b*x]]))`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

### Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} - \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} + \dots \right)}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} - \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}} + \dots \right)}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)}$

input

```
int(arctanh(tanh(b*x+a))^(3/2)/x^(13/2), x, method=_RETURNVERBOSE)
```

output

```
-2/11/(arctanh(tanh(b*x+a))-b*x)/x^(11/2)*arctanh(tanh(b*x+a))^(5/2)-12/11*b/(arctanh(tanh(b*x+a))-b*x)*(-1/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arctanh(tanh(b*x+a))^(5/2)-4/9*b/(arctanh(tanh(b*x+a))-b*x)*(-1/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(5/2)+2/35*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(5/2)*arctanh(tanh(b*x+a))^(5/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.45

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{2(16b^5x^5 - 8ab^4x^4 + 6a^2b^3x^3 - 5a^3b^2x^2 - 140a^4bx - 105a^5)\sqrt{bx+a}}{1155a^4x^{\frac{11}{2}}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(3/2)/x^(13/2), x, algorithm="fricas")
```

output  $2/1155*(16*b^5*x^5 - 8*a*b^4*x^4 + 6*a^2*b^3*x^3 - 5*a^3*b^2*x^2 - 140*a^4*b*x - 105*a^5)*\text{sqrt}(b*x + a)/(a^4*x^{(11/2)})$

### Sympy [F(-1)]

Timed out.

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(3/2)/x**(13/2),x)`

output Timed out

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.38

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{2(16b^4x^4 - 24ab^3x^3 + 30a^2b^2x^2 - 35a^3bx - 105a^4)(bx + a)^{\frac{3}{2}}}{1155a^4x^{\frac{11}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(13/2),x, algorithm="maxima")`

output  $2/1155*(16*b^4*x^4 - 24*a*b^3*x^3 + 30*a^2*b^2*x^2 - 35*a^3*b*x - 105*a^4)*(b*x + a)^{(3/2)}/(a^4*x^{(11/2)})$

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.67

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{\sqrt{2} \left( \frac{231\sqrt{2}b^5}{a} - 2 \left( \frac{99\sqrt{2}b^5}{a^2} + 4 \left( \frac{2\sqrt{2}(bx+a)b^5}{a^4} - \frac{11\sqrt{2}b^5}{a^3} \right) (bx + a) \right) (bx + a) \right) (bx + a)^{\frac{5}{2}} b^7}{1155 ((bx + a)b - ab)^{\frac{11}{2}} |b|}$$



input `integrate(arctanh(tanh(b*x+a))^(3/2)/x^(13/2),x, algorithm="giac")`

output 
$$-1/1155*\sqrt{2}*(231*\sqrt{2}*b^5/a - 2*(99*\sqrt{2})*b^5/a^2 + 4*(2*\sqrt{2})*(b*x + a)*b^5/a^4 - 11*\sqrt{2}*b^5/a^3)*(b*x + a)*(b*x + a)*(b*x + a)^{(5/2)}*b^7/(((b*x + a)*b - a*b)^{(11/2)}*abs(b))$$

### Mupad [B] (verification not implemented)

Time = 4.16 (sec) , antiderivative size = 348, normalized size of antiderivative = 2.35

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}{\frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{11} - \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{11} - \frac{2bx}{33} + \frac{1}{2}}}{x^{13/2}}$$

input `int(atanh(tanh(a + b*x))^(3/2)/x^(13/2),x)`

output 
$$\left(\frac{\log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right)}{2} - \log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right)\right)^{1/2} * \left(\frac{\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right)}{11} - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right)\right) / 11 - \frac{2*b*x}{33} + \frac{4*b^2*x^2}{231*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)} + \frac{16*b^3*x^3}{385*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)^2} + \frac{128*b^4*x^4}{1155*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)^3} + \frac{512*b^5*x^5}{1155*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)^4} \right) / x^{11/2}$$

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{13/2}} dx = \frac{-6\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a)) - 2\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{33\sqrt{x}x^5}$$

input `int(atanh(tanh(b*x+a))^(3/2)/x^(13/2),x)`

output

```
( - 6*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 2*sqrt(atanh(tanh(a + b*x)))*b*x + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**5),x)*b**2*x**5)/(33*sqrt(x)*x**5)
```

### 3.232 $\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$

Optimal result	1638
Mathematica [A] (verified)	1639
Rubi [A] (verified)	1639
Maple [A] (verified)	1641
Fricas [A] (verification not implemented)	1642
Sympy [F(-1)]	1642
Maxima [F]	1643
Giac [A] (verification not implemented)	1643
Mupad [F(-1)]	1644
Reduce [F]	1644

#### Optimal result

Integrand size = 17, antiderivative size = 174

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx =$$

$$-\frac{5 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))^4}{64b^{3/2}} + \frac{5}{32} x^{3/2} (bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{5\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{64b}$$

output

```
-5/64*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^4/b^(3/2)+5/32*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)-5/64*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^3*arctanh(tanh(b*x+a))^(1/2)/b-5/24*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(3/2)+1/4*x^(3/2)*arctanh(tanh(b*x+a))^(5/2)
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.70

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (15b^3 x^3 - 55b^2 x^2 \operatorname{arctanh}(\tanh(a + bx)) + 73bx \operatorname{arctanh}(\tanh(a + bx)))^{5/2}}{192b} - \frac{5(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4 \log\left(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)}{64b^{3/2}}$$

input `Integrate[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^3*x^3 - 55*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 73*b*x*ArcTanh[Tanh[a + b*x]]^2 + 15*ArcTanh[Tanh[a + b*x]]^3))/(192*b) - (5*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(64*b^(3/2))`

**Rubi [A] (verified)**

Time = 0.62 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2600, 2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx$$

$$\downarrow 2600$$

$$\frac{1}{4} x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2} - \frac{5}{8} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} dx$$

$$\downarrow 2600$$

$$\begin{aligned}
 & \frac{1}{4}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{8}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}dx \right) \\
 & \quad \downarrow 2600 \\
 & \frac{1}{4}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{8}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{2}x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{4} \right) \right) \\
 & \quad \downarrow 2600 \\
 & \frac{1}{4}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{8}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{2}x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{4} \right) \right) \\
 & \quad \downarrow 2596 \\
 & \frac{1}{4}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{8}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{1}{3}x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{1}{2} \left( \frac{1}{2}x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{4} \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) \right) \right)
 \end{aligned}$$

input `Int[Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(x^(3/2)*ArcTanh[Tanh[a + b*x]]^(5/2))/4 - (5*(b*x - ArcTanh[Tanh[a + b*x]])*(-1/2*((-1/4*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b*(b*x - ArcTanh[Tanh[a + b*x]]) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/2)*(b*x - ArcTanh[Tanh[a + b*x]]) + (x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2))/3))/8`

Defintions of rubi rules used

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Si
mplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v
]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u -
a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /;
PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n +
1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n,
-2]
```

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{4b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$
default	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{4b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$

```
input int(x^(1/2)*arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
1/4*x^(1/2)*arctanh(tanh(b*x+a))^(7/2)/b-1/4*(arctanh(tanh(b*x+a))-b*x)/b*
(1/6*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)+5/6*(arctanh(tanh(b*x+a))-b*x)*(1/
4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x
^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*l
n(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.91

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{15 a^4 \sqrt{b} \log(2bx - 2\sqrt{bx+a}\sqrt{b}\sqrt{x+a}) + 2(48b^4x^3 + 136ab^3x^2 + 118a^2b^2x + 15a^3b)\sqrt{bx+a}\sqrt{x}}{384b^2}$$

input

```
integrate(x^(1/2)*arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
[1/384*(15*a^4*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) +
2*(48*b^4*x^3 + 136*a*b^3*x^2 + 118*a^2*b^2*x + 15*a^3*b)*sqrt(b*x + a)*sq
rt(x))/b^2, 1/192*(15*a^4*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a))
+ (48*b^4*x^3 + 136*a*b^3*x^2 + 118*a^2*b^2*x + 15*a^3*b)*sqrt(b*x + a)*sq
rt(x))/b^2]
```

**Sympy [F(-1)]**

Timed out.

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \text{Timed out}$$

input

```
integrate(x**(1/2)*atanh(tanh(b*x+a))**(5/2),x)
```

output

Timed out

**Maxima [F]**

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \int \sqrt{x} \operatorname{artanh}(\tanh(bx + a))^{5/2} dx$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(x)*arctanh(tanh(b*x + a))^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.17

$$\int \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2} dx = \frac{1}{384} \sqrt{2} \left( 48 \sqrt{2} \left( \sqrt{bx + a} \left( 2x + \frac{a}{b} \right) \sqrt{x} + \frac{a^2 \log \left( \left| -\sqrt{b} \sqrt{x} + \sqrt{bx + a} \right| \right)}{b^{3/2}} \right) a^2 + 16 \sqrt{2} \left( \sqrt{bx + a} \right) \right)$$

input `integrate(x^(1/2)*arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/384*sqrt(2)*(48*sqrt(2)*(sqrt(b*x + a)*(2*x + a/b)*sqrt(x) + a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(3/2)))*a^2 + 16*sqrt(2)*(sqrt(b*x + a)*(2*(4*x + a/b)*x - 3*a^2/b^2)*sqrt(x) - 3*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2))*a*b + sqrt(2)*((2*(4*(6*x + a/b)*x - 5*a^2/b^2)*x + 15*a^3/b^3)*sqrt(b*x + a)*sqrt(x) + 15*a^4*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(7/2))*b^2)`



**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{x} \arctanh(\tanh(a + bx))^{5/2} dx = \int \sqrt{x} \operatorname{atanh}(\tanh(a + bx))^{5/2} dx$$

input `int(x^(1/2)*atanh(tanh(a + b*x))^(5/2), x)`output `int(x^(1/2)*atanh(tanh(a + b*x))^(5/2), x)`**Reduce [F]**

$$\int \sqrt{x} \arctanh(\tanh(a + bx))^{5/2} dx = \int \sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 dx$$

input `int(x^(1/2)*atanh(tanh(b*x+a))^(5/2), x)`output `int(sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2, x)`

### 3.233 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx$

Optimal result	1645
Mathematica [A] (verified)	1646
Rubi [A] (verified)	1646
Maple [A] (verified)	1648
Fricas [A] (verification not implemented)	1649
Sympy [F]	1649
Maxima [F]	1650
Giac [A] (verification not implemented)	1650
Mupad [F(-1)]	1650
Reduce [F]	1651

#### Optimal result

Integrand size = 17, antiderivative size = 136

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx =$$

$$\frac{5\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{8\sqrt{b}}$$

$$+ \frac{5}{8}\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}$$

$$- \frac{5}{12}\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^{3/2}$$

$$+ \frac{1}{3}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{5/2}$$

output

```
-5/8*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^3/b^(1/2)+5/8*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)-5/12*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(3/2)+1/3*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.74

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx = \frac{1}{24} \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} (15b^2x^2 - 40bx \operatorname{arctanh}(\tanh(a+bx)) + 33 \operatorname{arctanh}(\tanh(a+bx))^2) + \frac{5(-bx + \operatorname{arctanh}(\tanh(a+bx)))^3 \log(b\sqrt{x} + \sqrt{b} \sqrt{\operatorname{arctanh}(\tanh(a+bx))})}{8\sqrt{b}}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/Sqrt[x], x]
```

output

```
(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^2*x^2 - 40*b*x*ArcTanh[Tanh[a + b*x]] + 33*ArcTanh[Tanh[a + b*x]]^2))/24 + (5*(-(b*x) + ArcTanh[Tanh[a + b*x]])^3*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(8*Sqrt[b])
```

**Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{\sqrt{x}} dx$$

↓ 2600

$$\frac{1}{3} \sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{6} (bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{3/2}}{\sqrt{x}} dx$$

↓ 2600

$$\begin{aligned}
& \frac{1}{3}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \\
& \left( \frac{1}{2}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{3}{4}(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}} dx \right) \\
& \quad \downarrow \text{2600} \\
& \frac{1}{3}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \\
& \left( \frac{1}{2}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{3}{4}(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{1}{2}(bx - \right. \right. \\
& \quad \downarrow \text{2596} \\
& \left. \left. \frac{1}{3}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{5/2} - \frac{5}{6}(bx - \operatorname{arctanh}(\tanh(a+bx))) \right. \right. \\
& \left. \left. \left( \frac{1}{2}\sqrt{x}\operatorname{arctanh}(\tanh(a+bx))^{3/2} - \frac{3}{4} \left( \sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{b}} \right) (bx - \right. \right. \right.
\end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/Sqrt[x], x]`

output `(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(5/2))/3 - (5*(b*x - ArcTanh[Tanh[a + b*x]])*((-3*(-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/Sqrt[b]) + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/4 + (Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2))/2)/6`

## Definitions of rubi rules used

rule 2596

```
Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

## Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$
default	$\frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{3} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}{4} + \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx)}{\dots} \right)}{\dots}$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(1/2), x, method=_RETURNVERBOSE)
```

output

```
1/3*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)+5/3*(arctanh(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.01

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \left[ \frac{15 a^3 \sqrt{b} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2(8b^3x^2 + 26ab^2x + 33a^2b)\sqrt{bx+a}\sqrt{x}}{48b} - \frac{15 a^3 \sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) - (8b^3x^2 + 26ab^2x + 33a^2b)\sqrt{bx+a}\sqrt{x}}{24b} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(1/2),x, algorithm="fricas")`

output `[1/48*(15*a^3*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(8*b^3*x^2 + 26*a*b^2*x + 33*a^2*b)*sqrt(b*x + a)*sqrt(x))/b, -1/24*(15*a^3*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (8*b^3*x^2 + 26*a*b^2*x + 33*a^2*b)*sqrt(b*x + a)*sqrt(x))/b]`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}^{5/2}(\tanh(a + bx))}{\sqrt{x}} dx$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**(1/2),x)`

output `Integral(atanh(tanh(a + b*x))**(5/2)/sqrt(x), x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{\sqrt{x}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(1/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/sqrt(x), x)`

**Giac [A] (verification not implemented)**

Time = 73.07 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \frac{\sqrt{2} \left( \frac{15\sqrt{2}a^3 \log\left(\frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}}\right) - \sqrt{(bx+a)b-ab}\sqrt{bx+a} \left( 2(bx+a) \left( \frac{4\sqrt{2}(bx+a)}{b} + \frac{5\sqrt{2}a}{b} \right) + \frac{15\sqrt{2}a}{b} \right)}{48|b|} \right)}{48|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(1/2),x, algorithm="giac")`

output `-1/48*sqrt(2)*(15*sqrt(2)*a^3*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*b - a*b)))/sqrt(b) - sqrt((b*x + a)*b - a*b)*sqrt(b*x + a)*(2*(b*x + a)*(4*sqrt(2)*(b*x + a)/b + 5*sqrt(2)*a/b) + 15*sqrt(2)*a^2/b))*b/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(1/2),x)`

output `int(atanh(tanh(a + b*x))^(5/2)/x^(1/2), x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{x} dx$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(1/2), x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*atanh(tanh(a + b*x))**2)/x,x)`



### 3.234 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{3/2}} dx$

Optimal result	1652
Mathematica [A] (verified)	1653
Rubi [A] (verified)	1653
Maple [A] (verified)	1655
Fricas [A] (verification not implemented)	1656
Sympy [F]	1656
Maxima [F]	1657
Giac [A] (verification not implemented)	1657
Mupad [F(-1)]	1657
Reduce [F]	1658

#### Optimal result

Integrand size = 17, antiderivative size = 121

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \frac{15}{4} \sqrt{b} \operatorname{arctanh} \left( \frac{\sqrt{b} \sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx))) - \frac{15}{4} b \sqrt{x} (bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))} + \frac{5}{2} b \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}}$$

output

```
15/4*b^(1/2)*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^2-15/4*b*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)+5/2*b*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)-2*arctanh(tanh(b*x+a))^(5/2)/x^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx =$$

$$\frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))(15b^2x^2 - 25bx\operatorname{arctanh}(\tanh(a + bx)) + 8\operatorname{arctanh}(\tanh(a + bx))^2)}}{4\sqrt{x}}$$

$$+ \frac{15}{4}\sqrt{b}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(3/2), x]
```

output

```
-1/4*(Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^2*x^2 - 25*b*x*ArcTanh[Tanh[a + b*x]] + 8*ArcTanh[Tanh[a + b*x]]^2))/Sqrt[x] + (15*Sqrt[b]*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/4
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2599, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx$$

$$\downarrow \text{2599}$$

$$5b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}}$$

$$\downarrow \text{2600}$$

$$5b \left( \frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx \right) - \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}}$$

↓ 2600

$$5b \left( \frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{4} (bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2} (bx - \operatorname{arctanh}(\tanh(a + bx))) \right) \right) - \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}}$$

↓ 2596

$$5b \left( \frac{1}{2} \sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2} - \frac{3}{4} \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{\sqrt{b}} \right) \right) - \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{5/2}}{\sqrt{x}}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(3/2),x]`

output `(-2*ArcTanh[Tanh[a + b*x]]^(5/2))/Sqrt[x] + 5*b*((-3*(-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/Sqrt[b]) + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/4 + (Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2))/2`

**Defintions of rubi rules used**

rule 2596 `Int[1/(Sqrt[u]*Sqrt[v]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

### Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.28

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + 12b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))}{4} \right)}{\dots} \right)$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + 12b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))}{4} \right)}{\dots} \right)$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(3/2), x, method=_RETURNVERBOSE)
```

output

```
-2/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(tanh(b*x+a))^(7/2)+12*b/(arc
tanh(tanh(b*x+a))-b*x)*(1/6*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)+5/6*(arctan
h(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(t
anh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arct
anh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \left[ \frac{15 a^2 \sqrt{bx} \log(2bx + 2\sqrt{bx + a}\sqrt{bx} + a) + 2(2b^2x^2 + 9abx - 8a^2)}{8x} \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^(3/2),x, algorithm="fricas")
```

output

```
[1/8*(15*a^2*sqrt(b)*x*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) +
2*(2*b^2*x^2 + 9*a*b*x - 8*a^2)*sqrt(b*x + a)*sqrt(x))/x, -1/4*(15*a^2*sq
rt(-b)*x*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (2*b^2*x^2 + 9*a*b*x - 8*
a^2)*sqrt(b*x + a)*sqrt(x))/x]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \int \frac{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))}{x^{\frac{3}{2}}} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(5/2)/x**(3/2),x)
```

output

```
Integral(atanh(tanh(a + b*x))**(5/2)/x**(3/2), x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^{3/2}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(3/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 73.41 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \frac{\sqrt{2} \left( \frac{15\sqrt{2}a^2 \log\left(\left| \frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}} \right| \right) + \frac{(15\sqrt{2}a^2 - (bx+a)(2\sqrt{2}(bx+a) + 5\sqrt{2}a))\sqrt{bx+a}}{\sqrt{(bx+a)b-ab}} \right) b^2}{8|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(3/2),x, algorithm="giac")`

output `-1/8*sqrt(2)*(15*sqrt(2)*a^2*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*b - a*b)))/sqrt(b) + (15*sqrt(2)*a^2 - (b*x + a)*(2*sqrt(2)*(b*x + a) + 5*sqrt(2)*a))*sqrt(b*x + a)/sqrt((b*x + a)*b - a*b))*b^2/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(3/2),x)`

output `int(atanh(tanh(a + b*x))^(5/2)/x^(3/2), x)`

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2}{x^2} dx$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(3/2), x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*atanh(tanh(a + b*x))**2/x**2,x)`

### 3.235 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}} dx$

Optimal result	1659
Mathematica [A] (verified)	1659
Rubi [A] (verified)	1660
Maple [B] (verified)	1662
Fricas [A] (verification not implemented)	1663
Sympy [F]	1663
Maxima [F]	1664
Giac [A] (verification not implemented)	1664
Mupad [F(-1)]	1664
Reduce [F]	1665

#### Optimal result

Integrand size = 17, antiderivative size = 106

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}} dx = -5b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx))) + 5b^2 \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}$$

output

```
-5*b^(3/2)*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))+5*b^2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)-10/3*b*arctanh(tanh(b*x+a))^(3/2)/x^(1/2)-2/3*arctanh(tanh(b*x+a))^(5/2)/x^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{5/2}} dx = \frac{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}(15b^2x^2 - 10bx \operatorname{arctanh}(\tanh(a+bx)) - 2 \operatorname{arctanh}(\tanh(a+bx)))}{3x^{3/2}} + 5b^{3/2}(-bx + \operatorname{arctanh}(\tanh(a+bx))) \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(5/2),x]
```



output

```
(Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^2*x^2 - 10*b*x*ArcTanh[Tanh[a + b*x]]
- 2*ArcTanh[Tanh[a + b*x]]^2))/(3*x^(3/2)) + 5*b^(3/2)*(-(b*x) + ArcTanh[T
anh[a + b*x]])*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]]
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2599, 2599, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx$$

$$\downarrow 2599$$

$$\frac{5}{3}b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{3x^{3/2}}$$

$$\downarrow 2599$$

$$\frac{5}{3}b \left( 3b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{3x^{3/2}}$$

$$\downarrow 2600$$

$$\frac{5}{3}b \left( 3b \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} - \frac{1}{2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{3x^{3/2}} \right)$$

$$\downarrow 2596$$

$$\frac{5}{3}b \left( 3b \left( \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{\sqrt{b}} \right) - \frac{2\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{3x^{3/2}} \right)$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(5/2), x]`

output `(-2*ArcTanh[Tanh[a + b*x]]^(5/2))/(3*x^(3/2)) + (5*b*(3*b*(-((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])/Sqrt[b]) + Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]) - (2*ArcTanh[Tanh[a + b*x]]^(3/2))/Sqrt[x]))/3`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal.  $200$  vs.  $2(82) = 164$ .

Time =  $0.38$  (sec) , antiderivative size =  $201$ , normalized size of antiderivative =  $1.90$

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \left( 8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + 6b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a)))^{\frac{5}{2}}}{6} \right) \right) \right)$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \left( 8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + 6b \left( \frac{\sqrt{x} \operatorname{arctanh}(\tanh(bx+a))^{\frac{5}{2}}}{6} + \frac{5(\operatorname{arctanh}(\tanh(bx+a)))^{\frac{5}{2}}}{6} \right) \right) \right)$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(5/2), x, method=_RETURNVERBOSE)
```

output

```
-2/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh(tanh(b*x+a))^(7/2)+8/3*b/(
arctanh(tanh(b*x+a))-b*x)*(-1/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(t
anh(b*x+a))^(7/2)+6*b/(arctanh(tanh(b*x+a))-b*x)*(1/6*x^(1/2)*arctanh(tanh
(b*x+a))^(5/2)+5/6*(arctanh(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh(b*
x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a
))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh
(tanh(b*x+a))^(1/2))))))
```

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.27

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \left[ \frac{15 ab^{\frac{3}{2}} x^2 \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2(3b^2x^2 - 14abx - 2a^2)}{6x^2} \right]$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^(5/2),x, algorithm="fricas")
```

output

```
[1/6*(15*a*b^(3/2)*x^2*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) +
2*(3*b^2*x^2 - 14*a*b*x - 2*a^2)*sqrt(b*x + a)*sqrt(x))/x^2, -1/3*(15*a*sq
rt(-b)*b*x^2*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (3*b^2*x^2 - 14*a*b*
x - 2*a^2)*sqrt(b*x + a)*sqrt(x))/x^2]
```

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \int \frac{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))}{x^{\frac{5}{2}}} dx$$

input

```
integrate(atanh(tanh(b*x+a))**(5/2)/x**(5/2),x)
```

output

```
Integral(atanh(tanh(a + b*x))**(5/2)/x**(5/2), x)
```

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^{5/2}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(5/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 73.12 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \frac{\sqrt{2} \left( \frac{15\sqrt{2}a \log\left(\left| \frac{-\sqrt{bx+a}\sqrt{b} + \sqrt{(bx+a)b-ab}}{\sqrt{b}} \right| \right)}{\sqrt{b}} - \frac{(15\sqrt{2}a^2b + (3\sqrt{2}(bx+a)b - 20\sqrt{2}ab)(bx+a)\sqrt{bx+a})}{((bx+a)b-ab)^{3/2}} \right) b^3}{6|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(5/2),x, algorithm="giac")`

output `-1/6*sqrt(2)*(15*sqrt(2)*a*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*b - a*b)))/sqrt(b) - (15*sqrt(2)*a^2*b + (3*sqrt(2)*(b*x + a)*b - 20*sqrt(2)*a*b)*(b*x + a)*sqrt(b*x + a)/((b*x + a)*b - a*b)^(3/2))*b^3/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(5/2),x)`

output `int(atanh(tanh(a + b*x))^(5/2)/x^(5/2), x)`

### Reduce [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{5/2}} dx = \frac{-2\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 10\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{3x^{3/2}}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(5/2), x)`

output `( - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 10*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x + 15*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/x,x)*b**2*x**2)/(3*x**2)`

### 3.236 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{7/2}} dx$

Optimal result	1666
Mathematica [A] (verified)	1666
Rubi [A] (verified)	1667
Maple [B] (verified)	1668
Fricas [A] (verification not implemented)	1670
Sympy [F(-1)]	1670
Maxima [F]	1671
Giac [A] (verification not implemented)	1671
Mupad [F(-1)]	1672
Reduce [F]	1672

#### Optimal result

Integrand size = 17, antiderivative size = 93

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = 2b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right) - \frac{2b^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}}$$

output

```
2*b^(5/2)*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))-2*b^2*arctanh(tanh(b*x+a))^(1/2)/x^(1/2)-2/3*b*arctanh(tanh(b*x+a))^(3/2)/x^(3/2)-2/5*arctanh(tanh(b*x+a))^(5/2)/x^(5/2)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.02

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \frac{2\left(15b^2x^2\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + 5bx\operatorname{arctanh}(\tanh(a + bx))^{3/2} + 3\operatorname{arctanh}(\tanh(a + bx))^{5/2} - 15b^{5/2}\right)}{15x^{5/2}}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(7/2), x]
```

output

```
(-2*(15*b^2*x^2*Sqrt[ArcTanh[Tanh[a + b*x]]] + 5*b*x*ArcTanh[Tanh[a + b*x]]^(3/2) + 3*ArcTanh[Tanh[a + b*x]]^(5/2) - 15*b^(5/2)*x^(5/2)*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(15*x^(5/2))
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2599, 2599, 2599, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx$$

$$\downarrow 2599$$

$$b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{x^{5/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5x^{5/2}}$$

$$\downarrow 2599$$

$$b \left( b \int \frac{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{x^{3/2}} dx - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5x^{5/2}}$$

$$\downarrow 2599$$

$$b \left( b \left( b \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5x^{5/2}}$$

$$\downarrow 2596$$

$$b \left( b \left( 2\sqrt{b} \operatorname{arctanh} \left( \frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \right) - \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{3/2}}{3x^{3/2}} \right) - \frac{2\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{5x^{5/2}}$$



input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(7/2),x]`

output `(-2*ArcTanh[Tanh[a + b*x]]^(5/2))/(5*x^(5/2)) + b*(b*(2*Sqrt[b]*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]]) - (2*Sqrt[ArcTanh[Tanh[a + b*x]]])/Sqrt[x] - (2*ArcTanh[Tanh[a + b*x]]^(3/2))/(3*x^(3/2))`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs.  $2(69) = 138$ .

Time = 0.38 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.66

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \left( \frac{4b}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{6b\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} \right)$
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}} + \left( \frac{4b}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} + \frac{6b\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}} \right)$

input `int(arctanh(tanh(b*x+a))^(5/2)/x^(7/2),x,method=_RETURNVERBOSE)`

output `-2/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(7/2)+4/5*b/(arctanh(tanh(b*x+a))-b*x)*(-1/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh(tanh(b*x+a))^(7/2)+4/3*b/(arctanh(tanh(b*x+a))-b*x)*(-1/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(tanh(b*x+a))^(7/2)+6*b/(arctanh(tanh(b*x+a))-b*x))*(1/6*x^(1/2)*arctanh(tanh(b*x+a))^(5/2)+5/6*(arctanh(tanh(b*x+a))-b*x)*(1/4*x^(1/2)*arctanh(tanh(b*x+a))^(3/2)+3/4*(arctanh(tanh(b*x+a))-b*x)*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)+1/2/b^(1/2)*(arctanh(tanh(b*x+a))-b*x)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))))`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.44

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \left[ \frac{15 b^{5/2} x^3 \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x+a}) - 2(23b^2x^2 + 11abx + 3a^2)}{15x^3} \right]$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(7/2),x, algorithm="fricas")`

output `[1/15*(15*b^(5/2)*x^3*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) - 2*(23*b^2*x^2 + 11*a*b*x + 3*a^2)*sqrt(b*x + a)*sqrt(x))/x^3, -2/15*(15*sqrt(-b)*b^2*x^3*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (23*b^2*x^2 + 11*a*b*x + 3*a^2)*sqrt(b*x + a)*sqrt(x))/x^3]`

### Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**(7/2),x)`

output Timed out

### Maxima [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^{5/2}}{x^{7/2}} dx$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(7/2),x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^(5/2)/x^(7/2), x)`

### Giac [A] (verification not implemented)

Time = 73.14 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.19

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \frac{\sqrt{2} \left( 15 \sqrt{2} b^{5/2} \log \left( \left| -\sqrt{bx + a} \sqrt{b} + \sqrt{(bx + a)b - ab} \right| \right) + \frac{(15 \sqrt{2} a^2 b^5 + (23 \sqrt{2} (bx + a) b^5 - 35 \sqrt{2} ab^5) (bx + a) \sqrt{bx + a})}{((bx + a)b - ab)^{5/2}} \right) b}{15 |b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(7/2),x, algorithm="giac")`

output `-1/15*sqrt(2)*(15*sqrt(2)*b^(5/2)*log(abs(-sqrt(b*x + a)*sqrt(b) + sqrt((b*x + a)*b - a*b))) + (15*sqrt(2)*a^2*b^5 + (23*sqrt(2)*(b*x + a)*b^5 - 35*sqrt(2)*a*b^5)*(b*x + a))*sqrt(b*x + a)/((b*x + a)*b - a*b)^(5/2))*b/abs(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(7/2), x)`output `int(atanh(tanh(a + b*x))^(5/2)/x^(7/2), x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{7/2}} dx = \frac{-6\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 10\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x^{5/2}}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(7/2), x)`output `(-6*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 10*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 30*sqrt(atanh(tanh(a + b*x)))*b**2*x**2 + 15*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x), x)*b**3*x**2)/(15*sqrt(x)*x**2)`

$$3.237 \quad \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx$$

Optimal result	1673
Mathematica [A] (verified)	1673
Rubi [A] (verified)	1674
Maple [A] (verified)	1674
Fricas [A] (verification not implemented)	1675
Sympy [F(-1)]	1675
Maxima [A] (verification not implemented)	1676
Giac [A] (verification not implemented)	1676
Mupad [B] (verification not implemented)	1676
Reduce [F]	1677

### Optimal result

Integrand size = 17, antiderivative size = 35

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
2/7*arctanh(tanh(b*x+a))^(7/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))
```

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{x^{7/2}(7bx - 7\operatorname{arctanh}(\tanh(a+bx)))}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(9/2),x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(x^(7/2)*(7*b*x - 7*ArcTanh[Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(9/2),x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

**Maple [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}$	29
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}$	29

input `int(arctanh(tanh(b*x+a))^(5/2)/x^(9/2),x,method=_RETURNVERBOSE)`

output `-2/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(7/2)`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.20

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx = -\frac{2(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)\sqrt{bx+a}}{7ax^{\frac{7}{2}}}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(9/2),x, algorithm="fricas")`

output `-2/7*(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(b*x + a)/(a*x^(7/2))`

### Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**(9/2),x)`

output `Timed out`



**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.43

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx = -\frac{2(bx + a)^{7/2}}{7ax^{7/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(9/2),x, algorithm="maxima")`output `-2/7*(b*x + a)^(7/2)/(a*x^(7/2))`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx = -\frac{2(bx + a)^{7/2}b^8}{7((bx + a)b - ab)^{7/2}a|b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(9/2),x, algorithm="giac")`output `-2/7*(b*x + a)^(7/2)*b^8/(((b*x + a)*b - a*b)^(7/2)*a*abs(b))`**Mupad [B] (verification not implemented)**

Time = 3.81 (sec) , antiderivative size = 396, normalized size of antiderivative = 11.31

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx = \frac{\sqrt{2} \sqrt{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)} \left( \frac{\ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)^3}{14 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) - 14 \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 28bx} - \frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)^3}{14 \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) - 14 \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)} \right)}{2x^{7/2}}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(9/2),x)`

output

```

-(2^(1/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/
(exp(2*a)*exp(2*b*x) + 1)))^(1/2)*(log(1/(exp(2*a)*exp(2*b*x) + 1)))^3/(14*
log(1/(exp(2*a)*exp(2*b*x) + 1)) - 14*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 28*b*x) - log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x
) + 1))^3/(14*log(1/(exp(2*a)*exp(2*b*x) + 1)) - 14*log((exp(2*a)*exp(2*b*x
))/exp(2*a)*exp(2*b*x) + 1)) + 28*b*x) + (3*log(1/(exp(2*a)*exp(2*b*x) +
1))*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))^2/(14*log(1/(ex
p(2*a)*exp(2*b*x) + 1)) - 14*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x
) + 1)) + 28*b*x) - (3*log(1/(exp(2*a)*exp(2*b*x) + 1))^2*log((exp(2*a)*ex
p(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(14*log(1/(exp(2*a)*exp(2*b*x) + 1))
- 14*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 28*b*x))/(2*
x^(7/2))

```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx = \frac{-2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 2\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{9/2}}$$

input

```
int(atanh(tanh(b*x+a))^(5/2)/x^(9/2), x)
```

output

```

(- 2*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 2*sqrt(atanh(ta
nh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 2*sqrt(atanh(tanh(a + b*x)))*b**2
*x**2 + sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b
*x))*x**2), x)*b**3*x**3)/(7*sqrt(x)*x**3)

```

### 3.238 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx$

Optimal result	1678
Mathematica [A] (verified)	1678
Rubi [A] (verified)	1679
Maple [A] (verified)	1680
Fricas [A] (verification not implemented)	1680
Sympy [F(-1)]	1681
Maxima [A] (verification not implemented)	1681
Giac [A] (verification not implemented)	1681
Mupad [B] (verification not implemented)	1682
Reduce [F]	1682

#### Optimal result

Integrand size = 17, antiderivative size = 72

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx = \frac{4\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{63x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `4/63*b*arctanh(tanh(b*x+a))^(7/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/9*arctanh(tanh(b*x+a))^(7/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx = \frac{2(9bx - 7\operatorname{arctanh}(\tanh(a+bx)))\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{63x^{9/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(11/2),x]`

output `(2*(9*b*x - 7*ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(7/2))/(63*x^(9/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx$$

↓ 2602

$$\frac{2b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{9/2}} dx}{9(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{63x^{7/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(11/2), x]`

output `(4*b*ArcTanh[Tanh[a + b*x]]^(7/2))/(63*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(7/2))/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*(m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{63(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{7}{2}}}$	59
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \frac{4b \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{63(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{7}{2}}}$	59

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(11/2), x, method=_RETURNVERBOSE)
```

output

```
-2/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arctanh(tanh(b*x+a))^(7/2)+4/63*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(7/2)*arctanh(tanh(b*x+a))^(7/2)
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \frac{2(2b^4x^4 - ab^3x^3 - 15a^2b^2x^2 - 19a^3bx - 7a^4)\sqrt{bx + a}}{63a^2x^{\frac{9}{2}}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^(11/2), x, algorithm="fricas")
```

output

```
2/63*(2*b^4*x^4 - a*b^3*x^3 - 15*a^2*b^2*x^2 - 19*a^3*b*x - 7*a^4)*sqrt(b*x + a)/(a^2*x^(9/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**(11/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.47

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \frac{2(2b^2x^2 - 5abx - 7a^2)(bx + a)^{5/2}}{63a^2x^{9/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(11/2),x, algorithm="maxima")`

output `2/63*(2*b^2*x^2 - 5*a*b*x - 7*a^2)*(b*x + a)^(5/2)/(a^2*x^(9/2))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \frac{\sqrt{2} \left( \frac{2\sqrt{2}(bx+a)b^9}{a^2} - \frac{9\sqrt{2}b^9}{a} \right) (bx + a)^{7/2} b}{63((bx + a)b - ab)^{9/2} |b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(11/2),x, algorithm="giac")`

output `1/63*sqrt(2)*(2*sqrt(2)*(b*x + a)*b^9/a^2 - 9*sqrt(2)*b^9/a)*(b*x + a)^(7/2)*b/(((b*x + a)*b - a*b)^(9/2)*abs(b))`

**Mupad [B] (verification not implemented)**

Time = 3.79 (sec) , antiderivative size = 293, normalized size of antiderivative = 4.07

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}{\left(\frac{19bx\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{63}}\right)}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(11/2), x)`output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((19*b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/63 - (10*b^2*x^2)/21 - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/18 + (4*b^3*x^3)/(63*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (16*b^4*x^4)/(63*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(9/2)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx = \frac{-14\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 10\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x^{9/2}}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(11/2), x)`output `( - 14*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 10*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 6*sqrt(atanh(tanh(a + b*x)))*b*x**2 + 3*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**3), x)*b**3*x**4)/(63*sqrt(x)*x**4)`

### 3.239 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx$

Optimal result	1683
Mathematica [A] (verified)	1683
Rubi [A] (verified)	1684
Maple [A] (verified)	1685
Fricas [A] (verification not implemented)	1686
Sympy [F(-1)]	1686
Maxima [A] (verification not implemented)	1687
Giac [A] (verification not implemented)	1687
Mupad [B] (verification not implemented)	1688
Reduce [F]	1688

#### Optimal result

Integrand size = 17, antiderivative size = 110

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx = \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{693x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{8b \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{99x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
16/693*b^2*arctanh(tanh(b*x+a))^(7/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^3
+8/99*b*arctanh(tanh(b*x+a))^(7/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/
11*arctanh(tanh(b*x+a))^(7/2)/x^(11/2)/(b*x-arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.60

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx = \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{7/2} (99b^2x^2 - 154bx \operatorname{arctanh}(\tanh(a+bx)) + 63)}{693x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(13/2), x]
```



output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2)*(99*b^2*x^2 - 154*b*x*ArcTanh[Tanh[a + b*x]] + 63*ArcTanh[Tanh[a + b*x]]^2))/(693*x^(11/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{13/2}} dx$$

↓ 2602

$$\frac{4b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{11/2}} dx}{11(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2602

$$\frac{4b \left( \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{11(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

↓ 2598

$$\frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + 4b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{63x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) / 11(bx - \operatorname{arctanh}(\tanh(a+bx)))$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(13/2), x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(11*x^(11/2)*(b*x - ArcTanh[Tanh[a + b*x]
])) + (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(7/2))/(63*x^(7/2)*(b*x - ArcTanh[
Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(7/2))/(9*x^(9/2)*(b*x - Ar
cTanh[Tanh[a + b*x]]))))/(11*(b*x - ArcTanh[Tanh[a + b*x]]))
```

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /;
NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[
m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + S
imp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; Ne
Q[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m
, -1]
```

**Maple [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \frac{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{63(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{7}{2}}} \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	10
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{8b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \frac{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{63(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 x^{\frac{7}{2}}} \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	10

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(13/2), x, method=_RETURNVERBOSE)
```

output

```
-2/11/(arctanh(tanh(b*x+a))-b*x)/x^(11/2)*arctanh(tanh(b*x+a))^(7/2)-8/11*
b/(arctanh(tanh(b*x+a))-b*x)*(-1/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arct
anh(tanh(b*x+a))^(7/2)+2/63*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(7/2)*arctanh
(tanh(b*x+a))^(7/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.61

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = \frac{2(8b^5x^5 - 4ab^4x^4 + 3a^2b^3x^3 + 113a^3b^2x^2 + 161a^4bx + 63a^5)\sqrt{bx+a}}{693a^3x^{11/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^(13/2),x, algorithm="fricas")
```

output

```
-2/693*(8*b^5*x^5 - 4*a*b^4*x^4 + 3*a^2*b^3*x^3 + 113*a^3*b^2*x^2 + 161*a^
4*b*x + 63*a^5)*sqrt(b*x + a)/(a^3*x^(11/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = \text{Timed out}$$

input

```
integrate(atanh(tanh(b*x+a))**(5/2)/x**(13/2),x)
```

output

```
Timed out
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = -\frac{2(8b^3x^3 - 20ab^2x^2 + 35a^2bx + 63a^3)(bx + a)^{5/2}}{693a^3x^{11/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(13/2),x, algorithm="maxima")`output `-2/693*(8*b^3*x^3 - 20*a*b^2*x^2 + 35*a^2*b*x + 63*a^3)*(b*x + a)^(5/2)/(a^3*x^(11/2))`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.73

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = \frac{\sqrt{2} \left( \frac{99\sqrt{2}b^5}{a} + 4 \left( \frac{2\sqrt{2}(bx+a)b^5}{a^3} - \frac{11\sqrt{2}b^5}{a^2} \right) (bx + a) \right) (bx + a)^{7/2} b^7}{693 ((bx + a)b - ab)^{11/2} |b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(13/2),x, algorithm="giac")`output `-1/693*sqrt(2)*(99*sqrt(2)*b^5/a + 4*(2*sqrt(2)*(b*x + a)*b^5/a^3 - 11*sqrt(2)*b^5/a^2)*(b*x + a))*(b*x + a)^(7/2)*b^7/(((b*x + a)*b - a*b)^(11/2)*abs(b))`

**Mupad [B] (verification not implemented)**

Time = 3.80 (sec) , antiderivative size = 353, normalized size of antiderivative = 3.21

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{23bx \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)}{99} \right)}{x^{13/2}}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(13/2),x)`

output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((23*b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/99 - (226*b^2*x^2)/693 - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/22 + (4*b^3*x^3)/(231*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (32*b^4*x^4)/(693*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + (128*b^5*x^5)/(693*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3))/x^(11/2)
```

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx = \frac{-126\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 70\sqrt{\operatorname{atanh}(\tanh(bx + a))}}{x^{13/2}}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(13/2),x)`

output

```
( - 126*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 70*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 30*sqrt(atanh(tanh(a + b*x)))*b**2*x**2 + 15*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**4),x)*b**3*x**5)/(693*sqrt(x)*x**5)
```

$$3.240 \quad \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{15/2}} dx$$

Optimal result	1689
Mathematica [A] (verified)	1690
Rubi [A] (verified)	1690
Maple [A] (verified)	1692
Fricas [A] (verification not implemented)	1692
Sympy [F(-1)]	1693
Maxima [A] (verification not implemented)	1693
Giac [A] (verification not implemented)	1693
Mupad [B] (verification not implemented)	1694
Reduce [F]	1695

### Optimal result

Integrand size = 17, antiderivative size = 148

$$\int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{15/2}} dx = \frac{32b^3 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{3003x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{16b^2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{429x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{12b \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{143x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2 \operatorname{arctanh}(\tanh(a+bx))^{7/2}}{13x^{13/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
32/3003*b^3*arctanh(tanh(b*x+a))^(7/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))^4+16/429*b^2*arctanh(tanh(b*x+a))^(7/2)/x^(9/2)/(b*x-arctanh(tanh(b*x+a)))^3+12/143*b*arctanh(tanh(b*x+a))^(7/2)/x^(11/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/13*arctanh(tanh(b*x+a))^(7/2)/x^(13/2)/(b*x-arctanh(tanh(b*x+a)))
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.55

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2} (429b^3x^3 - 1001b^2x^2\operatorname{arctanh}(\tanh(a + bx)) - 3003x^{13/2}(-bx + \operatorname{arctanh}(\tanh(a + bx))))}{3003x^{13/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(15/2), x]
```

output

```
(2*ArcTanh[Tanh[a + b*x]]^(7/2)*(429*b^3*x^3 - 1001*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 819*b*x*ArcTanh[Tanh[a + b*x]]^2 - 231*ArcTanh[Tanh[a + b*x]]^3))/((3003*x^(13/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]]))^4)
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx \\ & \quad \downarrow 2602 \\ & \frac{6b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{13/2}} dx}{13(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{13x^{13/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2602 \\ & \frac{6b \left( \frac{4b \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{11/2}} dx}{11(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{13(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\operatorname{arctanh}(\tanh(a + bx))^{7/2}}{13x^{13/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \\ & \quad \downarrow 2602 \end{aligned}$$

$$\begin{aligned}
& 6b \left( \frac{4b \left( \frac{2b \int \frac{\operatorname{arctanh}(\tanh(a+bx))^{5/2}}{x^{9/2}} dx}{9(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{11(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \\
& \frac{13(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}} \\
& \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{13x^{13/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \\
& \quad \downarrow \text{2598} \\
& \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{13x^{13/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \\
& 6b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{11x^{11/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b \left( \frac{2\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{9x^{9/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\operatorname{arctanh}(\tanh(a+bx))^{7/2}}{63x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right)}{11(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{13(bx - \operatorname{arctanh}(\tanh(a+bx)))}{13(bx - \operatorname{arctanh}(\tanh(a+bx)))}
\end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^(5/2)/x^(15/2), x]`

output `(2*ArcTanh[Tanh[a + b*x]]^(7/2))/(13*x^(13/2)*(b*x - ArcTanh[Tanh[a + b*x]]) + (6*b*((2*ArcTanh[Tanh[a + b*x]]^(7/2))/(11*x^(11/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) + (4*b*((4*b*ArcTanh[Tanh[a + b*x]]^(7/2))/(63*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*ArcTanh[Tanh[a + b*x]]^(7/2))/(9*x^(9/2)*(b*x - ArcTanh[Tanh[a + b*x]]))))/(11*(b*x - ArcTanh[Tanh[a + b*x]])))/(13*(b*x - ArcTanh[Tanh[a + b*x]]))`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`



rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*(m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

### Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{13(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{13}{2}}} - \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \dots \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} \right)}{13(\operatorname{arctanh}(\tanh(bx+a))-bx)} \right)}{13(\operatorname{arctanh}(\tanh(bx+a))-bx)}$
default	$-\frac{2 \operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{13(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{13}{2}}} - \frac{12b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} - \frac{4b \left( -\frac{\operatorname{arctanh}(\tanh(bx+a))^{\frac{7}{2}}}{9(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{9}{2}}} + \dots \right)}{11(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{11}{2}}} \right)}{13(\operatorname{arctanh}(\tanh(bx+a))-bx)}$

input

```
int(arctanh(tanh(b*x+a))^(5/2)/x^(15/2), x, method=_RETURNVERBOSE)
```

output

```
-2/13/(arctanh(tanh(b*x+a))-b*x)/x^(13/2)*arctanh(tanh(b*x+a))^(7/2)-12/13*b/(arctanh(tanh(b*x+a))-b*x)*(-1/11/(arctanh(tanh(b*x+a))-b*x)/x^(11/2)*arctanh(tanh(b*x+a))^(7/2)-4/11*b/(arctanh(tanh(b*x+a))-b*x)*(-1/9/(arctanh(tanh(b*x+a))-b*x)/x^(9/2)*arctanh(tanh(b*x+a))^(7/2)+2/63*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(7/2)*arctanh(tanh(b*x+a))^(7/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.53

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{2(16b^6x^6 - 8ab^5x^5 + 6a^2b^4x^4 - 5a^3b^3x^3 - 371a^4b^2x^2 - 567a^5bx - 230a^6)}{3003a^4x^{13/2}}$$

input

```
integrate(arctanh(tanh(b*x+a))^(5/2)/x^(15/2), x, algorithm="fricas")
```

output  $2/3003*(16*b^6*x^6 - 8*a*b^5*x^5 + 6*a^2*b^4*x^4 - 5*a^3*b^3*x^3 - 371*a^4*b^2*x^2 - 567*a^5*b*x - 231*a^6)*\text{sqrt}(b*x + a)/(a^4*x^{(13/2)})$

### Sympy [F(-1)]

Timed out.

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \text{Timed out}$$

input `integrate(atanh(tanh(b*x+a))**(5/2)/x**(15/2),x)`

output Timed out

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.38

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{2(16b^4x^4 - 40ab^3x^3 + 70a^2b^2x^2 - 105a^3bx - 231a^4)(bx + a)^{5/2}}{3003a^4x^{13/2}}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(15/2),x, algorithm="maxima")`

output  $2/3003*(16*b^4*x^4 - 40*a*b^3*x^3 + 70*a^2*b^2*x^2 - 105*a^3*b*x - 231*a^4)*(b*x + a)^{(5/2)}/(a^4*x^{(13/2)})$

### Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.66

$$\int \frac{\text{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{\sqrt{2} \left( \frac{429\sqrt{2}b^{13}}{a} - 2 \left( \frac{143\sqrt{2}b^{13}}{a^2} + 4 \left( \frac{2\sqrt{2}(bx+a)b^{13}}{a^4} - \frac{13\sqrt{2}b^{13}}{a^3} \right) (bx + a) \right) (bx + a) \right) (bx + a)^{7/2} b}{3003((bx + a)b - ab)^{13/2} |b|}$$

input `integrate(arctanh(tanh(b*x+a))^(5/2)/x^(15/2),x, algorithm="giac")`

output `-1/3003*sqrt(2)*(429*sqrt(2)*b^13/a - 2*(143*sqrt(2)*b^13/a^2 + 4*(2*sqrt(2)*(b*x + a)*b^13/a^4 - 13*sqrt(2)*b^13/a^3)*(b*x + a))*(b*x + a)*(b*x + a)^(7/2)*b/(((b*x + a)*b - a*b)^(13/2)*abs(b))`

### Mupad [B] (verification not implemented)

Time = 3.93 (sec) , antiderivative size = 413, normalized size of antiderivative = 2.79

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{27bx \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)}{143} \right)}{143}$$

input `int(atanh(tanh(a + b*x))^(5/2)/x^(15/2),x)`

output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((27*b*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/143 - (10*6*b^2*x^2)/429 - (log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/26 + (20*b^3*x^3)/(3003*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (16*b^4*x^4)/(1001*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (128*b^5*x^5)/(3003*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (512*b^6*x^6)/(3003*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4))/x^(13/2)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^{5/2}}{x^{15/2}} dx = \frac{-66\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))^2 - 30\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{13/2}} - \frac{10\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{11/2}} + \frac{5\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{9/2}} + \frac{5\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{7/2}} + \frac{5\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{5/2}} + \frac{5\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{3/2}} + \frac{5\sqrt{\operatorname{atanh}(\tanh(bx + a))} \operatorname{atanh}(\tanh(bx + a))}{x^{1/2}}$$

input `int(atanh(tanh(b*x+a))^(5/2)/x^(15/2),x)`

output `( - 66*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))**2 - 30*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x))*b*x - 10*sqrt(atanh(tanh(a + b*x)))*b**2*x**2 + 5*sqrt(x)*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**5),x)*b**3*x**6)/(429*sqrt(x)*x**6)`

**3.241**  $\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1696
Mathematica [A] (verified)	1697
Rubi [A] (verified)	1697
Maple [A] (verified)	1699
Fricas [A] (verification not implemented)	1700
Sympy [F]	1700
Maxima [F]	1700
Giac [A] (verification not implemented)	1701
Mupad [F(-1)]	1701
Reduce [F]	1701

**Optimal result**

Integrand size = 17, antiderivative size = 145

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{5\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{8b^{7/2}} + \frac{x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b} + \frac{5x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{12b^2} + \frac{5\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{8b^3}$$

output

```
5/8*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^3/b^(7/2)+1/3*x^(5/2)*arctanh(tanh(b*x+a))^(1/2)/b+5/12*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b^2+5/8*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)/b^3
```

**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.72

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))(33b^2x^2 - 40bx\operatorname{arctanh}(\tanh(a+bx)) + 15(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx)))}}{24b^3} + \frac{5(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx)))}}{8b^{7/2}}$$

input `Integrate[x^(5/2)/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*(33*b^2*x^2 - 40*b*x*ArcTanh[Tanh[a + b*x]] + 15*ArcTanh[Tanh[a + b*x]]^2))/(24*b^3) + (5*(b*x - ArcTanh[Tanh[a + b*x]])^3*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(8*b^(7/2))`

**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

↓ 2600

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a+bx)))}{6b} \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \frac{x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b}$$

↓ 2600

$$5(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2b} \right) + \frac{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3b} \frac{6b}{3b}$$

↓ 2600

$$5(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{2b} + \sqrt{x} \sqrt{a} \right)}{4b} \right) + \frac{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3b} \frac{6b}{3b}$$

↓ 2596

$$5 \left( \frac{3 \left( \frac{\operatorname{arctanh} \left( \frac{\sqrt{b} \sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{b^{3/2}} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right) (bx - \operatorname{arctanh}(\tanh(a + bx)))}{4b} \right) + \frac{x^{3/2}}{3b} \frac{6b}{3b}$$

input `Int [x^(5/2)/Sqrt [ArcTanh [Tanh [a + b*x]]], x]`

output `(5*((3*((ArcTanh [(Sqrt [b]*Sqrt [x])/Sqrt [ArcTanh [Tanh [a + b*x]]]])*(b*x - ArcTanh [Tanh [a + b*x]]))/b^(3/2) + (Sqrt [x]*Sqrt [ArcTanh [Tanh [a + b*x]]])/b)*(b*x - ArcTanh [Tanh [a + b*x]]))/(4*b) + (x^(3/2)*Sqrt [ArcTanh [Tanh [a + b*x]]])/(2*b))*(b*x - ArcTanh [Tanh [a + b*x]])/(6*b) + (x^(5/2)*Sqrt [ArcTanh [Tanh [a + b*x]]])/(3*b)`

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

**Maple [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{x^{\frac{5}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3b} - \frac{5(\operatorname{arctanh}(\tanh(bx+a)) - bx) \left( \frac{x^{\frac{3}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{4b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx)}{3} \right)}{3b}$
default	$\frac{x^{\frac{5}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3b} - \frac{5(\operatorname{arctanh}(\tanh(bx+a)) - bx) \left( \frac{x^{\frac{3}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{4b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx)}{3} \right)}{3b}$

```
input int(x^(5/2)/arctanh(tanh(b*x+a))^(1/2), x, method=_RETURNVERBOSE)
```

```
output 1/3*x^(5/2)*arctanh(tanh(b*x+a))^(1/2)/b-5/3*(arctanh(tanh(b*x+a))-b*x)/b*(1/4*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)/b-3/4*(arctanh(tanh(b*x+a))-b*x)/b*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b-1/2*(arctanh(tanh(b*x+a))-b*x)/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```



**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.94

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \left[ \frac{15 a^3 \sqrt{b} \log(2bx - 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a) + 2(8b^3x^2 - 10ab^2x + 15a^2b)}{48b^4} \right]$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[1/48*(15*a^3*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(8*b^3*x^2 - 10*a*b^2*x + 15*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^4, 1/24*(15*a^3*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (8*b^3*x^2 - 10*a*b^2*x + 15*a^2*b)*sqrt(b*x + a)*sqrt(x))/b^4]`

**Sympy [F]**

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^{5/2}}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(x**(5/2)/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(x**(5/2)/sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [F]**

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^{5/2}}{\sqrt{\operatorname{artanh}(\tanh(bx + a))}} dx$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(x^(5/2)/sqrt(arctanh(tanh(b*x + a))), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.44

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{1}{24} \sqrt{bx+a} \left( 2x \left( \frac{4x}{b} - \frac{5a}{b^2} \right) + \frac{15a^2}{b^3} \right) \sqrt{x} + \frac{5a^3 \log \left( \left| -\sqrt{b}\sqrt{x} + \sqrt{bx+a} \right| \right)}{8b^{7/2}}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`output `1/24*sqrt(b*x + a)*(2*x*(4*x/b - 5*a/b^2) + 15*a^2/b^3)*sqrt(x) + 5/8*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(7/2)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{x^{5/2}}{\sqrt{\operatorname{atanh}(\tanh(a+bx))}} dx$$

input `int(x^(5/2)/atanh(tanh(a + b*x))^(1/2),x)`output `int(x^(5/2)/atanh(tanh(a + b*x))^(1/2), x)`**Reduce [F]**

$$\int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))} x^2}{\operatorname{atanh}(\tanh(bx+a))} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a))^(1/2),x)`output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x**2)/atanh(tanh(a + b*x)),x)`

**3.242**  $\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1702
Mathematica [A] (verified)	1703
Rubi [A] (verified)	1703
Maple [A] (verified)	1705
Fricas [A] (verification not implemented)	1705
Sympy [F]	1706
Maxima [F]	1706
Giac [A] (verification not implemented)	1706
Mupad [F(-1)]	1707
Reduce [F]	1707

**Optimal result**

Integrand size = 17, antiderivative size = 107

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{3\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{4b^{5/2}} + \frac{x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} + \frac{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{4b^2}$$

output

```
3/4*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^2/b^(5/2)+1/2*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)/b+3/4*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b^2
```

### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.82

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{\sqrt{b}\sqrt{x}(5bx - 3\operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + 3(-b\sqrt{x} + \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{4b^5}$$

input `Integrate[x^(3/2)/Sqrt[ArcTanh[Tanh[a + b*x]]], x]`

output `(Sqrt[b]*Sqrt[x]*(5*b*x - 3*ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]] + 3*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]])/(4*b^(5/2))`

### Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

$$\downarrow 2600$$

$$\frac{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2b}$$

$$\downarrow 2600$$

$$\frac{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{2b} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right)}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2b}$$

↓ 2596

$$3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right) (bx - \operatorname{arctanh}(\tanh(a+bx)))$$


---


$$\frac{x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b}$$

input `Int[x^(3/2)/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]])/b)*(b*x - ArcTanh[Tanh[a + b*x]]))/(4*b) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/(2*b)`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1)) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{x^{\frac{3}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2b} - \frac{(\operatorname{arctanh}(\tanh(bx+a)) - bx) \ln(\dots)}{2b} \right)}{2b}$
default	$\frac{x^{\frac{3}{2}} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2b} - \frac{3(\operatorname{arctanh}(\tanh(bx+a)) - bx) \left( \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{2b} - \frac{(\operatorname{arctanh}(\tanh(bx+a)) - bx) \ln(\dots)}{2b} \right)}{2b}$

input `int(x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)/b-3/2*(arctanh(tanh(b*x+a))-b*x)/b*(1/2*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b-1/2*(arctanh(tanh(b*x+a))-b*x)/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.08

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \left[ \frac{3a^2\sqrt{b} \log\left(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a\right) + 2(2b^2x - 3ab)\sqrt{bx+a}\sqrt{x}}{8b^3} - \frac{3a^2\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) - (2b^2x - 3ab)\sqrt{bx+a}\sqrt{x}}{4b^3} \right]$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[1/8*(3*a^2*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(2*b^2*x - 3*a*b)*sqrt(b*x + a)*sqrt(x))/b^3, -1/4*(3*a^2*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) - (2*b^2*x - 3*a*b)*sqrt(b*x + a)*sqrt(x))/b^3]`

**Sympy [F]**

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^{3/2}}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(x**(3/2)/atanh(tanh(b*x+a))**(1/2), x)`

output `Integral(x**(3/2)/sqrt(atanh(tanh(a + b*x))), x)`

**Maxima [F]**

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^{3/2}}{\sqrt{\operatorname{artanh}(\tanh(bx + a))}} dx$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(1/2), x, algorithm="maxima")`

output `integrate(x^(3/2)/sqrt(arctanh(tanh(b*x + a))), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.49

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{1}{4} \sqrt{bx + a} \sqrt{x} \left( \frac{2x}{b} - \frac{3a}{b^2} \right) - \frac{3a^2 \log \left( \left| -\sqrt{b} \sqrt{x} + \sqrt{bx + a} \right| \right)}{4b^{5/2}}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(1/2), x, algorithm="giac")`

output `1/4*sqrt(b*x + a)*sqrt(x)*(2*x/b - 3*a/b^2) - 3/4*a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{x^{3/2}}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `int(x^(3/2)/atanh(tanh(a + b*x))^(1/2),x)`output `int(x^(3/2)/atanh(tanh(a + b*x))^(1/2), x)`**Reduce [F]**

$$\int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(3/2)/atanh(tanh(b*x+a))^(1/2),x)`output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x)/atanh(tanh(a + b*x)),x)`



**3.243**  $\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1708
Mathematica [A] (verified)	1709
Rubi [A] (verified)	1709
Maple [A] (verified)	1710
Fricas [A] (verification not implemented)	1711
Sympy [F]	1711
Maxima [F]	1711
Giac [A] (verification not implemented)	1712
Mupad [F(-1)]	1712
Reduce [F]	1713

**Optimal result**

Integrand size = 17, antiderivative size = 63

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$= \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

output

```
arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))/b^(3/2)+x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$= \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

$$- \frac{(-bx + \operatorname{arctanh}(\tanh(a+bx))) \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{b^{3/2}}$$

input `Integrate[Sqrt[x]/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]])/b - ((-(b*x) + ArcTanh[Tanh[a + b*x]])*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b^(3/2)`

**Rubi [A] (verified)**

Time = 0.36 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$\downarrow 2600$$

$$\frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{2b} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

$$\downarrow 2596$$

$$\frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}$$

input `Int[Sqrt[x]/Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]]*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b`

**Defintions of rubi rules used**

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1)) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

**Maple [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))}\right)}{b^{\frac{3}{2}}}$	53
default	$\frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \ln\left(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))}\right)}{b^{\frac{3}{2}}}$	53

input `int(x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b-(arctanh(tanh(b*x+a))-b*x)/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.40

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$= \left[ \frac{a\sqrt{b} \log\left(2bx - 2\sqrt{bx+a}\sqrt{b}\sqrt{x} + a\right) + 2\sqrt{bx+a}ab\sqrt{x}}{2b^2}, \frac{a\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) + \sqrt{bx+a}ab\sqrt{x}}{b^2} \right]$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`output `[1/2*(a*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*sqrt(b*x + a)*b*sqrt(x))/b^2, (a*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + sqrt(b*x + a)*b*sqrt(x))/b^2]`**Sympy [F]**

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{\sqrt{x}}{\sqrt{\operatorname{atanh}(\tanh(a+bx))}} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a))**(1/2),x)`output `Integral(sqrt(x)/sqrt(atanh(tanh(a + b*x))), x)`**Maxima [F]**

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \int \frac{\sqrt{x}}{\sqrt{\operatorname{artanh}(\tanh(bx+a))}} dx$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x)/sqrt(arctanh(tanh(b*x + a))), x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.60

$$\int \frac{\sqrt{x}}{\sqrt{\arctanh(\tanh(a + bx))}} dx = \frac{a \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{b^{\frac{3}{2}}} + \frac{\sqrt{bx + a}\sqrt{x}}{b}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `a*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(3/2) + sqrt(b*x + a)*sqrt(x)/b`

### Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x}}{\sqrt{\arctanh(\tanh(a + bx))}} dx = \int \frac{\sqrt{x}}{\sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `int(x^(1/2)/atanh(tanh(a + b*x))^(1/2),x)`

output `int(x^(1/2)/atanh(tanh(a + b*x))^(1/2), x)`

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))} dx$$

input `int(x^(1/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/atanh(tanh(a + b*x)),x)`

**3.244**  $\int \frac{1}{\sqrt{x}\sqrt{\mathbf{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1714
Mathematica [A] (verified)	1714
Rubi [A] (verified)	1715
Maple [A] (verified)	1715
Fricas [A] (verification not implemented)	1716
Sympy [F]	1716
Maxima [F]	1717
Giac [A] (verification not implemented)	1717
Mupad [F(-1)]	1717
Reduce [F]	1718

**Optimal result**

Integrand size = 17, antiderivative size = 30

$$\int \frac{1}{\sqrt{x}\sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = \frac{2\mathbf{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\mathbf{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{b}}$$

output

```
2*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))/b^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{x}\sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = \frac{2 \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\mathbf{arctanh}(\tanh(a + bx))}\right)}{\sqrt{b}}$$

input

```
Integrate[1/(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]
```

output

```
(2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]])/Sqrt[b]
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2596

$$\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{\sqrt{b}}$$

input `Int[1/(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `(2*ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]])/Sqrt[b]`

**Defintions of rubi rules used**

rule 2596 `Int[1/(Sqrt[u]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

**Maple [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$\frac{2 \ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{\sqrt{b}}$	24
default	$\frac{2 \ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{\sqrt{b}}$	24



input `int(1/x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `2*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))/b^(1/2)`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.80

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \left[ \frac{\log\left(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x+a}\right)}{\sqrt{b}}, \right. \\ \left. - \frac{2\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right)}{b} \right]$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `[log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a)/sqrt(b), -2*sqrt(-b)*arc  
tan(sqrt(-b)*sqrt(x)/sqrt(b*x + a))/b]`

### Sympy [F]

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(1/x**(1/2)/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(1/(sqrt(x)*sqrt(atanh(tanh(a + b*x)))), x)`

**Maxima [F]**

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{\sqrt{x} \sqrt{\operatorname{artanh}(\tanh(bx + a))}} dx$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(x)*sqrt(arctanh(tanh(b*x + a)))), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -\frac{2 \log \left( \left| -\sqrt{b} \sqrt{x} + \sqrt{bx + a} \right| \right)}{\sqrt{b}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `-2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/sqrt(b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))^(1/2)),x)`

output `int(1/(x^(1/2)*atanh(tanh(a + b*x))^(1/2)), x)`

**Reduce [F]**

$$\int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x} dx$$

input `int(1/x^(1/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x),x)`

$$3.245 \quad \int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

Optimal result	1719
Mathematica [A] (verified)	1719
Rubi [A] (verified)	1720
Maple [A] (verified)	1720
Fricas [A] (verification not implemented)	1721
Sympy [F]	1721
Maxima [A] (verification not implemented)	1722
Giac [A] (verification not implemented)	1722
Mupad [B] (verification not implemented)	1722
Reduce [F]	1723

### Optimal result

Integrand size = 17, antiderivative size = 33

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output `2*arctanh(tanh(b*x+a))^(1/2)/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))`

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = -\frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{\sqrt{x}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}$$

input `Integrate[1/(x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `(-2*Sqrt[ArcTanh[Tanh[a + b*x]])/(Sqrt[x]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2598

$$\frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[1/(x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output `(2*Sqrt[ArcTanh[Tanh[a + b*x]])/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))`

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}}$	29
default	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{x}}$	29

input `int(1/x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output `-2/(arctanh(tanh(b*x+a))-b*x)/x^(1/2)*arctanh(tanh(b*x+a))^(1/2)`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -\frac{2\sqrt{bx+a}}{a\sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`

output `-2*sqrt(b*x + a)/(a*sqrt(x))`

### Sympy [F]

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^{3/2} \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(1/x**(3/2)/atanh(tanh(b*x+a))**(1/2),x)`

output `Integral(1/(x**(3/2)*sqrt(atanh(tanh(a + b*x)))), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -\frac{2\sqrt{bx+a}}{a\sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`output `-2*sqrt(b*x + a)/(a*sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{4\sqrt{b}}{(\sqrt{b}\sqrt{x} - \sqrt{bx+a})^2 - a}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`output `4*sqrt(b)/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)`**Mupad [B] (verification not implemented)**

Time = 3.74 (sec) , antiderivative size = 101, normalized size of antiderivative = 3.06

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{4\sqrt{\frac{\ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right)}{2}}}{\sqrt{x} \left( \ln\left(\frac{1}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)}$$

input `int(1/(x^(3/2)*atanh(tanh(a + b*x))^(1/2)),x)`

output  $(4 * (\log((\exp(2*a) * \exp(2*b*x)) / (\exp(2*a) * \exp(2*b*x) + 1))) / 2 - \log(1 / (\exp(2*a) * \exp(2*b*x) + 1))) / 2^{(1/2)} / (x^{(1/2)} * (\log(1 / (\exp(2*a) * \exp(2*b*x) + 1)) - \log((\exp(2*a) * \exp(2*b*x)) / (\exp(2*a) * \exp(2*b*x) + 1)) + 2*b*x))$

### Reduce [F]

$$\int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^2} dx$$

input `int(1/x^(3/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**2),x)`



**3.246**  $\int \frac{1}{x^{5/2} \sqrt{\mathbf{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1724
Mathematica [A] (verified)	1724
Rubi [A] (verified)	1725
Maple [A] (verified)	1726
Fricas [A] (verification not implemented)	1727
Sympy [F]	1727
Maxima [A] (verification not implemented)	1727
Giac [A] (verification not implemented)	1728
Mupad [B] (verification not implemented)	1728
Reduce [F]	1729

**Optimal result**

Integrand size = 17, antiderivative size = 72

$$\int \frac{1}{x^{5/2} \sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = \frac{4b \sqrt{\mathbf{arctanh}(\tanh(a + bx))}}{3\sqrt{x}(bx - \mathbf{arctanh}(\tanh(a + bx)))^2} + \frac{2\sqrt{\mathbf{arctanh}(\tanh(a + bx))}}{3x^{3/2}(bx - \mathbf{arctanh}(\tanh(a + bx)))}$$

output

$4/3*b*\mathbf{arctanh}(\mathbf{tanh}(b*x+a))^{(1/2)}/x^{(1/2)}/(b*x-\mathbf{arctanh}(\mathbf{tanh}(b*x+a)))^{2+2/3*}$   
 $\mathbf{arctanh}(\mathbf{tanh}(b*x+a))^{(1/2)}/x^{(3/2)}/(b*x-\mathbf{arctanh}(\mathbf{tanh}(b*x+a)))$

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.64

$$\int \frac{1}{x^{5/2} \sqrt{\mathbf{arctanh}(\tanh(a + bx))}} dx = \frac{2\sqrt{\mathbf{arctanh}(\tanh(a + bx))}(-3bx + \mathbf{arctanh}(\tanh(a + bx)))}{3x^{3/2}(-bx + \mathbf{arctanh}(\tanh(a + bx)))^2}$$

input

`Integrate[1/(x^(5/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]`

output

```
(-2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(-3*b*x + ArcTanh[Tanh[a + b*x]]))/(3*x^(3/2)*(-b*x + ArcTanh[Tanh[a + b*x]])^2)
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2602

$$\frac{2b \int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{4b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2}$$

input

```
Int[1/(x^(5/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]
```

output

```
(4*b*Sqrt[ArcTanh[Tanh[a + b*x]]]/(3*Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]]/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])))
```

## Definitions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

## Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}$	59
default	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}} + \frac{4b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}$	59

input `int(1/x^(5/2)/arctanh(tanh(b*x+a))^(1/2), x, method=_RETURNVERBOSE)`

output 
$$-2/3/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)/x^{3/2}*\operatorname{arctanh}(\tanh(b*x+a))^{1/2}+4/3*b/(\operatorname{arctanh}(\tanh(b*x+a))-b*x)^2/x^{1/2}*\operatorname{arctanh}(\tanh(b*x+a))^{1/2}$$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.32

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(2bx - a)\sqrt{bx + a}}{3a^2x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")`output `2/3*(2*b*x - a)*sqrt(b*x + a)/(a^2*x^(3/2))`**Sympy [F]**

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{1}{x^{5/2} \sqrt{\operatorname{atanh}(\tanh(a + bx))}} dx$$

input `integrate(1/x**(5/2)/atanh(tanh(b*x+a))**(1/2),x)`output `Integral(1/(x**(5/2)*sqrt(atanh(tanh(a + b*x))))), x)`**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.46

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(2b^2x^2 + abx - a^2)}{3\sqrt{bx + a}a^2x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`output `2/3*(2*b^2*x^2 + a*b*x - a^2)/(sqrt(b*x + a)*a^2*x^(3/2))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{8 \left( 3 \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a \right) b^{3/2}}{3 \left( \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a \right)^3}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `8/3*(3*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)*b^(3/2)/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^3`

**Mupad [B] (verification not implemented)**

Time = 3.82 (sec) , antiderivative size = 218, normalized size of antiderivative = 3.03

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{\sqrt{2} \left( \frac{4 \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right)}{3} - \frac{4 \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right)}{3} + \frac{8bx}{3^2} \right) + \frac{16bx}{3 \left( \ln\left(\frac{2}{e^{2a} e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx+1}}\right) \right)}}{2x^{3/2}}$$

input `int(1/(x^(5/2)*atanh(tanh(a + b*x))^(1/2)),x)`

output `(2^(1/2)*(((4*log(2/(exp(2*a)*exp(2*b*x) + 1)))/3 - (4*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/3 + (8*b*x)/3)/log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2 + (16*b*x)/(3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x^2))*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^(1/2))/(2*x^(3/2))`

**Reduce [F]**

$$\int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^3} dx$$

input `int(1/x^(5/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**3),x)`

**3.247**  $\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1730
Mathematica [A] (verified)	1730
Rubi [A] (verified)	1731
Maple [A] (verified)	1732
Fricas [A] (verification not implemented)	1733
Sympy [F(-1)]	1733
Maxima [A] (verification not implemented)	1734
Giac [A] (verification not implemented)	1734
Mupad [B] (verification not implemented)	1734
Reduce [F]	1735

**Optimal result**

Integrand size = 17, antiderivative size = 110

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{16b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{15\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{8b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
16/15*b^2*arctanh(tanh(b*x+a))^(1/2)/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3+
8/15*b*arctanh(tanh(b*x+a))^(1/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2+2/5
*arctanh(tanh(b*x+a))^(1/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.60

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}(15b^2x^2 - 10bx\operatorname{arctanh}(\tanh(a+bx)) + 3)}{15x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3}$$

input

```
Integrate[1/(x^(7/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]
```

output

```
(2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(15*b^2*x^2 - 10*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2))/(15*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3)
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx$$

↓ 2602

$$\frac{4b \int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2602

$$\frac{4b \left( \frac{2b \int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} \right)}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

↓ 2598

$$\frac{4b \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{4b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^2} \right)}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input

```
Int[1/(x^(7/2)*Sqrt[ArcTanh[Tanh[a + b*x]]],x]
```



output

```
(4*b*((4*b*Sqrt[ArcTanh[Tanh[a + b*x]]])/(3*Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])^2) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]])/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]]))) / (5*(b*x - ArcTanh[Tanh[a + b*x]]) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]]) / (5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])))
```

**Defintions of rubi rules used**

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{8b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}+\frac{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}\right)}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	105
default	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{8b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}}+\frac{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{x}}\right)}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}$	105

input

```
int(1/x^(7/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-2/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(1/2)-8/5*b/(
arctanh(tanh(b*x+a))-b*x)*(-1/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh
(tanh(b*x+a))^(1/2)+2/3*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(1/2)*arctanh(tan
h(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.31

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -\frac{2(8b^2x^2 - 4abx + 3a^2)\sqrt{bx + a}}{15a^3x^{5/2}}$$

input

```
integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
-2/15*(8*b^2*x^2 - 4*a*b*x + 3*a^2)*sqrt(b*x + a)/(a^3*x^(5/2))
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \text{Timed out}$$

input

```
integrate(1/x**(7/2)/atanh(tanh(b*x+a))**(1/2),x)
```

output

```
Timed out
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = -\frac{2(8b^3x^3 + 4ab^2x^2 - a^2bx + 3a^3)}{15\sqrt{bx + a}a^3x^{5/2}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output `-2/15*(8*b^3*x^3 + 4*a*b^2*x^2 - a^2*b*x + 3*a^3)/(sqrt(b*x + a)*a^3*x^(5/2))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.70

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{32 \left( 10 \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^4 - 5a \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 + a^2 \right) b^{5/2}}{15 \left( \left( \sqrt{b}\sqrt{x} - \sqrt{bx + a} \right)^2 - a \right)^5}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `32/15*(10*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 - 5*a*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 + a^2)*b^(5/2)/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^5`

**Mupad [B] (verification not implemented)**

Time = 3.59 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.06

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)}{2}}}{5 \left( \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx \right)}$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))^(1/2)),x)`

output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(4/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (128*b^2*x^2)/(15*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (32*b*x)/(15*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(5/2)`

### Reduce [F]

$$\int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^4} dx$$

input `int(1/x^(7/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**4),x)`

**3.248**  $\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$

Optimal result	1736
Mathematica [A] (verified)	1737
Rubi [A] (verified)	1737
Maple [A] (verified)	1739
Fricas [A] (verification not implemented)	1739
Sympy [F(-1)]	1740
Maxima [A] (verification not implemented)	1740
Giac [A] (verification not implemented)	1741
Mupad [B] (verification not implemented)	1741
Reduce [F]	1742

**Optimal result**

Integrand size = 17, antiderivative size = 148

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{32b^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{35\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^4} + \frac{16b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{35x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3} + \frac{12b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{35x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

output

```
32/35*b^3*arctanh(tanh(b*x+a))^(1/2)/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^4+
16/35*b^2*arctanh(tanh(b*x+a))^(1/2)/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^3+
12/35*b*arctanh(tanh(b*x+a))^(1/2)/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))^2+
7*arctanh(tanh(b*x+a))^(1/2)/x^(7/2)/(b*x-arctanh(tanh(b*x+a)))
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.55

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx = \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))(35b^3x^3 - 35b^2x^2\operatorname{arctanh}(\tanh(a+bx)) - 35x^{7/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}}{35x^{7/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))}$$

input

```
Integrate[1/(x^(9/2)*Sqrt[ArcTanh[Tanh[a + b*x]]]),x]
```

output

```
(2*Sqrt[ArcTanh[Tanh[a + b*x]]]*(35*b^3*x^3 - 35*b^2*x^2*ArcTanh[Tanh[a + b*x]] + 21*b*x*ArcTanh[Tanh[a + b*x]]^2 - 5*ArcTanh[Tanh[a + b*x]]^3))/(35*x^(7/2)*(-b*x) + ArcTanh[Tanh[a + b*x]])^4
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.24, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx$$

$$\downarrow 2602$$

$$6b \int \frac{1}{x^{7/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$$\downarrow 2602$$

$$6b \left( \frac{4b \int \frac{1}{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) + \frac{7(bx - \operatorname{arctanh}(\tanh(a+bx)))}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$$\downarrow 2602$$

$$\begin{aligned}
& 6b \left( \frac{4b \int \frac{1}{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{7(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
& \frac{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2598} \\
& 6b \left( \frac{4b \left( \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{4b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} \right) + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) \\
& \frac{7(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
& \frac{7x^{7/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2598}
\end{aligned}$$

input `Int[1/(x^(9/2)*Sqrt[ArcTanh[Tanh[a + b*x]]],x]`

output `(6*b*((4*b*((4*b*Sqrt[ArcTanh[Tanh[a + b*x]]])/(3*Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]]))^2) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]])/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]]))))/(5*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]])/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])))/(7*(b*x - ArcTanh[Tanh[a + b*x]])) + (2*Sqrt[ArcTanh[Tanh[a + b*x]]])/(7*x^(7/2)*(b*x - ArcTanh[Tanh[a + b*x]]))`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

### Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}-\frac{12b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{4b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}+\frac{2}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{1}{2}}}\right)}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}\right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}$
default	$-\frac{2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{7}{2}}}-\frac{12b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{5}{2}}}-\frac{4b\left(-\frac{\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{3}{2}}+\frac{2}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)x^{\frac{1}{2}}}\right)}{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}\right)}{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}$

input

```
int(1/x^(9/2)/arctanh(tanh(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-2/7/(arctanh(tanh(b*x+a))-b*x)/x^(7/2)*arctanh(tanh(b*x+a))^(1/2)-12/7*b/(arctanh(tanh(b*x+a))-b*x)*(-1/5/(arctanh(tanh(b*x+a))-b*x)/x^(5/2)*arctanh(tanh(b*x+a))^(1/2)-4/5*b/(arctanh(tanh(b*x+a))-b*x)*(-1/3/(arctanh(tanh(b*x+a))-b*x)/x^(3/2)*arctanh(tanh(b*x+a))^(1/2)+2/3*b/(arctanh(tanh(b*x+a))-b*x)^2/x^(1/2)*arctanh(tanh(b*x+a))^(1/2)))
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.30

$$\int \frac{1}{x^{9/2}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{2(16b^3x^3 - 8ab^2x^2 + 6a^2bx - 5a^3)\sqrt{bx + a}}{35a^4x^{\frac{7}{2}}}$$

input

```
integrate(1/x^(9/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="fricas")
```



output  $2/35*(16*b^3*x^3 - 8*a*b^2*x^2 + 6*a^2*b*x - 5*a^3)*\text{sqrt}(b*x + a)/(a^4*x^{(7/2)})$

### Sympy [F(-1)]

Timed out.

$$\int \frac{1}{x^{9/2} \sqrt{\text{arctanh}(\tanh(a + bx))}} dx = \text{Timed out}$$

input `integrate(1/x**(9/2)/atanh(tanh(b*x+a))**(1/2),x)`

output Timed out

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.37

$$\int \frac{1}{x^{9/2} \sqrt{\text{arctanh}(\tanh(a + bx))}} dx = \frac{2(16b^4x^4 + 8ab^3x^3 - 2a^2b^2x^2 + a^3bx - 5a^4)}{35\sqrt{bx + aa^4x^7}}$$

input `integrate(1/x^(9/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="maxima")`

output  $2/35*(16*b^4*x^4 + 8*a*b^3*x^3 - 2*a^2*b^2*x^2 + a^3*b*x - 5*a^4)/(\text{sqrt}(b*x + a)*a^4*x^{(7/2)})$

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.70

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{64 \left( 35 \left( \sqrt{b} \sqrt{x} - \sqrt{bx + a} \right)^6 - 21 a \left( \sqrt{b} \sqrt{x} - \sqrt{bx + a} \right)^4 + 7 a^2 \left( \sqrt{b} \sqrt{x} - \sqrt{bx + a} \right)^2 - a^3 \right) b^{7/2}}{35 \left( \left( \sqrt{b} \sqrt{x} - \sqrt{bx + a} \right)^2 - a \right)^7}$$

input `integrate(1/x^(9/2)/arctanh(tanh(b*x+a))^(1/2),x, algorithm="giac")`

output `64/35*(35*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 - 21*a*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 + 7*a^2*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a^3)*b^(7/2)/((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^7`

**Mupad [B] (verification not implemented)**

Time = 3.54 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.94

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{7 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^4}$$

input `int(1/(x^(9/2)*atanh(tanh(a + b*x))^(1/2)),x)`

output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(4/(7*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (128*b^2*x^2)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (512*b^3*x^3)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4) + (48*b*x)/(35*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(7/2)`

**Reduce [F]**

$$\int \frac{1}{x^{9/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) x^5} dx$$

input `int(1/x^(9/2)/atanh(tanh(b*x+a))^(1/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x**5),x)`

**3.249**  $\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1743
Mathematica [A] (verified)	1744
Rubi [A] (verified)	1744
Maple [A] (verified)	1747
Fricas [A] (verification not implemented)	1748
Sympy [F(-1)]	1748
Maxima [F]	1749
Giac [A] (verification not implemented)	1749
Mupad [F(-1)]	1749
Reduce [F]	1750

**Optimal result**

Integrand size = 17, antiderivative size = 166

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{35\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^3}{8b^{9/2}} - \frac{2x^{7/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{7x^{5/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b^2} + \frac{35x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{12b^3} + \frac{35\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{8b^4}$$

output

```
35/8*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^3/b^(9/2)-2*x^(7/2)/b/arctanh(tanh(b*x+a))^(1/2)+7/3*x^(5/2)*arctanh(tanh(b*x+a))^(1/2)/b^2+35/12*x^(3/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b^3+35/8*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))^2*arctanh(tanh(b*x+a))^(1/2)/b^4
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.73

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{\sqrt{x}(-48b^3x^3 + 231b^2x^2\operatorname{arctanh}(\tanh(a+bx)) - 280bx\operatorname{arctanh}(\tanh(a+bx))}{24b^4\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{35(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{8b^{9/2}}$$

input

```
Integrate[x^(7/2)/ArcTanh[Tanh[a + b*x]]^(3/2),x]
```

output

```
(Sqrt[x]*(-48*b^3*x^3 + 231*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 280*b*x*ArcTanh[Tanh[a + b*x]]^2 + 105*ArcTanh[Tanh[a + b*x]]^3)/(24*b^4*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (35*(b*x - ArcTanh[Tanh[a + b*x]])^3*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(8*b^(9/2))
```

**Rubi [A] (verified)**

Time = 0.41 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2599, 2600, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$$

↓ 2599

$$\frac{7 \int \frac{x^{5/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^{7/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2600

$$\begin{array}{c}
 7 \left( \frac{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \frac{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{3b}}{6b} \right) \\
 \hline
 \frac{b}{2x^{7/2}} \\
 \hline
 b \sqrt{\operatorname{arctanh}(\tanh(a+bx))} \\
 \downarrow 2600 \\
 7 \left( \frac{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{4b} + \frac{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{6b} \right) \\
 \hline
 \frac{2x^{7/2}}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \quad b \\
 \downarrow 2600 \\
 7 \left( \frac{5(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx + \sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{4b} + \frac{x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{4b} \right)}{6b} \right) \\
 \hline
 \frac{2x^{7/2}}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \quad b \\
 \downarrow 2596
 \end{array}$$

$$\frac{\left( \frac{\arctanh\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\arctanh(\tanh(a+bx))}}\right) (bx - \arctanh(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\arctanh(\tanh(a+bx))}}{b} \right) (bx - \arctanh(\tanh(a+bx)))}{4b} + \frac{x^{3/2}\sqrt{\arctanh(\tanh(a+bx))}}{6b} \right)}{b\sqrt{\arctanh(\tanh(a+bx))}}$$

```
input Int[x^(7/2)/ArcTanh[Tanh[a + b*x]]^(3/2), x]
```

```
output (7*((5*((3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]])/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b)*(b*x - ArcTanh[Tanh[a + b*x]]))/(4*b) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/(2*b))*(b*x - ArcTanh[Tanh[a + b*x]]))/(6*b) + (x^(5/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/(3*b)))/b - (2*x^(7/2))/(b*Sqrt[ArcTanh[Tanh[a + b*x]]])
```

**Defintions of rubi rules used**

```
rule 2596 Int[1/(Sqrt[u]*Sqrt[v]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

### Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{x^{\frac{7}{2}}}{3b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{5}{2}}}{4b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{3}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{1}{2}}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{1}{2b} \right)}{2b} \right)}{2b} \right)}{2b}$
default	$\frac{x^{\frac{7}{2}}}{3b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{5}{2}}}{4b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{3}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{1}{2}}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{1}{2b} \right)}{2b} \right)}{2b} \right)}{2b}$

input

```
int(x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```



output

```
1/3*x^(7/2)/b/arctanh(tanh(b*x+a))^(1/2)-7/3*(arctanh(tanh(b*x+a))-b*x)/b*
(1/4*x^(5/2)/b/arctanh(tanh(b*x+a))^(1/2)-5/4*(arctanh(tanh(b*x+a))-b*x)/b
*(1/2*x^(3/2)/b/arctanh(tanh(b*x+a))^(1/2)-3/2*(arctanh(tanh(b*x+a))-b*x)/
b*(-x^(1/2)/b/arctanh(tanh(b*x+a))^(1/2)+1/b^(3/2)*ln(b^(1/2)*x^(1/2)+arct
anh(tanh(b*x+a))^(1/2))))
```

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.16

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{105(a^3bx + a^4)\sqrt{b} \log(2bx - 2\sqrt{bx + a}\sqrt{b}\sqrt{x} + a) + 2(8b^4x^3 - 14a^2bx^2 + 35a^3b^2x + 105a^3b)\sqrt{b}\sqrt{x}}{48(b^6x + ab^5)}$$

input

```
integrate(x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
[1/48*(105*(a^3*b*x + a^4)*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt
t(x) + a) + 2*(8*b^4*x^3 - 14*a*b^3*x^2 + 35*a^2*b^2*x + 105*a^3*b)*sqrt(b
*x + a)*sqrt(x))/(b^6*x + a*b^5), 1/24*(105*(a^3*b*x + a^4)*sqrt(-b)*arcta
n(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (8*b^4*x^3 - 14*a*b^3*x^2 + 35*a^2*b^2
*x + 105*a^3*b)*sqrt(b*x + a)*sqrt(x))/(b^6*x + a*b^5)]
```

### Sympy [F(-1)]

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate(x**(7/2)/atanh(tanh(b*x+a))**(3/2),x)
```

output

```
Timed out
```

**Maxima [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{7/2}}{\operatorname{artanh}(\tanh(bx + a))^{3/2}} dx$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(x^(7/2)/arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.45

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\left( \left( 2x \left( \frac{4x}{b} - \frac{7a}{b^2} \right) + \frac{35a^2}{b^3} \right) x + \frac{105a^3}{b^4} \right) \sqrt{x}}{24 \sqrt{bx + a}} + \frac{35a^3 \log \left( \left| -\sqrt{b}\sqrt{x} + \sqrt{bx + a} \right| \right)}{8b^{9/2}}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/24*((2*x*(4*x/b - 7*a/b^2) + 35*a^2/b^3)*x + 105*a^3/b^4)*sqrt(x)/sqrt(b*x + a) + 35/8*a^3*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(9/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{7/2}}{\operatorname{atanh}(\tanh(a + bx))^{3/2}} dx$$

input `int(x^(7/2)/atanh(tanh(a + b*x))^(3/2),x)`

output `int(x^(7/2)/atanh(tanh(a + b*x))^(3/2), x)`

**Reduce [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^(7/2)/atanh(tanh(b*x+a))^(3/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x**3)/atanh(tanh(a + b*x))**2,x)`

**3.250**  $\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1751
Mathematica [A] (verified)	1752
Rubi [A] (verified)	1752
Maple [A] (verified)	1754
Fricas [A] (verification not implemented)	1755
Sympy [F]	1755
Maxima [F]	1756
Giac [A] (verification not implemented)	1756
Mupad [F(-1)]	1756
Reduce [F]	1757

**Optimal result**

Integrand size = 17, antiderivative size = 128

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{15\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{4b^{7/2}} - \frac{2x^{5/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{5x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b^2} + \frac{15\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{4b^3}$$

output

```
15/4*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))^2/b^(7/2)-2*x^(5/2)/b/arctanh(tanh(b*x+a))^(1/2)+5/2*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)/b^2+15/4*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b^3
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.81

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{\sqrt{x}(8b^2x^2 - 25bx\operatorname{arctanh}(\tanh(a+bx)) + 15\operatorname{arctanh}(\tanh(a+bx))^2)}{4b^3\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{15(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2 \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{4b^{7/2}}$$

input

```
Integrate[x^(5/2)/ArcTanh[Tanh[a + b*x]]^(3/2), x]
```

output

```
-1/4*(Sqrt[x]*(8*b^2*x^2 - 25*b*x*ArcTanh[Tanh[a + b*x]] + 15*ArcTanh[Tanh[a + b*x]]^2))/(b^3*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (15*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(4*b^(7/2))
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2599, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$$

↓ 2599

$$\frac{5 \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^{5/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

↓ 2600

$$\begin{aligned}
 & \frac{5 \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{\frac{b}{2x^{5/2}} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \\
 & \quad \downarrow 2600 \\
 & \frac{5 \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{2b} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{4b} \right)}{\frac{2x^{5/2}}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \\
 & \quad \downarrow 2596 \\
 & \frac{5 \left( \frac{3 \left( \frac{\operatorname{arctanh} \left( \frac{\sqrt{b} \sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{4b} \right)}{\frac{2x^{5/2}}{b \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b}
 \end{aligned}$$

input `Int [x^(5/2)/ArcTanh [Tanh [a + b*x]]^(3/2), x]`

output `(5*((3*((ArcTanh [(Sqrt [b]*Sqrt [x])/Sqrt [ArcTanh [Tanh [a + b*x]]]])*(b*x - ArcTanh [Tanh [a + b*x]]))/b^(3/2) + (Sqrt [x]*Sqrt [ArcTanh [Tanh [a + b*x]]])/b*(b*x - ArcTanh [Tanh [a + b*x]]))/(4*b) + (x^(3/2)*Sqrt [ArcTanh [Tanh [a + b*x]]])/(2*b))/b - (2*x^(5/2))/(b*Sqrt [ArcTanh [Tanh [a + b*x]]])`

Defintions of rubi rules used

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{x^{\frac{5}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{3}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{1}{b} \right)}{2b} \right)}{2b}$
default	$\frac{x^{\frac{5}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{x^{\frac{3}{2}}}{2b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{1}{b} \right)}{2b} \right)}{2b}$

```
input int(x^(5/2)/arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)
```

output  $\frac{1}{2}x^{5/2}/b/\operatorname{arctanh}(\tanh(bx+a))^{1/2}-5/2*(\operatorname{arctanh}(\tanh(bx+a))-bx)/b*(1/2x^{3/2}/b/\operatorname{arctanh}(\tanh(bx+a))^{1/2}-3/2*(\operatorname{arctanh}(\tanh(bx+a))-bx)/b*(-x^{1/2}/b/\operatorname{arctanh}(\tanh(bx+a))^{1/2}+1/b^{3/2}*\ln(b^{1/2}*x^{1/2}+\operatorname{arctanh}(\tanh(bx+a))^{1/2}))$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.34

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \left[ \frac{15(a^2bx+a^3)\sqrt{b}\log(2bx+2\sqrt{bx+a}\sqrt{b}\sqrt{x+a})+2(2b^3x^2-5ab^4)}{8(b^5x+ab^4)} \right]$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output  $[1/8*(15*(a^2*b*x+a^3)*\sqrt{b}*\log(2*b*x+2*\sqrt{b*x+a}*\sqrt{b}*\sqrt{x+a})+2*(2*b^3*x^2-5*a*b^2*x-15*a^2*b)*\sqrt{b*x+a}*\sqrt{x})/(b^5*x+a*b^4), -1/4*(15*(a^2*b*x+a^3)*\sqrt{-b}*\arctan(\sqrt{-b}*\sqrt{x}/\sqrt{b*x+a})-(2*b^3*x^2-5*a*b^2*x-15*a^2*b)*\sqrt{b*x+a}*\sqrt{x})/(b^5*x+a*b^4)]$

### Sympy [F]

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \int \frac{x^{5/2}}{\operatorname{atanh}^{3/2}(\tanh(a+bx))} dx$$

input `integrate(x**(5/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(x**(5/2)/atanh(tanh(a + b*x))**(3/2), x)`



**Maxima [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{5/2}}{\operatorname{artanh}(\tanh(bx + a))^{3/2}} dx$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(x^(5/2)/arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.49

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\left(x\left(\frac{2x}{b} - \frac{5a}{b^2}\right) - \frac{15a^2}{b^3}\right)\sqrt{x}}{4\sqrt{bx + a}} - \frac{15a^2 \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{4b^{7/2}}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `1/4*(x*(2*x/b - 5*a/b^2) - 15*a^2/b^3)*sqrt(x)/sqrt(b*x + a) - 15/4*a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(7/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{5/2}}{\operatorname{atanh}(\tanh(a + bx))^{3/2}} dx$$

input `int(x^(5/2)/atanh(tanh(a + b*x))^(3/2),x)`

output `int(x^(5/2)/atanh(tanh(a + b*x))^(3/2), x)`

**Reduce [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2}{\operatorname{atanh}(\tanh(bx + a))^2} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a))^(3/2),x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x**2)/atanh(tanh(a + b*x))**2,x)`

### 3.251 $\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1758
Mathematica [A] (verified)	1758
Rubi [A] (verified)	1759
Maple [A] (verified)	1761
Fricas [A] (verification not implemented)	1761
Sympy [F]	1762
Maxima [F]	1762
Giac [A] (verification not implemented)	1762
Mupad [F(-1)]	1763
Reduce [F]	1763

#### Optimal result

Integrand size = 17, antiderivative size = 86

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{3\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{5/2}} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{3\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^2}$$

output

```
3*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))/b^(5/2)-2*x^(3/2)/b/arctanh(tanh(b*x+a))^(1/2)+3*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b^2
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.94

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{\sqrt{x}(-2bx + 3\operatorname{arctanh}(\tanh(a+bx)))}{b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))\log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{b^{5/2}}$$

input `Integrate[x^(3/2)/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(Sqrt[x]*(-2*b*x + 3*ArcTanh[Tanh[a + b*x]]))/(b^2*Sqrt[ArcTanh[Tanh[a + b*x]]]) + (3*(b*x - ArcTanh[Tanh[a + b*x]])*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b^(5/2)`

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.05, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2599, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{3 \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{b} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \\
 & \quad \downarrow \text{2600} \\
 & \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a + bx))) \int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{2b} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right)}{b} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \\
 & \quad \downarrow \text{2596} \\
 & \frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a + bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{b} \right)}{b} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}
 \end{aligned}$$

input `Int[x^(3/2)/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output `(3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]/b))/b - (2*x^(3/2))/(b*Sqrt[ArcTanh[Tanh[a + b*x]]])`

### Defintions of rubi rules used

rule 2596 `Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

rule 2600 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]`

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{x^{\frac{3}{2}}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{\ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b^{\frac{3}{2}}} \right)}{b}$
default	$\frac{x^{\frac{3}{2}}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{\ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}{b^{\frac{3}{2}}} \right)}{b}$

input `int(x^(3/2)/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `x^(3/2)/b/arctanh(tanh(b*x+a))^(1/2)-3*(arctanh(tanh(b*x+a))-b*x)/b*(-x^(1/2)/b/arctanh(tanh(b*x+a))^(1/2)+1/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.65

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{3(abx+a^2)\sqrt{b}\log(2bx-2\sqrt{bx+a}\sqrt{b}\sqrt{x}+a)+2(b^2x+3ab)\sqrt{bx+a}}{2(b^4x+ab^3)}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[1/2*(3*(a*b*x + a^2)*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) + 2*(b^2*x + 3*a*b)*sqrt(b*x + a)*sqrt(x))/(b^4*x + a*b^3), (3*(a*b*x + a^2)*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (b^2*x + 3*a*b)*sqrt(b*x + a)*sqrt(x))/(b^4*x + a*b^3)]`

**Sympy [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{\frac{3}{2}}}{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(x**(3/2)/atanh(tanh(b*x+a))**(3/2), x)`

output `Integral(x**(3/2)/atanh(tanh(a + b*x))**(3/2), x)`

**Maxima [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{\frac{3}{2}}}{\operatorname{artanh}(\tanh(bx + a))^{\frac{3}{2}}} dx$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(3/2), x, algorithm="maxima")`

output `integrate(x^(3/2)/arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.56

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\sqrt{x}\left(\frac{x}{b} + \frac{3a}{b^2}\right)}{\sqrt{bx + a}} + \frac{3a \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{b^{\frac{5}{2}}}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(3/2), x, algorithm="giac")`

output `sqrt(x)*(x/b + 3*a/b^2)/sqrt(b*x + a) + 3*a*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{x^{3/2}}{\operatorname{atanh}(\tanh(a + bx))^{3/2}} dx$$

input `int(x^(3/2)/atanh(tanh(a + b*x))^(3/2),x)`output `int(x^(3/2)/atanh(tanh(a + b*x))^(3/2), x)`**Reduce [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{3 \operatorname{atanh}(\tanh(bx + a)) \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))} dx \right) - 2\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) b}$$

input `int(x^(3/2)/atanh(tanh(b*x+a))^(3/2),x)`output `(3*atanh(tanh(a + b*x))*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/atanh(tanh(a + b*x)),x) - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*x)/(atanh(tanh(a + b*x))*b)`



### 3.252 $\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1764
Mathematica [A] (verified)	1764
Rubi [A] (verified)	1765
Maple [A] (verified)	1766
Fricas [A] (verification not implemented)	1767
Sympy [F]	1767
Maxima [F]	1768
Giac [A] (verification not implemented)	1768
Mupad [F(-1)]	1768
Reduce [F]	1769

#### Optimal result

Integrand size = 17, antiderivative size = 52

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}} - \frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

$2*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)})/b^{(3/2)}-2*x^{(1/2)}/b/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = -\frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{2\log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{b^{3/2}}$$

input

`Integrate[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^(3/2),x]`

output

$$\frac{(-2\sqrt{x})/(b\sqrt{\text{ArcTanh}[\text{Tanh}[a + b*x]])} + (2\text{Log}[b\sqrt{x} + \sqrt{b} * \sqrt{\text{ArcTanh}[\text{Tanh}[a + b*x]])})/b^{3/2}}$$

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2599, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\text{arctanh}(\tanh(a + bx))^{3/2}} dx$$

$$\downarrow 2599$$

$$\frac{\int \frac{1}{\sqrt{x}\sqrt{\text{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2\sqrt{x}}{b\sqrt{\text{arctanh}(\tanh(a + bx))}}$$

$$\downarrow 2596$$

$$\frac{2\text{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\text{arctanh}(\tanh(a+bx))}}\right)}{b^{3/2}} - \frac{2\sqrt{x}}{b\sqrt{\text{arctanh}(\tanh(a + bx))}}$$

input

$$\text{Int}[\sqrt{x}/\text{ArcTanh}[\text{Tanh}[a + b*x]]^{3/2}, x]$$

output

$$\frac{(2\text{ArcTanh}[(\sqrt{b}*\sqrt{x})/\sqrt{\text{ArcTanh}[\text{Tanh}[a + b*x]])])}{b^{3/2}} - (2\sqrt{x})/(b\sqrt{\text{ArcTanh}[\text{Tanh}[a + b*x]])}$$

## Definitions of rubi rules used

rule 2596

```
Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

## Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{2\ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{b^{\frac{3}{2}}}$	42
default	$-\frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \frac{2\ln(\sqrt{b}\sqrt{x} + \sqrt{\operatorname{arctanh}(\tanh(bx+a))})}{b^{\frac{3}{2}}}$	42

input

```
int(x^(1/2)/arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)
```

output

```
-2*x^(1/2)/b/arctanh(tanh(b*x+a))^(1/2)+2/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.23

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \left[ \frac{(bx + a)\sqrt{b} \log(2bx + 2\sqrt{bx + a}\sqrt{b}\sqrt{x} + a) - 2\sqrt{bx + a}b\sqrt{x}}{b^3x + ab^2}, \right. \\ \left. - \frac{2\left((bx + a)\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{x}}{\sqrt{bx+a}}\right) + \sqrt{bx + a}b\sqrt{x}\right)}{b^3x + ab^2} \right]$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `[((b*x + a)*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) - 2*sqrt(b*x + a)*b*sqrt(x))/(b^3*x + a*b^2), -2*((b*x + a)*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + sqrt(b*x + a)*b*sqrt(x))/(b^3*x + a*b^2)]`

**Sympy [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(sqrt(x)/atanh(tanh(a + b*x))**(3/2), x)`

**Maxima [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x}}{\operatorname{artanh}(\tanh(bx + a))^{3/2}} dx$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(sqrt(x)/arctanh(tanh(b*x + a))^(3/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2 \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{b^{3/2}} - \frac{2\sqrt{x}}{\sqrt{bx + ab}}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `-2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(3/2) - 2*sqrt(x)/(sqrt(b*x + a)*b)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}(\tanh(a + bx))^{3/2}} dx$$

input `int(x^(1/2)/atanh(tanh(a + b*x))^(3/2),x)`

output `int(x^(1/2)/atanh(tanh(a + b*x))^(3/2), x)`

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\operatorname{atanh}(\tanh(bx + a)) \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x} dx \right) - 2\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a)) b}$$

input `int(x^(1/2)/atanh(tanh(b*x+a))^(3/2),x)`

output `(atanh(tanh(a + b*x))*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x),x) - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*b)`

**3.253**  $\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1770
Mathematica [A] (verified)	1770
Rubi [A] (verified)	1771
Maple [A] (verified)	1771
Fricas [A] (verification not implemented)	1772
Sympy [F]	1772
Maxima [F]	1773
Giac [A] (verification not implemented)	1773
Mupad [B] (verification not implemented)	1773
Reduce [F]	1774

**Optimal result**

Integrand size = 17, antiderivative size = 33

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2\sqrt{x}}{(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

output `-2*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)`

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}(-bx + \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Integrate[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(2*Sqrt[x])/(Sqrt[ArcTanh[Tanh[a + b*x]]]*(-(b*x) + ArcTanh[Tanh[a + b*x]]))`

### Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

↓ 2598

$$-\frac{2\sqrt{x}}{(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

input `Int[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-2*Sqrt[x])/((b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])`

#### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

### Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	29
default	$\frac{2\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	29



input `int(1/x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `2*x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2\sqrt{bx+a}\sqrt{x}}{abx+a^2}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `2*sqrt(b*x + a)*sqrt(x)/(a*b*x + a^2)`

### Sympy [F]

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**(1/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(1/(sqrt(x)*atanh(tanh(a + b*x))**(3/2)), x)`

**Maxima [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{\sqrt{x} \operatorname{artanh}(\tanh(bx + a))^{3/2}} dx$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(x)*arctanh(tanh(b*x + a))^(3/2)), x)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.45

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2\sqrt{x}}{\sqrt{bx + aa}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output `2*sqrt(x)/(sqrt(b*x + a)*a)`

**Mupad [B] (verification not implemented)**

Time = 3.68 (sec) , antiderivative size = 163, normalized size of antiderivative = 4.94

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{4x \sqrt{\frac{\ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2}}}{\left(\frac{\sqrt{x} \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right)}{2b} - \frac{\sqrt{x} \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2b}\right) \left(b \ln\left(\frac{1}{e^{2a} e^{2bx} + 1}\right) - b \ln\left(\frac{e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)\right)}$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))^(3/2)),x)`

output

```
(4*x*(log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(1/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/((x^(1/2)*log(1/(exp(2*a)*exp(2*b*x) + 1)))/(2*b) - (x^(1/2)*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/(2*b))*(b*log(1/(exp(2*a)*exp(2*b*x) + 1)) - b*log((exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b^2*x))
```

**Reduce [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x} dx$$

input

```
int(1/x^(1/2)/atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**2*x),x)
```

### 3.254 $\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1775
Mathematica [A] (verified)	1775
Rubi [A] (verified)	1776
Maple [A] (verified)	1777
Fricas [A] (verification not implemented)	1777
Sympy [F]	1778
Maxima [A] (verification not implemented)	1778
Giac [A] (verification not implemented)	1778
Mupad [B] (verification not implemented)	1779
Reduce [F]	1779

#### Optimal result

Integrand size = 17, antiderivative size = 68

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$-\frac{4b\sqrt{x}}{(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-4*b*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(1/2)+2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.63

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$-\frac{2(bx + \operatorname{arctanh}(\tanh(a+bx)))}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))} (-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$

input `Integrate[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-2*(b*x + ArcTanh[Tanh[a + b*x]]))/(Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]]*  
(-(b*x) + ArcTanh[Tanh[a + b*x]])^2)`

### Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

↓ 2602

$$\frac{2b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

↓ 2598

$$\frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

input `Int[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-4*b*Sqrt[x])/((b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]]) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])`

## Definitions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

## Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.87

method	result
derivativedivides	$-\frac{2}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{4b\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$
default	$-\frac{2}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{4b\sqrt{x}}{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$

input `int(1/x^(3/2)/arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)`

output `-2/x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-4*b/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2)`

## Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.50

$$\int \frac{1}{x^{3/2}\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2(2bx + a)\sqrt{bx + a}\sqrt{x}}{a^2bx^2 + a^3x}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(3/2), x, algorithm="fricas")`

output `-2*(2*b*x + a)*sqrt(b*x + a)*sqrt(x)/(a^2*b*x^2 + a^3*x)`

### Sympy [F]

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^{3/2} \operatorname{atanh}^{3/2}(\tanh(a + bx))} dx$$

input `integrate(1/x**(3/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(1/(x**(3/2)*atanh(tanh(a + b*x))**(3/2)), x)`

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.47

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2(2b^2x^2 + 3abx + a^2)}{(bx + a)^{3/2} a^2 \sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `-2*(2*b^2*x^2 + 3*a*b*x + a^2)/((b*x + a)^(3/2)*a^2*sqrt(x))`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2b\sqrt{x}}{\sqrt{bx + aa^2}} + \frac{4\sqrt{b}}{\left(\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2 - a\right)a}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`

output

$$-2*b*\sqrt{x}/(\sqrt{b*x + a})*a^2 + 4*\sqrt{b}/(((\sqrt{b})*\sqrt{x} - \sqrt{b*x + a})^2 - a)*a)$$
**Mupad [B] (verification not implemented)**

Time = 3.44 (sec) , antiderivative size = 281, normalized size of antiderivative = 4.13

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}} \left( \frac{16x}{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2} - \frac{8 \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - 8 \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 16bx}{2b \left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)} \right)}{x^{3/2} - \frac{\sqrt{x} \left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{2b}}$$

input

$$\text{int}(1/(x^{(3/2)}*\operatorname{atanh}(\tanh(a + b*x))^{(3/2)}), x)$$

output

$$-\left(\frac{\log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right)}{2} - \log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right)\right)^{(1/2)}*\left(\frac{16*x}{\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)^2} - \frac{8*\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - 8*\log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 16*b*x}{2*b*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right)}\right)/(x^{(3/2)} - (x^{(1/2)}*\left(\log\left(\frac{2}{\exp(2*a)*\exp(2*b*x) + 1}\right) - \log\left(\frac{2*\exp(2*a)*\exp(2*b*x)}{\exp(2*a)*\exp(2*b*x) + 1}\right) + 2*b*x\right))/(2*b))$$
**Reduce [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x^2} dx$$

input

$$\text{int}(1/x^{(3/2)}/\operatorname{atanh}(\tanh(b*x+a))^{(3/2)}, x)$$

output

$$\text{int}((\sqrt{x}*\sqrt{\operatorname{atanh}(\tanh(a + b*x))})/(\operatorname{atanh}(\tanh(a + b*x))^{**2*x**2}), x)$$



### 3.255 $\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1780
Mathematica [A] (verified)	1781
Rubi [A] (verified)	1781
Maple [A] (verified)	1783
Fricas [A] (verification not implemented)	1783
Sympy [F]	1784
Maxima [A] (verification not implemented)	1784
Giac [A] (verification not implemented)	1784
Mupad [B] (verification not implemented)	1785
Reduce [F]	1785

#### Optimal result

Integrand size = 17, antiderivative size = 110

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$-\frac{16b^2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{8b}$$

$$+ \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2}$$

output

```
-16/3*b^2*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(1/2)+
8/3*b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(1/2)+2/3/
x^(3/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.58

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2(-3b^2x^2 - 6bx \operatorname{arctanh}(\tanh(a+bx)) + \operatorname{arctanh}(\tanh(a+bx))^2)}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

input `Integrate[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`output `(2*(-3*b^2*x^2 - 6*b*x*ArcTanh[Tanh[a + b*x]] + ArcTanh[Tanh[a + b*x]]^2)) / (3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])^3*Sqrt[ArcTanh[Tanh[a + b*x]]])`**Rubi [A] (verified)**Time = 0.30 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.13, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$$

$$\downarrow 2602$$

$$\frac{4b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} +$$

$$\frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$\downarrow 2602$$

$$4b \left( \frac{2b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) +$$

$$\frac{2}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$$\frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$\downarrow 2598$$

$$\frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))} + 4b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{4b\sqrt{x}}{(bx - \operatorname{arctanh}(\tanh(a + bx)))^2\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} \right)}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

input `Int[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(4*b*((-4*b*Sqrt[x])/((b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]]) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))) / (3*(b*x - ArcTanh[Tanh[a + b*x]])) + 2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v))] Int[u^(m + 1)*v^n, x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

**Maple [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.95

method	result
derivativedivides	$-\frac{2}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{8b\left(-\frac{1}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}\right)}{3(\operatorname{arctanh}(\tanh(bx+a)))}$
default	$-\frac{2}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{8b\left(-\frac{1}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}\right)}{3(\operatorname{arctanh}(\tanh(bx+a)))}$

input `int(1/x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

output `-2/3/x^(3/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-8/3*b/(arctanh(tanh(b*x+a))-b*x)*(-1/x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-2*b/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.45

$$\int \frac{1}{x^{5/2}\operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx = \frac{2(8b^2x^2 + 4abx - a^2)\sqrt{bx+a}\sqrt{x}}{3(a^3bx^3 + a^4x^2)}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `2/3*(8*b^2*x^2 + 4*a*b*x - a^2)*sqrt(b*x + a)*sqrt(x)/(a^3*b*x^3 + a^4*x^2)`

**Sympy [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{1}{x^{5/2} \operatorname{atanh}^{\frac{3}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**(5/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Integral(1/(x**(5/2)*atanh(tanh(a + b*x))**(3/2)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.41

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(8b^3x^3 + 12ab^2x^2 + 3a^2bx - a^3)}{3(bx + a)^{\frac{3}{2}}a^3x^{\frac{3}{2}}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `2/3*(8*b^3*x^3 + 12*a*b^2*x^2 + 3*a^2*b*x - a^3)/((b*x + a)^(3/2)*a^3*x^(3/2))`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.97

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2b^2\sqrt{x}}{\sqrt{bx + a}a^3} - \frac{4\left(3b^{\frac{3}{2}}\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^4 - 12ab^{\frac{3}{2}}\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2 + 5a^2b^{\frac{3}{2}}\right)}{3\left(\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2 - a\right)^3 a^2}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`



**3.256**  $\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx$

Optimal result	1786
Mathematica [A] (verified)	1787
Rubi [A] (verified)	1787
Maple [A] (verified)	1789
Fricas [A] (verification not implemented)	1790
Sympy [F(-1)]	1790
Maxima [A] (verification not implemented)	1790
Giac [A] (verification not implemented)	1791
Mupad [B] (verification not implemented)	1791
Reduce [F]	1792

**Optimal result**

Integrand size = 17, antiderivative size = 148

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx =$$

$$-\frac{32b^3 \sqrt{x}}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))^4 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{16b^2}{5\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{4b}{5x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

$$+ \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-32/5*b^3*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^4/arctanh(tanh(b*x+a))^(1/2)+
16/5*b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(1/2)+
/5*b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(1/2)+2/5/x
^(5/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.54

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{2(5b^3x^3 + 15b^2x^2 \operatorname{arctanh}(\tanh(a + bx)) - 5bx \operatorname{arctanh}(\tanh(a + bx))^2 + \operatorname{arctanh}(\tanh(a + bx))^3)}{5x^{5/2} \sqrt{\operatorname{arctanh}(\tanh(a + bx))} (-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input `Integrate[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(-2*(5*b^3*x^3 + 15*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 5*b*x*ArcTanh[Tanh[a + b*x]]^2 + ArcTanh[Tanh[a + b*x]]^3)/(5*x^(5/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)`

**Rubi [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.22, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx$$

$$\downarrow \text{2602}$$

$$\frac{6b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{5(bx - \operatorname{arctanh}(\tanh(a + bx)))} +$$

$$\frac{1}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \sqrt{\operatorname{arctanh}(\tanh(a + bx))}}$$

$$\downarrow \text{2602}$$



$$6b \left( \frac{4b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right) +$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2}$$

$$\frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2}$$

↓ 2602

$$6b \left( \frac{4b \left( \frac{\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) +$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2}$$

$$\frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2}$$

↓ 2598

$$6b \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{4b \left( \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2}{(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right) +$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2}$$

$$\frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx)))\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}{2}$$

input `Int [1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^(3/2)),x]`

output `(6*b*((4*b*((-4*b*Sqrt[x])/((b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]])) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])))/(3*(b*x - ArcTanh[Tanh[a + b*x]])) + 2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(5*(b*x - ArcTanh[Tanh[a + b*x]])) + 2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])*Sqrt[ArcTanh[Tanh[a + b*x]]])`

Defintions of rubi rules used

```
rule 2598 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 2602 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2))/((m + 1)*(b*u - a*v))] Int[u^(m + 1)*v^n, x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

method	result
derivativedivides	$-\frac{2}{5x^{\frac{5}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{12b}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$
default	$-\frac{2}{5x^{\frac{5}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}} - \frac{12b}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$

```
input int(1/x^(7/2)/arctanh(tanh(b*x+a))^(3/2), x, method=_RETURNVERBOSE)
```

```
output -2/5/x^(5/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-12/5*b/(arctanh(tanh(b*x+a))-b*x)*(-1/3/x^(3/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-4/3*b/(arctanh(tanh(b*x+a))-b*x)*(-1/x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(1/2)-2*b/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.39

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2(16b^3x^3 + 8ab^2x^2 - 2a^2bx + a^3)\sqrt{bx+a}\sqrt{x}}{5(a^4bx^4 + a^5x^3)}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="fricas")`

output `-2/5*(16*b^3*x^3 + 8*a*b^2*x^2 - 2*a^2*b*x + a^3)*sqrt(b*x + a)*sqrt(x)/(a^4*b*x^4 + a^5*x^3)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \text{Timed out}$$

input `integrate(1/x**(7/2)/atanh(tanh(b*x+a))**(3/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.36

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2(16b^4x^4 + 24ab^3x^3 + 6a^2b^2x^2 - a^3bx + a^4)}{5(bx+a)^{3/2}a^4x^{5/2}}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="maxima")`

output `-2/5*(16*b^4*x^4 + 24*a*b^3*x^3 + 6*a^2*b^2*x^2 - a^3*b*x + a^4)/((b*x + a)^(3/2)*a^4*x^(5/2))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.09

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = -\frac{2b^3\sqrt{x}}{\sqrt{bx+a}a^4} + \frac{4\left(5b^{5/2}\left(\sqrt{b}\sqrt{x}-\sqrt{bx+a}\right)^8 - 30ab^{5/2}\left(\sqrt{b}\sqrt{x}-\sqrt{bx+a}\right)^6 + 80a^2b^{5/2}\left(\sqrt{b}\sqrt{x}-\sqrt{bx+a}\right)^4 - 50a^3b^{5/2}\left(\sqrt{b}\sqrt{x}-\sqrt{bx+a}\right)^2 + 11a^4b^{5/2}\right)}{5\left(\left(\sqrt{b}\sqrt{x}-\sqrt{bx+a}\right)^2 - a\right)^5 a^3}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(3/2),x, algorithm="giac")`output `-2*b^3*sqrt(x)/(sqrt(b*x + a)*a^4) + 4/5*(5*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^8 - 30*a*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 + 80*a^2*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 - 50*a^3*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 + 11*a^4*b^(5/2))/(((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^5*a^3)`**Mupad [B] (verification not implemented)**

Time = 3.75 (sec) , antiderivative size = 346, normalized size of antiderivative = 2.34

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)}{2}}}{5\left(\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx\right)^2} \left(\frac{16x}{5\left(\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx\right)^2}\right)$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))^(3/2)),x)`

output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2)^(1/2)*((16*x)/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + 4/(5*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (128*b*x^2)/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (512*b^2*x^3)/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)))/(x^(7/2) - (x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b))
```

**Reduce [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^2 x^4} dx$$

input

```
int(1/x^(7/2)/atanh(tanh(b*x+a))^(3/2),x)
```

output

```
int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**2*x**4),x)
```

**3.257**  $\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1793
Mathematica [A] (verified)	1794
Rubi [A] (verified)	1794
Maple [A] (verified)	1797
Fricas [A] (verification not implemented)	1798
Sympy [F(-1)]	1798
Maxima [F]	1799
Giac [A] (verification not implemented)	1799
Mupad [F(-1)]	1799
Reduce [F]	1800

**Optimal result**

Integrand size = 17, antiderivative size = 153

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{35\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))^2}{4b^{9/2}} - \frac{2x^{7/2}}{3\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{14x^{5/2}}{3b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{35x^{3/2}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{6b^3} + \frac{35\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{4b^4}$$

output

```
35/4*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh
(b*x+a)))^2/b^(9/2)-2/3*x^(7/2)/b/arctanh(tanh(b*x+a))^(3/2)-14/3*x^(5/2)/
b^2/arctanh(tanh(b*x+a))^(1/2)+35/6*x^(3/2)*arctanh(tanh(b*x+a))^(1/2)/b^3
+35/4*x^(1/2)*(b*x-arctanh(tanh(b*x+a)))*arctanh(tanh(b*x+a))^(1/2)/b^4
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.79

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx =$$

$$\frac{\sqrt{x}(8b^3x^3 + 56b^2x^2\operatorname{arctanh}(\tanh(a + bx)) - 175bx\operatorname{arctanh}(\tanh(a + bx))^2 + 105\operatorname{arctanh}(\tanh(a + bx)))}{12b^4\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

$$+ \frac{35(-bx + \operatorname{arctanh}(\tanh(a + bx)))^2 \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)}{4b^{9/2}}$$

input

```
Integrate[x^(7/2)/ArcTanh[Tanh[a + b*x]]^(5/2),x]
```

output

```
-1/12*(Sqrt[x]*(8*b^3*x^3 + 56*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 175*b*x*ArcTanh[Tanh[a + b*x]]^2 + 105*ArcTanh[Tanh[a + b*x]]^3)/(b^4*ArcTanh[Tanh[a + b*x]]^(3/2)) + (35*(-(b*x) + ArcTanh[Tanh[a + b*x]])^2*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/(4*b^(9/2))
```

**Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2599, 2599, 2600, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

$$\downarrow 2599$$

$$\frac{7 \int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{3b} - \frac{2x^{7/2}}{3b\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

$$\downarrow 2599$$

$$\begin{aligned}
 & \frac{7 \left( \frac{5 \int \frac{x^{3/2}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^{5/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^{7/2}}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2600 \\
 & \frac{7 \left( \frac{5 \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{b} - \frac{2x^{5/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{2x^{7/2} \cdot 3b} - \frac{2x^{7/2}}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2600 \\
 & \frac{7 \left( \frac{5 \left( \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{2b} + \frac{\sqrt{x} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{4b} + \frac{x^{3/2} \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{2b} \right)}{b} - \frac{2x^{5/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{2x^{7/2} \cdot 3b} - \frac{2x^{7/2}}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2596
 \end{aligned}$$



$$\frac{\left( \frac{\arctanh\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\arctanh(\tanh(a+bx))}}\right)(bx - \arctanh(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\arctanh(\tanh(a+bx))}}{b} \right)(bx - \arctanh(\tanh(a+bx)))}{4b} + \frac{x^{3/2}\sqrt{\arctanh(\tanh(a+bx))}}{b} \right)}{2x^{7/2}} \cdot \frac{1}{3b \arctanh(\tanh(a+bx))^{3/2}}$$

```
input Int [x^(7/2)/ArcTanh[Tanh[a + b*x]]^(5/2), x]
```

```
output (7*((5*((3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b)*(b*x - ArcTanh[Tanh[a + b*x]]))/(4*b) + (x^(3/2)*Sqrt[ArcTanh[Tanh[a + b*x]]])/(2*b)))/b - (2*x^(5/2))/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(3*b) - (2*x^(7/2))/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2))
```

**Defintions of rubi rules used**

```
rule 2596 Int [1/(Sqrt[u]*Sqrt[v]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

rule 2600

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

### Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{x^{\frac{7}{2}}}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} \left( \frac{x^{\frac{5}{2}}}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}{3} \right)$
default	$\frac{x^{\frac{7}{2}}}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{7(\operatorname{arctanh}(\tanh(bx+a))-bx)}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} \left( \frac{x^{\frac{5}{2}}}{2b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx)}{3} \right)$

input

```
int(x^(7/2)/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
1/2*x^(7/2)/b/arctanh(tanh(b*x+a))^(3/2)-7/2*(arctanh(tanh(b*x+a))-b*x)/b*
(1/2*x^(5/2)/b/arctanh(tanh(b*x+a))^(3/2)-5/2*(arctanh(tanh(b*x+a))-b*x)/b
*(-1/3*x^(3/2)/b/arctanh(tanh(b*x+a))^(3/2)+1/b*(-x^(1/2)/b/arctanh(tanh(b
*x+a))^(1/2)+1/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2))))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.56

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \left[ \frac{105(a^2 b^2 x^2 + 2a^3 bx + a^4) \sqrt{b} \log(2bx + 2\sqrt{bx + a} \sqrt{b} \sqrt{x + a}) + 2(6b^4 x^3 - 21a b^3 x^2 - 140a^2 b^2 x - 105a^3 b) \sqrt{b} \sqrt{x + a}}{24(b^7 x^2 + 2ab^6 x + a^2 b^5)} \right]$$

input

```
integrate(x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
[1/24*(105*(a^2*b^2*x^2 + 2*a^3*b*x + a^4)*sqrt(b)*log(2*b*x + 2*sqrt(b*x
+ a)*sqrt(b)*sqrt(x) + a) + 2*(6*b^4*x^3 - 21*a*b^3*x^2 - 140*a^2*b^2*x -
105*a^3*b)*sqrt(b*x + a)*sqrt(x))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5), -1/12*(
105*(a^2*b^2*x^2 + 2*a^3*b*x + a^4)*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(
b*x + a)) - (6*b^4*x^3 - 21*a*b^3*x^2 - 140*a^2*b^2*x - 105*a^3*b)*sqrt(b*
x + a)*sqrt(x))/(b^7*x^2 + 2*a*b^6*x + a^2*b^5)]
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate(x**(7/2)/atanh(tanh(b*x+a))**(5/2),x)
```

output

```
Timed out
```

**Maxima [F]**

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{7/2}}{\operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(x^(7/2)/arctanh(tanh(b*x + a))^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.49

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\left( \left( 3x \left( \frac{2x}{b} - \frac{7a}{b^2} \right) - \frac{140a^2}{b^3} \right) x - \frac{105a^3}{b^4} \right) \sqrt{x}}{12 (bx + a)^{3/2}} - \frac{35a^2 \log \left( \left| -\sqrt{b}\sqrt{x} + \sqrt{bx + a} \right| \right)}{4b^{9/2}}$$

input `integrate(x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/12*((3*x*(2*x/b - 7*a/b^2) - 140*a^2/b^3)*x - 105*a^3/b^4)*sqrt(x)/(b*x + a)^(3/2) - 35/4*a^2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(9/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{7/2}}{\operatorname{atanh}(\tanh(a + bx))^{5/2}} dx$$

input `int(x^(7/2)/atanh(tanh(a + b*x))^(5/2),x)`

output `int(x^(7/2)/atanh(tanh(a + b*x))^(5/2), x)`

### Reduce [F]

$$\int \frac{x^{7/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^3}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^(7/2)/atanh(tanh(b*x+a))^(5/2), x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x**3)/atanh(tanh(a + b*x))**3,x)`

### 3.258 $\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1801
Mathematica [A] (verified)	1802
Rubi [A] (verified)	1802
Maple [A] (verified)	1804
Fricas [A] (verification not implemented)	1805
Sympy [F]	1805
Maxima [F]	1806
Giac [A] (verification not implemented)	1806
Mupad [F(-1)]	1806
Reduce [F]	1807

#### Optimal result

Integrand size = 17, antiderivative size = 111

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{5 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right) (bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{7/2}} - \frac{2x^{5/2}}{3b \operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{10x^{3/2}}{3b^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{5\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b^3}$$

output

```
5*arctanh(b^(1/2)*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))*(b*x-arctanh(tanh(b*x+a)))/b^(7/2)-2/3*x^(5/2)/b/arctanh(tanh(b*x+a))^(3/2)-10/3*x^(3/2)/b^2/arctanh(tanh(b*x+a))^(1/2)+5*x^(1/2)*arctanh(tanh(b*x+a))^(1/2)/b^3
```

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.91

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\sqrt{x}(2b^2x^2 + 10bx\operatorname{arctanh}(\tanh(a + bx)) - 15\operatorname{arctanh}(\tanh(a + bx))^2)}{3b^3\operatorname{arctanh}(\tanh(a + bx))^{3/2}} + \frac{5(bx - \operatorname{arctanh}(\tanh(a + bx))) \log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}\right)}{b^{7/2}}$$

input

```
Integrate[x^(5/2)/ArcTanh[Tanh[a + b*x]]^(5/2),x]
```

output

```
-1/3*(Sqrt[x]*(2*b^2*x^2 + 10*b*x*ArcTanh[Tanh[a + b*x]] - 15*ArcTanh[Tanh[a + b*x]]^2))/(b^3*ArcTanh[Tanh[a + b*x]]^(3/2)) + (5*(b*x - ArcTanh[Tanh[a + b*x]])*Log[b*Sqrt[x] + Sqrt[b]*Sqrt[ArcTanh[Tanh[a + b*x]]]])/b^(7/2)
```

**Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2599, 2599, 2600, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2599

$$\frac{5 \int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{3b} - \frac{2x^{5/2}}{3b\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2599

$$\begin{aligned}
 & \frac{5 \left( \frac{3 \int \frac{\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{b} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b} - \frac{2x^{5/2}}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2600 \\
 & \frac{5 \left( \frac{3 \left( \frac{(bx - \operatorname{arctanh}(\tanh(a+bx))) \int \frac{1}{2b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} dx}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{b} \right)}{3b} - \frac{2x^{3/2}}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{2x^{5/2}} - \frac{3b}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}} \\
 & \quad \downarrow 2596 \\
 & \frac{5 \left( \frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)(bx - \operatorname{arctanh}(\tanh(a+bx)))}{b^{3/2}} + \frac{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}{b} \right)}{b} \right)}{2x^{5/2}} - \frac{3b}{b\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} \right)}{3b\operatorname{arctanh}(\tanh(a+bx))^{3/2}}
 \end{aligned}$$

input `Int [x^(5/2)/ArcTanh[Tanh[a + b*x]]^(5/2), x]`

output `(5*((3*((ArcTanh[(Sqrt[b]*Sqrt[x])/Sqrt[ArcTanh[Tanh[a + b*x]]])*(b*x - ArcTanh[Tanh[a + b*x]]))/b^(3/2) + (Sqrt[x]*Sqrt[ArcTanh[Tanh[a + b*x]]])/b)/b - (2*x^(3/2))/(b*Sqrt[ArcTanh[Tanh[a + b*x]]]))/(3*b) - (2*x^(5/2))/(3*b*ArcTanh[Tanh[a + b*x]]^(3/2))`



Defintions of rubi rules used

```
rule 2596 Int[1/(Sqrt[u_]*Sqrt[v_]), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(2/Rt[a*b, 2])*ArcTanh[Rt[a*b, 2]*(Sqrt[u]/(a*Sqrt[v]))]], x] /; NeQ[b*u - a*v, 0] && PosQ[a*b]] /; PiecewiseLinearQ[u, v, x]
```

```
rule 2599 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

```
rule 2600 Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + n + 1))), x] - Simp[n*((b*u - a*v)/(a*(m + n + 1))) Int[u^m*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || LtQ[0, m, n])) && !ILtQ[m + n, -2]
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{x^{\frac{5}{2}}}{b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{x^{\frac{3}{2}}}{3b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{-\frac{\sqrt{x}}{b \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \ln(\dots)}{b} \right)}{b}$
default	$\frac{x^{\frac{5}{2}}}{b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} - \frac{5(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( -\frac{x^{\frac{3}{2}}}{3b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{-\frac{\sqrt{x}}{b \sqrt{\operatorname{arctanh}(\tanh(bx+a))}} + \ln(\dots)}{b} \right)}{b}$

```
input int(x^(5/2)/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
x^(5/2)/b/arctanh(tanh(b*x+a))^(3/2)-5*(arctanh(tanh(b*x+a))-b*x)/b*(-1/3*
x^(3/2)/b/arctanh(tanh(b*x+a))^(3/2)+1/b*(-x^(1/2)/b/arctanh(tanh(b*x+a))^(
1/2)+1/b^(3/2)*ln(b^(1/2)*x^(1/2)+arctanh(tanh(b*x+a))^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.90

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \left[ \frac{15(ab^2x^2 + 2a^2bx + a^3)\sqrt{b} \log(2bx - 2\sqrt{bx+a}\sqrt{b}\sqrt{x+a}) + 2(3b^3x^2 + 20a^2bx + 15a^2b)\sqrt{b}\sqrt{x+a}}{6(b^6x^2 + 2ab^5x + a^2b^4)} \right]$$

input

```
integrate(x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
[1/6*(15*(a*b^2*x^2 + 2*a^2*b*x + a^3)*sqrt(b)*log(2*b*x - 2*sqrt(b*x + a)
*sqrt(b)*sqrt(x) + a) + 2*(3*b^3*x^2 + 20*a*b^2*x + 15*a^2*b)*sqrt(b*x + a)
*sqrt(x))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4), 1/3*(15*(a*b^2*x^2 + 2*a^2*b*x
+ a^3)*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (3*b^3*x^2 + 20*
a*b^2*x + 15*a^2*b)*sqrt(b*x + a)*sqrt(x))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4)
]
```

**Sympy [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{5/2}}{\operatorname{atanh}^{5/2}(\tanh(a + bx))} dx$$

input

```
integrate(x**(5/2)/atanh(tanh(b*x+a))**(5/2),x)
```

output

```
Integral(x**(5/2)/atanh(tanh(a + b*x))**(5/2), x)
```

**Maxima [F]**

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{5/2}}{\operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(x^(5/2)/arctanh(tanh(b*x + a))^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.55

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\left(x\left(\frac{3x}{b} + \frac{20a}{b^2}\right) + \frac{15a^2}{b^3}\right)\sqrt{x}}{3(bx + a)^{3/2}} + \frac{5a \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{b^{7/2}}$$

input `integrate(x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `1/3*(x*(3*x/b + 20*a/b^2) + 15*a^2/b^3)*sqrt(x)/(b*x + a)^(3/2) + 5*a*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(7/2)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{5/2}}{\operatorname{atanh}(\tanh(a + bx))^{5/2}} dx$$

input `int(x^(5/2)/atanh(tanh(a + b*x))^(5/2),x)`

output `int(x^(5/2)/atanh(tanh(a + b*x))^(5/2), x)`

### Reduce [F]

$$\int \frac{x^{5/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))} x^2}{\operatorname{atanh}(\tanh(bx + a))^3} dx$$

input `int(x^(5/2)/atanh(tanh(b*x+a))^(5/2), x)`

output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))*x**2)/atanh(tanh(a + b*x))**3,x)`

**3.259**  $\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1808
Mathematica [A] (verified)	1808
Rubi [A] (verified)	1809
Maple [A] (verified)	1810
Fricas [A] (verification not implemented)	1811
Sympy [F]	1811
Maxima [F]	1811
Giac [A] (verification not implemented)	1812
Mupad [F(-1)]	1812
Reduce [F]	1813

**Optimal result**

Integrand size = 17, antiderivative size = 75

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}\right)}{b^{5/2}} - \frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{2\sqrt{x}}{b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

$2*\operatorname{arctanh}(b^{(1/2)}*x^{(1/2)}/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)})/b^{(5/2)}-2/3*x^{(3/2)}/b/\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}-2*x^{(1/2)}/b^2/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}$

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.04

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a+bx))^{3/2}} - \frac{2\sqrt{x}}{b^2\sqrt{\operatorname{arctanh}(\tanh(a+bx))}} + \frac{2\log\left(b\sqrt{x} + \sqrt{b}\sqrt{\operatorname{arctanh}(\tanh(a+bx))}\right)}{b^{5/2}}$$

input `Integrate[x^(3/2)/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output 
$$\frac{(-2*x^{(3/2)})/(3*b*ArcTanh[Tanh[a + b*x]]^{(3/2)}) - (2*sqrt[x])/(b^2*sqrt[ArcTanh[Tanh[a + b*x]]]) + (2*Log[b*sqrt[x] + sqrt[b]*sqrt[ArcTanh[Tanh[a + b*x]]]])/b^{(5/2)}}{b^{(5/2)}}$$

### Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.07, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2599, 2599, 2596}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2599

$$\frac{\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{b} - \frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2599

$$\frac{\int \frac{1}{\sqrt{x}\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} dx}{b} - \frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2596

$$\frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{\operatorname{arctanh}(\tanh(a + bx))}}\right)}{b^{3/2}} - \frac{2\sqrt{x}}{b\sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input `Int[x^(3/2)/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output 
$$\left(\frac{2 \operatorname{ArcTanh}[\sqrt{b} \sqrt{x}]/\sqrt{\operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]}}{b^{3/2}} - \frac{2 \sqrt{x}}{b \sqrt{\operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]}}\right) / b - \frac{(2 x^{3/2})}{(3 b \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{3/2})}$$

**Defintions of rubi rules used**

rule 2596 
$$\operatorname{Int}[1/(\sqrt{u} \sqrt{v}), x\_Symbol] \rightarrow \operatorname{With}\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[(2/\operatorname{Rt}[a*b, 2]) \operatorname{ArcTanh}[\operatorname{Rt}[a*b, 2] * (\sqrt{u}/(a \sqrt{v}))]], x] /; \operatorname{NeQ}[b*u - a*v, 0] \&\& \operatorname{PosQ}[a*b] /; \operatorname{PiecewiseLinearQ}[u, v, x]$$

rule 2599 
$$\operatorname{Int}[(u)^{(m)} * (v)^{(n)}, x\_Symbol] \rightarrow \operatorname{With}\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)} * (v^n / (a * (m+1))), x] - \operatorname{Simp}[b * (n / (a * (m+1))) \operatorname{Int}[u^{(m+1)} * v^{(n-1)}, x], x] /; \operatorname{NeQ}[b*u - a*v, 0] /; \operatorname{FreeQ}\{m, n, x\} \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \&\& \operatorname{NeQ}[m, -1] \&\& ((\operatorname{LtQ}[m, -1] \&\& \operatorname{GtQ}[n, 0]) \&\& !(\operatorname{ILtQ}[m+n, -2] \&\& (\operatorname{FractionQ}[m] || \operatorname{GeQ}[2*n+m+1, 0]))) || (\operatorname{IGtQ}[n, 0] \&\& \operatorname{IGtQ}[m, 0] \&\& \operatorname{LeQ}[n, m]) || (\operatorname{IGtQ}[n, 0] \&\& !\operatorname{IntegerQ}[m]) || (\operatorname{ILtQ}[m, 0] \&\& !\operatorname{IntegerQ}[n]))$$

**Maple [A] (verified)**

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$-\frac{2x^{3/2}}{3b \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{3/2}} - \frac{2\sqrt{x}}{b^2 \sqrt{\operatorname{arctanh}(\operatorname{tanh}(bx+a))}} + \frac{2 \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\operatorname{tanh}(bx+a))})}{b^{5/2}}$	59
default	$-\frac{2x^{3/2}}{3b \operatorname{arctanh}(\operatorname{tanh}(bx+a))^{3/2}} - \frac{2\sqrt{x}}{b^2 \sqrt{\operatorname{arctanh}(\operatorname{tanh}(bx+a))}} + \frac{2 \ln(\sqrt{b} \sqrt{x} + \sqrt{\operatorname{arctanh}(\operatorname{tanh}(bx+a))})}{b^{5/2}}$	59

input 
$$\operatorname{int}(x^{3/2}/\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{5/2}, x, \operatorname{method}=\_RETURNVERBOSE)$$

output 
$$-2/3*x^{3/2}/b/\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{3/2}-2*x^{1/2}/b^2/\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{1/2}+2/b^{5/2}*\ln(b^{1/2}*x^{1/2}+\operatorname{arctanh}(\operatorname{tanh}(b*x+a))^{1/2})$$

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.44

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \left[ \frac{3(b^2x^2 + 2abx + a^2)\sqrt{b} \log(2bx + 2\sqrt{bx+a}\sqrt{b}\sqrt{x+a}) - 2(4b^2x + 3ab^2)}{3(b^5x^2 + 2ab^4x + a^2b^3)} \right]$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `[1/3*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(b)*log(2*b*x + 2*sqrt(b*x + a)*sqrt(b)*sqrt(x) + a) - 2*(4*b^2*x + 3*a*b)*sqrt(b*x + a)*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3), -2/3*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-b)*arctan(sqrt(-b)*sqrt(x)/sqrt(b*x + a)) + (4*b^2*x + 3*a*b)*sqrt(b*x + a)*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)]`

**Sympy [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{3/2}}{\operatorname{atanh}^{5/2}(\tanh(a + bx))} dx$$

input `integrate(x**(3/2)/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(x**(3/2)/atanh(tanh(a + b*x))**(5/2), x)`

**Maxima [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{3/2}}{\operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`



output `integrate(x^(3/2)/arctanh(tanh(b*x + a))^(5/2), x)`

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.65

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2\sqrt{x}\left(\frac{4x}{b} + \frac{3a}{b^2}\right)}{3(bx + a)^{3/2}} - \frac{2 \log\left(\left|-\sqrt{b}\sqrt{x} + \sqrt{bx + a}\right|\right)}{b^{5/2}}$$

input `integrate(x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-2/3*sqrt(x)*(4*x/b + 3*a/b^2)/(b*x + a)^(3/2) - 2*log(abs(-sqrt(b)*sqrt(x) + sqrt(b*x + a)))/b^(5/2)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{x^{3/2}}{\operatorname{atanh}(\tanh(a + bx))^{5/2}} dx$$

input `int(x^(3/2)/atanh(tanh(a + b*x))^(5/2),x)`

output `int(x^(3/2)/atanh(tanh(a + b*x))^(5/2), x)`

**Reduce [F]**

$$\int \frac{x^{3/2}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{3 \operatorname{atanh}(\tanh(bx + a))^2 \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))x} dx \right) - 6\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{3 \operatorname{atanh}(\tanh(bx + a))}$$

input `int(x^(3/2)/atanh(tanh(b*x+a))^(5/2),x)`

output `(3*atanh(tanh(a + b*x))**2*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))*x),x) - 6*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*atanh(tanh(a + b*x)) - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x)))*b*x)/(3*atanh(tanh(a + b*x))**2*b**2)`

**3.260**       $\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1814
Mathematica [A] (verified)	1814
Rubi [A] (verified)	1815
Maple [B] (verified)	1815
Fricas [A] (verification not implemented)	1816
Sympy [F]	1816
Maxima [F]	1817
Giac [A] (verification not implemented)	1817
Mupad [B] (verification not implemented)	1817
Reduce [F]	1818

**Optimal result**

Integrand size = 17, antiderivative size = 35

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2x^{3/2}}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

output

```
-2/3*x^(3/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2x^{3/2}}{3\operatorname{arctanh}(\tanh(a + bx))^{3/2}(-bx + \operatorname{arctanh}(\tanh(a + bx)))}$$

input

```
Integrate[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^(5/2),x]
```

output

```
(2*x^(3/2))/(3*ArcTanh[Tanh[a + b*x]]^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]]))
```

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$ , Rules used = {2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2598

$$\frac{2x^{3/2}}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))\operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input `Int[Sqrt[x]/ArcTanh[Tanh[a + b*x]]^(5/2),x]`

output `(-2*x^(3/2))/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))`

**Defintions of rubi rules used**

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 91 vs.  $2(29) = 58$ .

Time = 0.40 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.63

method	result
derivativedivides	$-\frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{1}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)} \right)}{b}$
default	$-\frac{\sqrt{x}}{b \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx) \left( \frac{\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{1}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)} \right)}{b}$

input `int(x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `-x^(1/2)/b/arctanh(tanh(b*x+a))^(3/2)+(arctanh(tanh(b*x+a))-b*x)/b*(1/3*x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)+2/3/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))`

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2\sqrt{bx+ax^{\frac{3}{2}}}}{3(ab^2x^2+2a^2bx+a^3)}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `2/3*sqrt(b*x + a)*x^(3/2)/(a*b^2*x^2 + 2*a^2*b*x + a^3)`

### Sympy [F]

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \int \frac{\sqrt{x}}{\operatorname{atanh}^{\frac{5}{2}}(\tanh(a+bx))} dx$$

input `integrate(x**(1/2)/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(sqrt(x)/atanh(tanh(a + b*x))**(5/2), x)`

**Maxima [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x}}{\operatorname{artanh}(\tanh(bx + a))^{5/2}} dx$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(x)/arctanh(tanh(b*x + a))^(5/2), x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.43

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2x^{3/2}}{3(bx + a)^{3/2}a}$$

input `integrate(x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `2/3*x^(3/2)/((b*x + a)^(3/2)*a)`

**Mupad [B] (verification not implemented)**

Time = 3.83 (sec) , antiderivative size = 229, normalized size of antiderivative = 6.54

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx =$$

$$\frac{4x^{3/2} \sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{3b^2 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right) \left( \frac{\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2}{4b^2} + x^2 - \frac{x \left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)\right)}{2} \right)}$$

input `int(x^(1/2)/atanh(tanh(a + b*x))^(5/2),x)`

output

```

-(4*x^(3/2)*(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2))/(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(4*b^2) + x^2 - (x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)/b))

```

**Reduce [F]**

$$\int \frac{\sqrt{x}}{\operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\operatorname{atanh}(\tanh(bx + a))^2 \left( \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx+a))}}{\operatorname{atanh}(\tanh(bx+a))^2 x} dx \right) - 2\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{3\operatorname{atanh}(\tanh(bx + a))^2 b}$$

input

```
int(x^(1/2)/atanh(tanh(b*x+a))^(5/2),x)
```

output

```

(atanh(tanh(a + b*x))**2*int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**2*x),x) - 2*sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(3*atanh(tanh(a + b*x))**2*b)

```

### 3.261 $\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1819
Mathematica [A] (verified)	1819
Rubi [A] (verified)	1820
Maple [A] (verified)	1821
Fricas [A] (verification not implemented)	1821
Sympy [F]	1822
Maxima [F]	1822
Giac [A] (verification not implemented)	1822
Mupad [B] (verification not implemented)	1823
Reduce [F]	1823

#### Optimal result

Integrand size = 17, antiderivative size = 71

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} + \frac{4\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

$$-2/3*x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))/\operatorname{arctanh}(\tanh(b*x+a))^{(3/2)}+4/3*x^{(1/2)}/(b*x-\operatorname{arctanh}(\tanh(b*x+a)))^2/\operatorname{arctanh}(\tanh(b*x+a))^{(1/2)}$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.66

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2\sqrt{x}(bx - 3\operatorname{arctanh}(\tanh(a+bx)))}{3\operatorname{arctanh}(\tanh(a+bx))^{3/2}(-bx + \operatorname{arctanh}(\tanh(a+bx)))^2}$$



input `Integrate[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(-2*Sqrt[x]*(b*x - 3*ArcTanh[Tanh[a + b*x]]))/(3*ArcTanh[Tanh[a + b*x]]^(3/2)*(-(b*x) + ArcTanh[Tanh[a + b*x]]^2)`

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2602

$$-\frac{2 \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2598

$$\frac{4\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a + bx))}} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input `Int[1/(Sqrt[x]*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(-2*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)) + (4*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]])`

### Defintions of rubi rules used

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

### Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.82

method	result	si
derivativedivides	$\frac{2\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{4\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	5
default	$\frac{2\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx) \operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}} + \frac{4\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 \sqrt{\operatorname{arctanh}(\tanh(bx+a))}}$	5

input

```
int(1/x^(1/2)/arctanh(tanh(b*x+a))^(5/2), x, method=_RETURNVERBOSE)
```

output

```
2/3*x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)+4/3/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2)
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.61

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(2bx + 3a)\sqrt{bx + a}\sqrt{x}}{3(a^2b^2x^2 + 2a^3bx + a^4)}$$

input

```
integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(5/2), x, algorithm="fricas")
```

output  $2/3*(2*b*x + 3*a)*sqrt(b*x + a)*sqrt(x)/(a^2*b^2*x^2 + 2*a^3*b*x + a^4)$

### Sympy [F]

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{\sqrt{x} \operatorname{atanh}^{\frac{5}{2}}(\tanh(a + bx))} dx$$

input `integrate(1/x**(1/2)/atanh(tanh(b*x+a))**(5/2),x)`

output `Integral(1/(sqrt(x)*atanh(tanh(a + b*x))**(5/2)), x)`

### Maxima [F]

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{1}{\sqrt{x} \operatorname{artanh}(\tanh(bx + a))^{\frac{5}{2}}} dx$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(x)*arctanh(tanh(b*x + a))^(5/2)), x)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.35

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2\sqrt{x}\left(\frac{2bx}{a^2} + \frac{3}{a}\right)}{3(bx + a)^{\frac{3}{2}}}$$

input `integrate(1/x^(1/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output  $2/3*sqrt(x)*(2*b*x/a^2 + 3/a)/(b*x + a)^(3/2)$

**Mupad [B] (verification not implemented)**

Time = 3.85 (sec) , antiderivative size = 346, normalized size of antiderivative = 4.87

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{x^{5/2} - \frac{x^{3/2}\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{b} + \sqrt{x}\left(\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) + 2bx\right)^2} \left( \frac{16x^2}{3b\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)} \right)$$

input `int(1/(x^(1/2)*atanh(tanh(a + b*x))^(5/2)),x)`output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*((16*x^2)/(3*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) - (x*(48*log(2/(exp(2*a)*exp(2*b*x) + 1)) - 48*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 96*b*x))/(12*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(5/2) - (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b + (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^2))`**Reduce [F]**

$$\int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x} dx$$

input `int(1/x^(1/2)/atanh(tanh(b*x+a))^(5/2),x)`output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**3*x),x)`

**3.262**  $\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1824
Mathematica [A] (verified)	1825
Rubi [A] (verified)	1825
Maple [A] (verified)	1826
Fricas [A] (verification not implemented)	1827
Sympy [F]	1827
Maxima [A] (verification not implemented)	1828
Giac [A] (verification not implemented)	1828
Mupad [B] (verification not implemented)	1829
Reduce [F]	1829

**Optimal result**

Integrand size = 17, antiderivative size = 106

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx =$$

$$-\frac{8b\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

$$+ \frac{16b\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-8/3*b*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(3/2)+2/x
^(1/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)+16/3*b*x^(1/2)
)/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(-b^2x^2 + 6bx \operatorname{arctanh}(\tanh(a + bx)) + 3 \operatorname{arctanh}(\tanh(a + bx))^2)}{3\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx)))^3 \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input

```
Integrate[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^(5/2)),x]
```

output

```
(2*(-b^2*x^2) + 6*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^2
)/ (3*Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])^3*ArcTanh[Tanh[a + b*x]]^(3/2)
)
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$ , Rules used = {2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

↓ 2602

$$\frac{4b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2602

$$\frac{4b \left( -\frac{2 \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a + bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a + bx)))} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} +$$

↓ 2598

$$\frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

$$4b \left( \frac{\frac{4\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

input `Int[1/(x^(3/2)*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(4*b*((-2*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)) + (4*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]]))) / (b*x - ArcTanh[Tanh[a + b*x]]) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))`

### Defintions of rubi rules used

rule 2598 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 2602 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]`

### Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.98

method	result
derivativedivides	$-\frac{2}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}-\frac{8b\left(\frac{\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}+\frac{1}{3\operatorname{arctanh}(\tanh(bx+a))}}\right)}{\operatorname{arctanh}(\tanh(bx+a))}$
default	$-\frac{2}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}-\frac{8b\left(\frac{\sqrt{x}}{3(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}+\frac{1}{3\operatorname{arctanh}(\tanh(bx+a))}}\right)}{\operatorname{arctanh}(\tanh(bx+a))}$

input `int(1/x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-8*b/(arctanh(tanh(b*x+a))-b*x)*(1/3*x^(1/2)/arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)+2/3/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2)`

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.55

$$\int \frac{1}{x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = -\frac{2(8b^2x^2+12abx+3a^2)\sqrt{bx+a}\sqrt{x}}{3(a^3b^2x^3+2a^4bx^2+a^5x)}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `-2/3*(8*b^2*x^2+12*a*b*x+3*a^2)*sqrt(b*x+a)*sqrt(x)/(a^3*b^2*x^3+2*a^4*b*x^2+a^5*x)`

### Sympy [F]

$$\int \frac{1}{x^{3/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \int \frac{1}{x^{\frac{3}{2}}\operatorname{atanh}^{\frac{5}{2}}(\tanh(a+bx))} dx$$

input `integrate(1/x**(3/2)/atanh(tanh(b*x+a))**(5/2),x)`



output `Integral(1/(x**(3/2)*atanh(tanh(a + b*x))**(5/2)), x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2(8b^3x^3 + 20ab^2x^2 + 15a^2bx + 3a^3)}{3(bx + a)^{5/2}a^3\sqrt{x}}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `-2/3*(8*b^3*x^3 + 20*a*b^2*x^2 + 15*a^2*b*x + 3*a^3)/((b*x + a)^(5/2)*a^3*sqrt(x))`

### Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.58

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2\sqrt{x}\left(\frac{5b^2x}{a^3} + \frac{6b}{a^2}\right)}{3(bx + a)^{3/2}} + \frac{4\sqrt{b}}{\left(\left(\sqrt{b}\sqrt{x} - \sqrt{bx + a}\right)^2 - a\right)a^2}$$

input `integrate(1/x^(3/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `-2/3*sqrt(x)*(5*b^2*x/a^3 + 6*b/a^2)/(b*x + a)^(3/2) + 4*sqrt(b)/(((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)*a^2)`

**Mupad [B] (verification not implemented)**

Time = 3.79 (sec) , antiderivative size = 348, normalized size of antiderivative = 3.28

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2}}}{x^{5/2} - \frac{x^{3/2}\left(\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx\right)}{b}}$$

input `int(1/(x^(3/2)*atanh(tanh(a + b*x))^(5/2)),x)`output `((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(4/(b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (128*x^2)/(3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) - (32*x)/(b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2))/x^(5/2) - (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b + (x^(1/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^2))`**Reduce [F]**

$$\int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^2} dx$$

input `int(1/x^(3/2)/atanh(tanh(b*x+a))^(5/2),x)`output `int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))^3*x**2),x)`

**3.263**  $\int \frac{1}{x^{5/2} \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1830
Mathematica [A] (verified)	1831
Rubi [A] (verified)	1831
Maple [A] (verified)	1833
Fricas [A] (verification not implemented)	1834
Sympy [F(-1)]	1834
Maxima [A] (verification not implemented)	1834
Giac [A] (verification not implemented)	1835
Mupad [B] (verification not implemented)	1835
Reduce [F]	1836

**Optimal result**

Integrand size = 17, antiderivative size = 146

$$\int \frac{1}{x^{5/2} \mathbf{arctanh}(\tanh(a+bx))^{5/2}} dx =$$

$$-\frac{16b^2 \sqrt{x}}{3(bx - \mathbf{arctanh}(\tanh(a+bx)))^3 \mathbf{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{\sqrt{x}(bx - \mathbf{arctanh}(\tanh(a+bx)))^2 \mathbf{arctanh}(\tanh(a+bx))^{3/2}}{4b}$$

$$+ \frac{3x^{3/2}(bx - \mathbf{arctanh}(\tanh(a+bx))) \mathbf{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

$$+ \frac{32b^2 \sqrt{x}}{3(bx - \mathbf{arctanh}(\tanh(a+bx)))^4 \sqrt{\mathbf{arctanh}(\tanh(a+bx))}}$$

output

```
-16/3*b^2*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(3/2)+
4*b/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(3/2)+2/3/x^(
3/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)+32/3*b^2*x^(1/
2)/(b*x-arctanh(tanh(b*x+a)))^4/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.54

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(b^3 x^3 - 9b^2 x^2 \operatorname{arctanh}(\tanh(a + bx)) - 9bx \operatorname{arctanh}(\tanh(a + bx))^2 + \operatorname{arctanh}(\tanh(a + bx))^3)}{3x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{3/2} (-bx + \operatorname{arctanh}(\tanh(a + bx)))^4}$$

input

```
Integrate[1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^(5/2)),x]
```

output

```
(-2*(b^3*x^3 - 9*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 9*b*x*ArcTanh[Tanh[a + b*x]]^2 + ArcTanh[Tanh[a + b*x]]^3)/(3*x^(3/2)*ArcTanh[Tanh[a + b*x]]^(3/2))*(-(b*x) + ArcTanh[Tanh[a + b*x]])^4)
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.21, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$ , Rules used = {2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx$$

$$\downarrow \text{2602}$$

$$\frac{2b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a + bx))} +$$

$$\frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}{2}$$

$$\downarrow \text{2602}$$

$$\begin{aligned}
 & \frac{2b \left( \frac{4b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \\
 & \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2} \\
 & \quad \downarrow \text{2602} \\
 & \frac{2b \left( \frac{4b \left( -\frac{2 \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \\
 & \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2} \\
 & \quad \downarrow \text{2598} \\
 & \frac{2b \left( \frac{4b \left( \frac{4\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \\
 & \frac{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}
 \end{aligned}$$

input `Int [1/(x^(5/2)*ArcTanh[Tanh[a + b*x]]^(5/2)),x]`

output `(2*b*((4*b*((-2*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)) + (4*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]])))/(b*x - ArcTanh[Tanh[a + b*x]]) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) + 2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))`

**Defintions of rubi rules used**

```
rule 2598 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 2602 Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v))], x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.03

method	result
derivativedivides	$-\frac{2}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}-\frac{4b}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$
default	$-\frac{2}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}-\frac{4b}{\sqrt{x}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$

```
input int(1/x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/3/x^(3/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-4*b/(arctanh(tanh(b*x+a))-b*x)*(-1/x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-4*b/(arctanh(tanh(b*x+a))-b*x)*(1/3*x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)+2/3/(arctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.49

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(16b^3x^3 + 24ab^2x^2 + 6a^2bx - a^3)\sqrt{bx+a}\sqrt{x}}{3(a^4b^2x^4 + 2a^5bx^3 + a^6x^2)}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")`

output `2/3*(16*b^3*x^3 + 24*a*b^2*x^2 + 6*a^2*b*x - a^3)*sqrt(b*x + a)*sqrt(x)/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2)`

**Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Timed out}$$

input `integrate(1/x**(5/2)/atanh(tanh(b*x+a))**(5/2),x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.38

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(16b^4x^4 + 40ab^3x^3 + 30a^2b^2x^2 + 5a^3bx - a^4)}{3(bx+a)^{5/2}a^4x^{3/2}}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")`

output `2/3*(16*b^4*x^4 + 40*a*b^3*x^3 + 30*a^2*b^2*x^2 + 5*a^3*b*x - a^4)/((b*x + a)^(5/2)*a^4*x^(3/2))`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.82

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2\sqrt{x} \left( \frac{8b^3x}{a^4} + \frac{9b^2}{a^3} \right)}{3(bx + a)^{3/2}} - \frac{8 \left( 3b^{3/2} (\sqrt{b}\sqrt{x} - \sqrt{bx + a})^4 - 9ab^{3/2} (\sqrt{b}\sqrt{x} - \sqrt{bx + a})^2 + 4a^2b^{3/2} \right)}{3 \left( (\sqrt{b}\sqrt{x} - \sqrt{bx + a})^2 - a \right)^3 a^3}$$

input `integrate(1/x^(5/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`

output `2/3*sqrt(x)*(8*b^3*x/a^4 + 9*b^2/a^3)/(b*x + a)^(3/2) - 8/3*(3*b^(3/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 - 9*a*b^(3/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 + 4*a^2*b^(3/2))/(((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^3*a^3)`

**Mupad [B] (verification not implemented)**

Time = 3.93 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.78

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)}{2}}}{3b^2 \left( \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx \right)} \left( \frac{4}{x^{7/2} - \frac{x^{5/2} \left( \ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) \right)}{e^{2a}e^{2bx+1}}} \right)$$

input `int(1/(x^(5/2)*atanh(tanh(a + b*x))^(5/2)),x)`



output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^(1/2)*(4/(3*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) - (128*x^2)/(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3 + (16*x)/(b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) + (512*b*x^3)/(3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)))/(x^(7/2) - (x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b + (x^(3/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^2))
```

**Reduce [F]**

$$\int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^3} dx$$

input

```
int(1/x^(5/2)/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**3*x**3),x)
```

**3.264**  $\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$

Optimal result	1837
Mathematica [A] (verified)	1838
Rubi [A] (verified)	1838
Maple [A] (verified)	1840
Fricas [A] (verification not implemented)	1841
Sympy [F(-1)]	1842
Maxima [A] (verification not implemented)	1842
Giac [A] (verification not implemented)	1843
Mupad [B] (verification not implemented)	1843
Reduce [F]	1844

**Optimal result**

Integrand size = 17, antiderivative size = 186

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx =$$

$$-\frac{128b^3 \sqrt{x}}{15(bx - \operatorname{arctanh}(\tanh(a+bx)))^4 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

$$+ \frac{5\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx)))^3 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{32b^2}$$

$$+ \frac{15x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{16b}$$

$$+ \frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

$$+ \frac{256b^3 \sqrt{x}}{15(bx - \operatorname{arctanh}(\tanh(a+bx)))^5 \sqrt{\operatorname{arctanh}(\tanh(a+bx))}}$$

output

```
-128/15*b^3*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^4/arctanh(tanh(b*x+a))^(3/2)
)+32/5*b^2/x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^3/arctanh(tanh(b*x+a))^(3/2)
+16/15*b/x^(3/2)/(b*x-arctanh(tanh(b*x+a)))^2/arctanh(tanh(b*x+a))^(3/2)+2
/5/x^(5/2)/(b*x-arctanh(tanh(b*x+a)))/arctanh(tanh(b*x+a))^(3/2)+256/15*b^3
*x^(1/2)/(b*x-arctanh(tanh(b*x+a)))^5/arctanh(tanh(b*x+a))^(1/2)
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.54

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx = \frac{2(-5b^4x^4 + 60b^3x^3 \operatorname{arctanh}(\tanh(a+bx)) + 90b^2x^2 \operatorname{arctanh}(\tanh(a+bx)) - 20bx \operatorname{arctanh}(\tanh(a+bx)) + 3 \operatorname{arctanh}(\tanh(a+bx)))}{15x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{3/2}}$$

input

```
Integrate[1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^(5/2)),x]
```

output

```
(2*(-5*b^4*x^4 + 60*b^3*x^3*ArcTanh[Tanh[a + b*x]] + 90*b^2*x^2*ArcTanh[Tanh[a + b*x]] - 20*b*x*ArcTanh[Tanh[a + b*x]] + 3*ArcTanh[Tanh[a + b*x]]^4)/(15*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])^5*ArcTanh[Tanh[a + b*x]]^(3/2))
```

**Rubi [A] (verified)**Time = 0.48 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.25, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {2602, 2602, 2602, 2602, 2598}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx$$

$$\downarrow 2602$$

$$\frac{8b \int \frac{1}{x^{5/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{5(bx - \operatorname{arctanh}(\tanh(a+bx)))^2} +$$

$$\frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

$$\downarrow 2602$$

$$8b \left( \frac{2b \int \frac{1}{x^{3/2} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a+bx)))}{2} +$$

$$\frac{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}{2}$$

↓ 2602

$$8b \left( \frac{2b \left( \frac{4b \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx}{bx - \operatorname{arctanh}(\tanh(a+bx))} + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx)))}$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2} \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2602

$$8b \left( \frac{2b \left( \frac{4b \left( -\frac{2 \int \frac{1}{\sqrt{x} \operatorname{arctanh}(\tanh(a+bx))^{3/2}} dx}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))} - \frac{2\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right) + \frac{2}{\sqrt{x}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}}}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2} \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

↓ 2598

$$8b \left( \frac{2}{3x^{3/2}(bx - \operatorname{arctanh}(\tanh(a+bx))) \operatorname{arctanh}(\tanh(a+bx))^{3/2}} + \frac{2b \left( \frac{4b \left( \frac{4\sqrt{x}}{3(bx - \operatorname{arctanh}(\tanh(a+bx)))^2 \sqrt{\operatorname{arctanh}(\tanh(a+bx))} - \frac{3(bx - \operatorname{arctanh}(\tanh(a+bx)))}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)}{bx - \operatorname{arctanh}(\tanh(a+bx))} \right)$$

$$\frac{5(bx - \operatorname{arctanh}(\tanh(a + bx)))}{2} \frac{2}{5x^{5/2}(bx - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{3/2}}$$

input

```
Int [1/(x^(7/2)*ArcTanh[Tanh[a + b*x]]^(5/2)), x]
```

output

```
(8*b*((2*b*((4*b*(-2*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)) + (4*Sqrt[x])/(3*(b*x - ArcTanh[Tanh[a + b*x]])^2*Sqrt[ArcTanh[Tanh[a + b*x]]]))))/(b*x - ArcTanh[Tanh[a + b*x]]) + 2/(Sqrt[x]*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(b*x - ArcTanh[Tanh[a + b*x]]) + 2/(3*x^(3/2)*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2)))/(5*(b*x - ArcTanh[Tanh[a + b*x]])) + 2/(5*x^(5/2)*(b*x - ArcTanh[Tanh[a + b*x]])*ArcTanh[Tanh[a + b*x]]^(3/2))
```

### Defintions of rubi rules used

rule 2598

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

rule 2602

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(-u^(m + 1))*(v^(n + 1))/((m + 1)*(b*u - a*v)), x] + Simp[b*((m + n + 2)/((m + 1)*(b*u - a*v)) Int[u^(m + 1)*v^n, x], x] /; NeQ[b*u - a*v, 0] /; PiecewiseLinearQ[u, v, x] && NeQ[m + n + 2, 0] && LtQ[m, -1]
```

### Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.05

method	result
derivativedivides	$\frac{2}{5x^{\frac{5}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$ $- \frac{16b}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))}$
default	$\frac{2}{5x^{\frac{5}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))^{\frac{3}{2}}}$ $- \frac{16b}{3x^{\frac{3}{2}}(\operatorname{arctanh}(\tanh(bx+a))-bx)\operatorname{arctanh}(\tanh(bx+a))}$

```
input int(1/x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/5/x^(5/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-16/5*b/
(arctanh(tanh(b*x+a))-b*x)*(-1/3/x^(3/2)/(arctanh(tanh(b*x+a))-b*x)/arctan
h(tanh(b*x+a))^(3/2)-2*b/(arctanh(tanh(b*x+a))-b*x)*(-1/x^(1/2)/(arctanh(t
anh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)-4*b/(arctanh(tanh(b*x+a))-b*x)
*(1/3*x^(1/2)/(arctanh(tanh(b*x+a))-b*x)/arctanh(tanh(b*x+a))^(3/2)+2/3/(a
rctanh(tanh(b*x+a))-b*x)^2*x^(1/2)/arctanh(tanh(b*x+a))^(1/2)))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.44

$$\int \frac{1}{x^{7/2}\operatorname{arctanh}(\tanh(a+bx))^{5/2}} dx =$$

$$-\frac{2(128b^4x^4 + 192ab^3x^3 + 48a^2b^2x^2 - 8a^3bx + 3a^4)\sqrt{bx+a}\sqrt{x}}{15(a^5b^2x^5 + 2a^6bx^4 + a^7x^3)}$$

```
input integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
-2/15*(128*b^4*x^4 + 192*a*b^3*x^3 + 48*a^2*b^2*x^2 - 8*a^3*b*x + 3*a^4)*s
qrt(b*x + a)*sqrt(x)/(a^5*b^2*x^5 + 2*a^6*b*x^4 + a^7*x^3)
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate(1/x**(7/2)/atanh(tanh(b*x+a))**(5/2),x)
```

output

Timed out

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.36

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{2(128b^5x^5 + 320ab^4x^4 + 240a^2b^3x^3 + 40a^3b^2x^2 - 5a^4bx + 3a^5)}{15(bx + a)^{5/2}a^5x^{5/2}}$$

input

```
integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="maxima")
```

output

```
-2/15*(128*b^5*x^5 + 320*a*b^4*x^4 + 240*a^2*b^3*x^3 + 40*a^3*b^2*x^2 - 5*
a^4*b*x + 3*a^5)/((b*x + a)^(5/2)*a^5*x^(5/2))
```

**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = -\frac{2\sqrt{x}\left(\frac{11b^4x}{a^5} + \frac{12b^3}{a^4}\right)}{3(bx+a)^{3/2}} + \frac{4\left(45b^{5/2}\left(\sqrt{b}\sqrt{x} - \sqrt{bx+a}\right)^8 - 240ab^{5/2}\left(\sqrt{b}\sqrt{x} - \sqrt{bx+a}\right)^6 + 490a^2b^{5/2}\left(\sqrt{b}\sqrt{x} - \sqrt{bx+a}\right)^4 - 320a^3\right)}{15\left(\left(\sqrt{b}\sqrt{x} - \sqrt{bx+a}\right)^2 - a\right)^5 a^4}$$

input `integrate(1/x^(7/2)/arctanh(tanh(b*x+a))^(5/2),x, algorithm="giac")`output `-2/3*sqrt(x)*(11*b^4*x/a^5 + 12*b^3/a^4)/(b*x + a)^(3/2) + 4/15*(45*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^8 - 240*a*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^6 + 490*a^2*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^4 - 320*a^3*b^(5/2)*(sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 + 73*a^4*b^(5/2))/(((sqrt(b)*sqrt(x) - sqrt(b*x + a))^2 - a)^5*a^4)`**Mupad [B] (verification not implemented)**

Time = 4.06 (sec) , antiderivative size = 466, normalized size of antiderivative = 2.51

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \frac{\sqrt{\frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right)}{2}}}{5b^2\left(\ln\left(\frac{2}{e^{2a}e^{2bx+1}}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx+1}}\right) + 2bx\right)^4}$$

input `int(1/(x^(7/2)*atanh(tanh(a + b*x))^(5/2)),x)`



output

```
((log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1)))/2^(1/2)*(4/(5*b^2*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)) + (256*x^2)/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3) + (64*x)/(15*b*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2) - (2048*b*x^3)/(5*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4) + (8192*b^2*x^4)/(15*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5)))/(x^(9/2) - (x^(7/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/b + (x^(5/2)*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2)/(4*b^2))
```

**Reduce [F]**

$$\int \frac{1}{x^{7/2} \operatorname{arctanh}(\tanh(a + bx))^{5/2}} dx = \int \frac{\sqrt{x} \sqrt{\operatorname{atanh}(\tanh(bx + a))}}{\operatorname{atanh}(\tanh(bx + a))^3 x^4} dx$$

input

```
int(1/x^(7/2)/atanh(tanh(b*x+a))^(5/2),x)
```

output

```
int((sqrt(x)*sqrt(atanh(tanh(a + b*x))))/(atanh(tanh(a + b*x))**3*x**4),x)
```

### 3.265 $\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1845
Mathematica [A] (verified)	1845
Rubi [A] (verified)	1846
Maple [F]	1847
Fricas [F]	1847
Sympy [F]	1847
Maxima [F]	1848
Giac [F]	1848
Mupad [F(-1)]	1848
Reduce [F]	1849

#### Optimal result

Integrand size = 13, antiderivative size = 79

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{x^m \left( \frac{bx}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)^{-m} \operatorname{arctanh}(\tanh(a + bx))^{1+n} \operatorname{Hypergeometric2F1} \left( -m, 1 + n, 2 + n, -\frac{\operatorname{arctanh}(\tanh(a + bx))}{bx - a} \right)}{b(1 + n)}$$

output

```
x^m*arctanh(tanh(b*x+a))^(1+n)*hypergeom([-m, 1+n],[2+n],-arctanh(tanh(b*x+a))/(b*x-arctanh(tanh(b*x+a))))/b/(1+n)/((b*x/(b*x-arctanh(tanh(b*x+a))))^m)
```

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.90

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{x^{1+m} \operatorname{arctanh}(\tanh(a + bx))^n \left( 1 + \frac{bx}{-bx + \operatorname{arctanh}(\tanh(a + bx))} \right)^{-n} \operatorname{Hypergeometric2F1} \left( 1 + m, -n, 2 + m, -\frac{\operatorname{arctanh}(\tanh(a + bx))}{bx - a} \right)}{1 + m}$$

input

```
Integrate[x^m*ArcTanh[Tanh[a + b*x]]^n,x]
```

output

```
(x^(1 + m)*ArcTanh[Tanh[a + b*x]]^n*Hypergeometric2F1[1 + m, -n, 2 + m, -(
(b*x)/(-(b*x) + ArcTanh[Tanh[a + b*x]])])]/((1 + m)*(1 + (b*x)/(-(b*x) + A
rcTanh[Tanh[a + b*x]]))^n)
```

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2604}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx$$

↓ 2604

$$\frac{x^m \left( \frac{bx}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)^{-m} \operatorname{arctanh}(\tanh(a + bx))^{n+1} \operatorname{Hypergeometric2F1} \left( -m, n + 1, n + 2, -\frac{\operatorname{arctanh}(\tanh(a + bx))}{bx - \operatorname{arctanh}(\tanh(a + bx))} \right)}{b(n + 1)}$$

input

```
Int[x^m*ArcTanh[Tanh[a + b*x]]^n,x]
```

output

```
(x^m*ArcTanh[Tanh[a + b*x]]^(1 + n)*Hypergeometric2F1[-m, 1 + n, 2 + n, -(
ArcTanh[Tanh[a + b*x]]/(b*x - ArcTanh[Tanh[a + b*x]])])]/(b*(1 + n)*((b*x)
/(b*x - ArcTanh[Tanh[a + b*x]]))^m)
```

**Defintions of rubi rules used**

rule 2604

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simp
lify[D[v, x]]}, Simp[u^m*(v^(n + 1)/(b*(n + 1)*(b*(u/(b*u - a*v)))^m))*Hype
rgeometric2F1[-m, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v,
0] /; PiecewiseLinearQ[u, v, x] && !IntegerQ[m] && !IntegerQ[n]
```

**Maple [F]**

$$\int x^m \operatorname{arctanh}(\tanh(bx + a))^n dx$$

input `int(x^m*arctanh(tanh(b*x+a))^n,x)`

output `int(x^m*arctanh(tanh(b*x+a))^n,x)`

**Fricas [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{artanh}(\tanh(bx + a))^n dx$$

input `integrate(x^m*arctanh(tanh(b*x+a))^n,x, algorithm="fricas")`

output `integral(x^m*arctanh(tanh(b*x + a))^n, x)`

**Sympy [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{atanh}^n(\tanh(a + bx)) dx$$

input `integrate(x**m*atanh(tanh(b*x+a))**n,x)`

output `Integral(x**m*atanh(tanh(a + b*x))**n, x)`

**Maxima [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{artanh}(\tanh(bx + a))^n dx$$

input `integrate(x^m*arctanh(tanh(b*x+a))^n,x, algorithm="maxima")`

output `integrate(x^m*arctanh(tanh(b*x + a))^n, x)`

**Giac [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{artanh}(\tanh(bx + a))^n dx$$

input `integrate(x^m*arctanh(tanh(b*x+a))^n,x, algorithm="giac")`

output `integrate(x^m*arctanh(tanh(b*x + a))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{atanh}(\tanh(a + bx))^n dx$$

input `int(x^m*atanh(tanh(a + b*x))^n,x)`

output `int(x^m*atanh(tanh(a + b*x))^n, x)`

**Reduce [F]**

$$\int x^m \operatorname{arctanh}(\tanh(a + bx))^n dx = \int x^m \operatorname{atanh}(\tanh(bx + a))^n dx$$

input `int(x^m*atanh(tanh(b*x+a))^n,x)`

output `int(x**m*atanh(tanh(a + b*x))**n,x)`

### 3.266 $\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1850
Mathematica [A] (verified)	1851
Rubi [A] (verified)	1851
Maple [B] (verified)	1853
Fricas [B] (verification not implemented)	1854
Sympy [F]	1855
Maxima [A] (verification not implemented)	1855
Giac [B] (verification not implemented)	1856
Mupad [B] (verification not implemented)	1857
Reduce [F]	1858

#### Optimal result

Integrand size = 13, antiderivative size = 165

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{x^4 \operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)} - \frac{4x^3 \operatorname{arctanh}(\tanh(a + bx))^{2+n}}{b^2(1+n)(2+n)} + \frac{12x^2 \operatorname{arctanh}(\tanh(a + bx))^{3+n}}{b^3(1+n)(2+n)(3+n)} - \frac{24x \operatorname{arctanh}(\tanh(a + bx))^{4+n}}{b^4(1+n)(2+n)(3+n)(4+n)} + \frac{24 \operatorname{arctanh}(\tanh(a + bx))^{5+n}}{b^5(1+n)(2+n)(3+n)(4+n)(5+n)}$$

output

```
x^4*arctanh(tanh(b*x+a))^(1+n)/b/(1+n)-4*x^3*arctanh(tanh(b*x+a))^(2+n)/b^2/(1+n)/(2+n)+12*x^2*arctanh(tanh(b*x+a))^(3+n)/b^3/(1+n)/(2+n)/(3+n)-24*x*arctanh(tanh(b*x+a))^(4+n)/b^4/(1+n)/(2+n)/(3+n)/(4+n)+24*arctanh(tanh(b*x+a))^(5+n)/b^5/(1+n)/(2+n)/(3+n)/(4+n)/(5+n)
```

### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.88

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n} (b^4(120 + 154n + 71n^2 + 14n^3 + n^4) x^4 - 4b^3(60 + 47n + 12n^2 + n^3) x^3 \operatorname{arctanh}(\tanh(a + bx)) + 12b^2(20 + 9n + n^2) x^2 \operatorname{arctanh}(\tanh(a + bx))^2 - 24b(5 + n) x \operatorname{arctanh}(\tanh(a + bx))^3 + 24 \operatorname{arctanh}(\tanh(a + bx))^4)}{b^5(1+n)(2+n)(3+n)(4+n)(5+n)}$$

input `Integrate[x^4*ArcTanh[Tanh[a + b*x]]^n,x]`

output `(ArcTanh[Tanh[a + b*x]]^(1 + n)*(b^4*(120 + 154*n + 71*n^2 + 14*n^3 + n^4)*x^4 - 4*b^3*(60 + 47*n + 12*n^2 + n^3)*x^3*ArcTanh[Tanh[a + b*x]] + 12*b^2*(20 + 9*n + n^2)*x^2*ArcTanh[Tanh[a + b*x]]^2 - 24*b*(5 + n)*x*ArcTanh[Tanh[a + b*x]]^3 + 24*ArcTanh[Tanh[a + b*x]]^4)/(b^5*(1 + n)*(2 + n)*(3 + n)*(4 + n)*(5 + n))`

### Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.91, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {2599, 2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$\downarrow 2599$$

$$\frac{x^4 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{4 \int x^3 \operatorname{arctanh}(\tanh(a + bx))^{n+1} dx}{b(n+1)}$$

$$\downarrow 2599$$

$$\frac{x^4 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b(n+2)} - \frac{3 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+2} dx}{b(n+2)} \right)}{b(n+1)}$$

$$\downarrow 2599$$



$$\begin{aligned}
 & \frac{x^4 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
 & 4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \int x \operatorname{arctanh}(\tanh(a+bx))^{n+3} dx}{b(n+3)} \right)}{b(n+2)} \right) \\
 & \frac{\phantom{4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \int x \operatorname{arctanh}(\tanh(a+bx))^{n+3} dx}{b(n+3)} \right)}{b(n+2)} \right)}}{b(n+1)} \\
 & \quad \downarrow \text{2599} \\
 & \frac{x^4 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
 & 4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+4} dx}{b(n+4)} \right)}{b(n+3)} \right)}{b(n+2)} \right) \\
 & \frac{\phantom{4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+4} dx}{b(n+4)} \right)}{b(n+3)} \right)}{b(n+2)} \right)}}{b(n+1)} \\
 & \quad \downarrow \text{2588} \\
 & \frac{x^4 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
 & 4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+4} dx \operatorname{arctanh}(\tanh(a+bx))}{b^2(n+4)} \right)}{b(n+3)} \right)}{b(n+2)} \right) \\
 & \frac{\phantom{4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+4} dx \operatorname{arctanh}(\tanh(a+bx))}{b^2(n+4)} \right)}{b(n+3)} \right)}{b(n+2)} \right)}}{b(n+1)} \\
 & \quad \downarrow \text{15} \\
 & \frac{x^4 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
 & 4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{n+5}}{b^2(n+4)(n+5)} \right)}{b(n+3)} \right)}{b(n+2)} \right) \\
 & \frac{\phantom{4 \left( \frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b(n+4)} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{n+5}}{b^2(n+4)(n+5)} \right)}{b(n+3)} \right)}{b(n+2)} \right)}}{b(n+1)}
 \end{aligned}$$

input

`Int [x^4*ArcTanh [Tanh [a + b*x]]^n,x]`

output

$$\frac{(x^4 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{(1+n)}) / (b(1+n)) - (4((x^3 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{(2+n)}) / (b(2+n)) - (3((x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{(3+n)}) / (b(3+n)) - (2((x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{(4+n)}) / (b(4+n)) - \operatorname{ArcTanh}[\operatorname{Tanh}[a + b x]]^{(5+n)} / (b^2(4+n)(5+n)))) / (b(3+n)))) / (b(2+n))) / (b(1+n))$$

### Defintions of rubi rules used

rule 15

$$\operatorname{Int}[(a_.)(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a(x^{(m+1)}) / (m+1), x] \;/; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$$

rule 2588

$$\operatorname{Int}[(u_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] \;/; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$$

rule 2599

$$\operatorname{Int}[(u_)^{(m_.)}(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}(v^n / (a(m+1))), x] - \operatorname{Simp}[b(n / (a(m+1))) \operatorname{Int}[u^{(m+1)}v^{(n-1)}, x], x] \;/; \operatorname{NeQ}[b u - a v, 0] \;/; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m+n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2n+m+1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs.  $2(165) = 330$ .

Time = 3.50 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.46

method	result
default	$\frac{x^5 e^{n \ln(\operatorname{arctanh}(\operatorname{tanh}(bx+a)))}}{5+n} + \frac{n(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)x^4 e^{n \ln(\operatorname{arctanh}(\operatorname{tanh}(bx+a)))}}{b(n^2+9n+20)} - \frac{4n(a^2+2a(\operatorname{arctanh}(\operatorname{tanh}(bx+a))))}{b^2(n^2+9n+20)}$
parallelrisch	$- \frac{120 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^n \operatorname{arctanh}(\operatorname{tanh}(bx+a))x^4 b^4 + 240 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^n \operatorname{arctanh}(\operatorname{tanh}(bx+a))^2 x^3 b^3 - 240 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^n}{b^2(n^2+9n+20)}$
risch	Expression too large to display

input

$$\operatorname{int}(x^4 \operatorname{arctanh}(\operatorname{tanh}(b x+a))^n, x, \operatorname{method}=\_RETURNVERBOSE)$$

output

```

1/(5+n)*x^5*exp(n*ln(arctanh(tanh(b*x+a))))+n/b*(arctanh(tanh(b*x+a))-b*x)
/(n^2+9*n+20)*x^4*exp(n*ln(arctanh(tanh(b*x+a))))-4*n*(a^2+2*a*(arctanh(ta
nh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/b^2/(n^3+12*n^2+47*n+60)
*x^3*exp(n*ln(arctanh(tanh(b*x+a))))+24*(arctanh(tanh(b*x+a))-b*x)*(a^2+2*
a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)/b^5/(n^3+
12*n^2+47*n+60)/(n^2+3*n+2)*exp(n*ln(arctanh(tanh(b*x+a))))-24*(a^2+2*a*(a
rctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*x+a))-b*x-a)^2)*n/b^4/(n^3+12
*n^2+47*n+60)/(n^2+3*n+2)*x*exp(n*ln(arctanh(tanh(b*x+a))))+12/b^3*(arctan
h(tanh(b*x+a))-b*x)*(a^2+2*a*(arctanh(tanh(b*x+a))-b*x-a)+(arctanh(tanh(b*
x+a))-b*x-a)^2)*n/(2+n)/(n^3+12*n^2+47*n+60)*x^2*exp(n*ln(arctanh(tanh(b*x
+a))))

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 374 vs.  $2(165) = 330$ .

Time = 0.08 (sec) , antiderivative size = 374, normalized size of antiderivative = 2.27

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{(24 a^4 b n x - (b^5 n^4 + 10 b^5 n^3 + 35 b^5 n^2 + 50 b^5 n + 24 b^5) x^5 - 24 a^5 - (a b^4 n^4 + 6 a b^4 n^3 + 11 a b^4 n^2 + 6 a b^4 n) x^4 + 4 (a^2 b^3 n^3 + 3 a^2 b^3 n^2 + 2 a^2 b^3 n) x^3 - 12 (a^3 b^2 n^2 + a^3 b^2 n) x^2) \cosh(n \log(b x + a)) + (24 a^4 b n x - (b^5 n^4 + 10 b^5 n^3 + 35 b^5 n^2 + 50 b^5 n + 24 b^5) x^5 - 24 a^5 - (a b^4 n^4 + 6 a b^4 n^3 + 11 a b^4 n^2 + 6 a b^4 n) x^4 + 4 (a^2 b^3 n^3 + 3 a^2 b^3 n^2 + 2 a^2 b^3 n) x^3 - 12 (a^3 b^2 n^2 + a^3 b^2 n) x^2) \sinh(n \log(b x + a))}{(b^5 n^5 + 15 b^5 n^4 + 85 b^5 n^3 + 225 b^5 n^2 + 274 b^5 n + 120 b^5)}$$

input

```
integrate(x^4*arctanh(tanh(b*x+a))^n,x, algorithm="fricas")
```

output

```

-((24*a^4*b*n*x - (b^5*n^4 + 10*b^5*n^3 + 35*b^5*n^2 + 50*b^5*n + 24*b^5)*
x^5 - 24*a^5 - (a*b^4*n^4 + 6*a*b^4*n^3 + 11*a*b^4*n^2 + 6*a*b^4*n)*x^4 +
4*(a^2*b^3*n^3 + 3*a^2*b^3*n^2 + 2*a^2*b^3*n)*x^3 - 12*(a^3*b^2*n^2 + a^3*
b^2*n)*x^2)*cosh(n*log(b*x + a)) + (24*a^4*b*n*x - (b^5*n^4 + 10*b^5*n^3 +
35*b^5*n^2 + 50*b^5*n + 24*b^5)*x^5 - 24*a^5 - (a*b^4*n^4 + 6*a*b^4*n^3 +
11*a*b^4*n^2 + 6*a*b^4*n)*x^4 + 4*(a^2*b^3*n^3 + 3*a^2*b^3*n^2 + 2*a^2*b^
3*n)*x^3 - 12*(a^3*b^2*n^2 + a^3*b^2*n)*x^2)*sinh(n*log(b*x + a)))/(b^5*n^
5 + 15*b^5*n^4 + 85*b^5*n^3 + 225*b^5*n^2 + 274*b^5*n + 120*b^5)

```

**Sympy [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx = \text{Too large to display}$$

input `integrate(x**4*atanh(tanh(b*x+a))**n,x)`

output `Piecewise((x**5*atanh(tanh(a))**n/5, Eq(b, 0)), (-x**4/(4*b*atanh(tanh(a + b*x))**4) - x**3/(3*b**2*atanh(tanh(a + b*x))**3) - x**2/(2*b**3*atanh(tanh(a + b*x))**2) - x/(b**4*atanh(tanh(a + b*x))) + log(atanh(tanh(a + b*x)))/b**5, Eq(n, -5)), (Integral(x**4/atanh(tanh(a + b*x))**4, x), Eq(n, -4)), (Integral(x**4/atanh(tanh(a + b*x))**3, x), Eq(n, -3)), (Integral(x**4/atanh(tanh(a + b*x))**2, x), Eq(n, -2)), (Integral(x**4/atanh(tanh(a + b*x))), x), Eq(n, -1)), (b**4*n**4*x**4*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) + 14*b**4*n**3*x**4*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) + 71*b**4*n**2*x**4*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) + 154*b**4*n*x**4*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) + 120*b**4*x**4*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) - 4*b**3*n**3*x**3*atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + 120*b**5) - 48*b**3*n**2*x**3*atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**5*n**5 + 15*b**5*n**4 + 85*b**5*n**3 + 225*b**5*n**2 + 274*b**5*n + ...`

**Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.84

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{((n^4 + 10n^3 + 35n^2 + 50n + 24)b^5x^5 + (n^4 + 6n^3 + 11n^2 + 6n)ab^4x^4 - 4(n^3 + 3n^2 + 2n)a^2b^3x^3 + 12n^2a^3b^2x^2 - 4n^2a^4b^2x - 4n^2a^5)x^5}{(n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)b^5}$$

input `integrate(x^4*arctanh(tanh(b*x+a))^n,x, algorithm="maxima")`

output

```
((n^4 + 10*n^3 + 35*n^2 + 50*n + 24)*b^5*x^5 + (n^4 + 6*n^3 + 11*n^2 + 6*n)
)*a*b^4*x^4 - 4*(n^3 + 3*n^2 + 2*n)*a^2*b^3*x^3 + 12*(n^2 + n)*a^3*b^2*x^2
- 24*a^4*b*n*x + 24*a^5)*(b*x + a)^n/((n^5 + 15*n^4 + 85*n^3 + 225*n^2 +
274*n + 120)*b^5)
```

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs.  $2(165) = 330$ .

Time = 0.12 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.01

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(bx + a)^n b^5 n^4 x^5 + (bx + a)^n a b^4 n^4 x^4 + 10 (bx + a)^n b^5 n^3 x^5 + 6 (bx + a)^n a b^4 n^3 x^4 + 35 (bx + a)^n b^5 n^2 x^5 -$$

input

```
integrate(x^4*arctanh(tanh(b*x+a))^n,x, algorithm="giac")
```

output

```
((b*x + a)^n*b^5*n^4*x^5 + (b*x + a)^n*a*b^4*n^4*x^4 + 10*(b*x + a)^n*b^5*
n^3*x^5 + 6*(b*x + a)^n*a*b^4*n^3*x^4 + 35*(b*x + a)^n*b^5*n^2*x^5 - 4*(b*
x + a)^n*a^2*b^3*n^3*x^3 + 11*(b*x + a)^n*a*b^4*n^2*x^4 + 50*(b*x + a)^n*b
^5*n*x^5 - 12*(b*x + a)^n*a^2*b^3*n^2*x^3 + 6*(b*x + a)^n*a*b^4*n*x^4 + 24
*(b*x + a)^n*b^5*x^5 + 12*(b*x + a)^n*a^3*b^2*n^2*x^2 - 8*(b*x + a)^n*a^2*
b^3*n*x^3 + 12*(b*x + a)^n*a^3*b^2*n*x^2 - 24*(b*x + a)^n*a^4*b*n*x + 24*(
b*x + a)^n*a^5)/(b^5*n^5 + 15*b^5*n^4 + 85*b^5*n^3 + 225*b^5*n^2 + 274*b^5
*n + 120*b^5)
```

**Mupad [B] (verification not implemented)**

Time = 3.86 (sec) , antiderivative size = 546, normalized size of antiderivative = 3.31

$$\begin{aligned}
& \int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx = \\
& - \left( \frac{\ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right)}{2} - \frac{\ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right)}{2} \right)^n \left( \frac{3 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^5}{4b^5 (n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)} \right. \\
& \quad - \frac{x^5 (n^4 + 10n^3 + 35n^2 + 50n + 24)}{n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120} \\
& \quad + \frac{3nx \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^4}{2b^4 (n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)} \\
& \quad + \frac{nx^4 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right) (n^3 + 6n^2 + 11n + 6)}{2b (n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)} \\
& \quad + \frac{3nx^2 (n + 1) \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^3}{2b^3 (n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)} \\
& \quad \left. + \frac{nx^3 \left( \ln\left(\frac{2}{e^{2a}e^{2bx}+1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}+1}\right) + 2bx \right)^2 (n^2 + 3n + 2)}{b^2 (n^5 + 15n^4 + 85n^3 + 225n^2 + 274n + 120)} \right)
\end{aligned}$$

input `int(x^4*atanh(tanh(a + b*x))^n,x)`

output

```

-(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^n*((3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^5)/(4*b^5*(274*n +
225*n^2 + 85*n^3 + 15*n^4 + n^5 + 120)) - (x^5*(50*n + 35*n^2 + 10*n^3 + n
^4 + 24))/(274*n + 225*n^2 + 85*n^3 + 15*n^4 + n^5 + 120) + (3*n*x*(log(2/
(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b
*x) + 1)) + 2*b*x)^4)/(2*b^4*(274*n + 225*n^2 + 85*n^3 + 15*n^4 + n^5 + 12
0)) + (n*x^4*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x
)))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x*(11*n + 6*n^2 + n^3 + 6))/(2*b*(274
*n + 225*n^2 + 85*n^3 + 15*n^4 + n^5 + 120)) + (3*n*x^2*(n + 1)*(log(2/(ex
p(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)
+ 1)) + 2*b*x)^3)/(2*b^3*(274*n + 225*n^2 + 85*n^3 + 15*n^4 + n^5 + 120))
+ (n*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/
(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2*(3*n + n^2 + 2))/(b^2*(274*n + 225*n
^2 + 85*n^3 + 15*n^4 + n^5 + 120)))

```

**Reduce [F]**

$$\int x^4 \operatorname{arctanh}(\tanh(a + bx))^n dx = \int \operatorname{atanh}(\tanh(bx + a))^n x^4 dx$$

input

```
int(x^4*atanh(tanh(b*x+a))^n,x)
```

output

```
int(atanh(tanh(a + b*x))^n*x**4,x)
```

### 3.267 $\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1859
Mathematica [A] (verified)	1860
Rubi [A] (verified)	1860
Maple [B] (verified)	1862
Fricas [B] (verification not implemented)	1863
Sympy [F]	1863
Maxima [A] (verification not implemented)	1864
Giac [A] (verification not implemented)	1865
Mupad [B] (verification not implemented)	1865
Reduce [F]	1866

#### Optimal result

Integrand size = 13, antiderivative size = 121

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)} - \frac{3x^2 \operatorname{arctanh}(\tanh(a + bx))^{2+n}}{b^2(1+n)(2+n)} + \frac{6x \operatorname{arctanh}(\tanh(a + bx))^{3+n}}{b^3(1+n)(2+n)(3+n)} - \frac{6 \operatorname{arctanh}(\tanh(a + bx))^{4+n}}{b^4(1+n)(2+n)(3+n)(4+n)}$$

output

```
x^3*arctanh(tanh(b*x+a))^(1+n)/b/(1+n)-3*x^2*arctanh(tanh(b*x+a))^(2+n)/b^2/(1+n)/(2+n)+6*x*arctanh(tanh(b*x+a))^(3+n)/b^3/(1+n)/(2+n)/(3+n)-6*arctanh(tanh(b*x+a))^(4+n)/b^4/(1+n)/(2+n)/(3+n)/(4+n)
```



**Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.88

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n} (b^3(24 + 26n + 9n^2 + n^3)x^3 - 3b^2(12 + 7n + n^2)x^2 \operatorname{arctanh}(\tanh(a + bx)) + a + b*x)]^2 - 6*\operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^3)}{b^4(1+n)(2+n)(3+n)(4+n)}$$

input `Integrate[x^3*ArcTanh[Tanh[a + b*x]]^n,x]`

output `(ArcTanh[Tanh[a + b*x]]^(1 + n)*(b^3*(24 + 26*n + 9*n^2 + n^3)*x^3 - 3*b^2*(12 + 7*n + n^2)*x^2*ArcTanh[Tanh[a + b*x]] + 6*b*(4 + n)*x*ArcTanh[Tanh[a + b*x]]^2 - 6*ArcTanh[Tanh[a + b*x]]^3))/(b^4*(1 + n)*(2 + n)*(3 + n)*(4 + n))`

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {2599, 2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$\downarrow 2599$$

$$\frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{3 \int x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+1} dx}{b(n+1)}$$

$$\downarrow 2599$$

$$\frac{x^3 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b(n+2)} - \frac{2 \int x \operatorname{arctanh}(\tanh(a + bx))^{n+2} dx}{b(n+2)} \right)}{b(n+1)}$$

$$\downarrow 2599$$

$$\begin{array}{c}
\frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
\frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+3} dx}{b(n+3)} \right)}{b(n+2)} \right)}{b(n+1)} \\
\downarrow 2588 \\
\frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
\frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{\int \operatorname{arctanh}(\tanh(a+bx))^{n+3} dx}{b^2(n+3)} \right)}{b(n+2)} \right)}{b(n+1)} \\
\downarrow 15 \\
\frac{x^3 \operatorname{arctanh}(\tanh(a+bx))^{n+1}}{b(n+1)} - \\
\frac{3 \left( \frac{x^2 \operatorname{arctanh}(\tanh(a+bx))^{n+2}}{b(n+2)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a+bx))^{n+3}}{b(n+3)} - \frac{\operatorname{arctanh}(\tanh(a+bx))^{n+4}}{b^2(n+3)(n+4)} \right)}{b(n+2)} \right)}{b(n+1)}
\end{array}$$

input `Int[x^3*ArcTanh[Tanh[a + b*x]]^n,x]`

output `(x^3*ArcTanh[Tanh[a + b*x]]^(1 + n))/(b*(1 + n)) - (3*((x^2*ArcTanh[Tanh[a + b*x]]^(2 + n))/(b*(2 + n)) - (2*((x*ArcTanh[Tanh[a + b*x]]^(3 + n))/(b*(3 + n)) - ArcTanh[Tanh[a + b*x]]^(4 + n)/(b^2*(3 + n)*(4 + n)))))/(b*(2 + n)))/(b*(1 + n))`

### Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 276 vs.  $2(121) = 242$ .

Time = 1.49 (sec) , antiderivative size = 277, normalized size of antiderivative = 2.29

method	result
parallelrisch	$-\frac{6 \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))^4 - 24 \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))^3 x b - 24 \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))^2 x^2 b^2 - x^3 \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))}{4+n} + \frac{n(\operatorname{arctanh}(\tanh(bx+a)) - bx)x^3 e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{b(n^2+7n+12)} - \frac{6(\operatorname{arctanh}(\tanh(bx+a)) - bx)}{b(n^2+7n+12)}$
default	$\frac{x^4 e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{4+n} + \frac{n(\operatorname{arctanh}(\tanh(bx+a)) - bx)x^3 e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{b(n^2+7n+12)} - \frac{6(\operatorname{arctanh}(\tanh(bx+a)) - bx)}{b(n^2+7n+12)}$
risch	Expression too large to display

input

```
int(x^3*arctanh(tanh(b*x+a))^n,x,method=_RETURNVERBOSE)
```

output

```
-(6*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))^4-24*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))^3*x*b-24*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))^2*x^2*b^2-x^3*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b^3*n^3-9*x^3*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b^3*n^2-26*x^3*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b^3*n+3*x^2*arctanh(tanh(b*x+a))^2*arctanh(tanh(b*x+a))^n*b^2*n^2+21*x^2*arctanh(tanh(b*x+a))^2*arctanh(tanh(b*x+a))^n*b^2*n-6*x*arctanh(tanh(b*x+a))^3*arctanh(tanh(b*x+a))^n*b*n)/(n^3+6*n^2+11*n+6)/(4+n)/b^4
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 255 vs.  $2(121) = 242$ .

Time = 0.08 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.11

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(6a^3bnx + (b^4n^3 + 6b^4n^2 + 11b^4n + 6b^4)x^4 - 6a^4 + (ab^3n^3 + 3ab^3n^2 + 2ab^3n)x^3 - 3(a^2b^2n^2 + a^2b^2n)) \cosh(n \log(bx + a)) + (6a^3bnx + (b^4n^3 + 6b^4n^2 + 11b^4n + 6b^4)x^4 - 6a^4 + (ab^3n^3 + 3ab^3n^2 + 2ab^3n)x^3 - 3(a^2b^2n^2 + a^2b^2n)) \sinh(n \log(bx + a))}{(b^4n^4 + 10b^4n^3 + 35b^4n^2 + 50b^4n + 24b^4)}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^n,x, algorithm="fricas")`

output `((6*a^3*b*n*x + (b^4*n^3 + 6*b^4*n^2 + 11*b^4*n + 6*b^4)*x^4 - 6*a^4 + (a*b^3*n^3 + 3*a*b^3*n^2 + 2*a*b^3*n)*x^3 - 3*(a^2*b^2*n^2 + a^2*b^2*n)*x^2)*cosh(n*log(b*x + a)) + (6*a^3*b*n*x + (b^4*n^3 + 6*b^4*n^2 + 11*b^4*n + 6*b^4)*x^4 - 6*a^4 + (a*b^3*n^3 + 3*a*b^3*n^2 + 2*a*b^3*n)*x^3 - 3*(a^2*b^2*n^2 + a^2*b^2*n)*x^2)*sinh(n*log(b*x + a))/(b^4*n^4 + 10*b^4*n^3 + 35*b^4*n^2 + 50*b^4*n + 24*b^4)`

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx = \text{Too large to display}$$

input `integrate(x**3*atanh(tanh(b*x+a))**n,x)`

output

```
Piecewise((x**4*atanh(tanh(a))**n/4, Eq(b, 0)), (-x**3/(3*b*atanh(tanh(a +
b*x))**3) - x**2/(2*b**2*atanh(tanh(a + b*x))**2) - x/(b**3*atanh(tanh(a
+ b*x))) + log(atanh(tanh(a + b*x)))/b**4, Eq(n, -4)), (Integral(x**3/atan
h(tanh(a + b*x))**3, x), Eq(n, -3)), (Integral(x**3/atanh(tanh(a + b*x))**
2, x), Eq(n, -2)), (Integral(x**3/atanh(tanh(a + b*x)), x), Eq(n, -1)), (b
**3*n**3*x**3*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10
*b**4*n**3 + 35*b**4*n**2 + 50*b**4*n + 24*b**4) + 9*b**3*n**2*x**3*atanh(
tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10*b**4*n**3 + 35*b**4
*n**2 + 50*b**4*n + 24*b**4) + 26*b**3*n*x**3*atanh(tanh(a + b*x))*atanh(t
anh(a + b*x))**n/(b**4*n**4 + 10*b**4*n**3 + 35*b**4*n**2 + 50*b**4*n + 24
*b**4) + 24*b**3*x**3*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**4*n
**4 + 10*b**4*n**3 + 35*b**4*n**2 + 50*b**4*n + 24*b**4) - 3*b**2*n**2*x**
2*atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10*b**4*n**
3 + 35*b**4*n**2 + 50*b**4*n + 24*b**4) - 21*b**2*n*x**2*atanh(tanh(a + b
*x))**2*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10*b**4*n**3 + 35*b**4*n**2 +
50*b**4*n + 24*b**4) - 36*b**2*x**2*atanh(tanh(a + b*x))**2*atanh(tanh(a +
b*x))**n/(b**4*n**4 + 10*b**4*n**3 + 35*b**4*n**2 + 50*b**4*n + 24*b**4)
+ 6*b*n*x*atanh(tanh(a + b*x))**3*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10*
b**4*n**3 + 35*b**4*n**2 + 50*b**4*n + 24*b**4) + 24*b*x*atanh(tanh(a + b*
x))**3*atanh(tanh(a + b*x))**n/(b**4*n**4 + 10*b**4*n**3 + 35*b**4*n**2...
```

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.83

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{((n^3 + 6n^2 + 11n + 6)b^4 x^4 + (n^3 + 3n^2 + 2n)ab^3 x^3 - 3(n^2 + n)a^2 b^2 x^2 + 6a^3 b n x - 6a^4)(bx + a)^n}{(n^4 + 10n^3 + 35n^2 + 50n + 24)b^4}$$

input

```
integrate(x^3*arctanh(tanh(b*x+a))^n,x, algorithm="maxima")
```

output

```
((n^3 + 6*n^2 + 11*n + 6)*b^4*x^4 + (n^3 + 3*n^2 + 2*n)*a*b^3*x^3 - 3*(n^2
+ n)*a^2*b^2*x^2 + 6*a^3*b*n*x - 6*a^4)*(b*x + a)^n/((n^4 + 10*n^3 + 35*n
^2 + 50*n + 24)*b^4)
```

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.87

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(bx + a)^n b^4 n^3 x^4 + (bx + a)^n a b^3 n^3 x^3 + 6 (bx + a)^n b^4 n^2 x^4 + 3 (bx + a)^n a b^3 n^2 x^3 + 11 (bx + a)^n b^4 n x^4 - 3 (bx + a)^n a b^3 n x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3}{b^4 n^4 + 10 b^4 n^3}$$

input `integrate(x^3*arctanh(tanh(b*x+a))^n,x, algorithm="giac")`output  $((bx + a)^n b^4 n^3 x^4 + (bx + a)^n a b^3 n^3 x^3 + 6 (bx + a)^n b^4 n^2 x^4 + 3 (bx + a)^n a b^3 n^2 x^3 + 11 (bx + a)^n b^4 n x^4 - 3 (bx + a)^n a b^3 n x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3 + 6 (bx + a)^n b^4 x^4 - 3 (bx + a)^n a b^3 x^3) / (b^4 n^4 + 10 b^4 n^3 + 35 b^4 n^2 + 50 b^4 n + 24 b^4)$ **Mupad [B] (verification not implemented)**

Time = 3.56 (sec) , antiderivative size = 418, normalized size of antiderivative = 3.45

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx =$$

$$- \left( \frac{\ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)}{2} - \frac{\ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right)}{2} \right)^n \left( \frac{3 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^4}{8b^4 (n^4 + 10n^3 + 35n^2 + 50n + 24)} \right.$$

$$- \frac{x^4 (n^3 + 6n^2 + 11n + 6)}{n^4 + 10n^3 + 35n^2 + 50n + 24} + \frac{3nx \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{4b^3 (n^4 + 10n^3 + 35n^2 + 50n + 24)}$$

$$+ \frac{nx^3 \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right) (n^2 + 3n + 2)}{2b (n^4 + 10n^3 + 35n^2 + 50n + 24)}$$

$$\left. + \frac{3nx^2 (n + 1) \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{4b^2 (n^4 + 10n^3 + 35n^2 + 50n + 24)} \right)$$

input `int(x^3*atanh(tanh(a + b*x))^n,x)`

output

```

-(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^n*((3*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*ex
p(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^4)/(8*b^4*(50*n + 3
5*n^2 + 10*n^3 + n^4 + 24)) - (x^4*(11*n + 6*n^2 + n^3 + 6))/(50*n + 35*n^
2 + 10*n^3 + n^4 + 24) + (3*n*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2
*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3)/(4*b^3*(50*n
+ 35*n^2 + 10*n^3 + n^4 + 24)) + (n*x^3*(log(2/(exp(2*a)*exp(2*b*x) + 1))
- log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)*(3*n + n
^2 + 2))/(2*b*(50*n + 35*n^2 + 10*n^3 + n^4 + 24)) + (3*n*x^2*(n + 1)*(log
(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(
2*b*x) + 1)) + 2*b*x)^2)/(4*b^2*(50*n + 35*n^2 + 10*n^3 + n^4 + 24)))

```

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(\tanh(a + bx))^n dx = \int \operatorname{atanh}(\tanh(bx + a))^n x^3 dx$$

input

```
int(x^3*atanh(tanh(b*x+a))^n,x)
```

output

```
int(atanh(tanh(a + b*x))^n*x**3,x)
```

### 3.268 $\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1867
Mathematica [A] (verified)	1867
Rubi [A] (verified)	1868
Maple [A] (verified)	1869
Fricas [B] (verification not implemented)	1870
Sympy [F]	1870
Maxima [A] (verification not implemented)	1871
Giac [A] (verification not implemented)	1872
Mupad [B] (verification not implemented)	1872
Reduce [F]	1873

#### Optimal result

Integrand size = 13, antiderivative size = 82

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)} - \frac{2x \operatorname{arctanh}(\tanh(a + bx))^{2+n}}{b^2(1+n)(2+n)} + \frac{2 \operatorname{arctanh}(\tanh(a + bx))^{3+n}}{b^3(1+n)(2+n)(3+n)}$$

output

$$x^2 \operatorname{arctanh}(\tanh(bx+a))^{1+n} / b / (1+n) - 2 * x * \operatorname{arctanh}(\tanh(bx+a))^{2+n} / b^2 / (1+n) / (2+n) + 2 * \operatorname{arctanh}(\tanh(bx+a))^{3+n} / b^3 / (1+n) / (2+n) / (3+n)$$

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.87

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n} (b^2(6 + 5n + n^2) x^2 - 2b(3 + n)x \operatorname{arctanh}(\tanh(a + bx)) + 2 \operatorname{arctanh}(\tanh(a + bx)))}{b^3(1+n)(2+n)(3+n)}$$

input

```
Integrate[x^2 * ArcTanh[Tanh[a + b*x]]^n, x]
```



output

```
(ArcTanh[Tanh[a + b*x]]^(1 + n)*(b^2*(6 + 5*n + n^2)*x^2 - 2*b*(3 + n)*x*ArcTanh[Tanh[a + b*x]] + 2*ArcTanh[Tanh[a + b*x]]^2))/(b^3*(1 + n)*(2 + n)*(3 + n))
```

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {2599, 2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$\downarrow 2599$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{2 \int x \operatorname{arctanh}(\tanh(a + bx))^{n+1} dx}{b(n+1)}$$

$$\downarrow 2599$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b(n+2)} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^{n+2} dx}{b(n+2)} \right)}{b(n+1)}$$

$$\downarrow 2588$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b(n+2)} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^{n+2} dx}{b^2(n+2)} \right)}{b(n+1)}$$

$$\downarrow 15$$

$$\frac{x^2 \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{2 \left( \frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b(n+2)} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{n+3}}{b^2(n+2)(n+3)} \right)}{b(n+1)}$$

input

```
Int [x^2*ArcTanh [Tanh [a + b*x]]^n, x]
```

output

$$(x^2 \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(1+n)}) / (b*(1+n)) - (2*((x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(2+n)}) / (b*(2+n)) - \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(3+n)} / (b^2*(2+n)*(3+n)))) / (b*(1+n))$$

### Defintions of rubi rules used

rule 15

$$\operatorname{Int}[(a_.)*(x_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a*(x^{(m+1)}) / (m+1), x] /; \operatorname{FreeQ}[\{a, m\}, x] \ \&\& \operatorname{NeQ}[m, -1]$$

rule 2588

$$\operatorname{Int}[(u_)^{(m_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{c = \operatorname{Simplify}[D[u, x]]\}, \operatorname{Simp}[1/c \operatorname{Subst}[\operatorname{Int}[x^m, x], x, u], x]] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, x]$$

rule 2599

$$\operatorname{Int}[(u_)^{(m_.)}*(v_)^{(n_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{a = \operatorname{Simplify}[D[u, x]], b = \operatorname{Simplify}[D[v, x]]\}, \operatorname{Simp}[u^{(m+1)}*(v^n / (a*(m+1))), x] - \operatorname{Simp}[b*(n / (a*(m+1))) \operatorname{Int}[u^{(m+1)}*v^{(n-1)}, x], x] /; \operatorname{NeQ}[b*u - a*v, 0] /; \operatorname{FreeQ}[\{m, n\}, x] \ \&\& \operatorname{PiecewiseLinearQ}[u, v, x] \ \&\& \operatorname{NeQ}[m, -1] \ \&\& ((\operatorname{LtQ}[m, -1] \ \&\& \operatorname{GtQ}[n, 0]) \ \&\& !(\operatorname{ILtQ}[m+n, -2] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0]))) \ || (\operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{LeQ}[n, m]) \ || (\operatorname{IGtQ}[n, 0] \ \&\& !\operatorname{IntegerQ}[m]) \ || (\operatorname{ILtQ}[m, 0] \ \&\& !\operatorname{IntegerQ}[n]))$$

### Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.99

method	result
parallelrisc	$\frac{-6 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^n \operatorname{arctanh}(\operatorname{tanh}(bx+a))x^2b^2 + 6 \operatorname{arctanh}(\operatorname{tanh}(bx+a))^n x \operatorname{arctanh}(\operatorname{tanh}(bx+a))^2 b - x^2 \operatorname{arctanh}(\operatorname{tanh}(bx+a))}{b^2}$
default	$\frac{x^3 e^{n \ln(\operatorname{arctanh}(\operatorname{tanh}(bx+a)))}}{3+n} + \frac{n(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)x^2 e^{n \ln(\operatorname{arctanh}(\operatorname{tanh}(bx+a)))}}{b(n^2+5n+6)} + \frac{2(\operatorname{arctanh}(\operatorname{tanh}(bx+a)) - bx)}{b}$
risc	Expression too large to display

input

$$\operatorname{int}(x^2 \operatorname{arctanh}(\operatorname{tanh}(b*x+a))^n, x, \operatorname{method}=\_RETURNVERBOSE)$$

output

```
-(-6*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))*x^2*b^2+6*arctanh(tanh(b*x+a))^n*x*arctanh(tanh(b*x+a))^2*b-x^2*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b^2*n^2-5*x^2*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b^2*n+2*x*arctanh(tanh(b*x+a))^2*arctanh(tanh(b*x+a))^n*b*n-2*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))^3)/b^3/(n^3+6*n^2+11*n+6)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs.  $2(82) = 164$ .

Time = 0.08 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.05

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{(2a^2bnx - (b^3n^2 + 3b^3n + 2b^3)x^3 - 2a^3 - (ab^2n^2 + ab^2n)x^2) \cosh(n \log(bx + a)) + (2a^2bnx - (b^3n^2 + 3b^3n + 2b^3)x^3 - 2a^3 - (ab^2n^2 + ab^2n)x^2) \sinh(n \log(bx + a))}{b^3n^3 + 6b^3n^2 + 11b^3n + 6b^3}$$

input

```
integrate(x^2*arctanh(tanh(b*x+a))^n,x, algorithm="fricas")
```

output

```
-((2*a^2*b*n*x - (b^3*n^2 + 3*b^3*n + 2*b^3)*x^3 - 2*a^3 - (a*b^2*n^2 + a*b^2*n)*x^2)*cosh(n*log(b*x + a)) + (2*a^2*b*n*x - (b^3*n^2 + 3*b^3*n + 2*b^3)*x^3 - 2*a^3 - (a*b^2*n^2 + a*b^2*n)*x^2)*sinh(n*log(b*x + a)))/(b^3*n^3 + 6*b^3*n^2 + 11*b^3*n + 6*b^3)
```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx = \begin{cases} \frac{x^3 \operatorname{atanh}^n(\tanh(a))}{3} \\ -\frac{x^2}{2b \operatorname{atanh}^2(\tanh(a+bx))} - \frac{x}{b^2 \operatorname{atanh}(\tanh(a+bx))} + \frac{\log(\operatorname{atanh}(\tanh(a+bx)))}{b^3} \\ \int \frac{x^2}{\operatorname{atanh}^2(\tanh(a+bx))} dx \\ \int \frac{x^2}{\operatorname{atanh}(\tanh(a+bx))} dx \\ \frac{b^2 n^2 x^2 \operatorname{atanh}(\tanh(a+bx)) \operatorname{atanh}^n(\tanh(a+bx))}{b^3 n^3 + 6b^3 n^2 + 11b^3 n + 6b^3} + \frac{5b^2 n x^2 \operatorname{atanh}(\tanh(a+bx)) \operatorname{atanh}^n(\tanh(a+bx))}{b^3 n^3 + 6b^3 n^2 + 11b^3 n + 6b^3} + \frac{6b^2 x^2 \operatorname{atanh}(\tanh(a+bx)) \operatorname{atanh}^n(\tanh(a+bx))}{b^3 n^3 + 6b^3 n^2 + 11b^3 n + 6b^3} \end{cases}$$

input `integrate(x**2*atanh(tanh(b*x+a))**n,x)`

output `Piecewise((x**3*atanh(tanh(a))**n/3, Eq(b, 0)), (-x**2/(2*b*atanh(tanh(a + b*x))**2) - x/(b**2*atanh(tanh(a + b*x))) + log(atanh(tanh(a + b*x)))/b**3, Eq(n, -3)), (Integral(x**2/atanh(tanh(a + b*x))**2, x), Eq(n, -2)), (Integral(x**2/atanh(tanh(a + b*x)), x), Eq(n, -1)), (b**2*n**2*x**2*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3) + 5*b**2*n*x**2*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3) + 6*b**2*x**2*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3) - 2*b*n*x*atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3) - 6*b*x*atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3) + 2*atanh(tanh(a + b*x))**3*atanh(tanh(a + b*x))**n/(b**3*n**3 + 6*b**3*n**2 + 11*b**3*n + 6*b**3), True))`

### Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{((n^2 + 3n + 2)b^3x^3 + (n^2 + n)ab^2x^2 - 2a^2bnx + 2a^3)(bx + a)^n}{(n^3 + 6n^2 + 11n + 6)b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^n,x, algorithm="maxima")`

output `((n^2 + 3*n + 2)*b^3*x^3 + (n^2 + n)*a*b^2*x^2 - 2*a^2*b*n*x + 2*a^3)*(b*x + a)^n/((n^3 + 6*n^2 + 11*n + 6)*b^3)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.71

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(bx + a)^n b^3 n^2 x^3 + (bx + a)^n a b^2 n^2 x^2 + 3(bx + a)^n b^3 n x^3 + (bx + a)^n a b^2 n x^2 + 2(bx + a)^n b^3 x^3 - 2(bx + a)^n a^2 b^2 x^2 + 2(bx + a)^n a^3 b}{b^3 n^3 + 6 b^3 n^2 + 11 b^3 n + 6 b^3}$$

input `integrate(x^2*arctanh(tanh(b*x+a))^n,x, algorithm="giac")`output `((b*x + a)^n*b^3*n^2*x^3 + (b*x + a)^n*a*b^2*n^2*x^2 + 3*(b*x + a)^n*b^3*n*x^3 + (b*x + a)^n*a*b^2*n*x^2 + 2*(b*x + a)^n*b^3*x^3 - 2*(b*x + a)^n*a^2*b^2*x^2 + 2*(b*x + a)^n*a^3*b)/(b^3*n^3 + 6*b^3*n^2 + 11*b^3*n + 6*b^3)`**Mupad [B] (verification not implemented)**

Time = 3.52 (sec) , antiderivative size = 304, normalized size of antiderivative = 3.71

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= - \left( \frac{\ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)}{2} - \frac{\ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right)}{2} \right)^n \left( \frac{\left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^3}{4b^3 (n^3 + 6n^2 + 11n + 6)} \right.$$

$$- \frac{x^3 (n^2 + 3n + 2)}{n^3 + 6n^2 + 11n + 6} + \frac{nx \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{2b^2 (n^3 + 6n^2 + 11n + 6)}$$

$$\left. + \frac{nx^2 (n + 1) \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{2b (n^3 + 6n^2 + 11n + 6)} \right)$$

input `int(x^2*atanh(tanh(a + b*x))^n,x)`

output

```

-(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 - log(2/(exp(2*
a)*exp(2*b*x) + 1))/2)^n*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2
*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^3/(4*b^3*(11*n + 6*n^2
+ n^3 + 6)) - (x^3*(3*n + n^2 + 2))/(11*n + 6*n^2 + n^3 + 6) + (n*x*(log(
2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2
*b*x) + 1)) + 2*b*x)^2)/(2*b^2*(11*n + 6*n^2 + n^3 + 6)) + (n*x^2*(n + 1)*
(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*
exp(2*b*x) + 1)) + 2*b*x))/(2*b*(11*n + 6*n^2 + n^3 + 6)))

```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\tanh(a + bx))^n dx = \int \operatorname{atanh}(\tanh(bx + a))^n x^2 dx$$

input

```
int(x^2*atanh(tanh(b*x+a))^n,x)
```

output

```
int(atanh(tanh(a + b*x))^n*x**2,x)
```

### 3.269 $\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1874
Mathematica [A] (verified)	1874
Rubi [A] (verified)	1875
Maple [A] (verified)	1876
Fricas [A] (verification not implemented)	1876
Sympy [F]	1877
Maxima [A] (verification not implemented)	1877
Giac [A] (verification not implemented)	1878
Mupad [B] (verification not implemented)	1878
Reduce [B] (verification not implemented)	1879

#### Optimal result

Integrand size = 11, antiderivative size = 48

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{x \operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{2+n}}{b^2(1+n)(2+n)}$$

output

$$x \operatorname{arctanh}(\tanh(bx+a))^{(1+n)} / b / (1+n) - \operatorname{arctanh}(\tanh(bx+a))^{(2+n)} / b^2 / (1+n) / (2+n)$$

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

$$\begin{aligned} & \int x \operatorname{arctanh}(\tanh(a + bx))^n dx \\ &= \frac{(b(2+n)x - \operatorname{arctanh}(\tanh(a + bx))) \operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b^2(1+n)(2+n)} \end{aligned}$$

input

$$\text{Integrate}[x \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^n, x]$$

output

$$((b*(2+n)*x - \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]) * \operatorname{ArcTanh}[\operatorname{Tanh}[a + b*x]]^{(1+n)}) / (b^2*(1+n)*(2+n))$$

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {2599, 2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$\downarrow 2599$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^{n+1} dx}{b(n+1)}$$

$$\downarrow 2588$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{\int \operatorname{arctanh}(\tanh(a + bx))^{n+1} d \operatorname{arctanh}(\tanh(a + bx))}{b^2(n+1)}$$

$$\downarrow 15$$

$$\frac{x \operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n+1)} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{n+2}}{b^2(n+1)(n+2)}$$

input `Int[x*ArcTanh[Tanh[a + b*x]]^n,x]`

output `(x*ArcTanh[Tanh[a + b*x]]^(1 + n))/(b*(1 + n)) - ArcTanh[Tanh[a + b*x]]^(2 + n)/(b^2*(1 + n)*(2 + n))`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`



rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.60

method	result
parallelsch	$-\frac{-2 \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))xb + \operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))^2 - x \operatorname{arctanh}(\tanh(bx+a))}{b^2(n^2+3n+2)}$
default	$\frac{x^2 e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{2+n} + \frac{n(\operatorname{arctanh}(\tanh(bx+a))-bx)x e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{b(n^2+3n+2)} - \frac{(\operatorname{arctanh}(\tanh(bx+a))-bx)^2 e^{n \ln(\operatorname{arctanh}(\tanh(bx+a)))}}{b^2(n^2+3n+2)}$
risch	$\frac{\left(\frac{1}{2}\right)^n \left(2 \ln(e^{bx+a}) - \frac{i\pi \operatorname{csgn}(ie^{2bx+2a})}{2} (-\operatorname{csgn}(ie^{2bx+2a}) + \operatorname{csgn}(ie^{bx+a}))^2 - \frac{i\pi \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)}{2} \left(-\operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) + \operatorname{csgn}\left(\frac{ie^{bx+a}}{e^{2bx+2a}+1}\right)\right)\right)}{2b(1+n)}$

input

```
int(x*arctanh(tanh(b*x+a))^n,x,method=_RETURNVERBOSE)
```

output

```
-((-2*arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))*x*b+arctanh(tanh(b*x+a))^n*arctanh(tanh(b*x+a))^2-x*arctanh(tanh(b*x+a))*arctanh(tanh(b*x+a))^n*b*n)/b^2/(n^2+3*n+2)
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.90

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(abnx + (b^2n + b^2)x^2 - a^2) \cosh(n \log(bx + a)) + (abnx + (b^2n + b^2)x^2 - a^2) \sinh(n \log(bx + a))}{b^2n^2 + 3b^2n + 2b^2}$$

input

```
integrate(x*arctanh(tanh(b*x+a))^n,x, algorithm="fricas")
```

output  $((a*b*n*x + (b^2*n + b^2)*x^2 - a^2)*\cosh(n*\log(b*x + a)) + (a*b*n*x + (b^2*n + b^2)*x^2 - a^2)*\sinh(n*\log(b*x + a)))/(b^2*n^2 + 3*b^2*n + 2*b^2)$

## Sympy [F]

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \begin{cases} \frac{x^2 \operatorname{atanh}^n(\tanh(a))}{2} \\ -\frac{x}{b \operatorname{atanh}(\tanh(a + bx))} + \frac{\log(\operatorname{atanh}(\tanh(a + bx)))}{b^2} \\ \int \frac{x}{\operatorname{atanh}(\tanh(a + bx))} dx \\ \frac{bnx \operatorname{atanh}(\tanh(a + bx)) \operatorname{atanh}^n(\tanh(a + bx))}{b^2 n^2 + 3b^2 n + 2b^2} + \frac{2bx \operatorname{atanh}(\tanh(a + bx)) \operatorname{atanh}^n(\tanh(a + bx))}{b^2 n^2 + 3b^2 n + 2b^2} - \frac{\operatorname{atanh}^2(\tanh(a + bx)) \operatorname{atanh}^n(\tanh(a + bx))}{b^2 n^2 + 3b^2 n + 2b^2} \end{cases}$$

input `integrate(x*atanh(tanh(b*x+a))**n,x)`

output `Piecewise((x**2*atanh(tanh(a))**n/2, Eq(b, 0)), (-x/(b*atanh(tanh(a + b*x))) + log(atanh(tanh(a + b*x)))/b**2, Eq(n, -2)), (Integral(x/atanh(tanh(a + b*x)), x), Eq(n, -1)), (b*n*x*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**2*n**2 + 3*b**2*n + 2*b**2) + 2*b*x*atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b**2*n**2 + 3*b**2*n + 2*b**2) - atanh(tanh(a + b*x))**2*atanh(tanh(a + b*x))**n/(b**2*n**2 + 3*b**2*n + 2*b**2), True))`

## Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.88

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{(b^2(n+1)x^2 + abnx - a^2)(bx + a)^n}{(n^2 + 3n + 2)b^2}$$

input `integrate(x*arctanh(tanh(b*x+a))^n,x, algorithm="maxima")`

output  $(b^2*(n + 1)*x^2 + a*b*n*x - a^2)*(b*x + a)^n/((n^2 + 3*n + 2)*b^2)$

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.58

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(bx + a)^n b^2 n x^2 + (bx + a)^n a b n x + (bx + a)^n b^2 x^2 - (bx + a)^n a^2}{b^2 n^2 + 3 b^2 n + 2 b^2}$$

input `integrate(x*arctanh(tanh(b*x+a))^n,x, algorithm="giac")`output `((b*x + a)^n*b^2*n*x^2 + (b*x + a)^n*a*b*n*x + (b*x + a)^n*b^2*x^2 - (b*x + a)^n*a^2)/(b^2*n^2 + 3*b^2*n + 2*b^2)`**Mupad [B] (verification not implemented)**

Time = 3.43 (sec) , antiderivative size = 205, normalized size of antiderivative = 4.27

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= - \left( \frac{\ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right)}{2} - \frac{\ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right)}{2} \right)^n \left( \frac{\left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)^2}{4b^2 (n^2 + 3n + 2)} \right.$$

$$\left. - \frac{x^2 (n + 1)}{n^2 + 3n + 2} + \frac{nx \left( \ln \left( \frac{2}{e^{2a} e^{2bx} + 1} \right) - \ln \left( \frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1} \right) + 2bx \right)}{2b (n^2 + 3n + 2)} \right)$$

input `int(x*atanh(tanh(a + b*x))^n,x)`output `-(log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)))/2 - log(2/(exp(2*a)*exp(2*b*x) + 1))/2)^n*((log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x)^2/(4*b^2*(3*n + n^2 + 2)) - (x^2*(n + 1))/(3*n + n^2 + 2) + (n*x*(log(2/(exp(2*a)*exp(2*b*x) + 1)) - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1)) + 2*b*x))/(2*b*(3*n + n^2 + 2)))`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int x \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{\operatorname{atanh}(\tanh(bx + a))^n \operatorname{atanh}(\tanh(bx + a)) (-\operatorname{atanh}(\tanh(bx + a)) + bnx + 2bx)}{b^2(n^2 + 3n + 2)}$$

input `int(x*atanh(tanh(b*x+a))^n,x)`output `(atanh(tanh(a + b*x))^n*atanh(tanh(a + b*x))*(- atanh(tanh(a + b*x)) + b  
*n*x + 2*b*x))/(b**2*(n**2 + 3*n + 2))`

### 3.270 $\int \operatorname{arctanh}(\tanh(a + bx))^n dx$

Optimal result	1880
Mathematica [A] (verified)	1880
Rubi [A] (verified)	1881
Maple [A] (verified)	1882
Fricas [A] (verification not implemented)	1882
Sympy [B] (verification not implemented)	1883
Maxima [A] (verification not implemented)	1883
Giac [A] (verification not implemented)	1884
Mupad [B] (verification not implemented)	1884
Reduce [B] (verification not implemented)	1885

#### Optimal result

Integrand size = 9, antiderivative size = 20

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)}$$

output

```
arctanh(tanh(b*x+a))^(1+n)/b/(1+n)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n}}{b(1+n)}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^n,x]
```

output

```
ArcTanh[Tanh[a + b*x]]^(1 + n)/(b*(1 + n))
```

**Rubi [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \operatorname{arctanh}(\tanh(a + bx))^n d\operatorname{arctanh}(\tanh(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{\operatorname{arctanh}(\tanh(a + bx))^{n+1}}{b(n + 1)}$$

input `Int[ArcTanh[Tanh[a + b*x]]^n,x]`

output `ArcTanh[Tanh[a + b*x]]^(1 + n)/(b*(1 + n))`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

method	result
derivativdivides	$\frac{\operatorname{arctanh}(\tanh(bx+a))^{1+n}}{b(1+n)}$
default	$\frac{\operatorname{arctanh}(\tanh(bx+a))^{1+n}}{b(1+n)}$
parallelrisch	$\frac{\operatorname{arctanh}(\tanh(bx+a))^n \operatorname{arctanh}(\tanh(bx+a))}{b(1+n)}$
risch	$\frac{\left(\frac{1}{2}\right)^n \left(2 \ln(e^{bx+a}) - \frac{i\pi \operatorname{csgn}(ie^{2bx+2a}) \left(-\operatorname{csgn}(ie^{2bx+2a}) + \operatorname{csgn}(ie^{bx+a})\right)^2}{2} - i\pi \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right) \left(-\operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}+1}\right)\right)\right)}{2b(1+n)}$

input `int(arctanh(tanh(b*x+a))^n,x,method=_RETURNVERBOSE)`output `arctanh(tanh(b*x+a))^(1+n)/b/(1+n)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.95

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \frac{(bx + a) \cosh(n \log(bx + a)) + (bx + a) \sinh(n \log(bx + a))}{bn + b}$$

input `integrate(arctanh(tanh(b*x+a))^n,x, algorithm="fricas")`output `((b*x + a)*cosh(n*log(b*x + a)) + (b*x + a)*sinh(n*log(b*x + a)))/(b*n + b)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs.  $2(15) = 30$ .

Time = 0.28 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \begin{cases} \frac{x}{\operatorname{atanh}(\tanh(a))} & \text{for } b = 0 \wedge n = -1 \\ x \operatorname{atanh}^n(\tanh(a)) & \text{for } b = 0 \\ \frac{\log(\operatorname{atanh}(\tanh(a + bx)))}{b} & \text{for } n = -1 \\ \frac{\operatorname{atanh}(\tanh(a + bx)) \operatorname{atanh}^n(\tanh(a + bx))}{bn + b} & \text{otherwise} \end{cases}$$

input `integrate(atanh(tanh(b*x+a))**n,x)`

output `Piecewise((x/atanh(tanh(a)), Eq(b, 0) & Eq(n, -1)), (x*atanh(tanh(a))**n, Eq(b, 0)), (log(atanh(tanh(a + b*x)))/b, Eq(n, -1)), (atanh(tanh(a + b*x))*atanh(tanh(a + b*x))**n/(b*n + b), True))`

**Maxima [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{(bx + a)(bx + a)^n}{b(n + 1)}$$

input `integrate(arctanh(tanh(b*x+a))^n,x, algorithm="maxima")`

output `(b*x + a)*(b*x + a)^n/(b*(n + 1))`



**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.40

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{(bx + a)^n bx + (bx + a)^n a}{bn + b}$$

input `integrate(arctanh(tanh(b*x+a))^n,x, algorithm="giac")`output `((b*x + a)^n*b*x + (b*x + a)^n*a)/(b*n + b)`**Mupad [B] (verification not implemented)**

Time = 3.36 (sec) , antiderivative size = 121, normalized size of antiderivative = 6.05

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx$$

$$= \left(\frac{1}{2}\right)^n \left( \frac{x}{n+1} - \frac{\frac{\ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right)}{2} - \frac{\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right)}{2} + bx}{b(n+1)} \right) \left( \ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} + 1}\right) - \ln\left(\frac{2}{e^{2a} e^{2bx} + 1}\right) \right)^n$$

input `int(atanh(tanh(a + b*x))^n,x)`output `(1/2)^n*(x/(n + 1) - (log(2/(exp(2*a)*exp(2*b*x) + 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1))/2 + b*x)/(b*(n + 1)))*log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) + 1) - log(2/(exp(2*a)*exp(2*b*x) + 1)))^n`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.25

$$\int \operatorname{arctanh}(\tanh(a + bx))^n dx = \frac{\operatorname{atanh}(\tanh(bx + a))^n \operatorname{atanh}(\tanh(bx + a))}{b(n + 1)}$$

input `int(atanh(tanh(b*x+a))^n,x)`

output `(atanh(tanh(a + b*x))**n*atanh(tanh(a + b*x)))/(b*(n + 1))`

### 3.271 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x} dx$

Optimal result	1886
Mathematica [A] (verified)	1886
Rubi [A] (verified)	1887
Maple [F]	1888
Fricas [F]	1888
Sympy [F]	1888
Maxima [F]	1889
Giac [F]	1889
Mupad [F(-1)]	1889
Reduce [F]	1890

#### Optimal result

Integrand size = 13, antiderivative size = 64

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx$$

$$= \frac{\operatorname{arctanh}(\tanh(a + bx))^{1+n} \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, -\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{(1 + n)(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output `arctanh(tanh(b*x+a))^(1+n)*hypergeom([1, 1+n], [2+n], -arctanh(tanh(b*x+a))/(b*x-arctanh(tanh(b*x+a))))/(1+n)/(b*x-arctanh(tanh(b*x+a)))`

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx$$

$$= \frac{\operatorname{arctanh}(\tanh(a + bx))^n \left(\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)^{-n} \operatorname{Hypergeometric2F1}\left(-n, -n, 1 - n, 1 - \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)}{n}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^n/x,x]`

output  $(\text{ArcTanh}[\text{Tanh}[a + b*x]]^n * \text{Hypergeometric2F1}[-n, -n, 1 - n, 1 - \text{ArcTanh}[\text{Tanh}[a + b*x]]/(b*x)]) / (n * (\text{ArcTanh}[\text{Tanh}[a + b*x]]/(b*x))^n)$

### Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\text{arctanh}(\tanh(a + bx))^n}{x} dx$$

↓ 2595

$$\frac{\text{arctanh}(\tanh(a + bx))^{n+1} \text{Hypergeometric2F1}\left(1, n + 1, n + 2, -\frac{\text{arctanh}(\tanh(a + bx))}{bx - \text{arctanh}(\tanh(a + bx))}\right)}{(n + 1)(bx - \text{arctanh}(\tanh(a + bx)))}$$

input  $\text{Int}[\text{ArcTanh}[\text{Tanh}[a + b*x]]^n/x, x]$

output  $(\text{ArcTanh}[\text{Tanh}[a + b*x]]^{(1 + n)} * \text{Hypergeometric2F1}[1, 1 + n, 2 + n, -(\text{ArcTanh}[\text{Tanh}[a + b*x]]/(b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]))]) / ((1 + n) * (b*x - \text{ArcTanh}[\text{Tanh}[a + b*x]]))$

### Defintions of rubi rules used

rule 2595  $\text{Int}[(v_)^n/(u_), x\_Symbol] \rightarrow \text{With}\{a = \text{Simplify}[D[u, x]], b = \text{Simplify}[D[v, x]]\}, \text{Simp}[(v^{(n + 1)})/((n + 1)*(b*u - a*v)) * \text{Hypergeometric2F1}[1, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; \text{NeQ}[b*u - a*v, 0] /; \text{PiecewiseLinearQ}[u, v, x] \&\& !\text{IntegerQ}[n]$

**Maple [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(bx + a))^n}{x} dx$$

input `int(arctanh(tanh(b*x+a))^n/x,x)`

output `int(arctanh(tanh(b*x+a))^n/x,x)`

**Fricas [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x,x, algorithm="fricas")`

output `integral(arctanh(tanh(b*x + a))^n/x, x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{atanh}^n(\tanh(a + bx))}{x} dx$$

input `integrate(atanh(tanh(b*x+a))**n/x,x)`

output `Integral(atanh(tanh(a + b*x))**n/x, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^n/x, x)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x,x, algorithm="giac")`

output `integrate(arctanh(tanh(b*x + a))^n/x, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^n}{x} dx$$

input `int(atanh(tanh(a + b*x))^n/x,x)`

output `int(atanh(tanh(a + b*x))^n/x, x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a))^n}{x} dx$$

input `int(atanh(tanh(b*x+a))^n/x,x)`

output `int(atanh(tanh(a + b*x))**n/x,x)`

### 3.272 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^2} dx$

Optimal result	1891
Mathematica [A] (verified)	1891
Rubi [A] (verified)	1892
Maple [F]	1893
Fricas [F]	1893
Sympy [F]	1894
Maxima [F]	1894
Giac [F]	1894
Mupad [F(-1)]	1895
Reduce [F]	1895

#### Optimal result

Integrand size = 13, antiderivative size = 71

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = -\frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x} + \frac{b \operatorname{arctanh}(\tanh(a + bx))^n \operatorname{Hypergeometric2F1}\left(1, n, 1 + n, -\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{bx - \operatorname{arctanh}(\tanh(a + bx))}$$

output

```
-arctanh(tanh(b*x+a))^n/x+b*arctanh(tanh(b*x+a))^n*hypergeom([1, n], [1+n],
-arctanh(tanh(b*x+a))/(b*x-arctanh(tanh(b*x+a))))/(b*x-arctanh(tanh(b*x+a)
))
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \frac{\operatorname{arctanh}(\tanh(a + bx))^n \left(\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)^{-n} \operatorname{Hypergeometric2F1}\left(1 - n, -n, 2 - n, 1 - \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)}{(-1 + n)x}$$

input

```
Integrate[ArcTanh[Tanh[a + b*x]]^n/x^2,x]
```



output

```
(ArcTanh[Tanh[a + b*x]]^n*Hypergeometric2F1[1 - n, -n, 2 - n, 1 - ArcTanh[
Tanh[a + b*x]]/(b*x)]/((-1 + n)*x*(ArcTanh[Tanh[a + b*x]]/(b*x))^n)
```

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {2599, 2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx$$

$$\downarrow 2599$$

$$bn \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{n-1}}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x}$$

$$\downarrow 2595$$

$$\frac{b \operatorname{arctanh}(\tanh(a + bx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, -\frac{\operatorname{arctanh}(\tanh(a + bx))}{bx - \operatorname{arctanh}(\tanh(a + bx))}\right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x}$$

input

```
Int[ArcTanh[Tanh[a + b*x]]^n/x^2,x]
```

output

```
-(ArcTanh[Tanh[a + b*x]]^n/x) + (b*ArcTanh[Tanh[a + b*x]]^n*Hypergeometric
2F1[1, n, 1 + n, -(ArcTanh[Tanh[a + b*x]]/(b*x - ArcTanh[Tanh[a + b*x]]))])
)/(b*x - ArcTanh[Tanh[a + b*x]])
```

## Definitions of rubi rules used

rule 2595 `Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[(v^(n + 1)/((n + 1)*(b*u - a*v))*Hypergeometric2F1[1, n + 1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x] && !IntegerQ[n]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

## Maple [F]

$$\int \frac{\operatorname{arctanh}(\tanh(bx + a))^n}{x^2} dx$$

input `int(arctanh(tanh(b*x+a))^n/x^2,x)`

output `int(arctanh(tanh(b*x+a))^n/x^2,x)`

## Fricas [F]

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^2} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^2,x, algorithm="fricas")`

output `integral(arctanh(tanh(b*x + a))^n/x^2, x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \int \frac{\operatorname{atanh}^n(\tanh(a + bx))}{x^2} dx$$

input `integrate(atanh(tanh(b*x+a))**n/x**2,x)`

output `Integral(atanh(tanh(a + b*x))**n/x**2, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^2} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^2,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^n/x^2, x)`

**Giac [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^2} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^2,x, algorithm="giac")`

output `integrate(arctanh(tanh(b*x + a))^n/x^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^n}{x^2} dx$$

input `int(atanh(tanh(a + b*x))^n/x^2,x)`output `int(atanh(tanh(a + b*x))^n/x^2, x)`**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^2} dx = \frac{-\operatorname{atanh}(\tanh(bx + a))^n + \left( \int \frac{\operatorname{atanh}(\tanh(bx+a))^n}{\operatorname{atanh}(\tanh(bx+a))x} dx \right) bnx}{x}$$

input `int(atanh(tanh(b*x+a))^n/x^2,x)`output `( - atanh(tanh(a + b*x))**n + int(atanh(tanh(a + b*x))**n/(atanh(tanh(a + b*x))*x),x)*b*n*x)/x`

### 3.273 $\int \frac{\operatorname{arctanh}(\tanh(a+bx))^n}{x^3} dx$

Optimal result	1896
Mathematica [A] (verified)	1896
Rubi [A] (verified)	1897
Maple [F]	1898
Fricas [F]	1899
Sympy [F]	1899
Maxima [F]	1899
Giac [F]	1900
Mupad [F(-1)]	1900
Reduce [F]	1900

#### Optimal result

Integrand size = 13, antiderivative size = 101

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx$$

$$= -\frac{bn \operatorname{arctanh}(\tanh(a + bx))^{-1+n}}{2x} - \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{2x^2}$$

$$+ \frac{b^2 n \operatorname{arctanh}(\tanh(a + bx))^{-1+n} \operatorname{Hypergeometric2F1}\left(1, -1 + n, n, -\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{2(bx - \operatorname{arctanh}(\tanh(a + bx)))}$$

output

```
-1/2*b*n*arctanh(tanh(b*x+a))(-1+n)/x-1/2*arctanh(tanh(b*x+a))n/x2+b2*n*arctanh(tanh(b*x+a))(-1+n)*hypergeom([1, -1+n], [n], -arctanh(tanh(b*x+a))/(b*x-arctanh(tanh(b*x+a))))/(2*b*x-2*arctanh(tanh(b*x+a)))
```

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.66

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx$$

$$= \frac{\operatorname{arctanh}(\tanh(a + bx))^n \left(\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)^{-n} \operatorname{Hypergeometric2F1}\left(2 - n, -n, 3 - n, 1 - \frac{\operatorname{arctanh}(\tanh(a+bx))}{bx}\right)}{(-2 + n)x^2}$$

input `Integrate[ArcTanh[Tanh[a + b*x]]^n/x^3,x]`

output `(ArcTanh[Tanh[a + b*x]]^n*Hypergeometric2F1[2 - n, -n, 3 - n, 1 - ArcTanh[Tanh[a + b*x]]/(b*x)]/((-2 + n)*x^2*(ArcTanh[Tanh[a + b*x]]/(b*x))^n)`

### Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.97, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {2599, 2599, 2595}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}bn \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{n-1}}{x^2} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{2x^2} \\
 & \quad \downarrow \text{2599} \\
 & \frac{1}{2}bn \left( -b(1-n) \int \frac{\operatorname{arctanh}(\tanh(a + bx))^{n-2}}{x} dx - \frac{\operatorname{arctanh}(\tanh(a + bx))^{n-1}}{x} \right) - \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{2x^2} \\
 & \quad \downarrow \text{2595} \\
 & \frac{1}{2}bn \left( \frac{b \operatorname{arctanh}(\tanh(a + bx))^{n-1} \operatorname{Hypergeometric2F1}\left(1, n-1, n, -\frac{\operatorname{arctanh}(\tanh(a+bx))}{bx - \operatorname{arctanh}(\tanh(a+bx))}\right)}{bx - \operatorname{arctanh}(\tanh(a + bx))} - \frac{\operatorname{arctanh}(\tanh(a + bx))^{n-1}}{x} \right) - \\
 & \quad \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{2x^2}
 \end{aligned}$$

input `Int[ArcTanh[Tanh[a + b*x]]^n/x^3,x]`

output

```
-1/2*ArcTanh[Tanh[a + b*x]]^n/x^2 + (b*n*(-(ArcTanh[Tanh[a + b*x]]^(-1 + n)
)/x) + (b*ArcTanh[Tanh[a + b*x]]^(-1 + n)*Hypergeometric2F1[1, -1 + n, n,
-(ArcTanh[Tanh[a + b*x]]/(b*x - ArcTanh[Tanh[a + b*x]])]))/(b*x - ArcTanh[
Tanh[a + b*x]])))/2
```

### Defintions of rubi rules used

rule 2595

```
Int[(v_)^(n_)/(u_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[
D[v, x]]}, Simp[(v^(n + 1)/((n + 1)*(b*u - a*v))*Hypergeometric2F1[1, n +
1, n + 2, (-a)*(v/(b*u - a*v))], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinea
rQ[u, v, x] && !IntegerQ[n]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Sim
plify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1
))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0]] /; FreeQ[{m, n]
, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0
] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ
[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILt
Q[m, 0] && !IntegerQ[n]))
```

### Maple [F]

$$\int \frac{\operatorname{arctanh}(\tanh(bx + a))^n}{x^3} dx$$

input

```
int(arctanh(tanh(b*x+a))^n/x^3,x)
```

output

```
int(arctanh(tanh(b*x+a))^n/x^3,x)
```

**Fricas [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^3} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^3,x, algorithm="fricas")`

output `integral(arctanh(tanh(b*x + a))^n/x^3, x)`

**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx = \int \frac{\operatorname{atanh}^n(\tanh(a + bx))}{x^3} dx$$

input `integrate(atanh(tanh(b*x+a))**n/x**3,x)`

output `Integral(atanh(tanh(a + b*x))**n/x**3, x)`

**Maxima [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^3} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^3,x, algorithm="maxima")`

output `integrate(arctanh(tanh(b*x + a))^n/x^3, x)`



**Giac [F]**

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx = \int \frac{\operatorname{artanh}(\tanh(bx + a))^n}{x^3} dx$$

input `integrate(arctanh(tanh(b*x+a))^n/x^3,x, algorithm="giac")`

output `integrate(arctanh(tanh(b*x + a))^n/x^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx = \int \frac{\operatorname{atanh}(\tanh(a + bx))^n}{x^3} dx$$

input `int(atanh(tanh(a + b*x))^n/x^3,x)`

output `int(atanh(tanh(a + b*x))^n/x^3, x)`

**Reduce [F]**

$$\begin{aligned} & \int \frac{\operatorname{arctanh}(\tanh(a + bx))^n}{x^3} dx \\ &= \frac{-\operatorname{atanh}(\tanh(bx + a))^n + \left( \int \frac{\operatorname{atanh}(\tanh(bx+a))^n}{\operatorname{atanh}(\tanh(bx+a))x^2} dx \right) b n x^2}{2x^2} \end{aligned}$$

input `int(atanh(tanh(b*x+a))^n/x^3,x)`

output `( - atanh(tanh(a + b*x))**n + int(atanh(tanh(a + b*x))**n/(atanh(tanh(a + b*x))*x**2), x)*b*n*x**2)/(2*x**2)`

### 3.274 $\int x^m \coth^{-1}(\tanh(a + bx)) dx$

Optimal result	1901
Mathematica [A] (verified)	1901
Rubi [A] (verified)	1902
Maple [A] (verified)	1903
Fricas [C] (verification not implemented)	1903
Sympy [F]	1904
Maxima [A] (verification not implemented)	1904
Giac [B] (verification not implemented)	1905
Mupad [B] (verification not implemented)	1905
Reduce [B] (verification not implemented)	1906

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx = -\frac{bx^{2+m}}{2 + 3m + m^2} + \frac{x^{1+m} \coth^{-1}(\tanh(a + bx))}{1 + m}$$

output

```
-b*x^(2+m)/(m^2+3*m+2)+x^(1+m)*arccoth(tanh(b*x+a))/(1+m)
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx = x^m \left( \frac{bx^2}{2 + m} + \frac{x(-bx + \coth^{-1}(\tanh(a + bx)))}{1 + m} \right)$$

input

```
Integrate[x^m*ArcCoth[Tanh[a + b*x]],x]
```

output

```
x^m*((b*x^2)/(2 + m) + (x*(-(b*x) + ArcCoth[Tanh[a + b*x]]))/(1 + m))
```

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{x^{m+1} \coth^{-1}(\tanh(a + bx))}{m + 1} - \frac{b \int x^{m+1} dx}{m + 1}$$

$$\downarrow \text{15}$$

$$\frac{x^{m+1} \coth^{-1}(\tanh(a + bx))}{m + 1} - \frac{bx^{m+2}}{(m + 1)(m + 2)}$$

input `Int[x^m*ArcCoth[Tanh[a + b*x]],x]`

output `-((b*x^(2 + m))/((1 + m)*(2 + m))) + (x^(1 + m)*ArcCoth[Tanh[a + b*x]])/(1 + m)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.32

method	result
parallelrisch	$-\frac{-x x^m \operatorname{arccoth}(\tanh(bx+a))m-2 \operatorname{arccoth}(\tanh(bx+a))x^m x+b x^m x^2}{m^2+3m+2}$
risch	$\frac{x x^m \ln(e^{bx+a})}{1+m} - \frac{x \left( 4i\pi+i\pi \operatorname{csgn}(ie^{2bx+2a})^3 m+4bx+i\pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}(ie^{2bx+2a})m+4i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a+1}}\right)^3 +2i\pi \right)}{m^2+3m+2}$

input `int(x^m*arccoth(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `-(-x*x^m*arccoth(tanh(b*x+a))*m-2*arccoth(tanh(b*x+a))*x^m*x+b*x^m*x^2)/(m^2+3*m+2)`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 81, normalized size of antiderivative = 2.19

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx$$

$$= \frac{(i\pi(m+2)x + 2(bm+b)x^2 + 2(am+2a)x) \cosh(m \log(x)) + (i\pi(m+2)x + 2(bm+b)x^2 + 2(am+2a)x) \sinh(m \log(x))}{2(m^2 + 3m + 2)}$$

input `integrate(x^m*arccoth(tanh(b*x+a)),x, algorithm="fricas")`

output `1/2*((I*pi*(m+2)*x + 2*(b*m + b)*x^2 + 2*(a*m + 2*a)*x)*cosh(m*log(x)) + (I*pi*(m+2)*x + 2*(b*m + b)*x^2 + 2*(a*m + 2*a)*x)*sinh(m*log(x)))/(m^2 + 3*m + 2)`

**Sympy [F]**

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx$$

$$= \begin{cases} b \log(x) - \frac{\operatorname{acoth}(\tanh(a+bx))}{x} & \text{for } m = -2 \\ \int \frac{\operatorname{acoth}(\tanh(a+bx))}{x} dx & \text{for } m = -1 \\ -\frac{bx^2x^m}{m^2+3m+2} + \frac{mxx^m \operatorname{acoth}(\tanh(a+bx))}{m^2+3m+2} + \frac{2xx^m \operatorname{acoth}(\tanh(a+bx))}{m^2+3m+2} & \text{otherwise} \end{cases}$$

input `integrate(x**m*acoth(tanh(b*x+a)), x)`

output `Piecewise((b*log(x) - acoth(tanh(a + b*x))/x, Eq(m, -2)), (Integral(acoth(tanh(a + b*x))/x, x), Eq(m, -1)), (-b*x**2*x**m/(m**2 + 3*m + 2) + m*x*x**m*acoth(tanh(a + b*x))/(m**2 + 3*m + 2) + 2*x*x**m*acoth(tanh(a + b*x))/(m**2 + 3*m + 2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx = -\frac{bx^2x^m}{(m+2)(m+1)} + \frac{x^{m+1} \operatorname{arccoth}(\tanh(bx + a))}{m+1}$$

input `integrate(x^m*arccoth(tanh(b*x+a)), x, algorithm="maxima")`

output `-b*x^2*x^m/((m + 2)*(m + 1)) + x^(m + 1)*arccoth(tanh(b*x + a))/(m + 1)`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. 2(37) = 74.

Time = 0.12 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.43

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx = \frac{x^{m+1} \log\left(-\frac{\frac{e^{(2bx+2a)+1}+1}{e^{(2bx+2a)-1}}}{\frac{e^{(2bx+2a)+1}-1}{e^{(2bx+2a)-1}}}\right)}{2(m+1)} - \frac{bx^{m+2}}{(m+2)(m+1)}$$

input `integrate(x^m*arccoth(tanh(b*x+a)),x, algorithm="giac")`

output `1/2*x^(m+1)*log(-((e^(2*b*x+2*a)+1)/(e^(2*b*x+2*a)-1)+1)/((e^(2*b*x+2*a)+1)/(e^(2*b*x+2*a)-1)-1))/(m+1)-b*x^(m+2)/((m+2)*(m+1))`

**Mupad [B] (verification not implemented)**

Time = 3.94 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.59

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx = \frac{2bx^m x^2(m+1)}{2m^2+6m+4} - \frac{xx^m(m+2)\left(\ln\left(-\frac{2}{e^{2a}e^{2bx}-1}\right) - \ln\left(\frac{2e^{2a}e^{2bx}}{e^{2a}e^{2bx}-1}\right) + 2bx\right)}{2m^2+6m+4}$$

input `int(x^m*acoth(tanh(a+b*x)),x)`

output `(2*b*x^m*x^2*(m+1))/(6*m+2*m^2+4)-(x*x^m*(m+2)*(log(-2/(exp(2*a)*exp(2*b*x)-1))-log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x)-1))+2*b*x))/(6*m+2*m^2+4)`

**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int x^m \coth^{-1}(\tanh(a + bx)) dx$$
$$= \frac{x^m x (\operatorname{acoth}(\tanh(bx + a)) m + 2 \operatorname{acoth}(\tanh(bx + a)) + bx)}{m^2 + 3m + 2}$$

input `int(x^m*acoth(tanh(b*x+a)),x)`output `(x**m*x*(acoth(tanh(a + b*x))*m + 2*acoth(tanh(a + b*x)) + b*x))/(m**2 + 3*m + 2)`

### 3.275 $\int x^2 \operatorname{arctanh}(\coth(a + bx)) dx$

Optimal result	1907
Mathematica [A] (verified)	1907
Rubi [A] (verified)	1908
Maple [A] (verified)	1909
Fricas [C] (verification not implemented)	1909
Sympy [B] (verification not implemented)	1910
Maxima [A] (verification not implemented)	1910
Giac [B] (verification not implemented)	1911
Mupad [B] (verification not implemented)	1911
Reduce [F]	1911

#### Optimal result

Integrand size = 11, antiderivative size = 23

$$\int x^2 \operatorname{arctanh}(\coth(a + bx)) dx = -\frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(\coth(a + bx))$$

output `-1/12*b*x^4+1/3*x^3*arctanh(coth(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^2 \operatorname{arctanh}(\coth(a + bx)) dx = -\frac{1}{12}x^3(bx - 4\operatorname{arctanh}(\coth(a + bx)))$$

input `Integrate[x^2*ArcTanh[Coth[a + b*x]],x]`

output `-1/12*(x^3*(b*x - 4*ArcTanh[Coth[a + b*x]]))`



**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{1}{3} x^3 \operatorname{arctanh}(\operatorname{coth}(a + bx)) - \frac{b \int x^3 dx}{3}$$

$$\downarrow \text{15}$$

$$\frac{1}{3} x^3 \operatorname{arctanh}(\operatorname{coth}(a + bx)) - \frac{bx^4}{12}$$

input `Int[x^2*ArcTanh[Coth[a + b*x]],x]`

output `-1/12*(b*x^4) + (x^3*ArcTanh[Coth[a + b*x]])/3`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

**Maple [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result
parallelrisch	$-\frac{x^3(bx-4 \operatorname{arctanh}(\coth(bx+a)))}{12}$
default	$-\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\coth(bx+a))}{3}$
parts	$-\frac{bx^4}{12} + \frac{x^3 \operatorname{arctanh}(\coth(bx+a))}{3}$
risch	$\frac{x^3 \ln(e^{bx+a})}{3} - \frac{bx^4}{12} + \frac{i\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}-1}\right)^2}{12} - \frac{i\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^3}{6} + \frac{i\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)}{6}$

input `int(x^2*arctanh(coth(b*x+a)),x,method=_RETURNVERBOSE)`

output `-1/12*x^3*(b*x-4*arctanh(coth(b*x+a)))`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\coth(a + bx)) dx = \frac{1}{4} bx^4 + \frac{1}{6} i \pi x^3 + \frac{1}{3} ax^3$$

input `integrate(x^2*arctanh(coth(b*x+a)),x, algorithm="fricas")`

output `1/4*b*x^4 + 1/6*I*pi*x^3 + 1/3*a*x^3`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs.  $2(19) = 38$ .

Time = 4.80 (sec) , antiderivative size = 76, normalized size of antiderivative = 3.30

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \begin{cases} \frac{x^3 \operatorname{atanh}(\operatorname{coth}(bx + \log(-e^{-bx})))}{3} & \text{for } a = \log(-e^{-bx}) \\ \frac{x^3 \operatorname{atanh}(\operatorname{coth}(bx + \log(e^{-bx})))}{3} & \text{for } a = \log(e^{-bx}) \\ -\frac{bx^4}{12} + \frac{x^3 \operatorname{atanh}\left(\frac{1}{\tanh(a + bx)}\right)}{3} & \text{otherwise} \end{cases}$$

input `integrate(x**2*atanh(coth(b*x+a)),x)`

output `Piecewise((x**3*atanh(coth(b*x + log(-exp(-b*x))))/3, Eq(a, log(-exp(-b*x)))), (x**3*atanh(coth(b*x + log(exp(-b*x))))/3, Eq(a, log(exp(-b*x))))), (-b*x**4/12 + x**3*atanh(1/tanh(a + b*x))/3, True))`

**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{12} bx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(\operatorname{coth}(bx + a))$$

input `integrate(x^2*arctanh(coth(b*x+a)),x, algorithm="maxima")`

output `-1/12*b*x^4 + 1/3*x^3*arctanh(coth(b*x + a))`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(19) = 38$ .

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 3.09

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{12} bx^4 + \frac{1}{6} x^3 \log\left(-\frac{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} + 1}{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} - 1}\right)$$

input `integrate(x^2*arctanh(coth(b*x+a)),x, algorithm="giac")`

output `-1/12*b*x^4 + 1/6*x^3*log(-((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) + 1)/((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \frac{x^3 \operatorname{atanh}(\operatorname{coth}(a + bx))}{3} - \frac{bx^4}{12}$$

input `int(x^2*atanh(coth(a + b*x)),x)`

output `(x^3*atanh(coth(a + b*x)))/3 - (b*x^4)/12`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a)) x^2 dx$$

input `int(x^2*atanh(coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x))*x**2,x)`

### 3.276 $\int x \operatorname{arctanh}(\coth(a + bx)) dx$

Optimal result	1912
Mathematica [A] (verified)	1912
Rubi [A] (verified)	1913
Maple [A] (verified)	1914
Fricas [C] (verification not implemented)	1914
Sympy [B] (verification not implemented)	1915
Maxima [A] (verification not implemented)	1915
Giac [B] (verification not implemented)	1916
Mupad [B] (verification not implemented)	1916
Reduce [F]	1916

#### Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x \operatorname{arctanh}(\coth(a + bx)) dx = -\frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(\coth(a + bx))$$

output `-1/6*b*x^3+1/2*x^2*arctanh(coth(b*x+a))`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x \operatorname{arctanh}(\coth(a + bx)) dx = -\frac{1}{6}x^2(bx - 3\operatorname{arctanh}(\coth(a + bx)))$$

input `Integrate[x*ArcTanh[Coth[a + b*x]],x]`

output `-1/6*(x^2*(b*x - 3*ArcTanh[Coth[a + b*x]]))`

**Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6795, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(\coth(a + bx)) dx$$

$$\downarrow 6795$$

$$\frac{1}{2}b \int -x^2 dx + \frac{1}{2}x^2 \operatorname{arctanh}(\coth(a + bx))$$

$$\downarrow 15$$

$$\frac{1}{2}x^2 \operatorname{arctanh}(\coth(a + bx)) - \frac{bx^3}{6}$$

input `Int[x*ArcTanh[Coth[a + b*x]],x]`

output `-1/6*(b*x^3) + (x^2*ArcTanh[Coth[a + b*x]])/2`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6795 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]]*(d_.)]*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result
parallelrisch	$-\frac{x^2(bx-3 \operatorname{arctanh}(\coth(bx+a)))}{6}$
default	$-\frac{bx^3}{6} + \frac{x^2 \operatorname{arctanh}(\coth(bx+a))}{2}$
parts	$-\frac{bx^3}{6} + \frac{x^2 \operatorname{arctanh}(\coth(bx+a))}{2}$
risch	$\frac{x^2 \ln(e^{bx+a})}{2} - \frac{bx^3}{6} + \frac{i\pi x^2 \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}-1}\right)^2 \operatorname{csgn}(ie^{2bx+2a})}{8} - \frac{i\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^3}{4} - \frac{i\pi x^2 \operatorname{csgn}(ie^{2bx+2a})^3}{8}$

input `int(x*arctanh(coth(b*x+a)),x,method=_RETURNVERBOSE)`output `-1/6*x^2*(b*x-3*arctanh(coth(b*x+a)))`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\coth(a + bx)) dx = \frac{1}{3} bx^3 + \frac{1}{4} i \pi x^2 + \frac{1}{2} ax^2$$

input `integrate(x*arctanh(coth(b*x+a)),x, algorithm="fricas")`output `1/3*b*x^3 + 1/4*I*pi*x^2 + 1/2*a*x^2`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs.  $2(19) = 38$ .

Time = 2.89 (sec) , antiderivative size = 76, normalized size of antiderivative = 3.30

$$\int x \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \begin{cases} \frac{x^2 \operatorname{atanh}(\operatorname{coth}(bx + \log(-e^{-bx})))}{2} & \text{for } a = \log(-e^{-bx}) \\ \frac{x^2 \operatorname{atanh}(\operatorname{coth}(bx + \log(e^{-bx})))}{2} & \text{for } a = \log(e^{-bx}) \\ -\frac{bx^3}{6} + \frac{x^2 \operatorname{atanh}\left(\frac{1}{\tanh(a + bx)}\right)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*atanh(coth(b*x+a)),x)`

output `Piecewise((x**2*atanh(coth(b*x + log(-exp(-b*x))))/2, Eq(a, log(-exp(-b*x)))), (x**2*atanh(coth(b*x + log(exp(-b*x))))/2, Eq(a, log(exp(-b*x))))), (-b*x**3/6 + x**2*atanh(1/tanh(a + b*x))/2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{6} bx^3 + \frac{1}{2} x^2 \operatorname{arctanh}(\operatorname{coth}(bx + a))$$

input `integrate(x*arctanh(coth(b*x+a)),x, algorithm="maxima")`

output `-1/6*b*x^3 + 1/2*x^2*arctanh(coth(b*x + a))`



**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(19) = 38$ .

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 3.09

$$\int x \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{6} bx^3 + \frac{1}{4} x^2 \log \left( -\frac{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} + 1}{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} - 1} \right)$$

input `integrate(x*arctanh(coth(b*x+a)),x, algorithm="giac")`

output `-1/6*b*x^3 + 1/4*x^2*log(-((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) + 1)/((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) - 1))`

**Mupad [B] (verification not implemented)**

Time = 3.50 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \frac{x^2 \operatorname{atanh}(\operatorname{coth}(a + bx))}{2} - \frac{bx^3}{6}$$

input `int(x*atanh(coth(a + b*x)),x)`

output `(x^2*atanh(coth(a + b*x)))/2 - (b*x^3)/6`

**Reduce [F]**

$$\int x \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a)) x dx$$

input `int(x*atanh(coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x))*x,x)`

### 3.277 $\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx$

Optimal result	1917
Mathematica [A] (verified)	1917
Rubi [A] (verified)	1918
Maple [A] (verified)	1919
Fricas [C] (verification not implemented)	1919
Sympy [B] (verification not implemented)	1920
Maxima [A] (verification not implemented)	1920
Giac [B] (verification not implemented)	1921
Mupad [B] (verification not implemented)	1921
Reduce [B] (verification not implemented)	1921

#### Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))^2}{2b}$$

output `1/2*arctanh(coth(b*x+a))^2/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{bx^2}{2} + x \operatorname{arctanh}(\operatorname{coth}(a + bx))$$

input `Integrate[ArcTanh[Coth[a + b*x]],x]`

output `-1/2*(b*x^2) + x*ArcTanh[Coth[a + b*x]]`

**Rubi [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx$$

$$\downarrow 2588$$

$$\frac{\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) d\operatorname{arctanh}(\operatorname{coth}(a + bx))}{b}$$

$$\downarrow 15$$

$$\frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))^2}{2b}$$

input `Int[ArcTanh[Coth[a + b*x]],x]`

output `ArcTanh[Coth[a + b*x]]^2/(2*b)`

**Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x]] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))^2}{2b}$
default	$\frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))^2}{2b}$
parallelrisc	$-\frac{x(bx-2 \operatorname{arctanh}(\operatorname{coth}(bx+a)))}{2}$
parts	$-\frac{x^2b}{2} + x \operatorname{arctanh}(\operatorname{coth}(bx+a))$
risc	$x \ln(e^{bx+a}) - \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^3 x}{2} + \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}-1}\right)^2 x}{4} - \frac{i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)}{2}$

input `int(arctanh(coth(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*arctanh(coth(b*x+a))^2/b`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\operatorname{coth}(a+bx)) dx = \frac{1}{2}bx^2 + \frac{1}{2}i\pi x + ax$$

input `integrate(arctanh(coth(b*x+a)),x, algorithm="fricas")`output `1/2*b*x^2 + 1/2*I*pi*x + a*x`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 68 vs.  $2(12) = 24$ .

Time = 1.59 (sec) , antiderivative size = 68, normalized size of antiderivative = 4.25

$$\int \operatorname{arctanh}(\operatorname{coth}(a+bx)) dx = \begin{cases} x \operatorname{atanh}(\operatorname{coth}(a)) & \text{for } b = 0 \\ x \operatorname{atanh}(\operatorname{coth}(bx + \log(-e^{-bx}))) & \text{for } a = \log(-e^{-bx}) \\ x \operatorname{atanh}(\operatorname{coth}(bx + \log(e^{-bx}))) & \text{for } a = \log(e^{-bx}) \\ \frac{\operatorname{atanh}^2\left(\frac{1}{\tanh(a+bx)}\right)}{2b} & \text{otherwise} \end{cases}$$

input `integrate(atanh(coth(b*x+a)),x)`

output `Piecewise((x*atanh(coth(a)), Eq(b, 0)), (x*atanh(coth(b*x + log(-exp(-b*x))))), Eq(a, log(-exp(-b*x))))), (x*atanh(coth(b*x + log(exp(-b*x))))), Eq(a, log(exp(-b*x))))), (atanh(1/tanh(a + b*x))**2/(2*b), True))`

**Maxima [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{2}bx^2 + x \operatorname{artanh}(\operatorname{coth}(bx + a))$$

input `integrate(arctanh(coth(b*x+a)),x, algorithm="maxima")`

output `-1/2*b*x^2 + x*arctanh(coth(b*x + a))`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 69 vs.  $2(14) = 28$ .

Time = 0.12 (sec) , antiderivative size = 69, normalized size of antiderivative = 4.31

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}x \log\left(-\frac{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} + 1}{\frac{e^{(2bx+2a)+1}}{e^{(2bx+2a)-1}} - 1}\right)$$

input `integrate(arctanh(coth(b*x+a)),x, algorithm="giac")`

output `-1/2*b*x^2 + 1/2*x*log(-((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) + 1)/  
((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) - 1))`

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = x \operatorname{atanh}(\operatorname{coth}(a + bx)) - \frac{bx^2}{2}$$

input `int(atanh(coth(a + b*x)),x)`

output `x*atanh(coth(a + b*x)) - (b*x^2)/2`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \operatorname{arctanh}(\operatorname{coth}(a + bx)) dx = \frac{\operatorname{atanh}(\operatorname{coth}(bx + a))^2}{2b}$$

input `int(atanh(coth(b*x+a)),x)`

output `atanh(coth(a + b*x))**2/(2*b)`

### 3.278 $\int \frac{\operatorname{arctanh}(\operatorname{coth}(a+bx))}{x} dx$

Optimal result	1922
Mathematica [A] (verified)	1922
Rubi [A] (verified)	1923
Maple [A] (verified)	1924
Fricas [C] (verification not implemented)	1924
Sympy [F]	1924
Maxima [A] (verification not implemented)	1925
Giac [C] (verification not implemented)	1925
Mupad [B] (verification not implemented)	1926
Reduce [F]	1926

#### Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a+bx))}{x} dx = bx - (bx - \operatorname{arctanh}(\operatorname{coth}(a+bx))) \log(x)$$

output `b*x-(b*x-arctanh(coth(b*x+a)))*ln(x)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a+bx))}{x} dx = bx + (-bx + \operatorname{arctanh}(\operatorname{coth}(a+bx))) \log(x)$$

input `Integrate[ArcTanh[Coth[a + b*x]]/x,x]`

output `b*x + (-b*x) + ArcTanh[Coth[a + b*x]]*Log[x]`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x} dx$$

$$\downarrow \text{2589}$$

$$bx - (bx - \operatorname{arctanh}(\operatorname{coth}(a + bx))) \int \frac{1}{x} dx$$

$$\downarrow \text{14}$$

$$bx - \log(x)(bx - \operatorname{arctanh}(\operatorname{coth}(a + bx)))$$

input `Int[ArcTanh[Coth[a + b*x]]/x,x]`

output `b*x - (b*x - ArcTanh[Coth[a + b*x]])*Log[x]`

**Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`



**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

method	result
default	$\ln(x) \operatorname{arctanh}(\coth(bx + a)) - b(x \ln(x) - x)$
parts	$\ln(x) \operatorname{arctanh}(\coth(bx + a)) - b(x \ln(x) - x)$
risch	$\ln(x) \ln(e^{bx+a}) - \ln(x)xb + bx + \frac{i\pi \left( 2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^2 - 2 \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^3 + \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right) \operatorname{csgn}\left(\frac{ie^{2bx+2a}}{e^{2bx+2a}-1}\right) \right)}{2}$

input `int(arctanh(coth(b*x+a))/x,x,method=_RETURNVERBOSE)`output `ln(x)*arctanh(coth(b*x+a))-b*(x*ln(x)-x)`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\coth(a + bx))}{x} dx = bx + \frac{1}{2} (i\pi + 2a) \log(x)$$

input `integrate(arctanh(coth(b*x+a))/x,x, algorithm="fricas")`output `b*x + 1/2*(I*pi + 2*a)*log(x)`**Sympy [F]**

$$\int \frac{\operatorname{arctanh}(\coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\coth(a + bx))}{x} dx$$

input `integrate(atanh(coth(b*x+a))/x,x)`

output `Integral(atanh(coth(a + b*x))/x, x)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x} dx = -b\left(x + \frac{a}{b}\right) \log(x) + b\left(x + \frac{a \log(x)}{b}\right) + \operatorname{arctanh}(\operatorname{coth}(bx + a)) \log(x)$$

input `integrate(arctanh(coth(b*x+a))/x,x, algorithm="maxima")`

output `-b*(x + a/b)*log(x) + b*(x + a*log(x)/b) + arctanh(coth(b*x + a))*log(x)`

### Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x} dx = bx + \frac{1}{2} (i\pi + 2a) \log(x)$$

input `integrate(arctanh(coth(b*x+a))/x,x, algorithm="giac")`

output `b*x + 1/2*(I*pi + 2*a)*log(x)`

**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.81

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x} dx = bx - \ln(x) \left( \frac{\ln\left(-\frac{2}{e^{2a} e^{2bx} - 1}\right)}{2} - \frac{\ln\left(\frac{2e^{2a} e^{2bx}}{e^{2a} e^{2bx} - 1}\right)}{2} + bx \right)$$

input `int(atanh(coth(a + b*x))/x,x)`

output `b*x - log(x)*(log(-2/(exp(2*a)*exp(2*b*x) - 1))/2 - log((2*exp(2*a)*exp(2*b*x))/(exp(2*a)*exp(2*b*x) - 1))/2 + b*x)`

**Reduce [F]**

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\operatorname{coth}(bx + a))}{x} dx$$

input `int(atanh(coth(b*x+a))/x,x)`

output `int(atanh(coth(a + b*x))/x,x)`

### 3.279 $\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^2} dx$

Optimal result	1927
Mathematica [A] (verified)	1927
Rubi [A] (verified)	1928
Maple [A] (verified)	1929
Fricas [C] (verification not implemented)	1929
Sympy [B] (verification not implemented)	1930
Maxima [A] (verification not implemented)	1930
Giac [B] (verification not implemented)	1931
Mupad [B] (verification not implemented)	1931
Reduce [B] (verification not implemented)	1931

#### Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^2} dx = -\frac{\operatorname{arctanh}(\coth(a+bx))}{x} + b \log(x)$$

output `-arctanh(coth(b*x+a))/x+b*ln(x)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^2} dx = b - \frac{\operatorname{arctanh}(\coth(a+bx))}{x} + b \log(x)$$

input `Integrate[ArcTanh[Coth[a + b*x]]/x^2,x]`

output `b - ArcTanh[Coth[a + b*x]]/x + b*Log[x]`

**Rubi [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx$$

↓ 2599

$$b \int \frac{1}{x} dx - \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x}$$

↓ 14

$$b \log(x) - \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x}$$

input

```
Int[ArcTanh[Coth[a + b*x]]/x^2,x]
```

output

```
-(ArcTanh[Coth[a + b*x]]/x) + b*Log[x]
```

**Defintions of rubi rules used**

rule 14

```
Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

method	result
default	$-\frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))}{x} + b \ln(x)$
parts	$-\frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))}{x} + b \ln(x)$
parallelrisch	$\frac{\ln(x)xb - \operatorname{arctanh}(\operatorname{coth}(bx+a))}{x}$
risch	$-\frac{\ln(e^{bx+a})}{x} + \frac{-2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 + 2i\pi - 2i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^2 + i\pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}(ie^{2bx+2a}) + 2i\pi}{2x}$

input `int(arctanh(coth(b*x+a))/x^2,x,method=_RETURNVERBOSE)`output `-arctanh(coth(b*x+a))/x+b*ln(x)`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = \frac{-i\pi + 2bx \log(x) - 2a}{2x}$$

input `integrate(arctanh(coth(b*x+a))/x^2,x, algorithm="fricas")`output `1/2*(-I*pi + 2*b*x*log(x) - 2*a)/x`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 68 vs.  $2(14) = 28$ .

Time = 3.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 4.00

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = \begin{cases} -\frac{\operatorname{atanh}(\operatorname{coth}(bx + \log(-e^{-bx})))}{x} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\operatorname{atanh}(\operatorname{coth}(bx + \log(e^{-bx})))}{x} & \text{for } a = \log(e^{-bx}) \\ b \log(x) - \frac{\operatorname{atanh}\left(\frac{1}{\tanh(a + bx)}\right)}{x} & \text{otherwise} \end{cases}$$

input `integrate(atanh(coth(b*x+a))/x**2,x)`

output `Piecewise((-atanh(coth(b*x + log(-exp(-b*x))))/x, Eq(a, log(-exp(-b*x)))), (-atanh(coth(b*x + log(exp(-b*x))))/x, Eq(a, log(exp(-b*x)))), (b*log(x) - atanh(1/tanh(a + b*x))/x, True))`

**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = b \log(x) - \frac{\operatorname{artanh}(\operatorname{coth}(bx + a))}{x}$$

input `integrate(arctanh(coth(b*x+a))/x^2,x, algorithm="maxima")`

output `b*log(x) - arctanh(coth(b*x + a))/x`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 70 vs.  $2(17) = 34$ .

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 4.12

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = b \log(|x|) - \frac{\log\left(-\frac{e^{\frac{2bx+2a}{-1}+1}}{e^{\frac{2bx+2a}{-1}-1}}\right)}{2x}$$

input `integrate(arctanh(coth(b*x+a))/x^2,x, algorithm="giac")`

output `b*log(abs(x)) - 1/2*log(-((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) + 1) / ((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) - 1))/x`

**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = b \ln(x) - \frac{\operatorname{atanh}(\operatorname{coth}(a + bx))}{x}$$

input `int(atanh(coth(a + b*x))/x^2,x)`

output `b*log(x) - atanh(coth(a + b*x))/x`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^2} dx = \frac{-\operatorname{atanh}(\operatorname{coth}(bx + a)) + \log(x) bx}{x}$$

input `int(atanh(coth(b*x+a))/x^2,x)`

output `( - atanh(coth(a + b*x)) + log(x)*b*x)/x`



### 3.280 $\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^3} dx$

Optimal result	1932
Mathematica [A] (verified)	1932
Rubi [A] (verified)	1933
Maple [A] (verified)	1934
Fricas [C] (verification not implemented)	1934
Sympy [B] (verification not implemented)	1935
Maxima [A] (verification not implemented)	1935
Giac [B] (verification not implemented)	1936
Mupad [B] (verification not implemented)	1936
Reduce [B] (verification not implemented)	1936

#### Optimal result

Integrand size = 11, antiderivative size = 23

$$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^3} dx = -\frac{b}{2x} - \frac{\operatorname{arctanh}(\coth(a+bx))}{2x^2}$$

output `-1/2*b/x-1/2*arctanh(coth(b*x+a))/x^2`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arctanh}(\coth(a+bx))}{x^3} dx = -\frac{bx + \operatorname{arctanh}(\coth(a+bx))}{2x^2}$$

input `Integrate[ArcTanh[Coth[a + b*x]]/x^3,x]`

output `-1/2*(b*x + ArcTanh[Coth[a + b*x]])/x^2`

**Rubi [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx$$

↓ 2599

$$\frac{1}{2}b \int \frac{1}{x^2} dx - \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{2x^2}$$

↓ 15

$$-\frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{2x^2} - \frac{b}{2x}$$

input

```
Int[ArcTanh[Coth[a + b*x]]/x^3,x]
```

output

```
-1/2*b/x - ArcTanh[Coth[a + b*x]]/(2*x^2)
```

**Defintions of rubi rules used**

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]
```

rule 2599

```
Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0] && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))
```

**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
default	$-\frac{b}{2x} - \frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))}{2x^2}$
parallelrisch	$\frac{-bx - \operatorname{arctanh}(\operatorname{coth}(bx+a))}{2x^2}$
parts	$-\frac{b}{2x} - \frac{\operatorname{arctanh}(\operatorname{coth}(bx+a))}{2x^2}$
risch	$-\frac{\ln(e^{bx+a})}{2x^2} - \frac{4bx + 2i\pi \operatorname{csgn}(ie^{bx+a}) \operatorname{csgn}(ie^{2bx+2a})^2 - 2i\pi + 2i\pi \operatorname{csgn}\left(\frac{i}{e^{2bx+2a}-1}\right)^2 - i\pi \operatorname{csgn}(ie^{bx+a})^2 \operatorname{csgn}(ie^{2bx+2a})}{2x^2}$

input `int(arctanh(coth(b*x+a))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*b/x-1/2*arctanh(coth(b*x+a))/x^2`

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = \frac{-i\pi - 4bx - 2a}{4x^2}$$

input `integrate(arctanh(coth(b*x+a))/x^3,x, algorithm="fricas")`

output `1/4*(-I*pi - 4*b*x - 2*a)/x^2`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 80 vs.  $2(19) = 38$ .

Time = 5.34 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.48

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = \begin{cases} -\frac{\operatorname{atanh}(\operatorname{coth}(bx + \log(-e^{-bx})))}{2x^2} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\operatorname{atanh}(\operatorname{coth}(bx + \log(e^{-bx})))}{2x^2} & \text{for } a = \log(e^{-bx}) \\ -\frac{b}{2x} - \frac{\operatorname{atanh}\left(\frac{1}{\tanh(a + bx)}\right)}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate(atanh(coth(b*x+a))/x**3,x)`

output `Piecewise((-atanh(coth(b*x + log(-exp(-b*x))))/(2*x**2), Eq(a, log(-exp(-b*x)))), (-atanh(coth(b*x + log(exp(-b*x))))/(2*x**2), Eq(a, log(exp(-b*x)))), (-b/(2*x) - atanh(1/tanh(a + b*x))/(2*x**2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = -\frac{b}{2x} - \frac{\operatorname{artanh}(\operatorname{coth}(bx + a))}{2x^2}$$

input `integrate(arctanh(coth(b*x+a))/x^3,x, algorithm="maxima")`

output `-1/2*b/x - 1/2*arctanh(coth(b*x + a))/x^2`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(19) = 38$ .

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 3.09

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = -\frac{b}{2x} - \frac{\log\left(-\frac{e^{\frac{(2bx+2a)+1}{e^{(2bx+2a)-1}}+1}}{e^{\frac{(2bx+2a)+1}{e^{(2bx+2a)-1}}-1}}\right)}{4x^2}$$

input `integrate(arctanh(coth(b*x+a))/x^3,x, algorithm="giac")`

output `-1/2*b/x - 1/4*log(-((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) + 1)/((e^(2*b*x + 2*a) + 1)/(e^(2*b*x + 2*a) - 1) - 1))/x^2`

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = -\frac{\operatorname{atanh}(\operatorname{coth}(a + bx)) + bx}{2x^2}$$

input `int(atanh(coth(a + b*x))/x^3,x)`

output `-(atanh(coth(a + b*x)) + b*x)/(2*x^2)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arctanh}(\operatorname{coth}(a + bx))}{x^3} dx = \frac{-\operatorname{atanh}(\operatorname{coth}(bx + a)) - bx}{2x^2}$$

input `int(atanh(coth(b*x+a))/x^3,x)`

output `( - (atanh(coth(a + b*x)) + b*x))/(2*x**2)`

### 3.281 $\int \operatorname{arctanh}(\cosh(x)) dx$

Optimal result	1937
Mathematica [A] (verified)	1937
Rubi [C] (verified)	1938
Maple [A] (verified)	1940
Fricas [B] (verification not implemented)	1940
Sympy [F]	1941
Maxima [A] (verification not implemented)	1941
Giac [F]	1941
Mupad [F(-1)]	1942
Reduce [F]	1942

#### Optimal result

Integrand size = 3, antiderivative size = 27

$$\int \operatorname{arctanh}(\cosh(x)) dx = -2x\operatorname{arctanh}(e^x) + x\operatorname{arctanh}(\cosh(x)) \\ - \operatorname{PolyLog}(2, -e^x) + \operatorname{PolyLog}(2, e^x)$$

output

```
-2*x*arctanh(exp(x))+x*arctanh(cosh(x))-polylog(2,-exp(x))+polylog(2,exp(x))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.44

$$\int \operatorname{arctanh}(\cosh(x)) dx = x\operatorname{arctanh}(\cosh(x)) + x(\log(1 - e^x) - \log(1 + e^x)) \\ - \operatorname{PolyLog}(2, -e^x) + \operatorname{PolyLog}(2, e^x)$$

input

```
Integrate[ArcTanh[Cosh[x]],x]
```

output

```
x*ArcTanh[Cosh[x]] + x*(Log[1 - E^x] - Log[1 + E^x]) - PolyLog[2, -E^x] + PolyLog[2, E^x]
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.48, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 2.333$ , Rules used = {6825, 25, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arctanh}(\cosh(x)) dx \\
 & \quad \downarrow 6825 \\
 & x \operatorname{arctanh}(\cosh(x)) - \int -x \operatorname{csch}(x) dx \\
 & \quad \downarrow 25 \\
 & \int x \operatorname{csch}(x) dx + x \operatorname{arctanh}(\cosh(x)) \\
 & \quad \downarrow 3042 \\
 & x \operatorname{arctanh}(\cosh(x)) + \int ix \operatorname{csc}(ix) dx \\
 & \quad \downarrow 26 \\
 & x \operatorname{arctanh}(\cosh(x)) + i \int x \operatorname{csc}(ix) dx \\
 & \quad \downarrow 4670 \\
 & x \operatorname{arctanh}(\cosh(x)) + i \left( i \int \log(1 - e^x) dx - i \int \log(1 + e^x) dx + 2ix \operatorname{arctanh}(e^x) \right) \\
 & \quad \downarrow 2715 \\
 & x \operatorname{arctanh}(\cosh(x)) + i \left( i \int e^{-x} \log(1 - e^x) de^x - i \int e^{-x} \log(1 + e^x) de^x + 2ix \operatorname{arctanh}(e^x) \right) \\
 & \quad \downarrow 2838 \\
 & x \operatorname{arctanh}(\cosh(x)) + i(2ix \operatorname{arctanh}(e^x) + i \operatorname{PolyLog}(2, -e^x) - i \operatorname{PolyLog}(2, e^x))
 \end{aligned}$$

input `Int[ArcTanh[Cosh[x]],x]`

output `x*ArcTanh[Cosh[x]] + I*((2*I)*x*ArcTanh[E^x] + I*PolyLog[2, -E^x] - I*PolyLog[2, E^x])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x) + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x]], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6825 `Int[ArcTanh[u_], x_Symbol] := Simp[x*ArcTanh[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/(1 - u^2)), x], x] /; InverseFunctionFreeQ[u, x]`



**Maple [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.33

method	result
default	$x \operatorname{arctanh}(\cosh(x)) + x \ln(1 - e^x) + \operatorname{polylog}(2, e^x) - x \ln(1 + e^x) - \operatorname{polylog}(2, -e^x)$
parts	$x \operatorname{arctanh}(\cosh(x)) + x \ln(1 - e^x) + \operatorname{polylog}(2, e^x) - x \ln(1 + e^x) - \operatorname{polylog}(2, -e^x)$
risch	$-\operatorname{dilog}(1 + e^x) - x \ln(e^x - 1) - \frac{i\pi \operatorname{csgn}(i(e^x - 1)^2) \operatorname{csgn}(ie^{-x}(e^x - 1)^2)^2 x}{4} + \frac{i\pi \operatorname{csgn}(ie^{-x}(e^x - 1)^2)^2 x}{2} + \frac{i\pi \operatorname{csgn}(ie^{-x}(e^x - 1)^2)^2 x}{4}$

input `int(arctanh(cosh(x)), x, method=_RETURNVERBOSE)`

output `x*arctanh(cosh(x))+x*ln(1-exp(x))+polylog(2,exp(x))-x*ln(1+exp(x))-polylog(2,-exp(x))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 58 vs.  $2(22) = 44$ .

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{2} x \log\left(-\frac{\cosh(x) + 1}{\cosh(x) - 1}\right) - x \log(\cosh(x) + \sinh(x) + 1) + x \log(-\cosh(x) - \sinh(x) + 1) + \operatorname{Li}_2(\cosh(x) + \sinh(x)) - \operatorname{Li}_2(-\cosh(x) - \sinh(x))$$

input `integrate(arctanh(cosh(x)), x, algorithm="fricas")`

output `1/2*x*log(-(cosh(x) + 1)/(cosh(x) - 1)) - x*log(cosh(x) + sinh(x) + 1) + x*log(-cosh(x) - sinh(x) + 1) + dilog(cosh(x) + sinh(x)) - dilog(-cosh(x) - sinh(x))`

**Sympy [F]**

$$\int \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{atanh}(\cosh(x)) dx$$

input `integrate(atanh(cosh(x)),x)`

output `Integral(atanh(cosh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22

$$\int \operatorname{arctanh}(\cosh(x)) dx = x \operatorname{artanh}(\cosh(x)) - x \log(e^x + 1) \\ + x \log(-e^x + 1) - \operatorname{Li}_2(-e^x) + \operatorname{Li}_2(e^x)$$

input `integrate(arctanh(cosh(x)),x, algorithm="maxima")`

output `x*arctanh(cosh(x)) - x*log(e^x + 1) + x*log(-e^x + 1) - dilog(-e^x) + dilog(e^x)`

**Giac [F]**

$$\int \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{artanh}(\cosh(x)) dx$$

input `integrate(arctanh(cosh(x)),x, algorithm="giac")`

output `integrate(arctanh(cosh(x)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{atanh}(\cosh(x)) dx$$

input `int(atanh(cosh(x)), x)`output `int(atanh(cosh(x)), x)`**Reduce [F]**

$$\int \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{atanh}(\cosh(x)) dx$$

input `int(atanh(cosh(x)), x)`output `int(atanh(cosh(x)), x)`

### 3.282 $\int x \operatorname{arctanh}(\cosh(x)) dx$

Optimal result	1943
Mathematica [A] (verified)	1943
Rubi [C] (verified)	1944
Maple [C] (warning: unable to verify)	1946
Fricas [B] (verification not implemented)	1947
Sympy [F]	1948
Maxima [A] (verification not implemented)	1948
Giac [F]	1948
Mupad [F(-1)]	1949
Reduce [F]	1949

#### Optimal result

Integrand size = 5, antiderivative size = 51

$$\int x \operatorname{arctanh}(\cosh(x)) dx = -x^2 \operatorname{arctanh}(e^x) + \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) - x \operatorname{PolyLog}(2, -e^x) + x \operatorname{PolyLog}(2, e^x) + \operatorname{PolyLog}(3, -e^x) - \operatorname{PolyLog}(3, e^x)$$

output

$-x^2 \operatorname{arctanh}(\exp(x)) + 1/2 x^2 \operatorname{arctanh}(\cosh(x)) - x \operatorname{polylog}(2, -\exp(x)) + x \operatorname{polylog}(2, \exp(x)) + \operatorname{polylog}(3, -\exp(x)) - \operatorname{polylog}(3, \exp(x))$

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.37

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) + \frac{1}{2} x^2 \log(1 - e^x) - \frac{1}{2} x^2 \log(1 + e^x) - x \operatorname{PolyLog}(2, -e^x) + x \operatorname{PolyLog}(2, e^x) + \operatorname{PolyLog}(3, -e^x) - \operatorname{PolyLog}(3, e^x)$$

input

`Integrate[x*ArcTanh[Cosh[x]], x]`

output

```
(x^2*ArcTanh[Cosh[x]])/2 + (x^2*Log[1 - E^x])/2 - (x^2*Log[1 + E^x])/2 - x
*PolyLog[2, -E^x] + x*PolyLog[2, E^x] + PolyLog[3, -E^x] - PolyLog[3, E^x]
```

**Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.35, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.600$ , Rules used = {6827, 25, 3042, 26, 4670, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{arctanh}(\cosh(x)) dx \\
 & \quad \downarrow \text{6827} \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) - \frac{1}{2} \int -x^2 \operatorname{csch}(x) dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \int x^2 \operatorname{csch}(x) dx + \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) + \frac{1}{2} \int i x^2 \csc(ix) dx \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) + \frac{1}{2} i \int x^2 \csc(ix) dx \\
 & \quad \downarrow \text{4670} \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) + \frac{1}{2} i \left( 2i \int x \log(1 - e^x) dx - 2i \int x \log(1 + e^x) dx + 2ix^2 \operatorname{arctanh}(e^x) \right) \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(\cosh(x)) + \\
 & \frac{1}{2} i \left( -2i \left( \int \operatorname{PolyLog}(2, -e^x) dx - x \operatorname{PolyLog}(2, -e^x) \right) + 2i \left( \int \operatorname{PolyLog}(2, e^x) dx - x \operatorname{PolyLog}(2, e^x) \right) + 2ix^2 \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 2720 \\
 & \frac{1}{2}x^2 \operatorname{arctanh}(\cosh(x)) + \\
 & \frac{1}{2}i \left( -2i \left( \int e^{-x} \operatorname{PolyLog}(2, -e^x) dx - x \operatorname{PolyLog}(2, -e^x) \right) + 2i \left( \int e^{-x} \operatorname{PolyLog}(2, e^x) dx - x \operatorname{PolyLog}(2, e^x) \right) \right) \\
 & \downarrow 7143 \\
 & \frac{1}{2}x^2 \operatorname{arctanh}(\cosh(x)) + \\
 & \frac{1}{2}i (2ix^2 \operatorname{arctanh}(e^x) - 2i(\operatorname{PolyLog}(3, -e^x) - x \operatorname{PolyLog}(2, -e^x)) + 2i(\operatorname{PolyLog}(3, e^x) - x \operatorname{PolyLog}(2, e^x)))
 \end{aligned}$$

input `Int[x*ArcTanh[Cosh[x]], x]`

output `(x^2*ArcTanh[Cosh[x]])/2 + (I/2)*((2*I)*x^2*ArcTanh[E^x] - (2*I)*(-(x*PolyLog[2, -E^x]) + PolyLog[3, -E^x]) + (2*I)*(-(x*PolyLog[2, E^x]) + PolyLog[3, E^x]))`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6827 `Int[((a_.) + ArcTanh[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcTanh[u])/(d*(m + 1))), x] - Simp[b/(d*(m + 1)) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(1 - u^2)), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && FalseQ[PowerVariableExpn[u, m + 1, x]]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

## Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.40 (sec) , antiderivative size = 400, normalized size of antiderivative = 7.84

method	result
risch	$-\frac{x^2 \ln(e^x - 1)}{2} + \frac{i\pi \left( 2 \operatorname{csgn}\left(ie^{-x}(e^x - 1)^2\right)^2 - \operatorname{csgn}(i(1+e^x))^2 \operatorname{csgn}\left(i(1+e^x)^2\right) + 2 \operatorname{csgn}(i(1+e^x)) \operatorname{csgn}\left(i(1+e^x)^2\right)^2 - \operatorname{csgn}\left(i(1+e^x)^2\right) \right)}{2}$

input `int(x*arctanh(cosh(x)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/2*x^2*\ln(\exp(x)-1)+1/8*I*Pi*(2*csgn(I*\exp(-x))*(\exp(x)-1)^2-csgn(I*(1 \\ & +\exp(x)))^2*csgn(I*(1+\exp(x))^2)+2*csgn(I*(1+\exp(x)))*csgn(I*(1+\exp(x))^2) \\ & ^2-csgn(I*(1+\exp(x))^2)^3-csgn(I*(1+\exp(x))^2)*csgn(I*\exp(-x))*csgn(I*\exp( \\ & -x)*(1+\exp(x))^2)+csgn(I*(1+\exp(x))^2)*csgn(I*\exp(-x)*(1+\exp(x))^2)^2+csgn \\ & (I*(\exp(x)-1))^2*csgn(I*(\exp(x)-1)^2)-2*csgn(I*(\exp(x)-1))*csgn(I*(\exp(x)- \\ & 1)^2)^2+csgn(I*(\exp(x)-1)^2)^3+csgn(I*(\exp(x)-1)^2)*csgn(I*\exp(-x))*csgn(I \\ & *\exp(-x)*(\exp(x)-1)^2)-csgn(I*(\exp(x)-1)^2)*csgn(I*\exp(-x)*(\exp(x)-1)^2)^2 \\ & +csgn(I*\exp(-x))*csgn(I*\exp(-x)*(1+\exp(x))^2)^2-csgn(I*\exp(-x))*csgn(I*\exp \\ & (-x)*(\exp(x)-1)^2)^2-csgn(I*\exp(-x)*(1+\exp(x))^2)^3-csgn(I*\exp(-x)*(\exp(x) \\ & -1)^2)^3-2)*x^2-x*polylog(2,-\exp(x))+polylog(3,-\exp(x))+1/2*x^2*\ln(1-\exp(x) \\ & ))+x*polylog(2,\exp(x))-polylog(3,\exp(x)) \end{aligned}$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 88 vs.  $2(42) = 84$ .

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.73

$$\begin{aligned} \int x \operatorname{arctanh}(\cosh(x)) dx &= \frac{1}{4} x^2 \log\left(-\frac{\cosh(x)+1}{\cosh(x)-1}\right) - \frac{1}{2} x^2 \log(\cosh(x) + \sinh(x) + 1) \\ &+ \frac{1}{2} x^2 \log(-\cosh(x) - \sinh(x) + 1) \\ &+ x \operatorname{Li}_2(\cosh(x) + \sinh(x)) - x \operatorname{Li}_2(-\cosh(x) - \sinh(x)) \\ &- \operatorname{polylog}(3, \cosh(x) + \sinh(x)) \\ &+ \operatorname{polylog}(3, -\cosh(x) - \sinh(x)) \end{aligned}$$

input `integrate(x*arctanh(cosh(x)),x, algorithm="fricas")`

output 
$$\begin{aligned} & 1/4*x^2*\log(-(\cosh(x) + 1)/(\cosh(x) - 1)) - 1/2*x^2*\log(\cosh(x) + \sinh(x) \\ & + 1) + 1/2*x^2*\log(-\cosh(x) - \sinh(x) + 1) + x*dilog(\cosh(x) + \sinh(x)) - \\ & x*dilog(-\cosh(x) - \sinh(x)) - \operatorname{polylog}(3, \cosh(x) + \sinh(x)) + \operatorname{polylog}(3, - \\ & \cosh(x) - \sinh(x)) \end{aligned}$$



**Sympy [F]**

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \int x \operatorname{atanh}(\cosh(x)) dx$$

input `integrate(x*atanh(cosh(x)),x)`

output `Integral(x*atanh(cosh(x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.10

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{2} x^2 \operatorname{artanh}(\cosh(x)) - \frac{1}{2} x^2 \log(e^x + 1) + \frac{1}{2} x^2 \log(-e^x + 1) - x \operatorname{Li}_2(-e^x) + x \operatorname{Li}_2(e^x) + \operatorname{Li}_3(-e^x) - \operatorname{Li}_3(e^x)$$

input `integrate(x*arctanh(cosh(x)),x, algorithm="maxima")`

output `1/2*x^2*arctanh(cosh(x)) - 1/2*x^2*log(e^x + 1) + 1/2*x^2*log(-e^x + 1) - x*dilog(-e^x) + x*dilog(e^x) + polylog(3, -e^x) - polylog(3, e^x)`

**Giac [F]**

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \int x \operatorname{artanh}(\cosh(x)) dx$$

input `integrate(x*arctanh(cosh(x)),x, algorithm="giac")`

output `integrate(x*arctanh(cosh(x)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \int x \operatorname{atanh}(\cosh(x)) dx$$

input `int(x*atanh(cosh(x)),x)`output `int(x*atanh(cosh(x)), x)`**Reduce [F]**

$$\int x \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{atanh}(\cosh(x)) x dx$$

input `int(x*atanh(cosh(x)),x)`output `int(atanh(cosh(x))*x,x)`

### 3.283 $\int x^2 \operatorname{arctanh}(\cosh(x)) dx$

Optimal result	1950
Mathematica [A] (verified)	1950
Rubi [C] (verified)	1951
Maple [C] (warning: unable to verify)	1954
Fricas [A] (verification not implemented)	1955
Sympy [F]	1955
Maxima [A] (verification not implemented)	1956
Giac [F]	1956
Mupad [F(-1)]	1956
Reduce [F]	1957

#### Optimal result

Integrand size = 7, antiderivative size = 77

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = -\frac{2}{3}x^3 \operatorname{arctanh}(e^x) + \frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) - x^2 \operatorname{PolyLog}(2, -e^x) + x^2 \operatorname{PolyLog}(2, e^x) + 2x \operatorname{PolyLog}(3, -e^x) - 2x \operatorname{PolyLog}(3, e^x) - 2 \operatorname{PolyLog}(4, -e^x) + 2 \operatorname{PolyLog}(4, e^x)$$

output

```
-2/3*x^3*arctanh(exp(x))+1/3*x^3*arctanh(cosh(x))-x^2*polylog(2,-exp(x))+x^2*polylog(2,exp(x))+2*x*polylog(3,-exp(x))-2*x*polylog(3,exp(x))-2*polylog(4,-exp(x))+2*polylog(4,exp(x))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{3}(x^3 \operatorname{arctanh}(\cosh(x)) + x^3 \log(1 - e^x) - x^3 \log(1 + e^x) - 3x^2 \operatorname{PolyLog}(2, -e^x) + 3x^2 \operatorname{PolyLog}(2, e^x) + 6x \operatorname{PolyLog}(3, -e^x) - 6x \operatorname{PolyLog}(3, e^x) - 6 \operatorname{PolyLog}(4, -e^x) + 6 \operatorname{PolyLog}(4, e^x))$$

input `Integrate[x^2*ArcTanh[Cosh[x]],x]`

output `(x^3*ArcTanh[Cosh[x]] + x^3*Log[1 - E^x] - x^3*Log[1 + E^x] - 3*x^2*PolyLog[2, -E^x] + 3*x^2*PolyLog[2, E^x] + 6*x*PolyLog[3, -E^x] - 6*x*PolyLog[3, E^x] - 6*PolyLog[4, -E^x] + 6*PolyLog[4, E^x])/3`

## Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.29, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.286$ , Rules used = {6827, 25, 3042, 26, 4670, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}(\cosh(x)) dx \\
 & \quad \downarrow \text{6827} \\
 & \frac{1}{3} x^3 \operatorname{arctanh}(\cosh(x)) - \frac{1}{3} \int -x^3 \operatorname{csch}(x) dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \int x^3 \operatorname{csch}(x) dx + \frac{1}{3} x^3 \operatorname{arctanh}(\cosh(x)) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3} \int i x^3 \csc(ix) dx \\
 & \quad \downarrow \text{26} \\
 & \frac{1}{3} x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3} i \int x^3 \csc(ix) dx \\
 & \quad \downarrow \text{4670}
 \end{aligned}$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3}i \left( 3i \int x^2 \log(1 - e^x) dx - 3i \int x^2 \log(1 + e^x) dx + 2ix^3 \operatorname{arctanh}(e^x) \right)$$

↓ 3011

$$\frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3}i \left( -3i \left( 2 \int x \operatorname{PolyLog}(2, -e^x) dx - x^2 \operatorname{PolyLog}(2, -e^x) \right) + 3i \left( 2 \int x \operatorname{PolyLog}(2, e^x) dx - x^2 \operatorname{PolyLog}(2, e^x) \right) \right)$$

↓ 7163

$$\frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3}i \left( -3i \left( 2 \left( x \operatorname{PolyLog}(3, -e^x) - \int \operatorname{PolyLog}(3, -e^x) dx \right) - x^2 \operatorname{PolyLog}(2, -e^x) \right) + 3i \left( 2 \left( x \operatorname{PolyLog}(3, e^x) - \int \operatorname{PolyLog}(3, e^x) dx \right) - x^2 \operatorname{PolyLog}(2, e^x) \right) \right)$$

↓ 2720

$$\frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3}i \left( -3i \left( 2 \left( x \operatorname{PolyLog}(3, -e^x) - \int e^{-x} \operatorname{PolyLog}(3, -e^x) dx \right) - x^2 \operatorname{PolyLog}(2, -e^x) \right) + 3i \left( 2 \left( x \operatorname{PolyLog}(3, e^x) - \int e^x \operatorname{PolyLog}(3, e^x) dx \right) - x^2 \operatorname{PolyLog}(2, e^x) \right) \right)$$

↓ 7143

$$\frac{1}{3}x^3 \operatorname{arctanh}(\cosh(x)) + \frac{1}{3}i \left( 2ix^3 \operatorname{arctanh}(e^x) - 3i \left( 2 \left( x \operatorname{PolyLog}(3, -e^x) - \operatorname{PolyLog}(4, -e^x) \right) - x^2 \operatorname{PolyLog}(2, -e^x) \right) + 3i \left( 2 \left( x \operatorname{PolyLog}(3, e^x) - \operatorname{PolyLog}(4, e^x) \right) - x^2 \operatorname{PolyLog}(2, e^x) \right) \right)$$

input `Int [x^2*ArcTanh[Cosh[x]], x]`

output `(x^3*ArcTanh[Cosh[x]])/3 + (I/3)*((2*I)*x^3*ArcTanh[E^x] - (3*I)*(-(x^2*PolyLog[2, -E^x]) + 2*(x*PolyLog[3, -E^x] - PolyLog[4, -E^x])) + (3*I)*(-(x^2*PolyLog[2, E^x]) + 2*(x*PolyLog[3, E^x] - PolyLog[4, E^x])))`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))]*(f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6827 `Int[((a_.) + ArcTanh[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcTanh[u])/(d*(m + 1))), x] - Simp[b/(d*(m + 1)) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(1 - u^2)), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && FalseQ[PowerVariableExpn[u, m + 1, x]]`

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol]
:> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x]
&& GtQ[m, 0]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.42 (sec) , antiderivative size = 422, normalized size of antiderivative = 5.48

method	result
risch	$-\frac{x^3 \ln(e^x - 1)}{3} + \frac{i\pi \left( 2 \operatorname{csgn}\left(ie^{-x}(e^x - 1)^2\right)^2 - \operatorname{csgn}(i(1+e^x))^2 \operatorname{csgn}(i(1+e^x)^2) + 2 \operatorname{csgn}(i(1+e^x)) \operatorname{csgn}(i(1+e^x)^2) - \operatorname{csgn}(i(1+e^x)^2) \right)}{3}$

input

```
int(x^2*arctanh(cosh(x)),x,method=_RETURNVERBOSE)
```

output

```
-1/3*x^3*ln(exp(x)-1)+1/12*I*Pi*(2*csgn(I*exp(-x)*(exp(x)-1)^2)-csgn(I*(1+exp(x)))^2*csgn(I*(1+exp(x))^2)+2*csgn(I*(1+exp(x)))*csgn(I*(1+exp(x))^2)-csgn(I*(1+exp(x))^2)^3-csgn(I*(1+exp(x))^2)*csgn(I*exp(-x))*csgn(I*exp(-x)*(1+exp(x))^2)+csgn(I*(1+exp(x))^2)*csgn(I*exp(-x)*(1+exp(x))^2)^2+csgn(I*(exp(x)-1))^2*csgn(I*(exp(x)-1)^2)-2*csgn(I*(exp(x)-1))*csgn(I*(exp(x)-1)^2)^2+csgn(I*(exp(x)-1)^2)^3+csgn(I*(exp(x)-1)^2)*csgn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)-1)^2)-csgn(I*(exp(x)-1)^2)*csgn(I*exp(-x)*(exp(x)-1)^2)^2+csgn(I*exp(-x))*csgn(I*exp(-x)*(1+exp(x))^2)^2-csgn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)-1)^2)^2-csgn(I*exp(-x)*(1+exp(x))^2)^3-csgn(I*exp(-x)*(exp(x)-1)^2)^3-2)*x^3-x^2*polylog(2,-exp(x))+2*x*polylog(3,-exp(x))-2*polylog(4,-exp(x))+1/3*x^3*ln(1-exp(x))+x^2*polylog(2,exp(x))-2*x*polylog(3,exp(x))+2*polylog(4,exp(x))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.53

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{6} x^3 \log\left(-\frac{\cosh(x)+1}{\cosh(x)-1}\right) - \frac{1}{3} x^3 \log(\cosh(x)+\sinh(x)+1) \\ + \frac{1}{3} x^3 \log(-\cosh(x)-\sinh(x)+1) \\ + x^2 \operatorname{Li}_2(\cosh(x)+\sinh(x)) - x^2 \operatorname{Li}_2(-\cosh(x)-\sinh(x)) \\ - 2x \operatorname{polylog}(3, \cosh(x)+\sinh(x)) \\ + 2x \operatorname{polylog}(3, -\cosh(x)-\sinh(x)) \\ + 2 \operatorname{polylog}(4, \cosh(x)+\sinh(x)) \\ - 2 \operatorname{polylog}(4, -\cosh(x)-\sinh(x))$$

input `integrate(x^2*arctanh(cosh(x)),x, algorithm="fricas")`

output `1/6*x^3*log(-(cosh(x) + 1)/(cosh(x) - 1)) - 1/3*x^3*log(cosh(x) + sinh(x) + 1) + 1/3*x^3*log(-cosh(x) - sinh(x) + 1) + x^2*dilog(cosh(x) + sinh(x)) - x^2*dilog(-cosh(x) - sinh(x)) - 2*x*polylog(3, cosh(x) + sinh(x)) + 2*x*polylog(3, -cosh(x) - sinh(x)) + 2*polylog(4, cosh(x) + sinh(x)) - 2*polylog(4, -cosh(x) - sinh(x))`

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \int x^2 \operatorname{atanh}(\cosh(x)) dx$$

input `integrate(x**2*atanh(cosh(x)),x)`

output `Integral(x**2*atanh(cosh(x)), x)`



**Maxima [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.01

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \frac{1}{3} x^3 \operatorname{artanh}(\cosh(x)) - \frac{1}{3} x^3 \log(e^x + 1) + \frac{1}{3} x^3 \log(-e^x + 1) - x^2 \operatorname{Li}_2(-e^x) + x^2 \operatorname{Li}_2(e^x) + 2x \operatorname{Li}_3(-e^x) - 2x \operatorname{Li}_3(e^x) - 2 \operatorname{Li}_4(-e^x) + 2 \operatorname{Li}_4(e^x)$$

input `integrate(x^2*arctanh(cosh(x)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(cosh(x)) - 1/3*x^3*log(e^x + 1) + 1/3*x^3*log(-e^x + 1) - x^2*dilog(-e^x) + x^2*dilog(e^x) + 2*x*polylog(3, -e^x) - 2*x*polylog(3, e^x) - 2*polylog(4, -e^x) + 2*polylog(4, e^x)`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \int x^2 \operatorname{artanh}(\cosh(x)) dx$$

input `integrate(x^2*arctanh(cosh(x)),x, algorithm="giac")`

output `integrate(x^2*arctanh(cosh(x)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \int x^2 \operatorname{atanh}(\cosh(x)) dx$$

input `int(x^2*atanh(cosh(x)),x)`

output `int(x^2*atanh(cosh(x)), x)`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(\cosh(x)) dx = \int \operatorname{atanh}(\cosh(x)) x^2 dx$$

input `int(x^2*atanh(cosh(x)),x)`

output `int(atanh(cosh(x))*x**2,x)`

### 3.284 $\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx$

Optimal result	1958
Mathematica [A] (verified)	1959
Rubi [A] (verified)	1959
Maple [C] (warning: unable to verify)	1964
Fricas [B] (verification not implemented)	1964
Sympy [F]	1965
Maxima [A] (verification not implemented)	1966
Giac [F]	1966
Mupad [F(-1)]	1967
Reduce [F]	1967

#### Optimal result

Integrand size = 15, antiderivative size = 307

$$\begin{aligned}
 \int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx &= \frac{1}{3} x^3 \operatorname{arctanh}(c + d \tanh(a + bx)) \\
 &+ \frac{1}{6} x^3 \log \left( 1 + \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right) \\
 &- \frac{1}{6} x^3 \log \left( 1 + \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right) \\
 &+ \frac{x^2 \operatorname{PolyLog} \left( 2, -\frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{4b} \\
 &- \frac{x^2 \operatorname{PolyLog} \left( 2, -\frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{4b} \\
 &- \frac{x \operatorname{PolyLog} \left( 3, -\frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{4b^2} \\
 &+ \frac{x \operatorname{PolyLog} \left( 3, -\frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{4b^2} \\
 &+ \frac{\operatorname{PolyLog} \left( 4, -\frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{8b^3} \\
 &- \frac{\operatorname{PolyLog} \left( 4, -\frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{8b^3}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{3}x^3 \operatorname{arctanh}(c+d \tanh(bx+a)) + \frac{1}{6}x^3 \ln(1+(1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & - \frac{1}{6}x^3 \ln(1+(1+c+d)\exp(2bx+2a)/(1+c-d)) + \frac{1}{4}x^2 \operatorname{polylog}(2, -(1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & / b - \frac{1}{4}x^2 \operatorname{polylog}(2, -(1+c+d)\exp(2bx+2a)/(1+c-d)) / b - \frac{1}{4}x \operatorname{polylog}(3, -(1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & / b^2 + \frac{1}{4}x \operatorname{polylog}(3, -(1+c+d)\exp(2bx+2a)/(1+c-d)) / b^2 + \frac{1}{8} \operatorname{polylog}(4, -(1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & / b^3 - \frac{1}{8} \operatorname{polylog}(4, -(1+c+d)\exp(2bx+2a)/(1+c-d)) / b^3 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.86

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \frac{1}{3}x^3 \operatorname{arctanh}(c + d \tanh(a + bx)) + \frac{4b^3 x^3 \log\left(1 + \frac{(-1+c-d)e^{-2(a+bx)}}{-1+c+d}\right) - 4b^3 x^3 \log\left(1 + \frac{(1+c-d)e^{-2(a+bx)}}{1+c+d}\right) - 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{(1-c+d)e^{-2(a+bx)}}{-1+c+d}\right)}{24b^3}$$

input

Integrate[x^2\*ArcTanh[c + d\*Tanh[a + b\*x]], x]

output

$$\begin{aligned} & \frac{(x^3 \operatorname{ArcTanh}[c + d \operatorname{Tanh}[a + b x]])}{3} + \frac{(4 b^3 x^3 \operatorname{Log}[1 + (-1 + c - d)/((-1 + c + d) E^{(2(a + b x))} - 4 b^3 x^3 \operatorname{Log}[1 + (1 + c - d)/((1 + c + d) E^{(2(a + b x))}] - 6 b^2 x^2 \operatorname{PolyLog}[2, (1 - c + d)/((-1 + c + d) E^{(2(a + b x))}] + 6 b^2 x^2 \operatorname{PolyLog}[2, (-1 - c + d)/((1 + c + d) E^{(2(a + b x))}]] - 6 b x \operatorname{PolyLog}[3, (1 - c + d)/((-1 + c + d) E^{(2(a + b x))}] + 6 b x \operatorname{PolyLog}[3, (-1 - c + d)/((1 + c + d) E^{(2(a + b x))}] - 3 \operatorname{PolyLog}[4, (1 - c + d)/((-1 + c + d) E^{(2(a + b x))}] + 3 \operatorname{PolyLog}[4, (-1 - c + d)/((1 + c + d) E^{(2(a + b x))}]])/(24 b^3)}{24 b^3} \end{aligned}$$
**Rubi [A] (verified)**Time = 1.42 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6797, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d \tanh(a + bx) + c) dx$$

↓ 6797

$$\frac{1}{3}b(-c - d + 1) \int \frac{e^{2a+2bx} x^3}{-c + (-c - d + 1)e^{2a+2bx} + d + 1} dx - \frac{1}{3}b(c + d + 1) \int \frac{e^{2a+2bx} x^3}{c + (c + d + 1)e^{2a+2bx} - d + 1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + c)$$

↓ 2620

$$\frac{1}{3}b(-c - d + 1) \left( \frac{x^3 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c - d + 1)} - \frac{3 \int x^2 \log\left(\frac{e^{2a+2bx}(-c-d+1)}{-c+d+1} + 1\right) dx}{2b(-c - d + 1)} \right) - \frac{1}{3}b(c + d + 1) \left( \frac{x^3 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c + d + 1)} - \frac{3 \int x^2 \log\left(\frac{e^{2a+2bx}(c+d+1)}{c-d+1} + 1\right) dx}{2b(c + d + 1)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + c)$$

↓ 3011

$$\frac{1}{3}b(-c - d + 1) \left( \frac{x^3 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c - d + 1)} - \frac{3 \left( \frac{\int x \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right)}{2b(-c - d + 1)} \right) - \frac{1}{3}b(c + d + 1) \left( \frac{x^3 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c + d + 1)} - \frac{3 \left( \frac{\int x \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)}{2b(c + d + 1)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + c)$$

↓ 7163

$$1) \left( \frac{x^3 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\frac{\frac{1}{3}b(-c-d+1)}{3} \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\int \operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right)}{2b(-c-d+1)} \right)$$

$$1) \left( \frac{x^3 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\frac{\frac{1}{3}b(c+d+1)}{3} \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\int \operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)}{2b(c+d+1)} \right)$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + c)$$

↓ 2720

$$1) \left( \frac{x^3 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\frac{\frac{1}{3}b(-c-d+1)}{3} \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{4b^2} \right)}{2b(-c-d+1)} \right)$$

$$1) \left( \frac{x^3 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\frac{\frac{1}{3}b(c+d+1)}{3} \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{4b^2} \right)}{2b(c+d+1)} \right)$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + c)$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + c) + \frac{1}{3}b(-c - d + \\
 1) & \left( \frac{x^3 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{3 \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{4b^2} \right)}{2b(-c-d+1)} \right) \\
 1) & \left( \frac{x^3 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{3 \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{4b^2} \right)}{2b(c+d+1)} \right)
 \end{aligned}$$

input `Int[x^2*ArcTanh[c + d*Tanh[a + b*x]],x]`

output `(x^3*ArcTanh[c + d*Tanh[a + b*x]])/3 + (b*(1 - c - d)*((x^3*Log[1 + ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)])/(2*b*(1 - c - d)) - (3*(-1/2*(x^2*PolyLog[2, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)))]/b + ((x*PolyLog[3, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)))]/(2*b) - PolyLog[4, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d))]/(4*b^2))/b))/(2*b*(1 - c - d)))/3 - (b*(1 + c + d)*((x^3*Log[1 + ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)])/(2*b*(1 + c + d)) - (3*(-1/2*(x^2*PolyLog[2, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)))]/b + ((x*PolyLog[3, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)))]/(2*b) - PolyLog[4, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d))]/(4*b^2))/b))/(2*b*(1 + c + d)))/3`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6797

```
Int[ArcTanh[(c_) + (d_)*Tanh[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m
_)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(
m + 1))), x] + (Simp[b*((1 - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E
^(2*a + 2*b*x))/(1 - c + d + (1 - c - d)*E^(2*a + 2*b*x))], x], x] - Simp[b*
((1 + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(1 + c -
d + (1 + c + d)*E^(2*a + 2*b*x))], x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& IGtQ[m, 0] && NeQ[(c - d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```



rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^(m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 13.13 (sec) , antiderivative size = 5316, normalized size of antiderivative = 17.32

method	result	size
risch	Expression too large to display	5316

input

```
int(x^2*arctanh(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 900 vs.  $2(263) = 526$ .

Time = 0.12 (sec) , antiderivative size = 900, normalized size of antiderivative = 2.93

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctanh(c+d*tanh(b*x+a)),x, algorithm="fricas")
```

output

```

1/6*(b^3*x^3*log(-((c + 1)*cosh(b*x + a) + d*sinh(b*x + a))/((c - 1)*cosh(
b*x + a) + d*sinh(b*x + a))) - 3*b^2*x^2*dilog(sqrt(-(c + d + 1)/(c - d +
1))*(cosh(b*x + a) + sinh(b*x + a))) - 3*b^2*x^2*dilog(-sqrt(-(c + d + 1)/
(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) + 3*b^2*x^2*dilog(sqrt(-(c +
d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + 3*b^2*x^2*dilog(-s
qrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + a^3*log(2
*(c + d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(b*x + a) + 2*(c - d + 1)*s
qrt(-(c + d + 1)/(c - d + 1))) + a^3*log(2*(c + d + 1)*cosh(b*x + a) + 2*(
c + d + 1)*sinh(b*x + a) - 2*(c - d + 1)*sqrt(-(c + d + 1)/(c - d + 1))) -
a^3*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) + 2*(c
- d - 1)*sqrt(-(c + d - 1)/(c - d - 1))) - a^3*log(2*(c + d - 1)*cosh(b*x
+ a) + 2*(c + d - 1)*sinh(b*x + a) - 2*(c - d - 1)*sqrt(-(c + d - 1)/(c -
d - 1))) + 6*b*x*polylog(3, sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a)
+ sinh(b*x + a))) + 6*b*x*polylog(3, -sqrt(-(c + d + 1)/(c - d + 1))*(cosh
(b*x + a) + sinh(b*x + a))) - 6*b*x*polylog(3, sqrt(-(c + d - 1)/(c - d -
1))*(cosh(b*x + a) + sinh(b*x + a))) - 6*b*x*polylog(3, -sqrt(-(c + d - 1)
/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) - (b^3*x^3 + a^3)*log(sqrt(
-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) - (b^3*x^3
+ a^3)*log(-sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))
+ 1) + (b^3*x^3 + a^3)*log(sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + ...

```

## Sympy [F]

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input

```
integrate(x**2*atanh(c+d*tanh(b*x+a)), x)
```

output

```
Integral(x**2*atanh(c + d*tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.92

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \frac{1}{3} x^3 \operatorname{artanh}(d \tanh(bx + a) + c) - \frac{1}{18} bd \left( \frac{4 b^3 x^3 \log\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + 6 b^2 x^2 \operatorname{Li}_2\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) - 6 bx \operatorname{Li}_3\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) + 3}{b^4 d} \right)$$

input `integrate(x^2*arctanh(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(d*tanh(b*x + a) + c) - 1/18*b*d*((4*b^3*x^3*log((c + d + 1)*e^(2*b*x + 2*a))/(c - d + 1) + 1) + 6*b^2*x^2*dilog(-(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) - 6*b*x*polylog(3, -(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) + 3*polylog(4, -(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^4*d) - (4*b^3*x^3*log((c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1) + 1) + 6*b^2*x^2*dilog(-(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) - 6*b*x*polylog(3, -(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) + 3*polylog(4, -(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)))/(b^4*d)`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x^2 \operatorname{artanh}(d \tanh(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(d*tanh(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input `int(x^2*atanh(c + d*tanh(a + b*x)),x)`output `int(x^2*atanh(c + d*tanh(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a)d + c) x^2 dx$$

input `int(x^2*atanh(c+d*tanh(b*x+a)),x)`output `int(atanh(tanh(a + b*x)*d + c)*x**2,x)`

### 3.285 $\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx$

Optimal result	1968
Mathematica [A] (verified)	1969
Rubi [A] (verified)	1969
Maple [C] (warning: unable to verify)	1973
Fricas [B] (verification not implemented)	1973
Sympy [F]	1974
Maxima [A] (verification not implemented)	1975
Giac [F]	1975
Mupad [F(-1)]	1976
Reduce [F]	1976

#### Optimal result

Integrand size = 13, antiderivative size = 231

$$\begin{aligned}
 \int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx &= \frac{1}{2} x^2 \operatorname{arctanh}(c + d \tanh(a + bx)) \\
 &+ \frac{1}{4} x^2 \log \left( 1 + \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right) \\
 &- \frac{1}{4} x^2 \log \left( 1 + \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right) \\
 &+ \frac{x \operatorname{PolyLog} \left( 2, -\frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{4b} \\
 &- \frac{x \operatorname{PolyLog} \left( 2, -\frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{4b} \\
 &- \frac{\operatorname{PolyLog} \left( 3, -\frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{8b^2} \\
 &+ \frac{\operatorname{PolyLog} \left( 3, -\frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\operatorname{arctanh}(c+d*\operatorname{tanh}(b*x+a))+1/4*x^2*\ln(1+(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))-1/4*x^2*\ln(1+(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))+1/4*x*\operatorname{polylog}(2,-(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))/b-1/4*x*\operatorname{polylog}(2,-(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))/b-1/8*\operatorname{polylog}(3,-(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))/b^2+1/8*\operatorname{polylog}(3,-(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))/b^2 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.86

$$\int x \operatorname{arctanh}(c + d \operatorname{tanh}(a + bx)) dx$$

$$= \frac{4b^2x^2 \operatorname{arctanh}(c + d \operatorname{tanh}(a + bx)) + 2b^2x^2 \log\left(1 + \frac{(-1+c-d)e^{-2(a+bx)}}{-1+c+d}\right) - 2b^2x^2 \log\left(1 + \frac{(1+c-d)e^{-2(a+bx)}}{1+c+d}\right)}{b^2}$$

input

Integrate[x\*ArcTanh[c + d\*Tanh[a + b\*x]],x]

output

$$\begin{aligned} & (4*b^2*x^2*ArcTanh[c + d*Tanh[a + b*x]] + 2*b^2*x^2*Log[1 + (-1 + c - d)/((-1 + c + d)*E^(2*(a + b*x)))] - 2*b^2*x^2*Log[1 + (1 + c - d)/((1 + c + d)*E^(2*(a + b*x)))] - 2*b*x*PolyLog[2, (1 - c + d)/((-1 + c + d)*E^(2*(a + b*x)))] + 2*b*x*PolyLog[2, (-1 - c + d)/((1 + c + d)*E^(2*(a + b*x)))] - PolyLog[3, (1 - c + d)/((-1 + c + d)*E^(2*(a + b*x)))] + PolyLog[3, (-1 - c + d)/((1 + c + d)*E^(2*(a + b*x)))])/(8*b^2) \end{aligned}$$

**Rubi [A] (verified)**Time = 1.18 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.32, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {6797, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \operatorname{tanh}(a + bx) + c) dx$$

$$\begin{aligned}
& \downarrow \text{6797} \\
& \frac{1}{2}b(-c-d+1) \int \frac{e^{2a+2bx}x^2}{-c+(-c-d+1)e^{2a+2bx}+d+1} dx - \frac{1}{2}b(c+d+1) \\
& 1) \int \frac{e^{2a+2bx}x^2}{c+(c+d+1)e^{2a+2bx}-d+1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx)+c) \\
& \downarrow \text{2620} \\
& \frac{1}{2}b(-c-d+1) \left( \frac{x^2 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\int x \log\left(\frac{e^{2a+2bx}(-c-d+1)}{-c+d+1} + 1\right) dx}{b(-c-d+1)} \right) - \frac{1}{2}b(c+d+1) \\
& \left( \frac{x^2 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\int x \log\left(\frac{e^{2a+2bx}(c+d+1)}{c-d+1} + 1\right) dx}{b(c+d+1)} \right) + \\
& \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx)+c) \\
& \downarrow \text{3011} \\
& \frac{1}{2}b(-c-d+1) \\
& 1) \left( \frac{x^2 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\int \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) dx}{2b} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right) - \\
& \frac{1}{2}b(c+d+1) \\
& 1) \left( \frac{x^2 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\int \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) dx}{2b} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right) + \\
& \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx)+c) \\
& \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
 & 1) \left( \frac{x^2 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right) \\
 & 1) \left( \frac{x^2 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right) + \\
 & \frac{1}{2} x^2 \operatorname{arctanh}(d \tanh(a + bx) + c)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{2} x^2 \operatorname{arctanh}(d \tanh(a + bx) + c) + \frac{1}{2} b(-c-d+1) \\
 & 1) \left( \frac{x^2 \log\left(\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1\right)}{2b(-c-d+1)} - \frac{\operatorname{PolyLog}\left(3, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right) - \\
 & 1) \left( \frac{x^2 \log\left(\frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1\right)}{2b(c+d+1)} - \frac{\operatorname{PolyLog}\left(3, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)
 \end{aligned}$$

input `Int[x*ArcTanh[c + d*Tanh[a + b*x]],x]`

output `(x^2*ArcTanh[c + d*Tanh[a + b*x]])/2 + (b*(1 - c - d)*((x^2*Log[1 + ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)])/(2*b*(1 - c - d)) - (-1/2*(x*PolyLog[2, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d))])/b + PolyLog[3, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d))]/(4*b^2)))/(b*(1 - c - d)))/2 - (b*(1 + c + d)*((x^2*Log[1 + ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)])/(2*b*(1 + c + d)) - (-1/2*(x*PolyLog[2, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d))])/b + PolyLog[3, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d))]/(4*b^2)))/(b*(1 + c + d)))/2`



## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6797

```
Int[ArcTanh[(c_) + (d_)*Tanh[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_))^(m
_), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(
m + 1))), x] + (Simp[b*((1 - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E
^(2*a + 2*b*x))/(1 - c + d + (1 - c - d)*E^(2*a + 2*b*x))], x], x] - Simp[b*
((1 + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(1 + c -
d + (1 + c + d)*E^(2*a + 2*b*x))], x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& IGtQ[m, 0] && NeQ[(c - d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.92 (sec) , antiderivative size = 5012, normalized size of antiderivative = 21.70

method	result	size
risch	Expression too large to display	5012

input `int(x*arctanh(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 746 vs.  $2(197) = 394$ .

Time = 0.13 (sec) , antiderivative size = 746, normalized size of antiderivative = 3.23

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctanh(c+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```

1/4*(b^2*x^2*log(-((c + 1)*cosh(b*x + a) + d*sinh(b*x + a))/((c - 1)*cosh(
b*x + a) + d*sinh(b*x + a))) - 2*b*x*dilog(sqrt(-(c + d + 1)/(c - d + 1))*
(cosh(b*x + a) + sinh(b*x + a))) - 2*b*x*dilog(-sqrt(-(c + d + 1)/(c - d +
1))*(cosh(b*x + a) + sinh(b*x + a))) + 2*b*x*dilog(sqrt(-(c + d - 1)/(c -
d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + 2*b*x*dilog(-sqrt(-(c + d - 1)
/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) - a^2*log(2*(c + d + 1)*cos
h(b*x + a) + 2*(c + d + 1)*sinh(b*x + a) + 2*(c - d + 1)*sqrt(-(c + d + 1)
/(c - d + 1))) - a^2*log(2*(c + d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(
b*x + a) - 2*(c - d + 1)*sqrt(-(c + d + 1)/(c - d + 1))) + a^2*log(2*(c +
d - 1)*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) + 2*(c - d - 1)*sqrt(-
(c + d - 1)/(c - d - 1))) + a^2*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d
- 1)*sinh(b*x + a) - 2*(c - d - 1)*sqrt(-(c + d - 1)/(c - d - 1))) - (b^2*
x^2 - a^2)*log(sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x +
a)) + 1) - (b^2*x^2 - a^2)*log(-sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x +
a) + sinh(b*x + a)) + 1) + (b^2*x^2 - a^2)*log(sqrt(-(c + d - 1)/(c - d -
1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b^2*x^2 - a^2)*log(-sqrt(-(c
+ d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 2*polylog(3,
sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) + 2*polylo
g(3, -sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 2*
polylog(3, sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a...

```

### Sympy [F]

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input

```
integrate(x*atanh(c+d*tanh(b*x+a)),x)
```

output

```
Integral(x*atanh(c + d*tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.93

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx =$$

$$-\frac{1}{8} bd \left( \frac{2b^2x^2 \log\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + 2bx \operatorname{Li}_2\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) - \operatorname{Li}_3\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right)}{b^3d} - \frac{2b^2x^2}{b^3d} \right)$$

$$+ \frac{1}{2} x^2 \operatorname{artanh}(d \tanh(bx + a) + c)$$

input `integrate(x*arctanh(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `-1/8*b*d*((2*b^2*x^2*log((c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1) + 1) + 2*b*x*dilog(-(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) - polylog(3, -(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^3*d) - (2*b^2*x^2*log((c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1) + 1) + 2*b*x*dilog(-(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) - polylog(3, -(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)))/(b^3*d)) + 1/2*x^2*arctanh(d*tanh(b*x + a) + c)`

**Giac [F]**

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x \operatorname{artanh}(d \tanh(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*tanh(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int x \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input `int(x*atanh(c + d*tanh(a + b*x)),x)`output `int(x*atanh(c + d*tanh(a + b*x)), x)`**Reduce [F]**

$$\int x \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a)d + c) x dx$$

input `int(x*atanh(c+d*tanh(b*x+a)),x)`output `int(atanh(tanh(a + b*x)*d + c)*x,x)`

### 3.286 $\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx$

Optimal result	1977
Mathematica [A] (verified)	1978
Rubi [A] (verified)	1978
Maple [B] (verified)	1980
Fricas [B] (verification not implemented)	1981
Sympy [F]	1982
Maxima [A] (verification not implemented)	1982
Giac [F]	1983
Mupad [F(-1)]	1983
Reduce [F]	1984

#### Optimal result

Integrand size = 11, antiderivative size = 150

$$\begin{aligned}
 \int \operatorname{arctanh}(c + d \tanh(a + bx)) dx &= x \operatorname{arctanh}(c + d \tanh(a + bx)) \\
 &+ \frac{1}{2} x \log \left( 1 + \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right) \\
 &- \frac{1}{2} x \log \left( 1 + \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right) \\
 &+ \frac{\operatorname{PolyLog} \left( 2, -\frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right)}{4b} \\
 &- \frac{\operatorname{PolyLog} \left( 2, -\frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right)}{4b}
 \end{aligned}$$

output

```

x*arctanh(c+d*tanh(b*x+a))+1/2*x*ln(1+(1-c-d)*exp(2*b*x+2*a)/(1-c+d))-1/2*
x*ln(1+(1+c+d)*exp(2*b*x+2*a)/(1+c-d))+1/4*polylog(2,-(1-c-d)*exp(2*b*x+2*
a)/(1-c+d))/b-1/4*polylog(2,-(1+c+d)*exp(2*b*x+2*a)/(1+c-d))/b

```

**Mathematica [A] (verified)**

Time = 3.34 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.87

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = x \operatorname{arctanh}(c + d \tanh(a + bx)) + \frac{2bx \left( \log \left( 1 + \frac{(-1+c+d)e^{2(a+bx)}}{-1+c-d} \right) - \log \left( 1 + \frac{(1+c+d)e^{2(a+bx)}}{1+c-d} \right) \right) + \operatorname{PolyLog} \left( 2, -\frac{(-1+c+d)e^{2(a+bx)}}{-1+c-d} \right) - \operatorname{PolyLog} \left( 2, \frac{(1+c+d)e^{2(a+bx)}}{1+c-d} \right)}{4b}$$

input `Integrate[ArcTanh[c + d*Tanh[a + b*x]], x]`

output

```
x*ArcTanh[c + d*Tanh[a + b*x]] + (2*b*x*(Log[1 + ((-1 + c + d)*E^(2*(a + b*x)))/(-1 + c - d)] - Log[1 + ((1 + c + d)*E^(2*(a + b*x)))/(1 + c - d)] + PolyLog[2, -(((1 + c + d)*E^(2*(a + b*x)))/(-1 + c - d))] - PolyLog[2, -(((1 + c + d)*E^(2*(a + b*x)))/(1 + c - d))])/(4*b)
```

**Rubi [A] (verified)**Time = 1.06 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.38, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6789, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \tanh(a + bx) + c) dx$$

$$\downarrow \text{6789}$$

$$b(-c - d + 1) \int \frac{e^{2a+2bx}}{-c + (-c - d + 1)e^{2a+2bx} + d + 1} dx - b(c + d + 1) \int \frac{e^{2a+2bx}}{c + (c + d + 1)e^{2a+2bx} - d + 1} dx + x \operatorname{arctanh}(d \tanh(a + bx) + c)$$

$$\downarrow \text{2620}$$

$$b(-c-d+1) \left( \frac{x \log \left( \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1 \right)}{2b(-c-d+1)} - \frac{\int \log \left( \frac{e^{2a+2bx}(-c-d+1)}{-c+d+1} + 1 \right) dx}{2b(-c-d+1)} \right) - b(c+d+1) \left( \frac{x \log \left( \frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1 \right)}{2b(c+d+1)} - \frac{\int \log \left( \frac{e^{2a+2bx}(c+d+1)}{c-d+1} + 1 \right) dx}{2b(c+d+1)} \right) + x \operatorname{arctanh}(d \tanh(a+bx) + c)$$

↓ 2715

$$b(-c-d+1) \left( \frac{x \log \left( \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1 \right)}{2b(-c-d+1)} - \frac{\int e^{-2a-2bx} \log \left( \frac{e^{2a+2bx}(-c-d+1)}{-c+d+1} + 1 \right) de^{2a+2bx}}{4b^2(-c-d+1)} \right) - b(c+d+1) \left( \frac{x \log \left( \frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1 \right)}{2b(c+d+1)} - \frac{\int e^{-2a-2bx} \log \left( \frac{e^{2a+2bx}(c+d+1)}{c-d+1} + 1 \right) de^{2a+2bx}}{4b^2(c+d+1)} \right) + x \operatorname{arctanh}(d \tanh(a+bx) + c)$$

↓ 2838

$$1) \left( \frac{\operatorname{PolyLog} \left( 2, -\frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{4b^2(-c-d+1)} + \frac{x \log \left( \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} + 1 \right)}{2b(-c-d+1)} \right) - b(c+d+1) \left( \frac{\operatorname{PolyLog} \left( 2, -\frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{4b^2(c+d+1)} + \frac{x \log \left( \frac{(c+d+1)e^{2a+2bx}}{c-d+1} + 1 \right)}{2b(c+d+1)} \right)$$

input `Int[ArcTanh[c + d*Tanh[a + b*x]],x]`

output `x*ArcTanh[c + d*Tanh[a + b*x]] + b*(1 - c - d)*((x*Log[1 + ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)])/(2*b*(1 - c - d)) + PolyLog[2, -(((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d))]/(4*b^2*(1 - c - d))] - b*(1 + c + d)*((x*Log[1 + ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)])/(2*b*(1 + c + d)) + PolyLog[2, -(((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d))]/(4*b^2*(1 + c + d))]`



Defintions of rubi rules used

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 6789 Int[ArcTanh[(c_) + (d_)*Tanh[(a_) + (b_)*(x_)]], x_Symbol] := Simp[x*Ar
cTanh[c + d*Tanh[a + b*x]], x] + (Simp[b*(1 - c - d) Int[x*(E^(2*a + 2*b*
x))/(1 - c + d + (1 - c - d)*E^(2*a + 2*b*x))), x], x] - Simp[b*(1 + c + d)
Int[x*(E^(2*a + 2*b*x))/(1 + c - d + (1 + c + d)*E^(2*a + 2*b*x))), x], x]
) /; FreeQ[{a, b, c, d}, x] && NeQ[(c - d)^2, 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 347 vs. 2(138) = 276.

Time = 3.83 (sec) , antiderivative size = 348, normalized size of antiderivative = 2.32

method	result
derivativedivides	$\frac{-\frac{\operatorname{arctanh}(c+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)+d)}{2}+\frac{\operatorname{arctanh}(c+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)-d)}{2}+d^2 \left(\frac{\operatorname{dilog}\left(\frac{-d \tanh (b x+a)}{1-c+d}\right)}{2}\right)}{2}$
default	$\frac{-\frac{\operatorname{arctanh}(c+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)+d)}{2}+\frac{\operatorname{arctanh}(c+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)-d)}{2}+d^2 \left(\frac{\operatorname{dilog}\left(\frac{-d \tanh (b x+a)}{1-c+d}\right)}{2}\right)}{2}$
risch	Expression too large to display

input `int(arctanh(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `1/b/d*(-1/2*arctanh(c+d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)+d)+1/2*arctanh(c+d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)-d)+1/2*d^2*(1/d*(1/2*dilog((-d*tanh(b*x+a)-c+1)/(1-c+d))+1/2*ln(-d*tanh(b*x+a)-d)*ln((-d*tanh(b*x+a)-c+1)/(1-c+d))-1/2*dilog((-d*tanh(b*x+a)-c-1)/(-1-c+d))-1/2*ln(-d*tanh(b*x+a)-d)*ln((-d*tanh(b*x+a)-c-1)/(-1-c+d)))-1/d*(-1/2*dilog((-d*tanh(b*x+a)-c-1)/(-1-c-d))-1/2*ln(-d*tanh(b*x+a)+d)*ln((-d*tanh(b*x+a)-c-1)/(-1-c-d))+1/2*dilog((-d*tanh(b*x+a)-c+1)/(1-c-d))+1/2*ln(-d*tanh(b*x+a)+d)*ln((-d*tanh(b*x+a)-c+1)/(1-c-d))))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 552 vs.  $2(128) = 256$ .

Time = 0.12 (sec) , antiderivative size = 552, normalized size of antiderivative = 3.68

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctanh(c+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```

1/2*(b*x*log(-(c + 1)*cosh(b*x + a) + d*sinh(b*x + a))/((c - 1)*cosh(b*x
+ a) + d*sinh(b*x + a))) + a*log(2*(c + d + 1)*cosh(b*x + a) + 2*(c + d +
1)*sinh(b*x + a) + 2*(c - d + 1)*sqrt(-(c + d + 1)/(c - d + 1))) + a*log(2
*(c + d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(b*x + a) - 2*(c - d + 1)*s
qrt(-(c + d + 1)/(c - d + 1))) - a*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c
+ d - 1)*sinh(b*x + a) + 2*(c - d - 1)*sqrt(-(c + d - 1)/(c - d - 1))) - a
*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) - 2*(c - d
- 1)*sqrt(-(c + d - 1)/(c - d - 1))) - (b*x + a)*log(sqrt(-(c + d + 1)/(c
- d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) - (b*x + a)*log(-sqrt(-(c +
d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b*x + a)*log(
sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b*x
+ a)*log(-sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))
+ 1) - dilog(sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)
)) - dilog(-sqrt(-(c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)
)) + dilog(sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a)))
+ dilog(-sqrt(-(c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))))/
b

```

**Sympy [F]**

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input

```
integrate(atanh(c+d*tanh(b*x+a)),x)
```

output

```
Integral(atanh(c + d*tanh(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.95

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx =$$

$$-\frac{1}{4}bd \left( \frac{2bx \log\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + \operatorname{Li}_2\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right)}{b^2d} - \frac{2bx \log\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1} + 1\right) + \operatorname{Li}_2\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1}\right)}{b^2d} \right)$$

$$+ x \operatorname{artanh}(d \tanh(bx + a) + c)$$

input `integrate(arctanh(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `-1/4*b*d*((2*b*x*log((c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1) + 1) + dilog(-  
(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^2*d) - (2*b*x*log((c + d - 1)  
) * e^(2*b*x + 2*a)/(c - d - 1) + 1) + dilog(-(c + d - 1)*e^(2*b*x + 2*a)/(c  
- d - 1)))/(b^2*d)) + x*arctanh(d*tanh(b*x + a) + c)`

### Giac [F]

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{artanh}(d \tanh(bx + a) + c) dx$$

input `integrate(arctanh(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*tanh(b*x + a) + c), x)`

### Mupad [F(-1)]

Timed out.

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{atanh}(c + d \tanh(a + bx)) dx$$

input `int(atanh(c + d*tanh(a + b*x)),x)`

output `int(atanh(c + d*tanh(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(c + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a) d + c) dx$$

input `int(atanh(c+d*tanh(b*x+a)),x)`

output `int(atanh(tanh(a + b*x)*d + c),x)`

### 3.287 $\int \frac{\operatorname{arctanh}(c+d \tanh(a+bx))}{x} dx$

Optimal result	1985
Mathematica [N/A]	1985
Rubi [N/A]	1986
Maple [N/A]	1986
Fricas [N/A]	1987
Sympy [N/A]	1987
Maxima [N/A]	1987
Giac [N/A]	1988
Mupad [N/A]	1988
Reduce [N/A]	1989

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(c+d*tanh(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 9.75 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx$$

input `Integrate[ArcTanh[c + d*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTanh[c + d*Tanh[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.47 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \tanh(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \tanh(a + bx) + c)}{x} dx$$

input `Int[ArcTanh[c + d*Tanh[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(c + d \tanh(bx + a))}{x} dx$$

input `int(arctanh(c+d*tanh(b*x+a))/x,x)`

output `int(arctanh(c+d*tanh(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*tanh(b*x + a) + c)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.95 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \tanh(a + bx))}{x} dx$$

input `integrate(atanh(c+d*tanh(b*x+a))/x,x)`

output `Integral(atanh(c + d*tanh(a + b*x))/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.75 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tanh(b*x+a))/x,x, algorithm="maxima")`



output `integrate(arctanh(d*tanh(b*x + a) + c)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.52 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tanh(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*tanh(b*x + a) + c)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.64 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \tanh(a + bx))}{x} dx$$

input `int(atanh(c + d*tanh(a + b*x))/x,x)`

output `int(atanh(c + d*tanh(a + b*x))/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a) d + c)}{x} dx$$

input `int(atanh(c+d*tanh(b*x+a))/x,x)`output `int(atanh(tanh(a + b*x)*d + c)/x,x)`

### 3.288 $\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$

Optimal result	1990
Mathematica [A] (verified)	1991
Rubi [A] (verified)	1991
Maple [C] (warning: unable to verify)	1995
Fricas [B] (verification not implemented)	1996
Sympy [F]	1996
Maxima [A] (verification not implemented)	1997
Giac [F]	1997
Mupad [F(-1)]	1997
Reduce [F]	1998

#### Optimal result

Integrand size = 16, antiderivative size = 155

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{bx^5}{20} + \frac{1}{4}x^4 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - \frac{1}{8}x^4 \log(1 + (1 + d)e^{2a+2bx}) - \frac{x^3 \operatorname{PolyLog}(2, -((1 + d)e^{2a+2bx}))}{4b} + \frac{3x^2 \operatorname{PolyLog}(3, -((1 + d)e^{2a+2bx}))}{8b^2} - \frac{3x \operatorname{PolyLog}(4, -((1 + d)e^{2a+2bx}))}{8b^3} + \frac{3 \operatorname{PolyLog}(5, -((1 + d)e^{2a+2bx}))}{16b^4}$$

output

```
1/20*b*x^5+1/4*x^4*arctanh(1+d*d*tanh(b*x+a))-1/8*x^4*ln(1+(1+d)*exp(2*b*x+2*a))-1/4*x^3*polylog(2,-(1+d)*exp(2*b*x+2*a))/b+3/8*x^2*polylog(3,-(1+d)*exp(2*b*x+2*a))/b^2-3/8*x*polylog(4,-(1+d)*exp(2*b*x+2*a))/b^3+3/16*polylog(5,-(1+d)*exp(2*b*x+2*a))/b^4
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{4b^4 x^4 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - 2b^4 x^4 \log\left(1 + \frac{e^{-2(a+bx)}}{1+d}\right) + 4b^3 x^3 \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{1+d}\right) + 6b^2}{16b^4}$$

input

```
Integrate[x^3*ArcTanh[1 + d + d*Tanh[a + b*x]],x]
```

output

```
(4*b^4*x^4*ArcTanh[1 + d + d*Tanh[a + b*x]] - 2*b^4*x^4*Log[1 + 1/((1 + d)*E^(2*(a + b*x)))] + 4*b^3*x^3*PolyLog[2, -(1/((1 + d)*E^(2*(a + b*x))))] + 6*b^2*x^2*PolyLog[3, -(1/((1 + d)*E^(2*(a + b*x))))] + 6*b*x*PolyLog[4, -(1/((1 + d)*E^(2*(a + b*x))))] + 3*PolyLog[5, -(1/((1 + d)*E^(2*(a + b*x))))])/(16*b^4)
```

**Rubi [A] (verified)**

Time = 1.34 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.28, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6793, 2615, 2620, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(d \tanh(a + bx) + d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{4}b \int \frac{x^4}{e^{2a+2bx}(d+1)+1} dx + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \int \frac{e^{2a+2bx}x^4}{e^{2a+2bx}(d+1)+1} dx \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \left( \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \int x^3 \log(e^{2a+2bx}(d+1) + 1) dx}{b(d+1)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \left( \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \left( \frac{3 \int x^2 \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx})) dx}{2b} - \frac{x^3 \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx})}{2b} \right)}{b(d+1)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \left( \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx})}{2b} \right) - \frac{\int x \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}) dx}{b}}{2b} \right)}{b(d+1)} \right) \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \left( \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx})}{2b} \right) - \frac{x \operatorname{PolyLog}(4, -((d+1)e^{2a+2bx})}{2b} \right) - \frac{\int \operatorname{PolyLog}(4, -((d+1)e^{2a+2bx}) dx}{b}}{2b} \right)}{b(d+1)} \right) \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 2720

$$\begin{aligned}
 & \left( \frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \right) \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{2b} - \frac{x \operatorname{PolyLog}(4, -((d+1)e^{2a+2bx}))}{2b} \right) - \frac{f e^{-2a}}{b} \right)}{2b} \right) \\
 & \qquad \qquad \qquad \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a + bx) + d + 1) \\
 & \qquad \qquad \qquad \downarrow \text{7143} \\
 & \frac{1}{4}x^4 \operatorname{arctanh}(d \tanh(a + bx) + d + 1) + \\
 & \left( \frac{1}{4}b \left( \frac{x^5}{5} - (d+1) \right) \frac{x^4 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{2b} - \frac{x \operatorname{PolyLog}(4, -((d+1)e^{2a+2bx}))}{2b} \right) - \frac{\operatorname{PolyLog}(5, -((d+1)e^{2a+2bx}))}{b} \right)}{2b} \right) \frac{1}{b(d+1)}
 \end{aligned}$$

input `Int[x^3*ArcTanh[1 + d + d*Tanh[a + b*x]],x]`

output `(x^4*ArcTanh[1 + d + d*Tanh[a + b*x]])/4 + (b*(x^5/5 - (1 + d)*((x^4*Log[1 + (1 + d)*E^(2*a + 2*b*x)]))/(2*b*(1 + d)) - (2*(-1/2*(x^3*PolyLog[2, -((1 + d)*E^(2*a + 2*b*x)])))/b + (3*((x^2*PolyLog[3, -((1 + d)*E^(2*a + 2*b*x)])))/(2*b) - ((x*PolyLog[4, -((1 + d)*E^(2*a + 2*b*x)])))/(2*b) - PolyLog[5, -((1 + d)*E^(2*a + 2*b*x)]/(4*b^2))/b)/(b*(1 + d)))/4`

## Definitions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 6793

```
Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.17 (sec) , antiderivative size = 1721, normalized size of antiderivative = 11.10

method	result	size
risch	Expression too large to display	1721

input

```
int(x^3*arctanh(1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/8*d/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*x^4+3/16/b^4*d/(1+d)*polylog(5,-(1
+d)*exp(2*b*x+2*a))+1/2/b^4*a^4/(1+d)*ln(1+exp(b*x+a)*(-d-1)^(1/2))+1/2/b^
4*a^4/(1+d)*ln(1-exp(b*x+a)*(-d-1)^(1/2))+1/2/b^4*a^3/(1+d)*dilog(1+exp(b*
x+a)*(-d-1)^(1/2))+1/2/b^4*a^3/(1+d)*dilog(1-exp(b*x+a)*(-d-1)^(1/2))-1/4/
b/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*x^3-3/8/b^4/(1+d)*ln(1+(1+d)*exp(
2*b*x+2*a))*a^4+3/8/b^2/(1+d)*polylog(3,-(1+d)*exp(2*b*x+2*a))*x^2-1/4/b^4
/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*a^3-3/8/b^3/(1+d)*polylog(4,-(1+d)
*exp(2*b*x+2*a))*x+1/8*x^4*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)+1)-1/8/b^4*a
^4/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)+1)+1/20*b*x^5-1/4*x^4*ln(exp(b
*x+a))-1/8/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*x^4+3/16/b^4/(1+d)*polylog(5,-
(1+d)*exp(2*b*x+2*a))+1/2/b^4*d*a^4/(1+d)*ln(1+exp(b*x+a)*(-d-1)^(1/2))+1/
2/b^4*d*a^4/(1+d)*ln(1-exp(b*x+a)*(-d-1)^(1/2))+1/2/b^4*d*a^3/(1+d)*dilog(
1+exp(b*x+a)*(-d-1)^(1/2))+1/2/b^4*d*a^3/(1+d)*dilog(1-exp(b*x+a)*(-d-1)^(
1/2))+1/2/b^3*a^3/(1+d)*ln(1+exp(b*x+a)*(-d-1)^(1/2))*x+1/2/b^3*a^3/(1+d)*
ln(1-exp(b*x+a)*(-d-1)^(1/2))*x-1/4/b*d/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2
*a))*x^3-1/16*(2*I*Pi+2*ln(d)-I*Pi*csgn(I*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)
+1))*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)+1))^2+I*Pi
*csgn(I/(exp(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2*csgn(I*d)-2*I*Pi*csgn(I/(ex
p(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2+I*Pi*csgn(I*exp(2*b*x+2*a))*csgn(I*exp
(2*b*x+2*a)/(exp(2*b*x+2*a)+1))^2-I*Pi*csgn(I*exp(b*x+a))^2*csgn(I*exp(...
```



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 451 vs.  $2(135) = 270$ .

Time = 0.10 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.91

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(x^3*arctanh(1+d+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```
1/40*(2*b^5*x^5 + 5*b^4*x^4*log(-((d + 2)*cosh(b*x + a) + d*sinh(b*x + a))
/(d*cosh(b*x + a) + d*sinh(b*x + a))) - 20*b^3*x^3*dilog(1/2*sqrt(-4*d - 4)
)*(cosh(b*x + a) + sinh(b*x + a))) - 20*b^3*x^3*dilog(-1/2*sqrt(-4*d - 4)*
(cosh(b*x + a) + sinh(b*x + a))) - 5*a^4*log(2*(d + 1)*cosh(b*x + a) + 2*(
d + 1)*sinh(b*x + a) + sqrt(-4*d - 4)) - 5*a^4*log(2*(d + 1)*cosh(b*x + a)
+ 2*(d + 1)*sinh(b*x + a) - sqrt(-4*d - 4)) + 60*b^2*x^2*polylog(3, 1/2*sq
rt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) + 60*b^2*x^2*polylog(3, -1/2
*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, 1/2
*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, -1/2
*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 5*(b^4*x^4 - a^4)*log(1
/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 5*(b^4*x^4 - a^4)
*log(-1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 120*polylo
g(5, 1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) + 120*polylog(5,
-1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^4
```

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^3 \operatorname{atanh}(d \tanh(a + bx) + d + 1) dx$$

input `integrate(x**3*atanh(1+d+d*tanh(b*x+a)),x)`

output `Integral(x**3*atanh(d*tanh(a + b*x) + d + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.96

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{1}{4} x^4 \operatorname{artanh}(d \tanh(bx + a) + d + 1) + \frac{1}{40} \left( \frac{2x^5}{d} - \frac{5(2b^4 x^4 \log((d+1)e^{(2bx+2a)} + 1) + 4b^3 x^3 \operatorname{Li}_2(-(d+1)e^{(2bx+2a)}) - 6b^2 x^2 \operatorname{Li}_3(-(d+1)e^{(2bx+2a)}) + 4b x \operatorname{Li}_4(-(d+1)e^{(2bx+2a)}) - \operatorname{Li}_5(-(d+1)e^{(2bx+2a)}))}{b^5 d} \right)$$

input `integrate(x^3*arctanh(1+d*d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/4*x^4*arctanh(d*tanh(b*x + a) + d + 1) + 1/40*(2*x^5/d - 5*(2*b^4*x^4*log((d + 1)*e^(2*b*x + 2*a) + 1) + 4*b^3*x^3*dilog(-(d + 1)*e^(2*b*x + 2*a)) - 6*b^2*x^2*polylog(3, -(d + 1)*e^(2*b*x + 2*a)) + 6*b*x*polylog(4, -(d + 1)*e^(2*b*x + 2*a)) - 3*polylog(5, -(d + 1)*e^(2*b*x + 2*a)))/(b^5*d))*b*d`

**Giac [F]**

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^3 \operatorname{artanh}(d \tanh(bx + a) + d + 1) dx$$

input `integrate(x^3*arctanh(1+d*d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x^3*arctanh(d*tanh(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^3 \operatorname{atanh}(d + d \tanh(a + bx) + 1) dx$$

input `int(x^3*atanh(d + d*tanh(a + b*x) + 1),x)`

output `int(x^3*atanh(d + d*tanh(a + b*x) + 1), x)`

### Reduce [F]

$$\int x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a) d + d + 1) x^3 dx$$

input `int(x^3*atanh(1+d+d*tanh(b*x+a)), x)`

output `int(atanh(tanh(a + b*x)*d + d + 1)*x**3, x)`

### 3.289 $\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$

Optimal result	1999
Mathematica [A] (verified)	2000
Rubi [A] (verified)	2000
Maple [C] (warning: unable to verify)	2003
Fricas [B] (verification not implemented)	2004
Sympy [F]	2005
Maxima [A] (verification not implemented)	2005
Giac [F]	2006
Mupad [F(-1)]	2006
Reduce [F]	2006

#### Optimal result

Integrand size = 16, antiderivative size = 128

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - \frac{1}{6}x^3 \log(1 + (1 + d)e^{2a+2bx}) - \frac{x^2 \operatorname{PolyLog}(2, -((1 + d)e^{2a+2bx}))}{4b} + \frac{x \operatorname{PolyLog}(3, -((1 + d)e^{2a+2bx}))}{4b^2} - \frac{\operatorname{PolyLog}(4, -((1 + d)e^{2a+2bx}))}{8b^3}$$

output

```
1/12*b*x^4+1/3*x^3*arctanh(1+d+d*tanh(b*x+a))-1/6*x^3*ln(1+(1+d)*exp(2*b*x+2*a))-1/4*x^2*polylog(2,-(1+d)*exp(2*b*x+2*a))/b+1/4*x*polylog(3,-(1+d)*exp(2*b*x+2*a))/b^2-1/8*polylog(4,-(1+d)*exp(2*b*x+2*a))/b^3
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - 4b^3 x^3 \log\left(1 + \frac{e^{-2(a+bx)}}{1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{1+d}\right) + 6bx}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 + d + d*Tanh[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTanh[1 + d + d*Tanh[a + b*x]] - 4*b^3*x^3*Log[1 + 1/((1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[2, -(1/((1 + d)*E^(2*(a + b*x))))] + 6*b*x*PolyLog[3, -(1/((1 + d)*E^(2*(a + b*x))))] + 3*PolyLog[4, -(1/((1 + d)*E^(2*(a + b*x))))])/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.82 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.29, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {6793, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d \tanh(a + bx) + d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{3}b \int \frac{x^3}{e^{2a+2bx}(d+1)+1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \int \frac{e^{2a+2bx} x^3}{e^{2a+2bx}(d+1)+1} dx \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \left( \frac{x^3 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{3 \int x^2 \log(e^{2a+2bx}(d+1) + 1) dx}{2b(d+1)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \left( \frac{x^3 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{3 \left( \frac{\int x \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx})) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx}))}{2b} \right)}{2b(d+1)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \left( \frac{x^3 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{2b} - \frac{\int \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx})) dx}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{4b^2} \right)}{2b(d+1)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 2720

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \left( \frac{x^3 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx})) de^{2bx}}{b}}{4b^2} \right)}{2b(d+1)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 7143

$$\frac{1}{3}b \left( \frac{x^4}{4} - (d+1) \left( \frac{x^3 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{2b} - \frac{\operatorname{PolyLog}(4, -((d+1)e^{2a+2bx}))}{4b^2}}{b} - \frac{x^2 \operatorname{PolyLog}(4, -((d+1)e^{2a+2bx}))}{4b^2} \right)}{2b(d+1)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tanh(a+bx) + d+1) +$$

input `Int[x^2*ArcTanh[1 + d + d*Tanh[a + b*x]],x]`

output

$$\frac{(x^3 \operatorname{ArcTanh}[1 + d + d \operatorname{Tanh}[a + b x]])}{3} + \frac{(b(x^4/4 - (1 + d)((x^3 \operatorname{Log}[1 + (1 + d)E^{2a + 2bx}]])/(2b(1 + d)) - (3(-1/2(x^2 \operatorname{PolyLog}[2, -((1 + d)E^{2a + 2bx}]])/b + ((x \operatorname{PolyLog}[3, -((1 + d)E^{2a + 2bx}]])/(2b) - \operatorname{PolyLog}[4, -((1 + d)E^{2a + 2bx}]]/(4b^2))/b))/(2b(1 + d))))}{3}$$

### Defintions of rubi rules used

rule 2615

$$\operatorname{Int}[\frac{(c + d x)^m}{(a + b(F^{g(e + f x)}))^{n+1}}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\frac{(c + d x)^{m+1}}{a d (m+1)}, x] - \operatorname{Simp}[\frac{b}{a} \operatorname{Int}[\frac{(c + d x)^m (F^{g(e + f x)})^n}{(a + b(F^{g(e + f x)}))^{n+1}}, x], x] /;$$

$$\operatorname{FreeQ}\{F, a, b, c, d, e, f, g, n, x\} \ \&\& \ \operatorname{IGtQ}[m, 0]$$

rule 2620

$$\operatorname{Int}[\frac{(F^{g(e + f x)})^n (c + d x)^m}{(a + b(F^{g(e + f x)})^n)}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\frac{(c + d x)^m (b f g n \operatorname{Log}[F]) \operatorname{Log}[1 + b(F^{g(e + f x)})^n/a]}{d(m/(b f g n \operatorname{Log}[F])) \operatorname{Int}[(c + d x)^{m-1} \operatorname{Log}[1 + b(F^{g(e + f x)})^n/a]}], x], x] /;$$

$$\operatorname{FreeQ}\{F, a, b, c, d, e, f, g, n, x\} \ \&\& \ \operatorname{IGtQ}[m, 0]$$

rule 2720

$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{With}\{v = \operatorname{FunctionOfExponential}[u, x]\}, \operatorname{Simp}[v/D[v, x] \operatorname{Subst}[\operatorname{Int}[\operatorname{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /;$$

$$\operatorname{FunctionOfExponentialQ}[u, x] \ \&\& \ \operatorname{!MatchQ}[u, (w_*)^{(a_*)^{(v_*)^{(n_*)^{(m_*)}}}} /;$$

$$\operatorname{FreeQ}\{a, m, n, x\} \ \&\& \ \operatorname{IntegerQ}[m n] \ \&\& \ \operatorname{!MatchQ}[u, E^{(c_*)^{(a_*)^{(b_*)^{(x_*)}}}} * (F_)[v_]] /;$$

$$\operatorname{FreeQ}\{a, b, c, x\} \ \&\& \ \operatorname{InverseFunctionQ}[F[x]]$$

rule 3011

$$\operatorname{Int}[\operatorname{Log}[1 + (e + d x)^m (F^{c(a + b x)})^n] * (f + g x)^m, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-f + g x)^m * (\operatorname{PolyLog}[2, (-e) * (F^{c(a + b x)})^n] / (b c n \operatorname{Log}[F]))], x] + \operatorname{Simp}[g * (m / (b c n \operatorname{Log}[F])) \operatorname{Int}[(f + g x)^{m-1} * \operatorname{PolyLog}[2, (-e) * (F^{c(a + b x)})^n], x], x] /;$$

$$\operatorname{FreeQ}\{F, a, b, c, e, f, g, n, x\} \ \&\& \ \operatorname{GtQ}[m, 0]$$

rule 6793

```
Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(
m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a
+ 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c
- d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.70 (sec) , antiderivative size = 1662, normalized size of antiderivative = 12.98

method	result	size
risch	Expression too large to display	1662

input

```
int(x^2*arctanh(1+d+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```



output

```

1/12*b*x^4-1/3*x^3*ln(exp(b*x+a))-1/6/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*x^3
-1/8/b^3/(1+d)*polylog(4,-(1+d)*exp(2*b*x+2*a))+1/6/b^3*a^3/(1+d)*ln(d*exp
(2*b*x+2*a)+exp(2*b*x+2*a)+1)-1/2/b^3*a^3/(1+d)*ln(1+exp(b*x+a)*(-d-1)^(1/
2))-1/2/b^3*a^3/(1+d)*ln(1-exp(b*x+a)*(-d-1)^(1/2))-1/2/b^3*a^2/(1+d)*dilo
g(1+exp(b*x+a)*(-d-1)^(1/2))-1/2/b^3*a^2/(1+d)*dilog(1-exp(b*x+a)*(-d-1)^(
1/2))-1/4/b/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*x^2+1/3/b^3/(1+d)*ln(1+
(1+d)*exp(2*b*x+2*a))*a^3+1/4/b^3/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*a
^2+1/4/b^2/(1+d)*polylog(3,-(1+d)*exp(2*b*x+2*a))*x-1/8/b^3*d/(1+d)*polylo
g(4,-(1+d)*exp(2*b*x+2*a))-1/6*d/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*x^3-1/2/
b^3*d*a^3/(1+d)*ln(1+exp(b*x+a)*(-d-1)^(1/2))-1/2/b^3*d*a^3/(1+d)*ln(1-exp
(b*x+a)*(-d-1)^(1/2))-1/2/b^3*d*a^2/(1+d)*dilog(1+exp(b*x+a)*(-d-1)^(1/2))
-1/2/b^3*d*a^2/(1+d)*dilog(1-exp(b*x+a)*(-d-1)^(1/2))+1/2/b^2/(1+d)*ln(1+(
1+d)*exp(2*b*x+2*a))*a^2*x-1/4/b*d/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*
x^2+1/3/b^3*d/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*a^3+1/4/b^3*d/(1+d)*polylog
(2,-(1+d)*exp(2*b*x+2*a))*a^2+1/4/b^2*d/(1+d)*polylog(3,-(1+d)*exp(2*b*x+2
*a))*x+1/6/b^3*d*a^3/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)+1)-1/2/b^2*a
^2/(1+d)*x*ln(1+exp(b*x+a)*(-d-1)^(1/2))-1/2/b^2*a^2/(1+d)*x*ln(1-exp(b*x+
a)*(-d-1)^(1/2))+1/2/b^2*d/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*a^2*x-1/2/b^2*
d*a^2/(1+d)*x*ln(1+exp(b*x+a)*(-d-1)^(1/2))-1/2/b^2*d*a^2/(1+d)*x*ln(1-exp
(b*x+a)*(-d-1)^(1/2))+1/6*x^3*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)+1)-1/1...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 382 vs.  $2(111) = 222$ .

Time = 0.10 (sec) , antiderivative size = 382, normalized size of antiderivative = 2.98

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{b^4 x^4 + 2 b^3 x^3 \log\left(-\frac{(d+2) \cosh(bx+a) + d \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) - 6 b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4d-4} (\cosh(bx+a) + \sinh(bx+a))\right)}{1}$$

input

```
integrate(x^2*arctanh(1+d*d*tanh(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(b^4*x^4 + 2*b^3*x^3*log(-((d + 2)*cosh(b*x + a) + d*sinh(b*x + a))/(
d*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b^2*x^2*dilog(1/2*sqrt(-4*d - 4)*(
cosh(b*x + a) + sinh(b*x + a))) - 6*b^2*x^2*dilog(-1/2*sqrt(-4*d - 4)*(cos
h(b*x + a) + sinh(b*x + a))) + 2*a^3*log(2*(d + 1)*cosh(b*x + a) + 2*(d +
1)*sinh(b*x + a) + sqrt(-4*d - 4)) + 2*a^3*log(2*(d + 1)*cosh(b*x + a) + 2
*(d + 1)*sinh(b*x + a) - sqrt(-4*d - 4)) + 12*b*x*polylog(3, 1/2*sqrt(-4*d
- 4)*(cosh(b*x + a) + sinh(b*x + a))) + 12*b*x*polylog(3, -1/2*sqrt(-4*d
- 4)*(cosh(b*x + a) + sinh(b*x + a))) - 2*(b^3*x^3 + a^3)*log(1/2*sqrt(-4*d
- 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 2*(b^3*x^3 + a^3)*log(-1/2*s
qrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 12*polylog(4, 1/2*sq
rt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 12*polylog(4, -1/2*sqrt(-4*
d - 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^3
```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^2 \operatorname{atanh}(d \tanh(a + bx) + d + 1) dx$$

input

```
integrate(x**2*atanh(1+d*d*tanh(b*x+a)),x)
```

output

```
Integral(x**2*atanh(d*tanh(a + b*x) + d + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.57 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.98

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{1}{3} x^3 \operatorname{artanh}(d \tanh(bx + a) + d + 1) + \frac{1}{36} \left( \frac{3x^4}{d} - \frac{2(4b^3x^3 \log((d+1)e^{2bx+2a}) + 1) + 6b^2x^2 \operatorname{Li}_2(-(d+1)e^{2bx+2a}) - 6bx \operatorname{Li}_3(-(d+1)e^{2bx+2a})}{b^4d} \right)$$

input

```
integrate(x^2*arctanh(1+d*d*tanh(b*x+a)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctanh(d*tanh(b*x + a) + d + 1) + 1/36*(3*x^4/d - 2*(4*b^3*x^3*log((d + 1)*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(-(d + 1)*e^(2*b*x + 2*a)) - 6*b*x*polylog(3, -(d + 1)*e^(2*b*x + 2*a)) + 3*polylog(4, -(d + 1)*e^(2*b*x + 2*a)))/(b^4*d))*b*d
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^2 \operatorname{artanh}(d \tanh(bx + a) + d + 1) dx$$

input

```
integrate(x^2*arctanh(1+d+d*tanh(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctanh(d*tanh(b*x + a) + d + 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x^2 \operatorname{atanh}(d + d \tanh(a + bx) + 1) dx$$

input

```
int(x^2*atanh(d + d*tanh(a + b*x) + 1),x)
```

output

```
int(x^2*atanh(d + d*tanh(a + b*x) + 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a) d + d + 1) x^2 dx$$

input

```
int(x^2*atanh(1+d+d*tanh(b*x+a)),x)
```

output `int(atanh(tanh(a + b*x)*d + d + 1)*x**2,x)`

### 3.290 $\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$

Optimal result	2008
Mathematica [A] (verified)	2009
Rubi [A] (verified)	2009
Maple [C] (warning: unable to verify)	2012
Fricas [B] (verification not implemented)	2013
Sympy [F]	2013
Maxima [A] (verification not implemented)	2014
Giac [F]	2014
Mupad [F(-1)]	2014
Reduce [F]	2015

#### Optimal result

Integrand size = 14, antiderivative size = 101

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - \frac{1}{4}x^2 \log(1 + (1 + d)e^{2a+2bx}) - \frac{x \operatorname{PolyLog}(2, -((1 + d)e^{2a+2bx}))}{4b} + \frac{\operatorname{PolyLog}(3, -((1 + d)e^{2a+2bx}))}{8b^2}$$

output

```
1/6*b*x^3+1/2*x^2*arctanh(1+d+d*tanh(b*x+a))-1/4*x^2*ln(1+(1+d)*exp(2*b*x+2*a))-1/4*x*polylog(2,-(1+d)*exp(2*b*x+2*a))/b+1/8*polylog(3,-(1+d)*exp(2*b*x+2*a))/b^2
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.90

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left( 2 \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - \log \left( 1 + \frac{e^{-2(a+bx)}}{1+d} \right) \right) + 2bx \operatorname{PolyLog} \left( 2, -\frac{e^{-2(a+bx)}}{1+d} \right) + \operatorname{PolyLog} \left( 3, -\frac{e^{-2(a+bx)}}{1+d} \right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 + d + d*Tanh[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTanh[1 + d + d*Tanh[a + b*x]] - Log[1 + 1/((1 + d)*E^(2*(a + b*x))])) + 2*b*x*PolyLog[2, -(1/((1 + d)*E^(2*(a + b*x)))] + PolyLog[3, -(1/((1 + d)*E^(2*(a + b*x)))])/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.66 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.30, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6793, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \tanh(a + bx) + d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{2}b \int \frac{x^2}{e^{2a+2bx}(d+1)+1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2}b \left( \frac{x^3}{3} - (d+1) \int \frac{e^{2a+2bx} x^2}{e^{2a+2bx}(d+1)+1} dx \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a + bx) + d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}b\left(\frac{x^3}{3} - (d+1)\left(\frac{x^2 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\int x \log(e^{2a+2bx}(d+1) + 1) dx}{b(d+1)}\right)\right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{2}b\left(\frac{x^3}{3} - (d+1)\left(\frac{x^2 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\int \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx})) dx}{2b} - \frac{x \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx}))}{2b}\right)\right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 2720

$$\frac{1}{2}b\left(\frac{x^3}{3} - (d+1)\left(\frac{x^2 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx})) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx}))}{2b}\right)\right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

↓ 7143

$$\frac{1}{2}b\left(\frac{x^3}{3} - (d+1)\left(\frac{x^2 \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\operatorname{PolyLog}(3, -((d+1)e^{2a+2bx}))}{4b^2} - \frac{x \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx}))}{2b}\right)\right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tanh(a+bx) + d+1)$$

input `Int[x*ArcTanh[1 + d + d*Tanh[a + b*x]], x]`

output `(x^2*ArcTanh[1 + d + d*Tanh[a + b*x]])/2 + (b*(x^3/3 - (1 + d)*((x^2*Log[1 + (1 + d)*E^(2*a + 2*b*x)])/(2*b*(1 + d)) - (-1/2*(x*PolyLog[2, -((1 + d)*E^(2*a + 2*b*x)])]/b + PolyLog[3, -((1 + d)*E^(2*a + 2*b*x)])/(4*b^2)))/(b*(1 + d))))/2`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6793 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`



**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.48 (sec) , antiderivative size = 1579, normalized size of antiderivative = 15.63

method	result	size
risch	Expression too large to display	1579

input `int(x*arctanh(1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/2*x^2*ln(exp(b*x+a))+1/6*b*x^3+1/2/b^2*a^2/(1+d)*ln(1+exp(b*x+a)*(-d-1)
^(1/2))+1/2/b^2*a^2/(1+d)*ln(1-exp(b*x+a)*(-d-1)^(1/2))+1/2/b^2*a/(1+d)*di
log(1+exp(b*x+a)*(-d-1)^(1/2))+1/2/b^2*a/(1+d)*dilog(1-exp(b*x+a)*(-d-1)^(
1/2))-1/4/b^2/(1+d)*polylog(2,-(1+d)*exp(2*b*x+2*a))*a+1/8/b^2*d/(1+d)*pol
ylog(3,-(1+d)*exp(2*b*x+2*a))+1/2/b*d*a/(1+d)*x*ln(1+exp(b*x+a)*(-d-1)^(1/
2))-1/4/b^2/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*a^2-1/4/b/(1+d)*polylog(2,-(1
+d)*exp(2*b*x+2*a))*x-1/4/b^2*a^2/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)
+1)-1/4*d/(1+d)*ln(1+(1+d)*exp(2*b*x+2*a))*x^2-1/4/(1+d)*ln(1+(1+d)*exp(2*
b*x+2*a))*x^2+1/8/b^2/(1+d)*polylog(3,-(1+d)*exp(2*b*x+2*a))+1/4*x^2*ln(d*
exp(2*b*x+2*a)+exp(2*b*x+2*a)+1)-1/8*(2*I*Pi+2*ln(d)-I*Pi*csgn(I*(d*exp(2*
b*x+2*a)+exp(2*b*x+2*a)+1))*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)+ex
p(2*b*x+2*a)+1))^2+I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2*csgn
(I*d)-2*I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2+I*Pi*csgn(I*exp
(2*b*x+2*a))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)+1))^2-I*Pi*csgn(I*exp(b
*x+a))^2*csgn(I*exp(2*b*x+2*a))-I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*
x+2*a)+exp(2*b*x+2*a)+1))^2*csgn(I/(exp(2*b*x+2*a)+1))+I*Pi*csgn(I/(exp(2*
b*x+2*a)+1)*d*exp(2*b*x+2*a))^3-I*Pi*csgn(I*exp(2*b*x+2*a))^3+2*I*Pi*csgn(
I*exp(b*x+a))*csgn(I*exp(2*b*x+2*a))^2-I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b
*x+2*a)+1))^3+I*Pi*csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*exp(2*b*x+2*a)/(exp(2
*b*x+2*a)+1))^2+I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)+exp(2*...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 323 vs.  $2(87) = 174$ .

Time = 0.10 (sec) , antiderivative size = 323, normalized size of antiderivative = 3.20

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{2b^3x^3 + 3b^2x^2 \log\left(-\frac{(d+2)\cosh(bx+a)+d\sinh(bx+a)}{d\cosh(bx+a)+d\sinh(bx+a)}\right) - 6bx \operatorname{Li}_2\left(\frac{1}{2}\sqrt{-4d-4}(\cosh(bx+a) + \sinh(bx+a))\right)}{b^2}$$

input `integrate(x*arctanh(1+d+d*tanh(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*b^3*x^3 + 3*b^2*x^2*log(-((d + 2)*cosh(b*x + a) + d*sinh(b*x + a)) / (d*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b*x*dilog(1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 6*b*x*dilog(-1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) - 3*a^2*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) + sqrt(-4*d - 4)) - 3*a^2*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) - sqrt(-4*d - 4)) - 3*(b^2*x^2 - a^2)*log(1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 3*(b^2*x^2 - a^2)*log(-1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 6*polylog(3, 1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))) + 6*polylog(3, -1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x \operatorname{atanh}(d \tanh(a + bx) + d + 1) dx$$

input `integrate(x*atanh(1+d+d*tanh(b*x+a)),x)`

output `Integral(x*atanh(d*tanh(a + b*x) + d + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{1}{24} \left( \frac{4x^3}{d} - \frac{3(2b^2x^2 \log((d+1)e^{2bx+2a}) + 1) + 2bx \operatorname{Li}_2(-(d+1)e^{2bx+2a}) - \operatorname{Li}_3(-(d+1)e^{2bx+2a}))}{b^3d} \right) + \frac{1}{2} x^2 \operatorname{artanh}(d \tanh(bx + a) + d + 1)$$

input `integrate(x*arctanh(1+d+d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/24*(4*x^3/d - 3*(2*b^2*x^2*log((d + 1)*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(-(d + 1)*e^(2*b*x + 2*a)) - polylog(3, -(d + 1)*e^(2*b*x + 2*a)))/(b^3*d)) * b*d + 1/2*x^2*arctanh(d*tanh(b*x + a) + d + 1)`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x \operatorname{artanh}(d \tanh(bx + a) + d + 1) dx$$

input `integrate(x*arctanh(1+d+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*tanh(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int x \operatorname{atanh}(d + d \tanh(a + bx) + 1) dx$$

input `int(x*atanh(d + d*tanh(a + b*x) + 1),x)`

output `int(x*atanh(d + d*tanh(a + b*x) + 1), x)`

### Reduce [F]

$$\int x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a) d + d + 1) x dx$$

input `int(x*atanh(1+d+d*tanh(b*x+a)),x)`

output `int(atanh(tanh(a + b*x)*d + d + 1)*x,x)`

### 3.291 $\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$

Optimal result	2016
Mathematica [A] (verified)	2016
Rubi [A] (verified)	2017
Maple [B] (verified)	2019
Fricas [B] (verification not implemented)	2019
Sympy [F]	2020
Maxima [A] (verification not implemented)	2020
Giac [F]	2021
Mupad [F(-1)]	2021
Reduce [F]	2021

#### Optimal result

Integrand size = 12, antiderivative size = 69

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \frac{bx^2}{2} + x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) - \frac{1}{2} x \log(1 + (1 + d)e^{2a+2bx}) - \frac{\operatorname{PolyLog}(2, -((1 + d)e^{2a+2bx}))}{4b}$$

output

```
1/2*b*x^2+x*arctanh(1+d+d*tanh(b*x+a))-1/2*x*ln(1+(1+d)*exp(2*b*x+2*a))-1/4*polylog(2,-(1+d)*exp(2*b*x+2*a))/b
```

#### Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.91

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = x \operatorname{arctanh}(1 + d + d \tanh(a + bx)) + \frac{-2bx \log\left(1 + \frac{e^{-2(a+bx)}}{1+d}\right) + \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{1+d}\right)}{4b}$$

input `Integrate[ArcTanh[1 + d + d*Tanh[a + b*x]], x]`

output `x*ArcTanh[1 + d + d*Tanh[a + b*x]] + (-2*b*x*Log[1 + 1/((1 + d)*E^(2*(a + b*x)))] + PolyLog[2, -(1/((1 + d)*E^(2*(a + b*x))))])/(4*b)`

### Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.30, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6785, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arctanh}(d \tanh(a + bx) + d + 1) dx \\
 & \quad \downarrow \text{6785} \\
 & b \int \frac{x}{e^{2a+2bx}(d+1) + 1} dx + x \operatorname{arctanh}(d \tanh(a + bx) + d + 1) \\
 & \quad \downarrow \text{2615} \\
 & b \left( \frac{x^2}{2} - (d+1) \int \frac{e^{2a+2bx} x}{e^{2a+2bx}(d+1) + 1} dx \right) + x \operatorname{arctanh}(d \tanh(a + bx) + d + 1) \\
 & \quad \downarrow \text{2620} \\
 & b \left( \frac{x^2}{2} - (d+1) \left( \frac{x \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\int \log(e^{2a+2bx}(d+1) + 1) dx}{2b(d+1)} \right) \right) + \\
 & \quad \quad \quad x \operatorname{arctanh}(d \tanh(a + bx) + d + 1) \\
 & \quad \downarrow \text{2715} \\
 & b \left( \frac{x^2}{2} - (d+1) \left( \frac{x \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} - \frac{\int e^{-2a-2bx} \log(e^{2a+2bx}(d+1) + 1) de^{2a+2bx}}{4b^2(d+1)} \right) \right) + \\
 & \quad \quad \quad x \operatorname{arctanh}(d \tanh(a + bx) + d + 1) \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

$$b \left( \frac{x^2}{2} - (d+1) \left( \frac{x \operatorname{arctanh}(d \tanh(a+bx) + d + 1) + \operatorname{PolyLog}(2, -((d+1)e^{2a+2bx}))}{4b^2(d+1)} + \frac{x \log((d+1)e^{2a+2bx} + 1)}{2b(d+1)} \right) \right)$$

input `Int[ArcTanh[1 + d + d*Tanh[a + b*x]], x]`

output `x*ArcTanh[1 + d + d*Tanh[a + b*x]] + b*(x^2/2 - (1 + d)*((x*Log[1 + (1 + d)*E^(2*a + 2*b*x)])/(2*b*(1 + d)) + PolyLog[2, -((1 + d)*E^(2*a + 2*b*x)]/(4*b^2*(1 + d)))))`

### Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int((((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_.) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_.))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_.) + (e_.)*(x_.)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 6785 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]], x_Symbol] := Simp[x*ArcTanh[c + d*Tanh[a + b*x]], x] + Simp[b Int[x/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, 1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(61) = 122.

Time = 2.30 (sec) , antiderivative size = 255, normalized size of antiderivative = 3.70

method	result
derivativedivides	$\frac{\operatorname{arctanh}(1+d+d \tanh(bx+a))d \ln(d+d \tanh(bx+a))}{2} - \frac{\operatorname{arctanh}(1+d+d \tanh(bx+a))d \ln(-d \tanh(bx+a)+d)}{2} - d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \tanh(bx+a)-2}{2}\right)}{2} \right)$
default	$\frac{\operatorname{arctanh}(1+d+d \tanh(bx+a))d \ln(d+d \tanh(bx+a))}{2} - \frac{\operatorname{arctanh}(1+d+d \tanh(bx+a))d \ln(-d \tanh(bx+a)+d)}{2} - d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \tanh(bx+a)-2}{2}\right)}{2} \right)$
risch	Expression too large to display

input

```
int(arctanh(1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/b/d*(1/2*arctanh(1+d*d*tanh(b*x+a))*d*ln(d+d*tanh(b*x+a))-1/2*arctanh(1+d*d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)+d)-1/2*d^2*(1/d*(-1/2*dilog((-d*tanh(b*x+a)-d-2)/(-2*d-2))-1/2*ln(-d*tanh(b*x+a)+d)*ln((-d*tanh(b*x+a)-d-2)/(-2*d-2))+1/2*dilog(-1/2*(-d*tanh(b*x+a)-d)/d)+1/2*ln(-d*tanh(b*x+a)+d)*ln(-1/2*(-d*tanh(b*x+a)-d)/d))-1/d*(-1/2*dilog(1/2*d*tanh(b*x+a)+1/2*d+1)-1/2*ln(d+d*tanh(b*x+a))*ln(1/2*d*tanh(b*x+a)+1/2*d+1)+1/4*ln(d+d*tanh(b*x+a))^2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(60) = 120.

Time = 0.09 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.46

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{b^2 x^2 + bx \log\left(-\frac{(d+2) \cosh(bx+a) + d \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) + a \log(2(d+1) \cosh(bx+a) + 2(d+1) \sinh(bx+a) + \sqrt{4(d+1)^2 - 1}}{2}}{2}$$

input

```
integrate(arctanh(1+d*d*tanh(b*x+a)),x, algorithm="fricas")
```



output

```
1/2*(b^2*x^2 + b*x*log(-((d + 2)*cosh(b*x + a) + d*sinh(b*x + a))/(d*cosh(
b*x + a) + d*sinh(b*x + a))) + a*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*s
inh(b*x + a) + sqrt(-4*d - 4)) + a*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)
*sinh(b*x + a) - sqrt(-4*d - 4)) - (b*x + a)*log(1/2*sqrt(-4*d - 4)*(cosh(
b*x + a) + sinh(b*x + a)) + 1) - (b*x + a)*log(-1/2*sqrt(-4*d - 4)*(cosh(b
*x + a) + sinh(b*x + a)) + 1) - dilog(1/2*sqrt(-4*d - 4)*(cosh(b*x + a) +
sinh(b*x + a))) - dilog(-1/2*sqrt(-4*d - 4)*(cosh(b*x + a) + sinh(b*x + a
))))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(d \tanh(a + bx) + d + 1) dx$$

input

```
integrate(atanh(1+d*d*tanh(b*x+a)),x)
```

output

```
Integral(atanh(d*tanh(a + b*x) + d + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx$$

$$= \frac{1}{4} bd \left( \frac{2x^2}{d} - \frac{2bx \log((d+1)e^{(2bx+2a)} + 1) + \operatorname{Li}_2(-(d+1)e^{(2bx+2a)})}{b^2 d} \right)$$

$$+ x \operatorname{artanh}(d \tanh(bx + a) + d + 1)$$

input

```
integrate(arctanh(1+d*d*tanh(b*x+a)),x, algorithm="maxima")
```

output

```
1/4*b*d*(2*x^2/d - (2*b*x*log((d + 1)*e^(2*b*x + 2*a) + 1) + dilog(-(d + 1)
)*e^(2*b*x + 2*a)))/(b^2*d) + x*arctanh(d*tanh(b*x + a) + d + 1)
```

**Giac [F]**

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{artanh}(d \tanh(bx + a) + d + 1) dx$$

input `integrate(arctanh(1+d+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*tanh(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(d + d \tanh(a + bx) + 1) dx$$

input `int(atanh(d + d*tanh(a + b*x) + 1),x)`

output `int(atanh(d + d*tanh(a + b*x) + 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 + d + d \tanh(a + bx)) dx = \int \operatorname{atanh}(\tanh(bx + a) d + d + 1) dx$$

input `int(atanh(1+d+d*tanh(b*x+a)),x)`

output `int(atanh(tanh(a + b*x)*d + d + 1),x)`

### 3.292 $\int \frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x} dx$

Optimal result	2022
Mathematica [N/A]	2022
Rubi [N/A]	2023
Maple [N/A]	2023
Fricas [N/A]	2024
Sympy [N/A]	2024
Maxima [N/A]	2024
Giac [N/A]	2025
Mupad [N/A]	2025
Reduce [N/A]	2026

#### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(1+d+d*tanh(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 3.55 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1+d+d \tanh(a+bx))}{x} dx$$

input `Integrate[ArcTanh[1 + d + d*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 + d + d*Tanh[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \tanh(a + bx) + d + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \tanh(a + bx) + d + 1)}{x} dx$$

input `Int[ArcTanh[1 + d + d*Tanh[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(bx + a))}{x} dx$$

input `int(arctanh(1+d+d*tanh(b*x+a))/x,x)`

output `int(arctanh(1+d+d*tanh(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*tanh(b*x + a) + d + 1)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.62 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \tanh(a + bx) + d + 1)}{x} dx$$

input `integrate(atanh(1+d+d*tanh(b*x+a))/x,x)`

output `Integral(atanh(d*tanh(a + b*x) + d + 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.63 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*tanh(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctanh(d*tanh(b*x + a) + d + 1)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\arctanh(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tanh(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*tanh(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*tanh(b*x + a) + d + 1)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.33 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\arctanh(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d + d \tanh(a + bx) + 1)}{x} dx$$

input `int(atanh(d + d*tanh(a + b*x) + 1)/x,x)`

output `int(atanh(d + d*tanh(a + b*x) + 1)/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tanh(bx + a) d + d + 1)}{x} dx$$

input `int(atanh(1+d+d*tanh(b*x+a))/x,x)`output `int(atanh(tanh(a + b*x)*d + d + 1)/x,x)`

### 3.293 $\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$

Optimal result	2027
Mathematica [A] (verified)	2028
Rubi [A] (verified)	2028
Maple [C] (warning: unable to verify)	2032
Fricas [B] (verification not implemented)	2033
Sympy [F]	2033
Maxima [A] (verification not implemented)	2034
Giac [F]	2034
Mupad [F(-1)]	2034
Reduce [F]	2035

#### Optimal result

Integrand size = 19, antiderivative size = 168

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \frac{bx^5}{20} + \frac{1}{4}x^4 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - \frac{1}{8}x^4 \log(1 + (1 - d)e^{2a+2bx}) - \frac{x^3 \operatorname{PolyLog}(2, -((1 - d)e^{2a+2bx}))}{4b} + \frac{3x^2 \operatorname{PolyLog}(3, -((1 - d)e^{2a+2bx}))}{8b^2} - \frac{3x \operatorname{PolyLog}(4, -((1 - d)e^{2a+2bx}))}{8b^3} + \frac{3 \operatorname{PolyLog}(5, -((1 - d)e^{2a+2bx}))}{16b^4}$$

output

```
1/20*b*x^5-1/4*x^4*arctanh(-1+d+d*tanh(b*x+a))-1/8*x^4*ln(1+(1-d)*exp(2*b*x+2*a))-1/4*x^3*polylog(2,-(1-d)*exp(2*b*x+2*a))/b+3/8*x^2*polylog(3,-(1-d)*exp(2*b*x+2*a))/b^2-3/8*x*polylog(4,-(1-d)*exp(2*b*x+2*a))/b^3+3/16*polylog(5,-(1-d)*exp(2*b*x+2*a))/b^4
```



**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.88

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{4b^4 x^4 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - 2b^4 x^4 \log\left(1 - \frac{e^{-2(a+bx)}}{-1+d}\right) + 4b^3 x^3 \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{-1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(3, \frac{e^{-2(a+bx)}}{-1+d}\right) + 6b x \operatorname{PolyLog}\left(4, \frac{e^{-2(a+bx)}}{-1+d}\right) + 3 \operatorname{PolyLog}\left(5, \frac{e^{-2(a+bx)}}{-1+d}\right)}{16b^4}$$

input

```
Integrate[x^3*ArcTanh[1 - d - d*Tanh[a + b*x]],x]
```

output

```
(4*b^4*x^4*ArcTanh[1 - d - d*Tanh[a + b*x]] - 2*b^4*x^4*Log[1 - 1/((-1 + d)*E^(2*(a + b*x)))] + 4*b^3*x^3*PolyLog[2, 1/((-1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[3, 1/((-1 + d)*E^(2*(a + b*x)))] + 6*b*x*PolyLog[4, 1/((-1 + d)*E^(2*(a + b*x)))] + 3*PolyLog[5, 1/((-1 + d)*E^(2*(a + b*x)))])/(16*b^4)
```

**Rubi [A] (verified)**

Time = 1.09 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.30, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {6793, 2615, 2620, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{4}b \int \frac{x^4}{e^{2a+2bx}(1-d)+1} dx + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \int \frac{e^{2a+2bx}x^4}{e^{2a+2bx}(1-d)+1} dx \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \int x^3 \log(e^{2a+2bx}(1-d) + 1) dx}{b(1-d)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \left( \frac{3 \int x^2 \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx})) dx}{2b} - \frac{x^3 \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{2b} - \frac{\int x \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx})) dx}{b} \right)}{2b} \right)}{b(1-d)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{2b} - \frac{x \operatorname{PolyLog}(4, -((1-d)e^{2a+2bx}))}{2b} \right) - \frac{\int \operatorname{PolyLog}(4, -((1-d)e^{2a+2bx})) dx}{b} \right)}{2b} \right)}{b(1-d)} \right) \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \right) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, -(1-d)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, -(1-d)e^{2a+2bx})}{2b} \right) - \frac{f e^{-2a}}{b} \right)}{2b} \right)$$

$\frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$

$\downarrow$  7143

$$\frac{1}{4}x^4 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1) + \frac{1}{4}b \left( \frac{x^5}{5} - (1-d) \right) \left( \frac{x^4 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, -(1-d)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, -(1-d)e^{2a+2bx})}{2b} \right) - \frac{\operatorname{PolyLog}(5, -(1-d)e^{2a+2bx})}{b} \right)}{2b} \right)$$

input `Int[x^3*ArcTanh[1 - d - d*Tanh[a + b*x]],x]`

output `(x^4*ArcTanh[1 - d - d*Tanh[a + b*x]])/4 + (b*(x^5/5 - (1 - d)*((x^4*Log[1 + (1 - d)*E^(2*a + 2*b*x)])/(2*b*(1 - d)) - (2*(-1/2*(x^3*PolyLog[2, -((1 - d)*E^(2*a + 2*b*x)])))/b + (3*((x^2*PolyLog[3, -((1 - d)*E^(2*a + 2*b*x)])))/(2*b) - ((x*PolyLog[4, -((1 - d)*E^(2*a + 2*b*x)])))/(2*b) - PolyLog[5, -((1 - d)*E^(2*a + 2*b*x)]/(4*b^2))/b)/(2*b))/(b*(1 - d)))/4`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6793 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)
)*(x_)))]^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.30 (sec) , antiderivative size = 1725, normalized size of antiderivative = 10.27

method	result	size
risch	Expression too large to display	1725

input

```
int(-x^3*arctanh(-1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/8*x^4*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)-1/16*(-I*Pi*csgn(I*(d*exp(2*
b*x+2*a)-exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)-e
p(2*b*x+2*a)-1))^2+2*ln(d)-2*I*Pi+I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*d*exp(2*b
*x+2*a))^2*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)+1))+I*Pi*csgn(I/(exp(2*b*
x+2*a)+1)*d*exp(2*b*x+2*a))^2*csgn(I*d)-I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*d*e
xp(2*b*x+2*a))^3+I*Pi*csgn(I*exp(2*b*x+2*a))*csgn(I*exp(2*b*x+2*a)/(exp(2*
b*x+2*a)+1))^2-I*Pi*csgn(I*exp(b*x+a))^2*csgn(I*exp(2*b*x+2*a))+I*Pi*csgn(
I*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*
b*x+2*a)-exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)+1))-I*Pi*csgn(I/(exp(2*
b*x+2*a)+1)*d*exp(2*b*x+2*a))*csgn(I*d)*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2
*a)+1))-I*Pi*csgn(I*exp(2*b*x+2*a))^3+2*I*Pi*csgn(I*exp(b*x+a))*csgn(I*exp
(2*b*x+2*a))^2-I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)+1))^3-I*Pi*csgn(
I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))^3+2*I*Pi*csgn(I/
(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))^2+I*Pi*csgn(I/(exp
(2*b*x+2*a)+1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)+1))^2-I*Pi*csgn(I/(e
xp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+
2*a)+1))-I*Pi*csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*exp(2*b*x+2*a))*csgn(I*exp
(2*b*x+2*a)/(exp(2*b*x+2*a)+1)))*x^4+1/20*b*x^5-1/4*x^4*ln(exp(b*x+a))+1/8
/b^4*a^4/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)+3/16/b^4*d/(d-1)*poly
log(5, (d-1)*exp(2*b*x+2*a))+1/4/b/(d-1)*polylog(2, (d-1)*exp(2*b*x+2*a))...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 424 vs.  $2(132) = 264$ .

Time = 0.09 (sec) , antiderivative size = 424, normalized size of antiderivative = 2.52

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{2b^5x^5 - 5b^4x^4 \log\left(-\frac{d \cosh(bx+a) + d \sinh(bx+a)}{(d-2) \cosh(bx+a) + d \sinh(bx+a)}\right) - 20b^3x^3 \operatorname{Li}_2(\sqrt{d-1}(\cosh(bx+a) + \sinh(bx+a))) - \dots}{\dots}$$

input `integrate(-x^3*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="fricas")`

output `1/40*(2*b^5*x^5 - 5*b^4*x^4*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/((d - 2)*cosh(b*x + a) + d*sinh(b*x + a))) - 20*b^3*x^3*dilog(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 20*b^3*x^3*dilog(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 5*a^4*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) + 2*sqrt(d - 1)) - 5*a^4*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) - 2*sqrt(d - 1)) + 60*b^2*x^2*polylog(3, sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) + 60*b^2*x^2*polylog(3, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 5*(b^4*x^4 - a^4)*log(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 5*(b^4*x^4 - a^4)*log(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 120*polylog(5, sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) + 120*polylog(5, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^4`

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \int x^3 \operatorname{atanh}(d \tanh(a + bx) + d - 1) dx$$

input `integrate(-x**3*atanh(-1+d+d*tanh(b*x+a)),x)`

output `-Integral(x**3*atanh(d*tanh(a + b*x) + d - 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.87

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = -\frac{1}{4} x^4 \operatorname{artanh}(d \tanh(bx + a) + d - 1) + \frac{1}{40} \left( \frac{2x^5}{d} - \frac{5(2b^4x^4 \log(-(d-1)e^{(2bx+2a)} + 1) + 4b^3x^3 \operatorname{Li}_2((d-1)e^{(2bx+2a)}) - 6b^2x^2 \operatorname{Li}_3((d-1)e^{(2bx+2a)})}{b^5d} \right)$$

input `integrate(-x^3*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="maxima")`

output `-1/4*x^4*arctanh(d*tanh(b*x + a) + d - 1) + 1/40*(2*x^5/d - 5*(2*b^4*x^4*log(-(d - 1)*e^(2*b*x + 2*a) + 1) + 4*b^3*x^3*dilog((d - 1)*e^(2*b*x + 2*a)) - 6*b^2*x^2*polylog(3, (d - 1)*e^(2*b*x + 2*a)) + 6*b*x*polylog(4, (d - 1)*e^(2*b*x + 2*a)) - 3*polylog(5, (d - 1)*e^(2*b*x + 2*a)))/(b^5*d))*b*d`

**Giac [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x^3 \operatorname{artanh}(d \tanh(bx + a) + d - 1) dx$$

input `integrate(-x^3*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(-x^3*arctanh(d*tanh(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x^3 \operatorname{atanh}(d + d \tanh(a + bx) - 1) dx$$

input `int(-x^3*atanh(d + d*tanh(a + b*x) - 1),x)`

output `int(-x^3*atanh(d + d*tanh(a + b*x) - 1), x)`

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \left( \int \operatorname{atanh}(\tanh(bx + a) d + d - 1) x^3 dx \right)$$

input `int(-x^3*atanh(-1+d+d*tanh(b*x+a)), x)`

output `- int(atanh(tanh(a + b*x)*d + d - 1)*x**3, x)`



### 3.294 $\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$

Optimal result	2036
Mathematica [A] (verified)	2037
Rubi [A] (verified)	2037
Maple [C] (warning: unable to verify)	2040
Fricas [B] (verification not implemented)	2041
Sympy [F]	2042
Maxima [A] (verification not implemented)	2042
Giac [F]	2043
Mupad [F(-1)]	2043
Reduce [F]	2043

#### Optimal result

Integrand size = 19, antiderivative size = 139

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - \frac{1}{6}x^3 \log(1 + (1 - d)e^{2a+2bx}) - \frac{x^2 \operatorname{PolyLog}(2, -((1 - d)e^{2a+2bx}))}{4b} + \frac{x \operatorname{PolyLog}(3, -((1 - d)e^{2a+2bx}))}{4b^2} - \frac{\operatorname{PolyLog}(4, -((1 - d)e^{2a+2bx}))}{8b^3}$$

output

```
1/12*b*x^4-1/3*x^3*arctanh(-1+d+d*tanh(b*x+a))-1/6*x^3*ln(1+(1-d)*exp(2*b*x+2*a))-1/4*x^2*polylog(2,-(1-d)*exp(2*b*x+2*a))/b+1/4*x*polylog(3,-(1-d)*exp(2*b*x+2*a))/b^2-1/8*polylog(4,-(1-d)*exp(2*b*x+2*a))/b^3
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.88

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - 4b^3 x^3 \log\left(1 - \frac{e^{-2(a+bx)}}{-1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{-1+d}\right) + 6bx \operatorname{PolyLog}\left(3, \frac{e^{-2(a+bx)}}{-1+d}\right) + 3 \operatorname{PolyLog}\left(4, \frac{e^{-2(a+bx)}}{-1+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 - d - d*Tanh[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTanh[1 - d - d*Tanh[a + b*x]] - 4*b^3*x^3*Log[1 - 1/((-1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[2, 1/((-1 + d)*E^(2*(a + b*x)))] + 6*b*x*PolyLog[3, 1/((-1 + d)*E^(2*(a + b*x)))] + 3*PolyLog[4, 1/((-1 + d)*E^(2*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.87 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.31, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6793, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{3}b \int \frac{x^3}{e^{2a+2bx}(1-d)+1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \int \frac{e^{2a+2bx} x^3}{e^{2a+2bx}(1-d)+1} dx \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \left( \frac{x^3 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{3 \int x^2 \log(e^{2a+2bx}(1-d) + 1) dx}{2b(1-d)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \left( \frac{x^3 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{3 \left( \frac{\int x \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx})) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx}))}{2b} \right)}{2b(1-d)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \left( \frac{x^3 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{2b} - \frac{\int \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx})) dx}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{4b^2} \right)}{2b(1-d)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \left( \frac{x^3 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{3 \left( \frac{\frac{\frac{x \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx})) dx}{b}}{4b^2}}{2b} - \frac{x^2 \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{4b^2} \right)}{2b(1-d)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 7143

$$\frac{1}{3}b \left( \frac{x^4}{4} - (1-d) \left( \frac{x^3 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{3 \left( \frac{\frac{\frac{x \operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{2b} - \frac{\operatorname{PolyLog}(4, -((1-d)e^{2a+2bx}))}{4b^2}}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(4, -((1-d)e^{2a+2bx}))}{4b^2} \right)}{2b(1-d)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1) +$$

input `Int[x^2*ArcTanh[1 - d - d*Tanh[a + b*x]],x]`

output

```
(x^3*ArcTanh[1 - d - d*Tanh[a + b*x]])/3 + (b*(x^4/4 - (1 - d)*((x^3*Log[1
+ (1 - d)*E^(2*a + 2*b*x)]))/(2*b*(1 - d)) - (3*(-1/2*(x^2*PolyLog[2, -((1
- d)*E^(2*a + 2*b*x)]))/b + ((x*PolyLog[3, -((1 - d)*E^(2*a + 2*b*x)]))/(
2*b) - PolyLog[4, -((1 - d)*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b*(1 - d)))
)/3
```

### Defintions of rubi rules used

rule 2615

```
Int[(((c_.) + (d_.)*(x_.))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x
_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[
b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x]
, x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/
((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_.))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6793

```
Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_)^(m
_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(
m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a
+ 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c
- d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_)^(m_.))*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.18 (sec) , antiderivative size = 1668, normalized size of antiderivative = 12.00

method	result	size
risch	Expression too large to display	1668

input

```
int(-x^2*arctanh(-1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/12*b*x^4-1/3*x^3*ln(exp(b*x+a))+1/8/b^3/(d-1)*polylog(4,(d-1)*exp(2*b*x+
2*a))+1/6/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*x^3-1/6*d/(d-1)*ln(1-(d-1)*exp(
2*b*x+2*a))*x^3-1/6/b^3*a^3/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)-1/
8/b^3*d/(d-1)*polylog(4,(d-1)*exp(2*b*x+2*a))+1/4/b/(d-1)*polylog(2,(d-1)*
exp(2*b*x+2*a))*x^2-1/3/b^3/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a^3-1/4/b^3/(
d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*a^2-1/4/b^2/(d-1)*polylog(3,(d-1)*exp
(2*b*x+2*a))*x+1/2/b^3*a^3/(d-1)*ln(1+exp(b*x+a)*(d-1)^(1/2))+1/2/b^3*a^3/
(d-1)*ln(1-exp(b*x+a)*(d-1)^(1/2))+1/2/b^3*a^2/(d-1)*dilog(1-exp(b*x+a)*(d
-1)^(1/2))+1/2/b^3*a^2/(d-1)*dilog(1+exp(b*x+a)*(d-1)^(1/2))+1/6/b^3*d*a^3
/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)-1/2/b^3*d*a^3/(d-1)*ln(1+exp(
b*x+a)*(d-1)^(1/2))-1/2/b^3*d*a^3/(d-1)*ln(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b
^3*d*a^2/(d-1)*dilog(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b^3*d*a^2/(d-1)*dilog(1
+exp(b*x+a)*(d-1)^(1/2))-1/4/b*d/(d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*x^2
+1/3/b^3*d/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a^3+1/4/b^3*d/(d-1)*polylog(2,
(d-1)*exp(2*b*x+2*a))*a^2+1/4/b^2*d/(d-1)*polylog(3,(d-1)*exp(2*b*x+2*a))*
x-1/2/b^2/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a^2*x+1/2/b^2*a^2/(d-1)*x*ln(1+
exp(b*x+a)*(d-1)^(1/2))+1/2/b^2*a^2/(d-1)*x*ln(1-exp(b*x+a)*(d-1)^(1/2))-1
/12*(-I*Pi*csgn(I*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2
*a)+1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))^2+2*ln(d)-2*I*Pi+I*Pi*csgn(I/(
exp(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 360 vs.  $2(109) = 218$ .

Time = 0.10 (sec) , antiderivative size = 360, normalized size of antiderivative = 2.59

$$\int x^2 \operatorname{arctanh}(1-d-d \tanh(a+bx)) dx$$

$$= \frac{b^4 x^4 - 2b^3 x^3 \log\left(-\frac{d \cosh(bx+a)+d \sinh(bx+a)}{(d-2) \cosh(bx+a)+d \sinh(bx+a)}\right) - 6b^2 x^2 \operatorname{Li}_2(\sqrt{d-1}(\cosh(bx+a)+\sinh(bx+a))) - 6b^2 x^2 \operatorname{Li}_2(\sqrt{d-1}(\cosh(bx+a)-\sinh(bx+a)))}{1}$$

input

```
integrate(-x^2*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(b^4*x^4 - 2*b^3*x^3*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/((d - 2)
)*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b^2*x^2*dilog(sqrt(d - 1)*(cosh(b*
x + a) + sinh(b*x + a))) - 6*b^2*x^2*dilog(-sqrt(d - 1)*(cosh(b*x + a) + s
inh(b*x + a))) + 2*a^3*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x +
a) + 2*sqrt(d - 1)) + 2*a^3*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b
*x + a) - 2*sqrt(d - 1)) + 12*b*x*polylog(3, sqrt(d - 1)*(cosh(b*x + a) +
sinh(b*x + a))) + 12*b*x*polylog(3, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x
+ a))) - 2*(b^3*x^3 + a^3)*log(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)
) + 1) - 2*(b^3*x^3 + a^3)*log(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)
) + 1) - 12*polylog(4, sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 12*p
olylog(4, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^3
```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \int x^2 \operatorname{atanh}(d \tanh(a + bx) + d - 1) dx$$

input

```
integrate(-x**2*atanh(-1+d*d*tanh(b*x+a)),x)
```

output

```
-Integral(x**2*atanh(d*tanh(a + b*x) + d - 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.62 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.88

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = -\frac{1}{3} x^3 \operatorname{artanh}(d \tanh(bx + a) + d - 1) + \frac{1}{36} \left( \frac{3x^4}{d} - \frac{2(4b^3x^3 \log(-(d-1)e^{(2bx+2a)}) + 1) + 6b^2x^2 \operatorname{Li}_2((d-1)e^{(2bx+2a)}) - 6bx \operatorname{Li}_3((d-1)e^{(2bx+2a)})}{b^4d} \right)$$

input

```
integrate(-x^2*arctanh(-1+d*d*tanh(b*x+a)),x, algorithm="maxima")
```

output

```
-1/3*x^3*arctanh(d*tanh(b*x + a) + d - 1) + 1/36*(3*x^4/d - 2*(4*b^3*x^3*1
og(-(d - 1)*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog((d - 1)*e^(2*b*x + 2*a)
) - 6*b*x*polylog(3, (d - 1)*e^(2*b*x + 2*a)) + 3*polylog(4, (d - 1)*e^(2*
b*x + 2*a)))/(b^4*d))*b*d
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x^2 \operatorname{artanh}(d \tanh(bx + a) + d - 1) dx$$

input

```
integrate(-x^2*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="giac")
```

output

```
integrate(-x^2*arctanh(d*tanh(b*x + a) + d - 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x^2 \operatorname{atanh}(d + d \tanh(a + bx) - 1) dx$$

input

```
int(-x^2*atanh(d + d*tanh(a + b*x) - 1),x)
```

output

```
int(-x^2*atanh(d + d*tanh(a + b*x) - 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \left( \int \operatorname{atanh}(\tanh(bx + a) d + d - 1) x^2 dx \right)$$

input

```
int(-x^2*atanh(-1+d+d*tanh(b*x+a)),x)
```



output `- int(atanh(tanh(a + b*x)*d + d - 1)*x**2,x)`

### 3.295 $\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$

Optimal result	2045
Mathematica [A] (verified)	2046
Rubi [A] (verified)	2046
Maple [C] (warning: unable to verify)	2049
Fricas [B] (verification not implemented)	2050
Sympy [F]	2050
Maxima [A] (verification not implemented)	2051
Giac [F]	2051
Mupad [F(-1)]	2051
Reduce [F]	2052

#### Optimal result

Integrand size = 17, antiderivative size = 110

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - \frac{1}{4}x^2 \log(1 + (1 - d)e^{2a+2bx}) - \frac{x \operatorname{PolyLog}(2, -((1 - d)e^{2a+2bx}))}{4b} + \frac{\operatorname{PolyLog}(3, -((1 - d)e^{2a+2bx}))}{8b^2}$$

output

```
1/6*b*x^3-1/2*x^2*arctanh(-1+d+d*tanh(b*x+a))-1/4*x^2*ln(1+(1-d)*exp(2*b*x+2*a))-1/4*x*polylog(2,-(1-d)*exp(2*b*x+2*a))/b+1/8*polylog(3,-(1-d)*exp(2*b*x+2*a))/b^2
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.85

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left( 2 \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - \log \left( 1 - \frac{e^{-2(a+bx)}}{-1+d} \right) \right) + 2bx \operatorname{PolyLog} \left( 2, \frac{e^{-2(a+bx)}}{-1+d} \right) + \operatorname{PolyLog} \left( 3, \frac{e^{-2(a+bx)}}{-1+d} \right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 - d - d*Tanh[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTanh[1 - d - d*Tanh[a + b*x]] - Log[1 - 1/((-1 + d)*E^(2*(a + b*x))])) + 2*b*x*PolyLog[2, 1/((-1 + d)*E^(2*(a + b*x))] + PolyLog[3, 1/((-1 + d)*E^(2*(a + b*x)))]/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.33, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$ , Rules used = {6793, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1) dx$$

$$\downarrow \text{6793}$$

$$\frac{1}{2}b \int \frac{x^2}{e^{2a+2bx}(1-d)+1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2}b \left( \frac{x^3}{3} - (1-d) \int \frac{e^{2a+2bx} x^2}{e^{2a+2bx}(1-d)+1} dx \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}b \left( \frac{x^3}{3} - (1-d) \left( \frac{x^2 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\int x \log(e^{2a+2bx}(1-d) + 1) dx}{b(1-d)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{2}b \left( \frac{x^3}{3} - (1-d) \left( \frac{x^2 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\int \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx})) dx}{2b} - \frac{x \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx}))}{2b} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{2}b \left( \frac{x^3}{3} - (1-d) \left( \frac{x^2 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx})) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx}))}{2b} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

↓ 7143

$$\frac{1}{2}b \left( \frac{x^3}{3} - (1-d) \left( \frac{x^2 \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\operatorname{PolyLog}(3, -((1-d)e^{2a+2bx}))}{4b^2} - \frac{x \operatorname{PolyLog}(2, -((1-d)e^{2a+2bx}))}{2b} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

input `Int[x*ArcTanh[1 - d - d*Tanh[a + b*x]], x]`

output `(x^2*ArcTanh[1 - d - d*Tanh[a + b*x]])/2 + (b*(x^3/3 - (1 - d)*((x^2*Log[1 + (1 - d)*E^(2*a + 2*b*x)])/(2*b*(1 - d)) - (-1/2*(x*PolyLog[2, -((1 - d)*E^(2*a + 2*b*x)])]/b + PolyLog[3, -((1 - d)*E^(2*a + 2*b*x)])/(4*b^2)))/(b*(1 - d))))/2`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6793 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.73 (sec) , antiderivative size = 1587, normalized size of antiderivative = 14.43

method	result	size
risch	Expression too large to display	1587

input `int(-x*arctanh(-1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/4/b^2*d/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a^2-1/4/b*d/(d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*x-1/4/b^2*d/(d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*a-1/4/b^2*d*a^2/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)+1/2/b/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a*x-1/2/b*a/(d-1)*x*ln(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b*a/(d-1)*x*ln(1+exp(b*x+a)*(d-1)^(1/2))+1/2/b^2*d*a^2/(d-1)*ln(1-exp(b*x+a)*(d-1)^(1/2))+1/2/b^2*d*a^2/(d-1)*ln(1+exp(b*x+a)*(d-1)^(1/2))+1/2/b^2*d*a/(d-1)*dilog(1-exp(b*x+a)*(d-1)^(1/2))+1/2/b^2*d*a/(d-1)*dilog(1+exp(b*x+a)*(d-1)^(1/2))-1/2*x^2*ln(exp(b*x+a))+1/6*b*x^3-1/4*d/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*x^2+1/4/b^2*a^2/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1)-1/2/b^2*a^2/(d-1)*ln(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b^2*a^2/(d-1)*ln(1+exp(b*x+a)*(d-1)^(1/2))-1/2/b^2*a/(d-1)*dilog(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b^2*a/(d-1)*dilog(1+exp(b*x+a)*(d-1)^(1/2))+1/8/b^2*d/(d-1)*polylog(3,(d-1)*exp(2*b*x+2*a))+1/4/b^2/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a^2+1/4/b/(d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*x+1/4/b^2/(d-1)*polylog(2,(d-1)*exp(2*b*x+2*a))*a+1/2/b*d*a/(d-1)*x*ln(1+exp(b*x+a)*(d-1)^(1/2))+1/2/b*d*a/(d-1)*x*ln(1-exp(b*x+a)*(d-1)^(1/2))-1/2/b*d/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*a*x+1/4/(d-1)*ln(1-(d-1)*exp(2*b*x+2*a))*x^2-1/8/b^2/(d-1)*polylog(3,(d-1)*exp(2*b*x+2*a))-1/8*(-I*Pi*csgn(I*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)+1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)-1))^2+2*ln(d)-2*I*Pi+I*Pi*csgn(I/(exp(2*b*x+2*a)+1)*d*exp(2*b*x+2*a))^2*csgn(I*exp(2*b*x+2*a))...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 306 vs.  $2(86) = 172$ .

Time = 0.09 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.78

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{2 b^3 x^3 - 3 b^2 x^2 \log\left(-\frac{d \cosh(bx+a) + d \sinh(bx+a)}{(d-2) \cosh(bx+a) + d \sinh(bx+a)}\right) - 6 b x \operatorname{Li}_2\left(\sqrt{d-1}(\cosh(bx+a) + \sinh(bx+a))\right) - 6 b x \operatorname{Li}_2\left(-\sqrt{d-1}(\cosh(bx+a) + \sinh(bx+a))\right)}{b^2}$$

input `integrate(-x*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```
1/12*(2*b^3*x^3 - 3*b^2*x^2*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/((d - 2)*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b*x*dilog(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 6*b*x*dilog(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - 3*a^2*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) + 2*sqrt(d - 1)) - 3*a^2*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) - 2*sqrt(d - 1)) - 3*(b^2*x^2 - a^2)*log(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 3*(b^2*x^2 - a^2)*log(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 6*polylog(3, sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a)))) + 6*polylog(3, -sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^2
```

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \int x \operatorname{atanh}(d \tanh(a + bx) + d - 1) dx$$

input `integrate(-x*atanh(-1+d+d*tanh(b*x+a)),x)`

output `-Integral(x*atanh(d*tanh(a + b*x) + d - 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.61 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.91

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$$

$$= \frac{1}{24} \left( \frac{4x^3}{d} - \frac{3(2b^2x^2 \log(-(d-1)e^{(2bx+2a)} + 1) + 2bx \operatorname{Li}_2((d-1)e^{(2bx+2a)}) - \operatorname{Li}_3((d-1)e^{(2bx+2a)}))}{b^3d} \right) - \frac{1}{2} x^2 \operatorname{artanh}(d \tanh(bx + a) + d - 1)$$

input `integrate(-x*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/24*(4*x^3/d - 3*(2*b^2*x^2*log(-(d - 1)*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog((d - 1)*e^(2*b*x + 2*a)) - polylog(3, (d - 1)*e^(2*b*x + 2*a)))/(b^3*d) *b*d - 1/2*x^2*arctanh(d*tanh(b*x + a) + d - 1)`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x \operatorname{artanh}(d \tanh(bx + a) + d - 1) dx$$

input `integrate(-x*arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctanh(d*tanh(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -x \operatorname{atanh}(d + d \tanh(a + bx) - 1) dx$$

input `int(-x*atanh(d + d*tanh(a + b*x) - 1),x)`



output `int(-x*atanh(d + d*tanh(a + b*x) - 1), x)`

### Reduce [F]

$$\int x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \left( \int \operatorname{atanh}(\tanh(bx + a) d + d - 1) x dx \right)$$

input `int(-x*atanh(-1+d+d*tanh(b*x+a)), x)`

output `- int(atanh(tanh(a + b*x)*d + d - 1)*x, x)`

### 3.296 $\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx$

Optimal result	2053
Mathematica [A] (verified)	2053
Rubi [A] (verified)	2054
Maple [B] (verified)	2056
Fricas [B] (verification not implemented)	2056
Sympy [F]	2057
Maxima [A] (verification not implemented)	2057
Giac [F]	2058
Mupad [F(-1)]	2058
Reduce [F]	2058

#### Optimal result

Integrand size = 15, antiderivative size = 76

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \frac{bx^2}{2} + x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) - \frac{1}{2} x \log(1 + (1 - d)e^{2a+2bx}) - \frac{\operatorname{PolyLog}(2, -((1 - d)e^{2a+2bx}))}{4b}$$

output

```
1/2*b*x^2-x*arctanh(-1+d+d*tanh(b*x+a))-1/2*x*ln(1+(1-d)*exp(2*b*x+2*a))-1/4*polylog(2,-(1-d)*exp(2*b*x+2*a))/b
```

#### Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = x \operatorname{arctanh}(1 - d - d \tanh(a + bx)) + \frac{-2bx \log\left(1 - \frac{e^{-2(a+bx)}}{-1+d}\right) + \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{-1+d}\right)}{4b}$$

input

```
Integrate[ArcTanh[1 - d - d*Tanh[a + b*x]], x]
```

output

```
x*ArcTanh[1 - d - d*Tanh[a + b*x]] + (-2*b*x*Log[1 - 1/((-1 + d)*E^(2*(a + b*x)))] + PolyLog[2, 1/((-1 + d)*E^(2*(a + b*x)))])/(4*b)
```

**Rubi [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.36, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6785, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1) dx$$

$$\downarrow 6785$$

$$b \int \frac{x}{e^{2a+2bx}(1-d)+1} dx + x \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

$$\downarrow 2615$$

$$b \left( \frac{x^2}{2} - (1-d) \int \frac{e^{2a+2bx} x}{e^{2a+2bx}(1-d)+1} dx \right) + x \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

$$\downarrow 2620$$

$$b \left( \frac{x^2}{2} - (1-d) \left( \frac{x \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\int \log(e^{2a+2bx}(1-d)+1) dx}{2b(1-d)} \right) \right) + x \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

$$\downarrow 2715$$

$$b \left( \frac{x^2}{2} - (1-d) \left( \frac{x \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} - \frac{\int e^{-2a-2bx} \log(e^{2a+2bx}(1-d)+1) de^{2a+2bx}}{4b^2(1-d)} \right) \right) + x \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

$$\downarrow 2838$$

$$b \left( \frac{x^2}{2} - (1-d) \left( \frac{\operatorname{PolyLog}(2, -((1-d)e^{2a+2bx}))}{4b^2(1-d)} + \frac{x \log((1-d)e^{2a+2bx} + 1)}{2b(1-d)} \right) \right) + x \operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)$$

input `Int[ArcTanh[1 - d - d*Tanh[a + b*x]],x]`

output `x*ArcTanh[1 - d - d*Tanh[a + b*x]] + b*(x^2/2 - (1 - d)*((x*Log[1 + (1 - d)*E^(2*a + 2*b*x)])/(2*b*(1 - d)) + PolyLog[2, -((1 - d)*E^(2*a + 2*b*x)]/(4*b^2*(1 - d))))`

### Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 6785 `Int[ArcTanh[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]], x_Symbol] := Simp[x*ArcTanh[c + d*Tanh[a + b*x]], x] + Simp[b Int[x/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, 1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(66) = 132.

Time = 2.56 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.58

method	result
derivativedivides	$-\frac{\operatorname{arctanh}(-1+d+d \tanh (b x+a)) d \ln (d+d \tanh (b x+a))}{2}-\frac{\operatorname{arctanh}(-1+d+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)+d)}{2}-\frac{d^2 \left(\operatorname{dilog}\left(-\frac{-d \tanh (b x+a)+d}{d}\right)\right)}{2}$
default	$-\frac{\operatorname{arctanh}(-1+d+d \tanh (b x+a)) d \ln (d+d \tanh (b x+a))}{2}-\frac{\operatorname{arctanh}(-1+d+d \tanh (b x+a)) d \ln (-d \tanh (b x+a)+d)}{2}-\frac{d^2 \left(\operatorname{dilog}\left(-\frac{-d \tanh (b x+a)+d}{d}\right)\right)}{2}$
risch	Expression too large to display

input

```
int(-arctanh(-1+d*d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/b/d*(1/2*arctanh(-1+d*d*tanh(b*x+a))*d*ln(d+d*tanh(b*x+a))-1/2*arctanh(-1+d*d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)+d)-1/2*d^2*(1/d*(-1/2*dilog(-1/2*(-d*tanh(b*x+a)-d)/d)-1/2*ln(-d*tanh(b*x+a)+d)*ln(-1/2*(-d*tanh(b*x+a)-d)/d)+1/2*dilog((-d*tanh(b*x+a)-d+2)/(-2*d+2))+1/2*ln(-d*tanh(b*x+a)+d)*ln((-d*tanh(b*x+a)-d+2)/(-2*d+2)))-1/d*(-1/4*ln(d+d*tanh(b*x+a))^2+1/2*(ln(d+d*tanh(b*x+a))-ln(1/2*d*tanh(b*x+a)+1/2*d))*ln(-1/2*d*tanh(b*x+a)-1/2*d+1)-1/2*dilog(1/2*d*tanh(b*x+a)+1/2*d)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(61) = 122.

Time = 0.09 (sec) , antiderivative size = 228, normalized size of antiderivative = 3.00

$$\int \operatorname{arctanh}(1-d-d \tanh (a+b x)) d x$$

$$= \frac{b^2 x^2 - b x \log \left( -\frac{d \cosh (b x+a)+d \sinh (b x+a)}{(d-2) \cosh (b x+a)+d \sinh (b x+a)} \right) + a \log (2(d-1) \cosh (b x+a)+2(d-1) \sinh (b x+a)+2)}{2}$$

input

```
integrate(-arctanh(-1+d*d*tanh(b*x+a)),x, algorithm="fricas")
```

output

```
1/2*(b^2*x^2 - b*x*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/((d - 2)*cosh(
b*x + a) + d*sinh(b*x + a))) + a*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*s
inh(b*x + a) + 2*sqrt(d - 1)) + a*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*
sinh(b*x + a) - 2*sqrt(d - 1)) - (b*x + a)*log(sqrt(d - 1)*(cosh(b*x + a)
+ sinh(b*x + a)) + 1) - (b*x + a)*log(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b
*x + a)) + 1) - dilog(sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))) - dilog
(-sqrt(d - 1)*(cosh(b*x + a) + sinh(b*x + a))))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = - \int \operatorname{atanh}(d \tanh(a + bx) + d - 1) dx$$

input

```
integrate(-atanh(-1+d+d*tanh(b*x+a)),x)
```

output

```
-Integral(atanh(d*tanh(a + b*x) + d - 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.59 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx \\ &= \frac{1}{4} bd \left( \frac{2x^2}{d} - \frac{2bx \log(-(d-1)e^{(2bx+2a)} + 1) + \operatorname{Li}_2((d-1)e^{(2bx+2a)})}{b^2 d} \right) \\ & \quad - x \operatorname{artanh}(d \tanh(bx + a) + d - 1) \end{aligned}$$

input

```
integrate(-arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="maxima")
```

output

```
1/4*b*d*(2*x^2/d - (2*b*x*log(-(d - 1)*e^(2*b*x + 2*a) + 1) + dilog((d - 1)
)*e^(2*b*x + 2*a)))/(b^2*d) - x*arctanh(d*tanh(b*x + a) + d - 1)
```

**Giac [F]**

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -\operatorname{artanh}(d \tanh(bx + a) + d - 1) dx$$

input `integrate(-arctanh(-1+d+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(-arctanh(d*tanh(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = \int -\operatorname{atanh}(d + d \tanh(a + bx) - 1) dx$$

input `int(-atanh(d + d*tanh(a + b*x) - 1),x)`

output `int(-atanh(d + d*tanh(a + b*x) - 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 - d - d \tanh(a + bx)) dx = -\left( \int \operatorname{atanh}(\tanh(bx + a) d + d - 1) dx \right)$$

input `int(-atanh(-1+d+d*tanh(b*x+a)),x)`

output `- int(atanh(tanh(a + b*x)*d + d - 1),x)`

### 3.297 $\int \frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x} dx$

Optimal result	2059
Mathematica [N/A]	2059
Rubi [N/A]	2060
Maple [N/A]	2060
Fricas [N/A]	2061
Sympy [N/A]	2061
Maxima [N/A]	2061
Giac [N/A]	2062
Mupad [N/A]	2062
Reduce [N/A]	2063

#### Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x}, x\right)$$

output `Defer(Int)(-arctanh(-1+d+d*tanh(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 4.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1-d-d \tanh(a+bx))}{x} dx$$

input `Integrate[ArcTanh[1 - d - d*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 - d - d*Tanh[a + b*x]]/x, x]`



**Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d(-\tanh(a+bx)) - d + 1)}{x} dx$$

input `Int[ArcTanh[1 - d - d*Tanh[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int -\frac{\operatorname{arctanh}(-1 + d + d \tanh(bx + a))}{x} dx$$

input `int(-arctanh(-1+d+d*tanh(b*x+a))/x,x)`

output `int(-arctanh(-1+d+d*tanh(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \tanh(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(-arctanh(d*tanh(b*x + a) + d - 1)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.63 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = - \int \frac{\operatorname{atanh}(d \tanh(a + bx) + d - 1)}{x} dx$$

input `integrate(-atanh(-1+d+d*tanh(b*x+a))/x,x)`

output `-Integral(atanh(d*tanh(a + b*x) + d - 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.72 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \tanh(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*tanh(b*x+a))/x,x, algorithm="maxima")`

output `-integrate(arctanh(d*tanh(b*x + a) + d - 1)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = \int -\frac{\operatorname{arctanh}(d \tanh(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*tanh(b*x+a))/x,x, algorithm="giac")`

output `integrate(-arctanh(d*tanh(b*x + a) + d - 1)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.35 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = \int -\frac{\operatorname{atanh}(d + d \tanh(a + bx) - 1)}{x} dx$$

input `int(-atanh(d + d*tanh(a + b*x) - 1)/x,x)`

output `int(-atanh(d + d*tanh(a + b*x) - 1)/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - d - d \tanh(a + bx))}{x} dx = - \left( \int \frac{\operatorname{atanh}(\tanh(bx + a) d + d - 1)}{x} dx \right)$$

input `int(-atanh(-1+d+d*tanh(b*x+a))/x,x)`output `- int(atanh(tanh(a + b*x)*d + d - 1)/x,x)`

### 3.298 $\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx$

Optimal result	2064
Mathematica [A] (verified)	2065
Rubi [A] (verified)	2065
Maple [C] (warning: unable to verify)	2070
Fricas [B] (verification not implemented)	2070
Sympy [F]	2071
Maxima [A] (verification not implemented)	2072
Giac [F]	2072
Mupad [F(-1)]	2073
Reduce [F]	2073

#### Optimal result

Integrand size = 15, antiderivative size = 303

$$\begin{aligned}
 \int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx = & \frac{1}{3} x^3 \operatorname{arctanh}(c + d \coth(a + bx)) \\
 & + \frac{1}{6} x^3 \log \left( 1 - \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right) \\
 & - \frac{1}{6} x^3 \log \left( 1 - \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right) \\
 & + \frac{x^2 \operatorname{PolyLog} \left( 2, \frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{4b} \\
 & - \frac{x^2 \operatorname{PolyLog} \left( 2, \frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{4b} \\
 & - \frac{x \operatorname{PolyLog} \left( 3, \frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{4b^2} \\
 & + \frac{x \operatorname{PolyLog} \left( 3, \frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{4b^2} \\
 & + \frac{\operatorname{PolyLog} \left( 4, \frac{(1-c-d)e^{2a+2bx}}{1-c+d} \right)}{8b^3} \\
 & - \frac{\operatorname{PolyLog} \left( 4, \frac{(1+c+d)e^{2a+2bx}}{1+c-d} \right)}{8b^3}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{3}x^3 \operatorname{arctanh}(c+d \operatorname{coth}(bx+a)) + \frac{1}{6}x^3 \ln(1-(1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & - \frac{1}{6}x^3 \ln(1-(1+c+d)\exp(2bx+2a)/(1+c-d)) + \frac{1}{4}x^2 \operatorname{polylog}(2, (1-c-d)\exp(2bx+2a)/(1-c+d)) \\ & /b - \frac{1}{4}x^2 \operatorname{polylog}(2, (1+c+d)\exp(2bx+2a)/(1+c-d)) /b - \frac{1}{4}x \operatorname{polylog}(3, (1-c-d)\exp(2bx+2a)/(1-c+d)) /b^2 \\ & + \frac{1}{4}x \operatorname{polylog}(3, (1+c+d)\exp(2bx+2a)/(1+c-d)) /b^2 + \frac{1}{8} \operatorname{polylog}(4, (1-c-d)\exp(2bx+2a)/(1-c+d)) /b^3 \\ & - \frac{1}{8} \operatorname{polylog}(4, (1+c+d)\exp(2bx+2a)/(1+c-d)) /b^3 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.87

$$\int x^2 \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \frac{1}{3}x^3 \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) - 4b^3 x^3 \log\left(1 + \frac{(1-c+d)e^{-2(a+bx)}}{-1+c+d}\right) + 4b^3 x^3 \log\left(1 + \frac{(-1-c+d)e^{-2(a+bx)}}{1+c+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{(-1+c-d)e^{-2(a+bx)}}{-1+c+d}\right) - 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{(-1-c+d)e^{-2(a+bx)}}{1+c+d}\right)$$

input

Integrate[x^2\*ArcTanh[c + d\*Coth[a + b\*x]],x]

output

$$\begin{aligned} & \frac{(x^3 \operatorname{ArcTanh}[c + d \operatorname{Coth}[a + b x]])}{3} - \frac{(-4 b^3 x^3 \operatorname{Log}[1 + (1 - c + d)/((-1 + c + d) E^{2(a + b x)})]}{3} + \frac{4 b^3 x^3 \operatorname{Log}[1 + (-1 - c + d)/((1 + c + d) E^{2(a + b x)})]}{3} \\ & + \frac{6 b^2 x^2 \operatorname{PolyLog}[2, (-1 + c - d)/((-1 + c + d) E^{2(a + b x)})]}{3} - \frac{6 b^2 x^2 \operatorname{PolyLog}[2, (1 + c - d)/((1 + c + d) E^{2(a + b x)})]}{3} \\ & + \frac{6 b x \operatorname{PolyLog}[3, (-1 + c - d)/((-1 + c + d) E^{2(a + b x)})]}{3} - \frac{6 b x \operatorname{PolyLog}[3, (1 + c - d)/((1 + c + d) E^{2(a + b x)})]}{3} \\ & + \frac{3 \operatorname{PolyLog}[4, (-1 + c - d)/((-1 + c + d) E^{2(a + b x)})]}{3} - \frac{3 \operatorname{PolyLog}[4, (1 + c - d)/((1 + c + d) E^{2(a + b x)})]}{3} \end{aligned}$$
**Rubi [A] (verified)**Time = 1.42 (sec) , antiderivative size = 391, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6799, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x^2 \operatorname{arctanh}(d \coth(a + bx) + c) dx \\
& \quad \downarrow \text{6799} \\
& -\frac{1}{3}b(-c-d+1) \int \frac{e^{2a+2bx} x^3}{-c - (-c-d+1)e^{2a+2bx} + d+1} dx + \frac{1}{3}b(c+d+1) \\
& 1) \int \frac{e^{2a+2bx} x^3}{c - (c+d+1)e^{2a+2bx} - d+1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& -\frac{1}{3}b(-c-d+1) \left( \frac{3 \int x^2 \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) dx}{2b(-c-d+1)} - \frac{x^3 \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + \\
& \frac{1}{3}b(c+d+1) \left( \frac{3 \int x^2 \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) dx}{2b(c+d+1)} - \frac{x^3 \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + \\
& \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + c) \\
& \quad \downarrow \text{3011} \\
& 1) \left( \frac{-\frac{1}{3}b(-c-d+1) \left( \frac{\int x \operatorname{PolyLog} \left( 2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left( 2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b} \right)}{2b(-c-d+1)} - \frac{x^3 \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + \\
& \frac{1}{3}b(c+d+1) \left( \frac{\int x \operatorname{PolyLog} \left( 2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left( 2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b} \right) - \frac{x^3 \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + \\
& \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + c) \\
& \quad \downarrow \text{7163}
\end{aligned}$$

$$1) \left( \frac{\frac{1}{3}b(-c-d+)}{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\int \operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right)}{2b(-c-d+1)} - \frac{x^3 \log\left(1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b(-c-d+1)} \right)$$

$$1) \left( \frac{\frac{1}{3}b(c+d+)}{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\int \operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) dx}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)}{2b(c+d+1)} - \frac{x^3 \log\left(1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b(c+d+1)} \right)$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + c)$$

↓ 2720

$$1) \left( \frac{\frac{1}{3}b(-c-d+)}{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right)}{2b(-c-d+1)} - \frac{x^3 \log\left(1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b(-c-d+1)} \right)$$

$$1) \left( \frac{\frac{1}{3}b(c+d+)}{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)}{2b(c+d+1)} - \frac{x^3 \log\left(1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b(c+d+1)} \right)$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + c)$$

↓ 7143



$$\begin{aligned}
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + c) - \frac{1}{3}b(-c - d + \\
 1) & \left( \frac{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{4b^2} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} \right)}{b} \right) - \frac{x^3 \log\left(1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b(-c-d+1)} \\
 & \frac{1}{3}b(c + d + \\
 1) & \left( \frac{3 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{4b^2} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} \right)}{b} \right) - \frac{x^3 \log\left(1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b(c+d+1)}
 \end{aligned}$$

input `Int[x^2*ArcTanh[c + d*Coth[a + b*x]],x]`

output `(x^3*ArcTanh[c + d*Coth[a + b*x]])/3 - (b*(1 - c - d)*(-1/2*(x^3*Log[1 - ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d]])/(b*(1 - c - d)) + (3*(-1/2*(x^2*PolyLog[2, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d]])/b + ((x*PolyLog[3, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d]])/(2*b) - PolyLog[4, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)]/(4*b^2))/b)/(2*b*(1 - c - d)))/3 + (b*(1 + c + d)*(-1/2*(x^3*Log[1 - ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d]])/(b*(1 + c + d)) + (3*(-1/2*(x^2*PolyLog[2, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d]])/b + ((x*PolyLog[3, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d]])/(2*b) - PolyLog[4, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)]/(4*b^2))/b)/(2*b*(1 + c + d)))/3`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6799

```
Int[ArcTanh[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_))^(m
_), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(
m + 1))), x] + (-Simp[b*((1 - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(
E^(2*a + 2*b*x)/(1 - c + d - (1 - c - d)*E^(2*a + 2*b*x))), x], x] + Simp[b
*((1 + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x)/(1 + c
- d - (1 + c + d)*E^(2*a + 2*b*x))), x], x]) /; FreeQ[{a, b, c, d, e, f}, x
] && IGtQ[m, 0] && NeQ[(c - d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 15.18 (sec) , antiderivative size = 5248, normalized size of antiderivative = 17.32

method	result	size
risch	Expression too large to display	5248

input

```
int(x^2*arctanh(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 880 vs. 2(259) = 518.

Time = 0.12 (sec) , antiderivative size = 880, normalized size of antiderivative = 2.90

$$\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctanh(c+d*coth(b*x+a)),x, algorithm="fricas")
```

output

```

1/6*(b^3*x^3*log(-(d*cosh(b*x + a) + (c + 1)*sinh(b*x + a))/(d*cosh(b*x +
a) + (c - 1)*sinh(b*x + a))) - 3*b^2*x^2*dilog(sqrt((c + d + 1)/(c - d + 1
))*(cosh(b*x + a) + sinh(b*x + a))) - 3*b^2*x^2*dilog(-sqrt((c + d + 1)/(c
- d + 1))*(cosh(b*x + a) + sinh(b*x + a))) + 3*b^2*x^2*dilog(sqrt((c + d
- 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + 3*b^2*x^2*dilog(-sqrt
((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + a^3*log(2*(c
+ d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(b*x + a) + 2*(c - d + 1)*sqrt(
(c + d + 1)/(c - d + 1))) + a^3*log(2*(c + d + 1)*cosh(b*x + a) + 2*(c + d
+ 1)*sinh(b*x + a) - 2*(c - d + 1)*sqrt((c + d + 1)/(c - d + 1))) - a^3*l
og(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) + 2*(c - d -
1)*sqrt((c + d - 1)/(c - d - 1))) - a^3*log(2*(c + d - 1)*cosh(b*x + a) +
2*(c + d - 1)*sinh(b*x + a) - 2*(c - d - 1)*sqrt((c + d - 1)/(c - d - 1)))
+ 6*b*x*polylog(3, sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x
+ a))) + 6*b*x*polylog(3, -sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a)
+ sinh(b*x + a))) - 6*b*x*polylog(3, sqrt((c + d - 1)/(c - d - 1))*(cosh(b
*x + a) + sinh(b*x + a))) - 6*b*x*polylog(3, -sqrt((c + d - 1)/(c - d - 1
))*(cosh(b*x + a) + sinh(b*x + a))) - (b^3*x^3 + a^3)*log(sqrt((c + d + 1)/
(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) - (b^3*x^3 + a^3)*log(-s
qrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b^3*x
^3 + a^3)*log(sqrt((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x +...

```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \coth(a + bx)) dx$$

input

```
integrate(x**2*atanh(c+d*coth(b*x+a)),x)
```

output

```
Integral(x**2*atanh(c + d*coth(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.91

$$\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx = \frac{1}{3} x^3 \operatorname{artanh}(d \coth(bx + a) + c) - \frac{1}{18} bd \left( \frac{4 b^3 x^3 \log\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + 6 b^2 x^2 \operatorname{Li}_2\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) - 6 bx \operatorname{Li}_3\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) + 3 \operatorname{Li}_4\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right)}{b^4 d} \right)$$

input `integrate(x^2*arctanh(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(d*coth(b*x + a) + c) - 1/18*b*d*((4*b^3*x^3*log(-(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1) + 1) + 6*b^2*x^2*dilog((c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) - 6*b*x*polylog(3, (c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) + 3*polylog(4, (c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^4*d) - (4*b^3*x^3*log(-(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1) + 1) + 6*b^2*x^2*dilog((c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) - 6*b*x*polylog(3, (c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) + 3*polylog(4, (c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)))/(b^4*d))`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int x^2 \operatorname{artanh}(d \coth(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(d*coth(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \operatorname{coth}(a + bx)) dx$$

input `int(x^2*atanh(c + d*coth(a + b*x)),x)`output `int(x^2*atanh(c + d*coth(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a)d + c) x^2 dx$$

input `int(x^2*atanh(c+d*coth(b*x+a)),x)`output `int(atanh(coth(a + b*x)*d + c)*x**2,x)`

### 3.299 $\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx$

Optimal result	2074
Mathematica [A] (verified)	2075
Rubi [A] (verified)	2075
Maple [C] (warning: unable to verify)	2079
Fricas [B] (verification not implemented)	2080
Sympy [F]	2081
Maxima [A] (verification not implemented)	2081
Giac [F]	2082
Mupad [F(-1)]	2082
Reduce [F]	2082

#### Optimal result

Integrand size = 13, antiderivative size = 229

$$\begin{aligned}
 \int x \operatorname{arctanh}(c + d \coth(a + bx)) dx &= \frac{1}{2} x^2 \operatorname{arctanh}(c + d \coth(a + bx)) \\
 &+ \frac{1}{4} x^2 \log \left( 1 - \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right) \\
 &- \frac{1}{4} x^2 \log \left( 1 - \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right) \\
 &+ \frac{x \operatorname{PolyLog} \left( 2, \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right)}{4b} \\
 &- \frac{x \operatorname{PolyLog} \left( 2, \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right)}{4b} \\
 &- \frac{\operatorname{PolyLog} \left( 3, \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d} \right)}{8b^2} \\
 &+ \frac{\operatorname{PolyLog} \left( 3, \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d} \right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\operatorname{arctanh}(c+d*\operatorname{coth}(b*x+a))+1/4*x^2*\ln(1-(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))-1/4*x^2*\ln(1-(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))+1/4*x*\operatorname{polylog}(2,(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))/b-1/4*x*\operatorname{polylog}(2,(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))/b \\ & -1/8*\operatorname{polylog}(3,(1-c-d)*\exp(2*b*x+2*a)/(1-c+d))/b^2+1/8*\operatorname{polylog}(3,(1+c+d)*\exp(2*b*x+2*a)/(1+c+d))/b^2 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.87

$$\int x \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx$$

$$= \frac{4b^2x^2 \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) + 2b^2x^2 \log\left(1 + \frac{(1-c+d)e^{-2(a+bx)}}{-1+c+d}\right) - 2b^2x^2 \log\left(1 + \frac{(-1-c+d)e^{-2(a+bx)}}{1+c+d}\right)}{1}$$

input

`Integrate[x*ArcTanh[c + d*Coth[a + b*x]],x]`

output

$$\begin{aligned} & (4*b^2*x^2*ArcTanh[c + d*Coth[a + b*x]] + 2*b^2*x^2*Log[1 + (1 - c + d)/((-1 + c + d)*E^(2*(a + b*x)))] - 2*b^2*x^2*Log[1 + (-1 - c + d)/((1 + c + d)*E^(2*(a + b*x)))] - 2*b*x*PolyLog[2, (-1 + c - d)/((-1 + c + d)*E^(2*(a + b*x)))] + 2*b*x*PolyLog[2, (1 + c - d)/((1 + c + d)*E^(2*(a + b*x)))] - PolyLog[3, (-1 + c - d)/((-1 + c + d)*E^(2*(a + b*x)))] + PolyLog[3, (1 + c - d)/((1 + c + d)*E^(2*(a + b*x)))])/(8*b^2) \end{aligned}$$
**Rubi [A] (verified)**Time = 1.05 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.31, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {6799, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \operatorname{coth}(a + bx) + c) dx$$



$$\begin{aligned}
& \downarrow \text{6799} \\
& -\frac{1}{2}b(-c-d+1) \int \frac{e^{2a+2bx}x^2}{-c-(-c-d+1)e^{2a+2bx}+d+1} dx + \frac{1}{2}b(c+d+1) \\
& 1) \int \frac{e^{2a+2bx}x^2}{c-(c+d+1)e^{2a+2bx}-d+1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + c) \\
& \downarrow \text{2620} \\
& -\frac{1}{2}b(-c-d+1) \left( \frac{\int x \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) dx}{b(-c-d+1)} - \frac{x^2 \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + \frac{1}{2}b(c+d+1) \\
& \left( \frac{\int x \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) dx}{b(c+d+1)} - \frac{x^2 \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + \\
& \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + c) \\
& \downarrow \text{3011} \\
& -\frac{1}{2}b(-c-d+1) \\
& 1) \left( \frac{\int \operatorname{PolyLog} \left( 2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) dx}{2b} - \frac{x \operatorname{PolyLog} \left( 2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b} - \frac{x^2 \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + \\
& \frac{1}{2}b(c+d+1) \\
& 1) \left( \frac{\int \operatorname{PolyLog} \left( 2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) dx}{2b} - \frac{x \operatorname{PolyLog} \left( 2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b} - \frac{x^2 \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + \\
& \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + c) \\
& \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{2}b(-c-d+ \\
 1) & \left( \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{x^2 \log\left(1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b(-c-d+1)} \right) + \\
 & \frac{1}{2}b(c+d+ \\
 1) & \left( \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{x^2 \log\left(1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b(c+d+1)} \right) + \\
 & \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + c)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + c) - \frac{1}{2}b(-c-d+ \\
 1) & \left( \frac{\operatorname{PolyLog}\left(3, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b} - \frac{x^2 \log\left(1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1}\right)}{2b(-c-d+1)} \right) + \\
 & \frac{1}{2}b(c+d+ \\
 1) & \left( \frac{\operatorname{PolyLog}\left(3, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b} - \frac{x^2 \log\left(1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1}\right)}{2b(c+d+1)} \right)
 \end{aligned}$$

input `Int[x*ArcTanh[c + d*Coth[a + b*x]],x]`

output `(x^2*ArcTanh[c + d*Coth[a + b*x]])/2 - (b*(1 - c - d)*(-1/2*(x^2*Log[1 - ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)])/(b*(1 - c - d)) + (-1/2*(x*PolyLog[2, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)]/b + PolyLog[3, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)]/(4*b^2)))/(b*(1 - c - d)))/2 + (b*(1 + c + d)*(-1/2*(x^2*Log[1 - ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)])/(b*(1 + c + d)) + (-1/2*(x*PolyLog[2, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)]/b + PolyLog[3, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)]/(4*b^2)))/(b*(1 + c + d)))/2`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6799

```
Int[ArcTanh[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_))^(m
_), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(
m + 1))), x] + (-Simp[b*((1 - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(
E^(2*a + 2*b*x)/(1 - c + d - (1 - c - d)*E^(2*a + 2*b*x))), x], x] + Simp[b
*((1 + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x)/(1 + c
- d - (1 + c + d)*E^(2*a + 2*b*x))), x], x]) /; FreeQ[{a, b, c, d, e, f}, x
] && IGtQ[m, 0] && NeQ[(c - d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.16 (sec) , antiderivative size = 4944, normalized size of antiderivative = 21.59

method	result	size
risch	Expression too large to display	4944

input `int(x*arctanh(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/4*x^2*ln((exp(2*b*x+2*a)-1)*c+(exp(2*b*x+2*a)+1)*d+exp(2*b*x+2*a)-1)-1/4
*x^2*ln((exp(2*b*x+2*a)-1)*c+(exp(2*b*x+2*a)+1)*d-exp(2*b*x+2*a)+1)+1/2/b^
2*c*a^2/(1+c+d)*ln((-c*exp(b*x+a)-exp(b*x+a)*d+((1+c-d)*(1+c+d))^(1/2)-exp
(b*x+a))/((1+c-d)*(1+c+d))^(1/2))+1/2/b^2*c*a^2/(1+c+d)*ln((c*exp(b*x+a)+e
xp(b*x+a)*d+((1+c-d)*(1+c+d))^(1/2)+exp(b*x+a))/((1+c-d)*(1+c+d))^(1/2))+1
/2/b^2*d*a^2/(1+c+d)*ln((-c*exp(b*x+a)-exp(b*x+a)*d+((1+c-d)*(1+c+d))^(1/2)
)-exp(b*x+a))/((1+c-d)*(1+c+d))^(1/2))+1/8/b^2*d/(1+c+d)*polylog(3,(1+c+d)
*exp(2*b*x+2*a)/(1+c-d))+1/2/b^2*a/(1+c+d)*dilog((-c*exp(b*x+a)-exp(b*x+a)
*d+((1+c-d)*(1+c+d))^(1/2)-exp(b*x+a))/((1+c-d)*(1+c+d))^(1/2))-1/4/b^2/(1
+c+d)*ln(1-(1+c+d)*exp(2*b*x+2*a)/(1+c-d))*a^2-1/4/b/(1+c+d)*polylog(2,(1+
c+d)*exp(2*b*x+2*a)/(1+c-d))*x-1/4/b^2*a^2/(1+c+d)*ln(c*exp(2*b*x+2*a)+d*e
xp(2*b*x+2*a)+exp(2*b*x+2*a)-c+d-1)+1/2/b^2*a/(1+c+d)*dilog((c*exp(b*x+a)+
exp(b*x+a)*d+((1+c-d)*(1+c+d))^(1/2)+exp(b*x+a))/((1+c-d)*(1+c+d))^(1/2))-
1/4/b^2/(1+c+d)*polylog(2,(1+c+d)*exp(2*b*x+2*a)/(1+c-d))*a+1/8/b^2*c/(1+c
+d)*polylog(3,(1+c+d)*exp(2*b*x+2*a)/(1+c-d))-1/4*c/(1+c+d)*ln(1-(1+c+d)*e
xp(2*b*x+2*a)/(1+c-d))*x^2-1/4*d/(1+c+d)*ln(1-(1+c+d)*exp(2*b*x+2*a)/(1+c-
d))*x^2+1/2/b^2*a/(c+d-1)*dilog((-c*exp(b*x+a)-exp(b*x+a)*d+((c-d-1)*(c+d-
1))^(1/2)+exp(b*x+a))/((c-d-1)*(c+d-1))^(1/2))+1/2/b^2*a/(c+d-1)*dilog((c*
exp(b*x+a)+exp(b*x+a)*d+((c-d-1)*(c+d-1))^(1/2)-exp(b*x+a))/((c-d-1)*(c+d-
1))^(1/2))-1/8/b^2*c/(c+d-1)*polylog(3,(c+d-1)*exp(2*b*x+2*a)/(c-d-1))-...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 730 vs.  $2(195) = 390$ .

Time = 0.12 (sec) , antiderivative size = 730, normalized size of antiderivative = 3.19

$$\int x \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctanh(c+d*coth(b*x+a)),x, algorithm="fricas")`

output

```
1/4*(b^2*x^2*log(-(d*cosh(b*x + a) + (c + 1)*sinh(b*x + a))/(d*cosh(b*x +
a) + (c - 1)*sinh(b*x + a))) - 2*b*x*dilog(sqrt((c + d + 1)/(c - d + 1))*(
cosh(b*x + a) + sinh(b*x + a))) - 2*b*x*dilog(-sqrt((c + d + 1)/(c - d + 1
))*(cosh(b*x + a) + sinh(b*x + a))) + 2*b*x*dilog(sqrt((c + d - 1)/(c - d
- 1))*(cosh(b*x + a) + sinh(b*x + a))) + 2*b*x*dilog(-sqrt((c + d - 1)/(c
- d - 1))*(cosh(b*x + a) + sinh(b*x + a))) - a^2*log(2*(c + d + 1)*cosh(b*
x + a) + 2*(c + d + 1)*sinh(b*x + a) + 2*(c - d + 1)*sqrt((c + d + 1)/(c -
d + 1))) - a^2*log(2*(c + d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(b*x +
a) - 2*(c - d + 1)*sqrt((c + d + 1)/(c - d + 1))) + a^2*log(2*(c + d - 1)
*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) + 2*(c - d - 1)*sqrt((c + d -
1)/(c - d - 1))) + a^2*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d - 1)*si
nh(b*x + a) - 2*(c - d - 1)*sqrt((c + d - 1)/(c - d - 1))) - (b^2*x^2 - a^
2)*log(sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1)
- (b^2*x^2 - a^2)*log(-sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh
(b*x + a)) + 1) + (b^2*x^2 - a^2)*log(sqrt((c + d - 1)/(c - d - 1))*(cosh(
b*x + a) + sinh(b*x + a)) + 1) + (b^2*x^2 - a^2)*log(-sqrt((c + d - 1)/(c
- d - 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 2*polylog(3, sqrt((c + d
+ 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) + 2*polylog(3, -sqrt((c
+ d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 2*polylog(3, sqr
t((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) - 2*polylog...
```

**Sympy [F]**

$$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int x \operatorname{atanh}(c + d \coth(a + bx)) dx$$

input `integrate(x*atanh(c+d*coth(b*x+a)),x)`

output `Integral(x*atanh(c + d*coth(a + b*x)), x)`

**Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.93

$$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx =$$

$$-\frac{1}{8}bd \left( \frac{2b^2x^2 \log\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + 2bx \operatorname{Li}_2\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right) - \operatorname{Li}_3\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right)}{b^3d} - \frac{2b^2x^2 \log\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1} + 1\right) + 2bx \operatorname{Li}_2\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1}\right) - \operatorname{Li}_3\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1}\right)}{b^3d} \right)$$

$$+ \frac{1}{2}x^2 \operatorname{artanh}(d \coth(bx + a) + c)$$

input `integrate(x*arctanh(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `-1/8*b*d*((2*b^2*x^2*log(-(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1) + 1) + 2*b*x*dilog((c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)) - polylog(3, (c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^3*d) - (2*b^2*x^2*log(-(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1) + 1) + 2*b*x*dilog((c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)) - polylog(3, (c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)))/(b^3*d) + 1/2*x^2*arctanh(d*coth(b*x + a) + c)`

**Giac [F]**

$$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int x \operatorname{artanh}(d \coth(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*coth(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int x \operatorname{atanh}(c + d \coth(a + bx)) dx$$

input `int(x*atanh(c + d*coth(a + b*x)),x)`

output `int(x*atanh(c + d*coth(a + b*x)), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int \operatorname{atanh}(\coth(bx + a) d + c) x dx$$

input `int(x*atanh(c+d*coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x)*d + c)*x,x)`

### 3.300 $\int \operatorname{arctanh}(c + d \coth(a + bx)) dx$

Optimal result	2083
Mathematica [A] (verified)	2084
Rubi [A] (verified)	2084
Maple [B] (verified)	2086
Fricas [B] (verification not implemented)	2087
Sympy [F]	2088
Maxima [A] (verification not implemented)	2088
Giac [F]	2089
Mupad [F(-1)]	2089
Reduce [F]	2090

#### Optimal result

Integrand size = 11, antiderivative size = 150

$$\int \operatorname{arctanh}(c + d \coth(a + bx)) dx = x \operatorname{arctanh}(c + d \coth(a + bx)) + \frac{1}{2}x \log\left(1 - \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d}\right) - \frac{1}{2}x \log\left(1 - \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d}\right) + \frac{\operatorname{PolyLog}\left(2, \frac{(1 - c - d)e^{2a+2bx}}{1 - c + d}\right)}{4b} - \frac{\operatorname{PolyLog}\left(2, \frac{(1 + c + d)e^{2a+2bx}}{1 + c - d}\right)}{4b}$$

output

```
x*arctanh(c+d*coth(b*x+a))+1/2*x*ln(1-(1-c-d)*exp(2*b*x+2*a)/(1-c+d))-1/2*x*ln(1-(1+c+d)*exp(2*b*x+2*a)/(1+c-d))+1/4*polylog(2,(1-c-d)*exp(2*b*x+2*a)/(1-c+d))/b-1/4*polylog(2,(1+c+d)*exp(2*b*x+2*a)/(1+c-d))/b
```



**Mathematica [A] (verified)**

Time = 3.11 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.87

$$\int \operatorname{arctanh}(c + d \coth(a + bx)) dx = x \operatorname{arctanh}(c + d \coth(a + bx)) - \frac{-2bx \left( \log \left( 1 - \frac{(-1+c+d)e^{2(a+bx)}}{-1+c-d} \right) - \log \left( 1 - \frac{(1+c+d)e^{2(a+bx)}}{1+c-d} \right) \right) - \operatorname{PolyLog} \left( 2, \frac{(-1+c+d)e^{2(a+bx)}}{-1+c-d} \right) + \operatorname{PolyLog} \left( 2, \frac{(1+c+d)e^{2(a+bx)}}{1+c-d} \right)}{4b}$$

input `Integrate[ArcTanh[c + d*Coth[a + b*x]], x]`

output

```
x*ArcTanh[c + d*Coth[a + b*x]] - (-2*b*x*(Log[1 - ((-1 + c + d)*E^(2*(a + b*x)))/(-1 + c - d)] - Log[1 - ((1 + c + d)*E^(2*(a + b*x)))/(1 + c - d)]) - PolyLog[2, ((-1 + c + d)*E^(2*(a + b*x)))/(-1 + c - d)] + PolyLog[2, ((1 + c + d)*E^(2*(a + b*x)))/(1 + c - d)]/(4*b)
```

**Rubi [A] (verified)**Time = 0.65 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.38, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6791, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \coth(a + bx) + c) dx$$

$$\downarrow 6791$$

$$-b(-c - d + 1) \int \frac{e^{2a+2bx} x}{-c - (-c - d + 1)e^{2a+2bx} + d + 1} dx + b(c + d + 1) \int \frac{e^{2a+2bx} x}{c - (c + d + 1)e^{2a+2bx} - d + 1} dx + x \operatorname{arctanh}(d \coth(a + bx) + c)$$

$$\downarrow 2620$$

$$-b(-c-d+1) \left( \frac{\int \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) dx}{2b(-c-d+1)} - \frac{x \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + b(c+d+1) \\ 1) \left( \frac{\int \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) dx}{2b(c+d+1)} - \frac{x \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + x \operatorname{arctanh}(d \coth(a+bx) + c)$$

↓ 2715

$$1) \left( \frac{\int e^{-2a-2bx} \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right) de^{2a+2bx}}{4b^2(-c-d+1)} - \frac{x \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + b(c+d+1) \\ 1) \left( \frac{\int e^{-2a-2bx} \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right) de^{2a+2bx}}{4b^2(c+d+1)} - \frac{x \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right) + \\ x \operatorname{arctanh}(d \coth(a+bx) + c)$$

↓ 2838

$$1) \left( -\frac{\operatorname{PolyLog} \left( 2, \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{4b^2(-c-d+1)} - \frac{x \log \left( 1 - \frac{(-c-d+1)e^{2a+2bx}}{-c+d+1} \right)}{2b(-c-d+1)} \right) + b(c+d+1) \\ 1) \left( -\frac{\operatorname{PolyLog} \left( 2, \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{4b^2(c+d+1)} - \frac{x \log \left( 1 - \frac{(c+d+1)e^{2a+2bx}}{c-d+1} \right)}{2b(c+d+1)} \right)$$

input `Int[ArcTanh[c + d*Coth[a + b*x]],x]`

output `x*ArcTanh[c + d*Coth[a + b*x]] - b*(1 - c - d)*(-1/2*(x*Log[1 - ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)])/(b*(1 - c - d)) - PolyLog[2, ((1 - c - d)*E^(2*a + 2*b*x))/(1 - c + d)]/(4*b^2*(1 - c - d)) + b*(1 + c + d)*(-1/2*(x*Log[1 - ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)])/(b*(1 + c + d)) - PolyLog[2, ((1 + c + d)*E^(2*a + 2*b*x))/(1 + c - d)]/(4*b^2*(1 + c + d))`

Defintions of rubi rules used

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 6791 Int[ArcTanh[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)], x_Symbol] := Simp[x*Ar
cTanh[c + d*Coth[a + b*x]], x] + (-Simp[b*(1 - c - d) Int[x*(E^(2*a + 2*b
*x))/(1 - c + d - (1 - c - d)*E^(2*a + 2*b*x))], x], x] + Simp[b*(1 + c + d)
Int[x*(E^(2*a + 2*b*x))/(1 + c - d - (1 + c + d)*E^(2*a + 2*b*x))], x], x
]) /; FreeQ[{a, b, c, d}, x] && NeQ[(c - d)^2, 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 347 vs. 2(138) = 276.

Time = 3.96 (sec) , antiderivative size = 348, normalized size of antiderivative = 2.32

method	result
derivativedivides	$\frac{-\frac{\operatorname{arctanh}(c+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} + \frac{\operatorname{arctanh}(c+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)-d)}{2} + d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \operatorname{coth}(bx+a)}{1-c+d}\right)}{2} \right)}{\dots}$
default	$\frac{-\frac{\operatorname{arctanh}(c+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} + \frac{\operatorname{arctanh}(c+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)-d)}{2} + d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \operatorname{coth}(bx+a)}{1-c+d}\right)}{2} \right)}{\dots}$
risch	Expression too large to display

input `int(arctanh(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output `1/b/d*(-1/2*arctanh(c+d*coth(b*x+a))*d*ln(-d*coth(b*x+a)+d)+1/2*arctanh(c+d*coth(b*x+a))*d*ln(-d*coth(b*x+a)-d)+1/2*d^2*(1/d*(1/2*dilog((-d*coth(b*x+a)-c+1)/(1-c+d))+1/2*ln(-d*coth(b*x+a)-d)*ln((-d*coth(b*x+a)-c+1)/(1-c+d))-1/2*dilog((-d*coth(b*x+a)-c-1)/(-1-c+d))-1/2*ln(-d*coth(b*x+a)-d)*ln((-d*coth(b*x+a)-c-1)/(-1-c+d)))-1/d*(1/2*dilog((-d*coth(b*x+a)-c+1)/(1-c-d))+1/2*ln(-d*coth(b*x+a)+d)*ln((-d*coth(b*x+a)-c+1)/(1-c-d))-1/2*dilog((-d*coth(b*x+a)-c-1)/(-1-c-d))-1/2*ln(-d*coth(b*x+a)+d)*ln((-d*coth(b*x+a)-c-1)/(-1-c-d))))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 540 vs.  $2(128) = 256$ .

Time = 0.12 (sec) , antiderivative size = 540, normalized size of antiderivative = 3.60

$$\int \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctanh(c+d*coth(b*x+a)),x, algorithm="fricas")`

output

```

1/2*(b*x*log(-(d*cosh(b*x + a) + (c + 1)*sinh(b*x + a))/(d*cosh(b*x + a) +
(c - 1)*sinh(b*x + a))) + a*log(2*(c + d + 1)*cosh(b*x + a) + 2*(c + d +
1)*sinh(b*x + a) + 2*(c - d + 1)*sqrt((c + d + 1)/(c - d + 1))) + a*log(2*
(c + d + 1)*cosh(b*x + a) + 2*(c + d + 1)*sinh(b*x + a) - 2*(c - d + 1)*sq
rt((c + d + 1)/(c - d + 1))) - a*log(2*(c + d - 1)*cosh(b*x + a) + 2*(c +
d - 1)*sinh(b*x + a) + 2*(c - d - 1)*sqrt((c + d - 1)/(c - d - 1))) - a*lo
g(2*(c + d - 1)*cosh(b*x + a) + 2*(c + d - 1)*sinh(b*x + a) - 2*(c - d - 1
)*sqrt((c + d - 1)/(c - d - 1))) - (b*x + a)*log(sqrt((c + d + 1)/(c - d +
1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) - (b*x + a)*log(-sqrt((c + d + 1
)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b*x + a)*log(sqrt((
c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (b*x + a)*l
og(-sqrt((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) - d
ilog(sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) - dilo
g(-sqrt((c + d + 1)/(c - d + 1))*(cosh(b*x + a) + sinh(b*x + a))) + dilog(
sqrt((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))) + dilog(-sq
rt((c + d - 1)/(c - d - 1))*(cosh(b*x + a) + sinh(b*x + a))))/b

```

**Sympy [F]**

$$\int \operatorname{arctanh}(c + d \coth(a + bx)) dx = \int \operatorname{atanh}(c + d \coth(a + bx)) dx$$

input

```
integrate(atanh(c+d*coth(b*x+a)),x)
```

output

```
Integral(atanh(c + d*coth(a + b*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.95

$$\int \operatorname{arctanh}(c + d \coth(a + bx)) dx =$$

$$-\frac{1}{4}bd \left( \frac{2bx \log\left(-\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1} + 1\right) + \operatorname{Li}_2\left(\frac{(c+d+1)e^{(2bx+2a)}}{c-d+1}\right)}{b^2d} - \frac{2bx \log\left(-\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1} + 1\right) + \operatorname{Li}_2\left(\frac{(c+d-1)e^{(2bx+2a)}}{c-d-1}\right)}{b^2d} \right)$$

$$+ x \operatorname{artanh}(d \coth(bx + a) + c)$$

input `integrate(arctanh(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `-1/4*b*d*((2*b*x*log(-(c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1) + 1) + dilog((c + d + 1)*e^(2*b*x + 2*a)/(c - d + 1)))/(b^2*d) - (2*b*x*log(-(c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1) + 1) + dilog((c + d - 1)*e^(2*b*x + 2*a)/(c - d - 1)))/(b^2*d)) + x*arctanh(d*coth(b*x + a) + c)`

### Giac [F]

$$\int \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \int \operatorname{artanh}(d \operatorname{coth}(bx + a) + c) dx$$

input `integrate(arctanh(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*coth(b*x + a) + c), x)`

### Mupad [F(-1)]

Timed out.

$$\int \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(c + d \operatorname{coth}(a + bx)) dx$$

input `int(atanh(c + d*coth(a + b*x)),x)`

output `int(atanh(c + d*coth(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(c + d \operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + c) dx$$

input `int(atanh(c+d*coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x)*d + c),x)`

### 3.301 $\int \frac{\operatorname{arctanh}(c+d \coth(a+bx))}{x} dx$

Optimal result	2091
Mathematica [N/A]	2091
Rubi [N/A]	2092
Maple [N/A]	2092
Fricas [N/A]	2093
Sympy [N/A]	2093
Maxima [N/A]	2093
Giac [N/A]	2094
Mupad [N/A]	2094
Reduce [N/A]	2095

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(c+d*coth(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 9.91 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx$$

input `Integrate[ArcTanh[c + d*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTanh[c + d*Coth[a + b*x]]/x, x]`



**Rubi [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \coth(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \coth(a + bx) + c)}{x} dx$$

input `Int[ArcTanh[c + d*Coth[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.35 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(c + d \coth(bx + a))}{x} dx$$

input `int(arctanh(c+d*coth(b*x+a))/x,x)`

output `int(arctanh(c+d*coth(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*coth(b*x + a) + c)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 2.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \coth(a + bx))}{x} dx$$

input `integrate(atanh(c+d*coth(b*x+a))/x,x)`

output `Integral(atanh(c + d*coth(a + b*x))/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.76 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*coth(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctanh(d*coth(b*x + a) + c)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*coth(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*coth(b*x + a) + c)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.38 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \coth(a + bx))}{x} dx$$

input `int(atanh(c + d*coth(a + b*x))/x,x)`

output `int(atanh(c + d*coth(a + b*x))/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \operatorname{coth}(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\operatorname{coth}(bx + a) d + c)}{x} dx$$

input `int(atanh(c+d*coth(b*x+a))/x,x)`output `int(atanh(coth(a + b*x)*d + c)/x,x)`

### 3.302 $\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$

Optimal result	2096
Mathematica [A] (verified)	2097
Rubi [A] (verified)	2097
Maple [C] (warning: unable to verify)	2101
Fricas [B] (verification not implemented)	2102
Sympy [F]	2102
Maxima [A] (verification not implemented)	2103
Giac [F]	2103
Mupad [F(-1)]	2103
Reduce [F]	2104

#### Optimal result

Integrand size = 16, antiderivative size = 152

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{bx^5}{20} + \frac{1}{4}x^4 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - \frac{1}{8}x^4 \log(1 - (1 + d)e^{2a+2bx}) - \frac{x^3 \operatorname{PolyLog}(2, (1 + d)e^{2a+2bx})}{4b} + \frac{3x^2 \operatorname{PolyLog}(3, (1 + d)e^{2a+2bx})}{8b^2} - \frac{3x \operatorname{PolyLog}(4, (1 + d)e^{2a+2bx})}{8b^3} + \frac{3 \operatorname{PolyLog}(5, (1 + d)e^{2a+2bx})}{16b^4}$$

output

```
1/20*b*x^5+1/4*x^4*arctanh(1+d*d*coth(b*x+a))-1/8*x^4*ln(1-(1+d)*exp(2*b*x+2*a))-1/4*x^3*polylog(2,(1+d)*exp(2*b*x+2*a))/b+3/8*x^2*polylog(3,(1+d)*exp(2*b*x+2*a))/b^2-3/8*x*polylog(4,(1+d)*exp(2*b*x+2*a))/b^3+3/16*polylog(5,(1+d)*exp(2*b*x+2*a))/b^4
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.95

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{4b^4 x^4 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - 2b^4 x^4 \log\left(1 - \frac{e^{-2(a+bx)}}{1+d}\right) + 4b^3 x^3 \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(3, \frac{e^{-2(a+bx)}}{1+d}\right) + 6b x \operatorname{PolyLog}\left(4, \frac{e^{-2(a+bx)}}{1+d}\right) + 3 \operatorname{PolyLog}\left(5, \frac{e^{-2(a+bx)}}{1+d}\right)}{16b^4}$$

input

```
Integrate[x^3*ArcTanh[1 + d + d*Coth[a + b*x]],x]
```

output

```
(4*b^4*x^4*ArcTanh[1 + d + d*Coth[a + b*x]] - 2*b^4*x^4*Log[1 - 1/((1 + d)*E^(2*(a + b*x)))] + 4*b^3*x^3*PolyLog[2, 1/((1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[3, 1/((1 + d)*E^(2*(a + b*x)))] + 6*b*x*PolyLog[4, 1/((1 + d)*E^(2*(a + b*x)))] + 3*PolyLog[5, 1/((1 + d)*E^(2*(a + b*x)))])/(16*b^4)
```

**Rubi [A] (verified)**

Time = 1.50 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.28, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6795, 2615, 2620, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(d \coth(a + bx) + d + 1) dx$$

$$\downarrow 6795$$

$$\frac{1}{4}b \int \frac{x^4}{1 - (d + 1)e^{2a+2bx}} dx + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2615$$

$$\frac{1}{4}b \left( (d + 1) \int \frac{e^{2a+2bx} x^4}{1 - (d + 1)e^{2a+2bx}} dx + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2620$$

$$\frac{1}{4}b \left( (d+1) \left( \frac{2 \int x^3 \log(1 - (d+1)e^{2a+2bx}) dx}{b(d+1)} - \frac{x^4 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{4}b \left( (d+1) \left( \frac{2 \left( \frac{3 \int x^2 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx}) dx}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{b(d+1)} - \frac{x^4 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{4}b \left( (d+1) \left( \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{\int x \operatorname{PolyLog}(3, (d+1)e^{2a+2bx}) dx}{b} \right)}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{b(d+1)} - \frac{x^4 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{4}b \left( (d+1) \left( \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, (d+1)e^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(4, (d+1)e^{2a+2bx}) dx}{2b} \right)}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{b(d+1)} - \frac{x^4 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 2720

$$\frac{1}{4}b \left( (d+1) \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, (d+1)e^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(4, (d+1)e^{2a+2bx}) de^{2a+2bx}}{b} \right)}{4b^2} \right)}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right) - \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 7143

$$\frac{1}{4}b \left( (d+1) \frac{2 \left( 3 \left( \frac{x^2 \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, (d+1)e^{2a+2bx})}{2b} - \frac{\operatorname{PolyLog}(5, (d+1)e^{2a+2bx})}{4b^2} \right)}{2b} \right) - \frac{x^3 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

input `Int[x^3*ArcTanh[1 + d + d*Coth[a + b*x]],x]`

output `(x^4*ArcTanh[1 + d + d*Coth[a + b*x]])/4 + (b*(x^5/5 + (1 + d)*(-1/2*(x^4*Log[1 - (1 + d)*E^(2*a + 2*b*x)]))/(b*(1 + d)) + (2*(-1/2*(x^3*PolyLog[2, (1 + d)*E^(2*a + 2*b*x)])/b + (3*((x^2*PolyLog[3, (1 + d)*E^(2*a + 2*b*x)])/(2*b) - ((x*PolyLog[4, (1 + d)*E^(2*a + 2*b*x)])/(2*b) - PolyLog[5, (1 + d)*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b)))/(b*(1 + d)))/4`



## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6795 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e._) + (f._)*(x._))^(m._)*PolyLog[n_, (d._)*((F_)^((c._)*((a._) + (b._)
)*(x._)))^(p._)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.67 (sec) , antiderivative size = 1693, normalized size of antiderivative = 11.14

method	result	size
risch	Expression too large to display	1693

input

```
int(x^3*arctanh(1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/16*(I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*
a)-1))^2+I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))^2*csgn(I*exp(2*b*x
+2*a))-2*I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2+2*I*Pi+2*ln(d)
+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2*csgn(I*d)-I*Pi*csgn(I/
(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))*csgn(I*exp(2
*b*x+2*a))-I*Pi*csgn(I*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1))*csgn(I/(exp(2*
b*x+2*a)-1)*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1))^2-I*Pi*csgn(I*exp(b*x+a))
^2*csgn(I*exp(2*b*x+2*a))+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)
+exp(2*b*x+2*a)-1))^3+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^3-I
*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)
+exp(2*b*x+2*a)-1))^2+I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))*csgn(
I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2+I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*c
sgn(I*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*ex
p(2*b*x+2*a)+exp(2*b*x+2*a)-1))-I*Pi*csgn(I*exp(2*b*x+2*a))^3+2*I*Pi*csgn(
I*exp(b*x+a))*csgn(I*exp(2*b*x+2*a))^2-I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b
*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))*csgn(I*d)-I*Picsg
n(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))^3)*x^4+1/2/b^4*a^4/(1+d)*ln(1+exp(b
*x+a)*(1+d)^(1/2))+1/2/b^4*a^3/(1+d)*dilog(1-exp(b*x+a)*(1+d)^(1/2))+1/2/b
^4*a^3/(1+d)*dilog(1+exp(b*x+a)*(1+d)^(1/2))-1/8*d/(1+d)*ln(1-(1+d)*exp(2*
b*x+2*a))*x^4+3/16/b^4/(1+d)*polylog(5,(1+d)*exp(2*b*x+2*a))-1/8/(1+d)*...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 424 vs.  $2(132) = 264$ .

Time = 0.11 (sec) , antiderivative size = 424, normalized size of antiderivative = 2.79

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{2b^5x^5 + 5b^4x^4 \log\left(-\frac{d \cosh(bx+a) + (d+2) \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) - 20b^3x^3 \operatorname{Li}_2(\sqrt{d+1}(\cosh(bx+a) + \sinh(bx+a))) - \dots}{b^4}$$

input `integrate(x^3*arctanh(1+d+d*coth(b*x+a)),x, algorithm="fricas")`

output `1/40*(2*b^5*x^5 + 5*b^4*x^4*log(-(d*cosh(b*x + a) + (d + 2)*sinh(b*x + a))/(d*cosh(b*x + a) + d*sinh(b*x + a))) - 20*b^3*x^3*dilog(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 20*b^3*x^3*dilog(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 5*a^4*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) + 2*sqrt(d + 1)) - 5*a^4*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) - 2*sqrt(d + 1)) + 60*b^2*x^2*polylog(3, sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) + 60*b^2*x^2*polylog(3, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 5*(b^4*x^4 - a^4)*log(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 5*(b^4*x^4 - a^4)*log(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 120*polylog(5, sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) + 120*polylog(5, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^4`

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^3 \operatorname{atanh}(d \coth(a + bx) + d + 1) dx$$

input `integrate(x**3*atanh(1+d+d*coth(b*x+a)),x)`

output `Integral(x**3*atanh(d*coth(a + b*x) + d + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.96

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{1}{4} x^4 \operatorname{artanh}(d \coth(bx + a) + d + 1) + \frac{1}{40} \left( \frac{2x^5}{d} - \frac{5(2b^4x^4 \log(-(d+1)e^{(2bx+2a)} + 1) + 4b^3x^3 \operatorname{Li}_2((d+1)e^{(2bx+2a)}) - 6b^2x^2 \operatorname{Li}_3((d+1)e^{(2bx+2a)})}{b^5d} \right)$$

input `integrate(x^3*arctanh(1+d+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/4*x^4*arctanh(d*coth(b*x + a) + d + 1) + 1/40*(2*x^5/d - 5*(2*b^4*x^4*log(-(d + 1)*e^(2*b*x + 2*a) + 1) + 4*b^3*x^3*dilog((d + 1)*e^(2*b*x + 2*a)) - 6*b^2*x^2*polylog(3, (d + 1)*e^(2*b*x + 2*a)) + 6*b*x*polylog(4, (d + 1)*e^(2*b*x + 2*a)) - 3*polylog(5, (d + 1)*e^(2*b*x + 2*a)))/(b^5*d))*b*d`

**Giac [F]**

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^3 \operatorname{artanh}(d \coth(bx + a) + d + 1) dx$$

input `integrate(x^3*arctanh(1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x^3*arctanh(d*coth(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^3 \operatorname{atanh}(d + d \coth(a + bx) + 1) dx$$

input `int(x^3*atanh(d + d*coth(a + b*x) + 1),x)`

output `int(x^3*atanh(d + d*coth(a + b*x) + 1), x)`

### Reduce [F]

$$\int x^3 \operatorname{arctanh}(1 + d + d \operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + d + 1) x^3 dx$$

input `int(x^3*atanh(1+d+d*coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x)*d + d + 1)*x**3,x)`

### 3.303 $\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$

Optimal result	2105
Mathematica [A] (verified)	2106
Rubi [A] (verified)	2106
Maple [C] (warning: unable to verify)	2109
Fricas [B] (verification not implemented)	2110
Sympy [F]	2111
Maxima [A] (verification not implemented)	2111
Giac [F]	2112
Mupad [F(-1)]	2112
Reduce [F]	2112

#### Optimal result

Integrand size = 16, antiderivative size = 126

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - \frac{1}{6}x^3 \log(1 - (1 + d)e^{2a+2bx}) - \frac{x^2 \operatorname{PolyLog}(2, (1 + d)e^{2a+2bx})}{4b} + \frac{x \operatorname{PolyLog}(3, (1 + d)e^{2a+2bx})}{4b^2} - \frac{\operatorname{PolyLog}(4, (1 + d)e^{2a+2bx})}{8b^3}$$

output

```
1/12*b*x^4+1/3*x^3*arctanh(1+d+d*coth(b*x+a))-1/6*x^3*ln(1-(1+d)*exp(2*b*x+2*a))-1/4*x^2*polylog(2,(1+d)*exp(2*b*x+2*a))/b+1/4*x*polylog(3,(1+d)*exp(2*b*x+2*a))/b^2-1/8*polylog(4,(1+d)*exp(2*b*x+2*a))/b^3
```

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.95

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - 4b^3 x^3 \log\left(1 - \frac{e^{-2(a+bx)}}{1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{1+d}\right) + 6bx \operatorname{PolyLog}\left(3, \frac{e^{-2(a+bx)}}{1+d}\right) + 6b \operatorname{PolyLog}\left(4, \frac{e^{-2(a+bx)}}{1+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 + d + d*Coth[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTanh[1 + d + d*Coth[a + b*x]] - 4*b^3*x^3*Log[1 - 1/((1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[2, 1/((1 + d)*E^(2*(a + b*x)))] + 6*b*x*PolyLog[3, 1/((1 + d)*E^(2*(a + b*x)))] + 3*PolyLog[4, 1/((1 + d)*E^(2*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**

Time = 1.28 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.29, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$ , Rules used = {6795, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d \coth(a + bx) + d + 1) dx$$

$$\downarrow \text{6795}$$

$$\frac{1}{3}b \int \frac{x^3}{1 - (d + 1)e^{2a+2bx}} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}b \left( (d + 1) \int \frac{e^{2a+2bx} x^3}{1 - (d + 1)e^{2a+2bx}} dx + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}b \left( (d+1) \left( \frac{3 \int x^2 \log(1 - (d+1)e^{2a+2bx}) dx}{2b(d+1)} - \frac{x^3 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{3}b \left( (d+1) \left( \frac{3 \left( \frac{\int x \operatorname{PolyLog}(2, (d+1)e^{2a+2bx}) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{2b(d+1)} - \frac{x^3 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 7163

$$\frac{1}{3}b \left( (d+1) \left( \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(3, (d+1)e^{2a+2bx}) dx}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{2b(d+1)} - \frac{x^3 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + d+1) \right)$$

↓ 2720

$$\frac{1}{3}b \left( (d+1) \left( \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, (d+1)e^{2a+2bx}) de^{2a+2bx}}{b}}{4b^2} - \frac{x^2 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{2b(d+1)} - \frac{x^3 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + d+1) \right)$$

↓ 7143

$$\frac{1}{3}b \left( (d+1) \left( \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{2b} - \frac{\operatorname{PolyLog}(4, (d+1)e^{2a+2bx})}{4b^2}}{b} - \frac{x^2 \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} \right)}{2b(d+1)} - \frac{x^3 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \coth(a+bx) + d+1) \right)$$

input `Int[x^2*ArcTanh[1 + d + d*Coth[a + b*x]], x]`



output

```
(x^3*ArcTanh[1 + d + d*Coth[a + b*x]])/3 + (b*(x^4/4 + (1 + d)*(-1/2*(x^3*
Log[1 - (1 + d)*E^(2*a + 2*b*x)]))/(b*(1 + d)) + (3*(-1/2*(x^2*PolyLog[2, (
1 + d)*E^(2*a + 2*b*x)])/b + ((x*PolyLog[3, (1 + d)*E^(2*a + 2*b*x)])/(2*b
) - PolyLog[4, (1 + d)*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b*(1 + d))))/3
```

### Defintions of rubi rules used

rule 2615

```
Int[(((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x
_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[
b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x]
, x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/
((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6795

```
Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(
m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a
+ 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c
- d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.22 (sec) , antiderivative size = 1636, normalized size of antiderivative = 12.98

method	result	size
risch	Expression too large to display	1636

input

```
int(x^2*arctanh(1+d+d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/12*b*x^4-1/4/b/(1+d)*polylog(2,(1+d)*exp(2*b*x+2*a))*x^2+1/3/b^3/(1+d)*l
n(1-(1+d)*exp(2*b*x+2*a))*a^3+1/4/b^3/(1+d)*polylog(2,(1+d)*exp(2*b*x+2*a)
)*a^2-1/3*x^3*ln(exp(b*x+a))+1/4/b^2/(1+d)*polylog(3,(1+d)*exp(2*b*x+2*a))
*x-1/8/b^3*d/(1+d)*polylog(4,(1+d)*exp(2*b*x+2*a))-1/2/b^3*a^3/(1+d)*ln(1-
exp(b*x+a)*(1+d)^(1/2))-1/2/b^3*a^3/(1+d)*ln(1+exp(b*x+a)*(1+d)^(1/2))-1/2
/b^3*a^2/(1+d)*dilog(1-exp(b*x+a)*(1+d)^(1/2))-1/2/b^3*a^2/(1+d)*dilog(1+e
xp(b*x+a)*(1+d)^(1/2))+1/6/b^3*a^3/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a
)-1)-1/6*d/(1+d)*ln(1-(1+d)*exp(2*b*x+2*a))*x^3-1/6/(1+d)*ln(1-(1+d)*exp(2
*b*x+2*a))*x^3-1/8/b^3/(1+d)*polylog(4,(1+d)*exp(2*b*x+2*a))-1/2/b^2*a^2/(
1+d)*x*ln(1-exp(b*x+a)*(1+d)^(1/2))-1/2/b^2*a^2/(1+d)*x*ln(1+exp(b*x+a)*(1
+d)^(1/2))-1/2/b^3*d*a^3/(1+d)*ln(1-exp(b*x+a)*(1+d)^(1/2))-1/2/b^3*d*a^3/
(1+d)*ln(1+exp(b*x+a)*(1+d)^(1/2))-1/2/b^3*d*a^2/(1+d)*dilog(1-exp(b*x+a)*
(1+d)^(1/2))-1/2/b^3*d*a^2/(1+d)*dilog(1+exp(b*x+a)*(1+d)^(1/2))+1/6/b^3*d
*a^3/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1)+1/2/b^2/(1+d)*ln(1-(1+d)*
exp(2*b*x+2*a))*a^2*x-1/4/b*d/(1+d)*polylog(2,(1+d)*exp(2*b*x+2*a))*x^2+1/
3/b^3*d/(1+d)*ln(1-(1+d)*exp(2*b*x+2*a))*a^3+1/4/b^3*d/(1+d)*polylog(2,(1+
d)*exp(2*b*x+2*a))*a^2+1/4/b^2*d/(1+d)*polylog(3,(1+d)*exp(2*b*x+2*a))*x-1
/2/b^2*d*a^2/(1+d)*x*ln(1-exp(b*x+a)*(1+d)^(1/2))-1/2/b^2*d*a^2/(1+d)*x*ln
(1+exp(b*x+a)*(1+d)^(1/2))+1/2/b^2*d/(1+d)*ln(1-(1+d)*exp(2*b*x+2*a))*a^2*
x-1/12*(I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 360 vs.  $2(109) = 218$ .

Time = 0.11 (sec) , antiderivative size = 360, normalized size of antiderivative = 2.86

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(ax + b)) dx$$

$$= \frac{b^4 x^4 + 2 b^3 x^3 \log\left(-\frac{d \cosh(bx+a) + (d+2) \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) - 6 b^2 x^2 \operatorname{Li}_2(\sqrt{d+1}(\cosh(bx+a) + \sinh(bx+a))) - 6 b^2 x \operatorname{Li}_2(\sqrt{d+1} \sinh(bx+a)) + 6 b^2 x \operatorname{Li}_2(\sqrt{d+1} \cosh(bx+a)) + 6 b^2 \operatorname{Li}_2(\sqrt{d+1})}{1}$$

input

```
integrate(x^2*arctanh(1+d+d*coth(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(b^4*x^4 + 2*b^3*x^3*log(-(d*cosh(b*x + a) + (d + 2)*sinh(b*x + a))/(
d*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b^2*x^2*dilog(sqrt(d + 1)*(cosh(b*
x + a) + sinh(b*x + a))) - 6*b^2*x^2*dilog(-sqrt(d + 1)*(cosh(b*x + a) + s
inh(b*x + a))) + 2*a^3*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x +
a) + 2*sqrt(d + 1)) + 2*a^3*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b
*x + a) - 2*sqrt(d + 1)) + 12*b*x*polylog(3, sqrt(d + 1)*(cosh(b*x + a) +
sinh(b*x + a))) + 12*b*x*polylog(3, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x
+ a))) - 2*(b^3*x^3 + a^3)*log(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)
) + 1) - 2*(b^3*x^3 + a^3)*log(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)
) + 1) - 12*polylog(4, sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 12*p
olylog(4, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^3
```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^2 \operatorname{atanh}(d \coth(a + bx) + d + 1) dx$$

input

```
integrate(x**2*atanh(1+d+d*coth(b*x+a)),x)
```

output

```
Integral(x**2*atanh(d*coth(a + b*x) + d + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.98

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{1}{3} x^3 \operatorname{artanh}(d \coth(bx + a) + d + 1) + \frac{1}{36} \left( \frac{3x^4}{d} - \frac{2(4b^3x^3 \log(-(d+1)e^{(2bx+2a)}) + 1) + 6b^2x^2 \operatorname{Li}_2((d+1)e^{(2bx+2a)}) - 6bx \operatorname{Li}_3((d+1)e^{(2bx+2a)})}{b^4d} \right)$$

input

```
integrate(x^2*arctanh(1+d+d*coth(b*x+a)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctanh(d*coth(b*x + a) + d + 1) + 1/36*(3*x^4/d - 2*(4*b^3*x^3*log(-(d + 1)*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog((d + 1)*e^(2*b*x + 2*a)) - 6*b*x*polylog(3, (d + 1)*e^(2*b*x + 2*a)) + 3*polylog(4, (d + 1)*e^(2*b*x + 2*a)))/(b^4*d))*b*d
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^2 \operatorname{artanh}(d \coth(bx + a) + d + 1) dx$$

input

```
integrate(x^2*arctanh(1+d+d*coth(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctanh(d*coth(b*x + a) + d + 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x^2 \operatorname{atanh}(d + d \coth(a + bx) + 1) dx$$

input

```
int(x^2*atanh(d + d*coth(a + b*x) + 1),x)
```

output

```
int(x^2*atanh(d + d*coth(a + b*x) + 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int \operatorname{atanh}(\coth(bx + a) d + d + 1) x^2 dx$$

input

```
int(x^2*atanh(1+d+d*coth(b*x+a)),x)
```

output `int(atanh(coth(a + b*x)*d + d + 1)*x**2,x)`

### 3.304 $\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$

Optimal result	2114
Mathematica [A] (verified)	2115
Rubi [A] (verified)	2115
Maple [C] (warning: unable to verify)	2118
Fricas [B] (verification not implemented)	2119
Sympy [F]	2119
Maxima [A] (verification not implemented)	2120
Giac [F]	2120
Mupad [F(-1)]	2120
Reduce [F]	2121

#### Optimal result

Integrand size = 14, antiderivative size = 100

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - \frac{1}{4}x^2 \log(1 - (1 + d)e^{2a+2bx}) - \frac{x \operatorname{PolyLog}(2, (1 + d)e^{2a+2bx})}{4b} + \frac{\operatorname{PolyLog}(3, (1 + d)e^{2a+2bx})}{8b^2}$$

output

```
1/6*b*x^3+1/2*x^2*arctanh(1+d+d*coth(b*x+a))-1/4*x^2*ln(1-(1+d)*exp(2*b*x+2*a))-1/4*x*polylog(2,(1+d)*exp(2*b*x+2*a))/b+1/8*polylog(3,(1+d)*exp(2*b*x+2*a))/b^2
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.90

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left( 2 \operatorname{arctanh}(1 + d + d \coth(a + bx)) - \log \left( 1 - \frac{e^{-2(a+bx)}}{1+d} \right) \right) + 2bx \operatorname{PolyLog} \left( 2, \frac{e^{-2(a+bx)}}{1+d} \right) + \operatorname{PolyLog} \left( 3, \frac{e^{-2(a+bx)}}{1+d} \right)}{8b^2}$$

input `Integrate[x*ArcTanh[1 + d + d*Coth[a + b*x]],x]`

output `(2*b^2*x^2*(2*ArcTanh[1 + d + d*Coth[a + b*x]] - Log[1 - 1/((1 + d)*E^(2*(a + b*x))])) + 2*b*x*PolyLog[2, 1/((1 + d)*E^(2*(a + b*x))]] + PolyLog[3, 1/((1 + d)*E^(2*(a + b*x))]])/(8*b^2)`

**Rubi [A] (verified)**

Time = 0.71 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.28, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6795, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \coth(a + bx) + d + 1) dx$$

$$\downarrow \text{6795}$$

$$\frac{1}{2} b \int \frac{x^2}{1 - (d + 1)e^{2a+2bx}} dx + \frac{1}{2} x^2 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} b \left( (d + 1) \int \frac{e^{2a+2bx} x^2}{1 - (d + 1)e^{2a+2bx}} dx + \frac{x^3}{3} \right) + \frac{1}{2} x^2 \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow \text{2620}$$



$$\frac{1}{2}b \left( (d+1) \left( \frac{\int x \log(1 - (d+1)e^{2a+2bx}) dx}{b(d+1)} - \frac{x^2 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^3}{3} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 3011

$$\frac{1}{2}b \left( (d+1) \left( \frac{\frac{\int \operatorname{PolyLog}(2, (d+1)e^{2a+2bx}) dx}{2b}}{b(d+1)} - \frac{x \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} - \frac{x^2 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^3}{3} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + d+1)$$

↓ 2720

$$\frac{1}{2}b \left( (d+1) \left( \frac{\frac{\int e^{-2a-2bx} \operatorname{PolyLog}(2, (d+1)e^{2a+2bx}) de^{2a+2bx}}{4b^2}}{b(d+1)} - \frac{x \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b} - \frac{x^2 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + d+1) \right) +$$

↓ 7143

$$\frac{1}{2}b \left( (d+1) \left( \frac{\frac{\frac{1}{2}x^2 \operatorname{arctanh}(d \coth(a+bx) + d+1) + \operatorname{PolyLog}(3, (d+1)e^{2a+2bx})}{4b^2} - \frac{x \operatorname{PolyLog}(2, (d+1)e^{2a+2bx})}{2b}}{b(d+1)} - \frac{x^2 \log(1 - (d+1)e^{2a+2bx})}{2b(d+1)} \right) + \frac{x^3}{3} \right)$$

input `Int[x*ArcTanh[1 + d + d*Coth[a + b*x]], x]`

output `(x^2*ArcTanh[1 + d + d*Coth[a + b*x]])/2 + (b*(x^3/3 + (1 + d)*(-1/2*(x^2*Log[1 - (1 + d)*E^(2*a + 2*b*x)]))/(b*(1 + d)) + (-1/2*(x*PolyLog[2, (1 + d)*E^(2*a + 2*b*x)])/b + PolyLog[3, (1 + d)*E^(2*a + 2*b*x)]/(4*b^2))/(b*(1 + d)))/2`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6795 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.76 (sec) , antiderivative size = 1555, normalized size of antiderivative = 15.55

method	result	size
risch	Expression too large to display	1555

input `int(x*arctanh(1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/2*x^2*ln(exp(b*x+a))+1/6*b*x^3+1/2/b^2*a/(1+d)*dilog(1+exp(b*x+a)*(1+d)
^(1/2))-1/4/b^2*a^2/(1+d)*ln(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1)-1/4*d/(1+d)
)*ln(1-(1+d)*exp(2*b*x+2*a))*x^2+1/8/b^2/(1+d)*polylog(3,(1+d)*exp(2*b*x+2
*a))-1/4/b^2/(1+d)*ln(1-(1+d)*exp(2*b*x+2*a))*a^2-1/4/b/(1+d)*polylog(2,(1
+d)*exp(2*b*x+2*a))*x-1/4/b^2/(1+d)*polylog(2,(1+d)*exp(2*b*x+2*a))*a+1/8/
b^2*d/(1+d)*polylog(3,(1+d)*exp(2*b*x+2*a))+1/2/b^2*a^2/(1+d)*ln(1-exp(b*x
+a)*(1+d)^(1/2))+1/2/b^2*a^2/(1+d)*ln(1+exp(b*x+a)*(1+d)^(1/2))+1/2/b^2*a/
(1+d)*dilog(1-exp(b*x+a)*(1+d)^(1/2))-1/4/(1+d)*ln(1-(1+d)*exp(2*b*x+2*a))
*x^2-1/8*(I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x
+2*a)-1))^2+I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))^2*csgn(I*exp(2*
b*x+2*a))-2*I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2+2*I*Pi+2*ln
(d)+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2*csgn(I*d)-I*Pi*csgn
(I/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))*csgn(I*ex
p(2*b*x+2*a))-I*Pi*csgn(I*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1))*csgn(I/(exp
(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)+exp(2*b*x+2*a)-1))^2-I*Pi*csgn(I*exp(b*x+
a))^2*csgn(I*exp(2*b*x+2*a))+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2
*a)+exp(2*b*x+2*a)-1))^3+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^
3-I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2
*a)+exp(2*b*x+2*a)-1))^2+I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))*cs
gn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2+I*Pi*csgn(I/(exp(2*b*x+2*a)...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 306 vs.  $2(86) = 172$ .

Time = 0.10 (sec) , antiderivative size = 306, normalized size of antiderivative = 3.06

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{2b^3x^3 + 3b^2x^2 \log\left(-\frac{d \cosh(bx+a) + (d+2) \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) - 6bx \operatorname{Li}_2(\sqrt{d+1}(\cosh(bx+a) + \sinh(bx+a))) - 6b}{b^2}$$

input `integrate(x*arctanh(1+d+d*coth(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*b^3*x^3 + 3*b^2*x^2*log(-(d*cosh(b*x + a) + (d + 2)*sinh(b*x + a)) / (d*cosh(b*x + a) + d*sinh(b*x + a))) - 6*b*x*dilog(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 6*b*x*dilog(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - 3*a^2*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) + 2*sqrt(d + 1)) - 3*a^2*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*sinh(b*x + a) - 2*sqrt(d + 1)) - 3*(b^2*x^2 - a^2)*log(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 3*(b^2*x^2 - a^2)*log(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 6*polylog(3, sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a)))) + 6*polylog(3, -sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x \operatorname{atanh}(d \coth(a + bx) + d + 1) dx$$

input `integrate(x*atanh(1+d+d*coth(b*x+a)),x)`

output `Integral(x*atanh(d*coth(a + b*x) + d + 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.57 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$$

$$= \frac{1}{24} \left( \frac{4x^3}{d} - \frac{3(2b^2x^2 \log(-(d+1)e^{(2bx+2a)} + 1) + 2bx \operatorname{Li}_2((d+1)e^{(2bx+2a)}) - \operatorname{Li}_3((d+1)e^{(2bx+2a)}))}{b^3d} \right) + \frac{1}{2} x^2 \operatorname{artanh}(d \coth(bx + a) + d + 1)$$

input `integrate(x*arctanh(1+d+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/24*(4*x^3/d - 3*(2*b^2*x^2*log(-(d + 1)*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog((d + 1)*e^(2*b*x + 2*a)) - polylog(3, (d + 1)*e^(2*b*x + 2*a)))/(b^3*d) *b*d + 1/2*x^2*arctanh(d*coth(b*x + a) + d + 1)`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x \operatorname{artanh}(d \coth(bx + a) + d + 1) dx$$

input `integrate(x*arctanh(1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*coth(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int x \operatorname{atanh}(d + d \coth(a + bx) + 1) dx$$

input `int(x*atanh(d + d*coth(a + b*x) + 1),x)`

output `int(x*atanh(d + d*coth(a + b*x) + 1), x)`

### Reduce [F]

$$\int x \operatorname{arctanh}(1 + d + d \operatorname{coth}(a + bx)) dx = \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + d + 1) x dx$$

input `int(x*atanh(1+d+d*coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x)*d + d + 1)*x,x)`

### 3.305 $\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx$

Optimal result	2122
Mathematica [A] (verified)	2122
Rubi [A] (verified)	2123
Maple [B] (verified)	2125
Fricas [B] (verification not implemented)	2125
Sympy [F]	2126
Maxima [A] (verification not implemented)	2126
Giac [F]	2127
Mupad [F(-1)]	2127
Reduce [F]	2127

#### Optimal result

Integrand size = 12, antiderivative size = 69

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \frac{bx^2}{2} + x \operatorname{arctanh}(1 + d + d \coth(a + bx)) - \frac{1}{2}x \log(1 - (1 + d)e^{2a+2bx}) - \frac{\operatorname{PolyLog}(2, (1 + d)e^{2a+2bx})}{4b}$$

output

```
1/2*b*x^2+x*arctanh(1+d+d*coth(b*x+a))-1/2*x*ln(1-(1+d)*exp(2*b*x+2*a))-1/4*polylog(2,(1+d)*exp(2*b*x+2*a))/b
```

#### Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.91

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = x \operatorname{arctanh}(1 + d + d \coth(a + bx)) + \frac{-2bx \log\left(1 - \frac{e^{-2(a+bx)}}{1+d}\right) + \operatorname{PolyLog}\left(2, \frac{e^{-2(a+bx)}}{1+d}\right)}{4b}$$

input

```
Integrate[ArcTanh[1 + d + d*Coth[a + b*x]],x]
```

output

```
x*ArcTanh[1 + d + d*Coth[a + b*x]] + (-2*b*x*Log[1 - 1/((1 + d)*E^(2*(a + b*x)))] + PolyLog[2, 1/((1 + d)*E^(2*(a + b*x)))])/(4*b)
```

**Rubi [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.29, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6787, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \coth(a + bx) + d + 1) dx$$

$$\downarrow 6787$$

$$b \int \frac{x}{1 - (d + 1)e^{2a + 2bx}} dx + x \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2615$$

$$b \left( (d + 1) \int \frac{e^{2a + 2bx} x}{1 - (d + 1)e^{2a + 2bx}} dx + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2620$$

$$b \left( (d + 1) \left( \frac{\int \log(1 - (d + 1)e^{2a + 2bx}) dx}{2b(d + 1)} - \frac{x \log(1 - (d + 1)e^{2a + 2bx})}{2b(d + 1)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2715$$

$$b \left( (d + 1) \left( \frac{\int e^{-2a - 2bx} \log(1 - (d + 1)e^{2a + 2bx}) de^{2a + 2bx}}{4b^2(d + 1)} - \frac{x \log(1 - (d + 1)e^{2a + 2bx})}{2b(d + 1)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$

$$\downarrow 2838$$

$$b \left( (d + 1) \left( -\frac{\operatorname{PolyLog}(2, (d + 1)e^{2a + 2bx})}{4b^2(d + 1)} - \frac{x \log(1 - (d + 1)e^{2a + 2bx})}{2b(d + 1)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d \coth(a + bx) + d + 1)$$



input `Int[ArcTanh[1 + d + d*Coth[a + b*x]],x]`

output `x*ArcTanh[1 + d + d*Coth[a + b*x]] + b*(x^2/2 + (1 + d)*(-1/2*(x*Log[1 - (1 + d)*E^(2*a + 2*b*x)])/(b*(1 + d)) - PolyLog[2, (1 + d)*E^(2*a + 2*b*x)]/(4*b^2*(1 + d))))`

### Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 6787 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)], x_Symbol] := Simp[x*ArcTanh[c + d*Coth[a + b*x]], x] + Simp[b Int[x/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, 1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(61) = 122.

Time = 2.77 (sec) , antiderivative size = 255, normalized size of antiderivative = 3.70

method	result
derivativedivides	$\frac{-\frac{\operatorname{arctanh}(1+d+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} + \frac{\operatorname{arctanh}(1+d+d \operatorname{coth}(bx+a))d \ln(d+d \operatorname{coth}(bx+a))}{2}}{d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \operatorname{coth}(bx+a)-d}{-2}\right)}{d^2} \right)}$
default	$\frac{-\frac{\operatorname{arctanh}(1+d+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} + \frac{\operatorname{arctanh}(1+d+d \operatorname{coth}(bx+a))d \ln(d+d \operatorname{coth}(bx+a))}{2}}{d^2 \left( \frac{\operatorname{dilog}\left(\frac{-d \operatorname{coth}(bx+a)-d}{-2}\right)}{d^2} \right)}$
risch	Expression too large to display

input

```
int(arctanh(1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/b/d*(-1/2*arctanh(1+d*d*coth(b*x+a))*d*ln(-d*coth(b*x+a)+d)+1/2*arctanh(1+d*d*coth(b*x+a))*d*ln(d+d*coth(b*x+a))-1/2*d^2*(1/d*(-1/2*dilog((-d*coth(b*x+a)-d-2)/(-2*d-2))-1/2*ln(-d*coth(b*x+a)+d)*ln((-d*coth(b*x+a)-d-2)/(-2*d-2))+1/2*dilog(-1/2*(-d*coth(b*x+a)-d)/d)+1/2*ln(-d*coth(b*x+a)+d)*ln(-1/2*(-d*coth(b*x+a)-d)/d))-1/d*(1/4*ln(d+d*coth(b*x+a))^2-1/2*dilog(1/2*d*coth(b*x+a)+1/2*d+1)-1/2*ln(d+d*coth(b*x+a))*ln(1/2*d*coth(b*x+a)+1/2*d+1)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(60) = 120.

Time = 0.11 (sec) , antiderivative size = 227, normalized size of antiderivative = 3.29

$$\int \operatorname{arctanh}(1 + d + d \operatorname{coth}(a + bx)) dx$$

$$= \frac{b^2 x^2 + bx \log\left(-\frac{d \cosh(bx+a) + (d+2) \sinh(bx+a)}{d \cosh(bx+a) + d \sinh(bx+a)}\right) + a \log\left(2(d+1) \cosh(bx+a) + 2(d+1) \sinh(bx+a) + 2\right)}{2}$$

input

```
integrate(arctanh(1+d*d*coth(b*x+a)),x, algorithm="fricas")
```

output

```
1/2*(b^2*x^2 + b*x*log(-(d*cosh(b*x + a) + (d + 2)*sinh(b*x + a))/(d*cosh(
b*x + a) + d*sinh(b*x + a))) + a*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*s
inh(b*x + a) + 2*sqrt(d + 1)) + a*log(2*(d + 1)*cosh(b*x + a) + 2*(d + 1)*
sinh(b*x + a) - 2*sqrt(d + 1)) - (b*x + a)*log(sqrt(d + 1)*(cosh(b*x + a)
+ sinh(b*x + a)) + 1) - (b*x + a)*log(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b
*x + a)) + 1) - dilog(sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))) - dilog
(-sqrt(d + 1)*(cosh(b*x + a) + sinh(b*x + a))))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int \operatorname{atanh}(d \coth(a + bx) + d + 1) dx$$

input

```
integrate(atanh(1+d+d*coth(b*x+a)),x)
```

output

```
Integral(atanh(d*coth(a + b*x) + d + 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.58 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

$$\begin{aligned} & \int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx \\ &= \frac{1}{4} bd \left( \frac{2x^2}{d} - \frac{2bx \log(-(d+1)e^{(2bx+2a)} + 1) + \operatorname{Li}_2((d+1)e^{(2bx+2a)})}{b^2 d} \right) \\ & \quad + x \operatorname{artanh}(d \coth(bx + a) + d + 1) \end{aligned}$$

input

```
integrate(arctanh(1+d+d*coth(b*x+a)),x, algorithm="maxima")
```

output

```
1/4*b*d*(2*x^2/d - (2*b*x*log(-(d + 1)*e^(2*b*x + 2*a) + 1) + dilog((d + 1
)*e^(2*b*x + 2*a)))/(b^2*d)) + x*arctanh(d*coth(b*x + a) + d + 1)
```

**Giac [F]**

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int \operatorname{artanh}(d \coth(bx + a) + d + 1) dx$$

input `integrate(arctanh(1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*coth(b*x + a) + d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int \operatorname{atanh}(d + d \coth(a + bx) + 1) dx$$

input `int(atanh(d + d*coth(a + b*x) + 1),x)`

output `int(atanh(d + d*coth(a + b*x) + 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 + d + d \coth(a + bx)) dx = \int \operatorname{atanh}(\coth(bx + a) d + d + 1) dx$$

input `int(atanh(1+d+d*coth(b*x+a)),x)`

output `int(atanh(coth(a + b*x)*d + d + 1),x)`

### 3.306 $\int \frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x} dx$

Optimal result	2128
Mathematica [N/A]	2128
Rubi [N/A]	2129
Maple [N/A]	2129
Fricas [N/A]	2130
Sympy [N/A]	2130
Maxima [N/A]	2130
Giac [N/A]	2131
Mupad [N/A]	2131
Reduce [N/A]	2132

#### Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(1+d*d*coth(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 3.52 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1+d+d \coth(a+bx))}{x} dx$$

input `Integrate[ArcTanh[1 + d + d*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 + d + d*Coth[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \coth(a + bx) + d + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \coth(a + bx) + d + 1)}{x} dx$$

input `Int[ArcTanh[1 + d + d*Coth[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.39 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 + d + d \coth(bx + a))}{x} dx$$

input `int(arctanh(1+d+d*coth(b*x+a))/x,x)`

output `int(arctanh(1+d+d*coth(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*coth(b*x + a) + d + 1)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 1.95 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\operatorname{arctanh}(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \coth(a + bx) + d + 1)}{x} dx$$

input `integrate(atanh(1+d+d*coth(b*x+a))/x,x)`

output `Integral(atanh(d*coth(a + b*x) + d + 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.66 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*coth(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctanh(d*coth(b*x + a) + d + 1)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\arctanh(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \coth(bx + a) + d + 1)}{x} dx$$

input `integrate(arctanh(1+d+d*coth(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*coth(b*x + a) + d + 1)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\arctanh(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d + d \coth(a + bx) + 1)}{x} dx$$

input `int(atanh(d + d*coth(a + b*x) + 1)/x,x)`

output `int(atanh(d + d*coth(a + b*x) + 1)/x, x)`



**Reduce [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\operatorname{arctanh}(1 + d + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\coth(bx + a) d + d + 1)}{x} dx$$

input `int(atanh(1+d+d*coth(b*x+a))/x,x)`output `int(atanh(coth(a + b*x)*d + d + 1)/x,x)`

### 3.307 $\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$

Optimal result	2133
Mathematica [A] (verified)	2134
Rubi [A] (verified)	2134
Maple [C] (warning: unable to verify)	2138
Fricas [B] (verification not implemented)	2139
Sympy [F]	2139
Maxima [A] (verification not implemented)	2140
Giac [F]	2140
Mupad [F(-1)]	2140
Reduce [F]	2141

#### Optimal result

Integrand size = 19, antiderivative size = 165

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \frac{bx^5}{20} + \frac{1}{4}x^4 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - \frac{1}{8}x^4 \log(1 - (1 - d)e^{2a+2bx}) - \frac{x^3 \operatorname{PolyLog}(2, (1 - d)e^{2a+2bx})}{4b} + \frac{3x^2 \operatorname{PolyLog}(3, (1 - d)e^{2a+2bx})}{8b^2} - \frac{3x \operatorname{PolyLog}(4, (1 - d)e^{2a+2bx})}{8b^3} + \frac{3 \operatorname{PolyLog}(5, (1 - d)e^{2a+2bx})}{16b^4}$$

output

```
1/20*b*x^5-1/4*x^4*arctanh(-1+d+d*coth(b*x+a))-1/8*x^4*ln(1-(1-d)*exp(2*b*x+2*a))-1/4*x^3*polylog(2,(1-d)*exp(2*b*x+2*a))/b+3/8*x^2*polylog(3,(1-d)*exp(2*b*x+2*a))/b^2-3/8*x*polylog(4,(1-d)*exp(2*b*x+2*a))/b^3+3/16*polylog(5,(1-d)*exp(2*b*x+2*a))/b^4
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.92

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{4b^4 x^4 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - 2b^4 x^4 \log\left(1 + \frac{e^{-2(a+bx)}}{-1+d}\right) + 4b^3 x^3 \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{-1+d}\right) + 6b^2}{16b^4}$$

input

```
Integrate[x^3*ArcTanh[1 - d - d*Coth[a + b*x]],x]
```

output

```
(4*b^4*x^4*ArcTanh[1 - d - d*Coth[a + b*x]] - 2*b^4*x^4*Log[1 + 1/((-1 + d)*E^(2*(a + b*x)))] + 4*b^3*x^3*PolyLog[2, -(1/((-1 + d)*E^(2*(a + b*x))))] + 6*b^2*x^2*PolyLog[3, -(1/((-1 + d)*E^(2*(a + b*x))))] + 6*b*x*PolyLog[4, -(1/((-1 + d)*E^(2*(a + b*x))))] + 3*PolyLog[5, -(1/((-1 + d)*E^(2*(a + b*x))))])/(16*b^4)
```

**Rubi [A] (verified)**

Time = 1.09 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.30, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {6795, 2615, 2620, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1) dx$$

$$\downarrow \text{6795}$$

$$\frac{1}{4}b \int \frac{x^4}{1 - (1 - d)e^{2a+2bx}} dx + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{4}b \left( (1 - d) \int \frac{e^{2a+2bx} x^4}{1 - (1 - d)e^{2a+2bx}} dx + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{4}b \left( (1-d) \left( \frac{2 \int x^3 \log(1 - (1-d)e^{2a+2bx}) dx}{b(1-d)} - \frac{x^4 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{4}b \left( (1-d) \left( \frac{2 \left( \frac{3 \int x^2 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx}) dx}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} - \frac{x^4 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^5}{5} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{4}b \left( (1-d) \left( \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{\int x \operatorname{PolyLog}(3, (1-d)e^{2a+2bx}) dx}{b} \right)}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} - \frac{x^4 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{4}b \left( (1-d) \left( \frac{2 \left( \frac{3 \left( \frac{x^2 \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{x \operatorname{PolyLog}(4, (1-d)e^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(4, (1-d)e^{2a+2bx}) dx}{2b} \right)}{2b} - \frac{x^3 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} + \frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{4}b(1-d) \left( \frac{2 \left( 3 \left( \frac{x^2 \text{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{x \text{PolyLog}(4, (1-d)e^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \text{PolyLog}(4, (1-d)e^{2a+2bx}) de^{2a+2bx}}{b} \right)}{2b} - \frac{x^3 \text{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} \right)$$

$$\frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 7143

$$\frac{1}{4}x^4 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1) +$$

$$\frac{1}{4}b(1-d) \left( \frac{2 \left( 3 \left( \frac{x^2 \text{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{x \text{PolyLog}(4, (1-d)e^{2a+2bx})}{2b} - \frac{\text{PolyLog}(5, (1-d)e^{2a+2bx})}{4b^2} \right)}{2b} - \frac{x^3 \text{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{b(1-d)} \right)$$

input `Int[x^3*ArcTanh[1 - d - d*Coth[a + b*x]],x]`

output `(x^4*ArcTanh[1 - d - d*Coth[a + b*x]])/4 + (b*(x^5/5 + (1 - d)*(-1/2*(x^4*Log[1 - (1 - d)*E^(2*a + 2*b*x)]))/(b*(1 - d)) + (2*(-1/2*(x^3*PolyLog[2, (1 - d)*E^(2*a + 2*b*x)])/b + (3*((x^2*PolyLog[3, (1 - d)*E^(2*a + 2*b*x)])/(2*b) - ((x*PolyLog[4, (1 - d)*E^(2*a + 2*b*x)])/(2*b) - PolyLog[5, (1 - d)*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b)))/(b*(1 - d)))/4`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6795 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.
)*(x_)))]^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.31 (sec) , antiderivative size = 1753, normalized size of antiderivative = 10.62

method	result	size
risch	Expression too large to display	1753

input

```
int(-x^3*arctanh(-1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/8/b^4*a^4/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)+1)+1/20*b*x^5-1/4*x^4
*ln(exp(b*x+a))-1/16*(-I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)-ex
p(2*b*x+2*a)+1))^3-2*I*Pi+2*ln(d)-I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b
*x+2*a))^3+I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*
x+2*a)-1))^2+I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))^2*csgn(I*exp(2
*b*x+2*a))+I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*(d*exp(2*b*x+2*a)-exp(2*
b*x+2*a)+1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)+1)
)-I*Pi*csgn(I*exp(b*x+a))^2*csgn(I*exp(2*b*x+2*a))-I*Pi*csgn(I/(exp(2*b*x+
2*a)-1))*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))*csgn(I*exp(2*b*x+2*a))-
I*Pi*csgn(I/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a
)-exp(2*b*x+2*a)+1))^2+I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^2*
csgn(I*d)+2*I*Pi*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a)-exp(2*b*x+2*a
)+1))^2-I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1))^3-I*Pi*csgn(I*(d*ex
p(2*b*x+2*a)-exp(2*b*x+2*a)+1))*csgn(I/(exp(2*b*x+2*a)-1)*(d*exp(2*b*x+2*a
)-exp(2*b*x+2*a)+1))^2-I*Pi*csgn(I*exp(2*b*x+2*a))^3+2*I*Pi*csgn(I*exp(b*x
+a))*csgn(I*exp(2*b*x+2*a))^2-I*Pi*csgn(I*exp(2*b*x+2*a)/(exp(2*b*x+2*a)-1
))*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))*csgn(I*d)+I*Pi*csgn(I*exp(2
*b*x+2*a)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1)*d*exp(2*b*x+2*a))^
2)*x^4-1/2/b^3*a^3/(d-1)*ln(1-exp(b*x+a)*(1-d)^(1/2))*x-1/2/b^4*a^3/(d-1)*
dilog(1+exp(b*x+a)*(1-d)^(1/2))-1/2/b^4*a^3/(d-1)*dilog(1-exp(b*x+a)*(1...
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 451 vs.  $2(135) = 270$ .

Time = 0.11 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.73

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \text{Too large to display}$$

input `integrate(-x^3*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="fricas")`

output `1/40*(2*b^5*x^5 - 5*b^4*x^4*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/(d*cosh(b*x + a) + (d - 2)*sinh(b*x + a))) - 20*b^3*x^3*dilog(1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 20*b^3*x^3*dilog(-1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 5*a^4*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) + sqrt(-4*d + 4)) - 5*a^4*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) - sqrt(-4*d + 4)) + 60*b^2*x^2*polylog(3, 1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) + 60*b^2*x^2*polylog(3, -1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, 1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 120*b*x*polylog(4, -1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 5*(b^4*x^4 - a^4)*log(1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 5*(b^4*x^4 - a^4)*log(-1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 120*polylog(5, 1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) + 120*polylog(5, -1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^4`

**Sympy [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = - \int x^3 \operatorname{atanh}(d \coth(a + bx) + d - 1) dx$$

input `integrate(-x**3*atanh(-1+d+d*coth(b*x+a)),x)`

output `-Integral(x**3*atanh(d*coth(a + b*x) + d - 1), x)`



**Maxima [A] (verification not implemented)**

Time = 0.57 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.90

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = -\frac{1}{4} x^4 \operatorname{artanh}(d \coth(bx + a) + d - 1) + \frac{1}{40} \left( \frac{2x^5}{d} - \frac{5(2b^4 x^4 \log((d-1)e^{(2bx+2a)} + 1) + 4b^3 x^3 \operatorname{Li}_2(-(d-1)e^{(2bx+2a)}) - 6b^2 x^2 \operatorname{Li}_3(-(d-1)e^{(2bx+2a)}) + 6b x \operatorname{Li}_4(-(d-1)e^{(2bx+2a)}) - 3 \operatorname{Li}_5(-(d-1)e^{(2bx+2a)}))}{b^5 d} \right)$$

input `integrate(-x^3*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="maxima")`

output `-1/4*x^4*arctanh(d*coth(b*x + a) + d - 1) + 1/40*(2*x^5/d - 5*(2*b^4*x^4*log((d - 1)*e^(2*b*x + 2*a) + 1) + 4*b^3*x^3*dilog(-(d - 1)*e^(2*b*x + 2*a)) - 6*b^2*x^2*polylog(3, -(d - 1)*e^(2*b*x + 2*a)) + 6*b*x*polylog(4, -(d - 1)*e^(2*b*x + 2*a)) - 3*polylog(5, -(d - 1)*e^(2*b*x + 2*a)))/(b^5*d))*d`

**Giac [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -x^3 \operatorname{artanh}(d \coth(bx + a) + d - 1) dx$$

input `integrate(-x^3*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(-x^3*arctanh(d*coth(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -x^3 \operatorname{atanh}(d + d \coth(a + bx) - 1) dx$$

input `int(-x^3*atanh(d + d*coth(a + b*x) - 1),x)`

output `int(-x^3*atanh(d + d*coth(a + b*x) - 1), x)`

**Reduce [F]**

$$\int x^3 \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx = - \left( \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + d - 1) x^3 dx \right)$$

input `int(-x^3*atanh(-1+d+d*coth(b*x+a)), x)`

output `- int(atanh(coth(a + b*x)*d + d - 1)*x**3, x)`

### 3.308 $\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$

Optimal result	2142
Mathematica [A] (verified)	2143
Rubi [A] (verified)	2143
Maple [C] (warning: unable to verify)	2146
Fricas [B] (verification not implemented)	2147
Sympy [F]	2148
Maxima [A] (verification not implemented)	2148
Giac [F]	2149
Mupad [F(-1)]	2149
Reduce [F]	2149

#### Optimal result

Integrand size = 19, antiderivative size = 137

$$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - \frac{1}{6}x^3 \log(1 - (1 - d)e^{2a+2bx}) - \frac{x^2 \operatorname{PolyLog}(2, (1 - d)e^{2a+2bx})}{4b} + \frac{x \operatorname{PolyLog}(3, (1 - d)e^{2a+2bx})}{4b^2} - \frac{\operatorname{PolyLog}(4, (1 - d)e^{2a+2bx})}{8b^3}$$

output

```
1/12*b*x^4-1/3*x^3*arctanh(-1+d+d*coth(b*x+a))-1/6*x^3*ln(1-(1-d)*exp(2*b*x+2*a))-1/4*x^2*polylog(2,(1-d)*exp(2*b*x+2*a))/b+1/4*x*polylog(3,(1-d)*exp(2*b*x+2*a))/b^2-1/8*polylog(4,(1-d)*exp(2*b*x+2*a))/b^3
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.91

$$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - 4b^3 x^3 \log\left(1 + \frac{e^{-2(a+bx)}}{-1+d}\right) + 6b^2 x^2 \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{-1+d}\right) + 6bx}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 - d - d*Coth[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTanh[1 - d - d*Coth[a + b*x]] - 4*b^3*x^3*Log[1 + 1/((-1 + d)*E^(2*(a + b*x)))] + 6*b^2*x^2*PolyLog[2, -(1/((-1 + d)*E^(2*(a + b*x))))] + 6*b*x*PolyLog[3, -(1/((-1 + d)*E^(2*(a + b*x)))))] + 3*PolyLog[4, -(1/((-1 + d)*E^(2*(a + b*x)))))]/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.86 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.31, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6795, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1) dx$$

$$\downarrow \text{6795}$$

$$\frac{1}{3}b \int \frac{x^3}{1 - (1 - d)e^{2a+2bx}} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}b \left( (1 - d) \int \frac{e^{2a+2bx} x^3}{1 - (1 - d)e^{2a+2bx}} dx + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}b \left( (1-d) \left( \frac{3 \int x^2 \log(1 - (1-d)e^{2a+2bx}) dx}{2b(1-d)} - \frac{x^3 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{3}b \left( (1-d) \left( \frac{3 \left( \frac{\int x \operatorname{PolyLog}(2, (1-d)e^{2a+2bx}) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{2b(1-d)} - \frac{x^3 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^4}{4} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 7163

$$\frac{1}{3}b \left( (1-d) \left( \frac{3 \left( \frac{\frac{x \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(3, (1-d)e^{2a+2bx}) dx}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{2b(1-d)} - \frac{x^3 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{3}b \left( (1-d) \left( \frac{3 \left( \frac{\frac{\frac{x \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, (1-d)e^{2a+2bx}) de^{2a+2bx}}{4b^2}}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{2b(1-d)} - \frac{x^3 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 7143

$$\frac{1}{3}b \left( (1-d) \left( \frac{3 \left( \frac{\frac{\frac{\frac{x \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{2b} - \frac{\operatorname{PolyLog}(4, (1-d)e^{2a+2bx})}{4b^2}}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} \right)}{2b(1-d)} - \frac{x^3 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1) \right) +$$

input `Int[x^2*ArcTanh[1 - d - d*Coth[a + b*x]], x]`

output

```
(x^3*ArcTanh[1 - d - d*Coth[a + b*x]])/3 + (b*(x^4/4 + (1 - d)*(-1/2*(x^3*
Log[1 - (1 - d)*E^(2*a + 2*b*x)]))/(b*(1 - d)) + (3*(-1/2*(x^2*PolyLog[2, (
1 - d)*E^(2*a + 2*b*x)])/b + ((x*PolyLog[3, (1 - d)*E^(2*a + 2*b*x)])/(2*b
) - PolyLog[4, (1 - d)*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b*(1 - d))))/3
```

### Defintions of rubi rules used

rule 2615

```
Int[(((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x
_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[
b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x]
, x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/
((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6795

```
Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(
m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a
+ 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c
- d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.26 (sec) , antiderivative size = 1694, normalized size of antiderivative = 12.36

method	result	size
risch	Expression too large to display	1694

input

```
int(-x^2*arctanh(-1+d+d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/12*b*x^4-1/3*x^3*ln(exp(b*x+a))+1/8/b^3/(d-1)*polylog(4,-(d-1)*exp(2*b*x
+2*a))+1/6/(d-1)*ln(1+(d-1)*exp(2*b*x+2*a))*x^3-1/6*d/(d-1)*ln(1+(d-1)*exp
(2*b*x+2*a))*x^3-1/6/b^3*a^3/(d-1)*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)+1)+1
/2/b^3*a^3/(d-1)*ln(1+exp(b*x+a)*(1-d)^(1/2))+1/2/b^3*a^3/(d-1)*ln(1-exp(b
*x+a)*(1-d)^(1/2))+1/2/b^3*a^2/(d-1)*dilog(1+exp(b*x+a)*(1-d)^(1/2))+1/2/b
^3*a^2/(d-1)*dilog(1-exp(b*x+a)*(1-d)^(1/2))-1/8/b^3*d/(d-1)*polylog(4,-(d
-1)*exp(2*b*x+2*a))+1/4/b/(d-1)*polylog(2,-(d-1)*exp(2*b*x+2*a))*x^2-1/3/b
^3/(d-1)*ln(1+(d-1)*exp(2*b*x+2*a))*a^3-1/4/b^3/(d-1)*polylog(2,-(d-1)*exp
(2*b*x+2*a))*a^2-1/4/b^2/(d-1)*polylog(3,-(d-1)*exp(2*b*x+2*a))*x+1/2/b^2*
d/(d-1)*ln(1+(d-1)*exp(2*b*x+2*a))*a^2*x-1/2/b^2*d*a^2/(d-1)*x*ln(1+exp(b*
x+a)*(1-d)^(1/2))-1/2/b^2*d*a^2/(d-1)*x*ln(1-exp(b*x+a)*(1-d)^(1/2))-1/4/b
*d/(d-1)*polylog(2,-(d-1)*exp(2*b*x+2*a))*x^2+1/3/b^3*d/(d-1)*ln(1+(d-1)*e
xp(2*b*x+2*a))*a^3+1/4/b^3*d/(d-1)*polylog(2,-(d-1)*exp(2*b*x+2*a))*a^2+1/
4/b^2*d/(d-1)*polylog(3,-(d-1)*exp(2*b*x+2*a))*x-1/2/b^2/(d-1)*ln(1+(d-1)*
exp(2*b*x+2*a))*a^2*x+1/2/b^2*a^2/(d-1)*x*ln(1+exp(b*x+a)*(1-d)^(1/2))+1/2
/b^2*a^2/(d-1)*x*ln(1-exp(b*x+a)*(1-d)^(1/2))+1/6/b^3*d*a^3/(d-1)*ln(d*exp
(2*b*x+2*a)-exp(2*b*x+2*a)+1)-1/2/b^3*d*a^3/(d-1)*ln(1+exp(b*x+a)*(1-d)^(1
/2))-1/2/b^3*d*a^3/(d-1)*ln(1-exp(b*x+a)*(1-d)^(1/2))-1/2/b^3*d*a^2/(d-1)*
dilog(1+exp(b*x+a)*(1-d)^(1/2))-1/2/b^3*d*a^2/(d-1)*dilog(1-exp(b*x+a)*(1-
d)^(1/2))+1/6*x^3*ln(d*exp(2*b*x+2*a)-exp(2*b*x+2*a)+1)-1/12*(-I*Pi*csg...

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 382 vs.  $2(111) = 222$ .

Time = 0.10 (sec) , antiderivative size = 382, normalized size of antiderivative = 2.79

$$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{b^4 x^4 - 2 b^3 x^3 \log\left(-\frac{d \cosh(bx+a) + d \sinh(bx+a)}{d \cosh(bx+a) + (d-2) \sinh(bx+a)}\right) - 6 b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4d+4} (\cosh(bx+a) + \sinh(bx+a))\right)}{b^4}$$

input

```
integrate(-x^2*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="fricas")
```



output

```
1/12*(b^4*x^4 - 2*b^3*x^3*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/(d*cosh
(b*x + a) + (d - 2)*sinh(b*x + a))) - 6*b^2*x^2*dilog(1/2*sqrt(-4*d + 4)*(
cosh(b*x + a) + sinh(b*x + a))) - 6*b^2*x^2*dilog(-1/2*sqrt(-4*d + 4)*(cos
h(b*x + a) + sinh(b*x + a))) + 2*a^3*log(2*(d - 1)*cosh(b*x + a) + 2*(d -
1)*sinh(b*x + a) + sqrt(-4*d + 4)) + 2*a^3*log(2*(d - 1)*cosh(b*x + a) + 2
*(d - 1)*sinh(b*x + a) - sqrt(-4*d + 4)) + 12*b*x*polylog(3, 1/2*sqrt(-4*d
+ 4)*(cosh(b*x + a) + sinh(b*x + a))) + 12*b*x*polylog(3, -1/2*sqrt(-4*d
+ 4)*(cosh(b*x + a) + sinh(b*x + a))) - 2*(b^3*x^3 + a^3)*log(1/2*sqrt(-4*d
+ 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 2*(b^3*x^3 + a^3)*log(-1/2*sqr
t(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 12*polylog(4, 1/2*sqr
t(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 12*polylog(4, -1/2*sqrt(-4*d
+ 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^3
```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = - \int x^2 \operatorname{atanh}(d \coth(a + bx) + d - 1) dx$$

input

```
integrate(-x**2*atanh(-1+d+d*coth(b*x+a)),x)
```

output

```
-Integral(x**2*atanh(d*coth(a + b*x) + d - 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.60 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.91

$$\int x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = -\frac{1}{3} x^3 \operatorname{artanh}(d \coth(bx + a) + d - 1) + \frac{1}{36} \left( \frac{3x^4}{d} - \frac{2(4b^3x^3 \log((d-1)e^{(2bx+2a)} + 1) + 6b^2x^2 \operatorname{Li}_2(-(d-1)e^{(2bx+2a)}) - 6bx \operatorname{Li}_3(-(d-1)e^{(2bx+2a)}))}{b^4d} \right)$$

input

```
integrate(-x^2*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="maxima")
```

output

```
-1/3*x^3*arctanh(d*coth(b*x + a) + d - 1) + 1/36*(3*x^4/d - 2*(4*b^3*x^3*log((d - 1)*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(-(d - 1)*e^(2*b*x + 2*a)) - 6*b*x*polylog(3, -(d - 1)*e^(2*b*x + 2*a)) + 3*polylog(4, -(d - 1)*e^(2*b*x + 2*a)))/(b^4*d))*b*d
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx = \int -x^2 \operatorname{artanh}(d \operatorname{coth}(bx + a) + d - 1) dx$$

input

```
integrate(-x^2*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="giac")
```

output

```
integrate(-x^2*arctanh(d*coth(b*x + a) + d - 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx = \int -x^2 \operatorname{atanh}(d + d \operatorname{coth}(a + bx) - 1) dx$$

input

```
int(-x^2*atanh(d + d*coth(a + b*x) - 1),x)
```

output

```
int(-x^2*atanh(d + d*coth(a + b*x) - 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx = - \left( \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + d - 1) x^2 dx \right)$$

input

```
int(-x^2*atanh(-1+d+d*coth(b*x+a)),x)
```

output `- int(atanh(coth(a + b*x)*d + d - 1)*x**2,x)`

### 3.309 $\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$

Optimal result	2151
Mathematica [A] (verified)	2152
Rubi [A] (verified)	2152
Maple [C] (warning: unable to verify)	2155
Fricas [B] (verification not implemented)	2156
Sympy [F]	2156
Maxima [A] (verification not implemented)	2157
Giac [F]	2157
Mupad [F(-1)]	2157
Reduce [F]	2158

#### Optimal result

Integrand size = 17, antiderivative size = 109

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - \frac{1}{4}x^2 \log(1 - (1 - d)e^{2a+2bx}) - \frac{x \operatorname{PolyLog}(2, (1 - d)e^{2a+2bx})}{4b} + \frac{\operatorname{PolyLog}(3, (1 - d)e^{2a+2bx})}{8b^2}$$

output

```
1/6*b*x^3-1/2*x^2*arctanh(-1+d+d*coth(b*x+a))-1/4*x^2*ln(1-(1-d)*exp(2*b*x+2*a))-1/4*x*polylog(2,(1-d)*exp(2*b*x+2*a))/b+1/8*polylog(3,(1-d)*exp(2*b*x+2*a))/b^2
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.86

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left( 2 \operatorname{arctanh}(1 - d - d \coth(a + bx)) - \log \left( 1 + \frac{e^{-2(a+bx)}}{-1+d} \right) \right) + 2bx \operatorname{PolyLog} \left( 2, -\frac{e^{-2(a+bx)}}{-1+d} \right) + \operatorname{PolyLog} \left( 3, -\frac{e^{-2(a+bx)}}{-1+d} \right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 - d - d*Coth[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTanh[1 - d - d*Coth[a + b*x]] - Log[1 + 1/((-1 + d)*E^(2*(a + b*x))])) + 2*b*x*PolyLog[2, -(1/((-1 + d)*E^(2*(a + b*x)))] + PolyLog[3, -(1/((-1 + d)*E^(2*(a + b*x)))])/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.69 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.31, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$ , Rules used = {6795, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1) dx$$

$$\downarrow \text{6795}$$

$$\frac{1}{2}b \int \frac{x^2}{1 - (1 - d)e^{2a+2bx}} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2}b \left( (1 - d) \int \frac{e^{2a+2bx} x^2}{1 - (1 - d)e^{2a+2bx}} dx + \frac{x^3}{3} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a + bx)) - d + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}b \left( (1-d) \left( \frac{\int x \log(1 - (1-d)e^{2a+2bx}) dx}{b(1-d)} - \frac{x^2 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^3}{3} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 3011

$$\frac{1}{2}b \left( (1-d) \left( \frac{\frac{\int \operatorname{PolyLog}(2, (1-d)e^{2a+2bx}) dx}{2b}}{b(1-d)} - \frac{x \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} - \frac{x^2 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^3}{3} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)$$

↓ 2720

$$\frac{1}{2}b \left( (1-d) \left( \frac{\frac{\int e^{-2a-2bx} \operatorname{PolyLog}(2, (1-d)e^{2a+2bx}) de^{2a+2bx}}{4b^2}}{b(1-d)} - \frac{x \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} - \frac{x^2 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1) \right) +$$

↓ 7143

$$\frac{1}{2}b \left( (1-d) \left( \frac{\frac{\frac{1}{2}x^2 \operatorname{arctanh}(d(-\coth(a+bx)) - d + 1) + \operatorname{PolyLog}(3, (1-d)e^{2a+2bx})}{4b^2}}{b(1-d)} - \frac{x \operatorname{PolyLog}(2, (1-d)e^{2a+2bx})}{2b} - \frac{x^2 \log(1 - (1-d)e^{2a+2bx})}{2b(1-d)} \right) + \frac{x^3}{3} \right) +$$

input `Int[x*ArcTanh[1 - d - d*Coth[a + b*x]], x]`

output `(x^2*ArcTanh[1 - d - d*Coth[a + b*x]])/2 + (b*(x^3/3 + (1 - d)*(-1/2*(x^2*Log[1 - (1 - d)*E^(2*a + 2*b*x)]))/(b*(1 - d)) + (-1/2*(x*PolyLog[2, (1 - d)*E^(2*a + 2*b*x)])/b + PolyLog[3, (1 - d)*E^(2*a + 2*b*x)]/(4*b^2))/(b*(1 - d))))/2`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6795 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.52 (sec) , antiderivative size = 1611, normalized size of antiderivative = 14.78

method	result	size
risch	Expression too large to display	1611

input `int(-x*arctanh(-1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/2*x^2*\ln(\exp(b*x+a))+1/6*b*x^3+1/2/b^2*d*a/(d-1)*\operatorname{dilog}(1+\exp(b*x+a)*(1-d)^{(1/2)})+1/2/b^2*d*a/(d-1)*\operatorname{dilog}(1-\exp(b*x+a)*(1-d)^{(1/2)})-1/4/b^2*d/(d-1) \\
 & *\ln(1+(d-1)*\exp(2*b*x+2*a))*a^2-1/4/b*d/(d-1)*\operatorname{polylog}(2,-(d-1)*\exp(2*b*x+2*a))*x-1/4/b^2*d/(d-1)*\operatorname{polylog}(2,-(d-1)*\exp(2*b*x+2*a))*a+1/2/b/(d-1)*\ln(1+(d-1)*\exp(2*b*x+2*a))*a*x-1/2/b*a/(d-1)*x*\ln(1+\exp(b*x+a)*(1-d)^{(1/2)})-1/2/b*a/(d-1)*x*\ln(1-\exp(b*x+a)*(1-d)^{(1/2)})+1/2/b^2*d*a^2/(d-1)*\ln(1+\exp(b*x+a)*(1-d)^{(1/2)})-1/4/b^2*d*a^2/(d-1)*\ln(d*\exp(2*b*x+2*a)-\exp(2*b*x+2*a)+1)+1/8/b^2*d/(d-1)*\operatorname{polylog}(3,-(d-1)*\exp(2*b*x+2*a))+1/4/b^2/(d-1)*\ln(1+(d-1)*\exp(2*b*x+2*a))*a^2+1/4/b/(d-1)*\operatorname{polylog}(2,-(d-1)*\exp(2*b*x+2*a))*x+1/4/b^2/(d-1)*\operatorname{polylog}(2,-(d-1)*\exp(2*b*x+2*a))*a-1/2/b^2*a^2/(d-1)*\ln(1+\exp(b*x+a)*(1-d)^{(1/2)})-1/2/b^2*a^2/(d-1)*\ln(1-\exp(b*x+a)*(1-d)^{(1/2)})-1/2/b^2*a/(d-1)*\operatorname{dilog}(1+\exp(b*x+a)*(1-d)^{(1/2)})-1/2/b^2*a/(d-1)*\operatorname{dilog}(1-\exp(b*x+a)*(1-d)^{(1/2)})+1/4/b^2*a^2/(d-1)*\ln(d*\exp(2*b*x+2*a)-\exp(2*b*x+2*a)+1)-1/4*d/(d-1)*\ln(1+(d-1)*\exp(2*b*x+2*a))*x^2+1/4/(d-1)*\ln(1+(d-1)*\exp(2*b*x+2*a))*x^2-1/8/b^2/(d-1)*\operatorname{polylog}(3,-(d-1)*\exp(2*b*x+2*a))-1/2/b*d/(d-1)*\ln(1+(d-1)*\exp(2*b*x+2*a))*a*x+1/2/b*d*a/(d-1)*x*\ln(1+\exp(b*x+a)*(1-d)^{(1/2)})+1/2/b*d*a/(d-1)*x*\ln(1-\exp(b*x+a)*(1-d)^{(1/2)})+1/4*x^2*\ln(d*\exp(2*b*x+2*a)-\exp(2*b*x+2*a)+1)-1/8*(-I*Pi*csgn(I/(\exp(2*b*x+2*a)-1))*(d*\exp(2*b*x+2*a)-\exp(2*b*x+2*a)+1))^3-2*I*Pi+2*\ln(d)-I*Pi*csgn(I/(\exp(2*b*x+2*a)-1)*d*\exp(2*b*x+2*a))^3+I*Pi*csgn(I/(\exp(2*b*x+2*a)-1))*csgn(I*\exp(2*b*x+2*a)/(\exp(2*b...
 \end{aligned}$$



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 323 vs.  $2(87) = 174$ .

Time = 0.10 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.96

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{2b^3x^3 - 3b^2x^2 \log\left(-\frac{d \cosh(bx+a) + d \sinh(bx+a)}{d \cosh(bx+a) + (d-2) \sinh(bx+a)}\right) - 6bx \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4d+4} (\cosh(bx+a) + \sinh(bx+a))\right)}{b^2}$$

input `integrate(-x*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="fricas")`

output

```
1/12*(2*b^3*x^3 - 3*b^2*x^2*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/(d*cosh(b*x + a) + (d - 2)*sinh(b*x + a))) - 6*b*x*dilog(1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 6*b*x*dilog(-1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) - 3*a^2*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) + sqrt(-4*d + 4)) - 3*a^2*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*sinh(b*x + a) - sqrt(-4*d + 4)) - 3*(b^2*x^2 - a^2)*log(1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - 3*(b^2*x^2 - a^2)*log(-1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + 6*polylog(3, 1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))) + 6*polylog(3, -1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a))))/b^2
```

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = - \int x \operatorname{atanh}(d \coth(a + bx) + d - 1) dx$$

input `integrate(-x*atanh(-1+d+d*coth(b*x+a)),x)`

output `-Integral(x*atanh(d*coth(a + b*x) + d - 1), x)`

**Maxima [A] (verification not implemented)**

Time = 0.60 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.93

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$$

$$= \frac{1}{24} \left( \frac{4x^3}{d} - \frac{3(2b^2x^2 \log((d-1)e^{2bx+2a}) + 1) + 2bx \operatorname{Li}_2(-(d-1)e^{2bx+2a}) - \operatorname{Li}_3(-(d-1)e^{2bx+2a}))}{b^3d} \right) - \frac{1}{2} x^2 \operatorname{artanh}(d \coth(bx + a) + d - 1)$$

input `integrate(-x*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/24*(4*x^3/d - 3*(2*b^2*x^2*log((d - 1)*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(-(d - 1)*e^(2*b*x + 2*a)) - polylog(3, -(d - 1)*e^(2*b*x + 2*a)))/(b^3*d)))*b*d - 1/2*x^2*arctanh(d*coth(b*x + a) + d - 1)`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -x \operatorname{artanh}(d \coth(bx + a) + d - 1) dx$$

input `integrate(-x*arctanh(-1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctanh(d*coth(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -x \operatorname{atanh}(d + d \coth(a + bx) - 1) dx$$

input `int(-x*atanh(d + d*coth(a + b*x) - 1),x)`

output `int(-x*atanh(d + d*coth(a + b*x) - 1), x)`

### Reduce [F]

$$\int x \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx = - \left( \int \operatorname{atanh}(\operatorname{coth}(bx + a) d + d - 1) x dx \right)$$

input `int(-x*atanh(-1+d+d*coth(b*x+a)),x)`

output `- int(atanh(coth(a + b*x)*d + d - 1)*x,x)`

### 3.310 $\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx$

Optimal result	2159
Mathematica [A] (verified)	2159
Rubi [A] (verified)	2160
Maple [B] (verified)	2162
Fricas [B] (verification not implemented)	2162
Sympy [F]	2163
Maxima [A] (verification not implemented)	2163
Giac [F]	2164
Mupad [F(-1)]	2164
Reduce [F]	2164

#### Optimal result

Integrand size = 15, antiderivative size = 76

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \frac{bx^2}{2} + x \operatorname{arctanh}(1 - d - d \coth(a + bx)) - \frac{1}{2} x \log(1 - (1 - d)e^{2a+2bx}) - \frac{\operatorname{PolyLog}(2, (1 - d)e^{2a+2bx})}{4b}$$

output

```
1/2*b*x^2-x*arctanh(-1+d+d*coth(b*x+a))-1/2*x*ln(1-(1-d)*exp(2*b*x+2*a))-1/4*polylog(2,(1-d)*exp(2*b*x+2*a))/b
```

#### Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = x \operatorname{arctanh}(1 - d - d \coth(a + bx)) + \frac{-2bx \log\left(1 + \frac{e^{-2(a+bx)}}{-1+d}\right) + \operatorname{PolyLog}\left(2, -\frac{e^{-2(a+bx)}}{-1+d}\right)}{4b}$$

input

```
Integrate[ArcTanh[1 - d - d*Coth[a + b*x]], x]
```

output

```
x*ArcTanh[1 - d - d*Coth[a + b*x]] + (-2*b*x*Log[1 + 1/((-1 + d)*E^(2*(a +
b*x)))] + PolyLog[2, -(1/((-1 + d)*E^(2*(a + b*x))))])/(4*b)
```

**Rubi [A] (verified)**

Time = 0.46 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.34, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6787, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1) dx$$

$$\downarrow 6787$$

$$b \int \frac{x}{1 - (1 - d)e^{2a+2bx}} dx + x \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1)$$

$$\downarrow 2615$$

$$b \left( (1 - d) \int \frac{e^{2a+2bx} x}{1 - (1 - d)e^{2a+2bx}} dx + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1)$$

$$\downarrow 2620$$

$$b \left( (1 - d) \left( \frac{\int \log(1 - (1 - d)e^{2a+2bx}) dx}{2b(1 - d)} - \frac{x \log(1 - (1 - d)e^{2a+2bx})}{2b(1 - d)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1)$$

$$\downarrow 2715$$

$$b \left( (1 - d) \left( \frac{\int e^{-2a-2bx} \log(1 - (1 - d)e^{2a+2bx}) de^{2a+2bx}}{4b^2(1 - d)} - \frac{x \log(1 - (1 - d)e^{2a+2bx})}{2b(1 - d)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1)$$

$$\downarrow 2838$$

$$b \left( (1 - d) \left( -\frac{\operatorname{PolyLog}(2, (1 - d)e^{2a+2bx})}{4b^2(1 - d)} - \frac{x \log(1 - (1 - d)e^{2a+2bx})}{2b(1 - d)} \right) + \frac{x^2}{2} \right) + x \operatorname{arctanh}(d(-\operatorname{coth}(a + bx)) - d + 1)$$

input `Int[ArcTanh[1 - d - d*Coth[a + b*x]],x]`

output `x*ArcTanh[1 - d - d*Coth[a + b*x]] + b*(x^2/2 + (1 - d)*(-1/2*(x*Log[1 - (1 - d)*E^(2*a + 2*b*x)])/(b*(1 - d)) - PolyLog[2, (1 - d)*E^(2*a + 2*b*x)]/(4*b^2*(1 - d))))`

### Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 6787 `Int[ArcTanh[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)], x_Symbol] := Simp[x*ArcTanh[c + d*Coth[a + b*x]], x] + Simp[b Int[x/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, 1]`

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(66) = 132.

Time = 2.57 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.58

method	result
derivativedivides	$-\frac{\operatorname{arctanh}(-1+d+d \operatorname{coth}(bx+a))d \ln(d+d \operatorname{coth}(bx+a))}{2} - \frac{\operatorname{arctanh}(-1+d+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} - \frac{d^2 \left( \frac{\operatorname{dilog}\left(-\frac{-d \operatorname{coth}(bx+a)-d}{d}\right)}{\dots} \right)}{\dots}$
default	$-\frac{\operatorname{arctanh}(-1+d+d \operatorname{coth}(bx+a))d \ln(d+d \operatorname{coth}(bx+a))}{2} - \frac{\operatorname{arctanh}(-1+d+d \operatorname{coth}(bx+a))d \ln(-d \operatorname{coth}(bx+a)+d)}{2} - \frac{d^2 \left( \frac{\operatorname{dilog}\left(-\frac{-d \operatorname{coth}(bx+a)-d}{d}\right)}{\dots} \right)}{\dots}$
risch	Expression too large to display

input

```
int(-arctanh(-1+d*d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/b/d*(1/2*arctanh(-1+d*d*coth(b*x+a))*d*ln(d+d*coth(b*x+a))-1/2*arctanh(-1+d*d*coth(b*x+a))*d*ln(-d*coth(b*x+a)+d)-1/2*d^2*(1/d*(-1/2*dilog(-1/2*(-d*coth(b*x+a)-d)/d)-1/2*ln(-d*coth(b*x+a)+d)*ln(-1/2*(-d*coth(b*x+a)-d)/d)+1/2*dilog((-d*coth(b*x+a)-d+2)/(-2*d+2))+1/2*ln(-d*coth(b*x+a)+d)*ln((-d*coth(b*x+a)-d+2)/(-2*d+2)))-1/d*(-1/4*ln(d+d*coth(b*x+a))^2+1/2*(ln(d+d*coth(b*x+a))-ln(1/2*d*coth(b*x+a)+1/2*d))*ln(-1/2*d*coth(b*x+a)-1/2*d+1)-1/2*dilog(1/2*d*coth(b*x+a)+1/2*d)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(61) = 122.

Time = 0.13 (sec) , antiderivative size = 240, normalized size of antiderivative = 3.16

$$\int \operatorname{arctanh}(1 - d - d \operatorname{coth}(a + bx)) dx$$

$$= \frac{b^2 x^2 - bx \log\left(-\frac{d \cosh(bx+a)+d \sinh(bx+a)}{d \cosh(bx+a)+(d-2) \sinh(bx+a)}\right) + a \log(2(d-1) \cosh(bx+a) + 2(d-1) \sinh(bx+a) + \dots)}{\dots}$$

input

```
integrate(-arctanh(-1+d*d*coth(b*x+a)),x, algorithm="fricas")
```

output

```
1/2*(b^2*x^2 - b*x*log(-(d*cosh(b*x + a) + d*sinh(b*x + a))/(d*cosh(b*x +
a) + (d - 2)*sinh(b*x + a))) + a*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)*s
inh(b*x + a) + sqrt(-4*d + 4)) + a*log(2*(d - 1)*cosh(b*x + a) + 2*(d - 1)
*sinh(b*x + a) - sqrt(-4*d + 4)) - (b*x + a)*log(1/2*sqrt(-4*d + 4)*(cosh(
b*x + a) + sinh(b*x + a)) + 1) - (b*x + a)*log(-1/2*sqrt(-4*d + 4)*(cosh(b
*x + a) + sinh(b*x + a)) + 1) - dilog(1/2*sqrt(-4*d + 4)*(cosh(b*x + a) +
sinh(b*x + a))) - dilog(-1/2*sqrt(-4*d + 4)*(cosh(b*x + a) + sinh(b*x + a
))))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = - \int \operatorname{atanh}(d \coth(a + bx) + d - 1) dx$$

input

```
integrate(-atanh(-1+d+d*coth(b*x+a)),x)
```

output

```
-Integral(atanh(d*coth(a + b*x) + d - 1), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.56 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx \\ &= \frac{1}{4} bd \left( \frac{2x^2}{d} - \frac{2bx \log((d-1)e^{(2bx+2a)} + 1) + \operatorname{Li}_2(-(d-1)e^{(2bx+2a)})}{b^2 d} \right) \\ & \quad - x \operatorname{artanh}(d \coth(bx + a) + d - 1) \end{aligned}$$

input

```
integrate(-arctanh(-1+d+d*coth(b*x+a)),x, algorithm="maxima")
```

output

```
1/4*b*d*(2*x^2/d - (2*b*x*log((d - 1)*e^(2*b*x + 2*a) + 1) + dilog(-(d - 1)
)*e^(2*b*x + 2*a)))/(b^2*d) - x*arctanh(d*coth(b*x + a) + d - 1)
```



**Giac [F]**

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -\operatorname{artanh}(d \coth(bx + a) + d - 1) dx$$

input `integrate(-arctanh(-1+d+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(-arctanh(d*coth(b*x + a) + d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = \int -\operatorname{atanh}(d + d \coth(a + bx) - 1) dx$$

input `int(-atanh(d + d*coth(a + b*x) - 1),x)`

output `int(-atanh(d + d*coth(a + b*x) - 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 - d - d \coth(a + bx)) dx = -\left( \int \operatorname{atanh}(\coth(bx + a) d + d - 1) dx \right)$$

input `int(-atanh(-1+d+d*coth(b*x+a)),x)`

output `- int(atanh(coth(a + b*x)*d + d - 1),x)`

### 3.311 $\int \frac{\operatorname{arctanh}(1-d-d \coth(a+bx))}{x} dx$

Optimal result	2165
Mathematica [N/A]	2165
Rubi [N/A]	2166
Maple [N/A]	2166
Fricas [N/A]	2167
Sympy [N/A]	2167
Maxima [N/A]	2167
Giac [N/A]	2168
Mupad [N/A]	2168
Reduce [N/A]	2169

#### Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\operatorname{arctanh}(1-d-d \coth(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1-d-d \coth(a+bx))}{x}, x\right)$$

output `Defer(Int)(-arctanh(-1+d+d*coth(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 3.60 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{arctanh}(1-d-d \coth(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1-d-d \coth(a+bx))}{x} dx$$

input `Integrate[ArcTanh[1 - d - d*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 - d - d*Coth[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d(-\coth(a+bx)) - d + 1)}{x} dx$$

input `Int[ArcTanh[1 - d - d*Coth[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int -\frac{\operatorname{arctanh}(-1 + d + d \coth(bx + a))}{x} dx$$

input `int(-arctanh(-1+d+d*coth(b*x+a))/x,x)`

output `int(-arctanh(-1+d+d*coth(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \coth(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(-arctanh(d*coth(b*x + a) + d - 1)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 1.81 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = - \int \frac{\operatorname{atanh}(d \coth(a + bx) + d - 1)}{x} dx$$

input `integrate(-atanh(-1+d+d*coth(b*x+a))/x,x)`

output `-Integral(atanh(d*coth(a + b*x) + d - 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 0.70 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \coth(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*coth(b*x+a))/x,x, algorithm="maxima")`

output `-integrate(arctanh(d*coth(b*x + a) + d - 1)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \coth(bx + a) + d - 1)}{x} dx$$

input `integrate(-arctanh(-1+d+d*coth(b*x+a))/x,x, algorithm="giac")`

output `integrate(-arctanh(d*coth(b*x + a) + d - 1)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = \int -\frac{\operatorname{atanh}(d + d \coth(a + bx) - 1)}{x} dx$$

input `int(-atanh(d + d*coth(a + b*x) - 1)/x,x)`

output `int(-atanh(d + d*coth(a + b*x) - 1)/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - d - d \coth(a + bx))}{x} dx = - \left( \int \frac{\operatorname{atanh}(\coth(bx + a) d + d - 1)}{x} dx \right)$$

input `int(-atanh(-1+d+d*coth(b*x+a))/x,x)`output `- int(atanh(coth(a + b*x)*d + d - 1)/x,x)`

### 3.312 $\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx$

Optimal result	2170
Mathematica [B] (verified)	2171
Rubi [A] (verified)	2172
Maple [C] (warning: unable to verify)	2176
Fricas [B] (verification not implemented)	2177
Sympy [F]	2178
Maxima [F]	2179
Giac [F]	2179
Mupad [F(-1)]	2180
Reduce [F]	2180

#### Optimal result

Integrand size = 15, antiderivative size = 302

$$\begin{aligned}
 \int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx = & \frac{i(e + fx)^4 \arctan(e^{2i(a+bx)})}{4f} \\
 & + \frac{(e + fx)^4 \operatorname{arctanh}(\tan(a + bx))}{4f} \\
 & - \frac{i(e + fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} \\
 & + \frac{i(e + fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} \\
 & + \frac{3f(e + fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{8b^2} \\
 & - \frac{3f(e + fx)^2 \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{8b^2} \\
 & + \frac{3if^2(e + fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{8b^3} \\
 & - \frac{3if^2(e + fx) \operatorname{PolyLog}(4, ie^{2i(a+bx)})}{8b^3} \\
 & - \frac{3f^3 \operatorname{PolyLog}(5, -ie^{2i(a+bx)})}{16b^4} \\
 & + \frac{3f^3 \operatorname{PolyLog}(5, ie^{2i(a+bx)})}{16b^4}
 \end{aligned}$$

output

```
1/4*I*(f*x+e)^4*arctan(exp(2*I*(b*x+a)))/f+1/4*(f*x+e)^4*arctanh(tan(b*x+a
))/f-1/4*I*(f*x+e)^3*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)^3*poly
log(2,I*exp(2*I*(b*x+a)))/b+3/8*f*(f*x+e)^2*polylog(3,-I*exp(2*I*(b*x+a)))
/b^2-3/8*f*(f*x+e)^2*polylog(3,I*exp(2*I*(b*x+a)))/b^2+3/8*I*f^2*(f*x+e)*p
olylog(4,-I*exp(2*I*(b*x+a)))/b^3-3/8*I*f^2*(f*x+e)*polylog(4,I*exp(2*I*(b
*x+a)))/b^3-3/16*f^3*polylog(5,-I*exp(2*I*(b*x+a)))/b^4+3/16*f^3*polylog(5
,I*exp(2*I*(b*x+a)))/b^4
```

### Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 654 vs.  $2(302) = 604$ .

Time = 0.98 (sec) , antiderivative size = 654, normalized size of antiderivative = 2.17

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx$$

$$= \frac{1}{4}x(4e^3 + 6e^2fx + 4ef^2x^2 + f^3x^3) \operatorname{arctanh}(\tan(a + bx))$$

$$+ \frac{-8b^4e^3x \log(1 - ie^{2i(a+bx)}) - 12b^4e^2fx^2 \log(1 - ie^{2i(a+bx)}) - 8b^4ef^2x^3 \log(1 - ie^{2i(a+bx)}) - 2b^4f^3x^4 \log(1 - ie^{2i(a+bx)})}{4}$$

input

```
Integrate[(e + f*x)^3*ArcTanh[Tan[a + b*x]],x]
```



output

```
(x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcTanh[Tan[a + b*x]])/4 +
(-8*b^4*e^3*x*Log[1 - I*E^((2*I)*(a + b*x))] - 12*b^4*e^2*f*x^2*Log[1 - I*
E^((2*I)*(a + b*x))] - 8*b^4*e*f^2*x^3*Log[1 - I*E^((2*I)*(a + b*x))] - 2*
b^4*f^3*x^4*Log[1 - I*E^((2*I)*(a + b*x))] + 8*b^4*e^3*x*Log[1 + I*E^((2*I)
)*(a + b*x))] + 12*b^4*e^2*f*x^2*Log[1 + I*E^((2*I)*(a + b*x))] + 8*b^4*e*
f^2*x^3*Log[1 + I*E^((2*I)*(a + b*x))] + 2*b^4*f^3*x^4*Log[1 + I*E^((2*I)*
(a + b*x))] - (4*I)*b^3*(e + f*x)^3*PolyLog[2, (-I)*E^((2*I)*(a + b*x))] +
(4*I)*b^3*(e + f*x)^3*PolyLog[2, I*E^((2*I)*(a + b*x))] + 6*b^2*e^2*f*Pol
yLog[3, (-I)*E^((2*I)*(a + b*x))] + 12*b^2*e*f^2*x*PolyLog[3, (-I)*E^((2*I)
)*(a + b*x))] + 6*b^2*f^3*x^2*PolyLog[3, (-I)*E^((2*I)*(a + b*x))] - 6*b^2
*e^2*f*PolyLog[3, I*E^((2*I)*(a + b*x))] - 12*b^2*e*f^2*x*PolyLog[3, I*E^((
2*I)*(a + b*x))] - 6*b^2*f^3*x^2*PolyLog[3, I*E^((2*I)*(a + b*x))] + (6*I)
*b*e*f^2*PolyLog[4, (-I)*E^((2*I)*(a + b*x))] + (6*I)*b*f^3*x*PolyLog[4,
(-I)*E^((2*I)*(a + b*x))] - (6*I)*b*e*f^2*PolyLog[4, I*E^((2*I)*(a + b*x))
] - (6*I)*b*f^3*x*PolyLog[4, I*E^((2*I)*(a + b*x))] - 3*f^3*PolyLog[5, (-I)
)*E^((2*I)*(a + b*x))] + 3*f^3*PolyLog[5, I*E^((2*I)*(a + b*x)))]/(16*b^4)
```

### Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$ , Rules used = {6805, 3042, 4669, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) \, dx \\
 & \quad \downarrow \text{6805} \\
 & \frac{(e + fx)^4 \operatorname{arctanh}(\tan(a + bx))}{4f} - \frac{b \int (e + fx)^4 \sec(2a + 2bx) \, dx}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^4 \operatorname{arctanh}(\tan(a + bx))}{4f} - \frac{b \int (e + fx)^4 \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx}{4f} \\
 & \quad \downarrow \text{4669}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(e+fx)^4 \operatorname{arctanh}(\tan(a+bx))}{4f} \\
 & b \left( -\frac{2f \int (e+fx)^3 \log(1-ie^{2i(a+bx)}) dx}{b} + \frac{2f \int (e+fx)^3 \log(1+ie^{2i(a+bx)}) dx}{b} - \frac{i(e+fx)^4 \operatorname{arctan}(e^{2i(a+bx)})}{b} \right) \\
 & \quad \downarrow 3011 \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\tan(a+bx))}{4f} \\
 & b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)}) dx}{2b} \right)}{b} \right) \\
 & \quad \downarrow 7163 \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\tan(a+bx))}{4f} \\
 & b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{if \int (e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) dx}{b} - \frac{i(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{2b} \right)}{b} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)}) dx}{2b} \right)}{b} \right) \\
 & \quad \downarrow 7163 \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\tan(a+bx))}{4f} \\
 & b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{if \int \operatorname{PolyLog}(4, -ie^{2i(a+bx)}) dx}{2b} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} - \frac{i(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{2b} \right)}{b} \\
 & \quad \downarrow 2720
 \end{aligned}$$

$$\frac{(e + fx)^4 \operatorname{arctanh}(\tan(a + bx))}{4f} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}(4, -ie^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{i(e+fx)^2}{b}$$

7143

$$\frac{(e + fx)^4 \operatorname{arctanh}(\tan(a + bx))}{4f} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{f \operatorname{PolyLog}(5, -ie^{2i(a+bx)})}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{i(e+fx)^4 \operatorname{arctan}(e^{2i(a+bx)})}{b} + \frac{i(e+fx)^2}{b}$$

input `Int[(e + f*x)^3*ArcTanh[Tan[a + b*x]],x]`

output

```
((e + f*x)^4*ArcTanh[Tan[a + b*x]]/(4*f) - (b*((( -I)*(e + f*x)^4*ArcTan[E
^((2*I)*(a + b*x))])/b + (2*f*(((I/2)*(e + f*x)^3*PolyLog[2, (-I)*E^((2*I)
*(a + b*x))])/b - (((3*I)/2)*f*((( -1/2*I)*(e + f*x)^2*PolyLog[3, (-I)*E^((
2*I)*(a + b*x))])/b + (I*f*((( -1/2*I)*(e + f*x)*PolyLog[4, (-I)*E^((2*I)*
(a + b*x))])/b + (f*PolyLog[5, (-I)*E^((2*I)*(a + b*x))]/(4*b^2))/b))/b))
/b - (2*f*(((I/2)*(e + f*x)^3*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (((3*
I)/2)*f*((( -1/2*I)*(e + f*x)^2*PolyLog[3, I*E^((2*I)*(a + b*x))])/b + (I*f
*((( -1/2*I)*(e + f*x)*PolyLog[4, I*E^((2*I)*(a + b*x))])/b + (f*PolyLog[5,
I*E^((2*I)*(a + b*x))]/(4*b^2))/b))/b))/b)/(4*f)
```

### Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6805 `Int[ArcTanh[Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTanh[Tan[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 16.67 (sec) , antiderivative size = 3640, normalized size of antiderivative = 12.05

method	result	size
risch	Expression too large to display	3640

input `int((f*x+e)^3*arctanh(tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

-3/16*f^3*polylog(5,-I*exp(2*I*(b*x+a)))/b^4+3/16*f^3*polylog(5,I*exp(2*I*
(b*x+a)))/b^4+3/2*f/b*e^2*ln(I*exp(2*I*(b*x+a))+1)*x*a-3/2*f^2/b^2*e*ln(I*
exp(2*I*(b*x+a))+1)*x*a^2-3/2*f/b*a*e^2*ln(1+exp(I*(b*x+a)))*(-1)^(3/4))*x-
3/2*f/b*a*e^2*ln(1-exp(I*(b*x+a)))*(-1)^(3/4))*x+3/2*f^2/b^2*a^2*e*ln(1+exp
(I*(b*x+a)))*(-1)^(3/4))*x-3/4*I*f/b^2*e^2*polylog(2,-I*exp(2*I*(b*x+a)))*a
+3/4*I*f^2/b*e*polylog(2,I*exp(2*I*(b*x+a)))*x^2-3/2*I*f/b^2*e^2*a*dilog((
(-I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))-3/2*I*f/b^2*e^2*a*dilog(((I)^(1/2)
+exp(I*(b*x+a)))/(-I)^(1/2))-3/2*I*f^2/b^3*a^2*e*dilog(1+exp(I*(b*x+a)))*(-
1)^(3/4))-3/2*I*f^2/b^3*a^2*e*dilog(1-exp(I*(b*x+a)))*(-1)^(3/4))-1/2*f^2*ln
(exp(2*I*(b*x+a))-I)*x^3*e+3/2*f/b*e^2*a*ln(((I)^(1/2)+exp(I*(b*x+a)))/(-
I)^(1/2))*x-3/2*f/b*e^2*ln(-I*exp(2*I*(b*x+a))+1)*x*a+3/2*f^2/b^2*e*a^2*ln
(-I*exp(2*I*(b*x+a))+1)*x-3/2*f^2/b^2*e*a^2*ln(((I)^(1/2)-exp(I*(b*x+a))
)/(-I)^(1/2))*x-3/2*f^2/b^2*e*a^2*ln(((I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2
))*x+3/2*f^2/b^2*a^2*e*ln(1-exp(I*(b*x+a)))*(-1)^(3/4))*x+3/4*I*f/b^2*e^2*p
olylog(2,I*exp(2*I*(b*x+a)))*a+3/2*I*f^2/b^3*e*a^2*dilog(((I)^(1/2)-exp(I
*(b*x+a)))/(-I)^(1/2))+3/2*I*f^2/b^3*e*a^2*dilog(((I)^(1/2)+exp(I*(b*x+a)
))/(-I)^(1/2))-3/4*I*f^2/b^3*e*a^2*polylog(2,I*exp(2*I*(b*x+a)))-3/4*f*ln(
exp(2*I*(b*x+a))-I)*x^2*e^2+3/8*f/b^2*e^2*polylog(3,-I*exp(2*I*(b*x+a)))-3
/8*f^3/b^2*polylog(3,I*exp(2*I*(b*x+a)))*x^2-3/8*f/b^2*e^2*polylog(3,I*exp
(2*I*(b*x+a)))-1/2*f^3/b^4*a^4*ln(1+exp(I*(b*x+a)))*(-1)^(3/4))-1/2*f^3/...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1809 vs.  $2(236) = 472$ .

Time = 0.15 (sec) , antiderivative size = 1809, normalized size of antiderivative = 5.99

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(ax + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^3*arctanh(tan(b*x+a)),x, algorithm="fricas")
```

output

```

-1/32*(3*f^3*polylog(5, (I*tan(b*x + a)^2 + 2*tan(b*x + a) - I)/(tan(b*x +
a)^2 + 1)) - 3*f^3*polylog(5, (I*tan(b*x + a)^2 - 2*tan(b*x + a) - I)/(ta
n(b*x + a)^2 + 1)) + 3*f^3*polylog(5, (-I*tan(b*x + a)^2 + 2*tan(b*x + a)
+ I)/(tan(b*x + a)^2 + 1)) - 3*f^3*polylog(5, (-I*tan(b*x + a)^2 - 2*tan(b
*x + a) + I)/(tan(b*x + a)^2 + 1)) + 4*(-I*b^3*f^3*x^3 - 3*I*b^3*e*f^2*x^2
- 3*I*b^3*e^2*f*x - I*b^3*e^3)*dilog(-((I + 1)*tan(b*x + a)^2 + 2*tan(b*x
+ a) - I + 1)/(tan(b*x + a)^2 + 1) + 1) + 4*(-I*b^3*f^3*x^3 - 3*I*b^3*e*f
^2*x^2 - 3*I*b^3*e^2*f*x - I*b^3*e^3)*dilog(-((I + 1)*tan(b*x + a)^2 - 2*t
an(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1) + 1) + 4*(I*b^3*f^3*x^3 + 3*I*b^
3*e*f^2*x^2 + 3*I*b^3*e^2*f*x + I*b^3*e^3)*dilog(-(-(I - 1)*tan(b*x + a)^2
+ 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) + 4*(I*b^3*f^3*x^3 +
3*I*b^3*e*f^2*x^2 + 3*I*b^3*e^2*f*x + I*b^3*e^3)*dilog(-(-(I - 1)*tan(b*x
+ a)^2 - 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) + 2*(b^4*f^3*x^
4 + 4*b^4*e*f^2*x^3 + 6*b^4*e^2*f*x^2 + 4*b^4*e^3*x + 4*a*b^3*e^3 - 6*a^2*
b^2*e^2*f + 4*a^3*b*e*f^2 - a^4*f^3)*log(((I + 1)*tan(b*x + a)^2 + 2*tan(b
*x + a) - I + 1)/(tan(b*x + a)^2 + 1)) - 2*(4*a*b^3*e^3 - 6*a^2*b^2*e^2*f
+ 4*a^3*b*e*f^2 - a^4*f^3)*log(((I + 1)*tan(b*x + a)^2 + 2*I*tan(b*x + a)
+ I - 1)/(tan(b*x + a)^2 + 1)) + 2*(4*a*b^3*e^3 - 6*a^2*b^2*e^2*f + 4*a^3*
b*e*f^2 - a^4*f^3)*log(((I + 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I - 1)
/(tan(b*x + a)^2 + 1)) - 2*(b^4*f^3*x^4 + 4*b^4*e*f^2*x^3 + 6*b^4*e^2*f...

```

### Sympy [F]

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx = \int (e + fx)^3 \operatorname{atanh}(\tan(a + bx)) dx$$

input

```
integrate((f*x+e)**3*atanh(tan(b*x+a)),x)
```

output

```
Integral((e + f*x)**3*atanh(tan(a + b*x)), x)
```

**Maxima [F]**

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e)^3 \operatorname{arctanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)^3*arctanh(tan(b*x+a)),x, algorithm="maxima")`

output `1/16*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/16*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(1/2*((b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

**Giac [F]**

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e)^3 \operatorname{arctanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)^3*arctanh(tan(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)^3*arctanh(tan(b*x + a)), x)`



**Mupad [F(-1)]**

Timed out.

$$\int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(a + bx)) (e + fx)^3 dx$$

input `int(atanh(tan(a + b*x))*(e + f*x)^3,x)`

output `int(atanh(tan(a + b*x))*(e + f*x)^3, x)`

**Reduce [F]**

$$\begin{aligned} \int (e + fx)^3 \operatorname{arctanh}(\tan(a + bx)) dx &= \left( \int \operatorname{atanh}(\tan(bx + a)) dx \right) e^3 \\ &+ \left( \int \operatorname{atanh}(\tan(bx + a)) x^3 dx \right) f^3 \\ &+ 3 \left( \int \operatorname{atanh}(\tan(bx + a)) x^2 dx \right) e f^2 \\ &+ 3 \left( \int \operatorname{atanh}(\tan(bx + a)) x dx \right) e^2 f \end{aligned}$$

input `int((f*x+e)^3*atanh(tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)),x)*e**3 + int(atanh(tan(a + b*x))*x**3,x)*f**3 + 3  
*int(atanh(tan(a + b*x))*x**2,x)*e*f**2 + 3*int(atanh(tan(a + b*x))*x,x)*e  
**2*f`

### 3.313 $\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx$

Optimal result	2181
Mathematica [A] (verified)	2182
Rubi [A] (verified)	2183
Maple [C] (warning: unable to verify)	2186
Fricas [B] (verification not implemented)	2187
Sympy [F]	2188
Maxima [F]	2189
Giac [F]	2189
Mupad [F(-1)]	2189
Reduce [F]	2190

#### Optimal result

Integrand size = 15, antiderivative size = 234

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \frac{i(e + fx)^3 \arctan(e^{2i(a+bx)})}{3f} + \frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{i(e + fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} + \frac{i(e + fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} + \frac{f(e + fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{4b^2} - \frac{f(e + fx) \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{4b^2} + \frac{if^2 \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{8b^3} - \frac{if^2 \operatorname{PolyLog}(4, ie^{2i(a+bx)})}{8b^3}$$

output

```
1/3*I*(f*x+e)^3*arctan(exp(2*I*(b*x+a)))/f+1/3*(f*x+e)^3*arctanh(tan(b*x+a
))/f-1/4*I*(f*x+e)^2*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)^2*poly
log(2,I*exp(2*I*(b*x+a)))/b+1/4*f*(f*x+e)*polylog(3,-I*exp(2*I*(b*x+a)))/b
^2-1/4*f*(f*x+e)*polylog(3,I*exp(2*I*(b*x+a)))/b^2+1/8*I*f^2*polylog(4,-I*
exp(2*I*(b*x+a)))/b^3-1/8*I*f^2*polylog(4,I*exp(2*I*(b*x+a)))/b^3
```

**Mathematica [A] (verified)**

Time = 0.51 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.75

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \frac{1}{3}x(3e^2 + 3efx + f^2x^2) \operatorname{arctanh}(\tan(a + bx)) \\ + \frac{-12b^3e^2x \log(1 - ie^{2i(a+bx)}) - 12b^3efx^2 \log(1 - ie^{2i(a+bx)}) - 4b^3f^2x^3 \log(1 - ie^{2i(a+bx)}) + 12b^3e^2x \log(1 + ie^{2i(a+bx)}) + 12b^3efx^2 \log(1 + ie^{2i(a+bx)}) + 4b^3f^2x^3 \log(1 + ie^{2i(a+bx)})}{24b^3}$$

input

```
Integrate[(e + f*x)^2*ArcTanh[Tan[a + b*x]],x]
```

output

```
(x*(3*e^2 + 3*e*f*x + f^2*x^2)*ArcTanh[Tan[a + b*x]])/3 + (-12*b^3*e^2*x*L
og[1 - I*E^((2*I)*(a + b*x))] - 12*b^3*e*f*x^2*Log[1 - I*E^((2*I)*(a + b*x
))] - 4*b^3*f^2*x^3*Log[1 - I*E^((2*I)*(a + b*x))] + 12*b^3*e^2*x*Log[1 +
I*E^((2*I)*(a + b*x))] + 12*b^3*e*f*x^2*Log[1 + I*E^((2*I)*(a + b*x))] + 4
*b^3*f^2*x^3*Log[1 + I*E^((2*I)*(a + b*x))] - (6*I)*b^2*(e + f*x)^2*PolyLo
g[2, (-I)*E^((2*I)*(a + b*x))] + (6*I)*b^2*(e + f*x)^2*PolyLog[2, I*E^((2*
I)*(a + b*x))] + 6*b*e*f*PolyLog[3, (-I)*E^((2*I)*(a + b*x))] + 6*b*f^2*x*
PolyLog[3, (-I)*E^((2*I)*(a + b*x))] - 6*b*e*f*PolyLog[3, I*E^((2*I)*(a +
b*x))] - 6*b*f^2*x*PolyLog[3, I*E^((2*I)*(a + b*x))] + (3*I)*f^2*PolyLog[4
, (-I)*E^((2*I)*(a + b*x))] - (3*I)*f^2*PolyLog[4, I*E^((2*I)*(a + b*x))])
/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.85 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.16, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {6805, 3042, 4669, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) \, dx \\
 & \quad \downarrow \text{6805} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{b \int (e + fx)^3 \sec(2a + 2bx) \, dx}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{b \int (e + fx)^3 \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx}{3f} \\
 & \quad \downarrow \text{4669} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \\
 & \frac{b \left( -\frac{3f \int (e + fx)^2 \log(1 - ie^{2i(a + bx)}) \, dx}{2b} + \frac{3f \int (e + fx)^2 \log(1 + ie^{2i(a + bx)}) \, dx}{2b} - \frac{i(e + fx)^3 \operatorname{arctan}(e^{2i(a + bx)})}{b} \right)}{3f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \\
 & b \left( \frac{3f \left( \frac{i(e + fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a + bx)})}{2b} - \frac{if \int (e + fx) \operatorname{PolyLog}(2, -ie^{2i(a + bx)}) \, dx}{b} \right)}{2b} - \frac{3f \left( \frac{i(e + fx)^2 \operatorname{PolyLog}(2, ie^{2i(a + bx)})}{2b} - \frac{if \int (e + fx) \operatorname{PolyLog}(2, ie^{2i(a + bx)}) \, dx}{b} \right)}{2b} \right) \\
 & \quad \downarrow \text{7163}
 \end{aligned}$$

$$\frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{b \left( \frac{3f \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{if \int \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) dx}{2b} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right)}{3f} - \frac{3f \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} \right)}{3f}$$

↓ 2720

$$\frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{b \left( \frac{3f \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right)}{3f} - \frac{3f \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} \right)}{3f}$$

↓ 7143

$$\frac{(e + fx)^3 \operatorname{arctanh}(\tan(a + bx))}{3f} - \frac{b \left( \frac{3f \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{f \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right)}{3f} + \frac{i(e+fx)^3 \operatorname{arctan}(e^{2i(a+bx)})}{b}$$

input

```
Int[(e + f*x)^2*ArcTanh[Tan[a + b*x]], x]
```

output

```
((e + f*x)^3*ArcTanh[Tan[a + b*x]])/(3*f) - (b*((( -I)*(e + f*x)^3*ArcTan[E
^((2*I)*(a + b*x))])/b + (3*f*(((I/2)*(e + f*x)^2*PolyLog[2, (-I)*E^((2*I)
*(a + b*x))])/b - (I*f*((( -1/2*I)*(e + f*x)*PolyLog[3, (-I)*E^((2*I)*(a +
b*x))])/b + (f*PolyLog[4, (-I)*E^((2*I)*(a + b*x))])/(4*b^2))/b))/(2*b) -
(3*f*(((I/2)*(e + f*x)^2*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (I*f*((( -
1/2*I)*(e + f*x)*PolyLog[3, I*E^((2*I)*(a + b*x))])/b + (f*PolyLog[4, I*E^
((2*I)*(a + b*x))])/(4*b^2))/b))/(2*b)))/(3*f)
```

### Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6805 `Int[ArcTanh[Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTanh[Tan[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 9.74 (sec) , antiderivative size = 2719, normalized size of antiderivative = 11.62

method	result	size
risch	Expression too large to display	2719

input `int((f*x+e)^2*arctanh(tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

1/2*I*f*e/b*polylog(2,I*exp(2*I*(b*x+a)))*x+1/2*I*f*e/b^2*polylog(2,I*exp(
2*I*(b*x+a)))*a-I*f/b^2*a*e*dilog((( -I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))-
I*f/b^2*a*e*dilog((( -I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))-1/2*f*e*ln(-I*ex
p(2*I*(b*x+a))+1)*x^2-1/4*f^2/b^2*polylog(3,I*exp(2*I*(b*x+a)))*x+1/3*f^2/
b^3*ln(-I*exp(2*I*(b*x+a))+1)*a^3-1/2*f^2*ln((( -I)^(1/2)-exp(I*(b*x+a)))/(-
I)^(1/2))/b^3*a^3-1/2*f^2*ln((( -I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))/b^3*
a^3+1/6*f^2/b^3*a^3*ln(exp(2*I*(b*x+a))+I)-1/4*f*e/b^2*polylog(3,I*exp(2*I
*(b*x+a)))-1/2*ln((( -I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))/b*a*e^2-1/2*ln((
(-I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))/b*a*e^2+1/2/b*a*e^2*ln(exp(2*I*(b*x
+a))+I)+1/2*I/b*e^2*dilog((( -I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))+1/2*I/b*
e^2*dilog((( -I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))+1/12*I*Pi*(csgn(I*(exp(2
*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))*csgn((1-I)*(exp(2*I*(b*x+a))-I)/(exp(
2*I*(b*x+a))+1))+csgn((1-I)*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2+c
sgn(I*(exp(2*I*(b*x+a))-I))*csgn(I*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+
1))*csgn(I/(exp(2*I*(b*x+a))+1))-csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(
2*I*(b*x+a))+I))*csgn(I*(exp(2*I*(b*x+a))+I)/(exp(2*I*(b*x+a))+1))-csgn(I*
(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2*csgn(I/(exp(2*I*(b*x+a))+1))+
csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))+I)/(exp(2*I*(b*x+a))
+1))^2-csgn(I*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2*csgn(I*(exp(2*I
*(b*x+a))-I))+csgn(I*(exp(2*I*(b*x+a))+I))*csgn(I*(exp(2*I*(b*x+a))+I)/...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1283 vs.  $2(180) = 360$ .

Time = 0.14 (sec) , antiderivative size = 1283, normalized size of antiderivative = 5.48

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(ax + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^2*arctanh(tan(b*x+a)),x, algorithm="fricas")
```



output

```

1/48*(3*I*f^2*polylog(4, (I*tan(b*x + a)^2 + 2*tan(b*x + a) - I)/(tan(b*x
+ a)^2 + 1)) + 3*I*f^2*polylog(4, (I*tan(b*x + a)^2 - 2*tan(b*x + a) - I)/
(tan(b*x + a)^2 + 1)) - 3*I*f^2*polylog(4, (-I*tan(b*x + a)^2 + 2*tan(b*x
+ a) + I)/(tan(b*x + a)^2 + 1)) - 3*I*f^2*polylog(4, (-I*tan(b*x + a)^2 -
2*tan(b*x + a) + I)/(tan(b*x + a)^2 + 1)) - 6*(-I*b^2*f^2*x^2 - 2*I*b^2*e*
f*x - I*b^2*e^2)*dilog(-((I + 1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/
(tan(b*x + a)^2 + 1) + 1) - 6*(-I*b^2*f^2*x^2 - 2*I*b^2*e*f*x - I*b^2*e^2)
*dilog(-((I + 1)*tan(b*x + a)^2 - 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2
+ 1) + 1) - 6*(I*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(-(-(I - 1)
*tan(b*x + a)^2 + 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) - 6*(I
*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(-(-(I - 1)*tan(b*x + a)^2
- 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) - 4*(b^3*f^2*x^3 + 3*b
^3*e*f*x^2 + 3*b^3*e^2*x + 3*a*b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(((I +
1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1)) + 4*(3*a
*b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(((I + 1)*tan(b*x + a)^2 + 2*I*tan(b*
x + a) + I - 1)/(tan(b*x + a)^2 + 1)) - 4*(3*a*b^2*e^2 - 3*a^2*b*e*f + a^3
*f^2)*log(((I + 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I - 1)/(tan(b*x + a
)^2 + 1)) + 4*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b^3*e^2*x + 3*a*b^2*e^2 - 3
*a^2*b*e*f + a^3*f^2)*log(((I + 1)*tan(b*x + a)^2 - 2*tan(b*x + a) - I + 1
)/(tan(b*x + a)^2 + 1)) - 4*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b^3*e^2*x ...

```

### Sympy [F]

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \int (e + fx)^2 \operatorname{atanh}(\tan(a + bx)) dx$$

input

```
integrate((f*x+e)**2*atanh(tan(b*x+a)), x)
```

output

```
Integral((e + f*x)**2*atanh(tan(a + b*x)), x)
```

**Maxima [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e)^2 \operatorname{artanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)^2*arctanh(tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/12*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(2/3*((b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

**Giac [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e)^2 \operatorname{artanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)^2*arctanh(tan(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)^2*arctanh(tan(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(a + bx)) (e + fx)^2 dx$$

input `int(atanh(tan(a + b*x))*(e + f*x)^2,x)`

output `int(atanh(tan(a + b*x))*(e + f*x)^2, x)`

**Reduce [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\tan(a + bx)) dx = \left( \int \operatorname{atanh}(\tan(bx + a)) dx \right) e^2$$

$$+ \left( \int \operatorname{atanh}(\tan(bx + a)) x^2 dx \right) f^2$$

$$+ 2 \left( \int \operatorname{atanh}(\tan(bx + a)) x dx \right) ef$$

input `int((f*x+e)^2*atanh(tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)),x)*e**2 + int(atanh(tan(a + b*x))*x**2,x)*f**2 + 2  
*int(atanh(tan(a + b*x))*x,x)*e*f`

### 3.314 $\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx$

Optimal result	2191
Mathematica [A] (verified)	2192
Rubi [A] (verified)	2192
Maple [C] (warning: unable to verify)	2195
Fricas [B] (verification not implemented)	2196
Sympy [F]	2197
Maxima [F]	2197
Giac [F]	2197
Mupad [F(-1)]	2198
Reduce [F]	2198

#### Optimal result

Integrand size = 13, antiderivative size = 162

$$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx = \frac{i(e + fx)^2 \arctan(e^{2i(a+bx)})}{2f} + \frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} + \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} + \frac{f \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{8b^2} - \frac{f \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{8b^2}$$

output `1/2*I*(f*x+e)^2*arctan(exp(2*I*(b*x+a)))/f+1/2*(f*x+e)^2*arctanh(tan(b*x+a))/f-1/4*I*(f*x+e)*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)*polylog(2,I*exp(2*I*(b*x+a)))/b+1/8*f*polylog(3,-I*exp(2*I*(b*x+a)))/b^2-1/8*f*polylog(3,I*exp(2*I*(b*x+a)))/b^2`

**Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.82

$$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx = ex \operatorname{arctanh}(\tan(a + bx)) + \frac{1}{2} f x^2 \operatorname{arctanh}(\tan(a + bx)) - \frac{e((-4a + \pi - 4bx) (\log(1 - ie^{-2i(a+bx)}) - \log(1 + ie^{-2i(a+bx)})) - (-4a + \pi) \log(\cot(a + \frac{\pi}{4} + bx)))}{8b} + \frac{f(4ib^2 x^2 \arctan(\cos(2(a + bx)) + i \sin(2(a + bx))) + 2ibx \operatorname{PolyLog}(2, i \cos(2(a + bx)) - \sin(2(a + bx)))}{8b}$$

input `Integrate[(e + f*x)*ArcTanh[Tan[a + b*x]], x]`

output `e*x*ArcTanh[Tan[a + b*x]] + (f*x^2*ArcTanh[Tan[a + b*x]])/2 - (e*((-4*a + Pi - 4*b*x)*(Log[1 - I/E^((2*I)*(a + b*x))] - Log[1 + I/E^((2*I)*(a + b*x))]) - (-4*a + Pi)*Log[Cot[a + Pi/4 + b*x]] + (2*I)*(PolyLog[2, (-I)/E^((2*I)*(a + b*x))] - PolyLog[2, I/E^((2*I)*(a + b*x))])))/(8*b) + (f*((4*I)*b^2*x^2*ArcTan[Cos[2*(a + b*x)] + I*Sin[2*(a + b*x)]] + (2*I)*b*x*PolyLog[2, I*Cos[2*(a + b*x)] - Sin[2*(a + b*x)]] - (2*I)*b*x*PolyLog[2, (-I)*Cos[2*(a + b*x)] + Sin[2*(a + b*x)]] - PolyLog[3, I*Cos[2*(a + b*x)] - Sin[2*(a + b*x)]] + PolyLog[3, (-I)*Cos[2*(a + b*x)] + Sin[2*(a + b*x)]]))/(8*b^2)`

**Rubi [A] (verified)**

Time = 0.60 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {6805, 3042, 4669, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx$$

$$\downarrow 6805$$

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \int (e + fx)^2 \sec(2a + 2bx) dx}{2f}$$

$$\downarrow 3042$$

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \int (e + fx)^2 \csc\left(2a + 2bx + \frac{\pi}{2}\right) dx}{2f}$$

↓ 4669

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \left( -\frac{f \int (e + fx) \log(1 - ie^{2i(a+bx)}) dx}{b} + \frac{f \int (e + fx) \log(1 + ie^{2i(a+bx)}) dx}{b} - \frac{i(e + fx)^2 \operatorname{arctan}(e^{2i(a+bx)})}{b} \right)}{2f}$$

↓ 3011

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \left( \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \int \operatorname{PolyLog}(2, -ie^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{if \int \operatorname{PolyLog}(2, ie^{2i(a+bx)}) dx}{2b} \right)}{b} \right)}{2f}$$


---

↓ 2720

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \left( \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -ie^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{f \int e^{-2i(a+bx)}}{b} \right)}{b} \right)}{2f}$$


---

↓ 7143

$$\frac{(e + fx)^2 \operatorname{arctanh}(\tan(a + bx))}{2f} - \frac{b \left( -\frac{i(e + fx)^2 \operatorname{arctan}(e^{2i(a+bx)})}{b} + \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{f \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{f \int e^{-2i(a+bx)}}{b} \right)}{b} \right)}{2f}$$

input

```
Int[(e + f*x)*ArcTanh[Tan[a + b*x]], x]
```

output

```
((e + f*x)^2*ArcTanh[Tan[a + b*x]])/(2*f) - (b*((( -I)*(e + f*x)^2*ArcTan[E
^((2*I)*(a + b*x))])/b + (f*(((I/2)*(e + f*x)*PolyLog[2, (-I)*E^((2*I)*(a
+ b*x))])/b - (f*PolyLog[3, (-I)*E^((2*I)*(a + b*x))])/(4*b^2)))/b - (f*((
(I/2)*(e + f*x)*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (f*PolyLog[3, I*E^((
2*I)*(a + b*x))])/(4*b^2)))/b))/(2*f)
```

### Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] :=> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6805

```
Int[ArcTanh[Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:=> Simp[(e + f*x)^(m + 1)*(ArcTanh[Tan[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.46 (sec) , antiderivative size = 1818, normalized size of antiderivative = 11.22

method	result	size
risch	Expression too large to display	1818

input

```
int((f*x+e)*arctanh(tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*f/b^2*a^2*ln(1+exp(I*(b*x+a)))*(-1)^(3/4))-1/2*f/b^2*a^2*ln(1-exp(I*(b*x+a)))*(-1)^(3/4))-1/2*e/b*ln((-I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))*a-1/2*e/b*ln((-I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))*a+1/2*I*e/b*dilog((-I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))+1/2*I*e/b*dilog((-I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))+1/2*f/b^2*a^2*ln((-I)^(1/2)-exp(I*(b*x+a)))/(-I)^(1/2))+1/2*f/b^2*a^2*ln((-I)^(1/2)+exp(I*(b*x+a)))/(-I)^(1/2))-1/4/b^2*f*a^2*ln(exp(2*I*(b*x+a))+I)+1/2/b*a*e*ln(exp(2*I*(b*x+a))+I)+1/2*(1/2*f*x^2+e*x)*ln(exp(2*I*(b*x+a))+I)-1/4*ln(exp(2*I*(b*x+a))-I)*f*x^2-1/2*ln(exp(2*I*(b*x+a))-I)*e*x+1/2*e/b*ln(1+exp(I*(b*x+a)))*(-1)^(3/4))*a+1/2*e/b*ln(1-exp(I*(b*x+a)))*(-1)^(3/4))*a-1/2*I*e/b*dilog(1+exp(I*(b*x+a)))*(-1)^(3/4))-1/2*I*e/b*dilog(1-exp(I*(b*x+a)))*(-1)^(3/4))+1/4/b^2*f*ln(I*exp(2*I*(b*x+a))+1)*a^2+1/8*f*polylog(3,-I*exp(2*I*(b*x+a)))/b^2-1/8*f*polylog(3,I*exp(2*I*(b*x+a)))/b^2-1/4/b^2*f*ln(-I*exp(2*I*(b*x+a))+1)*a^2-1/2*e/b*a*ln(-exp(2*I*(b*x+a))+I)+1/4*f/b^2*a^2*ln(-exp(2*I*(b*x+a))+I)+1/4*I*Pi*(csgn(I*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))*csgn((1-I)*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2+csgn(I*(exp(2*I*(b*x+a))-I))*csgn(I*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))*csgn(I/(exp(2*I*(b*x+a))+1))-csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))+I)/(exp(2*I*(b*x+a))+1))-csgn(I*(exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2*csgn(I/(exp(2*I*(b*x+a))+1))+cs...
```



**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 835 vs.  $2(130) = 260$ .

Time = 0.13 (sec) , antiderivative size = 835, normalized size of antiderivative = 5.15

$$\int (e + fx)\operatorname{arctanh}(\tan(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)*arctanh(tan(b*x+a)),x, algorithm="fricas")`

output

```
-1/16*(2*(-I*b*f*x - I*b*e)*dilog(-((I + 1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1) + 1) + 2*(-I*b*f*x - I*b*e)*dilog(-((I + 1)*tan(b*x + a)^2 - 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1) + 1) + 2*(I*b*f*x + I*b*e)*dilog(-((I - 1)*tan(b*x + a)^2 + 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) + 2*(I*b*f*x + I*b*e)*dilog(-((I - 1)*tan(b*x + a)^2 - 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) + 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log(((I + 1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1)) - 2*(2*a*b*e - a^2*f)*log(((I + 1)*tan(b*x + a)^2 + 2*I*tan(b*x + a) + I - 1)/(tan(b*x + a)^2 + 1)) + 2*(2*a*b*e - a^2*f)*log(((I + 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I - 1)/(tan(b*x + a)^2 + 1)) - 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log(((I + 1)*tan(b*x + a)^2 - 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1)) + 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log((- (I - 1)*tan(b*x + a)^2 + 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) - 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log((- (I - 1)*tan(b*x + a)^2 - 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) - 2*(2*a*b*e - a^2*f)*log(((I - 1)*tan(b*x + a)^2 + 2*I*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) + 2*(2*a*b*e - a^2*f)*log(((I - 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) - 4*(b^2*f*x^2 + 2*b^2*e*x)*log(- (tan(b*x + a) + 1)/(tan(b*x + a) - 1)) - f*polylog(3, (I*tan(b*x + a)^2 + 2*tan(b*x + a) - I)/(tan(b*x + a)^2 + 1)) + f*polyl...
```

**Sympy [F]**

$$\int (e + fx)\operatorname{arctanh}(\tan(a + bx)) dx = \int (e + fx) \operatorname{atanh}(\tan(a + bx)) dx$$

input `integrate((f*x+e)*atanh(tan(b*x+a)),x)`

output `Integral((e + f*x)*atanh(tan(a + b*x)), x)`

**Maxima [F]**

$$\int (e + fx)\operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e) \operatorname{artanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)*arctanh(tan(b*x+a)),x, algorithm="maxima")`

output `1/8*(f*x^2 + 2*e*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/8*(f*x^2 + 2*e*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(((b*f*x^2 + 2*b*e*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f*x^2 + 2*b*e*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f*x^2 + 2*b*e*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

**Giac [F]**

$$\int (e + fx)\operatorname{arctanh}(\tan(a + bx)) dx = \int (fx + e) \operatorname{artanh}(\tan(bx + a)) dx$$

input `integrate((f*x+e)*arctanh(tan(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)*arctanh(tan(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(a + bx)) (e + fx) dx$$

input `int(atanh(tan(a + b*x))*(e + f*x),x)`output `int(atanh(tan(a + b*x))*(e + f*x), x)`**Reduce [F]**

$$\int (e + fx) \operatorname{arctanh}(\tan(a + bx)) dx = \left( \int \operatorname{atanh}(\tan(bx + a)) dx \right) e + \left( \int \operatorname{atanh}(\tan(bx + a)) x dx \right) f$$

input `int((f*x+e)*atanh(tan(b*x+a)),x)`output `int(atanh(tan(a + b*x)),x)*e + int(atanh(tan(a + b*x))*x,x)*f`

### 3.315 $\int \operatorname{arctanh}(\tan(a + bx)) dx$

Optimal result . . . . .	2199
Mathematica [A] (verified) . . . . .	2199
Rubi [A] (verified) . . . . .	2200
Maple [A] (verified) . . . . .	2202
Fricas [B] (verification not implemented) . . . . .	2202
Sympy [F] . . . . .	2203
Maxima [B] (verification not implemented) . . . . .	2203
Giac [F] . . . . .	2204
Mupad [F(-1)] . . . . .	2204
Reduce [F] . . . . .	2205

#### Optimal result

Integrand size = 7, antiderivative size = 79

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = ix \arctan(e^{2i(a+bx)}) + x \operatorname{arctanh}(\tan(a + bx)) - \frac{i \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} + \frac{i \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b}$$

output I\*x\*arctan(exp(2\*I\*(b\*x+a)))+x\*arctanh(tan(b\*x+a))-1/4\*I\*polylog(2,-I\*exp(2\*I\*(b\*x+a)))/b+1/4\*I\*polylog(2,I\*exp(2\*I\*(b\*x+a)))/b

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.61

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = x \operatorname{arctanh}(\tan(a + bx)) - \frac{(-4a + \pi - 4bx) (\log(1 - ie^{-2i(a+bx)}) - \log(1 + ie^{-2i(a+bx)})) - (-4a + \pi) \log(\cot(a + \frac{\pi}{4} + bx))}{8b}$$

input Integrate[ArcTanh[Tan[a + b\*x]], x]

output

```
x*ArcTanh[Tan[a + b*x]] - ((-4*a + Pi - 4*b*x)*(Log[1 - I/E^((2*I)*(a + b*x))] - Log[1 + I/E^((2*I)*(a + b*x))]) - (-4*a + Pi)*Log[Cot[a + Pi/4 + b*x]] + (2*I)*(PolyLog[2, (-I)/E^((2*I)*(a + b*x))] - PolyLog[2, I/E^((2*I)*(a + b*x))]))/(8*b)
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6801, 3042, 4669, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arctanh}(\tan(a + bx)) \, dx \\
 & \quad \downarrow \text{6801} \\
 & x \operatorname{arctanh}(\tan(a + bx)) - b \int x \sec(2a + 2bx) \, dx \\
 & \quad \downarrow \text{3042} \\
 & x \operatorname{arctanh}(\tan(a + bx)) - b \int x \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx \\
 & \quad \downarrow \text{4669} \\
 & x \operatorname{arctanh}(\tan(a + bx)) - \\
 & b \left( -\frac{\int \log(1 - ie^{2i(a+bx)}) \, dx}{2b} + \frac{\int \log(1 + ie^{2i(a+bx)}) \, dx}{2b} - \frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} \right) \\
 & \quad \downarrow \text{2715} \\
 & b \left( \frac{i \int e^{-2i(a+bx)} \log(1 - ie^{2i(a+bx)}) \, de^{2i(a+bx)}}{4b^2} - \frac{i \int e^{-2i(a+bx)} \log(1 + ie^{2i(a+bx)}) \, de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} \right) \\
 & \quad \downarrow \text{2838} \\
 & b \left( -\frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} + \frac{x \operatorname{arctanh}(\tan(a + bx)) - i \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b^2} - \frac{i \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b^2} \right)
 \end{aligned}$$

input `Int[ArcTanh[Tan[a + b*x]],x]`

output `x*ArcTanh[Tan[a + b*x]] - b*((( -I)*x*ArcTan[E^((2*I)*(a + b*x))])/b + ((I/4)*PolyLog[2, (-I)*E^((2*I)*(a + b*x))])/b^2 - ((I/4)*PolyLog[2, I*E^((2*I)*(a + b*x))])/b^2)`

### Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4669 `Int[csc[(e_) + Pi*(k_) + (f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol]
:> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6801 `Int[ArcTanh[Tan[(a_) + (b_)*(x_)]], x_Symbol] :> Simp[x*ArcTanh[Tan[a + b*x]], x] - Simp[b Int[x*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b}, x]`

### Maple [A] (verified)

Time = 1.77 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.52

method	result
parts	$x \operatorname{arctanh}(\tan(bx + a)) - \frac{(bx+a) \ln(i e^{2i(bx+a)} + 1)}{2} + \frac{(bx+a) \ln(-i e^{2i(bx+a)} + 1)}{2} + \frac{i \operatorname{dilog}(i e^{2i(bx+a)} + 1)}{b} - \frac{i \operatorname{dilog}(-i e^{2i(bx+a)} + 1)}{b}$
derivativedivides	$\frac{\operatorname{arctan}(\tan(bx+a)) \operatorname{arctanh}(\tan(bx+a)) + \frac{\operatorname{arctan}(\tan(bx+a)) \ln\left(1 + \frac{i(1+i \tan(bx+a))^2}{1+\tan(bx+a)^2}\right)}{2}}{b} - \frac{\operatorname{arctan}(\tan(bx+a)) \ln\left(1 - \frac{i(1+i \tan(bx+a))^2}{1+\tan(bx+a)^2}\right)}{2}}{b}$
default	$\frac{\operatorname{arctan}(\tan(bx+a)) \operatorname{arctanh}(\tan(bx+a)) + \frac{\operatorname{arctan}(\tan(bx+a)) \ln\left(1 + \frac{i(1+i \tan(bx+a))^2}{1+\tan(bx+a)^2}\right)}{2}}{b} - \frac{\operatorname{arctan}(\tan(bx+a)) \ln\left(1 - \frac{i(1+i \tan(bx+a))^2}{1+\tan(bx+a)^2}\right)}{2}}{b}$
risch	Expression too large to display

```
input int(arctanh(tan(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output x*arctanh(tan(b*x+a))-1/b*(-1/2*(b*x+a)*ln(I*exp(2*I*(b*x+a))+1)+1/2*(b*x+a)*ln(-I*exp(2*I*(b*x+a))+1)+1/4*I*dilog(I*exp(2*I*(b*x+a))+1)-1/4*I*dilog(-I*exp(2*I*(b*x+a))+1)-1/2*a*ln(sec(2*b*x+2*a)+tan(2*b*x+2*a)))
```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 499 vs. 2(57) = 114.

Time = 0.12 (sec) , antiderivative size = 499, normalized size of antiderivative = 6.32

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = \text{Too large to display}$$

```
input integrate(arctanh(tan(b*x+a)), x, algorithm="fricas")
```

output

```

1/8*(4*b*x*log(-(tan(b*x + a) + 1)/(tan(b*x + a) - 1)) - 2*(b*x + a)*log((
(I + 1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/(tan(b*x + a)^2 + 1)) + 2
*a*log(((I + 1)*tan(b*x + a)^2 + 2*I*tan(b*x + a) + I - 1)/(tan(b*x + a)^2
+ 1)) - 2*a*log(((I + 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I - 1)/(tan(
b*x + a)^2 + 1)) + 2*(b*x + a)*log(((I + 1)*tan(b*x + a)^2 - 2*tan(b*x + a
) - I + 1)/(tan(b*x + a)^2 + 1)) - 2*(b*x + a)*log((-I - 1)*tan(b*x + a)^
2 + 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) + 2*(b*x + a)*log((-I -
1)*tan(b*x + a)^2 - 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1)) + 2*a*l
og(((I - 1)*tan(b*x + a)^2 + 2*I*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1
)) - 2*a*log(((I - 1)*tan(b*x + a)^2 - 2*I*tan(b*x + a) + I + 1)/(tan(b*x
+ a)^2 + 1)) + I*dilog(-((I + 1)*tan(b*x + a)^2 + 2*tan(b*x + a) - I + 1)/
(tan(b*x + a)^2 + 1) + 1) + I*dilog(-((I + 1)*tan(b*x + a)^2 - 2*tan(b*x +
a) - I + 1)/(tan(b*x + a)^2 + 1) + 1) - I*dilog(-(-(I - 1)*tan(b*x + a)^2
+ 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1) - I*dilog(-(-(I - 1)*
tan(b*x + a)^2 - 2*tan(b*x + a) + I + 1)/(tan(b*x + a)^2 + 1) + 1))/b

```

**Sympy [F]**

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(a + bx)) dx$$

input

```
integrate(atanh(tan(b*x+a)),x)
```

output

```
Integral(atanh(tan(a + b*x)), x)
```

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 182 vs.  $2(57) = 114$ .

Time = 0.14 (sec) , antiderivative size = 182, normalized size of antiderivative = 2.30

$$\int \operatorname{arctanh}(\tan(a + bx)) dx$$

$$= \frac{4(bx + a) \operatorname{artanh}(\tan(bx + a)) + \left(\operatorname{arctan}\left(\frac{1}{2} \tan(bx + a) + \frac{1}{2}\right), \frac{1}{2} \tan(bx + a) + \frac{1}{2}\right) - \operatorname{arctan}\left(\frac{1}{2} \tan(bx + a) - \frac{1}{2}\right)}{b}$$



input `integrate(arctanh(tan(b*x+a)),x, algorithm="maxima")`

output `1/4*(4*(b*x + a)*arctanh(tan(b*x + a)) + (arctan2(1/2*tan(b*x + a) + 1/2, 1/2*tan(b*x + a) + 1/2) - arctan2(1/2*tan(b*x + a) - 1/2, -1/2*tan(b*x + a) + 1/2))*log(tan(b*x + a)^2 + 1) - (b*x + a)*log(1/2*tan(b*x + a)^2 + tan(b*x + a) + 1/2) + (b*x + a)*log(1/2*tan(b*x + a)^2 - tan(b*x + a) + 1/2) - I*dilog((1/2*I + 1/2)*tan(b*x + a) - 1/2*I + 1/2) + I*dilog(-(1/2*I - 1/2)*tan(b*x + a) + 1/2*I + 1/2) + I*dilog((1/2*I - 1/2)*tan(b*x + a) + 1/2*I + 1/2) - I*dilog(-(1/2*I + 1/2)*tan(b*x + a) - 1/2*I + 1/2))/b`

### Giac [F]

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{artanh}(\tan(bx + a)) dx$$

input `integrate(arctanh(tan(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(tan(b*x + a)), x)`

### Mupad [F(-1)]

Timed out.

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(a + bx)) dx$$

input `int(atanh(tan(a + b*x)),x)`

output `int(atanh(tan(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(\tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a)) dx$$

input `int(atanh(tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)),x)`

### 3.316 $\int \frac{\operatorname{arctanh}(\tan(a+bx))}{e+fx} dx$

Optimal result	2206
Mathematica [N/A]	2206
Rubi [N/A]	2207
Maple [N/A]	2207
Fricas [N/A]	2208
Sympy [N/A]	2208
Maxima [N/A]	2208
Giac [N/A]	2209
Mupad [N/A]	2209
Reduce [N/A]	2210

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(\tan(a+bx))}{e+fx} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(\tan(a+bx))}{e+fx}, x\right)$$

output `Defer(Int)(arctanh(tan(b*x+a))/(f*x+e), x)`

#### Mathematica [N/A]

Not integrable

Time = 6.96 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\tan(a+bx))}{e+fx} dx = \int \frac{\operatorname{arctanh}(\tan(a+bx))}{e+fx} dx$$

input `Integrate[ArcTanh[Tan[a + b*x]]/(e + f*x), x]`

output `Integrate[ArcTanh[Tan[a + b*x]]/(e + f*x), x]`

**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx$$

input `Int[ArcTanh[Tan[a + b*x]]/(e + f*x),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\tan(bx + a))}{fx + e} dx$$

input `int(arctanh(tan(b*x+a))/(f*x+e),x)`

output `int(arctanh(tan(b*x+a))/(f*x+e),x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\tan(bx + a))}{fx + e} dx$$

input `integrate(arctanh(tan(b*x+a))/(f*x+e),x, algorithm="fricas")`

output `integral(arctanh(tan(b*x + a))/(f*x + e), x)`

**Sympy [N/A]**

Not integrable

Time = 0.60 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\tan(a + bx))}{e + fx} dx$$

input `integrate(atanh(tan(b*x+a))/(f*x+e),x)`

output `Integral(atanh(tan(a + b*x))/(e + f*x), x)`

**Maxima [N/A]**

Not integrable

Time = 1.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\tan(bx + a))}{fx + e} dx$$

input `integrate(arctanh(tan(b*x+a))/(f*x+e),x, algorithm="maxima")`

output `integrate(arctanh(tan(b*x + a))/(f*x + e), x)`

### Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\tan(bx + a))}{fx + e} dx$$

input `integrate(arctanh(tan(b*x+a))/(f*x+e),x, algorithm="giac")`

output `integrate(arctanh(tan(b*x + a))/(f*x + e), x)`

### Mupad [N/A]

Not integrable

Time = 3.35 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\tan(a + bx))}{e + fx} dx$$

input `int(atanh(tan(a + b*x))/(e + f*x),x)`

output `int(atanh(tan(a + b*x))/(e + f*x), x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\tan(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\tan(bx + a))}{fx + e} dx$$

input `int(atanh(tan(b*x+a))/(f*x+e),x)`output `int(atanh(tan(a + b*x))/(e + f*x),x)`

### 3.317 $\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx$

Optimal result	2211
Mathematica [A] (verified)	2212
Rubi [A] (verified)	2213
Maple [C] (warning: unable to verify)	2219
Fricas [B] (verification not implemented)	2219
Sympy [F]	2220
Maxima [F]	2221
Giac [F]	2221
Mupad [F(-1)]	2222
Reduce [F]	2222

#### Optimal result

Integrand size = 15, antiderivative size = 395

$$\begin{aligned}
 \int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx &= \frac{1}{3} x^3 \operatorname{arctanh}(c + d \tan(a + bx)) \\
 &+ \frac{1}{6} x^3 \log \left( 1 + \frac{(1 - c + id)e^{2ia + 2ibx}}{1 - c - id} \right) \\
 &- \frac{1}{6} x^3 \log \left( 1 + \frac{(1 + c - id)e^{2ia + 2ibx}}{1 + c + id} \right) \\
 &- \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(1 - c + id)e^{2ia + 2ibx}}{1 - c - id} \right)}{4b} \\
 &+ \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(1 + c - id)e^{2ia + 2ibx}}{1 + c + id} \right)}{4b} \\
 &+ \frac{x \operatorname{PolyLog} \left( 3, -\frac{(1 - c + id)e^{2ia + 2ibx}}{1 - c - id} \right)}{4b^2} \\
 &- \frac{x \operatorname{PolyLog} \left( 3, -\frac{(1 + c - id)e^{2ia + 2ibx}}{1 + c + id} \right)}{4b^2} \\
 &+ \frac{i \operatorname{PolyLog} \left( 4, -\frac{(1 - c + id)e^{2ia + 2ibx}}{1 - c - id} \right)}{8b^3} \\
 &- \frac{i \operatorname{PolyLog} \left( 4, -\frac{(1 + c - id)e^{2ia + 2ibx}}{1 + c + id} \right)}{8b^3}
 \end{aligned}$$



output

$$\begin{aligned} & \frac{1}{3}x^3 \operatorname{arctanh}(c+d \tan(bx+a)) + \frac{1}{6}x^3 \ln(1+(1-c+I*d) \exp(2I*a+2I*b*x) / \\ & (1-c-I*d)) - \frac{1}{6}x^3 \ln(1+(1+c-I*d) \exp(2I*a+2I*b*x) / (1+c+I*d)) - \frac{1}{4}I*x^2 * \\ & \operatorname{polylog}(2, -(1-c+I*d) \exp(2I*a+2I*b*x) / (1-c-I*d)) / b + \frac{1}{4}I*x^2 * \operatorname{polylog}(2, - \\ & (1+c-I*d) \exp(2I*a+2I*b*x) / (1+c+I*d)) / b + \frac{1}{4}x * \operatorname{polylog}(3, -(1-c+I*d) \exp(2 \\ & I*a+2I*b*x) / (1-c-I*d)) / b^2 - \frac{1}{4}x * \operatorname{polylog}(3, -(1+c-I*d) \exp(2I*a+2I*b*x) \\ & / (1+c+I*d)) / b^2 + \frac{1}{8}I * \operatorname{polylog}(4, -(1-c+I*d) \exp(2I*a+2I*b*x) / (1-c-I*d)) / b \\ & ^3 - \frac{1}{8}I * \operatorname{polylog}(4, -(1+c-I*d) \exp(2I*a+2I*b*x) / (1+c+I*d)) / b^3 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 3.20 (sec) , antiderivative size = 349, normalized size of antiderivative = 0.88

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(c + d \tan(a + bx)) + 4b^3 x^3 \log\left(1 + \frac{(-1+c+id)e^{-2i(a+bx)}}{-1+c-id}\right) - 4b^3 x^3 \log\left(1 + \frac{(1+c+id)e^{-2i(a+bx)}}{1+c-id}\right)}{24b^3}$$

input

`Integrate[x^2*ArcTanh[c + d*Tan[a + b*x]],x]`

output

$$\begin{aligned} & \frac{(8*b^3*x^3*ArcTanh[c + d*Tan[a + b*x]] + 4*b^3*x^3*Log[1 + (-1 + c + I*d) / \\ & ((-1 + c - I*d)*E^((2*I)*(a + b*x)))] - 4*b^3*x^3*Log[1 + (1 + c + I*d) / (( \\ & 1 + c - I*d)*E^((2*I)*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, (1 - c - I*d \\ & ) / ((-1 + c - I*d)*E^((2*I)*(a + b*x)))] - (6*I)*b^2*x^2*PolyLog[2, (-1 - c \\ & - I*d) / ((1 + c - I*d)*E^((2*I)*(a + b*x)))] + 6*b*x*PolyLog[3, (1 - c - I \\ & *d) / ((-1 + c - I*d)*E^((2*I)*(a + b*x)))] - 6*b*x*PolyLog[3, (-1 - c - I*d) / (( \\ & 1 + c - I*d)*E^((2*I)*(a + b*x)))] - (3*I)*PolyLog[4, (1 - c - I*d) / (( \\ & -1 + c - I*d)*E^((2*I)*(a + b*x)))] + (3*I)*PolyLog[4, (-1 - c - I*d) / ((1 \\ & + c - I*d)*E^((2*I)*(a + b*x)))] / (24*b^3) \end{aligned}$$

**Rubi [A] (verified)**

Time = 1.79 (sec) , antiderivative size = 522, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6821, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}(d \tan(a + bx) + c) dx \\
 & \quad \downarrow \text{6821} \\
 & -\frac{1}{3}b(ic + d + i) \int \frac{e^{2ia+2ibx} x^3}{c + (c - id + 1)e^{2ia+2ibx} + id + 1} dx + \frac{1}{3}b(-d + i(1 - \\
 & c)) \int \frac{e^{2ia+2ibx} x^3}{-c + (-c + id + 1)e^{2ia+2ibx} - id + 1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3}b(-d + i(1 - c)) \left( \frac{x^3 \log\left(1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b(-d + i(1 - c))} - \frac{3 \int x^2 \log\left(\frac{e^{2ia+2ibx}(-c+id+1)}{-c-id+1} + 1\right) dx}{2b(-d + i(1 - c))} \right) - \\
 & \frac{1}{3}b(ic + d + i) \left( \frac{x^3 \log\left(1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2(bd + i(bc + b))} - \frac{3 \int x^2 \log\left(\frac{e^{2ia+2ibx}(c-id+1)}{c+id+1} + 1\right) dx}{2(bd + i(bc + b))} \right) + \\
 & \quad \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\begin{aligned}
 & c) \left( \frac{x^3 \log \left( 1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b(-d+i(1-c))} - \frac{\frac{1}{3}b(-d+i(1-c)) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right) dx}{b} \right)}{2b(-d+i(1-c))} \right) \\
 & i) \left( \frac{x^3 \log \left( 1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2(bd+i(bc+b))} - \frac{\frac{1}{3}b(ic+d) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right) dx}{b} \right)}{2(bd+i(bc+b))} \right) + \\
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) + c)
 \end{aligned}$$

↓ 7163

$$\begin{aligned}
 & c) \left( \frac{x^3 \log \left( 1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b(-d+i(1-c))} - \frac{\frac{1}{3}b(-d+i(1-c)) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 3, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right) dx}{2b} - \frac{ix \operatorname{PolyLog} \left( 3, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{b} \right)}{2b(-d+i(1-c))} \right) \\
 & i) \left( \frac{x^3 \log \left( 1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2(bd+i(bc+b))} - \frac{\frac{1}{3}b(ic+d) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 3, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right) dx}{2b} - \frac{ix \operatorname{PolyLog} \left( 3, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{b} \right)}{2(bd+i(bc+b))} \right) \\
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) + c)
 \end{aligned}$$

↓ 2720

$$\begin{aligned}
 & c) \left( \frac{x^3 \log \left( 1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b(-d+i(1-c))} - \frac{\frac{1}{3}b(-d+i(1-c)) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b} - i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog} \left( 3, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{4b^2} \right)}{2b(-d+i(1-c))} \right)}{2b(-d+i(1-c))} \right) \\
 & i) \left( \frac{x^3 \log \left( 1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2(bd+i(bc+b))} - \frac{\frac{1}{3}b(ic+d) \left( \frac{ix^2 \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2b} - i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog} \left( 3, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right) de^2}{4b^2} \right)}{b} \right)}{2(bd+i(bc+b))} \right) \\
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) + c)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a + bx) + c) + \frac{1}{3}b(-d + i(1 - \\
 c)) & \left( \frac{x^3 \log\left(1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b(-d + i(1 - c))} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b} - i \frac{\operatorname{PolyLog}\left(4, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{b} \right)}{2b(-d + i(1 - c))} \right) \\
 i) & \left( \frac{x^3 \log\left(1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2(bd + i(bc + b))} - \frac{\frac{1}{3}b(ic + d + 3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2b} - i \frac{\operatorname{PolyLog}\left(4, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{b} \right)}{2(bd + i(bc + b))} \right)
 \end{aligned}$$

input `Int[x^2*ArcTanh[c + d*Tan[a + b*x]],x]`

output `(x^3*ArcTanh[c + d*Tan[a + b*x]])/3 + (b*(I*(1 - c) - d)*((x^3*Log[1 + ((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d)])/(2*b*(I*(1 - c) - d)) - (3*(((I/2)*x^2*PolyLog[2, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d))])/b - (I*(((1/2)*x*PolyLog[3, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d))])/b + PolyLog[4, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d)]/(4*b^2)))/b)/(2*b*(I*(1 - c) - d)))/3 - (b*(I + I*c + d)*((x^3*Log[1 + ((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d)])/(2*(I*(b + b*c) + b*d)) - (3*(((I/2)*x^2*PolyLog[2, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d))])/b - (I*(((1/2)*x*PolyLog[3, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d))])/b + PolyLog[4, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d)]/(4*b^2)))/b)/(2*(I*(b + b*c) + b*d)))/3`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6821

```
Int[ArcTanh[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m
+ 1))), x] + (-Simp[I*b*((1 + c - I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)
*(E^(2*I*a + 2*I*b*x))/(1 + c + I*d + (1 + c - I*d)*E^(2*I*a + 2*I*b*x))], x
], x] + Simp[I*b*((1 - c + I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*
I*a + 2*I*b*x))/(1 - c - I*d + (1 - c + I*d)*E^(2*I*a + 2*I*b*x))], x], x]
/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c + I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 46.01 (sec) , antiderivative size = 6917, normalized size of antiderivative = 17.51

method	result	size
risch	Expression too large to display	6917

input

```
int(x^2*arctanh(c+d*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2165 vs. 2(279) = 558.

Time = 0.17 (sec) , antiderivative size = 2165, normalized size of antiderivative = 5.48

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctanh(c+d*tan(b*x+a)),x, algorithm="fricas")
```



output

```

1/48*(8*b^3*x^3*log(-(d*tan(b*x + a) + c + 1)/(d*tan(b*x + a) + c - 1)) -
6*I*b^2*x^2*dilog(2*((I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c + 1)*
d + (I*c^2 - 2*(c + 1)*d - I*d^2 + 2*I*c + I)*tan(b*x + a) - 2*c - 1)/((c^
2 + d^2 + 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1) + 1) + 6*I*b^2*x^
2*dilog(2*((-I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 + I*(c + 1)*d + (-I*c
^2 - 2*(c + 1)*d + I*d^2 - 2*I*c - I)*tan(b*x + a) - 2*c - 1)/((c^2 + d^2
+ 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1) + 1) + 6*I*b^2*x^2*dilog(
2*((I*(c - 1)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c - 1)*d + (I*c^2 - 2*(c
- 1)*d - I*d^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)
*tan(b*x + a)^2 + c^2 + d^2 - 2*c + 1) + 1) - 6*I*b^2*x^2*dilog(2*((-I*(c
- 1)*d - d^2)*tan(b*x + a)^2 - c^2 + I*(c - 1)*d + (-I*c^2 - 2*(c - 1)*d +
I*d^2 + 2*I*c - I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)*tan(b*x
+ a)^2 + c^2 + d^2 - 2*c + 1) + 1) + 4*a^3*log(((I*(c + 1)*d + d^2)*tan(b
*x + a)^2 - c^2 + I*(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) -
2*c - 1)/(tan(b*x + a)^2 + 1)) + 4*a^3*log(((I*(c + 1)*d - d^2)*tan(b*x +
a)^2 + c^2 + I*(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) + 2*c
+ 1)/(tan(b*x + a)^2 + 1)) - 4*a^3*log(((I*(c - 1)*d + d^2)*tan(b*x + a)^
2 - c^2 + I*(c - 1)*d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1
)/(tan(b*x + a)^2 + 1)) - 4*a^3*log(((I*(c - 1)*d - d^2)*tan(b*x + a)^2 +
c^2 + I*(c - 1)*d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) - 2*c + 1)...

```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input

```
integrate(x**2*atanh(c+d*tan(b*x+a)),x)
```

output

```
Integral(x**2*atanh(c + d*tan(a + b*x)), x)
```

**Maxima [F]**

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x^2 \operatorname{artanh}(d \tan(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*tan(b*x+a)),x, algorithm="maxima")`

output

```
1/12*x^3*log((c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c + 1)*d*sin(2*
b*x + 2*a) + (c^2 + d^2 + 2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 + 2*(c^2
- d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 2*c + 1) - 1/12*x^3*log((c^2 + d^2 -
2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c - 1)*d*sin(2*b*x + 2*a) + (c^2 + d^2 -
2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 + 2*(c^2 - d^2 - 2*c + 1)*cos(2*b*
x + 2*a) - 2*c + 1) - 4*b*d*integrate(-1/3*(2*(c^2 + d^2 - 1)*x^3*cos(2*b*
x + 2*a)^2 + 2*c*d*x^3*sin(2*b*x + 2*a) + 2*(c^2 + d^2 - 1)*x^3*sin(2*b*x
+ 2*a)^2 + (c^2 - d^2 - 1)*x^3*cos(2*b*x + 2*a) - (2*c*d*x^3*sin(2*b*x + 2
*a) - (c^2 - d^2 - 1)*x^3*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x^3*
cos(2*b*x + 2*a) + (c^2 - d^2 - 1)*x^3*sin(2*b*x + 2*a))*sin(4*b*x + 4*a))
/(c^4 + d^4 + 2*(c^2 + 1)*d^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*
cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*cos(2*b*x
+ 2*a)^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*sin(4*b*x + 4*a)^2 +
4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*sin(2*b*x + 2*a)^2 - 2*c^2 +
2*(c^4 + d^4 - 2*(3*c^2 - 1)*d^2 - 2*c^2 + 2*(c^4 - d^4 - 2*c^2 + 1)*cos(2
*b*x + 2*a) - 4*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x + 4*
a) + 4*(c^4 - d^4 - 2*c^2 + 1)*cos(2*b*x + 2*a) - 4*(2*c*d^3 - 2*(c^3 - c)
*d - 2*(c*d^3 + (c^3 - c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 - 2*c^2 + 1)*si
n(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a
) + 1), x)
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x^2 \operatorname{artanh}(d \tan(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(d*tan(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input `int(x^2*atanh(c + d*tan(a + b*x)),x)`output `int(x^2*atanh(c + d*tan(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a)d + c) x^2 dx$$

input `int(x^2*atanh(c+d*tan(b*x+a)),x)`output `int(atanh(tan(a + b*x)*d + c)*x**2,x)`

### 3.318 $\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx$

Optimal result	2223
Mathematica [A] (verified)	2224
Rubi [A] (verified)	2224
Maple [C] (warning: unable to verify)	2228
Fricas [B] (verification not implemented)	2228
Sympy [F]	2229
Maxima [F]	2230
Giac [F]	2230
Mupad [F(-1)]	2231
Reduce [F]	2231

#### Optimal result

Integrand size = 13, antiderivative size = 295

$$\begin{aligned}
 \int x \operatorname{arctanh}(c + d \tan(a + bx)) dx &= \frac{1}{2} x^2 \operatorname{arctanh}(c + d \tan(a + bx)) \\
 &+ \frac{1}{4} x^2 \log \left( 1 + \frac{(1 - c + id)e^{2ia+2ibx}}{1 - c - id} \right) \\
 &- \frac{1}{4} x^2 \log \left( 1 + \frac{(1 + c - id)e^{2ia+2ibx}}{1 + c + id} \right) \\
 &- \frac{ix \operatorname{PolyLog} \left( 2, -\frac{(1-c+id)e^{2ia+2ibx}}{1-c-id} \right)}{4b} \\
 &+ \frac{ix \operatorname{PolyLog} \left( 2, -\frac{(1+c-id)e^{2ia+2ibx}}{1+c+id} \right)}{4b} \\
 &+ \frac{\operatorname{PolyLog} \left( 3, -\frac{(1-c+id)e^{2ia+2ibx}}{1-c-id} \right)}{8b^2} \\
 &- \frac{\operatorname{PolyLog} \left( 3, -\frac{(1+c-id)e^{2ia+2ibx}}{1+c+id} \right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{2}x^2 \operatorname{arctanh}(c+d \tan(bx+a)) + \frac{1}{4}x^2 \ln(1+(1-c+I*d) \exp(2I*a+2I*b*x) / \\ & (1-c-I*d)) - \frac{1}{4}x^2 \ln(1+(1+c-I*d) \exp(2I*a+2I*b*x) / (1+c+I*d)) - \frac{1}{4}I*x*po \\ & lylog(2, -(1-c+I*d) \exp(2I*a+2I*b*x) / (1-c-I*d)) / b + \frac{1}{4}I*x*polylog(2, -(1+c \\ & -I*d) \exp(2I*a+2I*b*x) / (1+c+I*d)) / b + \frac{1}{8}polylog(3, -(1-c+I*d) \exp(2I*a+2 \\ & *I*b*x) / (1-c-I*d)) / b^2 - \frac{1}{8}polylog(3, -(1+c-I*d) \exp(2I*a+2I*b*x) / (1+c+I* \\ & d)) / b^2 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 2.59 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.88

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx$$

$$= \frac{4b^2 x^2 \operatorname{arctanh}(c + d \tan(a + bx)) + 2b^2 x^2 \log\left(1 + \frac{(-1+c+id)e^{-2i(a+bx)}}{-1+c-id}\right) - 2b^2 x^2 \log\left(1 + \frac{(1+c+id)e^{-2i(a+bx)}}{1+c-id}\right)}{1}$$

input

Integrate[x\*ArcTanh[c + d\*Tan[a + b\*x]],x]

output

$$\begin{aligned} & (4*b^2*x^2*ArcTanh[c + d*Tan[a + b*x]] + 2*b^2*x^2*Log[1 + (-1 + c + I*d) / \\ & ((-1 + c - I*d)*E^((2*I)*(a + b*x)))] - 2*b^2*x^2*Log[1 + (1 + c + I*d) / (( \\ & 1 + c - I*d)*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, (1 - c - I*d) / (( \\ & -1 + c - I*d)*E^((2*I)*(a + b*x)))] - (2*I)*b*x*PolyLog[2, (-1 - c - I*d) / \\ & ((1 + c - I*d)*E^((2*I)*(a + b*x)))] + PolyLog[3, (1 - c - I*d) / ((-1 + c - \\ & I*d)*E^((2*I)*(a + b*x)))] - PolyLog[3, (-1 - c - I*d) / ((1 + c - I*d)*E^(( \\ & 2*I)*(a + b*x)))] / (8*b^2) \end{aligned}$$
**Rubi [A] (verified)**Time = 1.83 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.36, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {6821, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x \operatorname{arctanh}(d \tan(a + bx) + c) dx \\
& \quad \downarrow \text{6821} \\
& -\frac{1}{2}b(ic + d + i) \int \frac{e^{2ia+2ibx} x^2}{c + (c - id + 1)e^{2ia+2ibx} + id + 1} dx + \frac{1}{2}b(-d + i(1 - \\
c)) \int \frac{e^{2ia+2ibx} x^2}{-c + (-c + id + 1)e^{2ia+2ibx} - id + 1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& \frac{1}{2}b(-d + i(1 - c)) \left( \frac{x^2 \log \left( 1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b(-d + i(1 - c))} - \frac{\int x \log \left( \frac{e^{2ia+2ibx}(-c+id+1)}{-c-id+1} + 1 \right) dx}{b(-d + i(1 - c))} \right) - \\
& \frac{1}{2}b(ic + d + i) \left( \frac{x^2 \log \left( 1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2(bd + i(bc + b))} - \frac{\int x \log \left( \frac{e^{2ia+2ibx}(c-id+1)}{c+id+1} + 1 \right) dx}{bd + i(bc + b)} \right) + \\
& \quad \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a + bx) + c) \\
& \quad \downarrow \text{3011} \\
& \frac{1}{2}b(-d + i(1 - \\
c)) \left( \frac{x^2 \log \left( 1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b(-d + i(1 - c))} - \frac{ix \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1} \right) dx}{2b} \right) - \\
& \frac{1}{2}b(ic + d + \\
i) \left( \frac{x^2 \log \left( 1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2(bd + i(bc + b))} - \frac{ix \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1} \right) dx}{2b} \right) + \\
& \quad \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a + bx) + c) \\
& \quad \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
 c) & \left( \frac{x^2 \log\left(1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b(-d+i(1-c))} - \frac{\frac{1}{2}b(-d+i(1-c))}{2b} \frac{ix \operatorname{PolyLog}\left(2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right) dx}{4b^2} \right) \\
 i) & \left( \frac{x^2 \log\left(1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2(bd+i(bc+b))} - \frac{\frac{1}{2}b(ic+d)}{2b} \frac{ix \operatorname{PolyLog}\left(2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right) dx}{4b^2} \right) \\
 & \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) + c)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 c) & \left( \frac{x^2 \log\left(1 + \frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b(-d+i(1-c))} - \frac{\frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) + c) + \frac{1}{2}b(-d+i(1-c))}{2b} \frac{ix \operatorname{PolyLog}\left(2, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, -\frac{(-c+id+1)e^{2ia+2ibx}}{-c-id+1}\right)}{4b^2} \right) \\
 i) & \left( \frac{x^2 \log\left(1 + \frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2(bd+i(bc+b))} - \frac{\frac{1}{2}b(ic+d)}{2b} \frac{ix \operatorname{PolyLog}\left(2, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, -\frac{(c-id+1)e^{2ia+2ibx}}{c+id+1}\right)}{4b^2} \right)
 \end{aligned}$$

input `Int[x*ArcTanh[c + d*Tan[a + b*x]],x]`

output `(x^2*ArcTanh[c + d*Tan[a + b*x]])/2 + (b*(I*(1 - c) - d)*((x^2*Log[1 + ((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 - c - I*d)))/(2*b*(I*(1 - c) - d)) - (((I/2)*x*PolyLog[2, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 - c - I*d)))]/b - PolyLog[3, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 - c - I*d))]/(4*b^2)))/(b*(I*(1 - c) - d)))/2 - (b*(I + I*c + d)*((x^2*Log[1 + ((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 + c + I*d)))/(2*(I*(b + b*c) + b*d)) - (((I/2)*x*PolyLog[2, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 + c + I*d)))]/b - PolyLog[3, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x)]/(1 + c + I*d))]/(4*b^2)))/(I*(b + b*c) + b*d))/2`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6821

```
Int[ArcTanh[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m
+ 1))), x] + (-Simp[I*b*((1 + c - I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)
*(E^(2*I*a + 2*I*b*x))/(1 + c + I*d + (1 + c - I*d)*E^(2*I*a + 2*I*b*x))], x
], x] + Simp[I*b*((1 - c + I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*
I*a + 2*I*b*x))/(1 - c - I*d + (1 - c + I*d)*E^(2*I*a + 2*I*b*x))], x], x]
/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c + I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```



**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.25 (sec) , antiderivative size = 6525, normalized size of antiderivative = 22.12

method	result	size
risch	Expression too large to display	6525

input `int(x*arctanh(c+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1689 vs.  $2(209) = 418$ .

Time = 0.17 (sec) , antiderivative size = 1689, normalized size of antiderivative = 5.73

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctanh(c+d*tan(b*x+a)),x, algorithm="fricas")`

output

```

1/16*(4*b^2*x^2*log(-(d*tan(b*x + a) + c + 1)/(d*tan(b*x + a) + c - 1)) -
2*I*b*x*dilog(2*((I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c + 1)*d +
(I*c^2 - 2*(c + 1)*d - I*d^2 + 2*I*c + I)*tan(b*x + a) - 2*c - 1)/((c^2 +
d^2 + 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1) + 1) + 2*I*b*x*dilog(
2*((-I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 + I*(c + 1)*d + (-I*c^2 - 2*(
c + 1)*d + I*d^2 - 2*I*c - I)*tan(b*x + a) - 2*c - 1)/((c^2 + d^2 + 2*c +
1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1) + 1) + 2*I*b*x*dilog(2*((I*(c - 1
)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c - 1)*d + (I*c^2 - 2*(c - 1)*d - I*d
^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)*tan(b*x + a
)^2 + c^2 + d^2 - 2*c + 1) + 1) - 2*I*b*x*dilog(2*((-I*(c - 1)*d - d^2)*ta
n(b*x + a)^2 - c^2 + I*(c - 1)*d + (-I*c^2 - 2*(c - 1)*d + I*d^2 + 2*I*c -
I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)*tan(b*x + a)^2 + c^2 +
d^2 - 2*c + 1) + 1) - 2*a^2*log(((I*(c + 1)*d + d^2)*tan(b*x + a)^2 - c^2
+ I*(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) - 2*c - 1)/(tan(b
*x + a)^2 + 1)) - 2*a^2*log(((I*(c + 1)*d - d^2)*tan(b*x + a)^2 + c^2 + I*
(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) + 2*c + 1)/(tan(b*x +
a)^2 + 1)) + 2*a^2*log(((I*(c - 1)*d + d^2)*tan(b*x + a)^2 - c^2 + I*(c -
1)*d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1)/(tan(b*x + a)^
2 + 1)) + 2*a^2*log(((I*(c - 1)*d - d^2)*tan(b*x + a)^2 + c^2 + I*(c - 1)*
d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) - 2*c + 1)/(tan(b*x + a)^2...

```

### Sympy [F]

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input

```
integrate(x*atanh(c+d*tan(b*x+a)),x)
```

output

```
Integral(x*atanh(c + d*tan(a + b*x)), x)
```

**Maxima [F]**

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x \operatorname{artanh}(d \tan(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*tan(b*x+a)),x, algorithm="maxima")`

output

```
-2*b*d*integrate(-(2*(c^2 + d^2 - 1)*x^2*cos(2*b*x + 2*a)^2 + 2*c*d*x^2*si
n(2*b*x + 2*a) + 2*(c^2 + d^2 - 1)*x^2*sin(2*b*x + 2*a)^2 + (c^2 - d^2 - 1
)*x^2*cos(2*b*x + 2*a) - (2*c*d*x^2*sin(2*b*x + 2*a) - (c^2 - d^2 - 1)*x^2
*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x^2*cos(2*b*x + 2*a) + (c^2 -
d^2 - 1)*x^2*sin(2*b*x + 2*a))*sin(4*b*x + 4*a))/(c^4 + d^4 + 2*(c^2 + 1)
*d^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*cos(4*b*x + 4*a)^2 + 4*(c
^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*cos(2*b*x + 2*a)^2 + (c^4 + d^4 +
2*(c^2 + 1)*d^2 - 2*c^2 + 1)*sin(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 -
1)*d^2 - 2*c^2 + 1)*sin(2*b*x + 2*a)^2 - 2*c^2 + 2*(c^4 + d^4 - 2*(3*c^2 -
1)*d^2 - 2*c^2 + 2*(c^4 - d^4 - 2*c^2 + 1)*cos(2*b*x + 2*a) - 4*(c*d^3 +
(c^3 - c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x + 4*a) + 4*(c^4 - d^4 - 2*c^2
+ 1)*cos(2*b*x + 2*a) - 4*(2*c*d^3 - 2*(c^3 - c)*d - 2*(c*d^3 + (c^3 - c)
*d)*cos(2*b*x + 2*a) - (c^4 - d^4 - 2*c^2 + 1)*sin(2*b*x + 2*a))*sin(4*b*x
+ 4*a) + 8*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a) + 1), x) + 1/8*x^2*log(
(c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c + 1)*d*sin(2*b*x + 2*a) +
(c^2 + d^2 + 2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 + 2*(c^2 - d^2 + 2*c
+ 1)*cos(2*b*x + 2*a) + 2*c + 1) - 1/8*x^2*log((c^2 + d^2 - 2*c + 1)*cos(2
*b*x + 2*a)^2 + 4*(c - 1)*d*sin(2*b*x + 2*a) + (c^2 + d^2 - 2*c + 1)*sin(2
*b*x + 2*a)^2 + c^2 + d^2 + 2*(c^2 - d^2 - 2*c + 1)*cos(2*b*x + 2*a) - 2*c
+ 1)
```

**Giac [F]**

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x \operatorname{artanh}(d \tan(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*tan(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int x \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input `int(x*atanh(c + d*tan(a + b*x)),x)`output `int(x*atanh(c + d*tan(a + b*x)), x)`**Reduce [F]**

$$\int x \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a) d + c) x dx$$

input `int(x*atanh(c+d*tan(b*x+a)),x)`output `int(atanh(tan(a + b*x)*d + c)*x,x)`

### 3.319 $\int \operatorname{arctanh}(c + d \tan(a + bx)) dx$

Optimal result	2232
Mathematica [A] (warning: unable to verify)	2233
Rubi [A] (verified)	2233
Maple [B] (verified)	2236
Fricas [B] (verification not implemented)	2237
Sympy [F]	2238
Maxima [B] (verification not implemented)	2238
Giac [F]	2239
Mupad [F(-1)]	2239
Reduce [F]	2239

#### Optimal result

Integrand size = 11, antiderivative size = 194

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = x \operatorname{arctanh}(c + d \tan(a + bx)) + \frac{1}{2} x \log \left( 1 + \frac{(1 - c + id)e^{2ia+2ibx}}{1 - c - id} \right) - \frac{1}{2} x \log \left( 1 + \frac{(1 + c - id)e^{2ia+2ibx}}{1 + c + id} \right) - \frac{i \operatorname{PolyLog} \left( 2, -\frac{(1-c+id)e^{2ia+2ibx}}{1-c-id} \right)}{4b} + \frac{i \operatorname{PolyLog} \left( 2, -\frac{(1+c-id)e^{2ia+2ibx}}{1+c+id} \right)}{4b}$$

output

```
x*arctanh(c+d*tan(b*x+a))+1/2*x*ln(1+(1-c+I*d)*exp(2*I*a+2*I*b*x)/(1-c-I*d))
)-1/2*x*ln(1+(1+c-I*d)*exp(2*I*a+2*I*b*x)/(1+c+I*d))-1/4*I*polylog(2,-(1-
c+I*d)*exp(2*I*a+2*I*b*x)/(1-c-I*d))/b+1/4*I*polylog(2,-(1+c-I*d)*exp(2*I*
a+2*I*b*x)/(1+c+I*d))/b
```

**Mathematica [A] (warning: unable to verify)**

Time = 3.73 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.88

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = x \left( \operatorname{arctanh}(c + d \tan(a + bx)) \right. \\ \left. + \frac{2a \log(1 - c - d \tan(a + bx)) - i \log(1 - i \tan(a + bx)) \log\left(\frac{-1+c+d \tan(a+bx)}{-1+c-id}\right) + i \log(1 + i \tan(a + bx))}{1} \right)$$

input `Integrate[ArcTanh[c + d*Tan[a + b*x]],x]`

output

```
x*(ArcTanh[c + d*Tan[a + b*x]] + (2*a*Log[1 - c - d*Tan[a + b*x]] - I*Log[
1 - I*Tan[a + b*x]]*Log[(-1 + c + d*Tan[a + b*x])/(-1 + c - I*d)] + I*Log[
1 + I*Tan[a + b*x]]*Log[(-1 + c + d*Tan[a + b*x])/(-1 + c + I*d)] - 2*a*Lo
g[1 + c + d*Tan[a + b*x]] + I*Log[1 - I*Tan[a + b*x]]*Log[(1 + c + d*Tan[a
+ b*x])/(1 + c - I*d)] - I*Log[1 + I*Tan[a + b*x]]*Log[(1 + c + d*Tan[a +
b*x])/(1 + c + I*d)] + I*PolyLog[2, -((d*(-I + Tan[a + b*x]))/(-1 + c + I
*d))] - I*PolyLog[2, -((d*(-I + Tan[a + b*x]))/(1 + c + I*d))] - I*PolyLog
[2, -((d*(I + Tan[a + b*x]))/(-1 + c - I*d))] + I*PolyLog[2, -((d*(I + Tan
[a + b*x]))/(1 + c - I*d)))/(4*a - (2*I)*Log[1 - I*Tan[a + b*x]] + (2*I)*
Log[1 + I*Tan[a + b*x]]))
```

**Rubi [A] (verified)**Time = 0.73 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.48, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6813, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \tan(a + bx) + c) dx$$

↓ 6813



output

```
x*ArcTanh[c + d*Tan[a + b*x]] + b*(I*(1 - c) - d)*((x*Log[1 + ((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d)])/(2*b*(I*(1 - c) - d)) - ((I/4)*PolyLog[2, -(((1 - c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c - I*d))]/(b^2*(I*(1 - c) - d))) - b*(I + I*c + d)*((x*Log[1 + ((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d)])/(2*(I*(b + b*c) + b*d)) - ((I/4)*PolyLog[2, -(((1 + c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c + I*d))]/(b*(I*(b + b*c) + b*d)))
```

### Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 6813

```
Int[ArcTanh[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]], x_Symbol] := Simp[x*ArcTanh[c + d*Tan[a + b*x]], x] + (-Simp[I*b*(1 + c - I*d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 + c + I*d + (1 + c - I*d)*E^(2*I*a + 2*I*b*x))], x], x] + Simp[I*b*(1 - c + I*d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 - c - I*d + (1 - c + I*d)*E^(2*I*a + 2*I*b*x))], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[(c + I*d)^2, 1]
```



### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 556 vs.  $2(164) = 328$ .

Time = 3.42 (sec) , antiderivative size = 557, normalized size of antiderivative = 2.87

method	result
derivativedivides	$d \arctan(\tan(bx+a)) \operatorname{arctanh}(c+d \tan(bx+a)) - d^2 \left( \frac{\arctan\left(-\frac{c+d \tan(bx+a)}{d} + \frac{c}{d}\right) \ln\left(d\left(\frac{c+d \tan(bx+a)}{d} - \frac{c}{d}\right) + c-1\right)}{2d} - \arctan\left(\frac{c+d \tan(bx+a)}{d}\right) \right)$
default	$d \arctan(\tan(bx+a)) \operatorname{arctanh}(c+d \tan(bx+a)) - d^2 \left( \frac{\arctan\left(-\frac{c+d \tan(bx+a)}{d} + \frac{c}{d}\right) \ln\left(d\left(\frac{c+d \tan(bx+a)}{d} - \frac{c}{d}\right) + c-1\right)}{2d} - \arctan\left(\frac{c+d \tan(bx+a)}{d}\right) \right)$
risch	Expression too large to display

```
input int(arctanh(c+d*tan(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output 1/b/d*(d*arctan(tan(b*x+a))*arctanh(c+d*tan(b*x+a))-d^2*(1/2*arctan(-(c+d*
tan(b*x+a))/d+c/d)/d*ln(d*((c+d*tan(b*x+a))/d-c/d)+c-1)-1/2*arctan(-(c+d*t
an(b*x+a))/d+c/d)/d*ln(d*((c+d*tan(b*x+a))/d-c/d)+c+1)+1/4*I*ln(d*((c+d*ta
n(b*x+a))/d-c/d)+c+1)*(ln((I*d-d*((c+d*tan(b*x+a))/d-c/d))/(1+c+I*d))-ln((
I*d+d*((c+d*tan(b*x+a))/d-c/d))/(I*d-c-1)))/d+1/4*I*(dilog((I*d-d*((c+d*ta
n(b*x+a))/d-c/d))/(1+c+I*d))-dilog((I*d+d*((c+d*tan(b*x+a))/d-c/d))/(I*d-c
-1)))/d-1/4*I*ln(d*((c+d*tan(b*x+a))/d-c/d)+c-1)*(ln((I*d-d*((c+d*tan(b*x+
a))/d-c/d))/(I*d+c-1))-ln((I*d+d*((c+d*tan(b*x+a))/d-c/d))/(1-c+I*d)))/d-1
/4*I*(dilog((I*d-d*((c+d*tan(b*x+a))/d-c/d))/(I*d+c-1))-dilog((I*d+d*((c+d
*tan(b*x+a))/d-c/d))/(1-c+I*d)))/d)
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1185 vs.  $2(136) = 272$ .

Time = 0.15 (sec) , antiderivative size = 1185, normalized size of antiderivative = 6.11

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctanh(c+d*tan(b*x+a)),x, algorithm="fricas")`

output

```
1/8*(4*b*x*log(-(d*tan(b*x + a) + c + 1)/(d*tan(b*x + a) + c - 1)) - 2*(b*x + a)*log(-2*((I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c + 1)*d + (I*c^2 - 2*(c + 1)*d - I*d^2 + 2*I*c + I)*tan(b*x + a) - 2*c - 1)/((c^2 + d^2 + 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1)) - 2*(b*x + a)*log(-2*((-I*(c + 1)*d - d^2)*tan(b*x + a)^2 - c^2 + I*(c + 1)*d + (-I*c^2 - 2*(c + 1)*d + I*d^2 - 2*I*c - I)*tan(b*x + a) - 2*c - 1)/((c^2 + d^2 + 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*c + 1)) + 2*(b*x + a)*log(-2*((I*(c - 1)*d - d^2)*tan(b*x + a)^2 - c^2 - I*(c - 1)*d + (I*c^2 - 2*(c - 1)*d - I*d^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*c + 1)) + 2*(b*x + a)*log(-2*((-I*(c - 1)*d - d^2)*tan(b*x + a)^2 - c^2 + I*(c - 1)*d + (-I*c^2 - 2*(c - 1)*d + I*d^2 + 2*I*c - I)*tan(b*x + a) + 2*c - 1)/((c^2 + d^2 - 2*c + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*c + 1)) + 2*a*log(((I*(c + 1)*d + d^2)*tan(b*x + a)^2 - c^2 + I*(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) - 2*c - 1)/(tan(b*x + a)^2 + 1)) + 2*a*log(((I*(c + 1)*d - d^2)*tan(b*x + a)^2 + c^2 + I*(c + 1)*d + (I*c^2 + I*d^2 + 2*I*c + I)*tan(b*x + a) + 2*c + 1)/(tan(b*x + a)^2 + 1)) - 2*a*log(((I*(c - 1)*d + d^2)*tan(b*x + a)^2 - c^2 + I*(c - 1)*d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) + 2*c - 1)/(tan(b*x + a)^2 + 1)) - 2*a*log(((I*(c - 1)*d - d^2)*tan(b*x + a)^2 + c^2 + I*(c - 1)*d + (I*c^2 + I*d^2 - 2*I*c + I)*tan(b*x + a) - 2*c + 1)/(tan(b*x + a)^2 + 1)) - I*dilog(...
```

**Sympy [F]**

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input `integrate(atanh(c+d*tan(b*x+a)),x)`

output `Integral(atanh(c + d*tan(a + b*x)), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 372 vs.  $2(136) = 272$ .

Time = 0.20 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.92

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx$$

$$= \frac{4(bx + a) \operatorname{artanh}(d \tan(bx + a) + c) + \left( \operatorname{arctan}\left(\frac{d^2 \tan(bx+a) + (c+1)d}{c^2 + d^2 + 2c+1}, \frac{(c+1)d \tan(bx+a) + c^2 + 2c+1}{c^2 + d^2 + 2c+1}\right) - \operatorname{arctan}\left(\frac{d}{c+d}\right) \right)}{b}$$

input `integrate(arctanh(c+d*tan(b*x+a)),x, algorithm="maxima")`

output `1/4*(4*(b*x + a)*arctanh(d*tan(b*x + a) + c) + (arctan2((d^2*tan(b*x + a) + (c + 1)*d)/(c^2 + d^2 + 2*c + 1), ((c + 1)*d*tan(b*x + a) + c^2 + 2*c + 1)/(c^2 + d^2 + 2*c + 1)) - arctan2((d^2*tan(b*x + a) + (c - 1)*d)/(c^2 + d^2 - 2*c + 1), ((c - 1)*d*tan(b*x + a) + c^2 - 2*c + 1)/(c^2 + d^2 - 2*c + 1)))*log(tan(b*x + a)^2 + 1) - (b*x + a)*log((d^2*tan(b*x + a)^2 + 2*(c + 1)*d*tan(b*x + a) + c^2 + 2*c + 1)/(c^2 + d^2 + 2*c + 1)) + (b*x + a)*log((d^2*tan(b*x + a)^2 + 2*(c - 1)*d*tan(b*x + a) + c^2 - 2*c + 1)/(c^2 + d^2 - 2*c + 1)) - I*dilog(-(I*d*tan(b*x + a) - d)/(I*c + d + I)) + I*dilog(-(I*d*tan(b*x + a) - d)/(I*c + d - I)) - I*dilog((I*d*tan(b*x + a) + d)/(-I*c + d + I)) + I*dilog((I*d*tan(b*x + a) + d)/(-I*c + d - I)))/b`

**Giac [F]**

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{artanh}(d \tan(bx + a) + c) dx$$

input `integrate(arctanh(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*tan(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{atanh}(c + d \tan(a + bx)) dx$$

input `int(atanh(c + d*tan(a + b*x)),x)`

output `int(atanh(c + d*tan(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(c + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a)d + c) dx$$

input `int(atanh(c+d*tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)*d + c),x)`

### 3.320 $\int \frac{\operatorname{arctanh}(c+d \tan(a+bx))}{x} dx$

Optimal result	2240
Mathematica [N/A]	2240
Rubi [N/A]	2241
Maple [N/A]	2241
Fricas [N/A]	2242
Sympy [N/A]	2242
Maxima [N/A]	2242
Giac [N/A]	2243
Mupad [N/A]	2243
Reduce [N/A]	2244

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x}, x\right)$$

output

```
Defer(Int)(arctanh(c+d*tan(b*x+a))/x,x)
```

#### Mathematica [N/A]

Not integrable

Time = 2.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx$$

input

```
Integrate[ArcTanh[c + d*Tan[a + b*x]]/x,x]
```

output

```
Integrate[ArcTanh[c + d*Tan[a + b*x]]/x, x]
```

**Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \tan(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \tan(a + bx) + c)}{x} dx$$

input `Int[ArcTanh[c + d*Tan[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(c + d \tan(bx + a))}{x} dx$$

input `int(arctanh(c+d*tan(b*x+a))/x,x)`

output `int(arctanh(c+d*tan(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*tan(b*x + a) + c)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \tan(a + bx))}{x} dx$$

input `integrate(atanh(c+d*tan(b*x+a))/x,x)`

output `Integral(atanh(c + d*tan(a + b*x))/x, x)`

**Maxima [N/A]**

Not integrable

Time = 3.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tan(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctanh(d*tan(b*x + a) + c)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.72 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*tan(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*tan(b*x + a) + c)/x, x)`

### Mupad [N/A]

Not integrable

Time = 3.97 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \tan(a + bx))}{x} dx$$

input `int(atanh(c + d*tan(a + b*x))/x,x)`

output `int(atanh(c + d*tan(a + b*x))/x, x)`



**Reduce [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tan(bx + a) d + c)}{x} dx$$

input `int(atanh(c+d*tan(b*x+a))/x,x)`output `int(atanh(tan(a + b*x)*d + c)/x,x)`

### 3.321 $\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$

Optimal result	2245
Mathematica [A] (verified)	2246
Rubi [A] (verified)	2246
Maple [C] (warning: unable to verify)	2249
Fricas [B] (verification not implemented)	2250
Sympy [F]	2251
Maxima [B] (verification not implemented)	2251
Giac [F]	2252
Mupad [F(-1)]	2252
Reduce [F]	2253

#### Optimal result

Integrand size = 20, antiderivative size = 170

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(1 - id + d \tan(a + bx)) - \frac{1}{6} x^3 \log(1 + (1 - id)e^{2ia+2ibx}) + \frac{ix^2 \operatorname{PolyLog}(2, -((1 - id)e^{2ia+2ibx}))}{4b} - \frac{x \operatorname{PolyLog}(3, -((1 - id)e^{2ia+2ibx}))}{4b^2} - \frac{i \operatorname{PolyLog}(4, -((1 - id)e^{2ia+2ibx}))}{8b^3}$$

output

```
1/12*I*b*x^4+1/3*x^3*arctanh(1-I*d+d*tan(b*x+a))-1/6*x^3*ln(1+(1-I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x^2*polylog(2,-(1-I*d)*exp(2*I*a+2*I*b*x))/b-1/4*x*polylog(3,-(1-I*d)*exp(2*I*a+2*I*b*x))/b^2-1/8*I*polylog(4,-(1-I*d)*exp(2*I*a+2*I*b*x))/b^3
```

**Mathematica [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.91

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \frac{1}{3} x^3 \operatorname{arctanh}(1 - id + d \tan(a + bx)) - \frac{4b^3 x^3 \log\left(1 + \frac{ie^{-2i(a+bx)}}{i+d}\right) + 6ib^2 x^2 \operatorname{PolyLog}\left(2, -\frac{ie^{-2i(a+bx)}}{i+d}\right) + 6bx \operatorname{PolyLog}\left(3, -\frac{ie^{-2i(a+bx)}}{i+d}\right) - 3i \operatorname{PolyLog}\left(4, -\frac{ie^{-2i(a+bx)}}{i+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 - I*d + d*Tan[a + b*x]],x]
```

output

```
(x^3*ArcTanh[1 - I*d + d*Tan[a + b*x]])/3 - (4*b^3*x^3*Log[1 + I/((I + d)*E^((2*I)*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, (-I)/((I + d)*E^((2*I)*(a + b*x)))] + 6*b*x*PolyLog[3, (-I)/((I + d)*E^((2*I)*(a + b*x)))] - (3*I)*PolyLog[4, (-I)/((I + d)*E^((2*I)*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.89 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.28, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {6817, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d \tan(a + bx) - id + 1) dx$$

$$\downarrow \text{6817}$$

$$\frac{1}{3} ib \int \frac{x^3}{e^{2ia+2ibx}(1-id)+1} dx + \frac{1}{3} x^3 \operatorname{arctanh}(d \tan(a + bx) - id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3} ib \left( \frac{x^4}{4} - (1-id) \int \frac{e^{2ia+2ibx} x^3}{e^{2ia+2ibx}(1-id)+1} dx \right) + \frac{1}{3} x^3 \operatorname{arctanh}(d \tan(a + bx) - id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1-id) \left( \frac{x^3 \log(1+(1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{3 \int x^2 \log(e^{2ia+2ibx}(1-id)+1) dx}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 3011

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1-id) \left( \frac{x^3 \log(1+(1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{b} \right)}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 7163

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1-id) \left( \frac{x^3 \log(1+(1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b} - \frac{i \left( \frac{\int \operatorname{PolyLog}(3, -((1-id)e^{2ia+2ibx}))}{2b} \right)}{2b(d+i)} \right)}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 2720

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1-id) \left( \frac{x^3 \log(1+(1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b} - \frac{i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(3, -((1-id)e^{2ia+2ibx}))}{4} \right)}{2b(d+i)} \right)}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 7143

$$\frac{1}{3} x^3 \operatorname{arctanh}(d \tan(a + bx) - id + 1) + \frac{\frac{1}{3} ib \left( \frac{x^4}{4} - (1 - id) \left( \frac{x^3 \log(1 + (1 - id)e^{2ia + 2ibx}}{2b(d + i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((1 - id)e^{2ia + 2ibx})}{2b})}{2b(d + i)} - \frac{i \left( \frac{\operatorname{PolyLog}(4, -((1 - id)e^{2ia + 2ibx})}{4b^2})}{2b(d + i)} \right)}{2b(d + i)} \right)}{2b(d + i)} \right)}{2b(d + i)}$$

input `Int[x^2*ArcTanh[1 - I*d + d*Tan[a + b*x]],x]`

output `(x^3*ArcTanh[1 - I*d + d*Tan[a + b*x])/3 + (I/3)*b*(x^4/4 - (1 - I*d)*((x^3*Log[1 + (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(2*b*(I + d)) - (3*(((I/2)*x^2*PolyLog[2, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x)]))/b - (I*(((1/2*I)*x*PolyLog[3, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x)]))/b + PolyLog[4, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b))/(2*b*(I + d)))`

### Defintions of rubi rules used

rule 2615 `Int[(((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6817 `Int[ArcTanh[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.36 (sec) , antiderivative size = 2277, normalized size of antiderivative = 13.39

method	result	size
risch	Expression too large to display	2277

input `int(x^2*arctanh(1-I*d+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

1/12*I*b*x^4+1/2/b^2*d/(I+d)*ln(1-I*(I+d)*exp(2*I*(b*x+a)))*a^2*x-1/2/b^2*
d*a^2/(I+d)*ln(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x-1/2/b^2*d*a^2/(I+d)*
ln(1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x+1/2*I/b^3*d*a^2/(I+d)*dilog(1+I*
exp(I*(b*x+a))*(-I*(I+d))^(1/2))+1/2*I/b^2/(I+d)*ln(1-I*(I+d)*exp(2*I*(b*x
+a)))*a^2*x+1/4*I/b*d/(I+d)*polylog(2,I*(I+d)*exp(2*I*(b*x+a)))*x^2-1/2*I/
b^2*a^2/(I+d)*ln(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x-1/2*I/b^2*a^2/(I+d
)*ln(1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x-1/4*I/b^3*a^2*d/(I+d)*polylog(
2,I*(I+d)*exp(2*I*(b*x+a)))+1/2*I/b^3*d*a^2/(I+d)*dilog(1-I*exp(I*(b*x+a))
*(-I*(I+d))^(1/2))-1/3*x^3*ln(exp(I*(b*x+a)))+1/6*x^3*ln(I*exp(2*I*(b*x+a)
)+exp(2*I*(b*x+a))*d+I)-1/12*(2*I*Pi+2*ln(d)+I*Pi*csgn(I/(exp(2*I*(b*x+a)
+1))*csgn(I/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)))^2+I*Pi*csgn(1/(exp(2*I*
(b*x+a))+1)*exp(2*I*(b*x+a))*d)^3+I*Pi*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*I*
(b*x+a))*d+I)/(exp(2*I*(b*x+a))+1))*csgn((I*exp(2*I*(b*x+a))+exp(2*I*(b*x+
a))*d+I)/(exp(2*I*(b*x+a))+1))-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(I
*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(
1/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a))*d)^2-I*Pi*csgn(I*(I*exp(2*I*(b*x+a)
)+exp(2*I*(b*x+a))*d+I))*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)
/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*
d+I)/(exp(2*I*(b*x+a))+1))*csgn((I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)/
(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*exp(I*(b*x+a)))^2*csgn(I*exp(2*I*(b...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 345 vs.  $2(118) = 236$ .

Time = 0.10 (sec) , antiderivative size = 345, normalized size of antiderivative = 2.03

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$$

$$= \frac{ib^4x^4 + 2b^3x^3 \log\left(-\frac{((d+i)e^{2ibx+2ia}+i)e^{-2ibx-2ia}}{d}\right) + 6ib^2x^2 \operatorname{Li}_2\left(\frac{1}{2}\sqrt{4id-4}e^{(ibx+ia)}\right) + 6ib^2x^2 \operatorname{Li}_2\left(-\frac{1}{2}\sqrt{4id-4}e^{(ibx+ia)}\right)}{1}$$

input

```
integrate(x^2*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(I*b^4*x^4 + 2*b^3*x^3*log(-(d + I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I
*b*x - 2*I*a)/d) + 6*I*b^2*x^2*dilog(1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a))
+ 6*I*b^2*x^2*dilog(-1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - I*a^4 + 2*a^3*
log(1/2*(2*(d + I)*e^(I*b*x + I*a) + I*sqrt(4*I*d - 4))/(d + I)) + 2*a^3*
log(1/2*(2*(d + I)*e^(I*b*x + I*a) - I*sqrt(4*I*d - 4))/(d + I)) - 12*b*x*
polylog(3, 1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - 12*b*x*polylog(3, -1/2*
sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - 2*(b^3*x^3 + a^3)*log(1/2*sqrt(4*I*d - 4)
*e^(I*b*x + I*a) + 1) - 2*(b^3*x^3 + a^3)*log(-1/2*sqrt(4*I*d - 4)*e^(I*b*
x + I*a) + 1) - 12*I*polylog(4, 1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - 12*
I*polylog(4, -1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)))/b^3
```

## Sympy [F]

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(d \tan(a + bx) - id + 1) dx$$

input

```
integrate(x**2*atanh(1-I*d+d*tan(b*x+a)),x)
```

output

```
Integral(x**2*atanh(d*tan(a + b*x) - I*d + 1), x)
```

## Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 343 vs.  $2(118) = 236$ .

Time = 0.06 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.02

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$$

$$= \frac{12 \left( (bx+a)^3 - 3(bx+a)^2 a + 3(bx+a) a^2 \right) \operatorname{artanh}(d \tan(bx+a) - id + 1)}{b^2} - \frac{-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(-4i(bx+a)^3 + 9i(bx+a)^2 a - 3i a^3)}{b^2}$$

input

```
integrate(x^2*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="maxima")
```



output

```
1/36*(12*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctanh(d*tan(b
*x + a) - I*d + 1)/b^2 - (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*
*x + a)^2*a^2 - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a - 9*I*(b*x + a)*a^2
)*arctan2(-d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), d*sin(2*b*x + 2*a) + cos
(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog
((I*d - 1)*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*
*x + a)*a^2)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^
2 + 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog
(3, (I*d - 1)*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, (I*d - 1)*e^(2*I*b*x +
2*I*a)))/b^2)/b
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x^2 \operatorname{artanh}(d \tan(bx + a) - id + 1) dx$$

input

```
integrate(x^2*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctanh(d*tan(b*x + a) - I*d + 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(d \tan(a + bx) + 1 - d li) dx$$

input

```
int(x^2*atanh(d*tan(a + b*x) - d*1i + 1),x)
```

output

```
int(x^2*atanh(d*tan(a + b*x) - d*1i + 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a)) d - di + 1) x^2 dx$$

input `int(x^2*atanh(1-I*d+d*tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)*d - d*i + 1)*x**2,x)`

### 3.322 $\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$

Optimal result	2254
Mathematica [A] (verified)	2255
Rubi [A] (verified)	2255
Maple [C] (warning: unable to verify)	2258
Fricas [B] (verification not implemented)	2259
Sympy [F]	2259
Maxima [B] (verification not implemented)	2260
Giac [F]	2260
Mupad [F(-1)]	2261
Reduce [F]	2261

#### Optimal result

Integrand size = 18, antiderivative size = 133

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \frac{1}{6} ibx^3 + \frac{1}{2} x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) - \frac{1}{4} x^2 \log(1 + (1 - id)e^{2ia+2ibx}) + \frac{ix \operatorname{PolyLog}(2, -((1 - id)e^{2ia+2ibx}))}{4b} - \frac{\operatorname{PolyLog}(3, -((1 - id)e^{2ia+2ibx}))}{8b^2}$$

output

```
1/6*I*b*x^3+1/2*x^2*arctanh(1-I*d+d*tan(b*x+a))-1/4*x^2*ln(1+(1-I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x*polylog(2,-(1-I*d)*exp(2*I*a+2*I*b*x))/b-1/8*polylog(3,-(1-I*d)*exp(2*I*a+2*I*b*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.89

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \frac{1}{2} x^2 \operatorname{arctanh}(1 - id + d \tan(a + bx)) - \frac{2b^2 x^2 \log\left(1 + \frac{ie^{-2i(a+bx)}}{i+d}\right) + 2ibx \operatorname{PolyLog}\left(2, -\frac{ie^{-2i(a+bx)}}{i+d}\right) + \operatorname{PolyLog}\left(3, -\frac{ie^{-2i(a+bx)}}{i+d}\right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 - I*d + d*Tan[a + b*x]], x]
```

output

```
(x^2*ArcTanh[1 - I*d + d*Tan[a + b*x]])/2 - (2*b^2*x^2*Log[1 + I/((I + d)*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, (-I)/((I + d)*E^((2*I)*(a + b*x)))] + PolyLog[3, (-I)/((I + d)*E^((2*I)*(a + b*x)))])/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.70 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6817, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \tan(a + bx) - id + 1) dx$$

$$\downarrow \text{6817}$$

$$\frac{1}{2} ib \int \frac{x^2}{e^{2ia+2ibx}(1-id)+1} dx + \frac{1}{2} x^2 \operatorname{arctanh}(d \tan(a + bx) - id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} ib \left( \frac{x^3}{3} - (1-id) \int \frac{e^{2ia+2ibx} x^2}{e^{2ia+2ibx}(1-id)+1} dx \right) + \frac{1}{2} x^2 \operatorname{arctanh}(d \tan(a + bx) - id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1-id) \left( \frac{x^2 \log(1 + (1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{\int x \log(e^{2ia+2ibx}(1-id) + 1) dx}{b(d+i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 3011

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1-id) \left( \frac{x^2 \log(1 + (1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b}}{b(d+i)} - \frac{i \int \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 2720

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1-id) \left( \frac{x^2 \log(1 + (1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b}}{b(d+i)} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{4b^2} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) - id + 1)$$

↓ 7143

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1-id) \left( \frac{x^2 \log(1 + (1-id)e^{2ia+2ibx})}{2b(d+i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -((1-id)e^{2ia+2ibx}))}{2b}}{b(d+i)} - \frac{\frac{1}{2}x^2 \operatorname{arctanh}(d \tan(a+bx) - id + 1) + \operatorname{PolyLog}(3, -((1-id)e^{2ia+2ibx}))}{4b^2}}{b(d+i)} \right) \right)$$

input `Int[x*ArcTanh[1 - I*d + d*Tan[a + b*x]],x]`

output `(x^2*ArcTanh[1 - I*d + d*Tan[a + b*x])/2 + (I/2)*b*(x^3/3 - (1 - I*d)*((x^2*Log[1 + (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(2*b*(I + d)) - (((I/2)*x*PolyLog[2, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x)))]/b - PolyLog[3, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2))/(b*(I + d))))`

## Definitions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 6817

```
Int[ArcTanh[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.32 (sec) , antiderivative size = 2187, normalized size of antiderivative = 16.44

method	result	size
risch	Expression too large to display	2187

input `int(x*arctanh(1-I*d+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/2/b^2*d*a^2/(I+d)*ln(1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))-1/4/b^2*d/(I+d)
)*ln(1-I*(I+d)*exp(2*I*(b*x+a)))*a^2-1/4*I/b^2/(I+d)*ln(1-I*(I+d)*exp(2*I*
(b*x+a)))*a^2+1/2*I/b^2*a^2/(I+d)*ln(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))+
1/2*I/b^2*a^2/(I+d)*ln(1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))-1/4*I/b^2*a^2/
(I+d)*ln(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)+1/6*I*b*x^3-1/4/b^2*a^2*
d/(I+d)*ln(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)+1/2/b^2*d*a^2/(I+d)*ln
(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))+1/4*I/b*d/(I+d)*polylog(2,I*(I+d)*ex
p(2*I*(b*x+a)))*x+1/4*I/b^2*d/(I+d)*polylog(2,I*(I+d)*exp(2*I*(b*x+a)))*a+
1/2*I/b*a/(I+d)*ln(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x+1/2*I/b*a/(I+d)*
ln(1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))*x+1/2/b*d*a/(I+d)*ln(1+I*exp(I*(b*x+a))*(-I*(I+d))
^(1/2))*x-1/2/b*d/(I+d)*ln(1-I*(I+d)*exp(2*I*(b*x+a)))*a*x-1/2*I/b^2*d*a/(
I+d)*dilog(1-I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))-1/2*I/b^2*d*a/(I+d)*dilog(
1+I*exp(I*(b*x+a))*(-I*(I+d))^(1/2))-1/2*I/b/(I+d)*ln(1-I*(I+d)*exp(2*I*(b
*x+a)))*a*x-1/8*(2*I*Pi+2*ln(d)+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I/(
exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)))^2+I*Pi*csgn(1/(exp(2*I*(b*x+a))+1)*e
xp(2*I*(b*x+a))*d)^3+I*Pi*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)
/(exp(2*I*(b*x+a))+1))*csgn((I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d+I)/(exp
(2*I*(b*x+a))+1))-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(I*exp(2*I*(b*x
+a))+exp(2*I*(b*x+a))*d+I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(1/(exp(2*I...
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs.  $2(93) = 186$ .

Time = 0.10 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.20

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$$

$$= \frac{2i b^3 x^3 + 3 b^2 x^2 \log\left(-\frac{((d+i)e^{2i bx+2i a}+i)e^{(-2i bx-2i a)}}{d}\right) + 2i a^3 + 6i bx \operatorname{Li}_2\left(\frac{1}{2}\sqrt{4i d - 4}e^{(i bx+i a)}\right) + 6i bx \operatorname{Li}_2\left(-\frac{1}{2}\sqrt{4i d - 4}e^{(i bx+i a)}\right)}{b^2}$$

input `integrate(x*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*I*b^3*x^3 + 3*b^2*x^2*log(-((d + I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d) + 2*I*a^3 + 6*I*b*x*dilog(1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) + 6*I*b*x*dilog(-1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - 3*a^2*log(1/2*(2*(d + I)*e^(I*b*x + I*a) + I*sqrt(4*I*d - 4))/(d + I)) - 3*a^2*log(1/2*(2*(d + I)*e^(I*b*x + I*a) - I*sqrt(4*I*d - 4))/(d + I)) - 3*(b^2*x^2 - a^2)*log(1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a) + 1) - 3*(b^2*x^2 - a^2)*log(-1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a) + 1) - 6*polylog(3, 1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) - 6*polylog(3, -1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x \operatorname{atanh}(d \tan(a + bx) - id + 1) dx$$

input `integrate(x*atanh(1-I*d+d*tan(b*x+a)),x)`

output `Integral(x*atanh(d*tan(a + b*x) - I*d + 1), x)`



**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 248 vs.  $2(93) = 186$ .

Time = 0.05 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.86

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$$

$$= \frac{12((bx+a)^2 - 2(bx+a)a) \operatorname{artanh}(d \tan(bx+a) - id + 1)}{b} - \frac{-4i(bx+a)^3 + 12i(bx+a)^2 a - 6i b x \operatorname{Li}_2((i d - 1)e^{2i bx + 2i a}) - 6(-i(bx+a)^2 + 2i(bx+a)a)}{b}$$

input `integrate(x*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="maxima")`

output `1/24*(12*((b*x + a)^2 - 2*(b*x + a)*a)*arctanh(d*tan(b*x + a) - I*d + 1)/b - (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog((I*d - 1)*e^(2*I*b*x + 2*I*a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(-d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), d*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^2 + 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1) + 3*polylog(3, (I*d - 1)*e^(2*I*b*x + 2*I*a)))/b)/b`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x \operatorname{artanh}(d \tan(bx + a) - id + 1) dx$$

input `integrate(x*arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*tan(b*x + a) - I*d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int x \operatorname{atanh}(d \tan(a + bx) + 1 - d1i) dx$$

input `int(x*atanh(d*tan(a + b*x) - d*1i + 1),x)`

output `int(x*atanh(d*tan(a + b*x) - d*1i + 1), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a) d - di + 1) x dx$$

input `int(x*atanh(1-I*d+d*tan(b*x+a)),x)`

output `int(atanh(tan(a + b*x)*d - d*i + 1)*x,x)`

### 3.323 $\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$

Optimal result	2262
Mathematica [B] (warning: unable to verify)	2262
Rubi [A] (verified)	2263
Maple [B] (verified)	2266
Fricas [B] (verification not implemented)	2267
Sympy [F]	2267
Maxima [B] (verification not implemented)	2268
Giac [F]	2268
Mupad [F(-1)]	2269
Reduce [F]	2269

#### Optimal result

Integrand size = 16, antiderivative size = 93

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \frac{1}{2}ibx^2 + x \operatorname{arctanh}(1 - id + d \tan(a + bx)) - \frac{1}{2}x \log(1 + (1 - id)e^{2ia+2ibx}) + \frac{i \operatorname{PolyLog}(2, -((1 - id)e^{2ia+2ibx}))}{4b}$$

output

```
1/2*I*b*x^2+x*arctanh(1-I*d+d*tan(b*x+a))-1/2*x*ln(1+(1-I*d)*exp(2*I*a+2*I*b*x))+1/4*I*polylog(2,-(1-I*d)*exp(2*I*a+2*I*b*x))/b
```

#### Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 766 vs. 2(93) = 186.

Time = 7.50 (sec) , antiderivative size = 766, normalized size of antiderivative = 8.24

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = x \operatorname{arctanh}(1 - id + d \tan(a + bx)) + \frac{x \left( 2ibx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) - \log \left( \frac{\sec(bx)(\cos(a) - i \sin(a))((2i+d) \cos(a+bx) + id \sin(a+bx))}{2(i+d)} \right) \right) \log(1 - \dots)}{((2i + d) \cos(a + bx) + id \sin(a + bx)) \left( \frac{i \log(1 + i \tan(bx)) \sec(bx)(d \cos(a) + i(2i+d) \sin(a))}{(2i+d) \cos(a+bx) + id \sin(a+bx)} + \dots \right)}$$

input `Integrate[ArcTanh[1 - I*d + d*Tan[a + b*x]],x]`

output

```
x*ArcTanh[1 - I*d + d*Tan[a + b*x]] + (x*((2*I)*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] - Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]))/(2*(I + d))]*Log[1 - I*Tan[b*x]] + Log[(Sec[b*x]*((2 - I*d)*Cos[a + b*x] + d*Sin[a + b*x]))/(2*Cos[a] - (2*I)*Sin[a])]*Log[1 + I*Tan[b*x]] - PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] - PolyLog[2, (Sec[b*x]*(d*Cos[a] + I*(2*I + d)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*(I + d))] + PolyLog[2, -1/2*((Cos[a] + I*Sin[a])*(d*Cos[a] + I*(2*I + d)*Sin[a])*(-I + Tan[b*x]))]*Sec[a + b*x]^2*(Cos[b*x] + I*Sin[b*x])*(I*Cos[b*x] + Sin[b*x])*((2 - I*d)*Cos[a + b*x] + d*Sin[a + b*x])/((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x])*((I*Log[1 + I*Tan[b*x]]*Sec[b*x]*(d*Cos[a] + I*(2*I + d)*Sin[a]))/((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]) + (Log[1 - I*Tan[b*x]]*Sec[b*x]*((-I)*d*Cos[a] + (2*I + d)*Sin[a]))/((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]) + 2*b*x*(1 - I*Tan[b*x]) + (Log[(Sec[b*x]*((2 - I*d)*Cos[a + b*x] + d*Sin[a + b*x]))/(2*Cos[a] - (2*I)*Sin[a])]*Sec[b*x]^2)/(-I + Tan[b*x]) - (Log[1 + ((Cos[a] + I*Sin[a])*(d*Cos[a] + I*(2*I + d)*Sin[a])*(-I + Tan[b*x]))/2]*Sec[b*x]^2)/(-I + Tan[b*x]) + Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]))/(2*(I + d))]*(-I + Tan[b*x]) - (Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]))/(2*(I + d))]*Sec[b*x]^2)/(I + Tan[b*x]))*(-I + Tan[a + b*x])*(2 - I*d + d*Tan[a + b*x]))
```

### Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.32, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {6809, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \tan(a + bx) - id + 1) dx$$

↓ 6809

$$ib \int \frac{x}{e^{2ia+2ibx}(1-id)+1} dx + x \operatorname{arctanh}(d \tan(a + bx) - id + 1)$$

$$\begin{aligned}
& \downarrow 2615 \\
& ib \left( \frac{x^2}{2} - (1 - id) \int \frac{e^{2ia+2ibx} x}{e^{2ia+2ibx}(1-id) + 1} dx \right) + x \operatorname{arctanh}(d \tan(a + bx) - id + 1) \\
& \downarrow 2620 \\
& ib \left( \frac{x^2}{2} - (1 - id) \left( \frac{x \log(1 + (1 - id)e^{2ia+2ibx})}{2b(d+i)} - \frac{\int \log(e^{2ia+2ibx}(1-id) + 1) dx}{2b(d+i)} \right) \right) + \\
& \quad x \operatorname{arctanh}(d \tan(a + bx) - id + 1) \\
& \downarrow 2715 \\
& ib \left( \frac{x^2}{2} - (1 - id) \left( \frac{i \int e^{-2ia-2ibx} \log(e^{2ia+2ibx}(1-id) + 1) de^{2ia+2ibx}}{4b^2(d+i)} + \frac{x \log(1 + (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \\
& \quad x \operatorname{arctanh}(d \tan(a + bx) - id + 1) \\
& \downarrow 2838 \\
& ib \left( \frac{x^2}{2} - (1 - id) \left( \frac{x \operatorname{arctanh}(d \tan(a + bx) - id + 1) + x \log(1 + (1 - id)e^{2ia+2ibx})}{2b(d+i)} - \frac{i \operatorname{PolyLog}(2, -((1 - id)e^{2ia+2ibx}))}{4b^2(d+i)} \right) \right)
\end{aligned}$$

input `Int[ArcTanh[1 - I*d + d*Tan[a + b*x]],x]`

output `x*ArcTanh[1 - I*d + d*Tan[a + b*x]] + I*b*(x^2/2 - (1 - I*d)*((x*Log[1 + (1 - I*d)*E^((2*I)*a + (2*I)*b*x)])/(2*b*(I + d)) - ((I/4)*PolyLog[2, -((1 - I*d)*E^((2*I)*a + (2*I)*b*x))])/(b^2*(I + d))))`

## Definitions of rubi rules used

rule 2615  $\text{Int}[\left(\frac{(c_.) + (d_.)x^{(m_.)}}{(a_.) + (b_.)\left((F_.)^{(g_.)}\left((e_.) + (f_.)x\right)\right)}\right)^{(n_.)}], x\_Symbol] \rightarrow \text{Simp}[(c + dx)^{(m+1)}/(a*d*(m+1)), x] - \text{Simp}[b/a \int (c + dx)^m \left(\frac{F^{(g*(e+fx))}}{(a + b*(F^{(g*(e+fx)))})^n}\right)^n dx, x] /;$  FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

rule 2620  $\text{Int}[\left(\frac{\left(\frac{(F_.)^{(g_.)}\left((e_.) + (f_.)x\right)}{(a_.) + (b_.)\left((F_.)^{(g_.)}\left((e_.) + (f_.)x\right)\right)}\right)^{(n_.)}\left((c_.) + (d_.)x^{(m_.)}\right)}{\left((a_.) + (b_.)\left((F_.)^{(g_.)}\left((e_.) + (f_.)x\right)\right)}\right)^{(n_.)}], x\_Symbol] \rightarrow \text{Simp}[\left(\frac{(c + dx)^m}{(b*f*g*n*\text{Log}[F])}\right)*\text{Log}[1 + b*\left(\frac{F^{(g*(e+fx))}}{a}\right)^n], x] - \text{Simp}[d*\left(\frac{m}{(b*f*g*n*\text{Log}[F])}\right) \int (c + dx)^{(m-1)}*\text{Log}[1 + b*\left(\frac{F^{(g*(e+fx))}}{a}\right)^n] dx, x] /;$  FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

rule 2715  $\text{Int}[\text{Log}[(a_.) + (b_.)\left((F_.)^{(e_.)}\left((c_.) + (d_.)x\right)\right)^{(n_.)}], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\int \text{Log}[a + b*x]/x dx, x, (F^{(e*(c+dx))})^n], x] /;$  FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

rule 2838  $\text{Int}[\text{Log}[(c_.)\left((d_.) + (e_.)x^{(n_.)}\right)]/(x_)], x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /;$  FreeQ[{c, d, e, n}, x] && EqQ[c\*d, 1]

rule 6809  $\text{Int}[\text{ArcTanh}[(c_.) + (d_.)\text{Tan}[(a_.) + (b_.)x]], x\_Symbol] \rightarrow \text{Simp}[x*\text{ArcTanh}[c + d*\text{Tan}[a + b*x]], x] + \text{Simp}[I*b \int x/(c + I*d + c*E^{(2*I*a + 2*I*b*x)}) dx, x] /;$  FreeQ[{a, b, c, d}, x] && EqQ[(c + I\*d)^2, 1]

### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 306 vs.  $2(76) = 152$ .

Time = 3.37 (sec) , antiderivative size = 307, normalized size of antiderivative = 3.30

method	result
derivativedivides	$\frac{-\frac{i \operatorname{arctanh}(1-id+d \tan(bx+a))d \ln(-id+d \tan(bx+a))}{2} + \frac{i \operatorname{arctanh}(1-id+d \tan(bx+a))d \ln(id+d \tan(bx+a))}{2}}{d^2} - \left( \frac{i \operatorname{dilog}\left(\frac{i(-id+d \tan(bx+a))}{id+d \tan(bx+a)}\right)}{d^2} \right)$
default	$\frac{-\frac{i \operatorname{arctanh}(1-id+d \tan(bx+a))d \ln(-id+d \tan(bx+a))}{2} + \frac{i \operatorname{arctanh}(1-id+d \tan(bx+a))d \ln(id+d \tan(bx+a))}{2}}{d^2} - \left( \frac{i \operatorname{dilog}\left(\frac{i(-id+d \tan(bx+a))}{id+d \tan(bx+a)}\right)}{d^2} \right)$
risch	Expression too large to display

```
input int(arctanh(1-I*d+d*tan(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output 1/b/d*(-1/2*I*arctanh(1-I*d+d*tan(b*x+a))*d*ln(-I*d+d*tan(b*x+a))+1/2*I*arctanh(1-I*d+d*tan(b*x+a))*d*ln(I*d+d*tan(b*x+a))-1/2*d^2*(-I/d*(1/2*dilog(1/2*I*(-I*d+d*tan(b*x+a))/d)+1/2*ln(I*d+d*tan(b*x+a))*ln(1/2*I*(-I*d+d*tan(b*x+a))/d)-1/2*dilog(I*(I*d+d*tan(b*x+a)-I*(2*I+2*d))/(2*I+2*d))-1/2*ln(I*d+d*tan(b*x+a))*ln(I*(I*d+d*tan(b*x+a)-I*(2*I+2*d))/(2*I+2*d)))+I/d*(1/4*ln(-I*d+d*tan(b*x+a))^2-1/2*dilog(1-1/2*I*d+1/2*d*tan(b*x+a))-1/2*ln(-I*d+d*tan(b*x+a))*ln(1-1/2*I*d+1/2*d*tan(b*x+a))))
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs.  $2(65) = 130$ .

Time = 0.10 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.34

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx$$

$$= \frac{i b^2 x^2 + bx \log\left(-\frac{((d+i)e^{2i bx+2i a}+i)e^{(-2i bx-2i a)}}{d}\right) - i a^2 - (bx + a) \log\left(\frac{1}{2} \sqrt{4i d - 4} e^{(i bx+i a)} + 1\right) - (bx + a)}{1}$$

input `integrate(arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="fricas")`

output `1/2*(I*b^2*x^2 + b*x*log(-((d + I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d) - I*a^2 - (b*x + a)*log(1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a) + 1) - (b*x + a)*log(-1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a) + 1) + a*log(1/2*(2*(d + I)*e^(I*b*x + I*a) + I*sqrt(4*I*d - 4))/(d + I)) + a*log(1/2*(2*(d + I)*e^(I*b*x + I*a) - I*sqrt(4*I*d - 4))/(d + I)) + I*dilog(1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)) + I*dilog(-1/2*sqrt(4*I*d - 4)*e^(I*b*x + I*a)))/b`

**Sympy [F]**

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{atanh}(d \tan(a + bx) - id + 1) dx$$

input `integrate(atanh(1-I*d+d*tan(b*x+a)),x)`

output `Integral(atanh(d*tan(a + b*x) - I*d + 1), x)`



**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 260 vs.  $2(65) = 130$ .

Time = 0.12 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.80

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx =$$

$$\frac{4(bx + a)d \left( \frac{\log(d \tan(bx+a) - id + 2)}{d} - \frac{\log(\tan(bx+a) - i)}{d} \right) + d \left( -\frac{2i \left( \log(d \tan(bx+a) - id + 2) \log\left(-\frac{id \tan(bx+a) + d + 2i}{2(d+i)} + 1\right) + \dots \right)}{d} \right)}{1}$$

input `integrate(arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="maxima")`

output

```
-1/8*(4*(b*x + a)*d*(log(d*tan(b*x + a) - I*d + 2)/d - log(tan(b*x + a) - I)/d) + d*(-2*I*(log(d*tan(b*x + a) - I*d + 2)*log(-1/2*(I*d*tan(b*x + a) + d + 2*I)/(d + I) + 1) + dilog(1/2*(I*d*tan(b*x + a) + d + 2*I)/(d + I)))/d + (2*I*log(d*tan(b*x + a) - I*d + 2)*log(tan(b*x + a) - I) - I*log(tan(b*x + a) - I)^2)/d - 2*I*(log(1/2*d*tan(b*x + a) - 1/2*I*d + 1)*log(tan(b*x + a) - I) + dilog(-1/2*d*tan(b*x + a) + 1/2*I*d))/d + 2*I*(log(tan(b*x + a) - I)*log(-1/2*I*tan(b*x + a) + 1/2) + dilog(1/2*I*tan(b*x + a) + 1/2))/d) - 8*(b*x + a)*arctanh(d*tan(b*x + a) - I*d + 1))/b
```

**Giac [F]**

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{artanh}(d \tan(bx + a) - id + 1) dx$$

input `integrate(arctanh(1-I*d+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*tan(b*x + a) - I*d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{atanh}(d \tan(a + bx) + 1 - d i) dx$$

input `int(atanh(d*tan(a + b*x) - d*i + 1),x)`output `int(atanh(d*tan(a + b*x) - d*i + 1), x)`**Reduce [F]**

$$\int \operatorname{arctanh}(1 - id + d \tan(a + bx)) dx = \int \operatorname{atanh}(\tan(bx + a) d - di + 1) dx$$

input `int(atanh(1-I*d+d*tan(b*x+a)),x)`output `int(atanh(tan(a + b*x)*d - d*i + 1),x)`

$$3.324 \quad \int \frac{\operatorname{arctanh}(1-id+d \tan(a+bx))}{x} dx$$

Optimal result	2270
Mathematica [N/A]	2270
Rubi [N/A]	2271
Maple [N/A]	2271
Fricas [N/A]	2272
Sympy [N/A]	2272
Maxima [N/A]	2272
Giac [N/A]	2273
Mupad [N/A]	2273
Reduce [N/A]	2274

### Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{\operatorname{arctanh}(1-id+d \tan(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1-id+d \tan(a+bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(1-I*d+d*tan(b*x+a))/x,x)`

### Mathematica [N/A]

Not integrable

Time = 0.81 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(1-id+d \tan(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1-id+d \tan(a+bx))}{x} dx$$

input `Integrate[ArcTanh[1 - I*d + d*Tan[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 - I*d + d*Tan[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \tan(a + bx) - id + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \tan(a + bx) - id + 1)}{x} dx$$

input `Int[ArcTanh[1 - I*d + d*Tan[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.49 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(bx + a))}{x} dx$$

input `int(arctanh(1-I*d+d*tan(b*x+a))/x,x)`

output `int(arctanh(1-I*d+d*tan(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) - id + 1)}{x} dx$$

input `integrate(arctanh(1-I*d+d*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*log(-((d + I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.80 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \tan(a + bx) - id + 1)}{x} dx$$

input `integrate(atanh(1-I*d+d*tan(b*x+a))/x,x)`

output `Integral(atanh(d*tan(a + b*x) - I*d + 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 3.80 (sec) , antiderivative size = 144, normalized size of antiderivative = 7.20

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) - id + 1)}{x} dx$$

input `integrate(arctanh(1-I*d+d*tan(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x + 1/4*(I*pi - 4*I*a - 2*log(d))*log(x) - 1/2*I*integrate(arctan2(-d
*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) - cos(2*b*x + 2*
a) - 1)/x, x) + 1/4*integrate(log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)
*sin(2*b*x + 2*a)^2 + 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1)/x, x)
```

**Giac [N/A]**

Not integrable

Time = 0.46 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \tan(bx + a) - id + 1)}{x} dx$$

input

```
integrate(arctanh(1-I*d+d*tan(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctanh(d*tan(b*x + a) - I*d + 1)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.61 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \tan(a + bx) + 1 - d1i)}{x} dx$$

input

```
int(atanh(d*tan(a + b*x) - d*1i + 1)/x,x)
```

output

```
int(atanh(d*tan(a + b*x) - d*1i + 1)/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - id + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\tan(bx + a) d - di + 1)}{x} dx$$

input `int(atanh(1-I*d+d*tan(b*x+a))/x,x)`output `int(atanh(tan(a + b*x)*d - d*i + 1)/x,x)`

### 3.325 $\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$

Optimal result	2275
Mathematica [A] (verified)	2276
Rubi [A] (verified)	2276
Maple [C] (warning: unable to verify)	2279
Fricas [B] (verification not implemented)	2280
Sympy [F]	2281
Maxima [B] (verification not implemented)	2281
Giac [F]	2282
Mupad [F(-1)]	2282
Reduce [F]	2283

#### Optimal result

Integrand size = 21, antiderivative size = 171

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(1 + id - d \tan(a + bx)) - \frac{1}{6} x^3 \log(1 + (1 + id)e^{2ia+2ibx}) + \frac{ix^2 \operatorname{PolyLog}(2, -((1 + id)e^{2ia+2ibx}))}{4b} - \frac{x \operatorname{PolyLog}(3, -((1 + id)e^{2ia+2ibx}))}{4b^2} - \frac{i \operatorname{PolyLog}(4, -((1 + id)e^{2ia+2ibx}))}{8b^3}$$

output

```
1/12*I*b*x^4-1/3*x^3*arctanh(-1-I*d+d*tan(b*x+a))-1/6*x^3*ln(1+(1+I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x^2*polylog(2,-(1+I*d)*exp(2*I*a+2*I*b*x))/b-1/4*x*polylog(3,-(1+I*d)*exp(2*I*a+2*I*b*x))/b^2-1/8*I*polylog(4,-(1+I*d)*exp(2*I*a+2*I*b*x))/b^3
```



**Mathematica [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.91

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \frac{1}{3} x^3 \operatorname{arctanh}(1 + id - d \tan(a + bx)) - \frac{4b^3 x^3 \log\left(1 - \frac{ie^{-2i(a+bx)}}{-i+d}\right) + 6ib^2 x^2 \operatorname{PolyLog}\left(2, \frac{ie^{-2i(a+bx)}}{-i+d}\right) + 6bx \operatorname{PolyLog}\left(3, \frac{ie^{-2i(a+bx)}}{-i+d}\right) - 3i \operatorname{PolyLog}\left(4, \frac{ie^{-2i(a+bx)}}{-i+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 + I*d - d*Tan[a + b*x]],x]
```

output

```
(x^3*ArcTanh[1 + I*d - d*Tan[a + b*x]])/3 - (4*b^3*x^3*Log[1 - I/((-I + d)*E^((2*I)*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, I/((-I + d)*E^((2*I)*(a + b*x)))] + 6*b*x*PolyLog[3, I/((-I + d)*E^((2*I)*(a + b*x)))] - (3*I)*PolyLog[4, I/((-I + d)*E^((2*I)*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**

Time = 0.93 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.30, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6817, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1) dx$$

$$\downarrow \text{6817}$$

$$\frac{1}{3} ib \int \frac{x^3}{e^{2ia+2ibx}(id+1)+1} dx + \frac{1}{3} x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3} ib \left( \frac{x^4}{4} - (1 + id) \int \frac{e^{2ia+2ibx} x^3}{e^{2ia+2ibx}(id+1)+1} dx \right) + \frac{1}{3} x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1 + id) \left( \frac{x^3 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{3 \int x^2 \log(e^{2ia+2ibx}(id + 1) + 1) dx}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 3011

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1 + id) \left( \frac{x^3 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((id+1)e^{2ia+2ibx})}{2b}) - \frac{i \int x \operatorname{PolyLog}(2, -((id+1)e^{2ia+2ibx})}{b} \right)}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 7163

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1 + id) \left( \frac{x^3 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((id+1)e^{2ia+2ibx})}{2b}) - \frac{i \left( \frac{i \int \operatorname{PolyLog}(3, -((id+1)e^{2ia+2ibx})}{2b} \right)}{2b(-d + i)} \right)}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 2720

$$\frac{1}{3}ib \left( \frac{x^4}{4} - (1 + id) \left( \frac{x^3 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((id+1)e^{2ia+2ibx})}{2b}) - \frac{i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(3, -((id+1)e^{2ia+2ibx})}{4} \right)}{2b(-d + i)} \right)}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 7143

$$\frac{1}{3} x^3 \operatorname{arctanh}(d(-\tan(a+bx)) + id + 1) + \frac{1}{3} ib \left( \frac{x^4}{4} - (1+id) \frac{x^3 \log(1 + (1+id)e^{2ia+2ibx})}{2b(-d+i)} - \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, -((id+1)e^{2ia+2ibx})}{2b}) - i \left( \frac{\operatorname{PolyLog}(4, -((id+1)e^{2ia+2ibx})}{4b^2}) \right)}{2b(-d+i)} \right)}{2b(-d+i)} \right)$$

input `Int[x^2*ArcTanh[1 + I*d - d*Tan[a + b*x]],x]`

output `(x^3*ArcTanh[1 + I*d - d*Tan[a + b*x])/3 + (I/3)*b*(x^4/4 - (1 + I*d)*((x^3*Log[1 + (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(2*b*(I - d)) - (3*(((I/2)*x^2*PolyLog[2, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x)]))/b - (I*((( -1/2*I)*x*PolyLog[3, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x)]))/b + PolyLog[4, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b))/(2*b*(I - d)))`

### Defintions of rubi rules used

rule 2615 `Int[(((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6817 `Int[ArcTanh[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.48 (sec) , antiderivative size = 2387, normalized size of antiderivative = 13.96

method	result	size
risch	Expression too large to display	2387

input `int(-x^2*arctanh(-1-I*d+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

1/12*I*b*x^4+1/6*x^3*ln(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)-1/3*x^3*ln
n(exp(I*(b*x+a)))-1/12*(-2*I*Pi+2*ln(d)+I*Pi*csgn((exp(2*I*(b*x+a))*d-I*exp
p(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn((exp(2*I*(b*x+a))*d-I*
exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(1/(exp(2*I*(b*x+a))+
1)*exp(2*I*(b*x+a))*d)^2+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I/(exp(2*I
*(b*x+a))+1)*exp(2*I*(b*x+a)))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*
(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csg
n(I*(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I))*csgn(I*(exp(2*I*(b*x+a))*d-
I*exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(1/(exp(2*I*(b*x+a)
)+1)*exp(2*I*(b*x+a))*d)^3+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2
*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I))*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*I*
(b*x+a))-I)/(exp(2*I*(b*x+a))+1))-I*Pi*csgn(I*exp(I*(b*x+a)))^2*csgn(I*exp
(2*I*(b*x+a))-I*Pi*csgn(I*exp(2*I*(b*x+a)))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a)
))+1)*exp(2*I*(b*x+a)))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)
)*d)^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*exp(2*I*(b*x+a)))*)csgn(I/(
exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)))+2*I*Pi*csgn(I*exp(I*(b*x+a)))*)csgn(I
*exp(2*I*(b*x+a)))^2+I*Pi*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)
/(exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)
))*csgn(I/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a))*d)*csgn(I*d)+I*Pi*csgn(I*(e
xp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))*csgn((exp...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 345 vs.  $2(118) = 236$ .

Time = 0.10 (sec) , antiderivative size = 345, normalized size of antiderivative = 2.02

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$$

$$= \frac{ib^4x^4 - 2b^3x^3 \log\left(-\frac{de^{(2ibx+2ia)}}{(d-i)e^{(2ibx+2ia)}-i}\right) + 6ib^2x^2 \operatorname{Li}_2\left(\frac{1}{2}\sqrt{-4id-4}e^{(ibx+ia)}\right) + 6ib^2x^2 \operatorname{Li}_2\left(-\frac{1}{2}\sqrt{-4id-4}\right)}{1}$$

input

```
integrate(-x^2*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(I*b^4*x^4 - 2*b^3*x^3*log(-d*e^(2*I*b*x + 2*I*a)/((d - I)*e^(2*I*b*x
+ 2*I*a) - I)) + 6*I*b^2*x^2*dilog(1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a))
+ 6*I*b^2*x^2*dilog(-1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) - I*a^4 + 2*a^3
*log(1/2*(2*(d - I)*e^(I*b*x + I*a) + I*sqrt(-4*I*d - 4))/(d - I)) + 2*a^3
*log(1/2*(2*(d - I)*e^(I*b*x + I*a) - I*sqrt(-4*I*d - 4))/(d - I)) - 12*b*
x*polylog(3, 1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) - 12*b*x*polylog(3, -1/
2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) - 2*(b^3*x^3 + a^3)*log(1/2*sqrt(-4*I*
d - 4)*e^(I*b*x + I*a) + 1) - 2*(b^3*x^3 + a^3)*log(-1/2*sqrt(-4*I*d - 4)*
e^(I*b*x + I*a) + 1) - 12*I*polylog(4, 1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a
)) - 12*I*polylog(4, -1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)))/b^3
```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(-d \tan(a + bx) + id + 1) dx$$

input

```
integrate(-x**2*atanh(-1-I*d+d*tan(b*x+a)),x)
```

output

```
Integral(x**2*atanh(-d*tan(a + b*x) + I*d + 1), x)
```

### Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 342 vs.  $2(118) = 236$ .

Time = 0.06 (sec) , antiderivative size = 342, normalized size of antiderivative = 2.00

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx =$$

$$\frac{12((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2) \operatorname{arctanh}(d \tan(bx+a) - id - 1)}{b^2} + \frac{-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(-4i(bx+a)^3 + 9i a^3)}{b^2}$$

input

```
integrate(-x^2*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="maxima")
```

output

```
-1/36*(12*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctanh(d*tan(
b*x + a) - I*d - 1)/b^2 + (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b
*x + a)^2*a^2 - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a - 9*I*(b*x + a)*a^
2)*arctan2(d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) + co
s(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilo
g((-I*d - 1)*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(
b*x + a)*a^2)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a
)^2 - 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polyl
og(3, (-I*d - 1)*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, (-I*d - 1)*e^(2*I*b
*x + 2*I*a)))/b^2)/b
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int -x^2 \operatorname{artanh}(d \tan(bx + a) - id - 1) dx$$

input

```
integrate(-x^2*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="giac")
```

output

```
integrate(-x^2*arctanh(d*tan(b*x + a) - I*d - 1), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int x^2 \operatorname{atanh}(1 - d \tan(a + bx) + d li) dx$$

input

```
int(x^2*atanh(d*li - d*tan(a + b*x) + 1),x)
```

output

```
int(x^2*atanh(d*li - d*tan(a + b*x) + 1), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = - \left( \int \operatorname{atanh}(\tan(bx + a) d - di - 1) x^2 dx \right)$$

input `int(-x^2*atanh(-1-I*d+d*tan(b*x+a)),x)`

output `- int(atanh(tan(a + b*x)*d - d*i - 1)*x**2,x)`



### 3.326 $\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$

Optimal result	2284
Mathematica [A] (verified)	2285
Rubi [A] (verified)	2285
Maple [C] (warning: unable to verify)	2288
Fricas [B] (verification not implemented)	2289
Sympy [F]	2289
Maxima [B] (verification not implemented)	2290
Giac [F]	2290
Mupad [F(-1)]	2291
Reduce [F]	2291

#### Optimal result

Integrand size = 19, antiderivative size = 134

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \frac{1}{6} ibx^3 + \frac{1}{2} x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx)) - \frac{1}{4} x^2 \log(1 + (1 + id)e^{2ia+2ibx}) + \frac{ix \operatorname{PolyLog}(2, -((1 + id)e^{2ia+2ibx}))}{4b} - \frac{\operatorname{PolyLog}(3, -((1 + id)e^{2ia+2ibx}))}{8b^2}$$

output

```
1/6*I*b*x^3-1/2*x^2*arctanh(-1-I*d+d*tan(b*x+a))-1/4*x^2*ln(1+(1+I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x*polylog(2,-(1+I*d)*exp(2*I*a+2*I*b*x))/b-1/8*polylog(3,-(1+I*d)*exp(2*I*a+2*I*b*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.90

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$$

$$= \frac{1}{2} x^2 \operatorname{arctanh}(1 + id - d \tan(a + bx))$$

$$- \frac{2b^2 x^2 \log\left(1 - \frac{ie^{-2i(a+bx)}}{-i+d}\right) + 2ibx \operatorname{PolyLog}\left(2, \frac{ie^{-2i(a+bx)}}{-i+d}\right) + \operatorname{PolyLog}\left(3, \frac{ie^{-2i(a+bx)}}{-i+d}\right)}{8b^2}$$

input `Integrate[x*ArcTanh[1 + I*d - d*Tan[a + b*x]], x]`

output `(x^2*ArcTanh[1 + I*d - d*Tan[a + b*x]])/2 - (2*b^2*x^2*Log[1 - I/((-I + d)*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, I/((-I + d)*E^((2*I)*(a + b*x)))] + PolyLog[3, I/((-I + d)*E^((2*I)*(a + b*x)))])/(8*b^2)`

**Rubi [A] (verified)**

Time = 0.69 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.31, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6817, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1) dx$$

$$\downarrow 6817$$

$$\frac{1}{2} ib \int \frac{x^2}{e^{2ia+2ibx}(id+1)+1} dx + \frac{1}{2} x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

$$\downarrow 2615$$

$$\frac{1}{2} ib \left( \frac{x^3}{3} - (1 + id) \int \frac{e^{2ia+2ibx} x^2}{e^{2ia+2ibx}(id+1)+1} dx \right) + \frac{1}{2} x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

$$\downarrow 2620$$

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1 + id) \left( \frac{x^2 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{\int x \log(e^{2ia+2ibx}(id + 1) + 1) dx}{b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 3011

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1 + id) \left( \frac{x^2 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -(id+1)e^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -(id+1)e^{2ia+2ibx})}{2b}}{b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 2720

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1 + id) \left( \frac{x^2 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -(id+1)e^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, -(id+1))}{4b^2}}{b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1)$$

↓ 7143

$$\frac{1}{2}ib \left( \frac{x^3}{3} - (1 + id) \left( \frac{x^2 \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d + i)} - \frac{\frac{ix \operatorname{PolyLog}(2, -(id+1)e^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, -(id+1)e^{2ia+2ibx})}{4b^2}}{b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\tan(a + bx)) + id + 1) +$$

input `Int[x*ArcTanh[1 + I*d - d*Tan[a + b*x]],x]`

output `(x^2*ArcTanh[1 + I*d - d*Tan[a + b*x])/2 + (I/2)*b*(x^3/3 - (1 + I*d)*((x^2*Log[1 + (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(2*b*(I - d)) - (((I/2)*x*PolyLog[2, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x))])/b - PolyLog[3, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2))/(b*(I - d))))`

## Definitions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 6817

```
Int[ArcTanh[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Tan[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.81 (sec) , antiderivative size = 2289, normalized size of antiderivative = 17.08

method	result	size
risch	Expression too large to display	2289

input `int(-x*arctanh(-1-I*d+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/6*I*b*x^3-1/2/b*d*a/(I-d)*ln(1+I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))*x-1/2/
b*d*a/(I-d)*ln(1-I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))*x+1/2/b*d/(I-d)*ln(1-I
*exp(2*I*(b*x+a))*(I-d))*a*x+1/4/b^2*a^2*d/(I-d)*ln(I*exp(2*I*(b*x+a))-exp
(2*I*(b*x+a))*d+I)+1/4/b^2*d/(I-d)*ln(1-I*exp(2*I*(b*x+a))*(I-d))*a^2-1/2/
b^2*d*a^2/(I-d)*ln(1+I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))-1/2/b^2*d*a^2/(I-d
)*ln(1-I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))-1/4*I/b^2*a^2/(I-d)*ln(I*exp(2*I
*(b*x+a))-exp(2*I*(b*x+a))*d+I)-1/4*I/b^2/(I-d)*ln(1-I*exp(2*I*(b*x+a))*(I
-d))*a^2+1/2*I/b^2*a^2/(I-d)*ln(1+I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))+1/2*I
/b^2*a^2/(I-d)*ln(1-I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))-1/2*I/b/(I-d)*ln(1-
I*exp(2*I*(b*x+a))*(I-d))*a*x+1/2*I/b^2*d*a/(I-d)*dilog(1+I*exp(I*(b*x+a))
*(-I*(I-d))^(1/2))+1/2*I/b^2*d*a/(I-d)*dilog(1-I*exp(I*(b*x+a))*(-I*(I-d))
^(1/2))+1/2*I/b*a/(I-d)*ln(1+I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))*x+1/2*I/b*
a/(I-d)*ln(1-I*exp(I*(b*x+a))*(-I*(I-d))^(1/2))*x-1/4*I/b*d/(I-d)*polylog(
2,I*exp(2*I*(b*x+a))*(I-d))*x-1/4*I/b^2*d/(I-d)*polylog(2,I*exp(2*I*(b*x+a
))*(I-d))*a-1/2*x^2*ln(exp(I*(b*x+a)))-1/8*(-2*I*Pi+2*ln(d)+I*Pi*csgn((exp
(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn((e
xp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(
1/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a))*d)^2+I*Pi*csgn(I/(exp(2*I*(b*x+a))
+1))*csgn(I/(exp(2*I*(b*x+a))+1)*exp(2*I*(b*x+a)))^2-I*Pi*csgn(I/(exp(2*I*
(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))-I)/(exp(2*I*...
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs.  $2(93) = 186$ .

Time = 0.11 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.19

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$$

$$= \frac{2i b^3 x^3 - 3 b^2 x^2 \log\left(-\frac{d e^{(2i b x + 2i a)}}{(d-i)e^{(2i b x + 2i a)} - i}\right) + 2i a^3 + 6i b x \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4i d - 4} e^{(i b x + i a)}\right) + 6i b x \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{-4i d - 4} e^{(i b x + i a)}\right)}{b^2}$$

input `integrate(-x*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*I*b^3*x^3 - 3*b^2*x^2*log(-d*e^(2*I*b*x + 2*I*a)/((d - I)*e^(2*I*b*x + 2*I*a) - I)) + 2*I*a^3 + 6*I*b*x*dilog(1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) + 6*I*b*x*dilog(-1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) - 3*a^2*log(1/2*(2*(d - I)*e^(I*b*x + I*a) + I*sqrt(-4*I*d - 4))/(d - I)) - 3*a^2*log(1/2*(2*(d - I)*e^(I*b*x + I*a) - I*sqrt(-4*I*d - 4))/(d - I)) - 3*(b^2*x^2 - a^2)*log(1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a) + 1) - 3*(b^2*x^2 - a^2)*log(-1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a) + 1) - 6*polylog(3, 1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)) - 6*polylog(3, -1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int x \operatorname{atanh}(-d \tan(a + bx) + id + 1) dx$$

input `integrate(-x*atanh(-1-I*d+d*tan(b*x+a)),x)`

output `Integral(x*atanh(-d*tan(a + b*x) + I*d + 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 247 vs.  $2(93) = 186$ .

Time = 0.05 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.84

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \frac{12((bx+a)^2 - 2(bx+a)a) \operatorname{artanh}(d \tan(bx+a) - id - 1)}{b} + \frac{-4i(bx+a)^3 + 12i(bx+a)^2 a - 6i b x \operatorname{Li}_2((-id-1)e^{2i bx + 2i a}) - 6(-i(bx+a)^2 + 2i a)}{b}$$

input `integrate(-x*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="maxima")`

output `-1/24*(12*((b*x + a)^2 - 2*(b*x + a)*a)*arctanh(d*tan(b*x + a) - I*d - 1)/b + (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog((-I*d - 1)*e^(2*I*b*x + 2*I*a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^2 - 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1) + 3*polylog(3, (-I*d - 1)*e^(2*I*b*x + 2*I*a)))/b/b`

**Giac [F]**

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int -x \operatorname{artanh}(d \tan(bx + a) - id - 1) dx$$

input `integrate(-x*arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctanh(d*tan(b*x + a) - I*d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int x \operatorname{atanh}(1 - d \tan(a + bx) + d1i) dx$$

input `int(x*atanh(d*1i - d*tan(a + b*x) + 1),x)`

output `int(x*atanh(d*1i - d*tan(a + b*x) + 1), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = - \left( \int \operatorname{atanh}(\tan(bx + a) d - di - 1) x dx \right)$$

input `int(-x*atanh(-1-I*d+d*tan(b*x+a)),x)`

output `- int(atanh(tan(a + b*x)*d - d*i - 1)*x,x)`



### 3.327 $\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$

Optimal result	2292
Mathematica [B] (warning: unable to verify)	2292
Rubi [A] (verified)	2293
Maple [B] (verified)	2295
Fricas [B] (verification not implemented)	2296
Sympy [F]	2297
Maxima [B] (verification not implemented)	2297
Giac [F]	2298
Mupad [F(-1)]	2298
Reduce [F]	2298

#### Optimal result

Integrand size = 17, antiderivative size = 94

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \frac{1}{2}ibx^2 + x \operatorname{arctanh}(1 + id - d \tan(a + bx)) - \frac{1}{2}x \log(1 + (1 + id)e^{2ia+2ibx}) + \frac{i \operatorname{PolyLog}(2, -((1 + id)e^{2ia+2ibx}))}{4b}$$

output `1/2*I*b*x^2-x*arctanh(-1-I*d+d*tan(b*x+a))-1/2*x*ln(1+(1+I*d)*exp(2*I*a+2*I*b*x))+1/4*I*polylog(2,-(1+I*d)*exp(2*I*a+2*I*b*x))/b`

#### Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 723 vs. 2(94) = 188.

Time = 4.73 (sec) , antiderivative size = 723, normalized size of antiderivative = 7.69

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = x \operatorname{arctanh}(1 + id - d \tan(a + bx)) - \frac{x \left( -2ibx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) + \log \left( \frac{\sec(bx)(\cos(a) - i \sin(a))((-2i+d) \cos(a+bx) + id \sin(a+bx))}{2(-i+d)} \right) \right) \log \left( \frac{i \log(1 - i \tan(bx)) \sec(bx)(d \cos(a) + (2+id) \sin(a))}{(-2i+d) \cos(a+bx) + id \sin(a+bx)} + \frac{\log(1 + i \tan(bx))}{(-2i+d)} \right)}{((-2i + d) \cos(a + bx) + id \sin(a + bx))}$$

input `Integrate[ArcTanh[1 + I*d - d*Tan[a + b*x]],x]`

output

```
x*ArcTanh[1 + I*d - d*Tan[a + b*x]] - (x*((-2*I)*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] + Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((-2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]))/(2*(-I + d))]*Log[1 - I*Tan[b*x]] - Log[(Sec[b*x]*((2 + I*d)*Cos[a + b*x] - d*Sin[a + b*x]))/(2*Cos[a] - (2*I)*Sin[a])]*Log[1 + I*Tan[b*x]] + PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] + PolyLog[2, (Sec[b*x]*(d*Cos[a] + (2 + I*d)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*(-I + d))] - PolyLog[2, ((Cos[a] + I*Sin[a])*(d*Cos[a] + (2 + I*d)*Sin[a])*(-I + Tan[b*x]))/2])*Sec[a + b*x]*(Cos[b*x] + I*Sin[b*x])*(I*Cos[b*x] + Sin[b*x]))/(((2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x])*((I*Log[1 - I*Tan[b*x]]*Sec[b*x]*(d*Cos[a] + (2 + I*d)*Sin[a]))/((-2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]) + (Log[1 + I*Tan[b*x]]*Sec[b*x]*((-I)*d*Cos[a] + (-2*I + d)*Sin[a]))/((-2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]) - (Log[(Sec[b*x]*((2 + I*d)*Cos[a + b*x] - d*Sin[a + b*x]))/(2*Cos[a] - (2*I)*Sin[a])]*Sec[b*x]^2)/(-I + Tan[b*x]) + (Log[1 - ((Cos[a] + I*Sin[a])*(d*Cos[a] + (2 + I*d)*Sin[a])*(-I + Tan[b*x]))/2]*Sec[b*x]^2)/(-I + Tan[b*x]) - Log[1 - (Sec[b*x]*(d*Cos[a] + (2 + I*d)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*(-I + d))]*(-I + Tan[b*x]) + (Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((-2*I + d)*Cos[a + b*x] + I*d*Sin[a + b*x]))/(2*(-I + d))]*Sec[b*x]^2)/(I + Tan[b*x]) + (2*I)*b*x*(I + Tan[b*x]))*(-I + Tan[a + b*x]))
```

## Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.36, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {6809, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctanh(d(-\tan(a + bx)) + id + 1) dx$$

$$\downarrow 6809$$

$$ib \int \frac{x}{e^{2ia+2ibx}(id + 1) + 1} dx + x \arctanh(d(-\tan(a + bx)) + id + 1)$$

$$\downarrow 2615$$

$$\begin{aligned}
& ib \left( \frac{x^2}{2} - (1 + id) \int \frac{e^{2ia+2ibx} x}{e^{2ia+2ibx}(id+1)+1} dx \right) + x \operatorname{arctanh}(d(-\tan(a+bx)) + id + 1) \\
& \quad \downarrow \text{2620} \\
& ib \left( \frac{x^2}{2} - (1 + id) \left( \frac{x \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d+i)} - \frac{\int \log(e^{2ia+2ibx}(id+1)+1) dx}{2b(-d+i)} \right) \right) + \\
& \quad \quad \quad x \operatorname{arctanh}(d(-\tan(a+bx)) + id + 1) \\
& \quad \quad \quad \downarrow \text{2715} \\
& ib \left( \frac{x^2}{2} - (1 + id) \left( \frac{i \int e^{-2ia-2ibx} \log(e^{2ia+2ibx}(id+1)+1) de^{2ia+2ibx}}{4b^2(-d+i)} + \frac{x \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d+i)} \right) \right) + \\
& \quad \quad \quad x \operatorname{arctanh}(d(-\tan(a+bx)) + id + 1) \\
& \quad \quad \quad \downarrow \text{2838} \\
& ib \left( \frac{x^2}{2} - (1 + id) \left( \frac{x \operatorname{arctanh}(d(-\tan(a+bx)) + id + 1) + x \log(1 + (1 + id)e^{2ia+2ibx})}{2b(-d+i)} - \frac{i \operatorname{PolyLog}(2, -(id+1)e^{2ia+2ibx})}{4b^2(-d+i)} \right) \right)
\end{aligned}$$

input `Int[ArcTanh[1 + I*d - d*Tan[a + b*x]], x]`

output `x*ArcTanh[1 + I*d - d*Tan[a + b*x]] + I*b*(x^2/2 - (1 + I*d)*((x*Log[1 + (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/(2*b*(I - d)) - ((I/4)*PolyLog[2, -((1 + I*d)*E^((2*I)*a + (2*I)*b*x))])/(b^2*(I - d))))`

### Defintions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2620 Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 6809 Int[ArcTanh[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]], x_Symbol] := Simp[x*Arc
Tanh[c + d*Tan[a + b*x]], x] + Simp[I*b Int[x/(c + I*d + c*E^(2*I*a + 2*I
*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c + I*d)^2, 1]
```

### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 331 vs. 2(77) = 154.

Time = 3.73 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.53

method	result
derivativedivides	$-\frac{i \operatorname{arctanh}(-1-id+d \tan (bx+a)) d \ln (-id+d \tan (bx+a))}{2} + \frac{i \operatorname{arctanh}(-1-id+d \tan (bx+a)) d \ln (-id-d \tan (bx+a))}{2} - \frac{d^2 \left( \frac{i \left( -\frac{c}{d} \right)}{d^2} \right)}{d^2}$
default	$-\frac{i \operatorname{arctanh}(-1-id+d \tan (bx+a)) d \ln (-id+d \tan (bx+a))}{2} + \frac{i \operatorname{arctanh}(-1-id+d \tan (bx+a)) d \ln (-id-d \tan (bx+a))}{2} - \frac{d^2 \left( \frac{i \left( -\frac{c}{d} \right)}{d^2} \right)}{d^2}$
risch	Expression too large to display

```
input int(-arctanh(-1-I*d+d*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

```
output -1/b/d*(-1/2*I*arctanh(-1-I*d+d*tan(b*x+a))*d*ln(-I*d+d*tan(b*x+a))+1/2*I*
arctanh(-1-I*d+d*tan(b*x+a))*d*ln(-I*d-d*tan(b*x+a))-1/2*d^2*(-I/d*(-1/2*d
ilog(-1/2*I*(I*d-d*tan(b*x+a))/d)-1/2*ln(-I*d-d*tan(b*x+a))*ln(-1/2*I*(I*d
-d*tan(b*x+a))/d)+1/2*dilog(I*(-I*d-d*tan(b*x+a)-I*(2*I-2*d))/(2*I-2*d))+1
/2*ln(-I*d-d*tan(b*x+a))*ln(I*(-I*d-d*tan(b*x+a)-I*(2*I-2*d))/(2*I-2*d)))+
I/d*(1/2*(ln(-I*d+d*tan(b*x+a))-ln(-1/2*I*d+1/2*d*tan(b*x+a)))*ln(1+1/2*I*
d-1/2*d*tan(b*x+a))-1/2*dilog(-1/2*I*d+1/2*d*tan(b*x+a))-1/4*ln(-I*d+d*tan
(b*x+a))^2))
```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 219 vs.  $2(66) = 132$ .

Time = 0.10 (sec) , antiderivative size = 219, normalized size of antiderivative = 2.33

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx$$

$$= \frac{i b^2 x^2 - bx \log\left(-\frac{de^{(2i bx + 2i a)}}{(d-i)e^{(2i bx + 2i a)} - i}\right) - i a^2 - (bx + a) \log\left(\frac{1}{2} \sqrt{-4i d - 4} e^{(i bx + i a)} + 1\right) - (bx + a) \log\left(-\frac{1}{2}\right)}{1}$$

```
input integrate(-arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="fricas")
```

```
output 1/2*(I*b^2*x^2 - b*x*log(-d*e^(2*I*b*x + 2*I*a)/((d - I)*e^(2*I*b*x + 2*I*
a) - I)) - I*a^2 - (b*x + a)*log(1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a) + 1)
- (b*x + a)*log(-1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a) + 1) + a*log(1/2*(2
*(d - I)*e^(I*b*x + I*a) + I*sqrt(-4*I*d - 4))/(d - I)) + a*log(1/2*(2*(d
- I)*e^(I*b*x + I*a) - I*sqrt(-4*I*d - 4))/(d - I)) + I*dilog(1/2*sqrt(-4*
I*d - 4)*e^(I*b*x + I*a)) + I*dilog(-1/2*sqrt(-4*I*d - 4)*e^(I*b*x + I*a)
)/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int \operatorname{atanh}(-d \tan(a + bx) + id + 1) dx$$

input `integrate(-atanh(-1-I*d+d*tan(b*x+a)), x)`

output `Integral(atanh(-d*tan(a + b*x) + I*d + 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 262 vs.  $2(66) = 132$ .

Time = 0.12 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.79

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx =$$

$$\frac{4(bx + a)d \left( \frac{\log(d \tan(bx+a) - id - 2)}{d} - \frac{\log(\tan(bx+a) - i)}{d} \right) - d \left( \frac{2i \left( \log(d \tan(bx+a) - id - 2) \log\left(-\frac{id \tan(bx+a) + d - 2i}{2(d-i)} + 1\right) + \operatorname{Li}_2\left(\frac{d \tan(bx+a) - id - 2}{d}\right) \right)}{d} \right)}{1}$$

input `integrate(-arctanh(-1-I*d+d*tan(b*x+a)), x, algorithm="maxima")`

output `-1/8*(4*(b*x + a)*d*(log(d*tan(b*x + a) - I*d - 2)/d - log(tan(b*x + a) - I)/d) - d*(2*I*(log(d*tan(b*x + a) - I*d - 2)*log(-1/2*(I*d*tan(b*x + a) + d - 2*I)/(d - I) + 1) + dilog(1/2*(I*d*tan(b*x + a) + d - 2*I)/(d - I)))/d - (2*I*log(d*tan(b*x + a) - I*d - 2)*log(tan(b*x + a) - I) - I*log(tan(b*x + a) - I)^2)/d + 2*I*(log(-1/2*d*tan(b*x + a) + 1/2*I*d + 1)*log(tan(b*x + a) - I) + dilog(1/2*d*tan(b*x + a) - 1/2*I*d))/d - 2*I*(log(tan(b*x + a) - I)*log(-1/2*I*tan(b*x + a) + 1/2) + dilog(1/2*I*tan(b*x + a) + 1/2))/d + 8*(b*x + a)*arctanh(d*tan(b*x + a) - I*d - 1))/b`

**Giac [F]**

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int -\operatorname{artanh}(d \tan(bx + a) - id - 1) dx$$

input `integrate(-arctanh(-1-I*d+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(-arctanh(d*tan(b*x + a) - I*d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = \int \operatorname{atanh}(1 - d \tan(a + bx) + d li) dx$$

input `int(atanh(d*li - d*tan(a + b*x) + 1),x)`

output `int(atanh(d*li - d*tan(a + b*x) + 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 + id - d \tan(a + bx)) dx = -\left(\int \operatorname{atanh}(\tan(bx + a)d - di - 1) dx\right)$$

input `int(-atanh(-1-I*d+d*tan(b*x+a)),x)`

output `- int(atanh(tan(a + b*x)*d - d*i - 1),x)`

### 3.328 $\int \frac{\operatorname{arctanh}(1+id-d \tan(a+bx))}{x} dx$

Optimal result	2299
Mathematica [N/A]	2299
Rubi [N/A]	2300
Maple [N/A]	2300
Fricas [N/A]	2301
Sympy [N/A]	2301
Maxima [N/A]	2301
Giac [N/A]	2302
Mupad [N/A]	2302
Reduce [N/A]	2303

#### Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x}, x\right)$$

output

```
Defer(Int)(-arctanh(-1-I*d+d*tan(b*x+a))/x,x)
```

#### Mathematica [N/A]

Not integrable

Time = 0.79 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx$$

input

```
Integrate[ArcTanh[1 + I*d - d*Tan[a + b*x]]/x,x]
```

output

```
Integrate[ArcTanh[1 + I*d - d*Tan[a + b*x]]/x, x]
```



**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d(-\tan(a+bx)) + id + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d(-\tan(a+bx)) + id + 1)}{x} dx$$

input `Int[ArcTanh[1 + I*d - d*Tan[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int -\frac{\operatorname{arctanh}(-1 - id + d \tan(bx + a))}{x} dx$$

input `int(-arctanh(-1-I*d+d*tan(b*x+a))/x,x)`

output `int(-arctanh(-1-I*d+d*tan(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \tan(bx + a) - id - 1)}{x} dx$$

input `integrate(-arctanh(-1-I*d+d*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(-1/2*log(-d*e^(2*I*b*x + 2*I*a)/((d - I)*e^(2*I*b*x + 2*I*a) - I))/x, x)`

**Sympy [N/A]**

Not integrable

Time = 0.87 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(-d \tan(a + bx) + id + 1)}{x} dx$$

input `integrate(-atanh(-1-I*d+d*tan(b*x+a))/x,x)`

output `Integral(atanh(-d*tan(a + b*x) + I*d + 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 3.96 (sec) , antiderivative size = 143, normalized size of antiderivative = 6.81

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \tan(bx + a) - id - 1)}{x} dx$$

input `integrate(-arctanh(-1-I*d+d*tan(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x + 1/4*(-I*pi - 4*I*a - 2*log(-d))*log(x) + 1/2*I*integrate(arctan2(
d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) + cos(2*b*x + 2
*a) + 1)/x, x) + 1/4*integrate(log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1
)*sin(2*b*x + 2*a)^2 - 2*d*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) + 1)/x, x
)
```

**Giac [N/A]**

Not integrable

Time = 0.48 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \tan(bx + a) - id - 1)}{x} dx$$

input

```
integrate(-arctanh(-1-I*d+d*tan(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(-arctanh(d*tan(b*x + a) - I*d - 1)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.58 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(1 - d \tan(a + bx) + d li)}{x} dx$$

input

```
int(atanh(d*li - d*tan(a + b*x) + 1)/x,x)
```

output

```
int(atanh(d*li - d*tan(a + b*x) + 1)/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(1 + id - d \tan(a + bx))}{x} dx = - \left( \int \frac{\operatorname{atanh}(\tan(bx + a) d - di - 1)}{x} dx \right)$$

input

```
int(-atanh(-1-I*d+d*tan(b*x+a))/x,x)
```

output

```
- int(atanh(tan(a + b*x)*d - d*i - 1)/x,x)
```

### 3.329 $\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx$

Optimal result	2304
Mathematica [B] (verified)	2305
Rubi [A] (verified)	2306
Maple [C] (warning: unable to verify)	2310
Fricas [B] (verification not implemented)	2311
Sympy [F]	2312
Maxima [F]	2313
Giac [F]	2313
Mupad [F(-1)]	2314
Reduce [F]	2314

#### Optimal result

Integrand size = 15, antiderivative size = 302

$$\begin{aligned}
 \int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx = & \frac{i(e + fx)^4 \arctan(e^{2i(a+bx)})}{4f} \\
 & + \frac{(e + fx)^4 \operatorname{arctanh}(\cot(a + bx))}{4f} \\
 & - \frac{i(e + fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} \\
 & + \frac{i(e + fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} \\
 & + \frac{3f(e + fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{8b^2} \\
 & - \frac{3f(e + fx)^2 \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{8b^2} \\
 & + \frac{3if^2(e + fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{8b^3} \\
 & - \frac{3if^2(e + fx) \operatorname{PolyLog}(4, ie^{2i(a+bx)})}{8b^3} \\
 & - \frac{3f^3 \operatorname{PolyLog}(5, -ie^{2i(a+bx)})}{16b^4} \\
 & + \frac{3f^3 \operatorname{PolyLog}(5, ie^{2i(a+bx)})}{16b^4}
 \end{aligned}$$

output

```

1/4*I*(f*x+e)^4*arctan(exp(2*I*(b*x+a)))/f+1/4*(f*x+e)^4*arctanh(cot(b*x+a
))/f-1/4*I*(f*x+e)^3*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)^3*poly
log(2,I*exp(2*I*(b*x+a)))/b+3/8*f*(f*x+e)^2*polylog(3,-I*exp(2*I*(b*x+a)))
/b^2-3/8*f*(f*x+e)^2*polylog(3,I*exp(2*I*(b*x+a)))/b^2+3/8*I*f^2*(f*x+e)*p
olylog(4,-I*exp(2*I*(b*x+a)))/b^3-3/8*I*f^2*(f*x+e)*polylog(4,I*exp(2*I*(b
*x+a)))/b^3-3/16*f^3*polylog(5,-I*exp(2*I*(b*x+a)))/b^4+3/16*f^3*polylog(5
,I*exp(2*I*(b*x+a)))/b^4

```

### Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 654 vs.  $2(302) = 604$ .

Time = 0.19 (sec) , antiderivative size = 654, normalized size of antiderivative = 2.17

$$\begin{aligned}
& \int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx \\
&= \frac{1}{4} x (4e^3 + 6e^2 fx + 4ef^2 x^2 + f^3 x^3) \operatorname{arctanh}(\cot(a + bx)) \\
&+ \frac{-8b^4 e^3 x \log(1 - ie^{2i(a+bx)}) - 12b^4 e^2 f x^2 \log(1 - ie^{2i(a+bx)}) - 8b^4 e f^2 x^3 \log(1 - ie^{2i(a+bx)}) - 2b^4 f^3 x^4}{4}
\end{aligned}$$

input

```
Integrate[(e + f*x)^3*ArcTanh[Cot[a + b*x]],x]
```

output

```
(x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcTanh[Cot[a + b*x]])/4 +
(-8*b^4*e^3*x*Log[1 - I*E^((2*I)*(a + b*x))] - 12*b^4*e^2*f*x^2*Log[1 - I*
E^((2*I)*(a + b*x))] - 8*b^4*e*f^2*x^3*Log[1 - I*E^((2*I)*(a + b*x))] - 2*
b^4*f^3*x^4*Log[1 - I*E^((2*I)*(a + b*x))] + 8*b^4*e^3*x*Log[1 + I*E^((2*I)
*(a + b*x))] + 12*b^4*e^2*f*x^2*Log[1 + I*E^((2*I)*(a + b*x))] + 8*b^4*e*
f^2*x^3*Log[1 + I*E^((2*I)*(a + b*x))] + 2*b^4*f^3*x^4*Log[1 + I*E^((2*I)*
(a + b*x))] - (4*I)*b^3*(e + f*x)^3*PolyLog[2, (-I)*E^((2*I)*(a + b*x))] +
(4*I)*b^3*(e + f*x)^3*PolyLog[2, I*E^((2*I)*(a + b*x))] + 6*b^2*e^2*f*Pol
yLog[3, (-I)*E^((2*I)*(a + b*x))] + 12*b^2*e*f^2*x*PolyLog[3, (-I)*E^((2*I)
*(a + b*x))] + 6*b^2*f^3*x^2*PolyLog[3, (-I)*E^((2*I)*(a + b*x))] - 6*b^2
*e^2*f*PolyLog[3, I*E^((2*I)*(a + b*x))] - 12*b^2*e*f^2*x*PolyLog[3, I*E^((
2*I)*(a + b*x))] - 6*b^2*f^3*x^2*PolyLog[3, I*E^((2*I)*(a + b*x))] + (6*I)
*b*e*f^2*PolyLog[4, (-I)*E^((2*I)*(a + b*x))] + (6*I)*b*f^3*x*PolyLog[4,
(-I)*E^((2*I)*(a + b*x))] - (6*I)*b*e*f^2*PolyLog[4, I*E^((2*I)*(a + b*x))
] - (6*I)*b*f^3*x*PolyLog[4, I*E^((2*I)*(a + b*x))] - 3*f^3*PolyLog[5, (-I)
)*E^((2*I)*(a + b*x))] + 3*f^3*PolyLog[5, I*E^((2*I)*(a + b*x)))]/(16*b^4)
```

### Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$ , Rules used = {6807, 3042, 4669, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) \, dx \\
 & \quad \downarrow \text{6807} \\
 & \frac{(e + fx)^4 \operatorname{arctanh}(\cot(a + bx))}{4f} - \frac{b \int (e + fx)^4 \sec(2a + 2bx) \, dx}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^4 \operatorname{arctanh}(\cot(a + bx))}{4f} - \frac{b \int (e + fx)^4 \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx}{4f} \\
 & \quad \downarrow \text{4669}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(e+fx)^4 \operatorname{arctanh}(\cot(a+bx))}{4f} - \frac{b \left( -\frac{2f \int (e+fx)^3 \log(1-ie^{2i(a+bx)}) dx}{b} + \frac{2f \int (e+fx)^3 \log(1+ie^{2i(a+bx)}) dx}{b} - \frac{i(e+fx)^4 \operatorname{arctan}(e^{2i(a+bx)})}{b} \right)}{4f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\cot(a+bx))}{4f} - \frac{b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)}) dx}{2b} \right)}{b} \right)}{4f} \\
 & \quad \downarrow \text{7163} \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\cot(a+bx))}{4f} - \frac{b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{if \int (e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) dx}{b} - \frac{i(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{2b} \right)}{b} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} - \frac{3if \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)}) dx}{2b} \right)}{b} \right)}{4f} \\
 & \quad \downarrow \text{7163} \\
 & \frac{(e+fx)^4 \operatorname{arctanh}(\cot(a+bx))}{4f} - \frac{b \left( \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{if \int \operatorname{PolyLog}(4, -ie^{2i(a+bx)}) dx}{2b} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} - \frac{i(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{2b} \right)}{b} \\
 & \quad \downarrow \text{2720}
 \end{aligned}$$



$$\frac{(e + fx)^4 \operatorname{arctanh}(\cot(a + bx))}{4f} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}(4, -ie^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b}$$

7143

$$\frac{(e + fx)^4 \operatorname{arctanh}(\cot(a + bx))}{4f} - \frac{2f \left( \frac{i(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{3if \left( \frac{f \operatorname{PolyLog}(5, -ie^{2i(a+bx)})}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} + \frac{i(e+fx)^4 \arctan(e^{2i(a+bx)})}{b}$$

input `Int[(e + f*x)^3*ArcTanh[Cot[a + b*x]],x]`

output

```
((e + f*x)^4*ArcTanh[Cot[a + b*x]]/(4*f) - (b*((( -I)*(e + f*x)^4*ArcTan[E
^((2*I)*(a + b*x))])/b + (2*f*(((I/2)*(e + f*x)^3*PolyLog[2, (-I)*E^((2*I)
*(a + b*x))])/b - (((3*I)/2)*f*((( -1/2*I)*(e + f*x)^2*PolyLog[3, (-I)*E^((
2*I)*(a + b*x))])/b + (I*f*((( -1/2*I)*(e + f*x)*PolyLog[4, (-I)*E^((2*I)*
(a + b*x))])/b + (f*PolyLog[5, (-I)*E^((2*I)*(a + b*x))]/(4*b^2))/b))/b))
/b - (2*f*(((I/2)*(e + f*x)^3*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (((3*
I)/2)*f*((( -1/2*I)*(e + f*x)^2*PolyLog[3, I*E^((2*I)*(a + b*x))])/b + (I*f
*((( -1/2*I)*(e + f*x)*PolyLog[4, I*E^((2*I)*(a + b*x))])/b + (f*PolyLog[5,
I*E^((2*I)*(a + b*x))]/(4*b^2))/b))/b))/b)/(4*f)
```

### Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6807 `Int[ArcTanh[Cot[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTanh[Cot[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 14.02 (sec) , antiderivative size = 3640, normalized size of antiderivative = 12.05

method	result	size
risch	Expression too large to display	3640

input `int((f*x+e)^3*arctanh(cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

-3/16*f^3*polylog(5,-I*exp(2*I*(b*x+a)))/b^4+3/16*f^3*polylog(5,I*exp(2*I*
(b*x+a)))/b^4+3/2*f/b*e^2*ln(I*exp(2*I*(b*x+a))+1)*x*a-3/2*f^2/b^2*e*ln(I*
exp(2*I*(b*x+a))+1)*x*a^2-3/4*I*f/b^2*e^2*polylog(2,-I*exp(2*I*(b*x+a)))*a
+3/2*I*f^2/b^3*e*a^2*dilog(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))+3/2*f^2
/b^3*a^3*e*ln(1-exp(I*(b*x+a)))*(-I)^(3/4))-1/2*f^3/b^3*a^3*ln(1+exp(I*(b*x
+a)))*(-I)^(3/4))*x-1/2*f^3/b^3*a^3*ln(1-exp(I*(b*x+a)))*(-I)^(3/4))*x+3/4*I
*f^2/b*e*polylog(2,I*exp(2*I*(b*x+a)))*x^2-1/2*I/b*e^3*dilog(1+exp(I*(b*x+
a)))*(-I)^(3/4))-1/2*I/b*e^3*dilog(1-exp(I*(b*x+a)))*(-I)^(3/4))+1/2*I/b*e^3
*dilog(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))+1/2*I/b*e^3*dilog(((I)^(1/
2)+exp(I*(b*x+a)))/(I)^(1/2))+1/2*I*f^3/b^4*a^3*dilog(1-exp(I*(b*x+a)))*(-
I)^(3/4))-1/2*I*f^3/b^4*a^3*dilog(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))-
1/2*I*f^3/b^4*a^3*dilog(((I)^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))+1/2*I*f^3/
b^4*a^3*dilog(1+exp(I*(b*x+a)))*(-I)^(3/4))+3/2*I*f^2/b^3*e*a^2*dilog(((I)
^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))-3/2*I*f/b^2*e^2*a*dilog(((I)^(1/2)-exp
(I*(b*x+a)))/(I)^(1/2))-3/2*I*f/b^2*e^2*a*dilog(((I)^(1/2)+exp(I*(b*x+a)
)))/(I)^(1/2))-3/2*I*f^2/b^3*a^2*e*dilog(1+exp(I*(b*x+a)))*(-I)^(3/4))-3/2*
I*f^2/b^3*a^2*e*dilog(1-exp(I*(b*x+a)))*(-I)^(3/4))+3/2*I*f/b^2*a*e^2*dilog
(1+exp(I*(b*x+a)))*(-I)^(3/4))+3/2*I*f/b^2*a*e^2*dilog(1-exp(I*(b*x+a)))*(-1
)^(3/4))+3/2*f/b*e^2*a*ln(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))*x+3/2*f/
b*e^2*a*ln(((I)^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))*x-1/2*f^2*ln(exp(2*I...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1567 vs.  $2(236) = 472$ .

Time = 0.19 (sec) , antiderivative size = 1567, normalized size of antiderivative = 5.19

$$\int (e + fx)^3 \operatorname{arctanh}(\cot(ax + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^3*arctanh(cot(b*x+a)),x, algorithm="fricas")
```



**Maxima [F]**

$$\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e)^3 \operatorname{arctanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)^3*arctanh(cot(b*x+a)),x, algorithm="maxima")`

output `1/16*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/16*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(1/2*((b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f^3*x^4 + 4*b*e*f^2*x^3 + 6*b*e^2*f*x^2 + 4*b*e^3*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

**Giac [F]**

$$\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e)^3 \operatorname{arctanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)^3*arctanh(cot(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)^3*arctanh(cot(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(a + bx)) (e + fx)^3 dx$$

input `int(atanh(cot(a + b*x))*(e + f*x)^3,x)`

output `int(atanh(cot(a + b*x))*(e + f*x)^3, x)`

**Reduce [F]**

$$\begin{aligned} \int (e + fx)^3 \operatorname{arctanh}(\cot(a + bx)) dx &= \left( \int \operatorname{atanh}(\cot(bx + a)) dx \right) e^3 \\ &+ \left( \int \operatorname{atanh}(\cot(bx + a)) x^3 dx \right) f^3 \\ &+ 3 \left( \int \operatorname{atanh}(\cot(bx + a)) x^2 dx \right) e f^2 \\ &+ 3 \left( \int \operatorname{atanh}(\cot(bx + a)) x dx \right) e^2 f \end{aligned}$$

input `int((f*x+e)^3*atanh(cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)),x)*e**3 + int(atanh(cot(a + b*x))*x**3,x)*f**3 + 3  
*int(atanh(cot(a + b*x))*x**2,x)*e*f**2 + 3*int(atanh(cot(a + b*x))*x,x)*e  
**2*f`

### 3.330 $\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx$

Optimal result	2315
Mathematica [A] (verified)	2316
Rubi [A] (verified)	2317
Maple [C] (warning: unable to verify)	2320
Fricas [B] (verification not implemented)	2321
Sympy [F]	2322
Maxima [F]	2323
Giac [F]	2323
Mupad [F(-1)]	2323
Reduce [F]	2324

#### Optimal result

Integrand size = 15, antiderivative size = 234

$$\begin{aligned}
 \int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = & \frac{i(e + fx)^3 \arctan(e^{2i(a+bx)})}{3f} \\
 & + \frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} \\
 & - \frac{i(e + fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} \\
 & + \frac{i(e + fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} \\
 & + \frac{f(e + fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{4b^2} \\
 & - \frac{f(e + fx) \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{4b^2} \\
 & + \frac{if^2 \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{8b^3} \\
 & - \frac{if^2 \operatorname{PolyLog}(4, ie^{2i(a+bx)})}{8b^3}
 \end{aligned}$$



output

```
1/3*I*(f*x+e)^3*arctan(exp(2*I*(b*x+a)))/f+1/3*(f*x+e)^3*arctanh(cot(b*x+a
))/f-1/4*I*(f*x+e)^2*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)^2*poly
log(2,I*exp(2*I*(b*x+a)))/b+1/4*f*(f*x+e)*polylog(3,-I*exp(2*I*(b*x+a)))/b
^2-1/4*f*(f*x+e)*polylog(3,I*exp(2*I*(b*x+a)))/b^2+1/8*I*f^2*polylog(4,-I*
exp(2*I*(b*x+a)))/b^3-1/8*I*f^2*polylog(4,I*exp(2*I*(b*x+a)))/b^3
```

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.75

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \frac{1}{3}x(3e^2 + 3efx + f^2x^2) \operatorname{arctanh}(\cot(a + bx)) \\ + \frac{-12b^3e^2x \log(1 - ie^{2i(a+bx)}) - 12b^3efx^2 \log(1 - ie^{2i(a+bx)}) - 4b^3f^2x^3 \log(1 - ie^{2i(a+bx)}) + 12b^3e^2x \log(1 + ie^{2i(a+bx)}) + 12b^3efx^2 \log(1 + ie^{2i(a+bx)}) + 4b^3f^2x^3 \log(1 + ie^{2i(a+bx)})}{24b^3}$$

input

```
Integrate[(e + f*x)^2*ArcTanh[Cot[a + b*x]],x]
```

output

```
(x*(3*e^2 + 3*e*f*x + f^2*x^2)*ArcTanh[Cot[a + b*x]])/3 + (-12*b^3*e^2*x*L
og[1 - I*E^((2*I)*(a + b*x))] - 12*b^3*e*f*x^2*Log[1 - I*E^((2*I)*(a + b*x
))] - 4*b^3*f^2*x^3*Log[1 - I*E^((2*I)*(a + b*x))] + 12*b^3*e^2*x*Log[1 +
I*E^((2*I)*(a + b*x))] + 12*b^3*e*f*x^2*Log[1 + I*E^((2*I)*(a + b*x))] + 4
*b^3*f^2*x^3*Log[1 + I*E^((2*I)*(a + b*x))] - (6*I)*b^2*(e + f*x)^2*PolyLo
g[2, (-I)*E^((2*I)*(a + b*x))] + (6*I)*b^2*(e + f*x)^2*PolyLog[2, I*E^((2*
I)*(a + b*x))] + 6*b*e*f*PolyLog[3, (-I)*E^((2*I)*(a + b*x))] + 6*b*f^2*x*
PolyLog[3, (-I)*E^((2*I)*(a + b*x))] - 6*b*e*f*PolyLog[3, I*E^((2*I)*(a +
b*x))] - 6*b*f^2*x*PolyLog[3, I*E^((2*I)*(a + b*x))] + (3*I)*f^2*PolyLog[4
, (-I)*E^((2*I)*(a + b*x))] - (3*I)*f^2*PolyLog[4, I*E^((2*I)*(a + b*x))])
/(24*b^3)
```

### Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.16, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$ , Rules used = {6807, 3042, 4669, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) \, dx \\
 & \quad \downarrow \text{6807} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \frac{b \int (e + fx)^3 \sec(2a + 2bx) \, dx}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \frac{b \int (e + fx)^3 \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx}{3f} \\
 & \quad \downarrow \text{4669} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \\
 & \frac{b \left( -\frac{3f \int (e + fx)^2 \log(1 - ie^{2i(a + bx)}) \, dx}{2b} + \frac{3f \int (e + fx)^2 \log(1 + ie^{2i(a + bx)}) \, dx}{2b} - \frac{i(e + fx)^3 \operatorname{arctan}(e^{2i(a + bx)})}{b} \right)}{3f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \\
 & b \left( \frac{3f \left( \frac{i(e + fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a + bx)})}{2b} - \frac{if \int (e + fx) \operatorname{PolyLog}(2, -ie^{2i(a + bx)}) \, dx}{b} \right)}{2b} - \frac{3f \left( \frac{i(e + fx)^2 \operatorname{PolyLog}(2, ie^{2i(a + bx)})}{2b} - \frac{if \int (e + fx) \operatorname{PolyLog}(2, ie^{2i(a + bx)}) \, dx}{b} \right)}{2b} \right) \\
 & \quad \downarrow \text{7163}
 \end{aligned}$$

$$\frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \frac{b \left( \frac{3f}{2b} \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{if \int \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) dx}{2b} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right) \right)}{2b} - \frac{3f}{2b} \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} \right)}{3f}$$

↓ 2720

$$\frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \frac{b \left( \frac{3f}{2b} \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -ie^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right) \right)}{2b} - \frac{3f}{2b} \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{2b} \right)}{3f}$$

↓ 7143

$$\frac{(e + fx)^3 \operatorname{arctanh}(\cot(a + bx))}{3f} - \frac{b \left( \frac{3f}{2b} \left( \frac{i(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{2b} - \frac{if \left( \frac{f \operatorname{PolyLog}(4, -ie^{2i(a+bx)})}{4b^2} - \frac{i(e+fx) \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{2b} \right)}{b} \right) \right)}{2b} + \frac{i(e+fx)^3 \operatorname{arctan}(e^{2i(a+bx)})}{b}}{3f}$$

input `Int[(e + f*x)^2*ArcTanh[Cot[a + b*x]],x]`

output

```
((e + f*x)^3*ArcTanh[Cot[a + b*x]])/(3*f) - (b*((( -I)*(e + f*x)^3*ArcTan[E
^((2*I)*(a + b*x))])/b + (3*f*(((I/2)*(e + f*x)^2*PolyLog[2, (-I)*E^((2*I)
*(a + b*x))])/b - (I*f*((( -1/2*I)*(e + f*x)*PolyLog[3, (-I)*E^((2*I)*(a +
b*x))])/b + (f*PolyLog[4, (-I)*E^((2*I)*(a + b*x))])/(4*b^2))/b))/(2*b) -
(3*f*(((I/2)*(e + f*x)^2*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (I*f*((( -
1/2*I)*(e + f*x)*PolyLog[3, I*E^((2*I)*(a + b*x))])/b + (f*PolyLog[4, I*E^
((2*I)*(a + b*x))])/(4*b^2))/b))/(2*b)))/(3*f)
```

**Defintions of rubi rules used**

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6807 `Int[ArcTanh[Cot[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTanh[Cot[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 10.92 (sec) , antiderivative size = 2719, normalized size of antiderivative = 11.62

method	result	size
risch	Expression too large to display	2719

input `int((f*x+e)^2*arctanh(cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

1/8*I*f^2*polylog(4,-I*exp(2*I*(b*x+a)))/b^3-1/8*I*f^2*polylog(4,I*exp(2*I
*(b*x+a)))/b^3-f/b*a*e*ln(1+exp(I*(b*x+a)))*(-1)^(3/4)*x-f/b*a*e*ln(1-exp(
I*(b*x+a)))*(-1)^(3/4)*x+f*e/b*ln(I*exp(2*I*(b*x+a))+1)*a*x+I*f/b^2*a*e*di
log(1+exp(I*(b*x+a)))*(-1)^(3/4)+I*f/b^2*a*e*dilog(1-exp(I*(b*x+a)))*(-1)^(
3/4))-1/2*I*f*e/b*polylog(2,-I*exp(2*I*(b*x+a)))*x-1/2*I*f*e/b^2*polylog(2
,-I*exp(2*I*(b*x+a)))*a-1/4*f^2/b^2*polylog(3,I*exp(2*I*(b*x+a)))*x+1/3*f^
2/b^3*ln(-I*exp(2*I*(b*x+a))+1)*a^3-1/2*f*e*ln(-I*exp(2*I*(b*x+a))+1)*x^2-
1/2*f^2*ln(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))/b^3*a^3-1/2*f^2*ln(((I)
)^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))/b^3*a^3-1/4*f*e/b^2*polylog(3,I*exp(2*
I*(b*x+a)))+1/6*f^2/b^3*a^3*ln(exp(2*I*(b*x+a))+I)-1/2*ln(((I)^(1/2)-exp(
I*(b*x+a)))/(I)^(1/2))/b*a*e^2-1/2*ln(((I)^(1/2)+exp(I*(b*x+a)))/(I)^(1
/2))/b*a*e^2+1/2/b*a*e^2*ln(exp(2*I*(b*x+a))+I)+1/2*I/b*e^2*dilog(((I)^(1
/2)-exp(I*(b*x+a)))/(I)^(1/2))+1/2*I/b*e^2*dilog(((I)^(1/2)+exp(I*(b*x+a
)))/(I)^(1/2))+1/6/f*e^3*ln(-exp(2*I*(b*x+a))+I)+1/6*f^2*ln(I*exp(2*I*(b*
x+a))+1)*x^3+1/2*e^2*ln(1+exp(I*(b*x+a)))*(-1)^(3/4)*x+1/2*e^2*ln(1-exp(I*
(b*x+a)))*(-1)^(3/4)*x+f/b*a*e*ln(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))*
x+f/b*a*e*ln(((I)^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))*x-f*e/b*ln(-I*exp(2*I
*(b*x+a))+1)*a*x-I*f/b^2*a*e*dilog(((I)^(1/2)-exp(I*(b*x+a)))/(I)^(1/2))
-I*f/b^2*a*e*dilog(((I)^(1/2)+exp(I*(b*x+a)))/(I)^(1/2))+1/2*I*f*e/b*pol
ylog(2,I*exp(2*I*(b*x+a)))*x+1/2*I*f*e/b^2*polylog(2,I*exp(2*I*(b*x+a))...

```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1085 vs.  $2(180) = 360$ .

Time = 0.17 (sec) , antiderivative size = 1085, normalized size of antiderivative = 4.64

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(ax + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^2*arctanh(cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/48*(-3*I*f^2*polylog(4, I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) - 3*I*f^2
*polylog(4, I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) + 3*I*f^2*polylog(4, -I
*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) + 3*I*f^2*polylog(4, -I*cos(2*b*x +
2*a) - sin(2*b*x + 2*a)) - 6*(-I*b^2*f^2*x^2 - 2*I*b^2*e*f*x - I*b^2*e^2)*
dilog(I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) - 6*(-I*b^2*f^2*x^2 - 2*I*b^2
*e*f*x - I*b^2*e^2)*dilog(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) - 6*(I*b^
2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(-I*cos(2*b*x + 2*a) + sin(2*b
*x + 2*a)) - 6*(I*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(-I*cos(2*
b*x + 2*a) - sin(2*b*x + 2*a)) + 8*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b^3*e^
2*x)*log(-(cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/(cos(2*b*x + 2*a) - si
n(2*b*x + 2*a) + 1)) + 4*(3*a*b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(cos(2*b
*x + 2*a) + I*sin(2*b*x + 2*a) + I) - 4*(3*a*b^2*e^2 - 3*a^2*b*e*f + a^3*f
^2)*log(cos(2*b*x + 2*a) - I*sin(2*b*x + 2*a) + I) - 4*(b^3*f^2*x^3 + 3*b^
3*e*f*x^2 + 3*b^3*e^2*x + 3*a*b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(I*cos(2
*b*x + 2*a) + sin(2*b*x + 2*a) + 1) + 4*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b
^3*e^2*x + 3*a*b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(I*cos(2*b*x + 2*a) - s
in(2*b*x + 2*a) + 1) - 4*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b^3*e^2*x + 3*a*
b^2*e^2 - 3*a^2*b*e*f + a^3*f^2)*log(-I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a
) + 1) + 4*(b^3*f^2*x^3 + 3*b^3*e*f*x^2 + 3*b^3*e^2*x + 3*a*b^2*e^2 - 3*a^
2*b*e*f + a^3*f^2)*log(-I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a) + 1) + 4*...

```

### Sympy [F]

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \int (e + fx)^2 \operatorname{atanh}(\cot(a + bx)) dx$$

input

```
integrate((f*x+e)**2*atanh(cot(b*x+a)),x)
```

output

```
Integral((e + f*x)**2*atanh(cot(a + b*x)), x)
```

**Maxima [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e)^2 \operatorname{artanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)^2*arctanh(cot(b*x+a)),x, algorithm="maxima")`

output `1/12*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/12*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(2/3*((b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f^2*x^3 + 3*b*e*f*x^2 + 3*b*e^2*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

**Giac [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e)^2 \operatorname{artanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)^2*arctanh(cot(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)^2*arctanh(cot(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(a + bx)) (e + fx)^2 dx$$

input `int(atanh(cot(a + b*x))*(e + f*x)^2,x)`

output `int(atanh(cot(a + b*x))*(e + f*x)^2, x)`



**Reduce [F]**

$$\int (e + fx)^2 \operatorname{arctanh}(\cot(a + bx)) dx = \left( \int \operatorname{atanh}(\cot(bx + a)) dx \right) e^2$$

$$+ \left( \int \operatorname{atanh}(\cot(bx + a)) x^2 dx \right) f^2$$

$$+ 2 \left( \int \operatorname{atanh}(\cot(bx + a)) x dx \right) ef$$

input `int((f*x+e)^2*atanh(cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)),x)*e**2 + int(atanh(cot(a + b*x))*x**2,x)*f**2 + 2  
*int(atanh(cot(a + b*x))*x,x)*e*f`

### 3.331 $\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx$

Optimal result	2325
Mathematica [A] (verified)	2326
Rubi [A] (verified)	2326
Maple [C] (warning: unable to verify)	2329
Fricas [B] (verification not implemented)	2330
Sympy [F]	2330
Maxima [F]	2331
Giac [F]	2331
Mupad [F(-1)]	2332
Reduce [F]	2332

#### Optimal result

Integrand size = 13, antiderivative size = 162

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = \frac{i(e + fx)^2 \arctan(e^{2i(a+bx)})}{2f} + \frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} + \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b} + \frac{f \operatorname{PolyLog}(3, -ie^{2i(a+bx)})}{8b^2} - \frac{f \operatorname{PolyLog}(3, ie^{2i(a+bx)})}{8b^2}$$

output `1/2*I*(f*x+e)^2*arctan(exp(2*I*(b*x+a)))/f+1/2*(f*x+e)^2*arctanh(cot(b*x+a))/f-1/4*I*(f*x+e)*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*(f*x+e)*polylog(2,I*exp(2*I*(b*x+a)))/b+1/8*f*polylog(3,-I*exp(2*I*(b*x+a)))/b^2-1/8*f*polylog(3,I*exp(2*I*(b*x+a)))/b^2`

**Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.82

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = ex \operatorname{arctanh}(\cot(a + bx)) + \frac{1}{2} f x^2 \operatorname{arctanh}(\cot(a + bx)) - \frac{e((-4a + \pi - 4bx) (\log(1 - ie^{-2i(a+bx)}) - \log(1 + ie^{-2i(a+bx)})) - (-4a + \pi) \log(\cot(a + \frac{\pi}{4} + bx)))}{8b} + \frac{f(4ib^2 x^2 \arctan(\cos(2(a + bx)) + i \sin(2(a + bx))) + 2ibx \operatorname{PolyLog}(2, i \cos(2(a + bx)) - \sin(2(a + bx)))}{8b}$$

input `Integrate[(e + f*x)*ArcTanh[Cot[a + b*x]],x]`

output `e*x*ArcTanh[Cot[a + b*x]] + (f*x^2*ArcTanh[Cot[a + b*x]])/2 - (e*((-4*a + Pi - 4*b*x)*(Log[1 - I/E^((2*I)*(a + b*x))] - Log[1 + I/E^((2*I)*(a + b*x))]) - (-4*a + Pi)*Log[Cot[a + Pi/4 + b*x]] + (2*I)*(PolyLog[2, (-I)/E^((2*I)*(a + b*x))] - PolyLog[2, I/E^((2*I)*(a + b*x))])))/(8*b) + (f*((4*I)*b^2*x^2*ArcTan[Cos[2*(a + b*x)] + I*Sin[2*(a + b*x)]] + (2*I)*b*x*PolyLog[2, I*Cos[2*(a + b*x)] - Sin[2*(a + b*x)]] - (2*I)*b*x*PolyLog[2, (-I)*Cos[2*(a + b*x)] + Sin[2*(a + b*x)]] - PolyLog[3, I*Cos[2*(a + b*x)] - Sin[2*(a + b*x)]] + PolyLog[3, (-I)*Cos[2*(a + b*x)] + Sin[2*(a + b*x)]]))/(8*b^2)`

**Rubi [A] (verified)**

Time = 0.64 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {6807, 3042, 4669, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx$$

$$\downarrow 6807$$

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \int (e + fx)^2 \sec(2a + 2bx) dx}{2f}$$

$$\downarrow 3042$$

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \int (e + fx)^2 \csc\left(2a + 2bx + \frac{\pi}{2}\right) dx}{2f}$$

↓ 4669

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \left( -\frac{f \int (e + fx) \log(1 - ie^{2i(a+bx)}) dx}{b} + \frac{f \int (e + fx) \log(1 + ie^{2i(a+bx)}) dx}{b} - \frac{i(e + fx)^2 \arctan(e^{2i(a+bx)})}{b} \right)}{2f}$$

↓ 3011

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \left( \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, -ie^{2i(a+bx)}\right)}{2b} - \frac{if \int \operatorname{PolyLog}\left(2, -ie^{2i(a+bx)}\right) dx}{2b} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, ie^{2i(a+bx)}\right)}{2b} - \frac{if \int \operatorname{PolyLog}\left(2, ie^{2i(a+bx)}\right) dx}{2b} \right)}{b} \right)}{2f}$$


---

↓ 2720

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \left( \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, -ie^{2i(a+bx)}\right)}{2b} - \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}\left(2, -ie^{2i(a+bx)}\right) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, ie^{2i(a+bx)}\right)}{2b} - \frac{f \int e^{-2i(a+bx)} \operatorname{PolyLog}\left(2, ie^{2i(a+bx)}\right) de^{2i(a+bx)}}{4b^2} \right)}{b} \right)}{2f}$$


---

↓ 7143

$$\frac{(e + fx)^2 \operatorname{arctanh}(\cot(a + bx))}{2f} - \frac{b \left( -\frac{i(e + fx)^2 \arctan(e^{2i(a+bx)})}{b} + \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, -ie^{2i(a+bx)}\right)}{2b} - \frac{f \operatorname{PolyLog}\left(3, -ie^{2i(a+bx)}\right)}{4b^2} \right)}{b} - \frac{f \left( \frac{i(e + fx) \operatorname{PolyLog}\left(2, ie^{2i(a+bx)}\right)}{2b} - \frac{f \operatorname{PolyLog}\left(3, ie^{2i(a+bx)}\right)}{4b^2} \right)}{b} \right)}{2f}$$

input

```
Int[(e + f*x)*ArcTanh[Cot[a + b*x]], x]
```

output

```
((e + f*x)^2*ArcTanh[Cot[a + b*x]])/(2*f) - (b*((( -I)*(e + f*x)^2*ArcTan[E
^((2*I)*(a + b*x))])/b + (f*(((I/2)*(e + f*x)*PolyLog[2, (-I)*E^((2*I)*(a
+ b*x))])/b - (f*PolyLog[3, (-I)*E^((2*I)*(a + b*x))])/((4*b^2)))/b - (f*((
(I/2)*(e + f*x)*PolyLog[2, I*E^((2*I)*(a + b*x))])/b - (f*PolyLog[3, I*E^((
2*I)*(a + b*x))])/((4*b^2)))/b))/b)/(2*f)
```

## Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4669

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol
] :=> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Si
mp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x],
x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x
))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6807

```
Int[ArcTanh[Cot[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:=> Simp[(e + f*x)^(m + 1)*(ArcTanh[Cot[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b
, e, f}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.64 (sec) , antiderivative size = 1819, normalized size of antiderivative = 11.23

method	result	size
risch	Expression too large to display	1819

input

```
int((f*x+e)*arctanh(cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*f/b^2*a^2*ln(1+exp(I*(b*x+a))*(-1)^(3/4))-1/2*f/b^2*a^2*ln(1-exp(I*(b
*x+a))*(-1)^(3/4))-1/2*e/b*ln(((I)^(1/2)-exp(I*(b*x+a)))/((I)^(1/2)))*a-1/
2*e/b*ln(((I)^(1/2)+exp(I*(b*x+a)))/((I)^(1/2)))*a+1/2*I*e/b*dilog(((I)^(
1/2)-exp(I*(b*x+a)))/((I)^(1/2))+1/2*I*e/b*dilog(((I)^(1/2)+exp(I*(b*x+a)
))/((I)^(1/2))+1/2*f/b^2*a^2*ln(((I)^(1/2)-exp(I*(b*x+a)))/((I)^(1/2))+1/
2*f/b^2*a^2*ln(((I)^(1/2)+exp(I*(b*x+a)))/((I)^(1/2))-1/4/b^2*f*a^2*ln(ex
p(2*I*(b*x+a))+I)+1/2/b*a*e*ln(exp(2*I*(b*x+a))+I)-1/4*ln(exp(2*I*(b*x+a)
-I)*f*x^2-1/2*ln(exp(2*I*(b*x+a))-I)*e*x-1/2*(-1/2*f*x^2-e*x)*ln(exp(2*I*(
b*x+a))+I)+1/2*e/b*ln(1+exp(I*(b*x+a))*(-1)^(3/4))*a+1/2*e/b*ln(1-exp(I*(b
*x+a))*(-1)^(3/4))*a-1/2*I*e/b*dilog(1+exp(I*(b*x+a))*(-1)^(3/4))-1/2*I*e/
b*dilog(1-exp(I*(b*x+a))*(-1)^(3/4))+1/4/b^2*f*ln(I*exp(2*I*(b*x+a))+1)*a^
2+1/8*f*polylog(3,-I*exp(2*I*(b*x+a)))/b^2-1/8*f*polylog(3,I*exp(2*I*(b*x+
a)))/b^2-1/4/b^2*f*ln(-I*exp(2*I*(b*x+a))+1)*a^2-1/2*e/b*a*ln(-exp(2*I*(b*
x+a))+I)+1/4*f/b^2*a^2*ln(-exp(2*I*(b*x+a))+I)-1/4*I*Pi*(csgn(I*(exp(2*I*(
b*x+a))+I)/(-1+exp(2*I*(b*x+a))))*csgn((1-I)*(exp(2*I*(b*x+a))+I)/(-1+exp(
2*I*(b*x+a))))+csgn((1-I)*(exp(2*I*(b*x+a))+I)/(-1+exp(2*I*(b*x+a))))^2-cs
gn(I*(exp(2*I*(b*x+a))-I))*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I*(exp(2*I*(
b*x+a))-I)/(-1+exp(2*I*(b*x+a))))+csgn(I*(exp(2*I*(b*x+a))-I))*csgn(I*(exp
(2*I*(b*x+a))-I)/(-1+exp(2*I*(b*x+a))))^2+csgn(I*(exp(2*I*(b*x+a))+I))*csg
n(I/(-1+exp(2*I*(b*x+a))))*csgn(I*(exp(2*I*(b*x+a))+I)/(-1+exp(2*I*(b*x...
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 681 vs.  $2(130) = 260$ .

Time = 0.14 (sec) , antiderivative size = 681, normalized size of antiderivative = 4.20

$$\int (e + fx)\operatorname{arctanh}(\cot(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)*arctanh(cot(b*x+a)),x, algorithm="fricas")`

output

```
-1/16*(2*(-I*b*f*x - I*b*e)*dilog(I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) +
  2*(-I*b*f*x - I*b*e)*dilog(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) + 2*(I*
  b*f*x + I*b*e)*dilog(-I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) + 2*(I*b*f*x
  + I*b*e)*dilog(-I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) - 4*(b^2*f*x^2 + 2*
  b^2*e*x)*log(-(cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/(cos(2*b*x + 2*a)
  - sin(2*b*x + 2*a) + 1)) - 2*(2*a*b*e - a^2*f)*log(cos(2*b*x + 2*a) + I*si
  n(2*b*x + 2*a) + I) + 2*(2*a*b*e - a^2*f)*log(cos(2*b*x + 2*a) - I*sin(2*b
  *x + 2*a) + I) + 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log(I*cos(2*b
  *x + 2*a) + sin(2*b*x + 2*a) + 1) - 2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a
  ^2*f)*log(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a) + 1) + 2*(b^2*f*x^2 + 2*b^
  2*e*x + 2*a*b*e - a^2*f)*log(-I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1) -
  2*(b^2*f*x^2 + 2*b^2*e*x + 2*a*b*e - a^2*f)*log(-I*cos(2*b*x + 2*a) - sin
  (2*b*x + 2*a) + 1) - 2*(2*a*b*e - a^2*f)*log(-cos(2*b*x + 2*a) + I*sin(2*b
  *x + 2*a) + I) + 2*(2*a*b*e - a^2*f)*log(-cos(2*b*x + 2*a) - I*sin(2*b*x +
  2*a) + I) - f*polylog(3, I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) + f*polyl
  og(3, I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) - f*polylog(3, -I*cos(2*b*x +
  2*a) + sin(2*b*x + 2*a)) + f*polylog(3, -I*cos(2*b*x + 2*a) - sin(2*b*x +
  2*a)))/b^2
```

**Sympy [F]**

$$\int (e + fx)\operatorname{arctanh}(\cot(a + bx)) dx = \int (e + fx)\operatorname{atanh}(\cot(a + bx)) dx$$

input `integrate((f*x+e)*atanh(cot(b*x+a)),x)`

output `Integral((e + f*x)*atanh(cot(a + b*x)), x)`

### Maxima [F]

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e) \operatorname{artanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)*arctanh(cot(b*x+a)),x, algorithm="maxima")`

output `1/8*(f*x^2 + 2*e*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 + 4*sin(2*b*x + 2*a) + 2) - 1/8*(f*x^2 + 2*e*x)*log(2*cos(2*b*x + 2*a)^2 + 2*sin(2*b*x + 2*a)^2 - 4*sin(2*b*x + 2*a) + 2) - integrate(((b*f*x^2 + 2*b*e*x)*cos(4*b*x + 4*a)*cos(2*b*x + 2*a) + (b*f*x^2 + 2*b*e*x)*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + (b*f*x^2 + 2*b*e*x)*cos(2*b*x + 2*a))/(cos(4*b*x + 4*a)^2 + sin(4*b*x + 4*a)^2 + 2*cos(4*b*x + 4*a) + 1), x)`

### Giac [F]

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = \int (fx + e) \operatorname{artanh}(\cot(bx + a)) dx$$

input `integrate((f*x+e)*arctanh(cot(b*x+a)),x, algorithm="giac")`

output `integrate((f*x + e)*arctanh(cot(b*x + a)), x)`



**Mupad [F(-1)]**

Timed out.

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(a + bx)) (e + fx) dx$$

input `int(atanh(cot(a + b*x))*(e + f*x),x)`output `int(atanh(cot(a + b*x))*(e + f*x), x)`**Reduce [F]**

$$\int (e + fx) \operatorname{arctanh}(\cot(a + bx)) dx = \left( \int \operatorname{atanh}(\cot(bx + a)) dx \right) e + \left( \int \operatorname{atanh}(\cot(bx + a)) x dx \right) f$$

input `int((f*x+e)*atanh(cot(b*x+a)),x)`output `int(atanh(cot(a + b*x)),x)*e + int(atanh(cot(a + b*x))*x,x)*f`

### 3.332 $\int \operatorname{arctanh}(\cot(a + bx)) dx$

Optimal result	2333
Mathematica [A] (verified)	2333
Rubi [A] (verified)	2334
Maple [A] (verified)	2336
Fricas [B] (verification not implemented)	2336
Sympy [F]	2337
Maxima [B] (verification not implemented)	2337
Giac [F]	2338
Mupad [F(-1)]	2338
Reduce [F]	2339

#### Optimal result

Integrand size = 7, antiderivative size = 79

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = ix \arctan(e^{2i(a+bx)}) + x \operatorname{arctanh}(\cot(a + bx)) - \frac{i \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b} + \frac{i \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b}$$

output

```
I*x*arctan(exp(2*I*(b*x+a)))+x*arctanh(cot(b*x+a))-1/4*I*polylog(2,-I*exp(2*I*(b*x+a)))/b+1/4*I*polylog(2,I*exp(2*I*(b*x+a)))/b
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.61

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = x \operatorname{arctanh}(\cot(a + bx)) - \frac{(-4a + \pi - 4bx) (\log(1 - ie^{-2i(a+bx)}) - \log(1 + ie^{-2i(a+bx)})) - (-4a + \pi) \log(\cot(a + \frac{\pi}{4} + bx))}{8b}$$

input

```
Integrate[ArcTanh[Cot[a + b*x]],x]
```

output

```
x*ArcTanh[Cot[a + b*x]] - ((-4*a + Pi - 4*b*x)*(Log[1 - I/E^((2*I)*(a + b*x))] - Log[1 + I/E^((2*I)*(a + b*x))]) - (-4*a + Pi)*Log[Cot[a + Pi/4 + b*x]] + (2*I)*(PolyLog[2, (-I)/E^((2*I)*(a + b*x))] - PolyLog[2, I/E^((2*I)*(a + b*x))]))/(8*b)
```

**Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.09, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$ , Rules used = {6803, 3042, 4669, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arctanh}(\cot(a + bx)) \, dx \\
 & \quad \downarrow \text{6803} \\
 & x \operatorname{arctanh}(\cot(a + bx)) - b \int x \sec(2a + 2bx) \, dx \\
 & \quad \downarrow \text{3042} \\
 & x \operatorname{arctanh}(\cot(a + bx)) - b \int x \csc\left(2a + 2bx + \frac{\pi}{2}\right) \, dx \\
 & \quad \downarrow \text{4669} \\
 & x \operatorname{arctanh}(\cot(a + bx)) - \\
 & b \left( -\frac{\int \log(1 - ie^{2i(a+bx)}) \, dx}{2b} + \frac{\int \log(1 + ie^{2i(a+bx)}) \, dx}{2b} - \frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} \right) \\
 & \quad \downarrow \text{2715} \\
 & b \left( \frac{i \int e^{-2i(a+bx)} \log(1 - ie^{2i(a+bx)}) \, de^{2i(a+bx)}}{4b^2} - \frac{i \int e^{-2i(a+bx)} \log(1 + ie^{2i(a+bx)}) \, de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} \right) \\
 & \quad \downarrow \text{2838} \\
 & b \left( -\frac{ix \operatorname{arctan}(e^{2i(a+bx)})}{b} + \frac{x \operatorname{arctanh}(\cot(a + bx)) - i \operatorname{PolyLog}(2, -ie^{2i(a+bx)})}{4b^2} - \frac{i \operatorname{PolyLog}(2, ie^{2i(a+bx)})}{4b^2} \right)
 \end{aligned}$$

input `Int[ArcTanh[Cot[a + b*x]],x]`

output `x*ArcTanh[Cot[a + b*x]] - b*((( -I)*x*ArcTan[E^((2*I)*(a + b*x))])/b + ((I/4)*PolyLog[2, (-I)*E^((2*I)*(a + b*x))])/b^2 - ((I/4)*PolyLog[2, I*E^((2*I)*(a + b*x))])/b^2)`

### Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4669 `Int[csc[(e_) + Pi*(k_) + (f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol]
:> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6803 `Int[ArcTanh[Cot[(a_) + (b_)*(x_)]], x_Symbol] :> Simp[x*ArcTanh[Cot[a + b*x]], x] - Simp[b Int[x*Sec[2*a + 2*b*x], x], x] /; FreeQ[{a, b}, x]`

### Maple [A] (verified)

Time = 2.01 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.52

method	result
parts	$x \operatorname{arctanh}(\cot(bx + a)) - \frac{(bx+a) \ln(i e^{2i(bx+a)} + 1)}{2} + \frac{(bx+a) \ln(-i e^{2i(bx+a)} + 1)}{2} + \frac{i \operatorname{dilog}(i e^{2i(bx+a)} + 1)}{4} - \frac{i \operatorname{dilog}(-i e^{2i(bx+a)} + 1)}{4}$
derivativedivides	$\frac{i \operatorname{arctanh}(\cot(bx+a)) \left( -\ln\left(1 - \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right) + \ln\left(1 + \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right) \right)}{2} + \frac{i \operatorname{dilog}\left(1 + \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right)}{4} - \frac{i \operatorname{dilog}\left(1 - \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right)}{4}$
default	$\frac{i \operatorname{arctanh}(\cot(bx+a)) \left( -\ln\left(1 - \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right) + \ln\left(1 + \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right) \right)}{2} + \frac{i \operatorname{dilog}\left(1 + \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right)}{4} - \frac{i \operatorname{dilog}\left(1 - \frac{i(\cot(bx+a)+1)^2}{1-\cot(bx+a)^2}\right)}{4}$
risch	Expression too large to display

```
input int(arctanh(cot(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output x*arctanh(cot(b*x+a))-1/b*(-1/2*(b*x+a)*ln(I*exp(2*I*(b*x+a))+1)+1/2*(b*x+a)*ln(-I*exp(2*I*(b*x+a))+1)+1/4*I*dilog(I*exp(2*I*(b*x+a))+1)-1/4*I*dilog(-I*exp(2*I*(b*x+a))+1)-1/2*a*ln(sec(2*b*x+2*a)+tan(2*b*x+2*a)))
```

### Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 389 vs. 2(57) = 114.

Time = 0.13 (sec) , antiderivative size = 389, normalized size of antiderivative = 4.92

$$\int \operatorname{arctanh}(\cot(a + bx)) dx$$

$$= \frac{4bx \log\left(-\frac{\cos(2bx+2a)+\sin(2bx+2a)+1}{\cos(2bx+2a)-\sin(2bx+2a)+1}\right) + 2a \log(\cos(2bx+2a) + i \sin(2bx+2a) + i) - 2a \log(\cos(2bx+2a) - i \sin(2bx+2a) - i)}{2}$$

```
input integrate(arctanh(cot(b*x+a)), x, algorithm="fricas")
```

output

```
1/8*(4*b*x*log(-cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/(cos(2*b*x + 2*a)
) - sin(2*b*x + 2*a) + 1)) + 2*a*log(cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a)
+ I) - 2*a*log(cos(2*b*x + 2*a) - I*sin(2*b*x + 2*a) + I) - 2*(b*x + a)*l
og(I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1) + 2*(b*x + a)*log(I*cos(2*b*
x + 2*a) - sin(2*b*x + 2*a) + 1) - 2*(b*x + a)*log(-I*cos(2*b*x + 2*a) + s
in(2*b*x + 2*a) + 1) + 2*(b*x + a)*log(-I*cos(2*b*x + 2*a) - sin(2*b*x + 2
*a) + 1) + 2*a*log(-cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a) + I) - 2*a*log(-
cos(2*b*x + 2*a) - I*sin(2*b*x + 2*a) + I) + I*dilog(I*cos(2*b*x + 2*a) +
sin(2*b*x + 2*a)) + I*dilog(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a)) - I*dil
og(-I*cos(2*b*x + 2*a) + sin(2*b*x + 2*a)) - I*dilog(-I*cos(2*b*x + 2*a) -
sin(2*b*x + 2*a)))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(a + bx)) dx$$

input

```
integrate(atanh(cot(b*x+a)),x)
```

output

```
Integral(atanh(cot(a + b*x)), x)
```

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 184 vs.  $2(57) = 114$ .

Time = 0.16 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.33

$$\int \operatorname{arctanh}(\cot(a + bx)) dx$$

$$= \frac{4(bx + a) \operatorname{artanh}\left(\frac{1}{\tan(bx+a)}\right) + \left(\arctan\left(\frac{1}{2} \tan(bx + a) + \frac{1}{2}, \frac{1}{2} \tan(bx + a) + \frac{1}{2}\right) - \arctan\left(\frac{1}{2} \tan(bx + a) + \frac{1}{2}\right)\right)}{b}$$

input

```
integrate(arctanh(cot(b*x+a)),x, algorithm="maxima")
```

output

```
1/4*(4*(b*x + a)*arctanh(1/tan(b*x + a)) + (arctan2(1/2*tan(b*x + a) + 1/2
, 1/2*tan(b*x + a) + 1/2) - arctan2(1/2*tan(b*x + a) - 1/2, -1/2*tan(b*x +
a) + 1/2))*log(tan(b*x + a)^2 + 1) - (b*x + a)*log(1/2*tan(b*x + a)^2 + t
an(b*x + a) + 1/2) + (b*x + a)*log(1/2*tan(b*x + a)^2 - tan(b*x + a) + 1/2
) - I*dilog((1/2*I + 1/2)*tan(b*x + a) - 1/2*I + 1/2) + I*dilog(-(1/2*I -
1/2)*tan(b*x + a) + 1/2*I + 1/2) + I*dilog((1/2*I - 1/2)*tan(b*x + a) + 1/
2*I + 1/2) - I*dilog(-(1/2*I + 1/2)*tan(b*x + a) - 1/2*I + 1/2))/b
```

**Giac [F]**

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{artanh}(\cot(bx + a)) dx$$

input

```
integrate(arctanh(cot(b*x+a)),x, algorithm="giac")
```

output

```
integrate(arctanh(cot(b*x + a)), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(a + bx)) dx$$

input

```
int(atanh(cot(a + b*x)),x)
```

output

```
int(atanh(cot(a + b*x)), x)
```

**Reduce [F]**

$$\int \operatorname{arctanh}(\cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a)) dx$$

input `int(atanh(cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)),x)`



### 3.333 $\int \frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx} dx$

Optimal result	2340
Mathematica [N/A]	2340
Rubi [N/A]	2341
Maple [N/A]	2341
Fricas [N/A]	2342
Sympy [N/A]	2342
Maxima [N/A]	2342
Giac [N/A]	2343
Mupad [N/A]	2343
Reduce [N/A]	2344

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx}, x\right)$$

output `Defer(Int)(arctanh(cot(b*x+a))/(f*x+e), x)`

#### Mathematica [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx} dx = \int \frac{\operatorname{arctanh}(\cot(a+bx))}{e+fx} dx$$

input `Integrate[ArcTanh[Cot[a + b*x]]/(e + f*x), x]`

output `Integrate[ArcTanh[Cot[a + b*x]]/(e + f*x), x]`

**Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx$$

input `Int[ArcTanh[Cot[a + b*x]]/(e + f*x),x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(\cot(bx + a))}{fx + e} dx$$

input `int(arctanh(cot(b*x+a))/(f*x+e),x)`

output `int(arctanh(cot(b*x+a))/(f*x+e),x)`

**Fricas [N/A]**

Not integrable

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\cot(bx + a))}{fx + e} dx$$

input `integrate(arctanh(cot(b*x+a))/(f*x+e),x, algorithm="fricas")`

output `integral(arctanh(cot(b*x + a))/(f*x + e), x)`

**Sympy [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\cot(a + bx))}{e + fx} dx$$

input `integrate(atanh(cot(b*x+a))/(f*x+e),x)`

output `Integral(atanh(cot(a + b*x))/(e + f*x), x)`

**Maxima [N/A]**

Not integrable

Time = 1.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\cot(bx + a))}{fx + e} dx$$

input `integrate(arctanh(cot(b*x+a))/(f*x+e),x, algorithm="maxima")`

output `integrate(arctanh(cot(b*x + a))/(f*x + e), x)`

### Giac [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{artanh}(\cot(bx + a))}{fx + e} dx$$

input `integrate(arctanh(cot(b*x+a))/(f*x+e),x, algorithm="giac")`

output `integrate(arctanh(cot(b*x + a))/(f*x + e), x)`

### Mupad [N/A]

Not integrable

Time = 3.43 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctanh(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\cot(a + bx))}{e + fx} dx$$

input `int(atanh(cot(a + b*x))/(e + f*x),x)`

output `int(atanh(cot(a + b*x))/(e + f*x), x)`

**Reduce [N/A]**

Not integrable

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(\cot(a + bx))}{e + fx} dx = \int \frac{\operatorname{atanh}(\cot(bx + a))}{fx + e} dx$$

input `int(atanh(cot(b*x+a))/(f*x+e),x)`output `int(atanh(cot(a + b*x))/(e + f*x),x)`

### 3.334 $\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx$

Optimal result	2345
Mathematica [A] (verified)	2346
Rubi [A] (verified)	2347
Maple [C] (warning: unable to verify)	2353
Fricas [B] (verification not implemented)	2353
Sympy [F]	2354
Maxima [F]	2355
Giac [F]	2355
Mupad [F(-1)]	2356
Reduce [F]	2356

#### Optimal result

Integrand size = 15, antiderivative size = 391

$$\begin{aligned}
 \int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx &= \frac{1}{3} x^3 \operatorname{arctanh}(c + d \cot(a + bx)) \\
 &+ \frac{1}{6} x^3 \log \left( 1 - \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right) \\
 &- \frac{1}{6} x^3 \log \left( 1 - \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right) \\
 &- \frac{ix^2 \operatorname{PolyLog} \left( 2, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{4b} \\
 &+ \frac{ix^2 \operatorname{PolyLog} \left( 2, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{4b} \\
 &+ \frac{x \operatorname{PolyLog} \left( 3, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{4b^2} \\
 &- \frac{x \operatorname{PolyLog} \left( 3, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{4b^2} \\
 &+ \frac{i \operatorname{PolyLog} \left( 4, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{8b^3} \\
 &- \frac{i \operatorname{PolyLog} \left( 4, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{8b^3}
 \end{aligned}$$

output

$$\begin{aligned} & 1/3*x^3*\operatorname{arctanh}(c+d*\cot(b*x+a))+1/6*x^3*\ln(1-(1-c-I*d)*\exp(2*I*a+2*I*b*x)/ \\ & (1-c+I*d))-1/6*x^3*\ln(1-(1+c+I*d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))-1/4*I*x^2* \\ & \operatorname{polylog}(2,(1-c-I*d)*\exp(2*I*a+2*I*b*x)/(1-c+I*d))/b+1/4*I*x^2*\operatorname{polylog}(2,(1 \\ & +c+I*d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))/b+1/4*x*\operatorname{polylog}(3,(1-c-I*d)*\exp(2*I* \\ & a+2*I*b*x)/(1-c+I*d))/b^2-1/4*x*\operatorname{polylog}(3,(1+c+I*d)*\exp(2*I*a+2*I*b*x)/(1+ \\ & c-I*d))/b^2+1/8*I*\operatorname{polylog}(4,(1-c-I*d)*\exp(2*I*a+2*I*b*x)/(1-c+I*d))/b^3-1/ \\ & 8*I*\operatorname{polylog}(4,(1+c+I*d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))/b^3 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 3.33 (sec) , antiderivative size = 341, normalized size of antiderivative = 0.87

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx$$

$$= \frac{8b^3 x^3 \operatorname{arctanh}(c + d \cot(a + bx)) + 4b^3 x^3 \log\left(1 + \frac{(1-c+id)e^{-2i(a+bx)}}{-1+c+id}\right) - 4b^3 x^3 \log\left(1 + \frac{(-1-c+id)e^{-2i(a+bx)}}{1+c+id}\right)}{24b^3}$$

input

`Integrate[x^2*ArcTanh[c + d*Cot[a + b*x]],x]`

output

$$\begin{aligned} & (8*b^3*x^3*\operatorname{ArcTanh}[c + d*\operatorname{Cot}[a + b*x]] + 4*b^3*x^3*\operatorname{Log}[1 + (1 - c + I*d)/ \\ & (-1 + c + I*d)*E^{((2*I)*(a + b*x))}] - 4*b^3*x^3*\operatorname{Log}[1 + (-1 - c + I*d)/ \\ & (1 + c + I*d)*E^{((2*I)*(a + b*x))}] + (6*I)*b^2*x^2*\operatorname{PolyLog}[2, (-1 + c - I* \\ & d)/((-1 + c + I*d)*E^{((2*I)*(a + b*x))})] - (6*I)*b^2*x^2*\operatorname{PolyLog}[2, (1 + c \\ & - I*d)/((1 + c + I*d)*E^{((2*I)*(a + b*x))})] + 6*b*x*\operatorname{PolyLog}[3, (-1 + c - \\ & I*d)/((-1 + c + I*d)*E^{((2*I)*(a + b*x))})] - 6*b*x*\operatorname{PolyLog}[3, (1 + c - I*d) \\ & ]/((1 + c + I*d)*E^{((2*I)*(a + b*x))})] - (3*I)*\operatorname{PolyLog}[4, (-1 + c - I*d)/ \\ & (-1 + c + I*d)*E^{((2*I)*(a + b*x))}] + (3*I)*\operatorname{PolyLog}[4, (1 + c - I*d)/ \\ & (1 + c + I*d)*E^{((2*I)*(a + b*x))})])/(24*b^3) \end{aligned}$$

**Rubi [A] (verified)**

Time = 1.50 (sec) , antiderivative size = 514, normalized size of antiderivative = 1.31, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6823, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}(d \cot(a + bx) + c) dx \\
 & \quad \downarrow \text{6823} \\
 & -\frac{1}{3}b(-ic + d + i) \int \frac{e^{2ia+2ibx} x^3}{-c - (-c - id + 1)e^{2ia+2ibx} + id + 1} dx + \frac{1}{3}b(-d + i(c + 1)) \int \frac{e^{2ia+2ibx} x^3}{c - (c + id + 1)e^{2ia+2ibx} - id + 1} dx + \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & -\frac{1}{3}b(-ic + d + i) \left( \frac{3 \int x^2 \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) dx}{2b(d + i(1 - c))} - \frac{x^3 \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d + i(1 - c))} \right) + \\
 & \frac{1}{3}b(-d + i(c + 1)) \left( \frac{3 \int x^2 \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) dx}{2(-bd + i(bc + b))} - \frac{x^3 \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd + i(bc + b))} \right) + \\
 & \quad \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$



$$\begin{aligned}
& -\frac{1}{3}b(-ic + d + \\
i) & \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{i \int x \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right) dx}{b} \right)}{2b(d+i(1-c))} - \frac{x^3 \log\left(1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b(d+i(1-c))} \right) + \\
& \\
& \frac{1}{3}b(-d + i(c + \\
1)) & \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{i \int x \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right) dx}{b} \right)}{2(-bd+i(bc+b))} - \frac{x^3 \log\left(1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2(-bd+i(bc+b))} \right) + \\
& \\
& \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + c)
\end{aligned}$$

↓ 7163

$$i) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{-\frac{1}{3}b(-ic+d + i \left( \frac{i \int \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right) dx}{2b} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b}\right)}{b} \right)}{2b(d+i(1-c))} - \frac{x^3 \log(\dots)}{2} \right)$$

$$1)) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{\frac{1}{3}b(-d+i(c + i \left( \frac{i \int \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right) dx}{2b} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b}\right)}{b} \right)}{2(-bd+i(bc+b))} - \frac{x^3 \log(\dots)}{2} \right)$$

$$\frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a+bx) + c)$$

↓ 2720

$$\begin{aligned}
 & i) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{-\frac{1}{3}b(-ic+d+ \int e^{-2ia-2ibx} \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right) de^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b}}{b} \right)}{2b(d+i(1-c))} \right. \\
 & 1)) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{\frac{1}{3}b(-d+i(c+ \int e^{-2ia-2ibx} \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right) de^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b}}{b} \right)}{2(-bd+i(bc+b))} \right. \\
 & \left. \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a+bx) + c) \right)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + c) - \frac{1}{3}b(-ic + d + \\
 & \left. i \left( \frac{\operatorname{PolyLog}\left(4, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} \right) \right) \\
 i) & \frac{\left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{\left( \frac{\operatorname{PolyLog}\left(4, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} \right)}{b} \right)}{2b(d + i(1 - c))} - \frac{x^3 \log\left(1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2} \\
 & \frac{1}{3}b(-d + i(c + \\
 & \left. i \left( \frac{\operatorname{PolyLog}\left(4, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} \right) \right) \\
 1)) & \frac{\left( \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{\left( \frac{\operatorname{PolyLog}\left(4, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} \right)}{b} \right)}{2(-bd + i(bc + b))} - \frac{x^3 \log\left(1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2(-bd + i(bc + b))}
 \end{aligned}$$

input `Int[x^2*ArcTanh[c + d*Cot[a + b*x]],x]`

output `(x^3*ArcTanh[c + d*Cot[a + b*x]])/3 - (b*(I - I*c + d)*(-1/2*(x^3*Log[1 - ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)])/(b*(I*(1 - c) + d) + (3*(((I/2)*x^2*PolyLog[2, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)])/b - (I*(((1/2)*x*PolyLog[3, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)])/b + PolyLog[4, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)]/(4*b^2)))/b)/(2*b*(I*(1 - c) + d)))/3 + (b*(I*(1 + c) - d)*(-1/2*(x^3*Log[1 - ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)])/(I*(b + b*c) - b*d) + (3*(((I/2)*x^2*PolyLog[2, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)])/b - (I*(((1/2)*x*PolyLog[3, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)])/b + PolyLog[4, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)]/(4*b^2)))/b)/(2*(I*(b + b*c) - b*d)))/3`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6823

```
Int[ArcTanh[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_)^(m_
), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m
+ 1))), x] + (-Simp[I*b*((1 - c - I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)
*(E^(2*I*a + 2*I*b*x))/(1 - c + I*d - (1 - c - I*d)*E^(2*I*a + 2*I*b*x))], x
], x] + Simp[I*b*((1 + c + I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*
I*a + 2*I*b*x))/(1 + c - I*d - (1 + c + I*d)*E^(2*I*a + 2*I*b*x))], x], x]
/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 49.45 (sec) , antiderivative size = 6725, normalized size of antiderivative = 17.20

method	result	size
risch	Expression too large to display	6725

input

```
int(x^2*arctanh(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1799 vs.  $2(275) = 550$ .

Time = 0.27 (sec) , antiderivative size = 1799, normalized size of antiderivative = 4.60

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctanh(c+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/48*(8*b^3*x^3*log(-(d*cos(2*b*x + 2*a) + (c + 1)*sin(2*b*x + 2*a) + d)/(
d*cos(2*b*x + 2*a) + (c - 1)*sin(2*b*x + 2*a) + d)) + 6*I*b^2*x^2*dilog(-(
c^2 + d^2 - (c^2 + 2*I*(c + 1)*d - d^2 + 2*c + 1)*cos(2*b*x + 2*a) + (-I*c
^2 + 2*(c + 1)*d + I*d^2 - 2*I*c - I)*sin(2*b*x + 2*a) + 2*c + 1)/(c^2 + d
^2 + 2*c + 1) + 1) - 6*I*b^2*x^2*dilog(-(c^2 + d^2 - (c^2 - 2*I*(c + 1)*d
- d^2 + 2*c + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*(c + 1)*d - I*d^2 + 2*I*c +
I)*sin(2*b*x + 2*a) + 2*c + 1)/(c^2 + d^2 + 2*c + 1) + 1) - 6*I*b^2*x^2*d
ilog(-(c^2 + d^2 - (c^2 + 2*I*(c - 1)*d - d^2 - 2*c + 1)*cos(2*b*x + 2*a)
+ (-I*c^2 + 2*(c - 1)*d + I*d^2 + 2*I*c - I)*sin(2*b*x + 2*a) - 2*c + 1)/(
c^2 + d^2 - 2*c + 1) + 1) + 6*I*b^2*x^2*dilog(-(c^2 + d^2 - (c^2 - 2*I*(c
- 1)*d - d^2 - 2*c + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*(c - 1)*d - I*d^2 -
2*I*c + I)*sin(2*b*x + 2*a) - 2*c + 1)/(c^2 + d^2 - 2*c + 1) + 1) + 4*a^3*
log(1/2*c^2 + I*(c + 1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 + 2*c + 1)*cos(2*b*x
+ 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) + c + 1/2) - 4*a
^3*log(1/2*c^2 + I*(c - 1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 - 2*c + 1)*cos(2*b
*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) - c + 1/2) +
4*a^3*log(-1/2*c^2 + I*(c + 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 + 2*c + 1)*cos
(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) - c - 1/2
) - 4*a^3*log(-1/2*c^2 + I*(c - 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 - 2*c + 1)
*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) + ...

```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input

```
integrate(x**2*atanh(c+d*cot(b*x+a)), x)
```

output

```
Integral(x**2*atanh(c + d*cot(a + b*x)), x)
```

**Maxima [F]**

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x^2 \operatorname{artanh}(d \cot(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*cot(b*x+a)),x, algorithm="maxima")`

output

```
1/12*x^3*log((c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c + 1)*d*sin(2*
b*x + 2*a) + (c^2 + d^2 + 2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 - 2*(c^2
- d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 2*c + 1) - 1/12*x^3*log((c^2 + d^2 -
2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c - 1)*d*sin(2*b*x + 2*a) + (c^2 + d^2 -
2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 - 2*(c^2 - d^2 - 2*c + 1)*cos(2*b*
x + 2*a) - 2*c + 1) - 4*b*d*integrate(1/3*(2*(c^2 + d^2 - 1)*x^3*cos(2*b*x
+ 2*a)^2 + 2*c*d*x^3*sin(2*b*x + 2*a) + 2*(c^2 + d^2 - 1)*x^3*sin(2*b*x +
2*a)^2 - (c^2 - d^2 - 1)*x^3*cos(2*b*x + 2*a) - (2*c*d*x^3*sin(2*b*x + 2*
a) + (c^2 - d^2 - 1)*x^3*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x^3*c
os(2*b*x + 2*a) - (c^2 - d^2 - 1)*x^3*sin(2*b*x + 2*a))*sin(4*b*x + 4*a))/
(c^4 + d^4 + 2*(c^2 + 1)*d^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*c
os(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*cos(2*b*x
+ 2*a)^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*sin(4*b*x + 4*a)^2 +
4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*sin(2*b*x + 2*a)^2 - 2*c^2 + 2
*(c^4 + d^4 - 2*(3*c^2 - 1)*d^2 - 2*c^2 - 2*(c^4 - d^4 - 2*c^2 + 1)*cos(2*
b*x + 2*a) - 4*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x + 4*a
) - 4*(c^4 - d^4 - 2*c^2 + 1)*cos(2*b*x + 2*a) + 4*(2*c*d^3 - 2*(c^3 - c)*
d + 2*(c*d^3 + (c^3 - c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 - 2*c^2 + 1)*sin
(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a)
+ 1), x)
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x^2 \operatorname{artanh}(d \cot(bx + a) + c) dx$$

input `integrate(x^2*arctanh(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(d*cot(b*x + a) + c), x)`



**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x^2 \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input `int(x^2*atanh(c + d*cot(a + b*x)),x)`output `int(x^2*atanh(c + d*cot(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + c) x^2 dx$$

input `int(x^2*atanh(c+d*cot(b*x+a)),x)`output `int(atanh(cot(a + b*x)*d + c)*x**2,x)`

### 3.335 $\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx$

Optimal result	2357
Mathematica [A] (verified)	2358
Rubi [A] (verified)	2358
Maple [C] (warning: unable to verify)	2362
Fricas [B] (verification not implemented)	2362
Sympy [F]	2363
Maxima [F]	2364
Giac [F]	2364
Mupad [F(-1)]	2365
Reduce [F]	2365

#### Optimal result

Integrand size = 13, antiderivative size = 293

$$\begin{aligned}
 \int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = & \frac{1}{2} x^2 \operatorname{arctanh}(c + d \cot(a + bx)) \\
 & + \frac{1}{4} x^2 \log \left( 1 - \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right) \\
 & - \frac{1}{4} x^2 \log \left( 1 - \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right) \\
 & - \frac{ix \operatorname{PolyLog} \left( 2, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{4b} \\
 & + \frac{ix \operatorname{PolyLog} \left( 2, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{4b} \\
 & + \frac{\operatorname{PolyLog} \left( 3, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{8b^2} \\
 & - \frac{\operatorname{PolyLog} \left( 3, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\operatorname{arctanh}(c+d*\cot(b*x+a))+1/4*x^2*\ln(1-(1-c-I*d)*\exp(2*I*a+2*I*b*x)/ \\ & (1-c+I*d))-1/4*x^2*\ln(1-(1+c+I*d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))-1/4*I*x*po \\ & \operatorname{lylog}(2,(1-c-I*d)*\exp(2*I*a+2*I*b*x)/(1-c+I*d))/b+1/4*I*x*\operatorname{polylog}(2,(1+c+I \\ & *d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))/b+1/8*\operatorname{polylog}(3,(1-c-I*d)*\exp(2*I*a+2*I* \\ & b*x)/(1-c+I*d))/b^2-1/8*\operatorname{polylog}(3,(1+c+I*d)*\exp(2*I*a+2*I*b*x)/(1+c-I*d))/ \\ & b^2 \end{aligned}$$
**Mathematica [A] (verified)**

Time = 2.75 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.87

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx$$

$$= \frac{4b^2x^2 \operatorname{arctanh}(c + d \cot(a + bx)) + 2b^2x^2 \log\left(1 + \frac{(1-c+id)e^{-2i(a+bx)}}{-1+c+id}\right) - 2b^2x^2 \log\left(1 + \frac{(-1-c+id)e^{-2i(a+bx)}}{1+c+id}\right)}{1}$$

input

`Integrate[x*ArcTanh[c + d*Cot[a + b*x]],x]`

output

$$\begin{aligned} & (4*b^2*x^2*ArcTanh[c + d*Cot[a + b*x]] + 2*b^2*x^2*Log[1 + (1 - c + I*d)/ \\ & (-1 + c + I*d)*E^((2*I)*(a + b*x))]) - 2*b^2*x^2*Log[1 + (-1 - c + I*d)/(( \\ & 1 + c + I*d)*E^((2*I)*(a + b*x))]) + (2*I)*b*x*PolyLog[2, (-1 + c - I*d)/ \\ & (-1 + c + I*d)*E^((2*I)*(a + b*x))]) - (2*I)*b*x*PolyLog[2, (1 + c - I*d)/ \\ & ((1 + c + I*d)*E^((2*I)*(a + b*x)))] + PolyLog[3, (-1 + c - I*d)/((-1 + c \\ & + I*d)*E^((2*I)*(a + b*x)))] - PolyLog[3, (1 + c - I*d)/((1 + c + I*d)*E^(( \\ & (2*I)*(a + b*x)))])/(8*b^2) \end{aligned}$$
**Rubi [A] (verified)**Time = 1.10 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.34, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {6823, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x \operatorname{arctanh}(d \cot(a + bx) + c) dx \\
& \quad \downarrow \text{6823} \\
& -\frac{1}{2}b(-ic + d + i) \int \frac{e^{2ia+2ibx} x^2}{-c - (-c - id + 1)e^{2ia+2ibx} + id + 1} dx + \frac{1}{2}b(-d + i(c + \\
& \quad 1)) \int \frac{e^{2ia+2ibx} x^2}{c - (c + id + 1)e^{2ia+2ibx} - id + 1} dx + \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& -\frac{1}{2}b(-ic + d + i) \left( \frac{\int x \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) dx}{b(d+i(1-c))} - \frac{x^2 \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d+i(1-c))} \right) + \\
& \frac{1}{2}b(-d + i(c + 1)) \left( \frac{\int x \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) dx}{-bd + i(bc + b)} - \frac{x^2 \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd + i(bc + b))} \right) + \\
& \quad \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{3011} \\
& i) \left( \frac{-\frac{1}{2}b(-ic + d + \frac{ix \operatorname{PolyLog} \left( 2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) dx}{2b}}{b(d+i(1-c))} - \frac{x^2 \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d+i(1-c))} \right) + \\
& \quad \frac{1}{2}b(-d + i(c + \frac{ix \operatorname{PolyLog} \left( 2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left( 2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) dx}{2b}}{-bd + i(bc + b)} - \frac{x^2 \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd + i(bc + b))} \right) + \\
& \quad \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
 & i) \left( \frac{\frac{ix \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right) de^{2ia+2ibx}}{4b^2}}{b(d+i(1-c))} - \frac{x^2 \log\left(1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b(d+i(1-c))} \right. \\
 & 1)) \left( \frac{\frac{ix \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right) de^{2ia+2ibx}}{4b^2}}{-bd+i(bc+b)} - \frac{x^2 \log\left(1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2(-bd+i(bc+b))} \right. \\
 & \left. \left. \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a+bx) + c) \right)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & i) \left( \frac{\frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a+bx) + c) - \frac{1}{2}b(-ic+d + \frac{ix \operatorname{PolyLog}\left(2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{4b^2}}{b(d+i(1-c))} - \frac{x^2 \log\left(1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1}\right)}{2b(d+i(1-c))}}{\right) + \\
 & 1)) \left( \frac{\frac{1}{2}b(-d+i(c + \frac{ix \operatorname{PolyLog}\left(2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{4b^2}}{-bd+i(bc+b)} - \frac{x^2 \log\left(1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1}\right)}{2(-bd+i(bc+b))}}{\right)
 \end{aligned}$$

input `Int[x*ArcTanh[c + d*Cot[a + b*x]],x]`

output `(x^2*ArcTanh[c + d*Cot[a + b*x]])/2 - (b*(I - I*c + d)*(-1/2*(x^2*Log[1 - ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)])/(b*(I*(1 - c) + d)) + (((I/2)*x*PolyLog[2, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)]/b - PolyLog[3, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)]/(4*b^2))/(b*(I*(1 - c) + d)))/2 + (b*(I*(1 + c) - d)*(-1/2*(x^2*Log[1 - ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)])/(I*(b + b*c) - b*d) + (((I/2)*x*PolyLog[2, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)]/b - PolyLog[3, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)]/(4*b^2))/(I*(b + b*c) - b*d))/2`

## Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 6823

```
Int[ArcTanh[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m
+ 1))), x] + (-Simp[I*b*((1 - c - I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)
*(E^(2*I*a + 2*I*b*x))/(1 - c + I*d - (1 - c - I*d)*E^(2*I*a + 2*I*b*x))], x
], x] + Simp[I*b*((1 + c + I*d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*
I*a + 2*I*b*x))/(1 + c - I*d - (1 + c + I*d)*E^(2*I*a + 2*I*b*x))], x], x])
/; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - I*d)^2, 1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 6.41 (sec) , antiderivative size = 6375, normalized size of antiderivative = 21.76

method	result	size
risch	Expression too large to display	6375

input `int(x*arctanh(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1463 vs.  $2(207) = 414$ .

Time = 0.25 (sec) , antiderivative size = 1463, normalized size of antiderivative = 4.99

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctanh(c+d*cot(b*x+a)),x, algorithm="fricas")`

output

```

1/16*(4*b^2*x^2*log(-(d*cos(2*b*x + 2*a) + (c + 1)*sin(2*b*x + 2*a) + d)/(
d*cos(2*b*x + 2*a) + (c - 1)*sin(2*b*x + 2*a) + d)) + 2*I*b*x*dilog(-(c^2
+ d^2 - (c^2 + 2*I*(c + 1)*d - d^2 + 2*c + 1)*cos(2*b*x + 2*a) + (-I*c^2 +
2*(c + 1)*d + I*d^2 - 2*I*c - I)*sin(2*b*x + 2*a) + 2*c + 1)/(c^2 + d^2 +
2*c + 1) + 1) - 2*I*b*x*dilog(-(c^2 + d^2 - (c^2 - 2*I*(c + 1)*d - d^2 +
2*c + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*(c + 1)*d - I*d^2 + 2*I*c + I)*sin(
2*b*x + 2*a) + 2*c + 1)/(c^2 + d^2 + 2*c + 1) + 1) - 2*I*b*x*dilog(-(c^2 +
d^2 - (c^2 + 2*I*(c - 1)*d - d^2 - 2*c + 1)*cos(2*b*x + 2*a) + (-I*c^2 +
2*(c - 1)*d + I*d^2 + 2*I*c - I)*sin(2*b*x + 2*a) - 2*c + 1)/(c^2 + d^2 -
2*c + 1) + 1) + 2*I*b*x*dilog(-(c^2 + d^2 - (c^2 - 2*I*(c - 1)*d - d^2 - 2
*c + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*(c - 1)*d - I*d^2 - 2*I*c + I)*sin(2
*b*x + 2*a) - 2*c + 1)/(c^2 + d^2 - 2*c + 1) + 1) - 2*a^2*log(1/2*c^2 + I*
(c + 1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 1/2*(I*
c^2 + I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) + c + 1/2) + 2*a^2*log(1/2*c^2 +
I*(c - 1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 - 2*c + 1)*cos(2*b*x + 2*a) + 1/2*
(I*c^2 + I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) - c + 1/2) - 2*a^2*log(-1/2*c
^2 + I*(c + 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a) +
1/2*(I*c^2 + I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) - c - 1/2) + 2*a^2*log(-1
/2*c^2 + I*(c - 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 - 2*c + 1)*cos(2*b*x + 2*a
) + 1/2*(I*c^2 + I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) + c - 1/2) - 2*(b^...

```

SymPy [F]

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input

```
integrate(x*atanh(c+d*cot(b*x+a)), x)
```

output

```
Integral(x*atanh(c + d*cot(a + b*x)), x)
```



**Maxima [F]**

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x \operatorname{artanh}(d \cot(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*cot(b*x+a)),x, algorithm="maxima")`

output

```
-2*b*d*integrate((2*(c^2 + d^2 - 1)*x^2*cos(2*b*x + 2*a)^2 + 2*c*d*x^2*sin(2*b*x + 2*a) + 2*(c^2 + d^2 - 1)*x^2*sin(2*b*x + 2*a)^2 - (c^2 - d^2 - 1)*x^2*cos(2*b*x + 2*a) - (2*c*d*x^2*sin(2*b*x + 2*a) + (c^2 - d^2 - 1)*x^2*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x^2*cos(2*b*x + 2*a) - (c^2 - d^2 - 1)*x^2*sin(2*b*x + 2*a))*sin(4*b*x + 4*a))/(c^4 + d^4 + 2*(c^2 + 1)*d^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*cos(2*b*x + 2*a)^2 + (c^4 + d^4 + 2*(c^2 + 1)*d^2 - 2*c^2 + 1)*sin(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 - 1)*d^2 - 2*c^2 + 1)*sin(2*b*x + 2*a)^2 - 2*c^2 + 2*(c^4 + d^4 - 2*(3*c^2 - 1)*d^2 - 2*c^2 - 2*(c^4 - d^4 - 2*c^2 + 1)*cos(2*b*x + 2*a) - 4*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x + 4*a) - 4*(c^4 - d^4 - 2*c^2 + 1)*cos(2*b*x + 2*a) + 4*(2*c*d^3 - 2*(c^3 - c)*d + 2*(c*d^3 + (c^3 - c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 - 2*c^2 + 1)*sin(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 - c)*d)*sin(2*b*x + 2*a) + 1), x) + 1/8*x^2*log((c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c + 1)*d*sin(2*b*x + 2*a) + (c^2 + d^2 + 2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 - 2*(c^2 - d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 2*c + 1) - 1/8*x^2*log((c^2 + d^2 - 2*c + 1)*cos(2*b*x + 2*a)^2 + 4*(c - 1)*d*sin(2*b*x + 2*a) + (c^2 + d^2 - 2*c + 1)*sin(2*b*x + 2*a)^2 + c^2 + d^2 - 2*(c^2 - d^2 - 2*c + 1)*cos(2*b*x + 2*a) - 2*c + 1)
```

**Giac [F]**

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x \operatorname{artanh}(d \cot(bx + a) + c) dx$$

input `integrate(x*arctanh(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*cot(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int x \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input `int(x*atanh(c + d*cot(a + b*x)),x)`output `int(x*atanh(c + d*cot(a + b*x)), x)`**Reduce [F]**

$$\int x \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + c) x dx$$

input `int(x*atanh(c+d*cot(b*x+a)),x)`output `int(atanh(cot(a + b*x)*d + c)*x,x)`

### 3.336 $\int \operatorname{arctanh}(c + d \cot(a + bx)) dx$

Optimal result	2366
Mathematica [B] (warning: unable to verify)	2367
Rubi [A] (verified)	2367
Maple [B] (verified)	2370
Fricas [B] (verification not implemented)	2371
Sympy [F]	2372
Maxima [B] (verification not implemented)	2372
Giac [F]	2373
Mupad [F(-1)]	2373
Reduce [F]	2373

#### Optimal result

Integrand size = 11, antiderivative size = 194

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = x \operatorname{arctanh}(c + d \cot(a + bx)) + \frac{1}{2} x \log \left( 1 - \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right) - \frac{1}{2} x \log \left( 1 - \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right) - \frac{i \operatorname{PolyLog} \left( 2, \frac{(1 - c - id)e^{2ia+2ibx}}{1 - c + id} \right)}{4b} + \frac{i \operatorname{PolyLog} \left( 2, \frac{(1 + c + id)e^{2ia+2ibx}}{1 + c - id} \right)}{4b}$$

output

```
x*arctanh(c+d*cot(b*x+a))+1/2*x*ln(1-(1-c-I*d)*exp(2*I*a+2*I*b*x)/(1-c+I*d))
)-1/2*x*ln(1-(1+c+I*d)*exp(2*I*a+2*I*b*x)/(1+c-I*d))-1/4*I*polylog(2,(1-c-I*d)*exp(2*I*a+2*I*b*x)/(1-c+I*d))/b+1/4*I*polylog(2,(1+c+I*d)*exp(2*I*a+2*I*b*x)/(1+c-I*d))/b
```

**Mathematica [B] (warning: unable to verify)**

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 390 vs.  $2(194) = 388$ .

Time = 3.75 (sec) , antiderivative size = 390, normalized size of antiderivative = 2.01

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = x \left( \operatorname{arctanh}(c + d \cot(a + bx)) \right. \\ \left. + \frac{2a \log(d + (-1 + c) \tan(a + bx)) + i \log(1 + i \tan(a + bx)) \log\left(-\frac{i(d + (-1 + c) \tan(a + bx))}{-1 + c - id}\right) - i \log(1 - i \tan(a + bx))}{4a - (2 + i) \log(1 - i \tan(a + bx)) + (2 + i) \log(1 + i \tan(a + bx))} \right)$$

input `Integrate[ArcTanh[c + d*Cot[a + b*x]],x]`

output

```
x*(ArcTanh[c + d*Cot[a + b*x]] + (2*a*Log[d + (-1 + c)*Tan[a + b*x]] + I*Log[1 + I*Tan[a + b*x]]*Log[(-I)*(d + (-1 + c)*Tan[a + b*x])/(-1 + c - I*d)] - I*Log[1 - I*Tan[a + b*x]]*Log[(I*(d + (-1 + c)*Tan[a + b*x])/(-1 + c + I*d)] - 2*a*Log[d + (1 + c)*Tan[a + b*x]] + I*Log[1 - I*Tan[a + b*x]]*Log[(I*(d + (1 + c)*Tan[a + b*x])/(1 + c + I*d)] - I*Log[1 + I*Tan[a + b*x]]*Log[(d + (1 + c)*Tan[a + b*x])/(I*(1 + c) + d)] - I*PolyLog[2, ((-1 + c)*(1 - I*Tan[a + b*x])/(1 + c + I*d)] + I*PolyLog[2, ((1 + c)*(1 - I*Tan[a + b*x])/(1 + c + I*d)] + I*PolyLog[2, ((-1 + c)*(1 + I*Tan[a + b*x])/(1 + c - I*d)] - I*PolyLog[2, ((1 + c)*(1 + I*Tan[a + b*x])/(1 + c - I*d)]))/(4*a - (2*I)*Log[1 - I*Tan[a + b*x]] + (2*I)*Log[1 + I*Tan[a + b*x]]])
```

**Rubi [A] (verified)**

Time = 1.09 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.46, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$ , Rules used = {6815, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \operatorname{arctanh}(d \cot(a + bx) + c) dx \\
& \quad \downarrow \text{6815} \\
& -b(-ic + d + i) \int \frac{e^{2ia+2ibx} x}{-c - (-c - id + 1)e^{2ia+2ibx} + id + 1} dx + b(-d + i(c + \\
& 1)) \int \frac{e^{2ia+2ibx} x}{c - (c + id + 1)e^{2ia+2ibx} - id + 1} dx + x \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& -b(-ic + d + i) \left( \frac{\int \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) dx}{2b(d+i(1-c))} - \frac{x \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d+i(1-c))} \right) + b(-d + \\
& i(c+1)) \left( \frac{\int \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) dx}{2(-bd+i(bc+b))} - \frac{x \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd+i(bc+b))} \right) + \\
& \quad x \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{2715} \\
& i) \left( -\frac{-b(-ic + d + i) \int e^{-2ia-2ibx} \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) de^{2ia+2ibx}}{4b^2(d+i(1-c))} - \frac{x \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d+i(1-c))} \right) + \\
& 1)) \left( -\frac{b(-d + i(c + 1)) \int e^{-2ia-2ibx} \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) de^{2ia+2ibx}}{4b(-bd+i(bc+b))} - \frac{x \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd+i(bc+b))} \right) + \\
& \quad x \operatorname{arctanh}(d \cot(a + bx) + c) \\
& \quad \downarrow \text{2838} \\
& i) \left( \frac{x \operatorname{arctanh}(d \cot(a + bx) + c) - b(-ic + d + i) \int \operatorname{PolyLog} \left( 2, \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right) dx}{4b^2(d+i(1-c))} - \frac{x \log \left( 1 - \frac{(-c-id+1)e^{2ia+2ibx}}{-c+id+1} \right)}{2b(d+i(1-c))} \right) + b(-d + i(c + \\
& 1)) \left( \frac{i \operatorname{PolyLog} \left( 2, \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right) dx}{4b(-bd+i(bc+b))} - \frac{x \log \left( 1 - \frac{(c+id+1)e^{2ia+2ibx}}{c-id+1} \right)}{2(-bd+i(bc+b))} \right)
\end{aligned}$$

input `Int[ArcTanh[c + d*Cot[a + b*x]], x]`

output

```
x*ArcTanh[c + d*Cot[a + b*x]] - b*(I - I*c + d)*(-1/2*(x*Log[1 - ((1 - c -
I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)]/(b*(I*(1 - c) + d)) + ((I/4
)*PolyLog[2, ((1 - c - I*d)*E^((2*I)*a + (2*I)*b*x))/(1 - c + I*d)]/(b^2*
(I*(1 - c) + d))) + b*(I*(1 + c) - d)*(-1/2*(x*Log[1 - ((1 + c + I*d)*E^((
2*I)*a + (2*I)*b*x))/(1 + c - I*d)]/(I*(b + b*c) - b*d)) + ((I/4)*PolyLog[
2, ((1 + c + I*d)*E^((2*I)*a + (2*I)*b*x))/(1 + c - I*d)]/(b*(I*(b + b*c)
- b*d)))
```

### Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 6815

```
Int[ArcTanh[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)], x_Symbol] := Simp[x*Arc
Tanh[c + d*Cot[a + b*x]], x] + (-Simp[I*b*(1 - c - I*d) Int[x*(E^(2*I*a +
2*I*b*x))/(1 - c + I*d - (1 - c - I*d)*E^(2*I*a + 2*I*b*x))], x], x] + Simp
[I*b*(1 + c + I*d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 + c - I*d - (1 + c + I*d
)*E^(2*I*a + 2*I*b*x))], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[(c - I*d)
^2, 1]
```

### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 562 vs.  $2(164) = 328$ .

Time = 4.05 (sec) , antiderivative size = 563, normalized size of antiderivative = 2.90

method	result
derivativedivides	$-d\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(bx+a))\right) \operatorname{arctanh}(c+d \cot(bx+a))+d^2 \frac{\operatorname{arctan}\left(-\frac{c+d \cot(bx+a)}{d}+\frac{c}{d}\right) \ln\left(d\left(\frac{c+d \cot(bx+a)}{d}-\frac{c}{d}\right)+c-1\right)}{2d}$
default	$-d\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(bx+a))\right) \operatorname{arctanh}(c+d \cot(bx+a))+d^2 \frac{\operatorname{arctan}\left(-\frac{c+d \cot(bx+a)}{d}+\frac{c}{d}\right) \ln\left(d\left(\frac{c+d \cot(bx+a)}{d}-\frac{c}{d}\right)+c-1\right)}{2d}$
risch	Expression too large to display

input `int(arctanh(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output `1/b/d*(-d*(1/2*Pi-arccot(cot(b*x+a)))*arctanh(c+d*cot(b*x+a))+d^2*(1/2*arctan(-(c+d*cot(b*x+a))/d+c/d)/d*ln(d*((c+d*cot(b*x+a))/d-c/d)+c-1)-1/2*arctan(-(c+d*cot(b*x+a))/d+c/d)/d*ln(d*((c+d*cot(b*x+a))/d-c/d)+c+1)+1/4*I*ln(d*((c+d*cot(b*x+a))/d-c/d)+c+1)*(ln((I*d-d*((c+d*cot(b*x+a))/d-c/d))/(1+c+I*d))-ln((I*d+d*((c+d*cot(b*x+a))/d-c/d))/(I*d-c-1)))/d+1/4*I*(dilog((I*d-d*((c+d*cot(b*x+a))/d-c/d))/(1+c+I*d))-dilog((I*d+d*((c+d*cot(b*x+a))/d-c/d))/(I*d-c-1)))/d-1/4*I*ln(d*((c+d*cot(b*x+a))/d-c/d)+c-1)*(ln((I*d-d*((c+d*cot(b*x+a))/d-c/d))/(I*d+c-1))-ln((I*d+d*((c+d*cot(b*x+a))/d-c/d))/(1-c+I*d)))/d-1/4*I*(dilog((I*d-d*((c+d*cot(b*x+a))/d-c/d))/(I*d+c-1))-dilog((I*d+d*((c+d*cot(b*x+a))/d-c/d))/(1-c+I*d)))/d)`

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1099 vs.  $2(136) = 272$ .

Time = 0.22 (sec) , antiderivative size = 1099, normalized size of antiderivative = 5.66

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctanh(c+d*cot(b*x+a)),x, algorithm="fricas")`

output

```
1/8*(4*b*x*log(-(d*cos(2*b*x + 2*a) + (c + 1)*sin(2*b*x + 2*a) + d)/(d*cos
(2*b*x + 2*a) + (c - 1)*sin(2*b*x + 2*a) + d)) + 2*a*log(1/2*c^2 + I*(c +
1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 +
I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) + c + 1/2) - 2*a*log(1/2*c^2 + I*(c -
1)*d - 1/2*d^2 - 1/2*(c^2 + d^2 - 2*c + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2
+ I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) - c + 1/2) + 2*a*log(-1/2*c^2 + I*(c
+ 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 + 2*c + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^
2 + I*d^2 + 2*I*c + I)*sin(2*b*x + 2*a) - c - 1/2) - 2*a*log(-1/2*c^2 + I*
(c - 1)*d + 1/2*d^2 + 1/2*(c^2 + d^2 - 2*c + 1)*cos(2*b*x + 2*a) + 1/2*(I*
c^2 + I*d^2 - 2*I*c + I)*sin(2*b*x + 2*a) + c - 1/2) - 2*(b*x + a)*log((c^
2 + d^2 - (c^2 + 2*I*(c + 1)*d - d^2 + 2*c + 1)*cos(2*b*x + 2*a) + (-I*c^2
+ 2*(c + 1)*d + I*d^2 - 2*I*c - I)*sin(2*b*x + 2*a) + 2*c + 1)/(c^2 + d^2
+ 2*c + 1)) - 2*(b*x + a)*log((c^2 + d^2 - (c^2 - 2*I*(c + 1)*d - d^2 + 2
*c + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*(c + 1)*d - I*d^2 + 2*I*c + I)*sin(2
*b*x + 2*a) + 2*c + 1)/(c^2 + d^2 + 2*c + 1)) + 2*(b*x + a)*log((c^2 + d^2
- (c^2 + 2*I*(c - 1)*d - d^2 - 2*c + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*(c
- 1)*d + I*d^2 + 2*I*c - I)*sin(2*b*x + 2*a) - 2*c + 1)/(c^2 + d^2 - 2*c
+ 1)) + 2*(b*x + a)*log((c^2 + d^2 - (c^2 - 2*I*(c - 1)*d - d^2 - 2*c + 1)
*cos(2*b*x + 2*a) + (I*c^2 + 2*(c - 1)*d - I*d^2 - 2*I*c + I)*sin(2*b*x +
2*a) - 2*c + 1)/(c^2 + d^2 - 2*c + 1)) + I*dilog(-(c^2 + d^2 - (c^2 + 2...
```



**Sympy [F]**

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input `integrate(atanh(c+d*cot(b*x+a)),x)`

output `Integral(atanh(c + d*cot(a + b*x)), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 392 vs.  $2(136) = 272$ .

Time = 0.19 (sec) , antiderivative size = 392, normalized size of antiderivative = 2.02

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx$$

$$= \frac{4(bx + a) \operatorname{artanh}\left(c + \frac{d}{\tan(bx+a)}\right) + \left(\arctan\left(\frac{(c+1)d + (c^2+2c+1)\tan(bx+a)}{c^2+d^2+2c+1}\right), \frac{(c+1)d \tan(bx+a) + d^2}{c^2+d^2+2c+1}\right) - \arctan\left(\frac{(c-1)d + (c^2-2c+1)\tan(bx+a)}{c^2+d^2-2c+1}\right) - \arctan\left(\frac{(c-1)d \tan(bx+a) + d^2}{c^2+d^2-2c+1}\right)}{b}$$

input `integrate(arctanh(c+d*cot(b*x+a)),x, algorithm="maxima")`

output `1/4*(4*(b*x + a)*arctanh(c + d/tan(b*x + a)) + (arctan2(((c + 1)*d + (c^2 + 2*c + 1)*tan(b*x + a))/(c^2 + d^2 + 2*c + 1), ((c + 1)*d*tan(b*x + a) + d^2)/(c^2 + d^2 + 2*c + 1)) - arctan2(((c - 1)*d + (c^2 - 2*c + 1)*tan(b*x + a))/(c^2 + d^2 - 2*c + 1), ((c - 1)*d*tan(b*x + a) + d^2)/(c^2 + d^2 - 2*c + 1)))*log(tan(b*x + a)^2 + 1) - (b*x + a)*log((2*(c + 1)*d*tan(b*x + a) + (c^2 + 2*c + 1)*tan(b*x + a)^2 + d^2)/(c^2 + d^2 + 2*c + 1)) + (b*x + a)*log((2*(c - 1)*d*tan(b*x + a) + (c^2 - 2*c + 1)*tan(b*x + a)^2 + d^2)/(c^2 + d^2 - 2*c + 1)) + I*dilog(-((c + 1)*tan(b*x + a) - I*c - I)/(I*c + d + I)) - I*dilog(-((c - 1)*tan(b*x + a) - I*c + I)/(I*c + d - I)) + I*dilog(-((c - 1)*tan(b*x + a) + I*c - I)/(-I*c + d + I)) - I*dilog(-((c + 1)*tan(b*x + a) + I*c + I)/(-I*c + d - I)))/b`

**Giac [F]**

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{artanh}(d \cot(bx + a) + c) dx$$

input `integrate(arctanh(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*cot(b*x + a) + c), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{atanh}(c + d \cot(a + bx)) dx$$

input `int(atanh(c + d*cot(a + b*x)),x)`

output `int(atanh(c + d*cot(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(c + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + c) dx$$

input `int(atanh(c+d*cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)*d + c),x)`

### 3.337 $\int \frac{\operatorname{arctanh}(c+d \cot(a+bx))}{x} dx$

Optimal result	2374
Mathematica [N/A]	2374
Rubi [N/A]	2375
Maple [N/A]	2375
Fricas [N/A]	2376
Sympy [N/A]	2376
Maxima [N/A]	2376
Giac [N/A]	2377
Mupad [N/A]	2377
Reduce [N/A]	2378

#### Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x}, x\right)$$

output

```
Defer(Int)(arctanh(c+d*cot(b*x+a))/x,x)
```

#### Mathematica [N/A]

Not integrable

Time = 1.88 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx$$

input

```
Integrate[ArcTanh[c + d*Cot[a + b*x]]/x,x]
```

output

```
Integrate[ArcTanh[c + d*Cot[a + b*x]]/x, x]
```

**Rubi [N/A]**

Not integrable

Time = 0.47 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \cot(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \cot(a + bx) + c)}{x} dx$$

input `Int[ArcTanh[c + d*Cot[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.35 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(c + d \cot(bx + a))}{x} dx$$

input `int(arctanh(c+d*cot(b*x+a))/x,x)`

output `int(arctanh(c+d*cot(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctanh(d*cot(b*x + a) + c)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 1.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \cot(a + bx))}{x} dx$$

input `integrate(atanh(c+d*cot(b*x+a))/x,x)`

output `Integral(atanh(c + d*cot(a + b*x))/x, x)`

**Maxima [N/A]**

Not integrable

Time = 3.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*cot(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctanh(d*cot(b*x + a) + c)/x, x)`

### Giac [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + c)}{x} dx$$

input `integrate(arctanh(c+d*cot(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctanh(d*cot(b*x + a) + c)/x, x)`

### Mupad [N/A]

Not integrable

Time = 4.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(c + d \cot(a + bx))}{x} dx$$

input `int(atanh(c + d*cot(a + b*x))/x,x)`

output `int(atanh(c + d*cot(a + b*x))/x, x)`

**Reduce [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\operatorname{arctanh}(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\cot(bx + a) d + c)}{x} dx$$

input `int(atanh(c+d*cot(b*x+a))/x,x)`output `int(atanh(cot(a + b*x)*d + c)/x,x)`

### 3.338 $\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$

Optimal result	2379
Mathematica [A] (verified)	2380
Rubi [A] (verified)	2380
Maple [C] (warning: unable to verify)	2383
Fricas [A] (verification not implemented)	2384
Sympy [F]	2385
Maxima [B] (verification not implemented)	2385
Giac [F]	2386
Mupad [F(-1)]	2386
Reduce [F]	2386

#### Optimal result

Integrand size = 20, antiderivative size = 168

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(1 + id + d \cot(a + bx)) - \frac{1}{6} x^3 \log(1 - (1 + id)e^{2ia+2ibx}) + \frac{ix^2 \operatorname{PolyLog}(2, (1 + id)e^{2ia+2ibx})}{4b} - \frac{x \operatorname{PolyLog}(3, (1 + id)e^{2ia+2ibx})}{4b^2} - \frac{i \operatorname{PolyLog}(4, (1 + id)e^{2ia+2ibx})}{8b^3}$$

output

```
1/12*I*b*x^4+1/3*x^3*arctanh(1+I*d+d*cot(b*x+a))-1/6*x^3*ln(1-(1+I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x^2*polylog(2,(1+I*d)*exp(2*I*a+2*I*b*x))/b-1/4*x*polylog(3,(1+I*d)*exp(2*I*a+2*I*b*x))/b^2-1/8*I*polylog(4,(1+I*d)*exp(2*I*a+2*I*b*x))/b^3
```



**Mathematica [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.92

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \frac{1}{3} x^3 \operatorname{arctanh}(1 + id + d \cot(a + bx)) - \frac{4b^3 x^3 \log\left(1 + \frac{ie^{-2i(a+bx)}}{-i+d}\right) + 6ib^2 x^2 \operatorname{PolyLog}\left(2, -\frac{ie^{-2i(a+bx)}}{-i+d}\right) + 6bx \operatorname{PolyLog}\left(3, -\frac{ie^{-2i(a+bx)}}{-i+d}\right) - 3i \operatorname{PolyLog}\left(4, -\frac{ie^{-2i(a+bx)}}{-i+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 + I*d + d*Cot[a + b*x]],x]
```

output

```
(x^3*ArcTanh[1 + I*d + d*Cot[a + b*x]])/3 - (4*b^3*x^3*Log[1 + I/((-I + d)*E^((2*I)*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, (-I)/((-I + d)*E^((2*I)*(a + b*x)))] + 6*b*x*PolyLog[3, (-I)/((-I + d)*E^((2*I)*(a + b*x)))] - (3*I)*PolyLog[4, (-I)/((-I + d)*E^((2*I)*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**Time = 1.32 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.30, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {6819, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1) dx$$

$$\downarrow \text{6819}$$

$$\frac{1}{3} ib \int \frac{x^3}{1 - (id + 1)e^{2ia + 2ibx}} dx + \frac{1}{3} x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3} ib \left( \frac{x^4}{4} + (1 + id) \int \frac{e^{2ia + 2ibx} x^3}{1 - (id + 1)e^{2ia + 2ibx}} dx \right) + \frac{1}{3} x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 + id) \left( \frac{3 \int x^2 \log(1 - (id + 1)e^{2ia+2ibx}) dx}{2b(-d + i)} - \frac{x^3 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 3011

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 + id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx}) dx}{b} \right)}{2b(-d + i)} - \frac{x^3 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 7163

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 + id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{i \left( \frac{\int \operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx}) dx}{2b} - \frac{i x \operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx})}{2b} \right)}{b} \right)}{2b(-d + i)} - \frac{x^3 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 2720

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 + id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{i x \operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx})}{2b} \right)}{b} \right)}{2b(-d + i)} - \frac{x^3 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 7143

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 + id) \frac{\frac{1}{3}x^3 \operatorname{arctanh}(d \cot(a + bx) + id + 1) + 3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx}}{2b} - \frac{i \left( \frac{\operatorname{PolyLog}(4, (id+1)e^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx}}{2b} \right)}{b} \right)}{2b(-d+i)} \right)}{2b(-d+i)} - x^3 \right)$$

input `Int[x^2*ArcTanh[1 + I*d + d*Cot[a + b*x]],x]`

output `(x^3*ArcTanh[1 + I*d + d*Cot[a + b*x])/3 + (I/3)*b*(x^4/4 + (1 + I*d)*(-1/2*(x^3*Log[1 - (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/(b*(I - d)) + (3*(((I/2)*x^2*PolyLog[2, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/b - (I*(((1/2*I)*x*PolyLog[3, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/b + PolyLog[4, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b)/(2*b*(I - d)))`

### Defintions of rubi rules used

rule 2615 `Int[(((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6819 `Int[ArcTanh[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.77 (sec) , antiderivative size = 2387, normalized size of antiderivative = 14.21

method	result	size
risch	Expression too large to display	2387

input `int(x^2*arctanh(1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

1/8/b^3/(I-d)*polylog(4,-I*exp(2*I*(b*x+a))*(I-d))+1/12*I*b*x^4-1/2/b^2*d/
(I-d)*ln(1+I*exp(2*I*(b*x+a))*(I-d))*a^2*x+1/2/b^2*d*a^2/(I-d)*ln(1-I*exp(
I*(b*x+a))*(I*(I-d))^(1/2))*x+1/2/b^2*d*a^2/(I-d)*ln(1+I*exp(I*(b*x+a))*(I
*(I-d))^(1/2))*x+1/2*I/b^2/(I-d)*ln(1+I*exp(2*I*(b*x+a))*(I-d))*a^2*x+1/6*
x^3*ln(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))+I)-1/3*x^3*ln(exp(I*(b*x+a)))
-1/12*(-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*I*(b*x+a)))^3-2*I*Pi+2*ln(
d)-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*I
*(b*x+a))+I)/(-1+exp(2*I*(b*x+a))))^2-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*ex
p(2*I*(b*x+a))*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*I*(b*x+a))*d*csgn(I*d)
-I*Pi*csgn(I*exp(2*I*(b*x+a))*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I/(-1+ex
p(2*I*(b*x+a))*exp(2*I*(b*x+a)))-I*Pi*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*
I*(b*x+a))+I)/(-1+exp(2*I*(b*x+a))))*csgn((exp(2*I*(b*x+a))*d-I*exp(2*I*(b
*x+a))+I)/(-1+exp(2*I*(b*x+a))))^2-I*Pi*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2
*I*(b*x+a))+I))*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))+I)/(-1+exp(2
*I*(b*x+a))))^2+I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*I*(b*x+a))*d*csgn
(1/(-1+exp(2*I*(b*x+a))*exp(2*I*(b*x+a))*d)^2-I*Pi*csgn(I*exp(I*(b*x+a)))
^2*csgn(I*exp(2*I*(b*x+a)))+I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I*(exp
(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))+I))*csgn(I*(exp(2*I*(b*x+a))*d-I*exp(2*
I*(b*x+a))+I)/(-1+exp(2*I*(b*x+a))))+I*Pi*csgn(1/(-1+exp(2*I*(b*x+a))))*exp
(2*I*(b*x+a))*d)^2+I*Pi*csgn((exp(2*I*(b*x+a))*d-I*exp(2*I*(b*x+a))+I)/...

```

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.07

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$$

$$= \frac{2i b^4 x^4 + 4 b^3 x^3 \log\left(-\frac{((d-i)e^{(2i bx+2i a)}+i)e^{(-2i bx-2i a)}}{d}\right) + 6i b^2 x^2 \operatorname{Li}_2(-(-id-1)e^{(2i bx+2i a)}) - 2i a^4 + 4 a^3 \log\left(\frac{((d-i)e^{(2i bx+2i a)}+i)e^{(-2i bx-2i a)}}{d}\right)}{d}$$

input

```
integrate(x^2*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/24*(2*I*b^4*x^4 + 4*b^3*x^3*log(-((d - I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2
*I*b*x - 2*I*a)/d) + 6*I*b^2*x^2*dilog(-(-I*d - 1)*e^(2*I*b*x + 2*I*a)) -
2*I*a^4 + 4*a^3*log(((d - I)*e^(2*I*b*x + 2*I*a) + I)/(d - I)) - 6*b*x*pol
ylog(3, (I*d + 1)*e^(2*I*b*x + 2*I*a)) - 4*(b^3*x^3 + a^3)*log((-I*d - 1)*
e^(2*I*b*x + 2*I*a) + 1) - 3*I*polylog(4, (I*d + 1)*e^(2*I*b*x + 2*I*a)))/
b^3

```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x^2 \operatorname{atanh}(d \cot(a + bx) + id + 1) dx$$

input `integrate(x**2*atanh(1+I*d+d*cot(b*x+a)),x)`

output `Integral(x**2*atanh(d*cot(a + b*x) + I*d + 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 344 vs.  $2(119) = 238$ .

Time = 0.06 (sec) , antiderivative size = 344, normalized size of antiderivative = 2.05

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$$

$$= \frac{12 \left( (bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2 \right) \operatorname{artanh}(d \cot(bx+a) + id + 1)}{b^2} - \frac{-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(4i(bx+a)^3 - 9i(bx+a)a^2)}{b^2}$$

input `integrate(x^2*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output `1/36*(12*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctanh(d*cot(b*x + a) + I*d + 1)/b^2 - (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 - 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*I*(b*x + a)*a^2)*arctan2(d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), d*sin(2*b*x + 2*a) - cos(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog((I*d + 1)*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^2 + 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3, (I*d + 1)*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, (I*d + 1)*e^(2*I*b*x + 2*I*a)))/b^2)/b`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x^2 \operatorname{artanh}(d \cot(bx + a) + id + 1) dx$$

input `integrate(x^2*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(d*cot(b*x + a) + I*d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x^2 \operatorname{atanh}(d \cot(a + bx) + 1 + d li) dx$$

input `int(x^2*atanh(d*I*i + d*cot(a + b*x) + 1),x)`

output `int(x^2*atanh(d*I*i + d*cot(a + b*x) + 1), x)`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + di + 1) x^2 dx$$

input `int(x^2*atanh(1+I*d+d*cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)*d + d*i + 1)*x**2,x)`

### 3.339 $\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$

Optimal result	2387
Mathematica [A] (verified)	2388
Rubi [A] (verified)	2388
Maple [C] (warning: unable to verify)	2391
Fricas [A] (verification not implemented)	2392
Sympy [F]	2392
Maxima [B] (verification not implemented)	2392
Giac [F]	2393
Mupad [F(-1)]	2393
Reduce [F]	2394

#### Optimal result

Integrand size = 18, antiderivative size = 132

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \frac{1}{6} ibx^3 + \frac{1}{2} x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) - \frac{1}{4} x^2 \log(1 - (1 + id)e^{2ia+2ibx}) + \frac{ix \operatorname{PolyLog}(2, (1 + id)e^{2ia+2ibx})}{4b} - \frac{\operatorname{PolyLog}(3, (1 + id)e^{2ia+2ibx})}{8b^2}$$

output

```
1/6*I*b*x^3+1/2*x^2*arctanh(1+I*d+d*cot(b*x+a))-1/4*x^2*ln(1-(1+I*d)*exp(2
*I*a+2*I*b*x))+1/4*I*x*polylog(2,(1+I*d)*exp(2*I*a+2*I*b*x))/b-1/8*polylog
(3,(1+I*d)*exp(2*I*a+2*I*b*x))/b^2
```



**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.90

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \frac{1}{2} x^2 \operatorname{arctanh}(1 + id + d \cot(a + bx)) - \frac{2b^2 x^2 \log\left(1 + \frac{ie^{-2i(a+bx)}}{-i+d}\right) + 2ibx \operatorname{PolyLog}\left(2, -\frac{ie^{-2i(a+bx)}}{-i+d}\right) + \operatorname{PolyLog}\left(3, -\frac{ie^{-2i(a+bx)}}{-i+d}\right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 + I*d + d*Cot[a + b*x]], x]
```

output

```
(x^2*ArcTanh[1 + I*d + d*Cot[a + b*x]])/2 - (2*b^2*x^2*Log[1 + I/((-I + d)*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, (-I)/((-I + d)*E^((2*I)*(a + b*x)))] + PolyLog[3, (-I)/((-I + d)*E^((2*I)*(a + b*x)))])/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.77 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.30, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6819, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d \cot(a + bx) + id + 1) dx$$

$$\downarrow \text{6819}$$

$$\frac{1}{2} ib \int \frac{x^2}{1 - (id + 1)e^{2ia + 2ibx}} dx + \frac{1}{2} x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} ib \left( \frac{x^3}{3} + (1 + id) \int \frac{e^{2ia + 2ibx} x^2}{1 - (id + 1)e^{2ia + 2ibx}} dx \right) + \frac{1}{2} x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1 + id) \left( \frac{\int x \log(1 - (id + 1)e^{2ia+2ibx}) dx}{b(-d + i)} - \frac{x^2 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 3011

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1 + id) \left( \frac{\frac{ix \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx}) dx}{2b}}{b(-d + i)} - \frac{x^2 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 2720

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1 + id) \left( \frac{\frac{ix \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2}}{b(-d + i)} - \frac{x^2 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

↓ 7143

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1 + id) \left( \frac{\frac{1}{2}x^2 \operatorname{arctanh}(d \cot(a + bx) + id + 1) + \frac{ix \operatorname{PolyLog}(2, (id+1)e^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, (id+1)e^{2ia+2ibx})}{4b^2}}{b(-d + i)} - \frac{x^2 \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right)$$

input

```
Int[x*ArcTanh[1 + I*d + d*Cot[a + b*x]],x]
```

output

```
(x^2*ArcTanh[1 + I*d + d*Cot[a + b*x])/2 + (I/2)*b*(x^3/3 + (1 + I*d)*(-1/2*(x^2*Log[1 - (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/(b*(I - d)) + (((I/2)*x*PolyLog[2, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/(b*(I - d)))
```

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6819 `Int[ArcTanh[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.64 (sec) , antiderivative size = 2289, normalized size of antiderivative = 17.34

method	result	size
risch	Expression too large to display	2289

input `int(x*arctanh(1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/6*I*b*x^3+1/4/b^2*a^2*d/(I-d)*\ln(I*\exp(2*I*(b*x+a))- \exp(2*I*(b*x+a))*d-I) \\ & -1/2/b^2*d*a^2/(I-d)*\ln(1-I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})-1/2/b^2*d*a^2 \\ & / (I-d)*\ln(1+I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/4/b^2*d/(I-d)*\ln(1+I*\exp(2 \\ & *I*(b*x+a))*(I-d))*a^2-1/4*I/b^2*a^2/(I-d)*\ln(I*\exp(2*I*(b*x+a))- \exp(2*I*( \\ & b*x+a))*d-I)-1/4*I/b^2/(I-d)*\ln(1+I*\exp(2*I*(b*x+a))*(I-d))*a^2+1/2*I/b^2* \\ & a^2/(I-d)*\ln(1-I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/2*I/b^2*a^2/(I-d)*\ln(1+ \\ & I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/2/b*d/(I-d)*\ln(1+I*\exp(2*I*(b*x+a))*(I \\ & -d))*a*x-1/2/b*d*a/(I-d)*\ln(1-I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})*x-1/2/b*d* \\ & a/(I-d)*\ln(1+I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})*x+1/2*I/b^2*d*a/(I-d)*\operatorname{dilog} \\ & (1-I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/2*I/b^2*d*a/(I-d)*\operatorname{dilog}(1+I*\exp(I*( \\ & b*x+a))*(I*(I-d))^{(1/2)})-1/4*I/b*d/(I-d)*\operatorname{polylog}(2,-I*\exp(2*I*(b*x+a))*(I- \\ & d))*x-1/4*I/b^2*d/(I-d)*\operatorname{polylog}(2,-I*\exp(2*I*(b*x+a))*(I-d))*a-1/2*I/b/(I- \\ & d)*\ln(1+I*\exp(2*I*(b*x+a))*(I-d))*a*x+1/2*I/b*a/(I-d)*\ln(1-I*\exp(I*(b*x+a) \\ & ))*(I*(I-d))^{(1/2)}*x+1/2*I/b*a/(I-d)*\ln(1+I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2) \\ & )}*x+1/8/b^2*d/(I-d)*\operatorname{polylog}(3,-I*\exp(2*I*(b*x+a))*(I-d))-1/4/b/(I-d)*\operatorname{polyl} \\ & \operatorname{og}(2,-I*\exp(2*I*(b*x+a))*(I-d))*x-1/4/b^2/(I-d)*\operatorname{polylog}(2,-I*\exp(2*I*(b*x+ \\ & a))*(I-d))*a+1/2/b^2*a/(I-d)*\operatorname{dilog}(1-I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/2 \\ & /b^2*a/(I-d)*\operatorname{dilog}(1+I*\exp(I*(b*x+a))*(I*(I-d))^{(1/2)})+1/4*d/(I-d)*\ln(1+I* \\ & \exp(2*I*(b*x+a))*(I-d))*x^2-1/4*I/(I-d)*\ln(1+I*\exp(2*I*(b*x+a))*(I-d))*x^2 \\ & -1/8*I/b^2/(I-d)*\operatorname{polylog}(3,-I*\exp(2*I*(b*x+a))*(I-d))+1/4*x^2*\ln(\exp(2* \dots \end{aligned}$$

**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.19

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$$

$$= \frac{4i b^3 x^3 + 6 b^2 x^2 \log\left(-\frac{((d-i)e^{(2i bx+2i a)}+i)e^{(-2i bx-2i a)}}{d}\right) + 4i a^3 + 6i bx \operatorname{Li}_2(-(-i d - 1)e^{(2i bx+2i a)}) - 6 a^2 \log}{24 b^2}$$

input `integrate(x*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="fricas")`

output `1/24*(4*I*b^3*x^3 + 6*b^2*x^2*log(-((d - I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d) + 4*I*a^3 + 6*I*b*x*dilog(-(-I*d - 1)*e^(2*I*b*x + 2*I*a)) - 6*a^2*log(((d - I)*e^(2*I*b*x + 2*I*a) + I)/(d - I)) - 6*(b^2*x^2 - a^2)*log((-I*d - 1)*e^(2*I*b*x + 2*I*a) + 1) - 3*polylog(3, (I*d + 1)*e^(2*I*b*x + 2*I*a)))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x \operatorname{atanh}(d \cot(a + bx) + id + 1) dx$$

input `integrate(x*atanh(1+I*d+d*cot(b*x+a)),x)`

output `Integral(x*atanh(d*cot(a + b*x) + I*d + 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 249 vs. 2(94) = 188.

Time = 0.05 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.89

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$$

$$= \frac{12((bx+a)^2 - 2(bx+a)a) \operatorname{artanh}(d \cot(bx+a) + id + 1)}{b} - \frac{-4i(bx+a)^3 + 12i(bx+a)^2 a - 6i bx \operatorname{Li}_2((i d + 1)e^{(2i bx+2i a)}) - 6(i(bx+a)^2 - 2i(bx+a))}{b}$$

input `integrate(x*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output `1/24*(12*((b*x + a)^2 - 2*(b*x + a)*a)*arctanh(d*cot(b*x + a) + I*d + 1)/b - (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog((I*d + 1)*e^(2*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a)*arctan2(d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), d*sin(2*b*x + 2*a) - cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^2 + 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1) + 3*polylog(3, (I*d + 1)*e^(2*I*b*x + 2*I*a)))/b)/b`

### Giac [F]

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x \operatorname{artanh}(d \cot(bx + a) + id + 1) dx$$

input `integrate(x*arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(d*cot(b*x + a) + I*d + 1), x)`

### Mupad [F(-1)]

Timed out.

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int x \operatorname{atanh}(d \cot(a + bx) + 1 + d li) dx$$

input `int(x*atanh(d*1i + d*cot(a + b*x) + 1),x)`

output `int(x*atanh(d*1i + d*cot(a + b*x) + 1), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + di + 1) x dx$$

input `int(x*atanh(1+I*d+d*cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)*d + d*i + 1)*x,x)`

### 3.340 $\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$

Optimal result	2395
Mathematica [B] (warning: unable to verify)	2395
Rubi [A] (verified)	2396
Maple [B] (verified)	2398
Fricas [A] (verification not implemented)	2399
Sympy [F]	2399
Maxima [B] (verification not implemented)	2400
Giac [F]	2400
Mupad [F(-1)]	2401
Reduce [F]	2401

#### Optimal result

Integrand size = 16, antiderivative size = 93

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \frac{1}{2}ibx^2 + x\operatorname{arctanh}(1 + id + d \cot(a + bx)) - \frac{1}{2}x \log(1 - (1 + id)e^{2ia+2ibx}) + \frac{i \operatorname{PolyLog}(2, (1 + id)e^{2ia+2ibx})}{4b}$$

output

```
1/2*I*b*x^2+x*arctanh(1+I*d+d*cot(b*x+a))-1/2*x*ln(1-(1+I*d)*exp(2*I*a+2*I*b*x))+1/4*I*polylog(2,(1+I*d)*exp(2*I*a+2*I*b*x))/b
```

#### Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 709 vs. 2(93) = 186.

Time = 9.93 (sec) , antiderivative size = 709, normalized size of antiderivative = 7.62

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = x\operatorname{arctanh}(1 + id + d \cot(a + bx)) + \frac{x \csc^2(a + bx) \left( 2bx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) + (i + \cot(a + bx))(2 + id + d \cot(a + bx)) \left( 2ibx + \log\left(1 + \frac{1}{2} \sec(bx)((-2 - id) \cos(a) + d \sin(a))\right) \cos(a) \right) \right)}{(i + \cot(a + bx))(2 + id + d \cot(a + bx)) \left( 2ibx + \log\left(1 + \frac{1}{2} \sec(bx)((-2 - id) \cos(a) + d \sin(a))\right) \cos(a) \right)}$$



input `Integrate[ArcTanh[1 + I*d + d*Cot[a + b*x]],x]`

output

```
x*ArcTanh[1 + I*d + d*Cot[a + b*x]] + (x*Csc[a + b*x]^2*(2*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] + I*Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*(d*Cos[a + b*x] + (2 + I*d)*Sin[a + b*x]))/(2*(-I + d))]*Log[1 - I*Tan[b*x]] - I*Log[(Sec[b*x]*((-I)*Cos[a] + Sin[a])*(d*Cos[a + b*x] + (2 + I*d)*Sin[a + b*x]))/2]*Log[1 + I*Tan[b*x]] + I*PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] - I*PolyLog[2, (Sec[b*x]*((2 + I*d)*Cos[a] - d*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2] + I*PolyLog[2, ((Cos[a] - I*Sin[a])*(-2 - I*d)*Cos[a] + d*Sin[a])*(I + Tan[b*x]))/(2*(-I + d))])*(Cos[b*x] - I*Sin[b*x])*(Cos[b*x] + I*Sin[b*x]))/((I + Cot[a + b*x])*(2 + I*d + d*Cot[a + b*x])*((2*I)*b*x + Log[1 + (Sec[b*x]*((-2 - I*d)*Cos[a] + d*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2] - Log[(Sec[b*x]*((-I)*Cos[a] + Sin[a])*(d*Cos[a + b*x] + (2 + I*d)*Sin[a + b*x]))/2] + ((-2*I + d)*Cos[a + b*x]*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])))/(d*Cos[a + b*x] + (2 + I*d)*Sin[a + b*x]) + (d*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])*Sin[a + b*x])/((-I)*d*Cos[a + b*x] + (-2*I + d)*Sin[a + b*x]) + 2*b*x*Tan[b*x] - I*Log[1 + (Sec[b*x]*((-2 - I*d)*Cos[a] + d*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2]*Tan[b*x] + I*Log[(Sec[b*x]*((-I)*Cos[a] + Sin[a])*(d*Cos[a + b*x] + (2 + I*d)*Sin[a + b*x]))/2]*Tan[b*x] - I*Log[1 - I*Tan[b*x]]*Tan[b*x] + I*Log[1 + I*Tan[b*x]]*Tan[b*x]))
```

### Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.35, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$ , Rules used = {6811, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d \cot(a + bx) + id + 1) dx$$

$$\downarrow \text{6811}$$

$$ib \int \frac{x}{1 - (id + 1)e^{2ia + 2ibx}} dx + x \operatorname{arctanh}(d \cot(a + bx) + id + 1)$$

$$\downarrow \text{2615}$$

$$\begin{aligned}
& ib \left( \frac{x^2}{2} + (1 + id) \int \frac{e^{2ia+2ibx} x}{1 - (id + 1)e^{2ia+2ibx}} dx \right) + x \operatorname{arctanh}(d \cot(a + bx) + id + 1) \\
& \quad \downarrow \text{2620} \\
& ib \left( \frac{x^2}{2} + (1 + id) \left( \frac{\int \log(1 - (id + 1)e^{2ia+2ibx}) dx}{2b(-d + i)} - \frac{x \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \\
& \quad \quad \quad x \operatorname{arctanh}(d \cot(a + bx) + id + 1) \\
& \quad \downarrow \text{2715} \\
& ib \left( \frac{x^2}{2} + (1 + id) \left( -\frac{i \int e^{-2ia-2ibx} \log(1 - (id + 1)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2(-d + i)} - \frac{x \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right) + \\
& \quad \quad \quad x \operatorname{arctanh}(d \cot(a + bx) + id + 1) \\
& \quad \downarrow \text{2838} \\
& ib \left( \frac{x^2}{2} + (1 + id) \left( \frac{x \operatorname{arctanh}(d \cot(a + bx) + id + 1) + i \operatorname{PolyLog}(2, (id + 1)e^{2ia+2ibx})}{4b^2(-d + i)} - \frac{x \log(1 - (1 + id)e^{2ia+2ibx})}{2b(-d + i)} \right) \right)
\end{aligned}$$

input `Int[ArcTanh[1 + I*d + d*Cot[a + b*x]], x]`

output `x*ArcTanh[1 + I*d + d*Cot[a + b*x]] + I*b*(x^2/2 + (1 + I*d)*(-1/2*(x*Log[1 - (1 + I*d)*E^((2*I)*a + (2*I)*b*x)])/(b*(I - d)) + ((I/4)*PolyLog[2, (1 + I*d)*E^((2*I)*a + (2*I)*b*x)]/(b^2*(I - d))))`

### Defintions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```

rule 2620 Int[((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

rule 6811 Int[ArcTanh[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)], x_Symbol] := Simp[x*Arc
Tanh[c + d*Cot[a + b*x]], x] + Simp[I*b Int[x/(c - I*d - c*E^(2*I*a + 2*I
*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - I*d)^2, 1]
    
```

### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 306 vs. 2(76) = 152.

Time = 3.87 (sec) , antiderivative size = 307, normalized size of antiderivative = 3.30

method	result
derivativedivides	$-\frac{i \operatorname{arctanh}(1+id+d \cot (bx+a)) d \ln (id+d \cot (bx+a))}{2} + \frac{i \operatorname{arctanh}(1+id+d \cot (bx+a)) d \ln (-id+d \cot (bx+a))}{2} - \frac{d^2 \left( i \left( \frac{\operatorname{dilog}\left(-\frac{i(ic)}{\dots}{\dots}\right)}{\dots} \right)}{\dots} \right)}{\dots}$
default	$-\frac{i \operatorname{arctanh}(1+id+d \cot (bx+a)) d \ln (id+d \cot (bx+a))}{2} + \frac{i \operatorname{arctanh}(1+id+d \cot (bx+a)) d \ln (-id+d \cot (bx+a))}{2} - \frac{d^2 \left( i \left( \frac{\operatorname{dilog}\left(-\frac{i(ic)}{\dots}{\dots}\right)}{\dots} \right)}{\dots} \right)}{\dots}$
risch	Expression too large to display

input `int(arctanh(1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output `1/b/d*(-1/2*I*arctanh(1+I*d+d*cot(b*x+a))*d*ln(I*d+d*cot(b*x+a))+1/2*I*arctanh(1+I*d+d*cot(b*x+a))*d*ln(-I*d+d*cot(b*x+a))-1/2*d^2*(-I/d*(1/2*dilog(-1/2*I*(I*d+d*cot(b*x+a))/d)+1/2*ln(-I*d+d*cot(b*x+a))*ln(-1/2*I*(I*d+d*cot(b*x+a))/d)-1/2*dilog(I*(-I*d+d*cot(b*x+a)-I*(2*I-2*d))/(2*I-2*d))-1/2*ln(-I*d+d*cot(b*x+a))*ln(I*(-I*d+d*cot(b*x+a)-I*(2*I-2*d))/(2*I-2*d))+I/d*(1/4*ln(I*d+d*cot(b*x+a))^2-1/2*dilog(1+1/2*I*d+1/2*d*cot(b*x+a))-1/2*ln(I*d+d*cot(b*x+a))*ln(1+1/2*I*d+1/2*d*cot(b*x+a))))`

### Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.31

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx$$

$$= \frac{2i b^2 x^2 + 2bx \log\left(-\frac{((d-i)e^{2i bx+2i a}+i)e^{(-2i bx-2i a)}}{d}\right) - 2i a^2 - 2(bx + a) \log((-id - 1)e^{(2i bx+2i a)} + 1) + 2}{4b}$$

input `integrate(arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="fricas")`

output `1/4*(2*I*b^2*x^2 + 2*b*x*log(-((d - I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d) - 2*I*a^2 - 2*(b*x + a)*log((-I*d - 1)*e^(2*I*b*x + 2*I*a) + 1) + 2*a*log(((d - I)*e^(2*I*b*x + 2*I*a) + I)/(d - I)) + I*dilog(-(-I*d - 1)*e^(2*I*b*x + 2*I*a)))/b`

### Sympy [F]

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{atanh}(d \cot(a + bx) + id + 1) dx$$

input `integrate(atanh(1+I*d+d*cot(b*x+a)),x)`

output `Integral(atanh(d*cot(a + b*x) + I*d + 1), x)`

### Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 288 vs.  $2(66) = 132$ .

Time = 0.11 (sec) , antiderivative size = 288, normalized size of antiderivative = 3.10

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx =$$

$$\frac{4(bx + a)d \left( \frac{\log((id+2)\tan(bx+a)+d)}{d} - \frac{\log(i \tan(bx+a)+1)}{d} \right) - d \left( \frac{2i \left( \log((id+2)\tan(bx+a)+d) \log\left(\frac{(d-2i)\tan(bx+a)-id}{2id+2} + 1\right)}{d} \right)}{d} \right)}{d}$$

input `integrate(arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output `-1/8*(4*(b*x + a)*d*(log((I*d + 2)*tan(b*x + a) + d)/d - log(I*tan(b*x + a) + 1)/d) - d*(2*I*(log((I*d + 2)*tan(b*x + a) + d)*log(((d - 2*I)*tan(b*x + a) - I*d)/(2*I*d + 2) + 1) + dilog(-((d - 2*I)*tan(b*x + a) - I*d)/(2*I*d + 2)))/d + 2*I*(log(1/2*(d - 2*I)*tan(b*x + a) - 1/2*I*d)*log(I*tan(b*x + a) + 1) + dilog(-1/2*(d - 2*I)*tan(b*x + a) + 1/2*I*d + 1))/d - (2*I*log((I*d + 2)*tan(b*x + a) + d)*log(I*tan(b*x + a) + 1) - I*log(I*tan(b*x + a) + 1)^2)/d - 2*I*(log(I*tan(b*x + a) + 1)*log(-1/2*I*tan(b*x + a) + 1/2) + dilog(1/2*I*tan(b*x + a) + 1/2))/d - 8*(b*x + a)*arctanh(I*d + d/tan(b*x + a) + 1))/b`

### Giac [F]

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{artanh}(d \cot(bx + a) + id + 1) dx$$

input `integrate(arctanh(1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(d*cot(b*x + a) + I*d + 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{atanh}(d \cot(a + bx) + 1 + d i) dx$$

input `int(atanh(d*I*i + d*cot(a + b*x) + 1),x)`

output `int(atanh(d*I*i + d*cot(a + b*x) + 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 + id + d \cot(a + bx)) dx = \int \operatorname{atanh}(\cot(bx + a) d + di + 1) dx$$

input `int(atanh(1+I*d+d*cot(b*x+a)),x)`

output `int(atanh(cot(a + b*x)*d + d*i + 1),x)`

### 3.341 $\int \frac{\operatorname{arctanh}(1+id+d \cot(a+bx))}{x} dx$

Optimal result	2402
Mathematica [N/A]	2402
Rubi [N/A]	2403
Maple [N/A]	2403
Fricas [N/A]	2404
Sympy [N/A]	2404
Maxima [N/A]	2404
Giac [N/A]	2405
Mupad [N/A]	2405
Reduce [N/A]	2406

#### Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctanh(1+I*d+d*cot(b*x+a))/x,x)`

#### Mathematica [N/A]

Not integrable

Time = 0.80 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx$$

input `Integrate[ArcTanh[1 + I*d + d*Cot[a + b*x]]/x,x]`

output `Integrate[ArcTanh[1 + I*d + d*Cot[a + b*x]]/x, x]`

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d \cot(a + bx) + id + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d \cot(a + bx) + id + 1)}{x} dx$$

input `Int[ArcTanh[1 + I*d + d*Cot[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.58 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(bx + a))}{x} dx$$

input `int(arctanh(1+I*d+d*cot(b*x+a))/x,x)`

output `int(arctanh(1+I*d+d*cot(b*x+a))/x,x)`



**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + id + 1)}{x} dx$$

input `integrate(arctanh(1+I*d+d*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*log(-((d - I)*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/d)/x, x)`

**Sympy [N/A]**

Not integrable

Time = 1.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \cot(a + bx) + id + 1)}{x} dx$$

input `integrate(atanh(1+I*d+d*cot(b*x+a))/x,x)`

output `Integral(atanh(d*cot(a + b*x) + I*d + 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 4.32 (sec) , antiderivative size = 143, normalized size of antiderivative = 7.15

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + id + 1)}{x} dx$$

input `integrate(arctanh(1+I*d+d*cot(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x + 1/4*(-I*pi - 4*I*a - 2*log(-d))*log(x) + 1/2*I*integrate(arctan2(
d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) + cos(2*b*x + 2
*a) - 1)/x, x) + 1/4*integrate(log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1
)*sin(2*b*x + 2*a)^2 + 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1)/x, x
)
```

**Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{artanh}(d \cot(bx + a) + id + 1)}{x} dx$$

input

```
integrate(arctanh(1+I*d+d*cot(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctanh(d*cot(b*x + a) + I*d + 1)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 4.47 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(d \cot(a + bx) + 1 + d li)}{x} dx$$

input

```
int(atanh(d*1i + d*cot(a + b*x) + 1)/x,x)
```

output

```
int(atanh(d*1i + d*cot(a + b*x) + 1)/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 + id + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atanh}(\cot(bx + a) d + di + 1)}{x} dx$$

input `int(atanh(1+I*d+d*cot(b*x+a))/x,x)`output `int(atanh(cot(a + b*x)*d + d*i + 1)/x,x)`

### 3.342 $\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$

Optimal result	2407
Mathematica [A] (verified)	2408
Rubi [A] (verified)	2408
Maple [C] (warning: unable to verify)	2411
Fricas [A] (verification not implemented)	2412
Sympy [F]	2413
Maxima [B] (verification not implemented)	2413
Giac [F]	2414
Mupad [F(-1)]	2414
Reduce [F]	2414

#### Optimal result

Integrand size = 21, antiderivative size = 169

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \operatorname{arctanh}(1 - id - d \cot(a + bx)) - \frac{1}{6} x^3 \log(1 - (1 - id)e^{2ia+2ibx}) + \frac{ix^2 \operatorname{PolyLog}(2, (1 - id)e^{2ia+2ibx})}{4b} - \frac{x \operatorname{PolyLog}(3, (1 - id)e^{2ia+2ibx})}{4b^2} - \frac{i \operatorname{PolyLog}(4, (1 - id)e^{2ia+2ibx})}{8b^3}$$

output

```
1/12*I*b*x^4-1/3*x^3*arctanh(-1+I*d+d*cot(b*x+a))-1/6*x^3*ln(1-(1-I*d)*exp(2*I*a+2*I*b*x))+1/4*I*x^2*polylog(2,(1-I*d)*exp(2*I*a+2*I*b*x))/b-1/4*x*polylog(3,(1-I*d)*exp(2*I*a+2*I*b*x))/b^2-1/8*I*polylog(4,(1-I*d)*exp(2*I*a+2*I*b*x))/b^3
```

**Mathematica [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.92

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \frac{1}{3} x^3 \operatorname{arctanh}(1 - id - d \cot(a + bx)) - \frac{4b^3 x^3 \log\left(1 + \frac{e^{-2i(a+bx)}}{-1+id}\right) + 6ib^2 x^2 \operatorname{PolyLog}\left(2, \frac{ie^{-2i(a+bx)}}{i+d}\right) + 6bx \operatorname{PolyLog}\left(3, \frac{ie^{-2i(a+bx)}}{i+d}\right) - 3i \operatorname{PolyLog}\left(4, \frac{ie^{-2i(a+bx)}}{i+d}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTanh[1 - I*d - d*Cot[a + b*x]],x]
```

output

```
(x^3*ArcTanh[1 - I*d - d*Cot[a + b*x]])/3 - (4*b^3*x^3*Log[1 + 1/((-1 + I*d)*E^((2*I)*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, I/((I + d)*E^((2*I)*(a + b*x)))] + 6*b*x*PolyLog[3, I/((I + d)*E^((2*I)*(a + b*x)))] - (3*I)*PolyLog[4, I/((I + d)*E^((2*I)*(a + b*x)))])/(24*b^3)
```

**Rubi [A] (verified)**Time = 0.89 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.28, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6819, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1) dx$$

$$\downarrow \text{6819}$$

$$\frac{1}{3} ib \int \frac{x^3}{1 - (1 - id)e^{2ia+2ibx}} dx + \frac{1}{3} x^3 \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{3} ib \left( \frac{x^4}{4} + (1 - id) \int \frac{e^{2ia+2ibx} x^3}{1 - (1 - id)e^{2ia+2ibx}} dx \right) + \frac{1}{3} x^3 \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 - id) \left( \frac{3 \int x^2 \log(1 - (1 - id)e^{2ia+2ibx}) dx}{2b(d+i)} - \frac{x^3 \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d - \cot(a + bx)) - id + 1$$

↓ 3011

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 - id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx}) dx}{b} \right)}{2b(d+i)} - \frac{x^3 \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d - \cot(a + bx)) - id + 1$$

↓ 7163

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 - id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{i \left( \frac{\int \operatorname{PolyLog}(3, (1-id)e^{2ia+2ibx}) dx}{2b} - \frac{ix \operatorname{PolyLog}(3, (1-id)e^{2ia+2ibx})}{2b} \right)}{b} \right)}{2b(d+i)} - \frac{x^3 \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d - \cot(a + bx)) - id + 1$$

↓ 2720

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1 - id) \left( \frac{3 \left( \frac{ix^2 \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{i \left( \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(3, (1-id)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, (1-id)e^{2ia+2ibx})}{2b} \right)}{b} \right)}{2b(d+i)} - \frac{x^3 \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{3}x^3 \operatorname{arctanh}(d - \cot(a + bx)) - id + 1$$

↓ 7143

$$\frac{1}{3}ib \left( \frac{x^4}{4} + (1-id) \left( \frac{3 \left( \frac{ix^2 \text{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{i \left( \frac{\text{PolyLog}(4, (1-id)e^{2ia+2ibx})}{4b^2} - \frac{ix \text{PolyLog}(3, (1-id)e^{2ia+2ibx})}{2b} \right)}{b} \right)}{2b(d+i)} \right) - x^3 \right)$$

input `Int[x^2*ArcTanh[1 - I*d - d*Cot[a + b*x]],x]`

output

```
(x^3*ArcTanh[1 - I*d - d*Cot[a + b*x]])/3 + (I/3)*b*(x^4/4 + (1 - I*d)*(-1/2*(x^3*Log[1 - (1 - I*d)*E^((2*I)*a + (2*I)*b*x)])/(b*(I + d)) + (3*(((I/2)*x^2*PolyLog[2, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)])/b - (I*(((1/2*I)*x*PolyLog[3, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)])/b + PolyLog[4, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b)/(2*b*(I + d)))
```

### Defintions of rubi rules used

rule 2615

```
Int[(((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6819 `Int[ArcTanh[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.58 (sec) , antiderivative size = 2277, normalized size of antiderivative = 13.47

method	result	size
risch	Expression too large to display	2277

input `int(-x^2*arctanh(-1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`



output

```

1/6/b^3*a^3*d/(I+d)*ln(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I)-1/2*I/b^3*
a^3/(I+d)*ln(1+I*exp(I*(b*x+a))*(I*(I+d))^(1/2))-1/2*I/b^3*a^3/(I+d)*ln(1-
I*exp(I*(b*x+a))*(I*(I+d))^(1/2))-1/4*I/b^2/(I+d)*polylog(3,-I*(I+d)*exp(2
*I*(b*x+a)))*x+1/3*I/b^3/(I+d)*ln(1+I*(I+d)*exp(2*I*(b*x+a)))*a^3-1/8*I/b^
3*d/(I+d)*polylog(4,-I*(I+d)*exp(2*I*(b*x+a)))+1/6*I/b^3*a^3/(I+d)*ln(I*ex
p(2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I)+1/8/b^3/(I+d)*polylog(4,-I*(I+d)*exp(
2*I*(b*x+a)))+1/12*I*b*x^4-1/12*(-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*
I*(b*x+a)))^3+2*I*Pi+2*ln(d)-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*I*(b*
x+a))*csgn(I/(-1+exp(2*I*(b*x+a))))*exp(2*I*(b*x+a))*d)*csgn(I*d)-I*Pi*csg
n(I*exp(2*I*(b*x+a)))*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I/(-1+exp(2*I*(b*
x+a)))*exp(2*I*(b*x+a)))-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a))))*csgn(I*(I*exp(
2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I)/(-1+exp(2*I*(b*x+a))))^2-I*Pi*csgn(I*(I
*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I))*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*
I*(b*x+a))*d-I)/(-1+exp(2*I*(b*x+a))))^2+I*Pi*csgn(I/(-1+exp(2*I*(b*x+a)))
*exp(2*I*(b*x+a))*d)*csgn(1/(-1+exp(2*I*(b*x+a)))*exp(2*I*(b*x+a))*d)^2-I*
Pi*csgn(I*exp(I*(b*x+a)))^2*csgn(I*exp(2*I*(b*x+a)))+I*Pi*csgn(I/(-1+exp(2
*I*(b*x+a)))*csgn(I*(I*exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I))*csgn(I*(I*
exp(2*I*(b*x+a))+exp(2*I*(b*x+a))*d-I)/(-1+exp(2*I*(b*x+a))))-I*Pi*csgn(1/
(-1+exp(2*I*(b*x+a)))*exp(2*I*(b*x+a))*d)^2-I*Pi*csgn(I/(-1+exp(2*I*(b*x+a)
)))*exp(2*I*(b*x+a))*d)*csgn(1/(-1+exp(2*I*(b*x+a)))*exp(2*I*(b*x+a))*d...

```

### Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.07

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$$

$$= \frac{2i b^4 x^4 - 4 b^3 x^3 \log\left(-\frac{de^{(2i bx + 2i a)}}{(d+i)e^{(2i bx + 2i a)} - i}\right) + 6i b^2 x^2 \operatorname{Li}_2(-(id - 1)e^{(2i bx + 2i a)}) - 2i a^4 + 4a^3 \log\left(\frac{(d+i)e^{(2i bx + 2i a)}}{d+i}\right)}{1}$$

input

```
integrate(-x^2*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/24*(2*I*b^4*x^4 - 4*b^3*x^3*log(-d*e^(2*I*b*x + 2*I*a)/((d + I)*e^(2*I*b
*x + 2*I*a) - I)) + 6*I*b^2*x^2*dilog(-(I*d - 1)*e^(2*I*b*x + 2*I*a)) - 2*
I*a^4 + 4*a^3*log(((d + I)*e^(2*I*b*x + 2*I*a) - I)/(d + I)) - 6*b*x*polylo
g(3, (-I*d + 1)*e^(2*I*b*x + 2*I*a)) - 4*(b^3*x^3 + a^3)*log((I*d - 1)*e^
(2*I*b*x + 2*I*a) + 1) - 3*I*polylog(4, (-I*d + 1)*e^(2*I*b*x + 2*I*a))/b
^3

```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \int x^2 \operatorname{atanh}(d \cot(a + bx) + id - 1) dx$$

input `integrate(-x**2*atanh(-1+I*d+d*cot(b*x+a)),x)`

output `-Integral(x**2*atanh(d*cot(a + b*x) + I*d - 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 345 vs.  $2(119) = 238$ .

Time = 0.07 (sec) , antiderivative size = 345, normalized size of antiderivative = 2.04

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \frac{12((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2) \operatorname{artanh}(d \cot(bx+a) + id - 1)}{b^2} + \frac{-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(4i(bx+a)^3 - 9i(bx+a)^2 a + 3i a^2) \operatorname{arctan}(\frac{d \cot(bx+a) + id - 1}{d \cot(bx+a) + id - 1})}{b^2}$$

input `integrate(-x^2*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output `-1/36*(12*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctanh(d*cot(b*x + a) + I*d - 1)/b^2 + (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 - 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*I*(b*x + a)*a^2)*arctan2(-d*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) - cos(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog((-I*d + 1)*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)*sin(2*b*x + 2*a)^2 - 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3, (-I*d + 1)*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, (-I*d + 1)*e^(2*I*b*x + 2*I*a)))/b^2)/b`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -x^2 \operatorname{artanh}(d \cot(bx + a) + id - 1) dx$$

input `integrate(-x^2*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-x^2*arctanh(d*cot(b*x + a) + I*d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -x^2 \operatorname{atanh}(d \cot(a + bx) - 1 + di) dx$$

input `int(-x^2*atanh(d*Ii + d*cot(a + b*x) - 1),x)`

output `int(-x^2*atanh(d*Ii + d*cot(a + b*x) - 1), x)`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \left( \int \operatorname{atanh}(\cot(bx + a) d + di - 1) x^2 dx \right)$$

input `int(-x^2*atanh(-1+I*d+d*cot(b*x+a)),x)`

output `- int(atanh(cot(a + b*x)*d + d*i - 1)*x**2,x)`

### 3.343 $\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$

Optimal result	2415
Mathematica [A] (verified)	2416
Rubi [A] (verified)	2416
Maple [C] (warning: unable to verify)	2419
Fricas [A] (verification not implemented)	2420
Sympy [F]	2420
Maxima [B] (verification not implemented)	2420
Giac [F]	2421
Mupad [F(-1)]	2421
Reduce [F]	2422

#### Optimal result

Integrand size = 19, antiderivative size = 133

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \frac{1}{6} ibx^3 + \frac{1}{2} x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx)) - \frac{1}{4} x^2 \log(1 - (1 - id)e^{2ia+2ibx}) + \frac{ix \operatorname{PolyLog}(2, (1 - id)e^{2ia+2ibx})}{4b} - \frac{\operatorname{PolyLog}(3, (1 - id)e^{2ia+2ibx})}{8b^2}$$

output

```
1/6*I*b*x^3-1/2*x^2*arctanh(-1+I*d+d*cot(b*x+a))-1/4*x^2*ln(1-(1-I*d)*exp(
2*I*a+2*I*b*x))+1/4*I*x*polylog(2,(1-I*d)*exp(2*I*a+2*I*b*x))/b-1/8*polylo
g(3,(1-I*d)*exp(2*I*a+2*I*b*x))/b^2
```

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.89

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$$

$$= \frac{1}{2} x^2 \operatorname{arctanh}(1 - id - d \cot(a + bx))$$

$$- \frac{2b^2 x^2 \log\left(1 + \frac{e^{-2i(a+bx)}}{-1+id}\right) + 2ibx \operatorname{PolyLog}\left(2, \frac{ie^{-2i(a+bx)}}{i+d}\right) + \operatorname{PolyLog}\left(3, \frac{ie^{-2i(a+bx)}}{i+d}\right)}{8b^2}$$

input

```
Integrate[x*ArcTanh[1 - I*d - d*Cot[a + b*x]], x]
```

output

```
(x^2*ArcTanh[1 - I*d - d*Cot[a + b*x]])/2 - (2*b^2*x^2*Log[1 + 1/((-1 + I*d)*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, I/((I + d)*E^((2*I)*(a + b*x)))] + PolyLog[3, I/((I + d)*E^((2*I)*(a + b*x)))])/(8*b^2)
```

**Rubi [A] (verified)**

Time = 0.71 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.27, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$ , Rules used = {6819, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1) dx$$

$$\downarrow \text{6819}$$

$$\frac{1}{2} ib \int \frac{x^2}{1 - (1 - id)e^{2ia+2ibx}} dx + \frac{1}{2} x^2 \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} ib \left( \frac{x^3}{3} + (1 - id) \int \frac{e^{2ia+2ibx} x^2}{1 - (1 - id)e^{2ia+2ibx}} dx \right) + \frac{1}{2} x^2 \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1-id) \left( \frac{\int x \log(1 - (1-id)e^{2ia+2ibx}) dx}{b(d+i)} - \frac{x^2 \log(1 - (1-id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\cot(a+bx)) - id + 1)$$

↓ 3011

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1-id) \left( \frac{\frac{ix \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx}) dx}{2b}}{b(d+i)} - \frac{x^2 \log(1 - (1-id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\cot(a+bx)) - id + 1)$$

↓ 2720

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1-id) \left( \frac{\frac{ix \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2}}{b(d+i)} - \frac{x^2 \log(1 - (1-id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \frac{1}{2}x^2 \operatorname{arctanh}(d(-\cot(a+bx)) - id + 1)$$

↓ 7143

$$\frac{1}{2}ib \left( \frac{x^3}{3} + (1-id) \left( \frac{\frac{\frac{1}{2}x^2 \operatorname{arctanh}(d(-\cot(a+bx)) - id + 1) + \frac{ix \operatorname{PolyLog}(2, (1-id)e^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, (1-id)e^{2ia+2ibx})}{4b^2}}{b(d+i)} - \frac{x^2 \log(1 - (1-id)e^{2ia+2ibx})}{2b(d+i)}}{b(d+i)} \right) \right)$$

input `Int[x*ArcTanh[1 - I*d - d*Cot[a + b*x]],x]`

output `(x^2*ArcTanh[1 - I*d - d*Cot[a + b*x])/2 + (I/2)*b*(x^3/3 + (1 - I*d)*(-1/2*(x^2*Log[1 - (1 - I*d)*E^((2*I)*a + (2*I)*b*x)])/(b*(I + d)) + (((I/2)*x*PolyLog[2, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(4*b^2))/(b*(I + d))))`

## Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6819 `Int[ArcTanh[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTanh[c + d*Cot[a + b*x]]/(f*(m + 1))), x] + Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - I*d)^2, 1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.55 (sec) , antiderivative size = 2187, normalized size of antiderivative = 16.44

method	result	size
risch	Expression too large to display	2187

input `int(-x*arctanh(-1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & \frac{1}{2} \frac{b^2 d a}{(I+d)} \ln(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) * x + \frac{1}{2} \frac{b^2 d a}{(I+d)} \ln(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) * x \\ & - \frac{1}{2} \frac{I}{b} \frac{d}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * a * x - \frac{1}{2} \frac{I}{b^2} \frac{d a}{(I+d)} \operatorname{dilog}(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) \\ & - \frac{1}{2} \frac{I}{b^2} \frac{d a}{(I+d)} \operatorname{dilog}(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) + \frac{1}{4} \frac{I}{b} \frac{d}{(I+d)} \operatorname{polylog}(2, -I*(I+d) \exp(2*I(b*x+a))) * x \\ & + \frac{1}{4} \frac{I}{b^2} \frac{d}{(I+d)} \operatorname{polylog}(2, -I*(I+d) \exp(2*I(b*x+a))) * a + \frac{1}{2} \frac{I}{b} \frac{a}{(I+d)} \ln(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) * x \\ & + \frac{1}{2} \frac{I}{b} \frac{a}{(I+d)} \ln(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) * x + \frac{1}{6} I b^2 x^3 - \frac{1}{2} \frac{b^2 d}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * a * x - \frac{1}{4} \frac{b^2 a^2 d}{(I+d)} \ln(I \exp(2*I(b*x+a)) + \exp(2*I(b*x+a)) * d - I) \\ & - \frac{1}{4} \frac{b^2 d}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * a^2 + \frac{1}{2} \frac{I}{b^2} \frac{d a^2}{(I+d)} \ln(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) + \frac{1}{2} \frac{I}{b^2} \frac{d a^2}{(I+d)} \ln(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) \\ & + \frac{1}{2} \frac{I}{b^2} \frac{d a^2}{(I+d)} \ln(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) + \frac{1}{2} \frac{I}{b^2} \frac{d a^2}{(I+d)} \ln(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) - \frac{1}{4} \frac{I}{b^2} \frac{d}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * a^2 \\ & - \frac{1}{4} \frac{I}{b^2} \frac{d a^2}{(I+d)} \ln(I \exp(2*I(b*x+a)) + \exp(2*I(b*x+a)) * d - I) - \frac{1}{4} \frac{d}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * x^2 - \frac{1}{4} \frac{b}{(I+d)} \operatorname{polylog}(2, -I*(I+d) \exp(2*I(b*x+a))) * x \\ & - \frac{1}{4} \frac{b^2}{(I+d)} \operatorname{polylog}(2, -I*(I+d) \exp(2*I(b*x+a))) * a - \frac{1}{8} \frac{b^2 d}{(I+d)} \operatorname{polylog}(3, -I*(I+d) \exp(2*I(b*x+a))) + \frac{1}{2} \frac{b^2 a}{(I+d)} \operatorname{dilog}(1+I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) \\ & + \frac{1}{2} \frac{b^2 a}{(I+d)} \operatorname{dilog}(1-I \exp(I(b*x+a)) * (I*(I+d))^{1/2}) - \frac{1}{4} \frac{I}{(I+d)} \ln(1+I*(I+d) \exp(2*I(b*x+a))) * x^2 - \frac{1}{8} \frac{I}{b^2} \frac{d}{(I+d)} \operatorname{polylog}(3, -I*(I+d) \exp(2*I(b*x+a))) - \frac{1}{8} * (-I * \operatorname{Pi} * \operatorname{csgn}(I \dots \end{aligned}$$



**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.18

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$$

$$= \frac{4i b^3 x^3 - 6 b^2 x^2 \log\left(-\frac{d e^{(2i b x + 2i a)}}{(d+i)e^{(2i b x + 2i a)} - i}\right) + 4i a^3 + 6i b x \operatorname{Li}_2(-i d - 1) e^{(2i b x + 2i a)} - 6 a^2 \log\left(\frac{(d+i)e^{(2i b x + 2i a)}}{d+i}\right)}{24 b^2}$$

input `integrate(-x*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="fricas")`

output `1/24*(4*I*b^3*x^3 - 6*b^2*x^2*log(-d*e^(2*I*b*x + 2*I*a)/((d + I)*e^(2*I*b*x + 2*I*a) - I)) + 4*I*a^3 + 6*I*b*x*dilog(-(I*d - 1)*e^(2*I*b*x + 2*I*a)) - 6*a^2*log(((d + I)*e^(2*I*b*x + 2*I*a) - I)/(d + I)) - 6*(b^2*x^2 - a^2)*log((I*d - 1)*e^(2*I*b*x + 2*I*a) + 1) - 3*polylog(3, (-I*d + 1)*e^(2*I*b*x + 2*I*a)))/b^2`

**Sympy [F]**

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \int x \operatorname{atanh}(d \cot(a + bx) + id - 1) dx$$

input `integrate(-x*atanh(-1+I*d+d*cot(b*x+a)),x)`

output `-Integral(x*atanh(d*cot(a + b*x) + I*d - 1), x)`

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 250 vs. 2(94) = 188.

Time = 0.06 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.88

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx =$$

$$\frac{12((bx+a)^2 - 2(bx+a)a) \operatorname{artanh}(d \cot(bx+a) + id - 1)}{b} + \frac{-4i(bx+a)^3 + 12i(bx+a)^2 a - 6i b x \operatorname{Li}_2((-i d + 1) e^{(2i b x + 2i a)}) - 6(i(bx+a)^2 - 2i a)}{b}$$

input `integrate(-x*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output `-1/24*(12*((b*x + a)^2 - 2*(b*x + a)*a)*arctanh(d*cot(b*x + a) + I*d - 1)/  
b + (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog((-I*d + 1)*e^(2  
*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a)*arctan2(-d*cos(2*b*  
x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) - cos(2*b*x + 2*a) + 1) +  
3*((b*x + a)^2 - 2*(b*x + a)*a)*log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 +  
1)*sin(2*b*x + 2*a)^2 - 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1) +  
3*polylog(3, (-I*d + 1)*e^(2*I*b*x + 2*I*a)))/b/b`

### Giac [F]

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -x \operatorname{artanh}(d \cot(bx + a) + id - 1) dx$$

input `integrate(-x*arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctanh(d*cot(b*x + a) + I*d - 1), x)`

### Mupad [F(-1)]

Timed out.

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -x \operatorname{atanh}(d \cot(a + bx) - 1 + d li) dx$$

input `int(-x*atanh(d*1i + d*cot(a + b*x) - 1),x)`

output `int(-x*atanh(d*1i + d*cot(a + b*x) - 1), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \left( \int \operatorname{atanh}(\cot(bx + a) d + di - 1) x dx \right)$$

input `int(-x*atanh(-1+I*d+d*cot(b*x+a)),x)`

output `- int(atanh(cot(a + b*x)*d + d*i - 1)*x,x)`

### 3.344 $\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$

Optimal result	2423
Mathematica [B] (warning: unable to verify)	2423
Rubi [A] (verified)	2424
Maple [B] (verified)	2426
Fricas [A] (verification not implemented)	2427
Sympy [F]	2427
Maxima [B] (verification not implemented)	2428
Giac [F]	2428
Mupad [F(-1)]	2429
Reduce [F]	2429

#### Optimal result

Integrand size = 17, antiderivative size = 94

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \frac{1}{2}ibx^2 + x\operatorname{arctanh}(1 - id - d \cot(a + bx)) - \frac{1}{2}x \log(1 - (1 - id)e^{2ia+2ibx}) + \frac{i \operatorname{PolyLog}(2, (1 - id)e^{2ia+2ibx})}{4b}$$

output `1/2*I*b*x^2-x*arctanh(-1+I*d+d*cot(b*x+a))-1/2*x*ln(1-(1-I*d)*exp(2*I*a+2*I*b*x))+1/4*I*polylog(2,(1-I*d)*exp(2*I*a+2*I*b*x))/b`

#### Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 605 vs. 2(94) = 188.

Time = 8.98 (sec) , antiderivative size = 605, normalized size of antiderivative = 6.44

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = x\operatorname{arctanh}(1 - id - d \cot(a + bx)) + \frac{x \csc^2(a + bx) \left( 2bx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) + i \log \left( \frac{\sec(bx)(\cos(a) - i \sin(a))(d \cos(a + bx) + i(2i + d) \sin(a))}{2(i + d)} \right) \right)}{(i + \cot(a + bx))(-2 + id + d \cot(a + bx))}$$

input `Integrate[ArcTanh[1 - I*d - d*Cot[a + b*x]],x]`

output `x*ArcTanh[1 - I*d - d*Cot[a + b*x]] + (x*Csc[a + b*x]^2*(2*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] + I*Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*(d*Cos[a + b*x] + I*(2*I + d)*Sin[a + b*x])]/(2*(I + d))]*Log[1 - I*Tan[b*x]] - I*Log[(I*Sec[b*x]*(d*Cos[a + b*x] + I*(2*I + d)*Sin[a + b*x])]/(2*Cos[a] - (2*I)*Sin[a])]*Log[1 + I*Tan[b*x]] + I*PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] - I*PolyLog[2, (Sec[b*x]*((2 - I*d)*Cos[a] + d*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2] + I*PolyLog[2, ((Cos[a] - I*Sin[a])*((2 - I*d)*Cos[a] + d*Sin[a])*(I + Tan[b*x]))/(2*(I + d))])*(Cos[b*x] - I*Sin[b*x])*(Cos[b*x] + I*Sin[b*x]))/((I + Cot[a + b*x])*(-2 + I*d + d*Cot[a + b*x])*(-((Log[1 - I*Tan[b*x]]*Sec[b*x]*((2*I + d)*Cos[a] + I*d*Sin[a]))/(d*Cos[a + b*x] + I*(2*I + d)*Sin[a + b*x])) + (Log[1 + I*Tan[b*x]]*Sec[b*x]*((2*I + d)*Cos[a] + I*d*Sin[a]))/(d*Cos[a + b*x] + I*(2*I + d)*Sin[a + b*x]) + (Log[(I*Sec[b*x]*(d*Cos[a + b*x] + I*(2*I + d)*Sin[a + b*x])]/(2*Cos[a] - (2*I)*Sin[a]))*Sec[b*x]^2)/(1 + I*Tan[b*x]) - 2*b*x*(I + Tan[b*x]) + I*Log[1 - (Sec[b*x]*((2 - I*d)*Cos[a] + d*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2]*(I + Tan[b*x]))))`

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.31, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {6811, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1) dx$$

$$\downarrow 6811$$

$$ib \int \frac{x}{1 - (1 - id)e^{2ia + 2ibx}} dx + x \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\downarrow 2615$$

$$ib \left( \frac{x^2}{2} + (1 - id) \int \frac{e^{2ia + 2ibx} x}{1 - (1 - id)e^{2ia + 2ibx}} dx \right) + x \operatorname{arctanh}(d(-\cot(a + bx)) - id + 1)$$

$$\begin{aligned}
& \downarrow 2620 \\
& ib \left( \frac{x^2}{2} + (1 - id) \left( \frac{\int \log(1 - (1 - id)e^{2ia+2ibx}) dx}{2b(d+i)} - \frac{x \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \\
& \quad \quad \quad \text{arctanh}(d(-\cot(a+bx)) - id + 1) \\
& \downarrow 2715 \\
& ib \left( \frac{x^2}{2} + (1 - id) \left( -\frac{i \int e^{-2ia-2ibx} \log(1 - (1 - id)e^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2(d+i)} - \frac{x \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right) + \\
& \quad \quad \quad \text{arctanh}(d(-\cot(a+bx)) - id + 1) \\
& \downarrow 2838 \\
& ib \left( \frac{x^2}{2} + (1 - id) \left( \frac{\text{arctanh}(d(-\cot(a+bx)) - id + 1) + i \text{PolyLog}(2, (1 - id)e^{2ia+2ibx})}{4b^2(d+i)} - \frac{x \log(1 - (1 - id)e^{2ia+2ibx})}{2b(d+i)} \right) \right)
\end{aligned}$$

input `Int[ArcTanh[1 - I*d - d*Cot[a + b*x]],x]`

output `x*ArcTanh[1 - I*d - d*Cot[a + b*x]] + I*b*(x^2/2 + (1 - I*d)*(-1/2*(x*Log[1 - (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(b*(I + d)) + ((I/4)*PolyLog[2, (1 - I*d)*E^((2*I)*a + (2*I)*b*x)]/(b^2*(I + d))))`

### Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 6811 Int[ArcTanh[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)], x_Symbol] :> Simp[x*Arc
Tanh[c + d*Cot[a + b*x]], x] + Simp[I*b Int[x/(c - I*d - c*E^(2*I*a + 2*I
*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - I*d)^2, 1]
```

### Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 331 vs. 2(77) = 154.

Time = 4.68 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.53

method	result
derivativedivides	$-\frac{i \operatorname{arctanh}(-1+id+d \cot(\frac{bx+a}{2}))d \ln(id+d \cot(\frac{bx+a}{2}))}{2} + \frac{i \operatorname{arctanh}(-1+id+d \cot(\frac{bx+a}{2}))d \ln(id-d \cot(\frac{bx+a}{2}))}{2} - \frac{i \left( \operatorname{dilog}\left(\frac{d \cot(\frac{bx+a}{2}) + id - 1}{d \cot(\frac{bx+a}{2}) - id - 1}\right) \right)}{d^2}$
default	$-\frac{i \operatorname{arctanh}(-1+id+d \cot(\frac{bx+a}{2}))d \ln(id+d \cot(\frac{bx+a}{2}))}{2} + \frac{i \operatorname{arctanh}(-1+id+d \cot(\frac{bx+a}{2}))d \ln(id-d \cot(\frac{bx+a}{2}))}{2} - \frac{i \left( \operatorname{dilog}\left(\frac{d \cot(\frac{bx+a}{2}) + id - 1}{d \cot(\frac{bx+a}{2}) - id - 1}\right) \right)}{d^2}$
risch	Expression too large to display

```
input int(-arctanh(-1+I*d+d*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/b/d*(-1/2*I*arctanh(-1+I*d+d*cot(b*x+a))*d*ln(I*d+d*cot(b*x+a))+1/2*I*a
rctanh(-1+I*d+d*cot(b*x+a))*d*ln(I*d-d*cot(b*x+a))-1/2*d^2*(-I/d*(1/2*dilo
g(I*(I*d-d*cot(b*x+a)-I*(2*I+2*d))/(2*I+2*d))+1/2*ln(I*d-d*cot(b*x+a))*ln(
I*(I*d-d*cot(b*x+a)-I*(2*I+2*d))/(2*I+2*d))-1/2*dilog(1/2*I*(-I*d-d*cot(b*
x+a))/d)-1/2*ln(I*d-d*cot(b*x+a))*ln(1/2*I*(-I*d-d*cot(b*x+a))/d))+I/d*(-1
/4*ln(I*d+d*cot(b*x+a))^2+1/2*(ln(I*d+d*cot(b*x+a))-ln(1/2*I*d+1/2*d*cot(b
*x+a)))*ln(1-1/2*I*d-1/2*d*cot(b*x+a))-1/2*dilog(1/2*I*d+1/2*d*cot(b*x+a))
))
```

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.30

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx$$

$$= \frac{2i b^2 x^2 - 2bx \log\left(-\frac{de^{(2i bx + 2i a)}}{(d+i)e^{(2i bx + 2i a)} - i}\right) - 2i a^2 - 2(bx + a) \log((id - 1)e^{(2i bx + 2i a)} + 1) + 2a \log\left(\frac{(d+i)e^{(2i bx + 2i a)}}{(d+i)e^{(2i bx + 2i a)} - i}\right)}{4b}$$

input

```
integrate(-arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```
1/4*(2*I*b^2*x^2 - 2*b*x*log(-d*e^(2*I*b*x + 2*I*a)/((d + I)*e^(2*I*b*x +
2*I*a) - I)) - 2*I*a^2 - 2*(b*x + a)*log((I*d - 1)*e^(2*I*b*x + 2*I*a) + 1
) + 2*a*log(((d + I)*e^(2*I*b*x + 2*I*a) - I)/(d + I)) + I*dilog(-(I*d - 1
)*e^(2*I*b*x + 2*I*a)))/b
```

**Sympy [F]**

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \int \operatorname{atanh}(d \cot(a + bx) + id - 1) dx$$

input

```
integrate(-atanh(-1+I*d+d*cot(b*x+a)),x)
```

output

```
-Integral(atanh(d*cot(a + b*x) + I*d - 1), x)
```



**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs.  $2(67) = 134$ .

Time = 0.12 (sec) , antiderivative size = 286, normalized size of antiderivative = 3.04

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx =$$

$$\frac{4(bx + a)d \left( \frac{\log((id-2)\tan(bx+a)+d)}{d} - \frac{\log(i \tan(bx+a)+1)}{d} \right) + d \left( -\frac{2i \left( \log((id-2)\tan(bx+a)+d) \log\left(\frac{(d+2i)\tan(bx+a)-id}{2id-2} + 1\right)}{d} \right)}{d} \right)}{d}$$

input `integrate(-arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="maxima")`

output

$$\begin{aligned} & -1/8*(4*(b*x + a)*d*(\log((I*d - 2)*\tan(b*x + a) + d)/d - \log(I*\tan(b*x + a) \\ & + 1)/d) + d*(-2*I*(\log((I*d - 2)*\tan(b*x + a) + d)*\log(((d + 2*I)*\tan(b*x \\ & + a) - I*d)/(2*I*d - 2) + 1) + \operatorname{dilog}(-((d + 2*I)*\tan(b*x + a) - I*d)/(2* \\ & I*d - 2)))/d - 2*I*(\log(-1/2*(d + 2*I)*\tan(b*x + a) + 1/2*I*d)*\log(I*\tan(b \\ & *x + a) + 1) + \operatorname{dilog}(1/2*(d + 2*I)*\tan(b*x + a) - 1/2*I*d + 1))/d + (2*I*1 \\ & \log((I*d - 2)*\tan(b*x + a) + d)*\log(I*\tan(b*x + a) + 1) - I*\log(I*\tan(b*x + \\ & a) + 1)^2)/d + 2*I*(\log(I*\tan(b*x + a) + 1)*\log(-1/2*I*\tan(b*x + a) + 1/2 \\ & ) + \operatorname{dilog}(1/2*I*\tan(b*x + a) + 1/2))/d) + 8*(b*x + a)*\operatorname{arctanh}(I*d + d/\tan(b*x + a) - 1))/b \end{aligned}$$
**Giac [F]**

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -\operatorname{artanh}(d \cot(bx + a) + id - 1) dx$$

input `integrate(-arctanh(-1+I*d+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-arctanh(d*cot(b*x + a) + I*d - 1), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = \int -\operatorname{atanh}(d \cot(a + bx) - 1 + di) dx$$

input `int(-atanh(d*1i + d*cot(a + b*x) - 1), x)`

output `int(-atanh(d*1i + d*cot(a + b*x) - 1), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(1 - id - d \cot(a + bx)) dx = - \left( \int \operatorname{atanh}(\cot(bx + a) d + di - 1) dx \right)$$

input `int(-atanh(-1+I*d+d*cot(b*x+a)), x)`

output `- int(atanh(cot(a + b*x)*d + d*i - 1), x)`

### 3.345 $\int \frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x} dx$

Optimal result	2430
Mathematica [N/A]	2430
Rubi [N/A]	2431
Maple [N/A]	2431
Fricas [N/A]	2432
Sympy [N/A]	2432
Maxima [N/A]	2432
Giac [N/A]	2433
Mupad [N/A]	2433
Reduce [N/A]	2434

#### Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x} dx = \operatorname{Int}\left(\frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x}, x\right)$$

output

```
Defer(Int)(-arctanh(-1+I*d+d*cot(b*x+a))/x,x)
```

#### Mathematica [N/A]

Not integrable

Time = 0.78 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x} dx = \int \frac{\operatorname{arctanh}(1-id-d \cot(a+bx))}{x} dx$$

input

```
Integrate[ArcTanh[1 - I*d - d*Cot[a + b*x]]/x,x]
```

output

```
Integrate[ArcTanh[1 - I*d - d*Cot[a + b*x]]/x, x]
```

**Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arctanh}(d(-\cot(a+bx)) - id + 1)}{x} dx$$

↓ 7299

$$\int \frac{\operatorname{arctanh}(d(-\cot(a+bx)) - id + 1)}{x} dx$$

input `Int[ArcTanh[1 - I*d - d*Cot[a + b*x]]/x,x]`

output `$Aborted`

**Maple [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int -\frac{\operatorname{arctanh}(-1 + id + d \cot(bx + a))}{x} dx$$

input `int(-arctanh(-1+I*d+d*cot(b*x+a))/x,x)`

output `int(-arctanh(-1+I*d+d*cot(b*x+a))/x,x)`

**Fricas [N/A]**

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \cot(bx + a) + id - 1)}{x} dx$$

input `integrate(-arctanh(-1+I*d+d*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(-1/2*log(-d*e^(2*I*b*x + 2*I*a)/((d + I)*e^(2*I*b*x + 2*I*a) - I))/x, x)`

**Sympy [N/A]**

Not integrable

Time = 1.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = -\int \frac{\operatorname{atanh}(d \cot(a + bx) + id - 1)}{x} dx$$

input `integrate(-atanh(-1+I*d+d*cot(b*x+a))/x,x)`

output `-Integral(atanh(d*cot(a + b*x) + I*d - 1)/x, x)`

**Maxima [N/A]**

Not integrable

Time = 4.42 (sec) , antiderivative size = 144, normalized size of antiderivative = 6.86

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \cot(bx + a) + id - 1)}{x} dx$$

input `integrate(-arctanh(-1+I*d+d*cot(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x + 1/4*(I*pi - 4*I*a - 2*log(d))*log(x) - 1/2*I*integrate(arctan2(-d
*cos(2*b*x + 2*a) + sin(2*b*x + 2*a), -d*sin(2*b*x + 2*a) - cos(2*b*x + 2*
a) + 1)/x, x) + 1/4*integrate(log((d^2 + 1)*cos(2*b*x + 2*a)^2 + (d^2 + 1)
*sin(2*b*x + 2*a)^2 - 2*d*sin(2*b*x + 2*a) - 2*cos(2*b*x + 2*a) + 1)/x, x)
```

**Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = \int -\frac{\operatorname{artanh}(d \cot(bx + a) + id - 1)}{x} dx$$

input

```
integrate(-arctanh(-1+I*d+d*cot(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(-arctanh(d*cot(b*x + a) + I*d - 1)/x, x)
```

**Mupad [N/A]**

Not integrable

Time = 3.92 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = \int -\frac{\operatorname{atanh}(d \cot(a + bx) - 1 + d li)}{x} dx$$

input

```
int(-atanh(d*1i + d*cot(a + b*x) - 1)/x,x)
```

output

```
int(-atanh(d*1i + d*cot(a + b*x) - 1)/x, x)
```

**Reduce [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\operatorname{arctanh}(1 - id - d \cot(a + bx))}{x} dx = - \left( \int \frac{\operatorname{atanh}(\cot(bx + a) d + di - 1)}{x} dx \right)$$

input `int(-atanh(-1+I*d+d*cot(b*x+a))/x,x)`output `- int(atanh(cot(a + b*x)*d + d*i - 1)/x,x)`

### 3.346 $\int \operatorname{arctanh}(e^x) dx$

Optimal result	2435
Mathematica [B] (verified)	2435
Rubi [A] (verified)	2436
Maple [A] (verified)	2437
Fricas [B] (verification not implemented)	2437
Sympy [F]	2438
Maxima [B] (verification not implemented)	2438
Giac [F]	2438
Mupad [F(-1)]	2439
Reduce [F]	2439

#### Optimal result

Integrand size = 4, antiderivative size = 21

$$\int \operatorname{arctanh}(e^x) dx = -\frac{\operatorname{PolyLog}(2, -e^x)}{2} + \frac{\operatorname{PolyLog}(2, e^x)}{2}$$

output `-1/2*polylog(2, -exp(x))+1/2*polylog(2, exp(x))`

#### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs.  $2(21) = 42$ .

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.43

$$\int \operatorname{arctanh}(e^x) dx = x \operatorname{arctanh}(e^x) + \frac{1}{2} x \log(1 - e^x) - \frac{1}{2} x \log(1 + e^x) - \frac{\operatorname{PolyLog}(2, -e^x)}{2} + \frac{\operatorname{PolyLog}(2, e^x)}{2}$$

input `Integrate[ArcTanh[E^x], x]`

output `x*ArcTanh[E^x] + (x*Log[1 - E^x])/2 - (x*Log[1 + E^x])/2 - PolyLog[2, -E^x]/2 + PolyLog[2, E^x]/2`



**Rubi [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {2720, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(e^x) dx$$

↓ 2720

$$\int e^{-x} \operatorname{arctanh}(e^x) de^x$$

↓ 6446

$$\frac{\operatorname{PolyLog}(2, e^x)}{2} - \frac{\operatorname{PolyLog}(2, -e^x)}{2}$$

input `Int[ArcTanh[E^x], x]`

output `-1/2*PolyLog[2, -E^x] + PolyLog[2, E^x]/2`

**Defintions of rubi rules used**

rule 2720 `Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 6446 `Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] :=> Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /; FreeQ[{a, b, c}, x]`

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

method	result	size
risch	$-\frac{\operatorname{dilog}(1+e^x)}{2} + \frac{\operatorname{dilog}(1-e^x)}{2}$	18
derivativedivides	$\ln(e^x) \operatorname{arctanh}(e^x) - \frac{\operatorname{dilog}(e^x)}{2} - \frac{\operatorname{dilog}(1+e^x)}{2} - \frac{\ln(e^x) \ln(1+e^x)}{2}$	31
default	$\ln(e^x) \operatorname{arctanh}(e^x) - \frac{\operatorname{dilog}(e^x)}{2} - \frac{\operatorname{dilog}(1+e^x)}{2} - \frac{\ln(e^x) \ln(1+e^x)}{2}$	31
parts	$x \operatorname{arctanh}(e^x) + \frac{x \ln(1-e^x)}{2} + \frac{\operatorname{polylog}(2, e^x)}{2} - \frac{x \ln(1+e^x)}{2} - \frac{\operatorname{polylog}(2, -e^x)}{2}$	39

input `int(arctanh(exp(x)), x, method=_RETURNVERBOSE)`

output `-1/2*dilog(1+exp(x))+1/2*dilog(1-exp(x))`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(13) = 26$ .

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 3.10

$$\int \operatorname{arctanh}(e^x) dx = \frac{1}{2} x \log \left( -\frac{\cosh(x) + \sinh(x) + 1}{\cosh(x) + \sinh(x) - 1} \right) - \frac{1}{2} x \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{2} x \log(-\cosh(x) - \sinh(x) + 1) + \frac{1}{2} \operatorname{Li}_2(\cosh(x) + \sinh(x)) - \frac{1}{2} \operatorname{Li}_2(-\cosh(x) - \sinh(x))$$

input `integrate(arctanh(exp(x)), x, algorithm="fricas")`

output `1/2*x*log(-(cosh(x) + sinh(x) + 1)/(cosh(x) + sinh(x) - 1)) - 1/2*x*log(cosh(x) + sinh(x) + 1) + 1/2*x*log(-cosh(x) - sinh(x) + 1) + 1/2*dilog(cosh(x) + sinh(x)) - 1/2*dilog(-cosh(x) - sinh(x))`

**Sympy [F]**

$$\int \operatorname{arctanh}(e^x) dx = \int \operatorname{atanh}(e^x) dx$$

input `integrate(atanh(exp(x)), x)`

output `Integral(atanh(exp(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 58 vs.  $2(13) = 26$ .

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.76

$$\begin{aligned} \int \operatorname{arctanh}(e^x) dx &= -\frac{1}{2} x (\log(e^x + 1) - \log(e^x - 1)) + x \operatorname{artanh}(e^x) \\ &\quad + \frac{1}{2} \log(-e^x) \log(e^x + 1) - \frac{1}{2} x \log(e^x - 1) \\ &\quad + \frac{1}{2} \operatorname{Li}_2(e^x + 1) - \frac{1}{2} \operatorname{Li}_2(-e^x + 1) \end{aligned}$$

input `integrate(arctanh(exp(x)), x, algorithm="maxima")`

output `-1/2*x*(log(e^x + 1) - log(e^x - 1)) + x*arctanh(e^x) + 1/2*log(-e^x)*log(e^x + 1) - 1/2*x*log(e^x - 1) + 1/2*dilog(e^x + 1) - 1/2*dilog(-e^x + 1)`

**Giac [F]**

$$\int \operatorname{arctanh}(e^x) dx = \int \operatorname{artanh}(e^x) dx$$

input `integrate(arctanh(exp(x)), x, algorithm="giac")`

output `integrate(arctanh(e^x), x)`

### Mupad [F(-1)]

Timed out.

$$\int \operatorname{arctanh}(e^x) dx = \int \operatorname{atanh}(e^x) dx$$

input `int(atanh(exp(x)), x)`

output `int(atanh(exp(x)), x)`

### Reduce [F]

$$\int \operatorname{arctanh}(e^x) dx = \int \operatorname{atanh}(e^x) dx$$

input `int(atanh(exp(x)), x)`

output `int(atanh(e**x), x)`

### 3.347 $\int x \operatorname{arctanh}(e^x) dx$

Optimal result	2440
Mathematica [A] (verified)	2440
Rubi [A] (verified)	2441
Maple [A] (verified)	2442
Fricas [B] (verification not implemented)	2443
Sympy [F]	2444
Maxima [B] (verification not implemented)	2444
Giac [F]	2444
Mupad [F(-1)]	2445
Reduce [F]	2445

#### Optimal result

Integrand size = 6, antiderivative size = 43

$$\int x \operatorname{arctanh}(e^x) dx = -\frac{1}{2}x \operatorname{PolyLog}(2, -e^x) + \frac{x \operatorname{PolyLog}(2, e^x)}{2} + \frac{\operatorname{PolyLog}(3, -e^x)}{2} - \frac{\operatorname{PolyLog}(3, e^x)}{2}$$

output

`-1/2*x*polylog(2,-exp(x))+1/2*x*polylog(2,exp(x))+1/2*polylog(3,-exp(x))-1/2*polylog(3,exp(x))`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.65

$$\int x \operatorname{arctanh}(e^x) dx = \frac{1}{4}(2x^2 \operatorname{arctanh}(e^x) + x^2 \log(1 - e^x) - x^2 \log(1 + e^x) - 2x \operatorname{PolyLog}(2, -e^x) + 2x \operatorname{PolyLog}(2, e^x) + 2 \operatorname{PolyLog}(3, -e^x) - 2 \operatorname{PolyLog}(3, e^x))$$

input

`Integrate[x*ArcTanh[E^x],x]`

output

$$(2x^2 \operatorname{ArcTanh}[E^x] + x^2 \operatorname{Log}[1 - E^x] - x^2 \operatorname{Log}[1 + E^x] - 2x \operatorname{PolyLog}[2, -E^x] + 2x \operatorname{PolyLog}[2, E^x] + 2 \operatorname{PolyLog}[3, -E^x] - 2 \operatorname{PolyLog}[3, E^x])/4$$
**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6767, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \operatorname{arctanh}(e^x) dx \\ & \quad \downarrow \text{6767} \\ & \frac{1}{2} \int x \log(1 + e^x) dx - \frac{1}{2} \int x \log(1 - e^x) dx \\ & \quad \downarrow \text{3011} \\ & \frac{1}{2} \left( \int \operatorname{PolyLog}(2, -e^x) dx - x \operatorname{PolyLog}(2, -e^x) \right) + \\ & \quad \frac{1}{2} \left( x \operatorname{PolyLog}(2, e^x) - \int \operatorname{PolyLog}(2, e^x) dx \right) \\ & \quad \downarrow \text{2720} \\ & \frac{1}{2} \left( \int e^{-x} \operatorname{PolyLog}(2, -e^x) dx - x \operatorname{PolyLog}(2, -e^x) \right) + \\ & \quad \frac{1}{2} \left( x \operatorname{PolyLog}(2, e^x) - \int e^{-x} \operatorname{PolyLog}(2, e^x) dx \right) \\ & \quad \downarrow \text{7143} \\ & \frac{1}{2} (\operatorname{PolyLog}(3, -e^x) - x \operatorname{PolyLog}(2, -e^x)) + \frac{1}{2} (x \operatorname{PolyLog}(2, e^x) - \operatorname{PolyLog}(3, e^x)) \end{aligned}$$

input

$$\operatorname{Int}[x \operatorname{ArcTanh}[E^x], x]$$

output  $(-x \operatorname{PolyLog}[2, -E^x] + \operatorname{PolyLog}[3, -E^x])/2 + (x \operatorname{PolyLog}[2, E^x] - \operatorname{PolyLog}[3, E^x])/2$

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6767 `Int[ArcTanh[(a_.) + (b_.)*(f_)^(c_.) + (d_.)*(x_)]*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[x^m*Log[1 + a + b*f^(c + d*x)], x], x] - Simp[1/2 Int[x^m*Log[1 - a - b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

method	result
risch	$-\frac{x \operatorname{polylog}(2, -e^x)}{2} + \frac{x \operatorname{polylog}(2, e^x)}{2} + \frac{\operatorname{polylog}(3, -e^x)}{2} - \frac{\operatorname{polylog}(3, e^x)}{2}$
default	$\frac{x^2 \operatorname{arctanh}(e^x)}{2} + \frac{x^2 \ln(1 - e^x)}{4} + \frac{x \operatorname{polylog}(2, e^x)}{2} - \frac{\operatorname{polylog}(3, e^x)}{2} - \frac{x^2 \ln(1 + e^x)}{4} - \frac{x \operatorname{polylog}(2, -e^x)}{2} + \frac{\operatorname{polylog}(3, -e^x)}{2}$
parts	$\frac{x^2 \operatorname{arctanh}(e^x)}{2} + \frac{x^2 \ln(1 - e^x)}{4} + \frac{x \operatorname{polylog}(2, e^x)}{2} - \frac{\operatorname{polylog}(3, e^x)}{2} - \frac{x^2 \ln(1 + e^x)}{4} - \frac{x \operatorname{polylog}(2, -e^x)}{2} + \frac{\operatorname{polylog}(3, -e^x)}{2}$

input `int(x*arctanh(exp(x)),x,method=_RETURNVERBOSE)`

output `-1/2*x*polylog(2,-exp(x))+1/2*x*polylog(2,exp(x))+1/2*polylog(3,-exp(x))-1/2*polylog(3,exp(x))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs.  $2(29) = 58$ .

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.21

$$\int x \operatorname{arctanh}(e^x) dx = \frac{1}{4} x^2 \log \left( -\frac{\cosh(x) + \sinh(x) + 1}{\cosh(x) + \sinh(x) - 1} \right) - \frac{1}{4} x^2 \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{4} x^2 \log(-\cosh(x) - \sinh(x) + 1) + \frac{1}{2} x \operatorname{Li}_2(\cosh(x) + \sinh(x)) - \frac{1}{2} x \operatorname{Li}_2(-\cosh(x) - \sinh(x)) - \frac{1}{2} \operatorname{polylog}(3, \cosh(x) + \sinh(x)) + \frac{1}{2} \operatorname{polylog}(3, -\cosh(x) - \sinh(x))$$

input `integrate(x*arctanh(exp(x)),x, algorithm="fricas")`

output `1/4*x^2*log(-(cosh(x) + sinh(x) + 1)/(cosh(x) + sinh(x) - 1)) - 1/4*x^2*log(cosh(x) + sinh(x) + 1) + 1/4*x^2*log(-cosh(x) - sinh(x) + 1) + 1/2*x*dilog(cosh(x) + sinh(x)) - 1/2*x*dilog(-cosh(x) - sinh(x)) - 1/2*polylog(3, cosh(x) + sinh(x)) + 1/2*polylog(3, -cosh(x) - sinh(x))`



**Sympy [F]**

$$\int x \operatorname{arctanh}(e^x) dx = \int x \operatorname{atanh}(e^x) dx$$

input `integrate(x*atanh(exp(x)),x)`

output `Integral(x*atanh(exp(x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs.  $2(29) = 58$ .

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.37

$$\begin{aligned} \int x \operatorname{arctanh}(e^x) dx = & \frac{1}{2} x^2 \operatorname{artanh}(e^x) - \frac{1}{4} x^2 \log(e^x + 1) + \frac{1}{4} x^2 \log(-e^x + 1) \\ & - \frac{1}{2} x \operatorname{Li}_2(-e^x) + \frac{1}{2} x \operatorname{Li}_2(e^x) + \frac{1}{2} \operatorname{Li}_3(-e^x) - \frac{1}{2} \operatorname{Li}_3(e^x) \end{aligned}$$

input `integrate(x*arctanh(exp(x)),x, algorithm="maxima")`

output `1/2*x^2*arctanh(e^x) - 1/4*x^2*log(e^x + 1) + 1/4*x^2*log(-e^x + 1) - 1/2*x*dilog(-e^x) + 1/2*x*dilog(e^x) + 1/2*polylog(3, -e^x) - 1/2*polylog(3, e^x)`

**Giac [F]**

$$\int x \operatorname{arctanh}(e^x) dx = \int x \operatorname{artanh}(e^x) dx$$

input `integrate(x*arctanh(exp(x)),x, algorithm="giac")`

output `integrate(x*arctanh(e^x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(e^x) dx = \int x \operatorname{atanh}(e^x) dx$$

input `int(x*atanh(exp(x)), x)`output `int(x*atanh(exp(x)), x)`**Reduce [F]**

$$\int x \operatorname{arctanh}(e^x) dx = \int \operatorname{atanh}(e^x) x dx$$

input `int(x*atanh(exp(x)), x)`output `int(atanh(e**x)*x, x)`

### 3.348 $\int x^2 \operatorname{arctanh}(e^x) dx$

Optimal result	2446
Mathematica [A] (verified)	2446
Rubi [A] (verified)	2447
Maple [A] (verified)	2449
Fricas [B] (verification not implemented)	2449
Sympy [F]	2450
Maxima [A] (verification not implemented)	2450
Giac [F]	2451
Mupad [F(-1)]	2451
Reduce [F]	2451

#### Optimal result

Integrand size = 8, antiderivative size = 58

$$\int x^2 \operatorname{arctanh}(e^x) dx = -\frac{1}{2}x^2 \operatorname{PolyLog}(2, -e^x) + \frac{1}{2}x^2 \operatorname{PolyLog}(2, e^x) + x \operatorname{PolyLog}(3, -e^x) - x \operatorname{PolyLog}(3, e^x) - \operatorname{PolyLog}(4, -e^x) + \operatorname{PolyLog}(4, e^x)$$

output

```
-1/2*x^2*polylog(2,-exp(x))+1/2*x^2*polylog(2,exp(x))+x*polylog(3,-exp(x))
-x*polylog(3,exp(x))-polylog(4,-exp(x))+polylog(4,exp(x))
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.60

$$\int x^2 \operatorname{arctanh}(e^x) dx = \frac{1}{6}(2x^3 \operatorname{arctanh}(e^x) + x^3 \log(1 - e^x) - x^3 \log(1 + e^x) - 3x^2 \operatorname{PolyLog}(2, -e^x) + 3x^2 \operatorname{PolyLog}(2, e^x) + 6x \operatorname{PolyLog}(3, -e^x) - 6x \operatorname{PolyLog}(3, e^x) - 6 \operatorname{PolyLog}(4, -e^x) + 6 \operatorname{PolyLog}(4, e^x))$$

input

```
Integrate[x^2*ArcTanh[E^x],x]
```

output

$$(2x^3 \operatorname{ArcTanh}[E^x] + x^3 \operatorname{Log}[1 - E^x] - x^3 \operatorname{Log}[1 + E^x] - 3x^2 \operatorname{PolyLog}[2, -E^x] + 3x^2 \operatorname{PolyLog}[2, E^x] + 6x \operatorname{PolyLog}[3, -E^x] - 6x \operatorname{PolyLog}[3, E^x] - 6 \operatorname{PolyLog}[4, -E^x] + 6 \operatorname{PolyLog}[4, E^x]) / 6$$
**Rubi [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.21, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {6767, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(e^x) dx$$

$$\downarrow 6767$$

$$\frac{1}{2} \int x^2 \log(1 + e^x) dx - \frac{1}{2} \int x^2 \log(1 - e^x) dx$$

$$\downarrow 3011$$

$$\frac{1}{2} \left( 2 \int x \operatorname{PolyLog}(2, -e^x) dx - x^2 \operatorname{PolyLog}(2, -e^x) \right) + \frac{1}{2} \left( x^2 \operatorname{PolyLog}(2, e^x) - 2 \int x \operatorname{PolyLog}(2, e^x) dx \right)$$

$$\downarrow 7163$$

$$\frac{1}{2} \left( 2 \left( x \operatorname{PolyLog}(3, -e^x) - \int \operatorname{PolyLog}(3, -e^x) dx \right) - x^2 \operatorname{PolyLog}(2, -e^x) \right) + \frac{1}{2} \left( x^2 \operatorname{PolyLog}(2, e^x) - 2 \left( x \operatorname{PolyLog}(3, e^x) - \int \operatorname{PolyLog}(3, e^x) dx \right) \right)$$

$$\downarrow 2720$$

$$\frac{1}{2} \left( 2 \left( x \operatorname{PolyLog}(3, -e^x) - \int e^{-x} \operatorname{PolyLog}(3, -e^x) de^x \right) - x^2 \operatorname{PolyLog}(2, -e^x) \right) + \frac{1}{2} \left( x^2 \operatorname{PolyLog}(2, e^x) - 2 \left( x \operatorname{PolyLog}(3, e^x) - \int e^{-x} \operatorname{PolyLog}(3, e^x) de^x \right) \right)$$

$$\downarrow 7143$$

$$\frac{1}{2}(2(x \operatorname{PolyLog}(3, -e^x) - \operatorname{PolyLog}(4, -e^x)) - x^2 \operatorname{PolyLog}(2, -e^x)) + \frac{1}{2}(x^2 \operatorname{PolyLog}(2, e^x) - 2(x \operatorname{PolyLog}(3, e^x) - \operatorname{PolyLog}(4, e^x)))$$

input `Int[x^2*ArcTanh[E^x], x]`

output `(-(x^2*PolyLog[2, -E^x]) + 2*(x*PolyLog[3, -E^x] - PolyLog[4, -E^x]))/2 + (x^2*PolyLog[2, E^x] - 2*(x*PolyLog[3, E^x] - PolyLog[4, E^x]))/2`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6767 `Int[ArcTanh[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[x^m*Log[1 + a + b*f^(c + d*x)], x], x] - Simp[1/2 Int[x^m*Log[1 - a - b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^(m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

**Maple [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.84

method	result
risch	$-\frac{x^2 \operatorname{polylog}(2, -e^x)}{2} + \frac{x^2 \operatorname{polylog}(2, e^x)}{2} + x \operatorname{polylog}(3, -e^x) - x \operatorname{polylog}(3, e^x) - \operatorname{polylog}(4, -e^x) + \operatorname{polylog}(4, e^x)$
default	$\frac{x^3 \operatorname{arctanh}(e^x)}{3} + \frac{x^3 \ln(1-e^x)}{6} + \frac{x^2 \operatorname{polylog}(2, e^x)}{2} - x \operatorname{polylog}(3, e^x) + \operatorname{polylog}(4, e^x) - \frac{x^3 \ln(1+e^x)}{6} - \frac{x^2 \operatorname{polylog}(2, -e^x)}{2}$
parts	$\frac{x^3 \operatorname{arctanh}(e^x)}{3} + \frac{x^3 \ln(1-e^x)}{6} + \frac{x^2 \operatorname{polylog}(2, e^x)}{2} - x \operatorname{polylog}(3, e^x) + \operatorname{polylog}(4, e^x) - \frac{x^3 \ln(1+e^x)}{6} - \frac{x^2 \operatorname{polylog}(2, -e^x)}{2}$

input

```
int(x^2*arctanh(exp(x)), x, method=_RETURNVERBOSE)
```

output

```
-1/2*x^2*polylog(2, -exp(x))+1/2*x^2*polylog(2, exp(x))+x*polylog(3, -exp(x))
-x*polylog(3, exp(x))-polylog(4, -exp(x))+polylog(4, exp(x))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(46) = 92.

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.07

$$\int x^2 \operatorname{arctanh}(e^x) dx = \frac{1}{6} x^3 \log \left( -\frac{\cosh(x) + \sinh(x) + 1}{\cosh(x) + \sinh(x) - 1} \right) - \frac{1}{6} x^3 \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{6} x^3 \log(-\cosh(x) - \sinh(x) + 1) + \frac{1}{2} x^2 \operatorname{Li}_2(\cosh(x) + \sinh(x)) - \frac{1}{2} x^2 \operatorname{Li}_2(-\cosh(x) - \sinh(x)) - x \operatorname{polylog}(3, \cosh(x) + \sinh(x)) + x \operatorname{polylog}(3, -\cosh(x) - \sinh(x)) + \operatorname{polylog}(4, \cosh(x) + \sinh(x)) - \operatorname{polylog}(4, -\cosh(x) - \sinh(x))$$

input `integrate(x^2*arctanh(exp(x)),x, algorithm="fricas")`

output `1/6*x^3*log(-(cosh(x) + sinh(x) + 1)/(cosh(x) + sinh(x) - 1)) - 1/6*x^3*log(cosh(x) + sinh(x) + 1) + 1/6*x^3*log(-cosh(x) - sinh(x) + 1) + 1/2*x^2*dilog(cosh(x) + sinh(x)) - 1/2*x^2*dilog(-cosh(x) - sinh(x)) - x*polylog(3, cosh(x) + sinh(x)) + x*polylog(3, -cosh(x) - sinh(x)) + polylog(4, cosh(x) + sinh(x)) - polylog(4, -cosh(x) - sinh(x))`

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(e^x) dx = \int x^2 \operatorname{atanh}(e^x) dx$$

input `integrate(x**2*atanh(exp(x)),x)`

output `Integral(x**2*atanh(exp(x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.31

$$\begin{aligned} \int x^2 \operatorname{arctanh}(e^x) dx &= \frac{1}{3} x^3 \operatorname{artanh}(e^x) - \frac{1}{6} x^3 \log(e^x + 1) \\ &+ \frac{1}{6} x^3 \log(-e^x + 1) - \frac{1}{2} x^2 \operatorname{Li}_2(-e^x) + \frac{1}{2} x^2 \operatorname{Li}_2(e^x) \\ &+ x \operatorname{Li}_3(-e^x) - x \operatorname{Li}_3(e^x) - \operatorname{Li}_4(-e^x) + \operatorname{Li}_4(e^x) \end{aligned}$$

input `integrate(x^2*arctanh(exp(x)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(e^x) - 1/6*x^3*log(e^x + 1) + 1/6*x^3*log(-e^x + 1) - 1/2*x^2*dilog(-e^x) + 1/2*x^2*dilog(e^x) + x*polylog(3, -e^x) - x*polylog(3, e^x) - polylog(4, -e^x) + polylog(4, e^x)`

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(e^x) dx = \int x^2 \operatorname{artanh}(e^x) dx$$

input `integrate(x^2*arctanh(exp(x)),x, algorithm="giac")`

output `integrate(x^2*arctanh(e^x), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(e^x) dx = \int x^2 \operatorname{atanh}(e^x) dx$$

input `int(x^2*atanh(exp(x)),x)`

output `int(x^2*atanh(exp(x)), x)`

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(e^x) dx = \int \operatorname{atanh}(e^x) x^2 dx$$

input `int(x^2*atanh(exp(x)),x)`

output `int(atanh(e**x)*x**2,x)`



### 3.349 $\int \operatorname{arctanh}(e^{a+bx}) dx$

Optimal result	2452
Mathematica [A] (verified)	2452
Rubi [A] (verified)	2453
Maple [A] (verified)	2454
Fricas [B] (verification not implemented)	2454
Sympy [F]	2455
Maxima [B] (verification not implemented)	2455
Giac [F]	2456
Mupad [F(-1)]	2456
Reduce [F]	2456

#### Optimal result

Integrand size = 8, antiderivative size = 35

$$\int \operatorname{arctanh}(e^{a+bx}) dx = -\frac{\operatorname{PolyLog}(2, -e^{a+bx})}{2b} + \frac{\operatorname{PolyLog}(2, e^{a+bx})}{2b}$$

output `-1/2*polylog(2, -exp(b*x+a))/b+1/2*polylog(2, exp(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.94

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \frac{bx(2\operatorname{arctanh}(e^{a+bx}) + \log(1 - e^{a+bx}) - \log(1 + e^{a+bx})) - \operatorname{PolyLog}(2, -e^{a+bx}) + \operatorname{PolyLog}(2, e^{a+bx})}{2b}$$

input `Integrate[ArcTanh[E^(a + b*x)], x]`

output `(b*x*(2*ArcTanh[E^(a + b*x)] + Log[1 - E^(a + b*x)] - Log[1 + E^(a + b*x)]) - PolyLog[2, -E^(a + b*x)] + PolyLog[2, E^(a + b*x)])/(2*b)`

**Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {2720, 6446}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(e^{a+bx}) dx$$

$$\downarrow \text{2720}$$

$$\frac{\int e^{-a-bx} \operatorname{arctanh}(e^{a+bx}) de^{a+bx}}{b}$$

$$\downarrow \text{6446}$$

$$\frac{\frac{1}{2} \operatorname{PolyLog}(2, e^{a+bx}) - \frac{1}{2} \operatorname{PolyLog}(2, -e^{a+bx})}{b}$$

input

```
Int[ArcTanh[E^(a + b*x)], x]
```

output

```
(-1/2*PolyLog[2, -E^(a + b*x)] + PolyLog[2, E^(a + b*x)]/2)/b
```

**Defintions of rubi rules used**

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 6446

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x] + (-Simp[(b/2)*PolyLog[2, (-c)*x], x] + Simp[(b/2)*PolyLog[2, c*x], x]) /; FreeQ[{a, b, c}, x]
```

**Maple [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result
risch	$-\frac{\operatorname{dilog}(e^{bx+a}+1)}{2b} + \frac{\operatorname{dilog}(1-e^{bx+a})}{2b}$
derivativedivides	$\frac{\ln(e^{bx+a}) \operatorname{arctanh}(e^{bx+a}) - \frac{\operatorname{dilog}(e^{bx+a})}{2} - \frac{\operatorname{dilog}(e^{bx+a}+1)}{2} - \frac{\ln(e^{bx+a}) \ln(e^{bx+a}+1)}{2}}{b}$
default	$\frac{\ln(e^{bx+a}) \operatorname{arctanh}(e^{bx+a}) - \frac{\operatorname{dilog}(e^{bx+a})}{2} - \frac{\operatorname{dilog}(e^{bx+a}+1)}{2} - \frac{\ln(e^{bx+a}) \ln(e^{bx+a}+1)}{2}}{b}$
parts	$x \operatorname{arctanh}(e^{bx+a}) - \frac{(bx+a) \ln(1-e^{bx+a})}{2} - \frac{\operatorname{polylog}(2, e^{bx+a})}{2} + \frac{(bx+a) \ln(e^{bx+a}+1)}{2} + \frac{\operatorname{polylog}(2, -e^{bx+a})}{2} - a \operatorname{arctanh}(e^{bx+a})$

input `int(arctanh(exp(b*x+a)),x,method=_RETURNVERBOSE)`

output `-1/2/b*dilog(exp(b*x+a)+1)+1/2/b*dilog(1-exp(b*x+a))`

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 138 vs.  $2(27) = 54$ .

Time = 0.11 (sec) , antiderivative size = 138, normalized size of antiderivative = 3.94

$$\int \operatorname{arctanh}(e^{a+bx}) dx$$

$$= \frac{bx \log\left(-\frac{\cosh(bx+a)+\sinh(bx+a)+1}{\cosh(bx+a)+\sinh(bx+a)-1}\right) - bx \log(\cosh(bx+a) + \sinh(bx+a) + 1) - a \log(\cosh(bx+a) + \sinh(bx+a) + 1)}{b}$$

input `integrate(arctanh(exp(b*x+a)),x, algorithm="fricas")`

output `1/2*(b*x*log(-(cosh(b*x + a) + sinh(b*x + a) + 1)/(cosh(b*x + a) + sinh(b*x + a) - 1)) - b*x*log(cosh(b*x + a) + sinh(b*x + a) + 1) - a*log(cosh(b*x + a) + sinh(b*x + a) - 1) + (b*x + a)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) + dilog(cosh(b*x + a) + sinh(b*x + a)) - dilog(-cosh(b*x + a) - sinh(b*x + a)))/b`

**Sympy [F]**

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{atanh}(e^{a+bx}) dx$$

input `integrate(atanh(exp(b*x+a)),x)`

output `Integral(atanh(exp(a + b*x)), x)`

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 107 vs.  $2(27) = 54$ .

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 3.06

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \frac{(bx+a) \operatorname{artanh}(e^{(bx+a)})}{b} - \frac{(bx+a)(\log(e^{(bx+a)}+1) - \log(e^{(bx+a)}-1)) - \log(-e^{(bx+a)}) \log(e^{(bx+a)}+1) + (bx+a) \log(e^{(bx+a)})}{2b}$$

input `integrate(arctanh(exp(b*x+a)),x, algorithm="maxima")`

output `(b*x + a)*arctanh(e^(b*x + a))/b - 1/2*((b*x + a)*(log(e^(b*x + a) + 1) - log(e^(b*x + a) - 1)) - log(-e^(b*x + a))*log(e^(b*x + a) + 1) + (b*x + a)*log(e^(b*x + a) - 1) - dilog(e^(b*x + a) + 1) + dilog(-e^(b*x + a) + 1))/b`

**Giac [F]**

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{artanh}(e^{(bx+a)}) dx$$

input `integrate(arctanh(exp(b*x+a)),x, algorithm="giac")`

output `integrate(arctanh(e^(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{atanh}(e^{a+bx}) dx$$

input `int(atanh(exp(a + b*x)),x)`

output `int(atanh(exp(a + b*x)), x)`

**Reduce [F]**

$$\int \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{atanh}(e^{bx+a}) dx$$

input `int(atanh(exp(b*x+a)),x)`

output `int(atanh(e**(a + b*x)),x)`

### 3.350 $\int x \operatorname{arctanh}(e^{a+bx}) dx$

Optimal result	2457
Mathematica [A] (verified)	2457
Rubi [A] (verified)	2458
Maple [B] (verified)	2460
Fricas [B] (verification not implemented)	2460
Sympy [F]	2461
Maxima [A] (verification not implemented)	2461
Giac [F]	2462
Mupad [F(-1)]	2462
Reduce [F]	2462

#### Optimal result

Integrand size = 10, antiderivative size = 71

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = -\frac{x \operatorname{PolyLog}(2, -e^{a+bx})}{2b} + \frac{x \operatorname{PolyLog}(2, e^{a+bx})}{2b} + \frac{\operatorname{PolyLog}(3, -e^{a+bx})}{2b^2} - \frac{\operatorname{PolyLog}(3, e^{a+bx})}{2b^2}$$

output

```
-1/2*x*polylog(2,-exp(b*x+a))/b+1/2*x*polylog(2,exp(b*x+a))/b+1/2*polylog(3,-exp(b*x+a))/b^2-1/2*polylog(3,exp(b*x+a))/b^2
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.59

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \frac{2b^2 x^2 \operatorname{arctanh}(e^{a+bx}) + b^2 x^2 \log(1 - e^{a+bx}) - b^2 x^2 \log(1 + e^{a+bx}) - 2bx \operatorname{PolyLog}(2, -e^{a+bx}) + 2bx \operatorname{PolyLog}(2, e^{a+bx})}{4b^2}$$

input

```
Integrate[x*ArcTanh[E^(a + b*x)],x]
```

output

```
(2*b^2*x^2*ArcTanh[E^(a + b*x)] + b^2*x^2*Log[1 - E^(a + b*x)] - b^2*x^2*Log[1 + E^(a + b*x)] - 2*b*x*PolyLog[2, -E^(a + b*x)] + 2*b*x*PolyLog[2, E^(a + b*x)] + 2*PolyLog[3, -E^(a + b*x)] - 2*PolyLog[3, E^(a + b*x)])/(4*b^2)
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6767, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \operatorname{arctanh}(e^{a+bx}) dx \\
 & \quad \downarrow \text{6767} \\
 & \frac{1}{2} \int x \log(1 + e^{a+bx}) dx - \frac{1}{2} \int x \log(1 - e^{a+bx}) dx \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2} \left( \frac{\int \operatorname{PolyLog}(2, -e^{a+bx}) dx}{b} - \frac{x \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) + \\
 & \quad \frac{1}{2} \left( \frac{x \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{\int \operatorname{PolyLog}(2, e^{a+bx}) dx}{b} \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2} \left( \frac{\int e^{-a-bx} \operatorname{PolyLog}(2, -e^{a+bx}) de^{a+bx}}{b^2} - \frac{x \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) + \\
 & \quad \frac{1}{2} \left( \frac{x \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{\int e^{-a-bx} \operatorname{PolyLog}(2, e^{a+bx}) de^{a+bx}}{b^2} \right) \\
 & \quad \downarrow \text{7143} \\
 & \frac{1}{2} \left( \frac{\operatorname{PolyLog}(3, -e^{a+bx})}{b^2} - \frac{x \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) + \\
 & \quad \frac{1}{2} \left( \frac{x \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{\operatorname{PolyLog}(3, e^{a+bx})}{b^2} \right)
 \end{aligned}$$

input `Int[x*ArcTanh[E^(a + b*x)],x]`

output `((-(x*PolyLog[2, -E^(a + b*x)])/b) + PolyLog[3, -E^(a + b*x)]/b^2)/2 + ((x*PolyLog[2, E^(a + b*x)])/b - PolyLog[3, E^(a + b*x)]/b^2)/2`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6767 `Int[ArcTanh[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[x^m*Log[1 + a + b*f^(c + d*x)], x], x] - Simp[1/2 Int[x^m*Log[1 - a - b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`



**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 154 vs.  $2(59) = 118$ .

Time = 0.38 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.18

method	result
risch	$\frac{\ln(1-e^{bx+a})ax}{2b} + \frac{x \operatorname{polylog}(2, e^{bx+a})}{2b} + \frac{a^2 \ln(1-e^{bx+a})}{2b^2} + \frac{\operatorname{polylog}(2, e^{bx+a})a}{2b^2} + \frac{a \operatorname{dilog}(e^{bx+a})}{2b^2} - \frac{\operatorname{polylog}(3, e^{bx+a})}{2b^2} - \dots$
default	$\frac{x^2 \operatorname{arctanh}(e^{bx+a})}{2} - \frac{a^2 \operatorname{arctanh}(e^{bx+a})}{2} - \frac{(bx+a)^2 \ln(1-e^{bx+a})}{2} - (bx+a) \operatorname{polylog}(2, e^{bx+a}) + \operatorname{polylog}(3, e^{bx+a}) + \frac{(bx+a)^2 \ln(e^{bx+a})}{2}$
parts	$\frac{x^2 \operatorname{arctanh}(e^{bx+a})}{2} - \frac{a^2 \operatorname{arctanh}(e^{bx+a})}{2} - \frac{(bx+a)^2 \ln(1-e^{bx+a})}{2} - (bx+a) \operatorname{polylog}(2, e^{bx+a}) + \operatorname{polylog}(3, e^{bx+a}) + \frac{(bx+a)^2 \ln(e^{bx+a})}{2}$

input

```
int(x*arctanh(exp(b*x+a)), x, method=_RETURNVERBOSE)
```

output

```
1/2/b*ln(1-exp(b*x+a))*a*x+1/2*x*polylog(2, exp(b*x+a))/b+1/2/b^2*a^2*ln(1-
exp(b*x+a))+1/2/b^2*polylog(2, exp(b*x+a))*a+1/2/b^2*a*dilog(exp(b*x+a))-1/
2*polylog(3, exp(b*x+a))/b^2-1/2*x*polylog(2, -exp(b*x+a))/b+1/2/b^2*dilog(e
xp(b*x+a)+1)*a-1/2/b^2*polylog(2, -exp(b*x+a))*a+1/2*polylog(3, -exp(b*x+a))
/b^2
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 199 vs.  $2(57) = 114$ .

Time = 0.09 (sec) , antiderivative size = 199, normalized size of antiderivative = 2.80

$$\int x \operatorname{arctanh}(e^{a+bx}) dx$$

$$= \frac{b^2 x^2 \log\left(-\frac{\cosh(bx+a)+\sinh(bx+a)+1}{\cosh(bx+a)+\sinh(bx+a)-1}\right) - b^2 x^2 \log(\cosh(bx+a) + \sinh(bx+a) + 1) + 2bx \operatorname{Li}_2(\cosh(bx+a))}{b^2}$$

input

```
integrate(x*arctanh(exp(b*x+a)), x, algorithm="fricas")
```

output

```
1/4*(b^2*x^2*log(-cosh(b*x + a) + sinh(b*x + a) + 1)/(cosh(b*x + a) + sin
h(b*x + a) - 1)) - b^2*x^2*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 2*b*x*
dilog(cosh(b*x + a) + sinh(b*x + a)) - 2*b*x*dilog(-cosh(b*x + a) - sinh(b
*x + a)) + a^2*log(cosh(b*x + a) + sinh(b*x + a) - 1) + (b^2*x^2 - a^2)*lo
g(-cosh(b*x + a) - sinh(b*x + a) + 1) - 2*polylog(3, cosh(b*x + a) + sinh(
b*x + a)) + 2*polylog(3, -cosh(b*x + a) - sinh(b*x + a))/b^2
```

## Sympy [F]

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \int x \operatorname{atanh}(e^a e^{bx}) dx$$

input

```
integrate(x*atanh(exp(b*x+a)),x)
```

output

```
Integral(x*atanh(exp(a)*exp(b*x)), x)
```

## Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.52

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \frac{1}{2} x^2 \operatorname{artanh}(e^{(bx+a)}) - \frac{1}{4} b \left( \frac{b^2 x^2 \log(e^{(bx+a)} + 1) + 2 b x \operatorname{Li}_2(-e^{(bx+a)}) - 2 \operatorname{Li}_3(-e^{(bx+a)})}{b^3} - \frac{b^2 x^2 \log(-e^{(bx+a)} + 1) + 2 b x \operatorname{Li}_2(-e^{(bx+a)})}{b^3} \right)$$

input

```
integrate(x*arctanh(exp(b*x+a)),x, algorithm="maxima")
```

output

```
1/2*x^2*arctanh(e^(b*x + a)) - 1/4*b*((b^2*x^2*log(e^(b*x + a) + 1) + 2*b*
x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a)))/b^3 - (b^2*x^2*log(-e
^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))/b^3
)
```

**Giac [F]**

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \int x \operatorname{artanh}(e^{(bx+a)}) dx$$

input `integrate(x*arctanh(exp(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctanh(e^(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \int x \operatorname{atanh}(e^{a+bx}) dx$$

input `int(x*atanh(exp(a + b*x)),x)`

output `int(x*atanh(exp(a + b*x)), x)`

**Reduce [F]**

$$\int x \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{atanh}(e^{bx+a}) x dx$$

input `int(x*atanh(exp(b*x+a)),x)`

output `int(atanh(e**(a + b*x))*x,x)`

### 3.351 $\int x^2 \operatorname{arctanh}(e^{a+bx}) dx$

Optimal result	2463
Mathematica [A] (verified)	2463
Rubi [A] (verified)	2464
Maple [B] (verified)	2466
Fricas [B] (verification not implemented)	2467
Sympy [F]	2467
Maxima [A] (verification not implemented)	2468
Giac [F]	2468
Mupad [F(-1)]	2469
Reduce [F]	2469

#### Optimal result

Integrand size = 12, antiderivative size = 101

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = -\frac{x^2 \operatorname{PolyLog}(2, -e^{a+bx})}{2b} + \frac{x^2 \operatorname{PolyLog}(2, e^{a+bx})}{2b} + \frac{x \operatorname{PolyLog}(3, -e^{a+bx})}{b^2} - \frac{x \operatorname{PolyLog}(3, e^{a+bx})}{b^2} - \frac{\operatorname{PolyLog}(4, -e^{a+bx})}{b^3} + \frac{\operatorname{PolyLog}(4, e^{a+bx})}{b^3}$$

output

```
-1/2*x^2*polylog(2,-exp(b*x+a))/b+1/2*x^2*polylog(2,exp(b*x+a))/b+x*polylog(3,-exp(b*x+a))/b^2-x*polylog(3,exp(b*x+a))/b^2-polylog(4,-exp(b*x+a))/b^3+polylog(4,exp(b*x+a))/b^3
```

#### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.48

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \frac{2b^3 x^3 \operatorname{arctanh}(e^{a+bx}) + b^3 x^3 \log(1 - e^{a+bx}) - b^3 x^3 \log(1 + e^{a+bx}) - 3b^2 x^2 \operatorname{PolyLog}(2, -e^{a+bx}) + 3b^2 x^2 \operatorname{PolyLog}(2, e^{a+bx})}{b^3}$$

input `Integrate[x^2*ArcTanh[E^(a + b*x)],x]`

output  $(2*b^3*x^3*ArcTanh[E^(a + b*x)] + b^3*x^3*Log[1 - E^(a + b*x)] - b^3*x^3*Log[1 + E^(a + b*x)] - 3*b^2*x^2*PolyLog[2, -E^(a + b*x)] + 3*b^2*x^2*PolyLog[2, E^(a + b*x)] + 6*b*x*PolyLog[3, -E^(a + b*x)] - 6*b*x*PolyLog[3, E^(a + b*x)] - 6*PolyLog[4, -E^(a + b*x)] + 6*PolyLog[4, E^(a + b*x)])/(6*b^3)$

### Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6767, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \operatorname{arctanh}(e^{a+bx}) \, dx \\
 & \quad \downarrow \text{6767} \\
 & \frac{1}{2} \int x^2 \log(1 + e^{a+bx}) \, dx - \frac{1}{2} \int x^2 \log(1 - e^{a+bx}) \, dx \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2} \left( \frac{2 \int x \operatorname{PolyLog}(2, -e^{a+bx}) \, dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) + \\
 & \quad \frac{1}{2} \left( \frac{x^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{2 \int x \operatorname{PolyLog}(2, e^{a+bx}) \, dx}{b} \right) \\
 & \quad \downarrow \text{7163} \\
 & \frac{1}{2} \left( \frac{2 \left( \frac{x \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{\int \operatorname{PolyLog}(3, -e^{a+bx}) \, dx}{b} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) + \\
 & \quad \frac{1}{2} \left( \frac{x^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{2 \left( \frac{x \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{\int \operatorname{PolyLog}(3, e^{a+bx}) \, dx}{b} \right)}{b} \right) \\
 & \quad \downarrow \text{2720}
 \end{aligned}$$

$$\frac{1}{2} \left( \frac{2 \left( \frac{x \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{\int e^{-a-bx} \operatorname{PolyLog}(3, -e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) +$$

$$\frac{1}{2} \left( \frac{x^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{2 \left( \frac{x \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{\int e^{-a-bx} \operatorname{PolyLog}(3, e^{a+bx}) de^{a+bx}}{b^2} \right)}{b} \right)$$

↓ 7143

$$\frac{1}{2} \left( \frac{2 \left( \frac{x \operatorname{PolyLog}(3, -e^{a+bx})}{b} - \frac{\operatorname{PolyLog}(4, -e^{a+bx})}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, -e^{a+bx})}{b} \right) +$$

$$\frac{1}{2} \left( \frac{x^2 \operatorname{PolyLog}(2, e^{a+bx})}{b} - \frac{2 \left( \frac{x \operatorname{PolyLog}(3, e^{a+bx})}{b} - \frac{\operatorname{PolyLog}(4, e^{a+bx})}{b^2} \right)}{b} \right)$$

input `Int[x^2*ArcTanh[E^(a + b*x)],x]`

output `((-(x^2*PolyLog[2, -E^(a + b*x)])/b) + (2*((x*PolyLog[3, -E^(a + b*x)])/b - PolyLog[4, -E^(a + b*x)]/b^2))/b)/2 + ((x^2*PolyLog[2, E^(a + b*x)])/b - (2*((x*PolyLog[3, E^(a + b*x)])/b - PolyLog[4, E^(a + b*x)]/b^2))/b)/2`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(- (f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 6767

```
Int[ArcTanh[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol]
:> Simp[1/2 Int[x^m*Log[1 + a + b*f^(c + d*x)], x], x] - Simp[1/2 Int[x
^m*Log[1 - a - b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IGtQ[
m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_)^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs.  $2(91) = 182$ .

Time = 0.45 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.95

method	result
risch	$\frac{x^2 \operatorname{polylog}(2, e^{bx+a})}{2b} - \frac{\ln(1-e^{bx+a}) x a^2}{2b^2} - \frac{a^3 \ln(1-e^{bx+a})}{2b^3} - \frac{x \operatorname{polylog}(3, e^{bx+a})}{b^2} - \frac{\operatorname{polylog}(2, e^{bx+a}) a^2}{2b^3} - \frac{a^2 \operatorname{dilog}(e^{bx+a})}{2b^3}$
default	$\frac{x^3 \operatorname{arctanh}(e^{bx+a})}{3} - \frac{(bx+a)^3 \ln(1-e^{bx+a})}{2} - \frac{3(bx+a)^2 \operatorname{polylog}(2, e^{bx+a})}{2} + 3(bx+a) \operatorname{polylog}(3, e^{bx+a}) - 3 \operatorname{polylog}(4, e^{bx+a}) + \frac{(bx+a)^2 \operatorname{dilog}(e^{bx+a})}{2}$
parts	$\frac{x^3 \operatorname{arctanh}(e^{bx+a})}{3} - \frac{(bx+a)^3 \ln(1-e^{bx+a})}{2} - \frac{3(bx+a)^2 \operatorname{polylog}(2, e^{bx+a})}{2} + 3(bx+a) \operatorname{polylog}(3, e^{bx+a}) - 3 \operatorname{polylog}(4, e^{bx+a}) + \frac{(bx+a)^2 \operatorname{dilog}(e^{bx+a})}{2}$

input

```
int(x^2*arctanh(exp(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/2*x^2*polylog(2,exp(b*x+a))/b-1/2/b^2*ln(1-exp(b*x+a))*x*a^2-1/2/b^3*a^3
*ln(1-exp(b*x+a))-x*polylog(3,exp(b*x+a))/b^2-1/2/b^3*polylog(2,exp(b*x+a)
)*a^2-1/2/b^3*a^2*dilog(exp(b*x+a))+polylog(4,exp(b*x+a))/b^3-1/2*x^2*poly
log(2,-exp(b*x+a))/b+x*polylog(3,-exp(b*x+a))/b^2+1/2/b^3*polylog(2,-exp(b
*x+a))*a^2-1/2/b^3*dilog(exp(b*x+a)+1)*a^2-polylog(4,-exp(b*x+a))/b^3
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(89) = 178.

Time = 0.10 (sec) , antiderivative size = 248, normalized size of antiderivative = 2.46

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx$$

$$= \frac{b^3 x^3 \log\left(-\frac{\cosh(bx+a)+\sinh(bx+a)+1}{\cosh(bx+a)+\sinh(bx+a)-1}\right) - b^3 x^3 \log(\cosh(bx+a) + \sinh(bx+a) + 1) + 3b^2 x^2 \operatorname{Li}_2(\cosh(bx+a) + \sinh(bx+a) + 1)}{2}$$

input

```
integrate(x^2*arctanh(exp(b*x+a)),x, algorithm="fricas")
```

output

```
1/6*(b^3*x^3*log(-(cosh(b*x + a) + sinh(b*x + a) + 1)/(cosh(b*x + a) + sin
h(b*x + a) - 1)) - b^3*x^3*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 3*b^2*
x^2*dilog(cosh(b*x + a) + sinh(b*x + a)) - 3*b^2*x^2*dilog(-cosh(b*x + a)
- sinh(b*x + a)) - a^3*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 6*b*x*poly
log(3, cosh(b*x + a) + sinh(b*x + a)) + 6*b*x*polylog(3, -cosh(b*x + a) -
sinh(b*x + a)) + (b^3*x^3 + a^3)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) +
6*polylog(4, cosh(b*x + a) + sinh(b*x + a)) - 6*polylog(4, -cosh(b*x + a)
- sinh(b*x + a)))/b^3
```

### Sympy [F]

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \int x^2 \operatorname{atanh}(e^a e^{bx}) dx$$

input

```
integrate(x**2*atanh(exp(b*x+a)),x)
```



output `Integral(x**2*atanh(exp(a)*exp(b*x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.41

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \frac{1}{3} x^3 \operatorname{arctanh}(e^{(bx+a)}) - \frac{1}{6} b \left( \frac{b^3 x^3 \log(e^{(bx+a)} + 1) + 3 b^2 x^2 \operatorname{Li}_2(-e^{(bx+a)}) - 6 b x \operatorname{Li}_3(-e^{(bx+a)}) + 6 \operatorname{Li}_4(-e^{(bx+a)})}{b^4} - \frac{b^3 x^3 \log(-e^{(bx+a)})}{b^4} \right)$$

input `integrate(x^2*arctanh(exp(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctanh(e^(b*x + a)) - 1/6*b*((b^3*x^3*log(e^(b*x + a) + 1) + 3*b^2*x^2*dilog(-e^(b*x + a)) - 6*b*x*polylog(3, -e^(b*x + a)) + 6*polylog(4, -e^(b*x + a)))/b^4 - (b^3*x^3*log(-e^(b*x + a) + 1) + 3*b^2*x^2*dilog(e^(b*x + a)) - 6*b*x*polylog(3, e^(b*x + a)) + 6*polylog(4, e^(b*x + a)))/b^4)`

### Giac [F]

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \int x^2 \operatorname{arctanh}(e^{(bx+a)}) dx$$

input `integrate(x^2*arctanh(exp(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctanh(e^(b*x + a)), x)`

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \int x^2 \operatorname{atanh}(e^{a+bx}) dx$$

input `int(x^2*atanh(exp(a + b*x)),x)`output `int(x^2*atanh(exp(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(e^{a+bx}) dx = \int \operatorname{atanh}(e^{bx+a}) x^2 dx$$

input `int(x^2*atanh(exp(b*x+a)),x)`output `int(atanh(e**(a + b*x))*x**2,x)`

### 3.352 $\int \operatorname{arctanh}(a + bf^{c+dx}) dx$

Optimal result	2470
Mathematica [A] (verified)	2471
Rubi [A] (verified)	2471
Maple [A] (verified)	2474
Fricas [A] (verification not implemented)	2474
Sympy [F]	2475
Maxima [A] (verification not implemented)	2475
Giac [F(-2)]	2476
Mupad [F(-1)]	2476
Reduce [F]	2477

#### Optimal result

Integrand size = 12, antiderivative size = 168

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = -\frac{\operatorname{arctanh}(a + bf^{c+dx}) \log\left(\frac{2}{1+bf^{c+dx}}\right)}{d \log(f)} + \frac{\operatorname{arctanh}(a + bf^{c+dx}) \log\left(\frac{2bf^{c+dx}}{(1-a)(1+bf^{c+dx})}\right)}{d \log(f)} + \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2}{1+bf^{c+dx}}\right)}{2d \log(f)} - \frac{\operatorname{PolyLog}\left(2, 1 - \frac{2bf^{c+dx}}{(1-a)(1+bf^{c+dx})}\right)}{2d \log(f)}$$

output 
$$-\operatorname{arctanh}(a+b*f^{(d*x+c)})*\ln(2/(1+a+b*f^{(d*x+c)}))/d/\ln(f)+\operatorname{arctanh}(a+b*f^{(d*x+c)})*\ln(2*b*f^{(d*x+c)/(1-a)/(1+a+b*f^{(d*x+c)}))/d/\ln(f)+1/2*\operatorname{polylog}(2,1-2/(1+a+b*f^{(d*x+c)}))/d/\ln(f)-1/2*\operatorname{polylog}(2,1-2*b*f^{(d*x+c)/(1-a)/(1+a+b*f^{(d*x+c)}))/d/\ln(f)$$

**Mathematica [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.64

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$= \frac{dx \log(f) \left( 2\operatorname{arctanh}(a + bf^{c+dx}) + \log\left(\frac{-1+a+bf^{c+dx}}{-1+a}\right) - \log\left(\frac{1+a+bf^{c+dx}}{1+a}\right) \right) + \operatorname{PolyLog}\left(2, -\frac{bf^{c+dx}}{-1+a}\right) - \operatorname{PolyLog}\left(2, \frac{bf^{c+dx}}{1+a}\right)}{2d \log(f)}$$

input `Integrate[ArcTanh[a + b*f^(c + d*x)], x]`

output

```
(d*x*Log[f]*(2*ArcTanh[a + b*f^(c + d*x)] + Log[(-1 + a + b*f^(c + d*x))/(
-1 + a)] - Log[(1 + a + b*f^(c + d*x))/(1 + a)]) + PolyLog[2, -((b*f^(c +
d*x))/(-1 + a))] - PolyLog[2, -((b*f^(c + d*x))/(1 + a))]/(2*d*Log[f])
```

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {2720, 6661, 25, 27, 6472, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$\downarrow 2720$$

$$\frac{\int f^{-c-dx} \operatorname{arctanh}(bf^{c+dx} + a) df^{c+dx}}{d \log(f)}$$

$$\downarrow 6661$$

$$\frac{\int f^{-c-dx} \operatorname{arctanh}(bf^{c+dx} + a) d(bf^{c+dx} + a)}{bd \log(f)}$$

$$\downarrow 25$$

$$\frac{\int -f^{-c-dx} \operatorname{arctanh}(bf^{c+dx} + a) d(bf^{c+dx} + a)}{bd \log(f)}$$

$$\int \frac{-f^{-c-dx} \operatorname{arctanh}(bf^{c+dx}+a)}{b} d(bf^{c+dx}+a)$$

27

6472

$$\int \frac{\log\left(\frac{2}{bf^{c+dx}+a+1}\right)}{1-f^{2c+2dx}} d(bf^{c+dx}+a) + \int \frac{\log\left(\frac{2bf^{c+dx}}{(1-a)(bf^{c+dx}+a+1)}\right)}{1-f^{2c+2dx}} d(bf^{c+dx}+a) + \operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2}{a+bf^{c+dx}+1}\right)$$

$d \log(f)$

2849

$$\int \frac{\log\left(\frac{2}{bf^{c+dx}+a+1}\right)}{1-\frac{2}{bf^{c+dx}+a+1}} d\frac{1}{bf^{c+dx}+a+1} + \int \frac{\log\left(\frac{2bf^{c+dx}}{(1-a)(bf^{c+dx}+a+1)}\right)}{1-f^{2c+2dx}} d(bf^{c+dx}+a) + \operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2}{a+bf^{c+dx}+1}\right)$$

$d \log(f)$

2752

$$\int \frac{\log\left(\frac{2bf^{c+dx}}{(1-a)(bf^{c+dx}+a+1)}\right)}{1-f^{2c+2dx}} d(bf^{c+dx}+a) + \operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2}{a+bf^{c+dx}+1}\right) - \operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2}{(1-a)(bf^{c+dx}+a+1)}\right)$$

$d \log(f)$

2897

$$\operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2}{a+bf^{c+dx}+1}\right) - \operatorname{arctanh}(a+bf^{c+dx}) \log\left(\frac{2bf^{c+dx}}{(1-a)(a+bf^{c+dx}+1)}\right) - \frac{1}{2} \operatorname{PolyLog}\left(2, 1 - \frac{2}{a+bf^{c+dx}+1}\right)$$

$d \log(f)$

input `Int[ArcTanh[a + b*f^(c + d*x)], x]`

output `-((ArcTanh[a + b*f^(c + d*x)]*Log[2/(1 + a + b*f^(c + d*x))] - ArcTanh[a + b*f^(c + d*x)]*Log[(2*b*f^(c + d*x))/((1 - a)*(1 + a + b*f^(c + d*x)))] - PolyLog[2, 1 - 2/(1 + a + b*f^(c + d*x))]/2 + PolyLog[2, 1 - (2*b*f^(c + d*x))/((1 - a)*(1 + a + b*f^(c + d*x)))]/2)/(d*Log[f])`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]`
- rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`
- rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`
- rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`
- rule 6472 `Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x]))*(Log[2/(1 + c*x)]/e), x] + (Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x] + Simp[b*(c/e) Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Simp[b*(c/e) Int[Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]`

rule 6661

```
Int[((a_.) + ArcTanh[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(
m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*
ArcTanh[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IG
tQ[p, 0]
```

### Maple [A] (verified)

Time = 1.50 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.95

method	result
derivativedivides	$\frac{\ln(-b f^{dx+c}) \operatorname{arctanh}(a+b f^{dx+c}) - \frac{\operatorname{dilog}\left(\frac{-b f^{dx+c}-a-1}{-a-1}\right)}{2} - \frac{\ln(-b f^{dx+c}) \ln\left(\frac{-b f^{dx+c}-a-1}{-a-1}\right)}{d \ln(f)} + \frac{\operatorname{dilog}\left(\frac{1-a-b f^{dx+c}}{1-a}\right)}{2}}{2}$
default	$\frac{\ln(-b f^{dx+c}) \operatorname{arctanh}(a+b f^{dx+c}) - \frac{\operatorname{dilog}\left(\frac{-b f^{dx+c}-a-1}{-a-1}\right)}{2} - \frac{\ln(-b f^{dx+c}) \ln\left(\frac{-b f^{dx+c}-a-1}{-a-1}\right)}{d \ln(f)} + \frac{\operatorname{dilog}\left(\frac{1-a-b f^{dx+c}}{1-a}\right)}{2}}{2}$
risch	$\frac{x \ln(1+a+b f^{dx+c})}{2} - \frac{\operatorname{dilog}\left(\frac{1+a+b f^{dx+c}}{1+a}\right)}{2 \ln(f) d} - \frac{\ln\left(\frac{1+a+b f^{dx+c}}{1+a}\right) x}{2} - \frac{\ln\left(\frac{1+a+b f^{dx+c}}{1+a}\right) c}{2d} + \frac{c \ln(1+a+b f^{dx+c})}{2d}$

input

```
int(arctanh(a+b*f^(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/d/ln(f)*(ln(-b*f^(d*x+c))*arctanh(a+b*f^(d*x+c))-1/2*dilog((-b*f^(d*x+c)
-a-1)/(-a-1))-1/2*ln(-b*f^(d*x+c))*ln((-b*f^(d*x+c)-a-1)/(-a-1))+1/2*dilog
((1-a-b*f^(d*x+c))/(1-a))+1/2*ln(-b*f^(d*x+c))*ln((1-a-b*f^(d*x+c))/(1-a)
)
```

### Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.69

$$\int \operatorname{arctanh}(a + b f^{c+dx}) dx$$

$$= \frac{dx \log(f) \log\left(-\frac{b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a + 1}{b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a - 1}\right) + c \log(b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a + 1)}{2 \ln(f) d}$$

input

```
integrate(arctanh(a+b*f^(d*x+c)),x, algorithm="fricas")
```

output

```
1/2*(d*x*log(f)*log(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f))
+ a + 1)/(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)) +
c*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)*log(f)
- c*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)*log(f)
) - (d*x + c)*log(f)*log((b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)
+ a + 1)/(a + 1)) + (d*x + c)*log(f)*log((b*cosh((d*x + c)*log(f)) + b
*sinh((d*x + c)*log(f)) + a - 1)/(a - 1)) - dilog(-(b*cosh((d*x + c)*log(f)
)) + b*sinh((d*x + c)*log(f)) + a + 1)/(a + 1) + 1) + dilog(-(b*cosh((d*x
+ c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)/(a - 1) + 1))/(d*log(f))
```

**Sympy [F]**

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = \int \operatorname{atanh}(a + bf^{c+dx}) dx$$

input

```
integrate(atanh(a+b*f**(d*x+c)),x)
```

output

```
Integral(atanh(a + b*f**(c + d*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.20

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = \frac{(dx + c) \operatorname{artanh}(bf^{dx+c} + a)}{d} \\ - \frac{(dx + c)b \left( \frac{\log(bf^{dx+c+a+1})}{b} - \frac{\log(bf^{dx+c+a-1})}{b} \right) \log(f) - b \left( \frac{\log(bf^{dx+c+a+1}) \log\left(-\frac{bf^{dx+c+a+1}}{a+1} + 1\right) + \operatorname{Li}_2\left(\frac{bf^{dx+c+a+1}}{a+1}\right)}{b} \right)}{2d \log(f)}$$

input

```
integrate(arctanh(a+b*f^(d*x+c)),x, algorithm="maxima")
```



output

```
(d*x + c)*arctanh(b*f^(d*x + c) + a)/d - 1/2*((d*x + c)*b*(log(b*f^(d*x +
c) + a + 1)/b - log(b*f^(d*x + c) + a - 1)/b)*log(f) - b*((log(b*f^(d*x +
c) + a + 1)*log(-(b*f^(d*x + c) + a + 1)/(a + 1) + 1) + dilog((b*f^(d*x +
c) + a + 1)/(a + 1)))/b - (log(b*f^(d*x + c) + a - 1)*log(-(b*f^(d*x + c)
+ a - 1)/(a - 1) + 1) + dilog((b*f^(d*x + c) + a - 1)/(a - 1)))/b)/(d*log
(f))
```

**Giac [F(-2)]**

Exception generated.

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = \text{Exception raised: RuntimeError}$$

input

```
integrate(arctanh(a+b*f^(d*x+c)),x, algorithm="giac")
```

output

```
Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0
,1,2,0,0,0]%%}+%%{2,[0,1,1,1,1,0]%%}+%%{-2,[0,1,1,0,0,0]%%}+%%{1,[0,
1,0,2,0,1]%%
```

**Mupad [F(-1)]**

Timed out.

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = \int \operatorname{atanh}(a + bf^{c+dx}) dx$$

input

```
int(atanh(a + b*f^(c + d*x)),x)
```

output

```
int(atanh(a + b*f^(c + d*x)), x)
```

**Reduce [F]**

$$\int \operatorname{arctanh}(a + bf^{c+dx}) dx = \int \operatorname{atanh}(f^{dx+c}b + a) dx$$

input `int(atanh(a+b*f^(d*x+c)),x)`

output `int(atanh(f**(c + d*x)*b + a),x)`

### 3.353 $\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$

Optimal result	2478
Mathematica [A] (verified)	2479
Rubi [A] (verified)	2479
Maple [B] (verified)	2482
Fricas [B] (verification not implemented)	2482
Sympy [F]	2483
Maxima [A] (verification not implemented)	2483
Giac [F]	2484
Mupad [F(-1)]	2484
Reduce [F]	2485

#### Optimal result

Integrand size = 14, antiderivative size = 211

$$\begin{aligned} \int x \operatorname{arctanh}(a + bf^{c+dx}) dx &= -\frac{1}{4}x^2 \log(1 - a - bf^{c+dx}) + \frac{1}{4}x^2 \log(1 + a + bf^{c+dx}) \\ &+ \frac{1}{4}x^2 \log\left(1 - \frac{bf^{c+dx}}{1-a}\right) - \frac{1}{4}x^2 \log\left(1 + \frac{bf^{c+dx}}{1+a}\right) \\ &+ \frac{x \operatorname{PolyLog}\left(2, \frac{bf^{c+dx}}{1-a}\right)}{2d \log(f)} - \frac{x \operatorname{PolyLog}\left(2, -\frac{bf^{c+dx}}{1+a}\right)}{2d \log(f)} \\ &- \frac{\operatorname{PolyLog}\left(3, \frac{bf^{c+dx}}{1-a}\right)}{2d^2 \log^2(f)} + \frac{\operatorname{PolyLog}\left(3, -\frac{bf^{c+dx}}{1+a}\right)}{2d^2 \log^2(f)} \end{aligned}$$

output

```
-1/4*x^2*ln(1-a-b*f^(d*x+c))+1/4*x^2*ln(1+a+b*f^(d*x+c))+1/4*x^2*ln(1-b*f^(d*x+c)/(1-a))-1/4*x^2*ln(1+b*f^(d*x+c)/(1+a))+1/2*x*polylog(2,b*f^(d*x+c)/(1-a))/d/ln(f)-1/2*x*polylog(2,-b*f^(d*x+c)/(1+a))/d/ln(f)-1/2*polylog(3,b*f^(d*x+c)/(1-a))/d^2/ln(f)^2+1/2*polylog(3,-b*f^(d*x+c)/(1+a))/d^2/ln(f)^2
```

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.84

$$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$= \frac{2d^2x^2 \operatorname{arctanh}(a + bf^{c+dx}) \log^2(f) + d^2x^2 \log^2(f) \log\left(1 + \frac{bf^{c+dx}}{-1+a}\right) - d^2x^2 \log^2(f) \log\left(1 + \frac{bf^{c+dx}}{1+a}\right) + 2dx^2 \operatorname{arctanh}(a + bf^{c+dx})}{d}$$

input `Integrate[x*ArcTanh[a + b*f^(c + d*x)],x]`

output

```
(2*d^2*x^2*ArcTanh[a + b*f^(c + d*x)]*Log[f]^2 + d^2*x^2*Log[f]^2*Log[1 +
(b*f^(c + d*x))/(-1 + a)] - d^2*x^2*Log[f]^2*Log[1 + (b*f^(c + d*x))/(1 +
a)] + 2*d*x*Log[f]*PolyLog[2, -((b*f^(c + d*x))/(-1 + a))] - 2*d*x*Log[f]*
PolyLog[2, -((b*f^(c + d*x))/(1 + a))] - 2*PolyLog[3, -((b*f^(c + d*x))/(-
1 + a))] + 2*PolyLog[3, -((b*f^(c + d*x))/(1 + a))]/(4*d^2*Log[f]^2)
```

**Rubi [A] (verified)**

Time = 0.67 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {6767, 3012, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$\downarrow \text{6767}$$

$$\frac{1}{2} \int x \log(bf^{c+dx} + a + 1) dx - \frac{1}{2} \int x \log(-bf^{c+dx} - a + 1) dx$$

$$\downarrow \text{3012}$$

$$\frac{1}{2} \left( - \int x \log\left(1 - \frac{bf^{c+dx}}{1-a}\right) dx - \frac{1}{2} x^2 \log(-a - bf^{c+dx} + 1) + \frac{1}{2} x^2 \log\left(1 - \frac{bf^{c+dx}}{1-a}\right) \right) +$$

$$\frac{1}{2} \left( \int x \log\left(\frac{bf^{c+dx}}{a+1} + 1\right) dx + \frac{1}{2} x^2 \log(a + bf^{c+dx} + 1) - \frac{1}{2} x^2 \log\left(\frac{bf^{c+dx}}{a+1} + 1\right) \right)$$

↓ 3011

$$\frac{1}{2} \left( -\frac{\int \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right) dx}{d \log(f)} + \frac{x \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{2} x^2 \log(-a - bf^{c+dx} + 1) + \frac{1}{2} x^2 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) \right) \\ \frac{1}{2} \left( \frac{\int \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right) dx}{d \log(f)} - \frac{x \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{2} x^2 \log(a + bf^{c+dx} + 1) - \frac{1}{2} x^2 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) \right)$$

↓ 2720

$$\frac{1}{2} \left( -\frac{\int f^{-c-dx} \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right) df^{c+dx}}{d^2 \log^2(f)} + \frac{x \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{2} x^2 \log(-a - bf^{c+dx} + 1) + \frac{1}{2} x^2 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) \right) \\ \frac{1}{2} \left( \frac{\int f^{-c-dx} \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right) df^{c+dx}}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{2} x^2 \log(a + bf^{c+dx} + 1) - \frac{1}{2} x^2 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) \right)$$

↓ 7143

$$\frac{1}{2} \left( -\frac{\text{PolyLog} \left( 3, \frac{bf^{c+dx}}{1-a} \right)}{d^2 \log^2(f)} + \frac{x \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{2} x^2 \log(-a - bf^{c+dx} + 1) + \frac{1}{2} x^2 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) \right) + \\ \frac{1}{2} \left( \frac{\text{PolyLog} \left( 3, -\frac{bf^{c+dx}}{a+1} \right)}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{2} x^2 \log(a + bf^{c+dx} + 1) - \frac{1}{2} x^2 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) \right)$$

input `Int[x*ArcTanh[a + b*f^(c + d*x)],x]`

output `(-1/2*(x^2*Log[1 - a - b*f^(c + d*x)]) + (x^2*Log[1 - (b*f^(c + d*x))/(1 - a]]))/2 + (x*PolyLog[2, (b*f^(c + d*x))/(1 - a]])/(d*Log[f]) - PolyLog[3, (b*f^(c + d*x))/(1 - a]]/(d^2*Log[f]^2))/2 + ((x^2*Log[1 + a + b*f^(c + d*x)]))/2 - (x^2*Log[1 + (b*f^(c + d*x))/(1 + a]]))/2 - (x*PolyLog[2, -((b*f^(c + d*x))/(1 + a))])/(d*Log[f]) + PolyLog[3, -((b*f^(c + d*x))/(1 + a))]/(d^2*Log[f]^2))/2`

## Definitions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3012

```
Int[Log[(d_) + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g
_)*(x_))^(m_), x_Symbol] := Simp[(f + g*x)^(m + 1)*(Log[d + e*(F^(c*(a +
b*x)))^n]/(g*(m + 1))), x] + (Int[(f + g*x)^m*Log[1 + (e/d)*(F^(c*(a + b*x)
))^n], x] - Simp[(f + g*x)^(m + 1)*(Log[1 + (e/d)*(F^(c*(a + b*x)))^n]/(g*(
m + 1))), x]) /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && NeQ[
d, 1]
```

rule 6767

```
Int[ArcTanh[(a_) + (b_)*(f_)^((c_) + (d_)*(x_))]*(x_)^(m_), x_Symbol]
:= Simp[1/2 Int[x^m*Log[1 + a + b*f^(c + d*x)], x], x] - Simp[1/2 Int[x
^m*Log[1 - a - b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IGtQ[
m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 595 vs.  $2(195) = 390$ .

Time = 0.86 (sec) , antiderivative size = 596, normalized size of antiderivative = 2.82

method	result
risch	$\frac{x^2 \ln(1+a+bf^{dx+c})}{4} - \frac{x^2 \ln(1-a-bf^{dx+c})}{4} + \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)x^2}{4} + \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)cx}{2d} + \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)c^2}{4d^2} + \frac{\text{polylog}\left(\dots\right)}{2d}$

input

```
int(x*arctanh(a+b*f^(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/4*x^2*ln(1+a+b*f^(d*x+c))-1/4*x^2*ln(1-a-b*f^(d*x+c))+1/4*ln(1-b*f^(d*x)
*f^c/(1-a))*x^2+1/2/d*ln(1-b*f^(d*x)*f^c/(1-a))*c*x+1/4/d^2*ln(1-b*f^(d*x)
*f^c/(1-a))*c^2+1/2/ln(f)/d*polylog(2,b*f^(d*x)*f^c/(1-a))*x+1/2/ln(f)/d^2
*polylog(2,b*f^(d*x)*f^c/(1-a))*c-1/2/ln(f)^2/d^2*polylog(3,b*f^(d*x)*f^c/
(1-a))+1/4/d^2*c^2*ln(1-a-b*f^(d*x)*f^c)-1/2/ln(f)/d^2*c*dilog((b*f^(d*x)*
f^c+a-1)/(a-1))-1/2/d*c*ln((b*f^(d*x)*f^c+a-1)/(a-1))*x-1/2/d^2*c^2*ln((b*
f^(d*x)*f^c+a-1)/(a-1))-1/4*ln(1-b*f^(d*x)*f^c/(-a-1))*x^2-1/2/d*ln(1-b*f^
(d*x)*f^c/(-a-1))*c*x-1/4/d^2*ln(1-b*f^(d*x)*f^c/(-a-1))*c^2-1/2/ln(f)/d*p
olylog(2,b*f^(d*x)*f^c/(-a-1))*x-1/2/ln(f)/d^2*polylog(2,b*f^(d*x)*f^c/(-a
-1))*c+1/2/ln(f)^2/d^2*polylog(3,b*f^(d*x)*f^c/(-a-1))-1/4/d^2*c^2*ln(1+a+
b*f^(d*x)*f^c)+1/2/ln(f)/d^2*c*dilog((1+a+b*f^(d*x)*f^c)/(1+a))+1/2/d*c*ln
((1+a+b*f^(d*x)*f^c)/(1+a))*x+1/2/d^2*c^2*ln((1+a+b*f^(d*x)*f^c)/(1+a))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 396 vs.  $2(188) = 376$ .

Time = 0.10 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.88

$$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$= \frac{d^2 x^2 \log(f)^2 \log\left(\frac{b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a + 1}{b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a - 1}\right) - c^2 \log(b \cosh((dx+c) \log(f)) + b \sinh((dx+c) \log(f)) + a + 1)}{d^2}$$

input

```
integrate(x*arctanh(a+b*f^(d*x+c)),x, algorithm="fricas")
```

output

```
1/4*(d^2*x^2*log(f)^2*log(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)) - c^2*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)*log(f)^2 + c^2*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)*log(f)^2 - 2*d*x*dilog(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(a + 1) + 1)*log(f) + 2*d*x*dilog(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)/(a - 1) + 1)*log(f) - (d^2*x^2 - c^2)*log(f)^2*log((b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(a + 1)) + (d^2*x^2 - c^2)*log(f)^2*log((b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)/(a - 1)) + 2*polylog(3, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a + 1)) - 2*polylog(3, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a - 1)))/(d^2*log(f)^2)
```

### Sympy [F]

$$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx = \int x \operatorname{atanh}(a + bf^{c+dx}) dx$$

input

```
integrate(x*atanh(a+b*f**(d*x+c)),x)
```

output

```
Integral(x*atanh(a + b*f**(c + d*x)), x)
```

### Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.92

$$\int x \operatorname{arctanh}(a + bf^{c+dx}) dx =$$

$$-\frac{1}{4} bd \left( \frac{d^2 x^2 \log\left(\frac{bf^{dx} f^c}{a+1} + 1\right) \log(f)^2 + 2 dx \operatorname{Li}_2\left(-\frac{bf^{dx} f^c}{a+1}\right) \log(f) - 2 \operatorname{Li}_3\left(-\frac{bf^{dx} f^c}{a+1}\right)}{bd^3 \log(f)^3} - \frac{d^2 x^2 \log\left(\frac{bf^{dx} f^c}{a-1} + 1\right)}{bd^3 \log(f)^3} \right)$$

$$+ \frac{1}{2} x^2 \operatorname{artanh}(bf^{dx+c} + a)$$

input

```
integrate(x*arctanh(a+b*f^(d*x+c)),x, algorithm="maxima")
```



output

```
-1/4*b*d*((d^2*x^2*log(b*f^(d*x)*f^c/(a + 1) + 1)*log(f)^2 + 2*d*x*dilog(-
b*f^(d*x)*f^c/(a + 1))*log(f) - 2*polylog(3, -b*f^(d*x)*f^c/(a + 1)))/(b*d
^3*log(f)^3) - (d^2*x^2*log(b*f^(d*x)*f^c/(a - 1) + 1)*log(f)^2 + 2*d*x*di
log(-b*f^(d*x)*f^c/(a - 1))*log(f) - 2*polylog(3, -b*f^(d*x)*f^c/(a - 1))
/(b*d^3*log(f)^3))*log(f) + 1/2*x^2*arctanh(b*f^(d*x + c) + a)
```

**Giac [F]**

$$\int x \operatorname{arctanh}(a + b f^{c+dx}) dx = \int x \operatorname{artanh}(b f^{dx+c} + a) dx$$

input

```
integrate(x*arctanh(a+b*f^(d*x+c)),x, algorithm="giac")
```

output

```
integrate(x*arctanh(b*f^(d*x + c) + a), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x \operatorname{arctanh}(a + b f^{c+dx}) dx = \int x \operatorname{atanh}(a + b f^{c+dx}) dx$$

input

```
int(x*atanh(a + b*f^(c + d*x)),x)
```

output

```
int(x*atanh(a + b*f^(c + d*x)), x)
```

**Reduce [F]**

$$\int x \operatorname{arctanh}(a + b f^{c+dx}) dx = \int \operatorname{atanh}(f^{dx+c} b + a) x dx$$

input `int(x*atanh(a+b*f^(d*x+c)),x)`

output `int(atanh(f**(c + d*x)*b + a)*x,x)`

### 3.354 $\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx$

Optimal result	2486
Mathematica [A] (verified)	2487
Rubi [A] (verified)	2487
Maple [B] (verified)	2491
Fricas [A] (verification not implemented)	2491
Sympy [F]	2492
Maxima [A] (verification not implemented)	2492
Giac [F]	2493
Mupad [F(-1)]	2493
Reduce [F]	2494

#### Optimal result

Integrand size = 16, antiderivative size = 264

$$\begin{aligned} \int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx &= -\frac{1}{6}x^3 \log(1 - a - bf^{c+dx}) + \frac{1}{6}x^3 \log(1 + a + bf^{c+dx}) \\ &+ \frac{1}{6}x^3 \log\left(1 - \frac{bf^{c+dx}}{1-a}\right) - \frac{1}{6}x^3 \log\left(1 + \frac{bf^{c+dx}}{1+a}\right) \\ &+ \frac{x^2 \operatorname{PolyLog}\left(2, \frac{bf^{c+dx}}{1-a}\right)}{2d \log(f)} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{bf^{c+dx}}{1+a}\right)}{2d \log(f)} \\ &- \frac{x \operatorname{PolyLog}\left(3, \frac{bf^{c+dx}}{1-a}\right)}{d^2 \log^2(f)} + \frac{x \operatorname{PolyLog}\left(3, -\frac{bf^{c+dx}}{1+a}\right)}{d^2 \log^2(f)} \\ &+ \frac{\operatorname{PolyLog}\left(4, \frac{bf^{c+dx}}{1-a}\right)}{d^3 \log^3(f)} - \frac{\operatorname{PolyLog}\left(4, -\frac{bf^{c+dx}}{1+a}\right)}{d^3 \log^3(f)} \end{aligned}$$

output

```
-1/6*x^3*ln(1-a-b*f^(d*x+c))+1/6*x^3*ln(1+a+b*f^(d*x+c))+1/6*x^3*ln(1-b*f^(d*x+c)/(1-a))-1/6*x^3*ln(1+b*f^(d*x+c)/(1+a))+1/2*x^2*polylog(2,b*f^(d*x+c)/(1-a))/d/ln(f)-1/2*x^2*polylog(2,-b*f^(d*x+c)/(1+a))/d/ln(f)-x*polylog(3,b*f^(d*x+c)/(1-a))/d^2/ln(f)^2+x*polylog(3,-b*f^(d*x+c)/(1+a))/d^2/ln(f)^2+polylog(4,b*f^(d*x+c)/(1-a))/d^3/ln(f)^3-polylog(4,-b*f^(d*x+c)/(1+a))/d^3/ln(f)^3
```

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.89

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$= \frac{2d^3 x^3 \operatorname{arctanh}(a + bf^{c+dx}) \log^3(f) + d^3 x^3 \log^3(f) \log\left(1 + \frac{bf^{c+dx}}{-1+a}\right) - d^3 x^3 \log^3(f) \log\left(1 + \frac{bf^{c+dx}}{1+a}\right) + 3d^2$$

input

```
Integrate[x^2*ArcTanh[a + b*f^(c + d*x)],x]
```

output

```
(2*d^3*x^3*ArcTanh[a + b*f^(c + d*x)]*Log[f]^3 + d^3*x^3*Log[f]^3*Log[1 +
(b*f^(c + d*x))/(-1 + a)] - d^3*x^3*Log[f]^3*Log[1 + (b*f^(c + d*x))/(1 +
a)] + 3*d^2*x^2*Log[f]^2*PolyLog[2, -((b*f^(c + d*x))/(-1 + a))] - 3*d^2*x
^2*Log[f]^2*PolyLog[2, -((b*f^(c + d*x))/(1 + a))] - 6*d*x*Log[f]*PolyLog[
3, -((b*f^(c + d*x))/(-1 + a))] + 6*d*x*Log[f]*PolyLog[3, -((b*f^(c + d*x)
)/(1 + a))] + 6*PolyLog[4, -((b*f^(c + d*x))/(-1 + a))] - 6*PolyLog[4, -((
b*f^(c + d*x))/(1 + a))]/(6*d^3*Log[f]^3)
```

**Rubi [A] (verified)**

Time = 0.98 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {6767, 3012, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx$$

$$\downarrow \text{6767}$$

$$\frac{1}{2} \int x^2 \log(bf^{c+dx} + a + 1) dx - \frac{1}{2} \int x^2 \log(-bf^{c+dx} - a + 1) dx$$

$$\downarrow \text{3012}$$

$$\frac{1}{2} \left( - \int x^2 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) dx - \frac{1}{3} x^3 \log (-a - bf^{c+dx} + 1) + \frac{1}{3} x^3 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) \right) + \frac{1}{2} \left( \int x^2 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) dx + \frac{1}{3} x^3 \log (a + bf^{c+dx} + 1) - \frac{1}{3} x^3 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) \right)$$

↓ 3011

$$\frac{1}{2} \left( - \frac{2 \int x \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right) dx}{d \log(f)} + \frac{x^2 \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{3} x^3 \log (-a - bf^{c+dx} + 1) + \frac{1}{3} x^3 \log \left( 1 - \frac{bf^{c+dx}}{1-a} \right) \right) + \frac{1}{2} \left( \frac{2 \int x \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right) dx}{d \log(f)} - \frac{x^2 \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{3} x^3 \log (a + bf^{c+dx} + 1) - \frac{1}{3} x^3 \log \left( \frac{bf^{c+dx}}{a+1} + 1 \right) \right)$$

↓ 7163

$$\frac{1}{2} \left( \frac{2 \left( \frac{x \text{PolyLog} \left( 3, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{\int \text{PolyLog} \left( 3, \frac{bf^{c+dx}}{1-a} \right) dx}{d \log(f)} \right)}{d \log(f)} + \frac{x^2 \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{3} x^3 \log (-a - bf^{c+dx} + 1) \right) + \frac{1}{2} \left( \frac{2 \left( \frac{x \text{PolyLog} \left( 3, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} - \frac{\int \text{PolyLog} \left( 3, -\frac{bf^{c+dx}}{a+1} \right) dx}{d \log(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{3} x^3 \log (a + bf^{c+dx} + 1) \right)$$

↓ 2720

$$\frac{1}{2} \left( \frac{2 \left( \frac{x \text{PolyLog} \left( 3, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{\int f^{-c-dx} \text{PolyLog} \left( 3, \frac{bf^{c+dx}}{1-a} \right) df^{c+dx}}{d^2 \log^2(f)} \right)}{d \log(f)} + \frac{x^2 \text{PolyLog} \left( 2, \frac{bf^{c+dx}}{1-a} \right)}{d \log(f)} - \frac{1}{3} x^3 \log (-a - bf^{c+dx} + 1) \right) + \frac{1}{2} \left( \frac{2 \left( \frac{x \text{PolyLog} \left( 3, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} - \frac{\int f^{-c-dx} \text{PolyLog} \left( 3, -\frac{bf^{c+dx}}{a+1} \right) df^{c+dx}}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left( 2, -\frac{bf^{c+dx}}{a+1} \right)}{d \log(f)} + \frac{1}{3} x^3 \log (a + bf^{c+dx} + 1) \right)$$

↓ 7143

$$\frac{1}{2} \left( \frac{2 \left( \frac{x \operatorname{PolyLog}\left(3, \frac{bf^{c+dx}}{1-a}\right)}{d \log(f)} - \frac{\operatorname{PolyLog}\left(4, \frac{bf^{c+dx}}{1-a}\right)}{d^2 \log^2(f)} \right)}{d \log(f)} + \frac{x^2 \operatorname{PolyLog}\left(2, \frac{bf^{c+dx}}{1-a}\right)}{d \log(f)} - \frac{1}{3} x^3 \log(-a - bf^{c+dx} + 1) + \frac{1}{3} x^3 \right) \\ \frac{1}{2} \left( \frac{2 \left( \frac{x \operatorname{PolyLog}\left(3, -\frac{bf^{c+dx}}{a+1}\right)}{d \log(f)} - \frac{\operatorname{PolyLog}\left(4, -\frac{bf^{c+dx}}{a+1}\right)}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{bf^{c+dx}}{a+1}\right)}{d \log(f)} + \frac{1}{3} x^3 \log(a + bf^{c+dx} + 1) - \frac{1}{3} x^3 \right)$$

input `Int[x^2*ArcTanh[a + b*f^(c + d*x)],x]`

output `(-1/3*(x^3*Log[1 - a - b*f^(c + d*x)]) + (x^3*Log[1 - (b*f^(c + d*x))/(1 - a)])/3 + (x^2*PolyLog[2, (b*f^(c + d*x))/(1 - a)]/(d*Log[f]) - (2*((x*PolyLog[3, (b*f^(c + d*x))/(1 - a)]/(d*Log[f]) - PolyLog[4, (b*f^(c + d*x))/(1 - a)]/(d^2*Log[f]^2)))/(d*Log[f]))/2 + ((x^3*Log[1 + a + b*f^(c + d*x)])/3 - (x^3*Log[1 + (b*f^(c + d*x))/(1 + a)])/3 - (x^2*PolyLog[2, -((b*f^(c + d*x))/(1 + a))]/(d*Log[f]) + (2*((x*PolyLog[3, -((b*f^(c + d*x))/(1 + a)))]/(d*Log[f]) - PolyLog[4, -((b*f^(c + d*x))/(1 + a))]/(d^2*Log[f]^2)))/(d*Log[f]))/2`

### Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011  $\text{Int}[\text{Log}[1 + (e\_.) * ((F\_)^{(c\_.) * ((a\_.) + (b\_.) * (x\_))})^{(n\_.)}] * ((f\_.) + (g\_.) * (x\_))^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[(-f + g*x)^m * (\text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n] / (b*c*n*\text{Log}[F])), x] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \text{Int}[(f + g*x)^{m-1} * \text{PolyLog}[2, (-e) * (F^{(c*(a + b*x))})^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0]$

rule 3012  $\text{Int}[\text{Log}[(d\_.) + (e\_.) * ((F\_)^{(c\_.) * ((a\_.) + (b\_.) * (x\_))})^{(n\_.)}] * ((f\_.) + (g\_.) * (x\_))^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[(f + g*x)^{m+1} * (\text{Log}[d + e * (F^{(c*(a + b*x))})^n] / (g*(m+1))), x] + (\text{Int}[(f + g*x)^m * \text{Log}[1 + (e/d) * (F^{(c*(a + b*x))})^n], x] - \text{Simp}[(f + g*x)^{m+1} * (\text{Log}[1 + (e/d) * (F^{(c*(a + b*x))})^n] / (g*(m+1))), x]) /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0] \&\& \text{NeQ}[d, 1]$

rule 6767  $\text{Int}[\text{ArcTanh}[(a\_.) + (b\_.) * (f\_.)^{(c\_.) + (d\_.) * (x\_)}] * (x\_)^{(m\_.)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \text{Int}[x^m * \text{Log}[1 + a + b*f^{(c + d*x)}], x], x] - \text{Simp}[1/2 \text{Int}[x^m * \text{Log}[1 - a - b*f^{(c + d*x)}], x], x] /; \text{FreeQ}[\{a, b, c, d, f\}, x] \&\& \text{IGtQ}[m, 0]$

rule 7143  $\text{Int}[\text{PolyLog}[n\_., (c\_.) * ((a\_.) + (b\_.) * (x\_))^{(p\_.)}] / ((d\_.) + (e\_.) * (x\_)), x\_Symbol] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p] / (e*p), x] /; \text{FreeQ}[\{a, b, c, d, e, n, p\}, x] \&\& \text{EqQ}[b*d, a*e]$

rule 7163  $\text{Int}[(e\_.) + (f\_.) * (x\_)]^{(m\_.)} * \text{PolyLog}[n\_., (d\_.) * ((F\_)^{(c\_.) * ((a\_.) + (b\_.) * (x\_))})^{(p\_.)}], x\_Symbol] \rightarrow \text{Simp}[(e + f*x)^m * (\text{PolyLog}[n + 1, d * (F^{(c*(a + b*x))})^p] / (b*c*p*\text{Log}[F])), x] - \text{Simp}[f*(m/(b*c*p*\text{Log}[F])) \text{Int}[(e + f*x)^{m-1} * \text{PolyLog}[n + 1, d * (F^{(c*(a + b*x))})^p], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[m, 0]$

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 671 vs.  $2(252) = 504$ .

Time = 1.26 (sec) , antiderivative size = 672, normalized size of antiderivative = 2.55

method	result
risch	$\frac{x^3 \ln(1+a+bf^{dx+c})}{6} - \frac{x^3 \ln(1-a-bf^{dx+c})}{6} + \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)x^3}{6} - \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)xc^2}{2d^2} - \frac{\ln\left(1-\frac{bf^{dx}f^c}{1-a}\right)c^3}{3d^3} + \frac{\text{polylog}}{6}$

input `int(x^2*arctanh(a+b*f^(d*x+c)),x,method=_RETURNVERBOSE)`

output

```
1/6*x^3*ln(1+a+b*f^(d*x+c))-1/6*x^3*ln(1-a-b*f^(d*x+c))+1/6*ln(1-b*f^(d*x)
*f^c/(1-a))*x^3-1/2/d^2*ln(1-b*f^(d*x)*f^c/(1-a))*x*c^2-1/3/d^3*ln(1-b*f^(
d*x)*f^c/(1-a))*c^3+1/2/ln(f)/d*polylog(2,b*f^(d*x)*f^c/(1-a))*x^2-1/2/ln(
f)/d^3*polylog(2,b*f^(d*x)*f^c/(1-a))*c^2-1/ln(f)^2/d^2*polylog(3,b*f^(d*x)
*f^c/(1-a))*x+1/ln(f)^3/d^3*polylog(4,b*f^(d*x)*f^c/(1-a))-1/6/d^3*c^3*ln
(1-a-b*f^(d*x)*f^c)+1/2/ln(f)/d^3*c^2*di-log((b*f^(d*x)*f^c+a-1)/(a-1))+1/2
/d^2*c^2*ln((b*f^(d*x)*f^c+a-1)/(a-1))*x+1/2/d^3*c^3*ln((b*f^(d*x)*f^c+a-1
)/(a-1))-1/6*ln(1-b*f^(d*x)*f^c/(-a-1))*x^3+1/2/d^2*ln(1-b*f^(d*x)*f^c/(-a
-1))*x*c^2+1/3/d^3*ln(1-b*f^(d*x)*f^c/(-a-1))*c^3-1/2/ln(f)/d*polylog(2,b*
f^(d*x)*f^c/(-a-1))*x^2+1/2/ln(f)/d^3*polylog(2,b*f^(d*x)*f^c/(-a-1))*c^2+
1/ln(f)^2/d^2*polylog(3,b*f^(d*x)*f^c/(-a-1))*x-1/ln(f)^3/d^3*polylog(4,b*
f^(d*x)*f^c/(-a-1))+1/6/d^3*c^3*ln(1+a+b*f^(d*x)*f^c)-1/2/ln(f)/d^3*c^2*di
-log(((1+a+b*f^(d*x)*f^c)/(1+a))-1/2/d^2*c^2*ln(((1+a+b*f^(d*x)*f^c)/(1+a))*x
-1/2/d^3*c^3*ln((1+a+b*f^(d*x)*f^c)/(1+a)))
```

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.82

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx = \text{Too large to display}$$

input `integrate(x^2*arctanh(a+b*f^(d*x+c)),x, algorithm="fricas")`



output

```
1/6*(d^3*x^3*log(f)^3*log(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)) - 3*d^2*x^2*dilog(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(a + 1) + 1)*log(f)^2 + 3*d^2*x^2*dilog(-(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)/(a - 1) + 1)*log(f)^2 + c^3*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)*log(f)^3 - c^3*log(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)*log(f)^3 - (d^3*x^3 + c^3)*log(f)^3*log((b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a + 1)/(a + 1)) + (d^3*x^3 + c^3)*log(f)^3*log((b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)) + a - 1)/(a - 1)) + 6*d*x*log(f)*polylog(3, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a + 1)) - 6*d*x*log(f)*polylog(3, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a - 1)) - 6*polylog(4, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a + 1)) + 6*polylog(4, -(b*cosh((d*x + c)*log(f)) + b*sinh((d*x + c)*log(f)))/(a - 1)))/(d^3*log(f)^3)
```

**Sympy [F]**

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx = \int x^2 \operatorname{atanh}(a + bf^{c+dx}) dx$$

input

```
integrate(x**2*atanh(a+b*f**(d*x+c)),x)
```

output

```
Integral(x**2*atanh(a + b*f**(c + d*x)), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.96

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx = \frac{1}{3} x^3 \operatorname{artanh}(bf^{dx+c} + a) - \frac{1}{6} bd \left( \frac{d^3 x^3 \log\left(\frac{bf^{dx} f^c}{a+1} + 1\right) \log(f)^3 + 3 d^2 x^2 \operatorname{Li}_2\left(-\frac{bf^{dx} f^c}{a+1}\right) \log(f)^2 - 6 dx \log(f) \operatorname{Li}_3\left(-\frac{bf^{dx} f^c}{a+1}\right) + 6 \operatorname{Li}_4\left(-\frac{bf^{dx} f^c}{a+1}\right)}{bd^4 \log(f)^4} \right)$$

input

```
integrate(x^2*arctanh(a+b*f^(d*x+c)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctanh(b*f^(d*x + c) + a) - 1/6*b*d*((d^3*x^3*log(b*f^(d*x)*f^c/(a + 1) + 1)*log(f)^3 + 3*d^2*x^2*dilog(-b*f^(d*x)*f^c/(a + 1))*log(f)^2 - 6*d*x*log(f)*polylog(3, -b*f^(d*x)*f^c/(a + 1)) + 6*polylog(4, -b*f^(d*x)*f^c/(a + 1)))/(b*d^4*log(f)^4) - (d^3*x^3*log(b*f^(d*x)*f^c/(a - 1) + 1)*log(f)^3 + 3*d^2*x^2*dilog(-b*f^(d*x)*f^c/(a - 1))*log(f)^2 - 6*d*x*log(f)*polylog(3, -b*f^(d*x)*f^c/(a - 1)) + 6*polylog(4, -b*f^(d*x)*f^c/(a - 1)))/(b*d^4*log(f)^4))*log(f)
```

**Giac [F]**

$$\int x^2 \operatorname{arctanh}(a + b f^{c+dx}) dx = \int x^2 \operatorname{artanh}(b f^{dx+c} + a) dx$$

input

```
integrate(x^2*arctanh(a+b*f^(d*x+c)),x, algorithm="giac")
```

output

```
integrate(x^2*arctanh(b*f^(d*x + c) + a), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int x^2 \operatorname{arctanh}(a + b f^{c+dx}) dx = \int x^2 \operatorname{atanh}(a + b f^{c+dx}) dx$$

input

```
int(x^2*atanh(a + b*f^(c + d*x)),x)
```

output

```
int(x^2*atanh(a + b*f^(c + d*x)), x)
```

**Reduce [F]**

$$\int x^2 \operatorname{arctanh}(a + bf^{c+dx}) dx = \int \operatorname{atanh}(f^{dx+c}b + a) x^2 dx$$

input `int(x^2*atanh(a+b*f^(d*x+c)),x)`

output `int(atanh(f**(c + d*x)*b + a)*x**2,x)`

### 3.355 $\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$

Optimal result	2495
Mathematica [A] (verified)	2495
Rubi [A] (warning: unable to verify)	2496
Maple [C] (warning: unable to verify)	2498
Fricas [B] (verification not implemented)	2499
Sympy [F(-1)]	2500
Maxima [A] (verification not implemented)	2500
Giac [A] (verification not implemented)	2501
Mupad [B] (verification not implemented)	2501
Reduce [F]	2502

#### Optimal result

Integrand size = 20, antiderivative size = 112

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx = \frac{e^{ac+bcx} \operatorname{arctanh}(\sinh(c(a + bx)))}{bc} + \frac{(1 - \sqrt{2}) \log(3 - 2\sqrt{2} - e^{2c(a+bx)})}{2bc} + \frac{(1 + \sqrt{2}) \log(4 + 3\sqrt{2} - \sqrt{2}e^{2c(a+bx)})}{2bc}$$

output

```
exp(b*c*x+a*c)*arctanh(sinh(c*(b*x+a)))/b/c+1/2*(1-2^(1/2))*ln(3-2*2^(1/2)
-exp(2*c*(b*x+a))/b/c+1/2*(1+2^(1/2))*ln(4+3*2^(1/2)-2^(1/2)*exp(2*c*(b*x
+a)))/b/c
```

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.37

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx = \frac{-2e^{c(a+bx)} \operatorname{arctanh}\left(\frac{1}{2}e^{-c(a+bx)} - \frac{1}{2}e^{c(a+bx)}\right) - 2\sqrt{2} \operatorname{arctanh}\left(\frac{-1+e^{c(a+bx)}}{\sqrt{2}}\right) + 2\sqrt{2} \operatorname{arctanh}\left(\frac{1+e^{c(a+bx)}}{\sqrt{2}}\right) + \log}{2bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTanh[Sinh[a*c + b*c*x]],x]`

output `(-2*E^(c*(a + b*x))*ArcTanh[1/(2*E^(c*(a + b*x))) - E^(c*(a + b*x))/2] - 2*  
*Sqrt[2]*ArcTanh[(-1 + E^(c*(a + b*x)))/Sqrt[2]] + 2*Sqrt[2]*ArcTanh[(1 +  
E^(c*(a + b*x)))/Sqrt[2]] + Log[1 - 2*E^(c*(a + b*x)) - E^(2*c*(a + b*x))]  
+ Log[1 + 2*E^(c*(a + b*x)) - E^(2*c*(a + b*x))])/(2*b*c)`

### Rubi [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.81, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {7281, 6829, 2720, 27, 1576, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac+bcx)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bx} \operatorname{arctanh}(\sinh(ac+bcx)) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{6829} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx)) - \int \frac{e^{ac+bcx} \cosh(ac+bcx)}{1-\sinh^2(ac+bcx)} d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx)) - \int -\frac{2e^{ac+bcx}(1+e^{2ac+2bcx})}{1-6e^{2ac+2bcx}+e^{4ac+4bcx}} de^{ac+bcx}}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \int \frac{e^{ac+bcx}(1+e^{2ac+2bcx})}{1-6e^{2ac+2bcx}+e^{4ac+4bcx}} de^{ac+bcx} + e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx))}{bc} \\
 & \quad \downarrow \text{1576} \\
 & \frac{\int \frac{1+e^{2ac+2bcx}}{1-5e^{2ac+2bcx}} de^{2ac+2bcx} + e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx))}{bc}
 \end{aligned}$$

$$\int \left( -\frac{1+\sqrt{2}}{2(-ac-bxc+2\sqrt{2}+3)} - \frac{1-\sqrt{2}}{2(-ac-bxc-2\sqrt{2}+3)} \right) de^{2ac+2bxc} + e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx))$$

$\downarrow$  1141  
 $bc$   
 $\downarrow$  2009

$$\frac{e^{ac+bcx} \operatorname{arctanh}(\sinh(ac+bcx)) + \frac{1}{2}(1-\sqrt{2}) \log(-ac-bcx-2\sqrt{2}+3) + \frac{1}{2}(1+\sqrt{2}) \log(-ac-bcx+2\sqrt{2}+3)}{bc}$$

input `Int[E^(c*(a + b*x))*ArcTanh[Sinh[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTanh[Sinh[a*c + b*c*x]] + ((1 - Sqrt[2])*Log[3 - 2*Sqrt[2] - a*c - b*c*x])/2 + ((1 + Sqrt[2])*Log[3 + 2*Sqrt[2] - a*c - b*c*x])/2)/(b*c)`

### Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1576 `Int[(x_)*((d_.) + (e_.)*(x_)^2)^(q_)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 6829

```
Int[((a_.) + ArcTanh[u]*(b_.))*(v_), x_Symbol] :=> With[{w = IntHide[v, x]}
, Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u,
x]/(1 - u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x
] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; F
reeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]
```

rule 7281

```
Int[u_, x_Symbol] :=> With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]]
  Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]
```

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 5.08 (sec) , antiderivative size = 868, normalized size of antiderivative = 7.75

method	result	size
risch	Expression too large to display	868

input

```
int(exp(c*(b*x+a))*arctanh(sinh(b*c*x+a*c)),x,method=_RETURNVERBOSE)
```

output

```

1/2/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1)+1/4*I/b/c*P
i*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))^3*exp(c*(
b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1))*csgn(I*
exp(-c*(b*x+a))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1))^2*exp(c*(b*x+a))+1/
4*I/b/c*Pi*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))*csgn(I*exp(-c*(b
*x+a))*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))^2*exp(c*(b*x+a))+1/2*I/b/c*
Pi*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))^2*exp(c*
(b*x+a))-1/4*I/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1))*csgn(I
*exp(-c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a)
)-1))*exp(c*(b*x+a))-1/2*I/b/c*exp(c*(b*x+a))*Pi-1/4*I/b/c*Pi*csgn(I*exp(-
c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1)
)^2*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+
1))*csgn(I*exp(-c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))+2*ex
p(c*(b*x+a))+1))*exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*exp(-c*(b*x+a))*(exp(2
*c*(b*x+a))+2*exp(c*(b*x+a))-1))^3*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*exp(
-c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1)
)^2*exp(c*(b*x+a))-1/2/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))-2*exp(c*(b*x+
a))-1)+1/2/b/c*ln(exp(2*c*(b*x+a))-(1+2^(1/2))^2)*2^(1/2)-1/2/b/c*ln(exp(2
*c*(b*x+a))-(2^(1/2)-1)^2)*2^(1/2)-2*a/b+1/2/b/c*ln(exp(2*c*(b*x+a))-(1+2^
(1/2))^2)+1/2/b/c*ln(exp(2*c*(b*x+a))-(2^(1/2)-1)^2)

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs.  $2(93) = 186$ .

Time = 0.11 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.09

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$$

$$= \frac{(\cosh(bcx + ac) + \sinh(bcx + ac)) \log\left(-\frac{\sinh(bcx+ac)+1}{\sinh(bcx+ac)-1}\right) + \sqrt{2} \log\left(\frac{3(2\sqrt{2}+3) \cosh(bcx+ac)^2 - 4(3\sqrt{2}+4) \cosh(bcx+ac)}{\cosh(bcx+ac)}\right)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(sinh(b*c*x+a*c)),x, algorithm="fricas")
```



output

```
1/2*((cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*log(-(sinh(b*c*x + a*c) + 1)/
(sinh(b*c*x + a*c) - 1)) + sqrt(2)*log((3*(2*sqrt(2) + 3)*cosh(b*c*x + a*c)
)^2 - 4*(3*sqrt(2) + 4)*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) + 3*(2*sqrt(2)
+ 3)*sinh(b*c*x + a*c)^2 - 2*sqrt(2) - 3)/(cosh(b*c*x + a*c)^2 + sinh(b*c
*x + a*c)^2 - 3)) + log(2*(cosh(b*c*x + a*c)^2 + sinh(b*c*x + a*c)^2 - 3)/
(cosh(b*c*x + a*c)^2 - 2*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) + sinh(b*c*x
+ a*c)^2)))/(b*c)
```

**Sympy [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx = \text{Timed out}$$

input

```
integrate(exp(c*(b*x+a))*atanh(sinh(b*c*x+a*c)),x)
```

output

Timed out

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.64

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx = \frac{\operatorname{artanh}(\sinh(bc x + ac)) e^{((bx+a)c)}}{bc} + \frac{\sqrt{2} \log\left(-\frac{\sqrt{2}-e^{(bcx+ac)+1}}{\sqrt{2}+e^{(bcx+ac)-1}}\right)}{2bc} - \frac{\sqrt{2} \log\left(-\frac{\sqrt{2}-e^{(bcx+ac)-1}}{\sqrt{2}+e^{(bcx+ac)+1}}\right)}{2bc} + \frac{\log(e^{(2bcx+2ac)} + 2e^{(bcx+ac)} - 1)}{2bc} + \frac{\log(e^{(2bcx+2ac)} - 2e^{(bcx+ac)} - 1)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(sinh(b*c*x+a*c)),x, algorithm="maxima")
```

output

```
arctanh(sinh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + 1/2*sqrt(2)*log(-(sqrt(2) - e^(b*c*x + a*c) + 1)/(sqrt(2) + e^(b*c*x + a*c) - 1))/(b*c) - 1/2*sqrt(2)*log(-(sqrt(2) - e^(b*c*x + a*c) - 1)/(sqrt(2) + e^(b*c*x + a*c) + 1))/(b*c) + 1/2*log(e^(2*b*c*x + 2*a*c) + 2*e^(b*c*x + a*c) - 1)/(b*c) + 1/2*log(e^(2*b*c*x + 2*a*c) - 2*e^(b*c*x + a*c) - 1)/(b*c)
```

**Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.40

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$$

$$= \frac{e^{((bx+a)c)} \log\left(-\frac{e^{(bcx+ac)} - e^{(-bcx-ac)} + 2}{e^{(bcx+ac)} - e^{(-bcx-ac)} - 2}\right)}{2bc}$$

$$+ \frac{\sqrt{2} \log\left(\left|\frac{-4\sqrt{2} + 2e^{(2bcx+2ac)} - 6}{4\sqrt{2} + 2e^{(2bcx+2ac)} - 6}\right|\right) + \log\left(|e^{(4bcx+4ac)} - 6e^{(2bcx+2ac)} + 1|\right)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(sinh(b*c*x+a*c)),x, algorithm="giac")
```

output

```
1/2*e^((b*x + a)*c)*log(-(e^(b*c*x + a*c) - e^(-b*c*x - a*c) + 2)/(e^(b*c*x + a*c) - e^(-b*c*x - a*c) - 2))/(b*c) + 1/2*(sqrt(2)*log(abs(-4*sqrt(2) + 2*e^(2*b*c*x + 2*a*c) - 6)/abs(4*sqrt(2) + 2*e^(2*b*c*x + 2*a*c) - 6)) + log(abs(e^(4*b*c*x + 4*a*c) - 6*e^(2*b*c*x + 2*a*c) + 1)))/(b*c)
```

**Mupad [B] (verification not implemented)**

Time = 4.10 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.60

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx$$

$$= \frac{e^{ac+bcx} \ln\left(\frac{e^{bcx} e^{ac}}{2} - \frac{e^{-bcx} e^{-ac}}{2} + 1\right)}{2bc} - \frac{e^{ac+bcx} \ln\left(\frac{e^{-bcx} e^{-ac}}{2} - \frac{e^{bcx} e^{ac}}{2} + 1\right)}{2bc}$$

$$+ \frac{\ln(6\sqrt{2}e^{2c(a+bx)} - 2\sqrt{2} - 8e^{2c(a+bx)}) (\sqrt{2} + 1)}{2bc}$$

$$- \frac{\ln(2\sqrt{2} - 8e^{2c(a+bx)} - 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} - 1)}{2bc}$$

input `int(exp(c*(a + b*x))*atanh(sinh(a*c + b*c*x)),x)`

output `(exp(a*c + b*c*x)*log((exp(b*c*x)*exp(a*c))/2 - (exp(-b*c*x)*exp(-a*c))/2 + 1))/(2*b*c) - (exp(a*c + b*c*x)*log((exp(-b*c*x)*exp(-a*c))/2 - (exp(b*c*x)*exp(a*c))/2 + 1))/(2*b*c) + (log(6*2^(1/2)*exp(2*c*(a + b*x)) - 2*2^(1/2) - 8*exp(2*c*(a + b*x)))*(2^(1/2) + 1))/(2*b*c) - (log(2*2^(1/2) - 8*exp(2*c*(a + b*x)) - 6*2^(1/2)*exp(2*c*(a + b*x)))*(2^(1/2) - 1))/(2*b*c)`

### Reduce [F]

$$\int e^{c(a+bx)} \operatorname{arctanh}(\sinh(ac + bcx)) dx = e^{ac} \left( \int e^{bcx} \operatorname{atanh}(\sinh(bcx + ac)) dx \right)$$

input `int(exp(c*(b*x+a))*atanh(sinh(b*c*x+a*c)),x)`

output `e**(a*c)*int(e**(b*c*x)*atanh(sinh(a*c + b*c*x)),x)`

### 3.356 $\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx$

Optimal result	2503
Mathematica [A] (verified)	2503
Rubi [A] (verified)	2504
Maple [C] (warning: unable to verify)	2505
Fricas [A] (verification not implemented)	2506
Sympy [F]	2507
Maxima [A] (verification not implemented)	2507
Giac [A] (verification not implemented)	2508
Mupad [B] (verification not implemented)	2508
Reduce [F]	2509

#### Optimal result

Integrand size = 20, antiderivative size = 49

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = \frac{e^{ac+bcx} \operatorname{arctanh}(\cosh(c(a + bx)))}{bc} + \frac{\log(1 - e^{2c(a+bx)})}{bc}$$

output

```
exp(b*c*x+a*c)*arctanh(cosh(c*(b*x+a)))/b/c+ln(1-exp(2*c*(b*x+a)))/b/c
```

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.22

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = \frac{e^{c(a+bx)} \operatorname{arctanh}\left(\frac{1}{2}e^{-c(a+bx)}(1 + e^{2c(a+bx)})\right) + \log(1 - e^{2c(a+bx)})}{bc}$$

input

```
Integrate[E^(c*(a + b*x))*ArcTanh[Cosh[a*c + b*c*x]],x]
```

output

```
(E^(c*(a + b*x))*ArcTanh[(1 + E^(2*c*(a + b*x)))/(2*E^(c*(a + b*x))]) + Log[1 - E^(2*c*(a + b*x))]/(b*c)
```

**Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {7281, 6829, 25, 2720, 27, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac+bcx)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx)) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{6829} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx)) - \int -e^{ac+bcx} \operatorname{csch}(ac+bcx) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int e^{ac+bcx} \operatorname{csch}(ac+bcx) d(ac+bcx) + e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx))}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int -\frac{2e^{ac+bcx}}{1-e^{2ac+2bcx}} de^{ac+bcx} + e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx))}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx)) - 2 \int \frac{e^{ac+bcx}}{1-e^{2ac+2bcx}} de^{ac+bcx}}{bc} \\
 & \quad \downarrow \text{240} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\cosh(ac+bcx)) + \log(1 - e^{2ac+2bcx})}{bc}
 \end{aligned}$$

input

```
Int[E^(c*(a + b*x))*ArcTanh[Cosh[a*c + b*c*x]], x]
```

output

```
(E^(a*c + b*c*x)*ArcTanh[Cosh[a*c + b*c*x]] + Log[1 - E^(2*a*c + 2*b*c*x)])/(b*c)
```

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]`
- rule 6829 `Int[((a_) + ArcTanh[u]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 - u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]`
- rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

## Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.77 (sec) , antiderivative size = 887, normalized size of antiderivative = 18.10

method	result	size
risch	Expression too large to display	887

input `int(exp(c*(b*x+a))*arctanh(cosh(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/b/c*\exp(c*(b*x+a))*\ln(\exp(c*(b*x+a))+1)-1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))+1))^2*csgn(I*(\exp(c*(b*x+a))+1)^2*\exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))))*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))-1)^2)^2*\exp(c*(b*x+a))+1/2*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))-1)^2)^2*\exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))))*csgn(I*(\exp(c*(b*x+a))-1)^2)*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))-1)^2)*\exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))+1)^2)^3*\exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))+1)^2)^3*\exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))))*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))+1)^2)^2*\exp(c*(b*x+a))-1/2*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))-1))*csgn(I*(\exp(c*(b*x+a))-1)^2)^2*\exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))+1)^2)*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))+1)^2)^2*\exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))))*csgn(I*(\exp(c*(b*x+a))+1)^2)*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))+1)^2)*\exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))-1)^2)^3*\exp(c*(b*x+a))-1/2*I/b/c*\exp(c*(b*x+a))*Pi-1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))-1)^2)*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))-1)^2)^2*\exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))-1))^2*csgn(I*(\exp(c*(b*x+a))-1)^2)*\exp(c*(b*x+a))+1/2*I/b/c*Pi*csgn(I*(\exp(c*(b*x+a))+1))*csgn(I*(\exp(c*(b*x+a))+1)^2)^2*\exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*\exp(-c*(b*x+a))*(\exp(c*(b*x+a))-1)^2)^3*\exp(c*(b*x+a))-1/b/c*\exp(c*(b*x+a))*\ln(\exp(c*(b*x+a))-1)-2*a/b+1/b/c*\ln(-1+ex...$$

### Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.90

$$\begin{aligned} & \int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac+bcx)) dx \\ & = \frac{(\cosh(bcx+ac) + \sinh(bcx+ac)) \log\left(-\frac{\cosh(bcx+ac)+1}{\cosh(bcx+ac)-1}\right) + 2 \log\left(\frac{2 \sinh(bcx+ac)}{\cosh(bcx+ac)-\sinh(bcx+ac)}\right)}{2bc} \end{aligned}$$

input `integrate(exp(c*(b*x+a))*arctanh(cosh(b*c*x+a*c)),x, algorithm="fricas")`

output  $1/2*((\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c))*\log(-(\cosh(b*c*x + a*c) + 1)/(\cosh(b*c*x + a*c) - 1)) + 2*\log(2*\sinh(b*c*x + a*c)/(\cosh(b*c*x + a*c) - \sinh(b*c*x + a*c))))/(b*c)$

### Sympy [F]

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atanh}(\cosh(ac + bcx)) dx$$

input `integrate(exp(c*(b*x+a))*atanh(cosh(b*c*x+a*c)),x)`

output `exp(a*c)*Integral(exp(b*c*x)*atanh(cosh(a*c + b*c*x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.31

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = \frac{\operatorname{artanh}(\cosh(bcx + ac)) e^{((bx+a)c)}}{bc} + \frac{\log(e^{(bcx+ac)} + 1)}{bc} + \frac{\log(e^{(bcx+ac)} - 1)}{bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(cosh(b*c*x+a*c)),x, algorithm="maxima")`

output `arctanh(cosh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + log(e^(b*c*x + a*c) + 1)/(b*c) + log(e^(b*c*x + a*c) - 1)/(b*c)`



**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.92

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx$$

$$= \frac{\left( e^{(bcx)} \log \left( -\frac{e^{(2bcx+2ac)} + 2e^{(bcx+ac)} + 1}{e^{(2bcx+2ac)} - 2e^{(bcx+ac)} + 1} \right) + 2e^{(-ac)} \log \left( |e^{(2bcx+2ac)} - 1| \right) \right) e^{(ac)}}{2bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(cosh(b*c*x+a*c)),x, algorithm="giac")`

output `1/2*(e^(b*c*x)*log(-(e^(2*b*c*x + 2*a*c) + 2*e^(b*c*x + a*c) + 1)/(e^(2*b*c*x + 2*a*c) - 2*e^(b*c*x + a*c) + 1)) + 2*e^(-a*c)*log(abs(e^(2*b*c*x + 2*a*c) - 1)))*e^(a*c)/(b*c)`

**Mupad [B] (verification not implemented)**

Time = 3.93 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.27

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = \frac{\ln(e^{2bcx} e^{2ac} - 1)}{bc}$$

$$- \frac{e^{bcx} e^{ac} \ln \left( 1 - \frac{e^{-bcx} e^{-ac}}{2} - \frac{e^{bcx} e^{ac}}{2} \right)}{2bc}$$

$$+ \frac{e^{bcx} e^{ac} \ln \left( \frac{e^{bcx} e^{ac}}{2} + \frac{e^{-bcx} e^{-ac}}{2} + 1 \right)}{2bc}$$

input `int(exp(c*(a + b*x))*atanh(cosh(a*c + b*c*x)),x)`

output `log(exp(2*b*c*x)*exp(2*a*c) - 1)/(b*c) - (exp(b*c*x)*exp(a*c)*log(1 - (exp(-b*c*x)*exp(-a*c))/2 - (exp(b*c*x)*exp(a*c))/2))/(2*b*c) + (exp(b*c*x)*exp(a*c)*log((exp(b*c*x)*exp(a*c))/2 + (exp(-b*c*x)*exp(-a*c))/2 + 1))/(2*b*c)`

**Reduce [F]**

$$\int e^{c(a+bx)} \operatorname{arctanh}(\cosh(ac + bcx)) dx = e^{ac} \left( \int e^{bcx} \operatorname{atanh}(\cosh(bc x + ac)) dx \right)$$

input `int(exp(c*(b*x+a))*atanh(cosh(b*c*x+a*c)),x)`

output `e**(a*c)*int(e**(b*c*x)*atanh(cosh(a*c + b*c*x)),x)`

### 3.357 $\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx$

Optimal result	2510
Mathematica [A] (verified)	2510
Rubi [A] (verified)	2511
Maple [A] (verified)	2512
Fricas [A] (verification not implemented)	2512
Sympy [A] (verification not implemented)	2513
Maxima [A] (verification not implemented)	2513
Giac [A] (verification not implemented)	2514
Mupad [B] (verification not implemented)	2514
Reduce [B] (verification not implemented)	2514

#### Optimal result

Integrand size = 20, antiderivative size = 45

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = -\frac{e^{ac+bcx}}{bc} + \frac{e^{ac+bcx} \operatorname{arctanh}(\tanh(c(a + bx)))}{bc}$$

output `-exp(b*c*x+a*c)/b/c+exp(b*c*x+a*c)*arctanh(tanh(c*(b*x+a)))/b/c`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = \frac{e^{c(a+bx)} \left( -1 + \operatorname{arctanh} \left( \frac{-1 + e^{2c(a+bx)}}{1 + e^{2c(a+bx)}} \right) \right)}{bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTanh[Tanh[a*c + b*c*x]],x]`

output `(E^(c*(a + b*x))*(-1 + ArcTanh[(-1 + E^(2*c*(a + b*x))]/(1 + E^(2*c*(a + b*x)))))/(b*c)`

**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {7281, 6829, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac+bcx)) dx \\
 \downarrow 7281 \\
 \frac{\int e^{ac+bcx} \operatorname{arctanh}(\tanh(ac+bcx)) d(ac+bcx)}{bc} \\
 \downarrow 6829 \\
 \frac{e^{ac+bcx} \operatorname{arctanh}(\tanh(ac+bcx)) - \int e^{ac+bcx} d(ac+bcx)}{bc} \\
 \downarrow 2624 \\
 \frac{e^{ac+bcx} \operatorname{arctanh}(\tanh(ac+bcx)) - e^{ac+bcx}}{bc}
 \end{array}$$

input `Int[E^(c*(a + b*x))*ArcTanh[Tanh[a*c + b*c*x]],x]`

output `(-E^(a*c + b*c*x) + E^(a*c + b*c*x)*ArcTanh[Tanh[a*c + b*c*x]])/(b*c)`

**Defintions of rubi rules used**

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`  
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 6829

```
Int[((a_.) + ArcTanh[u]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}
, Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u,
x]/(1 - u^2)), x], x] /; InverseFunctionFreeQ[w, x]] /; FreeQ[{a, b}, x
] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.) /; F
reeQ[{c, d, m}, x]] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]
```

rule 7281

```
Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]]
Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]
```

### Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

method	result
paralrelrisch	$-\frac{-e^{c(bx+a)} \operatorname{arctanh}(\tanh(c(bx+a))) + e^{c(bx+a)}}{bc}$
default	$\frac{e^{bcx+ac}(bcx+ac) - e^{bcx+ac} + e^{bcx+ac}(\operatorname{arctanh}(\tanh(bcx+ac)) - bcx - ac)}{bc}$
risch	$\frac{e^{c(bx+a)} \ln(e^{c(bx+a)})}{bc} - \frac{i \left( \pi \operatorname{csgn} \left( \frac{i}{1+e^{2c(bx+a)}} \right) \operatorname{csgn}(ie^{2c(bx+a)}) \operatorname{csgn} \left( \frac{ie^{2c(bx+a)}}{1+e^{2c(bx+a)}} \right) - \pi \operatorname{csgn} \left( \frac{i}{1+e^{2c(bx+a)}} \right) \operatorname{csgn} \left( \frac{ie^{2c(bx+a)}}{1+e^{2c(bx+a)}} \right)}{bc}$

input

```
int(exp(c*(b*x+a))*arctanh(tanh(b*c*x+a*c)), x, method=_RETURNVERBOSE)
```

output

```
-(-exp(c*(b*x+a))*arctanh(tanh(c*(b*x+a)))+exp(c*(b*x+a)))/b/c
```

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx$$

$$= \frac{(bcx + ac - 1) \cosh(bc x + ac) + (bcx + ac - 1) \sinh(bc x + ac)}{bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(tanh(b*c*x+a*c)), x, algorithm="fricas")
```

output 
$$\frac{((b*c*x + a*c - 1)*\cosh(b*c*x + a*c) + (b*c*x + a*c - 1)*\sinh(b*c*x + a*c))}{(b*c)}$$

### Sympy [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.24

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx$$

$$= \begin{cases} 0 & \text{for } b = 0 \wedge c = 0 \\ x e^{ac} \operatorname{atanh}(\tanh(ac)) & \text{for } b = 0 \\ 0 & \text{for } c = 0 \\ \frac{e^{ac} e^{bcx} \operatorname{atanh}(\tanh(ac+bcx))}{bc} - \frac{e^{ac} e^{bcx}}{bc} & \text{otherwise} \end{cases}$$

input `integrate(exp(c*(b*x+a))*atanh(tanh(b*c*x+a*c)),x)`

output `Piecewise((0, Eq(b, 0) & Eq(c, 0)), (x*exp(a*c)*atanh(tanh(a*c)), Eq(b, 0)), (0, Eq(c, 0)), (exp(a*c)*exp(b*c*x)*atanh(tanh(a*c + b*c*x))/(b*c) - exp(a*c)*exp(b*c*x)/(b*c), True))`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = \frac{\operatorname{artanh}(\tanh(bcx + ac)) e^{((bx+a)c)}}{bc} - \frac{e^{(bcx+ac)}}{bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(tanh(b*c*x+a*c)),x, algorithm="maxima")`

output `arctanh(tanh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - e^(b*c*x + a*c)/(b*c)`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = \frac{(b^2c^2x + abc^2 - bc)e^{(bcx+ac)}}{b^2c^2}$$

input `integrate(exp(c*(b*x+a))*arctanh(tanh(b*c*x+a*c)),x, algorithm="giac")`

output `(b^2*c^2*x + a*b*c^2 - b*c)*e^(b*c*x + a*c)/(b^2*c^2)`

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.62

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = \frac{e^{ac+bcx} (\operatorname{atanh}(\tanh(ac + bcx)) - 1)}{bc}$$

input `int(exp(c*(a + b*x))*atanh(tanh(a*c + b*c*x)),x)`

output `(exp(a*c + b*c*x)*(atanh(tanh(a*c + b*c*x)) - 1))/(b*c)`

**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.64

$$\int e^{c(a+bx)} \operatorname{arctanh}(\tanh(ac + bcx)) dx = \frac{e^{bcx+ac} (\operatorname{atanh}(\tanh(bcx + ac)) - 1)}{bc}$$

input `int(exp(c*(b*x+a))*atanh(tanh(b*c*x+a*c)),x)`

output `(e**(a*c + b*c*x)*(atanh(tanh(a*c + b*c*x)) - 1))/(b*c)`

### 3.358 $\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx$

Optimal result	2515
Mathematica [A] (verified)	2515
Rubi [A] (verified)	2516
Maple [A] (verified)	2517
Fricas [C] (verification not implemented)	2517
Sympy [F]	2518
Maxima [A] (verification not implemented)	2518
Giac [A] (verification not implemented)	2518
Mupad [B] (verification not implemented)	2519
Reduce [B] (verification not implemented)	2519

#### Optimal result

Integrand size = 20, antiderivative size = 45

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = -\frac{e^{ac+bcx}}{bc} + \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{coth}(c(a + bx)))}{bc}$$

output `-exp(b*c*x+a*c)/b/c+exp(b*c*x+a*c)*arctanh(coth(c*(b*x+a)))/b/c`

#### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = \frac{e^{c(a+bx)} \left( -1 + \operatorname{arctanh} \left( \frac{1+e^{2c(a+bx)}}{-1+e^{2c(a+bx)}} \right) \right)}{bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTanh[Coth[a*c + b*c*x]],x]`

output `(E^(c*(a + b*x))*(-1 + ArcTanh[(1 + E^(2*c*(a + b*x)))/(-1 + E^(2*c*(a + b*x))]))/(b*c)`



**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.91, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$ , Rules used = {7281, 6829, 2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx \\
 \downarrow 7281 \\
 \frac{\int e^{ac+bcx} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) d(ac + bcx)}{bc} \\
 \downarrow 6829 \\
 \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) - \int e^{ac+bcx} d(ac + bcx)}{bc} \\
 \downarrow 2624 \\
 \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) - e^{ac+bcx}}{bc}
 \end{array}$$

input `Int[E^(c*(a + b*x))*ArcTanh[Coth[a*c + b*c*x]],x]`

output `(-E^(a*c + b*c*x) + E^(a*c + b*c*x)*ArcTanh[Coth[a*c + b*c*x]])/(b*c)`

**Defintions of rubi rules used**

rule 2624 `Int[((F_)^(v_))^(n_.), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`  
`FreeQ[{F, n}, x] && LinearQ[v, x]`

rule 6829 `Int[((a_.) + ArcTanh[u]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]},  
Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u,  
x]/(1 - u^2)), x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x  
&& InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.) /; F  
reeQ[{c, d, m}, x]] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]]  
Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

### Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.60

method	result
paralelrisch	$\frac{e^{c(bx+a)}(-1+\operatorname{arctanh}(\operatorname{coth}(c(bx+a))))}{bc}$
risch	$\frac{e^{c(bx+a)} \ln(e^{c(bx+a)})}{bc} - \frac{i \left( -2\pi \operatorname{csgn} \left( \frac{i}{-1+e^{2c(bx+a)}} \right)^2 + \pi \operatorname{csgn}(ie^{c(bx+a)})^2 \operatorname{csgn}(ie^{2c(bx+a)}) - 2\pi \operatorname{csgn}(ie^{c(bx+a)}) \operatorname{csgn}(ie^{2c(bx+a)}) \right)}{bc}$

input `int(exp(c*(b*x+a))*arctanh(coth(b*c*x+a*c)), x, method=_RETURNVERBOSE)`

output `exp(c*(b*x+a))*(-1+arctanh(coth(c*(b*x+a))))/b/c`

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.27

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx$$

$$= \frac{(i\pi + 2bcx + 2ac - 2) \cosh(bcx + ac) + (i\pi + 2bcx + 2ac - 2) \sinh(bcx + ac)}{2bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(coth(b*c*x+a*c)), x, algorithm="fricas")`

output  $1/2*((I*pi + 2*b*c*x + 2*a*c - 2)*cosh(b*c*x + a*c) + (I*pi + 2*b*c*x + 2*a*c - 2)*sinh(b*c*x + a*c))/(b*c)$

### Sympy [F]

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atanh}(\operatorname{coth}(ac + bcx)) dx$$

input `integrate(exp(c*(b*x+a))*atanh(coth(b*c*x+a*c)),x)`

output `exp(a*c)*Integral(exp(b*c*x)*atanh(coth(a*c + b*c*x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = \frac{\operatorname{artanh}(\operatorname{coth}(bcx + ac)) e^{((bx+a)c)}}{bc} - \frac{e^{(bcx+ac)}}{bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(coth(b*c*x+a*c)),x, algorithm="maxima")`

output `arctanh(coth(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - e^(b*c*x + a*c)/(b*c)`

### Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = \frac{(e^{(bcx)} \log(-e^{(2bcx+2ac)}) - 2e^{(bcx)})e^{(ac)}}{2bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(coth(b*c*x+a*c)),x, algorithm="giac")`

output `1/2*(e^(b*c*x)*log(-e^(2*b*c*x + 2*a*c)) - 2*e^(b*c*x))*e^(a*c)/(b*c)`

**Mupad [B] (verification not implemented)**

Time = 3.62 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.62

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = \frac{e^{ac+bcx} (\operatorname{atanh}(\operatorname{coth}(ac + bcx)) - 1)}{bc}$$

input `int(exp(c*(a + b*x))*atanh(coth(a*c + b*c*x)),x)`output `(exp(a*c + b*c*x)*(atanh(coth(a*c + b*c*x)) - 1))/(b*c)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.64

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{coth}(ac + bcx)) dx = \frac{e^{bcx+ac} (\operatorname{atanh}(\operatorname{coth}(bcx + ac)) - 1)}{bc}$$

input `int(exp(c*(b*x+a))*atanh(coth(b*c*x+a*c)),x)`output `(e**(a*c + b*c*x)*(atanh(coth(a*c + b*c*x)) - 1))/(b*c)`

### 3.359 $\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx$

Optimal result	2520
Mathematica [A] (verified)	2520
Rubi [A] (verified)	2521
Maple [C] (warning: unable to verify)	2522
Fricas [A] (verification not implemented)	2523
Sympy [F(-1)]	2524
Maxima [A] (verification not implemented)	2524
Giac [B] (verification not implemented)	2525
Mupad [B] (verification not implemented)	2525
Reduce [F]	2526

#### Optimal result

Integrand size = 20, antiderivative size = 49

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(c(a + bx)))}{bc} + \frac{\log(1 - e^{2c(a+bx)})}{bc}$$

output `exp(b*c*x+a*c)*arctanh(sech(c*(b*x+a)))/b/c+ln(1-exp(2*c*(b*x+a)))/b/c`

#### Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.20

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \frac{e^{c(a+bx)} \operatorname{arctanh}\left(\frac{2e^{c(a+bx)}}{1+e^{2c(a+bx)}}\right) + \log(1 - e^{2c(a+bx)})}{bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTanh[Sech[a*c + b*c*x]],x]`

output `(E^(c*(a + b*x))*ArcTanh[(2*E^(c*(a + b*x)))/(1 + E^(2*c*(a + b*x))]) + Log[1 - E^(2*c*(a + b*x))]/(b*c)`

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {7281, 6829, 25, 2720, 27, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac+bcx)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx)) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{6829} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx)) - \int -e^{ac+bcx} \operatorname{csch}(ac+bcx) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int e^{ac+bcx} \operatorname{csch}(ac+bcx) d(ac+bcx) + e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx))}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int -\frac{2e^{ac+bcx}}{1-e^{2ac+2bcx}} de^{ac+bcx} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx))}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx)) - 2 \int \frac{e^{ac+bcx}}{1-e^{2ac+2bcx}} de^{ac+bcx}}{bc} \\
 & \quad \downarrow \text{240} \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{sech}(ac+bcx)) + \log(1 - e^{2ac+2bcx})}{bc}
 \end{aligned}$$

input

$$\text{Int}[E^{(c*(a + b*x))*ArcTanh[Sech[a*c + b*c*x]], x]$$

output

$$\frac{(E^{(a*c + b*c*x))*ArcTanh[Sech[a*c + b*c*x]] + \text{Log}[1 - E^{(2*a*c + 2*b*c*x)}]}{(b*c)}$$

### Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]`
- rule 6829 `Int[((a_) + ArcTanh[u]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 - u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]`
- rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.72 (sec) , antiderivative size = 872, normalized size of antiderivative = 17.80

method	result	size
risch	Expression too large to display	872

input `int(exp(c*(b*x+a))*arctanh(sech(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))+1))^2*csgn(I*(exp(c*(b*x+a))+1)^2)*ex \\
 & p(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I/(1+exp(2*c*(b*x+a))))*csgn(I*(exp(c*(b*x+ \\
 & a))-1)^2/(1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))-1/2*I/b/c*Pi*csgn(I*(exp(c \\
 & *(b*x+a))-1))*csgn(I*(exp(c*(b*x+a))-1)^2)^2*exp(c*(b*x+a))+1/4*I/b/c*Pi*c \\
 & sgn(I*(exp(c*(b*x+a))+1)^2)*csgn(I*(exp(c*(b*x+a))+1)^2/(1+exp(2*c*(b*x+a) \\
 & )))^2*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))-1))^2*csgn(I*(exp \\
 & (c*(b*x+a))-1)^2)*exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))+1)^2) \\
 & ^3*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))-1)^2)*csgn(I/(1+exp( \\
 & 2*c*(b*x+a))))*csgn(I*(exp(c*(b*x+a))-1)^2/(1+exp(2*c*(b*x+a))))*exp(c*(b* \\
 & x+a))+1/4*I/b/c*Pi*csgn(I/(1+exp(2*c*(b*x+a))))*csgn(I*(exp(c*(b*x+a))+1)^ \\
 & 2/(1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a) \\
 & ))-1)^2/(1+exp(2*c*(b*x+a))))^3*exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*(exp(c* \\
 & (b*x+a))+1)^2/(1+exp(2*c*(b*x+a))))^3*exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*( \\
 & exp(c*(b*x+a))-1)^2)*csgn(I*(exp(c*(b*x+a))-1)^2/(1+exp(2*c*(b*x+a))))^2*e \\
 & xp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))-1)^2)^3*exp(c*(b*x+a))+1 \\
 & /2*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))+1))*csgn(I*(exp(c*(b*x+a))+1)^2)^2*exp( \\
 & c*(b*x+a))-1/4*I/b/c*Pi*csgn(I*(exp(c*(b*x+a))+1)^2)*csgn(I/(1+exp(2*c*(b* \\
 & x+a))))*csgn(I*(exp(c*(b*x+a))+1)^2/(1+exp(2*c*(b*x+a))))*exp(c*(b*x+a))-1 \\
 & /b/c*exp(c*(b*x+a))*ln(exp(c*(b*x+a))-1)+1/b/c*exp(c*(b*x+a))*ln(exp(c*(b* \\
 & x+a))+1)-2*a/b+1/b/c*ln(-1+exp(2*c*(b*x+a)))
 \end{aligned}$$

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.88

$$\begin{aligned}
 & \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx \\
 & = \frac{(\cosh(bcx + ac) + \sinh(bcx + ac)) \log\left(\frac{\cosh(bcx+ac)+1}{\cosh(bcx+ac)-1}\right) + 2 \log\left(\frac{2 \sinh(bcx+ac)}{\cosh(bcx+ac)-\sinh(bcx+ac)}\right)}{2bc}
 \end{aligned}$$

input `integrate(exp(c*(b*x+a))*arctanh(sech(b*c*x+a*c)),x, algorithm="fricas")`



output

```
1/2*((cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*log((cosh(b*c*x + a*c) + 1)/(
cosh(b*c*x + a*c) - 1)) + 2*log(2*sinh(b*c*x + a*c)/(cosh(b*c*x + a*c) - s
inh(b*c*x + a*c))))/(b*c)
```

**Sympy [F(-1)]**

Timed out.

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \text{Timed out}$$

input

```
integrate(exp(c*(b*x+a))*atanh(sech(b*c*x+a*c)),x)
```

output

Timed out

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.31

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \frac{\operatorname{artanh}(\operatorname{sech}(bcx + ac)) e^{((bx+a)c)}}{bc} + \frac{\log(e^{(bcx+ac)} + 1)}{bc} + \frac{\log(e^{(bcx+ac)} - 1)}{bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(sech(b*c*x+a*c)),x, algorithm="maxima")
```

output

```
arctanh(sech(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + log(e^(b*c*x + a*c) + 1
)/(b*c) + log(e^(b*c*x + a*c) - 1)/(b*c)
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 98 vs.  $2(47) = 94$ .

Time = 0.20 (sec) , antiderivative size = 98, normalized size of antiderivative = 2.00

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \frac{e^{((bx+a)c)} \log\left(-\frac{e^{(bcx+ac)} + e^{(-bcx-ac)} + 1}{e^{(bcx+ac)} + e^{(-bcx-ac)} - 1}\right)}{2bc} + \frac{\log(|e^{(2bcx+2ac)} - 1|)}{bc}$$

input `integrate(exp(c*(b*x+a))*arctanh(sech(b*c*x+a*c)),x, algorithm="giac")`

output `1/2*e^((b*x + a)*c)*log(-(2/(e^(b*c*x + a*c) + e^(-b*c*x - a*c)) + 1)/(2/(e^(b*c*x + a*c) + e^(-b*c*x - a*c)) - 1))/(b*c) + log(abs(e^(2*b*c*x + 2*a*c) - 1))/(b*c)`

**Mupad [B] (verification not implemented)**

Time = 4.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.43

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = \frac{\ln(e^{2bcx} e^{2ac} - 1)}{bc} - \frac{e^{ac+bcx} \ln\left(1 - \frac{1}{\frac{e^{bcx} e^{ac}}{2} + \frac{e^{-bcx} e^{-ac}}{2}}}\right)}{2bc} + \frac{\ln\left(\frac{1}{\frac{e^{bcx} e^{ac}}{2} + \frac{e^{-bcx} e^{-ac}}{2}} + 1\right) e^{ac+bcx}}{2bc}$$

input `int(atanh(1/cosh(a*c + b*c*x))*exp(c*(a + b*x)),x)`

output `log(exp(2*b*c*x)*exp(2*a*c) - 1)/(b*c) - (exp(a*c + b*c*x)*log(1 - 1/((exp(b*c*x)*exp(a*c))/2 + (exp(-b*c*x)*exp(-a*c))/2)))/(2*b*c) + (log(1/((exp(b*c*x)*exp(a*c))/2 + (exp(-b*c*x)*exp(-a*c))/2) + 1)*exp(a*c + b*c*x))/(2*b*c)`

**Reduce [F]**

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{sech}(ac + bcx)) dx = e^{ac} \left( \int e^{bcx} \operatorname{atanh}(\operatorname{sech}(bcx + ac)) dx \right)$$

input `int(exp(c*(b*x+a))*atanh(sech(b*c*x+a*c)),x)`

output `e**(a*c)*int(e**(b*c*x)*atanh(sech(a*c + b*c*x)),x)`

### 3.360 $\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx$

Optimal result	2527
Mathematica [A] (verified)	2527
Rubi [A] (warning: unable to verify)	2528
Maple [C] (warning: unable to verify)	2530
Fricas [B] (verification not implemented)	2531
Sympy [F]	2532
Maxima [B] (verification not implemented)	2532
Giac [A] (verification not implemented)	2533
Mupad [B] (verification not implemented)	2534
Reduce [F]	2534

#### Optimal result

Integrand size = 20, antiderivative size = 107

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx = \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(c(a + bx)))}{bc} + \frac{(1 - \sqrt{2}) \log(3 - 2\sqrt{2} - e^{2c(a+bx)})}{2bc} + \frac{(1 + \sqrt{2}) \log(3 + 2\sqrt{2} - e^{2c(a+bx)})}{2bc}$$

output

```
exp(b*c*x+a*c)*arctanh(csch(c*(b*x+a)))/b/c+1/2*(1-2^(1/2))*ln(3-2*2^(1/2)
-exp(2*c*(b*x+a)))/b/c+1/2*(1+2^(1/2))*ln(3+2*2^(1/2)-exp(2*c*(b*x+a)))/b/
c
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.40

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx = \frac{-2\sqrt{2} \operatorname{arctanh}\left(\frac{-1+e^{c(a+bx)}}{\sqrt{2}}\right) + 2\sqrt{2} \operatorname{arctanh}\left(\frac{1+e^{c(a+bx)}}{\sqrt{2}}\right) + 2e^{c(a+bx)} \operatorname{arctanh}\left(\frac{2e^{c(a+bx)}}{-1+e^{2c(a+bx)}}\right) + \log(1 - 2e^{c(a+bx)})}{2bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTanh[Csch[a*c + b*c*x]],x]`

output `(-2*Sqrt[2]*ArcTanh[(-1 + E^(c*(a + b*x)))/Sqrt[2]] + 2*Sqrt[2]*ArcTanh[(1 + E^(c*(a + b*x)))/Sqrt[2]] + 2*E^(c*(a + b*x))*ArcTanh[(2*E^(c*(a + b*x)))/(-1 + E^(2*c*(a + b*x)))] + Log[1 - 2*E^(c*(a + b*x)) - E^(2*c*(a + b*x))] + Log[1 + 2*E^(c*(a + b*x)) - E^(2*c*(a + b*x))]/(2*b*c)`

### Rubi [A] (warning: unable to verify)

Time = 0.50 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.85, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {7281, 6829, 25, 2720, 27, 1576, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) dx \\
 & \quad \downarrow 7281 \\
 & \int e^{ac+bx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) d(ac+bcx) \\
 & \quad \downarrow 6829 \\
 & \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) - \int -\frac{e^{ac+bcx} \operatorname{coth}(ac+bcx) \operatorname{csch}(ac+bcx)}{1-\operatorname{csch}^2(ac+bcx)} d(ac+bcx)}{bc} \\
 & \quad \downarrow 25 \\
 & \int \frac{e^{ac+bcx} \operatorname{coth}(ac+bcx) \operatorname{csch}(ac+bcx)}{1-\operatorname{csch}^2(ac+bcx)} d(ac+bcx) + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) \\
 & \quad \downarrow 2720 \\
 & \int \frac{2e^{ac+bcx}(1+e^{2ac+2bcx})}{1-6e^{2ac+2bcx}+e^{4ac+4bcx}} de^{ac+bcx} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) \\
 & \quad \downarrow 27 \\
 & \frac{2 \int \frac{e^{ac+bcx}(1+e^{2ac+2bcx})}{1-6e^{2ac+2bcx}+e^{4ac+4bcx}} de^{ac+bcx} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx))}{bc}
 \end{aligned}$$

$$\begin{array}{c}
 \int \frac{1+e^{2ac+2bxc}}{1-5e^{2ac+2bxc}} de^{2ac+2bxc} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) \\
 \downarrow 1576 \\
 \frac{\int \left( -\frac{1+\sqrt{2}}{2(-ac-bxc+2\sqrt{2}+3)} - \frac{1-\sqrt{2}}{2(-ac-bxc-2\sqrt{2}+3)} \right) de^{2ac+2bxc} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx))}{bc} \\
 \downarrow 1141 \\
 \frac{\int \left( -\frac{1+\sqrt{2}}{2(-ac-bxc+2\sqrt{2}+3)} - \frac{1-\sqrt{2}}{2(-ac-bxc-2\sqrt{2}+3)} \right) de^{2ac+2bxc} + e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx))}{bc} \\
 \downarrow 2009 \\
 \frac{e^{ac+bcx} \operatorname{arctanh}(\operatorname{csch}(ac+bcx)) + \frac{1}{2}(1-\sqrt{2}) \log(-ac-bcx-2\sqrt{2}+3) + \frac{1}{2}(1+\sqrt{2}) \log(-ac-bcx+2\sqrt{2}+3)}{bc}
 \end{array}$$

input `Int[E^(c*(a + b*x))*ArcTanh[Csch[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTanh[Csch[a*c + b*c*x]] + ((1 - Sqrt[2])*Log[3 - 2*Sqrt[2] - a*c - b*c*x])/2 + ((1 + Sqrt[2])*Log[3 + 2*Sqrt[2] - a*c - b*c*x])/2)/(b*c)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1141 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 6829 `Int[((a_) + ArcTanh[u]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTanh[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 - u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTanh[u]), x]]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 4.47 (sec) , antiderivative size = 842, normalized size of antiderivative = 7.87

method	result	size
risch	Expression too large to display	842

input `int(exp(c*(b*x+a))*arctanh(csch(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

```

1/2/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1)+1/4*I/b/c*P
i*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))*csgn(I/(-1+exp(2*c*(b*x+a)
))))*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1)/(-1+exp(2*c*(b*x+a))))*
exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1))
*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1)/(-1+exp(2*c*(b*x+a))))^2*ex
p(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I/(-1+exp(2*c*(b*x+a))))*csgn(I*(-exp(2*c*(
b*x+a))+2*exp(c*(b*x+a))+1)/(-1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))-1/4*I/
b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1))*csgn(I/(-1+exp(2*c*(b
*x+a))))*csgn(I/(-1+exp(2*c*(b*x+a)))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1
))*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I/(-1+exp(2*c*(b*x+a))))*csgn(I/(-1+ex
p(2*c*(b*x+a)))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1))^2*exp(c*(b*x+a))-1/
4*I/b/c*Pi*csgn(I*(-exp(2*c*(b*x+a))+2*exp(c*(b*x+a))+1)/(-1+exp(2*c*(b*x+
a))))^3*exp(c*(b*x+a))+1/4*I/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+2*exp(c*(b*x+
a))-1))*csgn(I/(-1+exp(2*c*(b*x+a)))*(exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-1
))^2*exp(c*(b*x+a))-1/4*I/b/c*Pi*csgn(I/(-1+exp(2*c*(b*x+a)))*(exp(2*c*(b*x
+a))+2*exp(c*(b*x+a))-1))^3*exp(c*(b*x+a))+1/2/b/c*ln(exp(2*c*(b*x+a))-(1+
2^(1/2))^2)*2^(1/2)-1/2/b/c*ln(exp(2*c*(b*x+a))-(2^(1/2)-1)^2)*2^(1/2)-1/2
/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))-2*exp(c*(b*x+a))-1)-2*a/b+1/2/b/c*
ln(exp(2*c*(b*x+a))-(1+2^(1/2))^2)+1/2/b/c*ln(exp(2*c*(b*x+a))-(2^(1/2)-1)
^2)

```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 233 vs. 2(90) = 180.

Time = 0.11 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.18

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx$$

$$= \frac{(\cosh(bcx + ac) + \sinh(bcx + ac)) \log\left(\frac{\sinh(bcx+ac)+1}{\sinh(bcx+ac)-1}\right) + \sqrt{2} \log\left(\frac{3(2\sqrt{2}+3) \cosh(bcx+ac)^2 - 4(3\sqrt{2}+4) \cosh(bcx+ac)}{\cosh(bcx+ac)^2}\right)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(csch(b*c*x+a*c)),x, algorithm="fricas")
```



output

```
1/2*((cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*log((sinh(b*c*x + a*c) + 1)/(sinh(b*c*x + a*c) - 1)) + sqrt(2)*log((3*(2*sqrt(2) + 3)*cosh(b*c*x + a*c)^2 - 4*(3*sqrt(2) + 4)*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) + 3*(2*sqrt(2) + 3)*sinh(b*c*x + a*c)^2 - 2*sqrt(2) - 3)/(cosh(b*c*x + a*c)^2 + sinh(b*c*x + a*c)^2 - 3)) + log(2*(cosh(b*c*x + a*c)^2 + sinh(b*c*x + a*c)^2 - 3)/(cosh(b*c*x + a*c)^2 - 2*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) + sinh(b*c*x + a*c)^2)))/(b*c)
```

**Sympy [F]**

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atanh}(\operatorname{csch}(ac + bcx)) dx$$

input

```
integrate(exp(c*(b*x+a))*atanh(csch(b*c*x+a*c)),x)
```

output

```
exp(a*c)*Integral(exp(b*c*x)*atanh(csch(a*c + b*c*x)), x)
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 184 vs.  $2(90) = 180$ .

Time = 0.12 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.72

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx = \frac{\operatorname{artanh}(\operatorname{csch}(bcx + ac)) e^{(bx+a)c}}{bc} + \frac{\sqrt{2} \log\left(-\frac{\sqrt{2}-e^{(bcx+ac)+1}}{\sqrt{2}+e^{(bcx+ac)-1}}\right)}{2bc} - \frac{\sqrt{2} \log\left(-\frac{\sqrt{2}-e^{(bcx+ac)-1}}{\sqrt{2}+e^{(bcx+ac)+1}}\right)}{2bc} + \frac{\log(e^{(2bcx+2ac)} + 2e^{(bcx+ac)} - 1)}{2bc} + \frac{\log(e^{(2bcx+2ac)} - 2e^{(bcx+ac)} - 1)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctanh(csch(b*c*x+a*c)),x, algorithm="maxima")
```

output

```

arctanh(csch(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + 1/2*sqrt(2)*log(-(sqrt(
2) - e^(b*c*x + a*c) + 1)/(sqrt(2) + e^(b*c*x + a*c) - 1))/(b*c) - 1/2*sqrt
(2)*log(-(sqrt(2) - e^(b*c*x + a*c) - 1)/(sqrt(2) + e^(b*c*x + a*c) + 1))
/(b*c) + 1/2*log(e^(2*b*c*x + 2*a*c) + 2*e^(b*c*x + a*c) - 1)/(b*c) + 1/2*
log(e^(2*b*c*x + 2*a*c) - 2*e^(b*c*x + a*c) - 1)/(b*c)

```

**Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.56

$$\begin{aligned}
& \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx \\
&= \frac{e^{((bx+a)c)} \log\left(-\frac{\frac{e^{(bcx+ac)} - e^{(-bcx-ac)}}{2} + 1}{\frac{e^{(bcx+ac)} - e^{(-bcx-ac)}}{2} - 1}\right)}{2bc} \\
&+ \frac{\sqrt{2} \log\left(\frac{|-4\sqrt{2} + 2e^{(2bcx+2ac)} - 6|}{|4\sqrt{2} + 2e^{(2bcx+2ac)} - 6|}\right) + \log(|e^{(4bcx+4ac)} - 6e^{(2bcx+2ac)} + 1|)}{2bc}
\end{aligned}$$

input

```

integrate(exp(c*(b*x+a))*arctanh(csch(b*c*x+a*c)),x, algorithm="giac")

```

output

```

1/2*e^((b*x + a)*c)*log(-(2/(e^(b*c*x + a*c) - e^(-b*c*x - a*c)) + 1)/(2/(
e^(b*c*x + a*c) - e^(-b*c*x - a*c)) - 1))/(b*c) + 1/2*(sqrt(2)*log(abs(-4*
sqrt(2) + 2*e^(2*b*c*x + 2*a*c) - 6)/abs(4*sqrt(2) + 2*e^(2*b*c*x + 2*a*c)
- 6)) + log(abs(e^(4*b*c*x + 4*a*c) - 6*e^(2*b*c*x + 2*a*c) + 1)))/(b*c)

```

**Mupad [B] (verification not implemented)**

Time = 4.01 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.75

$$\begin{aligned}
& \int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx \\
&= \frac{\ln(6\sqrt{2}e^{2c(a+bx)} - 2\sqrt{2} - 8e^{2c(a+bx)}) (\sqrt{2} + 1)}{2bc} \\
&\quad - \frac{e^{ac+bcx} \ln\left(1 - \frac{\frac{e^{bcx}e^{ac}}{2} - \frac{1}{e^{-bcx}e^{-ac}}}{2}\right)}{2bc} \\
&\quad - \frac{\ln(2\sqrt{2} - 8e^{2c(a+bx)} - 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} - 1)}{2bc} \\
&\quad + \frac{\ln\left(\frac{\frac{e^{bcx}e^{ac}}{2} - \frac{1}{e^{-bcx}e^{-ac}}}{2} + 1\right) e^{ac+bcx}}{2bc}
\end{aligned}$$

input `int(atanh(1/sinh(a*c + b*c*x))*exp(c*(a + b*x)),x)`output `(log(6*2^(1/2)*exp(2*c*(a + b*x)) - 2*2^(1/2) - 8*exp(2*c*(a + b*x)))*(2^(1/2) + 1)/(2*b*c) - (exp(a*c + b*c*x)*log(1 - 1/((exp(b*c*x)*exp(a*c))/2 - (exp(-b*c*x)*exp(-a*c))/2)))/(2*b*c) - (log(2*2^(1/2) - 8*exp(2*c*(a + b*x)) - 6*2^(1/2)*exp(2*c*(a + b*x)))*(2^(1/2) - 1))/(2*b*c) + (log(1/((exp(b*c*x)*exp(a*c))/2 - (exp(-b*c*x)*exp(-a*c))/2) + 1)*exp(a*c + b*c*x))/(2*b*c)`**Reduce [F]**

$$\int e^{c(a+bx)} \operatorname{arctanh}(\operatorname{csch}(ac + bcx)) dx = e^{ac} \left( \int e^{bcx} \operatorname{atanh}(\operatorname{csch}(bcx + ac)) dx \right)$$

input `int(exp(c*(b*x+a))*atanh(csch(b*c*x+a*c)),x)`output `e**(a*c)*int(e**(b*c*x)*atanh(csch(a*c + b*c*x)),x)`

**3.361** 
$$\int \frac{(a+b\operatorname{arctanh}(cx^n))(d+e\log(fx^m))}{x} dx$$

Optimal result	2535
Mathematica [C] (verified)	2536
Rubi [A] (verified)	2536
Maple [C] (warning: unable to verify)	2537
Fricas [B] (verification not implemented)	2538
Sympy [F(-1)]	2539
Maxima [F]	2539
Giac [F]	2539
Mupad [F(-1)]	2540
Reduce [F]	2540

**Optimal result**

Integrand size = 24, antiderivative size = 110

$$\begin{aligned} & \int \frac{(a + b\operatorname{arctanh}(cx^n))(d + e\log(fx^m))}{x} dx \\ &= \frac{a(d + e\log(fx^m))^2}{2em} - \frac{b(d + e\log(fx^m))\operatorname{PolyLog}(2, -cx^n)}{2n} \\ & \quad + \frac{b(d + e\log(fx^m))\operatorname{PolyLog}(2, cx^n)}{2n} \\ & \quad + \frac{bem\operatorname{PolyLog}(3, -cx^n)}{2n^2} - \frac{bem\operatorname{PolyLog}(3, cx^n)}{2n^2} \end{aligned}$$

output

```
1/2*a*(d+e*ln(f*x^m))^2/e/m-1/2*b*(d+e*ln(f*x^m))*polylog(2,-c*x^n)/n+1/2*
b*(d+e*ln(f*x^m))*polylog(2,c*x^n)/n+1/2*b*e*m*polylog(3,-c*x^n)/n^2-1/2*b
*e*m*polylog(3,c*x^n)/n^2
```

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.18 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.04

$$\int \frac{(a + \operatorname{barctanh}(cx^n))(d + e \log(fx^m))}{x} dx$$

$$= -\frac{bcemx^n {}_4F_3\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}, \frac{3}{2}; c^2x^{2n}\right)}{n^2}$$

$$+ \frac{bcx^n {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}; c^2x^{2n}\right)(d + e \log(fx^m))}{n}$$

$$+ \frac{1}{2}a \log(x)(2d - em \log(x) + 2e \log(fx^m))$$

input

```
Integrate[((a + b*ArcTanh[c*x^n])*(d + e*Log[f*x^m]))/x,x]
```

output

```
-((b*c*e*m*x^n*HypergeometricPFQ[{1/2, 1/2, 1/2, 1}, {3/2, 3/2, 3/2}, c^2*x^(2*n)])/n^2) + (b*c*x^n*HypergeometricPFQ[{1/2, 1/2, 1}, {3/2, 3/2}, c^2*x^(2*n)]*(d + e*Log[f*x^m]))/n + (a*Log[x]*(2*d - e*m*Log[x] + 2*e*Log[f*x^m]))/2
```

**Rubi [A] (verified)**

Time = 0.75 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.24, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$ , Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barctanh}(cx^n))(d + e \log(fx^m))}{x} dx$$

$$\downarrow 7293$$

$$\int \left( \frac{d(a + \operatorname{barctanh}(cx^n))}{x} + \frac{e \log(fx^m)(a + \operatorname{barctanh}(cx^n))}{x} \right) dx$$

$$\downarrow 2009$$

$$\frac{ad \log(x) + \frac{ae \log^2(fx^m)}{2m} - \frac{bd \text{PolyLog}(2, -cx^n)}{2n} + \frac{bd \text{PolyLog}(2, cx^n)}{2n} - \frac{be \text{PolyLog}(2, -cx^n) \log(fx^m)}{2n} + \frac{be \text{PolyLog}(2, cx^n) \log(fx^m)}{2n} + \frac{bem \text{PolyLog}(3, -cx^n)}{2n^2} - \frac{bem \text{PolyLog}(3, cx^n)}{2n^2}}{2n}$$

input `Int[((a + b*ArcTanh[c*x^n])*(d + e*Log[f*x^m]))/x,x]`

output `a*d*Log[x] + (a*e*Log[f*x^m]^2)/(2*m) - (b*d*PolyLog[2, -(c*x^n)])/(2*n) - (b*e*Log[f*x^m]*PolyLog[2, -(c*x^n)])/(2*n) + (b*d*PolyLog[2, c*x^n]/(2*n) + (b*e*Log[f*x^m]*PolyLog[2, c*x^n]/(2*n) + (b*e*m*PolyLog[3, -(c*x^n)])/(2*n^2) - (b*e*m*PolyLog[3, c*x^n]/(2*n^2)`

### Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

### Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 179.31 (sec) , antiderivative size = 327, normalized size of antiderivative = 2.97

method	result
risch	$\frac{\left(\frac{i\pi \operatorname{csgn}(ix^m) \operatorname{csgn}(ifx^m)^2}{4} - \frac{i\pi \operatorname{csgn}(ix^m) \operatorname{csgn}(ifx^m) \operatorname{csgn}(if)}{4} - \frac{i\pi \operatorname{csgn}(ifx^m)^3}{4} + \frac{i\pi \operatorname{csgn}(ifx^m)^2 \operatorname{csgn}(if)}{4} + \frac{e \ln(f) + \frac{d}{2}}{2}\right) (-b \operatorname{dilog}}{n}$

input `int((a+b*arctanh(c*x^n))*(d+e*ln(f*x^m))/x,x,method=_RETURNVERBOSE)`

output

```
(1/4*I*e*Pi*csgn(I*x^m)*csgn(I*f*x^m)^2-1/4*I*e*Pi*csgn(I*x^m)*csgn(I*f*x^
m)*csgn(I*f)-1/4*I*e*Pi*csgn(I*f*x^m)^3+1/4*I*e*Pi*csgn(I*f*x^m)^2*csgn(I*
f)+1/2*e*ln(f)+1/2*d)/n*(-b*dilog(c*x^n+1)+2*a*ln(x^n)+b*dilog(1-c*x^n))-1
/2*e*b*m/n*ln(x)*polylog(2,-c*x^n)+1/2*b*e*m*polylog(3,-c*x^n)/n^2+1/2*e*b
/n*dilog(c*x^n+1)*m*ln(x)-1/2*e*b/n*dilog(c*x^n+1)*ln(x^m)+1/2*e*a/m*ln(x^
m)^2+1/2*e*b*m/n*ln(x)*polylog(2,c*x^n)-1/2*b*e*m*polylog(3,c*x^n)/n^2+1/2
*e*b/n*ln(1-c*x^n)*ln(c*x^n)*m*ln(x)-1/2*e*b/n*ln(1-c*x^n)*ln(c*x^n)*ln(x^
m)+1/2*e*b/n*dilog(c*x^n)*m*ln(x)-1/2*e*b/n*dilog(c*x^n)*ln(x^m)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 327 vs.  $2(98) = 196$ .

Time = 0.10 (sec) , antiderivative size = 327, normalized size of antiderivative = 2.97

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx$$

$$= \frac{2 a e m n^2 \log(x)^2 - 2 b e m \operatorname{polylog}(3, c \cosh(n \log(x)) + c \sinh(n \log(x))) + 2 b e m \operatorname{polylog}(3, -c \cosh(n \log(x)) - c \sinh(n \log(x))) + 2*(b*e*m*n*\log(x) + b*e*n*\log(f) + b*d*n)*\operatorname{dilog}(c*\cosh(n*\log(x)) + c*\sinh(n*\log(x))) - 2*(b*e*m*n*\log(x) + b*e*n*\log(f) + b*d*n)*\operatorname{dilog}(-c*\cosh(n*\log(x)) - c*\sinh(n*\log(x))) - (b*e*m*n^2*\log(x)^2 + 2*(b*e*n^2*\log(f) + b*d*n^2)*\log(x))*\log(c*\cosh(n*\log(x)) + c*\sinh(n*\log(x)) + 1) + (b*e*m*n^2*\log(x)^2 + 2*(b*e*n^2*\log(f) + b*d*n^2)*\log(x))*\log(-c*\cosh(n*\log(x)) - c*\sinh(n*\log(x)) + 1) + 4*(a*e*n^2*\log(f) + a*d*n^2)*\log(x) + (b*e*m*n^2*\log(x)^2 + 2*(b*e*n^2*\log(f) + b*d*n^2)*\log(x))*\log(-(c*\cosh(n*\log(x)) + c*\sinh(n*\log(x)) + 1))/(c*\cosh(n*\log(x)) + c*\sinh(n*\log(x)) - 1)))/n^2}{}$$

input

```
integrate((a+b*arctanh(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="fricas")
```

output

```
1/4*(2*a*e*m*n^2*log(x)^2 - 2*b*e*m*polylog(3, c*cosh(n*log(x)) + c*sinh(n
*log(x))) + 2*b*e*m*polylog(3, -c*cosh(n*log(x)) - c*sinh(n*log(x))) + 2*(
b*e*m*n*log(x) + b*e*n*log(f) + b*d*n)*dilog(c*cosh(n*log(x)) + c*sinh(n*l
og(x))) - 2*(b*e*m*n*log(x) + b*e*n*log(f) + b*d*n)*dilog(-c*cosh(n*log(x)
) - c*sinh(n*log(x))) - (b*e*m*n^2*log(x)^2 + 2*(b*e*n^2*log(f) + b*d*n^2)
*log(x))*log(c*cosh(n*log(x)) + c*sinh(n*log(x)) + 1) + (b*e*m*n^2*log(x)^
2 + 2*(b*e*n^2*log(f) + b*d*n^2)*log(x))*log(-c*cosh(n*log(x)) - c*sinh(n*
log(x)) + 1) + 4*(a*e*n^2*log(f) + a*d*n^2)*log(x) + (b*e*m*n^2*log(x)^2 +
2*(b*e*n^2*log(f) + b*d*n^2)*log(x))*log(-(c*cosh(n*log(x)) + c*sinh(n*lo
g(x)) + 1))/(c*cosh(n*log(x)) + c*sinh(n*log(x)) - 1)))/n^2
```

**Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx = \text{Timed out}$$

input `integrate((a+b*atanh(c*x**n))*(d+e*ln(f*x**m))/x,x)`

output Timed out

**Maxima [F]**

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx \\ &= \int \frac{(b \operatorname{arctanh}(cx^n) + a)(e \log(fx^m) + d)}{x} dx \end{aligned}$$

input `integrate((a+b*arctanh(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="maxima")`

output `1/2*a*e*log(f*x^m)^2/m + a*d*log(x) - 1/4*(b*e*m*log(x)^2 - 2*b*e*log(x)*log(x^m) - 2*(e*log(f) + d)*b*log(x))*log(c*x^n + 1) + 1/4*(b*e*m*log(x)^2 - 2*b*e*log(x)*log(x^m) - 2*(e*log(f) + d)*b*log(x))*log(-c*x^n + 1) + integrate(1/2*(2*b*c*e*n*x^n*log(x)*log(x^m) - (b*c*e*m*n*log(x))^2 - 2*(e*n*log(f) + d*n)*b*c*log(x))*x^n)/(c^2*x*x^(2*n) - x), x)`

**Giac [F]**

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx \\ &= \int \frac{(b \operatorname{arctanh}(cx^n) + a)(e \log(fx^m) + d)}{x} dx \end{aligned}$$

input `integrate((a+b*arctanh(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="giac")`



output `integrate((b*arctanh(c*x^n) + a)*(e*log(f*x^m) + d)/x, x)`

### Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx = \int \frac{(a + b \operatorname{atanh}(cx^n))(d + e \ln(fx^m))}{x} dx$$

input `int(((a + b*atanh(c*x^n))*(d + e*log(f*x^m)))/x,x)`

output `int(((a + b*atanh(c*x^n))*(d + e*log(f*x^m)))/x, x)`

### Reduce [F]

$$\begin{aligned} & \int \frac{(a + b \operatorname{arctanh}(cx^n))(d + e \log(fx^m))}{x} dx \\ &= \frac{2 \left( \int \frac{\operatorname{atanh}(x^n c)}{x} dx \right) b d m + 2 \left( \int \frac{\operatorname{atanh}(x^n c) \log(x^m f)}{x} dx \right) b e m + \log(x^m f)^2 a e + 2 \log(x) a d m}{2m} \end{aligned}$$

input `int((a+b*atanh(c*x^n))*(d+e*log(f*x^m))/x,x)`

output `(2*int(atanh(x**n*c)/x,x)*b*d*m + 2*int((atanh(x**n*c)*log(x**m*f))/x,x)*b  
*e*m + log(x**m*f)**2*a*e + 2*log(x)*a*d*m)/(2*m)`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	2541
4.2	Links to plain text integration problems used in this report for each CAS .	2559

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
If [AppellFunctionQ [Head [expn]],
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
If [Head [expn] === RootSum,
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
If [Head [expn] === Integrate || Head [expn] === Int,
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
9]]]]]]]]]]]
```

```
ElementaryFunctionQ [func_] :=
```

```
MemberQ [ {
Exp, Log,
Sin, Cos, Tan, Cot, Sec, Csc,
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
```

```
}, func]
```

```
SpecialFunctionQ [func_] :=
```

```
MemberQ [ {
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
```

```
}, func]
```

```
HypergeometricFunctionQ [func_] :=
```

```
MemberQ [ {Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ [func_] :=
```

```
MemberQ [ {AppellF1}, func]
```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#                    if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#                    see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```



```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```



```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result/leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file